

## New York State Immunization Schedule for Pneumococcal Vaccine

**Pneumococcal Conjugate Vaccine (PCV13):** required for Pre-k and day care entrance/attendance for children born on or after 1/1/08

### Vaccine Schedule for Children Starting the Series on Time at 2 Months of Age

2 Months*	4 Months**	6 Months**	12-15 Months***
1 dose	1 dose	1 dose	1 dose

\*May be given as early as 6 weeks of age.

\*\*The minimum interval between the first and second doses, and the second and third doses, is 4 weeks.

\*\*\*Should be administered at least 8 weeks after the third dose, and not before 12 months of age.

### Vaccine Schedule for Children Starting the Series Between 3 and 59 Months of Age

Age:	Required Doses of Vaccine for Pre-k and Day Care Entrance/Attendance
3-6 months of age	3 doses, 4-8 weeks apart and a 4 <sup>th</sup> dose at 12-15 months
7-11 Months of Age	3 doses*
12-23 Months of Age	2 doses**
≥ 24 – 59 Months of Age	1 dose
*Two doses at least 4 weeks apart; third dose after 12 months of age, and at least 2 months after the second dose.	
**Two doses at least 8 weeks apart.	

### Recommended PCV Transition Schedule

**Recommended transition schedule from 7-valent pneumococcal conjugate vaccine (PCV7) to 13-valent vaccine (PCV13) vaccination among infants and children, according to number of previous PCV7 doses received --- Advisory Committee on Immunization Practices (ACIP), United States, 2010**

Infant series			Booster dose	Supplemental PCV13 dose
2 mos	4 mos	6 mos	≥12 mos*	14—59 mos**
PCV7	PCV13	PCV13	PCV13	---
PCV7	PCV7	PCV13	PCV13	---
PCV7	PCV7	PCV7	PCV13	---
PCV7	PCV7	PCV7	PCV7	PCV13

\*No additional PCV13 doses are indicated for children age 12—23 months who have received 2 or 3 doses of PCV before age 12 months and at least 1 dose of PCV13 at ≥12 months.

\*\*For children with underlying medical conditions, a single supplemental PCV13 dose is recommended through age 71 months.