



Sugar-Sweetened Beverages and the Link between Obesity and Cancer

Sugar-sweetened beverage consumption is linked to weight gain, overweight and obesity.

- ◆ A 2007 report by the World Cancer Research Fund and American Institute for Cancer Research concluded that “the epidemiological and mechanistic evidence that drinks containing added sugars, including sucrose and high-fructose corn syrup, are a cause of weight gain, overweight, and obesity. Like energy-dense foods and drinks, they have this effect by promoting excess energy intake¹.”
- ◆ A 2007 meta-analysis on the effects of sugar-sweetened beverage consumption concluded that²:
 - ◇ Individuals do not reduce their calorie intake from other foods for the extra calories in these beverages and therefore are consuming more calories overall.
 - ◇ Individuals who consume these beverages drink less milk and have less overall calcium consumption.
 - ◇ Consumption of these beverages is related to higher intake of carbohydrates, lower intakes of fruit and dietary fiber and lower intakes of a variety of macronutrients.
- ◆ One study concluded that women who consumed one or more servings of a soft drink per day were twice as likely to develop diabetes as women who consumed less than one serving a month³.

Sugar-sweetened beverages accounted for 13 percent of total daily caloric intake in adolescents in 2004⁴.

Overweight and obesity contribute to 14 to 20 percent of all cancer-related deaths in the U.S.⁵

- ◆ Overweight and obesity are associated with increased risk for several common cancers, including colon, esophagus, kidney, endometrial and breast cancer in postmenopausal women⁶.
 - ◇ Obese men and women have a two times greater risk of esophageal cancer than healthy weight individuals⁷.
 - ◇ Obese, postmenopausal women have a 1.5 times greater risk of breast cancer and greater risk of actually dying from their cancer than healthy weight women⁸.
 - ◇ Obese women have a 2 to 4 times greater risk of endometrial cancer than healthy weight women⁹.
- ◆ There is also suggestive evidence that overweight and obesity are associated with increased risk for other cancers, including the pancreas, aggressive prostate, gall bladder, thyroid, ovary, cervix, and for multiple myeloma and Hodgkin's disease¹⁰.
- ◆ The biological link between overweight and obesity to cancer is believed to be related to multiple effects on fat and sugar metabolism, immune function, hormone levels and proteins that affect hormone levels, and other factors related to cell proliferation and growth¹¹.

The Centers for Disease Control and Prevention, the U.S. Surgeon General, the Institute of Medicine, the World Cancer Research Fund and the American Institute for Cancer Research all recommend reducing consumption of sugar-sweetened beverages, such as soft drinks, as a critical strategy to reverse the epidemic of overweight and obesity in our nation, especially among our children¹⁴.

Obesity is a major public health crisis in the U.S. with nearly two-thirds of adults and 18 percent of teens overweight or obese¹². The health care costs associated with obesity are approximately \$147 billion each year¹³.

The American Cancer Society Cancer Action Network (ACS CAN) supports a broad range of evidence-based strategies to prevent and reduce obesity, foster healthy behaviors and remove barriers to healthy living in all populations in order to reduce cancer incidence and death in this country.

ACS CAN supports evidenced-based efforts that reduce the consumption of sugar-sweetened beverages and improves the overall nutritional quality of the foods and beverages Americans consume. In particular, ACS CAN supports the removal of sugar-sweetened beverages and other low-nutrient foods out of schools.

ACS CAN supports efforts that improve a community's ability to promote good nutrition and physical activity by providing adequate resources, building a public health infrastructure, providing easy to understand, culturally competent information about healthy living, and increasing access to and availability of healthy foods and safe and convenient places to be physically active. Behavioral research is critical, as is coverage for prevention services for all.

1. World Cancer Research Fund/American Institute for Cancer Research. Food, Nutrition, Physical Activity, and the Prevention of Cancer: a Global Perspective. Washington DC: AICR, 2007.
2. Vartanian L, Schwartz M and Brownell K. "Effects of Soft Drink Consumption on Nutrition and Health: A Systematic Review and Meta-analysis." American Journal of Public Health, 97(4): 667-675, April 2007. Malik V, Schulze M and Hu F. "Intake of Sugar-sweetened Beverages and Weight Gain: A systematic Review," American Journal of Clinical Nutrition, 84(2): 274-288, August 2006. Pereira M. "The Possible Role of Sugar-sweetened Beverages in Obesity Etiology: A review of the Evidence." International Journal of Obesity, 30(S3): S28-S36, December 2006. Forshee R, Anderson P, Storey M. "Sugar-sweetened beverages and Body Mass Index in Children and Adolescents: a Meta-Analysis." American Journal of Clinical Nutrition, 87(6): 1662-1671, June 2008. Striegel-Moore RH, Thompson D, Affenito SG, et al. Correlates of beverage intake in adolescent girls: The National Heart, Lung, and Blood Institute Growth and Health Study. Journal of Pediatrics 2006; 148(2): 183-187. Phillips S, Bandini L, Naumova E. et al. Energy-dense snack food intake in adolescence: longitudinal relationship to weight and fatness. Obesity Research 2004; 12(3): 461-472. Berkey C, Rockett H, Field A, Gillman M, Golditz G. Sugar-added beverages and adolescent weight change. Obesity Research 2004; 12(5): 778-788. Ludwig D, Peterson K, Gortmaker S. Relation between consumption of sugar-sweetened drinks and childhood obesity: a prospective, observational analysis. The Lancet 2001; 357(9255): 505-508.
3. Schulze M, Mason J, Ludwig D, Et al. Sugar-sweetened beverages, weight gain, and incidence of type 2 diabetes in young and middle-aged women. Journal of the American Medical Association 2004; 292(8): 927-934.
4. Wang Y, Bleich S and Gortmaker S. "Increasing Caloric Contribution from Sugar-sweetened Beverages and 100% Fruit Juices among US Children and Adolescents, 1988-2004." Pediatrics, 121(6): e1604-1614, June 2008.
5. Cokkinides V, Bandi P, Siegel R, Ward EM, Thun MJ. *Cancer Prevention & Early Detection Facts and Figures 2008*. Atlanta, GA: American Cancer Society, 2007. Vainio H, Bianchini F. Weight control and physical activity, vol. 6. Lyon, France: International Agency for Cancer Research Press; 2002. Calle EE, Rodriguez C, Walker-Thurmond K, Thun MJ. Overweight, obesity and mortality from cancer in a prospectively studied cohort of US adults. N Engl J Med 2003; 348: 1625-1638.
6. Cokkinides V, Bandi P, Siegel R, Ward EM, Thun MJ. *Cancer Prevention & Early Detection Facts and Figures 2008*. Atlanta, GA: American Cancer Society, 2007. Vainio H, Bianchini F. Weight control and physical activity, vol. 6. Lyon, France: International Agency for Cancer Research Press; 2002. Calle EE, Rodriguez C, Walker-Thurmond K, Thun MJ. Overweight, obesity and mortality from cancer in a prospectively studied cohort of US adults. N Engl J Med 2003; 348: 1625-1638
7. Vainio H, Bianchini F. Weight control and physical activity, vol. 6. Lyon, France: International Agency for Cancer Research Press; 2002. Brown LM, Swanson CA, Gridley G, et al. Adenocarcinoma of the esophagus: Role of obesity and diet. Journal of the National Cancer Institute 1995; 87(2): 104-109. Chow WH, Blot WJ, Vaughan TL, et al. Body mass index and risk of adenocarcinomas of the esophagus and gastric cardia. Journal of the National Cancer Institute 1998; 90(2): 150-155. Li SD, Mobarhan S. Association between body mass index and adenocarcinoma of the esophagus and gastric cardia. Nutrition Reviews 2000; 58(2 Pt1): 54-56. Lagergren J, Bergstrom R, Nyren O. Association between body mass and adenocarcinoma of the esophagus and gastric cardia. Annals of Internal Medicine 1999; 130(11): 883-890.
8. Bianchini F, Kaaks R, Vainio H. Overweight, obesity, and cancer risk. The Lancet Oncology 2002; 3(9): 565-574. American Cancer Society. *Cancer Facts and Figures 2008*. Atlanta, GA: American Cancer Society; 2008. Radimer KL, Ballard-Barbash R, Miller JS, et al. Weight change and the risk of late-onset breast cancer in the original Framingham cohort. Nutr Cancer 2004; 49:7-13. Trentham-Dietz A, Newcomb PA, Egan KM, et al. Weight change and risk of postmenopausal breast cancer (United States). Cancer Causes Control 2000; 11: 533-542. Carmichael AR, Bates T. Obesity and breast cancer: a review of the literature. Breast 2004; 13: 85-92. Stephenson GD, Rose DP. Breast cancer and obesity: an update. Nutr Cancer 2003; 45: 1-16. Swerdlow AJ, De Stavola BL, Floderus B, et al. Risk factors for breast cancer at young ages in twins: an international population-based study. J Natl Cancer Inst 2002; 94: 1238-1246. van den Brandt PA, Spiegelman D, Yaun SS, et al. Pooled analysis of prospective cohort studies on height, weight, and breast cancer risk. Am J Epidemiol 2000; 152: 514-527. Morimoto LM, White E, Chen Z, et al. Obesity, body size, and risk of postmenopausal breast cancer: the Women's Health Initiative (United States). Cancer Causes Control 2002; 13: 741-751. Feigelson HS, Patel A, Teras LR, et al. Adult weight gain and histopathologic characteristics of breast cancer among postmenopausal women. Cancer 2006; 107: 12-21. Huang Z, Hankinson SE, Cloditz GA, et al. Dual effects of weight and weight gain on breast cancer risk. Journal of the American Medical Association 1997; 278(17): 1407-1411. Trentham-Dietz A, Newcomb PA, Storer BE, et al. Body size and risk of breast cancer. American Journal of Epidemiology 1997; 145(11): 1011-1019. World Cancer Research Fund/American Institute for Cancer Research. Food, Nutrition, Physical Activity, and the Prevention of Cancer: A Global Perspective. Washington DC: AICR, 2007. Vainio H, Bianchini F. Weight control and physical activity, vol. 6. Lyon, France: International Agency for Cancer Research Press; 2002. Huang Z, Hankinson SE, Cloditz GA, et al. Dual effects of weight and weight gain on breast cancer risk. Journal of the American Medical Association 1997; 278(17): 1407-1411. Lahmann PH, Lissner L, Gullberg B, Olsson H, Berglund G. A prospective study of adiposity and postmenopausal breast cancer risk: The Malmo Diet and Cancer Study. International Journal of Cancer 2003; 103(2): 246-252. Yoo KY, Tajima K, Park S, et al. Postmenopausal obesity as a breast cancer risk factor according to estrogen and progesterone receptor status (Japan). Cancer Letters 2001; 167(1): 57-63. Tretli S. Height and weight in relation to breast cancer morbidity and mortality. A prospective study of 570,000 women in Norway. International Journal of Cancer 1989; 44(1): 23-30. Petrelli JM, Calle EE, Rodriguez C, Thun MJ. Body mass index, height, and postmenopausal breast cancer mortality in a prospective cohort of U.S. women. Cancer Causes and Control 2002; 13(4): 325-332. Cui Y, Whitman MK, Flaws JA, et al. Body mass and stage of breast cancer at diagnosis. International Journal of Cancer 2002; 98(2): 279-283.
9. Vainio H, Bianchini F. Weight control and physical activity, vol. 6. Lyon, France: International Agency for Cancer Research Press; 2002. Amant F, Moerman P, Neven P, et al. Endometrial cancer. Lancet 2005; 366: 491-505.

10. Cokkinides V, Bandi P, Siegel R, Ward EM, Thun MJ. *Cancer Prevention & Early Detection Facts and Figures 2008*. Atlanta, GA: American Cancer Society, 2007. Vainio H, Bianchini F. Weight control and physical activity, vol. 6. Lyon, France: International Agency for Cancer Research Press; 2002. Calle EE, Rodriguez C, Walker-Thurmond K, Thun MJ. Overweight, obesity and mortality from cancer in a prospectively studied cohort of US adults. *N Engl J Med* 2003; 348: 1625-1638.
11. Cokkinides V, Bandi P, Siegel R, Ward EM, Thun MJ. *Cancer Prevention & Early Detection Facts and Figures 2008*. Atlanta, GA: American Cancer Society, 2007. Vainio H, Bianchini F. Weight control and physical activity, vol. 6. Lyon, France: International Agency for Cancer Research Press; 2002. Calle EE, Rodriguez C, Walker-Thurmond K, Thun MJ. Overweight, obesity and mortality from cancer in a prospectively studied cohort of US adults. *N Engl J Med* 2003; 348: 1625-1638.
12. Flegal KM, Carroll MD, Ogden CL, et al. Prevalence and Trends in Obesity Among US Adults, 1999-2008. *JAMA*. 2010; 303(3): 235-241. Ogden CL, Carroll MD, Curtin LR, et al. Prevalence of High Body Mass Index in US Children and Adolescents, 2007-2008. *JAMA*. 2010; 303(3): 242-249.
13. Finkelstein EA, Trogon JG, Cohen JW, Dietz W. Annual Medical Spending Attributable to Obesity: Payer- and Service-Specific Estimates. *Health Affairs*. 2009; 28(5): w822-w831.
14. CDC. Recommended Community Strategies and Measurements to Prevent Obesity in the United States. *MMWR*. July 24, 2009; 58 (RR-7). U.S. Department of Health and Human Services. The Surgeon General's Vision for a Healthy and Fit Nation. Rockville, MD: U.S. Department of Health and Human Services, Office of the Surgeon General, January 2010. U.S. Department of Health and Human Services. The Surgeon General's Call to action to prevent and decrease overweight and obesity. Rockville, MD: U.S. Department of Health and Human Service, Office of the Surgeon General, 2001. Institute of Medicine and National Research Council. 2009. Local Government Actions to Prevent Childhood Obesity. Washington, DC: The National Academies Press. Institute of Medicine. 2005. Prevention Childhood Obesity: Health in the Balance. Washington, DC: The National Academies Press. World Cancer Research Fund/American Institute for Cancer Research. Food, Nutrition, Physical Activity, and the Prevention of Cancer: a Global Perspective. Washington DC: AICR, 2007. World Cancer Research Fund/American Institute for Cancer Research. Policy and Action for Cancer Prevention. Food, Nutrition, and Physical Activity: a Global Perspective. Washington DC: AICR, 2009.