



New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008



All Invasive Malignant Tumors



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008



Oral cavity and pharynx



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008



Esophagus



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008



Stomach



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008



Colorectal



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008



Liver / intrahepatic bile duct



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008



Pancreas



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008



Larynx



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008



Lung and bronchus



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008


Melanoma of the skin



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008


Female breast



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008


Cervix uteri



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008


Corpus uterus and NOS



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008


Ovary



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008


Prostate



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008



Testis



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008



Urinary bladder (incl. in situ)



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008



Kidney and renal pelvis



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008



Brain and other nervous system



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008



Thyroid



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008



Hodgkin lymphoma



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008



Non-Hodgkin lymphomas



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008



Multiple myeloma



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'

New York State Cancer Registry, NYSDOH Age-Adjusted Cancer Incidence and Mortality Rates, New York State, 1976-2008

Leukemias



Rates are age-adjusted to the 2000 US population standard.
 Regression lines are calculated using the Joinpoint Regression Program Version 3.4.3, April 2010, National Cancer Institute (<http://srab.cancer.gov/joinpoint/>).
 The data shown here are not identical to the data in the corresponding 2004-2008 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time.'