

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence for Selected Areas, by Gender, 2005-2009
Nassau County**

| Site of cancer | Males | | | Females | | |
|---------------------------------|---------------|------------------|--------------|---------------|------------------|--------------|
| | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) |
| All Invasive Malignant Tumors | 4221.4 | 588.0 | 8.0 | 4224.2 | 475.5 | 6.6 |
| Oral cavity and pharynx | 94.4 | 12.8 | 1.2 | 55.4 | 6.3 | 0.8 |
| Esophagus | 54.2 | 7.6 | 0.9 | 15.8 | 1.6 | 0.4 |
| Stomach | 67.2 | 9.4 | 1.0 | 49.8 | 5.4 | 0.7 |
| Colorectal | 368.8 | 51.7 | 2.4 | 382.8 | 39.9 | 1.8 |
| Colon excluding rect | 257.0 | 36.3 | 2.0 | 286.4 | 29.5 | 1.6 |
| Rectum & rectosigmoi | 111.8 | 15.5 | 1.3 | 96.4 | 10.4 | 0.9 |
| Liver / intrahepatic bile duct | 76.6 | 10.4 | 1.0 | 29.2 | 3.2 | 0.5 |
| Pancreas | 120.8 | 16.9 | 1.4 | 112.8 | 11.5 | 1.0 |
| Larynx | 39.8 | 5.5 | 0.8 | 12.0 | 1.3 | 0.3 |
| Lung and bronchus | 460.2 | 64.9 | 2.7 | 517.8 | 55.6 | 2.2 |
| Melanoma of the skin | 225.4 | 31.6 | 1.9 | 158.6 | 18.9 | 1.4 |
| Female breast | | | | 1265.2 | 146.0 | 3.7 |
| Cervix uteri | | | | 53.2 | 7.1 | 0.9 |
| Corpus uterus and NOS | | | | 284.4 | 31.7 | 1.7 |
| Ovary | | | | 132.8 | 15.0 | 1.2 |
| Prostate | 1257.2 | 170.9 | 4.3 | | | |
| Testis | 40.4 | 6.5 | 0.9 | | | |
| Urinary bladder (incl. in situ) | 329.8 | 46.3 | 2.3 | 123.2 | 12.8 | 1.0 |
| Kidney and renal pelvis | 187.6 | 25.9 | 1.7 | 102.6 | 11.5 | 1.0 |
| Brain and other nervous system | 52.4 | 7.7 | 0.9 | 50.0 | 6.1 | 0.8 |
| Thyroid | 68.0 | 9.6 | 1.0 | 186.8 | 25.1 | 1.7 |
| Hodgkin lymphoma | 28.4 | 4.1 | 0.7 | 21.0 | 3.0 | 0.6 |
| Non-Hodgkin lymphomas | 199.4 | 28.2 | 1.8 | 180.0 | 20.0 | 1.3 |
| Multiple myeloma | 64.0 | 8.9 | 1.0 | 54.2 | 5.7 | 0.7 |
| Leukemias | 135.0 | 19.1 | 1.5 | 93.0 | 10.5 | 1.0 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2011. Rates based on fewer than 4 cases per year are unstable and should be used with caution. NOS=Not otherwise specified.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence for Selected Areas, by Gender, 2005-2009
Hempstead and Long Beach
Nassau County**

| Site of cancer | Males | | | Females | | |
|---------------------------------|---------------|------------------|--------------|---------------|------------------|--------------|
| | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) |
| All Invasive Malignant Tumors | 2369.4 | 595.9 | 10.7 | 2373.6 | 470.5 | 8.5 |
| Oral cavity and pharynx | 55.4 | 13.4 | 1.6 | 30.2 | 6.0 | 1.0 |
| Esophagus | 29.2 | 7.4 | 1.2 | 8.8 | 1.6 | 0.5 |
| Stomach | 39.6 | 10.1 | 1.4 | 30.2 | 5.9 | 0.9 |
| Colorectal | 214.6 | 54.6 | 3.3 | 220.8 | 41.3 | 2.4 |
| Colon excluding rect | 148.8 | 38.4 | 2.8 | 165.4 | 30.7 | 2.1 |
| Rectum & rectosigmoi | 65.8 | 16.2 | 1.8 | 55.4 | 10.6 | 1.2 |
| Liver / intrahepatic bile duct | 46.4 | 11.2 | 1.4 | 17.2 | 3.3 | 0.7 |
| Pancreas | 72.8 | 18.7 | 1.9 | 64.8 | 11.8 | 1.3 |
| Larynx | 25.0 | 6.2 | 1.1 | 7.2 | 1.4 | 0.5 |
| Lung and bronchus | 270.4 | 69.2 | 3.7 | 296.2 | 56.3 | 2.9 |
| Melanoma of the skin | 121.0 | 30.6 | 2.4 | 79.6 | 16.5 | 1.6 |
| Female breast | | | | 707.6 | 143.4 | 4.7 |
| Cervix uteri | | | | 32.6 | 7.6 | 1.2 |
| Corpus uterus and NOS | | | | 160.2 | 31.1 | 2.2 |
| Ovary | | | | 73.6 | 14.6 | 1.5 |
| Prostate | 702.2 | 171.1 | 5.7 | | | |
| Testis | 22.4 | 6.3 | 1.2 | | | |
| Urinary bladder (incl. in situ) | 174.2 | 45.0 | 3.0 | 67.8 | 12.5 | 1.3 |
| Kidney and renal pelvis | 105.4 | 26.2 | 2.2 | 59.6 | 11.7 | 1.3 |
| Brain and other nervous system | 28.6 | 7.5 | 1.2 | 25.6 | 5.4 | 0.9 |
| Thyroid | 37.6 | 9.4 | 1.3 | 103.8 | 23.7 | 2.0 |
| Hodgkin lymphoma | 13.8 | 3.5 | 0.8 | 12.2 | 3.0 | 0.8 |
| Non-Hodgkin lymphomas | 110.2 | 28.2 | 2.4 | 102.2 | 20.3 | 1.8 |
| Multiple myeloma | 36.0 | 9.3 | 1.4 | 32.2 | 6.1 | 0.9 |
| Leukemias | 70.6 | 18.2 | 1.9 | 49.6 | 10.1 | 1.3 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2011. Rates based on fewer than 4 cases per year are unstable and should be used with caution. NOS=Not otherwise specified.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence for Selected Areas, by Gender, 2005-2009**

**North Hempstead
Nassau County**

| Site of cancer | Males | | | Females | | |
|---------------------------------|---------------|------------------|--------------|---------------|------------------|--------------|
| | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) |
| All Invasive Malignant Tumors | 760.6 | 580.5 | 18.4 | 730.4 | 469.2 | 15.2 |
| Oral cavity and pharynx | 17.0 | 12.8 | 2.7 | 12.6 | 8.7 | 2.1 |
| Esophagus | 9.8 | 7.5 | 2.1 | 3.8 | 2.1 | 0.9 |
| Stomach | 13.4 | 10.1 | 2.4 | 8.2 | 4.9 | 1.5 |
| Colorectal | 61.8 | 47.7 | 5.3 | 62.8 | 36.7 | 4.1 |
| Colon excluding rect | 42.6 | 32.2 | 4.3 | 47.0 | 27.0 | 3.5 |
| Rectum & rectosigmoi | 19.2 | 15.5 | 3.1 | 15.8 | 9.7 | 2.1 |
| Liver / intrahepatic bile duct | 12.8 | 9.6 | 2.4 | 5.8 | 3.5 | 1.3 |
| Pancreas | 17.6 | 13.4 | 2.8 | 22.0 | 12.7 | 2.4 |
| Larynx | 6.4 | 4.8 | 1.7 | 1.4 | 0.8 | 0.6 |
| Lung and bronchus | 70.8 | 54.0 | 5.6 | 77.4 | 45.1 | 4.5 |
| Melanoma of the skin | 47.2 | 36.4 | 4.6 | 31.4 | 22.4 | 3.5 |
| Female breast | | | | 224.2 | 150.0 | 8.8 |
| Cervix uteri | | | | 9.6 | 7.6 | 2.2 |
| Corpus uterus and NOS | | | | 46.4 | 29.1 | 3.7 |
| Ovary | | | | 22.8 | 14.8 | 2.7 |
| Prostate | 232.2 | 172.5 | 9.9 | | | |
| Testis | 6.8 | 7.0 | 2.4 | | | |
| Urinary bladder (incl. in situ) | 60.8 | 45.5 | 5.1 | 21.2 | 11.8 | 2.2 |
| Kidney and renal pelvis | 32.8 | 24.5 | 3.7 | 18.8 | 11.9 | 2.4 |
| Brain and other nervous system | 10.2 | 8.3 | 2.3 | 10.2 | 7.5 | 2.1 |
| Thyroid | 12.0 | 10.1 | 2.6 | 30.0 | 24.7 | 4.0 |
| Hodgkin lymphoma | 5.2 | 4.6 | 1.8 | 3.0 | 2.7 | 1.4 |
| Non-Hodgkin lymphomas | 34.4 | 26.8 | 4.0 | 29.4 | 17.8 | 2.9 |
| Multiple myeloma | 12.6 | 9.2 | 2.3 | 9.4 | 5.6 | 1.6 |
| Leukemias | 27.8 | 21.9 | 3.6 | 18.0 | 11.0 | 2.3 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2011. Rates based on fewer than 4 cases per year are unstable and should be used with caution. NOS=Not otherwise specified.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence for Selected Areas, by Gender, 2005-2009
Oyster Bay and Glen Cove
Nassau County**

| Site of cancer | Males | | | Females | | |
|---------------------------------|---------------|------------------|--------------|---------------|------------------|--------------|
| | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) |
| All Invasive Malignant Tumors | 1086.0 | 601.7 | 16.0 | 1116.4 | 507.1 | 13.3 |
| Oral cavity and pharynx | 22.0 | 12.2 | 2.3 | 12.4 | 5.5 | 1.4 |
| Esophagus | 15.2 | 8.6 | 1.9 | 3.2 | 1.3 | 0.6 |
| Stomach | 14.2 | 7.9 | 1.8 | 11.4 | 4.7 | 1.2 |
| Colorectal | 91.8 | 51.7 | 4.7 | 99.0 | 41.4 | 3.6 |
| Colon excluding rect | 65.0 | 36.8 | 4.0 | 73.8 | 30.5 | 3.1 |
| Rectum & rectosigmoi | 26.8 | 14.9 | 2.5 | 25.2 | 10.9 | 1.9 |
| Liver / intrahepatic bile duct | 17.4 | 9.6 | 2.0 | 6.2 | 2.8 | 1.0 |
| Pancreas | 30.2 | 16.5 | 2.6 | 26.0 | 10.7 | 1.8 |
| Larynx | 8.4 | 4.7 | 1.4 | 3.4 | 1.4 | 0.7 |
| Lung and bronchus | 119.0 | 66.6 | 5.4 | 144.0 | 61.6 | 4.5 |
| Melanoma of the skin | 56.6 | 32.1 | 3.7 | 47.6 | 22.9 | 2.9 |
| Female breast | | | | 332.2 | 154.5 | 7.4 |
| Cervix uteri | | | | 11.0 | 6.2 | 1.6 |
| Corpus uterus and NOS | | | | 77.4 | 34.7 | 3.5 |
| Ovary | | | | 36.4 | 16.6 | 2.4 |
| Prostate | 321.4 | 171.0 | 8.4 | | | |
| Testis | 11.2 | 7.5 | 2.0 | | | |
| Urinary bladder (incl. in situ) | 94.6 | 52.7 | 4.7 | 34.2 | 14.8 | 2.2 |
| Kidney and renal pelvis | 49.2 | 27.0 | 3.4 | 24.0 | 10.9 | 2.0 |
| Brain and other nervous system | 13.6 | 8.0 | 1.9 | 14.2 | 6.7 | 1.6 |
| Thyroid | 18.2 | 10.3 | 2.1 | 53.0 | 30.1 | 3.6 |
| Hodgkin lymphoma | 9.4 | 5.5 | 1.6 | 5.8 | 3.7 | 1.3 |
| Non-Hodgkin lymphomas | 54.2 | 30.3 | 3.6 | 48.4 | 21.6 | 2.7 |
| Multiple myeloma | 15.4 | 8.6 | 1.9 | 12.4 | 5.1 | 1.3 |
| Leukemias | 35.8 | 20.3 | 3.0 | 25.2 | 11.6 | 2.0 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2011. Rates based on fewer than 4 cases per year are unstable and should be used with caution. NOS=Not otherwise specified.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence for Selected Areas, by Gender, 2005-2009**

Rockland County

| Site of cancer | Males | | | Females | | |
|---------------------------------|---------------|------------------|--------------|---------------|------------------|--------------|
| | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) |
| All Invasive Malignant Tumors | 851.8 | 591.4 | 18.2 | 802.8 | 462.9 | 14.6 |
| Oral cavity and pharynx | 15.2 | 10.0 | 2.3 | 8.0 | 4.6 | 1.5 |
| Esophagus | 13.2 | 9.0 | 2.2 | 4.6 | 2.5 | 1.0 |
| Stomach | 18.6 | 13.7 | 2.8 | 11.2 | 6.4 | 1.7 |
| Colorectal | 72.4 | 53.2 | 5.6 | 67.4 | 37.9 | 4.1 |
| Colon excluding rect | 52.8 | 38.6 | 4.8 | 52.6 | 29.5 | 3.6 |
| Rectum & rectosigmoi | 19.6 | 14.6 | 2.9 | 14.8 | 8.4 | 1.9 |
| Liver / intrahepatic bile duct | 12.6 | 8.9 | 2.3 | 6.4 | 3.8 | 1.4 |
| Pancreas | 20.0 | 14.5 | 2.9 | 20.4 | 11.2 | 2.2 |
| Larynx | 8.2 | 5.7 | 1.8 | 0.8 | 0.4 | 0.4 |
| Lung and bronchus | 95.0 | 68.4 | 6.3 | 94.0 | 52.4 | 4.8 |
| Melanoma of the skin | 37.0 | 25.5 | 3.8 | 27.0 | 16.5 | 2.8 |
| Female breast | | | | 239.0 | 138.3 | 8.0 |
| Cervix uteri | | | | 11.2 | 7.4 | 2.0 |
| Corpus uterus and NOS | | | | 57.6 | 31.9 | 3.7 |
| Ovary | | | | 23.8 | 13.5 | 2.5 |
| Prostate | 296.6 | 192.7 | 10.0 | | | |
| Testis | 10.6 | 8.0 | 2.2 | | | |
| Urinary bladder (incl. in situ) | 57.0 | 42.4 | 5.0 | 21.6 | 11.6 | 2.2 |
| Kidney and renal pelvis | 33.4 | 22.9 | 3.6 | 18.0 | 10.5 | 2.2 |
| Brain and other nervous system | 10.6 | 7.2 | 2.0 | 11.6 | 7.0 | 1.8 |
| Thyroid | 13.4 | 9.6 | 2.3 | 43.0 | 29.2 | 4.1 |
| Hodgkin lymphoma | 4.6 | 3.3 | 1.4 | 5.4 | 3.9 | 1.5 |
| Non-Hodgkin lymphomas | 41.0 | 28.8 | 4.0 | 36.8 | 21.2 | 3.1 |
| Multiple myeloma | 11.4 | 7.8 | 2.1 | 11.0 | 6.4 | 1.7 |
| Leukemias | 25.0 | 18.2 | 3.3 | 22.0 | 12.3 | 2.3 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2011. Rates based on fewer than 4 cases per year are unstable and should be used with caution. NOS=Not otherwise specified.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence for Selected Areas, by Gender, 2005-2009**

**Ramapo
Rockland County**

| Site of cancer | Males | | | Females | | |
|---------------------------------|---------------|------------------|--------------|---------------|------------------|--------------|
| | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) |
| All Invasive Malignant Tumors | 259.0 | 535.8 | 29.2 | 267.6 | 452.9 | 24.3 |
| Oral cavity and pharynx | 4.8 | 9.6 | 3.8 | 2.6 | 4.4 | 2.4 |
| Esophagus | 3.2 | 6.0 | 2.9 | 1.2 | 1.9 | 1.5 |
| Stomach | 7.2 | 15.7 | 5.1 | 4.6 | 8.0 | 3.3 |
| Colorectal | 25.0 | 54.6 | 9.6 | 25.4 | 41.6 | 7.2 |
| Colon excluding rect | 17.2 | 36.6 | 7.7 | 19.6 | 32.1 | 6.4 |
| Rectum & rectosigmoi | 7.8 | 18.0 | 5.6 | 5.8 | 9.5 | 3.5 |
| Liver / intrahepatic bile duct | 3.4 | 6.6 | 3.1 | 0.4 | 0.8 | 1.1 |
| Pancreas | 5.4 | 11.4 | 4.3 | 7.8 | 12.7 | 4.0 |
| Larynx | 3.2 | 6.2 | 3.0 | 0.2 | 0.4 | 0.8 |
| Lung and bronchus | 23.6 | 51.0 | 9.2 | 31.4 | 51.2 | 8.0 |
| Melanoma of the skin | 7.8 | 16.3 | 5.1 | 7.2 | 12.6 | 4.1 |
| Female breast | | | | 74.2 | 128.4 | 13.1 |
| Cervix uteri | | | | 4.8 | 8.8 | 3.5 |
| Corpus uterus and NOS | | | | 19.0 | 31.3 | 6.3 |
| Ovary | | | | 7.0 | 11.7 | 3.9 |
| Prostate | 89.6 | 174.1 | 16.1 | | | |
| Testis | 3.2 | 6.2 | 3.0 | | | |
| Urinary bladder (incl. in situ) | 15.4 | 35.2 | 7.9 | 6.8 | 10.8 | 3.6 |
| Kidney and renal pelvis | 11.4 | 23.8 | 6.2 | 6.4 | 11.1 | 3.8 |
| Brain and other nervous system | 3.6 | 6.4 | 3.0 | 4.4 | 7.3 | 3.1 |
| Thyroid | 5.0 | 10.3 | 4.0 | 18.4 | 33.8 | 6.9 |
| Hodgkin lymphoma | 2.4 | 4.7 | 2.7 | 1.6 | 2.9 | 2.0 |
| Non-Hodgkin lymphomas | 15.0 | 32.4 | 7.3 | 11.8 | 20.8 | 5.3 |
| Multiple myeloma | 4.0 | 8.9 | 3.9 | 4.4 | 7.5 | 3.1 |
| Leukemias | 8.8 | 19.3 | 5.7 | 7.0 | 11.5 | 3.8 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2011. Rates based on fewer than 4 cases per year are unstable and should be used with caution. NOS=Not otherwise specified.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence for Selected Areas, by Gender, 2005-2009
Remainder of Rockland County
Rockland County**

| Site of cancer | Males | | | Females | | |
|---------------------------------|---------------|------------------|--------------|---------------|------------------|--------------|
| | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) |
| All Invasive Malignant Tumors | 591.6 | 687.7 | 24.8 | 534.6 | 507.1 | 19.2 |
| Oral cavity and pharynx | 10.4 | 11.2 | 3.0 | 5.4 | 5.2 | 2.0 |
| Esophagus | 9.8 | 11.4 | 3.2 | 3.4 | 3.1 | 1.5 |
| Stomach | 11.4 | 14.7 | 3.8 | 6.6 | 6.2 | 2.1 |
| Colorectal | 47.4 | 60.2 | 7.7 | 41.8 | 39.5 | 5.4 |
| Colon excluding rect | 35.6 | 45.3 | 6.7 | 32.8 | 31.2 | 4.8 |
| Rectum & rectosigmoi | 11.8 | 14.9 | 3.8 | 9.0 | 8.4 | 2.5 |
| Liver / intrahepatic bile duct | 9.2 | 11.6 | 3.4 | 6.0 | 5.9 | 2.1 |
| Pancreas | 14.6 | 18.2 | 4.2 | 12.6 | 11.6 | 2.9 |
| Larynx | 5.0 | 6.0 | 2.4 | 0.6 | 0.5 | 0.6 |
| Lung and bronchus | 71.2 | 86.5 | 9.0 | 62.6 | 57.0 | 6.3 |
| Melanoma of the skin | 29.2 | 33.1 | 5.4 | 19.8 | 19.9 | 3.9 |
| Female breast | | | | 164.6 | 154.9 | 10.6 |
| Cervix uteri | | | | 6.4 | 7.1 | 2.5 |
| Corpus uterus and NOS | | | | 38.6 | 34.2 | 4.8 |
| Ovary | | | | 16.8 | 16.0 | 3.4 |
| Prostate | 206.8 | 219.1 | 13.4 | | | |
| Testis | 7.4 | 9.7 | 3.1 | | | |
| Urinary bladder (incl. in situ) | 41.4 | 52.3 | 7.1 | 14.8 | 13.3 | 3.0 |
| Kidney and renal pelvis | 21.8 | 24.7 | 4.6 | 11.6 | 10.9 | 2.8 |
| Brain and other nervous system | 7.0 | 8.4 | 2.8 | 7.2 | 7.6 | 2.5 |
| Thyroid | 8.4 | 9.9 | 3.0 | 24.6 | 27.4 | 4.8 |
| Hodgkin lymphoma | 2.2 | 2.7 | 1.6 | 3.8 | 4.7 | 2.1 |
| Non-Hodgkin lymphomas | 26.0 | 30.2 | 5.2 | 25.0 | 23.6 | 4.1 |
| Multiple myeloma | 7.2 | 8.0 | 2.6 | 6.6 | 6.2 | 2.1 |
| Leukemias | 16.2 | 20.0 | 4.4 | 15.0 | 14.3 | 3.2 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2011. Rates based on fewer than 4 cases per year are unstable and should be used with caution. NOS=Not otherwise specified.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence for Selected Areas, by Gender, 2005-2009
Suffolk County**

| Site of cancer | Males | | | Females | | |
|---------------------------------|---------------|------------------|--------------|---------------|------------------|--------------|
| | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) |
| All Invasive Malignant Tumors | 4435.2 | 613.9 | 8.3 | 4339.2 | 487.8 | 6.6 |
| Oral cavity and pharynx | 116.6 | 14.9 | 1.2 | 52.0 | 5.8 | 0.7 |
| Esophagus | 68.4 | 9.5 | 1.0 | 23.2 | 2.6 | 0.5 |
| Stomach | 78.0 | 11.1 | 1.1 | 42.8 | 4.7 | 0.6 |
| Colorectal | 398.6 | 56.8 | 2.6 | 402.2 | 43.8 | 1.9 |
| Colon excluding rect | 283.2 | 41.0 | 2.2 | 302.4 | 32.8 | 1.7 |
| Rectum & rectosigmoi | 115.4 | 15.8 | 1.3 | 99.8 | 11.0 | 1.0 |
| Liver / intrahepatic bile duct | 80.8 | 10.5 | 1.0 | 28.8 | 3.1 | 0.5 |
| Pancreas | 112.0 | 15.8 | 1.3 | 126.8 | 13.7 | 1.1 |
| Larynx | 45.8 | 6.2 | 0.8 | 16.4 | 1.8 | 0.4 |
| Lung and bronchus | 558.0 | 79.9 | 3.0 | 616.8 | 68.5 | 2.4 |
| Melanoma of the skin | 229.0 | 31.5 | 1.9 | 150.6 | 17.8 | 1.3 |
| Female breast | | | | 1209.4 | 135.6 | 3.5 |
| Cervix uteri | | | | 67.6 | 8.3 | 0.9 |
| Corpus uterus and NOS | | | | 266.4 | 29.2 | 1.6 |
| Ovary | | | | 125.2 | 14.1 | 1.1 |
| Prostate | 1262.8 | 168.4 | 4.2 | | | |
| Testis | 47.0 | 6.5 | 0.8 | | | |
| Urinary bladder (incl. in situ) | 336.2 | 49.1 | 2.4 | 110.2 | 12.1 | 1.0 |
| Kidney and renal pelvis | 183.8 | 24.6 | 1.6 | 111.8 | 12.6 | 1.1 |
| Brain and other nervous system | 60.8 | 8.2 | 0.9 | 43.6 | 5.2 | 0.7 |
| Thyroid | 77.0 | 9.9 | 1.0 | 210.8 | 26.2 | 1.6 |
| Hodgkin lymphoma | 25.4 | 3.5 | 0.6 | 27.6 | 3.7 | 0.6 |
| Non-Hodgkin lymphomas | 194.2 | 27.3 | 1.8 | 168.6 | 19.0 | 1.3 |
| Multiple myeloma | 58.8 | 8.2 | 1.0 | 53.4 | 5.8 | 0.7 |
| Leukemias | 138.8 | 19.5 | 1.5 | 101.8 | 11.7 | 1.0 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2011. Rates based on fewer than 4 cases per year are unstable and should be used with caution. NOS=Not otherwise specified.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence for Selected Areas, by Gender, 2005-2009**

**Babylon
Suffolk County**

| Site of cancer | Males | | | Females | | |
|---------------------------------|---------------|------------------|--------------|---------------|------------------|--------------|
| | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) |
| All Invasive Malignant Tumors | 620.4 | 634.6 | 22.3 | 629.6 | 491.9 | 17.2 |
| Oral cavity and pharynx | 16.4 | 15.7 | 3.4 | 7.4 | 5.9 | 1.9 |
| Esophagus | 8.6 | 8.9 | 2.7 | 3.0 | 2.3 | 1.2 |
| Stomach | 12.2 | 12.7 | 3.2 | 8.8 | 6.8 | 2.0 |
| Colorectal | 62.8 | 65.7 | 7.3 | 62.2 | 47.1 | 5.2 |
| Colon excluding rect | 45.6 | 48.0 | 6.2 | 46.8 | 35.1 | 4.5 |
| Rectum & rectosigmoi | 17.2 | 17.7 | 3.7 | 15.4 | 12.0 | 2.7 |
| Liver / intrahepatic bile duct | 9.8 | 9.3 | 2.6 | 6.8 | 5.2 | 1.7 |
| Pancreas | 17.4 | 18.5 | 3.9 | 21.0 | 15.7 | 3.0 |
| Larynx | 7.2 | 7.2 | 2.4 | 3.6 | 2.8 | 1.3 |
| Lung and bronchus | 85.6 | 90.3 | 8.6 | 94.8 | 73.1 | 6.6 |
| Melanoma of the skin | 27.2 | 27.6 | 4.6 | 16.8 | 13.5 | 2.9 |
| Female breast | | | | 170.0 | 133.1 | 8.9 |
| Cervix uteri | | | | 11.4 | 9.5 | 2.5 |
| Corpus uterus and NOS | | | | 41.2 | 31.8 | 4.3 |
| Ovary | | | | 18.6 | 15.0 | 3.0 |
| Prostate | 175.2 | 173.0 | 11.5 | | | |
| Testis | 6.8 | 6.6 | 2.2 | | | |
| Urinary bladder (incl. in situ) | 51.2 | 56.1 | 6.9 | 16.6 | 12.6 | 2.7 |
| Kidney and renal pelvis | 24.6 | 23.8 | 4.2 | 16.8 | 13.2 | 2.8 |
| Brain and other nervous system | 6.6 | 6.5 | 2.2 | 5.6 | 4.6 | 1.7 |
| Thyroid | 7.0 | 6.7 | 2.2 | 22.8 | 19.3 | 3.5 |
| Hodgkin lymphoma | 3.8 | 3.7 | 1.7 | 4.0 | 3.8 | 1.7 |
| Non-Hodgkin lymphomas | 23.6 | 24.2 | 4.4 | 25.0 | 19.7 | 3.5 |
| Multiple myeloma | 7.4 | 7.5 | 2.4 | 7.2 | 5.4 | 1.8 |
| Leukemias | 17.2 | 17.8 | 3.8 | 13.2 | 10.7 | 2.6 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2011. Rates based on fewer than 4 cases per year are unstable and should be used with caution. NOS=Not otherwise specified.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence for Selected Areas, by Gender, 2005-2009**

**Brookhaven
Suffolk County**

| Site of cancer | Males | | | Females | | |
|---------------------------------|---------------|------------------|--------------|---------------|------------------|--------------|
| | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) |
| All Invasive Malignant Tumors | 1346.6 | 654.1 | 15.6 | 1351.2 | 525.9 | 12.5 |
| Oral cavity and pharynx | 29.6 | 13.3 | 2.1 | 16.6 | 6.5 | 1.4 |
| Esophagus | 20.6 | 9.8 | 1.9 | 7.4 | 2.9 | 0.9 |
| Stomach | 24.4 | 12.3 | 2.2 | 11.8 | 4.5 | 1.1 |
| Colorectal | 123.4 | 63.0 | 5.0 | 121.6 | 47.4 | 3.8 |
| Colon excluding rect | 86.2 | 45.2 | 4.3 | 93.2 | 36.4 | 3.3 |
| Rectum & rectosigmoi | 37.2 | 17.7 | 2.5 | 28.4 | 11.0 | 1.8 |
| Liver / intrahepatic bile duct | 31.0 | 14.3 | 2.3 | 7.8 | 3.0 | 0.9 |
| Pancreas | 30.8 | 15.0 | 2.4 | 35.2 | 13.8 | 2.0 |
| Larynx | 14.2 | 6.9 | 1.6 | 6.0 | 2.4 | 0.9 |
| Lung and bronchus | 185.0 | 94.7 | 6.1 | 200.2 | 78.2 | 4.8 |
| Melanoma of the skin | 66.8 | 32.0 | 3.4 | 44.6 | 17.6 | 2.3 |
| Female breast | | | | 362.6 | 139.6 | 6.4 |
| Cervix uteri | | | | 19.6 | 7.8 | 1.5 |
| Corpus uterus and NOS | | | | 82.4 | 30.9 | 3.0 |
| Ovary | | | | 39.6 | 15.1 | 2.1 |
| Prostate | 351.6 | 164.7 | 7.7 | | | |
| Testis | 17.4 | 7.3 | 1.5 | | | |
| Urinary bladder (incl. in situ) | 99.8 | 51.4 | 4.5 | 32.2 | 12.7 | 2.0 |
| Kidney and renal pelvis | 57.2 | 26.7 | 3.1 | 37.4 | 14.8 | 2.1 |
| Brain and other nervous system | 20.4 | 9.4 | 1.8 | 11.8 | 4.7 | 1.2 |
| Thyroid | 33.4 | 14.5 | 2.2 | 80.4 | 32.3 | 3.2 |
| Hodgkin lymphoma | 7.0 | 3.2 | 1.1 | 11.0 | 4.6 | 1.2 |
| Non-Hodgkin lymphomas | 60.0 | 29.3 | 3.3 | 51.8 | 20.4 | 2.5 |
| Multiple myeloma | 15.6 | 8.0 | 1.8 | 16.8 | 6.5 | 1.4 |
| Leukemias | 44.2 | 21.4 | 2.8 | 33.8 | 13.4 | 2.0 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2011. Rates based on fewer than 4 cases per year are unstable and should be used with caution. NOS=Not otherwise specified.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence for Selected Areas, by Gender, 2005-2009**

**Huntingdon
Suffolk County**

| Site of cancer | Males | | | Females | | |
|---------------------------------|---------------|------------------|--------------|---------------|------------------|--------------|
| | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) |
| All Invasive Malignant Tumors | 646.4 | 628.2 | 21.7 | 593.6 | 487.4 | 17.5 |
| Oral cavity and pharynx | 16.6 | 15.4 | 3.3 | 6.2 | 5.0 | 1.8 |
| Esophagus | 9.8 | 10.1 | 2.8 | 2.4 | 2.0 | 1.1 |
| Stomach | 10.2 | 10.0 | 2.7 | 5.8 | 4.7 | 1.7 |
| Colorectal | 55.2 | 54.8 | 6.5 | 50.6 | 40.3 | 5.0 |
| Colon excluding rect | 38.4 | 38.6 | 5.5 | 36.4 | 28.7 | 4.2 |
| Rectum & rectosigmoi | 16.8 | 16.2 | 3.5 | 14.2 | 11.6 | 2.7 |
| Liver / intrahepatic bile duct | 8.4 | 7.8 | 2.4 | 2.6 | 2.0 | 1.1 |
| Pancreas | 17.6 | 17.2 | 3.6 | 16.6 | 12.9 | 2.8 |
| Larynx | 4.6 | 4.4 | 1.8 | 1.4 | 1.2 | 0.9 |
| Lung and bronchus | 61.2 | 62.7 | 7.0 | 75.6 | 61.0 | 6.1 |
| Melanoma of the skin | 40.4 | 39.2 | 5.4 | 26.2 | 23.4 | 4.0 |
| Female breast | | | | 181.0 | 149.4 | 9.7 |
| Cervix uteri | | | | 5.2 | 4.9 | 1.9 |
| Corpus uterus and NOS | | | | 35.2 | 27.2 | 4.0 |
| Ovary | | | | 18.6 | 15.3 | 3.1 |
| Prostate | 207.2 | 191.6 | 11.7 | | | |
| Testis | 5.6 | 6.1 | 2.3 | | | |
| Urinary bladder (incl. in situ) | 49.8 | 50.4 | 6.3 | 16.0 | 12.3 | 2.7 |
| Kidney and renal pelvis | 24.0 | 22.7 | 4.1 | 11.8 | 9.7 | 2.5 |
| Brain and other nervous system | 9.2 | 9.3 | 2.7 | 5.8 | 5.1 | 1.9 |
| Thyroid | 12.8 | 12.7 | 3.1 | 30.0 | 27.3 | 4.4 |
| Hodgkin lymphoma | 4.6 | 4.6 | 1.9 | 3.2 | 3.1 | 1.5 |
| Non-Hodgkin lymphomas | 30.8 | 31.5 | 5.0 | 25.2 | 20.2 | 3.5 |
| Multiple myeloma | 8.6 | 8.3 | 2.5 | 8.4 | 6.7 | 2.0 |
| Leukemias | 18.2 | 17.6 | 3.6 | 11.8 | 9.8 | 2.5 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2011. Rates based on fewer than 4 cases per year are unstable and should be used with caution. NOS=Not otherwise specified.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence for Selected Areas, by Gender, 2005-2009**

**Islip
Suffolk County**

| Site of cancer | Males | | | Females | | |
|---------------------------------|---------------|------------------|--------------|---------------|------------------|--------------|
| | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) |
| All Invasive Malignant Tumors | 889.8 | 639.6 | 18.8 | 885.6 | 501.6 | 14.8 |
| Oral cavity and pharynx | 26.8 | 17.1 | 2.9 | 10.0 | 5.6 | 1.6 |
| Esophagus | 13.6 | 9.7 | 2.3 | 4.8 | 2.7 | 1.1 |
| Stomach | 18.4 | 14.0 | 2.9 | 9.0 | 5.2 | 1.5 |
| Colorectal | 87.0 | 66.5 | 6.2 | 86.6 | 49.6 | 4.7 |
| Colon excluding rect | 63.2 | 50.0 | 5.5 | 64.4 | 37.0 | 4.0 |
| Rectum & rectosigmoi | 23.8 | 16.5 | 3.0 | 22.2 | 12.6 | 2.3 |
| Liver / intrahepatic bile duct | 17.6 | 11.8 | 2.5 | 6.2 | 3.6 | 1.3 |
| Pancreas | 23.0 | 17.3 | 3.2 | 25.6 | 14.5 | 2.5 |
| Larynx | 10.8 | 7.8 | 2.1 | 3.4 | 1.8 | 0.9 |
| Lung and bronchus | 111.6 | 83.6 | 6.9 | 121.4 | 69.4 | 5.5 |
| Melanoma of the skin | 40.8 | 28.9 | 4.0 | 26.2 | 15.2 | 2.6 |
| Female breast | | | | 244.6 | 135.8 | 7.6 |
| Cervix uteri | | | | 19.2 | 11.1 | 2.2 |
| Corpus uterus and NOS | | | | 56.4 | 30.7 | 3.6 |
| Ovary | | | | 23.0 | 13.2 | 2.4 |
| Prostate | 246.4 | 170.5 | 9.5 | | | |
| Testis | 10.2 | 6.2 | 1.7 | | | |
| Urinary bladder (incl. in situ) | 63.0 | 49.8 | 5.5 | 21.2 | 12.0 | 2.3 |
| Kidney and renal pelvis | 44.6 | 30.4 | 4.0 | 22.8 | 12.9 | 2.4 |
| Brain and other nervous system | 11.0 | 7.0 | 1.9 | 11.0 | 6.4 | 1.7 |
| Thyroid | 12.0 | 7.1 | 1.8 | 47.4 | 27.3 | 3.5 |
| Hodgkin lymphoma | 6.8 | 4.1 | 1.4 | 5.0 | 3.0 | 1.2 |
| Non-Hodgkin lymphomas | 38.6 | 27.8 | 3.9 | 36.2 | 20.4 | 3.0 |
| Multiple myeloma | 12.4 | 8.8 | 2.2 | 9.0 | 5.1 | 1.5 |
| Leukemias | 28.2 | 20.3 | 3.4 | 20.2 | 11.8 | 2.3 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2011. Rates based on fewer than 4 cases per year are unstable and should be used with caution. NOS=Not otherwise specified.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence for Selected Areas, by Gender, 2005-2009**

**Smithtown
Suffolk County**

| Site of cancer | Males | | | Females | | |
|---------------------------------|---------------|------------------|--------------|---------------|------------------|--------------|
| | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) |
| All Invasive Malignant Tumors | 377.8 | 620.7 | 28.0 | 388.2 | 509.6 | 22.7 |
| Oral cavity and pharynx | 8.0 | 11.8 | 3.7 | 5.2 | 6.9 | 2.7 |
| Esophagus | 5.6 | 9.0 | 3.3 | 2.8 | 3.6 | 1.9 |
| Stomach | 5.2 | 9.3 | 3.6 | 3.8 | 5.0 | 2.2 |
| Colorectal | 29.4 | 50.2 | 8.1 | 33.6 | 40.6 | 6.1 |
| Colon excluding rect | 20.0 | 34.6 | 6.8 | 24.0 | 29.2 | 5.2 |
| Rectum & rectosigmoi | 9.4 | 15.7 | 4.5 | 9.6 | 11.3 | 3.2 |
| Liver / intrahepatic bile duct | 7.4 | 10.9 | 3.5 | 2.4 | 3.0 | 1.7 |
| Pancreas | 7.2 | 12.0 | 3.9 | 13.6 | 16.3 | 3.9 |
| Larynx | 2.8 | 4.0 | 2.1 | 0.6 | 0.8 | 0.9 |
| Lung and bronchus | 42.6 | 71.3 | 9.6 | 54.0 | 67.9 | 8.1 |
| Melanoma of the skin | 22.4 | 37.5 | 6.9 | 11.8 | 16.8 | 4.3 |
| Female breast | | | | 116.8 | 156.8 | 12.7 |
| Cervix uteri | | | | 3.6 | 5.8 | 2.7 |
| Corpus uterus and NOS | | | | 22.8 | 29.3 | 5.4 |
| Ovary | | | | 11.2 | 15.8 | 4.1 |
| Prostate | 123.6 | 196.2 | 15.5 | | | |
| Testis | 2.6 | 5.4 | 2.9 | | | |
| Urinary bladder (incl. in situ) | 31.0 | 52.6 | 8.3 | 10.8 | 13.6 | 3.6 |
| Kidney and renal pelvis | 15.6 | 25.4 | 5.6 | 9.6 | 12.7 | 3.6 |
| Brain and other nervous system | 5.4 | 8.9 | 3.4 | 4.8 | 6.6 | 2.6 |
| Thyroid | 7.6 | 11.4 | 3.6 | 17.2 | 27.0 | 5.7 |
| Hodgkin lymphoma | 1.6 | 3.1 | 2.1 | 2.0 | 3.6 | 2.2 |
| Non-Hodgkin lymphomas | 14.4 | 23.4 | 5.4 | 14.0 | 17.8 | 4.2 |
| Multiple myeloma | 5.4 | 9.1 | 3.4 | 4.4 | 5.5 | 2.3 |
| Leukemias | 12.6 | 22.2 | 5.5 | 10.0 | 13.1 | 3.6 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2011. Rates based on fewer than 4 cases per year are unstable and should be used with caution. NOS=Not otherwise specified.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence for Selected Areas, by Gender, 2005-2009**

**East End Towns
Suffolk County**

| Site of cancer | Males | | | Females | | |
|---------------------------------|---------------|------------------|--------------|---------------|------------------|--------------|
| | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) |
| All Invasive Malignant Tumors | 545.4 | 604.1 | 22.7 | 483.4 | 463.7 | 18.5 |
| Oral cavity and pharynx | 18.4 | 20.4 | 4.2 | 6.4 | 6.2 | 2.1 |
| Esophagus | 10.0 | 11.0 | 3.0 | 2.8 | 2.5 | 1.3 |
| Stomach | 7.4 | 8.2 | 2.6 | 3.6 | 3.1 | 1.4 |
| Colorectal | 40.6 | 45.2 | 6.2 | 46.8 | 40.3 | 5.2 |
| Colon excluding rect | 29.6 | 32.9 | 5.3 | 37.0 | 31.6 | 4.6 |
| Rectum & rectosigmoi | 11.0 | 12.2 | 3.2 | 9.8 | 8.7 | 2.4 |
| Liver / intrahepatic bile duct | 6.6 | 7.1 | 2.4 | 3.0 | 2.6 | 1.3 |
| Pancreas | 15.6 | 17.0 | 3.8 | 14.6 | 12.6 | 2.9 |
| Larynx | 6.2 | 6.6 | 2.3 | 1.4 | 1.2 | 0.9 |
| Lung and bronchus | 70.2 | 76.2 | 8.0 | 70.0 | 62.6 | 6.6 |
| Melanoma of the skin | 30.2 | 34.0 | 5.4 | 24.6 | 27.4 | 4.8 |
| Female breast | | | | 132.4 | 129.6 | 9.9 |
| Cervix uteri | | | | 8.4 | 10.5 | 3.2 |
| Corpus uterus and NOS | | | | 28.0 | 27.3 | 4.5 |
| Ovary | | | | 14.2 | 13.4 | 3.1 |
| Prostate | 157.2 | 166.8 | 11.7 | | | |
| Testis | 4.4 | 7.0 | 2.9 | | | |
| Urinary bladder (incl. in situ) | 40.8 | 44.8 | 6.1 | 13.4 | 11.6 | 2.8 |
| Kidney and renal pelvis | 17.8 | 19.8 | 4.1 | 13.2 | 12.2 | 2.9 |
| Brain and other nervous system | 8.0 | 9.8 | 3.0 | 4.2 | 4.8 | 2.1 |
| Thyroid | 4.2 | 5.7 | 2.4 | 12.6 | 16.7 | 4.1 |
| Hodgkin lymphoma | 1.6 | 2.3 | 1.6 | 2.0 | 2.3 | 1.4 |
| Non-Hodgkin lymphomas | 26.2 | 30.2 | 5.2 | 16.2 | 16.1 | 3.5 |
| Multiple myeloma | 9.2 | 9.9 | 2.9 | 7.6 | 6.7 | 2.1 |
| Leukemias | 18.4 | 21.7 | 4.4 | 12.6 | 12.8 | 3.2 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2011. Rates based on fewer than 4 cases per year are unstable and should be used with caution. NOS=Not otherwise specified.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence for Selected Areas, by Gender, 2005-2009
Westchester County**

| Site of cancer | Males | | | Females | | |
|---------------------------------|---------------|------------------|--------------|---------------|------------------|--------------|
| | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) |
| All Invasive Malignant Tumors | 2685.8 | 574.8 | 9.8 | 2752.6 | 459.1 | 7.8 |
| Oral cavity and pharynx | 67.8 | 13.9 | 1.5 | 33.8 | 5.6 | 0.9 |
| Esophagus | 35.4 | 7.5 | 1.1 | 16.6 | 2.6 | 0.6 |
| Stomach | 50.8 | 11.1 | 1.4 | 39.2 | 6.1 | 0.9 |
| Colorectal | 234.0 | 50.5 | 2.9 | 255.4 | 40.3 | 2.3 |
| Colon excluding rect | 166.0 | 36.2 | 2.5 | 196.2 | 30.7 | 2.0 |
| Rectum & rectosigmoi | 68.0 | 14.3 | 1.5 | 59.2 | 9.6 | 1.1 |
| Liver / intrahepatic bile duct | 58.2 | 11.9 | 1.4 | 19.6 | 3.2 | 0.6 |
| Pancreas | 70.8 | 15.3 | 1.6 | 77.6 | 11.8 | 1.2 |
| Larynx | 27.8 | 6.0 | 1.0 | 8.0 | 1.3 | 0.4 |
| Lung and bronchus | 291.4 | 63.9 | 3.3 | 325.8 | 52.8 | 2.6 |
| Melanoma of the skin | 119.0 | 25.5 | 2.1 | 89.4 | 15.5 | 1.5 |
| Female breast | | | | 815.0 | 138.8 | 4.3 |
| Cervix uteri | | | | 39.6 | 7.4 | 1.0 |
| Corpus uterus and NOS | | | | 179.4 | 30.1 | 2.0 |
| Ovary | | | | 82.0 | 14.0 | 1.4 |
| Prostate | 834.2 | 174.3 | 5.3 | | | |
| Testis | 25.6 | 5.9 | 1.0 | | | |
| Urinary bladder (incl. in situ) | 182.6 | 40.6 | 2.7 | 71.8 | 11.1 | 1.2 |
| Kidney and renal pelvis | 103.6 | 21.8 | 1.9 | 58.8 | 9.6 | 1.1 |
| Brain and other nervous system | 39.0 | 8.4 | 1.2 | 29.6 | 5.2 | 0.9 |
| Thyroid | 46.4 | 9.6 | 1.3 | 128.0 | 24.8 | 2.0 |
| Hodgkin lymphoma | 13.4 | 3.0 | 0.7 | 17.0 | 3.6 | 0.8 |
| Non-Hodgkin lymphomas | 129.0 | 27.9 | 2.2 | 121.8 | 20.0 | 1.6 |
| Multiple myeloma | 43.8 | 9.3 | 1.3 | 37.0 | 5.8 | 0.9 |
| Leukemias | 90.4 | 19.8 | 1.8 | 71.4 | 11.7 | 1.2 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2011. Rates based on fewer than 4 cases per year are unstable and should be used with caution. NOS=Not otherwise specified.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence for Selected Areas, by Gender, 2005-2009**

**Northwest
Westchester County**

| Site of cancer | Males | | | Females | | |
|---------------------------------|---------------|------------------|--------------|---------------|------------------|--------------|
| | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) |
| All Invasive Malignant Tumors | 413.2 | 601.2 | 25.9 | 409.4 | 475.5 | 20.6 |
| Oral cavity and pharynx | 11.6 | 15.2 | 3.9 | 5.2 | 6.2 | 2.4 |
| Esophagus | 7.6 | 10.8 | 3.4 | 3.2 | 3.7 | 1.8 |
| Stomach | 7.6 | 12.0 | 3.8 | 5.6 | 5.8 | 2.1 |
| Colorectal | 33.0 | 48.3 | 7.4 | 37.0 | 41.4 | 6.0 |
| Colon excluding rect | 25.4 | 38.3 | 6.7 | 31.0 | 34.9 | 5.5 |
| Rectum & rectosigmoi | 7.6 | 10.1 | 3.2 | 6.0 | 6.6 | 2.4 |
| Liver / intrahepatic bile duct | 10.6 | 14.7 | 4.0 | 2.6 | 2.8 | 1.5 |
| Pancreas | 9.6 | 14.0 | 4.0 | 11.2 | 11.5 | 3.0 |
| Larynx | 4.8 | 6.5 | 2.6 | 0.8 | 1.0 | 1.0 |
| Lung and bronchus | 45.8 | 69.2 | 9.0 | 51.4 | 58.6 | 7.2 |
| Melanoma of the skin | 18.2 | 26.1 | 5.4 | 12.0 | 14.9 | 3.8 |
| Female breast | | | | 125.6 | 147.7 | 11.6 |
| Cervix uteri | | | | 5.4 | 7.1 | 2.7 |
| Corpus uterus and NOS | | | | 27.2 | 31.7 | 5.3 |
| Ovary | | | | 11.8 | 13.8 | 3.5 |
| Prostate | 125.4 | 177.9 | 13.9 | | | |
| Testis | 3.4 | 4.7 | 2.2 | | | |
| Urinary bladder (incl. in situ) | 29.0 | 45.8 | 7.5 | 9.2 | 10.0 | 2.9 |
| Kidney and renal pelvis | 13.8 | 19.9 | 4.7 | 9.4 | 11.2 | 3.2 |
| Brain and other nervous system | 6.8 | 9.4 | 3.2 | 3.8 | 4.9 | 2.2 |
| Thyroid | 6.6 | 8.3 | 2.8 | 20.6 | 27.1 | 5.2 |
| Hodgkin lymphoma | 1.6 | 2.3 | 1.6 | 1.8 | 2.8 | 1.8 |
| Non-Hodgkin lymphomas | 21.8 | 31.9 | 6.0 | 13.6 | 15.5 | 3.7 |
| Multiple myeloma | 6.4 | 9.5 | 3.3 | 3.4 | 3.5 | 1.7 |
| Leukemias | 14.4 | 22.2 | 5.1 | 10.8 | 12.0 | 3.2 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2011. Rates based on fewer than 4 cases per year are unstable and should be used with caution. NOS=Not otherwise specified.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence for Selected Areas, by Gender, 2005-2009**

**Northeast
Westchester County**

| Site of cancer | Males | | | Females | | |
|---------------------------------|---------------|------------------|--------------|---------------|------------------|--------------|
| | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) |
| All Invasive Malignant Tumors | 406.0 | 581.1 | 25.3 | 411.2 | 488.5 | 21.1 |
| Oral cavity and pharynx | 10.0 | 13.3 | 3.7 | 4.8 | 5.9 | 2.4 |
| Esophagus | 4.8 | 7.2 | 2.9 | 1.4 | 1.6 | 1.2 |
| Stomach | 4.4 | 6.5 | 2.7 | 4.0 | 4.9 | 2.1 |
| Colorectal | 29.8 | 46.0 | 7.4 | 34.4 | 40.5 | 6.1 |
| Colon excluding rect | 19.8 | 30.5 | 6.0 | 26.4 | 30.8 | 5.3 |
| Rectum & rectosigmoi | 10.0 | 15.5 | 4.3 | 8.0 | 9.7 | 3.0 |
| Liver / intrahepatic bile duct | 5.0 | 6.8 | 2.7 | 3.4 | 3.9 | 1.9 |
| Pancreas | 12.8 | 19.1 | 4.7 | 7.4 | 8.3 | 2.7 |
| Larynx | 2.4 | 3.5 | 2.0 | 1.0 | 1.1 | 1.0 |
| Lung and bronchus | 36.8 | 55.1 | 8.0 | 47.8 | 56.3 | 7.1 |
| Melanoma of the skin | 21.0 | 30.1 | 5.8 | 15.4 | 18.6 | 4.2 |
| Female breast | | | | 134.8 | 156.9 | 11.8 |
| Cervix uteri | | | | 3.0 | 3.8 | 1.9 |
| Corpus uterus and NOS | | | | 25.4 | 29.7 | 5.2 |
| Ovary | | | | 14.4 | 17.3 | 4.0 |
| Prostate | 136.8 | 182.5 | 13.7 | | | |
| Testis | 5.0 | 9.3 | 3.6 | | | |
| Urinary bladder (incl. in situ) | 32.2 | 49.4 | 7.6 | 9.8 | 11.5 | 3.2 |
| Kidney and renal pelvis | 19.6 | 28.0 | 5.5 | 9.8 | 11.6 | 3.2 |
| Brain and other nervous system | 5.2 | 6.7 | 2.6 | 5.2 | 6.4 | 2.5 |
| Thyroid | 10.4 | 13.9 | 3.8 | 19.8 | 26.8 | 5.3 |
| Hodgkin lymphoma | 2.0 | 2.8 | 1.7 | 3.0 | 4.5 | 2.3 |
| Non-Hodgkin lymphomas | 20.2 | 29.6 | 5.8 | 21.4 | 25.4 | 4.8 |
| Multiple myeloma | 5.8 | 8.4 | 3.1 | 5.8 | 6.6 | 2.4 |
| Leukemias | 12.4 | 18.1 | 4.5 | 7.6 | 9.0 | 2.9 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2011. Rates based on fewer than 4 cases per year are unstable and should be used with caution. NOS=Not otherwise specified.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence for Selected Areas, by Gender, 2005-2009**

**West Central
Westchester County**

| Site of cancer | Males | | | Females | | |
|---------------------------------|---------------|------------------|--------------|---------------|------------------|--------------|
| | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) |
| All Invasive Malignant Tumors | 467.0 | 545.8 | 22.1 | 482.0 | 434.8 | 17.4 |
| Oral cavity and pharynx | 11.6 | 12.9 | 3.3 | 4.2 | 3.7 | 1.6 |
| Esophagus | 4.6 | 5.7 | 2.3 | 2.6 | 2.1 | 1.1 |
| Stomach | 9.4 | 10.9 | 3.1 | 8.4 | 6.9 | 2.1 |
| Colorectal | 39.6 | 46.4 | 6.5 | 38.8 | 33.0 | 4.6 |
| Colon excluding rect | 28.6 | 33.9 | 5.6 | 29.0 | 24.7 | 4.0 |
| Rectum & rectosigmoi | 11.0 | 12.4 | 3.3 | 9.8 | 8.3 | 2.3 |
| Liver / intrahepatic bile duct | 7.8 | 8.9 | 2.8 | 2.2 | 1.9 | 1.1 |
| Pancreas | 15.0 | 17.8 | 4.0 | 13.2 | 10.7 | 2.6 |
| Larynx | 3.8 | 4.6 | 2.1 | 1.2 | 1.1 | 0.9 |
| Lung and bronchus | 43.4 | 53.3 | 7.1 | 57.6 | 50.3 | 5.8 |
| Melanoma of the skin | 25.4 | 30.0 | 5.2 | 20.0 | 18.9 | 3.7 |
| Female breast | | | | 146.2 | 135.5 | 9.8 |
| Cervix uteri | | | | 7.4 | 7.6 | 2.4 |
| Corpus uterus and NOS | | | | 33.0 | 29.1 | 4.4 |
| Ovary | | | | 14.2 | 12.8 | 3.0 |
| Prostate | 143.4 | 160.3 | 11.7 | | | |
| Testis | 3.8 | 4.9 | 2.2 | | | |
| Urinary bladder (incl. in situ) | 29.8 | 36.6 | 5.9 | 11.0 | 9.3 | 2.5 |
| Kidney and renal pelvis | 20.2 | 23.0 | 4.5 | 9.2 | 8.0 | 2.3 |
| Brain and other nervous system | 8.6 | 10.4 | 3.1 | 3.8 | 3.4 | 1.5 |
| Thyroid | 8.4 | 10.0 | 3.0 | 28.0 | 28.7 | 4.8 |
| Hodgkin lymphoma | 2.4 | 3.2 | 1.8 | 2.2 | 2.6 | 1.5 |
| Non-Hodgkin lymphomas | 26.4 | 31.2 | 5.3 | 21.8 | 19.5 | 3.7 |
| Multiple myeloma | 6.8 | 7.6 | 2.6 | 6.0 | 5.1 | 1.8 |
| Leukemias | 18.2 | 21.4 | 4.4 | 11.8 | 10.7 | 2.7 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2011. Rates based on fewer than 4 cases per year are unstable and should be used with caution. NOS=Not otherwise specified.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence for Selected Areas, by Gender, 2005-2009**

**East Central
Westchester County**

| Site of cancer | Males | | | Females | | |
|---------------------------------|---------------|------------------|--------------|---------------|------------------|--------------|
| | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) |
| All Invasive Malignant Tumors | 308.6 | 542.3 | 27.1 | 330.6 | 463.9 | 22.4 |
| Oral cavity and pharynx | 8.0 | 13.2 | 4.1 | 5.2 | 7.4 | 2.8 |
| Esophagus | 3.6 | 6.1 | 2.8 | 2.0 | 2.7 | 1.7 |
| Stomach | 4.2 | 7.6 | 3.3 | 4.0 | 5.3 | 2.3 |
| Colorectal | 27.4 | 47.6 | 8.0 | 31.0 | 40.6 | 6.4 |
| Colon excluding rect | 20.0 | 35.1 | 6.9 | 24.4 | 31.7 | 5.6 |
| Rectum & rectosigmoi | 7.4 | 12.6 | 4.1 | 6.6 | 9.0 | 3.1 |
| Liver / intrahepatic bile duct | 4.4 | 7.7 | 3.2 | 2.0 | 2.6 | 1.6 |
| Pancreas | 9.6 | 17.0 | 4.8 | 12.2 | 15.7 | 3.9 |
| Larynx | 2.8 | 4.9 | 2.6 | 1.0 | 1.3 | 1.1 |
| Lung and bronchus | 32.4 | 58.5 | 9.0 | 34.2 | 45.6 | 6.8 |
| Melanoma of the skin | 18.6 | 32.6 | 6.6 | 13.0 | 19.5 | 4.7 |
| Female breast | | | | 99.6 | 144.1 | 12.7 |
| Cervix uteri | | | | 3.8 | 5.6 | 2.5 |
| Corpus uterus and NOS | | | | 18.4 | 25.6 | 5.2 |
| Ovary | | | | 9.0 | 13.1 | 3.8 |
| Prostate | 95.0 | 165.5 | 14.9 | | | |
| Testis | 3.4 | 6.5 | 3.1 | | | |
| Urinary bladder (incl. in situ) | 22.6 | 40.7 | 7.5 | 10.4 | 13.7 | 3.7 |
| Kidney and renal pelvis | 9.8 | 16.6 | 4.6 | 5.4 | 7.3 | 2.8 |
| Brain and other nervous system | 4.2 | 7.5 | 3.2 | 4.6 | 7.0 | 2.9 |
| Thyroid | 6.4 | 11.1 | 3.8 | 17.0 | 27.8 | 5.9 |
| Hodgkin lymphoma | 1.4 | 2.6 | 1.9 | 3.0 | 5.5 | 2.8 |
| Non-Hodgkin lymphomas | 16.2 | 28.9 | 6.3 | 17.0 | 23.1 | 4.9 |
| Multiple myeloma | 3.6 | 6.2 | 2.9 | 3.2 | 4.2 | 2.1 |
| Leukemias | 10.2 | 18.0 | 4.9 | 8.8 | 12.3 | 3.6 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2011. Rates based on fewer than 4 cases per year are unstable and should be used with caution. NOS=Not otherwise specified.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence for Selected Areas, by Gender, 2005-2009**

**Southwest
Westchester County**

| Site of cancer | Males | | | Females | | |
|---------------------------------|---------------|------------------|--------------|---------------|------------------|--------------|
| | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) |
| All Invasive Malignant Tumors | 546.4 | 565.5 | 21.2 | 552.4 | 428.6 | 16.0 |
| Oral cavity and pharynx | 14.6 | 15.0 | 3.4 | 7.2 | 5.5 | 1.8 |
| Esophagus | 8.6 | 8.8 | 2.6 | 3.6 | 2.5 | 1.2 |
| Stomach | 13.6 | 14.0 | 3.3 | 9.6 | 7.1 | 2.0 |
| Colorectal | 53.4 | 55.5 | 6.7 | 53.8 | 38.9 | 4.6 |
| Colon excluding rect | 35.8 | 37.3 | 5.5 | 40.0 | 28.7 | 4.0 |
| Rectum & rectosigmoi | 17.6 | 18.2 | 3.8 | 13.8 | 10.2 | 2.4 |
| Liver / intrahepatic bile duct | 17.4 | 17.7 | 3.7 | 5.2 | 4.0 | 1.5 |
| Pancreas | 10.2 | 10.6 | 2.9 | 17.4 | 12.2 | 2.6 |
| Larynx | 8.8 | 9.1 | 2.7 | 2.0 | 1.4 | 0.9 |
| Lung and bronchus | 71.4 | 74.2 | 7.7 | 72.0 | 53.2 | 5.5 |
| Melanoma of the skin | 16.8 | 17.5 | 3.7 | 13.4 | 10.7 | 2.6 |
| Female breast | | | | 146.0 | 117.6 | 8.5 |
| Cervix uteri | | | | 11.0 | 9.9 | 2.6 |
| Corpus uterus and NOS | | | | 38.8 | 31.4 | 4.4 |
| Ovary | | | | 17.0 | 13.8 | 2.9 |
| Prostate | 152.8 | 156.1 | 11.1 | | | |
| Testis | 4.4 | 4.7 | 2.0 | | | |
| Urinary bladder (incl. in situ) | 36.8 | 38.4 | 5.5 | 15.6 | 11.1 | 2.5 |
| Kidney and renal pelvis | 21.6 | 22.3 | 4.2 | 13.0 | 9.9 | 2.4 |
| Brain and other nervous system | 8.4 | 8.7 | 2.6 | 6.0 | 4.8 | 1.7 |
| Thyroid | 5.4 | 5.6 | 2.1 | 18.0 | 16.7 | 3.5 |
| Hodgkin lymphoma | 2.4 | 2.6 | 1.5 | 3.8 | 3.4 | 1.5 |
| Non-Hodgkin lymphomas | 21.6 | 22.7 | 4.3 | 23.4 | 18.0 | 3.3 |
| Multiple myeloma | 12.8 | 13.4 | 3.3 | 9.0 | 6.7 | 2.0 |
| Leukemias | 17.4 | 18.2 | 3.8 | 16.8 | 12.7 | 2.7 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2011. Rates based on fewer than 4 cases per year are unstable and should be used with caution. NOS=Not otherwise specified.

**New York State Department of Health, New York State Cancer Registry
Cancer Incidence for Selected Areas, by Gender, 2005-2009**

**Southeast
Westchester County**

| Site of cancer | Males | | | Females | | |
|---------------------------------|---------------|------------------|--------------|---------------|------------------|--------------|
| | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) | Avg Ann Cases | Rate per 100,000 | 95% CI (+/-) |
| All Invasive Malignant Tumors | 536.4 | 600.2 | 22.7 | 561.0 | 462.7 | 17.1 |
| Oral cavity and pharynx | 11.8 | 13.1 | 3.3 | 7.0 | 5.7 | 1.9 |
| Esophagus | 6.2 | 6.8 | 2.4 | 3.8 | 2.9 | 1.3 |
| Stomach | 11.6 | 13.5 | 3.5 | 7.4 | 5.7 | 1.8 |
| Colorectal | 50.2 | 56.7 | 7.0 | 60.0 | 47.0 | 5.3 |
| Colon excluding rect | 35.8 | 40.7 | 6.0 | 45.2 | 35.0 | 4.6 |
| Rectum & rectosigmoi | 14.4 | 16.0 | 3.7 | 14.8 | 12.1 | 2.8 |
| Liver / intrahepatic bile duct | 12.8 | 13.9 | 3.4 | 4.2 | 3.5 | 1.5 |
| Pancreas | 13.6 | 14.9 | 3.5 | 16.2 | 12.1 | 2.6 |
| Larynx | 5.2 | 5.8 | 2.2 | 2.0 | 1.6 | 1.0 |
| Lung and bronchus | 61.4 | 69.5 | 7.8 | 62.6 | 50.4 | 5.6 |
| Melanoma of the skin | 18.0 | 20.7 | 4.3 | 15.6 | 13.2 | 2.9 |
| Female breast | | | | 161.4 | 136.9 | 9.4 |
| Cervix uteri | | | | 8.8 | 8.1 | 2.4 |
| Corpus uterus and NOS | | | | 36.2 | 30.0 | 4.4 |
| Ovary | | | | 15.6 | 13.2 | 2.9 |
| Prostate | 178.2 | 195.3 | 12.8 | | | |
| Testis | 5.4 | 6.5 | 2.5 | | | |
| Urinary bladder (incl. in situ) | 31.4 | 36.3 | 5.7 | 15.8 | 11.5 | 2.5 |
| Kidney and renal pelvis | 18.6 | 20.6 | 4.2 | 12.0 | 9.6 | 2.4 |
| Brain and other nervous system | 5.8 | 6.6 | 2.4 | 6.2 | 5.3 | 1.9 |
| Thyroid | 9.0 | 10.1 | 3.0 | 24.4 | 23.4 | 4.2 |
| Hodgkin lymphoma | 3.6 | 4.2 | 1.9 | 3.2 | 3.3 | 1.6 |
| Non-Hodgkin lymphomas | 22.4 | 25.5 | 4.7 | 23.8 | 19.2 | 3.4 |
| Multiple myeloma | 8.2 | 9.2 | 2.8 | 9.6 | 7.4 | 2.1 |
| Leukemias | 17.6 | 20.1 | 4.2 | 15.0 | 12.0 | 2.7 |

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2011. Rates based on fewer than 4 cases per year are unstable and should be used with caution. NOS=Not otherwise specified.