Table 12C 2008 Pediatric Nutrition Surveillance (1) New York

Page: 1 Run Date: 5/1/2009

Summary of Trends in Growth and Anemia Indicators Children Aged < 5 Years (2)

Summary of Trends in Growth and Anemia Indicators (3)													
	Bi	rthweight (4)	Height and Weight								Anemia Low Hb/Hct	
Year	Low High				Short Stature (5)	stature weight		>_2 Yrs Overweight (6)			(7)		
	Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% <u>></u> 95th	Number 85	% 5th-<95th	% <u>></u> 95th	Number	%	
2008	130,568	8.9	6.7	481,925	4.2	4.4	13.8	209,713	17.4	14.6	355,319	12.2	
2007	128,809	8.7	6.7	458,800	4.1	4.7	13.5	198,041	17.4	14.6	331,509	11.9	
2006	127,389	8.9	6.7	456,877	4.2	4.4	14.0	199,608	17.3	14.7	333,878	11.1	
2005	124,905	8.9	6.9	457,355	4.2	4.8	13.9	201,411	16.9	14.9	335,985	11.2	
2004	125,083	8.7	7.2	452,156	4.3	5.1	14.3	197,169	17.2	15.9	331,591	12.3	
2003	124,246	8.5	7.6	434,263	4.4	6.1	14.2	186,284	16.7	16.6	317,010	13.0	
2002	121,199	8.9	7.6	424,134	4.5	6.6	13.9	178,615	16.1	16.6	309,850	13.7	
2001	53,340	7.3	8.5	410,422	4.7	6.9	13.8	186,238	16.1	16.3	313,295	14.8	
2000	86,940	8.8	8.3	419,669	4.5	7.0	13.6	189,836	16.2	16.2	320,455	14.4	
1999	78,096	9.1	8.1	382,459	4.3	7.2	12.9	177,489	15.8	15.4	307,425	15.0	

⁽¹⁾ Reporting period is January 1 through December 31.



⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the analysis.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and >= 95th percentile category identifies obese children.

⁽⁷⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 12C 2008 Pediatric Nutrition Surveillance (1) New York

Page: 2 Run Date: 5/1/2009

Summary of Trends in Growth and Anemia Indicators Children Aged < 5 Years (2)

Summary of Trends in Growth and Anemia Indicators (3)													
	Birthweight (4)				Anemia								
Year	Low High				Short Stature (5)	Stature weight		> 2 Yrs Overweight (6)			Low Hb/Hct (7)		
	Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% <u>></u> 95th	Number 85	% 5th-<95th	% <u>></u> 95th	Number	%	
1998	93,741	9.2	8.4	448,130	4.5	7.2	12.6	207,479	15.7	14.7	365,051	14.8	
1997	89,596	9.5	8.0	446,926	4.7	7.1	12.5	208,237	15.4	14.3	368,183	14.5	
1996	84,519	9.3	8.4	444,700	4.9	7.0	12.8	206,651	15.4	14.5	370,406	14.3	
1995	83,759	9.3	8.0	444,469	5.1	7.5	12.2	203,087	14.9	13.7	368,371	15.5	
1994	87,220	9.5	8.2	438,633	5.4	7.6	12.4	194,146	14.8	13.8	359,006	15.5	
1993	88,482	9.6	8.1	424,037	5.8	7.8	12.1	178,048	14.5	13.2	343,167	15.8	
1992	85,006	9.8	8.1	384,075	6.2	8.1	12.5	152,322	14.5	13.7	307,475	16.5	
1991	82,991	10.4	7.9	365,011	6.5	8.7	12.1	139,919	13.8	13.5	289,960	17.6	
1990	77,756	10.2	8.1	350,273	6.6	8.7	12.2	140,224	14.1	13.1	280,382	15.6	
1989	73,393	10.4	8.1	354,388	6.7	8.4	11.7	149,413	13.5	12.3	285,460	14.7	

⁽¹⁾ Reporting period is January 1 through December 31.



⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the analysis.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and >= 95th percentile category identifies obese children.

⁽⁷⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

Percentages are not calculated if < 100 records are available for analysis after exclusions.