	Sum	mary of T	<b>Frends</b> ir	n Growt	h and An	emia In	dicato	rs by F	Race/Eth	nicity <sup>(3)</sup>			
Race/Ethnicity and Year		E	Birthweight	(4)		Anemia							
			Low	High		Short Stature (5)	Under- weight (6)	Obese (6)		≥ 2 Yrs Overweight (6)		Low HI	o/Hct
		Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% <u>&gt;</u> 95th	Number	% 85th - <95th	% ∩ <u>≥</u> 95th	Number	%
White, Not Hispanic	2008 2007 2006 2005 2004 2003 2002 2001 2000 1999	34,578 33,683 33,842 32,993 31,638 31,127 29,797 14,816 19,935 17,146	7.4 7.3 7.6 7.7 7.4 7.3 7.4 6.7 7.4 7.4 7.8	8.9 8.7 8.8 9.2 9.6 10.0 9.9 10.8 11.1 10.5	136,271 129,972 130,312 129,786 125,830 120,616 117,222 114,234 119,267 104,910	5.5 5.3 5.2 5.1 5.2 5.1 5.2 5.1 5.2 5.1 5.0	3.8 4.1 3.9 4.1 4.4 5.2 5.7 6.2 6.1 6.1	12.4 12.1 12.2 12.1 12.9 13.1 12.4 11.7 12.0 10.6	61,449 58,502 59,514 59,713 57,658 55,145 53,302 55,096 57,275 52,370	17.1 17.3 16.6 16.5 16.9 16.5 16.1 15.6 15.8 15.0	12.3 12.2 12.2 13.2 13.5 13.0 12.5 12.5 12.5 11.3	102,804 95,984 96,198 95,527 92,763 89,120 87,149 85,883 89,965 87,216	10.6 11.2 10.5 10.0 10.4 10.7 10.9 12.5 12.3 10.8
Black, Not Hispanic	2008 2007 2006 2005 2004 2003 2002 2001 2000 1999	33,033 32,315 32,375 32,233 32,676 32,605 32,485 13,837 26,576 24,528	12.3 12.0 12.0 12.6 12.0 11.4 12.2 9.3 11.8 12.0	5.2 5.3 5.0 5.4 5.6 6.0 5.9 6.8 6.7 6.6	113,090 109,510 112,103 114,447 113,894 110,169 108,674 108,045 116,332 110,992	4.2 4.0 4.1 4.3 4.4 4.5 4.8 4.6 4.5	5.9 6.1 5.8 6.0 6.2 7.3 7.6 8.0 8.1 8.2	12.8 12.6 13.0 12.9 13.5 13.1 12.9 12.8 12.4 12.0	47,300 46,328 48,134 49,919 48,863 46,144 44,426 48,237 51,094 50,047	15.8 15.5 15.8 15.4 15.8 15.7 14.8 15.4 15.2 15.1	12.9 12.8 13.0 13.3 14.5 14.9 14.9 14.6 14.5 13.6	80,653 77,770 81,088 83,690 83,011 79,882 79,148 82,372 88,228 88,228 88,420	17.3 16.0 14.8 15.0 16.6 17.6 18.7 19.4 18.6 19.8

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and >= 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis. \* Percentages are not calculated if < 100 records are available for analysis after exclusions.

	Sum	mary of 1	<b>Frends</b> ir	n Growt	h and An	emia In	dicato	rs by F	Race/Eth	nicity <sup>(3)</sup>			
Race/Ethnicity and Year		E	Birthweight	(4)		Anemia							
			Low			Short Stature (5)	Under- weight (6)			≥ 2 Yrs Overweight (6)		Low Ht (7)	o/Hct
		Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% <u>≥</u> 95th	Number	% 85th - <95th	% ⊨ <u>≥</u> 95th	Number	%
Hispanic	2008 2007 2006 2005 2004 2003 2002 2001 2000 1999	47,729 47,402 46,906 45,325 42,953 41,617 39,977 18,239 29,008 26,632	7.9 7.7 7.8 7.5 7.3 7.2 7.5 6.2 7.4 7.5	6.5 6.7 6.6 6.9 7.2 7.6 7.9 8.4 8.3 8.3	182,216 172,739 170,864 166,636 157,418 146,121 139,443 138,229 140,199 129,404	3.6 3.4 3.6 3.8 3.8 4.0 4.3 4.1 3.8	3.4 3.7 3.4 3.8 4.1 5.0 5.7 5.8 6.1 6.4	16.6 16.3 17.1 17.0 17.3 17.2 16.9 17.4 17.1 16.5	80,365 74,900 74,462 72,514 68,299 61,881 57,704 61,893 63,402 60,118	19.3 19.4 19.5 18.8 19.0 18.2 17.8 17.9 18.1 17.8	18.1 18.3 18.8 19.1 20.3 21.6 22.0 21.9 21.9 21.4	135,997 126,141 126,191 124,014 117,816 108,263 102,577 108,516 111,045 104,506	11.1 10.6 9.8 10.0 11.3 12.1 12.9 13.8 13.2 14.9
American Indian/Alaskan Native	2008 2007 2006 2005 2004 2003 2002 2001 2000 1999	1,341 1,310 1,363 826 382 357 327 165 429 316	10.0 10.9 9.2 8.5 8.4 12.9 9.5 5.5 10.7 11.1	6.1 6.6 7.4 5.3 8.9 9.5 8.9 8.5 8.4 7.9	4,190 3,553 3,124 2,170 1,478 1,389 1,390 1,782 2,904 1,642	4.6 4.9 4.3 4.1 4.2 5.2 4.5 4.0 4.4 4.9	7.3 8.1 6.9 6.5 6.4 6.3 7.7 7.4 8.5 11.3	13.0 12.1 13.8 12.7 13.2 15.3 14.2 13.6 12.3 10.7	1,467 1,043 980 839 646 634 704 1,100 1,592 813	14.5 15.9 16.0 17.6 17.2 14.7 16.2 14.9 16.0 13.0	14.8 14.3 16.4 13.2 16.1 18.9 16.8 15.2 13.8 11.6	2,807 2,251 1,882 1,396 1,097 1,038 1,078 1,445 2,286 1,326	12.0 11.6 11.8 11.1 10.9 13.5 10.9 13.6 13.1 12.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and >= 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis. \* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page: 2 Run Date: 5 / 1 / 09

	Sumi	mary of 1	<b>Frends</b> ir	n Growt	h and Ar	nemia In	dicato	rs by F	Race/Eth	nicity <sup>(3)</sup>			
Race/Ethnicity and Year		Birthweight (4)				Anemia							
			Low	High		Short Stature (5)	Under- weight (6)			≥ 2 Yrs Overweight (6)		Low H (7	b/Hct
		Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% <u>≥</u> 95th	Number	% 85th - <95th	% n <u>≥</u> 95th	Number	%
Asian/Pacific Islander	2008 2007 2006 2005 2004 2003 2002 2001 2000 1999	11,558 11,827 10,349 10,644 10,557 9,804 9,183 2,790 6,467 5,549	7.2 7.3 7.5 7.6 7.5 7.8 7.8 7.1 6.8 7.1	4.4 4.5 5.2 4.5 5.1 5.1 5.2 5.8 6.0 5.7	37,845 35,585 33,603 33,386 31,579 28,775 27,097 24,499 25,418 22,928	3.2 2.9 3.2 3.3 3.5 3.4 3.5 3.8 3.9 3.6	7.2 7.4 7.0 7.6 8.3 9.3 10.0 10.0 10.9 11.2	8.8 8.7 8.9 8.7 8.7 8.8 8.9 9.1 8.6 8.4	15,838 14,659 14,077 13,665 12,509 11,156 10,278 10,567 11,297 10,233	14.1 14.3 13.9 13.9 14.0 13.7 13.4 12.8 13.1 12.5	10.8 10.2 10.3 10.9 12.1 12.4 12.6 12.2 11.7	26,755 24,061 23,932 23,501 21,936 20,121 18,878 18,618 18,874 17,459	7.9 7.2 7.5 8.1 9.4 9.6 10.3 11.1 10.9 12.0
Multiple Races	2008 2007 2006 2005 2004 2003 2002 2001 2000 1999	2,311 2,250 2,535 1,524 879 690 557 0 0 0	10.0 9.2 8.8 9.8 11.7 8.0 8.6 *	8.2 6.7 6.3 7.5 7.1 9.0 6.3 *	8,273 7,398 6,834 4,390 2,617 1,719 1,038 0 0 0	4.8 4.6 4.5 4.8 4.1 5.6 4.8 *	4.3 5.1 5.3 6.3 5.3 7.7 10.9 *	13.6 12.5 12.6 12.8 13.5 12.9 10.6 *	3,287 2,601 2,432 1,622 823 318 121 0 0 0	16.1 16.0 15.7 17.1 13.1 20.8 18.2 *	12.7 12.7 13.4 12.1 14.8 14.8 13.2 *	6,288 5,291 4,575 3,004 1,813 1,097 553 0 0 0	13.7 13.2 12.4 11.8 12.6 14.9 15.4 *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and >= 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis. \* Percentages are not calculated if < 100 records are available for analysis after exclusions.

	Sum	mary of T	rends ir	n Growtl	h and Ar	nemia In	dicato	rs by F	Race/Eth	nicity <sup>(3)</sup>			
Race/Ethnicity and Year		E	Birthweight	(4)		Anemia							
		Low		High		Short Stature (5)	Under- weight (6)		≥ 2 Yrs Overweight (6)			Low Ht (7)	o/Hct
		Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% <u>&gt;</u> 95th	Number	% 85th - <95th	% n <u>≥</u> 95th	Number	%
All Other/Unknown	2008 2007 2006 2005 2004 2003 2002 2001 2000 1999	18 22 19 1,360 5,998 8,046 8,873 3,493 4,525 3,925	* 7.3 9.5 9.1 8.6 7.2 8.9 9.4	* * 6.3 7.6 7.6 7.5 8.4 8.2 8.5	40 43 37 6,540 19,340 25,474 29,270 23,633 15,549 12,583	* * 4.6 4.3 4.7 4.7 4.9 4.8 4.8	* * 6.9 6.6 7.3 7.7 8.2 7.9 7.7	* 13.5 13.4 13.2 13.5 12.6 12.2 10.9	7 8 9 3,139 8,371 11,006 12,080 9,345 5,176 3,908	* 15.9 16.5 15.9 15.2 14.1 15.5 14.7	* * 15.0 15.0 15.6 16.0 15.2 14.6 13.5	15 11 12 4,853 13,155 17,489 20,467 16,461 10,057 8,498	* * 11.2 12.0 12.8 13.8 14.6 16.0 15.7
Total All Race/Ethnic Groups	2008 2007 2006 2005 2004 2003 2002 2001 2000 1999	130,568 128,809 127,389 124,905 125,083 124,246 121,199 53,340 86,940 78,096	8.9 8.7 8.9 8.7 8.7 8.5 8.9 7.3 8.8 9.1	6.7 6.7 6.9 7.2 7.6 7.6 8.5 8.3 8.1	481,925 458,800 456,877 457,355 452,156 434,263 424,134 410,422 419,669 382,459	4.2 4.1 4.2 4.3 4.4 4.5 4.7 4.5 4.7 4.5 4.3	4.4 4.7 4.4 5.1 6.1 6.6 6.9 7.0 7.2	13.8 13.5 14.0 13.9 14.3 14.2 13.9 13.8 13.6 12.9	209,713 198,041 199,608 201,411 197,169 186,284 178,615 186,238 189,836 177,489	17.4 17.3 16.9 17.2 16.7 16.1 16.1 16.2 15.8	14.6 14.7 14.9 15.9 16.6 16.6 16.3 16.2 15.4	355,319 331,509 333,878 335,985 331,591 317,010 309,850 313,295 320,455 307,425	12.2 11.9 11.1 11.2 12.3 13.0 13.7 14.8 14.4 15.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and >= 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis. \* Percentages are not calculated if < 100 records are available for analysis after exclusions.