

Weekly Influenza Surveillance Report

The New York State Department of Health (NYSDOH) collects, compiles, and analyzes information on influenza activity year round in New York State (NYS) and produces this weekly report during the influenza season (October through the following May).¹


During the week ending May 12, 2018

- Influenza activity level was categorized as geographically widespread². This is the 23rd consecutive week that widespread activity has been reported.
- There were 457 laboratory-confirmed influenza reports, a 41% decrease over last week.
- Of the 1,569 specimens submitted to WHO/NREVSS laboratories, 86 (5.48%) were positive for influenza.
- Of the 111 specimens tested at Wadsworth Center, 102 were positive for influenza. 6 were influenza A(H1),
 67 were influenza A(H3), 26 were influenza B(Yamagata), and 3 were influenza B(Victoria).
- Reports of percent of patient visits for influenza-like illness (ILI3) from ILINet providers was 1.71%, which is below the regional baseline of 3.10%.
- The number of patients hospitalized with laboratory-confirmed influenza was 99 a 38% decrease over last week.
- There were no influenza-associated pediatric deaths reported this week. There have been six influenza-associated
 pediatric deaths reported this season.
- Preliminary results for influenza vaccine effectiveness (VE) are published on CDC's website at https://www.cdc.gov/mmwr/volumes/67/wr/mm6706a2.htm?s_cid=mm6706a2_w

Laboratory Reports of Influenza (including NYC)

All clinical laboratories that perform testing on residents of NYS report all positive influenza test results to NYSDOH.

- 54 counties reported cases this week.
- Incidence ranged from 0-14.67 cases/100,000 population.



¹ Information about influenza monitoring in New York City (NYC) is available from the NYC Department of Health and Mental Hygiene website at http://www.dc.gov/flu/weekly/. National influenza surveillance data is available on CDC's FluView website at http://www.cdc.gov/flu/weekly/.

² No Activity: No laboratory-confirmed cases of influenza reported to the NYSDOH.


Sporadic: Small numbers of lab-confirmed cases of influenza reported.


Local: Increased or sustained numbers of lab-confirmed cases of influenza reported in a single region of New York State; sporadic in rest of state. Regional: Increased or sustained numbers of lab-confirmed cases of influenza reported in at least two regions but in fewer than 31 of 62 counties. Widespread: Increased or sustained numbers of lab-confirmed cases of influenza reported in greater than 31 of the 62 counties. Increased or sustained is defined as 2 or more cases of laboratory-confirmed influenza per 100,000 population.


3 ILI = influenza-like illness, defined as temperature 100° F with cough and/or sore throat in the absence of a known cause other than influenza


Laboratory Reports of Influenza (including NYC)

Test results may identify influenza Type A, influenza Type B, or influenza without specifying Type A or B. Some tests only give a positive or negative result and cannot identify influenza type (not specified).












Laboratory Reports of Influenza (including NYC)

Data shown in the table represents the number of laboratory-confirmed cases by county for the current week, previous two weeks, and season-to-date totals.

				1					
County	28-Apr	/eek Endir 5-May	12-May	Season-To-Date					
Albany	25	20	11	1704					
Allegany	2	0	1	204					
Broome	27	18	4	2211					
Cattaraugus	0	1	2	489					
	21	10	3	1167					
Chautaugua									
Chautauqua	23	27	19	1414					
Chemung	13	6	6	484					
Chenango	3	4	1	567					
Clinton	8	6	5	634					
Columbia	0	2	2	334					
Cortland	4	2	1	593					
Delaware	3	2	0	313					
Dutchess	7	12	4	1591					
Erie	27	13	9	5168					
Essex	0	0	2	168					
Franklin	10	1	1	234					
Fulton	1	4	0	344					
Genesee	6	6	1	727					
	3	3	2						
Greene	0	1	0	231					
Hamilton				28					
Herkimer	7	8	5	794					
Jefferson	10	4	3	1300					
Lewis	6	2	1	390					
Livingston	11	6	4	692					
Madison	9	5	5	641					
Monroe	103	80	58	7364					
Montgomery	3	2	1	464					
Nassau	46	30	30	7866					
Niagara	9	0	0	869					
Oneida	57	45	32	3730					
Onondaga	70	40	30	3286					
Ontario	14	20	7	1374					
Orange	22	17	2	2416					
Orleans	4	1	1	401					
	19	23	8	1331					
Oswego									
Otsego	5	4	3	408					
Putnam	5	5	7	656					
Rensselaer	7	3	3	780					
Rockland	7	9	3	1408					
Saratoga	28	22	16	2160					
Schenectady	5	25	11	1961					
Schoharie	2	3	1	169					
Schuyler	0	0	0	51					
Seneca	5	3	2	300					
St. Lawrence	6	3	5	1040					
Steuben	6	5	2	608					
Suffolk	61	37	23	7615					
Sullivan	4	1	1	552					
Tioga	4	6	2	613					
	14	14							
Tompkins			2	1138					
Ulster	4	2	0	674					
Warren	3	1	0	227					
Washington	0	2	0	278					
Wayne	25	21	9	1407					
Westchester	59	44	19	8628					
Wyoming	3	1	1	284					
Yates	3	4	2	217					
Upstate Total	829	636	373	82697					
Bronx	47	42	25	11731					
Kings	50	28	11	11589					
New York	27	24	16	6245					
	71	32	30	13502					
Queens									
Richmond	26	7	2	2256					
NYC Total	221	133	84	45323					
Total	1050	769	457	128020					

World Health Organization (WHO) and National Respiratory & Enteric Virus Surveillance System (NREVSS) Collaborating Laboratories


Clinical virology laboratories, including the Wadsworth Center, that are WHO and/or NREVSS collaborating laboratories for influenza surveillance report weekly the number of respiratory specimens tested and the number positive for influenza types A and B to CDC. Some labs also report the influenza A subtype (H1 or H3) and influenza B lineage (Victoria or Yamagata). Because denominator data is provided, the weekly percentage of specimens testing positive for influenza is calculated.





Influenza Virus Types and Subtypes Identified at Wadsworth Center (excluding NYC)

Wadsworth Center, the NYSDOH public health laboratory, tests specimens from sources including, outpatient healthcare providers (ILINet) and hospitals (FluSurv-NET).

There are 2 common subtypes of influenza A viruses – H1 and H3. Each subtype has a slightly different genetic makeup. Wadsworth also identifies the lineage of influenza B specimens –Yamagata or Victoria. Rarely, an influenza virus is unable to have its subtype or lineage identified by the laboratory.









Influenza Antiviral Resistance Testing

The Wadsworth Center Virology Laboratory performs surveillance testing for antiviral drug resistance. 4

NYS Antiviral Resistance Testing Results on Samples Collected Season to date, 2017-18

	Samples tested	Oseltamivir Resistant Viruses, Number (%)	Zanamivir Resistant Viruses, Number (%)				
Influenza A (H1N1pdm09) i	136	0 (0.0)	0 (0.0)				
Influenza A (H3N2) ⁱⁱ	262	1 (0.4)	1 (0.4)				
Influenza B ⁱⁱⁱ	21	1 (4.7)	0 (0.0)				


- i. All samples tested by pyrosequencing for the H275Y variant in the neuraminidase gene which confers resistance to oseltamivir, and a subset tested by NA dideoxy sequencing for other variations known to cause, or suspected of causing, resistance to neuraminidase inhibitor drugs including zanamivir and oseltamivir.
- ii. All samples tested for oseltamivir resistance by pyrosequencing for E119V, R292K, and N294S in the neuraminidase gene (NA), and a subset tested by NA dideoxy sequencing for other variations known to cause, or suspected of causing, resistance to neuraminidase inhibitor drugs including zanamivir and oseltamivir.
- iii. Samples tested by whole gene dideoxysequencing of the neuraminidase gene. Sequence data reviewed for variations known to cause, or suspected of causing, resistance to neuraminidase inhibitor drugs including zanamivir and oseltamivir.

Outpatient Influenza-like Illness Surveillance Network (ILINet) (excluding NYC)

The NYSDOH works with ILINet healthcare providers who report the total number of patients seen and the total number of those with complaints of influenza-like illness (ILI) every week in an outpatient setting.

The CDC uses trends from past years to determine a regional baseline rate of doctors' office visits for ILI. For NYS, the regional baseline is currently 3%. Numbers above this regional baseline suggest high levels of illness consistent with influenza in the state.

Note that surrounding holiday weeks, it is not uncommon to notice a fluctuation in the ILI rate. This is a result of the different pattern of patient visits for non-urgent needs.




Emergency Department Visits for ILI-Syndromic Surveillance (excluding NYC)

Hospitals around NYS report the number of patients seen in their emergency departments with complaints of ILI. This is called syndromic surveillance.

An increase in visits to hospital emergency departments for ILI can be one sign that influenza has arrived in that part of NYS.

Syndromic surveillance does not reveal the actual cause of illness, but is thought to correlate with emergency department visits for influenza.





⁴Additional information regarding national antiviral resistance testing, as well as recommendations for antiviral treatment and chemoprophylaxis of influenza virus infection, can be found at http://www.cdc.gov/flu/weekly/.




Patients Hospitalized with Laboratory-Confirmed Influenza (including NYC)

Hospitals in NYS and NYC report the number of hospitalized patients with laboratory-confirmed Influenza to NYSDOH. 172 (94%) of 183 hospitals reported this week.








Influenza Hospitalization Surveillance Network (FluSurv-NET)

As part of the CDC's FluSurv-NET, the NYS Emerging Infections Program (EIP) conducts enhanced surveillance for hospitalized cases of laboratory-confirmed influenza among residents of 15 counties.⁵ Underlying health conditions are assessed through medical chart reviews for cases identified during the season.⁶





⁵Counties include, in the Capital District: Albany, Columbia, Greene, Montgomery, Rensselaer, Saratoga, Schenectady, and Schoharie; in the Western Region: Genesee, Livingston, Monroe, Ontario, Orleans, Wayne, and Yates ⁶Data are based on medical record reviews for 632 of 3275 hospitalized cases currently under investigation and should be considered preliminary.

Healthcare-associated Influenza Activity (including NYC)

Hospitals and nursing homes in NYS report outbreaks of influenza to the State. An outbreak in these settings is defined as one or more healthcare facility-associated case(s) of confirmed influenza in a patient or resident or two or more cases of influenza-like illness among healthcare workers and patients/residents of a facility on the same unit within 7 days. Outbreaks are considered confirmed only with positive laboratory testing.⁷

Week-to-Date (CDC week - 19)	Capital Region			Central Region			Metro Region			Wes	tern Re	gion	Statewide (Total)		
5/6/18 through 5/12/18	ACF	LTCF	Total	ACF	LTCF	Total	ACF	LTCF	Total	ACF	LTCF	Total	ACF	LTCF	Total
# Outbreaks* Lab-confirmed Influenza (any type)			0		1	1	1	1	2		1	1	1	3	4
# Outbreaks* viral respiratory illness**			0			0			0			0	0	0	0
Total # Outbreaks	0	0	0	0	1	1	1	1	2	0	1	1	1	3	4

Season-to-Date (CDC week - 19) Capital Region		Central Region			Metro Region			Western Region			Statewide (Total)				
9/29/17 through 5/12/18	ACF	LTCF	Total	ACF	LTCF	Total	ACF	LTCF	Total	ACF	LTCF	Total	ACF	LTCF	Total
# Outbreaks* Lab-confirmed Influenza (any type)	36	81	117	33	126	159	421	404	825	39	156	195	529	767	1296
# Outbreaks* viral respiratory illness**		7	7		16	16		22	22		6	6	0	51	51
Total # Outbreaks	36	88	124	33	142	175	421	426	847	39	162	201	529	818	1347

ACF - Article 28 Acute Care Facility

LTCF - Article 28 Long Term Care Facility


For information about the flu mask regulation and the current status of the Commissioner's declaration, please visit www.health.ny.gov/FluMaskReg

Pediatric influenza-associated deaths reported (including NYC)

Local health departments report pediatric influenza-associated deaths to NYSDOH.

Flu-associated deaths in children younger than 18 years old are nationally notifiable. Influenza-associated deaths in persons 18 years and older are not notifiable.

All pediatric flu-associated deaths included in this report are laboratory-confirmed.





^{*}Outbreaks are reported based on the onset date of symptoms in the first case

^{**} Includes outbreaks of suspect influenza and/or other viral upper respiratory pathogens