


County Level Prevalence of Smoking Among Adults

County level prevalence of current smoking was obtained from the New York State Expanded Behavioral Risk Factor Surveillance System (eBRFSS), a random-digit dial telephone survey designed to generate representative estimates of various chronic disease indicators and related behaviors for all adult residents in each county. Data was collected from April 2013 through March 2014.

The prevalence of cigarette smoking by county in New York State ranged from a low of 10.2% in Rockland County to a high of 30.6% in Cayuga County (see Table on page 2 for county estimates). Smoking rates were lowest among downstate counties. Higher rates were observed among NY's more rural counties.

Prevalence of Current Smoking Among Adults in NYS by County, eBRFSS 2013-2014



Note: Prevalence estimates for individual counties in NYC are not available. Prevalence reported is for the NYC area as a whole.



Bureau of Tobacco Control StatShot Vol. 9, No.3/June 2016

Prevalence of Current Smoking Among Adults in NYS by County, eBRFSS 2013-2014

County	Estimated Percent of Current Smokers (%)	[95%CI]	County	Estimated Percent of Current Smokers (%)	[95%CI]
Albany	16.3	[13.3-19.7]	Niagara	20.8	[16.2-26.2]
Allegany	26.8	[20.7-33.8]	Oneida	22.0	[17.4-27.4]
Broome	22.3	[18.1-27.2]	Onondaga	20.3	[16.8-24.3]
Cattaraugus	28.4	[21.2-36.8]	Ontario	17.6	[12.4-24.5]
Cayuga	30.6	[21.2-41.9]	Orange	15.7	[12.1-20.2]
Chautauqua	24.7	[19.0-31.5]	Orleans	25.7	[18.6-24.3]
Chemung	24.8	[17.4-34.2]	Oswego	28.0	[22.1-34.8]
Chenango	18.9	[13.9-25.1]	Otsego	26.3	[18.7-35.7]
Clinton	22.6	[17.4-28.7]	Putnam	13.9	[9.6-19.7]
Columbia	21.0	[15.6-27.5]	Rensselaer	23.8	[18.3-30.3]
Cortland	21.4	[14.2-30.8]	Rockland	10.2	[7.3-14.1]
Delaware	22.9	[16.7-30.4]	Saratoga	17.7	[13.8-22.4]
Dutchess	16.1	[11.5-22.0]	Schenectady	19.3	[15.5-23.9]
Erie	18.8	[15.9-22.0]	Schoharie	19.3	[14.4-25.4]
Essex	16.6	[12.0-22.7]	Schuyler	22.3	[15.6-30.9]
Franklin	27.0	[20.0-35.3]	Seneca	13.7	[9.3-19.7]
Fulton	29.0	[23.2-35.5]	St. Lawrence	19.5	[14.5-25.6]
Genesee	25.8	[19.1-34.0]	Steuben	23.7	[18.9-29.2]
Greene	24.5	[17.3-33.4]	Suffolk	14.4	[10.8-18.8]
Hamilton	19.0	[13.1-26.7]	Sullivan	24.5	[18.6-31.4]
Herkimer	25.8	[19.7-33.1]	Tioga	22.8	[16.8-30.0]
Jefferson	22.1	[16.6-28.7]	Tompkins	14.1	[8.9-21.6]
Lewis	14.9	[10.5-20.8]	Ulster	21.1	[15.2-28.6]
Livingston	23.0	[16.8-30.8]	Warren	18.7	[14.4-24.0]
Madison	17.0	[11.3-24.9]	Washington	21.0	[16.3-26.6]
Monroe	14.5	[11.9-17.6]	Wayne	24.5	[17.7-33.0]
Montgomery	23.4	[18.1-29.6]	Westchester	11.7	[9.0-15.2]
Nassau	12.7	[8.4-18.7]	Wyoming	21.6	[16.0-28.4]
New York City	13.3	[9.9-17.6]	Yates	13.8	[10.2-18.6]

Note: Prevalence estimates for individual counties in NYC are not available. Prevalence reported is for the NYC area as a whole.