

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State, Colorectal, All Races**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	5,680	34.2	33.3-35.1	2,764	40.4	38.9-42.1	2,916	30.1	29.0-31.2
1977	5,930	35.5	34.6-36.5	2,871	42.4	40.8-44.0	3,059	31.3	30.2-32.4
1978	5,948	35.5	34.5-36.4	2,914	42.7	41.1-44.4	3,034	30.9	29.8-32.0
1979	6,315	37.6	36.6-38.5	3,005	43.9	42.3-45.6	3,310	33.5	32.3-34.6
1980	6,434	37.9	37.0-38.9	3,122	45.5	43.9-47.2	3,312	33.1	31.9-34.2
1981	6,574	38.5	37.5-39.4	3,259	47.6	45.9-49.3	3,315	32.8	31.6-33.9
1982	6,673	38.8	37.9-39.8	3,263	47.2	45.5-48.9	3,410	33.4	32.3-34.6
1983	6,592	38.0	37.0-38.9	3,254	46.6	45.0-48.3	3,338	32.5	31.4-33.6
1984	6,491	37.5	36.6-38.4	3,146	45.3	43.7-47.0	3,345	32.5	31.3-33.6
1985	6,540	37.4	36.5-38.3	3,240	46.4	44.7-48.0	3,300	31.8	30.7-32.9
1986	6,389	36.4	35.5-37.3	3,102	44.0	42.4-45.6	3,287	31.6	30.5-32.7
1987	6,145	34.7	33.8-35.6	2,985	42.0	40.5-43.6	3,160	29.9	28.9-31.0
1988	5,975	33.7	32.8-34.5	2,870	40.3	38.8-41.9	3,105	29.3	28.3-30.4
1989	5,664	31.9	31.0-32.7	2,793	39.0	37.5-40.5	2,871	27.2	26.2-28.2
1990	5,532	30.8	30.0-31.7	2,732	37.7	36.2-39.2	2,800	26.2	25.2-27.2
1991	5,605	31.2	30.4-32.0	2,768	38.2	36.8-39.7	2,837	26.5	25.5-27.5
1992	5,257	29.1	28.3-29.9	2,551	34.8	33.4-36.2	2,706	25.3	24.3-26.3
1993	5,250	28.8	28.0-29.6	2,481	33.3	31.9-34.6	2,769	25.6	24.7-26.6
1994	5,707	31.1	30.3-32.0	2,751	37.1	35.7-38.5	2,956	27.3	26.3-28.3
1995	5,602	30.4	29.6-31.2	2,726	36.5	35.1-37.9	2,876	26.4	25.4-27.4
1996	5,693	30.7	29.9-31.5	2,779	36.7	35.3-38.1	2,914	26.6	25.6-27.6
1997	5,982	32.0	31.2-32.9	2,892	37.6	36.2-39.0	3,090	28.0	27.0-29.0
1998	6,129	32.5	31.7-33.3	2,933	37.6	36.2-39.0	3,196	28.9	27.9-30.0
1999	6,131	32.0	31.2-32.8	2,888	36.5	35.1-37.9	3,243	28.9	27.9-29.9
2000	6,127	31.6	30.8-32.4	2,978	36.9	35.6-38.3	3,149	27.7	26.7-28.7
2001	5,874	29.9	29.2-30.7	2,849	35.1	33.8-36.4	3,025	26.4	25.4-27.3
2002	5,997	30.2	29.4-31.0	2,853	34.3	33.1-35.6	3,144	27.1	26.2-28.1
2003	5,939	29.6	28.8-30.3	2,888	34.3	33.1-35.6	3,051	26.2	25.2-27.1
2004	5,606	27.6	26.9-28.4	2,655	30.9	29.8-32.1	2,951	25.1	24.2-26.1
2005	5,419	26.5	25.8-27.2	2,632	30.6	29.4-31.8	2,787	23.6	22.7-24.5
2006	5,192	25.2	24.5-25.9	2,501	28.7	27.6-29.8	2,691	22.6	21.8-23.5
2007	5,215	25.1	24.4-25.8	2,551	28.6	27.5-29.8	2,664	22.4	21.5-23.3
2008	5,142	24.4	23.7-25.1	2,506	27.7	26.6-28.8	2,636	21.9	21.1-22.8
2009	4,971	23.2	22.6-23.9	2,463	26.6	25.6-27.7	2,508	20.5	19.7-21.3
2010	5,058	23.4	22.8-24.1	2,493	26.5	25.5-27.6	2,565	21.0	20.1-21.8
2011	4,946	22.6	21.9-23.2	2,482	26.0	25.0-27.1	2,464	19.8	19.1-20.7
2012	4,816	21.6	21.0-22.2	2,430	24.8	23.8-25.8	2,386	19.0	18.2-19.8
2013	4,957	21.9	21.3-22.5	2,500	25.1	24.1-26.1	2,457	19.3	18.6-20.1
2014	5,037	21.9	21.3-22.6	2,553	25.0	24.1-26.1	2,484	19.4	18.6-20.2
2015	4,857	21.0	20.4-21.6	2,462	23.8	22.9-24.8	2,395	18.6	17.8-19.3
2016	5,242	22.3	21.7-23.0	2,731	26.1	25.1-27.1	2,511	19.3	18.5-20.1
2017	5,212	22.1	21.5-22.7	2,726	25.6	24.7-26.6	2,486	19.2	18.4-20.0
2018	5,172	21.5	20.9-22.1	2,640	24.6	23.6-25.6	2,532	18.9	18.2-19.7
2019	5,273	21.8	21.2-22.5	2,802	25.7	24.8-26.7	2,471	18.6	17.9-19.4
2020	4,863	20.1	19.5-20.7	2,590	23.6	22.7-24.5	2,273	17.1	16.4-17.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State, Colorectal, White Individuals**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	5,262	34.7	33.8-35.7	2,579	41.3	39.7-43.0	2,683	30.4	29.2-31.6
1977	5,401	35.6	34.6-36.6	2,654	42.9	41.2-44.6	2,747	31.0	29.8-32.2
1978	5,391	35.4	34.4-36.3	2,677	43.1	41.4-44.9	2,714	30.4	29.2-31.6
1979	5,719	37.6	36.6-38.6	2,735	44.0	42.3-45.8	2,984	33.4	32.2-34.6
1980	5,846	38.2	37.2-39.2	2,861	46.2	44.4-48.0	2,985	33.1	31.9-34.3
1981	5,983	39.0	38.0-40.0	2,984	48.3	46.5-50.1	2,999	33.1	31.9-34.3
1982	6,060	39.4	38.4-40.4	2,975	47.9	46.2-49.8	3,085	33.7	32.5-35.0
1983	5,962	38.5	37.5-39.5	2,981	47.8	46.0-49.6	2,981	32.5	31.4-33.8
1984	5,895	38.3	37.3-39.3	2,874	46.4	44.7-48.2	3,021	33.0	31.8-34.2
1985	5,895	38.1	37.1-39.1	2,965	47.6	45.9-49.4	2,930	31.9	30.8-33.1
1986	5,708	36.9	35.9-37.9	2,789	44.7	43.0-46.4	2,919	31.8	30.7-33.0
1987	5,533	35.6	34.6-36.5	2,714	43.2	41.5-44.9	2,819	30.6	29.4-31.7
1988	5,332	34.3	33.4-35.2	2,584	41.3	39.6-42.9	2,748	29.7	28.6-30.9
1989	5,006	32.3	31.4-33.2	2,474	39.5	38.0-41.2	2,532	27.4	26.4-28.5
1990	4,893	31.4	30.5-32.3	2,428	38.2	36.7-39.8	2,465	26.6	25.6-27.7
1991	4,936	31.7	30.8-32.6	2,480	39.3	37.7-40.9	2,456	26.5	25.5-27.6
1992	4,605	29.5	28.7-30.4	2,273	35.7	34.3-37.3	2,332	25.3	24.3-26.4
1993	4,585	29.3	28.5-30.2	2,179	33.8	32.3-35.3	2,406	26.1	25.0-27.1
1994	4,963	31.7	30.8-32.6	2,394	37.5	35.9-39.0	2,569	27.9	26.8-29.0
1995	4,800	30.6	29.8-31.5	2,361	36.9	35.4-38.4	2,439	26.3	25.3-27.4
1996	4,826	30.7	29.9-31.6	2,370	36.7	35.2-38.2	2,456	26.5	25.4-27.6
1997	5,079	32.2	31.3-33.1	2,458	37.6	36.1-39.1	2,621	28.1	27.0-29.2
1998	5,195	32.8	31.9-33.7	2,526	38.2	36.7-39.8	2,669	28.9	27.8-30.0
1999	5,131	32.0	31.2-32.9	2,469	37.1	35.6-38.6	2,662	28.4	27.3-29.5
2000	5,199	32.2	31.4-33.1	2,563	38.0	36.5-39.5	2,636	27.9	26.8-29.0
2001	4,892	30.1	29.3-31.0	2,385	35.2	33.8-36.6	2,507	26.5	25.5-27.6
2002	4,988	30.5	29.6-31.3	2,407	34.9	33.5-36.3	2,581	27.0	26.0-28.1
2003	4,931	30.0	29.1-30.8	2,421	34.9	33.5-36.3	2,510	26.3	25.2-27.3
2004	4,562	27.6	26.8-28.4	2,171	30.8	29.6-32.2	2,391	25.0	24.0-26.1
2005	4,342	26.1	25.3-26.9	2,117	30.0	28.8-31.4	2,225	23.3	22.3-24.3
2006	4,155	25.0	24.2-25.8	2,041	28.8	27.5-30.1	2,114	22.1	21.1-23.1
2007	4,218	25.2	24.5-26.0	2,072	28.8	27.5-30.0	2,146	22.6	21.6-23.6
2008	4,091	24.3	23.5-25.1	2,018	27.7	26.5-29.0	2,073	21.6	20.7-22.6
2009	3,962	23.2	22.5-24.0	1,971	26.6	25.5-27.9	1,991	20.5	19.6-21.4
2010	4,026	23.5	22.8-24.3	1,973	26.3	25.2-27.6	2,053	21.2	20.3-22.2
2011	3,912	22.6	21.9-23.4	1,965	25.9	24.8-27.1	1,947	20.0	19.1-21.0
2012	3,795	21.8	21.1-22.6	1,908	24.7	23.6-25.9	1,887	19.4	18.5-20.3
2013	3,910	22.3	21.6-23.0	1,986	25.4	24.3-26.6	1,924	19.7	18.8-20.6
2014	3,845	21.6	20.9-22.3	1,983	24.9	23.8-26.0	1,862	18.9	18.0-19.8
2015	3,748	21.1	20.4-21.8	1,886	23.6	22.6-24.8	1,862	18.9	18.0-19.8
2016	3,993	22.3	21.6-23.0	2,067	25.6	24.5-26.7	1,926	19.5	18.6-20.4
2017	3,962	22.1	21.4-22.8	2,055	25.1	24.0-26.2	1,907	19.5	18.6-20.5
2018	3,921	21.6	20.9-22.3	2,017	24.5	23.4-25.6	1,904	19.1	18.2-20.0
2019	3,947	21.8	21.1-22.6	2,133	25.9	24.8-27.1	1,814	18.3	17.5-19.3
2020	3,626	20.0	19.3-20.7	1,935	23.2	22.2-24.3	1,691	17.2	16.3-18.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State, Colorectal, Black Individuals**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	355	26.2	23.3-29.4	152	27.4	22.6-32.7	203	25.3	21.6-29.3
1977	473	35.1	31.7-38.8	185	36.1	30.4-42.5	288	34.6	30.4-39.2
1978	496	36.4	32.9-40.0	198	36.2	30.6-42.4	298	36.4	32.1-41.1
1979	532	36.8	33.5-40.4	229	40.1	34.5-46.4	303	34.7	30.7-39.1
1980	534	35.0	31.9-38.4	227	38.2	32.8-44.3	307	33.3	29.4-37.4
1981	525	34.1	31.0-37.4	236	40.9	35.2-47.2	289	30.4	26.9-34.3
1982	547	33.2	30.3-36.3	253	40.2	34.9-46.1	294	29.1	25.8-32.8
1983	567	33.4	30.5-36.4	238	37.7	32.4-43.4	329	31.4	28.0-35.1
1984	532	31.3	28.5-34.2	236	35.5	30.7-40.8	296	28.6	25.4-32.2
1985	575	32.6	29.9-35.6	247	39.0	33.7-44.8	328	29.7	26.4-33.2
1986	614	33.9	31.1-36.9	269	38.4	33.5-43.8	345	31.2	27.9-34.8
1987	541	28.7	26.2-31.3	234	33.8	29.2-39.0	307	26.0	23.1-29.2
1988	574	30.2	27.7-32.9	245	33.9	29.4-38.9	329	28.1	25.0-31.3
1989	569	28.7	26.3-31.3	272	37.4	32.6-42.6	297	24.2	21.5-27.2
1990	557	28.2	25.8-30.7	249	34.0	29.5-38.9	308	24.9	22.1-27.9
1991	568	28.3	26.0-30.9	244	31.4	27.3-36.0	324	26.3	23.4-29.3
1992	559	26.7	24.5-29.1	226	28.8	24.8-33.2	333	25.8	23.1-28.8
1993	557	26.6	24.3-28.9	247	31.3	27.1-35.9	310	24.0	21.3-26.8
1994	612	28.7	26.4-31.1	276	34.5	30.2-39.2	336	25.4	22.7-28.3
1995	645	29.0	26.8-31.4	292	35.7	31.4-40.5	353	25.6	23.0-28.4
1996	722	32.2	29.8-34.7	325	38.1	33.7-42.9	397	28.8	26.0-31.8
1997	749	32.5	30.1-34.9	354	40.9	36.4-45.8	395	27.8	25.1-30.7
1998	775	32.9	30.6-35.4	321	35.4	31.3-39.8	454	31.4	28.6-34.5
1999	827	33.6	31.3-36.0	337	35.3	31.3-39.5	490	32.5	29.7-35.6
2000	719	28.4	26.3-30.6	315	33.0	29.2-37.1	404	25.9	23.5-28.6
2001	777	30.2	28.1-32.4	348	36.5	32.5-40.8	429	27.0	24.5-29.7
2002	798	30.4	28.2-32.6	332	33.1	29.4-37.1	466	28.9	26.3-31.7
2003	790	29.0	27.0-31.1	347	33.4	29.7-37.4	443	26.8	24.3-29.4
2004	810	29.2	27.2-31.4	365	34.0	30.3-37.9	445	26.4	24.0-29.0
2005	821	29.1	27.1-31.2	374	34.0	30.4-37.9	447	26.2	23.8-28.7
2006	794	27.6	25.7-29.7	343	30.1	26.8-33.7	451	26.0	23.6-28.5
2007	759	25.9	24.1-27.9	367	31.2	27.9-34.7	392	22.4	20.2-24.8
2008	810	27.1	25.2-29.1	374	31.4	28.1-35.0	436	24.3	22.0-26.7
2009	754	24.7	22.9-26.6	362	29.2	26.1-32.6	392	21.5	19.4-23.8
2010	758	24.2	22.5-26.0	361	29.3	26.2-32.7	397	21.2	19.2-23.5
2011	762	23.7	22.0-25.4	370	28.5	25.5-31.7	392	20.4	18.4-22.5
2012	750	22.6	21.0-24.3	377	28.2	25.3-31.3	373	18.9	17.0-21.0
2013	751	22.1	20.5-23.7	356	25.4	22.8-28.4	395	19.6	17.7-21.7
2014	845	24.4	22.8-26.2	376	26.4	23.7-29.3	469	23.1	21.0-25.3
2015	781	21.7	20.2-23.3	399	27.0	24.3-29.9	382	18.0	16.2-20.0
2016	900	24.7	23.0-26.4	473	31.9	29.0-35.0	427	19.8	18.0-21.8
2017	829	22.6	21.0-24.2	440	29.5	26.7-32.5	389	18.1	16.3-20.0
2018	837	22.0	20.5-23.6	404	25.9	23.3-28.6	433	19.3	17.5-21.3
2019	887	23.1	21.5-24.7	438	27.9	25.3-30.7	449	19.9	18.0-21.8
2020	786	20.2	18.8-21.7	400	24.6	22.2-27.2	386	17.1	15.4-19.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York City, Colorectal, All Races**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	2,453	33.2	31.8-34.6	1,175	39.4	37.0-41.8	1,278	29.3	27.6-31.0
1977	2,542	34.8	33.5-36.3	1,195	41.3	38.9-43.8	1,347	30.9	29.2-32.6
1978	2,564	35.1	33.7-36.5	1,242	42.8	40.3-45.3	1,322	30.5	28.8-32.2
1979	2,716	37.5	36.1-39.0	1,289	45.0	42.5-47.7	1,427	32.9	31.2-34.7
1980	2,785	38.5	37.1-40.0	1,339	47.0	44.5-49.7	1,446	33.4	31.6-35.2
1981	2,704	37.5	36.1-39.0	1,296	45.8	43.3-48.5	1,408	32.3	30.6-34.1
1982	2,754	38.2	36.7-39.6	1,328	47.4	44.8-50.1	1,426	32.6	30.9-34.4
1983	2,700	37.1	35.7-38.5	1,299	45.9	43.3-48.5	1,401	32.0	30.3-33.8
1984	2,580	35.7	34.3-37.2	1,233	44.1	41.6-46.7	1,347	30.8	29.1-32.5
1985	2,601	35.7	34.4-37.2	1,294	45.9	43.4-48.6	1,307	29.8	28.2-31.5
1986	2,652	36.5	35.1-37.9	1,313	46.3	43.7-48.9	1,339	30.5	28.8-32.2
1987	2,391	32.7	31.4-34.1	1,127	39.8	37.5-42.3	1,264	28.4	26.8-30.0
1988	2,366	32.7	31.4-34.1	1,105	39.5	37.2-42.0	1,261	28.5	27.0-30.2
1989	2,199	30.5	29.3-31.8	1,053	37.3	35.1-39.7	1,146	26.3	24.8-27.9
1990	2,153	29.9	28.6-31.2	1,025	36.3	34.1-38.7	1,128	25.7	24.2-27.2
1991	2,197	30.8	29.6-32.2	1,070	38.8	36.5-41.3	1,127	25.9	24.4-27.5
1992	2,021	28.5	27.2-29.7	902	32.5	30.4-34.7	1,119	26.1	24.5-27.7
1993	1,973	27.7	26.5-28.9	905	32.0	29.9-34.2	1,068	24.8	23.3-26.3
1994	2,272	31.8	30.5-33.1	1,049	37.3	35.1-39.7	1,223	28.3	26.7-30.0
1995	2,268	31.7	30.4-33.0	1,063	37.9	35.6-40.3	1,205	27.8	26.3-29.4
1996	2,299	32.0	30.7-33.4	1,097	38.3	36.0-40.7	1,202	27.8	26.2-29.4
1997	2,415	33.5	32.1-34.8	1,105	38.5	36.2-40.9	1,310	30.0	28.4-31.7
1998	2,498	34.2	32.9-35.6	1,134	39.0	36.7-41.3	1,364	31.2	29.6-32.9
1999	2,451	32.9	31.6-34.3	1,112	37.3	35.1-39.5	1,339	30.2	28.6-31.8
2000	2,448	32.3	31.1-33.7	1,175	38.5	36.3-40.8	1,273	28.1	26.6-29.7
2001	2,364	30.9	29.7-32.2	1,122	36.7	34.5-38.9	1,242	27.2	25.7-28.8
2002	2,397	30.9	29.7-32.2	1,114	35.3	33.2-37.4	1,283	27.8	26.3-29.4
2003	2,380	30.5	29.3-31.7	1,141	36.2	34.1-38.4	1,239	26.7	25.2-28.3
2004	2,232	28.3	27.2-29.5	1,041	32.4	30.4-34.5	1,191	25.5	24.1-27.0
2005	2,226	28.1	26.9-29.3	1,056	32.8	30.8-34.9	1,170	25.0	23.6-26.5
2006	2,112	26.6	25.4-27.7	985	30.2	28.3-32.2	1,127	24.1	22.7-25.5
2007	2,053	25.5	24.4-26.6	999	29.9	28.0-31.8	1,054	22.3	21.0-23.7
2008	2,017	24.9	23.8-26.0	972	28.6	26.8-30.5	1,045	22.1	20.8-23.5
2009	2,004	24.4	23.3-25.5	984	28.5	26.7-30.4	1,020	21.2	19.9-22.6
2010	1,950	23.4	22.3-24.5	972	27.6	25.8-29.4	978	20.3	19.0-21.6
2011	1,876	22.0	21.0-23.1	944	26.3	24.6-28.1	932	18.9	17.7-20.1
2012	1,849	21.2	20.2-22.2	937	25.2	23.6-26.9	912	18.2	17.0-19.4
2013	1,976	22.3	21.4-23.4	969	25.7	24.1-27.4	1,007	19.8	18.5-21.1
2014	1,978	21.9	20.9-22.9	1,002	25.5	23.9-27.2	976	19.1	17.9-20.4
2015	1,871	20.4	19.5-21.4	952	24.0	22.5-25.6	919	17.6	16.4-18.8
2016	2,053	22.1	21.1-23.0	1,093	27.2	25.6-28.9	960	18.2	17.1-19.4
2017	2,046	21.7	20.7-22.6	1,059	25.5	24.0-27.2	987	18.7	17.5-19.9
2018	1,995	20.7	19.8-21.7	1,024	24.5	23.0-26.1	971	17.7	16.6-18.9
2019	2,122	22.0	21.0-23.0	1,127	26.5	25.0-28.1	995	18.4	17.3-19.6
2020	1,868	19.3	18.4-20.2	987	23.0	21.6-24.5	881	16.2	15.2-17.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York City, Colorectal, White Individuals**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	2,141	34.2	32.7-35.7	1,030	40.5	37.9-43.1	1,111	30.1	28.3-32.0
1977	2,138	34.8	33.3-36.4	1,031	42.0	39.3-44.7	1,107	30.4	28.5-32.3
1978	2,146	35.1	33.6-36.6	1,071	43.9	41.2-46.8	1,075	29.6	27.8-31.5
1979	2,261	37.6	36.1-39.3	1,089	45.6	42.9-48.5	1,172	32.7	30.8-34.7
1980	2,344	39.6	37.9-41.2	1,147	49.0	46.1-52.0	1,197	33.8	31.9-35.8
1981	2,261	38.6	36.9-40.2	1,099	47.6	44.7-50.6	1,162	32.8	30.8-34.8
1982	2,290	39.4	37.7-41.1	1,105	48.7	45.8-51.8	1,185	33.7	31.8-35.8
1983	2,206	37.8	36.2-39.4	1,086	47.6	44.7-50.6	1,120	31.9	30.0-33.9
1984	2,122	37.1	35.6-38.8	1,024	46.1	43.2-49.1	1,098	31.9	30.0-33.9
1985	2,096	36.6	35.0-38.3	1,076	48.1	45.2-51.2	1,020	29.6	27.8-31.6
1986	2,112	37.4	35.8-39.1	1,068	48.2	45.3-51.3	1,044	30.5	28.7-32.5
1987	1,914	34.0	32.5-35.6	921	41.8	39.1-44.6	993	29.1	27.2-31.0
1988	1,888	34.3	32.7-35.9	897	41.8	39.1-44.7	991	29.6	27.7-31.6
1989	1,720	31.8	30.3-33.4	822	38.8	36.1-41.6	898	27.2	25.4-29.2
1990	1,660	30.8	29.3-32.3	796	37.4	34.8-40.1	864	26.2	24.5-28.1
1991	1,678	31.7	30.2-33.3	849	41.2	38.4-44.1	829	25.6	23.9-27.5
1992	1,541	29.7	28.3-31.3	710	34.7	32.2-37.4	831	26.5	24.7-28.5
1993	1,476	28.8	27.3-30.3	685	33.1	30.7-35.7	791	25.7	23.9-27.6
1994	1,719	33.6	32.0-35.2	795	39.0	36.3-41.8	924	30.0	28.1-32.1
1995	1,658	32.8	31.3-34.5	784	39.1	36.4-42.0	874	28.7	26.8-30.7
1996	1,651	33.0	31.4-34.6	798	39.5	36.8-42.4	853	28.3	26.4-30.3
1997	1,715	34.2	32.6-35.9	774	38.5	35.8-41.3	941	31.1	29.1-33.2
1998	1,792	35.9	34.2-37.6	833	41.2	38.5-44.1	959	32.4	30.4-34.6
1999	1,697	33.7	32.1-35.4	808	39.5	36.8-42.3	889	29.8	27.9-31.9
2000	1,748	34.6	33.0-36.3	868	42.0	39.2-44.9	880	29.1	27.1-31.1
2001	1,630	32.1	30.5-33.7	793	38.4	35.7-41.2	837	27.9	26.0-29.9
2002	1,659	32.6	31.0-34.2	786	37.5	34.9-40.2	873	28.7	26.8-30.7
2003	1,636	32.2	30.6-33.8	808	38.8	36.2-41.6	828	27.5	25.6-29.4
2004	1,462	28.7	27.3-30.3	713	34.1	31.6-36.7	749	24.9	23.1-26.8
2005	1,435	28.3	26.9-29.9	691	33.1	30.7-35.7	744	25.1	23.3-27.1
2006	1,340	26.8	25.4-28.3	659	31.7	29.3-34.2	681	23.3	21.5-25.1
2007	1,311	26.2	24.8-27.6	644	30.6	28.3-33.1	667	22.9	21.2-24.8
2008	1,252	25.2	23.8-26.7	626	29.6	27.3-32.1	626	21.9	20.2-23.7
2009	1,266	25.3	23.9-26.7	630	29.6	27.3-32.0	636	21.9	20.2-23.8
2010	1,199	24.0	22.6-25.4	604	28.2	26.0-30.6	595	20.6	18.9-22.4
2011	1,124	22.1	20.8-23.5	575	26.4	24.3-28.7	549	18.8	17.2-20.5
2012	1,090	21.3	20.0-22.6	546	24.5	22.4-26.6	544	18.6	17.0-20.3
2013	1,203	23.3	22.0-24.7	599	26.8	24.7-29.1	604	20.5	18.9-22.3
2014	1,129	21.6	20.4-23.0	602	26.1	24.0-28.3	527	18.1	16.5-19.7
2015	1,073	20.5	19.2-21.7	537	23.4	21.4-25.5	536	18.1	16.5-19.8
2016	1,124	21.3	20.0-22.6	603	25.9	23.9-28.1	521	17.8	16.3-19.5
2017	1,155	21.7	20.4-23.0	588	24.5	22.5-26.6	567	19.4	17.7-21.1
2018	1,115	20.7	19.5-22.0	582	24.3	22.3-26.4	533	17.8	16.3-19.5
2019	1,172	22.0	20.8-23.4	650	27.1	25.0-29.3	522	17.7	16.1-19.3
2020	1,012	18.9	17.7-20.1	540	22.3	20.4-24.3	472	15.9	14.5-17.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York City, Colorectal, Black Individuals**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	277	26.4	23.0-30.0	125	30.7	24.9-37.3	152	23.7	19.8-28.1
1977	372	36.3	32.3-40.6	148	38.7	31.8-46.5	224	34.9	30.1-40.2
1978	380	36.9	32.9-41.2	147	36.7	30.2-44.2	233	37.0	32.1-42.4
1979	405	36.5	32.7-40.5	165	39.2	32.7-46.5	240	35.0	30.4-40.0
1980	405	34.2	30.7-37.9	168	35.9	30.1-42.5	237	33.0	28.7-37.8
1981	391	32.5	29.1-36.1	169	38.0	31.7-45.0	222	29.7	25.7-34.0
1982	421	32.8	29.5-36.2	198	41.1	35.0-47.9	223	27.9	24.2-32.0
1983	447	33.6	30.4-37.1	189	39.6	33.5-46.5	258	30.9	27.1-35.1
1984	407	30.9	27.8-34.2	180	35.3	29.9-41.4	227	28.1	24.4-32.1
1985	452	32.8	29.6-36.1	195	40.0	33.9-46.7	257	29.4	25.8-33.3
1986	484	34.1	31.0-37.4	208	38.6	33.0-44.8	276	31.5	27.8-35.6
1987	415	28.4	25.6-31.4	173	33.3	28.1-39.2	242	26.0	22.7-29.5
1988	426	29.2	26.3-32.2	179	33.3	28.1-39.1	247	26.9	23.5-30.5
1989	413	27.0	24.4-29.9	194	35.1	29.9-40.9	219	22.9	19.9-26.2
1990	430	28.3	25.6-31.2	189	34.6	29.4-40.4	241	25.0	21.8-28.4
1991	437	28.6	25.9-31.5	186	32.6	27.6-38.0	251	26.2	23.0-29.7
1992	412	26.0	23.4-28.7	155	27.0	22.5-32.1	257	25.7	22.6-29.1
1993	417	26.5	23.9-29.2	182	32.5	27.4-38.0	235	23.5	20.6-26.8
1994	447	27.6	25.0-30.3	191	32.5	27.7-37.8	256	25.0	22.0-28.3
1995	482	28.6	26.1-31.4	217	35.9	30.9-41.4	265	24.9	22.0-28.2
1996	536	31.7	29.0-34.6	231	37.1	32.1-42.6	305	28.8	25.6-32.2
1997	573	33.3	30.6-36.2	261	42.0	36.6-47.9	312	28.8	25.7-32.2
1998	581	33.0	30.4-35.9	233	35.5	30.8-40.6	348	31.6	28.3-35.1
1999	616	33.3	30.7-36.1	239	34.8	30.2-39.7	377	32.6	29.3-36.0
2000	524	27.6	25.2-30.1	222	32.0	27.7-36.8	302	25.3	22.5-28.3
2001	574	29.8	27.4-32.4	241	34.4	30.0-39.4	333	27.6	24.7-30.7
2002	578	29.4	27.0-31.9	243	32.9	28.7-37.6	335	27.4	24.5-30.5
2003	576	28.5	26.2-30.9	237	31.5	27.4-36.0	339	27.0	24.2-30.0
2004	579	28.2	25.9-30.6	235	30.6	26.6-35.0	344	26.9	24.1-29.9
2005	594	28.5	26.2-30.9	257	32.8	28.7-37.3	337	26.1	23.4-29.1
2006	574	27.3	25.0-29.6	231	29.2	25.4-33.5	343	26.2	23.5-29.1
2007	548	25.7	23.5-27.9	263	31.8	27.8-36.1	285	21.8	19.3-24.5
2008	581	26.6	24.5-28.9	260	30.4	26.6-34.5	321	24.1	21.5-26.9
2009	539	24.5	22.4-26.7	253	29.2	25.6-33.2	286	21.3	18.9-24.0
2010	547	24.2	22.2-26.3	249	28.5	24.9-32.5	298	21.8	19.3-24.4
2011	541	23.1	21.2-25.2	253	27.6	24.1-31.4	288	20.2	17.9-22.7
2012	543	22.8	20.9-24.8	271	29.3	25.8-33.2	272	18.7	16.5-21.1
2013	548	22.3	20.5-24.3	252	25.6	22.4-29.1	296	20.1	17.8-22.5
2014	600	24.1	22.2-26.2	265	26.5	23.3-30.1	335	22.7	20.3-25.3
2015	544	21.3	19.5-23.2	277	27.3	24.0-30.8	267	17.3	15.3-19.6
2016	657	25.1	23.2-27.2	336	32.4	28.9-36.1	321	20.4	18.2-22.9
2017	568	21.7	19.9-23.6	289	27.9	24.7-31.5	279	17.7	15.6-20.0
2018	573	21.2	19.5-23.1	274	25.1	22.1-28.3	299	18.5	16.4-20.8
2019	614	22.6	20.8-24.5	299	27.7	24.5-31.1	315	19.3	17.2-21.7
2020	528	19.4	17.7-21.2	254	22.7	19.9-25.7	274	17.1	15.1-19.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State Excl New York City, Colorectal, All Races**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	3,227	34.9	33.7-36.2	1,589	41.2	39.1-43.4	1,638	30.7	29.2-32.3
1977	3,388	36.1	34.9-37.4	1,676	43.2	41.1-45.5	1,712	31.5	30.1-33.1
1978	3,384	35.8	34.6-37.0	1,672	42.7	40.5-44.9	1,712	31.2	29.7-32.7
1979	3,598	37.6	36.4-38.9	1,716	43.1	41.0-45.3	1,882	33.8	32.3-35.4
1980	3,649	37.5	36.3-38.7	1,783	44.3	42.2-46.5	1,866	32.9	31.4-34.4
1981	3,870	39.2	38.0-40.5	1,963	48.9	46.7-51.3	1,907	33.1	31.6-34.6
1982	3,919	39.4	38.1-40.6	1,935	47.0	44.8-49.2	1,984	34.0	32.6-35.6
1983	3,892	38.6	37.4-39.9	1,955	47.1	44.9-49.3	1,937	32.9	31.4-34.4
1984	3,911	38.7	37.5-40.0	1,913	46.1	43.9-48.3	1,998	33.7	32.2-35.2
1985	3,936	38.6	37.4-39.8	1,946	46.6	44.4-48.8	1,990	33.3	31.8-34.8
1986	3,737	36.4	35.3-37.6	1,789	42.4	40.4-44.5	1,948	32.4	31.0-33.9
1987	3,753	36.1	34.9-37.2	1,858	43.4	41.4-45.6	1,895	31.1	29.7-32.6
1988	3,609	34.3	33.2-35.4	1,765	40.7	38.8-42.8	1,844	29.9	28.6-31.4
1989	3,465	32.8	31.7-33.9	1,740	40.2	38.2-42.2	1,725	27.8	26.5-29.2
1990	3,379	31.4	30.4-32.5	1,707	38.6	36.7-40.5	1,672	26.6	25.3-27.9
1991	3,408	31.4	30.3-32.5	1,698	37.7	35.9-39.6	1,710	26.9	25.6-28.2
1992	3,236	29.5	28.5-30.5	1,649	36.2	34.4-38.1	1,587	24.8	23.6-26.0
1993	3,277	29.5	28.5-30.5	1,576	34.1	32.4-35.8	1,701	26.2	25.0-27.5
1994	3,435	30.8	29.7-31.8	1,702	37.1	35.3-38.9	1,733	26.6	25.3-27.9
1995	3,334	29.6	28.6-30.6	1,663	35.7	33.9-37.5	1,671	25.4	24.2-26.7
1996	3,394	29.9	28.9-31.0	1,682	35.8	34.1-37.6	1,712	25.8	24.6-27.1
1997	3,567	31.1	30.1-32.2	1,787	37.0	35.3-38.8	1,780	26.7	25.4-28.0
1998	3,631	31.4	30.4-32.4	1,799	36.7	35.0-38.5	1,832	27.4	26.1-28.7
1999	3,680	31.4	30.4-32.4	1,776	36.1	34.4-37.8	1,904	28.0	26.7-29.3
2000	3,679	31.0	30.0-32.1	1,803	36.0	34.3-37.7	1,876	27.3	26.1-28.6
2001	3,509	29.3	28.3-30.3	1,726	34.1	32.5-35.8	1,783	25.8	24.6-27.1
2002	3,600	29.7	28.7-30.7	1,739	33.7	32.1-35.3	1,861	26.6	25.4-27.9
2003	3,559	29.0	28.0-30.0	1,747	33.3	31.7-34.9	1,812	25.7	24.6-27.0
2004	3,374	27.2	26.3-28.1	1,614	30.1	28.6-31.6	1,760	24.8	23.7-26.0
2005	3,193	25.4	24.5-26.3	1,576	29.2	27.8-30.7	1,617	22.6	21.5-23.8
2006	3,080	24.4	23.5-25.2	1,516	27.8	26.4-29.2	1,564	21.6	20.6-22.8
2007	3,162	24.8	23.9-25.7	1,552	27.9	26.5-29.3	1,610	22.4	21.3-23.5
2008	3,125	24.0	23.2-24.9	1,534	27.2	25.8-28.6	1,591	21.7	20.6-22.8
2009	2,967	22.5	21.7-23.4	1,479	25.6	24.3-27.0	1,488	20.0	18.9-21.0
2010	3,108	23.4	22.6-24.2	1,521	25.9	24.6-27.3	1,587	21.3	20.3-22.4
2011	3,070	22.9	22.1-23.7	1,538	25.8	24.5-27.2	1,532	20.5	19.5-21.6
2012	2,967	21.8	21.0-22.6	1,493	24.5	23.2-25.8	1,474	19.5	18.5-20.6
2013	2,981	21.6	20.9-22.5	1,531	24.7	23.4-26.0	1,450	19.1	18.1-20.1
2014	3,059	22.0	21.2-22.8	1,551	24.8	23.6-26.1	1,508	19.6	18.6-20.6
2015	2,986	21.3	20.6-22.2	1,510	23.7	22.5-25.0	1,476	19.3	18.3-20.3
2016	3,188	22.5	21.7-23.3	1,637	25.5	24.2-26.8	1,551	20.0	18.9-21.0
2017	3,166	22.4	21.6-23.2	1,667	25.8	24.5-27.1	1,499	19.5	18.5-20.6
2018	3,176	22.1	21.3-22.9	1,615	24.7	23.4-25.9	1,561	19.8	18.8-20.9
2019	3,151	21.8	21.0-22.6	1,675	25.3	24.1-26.6	1,476	18.8	17.8-19.8
2020	2,988	20.6	19.9-21.4	1,599	23.9	22.7-25.2	1,389	17.7	16.8-18.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State Excl New York City, Colorectal, White Individuals**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	3,121	35.1	33.8-36.3	1,549	41.8	39.6-44.0	1,572	30.6	29.0-32.1
1977	3,263	36.1	34.8-37.4	1,623	43.5	41.3-45.8	1,640	31.4	29.8-32.9
1978	3,245	35.6	34.4-36.9	1,606	42.6	40.4-44.9	1,639	31.0	29.5-32.5
1979	3,457	37.6	36.3-38.9	1,646	43.1	40.9-45.3	1,811	33.8	32.2-35.4
1980	3,502	37.4	36.2-38.7	1,714	44.3	42.1-46.5	1,788	32.8	31.2-34.3
1981	3,722	39.3	38.0-40.6	1,885	48.9	46.6-51.2	1,837	33.2	31.7-34.8
1982	3,770	39.4	38.2-40.7	1,870	47.3	45.1-49.7	1,900	33.9	32.4-35.5
1983	3,756	38.9	37.7-40.2	1,895	47.7	45.5-50.0	1,861	33.0	31.5-34.5
1984	3,773	39.0	37.8-40.3	1,850	46.5	44.3-48.8	1,923	33.8	32.3-35.3
1985	3,799	38.9	37.7-40.2	1,889	47.2	45.0-49.5	1,910	33.4	31.9-35.0
1986	3,596	36.7	35.5-37.9	1,721	42.7	40.6-44.8	1,875	32.7	31.2-34.2
1987	3,618	36.4	35.3-37.7	1,793	43.9	41.8-46.1	1,825	31.4	30.0-32.9
1988	3,444	34.3	33.2-35.5	1,687	40.9	38.9-43.0	1,757	29.9	28.5-31.3
1989	3,286	32.7	31.5-33.8	1,652	40.1	38.1-42.1	1,634	27.6	26.2-29.0
1990	3,233	31.6	30.6-32.8	1,632	38.8	36.8-40.8	1,601	26.8	25.5-28.1
1991	3,258	31.6	30.5-32.7	1,631	38.2	36.3-40.1	1,627	27.0	25.6-28.3
1992	3,064	29.4	28.4-30.5	1,563	36.3	34.4-38.2	1,501	24.7	23.4-26.0
1993	3,109	29.6	28.6-30.7	1,494	34.2	32.5-36.0	1,615	26.3	25.0-27.6
1994	3,244	30.8	29.7-31.9	1,599	36.9	35.0-38.8	1,645	26.8	25.5-28.1
1995	3,142	29.6	28.6-30.6	1,577	35.8	34.0-37.7	1,565	25.2	24.0-26.5
1996	3,175	29.7	28.7-30.8	1,572	35.6	33.8-37.4	1,603	25.7	24.4-27.0
1997	3,364	31.3	30.2-32.3	1,684	37.2	35.4-39.0	1,680	26.7	25.4-28.1
1998	3,403	31.4	30.3-32.4	1,693	36.9	35.1-38.7	1,710	27.2	25.9-28.5
1999	3,434	31.3	30.2-32.4	1,661	36.0	34.3-37.8	1,773	27.8	26.5-29.1
2000	3,451	31.2	30.2-32.3	1,695	36.2	34.5-38.0	1,756	27.4	26.1-28.7
2001	3,262	29.3	28.3-30.3	1,592	33.8	32.1-35.5	1,670	25.9	24.7-27.2
2002	3,329	29.5	28.5-30.6	1,621	33.8	32.2-35.5	1,708	26.2	25.0-27.5
2003	3,295	29.0	28.0-30.0	1,613	33.2	31.6-34.8	1,682	25.7	24.5-27.0
2004	3,100	27.1	26.1-28.0	1,458	29.5	27.9-31.0	1,642	25.1	23.9-26.3
2005	2,907	25.1	24.2-26.0	1,426	28.7	27.2-30.3	1,481	22.4	21.3-23.6
2006	2,815	24.2	23.3-25.1	1,382	27.6	26.2-29.1	1,433	21.5	20.4-22.7
2007	2,907	24.8	23.9-25.7	1,428	28.0	26.6-29.5	1,479	22.3	21.2-23.5
2008	2,839	23.9	23.0-24.8	1,392	27.0	25.6-28.5	1,447	21.5	20.4-22.7
2009	2,696	22.5	21.6-23.3	1,341	25.5	24.2-27.0	1,355	19.9	18.8-21.0
2010	2,827	23.4	22.5-24.3	1,369	25.7	24.3-27.1	1,458	21.5	20.4-22.7
2011	2,788	22.9	22.0-23.8	1,390	25.7	24.4-27.2	1,398	20.6	19.5-21.8
2012	2,705	22.1	21.2-23.0	1,362	24.8	23.5-26.2	1,343	19.8	18.7-20.9
2013	2,707	21.9	21.0-22.7	1,387	24.8	23.5-26.2	1,320	19.4	18.3-20.5
2014	2,716	21.7	20.9-22.6	1,381	24.5	23.2-25.9	1,335	19.3	18.2-20.4
2015	2,675	21.5	20.6-22.3	1,349	23.8	22.5-25.2	1,326	19.3	18.3-20.5
2016	2,868	22.7	21.8-23.6	1,463	25.5	24.1-26.9	1,405	20.2	19.1-21.4
2017	2,807	22.3	21.5-23.2	1,467	25.4	24.1-26.8	1,340	19.7	18.6-20.8
2018	2,805	22.1	21.2-22.9	1,434	24.7	23.4-26.0	1,371	19.7	18.6-20.9
2019	2,775	21.8	20.9-22.6	1,483	25.4	24.1-26.8	1,292	18.6	17.5-19.7
2020	2,608	20.5	19.7-21.4	1,392	23.7	22.4-25.0	1,216	17.8	16.7-18.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.



**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State Excl New York City, Colorectal, Black Individuals**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	78	25.6	19.6-32.6	27	16.6	10.4-25.1	51	31.2	22.6-41.6
1977	101	31.2	25.0-38.5	37	28.2	19.0-39.9	64	33.9	25.7-43.7
1978	116	34.7	28.2-42.1	51	34.4	24.9-46.1	65	34.4	26.1-44.3
1979	127	37.9	31.1-45.6	64	42.9	31.9-56.2	63	33.8	25.5-43.7
1980	129	38.0	31.2-45.8	59	45.5	33.1-60.6	70	34.2	26.3-43.5
1981	134	40.0	33.1-47.8	67	50.2	37.8-65.0	67	33.5	25.7-42.9
1982	126	34.7	28.5-41.8	55	37.6	27.1-50.4	71	33.5	25.9-42.5
1983	120	32.7	26.8-39.4	49	31.5	22.3-42.7	71	33.4	25.9-42.3
1984	125	32.4	26.6-38.9	56	36.1	26.1-48.2	69	30.7	23.7-38.9
1985	123	32.3	26.5-38.9	52	35.9	25.7-48.3	71	30.7	23.8-39.0
1986	130	33.3	27.5-39.9	61	37.7	27.9-49.6	69	30.1	23.1-38.3
1987	126	29.7	24.5-35.6	61	35.2	25.9-46.5	65	26.3	20.2-33.7
1988	148	33.6	28.2-39.8	66	35.8	26.9-46.5	82	32.2	25.5-40.1
1989	156	34.6	29.1-40.8	78	44.4	33.9-56.8	78	29.4	23.1-36.8
1990	127	27.6	22.8-33.0	60	32.1	23.7-42.3	67	24.7	19.0-31.5
1991	131	27.4	22.7-32.7	58	27.9	20.6-36.8	73	26.5	20.7-33.4
1992	147	29.1	24.4-34.4	71	33.9	25.7-43.8	76	26.2	20.5-32.8
1993	140	26.7	22.3-31.7	65	27.4	20.6-35.6	75	25.4	19.9-31.9
1994	165	32.2	27.3-37.8	85	40.2	31.1-50.8	80	27.0	21.3-33.7
1995	163	30.2	25.5-35.4	75	35.3	26.9-45.3	88	27.6	22.1-34.1
1996	186	33.7	28.8-39.0	94	40.7	32.2-50.7	92	28.8	23.1-35.4
1997	176	29.5	25.1-34.3	93	37.5	29.6-46.7	83	24.2	19.2-30.0
1998	194	32.2	27.7-37.3	88	35.0	27.3-44.1	106	30.7	25.1-37.2
1999	211	34.3	29.7-39.5	98	36.5	28.9-45.3	113	32.6	26.8-39.2
2000	195	31.0	26.7-35.8	93	35.4	27.9-44.1	102	28.2	22.9-34.3
2001	203	31.0	26.7-35.7	107	42.3	34.0-51.8	96	24.6	19.8-30.1
2002	220	33.4	29.0-38.2	89	33.9	26.6-42.4	131	33.6	28.1-40.0
2003	214	30.5	26.4-35.0	110	38.6	31.0-47.3	104	26.1	21.2-31.6
2004	231	32.3	28.1-36.9	130	42.9	35.1-51.7	101	25.1	20.3-30.5
2005	227	30.7	26.7-35.1	117	37.1	30.1-45.1	110	26.2	21.5-31.7
2006	220	28.5	24.8-32.7	112	31.8	25.7-38.8	108	25.3	20.7-30.5
2007	211	26.5	22.9-30.4	104	29.6	23.8-36.3	107	24.2	19.8-29.3
2008	229	28.5	24.8-32.6	114	34.3	27.8-41.8	115	24.8	20.4-29.9
2009	215	25.2	21.8-28.9	109	29.2	23.6-35.6	106	22.0	18.0-26.7
2010	211	24.3	21.1-28.0	112	31.4	25.4-38.2	99	19.7	16.0-24.1
2011	221	25.2	21.9-28.9	117	31.0	25.2-37.5	104	21.0	17.1-25.5
2012	207	22.1	19.1-25.4	106	25.3	20.4-31.0	101	19.5	15.8-23.7
2013	203	21.3	18.4-24.5	104	25.0	20.2-30.6	99	18.3	14.8-22.4
2014	245	25.3	22.2-28.8	111	25.9	21.0-31.4	134	24.4	20.4-29.0
2015	237	22.8	19.9-26.0	122	26.7	21.9-32.2	115	19.8	16.3-23.8
2016	243	23.4	20.5-26.6	137	31.3	26.0-37.2	106	17.9	14.6-21.8
2017	261	25.0	22.0-28.3	151	33.5	28.1-39.6	110	19.1	15.7-23.1
2018	264	24.3	21.4-27.5	130	28.2	23.3-33.8	134	21.6	18.0-25.6
2019	273	24.0	21.2-27.1	139	28.4	23.6-33.7	134	21.0	17.5-24.9
2020	257	22.4	19.7-25.4	145	29.0	24.3-34.4	112	17.3	14.2-20.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State, Lung and Bronchus, All Races**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	5,693	32.4	31.6-33.3	4,031	54.0	52.3-55.8	1,662	16.9	16.1-17.7
1977	5,776	32.8	32.0-33.7	4,054	54.7	53.0-56.5	1,722	17.5	16.6-18.3
1978	5,859	33.3	32.5-34.2	4,141	55.9	54.1-57.7	1,718	17.5	16.6-18.3
1979	6,021	34.2	33.3-35.1	4,118	55.5	53.7-57.2	1,903	19.4	18.5-20.3
1980	6,535	37.0	36.1-37.9	4,525	60.7	58.9-62.6	2,010	20.2	19.3-21.1
1981	6,452	36.5	35.6-37.4	4,332	58.4	56.6-60.2	2,120	21.2	20.3-22.2
1982	6,676	37.6	36.7-38.5	4,378	58.6	56.8-60.4	2,298	23.0	22.0-23.9
1983	6,805	38.1	37.2-39.0	4,468	59.6	57.8-61.4	2,337	23.1	22.1-24.0
1984	6,872	38.5	37.6-39.5	4,477	59.5	57.7-61.4	2,395	23.8	22.9-24.8
1985	7,203	40.2	39.2-41.1	4,627	61.3	59.5-63.2	2,576	25.4	24.4-26.5
1986	7,143	39.8	38.9-40.8	4,525	60.1	58.3-61.9	2,618	25.8	24.8-26.8
1987	7,373	40.9	40.0-41.9	4,679	61.8	60.0-63.7	2,694	26.4	25.4-27.4
1988	7,096	39.3	38.4-40.2	4,346	57.4	55.7-59.2	2,750	26.9	25.8-27.9
1989	7,186	39.8	38.9-40.8	4,326	56.8	55.0-58.5	2,860	28.0	27.0-29.1
1990	7,134	39.4	38.5-40.3	4,382	57.9	56.1-59.6	2,752	26.6	25.6-27.7
1991	7,100	39.3	38.4-40.2	4,226	55.3	53.6-57.0	2,874	27.9	26.9-29.0
1992	6,871	37.8	36.9-38.7	4,117	53.7	52.0-55.4	2,754	26.7	25.7-27.7
1993	6,617	36.2	35.3-37.1	3,832	49.7	48.1-51.3	2,785	26.9	25.9-27.9
1994	6,956	37.9	37.0-38.8	3,951	50.9	49.3-52.5	3,005	28.9	27.9-30.0
1995	7,257	39.3	38.4-40.2	4,152	52.9	51.3-54.6	3,105	29.6	28.6-30.7
1996	7,472	40.4	39.5-41.3	4,247	54.2	52.6-55.9	3,225	30.7	29.7-31.8
1997	7,443	39.9	39.0-40.8	4,219	53.4	51.7-55.0	3,224	30.5	29.5-31.6
1998	7,375	39.2	38.3-40.1	4,094	50.8	49.2-52.4	3,281	30.7	29.7-31.8
1999	7,504	39.3	38.4-40.2	4,165	51.2	49.7-52.8	3,339	30.9	29.8-31.9
2000	7,867	40.6	39.7-41.5	4,281	51.9	50.3-53.5	3,586	32.7	31.6-33.7
2001	7,588	38.8	37.9-39.7	4,065	48.9	47.4-50.5	3,523	31.7	30.6-32.8
2002	8,080	40.8	39.9-41.7	4,377	52.1	50.6-53.7	3,703	33.0	32.0-34.1
2003	8,469	42.3	41.4-43.3	4,607	53.9	52.4-55.5	3,862	34.1	33.0-35.2
2004	9,331	46.1	45.2-47.1	4,965	57.5	55.9-59.2	4,366	38.0	36.9-39.1
2005	9,397	46.2	45.2-47.1	5,041	58.0	56.4-59.6	4,356	37.8	36.7-39.0
2006	9,567	46.8	45.8-47.7	4,935	56.2	54.7-57.8	4,632	40.2	39.0-41.3
2007	9,607	46.3	45.4-47.3	5,008	56.3	54.7-57.9	4,599	39.3	38.1-40.4
2008	9,847	46.8	45.9-47.8	5,099	56.4	54.9-58.0	4,748	40.1	38.9-41.2
2009	9,728	45.6	44.7-46.5	5,153	56.1	54.5-57.6	4,575	38.1	37.0-39.3
2010	9,846	45.4	44.5-46.3	5,143	54.7	53.2-56.3	4,703	38.8	37.6-39.9
2011	10,028	45.4	44.5-46.3	5,182	54.2	52.7-55.7	4,846	39.1	38.0-40.2
2012	10,036	44.6	43.7-45.5	5,167	52.8	51.3-54.3	4,869	38.7	37.7-39.9
2013	9,910	43.2	42.4-44.1	5,028	50.2	48.8-51.7	4,882	38.2	37.1-39.3
2014	9,682	41.3	40.5-42.1	5,012	48.9	47.5-50.3	4,670	35.7	34.7-36.7
2015	9,863	41.4	40.6-42.2	5,091	48.9	47.6-50.3	4,772	36.0	34.9-37.0
2016	9,268	38.1	37.3-38.9	4,720	44.4	43.1-45.7	4,548	33.6	32.6-34.6
2017	9,255	37.5	36.7-38.2	4,733	43.4	42.2-44.7	4,522	33.0	32.1-34.0
2018	8,962	35.6	34.9-36.4	4,551	40.9	39.7-42.1	4,411	31.6	30.7-32.6
2019	9,031	35.1	34.4-35.9	4,488	39.4	38.3-40.6	4,543	32.0	31.1-33.0
2020	7,916	30.4	29.7-31.1	3,957	34.3	33.2-35.4	3,959	27.5	26.6-28.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State, Lung and Bronchus, White Individuals**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	5,041	31.8	30.9-32.7	3,547	52.6	50.8-54.4	1,494	16.9	16.0-17.8
1977	5,090	32.1	31.2-33.1	3,541	52.9	51.1-54.7	1,549	17.7	16.8-18.6
1978	5,152	32.6	31.7-33.6	3,628	54.4	52.6-56.2	1,524	17.4	16.5-18.3
1979	5,263	33.4	32.5-34.3	3,577	53.5	51.7-55.4	1,686	19.4	18.4-20.4
1980	5,681	36.0	35.1-37.0	3,906	58.4	56.5-60.3	1,775	20.1	19.2-21.1
1981	5,617	35.8	34.9-36.8	3,727	56.2	54.4-58.1	1,890	21.6	20.6-22.6
1982	5,795	36.8	35.8-37.8	3,763	56.5	54.6-58.3	2,032	23.2	22.1-24.2
1983	5,944	37.8	36.8-38.8	3,850	57.9	56.1-59.9	2,094	23.6	22.6-24.7
1984	5,952	38.1	37.1-39.1	3,848	58.0	56.1-59.9	2,104	24.1	23.1-25.2
1985	6,239	39.7	38.7-40.7	3,955	59.4	57.5-61.4	2,284	26.1	25.0-27.2
1986	6,167	39.5	38.5-40.5	3,872	58.6	56.7-60.5	2,295	26.3	25.2-27.4
1987	6,367	40.8	39.8-41.8	3,992	60.3	58.4-62.3	2,375	27.2	26.1-28.4
1988	6,171	39.6	38.6-40.6	3,730	56.5	54.7-58.4	2,441	27.9	26.8-29.1
1989	6,181	39.8	38.8-40.9	3,660	55.4	53.5-57.2	2,521	29.0	27.8-30.2
1990	6,139	39.5	38.5-40.5	3,743	56.9	55.1-58.8	2,396	27.4	26.2-28.5
1991	6,129	39.7	38.7-40.8	3,590	54.4	52.6-56.3	2,539	29.2	28.0-30.4
1992	5,870	37.9	37.0-38.9	3,459	52.4	50.7-54.3	2,411	27.8	26.7-29.0
1993	5,628	36.4	35.4-37.4	3,206	48.4	46.8-50.2	2,422	28.0	26.9-29.2
1994	5,966	38.5	37.5-39.5	3,358	50.6	48.9-52.4	2,608	30.2	29.0-31.4
1995	6,218	40.1	39.1-41.1	3,517	52.7	51.0-54.5	2,701	31.0	29.8-32.2
1996	6,319	40.8	39.8-41.9	3,571	53.8	52.1-55.7	2,748	31.6	30.4-32.8
1997	6,283	40.4	39.4-41.4	3,531	52.8	51.1-54.6	2,752	31.6	30.4-32.8
1998	6,188	39.5	38.6-40.5	3,400	50.2	48.5-51.9	2,788	31.7	30.6-33.0
1999	6,420	40.7	39.7-41.7	3,524	51.7	50.0-53.5	2,896	32.8	31.6-34.0
2000	6,692	42.0	41.0-43.0	3,584	52.1	50.4-53.9	3,108	34.8	33.6-36.1
2001	6,449	40.2	39.2-41.2	3,399	49.3	47.6-50.9	3,050	33.8	32.6-35.1
2002	6,921	42.8	41.8-43.8	3,719	53.4	51.7-55.2	3,202	35.4	34.2-36.7
2003	7,167	44.1	43.1-45.1	3,882	55.1	53.4-56.9	3,285	36.1	34.8-37.4
2004	7,894	48.1	47.0-49.2	4,145	58.5	56.7-60.3	3,749	40.7	39.4-42.1
2005	7,937	48.3	47.2-49.3	4,211	59.1	57.3-61.0	3,726	40.5	39.2-41.8
2006	8,134	49.5	48.4-50.6	4,138	57.8	56.0-59.6	3,996	43.7	42.4-45.1
2007	8,075	48.6	47.5-49.7	4,169	57.6	55.9-59.4	3,906	42.1	40.8-43.5
2008	8,314	49.5	48.4-50.6	4,259	58.1	56.3-59.9	4,055	43.4	42.1-44.8
2009	8,156	48.0	47.0-49.1	4,270	57.5	55.7-59.2	3,886	41.2	39.9-42.6
2010	8,193	47.8	46.7-48.8	4,236	56.1	54.4-57.8	3,957	41.8	40.5-43.2
2011	8,298	47.6	46.6-48.7	4,235	55.3	53.6-57.0	4,063	42.2	40.9-43.5
2012	8,226	46.5	45.5-47.5	4,190	53.6	52.0-55.3	4,036	41.5	40.2-42.8
2013	8,209	45.9	44.9-46.9	4,114	51.7	50.1-53.4	4,095	41.6	40.3-42.9
2014	7,929	43.4	42.5-44.4	4,044	49.9	48.4-51.5	3,885	38.6	37.4-39.9
2015	8,061	43.7	42.8-44.7	4,078	49.9	48.4-51.5	3,983	39.4	38.1-40.7
2016	7,476	39.8	38.9-40.7	3,737	44.8	43.3-46.3	3,739	36.3	35.1-37.5
2017	7,468	39.4	38.5-40.3	3,742	44.0	42.6-45.5	3,726	35.9	34.7-37.1
2018	7,219	37.5	36.6-38.3	3,583	41.4	40.0-42.8	3,636	34.6	33.4-35.8
2019	7,198	36.6	35.8-37.5	3,510	39.7	38.4-41.1	3,688	34.5	33.3-35.6
2020	6,317	32.0	31.2-32.8	3,081	34.7	33.5-36.0	3,236	30.0	28.9-31.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State, Lung and Bronchus, Black Individuals**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	575	35.4	32.4-38.6	428	64.0	57.7-70.9	147	15.7	13.2-18.6
1977	630	39.1	35.9-42.5	465	72.8	65.6-80.5	165	17.0	14.4-19.9
1978	645	38.6	35.5-41.8	473	69.8	63.1-77.0	172	17.8	15.1-20.7
1979	694	41.4	38.2-44.8	500	74.3	67.3-81.8	194	19.9	17.0-23.0
1980	784	44.4	41.2-47.8	573	81.9	74.7-89.5	211	20.0	17.3-23.0
1981	774	42.2	39.2-45.4	558	77.5	70.6-84.8	216	19.3	16.8-22.1
1982	784	42.5	39.4-45.7	552	75.4	68.8-82.5	232	21.0	18.3-24.0
1983	789	41.4	38.5-44.5	569	76.0	69.4-83.0	220	19.3	16.8-22.1
1984	840	43.5	40.5-46.7	570	74.0	67.7-80.8	270	23.7	20.9-26.8
1985	879	44.2	41.2-47.3	612	77.8	71.3-84.6	267	22.6	19.9-25.6
1986	888	43.8	40.8-46.8	597	76.6	70.1-83.5	291	23.5	20.9-26.4
1987	906	43.5	40.7-46.6	621	76.3	70.0-83.0	285	22.9	20.2-25.7
1988	846	40.3	37.5-43.2	559	69.2	63.2-75.6	287	22.6	20.0-25.4
1989	914	42.3	39.5-45.2	606	72.9	66.8-79.3	308	23.6	21.0-26.4
1990	887	41.0	38.3-43.9	571	69.6	63.5-76.0	316	23.8	21.3-26.7
1991	868	39.8	37.1-42.6	568	68.6	62.6-74.9	300	22.5	20.0-25.3
1992	884	38.7	36.1-41.4	582	66.2	60.5-72.2	302	21.9	19.4-24.5
1993	811	35.8	33.3-38.5	508	59.7	54.2-65.6	303	21.9	19.5-24.5
1994	884	37.8	35.3-40.4	531	58.9	53.7-64.5	353	24.9	22.3-27.6
1995	891	37.8	35.3-40.4	535	58.0	52.9-63.5	356	25.1	22.5-27.9
1996	1,014	42.1	39.5-44.8	588	63.4	58.1-69.1	426	29.3	26.6-32.2
1997	990	40.8	38.3-43.5	586	63.2	57.8-68.8	404	27.4	24.8-30.3
1998	984	39.4	36.9-42.0	557	56.8	51.9-62.0	427	28.5	25.8-31.3
1999	919	36.2	33.8-38.6	539	54.8	50.0-59.9	380	24.8	22.3-27.4
2000	991	37.8	35.4-40.2	581	56.6	51.8-61.6	410	26.0	23.5-28.7
2001	954	36.1	33.8-38.5	540	52.7	48.1-57.6	414	25.9	23.4-28.5
2002	969	35.6	33.3-37.9	544	52.2	47.6-57.0	425	25.7	23.3-28.3
2003	1,048	37.7	35.4-40.1	556	51.4	46.9-56.0	492	29.3	26.8-32.1
2004	1,168	41.1	38.8-43.6	658	57.7	53.2-62.5	510	30.1	27.5-32.9
2005	1,194	41.9	39.5-44.4	653	59.6	54.9-64.6	541	31.5	28.9-34.3
2006	1,157	40.0	37.7-42.5	629	55.5	51.0-60.3	528	30.4	27.8-33.1
2007	1,169	39.4	37.1-41.7	630	53.9	49.5-58.5	539	30.2	27.7-32.9
2008	1,175	39.4	37.1-41.7	636	54.4	50.0-59.1	539	30.0	27.5-32.7
2009	1,216	39.4	37.1-41.7	661	54.2	49.9-58.7	555	30.1	27.6-32.7
2010	1,273	40.3	38.0-42.6	674	54.1	49.9-58.6	599	31.7	29.2-34.4
2011	1,312	40.4	38.2-42.8	693	54.4	50.2-58.8	619	31.8	29.3-34.4
2012	1,337	40.2	38.0-42.5	702	53.4	49.3-57.7	635	32.0	29.5-34.6
2013	1,215	35.4	33.4-37.5	638	46.9	43.2-50.9	577	28.0	25.8-30.5
2014	1,267	36.1	34.1-38.2	671	47.9	44.1-51.8	596	28.4	26.2-30.9
2015	1,258	34.6	32.7-36.6	683	47.3	43.7-51.2	575	26.5	24.3-28.8
2016	1,242	33.7	31.8-35.7	652	43.9	40.4-47.5	590	26.9	24.7-29.2
2017	1,210	31.8	30.0-33.7	654	43.0	39.7-46.6	556	24.8	22.7-26.9
2018	1,190	30.8	29.1-32.7	626	40.5	37.3-44.0	564	24.5	22.5-26.6
2019	1,198	30.4	28.6-32.2	620	38.7	35.6-42.0	578	24.8	22.8-26.9
2020	1,012	24.9	23.3-26.5	544	33.0	30.2-36.0	468	19.6	17.8-21.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York City, Lung and Bronchus, All Races**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	2,475	32.1	30.8-33.4	1,725	53.8	51.2-56.5	750	16.9	15.7-18.2
1977	2,473	32.5	31.2-33.9	1,714	55.2	52.5-57.9	759	17.2	15.9-18.5
1978	2,504	33.5	32.1-34.8	1,750	56.6	53.9-59.4	754	17.7	16.4-19.0
1979	2,505	33.6	32.2-35.0	1,695	55.6	52.9-58.4	810	18.7	17.4-20.1
1980	2,588	35.1	33.8-36.6	1,775	58.5	55.8-61.4	813	19.2	17.8-20.6
1981	2,535	34.4	33.0-35.8	1,674	56.1	53.4-58.9	861	19.7	18.4-21.1
1982	2,698	36.7	35.3-38.2	1,717	57.4	54.7-60.3	981	22.8	21.4-24.4
1983	2,632	35.7	34.3-37.1	1,709	57.4	54.6-60.2	923	21.2	19.8-22.6
1984	2,677	36.5	35.1-37.9	1,751	58.7	56.0-61.6	926	21.7	20.3-23.2
1985	2,753	37.3	35.9-38.8	1,783	59.3	56.5-62.2	970	22.5	21.1-24.0
1986	2,706	36.8	35.4-38.2	1,696	56.6	53.8-59.4	1,010	23.5	22.0-25.0
1987	2,855	38.7	37.3-40.2	1,810	60.6	57.7-63.5	1,045	24.0	22.5-25.5
1988	2,622	35.9	34.5-37.3	1,601	54.3	51.6-57.1	1,021	23.7	22.3-25.3
1989	2,597	36.0	34.6-37.4	1,532	51.9	49.3-54.6	1,065	25.3	23.8-26.9
1990	2,549	35.5	34.1-36.9	1,519	52.4	49.8-55.2	1,030	24.3	22.9-25.9
1991	2,547	35.8	34.4-37.2	1,497	51.4	48.8-54.1	1,050	25.1	23.6-26.7
1992	2,614	36.6	35.2-38.1	1,603	55.3	52.6-58.1	1,011	24.1	22.7-25.7
1993	2,434	34.2	32.8-35.5	1,442	49.8	47.2-52.5	992	23.9	22.4-25.4
1994	2,504	35.1	33.7-36.5	1,417	48.6	46.1-51.2	1,087	26.1	24.6-27.7
1995	2,566	35.8	34.4-37.2	1,469	50.0	47.4-52.6	1,097	25.9	24.4-27.5
1996	2,689	37.5	36.1-38.9	1,507	51.5	48.9-54.2	1,182	28.1	26.5-29.7
1997	2,606	36.2	34.8-37.6	1,460	49.6	47.1-52.3	1,146	27.1	25.6-28.8
1998	2,625	36.0	34.6-37.4	1,445	47.8	45.4-50.4	1,180	27.6	26.0-29.2
1999	2,511	33.8	32.4-35.1	1,394	45.8	43.4-48.3	1,117	25.6	24.2-27.2
2000	2,745	36.3	35.0-37.7	1,504	48.5	46.1-51.1	1,241	28.0	26.4-29.6
2001	2,539	33.2	32.0-34.6	1,388	44.3	42.0-46.7	1,151	25.6	24.1-27.1
2002	2,670	34.5	33.2-35.8	1,447	45.4	43.1-47.9	1,223	27.0	25.5-28.5
2003	2,725	35.0	33.7-36.4	1,517	47.4	45.0-49.9	1,208	26.5	25.0-28.0
2004	3,024	38.5	37.1-39.9	1,657	51.2	48.7-53.7	1,367	29.7	28.2-31.3
2005	3,022	38.2	36.8-39.6	1,651	50.7	48.3-53.3	1,371	29.8	28.2-31.4
2006	3,010	38.1	36.7-39.5	1,625	50.0	47.5-52.5	1,385	29.9	28.4-31.6
2007	3,113	39.0	37.6-40.4	1,663	50.4	47.9-52.9	1,450	31.1	29.5-32.8
2008	3,040	37.7	36.3-39.1	1,599	47.7	45.3-50.1	1,441	30.8	29.2-32.4
2009	3,019	36.9	35.6-38.2	1,607	47.4	45.1-49.8	1,412	29.7	28.2-31.3
2010	3,055	36.7	35.4-38.1	1,644	47.3	45.0-49.7	1,411	29.2	27.7-30.8
2011	3,153	37.1	35.8-38.5	1,682	47.7	45.4-50.1	1,471	29.8	28.3-31.4
2012	3,112	35.8	34.5-37.1	1,610	44.2	42.1-46.5	1,502	30.1	28.6-31.7
2013	3,011	33.9	32.7-35.2	1,579	42.4	40.3-44.6	1,432	28.0	26.6-29.6
2014	2,988	32.7	31.5-33.9	1,603	41.5	39.4-43.6	1,385	26.4	25.0-27.8
2015	2,951	31.8	30.6-33.0	1,619	41.2	39.2-43.3	1,332	25.1	23.8-26.5
2016	2,879	30.3	29.2-31.4	1,489	37.1	35.2-39.1	1,390	25.5	24.1-26.9
2017	2,894	30.1	29.0-31.2	1,555	38.0	36.1-39.9	1,339	24.4	23.1-25.7
2018	2,742	27.8	26.8-28.9	1,440	34.0	32.2-35.8	1,302	23.3	22.0-24.6
2019	2,813	28.1	27.1-29.2	1,432	33.3	31.6-35.1	1,381	24.3	23.0-25.6
2020	2,326	23.0	22.0-24.0	1,251	29.1	27.4-30.8	1,075	18.5	17.4-19.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York City, Lung and Bronchus, White Individuals**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	1,976	30.7	29.3-32.1	1,356	50.7	47.9-53.6	620	16.8	15.5-18.3
1977	1,984	31.5	30.1-33.0	1,355	52.4	49.6-55.4	629	17.4	16.0-18.9
1978	1,956	32.0	30.6-33.5	1,360	53.8	50.9-56.8	596	17.3	15.9-18.8
1979	1,936	31.8	30.4-33.3	1,293	51.8	48.9-54.8	643	18.4	17.0-20.0
1980	1,971	33.2	31.7-34.8	1,326	54.0	51.1-57.1	645	19.0	17.5-20.6
1981	1,928	32.8	31.3-34.4	1,245	52.0	49.1-55.1	683	19.8	18.3-21.4
1982	2,068	35.6	34.0-37.2	1,289	54.1	51.1-57.3	779	23.2	21.5-25.0
1983	1,997	34.7	33.2-36.3	1,259	54.0	50.9-57.1	738	21.9	20.3-23.6
1984	2,003	35.6	34.0-37.2	1,297	56.2	53.1-59.4	706	21.8	20.2-23.6
1985	2,041	36.1	34.5-37.7	1,292	55.5	52.5-58.7	749	22.9	21.3-24.7
1986	1,971	35.4	33.8-37.0	1,218	53.1	50.1-56.2	753	23.3	21.6-25.1
1987	2,067	37.4	35.8-39.1	1,281	56.5	53.4-59.8	786	24.4	22.7-26.3
1988	1,955	36.3	34.6-38.0	1,164	52.7	49.6-55.9	791	25.2	23.4-27.1
1989	1,884	35.8	34.2-37.5	1,066	48.9	45.9-52.0	818	26.9	25.1-28.9
1990	1,820	35.0	33.4-36.7	1,064	49.9	46.9-53.1	756	25.2	23.4-27.1
1991	1,853	36.5	34.8-38.2	1,043	49.5	46.5-52.6	810	27.3	25.4-29.4
1992	1,893	37.5	35.8-39.2	1,131	54.3	51.2-57.6	762	26.1	24.2-28.0
1993	1,725	34.7	33.0-36.4	1,003	48.4	45.4-51.5	722	25.4	23.6-27.4
1994	1,796	36.3	34.7-38.1	985	47.7	44.8-50.8	811	28.7	26.7-30.8
1995	1,817	36.8	35.2-38.6	1,016	49.5	46.5-52.7	801	27.7	25.8-29.8
1996	1,860	38.0	36.3-39.8	1,026	50.6	47.5-53.8	834	29.4	27.4-31.5
1997	1,778	36.4	34.8-38.2	982	48.4	45.4-51.5	796	28.2	26.2-30.2
1998	1,782	36.3	34.6-38.0	959	46.7	43.8-49.8	823	28.6	26.7-30.7
1999	1,746	35.3	33.6-37.0	936	45.4	42.6-48.5	810	28.4	26.4-30.4
2000	1,917	38.5	36.8-40.3	1,028	49.4	46.5-52.6	889	30.9	28.8-33.0
2001	1,749	35.0	33.4-36.7	931	44.6	41.8-47.6	818	28.1	26.2-30.1
2002	1,842	36.7	35.1-38.4	977	46.4	43.6-49.5	865	29.7	27.7-31.8
2003	1,818	36.3	34.6-38.0	1,018	48.6	45.6-51.6	800	27.3	25.4-29.3
2004	2,016	40.1	38.4-41.9	1,093	52.0	49.0-55.2	923	31.9	29.8-34.0
2005	2,005	40.1	38.3-41.9	1,084	51.7	48.6-54.9	921	32.0	30.0-34.2
2006	1,994	40.3	38.5-42.1	1,063	51.1	48.1-54.3	931	32.6	30.5-34.8
2007	2,026	41.0	39.2-42.8	1,077	51.7	48.6-54.9	949	33.2	31.1-35.5
2008	1,975	40.0	38.2-41.8	1,019	48.5	45.5-51.6	956	33.9	31.7-36.1
2009	1,935	38.7	37.0-40.5	1,014	47.9	45.0-51.0	921	32.2	30.1-34.4
2010	1,918	38.3	36.6-40.1	1,015	47.7	44.7-50.7	903	31.5	29.5-33.7
2011	1,920	37.8	36.1-39.6	1,014	47.0	44.2-50.1	906	31.3	29.2-33.4
2012	1,874	36.3	34.7-38.1	964	43.5	40.8-46.4	910	31.3	29.3-33.5
2013	1,858	35.5	33.9-37.2	963	42.8	40.1-45.7	895	30.2	28.2-32.3
2014	1,768	33.1	31.6-34.7	918	39.8	37.3-42.5	850	28.1	26.2-30.1
2015	1,724	32.1	30.6-33.7	935	40.5	37.9-43.2	789	26.1	24.3-28.1
2016	1,671	30.6	29.1-32.1	833	35.5	33.1-38.0	838	27.2	25.3-29.1
2017	1,636	29.9	28.4-31.4	852	35.9	33.5-38.4	784	25.3	23.5-27.2
2018	1,507	27.0	25.6-28.4	737	30.1	27.9-32.4	770	24.8	23.0-26.7
2019	1,529	27.1	25.7-28.5	760	30.7	28.5-33.0	769	24.3	22.6-26.2
2020	1,259	22.4	21.2-23.7	655	27.0	24.9-29.1	604	18.9	17.4-20.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York City, Lung and Bronchus, Black Individuals**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	441	35.0	31.6-38.6	324	63.8	56.4-71.8	117	16.0	13.1-19.3
1977	448	35.5	32.1-39.2	325	66.7	58.8-75.3	123	15.8	13.1-19.0
1978	497	38.0	34.5-41.6	356	68.1	60.6-76.2	141	18.5	15.5-22.0
1979	516	39.8	36.2-43.6	370	71.3	63.6-79.7	146	19.5	16.3-23.2
1980	564	41.3	37.8-45.0	414	78.1	70.1-86.8	150	18.2	15.3-21.5
1981	562	39.4	36.1-43.0	396	72.3	64.7-80.5	166	18.8	16.0-22.0
1982	564	39.4	36.1-43.0	382	68.5	61.3-76.2	182	21.1	18.1-24.5
1983	575	39.0	35.8-42.5	408	71.6	64.3-79.5	167	18.8	16.0-22.0
1984	610	40.6	37.3-44.0	405	69.6	62.5-77.3	205	22.6	19.5-26.0
1985	649	42.4	39.1-45.9	446	75.5	68.1-83.3	203	22.1	19.1-25.4
1986	661	41.9	38.6-45.3	430	72.3	65.1-80.1	231	23.7	20.7-27.1
1987	702	43.9	40.6-47.4	471	77.8	70.4-85.7	231	23.6	20.6-26.9
1988	616	38.0	35.0-41.2	399	65.8	59.0-73.1	217	21.7	18.9-24.9
1989	643	38.8	35.8-42.0	417	67.3	60.6-74.5	226	22.1	19.3-25.3
1990	649	38.7	35.7-41.8	402	64.3	57.7-71.4	247	23.8	20.9-27.0
1991	613	36.6	33.6-39.7	401	64.6	58.0-71.7	212	20.4	17.7-23.4
1992	634	36.3	33.5-39.3	412	62.3	56.0-69.0	222	20.8	18.1-23.8
1993	594	34.7	31.8-37.6	363	58.1	51.8-65.0	231	21.7	18.9-24.7
1994	622	35.3	32.5-38.3	379	57.3	51.3-63.7	243	22.3	19.6-25.4
1995	635	35.6	32.9-38.6	377	55.8	50.0-62.0	258	23.6	20.8-26.6
1996	720	39.8	36.9-42.8	410	60.2	54.2-66.7	310	27.9	24.9-31.2
1997	691	37.9	35.1-40.9	393	57.8	51.9-64.2	298	26.5	23.6-29.7
1998	679	36.2	33.5-39.1	375	52.1	46.7-57.9	304	26.6	23.7-29.8
1999	638	33.3	30.7-36.0	374	51.9	46.5-57.8	264	22.4	19.8-25.3
2000	681	34.8	32.2-37.5	381	51.4	46.1-57.1	300	24.9	22.2-27.9
2001	635	32.3	29.8-34.9	351	47.2	42.2-52.6	284	23.4	20.8-26.3
2002	671	32.9	30.4-35.6	372	49.1	44.0-54.6	299	23.8	21.2-26.7
2003	704	34.3	31.8-36.9	360	46.4	41.5-51.7	344	27.2	24.4-30.2
2004	790	37.9	35.3-40.7	428	52.4	47.4-57.9	362	28.5	25.6-31.6
2005	804	38.2	35.5-40.9	423	53.4	48.2-59.0	381	29.5	26.6-32.7
2006	798	37.3	34.7-40.0	420	51.2	46.2-56.5	378	28.7	25.9-31.8
2007	791	36.3	33.7-38.9	411	48.8	44.0-54.0	380	28.6	25.8-31.7
2008	790	36.3	33.8-39.0	416	50.0	45.0-55.3	374	28.2	25.4-31.2
2009	801	35.7	33.3-38.3	415	48.0	43.3-53.1	386	28.3	25.5-31.3
2010	833	36.3	33.9-38.9	433	49.2	44.5-54.4	400	28.7	25.9-31.7
2011	898	38.3	35.8-40.9	457	51.3	46.4-56.4	441	30.6	27.8-33.7
2012	872	36.3	33.9-38.9	428	46.4	41.9-51.2	444	30.4	27.6-33.4
2013	778	31.6	29.4-33.9	400	41.8	37.6-46.4	378	25.3	22.8-28.0
2014	834	33.1	30.8-35.5	433	44.3	40.0-48.8	401	26.2	23.6-28.9
2015	782	29.7	27.6-31.9	409	39.2	35.4-43.4	373	23.6	21.2-26.2
2016	788	29.6	27.6-31.8	395	37.4	33.7-41.4	393	24.5	22.1-27.1
2017	799	29.3	27.3-31.4	426	39.5	35.7-43.6	373	22.8	20.6-25.3
2018	783	28.4	26.4-30.5	414	38.0	34.3-42.0	369	22.1	19.9-24.5
2019	779	27.9	25.9-29.9	385	34.6	31.2-38.4	394	23.4	21.1-25.9
2020	632	22.0	20.3-23.8	336	29.9	26.7-33.4	296	17.0	15.1-19.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State Excl New York City, Lung and Bronchus, All Races**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	3,218	32.8	31.6-34.0	2,306	54.4	52.1-56.8	912	16.9	15.8-18.0
1977	3,303	33.1	32.0-34.3	2,340	54.5	52.3-56.9	963	17.7	16.5-18.8
1978	3,355	33.3	32.2-34.5	2,391	55.4	53.1-57.8	964	17.4	16.3-18.5
1979	3,516	34.6	33.5-35.8	2,423	55.4	53.1-57.7	1,093	19.8	18.6-21.0
1980	3,946	38.4	37.2-39.7	2,750	62.4	60.0-64.9	1,196	21.0	19.8-22.3
1981	3,916	38.1	36.9-39.3	2,657	60.1	57.8-62.6	1,259	22.3	21.1-23.6
1982	3,978	38.2	37.0-39.4	2,661	59.3	57.0-61.7	1,317	23.0	21.7-24.3
1983	4,173	39.8	38.6-41.1	2,759	61.0	58.6-63.4	1,414	24.4	23.1-25.8
1984	4,194	40.0	38.7-41.2	2,726	60.0	57.7-62.4	1,468	25.4	24.1-26.8
1985	4,448	42.2	40.9-43.5	2,842	62.7	60.3-65.1	1,606	27.6	26.3-29.0
1986	4,436	42.0	40.8-43.3	2,828	62.5	60.1-64.9	1,608	27.5	26.2-29.0
1987	4,518	42.4	41.2-43.7	2,869	62.7	60.3-65.1	1,649	28.1	26.7-29.5
1988	4,474	41.6	40.4-42.9	2,745	59.4	57.2-61.8	1,729	29.1	27.7-30.5
1989	4,589	42.4	41.2-43.7	2,794	59.9	57.6-62.2	1,795	30.0	28.6-31.4
1990	4,584	42.0	40.7-43.2	2,862	61.2	58.9-63.6	1,722	28.2	26.9-29.6
1991	4,552	41.5	40.3-42.7	2,728	57.6	55.4-59.9	1,824	29.9	28.5-31.3
1992	4,257	38.5	37.4-39.7	2,514	52.6	50.5-54.7	1,743	28.5	27.1-29.8
1993	4,183	37.5	36.4-38.7	2,390	49.5	47.5-51.6	1,793	29.0	27.6-30.4
1994	4,451	39.6	38.5-40.8	2,534	52.2	50.2-54.3	1,917	30.9	29.5-32.3
1995	4,691	41.5	40.3-42.7	2,683	54.6	52.5-56.7	2,008	32.1	30.7-33.6
1996	4,782	42.3	41.1-43.5	2,739	55.8	53.7-57.9	2,043	32.5	31.1-34.0
1997	4,837	42.3	41.2-43.6	2,759	55.6	53.5-57.7	2,078	32.8	31.4-34.3
1998	4,750	41.2	40.0-42.4	2,649	52.4	50.4-54.4	2,101	33.0	31.6-34.4
1999	4,993	42.8	41.7-44.0	2,771	54.5	52.4-56.5	2,222	34.4	33.0-35.9
2000	5,121	43.4	42.2-44.6	2,776	53.9	51.9-55.9	2,345	35.9	34.4-37.4
2001	5,048	42.4	41.2-43.6	2,676	51.7	49.7-53.7	2,372	35.9	34.5-37.4
2002	5,410	44.9	43.7-46.1	2,930	56.1	54.1-58.2	2,480	37.2	35.8-38.7
2003	5,744	47.1	45.9-48.3	3,090	57.9	55.8-60.0	2,654	39.4	37.9-40.9
2004	6,303	51.0	49.7-52.3	3,305	61.2	59.1-63.4	2,998	43.6	42.1-45.2
2005	6,375	51.2	49.9-52.5	3,390	62.2	60.1-64.4	2,985	43.2	41.7-44.8
2006	6,557	52.3	51.0-53.6	3,310	60.0	57.9-62.1	3,247	47.0	45.4-48.7
2007	6,494	50.9	49.7-52.2	3,345	59.7	57.6-61.7	3,149	44.7	43.1-46.3
2008	6,807	52.5	51.3-53.8	3,500	61.4	59.4-63.5	3,307	46.2	44.6-47.9
2009	6,709	51.1	49.8-52.3	3,546	61.0	59.0-63.1	3,163	43.7	42.2-45.3
2010	6,791	50.8	49.6-52.1	3,499	58.9	57.0-61.0	3,292	45.1	43.6-46.7
2011	6,875	50.5	49.3-51.8	3,500	57.9	56.0-59.9	3,375	45.3	43.7-46.9
2012	6,924	50.1	48.9-51.3	3,557	57.9	55.9-59.8	3,367	44.5	43.0-46.0
2013	6,899	49.2	48.0-50.4	3,449	54.9	53.0-56.8	3,450	44.9	43.4-46.5
2014	6,694	46.7	45.6-47.9	3,409	53.3	51.5-55.1	3,285	41.9	40.4-43.4
2015	6,912	47.6	46.4-48.7	3,472	53.6	51.8-55.4	3,440	43.3	41.8-44.8
2016	6,389	43.1	42.0-44.2	3,231	48.7	47.0-50.5	3,158	39.1	37.7-40.6
2017	6,361	42.2	41.2-43.3	3,178	46.7	45.0-48.4	3,183	38.9	37.5-40.3
2018	6,219	40.6	39.6-41.6	3,110	45.1	43.5-46.7	3,109	37.3	35.9-38.6
2019	6,217	39.7	38.7-40.7	3,055	43.1	41.6-44.7	3,162	37.2	35.9-38.6
2020	5,587	35.2	34.2-36.1	2,706	37.4	36.0-38.9	2,881	33.6	32.3-34.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.



**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State Excl New York City, Lung and Bronchus, White Individuals**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	3,065	32.5	31.3-33.7	2,191	53.8	51.5-56.2	874	16.8	15.7-18.0
1977	3,106	32.5	31.3-33.7	2,186	53.1	50.8-55.5	920	17.7	16.5-18.9
1978	3,196	33.1	31.9-34.3	2,268	54.7	52.4-57.1	928	17.4	16.3-18.6
1979	3,327	34.2	33.0-35.4	2,284	54.4	52.1-56.8	1,043	19.7	18.5-21.0
1980	3,709	37.7	36.5-39.0	2,580	61.1	58.7-63.6	1,129	20.8	19.6-22.1
1981	3,688	37.5	36.3-38.8	2,481	58.8	56.4-61.2	1,207	22.5	21.2-23.8
1982	3,727	37.4	36.2-38.7	2,474	57.7	55.4-60.1	1,253	23.0	21.7-24.3
1983	3,947	39.5	38.2-40.8	2,591	60.0	57.6-62.5	1,356	24.6	23.2-25.9
1984	3,948	39.5	38.2-40.8	2,551	58.9	56.5-61.3	1,397	25.5	24.1-26.9
1985	4,197	41.8	40.5-43.1	2,662	61.6	59.2-64.0	1,535	27.9	26.4-29.3
1986	4,195	41.8	40.5-43.1	2,653	61.6	59.2-64.1	1,542	27.9	26.5-29.4
1987	4,300	42.6	41.3-43.9	2,711	62.4	60.0-64.9	1,589	28.7	27.2-30.1
1988	4,216	41.3	40.1-42.6	2,566	58.5	56.2-60.9	1,650	29.3	27.9-30.8
1989	4,297	41.9	40.7-43.2	2,594	58.7	56.4-61.1	1,703	30.1	28.6-31.6
1990	4,318	41.7	40.5-43.0	2,678	60.3	58.0-62.7	1,640	28.5	27.1-29.9
1991	4,275	41.3	40.1-42.6	2,546	56.9	54.6-59.2	1,729	30.1	28.7-31.6
1992	3,977	38.1	37.0-39.4	2,328	51.5	49.4-53.7	1,649	28.6	27.3-30.1
1993	3,903	37.2	36.1-38.4	2,203	48.4	46.4-50.5	1,700	29.3	27.9-30.7
1994	4,170	39.5	38.3-40.8	2,373	52.0	49.9-54.2	1,797	30.9	29.5-32.4
1995	4,401	41.5	40.3-42.8	2,501	54.1	52.0-56.3	1,900	32.5	31.0-34.0
1996	4,458	42.1	40.9-43.4	2,544	55.3	53.1-57.5	1,914	32.7	31.2-34.2
1997	4,505	42.2	41.0-43.5	2,549	54.9	52.7-57.0	1,956	33.2	31.7-34.7
1998	4,406	41.0	39.8-42.2	2,441	51.6	49.6-53.7	1,965	33.2	31.7-34.7
1999	4,674	43.1	41.9-44.3	2,588	54.5	52.4-56.6	2,086	34.9	33.4-36.4
2000	4,774	43.6	42.4-44.8	2,555	53.3	51.2-55.4	2,219	36.7	35.2-38.2
2001	4,699	42.6	41.4-43.8	2,467	51.3	49.3-53.4	2,232	36.5	35.0-38.1
2002	5,079	45.5	44.3-46.8	2,742	56.5	54.4-58.7	2,337	38.1	36.5-39.7
2003	5,349	47.5	46.2-48.8	2,864	58.0	55.8-60.1	2,485	40.1	38.5-41.7
2004	5,876	51.6	50.3-52.9	3,051	61.2	59.0-63.4	2,825	44.7	43.1-46.4
2005	5,932	51.8	50.5-53.2	3,127	62.3	60.1-64.5	2,805	44.3	42.6-46.0
2006	6,140	53.4	52.1-54.8	3,075	60.5	58.4-62.7	3,065	48.6	46.9-50.4
2007	6,049	51.8	50.5-53.1	3,092	60.0	57.9-62.2	2,957	46.0	44.4-47.7
2008	6,339	53.5	52.2-54.8	3,240	61.9	59.8-64.1	3,099	47.6	45.9-49.3
2009	6,221	51.9	50.6-53.2	3,256	61.3	59.1-63.4	2,965	45.1	43.4-46.7
2010	6,275	51.7	50.4-53.0	3,221	59.4	57.3-61.5	3,054	46.2	44.6-48.0
2011	6,378	51.7	50.4-53.0	3,221	58.5	56.5-60.6	3,157	46.9	45.2-48.6
2012	6,352	50.7	49.4-52.0	3,226	57.7	55.7-59.8	3,126	45.8	44.1-47.4
2013	6,351	50.1	48.8-51.4	3,151	55.3	53.3-57.3	3,200	46.3	44.7-48.0
2014	6,161	47.6	46.4-48.9	3,126	53.9	52.0-55.9	3,035	43.0	41.5-44.6
2015	6,337	48.5	47.3-49.8	3,143	53.6	51.7-55.6	3,194	45.0	43.4-46.6
2016	5,805	43.5	42.4-44.7	2,904	48.4	46.7-50.3	2,901	40.2	38.7-41.7
2017	5,832	43.2	42.1-44.4	2,890	47.2	45.4-49.0	2,942	40.3	38.8-41.9
2018	5,711	41.7	40.6-42.8	2,845	45.9	44.2-47.7	2,866	38.6	37.2-40.1
2019	5,668	40.6	39.5-41.7	2,749	43.3	41.7-45.0	2,919	38.7	37.3-40.2
2020	5,055	35.8	34.8-36.8	2,426	37.7	36.2-39.3	2,629	34.5	33.1-35.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State Excl New York City, Lung and Bronchus, Black Individuals**

Year of diagnosis	Males and Females			Males			Females		
	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI
1976	134	37.0	30.7-44.1	104	65.1	52.5-79.7	30	14.7	9.7-21.2
1977	182	51.9	44.2-60.5	140	92.3	76.2-110.7	42	21.3	15.0-29.2
1978	148	40.6	34.0-48.0	117	75.3	60.9-91.8	31	14.9	10.0-21.3
1979	178	47.1	40.0-54.9	130	84.0	68.5-101.7	48	21.1	15.5-28.0
1980	220	55.3	47.9-63.4	159	93.8	78.6-110.9	61	26.9	20.4-34.7
1981	212	52.0	44.9-59.8	162	94.4	79.2-111.5	50	21.1	15.5-27.9
1982	220	53.1	46.0-60.9	170	97.5	82.2-114.7	50	20.9	15.4-27.6
1983	214	49.7	43.0-57.0	161	89.8	75.5-105.8	53	20.8	15.5-27.3
1984	230	53.7	46.7-61.4	165	87.9	74.1-103.4	65	27.9	21.3-35.8
1985	230	50.3	43.8-57.5	166	84.8	71.7-99.6	64	24.7	18.9-31.6
1986	227	50.6	44.0-57.9	167	90.4	76.0-106.5	60	22.9	17.4-29.6
1987	204	42.1	36.4-48.5	150	71.4	59.8-84.4	54	20.0	14.9-26.2
1988	230	47.9	41.7-54.7	160	79.6	66.9-93.9	70	25.5	19.8-32.3
1989	271	54.0	47.5-61.0	189	89.8	76.5-104.7	82	28.7	22.8-35.7
1990	238	48.7	42.5-55.6	169	85.8	72.0-101.1	69	24.1	18.7-30.6
1991	255	50.8	44.5-57.6	167	80.7	67.7-95.1	88	30.3	24.2-37.4
1992	250	46.4	40.7-52.8	170	77.7	65.4-91.6	80	25.4	20.1-31.7
1993	217	39.6	34.3-45.4	145	64.0	53.2-76.2	72	22.8	17.7-28.7
1994	262	45.6	40.1-51.6	152	63.8	53.3-75.5	110	33.2	27.2-40.1
1995	256	44.6	39.1-50.6	158	64.5	54.1-76.1	98	30.2	24.4-36.9
1996	294	49.2	43.6-55.3	178	72.5	61.4-84.9	116	33.8	27.8-40.5
1997	299	49.8	44.1-55.9	193	78.1	66.6-90.8	106	30.5	24.9-37.0
1998	305	49.0	43.5-55.0	182	69.9	59.3-81.7	123	34.6	28.7-41.3
1999	281	45.1	39.9-50.9	165	62.8	52.9-73.9	116	32.8	27.0-39.4
2000	310	46.7	41.5-52.3	200	70.6	60.5-81.9	110	29.6	24.2-35.7
2001	319	47.3	42.1-52.9	189	67.9	57.8-79.0	130	33.5	27.9-39.8
2002	298	43.8	38.8-49.2	172	60.8	51.5-71.3	126	32.2	26.7-38.3
2003	344	47.6	42.6-53.1	196	64.6	55.3-75.0	148	35.9	30.3-42.3
2004	376	50.0	44.9-55.5	228	71.3	61.7-81.8	148	34.6	29.2-40.8
2005	390	52.7	47.4-58.4	230	76.1	65.7-87.6	160	37.7	32.0-44.1
2006	359	48.2	43.2-53.6	209	67.3	57.8-77.7	150	35.5	30.0-41.7
2007	378	48.2	43.3-53.4	219	66.8	57.7-76.9	159	35.1	29.8-41.1
2008	385	47.8	43.0-53.0	220	66.1	57.0-76.0	165	35.5	30.2-41.4
2009	415	49.3	44.5-54.4	246	69.8	60.7-79.7	169	35.3	30.1-41.1
2010	440	50.9	46.1-56.0	241	66.1	57.5-75.6	199	40.4	34.9-46.5
2011	414	46.0	41.5-50.8	236	61.6	53.4-70.5	178	35.1	30.0-40.7
2012	465	50.6	45.9-55.5	274	70.3	61.7-79.7	191	36.5	31.4-42.1
2013	437	45.0	40.7-49.6	238	59.2	51.4-67.6	199	35.3	30.5-40.6
2014	433	44.1	39.9-48.6	238	56.4	49.1-64.6	195	34.9	30.1-40.3
2015	476	47.9	43.6-52.5	274	67.8	59.6-76.8	202	34.4	29.8-39.6
2016	454	44.6	40.5-49.0	257	59.8	52.4-68.0	197	33.7	29.1-38.9
2017	411	38.5	34.8-42.6	228	52.3	45.4-59.9	183	29.7	25.5-34.4
2018	407	37.3	33.7-41.3	212	47.1	40.6-54.2	195	30.9	26.6-35.6
2019	419	36.9	33.4-40.7	235	48.7	42.4-55.7	184	28.6	24.6-33.2
2020	380	32.0	28.8-35.5	208	40.5	34.9-46.7	172	26.1	22.3-30.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State, Female Breast, All Races**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	4,334	46.5	45.1-47.9
1977	4,327	46.7	45.3-48.1
1978	4,554	48.8	47.4-50.3
1979	4,366	46.8	45.3-48.2
1980	4,545	48.4	47.0-49.9
1981	4,630	49.4	48.0-50.9
1982	4,681	49.6	48.2-51.1
1983	4,814	50.7	49.3-52.2
1984	4,769	49.9	48.4-51.3
1985	4,852	50.7	49.2-52.2
1986	4,745	49.5	48.0-50.9
1987	4,692	48.5	47.1-49.9
1988	4,466	46.1	44.7-47.5
1989	4,054	41.9	40.6-43.2
1990	4,164	42.8	41.5-44.2
1991	4,183	42.7	41.4-44.0
1992	4,029	41.2	39.9-42.5
1993	3,875	39.3	38.1-40.6
1994	4,157	41.8	40.5-43.1
1995	4,266	42.5	41.2-43.8
1996	4,432	43.9	42.6-45.2
1997	4,423	43.5	42.2-44.8
1998	4,386	42.7	41.4-44.0
1999	4,369	41.9	40.6-43.1
2000	4,557	43.2	41.9-44.5
2001	4,368	40.9	39.7-42.2
2002	4,544	42.1	40.9-43.3
2003	4,770	43.9	42.6-45.1
2004	5,003	45.7	44.4-47.0
2005	4,969	44.9	43.7-46.2
2006	5,095	46.0	44.7-47.2
2007	5,091	45.5	44.2-46.7
2008	5,090	45.3	44.0-46.6
2009	5,204	45.7	44.5-47.0
2010	4,911	43.0	41.7-44.2
2011	5,168	44.7	43.5-46.0
2012	4,852	41.5	40.3-42.7
2013	5,077	43.2	41.9-44.4
2014	5,285	44.3	43.1-45.6
2015	5,203	43.8	42.5-45.0
2016	5,041	42.1	40.9-43.3
2017	5,022	42.1	40.9-43.3
2018	4,960	41.1	39.9-42.3
2019	5,079	41.7	40.5-42.9
2020	4,630	38.5	37.4-39.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State, Female Breast, White Individuals**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	3,904	47.1	45.6-48.6
1977	3,874	47.3	45.7-48.8
1978	4,081	49.5	47.9-51.1
1979	3,906	47.5	46.0-49.1
1980	4,020	48.9	47.3-50.5
1981	4,104	50.0	48.5-51.7
1982	4,103	50.1	48.6-51.7
1983	4,246	52.0	50.3-53.6
1984	4,176	50.7	49.1-52.3
1985	4,251	51.9	50.3-53.5
1986	4,117	50.4	48.8-52.0
1987	4,096	50.0	48.4-51.6
1988	3,841	47.0	45.5-48.5
1989	3,453	42.5	41.1-44.0
1990	3,544	43.6	42.1-45.1
1991	3,522	43.3	41.8-44.8
1992	3,379	42.0	40.5-43.4
1993	3,231	40.0	38.6-41.4
1994	3,457	42.5	41.0-43.9
1995	3,526	43.2	41.7-44.6
1996	3,641	44.7	43.2-46.2
1997	3,612	44.1	42.6-45.6
1998	3,556	43.2	41.8-44.7
1999	3,573	43.0	41.6-44.5
2000	3,694	44.3	42.8-45.7
2001	3,592	42.9	41.5-44.3
2002	3,631	42.9	41.5-44.3
2003	3,831	45.2	43.8-46.7
2004	3,950	46.3	44.9-47.8
2005	3,894	45.4	44.0-46.9
2006	3,981	46.5	45.1-48.0
2007	3,991	46.6	45.2-48.1
2008	3,921	45.7	44.3-47.2
2009	4,029	46.5	45.1-48.0
2010	3,762	43.6	42.2-45.1
2011	3,868	44.3	42.9-45.8
2012	3,654	41.8	40.4-43.2
2013	3,792	43.3	41.9-44.8
2014	3,922	44.4	42.9-45.8
2015	3,772	42.9	41.5-44.4
2016	3,624	41.2	39.8-42.7
2017	3,604	41.3	39.9-42.8
2018	3,576	40.4	39.0-41.8
2019	3,566	40.1	38.7-41.6
2020	3,190	36.3	35.0-37.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State, Female Breast, Black Individuals**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	382	40.8	36.6-45.4
1977	395	41.4	37.2-45.9
1978	439	44.9	40.6-49.4
1979	421	42.0	37.9-46.4
1980	480	45.7	41.6-50.1
1981	489	46.2	42.1-50.7
1982	523	48.2	44.0-52.6
1983	511	45.4	41.4-49.6
1984	533	45.7	41.8-49.9
1985	528	44.9	41.0-49.0
1986	549	44.4	40.7-48.4
1987	536	42.4	38.8-46.2
1988	570	44.3	40.7-48.2
1989	548	41.8	38.3-45.6
1990	541	40.1	36.7-43.7
1991	575	42.3	38.8-46.0
1992	559	40.1	36.8-43.6
1993	566	39.6	36.4-43.1
1994	591	40.4	37.2-43.9
1995	634	42.6	39.3-46.1
1996	662	44.1	40.7-47.6
1997	689	44.2	41.0-47.7
1998	690	43.7	40.4-47.1
1999	656	40.3	37.3-43.6
2000	695	41.7	38.6-44.9
2001	608	36.1	33.3-39.1
2002	752	43.5	40.4-46.7
2003	758	43.2	40.2-46.4
2004	841	47.6	44.4-50.9
2005	864	48.1	44.9-51.5
2006	883	48.5	45.3-51.8
2007	865	46.9	43.8-50.2
2008	932	50.2	47.0-53.5
2009	909	48.6	45.5-51.9
2010	874	45.7	42.7-48.9
2011	972	49.8	46.7-53.1
2012	927	46.7	43.7-49.9
2013	977	48.6	45.5-51.8
2014	1,028	50.0	46.9-53.2
2015	1,038	50.1	47.1-53.3
2016	1,029	48.9	45.9-52.0
2017	1,024	48.5	45.5-51.7
2018	1,000	47.0	44.1-50.1
2019	1,086	50.7	47.6-53.9
2020	929	43.5	40.6-46.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York City, Female Breast, All Races**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	2,013	48.9	46.7-51.1
1977	1,938	48.0	45.8-50.2
1978	1,945	48.2	46.0-50.5
1979	1,912	47.7	45.5-50.0
1980	1,920	47.9	45.7-50.2
1981	2,021	50.7	48.4-53.0
1982	2,010	50.4	48.2-52.7
1983	1,987	49.2	47.0-51.5
1984	2,058	51.4	49.1-53.7
1985	1,993	49.4	47.2-51.7
1986	1,918	47.6	45.4-49.8
1987	1,899	46.7	44.6-48.9
1988	1,781	44.3	42.2-46.5
1989	1,716	43.1	41.0-45.2
1990	1,702	42.5	40.5-44.6
1991	1,849	46.4	44.3-48.6
1992	1,699	42.9	40.9-45.1
1993	1,594	39.9	38.0-41.9
1994	1,759	43.6	41.6-45.7
1995	1,795	44.2	42.2-46.3
1996	1,863	45.7	43.6-47.9
1997	1,743	42.3	40.3-44.4
1998	1,809	43.3	41.4-45.4
1999	1,776	41.8	39.8-43.8
2000	1,898	44.2	42.2-46.2
2001	1,747	40.0	38.2-41.9
2002	1,810	41.0	39.2-43.0
2003	1,865	42.1	40.2-44.0
2004	1,986	44.7	42.7-46.7
2005	1,919	42.8	40.9-44.8
2006	2,012	45.0	43.0-47.0
2007	2,013	44.6	42.6-46.6
2008	2,005	44.0	42.1-46.0
2009	2,049	44.6	42.7-46.6
2010	1,947	42.2	40.3-44.2
2011	2,129	45.2	43.3-47.2
2012	1,969	41.2	39.4-43.1
2013	2,066	42.9	41.0-44.8
2014	2,172	44.4	42.5-46.4
2015	2,160	43.9	42.0-45.8
2016	2,110	42.4	40.6-44.3
2017	2,110	42.6	40.8-44.5
2018	2,092	41.9	40.0-43.8
2019	2,152	42.5	40.7-44.4
2020	1,923	39.1	37.3-40.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York City, Female Breast, White Individuals**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	1,678	50.9	48.4-53.5
1977	1,590	49.8	47.2-52.5
1978	1,576	49.3	46.7-51.9
1979	1,556	50.0	47.4-52.7
1980	1,521	49.1	46.5-51.8
1981	1,611	52.3	49.6-55.1
1982	1,561	52.0	49.3-54.9
1983	1,554	51.7	49.0-54.5
1984	1,589	53.7	50.9-56.5
1985	1,517	51.2	48.5-54.0
1986	1,437	49.3	46.6-52.0
1987	1,439	49.6	47.0-52.4
1988	1,315	46.6	44.0-49.3
1989	1,253	45.1	42.6-47.8
1990	1,244	45.2	42.6-47.9
1991	1,334	49.5	46.8-52.4
1992	1,204	45.8	43.2-48.6
1993	1,118	42.7	40.1-45.3
1994	1,229	46.2	43.5-48.9
1995	1,229	46.7	44.1-49.5
1996	1,240	47.7	45.0-50.5
1997	1,137	43.5	40.9-46.1
1998	1,171	44.8	42.3-47.5
1999	1,178	44.7	42.1-47.3
2000	1,232	46.9	44.3-49.7
2001	1,160	44.0	41.5-46.7
2002	1,150	43.2	40.7-45.8
2003	1,173	44.6	42.0-47.3
2004	1,238	47.3	44.7-50.1
2005	1,158	44.2	41.6-46.8
2006	1,202	46.5	43.9-49.3
2007	1,218	47.6	44.9-50.4
2008	1,168	45.1	42.5-47.8
2009	1,209	47.0	44.3-49.8
2010	1,131	44.3	41.6-47.0
2011	1,196	45.7	43.1-48.5
2012	1,088	41.9	39.4-44.6
2013	1,140	43.4	40.9-46.1
2014	1,193	45.1	42.5-47.8
2015	1,122	42.3	39.8-44.9
2016	1,108	42.0	39.5-44.6
2017	1,114	42.4	39.9-45.1
2018	1,052	39.4	36.9-41.9
2019	1,062	39.5	37.0-42.0
2020	907	34.8	32.5-37.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York City, Female Breast, Black Individuals**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	307	41.4	36.6-46.5
1977	307	40.4	35.8-45.4
1978	346	44.9	40.1-50.1
1979	336	42.4	37.8-47.4
1980	370	44.9	40.3-49.9
1981	383	46.1	41.5-51.2
1982	419	49.5	44.7-54.7
1983	396	45.0	40.6-49.9
1984	425	46.9	42.4-51.7
1985	420	45.5	41.1-50.2
1986	421	43.5	39.4-48.0
1987	421	42.3	38.3-46.6
1988	433	43.3	39.2-47.7
1989	427	41.8	37.9-46.1
1990	404	38.4	34.7-42.4
1991	451	43.1	39.1-47.3
1992	430	40.2	36.4-44.2
1993	416	37.7	34.1-41.5
1994	446	40.0	36.3-43.9
1995	493	43.4	39.7-47.5
1996	525	46.1	42.2-50.2
1997	515	43.5	39.8-47.4
1998	530	44.3	40.6-48.3
1999	484	39.3	35.8-42.9
2000	530	42.1	38.6-45.9
2001	451	35.6	32.4-39.1
2002	540	41.8	38.3-45.5
2003	557	42.5	39.0-46.2
2004	593	44.9	41.4-48.7
2005	606	45.2	41.6-48.9
2006	636	47.1	43.5-50.9
2007	621	45.7	42.1-49.4
2008	668	49.0	45.3-52.9
2009	646	47.1	43.5-51.0
2010	621	44.5	41.1-48.2
2011	698	49.1	45.5-53.0
2012	686	47.2	43.7-50.9
2013	709	48.3	44.8-52.1
2014	729	49.2	45.7-53.0
2015	745	49.9	46.3-53.7
2016	725	48.0	44.5-51.7
2017	705	46.7	43.2-50.4
2018	747	49.5	45.9-53.3
2019	787	51.6	48.0-55.5
2020	647	43.4	40.0-47.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.



**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State Excl New York City, Female Breast, All Races**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	2,319	44.6	42.8-46.5
1977	2,388	45.8	44.0-47.7
1978	2,609	49.4	47.5-51.4
1979	2,454	46.1	44.3-48.0
1980	2,625	48.7	46.9-50.7
1981	2,608	48.5	46.6-50.5
1982	2,670	49.1	47.2-51.0
1983	2,826	51.8	49.9-53.8
1984	2,711	48.8	47.0-50.8
1985	2,859	51.7	49.7-53.6
1986	2,826	50.9	49.0-52.8
1987	2,793	49.8	47.9-51.7
1988	2,685	47.3	45.5-49.2
1989	2,337	41.0	39.3-42.7
1990	2,459	42.9	41.2-44.7
1991	2,334	40.1	38.5-41.8
1992	2,330	40.0	38.4-41.7
1993	2,280	38.9	37.3-40.5
1994	2,398	40.6	39.0-42.3
1995	2,471	41.3	39.7-43.0
1996	2,569	42.8	41.1-44.5
1997	2,680	44.3	42.6-46.0
1998	2,576	42.1	40.5-43.8
1999	2,592	41.9	40.3-43.6
2000	2,658	42.5	40.9-44.2
2001	2,620	41.5	39.9-43.1
2002	2,734	42.9	41.3-44.5
2003	2,904	45.1	43.5-46.8
2004	3,016	46.3	44.6-48.0
2005	3,050	46.3	44.7-48.0
2006	3,083	46.6	44.9-48.3
2007	3,078	46.0	44.4-47.7
2008	3,085	46.1	44.5-47.8
2009	3,155	46.4	44.8-48.1
2010	2,964	43.4	41.8-45.0
2011	3,039	44.2	42.6-45.9
2012	2,883	41.6	40.1-43.2
2013	3,011	43.3	41.7-44.9
2014	3,113	44.2	42.6-45.8
2015	3,043	43.6	42.0-45.3
2016	2,931	41.8	40.2-43.5
2017	2,911	41.7	40.1-43.3
2018	2,867	40.4	38.9-42.1
2019	2,927	41.2	39.6-42.8
2020	2,704	38.0	36.5-39.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State Excl New York City, Female Breast, White Individuals**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	2,225	44.7	42.9-46.7
1977	2,283	45.8	43.9-47.8
1978	2,505	49.7	47.7-51.7
1979	2,350	46.2	44.3-48.2
1980	2,499	48.8	46.8-50.8
1981	2,493	48.7	46.8-50.7
1982	2,541	49.2	47.2-51.2
1983	2,691	52.1	50.1-54.2
1984	2,587	49.3	47.4-51.3
1985	2,734	52.3	50.3-54.3
1986	2,680	51.1	49.1-53.1
1987	2,657	50.3	48.3-52.3
1988	2,526	47.3	45.4-49.2
1989	2,199	41.1	39.4-42.9
1990	2,297	42.7	41.0-44.6
1991	2,188	40.3	38.6-42.0
1992	2,175	40.1	38.4-41.9
1993	2,112	38.7	37.1-40.4
1994	2,228	40.7	39.0-42.4
1995	2,297	41.5	39.8-43.3
1996	2,401	43.4	41.6-45.2
1997	2,475	44.4	42.6-46.2
1998	2,384	42.4	40.7-44.2
1999	2,395	42.3	40.6-44.0
2000	2,462	43.1	41.4-44.8
2001	2,431	42.3	40.6-44.0
2002	2,481	42.7	41.0-44.5
2003	2,658	45.5	43.8-47.3
2004	2,711	45.9	44.2-47.7
2005	2,736	46.0	44.2-47.7
2006	2,779	46.6	44.8-48.4
2007	2,773	46.2	44.5-48.0
2008	2,753	45.9	44.2-47.7
2009	2,820	46.3	44.6-48.1
2010	2,631	43.3	41.6-45.0
2011	2,672	43.6	41.9-45.4
2012	2,566	41.6	40.0-43.4
2013	2,652	43.1	41.4-44.9
2014	2,729	44.0	42.3-45.8
2015	2,650	43.3	41.5-45.1
2016	2,516	41.0	39.3-42.7
2017	2,490	40.9	39.2-42.7
2018	2,524	40.9	39.2-42.6
2019	2,504	40.5	38.8-42.3
2020	2,280	36.9	35.3-38.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State Excl New York City, Female Breast, Black Individuals**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	74	38.5	29.8-48.9
1977	88	45.1	35.6-56.2
1978	93	45.1	36.1-55.7
1979	85	40.4	31.9-50.4
1980	110	48.7	39.8-59.1
1981	106	46.3	37.6-56.4
1982	104	43.3	35.1-52.6
1983	115	46.2	37.9-55.7
1984	108	41.8	34.0-50.7
1985	108	42.5	34.6-51.6
1986	127	47.1	39.1-56.2
1987	115	42.6	35.0-51.3
1988	137	48.3	40.4-57.2
1989	121	41.6	34.3-50.0
1990	137	46.1	38.6-54.7
1991	124	39.8	33.0-47.6
1992	129	39.9	33.2-47.4
1993	150	46.2	39.0-54.3
1994	145	41.7	35.1-49.2
1995	141	39.8	33.4-47.0
1996	137	37.4	31.3-44.3
1997	174	46.7	39.9-54.3
1998	160	41.5	35.2-48.5
1999	172	43.8	37.4-50.9
2000	165	40.3	34.3-47.1
2001	157	37.3	31.6-43.7
2002	212	48.2	41.8-55.2
2003	200	45.1	39.0-51.9
2004	248	55.3	48.6-62.8
2005	258	56.7	49.9-64.2
2006	247	52.4	46.0-59.4
2007	244	50.4	44.2-57.2
2008	264	53.5	47.1-60.4
2009	263	52.7	46.5-59.6
2010	253	49.0	43.1-55.6
2011	274	51.6	45.6-58.2
2012	241	45.3	39.6-51.4
2013	268	49.5	43.7-55.9
2014	299	51.8	46.0-58.1
2015	293	51.0	45.2-57.3
2016	304	51.2	45.5-57.4
2017	319	53.7	47.8-60.0
2018	252	40.9	35.9-46.4
2019	299	48.8	43.3-54.7
2020	282	44.0	38.9-49.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State, Cervix, All Races**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	483	5.3	4.8-5.8
1977	461	5.0	4.6-5.5
1978	447	5.0	4.5-5.5
1979	445	4.9	4.4-5.4
1980	471	5.1	4.6-5.6
1981	438	4.7	4.3-5.2
1982	454	5.0	4.5-5.5
1983	424	4.6	4.2-5.1
1984	432	4.8	4.3-5.3
1985	395	4.2	3.8-4.6
1986	406	4.4	3.9-4.8
1987	448	4.7	4.3-5.2
1988	396	4.2	3.8-4.7
1989	406	4.2	3.8-4.6
1990	443	4.7	4.3-5.2
1991	414	4.4	4.0-4.8
1992	412	4.2	3.8-4.6
1993	396	4.1	3.7-4.5
1994	439	4.5	4.0-4.9
1995	429	4.3	3.9-4.8
1996	452	4.5	4.1-5.0
1997	421	4.2	3.8-4.6
1998	379	3.7	3.4-4.1
1999	389	3.8	3.4-4.2
2000	390	3.8	3.4-4.2
2001	376	3.6	3.2-4.0
2002	374	3.5	3.2-3.9
2003	379	3.5	3.2-3.9
2004	395	3.7	3.3-4.1
2005	426	4.0	3.6-4.4
2006	437	4.0	3.6-4.4
2007	442	4.0	3.7-4.4
2008	435	4.0	3.6-4.4
2009	477	4.3	3.9-4.7
2010	437	4.0	3.6-4.4
2011	423	3.7	3.4-4.1
2012	438	3.9	3.5-4.2
2013	450	4.0	3.6-4.3
2014	461	4.0	3.6-4.4
2015	441	3.8	3.4-4.2
2016	400	3.5	3.1-3.8
2017	449	3.9	3.5-4.3
2018	397	3.4	3.0-3.7
2019	408	3.6	3.3-4.0
2020	390	3.3	3.0-3.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State, Cervix, White Individuals**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	353	4.3	3.8-4.8
1977	318	3.9	3.5-4.4
1978	314	4.0	3.5-4.5
1979	297	3.8	3.4-4.3
1980	340	4.2	3.8-4.7
1981	325	4.1	3.6-4.6
1982	313	4.1	3.6-4.6
1983	310	4.0	3.5-4.4
1984	299	3.9	3.5-4.4
1985	285	3.6	3.2-4.0
1986	268	3.4	3.0-3.8
1987	320	4.0	3.6-4.5
1988	288	3.7	3.3-4.2
1989	287	3.6	3.2-4.1
1990	296	3.8	3.4-4.3
1991	290	3.8	3.3-4.2
1992	274	3.4	3.0-3.9
1993	259	3.3	2.9-3.8
1994	303	3.8	3.4-4.3
1995	309	4.0	3.5-4.4
1996	312	3.9	3.5-4.4
1997	292	3.7	3.3-4.2
1998	263	3.3	2.9-3.7
1999	272	3.4	3.0-3.9
2000	265	3.3	2.9-3.7
2001	245	3.0	2.6-3.4
2002	229	2.8	2.5-3.2
2003	256	3.1	2.8-3.5
2004	249	3.0	2.7-3.4
2005	269	3.3	2.9-3.7
2006	283	3.5	3.1-3.9
2007	284	3.5	3.1-3.9
2008	299	3.7	3.3-4.2
2009	322	3.9	3.5-4.4
2010	267	3.3	2.9-3.7
2011	271	3.3	2.9-3.7
2012	284	3.4	3.0-3.9
2013	279	3.4	3.0-3.9
2014	298	3.5	3.1-4.0
2015	279	3.4	3.0-3.8
2016	242	2.9	2.5-3.3
2017	268	3.3	2.9-3.7
2018	254	3.1	2.7-3.5
2019	237	3.0	2.6-3.4
2020	244	2.9	2.6-3.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State, Cervix, Black Individuals**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	120	12.8	10.4-15.4
1977	128	13.6	11.3-16.3
1978	121	12.6	10.4-15.2
1979	138	13.5	11.3-16.1
1980	122	12.0	9.9-14.5
1981	101	8.9	7.2-10.9
1982	127	11.5	9.5-13.8
1983	103	9.1	7.4-11.1
1984	116	10.1	8.3-12.1
1985	99	8.4	6.8-10.3
1986	123	10.4	8.6-12.4
1987	109	8.6	7.0-10.4
1988	96	7.5	6.0-9.2
1989	108	8.1	6.6-9.8
1990	133	9.5	7.9-11.2
1991	112	8.2	6.7-9.9
1992	117	8.1	6.7-9.7
1993	107	7.5	6.2-9.1
1994	107	7.5	6.1-9.0
1995	100	6.7	5.4-8.2
1996	114	7.4	6.1-8.9
1997	103	6.7	5.5-8.2
1998	92	5.7	4.6-7.0
1999	94	5.8	4.7-7.1
2000	106	6.5	5.3-7.8
2001	110	6.5	5.3-7.8
2002	116	6.8	5.6-8.2
2003	105	6.0	4.9-7.3
2004	116	6.6	5.4-7.9
2005	128	7.2	6.0-8.6
2006	117	6.5	5.4-7.8
2007	128	7.0	5.8-8.4
2008	112	6.0	4.9-7.2
2009	124	6.5	5.4-7.7
2010	131	6.9	5.8-8.2
2011	115	5.9	4.8-7.1
2012	117	6.0	5.0-7.2
2013	121	6.0	5.0-7.2
2014	118	5.8	4.8-7.0
2015	121	5.8	4.8-7.0
2016	122	5.8	4.8-7.0
2017	122	5.8	4.8-6.9
2018	104	4.8	3.9-5.9
2019	107	5.1	4.1-6.2
2020	100	4.6	3.7-5.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York City, Cervix, All Races**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	254	6.4	5.6-7.2
1977	257	6.6	5.8-7.5
1978	233	6.3	5.5-7.2
1979	248	6.5	5.7-7.3
1980	227	6.0	5.2-6.9
1981	245	6.4	5.6-7.3
1982	254	6.8	6.0-7.7
1983	211	5.4	4.7-6.2
1984	210	5.6	4.8-6.4
1985	193	4.9	4.2-5.6
1986	209	5.4	4.7-6.2
1987	231	5.9	5.1-6.7
1988	209	5.4	4.7-6.2
1989	200	5.0	4.3-5.7
1990	234	6.0	5.2-6.8
1991	218	5.7	5.0-6.6
1992	225	5.6	4.9-6.4
1993	218	5.6	4.9-6.4
1994	257	6.4	5.6-7.2
1995	223	5.5	4.8-6.3
1996	235	5.8	5.1-6.6
1997	228	5.5	4.8-6.3
1998	204	4.9	4.2-5.6
1999	207	4.9	4.2-5.6
2000	214	5.0	4.4-5.7
2001	201	4.6	4.0-5.3
2002	205	4.7	4.1-5.4
2003	203	4.6	4.0-5.3
2004	212	4.8	4.1-5.4
2005	225	5.1	4.4-5.8
2006	229	5.1	4.5-5.8
2007	222	4.9	4.3-5.7
2008	215	4.7	4.1-5.4
2009	230	5.1	4.4-5.8
2010	215	4.7	4.1-5.4
2011	221	4.8	4.1-5.4
2012	225	4.7	4.1-5.4
2013	214	4.4	3.8-5.0
2014	231	4.8	4.2-5.5
2015	219	4.4	3.9-5.1
2016	217	4.5	3.9-5.1
2017	246	5.1	4.4-5.8
2018	209	4.1	3.6-4.8
2019	217	4.5	3.9-5.1
2020	183	3.6	3.1-4.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York City, Cervix, White Individuals**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	150	4.6	3.9-5.5
1977	138	4.4	3.7-5.3
1978	124	4.5	3.7-5.4
1979	127	4.4	3.7-5.3
1980	125	4.5	3.7-5.5
1981	144	5.2	4.3-6.2
1982	136	5.2	4.3-6.2
1983	122	4.4	3.6-5.3
1984	104	3.9	3.1-4.8
1985	103	3.8	3.0-4.6
1986	91	3.3	2.6-4.1
1987	128	4.7	3.9-5.7
1988	116	4.4	3.6-5.4
1989	111	4.1	3.3-5.0
1990	120	4.7	3.9-5.7
1991	117	4.8	3.9-5.7
1992	112	4.3	3.5-5.2
1993	108	4.3	3.5-5.2
1994	145	5.8	4.9-6.8
1995	128	5.2	4.4-6.3
1996	128	5.1	4.3-6.1
1997	116	4.6	3.8-5.6
1998	107	4.3	3.5-5.2
1999	116	4.6	3.8-5.6
2000	110	4.3	3.6-5.2
2001	91	3.6	2.9-4.4
2002	89	3.6	2.9-4.4
2003	105	4.1	3.4-5.0
2004	94	3.6	2.9-4.4
2005	100	4.0	3.2-4.9
2006	103	4.1	3.3-4.9
2007	96	3.9	3.1-4.7
2008	107	4.2	3.5-5.1
2009	110	4.4	3.6-5.3
2010	84	3.4	2.7-4.3
2011	97	3.9	3.1-4.8
2012	106	4.1	3.4-5.0
2013	86	3.3	2.6-4.1
2014	103	4.0	3.3-4.9
2015	97	3.8	3.0-4.6
2016	92	3.6	2.9-4.5
2017	102	4.1	3.3-5.0
2018	98	3.6	2.9-4.5
2019	83	3.2	2.5-4.0
2020	83	3.1	2.5-3.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.



**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York City, Cervix, Black Individuals**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	97	12.9	10.4-15.9
1977	106	14.4	11.7-17.6
1978	102	13.3	10.7-16.3
1979	114	14.1	11.5-17.1
1980	95	11.8	9.5-14.6
1981	91	10.2	8.1-12.5
1982	105	12.4	10.0-15.1
1983	82	9.3	7.4-11.7
1984	93	10.4	8.3-12.8
1985	81	8.8	6.9-11.0
1986	105	11.4	9.3-13.9
1987	87	8.7	6.9-10.7
1988	83	8.3	6.6-10.3
1989	80	7.6	6.0-9.5
1990	103	9.4	7.7-11.5
1991	91	8.5	6.8-10.5
1992	95	8.7	7.0-10.6
1993	86	8.0	6.4-9.9
1994	91	8.3	6.7-10.3
1995	81	7.1	5.6-8.9
1996	85	7.2	5.7-8.9
1997	93	8.0	6.4-9.8
1998	75	6.1	4.8-7.6
1999	74	6.0	4.7-7.5
2000	89	7.1	5.7-8.8
2001	91	7.1	5.7-8.8
2002	94	7.3	5.9-9.0
2003	82	6.2	4.9-7.7
2004	91	6.9	5.5-8.4
2005	98	7.4	6.0-9.1
2006	95	7.1	5.7-8.7
2007	101	7.5	6.1-9.1
2008	88	6.4	5.2-8.0
2009	97	6.9	5.5-8.4
2010	100	7.3	5.9-8.9
2011	96	6.8	5.5-8.3
2012	90	6.4	5.1-7.9
2013	84	5.8	4.6-7.2
2014	91	6.3	5.1-7.8
2015	93	6.1	4.9-7.6
2016	92	6.2	4.9-7.6
2017	93	6.1	4.9-7.6
2018	79	5.1	4.0-6.4
2019	85	5.8	4.6-7.3
2020	71	4.5	3.5-5.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State Excl New York City, Cervix, All Races**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	229	4.4	3.9-5.1
1977	204	3.9	3.4-4.5
1978	214	4.1	3.5-4.7
1979	197	3.8	3.3-4.4
1980	244	4.4	3.9-5.0
1981	193	3.5	3.0-4.1
1982	200	3.7	3.2-4.3
1983	213	4.0	3.5-4.6
1984	222	4.2	3.6-4.8
1985	202	3.7	3.2-4.2
1986	197	3.6	3.1-4.2
1987	217	3.8	3.3-4.4
1988	187	3.4	2.9-3.9
1989	206	3.7	3.2-4.2
1990	208	3.8	3.3-4.3
1991	196	3.4	3.0-4.0
1992	187	3.2	2.8-3.7
1993	178	3.1	2.7-3.6
1994	182	3.1	2.7-3.6
1995	206	3.5	3.1-4.1
1996	217	3.7	3.2-4.2
1997	193	3.3	2.8-3.8
1998	175	2.9	2.5-3.4
1999	182	3.0	2.6-3.5
2000	176	2.9	2.5-3.4
2001	175	2.8	2.4-3.3
2002	169	2.7	2.3-3.2
2003	176	2.8	2.4-3.3
2004	182	2.9	2.5-3.4
2005	201	3.2	2.8-3.7
2006	208	3.3	2.8-3.7
2007	220	3.4	3.0-3.9
2008	220	3.5	3.0-4.0
2009	247	3.8	3.3-4.3
2010	222	3.5	3.1-4.0
2011	202	3.1	2.7-3.6
2012	213	3.3	2.8-3.8
2013	236	3.7	3.2-4.2
2014	230	3.4	3.0-3.9
2015	222	3.3	2.9-3.8
2016	183	2.8	2.4-3.2
2017	203	3.0	2.6-3.5
2018	188	2.9	2.4-3.3
2019	191	3.0	2.6-3.5
2020	207	3.2	2.8-3.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State Excl New York City, Cervix, White Individuals**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	203	4.1	3.5-4.7
1977	180	3.6	3.1-4.2
1978	190	3.8	3.3-4.4
1979	170	3.4	2.9-4.0
1980	215	4.1	3.6-4.7
1981	181	3.5	3.0-4.1
1982	177	3.5	3.0-4.1
1983	188	3.7	3.2-4.3
1984	195	3.9	3.4-4.5
1985	182	3.5	3.0-4.1
1986	177	3.4	2.9-4.0
1987	192	3.6	3.1-4.2
1988	172	3.3	2.8-3.9
1989	176	3.4	2.9-3.9
1990	176	3.4	2.9-4.0
1991	173	3.3	2.8-3.8
1992	162	3.0	2.6-3.6
1993	151	2.9	2.4-3.4
1994	158	2.9	2.5-3.4
1995	181	3.4	2.9-3.9
1996	184	3.4	2.9-3.9
1997	176	3.3	2.8-3.8
1998	156	2.9	2.4-3.4
1999	156	2.9	2.5-3.4
2000	155	2.8	2.4-3.3
2001	154	2.7	2.3-3.2
2002	140	2.5	2.1-3.0
2003	151	2.7	2.3-3.2
2004	155	2.8	2.4-3.3
2005	169	3.0	2.6-3.5
2006	180	3.2	2.7-3.7
2007	188	3.3	2.8-3.8
2008	192	3.5	3.0-4.0
2009	212	3.7	3.2-4.2
2010	183	3.3	2.8-3.8
2011	174	3.0	2.6-3.6
2012	178	3.1	2.7-3.6
2013	193	3.5	3.0-4.1
2014	195	3.3	2.9-3.9
2015	182	3.2	2.7-3.7
2016	150	2.6	2.2-3.1
2017	166	2.9	2.4-3.4
2018	156	2.8	2.4-3.3
2019	154	2.9	2.5-3.5
2020	161	2.9	2.5-3.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State Excl New York City, Cervix, Black Individuals**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	23	12.1	7.3-18.8
1977	22	11.0	6.6-17.0
1978	19	10.4	6.0-16.4
1979	24	11.3	7.0-17.2
1980	27	13.0	8.4-19.1
1981	10	4.5	2.1-8.4
1982	22	8.6	5.3-13.0
1983	21	8.1	5.0-12.5
1984	23	9.0	5.6-13.5
1985	18	7.0	4.0-11.1
1986	18	6.5	3.8-10.4
1987	22	8.3	5.1-12.6
1988	13	4.7	2.4-8.0
1989	28	9.8	6.4-14.1
1990	29	9.1	6.0-13.2
1991	21	7.2	4.4-11.1
1992	22	6.0	3.7-9.2
1993	21	6.0	3.6-9.2
1994	16	4.5	2.6-7.4
1995	19	5.2	3.1-8.2
1996	29	8.1	5.4-11.6
1997	10	2.7	1.3-5.0
1998	17	4.3	2.5-6.9
1999	20	5.5	3.3-8.4
2000	17	4.3	2.5-6.9
2001	19	4.6	2.7-7.2
2002	22	5.0	3.1-7.6
2003	23	5.6	3.5-8.4
2004	24	5.4	3.5-8.1
2005	30	6.7	4.5-9.6
2006	22	4.8	3.0-7.2
2007	27	5.7	3.8-8.3
2008	24	4.7	3.0-7.0
2009	27	5.5	3.6-8.1
2010	31	5.9	4.0-8.3
2011	19	3.5	2.1-5.5
2012	27	5.0	3.3-7.3
2013	37	6.7	4.7-9.3
2014	27	4.6	3.0-6.7
2015	28	4.9	3.3-7.2
2016	30	5.0	3.3-7.2
2017	29	4.9	3.3-7.1
2018	25	4.1	2.6-6.1
2019	22	3.4	2.1-5.2
2020	29	4.8	3.2-6.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State, Prostate, All Races**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	1,616	26.1	24.8-27.5
1977	1,619	25.5	24.2-26.9
1978	1,635	25.9	24.6-27.2
1979	1,717	27.8	26.4-29.2
1980	1,870	29.3	27.9-30.7
1981	1,929	30.4	29.0-31.8
1982	1,948	30.5	29.1-31.9
1983	1,880	28.6	27.3-30.0
1984	1,951	29.5	28.1-30.9
1985	2,048	30.3	29.0-31.7
1986	2,041	30.4	29.0-31.8
1987	2,077	30.5	29.1-31.9
1988	2,117	31.0	29.6-32.4
1989	2,080	30.1	28.8-31.5
1990	2,329	33.0	31.6-34.4
1991	2,565	35.9	34.5-37.3
1992	2,718	36.5	35.1-37.9
1993	2,471	33.1	31.8-34.5
1994	2,243	29.7	28.5-31.0
1995	2,081	27.1	26.0-28.4
1996	1,906	24.7	23.6-25.9
1997	1,883	24.4	23.3-25.6
1998	1,623	20.8	19.7-21.8
1999	1,605	20.1	19.1-21.1
2000	1,614	19.9	18.9-20.9
2001	1,470	17.7	16.8-18.7
2002	1,476	17.6	16.7-18.5
2003	1,473	17.3	16.4-18.2
2004	1,712	19.3	18.4-20.2
2005	1,773	19.5	18.6-20.4
2006	1,954	21.3	20.4-22.3
2007	2,004	21.4	20.5-22.4
2008	2,096	21.7	20.7-22.6
2009	2,209	22.3	21.3-23.2
2010	2,316	22.9	21.9-23.9
2011	2,311	22.5	21.5-23.4
2012	2,135	20.1	19.3-21.0
2013	2,333	21.7	20.8-22.6
2014	2,409	22.0	21.1-22.9
2015	2,582	23.1	22.2-24.1
2016	2,791	24.7	23.7-25.6
2017	2,956	25.6	24.7-26.5
2018	3,096	26.6	25.6-27.5
2019	3,253	27.3	26.3-28.3
2020	2,946	24.3	23.5-25.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State, Prostate, White Individuals**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	1,386	24.5	23.2-25.9
1977	1,361	23.6	22.3-25.0
1978	1,356	23.6	22.3-24.9
1979	1,435	25.6	24.2-27.0
1980	1,508	26.2	24.8-27.6
1981	1,597	27.9	26.5-29.3
1982	1,625	28.2	26.7-29.6
1983	1,559	26.4	25.1-27.8
1984	1,609	27.2	25.8-28.6
1985	1,654	27.5	26.2-28.9
1986	1,647	27.6	26.2-29.0
1987	1,695	28.1	26.7-29.5
1988	1,723	28.4	27.0-29.8
1989	1,696	27.8	26.5-29.2
1990	1,897	30.6	29.2-32.1
1991	2,101	33.5	32.1-35.0
1992	2,227	34.2	32.8-35.7
1993	2,015	31.1	29.7-32.5
1994	1,720	26.4	25.1-27.7
1995	1,567	23.9	22.7-25.1
1996	1,483	22.6	21.5-23.8
1997	1,399	21.4	20.3-22.6
1998	1,204	18.2	17.2-19.3
1999	1,194	17.8	16.8-18.9
2000	1,198	17.8	16.8-18.8
2001	1,104	16.0	15.1-17.0
2002	1,106	15.9	15.0-16.9
2003	1,133	16.1	15.2-17.1
2004	1,308	18.0	17.1-19.1
2005	1,385	18.7	17.7-19.7
2006	1,505	20.3	19.3-21.3
2007	1,543	20.4	19.4-21.4
2008	1,620	20.9	19.9-22.0
2009	1,680	21.1	20.1-22.2
2010	1,795	22.2	21.2-23.3
2011	1,745	21.3	20.3-22.3
2012	1,613	19.2	18.3-20.2
2013	1,720	20.3	19.3-21.3
2014	1,778	20.6	19.7-21.7
2015	1,905	21.8	20.8-22.8
2016	2,077	23.5	22.4-24.5
2017	2,216	24.7	23.7-25.8
2018	2,313	25.6	24.6-26.7
2019	2,368	25.7	24.7-26.8
2020	2,179	23.4	22.4-24.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State, Prostate, Black Individuals**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	216	47.8	40.7-55.6
1977	246	51.5	44.4-59.2
1978	273	57.3	49.8-65.6
1979	275	59.0	51.4-67.2
1980	352	70.6	62.6-79.4
1981	320	63.3	55.8-71.5
1982	314	61.4	54.1-69.3
1983	307	56.2	49.4-63.6
1984	326	57.6	50.8-64.9
1985	370	61.2	54.5-68.5
1986	374	62.6	55.8-70.0
1987	366	58.1	51.8-65.0
1988	372	61.6	54.9-68.8
1989	375	58.3	52.0-65.1
1990	416	61.5	55.2-68.3
1991	427	61.6	55.3-68.3
1992	443	60.8	54.8-67.3
1993	422	58.0	52.1-64.3
1994	475	62.9	56.9-69.4
1995	463	57.5	52.0-63.4
1996	381	45.4	40.6-50.5
1997	440	52.8	47.5-58.3
1998	376	43.9	39.3-48.9
1999	384	42.6	38.1-47.4
2000	383	39.8	35.6-44.2
2001	321	33.6	29.8-37.7
2002	337	33.9	30.2-38.0
2003	303	29.1	25.7-32.8
2004	367	33.8	30.2-37.7
2005	331	29.7	26.4-33.3
2006	390	33.5	30.1-37.2
2007	406	34.8	31.3-38.5
2008	434	34.0	30.7-37.5
2009	444	34.7	31.4-38.3
2010	458	34.8	31.5-38.3
2011	466	34.2	31.0-37.7
2012	437	30.7	27.8-33.9
2013	515	35.9	32.7-39.3
2014	523	35.1	32.0-38.4
2015	529	34.8	31.7-38.0
2016	571	37.0	33.9-40.4
2017	576	35.3	32.3-38.4
2018	608	37.1	34.1-40.3
2019	678	40.6	37.5-43.9
2020	569	33.1	30.3-36.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York City, Prostate, All Races**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	730	26.4	24.4-28.5
1977	720	25.8	23.8-27.8
1978	753	27.4	25.4-29.5
1979	801	29.8	27.7-32.1
1980	861	31.6	29.4-33.9
1981	839	31.2	29.0-33.5
1982	870	32.2	30.0-34.5
1983	811	29.7	27.6-31.9
1984	867	31.5	29.4-33.7
1985	893	32.1	29.9-34.3
1986	863	31.5	29.3-33.7
1987	852	30.5	28.4-32.7
1988	814	30.3	28.2-32.5
1989	845	31.1	29.0-33.3
1990	921	34.0	31.7-36.3
1991	984	35.8	33.5-38.1
1992	962	34.5	32.3-36.8
1993	888	31.9	29.8-34.2
1994	917	32.4	30.3-34.6
1995	847	29.6	27.6-31.7
1996	721	25.0	23.2-26.9
1997	749	26.0	24.2-28.0
1998	643	22.0	20.3-23.8
1999	670	22.6	20.9-24.4
2000	641	20.9	19.3-22.6
2001	598	19.2	17.7-20.8
2002	595	18.9	17.4-20.5
2003	534	16.8	15.4-18.3
2004	613	18.4	16.9-19.9
2005	658	19.6	18.1-21.1
2006	718	21.2	19.7-22.8
2007	754	22.0	20.4-23.7
2008	723	20.3	18.8-21.9
2009	818	22.5	21.0-24.1
2010	821	22.2	20.6-23.8
2011	875	23.2	21.6-24.8
2012	802	20.4	19.0-21.9
2013	924	23.3	21.8-24.9
2014	917	22.6	21.1-24.1
2015	991	23.8	22.3-25.3
2016	1,041	24.7	23.1-26.2
2017	1,113	25.8	24.3-27.4
2018	1,117	25.7	24.2-27.3
2019	1,195	26.8	25.3-28.4
2020	1,061	23.5	22.1-25.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.



**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York City, Prostate, White Individuals**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	548	23.5	21.4-25.6
1977	527	22.4	20.4-24.5
1978	545	23.4	21.4-25.5
1979	573	25.5	23.4-27.8
1980	568	25.2	23.1-27.4
1981	581	26.2	24.0-28.5
1982	614	27.6	25.4-29.9
1983	576	26.0	23.9-28.3
1984	598	26.9	24.8-29.3
1985	581	26.3	24.1-28.6
1986	568	26.2	24.0-28.5
1987	554	25.2	23.2-27.5
1988	521	24.6	22.5-26.9
1989	539	25.7	23.6-28.0
1990	585	28.4	26.1-30.8
1991	639	30.7	28.4-33.3
1992	598	28.8	26.5-31.2
1993	560	27.4	25.1-29.7
1994	535	26.0	23.8-28.3
1995	462	22.7	20.7-24.9
1996	422	20.8	18.8-22.8
1997	411	20.4	18.5-22.5
1998	341	16.9	15.1-18.8
1999	362	17.9	16.1-19.8
2000	340	16.5	14.8-18.4
2001	330	15.8	14.1-17.6
2002	334	16.0	14.4-17.9
2003	309	14.7	13.1-16.4
2004	340	15.9	14.3-17.7
2005	378	17.6	15.8-19.5
2006	412	19.2	17.4-21.2
2007	436	20.1	18.3-22.1
2008	404	18.5	16.7-20.4
2009	461	20.6	18.8-22.6
2010	460	20.4	18.6-22.4
2011	476	20.9	19.0-22.9
2012	430	18.4	16.7-20.3
2013	481	20.4	18.6-22.4
2014	483	20.1	18.4-22.1
2015	502	20.7	18.9-22.6
2016	550	22.4	20.5-24.4
2017	580	23.6	21.7-25.6
2018	595	24.0	22.1-26.1
2019	596	23.6	21.8-25.6
2020	553	21.9	20.1-23.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York City, Prostate, Black Individuals**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	171	48.3	40.2-57.2
1977	185	51.2	43.1-60.1
1978	205	57.7	49.0-67.3
1979	224	63.0	54.1-72.8
1980	285	76.7	67.0-87.3
1981	250	65.7	56.9-75.3
1982	252	66.0	57.3-75.5
1983	229	55.0	47.3-63.4
1984	255	59.3	51.5-67.9
1985	291	63.2	55.4-71.7
1986	276	60.9	53.2-69.2
1987	285	59.9	52.5-67.9
1988	277	62.1	54.3-70.5
1989	299	61.3	53.9-69.3
1990	325	63.6	56.2-71.5
1991	315	60.1	53.0-67.7
1992	329	61.3	54.3-68.9
1993	306	56.3	49.7-63.5
1994	348	61.9	55.1-69.3
1995	342	57.1	50.8-63.9
1996	265	43.2	37.8-49.2
1997	307	50.6	44.7-57.1
1998	264	42.0	36.7-47.7
1999	290	43.4	38.2-49.0
2000	275	38.9	34.2-44.1
2001	239	34.8	30.3-39.7
2002	237	32.1	27.9-36.6
2003	202	27.0	23.2-31.2
2004	247	31.3	27.4-35.7
2005	239	30.1	26.2-34.3
2006	265	32.6	28.6-37.0
2007	283	34.6	30.5-39.1
2008	292	32.7	28.9-36.9
2009	300	33.6	29.7-37.8
2010	314	34.5	30.6-38.7
2011	336	35.7	31.8-39.9
2012	311	31.1	27.6-34.9
2013	370	36.9	33.1-41.1
2014	357	34.6	30.9-38.5
2015	381	36.0	32.3-40.0
2016	388	36.4	32.8-40.4
2017	409	35.9	32.4-39.7
2018	406	35.6	32.1-39.4
2019	453	39.2	35.6-43.1
2020	381	32.5	29.2-36.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State Excl New York City, Prostate, All Races**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	886	25.9	24.2-27.8
1977	899	25.4	23.7-27.2
1978	882	24.8	23.1-26.6
1979	916	26.2	24.4-28.0
1980	1,009	27.6	25.9-29.5
1981	1,090	29.8	28.0-31.8
1982	1,078	29.3	27.4-31.2
1983	1,069	27.8	26.1-29.7
1984	1,084	28.1	26.3-29.9
1985	1,155	29.2	27.4-31.0
1986	1,177	29.7	27.9-31.5
1987	1,225	30.5	28.8-32.4
1988	1,303	31.4	29.6-33.2
1989	1,235	29.5	27.8-31.2
1990	1,408	32.4	30.6-34.2
1991	1,581	35.9	34.1-37.8
1992	1,756	37.5	35.7-39.4
1993	1,583	33.8	32.1-35.6
1994	1,326	28.1	26.6-29.7
1995	1,234	25.7	24.2-27.2
1996	1,185	24.6	23.2-26.1
1997	1,134	23.5	22.1-24.9
1998	980	20.0	18.8-21.4
1999	934	18.6	17.4-19.8
2000	973	19.4	18.1-20.6
2001	872	16.8	15.7-18.0
2002	881	16.8	15.7-17.9
2003	938	17.5	16.4-18.7
2004	1,099	19.8	18.7-21.1
2005	1,114	19.4	18.3-20.6
2006	1,236	21.4	20.2-22.7
2007	1,250	21.1	20.0-22.4
2008	1,373	22.5	21.3-23.7
2009	1,391	22.1	21.0-23.4
2010	1,495	23.3	22.1-24.6
2011	1,436	22.0	20.9-23.2
2012	1,333	20.0	18.9-21.2
2013	1,409	20.8	19.7-21.9
2014	1,492	21.7	20.6-22.9
2015	1,591	22.7	21.6-23.9
2016	1,750	24.6	23.5-25.9
2017	1,843	25.5	24.3-26.7
2018	1,979	27.1	25.9-28.3
2019	2,057	27.6	26.4-28.8
2020	1,880	24.8	23.7-26.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State Excl New York City, Prostate, White Individuals**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	838	25.4	23.6-27.3
1977	834	24.5	22.8-26.3
1978	811	23.8	22.1-25.5
1979	862	25.6	23.9-27.5
1980	940	26.9	25.2-28.8
1981	1,016	29.0	27.1-30.9
1982	1,011	28.6	26.8-30.6
1983	983	26.7	24.9-28.5
1984	1,011	27.3	25.6-29.2
1985	1,073	28.3	26.6-30.1
1986	1,078	28.4	26.7-30.3
1987	1,141	29.8	28.0-31.7
1988	1,202	30.4	28.6-32.2
1989	1,157	28.9	27.2-30.7
1990	1,312	31.6	29.8-33.4
1991	1,462	34.9	33.0-36.8
1992	1,629	36.7	34.9-38.6
1993	1,455	32.8	31.0-34.5
1994	1,185	26.6	25.1-28.2
1995	1,105	24.4	22.9-25.9
1996	1,061	23.5	22.1-25.0
1997	988	21.9	20.5-23.3
1998	863	18.9	17.6-20.2
1999	831	17.7	16.5-18.9
2000	858	18.4	17.2-19.7
2001	774	16.1	15.0-17.3
2002	772	15.8	14.7-17.0
2003	824	16.7	15.5-17.8
2004	968	19.0	17.8-20.2
2005	1,006	19.1	17.9-20.3
2006	1,093	20.7	19.5-22.0
2007	1,107	20.5	19.3-21.7
2008	1,216	21.8	20.6-23.1
2009	1,219	21.3	20.1-22.6
2010	1,335	22.9	21.7-24.2
2011	1,269	21.5	20.3-22.7
2012	1,183	19.5	18.4-20.7
2013	1,239	20.2	19.1-21.4
2014	1,295	20.9	19.7-22.1
2015	1,403	22.2	21.0-23.4
2016	1,527	23.9	22.7-25.1
2017	1,636	25.1	23.9-26.4
2018	1,718	26.2	25.0-27.5
2019	1,771	26.5	25.3-27.8
2020	1,623	24.0	22.8-25.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by Year of Diagnosis, 1976-2020  
New York State Excl New York City, Prostate, Black Individuals**

Year of diagnosis	Cases	Rate per 100,000	95% CI
1976	45	46.5	32.3-63.9
1977	61	52.5	38.6-69.2
1978	68	56.1	42.0-72.9
1979	51	45.8	32.3-62.2
1980	67	51.2	38.3-66.6
1981	70	55.9	42.2-72.1
1982	62	46.6	34.9-60.6
1983	78	60.1	46.0-76.6
1984	71	52.1	39.5-67.0
1985	79	54.7	42.0-69.6
1986	98	68.2	54.0-84.5
1987	81	52.3	40.5-66.0
1988	95	60.0	47.4-74.5
1989	76	49.0	37.6-62.4
1990	91	54.9	43.1-68.7
1991	112	66.2	53.3-80.9
1992	114	59.0	47.6-72.1
1993	116	62.8	50.8-76.6
1994	127	66.0	53.9-79.7
1995	121	58.9	47.8-71.5
1996	116	51.4	41.7-62.4
1997	133	58.4	48.1-70.0
1998	112	49.5	40.0-60.3
1999	94	40.5	32.0-50.4
2000	108	42.4	34.1-51.8
2001	82	29.9	23.3-37.6
2002	100	39.6	31.4-49.2
2003	101	34.7	27.7-42.7
2004	120	40.6	33.0-49.2
2005	92	28.6	22.5-35.7
2006	125	35.4	29.1-42.6
2007	123	34.7	28.4-41.9
2008	142	36.9	30.7-43.9
2009	144	37.1	30.9-44.2
2010	144	35.1	29.3-41.7
2011	130	30.3	25.1-36.4
2012	126	29.5	24.3-35.5
2013	145	33.0	27.6-39.2
2014	166	36.0	30.4-42.4
2015	148	31.2	26.1-37.0
2016	183	37.9	32.4-44.2
2017	167	33.8	28.6-39.6
2018	202	40.6	34.9-46.9
2019	225	43.0	37.3-49.3
2020	186	33.5	28.6-38.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Cancer diagnoses trends in 2020 may have been affected by the coronavirus disease 19 (COVID-19) pandemic. Rates based on fewer than 20 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by County, All Races, 2016-2020  
Colorectal**

County/Region	Males and Females			Males			Females		
	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI
New York State	5,152.4	21.5	21.3-21.8	2,697.8	25.1	24.7-25.5	2,454.6	18.6	18.3-18.9
New York City	2,016.8	21.1	20.7-21.5	1,058.0	25.3	24.6-26.0	958.8	17.8	17.3-18.4
New York State Excl New York City	3,133.8	21.8	21.5-22.2	1,638.6	25.0	24.4-25.5	1,495.2	19.2	18.7-19.6
Bronx	311.2	21.3	20.2-22.4	159.6	25.8	24.0-27.7	151.6	17.9	16.6-19.2
Kings (Brooklyn)	621.8	22.3	21.5-23.1	324.8	27.1	25.7-28.4	297.0	18.8	17.8-19.8
New York (Manhattan)	361.4	18.4	17.5-19.2	192.8	22.4	21.0-23.9	168.6	15.1	14.1-16.2
Queens	575.2	20.9	20.2-21.7	300.6	24.3	23.0-25.6	274.6	18.2	17.2-19.2
Richmond (Staten Island)	147.2	25.1	23.3-27.0	80.2	30.3	27.3-33.5	67.0	20.8	18.6-23.3
Albany	84.2	22.3	20.1-24.6	43.8	27.0	23.5-31.0	40.4	18.4	15.9-21.3
Allegany	15.6	27.5	21.4-34.9	9.0	33.3	23.9-45.3	6.6	22.1	14.7-32.1
Broome	50.4	19.5	17.1-22.3	26.4	22.4	18.6-26.8	24.0	17.2	14.1-20.8
Cattaraugus	24.8	24.8	20.4-29.9	12.8	27.1	20.5-35.3	12.0	22.5	16.9-29.6
Cayuga	20.8	20.3	16.4-24.9	9.8	20.7	15.1-27.7	11.0	19.1	14.1-25.6
Chautauqua	33.8	19.8	16.8-23.2	17.0	21.9	17.3-27.4	16.8	17.6	13.8-22.2
Chemung	19.8	18.2	14.6-22.5	13.0	24.9	19.0-32.1	6.8	12.7	8.5-18.3
Chenango	15.2	22.2	17.3-28.2	8.2	24.7	17.4-34.3	7.0	19.6	13.4-27.9
Clinton	24.4	25.0	20.6-30.1	13.2	27.8	21.3-35.6	11.2	22.3	16.5-29.6
Columbia	21.0	21.3	17.3-26.2	10.6	22.9	16.9-30.7	10.4	19.4	14.2-26.3
Cortland	15.2	27.0	21.1-34.2	8.2	32.7	23.2-44.8	7.0	21.9	15.0-31.1
Delaware	18.0	25.5	20.2-32.1	10.4	30.4	22.3-41.0	7.6	21.6	14.7-31.0
Dutchess	74.8	19.3	17.3-21.4	39.8	21.6	18.6-25.0	35.0	17.2	14.6-20.1
Erie	264.8	22.2	21.0-23.5	133.6	25.3	23.3-27.3	131.2	19.7	18.1-21.3
Essex	12.2	21.8	16.3-28.8	5.6	22.0	14.1-33.1	6.6	20.2	13.7-29.7
Franklin	15.0	23.8	18.6-30.1	6.0	18.5	12.3-26.8	9.0	27.9	19.9-38.2
Fulton	18.6	24.5	19.6-30.3	8.2	24.2	17.1-33.6	10.4	24.5	18.1-32.8
Genesee	18.2	23.6	18.8-29.5	10.8	29.0	21.5-38.4	7.4	18.8	12.8-26.9
Greene	19.0	26.6	21.2-33.1	11.2	31.9	23.8-42.3	7.8	22.2	15.3-31.6
Hamilton	1.4	12.4	4.8-37.8	0.6	11.3	2.1-61.0	0.8	13.3	3.6-62.1
Herkimer	22.6	26.2	21.3-31.8	12.6	31.8	24.1-41.3	10.0	20.4	14.9-27.5
Jefferson	32.0	28.3	24.0-33.2	15.2	28.8	22.6-36.3	16.8	28.1	22.3-35.1
Lewis	10.6	32.3	23.8-43.1	4.8	31.4	19.7-47.7	5.8	33.0	21.5-49.0
Livingston	17.4	21.2	16.9-26.5	10.6	26.7	19.8-35.3	6.8	17.7	12.0-25.3
Madison	21.0	22.9	18.6-28.1	11.4	27.6	20.6-36.4	9.6	18.8	13.7-25.4
Monroe	202.0	21.6	20.2-23.0	104.0	24.4	22.3-26.7	98.0	19.2	17.4-21.1
Montgomery	15.6	24.8	19.4-31.3	8.2	28.4	20.1-39.1	7.4	22.1	15.4-31.0
Nassau	381.8	21.5	20.5-22.5	193.2	24.2	22.7-25.8	188.6	19.3	18.0-20.6
Niagara	72.0	26.3	23.5-29.3	38.6	31.5	27.0-36.5	33.4	21.6	18.3-25.5
Oneida	59.0	19.4	17.2-21.9	31.2	22.5	19.0-26.4	27.8	17.1	14.3-20.5
Onondaga	120.6	21.0	19.2-22.8	58.8	22.6	20.0-25.4	61.8	19.5	17.3-22.0
Ontario	30.6	20.5	17.2-24.3	14.8	20.9	16.2-26.6	15.8	20.3	15.8-25.8
Orange	92.4	22.0	20.0-24.2	49.8	25.3	22.2-28.8	42.6	19.3	16.7-22.2
Orleans	15.6	28.7	22.4-36.4	8.6	32.3	22.9-44.5	7.0	25.4	17.3-36.4
Oswego	34.8	24.2	20.6-28.3	17.4	24.9	19.7-31.0	17.4	23.4	18.6-29.1
Otsego	22.0	27.6	22.3-33.9	13.0	34.4	26.0-44.8	9.0	21.5	15.1-29.8
Putnam	24.8	19.1	15.8-23.0	11.4	18.0	13.5-23.7	13.4	19.3	14.8-24.9
Rensselaer	38.4	19.9	17.1-23.1	20.0	23.2	18.7-28.4	18.4	17.1	13.7-21.2
Rockland	85.8	22.7	20.6-25.0	44.8	26.2	22.8-29.9	41.0	19.9	17.2-23.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Rates based on fewer than 4 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by County, All Races, 2016-2020  
Colorectal**

County/Region	Males and Females			Males			Females		
	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI
St. Lawrence	34.0	24.9	21.2-29.2	18.4	28.7	23.0-35.5	15.6	21.2	16.6-26.8
Saratoga	64.8	21.3	19.0-23.9	34.2	24.5	20.9-28.7	30.6	18.5	15.6-21.9
Schenectady	37.2	19.4	16.6-22.5	21.4	24.3	19.8-29.5	15.8	14.9	11.7-18.8
Schoharie	9.8	20.9	15.2-28.3	5.2	23.3	14.9-35.3	4.6	18.4	11.3-29.2
Schuyler	6.4	25.3	16.8-36.9	4.4	35.1	21.4-55.5	2.0	15.5	6.8-31.3
Seneca	10.2	22.9	16.8-30.7	5.8	25.0	16.4-36.8	4.4	21.4	12.9-33.6
Steuben	35.2	26.2	22.4-30.7	18.0	28.5	22.7-35.4	17.2	24.0	19.0-30.1
Suffolk	423.6	22.2	21.2-23.2	229.0	26.3	24.8-27.9	194.6	18.8	17.6-20.1
Sullivan	24.8	24.4	20.2-29.4	14.4	28.1	21.8-35.8	10.4	20.6	15.1-27.6
Tioga	15.2	22.5	17.4-28.7	8.2	24.8	17.4-34.5	7.0	20.5	13.8-29.6
Tompkins	16.2	15.0	11.8-18.9	8.8	17.7	12.7-24.1	7.4	12.7	8.8-17.8
Ulster	58.8	24.1	21.3-27.1	30.8	27.3	23.0-32.2	28.0	21.2	17.6-25.3
Warren	18.2	19.5	15.4-24.4	9.2	21.0	15.0-28.7	9.0	18.1	12.7-25.2
Washington	19.2	22.9	18.4-28.3	9.2	22.6	16.3-30.6	10.0	23.0	16.7-31.1
Wayne	32.8	27.0	22.8-31.7	17.6	30.3	24.0-37.8	15.2	23.9	18.6-30.4
Westchester	244.4	19.6	18.5-20.8	127.4	23.1	21.3-25.1	117.0	16.8	15.4-18.2
Wyoming	12.4	24.7	18.8-32.1	7.4	28.9	20.1-40.4	5.0	21.5	13.4-33.0
Yates	4.4	11.0	6.7-17.5	2.6	14.7	7.5-26.7	1.8	7.9	3.5-16.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Rates based on fewer than 4 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by County, All Races, 2016-2020  
Lung and Bronchus**

County/Region	Males and Females			Males			Females		
	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI
New York State	8,888.2	35.3	35.0-35.6	4,490.4	40.4	39.8-40.9	4,397.8	31.5	31.1-32.0
New York City	2,731.4	27.8	27.3-28.3	1,433.8	34.2	33.4-35.0	1,297.6	23.1	22.6-23.7
New York State Excl New York City	6,155.8	40.1	39.6-40.5	3,056.2	44.1	43.4-44.8	3,099.6	37.2	36.6-37.8
Bronx	446.6	29.6	28.4-30.9	237.2	38.2	36.0-40.5	209.4	23.6	22.2-25.1
Kings (Brooklyn)	782.0	27.6	26.7-28.4	420.4	35.1	33.6-36.7	361.6	22.2	21.2-23.3
New York (Manhattan)	510.6	24.9	23.9-25.9	242.8	27.8	26.2-29.4	267.8	22.8	21.6-24.1
Queens	757.8	26.7	25.8-27.6	416.4	33.7	32.2-35.2	341.4	21.4	20.4-22.5
Richmond (Staten Island)	234.4	38.9	36.7-41.3	117.0	44.1	40.5-48.0	117.4	35.2	32.4-38.2
Albany	178.2	44.5	41.6-47.6	86.6	49.3	44.6-54.4	91.6	41.5	37.7-45.6
Allegany	32.6	47.8	40.6-56.1	17.4	51.7	41.2-64.4	15.2	44.2	34.5-56.1
Broome	116.6	42.9	39.3-46.7	60.0	48.0	42.6-53.9	56.6	39.1	34.5-44.3
Cattaraugus	52.0	45.3	39.9-51.5	25.4	46.5	38.5-55.9	26.6	45.1	37.5-53.9
Cayuga	55.4	48.1	42.4-54.4	26.2	47.4	39.4-56.7	29.2	48.8	41.0-57.9
Chautauqua	89.2	46.8	42.4-51.5	43.6	48.4	42.0-55.7	45.6	45.9	39.9-52.6
Chemung	67.8	55.4	49.5-61.8	35.0	61.6	52.7-71.9	32.8	50.3	42.7-59.1
Chenango	34.4	46.6	39.6-54.6	16.8	44.6	35.3-55.8	17.6	49.2	38.9-61.8
Clinton	55.0	50.2	44.3-56.7	29.8	57.7	48.6-68.1	25.2	43.9	36.3-52.7
Columbia	39.8	39.7	34.1-46.1	20.4	41.2	33.4-50.8	19.4	38.6	30.9-48.2
Cortland	35.4	58.0	49.6-67.5	18.6	65.5	52.5-80.9	16.8	51.9	41.2-64.8
Delaware	33.6	43.0	36.4-50.8	18.2	47.2	37.3-59.4	15.4	39.1	30.4-50.0
Dutchess	159.4	38.8	36.1-41.7	78.6	40.8	36.7-45.2	80.8	37.5	33.9-41.5
Erie	566.4	44.2	42.6-45.9	269.2	48.2	45.6-51.0	297.2	41.5	39.4-43.8
Essex	25.8	39.7	32.9-47.8	13.2	44.5	34.0-57.7	12.6	37.2	28.3-48.9
Franklin	31.2	45.8	38.8-53.9	19.0	56.0	45.0-69.1	12.2	36.1	27.4-47.0
Fulton	42.6	53.3	46.3-61.3	19.6	52.0	42.0-64.0	23.0	55.2	45.3-66.9
Genesee	42.8	50.1	43.4-57.5	22.2	55.1	45.1-67.0	20.6	46.6	37.8-57.1
Greene	39.0	49.6	42.7-57.5	21.2	55.1	44.9-67.4	17.8	44.4	35.4-55.6
Hamilton	4.8	49.0	30.8-81.6	3.0	58.4	32.2-115.8	1.8	39.1	16.9-95.0
Herkimer	41.2	43.9	37.9-50.7	21.4	48.2	39.2-59.0	19.8	39.4	31.9-48.5
Jefferson	68.2	56.3	50.4-62.7	35.0	63.6	54.3-74.1	33.2	50.4	42.9-59.0
Lewis	13.4	35.8	27.6-46.1	7.2	39.8	27.5-56.2	6.2	31.9	21.4-46.4
Livingston	41.4	45.7	39.5-52.7	19.8	46.5	37.6-57.2	21.6	45.4	37.1-55.5
Madison	39.6	37.7	32.5-43.6	21.6	42.4	34.6-51.6	18.0	33.6	26.8-41.9
Monroe	400.6	40.3	38.5-42.2	205.6	47.0	44.1-50.0	195.0	35.6	33.3-38.0
Montgomery	37.8	53.6	46.0-62.1	19.6	59.3	47.8-72.9	18.2	49.2	39.3-61.0
Nassau	567.2	30.1	29.0-31.2	264.8	32.0	30.3-33.8	302.4	28.9	27.4-30.4
Niagara	147.2	47.0	43.6-50.7	73.0	51.0	45.7-56.7	74.2	44.8	40.3-49.9
Oneida	151.8	47.1	43.7-50.7	79.0	53.8	48.5-59.6	72.8	41.7	37.4-46.5
Onondaga	266.4	43.4	41.1-45.9	131.0	48.2	44.5-52.2	135.4	40.0	36.9-43.2
Ontario	66.6	40.3	35.9-45.0	33.2	43.5	36.9-51.0	33.4	38.1	32.4-44.7
Orange	177.6	41.0	38.3-43.8	86.2	43.7	39.5-48.2	91.4	39.3	35.8-43.2
Orleans	36.4	63.8	54.6-74.2	18.4	69.8	55.8-86.6	18.0	59.8	47.8-74.2
Oswego	97.6	62.5	56.9-68.5	53.4	73.2	64.4-83.1	44.2	53.0	46.1-60.7
Otsego	42.6	45.1	39.1-52.0	22.4	50.5	41.4-61.4	20.2	40.6	32.6-50.2
Putnam	52.0	38.4	33.8-43.6	26.6	42.6	35.4-50.9	25.4	35.3	29.3-42.3
Rensselaer	105.4	49.2	45.0-53.7	52.0	54.5	47.9-61.9	53.4	45.8	40.3-51.9
Rockland	122.2	31.1	28.7-33.7	59.0	34.2	30.4-38.4	63.2	28.7	25.5-32.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Rates based on fewer than 4 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.



**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by County, All Races, 2016-2020  
Lung and Bronchus**

County/Region	Males and Females			Males			Females		
	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI
St. Lawrence	73.8	50.3	45.1-55.8	37.6	54.2	46.5-62.9	36.2	47.3	40.5-55.0
Saratoga	140.0	42.7	39.5-46.1	72.8	48.8	43.8-54.2	67.2	38.3	34.3-42.8
Schenectady	94.2	45.1	41.0-49.5	46.8	52.0	45.4-59.4	47.4	40.6	35.5-46.3
Schoharie	22.8	45.7	37.3-55.7	12.2	50.3	38.1-66.0	10.6	41.0	30.2-55.3
Schuyler	15.4	53.3	41.7-67.8	8.6	61.4	43.9-84.9	6.8	46.5	31.6-67.4
Seneca	24.0	49.5	40.8-59.7	11.8	50.3	38.0-65.6	12.2	50.4	38.0-66.1
Steuben	62.2	43.7	38.9-49.1	33.2	49.1	41.7-57.6	29.0	38.8	32.7-46.0
Suffolk	752.4	37.8	36.5-39.0	368.0	41.7	39.8-43.7	384.4	35.0	33.5-36.7
Sullivan	47.4	43.0	37.6-49.1	24.4	45.9	37.9-55.2	23.0	41.0	33.7-49.6
Tioga	30.6	41.1	34.7-48.5	17.0	47.9	38.1-59.8	13.6	34.8	26.7-44.9
Tompkins	40.8	35.5	30.7-41.0	21.6	42.5	34.7-51.8	19.2	30.6	24.6-37.7
Ulster	117.0	42.8	39.3-46.5	58.8	46.0	40.7-51.8	58.2	40.0	35.4-45.1
Warren	49.0	45.1	39.5-51.4	27.0	52.4	43.8-62.7	22.0	38.7	31.6-47.3
Washington	44.8	48.5	42.2-55.6	19.4	44.6	35.9-54.9	25.4	52.8	43.8-63.4
Wayne	69.0	52.6	47.0-58.7	36.4	59.2	50.6-68.9	32.6	46.9	39.8-55.1
Westchester	354.6	27.0	25.8-28.3	165.2	29.1	27.2-31.2	189.4	25.5	23.8-27.2
Wyoming	23.8	41.6	34.3-50.1	13.4	47.1	36.1-60.6	10.4	36.4	27.0-48.6
Yates	16.8	45.6	36.0-57.3	8.8	51.2	36.5-70.3	8.0	40.8	28.7-57.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Rates based on fewer than 4 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by County, All Races, 2016-2020  
Female Breast**

County/Region	Avg Ann Cases	Rate per 100,000	95% CI
New York State	4,946.4	41.1	40.6-41.6
New York City	2,077.4	41.7	40.9-42.5
New York State Excl New York City	2,868.0	40.6	39.9-41.3
Bronx	326.8	40.0	38.1-42.1
Kings (Brooklyn)	662.2	44.8	43.3-46.4
New York (Manhattan)	387.2	39.7	37.9-41.5
Queens	571.2	40.7	39.2-42.3
Richmond (Staten Island)	130.0	43.2	39.8-46.8
Albany	76.8	41.2	37.0-45.9
Allegany	9.0	37.5	26.5-51.4
Broome	48.2	39.8	34.5-45.7
Cattaraugus	20.2	41.0	32.9-50.6
Cayuga	17.4	33.8	26.5-42.6
Chautauqua	34.6	40.4	34.2-47.5
Chemung	26.0	46.5	38.3-56.0
Chenango	12.4	37.4	28.1-49.0
Clinton	18.0	36.9	29.2-46.0
Columbia	20.6	46.2	36.9-57.5
Cortland	12.2	46.4	34.8-60.7
Delaware	12.4	38.5	28.6-51.0
Dutchess	70.6	38.4	34.2-42.9
Erie	250.6	42.1	39.7-44.7
Essex	10.2	35.6	25.8-48.6
Franklin	10.0	35.6	26.0-47.9
Fulton	12.4	32.1	24.1-42.1
Genesee	14.6	36.9	28.5-47.3
Greene	13.6	40.2	30.4-52.4
Hamilton	1.6	35.0	14.1-90.3
Herkimer	12.6	30.2	22.7-39.5
Jefferson	23.0	40.2	32.9-48.6
Lewis	6.4	37.7	25.1-54.8
Livingston	11.8	29.9	22.2-39.4
Madison	18.4	42.2	33.4-52.6
Monroe	188.8	40.3	37.6-43.1
Montgomery	12.8	40.7	30.8-53.0
Nassau	363.0	40.9	39.0-43.0
Niagara	56.0	38.8	34.1-44.0
Oneida	50.2	33.7	29.4-38.5
Onondaga	110.0	38.2	34.9-41.7
Ontario	26.6	36.2	29.8-43.6
Orange	90.6	42.9	38.9-47.2
Orleans	8.4	33.1	23.4-45.6
Oswego	25.8	39.4	32.5-47.3
Otsego	17.8	47.6	37.2-60.1
Putnam	28.2	43.2	36.0-51.6
Rensselaer	38.4	37.5	32.2-43.5
Rockland	71.4	39.1	34.9-43.6
St. Lawrence	21.6	32.5	26.3-39.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Rates based on fewer than 4 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by County, All Races, 2016-2020  
Female Breast**

County/Region	Avg Ann Cases	Rate per 100,000	95% CI
Saratoga	63.8	41.6	37.0-46.7
Schenectady	41.0	41.4	35.6-47.8
Schoharie	8.0	41.0	28.2-58.0
Schuyler	5.6	45.8	29.3-69.0
Seneca	7.4	38.5	26.2-54.6
Steuben	25.6	41.9	34.4-50.6
Suffolk	394.4	41.6	39.6-43.5
Sullivan	23.0	48.0	39.2-58.4
Tioga	15.0	46.8	36.0-60.0
Tompkins	23.0	43.7	35.7-53.0
Ulster	52.6	44.6	39.0-50.8
Warren	21.0	44.2	35.6-54.6
Washington	13.2	36.6	27.6-47.6
Wayne	27.4	47.3	39.1-56.7
Westchester	255.4	41.4	39.1-43.9
Wyoming	13.4	51.4	39.2-66.5
Yates	5.0	34.7	21.4-53.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Rates based on fewer than 4 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by County, All Races, 2016-2020  
Cervix**

County/Region	Avg Ann Cases	Rate per 100,000	95% CI
New York State	408.8	3.5	3.4-3.7
New York City	214.4	4.4	4.1-4.6
New York State Excl New York City	194.4	3.0	2.8-3.2
Bronx	41.2	5.1	4.4-5.9
Kings (Brooklyn)	70.2	4.7	4.2-5.3
New York (Manhattan)	33.8	3.5	3.0-4.1
Queens	59.4	4.4	3.9-4.9
Richmond (Staten Island)	9.8	3.3	2.4-4.4
Albany	4.8	3.1	1.9-4.6
Allegany	0.8	3.4	0.9-9.1
Broome	3.0	2.9	1.5-4.9
Cattaraugus	2.0	4.8	2.1-9.2
Cayuga	0.8	2.5	0.6-6.3
Chautauqua	2.2	2.6	1.2-4.9
Chemung	1.6	2.9	1.2-6.2
Chenango	1.6	4.7	1.9-10.0
Clinton	1.2	3.2	1.1-7.2
Columbia	1.2	2.5	0.9-6.2
Cortland	0.4	2.3	0.2-7.8
Delaware	0.8	3.6	0.8-10.0
Dutchess	5.2	2.9	1.8-4.3
Erie	17.0	3.4	2.6-4.2
Essex	1.0	5.2	1.4-13.3
Franklin	0.8	2.6	0.7-7.4
Fulton	1.0	3.0	0.9-7.6
Genesee	0.4	1.7	0.2-6.0
Greene	0.6	2.4	0.4-7.8
Hamilton	0.2	2.8	0.1-51.4
Herkimer	1.0	2.6	0.8-6.4
Jefferson	1.4	2.7	1.0-5.7
Lewis	0.8	5.4	1.2-15.0
Livingston	1.0	3.5	1.0-8.5
Madison	0.4	1.1	0.1-4.5
Monroe	12.6	2.9	2.2-3.7
Montgomery	0.8	2.7	0.7-7.5
Nassau	25.4	3.0	2.5-3.6
Niagara	4.6	3.4	2.1-5.4
Oneida	3.6	2.6	1.5-4.2
Onondaga	7.8	2.9	2.0-4.0
Ontario	2.0	2.9	1.3-5.6
Orange	8.8	4.5	3.2-6.0
Orleans	0.8	3.7	0.9-9.9
Oswego	1.8	2.8	1.2-5.5
Otsego	1.4	3.5	1.1-8.3
Putnam	2.2	4.4	2.1-8.2
Rensselaer	3.2	4.0	2.2-6.5
Rockland	4.2	2.0	1.2-3.1
St. Lawrence	0.8	1.4	0.4-3.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Rates based on fewer than 4 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by County, All Races, 2016-2020**  
**Cervix**

County/Region	Avg Ann Cases	Rate per 100,000	95% CI
Saratoga	3.4	2.6	1.5-4.2
Schenectady	2.6	3.1	1.6-5.4
Schoharie	0.6	2.3	0.5-8.4
Schuyler	0.4	4.3	0.3-17.1
Seneca	1.2	6.6	2.2-15.4
Steuben	2.2	3.4	1.6-6.4
Suffolk	25.0	2.8	2.3-3.3
Sullivan	3.2	6.6	3.6-11.2
Tioga	0.8	3.1	0.8-8.4
Tompkins	0.6	1.4	0.2-4.1
Ulster	4.0	4.3	2.5-6.8
Warren	0.6	1.0	0.2-3.8
Washington	1.6	5.6	2.3-11.3
Wayne	0.8	1.5	0.4-4.1
Westchester	15.0	2.5	1.9-3.2
Wyoming	0.8	2.7	0.7-8.1
Yates	0.4	3.1	0.3-12.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Rates based on fewer than 4 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by County, All Races, 2016-2020  
Prostate**

County/Region	Avg Ann Cases	Rate per 100,000	95% CI
New York State	3,008.4	25.7	25.3-26.1
New York City	1,105.4	25.3	24.6-26.0
New York State Excl New York City	1,901.8	25.9	25.4-26.5
Bronx	193.8	30.3	28.4-32.3
Kings (Brooklyn)	313.0	25.1	23.8-26.4
New York (Manhattan)	270.2	30.1	28.5-31.7
Queens	263.6	20.2	19.1-21.3
Richmond (Staten Island)	64.8	23.1	20.6-25.9
Albany	54.6	29.5	26.0-33.3
Allegany	10.8	32.2	24.0-42.6
Broome	45.0	34.1	29.7-39.0
Cattaraugus	14.4	24.0	18.6-30.6
Cayuga	12.8	22.4	17.2-29.0
Chautauqua	16.0	16.8	13.2-21.1
Chemung	13.6	22.1	17.1-28.4
Chenango	12.6	32.5	24.8-42.2
Clinton	11.4	20.5	15.4-26.9
Columbia	16.4	30.3	24.0-38.2
Cortland	6.8	24.2	16.6-34.3
Delaware	9.8	24.4	17.9-33.3
Dutchess	50.0	24.9	21.9-28.4
Erie	123.2	20.4	18.8-22.1
Essex	6.6	19.6	13.4-28.6
Franklin	10.0	29.1	21.5-38.9
Fulton	11.0	28.5	21.3-37.7
Genesee	12.4	29.1	22.2-37.7
Greene	8.0	19.3	13.8-27.0
Hamilton	2.2	45.7	21.8-102.6
Herkimer	13.0	27.4	21.0-35.4
Jefferson	14.4	24.3	18.9-30.8
Lewis	4.2	22.7	13.8-35.7
Livingston	19.8	43.5	35.1-53.4
Madison	11.0	20.5	15.3-27.2
Monroe	182.4	38.7	36.2-41.4
Montgomery	8.8	25.0	18.0-34.1
Nassau	190.4	21.5	20.1-22.9
Niagara	33.8	22.4	19.0-26.2
Oneida	37.4	24.3	20.9-28.2
Onondaga	61.2	20.7	18.4-23.3
Ontario	40.0	48.1	41.5-55.6
Orange	49.0	23.4	20.4-26.6
Orleans	7.6	27.7	19.4-38.8
Oswego	15.2	20.9	16.3-26.5
Otsego	11.8	25.6	19.4-33.6
Putnam	19.0	26.4	21.2-32.7
Rensselaer	25.2	24.8	20.5-29.8
Rockland	35.6	19.5	16.7-22.7
St. Lawrence	14.6	21.0	16.3-26.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Rates based on fewer than 4 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.

**New York State Department of Health, New York State Cancer Registry  
Regional/Distant Stage Cancer Incidence by County, All Races, 2016-2020  
Prostate**

County/Region	Avg Ann Cases	Rate per 100,000	95% CI
Saratoga	43.8	27.9	24.2-32.0
Schenectady	24.6	26.6	22.0-31.9
Schoharie	10.0	39.2	28.9-52.8
Schuyler	5.0	37.3	23.3-57.6
Seneca	10.8	41.6	31.0-55.1
Steuben	23.0	32.0	26.2-38.7
Suffolk	252.2	27.1	25.6-28.6
Sullivan	10.6	19.1	14.2-25.5
Tioga	9.8	25.5	18.8-34.3
Tompkins	16.0	29.0	22.8-36.5
Ulster	28.6	21.7	18.2-25.8
Warren	14.6	27.6	21.5-35.4
Washington	10.4	22.3	16.5-29.7
Wayne	27.0	41.5	34.6-49.6
Westchester	156.8	26.3	24.5-28.3
Wyoming	8.2	28.2	19.9-39.1
Yates	8.4	44.8	31.9-62.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022. Rates based on fewer than 4 cases in one year are unstable and should be used with caution. Regional/Distant stage indicates disease has spreaded beyond the limits of the organ of origin at diagnosis.