

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**  
**New York State**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	67,863.8	404.1	402.7-405.5	33,156.8	481.9	479.4-484.3	34,707.0	362.4	360.7-364.2	36,737.4	220.5	219.4-221.5	19,028.2	283.3	281.4-285.2	17,709.2	182.5	181.3-183.7
	1981-1985	73,548.0	426.2	424.8-427.6	35,888.6	510.1	507.7-512.6	37,659.4	381.7	379.9-383.4	37,507.8	218.3	217.3-219.3	19,184.8	281.0	279.1-282.8	18,323.0	181.6	180.4-182.8
	1986-1990	77,993.4	440.6	439.2-442.0	37,924.6	524.0	521.5-526.4	40,068.8	396.4	394.6-398.2	38,207.0	216.7	215.8-217.7	19,326.8	277.0	275.2-278.8	18,880.2	181.7	180.5-182.9
	1991-1995	86,767.0	476.4	475.0-477.8	43,443.2	574.7	572.2-577.2	43,323.8	418.9	417.2-420.7	38,574.6	212.7	211.8-213.7	19,247.6	268.4	266.6-270.1	19,327.0	180.7	179.6-181.9
	1996-2000	93,382.8	494.3	492.9-495.7	46,446.6	583.8	581.4-586.2	46,936.2	438.6	436.8-440.4	37,355.8	197.5	196.6-198.4	18,446.6	243.2	241.6-244.8	18,909.2	169.9	168.8-171.0
	2001-2005	98,214.0	491.4	490.0-492.8	49,516.6	580.4	578.1-582.7	48,697.4	433.5	431.8-435.2	36,088.2	179.6	178.7-180.4	17,726.4	217.6	216.2-219.1	18,361.8	156.2	155.2-157.2
	2006-2010	105,094.2	499.6	498.3-501.0	53,354.0	580.2	578.0-582.5	51,740.2	445.2	443.5-447.0	35,034.4	165.1	164.3-165.9	17,389.8	198.1	196.8-199.5	17,644.6	143.9	142.9-144.9
	2011-2015	109,009.4	480.9	479.6-482.2	54,030.0	532.3	530.2-534.3	54,979.4	448.2	446.4-449.9	34,969.2	151.8	151.1-152.5	17,470.2	179.9	178.7-181.1	17,499.0	133.0	132.1-133.9
2016-2020	112,274.8	463.6	462.4-464.9	56,043.0	504.8	502.9-506.7	56,231.8	437.4	435.7-439.1	33,634.8	134.5	133.9-135.2	16,814.8	156.8	155.7-157.9	16,820.0	119.4	118.6-120.2	
Oral cavity and pharynx	1976-1980	1,931.8	11.2	11.0-11.5	1,291.8	17.6	17.1-18.0	640.0	6.6	6.3-6.8	823.8	4.8	4.7-5.0	571.8	7.9	7.6-8.2	252.0	2.6	2.4-2.7
	1981-1985	2,055.6	11.8	11.6-12.1	1,388.2	18.7	18.3-19.2	667.4	6.8	6.5-7.0	758.4	4.3	4.2-4.5	524.0	7.2	6.9-7.4	234.4	2.3	2.2-2.5
	1986-1990	1,945.8	11.1	10.9-11.3	1,276.0	16.9	16.5-17.4	669.8	6.7	6.4-6.9	690.8	3.9	3.8-4.1	455.0	6.2	6.0-6.5	235.8	2.3	2.2-2.4
	1991-1995	1,977.0	11.0	10.8-11.2	1,293.2	16.7	16.2-17.1	683.8	6.7	6.5-6.9	600.4	3.3	3.2-3.5	402.0	5.3	5.1-5.5	198.4	1.9	1.8-2.0
	1996-2000	1,925.2	10.2	10.0-10.5	1,263.8	15.3	14.9-15.7	661.4	6.2	6.0-6.4	512.6	2.7	2.6-2.8	338.8	4.2	4.0-4.4	173.8	1.6	1.5-1.7
	2001-2005	1,976.0	9.8	9.7-10.0	1,302.4	14.6	14.2-14.9	673.6	6.0	5.8-6.2	480.0	2.4	2.3-2.5	318.4	3.7	3.5-3.8	161.6	1.4	1.3-1.5
	2006-2010	2,229.0	10.5	10.3-10.7	1,503.2	15.6	15.3-16.0	725.8	6.2	6.0-6.4	459.4	2.1	2.1-2.2	315.2	3.4	3.2-3.6	144.2	1.2	1.1-1.3
	2011-2015	2,485.0	10.8	10.6-11.0	1,700.8	16.2	15.8-16.5	784.2	6.3	6.1-6.5	491.2	2.1	2.0-2.2	338.4	3.3	3.1-3.5	152.8	1.1	1.1-1.2
2016-2020	2,772.8	11.3	11.1-11.5	1,888.6	16.7	16.4-17.1	884.2	6.7	6.5-6.9	541.2	2.1	2.1-2.2	374.0	3.3	3.2-3.5	167.2	1.2	1.1-1.3	
Esophagus	1976-1980	882.2	5.1	5.0-5.3	616.0	8.5	8.2-8.9	266.2	2.7	2.6-2.9	794.0	4.6	4.5-4.8	556.8	7.8	7.5-8.1	237.2	2.4	2.3-2.6
	1981-1985	891.2	5.1	4.9-5.2	615.2	8.4	8.1-8.7	276.0	2.7	2.6-2.9	796.2	4.5	4.4-4.7	567.0	7.8	7.5-8.1	229.2	2.2	2.1-2.4
	1986-1990	907.4	5.1	5.0-5.3	619.4	8.3	8.0-8.6	288.0	2.8	2.6-2.9	807.8	4.6	4.4-4.7	576.4	7.8	7.5-8.1	231.4	2.2	2.1-2.4
	1991-1995	958.6	5.3	5.1-5.4	658.6	8.6	8.3-8.9	300.0	2.8	2.7-3.0	837.6	4.6	4.5-4.7	601.0	7.9	7.6-8.2	236.6	2.2	2.1-2.3
	1996-2000	965.8	5.1	5.0-5.3	677.2	8.5	8.2-8.8	288.6	2.6	2.5-2.7	845.4	4.5	4.3-4.6	605.0	7.6	7.4-7.9	240.4	2.1	2.0-2.3
	2001-2005	1,021.0	5.1	4.9-5.2	740.2	8.6	8.3-8.9	280.8	2.4	2.3-2.5	883.8	4.4	4.3-4.5	654.6	7.7	7.5-8.0	229.2	1.9	1.8-2.0
	2006-2010	1,080.2	5.1	4.9-5.2	811.8	8.8	8.5-9.1	268.4	2.2	2.1-2.3	884.2	4.2	4.0-4.3	669.2	7.3	7.1-7.6	215.0	1.7	1.6-1.9
	2011-2015	1,055.6	4.5	4.4-4.7	794.0	7.7	7.5-8.0	261.6	2.0	1.9-2.1	861.6	3.7	3.6-3.8	656.2	6.5	6.3-6.8	205.4	1.5	1.4-1.6
2016-2020	1,080.0	4.3	4.2-4.4	817.0	7.3	7.0-7.5	263.0	1.9	1.8-2.0	828.8	3.3	3.2-3.4	644.6	5.8	5.6-6.0	184.2	1.3	1.2-1.4	
Stomach	1976-1980	2,103.2	12.9	12.6-13.1	1,206.6	18.2	17.7-18.7	896.6	9.3	9.0-9.6	1,578.6	9.8	9.5-10.0	890.6	13.8	13.4-14.2	688.0	7.1	6.9-7.4
	1981-1985	2,076.4	12.2	11.9-12.4	1,202.2	17.6	17.1-18.1	874.2	8.6	8.3-8.8	1,443.2	8.5	8.3-8.7	834.0	12.5	12.1-12.9	609.2	6.0	5.8-6.2
	1986-1990	1,997.6	11.4	11.2-11.6	1,175.6	16.7	16.3-17.1	822.0	7.8	7.5-8.0	1,344.6	7.7	7.5-7.9	778.8	11.3	10.9-11.7	565.8	5.3	5.1-5.5
	1991-1995	2,026.4	11.2	11.0-11.4	1,182.0	16.2	15.8-16.7	844.4	7.7	7.5-8.0	1,305.6	7.2	7.1-7.4	755.6	10.7	10.4-11.1	550.0	5.0	4.8-5.2
	1996-2000	1,985.8	10.5	10.3-10.7	1,169.4	15.2	14.8-15.6	816.4	7.2	7.0-7.4	1,141.6	6.0	5.9-6.2	636.8	8.4	8.1-8.7	504.8	4.4	4.2-4.6
	2001-2005	1,924.8	9.6	9.4-9.8	1,123.2	13.5	13.1-13.8	801.6	6.8	6.6-7.0	1,015.4	5.0	4.9-5.2	568.4	7.0	6.7-7.2	447.0	3.7	3.6-3.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York State**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	1,848.8	8.8	8.6-9.0	1,113.4	12.4	12.0-12.7	735.4	6.1	5.9-6.3	870.2	4.1	4.0-4.2	500.8	5.7	5.5-5.9	369.4	3.0	2.8-3.1
	2011-2015	1,993.0	8.8	8.6-9.0	1,199.4	12.1	11.8-12.4	793.6	6.3	6.1-6.5	839.2	3.7	3.6-3.8	493.6	5.1	4.9-5.3	345.6	2.6	2.5-2.8
	2016-2020	2,009.2	8.3	8.1-8.5	1,191.4	11.0	10.7-11.2	817.8	6.2	6.0-6.4	787.4	3.2	3.1-3.3	468.8	4.4	4.2-4.6	318.6	2.3	2.2-2.4
Colorectal	1976-1980	11,233.2	67.7	67.1-68.3	5,465.2	81.6	80.6-82.6	5,768.0	58.8	58.1-59.5	5,641.2	34.8	34.3-35.2	2,741.4	42.9	42.1-43.6	2,899.8	29.8	29.3-30.3
	1981-1985	11,802.2	68.7	68.1-69.3	5,787.4	84.6	83.6-85.6	6,014.8	58.6	58.0-59.3	5,471.6	32.4	32.0-32.8	2,655.8	40.4	39.7-41.1	2,815.8	27.4	27.0-27.9
	1986-1990	11,213.8	63.4	62.9-63.9	5,485.6	77.9	76.9-78.8	5,728.2	54.0	53.4-54.7	5,123.4	29.3	28.9-29.7	2,525.6	37.4	36.7-38.1	2,597.8	24.4	23.9-24.8
	1991-1995	10,961.2	60.3	59.8-60.8	5,317.2	73.0	72.1-74.0	5,644.0	51.9	51.3-52.5	4,724.6	26.1	25.8-26.5	2,265.2	32.5	31.9-33.1	2,459.4	22.2	21.8-22.6
	1996-2000	11,518.4	60.8	60.4-61.3	5,614.0	72.7	71.9-73.6	5,904.4	52.6	52.0-53.2	4,334.4	22.9	22.6-23.2	2,104.0	28.2	27.7-28.8	2,230.4	19.3	18.9-19.6
	2001-2005	10,989.8	54.7	54.3-55.2	5,366.4	64.3	63.5-65.0	5,623.4	47.8	47.3-48.4	3,879.0	19.2	18.9-19.5	1,867.6	23.2	22.7-23.7	2,011.4	16.5	16.2-16.8
	2006-2010	9,834.6	46.5	46.1-46.9	4,833.0	53.5	52.8-54.2	5,001.6	41.2	40.7-41.7	3,386.6	15.9	15.6-16.1	1,653.2	19.0	18.6-19.4	1,733.4	13.6	13.3-13.9
	2011-2015	9,033.8	39.8	39.4-40.2	4,576.4	46.0	45.4-46.6	4,457.4	35.0	34.5-35.4	3,132.6	13.5	13.3-13.7	1,560.0	16.1	15.7-16.4	1,572.6	11.5	11.3-11.8
	2016-2020	8,792.8	36.6	36.3-37.0	4,552.8	42.2	41.7-42.8	4,240.0	32.1	31.6-32.5	2,888.6	11.6	11.4-11.8	1,461.0	13.7	13.4-14.0	1,427.6	10.0	9.7-10.2
Colon excluding rectum	1976-1980	7,963.6	48.3	47.9-48.8	3,708.6	56.3	55.5-57.2	4,255.0	43.5	42.9-44.1	4,597.6	28.3	28.0-28.7	2,168.4	34.0	33.3-34.7	2,429.2	24.9	24.5-25.4
	1981-1985	8,457.0	49.5	49.0-50.0	3,961.8	58.8	57.9-59.7	4,495.2	43.8	43.2-44.4	4,633.6	27.4	27.1-27.8	2,212.2	33.7	33.1-34.4	2,421.4	23.6	23.2-24.0
	1986-1990	8,113.4	46.0	45.6-46.5	3,804.2	54.8	53.9-55.6	4,309.2	40.6	40.0-41.1	4,425.0	25.3	25.0-25.6	2,161.6	32.0	31.4-32.7	2,263.4	21.2	20.8-21.6
	1991-1995	8,023.8	44.2	43.7-44.6	3,743.0	52.0	51.2-52.8	4,280.8	39.2	38.6-39.7	4,067.8	22.5	22.2-22.8	1,923.8	27.7	27.1-28.3	2,144.0	19.3	19.0-19.7
	1996-2000	8,429.6	44.5	44.1-44.9	3,937.4	51.6	50.9-52.3	4,492.2	39.8	39.3-40.3	3,725.6	19.7	19.4-19.9	1,782.8	24.0	23.5-24.6	1,942.8	16.8	16.4-17.1
	2001-2005	8,011.0	39.9	39.5-40.3	3,758.0	45.5	44.8-46.2	4,253.0	35.9	35.4-36.4	3,290.4	16.3	16.0-16.5	1,551.8	19.4	18.9-19.8	1,738.6	14.2	13.9-14.5
	2006-2010	7,099.6	33.6	33.2-33.9	3,342.2	37.4	36.9-38.0	3,757.4	30.7	30.3-31.2	2,840.6	13.3	13.1-13.5	1,360.8	15.7	15.4-16.1	1,479.8	11.6	11.3-11.9
	2011-2015	6,411.8	28.2	27.9-28.6	3,093.2	31.5	31.0-32.0	3,318.6	25.8	25.4-26.2	2,580.2	11.1	10.9-11.3	1,247.4	12.9	12.6-13.2	1,332.8	9.7	9.5-10.0
	2016-2020	6,212.6	25.8	25.5-26.0	3,061.0	28.5	28.0-29.0	3,151.6	23.5	23.1-23.9	2,287.6	9.2	9.0-9.3	1,119.8	10.5	10.3-10.8	1,167.8	8.1	7.9-8.3
Rectum & rectosigmoid	1976-1980	3,269.6	19.3	19.0-19.6	1,756.6	25.2	24.7-25.8	1,513.0	15.4	15.0-15.7	1,043.6	6.4	6.2-6.6	573.0	8.9	8.5-9.2	470.6	4.9	4.7-5.1
	1981-1985	3,345.2	19.2	18.9-19.5	1,825.6	25.8	25.2-26.3	1,519.6	14.8	14.5-15.2	838.0	5.0	4.8-5.1	443.6	6.7	6.4-7.0	394.4	3.8	3.7-4.0
	1986-1990	3,100.4	17.4	17.1-17.7	1,681.4	23.1	22.6-23.6	1,419.0	13.5	13.2-13.8	698.4	4.0	3.9-4.1	364.0	5.4	5.1-5.6	334.4	3.1	3.0-3.3
	1991-1995	2,937.4	16.1	15.8-16.4	1,574.2	21.1	20.6-21.5	1,363.2	12.7	12.4-13.0	656.8	3.6	3.5-3.8	341.4	4.8	4.6-5.0	315.4	2.9	2.7-3.0
	1996-2000	3,088.8	16.3	16.1-16.6	1,676.6	21.2	20.7-21.6	1,412.2	12.8	12.5-13.1	608.8	3.2	3.1-3.3	321.2	4.2	4.0-4.4	287.6	2.5	2.4-2.6
	2001-2005	2,978.8	14.8	14.6-15.1	1,608.4	18.8	18.4-19.2	1,370.4	11.9	11.6-12.2	588.6	2.9	2.8-3.0	315.8	3.8	3.6-4.0	272.8	2.3	2.1-2.4
	2006-2010	2,735.0	12.9	12.7-13.1	1,490.8	16.1	15.7-16.4	1,244.2	10.5	10.2-10.8	546.0	2.6	2.5-2.7	292.4	3.3	3.1-3.5	253.6	2.0	1.9-2.1
	2011-2015	2,622.0	11.6	11.4-11.8	1,483.2	14.5	14.2-14.9	1,138.8	9.2	8.9-9.4	552.4	2.4	2.3-2.5	312.6	3.2	3.0-3.3	239.8	1.8	1.7-1.9
	2016-2020	2,580.2	10.9	10.7-11.1	1,491.8	13.7	13.4-14.0	1,088.4	8.5	8.3-8.8	601.0	2.4	2.4-2.5	341.2	3.2	3.0-3.3	259.8	1.9	1.8-2.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York State**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	571.6	3.4	3.3-3.5	346.0	4.9	4.7-5.1	225.6	2.3	2.2-2.4	559.0	3.3	3.2-3.4	320.4	4.6	4.3-4.8	238.6	2.5	2.3-2.6
	1981-1985	596.4	3.5	3.3-3.6	376.6	5.3	5.1-5.6	219.8	2.2	2.0-2.3	622.2	3.6	3.5-3.8	371.6	5.3	5.1-5.6	250.6	2.5	2.3-2.6
	1986-1990	688.8	3.9	3.8-4.0	439.2	6.0	5.8-6.3	249.6	2.4	2.3-2.5	702.2	4.0	3.9-4.1	420.6	5.9	5.6-6.2	281.6	2.7	2.5-2.8
	1991-1995	892.2	4.9	4.8-5.1	568.0	7.5	7.2-7.8	324.2	3.0	2.9-3.2	833.8	4.6	4.4-4.7	506.0	6.8	6.5-7.1	327.8	3.0	2.9-3.2
	1996-2000	1,140.2	6.1	5.9-6.2	757.0	9.4	9.1-9.7	383.2	3.5	3.3-3.6	883.6	4.7	4.5-4.8	551.6	6.9	6.7-7.2	332.0	2.9	2.8-3.1
	2001-2005	1,377.4	6.9	6.7-7.0	974.4	11.0	10.7-11.4	403.0	3.5	3.4-3.7	1,016.0	5.1	4.9-5.2	659.0	7.7	7.4-7.9	357.0	3.0	2.9-3.2
	2006-2010	1,747.0	8.1	8.0-8.3	1,262.4	13.0	12.7-13.4	484.6	4.1	3.9-4.2	1,209.6	5.7	5.5-5.8	807.4	8.6	8.4-8.9	402.2	3.3	3.1-3.4
	2011-2015	2,106.8	8.9	8.7-9.1	1,495.4	13.9	13.5-14.2	611.4	4.8	4.6-4.9	1,412.4	6.0	5.9-6.2	938.0	9.0	8.7-9.3	474.4	3.6	3.5-3.8
	2016-2020	2,204.0	8.7	8.5-8.8	1,547.2	13.4	13.1-13.7	656.8	4.7	4.6-4.9	1,413.8	5.6	5.4-5.7	941.0	8.3	8.1-8.6	472.8	3.3	3.2-3.5
Pancreas	1976-1980	2,161.8	13.0	12.7-13.2	1,095.0	16.0	15.6-16.5	1,066.8	10.8	10.5-11.1	2,030.4	12.2	11.9-12.4	1,025.0	15.1	14.7-15.5	1,005.4	10.2	9.9-10.5
	1981-1985	2,164.2	12.6	12.3-12.8	1,040.0	15.0	14.6-15.4	1,124.2	10.9	10.6-11.2	1,996.0	11.6	11.4-11.8	969.4	14.0	13.6-14.4	1,026.6	9.9	9.7-10.2
	1986-1990	2,193.6	12.4	12.2-12.7	1,022.0	14.5	14.1-14.9	1,171.6	11.0	10.7-11.3	1,989.6	11.3	11.0-11.5	940.6	13.4	13.0-13.8	1,049.0	9.8	9.5-10.1
	1991-1995	2,286.4	12.6	12.4-12.8	1,060.8	14.5	14.1-14.9	1,225.6	11.2	10.9-11.5	2,050.6	11.3	11.1-11.5	948.2	13.0	12.6-13.4	1,102.4	10.0	9.7-10.3
	1996-2000	2,406.0	12.7	12.5-12.9	1,123.2	14.5	14.1-14.9	1,282.8	11.4	11.1-11.7	2,123.4	11.2	11.0-11.4	994.2	13.0	12.7-13.4	1,129.2	9.9	9.7-10.2
	2001-2005	2,571.0	12.8	12.6-13.0	1,200.0	14.3	14.0-14.7	1,371.0	11.6	11.3-11.8	2,210.4	11.0	10.8-11.2	1,030.0	12.5	12.1-12.8	1,180.4	9.8	9.6-10.1
	2006-2010	2,886.2	13.6	13.4-13.8	1,397.0	15.5	15.1-15.9	1,489.2	12.1	11.8-12.3	2,356.6	11.0	10.8-11.2	1,132.4	12.7	12.3-13.0	1,224.2	9.8	9.5-10.0
	2011-2015	3,233.2	14.0	13.8-14.2	1,598.4	15.9	15.6-16.3	1,634.8	12.4	12.2-12.7	2,540.4	11.0	10.8-11.2	1,250.2	12.7	12.3-13.0	1,290.2	9.6	9.4-9.9
	2016-2020	3,603.2	14.4	14.2-14.6	1,809.4	16.4	16.1-16.8	1,793.8	12.8	12.5-13.1	2,716.6	10.8	10.6-11.0	1,350.4	12.3	12.0-12.6	1,366.2	9.5	9.3-9.7
Larynx	1976-1980	1,026.4	5.8	5.6-6.0	840.8	11.2	10.8-11.5	185.6	1.9	1.8-2.0	385.8	2.2	2.1-2.3	314.2	4.4	4.1-4.6	71.6	0.7	0.7-0.8
	1981-1985	1,077.6	6.1	5.9-6.2	857.4	11.3	10.9-11.6	220.2	2.2	2.1-2.4	364.6	2.0	1.9-2.1	288.6	3.9	3.7-4.1	76.0	0.7	0.7-0.8
	1986-1990	1,089.4	6.1	5.9-6.3	834.6	10.9	10.6-11.3	254.8	2.6	2.4-2.7	345.4	1.9	1.9-2.0	270.0	3.7	3.5-3.9	75.4	0.7	0.7-0.8
	1991-1995	993.8	5.5	5.3-5.6	764.4	9.8	9.5-10.1	229.4	2.3	2.1-2.4	360.0	2.0	1.9-2.1	276.6	3.7	3.5-3.9	83.4	0.8	0.7-0.9
	1996-2000	899.0	4.8	4.6-4.9	696.8	8.5	8.3-8.8	202.2	1.9	1.8-2.1	310.0	1.6	1.6-1.7	242.0	3.0	2.9-3.2	68.0	0.6	0.6-0.7
	2001-2005	840.6	4.2	4.1-4.3	660.2	7.5	7.3-7.8	180.4	1.6	1.5-1.7	270.4	1.3	1.3-1.4	213.2	2.5	2.4-2.7	57.2	0.5	0.4-0.6
	2006-2010	820.8	3.8	3.7-4.0	638.8	6.8	6.6-7.0	182.0	1.6	1.4-1.7	253.2	1.2	1.1-1.3	197.0	2.2	2.0-2.3	56.2	0.5	0.4-0.5
	2011-2015	786.2	3.3	3.2-3.5	626.0	6.0	5.8-6.2	160.2	1.3	1.2-1.3	243.0	1.0	1.0-1.1	191.2	1.9	1.8-2.0	51.8	0.4	0.3-0.4
	2016-2020	705.8	2.8	2.7-2.9	557.0	4.9	4.7-5.1	148.8	1.1	1.0-1.2	198.4	0.8	0.7-0.8	157.6	1.4	1.3-1.5	40.8	0.3	0.2-0.3
Lung and bronchus	1976-1980	9,869.2	56.7	56.2-57.2	6,949.2	96.5	95.4-97.5	2,920.0	29.6	29.1-30.1	8,038.4	46.4	45.9-46.9	5,764.2	81.1	80.1-82.0	2,274.2	23.0	22.5-23.4
	1981-1985	10,950.4	61.8	61.3-62.4	7,186.4	98.7	97.6-99.8	3,764.0	37.2	36.7-37.7	8,823.2	49.9	49.5-50.4	5,894.6	82.1	81.2-83.1	2,928.6	28.7	28.3-29.2
	1986-1990	11,752.6	65.4	64.8-65.9	7,211.0	97.6	96.5-98.6	4,541.6	44.0	43.4-44.5	9,513.8	53.0	52.5-53.4	5,959.0	81.8	80.8-82.7	3,554.8	34.2	33.7-34.7
	1991-1995	12,405.6	67.8	67.3-68.4	7,110.4	94.0	93.0-95.0	5,295.2	50.3	49.7-50.9	9,959.2	54.5	54.0-54.9	5,862.8	78.8	77.9-79.8	4,096.4	38.5	37.9-39.0
	1996-2000	12,782.4	67.6	67.1-68.1	6,966.4	88.1	87.2-89.0	5,816.0	53.6	53.0-54.2	9,618.4	50.8	50.4-51.3	5,397.4	69.2	68.4-70.0	4,221.0	38.3	37.8-38.9
	2001-2005	13,030.8	65.2	64.7-65.7	6,841.2	81.3	80.4-82.2	6,189.6	54.1	53.5-54.8	9,446.4	47.2	46.8-47.6	5,083.4	61.3	60.6-62.1	4,363.0	37.6	37.1-38.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York State**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	13,569.2	64.6	64.1-65.0	6,880.6	76.7	75.9-77.5	6,688.6	56.2	55.6-56.8	9,127.6	43.3	42.9-43.7	4,795.6	54.1	53.4-54.8	4,332.0	35.8	35.3-36.3
	2011-2015	13,820.0	60.3	59.9-60.8	6,860.8	69.1	68.4-69.9	6,959.2	54.2	53.6-54.8	8,760.4	38.2	37.8-38.5	4,564.0	46.6	46.0-47.2	4,196.4	32.2	31.7-32.6
	2016-2020	13,944.2	55.4	55.0-55.8	6,723.2	60.8	60.1-61.4	7,221.0	51.7	51.1-52.2	7,448.2	29.5	29.2-29.8	3,813.4	35.0	34.5-35.5	3,634.8	25.5	25.1-25.8
Melanoma of the skin	1976-1980	1,099.0	6.6	6.4-6.8	563.8	7.7	7.4-8.0	535.2	5.9	5.6-6.1	382.4	2.3	2.2-2.4	220.8	3.1	2.9-3.3	161.6	1.7	1.6-1.9
	1981-1985	1,326.6	7.8	7.6-8.0	692.2	9.3	9.0-9.6	634.4	6.8	6.6-7.1	409.2	2.4	2.3-2.5	238.4	3.3	3.1-3.5	170.8	1.8	1.7-1.9
	1986-1990	1,376.8	7.8	7.7-8.0	743.4	9.8	9.5-10.1	633.4	6.5	6.3-6.8	438.4	2.5	2.4-2.6	263.8	3.6	3.4-3.8	174.6	1.8	1.7-1.9
	1991-1995	1,429.2	7.8	7.7-8.0	816.6	10.4	10.1-10.7	612.6	6.0	5.8-6.3	415.6	2.3	2.2-2.4	247.8	3.3	3.1-3.5	167.8	1.6	1.5-1.7
	1996-2000	1,927.8	10.2	10.0-10.4	1,083.2	13.3	12.9-13.7	844.6	8.1	7.8-8.3	429.4	2.3	2.2-2.4	261.6	3.3	3.1-3.5	167.8	1.5	1.4-1.7
	2001-2005	2,738.4	13.8	13.6-14.0	1,532.6	17.7	17.3-18.1	1,205.8	11.1	10.9-11.4	442.8	2.2	2.1-2.3	257.4	3.1	2.9-3.2	185.4	1.6	1.5-1.7
	2006-2010	3,575.4	17.3	17.0-17.5	2,023.6	22.2	21.8-22.6	1,551.8	13.9	13.6-14.2	461.2	2.2	2.1-2.3	290.0	3.2	3.1-3.4	171.2	1.4	1.3-1.5
	2011-2015	4,108.2	18.4	18.2-18.7	2,346.2	23.7	23.2-24.1	1,762.0	14.8	14.5-15.2	467.4	2.0	2.0-2.1	295.0	3.0	2.9-3.2	172.4	1.3	1.2-1.4
2016-2020	4,216.0	17.7	17.5-18.0	2,430.0	22.6	22.2-23.0	1,786.0	14.3	14.0-14.6	387.8	1.6	1.5-1.6	243.2	2.3	2.2-2.4	144.6	1.0	0.9-1.1	
Female breast	1976-1980							9,417.0	101.3	100.4-102.2							3,643.2	38.2	37.7-38.8
	1981-1985							10,306.2	108.4	107.4-109.4							3,656.6	37.3	36.7-37.8
	1986-1990							11,649.4	119.4	118.4-120.4							3,760.6	37.4	36.8-37.9
	1991-1995							12,603.8	126.0	125.0-127.0							3,579.2	34.6	34.1-35.1
	1996-2000							14,118.4	135.4	134.4-136.4							3,258.6	30.0	29.5-30.4
	2001-2005							13,929.4	126.4	125.5-127.4							2,961.8	25.6	25.2-26.0
	2006-2010							14,689.6	128.4	127.5-129.4							2,678.0	22.1	21.7-22.5
	2011-2015							15,852.0	131.9	131.0-132.8							2,553.8	19.8	19.4-20.1
2016-2020							16,688.2	134.0	133.1-135.0							2,486.0	18.2	17.9-18.6	
Cervix uteri	1976-1980							1,223.8	13.7	13.3-14.0							457.6	4.9	4.7-5.2
	1981-1985							1,074.6	11.8	11.5-12.1							376.4	4.0	3.8-4.2
	1986-1990							1,089.2	11.6	11.3-12.0							338.8	3.5	3.3-3.7
	1991-1995							1,152.4	11.8	11.5-12.1							321.8	3.3	3.1-3.4
	1996-2000							1,095.4	10.8	10.5-11.1							322.0	3.1	3.0-3.3
	2001-2005							932.2	8.9	8.6-9.1							284.6	2.6	2.5-2.7
	2006-2010							896.8	8.4	8.2-8.7							264.2	2.3	2.2-2.4
	2011-2015							846.8	7.7	7.5-8.0							276.8	2.3	2.2-2.5
2016-2020							808.6	7.4	7.1-7.6							233.2	1.9	1.8-2.0	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York State**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							2,327.2	23.7	23.3-24.2							554.4	5.6	5.3-5.8
	1981-1985							2,318.2	23.2	22.8-23.6							522.0	5.0	4.8-5.2
	1986-1990							2,233.0	22.0	21.6-22.5							483.6	4.5	4.4-4.7
	1991-1995							2,488.0	24.5	24.0-24.9							529.0	4.9	4.7-5.1
	1996-2000							2,831.2	26.9	26.5-27.4							536.8	4.8	4.6-5.0
	2001-2005							3,182.8	28.6	28.2-29.1							566.8	4.9	4.7-5.1
	2006-2010							3,505.2	29.8	29.4-30.3							626.2	5.1	5.0-5.3
	2011-2015							3,995.6	31.5	31.1-32.0							733.2	5.6	5.4-5.8
2016-2020							4,168.8	31.2	30.7-31.6							842.6	6.0	5.8-6.2	
Ovary	1976-1980							1,534.0	16.1	15.7-16.4							1,035.4	10.6	10.3-10.9
	1981-1985							1,631.8	16.9	16.5-17.3							1,015.6	10.1	9.8-10.4
	1986-1990							1,617.0	16.4	16.0-16.8							1,019.8	9.9	9.6-10.2
	1991-1995							1,738.0	17.2	16.8-17.6							1,064.2	10.1	9.9-10.4
	1996-2000							1,666.4	15.9	15.5-16.2							998.4	9.2	8.9-9.4
	2001-2005							1,613.6	14.5	14.2-14.9							1,021.6	8.8	8.6-9.1
	2006-2010							1,543.0	13.3	13.0-13.6							967.2	8.0	7.8-8.2
	2011-2015							1,526.0	12.5	12.2-12.8							935.8	7.2	7.0-7.4
2016-2020							1,409.6	11.2	10.9-11.5							866.0	6.2	6.0-6.4	
Prostate	1976-1980				4,995.0	80.7	79.7-81.8							1,777.0	31.5	30.8-32.2			
	1981-1985				5,832.6	89.8	88.7-90.9							1,933.4	33.0	32.3-33.7			
	1986-1990				7,027.4	103.3	102.1-104.4							2,131.6	34.8	34.1-35.4			
	1991-1995				11,630.8	155.3	154.0-156.6							2,360.8	37.2	36.5-37.9			
	1996-2000				13,561.6	170.0	168.7-171.3							2,151.0	31.5	30.9-32.1			
	2001-2005				14,778.6	171.9	170.7-173.2							1,940.4	26.2	25.7-26.7			
	2006-2010				16,031.0	169.8	168.6-171.0							1,755.0	21.6	21.1-22.1			
	2011-2015				14,237.0	132.9	131.9-133.9							1,667.8	18.3	17.9-18.7			
2016-2020				15,455.6	130.3	129.4-131.2							1,706.4	16.6	16.2-16.9				
Testis	1976-1980				290.4	3.4	3.2-3.6							49.2	0.6	0.5-0.7			
	1981-1985				362.2	4.1	3.9-4.3							33.6	0.4	0.3-0.5			
	1986-1990				396.0	4.2	4.0-4.4							30.8	0.3	0.3-0.4			
	1991-1995				434.8	4.6	4.4-4.8							26.8	0.3	0.3-0.4			
	1996-2000				468.0	5.0	4.8-5.2							20.6	0.2	0.2-0.3			
	2001-2005				500.4	5.3	5.1-5.5							22.6	0.2	0.2-0.3			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York State**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				546.8	5.9	5.6-6.1							19.6	0.2	0.2-0.3			
	2011-2015				564.4	6.0	5.7-6.2							18.8	0.2	0.2-0.2			
	2016-2020				579.2	6.2	6.0-6.4							22.0	0.2	0.2-0.3			
Urinary bladder (incl. in situ)	1976-1980	3,539.6	21.2	20.9-21.6	2,570.2	38.6	37.9-39.3	969.4	9.9	9.6-10.2	1,013.8	6.4	6.3-6.6	696.4	11.6	11.2-12.0	317.4	3.3	3.2-3.5
	1981-1985	3,757.6	21.7	21.4-22.0	2,710.2	39.5	38.8-40.2	1,047.4	10.2	9.9-10.5	950.8	5.7	5.6-5.9	649.8	10.6	10.2-11.0	301.0	2.9	2.8-3.1
	1986-1990	3,797.8	21.3	21.0-21.6	2,718.2	38.5	37.8-39.2	1,079.6	10.2	9.9-10.5	897.6	5.2	5.0-5.4	594.6	9.5	9.1-9.8	303.0	2.8	2.7-3.0
	1991-1995	3,968.6	21.7	21.4-22.0	2,803.8	38.5	37.8-39.2	1,164.8	10.8	10.5-11.1	902.4	5.0	4.9-5.2	587.8	8.9	8.6-9.2	314.6	2.8	2.7-2.9
	1996-2000	4,499.6	23.8	23.5-24.1	3,220.8	42.0	41.3-42.7	1,278.8	11.4	11.2-11.7	928.2	4.9	4.8-5.0	616.2	8.7	8.4-9.0	312.0	2.6	2.5-2.8
	2001-2005	4,870.0	24.3	24.0-24.6	3,544.8	43.2	42.5-43.8	1,325.2	11.3	11.0-11.6	901.4	4.5	4.3-4.6	601.0	7.8	7.5-8.1	300.4	2.4	2.3-2.5
	2006-2010	5,056.2	24.0	23.7-24.3	3,752.0	42.7	42.1-43.4	1,304.2	10.7	10.4-11.0	959.4	4.5	4.3-4.6	660.2	7.9	7.6-8.2	299.2	2.3	2.2-2.4
	2011-2015	5,373.0	23.5	23.2-23.8	3,982.2	41.2	40.6-41.8	1,390.8	10.6	10.4-10.9	993.4	4.2	4.1-4.4	680.6	7.3	7.1-7.6	312.8	2.2	2.1-2.3
2016-2020	5,362.4	21.4	21.2-21.7	4,001.0	37.1	36.6-37.7	1,361.4	9.6	9.4-9.9	958.2	3.8	3.6-3.9	672.2	6.5	6.3-6.8	286.0	1.9	1.8-2.0	
Kidney and renal pelvis	1976-1980	1,248.6	7.3	7.1-7.5	773.4	10.7	10.3-11.0	475.2	4.9	4.7-5.1	655.6	3.9	3.7-4.0	411.6	5.8	5.6-6.1	244.0	2.5	2.3-2.6
	1981-1985	1,474.6	8.5	8.3-8.7	897.6	12.3	11.9-12.7	577.0	5.8	5.6-6.0	653.8	3.8	3.6-3.9	390.2	5.5	5.3-5.8	263.6	2.6	2.5-2.7
	1986-1990	1,685.0	9.5	9.3-9.7	1,018.0	13.7	13.3-14.1	667.0	6.6	6.3-6.8	700.4	4.0	3.8-4.1	416.0	5.8	5.5-6.1	284.4	2.7	2.6-2.9
	1991-1995	1,952.8	10.7	10.5-11.0	1,172.8	15.2	14.8-15.6	780.0	7.5	7.3-7.8	692.0	3.8	3.7-3.9	417.0	5.7	5.4-5.9	275.0	2.5	2.4-2.7
	1996-2000	2,366.4	12.6	12.3-12.8	1,432.8	17.7	17.3-18.1	933.6	8.7	8.5-9.0	687.4	3.6	3.5-3.8	415.2	5.4	5.1-5.6	272.2	2.4	2.3-2.6
	2001-2005	2,848.8	14.3	14.0-14.5	1,758.0	20.1	19.7-20.6	1,090.8	9.7	9.5-10.0	702.2	3.5	3.4-3.6	425.6	5.1	4.9-5.3	276.6	2.3	2.2-2.5
	2006-2010	3,407.6	16.2	16.0-16.4	2,138.8	22.8	22.4-23.3	1,268.8	10.9	10.7-11.2	713.4	3.3	3.2-3.5	449.8	5.0	4.8-5.3	263.6	2.1	2.0-2.2
	2011-2015	3,760.8	16.6	16.4-16.9	2,414.0	23.5	23.1-23.9	1,346.8	10.9	10.7-11.2	714.4	3.1	3.0-3.2	455.4	4.6	4.4-4.8	259.0	1.9	1.8-2.0
2016-2020	4,044.2	16.8	16.6-17.1	2,618.2	23.7	23.3-24.2	1,426.0	11.0	10.8-11.3	655.8	2.6	2.5-2.7	420.8	3.9	3.7-4.1	235.0	1.6	1.5-1.7	
Brain and other central nervous system	1976-1980	1,014.0	5.7	5.6-5.9	556.8	7.0	6.7-7.3	457.2	4.8	4.6-5.0	734.0	4.1	4.0-4.3	406.0	5.2	5.0-5.4	328.0	3.3	3.2-3.5
	1981-1985	1,092.0	6.2	6.0-6.4	577.6	7.4	7.1-7.6	514.4	5.3	5.1-5.5	732.0	4.1	4.0-4.3	384.2	5.0	4.8-5.3	347.8	3.5	3.3-3.6
	1986-1990	1,209.2	6.8	6.6-7.0	653.2	8.2	7.9-8.5	556.0	5.7	5.4-5.9	727.8	4.1	3.9-4.2	397.4	5.1	4.9-5.4	330.4	3.2	3.1-3.4
	1991-1995	1,316.8	7.2	7.0-7.4	693.8	8.5	8.2-8.8	623.0	6.2	6.0-6.4	736.8	4.1	3.9-4.2	393.8	5.0	4.8-5.2	343.0	3.3	3.2-3.5
	1996-2000	1,391.2	7.4	7.2-7.6	736.0	8.7	8.4-8.9	655.2	6.3	6.1-6.6	752.2	4.0	3.9-4.1	399.8	4.8	4.6-5.0	352.4	3.3	3.2-3.5
	2001-2005	1,363.4	7.0	6.8-7.1	734.6	8.3	8.0-8.5	628.8	5.9	5.7-6.1	740.0	3.7	3.6-3.9	404.4	4.6	4.4-4.8	335.6	3.0	2.9-3.2
	2006-2010	1,355.4	6.7	6.6-6.9	735.4	8.0	7.7-8.2	620.0	5.7	5.5-5.9	766.6	3.7	3.6-3.8	431.6	4.7	4.5-4.9	335.0	2.9	2.8-3.0
	2011-2015	1,415.4	6.7	6.5-6.8	772.8	7.8	7.6-8.1	642.6	5.7	5.5-5.9	831.2	3.7	3.6-3.8	444.0	4.4	4.2-4.6	387.2	3.1	3.0-3.3
2016-2020	1,416.8	6.4	6.3-6.6	790.8	7.7	7.4-7.9	626.0	5.4	5.2-5.6	900.4	3.8	3.7-3.9	510.0	4.7	4.5-4.9	390.4	3.0	2.9-3.2	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York State**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	598.8	3.6	3.4-3.7	176.4	2.4	2.2-2.5	422.4	4.6	4.4-4.8	109.2	0.7	0.6-0.7	41.2	0.6	0.5-0.7	68.0	0.7	0.6-0.8
	1981-1985	662.0	3.9	3.7-4.0	185.0	2.4	2.2-2.6	477.0	5.1	4.9-5.4	101.8	0.6	0.6-0.7	38.4	0.6	0.5-0.7	63.4	0.6	0.6-0.7
	1986-1990	709.4	4.0	3.9-4.2	198.4	2.5	2.4-2.7	511.0	5.4	5.2-5.6	92.0	0.5	0.5-0.6	30.6	0.4	0.4-0.5	61.4	0.6	0.5-0.6
	1991-1995	955.4	5.2	5.1-5.4	250.2	3.0	2.9-3.2	705.2	7.2	7.0-7.5	93.0	0.5	0.5-0.6	32.4	0.4	0.4-0.5	60.6	0.6	0.5-0.6
	1996-2000	1,418.0	7.5	7.3-7.7	353.2	4.1	3.9-4.3	1,064.8	10.7	10.4-11.0	102.4	0.5	0.5-0.6	35.8	0.5	0.4-0.5	66.6	0.6	0.5-0.7
	2001-2005	2,075.2	10.6	10.3-10.8	503.2	5.5	5.3-5.7	1,572.0	15.3	14.9-15.6	95.0	0.5	0.4-0.5	36.6	0.4	0.4-0.5	58.4	0.5	0.4-0.5
	2006-2010	3,288.4	16.3	16.1-16.6	786.8	8.2	8.0-8.5	2,501.6	23.9	23.4-24.3	112.4	0.5	0.5-0.6	46.6	0.5	0.5-0.6	65.8	0.5	0.5-0.6
	2011-2015	4,170.8	20.1	19.8-20.3	1,036.6	10.3	10.0-10.6	3,134.2	29.3	28.8-29.7	121.6	0.5	0.5-0.6	51.2	0.5	0.5-0.6	70.4	0.5	0.5-0.6
2016-2020	3,820.0	18.3	18.1-18.6	1,021.4	10.0	9.7-10.2	2,798.6	26.3	25.9-26.8	124.8	0.5	0.5-0.5	52.6	0.5	0.4-0.6	72.2	0.5	0.4-0.6	
Hodgkin lymphoma	1976-1980	624.4	3.4	3.3-3.5	340.0	4.1	3.9-4.3	284.4	2.9	2.7-3.0	221.4	1.3	1.2-1.4	125.2	1.6	1.5-1.8	96.2	1.0	0.9-1.1
	1981-1985	643.4	3.5	3.4-3.6	352.4	4.2	4.0-4.4	291.0	2.9	2.7-3.1	186.4	1.0	1.0-1.1	106.4	1.3	1.2-1.4	80.0	0.8	0.7-0.9
	1986-1990	627.0	3.3	3.2-3.5	347.4	4.0	3.8-4.2	279.6	2.8	2.6-2.9	167.0	0.9	0.9-1.0	104.2	1.3	1.2-1.4	62.8	0.6	0.6-0.7
	1991-1995	634.4	3.4	3.3-3.5	339.8	3.9	3.7-4.0	294.6	3.0	2.8-3.1	136.4	0.7	0.7-0.8	72.8	0.9	0.8-1.0	63.6	0.6	0.6-0.7
	1996-2000	610.8	3.2	3.1-3.3	327.4	3.6	3.4-3.8	283.4	2.9	2.7-3.0	102.8	0.5	0.5-0.6	52.8	0.6	0.5-0.7	50.0	0.5	0.4-0.5
	2001-2005	662.0	3.4	3.3-3.5	357.2	3.9	3.7-4.1	304.8	3.0	2.9-3.2	92.2	0.5	0.4-0.5	49.2	0.6	0.5-0.6	43.0	0.4	0.3-0.5
	2006-2010	656.6	3.3	3.2-3.4	357.4	3.8	3.6-4.0	299.2	2.9	2.8-3.1	85.4	0.4	0.4-0.5	47.0	0.5	0.4-0.6	38.4	0.3	0.3-0.4
	2011-2015	660.4	3.3	3.2-3.4	357.6	3.7	3.5-3.9	302.8	2.9	2.8-3.1	72.0	0.3	0.3-0.4	37.2	0.4	0.3-0.5	34.8	0.3	0.2-0.3
2016-2020	620.2	3.0	2.9-3.1	335.4	3.4	3.2-3.6	284.8	2.7	2.6-2.9	62.2	0.3	0.2-0.3	37.6	0.4	0.3-0.4	24.6	0.2	0.2-0.2	
Non-Hodgkin lymphomas	1976-1980	2,019.6	11.9	11.7-12.1	1,025.4	14.1	13.7-14.5	994.2	10.2	9.9-10.5	1,063.4	6.3	6.1-6.5	547.0	7.7	7.4-8.1	516.4	5.3	5.1-5.5
	1981-1985	2,437.4	14.1	13.8-14.3	1,238.6	17.0	16.5-17.4	1,198.8	11.9	11.6-12.2	1,188.4	6.9	6.7-7.1	610.0	8.6	8.3-9.0	578.4	5.7	5.5-5.9
	1986-1990	2,976.2	16.8	16.6-17.1	1,612.2	21.3	20.8-21.7	1,364.0	13.3	13.0-13.6	1,368.8	7.8	7.6-8.0	718.0	10.0	9.6-10.3	650.8	6.2	6.0-6.4
	1991-1995	3,547.6	19.5	19.2-19.8	1,950.0	24.8	24.3-25.3	1,597.6	15.3	14.9-15.6	1,501.8	8.3	8.1-8.5	777.0	10.5	10.2-10.8	724.8	6.7	6.5-6.9
	1996-2000	3,892.2	20.6	20.3-20.9	2,027.6	24.8	24.4-25.3	1,864.6	17.3	16.9-17.6	1,554.4	8.2	8.0-8.4	778.8	10.1	9.8-10.4	775.6	6.9	6.7-7.1
	2001-2005	4,144.2	20.8	20.5-21.1	2,170.0	25.1	24.6-25.6	1,974.2	17.5	17.1-17.8	1,349.6	6.7	6.6-6.9	706.6	8.6	8.3-8.9	643.0	5.4	5.2-5.6
	2006-2010	4,599.6	22.0	21.7-22.3	2,438.0	26.7	26.2-27.2	2,161.6	18.4	18.1-18.8	1,283.4	6.1	5.9-6.2	679.4	7.8	7.5-8.0	604.0	4.9	4.7-5.0
	2011-2015	4,884.6	21.8	21.6-22.1	2,645.4	26.8	26.3-27.2	2,239.2	18.0	17.6-18.3	1,220.6	5.4	5.2-5.5	665.0	7.0	6.8-7.3	555.6	4.1	4.0-4.3
2016-2020	5,006.0	21.0	20.7-21.3	2,681.6	25.1	24.7-25.5	2,324.4	17.8	17.4-18.1	1,171.6	4.7	4.6-4.8	650.4	6.2	6.0-6.4	521.2	3.6	3.5-3.8	
Myeloma	1976-1980	788.6	4.7	4.5-4.9	383.0	5.6	5.4-5.9	405.6	4.1	3.9-4.3	507.2	3.0	2.9-3.2	245.8	3.6	3.4-3.9	261.4	2.6	2.5-2.8
	1981-1985	899.4	5.2	5.1-5.4	445.2	6.5	6.2-6.7	454.2	4.4	4.3-4.6	576.6	3.3	3.2-3.5	278.2	4.1	3.9-4.3	298.4	2.9	2.7-3.0
	1986-1990	943.4	5.4	5.2-5.5	469.8	6.6	6.4-6.9	473.6	4.5	4.3-4.7	596.4	3.4	3.3-3.5	300.4	4.4	4.1-4.6	296.0	2.8	2.6-2.9
	1991-1995	1,044.8	5.7	5.6-5.9	511.0	7.0	6.7-7.2	533.8	5.0	4.8-5.2	676.6	3.7	3.6-3.8	323.2	4.5	4.3-4.7	353.4	3.2	3.1-3.4
	1996-2000	1,193.4	6.3	6.2-6.5	582.6	7.5	7.2-7.8	610.8	5.5	5.4-5.7	669.4	3.5	3.4-3.7	323.6	4.3	4.1-4.5	345.8	3.1	2.9-3.2
2001-2005	1,353.0	6.7	6.6-6.9	697.6	8.3	8.0-8.6	655.4	5.7	5.5-5.9	654.2	3.3	3.1-3.4	332.0	4.1	3.9-4.3	322.2	2.7	2.6-2.8	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York State**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	1,597.6	7.6	7.4-7.7	843.8	9.3	9.0-9.6	753.8	6.3	6.1-6.5	658.2	3.1	3.0-3.2	345.8	4.0	3.8-4.2	312.4	2.5	2.4-2.6
	2011-2015	1,915.8	8.4	8.2-8.5	1,038.0	10.3	10.1-10.6	877.8	6.9	6.7-7.1	690.0	3.0	2.9-3.1	359.4	3.7	3.5-3.9	330.6	2.5	2.4-2.6
	2016-2020	1,984.6	8.0	7.9-8.2	1,078.6	9.8	9.5-10.1	906.0	6.7	6.5-6.9	672.4	2.7	2.6-2.7	352.8	3.3	3.1-3.4	319.6	2.2	2.1-2.3
Leukemias	1976-1980	2,050.4	12.3	12.1-12.6	1,126.2	16.2	15.8-16.7	924.2	9.7	9.4-10.0	1,351.4	8.2	8.0-8.4	731.4	10.8	10.4-11.2	620.0	6.5	6.3-6.8
	1981-1985	2,112.0	12.4	12.1-12.6	1,165.2	16.5	16.1-16.9	946.8	9.6	9.3-9.9	1,360.0	8.0	7.8-8.2	729.6	10.7	10.3-11.0	630.4	6.4	6.1-6.6
	1986-1990	2,143.0	12.2	11.9-12.4	1,151.6	15.8	15.4-16.3	991.4	9.7	9.5-10.0	1,314.8	7.5	7.3-7.7	704.6	10.0	9.6-10.3	610.2	5.9	5.7-6.1
	1991-1995	2,326.2	12.8	12.6-13.1	1,261.0	16.7	16.3-17.1	1,065.2	10.2	9.9-10.5	1,337.4	7.4	7.2-7.6	723.2	10.0	9.7-10.4	614.2	5.7	5.5-5.9
	1996-2000	2,573.8	13.6	13.4-13.9	1,415.4	17.8	17.4-18.2	1,158.4	10.7	10.4-10.9	1,385.0	7.3	7.1-7.5	747.0	9.7	9.4-10.1	638.0	5.7	5.5-5.9
	2001-2005	2,832.4	14.3	14.0-14.5	1,563.0	18.5	18.1-18.9	1,269.4	11.2	10.9-11.5	1,385.4	6.9	6.8-7.1	757.8	9.4	9.1-9.7	627.6	5.3	5.1-5.5
	2006-2010	3,148.4	15.2	15.0-15.5	1,792.6	19.9	19.5-20.4	1,355.8	11.7	11.4-12.0	1,376.6	6.6	6.4-6.7	769.6	8.9	8.6-9.2	607.0	5.0	4.8-5.1
	2011-2015	3,715.6	16.8	16.5-17.0	2,131.4	21.8	21.4-22.3	1,584.2	12.9	12.6-13.2	1,413.6	6.3	6.1-6.4	803.0	8.5	8.3-8.8	610.6	4.6	4.5-4.8
2016-2020	3,612.2	15.3	15.1-15.6	2,106.2	19.9	19.6-20.3	1,506.0	11.7	11.4-12.0	1,343.4	5.5	5.4-5.6	759.8	7.4	7.1-7.6	583.6	4.2	4.0-4.3	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York City**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	28,751.2	397.7	395.6-399.8	13,852.2	474.6	470.9-478.3	14,899.0	356.2	353.5-358.8	15,899.8	221.0	219.4-222.6	8,135.6	284.1	281.2-287.0	7,764.2	182.9	181.0-184.8
	1981-1985	30,031.6	417.8	415.7-420.0	14,435.4	503.0	499.2-506.8	15,596.2	371.6	369.0-374.3	15,293.8	213.0	211.5-214.5	7,697.6	274.4	271.5-277.2	7,596.2	177.1	175.3-178.9
	1986-1990	30,739.6	426.7	424.6-428.9	14,824.0	514.6	510.8-518.4	15,915.6	377.9	375.3-380.6	14,922.8	207.9	206.4-209.4	7,403.4	266.4	263.6-269.2	7,519.4	174.1	172.3-175.9
	1991-1995	33,795.2	471.4	469.2-473.7	16,578.8	572.7	568.8-576.7	17,216.4	412.2	409.5-415.0	14,538.2	204.4	202.9-205.9	7,113.8	258.7	256.0-261.5	7,424.4	173.4	171.7-175.2
	1996-2000	34,636.0	468.4	466.1-470.6	16,801.2	558.1	554.2-561.9	17,834.8	414.4	411.6-417.1	13,595.4	185.4	184.0-186.8	6,516.0	227.6	225.1-230.1	7,079.4	160.8	159.1-162.5
	2001-2005	35,446.4	451.3	449.2-453.4	17,505.8	540.3	536.6-543.9	17,940.6	395.6	393.0-398.2	12,829.4	164.6	163.3-165.9	6,181.0	200.9	198.6-203.2	6,648.4	142.7	141.2-144.3
	2006-2010	38,015.0	464.3	462.2-466.4	18,873.2	547.2	543.6-550.7	19,141.8	411.3	408.6-413.9	12,331.4	152.0	150.8-153.2	6,005.4	185.3	183.1-187.4	6,326.0	131.6	130.1-133.0
	2011-2015	39,758.6	445.5	443.5-447.4	19,205.6	498.7	495.5-501.9	20,553.0	413.3	410.8-415.9	12,453.8	139.9	138.8-141.0	6,063.8	167.2	165.3-169.1	6,390.0	122.5	121.1-123.9
2016-2020	40,627.0	423.2	421.3-425.0	19,525.0	459.5	456.5-462.4	21,102.0	402.4	399.9-404.9	11,816.6	120.7	119.7-121.7	5,754.6	141.4	139.7-143.0	6,062.0	107.2	106.0-108.5	
Oral cavity and pharynx	1976-1980	844.4	11.7	11.3-12.0	571.6	18.9	18.2-19.7	272.8	6.5	6.1-6.8	377.4	5.2	5.0-5.5	268.6	9.0	8.5-9.5	108.8	2.6	2.3-2.8
	1981-1985	896.6	12.7	12.3-13.0	611.6	20.6	19.9-21.4	285.0	6.9	6.5-7.2	327.8	4.6	4.4-4.8	238.4	8.1	7.6-8.5	89.4	2.1	1.9-2.4
	1986-1990	835.0	11.8	11.5-12.2	556.8	18.8	18.1-19.6	278.2	6.7	6.3-7.1	278.6	4.0	3.8-4.2	190.6	6.6	6.2-7.1	88.0	2.1	1.9-2.3
	1991-1995	882.0	12.5	12.2-12.9	585.6	19.7	19.0-20.4	296.4	7.2	6.9-7.6	251.2	3.6	3.4-3.8	177.0	6.1	5.7-6.5	74.2	1.7	1.6-1.9
	1996-2000	775.6	10.5	10.1-10.8	513.4	16.2	15.5-16.8	262.2	6.1	5.8-6.5	209.0	2.8	2.7-3.0	142.2	4.6	4.3-5.0	66.8	1.5	1.4-1.7
	2001-2005	761.0	9.6	9.3-9.9	504.6	14.8	14.2-15.4	256.4	5.7	5.4-6.0	196.2	2.5	2.3-2.6	132.8	4.0	3.7-4.3	63.4	1.4	1.2-1.5
	2006-2010	810.8	9.8	9.5-10.1	533.4	14.8	14.2-15.4	277.4	5.9	5.6-6.3	185.4	2.2	2.1-2.4	130.2	3.7	3.5-4.0	55.2	1.1	1.0-1.3
	2011-2015	875.8	9.7	9.4-10.0	577.0	14.5	14.0-15.0	298.8	6.0	5.7-6.3	196.6	2.2	2.0-2.3	136.8	3.5	3.3-3.8	59.8	1.2	1.0-1.3
2016-2020	931.6	9.6	9.4-9.9	608.0	14.2	13.6-14.7	323.6	6.0	5.7-6.4	198.2	2.0	1.9-2.2	136.4	3.2	3.0-3.5	61.8	1.1	1.0-1.2	
Esophagus	1976-1980	456.8	6.2	5.9-6.5	311.4	10.2	9.6-10.7	145.4	3.4	3.2-3.7	406.2	5.5	5.3-5.8	274.6	9.0	8.5-9.6	131.6	3.1	2.9-3.3
	1981-1985	453.4	6.2	6.0-6.5	305.6	10.2	9.7-10.8	147.8	3.5	3.2-3.7	386.8	5.3	5.1-5.6	270.6	9.1	8.6-9.6	116.2	2.7	2.5-2.9
	1986-1990	424.6	5.9	5.7-6.2	278.6	9.6	9.1-10.1	146.0	3.4	3.2-3.7	359.2	5.0	4.8-5.3	248.0	8.6	8.1-9.1	111.2	2.6	2.4-2.8
	1991-1995	407.0	5.7	5.5-6.0	270.0	9.3	8.8-9.8	137.0	3.2	3.0-3.5	327.6	4.6	4.4-4.8	227.6	7.9	7.4-8.4	100.0	2.3	2.1-2.6
	1996-2000	370.8	5.0	4.8-5.3	247.0	8.2	7.8-8.7	123.8	2.8	2.6-3.0	309.0	4.2	4.0-4.4	210.8	7.1	6.6-7.5	98.2	2.2	2.0-2.4
	2001-2005	359.0	4.6	4.4-4.8	239.2	7.4	6.9-7.8	119.8	2.6	2.4-2.8	278.4	3.6	3.4-3.8	193.8	6.1	5.7-6.5	84.6	1.8	1.6-2.0
	2006-2010	333.4	4.1	3.9-4.3	234.4	6.8	6.4-7.3	99.0	2.1	1.9-2.2	271.2	3.3	3.1-3.5	188.6	5.6	5.2-6.0	82.6	1.7	1.5-1.9
	2011-2015	321.4	3.6	3.4-3.8	229.2	6.0	5.7-6.4	92.2	1.8	1.6-1.9	234.8	2.6	2.5-2.8	166.8	4.5	4.2-4.8	68.0	1.3	1.2-1.4
2016-2020	316.0	3.2	3.0-3.3	225.8	5.3	5.0-5.6	90.2	1.6	1.4-1.7	213.0	2.1	2.0-2.3	157.2	3.7	3.4-4.0	55.8	1.0	0.9-1.1	
Stomach	1976-1980	1,067.4	15.1	14.7-15.5	601.4	21.3	20.5-22.1	466.0	11.1	10.6-11.5	794.2	11.3	11.0-11.7	443.8	16.1	15.4-16.8	350.4	8.3	7.9-8.7
	1981-1985	1,020.4	14.3	13.9-14.7	558.4	20.0	19.2-20.8	462.0	10.7	10.2-11.1	697.0	9.8	9.5-10.2	390.2	14.3	13.7-15.0	306.8	7.1	6.7-7.4
	1986-1990	966.2	13.5	13.1-13.9	537.2	19.2	18.4-19.9	429.0	9.8	9.3-10.2	625.4	8.8	8.4-9.1	348.8	12.6	12.0-13.3	276.6	6.3	5.9-6.6
	1991-1995	1,006.6	14.2	13.8-14.6	556.8	20.0	19.3-20.8	449.8	10.3	9.9-10.8	639.6	9.0	8.7-9.4	351.2	13.0	12.4-13.6	288.4	6.6	6.3-6.9
	1996-2000	977.4	13.3	12.9-13.7	541.2	18.7	18.0-19.5	436.2	9.8	9.4-10.2	572.8	7.8	7.5-8.1	305.4	10.7	10.2-11.3	267.4	6.0	5.6-6.3
	2001-2005	957.4	12.3	11.9-12.6	531.0	16.9	16.2-17.6	426.4	9.1	8.7-9.5	507.2	6.5	6.2-6.8	273.4	8.9	8.4-9.4	233.8	4.9	4.7-5.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York City**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	935.0	11.5	11.2-11.9	535.6	16.0	15.4-16.7	399.4	8.4	8.0-8.7	440.8	5.4	5.2-5.7	242.8	7.5	7.1-8.0	198.0	4.1	3.8-4.3
	2011-2015	970.2	11.0	10.6-11.3	555.4	14.9	14.3-15.5	414.8	8.1	7.7-8.5	438.6	5.0	4.8-5.2	249.0	6.9	6.5-7.3	189.6	3.6	3.4-3.9
	2016-2020	960.8	10.0	9.7-10.3	549.8	13.2	12.7-13.7	411.0	7.6	7.3-8.0	416.6	4.3	4.1-4.5	250.4	6.1	5.8-6.5	166.2	3.0	2.8-3.2
Colorectal	1976-1980	4,770.4	66.1	65.2-67.0	2,311.8	81.3	79.7-82.8	2,458.6	56.8	55.8-57.9	2,434.8	34.5	33.9-35.2	1,201.6	44.1	42.9-45.3	1,233.2	28.8	28.1-29.6
	1981-1985	4,780.0	66.3	65.4-67.1	2,298.8	82.5	80.9-84.0	2,481.2	56.6	55.6-57.6	2,170.4	30.5	29.9-31.1	1,043.6	38.5	37.5-39.6	1,126.8	25.8	25.1-26.5
	1986-1990	4,349.8	60.2	59.4-61.0	2,084.2	74.7	73.2-76.1	2,265.6	51.3	50.4-52.3	1,949.2	27.3	26.7-27.8	940.4	35.0	34.0-36.0	1,008.8	22.8	22.1-23.4
	1991-1995	4,221.6	59.2	58.4-60.0	1,976.6	71.7	70.3-73.2	2,245.0	51.5	50.6-52.5	1,776.4	25.0	24.5-25.6	843.6	31.7	30.8-32.7	932.8	21.1	20.5-21.7
	1996-2000	4,322.8	59.0	58.2-59.8	2,030.6	70.2	68.8-71.6	2,292.2	51.7	50.7-52.6	1,645.2	22.5	22.0-23.0	775.4	27.7	26.8-28.6	869.8	19.2	18.7-19.8
	2001-2005	4,157.4	53.3	52.6-54.0	1,993.0	63.5	62.2-64.8	2,164.4	46.5	45.6-47.4	1,481.8	19.1	18.6-19.5	706.0	23.4	22.6-24.2	775.8	16.3	15.8-16.8
	2006-2010	3,839.8	47.2	46.5-47.9	1,866.6	55.5	54.3-56.6	1,973.2	41.3	40.5-42.2	1,336.6	16.5	16.1-16.9	651.6	20.4	19.7-21.1	685.0	13.9	13.5-14.4
	2011-2015	3,546.4	39.9	39.3-40.5	1,784.8	47.4	46.4-48.4	1,761.6	34.4	33.7-35.2	1,240.6	13.9	13.5-14.2	602.6	16.6	16.0-17.2	638.0	11.9	11.5-12.3
	2016-2020	3,423.8	35.8	35.2-36.3	1,772.0	42.4	41.5-43.3	1,651.8	30.7	30.0-31.4	1,120.2	11.5	11.2-11.8	560.4	13.8	13.3-14.3	559.8	9.8	9.4-10.1
Colon excluding rectum	1976-1980	3,413.0	47.6	46.8-48.3	1,600.0	57.1	55.7-58.4	1,813.0	41.9	41.1-42.8	2,008.6	28.5	27.9-29.1	966.4	35.6	34.6-36.7	1,042.2	24.3	23.7-25.0
	1981-1985	3,450.6	48.0	47.3-48.7	1,602.0	58.3	56.9-59.6	1,848.6	42.1	41.3-43.0	1,869.2	26.3	25.7-26.8	888.8	32.9	31.9-33.9	980.4	22.5	21.8-23.1
	1986-1990	3,217.4	44.6	43.9-45.3	1,487.6	53.8	52.6-55.1	1,729.8	39.1	38.3-40.0	1,714.0	24.0	23.5-24.5	823.4	30.7	29.7-31.6	890.6	20.1	19.5-20.7
	1991-1995	3,148.6	44.2	43.5-44.9	1,432.2	52.5	51.2-53.7	1,716.4	39.3	38.4-40.1	1,558.6	22.0	21.5-22.5	736.0	27.8	26.9-28.7	822.6	18.6	18.0-19.2
	1996-2000	3,201.0	43.7	43.0-44.4	1,454.2	50.8	49.7-52.0	1,746.8	39.2	38.4-40.1	1,445.6	19.8	19.3-20.2	677.2	24.3	23.5-25.1	768.4	17.0	16.5-17.5
	2001-2005	3,049.6	39.2	38.6-39.8	1,412.0	45.5	44.5-46.6	1,637.6	35.0	34.2-35.8	1,288.6	16.6	16.2-17.0	603.8	20.1	19.3-20.8	684.8	14.3	13.9-14.8
	2006-2010	2,753.4	34.0	33.4-34.5	1,290.2	38.9	38.0-39.9	1,463.2	30.5	29.8-31.2	1,149.2	14.2	13.8-14.6	554.2	17.4	16.8-18.1	595.0	12.1	11.7-12.5
	2011-2015	2,515.8	28.4	27.9-28.9	1,219.2	32.9	32.0-33.7	1,296.6	25.2	24.6-25.9	1,035.6	11.6	11.2-11.9	492.2	13.7	13.1-14.2	543.4	10.1	9.7-10.5
	2016-2020	2,410.2	25.1	24.6-25.5	1,197.6	28.8	28.0-29.5	1,212.6	22.3	21.7-22.9	907.8	9.3	9.0-9.5	440.2	10.8	10.4-11.3	467.6	8.1	7.8-8.5
Rectum & rectosigmoid	1976-1980	1,357.4	18.6	18.1-19.0	711.8	24.2	23.4-25.1	645.6	14.9	14.4-15.4	426.2	6.0	5.8-6.3	235.2	8.5	7.9-9.0	191.0	4.5	4.2-4.8
	1981-1985	1,329.4	18.3	17.8-18.7	696.8	24.2	23.4-25.0	632.6	14.5	14.0-15.0	301.2	4.2	4.0-4.5	154.8	5.7	5.3-6.1	146.4	3.4	3.1-3.6
	1986-1990	1,132.4	15.6	15.2-16.0	596.6	20.8	20.1-21.6	535.8	12.2	11.8-12.7	235.2	3.3	3.1-3.5	117.0	4.3	4.0-4.7	118.2	2.7	2.4-2.9
	1991-1995	1,073.0	15.0	14.6-15.4	544.4	19.3	18.5-20.0	528.6	12.3	11.8-12.7	217.8	3.1	2.9-3.3	107.6	4.0	3.6-4.3	110.2	2.5	2.3-2.7
	1996-2000	1,121.8	15.3	14.9-15.7	576.4	19.3	18.6-20.1	545.4	12.4	12.0-12.9	199.6	2.7	2.6-2.9	98.2	3.4	3.1-3.7	101.4	2.3	2.1-2.5
	2001-2005	1,107.8	14.1	13.8-14.5	581.0	18.0	17.3-18.6	526.8	11.5	11.0-11.9	193.2	2.5	2.3-2.6	102.2	3.3	3.0-3.6	91.0	1.9	1.8-2.1
	2006-2010	1,086.4	13.2	12.9-13.6	576.4	16.5	15.9-17.2	510.0	10.8	10.4-11.3	187.4	2.3	2.1-2.4	97.4	2.9	2.7-3.2	90.0	1.8	1.7-2.0
	2011-2015	1,030.6	11.5	11.2-11.9	565.6	14.5	14.0-15.1	465.0	9.2	8.8-9.6	205.0	2.3	2.2-2.4	110.4	3.0	2.7-3.2	94.6	1.8	1.6-2.0
	2016-2020	1,013.6	10.7	10.4-11.0	574.4	13.6	13.1-14.1	439.2	8.4	8.0-8.8	212.4	2.2	2.1-2.3	120.2	2.9	2.7-3.2	92.2	1.6	1.5-1.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York City**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	292.8	4.1	3.8-4.3	178.2	6.0	5.6-6.4	114.6	2.7	2.5-2.9	329.2	4.6	4.3-4.8	186.2	6.3	5.9-6.7	143.0	3.4	3.1-3.6
	1981-1985	308.2	4.3	4.1-4.5	197.0	6.8	6.4-7.3	111.2	2.6	2.4-2.8	367.4	5.1	4.9-5.4	213.8	7.5	7.0-7.9	153.6	3.5	3.3-3.8
	1986-1990	347.2	4.9	4.6-5.1	219.8	7.6	7.2-8.1	127.4	2.9	2.7-3.2	397.0	5.5	5.3-5.8	229.4	8.1	7.6-8.6	167.6	3.8	3.6-4.1
	1991-1995	454.2	6.4	6.1-6.7	292.8	10.0	9.5-10.5	161.4	3.8	3.5-4.0	437.4	6.1	5.9-6.4	264.8	9.2	8.7-9.8	172.6	4.0	3.7-4.2
	1996-2000	613.0	8.3	8.0-8.6	413.4	13.3	12.7-13.9	199.6	4.5	4.3-4.8	445.2	6.1	5.8-6.3	284.8	9.4	8.9-9.9	160.4	3.6	3.4-3.9
	2001-2005	735.4	9.3	9.0-9.6	526.2	15.5	14.9-16.1	209.2	4.6	4.3-4.9	507.4	6.5	6.2-6.7	329.4	10.0	9.5-10.5	178.0	3.8	3.6-4.1
	2006-2010	922.0	11.0	10.7-11.4	663.4	18.1	17.5-18.8	258.6	5.5	5.2-5.8	582.0	7.1	6.8-7.3	399.2	11.3	10.8-11.9	182.8	3.8	3.6-4.1
	2011-2015	1,068.4	11.6	11.3-11.9	745.2	18.3	17.7-18.9	323.2	6.2	5.9-6.5	646.6	7.1	6.9-7.4	427.2	10.9	10.4-11.3	219.4	4.2	3.9-4.4
	2016-2020	1,032.2	10.3	10.1-10.6	718.4	16.3	15.8-16.9	313.8	5.6	5.3-5.9	628.2	6.3	6.1-6.5	413.8	9.7	9.2-10.1	214.4	3.7	3.5-4.0
Pancreas	1976-1980	961.4	13.2	12.9-13.6	469.4	16.1	15.4-16.8	492.0	11.3	10.9-11.8	885.2	12.2	11.8-12.6	431.6	14.8	14.2-15.5	453.6	10.5	10.0-10.9
	1981-1985	947.6	13.1	12.7-13.5	443.0	15.6	14.9-16.3	504.6	11.4	10.9-11.8	833.6	11.5	11.2-11.9	397.4	14.0	13.4-14.7	436.2	9.8	9.4-10.3
	1986-1990	913.8	12.7	12.3-13.1	404.8	14.5	13.8-15.1	509.0	11.5	11.0-11.9	783.8	10.9	10.5-11.2	354.6	12.7	12.1-13.3	429.2	9.6	9.2-10.0
	1991-1995	906.6	12.7	12.4-13.1	397.0	14.3	13.7-15.0	509.6	11.6	11.2-12.1	776.0	10.9	10.6-11.3	342.0	12.4	11.8-13.0	434.0	9.8	9.4-10.3
	1996-2000	947.8	13.0	12.6-13.3	424.2	14.7	14.0-15.3	523.6	11.8	11.3-12.2	774.4	10.6	10.3-10.9	343.4	12.0	11.4-12.6	431.0	9.6	9.2-10.0
	2001-2005	963.4	12.4	12.0-12.7	432.4	13.8	13.2-14.4	531.0	11.3	10.9-11.7	799.8	10.3	10.0-10.6	361.6	11.7	11.2-12.3	438.2	9.2	8.8-9.6
	2006-2010	1,100.6	13.5	13.2-13.9	520.8	15.6	15.0-16.3	579.8	12.0	11.5-12.4	862.4	10.6	10.3-11.0	403.0	12.3	11.8-12.9	459.4	9.4	9.0-9.8
	2011-2015	1,212.4	13.6	13.2-13.9	579.6	15.4	14.9-16.0	632.8	12.2	11.7-12.6	933.0	10.5	10.2-10.8	442.4	12.0	11.5-12.6	490.6	9.3	8.9-9.7
	2016-2020	1,312.0	13.4	13.0-13.7	629.2	15.0	14.5-15.5	682.8	12.1	11.6-12.5	974.0	9.9	9.6-10.1	467.8	11.3	10.8-11.8	506.2	8.8	8.4-9.1
Larynx	1976-1980	445.2	6.0	5.7-6.2	362.8	11.7	11.1-12.2	82.4	2.0	1.8-2.2	187.2	2.5	2.4-2.7	152.4	5.0	4.6-5.4	34.8	0.8	0.7-1.0
	1981-1985	454.0	6.2	6.0-6.5	357.6	11.8	11.2-12.4	96.4	2.3	2.1-2.6	167.8	2.3	2.1-2.4	132.4	4.4	4.1-4.8	35.4	0.8	0.7-1.0
	1986-1990	465.0	6.5	6.3-6.8	352.8	11.8	11.2-12.4	112.2	2.8	2.6-3.1	161.4	2.3	2.1-2.4	127.4	4.4	4.0-4.7	34.0	0.8	0.7-1.0
	1991-1995	407.6	5.7	5.5-6.0	315.2	10.6	10.1-11.2	92.4	2.3	2.1-2.5	168.4	2.4	2.2-2.5	128.4	4.4	4.1-4.8	40.0	1.0	0.8-1.1
	1996-2000	361.4	4.9	4.7-5.1	282.4	9.2	8.7-9.7	79.0	1.9	1.7-2.1	143.2	2.0	1.8-2.1	115.0	3.8	3.5-4.1	28.2	0.7	0.6-0.8
	2001-2005	329.2	4.2	4.0-4.4	258.4	7.8	7.4-8.2	70.8	1.6	1.4-1.7	114.8	1.5	1.3-1.6	93.2	2.9	2.6-3.1	21.6	0.5	0.4-0.6
	2006-2010	310.0	3.7	3.6-3.9	241.2	6.9	6.5-7.3	68.8	1.5	1.3-1.6	107.6	1.3	1.2-1.4	85.2	2.5	2.3-2.8	22.4	0.5	0.4-0.6
	2011-2015	286.2	3.1	3.0-3.3	229.0	5.9	5.5-6.2	57.2	1.1	1.0-1.2	94.8	1.1	1.0-1.2	74.2	2.0	1.8-2.2	20.6	0.4	0.3-0.5
	2016-2020	243.4	2.5	2.3-2.6	192.4	4.5	4.2-4.8	51.0	0.9	0.8-1.1	72.6	0.7	0.6-0.8	58.8	1.4	1.2-1.6	13.8	0.2	0.2-0.3
Lung and bronchus	1976-1980	3,986.8	53.3	52.5-54.1	2,764.0	90.9	89.4-92.5	1,222.8	28.2	27.5-28.9	3,259.0	43.7	43.0-44.4	2,308.2	76.5	75.1-78.0	950.8	21.9	21.3-22.6
	1981-1985	4,153.6	56.5	55.7-57.3	2,688.8	91.5	89.9-93.1	1,464.8	33.7	33.0-34.5	3,310.2	45.1	44.4-45.8	2,171.4	74.6	73.2-76.1	1,138.8	26.1	25.4-26.8
	1986-1990	4,252.0	58.4	57.6-59.2	2,579.8	88.8	87.3-90.4	1,672.2	38.7	37.9-39.6	3,392.4	46.6	45.9-47.3	2,109.2	73.5	72.1-75.0	1,283.2	29.5	28.8-30.3
	1991-1995	4,337.4	60.7	59.9-61.6	2,491.4	86.9	85.4-88.5	1,846.0	43.6	42.7-44.5	3,391.0	47.5	46.8-48.2	1,994.0	70.6	69.2-72.0	1,397.0	32.7	31.9-33.5
	1996-2000	4,229.2	57.7	56.9-58.5	2,306.4	77.7	76.3-79.2	1,922.8	44.3	43.4-45.2	3,130.0	42.8	42.1-43.4	1,763.8	60.2	59.0-61.5	1,366.2	31.2	30.5-32.0
	2001-2005	4,170.2	53.6	52.9-54.4	2,238.8	70.7	69.4-72.1	1,931.4	42.1	41.3-43.0	2,965.0	38.2	37.6-38.8	1,631.6	52.3	51.1-53.4	1,333.4	28.8	28.1-29.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York City**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	4,244.4	52.6	51.9-53.3	2,205.2	66.4	65.2-67.7	2,039.2	43.2	42.4-44.1	2,779.8	34.5	33.9-35.0	1,492.8	45.7	44.7-46.8	1,287.0	26.9	26.2-27.6
	2011-2015	4,325.0	48.7	48.1-49.4	2,195.8	59.3	58.2-60.4	2,129.2	41.6	40.8-42.4	2,667.8	30.0	29.5-30.6	1,430.4	39.3	38.4-40.2	1,237.4	23.7	23.1-24.3
	2016-2020	4,310.2	43.9	43.3-44.5	2,143.6	51.5	50.5-52.5	2,166.6	38.6	37.8-39.3	2,197.8	22.3	21.9-22.7	1,164.0	28.4	27.6-29.1	1,033.8	18.0	17.5-18.5
Melanoma of the skin	1976-1980	400.2	5.6	5.3-5.8	192.6	6.3	5.9-6.7	207.6	5.1	4.8-5.4	143.2	2.0	1.8-2.2	77.4	2.6	2.3-2.9	65.8	1.6	1.4-1.8
	1981-1985	435.6	6.1	5.9-6.4	223.8	7.5	7.0-7.9	211.8	5.2	4.9-5.6	162.2	2.3	2.1-2.5	93.2	3.2	2.9-3.5	69.0	1.7	1.5-1.9
	1986-1990	428.0	5.9	5.7-6.2	222.2	7.5	7.1-8.0	205.8	4.9	4.6-5.2	140.0	2.0	1.8-2.1	80.0	2.8	2.5-3.1	60.0	1.4	1.3-1.6
	1991-1995	429.0	5.9	5.7-6.2	234.4	7.9	7.4-8.4	194.6	4.6	4.3-4.9	114.0	1.6	1.5-1.7	64.8	2.2	2.0-2.5	49.2	1.2	1.0-1.3
	1996-2000	551.0	7.3	7.1-7.6	292.0	9.6	9.1-10.1	259.0	6.0	5.6-6.3	125.6	1.7	1.6-1.8	69.4	2.3	2.1-2.6	56.2	1.3	1.1-1.4
	2001-2005	724.0	9.1	8.8-9.4	379.4	11.5	11.0-12.1	344.6	7.6	7.2-8.0	111.4	1.4	1.3-1.5	57.0	1.9	1.6-2.1	54.4	1.2	1.0-1.3
	2006-2010	930.4	11.3	11.0-11.6	495.8	14.3	13.8-14.9	434.6	9.3	8.9-9.7	114.6	1.4	1.3-1.5	67.0	2.0	1.8-2.3	47.6	1.0	0.9-1.1
	2011-2015	949.8	10.7	10.4-11.0	515.4	13.8	13.3-14.3	434.4	8.7	8.3-9.1	108.0	1.2	1.1-1.3	62.2	1.7	1.5-1.9	45.8	0.9	0.8-1.0
	2016-2020	911.4	9.6	9.3-9.9	506.2	12.4	11.9-12.9	405.2	7.7	7.4-8.1	84.4	0.9	0.8-1.0	49.4	1.2	1.1-1.4	35.0	0.6	0.5-0.7
Female breast	1976-1980							3,994.8	98.6	97.2-100.1							1,576.4	37.9	37.0-38.7
	1981-1985							4,183.8	104.0	102.6-105.5							1,490.8	35.9	35.1-36.8
	1986-1990							4,479.8	110.4	109.0-111.9							1,485.4	35.6	34.8-36.4
	1991-1995							4,929.8	121.8	120.3-123.3							1,437.4	34.6	33.8-35.4
	1996-2000							5,285.6	125.5	124.0-127.0							1,297.8	30.0	29.3-30.8
	2001-2005							5,033.8	112.8	111.4-114.2							1,194.4	25.9	25.3-26.6
	2006-2010							5,405.2	117.9	116.5-119.4							1,054.2	22.1	21.5-22.7
	2011-2015							5,928.8	121.5	120.1-123.0							1,027.0	20.0	19.5-20.6
	2016-2020							6,422.4	126.1	124.7-127.6							987.6	18.0	17.5-18.5
Cervix uteri	1976-1980							602.2	15.9	15.4-16.5							216.4	5.4	5.1-5.8
	1981-1985							539.6	14.2	13.7-14.8							174.4	4.4	4.1-4.7
	1986-1990							535.8	13.8	13.3-14.3							148.8	3.7	3.5-4.0
	1991-1995							588.2	14.8	14.2-15.3							157.2	3.9	3.6-4.2
	1996-2000							566.2	13.5	13.0-14.0							162.4	3.8	3.6-4.1
	2001-2005							472.8	10.8	10.4-11.2							142.6	3.2	3.0-3.5
	2006-2010							445.2	10.0	9.5-10.4							136.8	3.0	2.7-3.2
	2011-2015							428.2	9.1	8.8-9.5							139.2	2.8	2.6-3.0
	2016-2020							415.2	8.6	8.2-9.0							110.8	2.2	2.0-2.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York City**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							895.4	20.7	20.1-21.4							257.8	5.8	5.5-6.2
	1981-1985							871.8	20.4	19.7-21.0							218.4	4.9	4.6-5.2
	1986-1990							813.6	19.2	18.6-19.8							199.6	4.5	4.2-4.8
	1991-1995							943.8	22.8	22.2-23.5							223.6	5.2	4.9-5.5
	1996-2000							1,088.0	25.6	24.9-26.3							243.4	5.6	5.2-5.9
	2001-2005							1,222.4	27.1	26.5-27.8							246.2	5.4	5.1-5.7
	2006-2010							1,377.2	29.3	28.6-30.0							276.2	5.8	5.5-6.1
	2011-2015							1,566.8	30.7	30.0-31.4							335.2	6.4	6.1-6.7
	2016-2020							1,654.0	30.5	29.9-31.2							386.6	6.8	6.5-7.1
Ovary	1976-1980							629.8	15.1	14.6-15.7							424.8	9.9	9.5-10.3
	1981-1985							665.4	16.2	15.6-16.7							388.4	9.1	8.7-9.5
	1986-1990							617.0	15.0	14.5-15.5							383.0	8.9	8.5-9.3
	1991-1995							677.6	16.6	16.0-17.1							401.2	9.5	9.1-10.0
	1996-2000							621.2	14.7	14.1-15.2							361.2	8.4	8.0-8.7
	2001-2005							608.4	13.6	13.1-14.0							357.0	7.8	7.4-8.2
	2006-2010							582.6	12.6	12.1-13.0							339.8	7.1	6.7-7.4
	2011-2015							598.2	12.1	11.7-12.5							340.4	6.6	6.3-7.0
	2016-2020							568.0	11.1	10.7-11.5							331.6	6.0	5.7-6.3
Prostate	1976-1980				2,140.0	78.8	77.3-80.4							745.8	29.8	28.8-30.9			
	1981-1985				2,348.2	86.1	84.5-87.7							798.6	32.0	31.0-33.1			
	1986-1990				2,629.8	96.6	94.9-98.3							837.2	33.4	32.3-34.4			
	1991-1995				4,179.0	147.9	145.9-150.0							922.0	37.2	36.1-38.3			
	1996-2000				4,953.6	166.3	164.2-168.4							823.0	31.9	30.9-32.9			
	2001-2005				5,278.8	163.2	161.2-165.2							762.8	27.4	26.6-28.3			
	2006-2010				5,805.2	165.7	163.8-167.7							699.4	23.8	23.0-24.6			
	2011-2015				5,415.4	135.3	133.7-137.0							680.0	20.3	19.6-21.0			
	2016-2020				5,560.4	124.9	123.4-126.4							668.6	17.2	16.6-17.8			
Testis	1976-1980				86.2	2.5	2.3-2.8							17.6	0.5	0.4-0.6			
	1981-1985				108.4	3.0	2.7-3.3							10.6	0.3	0.2-0.4			
	1986-1990				122.0	3.2	2.9-3.4							10.6	0.3	0.2-0.4			
	1991-1995				134.4	3.4	3.1-3.7							9.6	0.3	0.2-0.4			
	1996-2000				143.0	3.5	3.3-3.8							6.2	0.2	0.1-0.2			
	2001-2005				165.4	4.0	3.7-4.3							9.0	0.2	0.2-0.3			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020  
New York City**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				194.4	4.7	4.4-5.0							8.8	0.2	0.2-0.3			
	2011-2015				205.0	4.8	4.5-5.1							7.4	0.2	0.1-0.2			
	2016-2020				225.4	5.3	5.0-5.6							7.0	0.2	0.1-0.2			
Urinary bladder (incl. in situ)	1976-1980	1,425.4	19.6	19.2-20.1	1,014.8	35.7	34.7-36.7	410.6	9.4	9.0-9.9	415.4	6.1	5.8-6.3	279.2	10.7	10.2-11.4	136.2	3.3	3.0-3.5
	1981-1985	1,423.8	19.6	19.1-20.0	1,000.4	35.7	34.7-36.8	423.4	9.6	9.2-10.0	356.6	5.1	4.8-5.3	233.8	9.1	8.5-9.6	122.8	2.8	2.6-3.0
	1986-1990	1,333.2	18.4	18.0-18.9	930.8	33.7	32.7-34.7	402.4	9.1	8.7-9.5	327.6	4.6	4.4-4.8	206.4	8.1	7.6-8.6	121.2	2.7	2.5-2.9
	1991-1995	1,370.0	19.2	18.7-19.6	948.0	34.7	33.7-35.7	422.0	9.7	9.3-10.1	323.8	4.6	4.4-4.8	208.2	8.2	7.7-8.7	115.6	2.6	2.4-2.8
	1996-2000	1,367.6	18.7	18.3-19.2	948.8	33.3	32.4-34.3	418.8	9.4	9.0-9.8	308.6	4.2	4.0-4.5	194.4	7.3	6.9-7.8	114.2	2.5	2.3-2.7
	2001-2005	1,399.6	18.1	17.6-18.5	982.4	32.3	31.4-33.2	417.2	8.9	8.5-9.3	287.2	3.7	3.5-3.9	183.8	6.4	6.0-6.9	103.4	2.1	1.9-2.3
	2006-2010	1,446.4	18.0	17.5-18.4	1,058.0	33.0	32.1-33.9	388.4	8.1	7.7-8.5	299.0	3.7	3.5-3.9	195.0	6.5	6.0-6.9	104.0	2.1	1.9-2.2
	2011-2015	1,508.4	17.1	16.7-17.5	1,103.8	30.8	30.0-31.6	404.6	7.7	7.4-8.1	317.2	3.6	3.4-3.7	210.0	6.2	5.8-6.6	107.2	1.9	1.8-2.1
2016-2020	1,542.2	15.8	15.4-16.1	1,131.4	27.8	27.1-28.6	410.8	7.3	7.0-7.6	294.6	3.0	2.8-3.1	198.6	5.1	4.8-5.5	96.0	1.6	1.5-1.8	
Kidney and renal pelvis	1976-1980	513.4	6.9	6.7-7.2	322.4	10.5	10.0-11.1	191.0	4.4	4.1-4.7	258.4	3.5	3.4-3.8	168.0	5.7	5.3-6.1	90.4	2.1	1.9-2.3
	1981-1985	555.8	7.7	7.4-8.0	336.4	11.4	10.8-12.0	219.4	5.2	4.8-5.5	238.0	3.3	3.1-3.5	141.0	4.9	4.5-5.3	97.0	2.2	2.0-2.4
	1986-1990	624.0	8.6	8.3-8.9	371.6	12.6	12.0-13.2	252.4	5.9	5.6-6.3	239.2	3.3	3.1-3.5	140.8	5.0	4.6-5.4	98.4	2.3	2.1-2.5
	1991-1995	709.4	9.9	9.6-10.3	429.6	14.6	14.0-15.2	279.8	6.7	6.3-7.0	225.8	3.2	3.0-3.4	137.2	4.9	4.5-5.2	88.6	2.1	1.9-2.3
	1996-2000	849.2	11.5	11.2-11.9	503.4	16.5	15.8-17.1	345.8	8.0	7.7-8.4	220.6	3.0	2.8-3.2	131.4	4.5	4.2-4.9	89.2	2.0	1.8-2.2
	2001-2005	1,004.2	12.8	12.5-13.2	617.4	18.5	17.9-19.2	386.8	8.5	8.2-8.9	214.8	2.8	2.6-2.9	130.0	4.1	3.8-4.5	84.8	1.8	1.6-2.0
	2006-2010	1,159.4	14.2	13.8-14.6	727.6	20.7	20.0-21.4	431.8	9.3	8.9-9.7	227.0	2.8	2.6-3.0	141.8	4.3	4.0-4.7	85.2	1.8	1.6-1.9
	2011-2015	1,275.0	14.3	13.9-14.7	808.8	20.5	19.9-21.2	466.2	9.4	9.0-9.8	219.8	2.5	2.3-2.6	138.0	3.7	3.5-4.0	81.8	1.5	1.4-1.7
2016-2020	1,388.6	14.5	14.2-14.9	886.4	20.8	20.2-21.4	502.2	9.5	9.1-9.9	213.4	2.2	2.1-2.3	139.8	3.4	3.2-3.7	73.6	1.3	1.2-1.4	
Brain and other central nervous system	1976-1980	400.8	5.4	5.2-5.6	209.6	6.5	6.1-6.9	191.2	4.6	4.3-4.9	301.8	4.0	3.8-4.2	162.4	5.1	4.7-5.5	139.4	3.2	3.0-3.5
	1981-1985	416.4	5.7	5.5-6.0	219.4	7.0	6.6-7.5	197.0	4.8	4.5-5.1	295.4	4.0	3.8-4.2	152.2	5.0	4.6-5.4	143.2	3.3	3.1-3.6
	1986-1990	455.8	6.3	6.1-6.6	240.0	7.7	7.3-8.2	215.8	5.3	5.0-5.6	259.8	3.6	3.4-3.8	139.4	4.7	4.3-5.1	120.4	2.8	2.6-3.1
	1991-1995	497.8	6.8	6.6-7.1	255.2	8.0	7.6-8.5	242.6	5.9	5.6-6.3	242.4	3.4	3.2-3.6	124.6	4.1	3.8-4.5	117.8	2.8	2.6-3.1
	1996-2000	520.8	6.9	6.6-7.1	260.4	7.8	7.3-8.2	260.4	6.2	5.8-6.5	247.6	3.3	3.2-3.5	120.0	3.8	3.5-4.1	127.6	3.0	2.7-3.2
	2001-2005	481.4	6.1	5.9-6.3	250.4	7.2	6.8-7.6	231.0	5.3	5.0-5.6	225.8	2.9	2.7-3.0	124.0	3.7	3.4-4.0	101.8	2.3	2.1-2.5
	2006-2010	483.2	6.0	5.7-6.2	254.0	7.0	6.6-7.4	229.2	5.1	4.8-5.5	243.2	3.0	2.8-3.2	129.6	3.7	3.4-4.0	113.6	2.4	2.2-2.7
	2011-2015	504.6	5.9	5.6-6.1	269.8	6.9	6.5-7.3	234.8	5.0	4.7-5.3	263.0	3.0	2.8-3.1	132.4	3.5	3.2-3.8	130.6	2.6	2.4-2.8
2016-2020	524.8	5.8	5.6-6.1	290.8	7.0	6.7-7.4	234.0	4.8	4.5-5.1	290.4	3.1	2.9-3.3	160.4	3.9	3.6-4.2	130.0	2.5	2.3-2.7	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York City**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	252.0	3.5	3.3-3.7	70.6	2.3	2.0-2.5	181.4	4.6	4.3-4.9	48.8	0.7	0.6-0.8	18.8	0.6	0.5-0.8	30.0	0.7	0.6-0.9
	1981-1985	278.4	4.0	3.7-4.2	76.6	2.5	2.2-2.7	201.8	5.2	4.8-5.5	47.4	0.7	0.6-0.7	16.6	0.6	0.5-0.7	30.8	0.7	0.6-0.8
	1986-1990	282.2	3.9	3.7-4.1	75.0	2.4	2.2-2.7	207.2	5.2	4.8-5.5	38.8	0.5	0.5-0.6	10.8	0.4	0.3-0.5	28.0	0.6	0.5-0.7
	1991-1995	390.8	5.3	5.0-5.5	96.4	3.0	2.7-3.3	294.4	7.2	6.8-7.6	38.4	0.5	0.5-0.6	11.6	0.4	0.3-0.5	26.8	0.6	0.5-0.7
	1996-2000	535.4	6.9	6.6-7.1	125.2	3.7	3.4-4.0	410.2	9.7	9.3-10.1	39.8	0.5	0.5-0.6	13.4	0.5	0.4-0.6	26.4	0.6	0.5-0.7
	2001-2005	791.0	9.7	9.4-10.0	188.8	5.2	4.9-5.6	602.2	13.7	13.2-14.2	40.4	0.5	0.4-0.6	14.8	0.5	0.4-0.6	25.6	0.5	0.4-0.6
	2006-2010	1,357.4	16.2	15.8-16.6	303.8	8.0	7.6-8.4	1,053.6	23.5	22.9-24.2	48.4	0.6	0.5-0.7	18.6	0.6	0.5-0.7	29.8	0.6	0.5-0.7
	2011-2015	1,821.4	20.8	20.3-21.2	410.8	10.2	9.7-10.6	1,410.6	30.3	29.6-31.0	48.6	0.5	0.5-0.6	18.2	0.5	0.4-0.6	30.4	0.6	0.5-0.7
2016-2020	1,730.0	19.4	19.0-19.9	427.0	10.2	9.8-10.7	1,303.0	27.8	27.1-28.5	47.8	0.5	0.4-0.6	18.2	0.5	0.4-0.6	29.6	0.5	0.4-0.6	
Hodgkin lymphoma	1976-1980	245.8	3.2	3.1-3.4	131.2	3.9	3.6-4.2	114.6	2.7	2.5-3.0	91.2	1.2	1.1-1.4	51.6	1.6	1.4-1.8	39.6	1.0	0.8-1.1
	1981-1985	245.6	3.3	3.1-3.4	143.6	4.3	3.9-4.6	102.0	2.4	2.2-2.6	77.6	1.0	0.9-1.2	43.4	1.3	1.1-1.5	34.2	0.8	0.7-0.9
	1986-1990	241.6	3.2	3.0-3.3	142.6	4.1	3.8-4.4	99.0	2.4	2.2-2.6	66.0	0.9	0.8-1.0	42.8	1.3	1.1-1.5	23.2	0.5	0.4-0.6
	1991-1995	247.6	3.2	3.0-3.4	137.0	3.9	3.6-4.2	110.6	2.6	2.4-2.9	56.6	0.8	0.7-0.8	31.2	0.9	0.8-1.1	25.4	0.6	0.5-0.7
	1996-2000	241.0	3.0	2.8-3.2	129.8	3.5	3.2-3.8	111.2	2.6	2.4-2.8	42.8	0.6	0.5-0.6	21.8	0.6	0.5-0.8	21.0	0.5	0.4-0.6
	2001-2005	264.6	3.2	3.0-3.4	141.6	3.7	3.5-4.0	123.0	2.8	2.6-3.0	34.6	0.4	0.4-0.5	20.2	0.6	0.5-0.7	14.4	0.3	0.3-0.4
	2006-2010	274.8	3.3	3.1-3.5	145.0	3.7	3.5-4.0	129.8	2.9	2.7-3.1	32.8	0.4	0.3-0.5	19.6	0.5	0.4-0.7	13.2	0.3	0.2-0.4
	2011-2015	279.4	3.2	3.0-3.4	146.8	3.6	3.3-3.9	132.6	2.9	2.7-3.1	32.2	0.4	0.3-0.4	17.0	0.5	0.4-0.6	15.2	0.3	0.2-0.4
2016-2020	251.0	2.9	2.7-3.0	136.0	3.3	3.1-3.6	115.0	2.5	2.3-2.7	22.6	0.2	0.2-0.3	13.0	0.3	0.2-0.4	9.6	0.2	0.1-0.3	
Non-Hodgkin Lymphomas	1976-1980	839.4	11.5	11.2-11.9	413.8	13.6	13.0-14.3	425.6	10.0	9.6-10.5	426.8	5.9	5.7-6.2	221.2	7.5	7.0-8.0	205.6	4.8	4.5-5.1
	1981-1985	1,037.4	14.4	14.0-14.8	517.8	17.5	16.8-18.2	519.6	12.1	11.6-12.6	484.6	6.8	6.5-7.1	239.2	8.3	7.8-8.8	245.4	5.7	5.4-6.0
	1986-1990	1,263.2	17.6	17.1-18.0	705.8	23.2	22.4-24.0	557.4	13.0	12.5-13.5	530.2	7.4	7.1-7.7	281.8	9.7	9.2-10.2	248.4	5.7	5.4-6.0
	1991-1995	1,565.4	21.6	21.1-22.1	898.4	28.8	27.9-29.7	667.0	15.8	15.2-16.3	548.2	7.7	7.4-7.9	287.0	9.9	9.3-10.4	261.2	6.0	5.7-6.4
	1996-2000	1,516.6	20.3	19.8-20.7	806.8	25.3	24.5-26.1	709.8	16.4	15.9-16.9	544.6	7.4	7.1-7.7	274.2	9.3	8.8-9.8	270.4	6.1	5.8-6.4
	2001-2005	1,543.2	19.6	19.2-20.0	805.0	24.0	23.3-24.8	738.2	16.3	15.7-16.8	479.2	6.1	5.9-6.4	252.6	8.0	7.5-8.4	226.6	4.8	4.5-5.1
	2006-2010	1,661.0	20.4	19.9-20.8	875.4	25.0	24.3-25.8	785.6	16.8	16.3-17.4	439.2	5.4	5.2-5.7	229.4	7.0	6.6-7.4	209.8	4.4	4.1-4.6
	2011-2015	1,790.4	20.3	19.8-20.7	950.8	24.9	24.2-25.7	839.6	16.7	16.2-17.2	422.6	4.8	4.6-5.0	228.2	6.4	6.0-6.7	194.4	3.7	3.5-4.0
2016-2020	1,844.2	19.5	19.1-19.9	970.2	23.4	22.8-24.1	874.0	16.5	16.0-17.0	405.8	4.2	4.0-4.4	226.6	5.6	5.3-6.0	179.2	3.1	2.9-3.3	
Myeloma	1976-1980	357.4	4.9	4.7-5.2	174.6	6.0	5.6-6.5	182.8	4.2	3.9-4.5	222.0	3.0	2.9-3.2	102.2	3.5	3.2-3.8	119.8	2.7	2.5-3.0
	1981-1985	397.0	5.5	5.3-5.8	186.0	6.6	6.2-7.1	211.0	4.8	4.6-5.2	239.0	3.3	3.1-3.5	109.4	3.9	3.6-4.2	129.6	2.9	2.7-3.2
	1986-1990	392.6	5.5	5.2-5.7	190.2	6.7	6.3-7.2	202.4	4.7	4.4-5.0	228.8	3.2	3.0-3.4	111.6	4.0	3.7-4.4	117.2	2.6	2.4-2.9
	1991-1995	423.6	5.9	5.7-6.2	192.8	6.8	6.4-7.2	230.8	5.4	5.1-5.7	254.8	3.6	3.4-3.8	117.2	4.2	3.9-4.6	137.6	3.2	2.9-3.4
	1996-2000	476.4	6.5	6.2-6.7	214.8	7.2	6.8-7.7	261.6	6.0	5.7-6.3	236.0	3.2	3.0-3.4	100.6	3.5	3.2-3.8	135.4	3.1	2.8-3.3
	2001-2005	566.0	7.2	7.0-7.5	266.8	8.4	7.9-8.9	299.2	6.5	6.2-6.9	239.2	3.1	2.9-3.3	108.6	3.5	3.2-3.9	130.6	2.8	2.6-3.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**New York City**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	668.0	8.2	7.9-8.5	343.8	10.2	9.7-10.7	324.2	6.9	6.5-7.2	234.0	2.9	2.7-3.1	121.6	3.8	3.5-4.1	112.4	2.3	2.2-2.6
	2011-2015	821.8	9.2	8.9-9.5	430.0	11.4	10.9-11.9	391.8	7.7	7.4-8.1	257.8	2.9	2.8-3.1	130.8	3.6	3.3-3.9	127.0	2.4	2.3-2.6
	2016-2020	846.2	8.7	8.4-8.9	435.2	10.3	9.9-10.8	411.0	7.5	7.1-7.8	262.0	2.6	2.5-2.8	136.6	3.3	3.1-3.6	125.4	2.2	2.0-2.3
Leukemias	1976-1980	801.2	11.3	10.9-11.7	429.6	14.6	13.9-15.2	371.6	9.0	8.6-9.5	553.0	7.9	7.6-8.2	298.4	10.4	9.8-10.9	254.6	6.2	5.9-6.6
	1981-1985	818.6	11.5	11.2-11.9	441.8	15.3	14.7-16.0	376.8	9.1	8.6-9.5	536.8	7.6	7.3-7.9	284.0	10.0	9.5-10.6	252.8	6.0	5.7-6.4
	1986-1990	819.4	11.4	11.1-11.8	419.4	14.4	13.8-15.1	400.0	9.5	9.1-9.9	483.0	6.7	6.4-7.0	247.0	8.7	8.2-9.2	236.0	5.5	5.2-5.8
	1991-1995	856.4	11.9	11.6-12.3	441.8	15.0	14.4-15.7	414.6	9.8	9.4-10.3	484.2	6.8	6.5-7.0	255.2	9.0	8.5-9.6	229.0	5.3	5.0-5.7
	1996-2000	889.6	12.0	11.6-12.3	465.6	15.2	14.6-15.8	424.0	9.8	9.3-10.2	463.4	6.3	6.0-6.5	240.0	8.1	7.7-8.6	223.4	5.1	4.8-5.4
	2001-2005	937.8	12.0	11.7-12.4	499.2	15.4	14.7-16.0	438.6	9.7	9.3-10.1	462.2	5.9	5.7-6.2	241.0	7.7	7.3-8.2	221.2	4.7	4.5-5.0
	2006-2010	1,050.4	13.1	12.7-13.4	572.6	16.8	16.2-17.5	477.8	10.4	10.0-10.8	456.6	5.7	5.4-5.9	238.4	7.3	6.9-7.7	218.2	4.6	4.3-4.9
	2011-2015	1,247.2	14.3	13.9-14.6	684.2	18.3	17.7-18.9	563.0	11.4	11.0-11.9	478.6	5.5	5.3-5.7	264.2	7.4	7.0-7.8	214.4	4.2	3.9-4.4
	2016-2020	1,210.6	13.0	12.7-13.4	679.0	16.7	16.1-17.3	531.6	10.2	9.8-10.6	459.2	4.8	4.6-5.0	245.8	6.2	5.9-6.6	213.4	3.9	3.6-4.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York State Excl New York City**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	39,106.2	409.2	407.4-411.1	19,302.0	487.8	484.6-491.1	19,804.2	367.3	365.0-369.6	20,837.6	220.5	219.2-221.9	10,892.6	283.4	280.9-285.9	9,945.0	182.5	180.8-184.1
	1981-1985	43,499.2	432.3	430.5-434.2	21,446.6	515.7	512.4-518.9	22,052.6	388.9	386.6-391.3	22,214.0	222.4	221.1-223.7	11,487.2	286.3	283.8-288.7	10,726.8	185.1	183.5-186.7
	1986-1990	47,243.2	450.1	448.2-451.9	23,095.8	530.2	527.0-533.4	24,147.4	409.6	407.3-412.0	23,284.2	222.9	221.6-224.2	11,923.4	284.3	281.9-286.7	11,360.8	187.1	185.6-188.7
	1991-1995	52,962.8	479.2	477.4-481.0	26,859.6	575.1	571.9-578.2	26,103.2	423.4	421.1-425.8	24,036.4	218.0	216.8-219.3	12,133.8	274.4	272.2-276.7	11,902.6	185.5	184.0-187.1
	1996-2000	58,714.8	510.5	508.7-512.4	29,628.8	598.4	595.3-601.5	29,086.0	455.0	452.6-457.3	23,760.4	204.9	203.8-206.1	11,930.6	252.6	250.5-254.6	11,829.8	175.8	174.4-177.2
	2001-2005	62,622.6	516.0	514.2-517.8	31,953.0	603.1	600.1-606.1	30,669.6	458.1	455.8-460.4	23,258.8	188.9	187.8-190.0	11,545.4	227.4	225.5-229.3	11,713.4	165.1	163.7-166.4
	2006-2010	67,069.6	521.8	520.0-523.6	34,475.0	599.2	596.3-602.1	32,594.6	468.0	465.7-470.3	22,703.0	173.1	172.1-174.1	11,384.4	205.4	203.7-207.1	11,318.6	151.9	150.6-153.2
	2011-2015	69,248.2	503.7	502.0-505.5	34,823.2	552.1	549.5-554.8	34,425.0	472.1	469.8-474.4	22,515.4	159.2	158.3-160.2	11,406.4	187.2	185.7-188.8	11,109.0	139.9	138.7-141.1
2016-2020	71,605.6	489.7	488.0-491.3	36,496.0	532.0	529.5-534.5	35,109.6	461.5	459.2-463.8	21,818.2	143.4	142.6-144.3	11,060.2	166.1	164.6-167.5	10,758.0	127.7	126.5-128.8	
Oral cavity and pharynx	1976-1980	1,087.4	11.0	10.7-11.3	720.2	16.7	16.2-17.3	367.2	6.7	6.4-7.0	446.4	4.5	4.3-4.7	303.2	7.2	6.9-7.6	143.2	2.6	2.4-2.8
	1981-1985	1,158.4	11.3	11.0-11.6	776.2	17.5	16.9-18.1	382.2	6.7	6.4-7.0	430.6	4.2	4.0-4.4	285.6	6.6	6.2-7.0	145.0	2.5	2.3-2.7
	1986-1990	1,110.4	10.6	10.3-10.9	718.8	15.7	15.2-16.3	391.6	6.6	6.3-6.9	412.2	3.9	3.8-4.1	264.4	6.0	5.7-6.3	147.8	2.4	2.3-2.6
	1991-1995	1,094.6	10.0	9.8-10.3	707.2	14.7	14.3-15.3	387.4	6.3	6.0-6.6	349.2	3.2	3.0-3.3	225.0	4.8	4.5-5.1	124.2	2.0	1.8-2.1
	1996-2000	1,149.6	10.1	9.8-10.3	750.4	14.7	14.3-15.2	399.2	6.3	6.0-6.6	303.6	2.6	2.5-2.8	196.6	4.0	3.7-4.2	107.0	1.6	1.5-1.7
	2001-2005	1,214.0	10.0	9.7-10.2	797.4	14.4	14.0-14.9	416.6	6.2	5.9-6.5	283.8	2.3	2.2-2.4	185.6	3.5	3.2-3.7	98.2	1.4	1.2-1.5
	2006-2010	1,418.0	10.9	10.6-11.1	969.8	16.1	15.6-16.5	448.2	6.4	6.1-6.7	274.0	2.1	2.0-2.2	185.0	3.1	2.9-3.4	89.0	1.2	1.1-1.3
	2011-2015	1,609.2	11.5	11.2-11.7	1,123.8	17.1	16.7-17.6	485.4	6.5	6.2-6.8	294.6	2.1	1.9-2.2	201.6	3.1	3.0-3.4	93.0	1.1	1.0-1.3
2016-2020	1,840.2	12.4	12.1-12.6	1,280.2	18.3	17.8-18.7	560.0	7.1	6.9-7.4	343.0	2.2	2.1-2.3	237.6	3.4	3.2-3.6	105.4	1.2	1.1-1.4	
Esophagus	1976-1980	425.4	4.4	4.2-4.6	304.6	7.5	7.1-7.9	120.8	2.2	2.0-2.4	387.8	4.0	3.8-4.2	282.2	6.9	6.6-7.3	105.6	1.9	1.7-2.1
	1981-1985	437.8	4.3	4.1-4.4	309.6	7.2	6.8-7.5	128.2	2.2	2.0-2.3	409.4	4.0	3.8-4.2	296.4	7.1	6.7-7.4	113.0	1.9	1.8-2.1
	1986-1990	482.8	4.6	4.4-4.7	340.8	7.5	7.1-7.9	142.0	2.3	2.1-2.5	448.6	4.2	4.1-4.4	328.4	7.3	7.0-7.7	120.2	2.0	1.8-2.1
	1991-1995	551.6	5.0	4.8-5.2	388.6	8.1	7.7-8.5	163.0	2.5	2.3-2.7	510.0	4.6	4.4-4.8	373.4	7.9	7.5-8.3	136.6	2.1	1.9-2.3
	1996-2000	594.6	5.1	5.0-5.3	430.0	8.6	8.2-9.0	164.6	2.4	2.3-2.6	536.4	4.6	4.5-4.8	394.2	8.0	7.6-8.3	142.2	2.1	1.9-2.2
	2001-2005	661.4	5.4	5.2-5.6	500.6	9.4	9.0-9.7	160.8	2.3	2.1-2.4	605.4	4.9	4.8-5.1	460.8	8.7	8.4-9.1	144.6	2.0	1.8-2.1
	2006-2010	746.6	5.7	5.5-5.9	577.4	9.9	9.6-10.3	169.2	2.3	2.1-2.5	613.0	4.7	4.5-4.8	480.6	8.4	8.0-8.7	132.4	1.8	1.6-1.9
	2011-2015	734.2	5.1	5.0-5.3	564.8	8.7	8.4-9.1	169.4	2.1	2.0-2.3	626.8	4.4	4.3-4.6	489.4	7.8	7.4-8.1	137.4	1.7	1.6-1.8
2016-2020	763.2	5.0	4.8-5.2	590.6	8.5	8.2-8.8	172.6	2.1	1.9-2.2	615.8	4.0	3.9-4.1	487.4	7.0	6.7-7.3	128.4	1.5	1.4-1.6	
Stomach	1976-1980	1,035.6	11.2	10.9-11.5	605.0	15.9	15.3-16.5	430.6	7.9	7.6-8.3	784.4	8.6	8.3-8.9	446.8	12.0	11.5-12.6	337.6	6.2	5.9-6.5
	1981-1985	1,055.4	10.6	10.3-10.9	643.6	15.9	15.3-16.5	411.8	7.0	6.7-7.3	746.2	7.6	7.3-7.8	443.8	11.2	10.7-11.7	302.4	5.2	4.9-5.4
	1986-1990	1,031.2	9.9	9.6-10.2	638.2	15.1	14.5-15.6	393.0	6.3	6.1-6.6	719.2	7.0	6.7-7.2	430.0	10.4	10.0-10.9	289.2	4.6	4.4-4.9
	1991-1995	1,019.6	9.3	9.0-9.5	625.0	13.9	13.4-14.4	394.6	6.0	5.7-6.3	666.0	6.1	5.9-6.3	404.4	9.3	8.9-9.7	261.6	3.9	3.7-4.2
	1996-2000	1,007.8	8.7	8.5-9.0	627.8	13.0	12.6-13.5	380.0	5.5	5.3-5.8	568.8	4.9	4.7-5.1	331.4	7.0	6.7-7.4	237.4	3.4	3.2-3.6
	2001-2005	966.4	7.9	7.7-8.1	591.2	11.4	11.0-11.8	375.2	5.3	5.1-5.6	508.2	4.1	4.0-4.3	295.0	5.8	5.5-6.2	213.2	2.9	2.7-3.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York State Excl New York City**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	913.6	7.0	6.8-7.3	577.6	10.2	9.8-10.6	336.0	4.6	4.4-4.8	429.4	3.3	3.1-3.4	258.0	4.6	4.4-4.9	171.4	2.3	2.1-2.4
	2011-2015	1,022.6	7.4	7.2-7.6	644.0	10.4	10.0-10.7	378.6	5.0	4.8-5.2	400.6	2.9	2.7-3.0	244.6	4.0	3.8-4.2	156.0	2.0	1.8-2.1
	2016-2020	1,047.6	7.2	7.0-7.4	641.0	9.5	9.2-9.9	406.6	5.2	5.0-5.5	370.8	2.5	2.4-2.6	218.4	3.3	3.1-3.5	152.4	1.9	1.7-2.0
Colorectal	1976-1980	6,462.2	68.9	68.1-69.6	3,153.4	81.7	80.4-83.1	3,308.8	60.4	59.4-61.3	3,206.4	35.0	34.4-35.5	1,539.8	41.9	41.0-42.9	1,666.6	30.6	29.9-31.2
	1981-1985	7,020.2	70.5	69.7-71.2	3,488.0	86.0	84.6-87.3	3,532.2	60.1	59.2-61.0	3,301.2	33.7	33.2-34.3	1,612.2	41.8	40.8-42.7	1,689.0	28.7	28.1-29.3
	1986-1990	6,863.4	65.6	64.9-66.3	3,401.4	80.0	78.8-81.3	3,462.0	56.0	55.2-56.9	3,174.2	30.7	30.2-31.2	1,585.2	39.0	38.1-39.9	1,589.0	25.5	24.9-26.1
	1991-1995	6,739.0	60.9	60.3-61.6	3,340.2	73.8	72.7-75.0	3,398.8	52.1	51.3-52.9	2,948.2	26.8	26.4-27.3	1,421.6	33.0	32.2-33.8	1,526.6	22.9	22.4-23.4
	1996-2000	7,192.6	62.0	61.3-62.6	3,581.6	74.2	73.1-75.4	3,611.0	53.1	52.4-53.9	2,689.2	23.1	22.7-23.5	1,328.6	28.6	27.9-29.3	1,360.6	19.3	18.8-19.7
	2001-2005	6,828.4	55.6	55.0-56.2	3,371.2	64.7	63.7-65.7	3,457.2	48.7	48.0-49.4	2,397.2	19.3	19.0-19.6	1,161.6	23.1	22.5-23.7	1,235.6	16.6	16.2-17.0
	2006-2010	5,994.2	46.0	45.5-46.6	2,966.2	52.3	51.5-53.2	3,028.0	41.0	40.4-41.7	2,050.0	15.5	15.2-15.8	1,001.6	18.3	17.8-18.8	1,048.4	13.4	13.0-13.8
	2011-2015	5,487.2	39.8	39.3-40.2	2,791.4	45.2	44.4-45.9	2,695.8	35.4	34.7-36.0	1,892.0	13.3	13.0-13.5	957.4	15.7	15.3-16.2	934.6	11.3	11.0-11.7
	2016-2020	5,366.4	37.2	36.8-37.7	2,778.8	42.1	41.4-42.9	2,587.6	33.1	32.5-33.7	1,768.4	11.7	11.5-12.0	900.6	13.7	13.3-14.1	867.8	10.1	9.8-10.5
Colon excluding rectum	1976-1980	4,550.0	48.9	48.3-49.6	2,108.6	55.7	54.6-56.9	2,441.4	44.6	43.8-45.4	2,589.0	28.2	27.7-28.7	1,202.0	32.8	31.9-33.7	1,387.0	25.4	24.8-26.0
	1981-1985	5,004.8	50.6	49.9-51.2	2,359.2	59.1	57.9-60.2	2,645.6	45.1	44.3-45.8	2,764.4	28.3	27.8-28.7	1,323.4	34.3	33.5-35.2	1,441.0	24.5	23.9-25.0
	1986-1990	4,895.6	47.0	46.4-47.6	2,316.6	55.4	54.3-56.5	2,579.0	41.6	40.9-42.3	2,711.0	26.2	25.8-26.7	1,338.2	33.0	32.1-33.8	1,372.8	22.0	21.5-22.5
	1991-1995	4,874.6	44.1	43.6-44.7	2,310.4	51.7	50.7-52.7	2,564.2	39.1	38.4-39.8	2,509.2	22.8	22.4-23.2	1,187.8	27.7	27.0-28.4	1,321.4	19.8	19.3-20.3
	1996-2000	5,226.0	45.0	44.4-45.5	2,481.6	52.0	51.1-52.9	2,744.4	40.1	39.4-40.8	2,280.0	19.6	19.2-19.9	1,105.6	23.9	23.3-24.5	1,174.4	16.6	16.2-17.0
	2001-2005	4,958.8	40.3	39.8-40.8	2,344.8	45.4	44.6-46.3	2,614.0	36.5	35.9-37.1	2,001.8	16.1	15.8-16.4	948.0	19.0	18.4-19.5	1,053.8	14.1	13.7-14.5
	2006-2010	4,345.8	33.3	32.9-33.8	2,051.8	36.6	35.9-37.3	2,294.0	30.8	30.2-31.3	1,691.4	12.8	12.5-13.0	806.6	14.8	14.3-15.2	884.8	11.3	10.9-11.6
	2011-2015	3,895.8	28.2	27.8-28.6	1,873.8	30.7	30.0-31.3	2,022.0	26.2	25.6-26.7	1,544.6	10.8	10.6-11.1	755.2	12.5	12.1-12.9	789.4	9.5	9.2-9.8
	2016-2020	3,801.2	26.2	25.8-26.6	1,862.8	28.4	27.8-29.0	1,938.4	24.4	23.9-24.9	1,379.8	9.1	8.9-9.4	679.6	10.4	10.0-10.7	700.2	8.1	7.8-8.4
Rectum & rectosigmoid	1976-1980	1,912.2	20.0	19.6-20.4	1,044.8	26.0	25.3-26.8	867.4	15.8	15.3-16.2	617.4	6.7	6.5-7.0	337.8	9.2	8.7-9.6	279.6	5.1	4.9-5.4
	1981-1985	2,015.4	19.9	19.5-20.3	1,128.8	26.9	26.2-27.6	886.6	15.1	14.6-15.5	536.8	5.5	5.3-5.7	288.8	7.4	7.0-7.9	248.0	4.2	4.0-4.4
	1986-1990	1,967.8	18.6	18.3-19.0	1,084.8	24.6	23.9-25.3	883.0	14.4	14.0-14.8	463.2	4.5	4.3-4.7	247.0	6.0	5.7-6.4	216.2	3.5	3.3-3.7
	1991-1995	1,864.4	16.8	16.5-17.1	1,029.8	22.1	21.5-22.8	834.6	13.0	12.6-13.4	439.0	4.0	3.8-4.2	233.8	5.3	5.0-5.6	205.2	3.1	2.9-3.3
	1996-2000	1,966.6	17.0	16.7-17.4	1,100.0	22.2	21.6-22.8	866.6	13.0	12.7-13.4	409.2	3.5	3.4-3.7	223.0	4.7	4.4-5.0	186.2	2.7	2.5-2.9
	2001-2005	1,869.6	15.3	15.0-15.6	1,026.4	19.2	18.7-19.8	843.2	12.2	11.8-12.6	395.4	3.2	3.1-3.3	213.6	4.2	3.9-4.4	181.8	2.5	2.3-2.6
	2006-2010	1,648.4	12.7	12.4-13.0	914.4	15.8	15.3-16.2	734.0	10.3	9.9-10.6	358.6	2.7	2.6-2.9	195.0	3.5	3.3-3.7	163.6	2.1	2.0-2.3
	2011-2015	1,591.4	11.6	11.3-11.9	917.6	14.5	14.1-14.9	673.8	9.2	8.9-9.5	347.4	2.5	2.3-2.6	202.2	3.3	3.1-3.5	145.2	1.8	1.7-1.9
	2016-2020	1,565.2	11.0	10.8-11.3	916.0	13.8	13.4-14.2	649.2	8.6	8.3-9.0	388.6	2.6	2.5-2.7	221.0	3.3	3.1-3.5	167.6	2.0	1.9-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York State Excl New York City**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	278.8	2.9	2.7-3.0	167.8	4.1	3.8-4.4	111.0	2.0	1.8-2.2	229.8	2.4	2.2-2.5	134.2	3.3	3.0-3.6	95.6	1.7	1.6-1.9
	1981-1985	288.2	2.9	2.7-3.0	179.6	4.3	4.0-4.6	108.6	1.9	1.7-2.0	254.8	2.5	2.4-2.7	157.8	3.9	3.6-4.2	97.0	1.7	1.5-1.8
	1986-1990	341.4	3.2	3.1-3.4	219.2	4.9	4.6-5.3	122.2	2.0	1.8-2.2	305.2	2.9	2.8-3.1	191.2	4.4	4.1-4.7	114.0	1.9	1.7-2.0
	1991-1995	438.0	4.0	3.8-4.1	275.2	6.0	5.7-6.3	162.8	2.5	2.3-2.7	396.4	3.6	3.4-3.7	241.2	5.3	5.0-5.6	155.2	2.4	2.2-2.5
	1996-2000	527.0	4.6	4.4-4.8	343.4	6.9	6.6-7.3	183.6	2.7	2.5-2.9	438.4	3.8	3.6-4.0	266.8	5.5	5.2-5.8	171.6	2.5	2.3-2.7
	2001-2005	641.6	5.3	5.1-5.4	447.8	8.3	7.9-8.6	193.8	2.8	2.6-3.0	508.6	4.1	4.0-4.3	329.6	6.2	5.9-6.5	179.0	2.5	2.3-2.7
	2006-2010	824.8	6.3	6.1-6.5	599.0	10.0	9.6-10.3	225.8	3.1	2.9-3.3	627.6	4.8	4.6-4.9	408.2	7.0	6.7-7.3	219.4	2.9	2.8-3.1
	2011-2015	1,038.4	7.2	7.0-7.4	750.2	11.1	10.8-11.5	288.2	3.8	3.6-4.0	765.8	5.3	5.2-5.5	510.8	7.8	7.5-8.2	255.0	3.2	3.0-3.4
	2016-2020	1,171.0	7.5	7.3-7.7	828.4	11.5	11.1-11.8	342.6	4.2	4.0-4.4	785.6	5.1	4.9-5.2	527.2	7.5	7.2-7.8	258.4	3.1	2.9-3.3
Pancreas	1976-1980	1,200.2	12.7	12.4-13.1	625.4	16.0	15.4-16.6	574.8	10.5	10.1-10.9	1,145.2	12.2	11.8-12.5	593.4	15.3	14.7-15.8	551.8	10.0	9.6-10.4
	1981-1985	1,216.6	12.2	11.8-12.5	597.0	14.6	14.0-15.1	619.6	10.5	10.2-10.9	1,162.4	11.6	11.3-11.9	572.0	14.0	13.5-14.6	590.4	10.0	9.6-10.4
	1986-1990	1,279.6	12.2	11.9-12.5	617.0	14.5	13.9-15.0	662.6	10.7	10.3-11.0	1,205.8	11.5	11.2-11.8	586.0	13.8	13.3-14.3	619.8	9.9	9.6-10.3
	1991-1995	1,379.6	12.5	12.2-12.8	663.8	14.5	14.0-15.0	715.8	10.9	10.5-11.3	1,274.6	11.5	11.3-11.8	606.2	13.4	12.9-13.9	668.4	10.1	9.8-10.5
	1996-2000	1,457.6	12.6	12.3-12.9	698.6	14.4	13.9-14.9	759.0	11.1	10.8-11.5	1,349.0	11.6	11.3-11.9	650.8	13.6	13.2-14.1	698.2	10.1	9.8-10.5
	2001-2005	1,606.6	13.0	12.8-13.3	767.4	14.6	14.2-15.1	839.2	11.7	11.4-12.1	1,410.6	11.4	11.2-11.7	668.4	12.9	12.5-13.4	742.2	10.2	9.9-10.6
	2006-2010	1,785.4	13.6	13.3-13.9	876.2	15.4	15.0-15.9	909.2	12.1	11.8-12.5	1,494.2	11.3	11.0-11.6	729.4	12.9	12.5-13.3	764.8	10.0	9.7-10.4
	2011-2015	2,020.8	14.3	14.0-14.6	1,018.8	16.2	15.8-16.7	1,002.0	12.6	12.3-13.0	1,607.4	11.3	11.0-11.6	807.8	13.0	12.6-13.5	799.6	9.8	9.5-10.2
	2016-2020	2,290.6	15.1	14.8-15.4	1,180.0	17.3	16.8-17.8	1,110.6	13.3	12.9-13.7	1,742.6	11.4	11.1-11.6	882.6	13.0	12.6-13.4	860.0	10.0	9.7-10.3
Larynx	1976-1980	581.0	5.7	5.5-5.9	477.8	10.9	10.4-11.3	103.2	1.8	1.7-2.0	198.6	2.0	1.9-2.1	161.8	3.9	3.6-4.2	36.8	0.7	0.6-0.8
	1981-1985	623.4	6.0	5.8-6.2	499.6	11.0	10.5-11.4	123.8	2.2	2.0-2.4	196.8	1.9	1.8-2.0	156.2	3.6	3.3-3.9	40.6	0.7	0.6-0.8
	1986-1990	624.0	5.8	5.6-6.0	481.4	10.3	9.9-10.8	142.6	2.4	2.2-2.6	184.0	1.7	1.6-1.8	142.6	3.2	3.0-3.5	41.4	0.7	0.6-0.8
	1991-1995	586.2	5.3	5.1-5.5	449.2	9.3	8.9-9.7	137.0	2.2	2.1-2.4	191.6	1.7	1.6-1.8	148.2	3.2	3.0-3.4	43.4	0.7	0.6-0.8
	1996-2000	537.6	4.7	4.5-4.9	414.4	8.1	7.8-8.5	123.2	2.0	1.8-2.1	166.8	1.4	1.3-1.5	127.0	2.6	2.4-2.8	39.8	0.6	0.5-0.7
	2001-2005	511.0	4.2	4.0-4.4	401.4	7.3	7.0-7.7	109.6	1.7	1.5-1.8	155.6	1.3	1.2-1.4	120.0	2.3	2.1-2.5	35.6	0.5	0.4-0.6
	2006-2010	510.6	3.9	3.8-4.1	397.4	6.8	6.5-7.1	113.2	1.6	1.5-1.7	145.6	1.1	1.0-1.2	111.8	2.0	1.8-2.1	33.8	0.5	0.4-0.5
	2011-2015	500.0	3.5	3.3-3.6	397.0	6.0	5.8-6.3	103.0	1.4	1.2-1.5	148.2	1.0	1.0-1.1	117.0	1.8	1.7-2.0	31.2	0.4	0.3-0.5
	2016-2020	462.4	3.0	2.9-3.1	364.6	5.1	4.9-5.4	97.8	1.2	1.1-1.4	125.8	0.8	0.7-0.9	98.8	1.4	1.3-1.5	27.0	0.3	0.3-0.4
Lung and bronchus	1976-1980	5,882.0	59.5	58.8-60.2	4,185.0	100.9	99.5-102.3	1,697.0	30.7	30.0-31.4	4,779.4	48.6	48.0-49.2	3,456.0	84.7	83.4-86.1	1,323.4	23.8	23.2-24.4
	1981-1985	6,795.2	65.7	65.0-66.4	4,496.6	103.9	102.5-105.3	2,298.6	39.7	39.0-40.5	5,513.0	53.5	52.9-54.2	3,723.2	87.6	86.3-88.9	1,789.8	30.7	30.1-31.3
	1986-1990	7,499.8	70.1	69.4-70.9	4,630.4	103.4	102.0-104.8	2,869.4	47.7	46.9-48.5	6,121.4	57.3	56.7-58.0	3,849.8	87.2	85.9-88.5	2,271.6	37.5	36.8-38.2
	1991-1995	8,067.0	72.3	71.6-73.1	4,618.2	98.2	96.9-99.6	3,448.8	54.8	54.0-55.7	6,568.2	58.9	58.3-59.6	3,868.8	83.8	82.6-85.0	2,699.4	42.4	41.6-43.1
	1996-2000	8,552.4	73.8	73.1-74.5	4,659.2	94.2	93.0-95.4	3,893.2	59.8	58.9-60.6	6,488.4	55.9	55.3-56.5	3,633.6	74.5	73.4-75.6	2,854.8	43.1	42.4-43.8
	2001-2005	8,858.0	72.5	71.8-73.2	4,600.8	87.5	86.4-88.7	4,257.2	62.3	61.4-63.1	6,481.4	52.9	52.3-53.5	3,451.8	66.7	65.7-67.7	3,029.6	43.5	42.8-44.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York State Excl New York City**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	9,324.6	72.0	71.4-72.7	4,675.2	82.6	81.5-83.7	4,649.4	64.8	64.0-65.7	6,347.8	48.8	48.2-49.3	3,302.8	58.9	58.0-59.8	3,045.0	41.7	41.0-42.4
	2011-2015	9,494.8	67.6	67.0-68.2	4,664.8	74.9	73.9-75.9	4,830.0	62.6	61.8-63.4	6,092.6	43.3	42.8-43.8	3,133.6	50.9	50.1-51.7	2,959.0	37.8	37.2-38.4
	2016-2020	9,631.8	62.7	62.2-63.3	4,579.0	66.3	65.4-67.2	5,052.8	60.5	59.7-61.2	5,250.4	34.1	33.7-34.5	2,649.4	39.0	38.3-39.7	2,601.0	30.5	30.0-31.1
Melanoma of the skin	1976-1980	698.6	7.3	7.1-7.6	371.0	8.6	8.2-9.0	327.6	6.4	6.1-6.7	239.2	2.5	2.4-2.7	143.4	3.4	3.1-3.7	95.8	1.9	1.7-2.0
	1981-1985	891.0	9.0	8.7-9.2	468.4	10.5	10.1-10.9	422.6	8.0	7.6-8.3	247.0	2.5	2.3-2.6	145.2	3.3	3.1-3.6	101.8	1.9	1.7-2.0
	1986-1990	947.8	9.1	8.9-9.4	520.6	11.3	10.8-11.7	427.2	7.7	7.3-8.0	298.4	2.9	2.7-3.0	183.8	4.1	3.8-4.4	114.6	2.0	1.9-2.2
	1991-1995	999.8	9.1	8.9-9.4	582.0	12.0	11.5-12.4	417.8	7.1	6.7-7.4	301.6	2.8	2.6-2.9	183.0	3.9	3.7-4.2	118.6	1.9	1.8-2.1
	1996-2000	1,373.2	12.1	11.8-12.4	789.2	15.6	15.1-16.1	584.0	9.6	9.3-10.0	303.8	2.6	2.5-2.8	192.2	3.9	3.6-4.1	111.6	1.7	1.6-1.9
	2001-2005	1,957.2	16.4	16.1-16.7	1,128.6	21.0	20.5-21.6	828.6	13.1	12.7-13.5	331.4	2.7	2.6-2.9	200.4	3.8	3.6-4.1	131.0	1.9	1.8-2.1
	2006-2010	2,644.2	21.1	20.8-21.5	1,527.6	26.9	26.3-27.5	1,116.6	17.1	16.7-17.6	346.6	2.7	2.6-2.8	223.0	4.0	3.7-4.2	123.6	1.7	1.6-1.9
	2011-2015	3,158.4	23.6	23.3-24.0	1,830.8	29.7	29.1-30.3	1,327.6	19.4	18.9-19.8	359.4	2.6	2.5-2.7	232.8	3.9	3.6-4.1	126.6	1.6	1.5-1.8
	2016-2020	3,302.4	23.3	22.9-23.6	1,922.4	29.0	28.4-29.6	1,380.0	19.2	18.7-19.7	303.4	2.0	1.9-2.1	193.8	3.0	2.8-3.2	109.6	1.3	1.2-1.4
Female breast	1976-1980							5,420.8	103.3	102.0-104.5							2,066.8	38.5	37.8-39.3
	1981-1985							6,119.2	111.6	110.3-112.9							2,165.8	38.3	37.6-39.1
	1986-1990							7,167.6	125.7	124.4-127.1							2,275.2	38.6	37.9-39.4
	1991-1995							7,672.6	128.8	127.5-130.1							2,141.8	34.6	33.9-35.2
	1996-2000							8,828.6	142.1	140.7-143.4							1,960.8	29.9	29.3-30.5
	2001-2005							8,863.6	135.1	133.8-136.3							1,767.4	25.4	24.8-25.9
	2006-2010							9,284.0	135.4	134.1-136.6							1,623.8	22.1	21.6-22.6
	2011-2015							9,922.8	138.9	137.6-140.2							1,526.8	19.6	19.1-20.0
	2016-2020							10,258.0	139.3	138.0-140.5							1,498.4	18.5	18.0-18.9
Cervix uteri	1976-1980							621.4	12.0	11.6-12.5							241.2	4.6	4.3-4.9
	1981-1985							534.6	10.0	9.7-10.4							202.0	3.7	3.4-3.9
	1986-1990							553.0	10.1	9.8-10.5							190.0	3.3	3.1-3.6
	1991-1995							563.8	9.8	9.4-10.2							164.6	2.8	2.6-3.0
	1996-2000							528.0	9.0	8.6-9.3							159.6	2.6	2.4-2.8
	2001-2005							455.6	7.6	7.2-7.9							142.0	2.2	2.0-2.4
	2006-2010							451.2	7.5	7.1-7.8							127.4	1.9	1.7-2.0
	2011-2015							418.2	6.8	6.5-7.1							137.6	2.0	1.8-2.2
	2016-2020							393.2	6.6	6.3-6.9							122.4	1.7	1.6-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York State Excl New York City**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							1,431.4	26.0	25.4-26.6							296.6	5.3	5.1-5.6
	1981-1985							1,445.8	25.3	24.7-25.9							303.6	5.1	4.9-5.4
	1986-1990							1,419.0	24.1	23.5-24.7							284.0	4.6	4.3-4.8
	1991-1995							1,544.0	25.6	25.0-26.1							305.4	4.7	4.4-4.9
	1996-2000							1,741.8	27.8	27.3-28.4							293.4	4.3	4.1-4.6
	2001-2005							1,956.0	29.5	29.0-30.1							320.6	4.5	4.3-4.8
	2006-2010							2,127.8	30.3	29.7-30.8							350.0	4.7	4.5-4.9
	2011-2015							2,428.8	32.1	31.5-32.7							398.0	5.0	4.8-5.3
2016-2020							2,513.8	31.6	31.0-32.1							456.0	5.4	5.2-5.6	
Ovary	1976-1980							904.2	16.8	16.3-17.3							610.6	11.1	10.7-11.5
	1981-1985							965.8	17.4	16.9-17.9							627.2	10.8	10.5-11.2
	1986-1990							1,000.0	17.4	16.9-17.9							636.8	10.6	10.2-11.0
	1991-1995							1,060.2	17.6	17.1-18.1							663.0	10.6	10.2-10.9
	1996-2000							1,044.6	16.7	16.2-17.2							637.2	9.7	9.3-10.0
	2001-2005							1,004.4	15.2	14.8-15.6							664.6	9.5	9.2-9.9
	2006-2010							960.4	13.8	13.4-14.2							627.4	8.6	8.3-8.9
	2011-2015							927.8	12.8	12.4-13.2							595.4	7.6	7.3-7.9
2016-2020							841.4	11.3	10.9-11.6							534.4	6.4	6.2-6.7	
Prostate	1976-1980				2,854.8	82.3	80.9-83.7							1,031.2	32.8	31.9-33.7			
	1981-1985				3,483.0	92.6	91.2-94.1							1,134.8	33.7	32.8-34.7			
	1986-1990				4,397.0	107.9	106.4-109.4							1,294.4	35.8	34.9-36.7			
	1991-1995				7,450.2	159.9	158.2-161.5							1,438.8	37.3	36.4-38.2			
	1996-2000				8,601.8	172.1	170.4-173.7							1,328.0	31.4	30.6-32.2			
	2001-2005				9,483.8	176.7	175.1-178.4							1,177.6	25.4	24.8-26.1			
	2006-2010				10,223.8	172.1	170.6-173.6							1,055.6	20.4	19.8-20.9			
	2011-2015				8,821.0	131.3	130.1-132.6							987.8	17.2	16.7-17.6			
2016-2020				9,885.0	133.4	132.2-134.6							1,037.8	16.2	15.8-16.7				
Testis	1976-1980				204.0	4.0	3.8-4.3							31.6	0.6	0.5-0.7			
	1981-1985				253.2	4.8	4.5-5.1							23.0	0.5	0.4-0.5			
	1986-1990				273.8	4.9	4.7-5.2							20.2	0.4	0.3-0.5			
	1991-1995				300.4	5.4	5.1-5.7							17.2	0.3	0.3-0.4			
	1996-2000				325.0	6.0	5.7-6.3							14.4	0.3	0.2-0.3			
	2001-2005				334.8	6.4	6.1-6.7							13.6	0.3	0.2-0.3			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020  
New York State Excl New York City**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				352.4	6.8	6.5-7.1							10.8	0.2	0.1-0.3			
	2011-2015				359.4	6.9	6.6-7.3							11.4	0.2	0.2-0.3			
	2016-2020				353.8	6.9	6.6-7.3							15.0	0.3	0.2-0.3			
Urinary bladder (incl. in situ)	1976-1980	2,113.8	22.5	22.0-22.9	1,555.0	40.7	39.8-41.7	558.8	10.2	9.9-10.6	598.4	6.7	6.5-7.0	417.2	12.3	11.7-12.9	181.2	3.4	3.2-3.6
	1981-1985	2,333.4	23.2	22.8-23.6	1,709.4	42.1	41.1-43.0	624.0	10.7	10.3-11.1	594.2	6.2	6.0-6.4	416.0	11.7	11.2-12.3	178.2	3.0	2.8-3.2
	1986-1990	2,464.2	23.3	22.9-23.7	1,787.2	41.6	40.7-42.5	677.0	11.0	10.6-11.4	570.0	5.6	5.4-5.8	388.2	10.4	9.9-10.9	181.8	2.9	2.7-3.1
	1991-1995	2,598.4	23.4	23.0-23.8	1,855.6	40.8	40.0-41.7	742.8	11.5	11.1-11.9	578.6	5.3	5.1-5.5	379.6	9.4	8.9-9.8	199.0	2.9	2.7-3.1
	1996-2000	3,131.4	27.0	26.6-27.4	2,271.8	47.2	46.3-48.1	859.6	12.8	12.4-13.2	619.6	5.3	5.1-5.5	421.8	9.6	9.1-10.0	197.8	2.7	2.6-2.9
	2001-2005	3,467.8	28.3	27.9-28.7	2,560.6	49.6	48.8-50.5	907.2	12.9	12.5-13.3	614.2	4.9	4.7-5.1	417.2	8.6	8.3-9.0	197.0	2.6	2.4-2.8
	2006-2010	3,609.6	27.7	27.3-28.1	2,693.8	48.4	47.6-49.2	915.8	12.4	12.0-12.8	660.4	4.9	4.8-5.1	465.2	8.8	8.4-9.1	195.2	2.4	2.3-2.6
	2011-2015	3,864.4	27.6	27.2-28.0	2,878.2	47.4	46.6-48.2	986.2	12.5	12.2-12.9	676.2	4.7	4.5-4.8	470.6	8.0	7.7-8.3	205.6	2.4	2.3-2.6
2016-2020	3,819.0	25.1	24.7-25.5	2,868.6	42.8	42.1-43.5	950.4	11.2	10.9-11.6	663.6	4.3	4.1-4.4	473.6	7.4	7.1-7.7	190.0	2.1	2.0-2.2	
Kidney and renal pelvis	1976-1980	735.2	7.6	7.3-7.8	451.0	10.8	10.3-11.2	284.2	5.2	4.9-5.5	397.2	4.1	3.9-4.3	243.6	6.0	5.6-6.3	153.6	2.8	2.6-3.0
	1981-1985	918.0	9.1	8.8-9.3	560.8	12.9	12.4-13.4	357.2	6.2	5.9-6.5	415.8	4.1	4.0-4.3	249.2	5.9	5.6-6.3	166.6	2.9	2.7-3.1
	1986-1990	1,061.0	10.1	9.8-10.4	646.4	14.4	13.9-14.9	414.6	7.0	6.7-7.3	461.2	4.4	4.2-4.6	275.2	6.3	6.0-6.7	186.0	3.0	2.8-3.2
	1991-1995	1,243.4	11.3	11.0-11.6	743.2	15.5	15.0-16.0	500.2	8.1	7.8-8.4	466.2	4.2	4.1-4.4	279.8	6.2	5.9-6.5	186.4	2.9	2.7-3.1
	1996-2000	1,516.8	13.2	12.9-13.5	929.2	18.4	17.9-19.0	587.6	9.2	8.8-9.5	466.8	4.0	3.9-4.2	283.8	5.9	5.6-6.2	183.0	2.7	2.5-2.9
	2001-2005	1,844.2	15.2	14.9-15.5	1,140.4	21.1	20.5-21.6	703.8	10.5	10.2-10.9	487.4	4.0	3.8-4.1	295.6	5.7	5.4-6.0	191.8	2.7	2.5-2.9
	2006-2010	2,248.2	17.5	17.2-17.8	1,411.2	24.1	23.6-24.7	837.0	12.0	11.7-12.4	486.4	3.7	3.5-3.8	308.0	5.5	5.2-5.7	178.4	2.4	2.2-2.5
	2011-2015	2,485.8	18.1	17.8-18.5	1,605.2	25.3	24.7-25.8	880.6	12.0	11.6-12.4	494.6	3.5	3.3-3.6	317.4	5.1	4.9-5.4	177.2	2.2	2.0-2.3
2016-2020	2,655.0	18.4	18.1-18.7	1,731.2	25.6	25.0-26.1	923.8	12.2	11.8-12.5	442.4	2.9	2.8-3.0	281.0	4.2	3.9-4.4	161.4	1.9	1.8-2.0	
Brain and other central nervous system	1976-1980	613.0	6.0	5.8-6.2	347.0	7.4	7.0-7.7	266.0	4.9	4.6-5.2	432.2	4.2	4.0-4.4	243.6	5.2	4.9-5.6	188.6	3.4	3.2-3.6
	1981-1985	675.6	6.5	6.3-6.8	358.2	7.6	7.3-8.0	317.4	5.7	5.4-6.0	436.6	4.2	4.0-4.4	232.0	5.1	4.8-5.4	204.6	3.6	3.3-3.8
	1986-1990	753.2	7.1	6.9-7.3	413.2	8.5	8.2-8.9	340.0	5.9	5.6-6.2	468.0	4.4	4.2-4.6	258.0	5.4	5.1-5.8	210.0	3.5	3.3-3.7
	1991-1995	819.0	7.5	7.2-7.7	438.6	8.8	8.4-9.2	380.4	6.4	6.1-6.7	494.4	4.5	4.3-4.7	269.2	5.5	5.2-5.8	225.2	3.7	3.5-3.9
	1996-2000	869.0	7.7	7.5-8.0	474.6	9.2	8.8-9.5	394.4	6.5	6.2-6.8	504.6	4.4	4.3-4.6	279.8	5.5	5.2-5.8	224.8	3.5	3.3-3.8
	2001-2005	881.4	7.6	7.4-7.8	484.2	9.0	8.6-9.4	397.2	6.4	6.1-6.7	514.2	4.3	4.1-4.5	280.4	5.2	4.9-5.5	233.8	3.6	3.4-3.8
	2006-2010	871.8	7.2	7.0-7.4	481.0	8.6	8.2-8.9	390.8	6.1	5.8-6.4	523.4	4.1	4.0-4.3	302.0	5.3	5.0-5.6	221.4	3.2	3.0-3.4
	2011-2015	910.8	7.2	7.0-7.5	503.0	8.4	8.1-8.8	407.8	6.2	5.9-6.5	568.2	4.2	4.0-4.3	311.6	5.0	4.7-5.3	256.6	3.5	3.3-3.7
2016-2020	891.2	6.9	6.6-7.1	499.8	8.1	7.7-8.4	391.4	5.8	5.5-6.1	610.0	4.3	4.1-4.4	349.6	5.3	5.0-5.5	260.4	3.4	3.2-3.6	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York State Excl New York City**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	346.6	3.6	3.4-3.8	105.8	2.4	2.2-2.6	240.8	4.6	4.4-4.9	60.4	0.7	0.6-0.7	22.4	0.6	0.5-0.7	38.0	0.7	0.6-0.8
	1981-1985	383.6	3.8	3.6-4.0	108.4	2.3	2.1-2.6	275.2	5.1	4.9-5.4	54.4	0.6	0.5-0.6	21.8	0.5	0.4-0.7	32.6	0.6	0.5-0.7
	1986-1990	427.0	4.1	3.9-4.3	123.4	2.6	2.4-2.8	303.6	5.5	5.3-5.8	53.2	0.5	0.5-0.6	19.8	0.5	0.4-0.6	33.4	0.5	0.5-0.6
	1991-1995	564.4	5.2	5.0-5.4	153.8	3.1	2.8-3.3	410.6	7.2	6.9-7.6	54.6	0.5	0.4-0.6	20.8	0.5	0.4-0.6	33.8	0.5	0.4-0.6
	1996-2000	881.4	7.9	7.7-8.2	227.8	4.3	4.1-4.6	653.6	11.4	11.0-11.8	62.6	0.5	0.5-0.6	22.4	0.5	0.4-0.6	40.2	0.6	0.5-0.7
	2001-2005	1,282.6	11.2	10.9-11.5	314.0	5.6	5.4-5.9	968.6	16.6	16.1-17.0	54.6	0.4	0.4-0.5	21.8	0.4	0.3-0.5	32.8	0.5	0.4-0.5
	2006-2010	1,930.2	16.5	16.1-16.8	482.4	8.4	8.1-8.8	1,447.8	24.3	23.7-24.8	64.0	0.5	0.4-0.5	28.0	0.5	0.4-0.6	36.0	0.5	0.4-0.5
	2011-2015	2,349.2	19.7	19.3-20.1	625.8	10.4	10.0-10.8	1,723.4	28.8	28.2-29.4	73.0	0.5	0.5-0.6	33.0	0.5	0.5-0.6	40.0	0.5	0.4-0.6
2016-2020	2,087.2	17.5	17.2-17.9	594.0	9.8	9.4-10.1	1,493.2	25.3	24.7-25.9	77.0	0.5	0.5-0.6	34.4	0.5	0.5-0.6	42.6	0.5	0.4-0.6	
Hodgkin lymphoma	1976-1980	378.2	3.5	3.4-3.7	208.6	4.1	3.9-4.4	169.6	2.9	2.7-3.2	130.2	1.3	1.2-1.4	73.6	1.6	1.5-1.8	56.6	1.0	0.9-1.1
	1981-1985	397.6	3.6	3.5-3.8	208.8	4.1	3.8-4.4	188.8	3.2	3.0-3.4	108.8	1.0	1.0-1.1	63.0	1.3	1.2-1.5	45.8	0.8	0.7-0.9
	1986-1990	385.0	3.5	3.3-3.6	204.6	3.9	3.6-4.1	180.4	3.1	2.9-3.3	101.0	1.0	0.9-1.0	61.4	1.3	1.1-1.4	39.6	0.7	0.6-0.8
	1991-1995	386.8	3.5	3.4-3.7	202.8	3.8	3.6-4.1	184.0	3.2	3.0-3.4	79.8	0.7	0.7-0.8	41.6	0.8	0.7-1.0	38.2	0.6	0.5-0.7
	1996-2000	369.6	3.4	3.2-3.6	197.6	3.7	3.5-4.0	172.0	3.1	2.9-3.3	60.0	0.5	0.5-0.6	31.0	0.6	0.5-0.7	29.0	0.5	0.4-0.6
	2001-2005	397.2	3.6	3.4-3.8	215.4	4.0	3.8-4.2	181.8	3.3	3.0-3.5	57.6	0.5	0.4-0.6	29.0	0.5	0.5-0.6	28.6	0.5	0.4-0.5
	2006-2010	381.8	3.4	3.2-3.5	212.4	3.8	3.6-4.1	169.4	3.0	2.8-3.2	52.6	0.4	0.4-0.5	27.4	0.5	0.4-0.6	25.2	0.4	0.3-0.5
	2011-2015	381.0	3.3	3.2-3.5	210.8	3.8	3.5-4.0	170.2	3.0	2.8-3.2	39.8	0.3	0.3-0.3	20.2	0.3	0.3-0.4	19.6	0.3	0.2-0.3
2016-2020	368.6	3.2	3.0-3.3	199.2	3.5	3.2-3.7	169.4	2.9	2.7-3.1	39.6	0.3	0.2-0.3	24.6	0.4	0.3-0.5	15.0	0.2	0.2-0.2	
Non-Hodgkin lymphomas	1976-1980	1,179.8	12.2	11.8-12.5	611.4	14.4	13.9-15.0	568.4	10.4	10.0-10.8	636.6	6.6	6.4-6.9	325.8	8.0	7.6-8.4	310.8	5.7	5.4-5.9
	1981-1985	1,399.0	13.8	13.5-14.1	720.6	16.6	16.0-17.1	678.4	11.7	11.3-12.1	703.8	7.0	6.8-7.3	370.8	8.9	8.5-9.3	333.0	5.7	5.4-6.0
	1986-1990	1,712.6	16.3	16.0-16.7	906.2	19.9	19.3-20.5	806.4	13.5	13.1-13.9	838.6	8.1	7.8-8.3	436.2	10.1	9.7-10.6	402.4	6.6	6.3-6.9
	1991-1995	1,981.8	18.1	17.7-18.4	1,051.4	22.1	21.5-22.7	930.4	14.9	14.5-15.4	953.6	8.7	8.4-8.9	490.0	10.9	10.4-11.3	463.6	7.1	6.8-7.4
	1996-2000	2,375.2	20.7	20.4-21.1	1,220.4	24.4	23.8-25.0	1,154.8	17.9	17.4-18.3	1,009.8	8.7	8.5-9.0	504.6	10.5	10.1-11.0	505.2	7.4	7.1-7.7
	2001-2005	2,597.6	21.5	21.1-21.9	1,363.6	25.7	25.1-26.3	1,234.0	18.3	17.8-18.7	870.4	7.1	6.9-7.3	454.0	8.9	8.6-9.3	416.4	5.7	5.5-6.0
	2006-2010	2,938.0	23.0	22.6-23.4	1,562.0	27.5	26.9-28.2	1,376.0	19.4	19.0-19.9	844.2	6.4	6.2-6.6	450.0	8.2	7.9-8.5	394.2	5.2	4.9-5.4
	2011-2015	3,094.2	22.7	22.4-23.1	1,694.6	27.7	27.1-28.4	1,399.6	18.8	18.3-19.2	798.0	5.7	5.5-5.9	436.8	7.4	7.1-7.7	361.2	4.4	4.2-4.6
2016-2020	3,160.0	21.9	21.5-22.3	1,710.6	26.0	25.4-26.6	1,449.4	18.6	18.2-19.1	765.8	5.1	4.9-5.2	423.8	6.5	6.3-6.8	342.0	4.0	3.8-4.2	
Myeloma	1976-1980	431.2	4.5	4.3-4.7	208.4	5.4	5.0-5.7	222.8	4.0	3.8-4.3	285.2	3.0	2.9-3.2	143.6	3.7	3.5-4.0	141.6	2.6	2.4-2.8
	1981-1985	502.4	5.0	4.8-5.2	259.2	6.4	6.0-6.7	243.2	4.1	3.9-4.4	337.6	3.4	3.2-3.5	168.8	4.2	3.9-4.6	168.8	2.8	2.6-3.0
	1986-1990	550.6	5.3	5.1-5.5	279.4	6.6	6.2-6.9	271.2	4.4	4.2-4.7	367.6	3.5	3.4-3.7	188.8	4.6	4.3-4.9	178.8	2.9	2.7-3.1
	1991-1995	621.2	5.6	5.4-5.8	318.2	7.1	6.7-7.4	303.0	4.7	4.4-4.9	421.8	3.8	3.7-4.0	206.0	4.7	4.4-5.0	215.8	3.3	3.1-3.5
	1996-2000	716.8	6.2	6.0-6.4	367.6	7.6	7.2-7.9	349.2	5.2	5.0-5.5	433.4	3.7	3.6-3.9	223.0	4.8	4.5-5.1	210.4	3.0	2.9-3.2
2001-2005	786.2	6.4	6.2-6.6	430.4	8.2	7.9-8.6	355.8	5.1	4.8-5.3	415.0	3.4	3.2-3.5	223.4	4.4	4.2-4.7	191.6	2.6	2.5-2.8	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020  
New York State Excl New York City**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	929.6	7.1	6.9-7.3	500.0	8.8	8.4-9.1	429.6	5.9	5.6-6.1	424.2	3.2	3.1-3.4	224.2	4.1	3.8-4.3	200.0	2.6	2.5-2.8
	2011-2015	1,093.8	7.8	7.6-8.0	608.0	9.7	9.4-10.1	485.8	6.3	6.0-6.5	432.2	3.0	2.9-3.2	228.6	3.8	3.5-4.0	203.6	2.5	2.3-2.7
	2016-2020	1,138.2	7.6	7.4-7.8	643.2	9.5	9.1-9.8	495.0	6.1	5.9-6.4	410.4	2.7	2.6-2.8	216.2	3.3	3.1-3.5	194.2	2.2	2.1-2.4
Leukemias	1976-1980	1,249.0	13.2	12.8-13.5	696.6	17.5	16.9-18.1	552.4	10.2	9.9-10.6	798.4	8.5	8.2-8.8	433.0	11.1	10.6-11.6	365.4	6.8	6.5-7.1
	1981-1985	1,293.2	13.0	12.7-13.3	723.2	17.3	16.7-17.9	570.0	10.0	9.6-10.4	823.2	8.4	8.1-8.6	445.6	11.2	10.7-11.7	377.6	6.6	6.3-6.9
	1986-1990	1,323.2	12.7	12.4-13.0	732.2	16.8	16.2-17.3	591.0	9.9	9.6-10.3	831.8	8.1	7.8-8.3	457.6	10.8	10.4-11.3	374.2	6.2	5.9-6.5
	1991-1995	1,469.6	13.4	13.1-13.7	819.2	17.7	17.2-18.3	650.4	10.5	10.1-10.8	853.2	7.8	7.6-8.0	468.0	10.7	10.2-11.1	385.2	6.0	5.7-6.3
	1996-2000	1,682.4	14.7	14.4-15.0	949.2	19.3	18.8-19.9	733.2	11.2	10.9-11.6	921.6	8.0	7.7-8.2	507.0	10.7	10.2-11.1	414.6	6.1	5.9-6.4
	2001-2005	1,893.8	15.7	15.4-16.1	1,063.2	20.4	19.8-20.9	830.6	12.3	11.9-12.7	923.2	7.5	7.3-7.8	516.8	10.3	9.9-10.7	406.4	5.7	5.4-5.9
	2006-2010	2,096.8	16.6	16.3-16.9	1,219.2	21.7	21.2-22.3	877.6	12.6	12.2-13.0	920.0	7.1	6.9-7.3	531.2	9.8	9.4-10.2	388.8	5.2	5.0-5.4
	2011-2015	2,468.4	18.4	18.0-18.7	1,447.2	24.0	23.4-24.6	1,021.2	13.9	13.5-14.3	935.0	6.7	6.5-6.9	538.8	9.2	8.8-9.5	396.2	5.0	4.7-5.2
2016-2020	2,400.2	16.8	16.5-17.1	1,426.4	21.9	21.4-22.4	973.8	12.6	12.3-13.0	884.2	5.9	5.8-6.1	514.0	8.1	7.8-8.4	370.2	4.3	4.1-4.6	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Bronx**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	4,569.4	393.2	388.0-398.4	2,199.4	480.8	471.6-490.2	2,370.0	344.7	338.4-351.1	2,660.0	229.1	225.1-233.1	1,347.2	298.4	291.1-305.8	1,312.8	187.9	183.3-192.6
	1981-1985	4,416.0	401.1	395.7-406.5	2,078.6	488.1	478.5-497.7	2,337.4	353.2	346.7-359.8	2,360.0	213.8	209.9-217.7	1,183.8	281.5	274.2-288.9	1,176.2	174.6	170.1-179.2
	1986-1990	4,418.8	405.6	400.2-411.0	2,075.6	497.4	487.7-507.3	2,343.2	356.7	350.2-363.4	2,304.0	211.9	208.0-215.8	1,136.6	280.4	273.0-287.9	1,167.4	173.7	169.3-178.3
	1991-1995	4,799.4	452.6	446.8-458.4	2,292.6	559.5	549.1-570.1	2,506.8	393.1	386.2-400.1	2,310.6	219.8	215.8-223.9	1,125.0	287.7	280.1-295.5	1,185.6	182.6	177.9-187.3
	1996-2000	4,976.0	458.6	452.9-464.3	2,399.4	567.0	556.7-577.5	2,576.6	395.2	388.4-402.1	2,090.4	195.4	191.6-199.2	975.8	241.7	234.8-248.7	1,114.6	169.3	164.8-173.8
	2001-2005	5,360.2	462.3	456.7-467.9	2,650.2	577.2	567.2-587.4	2,710.0	392.5	385.9-399.2	2,055.6	180.6	177.1-184.2	980.0	226.3	219.8-232.9	1,075.6	154.2	150.1-158.4
	2006-2010	5,726.8	463.3	457.9-468.8	2,873.0	574.5	564.8-584.2	2,853.8	393.6	387.1-400.2	2,004.2	166.7	163.4-170.0	968.8	210.5	204.5-216.7	1,035.4	141.7	137.8-145.7
	2011-2015	6,160.4	450.0	444.9-455.1	3,001.6	527.9	519.3-536.7	3,158.8	401.3	395.0-407.7	2,040.8	152.7	149.7-155.7	999.8	191.3	185.9-196.8	1,041.0	129.6	126.0-133.2
	2016-2020	6,204.4	414.7	410.1-419.5	3,038.4	476.9	469.1-484.7	3,166.0	375.2	369.2-381.2	2,009.4	134.9	132.2-137.6	990.2	165.5	160.8-170.3	1,019.2	115.6	112.4-118.9
Oral cavity and pharynx	1976-1980	141.6	12.3	11.4-13.3	97.2	20.7	18.9-22.7	44.4	6.5	5.7-7.5	74.2	6.6	5.9-7.3	52.4	11.4	10.0-12.9	21.8	3.2	2.6-3.9
	1981-1985	147.8	13.9	12.9-14.9	101.0	23.4	21.3-25.5	46.8	7.3	6.3-8.3	57.8	5.3	4.7-6.0	44.2	10.1	8.8-11.5	13.6	2.1	1.6-2.7
	1986-1990	139.0	13.2	12.2-14.2	91.0	21.5	19.6-23.6	48.0	7.5	6.6-8.5	52.0	4.8	4.3-5.5	36.2	8.6	7.4-9.9	15.8	2.4	1.9-3.0
	1991-1995	145.4	14.1	13.1-15.2	96.6	23.0	21.0-25.2	48.8	7.9	6.9-8.9	44.8	4.3	3.8-4.9	32.8	7.9	6.7-9.2	12.0	1.9	1.4-2.4
	1996-2000	118.2	10.8	10.0-11.7	78.6	17.5	15.8-19.3	39.6	6.1	5.3-7.0	37.0	3.4	3.0-4.0	25.4	5.9	4.9-7.1	11.6	1.8	1.4-2.3
	2001-2005	129.0	10.9	10.0-11.7	90.2	18.2	16.5-20.0	38.8	5.6	4.8-6.5	37.0	3.1	2.7-3.6	27.4	5.7	4.8-6.7	9.6	1.3	1.0-1.8
	2006-2010	133.8	10.7	9.9-11.5	89.6	17.3	15.7-19.0	44.2	6.0	5.2-6.8	31.4	2.6	2.2-3.0	25.0	5.1	4.2-6.1	6.4	0.9	0.6-1.3
	2011-2015	141.4	10.1	9.4-10.9	95.2	16.0	14.6-17.5	46.2	5.8	5.1-6.6	33.6	2.4	2.1-2.8	23.6	4.1	3.4-4.9	10.0	1.2	0.9-1.6
	2016-2020	150.4	10.0	9.3-10.7	101.2	15.7	14.3-17.2	49.2	5.7	5.0-6.5	39.6	2.7	2.3-3.1	28.2	4.5	3.8-5.4	11.4	1.3	1.0-1.7
Esophagus	1976-1980	75.4	6.4	5.8-7.1	48.8	10.1	8.8-11.5	26.6	3.9	3.3-4.6	70.2	6.0	5.4-6.7	43.8	9.1	7.9-10.5	26.4	3.9	3.2-4.6
	1981-1985	87.6	8.1	7.3-8.9	57.2	13.4	11.9-15.1	30.4	4.5	3.8-5.3	73.8	6.8	6.1-7.5	50.6	11.8	10.4-13.4	23.2	3.4	2.8-4.1
	1986-1990	78.6	7.3	6.6-8.0	52.0	12.3	10.8-13.9	26.6	4.0	3.3-4.7	62.8	5.8	5.2-6.5	43.2	10.2	8.9-11.7	19.6	2.9	2.4-3.6
	1991-1995	78.6	7.6	6.8-8.4	51.8	12.6	11.1-14.2	26.8	4.2	3.5-4.9	66.6	6.4	5.8-7.2	46.4	11.5	10.1-13.1	20.2	3.2	2.6-3.9
	1996-2000	75.4	7.1	6.4-7.8	48.4	11.5	10.1-13.1	27.0	4.2	3.5-4.9	60.4	5.7	5.0-6.3	40.2	9.6	8.3-11.0	20.2	3.1	2.5-3.7
	2001-2005	59.6	5.2	4.6-5.8	42.0	9.2	8.0-10.6	17.6	2.5	2.0-3.1	49.4	4.3	3.8-4.9	32.2	7.2	6.1-8.5	17.2	2.5	2.0-3.0
	2006-2010	58.8	4.8	4.3-5.4	41.6	8.4	7.3-9.7	17.2	2.3	1.9-2.9	44.6	3.7	3.2-4.2	31.6	6.5	5.5-7.7	13.0	1.8	1.4-2.3
	2011-2015	53.4	3.9	3.4-4.4	36.8	6.7	5.7-7.7	16.6	2.0	1.6-2.5	45.6	3.4	2.9-3.8	32.6	6.0	5.1-7.0	13.0	1.6	1.2-2.0
	2016-2020	48.0	3.2	2.8-3.6	35.2	5.7	4.8-6.6	12.8	1.4	1.1-1.8	32.2	2.1	1.8-2.4	22.6	3.6	2.9-4.3	9.6	1.1	0.8-1.4
Stomach	1976-1980	177.6	15.3	14.3-16.4	96.8	21.6	19.6-23.6	80.8	11.5	10.4-12.7	139.6	12.2	11.3-13.1	73.6	16.7	15.0-18.6	66.0	9.3	8.3-10.4
	1981-1985	166.4	15.1	14.1-16.2	92.8	22.0	20.0-24.2	73.6	10.6	9.5-11.8	112.8	10.3	9.5-11.2	64.2	15.7	14.0-17.5	48.6	7.0	6.2-8.0
	1986-1990	142.2	13.1	12.1-14.1	71.6	17.6	15.8-19.5	70.6	10.3	9.2-11.4	101.4	9.3	8.5-10.2	51.8	12.9	11.4-14.6	49.6	7.1	6.3-8.1
	1991-1995	145.8	13.9	12.9-15.0	75.8	19.4	17.5-21.5	70.0	10.5	9.4-11.7	102.2	9.7	8.9-10.6	52.4	13.6	12.0-15.4	49.8	7.4	6.5-8.4
	1996-2000	144.0	13.5	12.5-14.5	75.4	18.7	16.8-20.7	68.6	10.3	9.2-11.5	96.4	9.0	8.2-9.9	49.8	12.5	11.0-14.2	46.6	6.9	6.1-7.9
	2001-2005	142.0	12.5	11.6-13.4	72.8	16.7	15.0-18.6	69.2	9.9	8.8-11.0	82.6	7.3	6.6-8.0	42.2	9.8	8.5-11.2	40.4	5.7	5.0-6.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Bronx**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	141.4	11.7	10.9-12.6	74.0	15.5	13.9-17.2	67.4	9.2	8.3-10.3	68.8	5.7	5.1-6.4	33.2	7.2	6.1-8.4	35.6	4.8	4.1-5.6
	2011-2015	150.0	11.3	10.5-12.1	77.2	14.5	13.1-16.1	72.8	9.2	8.3-10.2	68.6	5.2	4.7-5.8	35.8	7.1	6.1-8.3	32.8	4.1	3.5-4.8
	2016-2020	144.8	10.0	9.2-10.7	80.6	13.3	12.0-14.7	64.2	7.7	6.8-8.6	57.6	4.0	3.5-4.5	35.6	6.0	5.1-7.0	22.0	2.6	2.1-3.1
Colorectal	1976-1980	783.4	67.0	64.9-69.1	374.2	83.6	79.7-87.6	409.2	57.2	54.7-59.8	417.4	36.2	34.7-37.9	203.0	46.4	43.5-49.4	214.4	30.3	28.5-32.2
	1981-1985	688.0	61.9	59.8-64.0	321.2	77.9	74.0-81.8	366.8	52.9	50.5-55.4	337.2	30.4	29.0-31.9	158.0	38.7	36.0-41.6	179.2	25.8	24.1-27.6
	1986-1990	635.0	58.0	56.0-60.1	294.2	72.9	69.1-76.8	340.8	49.5	47.2-51.9	302.6	27.7	26.3-29.1	146.0	37.1	34.4-40.0	156.6	22.6	21.0-24.2
	1991-1995	593.8	56.7	54.6-58.7	261.0	67.1	63.5-70.9	332.8	50.5	48.1-53.0	282.4	26.8	25.4-28.2	126.6	33.4	30.8-36.1	155.8	23.1	21.5-24.8
	1996-2000	623.2	58.4	56.3-60.5	275.2	68.5	64.9-72.3	348.0	52.4	49.9-54.9	249.6	23.4	22.1-24.8	109.4	27.7	25.4-30.1	140.2	20.9	19.3-22.5
	2001-2005	622.8	54.8	52.9-56.8	289.6	65.9	62.4-69.4	333.2	48.2	45.9-50.5	238.6	21.2	20.0-22.4	105.6	25.1	23.0-27.4	133.0	18.8	17.4-20.3
	2006-2010	597.6	49.2	47.4-51.0	284.4	58.8	55.7-62.0	313.2	42.9	40.8-45.1	219.0	18.4	17.3-19.6	106.6	23.9	21.9-26.1	112.4	15.3	14.0-16.6
	2011-2015	543.6	40.3	38.8-41.9	270.6	49.2	46.5-51.9	273.0	34.1	32.3-36.0	210.6	15.8	14.9-16.8	101.6	19.4	17.7-21.2	109.0	13.4	12.2-14.5
2016-2020	521.2	35.4	34.0-36.8	267.2	43.0	40.7-45.4	254.0	29.9	28.3-31.6	180.4	12.2	11.4-13.1	90.2	15.3	13.9-16.8	90.2	10.1	9.2-11.1	
Colon excluding rectum	1976-1980	578.4	49.6	47.8-51.5	267.4	60.5	57.2-63.9	311.0	43.4	41.3-45.7	345.0	30.0	28.6-31.5	163.8	37.7	35.1-40.4	181.2	25.6	23.9-27.3
	1981-1985	497.2	44.7	42.9-46.5	223.2	54.5	51.3-57.8	274.0	39.4	37.3-41.6	286.8	25.8	24.5-27.2	132.8	32.6	30.1-35.2	154.0	22.1	20.6-23.8
	1986-1990	477.2	43.5	41.8-45.3	218.6	54.6	51.4-58.0	258.6	37.3	35.3-39.4	264.8	24.2	22.9-25.5	129.6	33.0	30.5-35.7	135.2	19.4	18.0-21.0
	1991-1995	442.2	42.1	40.4-43.9	191.2	49.7	46.5-53.0	251.0	37.9	35.8-40.1	242.2	23.0	21.7-24.3	108.8	28.8	26.4-31.3	133.4	19.8	18.3-21.4
	1996-2000	466.8	43.8	42.0-45.6	198.8	50.1	47.0-53.3	268.0	40.3	38.1-42.5	219.4	20.6	19.4-21.8	96.0	24.5	22.3-26.8	123.4	18.4	17.0-19.9
	2001-2005	464.2	41.0	39.4-42.7	209.6	48.4	45.5-51.5	254.6	36.7	34.7-38.8	206.2	18.3	17.2-19.5	89.6	21.4	19.5-23.5	116.6	16.5	15.1-17.9
	2006-2010	427.8	35.5	34.0-37.0	199.0	42.0	39.3-44.7	228.8	31.4	29.6-33.3	188.0	15.9	14.9-16.9	88.8	20.2	18.3-22.2	99.2	13.5	12.3-14.7
	2011-2015	389.4	29.1	27.8-30.5	189.8	35.3	33.0-37.7	199.6	25.0	23.4-26.6	177.2	13.3	12.4-14.2	82.8	16.0	14.4-17.6	94.4	11.5	10.5-12.6
2016-2020	365.4	24.9	23.7-26.0	181.6	29.4	27.5-31.4	183.8	21.6	20.2-23.1	148.6	10.1	9.3-10.8	70.0	11.9	10.6-13.2	78.6	8.8	8.0-9.8	
Rectum & rectosigmoid	1976-1980	205.0	17.4	16.3-18.5	106.8	23.1	21.1-25.2	98.2	13.8	12.6-15.1	72.4	6.3	5.6-7.0	39.2	8.7	7.5-10.1	33.2	4.7	4.0-5.5
	1981-1985	190.8	17.2	16.1-18.4	98.0	23.4	21.3-25.6	92.8	13.5	12.3-14.8	50.4	4.6	4.0-5.2	25.2	6.1	5.1-7.3	25.2	3.6	3.0-4.4
	1986-1990	157.8	14.5	13.5-15.6	75.6	18.3	16.5-20.3	82.2	12.2	11.0-13.4	37.8	3.5	3.0-4.0	16.4	4.1	3.3-5.1	21.4	3.1	2.5-3.8
	1991-1995	151.6	14.5	13.5-15.6	69.8	17.4	15.6-19.4	81.8	12.6	11.4-13.9	40.2	3.8	3.3-4.4	17.8	4.6	3.7-5.7	22.4	3.3	2.7-4.0
	1996-2000	156.4	14.6	13.6-15.6	76.4	18.4	16.6-20.4	80.0	12.1	11.0-13.4	30.2	2.8	2.4-3.3	13.4	3.2	2.5-4.0	16.8	2.5	2.0-3.1
	2001-2005	158.6	13.8	12.8-14.8	80.0	17.5	15.7-19.3	78.6	11.4	10.3-12.6	32.4	2.8	2.4-3.3	16.0	3.7	2.9-4.6	16.4	2.3	1.8-2.9
	2006-2010	169.8	13.7	12.8-14.7	85.4	16.8	15.2-18.6	84.4	11.5	10.4-12.6	31.0	2.6	2.2-3.0	17.8	3.7	3.0-4.6	13.2	1.8	1.4-2.2
	2011-2015	154.2	11.2	10.4-12.0	80.8	13.8	12.5-15.3	73.4	9.2	8.3-10.2	33.4	2.5	2.1-2.9	18.8	3.4	2.7-4.2	14.6	1.8	1.4-2.3
2016-2020	155.8	10.5	9.8-11.3	85.6	13.6	12.3-15.0	70.2	8.3	7.5-9.3	31.8	2.2	1.8-2.5	20.2	3.4	2.8-4.2	11.6	1.3	1.0-1.7	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Bronx**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	46.0	3.9	3.4-4.5	27.8	6.0	5.0-7.1	18.2	2.6	2.1-3.2	52.6	4.5	4.0-5.1	28.2	6.1	5.1-7.2	24.4	3.5	2.9-4.2
	1981-1985	51.6	4.7	4.1-5.3	33.0	7.6	6.5-8.9	18.6	2.8	2.2-3.4	51.6	4.6	4.1-5.3	29.0	6.6	5.6-7.8	22.6	3.3	2.7-4.0
	1986-1990	51.8	4.8	4.2-5.4	32.4	7.7	6.5-9.0	19.4	2.9	2.3-3.5	55.4	5.1	4.5-5.8	30.2	7.4	6.2-8.7	25.2	3.8	3.1-4.5
	1991-1995	73.4	7.0	6.3-7.8	49.8	12.0	10.5-13.6	23.6	3.6	3.0-4.3	69.4	6.6	6.0-7.4	44.0	10.8	9.4-12.3	25.4	3.9	3.2-4.6
	1996-2000	93.2	8.6	7.9-9.5	64.2	14.2	12.7-15.9	29.0	4.4	3.7-5.2	62.6	5.9	5.2-6.5	39.4	9.1	7.8-10.4	23.2	3.5	2.9-4.2
	2001-2005	144.0	12.3	11.4-13.2	103.0	20.9	19.1-22.8	41.0	6.0	5.2-6.9	94.0	8.2	7.4-8.9	61.0	12.8	11.4-14.4	33.0	4.7	4.0-5.5
	2006-2010	196.4	15.4	14.5-16.4	144.6	26.8	24.9-28.9	51.8	7.2	6.3-8.1	115.8	9.3	8.5-10.1	82.0	15.8	14.2-17.4	33.8	4.7	4.0-5.5
	2011-2015	243.2	17.0	16.1-18.0	173.0	28.4	26.5-30.4	70.2	8.7	7.8-9.6	135.0	9.6	8.9-10.4	93.4	15.9	14.4-17.4	41.6	5.1	4.4-5.8
	2016-2020	218.2	14.0	13.2-14.9	151.2	22.8	21.2-24.5	67.0	7.5	6.7-8.3	135.2	8.8	8.2-9.5	89.8	14.1	12.8-15.5	45.4	5.0	4.4-5.7
Pancreas	1976-1980	146.8	12.6	11.7-13.5	75.6	16.7	15.0-18.5	71.2	10.0	9.0-11.1	138.8	11.8	11.0-12.8	69.2	15.2	13.6-16.9	69.6	9.7	8.7-10.8
	1981-1985	133.2	12.0	11.1-12.9	60.8	14.3	12.7-16.0	72.4	10.5	9.4-11.6	120.2	10.7	9.9-11.7	59.8	13.9	12.3-15.6	60.4	8.7	7.8-9.8
	1986-1990	130.8	12.0	11.1-12.9	56.4	13.7	12.2-15.5	74.4	10.6	9.6-11.8	114.2	10.4	9.6-11.3	50.2	12.3	10.8-14.0	64.0	9.1	8.2-10.2
	1991-1995	129.6	12.4	11.5-13.4	51.6	13.2	11.6-14.9	78.0	11.7	10.6-13.0	117.8	11.3	10.4-12.3	47.8	12.3	10.8-14.0	70.0	10.5	9.4-11.6
	1996-2000	136.0	12.8	11.9-13.8	54.0	13.3	11.8-15.0	82.0	12.3	11.2-13.6	110.4	10.4	9.5-11.3	43.0	10.8	9.4-12.4	67.4	10.0	9.0-11.2
	2001-2005	148.4	13.1	12.2-14.1	64.0	14.3	12.8-16.0	84.4	12.1	10.9-13.3	116.6	10.3	9.5-11.2	50.8	11.6	10.2-13.1	65.8	9.3	8.3-10.4
	2006-2010	150.4	12.5	11.6-13.4	64.6	13.4	11.9-15.0	85.8	11.8	10.7-13.0	120.6	10.1	9.3-11.0	51.2	10.8	9.5-12.3	69.4	9.5	8.5-10.6
	2011-2015	194.2	14.4	13.5-15.4	88.2	16.0	14.5-17.6	106.0	13.3	12.2-14.5	142.6	10.6	9.9-11.5	64.8	12.1	10.8-13.5	77.8	9.7	8.7-10.7
	2016-2020	200.8	13.4	12.5-14.2	103.8	16.7	15.3-18.2	97.0	11.0	10.0-12.0	147.4	9.8	9.1-10.6	75.8	12.5	11.3-13.9	71.6	8.0	7.2-8.9
Larynx	1976-1980	67.6	5.8	5.2-6.4	55.2	11.6	10.2-13.0	12.4	1.9	1.5-2.5	29.6	2.6	2.2-3.0	23.8	5.2	4.3-6.2	5.8	0.9	0.6-1.3
	1981-1985	77.8	7.2	6.5-8.0	61.2	14.2	12.6-15.9	16.6	2.6	2.1-3.3	29.0	2.6	2.2-3.1	22.6	5.0	4.1-6.1	6.4	1.0	0.7-1.5
	1986-1990	76.6	7.2	6.5-8.0	55.0	12.9	11.4-14.5	21.6	3.6	2.9-4.3	28.6	2.7	2.3-3.2	21.8	5.2	4.2-6.3	6.8	1.1	0.8-1.5
	1991-1995	71.8	6.9	6.2-7.7	53.6	12.8	11.3-14.4	18.2	3.0	2.4-3.7	31.2	3.0	2.5-3.5	24.4	5.8	4.8-7.0	6.8	1.1	0.8-1.5
	1996-2000	65.8	6.1	5.5-6.8	49.8	11.4	10.0-12.9	16.0	2.5	2.0-3.1	33.0	3.1	2.6-3.6	27.0	6.2	5.2-7.4	6.0	1.0	0.6-1.4
	2001-2005	59.4	5.1	4.5-5.7	44.2	9.3	8.1-10.6	15.2	2.2	1.8-2.8	26.4	2.3	1.9-2.7	21.8	4.7	3.9-5.7	4.6	0.7	0.4-1.0
	2006-2010	59.0	4.7	4.2-5.3	45.4	8.7	7.6-10.0	13.6	1.9	1.4-2.4	22.0	1.8	1.4-2.1	17.6	3.5	2.8-4.3	4.4	0.6	0.4-0.9
	2011-2015	57.6	4.1	3.7-4.7	42.8	7.5	6.5-8.6	14.8	1.8	1.4-2.3	24.0	1.7	1.4-2.1	18.2	3.3	2.7-4.1	5.8	0.7	0.5-1.0
	2016-2020	55.4	3.7	3.2-4.1	41.6	6.6	5.7-7.6	13.8	1.6	1.3-2.1	16.6	1.1	0.9-1.4	13.0	2.1	1.6-2.7	3.6	0.4	0.2-0.6
Lung and bronchus	1976-1980	629.8	53.2	51.3-55.1	449.0	95.9	91.8-100.0	180.8	25.8	24.1-27.6	522.8	44.1	42.4-45.9	384.0	82.3	78.6-86.2	138.8	19.7	18.2-21.3
	1981-1985	624.0	56.1	54.1-58.1	402.6	93.3	89.2-97.6	221.4	32.7	30.8-34.8	502.6	45.3	43.5-47.1	334.2	78.3	74.5-82.2	168.4	25.0	23.3-26.7
	1986-1990	638.6	58.8	56.7-60.8	388.8	93.6	89.4-97.9	249.8	37.5	35.5-39.7	517.8	47.6	45.8-49.5	317.6	76.9	73.2-80.9	200.2	29.9	28.0-31.8
	1991-1995	616.2	59.0	56.9-61.1	346.2	85.8	81.7-89.9	270.0	42.7	40.4-45.1	505.6	48.4	46.5-50.4	297.4	74.3	70.6-78.2	208.2	32.6	30.7-34.7
	1996-2000	599.4	56.3	54.3-58.4	330.8	80.1	76.2-84.1	268.6	41.6	39.4-43.9	468.4	44.2	42.4-46.0	262.8	64.4	60.9-68.0	205.6	31.5	29.6-33.5
	2001-2005	608.8	53.7	51.8-55.6	323.2	72.6	69.0-76.3	285.6	41.8	39.6-44.0	461.0	40.9	39.2-42.6	247.6	56.7	53.6-60.0	213.4	31.1	29.2-33.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Bronx**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	651.0	54.2	52.3-56.1	343.2	72.1	68.6-75.6	307.8	42.8	40.7-45.0	439.0	36.7	35.2-38.3	236.6	50.8	47.9-53.9	202.4	28.0	26.3-29.8
	2011-2015	650.8	48.5	46.8-50.2	330.0	61.4	58.4-64.5	320.8	40.3	38.3-42.3	416.2	31.3	29.9-32.7	225.2	42.7	40.2-45.3	191.0	24.0	22.5-25.6
	2016-2020	644.2	42.8	41.4-44.4	325.2	52.9	50.3-55.6	319.0	36.0	34.3-37.9	374.4	24.9	23.8-26.1	199.6	32.7	30.7-34.8	174.8	19.5	18.2-20.9
Melanoma of the skin	1976-1980	51.4	4.5	4.0-5.1	26.8	5.7	4.8-6.8	24.6	3.8	3.1-4.5	20.4	1.8	1.5-2.2	12.2	2.7	2.1-3.5	8.2	1.2	0.9-1.7
	1981-1985	51.2	4.7	4.1-5.3	25.4	5.9	4.9-7.1	25.8	4.0	3.3-4.8	19.2	1.8	1.5-2.2	12.2	3.0	2.2-3.8	7.0	1.1	0.8-1.6
	1986-1990	37.4	3.4	2.9-4.0	20.4	4.9	4.0-6.0	17.0	2.6	2.0-3.2	18.2	1.7	1.4-2.1	11.6	3.0	2.2-3.8	6.6	1.0	0.7-1.4
	1991-1995	42.0	3.9	3.4-4.5	22.0	5.3	4.3-6.4	20.0	3.1	2.5-3.8	16.4	1.6	1.2-1.9	8.6	2.1	1.5-2.9	7.8	1.2	0.8-1.6
	1996-2000	42.2	3.9	3.3-4.4	21.6	5.2	4.3-6.3	20.6	3.0	2.4-3.6	14.2	1.3	1.0-1.7	7.2	1.7	1.2-2.4	7.0	1.1	0.7-1.5
	2001-2005	53.0	4.6	4.1-5.2	27.6	6.3	5.3-7.4	25.4	3.6	3.0-4.3	13.0	1.1	0.9-1.4	7.0	1.6	1.1-2.2	6.0	0.8	0.6-1.2
	2006-2010	54.4	4.5	4.0-5.1	27.6	6.0	5.0-7.2	26.8	3.7	3.1-4.4	12.6	1.1	0.8-1.3	8.0	1.8	1.3-2.4	4.6	0.6	0.4-0.9
	2011-2015	60.4	4.5	4.0-5.1	32.4	6.1	5.2-7.1	28.0	3.5	3.0-4.2	12.0	0.9	0.7-1.2	6.8	1.3	0.9-1.9	5.2	0.6	0.4-0.9
2016-2020	58.4	4.0	3.6-4.5	32.2	5.5	4.7-6.5	26.2	3.1	2.6-3.7	6.6	0.4	0.3-0.6	4.0	0.7	0.4-1.0	2.6	0.3	0.1-0.5	
Female breast	1976-1980							613.0	91.8	88.5-95.2							262.2	38.3	36.2-40.5
	1981-1985							592.4	93.3	89.9-96.9							229.0	35.1	33.0-37.2
	1986-1990							628.2	99.8	96.3-103.4							220.8	34.0	32.0-36.1
	1991-1995							680.2	110.0	106.3-113.8							220.6	35.0	32.9-37.1
	1996-2000							716.8	112.1	108.5-115.9							197.6	30.5	28.6-32.5
	2001-2005							732.6	106.9	103.5-110.5							182.8	26.3	24.6-28.1
	2006-2010							780.2	107.9	104.5-111.4							171.8	23.3	21.8-25.0
	2011-2015							882.0	112.9	109.6-116.3							170.2	21.4	20.0-22.9
2016-2020							925.0	111.6	108.3-114.9							166.4	19.3	18.0-20.7	
Cervix uteri	1976-1980							109.0	18.1	16.6-19.8							45.2	7.1	6.1-8.1
	1981-1985							108.6	18.4	16.9-20.1							36.2	5.8	5.0-6.8
	1986-1990							100.6	16.4	15.0-18.0							29.2	4.8	4.0-5.7
	1991-1995							104.0	16.9	15.5-18.4							30.8	4.9	4.2-5.8
	1996-2000							113.6	17.6	16.1-19.1							38.6	6.0	5.2-6.9
	2001-2005							92.4	13.5	12.2-14.7							30.8	4.5	3.8-5.3
	2006-2010							81.4	11.4	10.3-12.6							28.2	3.9	3.3-4.6
	2011-2015							74.4	9.8	8.8-10.9							25.2	3.2	2.7-3.9
	2016-2020							72.0	9.3	8.3-10.3							22.4	2.8	2.3-3.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Bronx**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							135.0	19.2	17.8-20.8							45.6	6.4	5.6-7.3
	1981-1985							122.0	18.2	16.7-19.7							34.6	5.0	4.3-5.8
	1986-1990							113.8	17.4	16.0-19.0							30.0	4.4	3.8-5.2
	1991-1995							127.4	20.2	18.6-21.8							37.4	5.7	4.9-6.6
	1996-2000							147.8	23.0	21.4-24.8							38.0	5.8	5.0-6.7
	2001-2005							177.8	26.0	24.3-27.8							37.0	5.4	4.7-6.2
	2006-2010							215.0	29.6	27.9-31.4							51.2	7.1	6.3-8.1
	2011-2015							249.4	31.0	29.3-32.8							60.0	7.5	6.7-8.5
	2016-2020							265.8	30.6	29.0-32.3							76.8	8.7	7.8-9.6
Ovary	1976-1980							97.6	14.1	12.9-15.5							65.0	9.2	8.2-10.3
	1981-1985							88.8	13.8	12.5-15.2							57.4	8.6	7.7-9.7
	1986-1990							72.6	11.2	10.1-12.4							47.4	7.1	6.2-8.1
	1991-1995							84.4	13.3	12.1-14.7							53.6	8.4	7.4-9.4
	1996-2000							79.4	12.4	11.2-13.7							48.6	7.6	6.6-8.6
	2001-2005							86.4	12.5	11.3-13.7							50.6	7.3	6.5-8.3
	2006-2010							82.8	11.3	10.3-12.5							48.6	6.6	5.8-7.4
	2011-2015							83.0	10.5	9.5-11.5							40.8	5.1	4.4-5.9
	2016-2020							89.0	10.6	9.7-11.7							46.2	5.3	4.6-6.0
Prostate	1976-1980				345.4	79.6	75.8-83.6							129.4	31.6	29.2-34.3			
	1981-1985				339.0	82.1	78.2-86.2							121.2	31.0	28.6-33.7			
	1986-1990				376.2	94.9	90.6-99.3							137.6	36.6	33.9-39.5			
	1991-1995				574.2	146.1	140.8-151.6							160.8	45.5	42.4-48.8			
	1996-2000				742.2	180.5	174.7-186.5							131.4	36.4	33.6-39.3			
	2001-2005				870.8	193.4	187.6-199.4							137.4	36.1	33.4-39.0			
	2006-2010				943.2	187.6	182.1-193.2							127.8	32.1	29.7-34.8			
	2011-2015				944.6	160.9	156.2-165.7							119.2	26.1	24.0-28.3			
	2016-2020				937.0	140.8	136.7-145.0							124.2	22.6	20.8-24.5			
Testis	1976-1980				9.6	1.8	1.3-2.4							1.8	0.3	0.1-0.6			
	1981-1985				9.8	1.8	1.3-2.4							0.4	0.1	0.0-0.3			
	1986-1990				11.8	1.9	1.4-2.5							1.2	0.2	0.1-0.5			
	1991-1995				15.0	2.4	1.9-3.1							0.6	0.1	0.0-0.4			
	1996-2000				14.8	2.4	1.9-3.0							0.8	0.1	0.0-0.3			
	2001-2005				16.2	2.5	2.0-3.2							2.0	0.3	0.2-0.6			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Bronx**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				25.4	4.1	3.4-4.8							2.2	0.4	0.2-0.7			
	2011-2015				24.8	3.7	3.1-4.5							1.6	0.3	0.1-0.5			
	2016-2020				21.4	3.2	2.6-3.8							2.2	0.4	0.2-0.7			
Urinary bladder (incl. in situ)	1976-1980	207.6	17.6	16.6-18.8	143.2	32.1	29.7-34.6	64.4	8.9	7.9-9.9	67.6	5.9	5.3-6.6	44.2	10.7	9.3-12.2	23.4	3.3	2.7-3.9
	1981-1985	186.6	16.7	15.6-17.8	127.4	30.8	28.4-33.3	59.2	8.4	7.5-9.4	50.4	4.5	4.0-5.1	33.4	8.4	7.1-9.8	17.0	2.4	1.9-3.0
	1986-1990	174.6	15.9	14.9-17.0	118.6	29.8	27.4-32.3	56.0	8.1	7.1-9.1	49.4	4.5	4.0-5.1	30.0	8.0	6.7-9.4	19.4	2.8	2.2-3.4
	1991-1995	155.6	14.9	13.8-16.0	105.6	28.0	25.7-30.5	50.0	7.6	6.7-8.6	51.8	5.0	4.4-5.6	33.0	9.2	7.9-10.7	18.8	2.8	2.3-3.4
	1996-2000	159.2	15.0	14.0-16.1	102.2	26.0	23.8-28.4	57.0	8.5	7.5-9.6	45.6	4.3	3.8-4.9	24.2	6.5	5.3-7.7	21.4	3.1	2.6-3.8
	2001-2005	172.0	15.4	14.4-16.5	113.0	27.1	24.9-29.5	59.0	8.4	7.4-9.4	43.8	3.9	3.4-4.5	24.8	6.6	5.4-7.8	19.0	2.6	2.1-3.2
	2006-2010	158.0	13.5	12.5-14.4	115.0	26.1	23.9-28.3	43.0	5.9	5.1-6.7	41.4	3.5	3.1-4.1	24.8	6.1	5.0-7.2	16.6	2.2	1.7-2.7
	2011-2015	182.8	13.9	13.0-14.8	126.8	24.9	22.9-26.9	56.0	7.0	6.2-7.8	42.2	3.3	2.8-3.7	27.4	5.9	5.0-7.0	14.8	1.8	1.4-2.3
2016-2020	193.4	13.1	12.2-13.9	137.8	23.3	21.5-25.1	55.6	6.3	5.6-7.1	44.2	3.0	2.6-3.5	28.0	5.0	4.2-6.0	16.2	1.8	1.4-2.2	
Kidney and renal pelvis	1976-1980	80.4	6.8	6.1-7.5	48.0	10.2	8.9-11.6	32.4	4.6	3.9-5.3	45.0	3.8	3.3-4.3	27.4	5.9	4.9-7.0	17.6	2.4	1.9-3.0
	1981-1985	74.4	6.7	6.0-7.4	40.2	9.2	8.0-10.6	34.2	5.1	4.3-5.9	32.4	2.9	2.5-3.4	18.6	4.4	3.5-5.4	13.8	2.0	1.6-2.6
	1986-1990	82.2	7.5	6.8-8.3	48.8	11.3	9.9-12.8	33.4	5.1	4.3-5.9	37.2	3.4	2.9-3.9	21.8	5.3	4.3-6.4	15.4	2.3	1.8-2.8
	1991-1995	89.8	8.5	7.7-9.3	54.2	12.9	11.4-14.6	35.6	5.6	4.8-6.5	36.2	3.4	2.9-4.0	20.6	5.1	4.2-6.3	15.6	2.4	1.9-3.0
	1996-2000	107.6	10.0	9.2-10.9	60.4	14.1	12.5-15.8	47.2	7.3	6.4-8.3	30.0	2.8	2.4-3.3	17.6	4.2	3.4-5.2	12.4	1.9	1.4-2.4
	2001-2005	145.6	12.5	11.6-13.4	79.8	16.5	14.9-18.2	65.8	9.6	8.6-10.7	33.8	2.9	2.5-3.4	20.8	4.6	3.7-5.6	13.0	1.8	1.4-2.3
	2006-2010	178.0	14.4	13.5-15.4	105.0	20.5	18.8-22.4	73.0	10.1	9.1-11.2	36.6	3.1	2.6-3.5	22.2	4.8	4.0-5.8	14.4	2.0	1.5-2.5
	2011-2015	226.4	16.6	15.7-17.7	139.0	24.1	22.3-26.0	87.4	11.1	10.1-12.2	39.8	3.0	2.6-3.4	26.0	5.0	4.2-6.0	13.8	1.6	1.3-2.1
2016-2020	226.8	15.0	14.1-15.9	139.0	21.7	20.1-23.4	87.8	10.2	9.3-11.3	41.6	2.8	2.5-3.3	26.0	4.4	3.7-5.2	15.6	1.7	1.4-2.2	
Brain and other central nervous system	1976-1980	61.4	5.1	4.5-5.7	27.4	5.4	4.5-6.4	34.0	5.0	4.3-5.8	42.8	3.5	3.0-4.0	19.0	3.7	3.0-4.6	23.8	3.4	2.8-4.0
	1981-1985	58.8	5.2	4.6-5.8	31.2	6.6	5.6-7.8	27.6	4.1	3.4-4.8	40.0	3.5	3.1-4.1	18.4	4.1	3.3-5.0	21.6	3.1	2.6-3.8
	1986-1990	68.4	6.1	5.4-6.7	34.4	7.4	6.3-8.6	34.0	5.2	4.4-6.0	40.2	3.6	3.1-4.2	20.4	4.6	3.7-5.6	19.8	2.9	2.4-3.6
	1991-1995	72.4	6.4	5.8-7.1	38.0	7.9	6.8-9.2	34.4	5.3	4.5-6.2	38.0	3.5	3.0-4.1	19.4	4.5	3.6-5.4	18.6	2.9	2.3-3.6
	1996-2000	70.8	6.0	5.4-6.7	34.2	6.8	5.8-8.0	36.6	5.4	4.6-6.2	35.4	3.2	2.7-3.7	15.2	3.1	2.5-4.0	20.2	3.1	2.5-3.7
	2001-2005	64.0	5.2	4.6-5.8	33.0	6.2	5.2-7.2	31.0	4.4	3.8-5.2	30.4	2.6	2.2-3.0	17.6	3.6	2.8-4.4	12.8	1.8	1.4-2.4
	2006-2010	65.2	5.0	4.5-5.6	34.6	6.3	5.4-7.3	30.6	4.1	3.5-4.9	35.4	2.8	2.4-3.3	17.8	3.4	2.7-4.3	17.6	2.4	1.9-3.0
	2011-2015	71.0	5.2	4.6-5.7	35.0	5.7	4.9-6.6	36.0	4.7	4.1-5.5	35.0	2.6	2.2-3.0	16.0	2.9	2.3-3.6	19.0	2.4	1.9-3.0
2016-2020	75.4	5.2	4.7-5.7	42.4	6.6	5.7-7.6	33.0	4.0	3.4-4.7	45.4	3.1	2.7-3.5	25.0	4.0	3.4-4.8	20.4	2.4	2.0-3.0	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Bronx**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	35.6	3.1	2.7-3.6	8.4	1.8	1.3-2.4	27.2	4.1	3.4-4.9	6.6	0.6	0.4-0.9	1.8	0.4	0.2-0.7	4.8	0.8	0.5-1.2
	1981-1985	37.4	3.6	3.1-4.2	9.8	2.3	1.7-3.0	27.6	4.5	3.8-5.4	5.4	0.5	0.3-0.7	1.4	0.3	0.1-0.7	4.0	0.6	0.3-0.9
	1986-1990	31.0	2.8	2.4-3.3	6.8	1.5	1.0-2.1	24.2	3.9	3.2-4.6	7.0	0.7	0.5-0.9	1.8	0.4	0.2-0.8	5.2	0.8	0.5-1.1
	1991-1995	43.6	3.9	3.4-4.4	7.6	1.7	1.2-2.3	36.0	5.6	4.8-6.5	6.8	0.6	0.4-0.9	1.4	0.3	0.1-0.7	5.4	0.8	0.5-1.2
	1996-2000	60.4	5.2	4.6-5.8	12.0	2.6	1.9-3.3	48.4	7.3	6.4-8.3	3.4	0.3	0.2-0.5	1.0	0.3	0.1-0.6	2.4	0.4	0.2-0.6
	2001-2005	95.6	7.7	7.0-8.4	21.2	4.1	3.3-4.9	74.4	10.6	9.6-11.7	5.6	0.5	0.3-0.7	2.2	0.5	0.3-0.9	3.4	0.5	0.3-0.8
	2006-2010	140.0	10.8	10.0-11.6	28.0	5.1	4.3-6.0	112.0	15.5	14.3-16.9	5.0	0.4	0.3-0.6	1.6	0.4	0.2-0.7	3.4	0.5	0.3-0.8
	2011-2015	229.0	16.7	15.7-17.7	36.0	6.0	5.2-7.0	193.0	25.4	23.8-27.1	7.2	0.5	0.4-0.8	2.4	0.5	0.2-0.8	4.8	0.6	0.4-0.9
2016-2020	203.8	14.3	13.5-15.3	41.8	6.6	5.7-7.6	162.0	21.0	19.5-22.5	7.0	0.5	0.3-0.7	3.0	0.5	0.3-0.9	4.0	0.5	0.3-0.7	
Hodgkin lymphoma	1976-1980	37.8	3.0	2.6-3.5	19.6	3.6	2.9-4.4	18.2	2.6	2.1-3.3	14.0	1.2	0.9-1.5	7.8	1.5	1.1-2.1	6.2	0.9	0.6-1.3
	1981-1985	32.8	2.8	2.4-3.3	18.0	3.5	2.8-4.4	14.8	2.1	1.7-2.7	11.4	1.0	0.8-1.3	5.6	1.1	0.7-1.6	5.8	0.9	0.6-1.3
	1986-1990	36.6	3.0	2.6-3.5	22.8	4.3	3.5-5.2	13.8	2.1	1.6-2.6	8.4	0.7	0.5-1.0	6.2	1.2	0.8-1.8	2.2	0.3	0.2-0.6
	1991-1995	41.6	3.5	3.0-4.0	24.2	4.5	3.7-5.5	17.4	2.6	2.0-3.2	13.2	1.2	0.9-1.5	7.2	1.5	1.1-2.1	6.0	0.9	0.6-1.3
	1996-2000	39.4	3.2	2.7-3.6	25.0	4.6	3.8-5.5	14.4	2.1	1.6-2.7	10.0	0.9	0.6-1.1	5.4	1.1	0.7-1.6	4.6	0.7	0.4-1.0
	2001-2005	45.4	3.5	3.0-4.0	24.4	4.1	3.4-4.9	21.0	3.0	2.4-3.6	6.2	0.5	0.3-0.7	4.4	0.8	0.5-1.3	1.8	0.2	0.1-0.5
	2006-2010	43.4	3.2	2.8-3.7	21.2	3.5	2.8-4.2	22.2	3.0	2.4-3.6	6.2	0.5	0.3-0.7	3.6	0.7	0.4-1.1	2.6	0.4	0.2-0.6
	2011-2015	52.4	3.6	3.2-4.1	26.0	3.9	3.3-4.7	26.4	3.4	2.8-4.0	5.8	0.4	0.3-0.6	2.8	0.5	0.3-0.8	3.0	0.4	0.2-0.6
2016-2020	41.2	2.9	2.5-3.3	22.2	3.4	2.8-4.1	19.0	2.5	2.0-3.0	4.6	0.3	0.2-0.5	3.0	0.5	0.3-0.8	1.6	0.2	0.1-0.4	
Non-Hodgkin lymphomas	1976-1980	130.4	11.0	10.1-11.9	64.2	13.2	11.7-14.8	66.2	9.4	8.4-10.5	72.4	6.1	5.5-6.7	35.8	7.5	6.4-8.7	36.6	5.1	4.4-5.9
	1981-1985	148.0	13.2	12.3-14.2	67.6	15.4	13.8-17.2	80.4	11.8	10.6-13.0	69.2	6.3	5.6-7.0	33.6	7.9	6.8-9.3	35.6	5.3	4.5-6.1
	1986-1990	175.2	15.9	14.8-17.0	91.6	20.3	18.4-22.3	83.6	12.5	11.3-13.8	82.6	7.6	6.9-8.4	41.6	9.8	8.5-11.3	41.0	6.1	5.2-7.0
	1991-1995	224.4	20.6	19.4-21.8	120.6	26.5	24.3-28.7	103.8	16.0	14.7-17.5	83.8	7.8	7.1-8.6	42.0	10.1	8.7-11.5	41.8	6.4	5.5-7.3
	1996-2000	234.6	20.8	19.6-22.1	119.8	25.1	23.1-27.2	114.8	17.5	16.0-18.9	83.0	7.6	6.9-8.4	38.4	8.8	7.5-10.1	44.6	6.6	5.8-7.6
	2001-2005	235.0	19.8	18.7-21.0	119.0	24.0	22.0-26.0	116.0	16.7	15.4-18.2	69.4	6.0	5.4-6.7	32.8	6.8	5.8-8.0	36.6	5.2	4.5-6.0
	2006-2010	262.4	20.9	19.8-22.1	139.8	26.0	24.1-28.1	122.6	16.9	15.6-18.3	70.4	5.8	5.2-6.4	34.6	6.9	5.9-8.1	35.8	4.9	4.2-5.7
	2011-2015	263.8	19.3	18.2-20.4	138.0	24.1	22.3-26.0	125.8	16.0	14.7-17.3	70.8	5.4	4.8-6.0	38.8	7.5	6.5-8.7	32.0	4.0	3.4-4.7
2016-2020	277.2	18.9	17.9-19.9	143.8	23.0	21.3-24.8	133.4	15.9	14.7-17.2	68.8	4.7	4.2-5.2	39.2	6.5	5.6-7.5	29.6	3.4	2.9-4.0	
Myeloma	1976-1980	60.8	5.2	4.6-5.8	31.2	6.9	5.8-8.1	29.6	4.1	3.5-4.9	39.8	3.4	2.9-3.9	19.2	4.2	3.4-5.1	20.6	2.9	2.3-3.5
	1981-1985	66.4	6.1	5.4-6.8	29.0	7.0	5.9-8.2	37.4	5.6	4.8-6.5	37.8	3.4	2.9-3.9	18.2	4.3	3.4-5.3	19.6	2.9	2.3-3.5
	1986-1990	60.0	5.5	4.9-6.2	27.0	6.5	5.5-7.7	33.0	5.0	4.2-5.8	36.4	3.4	2.9-3.9	16.4	4.1	3.3-5.1	20.0	2.9	2.4-3.6
	1991-1995	71.8	6.9	6.2-7.6	32.0	7.9	6.7-9.3	39.8	6.2	5.4-7.2	44.6	4.2	3.7-4.8	18.2	4.6	3.7-5.6	26.4	4.1	3.4-4.9
	1996-2000	83.2	7.8	7.1-8.6	37.2	9.0	7.7-10.4	46.0	7.1	6.2-8.1	45.4	4.3	3.7-4.9	20.8	5.0	4.1-6.1	24.6	3.8	3.1-4.5
	2001-2005	109.4	9.6	8.8-10.4	46.6	10.2	8.9-11.6	62.8	9.1	8.1-10.2	45.2	4.0	3.5-4.5	18.4	4.2	3.4-5.2	26.8	3.8	3.2-4.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Bronx**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	128.8	10.6	9.8-11.5	63.6	13.3	11.8-14.8	65.2	9.1	8.1-10.1	45.6	3.9	3.4-4.4	23.8	5.2	4.3-6.3	21.8	3.0	2.5-3.7
	2011-2015	149.6	11.1	10.3-12.0	78.0	14.1	12.7-15.6	71.6	9.2	8.2-10.2	45.6	3.4	3.0-3.9	21.8	4.2	3.4-5.1	23.8	3.0	2.4-3.5
	2016-2020	171.4	11.3	10.6-12.1	85.0	13.4	12.1-14.8	86.4	10.0	9.1-11.0	52.8	3.5	3.1-4.0	29.2	4.8	4.0-5.6	23.6	2.6	2.2-3.2
Leukemias	1976-1980	127.0	10.8	9.9-11.7	66.4	14.1	12.6-15.8	60.6	8.7	7.7-9.7	95.0	8.2	7.4-9.0	49.4	10.6	9.3-12.1	45.6	6.6	5.8-7.6
	1981-1985	128.4	11.4	10.5-12.3	71.4	16.1	14.4-17.9	57.0	8.4	7.4-9.5	85.6	7.6	6.9-8.4	46.4	10.6	9.2-12.1	39.2	5.8	5.0-6.6
	1986-1990	126.0	11.2	10.3-12.1	59.4	13.6	12.0-15.3	66.6	9.8	8.8-10.9	71.8	6.4	5.7-7.1	33.0	7.8	6.6-9.1	38.8	5.6	4.8-6.5
	1991-1995	126.8	11.5	10.6-12.4	62.4	14.5	12.9-16.3	64.4	9.7	8.7-10.8	75.6	6.9	6.2-7.7	40.4	9.8	8.5-11.3	35.2	5.2	4.5-6.1
	1996-2000	134.0	11.8	10.9-12.7	70.8	15.8	14.1-17.6	63.2	9.3	8.3-10.4	68.6	6.3	5.6-7.0	37.0	8.8	7.5-10.2	31.6	4.8	4.0-5.6
	2001-2005	151.4	12.7	11.8-13.7	78.8	16.6	14.9-18.4	72.6	10.3	9.3-11.5	70.8	6.1	5.4-6.7	35.2	7.8	6.7-9.1	35.6	5.0	4.3-5.8
	2006-2010	156.0	12.6	11.7-13.5	83.4	16.4	14.8-18.1	72.6	10.0	9.0-11.1	63.4	5.2	4.7-5.9	30.8	6.4	5.4-7.5	32.6	4.5	3.8-5.2
	2011-2015	182.4	13.4	12.6-14.3	93.8	16.7	15.2-18.4	88.6	11.3	10.3-12.4	72.0	5.4	4.8-6.0	39.8	7.4	6.4-8.5	32.2	4.0	3.4-4.7
	2016-2020	174.2	11.8	11.0-12.6	92.6	14.7	13.4-16.1	81.6	9.7	8.8-10.8	71.8	4.8	4.4-5.4	39.0	6.5	5.6-7.5	32.8	3.8	3.2-4.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Kings (Brooklyn)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	8,375.0	388.4	384.6-392.3	4,037.8	458.9	452.3-465.6	4,337.2	347.6	342.8-352.4	4,629.6	216.5	213.6-219.4	2,390.2	276.5	271.3-281.8	2,239.4	178.4	175.0-181.9
	1981-1985	8,814.4	411.5	407.6-415.5	4,231.6	493.8	486.9-500.7	4,582.8	366.0	361.2-370.9	4,522.0	212.3	209.5-215.2	2,271.6	270.1	265.0-275.4	2,250.4	177.6	174.2-181.0
	1986-1990	8,985.6	416.9	413.0-420.8	4,345.4	506.6	499.7-513.6	4,640.2	366.4	361.6-371.2	4,403.8	206.2	203.5-209.0	2,211.2	265.2	260.1-270.3	2,192.6	171.2	168.0-174.5
	1991-1995	9,775.8	457.7	453.6-461.8	4,752.0	556.8	549.6-564.1	5,023.8	400.3	395.3-405.3	4,295.4	203.3	200.6-206.1	2,127.6	259.9	254.8-265.0	2,167.8	170.4	167.2-173.7
	1996-2000	10,288.2	466.5	462.5-470.6	4,981.0	561.1	554.1-568.2	5,307.2	409.4	404.5-414.4	4,083.0	186.8	184.2-189.4	1,949.2	228.6	224.0-233.2	2,133.8	162.1	159.0-165.2
	2001-2005	10,372.8	442.4	438.6-446.2	5,113.2	535.2	528.6-541.9	5,259.6	384.4	379.8-389.1	3,765.4	161.7	159.4-164.0	1,802.6	197.2	193.1-201.4	1,962.8	140.6	137.8-143.4
	2006-2010	11,036.4	456.9	453.0-460.7	5,421.6	538.8	532.3-545.4	5,614.8	404.7	399.9-409.5	3,565.6	149.0	146.8-151.2	1,717.4	180.2	176.4-184.1	1,848.2	129.9	127.2-132.6
	2011-2015	11,657.4	445.0	441.3-448.6	5,562.8	497.7	491.8-503.8	6,094.6	413.0	408.3-417.8	3,757.4	144.9	142.8-147.0	1,800.8	171.9	168.3-175.5	1,956.6	127.8	125.2-130.4
	2016-2020	11,806.2	423.0	419.6-426.5	5,633.0	463.0	457.5-468.5	6,173.2	399.6	395.0-404.2	3,465.4	123.0	121.1-124.8	1,656.4	143.0	139.9-146.2	1,809.0	110.2	107.9-112.6
Oral cavity and pharynx	1976-1980	221.4	10.3	9.6-10.9	153.6	17.0	15.7-18.3	67.8	5.5	4.9-6.1	92.4	4.2	3.8-4.6	67.4	7.3	6.5-8.1	25.0	2.0	1.6-2.4
	1981-1985	242.8	11.7	11.0-12.4	167.8	19.2	17.8-20.5	75.0	6.2	5.6-6.9	89.2	4.3	3.9-4.7	65.2	7.6	6.7-8.5	24.0	2.0	1.6-2.4
	1986-1990	230.4	10.9	10.3-11.6	154.8	17.7	16.4-19.0	75.6	6.0	5.4-6.7	75.6	3.7	3.3-4.1	54.2	6.4	5.6-7.2	21.4	1.7	1.4-2.1
	1991-1995	244.2	11.7	11.0-12.3	167.2	18.9	17.7-20.3	77.0	6.3	5.7-6.9	80.6	3.9	3.5-4.3	57.2	6.7	5.9-7.5	23.4	1.8	1.5-2.2
	1996-2000	231.8	10.5	9.9-11.1	154.6	16.6	15.5-17.9	77.2	6.0	5.4-6.6	68.6	3.1	2.8-3.5	47.8	5.2	4.6-5.9	20.8	1.6	1.3-2.0
	2001-2005	216.6	9.2	8.6-9.7	140.0	13.8	12.8-14.8	76.6	5.6	5.1-6.2	63.8	2.7	2.4-3.0	44.2	4.4	3.8-5.0	19.6	1.4	1.2-1.7
	2006-2010	227.6	9.3	8.8-9.9	147.2	13.9	12.9-14.9	80.4	5.8	5.2-6.4	58.6	2.4	2.1-2.7	41.0	3.9	3.4-4.5	17.6	1.2	1.0-1.5
	2011-2015	240.8	9.2	8.7-9.7	159.4	13.9	12.9-14.9	81.4	5.5	5.0-6.1	59.2	2.2	2.0-2.5	41.6	3.7	3.2-4.2	17.6	1.1	0.9-1.4
	2016-2020	254.8	9.1	8.6-9.7	165.0	13.6	12.7-14.6	89.8	5.7	5.1-6.2	53.8	1.9	1.7-2.2	37.4	3.2	2.7-3.7	16.4	1.0	0.8-1.2
Esophagus	1976-1980	131.2	6.0	5.5-6.5	93.0	10.2	9.2-11.2	38.2	3.1	2.6-3.5	118.6	5.5	5.0-6.0	85.0	9.4	8.5-10.4	33.6	2.7	2.3-3.2
	1981-1985	132.8	6.1	5.7-6.6	89.6	10.0	9.0-11.0	43.2	3.4	3.0-3.9	122.4	5.7	5.2-6.1	85.4	9.6	8.7-10.6	37.0	2.9	2.5-3.4
	1986-1990	126.2	6.0	5.5-6.5	86.4	10.0	9.0-11.0	39.8	3.1	2.7-3.6	105.2	5.0	4.6-5.4	75.6	8.8	8.0-9.8	29.6	2.3	2.0-2.8
	1991-1995	122.4	5.9	5.4-6.3	81.0	9.5	8.6-10.5	41.4	3.3	2.9-3.8	97.0	4.6	4.2-5.0	66.4	7.7	6.9-8.6	30.6	2.4	2.1-2.8
	1996-2000	107.0	4.9	4.5-5.3	70.2	7.8	7.0-8.7	36.8	2.8	2.4-3.2	89.8	4.1	3.7-4.5	61.6	6.9	6.1-7.7	28.2	2.1	1.8-2.5
	2001-2005	100.6	4.3	3.9-4.7	69.2	7.2	6.4-8.0	31.4	2.3	1.9-2.6	73.8	3.1	2.8-3.5	50.8	5.3	4.6-6.0	23.0	1.6	1.3-2.0
	2006-2010	91.2	3.8	3.4-4.1	63.2	6.3	5.6-7.1	28.0	2.0	1.6-2.3	75.8	3.1	2.8-3.4	53.8	5.4	4.8-6.1	22.0	1.5	1.2-1.8
	2011-2015	91.0	3.5	3.2-3.8	64.0	5.9	5.2-6.6	27.0	1.8	1.5-2.1	61.4	2.4	2.1-2.7	43.6	4.1	3.6-4.7	17.8	1.1	0.9-1.4
	2016-2020	83.8	2.9	2.6-3.2	59.0	4.8	4.3-5.4	24.8	1.5	1.3-1.8	55.0	1.9	1.7-2.1	42.4	3.5	3.0-4.0	12.6	0.8	0.6-1.0
Stomach	1976-1980	334.6	16.2	15.4-17.0	188.4	22.4	20.9-24.0	146.2	12.0	11.1-12.9	241.6	11.7	11.0-12.4	136.4	16.5	15.2-17.9	105.2	8.5	7.8-9.3
	1981-1985	332.4	15.6	14.8-16.4	185.4	22.0	20.6-23.6	147.0	11.6	10.7-12.4	218.8	10.4	9.8-11.0	122.2	14.7	13.5-16.0	96.6	7.7	7.0-8.4
	1986-1990	311.8	14.7	14.0-15.5	174.4	20.9	19.5-22.4	137.4	10.6	9.8-11.5	205.2	9.6	9.1-10.3	119.8	14.6	13.4-15.9	85.4	6.5	5.9-7.1
	1991-1995	316.4	15.0	14.3-15.8	177.2	21.7	20.3-23.2	139.2	10.8	10.0-11.6	205.6	9.9	9.3-10.5	115.8	14.6	13.4-15.8	89.8	7.0	6.3-7.6
	1996-2000	309.4	14.2	13.5-14.9	175.2	20.6	19.2-22.0	134.2	10.1	9.3-10.9	182.8	8.4	7.8-8.9	101.2	12.0	11.0-13.2	81.6	6.1	5.6-6.8
	2001-2005	316.6	13.6	12.9-14.3	170.6	18.4	17.2-19.7	146.0	10.4	9.6-11.2	161.4	6.9	6.5-7.4	84.4	9.3	8.4-10.2	77.0	5.4	4.9-6.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020  
Kings (Brooklyn)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	298.6	12.5	11.9-13.2	169.0	17.6	16.4-18.9	129.6	9.2	8.5-9.9	142.8	6.0	5.6-6.5	82.8	8.9	8.0-9.8	60.0	4.1	3.7-4.6
	2011-2015	310.6	12.0	11.4-12.7	176.6	16.5	15.4-17.7	134.0	8.8	8.2-9.5	143.4	5.6	5.2-6.0	82.8	8.0	7.2-8.8	60.6	3.9	3.5-4.4
	2016-2020	301.8	10.7	10.2-11.3	175.0	14.6	13.6-15.6	126.8	7.9	7.3-8.6	141.4	5.1	4.7-5.5	83.6	7.3	6.6-8.0	57.8	3.5	3.1-4.0
Colorectal	1976-1980	1,393.0	65.2	63.6-66.8	681.0	79.8	77.0-82.7	712.0	56.0	54.2-58.0	709.0	34.3	33.1-35.5	353.6	43.6	41.5-45.8	355.4	28.5	27.2-29.9
	1981-1985	1,411.6	66.3	64.7-67.9	684.8	82.6	79.8-85.6	726.8	56.4	54.6-58.3	628.8	30.0	29.0-31.1	297.6	37.1	35.2-39.1	331.2	25.9	24.7-27.2
	1986-1990	1,295.6	60.4	58.9-61.9	634.8	76.1	73.4-78.8	660.8	50.6	48.9-52.4	568.8	27.0	26.0-28.0	278.0	34.5	32.7-36.4	290.8	22.5	21.3-23.7
	1991-1995	1,305.8	61.7	60.2-63.3	618.6	75.9	73.2-78.7	687.2	53.2	51.4-55.0	529.8	25.3	24.3-26.2	256.0	32.4	30.6-34.2	273.8	21.1	20.0-22.2
	1996-2000	1,360.6	62.4	60.9-63.9	649.0	75.9	73.3-78.6	711.6	53.6	51.9-55.4	498.8	23.0	22.1-23.9	236.8	28.4	26.7-30.0	262.0	19.5	18.4-20.6
	2001-2005	1,326.4	57.0	55.6-58.4	643.0	69.5	67.1-72.0	683.4	48.8	47.2-50.5	469.4	20.2	19.4-21.1	226.8	25.3	23.9-26.9	242.6	17.1	16.1-18.1
	2006-2010	1,210.6	50.5	49.3-51.8	584.6	59.7	57.5-61.9	626.0	44.3	42.8-45.9	417.6	17.5	16.8-18.3	208.6	22.2	20.8-23.6	209.0	14.5	13.6-15.4
	2011-2015	1,088.8	41.8	40.7-42.9	544.0	49.7	47.8-51.7	544.8	36.2	34.8-37.6	377.2	14.4	13.8-15.1	177.4	16.8	15.7-18.0	199.8	12.7	11.9-13.5
	2016-2020	1,047.2	37.7	36.6-38.7	538.4	45.0	43.3-46.8	508.8	32.1	30.9-33.4	350.0	12.4	11.8-13.0	172.0	14.8	13.9-15.9	178.0	10.7	10.0-11.4
Colon excluding rectum	1976-1980	994.0	46.8	45.5-48.2	471.2	56.3	53.9-58.7	522.8	41.1	39.5-42.8	582.6	28.2	27.1-29.2	284.2	35.3	33.4-37.3	298.4	23.8	22.6-25.1
	1981-1985	1,046.4	49.4	48.0-50.8	486.6	59.6	57.2-62.1	559.8	43.4	41.8-45.1	546.0	26.1	25.1-27.1	251.8	31.4	29.7-33.3	294.2	23.0	21.8-24.2
	1986-1990	971.0	45.4	44.1-46.7	462.8	56.0	53.6-58.3	508.2	38.9	37.4-40.5	498.4	23.6	22.7-24.6	244.0	30.3	28.5-32.1	254.4	19.7	18.6-20.8
	1991-1995	994.6	47.1	45.8-48.4	459.6	56.7	54.4-59.2	535.0	41.3	39.7-42.9	472.6	22.5	21.6-23.4	226.8	28.6	27.0-30.4	245.8	18.9	17.8-20.0
	1996-2000	1,031.8	47.4	46.1-48.7	479.2	56.7	54.4-59.0	552.6	41.5	40.0-43.1	442.4	20.4	19.5-21.2	208.6	25.1	23.6-26.7	233.8	17.4	16.4-18.4
	2001-2005	974.4	41.9	40.8-43.1	457.6	50.2	48.1-52.3	516.8	36.8	35.4-38.2	410.6	17.7	17.0-18.5	196.2	22.0	20.6-23.4	214.4	15.1	14.2-16.0
	2006-2010	877.8	36.7	35.6-37.8	404.4	41.9	40.0-43.8	473.4	33.4	32.0-34.8	363.0	15.3	14.6-16.0	180.0	19.3	18.0-20.6	183.0	12.7	11.8-13.5
	2011-2015	775.0	29.8	28.9-30.8	370.0	34.3	32.8-36.0	405.0	26.8	25.6-28.0	315.0	12.0	11.4-12.7	144.2	13.8	12.8-14.8	170.8	10.8	10.1-11.6
	2016-2020	740.8	26.6	25.7-27.4	363.4	30.5	29.1-32.0	377.4	23.7	22.6-24.8	286.4	10.1	9.6-10.6	138.6	12.0	11.1-12.9	147.8	8.8	8.2-9.5
Rectum & rectosigmoid	1976-1980	399.0	18.4	17.6-19.2	209.8	23.6	22.1-25.1	189.2	14.9	13.9-15.9	126.4	6.1	5.6-6.6	69.4	8.3	7.4-9.3	57.0	4.7	4.1-5.3
	1981-1985	365.2	16.9	16.1-17.7	198.2	23.0	21.6-24.5	167.0	13.0	12.1-13.9	82.8	4.0	3.6-4.4	45.8	5.6	4.9-6.5	37.0	2.9	2.5-3.4
	1986-1990	324.6	15.0	14.3-15.7	172.0	20.1	18.7-21.5	152.6	11.7	10.9-12.6	70.4	3.3	3.0-3.7	34.0	4.3	3.6-5.0	36.4	2.8	2.4-3.2
	1991-1995	311.2	14.6	13.9-15.4	159.0	19.1	17.8-20.5	152.2	11.9	11.0-12.7	57.2	2.7	2.4-3.1	29.2	3.7	3.1-4.4	28.0	2.2	1.8-2.6
	1996-2000	328.8	15.0	14.3-15.8	169.8	19.3	18.0-20.6	159.0	12.1	11.3-13.0	56.4	2.6	2.3-2.9	28.2	3.3	2.8-3.9	28.2	2.1	1.8-2.5
	2001-2005	352.0	15.1	14.4-15.8	185.4	19.4	18.1-20.7	166.6	12.0	11.2-12.9	58.8	2.5	2.2-2.8	30.6	3.3	2.8-3.9	28.2	2.0	1.7-2.4
	2006-2010	332.8	13.8	13.1-14.5	180.2	17.8	16.6-19.0	152.6	11.0	10.2-11.8	54.6	2.3	2.0-2.5	28.6	2.9	2.4-3.4	26.0	1.8	1.5-2.2
	2011-2015	313.8	12.0	11.4-12.6	174.0	15.4	14.4-16.5	139.8	9.4	8.7-10.1	62.2	2.4	2.1-2.7	33.2	3.0	2.6-3.5	29.0	1.9	1.6-2.2
	2016-2020	306.4	11.1	10.5-11.7	175.0	14.5	13.6-15.6	131.4	8.5	7.8-9.2	63.6	2.3	2.1-2.6	33.4	2.9	2.5-3.4	30.2	1.9	1.6-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Kings (Brooklyn)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	76.8	3.6	3.3-4.0	43.0	4.9	4.2-5.6	33.8	2.7	2.3-3.2	94.2	4.4	4.0-4.9	48.2	5.4	4.7-6.2	46.0	3.7	3.3-4.3
	1981-1985	82.0	3.8	3.5-4.2	49.8	5.8	5.1-6.6	32.2	2.5	2.1-2.9	109.6	5.2	4.7-5.6	62.4	7.4	6.6-8.3	47.2	3.7	3.2-4.2
	1986-1990	94.8	4.5	4.1-4.9	57.0	6.6	5.8-7.4	37.8	3.0	2.5-3.4	125.2	5.9	5.5-6.4	67.4	8.0	7.1-8.9	57.8	4.5	4.0-5.1
	1991-1995	125.6	5.9	5.4-6.4	77.4	8.9	8.0-9.9	48.2	3.8	3.3-4.3	120.0	5.6	5.2-6.1	71.4	8.3	7.5-9.2	48.6	3.7	3.3-4.2
	1996-2000	176.4	8.0	7.5-8.6	115.8	12.7	11.7-13.8	60.6	4.6	4.1-5.2	133.8	6.1	5.6-6.6	84.2	9.3	8.4-10.2	49.6	3.7	3.3-4.2
	2001-2005	200.6	8.5	8.0-9.1	140.4	14.1	13.1-15.2	60.2	4.4	3.9-4.9	145.2	6.2	5.8-6.7	91.8	9.4	8.6-10.3	53.4	3.8	3.4-4.3
	2006-2010	235.6	9.6	9.0-10.2	167.2	15.7	14.7-16.9	68.4	4.9	4.4-5.4	160.6	6.6	6.1-7.1	107.4	10.4	9.6-11.4	53.2	3.7	3.3-4.2
	2011-2015	288.0	10.7	10.2-11.3	200.4	17.0	16.0-18.1	87.6	5.8	5.3-6.4	182.4	6.9	6.5-7.4	119.2	10.5	9.7-11.5	63.2	4.1	3.7-4.6
2016-2020	276.4	9.6	9.1-10.2	194.2	15.6	14.6-16.6	82.2	5.0	4.5-5.5	167.8	5.8	5.5-6.3	110.2	9.0	8.2-9.8	57.6	3.4	3.0-3.9	
Pancreas	1976-1980	293.2	13.6	12.8-14.3	140.8	15.7	14.5-17.0	152.4	12.0	11.2-12.9	268.0	12.4	11.8-13.1	129.8	14.6	13.5-15.9	138.2	10.9	10.1-11.8
	1981-1985	286.2	13.3	12.6-14.0	133.8	15.9	14.7-17.2	152.4	11.6	10.8-12.5	248.2	11.6	10.9-12.2	120.4	14.3	13.2-15.6	127.8	9.7	9.0-10.5
	1986-1990	262.4	12.3	11.6-13.0	117.6	14.3	13.1-15.5	144.8	11.0	10.2-11.8	234.2	10.9	10.3-11.5	106.6	12.9	11.8-14.1	127.6	9.5	8.8-10.3
	1991-1995	274.0	13.0	12.3-13.7	120.2	14.7	13.5-15.9	153.8	11.8	11.0-12.7	231.4	10.9	10.3-11.6	104.0	12.8	11.7-13.9	127.4	9.8	9.0-10.6
	1996-2000	275.6	12.7	12.0-13.3	123.4	14.6	13.5-15.8	152.2	11.4	10.6-12.3	227.6	10.5	9.9-11.1	98.0	11.7	10.7-12.8	129.6	9.7	8.9-10.5
	2001-2005	280.6	12.1	11.5-12.7	124.2	13.5	12.4-14.6	156.4	11.1	10.3-11.9	224.0	9.7	9.1-10.2	97.8	10.7	9.8-11.7	126.2	8.9	8.2-9.6
	2006-2010	333.8	13.9	13.2-14.6	156.0	16.1	15.0-17.3	177.8	12.3	11.5-13.2	251.8	10.5	9.9-11.1	116.6	12.2	11.2-13.2	135.2	9.3	8.6-10.1
	2011-2015	362.8	14.0	13.3-14.7	169.6	15.7	14.6-16.8	193.2	12.7	11.9-13.5	285.6	11.1	10.5-11.7	132.0	12.5	11.5-13.5	153.6	10.0	9.3-10.8
2016-2020	384.6	13.6	13.0-14.2	180.2	14.9	14.0-16.0	204.4	12.5	11.7-13.3	282.2	10.0	9.5-10.5	134.8	11.5	10.6-12.4	147.4	8.9	8.2-9.5	
Larynx	1976-1980	130.4	5.8	5.4-6.3	107.0	11.4	10.4-12.4	23.4	1.9	1.6-2.3	56.6	2.6	2.3-2.9	46.0	5.0	4.3-5.7	10.6	0.9	0.6-1.1
	1981-1985	126.4	5.8	5.4-6.3	99.8	11.1	10.1-12.2	26.6	2.2	1.8-2.6	49.6	2.3	2.0-2.6	39.0	4.5	3.9-5.2	10.6	0.8	0.6-1.0
	1986-1990	131.4	6.2	5.7-6.7	103.2	11.6	10.6-12.7	28.2	2.3	1.9-2.7	52.6	2.5	2.2-2.8	41.6	4.9	4.2-5.6	11.0	0.9	0.6-1.1
	1991-1995	110.2	5.2	4.8-5.7	87.6	10.1	9.2-11.1	22.6	1.9	1.5-2.3	45.0	2.2	1.9-2.5	36.4	4.3	3.7-5.0	8.6	0.7	0.5-0.9
	1996-2000	103.6	4.7	4.3-5.1	81.0	9.0	8.2-10.0	22.6	1.8	1.5-2.1	43.0	2.0	1.7-2.2	34.8	3.9	3.3-4.5	8.2	0.6	0.5-0.9
	2001-2005	101.4	4.3	3.9-4.7	79.6	8.1	7.3-8.9	21.8	1.6	1.3-1.9	34.8	1.5	1.3-1.7	28.4	3.0	2.5-3.5	6.4	0.5	0.3-0.6
	2006-2010	83.6	3.4	3.1-3.7	64.8	6.3	5.6-7.0	18.8	1.3	1.1-1.7	31.0	1.3	1.1-1.5	24.0	2.5	2.0-3.0	7.0	0.5	0.3-0.7
	2011-2015	77.4	2.9	2.6-3.2	63.4	5.6	5.0-6.3	14.0	0.9	0.7-1.1	24.4	0.9	0.8-1.1	19.4	1.8	1.4-2.2	5.0	0.3	0.2-0.5
2016-2020	65.8	2.3	2.0-2.6	51.2	4.2	3.7-4.8	14.6	0.9	0.7-1.1	21.6	0.8	0.6-0.9	19.2	1.7	1.3-2.0	2.4	0.1	0.1-0.2	
Lung and bronchus	1976-1980	1,138.8	50.9	49.5-52.3	795.6	86.4	83.7-89.3	343.2	26.6	25.3-27.9	938.2	42.0	40.8-43.3	672.8	73.5	70.9-76.1	265.4	20.6	19.5-21.8
	1981-1985	1,175.8	53.5	52.1-54.9	783.6	88.7	85.9-91.6	392.2	30.3	29.0-31.7	954.4	43.5	42.3-44.8	645.8	73.7	71.1-76.3	308.6	23.7	22.5-25.0
	1986-1990	1,220.8	55.9	54.5-57.3	769.6	88.0	85.2-90.9	451.2	34.8	33.4-36.3	983.2	45.2	43.9-46.5	636.8	73.7	71.1-76.4	346.4	26.8	25.5-28.1
	1991-1995	1,267.8	59.7	58.2-61.1	760.6	89.7	86.8-92.6	507.2	40.1	38.6-41.7	1,010.0	47.5	46.2-48.9	625.0	74.6	72.0-77.3	385.0	30.2	28.8-31.5
	1996-2000	1,218.0	55.6	54.2-57.0	683.4	77.4	74.8-80.1	534.6	41.0	39.5-42.6	916.8	41.9	40.7-43.1	531.0	60.7	58.4-63.1	385.8	29.4	28.1-30.8
	2001-2005	1,161.0	49.9	48.6-51.2	641.6	68.3	66.0-70.8	519.4	37.5	36.1-39.0	828.8	35.6	34.6-36.7	478.4	51.5	49.5-53.7	350.4	25.2	24.0-26.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**Kings (Brooklyn)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	1,186.6	49.8	48.6-51.1	636.8	65.7	63.4-68.1	549.8	39.2	37.7-40.7	775.8	32.6	31.6-33.7	424.8	44.5	42.6-46.5	351.0	24.7	23.6-25.9
	2011-2015	1,241.4	48.0	46.8-49.2	666.4	62.4	60.2-64.6	575.0	38.3	36.9-39.7	789.8	30.6	29.7-31.6	440.4	42.1	40.3-43.9	349.4	22.9	21.8-24.0
	2016-2020	1,209.0	42.7	41.6-43.8	627.6	52.9	51.0-54.8	581.4	35.6	34.3-36.9	630.4	22.2	21.4-23.0	349.4	29.9	28.5-31.4	281.0	16.9	16.0-17.8
Melanoma of the skin	1976-1980	105.0	4.8	4.4-5.2	49.8	5.4	4.7-6.1	55.2	4.3	3.8-4.9	45.4	2.1	1.8-2.4	24.6	2.7	2.3-3.3	20.8	1.6	1.3-2.0
	1981-1985	112.8	5.2	4.8-5.7	53.8	5.9	5.2-6.7	59.0	4.8	4.3-5.4	44.0	2.0	1.8-2.3	23.8	2.7	2.2-3.2	20.2	1.6	1.3-1.9
	1986-1990	116.0	5.4	4.9-5.8	57.0	6.4	5.6-7.2	59.0	4.7	4.2-5.3	37.6	1.7	1.5-2.0	20.6	2.3	1.8-2.8	17.0	1.3	1.1-1.7
	1991-1995	107.0	5.0	4.6-5.4	54.2	6.3	5.6-7.1	52.8	4.1	3.6-4.6	30.8	1.4	1.2-1.7	18.0	2.1	1.7-2.6	12.8	1.0	0.8-1.3
	1996-2000	139.4	6.3	5.8-6.8	70.4	7.8	7.0-8.7	69.0	5.3	4.7-5.9	31.8	1.5	1.2-1.7	17.2	1.9	1.5-2.4	14.6	1.1	0.9-1.4
	2001-2005	161.6	6.9	6.4-7.4	81.2	8.4	7.6-9.3	80.4	5.9	5.4-6.5	29.8	1.3	1.1-1.5	13.6	1.5	1.1-1.9	16.2	1.2	0.9-1.4
	2006-2010	229.4	9.4	8.9-10.0	125.2	12.5	11.5-13.5	104.2	7.5	6.8-8.2	33.8	1.4	1.2-1.6	18.0	1.8	1.5-2.2	15.8	1.1	0.9-1.4
	2011-2015	227.4	8.7	8.2-9.3	116.0	10.6	9.7-11.5	111.4	7.6	7.0-8.3	28.8	1.1	0.9-1.3	18.2	1.7	1.4-2.1	10.6	0.7	0.5-1.0
	2016-2020	230.2	8.4	7.9-8.9	122.2	10.4	9.6-11.3	108.0	7.0	6.4-7.6	21.6	0.8	0.6-1.0	12.2	1.1	0.8-1.4	9.4	0.6	0.4-0.8
Female breast	1976-1980							1,158.2	94.8	92.3-97.4							462.4	37.1	35.5-38.7
	1981-1985							1,216.4	100.6	98.0-103.3							444.2	36.1	34.6-37.7
	1986-1990							1,282.2	104.2	101.6-106.8							435.4	35.0	33.5-36.5
	1991-1995							1,443.2	117.9	115.2-120.7							430.8	34.7	33.2-36.2
	1996-2000							1,545.4	121.4	118.7-124.1							402.4	31.1	29.8-32.5
	2001-2005							1,446.2	107.4	105.0-109.9							369.2	26.8	25.6-28.1
	2006-2010							1,508.0	110.5	108.0-113.0							324.0	22.9	21.8-24.1
	2011-2015							1,725.6	119.3	116.7-121.9							330.6	22.1	21.0-23.2
	2016-2020							1,858.8	123.9	121.4-126.5							320.8	20.0	19.0-21.0
Cervix uteri	1976-1980							198.6	17.5	16.4-18.7							67.0	5.7	5.1-6.4
	1981-1985							177.4	15.4	14.4-16.5							51.4	4.5	3.9-5.1
	1986-1990							172.8	14.8	13.8-15.8							49.0	4.2	3.7-4.7
	1991-1995							187.8	15.6	14.6-16.6							52.4	4.3	3.8-4.9
	1996-2000							185.4	14.7	13.8-15.7							51.0	4.0	3.5-4.6
	2001-2005							150.4	11.4	10.6-12.2							44.8	3.3	2.9-3.8
	2006-2010							148.6	11.1	10.3-11.9							45.4	3.3	2.9-3.8
	2011-2015							138.6	9.8	9.0-10.5							45.4	3.1	2.7-3.5
	2016-2020							134.4	9.3	8.6-10.0							34.6	2.3	2.0-2.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**Kings (Brooklyn)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							261.2	20.1	19.0-21.3							67.2	5.1	4.5-5.7
	1981-1985							271.2	21.1	19.9-22.3							69.8	5.2	4.7-5.8
	1986-1990							260.8	20.3	19.2-21.5							64.0	4.8	4.3-5.4
	1991-1995							292.2	23.4	22.2-24.7							69.4	5.5	4.9-6.1
	1996-2000							343.8	26.8	25.5-28.1							83.0	6.3	5.7-7.0
	2001-2005							379.8	27.9	26.7-29.2							81.0	5.9	5.3-6.5
	2006-2010							428.2	30.5	29.2-31.8							90.6	6.4	5.8-7.0
	2011-2015							491.8	32.4	31.1-33.8							111.2	7.2	6.6-7.8
	2016-2020							505.2	31.7	30.5-33.0							129.2	7.8	7.2-8.4
Ovary	1976-1980							182.8	14.5	13.6-15.6							125.0	9.8	9.0-10.6
	1981-1985							190.2	15.2	14.3-16.3							113.2	9.0	8.2-9.8
	1986-1990							174.8	14.0	13.1-15.0							109.0	8.5	7.8-9.3
	1991-1995							204.4	16.5	15.5-17.6							119.0	9.4	8.7-10.2
	1996-2000							179.0	13.9	13.0-14.9							114.4	8.8	8.1-9.6
	2001-2005							164.2	12.1	11.3-13.0							103.0	7.5	6.8-8.1
	2006-2010							169.8	12.3	11.5-13.1							94.2	6.6	6.0-7.2
	2011-2015							174.4	11.8	11.0-12.6							105.8	7.0	6.4-7.6
	2016-2020							166.6	11.0	10.2-11.8							106.2	6.6	6.0-7.2
Prostate	1976-1980				612.0	75.7	72.9-78.6							220.6	29.6	27.7-31.5			
	1981-1985				707.2	86.7	83.8-89.8							225.6	30.2	28.4-32.1			
	1986-1990				794.8	97.7	94.6-100.9							241.0	31.9	30.0-33.8			
	1991-1995				1,198.0	142.7	139.0-146.4							260.2	35.0	33.1-37.0			
	1996-2000				1,491.0	169.3	165.5-173.2							250.2	32.3	30.5-34.2			
	2001-2005				1,588.8	166.5	162.8-170.2							228.2	27.6	26.0-29.2			
	2006-2010				1,679.0	164.1	160.6-167.7							200.0	22.9	21.5-24.4			
	2011-2015				1,579.8	136.7	133.6-139.8							216.0	22.2	20.9-23.6			
	2016-2020				1,630.8	128.9	126.0-131.8							194.4	17.7	16.6-18.8			
Testis	1976-1980				23.6	2.2	1.8-2.6							4.8	0.5	0.3-0.7			
	1981-1985				27.2	2.6	2.1-3.1							3.0	0.3	0.2-0.5			
	1986-1990				32.8	2.9	2.5-3.4							3.0	0.3	0.1-0.4			
	1991-1995				31.2	2.7	2.2-3.1							3.0	0.3	0.2-0.5			
	1996-2000				36.8	3.2	2.7-3.7							1.6	0.2	0.1-0.3			
	2001-2005				38.8	3.2	2.8-3.7							1.2	0.1	0.0-0.2			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020  
Kings (Brooklyn)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				53.4	4.4	3.9-5.0							2.8	0.3	0.1-0.4			
	2011-2015				60.6	4.7	4.1-5.2							2.4	0.2	0.1-0.3			
	2016-2020				70.0	5.4	4.8-6.0							0.8	0.1	0.0-0.2			
Urinary bladder (incl. in situ)	1976-1980	441.8	20.5	19.7-21.5	320.8	37.3	35.4-39.3	121.0	9.6	8.8-10.4	113.6	5.7	5.2-6.2	79.6	10.2	9.2-11.3	34.0	2.9	2.4-3.3
	1981-1985	436.8	20.4	19.5-21.3	310.2	37.4	35.4-39.3	126.6	9.8	9.0-10.6	106.8	5.2	4.8-5.7	71.0	9.3	8.4-10.4	35.8	2.8	2.4-3.3
	1986-1990	421.0	19.6	18.7-20.4	303.2	37.1	35.2-39.1	117.8	9.0	8.2-9.7	99.2	4.8	4.4-5.2	64.4	8.4	7.5-9.4	34.8	2.7	2.3-3.1
	1991-1995	397.6	18.7	17.9-19.6	285.4	35.1	33.3-37.0	112.2	8.7	8.0-9.5	88.4	4.3	3.9-4.7	58.8	7.8	6.9-8.8	29.6	2.3	1.9-2.7
	1996-2000	403.8	18.5	17.7-19.3	290.4	34.5	32.7-36.3	113.4	8.5	7.8-9.3	81.4	3.8	3.4-4.2	50.6	6.5	5.7-7.4	30.8	2.2	1.9-2.6
	2001-2005	400.8	17.3	16.5-18.1	279.4	30.9	29.3-32.6	121.4	8.6	7.9-9.3	74.0	3.2	2.9-3.5	48.8	5.8	5.1-6.6	25.2	1.7	1.4-2.1
	2006-2010	406.6	17.1	16.4-17.9	292.2	31.0	29.4-32.7	114.4	8.0	7.4-8.7	71.0	3.0	2.7-3.3	42.4	4.9	4.3-5.7	28.6	1.9	1.6-2.3
	2011-2015	425.4	16.4	15.7-17.1	312.6	30.1	28.6-31.7	112.8	7.3	6.7-7.9	92.8	3.5	3.2-3.9	59.0	5.9	5.2-6.6	33.8	2.1	1.8-2.4
2016-2020	427.8	15.2	14.6-15.9	320.2	27.6	26.2-29.0	107.6	6.7	6.1-7.3	80.8	2.8	2.6-3.1	54.6	5.0	4.4-5.6	26.2	1.5	1.3-1.8	
Kidney and renal pelvis	1976-1980	141.4	6.3	5.8-6.8	90.8	9.7	8.8-10.7	50.6	3.9	3.4-4.4	73.0	3.3	3.0-3.7	48.4	5.3	4.6-6.0	24.6	1.9	1.6-2.3
	1981-1985	169.6	7.7	7.2-8.3	101.2	11.2	10.2-12.2	68.4	5.4	4.8-6.0	75.2	3.4	3.1-3.8	42.0	4.7	4.1-5.4	33.2	2.6	2.2-3.0
	1986-1990	190.2	8.7	8.1-9.3	111.2	12.4	11.4-13.5	79.0	6.1	5.5-6.8	66.4	3.1	2.7-3.4	39.2	4.6	3.9-5.3	27.2	2.1	1.7-2.4
	1991-1995	219.8	10.2	9.6-10.8	139.2	15.9	14.8-17.2	80.6	6.3	5.7-6.9	68.4	3.2	2.9-3.6	43.6	5.2	4.5-6.0	24.8	1.9	1.6-2.3
	1996-2000	269.8	12.2	11.6-12.9	157.4	17.3	16.1-18.5	112.4	8.6	7.9-9.4	63.8	2.9	2.6-3.2	38.2	4.4	3.7-5.0	25.6	1.9	1.6-2.3
	2001-2005	298.4	12.7	12.1-13.4	180.0	18.6	17.3-19.8	118.4	8.6	8.0-9.4	66.4	2.9	2.6-3.2	38.4	4.1	3.6-4.8	28.0	2.0	1.7-2.4
	2006-2010	339.0	14.1	13.4-14.7	210.2	20.6	19.3-21.9	128.8	9.3	8.6-10.1	67.4	2.8	2.5-3.1	43.0	4.5	3.9-5.2	24.4	1.7	1.4-2.0
	2011-2015	373.2	14.2	13.6-14.9	227.4	19.9	18.8-21.1	145.8	9.8	9.1-10.6	64.2	2.4	2.2-2.7	36.8	3.5	3.0-4.0	27.4	1.7	1.4-2.0
2016-2020	404.0	14.5	13.8-15.1	256.8	20.9	19.8-22.1	147.2	9.4	8.7-10.2	57.6	2.0	1.8-2.3	40.0	3.4	2.9-3.9	17.6	1.1	0.8-1.3	
Brain and other central nervous system	1976-1980	112.4	4.9	4.5-5.3	61.4	6.1	5.4-6.9	51.0	4.0	3.5-4.5	89.8	3.9	3.6-4.3	50.8	5.3	4.7-6.1	39.0	2.9	2.5-3.4
	1981-1985	122.4	5.5	5.1-6.0	65.4	6.9	6.1-7.7	57.0	4.4	3.9-5.0	90.8	4.2	3.8-4.6	49.0	5.4	4.7-6.1	41.8	3.3	2.8-3.8
	1986-1990	136.6	6.1	5.7-6.6	73.2	7.7	6.9-8.6	63.4	5.0	4.5-5.6	78.6	3.6	3.3-4.0	41.6	4.7	4.0-5.4	37.0	2.9	2.5-3.3
	1991-1995	144.0	6.4	6.0-6.9	71.8	7.4	6.6-8.2	72.2	5.7	5.1-6.3	73.6	3.4	3.1-3.8	35.4	3.9	3.3-4.5	38.2	3.0	2.6-3.5
	1996-2000	152.0	6.6	6.2-7.1	73.0	7.3	6.5-8.1	79.0	6.1	5.5-6.7	70.0	3.1	2.8-3.5	35.0	3.8	3.2-4.4	35.0	2.7	2.3-3.1
	2001-2005	143.6	6.0	5.6-6.5	71.2	6.8	6.1-7.6	72.4	5.4	4.9-6.0	62.0	2.6	2.3-2.9	34.2	3.4	2.9-3.9	27.8	2.0	1.7-2.4
	2006-2010	144.8	5.9	5.5-6.4	74.2	6.8	6.1-7.5	70.6	5.2	4.7-5.8	64.6	2.7	2.4-3.0	33.2	3.2	2.7-3.7	31.4	2.3	2.0-2.7
	2011-2015	153.2	5.8	5.4-6.3	84.0	7.1	6.4-7.8	69.2	4.8	4.3-5.4	79.4	3.0	2.7-3.3	40.4	3.5	3.0-4.0	39.0	2.6	2.2-3.0
2016-2020	164.0	6.1	5.7-6.5	88.2	7.2	6.5-7.9	75.8	5.2	4.6-5.7	81.6	3.0	2.7-3.3	44.8	3.7	3.3-4.3	36.8	2.4	2.1-2.8	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020  
Kings (Brooklyn)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	80.0	3.6	3.3-4.0	25.0	2.7	2.2-3.2	55.0	4.4	3.9-5.0	16.6	0.8	0.6-1.0	7.2	0.8	0.6-1.1	9.4	0.8	0.6-1.0
	1981-1985	86.8	4.1	3.7-4.5	20.0	2.2	1.8-2.7	66.8	5.6	5.0-6.2	17.4	0.8	0.6-1.0	6.4	0.8	0.5-1.1	11.0	0.8	0.6-1.1
	1986-1990	86.6	3.9	3.5-4.3	22.8	2.5	2.0-3.0	63.8	5.2	4.6-5.8	8.4	0.4	0.3-0.5	3.0	0.4	0.2-0.6	5.4	0.4	0.3-0.6
	1991-1995	113.0	5.1	4.7-5.5	28.2	3.0	2.5-3.6	84.8	6.8	6.1-7.5	9.6	0.5	0.3-0.6	3.2	0.4	0.2-0.6	6.4	0.5	0.3-0.7
	1996-2000	156.4	6.8	6.3-7.3	33.6	3.4	2.9-4.0	122.8	9.6	8.9-10.4	13.4	0.6	0.5-0.8	4.6	0.5	0.3-0.8	8.8	0.7	0.5-0.9
	2001-2005	222.6	9.2	8.7-9.8	50.2	4.8	4.2-5.4	172.4	13.0	12.1-13.9	13.0	0.6	0.4-0.7	3.4	0.4	0.2-0.6	9.6	0.7	0.5-0.9
	2006-2010	421.0	17.1	16.3-17.8	86.2	7.8	7.1-8.6	334.8	24.9	23.7-26.1	15.2	0.6	0.5-0.8	6.6	0.7	0.5-1.0	8.6	0.6	0.4-0.8
	2011-2015	583.8	22.3	21.5-23.1	125.4	10.6	9.8-11.5	458.4	32.4	31.1-33.8	16.6	0.6	0.5-0.8	5.6	0.5	0.3-0.8	11.0	0.7	0.5-0.9
	2016-2020	543.0	20.3	19.5-21.1	131.6	10.7	9.8-11.5	411.4	28.7	27.5-30.0	14.8	0.5	0.4-0.7	4.2	0.4	0.2-0.6	10.6	0.6	0.5-0.8
Hodgkin lymphoma	1976-1980	74.6	3.2	2.9-3.6	37.0	3.7	3.1-4.3	37.6	2.9	2.5-3.3	31.2	1.4	1.2-1.7	18.0	1.9	1.5-2.4	13.2	1.1	0.8-1.4
	1981-1985	71.0	3.0	2.7-3.4	44.2	4.2	3.6-4.8	26.8	2.1	1.7-2.5	24.8	1.1	0.9-1.3	14.4	1.5	1.2-1.9	10.4	0.8	0.6-1.1
	1986-1990	72.4	3.1	2.8-3.4	43.4	4.2	3.6-4.8	29.0	2.3	1.9-2.7	22.0	1.0	0.8-1.2	14.2	1.5	1.1-1.9	7.8	0.6	0.4-0.8
	1991-1995	63.8	2.7	2.4-3.0	34.6	3.3	2.8-3.8	29.2	2.3	1.9-2.7	15.0	0.7	0.5-0.9	8.0	0.8	0.6-1.1	7.0	0.6	0.4-0.8
	1996-2000	68.4	2.8	2.5-3.2	33.0	3.0	2.6-3.6	35.4	2.7	2.3-3.1	13.4	0.6	0.4-0.7	7.2	0.7	0.5-1.0	6.2	0.5	0.3-0.7
	2001-2005	83.4	3.4	3.1-3.7	45.4	4.0	3.5-4.6	38.0	2.8	2.4-3.3	9.6	0.4	0.3-0.5	5.0	0.5	0.3-0.7	4.6	0.3	0.2-0.5
	2006-2010	82.2	3.2	2.9-3.6	42.2	3.7	3.2-4.2	40.0	2.9	2.5-3.3	12.0	0.5	0.4-0.6	7.8	0.7	0.5-1.0	4.2	0.3	0.2-0.5
	2011-2015	82.8	3.1	2.8-3.5	42.8	3.5	3.0-4.0	40.0	2.8	2.5-3.3	7.4	0.3	0.2-0.4	3.4	0.3	0.2-0.5	4.0	0.3	0.2-0.4
	2016-2020	82.0	3.1	2.8-3.4	44.4	3.6	3.2-4.2	37.6	2.6	2.3-3.0	6.4	0.2	0.2-0.3	3.4	0.3	0.2-0.5	3.0	0.2	0.1-0.3
Non-Hodgkin lymphomas	1976-1980	247.0	11.4	10.8-12.1	121.8	13.6	12.5-14.7	125.2	9.9	9.1-10.7	119.2	5.6	5.1-6.1	64.0	7.2	6.4-8.1	55.2	4.4	3.9-5.0
	1981-1985	297.6	13.7	13.0-14.4	150.6	16.9	15.7-18.2	147.0	11.4	10.5-12.2	141.8	6.6	6.1-7.1	72.2	8.3	7.4-9.2	69.6	5.4	4.9-6.0
	1986-1990	356.6	16.4	15.7-17.2	187.0	20.7	19.4-22.1	169.6	13.2	12.3-14.1	150.0	7.0	6.5-7.5	80.0	9.2	8.3-10.2	70.0	5.5	4.9-6.1
	1991-1995	413.8	19.2	18.4-20.0	217.8	24.0	22.5-25.5	196.0	15.4	14.4-16.4	159.4	7.5	6.9-8.0	82.2	9.4	8.5-10.4	77.2	6.0	5.4-6.7
	1996-2000	433.2	19.5	18.6-20.3	216.4	23.3	21.9-24.8	216.8	16.6	15.6-17.6	162.8	7.4	6.9-7.9	72.8	8.3	7.5-9.2	90.0	6.8	6.2-7.4
	2001-2005	428.0	18.2	17.4-19.0	223.2	22.6	21.3-24.0	204.8	15.0	14.1-15.9	132.4	5.7	5.2-6.1	70.2	7.5	6.7-8.3	62.2	4.4	4.0-5.0
	2006-2010	466.2	19.4	18.6-20.2	238.4	23.3	22.0-24.7	227.8	16.4	15.4-17.3	126.6	5.3	4.9-5.8	61.0	6.4	5.7-7.2	65.6	4.6	4.1-5.2
	2011-2015	504.2	19.3	18.5-20.1	261.0	23.2	21.9-24.5	243.2	16.3	15.4-17.2	120.0	4.6	4.3-5.0	59.8	5.7	5.1-6.4	60.2	3.9	3.5-4.4
	2016-2020	511.4	18.6	17.9-19.4	272.0	22.8	21.6-24.1	239.4	15.4	14.5-16.3	111.0	4.0	3.7-4.3	57.0	4.9	4.4-5.6	54.0	3.3	2.9-3.7
Myeloma	1976-1980	99.8	4.6	4.2-5.0	48.0	5.5	4.8-6.2	51.8	4.0	3.5-4.5	61.0	2.7	2.4-3.1	27.6	3.1	2.6-3.7	33.4	2.5	2.1-2.9
	1981-1985	116.8	5.5	5.0-6.0	50.2	6.1	5.3-6.9	66.6	5.2	4.6-5.8	73.8	3.4	3.1-3.8	28.4	3.3	2.7-3.9	45.4	3.5	3.0-4.0
	1986-1990	115.4	5.4	5.0-5.9	55.2	6.5	5.7-7.3	60.2	4.7	4.2-5.3	63.2	3.0	2.6-3.3	29.0	3.5	2.9-4.1	34.2	2.6	2.2-3.1
	1991-1995	130.0	6.1	5.6-6.6	57.2	6.7	6.0-7.6	72.8	5.7	5.2-6.4	77.8	3.7	3.3-4.1	34.4	4.2	3.6-4.9	43.4	3.4	3.0-3.9
	1996-2000	145.4	6.6	6.2-7.1	61.0	6.8	6.1-7.7	84.4	6.4	5.8-7.1	71.2	3.3	2.9-3.6	29.2	3.4	2.9-4.0	42.0	3.2	2.8-3.6
	2001-2005	163.0	7.0	6.5-7.5	74.6	7.9	7.1-8.8	88.4	6.4	5.8-7.1	67.2	2.9	2.6-3.2	32.2	3.5	3.0-4.1	35.0	2.5	2.2-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Kings (Brooklyn)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	198.0	8.3	7.8-8.8	98.2	9.9	9.0-10.8	99.8	7.1	6.5-7.8	61.4	2.6	2.3-2.9	31.0	3.1	2.7-3.7	30.4	2.1	1.8-2.5
	2011-2015	251.2	9.6	9.1-10.2	125.6	11.5	10.6-12.5	125.6	8.3	7.7-9.0	77.2	3.0	2.7-3.3	37.8	3.6	3.1-4.1	39.4	2.6	2.2-3.0
	2016-2020	248.6	8.9	8.4-9.4	123.0	10.4	9.6-11.2	125.6	7.9	7.3-8.5	78.6	2.7	2.5-3.0	39.8	3.4	3.0-3.9	38.8	2.3	2.0-2.6
Leukemias	1976-1980	241.6	11.2	10.6-11.9	125.8	13.8	12.7-15.0	115.8	9.4	8.6-10.2	163.0	7.7	7.1-8.2	87.4	9.7	8.8-10.8	75.6	6.2	5.6-6.9
	1981-1985	246.8	11.5	10.8-12.2	132.2	15.1	14.0-16.4	114.6	9.2	8.4-10.0	164.2	7.7	7.2-8.3	87.0	10.1	9.1-11.1	77.2	6.2	5.6-6.9
	1986-1990	232.0	10.6	10.0-11.3	112.8	12.8	11.7-13.9	119.2	9.3	8.6-10.1	141.4	6.5	6.0-7.0	71.0	8.2	7.4-9.2	70.4	5.4	4.9-6.0
	1991-1995	254.6	11.8	11.2-12.5	133.2	15.1	14.0-16.4	121.4	9.6	8.8-10.4	136.8	6.4	5.9-6.9	70.8	8.3	7.5-9.3	66.0	5.2	4.6-5.8
	1996-2000	258.8	11.6	10.9-12.2	132.4	14.6	13.4-15.7	126.4	9.6	8.8-10.3	134.8	6.1	5.6-6.6	66.0	7.5	6.7-8.4	68.8	5.2	4.6-5.8
	2001-2005	272.0	11.5	10.9-12.2	142.4	14.7	13.6-15.9	129.6	9.4	8.7-10.2	131.6	5.6	5.2-6.1	70.0	7.5	6.8-8.4	61.6	4.4	3.9-4.9
	2006-2010	300.0	12.5	11.8-13.1	159.6	15.9	14.8-17.1	140.4	10.1	9.3-10.9	126.0	5.2	4.8-5.7	66.0	6.8	6.1-7.6	60.0	4.2	3.8-4.7
	2011-2015	356.6	13.7	13.0-14.3	191.6	17.2	16.1-18.3	165.0	11.1	10.3-11.9	136.8	5.3	4.9-5.7	73.8	7.0	6.3-7.8	63.0	4.1	3.6-4.6
	2016-2020	342.0	12.6	12.0-13.2	187.8	16.0	15.0-17.1	154.2	10.0	9.3-10.8	125.6	4.6	4.2-5.0	65.6	5.7	5.1-6.4	60.0	3.8	3.4-4.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020  
New York (Manhattan)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	6,491.4	425.2	420.4-429.9	3,096.2	505.8	497.6-514.1	3,395.2	384.9	378.9-390.9	3,518.8	232.1	228.6-235.6	1,772.8	296.1	289.7-302.6	1,746.0	194.4	190.2-198.7
	1981-1985	6,666.2	442.7	437.9-447.5	3,252.0	536.1	527.6-544.6	3,414.2	390.8	384.8-396.9	3,358.8	223.7	220.3-227.2	1,664.0	285.1	278.9-291.5	1,694.8	187.7	183.6-191.8
	1986-1990	6,805.4	447.7	442.9-452.6	3,439.6	553.0	544.5-561.6	3,365.8	383.7	377.8-389.6	3,232.2	214.5	211.1-217.8	1,598.8	275.6	269.5-281.9	1,633.4	179.1	175.2-183.1
	1991-1995	7,763.4	517.3	512.1-522.5	3,903.4	626.9	617.9-636.0	3,860.0	448.4	442.1-454.9	3,088.2	208.7	205.4-212.0	1,484.2	260.4	254.4-266.5	1,604.0	179.4	175.5-183.4
	1996-2000	7,500.4	494.7	489.7-499.8	3,638.0	577.2	568.6-585.8	3,862.4	443.7	437.4-450.1	2,872.8	192.3	189.1-195.5	1,366.6	235.2	229.5-240.9	1,506.2	167.4	163.6-171.3
	2001-2005	7,644.2	475.5	470.7-480.3	3,754.6	554.9	546.9-563.0	3,889.6	425.0	419.0-431.1	2,729.0	171.4	168.5-174.3	1,304.4	207.6	202.5-212.8	1,424.6	149.4	145.9-152.9
	2006-2010	8,222.0	485.3	480.6-490.1	4,075.8	561.5	553.7-569.4	4,146.2	437.1	431.0-443.2	2,630.2	155.1	152.4-157.8	1,297.4	191.0	186.3-195.7	1,332.8	132.6	129.4-135.9
	2011-2015	8,323.2	452.6	448.2-457.1	3,996.2	499.0	492.0-506.1	4,327.0	426.9	421.0-432.8	2,496.0	133.4	131.0-135.8	1,193.6	155.9	151.9-159.9	1,302.4	118.9	115.9-121.9
	2016-2020	8,409.0	426.6	422.4-430.8	4,023.0	458.9	452.5-465.4	4,386.0	410.4	404.7-416.1	2,303.2	111.2	109.1-113.2	1,103.8	127.8	124.4-131.2	1,199.4	100.4	97.8-103.1
Oral cavity and pharynx	1976-1980	226.0	15.0	14.1-15.9	152.6	23.8	22.1-25.6	73.4	8.4	7.6-9.4	102.4	6.7	6.2-7.4	73.6	11.5	10.3-12.8	28.8	3.3	2.7-3.9
	1981-1985	232.0	15.6	14.7-16.6	159.0	25.0	23.3-26.9	73.0	8.5	7.6-9.5	85.2	5.8	5.2-6.4	59.8	9.6	8.5-10.8	25.4	2.9	2.4-3.5
	1986-1990	205.6	13.8	13.0-14.7	137.4	21.6	20.0-23.3	68.2	8.0	7.1-8.9	67.8	4.6	4.1-5.1	46.8	7.5	6.6-8.6	21.0	2.4	2.0-2.9
	1991-1995	239.6	16.2	15.3-17.2	159.6	25.1	23.4-27.0	80.0	9.4	8.5-10.4	58.0	3.9	3.5-4.4	39.0	6.5	5.6-7.5	19.0	2.2	1.8-2.7
	1996-2000	195.0	12.8	12.0-13.6	132.4	19.8	18.3-21.4	62.6	7.2	6.4-8.0	52.2	3.5	3.1-3.9	36.0	5.6	4.8-6.5	16.2	1.8	1.4-2.2
	2001-2005	173.2	10.7	10.0-11.5	116.0	16.5	15.2-17.9	57.2	6.2	5.5-7.0	42.6	2.7	2.3-3.0	25.6	3.9	3.2-4.6	17.0	1.8	1.4-2.2
	2006-2010	200.8	11.9	11.2-12.7	138.2	18.6	17.2-20.1	62.6	6.7	6.0-7.5	41.0	2.4	2.1-2.8	29.0	4.1	3.5-4.8	12.0	1.2	0.9-1.6
	2011-2015	202.2	10.9	10.2-11.6	133.4	16.4	15.1-17.7	68.8	6.5	5.8-7.3	45.4	2.4	2.1-2.8	31.6	4.0	3.4-4.7	13.8	1.2	1.0-1.6
	2016-2020	217.0	10.9	10.2-11.5	144.8	16.3	15.1-17.5	72.2	6.5	5.8-7.3	40.8	2.0	1.7-2.2	27.8	3.2	2.7-3.8	13.0	1.0	0.8-1.3
Esophagus	1976-1980	128.4	8.3	7.7-9.0	84.4	13.2	11.9-14.5	44.0	4.9	4.3-5.6	109.0	7.1	6.5-7.8	72.6	11.6	10.4-12.9	36.4	4.1	3.5-4.8
	1981-1985	122.4	8.1	7.5-8.8	84.2	13.4	12.1-14.8	38.2	4.4	3.8-5.1	90.0	6.0	5.4-6.5	62.2	9.9	8.8-11.0	27.8	3.2	2.7-3.8
	1986-1990	107.0	7.1	6.5-7.7	65.6	10.5	9.4-11.8	41.4	4.6	4.0-5.3	92.4	6.1	5.6-6.7	60.4	9.9	8.8-11.1	32.0	3.6	3.0-4.2
	1991-1995	98.4	6.6	6.1-7.2	61.8	10.1	8.9-11.3	36.6	4.2	3.6-4.8	76.2	5.1	4.6-5.7	49.4	8.3	7.3-9.4	26.8	3.0	2.5-3.5
	1996-2000	89.8	6.0	5.4-6.6	59.4	9.6	8.5-10.8	30.4	3.4	2.9-4.0	74.0	4.9	4.4-5.5	48.6	7.9	6.9-9.0	25.4	2.8	2.4-3.4
	2001-2005	81.2	5.0	4.6-5.6	52.0	7.7	6.8-8.7	29.2	3.1	2.6-3.6	65.6	4.1	3.7-4.6	44.8	6.8	5.9-7.7	20.8	2.2	1.8-2.6
	2006-2010	79.2	4.7	4.2-5.2	54.0	7.6	6.7-8.5	25.2	2.5	2.1-3.0	62.8	3.7	3.3-4.2	41.6	5.9	5.1-6.8	21.2	2.1	1.7-2.6
	2011-2015	70.8	3.8	3.4-4.2	48.6	6.1	5.4-6.9	22.2	2.1	1.7-2.5	53.4	2.9	2.6-3.3	36.0	4.6	4.0-5.4	17.4	1.6	1.3-2.0
	2016-2020	63.8	3.1	2.8-3.5	41.8	4.8	4.1-5.5	22.0	1.8	1.5-2.2	44.0	2.1	1.8-2.4	29.6	3.4	2.8-4.0	14.4	1.2	0.9-1.5
Stomach	1976-1980	229.2	15.1	14.2-16.0	133.6	22.4	20.6-24.2	95.6	10.4	9.5-11.4	169.2	11.3	10.5-12.1	97.0	16.5	15.0-18.2	72.2	7.9	7.1-8.8
	1981-1985	197.2	13.2	12.4-14.1	98.2	17.0	15.5-18.7	99.0	10.9	10.0-11.9	138.6	9.3	8.6-10.0	74.6	13.2	11.9-14.7	64.0	7.0	6.2-7.8
	1986-1990	186.8	12.4	11.6-13.2	107.8	18.4	16.8-20.0	79.0	8.5	7.7-9.4	128.0	8.5	7.8-9.1	71.2	12.3	11.1-13.7	56.8	6.0	5.3-6.8
	1991-1995	206.4	13.9	13.1-14.8	109.6	18.8	17.2-20.5	96.8	10.8	9.8-11.8	121.8	8.2	7.6-8.9	61.0	11.0	9.7-12.3	60.8	6.7	5.9-7.5
	1996-2000	178.4	11.9	11.1-12.7	97.4	16.3	14.8-17.8	81.0	9.0	8.1-9.9	112.2	7.5	6.9-8.1	59.2	10.2	9.1-11.5	53.0	5.7	5.1-6.5
	2001-2005	169.8	10.7	10.0-11.4	97.4	14.8	13.5-16.2	72.4	7.7	6.9-8.5	91.4	5.7	5.2-6.3	50.8	8.0	7.0-9.1	40.6	4.2	3.6-4.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York (Manhattan)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	163.8	9.7	9.1-10.4	95.4	13.4	12.2-14.7	68.4	7.0	6.3-7.8	76.4	4.5	4.1-5.0	42.6	6.3	5.4-7.2	33.8	3.3	2.8-3.8
	2011-2015	153.6	8.3	7.7-8.9	93.6	11.9	10.8-13.0	60.0	5.6	5.0-6.3	69.0	3.7	3.3-4.1	38.8	5.0	4.3-5.8	30.2	2.7	2.3-3.2
	2016-2020	142.8	7.2	6.6-7.7	79.8	9.2	8.3-10.1	63.0	5.6	5.0-6.3	67.8	3.3	3.0-3.7	41.8	4.8	4.1-5.5	26.0	2.2	1.8-2.6
Colorectal	1976-1980	1,013.0	66.2	64.3-68.0	481.6	80.6	77.3-84.1	531.4	57.5	55.3-59.8	506.0	33.8	32.5-35.2	247.6	43.4	40.9-46.0	258.4	28.3	26.7-29.9
	1981-1985	961.4	63.5	61.7-65.4	455.4	79.1	75.8-82.5	506.0	54.7	52.5-56.9	449.0	30.1	28.9-31.4	214.2	38.0	35.7-40.4	234.8	25.3	23.8-26.8
	1986-1990	827.6	54.6	52.9-56.3	382.0	65.5	62.5-68.6	445.6	48.1	46.1-50.2	391.2	26.0	24.8-27.1	178.4	32.1	30.0-34.3	212.8	22.6	21.2-24.0
	1991-1995	825.2	55.7	54.0-57.4	376.6	65.5	62.5-68.6	448.6	49.9	47.9-52.1	350.2	23.6	22.5-24.8	161.8	29.7	27.7-31.9	188.4	20.4	19.1-21.7
	1996-2000	805.2	53.8	52.1-55.4	369.8	62.0	59.1-64.9	435.4	48.2	46.2-50.3	322.8	21.6	20.6-22.7	141.8	25.2	23.4-27.2	181.0	19.4	18.2-20.8
	2001-2005	770.6	48.3	46.7-49.8	366.6	56.2	53.7-58.9	404.0	42.7	40.9-44.7	284.0	17.8	16.9-18.8	137.0	22.2	20.5-23.9	147.0	15.0	13.9-16.2
	2006-2010	673.8	39.9	38.5-41.3	327.4	46.1	43.9-48.5	346.4	35.2	33.5-36.9	250.0	14.7	13.9-15.5	116.8	17.3	15.9-18.8	133.2	12.8	11.8-13.8
	2011-2015	645.6	35.2	34.0-36.5	325.6	41.4	39.4-43.5	320.0	30.4	28.9-32.0	223.4	11.9	11.2-12.6	107.0	13.9	12.7-15.1	116.4	10.3	9.5-11.2
	2016-2020	614.6	31.1	30.0-32.3	316.8	36.7	34.9-38.6	297.8	26.8	25.4-28.2	195.0	9.5	8.9-10.1	102.6	12.0	11.0-13.1	92.4	7.6	6.9-8.4
Colon excluding rectum	1976-1980	743.6	48.6	47.0-50.2	345.0	58.1	55.3-61.1	398.6	43.0	41.1-45.0	417.8	28.0	26.7-29.2	199.2	35.2	32.9-37.5	218.6	23.9	22.5-25.4
	1981-1985	700.6	46.3	44.8-47.9	321.4	56.7	53.9-59.7	379.2	40.6	38.8-42.5	381.2	25.5	24.4-26.7	181.2	32.3	30.1-34.5	200.0	21.5	20.2-22.9
	1986-1990	616.8	40.8	39.3-42.2	273.0	47.3	44.7-49.9	343.8	37.1	35.4-38.9	342.8	22.7	21.7-23.8	154.0	27.6	25.7-29.7	188.8	20.0	18.7-21.3
	1991-1995	621.4	42.0	40.5-43.5	276.6	48.6	46.0-51.2	344.8	38.2	36.4-40.1	302.2	20.4	19.4-21.5	138.0	25.4	23.5-27.4	164.2	17.8	16.6-19.1
	1996-2000	598.2	40.0	38.6-41.5	264.0	45.2	42.7-47.8	334.2	36.7	35.0-38.6	281.8	18.9	17.9-19.9	123.0	22.1	20.4-24.0	158.8	17.0	15.9-18.3
	2001-2005	572.8	35.9	34.6-37.3	263.0	40.9	38.7-43.2	309.8	32.5	30.9-34.2	242.6	15.2	14.4-16.1	116.0	18.8	17.3-20.4	126.6	12.9	11.9-14.0
	2006-2010	486.4	28.8	27.7-30.0	230.0	33.0	31.1-35.0	256.4	25.9	24.4-27.3	207.4	12.2	11.4-12.9	94.2	14.0	12.8-15.4	113.2	10.9	10.0-11.8
	2011-2015	454.6	24.7	23.7-25.7	221.0	28.4	26.7-30.1	233.6	21.9	20.7-23.3	180.6	9.6	8.9-10.2	84.8	11.1	10.0-12.2	95.8	8.4	7.7-9.2
	2016-2020	421.4	21.1	20.2-22.1	207.2	24.1	22.6-25.6	214.2	18.9	17.7-20.1	146.2	7.1	6.5-7.6	75.2	8.8	7.9-9.7	71.0	5.8	5.2-6.5
Rectum & rectosigmoid	1976-1980	269.4	17.5	16.6-18.5	136.6	22.5	20.8-24.3	132.8	14.5	13.4-15.6	88.2	5.9	5.3-6.4	48.4	8.2	7.2-9.4	39.8	4.4	3.8-5.0
	1981-1985	260.8	17.2	16.3-18.2	134.0	22.3	20.6-24.1	126.8	14.0	12.9-15.2	67.8	4.6	4.1-5.1	33.0	5.8	4.9-6.8	34.8	3.8	3.2-4.4
	1986-1990	210.8	13.8	13.0-14.7	109.0	18.3	16.7-19.9	101.8	11.0	10.0-12.0	48.4	3.2	2.8-3.7	24.4	4.5	3.7-5.4	24.0	2.6	2.1-3.1
	1991-1995	203.8	13.7	12.9-14.6	100.0	16.9	15.4-18.5	103.8	11.7	10.7-12.7	48.0	3.2	2.8-3.6	23.8	4.3	3.6-5.2	24.2	2.6	2.1-3.1
	1996-2000	207.0	13.7	12.9-14.6	105.8	16.8	15.4-18.3	101.2	11.5	10.5-12.5	41.0	2.7	2.4-3.1	18.8	3.1	2.5-3.8	22.2	2.4	2.0-2.9
	2001-2005	197.8	12.4	11.6-13.2	103.6	15.4	14.1-16.8	94.2	10.2	9.3-11.2	41.4	2.6	2.3-3.0	21.0	3.3	2.7-4.0	20.4	2.1	1.7-2.6
	2006-2010	187.4	11.0	10.3-11.8	97.4	13.2	12.0-14.4	90.0	9.4	8.5-10.3	42.6	2.5	2.2-2.9	22.6	3.3	2.7-3.9	20.0	1.9	1.6-2.4
	2011-2015	191.0	10.5	9.9-11.2	104.6	13.0	11.9-14.2	86.4	8.5	7.7-9.3	42.8	2.3	2.0-2.7	22.2	2.8	2.3-3.4	20.6	1.9	1.5-2.3
	2016-2020	193.2	10.0	9.4-10.7	109.6	12.7	11.6-13.8	83.6	7.9	7.1-8.7	48.8	2.4	2.1-2.8	27.4	3.2	2.7-3.8	21.4	1.8	1.5-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York (Manhattan)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	76.4	4.9	4.4-5.5	53.4	8.5	7.5-9.6	23.0	2.5	2.0-3.0	75.6	5.0	4.5-5.5	49.4	8.0	7.0-9.1	26.2	2.9	2.4-3.5
	1981-1985	80.4	5.3	4.8-5.9	55.2	8.9	7.9-10.1	25.2	2.8	2.3-3.3	84.2	5.5	5.0-6.1	51.2	8.4	7.4-9.5	33.0	3.5	3.0-4.1
	1986-1990	93.8	6.2	5.7-6.8	62.0	10.3	9.1-11.5	31.8	3.5	2.9-4.1	90.0	5.9	5.4-6.5	56.8	9.4	8.4-10.6	33.2	3.6	3.1-4.2
	1991-1995	104.8	7.1	6.5-7.7	67.6	10.9	9.7-12.1	37.2	4.3	3.7-4.9	94.8	6.4	5.9-7.1	58.6	9.8	8.7-11.0	36.2	4.1	3.5-4.7
	1996-2000	144.6	9.6	8.9-10.3	100.6	15.6	14.2-17.0	44.0	4.9	4.3-5.6	101.8	6.8	6.2-7.4	65.8	10.6	9.5-11.8	36.0	4.0	3.4-4.6
	2001-2005	165.2	10.3	9.6-11.1	122.0	17.4	16.0-18.8	43.2	4.7	4.1-5.4	109.2	6.8	6.3-7.4	73.4	10.8	9.7-12.0	35.8	3.8	3.2-4.4
	2006-2010	212.0	12.5	11.7-13.2	152.8	20.3	18.9-21.8	59.2	6.1	5.4-6.8	126.0	7.4	6.9-8.0	89.8	12.3	11.2-13.5	36.2	3.7	3.2-4.3
	2011-2015	211.2	11.1	10.5-11.8	143.2	17.2	16.0-18.6	68.0	6.1	5.5-6.8	128.8	6.8	6.3-7.4	84.0	10.4	9.4-11.4	44.8	4.0	3.5-4.6
	2016-2020	197.0	9.6	9.0-10.3	131.0	14.6	13.5-15.8	66.0	5.6	5.0-6.3	115.0	5.5	5.0-6.0	73.4	8.3	7.4-9.2	41.6	3.4	2.9-3.9
Pancreas	1976-1980	215.4	14.1	13.2-15.0	102.2	17.3	15.7-18.9	113.2	12.1	11.1-13.1	200.4	13.1	12.3-14.0	95.0	16.0	14.5-17.5	105.4	11.3	10.4-12.4
	1981-1985	212.2	14.0	13.2-14.9	99.8	16.8	15.3-18.4	112.4	12.0	11.0-13.1	180.2	11.9	11.2-12.7	83.0	14.3	12.9-15.7	97.2	10.3	9.4-11.3
	1986-1990	194.4	13.0	12.1-13.8	86.6	15.0	13.6-16.5	107.8	11.6	10.7-12.7	164.8	11.0	10.2-11.7	73.0	12.7	11.4-14.1	91.8	9.9	9.0-10.8
	1991-1995	190.0	12.9	12.1-13.7	88.8	15.2	13.8-16.7	101.2	11.1	10.2-12.2	160.0	10.9	10.1-11.6	72.2	12.6	11.3-14.0	87.8	9.5	8.7-10.5
	1996-2000	200.8	13.5	12.7-14.4	87.8	14.8	13.4-16.3	113.0	12.6	11.5-13.6	166.4	11.1	10.4-11.9	75.0	12.9	11.6-14.3	91.4	10.0	9.1-10.9
	2001-2005	206.0	12.9	12.2-13.8	94.2	14.6	13.3-16.0	111.8	11.7	10.7-12.7	176.6	11.1	10.4-11.9	77.0	12.2	11.0-13.5	99.6	10.2	9.3-11.1
	2006-2010	241.2	14.2	13.4-15.1	117.8	16.8	15.5-18.3	123.4	12.3	11.3-13.3	195.8	11.5	10.8-12.3	91.6	13.3	12.1-14.6	104.2	10.2	9.3-11.1
	2011-2015	246.4	13.2	12.5-14.0	125.6	16.0	14.7-17.3	120.8	11.1	10.2-12.0	190.6	10.2	9.5-10.8	93.0	12.0	10.9-13.2	97.6	8.7	8.0-9.6
	2016-2020	263.2	12.8	12.1-13.5	129.0	14.8	13.7-16.0	134.2	11.3	10.4-12.2	203.2	9.7	9.1-10.3	97.6	11.2	10.3-12.3	105.6	8.6	7.9-9.4
Larynx	1976-1980	108.0	7.0	6.4-7.7	87.4	13.6	12.3-14.9	20.6	2.4	1.9-2.9	47.0	3.0	2.6-3.4	38.4	6.0	5.2-6.9	8.6	1.0	0.7-1.4
	1981-1985	106.0	7.1	6.5-7.7	84.8	13.2	11.9-14.5	21.2	2.4	2.0-3.0	41.0	2.7	2.4-3.1	32.6	5.1	4.4-6.0	8.4	1.0	0.7-1.3
	1986-1990	123.0	8.4	7.7-9.1	88.4	13.9	12.6-15.3	34.6	4.4	3.8-5.2	41.6	2.8	2.4-3.2	32.2	5.2	4.4-6.1	9.4	1.1	0.8-1.4
	1991-1995	101.4	6.8	6.2-7.5	73.8	11.8	10.6-13.1	27.6	3.3	2.8-3.9	44.6	3.0	2.6-3.4	32.6	5.3	4.5-6.2	12.0	1.4	1.1-1.9
	1996-2000	80.6	5.3	4.8-5.9	63.0	9.9	8.8-11.1	17.6	2.1	1.6-2.5	32.0	2.1	1.8-2.5	25.0	4.1	3.4-4.9	7.0	0.8	0.6-1.1
	2001-2005	63.6	4.0	3.6-4.4	50.0	7.5	6.5-8.5	13.6	1.5	1.2-2.0	23.2	1.4	1.2-1.7	18.0	2.7	2.2-3.3	5.2	0.5	0.3-0.8
	2006-2010	61.4	3.6	3.2-4.0	48.4	6.6	5.8-7.5	13.0	1.3	1.0-1.7	21.4	1.3	1.0-1.5	16.6	2.4	1.9-3.0	4.8	0.5	0.3-0.7
	2011-2015	56.0	3.0	2.6-3.4	45.0	5.5	4.8-6.3	11.0	1.0	0.8-1.4	17.0	0.9	0.7-1.1	13.2	1.7	1.3-2.2	3.8	0.4	0.2-0.6
	2016-2020	44.0	2.2	1.9-2.5	34.6	3.9	3.3-4.5	9.4	0.9	0.6-1.2	13.6	0.6	0.5-0.8	10.4	1.1	0.8-1.5	3.2	0.2	0.1-0.4
Lung and bronchus	1976-1980	877.8	56.2	54.5-57.9	579.8	91.2	87.8-94.7	298.0	32.6	30.9-34.3	686.0	44.0	42.6-45.6	459.4	73.1	70.1-76.3	226.6	24.7	23.2-26.2
	1981-1985	913.8	59.9	58.2-61.7	561.6	91.5	88.1-95.0	352.2	39.1	37.2-41.0	714.2	46.9	45.4-48.5	435.0	71.7	68.7-74.9	279.2	30.7	29.1-32.4
	1986-1990	882.2	58.2	56.5-59.9	521.2	85.3	82.0-88.7	361.0	40.0	38.1-41.9	722.2	47.6	46.1-49.2	431.2	71.9	68.9-75.1	291.0	31.9	30.3-33.7
	1991-1995	939.8	63.4	61.6-65.3	518.4	85.9	82.6-89.3	421.4	48.0	46.0-50.2	713.4	48.1	46.5-49.7	398.2	67.6	64.6-70.7	315.2	35.4	33.6-37.2
	1996-2000	892.0	59.7	58.0-61.5	458.6	74.7	71.6-77.9	433.4	49.3	47.2-51.4	637.0	42.8	41.3-44.3	335.6	56.0	53.3-58.8	301.4	33.8	32.1-35.6
	2001-2005	907.4	57.2	55.6-58.9	458.0	69.8	67.0-72.8	449.4	48.4	46.4-50.5	636.8	40.2	38.8-41.6	327.8	51.0	48.5-53.6	309.0	32.6	31.0-34.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York (Manhattan)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	891.2	52.8	51.3-54.4	435.2	62.1	59.4-64.7	456.0	46.5	44.5-48.4	586.0	34.6	33.4-35.9	300.0	43.5	41.3-45.8	286.0	28.5	27.0-30.0
	2011-2015	916.0	49.2	47.7-50.6	411.8	52.9	50.6-55.3	504.2	47.0	45.2-48.9	538.0	28.7	27.6-29.8	255.8	33.3	31.5-35.2	282.2	25.6	24.2-27.0
	2016-2020	882.2	42.8	41.6-44.1	394.8	45.3	43.3-47.4	487.4	41.1	39.5-42.8	422.8	20.2	19.3-21.1	198.0	22.8	21.3-24.2	224.8	18.3	17.2-19.4
Melanoma of the skin	1976-1980	103.0	6.7	6.1-7.3	53.4	8.0	7.1-9.1	49.6	5.9	5.1-6.7	30.2	2.0	1.7-2.3	17.2	2.7	2.2-3.4	13.0	1.5	1.1-1.9
	1981-1985	108.0	7.1	6.5-7.7	58.2	9.0	8.0-10.1	49.8	5.8	5.1-6.6	38.6	2.6	2.2-3.0	22.2	3.5	2.9-4.3	16.4	1.9	1.5-2.3
	1986-1990	101.2	6.5	5.9-7.1	55.4	8.6	7.6-9.7	45.8	5.1	4.4-5.8	30.6	2.0	1.7-2.4	18.2	3.0	2.4-3.8	12.4	1.4	1.1-1.8
	1991-1995	119.8	7.8	7.2-8.5	71.0	10.9	9.8-12.1	48.8	5.6	4.9-6.3	26.6	1.8	1.5-2.1	15.4	2.4	1.9-3.1	11.2	1.2	0.9-1.6
	1996-2000	178.6	11.3	10.6-12.1	103.8	16.1	14.7-17.6	74.8	8.2	7.4-9.1	31.2	2.1	1.8-2.4	18.6	3.1	2.5-3.9	12.6	1.4	1.1-1.8
	2001-2005	275.4	16.4	15.5-17.3	147.0	20.9	19.4-22.5	128.4	13.7	12.6-14.8	27.8	1.7	1.5-2.1	15.4	2.5	1.9-3.1	12.4	1.3	1.0-1.7
	2006-2010	359.2	20.6	19.7-21.6	195.4	25.9	24.3-27.7	163.8	17.1	15.9-18.3	30.2	1.8	1.5-2.1	19.6	2.9	2.3-3.5	10.6	1.1	0.8-1.5
	2011-2015	343.8	18.5	17.6-19.4	196.6	24.7	23.2-26.3	147.2	14.1	13.1-15.2	25.2	1.3	1.1-1.6	14.2	1.8	1.4-2.3	11.0	1.0	0.7-1.3
2016-2020	340.2	17.2	16.3-18.0	195.2	22.5	21.1-24.0	145.0	13.3	12.4-14.4	22.8	1.1	0.9-1.3	14.0	1.6	1.3-2.1	8.8	0.7	0.5-1.0	
Female breast	1976-1980							900.2	106.8	103.6-110.1							343.8	39.4	37.5-41.4
	1981-1985							946.2	114.1	110.7-117.5							328.2	37.7	35.8-39.6
	1986-1990							992.8	117.9	114.6-121.3							324.6	36.6	34.8-38.4
	1991-1995							1,119.6	134.7	131.1-138.3							308.8	35.5	33.7-37.3
	1996-2000							1,213.0	142.6	139.0-146.3							270.2	30.4	28.8-32.1
	2001-2005							1,148.4	128.4	125.1-131.8							244.4	25.9	24.5-27.4
	2006-2010							1,268.2	137.7	134.3-141.2							218.6	22.1	20.8-23.4
	2011-2015							1,352.6	138.6	135.2-142.1							206.8	19.4	18.2-20.7
2016-2020							1,399.6	138.1	134.8-141.5							185.8	16.3	15.2-17.4	
Cervix uteri	1976-1980							141.0	17.4	16.1-18.8							52.0	6.1	5.4-7.0
	1981-1985							111.2	13.8	12.6-15.0							38.2	4.6	3.9-5.3
	1986-1990							99.6	11.9	10.8-13.0							29.6	3.5	2.9-4.1
	1991-1995							112.2	13.5	12.4-14.6							30.6	3.6	3.1-4.3
	1996-2000							100.0	11.6	10.6-12.7							30.8	3.6	3.0-4.2
	2001-2005							82.6	9.3	8.4-10.2							28.6	3.2	2.7-3.8
	2006-2010							63.8	7.1	6.4-8.0							25.8	2.8	2.3-3.3
	2011-2015							68.4	7.1	6.3-7.9							21.8	2.2	1.8-2.7
	2016-2020							62.0	6.5	5.8-7.3							17.4	1.6	1.3-2.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.  
Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.  
The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York (Manhattan)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							206.4	22.7	21.3-24.2							65.6	7.0	6.2-7.8
	1981-1985							170.4	19.3	18.0-20.7							46.8	4.9	4.3-5.6
	1986-1990							145.2	16.6	15.4-17.9							35.6	3.7	3.2-4.3
	1991-1995							180.2	21.4	20.0-22.8							44.2	4.9	4.3-5.6
	1996-2000							213.6	24.8	23.4-26.4							50.0	5.6	5.0-6.4
	2001-2005							244.2	27.0	25.5-28.5							46.2	5.0	4.4-5.7
	2006-2010							270.4	28.0	26.5-29.5							53.6	5.3	4.7-6.0
	2011-2015							298.6	28.9	27.4-30.5							65.6	6.0	5.3-6.7
	2016-2020							315.6	29.3	27.8-30.8							59.2	5.0	4.5-5.7
Ovary	1976-1980							139.0	16.2	15.0-17.5							84.8	9.5	8.6-10.5
	1981-1985							150.4	17.6	16.3-18.9							85.4	9.5	8.6-10.5
	1986-1990							127.8	14.9	13.7-16.1							84.2	9.5	8.6-10.4
	1991-1995							151.0	17.9	16.6-19.2							89.4	10.3	9.3-11.3
	1996-2000							137.2	16.1	14.9-17.3							77.8	8.9	8.0-9.8
	2001-2005							137.8	15.4	14.3-16.6							74.6	8.1	7.3-9.0
	2006-2010							126.2	13.5	12.4-14.6							74.6	7.6	6.8-8.4
	2011-2015							129.4	13.1	12.1-14.2							75.2	7.2	6.5-8.0
	2016-2020							105.6	10.0	9.2-11.0							72.8	6.3	5.6-7.0
Prostate	1976-1980				520.2	89.3	85.8-93.0							170.6	31.7	29.5-34.0			
	1981-1985				543.2	95.2	91.6-99.0							197.0	38.0	35.6-40.5			
	1986-1990				578.2	101.5	97.7-105.3							200.8	38.7	36.3-41.3			
	1991-1995				895.8	151.6	147.1-156.2							211.4	41.3	38.8-43.9			
	1996-2000				1,060.0	170.6	166.0-175.3							183.0	35.2	32.9-37.5			
	2001-2005				1,070.4	157.9	153.7-162.3							174.2	30.8	28.8-32.9			
	2006-2010				1,198.8	163.0	158.8-167.2							166.6	26.8	25.0-28.7			
	2011-2015				1,068.4	129.0	125.5-132.6							143.8	19.6	18.2-21.1			
	2016-2020				1,145.6	126.3	123.0-129.6							137.4	16.0	14.8-17.2			
Testis	1976-1980				22.4	2.9	2.4-3.6							4.4	0.6	0.4-0.9			
	1981-1985				32.2	3.8	3.2-4.5							2.6	0.4	0.2-0.7			
	1986-1990				35.8	4.2	3.6-4.9							2.2	0.3	0.2-0.6			
	1991-1995				37.4	4.3	3.7-5.0							2.0	0.3	0.1-0.6			
	1996-2000				41.8	4.6	3.9-5.3							1.8	0.2	0.1-0.5			
	2001-2005				51.4	5.6	4.9-6.3							1.6	0.2	0.1-0.4			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020  
New York (Manhattan)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				56.4	6.2	5.5-7.0							0.6	0.1	0.0-0.2			
	2011-2015				59.0	6.4	5.6-7.2							1.0	0.1	0.0-0.3			
	2016-2020				65.4	7.2	6.4-8.0							1.0	0.1	0.0-0.2			
Urinary bladder (incl. in situ)	1976-1980	294.0	19.1	18.2-20.2	199.4	33.9	31.7-36.2	94.6	10.1	9.2-11.1	92.2	6.4	5.8-7.0	59.4	10.9	9.7-12.3	32.8	3.7	3.2-4.4
	1981-1985	279.6	18.4	17.5-19.4	186.4	32.2	30.2-34.4	93.2	9.9	9.0-10.9	79.2	5.3	4.8-5.9	47.6	8.9	7.8-10.2	31.6	3.4	2.9-3.9
	1986-1990	247.6	16.4	15.5-17.3	166.2	29.1	27.1-31.2	81.4	8.8	7.9-9.7	61.8	4.1	3.7-4.6	37.2	7.1	6.1-8.2	24.6	2.5	2.1-3.1
	1991-1995	284.6	19.2	18.2-20.3	188.0	33.2	31.0-35.4	96.6	10.7	9.7-11.7	65.8	4.4	4.0-5.0	36.4	6.9	5.9-7.9	29.4	3.1	2.6-3.6
	1996-2000	277.8	18.7	17.7-19.7	186.0	31.5	29.5-33.6	91.8	10.3	9.3-11.3	66.6	4.5	4.0-5.0	45.4	8.5	7.4-9.7	21.2	2.2	1.8-2.7
	2001-2005	284.4	17.9	17.0-18.9	199.0	31.3	29.3-33.3	85.4	9.0	8.1-9.9	64.6	4.1	3.6-4.5	38.6	6.5	5.6-7.5	26.0	2.5	2.1-3.0
	2006-2010	309.4	18.3	17.4-19.2	227.2	33.3	31.3-35.3	82.2	8.2	7.4-9.1	72.4	4.2	3.8-4.7	51.0	7.9	6.9-8.9	21.4	2.0	1.6-2.4
	2011-2015	319.6	17.2	16.3-18.1	235.8	31.0	29.2-32.9	83.8	7.6	6.9-8.4	65.0	3.4	3.0-3.8	44.8	6.1	5.3-7.0	20.2	1.7	1.4-2.1
2016-2020	345.0	16.7	15.9-17.5	245.0	28.4	26.8-30.1	100.0	8.4	7.7-9.2	64.6	2.9	2.6-3.3	41.8	4.9	4.2-5.6	22.8	1.7	1.4-2.1	
Kidney and renal pelvis	1976-1980	107.8	7.0	6.4-7.6	67.8	10.6	9.5-11.9	40.0	4.4	3.8-5.1	49.8	3.3	2.9-3.7	33.8	5.4	4.6-6.4	16.0	1.8	1.4-2.2
	1981-1985	113.6	7.6	7.0-8.3	71.8	11.8	10.6-13.1	41.8	4.7	4.1-5.4	46.0	3.0	2.6-3.4	28.2	4.7	4.0-5.6	17.8	1.9	1.5-2.4
	1986-1990	126.8	8.4	7.8-9.1	78.6	12.9	11.6-14.2	48.2	5.5	4.8-6.2	50.8	3.4	3.0-3.8	29.0	5.1	4.3-6.0	21.8	2.4	1.9-2.9
	1991-1995	144.8	9.8	9.1-10.6	85.8	14.1	12.7-15.5	59.0	6.8	6.1-7.7	43.0	2.9	2.5-3.4	26.0	4.5	3.7-5.3	17.0	1.9	1.5-2.4
	1996-2000	169.4	11.3	10.6-12.1	101.4	16.2	14.8-17.7	68.0	7.9	7.1-8.8	45.8	3.1	2.7-3.5	26.0	4.3	3.6-5.2	19.8	2.2	1.8-2.7
	2001-2005	198.8	12.5	11.7-13.3	126.2	18.3	16.9-19.8	72.6	7.9	7.1-8.8	38.4	2.4	2.1-2.8	23.8	3.6	3.0-4.3	14.6	1.5	1.2-1.9
	2006-2010	218.2	13.1	12.3-13.9	142.0	19.7	18.3-21.2	76.2	8.1	7.2-8.9	40.8	2.4	2.1-2.8	25.4	3.7	3.1-4.5	15.4	1.5	1.2-1.9
	2011-2015	232.4	12.8	12.1-13.6	151.2	18.7	17.4-20.1	81.2	8.1	7.3-8.9	37.0	2.0	1.7-2.3	25.8	3.3	2.7-3.9	11.2	1.0	0.8-1.3
2016-2020	238.0	12.3	11.6-13.0	153.8	17.7	16.5-19.1	84.2	8.0	7.2-8.8	36.2	1.7	1.5-2.0	22.0	2.5	2.1-3.1	14.2	1.1	0.9-1.5	
Brain and other central nervous system	1976-1980	87.4	5.8	5.2-6.3	47.4	7.2	6.3-8.2	40.0	4.6	4.0-5.4	69.2	4.5	4.0-5.0	37.0	5.6	4.8-6.5	32.2	3.6	3.1-4.3
	1981-1985	89.2	6.2	5.6-6.8	45.2	7.0	6.1-8.0	44.0	5.6	4.8-6.4	64.8	4.3	3.8-4.8	35.2	5.6	4.8-6.5	29.6	3.3	2.8-3.9
	1986-1990	99.2	6.7	6.1-7.3	52.4	8.0	7.0-9.1	46.8	5.7	4.9-6.5	52.6	3.5	3.1-4.0	29.8	4.8	4.0-5.6	22.8	2.6	2.2-3.2
	1991-1995	112.4	7.6	7.0-8.3	60.8	9.2	8.1-10.3	51.6	6.4	5.6-7.3	50.6	3.4	3.0-3.9	27.0	4.3	3.6-5.2	23.6	2.8	2.3-3.3
	1996-2000	124.4	8.3	7.6-9.0	65.0	9.4	8.4-10.5	59.4	7.3	6.5-8.2	53.0	3.6	3.1-4.0	27.6	4.3	3.6-5.1	25.4	3.0	2.5-3.5
	2001-2005	110.8	7.0	6.4-7.6	57.2	8.2	7.2-9.2	53.6	6.2	5.5-7.0	52.2	3.3	2.9-3.7	28.0	4.1	3.4-4.9	24.2	2.7	2.2-3.2
	2006-2010	103.6	6.5	5.9-7.1	53.0	7.4	6.5-8.3	50.6	5.8	5.0-6.6	55.4	3.3	2.9-3.7	29.6	4.0	3.4-4.7	25.8	2.6	2.2-3.1
	2011-2015	113.8	6.7	6.2-7.3	62.0	8.2	7.3-9.2	51.8	5.6	4.9-6.3	55.0	3.0	2.7-3.4	28.2	3.7	3.1-4.4	26.8	2.5	2.1-3.0
2016-2020	117.2	6.5	6.0-7.1	68.6	8.3	7.4-9.2	48.6	5.0	4.4-5.8	64.8	3.4	3.0-3.8	37.8	4.5	3.8-5.2	27.0	2.6	2.1-3.1	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020  
New York (Manhattan)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	57.0	3.7	3.3-4.2	15.6	2.3	1.8-2.9	41.4	4.8	4.2-5.6	10.6	0.7	0.5-0.9	4.0	0.7	0.4-1.1	6.6	0.7	0.5-1.0
	1981-1985	60.0	4.0	3.5-4.5	19.8	2.8	2.3-3.5	40.2	5.0	4.3-5.7	8.8	0.6	0.4-0.8	2.6	0.5	0.3-0.9	6.2	0.7	0.5-1.0
	1986-1990	67.8	4.3	3.8-4.8	19.8	2.9	2.3-3.5	48.0	5.6	4.9-6.4	9.0	0.6	0.4-0.8	1.8	0.3	0.1-0.6	7.2	0.8	0.5-1.1
	1991-1995	93.4	5.9	5.3-6.4	25.8	3.7	3.1-4.4	67.6	7.9	7.0-8.8	6.8	0.5	0.3-0.6	2.0	0.3	0.2-0.6	4.8	0.5	0.3-0.8
	1996-2000	136.0	8.2	7.6-8.8	35.6	4.7	4.0-5.4	100.4	11.4	10.4-12.4	8.4	0.6	0.4-0.8	2.8	0.5	0.3-0.8	5.6	0.6	0.4-0.9
	2001-2005	187.2	11.0	10.3-11.8	49.4	6.5	5.7-7.4	137.8	15.1	14.0-16.3	9.2	0.6	0.4-0.8	4.2	0.7	0.4-1.0	5.0	0.5	0.3-0.7
	2006-2010	315.8	18.1	17.2-19.0	81.2	10.1	9.1-11.1	234.6	25.5	24.0-27.0	10.2	0.6	0.4-0.8	4.8	0.7	0.5-1.1	5.4	0.5	0.4-0.8
	2011-2015	352.4	19.7	18.7-20.6	96.2	11.5	10.5-12.6	256.2	27.4	25.8-29.0	10.0	0.5	0.4-0.7	4.6	0.6	0.4-0.9	5.4	0.5	0.3-0.7
2016-2020	326.4	17.9	17.0-18.8	89.4	10.4	9.4-11.4	237.0	24.9	23.5-26.5	7.8	0.4	0.3-0.5	2.8	0.3	0.2-0.6	5.0	0.4	0.3-0.6	
Hodgkin lymphoma	1976-1980	53.6	3.3	2.9-3.8	27.6	3.9	3.2-4.6	26.0	2.9	2.4-3.5	17.0	1.1	0.9-1.4	8.8	1.3	0.9-1.7	8.2	0.9	0.7-1.3
	1981-1985	56.8	3.5	3.1-3.9	32.0	4.4	3.8-5.2	24.8	2.7	2.2-3.3	17.2	1.1	0.9-1.3	9.2	1.3	0.9-1.7	8.0	0.9	0.6-1.2
	1986-1990	52.8	3.3	2.9-3.7	34.8	4.8	4.1-5.6	18.0	2.0	1.6-2.5	15.0	1.0	0.7-1.2	10.6	1.6	1.2-2.1	4.4	0.5	0.3-0.7
	1991-1995	61.4	3.9	3.4-4.3	33.4	4.5	3.9-5.3	28.0	3.2	2.7-3.9	9.6	0.6	0.4-0.8	5.2	0.8	0.5-1.1	4.4	0.5	0.3-0.7
	1996-2000	54.8	3.3	2.9-3.8	29.4	3.8	3.2-4.5	25.4	2.9	2.4-3.5	8.0	0.5	0.4-0.7	3.6	0.5	0.3-0.8	4.4	0.5	0.3-0.7
	2001-2005	59.6	3.5	3.1-3.9	31.6	4.0	3.4-4.7	28.0	3.0	2.5-3.6	8.2	0.5	0.4-0.7	5.0	0.7	0.5-1.1	3.2	0.4	0.2-0.6
	2006-2010	62.0	3.5	3.1-4.0	35.2	4.4	3.7-5.1	26.8	2.7	2.3-3.3	5.4	0.3	0.2-0.4	2.6	0.3	0.2-0.6	2.8	0.3	0.1-0.5
	2011-2015	62.8	3.5	3.1-3.9	32.0	3.8	3.2-4.5	30.8	3.2	2.7-3.7	9.6	0.5	0.4-0.7	5.4	0.7	0.5-1.0	4.2	0.4	0.2-0.6
2016-2020	51.0	2.8	2.5-3.2	25.6	3.0	2.5-3.6	25.4	2.6	2.2-3.2	4.2	0.2	0.1-0.3	2.4	0.3	0.1-0.5	1.8	0.1	0.1-0.3	
Non-Hodgkin Lymphomas	1976-1980	190.0	12.4	11.6-13.3	89.6	14.3	13.0-15.8	100.4	11.0	10.1-12.1	94.0	6.3	5.7-6.9	46.2	7.6	6.6-8.7	47.8	5.3	4.6-6.0
	1981-1985	238.0	16.0	15.0-16.9	129.8	20.4	18.8-22.1	108.2	12.1	11.1-13.2	109.0	7.3	6.7-8.0	55.4	9.1	8.0-10.2	53.6	5.9	5.2-6.7
	1986-1990	329.6	21.6	20.5-22.7	219.2	32.2	30.3-34.3	110.4	12.5	11.4-13.6	127.8	8.5	7.9-9.2	73.0	11.7	10.5-13.0	54.8	5.9	5.2-6.7
	1991-1995	447.6	29.0	27.8-30.3	300.0	43.1	40.9-45.5	147.6	16.9	15.7-18.2	126.8	8.5	7.8-9.2	68.8	11.2	10.0-12.5	58.0	6.3	5.6-7.1
	1996-2000	364.0	23.5	22.4-24.6	212.8	31.2	29.3-33.2	151.2	17.2	16.0-18.5	115.2	7.6	7.0-8.3	65.4	10.9	9.7-12.2	49.8	5.5	4.8-6.2
	2001-2005	371.4	23.2	22.2-24.3	204.4	29.7	27.9-31.6	167.0	18.3	17.0-19.6	109.6	6.8	6.3-7.5	62.0	9.7	8.7-10.9	47.6	4.9	4.3-5.6
	2006-2010	392.4	23.3	22.3-24.4	217.8	29.9	28.1-31.8	174.6	18.2	16.9-19.4	100.4	6.0	5.5-6.5	58.0	8.4	7.5-9.5	42.4	4.2	3.7-4.9
	2011-2015	413.0	22.8	21.8-23.8	225.6	28.6	26.9-30.3	187.4	18.3	17.1-19.6	86.4	4.7	4.2-5.1	48.2	6.4	5.6-7.2	38.2	3.4	2.9-4.0
2016-2020	424.8	21.7	20.7-22.6	218.2	25.5	24.0-27.1	206.6	18.8	17.6-20.0	87.0	4.2	3.8-4.6	49.2	5.8	5.1-6.5	37.8	3.0	2.6-3.5	
Myeloma	1976-1980	85.2	5.7	5.1-6.3	41.4	6.9	6.0-8.0	43.8	4.9	4.3-5.6	50.4	3.4	3.0-3.8	22.6	3.9	3.2-4.7	27.8	3.1	2.6-3.7
	1981-1985	83.8	5.5	5.0-6.1	39.4	6.8	5.8-7.8	44.4	4.7	4.1-5.4	50.6	3.4	3.0-3.8	23.8	4.3	3.5-5.1	26.8	2.9	2.4-3.5
	1986-1990	89.0	5.9	5.4-6.5	45.0	7.9	6.9-9.0	44.0	4.8	4.2-5.5	50.2	3.3	2.9-3.8	25.0	4.5	3.7-5.4	25.2	2.6	2.2-3.2
	1991-1995	93.8	6.3	5.7-6.9	42.6	7.2	6.2-8.2	51.2	5.8	5.1-6.6	52.2	3.5	3.1-4.0	25.6	4.5	3.7-5.3	26.6	2.9	2.4-3.4
	1996-2000	100.2	6.7	6.1-7.3	46.2	7.7	6.7-8.8	54.0	6.1	5.4-6.9	49.0	3.3	2.9-3.7	20.2	3.5	2.9-4.3	28.8	3.2	2.7-3.8
	2001-2005	107.6	6.7	6.2-7.3	53.6	8.3	7.3-9.3	54.0	5.8	5.1-6.6	52.2	3.3	2.9-3.7	22.4	3.7	3.1-4.5	29.8	3.1	2.6-3.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**New York (Manhattan)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	132.6	7.9	7.3-8.5	68.4	9.6	8.6-10.7	64.2	6.6	5.9-7.4	52.2	3.1	2.7-3.5	27.2	4.0	3.4-4.8	25.0	2.5	2.1-3.0
	2011-2015	156.4	8.4	7.8-9.0	86.8	10.9	9.9-12.0	69.6	6.5	5.8-7.3	53.4	2.9	2.5-3.2	28.6	3.7	3.1-4.4	24.8	2.2	1.9-2.7
	2016-2020	155.2	7.7	7.1-8.2	86.2	9.9	9.0-10.9	69.0	6.0	5.4-6.7	47.4	2.3	2.0-2.6	25.4	2.9	2.4-3.5	22.0	1.8	1.5-2.2
Leukemias	1976-1980	168.2	11.5	10.7-12.3	88.0	14.6	13.2-16.1	80.2	9.3	8.4-10.3	124.4	8.5	7.9-9.3	63.6	10.9	9.7-12.2	60.8	7.1	6.3-8.0
	1981-1985	171.0	11.7	11.0-12.6	88.6	15.2	13.7-16.7	82.4	9.5	8.6-10.5	116.6	8.1	7.4-8.8	58.6	10.4	9.2-11.7	58.0	6.7	5.9-7.5
	1986-1990	184.4	12.6	11.8-13.5	95.4	16.2	14.7-17.8	89.0	10.3	9.4-11.4	107.8	7.3	6.6-7.9	53.4	9.3	8.2-10.5	54.4	6.0	5.3-6.7
	1991-1995	185.8	12.7	11.9-13.6	95.2	15.6	14.2-17.2	90.6	10.5	9.6-11.6	107.0	7.3	6.7-8.0	55.0	9.4	8.3-10.6	52.0	5.8	5.1-6.6
	1996-2000	187.0	12.6	11.8-13.5	96.0	15.4	14.1-16.9	91.0	10.6	9.6-11.7	97.6	6.6	6.0-7.2	49.4	8.4	7.3-9.5	48.2	5.4	4.7-6.1
	2001-2005	192.4	12.4	11.6-13.3	105.2	16.2	14.8-17.7	87.2	9.7	8.7-10.6	102.8	6.5	6.0-7.1	49.8	8.0	7.0-9.0	53.0	5.6	4.9-6.3
	2006-2010	228.0	14.1	13.3-15.0	128.2	18.3	16.9-19.8	99.8	10.9	9.9-12.0	103.4	6.2	5.7-6.8	55.6	8.1	7.2-9.2	47.8	4.9	4.3-5.6
	2011-2015	268.8	15.2	14.4-16.0	155.2	20.5	19.0-22.0	113.6	11.4	10.4-12.4	105.2	5.9	5.4-6.4	55.4	7.6	6.7-8.6	49.8	4.7	4.1-5.3
	2016-2020	266.0	13.9	13.2-14.7	150.4	18.1	16.8-19.5	115.6	10.7	9.8-11.7	96.6	4.8	4.3-5.2	49.4	6.0	5.3-6.8	47.2	4.0	3.5-4.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Queens**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	8,139.8	386.8	382.9-390.7	3,926.8	458.6	451.8-465.4	4,213.0	349.4	344.6-354.4	4,461.6	211.9	209.1-214.8	2,289.0	272.8	267.5-278.2	2,172.6	175.8	172.4-179.3
	1981-1985	8,738.6	409.7	405.7-413.6	4,174.0	486.6	479.7-493.5	4,564.6	370.1	365.2-375.1	4,352.6	203.4	200.6-206.2	2,203.8	262.8	257.6-267.9	2,148.8	169.3	166.1-172.7
	1986-1990	8,873.8	419.8	415.8-423.7	4,188.4	490.9	484.1-497.8	4,685.4	383.8	378.8-388.9	4,210.8	197.9	195.2-200.6	2,055.8	247.5	242.6-252.5	2,155.0	170.3	167.1-173.6
	1991-1995	9,601.4	453.9	449.8-458.0	4,718.0	546.7	539.6-553.9	4,883.4	401.1	396.0-406.3	4,015.6	189.7	187.1-192.4	1,963.8	237.2	232.4-242.0	2,051.8	163.0	159.8-166.2
	1996-2000	9,731.0	446.2	442.3-450.2	4,723.4	526.2	519.4-533.0	5,007.6	399.0	394.0-404.0	3,707.6	170.1	167.7-172.6	1,806.4	209.3	205.0-213.8	1,901.2	146.9	144.0-149.9
	2001-2005	9,742.6	426.4	422.6-430.2	4,826.0	506.7	500.2-513.2	4,916.6	377.2	372.5-382.0	3,444.4	150.5	148.2-152.7	1,688.2	183.9	180.0-187.9	1,756.2	130.0	127.3-132.8
	2006-2010	10,416.0	444.1	440.3-448.0	5,202.8	520.4	514.0-526.9	5,213.2	395.6	390.7-400.5	3,303.4	141.0	138.8-143.2	1,610.8	169.7	166.0-173.5	1,692.6	123.0	120.4-125.7
	2011-2015	10,804.2	422.8	419.2-426.4	5,263.2	469.4	463.6-475.2	5,541.0	395.5	390.8-400.3	3,342.4	130.2	128.2-132.2	1,653.4	155.7	152.3-159.1	1,689.0	113.9	111.4-116.4
	2016-2020	11,239.6	407.6	404.2-411.1	5,392.8	431.2	426.0-436.5	5,846.8	397.0	392.3-401.7	3,194.4	113.1	111.3-114.9	1,593.0	133.0	130.1-136.1	1,601.4	99.9	97.6-102.1
Oral cavity and pharynx	1976-1980	217.2	10.2	9.6-10.8	141.0	16.0	14.8-17.3	76.2	6.0	5.4-6.7	90.8	4.3	3.9-4.8	63.8	7.5	6.6-8.4	27.0	2.2	1.8-2.6
	1981-1985	234.0	11.0	10.4-11.7	158.2	18.2	16.9-19.5	75.8	6.1	5.5-6.8	82.8	3.8	3.4-4.2	60.2	6.7	5.9-7.5	22.6	1.8	1.5-2.2
	1986-1990	211.0	10.3	9.6-10.9	139.4	16.0	14.8-17.3	71.6	5.9	5.3-6.6	69.4	3.4	3.0-3.7	43.6	5.2	4.5-6.0	25.8	2.1	1.7-2.5
	1991-1995	214.0	10.4	9.8-11.0	135.0	15.4	14.3-16.6	79.0	6.6	5.9-7.3	57.6	2.8	2.5-3.1	40.6	4.6	4.0-5.3	17.0	1.4	1.1-1.7
	1996-2000	191.8	8.9	8.3-9.4	121.8	12.9	11.9-14.0	70.0	5.6	5.0-6.2	43.6	2.0	1.8-2.3	28.8	3.1	2.6-3.7	14.8	1.2	0.9-1.5
	2001-2005	195.6	8.5	8.0-9.1	129.8	13.1	12.1-14.1	65.8	5.1	4.6-5.7	43.2	1.9	1.6-2.1	30.2	3.1	2.6-3.6	13.0	1.0	0.7-1.2
	2006-2010	206.8	8.7	8.2-9.3	130.4	12.4	11.5-13.4	76.4	5.8	5.2-6.4	44.2	1.9	1.6-2.1	28.8	2.9	2.4-3.4	15.4	1.1	0.9-1.4
	2011-2015	234.6	9.1	8.6-9.6	150.4	12.8	11.9-13.8	84.2	6.0	5.5-6.6	49.2	1.9	1.7-2.2	34.0	3.0	2.5-3.5	15.2	1.1	0.8-1.3
	2016-2020	249.0	9.0	8.5-9.5	155.0	12.2	11.3-13.1	94.0	6.3	5.8-7.0	52.4	1.9	1.6-2.1	34.8	2.8	2.4-3.2	17.6	1.1	0.9-1.4
Esophagus	1976-1980	105.8	4.9	4.5-5.3	74.4	8.3	7.4-9.2	31.4	2.5	2.1-2.9	95.6	4.3	3.9-4.7	64.2	7.0	6.2-7.8	31.4	2.4	2.1-2.9
	1981-1985	97.4	4.4	4.0-4.9	65.8	7.2	6.4-8.1	31.6	2.5	2.1-2.9	89.8	4.1	3.7-4.5	64.2	7.0	6.2-7.8	25.6	2.0	1.7-2.4
	1986-1990	97.2	4.6	4.2-5.1	63.2	7.4	6.6-8.3	34.0	2.7	2.3-3.2	84.0	4.0	3.6-4.4	58.2	6.8	6.1-7.7	25.8	2.0	1.7-2.4
	1991-1995	92.4	4.4	4.0-4.8	64.2	7.4	6.6-8.2	28.2	2.3	1.9-2.7	73.0	3.4	3.1-3.8	54.0	6.2	5.5-7.0	19.0	1.5	1.2-1.9
	1996-2000	84.2	3.9	3.5-4.3	58.2	6.5	5.8-7.3	26.0	2.0	1.7-2.4	71.4	3.3	3.0-3.6	49.8	5.7	5.0-6.4	21.6	1.6	1.3-2.0
	2001-2005	98.6	4.3	3.9-4.7	64.2	6.7	6.0-7.5	34.4	2.5	2.2-2.9	71.4	3.1	2.8-3.5	51.8	5.5	4.9-6.3	19.6	1.4	1.1-1.7
	2006-2010	81.6	3.5	3.1-3.8	58.8	5.9	5.3-6.6	22.8	1.7	1.4-2.0	68.8	2.9	2.6-3.3	48.0	5.0	4.3-5.6	20.8	1.5	1.2-1.9
	2011-2015	88.2	3.4	3.1-3.7	65.4	5.8	5.2-6.5	22.8	1.5	1.2-1.8	57.6	2.2	2.0-2.5	41.4	3.7	3.2-4.2	16.2	1.1	0.8-1.3
	2016-2020	92.0	3.2	2.9-3.5	67.6	5.3	4.7-5.9	24.4	1.5	1.2-1.8	64.2	2.2	2.0-2.5	48.6	3.9	3.4-4.5	15.6	0.9	0.7-1.2
Stomach	1976-1980	290.6	14.2	13.4-15.0	163.2	19.7	18.3-21.2	127.4	10.6	9.7-11.4	215.4	10.6	9.9-11.3	121.4	15.0	13.7-16.3	94.0	7.8	7.1-8.5
	1981-1985	289.2	13.7	13.0-14.5	163.0	19.7	18.3-21.1	126.2	9.8	9.0-10.6	202.2	9.5	8.9-10.2	115.2	14.2	13.0-15.5	87.0	6.6	6.0-7.3
	1986-1990	285.8	13.5	12.8-14.2	160.4	19.0	17.6-20.4	125.4	9.8	9.0-10.6	164.6	7.8	7.3-8.4	92.2	11.0	10.0-12.1	72.4	5.7	5.1-6.3
	1991-1995	297.4	14.1	13.4-14.8	170.6	20.2	18.8-21.6	126.8	9.9	9.1-10.7	184.2	8.7	8.2-9.3	107.6	13.0	11.9-14.2	76.6	6.0	5.4-6.6
	1996-2000	299.6	13.7	13.0-14.4	168.4	19.4	18.1-20.7	131.2	10.0	9.2-10.8	159.6	7.3	6.8-7.9	82.2	9.5	8.6-10.5	77.4	5.8	5.3-6.5
	2001-2005	282.0	12.3	11.6-12.9	162.6	17.3	16.2-18.6	119.4	8.9	8.2-9.7	148.8	6.5	6.0-6.9	84.8	9.2	8.4-10.2	64.0	4.7	4.2-5.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Queens**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	288.0	12.3	11.6-12.9	171.0	17.3	16.1-18.5	117.0	8.6	7.9-9.3	132.8	5.7	5.2-6.1	73.6	7.6	6.9-8.5	59.2	4.3	3.8-4.9
	2011-2015	311.2	12.2	11.6-12.8	183.0	16.7	15.6-17.8	128.2	8.8	8.1-9.5	137.4	5.4	5.0-5.8	79.2	7.4	6.7-8.2	58.2	4.0	3.5-4.5
	2016-2020	324.4	11.8	11.2-12.4	185.4	15.2	14.2-16.2	139.0	9.3	8.6-10.0	127.8	4.6	4.2-5.0	77.2	6.4	5.8-7.1	50.6	3.2	2.8-3.7
Colorectal	1976-1980	1,383.4	65.4	63.8-67.0	680.2	80.9	78.1-83.8	703.2	55.7	53.8-57.6	703.6	33.9	32.8-35.1	349.0	43.5	41.3-45.7	354.6	28.2	26.9-29.6
	1981-1985	1,484.4	68.7	67.1-70.3	716.8	83.9	81.1-86.8	767.6	59.1	57.2-61.0	656.6	30.9	29.8-32.0	325.4	39.7	37.7-41.8	331.2	25.6	24.3-26.9
	1986-1990	1,362.6	63.5	62.0-65.1	663.6	79.2	76.4-82.0	699.0	53.6	51.9-55.5	584.8	27.5	26.5-28.5	288.6	35.7	33.8-37.7	296.2	22.7	21.5-23.9
	1991-1995	1,268.8	59.6	58.1-61.1	601.4	72.0	69.4-74.7	667.4	51.5	49.8-53.3	518.0	24.4	23.5-25.4	250.0	30.9	29.1-32.7	268.0	20.4	19.3-21.5
	1996-2000	1,261.6	57.8	56.4-59.3	600.0	68.4	66.0-70.9	661.6	50.6	48.9-52.4	465.2	21.3	20.4-22.2	233.4	27.5	25.9-29.1	231.8	17.4	16.4-18.4
	2001-2005	1,185.4	51.7	50.4-53.1	566.6	60.7	58.4-63.0	618.8	45.6	44.0-47.2	397.4	17.3	16.6-18.1	192.2	21.1	19.8-22.5	205.2	14.7	13.8-15.7
	2006-2010	1,118.8	47.5	46.3-48.8	548.2	55.5	53.4-57.7	570.6	41.6	40.1-43.2	363.6	15.4	14.7-16.1	173.0	18.4	17.2-19.7	190.6	13.4	12.5-14.3
	2011-2015	1,034.8	40.5	39.4-41.6	528.2	48.1	46.2-50.0	506.6	34.8	33.5-36.3	344.6	13.3	12.7-13.9	176.2	16.7	15.6-17.9	168.4	10.9	10.2-11.7
2016-2020	1,000.2	36.3	35.3-37.4	519.4	42.0	40.3-43.6	480.8	31.8	30.5-33.1	314.2	11.1	10.6-11.7	153.0	12.7	11.8-13.6	161.2	9.9	9.2-10.6	
Colon excluding rectum	1976-1980	953.8	45.5	44.1-46.8	450.6	54.3	52.0-56.7	503.2	40.0	38.4-41.7	580.2	28.0	26.9-29.0	280.4	34.9	33.0-36.9	299.8	23.8	22.6-25.1
	1981-1985	1,043.2	48.5	47.2-49.9	487.4	57.8	55.4-60.3	555.8	42.9	41.3-44.5	567.8	26.7	25.7-27.7	280.2	34.2	32.4-36.1	287.6	22.2	21.1-23.4
	1986-1990	991.6	46.3	45.0-47.6	463.6	55.9	53.5-58.3	528.0	40.4	38.9-42.0	516.6	24.3	23.4-25.3	252.2	31.3	29.5-33.2	264.4	20.2	19.2-21.4
	1991-1995	925.2	43.5	42.2-44.7	420.4	51.0	48.8-53.2	504.8	38.8	37.3-40.3	453.6	21.4	20.5-22.3	216.4	26.9	25.3-28.6	237.2	18.0	17.0-19.1
	1996-2000	908.2	41.6	40.4-42.8	418.6	48.1	46.1-50.3	489.6	37.3	35.9-38.8	407.2	18.6	17.8-19.5	202.8	24.0	22.5-25.5	204.4	15.3	14.4-16.3
	2001-2005	854.0	37.3	36.2-38.4	394.8	42.6	40.7-44.5	459.2	33.6	32.2-35.0	348.2	15.2	14.4-15.9	164.8	18.2	17.0-19.5	183.4	13.1	12.3-14.0
	2006-2010	787.8	33.6	32.5-34.6	370.0	38.0	36.3-39.8	417.8	30.2	28.9-31.6	318.2	13.5	12.8-14.2	151.4	16.1	14.9-17.3	166.8	11.7	10.9-12.5
	2011-2015	729.8	28.7	27.7-29.6	358.6	33.2	31.6-34.8	371.2	25.4	24.3-26.6	291.2	11.2	10.6-11.8	146.8	13.9	12.9-15.0	144.4	9.4	8.7-10.1
2016-2020	705.2	25.5	24.7-26.4	352.4	28.6	27.3-30.0	352.8	23.0	21.9-24.2	260.2	9.2	8.7-9.7	122.8	10.3	9.5-11.1	137.4	8.4	7.8-9.1	
Rectum & rectosigmoid	1976-1980	429.6	20.0	19.1-20.8	229.6	26.6	25.0-28.3	200.0	15.7	14.7-16.7	123.4	6.0	5.5-6.5	68.6	8.5	7.6-9.5	54.8	4.4	3.9-5.0
	1981-1985	441.2	20.1	19.3-21.0	229.4	26.1	24.6-27.7	211.8	16.2	15.2-17.2	88.8	4.2	3.8-4.6	45.2	5.5	4.8-6.3	43.6	3.3	2.9-3.8
	1986-1990	371.0	17.2	16.5-18.1	200.0	23.3	21.8-24.8	171.0	13.2	12.3-14.2	68.2	3.2	2.8-3.5	36.4	4.4	3.7-5.1	31.8	2.4	2.0-2.8
	1991-1995	343.6	16.1	15.4-16.9	181.0	21.1	19.7-22.5	162.6	12.8	11.9-13.7	64.4	3.1	2.7-3.4	33.6	4.0	3.4-4.6	30.8	2.4	2.0-2.8
	1996-2000	353.4	16.2	15.5-17.0	181.4	20.3	19.0-21.6	172.0	13.3	12.4-14.2	58.0	2.6	2.4-3.0	30.6	3.5	2.9-4.1	27.4	2.1	1.7-2.5
	2001-2005	331.4	14.5	13.8-15.2	171.8	18.1	16.9-19.4	159.6	12.0	11.1-12.8	49.2	2.2	1.9-2.4	27.4	2.9	2.5-3.5	21.8	1.6	1.3-1.9
	2006-2010	331.0	14.0	13.3-14.6	178.2	17.5	16.3-18.7	152.8	11.4	10.6-12.2	45.4	1.9	1.7-2.2	21.6	2.3	1.9-2.8	23.8	1.7	1.4-2.0
	2011-2015	305.0	11.8	11.2-12.4	169.6	14.9	13.9-15.9	135.4	9.4	8.7-10.2	53.4	2.1	1.8-2.4	29.4	2.8	2.3-3.3	24.0	1.6	1.3-1.9
2016-2020	295.0	10.8	10.2-11.4	167.0	13.3	12.4-14.3	128.0	8.8	8.1-9.5	54.0	1.9	1.7-2.1	30.2	2.4	2.0-2.8	23.8	1.5	1.2-1.8	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Queens**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	81.2	3.9	3.5-4.3	47.0	5.5	4.8-6.3	34.2	2.8	2.4-3.3	93.8	4.4	4.0-4.8	53.0	5.9	5.2-6.8	40.8	3.3	2.8-3.8
	1981-1985	81.4	3.9	3.5-4.3	52.0	6.1	5.4-6.9	29.4	2.4	2.0-2.8	104.2	4.9	4.5-5.4	61.4	7.3	6.5-8.2	42.8	3.4	2.9-3.9
	1986-1990	91.4	4.3	3.9-4.8	59.0	6.8	6.1-7.7	32.4	2.6	2.2-3.0	105.2	4.9	4.5-5.4	62.0	7.3	6.5-8.2	43.2	3.4	2.9-3.9
	1991-1995	133.4	6.4	5.9-6.9	87.0	10.0	9.0-11.0	46.4	3.7	3.2-4.2	130.6	6.2	5.7-6.7	77.4	9.1	8.2-10.1	53.2	4.2	3.7-4.7
	1996-2000	171.8	7.9	7.4-8.5	113.8	12.4	11.3-13.4	58.0	4.5	4.0-5.0	124.6	5.7	5.3-6.2	81.0	9.0	8.1-9.9	43.6	3.3	2.9-3.8
	2001-2005	190.8	8.3	7.8-8.9	134.6	13.5	12.5-14.6	56.2	4.2	3.8-4.8	130.6	5.7	5.3-6.2	84.4	8.7	7.9-9.6	46.2	3.4	3.0-3.9
	2006-2010	225.4	9.5	8.9-10.0	160.8	15.1	14.1-16.2	64.6	4.8	4.3-5.4	145.4	6.2	5.7-6.6	98.6	9.7	8.8-10.6	46.8	3.4	3.0-3.9
	2011-2015	263.2	10.0	9.4-10.5	182.8	15.4	14.4-16.5	80.4	5.5	5.0-6.1	160.8	6.2	5.8-6.6	101.4	8.9	8.1-9.7	59.4	4.1	3.6-4.6
2016-2020	275.4	9.6	9.1-10.2	194.8	15.1	14.1-16.1	80.6	5.2	4.7-5.7	171.0	6.0	5.6-6.4	115.0	9.1	8.4-9.9	56.0	3.5	3.1-4.0	
Pancreas	1976-1980	276.0	12.9	12.2-13.6	134.4	15.4	14.2-16.7	141.6	11.2	10.3-12.0	248.6	11.7	11.0-12.3	122.0	14.1	12.9-15.3	126.6	10.0	9.2-10.8
	1981-1985	280.2	13.0	12.3-13.7	132.0	15.4	14.2-16.7	148.2	11.3	10.5-12.2	252.2	11.7	11.0-12.4	119.4	14.0	12.8-15.2	132.8	10.1	9.3-10.9
	1986-1990	277.6	13.0	12.3-13.7	121.4	14.2	13.1-15.4	156.2	11.9	11.1-12.8	229.2	10.7	10.1-11.3	103.2	12.1	11.1-13.2	126.0	9.6	8.8-10.4
	1991-1995	261.0	12.3	11.6-12.9	112.0	13.6	12.4-14.7	149.0	11.5	10.6-12.3	220.6	10.3	9.7-10.9	95.8	11.5	10.5-12.6	124.8	9.5	8.7-10.3
	1996-2000	273.0	12.5	11.9-13.2	127.2	14.5	13.3-15.6	145.8	11.1	10.3-11.9	216.8	9.9	9.3-10.5	101.4	11.6	10.6-12.7	115.4	8.7	8.0-9.4
	2001-2005	268.0	11.7	11.1-12.3	122.6	13.2	12.2-14.3	145.4	10.6	9.8-11.4	227.4	9.9	9.3-10.5	107.2	11.6	10.7-12.7	120.2	8.7	8.0-9.4
	2006-2010	300.8	12.8	12.2-13.5	147.0	15.1	14.0-16.3	153.8	11.0	10.3-11.9	234.8	10.0	9.4-10.6	115.0	12.1	11.1-13.1	119.8	8.5	7.8-9.2
	2011-2015	333.0	12.9	12.3-13.6	157.4	14.4	13.4-15.4	175.6	11.8	11.0-12.6	256.6	10.0	9.4-10.5	124.8	11.7	10.8-12.6	131.8	8.7	8.1-9.4
2016-2020	369.2	13.0	12.4-13.6	169.6	13.7	12.8-14.7	199.6	12.3	11.5-13.1	267.8	9.3	8.8-9.9	124.6	10.0	9.2-10.9	143.2	8.6	8.0-9.3	
Larynx	1976-1980	116.0	5.2	4.8-5.7	94.6	10.2	9.2-11.2	21.4	1.7	1.4-2.1	45.2	2.0	1.8-2.3	36.8	4.1	3.5-4.8	8.4	0.6	0.4-0.9
	1981-1985	118.6	5.3	4.9-5.8	92.0	9.8	8.9-10.8	26.6	2.2	1.8-2.6	40.0	1.8	1.5-2.0	32.4	3.5	3.0-4.1	7.6	0.6	0.4-0.8
	1986-1990	108.6	5.1	4.7-5.6	86.6	9.7	8.8-10.7	22.0	1.9	1.5-2.3	30.8	1.4	1.2-1.7	24.8	2.8	2.4-3.4	6.0	0.5	0.3-0.7
	1991-1995	103.8	5.0	4.5-5.4	84.6	9.5	8.6-10.5	19.2	1.6	1.3-2.0	40.4	1.9	1.7-2.2	30.2	3.5	2.9-4.1	10.2	0.8	0.6-1.1
	1996-2000	92.0	4.2	3.9-4.7	73.6	8.0	7.2-8.9	18.4	1.5	1.2-1.8	27.8	1.3	1.1-1.5	22.4	2.5	2.1-3.0	5.4	0.4	0.3-0.6
	2001-2005	82.0	3.6	3.2-3.9	66.6	6.9	6.1-7.6	15.4	1.1	0.9-1.4	24.6	1.1	0.9-1.3	19.8	2.1	1.7-2.5	4.8	0.3	0.2-0.5
	2006-2010	80.6	3.4	3.1-3.7	62.8	6.2	5.6-7.0	17.8	1.3	1.1-1.6	24.6	1.0	0.9-1.2	19.8	2.0	1.6-2.4	4.8	0.3	0.2-0.5
	2011-2015	72.2	2.8	2.5-3.1	60.0	5.3	4.7-5.9	12.2	0.8	0.6-1.1	23.6	0.9	0.8-1.1	18.8	1.7	1.4-2.1	4.8	0.3	0.2-0.5
2016-2020	60.6	2.1	1.9-2.4	51.0	4.0	3.5-4.6	9.6	0.6	0.4-0.8	16.6	0.6	0.5-0.7	13.8	1.1	0.9-1.4	2.8	0.2	0.1-0.3	
Lung and bronchus	1976-1980	1,154.6	52.3	50.9-53.7	809.2	90.2	87.3-93.2	345.4	27.3	26.0-28.6	959.6	43.6	42.3-44.9	681.2	76.6	73.9-79.4	278.4	22.0	20.8-23.2
	1981-1985	1,210.4	55.0	53.5-56.4	782.4	88.8	85.9-91.7	428.0	33.2	31.7-34.6	948.6	43.1	41.8-44.4	626.0	71.8	69.2-74.5	322.6	24.9	23.7-26.2
	1986-1990	1,254.0	58.0	56.6-59.5	739.6	85.4	82.6-88.3	514.4	40.8	39.2-42.4	970.2	44.7	43.4-46.0	595.2	69.6	67.1-72.2	375.0	29.3	27.9-30.7
	1991-1995	1,217.4	57.1	55.7-58.6	699.2	81.1	78.4-83.9	518.2	41.3	39.8-43.0	930.8	43.7	42.4-45.0	538.2	63.3	60.9-65.8	392.6	31.1	29.8-32.6
	1996-2000	1,212.0	55.6	54.2-57.0	669.4	75.2	72.7-77.8	542.6	42.3	40.7-43.9	872.6	40.0	38.8-41.2	503.6	57.2	55.0-59.5	369.0	28.5	27.2-29.8
	2001-2005	1,171.0	51.3	50.0-52.7	642.4	68.7	66.3-71.2	528.6	39.5	38.0-41.1	799.2	35.0	33.9-36.1	443.4	47.9	45.9-50.0	355.8	26.4	25.2-27.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Queens**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	1,173.8	50.6	49.3-51.9	613.8	63.5	61.3-65.9	560.0	41.6	40.1-43.2	741.4	31.8	30.8-32.9	406.6	42.7	40.9-44.6	334.8	24.4	23.3-25.6
	2011-2015	1,161.2	45.4	44.2-46.6	605.8	55.9	53.9-57.9	555.4	38.1	36.7-39.6	704.2	27.5	26.6-28.4	391.8	36.7	35.1-38.4	312.4	20.9	19.9-22.0
	2016-2020	1,190.0	41.9	40.9-43.0	614.0	50.0	48.2-51.9	576.0	36.1	34.8-37.5	573.2	20.1	19.4-20.9	317.6	26.5	25.1-27.8	255.6	15.5	14.7-16.4
Melanoma of the skin	1976-1980	119.6	5.8	5.4-6.3	52.0	5.9	5.2-6.7	67.6	5.9	5.2-6.5	42.0	2.0	1.7-2.3	20.6	2.3	1.9-2.9	21.4	1.8	1.5-2.2
	1981-1985	137.8	6.6	6.1-7.1	71.6	8.0	7.2-8.9	66.2	5.6	5.0-6.3	51.6	2.5	2.2-2.8	29.8	3.5	2.9-4.1	21.8	1.8	1.4-2.1
	1986-1990	139.6	6.7	6.2-7.3	70.6	8.2	7.3-9.1	69.0	5.8	5.2-6.4	46.8	2.3	2.0-2.6	24.6	2.9	2.4-3.5	22.2	1.9	1.5-2.3
	1991-1995	130.0	6.1	5.6-6.6	68.6	7.8	7.0-8.7	61.4	5.0	4.4-5.6	33.6	1.6	1.3-1.8	18.4	2.1	1.7-2.6	15.2	1.2	1.0-1.5
	1996-2000	150.8	6.9	6.4-7.4	75.2	8.4	7.6-9.3	75.6	6.0	5.4-6.7	39.6	1.8	1.6-2.1	21.8	2.4	2.0-3.0	17.8	1.4	1.1-1.7
	2001-2005	176.4	7.7	7.2-8.2	93.0	9.8	9.0-10.8	83.4	6.4	5.8-7.1	32.4	1.4	1.2-1.7	17.2	1.9	1.5-2.4	15.2	1.2	0.9-1.4
	2006-2010	215.8	9.2	8.7-9.8	108.6	10.9	10.0-11.9	107.2	8.1	7.4-8.8	29.2	1.3	1.1-1.5	15.0	1.6	1.2-2.0	14.2	1.1	0.8-1.4
	2011-2015	218.2	8.6	8.1-9.1	113.6	10.6	9.7-11.5	104.6	7.4	6.8-8.1	32.4	1.3	1.1-1.5	16.8	1.6	1.2-1.9	15.6	1.0	0.8-1.3
	2016-2020	198.4	7.2	6.8-7.7	111.0	9.3	8.5-10.1	87.4	5.9	5.3-6.5	25.0	0.9	0.8-1.1	15.0	1.3	1.0-1.6	10.0	0.7	0.5-0.9
Female breast	1976-1980							1,165.0	100.3	97.7-103.1							447.0	37.2	35.6-38.8
	1981-1985							1,236.0	104.8	102.1-107.6							431.0	35.4	33.9-37.0
	1986-1990							1,314.6	112.4	109.6-115.3							434.2	35.9	34.4-37.5
	1991-1995							1,407.0	120.0	117.1-122.8							397.0	32.9	31.4-34.4
	1996-2000							1,489.6	121.9	119.2-124.8							352.8	27.9	26.6-29.3
	2001-2005							1,372.6	107.6	105.1-110.2							320.0	24.1	22.9-25.3
	2006-2010							1,484.8	114.6	112.0-117.3							272.4	20.1	19.0-21.2
	2011-2015							1,570.2	114.2	111.6-116.8							262.4	17.9	17.0-18.9
	2016-2020							1,808.4	125.6	122.9-128.3							254.2	16.3	15.4-17.3
Cervix uteri	1976-1980							137.4	12.7	11.8-13.8							47.2	4.0	3.5-4.6
	1981-1985							125.0	11.7	10.8-12.7							44.4	3.9	3.4-4.4
	1986-1990							136.0	12.7	11.7-13.7							35.8	3.1	2.6-3.6
	1991-1995							155.8	14.0	13.0-15.1							37.4	3.4	2.9-3.9
	1996-2000							142.2	12.0	11.1-12.9							34.6	2.8	2.4-3.3
	2001-2005							122.0	10.0	9.2-10.8							32.6	2.6	2.2-3.0
	2006-2010							128.6	10.2	9.5-11.1							32.4	2.4	2.1-2.8
	2011-2015							127.2	9.9	9.2-10.7							39.4	2.8	2.4-3.3
	2016-2020							124.6	9.3	8.6-10.1							31.0	2.1	1.8-2.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Queens**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							260.2	21.0	19.8-22.2							69.4	5.4	4.8-6.0
	1981-1985							268.4	21.2	20.1-22.5							61.0	4.7	4.2-5.3
	1986-1990							243.4	19.7	18.6-20.9							61.2	4.7	4.2-5.2
	1991-1995							288.8	23.9	22.7-25.2							59.8	4.7	4.2-5.2
	1996-2000							315.0	25.5	24.2-26.8							61.8	4.8	4.3-5.3
	2001-2005							344.0	26.6	25.4-27.9							66.8	5.0	4.5-5.6
	2006-2010							370.6	28.0	26.7-29.3							65.8	4.8	4.3-5.3
	2011-2015							418.4	29.1	27.8-30.4							81.0	5.5	5.0-6.1
	2016-2020							451.0	29.5	28.2-30.7							99.4	6.2	5.6-6.8
Ovary	1976-1980							187.6	15.8	14.7-16.9							133.6	10.6	9.8-11.5
	1981-1985							207.2	17.3	16.2-18.4							118.0	9.4	8.6-10.2
	1986-1990							202.0	17.2	16.1-18.3							122.4	9.7	9.0-10.6
	1991-1995							193.4	16.4	15.4-17.5							114.2	9.3	8.5-10.1
	1996-2000							184.4	15.0	14.1-16.0							100.2	7.9	7.2-8.6
	2001-2005							187.4	14.6	13.7-15.6							103.4	7.8	7.1-8.5
	2006-2010							166.2	12.7	11.9-13.6							96.8	7.1	6.5-7.8
	2011-2015							173.0	12.5	11.7-13.4							95.6	6.6	6.0-7.2
	2016-2020							161.2	11.4	10.6-12.3							83.6	5.3	4.8-5.9
Prostate	1976-1980				583.0	73.9	71.0-76.8							200.4	28.1	26.3-30.0			
	1981-1985				670.6	82.4	79.5-85.4							219.6	29.9	28.0-31.8			
	1986-1990				774.2	94.1	91.1-97.3							220.8	29.2	27.4-31.0			
	1991-1995				1,300.4	151.3	147.5-155.1							247.4	32.8	30.9-34.7			
	1996-2000				1,366.2	152.7	149.0-156.4							213.6	27.2	25.6-28.9			
	2001-2005				1,433.0	150.2	146.7-153.8							186.0	21.9	20.6-23.4			
	2006-2010				1,620.6	159.5	156.0-163.1							172.4	19.7	18.4-21.1			
	2011-2015				1,492.4	127.4	124.4-130.4							165.4	16.7	15.6-17.9			
	2016-2020				1,499.6	113.1	110.5-115.7							176.8	15.5	14.5-16.6			
Testis	1976-1980				25.0	2.8	2.3-3.4							6.0	0.6	0.4-0.9			
	1981-1985				30.8	3.1	2.6-3.6							4.0	0.4	0.3-0.7			
	1986-1990				32.8	3.1	2.6-3.7							3.6	0.4	0.2-0.6			
	1991-1995				37.8	3.5	3.0-4.0							3.2	0.3	0.2-0.5			
	1996-2000				34.6	3.0	2.6-3.5							1.6	0.1	0.1-0.3			
	2001-2005				45.4	3.9	3.4-4.4							3.4	0.3	0.2-0.5			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Queens**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				45.2	3.9	3.4-4.5							2.6	0.2	0.1-0.4			
	2011-2015				46.6	3.9	3.4-4.5							2.0	0.2	0.1-0.3			
	2016-2020				54.4	4.8	4.2-5.4							2.0	0.2	0.1-0.3			
Urinary bladder (incl. in situ)	1976-1980	415.8	19.4	18.6-20.3	303.4	36.1	34.2-38.0	112.4	8.8	8.1-9.6	124.2	6.2	5.7-6.7	84.2	11.1	10.0-12.2	40.0	3.3	2.8-3.8
	1981-1985	444.4	20.2	19.4-21.1	318.8	37.2	35.3-39.2	125.6	9.6	8.9-10.4	103.4	4.9	4.5-5.3	70.2	9.0	8.1-10.1	33.2	2.5	2.2-3.0
	1986-1990	406.8	18.8	17.9-19.6	283.4	33.5	31.7-35.3	123.4	9.4	8.7-10.2	97.6	4.6	4.2-5.0	61.2	7.9	7.0-8.9	36.4	2.7	2.3-3.2
	1991-1995	441.2	20.5	19.7-21.4	303.6	36.4	34.5-38.3	137.6	10.6	9.8-11.5	92.8	4.4	4.0-4.8	62.4	8.2	7.3-9.2	30.4	2.3	1.9-2.7
	1996-2000	423.0	19.4	18.5-20.2	295.2	34.5	32.7-36.3	127.8	9.6	8.9-10.4	89.8	4.1	3.7-4.5	57.8	7.2	6.4-8.1	32.0	2.3	2.0-2.7
	2001-2005	420.6	18.4	17.6-19.2	303.2	33.5	31.8-35.2	117.4	8.6	7.9-9.3	82.2	3.6	3.2-3.9	56.6	6.5	5.8-7.3	25.6	1.8	1.5-2.1
	2006-2010	442.4	18.9	18.2-19.8	326.8	34.8	33.1-36.6	115.6	8.4	7.7-9.1	93.2	3.9	3.6-4.3	61.8	6.8	6.0-7.6	31.4	2.2	1.9-2.6
	2011-2015	438.0	17.3	16.6-18.1	317.0	30.3	28.8-31.8	121.0	8.2	7.6-8.9	90.0	3.5	3.2-3.8	58.6	5.9	5.2-6.6	31.4	2.0	1.7-2.3
2016-2020	431.6	15.3	14.6-15.9	322.2	26.9	25.6-28.3	109.4	6.7	6.2-7.4	78.2	2.7	2.5-3.0	55.6	4.9	4.3-5.5	22.6	1.4	1.1-1.7	
Kidney and renal pelvis	1976-1980	158.8	7.4	6.9-7.9	100.2	11.2	10.2-12.2	58.6	4.8	4.2-5.4	80.0	3.8	3.4-4.2	52.0	6.2	5.4-7.1	28.0	2.2	1.9-2.6
	1981-1985	171.6	8.0	7.5-8.6	106.6	12.2	11.1-13.3	65.0	5.2	4.6-5.8	73.0	3.4	3.0-3.8	45.0	5.3	4.6-6.0	28.0	2.2	1.8-2.6
	1986-1990	194.2	9.1	8.5-9.7	114.2	13.2	12.1-14.3	80.0	6.5	5.9-7.2	71.2	3.4	3.0-3.8	42.8	5.2	4.5-6.0	28.4	2.3	1.9-2.7
	1991-1995	212.8	10.1	9.5-10.7	125.2	14.3	13.2-15.5	87.6	7.2	6.5-7.9	66.0	3.1	2.8-3.4	39.6	4.6	4.0-5.3	26.4	2.0	1.7-2.4
	1996-2000	251.8	11.6	10.9-12.2	154.6	17.0	15.8-18.3	97.2	7.7	7.0-8.4	65.0	3.0	2.7-3.3	41.8	4.8	4.2-5.6	23.2	1.8	1.5-2.1
	2001-2005	282.0	12.4	11.8-13.1	179.0	18.3	17.1-19.5	103.0	7.9	7.3-8.7	61.0	2.7	2.4-3.0	38.4	4.1	3.6-4.8	22.6	1.7	1.4-2.0
	2006-2010	333.4	14.2	13.5-14.9	212.4	20.7	19.5-22.0	121.0	9.2	8.5-10.0	65.0	2.7	2.5-3.1	40.0	4.1	3.5-4.7	25.0	1.8	1.5-2.2
	2011-2015	338.0	13.1	12.5-13.8	221.0	19.1	18.0-20.3	117.0	8.3	7.6-9.0	58.6	2.3	2.0-2.5	37.4	3.5	3.0-4.0	21.2	1.4	1.1-1.7
2016-2020	393.8	14.4	13.7-15.1	251.4	20.2	19.0-21.3	142.4	9.6	8.8-10.3	59.2	2.1	1.9-2.4	38.6	3.2	2.8-3.7	20.6	1.3	1.1-1.6	
Brain and other central nervous system	1976-1980	120.4	5.7	5.2-6.2	61.8	6.7	6.0-7.5	58.6	5.0	4.4-5.6	86.8	4.0	3.6-4.4	47.2	5.1	4.4-5.8	39.6	3.3	2.8-3.8
	1981-1985	123.6	6.0	5.6-6.5	66.0	7.4	6.6-8.3	57.6	4.9	4.3-5.5	85.0	3.9	3.5-4.3	42.6	4.7	4.1-5.4	42.4	3.3	2.9-3.8
	1986-1990	125.4	6.2	5.7-6.7	65.6	7.4	6.6-8.3	59.8	5.3	4.7-5.9	73.6	3.5	3.1-3.9	39.8	4.5	3.9-5.2	33.8	2.7	2.3-3.2
	1991-1995	140.0	6.8	6.3-7.3	67.6	7.5	6.7-8.3	72.4	6.2	5.6-6.9	64.6	3.1	2.7-3.4	33.6	3.8	3.2-4.4	31.0	2.5	2.1-2.9
	1996-2000	141.6	6.6	6.1-7.1	73.2	7.6	6.9-8.5	68.4	5.7	5.1-6.3	73.6	3.4	3.1-3.8	35.6	3.8	3.3-4.4	38.0	3.1	2.7-3.6
	2001-2005	133.2	6.0	5.5-6.4	72.0	7.3	6.5-8.0	61.2	5.0	4.4-5.6	65.2	2.9	2.6-3.2	35.8	3.6	3.1-4.2	29.4	2.3	1.9-2.7
	2006-2010	135.2	6.0	5.5-6.4	75.0	7.4	6.6-8.2	60.2	4.9	4.3-5.5	69.8	3.0	2.7-3.4	39.2	3.9	3.4-4.5	30.6	2.4	2.0-2.8
	2011-2015	131.4	5.5	5.0-5.9	70.6	6.4	5.7-7.1	60.8	4.7	4.2-5.3	72.4	2.9	2.6-3.2	36.4	3.3	2.8-3.8	36.0	2.6	2.2-3.0
2016-2020	132.0	5.2	4.8-5.6	72.6	6.1	5.5-6.8	59.4	4.4	3.9-5.0	77.0	2.9	2.6-3.2	41.4	3.5	3.0-4.0	35.6	2.4	2.0-2.8	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Queens**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	66.2	3.4	3.0-3.8	18.8	2.1	1.7-2.6	47.4	4.5	3.9-5.1	14.0	0.7	0.5-0.8	5.4	0.6	0.4-0.9	8.6	0.7	0.5-0.9
	1981-1985	82.6	4.1	3.7-4.5	22.6	2.5	2.0-3.0	60.0	5.4	4.8-6.1	14.2	0.7	0.5-0.8	5.4	0.6	0.4-0.9	8.8	0.7	0.5-0.9
	1986-1990	80.6	4.0	3.6-4.4	20.8	2.4	1.9-2.9	59.8	5.4	4.8-6.1	12.0	0.5	0.4-0.7	3.2	0.4	0.2-0.6	8.8	0.7	0.5-0.9
	1991-1995	114.2	5.5	5.0-6.0	28.0	3.0	2.5-3.5	86.2	7.7	7.0-8.5	13.8	0.6	0.5-0.8	4.6	0.5	0.3-0.8	9.2	0.7	0.5-0.9
	1996-2000	154.0	7.0	6.5-7.5	34.6	3.5	3.0-4.0	119.4	10.1	9.3-10.9	13.4	0.6	0.5-0.8	4.8	0.6	0.4-0.8	8.6	0.7	0.5-0.9
	2001-2005	220.8	9.6	9.1-10.2	51.6	5.0	4.4-5.6	169.2	13.9	13.0-14.9	10.2	0.4	0.3-0.6	4.4	0.5	0.3-0.7	5.8	0.4	0.3-0.6
	2006-2010	342.4	14.6	13.9-15.3	74.4	6.7	6.1-7.5	268.0	21.8	20.7-23.1	15.0	0.7	0.5-0.8	4.6	0.5	0.3-0.8	10.4	0.8	0.6-1.0
	2011-2015	486.6	19.8	19.0-20.6	108.2	9.2	8.4-10.0	378.4	29.7	28.4-31.1	12.6	0.5	0.4-0.6	4.4	0.4	0.3-0.6	8.2	0.6	0.4-0.8
	2016-2020	492.4	20.1	19.3-20.9	115.6	9.7	8.9-10.6	376.8	30.0	28.6-31.4	15.2	0.5	0.4-0.7	6.6	0.5	0.4-0.8	8.6	0.5	0.4-0.7
Hodgkin lymphoma	1976-1980	66.2	3.2	2.9-3.6	38.2	4.1	3.5-4.7	28.0	2.6	2.1-3.0	24.0	1.2	1.0-1.4	13.2	1.5	1.1-1.9	10.8	0.9	0.7-1.2
	1981-1985	67.8	3.3	2.9-3.6	39.2	4.2	3.6-4.9	28.6	2.5	2.1-3.0	19.8	0.9	0.8-1.1	12.2	1.3	1.0-1.7	7.6	0.6	0.4-0.9
	1986-1990	60.6	2.9	2.6-3.2	32.6	3.3	2.8-3.9	28.0	2.5	2.1-2.9	16.8	0.8	0.6-1.0	9.2	0.9	0.7-1.3	7.6	0.6	0.4-0.9
	1991-1995	64.6	3.0	2.7-3.4	36.8	3.8	3.2-4.4	27.8	2.4	2.0-2.8	15.2	0.7	0.5-0.9	9.0	0.9	0.7-1.3	6.2	0.5	0.3-0.7
	1996-2000	61.4	2.7	2.4-3.1	33.8	3.1	2.7-3.7	27.6	2.3	2.0-2.8	9.2	0.4	0.3-0.5	4.2	0.4	0.3-0.7	5.0	0.4	0.2-0.6
	2001-2005	60.6	2.7	2.4-3.0	31.8	3.0	2.5-3.5	28.8	2.4	2.0-2.8	9.2	0.4	0.3-0.5	4.8	0.5	0.3-0.7	4.4	0.3	0.2-0.5
	2006-2010	70.2	3.1	2.8-3.4	36.8	3.4	2.9-3.9	33.4	2.9	2.4-3.3	7.8	0.3	0.2-0.5	4.4	0.4	0.3-0.7	3.4	0.3	0.2-0.4
	2011-2015	62.4	2.7	2.4-3.0	36.0	3.2	2.7-3.7	26.4	2.2	1.8-2.6	7.6	0.3	0.2-0.4	4.4	0.4	0.2-0.6	3.2	0.2	0.1-0.4
	2016-2020	60.8	2.6	2.3-2.9	34.6	3.0	2.6-3.5	26.2	2.2	1.8-2.6	4.8	0.2	0.1-0.3	2.6	0.2	0.1-0.3	2.2	0.2	0.1-0.3
Non-Hodgkin lymphomas	1976-1980	240.4	11.4	10.8-12.1	119.6	13.3	12.2-14.4	120.8	10.1	9.3-11.0	126.0	6.0	5.5-6.5	65.6	7.6	6.8-8.5	60.4	4.9	4.4-5.5
	1981-1985	309.4	14.7	14.0-15.5	147.8	17.0	15.7-18.3	161.6	13.0	12.1-13.9	140.6	6.7	6.2-7.2	67.2	7.8	7.0-8.7	73.4	5.8	5.2-6.4
	1986-1990	342.0	16.3	15.5-17.1	177.6	20.1	18.8-21.6	164.4	13.2	12.3-14.1	148.2	7.0	6.5-7.5	75.0	8.6	7.8-9.6	73.2	5.7	5.1-6.3
	1991-1995	401.8	19.1	18.3-20.0	220.0	24.6	23.1-26.1	181.8	14.7	13.8-15.7	149.0	7.1	6.6-7.6	77.6	9.1	8.2-10.1	71.4	5.6	5.0-6.2
	1996-2000	390.0	17.9	17.1-18.7	207.4	22.3	21.0-23.7	182.6	14.4	13.5-15.4	150.0	6.9	6.4-7.4	79.4	9.0	8.1-9.9	70.6	5.5	4.9-6.1
	2001-2005	400.4	17.5	16.8-18.3	200.8	20.4	19.2-21.7	199.6	15.2	14.3-16.2	134.6	5.9	5.4-6.3	71.2	7.6	6.8-8.4	63.4	4.6	4.1-5.2
	2006-2010	426.4	18.2	17.5-19.0	220.0	21.9	20.6-23.3	206.4	15.5	14.6-16.5	112.4	4.8	4.4-5.2	59.4	6.2	5.5-7.0	53.0	3.8	3.3-4.3
	2011-2015	480.8	19.1	18.3-19.9	258.0	23.5	22.2-24.9	222.8	15.6	14.7-16.6	120.2	4.7	4.3-5.1	66.2	6.2	5.5-6.9	54.0	3.6	3.2-4.1
	2016-2020	492.4	18.1	17.3-18.8	264.0	21.7	20.5-22.9	228.4	15.2	14.3-16.2	113.2	4.0	3.7-4.4	66.6	5.7	5.1-6.3	46.6	2.8	2.4-3.2
Myeloma	1976-1980	100.4	4.7	4.3-5.1	49.2	5.7	5.0-6.6	51.2	4.0	3.5-4.5	64.0	3.0	2.7-3.3	30.2	3.4	2.9-4.1	33.8	2.6	2.3-3.1
	1981-1985	116.2	5.4	4.9-5.8	59.4	6.9	6.1-7.8	56.8	4.4	3.9-5.0	65.8	3.0	2.7-3.3	33.4	3.9	3.3-4.6	32.4	2.4	2.1-2.8
	1986-1990	110.4	5.2	4.7-5.6	55.2	6.5	5.8-7.4	55.2	4.3	3.8-4.9	69.0	3.2	2.9-3.6	36.6	4.3	3.7-5.0	32.4	2.5	2.1-2.9
	1991-1995	113.0	5.3	4.9-5.8	54.0	6.4	5.6-7.2	59.0	4.6	4.1-5.2	69.0	3.2	2.9-3.6	34.4	4.1	3.5-4.8	34.6	2.7	2.3-3.1
	1996-2000	121.6	5.6	5.2-6.0	57.0	6.4	5.6-7.1	64.6	5.0	4.5-5.6	59.2	2.7	2.4-3.0	25.6	2.9	2.4-3.5	33.6	2.6	2.2-3.0
	2001-2005	153.6	6.7	6.3-7.2	75.4	8.0	7.2-8.8	78.2	5.9	5.3-6.5	61.0	2.7	2.4-3.0	29.4	3.2	2.7-3.8	31.6	2.3	2.0-2.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Queens**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	167.6	7.1	6.7-7.6	92.0	9.4	8.5-10.3	75.6	5.5	5.0-6.1	63.8	2.7	2.4-3.1	33.6	3.6	3.1-4.2	30.2	2.2	1.8-2.6
	2011-2015	215.2	8.4	7.9-9.0	110.0	10.0	9.2-10.9	105.2	7.4	6.7-8.0	69.2	2.7	2.4-3.0	36.6	3.4	2.9-3.9	32.6	2.2	1.9-2.6
	2016-2020	216.0	7.6	7.2-8.1	111.6	8.9	8.1-9.7	104.4	6.6	6.1-7.3	68.4	2.4	2.1-2.6	33.6	2.8	2.4-3.3	34.8	2.1	1.8-2.4
Leukemias	1976-1980	233.8	11.6	10.9-12.3	132.4	15.7	14.4-17.0	101.4	8.8	8.0-9.6	152.2	7.6	7.1-8.2	87.0	10.6	9.6-11.7	65.2	5.6	5.0-6.3
	1981-1985	236.6	11.5	10.8-12.2	128.6	15.2	14.0-16.4	108.0	9.0	8.3-9.9	144.6	6.9	6.4-7.5	78.2	9.2	8.3-10.2	66.4	5.4	4.8-6.0
	1986-1990	233.2	11.3	10.6-12.0	127.8	15.1	13.9-16.3	105.4	8.7	8.0-9.5	138.0	6.6	6.1-7.1	77.6	9.2	8.3-10.2	60.4	4.8	4.3-5.4
	1991-1995	237.0	11.4	10.8-12.1	119.4	13.9	12.8-15.1	117.6	9.8	9.0-10.6	136.8	6.5	6.0-7.0	72.6	8.7	7.8-9.7	64.2	5.2	4.6-5.8
	1996-2000	254.0	11.7	11.1-12.4	136.2	15.0	13.9-16.2	117.8	9.4	8.6-10.2	133.0	6.1	5.6-6.6	69.2	7.8	7.0-8.7	63.8	5.0	4.4-5.6
	2001-2005	263.6	11.7	11.1-12.3	143.2	15.1	14.0-16.2	120.4	9.2	8.5-10.0	130.2	5.7	5.3-6.2	72.4	7.8	7.0-8.6	57.8	4.3	3.8-4.8
	2006-2010	284.0	12.5	11.8-13.1	155.8	16.0	14.9-17.2	128.2	10.0	9.2-10.8	132.0	5.7	5.3-6.2	69.8	7.4	6.6-8.2	62.2	4.6	4.1-5.2
	2011-2015	339.8	13.7	13.0-14.4	188.4	17.5	16.4-18.7	151.4	10.9	10.1-11.7	135.8	5.4	5.0-5.9	79.6	7.7	7.0-8.6	56.2	3.8	3.4-4.3
	2016-2020	332.0	12.7	12.1-13.3	192.4	16.4	15.4-17.5	139.6	9.7	9.0-10.5	128.0	4.6	4.3-5.0	73.0	6.3	5.6-7.0	55.0	3.5	3.1-4.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020  
Richmond (Staten Island)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	1,175.6	421.0	410.0-432.1	592.0	520.6	500.8-541.0	583.6	366.8	353.4-380.6	629.8	230.0	221.9-238.4	336.4	306.1	290.6-322.0	293.4	184.9	175.5-194.8
	1981-1985	1,396.4	451.9	441.2-462.9	699.2	558.4	538.8-578.4	697.2	395.2	382.0-408.7	700.4	230.6	222.9-238.5	374.4	311.6	296.7-326.9	326.0	183.7	174.8-192.9
	1986-1990	1,656.0	497.4	486.6-508.3	775.0	571.5	552.7-590.9	881.0	461.5	447.9-475.5	772.0	237.3	229.7-245.0	401.0	308.8	294.7-323.5	371.0	194.4	185.6-203.5
	1991-1995	1,855.2	507.4	497.1-517.9	912.8	608.1	589.8-626.8	942.4	452.6	439.7-465.8	828.4	232.4	225.3-239.7	413.2	293.7	280.5-307.2	415.2	198.3	189.8-207.0
	1996-2000	2,140.4	519.4	509.6-529.4	1,059.4	611.4	594.6-628.6	1,081.0	463.9	451.6-476.4	841.6	208.6	202.4-215.1	418.0	259.5	248.1-271.2	423.6	179.1	171.5-186.9
	2001-2005	2,326.6	508.7	499.4-518.1	1,161.8	591.3	575.8-607.1	1,164.8	452.7	441.1-464.5	835.0	186.0	180.4-191.8	405.8	221.3	211.5-231.4	429.2	164.1	157.2-171.2
	2006-2010	2,613.8	527.1	517.9-536.3	1,300.0	599.3	584.4-614.5	1,313.8	479.7	468.0-491.6	828.0	171.0	165.8-176.4	411.0	205.4	196.4-214.8	417.0	149.2	142.8-155.9
	2011-2015	2,813.4	518.0	509.3-526.9	1,381.8	566.9	553.2-580.9	1,431.6	487.2	475.7-498.9	817.2	151.3	146.6-156.1	416.2	183.0	174.9-191.2	401.0	130.0	124.2-135.9
	2016-2020	2,967.8	504.2	495.9-512.6	1,437.8	533.6	521.0-546.5	1,530.0	486.8	475.5-498.2	844.2	141.6	137.2-146.0	411.2	161.8	154.7-169.2	433.0	127.5	122.0-133.1
Oral cavity and pharynx	1976-1980	38.2	13.6	11.7-15.7	27.2	23.1	19.1-27.6	11.0	7.0	5.3-9.1	17.6	6.2	5.0-7.7	11.4	9.9	7.4-13.1	6.2	3.8	2.6-5.4
	1981-1985	40.0	12.3	10.6-14.2	25.6	18.2	15.0-21.8	14.4	7.8	6.1-9.9	12.8	4.0	3.1-5.1	9.0	6.7	4.8-9.2	3.8	2.1	1.3-3.3
	1986-1990	49.0	14.4	12.6-16.3	34.2	23.8	20.2-27.8	14.8	7.5	5.9-9.4	13.8	4.2	3.2-5.3	9.8	7.3	5.3-9.8	4.0	2.0	1.2-3.2
	1991-1995	38.8	10.6	9.2-12.2	27.2	17.4	14.5-20.7	11.6	5.6	4.3-7.3	10.2	2.8	2.1-3.7	7.4	4.9	3.4-6.9	2.8	1.3	0.7-2.2
	1996-2000	38.8	9.3	8.1-10.8	26.0	14.3	11.9-17.1	12.8	5.5	4.2-7.0	7.6	1.9	1.3-2.5	4.2	2.5	1.5-3.8	3.4	1.4	0.8-2.3
	2001-2005	46.6	10.0	8.7-11.4	28.6	13.9	11.6-16.4	18.0	7.0	5.6-8.6	9.6	2.1	1.6-2.8	5.4	2.9	1.9-4.3	4.2	1.6	1.0-2.4
	2006-2010	41.8	8.2	7.1-9.4	28.0	12.0	10.0-14.2	13.8	4.9	3.8-6.3	10.2	2.0	1.5-2.7	6.4	2.7	1.8-3.9	3.8	1.4	0.8-2.2
	2011-2015	56.8	10.1	8.9-11.4	38.6	14.9	12.8-17.3	18.2	6.1	4.9-7.5	9.2	1.6	1.2-2.2	6.0	2.4	1.6-3.4	3.2	1.0	0.6-1.7
	2016-2020	60.4	10.0	8.9-11.2	42.0	15.1	13.0-17.4	18.4	5.6	4.5-6.9	11.6	1.8	1.4-2.4	8.2	2.9	2.0-3.9	3.4	0.9	0.5-1.5
Esophagus	1976-1980	16.0	5.8	4.6-7.2	10.8	9.7	7.1-12.8	5.2	3.3	2.1-4.8	12.8	4.7	3.6-6.0	9.0	8.4	5.9-11.3	3.8	2.4	1.4-3.7
	1981-1985	13.2	4.4	3.4-5.6	8.8	7.4	5.2-10.1	4.4	2.5	1.6-3.8	10.8	3.6	2.7-4.7	8.2	7.3	5.1-10.0	2.6	1.5	0.8-2.5
	1986-1990	15.6	4.7	3.7-5.9	11.4	8.4	6.2-11.0	4.2	2.2	1.4-3.4	14.8	4.5	3.5-5.6	10.6	8.0	5.9-10.6	4.2	2.2	1.4-3.4
	1991-1995	15.2	4.2	3.3-5.2	11.2	7.1	5.3-9.3	4.0	1.9	1.2-2.9	14.8	4.1	3.2-5.1	11.4	7.4	5.6-9.7	3.4	1.6	0.9-2.5
	1996-2000	14.4	3.6	2.8-4.5	10.8	6.4	4.8-8.4	3.6	1.5	0.9-2.4	13.4	3.3	2.6-4.2	10.6	6.3	4.7-8.3	2.8	1.2	0.6-2.0
	2001-2005	19.0	4.2	3.4-5.1	11.8	6.1	4.6-7.9	7.2	2.8	2.0-3.9	18.2	4.0	3.2-4.9	14.2	7.2	5.6-9.2	4.0	1.6	0.9-2.4
	2006-2010	22.6	4.4	3.6-5.3	16.8	7.3	5.8-9.1	5.8	2.1	1.4-3.0	19.2	3.8	3.0-4.6	13.6	6.1	4.7-7.7	5.6	2.0	1.3-2.9
	2011-2015	18.0	3.3	2.6-4.1	14.4	6.0	4.6-7.6	3.6	1.2	0.7-1.9	16.8	3.1	2.5-3.9	13.2	5.4	4.2-7.0	3.6	1.3	0.8-2.1
	2016-2020	28.4	4.5	3.8-5.3	22.2	7.7	6.3-9.4	6.2	1.9	1.3-2.7	17.6	2.8	2.2-3.5	14.0	4.9	3.8-6.2	3.6	1.1	0.6-1.7
Stomach	1976-1980	35.4	13.7	11.7-16.0	19.4	19.5	15.5-24.0	16.0	10.4	8.2-12.9	28.4	11.4	9.6-13.5	15.4	16.8	13.1-21.2	13.0	8.6	6.6-11.0
	1981-1985	35.2	11.8	10.1-13.7	19.0	16.7	13.3-20.6	16.2	9.0	7.1-11.2	24.6	8.5	7.0-10.1	14.0	13.0	10.0-16.6	10.6	5.9	4.4-7.7
	1986-1990	39.6	12.4	10.7-14.2	23.0	17.1	14.0-20.7	16.6	8.6	6.8-10.7	26.2	8.4	7.0-10.0	13.8	10.6	8.1-13.5	12.4	6.5	5.0-8.3
	1991-1995	40.6	11.5	10.0-13.2	23.6	16.6	13.6-20.1	17.0	8.1	6.4-10.0	25.8	7.4	6.2-8.8	14.4	10.6	8.2-13.5	11.4	5.4	4.1-7.0
	1996-2000	46.0	11.4	10.0-13.0	24.8	14.7	12.1-17.6	21.2	8.8	7.2-10.7	21.8	5.5	4.5-6.6	13.0	8.1	6.2-10.3	8.8	3.7	2.7-4.9
	2001-2005	47.0	10.5	9.2-11.9	27.6	14.5	12.1-17.2	19.4	7.4	6.0-9.1	23.0	5.1	4.2-6.1	11.2	5.8	4.3-7.6	11.8	4.4	3.3-5.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020  
Richmond (Staten Island)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	43.2	8.9	7.7-10.2	26.2	12.8	10.6-15.2	17.0	6.0	4.8-7.5	20.0	4.2	3.4-5.1	10.6	5.7	4.2-7.5	9.4	3.3	2.4-4.4
	2011-2015	44.8	8.4	7.3-9.6	25.0	10.5	8.7-12.6	19.8	6.6	5.4-8.1	20.2	3.5	2.9-4.3	12.4	5.1	3.9-6.7	7.8	2.3	1.6-3.2
	2016-2020	47.0	8.1	7.1-9.3	29.0	11.5	9.6-13.6	18.0	5.5	4.4-6.8	22.0	3.8	3.1-4.6	12.2	4.8	3.6-6.2	9.8	3.0	2.2-4.1
Colorectal	1976-1980	197.6	73.3	68.7-78.2	94.8	86.2	78.1-94.9	102.8	65.5	59.9-71.4	98.8	38.0	34.6-41.6	48.4	47.1	41.0-53.8	50.4	32.5	28.6-36.8
	1981-1985	234.6	78.6	74.0-83.3	120.6	101.7	93.3-110.7	114.0	64.4	59.2-70.0	98.8	33.8	30.8-36.9	48.4	42.4	36.9-48.4	50.4	28.7	25.2-32.5
	1986-1990	229.0	71.1	67.0-75.4	109.6	83.5	76.3-91.1	119.4	62.3	57.4-67.6	101.8	32.2	29.4-35.2	49.4	39.1	34.1-44.6	52.4	27.5	24.3-31.1
	1991-1995	228.0	64.1	60.4-68.0	119.0	83.1	76.2-90.3	109.0	51.5	47.2-56.0	96.0	27.9	25.4-30.5	49.2	37.6	32.9-42.8	46.8	22.3	19.5-25.3
	1996-2000	272.2	67.5	63.9-71.2	136.6	83.1	76.7-89.7	135.6	57.2	52.9-61.7	108.8	27.3	25.1-29.7	54.0	34.2	30.2-38.7	54.8	22.8	20.2-25.7
	2001-2005	252.2	56.5	53.4-59.7	127.2	67.4	62.1-73.0	125.0	47.8	44.1-51.8	92.4	21.0	19.2-23.0	44.4	25.3	22.0-28.9	48.0	18.1	15.9-20.6
	2006-2010	239.0	49.3	46.5-52.2	122.0	59.1	54.4-64.1	117.0	42.0	38.6-45.6	86.4	18.1	16.4-19.9	46.6	24.4	21.3-27.9	39.8	14.1	12.2-16.2
	2011-2015	233.6	43.3	40.8-45.9	116.4	49.1	45.1-53.4	117.2	38.4	35.3-41.8	84.8	15.9	14.4-17.5	40.4	18.1	15.6-20.8	44.4	14.2	12.4-16.3
2016-2020	240.6	41.0	38.7-43.4	130.2	49.3	45.5-53.4	110.4	34.0	31.2-37.1	80.6	13.4	12.1-14.9	42.6	16.6	14.4-19.1	38.0	10.9	9.4-12.6	
Colon excluding rectum	1976-1980	143.2	53.9	49.9-58.0	65.8	61.4	54.5-68.8	77.4	49.5	44.7-54.8	83.0	31.7	28.7-35.0	38.8	37.2	31.8-43.2	44.2	28.4	24.8-32.5
	1981-1985	163.2	55.3	51.5-59.3	83.4	71.2	64.1-78.8	79.8	45.6	41.2-50.3	87.4	29.8	27.1-32.8	42.8	37.2	32.1-42.9	44.6	25.5	22.2-29.1
	1986-1990	160.8	50.3	46.8-53.9	69.6	54.0	48.2-60.3	91.2	47.6	43.3-52.1	91.4	28.9	26.3-31.8	43.6	34.6	29.9-39.8	47.8	25.1	22.0-28.5
	1991-1995	165.2	46.7	43.5-50.0	84.4	59.5	53.7-65.7	80.8	38.2	34.6-42.2	88.0	25.5	23.2-28.1	46.0	35.3	30.6-40.3	42.0	20.0	17.4-22.9
	1996-2000	196.0	48.7	45.7-51.9	93.6	57.3	52.0-62.9	102.4	43.1	39.4-47.0	94.8	23.8	21.7-26.0	46.8	29.4	25.6-33.5	48.0	20.0	17.5-22.7
	2001-2005	184.2	41.4	38.8-44.2	87.0	46.6	42.2-51.3	97.2	37.1	33.9-40.6	81.0	18.5	16.7-20.4	37.2	21.2	18.2-24.6	43.8	16.6	14.4-19.0
	2006-2010	173.6	35.9	33.5-38.4	86.8	42.3	38.3-46.6	86.8	31.1	28.2-34.2	72.6	15.2	13.7-16.9	39.8	21.0	18.1-24.2	32.8	11.6	9.9-13.5
	2011-2015	167.0	31.1	29.0-33.4	79.8	34.2	30.8-37.9	87.2	28.5	25.8-31.4	71.6	13.5	12.1-15.0	33.6	15.2	12.9-17.8	38.0	12.2	10.5-14.1
2016-2020	177.4	30.1	28.1-32.2	93.0	35.4	32.2-38.9	84.4	25.7	23.3-28.4	66.4	11.1	9.9-12.4	33.6	13.1	11.2-15.4	32.8	9.5	8.1-11.1	
Rectum & rectosigmoid	1976-1980	54.4	19.5	17.2-22.0	29.0	24.8	20.7-29.5	25.4	15.9	13.3-19.0	15.8	6.3	4.9-7.8	9.6	9.9	7.1-13.2	6.2	4.1	2.8-5.8
	1981-1985	71.4	23.3	20.9-25.9	37.2	30.5	26.0-35.6	34.2	18.8	16.1-21.9	11.4	3.9	2.9-5.1	5.6	5.2	3.3-7.5	5.8	3.2	2.1-4.6
	1986-1990	68.2	20.8	18.6-23.2	40.0	29.5	25.4-34.0	28.2	14.8	12.4-17.4	10.4	3.3	2.4-4.3	5.8	4.5	3.0-6.6	4.6	2.4	1.5-3.6
	1991-1995	62.8	17.4	15.5-19.5	34.6	23.6	20.0-27.5	28.2	13.2	11.1-15.6	8.0	2.3	1.6-3.1	3.2	2.4	1.3-3.9	4.8	2.2	1.4-3.3
	1996-2000	76.2	18.7	16.9-20.7	43.0	25.8	22.3-29.6	33.2	14.1	12.0-16.4	14.0	3.5	2.8-4.5	7.2	4.8	3.3-6.7	6.8	2.8	2.0-4.0
	2001-2005	68.0	15.1	13.5-16.8	40.2	20.8	17.9-23.9	27.8	10.7	9.0-12.6	11.4	2.5	1.9-3.3	7.2	4.0	2.8-5.6	4.2	1.5	0.9-2.3
	2006-2010	65.4	13.4	12.0-14.9	35.2	16.8	14.4-19.6	30.2	10.9	9.2-12.8	13.8	2.9	2.2-3.6	6.8	3.5	2.4-4.9	7.0	2.5	1.7-3.5
	2011-2015	66.6	12.1	10.8-13.6	36.6	14.9	12.8-17.3	30.0	9.9	8.4-11.7	13.2	2.4	1.8-3.0	6.8	2.9	2.0-4.1	6.4	2.0	1.4-2.9
2016-2020	63.2	10.9	9.7-12.2	37.2	13.9	11.9-16.1	26.0	8.3	6.9-9.9	14.2	2.3	1.8-3.0	9.0	3.5	2.5-4.7	5.2	1.4	0.9-2.1	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020  
Richmond (Staten Island)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	12.4	4.5	3.4-5.8	7.0	5.8	4.0-8.2	5.4	3.4	2.2-4.9	13.0	4.9	3.8-6.3	7.4	6.7	4.6-9.3	5.6	3.5	2.3-5.1
	1981-1985	12.8	4.3	3.3-5.5	7.0	5.8	3.9-8.2	5.8	3.3	2.2-4.8	17.8	6.1	4.9-7.5	9.8	8.3	6.0-11.1	8.0	4.7	3.3-6.4
	1986-1990	15.4	4.8	3.8-6.0	9.4	6.8	4.9-9.1	6.0	3.2	2.2-4.6	21.2	6.7	5.4-8.1	13.0	10.4	7.9-13.4	8.2	4.3	3.1-5.9
	1991-1995	17.0	4.7	3.7-5.8	11.0	7.0	5.2-9.1	6.0	2.8	1.9-4.0	22.6	6.3	5.2-7.6	13.4	8.9	6.8-11.3	9.2	4.4	3.2-5.8
	1996-2000	27.0	6.6	5.6-7.8	19.0	10.9	8.8-13.4	8.0	3.3	2.4-4.6	22.4	5.6	4.6-6.7	14.4	8.6	6.7-10.8	8.0	3.3	2.4-4.5
	2001-2005	34.8	7.6	6.5-8.8	26.2	13.3	11.0-15.8	8.6	3.3	2.4-4.4	28.4	6.3	5.3-7.4	18.8	9.7	7.8-11.9	9.6	3.7	2.7-4.9
	2006-2010	52.6	10.2	9.0-11.5	38.0	16.1	13.8-18.6	14.6	5.1	4.0-6.5	34.2	6.9	5.9-8.0	21.4	9.7	7.9-11.9	12.8	4.6	3.6-6.0
	2011-2015	62.8	10.9	9.7-12.2	45.8	17.4	15.1-19.9	17.0	5.6	4.4-6.9	39.6	7.0	6.1-8.1	29.2	11.5	9.7-13.7	10.4	3.5	2.6-4.6
	2016-2020	65.2	10.3	9.2-11.5	47.2	16.4	14.3-18.7	18.0	5.3	4.2-6.5	39.2	6.3	5.5-7.3	25.4	9.1	7.6-11.0	13.8	4.0	3.1-5.1
Pancreas	1976-1980	30.0	10.9	9.2-12.8	16.4	14.1	11.1-17.8	13.6	8.6	6.7-10.9	29.4	10.8	9.1-12.7	15.6	13.8	10.7-17.4	13.8	8.7	6.8-11.0
	1981-1985	35.8	12.3	10.5-14.2	16.6	14.1	11.1-17.7	19.2	11.1	9.0-13.6	32.8	11.1	9.4-13.0	14.8	12.4	9.6-15.7	18.0	10.3	8.3-12.7
	1986-1990	48.6	15.1	13.2-17.1	22.8	17.6	14.4-21.4	25.8	13.4	11.2-15.9	41.4	12.8	11.1-14.7	21.6	16.8	13.6-20.4	19.8	10.3	8.4-12.6
	1991-1995	52.0	14.7	13.0-16.7	24.4	16.6	13.7-20.0	27.6	13.0	10.9-15.4	46.2	13.1	11.4-14.9	22.2	15.0	12.2-18.1	24.0	11.3	9.4-13.5
	1996-2000	62.4	15.4	13.7-17.2	31.8	18.6	15.7-21.8	30.6	12.8	10.9-15.0	53.2	13.2	11.6-14.9	26.0	15.3	12.7-18.2	27.2	11.3	9.5-13.4
	2001-2005	60.4	13.4	11.9-15.0	27.4	14.2	11.8-16.8	33.0	12.6	10.7-14.7	55.2	12.3	10.9-13.9	28.8	15.5	13.0-18.3	26.4	10.0	8.3-11.8
	2006-2010	74.4	15.1	13.6-16.7	35.4	16.4	14.0-19.1	39.0	13.8	11.9-15.9	59.4	12.1	10.7-13.6	28.6	13.5	11.3-16.0	30.8	10.8	9.1-12.7
	2011-2015	76.0	13.9	12.5-15.4	38.8	15.9	13.7-18.4	37.2	12.1	10.4-14.1	57.6	10.6	9.4-11.9	27.8	11.7	9.7-13.8	29.8	9.7	8.2-11.4
	2016-2020	94.2	15.5	14.1-17.0	46.6	17.0	14.8-19.5	47.6	14.1	12.3-16.1	73.4	12.0	10.8-13.3	35.0	13.2	11.2-15.4	38.4	11.1	9.6-12.9
Larynx	1976-1980	23.2	7.8	6.4-9.4	18.6	14.8	11.8-18.3	4.6	2.8	1.8-4.2	8.8	3.0	2.1-4.0	7.4	5.9	4.0-8.3	1.4	0.9	0.4-1.9
	1981-1985	25.2	8.0	6.7-9.6	19.8	15.2	12.3-18.7	5.4	3.0	2.0-4.4	8.2	2.7	1.9-3.7	5.8	4.7	3.1-6.8	2.4	1.4	0.7-2.5
	1986-1990	25.4	7.4	6.1-8.8	19.6	13.5	10.8-16.5	5.8	3.2	2.1-4.6	7.8	2.3	1.6-3.1	7.0	5.0	3.4-7.1	0.8	0.4	0.1-1.1
	1991-1995	20.4	5.5	4.5-6.7	15.6	9.8	7.7-12.3	4.8	2.3	1.5-3.4	7.2	1.9	1.3-2.7	4.8	3.2	2.0-4.8	2.4	1.2	0.6-2.0
	1996-2000	19.4	4.6	3.7-5.6	15.0	7.9	6.2-10.0	4.4	1.9	1.2-2.9	7.4	1.8	1.3-2.5	5.8	3.6	2.3-5.1	1.6	0.7	0.3-1.3
	2001-2005	22.8	4.8	4.0-5.8	18.0	8.8	7.0-10.9	4.8	1.9	1.2-2.8	5.8	1.3	0.8-1.8	5.2	2.7	1.8-4.1	0.6	0.2	0.0-0.7
	2006-2010	25.4	5.0	4.2-6.0	19.8	8.7	7.0-10.6	5.6	2.0	1.4-3.0	8.6	1.7	1.2-2.3	7.2	3.3	2.3-4.5	1.4	0.5	0.2-1.1
	2011-2015	23.0	4.0	3.3-4.9	17.8	7.1	5.6-8.8	5.2	1.7	1.1-2.6	5.8	1.1	0.7-1.6	4.6	2.1	1.3-3.2	1.2	0.4	0.1-0.9
	2016-2020	17.6	2.9	2.3-3.6	14.0	5.2	4.0-6.6	3.6	1.1	0.6-1.8	4.2	0.7	0.4-1.0	2.4	0.9	0.4-1.5	1.8	0.5	0.2-1.0
Lung and bronchus	1976-1980	185.8	64.8	60.6-69.2	130.4	111.3	102.4-120.6	55.4	33.7	29.8-38.0	152.4	53.3	49.5-57.3	110.8	94.7	86.5-103.3	41.6	25.4	22.0-29.1
	1981-1985	229.6	73.1	68.9-77.6	158.6	124.3	115.3-133.7	71.0	39.3	35.3-43.7	190.4	60.7	56.8-64.7	130.4	102.4	94.2-111.0	60.0	33.2	29.5-37.2
	1986-1990	256.4	76.9	72.7-81.3	160.6	118.5	110.0-127.3	95.8	49.7	45.3-54.4	199.0	59.4	55.7-63.3	128.4	94.4	86.9-102.4	70.6	36.4	32.6-40.4
	1991-1995	296.2	81.3	77.2-85.5	167.0	110.6	103.0-118.6	129.2	62.2	57.4-67.2	231.2	64.0	60.3-67.8	135.2	93.0	85.9-100.6	96.0	46.0	42.0-50.4
	1996-2000	307.8	75.2	71.5-79.1	164.2	97.1	90.4-104.1	143.6	61.7	57.3-66.4	235.2	58.0	54.7-61.4	130.8	79.5	73.4-86.0	104.4	44.6	40.9-48.6
	2001-2005	322.0	70.7	67.3-74.3	173.6	90.1	84.1-96.4	148.4	57.4	53.3-61.7	239.2	53.0	50.0-56.1	134.4	71.2	65.8-77.0	104.8	40.6	37.2-44.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**Richmond (Staten Island)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	341.8	70.0	66.7-73.4	176.2	85.2	79.5-91.2	165.6	59.8	55.7-64.0	237.6	49.2	46.4-52.1	124.8	62.2	57.3-67.4	112.8	40.6	37.3-44.1
	2011-2015	355.6	65.2	62.2-68.4	181.8	76.5	71.4-81.8	173.8	57.1	53.3-61.1	219.6	40.4	38.0-42.9	117.2	50.6	46.4-55.0	102.4	33.3	30.4-36.3
	2016-2020	384.8	63.6	60.8-66.6	182.0	69.1	64.6-73.9	202.8	60.3	56.6-64.2	197.0	33.1	31.0-35.3	99.4	39.3	35.8-43.0	97.6	28.8	26.3-31.6
Melanoma of the skin	1976-1980	21.2	7.2	5.9-8.7	10.6	8.0	5.9-10.6	10.6	6.6	4.9-8.7	5.2	1.9	1.2-2.8	2.8	2.2	1.1-3.7	2.4	1.6	0.8-2.8
	1981-1985	25.8	8.3	6.9-9.9	14.8	11.0	8.5-13.9	11.0	6.3	4.8-8.3	8.8	2.8	2.0-3.8	5.2	3.4	2.2-5.1	3.6	2.3	1.3-3.6
	1986-1990	33.8	9.9	8.4-11.5	18.8	13.0	10.4-16.1	15.0	7.7	6.1-9.7	6.8	2.1	1.4-2.9	5.0	3.3	2.1-5.0	1.8	1.0	0.5-1.9
	1991-1995	30.2	8.0	6.8-9.4	18.6	11.5	9.2-14.1	11.6	5.6	4.3-7.3	6.6	1.8	1.2-2.5	4.4	2.9	1.8-4.4	2.2	1.1	0.5-1.9
	1996-2000	40.0	9.5	8.2-10.9	21.0	11.1	9.0-13.5	19.0	8.2	6.6-10.0	8.8	2.1	1.5-2.8	4.6	2.3	1.5-3.5	4.2	1.8	1.1-2.8
	2001-2005	57.6	12.4	11.0-14.0	30.6	14.8	12.5-17.4	27.0	10.7	8.9-12.6	8.4	1.8	1.3-2.4	3.8	1.8	1.1-2.9	4.6	1.8	1.1-2.7
	2006-2010	71.6	14.6	13.1-16.2	39.0	17.8	15.3-20.5	32.6	12.3	10.5-14.4	8.8	1.8	1.3-2.4	6.4	3.0	2.0-4.3	2.4	0.9	0.4-1.5
	2011-2015	100.0	18.6	16.9-20.3	56.8	24.0	21.2-27.1	43.2	15.0	13.0-17.2	9.6	1.8	1.3-2.4	6.2	2.7	1.8-3.9	3.4	1.2	0.7-1.9
	2016-2020	84.2	14.4	13.0-15.9	45.6	17.5	15.2-20.0	38.6	12.1	10.4-14.1	8.4	1.5	1.1-2.1	4.2	1.9	1.2-2.9	4.2	1.3	0.8-2.1
Female breast	1976-1980							158.4	101.6	94.5-109.0							61.0	38.4	34.2-43.1
	1981-1985							192.8	111.8	104.8-119.2							58.4	33.4	29.7-37.6
	1986-1990							262.0	139.5	132.0-147.3							70.4	37.4	33.6-41.6
	1991-1995							279.8	136.5	129.4-143.8							80.2	38.5	34.8-42.5
	1996-2000							320.8	138.7	132.0-145.7							74.8	31.9	28.7-35.3
	2001-2005							334.0	130.0	123.8-136.5							78.0	29.6	26.7-32.7
	2006-2010							364.0	131.9	125.8-138.2							67.4	23.9	21.4-26.7
	2011-2015							398.4	135.5	129.5-141.8							57.0	18.0	15.9-20.3
	2016-2020							430.6	139.2	133.1-145.4							60.4	17.6	15.6-19.8
Cervix uteri	1976-1980							16.2	10.2	8.0-12.7							5.0	3.2	2.1-4.8
	1981-1985							17.4	9.7	7.7-12.0							4.2	2.2	1.4-3.5
	1986-1990							26.8	13.9	11.6-16.5							5.2	2.8	1.8-4.1
	1991-1995							28.4	13.5	11.4-16.0							6.0	2.9	2.0-4.2
	1996-2000							25.0	10.8	9.0-12.9							7.4	3.2	2.3-4.4
	2001-2005							25.4	10.1	8.4-12.0							5.8	2.2	1.5-3.2
	2006-2010							22.8	9.0	7.4-10.8							5.0	1.8	1.2-2.7
	2011-2015							19.6	7.3	5.9-9.0							7.4	2.5	1.7-3.5
	2016-2020							22.2	7.9	6.4-9.6							5.4	1.7	1.1-2.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**Richmond (Staten Island)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							32.6	19.9	16.9-23.2							10.0	6.2	4.6-8.3
	1981-1985							39.8	22.3	19.3-25.7							6.2	3.5	2.4-5.0
	1986-1990							50.4	25.8	22.7-29.2							8.8	4.5	3.3-6.1
	1991-1995							55.2	26.9	23.8-30.3							12.8	6.0	4.6-7.7
	1996-2000							67.8	29.2	26.2-32.5							10.6	4.5	3.4-5.9
	2001-2005							76.6	29.3	26.4-32.4							15.2	5.8	4.6-7.3
	2006-2010							93.0	32.6	29.7-35.8							15.0	5.5	4.3-6.9
	2011-2015							108.6	35.4	32.4-38.5							17.4	5.6	4.5-7.0
	2016-2020							116.4	34.9	32.0-37.9							22.0	6.5	5.3-7.8
Ovary	1976-1980							22.8	13.9	11.4-16.7							16.4	10.1	8.0-12.5
	1981-1985							28.8	16.3	13.7-19.3							14.4	8.1	6.4-10.3
	1986-1990							39.8	20.9	18.1-24.0							20.0	10.3	8.3-12.5
	1991-1995							44.4	21.6	18.8-24.6							25.0	12.1	10.0-14.4
	1996-2000							41.2	17.9	15.5-20.5							20.2	8.7	7.1-10.6
	2001-2005							32.6	12.6	10.7-14.7							25.4	9.7	8.1-11.6
	2006-2010							37.6	13.6	11.7-15.7							25.6	9.0	7.5-10.8
	2011-2015							38.4	13.0	11.2-15.1							23.0	7.4	6.1-8.9
	2016-2020							45.6	15.0	13.0-17.1							22.8	6.6	5.4-8.0
Prostate	1976-1980				79.4	81.2	72.9-90.0							24.8	27.3	22.5-32.7			
	1981-1985				88.2	78.5	70.9-86.6							35.2	34.2	29.1-39.9			
	1986-1990				106.4	88.6	80.8-96.9							37.0	34.3	29.3-39.8			
	1991-1995				210.6	146.6	137.6-156.1							42.2	36.1	31.1-41.5			
	1996-2000				294.2	170.0	161.2-179.1							44.8	32.4	28.2-37.0			
	2001-2005				315.8	158.5	150.7-166.7							37.0	23.8	20.4-27.6			
	2006-2010				363.6	162.2	154.6-170.1							32.6	18.7	15.9-21.8			
	2011-2015				330.2	126.9	120.7-133.4							35.6	18.0	15.4-20.9			
	2016-2020				347.4	119.5	113.8-125.4							35.8	15.6	13.4-18.2			
Testis	1976-1980				5.6	3.1	2.0-4.6							0.6	0.6	0.1-1.7			
	1981-1985				8.4	4.3	3.0-5.8							0.6	0.3	0.1-1.0			
	1986-1990				8.8	4.2	3.1-5.8							0.6	0.3	0.1-0.9			
	1991-1995				13.0	6.1	4.7-7.8							0.8	0.5	0.1-1.2			
	1996-2000				15.0	6.8	5.4-8.6							0.4	0.2	0.0-0.7			
	2001-2005				13.6	6.1	4.7-7.7							0.8	0.4	0.1-1.0			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**Richmond (Staten Island)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				14.0	6.4	4.9-8.0							0.6	0.2	0.0-0.8			
	2011-2015				14.0	6.3	4.9-8.0							0.4	0.2	0.0-0.6			
	2016-2020				14.2	6.6	5.1-8.3							1.0	0.4	0.1-1.0			
Urinary bladder (incl. in situ)	1976-1980	66.2	24.5	21.8-27.3	48.0	43.3	37.7-49.5	18.2	11.7	9.4-14.4	17.8	7.1	5.7-8.7	11.8	11.6	8.7-15.1	6.0	4.1	2.8-5.8
	1981-1985	76.4	24.9	22.4-27.6	57.6	47.1	41.5-53.2	18.8	10.4	8.4-12.7	16.8	5.8	4.6-7.3	11.6	11.0	8.2-14.4	5.2	2.9	1.9-4.3
	1986-1990	83.2	25.5	23.1-28.1	59.4	45.3	40.0-51.1	23.8	12.5	10.4-15.0	19.6	6.5	5.2-7.9	13.6	12.1	9.2-15.4	6.0	3.2	2.2-4.6
	1991-1995	91.0	25.3	23.0-27.8	65.4	45.2	40.2-50.6	25.6	12.2	10.1-14.5	25.0	7.2	5.9-8.5	17.6	13.4	10.7-16.6	7.4	3.4	2.4-4.8
	1996-2000	103.8	25.5	23.4-27.8	75.0	44.9	40.3-49.8	28.8	12.2	10.3-14.3	25.2	6.4	5.3-7.6	16.4	10.7	8.5-13.4	8.8	3.6	2.6-4.8
	2001-2005	121.8	27.3	25.2-29.6	87.8	47.7	43.3-52.5	34.0	13.1	11.2-15.3	22.6	5.2	4.3-6.2	15.0	9.0	7.0-11.3	7.6	2.9	2.1-4.1
	2006-2010	130.0	26.8	24.8-29.0	96.8	47.0	42.8-51.5	33.2	11.9	10.1-13.9	21.0	4.4	3.6-5.3	15.0	7.8	6.1-9.9	6.0	2.1	1.4-3.0
	2011-2015	142.6	26.2	24.3-28.3	111.6	48.1	44.1-52.4	31.0	10.1	8.5-11.8	27.2	5.1	4.3-6.1	20.2	9.5	7.7-11.5	7.0	2.1	1.5-3.0
	2016-2020	144.4	24.0	22.2-25.8	106.2	40.7	37.2-44.4	38.2	11.2	9.6-12.9	26.8	4.6	3.8-5.4	18.6	7.9	6.4-9.8	8.2	2.4	1.7-3.3
Kidney and renal pelvis	1976-1980	25.0	8.7	7.2-10.4	15.6	12.8	10.0-16.2	9.4	5.8	4.3-7.8	10.6	3.8	2.9-5.0	6.4	5.8	3.9-8.3	4.2	2.6	1.6-3.9
	1981-1985	26.6	8.4	7.0-10.0	16.6	12.5	9.8-15.6	10.0	5.4	4.0-7.2	11.4	3.8	2.8-4.9	7.2	5.9	4.0-8.2	4.2	2.3	1.4-3.5
	1986-1990	30.6	9.1	7.7-10.6	18.8	13.1	10.5-16.2	11.8	6.2	4.7-8.0	13.6	4.0	3.1-5.1	8.0	5.7	4.0-7.8	5.6	2.9	1.9-4.2
	1991-1995	42.2	11.3	9.8-12.9	25.2	15.5	12.9-18.5	17.0	8.1	6.4-10.0	12.2	3.3	2.5-4.3	7.4	4.8	3.3-6.7	4.8	2.2	1.4-3.3
	1996-2000	50.6	12.2	10.7-13.8	29.6	16.5	13.9-19.4	21.0	9.0	7.4-10.9	16.0	4.0	3.2-5.0	7.8	4.9	3.4-6.7	8.2	3.4	2.5-4.7
	2001-2005	79.4	17.2	15.5-19.0	52.4	25.1	22.1-28.4	27.0	10.6	8.9-12.5	15.2	3.4	2.7-4.3	8.6	4.5	3.2-6.0	6.6	2.5	1.7-3.6
	2006-2010	90.8	18.0	16.4-19.8	58.0	25.0	22.1-28.2	32.8	12.1	10.3-14.1	17.2	3.5	2.8-4.4	11.2	5.3	3.9-6.9	6.0	2.1	1.4-3.1
	2011-2015	105.0	19.0	17.4-20.7	70.2	27.4	24.5-30.5	34.8	11.9	10.1-13.8	20.2	3.8	3.1-4.6	12.0	5.0	3.8-6.5	8.2	2.8	2.0-3.8
	2016-2020	126.0	21.2	19.5-23.0	85.4	31.4	28.4-34.6	40.6	12.6	10.9-14.5	18.8	3.1	2.5-3.9	13.2	5.1	4.0-6.6	5.6	1.6	1.1-2.4
Brain and other central nervous system	1976-1980	19.2	6.1	4.9-7.5	11.6	7.9	5.9-10.4	7.6	4.6	3.2-6.3	13.2	4.3	3.3-5.5	8.4	6.1	4.3-8.3	4.8	2.9	1.8-4.3
	1981-1985	22.4	6.6	5.4-7.9	11.6	7.7	5.7-10.0	10.8	5.9	4.4-7.7	14.8	4.5	3.5-5.6	7.0	5.2	3.6-7.4	7.8	4.1	2.9-5.7
	1986-1990	26.2	7.5	6.2-8.9	14.4	9.1	7.0-11.5	11.8	6.1	4.6-7.8	14.8	4.4	3.5-5.6	7.8	5.5	3.9-7.7	7.0	3.7	2.5-5.1
	1991-1995	29.0	7.7	6.5-9.1	17.0	10.4	8.2-12.9	12.0	5.7	4.3-7.3	15.6	4.3	3.4-5.4	9.2	5.9	4.3-8.0	6.4	3.1	2.1-4.4
	1996-2000	32.0	7.6	6.5-8.9	15.0	8.0	6.3-10.1	17.0	7.5	6.0-9.3	15.6	3.8	3.0-4.7	6.6	3.7	2.5-5.2	9.0	3.9	2.8-5.2
	2001-2005	29.8	6.5	5.5-7.6	17.0	8.1	6.4-10.0	12.8	5.2	4.0-6.6	16.0	3.5	2.8-4.4	8.4	4.1	3.0-5.6	7.6	3.0	2.1-4.1
	2006-2010	34.4	7.2	6.1-8.3	17.2	7.6	6.0-9.4	17.2	6.9	5.5-8.5	18.0	3.6	2.9-4.5	9.8	4.3	3.2-5.8	8.2	3.0	2.1-4.1
	2011-2015	35.2	7.0	6.0-8.1	18.2	7.9	6.3-9.7	17.0	6.2	4.9-7.7	21.2	4.0	3.3-4.9	11.4	4.9	3.7-6.4	9.8	3.3	2.4-4.4
	2016-2020	36.2	6.7	5.7-7.8	19.0	7.5	6.1-9.3	17.2	5.9	4.6-7.3	21.6	3.7	3.0-4.5	11.4	4.3	3.2-5.6	10.2	3.1	2.3-4.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020  
Richmond (Staten Island)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	13.2	4.4	3.4-5.6	2.8	2.1	1.1-3.5	10.4	6.2	4.6-8.2	1.0	0.4	0.1-0.9	0.4	0.3	0.0-1.1	0.6	0.4	0.1-1.2
	1981-1985	11.6	3.4	2.6-4.4	4.4	2.7	1.7-4.1	7.2	4.0	2.8-5.5	1.6	0.5	0.2-1.1	0.8	0.7	0.2-1.8	0.8	0.4	0.1-1.1
	1986-1990	16.2	4.5	3.6-5.6	4.8	2.8	1.8-4.2	11.4	5.8	4.4-7.6	2.4	0.8	0.4-1.3	1.0	0.6	0.2-1.4	1.4	0.7	0.3-1.5
	1991-1995	26.6	6.7	5.6-8.0	6.8	3.8	2.6-5.4	19.8	9.2	7.5-11.3	1.4	0.4	0.2-0.8	0.4	0.2	0.0-0.9	1.0	0.5	0.2-1.1
	1996-2000	28.6	6.6	5.6-7.8	9.4	4.6	3.4-6.1	19.2	8.4	6.8-10.3	1.2	0.3	0.1-0.6	0.2	0.1	0.0-0.5	1.0	0.4	0.1-1.0
	2001-2005	64.8	13.7	12.3-15.3	16.4	7.3	5.8-9.1	48.4	19.5	17.2-22.2	2.4	0.6	0.3-1.0	0.6	0.3	0.1-0.9	1.8	0.7	0.3-1.3
	2006-2010	138.2	27.9	25.8-30.1	34.0	14.1	12.1-16.5	104.2	40.5	37.0-44.2	3.0	0.6	0.3-1.0	1.0	0.4	0.1-1.0	2.0	0.7	0.3-1.3
	2011-2015	169.6	33.2	31.0-35.6	45.0	18.3	15.9-20.9	124.6	47.2	43.4-51.1	2.2	0.4	0.2-0.7	1.2	0.4	0.1-1.0	1.0	0.4	0.1-0.9
	2016-2020	164.4	31.8	29.6-34.2	48.6	19.2	16.8-21.9	115.8	43.7	40.1-47.5	3.0	0.5	0.3-0.8	1.6	0.6	0.3-1.3	1.4	0.4	0.1-0.8
Hodgkin lymphoma	1976-1980	13.6	4.0	3.1-5.1	8.8	6.0	4.2-8.2	4.8	2.5	1.6-3.8	5.0	1.5	1.0-2.3	3.8	2.7	1.6-4.4	1.2	0.7	0.2-1.5
	1981-1985	17.2	4.8	3.8-5.9	10.2	5.8	4.2-7.7	7.0	3.8	2.6-5.3	4.4	1.3	0.8-1.9	2.0	1.3	0.6-2.5	2.4	1.2	0.6-2.2
	1986-1990	19.2	4.9	4.0-6.0	9.0	4.8	3.5-6.6	10.2	4.9	3.6-6.4	3.8	1.1	0.6-1.7	2.6	1.4	0.7-2.5	1.2	0.6	0.2-1.4
	1991-1995	16.2	4.0	3.2-5.0	8.0	4.3	3.0-5.9	8.2	3.9	2.8-5.2	3.6	0.9	0.5-1.4	1.8	0.9	0.4-1.8	1.8	0.9	0.4-1.7
	1996-2000	17.0	4.0	3.2-4.9	8.6	4.1	3.0-5.5	8.4	3.7	2.7-5.1	2.2	0.5	0.2-0.9	1.4	0.7	0.3-1.4	0.8	0.3	0.1-0.9
	2001-2005	15.6	3.4	2.7-4.3	8.4	4.0	2.8-5.4	7.2	3.1	2.1-4.2	1.4	0.3	0.1-0.6	1.0	0.5	0.2-1.2	0.4	0.2	0.0-0.6
	2006-2010	17.0	3.7	2.9-4.5	9.6	4.2	3.1-5.6	7.4	3.2	2.2-4.4	1.4	0.3	0.1-0.6	1.2	0.5	0.2-1.1	0.2	0.1	0.0-0.5
	2011-2015	19.0	4.0	3.2-4.8	10.0	4.2	3.1-5.6	9.0	3.7	2.7-5.0	1.8	0.3	0.2-0.7	1.0	0.5	0.2-1.1	0.8	0.3	0.1-0.7
	2016-2020	16.0	3.3	2.6-4.1	9.2	3.9	2.8-5.3	6.8	2.7	1.8-3.8	2.6	0.4	0.2-0.8	1.6	0.6	0.2-1.2	1.0	0.4	0.1-0.9
Non-Hodgkin Lymphomas	1976-1980	31.6	10.9	9.2-12.7	18.6	14.1	11.3-17.5	13.0	8.0	6.2-10.2	15.2	5.5	4.3-6.9	9.6	8.2	5.9-11.1	5.6	3.6	2.4-5.2
	1981-1985	44.4	13.8	12.0-15.8	22.0	14.8	12.1-18.0	22.4	12.6	10.3-15.1	24.0	7.6	6.3-9.1	10.8	7.7	5.6-10.1	13.2	7.3	5.6-9.3
	1986-1990	59.8	17.7	15.8-19.9	30.4	21.2	17.7-25.0	29.4	15.2	12.8-17.9	21.6	6.5	5.3-7.9	12.2	8.4	6.3-11.0	9.4	4.9	3.6-6.5
	1991-1995	77.8	20.8	18.8-23.0	40.0	24.0	20.7-27.7	37.8	18.1	15.6-20.9	29.2	7.9	6.7-9.3	16.4	10.2	8.0-12.8	12.8	6.1	4.7-7.8
	1996-2000	94.8	22.8	20.8-25.0	50.4	27.7	24.3-31.5	44.4	19.1	16.6-21.7	33.6	8.3	7.1-9.7	18.2	10.7	8.6-13.2	15.4	6.5	5.1-8.1
	2001-2005	108.4	23.7	21.7-25.8	57.6	28.6	25.4-32.2	50.8	19.9	17.5-22.6	33.2	7.4	6.3-8.6	16.4	8.6	6.8-10.7	16.8	6.5	5.2-8.1
	2006-2010	113.6	23.4	21.5-25.5	59.4	27.8	24.7-31.3	54.2	20.2	17.8-22.8	29.4	6.2	5.3-7.4	16.4	8.0	6.3-10.0	13.0	4.9	3.8-6.3
	2011-2015	128.6	24.1	22.3-26.1	68.2	29.0	25.9-32.4	60.4	20.4	18.1-22.9	25.2	4.8	4.0-5.7	15.2	7.0	5.5-8.8	10.0	3.3	2.4-4.4
	2016-2020	138.4	23.9	22.1-25.9	72.2	27.9	25.0-31.1	66.2	20.7	18.4-23.1	25.8	4.5	3.8-5.4	14.6	6.0	4.7-7.6	11.2	3.3	2.5-4.3
Myeloma	1976-1980	11.2	4.1	3.1-5.4	4.8	4.3	2.6-6.5	6.4	4.0	2.8-5.7	6.8	2.4	1.7-3.4	2.6	2.1	1.1-3.7	4.2	2.6	1.6-4.0
	1981-1985	13.8	4.6	3.6-5.9	8.0	6.7	4.7-9.3	5.8	3.3	2.2-4.7	11.0	3.7	2.7-4.8	5.6	4.7	3.0-6.8	5.4	3.0	2.0-4.4
	1986-1990	17.8	5.3	4.2-6.5	7.8	5.2	3.6-7.2	10.0	5.3	3.9-7.0	10.0	3.1	2.3-4.1	4.6	3.4	2.1-5.2	5.4	2.8	1.9-4.1
	1991-1995	15.0	4.1	3.2-5.2	7.0	4.9	3.4-6.9	8.0	3.8	2.7-5.1	11.2	3.1	2.4-4.1	4.6	3.3	2.1-5.0	6.6	3.1	2.1-4.4
	1996-2000	26.0	6.3	5.3-7.5	13.4	7.9	6.1-10.1	12.6	5.4	4.1-6.9	11.2	2.8	2.1-3.6	4.8	3.1	1.9-4.6	6.4	2.7	1.8-3.8
	2001-2005	32.4	7.2	6.1-8.3	16.6	8.9	7.1-11.1	15.8	6.0	4.7-7.5	13.6	3.0	2.4-3.8	6.2	3.5	2.3-4.9	7.4	2.8	1.9-3.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**Richmond (Staten Island)**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	41.0	8.4	7.3-9.7	21.6	10.4	8.4-12.6	19.4	7.0	5.7-8.6	11.0	2.3	1.7-3.0	6.0	3.1	2.0-4.4	5.0	1.8	1.2-2.7
	2011-2015	49.4	9.0	7.9-10.2	29.6	12.6	10.6-14.9	19.8	6.3	5.1-7.7	12.4	2.4	1.8-3.0	6.0	2.7	1.8-3.9	6.4	2.1	1.4-3.0
	2016-2020	55.0	9.2	8.1-10.4	29.4	11.2	9.4-13.2	25.6	7.7	6.4-9.2	14.8	2.4	1.9-3.1	8.6	3.2	2.3-4.3	6.2	1.8	1.2-2.6
Leukemias	1976-1980	30.6	10.8	9.1-12.7	17.0	14.7	11.5-18.4	13.6	8.5	6.5-10.7	18.4	6.9	5.5-8.4	11.0	10.6	7.8-14.0	7.4	4.6	3.2-6.3
	1981-1985	35.8	11.8	10.1-13.7	21.0	17.6	14.2-21.6	14.8	8.5	6.6-10.7	25.8	8.6	7.1-10.2	13.8	11.7	8.9-15.0	12.0	6.8	5.2-8.8
	1986-1990	43.8	13.2	11.5-15.1	24.0	16.6	13.6-20.0	19.8	10.5	8.5-12.8	24.0	7.4	6.1-8.8	12.0	8.7	6.5-11.4	12.0	6.4	4.9-8.2
	1991-1995	52.2	14.2	12.5-16.0	31.6	20.3	17.2-23.9	20.6	9.6	7.9-11.7	28.0	7.9	6.6-9.3	16.4	11.5	9.1-14.4	11.6	5.5	4.2-7.1
	1996-2000	55.8	13.6	12.0-15.3	30.2	17.3	14.5-20.3	25.6	11.1	9.2-13.2	29.4	7.3	6.1-8.6	18.4	10.8	8.7-13.4	11.0	4.6	3.5-6.0
	2001-2005	58.4	13.0	11.6-14.6	29.6	15.1	12.7-17.8	28.8	11.3	9.6-13.4	26.8	6.0	5.0-7.1	13.6	7.2	5.5-9.1	13.2	5.0	3.9-6.4
	2006-2010	82.4	17.0	15.4-18.8	45.6	21.7	18.9-24.8	36.8	13.6	11.7-15.8	31.8	6.7	5.7-7.8	16.2	8.5	6.7-10.6	15.6	5.7	4.5-7.1
	2011-2015	99.6	18.9	17.2-20.7	55.2	23.6	20.8-26.7	44.4	15.8	13.7-18.1	28.8	5.5	4.6-6.5	15.6	7.0	5.5-8.7	13.2	4.6	3.5-5.9
2016-2020	96.4	17.1	15.6-18.8	55.8	21.7	19.1-24.4	40.6	13.3	11.5-15.4	37.2	6.5	5.5-7.5	18.8	7.7	6.2-9.5	18.4	5.5	4.4-6.9	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Albany**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	1,197.8	419.3	408.5-430.3	599.8	516.1	496.9-535.8	598.0	366.2	352.8-380.0	625.6	217.4	209.7-225.3	328.8	288.3	273.9-303.3	296.8	174.5	165.6-183.9
	1981-1985	1,346.6	452.1	441.1-463.2	663.2	558.0	538.5-578.1	683.4	398.2	384.5-412.4	666.4	222.0	214.4-229.8	343.0	296.6	282.1-311.5	323.4	181.5	172.5-190.8
	1986-1990	1,392.6	455.6	444.7-466.6	667.4	550.3	531.2-569.9	725.2	411.3	397.5-425.4	704.8	227.5	219.9-235.3	342.4	291.1	277.0-305.7	362.4	195.6	186.4-205.1
	1991-1995	1,441.8	453.8	443.3-464.5	722.4	558.2	539.8-577.1	719.4	394.7	381.5-408.2	696.2	215.3	208.1-222.6	338.2	272.4	259.3-286.0	358.0	183.7	175.1-192.7
	1996-2000	1,578.4	494.3	483.4-505.4	793.2	597.2	578.6-616.2	785.2	432.0	418.3-446.1	696.6	211.0	204.1-218.2	349.6	271.4	258.7-284.6	347.0	175.4	167.0-184.1
	2001-2005	1,639.4	498.6	487.8-509.6	825.4	593.1	575.1-611.7	814.0	438.5	424.9-452.5	633.2	186.8	180.4-193.5	309.6	228.5	217.2-240.3	323.6	162.5	154.5-170.8
	2006-2010	1,759.0	512.5	501.6-523.5	913.2	611.3	593.3-629.6	845.8	444.5	430.8-458.5	623.6	177.0	170.7-183.4	309.2	215.0	204.3-226.2	314.4	152.7	145.0-160.7
	2011-2015	1,854.0	513.9	503.2-524.8	907.2	560.7	543.9-577.7	946.8	486.5	472.2-501.2	625.2	169.3	163.3-175.5	317.8	205.7	195.4-216.3	307.4	144.3	136.9-152.0
	2016-2020	1,943.8	507.1	496.7-517.6	976.4	559.1	543.0-575.6	967.4	474.5	460.5-488.8	597.8	149.8	144.3-155.4	299.6	177.9	168.8-187.4	298.2	131.4	124.6-138.6
Oral cavity and pharynx	1976-1980	46.2	15.7	13.7-17.9	32.6	26.2	22.1-30.7	13.6	8.0	6.2-10.2	16.6	5.6	4.4-7.0	11.4	9.4	7.0-12.4	5.2	3.0	1.9-4.4
	1981-1985	43.8	14.8	12.9-17.0	27.0	22.3	18.6-26.6	16.8	9.8	7.8-12.2	16.8	5.5	4.4-6.9	11.0	9.2	6.9-12.1	5.8	3.1	2.1-4.5
	1986-1990	41.6	13.9	12.0-16.0	24.2	19.7	16.2-23.7	17.4	10.0	8.0-12.5	14.2	4.5	3.5-5.7	8.6	7.1	5.1-9.7	5.6	2.9	1.9-4.3
	1991-1995	30.6	10.1	8.5-11.9	19.2	15.2	12.2-18.6	11.4	6.4	4.8-8.4	10.6	3.4	2.5-4.5	6.4	5.3	3.6-7.5	4.2	2.1	1.3-3.3
	1996-2000	41.6	13.3	11.5-15.2	27.8	20.8	17.5-24.6	13.8	7.6	5.9-9.7	11.0	3.4	2.6-4.5	5.0	3.9	2.5-5.8	6.0	3.2	2.1-4.6
	2001-2005	31.4	9.6	8.1-11.2	18.6	12.7	10.2-15.6	12.8	6.9	5.3-8.8	8.2	2.4	1.7-3.3	4.2	2.9	1.8-4.5	4.0	1.9	1.2-3.1
	2006-2010	42.8	12.3	10.7-14.1	31.2	19.8	16.7-23.2	11.6	6.3	4.7-8.2	7.4	2.0	1.4-2.8	5.6	3.6	2.4-5.3	1.8	0.9	0.4-1.7
	2011-2015	40.8	10.9	9.4-12.6	29.6	17.1	14.4-20.2	11.2	5.5	4.1-7.3	8.6	2.2	1.6-3.0	6.0	3.3	2.2-4.8	2.6	1.2	0.6-2.2
	2016-2020	57.6	14.8	13.1-16.7	42.4	23.7	20.5-27.3	15.2	7.3	5.7-9.2	10.4	2.5	1.9-3.3	8.0	4.3	3.1-6.0	2.4	1.0	0.5-1.9
Esophagus	1976-1980	13.8	4.5	3.5-5.7	10.2	8.4	6.2-11.1	3.6	1.9	1.1-3.1	14.0	4.6	3.6-5.9	9.8	8.3	6.1-11.1	4.2	2.4	1.4-3.7
	1981-1985	16.2	5.5	4.3-6.8	11.4	9.6	7.2-12.5	4.8	2.7	1.7-4.1	14.4	4.8	3.8-6.1	9.2	7.9	5.7-10.7	5.2	2.8	1.8-4.2
	1986-1990	17.8	5.7	4.5-7.0	12.2	9.5	7.2-12.3	5.6	2.8	1.9-4.1	14.6	4.6	3.6-5.9	9.6	7.9	5.8-10.5	5.0	2.4	1.6-3.6
	1991-1995	18.2	5.6	4.5-6.9	13.0	9.8	7.5-12.5	5.2	2.7	1.7-4.0	15.0	4.6	3.6-5.8	11.6	8.6	6.5-11.2	3.4	1.7	1.0-2.8
	1996-2000	18.8	5.9	4.7-7.2	13.6	10.3	8.0-13.1	5.2	2.7	1.7-4.0	16.4	5.1	4.0-6.3	12.8	9.9	7.6-12.7	3.6	1.8	1.0-2.9
	2001-2005	19.6	5.9	4.8-7.2	13.0	9.2	7.1-11.8	6.6	3.3	2.3-4.7	16.0	4.8	3.8-6.0	11.4	8.4	6.4-10.9	4.6	2.3	1.4-3.5
	2006-2010	21.8	6.3	5.1-7.6	17.6	11.5	9.2-14.3	4.2	2.1	1.3-3.2	17.2	4.8	3.9-6.0	12.8	8.4	6.5-10.8	4.4	2.1	1.3-3.2
	2011-2015	18.4	4.8	3.9-6.0	13.8	8.3	6.4-10.6	4.6	2.1	1.3-3.2	17.8	4.8	3.8-5.9	14.4	9.2	7.2-11.7	3.4	1.5	0.9-2.5
	2016-2020	22.6	5.6	4.6-6.7	16.4	9.3	7.3-11.6	6.2	2.7	1.8-3.9	16.0	3.9	3.1-4.9	12.8	7.2	5.5-9.3	3.2	1.4	0.8-2.3
Stomach	1976-1980	30.0	10.7	9.0-12.6	15.6	14.0	11.0-17.6	14.4	8.7	6.7-11.0	24.2	8.7	7.2-10.4	13.6	12.5	9.6-16.0	10.6	6.3	4.7-8.3
	1981-1985	33.2	10.9	9.2-12.7	19.8	16.4	13.2-20.1	13.4	7.0	5.4-8.9	21.0	6.8	5.6-8.3	12.2	10.0	7.6-12.9	8.8	4.6	3.3-6.2
	1986-1990	27.6	8.9	7.4-10.5	16.2	14.1	11.1-17.7	11.4	5.9	4.5-7.8	21.6	6.9	5.6-8.3	11.2	9.7	7.2-12.7	10.4	5.2	3.8-6.9
	1991-1995	26.8	8.1	6.8-9.6	15.2	12.2	9.5-15.3	11.6	5.7	4.3-7.4	18.6	5.6	4.5-6.9	9.8	8.1	5.9-10.8	8.8	4.3	3.1-5.9
	1996-2000	27.0	8.1	6.8-9.6	17.0	13.5	10.7-16.7	10.0	4.8	3.6-6.5	18.6	5.5	4.5-6.8	12.2	9.6	7.3-12.4	6.4	3.0	2.0-4.4
	2001-2005	23.2	6.9	5.7-8.3	14.4	10.6	8.3-13.3	8.8	4.4	3.1-5.9	11.6	3.4	2.5-4.4	7.0	5.2	3.6-7.3	4.6	2.2	1.4-3.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Albany**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	22.4	6.6	5.4-7.9	14.2	9.8	7.7-12.5	8.2	4.0	2.8-5.5	10.2	3.0	2.2-3.9	6.4	4.5	3.1-6.4	3.8	1.8	1.1-2.9
	2011-2015	27.0	7.8	6.5-9.3	17.4	11.7	9.3-14.5	9.6	5.0	3.7-6.7	11.6	3.3	2.5-4.3	6.4	4.2	2.8-6.0	5.2	2.5	1.6-3.8
	2016-2020	28.0	7.2	6.0-8.6	17.8	10.1	8.1-12.6	10.2	5.0	3.7-6.7	10.2	2.7	2.0-3.6	5.8	3.7	2.4-5.3	4.4	2.1	1.3-3.2
Colorectal	1976-1980	214.8	74.7	70.2-79.4	101.8	90.0	82.0-98.6	113.0	65.1	59.8-70.8	104.6	37.1	34.0-40.5	47.2	44.0	38.3-50.3	57.4	33.1	29.3-37.2
	1981-1985	234.6	77.5	73.1-82.2	118.8	102.4	94.0-111.3	115.8	62.1	57.1-67.5	100.6	33.5	30.6-36.6	53.2	47.1	41.4-53.4	47.4	25.3	22.1-28.8
	1986-1990	213.2	68.1	64.1-72.4	98.2	84.0	76.5-92.1	115.0	59.3	54.4-64.5	89.4	28.3	25.7-31.1	39.2	34.2	29.4-39.5	50.2	25.2	22.1-28.7
	1991-1995	193.8	59.5	55.8-63.4	89.6	70.7	64.2-77.7	104.2	51.7	47.3-56.5	86.6	26.4	24.0-29.1	37.0	30.2	26.0-35.0	49.6	23.7	20.7-26.9
	1996-2000	188.0	57.3	53.6-61.1	90.4	69.6	63.2-76.3	97.6	49.9	45.5-54.7	81.6	24.2	21.9-26.6	38.0	29.9	25.8-34.5	43.6	20.6	17.9-23.6
	2001-2005	191.6	56.6	53.0-60.3	88.4	64.4	58.5-70.7	103.2	50.8	46.4-55.5	64.2	18.5	16.5-20.7	30.6	22.8	19.3-26.8	33.6	15.3	13.1-18.0
	2006-2010	157.0	44.8	41.7-48.1	80.0	54.0	48.8-59.7	77.0	37.9	34.1-42.0	53.0	14.7	13.0-16.6	24.4	17.2	14.2-20.6	28.6	12.7	10.7-15.2
	2011-2015	146.6	40.4	37.4-43.5	75.4	46.8	42.1-52.0	71.2	34.8	31.1-38.8	52.2	13.8	12.1-15.6	22.2	14.4	11.8-17.4	30.0	13.1	11.0-15.5
2016-2020	140.0	37.0	34.2-39.9	69.4	42.1	37.7-47.0	70.6	32.8	29.3-36.6	42.0	10.5	9.1-12.1	19.4	11.6	9.4-14.3	22.6	9.6	7.9-11.7	
Colon excluding rectum	1976-1980	153.8	53.6	49.8-57.6	68.8	61.2	54.6-68.4	85.0	48.9	44.3-53.9	88.6	31.3	28.4-34.5	40.8	37.8	32.5-43.6	47.8	27.6	24.2-31.5
	1981-1985	170.2	56.7	52.9-60.7	81.0	71.4	64.4-79.0	89.2	48.0	43.6-52.8	86.4	28.9	26.2-31.8	44.8	40.2	34.9-46.0	41.6	22.3	19.3-25.6
	1986-1990	148.4	47.3	44.0-50.9	63.0	54.1	48.1-60.6	85.4	43.4	39.3-47.9	71.4	22.6	20.3-25.1	30.6	26.5	22.3-31.2	40.8	20.4	17.6-23.5
	1991-1995	145.6	44.4	41.2-47.8	66.2	52.3	46.7-58.4	79.4	39.1	35.2-43.2	69.8	21.1	18.9-23.4	29.6	24.2	20.4-28.5	40.2	18.7	16.1-21.5
	1996-2000	138.6	41.9	38.9-45.2	65.6	50.7	45.3-56.5	73.0	36.8	33.1-40.9	65.0	19.1	17.1-21.3	30.0	23.7	20.0-27.8	35.0	16.2	13.8-18.9
	2001-2005	140.8	41.4	38.3-44.6	60.4	44.3	39.4-49.6	80.4	39.5	35.6-43.6	52.4	15.0	13.3-17.0	24.0	18.0	14.9-21.6	28.4	12.9	10.8-15.3
	2006-2010	111.8	31.8	29.2-34.6	53.2	36.3	32.0-41.0	58.6	28.2	25.0-31.8	41.0	11.3	9.8-13.0	18.4	12.9	10.4-15.9	22.6	9.9	8.1-12.1
	2011-2015	105.4	29.1	26.6-31.8	51.0	32.7	28.7-37.1	54.4	26.1	22.9-29.6	42.2	11.2	9.7-12.9	17.0	11.1	8.8-13.8	25.2	11.3	9.3-13.6
2016-2020	102.2	26.3	24.0-28.8	48.0	28.7	25.1-32.7	54.2	24.6	21.6-27.9	32.2	8.1	6.9-9.5	14.4	8.8	6.9-11.2	17.8	7.6	6.0-9.5	
Rectum & rectosigmoid	1976-1980	61.0	21.0	18.7-23.6	33.0	28.8	24.4-33.8	28.0	16.2	13.6-19.2	16.0	5.8	4.6-7.2	6.4	6.3	4.2-8.9	9.6	5.4	4.0-7.3
	1981-1985	64.4	20.8	18.6-23.3	37.8	30.9	26.6-35.9	26.6	14.1	11.8-16.8	14.2	4.6	3.6-5.8	8.4	7.0	4.9-9.5	5.8	3.0	2.0-4.4
	1986-1990	64.8	20.8	18.6-23.2	35.2	29.9	25.5-34.8	29.6	15.8	13.3-18.7	18.0	5.7	4.6-7.1	8.6	7.7	5.5-10.4	9.4	4.9	3.5-6.5
	1991-1995	48.2	15.1	13.3-17.2	23.4	18.4	15.2-22.1	24.8	12.7	10.5-15.2	16.8	5.4	4.3-6.7	7.4	6.0	4.2-8.3	9.4	5.0	3.6-6.7
	1996-2000	49.4	15.3	13.5-17.4	24.8	18.9	15.7-22.5	24.6	13.1	10.9-15.7	16.6	5.1	4.0-6.3	8.0	6.3	4.5-8.5	8.6	4.4	3.1-5.9
	2001-2005	50.8	15.2	13.4-17.2	28.0	20.0	16.8-23.7	22.8	11.3	9.3-13.7	11.8	3.5	2.6-4.5	6.6	4.8	3.3-6.8	5.2	2.5	1.6-3.7
	2006-2010	45.2	13.0	11.3-14.8	26.8	17.7	14.8-21.0	18.4	9.6	7.7-11.9	12.0	3.4	2.6-4.4	6.0	4.2	2.9-6.1	6.0	2.8	1.9-4.1
	2011-2015	41.2	11.2	9.7-12.9	24.4	14.1	11.7-17.0	16.8	8.7	6.9-10.9	10.0	2.6	1.9-3.4	5.2	3.3	2.1-5.0	4.8	1.8	1.1-2.8
2016-2020	37.8	10.6	9.1-12.3	21.4	13.4	10.9-16.3	16.4	8.2	6.4-10.3	9.8	2.4	1.7-3.2	5.0	2.8	1.8-4.2	4.8	2.0	1.3-3.1	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Albany**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	9.6	3.3	2.4-4.4	6.2	5.2	3.4-7.4	3.4	1.9	1.1-3.1	8.8	2.9	2.1-4.0	5.0	4.2	2.7-6.4	3.8	2.1	1.3-3.4
	1981-1985	10.2	3.4	2.5-4.5	7.0	6.2	4.3-8.7	3.2	1.7	1.0-2.8	7.6	2.4	1.7-3.4	4.6	3.7	2.3-5.7	3.0	1.6	0.9-2.8
	1986-1990	10.8	3.6	2.7-4.7	7.4	5.8	4.1-8.1	3.4	2.0	1.1-3.2	8.2	2.6	1.9-3.6	5.4	4.3	2.8-6.3	2.8	1.3	0.7-2.3
	1991-1995	11.8	3.8	2.8-4.9	9.2	7.5	5.4-10.0	2.6	1.3	0.7-2.3	11.2	3.5	2.6-4.5	8.4	7.2	5.1-9.7	2.8	1.3	0.7-2.3
	1996-2000	18.0	5.6	4.5-6.9	14.2	10.5	8.2-13.2	3.8	1.9	1.1-3.1	13.6	4.3	3.3-5.4	11.2	8.5	6.4-11.0	2.4	1.2	0.6-2.2
	2001-2005	20.8	6.3	5.1-7.6	14.0	10.1	7.9-12.8	6.8	3.3	2.3-4.7	19.0	5.6	4.5-6.8	13.2	9.6	7.4-12.3	5.8	2.6	1.8-3.9
	2006-2010	27.4	7.9	6.6-9.4	18.2	11.8	9.5-14.6	9.2	4.8	3.5-6.4	17.2	4.9	3.9-6.1	9.6	6.3	4.6-8.5	7.6	3.7	2.6-5.2
	2011-2015	37.2	9.8	8.4-11.3	28.0	15.7	13.2-18.7	9.2	4.9	3.6-6.6	24.0	6.3	5.2-7.6	16.8	9.8	7.8-12.3	7.2	3.4	2.3-4.8
	2016-2020	37.8	9.2	7.9-10.7	27.8	15.3	12.8-18.2	10.0	4.4	3.2-5.9	25.8	6.4	5.3-7.7	18.4	10.0	8.0-12.4	7.4	3.5	2.4-5.0
Pancreas	1976-1980	35.6	12.4	10.6-14.4	19.6	16.7	13.5-20.5	16.0	9.5	7.5-11.9	35.4	12.4	10.6-14.4	18.6	16.1	12.8-19.9	16.8	10.0	7.9-12.4
	1981-1985	31.8	10.4	8.8-12.2	15.8	14.0	11.0-17.6	16.0	8.4	6.7-10.6	32.4	10.6	9.0-12.3	15.8	13.6	10.7-17.1	16.6	8.8	7.0-11.0
	1986-1990	38.2	12.2	10.5-14.1	19.0	16.5	13.2-20.3	19.2	9.8	7.9-12.0	36.4	11.4	9.8-13.2	17.8	14.9	11.8-18.4	18.6	9.3	7.5-11.5
	1991-1995	33.6	10.4	8.9-12.1	15.2	12.0	9.5-15.1	18.4	9.1	7.3-11.2	31.4	9.6	8.2-11.3	12.6	10.0	7.6-12.8	18.8	9.3	7.5-11.4
	1996-2000	34.2	10.5	9.0-12.2	17.0	12.9	10.3-16.0	17.2	8.8	7.0-11.0	37.0	11.2	9.6-12.9	16.8	12.9	10.3-16.0	20.2	10.0	8.1-12.2
	2001-2005	41.2	12.2	10.6-14.0	19.6	14.0	11.4-17.1	21.6	10.7	8.7-13.0	36.2	10.6	9.1-12.3	17.4	12.6	10.1-15.6	18.8	8.9	7.1-11.0
	2006-2010	46.4	13.3	11.6-15.1	24.8	17.2	14.3-20.6	21.6	10.4	8.5-12.6	38.8	11.1	9.5-12.8	18.6	13.0	10.5-16.0	20.2	9.7	7.8-11.8
	2011-2015	52.8	14.6	12.8-16.5	27.6	17.5	14.6-20.8	25.2	12.2	10.1-14.7	38.2	10.4	9.0-12.1	19.6	12.4	10.0-15.2	18.6	8.7	7.0-10.8
	2016-2020	64.0	15.9	14.2-17.9	29.8	17.4	14.6-20.5	34.2	14.8	12.6-17.3	46.4	11.5	10.1-13.2	21.0	12.5	10.1-15.2	25.4	10.8	9.0-13.0
Larynx	1976-1980	20.4	6.8	5.5-8.3	17.2	13.8	10.9-17.2	3.2	1.8	1.0-3.0	6.2	2.0	1.4-2.9	5.0	4.1	2.5-6.2	1.2	0.7	0.2-1.5
	1981-1985	19.2	6.3	5.1-7.8	14.6	11.5	9.0-14.6	4.6	2.9	1.8-4.4	6.4	2.1	1.4-3.0	5.4	4.3	2.7-6.3	1.0	0.8	0.2-1.8
	1986-1990	16.4	5.2	4.1-6.5	11.2	8.4	6.3-11.0	5.2	2.9	1.9-4.3	6.6	2.1	1.4-2.9	4.6	3.6	2.2-5.4	2.0	1.1	0.5-2.1
	1991-1995	18.6	6.0	4.9-7.4	14.2	10.8	8.4-13.7	4.4	2.4	1.5-3.7	5.6	1.8	1.2-2.6	4.0	3.0	1.8-4.7	1.6	0.9	0.4-1.7
	1996-2000	16.6	5.3	4.2-6.6	13.6	10.1	7.8-12.8	3.0	1.7	0.9-2.8	7.0	2.1	1.5-3.0	5.4	4.3	2.8-6.2	1.6	0.8	0.3-1.7
	2001-2005	14.4	4.4	3.4-5.5	11.8	8.1	6.2-10.5	2.6	1.4	0.7-2.4	4.8	1.5	0.9-2.2	4.6	3.4	2.1-5.0	0.2	0.1	0.0-0.7
	2006-2010	13.0	3.9	3.0-5.0	9.6	6.5	4.7-8.6	3.4	1.9	1.1-3.0	4.6	1.3	0.8-2.0	2.8	1.9	1.0-3.2	1.8	0.9	0.4-1.7
	2011-2015	14.8	3.8	3.0-4.9	10.0	6.0	4.4-8.0	4.8	2.4	1.5-3.6	4.6	1.2	0.8-1.9	3.4	2.0	1.2-3.3	1.2	0.7	0.2-1.5
	2016-2020	12.8	3.3	2.5-4.2	9.4	5.3	3.8-7.1	3.4	1.6	0.9-2.6	3.0	0.7	0.4-1.2	1.8	0.9	0.4-1.9	1.2	0.5	0.2-1.1
Lung and bronchus	1976-1980	180.8	61.7	57.6-65.9	131.4	110.2	101.6-119.3	49.4	29.6	25.9-33.6	144.8	49.0	45.4-52.8	105.4	89.1	81.4-97.4	39.4	23.1	19.9-26.7
	1981-1985	210.0	69.3	65.1-73.7	141.4	113.6	105.1-122.6	68.6	40.9	36.5-45.6	162.4	53.2	49.5-57.1	114.0	93.6	85.8-101.9	48.4	27.5	24.1-31.4
	1986-1990	229.8	74.9	70.5-79.4	140.6	113.5	105.0-122.4	89.2	50.2	45.5-55.3	191.8	62.1	58.2-66.3	112.8	91.7	84.0-99.8	79.0	44.1	39.7-48.8
	1991-1995	241.2	75.2	71.0-79.6	133.4	102.1	94.3-110.2	107.8	57.8	52.9-63.1	200.8	62.4	58.5-66.4	112.4	87.4	80.2-95.0	88.4	46.3	42.0-51.0
	1996-2000	252.4	78.2	73.9-82.7	135.0	101.6	94.0-109.6	117.4	63.0	57.9-68.4	190.6	58.5	54.9-62.4	102.2	77.5	70.9-84.6	88.4	46.3	42.0-50.9
	2001-2005	246.6	75.1	70.9-79.4	120.6	87.2	80.3-94.4	126.0	67.5	62.2-73.1	185.2	55.8	52.2-59.5	94.2	68.5	62.4-75.0	91.0	47.3	43.0-51.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Albany**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	260.8	76.0	71.9-80.3	132.6	90.4	83.6-97.7	128.2	66.5	61.3-72.0	186.2	53.6	50.1-57.2	92.0	63.5	57.7-69.7	94.2	46.8	42.5-51.3
	2011-2015	263.4	72.5	68.6-76.6	132.4	84.1	77.7-91.0	131.0	64.0	59.1-69.3	173.4	48.0	44.8-51.4	92.8	60.3	54.8-66.2	80.6	38.9	35.1-43.1
	2016-2020	277.2	69.2	65.5-73.1	130.8	74.5	68.7-80.6	146.4	66.0	61.2-71.1	149.0	37.1	34.4-39.9	75.6	43.4	39.0-48.2	73.4	32.6	29.3-36.3
Melanoma of the skin	1976-1980	20.2	7.4	6.0-9.0	12.0	10.0	7.5-13.0	8.2	5.2	3.7-7.1	5.6	2.1	1.4-3.1	3.0	2.6	1.4-4.4	2.6	1.6	0.8-2.8
	1981-1985	25.2	9.3	7.7-11.1	13.0	10.8	8.3-13.9	12.2	8.1	6.1-10.6	7.0	2.4	1.7-3.4	3.8	3.0	1.7-4.7	3.2	2.1	1.1-3.4
	1986-1990	27.2	9.1	7.6-10.8	16.0	12.3	9.7-15.4	11.2	7.3	5.4-9.5	8.8	3.0	2.2-4.1	5.8	5.1	3.4-7.3	3.0	1.7	0.9-2.9
	1991-1995	22.6	7.3	6.0-8.8	14.4	11.1	8.6-14.0	8.2	4.6	3.3-6.4	8.6	2.7	1.9-3.6	5.6	4.2	2.8-6.1	3.0	1.6	0.9-2.7
	1996-2000	29.0	9.4	7.9-11.1	17.8	13.0	10.5-16.1	11.2	6.5	4.9-8.5	9.4	2.9	2.1-3.9	6.2	4.7	3.2-6.6	3.2	1.6	0.9-2.6
	2001-2005	40.8	12.7	11.0-14.6	20.2	14.1	11.5-17.2	20.6	11.7	9.5-14.3	9.2	2.8	2.0-3.7	5.8	4.0	2.7-5.8	3.4	1.8	1.0-2.9
	2006-2010	52.8	15.6	13.7-17.6	31.6	21.2	17.9-24.8	21.2	11.6	9.5-14.2	10.4	3.0	2.2-4.0	5.6	3.8	2.5-5.5	4.8	2.5	1.6-3.8
	2011-2015	77.6	22.1	19.9-24.5	44.6	28.6	24.9-32.8	33.0	17.6	14.9-20.7	11.6	3.3	2.5-4.3	6.4	4.2	2.8-5.9	5.2	2.6	1.6-3.8
	2016-2020	87.0	23.4	21.2-25.9	50.0	30.0	26.3-34.1	37.0	19.1	16.3-22.3	9.0	2.2	1.6-2.9	5.6	3.2	2.1-4.7	3.4	1.5	0.8-2.4
Female breast	1976-1980							162.2	104.0	96.6-111.7							56.6	34.1	30.2-38.5
	1981-1985							191.8	116.5	108.9-124.5							66.8	39.1	34.8-43.7
	1986-1990							217.0	129.6	121.7-137.9							76.0	42.0	37.7-46.6
	1991-1995							197.8	116.4	109.0-124.1							57.2	31.9	28.2-36.0
	1996-2000							239.4	138.4	130.5-146.6							53.8	28.5	25.1-32.2
	2001-2005							230.8	129.3	121.9-137.2							48.2	24.5	21.5-28.0
	2006-2010							244.0	131.2	123.8-139.0							42.0	20.7	17.9-23.9
	2011-2015							284.0	151.0	143.0-159.5							43.0	21.1	18.3-24.3
	2016-2020							285.2	146.0	138.1-154.2							43.0	19.8	17.1-22.8
Cervix uteri	1976-1980							15.4	10.5	8.1-13.2							6.0	3.6	2.4-5.3
	1981-1985							14.8	10.0	7.7-12.6							7.4	4.6	3.2-6.4
	1986-1990							12.4	8.7	6.6-11.2							6.4	4.2	2.8-6.0
	1991-1995							11.8	7.6	5.7-9.8							3.4	2.1	1.2-3.3
	1996-2000							12.2	7.5	5.7-9.7							4.6	2.7	1.7-4.0
	2001-2005							10.8	6.7	5.0-8.8							4.0	2.2	1.4-3.5
	2006-2010							8.0	5.3	3.8-7.3							2.2	1.1	0.6-2.1
	2011-2015							10.0	5.6	4.1-7.5							2.2	1.1	0.5-2.1
	2016-2020							9.4	6.3	4.5-8.4							3.2	1.8	1.0-3.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Albany**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							32.8	20.8	17.6-24.4							6.8	3.8	2.7-5.4
	1981-1985							35.2	20.7	17.6-24.1							7.8	4.3	3.0-6.0
	1986-1990							38.6	22.1	19.0-25.6							6.8	3.7	2.5-5.2
	1991-1995							35.0	19.7	16.8-23.0							7.8	3.9	2.7-5.4
	1996-2000							41.0	23.5	20.4-27.0							8.4	4.1	2.9-5.6
	2001-2005							41.2	22.8	19.7-26.1							6.2	3.2	2.2-4.6
	2006-2010							48.6	25.6	22.4-29.1							7.6	3.8	2.6-5.3
	2011-2015							62.6	30.5	27.1-34.2							12.6	5.9	4.5-7.6
	2016-2020							66.8	31.6	28.2-35.4							11.2	5.1	3.8-6.8
Ovary	1976-1980							24.4	15.7	12.9-18.9							16.4	10.2	8.0-12.7
	1981-1985							24.6	14.8	12.2-17.8							18.6	10.8	8.6-13.3
	1986-1990							24.6	14.7	12.1-17.6							17.4	10.1	8.0-12.6
	1991-1995							30.8	17.7	15.0-20.9							19.8	10.4	8.4-12.8
	1996-2000							25.8	14.9	12.4-17.7							14.6	7.5	5.9-9.5
	2001-2005							21.4	11.6	9.5-14.1							18.6	10.0	8.1-12.4
	2006-2010							20.4	10.6	8.6-12.9							14.4	7.4	5.8-9.4
	2011-2015							22.8	11.3	9.2-13.7							14.0	6.6	5.1-8.4
	2016-2020							19.0	10.0	8.0-12.4							14.0	6.3	4.8-8.1
Prostate	1976-1980				78.4	74.4	66.9-82.5							31.0	30.7	25.9-36.2			
	1981-1985				102.4	92.7	84.5-101.4							35.4	36.4	31.0-42.3			
	1986-1990				124.8	104.9	96.6-113.8							37.8	35.9	30.8-41.5			
	1991-1995				196.0	150.5	141.1-160.4							43.6	37.8	32.8-43.3			
	1996-2000				222.2	166.5	156.8-176.7							38.0	31.0	26.7-35.8			
	2001-2005				241.2	171.3	161.7-181.3							30.2	23.8	20.1-27.9			
	2006-2010				248.6	161.3	152.3-170.7							27.0	19.8	16.6-23.5			
	2011-2015				220.8	126.4	118.8-134.4							26.8	18.1	15.1-21.5			
	2016-2020				247.0	130.7	123.3-138.5							29.0	18.1	15.2-21.3			
Testis	1976-1980				6.2	4.4	2.9-6.3							0.8	0.6	0.1-1.6			
	1981-1985				7.8	5.0	3.5-6.9							0.6	0.3	0.1-1.1			
	1986-1990				5.2	3.5	2.2-5.2							0.0	0.0	0.0-0.6			
	1991-1995				7.4	4.8	3.4-6.7							0.4	0.4	0.0-1.3			
	1996-2000				6.8	4.5	3.1-6.4							0.2	0.2	0.0-0.8			
	2001-2005				6.6	4.5	3.1-6.4							0.0	0.0	0.0-0.5			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Albany**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				10.0	6.9	5.1-9.2							0.2	0.1	0.0-0.8			
	2011-2015				9.2	6.8	4.9-9.1							0.2	0.2	0.0-0.9			
	2016-2020				10.4	7.1	5.3-9.4							0.8	0.5	0.1-1.4			
Urinary bladder (incl. in situ)	1976-1980	65.2	22.7	20.3-25.4	43.8	38.4	33.3-44.1	21.4	12.5	10.2-15.1	15.8	5.6	4.4-7.0	9.4	8.4	6.1-11.3	6.4	3.7	2.5-5.3
	1981-1985	74.2	24.7	22.2-27.4	49.4	41.0	35.9-46.6	24.8	13.8	11.4-16.5	14.6	5.0	3.9-6.3	8.6	7.8	5.5-10.5	6.0	3.1	2.1-4.5
	1986-1990	70.4	22.7	20.4-25.3	48.2	41.2	36.0-46.9	22.2	11.6	9.5-14.1	19.4	6.4	5.1-7.8	13.4	12.7	9.8-16.2	6.0	3.1	2.1-4.5
	1991-1995	72.0	22.1	19.8-24.5	51.0	40.2	35.4-45.6	21.0	10.6	8.6-13.0	16.6	4.9	3.9-6.1	10.0	8.5	6.2-11.2	6.6	3.0	2.0-4.2
	1996-2000	87.2	26.5	24.0-29.1	58.0	44.6	39.6-50.1	29.2	14.9	12.6-17.7	20.8	6.0	4.9-7.3	13.6	11.0	8.5-14.0	7.2	3.0	2.1-4.3
	2001-2005	107.0	31.9	29.2-34.7	79.2	58.3	52.6-64.3	27.8	14.0	11.7-16.6	13.6	3.9	3.0-5.0	9.6	7.1	5.3-9.5	4.0	1.8	1.1-2.9
	2006-2010	98.4	28.2	25.7-30.8	72.8	50.6	45.4-56.1	25.6	12.3	10.2-14.8	14.2	3.9	3.1-5.0	10.2	7.3	5.4-9.6	4.0	1.6	1.0-2.6
	2011-2015	100.6	27.4	25.0-30.0	73.8	47.5	42.7-52.8	26.8	12.8	10.6-15.3	16.4	4.4	3.5-5.6	12.2	8.5	6.4-10.9	4.2	2.0	1.2-3.1
2016-2020	99.8	25.4	23.1-27.8	75.6	44.9	40.4-49.8	24.2	10.8	8.9-13.0	16.0	3.9	3.1-4.9	11.8	7.1	5.4-9.2	4.2	1.7	1.0-2.7	
Kidney and renal pelvis	1976-1980	19.6	6.8	5.5-8.3	11.8	10.2	7.6-13.3	7.8	4.6	3.2-6.4	9.4	3.2	2.4-4.4	6.0	5.4	3.5-7.7	3.4	1.8	1.0-2.9
	1981-1985	24.8	8.5	7.1-10.2	15.6	13.0	10.2-16.3	9.2	5.3	3.8-7.1	11.2	3.5	2.7-4.6	5.8	4.8	3.1-6.9	5.4	2.8	1.9-4.2
	1986-1990	29.6	9.6	8.1-11.4	19.2	15.5	12.5-19.1	10.4	5.6	4.1-7.4	11.4	3.8	2.9-5.0	7.6	6.9	4.8-9.5	3.8	2.0	1.2-3.2
	1991-1995	39.8	12.6	10.9-14.5	21.2	16.1	13.1-19.5	18.6	9.9	8.0-12.2	13.4	4.1	3.2-5.3	6.6	5.2	3.5-7.3	6.8	3.3	2.3-4.7
	1996-2000	43.2	13.7	11.9-15.6	28.0	20.9	17.6-24.7	15.2	8.3	6.5-10.5	14.4	4.5	3.5-5.7	10.0	7.8	5.8-10.3	4.4	2.3	1.4-3.5
	2001-2005	39.4	12.1	10.5-14.0	23.0	16.5	13.6-19.8	16.4	9.2	7.3-11.5	12.2	3.7	2.8-4.7	7.6	5.6	3.9-7.6	4.6	2.4	1.5-3.7
	2006-2010	58.4	17.1	15.2-19.3	37.8	24.5	21.1-28.3	20.6	11.5	9.4-14.1	13.0	3.8	2.9-4.9	8.0	5.6	4.0-7.7	5.0	2.4	1.5-3.6
	2011-2015	64.8	18.2	16.3-20.4	40.0	25.1	21.7-29.0	24.8	12.7	10.5-15.2	15.4	4.2	3.3-5.3	9.2	6.0	4.3-8.0	6.2	3.0	2.0-4.4
2016-2020	76.0	20.3	18.2-22.5	49.6	28.7	25.1-32.7	26.4	13.1	10.8-15.7	12.2	3.1	2.4-4.1	8.2	5.3	3.7-7.2	4.0	1.6	1.0-2.5	
Brain and other central nervous system	1976-1980	17.0	5.8	4.6-7.2	9.2	6.8	4.9-9.1	7.8	5.0	3.5-6.9	10.0	3.3	2.4-4.4	4.8	3.3	2.1-5.0	5.2	3.3	2.1-5.0
	1981-1985	21.8	7.5	6.1-9.1	11.0	8.9	6.6-11.6	10.8	6.8	5.1-9.0	12.2	4.0	3.0-5.1	6.6	5.3	3.6-7.5	5.6	3.2	2.1-4.7
	1986-1990	20.0	7.1	5.7-8.7	12.6	9.8	7.5-12.6	7.4	4.9	3.4-6.8	10.8	3.6	2.7-4.7	6.8	5.4	3.7-7.6	4.0	2.2	1.3-3.4
	1991-1995	22.2	7.3	6.0-8.8	11.6	8.7	6.5-11.3	10.6	6.3	4.7-8.3	13.2	4.3	3.3-5.5	6.2	4.8	3.3-6.9	7.0	3.9	2.7-5.5
	1996-2000	21.2	6.7	5.5-8.1	9.4	7.0	5.2-9.4	11.8	6.4	4.8-8.3	16.4	5.2	4.1-6.4	8.2	6.0	4.3-8.2	8.2	4.2	3.0-5.7
	2001-2005	23.4	7.4	6.1-8.9	14.0	9.9	7.7-12.6	9.4	5.7	4.2-7.6	15.6	4.8	3.8-6.0	8.8	6.2	4.5-8.3	6.8	3.9	2.7-5.5
	2006-2010	23.2	7.1	5.9-8.6	12.8	8.8	6.8-11.3	10.4	5.6	4.2-7.5	16.4	4.7	3.7-5.9	10.8	7.0	5.3-9.2	5.6	2.8	1.8-4.1
	2011-2015	22.0	6.7	5.4-8.1	10.8	6.9	5.1-9.0	11.2	6.6	4.9-8.7	13.8	3.8	2.9-4.8	7.6	4.7	3.3-6.5	6.2	3.0	2.0-4.3
2016-2020	21.8	5.8	4.8-7.1	12.0	6.8	5.2-8.9	9.8	5.1	3.7-6.9	15.4	4.1	3.2-5.2	8.4	4.9	3.5-6.7	7.0	3.4	2.4-4.9	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Albany**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	8.4	3.2	2.3-4.4	4.0	3.7	2.2-5.8	4.4	3.0	1.8-4.6	1.2	0.4	0.1-0.9	0.8	0.6	0.2-1.7	0.4	0.2	0.0-0.9
	1981-1985	11.2	3.9	2.9-5.1	2.6	2.1	1.1-3.7	8.6	5.4	3.8-7.3	2.4	0.9	0.4-1.5	0.8	0.7	0.2-1.9	1.6	0.8	0.4-1.7
	1986-1990	13.0	4.6	3.5-5.9	4.2	3.4	2.1-5.2	8.8	5.7	4.1-7.7	1.6	0.5	0.2-1.0	0.8	0.6	0.2-1.5	0.8	0.4	0.1-1.1
	1991-1995	15.2	4.9	3.9-6.2	4.0	2.9	1.8-4.6	11.2	6.9	5.2-9.0	1.2	0.4	0.1-0.8	0.6	0.4	0.1-1.2	0.6	0.4	0.1-1.1
	1996-2000	24.6	8.2	6.8-9.7	5.8	4.2	2.8-6.1	18.8	11.9	9.6-14.5	2.2	0.6	0.3-1.2	0.6	0.5	0.1-1.4	1.6	0.7	0.3-1.5
	2001-2005	28.0	9.3	7.8-11.0	6.2	4.2	2.9-6.0	21.8	14.0	11.5-17.0	1.6	0.4	0.2-0.9	0.2	0.2	0.0-0.8	1.4	0.6	0.2-1.3
	2006-2010	46.6	14.7	12.8-16.8	10.8	7.2	5.4-9.4	35.8	21.7	18.6-25.2	1.6	0.5	0.2-1.0	1.2	0.9	0.3-1.9	0.4	0.2	0.0-0.9
	2011-2015	61.0	19.1	17.0-21.5	13.6	8.8	6.8-11.3	47.4	29.2	25.4-33.3	1.8	0.5	0.2-1.0	0.8	0.5	0.1-1.3	1.0	0.5	0.2-1.3
2016-2020	53.0	16.3	14.3-18.5	11.6	6.6	4.9-8.6	41.4	25.5	21.9-29.4	1.8	0.5	0.2-0.9	1.0	0.6	0.2-1.4	0.8	0.3	0.1-1.0	
Hodgkin lymphoma	1976-1980	9.6	2.9	2.1-3.9	5.8	3.7	2.4-5.4	3.8	2.2	1.3-3.5	3.6	1.3	0.7-2.0	2.8	2.1	1.1-3.6	0.8	0.5	0.1-1.3
	1981-1985	10.4	3.2	2.4-4.3	5.8	3.9	2.6-5.7	4.6	2.7	1.7-4.1	2.2	0.7	0.4-1.3	0.6	0.5	0.1-1.4	1.6	0.9	0.4-1.9
	1986-1990	11.2	3.4	2.6-4.5	5.2	3.5	2.2-5.2	6.0	3.3	2.2-4.8	3.4	1.2	0.7-1.9	1.4	1.1	0.4-2.3	2.0	1.1	0.5-2.1
	1991-1995	7.2	2.3	1.6-3.1	3.6	2.5	1.5-4.0	3.6	2.2	1.3-3.4	2.2	0.7	0.3-1.2	0.8	0.6	0.2-1.5	1.4	0.8	0.3-1.7
	1996-2000	8.2	2.6	1.9-3.6	4.4	3.0	1.9-4.6	3.8	2.3	1.4-3.6	1.6	0.5	0.2-1.0	1.0	0.7	0.2-1.7	0.6	0.3	0.1-1.0
	2001-2005	14.2	4.7	3.7-5.9	9.6	6.6	4.9-8.8	4.6	3.0	1.9-4.5	2.2	0.7	0.3-1.3	1.2	0.8	0.3-1.8	1.0	0.6	0.2-1.4
	2006-2010	12.0	3.8	2.9-4.9	6.4	4.3	2.9-6.2	5.6	3.4	2.3-5.0	1.0	0.3	0.1-0.7	0.4	0.3	0.0-1.0	0.6	0.3	0.1-1.1
	2011-2015	8.6	2.7	1.9-3.7	5.4	3.7	2.4-5.4	3.2	1.8	1.0-3.0	1.2	0.3	0.1-0.8	0.4	0.2	0.0-1.0	0.8	0.4	0.1-1.1
2016-2020	8.6	2.9	2.1-4.0	4.6	3.2	2.0-4.9	4.0	2.6	1.6-4.2	0.4	0.1	0.0-0.4	0.2	0.1	0.0-0.8	0.2	0.1	0.0-0.6	
Non-Hodgkin lymphomas	1976-1980	36.4	12.9	11.1-15.0	18.2	15.2	12.1-18.7	18.2	11.1	8.9-13.7	19.0	6.6	5.3-8.1	10.2	8.4	6.2-11.2	8.8	5.3	3.8-7.2
	1981-1985	43.2	14.4	12.5-16.5	21.0	17.4	14.1-21.2	22.2	12.3	10.0-14.9	19.2	6.6	5.3-8.0	9.8	8.9	6.5-11.9	9.4	5.3	3.9-7.2
	1986-1990	49.4	16.0	14.0-18.2	27.0	21.5	17.9-25.6	22.4	12.0	9.8-14.5	26.4	8.4	7.0-10.0	12.6	10.2	7.7-13.1	13.8	7.1	5.5-9.1
	1991-1995	54.0	17.1	15.1-19.3	25.2	19.4	16.1-23.2	28.8	15.3	12.8-18.1	26.8	8.2	6.8-9.7	12.2	9.9	7.5-12.8	14.6	6.8	5.3-8.6
	1996-2000	60.8	19.2	17.1-21.5	32.6	24.2	20.6-28.2	28.2	15.2	12.8-18.1	31.0	9.2	7.8-10.8	15.0	11.4	9.0-14.4	16.0	7.8	6.2-9.8
	2001-2005	67.4	20.2	18.1-22.5	35.8	26.4	22.6-30.5	31.6	16.4	13.9-19.3	23.6	6.7	5.5-8.0	13.2	9.9	7.7-12.7	10.4	4.6	3.4-6.1
	2006-2010	85.0	25.0	22.7-27.6	43.8	29.5	25.7-33.8	41.2	21.4	18.5-24.6	21.2	5.9	4.8-7.1	10.8	7.6	5.7-9.9	10.4	4.7	3.5-6.3
	2011-2015	84.0	23.2	21.0-25.6	46.6	29.6	25.8-33.8	37.4	18.4	15.8-21.3	23.0	6.4	5.2-7.7	13.2	9.1	7.0-11.6	9.8	4.5	3.3-6.1
2016-2020	85.2	22.7	20.5-25.0	45.8	26.9	23.4-30.7	39.4	19.6	16.8-22.7	18.6	4.7	3.8-5.8	10.0	6.3	4.6-8.3	8.6	3.5	2.5-4.9	
Myeloma	1976-1980	13.6	4.6	3.6-5.9	5.6	4.7	3.1-6.9	8.0	4.6	3.3-6.3	7.4	2.6	1.8-3.6	3.8	3.2	1.9-5.1	3.6	2.0	1.2-3.2
	1981-1985	16.4	5.4	4.3-6.7	8.0	6.8	4.8-9.4	8.4	4.5	3.2-6.2	11.0	3.6	2.7-4.7	4.6	3.9	2.4-6.0	6.4	3.5	2.3-5.0
	1986-1990	17.4	5.7	4.6-7.1	7.4	6.2	4.4-8.6	10.0	5.7	4.2-7.6	11.6	3.7	2.8-4.7	5.8	5.1	3.3-7.3	5.8	3.0	2.0-4.3
	1991-1995	15.2	4.7	3.7-6.0	7.2	6.1	4.3-8.5	8.0	3.9	2.8-5.4	12.6	3.9	3.0-5.0	5.4	4.9	3.2-7.2	7.2	3.6	2.5-5.0
	1996-2000	18.2	5.5	4.5-6.8	9.8	7.5	5.5-9.9	8.4	4.2	3.0-5.8	9.2	2.7	2.0-3.6	5.2	4.2	2.7-6.2	4.0	1.9	1.1-3.0
	2001-2005	19.6	5.8	4.7-7.1	10.4	7.2	5.4-9.5	9.2	4.8	3.5-6.5	8.8	2.5	1.8-3.4	4.0	2.9	1.8-4.5	4.8	2.2	1.4-3.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Albany**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	22.0	6.3	5.1-7.6	12.8	8.7	6.7-11.2	9.2	4.5	3.3-6.1	11.6	3.3	2.5-4.3	6.0	4.2	2.8-6.1	5.6	2.7	1.8-4.0
	2011-2015	34.6	9.3	7.9-10.8	17.6	10.7	8.5-13.3	17.0	8.3	6.6-10.3	12.4	3.3	2.5-4.2	6.4	4.5	3.0-6.4	6.0	2.7	1.8-3.9
	2016-2020	33.0	8.3	7.1-9.8	17.4	10.5	8.4-13.1	15.6	6.7	5.3-8.5	11.0	2.7	2.0-3.6	4.8	2.9	1.8-4.4	6.2	2.4	1.6-3.6
Leukemias	1976-1980	38.8	13.9	11.9-16.0	22.0	19.7	16.0-24.0	16.8	10.3	8.1-12.8	23.2	8.1	6.6-9.7	11.4	10.1	7.5-13.2	11.8	6.8	5.2-8.9
	1981-1985	39.6	13.6	11.7-15.7	23.2	20.1	16.5-24.3	16.4	9.8	7.7-12.2	25.8	9.0	7.5-10.7	13.6	11.9	9.2-15.3	12.2	7.2	5.5-9.4
	1986-1990	37.4	12.2	10.5-14.1	21.0	17.8	14.4-21.6	16.4	9.1	7.2-11.4	25.0	8.2	6.8-9.8	13.6	11.5	8.8-14.7	11.4	6.6	4.9-8.6
	1991-1995	33.8	10.7	9.2-12.5	18.4	14.1	11.3-17.3	15.4	8.4	6.6-10.6	20.0	6.2	5.1-7.6	11.8	9.3	7.0-12.0	8.2	4.2	3.0-5.8
	1996-2000	44.6	13.9	12.1-15.9	24.8	18.6	15.4-22.2	19.8	10.3	8.3-12.6	24.8	7.4	6.2-8.9	14.2	11.3	8.8-14.3	10.6	5.2	3.9-6.9
	2001-2005	47.0	14.3	12.5-16.3	25.8	18.8	15.7-22.3	21.2	10.8	8.8-13.2	25.2	7.3	6.1-8.7	12.8	9.9	7.6-12.7	12.4	5.8	4.4-7.5
	2006-2010	53.2	15.6	13.8-17.7	33.6	23.0	19.6-26.8	19.6	10.5	8.5-12.9	24.4	6.9	5.7-8.3	16.0	11.3	8.9-14.0	8.4	4.2	3.0-5.8
	2011-2015	58.0	16.6	14.7-18.7	32.0	21.0	17.8-24.6	26.0	13.6	11.3-16.4	22.8	5.8	4.7-7.0	13.2	8.3	6.4-10.6	9.6	4.2	3.1-5.7
2016-2020	64.0	17.0	15.1-19.0	38.6	22.9	19.6-26.5	25.4	12.1	10.0-14.6	22.8	5.7	4.7-6.9	13.8	8.3	6.4-10.6	9.0	3.9	2.8-5.3	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Allegany**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	174.2	399.8	373.2-427.9	88.0	467.5	423.1-515.0	86.2	360.8	326.7-397.5	93.0	215.3	195.8-236.1	51.8	284.2	249.4-322.3	41.2	167.5	145.0-192.5
	1981-1985	177.8	379.8	354.8-406.1	89.8	435.8	395.3-479.2	88.0	344.1	311.9-378.7	94.2	204.1	185.8-223.7	52.0	267.4	234.9-302.9	42.2	160.5	139.1-184.4
	1986-1990	206.0	423.9	398.1-451.1	100.0	464.8	424.0-508.5	106.0	407.4	372.6-444.5	104.2	213.5	195.3-232.9	52.2	251.8	221.4-285.1	52.0	191.2	168.2-216.6
	1991-1995	229.6	449.8	423.9-476.8	121.0	527.4	485.4-571.9	108.6	401.6	367.9-437.6	122.8	237.7	219.1-257.4	64.8	290.1	258.7-324.1	58.0	202.6	179.6-227.9
	1996-2000	254.8	488.8	462.2-516.6	131.0	547.8	506.3-591.9	123.8	450.6	415.3-488.1	124.4	235.8	217.5-255.2	69.0	296.4	265.7-329.7	55.4	193.0	170.6-217.6
	2001-2005	277.8	510.8	484.1-538.6	152.4	610.7	567.8-656.2	125.4	437.5	403.3-473.9	112.6	202.0	185.6-219.6	61.2	250.0	222.5-279.9	51.4	168.3	148.0-190.7
	2006-2010	270.4	478.7	453.1-505.5	142.6	527.2	488.7-568.1	127.8	443.6	408.9-480.6	97.6	169.8	154.9-185.8	53.8	204.9	180.9-231.3	43.8	145.6	126.5-166.9
	2011-2015	303.8	511.4	485.2-538.8	169.0	585.9	545.9-628.2	134.8	446.2	411.7-483.0	110.2	178.8	163.9-194.8	60.6	213.4	189.6-239.5	49.6	150.4	131.7-171.3
	2016-2020	291.8	471.7	446.5-498.1	157.0	504.7	468.5-543.0	134.8	444.1	408.8-481.8	105.4	162.4	148.4-177.5	55.2	180.4	159.1-204.0	50.2	147.0	128.7-167.6
Oral cavity and pharynx	1976-1980	4.4	9.9	6.1-15.0	3.0	14.6	8.0-24.4	1.4	5.5	2.1-11.7	2.4	5.4	2.8-9.6	1.8	8.1	3.6-15.5	0.6	2.4	0.5-7.2
	1981-1985	3.8	8.1	4.8-12.7	2.0	9.7	4.5-18.0	1.8	6.7	3.0-13.0	1.4	2.8	1.1-5.9	0.6	2.9	0.6-8.6	0.8	2.9	0.8-7.6
	1986-1990	4.2	8.6	5.3-13.3	2.6	11.3	6.0-19.5	1.6	5.8	2.4-11.7	1.6	3.0	1.3-6.0	0.8	3.2	0.9-8.4	0.8	2.8	0.7-7.4
	1991-1995	5.0	10.1	6.5-15.0	3.0	13.6	7.5-22.5	2.0	6.6	3.1-12.4	2.0	4.1	2.0-7.6	1.0	4.7	1.5-10.9	1.0	3.3	1.1-8.0
	1996-2000	5.8	11.1	7.4-16.0	3.6	14.8	8.7-23.4	2.2	8.0	3.9-14.5	1.4	2.7	1.1-5.6	1.2	5.2	1.9-11.4	0.2	0.7	0.0-4.1
	2001-2005	5.2	9.6	6.2-14.1	4.0	16.4	10.0-25.4	1.2	3.6	1.3-8.3	1.2	2.2	0.8-4.8	1.0	3.8	1.2-9.1	0.2	0.7	0.0-4.3
	2006-2010	8.6	15.3	11.0-20.8	6.8	24.8	17.1-35.0	1.8	6.1	2.7-12.1	2.0	3.3	1.6-6.3	1.4	5.1	2.0-10.9	0.6	1.8	0.4-5.9
	2011-2015	6.6	10.7	7.3-15.4	4.8	16.0	10.0-24.5	1.8	5.3	2.3-10.7	1.0	1.4	0.5-3.7	0.8	2.5	0.7-7.0	0.2	0.6	0.0-4.2
	2016-2020	8.0	13.0	9.0-18.2	6.6	21.7	14.5-31.5	1.4	4.3	1.6-9.8	1.6	2.7	1.1-5.6	1.6	5.5	2.3-11.5	0.0	0.0	0.0-3.5
Esophagus	1976-1980	1.4	2.9	1.2-6.1	0.8	3.8	1.0-9.9	0.6	2.1	0.4-6.5	1.8	4.5	2.0-8.4	1.2	6.6	2.4-14.4	0.6	2.4	0.5-7.1
	1981-1985	0.6	1.3	0.3-3.8	0.4	1.9	0.2-7.0	0.2	0.8	0.0-4.5	1.2	2.4	0.9-5.3	1.0	4.4	1.4-10.4	0.2	0.8	0.0-4.5
	1986-1990	2.2	4.5	2.2-8.1	2.0	9.5	4.4-17.6	0.2	0.6	0.0-4.0	2.2	4.6	2.3-8.2	2.0	9.9	4.6-18.2	0.2	0.6	0.0-4.0
	1991-1995	1.8	3.4	1.6-6.6	1.0	4.5	1.4-10.6	0.8	2.4	0.7-6.6	2.2	4.1	2.0-7.4	1.6	7.0	2.9-13.9	0.6	2.0	0.4-6.1
	1996-2000	3.6	7.0	4.1-11.1	3.2	13.4	7.6-21.9	0.4	1.5	0.2-5.6	3.6	7.1	4.2-11.2	3.4	14.7	8.6-23.6	0.2	0.8	0.0-4.5
	2001-2005	3.2	5.8	3.3-9.5	2.8	11.0	6.0-18.6	0.4	1.4	0.2-5.3	2.8	5.2	2.8-8.7	2.4	9.3	4.8-16.4	0.4	1.3	0.2-5.1
	2006-2010	3.4	5.8	3.3-9.4	2.6	9.3	4.9-16.2	0.8	2.6	0.7-7.2	2.6	4.3	2.3-7.5	2.4	8.3	4.2-14.8	0.2	0.7	0.0-4.4
	2011-2015	5.4	9.2	6.0-13.7	4.2	14.8	9.1-23.1	1.2	4.4	1.5-10.1	5.0	8.4	5.4-12.7	3.6	12.7	7.4-20.5	1.4	4.9	1.8-10.7
	2016-2020	3.0	4.3	2.4-7.5	2.2	6.6	3.2-12.5	0.8	2.3	0.6-6.8	2.8	4.2	2.3-7.4	2.4	8.1	4.1-14.7	0.4	0.9	0.1-4.7
Stomach	1976-1980	3.2	7.9	4.4-12.9	2.0	12.3	5.7-22.7	1.2	5.0	1.8-11.1	2.4	5.7	2.9-10.0	1.4	9.0	3.5-18.3	1.0	3.9	1.3-9.2
	1981-1985	2.8	6.0	3.3-10.1	2.4	11.4	5.8-20.1	0.4	1.6	0.2-5.8	2.2	4.8	2.4-8.6	1.8	9.4	4.2-17.8	0.4	1.4	0.2-5.4
	1986-1990	3.4	7.2	4.2-11.6	2.0	9.4	4.5-17.3	1.4	4.9	1.9-10.3	2.0	4.3	2.0-7.9	1.0	4.9	1.6-11.4	1.0	3.5	1.1-8.4
	1991-1995	4.0	7.7	4.7-11.9	3.0	13.2	7.3-21.9	1.0	3.4	1.1-8.3	3.6	6.8	4.0-10.9	3.0	13.2	7.3-21.9	0.6	1.9	0.4-6.0
	1996-2000	3.4	6.3	3.7-10.2	2.4	10.1	5.2-17.7	1.0	3.3	1.1-7.9	1.2	2.1	0.8-4.7	0.4	1.8	0.2-6.3	0.8	2.5	0.7-6.8
	2001-2005	3.2	5.9	3.4-9.7	2.8	11.2	6.1-19.0	0.4	1.3	0.2-5.1	1.6	2.9	1.2-5.9	1.0	4.2	1.3-9.8	0.6	1.9	0.4-6.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Allegany**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	3.8	6.7	4.0-10.7	2.6	9.8	5.2-17.0	1.2	4.4	1.5-10.0	2.0	3.3	1.6-6.3	1.6	5.3	2.3-11.0	0.4	1.2	0.1-5.0
	2011-2015	3.6	5.4	3.2-8.8	2.4	8.4	4.3-15.0	1.2	3.4	1.2-8.0	1.8	2.8	1.3-5.6	1.2	4.3	1.6-9.8	0.6	1.9	0.4-6.3
	2016-2020	5.4	8.8	5.7-13.2	3.2	10.2	5.7-17.1	2.2	7.7	3.7-14.5	1.0	1.5	0.5-3.9	0.6	2.2	0.5-7.0	0.4	0.9	0.1-4.7
Colorectal	1976-1980	29.6	67.5	56.9-79.4	14.2	75.4	58.3-95.6	15.4	62.0	48.7-77.9	15.6	36.7	28.9-45.8	7.4	40.7	28.3-56.3	8.2	32.8	23.5-44.7
	1981-1985	27.2	57.8	48.4-68.5	13.4	65.3	50.2-83.4	13.8	51.4	39.9-65.3	13.2	28.9	22.3-36.8	5.6	29.3	19.1-42.6	7.6	28.5	20.1-39.3
	1986-1990	28.0	57.2	48.0-67.6	13.0	64.0	49.0-82.0	15.0	53.5	41.9-67.4	13.0	26.3	20.3-33.6	6.4	32.7	22.0-46.4	6.6	23.6	16.1-33.4
	1991-1995	30.0	58.3	49.3-68.5	16.2	70.0	55.3-87.2	13.8	48.6	37.7-61.8	13.6	26.7	20.7-33.9	7.2	33.5	23.3-46.5	6.4	21.3	14.5-30.4
	1996-2000	35.8	67.9	58.2-78.6	17.6	76.0	60.9-93.8	18.2	61.4	49.3-75.8	16.4	30.7	24.4-38.2	7.6	33.2	23.4-45.6	8.8	28.3	20.4-38.3
	2001-2005	33.8	60.7	51.9-70.6	17.8	73.2	58.7-90.3	16.0	53.1	42.0-66.4	15.4	27.5	21.7-34.4	8.4	34.1	24.5-46.2	7.0	22.6	15.6-31.9
	2006-2010	29.8	51.8	43.7-61.0	16.6	61.8	49.0-77.0	13.2	42.9	32.9-55.1	10.0	16.9	12.5-22.5	6.4	25.0	17.1-35.5	3.6	10.9	6.4-17.6
	2011-2015	30.0	49.9	42.1-58.9	17.0	60.6	48.1-75.5	13.0	40.8	31.3-52.6	11.0	17.9	13.4-23.5	5.8	21.6	14.3-31.4	5.2	14.9	9.6-22.4
	2016-2020	26.0	43.6	36.1-52.4	15.0	52.9	41.1-67.1	11.0	35.2	25.9-47.0	10.4	16.6	12.3-22.2	4.2	14.3	8.7-22.5	6.2	18.2	12.1-26.8
Colon excluding rectum	1976-1980	20.2	46.3	37.6-56.4	8.8	47.7	34.3-64.4	11.4	45.6	34.3-59.3	12.8	30.0	23.0-38.3	6.0	32.4	21.6-46.5	6.8	27.5	18.9-38.6
	1981-1985	21.2	45.3	37.0-54.9	9.6	47.2	34.4-62.9	11.6	43.2	32.7-56.1	11.8	26.2	19.9-33.8	4.6	24.9	15.6-37.5	7.2	27.0	18.8-37.6
	1986-1990	21.2	43.3	35.4-52.4	9.4	46.8	34.0-62.7	11.8	42.0	31.8-54.5	11.4	23.1	17.5-30.1	5.8	30.1	19.8-43.4	5.6	20.3	13.4-29.7
	1991-1995	22.6	43.8	36.0-52.7	11.4	50.4	37.9-65.6	11.2	38.2	28.7-49.9	12.2	23.9	18.2-30.7	6.2	29.0	19.5-41.3	6.0	20.1	13.4-29.0
	1996-2000	27.4	51.7	43.4-61.2	12.6	54.2	41.5-69.5	14.8	49.3	38.5-62.2	14.8	27.7	21.8-34.9	6.2	27.0	18.3-38.4	8.6	27.6	19.8-37.5
	2001-2005	26.6	47.7	39.9-56.7	13.8	57.1	44.3-72.4	12.8	42.0	32.3-54.0	13.4	23.8	18.5-30.4	7.6	30.6	21.6-42.2	5.8	18.4	12.2-26.8
	2006-2010	21.6	38.3	31.3-46.4	10.8	41.5	31.0-54.4	10.8	35.4	26.4-46.8	9.2	15.6	11.4-21.0	5.8	22.9	15.3-33.1	3.4	10.2	5.9-16.8
	2011-2015	18.2	30.0	24.1-37.1	9.6	34.7	25.4-46.4	8.6	26.5	19.0-36.2	8.8	14.3	10.3-19.5	4.4	16.4	10.2-25.3	4.4	12.7	7.9-19.8
	2016-2020	18.6	31.6	25.1-39.3	10.2	36.7	26.9-49.0	8.4	27.5	19.2-38.3	9.0	14.7	10.6-20.1	3.6	12.8	7.4-20.7	5.4	16.2	10.3-24.5
Rectum & rectosigmoid	1976-1980	9.4	21.2	15.5-28.3	5.4	27.7	17.9-40.6	4.0	16.5	9.9-25.6	2.8	6.7	3.6-11.2	1.4	8.2	3.2-16.9	1.4	5.4	2.2-11.2
	1981-1985	6.0	12.5	8.4-18.0	3.8	18.1	10.7-28.6	2.2	8.2	4.0-14.9	1.4	2.7	1.1-5.7	1.0	4.4	1.3-10.6	0.4	1.5	0.2-5.6
	1986-1990	6.8	13.9	9.6-19.6	3.6	17.2	10.1-27.3	3.2	11.5	6.5-18.9	1.6	3.2	1.4-6.3	0.6	2.6	0.5-7.8	1.0	3.2	1.0-7.9
	1991-1995	7.4	14.5	10.2-20.1	4.8	19.6	12.5-29.2	2.6	10.4	5.5-17.8	1.4	2.8	1.1-5.8	1.0	4.4	1.4-10.3	0.4	1.3	0.2-5.0
	1996-2000	8.4	16.2	11.6-21.9	5.0	21.9	14.1-32.3	3.4	12.2	7.0-19.7	1.6	3.0	1.3-5.9	1.4	6.1	2.4-12.7	0.2	0.7	0.0-4.3
	2001-2005	7.2	13.0	9.1-18.0	4.0	16.2	9.8-25.1	3.2	11.1	6.3-18.2	2.0	3.6	1.7-6.8	0.8	3.4	0.9-8.8	1.2	4.2	1.5-9.5
	2006-2010	8.2	13.5	9.7-18.5	5.8	20.3	13.5-29.5	2.4	7.4	3.8-13.4	0.8	1.3	0.4-3.6	0.6	2.1	0.4-6.6	0.2	0.7	0.0-4.2
	2011-2015	11.8	19.9	15.0-26.0	7.4	25.9	18.0-36.3	4.4	14.3	8.8-22.3	2.2	3.5	1.8-6.6	1.4	5.1	2.0-10.9	0.8	2.2	0.6-6.5
	2016-2020	7.4	12.1	8.4-17.0	4.8	16.2	10.1-24.8	2.6	7.7	4.1-13.9	1.4	1.9	0.8-4.4	0.6	1.5	0.3-5.5	0.8	2.0	0.5-6.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Allegany**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	0.4	0.9	0.1-3.2	0.2	0.8	0.0-5.0	0.2	0.8	0.0-4.6	0.4	0.7	0.1-2.8	0.2	0.7	0.0-4.8	0.2	0.6	0.0-4.2
	1981-1985	0.4	1.1	0.1-3.7	0.2	1.3	0.0-6.3	0.2	1.3	0.0-5.9	0.4	1.1	0.1-3.7	0.2	1.3	0.0-6.3	0.2	1.3	0.0-5.9
	1986-1990	1.0	1.9	0.6-4.6	0.4	1.9	0.2-6.7	0.6	2.1	0.4-6.5	0.8	1.6	0.4-4.2	0.2	1.0	0.0-5.4	0.6	2.2	0.4-6.6
	1991-1995	2.0	3.6	1.7-6.7	1.4	5.8	2.3-12.0	0.6	1.9	0.4-6.0	2.0	3.7	1.8-6.8	1.4	6.6	2.6-13.6	0.6	1.9	0.4-5.9
	1996-2000	1.2	2.3	0.8-5.0	0.6	2.4	0.5-7.1	0.6	2.4	0.5-6.9	1.2	2.3	0.8-5.0	0.6	2.9	0.6-8.2	0.6	2.4	0.5-7.0
	2001-2005	2.2	3.9	2.0-7.1	1.6	6.5	2.8-12.9	0.6	2.0	0.4-6.2	2.4	4.3	2.2-7.6	1.4	5.8	2.3-12.1	1.0	3.0	1.0-7.5
	2006-2010	2.8	5.0	2.7-8.6	2.4	9.1	4.6-16.3	0.4	1.3	0.2-5.1	2.0	3.5	1.7-6.6	1.6	5.8	2.5-11.7	0.4	1.3	0.2-5.1
	2011-2015	2.2	3.6	1.8-6.6	1.6	5.0	2.1-10.5	0.6	2.2	0.4-6.8	1.0	1.5	0.5-3.8	0.4	1.3	0.1-5.3	0.6	1.6	0.3-5.6
2016-2020	3.6	5.2	3.1-8.6	2.4	7.1	3.6-13.0	1.2	3.6	1.3-8.6	2.6	4.0	2.1-7.1	1.6	5.1	2.2-10.7	1.0	3.2	1.0-8.2	
Pancreas	1976-1980	5.6	13.2	8.7-19.0	2.2	12.1	5.9-21.8	3.4	13.3	7.7-21.4	5.8	13.5	9.0-19.4	2.6	14.6	7.6-25.1	3.2	12.5	7.1-20.4
	1981-1985	6.0	12.3	8.3-17.6	3.4	16.8	9.6-27.0	2.6	9.0	4.8-15.5	4.6	9.4	5.9-14.2	2.8	13.5	7.3-22.9	1.8	6.2	2.8-12.1
	1986-1990	5.8	11.9	7.9-17.1	3.4	16.9	9.6-27.2	2.4	8.7	4.4-15.4	6.0	12.4	8.3-17.8	3.2	17.4	9.7-28.4	2.8	10.1	5.5-17.1
	1991-1995	6.6	12.7	8.7-17.9	2.8	11.8	6.3-19.9	3.8	13.5	8.0-21.3	5.6	10.6	7.0-15.4	2.8	11.3	6.1-19.2	2.8	9.6	5.2-16.5
	1996-2000	4.4	8.4	5.2-12.7	2.6	10.7	5.7-18.5	1.8	6.5	2.9-12.5	4.8	9.1	5.8-13.7	3.0	12.7	7.0-21.0	1.8	6.6	3.0-12.8
	2001-2005	6.0	10.6	7.2-15.3	2.2	8.8	4.4-15.9	3.8	12.2	7.3-19.4	4.6	8.1	5.1-12.2	1.6	6.9	2.9-13.6	3.0	9.7	5.4-16.3
	2006-2010	4.8	8.2	5.2-12.3	2.6	9.8	5.2-16.9	2.2	6.6	3.3-12.4	4.8	8.3	5.3-12.5	2.2	8.7	4.3-15.7	2.6	8.2	4.3-14.5
	2011-2015	8.6	13.9	10.0-19.0	3.8	13.0	7.7-20.7	4.8	14.1	9.0-21.6	5.4	8.6	5.7-12.8	2.8	8.9	4.8-15.6	2.6	7.7	4.1-13.7
2016-2020	9.0	12.9	9.4-17.6	4.6	13.1	8.2-20.3	4.4	12.6	7.8-19.7	6.2	9.4	6.3-13.7	2.8	8.1	4.4-14.3	3.4	10.4	5.9-17.4	
Larynx	1976-1980	2.2	5.1	2.5-9.2	1.8	8.7	3.9-16.6	0.4	2.2	0.2-7.5	1.2	2.6	0.9-5.7	1.2	6.6	2.2-14.5	0.0	0.0	0.0-3.1
	1981-1985	2.0	4.6	2.2-8.6	2.0	9.8	4.6-18.2	0.0	0.0	0.0-3.1	0.8	1.7	0.4-4.3	0.4	1.8	0.2-6.8	0.4	1.4	0.2-5.3
	1986-1990	2.6	5.2	2.7-9.0	2.0	9.0	4.2-16.8	0.6	2.6	0.5-7.7	1.0	2.4	0.8-5.6	0.6	3.2	0.6-8.9	0.4	1.9	0.2-6.7
	1991-1995	3.4	6.5	3.8-10.5	3.0	12.6	7.0-21.0	0.4	1.4	0.2-5.4	1.0	1.9	0.6-4.4	1.0	4.8	1.5-11.0	0.0	0.0	0.0-2.9
	1996-2000	4.2	8.1	5.0-12.4	2.8	11.2	6.1-18.9	1.4	5.2	2.1-10.9	1.0	2.0	0.6-4.7	0.6	2.4	0.5-7.2	0.4	1.8	0.2-6.2
	2001-2005	2.0	3.5	1.7-6.6	2.0	7.4	3.5-13.8	0.0	0.0	0.0-3.1	0.8	1.4	0.4-3.8	0.8	3.1	0.8-8.1	0.0	0.0	0.0-3.1
	2006-2010	2.2	4.0	1.9-7.3	2.0	7.3	3.4-13.8	0.2	0.7	0.0-4.4	1.2	1.8	0.7-4.3	1.2	4.0	1.4-9.1	0.0	0.0	0.0-3.2
	2011-2015	3.4	6.1	3.5-10.1	2.8	9.8	5.2-17.0	0.6	2.6	0.5-7.8	0.8	1.4	0.4-3.7	0.8	2.9	0.8-7.7	0.0	0.0	0.0-3.3
2016-2020	2.8	4.0	2.2-7.0	2.6	7.3	3.9-13.2	0.2	0.6	0.0-4.3	1.0	1.4	0.4-3.7	0.8	2.4	0.6-6.8	0.2	0.5	0.0-4.1	
Lung and bronchus	1976-1980	27.0	61.5	51.4-73.0	19.8	100.1	80.8-122.4	7.2	30.3	21.0-42.2	18.4	41.6	33.4-51.1	15.0	76.1	59.4-95.9	3.4	15.0	8.6-24.2
	1981-1985	35.6	75.6	64.7-87.9	23.2	111.0	91.2-133.7	12.4	47.2	35.9-60.9	28.2	60.4	50.7-71.5	19.2	94.4	76.0-115.8	9.0	33.1	24.0-44.5
	1986-1990	33.4	66.6	56.7-77.7	22.2	101.4	83.0-122.6	11.2	39.6	29.8-51.6	29.6	60.2	50.7-70.8	20.8	96.9	78.8-117.8	8.8	30.5	22.1-41.2
	1991-1995	38.4	74.7	64.4-86.1	22.8	98.2	80.8-118.3	15.6	56.5	44.5-70.9	33.4	64.1	54.7-74.7	20.6	88.3	71.9-107.4	12.8	45.2	34.7-58.0
	1996-2000	43.0	81.5	70.9-93.2	25.0	103.3	85.8-123.3	18.0	64.9	52.1-80.0	33.2	62.1	53.0-72.4	21.6	88.6	72.6-107.2	11.6	41.1	31.2-53.4
	2001-2005	46.2	83.5	73.0-95.0	27.8	111.1	93.2-131.4	18.4	61.5	49.5-75.8	36.0	64.9	55.7-75.2	21.8	87.8	72.0-106.1	14.2	46.5	36.2-58.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Allegany**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	41.0	71.7	62.2-82.5	21.0	78.4	64.0-95.3	20.0	67.5	54.8-82.5	25.0	43.3	36.0-51.8	13.6	50.3	38.9-64.1	11.4	38.2	28.8-49.9
	2011-2015	48.6	78.4	68.7-89.2	27.6	93.9	78.6-111.5	21.0	64.8	52.8-78.9	32.6	52.5	44.7-61.5	19.4	66.6	53.8-81.6	13.2	40.2	30.9-51.6
	2016-2020	45.6	67.6	59.0-77.4	23.6	72.0	59.3-86.9	22.0	63.9	52.3-77.8	28.2	41.3	34.6-49.0	16.0	49.7	39.2-62.5	12.2	33.7	25.7-43.9
Melanoma of the skin	1976-1980	3.0	7.2	4.0-11.9	1.8	8.9	4.0-17.0	1.2	5.5	1.9-11.9	1.4	3.7	1.5-7.5	1.0	6.5	2.1-14.6	0.4	1.9	0.2-6.7
	1981-1985	2.8	6.6	3.6-11.2	1.0	5.6	1.8-12.9	1.8	7.2	3.2-13.8	0.4	1.1	0.1-3.8	0.0	0.0	0.0-3.5	0.4	1.9	0.2-6.8
	1986-1990	3.2	7.1	4.0-11.6	1.8	7.9	3.5-15.3	1.4	6.4	2.5-13.2	1.0	2.4	0.8-5.5	0.4	2.0	0.2-7.1	0.6	2.5	0.5-7.5
	1991-1995	3.6	7.6	4.5-12.0	2.6	11.5	6.1-19.6	1.0	4.1	1.3-9.5	1.2	2.4	0.9-5.3	1.0	4.4	1.4-10.2	0.2	0.7	0.0-4.0
	1996-2000	7.2	14.6	10.2-20.3	4.6	19.5	12.3-29.3	2.6	9.6	5.1-16.6	2.4	4.9	2.5-8.5	1.6	6.9	3.0-13.7	0.8	2.6	0.7-7.1
	2001-2005	10.4	20.0	14.8-26.3	6.4	26.4	17.9-37.5	4.0	15.4	9.3-24.0	1.0	1.8	0.6-4.3	0.6	2.1	0.4-6.6	0.4	1.3	0.2-5.1
	2006-2010	11.0	20.0	14.9-26.2	5.6	20.8	13.8-30.3	5.4	19.9	12.9-29.4	1.4	2.8	1.1-5.8	0.6	2.6	0.5-7.5	0.8	3.2	0.8-8.4
	2011-2015	12.0	21.1	15.9-27.6	8.2	29.3	20.6-40.4	3.8	13.6	7.9-21.8	2.0	3.1	1.5-6.0	1.6	5.2	2.2-10.7	0.4	1.3	0.2-5.3
	2016-2020	16.6	30.0	23.5-37.7	10.2	35.4	26.1-47.2	6.4	26.1	17.2-37.9	1.8	2.8	1.2-5.6	1.4	4.5	1.8-10.0	0.4	1.5	0.2-5.9
Female breast	1976-1980							22.4	97.1	79.4-117.3							8.8	37.0	26.6-50.0
	1981-1985							21.2	90.0	73.2-109.4							7.8	31.7	22.3-43.8
	1986-1990							28.8	117.7	98.7-139.1							11.0	43.7	32.6-57.3
	1991-1995							29.8	113.6	95.8-133.8							11.4	41.5	31.2-54.0
	1996-2000							35.4	133.8	114.6-155.3							10.0	35.8	26.5-47.5
	2001-2005							35.0	124.6	106.6-145.0							5.8	19.9	13.2-28.9
	2006-2010							36.6	128.2	109.8-149.0							6.4	22.3	15.0-32.0
	2011-2015							37.0	125.8	107.4-146.5							5.4	18.7	12.0-27.9
	2016-2020							37.0	130.1	110.5-152.3							7.2	21.6	14.9-30.6
Cervix uteri	1976-1980							5.2	25.0	16.1-36.8							1.4	5.9	2.3-12.2
	1981-1985							3.2	13.2	7.4-21.6							1.2	5.9	2.1-12.9
	1986-1990							2.2	10.8	5.3-19.3							1.0	4.4	1.4-10.2
	1991-1995							3.4	14.6	8.5-23.4							1.2	5.1	1.8-11.0
	1996-2000							1.8	8.1	3.7-15.3							1.2	4.8	1.8-10.5
	2001-2005							2.8	12.0	6.4-20.3							1.2	4.6	1.6-10.3
	2006-2010							3.0	12.7	6.9-21.2							1.4	5.3	2.1-11.2
	2011-2015							2.0	8.8	3.9-16.6							0.4	1.0	0.1-4.8
	2016-2020							1.6	6.5	2.6-13.4							1.2	4.7	1.5-11.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Allegany**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							3.4	13.1	7.6-21.1							1.0	4.3	1.3-10.1
	1981-1985							5.4	20.8	13.6-30.6							0.8	2.7	0.7-7.3
	1986-1990							8.8	33.8	24.4-45.7							0.8	2.7	0.7-7.2
	1991-1995							5.6	21.0	13.8-30.5							0.6	1.7	0.3-5.4
	1996-2000							7.8	29.0	20.6-39.8							1.0	3.6	1.2-8.6
	2001-2005							9.8	34.6	25.5-46.0							1.0	3.3	1.0-8.0
	2006-2010							7.0	24.6	17.0-34.5							0.4	1.3	0.2-5.3
	2011-2015							9.2	31.8	22.9-43.1							1.4	4.4	1.8-9.6
	2016-2020							11.6	36.6	27.2-48.4							2.0	6.1	2.8-12.1
Ovary	1976-1980							3.4	14.1	8.1-22.8							2.4	9.5	4.8-16.8
	1981-1985							3.6	14.0	8.2-22.5							1.2	4.6	1.6-10.2
	1986-1990							5.0	18.3	11.7-27.3							3.2	11.2	6.3-18.5
	1991-1995							6.0	22.7	15.2-32.6							3.6	12.8	7.5-20.4
	1996-2000							5.6	20.5	13.6-29.9							4.0	14.2	8.6-22.1
	2001-2005							4.0	13.6	8.3-21.4							2.2	7.4	3.6-13.5
	2006-2010							3.6	12.3	7.1-19.8							2.4	8.3	4.2-15.0
	2011-2015							6.4	21.2	14.3-30.5							4.2	13.8	8.3-21.7
	2016-2020							3.2	9.6	5.3-16.4							3.6	10.4	6.0-17.2
Prostate	1976-1980				16.2	95.2	74.9-118.9							5.8	36.1	23.8-52.0			
	1981-1985				16.0	79.9	62.9-99.9							5.8	34.1	22.6-48.9			
	1986-1990				20.6	95.4	77.5-116.1							5.4	27.4	17.8-40.2			
	1991-1995				33.8	147.9	126.0-172.4							8.4	41.3	29.5-55.9			
	1996-2000				32.2	132.7	112.8-155.1							7.8	35.7	25.3-48.8			
	2001-2005				43.0	167.2	145.4-191.3							5.8	26.4	17.6-37.8			
	2006-2010				34.8	124.0	106.1-144.2							4.2	17.6	10.9-27.0			
	2011-2015				38.6	122.2	105.3-141.4							6.0	21.6	14.5-31.2			
	2016-2020				38.8	115.3	99.2-133.6							5.2	17.6	11.4-26.3			
Testis	1976-1980				0.6	2.7	0.4-8.4							0.0	0.0	0.0-3.7			
	1981-1985				0.4	1.6	0.2-6.2							0.0	0.0	0.0-3.5			
	1986-1990				1.2	5.9	2.1-12.8							0.0	0.0	0.0-3.4			
	1991-1995				1.6	7.1	3.0-14.0							0.0	0.0	0.0-3.2			
	1996-2000				1.0	4.6	1.5-10.7							0.4	1.9	0.2-6.6			
	2001-2005				1.2	6.0	2.1-12.9							0.0	0.0	0.0-3.2			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Allegany**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				0.8	2.8	0.7-7.6							0.0	0.0	0.0-3.2			
	2011-2015				2.4	12.7	6.4-22.2							0.2	1.0	0.0-5.3			
	2016-2020				1.2	4.6	1.5-10.5							0.0	0.0	0.0-3.4			
Urinary bladder (incl. in situ)	1976-1980	8.6	19.9	14.3-26.8	6.8	38.8	26.4-54.6	1.8	7.1	3.2-13.5	3.4	8.1	4.7-12.9	2.8	17.3	9.2-28.9	0.6	2.3	0.5-6.9
	1981-1985	10.2	21.1	15.6-27.9	8.2	38.1	27.1-52.1	2.0	7.3	3.4-13.7	2.0	4.4	2.1-8.0	1.6	8.9	3.7-17.5	0.4	1.4	0.2-5.3
	1986-1990	10.4	20.7	15.4-27.3	8.4	39.3	28.1-53.3	2.0	6.9	3.3-13.0	2.0	3.9	1.9-7.3	1.6	7.6	3.3-14.8	0.4	1.2	0.1-5.0
	1991-1995	8.4	16.4	11.8-22.3	5.6	24.8	16.4-36.0	2.8	9.9	5.3-16.9	3.2	6.1	3.5-10.0	2.0	9.2	4.3-16.9	1.2	4.3	1.6-9.6
	1996-2000	13.6	25.2	19.5-32.0	10.4	44.3	33.0-58.3	3.2	11.2	6.3-18.4	4.2	7.8	4.8-12.1	3.2	13.7	7.8-22.4	1.0	3.1	1.0-7.6
	2001-2005	14.4	25.8	20.1-32.5	12.0	48.4	36.8-62.4	2.4	7.5	3.9-13.6	3.8	6.8	4.1-10.7	2.8	11.7	6.4-19.7	1.0	3.1	1.0-7.7
	2006-2010	16.8	28.6	22.8-35.6	14.2	51.4	40.0-65.1	2.6	8.4	4.5-14.8	3.2	5.5	3.1-9.0	2.0	8.0	3.8-14.8	1.2	3.7	1.3-8.5
	2011-2015	18.2	29.9	24.0-37.0	15.2	54.2	42.5-68.2	3.0	8.9	4.9-15.2	3.4	5.3	3.1-8.7	3.0	11.0	6.1-18.4	0.4	1.3	0.2-5.3
2016-2020	15.8	25.4	19.9-32.1	12.8	42.5	32.4-55.0	3.0	9.6	5.1-16.7	3.0	4.8	2.6-8.2	2.2	8.1	3.9-15.0	0.8	2.2	0.6-6.5	
Kidney and renal pelvis	1976-1980	3.0	7.3	4.0-12.1	1.8	9.3	4.1-17.8	1.2	5.0	1.8-11.0	1.8	4.4	2.0-8.5	1.2	6.6	2.3-14.4	0.6	2.2	0.5-6.8
	1981-1985	4.0	8.3	5.0-12.9	1.8	8.7	3.9-16.6	2.2	8.1	4.0-14.7	1.6	3.3	1.4-6.5	0.6	3.3	0.7-9.5	1.0	3.6	1.2-8.7
	1986-1990	5.6	11.8	7.8-17.2	3.2	14.5	8.2-23.8	2.4	10.2	5.2-18.0	3.8	8.1	4.8-12.7	1.6	6.8	2.9-13.5	2.2	9.3	4.5-16.8
	1991-1995	5.0	9.7	6.3-14.4	2.2	9.4	4.6-16.9	2.8	10.2	5.5-17.3	2.2	4.1	2.1-7.5	1.2	4.8	1.7-10.6	1.0	3.2	1.0-7.7
	1996-2000	5.4	10.9	7.1-15.8	2.2	9.2	4.6-16.5	3.2	12.3	7.0-20.0	1.8	3.5	1.6-6.7	1.2	5.0	1.8-11.0	0.6	2.1	0.4-6.3
	2001-2005	10.8	20.7	15.5-27.1	6.6	26.8	18.4-37.8	4.2	16.1	9.9-24.8	2.2	3.8	1.9-7.0	1.2	5.0	1.8-10.9	1.0	3.4	1.1-8.2
	2006-2010	8.4	15.0	10.7-20.4	3.8	14.0	8.4-22.2	4.6	16.1	10.0-24.5	1.8	3.3	1.5-6.4	1.4	5.6	2.2-11.8	0.4	1.3	0.2-5.3
	2011-2015	9.6	16.4	12.0-22.0	5.6	19.6	12.8-28.8	4.0	13.4	8.0-21.2	4.2	6.5	4.0-10.2	2.8	10.3	5.6-17.6	1.4	3.9	1.5-8.8
2016-2020	10.4	16.9	12.4-22.7	6.4	20.4	13.7-29.5	4.0	13.6	7.9-22.1	2.6	4.3	2.2-7.7	2.2	7.8	3.7-14.5	0.4	1.1	0.1-5.1	
Brain and other central nervous system	1976-1980	3.8	8.4	4.9-13.3	1.8	8.6	3.9-16.5	2.0	8.3	3.8-15.5	2.0	3.6	1.7-6.9	1.0	3.9	1.3-9.6	1.0	3.5	1.1-8.6
	1981-1985	2.4	4.5	2.3-8.0	1.4	5.3	2.1-11.2	1.0	3.9	1.2-9.4	2.4	4.9	2.5-8.6	2.0	8.7	4.1-16.3	0.4	1.4	0.2-5.4
	1986-1990	4.2	8.6	5.3-13.2	2.0	9.0	4.2-16.9	2.2	8.1	4.0-14.7	2.4	4.7	2.4-8.3	0.4	2.2	0.2-7.7	2.0	7.0	3.3-13.0
	1991-1995	3.8	7.9	4.7-12.4	1.8	8.4	3.8-16.0	2.0	7.6	3.6-14.2	3.0	6.1	3.4-10.1	2.0	9.5	4.5-17.4	1.0	3.6	1.2-8.5
	1996-2000	4.4	8.7	5.4-13.3	2.8	11.3	6.1-19.1	1.6	6.1	2.6-12.1	2.6	4.9	2.6-8.4	1.4	5.5	2.2-11.4	1.2	4.1	1.5-9.2
	2001-2005	3.4	7.1	4.1-11.5	1.8	7.8	3.4-14.9	1.6	6.4	2.7-12.8	1.8	3.1	1.4-6.1	1.2	4.2	1.5-9.4	0.6	1.9	0.4-6.1
	2006-2010	5.2	10.2	6.6-15.2	3.4	13.6	7.8-22.1	1.8	6.7	2.9-13.2	3.4	6.5	3.7-10.6	2.2	8.0	4.0-14.7	1.2	4.8	1.6-10.8
	2011-2015	4.2	7.7	4.7-12.1	3.0	11.0	5.9-18.7	1.2	4.5	1.6-10.2	2.4	3.6	1.9-6.6	1.4	4.2	1.7-9.4	1.0	3.0	1.0-7.7
2016-2020	3.0	5.0	2.7-8.6	1.8	6.6	2.8-13.2	1.2	3.2	1.2-7.8	2.2	3.3	1.6-6.3	1.4	3.8	1.5-8.7	0.8	2.8	0.7-7.9	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Allegany**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	1.6	3.8	1.6-7.5	0.6	3.5	0.7-9.9	1.0	4.4	1.4-10.3	0.6	1.6	0.3-4.4	0.2	1.5	0.0-7.0	0.4	1.6	0.2-5.8
	1981-1985	1.0	1.9	0.6-4.6	0.4	1.7	0.2-6.5	0.6	2.0	0.4-6.3	0.0	0.0	0.0-1.6	0.0	0.0	0.0-3.5	0.0	0.0	0.0-3.1
	1986-1990	1.0	2.3	0.7-5.5	0.4	2.2	0.3-7.5	0.6	2.7	0.5-7.8	0.2	0.4	0.0-2.3	0.2	1.0	0.0-5.4	0.0	0.0	0.0-3.0
	1991-1995	1.6	3.1	1.3-6.2	0.6	2.6	0.5-7.5	1.0	3.9	1.2-9.2	0.4	0.8	0.1-2.9	0.4	1.6	0.2-6.1	0.0	0.0	0.0-2.9
	1996-2000	3.2	6.9	4.0-11.3	0.8	3.2	0.9-8.4	2.4	10.7	5.5-18.6	0.0	0.0	0.0-1.5	0.0	0.0	0.0-3.2	0.0	0.0	0.0-3.0
	2001-2005	3.6	7.5	4.4-12.0	0.8	3.0	0.8-7.9	2.8	11.8	6.3-20.0	0.6	1.1	0.2-3.3	0.2	0.8	0.0-4.5	0.4	1.2	0.1-4.9
	2006-2010	4.8	9.4	5.9-14.2	1.0	3.9	1.2-9.5	3.8	14.9	8.8-23.6	0.4	0.7	0.1-2.7	0.2	0.7	0.0-4.3	0.2	0.7	0.0-4.4
	2011-2015	6.6	14.2	9.5-20.3	2.0	7.6	3.5-14.4	4.6	21.0	12.9-32.1	0.4	0.6	0.1-2.4	0.0	0.0	0.0-3.3	0.4	1.1	0.1-5.0
2016-2020	4.4	9.9	6.1-15.2	1.4	7.3	2.9-15.0	3.0	12.6	6.8-21.2	0.0	0.0	0.0-1.7	0.0	0.0	0.0-3.4	0.0	0.0	0.0-3.5	
Hodgkin lymphoma	1976-1980	1.6	3.7	1.5-7.4	0.8	4.0	1.0-10.3	0.8	3.3	0.8-8.5	0.4	1.1	0.1-3.8	0.4	2.4	0.3-8.2	0.0	0.0	0.0-3.1
	1981-1985	0.6	0.8	0.2-2.8	0.4	1.2	0.1-5.2	0.2	0.5	0.0-3.8	0.2	0.2	0.0-1.9	0.2	0.4	0.0-4.0	0.0	0.0	0.0-3.1
	1986-1990	2.4	4.2	2.1-7.6	1.0	2.9	0.9-7.8	1.4	5.7	2.2-11.9	0.6	1.4	0.3-3.9	0.4	1.9	0.2-6.7	0.2	0.6	0.0-4.0
	1991-1995	1.0	1.9	0.6-4.4	0.4	1.4	0.2-5.7	0.6	2.2	0.4-6.7	0.2	0.4	0.0-2.2	0.0	0.0	0.0-3.2	0.2	0.6	0.0-3.8
	1996-2000	0.4	1.0	0.1-3.3	0.2	1.0	0.0-5.3	0.2	0.9	0.0-4.9	0.0	0.0	0.0-1.5	0.0	0.0	0.0-3.2	0.0	0.0	0.0-3.0
	2001-2005	0.6	1.4	0.3-4.1	0.4	1.7	0.2-6.2	0.2	1.1	0.0-5.4	0.6	1.2	0.2-3.7	0.6	2.5	0.5-7.5	0.0	0.0	0.0-3.1
	2006-2010	0.8	1.4	0.4-3.8	0.6	2.5	0.5-7.3	0.2	0.6	0.0-4.1	0.2	0.4	0.0-2.4	0.2	0.8	0.0-4.8	0.0	0.0	0.0-3.2
	2011-2015	2.0	4.0	1.8-7.5	1.4	5.7	2.1-12.1	0.6	2.3	0.4-7.1	0.0	0.0	0.0-1.7	0.0	0.0	0.0-3.3	0.0	0.0	0.0-3.3
2016-2020	2.0	4.1	1.8-7.7	0.8	3.3	0.8-8.8	1.2	4.9	1.6-11.2	0.2	0.3	0.0-2.3	0.2	0.7	0.0-4.6	0.0	0.0	0.0-3.5	
Non-Hodgkin lymphomas	1976-1980	7.8	17.2	12.1-23.6	3.2	14.9	8.3-24.6	4.6	19.1	12.0-28.8	4.2	9.5	5.9-14.6	2.4	12.5	6.3-21.9	1.8	7.4	3.4-14.1
	1981-1985	4.6	10.2	6.4-15.3	1.4	6.0	2.4-12.6	3.2	13.2	7.4-21.6	2.8	6.0	3.3-10.2	1.4	6.9	2.7-14.4	1.4	5.6	2.2-11.7
	1986-1990	10.4	21.8	16.2-28.7	5.0	23.1	14.9-34.2	5.4	21.0	13.7-30.7	4.4	8.8	5.5-13.3	1.8	7.9	3.6-15.2	2.6	9.2	4.8-16.0
	1991-1995	7.6	14.9	10.5-20.5	3.4	15.0	8.6-24.1	4.2	14.1	8.7-21.9	5.4	10.1	6.6-14.8	1.8	7.9	3.6-15.1	3.6	11.3	6.7-18.1
	1996-2000	8.8	17.3	12.5-23.3	5.4	23.0	15.1-33.6	3.4	12.4	7.2-20.1	6.0	11.9	8.0-17.1	4.4	19.1	11.9-28.9	1.6	5.9	2.5-11.8
	2001-2005	9.4	17.2	12.6-23.0	4.0	16.2	9.8-25.2	5.4	18.5	12.1-27.2	4.2	7.4	4.6-11.4	2.0	8.3	3.9-15.4	2.2	7.1	3.5-13.1
	2006-2010	10.6	19.1	14.3-25.2	5.8	21.9	14.6-31.7	4.8	16.7	10.5-25.4	4.2	7.4	4.6-11.5	2.8	11.3	6.2-19.1	1.4	4.3	1.7-9.3
	2011-2015	11.8	19.4	14.7-25.3	6.4	21.9	14.8-31.4	5.4	17.1	11.0-25.5	2.4	4.0	2.0-7.3	1.2	4.9	1.7-11.0	1.2	3.1	1.1-7.5
2016-2020	10.6	17.2	12.6-23.0	5.8	19.8	12.9-29.3	4.8	15.1	9.4-23.4	3.0	4.9	2.6-8.4	1.6	6.0	2.4-12.5	1.4	3.9	1.5-8.8	
Myeloma	1976-1980	1.4	3.1	1.2-6.5	0.6	3.9	0.7-11.0	0.8	3.2	0.9-8.4	0.6	1.4	0.3-4.1	0.4	3.3	0.4-10.5	0.2	0.6	0.0-4.2
	1981-1985	3.2	6.4	3.6-10.4	2.4	12.0	6.1-21.2	0.8	2.7	0.7-7.3	1.6	3.3	1.4-6.6	1.4	7.6	3.0-15.5	0.2	0.6	0.0-4.1
	1986-1990	2.0	4.0	1.9-7.4	0.4	1.8	0.2-6.6	1.6	5.4	2.3-11.0	2.0	3.9	1.9-7.2	0.6	2.3	0.5-7.1	1.4	4.8	1.9-10.2
	1991-1995	2.6	5.0	2.7-8.7	1.6	7.5	3.2-14.8	1.0	3.5	1.1-8.4	2.0	3.9	1.9-7.2	1.2	5.7	2.0-12.3	0.8	2.8	0.8-7.4
	1996-2000	3.2	6.0	3.4-9.9	2.0	8.6	4.1-15.8	1.2	4.0	1.5-9.1	2.8	5.5	3.0-9.2	1.8	8.4	3.8-15.9	1.0	3.6	1.1-8.6
	2001-2005	3.2	5.7	3.2-9.3	1.8	7.6	3.5-14.6	1.4	4.4	1.8-9.5	1.8	3.2	1.5-6.1	1.0	4.2	1.3-9.9	0.8	2.7	0.7-7.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Allegany**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	3.0	5.3	2.9-8.8	2.0	8.1	3.8-15.0	1.0	3.5	1.1-8.5	1.0	1.8	0.6-4.3	0.8	3.3	0.9-8.6	0.2	0.7	0.0-4.4
	2011-2015	5.2	8.6	5.6-12.9	3.4	12.4	7.1-20.2	1.8	5.2	2.3-10.5	1.6	2.5	1.1-5.2	0.8	2.8	0.7-7.6	0.8	2.3	0.6-6.6
	2016-2020	2.0	2.8	1.3-5.5	1.6	4.5	1.9-9.7	0.4	0.9	0.1-4.7	1.0	1.6	0.5-4.0	0.6	2.0	0.4-6.5	0.4	0.9	0.1-4.7
Leukemias	1976-1980	6.0	12.5	8.3-18.0	3.6	16.4	9.4-26.4	2.4	9.7	5.0-17.2	3.4	7.5	4.3-12.1	2.2	12.0	5.8-21.7	1.2	4.7	1.7-10.3
	1981-1985	7.4	16.6	11.6-22.9	4.4	22.2	13.7-33.7	3.0	12.5	6.9-20.8	6.4	13.8	9.4-19.6	3.4	16.8	9.6-27.2	3.0	11.6	6.3-19.4
	1986-1990	4.8	10.1	6.4-15.1	1.6	6.8	2.9-13.6	3.2	12.6	7.1-20.7	3.0	5.7	3.2-9.5	1.8	8.0	3.5-15.4	1.2	4.2	1.5-9.4
	1991-1995	7.0	13.8	9.6-19.3	4.0	17.9	10.7-27.8	3.0	11.9	6.6-19.7	4.4	8.9	5.5-13.5	1.6	7.4	3.1-14.6	2.8	10.2	5.5-17.4
	1996-2000	7.6	14.4	10.2-19.9	3.4	14.2	8.2-22.8	4.2	13.8	8.5-21.5	4.6	8.8	5.6-13.2	2.0	9.4	4.5-17.1	2.6	8.2	4.3-14.3
	2001-2005	9.4	17.7	13.0-23.7	6.0	23.9	16.0-34.2	3.4	11.6	6.7-19.0	3.8	6.9	4.2-11.0	2.4	9.7	5.0-17.1	1.4	4.7	1.8-10.1
	2006-2010	10.6	19.6	14.6-25.9	6.8	26.9	18.5-38.0	3.8	12.5	7.4-20.1	4.8	8.6	5.5-12.9	2.8	11.1	6.0-18.7	2.0	7.0	3.3-13.1
	2011-2015	10.0	16.4	12.1-22.0	7.2	25.7	17.8-36.1	2.8	8.7	4.5-15.3	3.4	5.4	3.1-8.9	2.4	8.9	4.5-16.0	1.0	2.5	0.8-6.7
	2016-2020	7.0	11.5	7.9-16.3	4.6	16.3	10.1-25.0	2.4	7.0	3.5-13.1	3.6	6.2	3.6-10.0	2.0	7.5	3.6-14.2	1.6	4.6	2.0-9.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Broome**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	868.8	410.4	398.0-423.1	431.6	502.5	480.4-525.3	437.2	365.1	349.6-381.2	440.4	208.9	200.1-218.0	232.0	278.1	261.5-295.5	208.4	169.9	159.5-180.7
	1981-1985	940.0	427.6	415.2-440.3	461.8	519.7	497.8-542.4	478.2	382.3	366.6-398.4	475.8	215.1	206.4-224.1	250.6	292.1	275.4-309.5	225.2	172.4	162.2-183.1
	1986-1990	1,095.2	483.9	470.9-497.1	525.0	569.5	547.1-592.6	570.2	445.0	428.3-462.3	481.8	210.3	201.9-219.0	251.0	283.6	267.5-300.4	230.8	170.0	160.0-180.4
	1991-1995	1,160.6	491.2	478.5-504.2	599.2	603.5	581.6-626.0	561.4	424.4	408.4-440.9	492.8	204.9	196.8-213.3	250.2	263.1	248.3-278.5	242.6	171.6	161.8-181.8
	1996-2000	1,163.2	491.7	479.0-504.6	578.0	571.4	550.6-592.7	585.2	443.3	426.9-460.2	508.2	206.1	198.1-214.3	253.0	254.9	241.0-269.5	255.2	175.3	165.5-185.5
	2001-2005	1,205.4	500.8	488.1-513.8	632.2	604.6	583.6-626.1	573.2	430.3	414.3-446.9	467.2	183.6	176.2-191.3	234.8	227.5	214.6-240.9	232.4	157.0	147.8-166.7
	2006-2010	1,267.0	512.2	499.4-525.2	665.4	605.2	584.6-626.4	601.6	446.5	430.0-463.5	459.0	173.7	166.5-181.1	232.2	213.1	201.0-225.9	226.8	148.0	139.2-157.3
	2011-2015	1,213.6	480.7	468.2-493.4	596.6	511.0	492.4-530.1	617.0	464.9	447.6-482.6	430.4	157.6	150.9-164.7	215.2	184.8	173.8-196.4	215.2	139.3	130.7-148.3
	2016-2020	1,271.6	493.0	480.3-505.9	649.6	527.8	509.2-547.1	622.0	470.6	453.0-488.9	418.4	149.7	143.1-156.6	216.8	175.6	165.1-186.7	201.6	132.4	123.8-141.4
Oral cavity and pharynx	1976-1980	19.6	8.6	6.9-10.5	12.6	13.3	10.1-17.3	7.0	5.4	3.8-7.6	9.0	3.9	2.8-5.2	6.2	6.4	4.3-9.2	2.8	2.2	1.2-3.7
	1981-1985	21.6	10.1	8.2-12.3	15.0	17.0	13.2-21.5	6.6	5.6	3.8-8.0	10.8	4.9	3.7-6.5	7.4	8.1	5.6-11.3	3.4	3.0	1.7-4.8
	1986-1990	27.2	12.1	10.1-14.4	17.0	17.5	13.9-21.8	10.2	8.2	6.0-10.9	9.2	3.8	2.8-5.1	6.2	6.5	4.3-9.3	3.0	2.2	1.2-3.7
	1991-1995	25.8	11.4	9.5-13.6	15.2	15.1	11.8-18.9	10.6	8.0	5.9-10.5	6.6	2.7	1.9-3.8	4.2	4.3	2.6-6.7	2.4	1.7	0.8-3.0
	1996-2000	21.2	9.3	7.6-11.3	14.6	14.4	11.2-18.1	6.6	5.2	3.5-7.4	7.0	2.9	2.0-4.0	3.8	3.8	2.3-6.0	3.2	2.3	1.3-3.8
	2001-2005	27.2	11.7	9.8-13.9	15.0	14.3	11.2-17.9	12.2	9.3	7.0-12.0	6.0	2.5	1.7-3.6	3.6	3.4	2.0-5.5	2.4	1.7	0.9-3.2
	2006-2010	29.0	11.8	9.9-13.9	21.4	19.5	15.9-23.6	7.6	5.4	3.7-7.5	6.6	2.5	1.7-3.5	4.0	3.5	2.1-5.5	2.6	1.5	0.8-2.8
	2011-2015	32.2	12.7	10.7-14.9	23.2	19.7	16.2-23.8	9.0	6.5	4.7-9.0	8.6	3.2	2.3-4.4	5.2	4.3	2.8-6.5	3.4	2.2	1.3-3.7
	2016-2020	35.4	14.3	12.2-16.8	26.4	22.9	19.0-27.5	9.0	6.7	4.8-9.1	8.0	3.0	2.1-4.1	6.2	5.2	3.5-7.5	1.8	1.2	0.5-2.5
Esophagus	1976-1980	10.0	4.5	3.3-5.9	7.0	7.4	5.1-10.3	3.0	2.3	1.3-3.9	8.0	3.6	2.6-4.9	6.2	6.6	4.4-9.5	1.8	1.4	0.6-2.7
	1981-1985	7.8	3.5	2.5-4.8	5.4	5.9	3.8-8.7	2.4	1.9	0.9-3.5	8.0	3.6	2.5-4.9	4.6	5.2	3.3-7.8	3.4	2.6	1.5-4.2
	1986-1990	12.6	5.5	4.2-7.1	10.0	10.7	7.9-14.3	2.6	1.8	1.0-3.2	12.0	5.4	4.1-7.0	9.6	10.9	7.9-14.6	2.4	1.8	0.9-3.3
	1991-1995	11.4	4.7	3.6-6.1	8.4	8.2	5.9-11.2	3.0	2.2	1.2-3.7	12.2	5.2	3.9-6.7	9.6	9.8	7.2-13.1	2.6	1.8	1.0-3.3
	1996-2000	12.4	5.1	3.9-6.6	7.6	7.5	5.3-10.3	4.8	3.3	2.1-5.0	10.6	4.2	3.2-5.6	6.6	6.6	4.5-9.3	4.0	2.6	1.6-4.2
	2001-2005	13.2	5.5	4.2-7.0	10.6	10.0	7.5-13.2	2.6	1.8	0.9-3.2	11.6	4.7	3.5-6.1	8.0	7.7	5.5-10.5	3.6	2.4	1.4-3.9
	2006-2010	16.6	6.7	5.3-8.3	13.4	12.5	9.7-16.0	3.2	2.3	1.3-3.9	14.0	5.5	4.3-7.0	11.4	10.5	8.0-13.7	2.6	1.8	0.9-3.2
	2011-2015	18.2	6.8	5.5-8.5	13.6	11.5	8.9-14.7	4.6	3.0	1.9-4.7	14.8	5.2	4.0-6.6	11.2	9.2	6.9-12.0	3.6	2.1	1.2-3.5
	2016-2020	15.6	5.9	4.7-7.5	12.0	9.8	7.4-12.7	3.6	2.7	1.6-4.5	11.2	3.9	3.0-5.2	9.8	7.6	5.6-10.2	1.4	0.9	0.4-2.1
Stomach	1976-1980	24.2	11.4	9.4-13.6	13.8	15.9	12.2-20.3	10.4	8.2	6.1-10.7	20.0	9.3	7.6-11.4	11.4	12.5	9.3-16.3	8.6	7.0	5.0-9.5
	1981-1985	20.0	8.8	7.1-10.7	11.6	12.1	9.0-15.7	8.4	6.2	4.4-8.4	13.6	6.1	4.7-7.8	6.8	7.2	4.9-10.2	6.8	4.9	3.4-7.0
	1986-1990	25.2	10.9	9.1-13.0	17.0	18.7	14.8-23.3	8.2	6.0	4.3-8.3	16.4	7.0	5.5-8.7	11.4	12.1	9.0-15.8	5.0	3.4	2.2-5.2
	1991-1995	20.6	8.5	6.9-10.3	11.6	12.3	9.3-16.0	9.0	5.9	4.3-7.9	12.2	5.0	3.8-6.5	6.2	6.4	4.3-9.2	6.0	4.0	2.7-5.9
	1996-2000	17.2	6.8	5.5-8.5	10.0	10.0	7.4-13.2	7.2	4.5	3.1-6.3	9.6	3.8	2.8-5.0	5.4	5.5	3.6-8.0	4.2	2.5	1.5-4.0
	2001-2005	14.4	5.9	4.6-7.5	9.2	9.2	6.7-12.2	5.2	3.7	2.4-5.5	7.6	3.0	2.1-4.1	3.6	3.5	2.1-5.5	4.0	2.6	1.6-4.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Broome**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	18.4	7.0	5.6-8.7	11.8	10.4	7.9-13.5	6.6	4.3	2.9-6.2	8.2	3.0	2.2-4.2	4.0	3.7	2.2-5.7	4.2	2.5	1.5-4.0
	2011-2015	17.2	6.6	5.2-8.2	11.6	10.0	7.6-13.1	5.6	3.7	2.5-5.6	7.0	2.6	1.8-3.7	4.2	3.8	2.3-5.9	2.8	1.7	0.9-3.0
	2016-2020	14.4	5.3	4.1-6.7	10.4	8.5	6.3-11.3	4.0	2.6	1.5-4.1	6.0	2.3	1.5-3.4	4.2	3.5	2.1-5.5	1.8	1.3	0.5-2.6
Colorectal	1976-1980	152.6	71.9	66.8-77.3	76.4	89.0	79.8-98.9	76.2	61.0	54.9-67.6	70.2	33.8	30.3-37.6	37.0	47.1	40.2-54.8	33.2	26.2	22.3-30.5
	1981-1985	162.0	72.8	67.8-78.1	77.2	88.9	79.9-98.7	84.8	63.1	57.1-69.5	77.0	34.8	31.4-38.5	38.2	45.5	38.9-52.7	38.8	28.2	24.4-32.6
	1986-1990	157.2	67.5	62.8-72.5	80.8	87.4	78.8-96.7	76.4	54.2	48.8-60.0	65.2	28.3	25.2-31.5	34.4	39.8	33.9-46.5	30.8	21.6	18.3-25.4
	1991-1995	138.4	57.5	53.2-62.0	65.8	68.4	61.0-76.4	72.6	49.3	44.2-54.8	55.2	22.7	20.1-25.6	24.2	25.1	20.8-30.2	31.0	19.9	16.9-23.5
	1996-2000	137.6	55.9	51.8-60.3	63.4	63.1	56.3-70.4	74.2	50.8	45.6-56.4	50.8	20.3	17.8-23.0	25.2	25.8	21.5-30.8	25.6	16.7	13.9-20.1
	2001-2005	134.4	53.8	49.7-58.1	68.0	65.2	58.5-72.6	66.4	45.2	40.4-50.6	41.6	16.0	13.9-18.4	21.6	21.2	17.4-25.6	20.0	12.7	10.2-15.6
	2006-2010	130.8	50.6	46.8-54.8	65.8	60.4	54.0-67.4	65.0	43.2	38.5-48.5	44.2	16.4	14.3-18.8	19.8	18.0	14.6-22.0	24.4	15.2	12.5-18.4
	2011-2015	95.4	37.1	33.7-40.7	46.0	40.2	35.0-45.9	49.4	34.0	29.7-38.9	36.8	13.5	11.6-15.7	17.0	15.2	12.1-19.0	19.8	11.9	9.6-14.8
2016-2020	92.6	35.7	32.4-39.4	50.0	42.4	37.1-48.4	42.6	30.0	25.9-34.7	33.6	12.4	10.5-14.5	16.0	13.3	10.5-16.8	17.6	11.6	9.1-14.6	
Colon excluding rectum	1976-1980	109.6	52.4	48.0-57.1	49.2	60.3	52.6-68.8	60.4	48.5	43.1-54.4	60.8	29.3	26.1-32.9	30.6	39.2	32.9-46.2	30.2	23.8	20.1-27.9
	1981-1985	116.2	52.3	48.1-56.8	52.8	62.2	54.6-70.6	63.4	46.9	41.8-52.4	68.0	30.7	27.5-34.2	33.0	39.1	33.1-45.9	35.0	25.6	22.0-29.8
	1986-1990	112.4	48.1	44.2-52.3	56.4	60.8	53.6-68.6	56.0	39.9	35.3-45.0	52.8	22.8	20.1-25.8	28.0	33.0	27.6-39.2	24.8	17.3	14.3-20.7
	1991-1995	101.6	42.1	38.5-46.0	46.4	48.4	42.2-55.2	55.2	37.6	33.2-42.5	46.0	18.9	16.5-21.5	20.6	21.7	17.6-26.4	25.4	16.4	13.6-19.6
	1996-2000	102.0	41.3	37.7-45.1	47.4	47.7	41.8-54.2	54.6	37.0	32.6-41.8	40.6	16.0	13.9-18.4	19.8	20.4	16.6-24.9	20.8	13.3	10.8-16.2
	2001-2005	93.4	36.8	33.5-40.3	44.8	43.1	37.6-49.1	48.6	32.1	28.0-36.5	32.4	12.4	10.5-14.5	16.2	16.0	12.7-19.9	16.2	10.1	8.0-12.7
	2006-2010	95.2	36.4	33.1-39.9	46.8	43.1	37.7-49.1	48.4	31.3	27.3-35.8	35.6	13.2	11.3-15.4	16.2	14.9	11.8-18.6	19.4	12.0	9.6-14.8
	2011-2015	67.8	25.7	22.9-28.7	30.4	26.2	22.1-30.8	37.4	25.1	21.5-29.3	29.0	10.5	8.8-12.5	12.0	10.7	8.1-13.9	17.0	10.3	8.1-12.9
2016-2020	63.6	24.3	21.5-27.3	31.8	27.4	23.1-32.2	31.8	21.7	18.2-25.7	26.4	9.6	8.0-11.5	12.2	10.2	7.8-13.3	14.2	9.2	7.0-11.9	
Rectum & rectosigmoid	1976-1980	43.0	19.5	16.9-22.3	27.2	28.6	23.8-34.1	15.8	12.5	9.9-15.7	9.4	4.5	3.3-6.0	6.4	8.0	5.3-11.4	3.0	2.4	1.3-4.0
	1981-1985	45.8	20.5	17.9-23.4	24.4	26.7	22.0-32.1	21.4	16.2	13.2-19.7	9.0	4.1	3.0-5.5	5.2	6.4	4.1-9.4	3.8	2.6	1.6-4.1
	1986-1990	44.8	19.4	16.9-22.2	24.4	26.6	21.9-32.0	20.4	14.3	11.6-17.5	12.4	5.5	4.2-7.0	6.4	6.8	4.6-9.6	6.0	4.3	2.9-6.2
	1991-1995	36.8	15.4	13.2-17.8	19.4	20.0	16.2-24.6	17.4	11.6	9.3-14.5	9.2	3.8	2.8-5.1	3.6	3.5	2.0-5.5	5.6	3.6	2.4-5.3
	1996-2000	35.6	14.7	12.6-17.0	16.0	15.4	12.2-19.2	19.6	13.8	11.1-16.9	10.2	4.3	3.2-5.7	5.4	5.4	3.5-7.8	4.8	3.5	2.2-5.3
	2001-2005	41.0	17.0	14.7-19.5	23.2	22.2	18.3-26.6	17.8	13.2	10.5-16.4	9.2	3.7	2.7-4.9	5.4	5.2	3.4-7.6	3.8	2.6	1.5-4.1
	2006-2010	35.6	14.2	12.2-16.6	19.0	17.3	13.9-21.2	16.6	11.9	9.4-15.0	8.6	3.2	2.3-4.4	3.6	3.1	1.8-5.0	5.0	3.2	2.0-5.0
	2011-2015	27.6	11.4	9.5-13.6	15.6	14.0	11.0-17.6	12.0	8.9	6.6-11.8	7.8	3.0	2.1-4.2	5.0	4.5	2.9-6.8	2.8	1.6	0.9-3.0
2016-2020	29.0	11.5	9.6-13.6	18.2	15.1	12.0-18.8	10.8	8.3	6.2-11.1	7.2	2.7	1.9-3.9	3.8	3.1	1.8-5.0	3.4	2.4	1.3-4.1	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Broome**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	6.0	2.8	1.9-4.0	3.8	4.1	2.4-6.5	2.2	1.8	0.9-3.2	3.6	1.7	1.0-2.7	2.2	2.4	1.2-4.4	1.4	1.1	0.4-2.4
	1981-1985	5.2	2.2	1.4-3.3	3.2	3.6	2.0-5.9	2.0	1.5	0.7-2.8	6.8	3.0	2.1-4.3	4.4	5.5	3.4-8.4	2.4	1.8	0.9-3.2
	1986-1990	5.6	2.4	1.6-3.5	4.0	4.1	2.5-6.4	1.6	1.1	0.5-2.2	6.8	2.9	2.0-4.1	4.4	5.2	3.2-7.9	2.4	1.7	0.9-3.1
	1991-1995	7.8	3.4	2.4-4.6	5.0	5.1	3.3-7.6	2.8	2.2	1.2-3.8	7.0	2.9	2.0-4.1	4.2	4.4	2.7-6.8	2.8	1.9	1.0-3.3
	1996-2000	11.4	4.8	3.6-6.2	6.4	6.2	4.2-8.8	5.0	3.6	2.3-5.4	10.0	4.1	3.0-5.5	5.6	5.6	3.7-8.1	4.4	2.9	1.8-4.5
	2001-2005	11.8	5.0	3.8-6.5	8.4	8.0	5.7-10.8	3.4	2.6	1.5-4.3	10.8	4.5	3.4-5.9	7.6	7.3	5.1-10.0	3.2	2.4	1.4-4.0
	2006-2010	14.0	5.6	4.4-7.2	10.6	9.2	6.9-12.1	3.4	2.6	1.5-4.3	9.6	3.7	2.7-5.0	6.4	5.6	3.8-7.9	3.2	2.2	1.2-3.7
	2011-2015	14.2	5.7	4.4-7.3	11.2	9.9	7.4-13.1	3.0	2.0	1.1-3.5	14.0	5.2	4.0-6.6	9.6	7.8	5.7-10.5	4.4	2.9	1.8-4.6
2016-2020	24.0	8.7	7.2-10.5	17.2	13.3	10.6-16.6	6.8	4.6	3.2-6.6	15.4	5.6	4.4-7.1	10.4	8.2	6.1-11.0	5.0	3.3	2.1-5.0	
Pancreas	1976-1980	22.6	10.9	8.9-13.1	12.6	14.9	11.3-19.3	10.0	8.0	6.0-10.7	23.8	11.4	9.5-13.7	14.0	16.5	12.7-21.0	9.8	8.0	5.9-10.6
	1981-1985	24.0	10.8	8.9-12.9	11.2	12.8	9.5-16.8	12.8	9.7	7.4-12.4	20.8	9.3	7.5-11.2	10.0	11.6	8.4-15.4	10.8	8.0	6.0-10.6
	1986-1990	30.2	13.4	11.3-15.8	13.0	15.5	11.9-19.9	17.2	12.3	9.8-15.3	24.0	10.9	9.0-13.1	9.8	12.5	9.2-16.6	14.2	10.5	8.2-13.4
	1991-1995	29.2	12.0	10.2-14.2	15.2	15.4	12.1-19.3	14.0	9.6	7.4-12.2	28.2	11.5	9.7-13.6	14.8	14.9	11.7-18.8	13.4	9.0	6.9-11.5
	1996-2000	26.8	11.1	9.3-13.2	14.6	14.8	11.6-18.7	12.2	8.3	6.3-10.7	25.6	10.3	8.6-12.3	12.6	12.9	9.9-16.6	13.0	8.6	6.6-11.1
	2001-2005	28.2	11.3	9.5-13.4	15.6	15.0	11.9-18.8	12.6	8.6	6.5-11.1	26.0	10.3	8.6-12.2	14.0	13.4	10.4-17.0	12.0	7.7	5.9-10.1
	2006-2010	27.0	10.3	8.6-12.3	12.4	11.3	8.6-14.6	14.6	9.2	7.2-11.8	23.2	8.5	7.0-10.3	9.0	8.2	5.9-11.0	14.2	8.6	6.7-11.1
	2011-2015	38.0	14.1	12.1-16.4	17.0	14.9	11.8-18.5	21.0	13.2	10.7-16.2	32.2	11.7	9.9-13.7	14.4	12.5	9.7-15.8	17.8	10.9	8.7-13.6
2016-2020	45.8	16.1	14.0-18.5	22.0	17.2	14.0-20.9	23.8	15.2	12.5-18.5	37.0	12.7	10.9-14.8	18.8	14.5	11.7-17.9	18.2	11.1	8.9-13.8	
Larynx	1976-1980	10.2	4.5	3.3-6.0	8.6	9.0	6.4-12.4	1.6	1.4	0.6-2.8	2.8	1.3	0.7-2.2	2.6	3.1	1.6-5.3	0.2	0.2	0.0-1.0
	1981-1985	14.2	6.4	5.0-8.1	11.2	11.6	8.7-15.3	3.0	2.5	1.4-4.2	4.4	1.9	1.2-2.9	3.2	3.4	1.9-5.6	1.2	0.8	0.3-1.9
	1986-1990	15.2	6.3	5.0-8.0	11.4	11.2	8.4-14.6	3.8	2.7	1.6-4.2	3.4	1.6	0.9-2.5	2.4	3.0	1.5-5.3	1.0	0.7	0.2-1.8
	1991-1995	11.6	5.0	3.8-6.5	8.4	8.7	6.2-11.8	3.2	2.4	1.3-3.9	4.0	1.7	1.0-2.6	3.4	3.2	1.9-5.2	0.6	0.4	0.1-1.2
	1996-2000	9.6	4.0	3.0-5.4	6.8	6.6	4.5-9.2	2.8	2.1	1.1-3.6	4.4	1.9	1.2-2.9	3.6	3.6	2.1-5.8	0.8	0.5	0.1-1.4
	2001-2005	10.2	4.4	3.2-5.8	8.4	7.9	5.7-10.7	1.8	1.3	0.6-2.6	2.8	1.1	0.6-1.9	2.0	1.9	0.9-3.6	0.8	0.5	0.1-1.5
	2006-2010	10.8	4.4	3.3-5.8	9.2	8.6	6.3-11.5	1.6	1.1	0.5-2.4	3.0	1.1	0.6-1.9	2.4	2.2	1.1-3.9	0.6	0.4	0.1-1.4
	2011-2015	9.8	4.0	3.0-5.4	6.4	5.6	3.8-8.0	3.4	2.7	1.5-4.4	3.0	1.2	0.6-2.0	2.2	1.8	0.9-3.3	0.8	0.7	0.2-2.0
2016-2020	5.4	1.9	1.2-2.8	4.6	3.3	2.1-5.1	0.8	0.6	0.1-1.7	1.8	0.6	0.3-1.3	1.0	0.8	0.3-2.0	0.8	0.5	0.1-1.5	
Lung and bronchus	1976-1980	129.4	59.1	54.6-64.0	89.8	99.8	90.3-109.9	39.6	31.7	27.4-36.6	100.0	45.7	41.7-49.9	71.6	80.8	72.3-90.1	28.4	22.3	18.7-26.4
	1981-1985	141.2	62.2	57.6-67.1	91.4	98.0	88.8-107.7	49.8	38.6	33.8-43.8	110.0	48.1	44.1-52.4	75.4	81.9	73.5-91.0	34.6	26.4	22.5-30.8
	1986-1990	156.8	67.6	62.9-72.6	96.4	101.4	92.2-111.2	60.4	45.4	40.3-51.0	123.2	53.1	49.0-57.6	76.0	80.7	72.5-89.6	47.2	35.1	30.7-40.1
	1991-1995	170.0	70.5	65.8-75.5	98.4	97.1	88.5-106.2	71.6	51.8	46.4-57.6	130.8	54.3	50.2-58.7	80.2	82.1	74.1-90.7	50.6	36.3	31.8-41.2
	1996-2000	184.8	76.9	72.0-82.0	102.2	100.1	91.6-109.2	82.6	60.8	55.0-67.2	138.8	57.0	52.8-61.5	78.2	76.8	69.3-84.9	60.6	43.1	38.3-48.4
	2001-2005	176.4	71.6	66.9-76.6	93.8	89.3	81.4-97.8	82.6	58.8	53.1-64.9	127.4	50.4	46.5-54.5	68.0	65.2	58.4-72.5	59.4	40.4	35.8-45.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Broome**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	160.6	62.5	58.2-67.1	85.4	76.7	69.6-84.5	75.2	52.1	46.8-57.8	125.0	48.2	44.4-52.2	65.4	59.6	53.3-66.5	59.6	40.3	35.7-45.3
	2011-2015	167.8	64.4	60.0-69.1	83.0	71.0	64.2-78.4	84.8	60.2	54.4-66.5	109.2	40.8	37.4-44.4	56.8	48.5	43.0-54.7	52.4	35.3	31.0-40.0
	2016-2020	170.2	61.6	57.4-66.0	86.4	68.4	61.9-75.4	83.8	56.6	51.1-62.7	104.2	37.3	34.1-40.8	52.8	42.0	37.0-47.6	51.4	34.2	30.0-38.9
Melanoma of the skin	1976-1980	15.4	7.8	6.1-9.8	8.4	9.7	6.9-13.2	7.0	6.5	4.4-9.1	5.2	2.8	1.8-4.1	3.6	4.2	2.4-6.7	1.6	1.6	0.7-3.2
	1981-1985	19.0	9.2	7.4-11.3	10.2	11.0	8.1-14.6	8.8	8.0	5.7-10.8	5.2	2.2	1.5-3.3	3.0	3.1	1.7-5.2	2.2	1.7	0.8-3.1
	1986-1990	23.0	10.9	9.0-13.2	12.0	12.5	9.4-16.2	11.0	9.9	7.4-13.0	5.8	2.8	1.8-4.0	2.8	2.9	1.5-5.0	3.0	2.8	1.6-4.7
	1991-1995	24.4	10.8	8.9-12.9	14.0	14.0	10.9-17.8	10.4	7.9	5.8-10.5	6.6	2.8	1.9-3.9	4.2	4.2	2.6-6.5	2.4	1.8	0.9-3.2
	1996-2000	33.8	15.2	13.0-17.7	20.8	21.1	17.2-25.6	13.0	11.3	8.7-14.5	7.4	3.2	2.2-4.4	5.0	4.9	3.2-7.3	2.4	1.9	0.9-3.4
	2001-2005	40.6	17.7	15.3-20.3	24.2	23.4	19.4-27.9	16.4	13.7	10.8-17.2	8.2	3.4	2.4-4.6	4.8	4.7	3.0-7.0	3.4	2.6	1.5-4.3
	2006-2010	50.8	21.5	18.8-24.4	31.4	29.2	24.8-34.3	19.4	16.0	12.8-19.8	6.8	2.6	1.8-3.7	4.8	4.4	2.8-6.6	2.0	1.4	0.6-2.7
	2011-2015	65.0	26.7	23.8-30.0	38.8	33.4	28.8-38.6	26.2	21.9	18.1-26.3	4.6	1.7	1.1-2.7	2.4	2.1	1.1-3.8	2.2	1.3	0.6-2.6
2016-2020	63.2	25.4	22.5-28.6	37.0	31.4	26.9-36.6	26.2	20.8	17.1-25.1	4.6	1.9	1.1-3.0	3.2	3.1	1.7-5.2	1.4	0.9	0.3-2.0	
Female breast	1976-1980							113.4	98.5	90.3-107.3							47.0	40.0	34.9-45.7
	1981-1985							127.6	108.0	99.5-117.1							45.0	37.2	32.4-42.6
	1986-1990							162.6	132.7	123.4-142.6							40.6	31.5	27.2-36.3
	1991-1995							163.8	130.6	121.5-140.2							44.4	33.2	28.9-38.1
	1996-2000							169.0	135.2	126.0-144.9							38.4	27.1	23.3-31.5
	2001-2005							155.6	121.9	113.3-131.0							33.6	23.4	19.9-27.4
	2006-2010							162.6	125.2	116.4-134.5							29.6	19.7	16.5-23.4
	2011-2015							177.2	137.7	128.3-147.6							27.8	19.7	16.4-23.5
2016-2020							184.4	145.3	135.4-155.8							23.6	16.2	13.1-19.8	
Cervix uteri	1976-1980							14.6	13.7	10.7-17.4							5.6	5.0	3.3-7.3
	1981-1985							12.0	10.9	8.2-14.1							3.2	2.7	1.5-4.4
	1986-1990							17.0	15.1	12.0-18.8							3.8	3.2	1.9-5.0
	1991-1995							10.2	9.1	6.7-12.0							4.2	3.2	2.0-5.0
	1996-2000							11.0	10.3	7.7-13.4							2.8	2.3	1.2-4.0
	2001-2005							7.2	6.3	4.4-8.9							2.6	2.3	1.2-4.0
	2006-2010							7.8	7.0	4.9-9.7							2.2	1.6	0.8-2.9
	2011-2015							6.8	5.7	3.9-8.2							2.8	2.0	1.1-3.5
2016-2020							7.8	8.4	5.8-11.7							2.6	2.2	1.1-3.9	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Broome**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							33.2	28.0	23.8-32.8							4.6	3.5	2.2-5.4
	1981-1985							34.8	29.0	24.7-33.8							6.6	4.9	3.4-7.0
	1986-1990							41.0	33.2	28.6-38.2							5.8	4.1	2.7-6.0
	1991-1995							36.0	28.8	24.7-33.5							4.4	3.0	1.8-4.6
	1996-2000							35.2	28.5	24.3-33.1							4.4	2.8	1.8-4.4
	2001-2005							39.0	30.9	26.6-35.7							7.6	5.0	3.5-7.0
	2006-2010							44.0	34.4	29.9-39.5							6.0	3.9	2.6-5.7
	2011-2015							48.2	36.4	31.7-41.6							6.2	4.2	2.8-6.1
	2016-2020							44.2	33.7	29.1-38.8							7.8	5.3	3.7-7.5
Ovary	1976-1980							20.8	17.8	14.5-21.7							13.2	10.5	8.1-13.4
	1981-1985							21.8	18.3	14.9-22.2							15.0	11.3	8.9-14.3
	1986-1990							27.6	22.3	18.6-26.5							16.4	12.1	9.5-15.1
	1991-1995							26.0	19.8	16.5-23.7							14.8	10.1	7.8-12.7
	1996-2000							22.4	16.1	13.2-19.6							15.6	10.5	8.3-13.3
	2001-2005							19.0	14.7	11.9-18.1							16.4	11.6	9.1-14.5
	2006-2010							19.0	14.1	11.3-17.4							15.2	10.5	8.2-13.2
	2011-2015							15.4	12.3	9.6-15.6							9.0	5.6	4.0-7.6
	2016-2020							20.2	16.1	12.9-19.8							11.4	7.8	5.8-10.3
Prostate	1976-1980				76.0	96.4	86.4-107.1							21.2	30.4	24.7-37.0			
	1981-1985				78.8	94.3	84.8-104.5							23.8	32.9	27.0-39.6			
	1986-1990				105.0	118.4	108.0-129.4							25.6	31.7	26.2-37.9			
	1991-1995				188.2	185.2	173.3-197.6							30.6	36.0	30.4-42.4			
	1996-2000				160.0	154.5	144.0-165.7							30.4	32.0	27.1-37.6			
	2001-2005				204.8	192.9	181.2-205.1							25.0	25.1	20.9-30.0			
	2006-2010				206.6	182.6	171.5-194.2							25.6	24.0	20.0-28.5			
	2011-2015				133.2	106.5	98.5-115.1							19.4	17.0	13.7-20.8			
	2016-2020				167.8	125.4	116.9-134.5							26.6	21.2	17.7-25.2			
Testis	1976-1980				4.2	3.7	2.2-5.8							0.4	0.4	0.0-1.5			
	1981-1985				5.8	5.8	3.8-8.5							1.0	0.8	0.3-2.1			
	1986-1990				7.4	6.6	4.6-9.3							0.6	0.5	0.1-1.6			
	1991-1995				7.2	7.1	4.9-9.8							0.2	0.2	0.0-1.0			
	1996-2000				6.2	6.5	4.4-9.3							0.4	0.4	0.1-1.5			
	2001-2005				6.6	7.4	5.0-10.3							0.0	0.0	0.0-0.8			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Broome**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				6.2	6.5	4.3-9.3							0.2	0.3	0.0-1.4			
	2011-2015				6.0	6.7	4.4-9.6							1.2	1.3	0.5-3.0			
	2016-2020				7.8	9.5	6.7-13.1							0.2	0.3	0.0-1.4			
Urinary bladder (incl. in situ)	1976-1980	45.8	21.9	19.1-25.0	32.6	40.8	34.4-47.9	13.2	10.7	8.3-13.7	13.8	6.8	5.3-8.6	9.6	13.4	9.7-18.0	4.2	3.3	2.1-5.2
	1981-1985	46.8	21.1	18.5-24.1	33.6	39.7	33.7-46.5	13.2	9.9	7.6-12.7	13.4	6.3	4.9-8.1	10.2	14.1	10.3-18.6	3.2	2.4	1.3-3.9
	1986-1990	54.0	23.6	20.8-26.6	40.4	44.4	38.3-51.2	13.6	9.7	7.5-12.4	15.6	6.8	5.4-8.5	10.8	13.6	10.1-17.9	4.8	3.3	2.1-5.0
	1991-1995	56.6	23.5	20.8-26.5	40.6	41.5	35.8-47.7	16.0	11.6	9.1-14.5	11.4	4.7	3.6-6.1	8.4	9.5	6.8-12.9	3.0	2.0	1.1-3.4
	1996-2000	65.0	26.5	23.7-29.6	48.0	47.8	41.9-54.3	17.0	12.0	9.5-14.9	12.6	4.9	3.8-6.3	9.8	10.0	7.4-13.3	2.8	1.8	1.0-3.1
	2001-2005	57.2	22.7	20.1-25.5	44.0	42.4	36.9-48.4	13.2	9.4	7.2-12.1	10.6	4.1	3.0-5.4	7.2	7.0	4.9-9.7	3.4	2.0	1.2-3.4
	2006-2010	65.4	25.1	22.5-28.1	48.6	44.5	39.0-50.5	16.8	11.4	9.0-14.3	14.0	4.9	3.8-6.3	10.8	10.2	7.6-13.3	3.2	1.9	1.1-3.3
	2011-2015	70.4	26.3	23.5-29.3	54.4	47.2	41.6-53.2	16.0	11.4	9.0-14.3	14.2	4.9	3.8-6.3	9.4	8.2	6.0-11.0	4.8	2.9	1.8-4.4
2016-2020	64.4	23.1	20.6-25.9	49.4	39.8	34.9-45.2	15.0	10.1	7.8-12.9	12.0	3.8	2.9-5.0	9.0	7.2	5.2-9.7	3.0	1.7	0.9-3.0	
Kidney and renal pelvis	1976-1980	17.6	7.9	6.3-9.8	10.0	10.1	7.4-13.4	7.6	6.2	4.3-8.5	9.6	4.2	3.1-5.7	5.4	5.7	3.7-8.5	4.2	3.3	2.0-5.2
	1981-1985	18.0	8.0	6.4-9.9	11.6	12.7	9.5-16.6	6.4	4.7	3.2-6.7	9.2	4.1	3.0-5.5	6.2	7.5	5.0-10.7	3.0	2.1	1.2-3.6
	1986-1990	25.6	11.5	9.6-13.8	13.8	15.3	11.8-19.5	11.8	9.1	6.8-11.8	9.4	4.1	3.0-5.5	5.0	5.5	3.5-8.1	4.4	3.0	1.9-4.6
	1991-1995	27.0	11.5	9.6-13.6	15.0	15.3	11.9-19.2	12.0	8.9	6.8-11.6	9.2	3.8	2.8-5.1	5.4	5.5	3.6-8.0	3.8	2.6	1.6-4.2
	1996-2000	31.0	13.1	11.1-15.4	19.4	19.4	15.7-23.7	11.6	7.9	5.9-10.3	12.4	5.1	3.9-6.6	6.4	6.3	4.3-9.0	6.0	4.0	2.6-5.8
	2001-2005	32.4	13.5	11.5-15.8	19.0	18.2	14.7-22.2	13.4	10.2	7.8-13.1	9.6	3.7	2.7-4.9	5.0	4.8	3.1-7.1	4.6	2.9	1.8-4.6
	2006-2010	40.6	17.0	14.7-19.6	24.8	23.0	19.1-27.5	15.8	12.5	9.8-15.8	11.4	4.3	3.3-5.7	7.2	6.6	4.6-9.2	4.2	2.7	1.7-4.4
	2011-2015	38.4	15.4	13.2-17.9	25.0	21.7	17.9-26.0	13.4	10.2	7.8-13.2	12.0	4.4	3.4-5.8	7.4	6.4	4.5-8.9	4.6	3.0	1.9-4.7
2016-2020	49.4	19.4	16.9-22.2	31.4	27.0	22.7-31.9	18.0	12.4	9.7-15.6	10.2	3.4	2.5-4.5	6.2	4.7	3.2-6.8	4.0	2.3	1.3-3.8	
Brain and other central nervous system	1976-1980	10.6	5.0	3.7-6.6	5.4	5.4	3.5-8.0	5.2	4.5	2.9-6.7	6.0	2.9	1.9-4.2	2.6	2.5	1.3-4.4	3.4	3.1	1.8-5.1
	1981-1985	15.0	7.2	5.6-9.0	8.8	9.0	6.5-12.2	6.2	5.5	3.7-8.0	11.2	5.3	4.0-6.9	6.6	7.2	4.9-10.2	4.6	3.6	2.3-5.5
	1986-1990	16.6	7.6	6.1-9.5	7.0	7.2	5.0-10.2	9.6	8.0	5.9-10.8	8.8	3.7	2.7-5.1	4.4	4.8	2.9-7.3	4.4	3.1	1.9-4.8
	1991-1995	21.2	9.5	7.8-11.6	10.0	10.1	7.5-13.4	11.2	9.2	6.9-12.0	12.4	5.4	4.1-7.0	5.2	5.4	3.5-8.0	7.2	5.8	4.0-8.1
	1996-2000	18.2	8.3	6.7-10.2	9.8	9.8	7.2-13.0	8.4	6.7	4.7-9.1	13.0	5.8	4.5-7.4	6.4	6.4	4.4-9.0	6.6	5.1	3.5-7.3
	2001-2005	16.4	7.4	5.9-9.2	8.8	8.6	6.3-11.6	7.6	6.3	4.4-8.8	10.6	4.6	3.4-6.0	6.2	5.9	4.0-8.5	4.4	3.4	2.1-5.2
	2006-2010	17.6	7.8	6.2-9.7	10.4	9.8	7.2-12.9	7.2	6.2	4.2-8.7	11.2	4.5	3.4-5.9	6.6	6.0	4.1-8.5	4.6	3.1	1.9-4.8
	2011-2015	12.0	5.0	3.8-6.6	6.0	5.4	3.6-7.8	6.0	4.7	3.0-6.9	9.8	4.0	2.9-5.4	4.0	3.5	2.1-5.6	5.8	4.4	2.8-6.5
2016-2020	17.6	7.9	6.2-9.9	9.6	8.9	6.5-12.1	8.0	6.8	4.7-9.6	9.8	4.1	3.0-5.5	6.2	5.8	3.9-8.4	3.6	2.6	1.5-4.4	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Broome**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	5.6	2.8	1.8-4.1	2.2	2.2	1.1-4.0	3.4	3.4	1.9-5.5	0.4	0.2	0.0-0.7	0.0	0.0	0.0-0.9	0.4	0.3	0.0-1.3
	1981-1985	6.8	3.1	2.1-4.4	2.2	2.2	1.1-4.1	4.6	3.8	2.4-5.8	0.4	0.1	0.0-0.6	0.2	0.2	0.0-1.1	0.2	0.1	0.0-0.9
	1986-1990	9.6	4.9	3.6-6.5	2.8	3.0	1.6-5.0	6.8	6.8	4.6-9.5	1.4	0.6	0.3-1.3	1.0	1.3	0.4-3.0	0.4	0.3	0.0-1.1
	1991-1995	8.6	4.0	2.9-5.4	2.2	2.2	1.1-4.0	6.4	5.8	3.9-8.2	0.4	0.2	0.0-0.6	0.2	0.2	0.0-1.2	0.2	0.1	0.0-0.9
	1996-2000	14.4	7.0	5.5-8.8	3.8	3.9	2.3-6.1	10.6	10.2	7.6-13.4	1.2	0.5	0.2-1.1	0.4	0.4	0.0-1.6	0.8	0.5	0.1-1.5
	2001-2005	23.2	11.2	9.2-13.5	5.2	5.0	3.3-7.4	18.0	17.2	13.8-21.2	1.2	0.5	0.2-1.1	0.4	0.4	0.0-1.4	0.8	0.6	0.2-1.6
	2006-2010	44.0	21.4	18.6-24.5	10.8	10.7	8.0-14.0	33.2	31.9	27.1-37.4	1.0	0.3	0.1-0.9	0.4	0.3	0.0-1.4	0.6	0.4	0.1-1.3
	2011-2015	42.8	21.6	18.7-24.9	11.6	11.0	8.3-14.3	31.2	32.2	27.0-37.9	1.0	0.4	0.1-0.9	0.2	0.1	0.0-1.0	0.8	0.6	0.2-1.6
2016-2020	33.4	16.8	14.2-19.8	8.4	8.3	5.8-11.5	25.0	25.4	20.8-30.6	1.6	0.5	0.2-1.1	1.0	0.7	0.2-1.8	0.6	0.4	0.1-1.3	
Hodgkin lymphoma	1976-1980	9.0	4.1	2.9-5.5	4.2	4.5	2.7-7.0	4.8	3.9	2.5-5.9	2.4	1.2	0.6-2.1	1.8	2.0	0.9-3.9	0.6	0.5	0.1-1.5
	1981-1985	6.6	2.9	2.0-4.1	4.4	4.1	2.5-6.4	2.2	1.7	0.8-3.2	2.2	1.1	0.5-2.0	1.4	1.5	0.6-3.2	0.8	0.7	0.2-1.9
	1986-1990	7.8	3.6	2.5-5.0	5.6	5.2	3.4-7.6	2.2	2.0	1.0-3.7	1.6	0.7	0.3-1.5	1.4	1.4	0.5-2.9	0.2	0.1	0.0-0.9
	1991-1995	7.6	3.5	2.4-4.8	4.4	4.3	2.7-6.6	3.2	2.8	1.6-4.6	0.4	0.2	0.0-0.7	0.2	0.2	0.0-1.1	0.2	0.2	0.0-1.1
	1996-2000	7.4	3.6	2.5-5.0	4.4	4.5	2.8-6.8	3.0	2.8	1.6-4.7	1.4	0.7	0.3-1.4	0.6	0.6	0.1-1.7	0.8	0.7	0.2-1.9
	2001-2005	7.2	3.4	2.4-4.7	3.4	3.4	2.0-5.5	3.8	3.3	2.0-5.2	0.8	0.4	0.1-0.9	0.0	0.0	0.0-0.8	0.8	0.7	0.2-1.8
	2006-2010	4.8	2.4	1.5-3.6	3.2	3.0	1.7-4.9	1.6	1.8	0.7-3.5	0.6	0.2	0.0-0.7	0.0	0.0	0.0-0.8	0.6	0.4	0.1-1.3
	2011-2015	5.2	2.5	1.6-3.7	3.0	3.1	1.7-5.2	2.2	1.9	0.9-3.6	1.0	0.4	0.1-1.0	0.4	0.4	0.0-1.4	0.6	0.4	0.1-1.4
2016-2020	6.4	3.2	2.1-4.7	3.2	3.0	1.6-5.1	3.2	3.4	1.9-5.7	0.8	0.3	0.1-1.0	0.6	0.6	0.1-1.8	0.2	0.1	0.0-1.0	
Non-Hodgkin lymphomas	1976-1980	25.6	12.1	10.1-14.5	11.0	13.1	9.7-17.2	14.6	11.9	9.3-15.1	14.6	6.9	5.4-8.6	7.6	8.8	6.1-12.3	7.0	5.7	4.0-8.0
	1981-1985	31.6	14.0	11.9-16.4	17.4	17.9	14.2-22.2	14.2	10.7	8.3-13.6	17.4	7.8	6.2-9.7	9.4	10.5	7.6-14.0	8.0	5.9	4.2-8.1
	1986-1990	43.4	19.2	16.7-22.0	22.4	23.5	19.2-28.4	21.0	15.8	12.9-19.3	18.2	7.8	6.3-9.6	10.6	11.7	8.6-15.4	7.6	5.3	3.8-7.4
	1991-1995	45.0	19.6	17.1-22.4	25.8	26.8	22.2-31.9	19.2	14.6	11.8-18.0	20.4	8.7	7.1-10.6	12.0	13.0	9.8-16.8	8.4	6.1	4.3-8.3
	1996-2000	45.4	19.4	16.9-22.1	22.2	22.1	18.1-26.6	23.2	17.2	14.1-20.8	21.8	8.8	7.2-10.7	11.4	11.2	8.5-14.6	10.4	7.0	5.2-9.3
	2001-2005	45.2	18.6	16.2-21.2	22.6	21.6	17.8-26.0	22.6	16.4	13.4-19.9	16.8	6.5	5.1-8.1	10.2	10.0	7.4-13.1	6.6	4.1	2.8-5.9
	2006-2010	52.0	20.5	18.0-23.2	27.6	26.0	21.8-30.8	24.4	15.7	12.9-18.9	16.4	6.0	4.7-7.5	8.6	7.9	5.7-10.6	7.8	4.6	3.2-6.4
	2011-2015	48.6	19.1	16.7-21.8	26.6	23.8	19.9-28.4	22.0	15.2	12.4-18.6	13.8	4.8	3.8-6.2	7.4	6.2	4.4-8.7	6.4	3.9	2.6-5.6
2016-2020	50.4	19.6	17.1-22.4	26.6	22.9	19.0-27.4	23.8	17.0	13.9-20.7	14.2	5.0	3.9-6.4	7.4	6.1	4.3-8.5	6.8	4.4	3.0-6.5	
Myeloma	1976-1980	10.0	4.8	3.5-6.3	4.2	5.5	3.3-8.4	5.8	4.4	3.0-6.4	5.6	2.7	1.8-3.9	0.8	1.0	0.3-2.6	4.8	3.7	2.4-5.5
	1981-1985	15.6	7.0	5.5-8.7	7.8	9.2	6.4-12.8	7.8	5.6	4.0-7.7	9.4	4.4	3.2-5.8	5.4	6.7	4.3-9.8	4.0	2.9	1.8-4.6
	1986-1990	18.2	7.8	6.3-9.6	9.2	10.3	7.4-13.9	9.0	6.4	4.7-8.7	9.8	4.1	3.0-5.4	6.0	7.1	4.7-10.3	3.8	2.6	1.6-4.1
	1991-1995	15.4	6.3	5.0-7.9	7.4	7.3	5.1-10.1	8.0	5.7	4.0-7.8	11.4	4.6	3.5-6.0	6.4	6.6	4.5-9.3	5.0	3.3	2.2-5.0
	1996-2000	15.6	6.3	5.0-7.9	7.2	7.5	5.2-10.4	8.4	6.0	4.3-8.2	12.0	4.7	3.6-6.0	5.6	5.7	3.8-8.3	6.4	4.3	2.9-6.1
	2001-2005	14.4	5.6	4.4-7.2	6.4	6.3	4.3-8.9	8.0	5.3	3.7-7.3	8.2	3.1	2.2-4.3	3.6	3.5	2.0-5.5	4.6	3.0	1.9-4.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Broome**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	16.2	6.3	5.0-7.9	7.6	6.9	4.9-9.5	8.6	6.0	4.3-8.2	6.2	2.3	1.5-3.3	2.6	2.4	1.3-4.1	3.6	2.2	1.3-3.7
	2011-2015	17.0	6.6	5.2-8.2	10.0	8.5	6.2-11.3	7.0	5.0	3.4-7.2	6.4	2.1	1.5-3.1	3.4	2.9	1.7-4.7	3.0	1.6	0.8-2.8
	2016-2020	22.0	8.6	7.0-10.5	11.8	9.4	7.1-12.2	10.2	7.9	5.7-10.7	8.8	3.0	2.2-4.1	3.8	3.0	1.8-4.8	5.0	3.0	1.9-4.6
Leukemias	1976-1980	33.2	16.0	13.6-18.7	19.8	22.9	18.3-28.1	13.4	11.2	8.7-14.4	20.0	9.8	8.0-12.0	11.8	13.9	10.3-18.3	8.2	7.3	5.2-10.0
	1981-1985	30.8	14.2	12.0-16.7	17.6	20.2	16.0-25.1	13.2	10.7	8.2-13.7	18.4	8.4	6.8-10.4	11.2	13.4	10.0-17.6	7.2	5.8	4.0-8.1
	1986-1990	31.2	14.0	11.9-16.4	15.8	18.7	14.6-23.5	15.4	11.4	8.9-14.3	19.0	8.5	6.8-10.4	9.8	11.1	8.1-14.8	9.2	6.6	4.8-8.9
	1991-1995	37.0	16.2	13.9-18.8	20.8	21.6	17.6-26.3	16.2	12.5	9.9-15.7	17.6	7.4	6.0-9.2	7.6	8.4	5.9-11.6	10.0	7.3	5.4-9.8
	1996-2000	35.2	14.9	12.7-17.3	18.8	18.9	15.3-23.2	16.4	12.1	9.6-15.2	22.8	9.1	7.5-11.0	11.2	11.4	8.6-14.9	11.6	7.8	5.9-10.2
	2001-2005	33.8	13.8	11.8-16.1	18.4	18.0	14.5-22.1	15.4	11.2	8.8-14.2	18.6	7.3	5.8-8.9	13.0	12.9	10.0-16.5	5.6	3.8	2.5-5.6
	2006-2010	35.6	14.3	12.2-16.7	21.0	20.0	16.3-24.3	14.6	9.8	7.6-12.6	18.6	7.2	5.8-8.9	10.8	10.2	7.6-13.3	7.8	5.0	3.5-7.1
	2011-2015	42.6	16.9	14.7-19.5	26.2	23.0	19.2-27.4	16.4	12.3	9.6-15.5	21.4	7.5	6.1-9.2	12.0	10.3	7.8-13.3	9.4	5.8	4.2-7.8
2016-2020	41.2	15.7	13.5-18.1	22.6	19.2	15.7-23.3	18.6	12.7	10.1-15.9	15.0	5.5	4.3-7.0	7.8	6.8	4.8-9.5	7.2	4.6	3.1-6.6	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cattaraugus**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	321.6	396.0	376.5-416.2	160.2	459.2	426.7-493.5	161.4	363.3	338.1-389.9	185.2	227.1	212.4-242.4	93.6	272.9	247.7-299.8	91.6	199.3	181.2-218.7
	1981-1985	344.2	409.1	389.6-429.3	180.2	507.0	473.1-542.5	164.0	349.2	325.0-374.7	185.6	218.4	204.3-233.1	95.2	275.7	250.5-302.7	90.4	186.6	169.4-205.1
	1986-1990	347.8	405.8	386.6-425.7	176.6	484.5	451.9-518.7	171.2	363.6	338.9-389.6	178.2	207.0	193.5-221.3	95.2	266.1	241.9-292.0	83.0	168.3	152.1-185.7
	1991-1995	392.0	437.3	418.0-457.3	208.0	544.4	510.9-579.4	184.0	374.2	350.0-399.8	180.6	200.2	187.2-213.8	93.8	257.3	233.8-282.5	86.8	170.6	154.6-187.8
	1996-2000	462.6	503.8	483.4-524.9	237.2	592.6	558.7-627.9	225.4	450.4	424.1-478.0	195.8	210.1	197.1-223.7	98.6	256.0	233.4-280.2	97.2	183.6	167.4-201.0
	2001-2005	472.4	499.9	479.9-520.6	239.8	558.0	526.5-590.9	232.6	459.9	433.4-487.6	186.8	194.9	182.5-207.9	97.4	234.1	213.5-256.2	89.4	167.5	152.2-184.1
	2006-2010	498.2	510.6	490.5-531.4	269.0	593.2	561.3-626.5	229.2	446.3	420.3-473.7	173.6	174.0	162.5-186.1	92.6	208.5	189.6-228.9	81.0	147.5	133.3-163.0
	2011-2015	533.4	524.8	504.5-545.9	276.0	571.9	541.3-603.9	257.4	492.2	464.5-521.2	166.6	160.6	149.7-172.2	89.4	187.9	170.5-206.7	77.2	138.2	124.4-153.3
	2016-2020	519.2	488.4	468.8-508.6	280.6	541.6	512.3-572.3	238.6	445.6	419.2-473.5	166.6	149.7	139.4-160.6	86.8	167.8	151.9-185.1	79.8	135.4	122.0-150.1
Oral cavity and pharynx	1976-1980	9.8	11.9	8.7-15.8	8.2	22.9	16.2-31.4	1.6	3.3	1.4-6.7	4.2	5.1	3.1-7.9	2.8	7.7	4.1-13.0	1.4	2.9	1.2-6.1
	1981-1985	8.8	10.7	7.7-14.5	6.0	16.9	11.2-24.3	2.8	5.3	2.9-9.0	2.8	3.2	1.7-5.4	1.6	4.3	1.8-8.6	1.2	2.2	0.8-5.0
	1986-1990	8.6	9.9	7.1-13.4	6.4	16.5	11.2-23.5	2.2	4.2	2.1-7.6	2.4	2.8	1.4-5.0	1.8	4.4	2.0-8.6	0.6	1.1	0.2-3.4
	1991-1995	8.4	9.4	6.8-12.8	6.2	15.8	10.7-22.6	2.2	3.9	2.0-7.2	2.6	2.7	1.4-4.6	2.0	4.9	2.3-9.2	0.6	1.1	0.2-3.3
	1996-2000	7.6	8.4	6.0-11.6	6.6	16.4	11.2-23.1	1.0	1.8	0.6-4.5	2.2	2.4	1.2-4.4	1.6	4.3	1.8-8.4	0.6	1.0	0.2-3.2
	2001-2005	8.6	9.2	6.6-12.4	5.8	13.4	8.9-19.5	2.8	5.9	3.2-10.0	2.0	2.1	1.0-3.9	1.0	2.5	0.8-5.9	1.0	2.0	0.6-4.8
	2006-2010	14.0	14.3	11.1-18.2	9.4	20.5	15.0-27.5	4.6	8.7	5.5-13.3	4.2	3.9	2.4-6.1	3.4	6.7	3.9-11.0	0.8	1.4	0.4-3.9
	2011-2015	14.4	13.8	10.7-17.7	9.8	19.9	14.5-26.7	4.6	8.3	5.1-12.8	3.4	3.3	1.9-5.5	2.8	5.6	3.0-9.7	0.6	1.4	0.3-4.2
	2016-2020	16.8	15.5	12.3-19.5	11.2	22.1	16.5-29.2	5.6	8.9	5.8-13.3	4.8	3.9	2.5-6.1	2.8	4.9	2.6-8.7	2.0	2.9	1.4-5.8
Esophagus	1976-1980	2.6	3.1	1.7-5.4	1.8	5.7	2.5-10.9	0.8	1.6	0.4-4.3	3.0	3.6	2.0-6.0	2.2	6.7	3.3-12.2	0.8	1.6	0.4-4.3
	1981-1985	3.2	3.6	2.0-5.9	2.0	5.2	2.4-9.8	1.2	2.2	0.8-5.0	3.6	4.2	2.5-6.7	2.2	6.3	3.1-11.4	1.4	2.6	1.1-5.6
	1986-1990	2.2	2.6	1.3-4.8	1.8	4.5	2.0-8.7	0.4	1.0	0.1-3.7	3.2	3.7	2.1-6.1	3.2	8.5	4.7-14.0	0.0	0.0	0.0-1.7
	1991-1995	5.4	5.7	3.8-8.4	4.0	9.3	5.7-14.5	1.4	2.5	1.0-5.4	4.4	4.7	2.9-7.2	3.6	8.3	4.9-13.3	0.8	1.4	0.4-3.9
	1996-2000	5.8	6.2	4.2-9.0	4.8	11.8	7.5-17.7	1.0	2.1	0.7-4.9	5.8	6.2	4.1-8.9	4.0	9.5	5.8-14.7	1.8	3.5	1.6-6.8
	2001-2005	6.6	6.8	4.7-9.6	5.2	12.5	8.1-18.5	1.4	2.5	1.0-5.3	6.6	6.8	4.7-9.6	5.2	11.9	7.8-17.6	1.4	2.5	1.0-5.5
	2006-2010	6.4	6.5	4.4-9.3	5.2	11.5	7.4-17.0	1.2	2.1	0.8-4.8	4.8	4.7	3.0-7.0	3.6	7.9	4.6-12.7	1.2	2.0	0.7-4.6
	2011-2015	6.6	5.9	4.0-8.4	5.8	10.9	7.2-16.0	0.8	1.3	0.4-3.8	5.4	4.9	3.2-7.3	4.4	8.7	5.4-13.5	1.0	1.6	0.5-4.2
	2016-2020	6.2	5.8	3.8-8.5	5.8	10.9	7.1-16.2	0.4	0.8	0.1-3.3	6.0	5.4	3.6-7.9	5.8	10.7	7.1-15.8	0.2	0.3	0.0-2.5
Stomach	1976-1980	5.6	7.0	4.6-10.1	3.2	8.7	4.8-14.3	2.4	5.5	2.8-9.7	5.0	6.5	4.2-9.7	2.6	7.1	3.7-12.3	2.4	5.4	2.8-9.4
	1981-1985	9.8	11.6	8.5-15.4	6.4	18.5	12.4-26.4	3.4	6.8	3.9-11.0	6.6	7.9	5.4-11.1	4.0	12.5	7.4-19.4	2.6	5.1	2.7-8.9
	1986-1990	7.0	8.7	6.0-12.2	4.2	12.4	7.6-19.0	2.8	5.6	3.0-9.6	5.2	6.1	4.0-9.1	3.8	10.3	6.2-16.2	1.4	2.6	1.0-5.5
	1991-1995	5.2	5.6	3.7-8.3	3.2	7.8	4.4-12.8	2.0	4.2	2.0-7.8	2.8	3.2	1.7-5.4	1.6	4.5	1.9-9.0	1.2	2.4	0.8-5.3
	1996-2000	6.8	7.4	5.1-10.4	4.8	12.3	7.8-18.5	2.0	3.6	1.7-6.9	3.6	3.8	2.3-6.1	2.2	6.3	3.1-11.4	1.4	2.4	1.0-5.2
	2001-2005	6.8	7.0	4.9-9.9	4.8	12.2	7.7-18.2	2.0	3.8	1.8-7.2	2.6	2.7	1.4-4.6	1.6	4.0	1.7-8.0	1.0	1.9	0.6-4.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cattaraugus**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	5.6	5.6	3.7-8.2	3.4	7.1	4.1-11.5	2.2	4.1	2.0-7.7	2.4	2.3	1.2-4.1	1.6	3.5	1.5-7.0	0.8	1.3	0.4-3.8
	2011-2015	6.6	6.1	4.1-8.7	5.0	10.1	6.5-15.2	1.6	2.8	1.2-5.9	2.8	2.5	1.4-4.4	1.8	3.8	1.7-7.5	1.0	1.5	0.5-4.0
	2016-2020	5.8	5.2	3.4-7.7	4.0	7.1	4.3-11.5	1.8	3.2	1.3-6.6	0.6	0.7	0.1-2.2	0.4	1.1	0.1-4.0	0.2	0.4	0.0-2.6
Colorectal	1976-1980	59.0	71.9	63.8-80.8	27.0	75.7	63.0-90.2	32.0	68.9	58.5-80.6	31.2	38.6	32.7-45.2	15.0	42.0	32.7-53.0	16.2	35.6	28.2-44.4
	1981-1985	50.2	60.1	52.8-68.1	26.4	76.4	63.5-91.0	23.8	48.1	39.7-57.8	27.2	32.2	27.0-38.2	14.2	43.7	33.7-55.4	13.0	25.9	19.9-33.3
	1986-1990	50.0	58.3	51.2-66.1	26.0	73.9	61.3-88.4	24.0	48.2	39.8-57.8	25.6	30.1	25.1-35.9	12.8	37.2	28.4-47.8	12.8	25.0	19.2-32.1
	1991-1995	58.8	65.1	57.8-73.1	30.8	82.1	69.3-96.6	28.0	52.5	44.0-62.2	23.6	26.3	21.7-31.5	12.6	34.6	26.4-44.5	11.0	20.4	15.3-26.8
	1996-2000	58.2	62.3	55.3-69.9	30.8	78.7	66.5-92.5	27.4	50.4	42.2-59.8	20.8	22.3	18.2-27.0	10.6	28.3	21.0-37.2	10.2	18.8	13.9-24.9
	2001-2005	53.6	56.1	49.6-63.3	26.2	60.7	50.7-72.2	27.4	51.2	42.8-60.7	22.2	23.1	19.0-27.8	11.8	28.9	21.9-37.4	10.4	19.7	14.7-26.1
	2006-2010	55.6	56.2	49.7-63.3	28.4	62.4	52.4-73.9	27.2	50.0	41.8-59.4	15.8	15.8	12.5-19.8	8.2	19.5	13.9-26.6	7.6	13.0	9.2-18.1
	2011-2015	47.0	47.4	41.3-54.1	22.0	47.6	38.9-57.7	25.0	47.8	39.5-57.6	15.0	15.1	11.8-19.1	8.0	17.1	12.1-23.5	7.0	13.3	9.1-18.9
2016-2020	38.8	37.8	32.4-43.9	21.6	43.8	35.5-53.6	17.2	32.0	25.2-40.1	14.2	13.2	10.2-17.0	6.8	13.2	9.0-18.9	7.4	13.1	8.9-18.7	
Colon excluding rectum	1976-1980	38.2	47.0	40.5-54.2	16.0	46.5	36.5-58.2	22.2	47.2	38.7-57.0	23.8	29.5	24.4-35.4	11.0	31.2	23.3-40.9	12.8	27.7	21.3-35.4
	1981-1985	36.0	43.2	37.0-50.1	18.4	53.7	42.9-66.2	17.6	35.4	28.3-43.8	21.2	25.0	20.4-30.3	10.6	32.3	23.9-42.7	10.6	21.4	15.9-28.1
	1986-1990	37.4	43.8	37.6-50.6	18.4	52.8	42.2-65.2	19.0	37.4	30.2-46.0	22.8	26.8	22.1-32.2	12.0	34.8	26.3-45.0	10.8	20.7	15.5-27.2
	1991-1995	45.2	50.2	43.8-57.2	22.0	59.1	48.3-71.6	23.2	43.3	35.7-52.2	21.2	23.6	19.3-28.6	11.0	29.6	22.1-38.7	10.2	18.7	13.9-24.8
	1996-2000	45.4	48.8	42.6-55.6	23.0	59.4	48.8-71.5	22.4	40.9	33.6-49.5	19.0	20.4	16.5-24.9	9.4	25.6	18.7-34.3	9.6	17.6	12.9-23.6
	2001-2005	35.2	36.7	31.5-42.6	16.0	36.8	29.1-45.9	19.2	35.9	29.0-44.0	18.8	19.5	15.7-23.9	9.8	23.9	17.6-31.7	9.0	16.7	12.1-22.6
	2006-2010	41.0	41.3	35.8-47.4	18.6	40.5	32.6-49.9	22.4	40.7	33.4-49.2	13.2	13.3	10.2-16.9	6.4	15.1	10.3-21.5	6.8	11.6	8.0-16.5
	2011-2015	32.4	33.0	28.0-38.7	13.6	30.6	23.6-39.1	18.8	35.5	28.4-44.0	11.8	12.0	9.1-15.7	6.0	13.0	8.7-18.9	5.8	11.1	7.3-16.4
2016-2020	26.6	25.8	21.4-31.0	14.6	29.5	22.8-37.6	12.0	22.1	16.6-29.1	12.6	11.7	8.8-15.2	5.6	10.9	7.1-16.2	7.0	12.0	8.1-17.3	
Rectum & rectosigmoid	1976-1980	20.8	25.0	20.3-30.4	11.0	29.2	21.7-38.4	9.8	21.7	16.0-28.8	7.4	9.1	6.3-12.6	4.0	10.8	6.4-16.9	3.4	7.9	4.5-12.8
	1981-1985	14.2	16.9	13.2-21.4	8.0	22.7	16.1-31.1	6.2	12.7	8.5-18.2	6.0	7.2	4.9-10.4	3.6	11.3	6.6-17.9	2.4	4.6	2.4-8.1
	1986-1990	12.6	14.6	11.2-18.7	7.6	21.1	14.7-29.4	5.0	10.7	6.9-16.0	2.8	3.3	1.8-5.6	0.8	2.4	0.6-6.3	2.0	4.3	2.0-8.0
	1991-1995	13.6	15.0	11.6-19.0	8.8	23.0	16.5-31.1	4.8	9.2	5.8-13.8	2.4	2.7	1.4-4.8	1.6	5.0	2.1-9.9	0.8	1.7	0.4-4.4
	1996-2000	12.8	13.5	10.4-17.3	7.8	19.3	13.7-26.6	5.0	9.5	6.1-14.1	1.8	1.9	0.9-3.6	1.2	2.7	1.0-5.9	0.6	1.2	0.2-3.6
	2001-2005	18.4	19.4	15.6-23.9	10.2	24.0	17.8-31.7	8.2	15.3	10.9-20.9	3.4	3.6	2.1-5.8	2.0	5.0	2.4-9.3	1.4	3.0	1.2-6.3
	2006-2010	14.6	14.9	11.7-18.9	9.8	21.9	16.1-29.2	4.8	9.3	5.9-14.1	2.6	2.6	1.4-4.5	1.8	4.3	1.9-8.4	0.8	1.4	0.4-3.9
	2011-2015	14.6	14.4	11.2-18.3	8.4	16.9	12.1-23.2	6.2	12.3	8.2-17.9	3.2	3.1	1.7-5.1	2.0	4.0	1.9-7.7	1.2	2.2	0.8-5.1
2016-2020	12.2	11.9	9.0-15.6	7.0	14.4	9.7-20.5	5.2	9.9	6.3-14.9	1.6	1.6	0.6-3.3	1.2	2.3	0.8-5.4	0.4	1.1	0.1-4.1	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cattaraugus**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	1.4	1.7	0.7-3.5	1.2	3.3	1.2-7.2	0.2	0.5	0.0-2.8	0.8	1.0	0.3-2.5	0.2	0.4	0.0-2.8	0.6	1.4	0.3-4.1
	1981-1985	2.8	3.1	1.7-5.3	1.4	3.7	1.4-7.7	1.4	2.9	1.2-6.1	2.8	3.1	1.7-5.3	1.8	4.7	2.1-9.1	1.0	1.9	0.6-4.6
	1986-1990	2.4	2.9	1.5-5.1	0.8	2.0	0.5-5.1	1.6	3.7	1.5-7.4	2.0	2.3	1.1-4.3	0.8	2.0	0.5-5.2	1.2	2.7	0.9-6.0
	1991-1995	2.2	2.4	1.2-4.3	1.2	3.3	1.2-7.1	1.0	1.9	0.6-4.6	1.2	1.3	0.5-2.9	0.6	1.8	0.4-5.1	0.6	1.2	0.2-3.5
	1996-2000	3.4	3.6	2.1-5.9	2.4	6.3	3.2-11.2	1.0	2.0	0.6-4.7	3.6	3.8	2.3-6.1	2.8	7.7	4.1-13.0	0.8	1.4	0.4-3.9
	2001-2005	4.4	4.6	2.9-7.0	2.4	5.7	2.9-10.1	2.0	3.8	1.8-7.1	3.8	3.9	2.4-6.2	2.4	5.5	2.8-9.6	1.4	2.6	1.0-5.6
	2006-2010	3.4	3.5	2.0-5.6	2.0	4.2	2.0-8.0	1.4	2.7	1.1-5.7	3.8	3.8	2.3-6.1	1.6	3.6	1.5-7.3	2.2	4.0	2.0-7.3
	2011-2015	7.6	7.1	5.0-9.9	5.8	11.1	7.4-16.2	1.8	3.4	1.5-6.8	5.0	4.6	3.0-7.0	3.2	6.1	3.4-10.3	1.8	3.1	1.4-6.3
2016-2020	6.2	5.7	3.8-8.4	5.2	10.0	6.4-15.1	1.0	2.0	0.6-5.1	3.2	3.1	1.7-5.2	2.4	4.7	2.4-8.6	0.8	1.6	0.4-4.5	
Pancreas	1976-1980	8.2	10.4	7.4-14.1	5.0	15.0	9.6-22.3	3.2	6.6	3.8-10.8	8.2	10.5	7.5-14.3	4.6	14.0	8.7-21.2	3.6	7.9	4.6-12.6
	1981-1985	11.6	13.8	10.4-17.9	5.8	17.5	11.5-25.4	5.8	12.1	8.0-17.5	9.8	11.4	8.4-15.2	4.8	13.8	8.7-20.6	5.0	9.9	6.4-14.7
	1986-1990	8.2	9.1	6.5-12.3	4.6	11.9	7.4-18.1	3.6	7.2	4.2-11.5	8.2	8.9	6.4-12.2	4.0	10.2	6.1-16.0	4.2	8.1	5.0-12.6
	1991-1995	10.0	11.1	8.2-14.6	4.2	10.5	6.4-16.2	5.8	11.4	7.6-16.5	8.6	9.4	6.8-12.8	3.4	9.2	5.2-14.9	5.2	10.1	6.5-14.9
	1996-2000	12.6	13.3	10.2-17.1	6.2	15.2	10.2-21.6	6.4	11.7	8.0-16.7	10.6	11.2	8.4-14.7	5.0	12.7	8.1-18.8	5.6	10.3	6.8-15.0
	2001-2005	7.0	7.3	5.1-10.2	4.4	10.1	6.3-15.5	2.6	5.2	2.7-9.0	8.6	8.9	6.5-12.1	5.4	12.5	8.2-18.4	3.2	5.9	3.4-9.9
	2006-2010	13.6	13.4	10.4-17.0	6.2	13.3	8.9-19.1	7.4	13.5	9.4-18.8	10.6	10.4	7.8-13.7	5.0	10.7	6.8-16.0	5.6	10.1	6.7-14.8
	2011-2015	9.6	9.8	7.1-13.1	4.8	10.9	6.8-16.5	4.8	8.5	5.3-13.0	10.0	10.0	7.3-13.3	5.4	12.4	8.1-18.3	4.6	7.7	4.8-11.9
2016-2020	14.2	13.5	10.4-17.3	8.8	17.1	12.2-23.5	5.4	10.6	6.8-15.8	8.8	7.6	5.5-10.3	5.4	9.7	6.3-14.5	3.4	6.1	3.5-10.1	
Larynx	1976-1980	6.8	7.9	5.4-11.1	5.8	14.5	9.6-21.0	1.0	2.6	0.8-6.0	2.0	2.2	1.1-4.2	1.6	4.0	1.7-8.0	0.4	0.8	0.1-3.1
	1981-1985	6.4	7.1	4.8-10.2	5.8	14.7	9.7-21.3	0.6	1.4	0.3-4.3	2.0	2.3	1.1-4.3	1.4	4.5	1.7-9.4	0.6	1.1	0.2-3.4
	1986-1990	3.6	4.2	2.5-6.8	2.8	7.2	3.9-12.2	0.8	1.9	0.5-5.0	2.4	2.6	1.3-4.5	2.0	4.8	2.3-9.0	0.4	0.9	0.1-3.2
	1991-1995	3.6	4.0	2.3-6.3	2.2	5.2	2.5-9.4	1.4	3.1	1.2-6.5	1.6	1.7	0.8-3.5	1.0	2.6	0.8-6.2	0.6	1.1	0.2-3.5
	1996-2000	4.2	4.6	2.9-7.1	2.8	6.8	3.6-11.5	1.4	2.9	1.1-6.0	1.6	1.8	0.8-3.5	1.6	4.4	1.8-8.8	0.0	0.0	0.0-1.7
	2001-2005	4.4	4.6	2.9-7.0	3.2	7.2	4.1-11.8	1.2	2.4	0.9-5.3	1.0	1.0	0.3-2.5	0.6	1.4	0.3-4.2	0.4	0.8	0.1-3.1
	2006-2010	6.2	6.3	4.2-9.1	4.0	8.4	5.1-13.2	2.2	4.2	2.1-7.8	1.8	1.7	0.8-3.4	1.4	3.1	1.2-6.5	0.4	0.6	0.1-2.7
	2011-2015	4.6	4.2	2.6-6.5	3.6	7.5	4.3-12.2	1.0	1.5	0.5-4.0	1.2	1.3	0.4-2.9	0.8	1.8	0.4-5.0	0.4	0.7	0.1-3.1
2016-2020	3.4	2.7	1.5-4.5	2.8	4.6	2.5-8.2	0.6	0.8	0.2-3.0	0.6	0.5	0.1-1.7	0.4	0.7	0.1-3.2	0.2	0.3	0.0-2.3	
Lung and bronchus	1976-1980	48.6	57.9	50.8-65.8	36.2	102.3	87.4-119.0	12.4	26.8	20.5-34.6	39.8	47.7	41.2-55.0	28.0	81.0	67.6-96.1	11.8	24.7	18.7-31.9
	1981-1985	55.0	63.3	55.9-71.4	35.6	95.5	81.5-111.2	19.4	41.0	33.0-50.3	48.8	55.7	48.8-63.3	31.6	85.3	72.0-100.3	17.2	35.6	28.3-44.2
	1986-1990	59.2	67.8	60.2-76.2	37.8	101.1	86.8-117.0	21.4	43.6	35.5-53.0	46.2	52.8	46.1-60.3	29.4	80.2	67.4-94.6	16.8	34.3	27.2-42.8
	1991-1995	64.2	69.7	62.2-77.8	40.4	101.3	87.5-116.7	23.8	46.4	38.3-55.8	54.4	59.4	52.5-67.0	34.2	88.3	75.2-102.9	20.2	40.1	32.6-49.0
	1996-2000	76.6	81.9	73.9-90.6	40.8	100.7	87.2-115.8	35.8	68.8	59.0-79.8	54.4	58.0	51.3-65.3	29.6	74.8	63.0-88.2	24.8	47.5	39.5-56.8
	2001-2005	69.6	72.5	65.0-80.6	35.8	82.3	70.6-95.5	33.8	64.8	55.3-75.5	47.8	49.7	43.6-56.5	26.0	60.7	50.6-72.3	21.8	40.7	33.4-49.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cattaraugus**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	76.6	77.3	69.7-85.6	44.4	97.0	84.4-111.0	32.2	61.3	52.1-71.8	51.6	52.1	45.9-59.0	29.4	64.2	54.1-75.8	22.2	41.8	34.3-50.6
	2011-2015	78.4	74.3	67.0-82.3	40.8	82.2	71.1-94.7	37.6	68.4	58.7-79.5	53.8	51.9	45.8-58.7	29.6	61.4	51.7-72.5	24.2	43.4	35.8-52.2
	2016-2020	78.2	67.8	61.1-75.2	38.6	70.4	60.5-81.6	39.6	66.4	57.2-76.9	46.0	40.5	35.4-46.4	23.0	42.9	35.2-52.0	23.0	38.7	31.8-46.8
Melanoma of the skin	1976-1980	4.0	5.0	3.0-7.8	2.0	5.8	2.7-10.8	2.0	4.5	2.1-8.5	2.2	2.9	1.4-5.3	1.2	3.7	1.3-8.1	1.0	2.2	0.7-5.3
	1981-1985	4.4	5.7	3.5-8.8	2.4	6.4	3.2-11.3	2.0	5.0	2.3-9.3	1.0	1.3	0.4-3.2	0.8	2.1	0.5-5.4	0.2	0.7	0.0-3.3
	1986-1990	4.4	5.5	3.4-8.4	2.2	5.5	2.7-9.9	2.2	5.6	2.8-10.1	1.0	1.1	0.4-2.6	0.2	0.5	0.0-3.0	0.8	1.6	0.4-4.2
	1991-1995	4.0	4.8	2.9-7.4	2.8	7.7	4.1-13.0	1.2	2.8	1.0-6.1	1.6	1.9	0.8-3.8	0.8	2.4	0.6-6.2	0.8	1.9	0.5-4.8
	1996-2000	14.0	16.1	12.5-20.4	5.8	14.6	9.7-21.1	8.2	18.4	13.1-25.0	2.4	2.7	1.4-4.7	1.4	3.5	1.4-7.2	1.0	2.0	0.6-4.7
	2001-2005	20.8	22.4	18.3-27.2	12.8	29.8	22.9-38.2	8.0	17.0	12.0-23.3	3.0	3.2	1.8-5.3	1.6	3.6	1.5-7.1	1.4	2.6	1.0-5.6
	2006-2010	15.8	16.6	13.0-20.8	10.4	23.2	17.2-30.7	5.4	11.7	7.6-17.2	2.2	2.0	1.0-3.8	1.0	1.9	0.6-4.7	1.2	2.1	0.7-4.8
	2011-2015	21.8	22.0	17.9-26.8	15.2	32.1	25.1-40.6	6.6	14.0	9.4-20.1	2.0	1.7	0.8-3.3	1.8	3.4	1.5-6.8	0.2	0.3	0.0-2.3
	2016-2020	28.8	29.2	24.4-34.8	17.0	35.9	28.4-45.0	11.8	25.6	19.1-33.7	2.0	2.0	0.9-4.0	1.8	4.0	1.7-7.9	0.2	0.3	0.0-2.3
Female breast	1976-1980							37.4	86.4	74.2-100.1							19.2	42.4	34.2-51.9
	1981-1985							45.2	99.9	86.9-114.3							18.8	40.7	32.7-50.1
	1986-1990							49.2	109.0	95.4-124.1							18.8	40.9	32.8-50.4
	1991-1995							51.4	109.2	96.0-123.7							15.2	30.9	24.2-38.9
	1996-2000							58.6	123.1	109.2-138.2							14.8	29.0	22.7-36.6
	2001-2005							66.2	133.2	119.1-148.6							14.4	26.5	20.7-33.6
	2006-2010							62.2	123.4	109.7-138.3							11.8	22.4	17.0-29.2
	2011-2015							71.8	138.2	123.7-154.0							9.2	17.5	12.7-23.7
	2016-2020							61.8	116.9	103.4-131.8							9.8	17.6	12.8-23.8
Cervix uteri	1976-1980							8.2	21.0	14.9-28.7							3.8	9.2	5.4-14.5
	1981-1985							3.0	8.5	4.7-14.0							1.6	3.8	1.6-7.6
	1986-1990							4.4	10.6	6.5-16.1							1.6	3.3	1.4-6.7
	1991-1995							5.4	12.2	8.0-17.9							2.2	4.6	2.2-8.3
	1996-2000							6.2	14.6	9.9-20.8							2.4	5.1	2.6-8.9
	2001-2005							5.2	11.8	7.6-17.4							3.0	6.7	3.7-11.1
	2006-2010							2.2	4.3	2.1-8.0							1.0	1.9	0.6-4.6
	2011-2015							3.8	9.2	5.3-14.7							0.8	1.6	0.4-4.3
	2016-2020							3.4	8.5	4.7-14.1							0.6	1.0	0.2-3.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cattaraugus**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							13.0	29.0	22.2-37.2							2.6	5.4	2.9-9.3
	1981-1985							11.2	23.0	17.2-30.0							2.4	4.8	2.5-8.6
	1986-1990							13.4	28.6	21.9-36.6							1.8	3.5	1.6-6.7
	1991-1995							11.4	26.0	19.5-33.8							3.2	6.4	3.6-10.6
	1996-2000							16.6	33.5	26.6-41.7							2.6	5.0	2.7-8.7
	2001-2005							18.8	37.9	30.6-46.5							1.8	3.3	1.5-6.5
	2006-2010							13.6	25.7	19.8-32.8							1.4	2.3	0.9-5.1
	2011-2015							20.0	37.5	30.2-46.1							2.4	3.9	2.0-7.3
	2016-2020							18.2	34.2	27.1-42.6							3.0	5.6	3.0-9.7
Ovary	1976-1980							10.4	24.2	17.9-32.0							5.6	11.9	7.9-17.3
	1981-1985							6.2	13.7	9.2-19.7							6.0	12.0	8.0-17.2
	1986-1990							8.4	19.3	13.7-26.3							4.8	10.3	6.5-15.5
	1991-1995							7.8	16.3	11.5-22.4							4.4	8.7	5.4-13.3
	1996-2000							6.0	12.2	8.2-17.5							4.6	8.9	5.6-13.5
	2001-2005							6.0	10.9	7.3-15.8							5.2	9.9	6.4-14.8
	2006-2010							8.0	15.5	11.0-21.4							4.8	8.7	5.6-13.2
	2011-2015							8.0	15.5	11.0-21.5							5.6	9.9	6.6-14.6
	2016-2020							6.8	13.1	8.8-18.9							5.0	8.4	5.4-12.9
Prostate	1976-1980				27.0	85.1	70.7-101.3							10.6	35.1	25.9-46.2			
	1981-1985				36.0	103.6	88.4-120.6							10.2	34.0	25.0-44.9			
	1986-1990				36.6	105.0	89.8-122.0							10.8	33.0	24.4-43.5			
	1991-1995				55.2	146.9	129.5-165.8							9.0	29.2	21.0-39.2			
	1996-2000				62.4	154.7	137.7-173.2							10.4	29.8	22.1-39.3			
	2001-2005				66.4	151.1	135.1-168.5							10.0	26.4	19.5-34.9			
	2006-2010				74.4	158.4	142.5-175.8							9.2	23.3	17.0-31.2			
	2011-2015				72.8	139.3	125.0-154.8							4.6	10.2	6.4-15.5			
	2016-2020				73.8	127.6	114.6-141.9							7.8	16.2	11.5-22.5			
Testis	1976-1980				1.0	2.3	0.7-5.7							0.2	0.4	0.0-2.6			
	1981-1985				1.2	3.1	1.1-6.8							0.4	1.0	0.1-3.9			
	1986-1990				1.4	3.9	1.5-8.0							0.0	0.0	0.0-2.0			
	1991-1995				3.2	8.0	4.5-13.0							0.0	0.0	0.0-1.9			
	1996-2000				2.8	7.2	4.0-12.1							0.2	0.4	0.0-2.6			
	2001-2005				3.0	8.1	4.5-13.4							0.2	0.5	0.0-2.9			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cattaraugus**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				3.0	8.5	4.6-14.0							0.2	0.5	0.0-2.8			
	2011-2015				3.2	9.4	5.3-15.3							0.4	0.8	0.1-3.3			
	2016-2020				2.6	8.5	4.4-14.5							0.0	0.0	0.0-2.0			
Urinary bladder (incl. in situ)	1976-1980	13.4	16.1	12.4-20.5	9.6	26.9	19.6-35.9	3.8	7.7	4.6-12.2	6.0	7.3	4.9-10.4	4.6	14.4	9.0-21.9	1.4	2.8	1.1-5.9
	1981-1985	20.2	24.2	19.6-29.5	15.0	45.3	35.3-57.2	5.2	10.7	6.9-15.8	4.2	4.8	3.0-7.4	3.4	10.0	5.7-16.1	0.8	1.6	0.4-4.3
	1986-1990	20.2	22.8	18.6-27.8	16.0	44.2	34.7-55.4	4.2	8.3	5.1-12.8	6.4	7.3	5.0-10.3	4.2	11.9	7.2-18.4	2.2	4.2	2.1-7.6
	1991-1995	20.8	22.9	18.7-27.9	15.6	41.6	32.6-52.1	5.2	10.0	6.5-14.8	3.8	4.3	2.6-6.7	2.6	7.2	3.7-12.4	1.2	2.7	0.9-5.9
	1996-2000	24.0	25.7	21.3-30.7	16.6	41.4	32.8-51.5	7.4	13.7	9.6-19.1	4.4	4.7	2.9-7.1	3.0	7.3	4.0-12.0	1.4	2.5	1.0-5.4
	2001-2005	29.0	30.2	25.5-35.6	22.8	53.4	44.0-64.3	6.2	11.7	7.9-16.8	3.6	3.8	2.2-6.1	2.0	5.1	2.4-9.4	1.6	2.9	1.2-5.9
	2006-2010	24.2	24.8	20.5-29.8	19.8	45.4	36.8-55.5	4.4	8.1	5.0-12.5	3.4	3.4	2.0-5.5	2.6	6.0	3.2-10.5	0.8	1.3	0.3-3.7
	2011-2015	32.2	30.0	25.5-35.1	24.0	50.0	41.3-60.2	8.2	13.5	9.6-18.6	5.2	4.9	3.2-7.3	3.0	6.5	3.6-10.9	2.2	3.5	1.7-6.7
	2016-2020	28.8	25.5	21.4-30.2	21.0	41.5	33.7-50.8	7.8	12.4	8.8-17.4	4.6	4.1	2.6-6.4	3.0	6.3	3.5-10.7	1.6	2.4	1.0-5.2
Kidney and renal pelvis	1976-1980	7.8	9.3	6.6-12.8	4.4	11.7	7.2-18.0	3.4	7.8	4.5-12.6	3.0	3.6	2.0-6.0	1.2	3.4	1.2-7.6	1.8	4.1	1.8-7.8
	1981-1985	5.8	7.0	4.6-10.0	3.0	9.3	5.0-15.5	2.8	5.9	3.2-10.1	3.2	3.7	2.1-6.0	1.4	4.1	1.6-8.7	1.8	3.4	1.6-6.6
	1986-1990	8.4	9.5	6.8-12.9	5.6	15.1	9.8-22.0	2.8	6.0	3.2-10.2	1.8	2.2	1.0-4.2	1.4	4.4	1.7-9.0	0.4	0.8	0.1-3.1
	1991-1995	9.2	10.3	7.5-13.8	5.2	14.1	9.1-20.9	4.0	8.1	4.9-12.7	4.0	4.4	2.7-6.9	2.8	8.5	4.6-14.4	1.2	2.3	0.8-5.2
	1996-2000	9.0	9.8	7.1-13.1	5.8	14.0	9.3-20.2	3.2	6.3	3.5-10.3	3.4	3.7	2.2-6.0	2.6	6.4	3.4-11.0	0.8	1.7	0.5-4.5
	2001-2005	9.2	9.9	7.2-13.3	4.6	10.5	6.6-15.8	4.6	9.5	6.0-14.4	4.0	4.2	2.6-6.6	2.2	5.2	2.6-9.3	1.8	3.3	1.5-6.4
	2006-2010	17.4	17.7	14.1-21.9	10.2	22.3	16.5-29.6	7.2	13.8	9.6-19.4	4.0	3.8	2.3-6.0	2.6	5.6	2.9-9.8	1.4	2.4	0.9-5.2
	2011-2015	17.2	17.1	13.6-21.3	9.8	20.9	15.3-27.9	7.4	14.5	10.1-20.2	3.0	2.8	1.5-4.7	1.8	3.4	1.5-6.9	1.2	2.1	0.8-5.0
	2016-2020	21.0	20.6	16.6-25.3	14.6	30.1	23.2-38.6	6.4	11.7	7.9-16.9	4.2	3.5	2.1-5.5	3.0	5.5	3.0-9.4	1.2	2.0	0.7-4.8
Brain and other central nervous system	1976-1980	5.6	6.8	4.5-9.9	3.0	7.6	4.1-12.7	2.6	6.0	3.1-10.3	4.8	5.6	3.6-8.4	3.2	7.5	4.3-12.5	1.6	3.7	1.6-7.4
	1981-1985	5.0	6.3	4.0-9.4	2.4	6.9	3.5-12.0	2.6	5.3	2.8-9.2	2.2	2.5	1.2-4.5	1.0	2.8	0.9-6.5	1.2	2.2	0.8-5.0
	1986-1990	4.6	5.0	3.2-7.6	2.2	4.9	2.4-9.0	2.4	5.1	2.6-9.0	3.8	4.2	2.5-6.7	1.8	4.5	2.0-8.7	2.0	3.8	1.8-7.1
	1991-1995	8.2	9.1	6.5-12.4	4.6	11.1	7.0-16.8	3.6	7.3	4.3-11.6	4.4	5.0	3.1-7.7	2.4	6.3	3.2-11.1	2.0	4.1	1.9-7.6
	1996-2000	8.8	9.9	7.2-13.3	4.4	10.3	6.5-15.7	4.4	9.3	5.8-14.2	5.6	6.1	4.0-8.8	2.8	6.5	3.6-11.0	2.8	5.5	3.0-9.4
	2001-2005	8.4	9.5	6.8-12.9	4.0	9.6	5.8-14.9	4.4	9.1	5.6-14.0	5.4	5.9	3.9-8.6	3.2	7.3	4.1-12.0	2.2	4.3	2.1-7.8
	2006-2010	6.6	7.6	5.2-10.8	3.8	8.9	5.3-14.1	2.8	6.5	3.5-11.1	2.8	2.8	1.5-4.9	2.0	4.4	2.0-8.3	0.8	1.4	0.4-4.0
	2011-2015	7.0	7.6	5.2-10.7	4.4	10.8	6.6-16.6	2.6	4.6	2.4-8.1	4.6	4.4	2.8-6.8	3.0	5.9	3.2-10.0	1.6	3.1	1.3-6.3
	2016-2020	7.6	8.9	6.2-12.4	5.2	12.5	7.9-18.6	2.4	5.5	2.7-9.9	4.4	4.5	2.8-7.0	2.6	5.5	2.8-9.8	1.8	3.5	1.5-7.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cattaraugus**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	3.0	3.9	2.1-6.5	0.6	1.7	0.3-5.0	2.4	5.9	2.9-10.4	0.2	0.2	0.0-1.3	0.2	0.4	0.0-2.8	0.0	0.0	0.0-1.8
	1981-1985	2.4	3.4	1.7-6.0	0.2	0.6	0.0-3.3	2.2	5.9	2.9-10.6	0.0	0.0	0.0-0.9	0.0	0.0	0.0-2.0	0.0	0.0	0.0-1.7
	1986-1990	3.0	4.0	2.2-6.6	1.0	2.8	0.9-6.6	2.0	4.8	2.2-9.0	1.2	1.5	0.5-3.2	0.2	0.7	0.0-3.5	1.0	1.9	0.6-4.6
	1991-1995	2.8	3.2	1.7-5.4	1.0	2.6	0.8-6.0	1.8	3.8	1.7-7.3	1.0	1.1	0.3-2.6	0.4	1.1	0.1-3.9	0.6	1.1	0.2-3.4
	1996-2000	6.2	7.4	5.0-10.5	1.4	3.5	1.4-7.3	4.8	11.3	7.2-16.9	0.4	0.4	0.1-1.6	0.0	0.0	0.0-1.9	0.4	0.7	0.1-2.8
	2001-2005	5.6	6.2	4.1-9.0	1.4	3.2	1.3-6.7	4.2	9.1	5.6-14.0	0.2	0.2	0.0-1.2	0.0	0.0	0.0-1.9	0.2	0.3	0.0-2.1
	2006-2010	7.6	8.7	6.1-12.1	1.6	3.8	1.6-7.7	6.0	13.6	9.0-19.7	0.2	0.2	0.0-1.2	0.2	0.3	0.0-2.4	0.0	0.0	0.0-1.8
	2011-2015	12.6	15.1	11.4-19.5	1.8	3.9	1.7-7.7	10.8	25.8	19.0-34.2	0.2	0.2	0.0-1.3	0.0	0.0	0.0-2.0	0.2	0.4	0.0-2.6
2016-2020	11.2	14.1	10.4-18.6	2.8	6.3	3.2-11.0	8.4	22.0	15.5-30.1	0.2	0.1	0.0-1.2	0.2	0.3	0.0-2.4	0.0	0.0	0.0-2.0	
Hodgkin lymphoma	1976-1980	3.6	4.2	2.4-6.7	1.4	3.4	1.3-7.3	2.2	4.6	2.2-8.4	1.6	1.9	0.8-3.9	0.6	1.4	0.3-4.4	1.0	2.1	0.7-5.0
	1981-1985	1.4	1.8	0.7-3.7	0.6	1.8	0.4-5.3	0.8	1.9	0.5-4.8	0.4	0.6	0.1-2.0	0.0	0.0	0.0-2.0	0.4	1.1	0.1-4.0
	1986-1990	2.2	2.5	1.2-4.5	1.2	2.9	1.1-6.5	1.0	2.0	0.7-4.8	0.4	0.5	0.1-1.8	0.4	1.0	0.1-3.7	0.0	0.0	0.0-1.7
	1991-1995	1.4	1.7	0.7-3.5	1.2	3.2	1.2-7.0	0.2	0.4	0.0-2.5	0.8	1.0	0.3-2.4	0.6	1.5	0.3-4.4	0.2	0.3	0.0-2.1
	1996-2000	2.0	2.4	1.2-4.4	1.0	2.4	0.8-5.6	1.0	2.3	0.7-5.5	0.4	0.4	0.1-1.6	0.2	0.5	0.0-2.8	0.2	0.3	0.0-2.1
	2001-2005	2.4	2.9	1.5-5.1	1.2	2.9	1.0-6.3	1.2	2.9	1.0-6.3	0.4	0.4	0.0-1.6	0.2	0.4	0.0-2.6	0.2	0.4	0.0-2.5
	2006-2010	1.8	2.1	0.9-4.1	0.6	1.4	0.3-4.1	1.2	2.9	1.0-6.5	0.4	0.4	0.1-1.6	0.0	0.0	0.0-1.9	0.4	0.8	0.1-3.1
	2011-2015	3.2	4.5	2.5-7.3	1.2	3.3	1.1-7.2	2.0	5.7	2.7-10.5	0.4	0.4	0.0-1.5	0.2	0.5	0.0-2.8	0.2	0.3	0.0-2.3
2016-2020	2.0	2.3	1.1-4.4	1.0	2.4	0.8-5.8	1.0	2.2	0.6-5.5	0.8	0.7	0.2-2.0	0.4	0.7	0.1-3.2	0.4	0.5	0.1-2.7	
Non-Hodgkin lymphomas	1976-1980	8.0	9.9	7.0-13.5	3.8	10.3	6.1-16.2	4.2	9.3	5.7-14.3	5.4	6.3	4.1-9.3	2.8	7.4	3.9-12.7	2.6	5.6	2.9-9.6
	1981-1985	13.8	16.6	12.8-21.1	8.6	22.3	16.0-30.3	5.2	11.5	7.4-17.0	5.2	6.2	4.0-9.1	2.8	7.3	3.9-12.3	2.4	5.4	2.7-9.5
	1986-1990	11.0	13.1	9.8-17.1	6.8	18.8	12.9-26.4	4.2	8.2	5.0-12.7	6.8	8.1	5.6-11.5	4.2	11.4	7.0-17.5	2.6	5.1	2.7-8.9
	1991-1995	10.8	12.3	9.2-16.0	6.4	16.5	11.2-23.4	4.4	9.0	5.6-13.8	5.6	6.1	4.0-8.9	3.4	9.4	5.4-15.1	2.2	3.9	1.9-7.2
	1996-2000	20.0	22.0	17.9-26.8	10.6	26.6	19.8-34.9	9.4	18.2	13.3-24.4	6.8	7.4	5.1-10.4	3.4	9.0	5.2-14.5	3.4	6.3	3.7-10.3
	2001-2005	16.6	18.2	14.4-22.6	7.2	17.1	11.9-23.7	9.4	19.2	14.0-25.7	7.4	7.8	5.5-10.8	4.2	10.0	6.2-15.3	3.2	5.9	3.3-9.8
	2006-2010	23.8	25.0	20.6-30.0	12.2	27.3	20.7-35.3	11.6	23.1	17.4-30.1	6.8	6.8	4.7-9.6	3.2	6.8	3.9-11.3	3.6	6.5	3.8-10.5
	2011-2015	23.2	23.6	19.4-28.6	12.0	27.6	20.8-35.8	11.2	19.8	14.8-26.0	6.4	6.0	4.1-8.6	4.0	8.1	4.9-12.8	2.4	4.2	2.2-7.6
2016-2020	22.2	21.5	17.5-26.1	12.2	25.3	19.1-32.9	10.0	17.4	12.8-23.4	6.4	5.6	3.8-8.1	3.4	6.5	3.7-10.7	3.0	4.7	2.6-8.2	
Myeloma	1976-1980	4.0	5.0	3.0-7.7	2.4	7.2	3.7-12.6	1.6	3.3	1.4-6.7	4.2	5.3	3.2-8.2	2.4	7.8	3.9-13.7	1.8	3.7	1.7-7.2
	1981-1985	5.0	6.0	3.8-8.8	3.4	9.3	5.3-14.9	1.6	3.1	1.3-6.2	2.4	2.9	1.5-5.1	1.4	3.9	1.5-8.1	1.0	1.9	0.6-4.7
	1986-1990	4.0	4.9	3.0-7.7	2.2	6.8	3.3-12.2	1.8	3.5	1.6-6.8	3.2	3.9	2.2-6.3	2.0	6.5	3.0-11.9	1.2	2.3	0.9-5.2
	1991-1995	4.6	5.1	3.3-7.7	1.4	3.9	1.5-8.1	3.2	5.9	3.4-9.8	2.2	2.5	1.2-4.4	0.8	2.4	0.6-6.1	1.4	2.8	1.1-5.9
	1996-2000	6.0	6.4	4.3-9.2	3.6	9.6	5.6-15.3	2.4	4.5	2.3-8.0	4.2	4.5	2.8-6.9	2.0	5.4	2.5-9.9	2.2	3.8	1.9-7.1
	2001-2005	6.0	6.1	4.1-8.8	3.8	9.3	5.5-14.7	2.2	4.0	2.0-7.4	3.2	3.3	1.9-5.4	2.0	5.2	2.5-9.6	1.2	2.3	0.8-5.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cattaraugus**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	7.4	7.7	5.4-10.7	4.0	9.2	5.6-14.3	3.4	6.4	3.7-10.5	2.8	2.8	1.5-4.8	1.8	4.3	2.0-8.3	1.0	1.7	0.5-4.2
	2011-2015	6.4	6.4	4.3-9.1	3.6	7.9	4.6-12.7	2.8	5.2	2.8-9.0	1.6	1.5	0.6-3.0	1.0	2.1	0.7-5.2	0.6	0.8	0.2-3.1
	2016-2020	7.0	6.5	4.5-9.2	4.2	8.0	4.9-12.7	2.8	5.0	2.7-8.8	2.6	2.4	1.3-4.3	1.2	2.4	0.9-5.5	1.4	2.2	0.9-5.0
Leukemias	1976-1980	9.6	11.8	8.6-15.7	6.2	17.9	11.9-25.6	3.4	7.7	4.4-12.4	7.2	8.5	5.9-11.8	3.4	9.2	5.2-15.0	3.8	8.4	5.0-13.2
	1981-1985	12.4	14.6	11.1-18.8	7.2	19.3	13.3-27.1	5.2	10.2	6.6-15.1	7.0	8.4	5.8-11.7	2.6	7.4	3.8-12.8	4.4	8.7	5.4-13.3
	1986-1990	11.4	12.8	9.7-16.7	6.4	16.6	11.2-23.7	5.0	9.8	6.2-14.6	7.8	8.9	6.3-12.2	5.2	14.4	9.3-21.2	2.6	4.6	2.5-8.1
	1991-1995	10.2	11.7	8.7-15.5	5.4	14.8	9.6-21.6	4.8	9.5	6.0-14.3	8.0	9.2	6.6-12.6	4.8	14.9	9.4-22.2	3.2	6.2	3.5-10.2
	1996-2000	13.6	14.9	11.5-18.9	8.2	20.7	14.8-28.1	5.4	10.0	6.5-14.7	7.0	7.6	5.3-10.7	3.6	9.5	5.6-15.1	3.4	6.0	3.4-9.8
	2001-2005	14.0	15.1	11.8-19.2	8.2	20.1	14.3-27.3	5.8	11.8	7.8-17.1	7.4	7.8	5.5-10.8	4.8	11.7	7.4-17.5	2.6	4.8	2.5-8.4
	2006-2010	14.2	14.7	11.4-18.6	9.0	20.2	14.6-27.2	5.2	10.4	6.7-15.5	6.2	6.4	4.3-9.2	4.0	9.3	5.6-14.5	2.2	4.1	2.0-7.7
	2011-2015	15.0	15.0	11.7-19.0	9.8	21.7	16.0-29.0	5.2	9.6	6.2-14.5	4.2	4.0	2.5-6.3	2.8	6.3	3.4-10.8	1.4	2.5	1.0-5.5
	2016-2020	16.4	15.5	12.2-19.6	9.2	18.7	13.5-25.5	7.2	12.7	8.7-18.0	5.0	4.4	2.9-6.7	2.6	5.7	3.0-10.0	2.4	3.7	1.9-6.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cayuga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	296.2	383.6	363.9-404.1	138.6	414.3	382.9-447.6	157.6	373.1	346.6-401.0	151.0	195.7	181.8-210.5	76.0	236.2	212.3-262.1	75.0	171.3	154.0-190.0
	1981-1985	348.2	434.8	414.1-456.1	175.8	513.5	478.8-549.9	172.4	393.1	366.3-421.3	167.0	206.8	192.8-221.6	89.0	269.7	244.2-296.9	78.0	170.1	153.1-188.4
	1986-1990	304.6	372.9	354.0-392.5	153.8	445.4	413.4-479.2	150.8	332.0	307.9-357.5	166.0	200.0	186.5-214.3	92.0	274.2	248.8-301.5	74.0	155.2	139.4-172.5
	1991-1995	354.6	414.3	395.1-434.3	176.6	482.9	450.8-516.6	178.0	375.8	350.9-402.1	176.4	203.5	190.2-217.6	92.0	262.4	238.2-288.3	84.4	168.6	152.5-186.1
	1996-2000	401.0	456.5	436.7-477.0	192.2	498.7	467.3-531.5	208.8	439.7	412.9-467.9	160.8	178.4	166.2-191.2	81.6	219.7	198.6-242.4	79.2	154.2	139.1-170.7
	2001-2005	453.4	495.6	475.3-516.5	235.2	575.0	542.4-609.1	218.2	451.5	424.6-479.7	178.2	189.5	177.2-202.5	92.4	233.2	212.3-255.7	85.8	163.6	148.2-180.3
	2006-2010	499.8	517.8	497.4-538.9	264.8	594.3	562.2-627.8	235.0	466.7	439.5-495.3	169.6	170.6	159.1-182.7	89.0	207.0	188.0-227.5	80.6	146.9	132.4-162.6
	2011-2015	500.6	491.4	471.7-511.7	274.4	570.1	539.5-602.0	226.2	435.6	409.3-463.2	174.4	166.1	155.0-177.9	95.2	203.5	185.2-223.1	79.2	140.7	126.6-156.1
	2016-2020	540.8	501.7	482.2-522.0	281.4	530.8	502.4-560.4	259.4	483.6	455.9-512.7	169.8	149.4	139.3-160.2	85.2	162.1	146.7-178.9	84.6	141.2	127.6-156.1
Oral cavity and pharynx	1976-1980	7.2	9.1	6.3-12.7	5.0	14.4	9.2-21.6	2.2	4.9	2.4-9.1	3.0	3.7	2.1-6.2	2.2	6.0	2.9-11.0	0.8	1.7	0.5-4.5
	1981-1985	8.6	10.6	7.6-14.5	5.4	14.5	9.4-21.4	3.2	7.3	4.0-12.1	3.0	3.8	2.1-6.3	1.6	4.5	1.9-9.0	1.4	2.9	1.2-6.1
	1986-1990	6.2	8.3	5.6-11.9	3.2	9.5	5.4-15.5	3.0	6.8	3.7-11.5	2.0	2.6	1.2-4.8	0.6	1.8	0.4-5.1	1.4	3.1	1.2-6.6
	1991-1995	6.8	8.1	5.6-11.4	4.2	11.1	6.7-17.1	2.6	5.7	3.0-9.9	3.6	4.4	2.6-6.9	2.0	5.4	2.6-10.0	1.6	2.9	1.3-6.0
	1996-2000	9.4	11.2	8.2-14.9	5.6	13.5	9.0-19.6	3.8	8.7	5.2-13.7	2.6	3.0	1.6-5.2	1.2	3.0	1.1-6.6	1.4	3.1	1.2-6.5
	2001-2005	8.6	9.6	7.0-13.0	5.0	11.4	7.4-16.9	3.6	7.8	4.6-12.6	1.8	2.1	1.0-4.0	1.2	2.8	1.0-6.2	0.6	1.3	0.3-4.0
	2006-2010	10.8	10.5	7.9-13.8	8.2	17.2	12.3-23.5	2.6	4.5	2.4-8.2	1.8	1.7	0.8-3.4	1.2	2.9	1.0-6.3	0.6	0.9	0.2-3.2
	2011-2015	12.0	10.9	8.3-14.2	8.0	15.2	10.8-21.1	4.0	6.3	3.7-10.2	3.0	2.6	1.5-4.5	1.8	3.5	1.6-6.9	1.2	1.7	0.6-4.3
	2016-2020	11.4	10.9	8.1-14.3	7.4	13.9	9.7-19.6	4.0	7.6	4.5-12.2	1.6	1.3	0.6-2.8	0.8	1.3	0.4-3.9	0.8	1.3	0.3-3.8
Esophagus	1976-1980	2.2	3.0	1.5-5.3	2.2	7.2	3.5-12.9	0.0	0.0	0.0-1.9	2.0	2.6	1.2-4.7	1.8	5.8	2.6-11.1	0.2	0.6	0.0-3.0
	1981-1985	2.6	2.9	1.5-5.1	2.4	6.5	3.2-11.5	0.2	0.4	0.0-2.5	3.4	3.8	2.2-6.2	2.4	6.7	3.3-11.9	1.0	1.9	0.6-4.7
	1986-1990	4.2	5.2	3.1-8.0	3.0	8.0	4.4-13.3	1.2	2.6	0.9-5.9	4.4	5.3	3.3-8.1	3.6	9.9	5.7-15.9	0.8	1.9	0.5-5.1
	1991-1995	4.0	4.6	2.8-7.1	3.6	9.3	5.5-14.7	0.4	0.7	0.1-2.9	3.8	4.3	2.6-6.8	3.4	8.5	4.9-13.6	0.4	0.7	0.1-2.9
	1996-2000	3.0	3.3	1.8-5.5	2.0	5.0	2.4-9.3	1.0	2.0	0.6-4.9	3.6	3.9	2.3-6.2	2.6	6.5	3.5-11.1	1.0	1.9	0.6-4.7
	2001-2005	4.8	5.1	3.3-7.7	4.2	9.8	6.0-15.0	0.6	1.2	0.3-3.8	5.4	5.7	3.8-8.4	4.2	10.2	6.3-15.6	1.2	2.4	0.9-5.4
	2006-2010	5.6	5.6	3.7-8.1	4.8	10.8	6.9-16.2	0.8	1.7	0.4-4.6	4.8	4.7	3.0-7.1	4.2	9.6	5.9-14.9	0.6	1.1	0.2-3.6
	2011-2015	5.6	5.1	3.4-7.5	5.0	9.6	6.1-14.5	0.6	1.0	0.2-3.4	5.6	5.1	3.3-7.5	5.0	9.4	6.0-14.3	0.6	1.1	0.2-3.6
	2016-2020	7.8	6.6	4.7-9.3	6.2	10.6	7.2-15.4	1.6	3.1	1.3-6.6	7.2	6.3	4.4-8.9	5.8	10.5	7.0-15.5	1.4	2.6	1.0-5.9
Stomach	1976-1980	3.6	4.5	2.6-7.1	2.0	6.2	2.8-11.5	1.6	3.2	1.4-6.4	2.6	3.3	1.7-5.8	1.2	3.7	1.3-8.1	1.4	3.3	1.2-7.0
	1981-1985	7.2	8.6	6.0-11.9	4.6	14.2	8.8-21.6	2.6	5.2	2.8-9.1	3.2	4.0	2.3-6.6	1.6	5.0	2.0-9.9	1.6	3.6	1.5-7.2
	1986-1990	4.2	5.3	3.2-8.2	2.8	9.0	4.7-15.2	1.4	3.2	1.2-6.8	3.6	4.4	2.6-7.0	2.6	8.1	4.1-14.0	1.0	2.1	0.7-5.0
	1991-1995	6.4	7.2	4.9-10.3	3.6	9.7	5.7-15.4	2.8	5.5	3.0-9.4	3.6	4.1	2.4-6.5	1.8	5.0	2.2-9.4	1.8	3.6	1.6-6.9
	1996-2000	7.4	8.3	5.9-11.5	3.6	9.0	5.3-14.2	3.8	7.5	4.5-11.9	3.8	4.2	2.5-6.5	1.8	4.6	2.1-8.8	2.0	4.0	1.9-7.5
	2001-2005	7.2	7.6	5.3-10.6	5.2	13.2	8.6-19.5	2.0	3.7	1.8-7.1	2.6	2.7	1.4-4.7	2.0	5.4	2.6-10.0	0.6	1.1	0.2-3.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cayuga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	6.2	6.3	4.2-9.1	4.0	9.1	5.5-14.2	2.2	4.3	2.1-8.1	2.8	2.8	1.5-4.8	1.6	3.6	1.5-7.2	1.2	2.4	0.8-5.7
	2011-2015	7.8	7.7	5.4-10.7	6.0	12.6	8.4-18.2	1.8	3.5	1.5-7.0	2.2	2.1	1.0-4.0	1.4	3.2	1.2-6.7	0.8	1.3	0.3-3.8
	2016-2020	9.0	8.4	6.1-11.5	5.8	11.5	7.6-16.8	3.2	5.6	3.2-9.6	2.4	2.2	1.1-4.1	2.0	4.3	2.0-8.1	0.4	0.5	0.1-2.8
Colorectal	1976-1980	54.0	68.0	60.0-76.8	24.6	72.4	59.7-87.0	29.4	64.2	54.0-75.7	26.8	35.4	29.6-42.0	11.6	38.7	29.0-50.3	15.2	33.9	26.6-42.6
	1981-1985	65.8	80.2	71.7-89.5	32.0	91.5	77.4-107.3	33.8	70.9	60.4-82.7	28.4	34.4	28.9-40.6	12.8	37.4	28.5-48.2	15.6	31.9	25.1-40.1
	1986-1990	47.8	57.5	50.3-65.3	21.6	65.0	52.8-79.1	26.2	53.9	44.8-64.3	23.0	27.9	23.0-33.5	12.6	38.4	29.2-49.6	10.4	21.5	15.9-28.5
	1991-1995	49.8	57.8	50.8-65.5	22.0	63.0	51.4-76.2	27.8	54.3	45.4-64.4	25.0	28.0	23.3-33.4	10.8	30.3	22.5-39.9	14.2	26.2	20.4-33.3
	1996-2000	52.2	58.3	51.4-65.8	26.6	70.6	59.0-83.9	25.6	49.6	41.2-59.4	18.4	20.3	16.4-25.0	9.0	24.4	17.7-32.7	9.4	17.6	12.8-23.6
	2001-2005	48.6	51.7	45.4-58.7	23.2	57.4	47.4-69.0	25.4	48.3	40.1-57.9	18.8	19.8	16.0-24.3	8.4	21.1	15.1-28.5	10.4	18.9	14.0-25.1
	2006-2010	42.4	43.2	37.5-49.5	22.6	52.1	42.8-62.8	19.8	36.5	29.5-44.9	15.0	14.5	11.4-18.3	7.6	18.1	12.8-25.0	7.4	12.3	8.5-17.3
	2011-2015	38.6	37.9	32.6-43.8	20.6	44.4	36.1-54.2	18.0	32.7	26.1-40.7	11.4	10.7	8.0-14.0	7.2	16.4	11.5-22.9	4.2	7.2	4.3-11.5
2016-2020	37.4	35.5	30.3-41.3	19.0	37.3	29.9-46.1	18.4	32.6	25.8-40.9	14.4	12.4	9.7-15.8	7.6	14.5	10.2-20.2	6.8	10.4	7.1-14.9	
Colon excluding rectum	1976-1980	38.8	48.7	42.0-56.1	16.8	50.4	39.8-62.9	22.0	47.1	38.6-56.9	21.2	27.9	22.8-33.9	8.4	27.9	19.8-37.9	12.8	28.3	21.7-36.3
	1981-1985	49.4	60.5	53.1-68.6	23.0	65.0	53.3-78.4	26.4	55.2	46.0-65.8	23.2	28.0	23.0-33.6	10.0	28.2	20.7-37.6	13.2	26.7	20.6-34.2
	1986-1990	40.6	48.8	42.2-56.1	18.4	54.1	43.2-66.8	22.2	46.1	37.7-55.9	19.8	24.0	19.4-29.3	11.2	33.7	25.1-44.1	8.6	17.5	12.6-23.9
	1991-1995	38.2	44.0	37.9-50.8	17.6	51.2	40.7-63.4	20.6	39.3	32.0-48.0	22.6	25.3	20.8-30.5	9.6	27.4	19.9-36.6	13.0	24.1	18.5-31.0
	1996-2000	38.2	42.4	36.6-48.9	19.6	51.8	42.0-63.2	18.6	35.3	28.4-43.6	17.2	18.9	15.1-23.4	8.4	22.9	16.4-31.0	8.8	16.2	11.7-22.1
	2001-2005	35.2	37.3	31.9-43.3	15.4	38.4	30.2-48.1	19.8	37.1	30.0-45.6	15.2	16.0	12.6-20.1	7.2	18.0	12.6-25.1	8.0	14.5	10.3-20.1
	2006-2010	32.2	32.7	27.8-38.3	15.8	36.6	28.9-45.8	16.4	29.9	23.6-37.6	11.4	11.0	8.3-14.4	5.2	12.6	8.2-18.5	6.2	10.3	6.9-15.0
	2011-2015	25.8	25.3	21.0-30.3	12.0	26.8	20.3-34.8	13.8	24.6	18.9-31.6	8.4	7.6	5.5-10.5	5.4	12.1	7.9-17.8	3.0	4.8	2.6-8.5
2016-2020	25.8	23.8	19.7-28.6	10.6	20.4	15.1-27.1	15.2	26.3	20.4-33.7	12.0	10.3	7.8-13.4	5.8	11.1	7.3-16.2	6.2	9.4	6.3-13.8	
Rectum & rectosigmoid	1976-1980	15.2	19.3	15.2-24.3	7.8	22.0	15.4-30.4	7.4	17.0	11.9-23.7	5.6	7.5	4.9-10.9	3.2	10.8	6.0-17.6	2.4	5.6	2.8-10.0
	1981-1985	16.4	19.7	15.6-24.6	9.0	26.5	19.1-35.7	7.4	15.7	10.9-21.8	5.2	6.4	4.2-9.5	2.8	9.2	4.9-15.5	2.4	5.2	2.6-9.3
	1986-1990	7.2	8.7	6.0-12.1	3.2	10.9	6.0-17.9	4.0	7.8	4.7-12.2	3.2	3.9	2.2-6.4	1.4	4.8	1.8-9.9	1.8	4.0	1.8-7.8
	1991-1995	11.6	13.8	10.4-17.9	4.4	11.8	7.3-17.9	7.2	14.9	10.3-20.9	2.4	2.7	1.4-4.8	1.2	3.0	1.1-6.5	1.2	2.1	0.8-4.9
	1996-2000	14.0	15.9	12.4-20.1	7.0	18.9	13.0-26.3	7.0	14.3	9.9-20.2	1.2	1.4	0.5-3.1	0.6	1.5	0.3-4.4	0.6	1.3	0.3-4.0
	2001-2005	13.4	14.4	11.2-18.4	7.8	19.0	13.5-26.1	5.6	11.2	7.4-16.4	3.6	3.8	2.2-6.1	1.2	3.0	1.1-6.6	2.4	4.4	2.2-8.0
	2006-2010	10.2	10.5	7.8-13.9	6.8	15.5	10.7-21.8	3.4	6.6	3.8-10.8	3.6	3.5	2.1-5.6	2.4	5.5	2.8-9.8	1.2	2.0	0.7-4.8
	2011-2015	12.8	12.6	9.6-16.2	8.6	17.5	12.6-23.9	4.2	8.1	4.9-12.7	3.0	3.1	1.7-5.2	1.8	4.3	2.0-8.3	1.2	2.3	0.8-5.6
2016-2020	11.6	11.6	8.7-15.3	8.4	16.9	12.0-23.2	3.2	6.3	3.4-10.8	2.4	2.2	1.1-4.0	1.8	3.4	1.5-6.9	0.6	1.0	0.2-3.5	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cayuga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	1.0	1.2	0.4-2.9	0.4	1.2	0.1-4.4	0.6	1.3	0.3-3.9	1.2	1.5	0.5-3.3	0.6	1.7	0.3-5.2	0.6	1.3	0.3-3.9
	1981-1985	1.8	2.0	0.9-3.9	1.4	3.7	1.4-7.7	0.4	0.9	0.1-3.4	0.8	0.9	0.2-2.5	0.6	1.3	0.3-4.1	0.2	0.7	0.0-3.3
	1986-1990	2.2	2.6	1.3-4.7	1.0	2.5	0.8-6.1	1.2	2.4	0.9-5.4	1.8	2.0	0.9-3.8	1.2	2.9	1.1-6.5	0.6	1.1	0.2-3.6
	1991-1995	2.2	2.3	1.2-4.2	1.6	3.8	1.6-7.7	0.6	1.2	0.2-3.6	2.6	2.9	1.5-5.0	1.2	2.9	1.0-6.4	1.4	2.8	1.1-5.9
	1996-2000	4.2	4.5	2.8-6.9	2.8	7.1	3.9-12.0	1.4	2.4	1.0-5.3	3.8	4.2	2.5-6.5	2.0	5.1	2.5-9.4	1.8	3.3	1.5-6.5
	2001-2005	3.4	3.7	2.1-5.9	2.2	5.3	2.6-9.6	1.2	2.2	0.8-5.2	2.4	2.5	1.3-4.4	1.4	3.6	1.4-7.4	1.0	1.8	0.6-4.5
	2006-2010	5.8	6.0	4.0-8.7	4.2	9.3	5.7-14.4	1.6	3.2	1.4-6.5	4.2	4.3	2.6-6.6	2.6	6.0	3.2-10.5	1.6	2.8	1.2-5.9
	2011-2015	7.6	6.9	4.8-9.6	6.0	11.8	7.9-17.2	1.6	2.9	1.3-6.1	7.0	6.4	4.4-9.1	4.8	9.7	6.1-14.8	2.2	3.5	1.7-6.7
2016-2020	5.8	5.2	3.4-7.6	3.6	6.5	3.8-10.6	2.2	4.3	2.0-8.3	5.0	4.3	2.8-6.5	4.2	7.7	4.7-12.1	0.8	1.3	0.4-4.0	
Pancreas	1976-1980	11.8	15.1	11.5-19.6	6.0	18.6	12.3-26.8	5.8	13.1	8.6-19.0	10.0	12.6	9.3-16.7	5.2	16.3	10.4-24.1	4.8	10.5	6.7-15.9
	1981-1985	8.8	11.0	7.9-14.8	3.8	11.5	6.7-18.2	5.0	10.8	6.9-16.1	10.8	13.2	9.9-17.3	4.6	13.9	8.6-21.1	6.2	13.1	8.8-18.8
	1986-1990	9.0	11.0	8.0-14.8	5.4	15.7	10.2-23.1	3.6	7.5	4.3-12.1	10.4	12.5	9.3-16.5	5.2	15.4	9.9-22.8	5.2	10.7	6.9-15.9
	1991-1995	7.8	9.3	6.6-12.8	4.2	11.9	7.2-18.3	3.6	7.3	4.2-11.8	7.6	9.0	6.4-12.4	4.2	12.5	7.6-19.3	3.4	6.3	3.6-10.3
	1996-2000	9.4	10.7	7.8-14.2	4.6	12.1	7.6-18.2	4.8	9.5	6.0-14.3	9.0	10.1	7.4-13.6	4.4	11.5	7.2-17.5	4.6	8.9	5.6-13.6
	2001-2005	9.6	10.2	7.5-13.6	4.4	10.7	6.7-16.3	5.2	9.4	6.1-14.1	8.8	9.5	6.9-12.9	4.0	10.0	6.1-15.5	4.8	9.1	5.7-13.7
	2006-2010	13.8	13.8	10.7-17.5	7.2	15.9	11.0-22.1	6.6	11.2	7.6-16.2	10.8	10.5	7.9-13.8	5.8	13.1	8.7-18.9	5.0	7.9	5.0-12.1
	2011-2015	11.4	10.3	7.8-13.6	5.6	11.5	7.6-16.9	5.8	10.1	6.7-14.8	10.6	9.7	7.2-12.9	5.8	12.4	8.3-18.0	4.8	7.9	4.9-12.2
2016-2020	13.4	11.6	9.0-15.0	5.8	10.4	6.9-15.3	7.6	12.5	8.7-17.6	9.0	7.7	5.6-10.5	3.4	6.1	3.5-10.2	5.6	8.9	5.9-13.3	
Larynx	1976-1980	4.2	5.3	3.3-8.2	3.6	10.1	5.9-16.1	0.6	1.9	0.4-5.3	2.4	3.1	1.6-5.4	2.0	6.1	2.8-11.3	0.4	1.1	0.1-4.1
	1981-1985	5.2	6.1	3.9-9.0	4.2	10.8	6.6-16.7	1.0	2.2	0.7-5.4	1.4	1.8	0.7-3.7	1.2	4.1	1.5-8.8	0.2	0.4	0.0-2.5
	1986-1990	5.4	6.7	4.4-9.9	4.6	13.1	8.1-19.8	0.8	1.7	0.4-4.5	1.4	1.6	0.6-3.3	1.0	2.5	0.8-5.9	0.4	0.8	0.1-3.1
	1991-1995	5.2	6.0	3.9-8.9	4.6	13.0	8.1-19.7	0.6	1.5	0.3-4.5	1.8	1.9	0.9-3.7	1.6	4.9	2.0-9.8	0.2	0.4	0.0-2.4
	1996-2000	3.6	4.2	2.5-6.6	3.6	9.7	5.7-15.4	0.0	0.0	0.0-1.8	1.0	1.2	0.4-2.7	0.6	1.8	0.3-5.1	0.4	1.0	0.1-3.5
	2001-2005	2.4	2.6	1.4-4.6	2.4	5.6	2.9-9.8	0.0	0.0	0.0-1.9	1.0	1.1	0.4-2.6	0.8	2.1	0.6-5.4	0.2	0.4	0.0-2.6
	2006-2010	4.8	5.0	3.2-7.5	3.8	8.3	5.0-13.2	1.0	2.1	0.7-5.2	1.4	1.4	0.5-2.9	0.8	1.6	0.4-4.4	0.6	1.2	0.2-3.8
	2011-2015	6.8	6.2	4.3-8.8	5.6	10.6	7.0-15.6	1.2	2.2	0.8-5.2	1.2	1.0	0.4-2.5	1.2	2.1	0.8-5.0	0.0	0.0	0.0-2.0
2016-2020	4.0	3.2	1.9-5.1	3.2	5.1	2.9-8.8	0.8	1.4	0.4-4.1	1.0	0.8	0.3-2.1	0.8	1.3	0.3-3.8	0.2	0.3	0.0-2.6	
Lung and bronchus	1976-1980	44.4	57.0	49.6-65.3	30.4	88.1	74.2-103.8	14.0	35.6	27.4-45.4	33.8	42.8	36.5-50.0	25.0	71.5	59.1-85.6	8.8	21.5	15.4-29.2
	1981-1985	49.4	59.9	52.5-68.0	35.0	99.2	84.5-115.6	14.4	32.7	25.3-41.5	42.8	52.1	45.2-59.8	30.0	86.0	72.3-101.4	12.8	29.3	22.4-37.8
	1986-1990	55.4	67.0	59.2-75.5	36.6	105.5	90.4-122.4	18.8	40.5	32.5-49.9	45.4	54.2	47.3-61.9	32.4	94.4	80.0-110.6	13.0	27.0	20.7-34.7
	1991-1995	64.6	74.7	66.7-83.4	41.2	112.4	97.2-129.2	23.4	47.2	38.8-56.9	51.4	59.0	52.0-66.8	33.2	91.9	78.1-107.4	18.2	36.6	29.3-45.3
	1996-2000	64.0	72.1	64.4-80.5	33.8	86.2	73.6-100.3	30.2	62.0	52.3-73.0	45.2	50.5	44.1-57.6	24.8	63.8	53.0-76.2	20.4	40.2	32.6-49.1
	2001-2005	77.2	84.4	76.2-93.3	41.2	100.5	87.2-115.3	36.0	74.6	64.0-86.6	56.4	60.9	53.9-68.5	29.8	74.1	62.7-87.1	26.6	53.4	44.6-63.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cayuga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	78.8	79.2	71.5-87.5	42.8	95.3	82.8-109.2	36.0	68.0	58.2-79.0	53.0	53.8	47.5-60.8	29.2	66.9	56.4-78.9	23.8	44.5	36.7-53.6
	2011-2015	83.0	80.0	72.4-88.4	44.8	94.5	82.3-108.2	38.2	69.1	59.3-80.1	53.8	51.7	45.5-58.4	30.2	63.9	53.9-75.3	23.6	42.0	34.5-50.8
	2016-2020	91.2	78.9	71.7-86.8	44.4	80.6	70.0-92.4	46.8	78.2	68.3-89.4	47.8	41.4	36.2-47.2	22.6	41.4	33.9-50.2	25.2	42.0	34.8-50.5
Melanoma of the skin	1976-1980	4.4	6.8	4.1-10.3	2.4	8.2	4.2-14.3	2.0	6.2	2.8-11.6	1.2	1.4	0.5-3.2	0.4	0.9	0.1-3.5	0.8	1.9	0.5-5.1
	1981-1985	6.0	7.4	5.0-10.7	4.4	12.1	7.5-18.5	1.6	3.5	1.5-7.2	1.4	1.8	0.7-3.7	1.0	3.3	1.0-7.8	0.4	1.1	0.1-3.9
	1986-1990	3.6	4.9	2.9-7.8	2.2	6.2	3.0-11.2	1.4	4.1	1.6-8.3	1.8	2.2	1.0-4.2	1.6	4.6	2.0-9.2	0.2	0.4	0.0-2.5
	1991-1995	6.2	7.6	5.1-10.8	3.6	9.6	5.6-15.1	2.6	5.9	3.1-10.3	1.8	2.2	1.0-4.2	1.4	4.4	1.7-9.0	0.4	0.7	0.1-3.0
	1996-2000	10.2	12.0	8.9-15.8	5.4	13.5	8.9-19.7	4.8	11.1	7.0-16.5	1.2	1.4	0.5-3.0	1.0	2.4	0.8-5.7	0.2	0.4	0.0-2.4
	2001-2005	13.0	14.8	11.4-18.9	7.6	18.2	12.9-25.1	5.4	12.4	8.1-18.2	2.4	2.6	1.3-4.6	2.2	5.4	2.7-9.7	0.2	0.4	0.0-2.6
	2006-2010	23.0	26.0	21.4-31.4	11.6	27.1	20.5-35.2	11.4	26.3	19.7-34.5	1.6	1.6	0.7-3.3	1.6	3.5	1.5-7.1	0.0	0.0	0.0-1.9
	2011-2015	29.2	30.2	25.3-35.7	15.0	31.2	24.4-39.4	14.2	30.9	23.8-39.5	2.4	2.4	1.2-4.4	1.8	3.8	1.7-7.4	0.6	1.0	0.2-3.5
2016-2020	23.4	24.0	19.7-29.1	11.6	23.0	17.3-30.1	11.8	26.5	19.8-34.8	3.2	3.4	1.9-5.7	2.4	5.2	2.7-9.3	0.8	1.8	0.4-5.1	
Female breast	1976-1980							38.0	93.3	80.0-108.2							12.0	27.0	20.5-35.0
	1981-1985							47.4	113.4	98.8-129.6							10.4	24.3	17.9-32.1
	1986-1990							41.6	95.1	82.1-109.6							14.4	31.0	24.0-39.4
	1991-1995							53.8	115.7	101.9-130.9							15.6	33.1	25.9-41.6
	1996-2000							61.4	134.9	119.9-151.1							10.8	21.0	15.7-27.7
	2001-2005							59.8	124.1	110.2-139.3							7.8	14.7	10.4-20.4
	2006-2010							65.0	130.7	116.5-146.2							10.4	19.2	14.1-25.5
	2011-2015							57.6	115.0	101.3-130.2							11.4	20.1	15.0-26.6
2016-2020							71.8	137.4	122.5-153.8							10.6	18.3	13.5-24.5	
Cervix uteri	1976-1980							5.4	14.2	9.2-20.8							2.4	6.5	3.3-11.5
	1981-1985							6.2	16.6	11.1-23.7							1.6	3.2	1.4-6.5
	1986-1990							3.4	8.6	4.9-13.9							1.2	2.6	1.0-5.9
	1991-1995							3.2	8.0	4.5-13.0							1.2	3.0	1.1-6.5
	1996-2000							2.8	6.7	3.6-11.3							0.8	1.7	0.4-4.5
	2001-2005							4.8	11.7	7.4-17.4							0.8	1.9	0.5-4.9
	2006-2010							5.4	12.8	8.3-18.8							1.0	2.0	0.6-4.9
	2011-2015							3.2	5.9	3.3-9.9							1.6	3.6	1.5-7.4
2016-2020							2.2	7.2	3.6-12.8							1.2	2.5	0.8-6.1	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cayuga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							14.2	34.1	26.4-43.4							1.8	3.8	1.8-7.4
	1981-1985							11.2	25.2	18.8-33.1							2.2	4.3	2.1-7.9
	1986-1990							9.4	20.7	15.0-27.8							1.6	3.1	1.4-6.4
	1991-1995							9.6	21.3	15.6-28.5							2.0	4.4	2.0-8.2
	1996-2000							13.0	27.3	20.9-35.0							2.0	3.6	1.7-7.0
	2001-2005							13.2	29.0	22.3-37.0							2.2	4.7	2.3-8.5
	2006-2010							13.8	26.9	20.8-34.3							0.8	1.3	0.3-3.8
	2011-2015							16.0	30.4	23.8-38.3							2.6	4.4	2.3-8.0
	2016-2020							22.0	38.3	31.1-46.8							2.8	4.2	2.3-7.6
Ovary	1976-1980							6.8	16.8	11.5-23.7							4.8	11.6	7.3-17.4
	1981-1985							7.2	17.9	12.3-25.2							4.6	10.5	6.5-16.0
	1986-1990							5.8	13.3	8.8-19.4							3.8	8.3	4.9-13.2
	1991-1995							6.4	15.4	10.4-21.9							3.6	8.2	4.8-13.2
	1996-2000							5.2	11.1	7.2-16.4							3.0	5.9	3.3-10.0
	2001-2005							4.4	8.3	5.2-12.9							4.0	7.3	4.4-11.5
	2006-2010							6.8	13.1	8.9-18.7							2.8	4.7	2.5-8.2
	2011-2015							5.2	10.0	6.4-15.1							3.6	6.5	3.8-10.6
	2016-2020							5.0	8.8	5.5-13.5							3.2	5.7	3.1-9.7
Prostate	1976-1980				19.2	62.8	50.5-77.0							7.8	27.3	19.3-37.4			
	1981-1985				30.2	95.8	80.6-112.8							9.4	33.1	24.1-44.1			
	1986-1990				30.2	87.8	73.9-103.4							8.8	28.5	20.4-38.6			
	1991-1995				34.4	96.2	81.9-112.1							10.0	34.1	25.0-45.2			
	1996-2000				44.0	114.8	100.0-131.2							9.8	30.2	22.2-40.0			
	2001-2005				70.4	170.4	153.0-189.3							6.6	17.9	12.3-25.2			
	2006-2010				73.4	158.8	142.7-176.3							7.8	20.0	14.2-27.4			
	2011-2015				67.4	132.3	118.2-147.7							8.2	18.6	13.3-25.4			
	2016-2020				78.8	138.2	124.5-153.1							6.6	13.4	9.2-19.1			
Testis	1976-1980				2.0	4.9	2.2-9.3							0.0	0.0	0.0-2.2			
	1981-1985				1.6	3.8	1.6-7.8							0.4	0.9	0.1-3.6			
	1986-1990				0.6	1.4	0.3-4.3							0.0	0.0	0.0-2.1			
	1991-1995				1.4	3.2	1.3-6.9							0.0	0.0	0.0-2.0			
	1996-2000				2.6	5.9	3.2-10.3							0.4	0.9	0.1-3.3			
	2001-2005				1.2	3.0	1.1-6.5							0.2	0.5	0.0-2.8			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cayuga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				2.4	6.0	3.1-10.5							0.0	0.0	0.0-1.9			
	2011-2015				2.8	7.6	4.1-12.7							0.0	0.0	0.0-1.9			
	2016-2020				3.0	8.1	4.5-13.5							0.2	0.5	0.0-3.0			
Urinary bladder (incl. in situ)	1976-1980	11.8	14.5	11.0-18.8	8.8	25.8	18.5-34.9	3.0	6.3	3.5-10.6	5.0	6.8	4.4-10.0	3.0	10.6	5.8-17.4	2.0	4.3	2.1-8.1
	1981-1985	19.0	24.2	19.5-29.7	13.0	39.4	30.1-50.5	6.0	13.6	9.0-19.6	5.6	7.2	4.8-10.5	4.0	13.0	7.8-20.2	1.6	3.9	1.6-7.8
	1986-1990	13.2	15.8	12.1-20.2	9.4	26.1	19.0-34.9	3.8	7.9	4.7-12.6	5.2	6.0	3.9-8.8	4.2	12.7	7.7-19.5	1.0	1.9	0.6-4.7
	1991-1995	16.8	19.0	15.1-23.6	12.8	36.0	27.5-46.2	4.0	7.6	4.6-11.9	3.8	4.2	2.5-6.5	2.6	7.5	3.9-12.9	1.2	2.2	0.8-5.0
	1996-2000	20.2	22.3	18.2-27.2	14.4	40.2	31.3-50.8	5.8	11.1	7.4-16.2	4.2	4.5	2.8-6.9	3.4	10.4	6.0-16.6	0.8	1.3	0.3-3.7
	2001-2005	24.2	26.0	21.6-31.2	16.4	41.6	33.1-51.8	7.8	15.8	11.2-21.8	6.8	6.9	4.8-9.7	4.8	12.8	8.2-19.1	2.0	3.4	1.6-6.6
	2006-2010	23.2	23.8	19.6-28.6	17.0	39.2	31.2-48.7	6.2	11.2	7.5-16.2	3.6	3.4	2.0-5.4	3.2	7.9	4.5-12.8	0.4	0.5	0.1-2.6
	2011-2015	31.2	29.5	25.0-34.8	24.4	52.0	42.9-62.4	6.8	11.5	7.9-16.4	4.0	3.8	2.3-6.0	2.8	6.3	3.4-10.7	1.2	1.9	0.7-4.6
	2016-2020	31.2	28.9	24.4-34.1	24.2	47.2	39.0-56.8	7.0	13.0	8.8-18.7	5.2	4.5	2.9-6.8	2.8	5.0	2.7-8.8	2.4	4.0	2.0-7.6
Kidney and renal pelvis	1976-1980	6.4	8.7	5.9-12.3	4.4	13.7	8.4-20.9	2.0	4.5	2.2-8.4	3.0	3.7	2.0-6.1	2.2	6.3	3.1-11.5	0.8	1.7	0.4-4.4
	1981-1985	7.2	8.9	6.2-12.4	5.2	14.7	9.4-21.8	2.0	4.4	2.0-8.3	3.0	4.1	2.3-6.8	2.2	7.5	3.7-13.3	0.8	1.6	0.4-4.3
	1986-1990	8.8	10.9	7.9-14.7	5.2	15.8	10.1-23.3	3.6	8.1	4.7-13.0	3.6	4.9	2.9-7.8	2.8	8.3	4.5-14.0	0.8	2.1	0.5-5.5
	1991-1995	8.8	10.2	7.4-13.8	6.8	17.6	12.1-24.6	2.0	4.0	1.9-7.6	4.4	5.1	3.2-7.7	2.8	7.4	4.0-12.4	1.6	2.8	1.2-5.8
	1996-2000	10.6	12.2	9.1-15.9	6.4	16.1	11.0-22.8	4.2	9.2	5.6-14.1	2.4	2.7	1.4-4.7	1.4	3.6	1.4-7.4	1.0	2.1	0.7-5.1
	2001-2005	16.0	17.6	14.0-22.0	10.8	26.1	19.6-34.1	5.2	10.8	7.0-16.1	4.4	4.7	2.9-7.1	3.6	9.1	5.4-14.5	0.8	1.4	0.4-4.0
	2006-2010	18.8	19.7	15.8-24.2	11.2	25.2	19.0-32.9	7.6	14.1	9.8-19.7	4.4	4.5	2.8-7.0	1.8	4.1	1.9-7.9	2.6	4.7	2.4-8.4
	2011-2015	15.4	15.5	12.1-19.5	9.6	19.3	14.1-26.0	5.8	11.7	7.7-17.2	5.0	4.7	3.0-7.1	2.8	6.2	3.3-10.6	2.2	3.6	1.8-6.9
	2016-2020	20.4	19.9	16.0-24.6	13.2	26.1	19.9-33.8	7.2	14.1	9.6-20.1	3.4	2.9	1.7-4.9	2.4	4.2	2.1-7.7	1.0	1.7	0.6-4.6
Brain and other central nervous system	1976-1980	4.4	5.7	3.5-8.7	3.0	8.3	4.6-13.9	1.4	3.5	1.4-7.2	4.4	6.1	3.7-9.3	3.0	8.8	4.8-14.7	1.4	3.8	1.4-7.9
	1981-1985	5.8	7.1	4.7-10.3	2.6	7.0	3.6-12.1	3.2	7.7	4.3-12.7	3.4	4.2	2.4-6.9	1.6	4.9	2.0-9.7	1.8	4.0	1.8-7.7
	1986-1990	6.2	7.5	5.0-10.7	2.4	6.0	3.1-10.6	3.8	8.8	5.2-13.9	4.6	5.2	3.3-7.9	1.2	3.0	1.1-6.7	3.4	6.9	3.9-11.2
	1991-1995	4.4	5.2	3.2-7.9	2.0	5.1	2.5-9.5	2.4	5.2	2.6-9.2	2.8	3.3	1.8-5.5	1.2	3.0	1.1-6.6	1.6	3.5	1.5-7.0
	1996-2000	4.6	5.3	3.3-7.9	2.2	5.5	2.7-9.8	2.4	5.2	2.7-9.2	2.4	2.7	1.4-4.8	1.0	2.6	0.8-6.0	1.4	3.2	1.3-6.6
	2001-2005	6.4	7.5	5.1-10.6	4.2	10.5	6.4-16.0	2.2	4.9	2.4-8.9	5.6	6.3	4.1-9.1	3.6	8.7	5.1-13.7	2.0	4.5	2.1-8.4
	2006-2010	7.6	8.7	6.1-12.1	4.4	10.3	6.4-15.7	3.2	7.5	4.2-12.5	3.8	4.0	2.4-6.4	2.6	5.9	3.1-10.2	1.2	2.1	0.7-5.0
	2011-2015	5.2	6.0	3.8-8.9	3.2	7.9	4.4-13.0	2.0	3.9	1.7-7.7	4.6	4.7	2.9-7.2	1.8	4.1	1.8-7.9	2.8	5.4	2.8-9.6
	2016-2020	6.6	8.0	5.3-11.4	3.2	7.6	4.2-12.7	3.4	8.6	4.7-14.2	3.4	3.2	1.8-5.4	1.6	3.3	1.4-6.9	1.8	3.2	1.4-6.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cayuga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	2.4	3.1	1.6-5.5	0.6	1.4	0.3-4.4	1.8	4.7	2.1-9.1	0.0	0.0	0.0-1.0	0.0	0.0	0.0-2.2	0.0	0.0	0.0-1.9
	1981-1985	2.6	3.8	2.0-6.5	1.0	2.9	0.9-6.8	1.6	4.8	2.0-9.4	0.4	0.4	0.1-1.7	0.0	0.0	0.0-2.1	0.4	0.8	0.1-3.2
	1986-1990	1.2	1.6	0.6-3.4	0.4	1.2	0.1-4.4	0.8	2.0	0.5-5.1	0.2	0.3	0.0-1.5	0.2	0.6	0.0-3.1	0.0	0.0	0.0-1.8
	1991-1995	3.6	4.4	2.6-7.0	0.4	1.2	0.1-4.1	3.2	8.0	4.6-13.1	0.2	0.2	0.0-1.3	0.2	0.6	0.0-3.1	0.0	0.0	0.0-1.8
	1996-2000	6.6	8.0	5.5-11.2	1.8	4.2	1.9-8.1	4.8	12.3	7.9-18.3	0.2	0.2	0.0-1.3	0.2	0.5	0.0-2.8	0.0	0.0	0.0-1.8
	2001-2005	9.4	11.2	8.2-15.0	2.4	5.6	2.9-9.8	7.0	17.2	11.9-24.0	0.4	0.4	0.0-1.6	0.0	0.0	0.0-1.9	0.4	0.6	0.1-2.8
	2006-2010	12.0	14.0	10.6-18.2	2.6	6.1	3.2-10.5	9.4	22.9	16.6-30.8	0.4	0.4	0.0-1.5	0.2	0.5	0.0-2.8	0.2	0.3	0.0-2.5
	2011-2015	11.4	13.4	10.0-17.6	3.4	7.1	4.0-11.6	8.0	20.7	14.5-28.6	0.2	0.2	0.0-1.3	0.2	0.4	0.0-2.7	0.0	0.0	0.0-2.0
	2016-2020	12.2	14.2	10.7-18.5	5.2	11.6	7.4-17.3	7.0	17.5	11.8-24.9	0.2	0.1	0.0-1.2	0.2	0.3	0.0-2.3	0.0	0.0	0.0-2.1
Hodgkin lymphoma	1976-1980	3.8	4.6	2.7-7.4	1.6	4.1	1.7-8.3	2.2	5.0	2.5-9.2	0.8	1.2	0.3-3.0	0.2	0.4	0.0-2.9	0.6	1.7	0.3-4.9
	1981-1985	2.2	2.5	1.3-4.6	1.4	3.0	1.2-6.5	0.8	2.1	0.5-5.4	0.8	1.0	0.2-2.5	0.8	1.9	0.5-5.2	0.0	0.0	0.0-1.9
	1986-1990	2.6	3.1	1.6-5.4	2.4	5.6	2.8-9.9	0.2	0.5	0.0-2.9	0.0	0.0	0.0-0.9	0.0	0.0	0.0-2.1	0.0	0.0	0.0-1.8
	1991-1995	1.6	2.0	0.9-4.0	1.2	3.0	1.1-6.6	0.4	0.9	0.1-3.4	0.4	0.4	0.1-1.6	0.2	0.4	0.0-2.6	0.2	0.4	0.0-2.4
	1996-2000	2.6	3.2	1.7-5.6	1.8	4.5	2.0-8.5	0.8	2.0	0.5-5.1	0.4	0.5	0.1-1.7	0.2	0.5	0.0-2.9	0.2	0.5	0.0-2.8
	2001-2005	2.2	2.5	1.3-4.5	0.8	1.8	0.5-4.8	1.4	3.5	1.4-7.3	0.6	0.7	0.1-2.0	0.4	1.0	0.1-3.6	0.2	0.4	0.0-2.6
	2006-2010	3.0	3.6	2.0-6.1	2.4	5.6	2.8-9.8	0.6	1.6	0.3-4.6	0.8	1.1	0.3-2.7	0.2	0.5	0.0-2.8	0.6	1.7	0.3-4.9
	2011-2015	1.8	2.0	0.9-3.9	1.4	3.0	1.2-6.4	0.4	1.0	0.1-3.8	0.2	0.1	0.0-1.2	0.2	0.5	0.0-2.8	0.0	0.0	0.0-2.0
	2016-2020	2.2	2.5	1.2-4.6	1.8	4.3	1.9-8.4	0.4	0.6	0.1-3.0	0.4	0.4	0.1-1.7	0.4	0.9	0.1-3.5	0.0	0.0	0.0-2.1
Non-Hodgkin lymphomas	1976-1980	9.6	12.5	9.1-16.6	4.6	13.3	8.2-20.2	5.0	11.7	7.5-17.6	3.8	4.9	2.9-7.6	2.0	6.5	3.0-12.0	1.8	3.9	1.8-7.6
	1981-1985	10.2	12.6	9.3-16.7	5.4	16.4	10.6-24.1	4.8	10.0	6.3-15.0	5.8	7.3	4.8-10.5	3.0	9.7	5.3-16.0	2.8	5.9	3.2-10.2
	1986-1990	13.8	17.1	13.2-21.7	7.2	20.8	14.5-29.0	6.6	13.2	9.0-18.7	7.4	8.8	6.1-12.2	2.8	7.9	4.2-13.5	4.6	9.5	6.0-14.5
	1991-1995	15.0	18.4	14.4-23.1	7.6	20.6	14.5-28.3	7.4	16.8	11.7-23.3	7.6	8.9	6.3-12.3	3.6	10.6	6.2-16.9	4.0	8.1	4.8-12.7
	1996-2000	19.2	22.2	18.0-27.1	8.0	20.8	14.8-28.4	11.2	24.2	18.2-31.7	8.0	9.1	6.5-12.4	3.0	8.1	4.5-13.4	5.0	10.3	6.6-15.5
	2001-2005	21.0	22.6	18.5-27.5	10.8	26.6	19.9-34.8	10.2	20.0	14.8-26.6	7.6	8.1	5.7-11.2	3.8	9.4	5.7-14.7	3.8	7.2	4.3-11.5
	2006-2010	20.8	20.8	17.0-25.4	10.6	23.5	17.5-30.9	10.2	18.6	13.6-24.9	7.2	7.0	4.9-9.8	2.6	5.4	2.9-9.5	4.6	7.7	4.8-12.0
	2011-2015	21.4	22.0	17.9-26.8	12.2	26.4	20.0-34.2	9.2	18.1	13.0-24.6	5.2	5.2	3.4-7.8	3.2	6.5	3.6-10.8	2.0	4.0	1.9-7.7
	2016-2020	20.4	18.3	14.8-22.5	10.6	20.5	15.2-27.2	9.8	16.7	12.2-22.7	7.2	6.0	4.2-8.6	3.4	6.3	3.6-10.4	3.8	6.0	3.5-10.0
Myeloma	1976-1980	3.6	4.8	2.8-7.6	2.0	6.8	3.2-12.5	1.6	3.4	1.5-6.8	2.4	3.4	1.7-5.9	1.0	3.8	1.2-8.5	1.4	3.0	1.2-6.4
	1981-1985	4.6	5.9	3.7-9.0	1.4	4.0	1.5-8.3	3.2	7.4	4.1-12.2	2.4	3.2	1.6-5.6	1.2	4.1	1.4-8.9	1.2	3.0	1.0-6.6
	1986-1990	2.2	2.7	1.3-4.9	1.6	5.4	2.3-10.7	0.6	1.1	0.2-3.6	2.4	2.9	1.5-5.1	0.8	2.6	0.7-6.5	1.6	3.2	1.4-6.4
	1991-1995	5.0	5.9	3.8-8.8	3.0	8.4	4.7-13.9	2.0	3.7	1.7-7.0	2.0	2.2	1.1-4.1	1.0	2.6	0.9-6.2	1.0	1.7	0.6-4.3
	1996-2000	5.4	6.0	4.0-8.8	2.2	5.6	2.8-10.0	3.2	6.1	3.5-10.1	2.8	3.0	1.6-5.0	1.8	5.3	2.4-9.9	1.0	1.8	0.6-4.5
	2001-2005	4.4	4.8	3.0-7.3	2.2	5.4	2.7-9.7	2.2	4.4	2.2-8.2	4.0	4.2	2.6-6.6	1.4	3.5	1.4-7.1	2.6	4.6	2.4-8.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cayuga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	7.6	7.7	5.4-10.7	4.0	9.2	5.6-14.4	3.6	6.5	3.8-10.6	4.0	4.1	2.5-6.5	2.2	5.2	2.6-9.5	1.8	3.4	1.5-6.8
	2011-2015	7.2	6.7	4.7-9.4	3.8	7.5	4.5-12.0	3.4	5.9	3.4-9.8	2.6	2.4	1.3-4.3	1.6	3.3	1.4-6.8	1.0	1.6	0.5-4.3
	2016-2020	6.0	5.7	3.8-8.2	3.0	5.9	3.3-10.1	3.0	5.3	2.9-9.2	2.0	1.9	0.9-3.7	1.2	2.3	0.8-5.4	0.8	1.5	0.4-4.3
Leukemias	1976-1980	10.8	14.1	10.5-18.4	4.4	13.5	8.3-20.6	6.4	14.3	9.7-20.3	8.2	10.7	7.6-14.5	3.0	10.2	5.6-16.8	5.2	11.3	7.3-16.6
	1981-1985	10.2	12.9	9.6-17.1	5.6	16.3	10.6-23.9	4.6	10.2	6.4-15.6	7.0	8.6	6.0-12.1	3.6	10.3	6.0-16.6	3.4	7.0	4.0-11.4
	1986-1990	7.8	9.6	6.8-13.1	3.6	10.6	6.1-16.9	4.2	8.9	5.4-13.8	6.0	7.4	5.0-10.7	3.2	10.5	5.8-17.4	2.8	6.5	3.4-11.0
	1991-1995	8.8	10.1	7.3-13.5	3.8	9.8	5.8-15.4	5.0	10.6	6.8-15.8	6.0	7.0	4.7-10.0	3.2	9.2	5.1-15.1	2.8	5.9	3.1-10.1
	1996-2000	11.8	13.4	10.2-17.3	7.2	19.6	13.7-27.2	4.6	9.2	5.8-14.0	5.8	6.4	4.3-9.2	3.8	11.1	6.6-17.4	2.0	3.7	1.8-7.1
	2001-2005	12.8	14.1	10.8-18.1	6.0	15.1	10.2-21.7	6.8	14.1	9.7-20.0	6.2	6.6	4.4-9.4	3.0	7.9	4.4-13.0	3.2	5.9	3.3-9.9
	2006-2010	16.8	17.8	14.1-22.2	10.0	24.1	17.8-31.9	6.8	13.8	9.4-19.7	8.0	8.6	6.1-11.8	3.2	7.4	4.2-12.2	4.8	9.6	6.0-14.7
	2011-2015	17.6	17.7	14.1-22.1	11.6	25.0	18.8-32.6	6.0	11.3	7.5-16.7	6.2	6.2	4.1-8.9	3.8	8.2	4.9-13.0	2.4	4.8	2.4-8.8
2016-2020	17.8	16.8	13.4-21.0	11.6	22.5	16.9-29.6	6.2	11.5	7.6-16.9	5.4	5.0	3.2-7.4	3.0	5.8	3.2-9.9	2.4	4.2	2.1-7.9	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chautauqua**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	598.4	395.0	380.7-409.8	315.2	492.1	467.3-517.8	283.2	339.4	321.3-358.3	298.6	195.0	185.0-205.3	157.8	252.7	234.7-271.6	140.8	160.5	148.6-173.2
	1981-1985	654.6	419.1	404.5-434.1	323.0	487.4	463.2-512.5	331.6	377.9	359.2-397.4	316.6	200.8	190.8-211.2	161.4	250.2	232.7-268.6	155.2	171.3	159.1-184.3
	1986-1990	680.6	433.6	418.8-448.7	330.4	501.3	476.9-526.7	350.2	404.3	384.8-424.5	330.2	206.8	196.8-217.3	174.6	269.3	251.3-288.3	155.6	170.1	157.9-183.0
	1991-1995	746.4	460.5	445.6-475.7	381.8	549.9	525.1-575.5	364.6	409.9	390.7-429.9	349.2	211.2	201.3-221.4	178.8	266.1	248.6-284.4	170.4	179.2	167.0-192.1
	1996-2000	831.4	510.5	495.0-526.4	426.8	601.8	576.4-628.1	404.6	452.2	432.2-473.0	322.6	191.2	181.9-200.9	164.2	238.1	222.0-255.1	158.4	161.4	150.1-173.5
	2001-2005	866.0	524.3	508.7-540.3	465.4	641.6	615.7-668.3	400.6	445.1	425.4-465.6	330.6	191.9	182.7-201.5	171.4	241.4	225.5-258.3	159.2	161.3	150.0-173.3
	2006-2010	907.4	535.8	520.0-551.9	505.2	652.8	627.3-679.2	402.2	448.3	428.2-469.2	309.2	173.3	164.7-182.4	165.2	216.7	202.0-232.2	144.0	143.8	133.1-155.1
	2011-2015	942.4	534.8	519.1-550.9	514.8	616.3	592.2-641.3	427.6	472.5	451.5-494.2	298.2	160.8	152.5-169.4	154.4	189.5	176.1-203.6	143.8	140.9	130.4-152.1
	2016-2020	933.0	517.2	501.7-533.1	494.8	568.8	545.6-592.7	438.2	477.2	456.0-499.2	283.8	148.9	141.0-157.2	145.6	170.6	158.1-184.0	138.2	131.7	121.6-142.5
Oral cavity and pharynx	1976-1980	16.0	10.3	8.2-12.9	11.0	16.8	12.5-22.0	5.0	5.4	3.5-8.1	5.2	3.2	2.1-4.8	3.4	5.4	3.1-8.8	1.8	1.9	0.8-3.7
	1981-1985	20.0	12.6	10.2-15.4	12.6	18.1	13.8-23.3	7.4	7.8	5.5-10.9	6.4	3.9	2.6-5.5	4.0	5.9	3.5-9.2	2.4	2.6	1.3-4.7
	1986-1990	16.2	10.2	8.1-12.8	9.4	13.7	10.0-18.4	6.8	7.5	5.1-10.6	6.6	4.1	2.8-5.8	4.2	6.2	3.8-9.6	2.4	2.5	1.3-4.6
	1991-1995	15.4	9.6	7.5-12.1	9.2	13.7	10.0-18.4	6.2	6.5	4.4-9.4	5.6	3.2	2.1-4.7	4.0	5.4	3.2-8.3	1.6	1.5	0.6-3.1
	1996-2000	14.2	9.0	7.0-11.4	8.8	12.3	8.9-16.5	5.4	6.1	4.0-9.0	3.0	1.8	1.0-3.0	2.0	3.0	1.4-5.5	1.0	1.1	0.3-2.7
	2001-2005	18.8	11.7	9.5-14.4	13.2	18.0	13.9-23.0	5.6	6.4	4.2-9.3	4.0	2.3	1.4-3.7	3.0	4.2	2.3-6.9	1.0	0.9	0.3-2.4
	2006-2010	18.8	11.2	9.0-13.7	13.0	16.6	12.8-21.3	5.8	6.5	4.3-9.5	5.0	3.0	1.9-4.4	3.8	4.9	2.9-7.7	1.2	1.2	0.4-2.9
	2011-2015	25.4	14.2	11.8-17.1	15.6	18.1	14.2-22.9	9.8	10.9	8.0-14.7	5.2	2.7	1.8-4.1	3.8	4.3	2.6-6.9	1.4	1.3	0.5-3.0
	2016-2020	30.2	16.9	14.2-20.1	21.0	24.9	20.1-30.5	9.2	9.6	6.8-13.2	5.2	2.6	1.7-3.9	3.4	3.7	2.1-6.1	1.8	1.6	0.7-3.4
Esophagus	1976-1980	4.2	2.7	1.7-4.1	3.2	5.1	2.9-8.4	1.0	1.1	0.3-2.7	4.2	2.6	1.6-4.0	2.4	3.8	1.9-6.6	1.8	1.9	0.8-3.6
	1981-1985	5.6	3.4	2.3-5.0	4.2	6.2	3.8-9.6	1.4	1.4	0.5-3.0	3.6	2.3	1.3-3.6	3.0	4.5	2.5-7.5	0.6	0.6	0.1-1.9
	1986-1990	6.2	3.8	2.5-5.4	4.6	6.8	4.2-10.3	1.6	1.7	0.7-3.5	8.2	5.0	3.6-6.8	6.6	9.6	6.5-13.6	1.6	1.9	0.8-3.9
	1991-1995	6.6	4.1	2.8-5.8	3.8	5.1	3.0-8.0	2.8	3.1	1.6-5.3	7.4	4.7	3.3-6.5	5.0	6.9	4.5-10.3	2.4	2.7	1.4-4.9
	1996-2000	8.8	5.4	3.9-7.3	7.2	9.9	7.0-13.8	1.6	1.6	0.7-3.3	6.6	4.1	2.8-5.8	5.4	7.4	4.9-10.9	1.2	1.2	0.4-2.7
	2001-2005	11.6	6.9	5.3-9.0	9.0	12.0	8.8-16.1	2.6	2.4	1.3-4.3	11.6	6.9	5.3-9.0	8.8	11.9	8.7-16.1	2.8	2.6	1.4-4.6
	2006-2010	11.0	6.3	4.8-8.3	10.0	12.9	9.5-17.1	1.0	1.0	0.3-2.5	9.6	5.5	4.0-7.4	8.0	10.3	7.3-14.1	1.6	1.8	0.8-3.8
	2011-2015	10.2	5.6	4.2-7.5	7.6	9.3	6.5-13.0	2.6	2.4	1.2-4.3	8.2	4.5	3.2-6.3	6.8	8.2	5.6-11.6	1.4	1.3	0.5-3.0
	2016-2020	14.0	8.0	6.1-10.3	10.2	12.9	9.4-17.2	3.8	3.6	2.1-6.0	10.4	6.3	4.6-8.4	7.8	9.9	7.0-13.8	2.6	3.0	1.5-5.6
Stomach	1976-1980	12.8	8.5	6.5-10.9	9.2	14.5	10.5-19.6	3.6	4.0	2.4-6.5	11.8	7.6	5.8-9.9	8.4	13.2	9.4-18.0	3.4	3.6	2.1-5.9
	1981-1985	14.6	9.5	7.4-12.0	9.8	15.1	11.0-20.1	4.8	5.5	3.4-8.3	10.0	6.6	4.9-8.8	6.8	10.6	7.3-14.9	3.2	3.7	2.1-6.2
	1986-1990	8.6	5.4	3.9-7.4	6.2	9.7	6.5-13.9	2.4	2.3	1.2-4.1	5.6	3.5	2.3-5.1	3.8	6.1	3.6-9.5	1.8	1.7	0.8-3.4
	1991-1995	9.4	5.8	4.2-7.8	5.4	8.2	5.4-12.0	4.0	4.2	2.5-6.7	7.0	4.3	3.0-6.0	3.8	5.8	3.5-9.1	3.2	3.3	1.9-5.6
	1996-2000	10.8	6.3	4.7-8.3	6.6	9.2	6.3-13.0	4.2	3.8	2.3-5.9	5.0	2.9	1.9-4.4	3.0	4.1	2.3-6.8	2.0	1.9	0.9-3.6
	2001-2005	12.2	7.3	5.6-9.5	6.2	8.8	6.0-12.5	6.0	6.3	4.2-9.1	6.2	3.6	2.5-5.2	3.0	4.2	2.3-7.0	3.2	3.2	1.8-5.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chautauqua**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	10.2	5.8	4.3-7.6	7.8	9.9	7.0-13.6	2.4	2.5	1.3-4.6	4.6	2.6	1.6-4.0	3.0	3.7	2.1-6.3	1.6	1.8	0.7-3.7
	2011-2015	11.0	6.0	4.5-7.9	8.4	10.2	7.3-13.9	2.6	2.5	1.3-4.5	4.4	2.4	1.5-3.7	3.8	4.6	2.8-7.4	0.6	0.6	0.1-2.1
	2016-2020	11.8	6.9	5.2-9.1	8.6	10.6	7.6-14.6	3.2	3.8	2.0-6.5	4.0	2.0	1.2-3.2	2.6	3.0	1.6-5.4	1.4	1.2	0.5-2.9
Colorectal	1976-1980	102.8	66.9	61.1-73.0	51.0	78.7	69.0-89.3	51.8	57.6	50.6-65.2	49.0	32.1	28.2-36.5	23.2	37.2	30.5-45.0	25.8	28.4	23.7-34.0
	1981-1985	99.0	62.4	56.9-68.3	45.8	70.1	61.0-80.0	53.2	56.3	49.6-63.7	48.2	29.9	26.2-34.0	21.6	33.2	27.1-40.4	26.6	27.3	22.8-32.5
	1986-1990	101.2	62.7	57.2-68.5	45.2	70.2	61.1-80.3	56.0	57.7	51.0-65.1	46.2	28.4	24.8-32.4	23.0	36.6	30.1-44.1	23.2	23.3	19.2-28.2
	1991-1995	105.0	63.0	57.7-68.7	51.0	74.0	65.0-83.7	54.0	54.4	47.9-61.5	45.6	27.2	23.7-31.0	19.4	28.5	23.0-34.9	26.2	25.8	21.4-30.9
	1996-2000	114.2	67.4	62.0-73.3	54.8	77.2	68.2-86.9	59.4	59.3	52.6-66.7	39.6	22.9	19.8-26.3	18.4	26.6	21.4-32.6	21.2	19.8	16.1-24.2
	2001-2005	82.8	48.1	43.6-53.0	39.8	55.4	47.9-63.7	43.0	43.2	37.5-49.6	28.4	16.1	13.6-19.1	14.0	19.7	15.4-25.0	14.4	13.8	10.7-17.6
	2006-2010	79.2	45.7	41.2-50.5	38.2	50.5	43.6-58.4	41.0	41.5	35.8-48.0	27.0	14.9	12.4-17.7	11.8	16.0	12.1-20.7	15.2	14.0	10.9-17.8
	2011-2015	76.2	42.7	38.4-47.4	39.0	48.1	41.4-55.6	37.2	38.0	32.4-44.3	28.2	15.2	12.7-18.1	14.6	18.8	14.7-23.9	13.6	12.3	9.5-15.9
2016-2020	57.8	32.9	29.1-37.2	30.0	37.7	31.6-44.6	27.8	28.5	23.7-34.1	21.8	11.9	9.7-14.6	10.2	12.2	9.0-16.3	11.6	11.9	8.8-15.7	
Colon excluding rectum	1976-1980	72.8	47.7	42.9-53.0	35.4	55.0	46.9-64.0	37.4	41.5	35.7-48.1	41.6	27.4	23.8-31.5	19.8	31.9	25.7-39.1	21.8	24.0	19.6-29.1
	1981-1985	69.2	43.3	38.8-48.2	30.2	46.4	39.0-54.6	39.0	40.3	34.7-46.5	39.0	24.2	20.9-27.9	16.8	25.8	20.4-32.2	22.2	22.5	18.5-27.3
	1986-1990	70.2	43.0	38.6-47.9	27.8	43.6	36.5-51.8	42.4	43.4	37.6-49.9	40.0	24.5	21.2-28.2	19.0	30.0	24.2-36.9	21.0	21.3	17.3-25.9
	1991-1995	74.6	44.6	40.2-49.5	32.6	47.5	40.4-55.4	42.0	42.0	36.4-48.4	39.0	23.2	20.1-26.8	15.2	22.4	17.6-28.2	23.8	23.3	19.2-28.1
	1996-2000	80.6	47.4	42.8-52.3	34.0	48.0	41.0-55.9	46.6	46.5	40.6-53.2	35.0	20.2	17.3-23.5	16.0	23.1	18.3-28.7	19.0	17.7	14.2-21.8
	2001-2005	61.4	35.3	31.5-39.6	28.4	39.9	33.6-47.1	33.0	32.8	27.9-38.4	24.6	14.0	11.6-16.7	11.2	15.9	12.0-20.7	13.4	12.9	9.9-16.6
	2006-2010	60.0	33.9	30.1-38.0	27.8	36.5	30.6-43.2	32.2	31.6	26.8-37.2	24.0	13.1	10.9-15.8	10.8	14.5	10.9-19.0	13.2	12.2	9.3-15.7
	2011-2015	56.2	31.4	27.7-35.5	27.2	33.8	28.2-40.2	29.0	29.3	24.5-34.8	24.2	12.8	10.6-15.5	12.0	15.2	11.5-19.7	12.2	10.8	8.2-14.1
2016-2020	43.2	24.3	21.0-28.0	21.6	27.7	22.5-33.8	21.6	21.1	17.1-25.8	18.0	9.9	7.8-12.3	8.4	10.0	7.1-13.8	9.6	10.0	7.2-13.6	
Rectum & rectosigmoid	1976-1980	30.0	19.1	16.1-22.5	15.6	23.7	18.6-29.7	14.4	16.1	12.5-20.4	7.4	4.7	3.3-6.5	3.4	5.4	3.0-8.7	4.0	4.5	2.7-7.0
	1981-1985	29.8	19.1	16.1-22.5	15.6	23.7	18.6-29.7	14.2	16.0	12.4-20.4	9.2	5.7	4.2-7.7	4.8	7.4	4.7-11.1	4.4	4.8	2.9-7.4
	1986-1990	31.0	19.6	16.6-23.1	17.4	26.6	21.2-32.9	13.6	14.3	11.0-18.3	6.2	4.0	2.7-5.7	4.0	6.6	4.0-10.3	2.2	2.1	1.0-3.9
	1991-1995	30.4	18.3	15.5-21.6	18.4	26.5	21.3-32.6	12.0	12.4	9.4-16.1	6.6	3.9	2.7-5.6	4.2	6.1	3.7-9.4	2.4	2.5	1.3-4.6
	1996-2000	33.6	20.1	17.1-23.4	20.8	29.1	23.8-35.4	12.8	12.8	9.7-16.5	4.6	2.6	1.7-4.0	2.4	3.5	1.8-6.2	2.2	2.1	1.0-4.0
	2001-2005	21.4	12.8	10.4-15.5	11.4	15.5	11.7-20.1	10.0	10.4	7.6-13.9	3.8	2.2	1.3-3.5	2.8	3.9	2.1-6.5	1.0	0.9	0.3-2.4
	2006-2010	19.2	11.8	9.5-14.5	10.4	14.0	10.4-18.5	8.8	9.9	7.0-13.5	3.0	1.7	0.9-3.0	1.0	1.5	0.4-3.5	2.0	1.9	0.9-3.7
	2011-2015	20.0	11.3	9.1-13.9	11.8	14.3	10.8-18.6	8.2	8.7	6.1-12.1	4.0	2.3	1.4-3.7	2.6	3.7	1.9-6.4	1.4	1.5	0.6-3.3
2016-2020	14.6	8.6	6.6-11.0	8.4	9.9	7.1-13.7	6.2	7.4	4.9-10.9	3.8	2.1	1.2-3.3	1.8	2.2	1.0-4.3	2.0	1.9	0.9-3.8	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chautauqua**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	2.6	1.7	0.9-2.9	1.6	2.3	1.0-4.7	1.0	1.1	0.4-2.7	2.4	1.5	0.8-2.6	1.6	2.5	1.1-5.0	0.8	0.9	0.2-2.4
	1981-1985	4.0	2.5	1.5-3.9	2.8	4.1	2.2-7.0	1.2	1.3	0.5-2.9	3.2	2.1	1.2-3.5	2.0	3.1	1.5-5.8	1.2	1.5	0.5-3.3
	1986-1990	4.2	2.6	1.6-4.0	3.0	4.6	2.5-7.6	1.2	1.1	0.4-2.7	3.6	2.2	1.3-3.5	2.0	3.2	1.5-6.0	1.6	1.5	0.6-3.1
	1991-1995	7.4	4.5	3.1-6.2	3.4	5.0	2.9-8.1	4.0	4.2	2.5-6.6	7.8	4.7	3.3-6.4	4.4	6.1	3.8-9.3	3.4	3.7	2.1-6.1
	1996-2000	5.4	3.1	2.1-4.6	2.8	4.1	2.2-6.9	2.6	2.7	1.4-4.7	5.0	2.9	1.9-4.3	2.8	4.0	2.2-6.7	2.2	2.0	1.0-3.8
	2001-2005	8.8	5.2	3.8-7.0	6.8	9.5	6.6-13.4	2.0	2.2	1.0-4.1	8.8	5.0	3.6-6.8	5.2	7.3	4.8-10.8	3.6	3.6	2.1-5.9
	2006-2010	7.8	4.4	3.1-6.1	5.2	6.4	4.2-9.5	2.6	2.3	1.2-4.2	7.6	4.2	2.9-5.8	4.8	6.0	3.9-9.1	2.8	2.4	1.3-4.3
	2011-2015	12.8	6.7	5.2-8.7	9.0	10.0	7.2-13.5	3.8	3.7	2.2-5.9	10.2	5.2	3.9-7.0	6.2	6.9	4.7-10.0	4.0	3.7	2.2-5.9
	2016-2020	12.0	6.1	4.6-7.9	9.4	9.8	7.1-13.3	2.6	2.7	1.4-4.8	10.0	5.0	3.7-6.7	8.4	8.8	6.3-12.1	1.6	1.6	0.7-3.5
Pancreas	1976-1980	16.0	10.3	8.1-12.8	9.6	15.1	11.0-20.2	6.4	6.7	4.6-9.6	12.8	8.4	6.5-10.8	7.8	12.8	9.0-17.6	5.0	5.4	3.5-8.0
	1981-1985	16.6	10.4	8.2-12.9	6.0	9.4	6.3-13.5	10.6	11.0	8.2-14.5	14.4	9.2	7.1-11.6	6.2	9.6	6.5-13.7	8.2	8.9	6.3-12.2
	1986-1990	21.8	13.7	11.2-16.6	10.2	16.0	11.8-21.1	11.6	12.2	9.2-16.0	16.2	10.0	7.9-12.5	8.0	12.8	9.1-17.5	8.2	8.4	6.0-11.6
	1991-1995	19.0	11.1	9.0-13.6	10.0	14.1	10.4-18.7	9.0	9.1	6.6-12.3	19.4	11.5	9.3-14.0	9.8	13.9	10.2-18.4	9.6	9.7	7.1-13.0
	1996-2000	19.4	11.5	9.3-14.1	9.6	14.1	10.4-18.7	9.8	9.7	7.1-13.0	18.4	10.8	8.7-13.3	9.6	14.1	10.4-18.7	8.8	8.5	6.1-11.6
	2001-2005	21.2	12.6	10.3-15.3	8.6	12.2	8.8-16.5	12.6	13.0	9.9-16.8	18.4	10.9	8.8-13.5	8.2	11.5	8.2-15.6	10.2	10.3	7.6-13.7
	2006-2010	21.6	12.2	10.0-14.8	11.6	15.1	11.4-19.6	10.0	9.9	7.3-13.2	19.8	11.0	8.9-13.5	9.8	12.6	9.3-16.8	10.0	9.9	7.3-13.3
	2011-2015	25.0	13.3	11.0-15.9	13.6	15.7	12.2-20.1	11.4	10.7	8.0-14.1	21.4	11.4	9.3-13.8	10.8	12.8	9.6-16.9	10.6	10.0	7.4-13.3
	2016-2020	25.2	13.4	11.1-16.2	12.6	14.2	10.7-18.4	12.6	12.8	9.7-16.7	17.6	9.3	7.4-11.6	7.0	8.0	5.5-11.4	10.6	10.1	7.5-13.6
Larynx	1976-1980	8.0	5.1	3.6-7.0	6.6	9.6	6.6-13.6	1.4	1.8	0.7-3.7	1.4	0.9	0.3-1.8	1.2	1.8	0.6-4.0	0.2	0.2	0.0-1.4
	1981-1985	7.6	4.8	3.3-6.6	5.6	7.9	5.2-11.5	2.0	2.3	1.1-4.4	1.8	1.0	0.5-2.0	1.2	1.6	0.6-3.7	0.6	0.6	0.1-1.8
	1986-1990	7.0	4.7	3.2-6.5	4.4	6.5	4.0-9.9	2.6	3.2	1.7-5.5	1.6	0.9	0.4-1.8	1.2	1.4	0.5-3.2	0.4	0.4	0.0-1.6
	1991-1995	7.0	4.2	2.9-5.9	5.0	7.1	4.6-10.5	2.0	2.0	0.9-3.9	4.2	2.6	1.6-4.1	3.4	5.0	2.9-8.0	0.8	0.9	0.2-2.4
	1996-2000	8.8	5.5	4.0-7.4	6.4	9.0	6.2-12.8	2.4	2.8	1.4-5.0	3.0	1.9	1.0-3.1	2.4	3.3	1.7-5.8	0.6	0.7	0.1-2.2
	2001-2005	7.6	4.8	3.4-6.6	5.2	7.0	4.6-10.3	2.4	2.8	1.5-5.1	2.4	1.4	0.7-2.5	2.0	2.7	1.3-5.1	0.4	0.4	0.1-1.8
	2006-2010	6.8	4.0	2.8-5.6	5.8	7.1	4.7-10.3	1.0	1.3	0.4-3.1	2.0	1.1	0.5-2.1	1.8	2.0	0.9-4.0	0.2	0.2	0.0-1.5
	2011-2015	9.6	5.2	3.8-7.0	7.6	8.8	6.2-12.3	2.0	2.1	1.0-4.2	2.2	1.1	0.6-2.2	1.6	2.0	0.8-4.0	0.6	0.6	0.1-2.1
	2016-2020	5.6	2.9	1.9-4.3	4.0	4.3	2.6-6.9	1.6	1.7	0.7-3.6	2.4	1.1	0.6-2.0	2.2	2.4	1.2-4.5	0.2	0.2	0.0-1.4
Lung and bronchus	1976-1980	79.4	50.3	45.4-55.7	55.8	81.0	71.5-91.4	23.6	28.3	23.2-34.1	62.8	39.8	35.4-44.5	46.4	70.0	61.0-79.8	16.4	18.9	14.9-23.6
	1981-1985	91.2	57.5	52.2-63.2	62.8	91.9	81.7-102.9	28.4	32.4	27.1-38.5	70.6	44.4	39.8-49.4	48.0	71.6	62.6-81.6	22.6	25.1	20.5-30.4
	1986-1990	98.6	62.7	57.1-68.6	58.6	84.9	75.3-95.5	40.0	46.9	40.3-54.2	80.4	50.5	45.6-55.9	50.0	71.2	62.5-80.8	30.4	35.6	29.9-42.1
	1991-1995	108.6	66.3	60.8-72.2	67.2	97.5	87.2-108.7	41.4	45.8	39.6-52.6	88.0	53.8	48.8-59.1	53.8	79.2	69.9-89.5	34.2	37.2	31.7-43.4
	1996-2000	109.4	67.3	61.7-73.2	61.0	85.6	76.2-95.9	48.4	54.8	48.0-62.3	86.4	52.2	47.4-57.4	48.8	69.1	60.6-78.3	37.6	41.1	35.3-47.6
	2001-2005	119.8	70.8	65.2-76.8	64.8	88.7	79.3-99.0	55.0	57.8	51.0-65.3	92.4	54.0	49.2-59.2	52.4	72.3	63.8-81.7	40.0	40.9	35.4-47.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chautauqua**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	123.0	70.5	65.0-76.4	66.4	86.2	77.1-96.1	56.6	59.5	52.7-67.1	91.2	51.3	46.6-56.3	50.0	64.8	56.9-73.4	41.2	41.7	36.1-48.1
	2011-2015	123.8	68.5	63.1-74.3	66.0	80.1	71.6-89.5	57.8	60.0	53.1-67.6	78.8	43.0	38.7-47.5	42.0	50.9	44.2-58.5	36.8	37.2	31.9-43.3
	2016-2020	135.2	70.7	65.3-76.5	64.0	71.4	63.6-80.1	71.2	71.2	63.7-79.5	83.0	43.4	39.2-47.9	42.6	48.7	42.2-56.1	40.4	38.8	33.5-44.9
Melanoma of the skin	1976-1980	9.6	7.4	5.4-9.9	5.6	9.9	6.5-14.4	4.0	5.9	3.5-9.2	2.6	1.8	1.0-3.2	1.6	2.5	1.0-5.0	1.0	1.3	0.4-3.2
	1981-1985	13.6	9.7	7.4-12.4	5.4	8.9	5.8-13.0	8.2	10.3	7.2-14.2	3.0	2.3	1.3-3.8	1.6	2.8	1.2-5.5	1.4	1.7	0.7-3.7
	1986-1990	12.2	8.1	6.2-10.5	7.2	10.7	7.4-14.9	5.0	6.4	4.0-9.5	2.4	1.7	0.8-2.9	1.2	1.9	0.7-4.1	1.2	1.7	0.6-3.7
	1991-1995	18.6	12.3	9.9-15.1	10.2	15.2	11.2-20.0	8.4	10.5	7.5-14.4	2.8	1.7	0.9-2.9	2.4	3.5	1.8-6.1	0.4	0.4	0.1-1.7
	1996-2000	21.2	13.5	11.1-16.4	11.6	16.6	12.6-21.5	9.6	11.6	8.5-15.6	2.8	1.7	0.9-2.9	2.2	3.2	1.6-5.7	0.6	0.6	0.1-2.0
	2001-2005	27.8	17.8	14.9-21.1	15.2	21.7	17.0-27.2	12.6	15.2	11.6-19.6	5.6	3.3	2.2-4.9	4.2	5.7	3.5-8.7	1.4	1.5	0.6-3.3
	2006-2010	35.0	22.4	19.1-26.1	20.6	27.7	22.5-33.7	14.4	18.2	14.0-23.3	5.2	3.1	2.0-4.6	3.2	4.6	2.6-7.5	2.0	2.1	1.0-4.1
	2011-2015	41.0	24.1	20.8-27.8	23.4	28.9	23.8-34.9	17.6	20.4	16.1-25.6	4.8	2.6	1.6-3.9	2.6	2.9	1.5-5.2	2.2	2.3	1.1-4.5
	2016-2020	48.4	29.7	25.8-34.0	27.8	34.4	28.7-41.1	20.6	26.7	21.3-33.0	4.0	2.1	1.2-3.3	2.4	2.9	1.5-5.4	1.6	1.4	0.6-3.1
Female breast	1976-1980							78.2	97.9	88.0-108.7							29.8	36.0	30.3-42.6
	1981-1985							90.4	105.6	95.6-116.3							36.0	42.5	36.2-49.5
	1986-1990							101.2	126.5	115.2-138.6							30.4	34.6	29.1-40.8
	1991-1995							105.8	124.3	113.6-135.8							28.2	32.8	27.4-39.0
	1996-2000							126.8	147.8	136.2-160.1							23.6	23.7	19.5-28.6
	2001-2005							118.0	135.1	124.2-146.8							27.2	28.9	24.1-34.5
	2006-2010							115.0	131.6	120.7-143.3							17.0	17.7	14.0-22.1
	2011-2015							116.0	130.6	119.5-142.5							19.2	19.1	15.3-23.7
	2016-2020							121.8	133.0	122.0-144.9							16.0	14.9	11.7-19.0
Cervix uteri	1976-1980							9.8	13.4	9.8-17.8							2.2	2.9	1.4-5.3
	1981-1985							9.2	11.7	8.5-15.9							3.8	4.6	2.7-7.4
	1986-1990							9.0	12.0	8.7-16.2							2.6	2.9	1.5-5.1
	1991-1995							7.4	9.8	6.8-13.7							1.8	2.1	0.9-4.1
	1996-2000							8.0	10.6	7.6-14.6							2.6	3.1	1.6-5.5
	2001-2005							4.8	6.5	4.1-9.8							0.4	0.5	0.1-1.9
	2006-2010							6.8	9.0	6.1-12.8							1.2	1.2	0.4-2.9
	2011-2015							5.8	8.3	5.4-12.3							2.0	2.0	0.9-3.8
	2016-2020							5.4	7.9	5.0-11.9							2.2	2.1	1.0-4.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chautauqua**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							21.8	26.5	21.6-32.3							5.6	6.1	4.0-8.9
	1981-1985							23.8	27.1	22.2-32.6							4.6	4.6	2.9-7.0
	1986-1990							21.4	24.6	20.0-30.0							3.2	3.1	1.8-5.2
	1991-1995							24.8	29.7	24.5-35.6							5.2	4.6	3.0-7.0
	1996-2000							23.6	27.2	22.4-32.8							3.8	3.9	2.3-6.2
	2001-2005							26.0	30.6	25.5-36.4							4.0	4.2	2.5-6.6
	2006-2010							27.8	31.8	26.5-37.7							6.8	6.9	4.7-9.9
	2011-2015							30.8	33.3	28.0-39.4							6.6	6.6	4.5-9.4
	2016-2020							32.2	35.3	29.7-41.7							5.2	5.1	3.3-7.8
Ovary	1976-1980							16.8	20.4	16.1-25.5							11.4	12.9	9.7-16.8
	1981-1985							17.8	21.3	16.9-26.5							8.4	9.7	6.8-13.3
	1986-1990							14.8	17.9	13.9-22.7							7.4	7.9	5.5-11.1
	1991-1995							15.8	18.1	14.2-22.8							9.8	10.0	7.3-13.4
	1996-2000							14.4	16.5	12.8-21.0							8.6	9.1	6.5-12.4
	2001-2005							15.4	17.8	14.0-22.4							12.0	12.6	9.6-16.4
	2006-2010							12.6	14.4	11.0-18.7							10.4	10.8	8.0-14.3
	2011-2015							11.8	13.3	10.0-17.5							7.2	8.2	5.6-11.5
	2016-2020							9.6	10.0	7.2-13.7							7.0	6.8	4.7-9.7
Prostate	1976-1980				72.2	121.2	108.6-134.7							21.0	37.7	30.6-45.8			
	1981-1985				67.8	105.9	94.6-118.2							22.8	39.5	32.4-47.7			
	1986-1990				76.8	118.9	107.0-131.7							27.2	46.3	38.7-54.9			
	1991-1995				116.0	163.2	150.0-177.3							28.0	45.2	37.9-53.5			
	1996-2000				140.6	195.7	181.4-210.8							20.6	32.6	26.6-39.6			
	2001-2005				163.4	222.1	207.0-237.9							18.0	27.0	21.7-33.2			
	2006-2010				188.8	237.8	222.7-253.6							16.0	22.0	17.5-27.5			
	2011-2015				181.0	206.7	193.2-220.9							11.0	13.8	10.4-18.1			
	2016-2020				148.8	155.0	143.8-167.0							12.6	14.6	11.2-18.9			
Testis	1976-1980				2.6	4.0	2.0-6.9							0.8	1.2	0.3-3.2			
	1981-1985				2.8	4.0	2.2-6.9							0.2	0.2	0.0-1.5			
	1986-1990				3.0	4.1	2.3-6.8							0.6	0.7	0.2-2.3			
	1991-1995				4.8	7.0	4.5-10.4							0.2	0.3	0.0-1.6			
	1996-2000				4.4	6.7	4.2-10.1							0.0	0.0	0.0-1.1			
	2001-2005				2.2	3.6	1.8-6.4							0.2	0.4	0.0-1.9			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chautauqua**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				3.4	5.8	3.3-9.3							0.2	0.3	0.0-1.7			
	2011-2015				4.6	7.2	4.5-10.9							0.0	0.0	0.0-1.2			
	2016-2020				3.2	6.0	3.4-9.8							0.6	1.0	0.2-2.9			
Urinary bladder (incl. in situ)	1976-1980	29.6	19.0	16.0-22.4	22.6	34.8	28.5-42.1	7.0	8.0	5.5-11.2	7.4	5.0	3.5-6.9	5.6	9.8	6.4-14.3	1.8	2.0	0.9-4.0
	1981-1985	34.8	21.5	18.3-25.0	26.8	40.5	33.8-48.2	8.0	7.9	5.7-10.9	8.4	5.4	3.8-7.3	7.2	11.7	8.1-16.2	1.2	1.2	0.4-2.7
	1986-1990	40.4	24.4	21.1-28.1	31.2	47.0	39.7-55.2	9.2	9.2	6.7-12.4	7.6	4.6	3.2-6.3	4.4	7.9	4.9-11.9	3.2	3.0	1.7-5.0
	1991-1995	36.8	22.0	18.9-25.5	27.2	39.3	32.9-46.6	9.6	9.4	6.9-12.6	9.8	5.6	4.2-7.5	6.6	10.3	7.0-14.5	3.2	3.0	1.7-5.1
	1996-2000	44.4	26.2	22.8-29.9	33.0	47.2	40.2-55.0	11.4	11.8	8.8-15.4	9.2	5.4	3.9-7.2	6.2	9.4	6.4-13.4	3.0	2.9	1.6-4.9
	2001-2005	52.4	31.0	27.3-35.0	38.8	53.8	46.5-62.0	13.6	14.2	10.9-18.1	8.8	5.0	3.6-6.8	7.2	10.2	7.1-14.1	1.6	1.5	0.6-3.1
	2006-2010	51.8	29.3	25.8-33.1	37.2	48.8	42.0-56.5	14.6	14.9	11.6-19.0	10.6	5.8	4.3-7.7	6.6	8.8	6.1-12.5	4.0	3.6	2.1-5.8
	2011-2015	59.2	32.4	28.7-36.5	42.2	52.1	45.2-59.9	17.0	17.1	13.5-21.5	11.0	5.7	4.3-7.5	6.8	8.6	5.9-12.1	4.2	3.7	2.3-6.0
	2016-2020	54.8	28.6	25.2-32.3	41.4	48.0	41.6-55.3	13.4	12.4	9.5-16.0	6.8	3.4	2.4-4.9	5.2	6.1	4.0-9.1	1.6	1.4	0.6-3.0
Kidney and renal pelvis	1976-1980	9.4	6.1	4.4-8.1	6.2	9.2	6.2-13.3	3.2	3.9	2.2-6.4	5.0	3.1	2.0-4.6	1.8	2.6	1.2-5.0	3.2	3.4	1.9-5.6
	1981-1985	13.2	8.3	6.3-10.6	7.4	10.2	7.1-14.3	5.8	6.4	4.2-9.3	5.6	3.3	2.2-4.8	3.6	5.1	3.0-8.3	2.0	2.1	1.0-4.0
	1986-1990	13.8	8.7	6.7-11.1	9.0	13.0	9.4-17.4	4.8	5.1	3.2-7.7	8.6	5.5	3.9-7.4	5.8	8.7	5.8-12.6	2.8	2.7	1.5-4.7
	1991-1995	15.6	10.1	7.9-12.6	8.8	12.9	9.3-17.4	6.8	8.2	5.6-11.6	5.0	2.9	1.9-4.3	2.6	3.5	1.8-6.0	2.4	2.3	1.2-4.3
	1996-2000	16.2	10.0	7.9-12.5	9.8	13.8	10.2-18.3	6.4	7.2	4.9-10.4	6.4	3.7	2.5-5.3	3.6	5.3	3.1-8.4	2.8	2.9	1.5-4.9
	2001-2005	22.2	13.5	11.1-16.3	15.6	21.1	16.7-26.4	6.6	7.0	4.7-10.0	7.0	4.1	2.8-5.7	4.8	6.7	4.3-10.1	2.2	2.2	1.0-4.1
	2006-2010	30.8	18.2	15.4-21.4	18.4	23.8	19.2-29.4	12.4	13.5	10.3-17.5	6.4	3.5	2.4-5.0	4.8	6.2	3.9-9.3	1.6	1.4	0.6-3.1
	2011-2015	29.6	16.4	13.8-19.4	17.8	20.9	16.7-26.0	11.8	12.9	9.7-16.9	7.2	3.9	2.7-5.5	3.8	4.8	2.8-7.6	3.4	3.3	1.9-5.5
	2016-2020	33.6	20.1	17.0-23.7	23.6	30.0	24.5-36.4	10.0	10.9	7.9-14.8	5.6	2.8	1.8-4.2	4.0	4.8	2.9-7.7	1.6	1.2	0.5-2.7
Brain and other central nervous system	1976-1980	8.6	5.9	4.2-8.0	4.4	6.2	3.8-9.5	4.2	5.5	3.3-8.6	7.4	5.1	3.5-7.1	4.2	6.5	4.0-10.1	3.2	3.7	2.1-6.2
	1981-1985	9.6	6.8	5.0-9.1	4.8	7.5	4.7-11.3	4.8	6.3	3.9-9.4	6.4	4.4	3.0-6.3	2.6	4.0	2.1-7.0	3.8	4.5	2.7-7.2
	1986-1990	9.2	5.7	4.2-7.7	6.2	8.7	5.9-12.4	3.0	3.4	1.9-5.8	7.0	4.6	3.2-6.4	4.4	6.3	3.9-9.6	2.6	3.2	1.6-5.6
	1991-1995	10.4	6.8	5.0-8.9	5.0	7.1	4.6-10.5	5.4	6.7	4.3-9.8	5.6	3.5	2.3-5.1	3.2	4.6	2.6-7.5	2.4	2.6	1.3-4.6
	1996-2000	11.8	7.8	5.9-10.1	7.0	10.0	6.9-13.9	4.8	6.1	3.9-9.2	8.6	5.6	4.0-7.6	5.0	7.0	4.5-10.4	3.6	4.3	2.5-6.9
	2001-2005	10.2	6.7	4.9-8.8	5.2	7.5	4.9-11.0	5.0	6.1	3.9-9.2	5.6	3.4	2.3-5.0	2.8	3.8	2.1-6.4	2.8	3.0	1.6-5.2
	2006-2010	10.6	7.2	5.4-9.5	6.2	8.5	5.7-12.2	4.4	5.9	3.7-9.1	7.4	4.6	3.2-6.4	4.2	5.5	3.4-8.6	3.2	3.6	2.0-6.0
	2011-2015	11.2	7.5	5.6-9.9	6.0	8.7	5.8-12.5	5.2	6.5	4.1-9.8	6.4	3.8	2.6-5.6	3.4	4.9	2.7-8.0	3.0	3.2	1.7-5.5
	2016-2020	12.6	8.4	6.3-10.9	6.2	8.7	5.7-12.6	6.4	8.1	5.4-11.8	7.4	4.4	3.0-6.3	4.8	6.3	3.8-9.7	2.6	2.7	1.3-4.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chautauqua**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	5.0	3.6	2.3-5.4	1.8	3.4	1.5-6.4	3.2	4.1	2.3-6.7	1.4	0.9	0.4-1.9	0.6	1.1	0.2-3.2	0.8	0.9	0.2-2.4
	1981-1985	6.0	4.3	2.8-6.1	1.8	2.6	1.2-5.1	4.2	5.7	3.5-8.9	0.8	0.5	0.1-1.3	0.2	0.4	0.0-2.0	0.6	0.6	0.1-2.0
	1986-1990	4.8	3.3	2.1-5.0	1.8	3.0	1.3-5.7	3.0	3.8	2.1-6.3	1.2	0.7	0.3-1.6	0.2	0.2	0.0-1.5	1.0	1.0	0.3-2.5
	1991-1995	5.8	4.1	2.7-5.9	1.4	2.2	0.9-4.6	4.4	6.1	3.8-9.3	0.8	0.5	0.1-1.3	0.2	0.2	0.0-1.5	0.6	0.6	0.1-1.8
	1996-2000	7.6	5.4	3.8-7.5	2.2	3.3	1.6-5.9	5.4	7.8	5.1-11.3	1.0	0.6	0.2-1.4	0.6	0.9	0.2-2.7	0.4	0.4	0.1-1.7
	2001-2005	12.4	8.4	6.4-10.8	2.6	3.6	1.9-6.1	9.8	13.3	9.8-17.6	1.4	0.8	0.3-1.7	0.4	0.6	0.1-2.2	1.0	1.0	0.3-2.6
	2006-2010	16.6	12.0	9.5-15.0	3.8	5.5	3.2-8.6	12.8	18.6	14.1-23.9	0.4	0.2	0.0-0.8	0.2	0.2	0.0-1.5	0.2	0.1	0.0-1.3
	2011-2015	23.4	17.2	14.1-20.8	5.0	6.7	4.2-10.1	18.4	27.5	21.8-34.1	0.4	0.2	0.0-0.9	0.2	0.2	0.0-1.5	0.2	0.2	0.0-1.4
	2016-2020	23.4	16.4	13.4-19.9	4.8	6.1	3.8-9.4	18.6	26.6	21.1-33.1	1.4	0.7	0.3-1.5	0.4	0.4	0.1-1.9	1.0	0.8	0.2-2.3
Hodgkin lymphoma	1976-1980	4.4	2.8	1.8-4.4	2.6	3.4	1.7-6.0	1.8	2.3	1.0-4.5	2.0	1.4	0.7-2.6	1.2	2.2	0.8-4.8	0.8	1.0	0.3-2.6
	1981-1985	5.6	3.5	2.3-5.1	3.2	4.4	2.5-7.3	2.4	2.8	1.4-5.0	1.6	1.0	0.4-1.9	0.8	1.1	0.3-2.9	0.8	0.8	0.2-2.3
	1986-1990	3.6	2.4	1.4-3.9	1.8	2.6	1.2-5.0	1.8	2.4	1.1-4.7	1.8	1.3	0.6-2.4	1.2	1.8	0.7-4.0	0.6	0.8	0.2-2.4
	1991-1995	3.6	2.5	1.5-3.9	1.8	2.6	1.2-5.0	1.8	2.4	1.1-4.7	0.6	0.4	0.1-1.1	0.6	0.8	0.2-2.4	0.0	0.0	0.0-1.0
	1996-2000	4.6	3.2	2.0-4.9	1.8	2.6	1.2-4.9	2.8	4.0	2.2-6.7	0.4	0.3	0.0-1.0	0.2	0.3	0.0-1.7	0.2	0.3	0.0-1.5
	2001-2005	4.6	3.2	2.0-4.8	2.4	3.5	1.8-6.2	2.2	3.1	1.5-5.6	1.0	0.6	0.2-1.5	0.4	0.7	0.1-2.4	0.6	0.7	0.1-2.3
	2006-2010	5.0	3.5	2.2-5.2	3.0	4.4	2.4-7.3	2.0	2.5	1.2-4.8	0.4	0.2	0.0-0.9	0.2	0.2	0.0-1.5	0.2	0.2	0.0-1.4
	2011-2015	4.2	3.5	2.1-5.4	1.6	2.6	1.1-5.3	2.6	4.3	2.2-7.4	1.2	0.7	0.2-1.7	0.6	0.8	0.1-2.5	0.6	0.6	0.1-2.0
	2016-2020	2.8	2.4	1.2-4.0	0.6	0.9	0.2-2.8	2.2	3.8	1.8-6.9	0.6	0.3	0.1-1.1	0.2	0.3	0.0-1.7	0.4	0.3	0.0-1.6
Non-Hodgkin lymphomas	1976-1980	17.8	11.4	9.1-14.1	10.4	15.9	11.8-21.1	7.4	8.2	5.8-11.5	9.4	6.1	4.5-8.2	5.2	9.0	5.8-13.4	4.2	4.6	2.8-7.1
	1981-1985	20.8	13.4	10.9-16.3	10.0	14.7	10.8-19.5	10.8	12.4	9.2-16.3	9.4	5.8	4.3-7.8	4.4	6.6	4.1-10.1	5.0	5.4	3.5-8.1
	1986-1990	22.0	14.0	11.5-17.0	11.8	17.8	13.5-23.0	10.2	11.1	8.2-14.7	11.8	7.7	5.8-9.9	7.4	11.7	8.2-16.2	4.4	5.0	3.1-7.7
	1991-1995	24.0	15.1	12.5-18.1	11.2	16.4	12.3-21.3	12.8	14.0	10.7-18.1	12.0	7.1	5.4-9.2	4.4	6.7	4.2-10.2	7.6	7.5	5.2-10.4
	1996-2000	33.0	20.4	17.3-23.8	15.6	22.0	17.4-27.6	17.4	18.9	15.1-23.5	12.6	7.4	5.7-9.5	5.8	8.5	5.6-12.2	6.8	6.6	4.5-9.4
	2001-2005	30.2	18.3	15.5-21.5	16.6	23.3	18.5-28.9	13.6	14.7	11.3-18.8	11.2	6.3	4.8-8.3	7.0	10.0	7.0-14.0	4.2	3.8	2.3-5.9
	2006-2010	31.6	18.6	15.8-21.8	17.2	22.4	17.9-27.8	14.4	15.6	12.1-20.0	10.8	6.1	4.5-8.0	6.4	8.5	5.8-12.2	4.4	4.5	2.7-7.0
	2011-2015	31.6	17.9	15.2-21.1	15.8	19.9	15.6-24.9	15.8	16.8	13.1-21.2	10.4	5.9	4.3-7.8	5.0	6.7	4.3-10.0	5.4	5.3	3.4-8.1
	2016-2020	39.0	21.9	18.8-25.5	22.4	26.8	21.9-32.6	16.6	17.8	13.9-22.4	10.4	5.4	4.0-7.2	5.2	6.2	4.0-9.2	5.2	4.8	3.1-7.3
Myeloma	1976-1980	8.8	5.8	4.2-7.8	5.4	9.3	6.0-13.5	3.4	3.7	2.1-6.0	4.4	2.8	1.8-4.3	3.0	4.9	2.7-8.1	1.4	1.5	0.6-3.1
	1981-1985	9.8	6.2	4.5-8.2	5.8	9.1	6.0-13.2	4.0	4.1	2.5-6.4	5.0	3.2	2.0-4.7	2.4	4.2	2.1-7.3	2.6	2.6	1.4-4.5
	1986-1990	9.6	6.1	4.5-8.2	4.8	7.7	4.9-11.5	4.8	5.5	3.4-8.4	5.8	3.5	2.4-5.1	3.0	4.9	2.7-8.2	2.8	3.1	1.6-5.3
	1991-1995	8.4	5.1	3.7-7.0	5.4	8.1	5.3-11.8	3.0	3.0	1.7-5.2	5.4	3.3	2.2-4.9	3.2	4.7	2.7-7.7	2.2	2.4	1.1-4.4
	1996-2000	11.2	6.6	5.0-8.7	6.0	8.5	5.7-12.1	5.2	5.2	3.3-7.8	6.2	3.6	2.4-5.1	2.4	3.3	1.7-5.8	3.8	3.3	2.0-5.4
	2001-2005	12.8	7.6	5.8-9.8	8.0	10.9	7.8-14.9	4.8	4.9	3.1-7.4	6.2	3.6	2.4-5.1	4.0	5.6	3.4-8.8	2.2	2.3	1.1-4.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chautauqua**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	12.4	7.1	5.4-9.2	6.2	8.2	5.5-11.8	6.2	6.1	4.1-8.8	6.6	3.6	2.5-5.2	3.8	5.0	3.0-7.9	2.8	2.6	1.4-4.6
	2011-2015	11.4	6.5	4.9-8.5	5.2	6.1	3.9-9.1	6.2	6.6	4.4-9.6	5.8	3.0	2.0-4.5	3.0	3.6	2.0-6.1	2.8	2.7	1.5-4.8
	2016-2020	14.0	7.3	5.6-9.4	8.0	9.2	6.4-12.7	6.0	5.8	3.9-8.6	5.0	2.5	1.6-3.9	1.6	2.0	0.8-4.0	3.4	3.0	1.7-5.1
Leukemias	1976-1980	18.4	12.1	9.7-14.9	10.8	15.7	11.7-20.7	7.6	9.0	6.3-12.5	12.0	7.9	6.0-10.3	6.2	9.0	6.0-12.9	5.8	7.0	4.6-10.2
	1981-1985	23.6	15.1	12.4-18.1	14.6	21.3	16.6-27.0	9.0	10.7	7.7-14.5	12.6	8.1	6.2-10.4	8.2	12.3	8.7-16.8	4.4	5.2	3.2-8.0
	1986-1990	24.6	16.2	13.4-19.5	13.0	21.2	16.2-27.2	11.6	13.1	9.8-17.1	16.0	10.3	8.1-12.8	8.6	13.8	9.9-18.7	7.4	7.8	5.4-10.9
	1991-1995	19.6	11.9	9.7-14.6	10.6	15.4	11.5-20.2	9.0	9.6	6.9-13.1	15.4	9.4	7.4-11.8	7.4	11.4	8.0-15.8	8.0	8.4	5.9-11.6
	1996-2000	23.6	14.5	12.0-17.5	12.4	17.7	13.6-22.7	11.2	11.9	8.9-15.7	15.2	9.0	7.1-11.3	7.2	10.3	7.2-14.4	8.0	7.9	5.6-11.0
	2001-2005	21.0	13.0	10.6-15.8	11.2	16.1	12.1-20.9	9.8	10.6	7.7-14.3	9.8	5.8	4.3-7.7	5.2	7.5	4.9-11.0	4.6	4.6	2.8-7.1
	2006-2010	26.4	15.7	13.1-18.8	14.8	19.6	15.3-24.7	11.6	12.6	9.5-16.6	10.2	5.6	4.2-7.5	6.2	8.5	5.7-12.1	4.0	3.7	2.2-5.9
	2011-2015	29.4	16.7	14.0-19.8	16.0	20.2	15.9-25.3	13.4	14.0	10.6-18.1	11.4	6.0	4.5-7.9	7.0	8.7	6.0-12.3	4.4	3.9	2.4-6.3
2016-2020	28.6	16.2	13.5-19.3	16.2	19.5	15.3-24.6	12.4	13.1	9.9-17.2	9.2	4.7	3.4-6.4	5.2	6.2	4.0-9.3	4.0	3.3	2.0-5.4	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chemung**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	390.2	409.6	391.2-428.5	187.8	492.3	459.9-526.3	202.4	364.0	341.5-387.7	209.8	220.0	206.6-233.9	105.4	276.5	252.4-302.3	104.4	185.1	169.3-202.0
	1981-1985	423.8	437.9	419.0-457.3	205.8	529.0	495.8-563.8	218.0	391.5	367.8-416.3	214.2	219.6	206.4-233.4	111.8	295.8	270.7-322.5	102.4	175.3	160.1-191.6
	1986-1990	495.6	494.3	474.7-514.5	235.6	581.5	547.5-617.1	260.0	456.2	430.9-482.6	238.2	236.5	223.1-250.5	124.8	315.5	290.2-342.4	113.4	189.9	174.2-206.7
	1991-1995	539.0	515.6	496.1-535.8	279.8	644.3	610.1-679.9	259.2	442.5	418.0-468.2	240.2	227.0	214.2-240.3	117.4	283.0	259.8-307.6	122.8	196.3	180.7-213.0
	1996-2000	538.6	516.2	496.8-536.2	279.6	627.8	594.9-662.1	259.0	445.8	421.3-471.4	244.2	228.3	215.6-241.6	119.8	277.3	255.2-300.9	124.4	198.7	183.0-215.4
	2001-2005	555.8	522.2	502.9-542.2	281.2	609.5	577.7-642.5	274.6	467.0	442.0-493.2	224.2	203.9	192.1-216.4	108.4	242.3	222.0-263.9	115.8	181.2	166.3-197.1
	2006-2010	587.0	537.9	518.3-558.1	299.8	607.4	576.7-639.4	287.2	492.9	466.8-520.1	226.8	198.0	186.5-210.1	111.2	229.3	210.5-249.4	115.6	177.4	162.8-193.1
	2011-2015	546.4	487.3	468.6-506.5	287.6	557.6	528.6-587.9	258.8	438.6	413.9-464.4	203.4	170.9	160.3-182.0	107.0	207.7	190.2-226.4	96.4	143.4	130.5-157.5
	2016-2020	555.4	482.1	463.6-501.2	275.6	501.7	474.8-529.8	279.8	475.7	449.6-503.1	198.2	163.0	152.8-173.9	101.6	191.3	174.7-209.2	96.6	142.8	129.9-156.9
Oral cavity and pharynx	1976-1980	12.0	12.2	9.3-15.8	8.4	21.2	15.0-29.1	3.6	5.9	3.5-9.5	3.6	3.7	2.2-6.0	2.0	4.6	2.2-8.6	1.6	2.7	1.2-5.5
	1981-1985	10.4	10.8	8.0-14.3	6.8	16.1	10.9-22.8	3.6	6.8	3.9-10.9	5.4	5.6	3.7-8.2	3.2	8.1	4.5-13.4	2.2	4.1	2.0-7.4
	1986-1990	10.6	10.4	7.7-13.7	7.6	18.5	12.9-25.6	3.0	4.9	2.7-8.3	2.8	2.7	1.5-4.6	2.2	5.9	2.8-10.8	0.6	0.9	0.2-2.9
	1991-1995	12.0	12.4	9.4-16.0	7.6	18.4	12.9-25.4	4.4	8.0	4.9-12.3	4.0	3.7	2.3-5.8	2.4	5.2	2.7-9.2	1.6	2.6	1.1-5.4
	1996-2000	9.8	9.6	7.1-12.8	6.4	14.4	9.8-20.4	3.4	5.6	3.2-9.2	2.6	2.4	1.3-4.2	1.4	2.9	1.2-6.1	1.2	1.9	0.7-4.3
	2001-2005	11.4	10.9	8.2-14.1	7.6	15.6	11.1-21.5	3.8	6.4	3.8-10.2	1.6	1.4	0.6-2.9	1.0	2.0	0.6-4.8	0.6	0.8	0.2-2.7
	2006-2010	14.0	12.4	9.6-15.8	9.4	17.9	13.1-24.0	4.6	7.1	4.4-10.9	2.4	2.2	1.1-4.0	2.0	4.1	1.9-7.6	0.4	0.6	0.1-2.7
	2011-2015	12.0	10.9	8.2-14.2	10.0	19.4	14.3-25.9	2.0	3.0	1.4-6.0	2.0	1.9	0.9-3.5	1.8	3.7	1.7-7.1	0.2	0.2	0.0-2.0
	2016-2020	14.4	13.6	10.5-17.4	9.8	19.3	14.1-25.9	4.6	8.5	5.2-13.2	3.0	2.6	1.4-4.5	1.6	3.3	1.4-6.7	1.4	1.9	0.7-4.3
Esophagus	1976-1980	4.6	4.7	3.0-7.1	3.0	6.9	3.8-11.6	1.6	3.0	1.3-5.9	6.0	6.1	4.1-8.7	3.6	8.0	4.7-12.9	2.4	4.3	2.2-7.6
	1981-1985	4.0	4.2	2.5-6.5	2.0	5.3	2.5-9.7	2.0	3.3	1.6-6.3	3.8	3.7	2.2-5.8	2.4	5.3	2.7-9.5	1.4	2.2	0.9-4.6
	1986-1990	6.2	6.2	4.2-8.9	4.0	9.6	5.8-15.0	2.2	3.3	1.6-6.1	5.2	5.2	3.4-7.6	3.6	8.7	5.1-13.8	1.6	2.5	1.1-5.1
	1991-1995	6.4	6.0	4.1-8.6	4.6	10.6	6.6-16.0	1.8	2.8	1.3-5.5	4.8	4.5	2.9-6.8	3.8	8.4	5.0-13.3	1.0	1.6	0.5-4.0
	1996-2000	6.4	6.1	4.2-8.6	4.8	10.5	6.7-15.7	1.6	2.4	1.0-5.0	7.6	6.9	4.9-9.6	5.0	11.2	7.2-16.6	2.6	3.7	1.9-6.6
	2001-2005	5.4	5.2	3.4-7.7	4.2	8.7	5.4-13.4	1.2	1.9	0.7-4.5	5.2	5.0	3.2-7.3	4.2	8.8	5.4-13.5	1.0	1.4	0.4-3.7
	2006-2010	6.2	5.5	3.7-7.8	4.6	8.7	5.5-13.2	1.6	2.6	1.1-5.4	5.0	4.6	2.9-6.8	3.6	7.2	4.2-11.5	1.4	2.6	1.0-5.7
	2011-2015	7.4	6.2	4.4-8.7	5.8	11.1	7.4-16.1	1.6	2.4	1.0-5.0	7.6	6.3	4.4-8.7	6.6	11.9	8.2-17.0	1.0	1.5	0.5-3.9
	2016-2020	8.2	6.8	4.8-9.3	6.4	11.1	7.5-16.0	1.8	2.9	1.3-5.9	5.2	4.2	2.7-6.3	3.6	6.3	3.7-10.2	1.6	2.5	1.0-5.4
Stomach	1976-1980	7.8	8.5	6.0-11.6	4.8	13.1	8.3-19.7	3.0	5.7	3.1-9.5	6.0	6.8	4.6-9.7	3.2	9.4	5.3-15.3	2.8	5.3	2.9-9.0
	1981-1985	8.6	8.9	6.4-12.0	5.6	13.9	9.1-20.3	3.0	5.0	2.8-8.4	6.6	6.7	4.6-9.5	4.8	11.6	7.3-17.4	1.8	2.9	1.3-5.7
	1986-1990	8.8	9.1	6.6-12.2	4.6	12.3	7.6-18.6	4.2	6.8	4.1-10.5	7.0	7.4	5.1-10.3	3.2	9.0	5.0-14.7	3.8	6.2	3.7-9.9
	1991-1995	6.6	6.7	4.6-9.5	4.2	11.6	7.1-17.8	2.4	3.9	2.0-7.1	5.8	5.8	3.8-8.3	4.2	11.2	6.8-17.2	1.6	2.5	1.1-5.3
	1996-2000	7.2	6.8	4.8-9.5	4.6	10.5	6.6-15.8	2.6	4.3	2.2-7.5	3.8	3.6	2.1-5.6	2.2	5.3	2.6-9.6	1.6	2.6	1.1-5.4
	2001-2005	7.2	6.6	4.6-9.2	5.8	12.8	8.6-18.5	1.4	2.1	0.8-4.6	3.6	3.3	2.0-5.3	3.0	7.1	3.9-11.7	0.6	0.8	0.2-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chemung**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	8.4	7.8	5.6-10.6	5.0	10.5	6.7-15.6	3.4	5.9	3.4-9.6	4.0	3.6	2.2-5.7	2.4	5.1	2.6-9.0	1.6	2.7	1.2-5.5
	2011-2015	5.8	5.0	3.3-7.3	4.0	7.3	4.4-11.5	1.8	3.1	1.4-6.2	2.8	2.5	1.4-4.3	1.6	3.2	1.4-6.5	1.2	2.2	0.8-4.9
	2016-2020	8.2	6.8	4.8-9.4	4.8	9.0	5.7-13.7	3.4	4.9	2.9-8.3	2.6	2.1	1.1-3.7	1.8	3.2	1.4-6.4	0.8	1.0	0.3-3.2
Colorectal	1976-1980	66.8	71.4	63.8-79.6	29.8	77.3	64.9-91.4	37.0	66.4	57.0-76.9	33.2	35.4	30.2-41.3	14.4	37.5	29.0-47.6	18.8	33.6	27.0-41.2
	1981-1985	71.4	72.5	65.0-80.5	33.4	86.3	73.2-101.0	38.0	63.5	54.6-73.4	29.0	29.9	25.2-35.3	13.4	36.2	27.7-46.3	15.6	25.5	20.1-32.0
	1986-1990	70.4	69.6	62.4-77.4	33.2	85.2	72.2-99.8	37.2	60.4	51.9-70.0	28.4	28.3	23.8-33.4	13.8	35.2	27.0-44.9	14.6	23.3	18.2-29.5
	1991-1995	57.8	53.1	47.1-59.7	28.2	63.8	53.4-75.5	29.6	46.1	38.9-54.5	26.0	24.1	20.1-28.7	11.8	29.2	22.0-37.9	14.2	21.5	16.7-27.3
	1996-2000	65.6	60.7	54.3-67.7	34.0	75.5	64.5-87.9	31.6	49.5	42.0-58.2	24.6	22.5	18.7-26.9	12.4	27.2	20.8-35.0	12.2	17.8	13.6-23.1
	2001-2005	54.0	49.1	43.4-55.4	28.0	61.2	51.3-72.3	26.0	39.8	33.1-47.7	21.4	19.0	15.5-23.0	9.2	21.0	15.3-28.1	12.2	17.5	13.3-22.8
	2006-2010	52.2	46.8	41.2-53.0	28.0	56.5	47.4-66.8	24.2	38.3	31.6-46.3	17.6	15.0	12.0-18.5	8.4	17.1	12.3-23.2	9.2	13.5	9.8-18.3
	2011-2015	42.2	38.6	33.4-44.4	24.2	48.7	40.3-58.4	18.0	30.8	24.5-38.4	16.0	13.3	10.5-16.7	8.4	17.0	12.2-23.1	7.6	10.2	7.1-14.4
2016-2020	35.4	32.0	27.2-37.4	20.6	39.1	31.7-47.8	14.8	26.2	20.2-33.4	11.6	9.7	7.3-12.8	7.0	13.8	9.5-19.4	4.6	6.4	4.0-10.1	
Colon excluding rectum	1976-1980	50.0	54.1	47.5-61.4	21.0	56.3	45.6-68.7	29.0	52.0	43.7-61.3	28.6	30.6	25.7-36.1	12.0	32.3	24.4-42.0	16.6	29.4	23.3-36.6
	1981-1985	48.8	49.3	43.2-55.9	19.6	50.8	40.9-62.3	29.2	48.1	40.5-56.7	24.2	24.6	20.4-29.5	10.6	27.1	20.1-35.8	13.6	22.3	17.3-28.4
	1986-1990	48.6	48.2	42.3-54.8	21.2	55.3	44.8-67.3	27.4	45.2	37.8-53.7	24.2	24.1	20.0-28.9	11.8	30.4	22.8-39.6	12.4	20.0	15.2-25.8
	1991-1995	42.2	38.6	33.6-44.3	19.8	45.1	36.4-55.2	22.4	34.5	28.3-41.7	19.4	18.0	14.5-22.0	8.8	22.6	16.2-30.5	10.6	15.9	11.9-21.0
	1996-2000	47.8	43.8	38.4-49.8	23.6	52.1	43.0-62.5	24.2	37.5	31.0-45.0	17.8	16.2	13.0-20.0	8.0	17.3	12.3-23.6	9.8	14.4	10.6-19.2
	2001-2005	39.0	35.0	30.3-40.4	21.2	46.5	38.0-56.4	17.8	26.5	21.2-33.0	15.0	13.3	10.4-16.7	6.4	15.0	10.2-21.3	8.6	12.5	8.9-17.1
	2006-2010	38.6	34.2	29.5-39.5	19.6	39.9	32.4-48.8	19.0	29.4	23.6-36.4	14.8	12.6	9.9-16.0	6.8	13.9	9.6-19.6	8.0	11.6	8.2-16.1
	2011-2015	30.2	27.7	23.3-32.6	16.6	33.9	26.9-42.2	13.6	23.5	18.0-30.3	13.0	10.6	8.2-13.7	7.0	13.9	9.6-19.5	6.0	8.0	5.3-11.9
2016-2020	26.0	23.0	19.1-27.6	14.0	26.2	20.2-33.5	12.0	20.6	15.4-27.0	9.4	8.1	5.9-10.9	5.8	11.3	7.5-16.5	3.6	5.3	3.1-8.8	
Rectum & rectosigmoid	1976-1980	16.8	17.3	13.7-21.5	8.8	21.0	15.0-28.6	8.0	14.4	10.3-19.8	4.6	4.8	3.0-7.3	2.4	5.2	2.6-9.2	2.2	4.1	2.0-7.5
	1981-1985	22.6	23.2	19.0-28.0	13.8	35.5	27.3-45.3	8.8	15.4	11.1-20.8	4.8	5.3	3.4-7.9	2.8	9.0	4.9-15.1	2.0	3.2	1.5-6.0
	1986-1990	21.8	21.3	17.5-25.8	12.0	29.9	22.5-38.9	9.8	15.2	11.2-20.3	4.2	4.2	2.6-6.4	2.0	4.8	2.2-8.9	2.2	3.3	1.6-6.1
	1991-1995	15.6	14.5	11.4-18.1	8.4	18.7	13.4-25.4	7.2	11.7	8.1-16.4	6.6	6.1	4.2-8.7	3.0	6.6	3.6-11.1	3.6	5.6	3.3-9.0
	1996-2000	17.8	16.9	13.6-20.9	10.4	23.5	17.4-30.9	7.4	12.1	8.4-16.8	6.8	6.3	4.4-8.9	4.4	9.9	6.2-15.1	2.4	3.5	1.8-6.3
	2001-2005	15.0	14.0	11.0-17.7	6.8	14.7	10.1-20.6	8.2	13.3	9.4-18.4	6.4	5.7	3.9-8.1	2.8	6.0	3.2-10.1	3.6	5.0	2.9-8.3
	2006-2010	13.6	12.6	9.7-16.0	8.4	16.5	11.9-22.5	5.2	8.9	5.7-13.4	2.8	2.3	1.3-4.0	1.6	3.1	1.3-6.3	1.2	1.9	0.7-4.5
	2011-2015	12.0	11.0	8.3-14.3	7.6	14.8	10.4-20.6	4.4	7.3	4.5-11.4	3.0	2.7	1.5-4.5	1.4	3.1	1.2-6.5	1.6	2.1	0.9-4.7
2016-2020	9.4	9.0	6.5-12.2	6.6	12.9	8.8-18.4	2.8	5.6	2.9-9.8	2.2	1.7	0.8-3.2	1.2	2.4	0.9-5.5	1.0	1.1	0.4-3.3	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chemung**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	1.4	1.4	0.6-3.0	0.4	1.0	0.1-3.7	1.0	1.7	0.5-4.1	2.0	2.1	1.0-3.9	1.0	2.6	0.8-6.2	1.0	1.7	0.5-4.0
	1981-1985	1.0	0.9	0.3-2.2	0.6	1.2	0.3-3.8	0.4	0.6	0.1-2.5	1.6	1.6	0.7-3.2	0.4	0.8	0.1-3.1	1.2	2.2	0.8-5.0
	1986-1990	2.6	2.6	1.4-4.5	1.8	4.5	2.0-8.6	0.8	1.2	0.3-3.4	2.4	2.5	1.3-4.4	1.8	4.9	2.2-9.4	0.6	0.9	0.2-3.0
	1991-1995	6.0	5.8	3.9-8.3	4.6	11.4	7.1-17.3	1.4	2.0	0.8-4.3	5.0	4.7	3.0-7.0	3.2	8.3	4.6-13.6	1.8	2.7	1.2-5.4
	1996-2000	4.0	3.9	2.4-6.1	2.0	4.7	2.2-8.7	2.0	3.4	1.6-6.5	3.4	3.2	1.9-5.1	2.2	5.0	2.5-9.1	1.2	1.9	0.7-4.3
	2001-2005	5.8	5.3	3.6-7.7	4.0	8.5	5.2-13.2	1.8	2.7	1.2-5.4	4.2	3.8	2.4-5.9	2.8	6.2	3.4-10.5	1.4	2.2	0.9-4.7
	2006-2010	4.8	4.4	2.8-6.6	4.0	8.2	5.0-12.8	0.8	1.1	0.3-3.3	4.0	3.5	2.1-5.5	2.8	5.6	3.0-9.5	1.2	1.7	0.6-4.0
	2011-2015	6.4	5.6	3.8-8.0	5.2	10.0	6.5-14.8	1.2	1.8	0.7-4.3	5.8	5.0	3.3-7.3	3.8	7.5	4.5-11.8	2.0	2.9	1.4-5.6
	2016-2020	6.6	5.1	3.5-7.4	4.6	7.6	4.8-11.7	2.0	3.1	1.4-6.1	5.8	4.4	2.9-6.4	4.2	7.2	4.4-11.4	1.6	2.1	0.9-4.6
Pancreas	1976-1980	10.4	10.9	8.1-14.4	5.8	14.3	9.4-20.8	4.6	8.3	5.2-12.5	9.6	10.0	7.3-13.3	5.0	12.3	7.9-18.4	4.6	8.0	5.1-12.1
	1981-1985	11.2	11.7	8.8-15.3	5.4	15.3	9.9-22.4	5.8	9.4	6.3-13.7	11.6	12.0	9.1-15.6	5.4	14.7	9.5-21.6	6.2	10.9	7.3-15.7
	1986-1990	12.2	12.0	9.2-15.5	5.4	15.0	9.7-22.1	6.8	10.6	7.3-15.1	12.6	12.7	9.7-16.3	6.8	18.0	12.2-25.4	5.8	9.4	6.2-13.7
	1991-1995	11.2	10.5	7.9-13.7	5.6	12.7	8.3-18.5	5.6	7.9	5.3-11.7	10.4	9.4	7.0-12.4	4.4	10.2	6.3-15.7	6.0	8.4	5.7-12.2
	1996-2000	16.0	14.9	11.8-18.6	8.4	18.6	13.4-25.3	7.6	11.8	8.3-16.5	16.0	14.8	11.8-18.5	8.2	18.2	13.0-24.8	7.8	11.9	8.4-16.5
	2001-2005	17.0	15.4	12.3-19.1	5.0	11.3	7.2-16.7	12.0	18.7	14.2-24.4	16.6	14.9	11.9-18.6	5.6	13.0	8.6-18.9	11.0	16.7	12.5-22.0
	2006-2010	15.8	13.6	10.8-17.1	7.8	15.6	11.0-21.4	8.0	12.1	8.6-16.8	11.6	10.0	7.6-13.0	4.2	8.5	5.3-13.2	7.4	11.1	7.7-15.6
	2011-2015	20.2	16.8	13.7-20.6	10.4	19.8	14.7-26.2	9.8	14.4	10.6-19.4	14.4	11.9	9.3-15.1	6.8	13.3	9.1-18.7	7.6	10.9	7.6-15.3
	2016-2020	15.0	12.4	9.6-15.7	6.8	11.9	8.2-16.8	8.2	13.1	9.2-18.3	13.4	10.7	8.3-13.8	6.8	12.3	8.5-17.5	6.6	9.4	6.4-13.7
Larynx	1976-1980	5.6	6.0	3.9-8.7	4.2	10.2	6.2-15.8	1.4	2.4	1.0-5.1	2.6	2.5	1.3-4.4	1.8	4.1	1.8-8.0	0.8	1.3	0.4-3.5
	1981-1985	7.4	7.5	5.2-10.4	6.2	15.1	10.1-21.7	1.2	1.9	0.7-4.3	3.4	3.3	1.9-5.4	3.0	6.7	3.7-11.2	0.4	0.8	0.1-3.0
	1986-1990	7.8	7.7	5.4-10.6	6.0	13.6	9.0-19.7	1.8	3.0	1.4-6.0	3.0	3.2	1.8-5.2	2.2	6.0	2.9-10.7	0.8	1.3	0.3-3.5
	1991-1995	3.4	3.1	1.8-5.0	2.4	5.0	2.6-8.9	1.0	1.5	0.5-3.7	2.4	2.3	1.2-4.0	2.0	4.4	2.1-8.1	0.4	0.7	0.1-2.6
	1996-2000	5.8	5.7	3.8-8.3	3.8	8.2	4.9-12.8	2.0	3.5	1.6-6.6	1.2	1.1	0.4-2.5	0.8	1.7	0.5-4.5	0.4	0.6	0.1-2.5
	2001-2005	5.2	4.8	3.1-7.1	4.2	8.6	5.3-13.2	1.0	1.5	0.5-3.8	1.0	0.9	0.3-2.2	0.8	1.7	0.5-4.4	0.2	0.2	0.0-1.9
	2006-2010	5.8	5.4	3.6-7.8	4.2	8.3	5.1-12.9	1.6	2.5	1.1-5.2	1.0	0.9	0.3-2.2	0.6	1.3	0.3-3.8	0.4	0.7	0.1-2.8
	2011-2015	4.8	4.1	2.6-6.2	4.2	7.5	4.6-11.7	0.6	1.0	0.2-3.3	1.0	0.9	0.3-2.3	0.6	1.3	0.3-3.9	0.4	0.7	0.1-2.7
	2016-2020	4.8	4.0	2.5-6.2	3.6	6.6	3.8-10.8	1.2	1.7	0.6-4.1	1.0	0.9	0.3-2.2	1.0	1.9	0.6-4.8	0.0	0.0	0.0-1.8
Lung and bronchus	1976-1980	67.6	68.5	61.3-76.3	46.0	118.1	102.7-135.0	21.6	37.4	30.5-45.3	52.8	54.0	47.6-61.0	36.4	94.9	81.0-110.4	16.4	28.5	22.5-35.5
	1981-1985	74.4	74.4	66.9-82.5	50.4	126.3	110.5-143.5	24.0	41.3	34.1-49.7	60.0	59.6	53.0-66.9	40.2	102.2	87.9-118.0	19.8	33.3	26.9-40.8
	1986-1990	91.8	89.5	81.3-98.3	58.4	135.9	120.3-153.0	33.4	58.6	49.8-68.6	66.8	65.0	58.1-72.6	45.6	109.9	95.6-125.8	21.2	36.1	29.4-44.0
	1991-1995	83.6	79.8	72.2-87.9	48.8	112.3	98.4-127.7	34.8	56.4	48.1-65.7	69.4	65.6	58.8-73.0	40.6	93.5	80.8-107.6	28.8	46.8	39.3-55.4
	1996-2000	92.6	87.1	79.3-95.5	50.4	113.0	99.3-128.1	42.2	69.5	60.3-79.9	74.4	69.5	62.6-77.0	40.4	91.8	79.4-105.6	34.0	54.9	46.8-64.1
	2001-2005	86.2	79.2	71.9-87.1	44.8	98.4	85.8-112.3	41.4	68.4	59.2-78.7	67.2	61.4	55.0-68.5	36.8	81.4	70.0-94.3	30.4	50.4	42.6-59.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chemung**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	97.2	86.2	78.6-94.4	50.2	102.2	89.8-115.8	47.0	75.2	65.7-85.8	76.0	66.3	59.7-73.5	40.8	83.6	72.4-96.0	35.2	53.6	45.8-62.5
	2011-2015	84.4	72.9	65.9-80.3	44.2	84.3	73.4-96.5	40.2	64.3	55.5-74.3	56.2	46.8	41.4-52.7	29.4	55.3	46.6-65.3	26.8	40.7	34.0-48.6
	2016-2020	92.8	75.7	68.8-83.1	48.0	86.4	75.6-98.4	44.8	67.6	58.8-77.6	57.4	46.7	41.4-52.7	31.2	57.3	48.5-67.4	26.2	38.7	32.2-46.3
Melanoma of the skin	1976-1980	7.4	8.6	6.0-11.9	4.0	11.6	7.0-18.0	3.4	6.6	3.8-10.7	1.0	1.0	0.3-2.5	0.8	1.7	0.4-4.7	0.2	0.4	0.0-2.1
	1981-1985	7.6	8.8	6.1-12.1	4.2	10.0	6.0-15.6	3.4	7.8	4.4-12.7	3.2	3.1	1.8-5.2	1.8	4.6	2.0-9.0	1.4	2.6	1.0-5.4
	1986-1990	8.6	8.9	6.4-12.1	3.8	8.7	5.2-13.7	4.8	9.7	6.1-14.5	2.2	2.3	1.1-4.2	1.4	3.3	1.3-6.8	0.8	1.5	0.4-4.0
	1991-1995	12.8	12.6	9.7-16.1	7.4	16.6	11.6-23.0	5.4	9.7	6.3-14.3	2.2	2.2	1.1-4.0	1.2	2.8	1.0-6.2	1.0	1.8	0.6-4.4
	1996-2000	16.0	16.4	13.0-20.4	7.4	16.6	11.6-23.0	8.6	17.4	12.5-23.6	2.6	2.5	1.3-4.4	1.6	3.5	1.5-6.9	1.0	2.0	0.6-4.8
	2001-2005	17.8	17.8	14.3-22.0	10.0	21.3	15.8-28.2	7.8	15.1	10.7-20.9	2.8	2.7	1.5-4.6	1.6	3.9	1.7-7.7	1.2	2.2	0.8-5.0
	2006-2010	24.8	24.1	19.9-28.9	12.2	24.8	18.9-32.0	12.6	24.6	18.6-31.9	3.2	3.2	1.8-5.3	2.0	4.3	2.0-7.9	1.2	2.5	0.9-5.8
	2011-2015	23.6	22.0	18.1-26.6	14.2	28.2	21.9-35.9	9.4	17.5	12.6-23.8	2.6	2.1	1.1-3.7	1.8	3.2	1.5-6.3	0.8	1.1	0.3-3.3
2016-2020	23.2	21.1	17.3-25.6	15.4	28.4	22.3-35.9	7.8	16.0	11.1-22.3	3.4	2.9	1.7-4.8	2.4	4.4	2.2-7.9	1.0	1.6	0.5-4.2	
Female breast	1976-1980							47.8	89.3	78.0-101.7							16.0	30.0	23.7-37.6
	1981-1985							53.8	102.2	89.8-115.7							15.6	27.7	21.8-34.9
	1986-1990							72.2	133.1	119.2-148.2							18.2	32.7	26.1-40.4
	1991-1995							76.0	137.2	123.4-152.3							20.6	35.2	28.6-43.1
	1996-2000							68.0	122.2	109.3-136.3							20.0	34.1	27.6-41.7
	2001-2005							78.2	138.6	124.9-153.4							14.6	22.9	17.8-29.2
	2006-2010							74.2	130.2	116.9-144.7							13.2	20.6	15.8-26.7
	2011-2015							73.4	124.1	111.2-138.1							13.0	20.0	15.2-26.0
2016-2020							81.0	140.8	126.5-156.3							11.4	16.3	12.2-21.6	
Cervix uteri	1976-1980							6.6	13.9	9.4-19.8							4.2	7.9	4.8-12.2
	1981-1985							6.2	11.5	7.7-16.5							4.2	8.1	4.9-12.7
	1986-1990							6.2	12.6	8.4-18.1							2.0	3.6	1.7-6.8
	1991-1995							7.2	14.3	10.0-20.0							3.2	5.9	3.3-9.7
	1996-2000							5.2	11.3	7.3-16.6							2.2	4.4	2.2-7.9
	2001-2005							3.4	7.7	4.4-12.4							0.6	0.9	0.2-3.1
	2006-2010							3.2	7.2	4.0-11.8							0.8	1.5	0.4-4.0
	2011-2015							3.0	7.9	4.3-13.0							1.4	2.8	1.0-6.0
2016-2020							2.6	5.4	2.7-9.6							1.4	2.7	1.0-6.0	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chemung**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							16.8	29.9	23.7-37.1							3.0	5.2	2.9-8.6
	1981-1985							19.8	34.2	27.6-41.9							4.8	7.9	5.0-11.9
	1986-1990							19.2	33.0	26.5-40.6							3.8	6.1	3.6-9.7
	1991-1995							17.2	30.6	24.3-38.1							4.8	7.4	4.7-11.2
	1996-2000							14.0	26.0	20.2-33.0							2.0	3.2	1.5-6.2
	2001-2005							14.6	24.9	19.4-31.6							3.0	4.4	2.4-7.6
	2006-2010							18.4	32.2	25.7-39.8							2.8	4.0	2.2-7.0
	2011-2015							17.2	29.0	23.0-36.3							3.2	4.6	2.6-7.8
	2016-2020							20.4	33.1	26.7-40.8							2.8	4.5	2.4-8.0
Ovary	1976-1980							11.4	19.6	14.8-25.6							7.6	12.9	9.1-17.8
	1981-1985							9.2	18.9	13.6-25.5							5.8	10.2	6.8-14.9
	1986-1990							9.0	16.5	11.8-22.3							4.8	7.7	4.9-11.7
	1991-1995							9.6	17.0	12.4-22.8							6.8	11.3	7.7-16.0
	1996-2000							10.0	17.1	12.6-22.8							9.0	13.7	9.9-18.5
	2001-2005							9.4	16.6	12.1-22.3							5.6	8.8	5.8-13.0
	2006-2010							11.8	20.8	15.7-27.1							8.6	13.5	9.7-18.5
	2011-2015							9.0	14.8	10.6-20.2							8.2	12.7	9.0-17.6
	2016-2020							6.0	9.8	6.4-14.5							4.8	6.6	4.2-10.2
Prostate	1976-1980				31.2	88.4	74.6-104.0							12.2	36.4	27.4-47.1			
	1981-1985				35.6	98.9	84.3-115.2							11.4	36.1	27.1-47.0			
	1986-1990				41.4	106.2	91.6-122.4							12.8	36.6	27.8-47.1			
	1991-1995				91.8	210.1	190.8-230.9							17.0	45.5	36.0-56.6			
	1996-2000				77.8	175.8	158.4-194.5							10.0	27.1	19.9-35.9			
	2001-2005				79.2	170.5	154.0-188.3							11.0	26.3	19.7-34.4			
	2006-2010				79.6	157.9	142.6-174.4							11.2	24.6	18.6-32.1			
	2011-2015				62.2	113.2	100.7-126.9							10.4	20.9	15.6-27.6			
	2016-2020				62.6	104.2	92.7-116.8							7.6	14.2	10.0-19.7			
Testis	1976-1980				1.6	3.8	1.5-7.6							0.0	0.0	0.0-1.9			
	1981-1985				2.8	7.0	3.7-11.9							0.2	0.5	0.0-2.9			
	1986-1990				2.2	4.5	2.2-8.2							0.0	0.0	0.0-1.8			
	1991-1995				3.2	6.9	3.9-11.2							0.2	0.4	0.0-2.3			
	1996-2000				2.8	6.1	3.3-10.3							0.0	0.0	0.0-1.7			
	2001-2005				3.4	8.0	4.7-12.8							0.2	0.5	0.0-2.6			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chemung**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				2.2	5.3	2.6-9.4							0.2	0.5	0.0-2.6			
	2011-2015				3.8	9.0	5.4-14.1							0.8	2.0	0.5-5.0			
	2016-2020				2.2	5.0	2.4-9.1							0.0	0.0	0.0-1.8			
Urinary bladder (incl. in situ)	1976-1980	16.2	16.9	13.4-21.1	11.4	31.7	23.7-41.5	4.8	8.1	5.2-12.1	5.8	6.3	4.2-9.1	3.6	10.4	6.0-16.7	2.2	3.8	1.9-6.9
	1981-1985	19.6	20.0	16.2-24.5	15.0	37.2	29.0-47.0	4.6	8.2	5.2-12.5	3.4	3.5	2.0-5.6	2.6	7.1	3.6-12.3	0.8	1.3	0.4-3.6
	1986-1990	25.8	24.8	20.6-29.5	18.2	44.8	35.8-55.3	7.6	11.9	8.4-16.6	6.0	5.8	3.9-8.3	4.4	11.5	7.0-17.5	1.6	2.4	1.1-5.0
	1991-1995	28.0	26.3	22.1-31.1	20.0	46.2	37.4-56.4	8.0	12.2	8.7-16.9	6.6	6.0	4.1-8.5	3.4	7.5	4.3-12.2	3.2	4.5	2.6-7.6
	1996-2000	33.0	31.0	26.4-36.1	23.0	52.5	43.2-63.3	10.0	16.6	12.2-22.1	5.4	4.8	3.2-7.0	3.6	9.2	5.3-14.6	1.8	2.5	1.1-5.0
	2001-2005	33.0	29.9	25.5-34.9	26.8	59.2	49.5-70.3	6.2	10.0	6.7-14.4	3.8	3.2	1.9-5.1	2.6	6.8	3.6-11.7	1.2	1.6	0.6-3.9
	2006-2010	35.6	31.7	27.1-36.8	29.2	60.5	51.0-71.3	6.4	10.4	7.1-14.9	7.4	6.5	4.6-9.0	5.4	11.1	7.3-16.3	2.0	3.1	1.5-6.0
	2011-2015	36.6	31.1	26.7-36.1	27.6	52.5	43.9-62.3	9.0	14.0	10.1-19.0	7.0	5.4	3.7-7.6	3.8	7.1	4.3-11.3	3.2	4.3	2.4-7.4
	2016-2020	29.4	24.0	20.2-28.4	20.8	38.1	31.0-46.5	8.6	13.1	9.4-18.1	6.6	5.3	3.6-7.6	4.2	8.0	4.9-12.5	2.4	3.7	1.9-6.8
Kidney and renal pelvis	1976-1980	5.4	5.3	3.5-7.7	4.8	11.1	7.0-16.8	0.6	0.9	0.2-3.0	3.8	3.7	2.2-5.9	2.8	6.9	3.7-11.8	1.0	1.6	0.5-4.0
	1981-1985	10.4	10.9	8.0-14.3	6.4	16.1	10.8-23.0	4.0	7.5	4.5-11.9	3.4	3.5	2.0-5.6	2.4	7.0	3.5-12.2	1.0	1.6	0.5-3.9
	1986-1990	12.0	11.8	8.9-15.2	5.4	13.0	8.4-19.0	6.6	10.8	7.3-15.4	5.8	5.4	3.6-7.9	2.2	5.2	2.5-9.5	3.6	5.7	3.3-9.2
	1991-1995	12.8	12.5	9.6-16.1	8.6	19.7	14.2-26.6	4.2	7.2	4.4-11.3	5.0	4.5	2.9-6.7	3.2	7.5	4.1-12.3	1.8	2.7	1.2-5.3
	1996-2000	17.2	16.9	13.5-20.9	9.4	21.3	15.6-28.4	7.8	13.1	9.3-18.2	5.0	4.9	3.1-7.2	2.6	6.0	3.2-10.4	2.4	3.7	1.9-6.7
	2001-2005	15.6	14.9	11.7-18.6	9.8	21.0	15.5-27.8	5.8	9.4	6.2-13.7	5.2	4.8	3.2-7.2	3.2	6.6	3.8-10.9	2.0	3.0	1.4-5.8
	2006-2010	18.8	17.1	13.8-21.1	10.6	20.6	15.4-27.1	8.2	14.5	10.3-20.1	3.4	3.0	1.7-4.8	1.6	3.3	1.4-6.6	1.8	2.6	1.2-5.3
	2011-2015	19.4	17.4	14.0-21.4	11.6	22.4	16.9-29.3	7.8	12.7	8.9-17.6	6.2	5.0	3.4-7.3	4.4	8.0	5.0-12.3	1.8	2.4	1.1-5.0
	2016-2020	19.6	17.2	13.8-21.2	12.8	23.3	17.8-30.1	6.8	12.0	8.0-17.3	4.6	4.1	2.6-6.3	2.4	4.9	2.5-8.8	2.2	3.6	1.7-7.0
Brain and other central nervous system	1976-1980	9.8	10.5	7.7-14.0	6.0	14.9	10.0-21.5	3.8	6.6	3.9-10.4	6.8	7.0	4.8-9.9	4.2	10.0	6.1-15.5	2.6	4.4	2.3-7.6
	1981-1985	7.6	8.2	5.7-11.3	4.0	9.4	5.6-14.7	3.6	7.2	4.2-11.5	3.6	4.0	2.3-6.4	2.4	6.1	3.0-10.8	1.2	2.3	0.8-5.1
	1986-1990	6.6	6.8	4.7-9.7	2.6	5.8	3.0-10.2	4.0	7.9	4.8-12.3	4.6	4.5	2.8-6.7	2.4	5.1	2.6-9.1	2.2	3.8	1.9-6.9
	1991-1995	7.8	8.0	5.6-10.9	3.0	7.1	3.9-11.7	4.8	9.6	6.1-14.5	6.0	5.9	4.0-8.5	2.2	5.1	2.5-9.3	3.8	7.0	4.1-11.1
	1996-2000	6.8	6.9	4.8-9.7	4.0	8.7	5.3-13.5	2.8	4.9	2.6-8.5	4.4	4.5	2.8-6.9	2.4	5.3	2.7-9.3	2.0	3.9	1.8-7.3
	2001-2005	10.0	10.2	7.5-13.5	6.0	13.1	8.8-18.8	4.0	7.8	4.7-12.3	6.0	6.0	4.0-8.6	3.0	6.4	3.6-10.6	3.0	5.7	3.2-9.5
	2006-2010	9.2	9.1	6.6-12.2	4.6	9.7	6.1-14.7	4.6	8.4	5.2-12.9	5.8	5.4	3.6-7.8	2.8	6.0	3.2-10.1	3.0	5.0	2.7-8.4
	2011-2015	10.0	10.5	7.7-14.0	6.4	13.2	8.9-18.9	3.6	8.3	4.8-13.1	5.4	5.0	3.2-7.4	4.0	7.5	4.5-11.9	1.4	2.6	1.0-5.7
	2016-2020	6.6	6.5	4.4-9.3	4.2	8.6	5.2-13.4	2.4	4.5	2.2-8.2	5.6	5.1	3.3-7.5	3.6	7.3	4.2-11.8	2.0	3.0	1.4-6.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chemung**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	3.2	3.1	1.7-5.1	0.4	1.0	0.1-3.7	2.8	4.7	2.6-8.0	0.8	0.8	0.2-2.0	0.0	0.0	0.0-1.9	0.8	1.3	0.4-3.5
	1981-1985	3.4	3.7	2.1-6.0	0.8	1.5	0.4-4.2	2.6	5.4	2.8-9.3	0.6	0.7	0.1-2.0	0.2	0.6	0.0-3.1	0.4	0.7	0.1-2.6
	1986-1990	1.6	1.6	0.7-3.2	0.4	0.8	0.1-3.1	1.2	2.3	0.8-5.1	0.4	0.3	0.0-1.3	0.2	0.4	0.0-2.4	0.2	0.3	0.0-2.0
	1991-1995	3.8	3.8	2.3-6.0	0.4	1.0	0.1-3.6	3.4	6.3	3.6-10.2	0.4	0.4	0.0-1.4	0.0	0.0	0.0-1.7	0.4	0.7	0.1-2.7
	1996-2000	4.6	4.9	3.1-7.4	1.4	3.0	1.2-6.2	3.2	7.0	4.0-11.4	0.6	0.5	0.1-1.6	0.2	0.4	0.0-2.4	0.4	0.6	0.1-2.5
	2001-2005	10.2	10.8	8.0-14.3	2.0	4.0	1.9-7.5	8.2	17.6	12.5-24.0	0.2	0.2	0.0-1.1	0.0	0.0	0.0-1.7	0.2	0.3	0.0-2.1
	2006-2010	13.2	14.4	11.1-18.4	1.8	4.0	1.8-7.6	11.4	25.3	19.0-33.0	0.4	0.3	0.0-1.4	0.0	0.0	0.0-1.7	0.4	0.6	0.1-2.5
	2011-2015	13.6	14.9	11.5-19.1	3.2	7.2	4.1-11.8	10.4	23.0	16.9-30.4	0.0	0.0	0.0-0.9	0.0	0.0	0.0-1.7	0.0	0.0	0.0-1.7
2016-2020	16.6	18.8	14.8-23.5	4.4	9.8	6.0-15.0	12.2	28.0	21.1-36.6	0.8	0.6	0.2-1.7	0.0	0.0	0.0-1.8	0.8	0.9	0.2-3.0	
Hodgkin lymphoma	1976-1980	2.6	2.2	1.2-3.9	1.4	2.6	1.0-5.7	1.2	2.0	0.7-4.5	1.4	1.2	0.5-2.6	1.2	2.3	0.8-5.2	0.2	0.3	0.0-2.0
	1981-1985	2.6	2.3	1.2-4.1	1.2	2.1	0.8-5.0	1.4	2.4	1.0-5.0	0.6	0.5	0.1-1.7	0.4	0.8	0.1-3.1	0.2	0.3	0.0-2.1
	1986-1990	3.4	3.4	2.0-5.5	1.4	2.7	1.1-5.8	2.0	4.1	2.0-7.7	0.4	0.4	0.0-1.5	0.2	0.4	0.0-2.4	0.2	0.3	0.0-2.1
	1991-1995	2.4	2.7	1.4-4.7	1.4	3.1	1.2-6.5	1.0	2.4	0.8-5.4	0.6	0.7	0.1-1.9	0.4	0.9	0.1-3.2	0.2	0.5	0.0-2.5
	1996-2000	1.8	2.0	0.9-3.8	1.2	2.6	1.0-5.8	0.6	1.3	0.3-3.8	0.2	0.2	0.0-1.1	0.0	0.0	0.0-1.7	0.2	0.3	0.0-2.0
	2001-2005	2.4	2.5	1.3-4.4	1.4	3.0	1.2-6.3	1.0	2.0	0.6-4.8	0.4	0.4	0.0-1.5	0.4	0.9	0.1-3.2	0.0	0.0	0.0-1.6
	2006-2010	3.4	4.0	2.3-6.4	2.0	4.8	2.3-8.8	1.4	3.1	1.2-6.5	0.6	0.6	0.1-1.8	0.4	0.9	0.1-3.2	0.2	0.3	0.0-2.2
	2011-2015	3.0	3.7	2.0-6.1	2.0	5.0	2.4-9.2	1.0	2.2	0.7-5.2	0.2	0.2	0.0-1.1	0.0	0.0	0.0-1.7	0.2	0.3	0.0-2.2
2016-2020	2.0	2.2	1.0-4.2	1.2	2.7	1.0-6.1	0.8	1.8	0.4-4.8	0.4	0.3	0.0-1.3	0.2	0.3	0.0-2.3	0.2	0.3	0.0-2.2	
Non-Hodgkin lymphomas	1976-1980	10.4	10.9	8.1-14.4	5.6	13.5	8.9-19.8	4.8	8.2	5.2-12.3	5.8	6.2	4.1-9.0	2.6	6.5	3.3-11.4	3.2	5.8	3.3-9.6
	1981-1985	10.6	10.7	8.0-14.1	4.4	10.2	6.3-15.7	6.2	10.7	7.2-15.4	4.8	5.2	3.3-7.8	2.2	5.4	2.6-9.8	2.6	4.5	2.3-7.9
	1986-1990	17.8	18.2	14.6-22.5	10.0	24.7	18.1-32.8	7.8	14.0	9.8-19.5	8.4	8.6	6.2-11.7	4.6	11.5	7.2-17.5	3.8	6.7	4.0-10.7
	1991-1995	17.2	17.3	13.8-21.4	7.4	17.0	11.9-23.5	9.8	16.9	12.4-22.7	7.8	7.4	5.2-10.2	2.8	6.8	3.6-11.5	5.0	8.0	5.1-12.0
	1996-2000	19.2	18.6	15.1-22.8	11.0	24.6	18.5-32.2	8.2	13.5	9.6-18.5	8.8	8.3	6.0-11.1	4.8	11.3	7.2-16.9	4.0	5.9	3.6-9.4
	2001-2005	22.4	21.5	17.7-25.9	12.4	26.2	20.1-33.6	10.0	17.1	12.5-22.8	8.8	8.1	5.9-11.0	4.0	8.4	5.1-13.0	4.8	7.7	4.9-11.7
	2006-2010	25.4	23.2	19.3-27.7	14.4	29.6	23.1-37.4	11.0	18.0	13.5-23.7	9.4	8.0	5.9-10.7	6.0	11.7	7.9-16.9	3.4	5.0	2.8-8.4
	2011-2015	23.0	20.3	16.6-24.5	12.0	24.2	18.3-31.4	11.0	16.2	12.2-21.5	8.0	6.4	4.5-8.8	4.0	8.1	4.9-12.7	4.0	4.8	2.9-7.9
2016-2020	24.2	21.5	17.6-25.9	13.4	25.7	19.8-33.0	10.8	18.0	13.3-24.0	7.6	6.4	4.5-8.9	4.6	9.4	5.9-14.3	3.0	4.5	2.5-7.7	
Myeloma	1976-1980	4.6	4.7	3.0-7.1	2.0	5.1	2.4-9.4	2.6	4.4	2.3-7.6	2.0	2.2	1.1-4.0	1.0	2.7	0.9-6.3	1.0	1.8	0.6-4.3
	1981-1985	4.2	4.3	2.6-6.6	2.6	6.7	3.5-11.7	1.6	2.6	1.1-5.3	2.8	3.0	1.6-5.1	1.6	4.5	1.9-8.9	1.2	1.9	0.7-4.3
	1986-1990	4.4	4.4	2.8-6.7	2.0	5.5	2.5-10.2	2.4	3.6	1.9-6.5	4.2	4.3	2.6-6.5	1.6	4.4	1.8-8.7	2.6	4.0	2.1-7.0
	1991-1995	8.0	7.6	5.4-10.4	4.4	9.9	6.1-15.0	3.6	5.7	3.3-9.3	3.0	2.8	1.6-4.7	1.8	4.8	2.1-9.2	1.2	1.9	0.7-4.4
	1996-2000	5.4	5.1	3.3-7.4	3.8	8.3	5.0-13.1	1.6	2.5	1.1-5.1	4.8	4.4	2.8-6.6	2.8	6.4	3.5-10.8	2.0	2.7	1.3-5.3
	2001-2005	7.0	6.5	4.5-9.1	4.0	8.2	5.0-12.7	3.0	4.9	2.7-8.3	4.4	4.0	2.5-6.2	2.4	4.9	2.5-8.6	2.0	3.3	1.5-6.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chemung**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	6.6	6.0	4.1-8.6	3.4	7.0	4.1-11.4	3.2	5.1	2.9-8.5	2.8	2.4	1.3-4.1	1.4	2.8	1.1-5.9	1.4	1.8	0.7-4.1
	2011-2015	7.6	7.0	4.9-9.8	4.4	8.6	5.3-13.3	3.2	5.6	3.1-9.4	2.8	2.3	1.2-4.1	1.6	2.6	1.1-5.4	1.2	2.2	0.7-5.0
	2016-2020	10.2	8.2	6.1-11.0	5.2	9.2	6.0-13.8	5.0	7.6	4.8-11.6	3.2	2.6	1.4-4.4	1.4	3.1	1.2-6.5	1.8	2.3	1.0-4.8
Leukemias	1976-1980	12.0	12.3	9.3-15.9	5.6	15.9	10.2-23.3	6.4	11.3	7.6-16.1	6.6	6.9	4.7-9.8	3.8	9.7	5.7-15.4	2.8	5.1	2.7-8.7
	1981-1985	13.6	13.9	10.8-17.7	6.0	15.8	10.5-22.9	7.6	12.8	9.0-17.7	8.0	8.4	6.0-11.5	4.2	11.6	7.0-18.0	3.8	6.4	3.8-10.1
	1986-1990	15.4	15.8	12.4-19.8	7.4	19.8	13.7-27.6	8.0	14.2	10.0-19.5	10.4	10.4	7.7-13.7	4.6	11.8	7.4-17.9	5.8	9.3	6.2-13.6
	1991-1995	15.2	14.3	11.3-18.0	8.6	20.2	14.4-27.4	6.6	10.2	7.0-14.6	9.0	8.5	6.2-11.4	4.2	9.8	6.0-15.1	4.8	7.1	4.5-10.8
	1996-2000	15.2	14.5	11.4-18.2	8.0	18.0	12.8-24.7	7.2	12.5	8.6-17.5	9.4	8.7	6.4-11.7	4.8	11.0	7.0-16.5	4.6	7.3	4.6-11.2
	2001-2005	19.0	17.5	14.1-21.5	7.4	16.7	11.7-23.1	11.6	18.4	13.8-24.2	9.6	8.5	6.2-11.3	3.6	7.9	4.6-12.5	6.0	8.4	5.6-12.3
	2006-2010	17.0	16.0	12.8-20.0	8.2	17.8	12.7-24.3	8.8	14.5	10.4-19.8	9.0	8.0	5.8-10.8	3.6	7.9	4.7-12.6	5.4	8.1	5.2-12.2
	2011-2015	18.8	17.2	13.8-21.2	10.6	21.3	15.9-28.1	8.2	14.2	10.0-19.6	6.2	5.3	3.6-7.7	4.0	8.1	4.9-12.7	2.2	3.1	1.5-6.0
2016-2020	19.6	17.8	14.3-21.9	10.6	20.8	15.4-27.5	9.0	15.4	11.0-21.0	9.2	8.0	5.8-10.9	4.2	8.6	5.3-13.3	5.0	7.9	5.0-12.1	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chenango**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	198.2	430.5	403.7-458.7	94.6	478.1	434.2-525.0	103.6	412.5	377.0-450.5	93.2	200.5	182.4-219.8	43.4	221.5	191.8-254.3	49.8	190.4	167.1-216.2
	1981-1985	210.8	426.4	400.6-453.4	111.2	511.5	468.5-557.4	99.6	369.5	337.0-404.4	105.0	213.0	194.9-232.3	58.2	269.9	238.8-303.8	46.8	166.2	145.2-189.5
	1986-1990	245.0	465.8	439.8-492.9	120.4	518.5	476.9-562.7	124.6	436.0	401.7-472.4	113.4	215.2	197.7-233.9	60.2	268.1	238.0-301.0	53.2	177.2	156.2-200.4
	1991-1995	242.4	440.2	415.6-465.9	125.2	509.6	469.9-551.6	117.2	393.4	361.7-427.3	126.0	226.1	208.7-244.6	63.0	263.4	234.6-294.6	63.0	200.2	178.3-224.1
	1996-2000	269.8	470.1	445.3-496.0	134.6	519.8	480.9-560.9	135.2	444.0	410.7-479.3	113.0	193.2	177.6-209.9	62.0	245.2	218.5-274.4	51.0	156.5	137.6-177.4
	2001-2005	300.2	499.3	474.1-525.5	156.8	566.0	526.7-607.5	143.4	450.9	417.9-486.0	122.6	198.7	183.2-215.3	63.4	236.5	211.0-264.4	59.2	171.6	152.4-192.8
	2006-2010	324.6	504.2	479.4-530.0	172.2	563.2	525.3-603.1	152.4	457.2	424.3-492.2	119.2	183.1	168.5-198.7	65.0	222.1	198.2-248.2	54.2	154.8	136.5-175.1
	2011-2015	353.2	521.1	496.2-547.1	177.8	529.9	494.5-567.3	175.4	520.3	484.7-557.9	120.4	171.5	157.8-186.3	62.4	191.4	170.3-214.6	58.0	158.9	140.5-179.2
	2016-2020	340.0	496.0	471.4-521.7	177.2	511.6	477.0-548.3	162.8	486.5	451.3-524.0	117.2	163.1	149.7-177.4	61.2	181.9	161.4-204.5	56.0	148.2	130.7-167.8
Oral cavity and pharynx	1976-1980	4.6	9.8	6.2-14.8	2.6	10.9	5.8-19.0	2.0	8.1	3.8-15.0	1.8	3.6	1.6-6.8	1.0	4.3	1.4-10.3	0.8	2.8	0.8-7.5
	1981-1985	5.6	11.9	7.8-17.3	3.8	17.7	10.4-27.9	1.8	7.3	3.2-13.9	1.6	3.3	1.4-6.6	1.4	6.5	2.5-13.6	0.2	0.7	0.0-4.0
	1986-1990	3.2	6.0	3.4-9.8	1.8	8.6	3.8-16.2	1.4	4.4	1.8-9.3	0.6	1.2	0.2-3.4	0.0	0.0	0.0-3.2	0.6	1.9	0.4-5.9
	1991-1995	5.2	9.8	6.4-14.4	3.0	12.8	7.2-21.2	2.2	7.6	3.8-13.8	1.6	3.0	1.3-5.9	1.0	3.8	1.2-9.0	0.6	2.2	0.4-6.5
	1996-2000	7.2	12.8	8.9-17.8	5.4	20.4	13.4-29.7	1.8	5.8	2.6-11.3	2.8	4.9	2.7-8.3	2.0	7.7	3.7-14.3	0.8	2.5	0.7-6.8
	2001-2005	9.2	15.3	11.2-20.6	5.8	20.1	13.4-29.1	3.4	10.9	6.3-17.8	1.4	2.4	0.9-5.0	1.0	3.7	1.2-8.8	0.4	1.1	0.1-4.5
	2006-2010	8.0	12.7	9.0-17.5	6.8	22.1	15.2-31.4	1.2	3.6	1.3-8.4	1.6	2.3	1.0-4.9	1.0	3.0	0.9-7.5	0.6	1.7	0.3-5.6
	2011-2015	7.4	9.9	6.9-13.9	6.0	16.3	10.9-23.9	1.4	3.4	1.4-7.8	1.8	2.3	1.1-4.8	1.6	4.2	1.8-8.9	0.2	0.6	0.0-4.0
	2016-2020	10.4	15.5	11.4-20.8	7.0	21.1	14.4-30.2	3.4	10.4	5.7-17.5	1.8	2.5	1.1-5.0	1.2	3.3	1.2-8.0	0.6	1.8	0.3-6.0
Esophagus	1976-1980	1.0	2.1	0.7-4.9	0.6	2.7	0.5-8.2	0.4	1.4	0.2-5.3	1.4	3.1	1.2-6.4	1.2	5.5	1.9-12.2	0.2	0.7	0.0-4.3
	1981-1985	1.8	3.9	1.8-7.4	1.2	7.7	2.8-16.1	0.6	2.1	0.4-6.5	1.6	3.1	1.3-6.2	1.2	6.2	2.2-13.5	0.4	1.3	0.2-5.0
	1986-1990	1.8	3.2	1.5-6.2	1.2	4.7	1.7-10.4	0.6	1.9	0.4-5.9	1.4	2.5	1.0-5.2	0.8	3.6	0.9-9.4	0.6	1.9	0.4-5.9
	1991-1995	2.0	3.5	1.7-6.5	1.6	6.5	2.8-12.9	0.4	1.1	0.1-4.5	2.0	3.6	1.7-6.6	1.4	5.7	2.3-11.7	0.6	1.8	0.4-5.5
	1996-2000	1.8	3.1	1.4-5.9	1.6	6.1	2.6-12.2	0.2	0.7	0.0-3.9	1.8	3.2	1.4-6.1	1.2	4.3	1.6-9.5	0.6	2.0	0.4-6.2
	2001-2005	6.0	9.8	6.6-14.2	4.4	14.9	9.3-22.8	1.6	5.0	2.1-10.2	5.6	8.9	5.9-13.0	3.6	12.7	7.5-20.3	2.0	5.5	2.6-10.6
	2006-2010	4.6	6.8	4.3-10.4	4.0	12.6	7.6-19.9	0.6	1.8	0.4-5.8	5.8	8.8	5.8-12.8	5.4	17.5	11.4-25.8	0.4	1.1	0.1-4.7
	2011-2015	4.4	6.0	3.7-9.4	3.6	10.4	6.1-17.1	0.8	1.9	0.5-5.7	2.6	3.6	1.9-6.5	2.0	5.8	2.7-11.2	0.6	1.7	0.3-5.7
	2016-2020	3.8	5.2	3.1-8.5	3.0	8.3	4.5-14.4	0.8	2.2	0.6-6.5	2.6	3.4	1.8-6.1	1.6	4.2	1.8-9.1	1.0	2.5	0.8-6.7
Stomach	1976-1980	3.0	6.3	3.5-10.5	2.0	10.2	4.7-19.0	1.0	3.7	1.2-8.7	2.8	5.8	3.2-9.8	1.6	8.4	3.4-16.8	1.2	4.4	1.6-9.8
	1981-1985	3.0	5.9	3.3-9.8	2.0	9.5	4.3-17.7	1.0	3.2	1.0-7.9	1.8	3.6	1.6-6.9	1.2	5.9	2.0-13.0	0.6	2.0	0.4-6.2
	1986-1990	3.0	5.8	3.2-9.6	2.0	9.3	4.3-17.2	1.0	3.4	1.1-8.3	2.8	5.4	2.9-9.1	2.0	9.7	4.5-17.8	0.8	2.8	0.7-7.5
	1991-1995	3.6	6.7	4.0-10.7	2.4	11.2	5.7-19.5	1.2	3.9	1.4-8.7	2.2	4.2	2.1-7.6	1.6	7.8	3.3-15.2	0.6	2.0	0.4-6.2
	1996-2000	3.4	5.8	3.4-9.4	2.4	9.3	4.8-16.4	1.0	3.1	1.0-7.7	2.0	3.3	1.6-6.2	1.8	7.1	3.2-13.6	0.2	0.5	0.0-3.5
	2001-2005	4.4	7.4	4.6-11.3	3.2	11.6	6.6-19.0	1.2	3.8	1.4-8.6	1.0	1.8	0.6-4.2	0.8	3.0	0.8-7.8	0.2	0.7	0.0-4.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chenango**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	2.6	4.2	2.2-7.3	2.4	8.5	4.4-15.1	0.2	0.5	0.0-3.7	1.2	1.9	0.7-4.3	1.0	3.7	1.2-8.9	0.2	0.5	0.0-3.7
	2011-2015	3.6	5.3	3.1-8.7	2.4	7.3	3.6-13.3	1.2	3.4	1.2-8.1	1.4	1.9	0.7-4.2	1.0	2.5	0.8-6.7	0.4	1.2	0.1-5.0
	2016-2020	2.8	4.6	2.4-8.1	1.0	3.6	1.0-9.0	1.8	5.6	2.4-11.4	1.6	2.6	1.0-5.5	1.2	3.9	1.3-9.3	0.4	1.3	0.2-5.4
Colorectal	1976-1980	35.8	77.4	66.3-89.7	15.6	80.6	63.0-101.4	20.2	76.9	62.4-93.8	16.4	36.0	28.6-44.8	6.6	35.7	24.1-50.7	9.8	37.6	27.6-49.9
	1981-1985	33.8	67.9	57.9-79.1	16.8	78.2	61.8-97.3	17.0	58.8	46.7-73.0	18.2	36.8	29.6-45.3	9.8	45.5	33.3-60.6	8.4	28.7	20.6-39.1
	1986-1990	34.6	64.6	55.3-75.1	16.2	68.7	54.2-85.7	18.4	59.4	47.7-73.1	14.2	26.6	20.7-33.6	7.8	34.4	24.3-47.3	6.4	20.5	13.9-29.2
	1991-1995	33.2	60.0	51.1-69.9	17.6	73.6	58.8-90.8	15.6	48.7	38.3-61.0	16.6	29.5	23.5-36.6	7.6	32.9	23.1-45.3	9.0	27.0	19.6-36.4
	1996-2000	29.4	50.0	42.2-58.9	14.0	55.6	43.3-70.4	15.4	46.9	36.9-58.9	13.0	21.8	16.8-27.8	6.4	25.7	17.5-36.4	6.6	18.9	12.9-26.9
	2001-2005	32.6	53.3	45.4-62.3	16.6	60.0	47.7-74.6	16.0	46.8	36.9-58.7	13.2	20.8	16.1-26.6	5.8	21.9	14.6-31.6	7.4	20.4	14.3-28.5
	2006-2010	33.2	50.8	43.2-59.3	14.6	48.8	38.1-61.8	18.6	51.9	41.6-64.1	10.8	17.2	12.8-22.6	5.8	21.1	14.0-30.6	5.0	13.3	8.6-20.2
	2011-2015	33.0	49.4	42.0-58.0	17.2	54.0	42.9-67.3	15.8	45.4	35.6-57.3	10.2	14.5	10.7-19.3	3.4	10.4	6.0-17.1	6.8	18.0	12.3-25.7
2016-2020	28.6	41.4	34.6-49.2	15.0	45.2	35.2-57.4	13.6	37.6	28.8-48.5	9.2	13.7	9.8-18.7	4.2	13.9	8.3-22.0	5.0	13.2	8.5-20.2	
Colon excluding rectum	1976-1980	22.8	50.0	41.1-60.2	8.4	45.6	32.3-62.2	14.4	55.3	43.1-69.9	13.0	28.6	22.0-36.6	5.2	28.2	18.0-41.8	7.8	29.9	21.2-41.1
	1981-1985	21.0	41.8	34.1-50.7	9.0	41.7	30.0-56.2	12.0	41.4	31.5-53.7	14.4	29.3	22.9-37.0	7.0	32.8	22.5-46.0	7.4	25.4	17.8-35.3
	1986-1990	24.0	44.9	37.1-53.7	9.0	38.8	28.0-52.3	15.0	48.4	38.0-61.0	12.0	22.5	17.1-29.0	6.6	29.6	20.1-41.8	5.4	17.5	11.4-25.6
	1991-1995	23.0	41.1	33.9-49.5	11.2	47.5	35.7-61.9	11.8	36.5	27.7-47.3	13.0	23.0	17.7-29.4	6.0	25.8	17.2-36.9	7.0	21.6	15.0-30.3
	1996-2000	22.0	37.2	30.6-44.9	9.8	38.9	28.7-51.6	12.2	36.3	27.7-47.0	11.2	18.8	14.2-24.5	5.4	21.5	14.1-31.5	5.8	16.7	11.1-24.3
	2001-2005	22.6	36.6	30.2-44.2	10.6	38.6	28.8-50.7	12.0	34.2	26.0-44.5	10.4	16.4	12.2-21.6	4.4	16.6	10.4-25.3	6.0	16.2	10.9-23.6
	2006-2010	25.0	38.0	31.5-45.5	10.2	34.6	25.6-45.9	14.8	40.9	31.9-51.9	8.8	13.8	9.9-18.7	4.8	17.3	11.0-26.0	4.0	10.3	6.2-16.4
	2011-2015	24.8	37.3	30.8-44.8	12.2	38.1	28.8-49.7	12.6	36.5	27.7-47.4	9.2	13.1	9.5-17.7	3.2	9.8	5.5-16.4	6.0	16.0	10.7-23.4
2016-2020	21.4	30.1	24.5-36.8	10.8	31.9	23.7-42.3	10.6	28.4	21.1-37.8	8.0	12.0	8.4-16.7	3.4	11.4	6.3-19.1	4.6	12.1	7.6-18.8	
Rectum & rectosigmoid	1976-1980	13.0	27.4	21.0-35.0	7.2	35.0	24.1-48.9	5.8	21.7	14.4-31.3	3.4	7.4	4.3-12.0	1.4	7.5	2.9-15.6	2.0	7.6	3.6-14.3
	1981-1985	12.8	26.1	20.0-33.5	7.8	36.5	25.6-50.3	5.0	17.3	11.1-25.8	3.8	7.5	4.5-11.8	2.8	12.7	6.8-21.6	1.0	3.3	1.1-8.1
	1986-1990	10.6	19.7	14.7-25.9	7.2	29.8	20.7-41.5	3.4	11.0	6.3-17.8	2.2	4.1	2.0-7.3	1.2	4.9	1.8-10.7	1.0	3.0	1.0-7.4
	1991-1995	10.2	18.8	14.0-24.8	6.4	26.1	17.8-36.9	3.8	12.2	7.3-19.3	3.6	6.5	3.8-10.3	1.6	7.1	3.0-13.9	2.0	5.4	2.6-10.3
	1996-2000	7.4	12.8	9.0-17.7	4.2	16.7	10.3-25.6	3.2	10.6	6.0-17.4	1.8	3.0	1.4-5.8	1.0	4.2	1.3-9.8	0.8	2.2	0.6-6.2
	2001-2005	10.0	16.6	12.3-22.1	6.0	21.4	14.4-30.8	4.0	12.6	7.6-19.8	2.8	4.5	2.4-7.6	1.4	5.2	2.1-11.0	1.4	4.2	1.7-8.9
	2006-2010	8.2	12.8	9.1-17.5	4.4	14.2	8.8-21.9	3.8	10.9	6.5-17.6	2.0	3.4	1.6-6.4	1.0	3.8	1.2-9.0	1.0	3.1	1.0-7.6
	2011-2015	8.2	12.2	8.7-16.7	5.0	15.9	10.2-23.8	3.2	8.9	5.0-15.0	1.0	1.4	0.4-3.5	0.2	0.6	0.0-4.0	0.8	2.0	0.5-6.0
2016-2020	7.2	11.3	7.7-16.0	4.2	13.3	8.1-21.0	3.0	9.2	4.9-16.0	1.2	1.7	0.6-4.1	0.8	2.5	0.7-7.1	0.4	1.2	0.1-5.1	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chenango**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	0.8	1.8	0.5-4.7	0.4	2.0	0.2-7.3	0.4	1.5	0.2-5.6	0.6	1.4	0.3-3.9	0.2	0.8	0.0-4.9	0.4	1.5	0.2-5.6
	1981-1985	1.2	2.5	0.9-5.4	0.4	1.6	0.2-6.0	0.8	2.9	0.8-7.5	0.6	1.3	0.3-3.8	0.2	1.1	0.0-5.5	0.4	1.3	0.2-5.1
	1986-1990	2.4	4.2	2.2-7.5	2.0	8.0	3.8-14.9	0.4	1.2	0.1-4.7	1.8	3.5	1.6-6.6	1.0	4.1	1.3-9.7	0.8	2.9	0.7-7.6
	1991-1995	2.2	3.9	2.0-7.1	1.2	4.7	1.7-10.4	1.0	3.3	1.0-7.9	2.0	3.7	1.7-6.8	1.0	4.1	1.3-9.7	1.0	3.2	1.0-7.7
	1996-2000	2.0	3.4	1.6-6.4	1.6	6.2	2.6-12.2	0.4	1.4	0.2-5.3	1.8	3.0	1.4-5.8	1.4	5.4	2.2-11.3	0.4	1.2	0.1-4.7
	2001-2005	2.8	4.9	2.6-8.3	2.0	7.3	3.5-13.6	0.8	2.1	0.6-6.1	3.0	5.1	2.8-8.5	1.4	5.1	2.0-10.6	1.6	4.9	2.0-10.1
	2006-2010	2.2	3.7	1.8-6.9	1.2	3.9	1.4-8.7	1.0	3.8	1.2-9.2	2.2	3.5	1.7-6.4	1.2	3.7	1.4-8.6	1.0	3.3	1.0-8.1
	2011-2015	2.8	3.6	2.0-6.3	1.6	4.2	1.8-8.9	1.2	3.2	1.2-7.6	2.4	3.2	1.6-5.9	1.2	3.2	1.2-7.8	1.2	3.2	1.2-7.6
2016-2020	4.8	6.1	3.9-9.4	2.8	7.3	4.0-13.0	2.0	4.9	2.3-9.9	4.0	5.4	3.2-8.8	2.2	6.0	2.9-11.4	1.8	4.9	2.0-10.4	
Pancreas	1976-1980	4.2	8.7	5.4-13.4	2.0	9.1	4.2-17.2	2.2	7.7	3.8-14.0	4.8	9.6	6.1-14.4	2.2	9.5	4.5-17.5	2.6	9.3	4.9-16.1
	1981-1985	5.4	11.5	7.5-16.7	3.2	15.7	8.8-25.7	2.2	7.7	3.8-14.1	4.6	9.5	6.0-14.4	2.6	11.7	6.1-20.3	2.0	7.1	3.3-13.4
	1986-1990	4.8	9.4	6.0-14.0	2.4	11.6	5.9-20.2	2.4	8.3	4.2-14.6	5.2	10.4	6.7-15.2	3.6	17.7	10.4-28.0	1.6	5.8	2.5-11.6
	1991-1995	6.8	12.1	8.4-17.0	3.6	14.7	8.7-23.4	3.2	9.7	5.5-16.0	6.0	10.7	7.2-15.4	2.8	11.6	6.3-19.5	3.2	10.1	5.7-16.7
	1996-2000	5.4	9.1	6.0-13.3	1.8	7.0	3.2-13.3	3.6	10.7	6.3-17.2	5.8	9.7	6.5-14.1	2.0	8.1	3.9-15.0	3.8	10.8	6.4-17.2
	2001-2005	7.2	11.5	8.0-16.0	3.2	11.0	6.3-18.1	4.0	11.8	7.2-18.6	8.0	12.9	9.2-17.7	3.4	12.0	6.9-19.4	4.6	13.3	8.4-20.3
	2006-2010	8.4	12.5	9.0-17.1	3.2	10.7	6.1-17.7	5.2	14.0	9.1-20.9	7.8	11.9	8.4-16.4	3.4	11.4	6.6-18.6	4.4	12.0	7.5-18.6
	2011-2015	10.2	14.3	10.6-19.0	4.6	13.2	8.3-20.4	5.6	15.3	10.1-22.6	6.8	9.4	6.5-13.4	3.0	8.8	4.9-15.1	3.8	10.0	5.9-16.1
2016-2020	11.0	15.2	11.4-20.2	6.2	16.6	11.1-24.3	4.8	13.7	8.7-21.1	11.0	15.0	11.2-19.8	6.4	18.4	12.4-26.6	4.6	11.9	7.5-18.6	
Larynx	1976-1980	2.2	4.4	2.1-7.9	2.0	8.4	4.0-15.9	0.2	0.6	0.0-4.1	0.6	1.2	0.2-3.6	0.4	2.1	0.2-7.5	0.2	0.6	0.0-4.1
	1981-1985	1.6	3.0	1.3-5.9	1.4	5.7	2.3-12.0	0.2	0.7	0.0-4.1	0.8	1.6	0.4-4.1	0.8	3.3	0.9-8.8	0.0	0.0	0.0-2.9
	1986-1990	2.0	3.9	1.8-7.2	1.0	4.1	1.3-9.8	1.0	4.0	1.2-9.3	0.0	0.0	0.0-1.5	0.0	0.0	0.0-3.2	0.0	0.0	0.0-2.8
	1991-1995	2.2	3.9	1.9-7.0	0.6	2.1	0.4-6.4	1.6	5.5	2.3-10.9	0.8	1.3	0.4-3.4	0.6	2.1	0.4-6.3	0.2	0.6	0.0-3.7
	1996-2000	1.4	2.4	1.0-5.0	0.8	3.0	0.8-7.8	0.6	2.0	0.4-6.1	0.4	0.7	0.1-2.5	0.2	0.9	0.0-4.7	0.2	0.7	0.0-3.9
	2001-2005	2.6	4.3	2.3-7.5	2.0	7.2	3.5-13.5	0.6	1.9	0.4-6.0	1.2	2.0	0.7-4.5	1.0	3.8	1.2-8.9	0.2	0.6	0.0-4.0
	2006-2010	3.0	4.1	2.3-7.0	2.4	6.5	3.4-11.9	0.6	1.7	0.3-5.5	0.0	0.0	0.0-1.5	0.0	0.0	0.0-3.0	0.0	0.0	0.0-3.0
	2011-2015	2.6	3.2	1.7-5.8	2.2	5.4	2.7-10.3	0.4	1.0	0.1-4.6	0.6	0.8	0.2-2.7	0.4	1.0	0.1-4.6	0.2	0.6	0.0-4.0
2016-2020	2.8	3.9	2.1-7.0	1.8	5.4	2.4-10.9	1.0	2.7	0.8-7.3	0.6	0.7	0.1-2.6	0.6	1.5	0.3-5.5	0.0	0.0	0.0-3.4	
Lung and bronchus	1976-1980	24.2	51.8	42.8-62.1	19.0	91.9	73.8-112.9	5.2	20.0	12.9-29.5	18.4	39.2	31.4-48.2	14.0	69.5	53.6-88.3	4.4	16.9	10.4-25.8
	1981-1985	36.4	72.4	62.1-84.0	24.4	106.1	87.6-127.3	12.0	43.8	33.2-56.7	25.0	49.9	41.4-59.7	18.0	81.3	64.9-100.5	7.0	24.0	16.6-33.7
	1986-1990	42.4	79.0	68.6-90.6	28.2	117.6	98.6-139.3	14.2	48.7	37.8-61.8	30.4	56.6	47.8-66.5	20.2	85.2	69.0-104.0	10.2	33.9	25.1-44.9
	1991-1995	42.0	75.4	65.5-86.4	24.8	100.5	83.4-120.0	17.2	56.9	45.4-70.6	35.4	63.0	54.0-73.1	20.4	82.6	67.1-100.5	15.0	48.4	37.9-61.0
	1996-2000	38.6	66.3	57.2-76.4	23.2	89.1	73.5-107.1	15.4	49.5	39.0-62.1	29.8	50.9	43.0-59.8	19.4	75.1	60.8-91.8	10.4	32.0	23.8-42.2
	2001-2005	40.2	65.8	57.0-75.7	22.0	78.9	64.7-95.3	18.2	56.2	45.1-69.3	29.2	48.1	40.6-56.7	17.4	62.8	50.2-77.7	11.8	36.1	27.4-46.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chenango**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	52.0	78.5	69.1-88.9	30.8	100.8	85.2-118.5	21.2	61.4	50.1-74.7	36.4	55.2	47.4-64.1	20.8	68.2	55.5-83.0	15.6	45.5	35.8-57.3
	2011-2015	51.0	71.8	63.1-81.5	27.0	79.7	66.5-95.1	24.0	65.2	53.8-78.6	33.8	48.1	41.0-56.3	18.0	54.5	43.7-67.6	15.8	43.0	33.9-54.2
	2016-2020	50.8	68.7	60.2-78.2	25.6	69.4	57.7-83.2	25.2	69.1	56.9-83.4	31.8	43.2	36.5-50.8	16.6	45.3	35.9-56.8	15.2	41.3	32.1-52.6
Melanoma of the skin	1976-1980	3.2	7.1	4.0-11.7	1.8	7.9	3.5-15.3	1.4	6.3	2.4-13.0	1.4	3.3	1.3-6.7	0.6	2.9	0.6-8.6	0.8	3.6	0.9-9.0
	1981-1985	3.6	8.3	4.8-13.1	1.2	5.5	1.9-12.2	2.4	11.5	5.8-20.0	0.4	0.8	0.1-3.1	0.4	1.7	0.2-6.5	0.0	0.0	0.0-2.9
	1986-1990	3.8	7.7	4.6-12.0	2.0	8.9	4.2-16.5	1.8	6.9	3.1-13.1	1.2	2.8	1.0-5.9	0.4	2.0	0.2-6.9	0.8	3.5	1.0-8.8
	1991-1995	5.0	8.9	5.8-13.2	3.4	12.8	7.4-20.6	1.6	6.2	2.6-12.1	0.8	1.3	0.4-3.5	0.2	0.9	0.0-4.7	0.6	1.8	0.4-5.5
	1996-2000	7.6	13.7	9.7-18.9	4.4	16.6	10.4-25.2	3.2	11.6	6.6-19.0	1.2	2.1	0.8-4.6	1.0	3.6	1.2-8.6	0.2	0.8	0.0-4.3
	2001-2005	10.0	16.9	12.5-22.5	5.4	19.6	12.9-28.8	4.6	15.5	9.7-23.6	2.6	4.3	2.3-7.4	1.6	6.0	2.6-12.0	1.0	3.4	1.1-8.1
	2006-2010	13.2	21.0	16.1-27.0	8.6	28.5	20.4-38.9	4.6	14.9	9.2-22.8	1.2	2.1	0.7-4.8	0.8	2.8	0.7-7.4	0.4	1.5	0.1-5.9
	2011-2015	17.8	27.3	21.7-34.0	9.8	29.1	21.3-39.1	8.0	26.6	18.6-37.0	1.6	2.4	1.0-5.0	0.8	2.9	0.8-7.7	0.8	2.2	0.6-6.4
	2016-2020	17.0	26.1	20.6-32.9	10.2	33.9	24.8-45.5	6.8	19.0	12.8-27.4	1.2	1.4	0.5-3.6	0.6	1.6	0.3-5.5	0.6	1.3	0.3-5.1
Female breast	1976-1980							29.2	122.6	103.0-144.7							11.4	46.3	34.8-60.4
	1981-1985							19.8	73.5	59.4-89.9							9.4	34.3	25.0-46.0
	1986-1990							38.0	141.6	121.7-163.7							9.6	32.0	23.4-42.7
	1991-1995							34.4	120.6	103.0-140.4							12.0	39.8	30.2-51.5
	1996-2000							42.2	141.9	123.2-162.8							9.0	29.0	21.1-39.1
	2001-2005							33.6	106.3	90.6-124.1							7.6	21.5	15.1-30.0
	2006-2010							37.6	112.8	96.8-130.9							7.8	21.2	15.0-29.6
	2011-2015							45.6	134.5	116.8-154.4							8.2	24.6	17.3-34.3
	2016-2020							40.8	123.2	105.7-142.9							6.0	14.5	9.6-21.4
Cervix uteri	1976-1980							5.2	22.1	14.3-32.5							1.4	5.0	2.0-10.5
	1981-1985							4.6	20.4	12.8-30.8							1.6	6.5	2.8-12.9
	1986-1990							5.0	19.4	12.4-28.9							1.8	6.6	2.9-12.8
	1991-1995							2.8	9.8	5.3-16.6							1.0	3.3	1.0-7.9
	1996-2000							3.0	11.5	6.4-19.1							0.8	2.8	0.8-7.3
	2001-2005							3.0	10.4	5.8-17.4							1.2	3.7	1.4-8.5
	2006-2010							2.8	11.0	5.8-18.7							0.4	1.1	0.1-4.6
	2011-2015							2.4	10.7	5.4-18.8							0.6	2.6	0.5-7.7
	2016-2020							3.0	10.8	5.7-18.7							1.2	3.5	1.2-8.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chenango**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							10.2	41.0	30.3-54.3							1.6	5.5	2.4-11.0
	1981-1985							8.6	32.3	23.1-43.9							1.4	5.5	2.1-11.5
	1986-1990							7.6	27.4	19.3-37.9							1.4	4.1	1.7-8.8
	1991-1995							7.8	28.1	19.8-38.6							2.0	6.5	3.1-12.2
	1996-2000							8.0	27.4	19.5-37.5							1.4	4.6	1.8-9.8
	2001-2005							11.6	36.2	27.4-47.1							2.0	5.5	2.6-10.6
	2006-2010							10.8	31.0	23.1-40.8							1.0	2.9	0.9-7.3
	2011-2015							15.2	44.9	34.9-57.1							2.0	4.7	2.3-9.4
	2016-2020							14.4	45.1	34.5-58.2							1.8	4.6	2.1-9.6
Ovary	1976-1980							5.2	22.1	14.3-32.6							3.2	12.6	7.1-20.8
	1981-1985							5.0	19.5	12.5-29.1							4.0	13.8	8.3-21.5
	1986-1990							5.4	18.8	12.3-27.6							3.2	10.4	5.9-17.2
	1991-1995							3.6	11.7	6.9-18.8							3.2	10.1	5.7-16.6
	1996-2000							4.4	15.6	9.7-23.8							2.6	8.2	4.3-14.3
	2001-2005							4.6	14.1	8.8-21.7							2.6	7.1	3.8-12.7
	2006-2010							5.8	17.0	11.2-24.9							3.4	8.8	5.1-14.6
	2011-2015							4.4	12.9	7.9-20.2							2.2	5.8	2.8-11.0
	2016-2020							4.4	14.6	8.6-23.1							2.4	5.8	3.0-10.9
Prostate	1976-1980				21.8	125.3	102.1-151.8							4.8	28.3	17.8-42.3			
	1981-1985				21.0	101.7	82.5-123.8							7.2	34.3	23.6-47.9			
	1986-1990				23.6	102.6	84.5-123.3							5.6	28.5	18.7-41.2			
	1991-1995				32.6	132.1	112.3-154.3							8.0	36.1	25.6-49.2			
	1996-2000				35.8	138.1	118.4-160.1							7.8	32.9	23.3-45.1			
	2001-2005				46.2	164.6	143.9-187.6							9.6	38.5	28.3-51.2			
	2006-2010				41.4	127.9	110.8-147.2							4.8	17.5	11.1-26.2			
	2011-2015				37.8	103.3	88.9-119.8							7.8	25.6	18.1-35.3			
	2016-2020				43.0	111.2	96.5-127.9							6.8	21.2	14.6-30.1			
Testis	1976-1980				1.0	4.0	1.2-9.8							0.4	2.0	0.2-7.3			
	1981-1985				1.2	5.0	1.8-11.2							0.0	0.0	0.0-3.4			
	1986-1990				0.4	1.3	0.2-5.3							0.2	0.8	0.0-4.6			
	1991-1995				1.4	5.4	2.2-11.3							0.0	0.0	0.0-3.0			
	1996-2000				1.4	5.6	2.2-11.5							0.0	0.0	0.0-3.0			
	2001-2005				1.4	6.4	2.5-13.0							0.0	0.0	0.0-3.0			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chenango**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				2.0	9.0	4.2-16.6							0.0	0.0	0.0-3.0			
	2011-2015				1.8	8.2	3.7-15.6							0.0	0.0	0.0-3.1			
	2016-2020				1.4	8.1	3.2-16.2							0.2	0.5	0.0-4.0			
Urinary bladder (incl. in situ)	1976-1980	10.8	22.4	16.7-29.3	8.2	40.0	28.2-54.8	2.6	9.4	5.0-16.3	1.8	3.8	1.7-7.2	1.4	7.9	3.0-16.3	0.4	1.4	0.2-5.5
	1981-1985	12.2	24.5	18.7-31.6	9.8	47.7	34.9-63.5	2.4	8.2	4.2-14.6	3.0	6.0	3.4-10.0	2.4	12.1	6.1-21.2	0.6	1.9	0.4-6.0
	1986-1990	15.2	28.0	22.0-35.1	10.6	46.4	34.4-61.0	4.6	14.6	9.2-22.2	3.2	5.9	3.3-9.6	2.4	10.0	5.1-17.5	0.8	2.5	0.7-6.7
	1991-1995	9.6	17.5	12.9-23.2	6.8	28.2	19.4-39.4	2.8	8.6	4.6-14.7	3.6	6.4	3.8-10.2	2.2	9.8	4.8-17.5	1.4	4.1	1.6-8.7
	1996-2000	14.6	24.8	19.4-31.3	10.8	41.3	31.0-54.1	3.8	11.1	6.6-17.7	2.8	4.7	2.6-8.0	2.4	10.0	5.1-17.5	0.4	1.1	0.1-4.5
	2001-2005	17.2	27.8	22.2-34.5	13.2	47.2	36.4-60.3	4.0	11.5	7.0-18.2	4.2	6.8	4.2-10.6	3.0	11.5	6.4-19.1	1.2	3.2	1.2-7.6
	2006-2010	16.6	25.4	20.2-31.6	12.2	40.5	30.8-52.5	4.4	12.0	7.5-18.7	5.2	7.8	5.1-11.7	3.8	13.7	8.2-21.6	1.4	3.9	1.5-8.6
	2011-2015	19.4	27.1	21.9-33.4	16.6	49.9	39.6-62.4	2.8	7.5	4.1-13.1	4.0	5.6	3.4-8.9	3.4	10.5	6.1-17.3	0.6	1.6	0.3-5.5
	2016-2020	18.6	25.1	20.2-31.1	16.2	46.2	36.5-58.1	2.4	6.7	3.4-12.5	4.0	5.5	3.3-8.8	3.0	8.9	5.0-15.3	1.0	2.7	0.8-7.1
Kidney and renal pelvis	1976-1980	3.4	7.1	4.1-11.5	1.8	9.2	4.1-17.5	1.6	6.1	2.6-12.1	2.2	4.7	2.3-8.4	1.2	5.5	2.0-12.2	1.0	3.8	1.2-9.0
	1981-1985	4.8	9.2	5.9-13.7	3.6	16.3	9.5-26.1	1.2	4.1	1.5-9.1	2.2	4.3	2.1-7.8	1.8	8.8	3.9-16.8	0.4	1.4	0.2-5.3
	1986-1990	4.8	9.4	6.0-14.1	3.6	16.7	9.8-26.5	1.2	3.9	1.4-8.8	2.4	4.4	2.3-7.7	1.6	7.6	3.2-15.1	0.8	2.4	0.6-6.5
	1991-1995	6.2	11.6	7.9-16.5	3.2	13.2	7.5-21.6	3.0	10.0	5.5-16.7	3.8	7.1	4.3-11.1	2.4	9.9	5.0-17.3	1.4	4.7	1.8-9.9
	1996-2000	7.0	12.5	8.7-17.4	4.2	15.8	9.8-24.2	2.8	9.6	5.2-16.3	3.4	5.9	3.5-9.6	2.6	9.8	5.2-16.8	0.8	2.4	0.6-6.5
	2001-2005	8.4	14.2	10.2-19.4	4.6	16.4	10.3-24.8	3.8	12.6	7.5-20.0	3.0	4.9	2.7-8.2	1.2	4.2	1.5-9.5	1.8	5.3	2.4-10.4
	2006-2010	12.6	19.8	15.1-25.6	8.0	26.1	18.5-36.1	4.6	13.7	8.5-21.2	2.0	3.0	1.4-5.7	1.2	3.9	1.4-8.9	0.8	2.1	0.6-6.0
	2011-2015	13.2	18.7	14.4-24.1	8.0	22.3	15.8-30.9	5.2	15.4	9.9-23.1	2.6	3.5	1.8-6.3	1.8	5.5	2.5-10.9	0.8	2.2	0.6-6.3
	2016-2020	14.0	20.9	16.1-26.9	8.6	24.7	17.7-34.0	5.4	17.4	11.0-26.3	2.6	3.5	1.9-6.3	1.8	5.2	2.4-10.5	0.8	1.9	0.5-5.9
Brain and other central nervous system	1976-1980	3.6	7.8	4.5-12.4	1.6	6.7	2.8-13.7	2.0	9.3	4.3-17.0	2.6	5.3	2.8-9.2	1.2	5.6	2.0-12.5	1.4	5.7	2.2-11.9
	1981-1985	5.2	10.4	6.8-15.4	1.8	8.2	3.7-15.8	3.4	13.0	7.5-21.1	3.0	6.4	3.5-10.6	0.8	3.4	0.9-8.9	2.2	9.1	4.4-16.5
	1986-1990	4.2	8.1	5.0-12.4	2.6	10.5	5.5-18.2	1.6	5.6	2.4-11.2	3.2	6.7	3.8-10.9	1.8	8.0	3.6-15.2	1.4	4.8	1.9-10.2
	1991-1995	4.2	8.0	4.9-12.3	2.6	10.1	5.4-17.4	1.6	5.8	2.5-11.5	3.0	5.5	3.1-9.1	1.8	6.9	3.1-13.2	1.2	4.0	1.5-9.0
	1996-2000	4.2	7.6	4.7-11.6	2.0	7.6	3.6-14.1	2.2	7.5	3.7-13.6	2.6	4.8	2.5-8.2	1.6	6.0	2.6-11.9	1.0	3.5	1.1-8.2
	2001-2005	3.8	7.0	4.2-11.1	1.2	4.7	1.7-10.3	2.6	9.2	4.8-15.9	1.0	1.9	0.6-4.5	0.4	1.7	0.2-6.1	0.6	2.3	0.5-6.7
	2006-2010	5.0	8.8	5.6-13.2	3.4	12.2	7.0-19.9	1.6	5.7	2.4-11.6	3.6	5.8	3.4-9.4	2.2	7.6	3.7-14.0	1.4	4.8	1.9-10.1
	2011-2015	5.0	9.0	5.7-13.6	2.4	9.4	4.6-16.9	2.6	8.3	4.2-14.9	2.4	3.9	1.9-7.1	1.6	5.2	2.2-10.8	0.8	2.6	0.6-7.4
	2016-2020	5.8	10.3	6.7-15.2	3.2	11.5	6.3-19.2	2.6	9.4	4.7-16.8	3.8	6.8	4.0-11.0	2.0	7.5	3.5-14.2	1.8	6.3	2.6-12.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chenango**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	1.0	2.0	0.6-4.7	0.2	0.7	0.0-4.7	0.8	2.7	0.7-7.3	0.2	0.4	0.0-2.3	0.0	0.0	0.0-3.6	0.2	0.7	0.0-4.3
	1981-1985	1.0	2.0	0.6-4.7	0.2	1.1	0.0-5.5	0.8	2.9	0.8-7.7	0.0	0.0	0.0-1.5	0.0	0.0	0.0-3.4	0.0	0.0	0.0-2.9
	1986-1990	2.2	4.1	2.1-7.4	0.0	0.0	0.0-3.2	2.2	7.4	3.7-13.5	0.6	1.2	0.2-3.5	0.0	0.0	0.0-3.2	0.6	1.7	0.4-5.5
	1991-1995	1.6	2.8	1.2-5.6	0.6	2.2	0.4-6.5	1.0	3.5	1.1-8.2	0.4	0.7	0.1-2.5	0.0	0.0	0.0-3.0	0.4	1.3	0.2-4.9
	1996-2000	4.2	8.1	5.0-12.4	0.6	2.2	0.4-6.6	3.6	13.8	8.1-21.8	0.0	0.0	0.0-1.4	0.0	0.0	0.0-3.0	0.0	0.0	0.0-2.8
	2001-2005	7.2	13.7	9.5-19.1	1.2	4.6	1.6-10.1	6.0	22.6	15.1-32.6	0.2	0.3	0.0-1.9	0.0	0.0	0.0-3.0	0.2	0.4	0.0-3.5
	2006-2010	8.6	17.0	12.2-23.1	1.8	7.0	3.2-13.4	6.8	27.2	18.5-38.4	0.6	0.8	0.2-2.8	0.4	1.1	0.1-4.8	0.2	0.5	0.0-3.8
	2011-2015	11.6	22.2	16.5-29.2	2.2	7.6	3.6-14.3	9.4	36.7	26.3-49.6	0.2	0.2	0.0-1.9	0.0	0.0	0.0-3.1	0.2	0.5	0.0-3.8
	2016-2020	11.2	21.0	15.5-27.9	3.2	10.8	5.9-18.3	8.0	31.9	22.2-44.5	0.4	0.6	0.1-2.5	0.4	1.4	0.2-5.5	0.0	0.0	0.0-3.4
Hodgkin lymphoma	1976-1980	1.6	3.0	1.3-6.0	0.2	1.0	0.0-5.4	1.4	4.8	1.9-10.1	0.6	1.4	0.3-4.1	0.2	1.0	0.0-5.4	0.4	1.5	0.2-5.6
	1981-1985	0.4	0.7	0.1-2.7	0.2	0.7	0.0-4.5	0.2	0.7	0.0-4.2	0.2	0.3	0.0-2.1	0.0	0.0	0.0-3.4	0.2	0.7	0.0-4.1
	1986-1990	2.4	4.6	2.4-8.1	1.8	7.6	3.4-14.6	0.6	2.1	0.4-6.2	0.4	0.7	0.1-2.7	0.0	0.0	0.0-3.2	0.4	1.3	0.2-5.1
	1991-1995	1.6	3.3	1.4-6.5	0.4	1.7	0.2-6.0	1.2	4.9	1.8-10.5	0.2	0.4	0.0-2.1	0.2	0.7	0.0-4.4	0.0	0.0	0.0-2.7
	1996-2000	1.6	3.1	1.3-6.1	1.2	4.8	1.7-10.4	0.4	1.7	0.2-5.8	0.2	0.4	0.0-2.2	0.2	0.8	0.0-4.4	0.0	0.0	0.0-2.8
	2001-2005	1.0	1.7	0.5-4.0	0.6	2.0	0.4-6.2	0.4	1.2	0.1-4.8	0.8	1.4	0.4-3.8	0.4	1.4	0.2-5.4	0.4	1.5	0.2-5.4
	2006-2010	2.6	4.3	2.2-7.6	2.0	6.6	3.0-12.6	0.6	2.2	0.4-6.5	0.4	0.7	0.1-2.7	0.2	0.7	0.0-4.3	0.2	0.7	0.0-4.1
	2011-2015	2.0	4.3	2.0-7.9	1.4	6.3	2.5-13.1	0.6	2.3	0.4-6.9	0.0	0.0	0.0-1.6	0.0	0.0	0.0-3.1	0.0	0.0	0.0-3.1
	2016-2020	0.6	0.7	0.1-2.6	0.4	1.0	0.1-4.7	0.2	0.4	0.0-3.9	0.0	0.0	0.0-1.7	0.0	0.0	0.0-3.3	0.0	0.0	0.0-3.4
Non-Hodgkin lymphomas	1976-1980	6.8	14.5	10.0-20.4	3.2	15.1	8.4-24.9	3.6	13.0	7.7-20.7	2.8	5.7	3.1-9.7	0.6	2.3	0.4-7.2	2.2	8.3	4.1-15.0
	1981-1985	8.0	16.2	11.5-22.2	3.2	14.3	8.0-23.5	4.8	18.0	11.4-27.0	3.2	6.5	3.7-10.6	1.4	6.4	2.5-13.5	1.8	6.2	2.8-12.0
	1986-1990	9.8	18.8	13.9-24.9	5.0	21.3	13.7-31.5	4.8	15.5	9.9-23.3	5.4	10.3	6.8-15.0	2.2	9.8	4.8-17.6	3.2	10.3	5.8-17.0
	1991-1995	8.4	15.2	10.9-20.6	5.2	21.0	13.6-30.8	3.2	10.0	5.7-16.5	4.6	8.1	5.1-12.2	1.6	6.5	2.8-12.9	3.0	9.1	5.1-15.2
	1996-2000	11.4	19.8	15.0-25.7	5.8	22.6	15.1-32.6	5.6	17.6	11.6-25.7	5.4	9.2	6.1-13.5	2.6	10.3	5.4-17.7	2.8	8.3	4.5-14.2
	2001-2005	10.8	18.2	13.6-23.8	5.8	22.0	14.7-31.7	5.0	15.6	10.0-23.3	5.0	8.1	5.2-12.0	2.6	9.7	5.1-16.7	2.4	6.6	3.4-12.0
	2006-2010	11.6	18.8	14.2-24.6	7.0	24.1	16.6-33.9	4.6	14.1	8.8-21.7	3.2	5.0	2.8-8.2	2.2	7.7	3.8-14.0	1.0	2.9	0.9-7.3
	2011-2015	15.6	23.5	18.4-29.7	9.6	30.8	22.4-41.5	6.0	17.1	11.4-25.1	4.2	5.6	3.4-8.8	2.4	6.9	3.5-12.7	1.8	4.6	2.1-9.4
	2016-2020	11.2	16.2	12.0-21.4	6.6	20.1	13.5-28.9	4.6	12.6	7.8-19.7	3.4	5.1	2.9-8.5	2.0	6.6	3.1-12.6	1.4	4.1	1.6-9.2
Myeloma	1976-1980	3.6	7.6	4.4-12.1	2.4	10.7	5.4-18.9	1.2	4.9	1.7-10.9	2.4	5.4	2.7-9.5	1.8	9.9	4.4-18.7	0.6	2.0	0.4-6.2
	1981-1985	4.6	8.7	5.5-13.2	2.6	11.6	6.1-20.1	2.0	6.4	3.1-12.1	2.6	5.5	2.9-9.4	1.6	7.8	3.4-15.4	1.0	3.3	1.1-7.9
	1986-1990	5.6	10.7	7.1-15.5	2.8	13.0	6.9-22.0	2.8	9.5	5.2-16.2	3.0	5.8	3.2-9.6	2.0	9.4	4.4-17.4	1.0	3.5	1.1-8.4
	1991-1995	2.6	4.6	2.4-7.9	1.2	4.7	1.7-10.3	1.4	4.4	1.7-9.2	3.0	5.4	3.0-9.0	1.2	4.7	1.7-10.3	1.8	5.6	2.5-10.8
	1996-2000	4.4	7.4	4.6-11.3	2.8	10.5	5.7-17.7	1.6	4.5	1.9-9.2	2.8	4.7	2.6-8.0	1.6	6.2	2.7-12.2	1.2	3.6	1.3-8.2
	2001-2005	3.8	6.2	3.7-9.8	2.0	7.1	3.4-13.3	1.8	5.0	2.3-9.9	2.2	3.4	1.7-6.3	1.0	3.6	1.2-8.6	1.2	2.8	1.0-6.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Chenango**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	5.0	7.8	5.0-11.7	3.0	10.1	5.6-17.1	2.0	5.8	2.7-11.1	2.0	3.6	1.7-6.7	1.0	3.9	1.2-9.4	1.0	3.1	1.0-7.6
	2011-2015	4.2	6.1	3.7-9.5	2.6	8.4	4.4-14.8	1.6	4.3	1.9-9.1	1.6	2.4	1.0-5.0	0.6	1.8	0.4-5.9	1.0	2.9	0.9-7.3
	2016-2020	3.0	4.3	2.3-7.5	2.0	5.5	2.4-11.1	1.0	3.0	0.9-7.9	1.0	1.3	0.4-3.5	1.0	2.9	0.9-7.5	0.0	0.0	0.0-3.4
Leukemias	1976-1980	5.4	11.3	7.4-16.6	3.0	14.9	8.2-24.8	2.4	8.6	4.4-15.3	3.4	7.2	4.1-11.5	2.0	9.8	4.5-18.3	1.4	4.8	1.9-10.1
	1981-1985	7.2	15.0	10.4-20.9	5.2	24.1	15.5-35.6	2.0	7.4	3.4-13.8	4.8	9.9	6.3-14.8	2.8	13.3	7.1-22.4	2.0	7.1	3.3-13.3
	1986-1990	6.2	11.8	8.0-16.8	4.2	19.0	11.6-29.2	2.0	6.3	3.0-11.8	4.8	9.5	6.0-14.2	3.4	15.3	8.8-24.5	1.4	4.5	1.8-9.6
	1991-1995	8.0	14.6	10.4-19.9	5.0	20.4	13.1-30.3	3.0	9.9	5.5-16.5	3.8	7.0	4.2-11.0	3.2	14.0	7.9-22.8	0.6	2.0	0.4-6.0
	1996-2000	9.2	16.1	11.8-21.6	5.2	20.2	13.2-29.7	4.0	13.1	7.9-20.5	4.2	7.2	4.5-11.2	2.4	9.7	4.9-17.0	1.8	5.9	2.7-11.5
	2001-2005	13.0	22.5	17.3-28.9	7.6	29.1	20.5-40.1	5.4	17.9	11.7-26.5	4.8	7.8	5.0-11.8	2.8	10.5	5.7-17.8	2.0	5.7	2.7-10.9
	2006-2010	11.0	17.1	12.8-22.5	6.6	21.7	14.8-30.9	4.4	13.3	8.2-20.6	3.6	6.0	3.5-9.7	2.0	7.6	3.6-14.1	1.6	5.5	2.3-11.3
	2011-2015	13.8	20.1	15.5-25.8	8.2	24.0	17.1-33.1	5.6	16.6	10.8-24.7	5.4	7.5	4.9-11.1	3.6	10.9	6.3-17.7	1.8	4.3	1.9-8.9
2016-2020	13.2	20.4	15.5-26.4	7.8	25.5	17.7-35.6	5.4	15.9	10.1-24.0	4.0	6.0	3.6-9.6	1.8	5.9	2.6-11.7	2.2	6.1	2.9-11.9	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Clinton**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	207.6	377.5	354.3-401.7	107.0	472.2	429.7-517.4	100.6	327.7	299.3-358.0	116.8	220.3	202.4-239.3	64.6	303.7	268.6-341.7	52.2	172.6	152.1-195.0
	1981-1985	238.0	398.4	375.6-422.1	120.2	478.6	438.8-520.8	117.8	352.7	324.3-382.8	122.8	210.9	194.3-228.5	67.6	289.3	257.3-323.8	55.2	165.1	146.0-186.0
	1986-1990	301.4	459.9	436.6-484.0	164.4	606.5	563.3-651.8	137.0	375.5	347.5-405.0	134.4	212.1	196.2-228.9	74.4	295.5	264.6-328.7	60.0	164.7	146.4-184.6
	1991-1995	379.4	534.3	510.3-559.1	206.4	691.9	648.2-737.7	173.0	443.2	413.9-474.0	141.8	204.7	189.8-220.4	70.2	246.9	220.4-275.5	71.6	180.7	162.3-200.5
	1996-2000	389.4	522.3	499.3-546.1	210.8	648.6	608.5-690.6	178.6	438.6	410.1-468.5	150.4	206.2	191.6-221.5	80.6	270.7	243.7-299.7	69.8	167.5	150.3-186.1
	2001-2005	422.8	521.2	499.1-544.0	224.8	611.3	574.8-649.3	198.0	450.0	422.3-479.1	153.2	193.8	180.3-208.1	76.6	229.8	206.5-255.0	76.6	169.9	153.3-188.0
	2006-2010	472.6	534.6	513.0-556.8	259.8	644.7	609.0-682.0	212.8	456.1	428.7-484.8	165.6	189.4	176.7-202.9	90.8	240.6	218.1-264.7	74.8	155.5	140.0-172.3
	2011-2015	499.8	512.9	492.6-533.9	257.4	556.4	525.6-588.6	242.4	486.3	458.4-515.4	164.2	170.1	158.6-182.4	88.6	206.4	187.2-227.1	75.6	144.3	129.9-159.9
	2016-2020	508.4	488.9	469.5-509.1	265.2	523.2	494.7-553.1	243.2	467.1	439.7-495.8	166.6	155.6	145.0-166.9	88.6	179.6	162.9-197.7	78.0	137.2	123.5-152.1
Oral cavity and pharynx	1976-1980	3.4	6.2	3.5-9.9	2.0	9.0	3.9-17.1	1.4	4.9	2.0-9.9	1.2	2.0	0.7-4.3	0.6	2.0	0.4-6.1	0.6	2.1	0.4-6.0
	1981-1985	6.0	9.9	6.6-14.2	3.4	11.9	6.9-19.1	2.6	7.3	3.8-12.6	3.0	4.8	2.6-7.8	2.8	10.8	5.7-18.3	0.2	0.6	0.0-3.2
	1986-1990	5.4	8.4	5.5-12.2	3.2	11.5	6.3-19.1	2.2	6.3	3.1-11.3	2.6	4.2	2.2-7.1	1.2	4.3	1.5-9.2	1.4	3.7	1.5-7.6
	1991-1995	7.4	10.1	7.1-14.0	3.8	11.1	6.6-17.5	3.6	9.5	5.6-15.0	3.8	5.4	3.3-8.5	1.8	5.8	2.6-11.0	2.0	5.2	2.5-9.5
	1996-2000	7.2	9.4	6.6-13.1	5.0	14.2	9.1-21.1	2.2	5.2	2.6-9.3	1.6	2.2	0.9-4.3	1.2	4.0	1.4-8.8	0.4	1.0	0.1-3.6
	2001-2005	8.0	9.7	6.9-13.2	5.4	13.5	8.8-19.8	2.6	5.9	3.1-10.2	1.8	2.2	1.0-4.3	1.0	2.9	0.8-6.9	0.8	1.7	0.5-4.6
	2006-2010	9.0	10.2	7.4-13.7	6.2	14.8	9.9-21.3	2.8	6.2	3.3-10.6	2.0	2.3	1.1-4.3	0.8	2.1	0.5-5.6	1.2	2.7	1.0-6.0
	2011-2015	11.8	11.7	8.9-15.3	8.0	16.5	11.7-22.8	3.8	7.0	4.2-11.2	2.2	2.3	1.1-4.2	1.2	2.5	0.9-5.8	1.0	1.9	0.6-4.7
	2016-2020	10.6	10.2	7.6-13.6	7.2	14.3	9.9-20.1	3.4	6.6	3.7-10.9	2.4	2.1	1.1-3.9	2.0	4.0	1.9-7.7	0.4	0.6	0.1-2.8
Esophagus	1976-1980	2.0	3.8	1.8-6.9	1.6	6.8	2.9-13.3	0.4	1.4	0.2-4.8	2.0	3.6	1.7-6.6	1.6	6.7	2.8-13.2	0.4	1.3	0.2-4.6
	1981-1985	3.2	5.3	3.0-8.7	2.4	8.9	4.5-15.7	0.8	2.5	0.7-6.4	3.6	5.9	3.5-9.3	3.0	11.6	6.1-19.7	0.6	2.0	0.4-5.6
	1986-1990	3.0	4.7	2.6-7.8	2.0	7.6	3.6-13.9	1.0	2.9	0.9-6.6	3.0	4.5	2.5-7.4	2.4	8.3	4.2-14.6	0.6	1.9	0.4-5.3
	1991-1995	2.6	3.7	1.9-6.3	1.4	4.0	1.6-8.4	1.2	3.1	1.1-6.7	2.6	3.8	2.0-6.4	1.6	5.9	2.4-11.7	1.0	2.4	0.8-5.6
	1996-2000	2.2	2.8	1.4-5.1	2.0	5.5	2.6-10.3	0.2	0.6	0.0-3.0	1.4	1.8	0.7-3.7	1.2	3.6	1.3-7.9	0.2	0.5	0.0-2.8
	2001-2005	5.0	6.2	4.0-9.2	3.8	10.7	6.4-16.9	1.2	2.7	1.0-6.1	3.8	4.8	2.9-7.5	3.0	8.4	4.6-13.9	0.8	1.8	0.5-4.7
	2006-2010	8.4	9.5	6.8-12.8	6.6	17.0	11.6-24.2	1.8	3.8	1.7-7.4	5.8	6.7	4.5-9.6	4.6	12.5	7.8-19.0	1.2	2.5	0.9-5.6
	2011-2015	6.4	6.3	4.3-9.1	5.6	11.8	7.7-17.3	0.8	1.5	0.4-4.3	5.6	5.5	3.6-8.1	4.6	9.9	6.2-15.2	1.0	2.0	0.6-5.0
	2016-2020	4.6	4.3	2.7-6.6	3.6	7.0	4.1-11.3	1.0	1.8	0.6-4.6	3.8	3.5	2.1-5.6	2.8	5.0	2.7-8.8	1.0	1.8	0.6-4.6
Stomach	1976-1980	5.4	9.7	6.3-14.1	3.4	13.9	7.8-22.7	2.0	6.2	2.9-11.4	3.6	7.0	4.1-11.0	2.6	12.2	6.1-21.4	1.0	3.3	1.1-7.7
	1981-1985	5.4	9.5	6.2-13.8	4.0	17.9	10.4-28.3	1.4	4.2	1.7-8.5	2.6	4.6	2.4-7.8	1.4	5.6	2.1-11.6	1.2	3.5	1.3-7.6
	1986-1990	4.4	7.1	4.4-10.7	3.0	10.9	6.0-18.0	1.4	3.8	1.5-7.7	2.6	4.0	2.1-6.8	1.8	6.9	3.1-13.0	0.8	2.1	0.6-5.4
	1991-1995	7.6	11.2	7.9-15.4	4.4	16.2	9.9-24.7	3.2	8.2	4.7-13.3	3.6	5.4	3.2-8.5	1.8	6.5	2.9-12.3	1.8	4.6	2.1-8.7
	1996-2000	5.6	7.8	5.2-11.3	2.6	8.4	4.3-14.4	3.0	7.0	3.9-11.6	3.2	4.6	2.6-7.4	1.6	5.3	2.2-10.6	1.6	3.9	1.7-7.8
	2001-2005	8.0	10.0	7.1-13.7	4.8	12.8	8.1-19.3	3.2	7.0	4.0-11.5	3.0	3.8	2.1-6.3	1.2	3.5	1.2-7.6	1.8	4.0	1.8-7.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Clinton**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	6.2	7.1	4.8-10.1	3.4	8.9	5.0-14.5	2.8	5.6	3.1-9.6	2.8	3.2	1.8-5.4	1.6	4.0	1.7-8.2	1.2	2.5	0.9-5.7
	2011-2015	6.8	7.0	4.8-9.9	4.0	8.7	5.2-13.7	2.8	5.6	3.0-9.8	2.0	2.2	1.0-4.1	1.6	4.0	1.7-8.0	0.4	0.9	0.1-3.4
	2016-2020	7.0	6.7	4.6-9.6	5.8	11.6	7.7-17.0	1.2	2.4	0.8-5.8	3.6	3.4	2.0-5.6	3.0	5.8	3.1-9.9	0.6	1.3	0.3-4.2
Colorectal	1976-1980	33.6	63.5	54.1-74.1	15.2	68.0	52.6-86.2	18.4	60.2	48.5-73.9	18.6	38.5	30.9-47.3	8.2	45.3	31.4-62.5	10.4	35.2	26.3-46.2
	1981-1985	34.6	61.3	52.4-71.2	16.4	69.6	54.5-87.2	18.2	55.5	44.6-68.2	16.8	29.7	23.6-36.8	9.8	42.9	31.1-57.5	7.0	21.3	14.8-29.7
	1986-1990	38.4	59.8	51.5-68.9	20.4	77.4	62.4-94.7	18.0	48.4	38.8-59.5	17.4	28.1	22.5-34.7	8.6	35.5	25.2-48.2	8.8	24.7	17.9-33.2
	1991-1995	55.0	79.9	70.7-89.9	29.0	106.0	88.6-125.6	26.0	64.4	53.8-76.5	21.4	31.5	25.7-38.0	11.2	41.6	30.9-54.6	10.2	25.3	18.8-33.3
	1996-2000	43.4	59.0	51.4-67.4	22.2	72.1	58.8-87.4	21.2	49.5	40.5-59.9	17.2	24.0	19.2-29.6	9.2	32.2	23.2-43.2	8.0	18.3	13.1-25.0
	2001-2005	53.8	67.0	59.2-75.5	27.0	74.6	62.1-88.9	26.8	60.1	50.3-71.3	16.0	20.3	16.1-25.3	8.2	23.6	16.7-32.4	7.8	17.0	12.1-23.4
	2006-2010	41.0	46.8	40.6-53.8	21.0	52.2	42.4-63.5	20.0	41.7	33.8-50.8	14.6	17.0	13.3-21.5	7.4	19.7	13.7-27.5	7.2	14.9	10.4-20.8
	2011-2015	38.0	39.7	34.2-45.9	20.0	44.4	35.9-54.2	18.0	34.6	27.7-42.9	12.4	13.1	10.0-16.9	7.8	17.9	12.7-24.7	4.6	8.7	5.5-13.3
2016-2020	39.6	39.9	34.4-46.2	21.8	45.5	37.1-55.2	17.8	34.5	27.3-43.2	13.6	12.9	10.0-16.6	8.6	18.0	13.0-24.5	5.0	8.4	5.4-12.9	
Colon excluding rectum	1976-1980	23.2	43.7	35.9-52.5	9.6	41.6	30.0-55.9	13.6	44.3	34.3-56.1	13.6	28.1	21.7-35.7	5.6	31.8	20.0-47.0	8.0	27.2	19.4-36.9
	1981-1985	24.8	43.9	36.4-52.5	11.6	50.5	37.6-66.1	13.2	39.8	30.7-50.7	14.2	24.8	19.3-31.3	8.4	37.0	25.9-50.7	5.8	17.5	11.7-25.2
	1986-1990	26.6	41.4	34.6-49.1	12.0	46.3	34.7-60.2	14.6	38.7	30.3-48.7	13.8	22.0	17.1-27.8	6.8	27.9	18.9-39.4	7.0	19.8	13.7-27.6
	1991-1995	37.6	54.5	46.9-62.9	19.0	68.6	54.8-84.4	18.6	46.1	37.2-56.6	18.2	26.8	21.5-32.9	9.4	34.8	25.0-46.8	8.8	22.0	15.9-29.5
	1996-2000	33.4	45.8	39.1-53.3	15.2	51.0	39.7-64.3	18.2	42.5	34.2-52.3	14.2	19.7	15.4-24.9	6.8	23.7	16.2-33.3	7.4	17.0	11.9-23.5
	2001-2005	41.2	51.3	44.5-58.8	19.0	52.2	41.8-64.2	22.2	49.3	40.6-59.5	12.4	15.8	12.1-20.3	5.8	17.0	11.2-24.8	6.6	14.3	9.8-20.2
	2006-2010	30.2	34.9	29.5-41.0	14.8	37.5	29.3-47.4	15.4	32.0	25.2-40.2	10.6	12.7	9.5-16.6	4.6	12.8	8.0-19.3	6.0	12.4	8.4-17.9
	2011-2015	28.8	30.2	25.4-35.6	14.2	31.8	24.7-40.3	14.6	28.1	21.9-35.7	10.6	11.3	8.4-14.8	6.4	14.5	9.8-20.7	4.2	8.0	4.9-12.5
2016-2020	26.6	26.4	22.0-31.5	14.0	29.7	23.0-37.8	12.6	22.7	17.3-29.6	11.2	10.6	8.0-13.9	7.6	16.1	11.3-22.4	3.6	5.7	3.4-9.5	
Rectum & rectosigmoid	1976-1980	10.4	19.9	14.7-26.1	5.6	26.4	17.0-38.8	4.8	16.0	10.2-23.7	5.0	10.4	6.7-15.3	2.6	13.5	7.0-23.0	2.4	8.1	4.1-14.0
	1981-1985	9.8	17.3	12.8-22.9	4.8	19.0	12.0-28.5	5.0	15.6	10.1-23.0	2.6	4.9	2.6-8.3	1.4	6.0	2.3-12.2	1.2	3.8	1.4-8.2
	1986-1990	11.8	18.4	13.9-23.7	8.4	31.1	22.1-42.3	3.4	9.7	5.6-15.5	3.6	6.2	3.6-9.7	1.8	7.5	3.2-14.3	1.8	4.9	2.2-9.2
	1991-1995	17.4	25.4	20.3-31.3	10.0	37.5	27.3-49.9	7.4	18.3	12.8-25.2	3.2	4.7	2.7-7.6	1.8	6.9	3.0-13.0	1.4	3.4	1.3-7.0
	1996-2000	10.0	13.2	9.8-17.4	7.0	21.1	14.5-29.6	3.0	7.0	3.9-11.6	3.0	4.2	2.4-7.0	2.4	8.5	4.2-14.9	0.6	1.3	0.3-4.0
	2001-2005	12.6	15.7	12.0-20.1	8.0	22.4	15.8-30.9	4.6	10.7	6.8-16.2	3.6	4.4	2.6-7.1	2.4	6.6	3.4-11.6	1.2	2.8	1.0-6.1
	2006-2010	10.8	11.9	8.9-15.6	6.2	14.7	9.8-21.0	4.6	9.6	6.1-14.6	4.0	4.4	2.7-6.8	2.8	7.0	3.7-12.0	1.2	2.5	0.9-5.6
	2011-2015	9.2	9.5	6.9-12.8	5.8	12.6	8.4-18.3	3.4	6.5	3.7-10.8	1.8	1.8	0.8-3.6	1.4	3.4	1.3-7.2	0.4	0.7	0.1-3.0
2016-2020	13.0	13.6	10.4-17.5	7.8	15.8	11.1-21.9	5.2	11.8	7.4-17.8	2.4	2.3	1.2-4.2	1.0	1.9	0.6-4.8	1.4	2.7	1.0-6.0	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Clinton**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	1.6	2.4	1.0-4.8	0.6	2.0	0.4-6.1	1.0	2.8	0.9-6.5	0.6	0.9	0.2-2.7	0.2	0.7	0.0-4.2	0.4	1.1	0.1-4.0
	1981-1985	2.2	3.6	1.8-6.5	1.0	3.7	1.2-8.7	1.2	3.1	1.1-7.0	1.4	2.3	0.9-4.6	1.2	4.1	1.5-9.0	0.2	0.6	0.0-3.2
	1986-1990	1.2	2.0	0.7-4.3	0.6	2.5	0.5-7.1	0.6	1.7	0.3-4.9	1.0	1.6	0.5-3.6	0.6	2.5	0.5-7.1	0.4	1.1	0.1-4.0
	1991-1995	2.0	2.8	1.4-5.2	1.0	3.2	1.0-7.4	1.0	2.5	0.8-5.8	1.6	2.3	1.0-4.5	0.8	2.6	0.7-6.6	0.8	1.9	0.5-4.9
	1996-2000	2.0	2.8	1.3-5.1	1.2	3.8	1.3-8.2	0.8	1.9	0.5-5.0	2.2	2.9	1.4-5.2	0.8	2.1	0.6-5.6	1.4	3.4	1.4-7.1
	2001-2005	5.6	7.0	4.6-10.1	4.2	12.0	7.3-18.6	1.4	3.3	1.3-6.8	3.2	4.2	2.4-6.8	2.2	7.0	3.4-12.7	1.0	2.2	0.7-5.4
	2006-2010	7.6	8.4	5.9-11.6	5.8	13.5	9.0-19.6	1.8	3.7	1.7-7.2	4.8	5.5	3.5-8.2	3.8	9.7	5.8-15.2	1.0	2.0	0.6-4.9
	2011-2015	7.4	7.0	4.9-9.8	5.8	12.0	7.9-17.6	1.6	2.9	1.2-6.0	5.0	4.9	3.2-7.4	3.2	6.6	3.7-11.1	1.8	3.5	1.6-6.9
	2016-2020	10.6	10.0	7.4-13.3	7.2	14.0	9.7-19.8	3.4	5.9	3.3-9.9	5.6	5.2	3.4-7.7	3.4	6.6	3.8-10.9	2.2	3.9	1.8-7.5
Pancreas	1976-1980	5.0	9.2	5.9-13.5	2.0	8.6	4.0-15.8	3.0	9.9	5.5-16.3	4.8	8.6	5.5-12.8	1.6	6.3	2.7-12.4	3.2	10.4	5.9-16.9
	1981-1985	7.8	13.6	9.6-18.6	3.8	16.1	9.5-25.2	4.0	11.4	6.9-17.6	7.6	13.6	9.5-18.6	3.4	15.2	8.7-24.4	4.2	12.2	7.5-18.6
	1986-1990	7.0	10.8	7.5-15.1	4.4	16.8	10.1-25.8	2.6	7.0	3.7-11.9	6.4	10.4	7.1-14.6	4.2	18.6	11.0-28.8	2.2	6.0	3.0-10.8
	1991-1995	6.6	9.7	6.7-13.7	3.6	12.9	7.4-20.7	3.0	7.5	4.2-12.4	5.6	8.2	5.5-11.9	2.4	8.8	4.3-15.6	3.2	8.0	4.6-13.0
	1996-2000	7.4	10.2	7.2-14.1	3.8	13.6	7.9-21.4	3.6	8.5	5.0-13.5	8.0	10.9	7.8-14.8	5.4	18.2	11.7-26.9	2.6	6.0	3.2-10.3
	2001-2005	12.2	15.4	11.8-19.8	6.4	18.8	12.7-26.8	5.8	12.8	8.5-18.4	11.2	14.1	10.7-18.4	5.8	17.4	11.5-25.2	5.4	11.8	7.8-17.3
	2006-2010	10.8	12.0	9.0-15.8	6.2	14.9	10.0-21.3	4.6	9.0	5.7-13.7	8.8	9.7	7.0-13.0	5.2	12.0	7.7-17.8	3.6	7.3	4.3-11.7
	2011-2015	13.4	13.1	10.2-16.8	7.0	14.2	9.8-20.0	6.4	11.7	8.0-16.8	10.4	10.3	7.7-13.7	5.4	11.5	7.5-16.9	5.0	8.9	5.7-13.4
	2016-2020	16.0	14.2	11.2-17.9	8.0	14.7	10.4-20.3	8.0	13.6	9.7-18.9	12.2	11.1	8.5-14.4	6.4	12.4	8.4-17.8	5.8	10.1	6.7-14.9
Larynx	1976-1980	3.0	5.6	3.1-9.3	2.4	11.5	5.6-20.4	0.6	2.1	0.4-6.0	1.6	3.2	1.3-6.2	1.2	6.2	2.0-13.8	0.4	1.3	0.2-4.6
	1981-1985	4.0	6.5	3.9-10.0	3.6	13.4	7.8-21.3	0.4	1.2	0.1-4.4	0.8	1.3	0.3-3.2	0.6	2.3	0.4-6.6	0.2	0.5	0.0-3.0
	1986-1990	5.2	7.8	5.1-11.4	3.8	13.6	8.1-21.4	1.4	3.7	1.5-7.6	0.6	0.8	0.2-2.5	0.2	0.6	0.0-3.5	0.4	1.0	0.1-3.7
	1991-1995	5.4	7.7	5.1-11.2	3.8	11.7	7.0-18.3	1.6	4.1	1.8-8.1	2.0	2.8	1.3-5.1	1.2	3.9	1.4-8.6	0.8	1.9	0.5-4.8
	1996-2000	4.4	5.6	3.5-8.5	3.0	8.7	4.8-14.5	1.4	3.4	1.3-7.0	1.2	1.6	0.6-3.5	1.2	3.7	1.3-8.2	0.0	0.0	0.0-1.9
	2001-2005	6.6	8.1	5.6-11.4	5.4	14.4	9.3-21.1	1.2	2.5	0.9-5.7	1.0	1.3	0.4-3.0	0.6	1.7	0.3-4.9	0.4	0.9	0.1-3.4
	2006-2010	5.4	5.9	3.9-8.7	3.8	8.3	5.0-13.2	1.6	3.5	1.5-7.1	0.8	0.9	0.3-2.4	0.6	1.4	0.3-4.3	0.2	0.5	0.0-2.7
	2011-2015	5.8	5.6	3.7-8.2	4.4	9.2	5.7-14.2	1.4	2.7	1.1-5.8	0.2	0.2	0.0-1.4	0.0	0.0	0.0-1.9	0.2	0.5	0.0-2.9
	2016-2020	5.2	4.5	2.9-6.8	3.8	6.7	4.0-10.9	1.4	2.1	0.9-4.9	1.0	0.9	0.3-2.3	0.8	1.4	0.4-4.0	0.2	0.4	0.0-2.7
Lung and bronchus	1976-1980	38.4	66.9	57.7-77.2	28.4	116.6	97.2-138.4	10.0	31.7	23.4-41.8	33.0	57.8	49.2-67.4	25.4	105.7	87.1-127.0	7.6	24.8	17.5-34.0
	1981-1985	40.4	67.0	58.0-77.0	29.0	117.9	98.3-139.9	11.4	33.8	25.5-43.8	30.6	51.6	43.7-60.6	22.8	99.2	80.8-120.3	7.8	22.4	15.9-30.7
	1986-1990	62.0	95.0	84.7-106.3	41.2	150.5	129.8-173.5	20.8	56.5	46.1-68.6	44.6	68.7	59.9-78.4	31.2	118.2	99.5-139.1	13.4	36.2	28.0-46.0
	1991-1995	56.8	79.9	70.8-89.8	33.6	107.8	91.6-125.8	23.2	58.6	48.4-70.4	36.8	52.3	45.0-60.4	21.2	69.0	56.0-84.0	15.6	39.7	31.4-49.6
	1996-2000	63.2	85.7	76.4-95.7	36.6	115.3	98.6-133.9	26.6	66.0	55.3-78.3	44.8	60.7	53.0-69.3	26.4	83.3	69.2-99.2	18.4	45.2	36.4-55.5
	2001-2005	70.6	87.8	78.8-97.5	39.8	112.8	97.1-130.3	30.8	70.3	59.6-82.4	45.0	56.5	49.3-64.4	25.6	74.8	61.9-89.5	19.4	43.8	35.5-53.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Clinton**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	82.4	93.5	84.6-103.1	44.4	115.3	100.2-132.1	38.0	80.4	69.3-92.8	56.2	64.2	56.8-72.2	32.0	84.4	71.4-99.0	24.2	50.7	42.0-60.7
	2011-2015	91.8	93.5	85.1-102.7	47.2	104.2	91.0-118.8	44.6	86.8	75.6-99.3	50.6	52.0	45.7-58.9	26.6	60.9	50.8-72.5	24.0	46.5	38.5-55.9
	2016-2020	106.6	97.4	89.1-106.3	53.0	102.1	89.9-115.6	53.6	93.9	82.7-106.4	49.6	45.1	39.6-51.3	25.8	50.3	41.8-60.1	23.8	40.6	33.6-49.0
Melanoma of the skin	1976-1980	2.6	4.3	2.2-7.4	1.4	4.6	1.7-10.0	1.2	4.1	1.4-8.8	0.8	1.4	0.4-3.7	0.6	2.2	0.4-6.9	0.2	0.9	0.0-4.1
	1981-1985	2.8	4.4	2.3-7.4	1.0	3.2	1.0-7.7	1.8	5.2	2.3-9.9	1.0	1.3	0.4-3.2	0.6	1.9	0.4-5.8	0.4	0.8	0.1-3.3
	1986-1990	3.8	5.8	3.5-9.1	2.0	7.9	3.5-14.6	1.8	5.0	2.2-9.5	0.4	0.6	0.1-2.1	0.2	0.8	0.0-4.1	0.2	0.5	0.0-2.8
	1991-1995	6.2	8.4	5.6-11.9	4.0	12.0	7.1-18.7	2.2	5.8	2.9-10.4	2.0	2.6	1.2-4.8	1.4	3.8	1.4-8.0	0.6	1.5	0.3-4.3
	1996-2000	6.6	8.6	5.9-12.0	4.2	11.9	7.3-18.5	2.4	6.2	3.2-10.9	1.8	2.5	1.1-4.7	1.4	4.3	1.7-9.0	0.4	1.0	0.1-3.6
	2001-2005	12.0	14.4	11.0-18.6	7.2	18.7	12.9-26.2	4.8	11.6	7.4-17.3	2.4	2.9	1.5-5.2	1.6	4.1	1.8-8.3	0.8	1.8	0.5-4.8
	2006-2010	15.2	17.5	13.7-21.9	9.2	22.4	16.2-30.0	6.0	13.4	9.0-19.4	2.2	2.6	1.3-4.7	1.6	3.8	1.6-7.7	0.6	1.3	0.3-3.9
	2011-2015	23.2	24.5	20.1-29.5	12.0	25.7	19.5-33.4	11.2	23.9	17.8-31.5	1.8	1.9	0.9-3.8	1.4	3.4	1.3-7.1	0.4	0.9	0.1-3.4
2016-2020	21.2	21.7	17.6-26.5	11.4	23.7	17.8-31.0	9.8	21.2	15.3-28.6	1.4	1.3	0.5-2.8	1.0	2.2	0.7-5.4	0.4	0.6	0.1-3.0	
Female breast	1976-1980							21.4	72.5	59.3-87.7							7.8	25.6	18.2-35.0
	1981-1985							34.2	107.3	91.6-124.8							12.6	39.0	29.9-49.9
	1986-1990							42.4	119.9	104.1-137.3							12.4	35.4	27.1-45.5
	1991-1995							48.4	128.5	112.7-145.8							10.4	27.4	20.4-36.0
	1996-2000							50.8	126.9	111.7-143.6							9.2	22.5	16.5-30.1
	2001-2005							51.8	118.4	104.3-133.8							8.0	17.5	12.5-24.0
	2006-2010							55.0	118.8	105.0-133.9							7.8	16.2	11.5-22.4
	2011-2015							56.4	115.5	102.0-130.4							7.6	14.8	10.3-20.6
2016-2020							59.4	118.8	104.8-134.2							8.8	15.9	11.4-21.8	
Cervix uteri	1976-1980							6.0	20.0	13.4-28.6							1.2	4.0	1.4-8.6
	1981-1985							2.8	7.5	3.9-12.8							0.8	2.5	0.7-6.3
	1986-1990							3.6	9.5	5.6-15.0							1.2	3.2	1.2-6.9
	1991-1995							2.2	5.6	2.7-10.0							0.8	1.8	0.5-4.8
	1996-2000							3.8	9.6	5.8-15.0							0.4	1.1	0.1-3.8
	2001-2005							2.0	4.8	2.3-8.9							1.2	2.8	1.0-6.1
	2006-2010							2.6	6.4	3.4-11.0							0.6	1.3	0.3-4.0
	2011-2015							3.2	8.3	4.6-13.6							1.6	3.6	1.5-7.4
	2016-2020							2.6	7.4	3.8-12.8							0.8	1.9	0.4-5.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Clinton**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							7.8	25.3	18.0-34.6							1.8	6.0	2.7-11.3
	1981-1985							5.8	17.7	11.8-25.5							1.2	3.4	1.3-7.5
	1986-1990							7.4	19.7	13.8-27.2							1.2	3.2	1.2-6.9
	1991-1995							10.6	28.2	21.1-36.9							0.8	2.1	0.6-5.3
	1996-2000							10.8	27.1	20.3-35.4							0.8	1.9	0.5-4.9
	2001-2005							9.6	22.0	16.2-29.2							2.4	5.3	2.7-9.4
	2006-2010							12.4	26.7	20.4-34.5							1.6	3.3	1.4-6.8
	2011-2015							16.2	31.5	24.8-39.5							2.4	4.8	2.5-8.6
	2016-2020							11.8	19.9	15.0-26.2							1.8	3.3	1.4-6.8
Ovary	1976-1980							5.2	16.1	10.4-23.6							2.6	8.5	4.5-14.4
	1981-1985							5.4	15.6	10.1-22.8							4.6	13.9	8.8-20.9
	1986-1990							5.6	15.2	10.0-22.1							2.2	6.1	3.0-10.8
	1991-1995							6.8	18.0	12.4-25.1							6.2	15.8	10.7-22.5
	1996-2000							6.0	14.9	10.1-21.4							4.0	9.6	5.8-14.9
	2001-2005							5.8	13.2	8.8-19.1							4.4	10.0	6.2-15.2
	2006-2010							6.2	12.9	8.7-18.6							4.8	10.0	6.4-15.1
	2011-2015							6.2	13.0	8.7-18.9							3.2	5.5	3.2-9.3
	2016-2020							5.4	10.4	6.7-15.7							3.2	5.6	3.2-9.5
Prostate	1976-1980				15.8	91.0	70.5-114.9							6.2	42.2	27.7-60.6			
	1981-1985				18.4	80.6	64.0-99.7							4.4	23.0	14.0-35.1			
	1986-1990				34.8	147.8	125.7-172.3							7.8	35.5	24.8-48.9			
	1991-1995				64.8	229.9	204.4-257.4							7.6	33.3	23.2-45.8			
	1996-2000				66.8	202.6	180.8-226.2							6.8	28.4	19.3-39.7			
	2001-2005				50.6	133.3	117.0-151.3							7.2	27.3	18.8-37.8			
	2006-2010				72.2	169.7	152.2-188.7							5.0	15.9	10.1-23.5			
	2011-2015				44.8	90.8	79.0-103.9							6.4	16.8	11.4-23.8			
	2016-2020				51.6	92.9	81.6-105.4							6.8	14.4	9.9-20.4			
Testis	1976-1980				1.8	4.8	2.1-9.8							0.0	0.0	0.0-3.0			
	1981-1985				1.6	4.5	1.8-9.5							0.0	0.0	0.0-2.8			
	1986-1990				2.2	3.8	1.8-7.6							0.2	0.7	0.0-3.8			
	1991-1995				2.4	4.3	2.2-8.0							0.0	0.0	0.0-2.3			
	1996-2000				3.6	8.2	4.8-13.2							0.0	0.0	0.0-2.2			
	2001-2005				2.6	5.7	3.0-9.9							0.0	0.0	0.0-2.1			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Clinton**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				4.0	9.6	5.8-15.0							0.2	0.3	0.0-2.5			
	2011-2015				2.8	6.9	3.7-11.7							0.0	0.0	0.0-1.9			
	2016-2020				3.4	8.0	4.6-12.9							0.0	0.0	0.0-1.9			
Urinary bladder (incl. in situ)	1976-1980	12.2	22.5	17.1-29.0	8.8	38.2	27.0-52.2	3.4	11.3	6.6-18.0	2.0	4.3	2.0-7.8	1.0	5.2	1.4-12.4	1.0	3.6	1.2-8.1
	1981-1985	13.4	22.9	17.7-29.1	9.8	38.1	27.9-50.6	3.6	10.7	6.4-16.9	3.0	5.2	2.9-8.5	2.4	10.6	5.3-18.5	0.6	1.8	0.4-5.1
	1986-1990	17.8	27.2	21.8-33.5	13.2	47.9	36.5-61.5	4.6	12.1	7.7-18.2	5.0	8.0	5.2-11.8	3.8	15.5	9.1-24.4	1.2	3.1	1.1-6.8
	1991-1995	23.4	33.0	27.2-39.6	17.8	62.2	49.2-77.4	5.6	13.7	9.1-19.9	5.0	7.3	4.7-10.8	3.8	14.5	8.4-22.9	1.2	3.0	1.1-6.6
	1996-2000	24.0	32.7	27.1-39.1	17.8	56.0	44.5-69.4	6.2	15.1	10.2-21.5	5.8	7.9	5.3-11.4	3.4	12.8	7.2-20.8	2.4	5.6	2.9-9.8
	2001-2005	27.4	33.9	28.4-40.1	20.6	57.1	46.3-69.6	6.8	15.1	10.5-21.2	5.2	6.7	4.4-9.8	3.2	9.1	5.1-14.8	2.0	4.2	2.0-7.9
	2006-2010	29.0	32.6	27.5-38.5	20.2	52.5	42.4-64.3	8.8	19.2	13.9-25.9	5.6	6.5	4.3-9.5	4.4	13.2	8.1-20.2	1.2	2.3	0.8-5.3
	2011-2015	27.2	28.0	23.4-33.2	20.6	46.5	37.8-56.8	6.6	12.3	8.5-17.6	4.4	4.7	2.9-7.1	2.8	7.0	3.8-11.9	1.6	3.1	1.3-6.4
	2016-2020	25.4	23.6	19.6-28.2	20.2	40.2	32.5-49.2	5.2	8.9	5.7-13.4	4.6	4.3	2.7-6.6	3.4	6.9	4.0-11.3	1.2	1.9	0.7-4.6
Kidney and renal pelvis	1976-1980	4.2	7.4	4.5-11.4	1.6	6.4	2.7-12.9	2.6	8.3	4.3-14.2	2.8	5.1	2.8-8.6	1.0	4.4	1.3-10.3	1.8	5.8	2.6-11.0
	1981-1985	7.4	12.1	8.5-16.7	4.4	15.7	9.7-24.0	3.0	8.9	4.9-14.6	3.0	4.8	2.7-7.9	1.6	6.0	2.5-11.9	1.4	4.0	1.6-8.2
	1986-1990	5.2	7.7	5.0-11.3	3.4	12.1	6.8-19.7	1.8	4.5	2.0-8.6	2.6	3.8	2.0-6.5	1.4	5.4	1.9-11.3	1.2	3.0	1.1-6.6
	1991-1995	9.4	13.3	9.8-17.7	5.0	15.7	9.9-23.6	4.4	11.9	7.4-17.9	2.6	4.0	2.1-6.7	1.6	6.0	2.4-11.9	1.0	2.5	0.8-5.8
	1996-2000	13.4	17.9	13.8-22.7	7.8	23.3	16.4-32.1	5.6	13.6	9.0-19.7	4.4	5.9	3.7-8.9	2.6	8.1	4.1-14.0	1.8	4.1	1.9-7.9
	2001-2005	16.6	20.3	16.2-25.2	9.2	25.2	18.3-34.0	7.4	16.9	11.9-23.4	3.0	3.7	2.0-6.1	1.6	4.4	1.8-8.7	1.4	3.1	1.3-6.6
	2006-2010	22.2	24.8	20.3-29.9	13.4	32.6	25.0-41.8	8.8	18.3	13.2-24.8	4.4	4.8	3.0-7.3	2.4	5.2	2.7-9.3	2.0	4.0	1.9-7.6
	2011-2015	26.8	27.6	23.1-32.9	18.2	38.2	30.6-47.3	8.6	18.2	13.0-24.9	3.0	3.1	1.7-5.3	2.2	4.8	2.4-8.9	0.8	1.6	0.4-4.4
	2016-2020	26.0	25.7	21.3-30.8	16.4	32.2	25.4-40.4	9.6	18.9	13.7-25.7	5.0	4.8	3.1-7.2	2.8	5.7	3.1-9.9	2.2	4.0	2.0-7.6
Brain and other central nervous system	1976-1980	4.0	6.1	3.7-9.6	2.6	8.7	4.5-15.3	1.4	4.0	1.6-8.4	3.4	5.2	3.0-8.5	2.2	7.0	3.4-12.9	1.2	3.6	1.3-7.9
	1981-1985	3.0	4.1	2.2-6.8	1.0	3.1	0.9-7.7	2.0	5.3	2.5-10.0	2.0	3.0	1.4-5.6	1.2	4.5	1.6-9.9	0.8	1.8	0.5-4.9
	1986-1990	7.0	9.7	6.7-13.5	5.0	13.7	8.7-20.6	2.0	5.7	2.7-10.5	2.2	2.9	1.4-5.2	1.6	4.6	2.0-9.1	0.6	1.5	0.3-4.5
	1991-1995	4.6	6.0	3.8-9.1	2.6	6.6	3.4-11.6	2.0	5.1	2.4-9.3	3.0	4.3	2.4-7.0	1.0	3.0	0.9-7.1	2.0	5.3	2.5-9.7
	1996-2000	6.6	8.6	5.9-12.1	3.6	9.3	5.5-14.9	3.0	7.4	4.1-12.2	2.4	3.1	1.6-5.5	2.0	5.1	2.4-9.5	0.4	1.0	0.1-3.7
	2001-2005	6.0	7.5	5.0-10.7	3.4	8.2	4.7-13.3	2.6	5.9	3.1-10.2	3.2	4.0	2.3-6.5	1.2	2.8	1.0-6.2	2.0	4.3	2.1-8.1
	2006-2010	6.2	7.5	5.0-10.7	3.6	9.5	5.5-15.2	2.6	5.7	3.0-9.8	4.0	4.3	2.6-6.7	2.8	5.8	3.1-9.9	1.2	2.6	1.0-5.9
	2011-2015	6.8	7.2	4.9-10.2	3.6	8.2	4.7-13.2	3.2	6.0	3.4-10.1	4.6	5.1	3.2-7.8	2.8	6.6	3.5-11.2	1.8	3.4	1.5-6.7
	2016-2020	6.4	6.4	4.3-9.2	3.6	7.8	4.5-12.6	2.8	5.2	2.7-9.2	3.6	3.5	2.0-5.7	1.8	3.7	1.6-7.3	1.8	3.1	1.4-6.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Clinton**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	1.2	1.7	0.5-3.8	0.4	1.2	0.1-5.0	0.8	2.0	0.5-5.5	0.8	1.5	0.4-3.7	0.4	1.7	0.2-6.1	0.4	1.2	0.1-4.4
	1981-1985	2.0	2.5	1.1-4.8	0.2	0.8	0.0-4.3	1.8	4.5	1.9-8.8	0.4	0.6	0.1-2.3	0.2	0.8	0.0-4.3	0.2	0.6	0.0-3.2
	1986-1990	1.6	2.2	0.9-4.5	0.2	0.7	0.0-3.9	1.4	3.7	1.4-7.9	0.0	0.0	0.0-1.0	0.0	0.0	0.0-2.5	0.0	0.0	0.0-1.9
	1991-1995	1.6	2.1	0.9-4.1	0.4	0.9	0.1-3.6	1.2	3.2	1.2-7.0	0.0	0.0	0.0-1.0	0.0	0.0	0.0-2.3	0.0	0.0	0.0-1.8
	1996-2000	3.2	4.0	2.3-6.5	1.0	2.5	0.8-6.0	2.2	5.7	2.8-10.2	0.0	0.0	0.0-1.0	0.0	0.0	0.0-2.2	0.0	0.0	0.0-1.9
	2001-2005	5.2	6.3	4.1-9.2	1.4	3.2	1.3-6.8	3.8	9.4	5.6-14.8	0.2	0.3	0.0-1.5	0.0	0.0	0.0-2.1	0.2	0.4	0.0-2.5
	2006-2010	9.4	10.9	8.0-14.6	3.0	6.5	3.6-10.9	6.4	15.6	10.6-22.2	0.2	0.2	0.0-1.4	0.0	0.0	0.0-2.0	0.2	0.4	0.0-2.6
	2011-2015	14.6	16.2	12.6-20.6	4.4	8.8	5.5-13.7	10.2	24.7	18.1-32.9	0.4	0.5	0.1-1.7	0.2	0.6	0.0-3.2	0.2	0.4	0.0-2.7
	2016-2020	11.2	13.6	10.1-17.9	3.0	6.7	3.7-11.4	8.2	21.4	15.0-29.5	0.6	0.7	0.1-2.0	0.2	0.5	0.0-2.8	0.4	0.9	0.1-3.5
Hodgkin lymphoma	1976-1980	0.8	0.9	0.2-2.7	0.6	0.9	0.2-4.1	0.2	0.7	0.0-3.6	0.2	0.3	0.0-1.8	0.0	0.0	0.0-3.0	0.2	0.6	0.0-3.4
	1981-1985	3.6	5.0	2.9-8.1	1.8	5.0	2.2-10.0	1.8	4.5	2.0-8.8	1.0	1.5	0.5-3.7	0.4	1.2	0.1-5.1	0.6	1.8	0.4-5.2
	1986-1990	2.6	3.1	1.6-5.4	1.4	2.7	1.0-6.3	1.2	3.0	1.1-6.6	0.4	0.7	0.1-2.3	0.4	1.3	0.2-4.8	0.0	0.0	0.0-1.9
	1991-1995	2.0	2.2	1.0-4.1	1.2	2.4	0.9-5.7	0.8	1.7	0.5-4.6	0.2	0.2	0.0-1.2	0.2	0.3	0.0-2.7	0.0	0.0	0.0-1.8
	1996-2000	2.6	3.2	1.7-5.5	1.6	3.5	1.5-7.2	1.0	2.6	0.8-6.0	0.4	0.5	0.1-1.7	0.0	0.0	0.0-2.2	0.4	0.9	0.1-3.4
	2001-2005	2.0	2.3	1.1-4.3	1.4	3.1	1.2-6.6	0.6	1.4	0.3-4.2	0.2	0.2	0.0-1.4	0.0	0.0	0.0-2.1	0.2	0.4	0.0-2.7
	2006-2010	1.8	2.3	1.0-4.3	1.0	2.2	0.7-5.4	0.8	2.4	0.6-6.0	0.4	0.5	0.1-1.9	0.0	0.0	0.0-2.0	0.4	1.1	0.1-3.9
	2011-2015	3.0	3.3	1.8-5.6	1.2	2.8	1.0-6.2	1.8	3.8	1.7-7.5	0.0	0.0	0.0-1.0	0.0	0.0	0.0-1.9	0.0	0.0	0.0-2.0
	2016-2020	2.6	3.1	1.6-5.4	1.2	2.7	0.9-6.0	1.4	3.5	1.3-7.5	0.0	0.0	0.0-1.0	0.0	0.0	0.0-1.9	0.0	0.0	0.0-2.1
Non-Hodgkin lymphomas	1976-1980	7.6	14.0	9.8-19.2	4.2	17.5	10.4-27.4	3.4	11.4	6.6-18.2	4.8	9.3	5.9-13.8	2.0	8.7	4.1-16.0	2.8	9.6	5.2-16.0
	1981-1985	6.8	11.0	7.6-15.5	3.0	10.1	5.5-17.1	3.8	11.2	6.6-17.5	3.4	6.0	3.5-9.6	1.6	6.9	2.8-13.7	1.8	5.2	2.3-9.8
	1986-1990	12.2	17.5	13.3-22.7	7.2	22.1	14.9-31.3	5.0	13.7	8.8-20.3	5.4	8.7	5.7-12.6	2.2	9.7	4.6-17.2	3.2	8.6	4.9-13.9
	1991-1995	13.8	18.7	14.5-23.8	7.6	23.5	16.1-33.0	6.2	15.1	10.2-21.4	4.8	6.6	4.2-9.8	3.0	9.0	4.9-14.9	1.8	4.3	2.0-8.3
	1996-2000	16.6	22.3	17.8-27.7	9.6	29.7	21.6-39.8	7.0	17.3	12.0-24.1	6.8	9.4	6.5-13.2	3.6	12.7	7.3-20.2	3.2	7.7	4.4-12.5
	2001-2005	18.6	22.8	18.4-28.0	9.0	23.0	16.6-31.0	9.6	21.4	15.7-28.4	6.6	8.4	5.8-11.8	3.8	10.5	6.2-16.6	2.8	6.1	3.4-10.4
	2006-2010	16.8	19.5	15.5-24.1	9.4	23.3	17.0-31.2	7.4	16.1	11.3-22.3	4.2	4.9	3.0-7.6	2.0	5.2	2.4-9.6	2.2	4.5	2.2-8.2
	2011-2015	26.6	27.3	22.8-32.5	14.6	32.5	25.3-41.2	12.0	23.6	17.9-30.7	7.8	8.2	5.8-11.3	5.0	12.6	8.1-18.8	2.8	5.4	2.9-9.4
	2016-2020	22.4	21.8	17.8-26.4	12.2	25.3	19.2-32.9	10.2	19.8	14.4-26.6	6.2	6.0	4.1-8.6	4.0	8.6	5.2-13.5	2.2	4.2	2.0-7.9
Myeloma	1976-1980	2.2	4.1	2.0-7.4	1.4	7.1	2.5-15.0	0.8	2.7	0.7-6.8	1.2	2.6	0.9-5.5	0.6	4.2	0.7-11.6	0.6	2.1	0.4-6.0
	1981-1985	2.0	3.1	1.5-5.8	1.4	5.1	1.9-10.8	0.6	1.9	0.4-5.4	1.6	2.8	1.2-5.4	1.0	4.3	1.3-9.9	0.6	2.0	0.4-5.5
	1986-1990	4.0	6.3	3.8-9.7	1.8	6.9	3.1-13.0	2.2	6.1	3.0-10.9	2.0	3.2	1.5-5.9	0.6	2.7	0.5-7.3	1.4	3.6	1.4-7.4
	1991-1995	6.4	9.0	6.1-12.7	3.6	11.1	6.5-17.6	2.8	6.7	3.6-11.2	3.8	5.7	3.4-8.8	1.2	4.7	1.7-9.9	2.6	6.5	3.5-11.1
	1996-2000	5.2	7.2	4.7-10.5	2.6	8.4	4.3-14.5	2.6	6.1	3.3-10.6	2.8	3.9	2.1-6.5	1.6	5.5	2.3-10.9	1.2	2.7	1.0-6.0
	2001-2005	5.8	7.2	4.8-10.4	3.4	9.8	5.6-15.9	2.4	5.4	2.8-9.6	2.8	3.6	1.9-6.0	1.4	4.2	1.6-8.6	1.4	3.2	1.3-6.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Clinton**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	7.6	8.6	6.1-11.9	4.0	11.5	6.8-17.9	3.6	7.4	4.4-11.9	4.6	5.3	3.3-7.9	2.6	7.2	3.7-12.5	2.0	4.1	2.0-7.8
	2011-2015	9.2	9.1	6.7-12.3	5.8	11.7	7.8-17.1	3.4	6.8	3.9-11.1	4.0	4.1	2.5-6.5	2.6	6.2	3.2-10.8	1.4	2.8	1.1-6.0
	2016-2020	11.2	11.4	8.5-15.0	7.4	14.7	10.3-20.7	3.8	8.7	5.1-14.0	4.4	4.3	2.6-6.6	2.0	4.5	2.1-8.4	2.4	4.0	2.1-7.5
Leukemias	1976-1980	8.0	14.2	10.0-19.5	4.8	19.8	12.4-29.8	3.2	9.4	5.3-15.4	5.4	9.5	6.2-14.0	3.0	12.9	6.8-21.6	2.4	7.2	3.7-12.7
	1981-1985	8.2	13.2	9.3-18.0	5.0	20.0	12.1-30.5	3.2	9.2	5.2-15.0	4.4	8.0	5.0-12.0	3.2	16.8	9.1-27.6	1.2	3.6	1.3-7.7
	1986-1990	7.4	10.7	7.5-14.8	3.2	10.4	5.9-17.1	4.2	10.8	6.6-16.5	3.6	5.5	3.2-8.7	1.0	3.5	1.1-8.1	2.6	7.2	3.8-12.3
	1991-1995	10.0	13.3	9.8-17.6	5.6	16.6	10.8-24.2	4.4	10.7	6.7-16.3	4.8	6.8	4.3-10.1	1.8	5.7	2.5-10.9	3.0	6.9	3.9-11.5
	1996-2000	11.4	15.5	11.7-20.1	6.4	20.7	13.9-29.6	5.0	12.0	7.8-17.8	6.6	9.0	6.2-12.6	2.8	9.0	4.8-15.3	3.8	9.1	5.5-14.4
	2001-2005	10.2	13.2	9.8-17.4	5.6	16.0	10.5-23.2	4.6	10.5	6.6-16.0	3.8	4.8	2.9-7.5	2.0	5.7	2.7-10.5	1.8	4.2	1.9-8.0
	2006-2010	13.8	16.0	12.4-20.3	9.2	24.4	17.6-32.9	4.6	10.0	6.3-15.2	6.0	7.0	4.7-10.1	3.2	9.4	5.2-15.4	2.8	5.8	3.2-9.9
	2011-2015	17.8	18.6	14.9-23.1	11.2	25.8	19.3-33.8	6.6	12.4	8.5-17.8	6.0	6.2	4.2-9.0	3.6	8.4	4.9-13.5	2.4	4.5	2.3-8.2
2016-2020	16.0	16.4	12.9-20.7	11.2	24.6	18.4-32.3	4.8	8.7	5.5-13.4	6.2	6.3	4.2-9.0	4.2	9.7	6.0-14.9	2.0	3.8	1.8-7.4	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Columbia**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	251.2	381.1	359.8-403.4	133.6	461.1	425.5-499.0	117.6	323.5	297.1-351.8	135.8	203.8	188.4-220.2	72.0	247.6	221.6-275.9	63.8	172.5	153.5-193.3
	1981-1985	299.0	429.0	407.0-452.0	154.8	502.6	466.7-540.6	144.2	381.0	352.5-411.2	157.0	221.9	206.3-238.4	84.2	275.1	248.6-303.6	72.8	182.0	163.2-202.5
	1986-1990	309.6	425.3	404.0-447.4	157.0	489.8	455.5-526.1	152.6	387.1	359.2-416.7	155.2	209.8	195.1-225.4	81.0	255.4	230.6-282.1	74.2	178.6	160.4-198.4
	1991-1995	319.2	429.0	408.0-450.9	171.0	514.7	480.3-550.9	148.2	369.1	342.4-397.5	167.6	219.2	204.5-234.8	90.6	278.1	252.7-305.4	77.0	178.4	160.7-197.7
	1996-2000	391.0	507.5	485.1-530.7	202.8	578.3	542.9-615.4	188.2	455.4	426.2-486.2	164.2	207.6	193.6-222.4	80.0	233.4	210.9-257.8	84.2	192.8	174.5-212.6
	2001-2005	387.8	483.5	462.0-505.8	203.6	552.4	518.6-587.9	184.2	433.3	405.0-463.2	156.0	190.2	177.0-204.2	81.6	228.5	206.6-252.1	74.4	163.7	147.1-181.9
	2006-2010	430.4	504.4	482.8-526.9	237.0	580.4	547.1-615.4	193.4	442.0	413.6-472.0	152.0	173.5	161.2-186.6	78.8	198.8	179.3-220.0	73.2	154.9	139.1-172.3
	2011-2015	433.2	471.7	451.1-493.1	222.0	495.6	465.7-527.1	211.2	457.3	428.4-487.8	152.8	159.5	148.1-171.6	86.0	193.2	174.9-213.1	66.8	131.9	117.7-147.7
	2016-2020	469.2	481.8	461.1-503.2	242.0	499.7	470.4-530.6	227.2	469.4	440.1-500.4	142.8	138.2	127.9-149.3	75.6	156.3	140.4-173.8	67.2	124.0	110.5-139.2
Oral cavity and pharynx	1976-1980	7.6	11.4	8.0-15.8	3.8	13.1	7.7-20.8	3.8	9.9	5.9-15.9	3.2	4.9	2.7-8.1	1.8	5.6	2.5-10.9	1.4	4.0	1.5-8.7
	1981-1985	8.8	13.1	9.4-17.8	6.8	23.2	15.8-32.7	2.0	5.5	2.5-10.4	2.8	4.1	2.2-7.0	2.2	7.7	3.7-14.0	0.6	1.4	0.3-4.6
	1986-1990	4.8	6.5	4.1-9.7	3.0	9.6	5.3-16.0	1.8	4.3	1.9-8.5	2.0	2.5	1.2-4.8	1.2	3.8	1.3-8.5	0.8	1.7	0.5-4.9
	1991-1995	7.6	10.9	7.7-15.0	5.0	16.0	10.3-23.7	2.6	6.4	3.3-11.2	2.8	3.9	2.1-6.7	2.0	6.9	3.3-12.7	0.8	1.6	0.4-4.6
	1996-2000	9.8	13.0	9.6-17.3	5.6	16.2	10.7-23.5	4.2	10.0	6.1-15.7	3.0	3.9	2.2-6.6	1.8	5.4	2.4-10.3	1.2	2.8	1.0-6.5
	2001-2005	8.6	10.5	7.6-14.3	6.4	16.5	11.2-23.5	2.2	4.6	2.3-8.7	2.0	2.4	1.1-4.5	1.6	4.3	1.8-8.8	0.4	0.7	0.1-3.4
	2006-2010	11.0	12.5	9.4-16.6	8.2	19.2	13.6-26.5	2.8	5.8	3.2-10.3	2.0	2.4	1.1-4.6	1.2	3.0	1.0-7.1	0.8	1.6	0.4-4.7
	2011-2015	8.8	8.8	6.3-12.1	6.6	14.3	9.8-20.6	2.2	4.1	2.0-8.1	1.0	1.0	0.3-2.6	1.0	2.1	0.7-5.5	0.0	0.0	0.0-2.7
	2016-2020	14.4	14.7	11.3-18.9	9.8	21.0	15.2-28.5	4.6	8.7	5.3-13.9	2.6	2.9	1.5-5.3	1.8	4.3	1.9-8.8	0.8	1.6	0.4-5.0
Esophagus	1976-1980	0.8	1.1	0.3-3.1	0.4	1.5	0.2-5.3	0.4	0.9	0.1-4.0	1.0	1.3	0.4-3.3	0.6	1.8	0.4-5.7	0.4	0.9	0.1-4.0
	1981-1985	2.2	3.0	1.5-5.5	1.8	5.6	2.5-10.9	0.4	0.9	0.1-3.8	3.8	4.8	2.9-7.7	3.4	9.8	5.7-16.0	0.4	0.9	0.1-3.8
	1986-1990	3.2	4.4	2.5-7.2	2.8	8.6	4.6-14.6	0.4	1.0	0.1-4.0	2.8	3.9	2.1-6.6	2.0	6.4	3.0-11.9	0.8	1.9	0.5-5.2
	1991-1995	2.8	3.8	2.1-6.4	2.6	7.5	4.0-13.0	0.2	0.4	0.0-2.8	2.2	3.1	1.5-5.6	2.0	6.0	2.8-11.1	0.2	0.6	0.0-3.5
	1996-2000	5.4	6.9	4.5-10.1	4.4	12.2	7.6-18.7	1.0	2.0	0.6-5.2	3.8	4.8	2.9-7.5	3.2	9.5	5.4-15.6	0.6	1.2	0.2-4.1
	2001-2005	4.6	5.5	3.5-8.4	4.0	10.2	6.2-16.1	0.6	1.2	0.2-4.1	4.8	5.8	3.7-8.7	3.8	10.0	6.0-15.9	1.0	1.9	0.6-5.1
	2006-2010	5.2	6.0	3.9-9.0	4.6	11.0	6.8-16.9	0.6	1.3	0.3-4.5	4.6	5.5	3.4-8.5	4.0	10.0	6.0-15.9	0.6	1.3	0.3-4.5
	2011-2015	4.6	4.4	2.8-6.9	3.8	7.9	4.7-12.9	0.8	1.3	0.4-4.3	3.4	3.5	2.0-5.8	3.2	6.9	3.9-11.8	0.2	0.5	0.0-3.4
	2016-2020	4.6	4.5	2.7-7.2	3.6	7.5	4.3-12.7	1.0	1.7	0.5-5.0	4.6	4.5	2.8-7.2	3.6	7.5	4.2-12.7	1.0	1.8	0.6-5.1
Stomach	1976-1980	7.4	11.4	8.0-15.8	4.4	14.9	9.2-22.8	3.0	8.0	4.4-13.5	4.6	6.9	4.4-10.5	2.4	8.0	4.0-14.4	2.2	6.0	3.0-11.1
	1981-1985	6.6	9.2	6.3-13.1	3.8	12.0	7.1-19.1	2.8	7.0	3.7-12.1	4.4	6.3	3.9-9.7	2.8	9.2	4.9-15.7	1.6	4.0	1.6-8.3
	1986-1990	5.8	7.7	5.1-11.1	2.6	8.2	4.3-14.2	3.2	7.0	4.0-11.7	4.8	6.2	3.9-9.3	2.6	7.8	4.1-13.5	2.2	4.8	2.4-8.9
	1991-1995	4.0	5.2	3.2-8.2	2.8	8.0	4.4-13.6	1.2	2.6	0.9-6.0	4.4	5.9	3.7-9.0	2.4	7.0	3.6-12.4	2.0	4.4	2.1-8.4
	1996-2000	4.8	6.0	3.9-9.1	3.2	9.4	5.3-15.4	1.6	3.6	1.6-7.5	2.2	2.8	1.4-5.1	1.4	4.1	1.6-8.5	0.8	1.8	0.5-5.0
	2001-2005	4.0	4.9	3.0-7.7	2.4	6.6	3.4-11.7	1.6	3.3	1.4-7.0	3.4	4.0	2.3-6.5	1.4	3.6	1.5-7.8	2.0	3.9	1.8-7.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Columbia**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	5.8	6.9	4.6-10.1	4.4	10.9	6.8-16.8	1.4	3.6	1.4-7.7	3.8	4.5	2.7-7.2	2.8	7.0	3.8-12.0	1.0	2.4	0.7-6.1
	2011-2015	4.8	5.4	3.4-8.3	4.0	9.5	5.7-15.3	0.8	1.5	0.4-4.7	2.4	2.5	1.3-4.7	2.2	5.2	2.6-9.7	0.2	0.4	0.0-3.3
	2016-2020	6.2	6.1	4.1-8.9	3.8	7.6	4.5-12.5	2.4	5.0	2.5-9.5	2.2	2.3	1.1-4.5	1.8	3.8	1.7-7.9	0.4	1.2	0.1-4.8
Colorectal	1976-1980	41.8	62.5	54.2-71.8	22.0	76.5	62.3-92.8	19.8	51.9	42.0-63.6	21.2	31.2	25.5-37.9	10.8	37.3	27.7-49.1	10.4	26.5	19.7-35.0
	1981-1985	50.2	69.4	61.0-78.8	24.2	76.4	63.0-91.9	26.0	62.7	52.2-74.9	25.4	35.8	29.7-42.8	13.4	44.8	34.3-57.4	12.0	28.2	21.4-36.6
	1986-1990	56.4	76.4	67.6-86.0	29.0	90.1	75.8-106.4	27.4	66.8	55.8-79.5	25.2	34.6	28.7-41.3	13.0	41.8	32.1-53.6	12.2	29.6	22.4-38.4
	1991-1995	40.4	52.1	45.1-59.9	19.4	60.3	48.7-73.8	21.0	46.4	37.8-56.5	20.8	26.3	21.5-32.0	10.2	31.7	23.5-41.9	10.6	22.3	16.7-29.4
	1996-2000	47.8	60.2	52.8-68.5	22.4	63.7	52.3-76.8	25.4	56.4	46.9-67.6	22.6	27.8	22.9-33.5	11.0	31.7	23.8-41.4	11.6	23.9	18.1-31.4
	2001-2005	43.0	52.1	45.3-59.7	18.2	49.2	39.5-60.7	24.8	53.9	44.6-64.8	14.2	16.9	13.2-21.4	5.8	17.2	11.5-24.9	8.4	17.3	12.3-23.9
	2006-2010	43.2	50.4	43.8-57.9	21.0	52.3	42.5-63.7	22.2	49.1	40.1-59.7	13.8	15.7	12.2-20.0	6.2	15.4	10.4-22.2	7.6	16.4	11.5-23.0
	2011-2015	39.8	42.1	36.3-48.7	20.4	45.6	37.0-55.9	19.4	39.5	31.8-49.0	15.8	16.8	13.2-21.2	8.4	19.3	13.8-26.6	7.4	14.4	10.0-20.6
	2016-2020	35.0	35.5	30.2-41.6	17.4	37.4	29.7-46.9	17.6	33.0	26.2-41.6	13.0	13.5	10.2-17.8	6.4	13.1	8.8-19.3	6.6	13.7	9.0-20.4
Colon excluding rectum	1976-1980	30.4	45.5	38.4-53.5	14.2	49.2	38.1-62.6	16.2	42.5	33.6-53.2	17.2	25.1	20.0-31.1	8.6	29.0	20.8-39.4	8.6	21.6	15.6-29.4
	1981-1985	33.4	46.2	39.3-53.9	15.0	48.0	37.5-60.7	18.4	43.8	35.2-54.1	20.6	28.7	23.4-35.0	10.0	33.7	24.7-45.0	10.6	24.5	18.3-32.4
	1986-1990	39.4	53.4	46.1-61.6	19.2	61.0	49.2-74.7	20.2	48.6	39.3-59.5	22.8	31.3	25.7-37.7	11.8	38.6	29.2-50.0	11.0	26.2	19.6-34.5
	1991-1995	29.8	38.3	32.4-45.1	14.4	45.3	35.3-57.2	15.4	33.7	26.5-42.4	18.0	22.8	18.3-28.2	9.4	29.3	21.4-39.2	8.6	18.3	13.2-25.0
	1996-2000	35.6	44.7	38.3-51.8	15.0	43.1	33.8-54.3	20.6	46.1	37.5-56.3	19.4	23.8	19.3-29.2	9.8	28.2	20.8-37.5	9.6	19.8	14.5-26.6
	2001-2005	31.0	37.2	31.5-43.7	11.4	31.3	23.7-40.8	19.6	41.4	33.4-50.9	12.2	14.5	11.0-18.7	4.8	14.5	9.2-21.8	7.4	15.4	10.7-21.7
	2006-2010	29.4	34.5	29.0-40.7	14.0	36.1	27.9-46.0	15.4	33.2	26.0-42.0	10.4	12.0	9.0-16.0	4.6	11.8	7.4-18.0	5.8	12.5	8.2-18.4
	2011-2015	27.6	29.2	24.4-34.8	13.8	31.7	24.5-40.6	13.8	27.7	21.3-35.7	12.8	13.7	10.5-17.7	6.8	15.6	10.7-22.3	6.0	11.9	7.9-17.6
	2016-2020	23.2	23.5	19.3-28.6	10.8	23.3	17.3-31.1	12.4	23.3	17.6-30.7	10.8	11.5	8.4-15.5	5.0	10.3	6.5-16.0	5.8	12.3	7.8-18.8
Rectum & rectosigmoid	1976-1980	11.4	17.0	12.8-22.2	7.8	27.2	19.1-37.7	3.6	9.4	5.5-15.2	4.0	6.2	3.7-9.7	2.2	8.3	4.0-15.1	1.8	4.9	2.2-9.6
	1981-1985	16.8	23.3	18.5-29.0	9.2	28.4	20.6-38.2	7.6	18.9	13.2-26.3	4.8	7.1	4.5-10.7	3.4	11.1	6.3-18.1	1.4	3.6	1.4-7.8
	1986-1990	17.0	22.9	18.2-28.5	9.8	29.1	21.4-38.8	7.2	18.3	12.6-25.7	2.4	3.3	1.7-5.8	1.2	3.2	1.2-7.4	1.2	3.4	1.2-7.5
	1991-1995	10.6	13.7	10.3-18.1	5.0	15.0	9.7-22.3	5.6	12.7	8.4-18.7	2.8	3.5	1.9-6.0	0.8	2.4	0.6-6.2	2.0	3.9	1.9-7.7
	1996-2000	12.2	15.6	11.9-20.1	7.4	20.5	14.4-28.5	4.8	10.4	6.6-15.8	3.2	4.0	2.3-6.5	1.2	3.4	1.3-7.6	2.0	4.2	2.0-8.1
	2001-2005	12.0	14.9	11.3-19.3	6.8	17.9	12.3-25.3	5.2	12.6	8.1-18.9	2.0	2.4	1.1-4.6	1.0	2.7	0.9-6.6	1.0	1.9	0.6-5.1
	2006-2010	13.8	16.0	12.3-20.4	7.0	16.2	11.2-22.9	6.8	15.9	10.8-22.8	3.4	3.7	2.1-6.1	1.6	3.6	1.5-7.6	1.8	3.9	1.8-7.9
	2011-2015	12.2	12.9	9.7-16.8	6.6	13.9	9.5-20.1	5.6	11.9	7.7-17.9	3.0	3.1	1.7-5.4	1.6	3.7	1.6-7.9	1.4	2.6	1.0-6.1
	2016-2020	11.8	12.0	9.0-15.8	6.6	14.1	9.6-20.5	5.2	9.7	6.3-15.1	2.2	2.1	1.0-4.1	1.4	2.8	1.0-6.7	0.8	1.4	0.4-4.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Columbia**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	1.8	2.7	1.2-5.2	0.8	2.1	0.6-5.9	1.0	3.0	0.9-7.2	1.4	2.4	0.9-5.0	0.8	3.1	0.8-8.2	0.6	1.9	0.4-5.7
	1981-1985	1.2	2.2	0.8-4.7	0.8	3.0	0.8-7.7	0.4	1.3	0.1-4.9	0.8	1.3	0.3-3.4	0.6	2.0	0.4-5.9	0.2	0.9	0.0-4.4
	1986-1990	1.8	2.3	1.0-4.4	1.2	3.7	1.3-8.2	0.6	1.3	0.3-4.2	2.2	3.0	1.5-5.5	1.8	6.0	2.7-11.5	0.4	0.9	0.1-3.7
	1991-1995	4.6	6.0	3.8-9.1	3.0	8.9	4.9-14.9	1.6	4.0	1.7-8.1	2.2	2.8	1.4-5.1	1.8	5.4	2.4-10.4	0.4	1.1	0.1-4.1
	1996-2000	4.6	5.8	3.7-8.8	2.0	5.5	2.6-10.3	2.6	6.1	3.2-10.7	3.0	3.7	2.1-6.3	1.6	4.3	1.9-8.8	1.4	3.0	1.2-6.6
	2001-2005	2.4	2.8	1.5-5.1	1.8	4.8	2.2-9.3	0.6	1.3	0.3-4.4	2.6	3.2	1.7-5.6	2.0	5.5	2.6-10.4	0.6	1.3	0.3-4.4
	2006-2010	4.6	5.6	3.5-8.6	3.0	7.3	4.0-12.5	1.6	4.0	1.7-8.5	4.4	5.2	3.2-8.0	2.6	6.1	3.2-10.8	1.8	4.4	1.9-8.9
	2011-2015	6.4	6.2	4.3-9.1	5.0	10.2	6.5-15.5	1.4	2.6	1.0-6.2	4.8	4.5	2.9-6.9	3.2	6.2	3.5-10.7	1.6	2.9	1.2-6.5
2016-2020	8.6	8.4	5.9-11.8	6.6	13.9	9.3-20.4	2.0	3.2	1.5-6.8	5.6	5.1	3.4-7.7	3.6	6.8	4.0-11.5	2.0	3.7	1.7-7.7	
Pancreas	1976-1980	7.2	11.5	8.0-16.0	3.8	15.0	8.7-23.8	3.4	9.2	5.3-15.0	7.0	10.7	7.4-15.0	4.4	16.1	9.8-25.0	2.6	6.7	3.6-11.8
	1981-1985	11.4	15.9	11.9-20.8	6.2	20.5	13.8-29.5	5.2	11.7	7.6-17.4	10.8	15.3	11.4-20.1	6.0	20.2	13.5-29.2	4.8	10.8	6.9-16.4
	1986-1990	7.2	9.5	6.6-13.3	3.2	10.6	6.0-17.3	4.0	9.2	5.5-14.5	7.0	9.4	6.5-13.1	3.6	12.5	7.3-19.9	3.4	7.7	4.5-12.7
	1991-1995	9.2	12.0	8.8-16.1	4.4	13.5	8.4-20.6	4.8	9.9	6.3-15.0	10.0	13.0	9.7-17.3	4.8	15.1	9.6-22.6	5.2	11.0	7.1-16.4
	1996-2000	9.4	12.0	8.8-16.0	3.4	9.8	5.7-15.9	6.0	13.6	9.1-19.7	10.6	13.4	10.0-17.6	3.8	11.0	6.6-17.3	6.8	15.3	10.5-21.8
	2001-2005	12.0	14.5	11.1-18.9	5.4	14.6	9.6-21.5	6.6	14.1	9.6-20.3	9.4	11.4	8.3-15.3	3.8	10.5	6.3-16.6	5.6	11.9	7.9-17.7
	2006-2010	11.8	13.5	10.2-17.5	6.0	14.5	9.7-21.1	5.8	12.3	8.2-18.1	9.8	11.2	8.2-14.9	5.0	12.7	8.2-19.1	4.8	10.1	6.4-15.5
	2011-2015	12.2	12.9	9.8-16.8	6.6	15.1	10.3-21.7	5.6	10.6	7.0-16.0	9.8	10.1	7.4-13.6	6.0	13.3	8.9-19.4	3.8	6.9	4.1-11.4
2016-2020	13.8	13.9	10.6-18.1	7.2	14.3	9.8-20.6	6.6	13.5	8.9-19.9	11.0	10.7	8.0-14.3	5.2	10.4	6.6-16.0	5.8	10.7	7.1-16.2	
Larynx	1976-1980	3.2	4.9	2.7-8.1	2.8	8.9	4.8-15.4	0.4	1.5	0.2-5.4	0.8	1.2	0.3-3.1	0.4	1.1	0.1-4.6	0.4	1.1	0.1-4.2
	1981-1985	2.8	3.9	2.1-6.7	2.4	8.4	4.2-14.9	0.4	0.9	0.1-3.7	0.8	1.4	0.4-3.6	0.8	3.1	0.8-7.8	0.0	0.0	0.0-2.4
	1986-1990	3.4	4.8	2.7-7.7	2.6	8.0	4.2-13.9	0.8	2.0	0.5-5.3	0.8	1.1	0.3-3.0	0.6	1.9	0.4-5.6	0.2	0.5	0.0-3.1
	1991-1995	3.8	5.0	3.0-7.9	2.8	8.3	4.5-14.1	1.0	2.4	0.8-5.9	1.6	2.1	0.9-4.2	1.6	4.5	1.9-9.0	0.0	0.0	0.0-2.3
	1996-2000	3.6	4.7	2.8-7.6	2.8	8.0	4.4-13.6	0.8	2.0	0.5-5.5	1.2	1.5	0.5-3.4	0.8	2.1	0.6-5.7	0.4	0.9	0.1-3.8
	2001-2005	2.8	3.4	1.9-5.9	2.2	5.8	2.9-10.7	0.6	1.5	0.3-4.7	0.2	0.3	0.0-1.7	0.2	0.5	0.0-3.4	0.0	0.0	0.0-2.5
	2006-2010	3.4	3.9	2.3-6.4	3.0	7.2	4.0-12.2	0.4	1.0	0.1-4.1	0.4	0.4	0.0-1.8	0.2	0.6	0.0-3.5	0.2	0.3	0.0-3.0
	2011-2015	3.2	3.1	1.7-5.3	2.8	5.4	2.9-9.7	0.4	0.8	0.1-3.8	0.6	0.5	0.1-1.9	0.6	1.0	0.2-3.9	0.0	0.0	0.0-2.7
2016-2020	3.8	4.3	2.5-7.2	3.0	6.4	3.5-11.3	0.8	2.5	0.6-6.9	1.2	1.0	0.4-2.7	1.0	1.8	0.6-5.1	0.2	0.3	0.0-3.3	
Lung and bronchus	1976-1980	41.8	61.3	53.0-70.6	31.6	103.5	87.5-121.6	10.2	27.4	20.2-36.5	32.2	45.6	38.7-53.5	23.0	71.3	58.5-86.1	9.2	24.7	17.8-33.5
	1981-1985	47.0	64.4	56.2-73.5	31.0	94.8	80.1-111.6	16.0	40.9	32.1-51.4	39.2	53.9	46.5-62.3	26.4	83.0	69.0-98.9	12.8	32.2	24.6-41.5
	1986-1990	52.4	70.0	61.7-79.3	29.8	89.3	75.3-105.2	22.6	55.6	45.5-67.4	44.2	59.0	51.4-67.5	26.8	82.3	68.7-97.9	17.4	42.1	33.5-52.4
	1991-1995	55.8	73.9	65.4-83.2	34.2	102.0	87.1-118.7	21.6	51.4	42.0-62.4	50.6	66.4	58.4-75.3	29.8	88.4	74.6-104.0	20.8	48.8	39.7-59.5
	1996-2000	57.0	72.4	64.2-81.4	30.6	87.1	73.8-102.3	26.4	63.0	52.6-75.0	40.0	50.5	43.8-58.1	22.8	64.7	53.3-77.9	17.2	40.7	32.5-50.6
	2001-2005	65.8	79.3	70.9-88.4	36.8	99.7	85.6-115.5	29.0	64.5	54.3-76.2	49.2	59.6	52.4-67.7	29.0	79.7	67.2-94.1	20.2	45.0	36.6-55.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Columbia**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	68.6	78.9	70.6-87.9	38.8	96.2	82.9-111.1	29.8	64.1	54.1-75.7	47.4	54.5	47.7-62.2	26.6	65.8	54.9-78.4	20.8	45.4	37.0-55.4
	2011-2015	77.6	80.1	72.2-88.8	39.6	86.3	74.3-99.8	38.0	76.0	65.4-88.1	47.6	48.8	42.7-55.7	27.4	60.2	50.4-71.7	20.2	39.5	32.0-48.5
	2016-2020	68.6	66.7	59.5-74.6	32.2	64.4	54.6-75.9	36.4	69.4	59.1-81.3	35.6	34.7	29.6-40.6	18.2	37.9	30.2-47.2	17.4	31.8	25.3-40.0
Melanoma of the skin	1976-1980	2.6	4.3	2.2-7.5	1.6	5.6	2.3-11.2	1.0	3.3	1.0-7.8	1.2	1.9	0.7-4.4	1.0	3.6	1.1-8.7	0.2	0.5	0.0-3.2
	1981-1985	3.8	5.9	3.4-9.3	2.4	8.2	4.1-14.6	1.4	3.6	1.4-7.9	2.0	2.9	1.4-5.5	0.8	2.2	0.6-6.0	1.2	3.6	1.2-8.1
	1986-1990	4.6	6.7	4.2-10.1	3.8	11.6	6.9-18.3	0.8	2.8	0.7-7.0	2.6	3.8	2.0-6.5	2.2	7.4	3.6-13.4	0.4	1.1	0.1-4.2
	1991-1995	5.0	7.1	4.5-10.5	3.0	9.0	5.0-15.0	2.0	5.7	2.7-10.7	1.2	1.5	0.6-3.5	1.0	3.0	0.9-7.1	0.2	0.7	0.0-3.6
	1996-2000	9.4	12.5	9.2-16.8	4.8	14.1	9.0-21.1	4.6	11.4	7.2-17.4	2.2	2.8	1.4-5.1	1.6	4.2	1.8-8.5	0.6	1.4	0.3-4.5
	2001-2005	8.6	11.8	8.5-16.1	6.0	17.9	12.0-25.8	2.6	7.2	3.7-12.7	3.0	4.0	2.2-6.8	1.8	5.4	2.4-10.4	1.2	2.7	1.0-6.3
	2006-2010	16.8	21.0	16.6-26.3	9.6	23.9	17.5-32.0	7.2	18.8	12.9-26.6	3.0	3.6	2.0-6.1	2.6	6.6	3.5-11.6	0.4	0.8	0.1-3.7
	2011-2015	17.2	20.6	16.2-26.0	11.0	26.1	19.2-34.8	6.2	16.2	10.6-23.8	2.4	3.1	1.5-5.7	1.8	5.3	2.3-10.5	0.6	1.1	0.2-4.2
2016-2020	17.8	19.0	15.0-23.9	9.8	20.8	15.0-28.3	8.0	17.6	12.1-25.0	3.6	3.5	2.1-5.9	2.4	5.1	2.6-9.5	1.2	2.2	0.8-5.8	
Female breast	1976-1980							34.2	95.6	81.3-111.7							14.0	38.7	29.9-49.4
	1981-1985							35.6	103.2	87.9-120.4							14.2	38.0	29.3-48.5
	1986-1990							39.0	104.9	90.2-121.3							10.2	26.8	19.7-35.6
	1991-1995							39.4	103.2	88.9-119.1							10.4	26.4	19.6-35.0
	1996-2000							50.0	124.7	109.5-141.6							14.8	34.5	27.0-43.7
	2001-2005							50.8	121.6	106.9-138.0							8.2	17.4	12.4-24.1
	2006-2010							47.8	111.6	97.5-127.4							6.6	14.3	9.8-20.5
	2011-2015							56.6	124.7	109.6-141.4							7.2	15.4	10.5-22.0
2016-2020							65.8	140.2	124.2-158.0							8.0	15.3	10.7-21.7	
Cervix uteri	1976-1980							4.4	14.2	8.8-21.8							2.6	8.6	4.4-15.0
	1981-1985							5.0	16.4	10.4-24.5							2.6	8.1	4.2-14.2
	1986-1990							4.0	10.8	6.5-16.9							0.8	2.1	0.5-5.5
	1991-1995							6.8	20.8	14.4-29.2							1.2	3.4	1.2-7.6
	1996-2000							3.6	11.2	6.5-17.9							1.6	4.2	1.8-8.5
	2001-2005							3.8	10.9	6.4-17.4							0.8	2.2	0.6-5.9
	2006-2010							3.4	9.2	5.2-15.3							0.8	2.0	0.6-5.6
	2011-2015							3.2	9.3	5.0-15.8							0.8	2.2	0.5-6.1
	2016-2020							2.6	7.7	3.8-14.0							1.6	3.4	1.3-7.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Columbia**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							6.2	17.3	11.6-24.9							1.8	4.5	2.0-8.9
	1981-1985							8.8	24.0	17.1-32.7							1.6	3.5	1.5-7.4
	1986-1990							8.6	21.6	15.5-29.5							2.2	5.2	2.5-9.6
	1991-1995							9.2	22.7	16.4-30.6							2.0	4.7	2.2-9.0
	1996-2000							11.6	27.3	20.7-35.6							2.4	5.0	2.6-9.2
	2001-2005							10.4	24.6	18.3-32.5							1.6	3.3	1.4-7.0
	2006-2010							13.4	28.8	22.2-37.1							2.4	4.6	2.3-8.7
	2011-2015							15.4	30.8	23.9-39.3							2.4	4.2	2.1-8.1
	2016-2020							15.8	29.7	23.1-38.1							3.0	4.9	2.7-8.9
Ovary	1976-1980							6.0	16.2	10.8-23.5							4.6	12.5	7.8-19.2
	1981-1985							8.2	22.6	15.9-31.1							4.4	10.7	6.6-16.5
	1986-1990							8.0	20.6	14.5-28.4							5.6	13.5	8.8-19.8
	1991-1995							7.2	19.2	13.3-26.9							5.4	12.7	8.3-18.8
	1996-2000							6.8	16.9	11.6-24.0							3.8	9.2	5.5-14.7
	2001-2005							8.4	19.6	14.0-26.8							3.8	8.7	5.2-14.0
	2006-2010							5.4	13.1	8.5-19.4							4.2	8.8	5.4-13.9
	2011-2015							3.0	7.8	4.1-13.5							2.8	5.6	3.0-10.1
	2016-2020							4.4	10.0	6.0-16.0							2.2	3.8	1.9-7.7
Prostate	1976-1980				21.2	73.7	59.9-89.8							7.0	28.1	19.1-39.6			
	1981-1985				29.2	101.0	84.7-119.5							7.0	26.1	18.0-36.5			
	1986-1990				36.6	114.6	98.3-132.9							8.4	28.3	20.2-38.4			
	1991-1995				47.2	139.7	122.2-159.0							12.2	40.5	30.9-52.3			
	1996-2000				63.4	176.1	157.2-196.9							9.0	27.6	20.1-37.1			
	2001-2005				59.8	157.2	139.8-176.4							9.4	27.7	20.3-37.0			
	2006-2010				63.4	145.8	129.9-163.3							8.2	22.3	15.9-30.4			
	2011-2015				48.2	98.2	85.9-112.0							7.6	17.5	12.3-24.3			
	2016-2020				62.8	116.4	103.7-130.7							5.4	11.6	7.6-17.4			
Testis	1976-1980				1.2	4.6	1.7-10.2							0.6	2.3	0.4-7.0			
	1981-1985				1.8	6.0	2.7-11.5							0.2	0.9	0.0-4.7			
	1986-1990				1.0	3.1	1.0-7.3							0.2	0.7	0.0-3.8			
	1991-1995				1.8	6.1	2.8-11.6							0.0	0.0	0.0-2.4			
	1996-2000				3.0	9.8	5.5-16.2							0.4	1.3	0.2-4.8			
	2001-2005				1.8	6.1	2.8-11.7							0.2	0.7	0.0-3.9			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Columbia**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				3.0	9.4	5.2-15.8							0.0	0.0	0.0-2.5			
	2011-2015				1.8	7.2	3.3-13.7							0.2	0.9	0.0-4.6			
	2016-2020				2.2	7.9	3.7-14.5							0.0	0.0	0.0-2.8			
Urinary bladder (incl. in situ)	1976-1980	16.8	25.7	20.3-32.0	12.8	47.3	35.9-61.0	4.0	10.4	6.3-16.3	4.0	6.1	3.7-9.5	2.8	11.3	5.9-19.3	1.2	2.9	1.1-6.8
	1981-1985	19.8	27.6	22.3-33.8	15.6	50.0	39.2-62.8	4.2	9.9	6.1-15.5	5.4	7.6	4.9-11.1	4.0	13.5	8.1-21.1	1.4	3.3	1.3-7.1
	1986-1990	16.0	21.9	17.2-27.4	12.0	38.2	29.0-49.5	4.0	9.3	5.6-14.8	4.4	5.9	3.6-9.0	3.0	9.6	5.2-16.0	1.4	3.0	1.2-6.5
	1991-1995	15.2	19.6	15.4-24.6	11.0	33.4	25.1-43.7	4.2	10.3	6.3-16.0	4.2	5.4	3.3-8.3	2.4	7.2	3.7-12.7	1.8	4.0	1.8-7.9
	1996-2000	22.8	29.2	24.1-35.2	15.4	44.6	35.1-55.9	7.4	17.2	12.1-24.0	4.6	5.9	3.7-9.0	3.6	10.7	6.3-17.0	1.0	2.6	0.8-6.4
	2001-2005	20.4	25.0	20.3-30.4	15.0	41.4	32.5-52.1	5.4	11.8	7.7-17.5	3.8	4.6	2.8-7.3	2.8	8.5	4.6-14.4	1.0	2.3	0.7-5.8
	2006-2010	26.0	29.2	24.3-34.9	19.4	48.1	38.8-59.1	6.6	13.6	9.3-19.6	4.6	5.2	3.3-8.1	2.8	7.6	4.1-13.0	1.8	3.5	1.6-7.2
	2011-2015	24.2	25.1	20.8-30.3	18.0	40.2	32.1-49.9	6.2	12.0	8.1-17.5	2.0	2.1	1.0-4.2	1.2	2.7	1.0-6.4	0.8	1.7	0.4-5.2
	2016-2020	26.2	25.3	21.0-30.4	20.2	40.9	33.1-50.4	6.0	11.5	7.6-17.3	4.2	3.9	2.4-6.3	2.6	5.4	2.9-9.9	1.6	2.7	1.2-6.3
Kidney and renal pelvis	1976-1980	5.2	8.0	5.2-12.0	4.0	13.4	8.1-21.1	1.2	3.3	1.1-7.5	3.6	5.2	3.0-8.3	3.0	9.4	5.2-15.9	0.6	1.7	0.3-5.3
	1981-1985	6.8	10.2	7.0-14.5	4.0	13.7	8.3-21.4	2.8	6.6	3.6-11.5	3.2	4.9	2.8-8.1	2.4	8.0	4.0-14.2	0.8	2.1	0.5-5.7
	1986-1990	8.0	11.1	7.8-15.2	5.0	15.8	10.2-23.6	3.0	7.4	4.0-12.5	3.4	4.3	2.5-7.0	2.0	6.0	2.9-11.2	1.4	3.1	1.2-6.7
	1991-1995	6.8	9.2	6.4-13.0	4.0	12.2	7.4-19.0	2.8	6.9	3.7-11.8	4.0	5.1	3.1-8.0	2.8	8.3	4.5-14.1	1.2	2.5	0.9-5.9
	1996-2000	11.0	14.4	10.8-18.9	7.0	19.7	13.7-27.6	4.0	10.3	6.2-16.3	4.2	5.3	3.3-8.2	2.8	8.0	4.3-13.6	1.4	3.1	1.2-6.7
	2001-2005	12.2	15.3	11.7-19.7	7.2	19.2	13.4-26.8	5.0	11.9	7.7-18.0	3.0	3.7	2.1-6.2	1.2	3.5	1.3-7.7	1.8	4.1	1.9-8.2
	2006-2010	11.6	13.2	9.9-17.3	7.4	17.7	12.4-24.8	4.2	9.0	5.4-14.4	3.2	3.5	2.0-5.9	2.2	5.3	2.6-9.8	1.0	1.9	0.6-5.2
	2011-2015	16.4	18.7	14.6-23.6	8.2	18.9	13.4-26.3	8.2	18.1	12.6-25.5	5.2	5.4	3.5-8.1	2.6	5.9	3.1-10.7	2.6	4.9	2.6-9.0
	2016-2020	18.2	18.8	14.9-23.6	11.2	24.0	17.8-32.0	7.0	14.0	9.4-20.5	3.8	3.7	2.2-6.1	1.6	3.5	1.5-7.7	2.2	3.7	1.8-7.4
Brain and other central nervous system	1976-1980	4.0	6.8	4.1-10.7	2.6	9.8	5.1-17.1	1.4	3.7	1.5-8.0	2.2	3.6	1.7-6.5	1.4	5.5	2.1-11.4	0.8	2.0	0.5-5.5
	1981-1985	3.4	5.2	3.0-8.4	2.4	8.6	4.3-15.2	1.0	2.6	0.8-6.3	3.0	4.1	2.3-6.9	1.6	4.9	2.1-10.0	1.4	3.3	1.3-7.2
	1986-1990	4.8	7.3	4.6-10.9	2.4	7.6	3.9-13.5	2.4	6.9	3.4-12.2	4.4	6.0	3.7-9.1	2.2	6.7	3.3-12.2	2.2	5.5	2.7-10.1
	1991-1995	3.8	5.5	3.3-8.6	2.6	7.7	4.1-13.3	1.2	3.5	1.3-7.8	1.6	2.1	0.9-4.2	1.0	2.7	0.9-6.5	0.6	1.6	0.3-4.9
	1996-2000	6.4	9.4	6.4-13.3	3.0	9.0	5.0-15.0	3.4	10.0	5.7-16.2	3.4	4.6	2.7-7.4	2.2	6.8	3.4-12.2	1.2	3.1	1.1-7.0
	2001-2005	4.6	6.5	4.1-9.9	2.0	5.9	2.8-11.1	2.6	7.2	3.7-12.7	3.2	4.6	2.6-7.6	1.0	2.7	0.9-6.6	2.2	6.9	3.3-12.5
	2006-2010	6.2	8.2	5.5-11.9	3.4	9.6	5.5-15.6	2.8	6.9	3.7-12.1	3.0	3.4	1.9-5.9	1.2	2.9	1.0-6.7	1.8	3.9	1.8-8.0
	2011-2015	4.8	6.4	3.9-9.9	2.0	5.4	2.5-10.5	2.8	7.3	3.7-13.0	3.2	3.3	1.9-5.7	1.2	2.6	0.9-6.3	2.0	3.8	1.8-7.8
	2016-2020	6.4	7.7	5.0-11.5	3.8	8.9	5.0-14.8	2.6	6.9	3.2-12.9	3.2	3.0	1.7-5.3	2.0	3.6	1.7-7.3	1.2	2.7	0.9-6.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Columbia**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	1.0	1.5	0.5-3.5	0.2	0.5	0.0-3.6	0.8	2.3	0.6-6.1	0.2	0.2	0.0-1.7	0.2	0.5	0.0-3.6	0.0	0.0	0.0-2.5
	1981-1985	2.2	3.4	1.7-6.1	1.0	3.3	1.0-7.8	1.2	3.3	1.2-7.5	1.0	1.4	0.5-3.4	0.2	0.7	0.0-3.9	0.8	1.8	0.5-5.1
	1986-1990	1.4	2.1	0.9-4.5	0.6	2.1	0.4-6.2	0.8	2.1	0.5-5.6	0.2	0.3	0.0-1.7	0.0	0.0	0.0-2.5	0.2	0.4	0.0-2.9
	1991-1995	3.0	4.6	2.6-7.6	1.2	3.5	1.3-7.7	1.8	5.8	2.6-10.9	0.4	0.5	0.1-2.0	0.0	0.0	0.0-2.4	0.4	0.9	0.1-3.7
	1996-2000	4.0	6.3	3.8-9.8	1.6	5.2	2.2-10.3	2.4	7.2	3.6-12.8	0.0	0.0	0.0-1.2	0.0	0.0	0.0-2.4	0.0	0.0	0.0-2.4
	2001-2005	6.0	9.1	6.1-13.2	2.6	7.8	4.1-13.5	3.4	10.5	6.0-17.2	0.4	0.5	0.1-2.0	0.2	0.5	0.0-3.3	0.2	0.5	0.0-3.2
	2006-2010	6.6	8.9	6.0-12.8	1.0	2.4	0.8-6.0	5.6	15.6	10.2-23.1	0.4	0.4	0.0-1.8	0.0	0.0	0.0-2.5	0.4	0.7	0.1-3.5
	2011-2015	11.6	16.4	12.1-21.6	3.0	7.3	3.9-12.6	8.6	25.6	18.1-35.3	0.2	0.2	0.0-1.5	0.2	0.3	0.0-3.0	0.0	0.0	0.0-2.7
	2016-2020	15.0	21.3	16.4-27.2	4.6	10.0	6.2-15.7	10.4	33.1	24.1-44.4	0.4	0.3	0.0-1.8	0.0	0.0	0.0-2.8	0.4	0.6	0.1-3.6
Hodgkin lymphoma	1976-1980	2.0	3.3	1.6-6.1	1.2	4.0	1.5-9.0	0.8	2.5	0.7-6.5	0.4	0.8	0.1-2.8	0.2	0.7	0.0-4.1	0.2	0.9	0.0-4.6
	1981-1985	1.4	2.6	1.0-5.2	1.0	3.5	1.1-8.3	0.4	1.4	0.1-5.0	0.8	1.3	0.3-3.4	0.6	2.1	0.4-6.2	0.2	0.4	0.0-3.0
	1986-1990	1.2	2.0	0.7-4.4	0.6	2.0	0.4-5.9	0.6	2.1	0.4-5.9	0.2	0.4	0.0-1.9	0.2	0.7	0.0-3.9	0.0	0.0	0.0-2.3
	1991-1995	2.0	3.2	1.5-5.8	1.0	3.4	1.1-8.0	1.0	2.7	0.9-6.6	0.6	0.9	0.2-2.7	0.2	0.7	0.0-3.7	0.4	1.0	0.1-3.9
	1996-2000	1.0	1.4	0.4-3.3	0.8	2.5	0.7-6.4	0.2	0.8	0.0-4.0	0.2	0.2	0.0-1.5	0.2	0.8	0.0-4.1	0.0	0.0	0.0-2.4
	2001-2005	2.0	3.5	1.7-6.5	1.2	4.0	1.4-8.8	0.8	3.0	0.8-7.6	0.2	0.2	0.0-1.6	0.2	0.4	0.0-3.2	0.0	0.0	0.0-2.5
	2006-2010	2.6	4.2	2.2-7.2	1.4	4.8	1.9-10.0	1.2	3.6	1.2-8.1	0.2	0.2	0.0-1.6	0.0	0.0	0.0-2.5	0.2	0.4	0.0-3.1
	2011-2015	2.8	5.6	3.0-9.5	1.6	6.3	2.6-12.5	1.2	5.0	1.7-10.8	0.2	0.2	0.0-1.6	0.2	0.4	0.0-3.2	0.0	0.0	0.0-2.7
	2016-2020	2.4	4.1	2.0-7.4	1.8	6.6	2.8-12.7	0.6	1.6	0.2-5.4	0.6	0.5	0.1-2.1	0.4	0.8	0.1-3.8	0.2	0.4	0.0-3.4
Non-Hodgkin lymphomas	1976-1980	6.0	9.0	6.0-12.9	3.2	10.3	5.8-17.2	2.8	8.1	4.3-13.9	3.4	5.2	3.0-8.4	2.4	8.8	4.3-15.7	1.0	2.9	0.9-7.1
	1981-1985	9.8	13.6	10.0-18.1	3.6	11.3	6.6-18.1	6.2	15.3	10.3-22.1	2.2	3.0	1.5-5.5	0.8	2.3	0.6-6.2	1.4	3.3	1.3-7.2
	1986-1990	9.8	15.0	11.0-19.9	6.2	20.6	13.9-29.5	3.6	9.8	5.6-15.8	6.2	8.3	5.6-11.9	3.2	10.1	5.7-16.6	3.0	7.1	3.9-12.1
	1991-1995	10.2	13.9	10.3-18.3	7.4	22.1	15.5-30.6	2.8	7.3	4.0-12.5	5.6	7.5	5.0-10.9	4.0	12.8	7.7-19.8	1.6	4.1	1.7-8.4
	1996-2000	14.8	19.5	15.3-24.6	9.6	27.1	20.0-36.1	5.2	12.2	7.9-18.2	6.2	7.9	5.4-11.3	3.4	9.5	5.5-15.4	2.8	6.8	3.7-11.7
	2001-2005	17.4	22.3	17.8-27.7	10.8	28.8	21.5-37.8	6.6	16.3	11.0-23.4	4.0	4.7	2.9-7.5	2.6	7.0	3.7-12.3	1.4	2.6	1.0-6.0
	2006-2010	20.2	25.2	20.4-31.0	10.8	28.2	21.0-37.3	9.4	22.3	16.1-30.3	4.6	5.3	3.3-8.2	2.2	5.6	2.7-10.3	2.4	5.4	2.6-10.0
	2011-2015	17.8	19.4	15.4-24.2	10.6	25.4	18.8-33.9	7.2	14.4	10.0-20.6	7.4	7.8	5.5-11.0	3.8	8.7	5.2-14.0	3.6	6.9	4.1-11.5
	2016-2020	20.8	20.9	16.9-25.8	12.2	26.8	20.1-35.3	8.6	16.0	11.4-22.4	6.6	6.2	4.2-9.1	4.6	10.0	6.3-15.6	2.0	3.4	1.6-7.1
Myeloma	1976-1980	2.4	3.5	1.8-6.2	1.0	3.4	1.1-8.3	1.4	3.6	1.4-7.8	1.2	1.6	0.6-3.7	0.6	1.8	0.4-5.7	0.6	1.5	0.3-4.8
	1981-1985	4.2	6.0	3.7-9.4	2.4	7.5	3.8-13.3	1.8	5.1	2.2-10.1	2.4	3.1	1.6-5.6	1.4	4.3	1.7-9.0	1.0	2.2	0.7-5.5
	1986-1990	2.8	3.6	1.9-6.2	1.4	4.2	1.6-8.9	1.4	3.4	1.3-7.3	2.8	3.5	1.9-6.0	1.4	4.0	1.6-8.4	1.4	3.0	1.2-6.5
	1991-1995	2.6	3.4	1.8-5.8	1.6	5.0	2.1-9.8	1.0	2.4	0.7-5.9	3.0	3.8	2.1-6.3	1.8	5.8	2.6-11.0	1.2	2.6	1.0-6.1
	1996-2000	4.8	6.1	3.9-9.1	2.4	6.9	3.6-12.3	2.4	5.6	2.9-10.1	4.0	5.1	3.1-8.0	1.0	3.3	1.1-7.8	3.0	7.0	3.9-11.9
	2001-2005	4.2	5.0	3.1-7.7	2.4	6.8	3.5-12.1	1.8	4.1	1.9-8.2	1.6	1.9	0.8-3.9	1.0	2.8	0.9-6.9	0.6	1.4	0.3-4.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Columbia**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	6.6	7.5	5.1-10.7	4.0	10.0	6.1-15.7	2.6	5.4	2.8-9.7	3.8	4.2	2.5-6.8	1.8	4.8	2.2-9.4	2.0	3.8	1.8-7.7
	2011-2015	5.4	5.3	3.5-7.9	3.6	7.3	4.3-12.0	1.8	3.4	1.5-7.1	2.4	2.4	1.3-4.6	1.4	3.2	1.3-7.1	1.0	1.9	0.6-5.2
	2016-2020	6.4	6.1	4.1-9.0	3.8	7.7	4.4-12.9	2.6	4.5	2.4-8.5	2.4	2.3	1.2-4.4	1.2	2.9	1.0-6.8	1.2	1.8	0.7-5.1
Leukemias	1976-1980	10.0	15.4	11.3-20.6	6.2	20.6	13.7-29.8	3.8	10.4	6.1-16.6	6.8	10.5	7.2-14.9	3.6	11.6	6.7-18.9	3.2	8.6	4.8-14.3
	1981-1985	7.6	10.1	7.1-14.1	3.8	11.3	6.7-18.0	3.8	9.4	5.6-15.1	5.4	7.6	5.0-11.2	2.8	8.3	4.5-14.1	2.6	6.9	3.6-12.1
	1986-1990	6.6	9.1	6.2-12.9	2.6	7.7	4.1-13.4	4.0	10.7	6.4-16.8	3.4	4.7	2.7-7.6	1.4	3.8	1.5-8.1	2.0	5.1	2.4-9.7
	1991-1995	9.2	12.4	9.0-16.6	4.8	14.5	9.3-21.8	4.4	9.9	6.2-15.3	6.2	8.0	5.4-11.5	2.8	8.6	4.6-14.5	3.4	7.5	4.3-12.3
	1996-2000	9.4	12.3	9.0-16.5	4.8	14.5	9.3-21.8	4.6	11.5	7.2-17.5	4.2	5.6	3.5-8.7	2.2	7.1	3.5-12.7	2.0	5.0	2.3-9.5
	2001-2005	10.2	13.7	10.1-18.3	6.0	17.4	11.6-25.0	4.2	10.9	6.5-17.2	6.2	8.0	5.4-11.5	3.0	8.7	4.8-14.4	3.2	7.8	4.3-13.1
	2006-2010	13.6	16.2	12.4-20.8	7.4	18.1	12.6-25.4	6.2	14.5	9.6-21.2	4.8	5.8	3.7-8.8	2.2	5.8	2.9-10.7	2.6	5.6	2.9-10.2
	2011-2015	14.4	16.5	12.7-21.2	8.4	20.4	14.4-28.2	6.0	12.8	8.4-19.0	5.4	5.4	3.5-8.1	4.0	8.5	5.1-13.6	1.4	2.6	1.0-6.2
2016-2020	16.0	16.5	12.9-21.1	10.0	21.9	15.9-29.8	6.0	11.6	7.5-17.4	3.4	3.5	1.9-6.0	2.0	4.1	1.9-8.2	1.4	3.0	1.0-7.2	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cortland**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	169.8	417.8	389.7-447.4	88.2	525.1	475.3-578.4	81.6	358.7	323.9-396.2	78.6	194.9	175.8-215.5	45.2	275.2	239.0-315.0	33.4	148.5	126.3-173.5
	1981-1985	193.4	462.6	433.5-493.2	91.4	533.4	483.8-586.3	102.0	428.6	391.3-468.6	86.2	206.6	187.4-227.3	45.0	267.4	232.5-305.9	41.2	168.4	145.8-193.6
	1986-1990	204.4	472.5	443.6-502.7	105.8	595.0	543.8-649.5	98.6	398.4	363.2-436.0	97.4	226.2	206.3-247.4	52.6	305.4	268.3-345.9	44.8	181.6	158.1-207.6
	1991-1995	207.6	455.9	428.5-484.7	102.0	532.9	486.7-582.2	105.6	412.3	377.3-449.6	89.8	197.2	179.3-216.4	41.0	228.0	196.9-262.3	48.8	182.8	160.3-207.7
	1996-2000	223.2	473.3	445.9-501.9	111.2	543.5	498.8-591.1	112.0	431.2	395.9-468.8	97.6	204.4	186.6-223.4	50.2	258.2	226.8-292.7	47.4	170.7	149.4-194.2
	2001-2005	254.8	515.1	487.1-544.4	134.6	611.7	565.8-660.3	120.2	446.8	411.3-484.6	96.0	193.1	176.1-211.3	48.6	231.8	203.2-263.3	47.4	168.9	147.7-192.4
	2006-2010	276.8	524.8	497.1-553.7	143.2	605.8	561.4-652.8	133.6	469.2	433.4-507.2	103.4	195.8	179.1-213.7	51.6	227.2	199.9-257.2	51.8	176.5	155.2-200.0
	2011-2015	283.2	509.9	483.0-538.0	144.6	558.0	517.0-601.5	138.6	479.6	443.2-518.3	95.4	169.6	154.5-185.9	49.0	201.8	176.7-229.4	46.4	149.3	130.2-170.5
2016-2020	304.8	527.3	500.2-555.4	158.0	572.8	532.3-615.6	146.8	491.6	455.0-530.5	91.8	154.7	140.6-170.0	47.6	177.8	155.4-202.7	44.2	136.5	118.7-156.6	
Oral cavity and pharynx	1976-1980	3.6	8.3	4.9-13.2	2.0	10.5	5.0-19.4	1.6	6.8	2.9-13.5	2.0	4.9	2.3-9.1	1.2	6.7	2.4-14.6	0.8	3.4	0.9-8.9
	1981-1985	4.2	9.5	5.9-14.6	3.2	18.0	9.9-29.7	1.0	3.9	1.3-9.4	1.2	2.8	1.0-6.1	0.4	3.4	0.4-11.1	0.8	3.3	0.9-8.7
	1986-1990	3.8	8.9	5.3-13.9	2.2	10.7	5.3-19.3	1.6	7.4	3.1-14.6	2.0	4.4	2.1-8.2	1.2	6.6	2.2-14.7	0.8	3.1	0.8-8.1
	1991-1995	4.6	10.5	6.7-15.8	2.4	12.8	6.5-22.4	2.2	9.7	4.8-17.3	1.2	2.7	1.0-5.9	0.8	4.7	1.2-11.9	0.4	2.1	0.2-6.9
	1996-2000	3.4	7.2	4.2-11.5	1.8	9.1	4.1-17.2	1.6	6.2	2.7-12.3	0.4	0.9	0.1-3.2	0.4	1.8	0.2-6.6	0.0	0.0	0.0-3.0
	2001-2005	4.6	9.2	5.8-13.9	3.0	12.2	6.8-20.4	1.6	6.2	2.7-12.3	1.4	2.9	1.1-6.0	0.8	3.5	0.9-9.0	0.6	2.2	0.4-6.8
	2006-2010	6.6	12.0	8.2-17.1	4.6	18.3	11.5-27.8	2.0	6.7	3.1-12.7	1.0	1.9	0.6-4.6	0.6	2.3	0.5-7.0	0.4	1.4	0.2-5.6
	2011-2015	7.0	12.8	8.8-18.1	5.0	18.1	11.5-27.2	2.0	8.1	3.7-15.2	0.6	1.0	0.2-3.3	0.2	0.5	0.0-4.2	0.4	1.5	0.2-5.7
2016-2020	8.6	15.5	11.1-21.2	6.0	21.4	14.3-31.1	2.6	10.0	5.0-17.7	0.8	1.4	0.4-3.8	0.4	1.4	0.2-5.7	0.4	1.2	0.1-5.3	
Esophagus	1976-1980	1.4	3.2	1.3-6.6	1.0	4.9	1.6-11.6	0.4	1.5	0.2-5.8	1.2	2.8	1.0-6.2	0.8	4.3	1.1-11.1	0.4	1.5	0.2-5.7
	1981-1985	1.2	2.6	0.9-5.7	1.0	5.0	1.6-11.9	0.2	0.7	0.0-4.5	1.0	2.2	0.7-5.1	1.0	4.9	1.6-11.7	0.0	0.0	0.0-3.2
	1986-1990	2.2	5.5	2.7-9.8	1.6	10.6	4.3-20.8	0.6	2.0	0.4-6.3	2.8	6.4	3.5-10.8	2.2	13.5	6.5-24.4	0.6	2.1	0.4-6.5
	1991-1995	2.8	6.3	3.4-10.5	2.0	11.7	5.4-21.4	0.8	2.8	0.7-7.3	1.4	3.0	1.2-6.2	0.8	4.9	1.3-12.2	0.6	2.1	0.4-6.4
	1996-2000	4.0	8.5	5.2-13.1	2.8	13.4	7.3-22.4	1.2	4.3	1.6-9.6	4.0	8.4	5.1-13.0	2.6	13.0	6.9-22.1	1.4	4.9	1.9-10.3
	2001-2005	4.2	8.6	5.3-13.2	4.0	18.2	11.0-28.2	0.2	0.5	0.0-3.8	4.4	9.0	5.6-13.7	4.0	18.4	11.2-28.5	0.4	1.0	0.1-4.5
	2006-2010	4.8	9.5	6.0-14.2	4.0	17.1	10.3-26.6	0.8	2.9	0.8-7.6	4.2	8.0	4.9-12.4	3.6	15.1	8.8-24.1	0.6	2.2	0.4-6.7
	2011-2015	4.0	7.4	4.5-11.5	3.0	12.9	7.2-21.6	1.0	3.2	1.0-8.1	4.4	8.1	5.0-12.4	3.6	14.6	8.5-23.4	0.8	2.6	0.7-7.3
2016-2020	3.6	5.6	3.3-9.1	2.6	9.0	4.7-15.9	1.0	2.8	0.9-7.3	3.4	5.6	3.2-9.2	2.4	8.1	4.1-14.7	1.0	3.7	1.2-9.2	
Stomach	1976-1980	3.6	9.4	5.5-14.8	2.6	18.4	9.6-31.3	1.0	4.3	1.3-10.3	1.6	4.4	1.9-8.6	1.2	8.7	3.1-18.4	0.4	2.1	0.2-7.3
	1981-1985	3.4	8.0	4.6-12.7	2.0	11.7	5.4-21.7	1.4	5.4	2.2-11.3	1.8	4.3	1.9-8.1	1.2	8.3	2.9-17.7	0.6	2.3	0.5-6.9
	1986-1990	2.0	4.5	2.2-8.3	1.4	7.3	2.9-15.1	0.6	2.2	0.5-6.7	2.0	4.5	2.1-8.2	1.4	7.3	2.9-15.1	0.6	2.1	0.4-6.5
	1991-1995	3.6	8.1	4.8-12.7	2.6	14.4	7.4-24.7	1.0	3.8	1.2-9.1	1.8	4.2	1.9-7.9	1.6	10.5	4.4-20.4	0.2	1.0	0.0-5.0
	1996-2000	2.6	5.5	2.9-9.4	2.0	10.0	4.7-18.4	0.6	2.1	0.4-6.5	1.6	3.3	1.4-6.5	1.4	7.3	2.9-15.0	0.2	0.6	0.0-3.9
	2001-2005	3.0	6.1	3.4-10.1	2.2	9.6	4.7-17.4	0.8	3.1	0.8-8.2	1.4	2.8	1.1-5.9	1.0	4.1	1.3-9.9	0.4	1.5	0.2-5.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cortland**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	3.0	5.3	2.9-8.8	2.0	7.7	3.7-14.5	1.0	3.0	1.0-7.6	1.4	2.4	1.0-5.2	1.0	4.1	1.3-9.8	0.4	1.1	0.1-4.7
	2011-2015	3.4	6.3	3.6-10.2	2.8	11.0	5.9-18.9	0.6	2.1	0.4-6.6	1.0	1.7	0.5-4.3	0.6	2.5	0.5-7.6	0.4	1.1	0.1-4.9
	2016-2020	4.4	7.1	4.4-11.1	3.0	9.9	5.4-16.8	1.4	4.6	1.8-10.2	1.8	2.9	1.3-5.7	1.0	3.6	1.1-8.9	0.8	2.1	0.6-6.3
Colorectal	1976-1980	27.8	69.0	57.9-81.6	12.2	76.1	57.5-98.4	15.6	64.9	51.2-81.2	14.2	35.0	27.3-44.2	6.6	42.5	28.6-60.3	7.6	31.4	22.2-43.3
	1981-1985	32.6	78.4	66.7-91.6	14.2	83.4	64.7-105.7	18.4	72.2	58.0-88.8	17.4	42.5	33.9-52.5	9.0	53.6	38.7-72.1	8.4	32.3	23.2-43.8
	1986-1990	24.0	54.5	45.1-65.3	11.2	64.3	48.0-84.1	12.8	49.2	37.7-63.1	11.2	25.9	19.5-33.7	6.2	39.7	26.5-56.8	5.0	18.6	12.0-27.7
	1991-1995	23.8	51.6	42.7-61.8	9.2	49.7	36.0-66.5	14.6	52.6	41.1-66.3	10.8	23.2	17.4-30.3	3.8	20.6	12.2-32.3	7.0	25.1	17.5-35.1
	1996-2000	28.8	60.6	51.1-71.4	14.2	69.0	53.8-87.1	14.6	54.1	42.3-68.3	12.0	25.1	19.2-32.3	5.6	28.0	18.5-40.4	6.4	23.0	15.7-32.7
	2001-2005	31.6	62.7	53.3-73.4	14.6	71.0	55.5-89.5	17.0	59.1	47.0-73.5	13.0	25.6	19.8-32.8	5.8	28.6	19.0-41.1	7.2	24.5	17.1-34.4
	2006-2010	28.0	52.6	44.1-62.2	11.8	51.7	39.2-66.9	16.2	52.9	41.8-66.4	8.4	15.8	11.4-21.5	4.0	17.5	10.6-27.2	4.4	13.9	8.6-21.6
	2011-2015	23.4	41.9	34.5-50.5	10.4	42.2	31.3-55.8	13.0	41.2	31.5-53.1	8.6	15.1	10.9-20.6	4.6	19.6	12.2-29.7	4.0	11.8	7.1-18.7
	2016-2020	26.2	46.3	38.5-55.3	12.8	49.4	37.7-63.7	13.4	42.5	32.7-54.7	8.0	13.4	9.5-18.6	3.6	14.2	8.2-23.0	4.4	12.4	7.7-19.3
Colon excluding rectum	1976-1980	21.2	52.5	42.9-63.6	8.4	52.5	37.3-71.5	12.8	53.0	40.7-67.8	12.2	30.0	22.9-38.6	5.4	33.9	21.8-49.9	6.8	28.3	19.5-39.7
	1981-1985	24.2	57.9	47.9-69.2	9.8	58.1	42.6-77.3	14.4	55.3	43.2-69.9	15.0	36.1	28.3-45.4	7.8	46.2	32.5-63.5	7.2	27.2	19.0-37.9
	1986-1990	17.8	40.3	32.3-49.6	8.4	48.0	34.0-65.5	9.4	35.9	26.3-47.9	8.8	20.2	14.6-27.1	5.2	32.6	20.8-48.2	3.6	13.6	8.0-21.7
	1991-1995	19.0	41.2	33.3-50.3	6.8	36.9	25.3-51.8	12.2	44.0	33.5-56.7	9.8	21.0	15.5-27.8	3.4	17.8	10.2-28.6	6.4	23.0	15.7-32.6
	1996-2000	18.6	38.9	31.4-47.7	9.8	47.5	35.1-62.9	8.8	31.8	23.0-42.9	9.2	19.2	14.1-25.6	4.2	21.2	13.0-32.4	5.0	18.0	11.6-26.8
	2001-2005	24.6	48.7	40.5-58.2	11.2	54.7	41.1-71.1	13.4	46.5	35.9-59.4	10.4	20.8	15.5-27.4	3.8	18.3	10.9-28.6	6.6	23.0	15.7-32.6
	2006-2010	21.4	40.4	33.0-48.9	8.8	39.0	28.2-52.5	12.6	41.5	31.7-53.6	7.0	13.4	9.3-18.7	3.4	15.1	8.7-24.4	3.6	11.5	6.7-18.6
	2011-2015	18.0	32.2	25.8-39.8	8.0	33.0	23.4-45.4	10.0	31.0	22.8-41.6	7.4	13.1	9.1-18.3	3.8	16.7	9.9-26.3	3.6	10.8	6.3-17.7
	2016-2020	19.2	34.1	27.5-42.0	9.6	37.6	27.5-50.4	9.6	30.4	22.2-41.0	6.0	10.1	6.7-14.7	2.2	9.1	4.3-16.7	3.8	10.6	6.4-17.2
Rectum & rectosigmoid	1976-1980	6.6	16.5	11.3-23.3	3.8	23.6	13.9-37.0	2.8	12.0	6.4-20.2	2.0	5.0	2.4-9.2	1.2	8.6	2.9-18.6	0.8	3.1	0.9-8.2
	1981-1985	8.4	20.6	14.8-27.9	4.4	25.3	15.7-38.4	4.0	16.8	10.2-26.2	2.4	6.4	3.2-11.1	1.2	7.4	2.7-16.0	1.2	5.0	1.8-11.1
	1986-1990	6.2	14.2	9.6-20.2	2.8	16.3	8.7-27.5	3.4	13.3	7.7-21.6	2.4	5.8	3.0-10.0	1.0	7.1	2.2-16.1	1.4	5.0	2.0-10.6
	1991-1995	4.8	10.5	6.7-15.6	2.4	12.7	6.4-22.3	2.4	8.6	4.4-15.3	1.0	2.2	0.7-5.1	0.4	2.8	0.3-9.3	0.6	2.2	0.4-6.5
	1996-2000	10.2	21.7	16.1-28.5	4.4	21.5	13.4-32.6	5.8	22.3	14.9-32.2	2.8	5.9	3.2-9.9	1.4	6.8	2.7-13.9	1.4	5.0	2.0-10.5
	2001-2005	7.0	14.0	9.7-19.5	3.4	16.4	9.5-26.2	3.6	12.6	7.4-20.2	2.6	4.8	2.6-8.3	2.0	10.3	4.9-18.9	0.6	1.6	0.3-5.3
	2006-2010	6.6	12.2	8.4-17.3	3.0	12.7	7.1-21.1	3.6	11.5	6.6-18.6	1.4	2.5	1.0-5.3	0.6	2.4	0.5-7.3	0.8	2.4	0.6-6.8
	2011-2015	5.4	9.7	6.3-14.3	2.4	9.1	4.6-16.4	3.0	10.1	5.6-17.2	1.2	2.1	0.7-4.7	0.8	2.9	0.8-7.9	0.4	0.9	0.1-4.5
	2016-2020	7.0	12.2	8.4-17.2	3.2	11.8	6.6-19.7	3.8	12.1	7.2-19.4	2.0	3.3	1.6-6.3	1.4	5.1	2.0-11.0	0.6	1.7	0.3-5.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cortland**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	0.6	1.5	0.3-4.3	0.4	2.4	0.3-8.5	0.2	0.8	0.0-4.8	0.6	1.5	0.3-4.3	0.4	2.9	0.3-10.1	0.2	0.8	0.0-4.8
	1981-1985	0.6	1.3	0.3-3.9	0.6	3.1	0.6-9.2	0.0	0.0	0.0-3.2	0.6	1.4	0.3-4.0	0.4	2.0	0.2-7.4	0.2	0.7	0.0-4.4
	1986-1990	0.4	1.0	0.1-3.4	0.2	1.2	0.0-6.3	0.2	0.7	0.0-4.3	0.6	1.3	0.2-3.8	0.2	1.2	0.0-6.3	0.4	1.2	0.1-5.0
	1991-1995	1.4	2.9	1.2-6.0	1.2	6.0	2.2-13.1	0.2	0.7	0.0-4.2	2.2	4.6	2.3-8.3	1.4	7.1	2.8-14.6	0.8	2.8	0.8-7.4
	1996-2000	1.2	2.5	0.9-5.5	0.6	2.7	0.6-8.0	0.6	2.4	0.5-6.9	1.0	2.1	0.7-4.8	0.4	2.0	0.2-7.2	0.6	2.3	0.5-6.8
	2001-2005	2.2	4.5	2.2-8.0	1.2	5.7	2.1-12.3	1.0	3.1	1.0-7.6	2.0	4.3	2.0-7.8	1.4	7.3	2.9-14.8	0.6	2.4	0.5-7.0
	2006-2010	2.0	3.6	1.7-6.8	1.2	4.7	1.7-10.5	0.8	2.8	0.7-7.5	2.2	4.2	2.1-7.7	1.0	4.5	1.4-10.5	1.2	4.1	1.5-9.3
	2011-2015	4.0	7.2	4.4-11.4	3.2	12.5	7.1-20.7	0.8	3.1	0.8-8.2	3.6	6.2	3.6-10.0	2.4	9.1	4.6-16.3	1.2	4.0	1.4-9.2
	2016-2020	6.4	10.6	7.2-15.3	5.2	18.8	12.1-28.2	1.2	3.7	1.4-8.8	4.0	6.7	4.0-10.6	2.8	10.1	5.4-17.6	1.2	3.7	1.3-8.6
Pancreas	1976-1980	4.0	9.4	5.7-14.6	2.4	15.6	7.8-27.4	1.6	6.3	2.7-12.6	3.2	7.6	4.3-12.4	2.2	13.3	6.4-24.0	1.0	4.1	1.3-9.6
	1981-1985	4.0	10.1	6.1-15.5	2.0	12.1	5.6-22.4	2.0	8.4	4.0-15.7	4.0	9.9	6.0-15.3	2.2	13.2	6.5-23.5	1.8	7.3	3.3-14.0
	1986-1990	5.0	11.4	7.3-16.8	2.0	10.1	4.8-18.8	3.0	11.0	6.1-18.3	6.2	14.1	9.5-20.0	2.8	14.6	7.9-24.6	3.4	12.5	7.3-20.2
	1991-1995	3.6	8.1	4.8-12.8	1.8	9.2	4.2-17.4	1.8	6.7	3.0-13.0	3.4	7.6	4.4-12.2	1.8	9.3	4.2-17.6	1.6	6.3	2.7-12.5
	1996-2000	6.8	14.2	9.9-19.9	2.2	11.7	5.7-20.9	4.6	16.5	10.4-24.9	6.8	14.2	9.8-19.9	2.6	14.0	7.3-23.9	4.2	14.4	8.8-22.2
	2001-2005	5.8	11.7	7.8-16.9	2.8	13.6	7.4-22.8	3.0	10.9	6.0-18.1	4.8	9.6	6.1-14.3	1.8	9.5	4.3-17.9	3.0	10.8	6.0-18.1
	2006-2010	9.6	18.8	13.8-25.1	3.8	17.6	10.5-27.5	5.8	20.3	13.5-29.5	8.2	15.4	11.0-21.0	3.6	16.0	9.4-25.4	4.6	15.5	9.8-23.7
	2011-2015	6.2	11.1	7.5-16.0	3.0	11.9	6.6-20.1	3.2	11.1	6.3-18.5	5.2	9.2	6.0-13.7	2.4	9.1	4.6-16.3	2.8	9.1	4.9-15.8
	2016-2020	8.6	14.4	10.3-19.7	4.8	17.1	10.8-26.0	3.8	11.9	7.1-19.1	6.8	11.0	7.6-15.6	3.8	12.8	7.6-20.6	3.0	9.2	5.1-15.7
Larynx	1976-1980	3.4	7.8	4.5-12.6	2.6	12.8	6.8-22.2	0.8	3.3	0.9-8.6	0.4	0.8	0.1-3.1	0.4	1.7	0.2-6.8	0.0	0.0	0.0-3.3
	1981-1985	1.2	2.7	1.0-6.0	1.0	6.0	1.8-14.1	0.2	0.8	0.0-4.6	0.6	1.2	0.2-3.6	0.6	2.6	0.5-8.1	0.0	0.0	0.0-3.2
	1986-1990	2.0	4.6	2.2-8.4	1.6	9.0	3.8-17.7	0.4	1.7	0.2-6.1	0.4	0.9	0.1-3.2	0.4	1.9	0.2-7.0	0.0	0.0	0.0-3.1
	1991-1995	2.2	5.2	2.6-9.3	2.0	9.7	4.6-17.9	0.2	1.0	0.0-5.0	0.6	1.2	0.2-3.6	0.4	2.6	0.3-8.9	0.2	0.7	0.0-4.2
	1996-2000	1.6	3.5	1.5-6.9	1.2	5.4	2.0-11.9	0.4	1.7	0.2-6.0	0.4	0.8	0.1-3.0	0.2	1.0	0.0-5.3	0.2	0.6	0.0-4.0
	2001-2005	2.0	3.9	1.9-7.2	1.8	7.4	3.4-14.2	0.2	0.8	0.0-4.5	0.0	0.0	0.0-1.6	0.0	0.0	0.0-3.4	0.0	0.0	0.0-3.0
	2006-2010	1.8	3.1	1.4-6.1	1.2	4.8	1.7-10.6	0.6	2.0	0.4-6.2	1.2	2.3	0.8-5.1	0.8	3.2	0.8-8.4	0.4	1.3	0.1-5.4
	2011-2015	3.4	5.7	3.2-9.3	2.6	9.0	4.7-16.0	0.8	2.4	0.6-6.8	0.6	1.0	0.2-3.1	0.4	1.6	0.2-6.1	0.2	0.5	0.0-4.0
	2016-2020	2.0	3.5	1.7-6.6	1.6	5.9	2.5-12.1	0.4	1.5	0.2-5.8	0.8	1.4	0.4-3.9	0.4	1.8	0.2-6.6	0.4	1.3	0.2-5.5
Lung and bronchus	1976-1980	25.6	61.7	51.3-73.5	18.6	103.6	83.0-127.5	7.0	31.6	21.8-44.2	18.6	44.6	35.9-54.8	14.4	84.1	65.2-106.6	4.2	17.8	10.9-27.4
	1981-1985	24.2	55.6	46.0-66.5	15.8	84.7	66.7-106.0	8.4	34.4	24.6-46.7	18.0	41.9	33.6-51.7	12.8	69.0	52.8-88.4	5.2	22.4	14.5-33.1
	1986-1990	33.6	76.7	65.5-89.4	22.6	123.0	101.0-148.3	11.0	43.6	32.6-57.0	26.8	61.9	51.8-73.4	18.6	104.0	83.4-127.8	8.2	33.6	23.9-45.9
	1991-1995	29.6	65.5	55.3-77.0	16.8	90.0	71.5-111.7	12.8	50.2	38.5-64.4	25.4	56.5	47.0-67.2	13.2	71.9	55.3-91.6	12.2	46.3	35.3-59.7
	1996-2000	35.8	75.9	65.2-87.9	21.0	101.5	82.8-123.0	14.8	58.1	45.6-73.0	28.4	59.8	50.4-70.5	16.8	83.2	66.2-103.2	11.6	43.5	32.9-56.4
	2001-2005	36.6	73.8	63.4-85.4	19.4	88.7	71.7-108.4	17.2	64.1	51.2-79.4	25.0	50.0	41.6-59.7	14.4	68.4	53.3-86.4	10.6	39.7	29.6-52.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cortland**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	46.0	86.3	75.4-98.4	25.6	108.5	90.2-129.4	20.4	70.4	57.2-85.8	30.6	58.0	49.1-68.2	17.4	76.7	61.2-94.9	13.2	45.9	35.3-58.8
	2011-2015	43.6	77.9	67.7-89.2	22.8	90.3	74.1-109.0	20.8	68.9	56.1-84.0	27.4	48.0	40.2-57.0	13.8	54.7	42.3-69.7	13.6	42.2	32.6-54.0
	2016-2020	50.2	83.3	73.2-94.7	25.2	90.9	75.3-108.9	25.0	77.6	64.4-93.1	28.2	47.2	39.6-56.0	15.2	56.5	44.3-71.2	13.0	39.7	30.5-51.1
Melanoma of the skin	1976-1980	4.0	9.8	5.9-15.2	2.2	11.6	5.7-21.0	1.8	8.6	3.7-16.5	0.8	2.6	0.7-6.3	0.6	4.0	0.8-11.1	0.2	1.5	0.0-6.7
	1981-1985	2.8	7.2	3.9-12.0	1.4	7.3	2.9-15.3	1.4	6.7	2.6-14.0	0.6	1.4	0.3-4.0	0.4	1.8	0.2-7.0	0.2	0.7	0.0-4.5
	1986-1990	2.4	5.3	2.7-9.4	1.2	6.7	2.4-14.5	1.2	4.3	1.6-9.7	1.2	3.0	1.1-6.4	0.4	2.6	0.3-8.7	0.8	3.8	1.0-9.7
	1991-1995	4.8	10.4	6.6-15.5	2.2	10.2	5.1-18.4	2.6	10.7	5.6-18.2	1.4	3.0	1.2-6.1	0.2	1.1	0.0-5.7	1.2	4.8	1.7-10.5
	1996-2000	5.4	11.5	7.5-16.7	2.6	12.0	6.4-20.6	2.8	10.5	5.7-17.8	1.2	2.5	0.9-5.5	0.6	3.0	0.6-8.6	0.6	2.4	0.5-7.0
	2001-2005	6.0	12.4	8.3-17.7	3.6	16.1	9.4-25.5	2.4	9.3	4.8-16.4	0.4	0.9	0.1-3.2	0.2	1.0	0.0-5.3	0.2	0.7	0.0-4.4
	2006-2010	8.6	17.6	12.6-23.8	4.2	19.1	11.7-29.3	4.4	17.0	10.5-26.1	1.8	3.3	1.5-6.5	1.0	4.3	1.3-10.2	0.8	2.6	0.7-7.0
	2011-2015	8.4	16.9	12.0-23.1	4.4	18.2	11.2-27.9	4.0	16.4	9.7-25.9	1.4	2.4	1.0-5.2	1.0	4.0	1.2-9.6	0.4	1.3	0.2-5.3
2016-2020	15.0	27.5	21.4-34.9	8.6	33.2	23.6-45.4	6.4	22.8	15.2-33.1	1.0	1.9	0.6-4.8	0.0	0.0	0.0-3.5	1.0	3.6	1.0-9.2	
Female breast	1976-1980							21.8	100.0	81.6-121.2							7.2	34.3	23.7-47.8
	1981-1985							27.2	120.5	100.5-143.2							7.2	30.9	21.4-43.1
	1986-1990							33.4	143.4	121.9-167.5							9.2	42.8	31.0-57.4
	1991-1995							28.4	116.0	97.5-137.1							7.0	27.1	18.7-37.8
	1996-2000							29.8	117.2	98.9-137.8							5.0	17.5	11.2-26.1
	2001-2005							35.6	136.9	117.3-158.8							6.0	21.3	14.2-30.7
	2006-2010							28.8	102.6	86.1-121.3							5.6	19.3	12.7-28.4
	2011-2015							39.6	139.4	119.9-161.3							5.6	18.9	12.4-27.9
2016-2020							40.2	135.9	116.9-157.3							5.4	17.7	11.3-26.5	
Cervix uteri	1976-1980							3.0	14.2	7.7-23.7							1.0	6.4	2.0-14.3
	1981-1985							2.6	11.7	6.0-20.1							1.2	5.5	2.0-12.1
	1986-1990							2.8	12.1	6.5-20.4							1.4	5.1	2.1-10.8
	1991-1995							4.8	19.0	12.1-28.5							1.4	5.0	2.0-10.5
	1996-2000							3.2	12.6	7.2-20.6							1.2	4.9	1.8-10.7
	2001-2005							2.6	11.6	6.2-19.8							0.6	2.5	0.5-7.3
	2006-2010							1.0	4.2	1.3-10.0							0.8	3.3	0.9-8.6
	2011-2015							0.6	2.3	0.5-7.0							0.2	0.5	0.0-4.0
	2016-2020							1.4	8.2	3.2-16.5							0.0	0.0	0.0-3.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cortland**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							7.4	31.9	22.3-44.1							0.8	3.9	1.0-10.0
	1981-1985							6.6	28.4	19.4-40.1							1.0	4.0	1.3-9.4
	1986-1990							5.4	21.1	13.8-30.9							0.8	3.3	0.9-8.4
	1991-1995							6.2	25.0	16.9-35.7							1.2	4.7	1.7-10.5
	1996-2000							8.2	32.1	22.9-43.6							0.8	2.6	0.7-7.0
	2001-2005							9.6	35.1	25.7-46.8							2.2	7.6	3.7-13.9
	2006-2010							10.6	37.2	27.7-49.0							1.2	4.9	1.8-10.7
	2011-2015							14.2	49.2	38.1-62.8							1.6	5.3	2.3-11.0
	2016-2020							10.2	35.4	25.8-47.6							1.6	4.7	2.0-10.0
Ovary	1976-1980							1.4	6.0	2.3-12.5							2.0	9.3	4.3-17.1
	1981-1985							3.8	18.1	10.7-28.4							2.2	8.9	4.4-16.1
	1986-1990							4.8	18.9	11.9-28.4							3.0	12.2	6.7-20.3
	1991-1995							4.8	17.8	11.3-26.6							2.8	10.9	5.9-18.4
	1996-2000							4.4	17.1	10.7-26.0							3.4	12.2	7.1-19.8
	2001-2005							3.4	12.7	7.3-20.6							2.2	8.5	4.2-15.4
	2006-2010							5.2	17.4	11.3-25.9							3.2	11.1	6.3-18.4
	2011-2015							3.6	11.8	6.9-19.1							3.2	10.6	6.0-17.7
	2016-2020							4.6	14.8	9.2-22.9							2.2	7.5	3.6-13.9
Prostate	1976-1980				16.6	107.2	84.4-133.8							4.8	32.2	20.3-48.1			
	1981-1985				21.6	135.3	110.2-164.0							4.0	28.1	16.9-43.4			
	1986-1990				31.4	187.4	158.3-220.0							6.8	44.0	30.1-61.7			
	1991-1995				34.6	177.6	151.7-206.5							5.2	34.6	22.2-50.8			
	1996-2000				32.4	158.4	134.7-184.9							4.8	26.5	16.8-39.3			
	2001-2005				45.0	200.4	174.8-228.6							4.8	24.0	15.3-35.9			
	2006-2010				43.0	173.7	150.8-199.1							4.2	19.9	12.2-30.5			
	2011-2015				37.8	133.1	114.4-154.2							4.0	17.9	10.9-27.8			
	2016-2020				37.8	124.2	106.8-144.0							2.0	8.0	3.8-15.1			
Testis	1976-1980				0.4	2.0	0.2-7.6							0.2	1.2	0.0-6.4			
	1981-1985				0.8	4.1	1.0-10.8							0.0	0.0	0.0-4.0			
	1986-1990				1.0	4.2	1.3-10.3							0.2	1.3	0.0-6.5			
	1991-1995				0.8	3.8	1.0-9.8							0.0	0.0	0.0-3.6			
	1996-2000				1.2	5.4	2.0-11.7							0.2	0.9	0.0-5.1			
	2001-2005				1.8	8.6	3.9-16.3							0.0	0.0	0.0-3.4			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cortland**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				1.2	4.6	1.6-10.4							0.0	0.0	0.0-3.3			
	2011-2015				1.2	5.1	1.8-11.5							0.0	0.0	0.0-3.4			
	2016-2020				1.6	6.8	2.7-13.9							0.0	0.0	0.0-3.5			
Urinary bladder (incl. in situ)	1976-1980	8.8	22.1	16.0-29.8	7.8	47.9	33.6-65.8	1.0	3.9	1.3-9.4	2.6	6.5	3.4-11.1	1.8	11.9	5.3-22.6	0.8	3.0	0.8-8.1
	1981-1985	13.0	32.6	25.0-41.7	8.6	51.8	37.0-70.2	4.4	20.3	12.4-31.0	3.4	8.1	4.7-13.0	2.6	16.3	8.3-28.0	0.8	3.5	0.9-9.0
	1986-1990	12.0	27.2	20.7-35.1	8.8	48.4	34.8-65.3	3.2	11.9	6.7-19.6	2.6	6.3	3.4-10.8	2.0	13.1	6.1-24.1	0.6	2.0	0.4-6.3
	1991-1995	12.0	25.9	19.7-33.3	7.8	42.6	29.9-58.6	4.2	15.8	9.7-24.4	1.0	2.3	0.8-5.4	0.6	3.1	0.6-8.9	0.4	1.7	0.2-6.2
	1996-2000	11.0	23.3	17.6-30.4	7.0	34.0	23.6-47.3	4.0	15.0	9.1-23.3	2.6	5.4	2.9-9.2	2.0	10.4	4.9-18.9	0.6	2.2	0.4-6.6
	2001-2005	11.6	22.8	17.3-29.5	9.0	41.6	30.2-55.8	2.6	9.2	4.8-16.0	0.8	1.5	0.4-3.9	0.6	3.2	0.6-9.0	0.2	0.8	0.0-4.5
	2006-2010	17.4	33.0	26.4-40.9	12.2	54.9	41.8-70.8	5.2	18.2	11.8-27.0	2.8	5.3	2.9-9.0	2.0	9.2	4.3-16.9	0.8	2.7	0.7-7.4
	2011-2015	16.2	29.2	23.0-36.5	13.2	53.5	41.1-68.6	3.0	9.3	5.2-15.9	2.8	5.1	2.8-8.8	2.0	9.5	4.5-17.5	0.8	2.4	0.6-6.7
2016-2020	18.2	30.6	24.5-37.8	13.8	52.4	40.5-66.8	4.4	13.2	8.3-20.6	2.8	5.2	2.7-9.0	2.4	10.1	5.1-18.1	0.4	1.3	0.2-5.4	
Kidney and renal pelvis	1976-1980	2.8	7.6	4.1-12.7	1.8	10.8	4.8-20.5	1.0	5.4	1.7-12.3	1.4	3.9	1.5-7.9	1.2	7.6	2.7-16.4	0.2	1.3	0.0-6.2
	1981-1985	4.6	11.4	7.1-17.1	2.4	14.3	7.2-25.1	2.2	9.3	4.6-16.9	2.2	5.5	2.7-9.8	1.2	7.4	2.7-15.8	1.0	4.0	1.3-9.4
	1986-1990	4.0	9.7	5.8-14.9	2.0	10.9	5.2-20.0	2.0	8.0	3.8-14.9	0.8	2.0	0.5-5.0	0.2	1.3	0.0-6.5	0.6	2.2	0.5-6.7
	1991-1995	3.2	7.2	4.1-11.6	1.4	7.5	3.0-15.3	1.8	7.6	3.4-14.4	1.6	3.5	1.5-6.8	1.2	6.1	2.2-13.2	0.4	1.3	0.2-5.0
	1996-2000	5.8	12.4	8.3-17.8	3.4	16.0	9.3-25.7	2.4	9.1	4.6-16.0	2.0	4.2	2.0-7.8	1.2	7.2	2.5-15.3	0.8	3.2	0.9-8.2
	2001-2005	5.8	12.0	8.0-17.3	4.4	19.9	12.4-30.3	1.4	5.2	2.0-10.9	1.2	2.5	0.9-5.5	0.4	1.8	0.2-6.5	0.8	2.8	0.7-7.7
	2006-2010	5.2	9.7	6.3-14.3	3.0	11.3	6.2-19.1	2.2	8.4	4.1-15.3	1.0	2.1	0.7-4.9	0.0	0.0	0.0-3.3	1.0	4.0	1.3-9.4
	2011-2015	8.2	15.2	10.8-20.9	5.6	22.4	14.7-32.8	2.6	8.8	4.5-15.7	1.2	2.1	0.8-4.8	0.4	1.7	0.2-6.4	0.8	2.7	0.7-7.4
2016-2020	9.2	17.3	12.5-23.5	5.2	20.6	13.2-30.7	4.0	14.3	8.5-22.7	1.8	3.3	1.4-6.6	0.8	2.9	0.8-8.0	1.0	3.8	1.1-9.5	
Brain and other central nervous system	1976-1980	2.0	4.4	2.1-8.3	0.8	3.9	1.0-10.4	1.2	4.8	1.7-10.7	1.8	4.3	1.9-8.2	1.0	5.4	1.7-12.7	0.8	3.5	0.9-9.0
	1981-1985	3.2	6.8	3.8-11.2	1.6	7.9	3.2-16.0	1.6	6.4	2.7-12.7	1.8	4.5	2.0-8.5	0.4	2.4	0.3-8.4	1.4	5.9	2.3-12.3
	1986-1990	4.4	9.8	6.1-14.9	2.4	11.6	5.9-20.5	2.0	7.8	3.7-14.5	1.8	4.0	1.8-7.6	0.4	1.7	0.2-6.6	1.4	5.6	2.2-11.6
	1991-1995	2.8	6.2	3.4-10.4	1.8	8.3	3.8-15.9	1.0	4.3	1.4-10.0	2.2	4.9	2.4-8.7	1.6	7.5	3.2-14.9	0.6	2.5	0.5-7.2
	1996-2000	3.0	6.4	3.6-10.5	2.2	10.4	5.2-18.7	0.8	3.3	0.9-8.4	2.6	5.6	3.0-9.5	1.8	8.5	3.9-16.1	0.8	3.3	0.9-8.5
	2001-2005	2.6	5.3	2.8-9.2	1.6	6.4	2.7-13.0	1.0	3.8	1.2-9.1	2.4	4.9	2.5-8.7	1.2	5.2	1.9-11.4	1.2	4.6	1.6-10.2
	2006-2010	2.0	3.8	1.8-7.1	0.8	3.4	0.9-8.8	1.2	4.3	1.5-9.7	1.6	3.4	1.5-6.7	1.2	5.4	2.0-11.8	0.4	1.7	0.2-6.2
	2011-2015	4.0	7.7	4.6-12.2	2.4	9.5	4.7-17.1	1.6	6.0	2.5-12.2	2.4	4.7	2.4-8.3	1.2	5.3	1.9-11.8	1.2	4.2	1.5-9.6
2016-2020	3.6	6.3	3.6-10.2	1.8	6.6	2.9-13.0	1.8	6.1	2.6-12.2	1.8	3.1	1.4-6.2	1.2	4.7	1.7-10.5	0.6	1.8	0.4-6.0	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cortland**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	1.2	2.9	1.0-6.3	0.0	0.0	0.0-4.1	1.2	5.3	1.8-11.8	0.0	0.0	0.0-1.8	0.0	0.0	0.0-4.1	0.0	0.0	0.0-3.3
	1981-1985	1.4	2.7	1.1-5.8	0.4	1.7	0.2-6.6	1.0	3.5	1.1-8.6	0.2	0.5	0.0-2.7	0.0	0.0	0.0-4.0	0.2	0.8	0.0-4.8
	1986-1990	0.4	0.6	0.1-2.6	0.0	0.0	0.0-3.8	0.4	1.2	0.1-5.0	0.0	0.0	0.0-1.7	0.0	0.0	0.0-3.8	0.0	0.0	0.0-3.1
	1991-1995	1.6	3.3	1.4-6.6	0.2	1.0	0.0-5.4	1.4	5.5	2.2-11.5	0.2	0.4	0.0-2.2	0.0	0.0	0.0-3.6	0.2	0.7	0.0-4.2
	1996-2000	2.2	4.8	2.4-8.6	0.6	2.8	0.6-8.1	1.6	6.9	3.0-13.6	0.0	0.0	0.0-1.6	0.0	0.0	0.0-3.4	0.0	0.0	0.0-3.0
	2001-2005	3.8	7.8	4.7-12.3	1.2	4.9	1.8-11.0	2.6	10.8	5.7-18.5	0.4	0.9	0.1-3.3	0.4	2.0	0.2-7.0	0.0	0.0	0.0-3.0
	2006-2010	5.4	12.3	8.0-17.9	1.0	4.8	1.5-11.2	4.4	19.0	11.7-29.0	0.6	1.0	0.2-3.2	0.0	0.0	0.0-3.3	0.6	1.6	0.3-5.4
	2011-2015	9.4	18.0	13.0-24.3	1.4	5.6	2.1-11.9	8.0	30.0	21.0-41.6	0.0	0.0	0.0-1.7	0.0	0.0	0.0-3.4	0.0	0.0	0.0-3.3
2016-2020	8.4	17.8	12.6-24.4	2.6	10.8	5.6-18.9	5.8	24.7	16.0-36.2	0.0	0.0	0.0-1.7	0.0	0.0	0.0-3.5	0.0	0.0	0.0-3.4	
Hodgkin lymphoma	1976-1980	2.0	4.7	2.2-8.8	1.4	6.2	2.4-13.4	0.6	3.2	0.6-9.1	0.4	1.1	0.1-3.8	0.4	2.2	0.2-7.9	0.0	0.0	0.0-3.3
	1981-1985	3.0	5.7	3.1-9.7	1.2	6.6	2.1-15.0	1.8	5.8	2.6-11.7	0.2	0.5	0.0-2.6	0.0	0.0	0.0-4.0	0.2	0.8	0.0-4.6
	1986-1990	1.4	2.7	1.0-5.7	0.8	3.4	0.9-9.3	0.6	2.2	0.4-6.9	1.6	2.7	1.1-5.6	1.2	4.4	1.6-10.2	0.4	1.2	0.1-5.0
	1991-1995	1.2	2.8	1.0-6.0	0.8	3.7	1.0-9.5	0.4	1.6	0.2-5.8	0.2	0.4	0.0-2.3	0.2	1.2	0.0-6.1	0.0	0.0	0.0-3.0
	1996-2000	1.2	2.4	0.9-5.3	1.0	4.1	1.3-9.9	0.2	0.9	0.0-4.7	0.4	0.7	0.1-2.7	0.4	1.5	0.2-5.8	0.0	0.0	0.0-3.0
	2001-2005	1.6	3.0	1.3-6.0	0.8	2.8	0.7-7.7	0.8	3.1	0.8-8.2	0.2	0.5	0.0-2.7	0.2	1.1	0.0-5.6	0.0	0.0	0.0-3.0
	2006-2010	1.2	2.3	0.8-5.2	0.8	2.9	0.7-7.9	0.4	1.7	0.2-6.3	0.2	0.4	0.0-2.3	0.2	0.8	0.0-4.8	0.0	0.0	0.0-3.1
	2011-2015	1.6	3.3	1.3-6.7	1.0	4.7	1.5-11.1	0.6	1.7	0.3-5.8	0.2	0.4	0.0-2.3	0.2	0.7	0.0-4.6	0.0	0.0	0.0-3.3
2016-2020	1.2	2.7	0.9-6.0	1.0	4.9	1.5-11.5	0.2	0.5	0.0-4.1	0.2	0.5	0.0-2.8	0.2	1.1	0.0-5.6	0.0	0.0	0.0-3.4	
Non-Hodgkin lymphomas	1976-1980	6.0	16.1	10.7-23.0	3.2	21.4	12.1-34.7	2.8	12.6	6.8-21.4	3.4	8.2	4.7-13.2	1.8	11.5	5.0-22.0	1.6	6.9	2.9-13.7
	1981-1985	3.8	9.3	5.6-14.6	1.6	9.9	4.1-19.6	2.2	9.7	4.7-17.5	2.2	6.0	3.0-10.7	1.0	6.6	2.1-14.8	1.2	5.3	1.8-11.6
	1986-1990	4.8	11.7	7.4-17.4	2.8	15.8	8.5-26.7	2.0	8.8	4.1-16.3	2.8	6.7	3.6-11.2	1.4	7.5	2.9-15.5	1.4	5.7	2.2-11.8
	1991-1995	7.2	15.7	11.0-21.8	3.6	18.9	11.0-30.1	3.6	13.8	8.1-21.9	2.6	5.6	3.0-9.6	1.8	10.3	4.5-19.6	0.8	2.8	0.8-7.4
	1996-2000	10.0	21.2	15.7-28.0	4.6	23.3	14.6-35.1	5.4	20.4	13.4-29.8	3.0	6.3	3.5-10.4	0.8	3.6	1.0-9.4	2.2	7.9	3.9-14.3
	2001-2005	8.2	16.2	11.6-22.0	3.8	16.4	9.8-25.8	4.4	15.6	9.7-23.9	4.6	9.2	5.8-13.8	2.4	10.7	5.5-18.8	2.2	6.9	3.4-12.9
	2006-2010	9.8	19.2	14.1-25.5	4.6	20.7	13.1-31.3	5.2	18.6	12.0-27.6	4.6	8.8	5.5-13.3	2.4	10.3	5.2-18.1	2.2	7.9	3.9-14.4
	2011-2015	10.6	19.0	14.1-25.1	6.2	24.3	16.3-35.0	4.4	14.7	9.0-22.7	2.8	5.0	2.7-8.5	1.6	7.2	3.1-14.3	1.2	3.5	1.3-8.4
2016-2020	10.4	18.0	13.4-23.9	6.4	23.9	16.2-34.3	4.0	12.9	7.7-20.6	1.4	2.2	0.9-4.8	0.8	2.3	0.6-6.7	0.6	1.9	0.4-6.2	
Myeloma	1976-1980	2.0	4.7	2.2-8.7	1.0	5.2	1.6-12.4	1.0	4.0	1.3-9.6	1.6	3.9	1.7-7.7	1.2	6.7	2.4-14.7	0.4	1.7	0.2-6.2
	1981-1985	2.8	6.2	3.4-10.4	1.0	4.7	1.5-11.2	1.8	6.8	3.1-13.1	2.0	4.6	2.2-8.5	0.4	1.7	0.2-6.8	1.6	6.1	2.6-12.2
	1986-1990	2.6	6.7	3.5-11.4	1.4	8.1	3.2-16.5	1.2	5.3	1.9-11.7	1.6	3.7	1.6-7.3	1.2	6.3	2.3-13.8	0.4	1.5	0.2-5.7
	1991-1995	4.0	8.8	5.4-13.7	1.4	7.6	3.0-15.5	2.6	10.0	5.3-17.2	3.0	6.4	3.6-10.6	0.6	2.9	0.6-8.5	2.4	8.7	4.5-15.4
	1996-2000	3.2	6.7	3.8-10.9	1.4	8.2	3.2-16.6	1.8	6.7	3.0-12.8	2.0	4.2	2.0-7.7	1.4	8.1	3.2-16.4	0.6	2.0	0.4-6.3
	2001-2005	4.0	8.4	5.1-13.0	2.6	12.5	6.6-21.4	1.4	5.3	2.1-11.1	2.2	4.6	2.3-8.2	1.6	7.7	3.3-15.2	0.6	1.9	0.4-5.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Cortland**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	3.8	7.3	4.4-11.5	2.0	8.0	3.8-15.0	1.8	6.7	3.0-12.9	2.2	4.0	2.0-7.3	1.4	6.3	2.5-13.0	0.8	2.6	0.7-7.1
	2011-2015	3.4	5.9	3.4-9.6	2.2	8.7	4.3-16.0	1.2	4.2	1.5-9.7	2.2	3.8	1.9-7.0	1.2	5.0	1.8-11.2	1.0	3.3	1.1-8.3
	2016-2020	2.6	4.5	2.3-8.1	1.6	6.1	2.5-12.6	1.0	3.2	1.0-8.2	1.4	2.5	1.0-5.4	0.8	3.1	0.8-8.4	0.6	1.9	0.4-6.3
Leukemias	1976-1980	8.0	19.1	13.6-26.2	4.2	24.2	14.5-37.7	3.8	16.1	9.6-25.4	3.8	8.0	4.8-12.7	2.2	11.0	5.3-20.2	1.6	6.0	2.5-12.0
	1981-1985	7.6	17.8	12.6-24.5	3.4	23.1	13.1-37.1	4.2	15.5	9.5-24.0	5.8	13.7	9.2-19.7	3.2	20.9	11.5-34.2	2.6	9.9	5.2-17.1
	1986-1990	5.6	12.6	8.3-18.2	3.4	19.2	10.9-30.8	2.2	8.3	4.1-15.1	3.2	7.3	4.1-11.8	2.4	14.0	7.1-24.6	0.8	2.9	0.7-7.7
	1991-1995	4.6	10.0	6.3-15.0	3.4	16.8	9.6-27.1	1.2	3.9	1.4-8.9	3.6	7.8	4.6-12.3	2.2	11.2	5.4-20.2	1.4	4.6	1.8-9.8
	1996-2000	5.2	11.0	7.2-16.1	2.2	11.8	5.8-21.0	3.0	11.2	6.2-18.7	4.2	8.7	5.4-13.3	2.0	12.4	5.8-22.4	2.2	7.4	3.7-13.6
	2001-2005	8.0	15.9	11.3-21.7	5.0	21.7	13.9-32.3	3.0	10.9	6.0-18.2	4.4	8.9	5.5-13.5	2.6	12.0	6.3-20.6	1.8	6.2	2.8-12.0
	2006-2010	9.8	19.6	14.4-26.1	6.0	27.2	18.2-39.1	3.8	14.1	8.3-22.4	4.2	7.9	4.8-12.1	2.2	10.1	5.0-18.1	2.0	6.3	3.0-12.1
	2011-2015	8.4	15.6	11.1-21.2	5.4	21.8	14.2-32.0	3.0	10.2	5.6-17.5	3.6	6.6	3.9-10.7	2.4	10.7	5.4-18.9	1.2	4.1	1.5-9.5
2016-2020	9.4	16.1	11.7-21.7	5.2	19.6	12.6-29.2	4.2	13.6	8.2-21.4	4.0	6.6	4.0-10.5	2.8	10.9	5.8-18.8	1.2	3.4	1.2-8.2	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Delaware**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	193.6	387.3	362.8-413.2	96.0	434.9	395.7-477.0	97.6	360.7	328.5-395.3	95.6	189.3	172.4-207.4	47.2	219.6	191.7-250.4	48.4	167.5	146.8-190.4
	1981-1985	206.8	395.2	371.0-420.7	110.4	488.6	447.3-532.5	96.4	336.7	306.2-369.5	112.6	215.1	197.3-234.1	64.4	293.3	261.1-328.4	48.2	162.3	141.8-185.0
	1986-1990	237.2	441.2	415.9-467.7	128.0	554.6	511.0-600.9	109.2	373.4	341.5-407.6	118.2	215.5	198.2-234.0	64.8	292.4	260.2-327.5	53.4	172.9	152.1-195.9
	1991-1995	236.4	403.3	380.2-427.4	125.2	481.8	443.9-522.2	111.2	345.2	316.2-376.3	123.0	204.5	188.4-221.6	61.8	250.5	222.5-281.1	61.2	177.9	158.0-199.8
	1996-2000	273.6	431.2	408.3-455.1	142.2	496.4	459.9-535.1	131.4	391.0	360.6-423.4	125.8	190.2	175.4-205.9	64.8	231.2	206.3-258.5	61.0	161.3	143.2-181.3
	2001-2005	317.8	488.1	463.9-513.4	165.8	540.9	504.0-580.0	152.0	448.0	415.6-482.5	125.6	185.0	170.7-200.4	64.2	214.3	191.1-239.7	61.4	165.6	147.2-186.1
	2006-2010	319.6	469.4	445.9-494.0	177.6	537.3	501.5-575.1	142.0	418.9	387.1-452.9	117.0	166.3	152.8-180.8	64.6	199.5	177.8-223.3	52.4	139.1	122.3-157.9
	2011-2015	323.8	458.4	435.0-482.9	163.2	467.9	434.7-503.3	160.6	452.3	419.2-487.5	118.4	158.3	145.5-172.2	63.2	181.0	161.0-203.2	55.2	139.4	122.8-158.1
	2016-2020	330.2	456.5	432.7-481.5	170.6	466.4	433.3-501.7	159.6	450.6	416.3-487.2	115.8	146.9	134.6-160.3	63.2	168.4	149.4-189.6	52.6	127.1	111.5-144.9
Oral cavity and pharynx	1976-1980	3.2	6.8	3.8-11.2	1.8	8.0	3.5-15.5	1.4	5.7	2.2-11.9	2.6	5.2	2.8-9.1	1.6	7.3	3.1-14.6	1.0	3.6	1.2-8.8
	1981-1985	5.6	11.1	7.3-16.3	3.8	18.2	10.8-28.7	1.8	6.3	2.8-12.5	3.4	6.4	3.7-10.4	2.4	10.8	5.4-19.1	1.0	3.4	1.0-8.4
	1986-1990	6.2	12.7	8.6-18.2	5.2	24.8	16.0-36.7	1.0	3.8	1.2-9.3	1.4	2.3	0.9-5.0	1.2	5.0	1.7-11.4	0.2	0.6	0.0-3.9
	1991-1995	5.4	9.7	6.3-14.3	3.6	14.2	8.3-22.8	1.8	5.9	2.6-11.5	2.2	3.6	1.8-6.5	1.4	5.6	2.2-11.8	0.8	2.5	0.6-6.7
	1996-2000	4.4	6.8	4.2-10.5	3.2	11.9	6.7-19.6	1.2	3.2	1.2-7.6	1.6	2.6	1.1-5.3	1.2	4.4	1.6-9.8	0.4	1.0	0.1-4.5
	2001-2005	8.4	14.5	10.3-20.0	6.0	22.4	14.9-32.5	2.4	7.4	3.6-13.7	1.2	1.9	0.7-4.5	1.0	3.6	1.2-8.8	0.2	0.6	0.0-4.1
	2006-2010	6.8	10.7	7.3-15.2	5.0	16.1	10.2-24.4	1.8	4.9	2.2-10.0	1.2	1.7	0.6-4.0	0.2	0.6	0.0-4.2	1.0	2.5	0.8-6.8
	2011-2015	10.2	15.2	11.1-20.6	7.4	21.3	14.7-30.3	2.8	9.4	4.8-16.9	2.0	2.7	1.3-5.3	1.2	3.3	1.2-8.1	0.8	2.4	0.6-6.9
	2016-2020	9.4	13.0	9.3-17.9	7.0	18.7	12.8-27.1	2.4	7.9	3.7-15.0	2.0	2.9	1.3-5.8	1.6	4.5	1.8-10.0	0.4	1.2	0.1-5.7
Esophagus	1976-1980	1.0	1.8	0.6-4.4	1.0	4.2	1.3-10.2	0.0	0.0	0.0-3.2	1.6	3.0	1.3-6.0	1.0	4.0	1.2-9.7	0.6	2.1	0.4-6.5
	1981-1985	1.4	2.4	1.0-5.2	1.4	5.6	2.2-12.0	0.0	0.0	0.0-3.2	1.2	2.3	0.8-5.2	1.2	5.7	2.0-12.7	0.0	0.0	0.0-3.2
	1986-1990	1.4	2.3	0.9-4.9	0.8	3.0	0.8-8.1	0.6	1.6	0.3-5.5	2.2	3.8	1.9-6.9	1.4	6.4	2.5-13.5	0.8	2.4	0.6-6.6
	1991-1995	2.6	4.1	2.2-7.3	1.6	5.4	2.3-11.1	1.0	2.6	0.8-6.7	2.4	3.9	2.0-6.9	1.2	4.3	1.6-9.6	1.2	3.0	1.1-7.2
	1996-2000	2.8	4.4	2.4-7.6	1.8	6.2	2.8-12.1	1.0	2.4	0.8-6.5	2.0	3.0	1.4-5.8	1.4	4.4	1.7-9.5	0.6	1.8	0.4-5.9
	2001-2005	4.4	6.4	4.0-9.9	3.4	10.3	6.0-17.0	1.0	2.9	0.9-7.5	4.2	6.0	3.7-9.5	2.8	8.6	4.7-15.0	1.4	3.8	1.5-8.5
	2006-2010	6.4	8.8	6.0-12.7	4.8	13.5	8.6-20.7	1.6	4.2	1.8-9.1	4.6	6.8	4.3-10.5	4.0	12.2	7.4-19.4	0.6	1.6	0.3-5.5
	2011-2015	5.2	6.7	4.4-10.3	4.6	12.1	7.6-18.9	0.6	1.5	0.3-5.7	4.8	6.4	4.1-9.9	3.4	9.6	5.5-16.1	1.4	3.3	1.3-7.8
	2016-2020	3.6	4.1	2.4-7.0	2.6	6.2	3.3-11.7	1.0	2.1	0.7-6.4	3.6	4.2	2.5-7.2	3.0	7.5	4.2-13.3	0.6	1.3	0.3-5.5
Stomach	1976-1980	4.4	8.9	5.6-13.6	3.2	15.5	8.8-25.3	1.2	3.9	1.4-8.9	2.4	4.5	2.3-8.0	2.0	9.3	4.4-17.3	0.4	1.1	0.1-4.9
	1981-1985	5.6	10.3	6.8-15.1	3.8	18.1	10.6-28.6	1.8	5.3	2.4-10.6	4.0	7.6	4.6-11.9	2.6	12.8	6.7-22.1	1.4	4.3	1.7-9.3
	1986-1990	4.4	8.1	5.0-12.4	3.6	16.9	9.7-27.2	0.8	2.2	0.6-6.3	3.6	6.5	3.8-10.4	2.8	12.4	6.5-21.2	0.8	2.3	0.6-6.4
	1991-1995	3.0	5.0	2.8-8.3	2.2	8.9	4.4-16.1	0.8	2.3	0.6-6.5	2.4	3.7	1.9-6.7	1.4	5.0	2.0-10.6	1.0	2.8	0.9-7.0
	1996-2000	3.8	5.9	3.5-9.4	2.4	8.0	4.1-14.4	1.4	4.0	1.6-8.8	3.4	5.1	3.0-8.4	1.6	6.0	2.5-12.2	1.8	4.7	2.1-9.5
	2001-2005	3.4	5.2	3.0-8.5	2.4	8.3	4.2-15.0	1.0	2.5	0.8-6.7	2.6	3.9	2.1-6.9	1.8	6.0	2.7-11.9	0.8	1.9	0.5-5.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Delaware**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	4.2	6.0	3.7-9.5	3.0	9.3	5.1-15.8	1.2	3.3	1.2-7.9	3.2	4.8	2.7-8.1	2.0	6.0	2.8-11.6	1.2	3.8	1.3-9.0
	2011-2015	3.8	5.2	3.0-8.5	2.2	5.4	2.7-10.7	1.6	4.9	1.9-10.5	1.2	1.6	0.6-3.9	0.8	2.1	0.6-6.4	0.4	1.0	0.1-4.9
	2016-2020	4.2	5.1	3.1-8.4	2.8	6.6	3.6-12.1	1.4	3.8	1.4-9.1	1.2	1.4	0.5-3.7	0.8	1.9	0.5-6.3	0.4	0.8	0.1-4.8
Colorectal	1976-1980	34.8	68.3	58.4-79.5	16.0	73.9	58.1-92.6	18.8	65.3	52.6-80.4	19.0	37.3	30.1-45.8	7.4	34.2	23.9-47.5	11.6	39.9	30.2-51.9
	1981-1985	37.4	69.9	60.0-80.9	17.0	73.8	58.5-92.0	20.4	67.1	54.4-82.0	17.8	33.4	26.7-41.2	9.0	40.7	29.4-54.9	8.8	28.2	20.4-38.3
	1986-1990	34.2	62.2	53.1-72.5	15.2	64.8	50.7-81.8	19.0	58.8	47.3-72.4	19.0	35.1	28.3-43.1	7.4	34.8	24.1-48.6	11.6	36.7	27.6-48.1
	1991-1995	31.2	50.6	42.9-59.4	15.2	59.4	46.4-74.9	16.0	46.1	36.3-57.9	17.8	28.8	23.1-35.6	8.8	34.6	24.8-47.0	9.0	24.8	18.0-33.7
	1996-2000	36.2	54.7	46.9-63.4	20.0	70.7	57.3-86.4	16.2	42.5	33.6-53.5	16.4	24.1	19.2-30.2	8.4	29.7	21.3-40.6	8.0	19.4	13.7-27.1
	2001-2005	41.2	60.3	52.2-69.4	20.6	66.8	54.3-81.5	20.6	52.9	43.0-64.9	14.4	20.9	16.3-26.6	7.4	24.2	17.0-33.7	7.0	17.2	11.9-24.7
	2006-2010	31.6	45.3	38.3-53.3	16.0	47.9	37.7-60.2	15.6	42.3	33.0-53.7	11.8	16.3	12.4-21.2	6.2	19.0	12.8-27.4	5.6	13.3	8.7-19.9
	2011-2015	32.2	45.6	38.5-53.8	16.6	48.3	38.0-60.9	15.6	41.9	32.7-53.4	11.0	14.1	10.6-18.7	5.8	16.3	10.8-24.2	5.2	11.8	7.7-18.1
2016-2020	27.4	38.7	32.1-46.5	15.6	46.1	35.9-58.6	11.8	32.7	24.2-43.7	8.8	11.1	8.0-15.4	5.2	12.9	8.4-19.8	3.6	9.4	5.5-15.9	
Colon excluding rectum	1976-1980	24.4	48.0	39.8-57.5	10.2	48.9	36.0-64.7	14.2	48.9	38.0-62.0	14.4	28.1	22.0-35.5	4.6	21.4	13.4-32.4	9.8	33.2	24.5-44.2
	1981-1985	25.0	46.5	38.5-55.6	10.2	45.2	33.3-60.1	14.8	48.3	37.6-61.1	14.6	27.6	21.5-34.9	6.8	31.5	21.5-44.4	7.8	25.1	17.7-34.8
	1986-1990	23.2	42.5	35.0-51.2	9.6	40.7	29.8-54.5	13.6	42.1	32.5-53.9	16.0	29.3	23.1-36.7	6.0	28.7	19.0-41.4	10.0	31.4	23.0-41.9
	1991-1995	22.0	35.5	29.2-43.0	10.6	41.8	31.0-55.2	11.4	32.4	24.4-42.4	15.2	24.5	19.2-30.8	7.4	28.7	19.9-40.1	7.8	21.5	15.2-29.8
	1996-2000	25.8	39.0	32.5-46.5	14.4	51.2	39.9-64.9	11.4	29.2	22.0-38.5	12.8	18.8	14.5-24.2	6.4	22.7	15.4-32.4	6.4	15.4	10.4-22.4
	2001-2005	30.4	44.0	37.2-51.8	14.4	46.9	36.6-59.5	16.0	40.5	32.0-51.1	11.8	16.9	12.9-22.1	5.8	18.6	12.5-27.2	6.0	14.3	9.5-21.2
	2006-2010	22.8	32.2	26.4-38.9	10.8	32.0	24.0-42.3	12.0	32.1	24.2-42.1	10.4	14.3	10.6-19.0	5.2	16.2	10.5-24.3	5.2	12.2	7.9-18.6
	2011-2015	22.0	31.0	25.3-37.9	10.4	31.6	23.2-42.4	11.6	29.7	22.4-39.2	8.6	11.0	7.9-15.2	4.0	11.4	6.8-18.3	4.6	10.3	6.5-16.4
2016-2020	19.0	27.2	21.6-34.0	10.8	32.3	23.8-43.2	8.2	23.3	16.1-33.1	6.0	7.8	5.2-11.6	3.2	8.0	4.5-13.9	2.8	7.6	4.0-13.8	
Rectum & rectosigmoid	1976-1980	10.4	20.3	15.0-26.8	5.8	25.0	16.5-36.4	4.6	16.4	10.3-25.0	4.6	9.2	5.8-14.0	2.8	12.8	6.9-21.8	1.8	6.6	2.9-13.0
	1981-1985	12.4	23.4	17.8-30.2	6.8	28.6	19.6-40.4	5.6	18.8	12.4-27.6	3.2	5.8	3.3-9.5	2.2	9.2	4.6-16.8	1.0	3.1	1.0-7.7
	1986-1990	11.0	19.7	14.7-25.8	5.6	24.1	15.7-35.3	5.4	16.7	10.9-24.8	3.0	5.8	3.2-9.7	1.4	6.2	2.3-13.1	1.6	5.4	2.2-11.0
	1991-1995	9.2	15.1	11.0-20.3	4.6	17.6	11.0-26.8	4.6	13.7	8.5-21.1	2.6	4.3	2.3-7.5	1.4	5.9	2.3-12.3	1.2	3.4	1.2-7.9
	1996-2000	10.4	15.7	11.7-20.7	5.6	19.5	12.9-28.6	4.8	13.3	8.4-20.4	3.6	5.3	3.2-8.7	2.0	7.0	3.3-13.3	1.6	4.1	1.7-8.7
	2001-2005	10.8	16.3	12.2-21.5	6.2	19.9	13.4-28.7	4.6	12.4	7.7-19.4	2.6	4.0	2.1-7.0	1.6	5.5	2.3-11.3	1.0	2.9	0.9-7.5
	2006-2010	8.8	13.2	9.4-18.0	5.2	15.8	10.1-23.9	3.6	10.2	5.8-17.0	1.4	2.0	0.8-4.4	1.0	2.8	0.9-7.2	0.4	1.1	0.1-4.8
	2011-2015	10.2	14.6	10.6-19.7	6.2	16.8	11.2-24.6	4.0	12.2	7.0-20.0	2.4	3.1	1.6-5.8	1.8	5.0	2.2-10.2	0.6	1.5	0.3-5.5
2016-2020	8.4	11.5	8.1-16.1	4.8	13.8	8.6-21.5	3.6	9.4	5.4-16.1	2.8	3.3	1.8-6.1	2.0	4.9	2.3-10.1	0.8	1.8	0.5-6.2	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020  
Delaware**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	2.0	3.9	1.9-7.3	0.8	3.0	0.8-8.2	1.2	4.1	1.5-9.3	1.4	2.9	1.1-6.0	0.6	2.6	0.5-7.8	0.8	2.8	0.8-7.5
	1981-1985	3.2	6.0	3.4-9.9	1.8	8.5	3.8-16.4	1.4	4.4	1.8-9.5	3.0	5.6	3.1-9.4	1.8	8.1	3.6-15.8	1.2	3.8	1.4-8.8
	1986-1990	2.2	4.0	2.0-7.3	1.8	7.2	3.3-13.9	0.4	1.6	0.2-5.8	2.8	5.3	2.9-9.0	2.2	9.0	4.4-16.4	0.6	2.1	0.4-6.5
	1991-1995	2.6	4.4	2.3-7.6	2.0	7.8	3.7-14.5	0.6	1.6	0.3-5.3	3.4	5.7	3.3-9.4	1.4	5.1	2.0-10.8	2.0	6.2	2.9-11.9
	1996-2000	2.0	3.5	1.6-6.6	1.4	5.3	2.1-11.2	0.6	2.2	0.4-6.7	2.8	4.3	2.3-7.4	1.6	5.3	2.3-10.9	1.2	3.6	1.3-8.4
	2001-2005	2.8	4.0	2.2-7.0	2.2	6.5	3.2-12.1	0.6	1.7	0.3-5.7	2.2	3.2	1.6-6.0	1.4	4.5	1.8-9.8	0.8	2.1	0.6-6.3
	2006-2010	1.8	2.4	1.1-4.9	1.0	3.0	1.0-7.7	0.8	2.0	0.5-6.0	2.4	3.3	1.7-6.1	1.6	4.7	2.0-10.0	0.8	2.4	0.7-6.9
	2011-2015	4.0	5.3	3.2-8.6	3.4	9.0	5.1-15.4	0.6	1.4	0.3-5.4	3.0	3.6	2.0-6.4	2.4	6.0	3.1-11.4	0.6	1.4	0.3-5.4
2016-2020	4.8	7.2	4.3-11.5	3.4	8.4	4.8-14.4	1.4	6.2	2.2-13.8	3.6	5.0	2.8-8.5	2.4	7.1	3.3-13.7	1.2	3.0	1.1-7.9	
Pancreas	1976-1980	4.8	9.2	5.8-13.8	1.8	7.8	3.5-15.2	3.0	10.0	5.6-16.8	5.2	9.9	6.5-14.7	2.0	8.3	3.9-15.7	3.2	11.0	6.3-18.1
	1981-1985	6.8	13.1	9.0-18.5	2.8	12.9	6.8-22.0	4.0	12.6	7.6-20.0	5.6	10.9	7.2-15.9	2.4	11.0	5.5-19.5	3.2	10.2	5.7-17.2
	1986-1990	6.0	11.2	7.5-16.1	3.6	15.5	9.1-24.7	2.4	7.4	3.7-13.5	4.8	9.0	5.7-13.5	2.6	11.1	5.8-19.2	2.2	6.9	3.3-12.9
	1991-1995	5.2	8.2	5.3-12.2	2.0	7.8	3.6-14.7	3.2	8.4	4.8-14.2	5.8	9.6	6.4-13.9	2.4	10.0	5.0-17.7	3.4	9.6	5.6-15.9
	1996-2000	5.2	7.8	5.1-11.6	1.8	5.8	2.6-11.4	3.4	9.1	5.3-15.1	4.8	7.0	4.5-10.6	1.6	5.3	2.3-10.8	3.2	8.2	4.6-13.9
	2001-2005	7.6	10.9	7.7-15.2	2.8	8.6	4.7-14.9	4.8	12.4	7.9-19.1	6.0	8.4	5.7-12.3	2.2	6.8	3.4-12.6	3.8	9.6	5.8-15.7
	2006-2010	8.6	12.1	8.7-16.6	5.2	15.8	10.2-23.8	3.4	9.4	5.4-15.6	8.0	11.0	7.9-15.3	4.6	13.5	8.5-20.8	3.4	9.1	5.3-15.1
	2011-2015	7.4	9.9	6.9-14.1	3.8	9.9	5.9-16.2	3.6	9.5	5.4-16.0	5.8	7.6	5.0-11.2	3.4	9.5	5.5-15.9	2.4	5.9	3.0-11.2
2016-2020	11.2	14.7	10.9-19.7	5.8	17.6	11.4-26.4	5.4	12.0	7.9-18.5	7.2	9.7	6.6-14.0	3.4	10.9	6.0-18.6	3.8	8.6	5.1-14.5	
Larynx	1976-1980	2.4	4.3	2.2-7.8	1.8	6.3	2.9-12.5	0.6	2.5	0.4-7.6	0.4	1.0	0.1-3.5	0.2	0.8	0.0-4.8	0.2	1.3	0.0-6.0
	1981-1985	3.2	6.5	3.6-10.7	1.8	7.8	3.4-15.2	1.4	5.5	2.1-11.6	0.6	1.0	0.2-3.2	0.4	1.3	0.2-5.5	0.2	0.7	0.0-4.4
	1986-1990	2.6	4.4	2.3-7.7	1.8	7.1	3.1-13.8	0.8	2.3	0.6-6.5	0.8	1.6	0.4-4.3	0.8	4.1	1.0-10.4	0.0	0.0	0.0-3.1
	1991-1995	3.4	6.2	3.6-10.1	3.0	11.7	6.5-19.5	0.4	1.6	0.2-5.8	0.6	0.9	0.2-2.9	0.4	1.4	0.2-5.6	0.2	0.5	0.0-3.8
	1996-2000	2.8	4.1	2.2-7.2	1.6	4.9	2.1-10.2	1.2	3.4	1.2-8.0	1.0	1.5	0.5-3.7	0.8	2.5	0.7-7.0	0.2	0.5	0.0-3.9
	2001-2005	1.0	1.7	0.5-4.1	0.6	2.3	0.5-7.0	0.4	1.1	0.1-4.8	0.4	0.6	0.1-2.4	0.4	1.5	0.2-5.7	0.0	0.0	0.0-3.3
	2006-2010	3.2	4.4	2.5-7.5	2.6	8.0	4.2-14.2	0.6	1.6	0.3-5.6	0.8	1.0	0.3-3.1	0.2	0.5	0.0-4.1	0.6	1.6	0.3-5.5
	2011-2015	3.6	5.1	2.9-8.4	3.2	8.8	4.9-15.1	0.4	1.3	0.2-5.5	0.8	1.0	0.3-3.2	0.8	2.0	0.5-6.3	0.0	0.0	0.0-3.6
2016-2020	3.4	4.0	2.3-7.0	2.6	6.2	3.3-11.7	0.8	2.1	0.5-6.7	1.0	1.2	0.4-3.5	1.0	2.6	0.8-7.3	0.0	0.0	0.0-3.9	
Lung and bronchus	1976-1980	21.8	42.2	34.4-51.1	16.0	69.0	54.2-86.6	5.8	20.3	13.5-29.5	17.6	34.3	27.3-42.5	11.8	51.6	38.9-67.1	5.8	20.4	13.6-29.7
	1981-1985	31.0	58.1	49.1-68.3	22.0	93.8	76.5-113.7	9.0	30.6	22.1-41.5	28.2	53.9	45.2-63.9	20.0	88.1	71.2-107.8	8.2	28.1	19.9-38.6
	1986-1990	37.2	66.1	56.8-76.6	23.2	96.8	79.5-116.8	14.0	45.5	35.2-58.1	29.6	53.1	44.8-62.7	18.6	80.8	64.7-99.6	11.0	34.8	26.0-45.9
	1991-1995	37.8	63.8	54.9-73.9	23.2	88.1	72.4-106.1	14.6	43.3	33.7-54.9	30.8	51.8	43.8-60.8	17.4	66.1	52.6-82.0	13.4	40.2	30.9-51.6
	1996-2000	45.4	69.3	60.5-79.2	25.6	87.6	72.8-104.6	19.8	56.3	45.5-69.2	32.6	49.0	41.8-57.3	20.2	68.7	55.8-83.8	12.4	33.1	25.3-43.0
	2001-2005	45.2	67.5	58.9-77.1	27.2	87.8	73.4-104.3	18.0	51.2	41.0-63.5	34.0	51.0	43.6-59.6	18.8	60.3	48.6-74.3	15.2	44.6	34.9-56.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Delaware**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	54.4	75.8	67.0-85.7	33.0	96.5	82.1-113.0	21.4	58.6	47.8-71.4	33.8	48.5	41.3-56.7	20.8	62.8	51.1-76.8	13.0	35.7	27.4-46.1
	2011-2015	52.2	68.3	60.1-77.5	29.0	78.3	65.9-92.8	23.2	59.7	49.1-72.4	35.0	46.2	39.5-54.0	20.0	54.8	44.3-67.4	15.0	39.0	30.4-49.7
	2016-2020	49.6	62.5	54.6-71.5	25.6	65.7	54.2-79.5	24.0	59.8	49.1-72.8	29.8	37.8	31.7-44.9	16.0	41.9	32.8-53.2	13.8	33.8	25.9-44.0
Melanoma of the skin	1976-1980	4.8	10.8	6.8-16.3	3.4	17.1	9.8-27.6	1.4	5.5	2.2-11.6	1.8	4.1	1.8-7.8	1.6	8.2	3.5-16.2	0.2	0.7	0.0-4.3
	1981-1985	4.2	9.5	5.8-14.7	2.4	11.4	5.7-20.2	1.8	8.9	3.9-16.8	1.2	2.6	0.9-5.8	0.8	3.6	0.9-9.5	0.4	1.7	0.2-6.2
	1986-1990	5.4	10.7	6.9-15.7	3.6	15.9	9.2-25.4	1.8	7.0	3.1-13.6	1.2	2.3	0.8-5.1	1.2	4.8	1.7-10.9	0.0	0.0	0.0-3.1
	1991-1995	3.4	6.3	3.6-10.2	2.4	9.7	4.9-17.0	1.0	3.2	1.0-7.9	1.0	1.6	0.5-4.0	0.8	3.5	0.9-9.0	0.2	0.6	0.0-3.9
	1996-2000	6.6	11.5	7.8-16.4	4.6	16.6	10.4-25.3	2.0	7.1	3.3-13.3	1.2	2.2	0.8-5.0	1.0	4.2	1.3-9.9	0.2	0.4	0.0-3.6
	2001-2005	9.2	15.7	11.4-21.3	5.0	17.8	11.3-26.8	4.2	14.1	8.5-22.1	1.8	3.1	1.4-6.1	0.8	3.1	0.8-8.3	1.0	3.0	0.9-7.7
	2006-2010	15.0	22.0	17.2-27.8	9.0	27.1	19.6-36.8	6.0	17.4	11.6-25.5	2.0	3.5	1.6-6.8	1.4	5.4	2.0-11.7	0.6	1.6	0.3-5.6
	2011-2015	13.6	20.8	15.8-27.0	7.2	23.7	16.0-34.0	6.4	18.0	11.9-26.5	1.8	2.4	1.1-5.0	1.4	4.4	1.7-9.7	0.4	0.9	0.1-4.7
2016-2020	15.2	24.7	18.8-32.0	9.4	27.6	19.9-37.7	5.8	22.8	14.2-34.6	1.0	2.0	0.5-5.2	0.6	2.6	0.4-8.5	0.4	1.5	0.1-6.5	
Female breast	1976-1980							27.4	107.5	89.7-127.9							8.2	27.9	19.9-38.1
	1981-1985							21.0	75.4	61.1-92.0							6.0	22.8	15.0-33.1
	1986-1990							28.8	103.2	86.3-122.5							9.0	30.5	21.9-41.5
	1991-1995							32.0	107.5	90.9-126.3							9.8	29.5	21.6-39.6
	1996-2000							34.4	110.1	93.9-128.6							8.8	26.0	18.7-35.5
	2001-2005							39.4	121.7	104.8-140.8							7.2	19.2	13.3-27.2
	2006-2010							36.6	112.4	95.9-131.1							7.0	18.7	13.0-26.6
	2011-2015							38.4	106.7	91.0-124.5							7.4	18.5	12.9-26.3
2016-2020							44.0	121.6	104.8-140.8							7.8	18.9	13.2-26.9	
Cervix uteri	1976-1980							4.0	16.1	9.7-25.3							1.2	4.0	1.4-9.2
	1981-1985							2.0	8.4	3.9-15.7							1.2	4.5	1.5-10.3
	1986-1990							3.4	12.4	7.0-20.2							1.2	4.4	1.5-9.9
	1991-1995							2.8	10.3	5.5-17.5							1.0	2.9	0.9-7.4
	1996-2000							2.4	9.4	4.8-16.8							1.2	3.9	1.4-9.0
	2001-2005							3.0	12.6	6.8-21.2							1.2	3.0	1.1-7.5
	2006-2010							1.4	5.5	2.1-11.7							0.2	0.5	0.0-3.9
	2011-2015							1.0	3.4	1.1-8.6							0.2	0.4	0.0-4.0
	2016-2020							2.0	11.1	5.0-20.8							0.2	0.4	0.0-4.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Delaware**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							9.0	32.5	23.4-43.9							2.8	9.4	5.1-16.0
	1981-1985							7.6	27.7	19.3-38.7							1.6	5.1	2.2-10.4
	1986-1990							6.2	23.4	15.6-33.8							1.2	4.0	1.4-9.2
	1991-1995							7.4	23.8	16.5-33.3							1.8	5.1	2.3-10.1
	1996-2000							8.6	25.8	18.5-35.3							1.8	4.0	1.8-8.4
	2001-2005							11.2	32.2	24.2-42.4							2.8	7.5	4.0-13.3
	2006-2010							10.8	33.9	25.0-45.2							0.8	1.9	0.5-5.9
	2011-2015							13.2	36.8	27.8-48.1							1.0	2.5	0.8-6.9
	2016-2020							10.6	30.5	22.2-41.4							2.8	7.4	3.8-13.7
Ovary	1976-1980							3.8	15.5	9.1-24.5							3.0	12.0	6.6-20.1
	1981-1985							4.0	15.0	8.9-23.8							3.4	11.0	6.3-18.0
	1986-1990							4.2	14.1	8.5-22.0							4.4	14.0	8.6-21.7
	1991-1995							4.6	13.4	8.4-20.5							3.0	8.8	4.8-15.0
	1996-2000							4.2	12.4	7.5-19.6							3.8	10.9	6.5-17.6
	2001-2005							6.8	20.7	14.2-29.4							4.2	11.7	7.2-18.4
	2006-2010							3.8	10.3	6.1-16.8							3.6	9.0	5.3-14.9
	2011-2015							6.6	18.8	12.5-27.6							4.4	11.4	6.9-18.4
	2016-2020							3.6	8.9	5.2-15.2							3.0	6.7	3.7-12.2
Prostate	1976-1980				18.4	87.0	69.7-107.4							5.0	24.8	15.8-36.9			
	1981-1985				22.4	99.6	81.5-120.5							8.2	38.9	27.6-53.2			
	1986-1990				29.6	125.7	105.5-148.6							9.4	43.9	31.8-58.9			
	1991-1995				33.0	120.2	102.1-140.7							9.4	42.1	30.5-56.4			
	1996-2000				35.8	121.4	104.0-141.0							7.8	30.9	21.7-42.6			
	2001-2005				43.8	137.0	119.3-156.9							8.0	29.7	21.1-40.8			
	2006-2010				41.8	120.2	104.2-138.3							4.8	15.7	10.0-23.9			
	2011-2015				32.4	85.3	72.4-100.3							5.8	18.0	12.0-26.4			
	2016-2020				34.6	83.5	71.2-97.9							6.0	16.1	10.8-23.8			
Testis	1976-1980				0.6	2.5	0.5-7.7							0.0	0.0	0.0-3.6			
	1981-1985				0.2	0.9	0.0-5.2							0.0	0.0	0.0-3.5			
	1986-1990				2.2	10.7	5.3-19.1							0.2	0.8	0.0-4.8			
	1991-1995				1.8	8.6	3.9-16.1							0.0	0.0	0.0-3.2			
	1996-2000				1.0	4.3	1.4-10.2							0.0	0.0	0.0-3.2			
	2001-2005				1.0	4.6	1.5-10.8							0.0	0.0	0.0-3.3			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Delaware**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				1.0	4.0	1.2-9.7							0.0	0.0	0.0-3.4			
	2011-2015				1.4	7.1	2.7-14.9							0.0	0.0	0.0-3.6			
	2016-2020				1.8	10.8	4.9-20.4							0.0	0.0	0.0-3.8			
Urinary bladder (incl. in situ)	1976-1980	13.2	25.3	19.5-32.3	11.0	47.5	35.5-62.3	2.2	8.5	4.1-15.5	2.8	5.6	3.0-9.5	2.2	10.4	5.1-18.8	0.6	2.0	0.4-6.3
	1981-1985	12.0	21.9	16.7-28.4	9.6	40.2	29.4-53.9	2.4	7.6	3.9-13.6	4.0	7.7	4.7-12.0	2.6	12.9	6.7-22.3	1.4	4.9	1.9-10.4
	1986-1990	12.4	21.8	16.6-28.0	9.0	38.0	27.4-51.3	3.4	11.0	6.3-18.1	2.6	4.7	2.5-8.1	2.2	12.1	5.9-21.7	0.4	1.2	0.1-5.0
	1991-1995	13.6	21.7	16.8-27.7	10.6	38.5	28.6-50.8	3.0	8.1	4.5-13.7	4.4	7.1	4.4-10.9	3.8	14.7	8.8-23.3	0.6	1.5	0.3-5.1
	1996-2000	18.4	28.0	22.5-34.6	12.2	43.7	33.2-56.6	6.2	17.0	11.5-24.8	4.6	6.7	4.2-10.2	3.0	11.8	6.6-19.8	1.6	3.8	1.6-8.3
	2001-2005	22.2	32.8	26.9-39.7	17.2	55.8	44.5-69.5	5.0	13.6	8.7-20.6	4.8	6.8	4.4-10.4	3.6	11.9	7.0-19.3	1.2	2.8	1.0-7.0
	2006-2010	19.8	28.0	22.6-34.3	15.6	48.7	38.3-61.3	4.2	12.3	7.4-19.5	4.6	6.4	4.0-9.9	3.4	10.8	6.2-17.7	1.2	3.0	1.1-7.5
	2011-2015	19.2	26.1	21.0-32.3	14.6	42.2	32.8-54.0	4.6	11.5	7.2-18.1	4.0	5.4	3.3-8.7	3.0	8.5	4.7-14.8	1.0	2.6	0.8-7.3
2016-2020	20.6	24.6	20.0-30.2	15.0	37.1	29.0-47.4	5.6	12.3	8.2-18.8	3.8	4.5	2.7-7.6	3.2	8.2	4.7-14.3	0.6	1.3	0.3-5.5	
Kidney and renal pelvis	1976-1980	3.4	6.8	3.9-11.1	1.6	6.9	2.9-13.9	1.8	6.4	2.9-12.5	2.0	3.7	1.7-7.0	1.2	4.7	1.7-10.7	0.8	2.8	0.7-7.5
	1981-1985	3.2	6.5	3.7-10.7	1.6	7.9	3.3-15.6	1.6	5.5	2.3-11.2	2.2	4.0	2.0-7.3	1.6	7.4	3.1-14.8	0.6	1.8	0.4-5.9
	1986-1990	4.6	8.8	5.6-13.4	3.4	15.3	8.8-24.7	1.2	4.3	1.5-9.7	2.6	4.8	2.5-8.3	1.6	6.5	2.8-13.1	1.0	3.3	1.0-8.2
	1991-1995	4.6	8.4	5.3-12.7	2.4	10.0	5.1-17.6	2.2	6.9	3.3-12.9	2.4	4.2	2.1-7.5	1.2	5.5	2.0-11.9	1.2	3.4	1.2-8.1
	1996-2000	6.8	11.1	7.6-15.7	4.0	14.7	8.9-23.0	2.8	8.2	4.4-14.4	2.8	4.6	2.5-7.8	1.6	6.2	2.6-12.4	1.2	3.2	1.1-7.7
	2001-2005	7.6	12.2	8.6-17.1	3.8	12.7	7.5-20.3	3.8	11.4	6.6-18.6	2.0	3.1	1.5-5.9	1.2	3.9	1.4-8.9	0.8	2.1	0.5-6.2
	2006-2010	9.2	12.5	9.1-17.0	6.6	18.7	12.8-26.8	2.6	6.9	3.6-12.6	1.6	2.2	1.0-4.7	0.8	2.4	0.6-6.8	0.8	1.8	0.5-5.7
	2011-2015	11.2	16.6	12.2-22.1	6.2	18.7	12.3-27.7	5.0	14.1	8.7-21.8	2.4	2.9	1.5-5.5	1.4	3.3	1.3-7.9	1.0	2.2	0.7-6.4
2016-2020	10.8	15.9	11.4-21.6	6.4	18.7	12.2-27.8	4.4	13.3	7.7-21.8	2.0	2.8	1.3-5.8	0.8	2.0	0.5-6.4	1.2	3.7	1.2-9.3	
Brain and other central nervous system	1976-1980	3.2	6.8	3.8-11.1	1.6	6.8	2.8-13.7	1.6	6.5	2.8-13.0	1.8	3.3	1.5-6.5	1.2	4.7	1.7-10.5	0.6	2.0	0.4-6.4
	1981-1985	2.6	4.9	2.5-8.5	1.6	7.1	2.9-14.3	1.0	3.6	1.2-8.8	1.8	3.5	1.6-6.8	1.0	4.0	1.3-9.7	0.8	3.3	0.9-8.6
	1986-1990	2.8	5.2	2.8-8.8	1.8	6.3	2.9-12.4	1.0	4.0	1.2-9.5	2.6	5.1	2.6-9.0	1.8	7.4	3.3-14.5	0.8	3.1	0.8-8.2
	1991-1995	2.8	5.1	2.8-8.7	1.4	5.8	2.3-12.1	1.4	4.7	1.8-10.1	2.0	3.5	1.6-6.6	0.6	2.5	0.5-7.5	1.4	4.5	1.7-9.7
	1996-2000	5.4	8.9	5.8-13.2	4.0	13.2	8.0-20.8	1.4	5.4	2.1-11.4	3.2	5.0	2.8-8.3	2.0	6.3	3.0-12.0	1.2	3.9	1.4-8.9
	2001-2005	3.8	6.6	3.9-10.6	2.2	8.3	4.0-15.2	1.6	4.9	2.1-10.3	2.4	3.7	1.9-6.8	1.6	5.4	2.3-11.3	0.8	2.1	0.6-6.3
	2006-2010	4.8	8.4	5.2-12.9	2.2	8.5	4.0-15.8	2.6	8.4	4.3-15.0	2.6	3.9	2.0-7.0	1.2	3.9	1.3-9.2	1.4	3.7	1.5-8.4
	2011-2015	4.2	7.7	4.6-12.2	1.6	5.5	2.1-11.8	2.6	10.0	5.1-17.8	2.8	4.8	2.5-8.6	1.6	6.2	2.4-13.1	1.2	3.3	1.1-8.2
2016-2020	2.6	5.7	2.7-10.4	1.2	5.0	1.5-12.0	1.4	6.2	2.1-14.0	2.2	2.9	1.4-5.8	0.8	1.8	0.5-6.1	1.4	4.0	1.5-9.4	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Delaware**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	2.0	3.8	1.8-7.1	1.2	5.1	1.8-11.4	0.8	2.6	0.7-7.2	0.2	0.3	0.0-2.1	0.0	0.0	0.0-3.6	0.2	0.6	0.0-4.0
	1981-1985	0.8	1.4	0.4-3.7	0.0	0.0	0.0-3.5	0.8	2.7	0.7-7.4	0.2	0.3	0.0-2.1	0.2	0.6	0.0-4.5	0.0	0.0	0.0-3.2
	1986-1990	1.6	3.5	1.5-6.9	0.4	1.5	0.2-5.9	1.2	5.4	1.9-11.7	0.0	0.0	0.0-1.6	0.0	0.0	0.0-3.4	0.0	0.0	0.0-3.1
	1991-1995	1.2	2.1	0.7-4.6	0.6	2.1	0.4-6.7	0.6	2.2	0.4-6.6	0.2	0.3	0.0-2.0	0.0	0.0	0.0-3.2	0.2	0.5	0.0-3.7
	1996-2000	2.6	5.0	2.6-8.6	0.4	1.3	0.2-5.2	2.2	8.6	4.2-15.7	0.4	0.6	0.1-2.4	0.0	0.0	0.0-3.2	0.4	0.8	0.1-4.1
	2001-2005	4.8	9.4	6.0-14.2	1.2	4.7	1.6-10.6	3.6	14.1	8.2-22.6	0.2	0.3	0.0-2.1	0.0	0.0	0.0-3.3	0.2	0.6	0.0-4.1
	2006-2010	7.8	15.2	10.5-21.2	1.2	4.8	1.6-11.0	6.6	25.3	16.9-36.4	0.8	1.1	0.3-3.1	0.0	0.0	0.0-3.4	0.8	1.7	0.4-5.6
	2011-2015	8.4	17.2	12.0-23.9	2.8	10.9	5.5-19.3	5.6	23.9	15.2-35.7	0.0	0.0	0.0-1.8	0.0	0.0	0.0-3.6	0.0	0.0	0.0-3.6
	2016-2020	6.8	13.1	8.5-19.2	2.0	5.7	2.6-11.5	4.8	20.4	12.2-31.9	0.8	1.0	0.3-3.3	0.6	1.7	0.3-6.1	0.2	0.5	0.0-4.6
Hodgkin lymphoma	1976-1980	1.6	3.3	1.4-6.7	0.6	3.0	0.6-8.8	1.0	3.5	1.1-8.5	0.6	1.5	0.3-4.4	0.4	2.3	0.2-8.0	0.2	0.8	0.0-4.6
	1981-1985	1.6	3.2	1.4-6.6	0.2	1.1	0.0-5.7	1.4	5.5	2.1-11.5	0.0	0.0	0.0-1.6	0.0	0.0	0.0-3.5	0.0	0.0	0.0-3.2
	1986-1990	2.8	6.0	3.2-10.1	1.8	8.0	3.6-15.3	1.0	3.9	1.2-9.5	0.4	0.8	0.1-2.9	0.2	0.9	0.0-5.1	0.2	0.7	0.0-4.4
	1991-1995	1.4	3.1	1.2-6.3	1.2	5.6	2.0-12.0	0.2	0.6	0.0-3.9	0.4	0.6	0.1-2.5	0.0	0.0	0.0-3.2	0.4	1.1	0.1-4.7
	1996-2000	1.4	3.0	1.2-6.3	1.4	6.1	2.4-12.8	0.0	0.0	0.0-3.1	0.2	0.5	0.0-2.6	0.2	1.0	0.0-5.3	0.0	0.0	0.0-3.1
	2001-2005	2.0	4.1	1.9-7.8	1.2	4.9	1.7-11.0	0.8	3.5	0.9-9.0	0.0	0.0	0.0-1.6	0.0	0.0	0.0-3.3	0.0	0.0	0.0-3.3
	2006-2010	1.6	3.2	1.3-6.5	1.2	3.9	1.4-9.1	0.4	2.4	0.3-7.9	0.4	0.6	0.1-2.5	0.2	0.5	0.0-4.1	0.2	0.6	0.0-4.2
	2011-2015	1.4	2.3	0.8-5.2	0.4	1.0	0.1-5.0	1.0	3.7	1.1-9.5	0.6	1.0	0.2-3.5	0.0	0.0	0.0-3.6	0.6	2.0	0.3-6.9
	2016-2020	2.0	4.3	1.9-8.3	1.2	5.4	2.0-12.1	0.8	2.9	0.6-8.7	0.2	0.2	0.0-2.2	0.0	0.0	0.0-3.8	0.2	0.4	0.0-4.5
Non-Hodgkin lymphomas	1976-1980	5.2	11.0	7.0-16.3	2.8	14.3	7.7-24.1	2.4	9.5	4.6-17.1	3.2	7.2	4.0-11.8	2.2	11.5	5.6-20.6	1.0	3.8	1.2-9.1
	1981-1985	7.0	13.4	9.3-18.9	4.4	20.0	12.4-30.7	2.6	9.2	4.8-16.3	5.2	10.2	6.5-15.1	2.6	11.2	5.8-19.6	2.6	9.4	4.8-16.7
	1986-1990	8.4	15.6	11.2-21.3	5.8	25.7	17.0-37.3	2.6	7.7	4.1-13.6	2.4	4.2	2.2-7.4	1.6	6.9	2.9-13.8	0.8	2.4	0.7-6.7
	1991-1995	10.0	18.8	13.9-25.0	5.8	23.9	15.9-34.5	4.2	14.1	8.6-22.0	3.2	5.5	3.1-9.0	1.8	7.3	3.3-14.1	1.4	4.3	1.7-9.4
	1996-2000	11.0	17.2	12.9-22.7	6.2	22.4	15.1-32.1	4.8	13.3	8.4-20.5	6.6	10.2	7.0-14.5	2.8	10.3	5.6-17.6	3.8	10.1	5.9-16.5
	2001-2005	14.8	21.8	17.1-27.6	8.4	27.2	19.5-37.2	6.4	17.4	11.8-25.1	3.8	5.4	3.3-8.7	2.2	6.5	3.2-12.1	1.6	4.1	1.7-8.9
	2006-2010	13.0	20.5	15.5-26.6	7.4	24.2	16.7-34.0	5.6	17.1	10.9-25.7	4.6	6.9	4.3-10.7	2.8	9.2	5.0-15.8	1.8	5.1	2.2-10.5
	2011-2015	14.0	19.5	15.0-25.1	6.4	18.6	12.4-27.4	7.6	19.9	13.9-28.1	4.0	5.2	3.2-8.4	1.8	5.2	2.4-10.7	2.2	5.1	2.5-10.1
	2016-2020	15.8	20.9	16.3-26.8	9.2	26.2	18.5-36.4	6.6	15.7	10.7-23.0	4.2	5.2	3.2-8.4	3.4	8.5	4.9-14.6	0.8	1.9	0.5-6.2
Myeloma	1976-1980	2.4	4.5	2.3-8.0	0.6	3.0	0.6-8.9	1.8	5.9	2.7-11.5	1.0	1.9	0.6-4.7	0.4	2.5	0.3-8.5	0.6	1.9	0.4-6.1
	1981-1985	2.2	4.1	2.1-7.5	1.6	7.3	3.1-14.6	0.6	1.8	0.4-6.0	1.4	2.6	1.0-5.5	0.8	4.0	1.1-10.3	0.6	1.7	0.4-5.7
	1986-1990	2.4	4.1	2.1-7.4	1.0	3.7	1.2-9.0	1.4	4.6	1.8-9.9	3.2	6.2	3.5-10.2	2.2	10.5	5.1-19.1	1.0	3.3	1.0-8.2
	1991-1995	4.0	6.4	3.9-10.1	1.6	5.9	2.5-11.9	2.4	6.5	3.3-11.8	2.0	3.2	1.5-6.0	0.8	3.3	0.8-8.7	1.2	3.3	1.2-7.6
	1996-2000	1.8	2.8	1.3-5.6	1.0	3.3	1.1-8.2	0.8	2.4	0.6-6.8	1.2	1.9	0.7-4.3	0.8	2.9	0.8-7.7	0.4	1.2	0.1-5.1
	2001-2005	0.8	1.1	0.3-3.2	0.2	0.7	0.0-4.4	0.6	1.3	0.3-4.9	0.4	0.7	0.1-2.9	0.4	1.5	0.2-5.9	0.0	0.0	0.0-3.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Delaware**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	4.2	6.3	3.8-9.9	2.8	9.2	4.9-16.0	1.4	3.8	1.5-8.6	1.8	2.5	1.2-5.1	1.2	3.7	1.3-8.6	0.6	1.7	0.4-5.8
	2011-2015	5.4	7.1	4.6-10.7	2.8	7.9	4.2-14.0	2.6	6.2	3.3-11.5	2.0	2.8	1.3-5.5	0.6	2.0	0.4-6.5	1.4	3.6	1.4-8.4
	2016-2020	6.0	7.5	5.0-11.3	4.0	10.4	6.2-17.2	2.0	4.5	2.1-9.5	2.2	2.8	1.3-5.7	1.4	3.9	1.4-9.2	0.8	1.6	0.4-5.8
Leukemias	1976-1980	6.8	13.9	9.6-19.5	4.4	21.2	13.1-32.4	2.4	8.8	4.5-15.7	3.6	7.3	4.3-11.6	2.4	12.0	6.1-21.0	1.2	4.2	1.5-9.4
	1981-1985	7.0	13.5	9.3-19.0	3.0	13.4	7.4-22.4	4.0	13.7	8.2-21.7	3.2	6.3	3.6-10.4	1.6	7.8	3.3-15.5	1.6	5.3	2.2-11.0
	1986-1990	8.4	16.4	11.8-22.3	5.0	23.5	15.0-35.0	3.4	12.0	6.9-19.6	4.6	8.5	5.4-12.9	2.6	11.8	6.2-20.5	2.0	6.9	3.2-13.1
	1991-1995	8.2	14.3	10.2-19.5	5.0	21.5	13.7-32.1	3.2	10.1	5.6-16.9	6.6	11.3	7.7-16.0	4.0	18.5	11.1-28.8	2.6	7.0	3.7-12.5
	1996-2000	12.0	18.7	14.2-24.3	5.2	18.3	11.9-27.1	6.8	19.4	13.2-27.8	5.0	7.4	4.8-11.1	2.6	9.4	4.9-16.4	2.4	5.8	3.0-10.9
	2001-2005	12.6	19.5	14.9-25.2	6.8	22.5	15.4-32.0	5.8	17.4	11.4-25.6	7.0	10.6	7.3-15.1	3.0	10.6	5.8-18.1	4.0	11.1	6.6-17.9
	2006-2010	11.4	16.4	12.3-21.5	7.6	23.0	16.1-32.2	3.8	9.9	5.9-16.1	4.8	6.3	4.1-9.7	3.2	9.0	5.1-15.2	1.6	4.2	1.8-9.0
	2011-2015	11.6	18.1	13.3-24.0	6.6	21.7	14.4-31.5	5.0	15.1	9.3-23.4	4.8	7.2	4.4-11.2	3.0	9.5	5.1-16.5	1.8	4.9	2.0-10.6
	2016-2020	13.6	19.3	14.5-25.4	7.4	22.3	15.0-32.4	6.2	16.3	10.5-24.5	6.4	8.6	5.6-12.8	3.8	11.1	6.2-18.7	2.6	5.9	3.1-11.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Dutchess**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	810.2	387.1	375.1-399.4	406.6	457.5	436.8-478.9	403.6	349.4	334.1-365.2	412.2	199.0	190.4-208.0	215.0	247.2	231.8-263.3	197.2	169.9	159.3-180.9
	1981-1985	945.0	423.3	411.2-435.7	477.8	514.9	493.4-537.0	467.2	376.2	360.9-392.0	454.8	206.3	197.8-215.0	238.2	264.7	249.2-280.9	216.6	170.8	160.7-181.4
	1986-1990	1,056.2	454.0	441.7-466.5	515.8	532.8	511.5-554.7	540.4	416.9	401.1-433.0	501.0	217.9	209.4-226.7	263.8	285.2	269.3-301.7	237.2	178.6	168.5-189.1
	1991-1995	1,179.4	479.2	467.0-491.6	608.6	582.4	561.1-604.2	570.8	419.7	404.4-435.5	529.0	218.7	210.4-227.2	267.4	273.2	258.2-288.9	261.6	187.2	177.1-197.7
	1996-2000	1,319.6	495.1	483.2-507.2	675.0	577.9	558.1-598.3	644.6	441.4	426.2-456.9	555.2	212.0	204.2-220.1	282.0	261.8	247.9-276.3	273.2	182.2	172.6-192.1
	2001-2005	1,461.0	493.9	482.6-505.4	753.6	569.4	550.9-588.3	707.4	442.6	428.0-457.5	543.6	187.4	180.4-194.6	277.4	228.3	216.2-241.0	266.2	162.2	153.6-171.2
	2006-2010	1,614.8	494.5	483.7-505.5	817.2	550.5	533.4-568.0	797.6	459.5	445.1-474.2	550.8	170.1	163.7-176.6	278.4	199.5	189.0-210.5	272.4	151.0	143.0-159.4
	2011-2015	1,734.0	478.4	468.1-488.8	860.8	509.1	493.6-524.9	873.2	461.7	447.7-476.1	546.6	150.3	144.6-156.1	288.6	181.8	172.3-191.6	258.0	128.1	121.1-135.5
2016-2020	1,782.0	453.2	443.5-463.1	929.8	493.9	479.3-508.8	852.2	424.5	411.2-438.1	540.0	133.3	128.2-138.5	274.8	150.9	142.9-159.3	265.2	120.5	113.9-127.4	
Oral cavity and pharynx	1976-1980	23.0	10.6	8.7-12.7	16.4	16.1	12.7-20.0	6.6	5.6	3.9-7.9	10.8	5.1	3.8-6.6	7.0	7.3	5.1-10.2	3.8	3.1	1.8-4.8
	1981-1985	26.6	11.8	9.9-14.0	18.8	19.3	15.4-23.8	7.8	6.2	4.4-8.5	10.0	4.5	3.3-5.9	7.0	7.3	5.0-10.2	3.0	2.3	1.3-3.9
	1986-1990	27.2	11.4	9.6-13.6	17.2	16.1	12.8-20.0	10.0	7.9	5.8-10.4	13.0	5.5	4.2-7.0	7.8	7.5	5.3-10.3	5.2	3.9	2.6-5.8
	1991-1995	24.6	9.9	8.2-11.8	16.0	14.4	11.3-18.0	8.6	6.1	4.4-8.3	10.2	4.1	3.1-5.4	7.2	6.6	4.5-9.2	3.0	2.2	1.2-3.7
	1996-2000	28.6	10.6	9.0-12.5	19.4	15.9	12.9-19.5	9.2	6.3	4.6-8.4	10.0	3.7	2.8-4.9	7.2	6.0	4.2-8.4	2.8	1.9	1.0-3.2
	2001-2005	31.6	10.5	8.9-12.2	21.8	15.3	12.5-18.6	9.8	6.2	4.6-8.3	7.8	2.6	1.9-3.6	5.8	4.5	3.0-6.5	2.0	1.2	0.6-2.3
	2006-2010	29.4	8.8	7.4-10.4	21.2	13.4	10.9-16.3	8.2	4.7	3.4-6.4	6.4	2.0	1.4-2.8	4.8	3.4	2.1-5.1	1.6	1.0	0.4-1.9
	2011-2015	37.0	9.9	8.5-11.5	25.8	14.4	12.0-17.2	11.2	6.0	4.5-7.9	5.2	1.5	0.9-2.2	4.0	2.4	1.5-3.8	1.2	0.7	0.2-1.6
2016-2020	48.2	12.1	10.6-13.8	34.8	17.8	15.2-20.9	13.4	6.8	5.2-8.8	8.4	2.0	1.4-2.8	6.2	3.1	2.1-4.5	2.2	1.0	0.5-1.9	
Esophagus	1976-1980	9.8	4.7	3.5-6.3	6.6	7.4	5.0-10.6	3.2	2.8	1.6-4.5	8.2	4.1	2.9-5.5	5.8	6.9	4.5-10.0	2.4	2.1	1.1-3.7
	1981-1985	9.4	4.2	3.1-5.6	7.4	7.6	5.3-10.6	2.0	1.7	0.8-3.1	7.2	3.2	2.2-4.4	6.0	6.3	4.1-9.1	1.2	1.1	0.4-2.3
	1986-1990	12.0	5.3	4.0-6.8	7.8	8.1	5.7-11.2	4.2	3.3	2.0-5.0	11.8	5.2	3.9-6.7	7.8	7.9	5.5-10.9	4.0	3.0	1.8-4.7
	1991-1995	11.4	4.7	3.5-6.1	8.2	7.8	5.5-10.7	3.2	2.2	1.2-3.6	10.8	4.4	3.3-5.8	8.0	7.6	5.3-10.4	2.8	2.0	1.1-3.4
	1996-2000	10.8	4.1	3.1-5.3	6.8	6.0	4.1-8.4	4.0	2.6	1.6-4.1	10.8	4.1	3.1-5.3	7.2	6.3	4.4-8.8	3.6	2.3	1.4-3.7
	2001-2005	19.0	6.5	5.3-8.0	13.6	10.4	8.0-13.2	5.4	3.3	2.2-4.8	15.0	5.1	4.0-6.4	11.6	9.0	6.8-11.7	3.4	2.1	1.2-3.3
	2006-2010	16.2	4.8	3.8-6.0	12.6	8.2	6.2-10.5	3.6	1.9	1.1-3.1	13.8	4.2	3.3-5.3	11.4	7.6	5.7-9.9	2.4	1.4	0.7-2.5
	2011-2015	18.0	4.8	3.9-6.0	15.2	8.9	6.9-11.2	2.8	1.4	0.8-2.4	15.6	4.3	3.4-5.4	12.6	7.8	6.0-10.1	3.0	1.5	0.8-2.5
2016-2020	20.0	5.1	4.1-6.3	14.6	8.2	6.4-10.4	5.4	2.5	1.6-3.7	14.8	3.6	2.8-4.6	10.6	5.9	4.4-7.8	4.2	1.8	1.1-2.8	
Stomach	1976-1980	22.4	11.1	9.1-13.4	12.8	15.5	11.8-20.0	9.6	8.2	6.0-10.9	17.0	8.6	6.8-10.7	9.6	12.8	9.2-17.1	7.4	6.3	4.4-8.8
	1981-1985	24.8	11.4	9.4-13.6	16.8	19.2	15.1-24.0	8.0	6.3	4.5-8.6	16.8	7.7	6.1-9.5	10.8	12.4	9.2-16.4	6.0	4.8	3.2-6.8
	1986-1990	26.6	11.7	9.8-13.9	16.2	17.2	13.5-21.6	10.4	7.7	5.7-10.1	16.2	7.3	5.8-9.0	10.2	11.9	8.8-15.8	6.0	4.3	2.9-6.2
	1991-1995	21.2	8.8	7.2-10.6	12.8	13.0	9.9-16.7	8.4	5.8	4.2-7.9	14.0	5.8	4.5-7.3	7.6	7.6	5.3-10.5	6.4	4.5	3.1-6.4
	1996-2000	23.8	9.0	7.5-10.8	15.6	14.4	11.2-18.0	8.2	5.4	3.9-7.4	15.4	5.9	4.7-7.4	9.8	9.2	6.7-12.3	5.6	3.6	2.4-5.2
	2001-2005	22.0	7.5	6.2-9.1	13.6	10.4	8.0-13.3	8.4	5.2	3.8-7.1	11.0	3.8	2.9-5.0	6.6	5.7	3.9-8.0	4.4	2.7	1.7-4.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Dutchess**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	21.2	6.5	5.3-7.8	13.6	9.2	7.1-11.7	7.6	4.4	3.1-6.1	10.6	3.2	2.4-4.2	7.0	4.7	3.3-6.7	3.6	2.0	1.2-3.2
	2011-2015	24.8	6.7	5.6-8.1	14.0	8.3	6.4-10.5	10.8	5.5	4.1-7.3	9.8	2.7	2.0-3.6	6.2	3.7	2.5-5.4	3.6	1.8	1.1-3.0
	2016-2020	25.6	6.6	5.4-7.9	16.4	9.0	7.1-11.3	9.2	4.5	3.2-6.1	10.0	2.5	1.9-3.4	6.2	3.2	2.2-4.7	3.8	2.0	1.2-3.2
Colorectal	1976-1980	131.0	63.8	58.9-69.0	61.8	70.9	62.8-79.6	69.2	58.6	52.5-65.2	61.8	30.8	27.4-34.4	30.0	36.1	30.2-42.7	31.8	27.3	23.2-32.0
	1981-1985	144.8	66.2	61.4-71.2	74.8	84.8	76.0-94.3	70.0	54.3	48.7-60.3	61.6	28.2	25.1-31.5	33.0	37.9	32.1-44.4	28.6	22.1	18.6-26.0
	1986-1990	155.8	68.8	64.0-73.9	76.4	82.9	74.4-92.0	79.4	59.5	53.7-65.7	67.0	30.0	26.8-33.4	34.6	40.1	34.0-46.7	32.4	23.9	20.3-27.9
	1991-1995	152.4	63.0	58.6-67.7	75.4	73.0	65.6-81.0	77.0	55.0	49.6-60.8	67.8	28.4	25.4-31.6	32.0	33.2	28.0-39.0	35.8	25.0	21.4-29.0
	1996-2000	151.6	57.9	53.8-62.1	73.8	65.1	58.5-72.3	77.8	51.9	46.8-57.3	61.8	23.8	21.2-26.6	31.8	29.0	24.5-34.0	30.0	19.4	16.4-22.8
	2001-2005	151.2	51.6	48.0-55.4	78.2	61.1	55.0-67.6	73.0	44.3	39.9-49.2	54.4	18.7	16.6-21.1	25.2	20.5	17.0-24.6	29.2	17.2	14.5-20.3
	2006-2010	156.4	48.2	44.9-51.8	80.6	56.3	50.8-62.2	75.8	42.4	38.2-47.0	54.6	17.2	15.2-19.4	28.2	20.9	17.6-24.7	26.4	14.2	11.9-16.9
	2011-2015	135.4	37.4	34.6-40.4	69.0	41.4	37.1-46.2	66.4	34.1	30.5-38.1	47.8	12.9	11.3-14.7	23.4	14.3	11.8-17.2	24.4	11.7	9.7-14.1
2016-2020	130.4	33.8	31.1-36.6	69.2	38.0	34.0-42.4	61.2	30.1	26.6-33.9	40.4	9.9	8.6-11.4	20.6	11.2	9.1-13.7	19.8	8.8	7.1-10.8	
Colon excluding rectum	1976-1980	90.8	44.6	40.5-48.9	40.2	46.5	40.0-53.7	50.6	42.9	37.8-48.6	49.0	24.6	21.5-27.9	23.8	28.2	23.0-34.0	25.2	21.9	18.2-26.1
	1981-1985	101.0	46.2	42.2-50.4	50.8	57.5	50.3-65.4	50.2	38.9	34.2-44.0	49.6	22.5	19.8-25.5	26.4	30.1	24.9-35.9	23.2	17.8	14.7-21.4
	1986-1990	114.4	50.6	46.5-55.0	54.8	60.7	53.4-68.7	59.6	44.6	39.7-50.0	57.4	25.6	22.7-28.8	29.6	34.1	28.6-40.3	27.8	20.5	17.2-24.2
	1991-1995	114.8	47.7	43.9-51.8	54.8	53.9	47.5-60.9	60.0	42.7	38.0-47.9	58.0	24.3	21.6-27.3	26.6	27.6	22.9-32.9	31.4	22.0	18.6-25.7
	1996-2000	111.6	42.7	39.3-46.4	52.6	47.2	41.5-53.4	59.0	39.2	34.8-44.0	51.0	19.6	17.3-22.2	25.8	23.9	19.8-28.5	25.2	16.3	13.6-19.5
	2001-2005	111.6	38.2	35.1-41.6	56.4	45.0	39.8-50.7	55.2	33.5	29.7-37.8	45.4	15.7	13.7-17.9	20.6	17.0	13.8-20.7	24.8	14.6	12.2-17.5
	2006-2010	117.4	36.3	33.4-39.4	58.0	40.8	36.2-45.9	59.4	33.5	29.7-37.6	45.6	14.4	12.6-16.4	23.4	17.6	14.5-21.1	22.2	11.9	9.8-14.5
	2011-2015	96.2	26.7	24.3-29.2	46.0	28.2	24.6-32.2	50.2	25.7	22.6-29.2	40.0	10.9	9.4-12.5	19.2	12.0	9.6-14.7	20.8	10.2	8.3-12.4
2016-2020	94.2	24.4	22.1-26.8	48.0	26.3	23.0-30.0	46.2	22.7	19.7-26.0	30.2	7.4	6.3-8.7	14.6	8.1	6.3-10.3	15.6	6.8	5.4-8.6	
Rectum & rectosigmoid	1976-1980	40.2	19.2	16.6-22.1	21.6	24.4	19.8-29.6	18.6	15.6	12.6-19.2	12.8	6.2	4.7-7.9	6.2	7.9	5.2-11.4	6.6	5.5	3.8-7.7
	1981-1985	43.8	20.0	17.4-22.8	24.0	27.3	22.4-32.8	19.8	15.4	12.5-18.8	12.0	5.7	4.3-7.3	6.6	7.9	5.3-11.1	5.4	4.2	2.8-6.2
	1986-1990	41.4	18.2	15.8-20.9	21.6	22.1	18.0-26.9	19.8	14.9	12.1-18.1	9.6	4.3	3.2-5.7	5.0	6.0	3.8-8.9	4.6	3.4	2.1-5.1
	1991-1995	37.6	15.3	13.2-17.7	20.6	19.1	15.5-23.2	17.0	12.3	9.8-15.2	9.8	4.0	3.0-5.3	5.4	5.6	3.6-8.2	4.4	3.0	1.9-4.6
	1996-2000	40.0	15.1	13.1-17.4	21.2	17.9	14.6-21.7	18.8	12.7	10.2-15.6	10.8	4.2	3.1-5.4	6.0	5.1	3.4-7.3	4.8	3.0	1.9-4.6
	2001-2005	39.6	13.4	11.6-15.4	21.8	16.1	13.1-19.5	17.8	10.8	8.7-13.4	9.0	3.0	2.2-4.1	4.6	3.6	2.2-5.4	4.4	2.6	1.6-3.9
	2006-2010	39.0	11.9	10.3-13.7	22.6	15.4	12.7-18.6	16.4	9.0	7.1-11.2	9.0	2.8	2.0-3.8	4.8	3.3	2.1-5.0	4.2	2.3	1.4-3.5
	2011-2015	39.2	10.7	9.3-12.4	23.0	13.3	10.9-16.0	16.2	8.4	6.6-10.5	7.8	2.0	1.4-2.8	4.2	2.3	1.4-3.7	3.6	1.5	0.9-2.5
2016-2020	36.2	9.4	8.0-10.9	21.2	11.8	9.6-14.4	15.0	7.4	5.7-9.4	10.2	2.5	1.8-3.3	6.0	3.1	2.1-4.6	4.2	2.0	1.2-3.1	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Dutchess**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	5.2	2.5	1.6-3.7	2.8	3.0	1.6-5.0	2.4	2.1	1.1-3.6	3.4	1.7	1.0-2.7	2.0	2.1	1.0-3.8	1.4	1.3	0.5-2.6
	1981-1985	6.8	3.1	2.1-4.3	5.2	5.5	3.5-8.0	1.6	1.3	0.5-2.6	4.4	1.9	1.2-2.9	3.2	3.5	2.0-5.8	1.2	0.9	0.3-1.9
	1986-1990	8.4	3.6	2.6-4.9	5.4	5.0	3.3-7.4	3.0	2.2	1.2-3.7	6.0	2.5	1.7-3.6	4.4	4.0	2.5-6.1	1.6	1.2	0.5-2.4
	1991-1995	8.6	3.6	2.6-4.8	5.2	4.8	3.1-7.1	3.4	2.4	1.4-3.8	8.8	3.7	2.6-4.9	4.6	4.6	2.9-6.9	4.2	2.9	1.8-4.5
	1996-2000	11.6	4.4	3.3-5.7	7.2	6.0	4.2-8.3	4.4	2.9	1.8-4.4	9.4	3.5	2.6-4.7	5.8	4.6	3.1-6.7	3.6	2.4	1.4-3.8
	2001-2005	19.0	6.3	5.1-7.7	13.0	9.3	7.1-11.9	6.0	3.7	2.5-5.3	13.2	4.5	3.5-5.7	8.2	6.3	4.5-8.6	5.0	3.0	2.0-4.5
	2006-2010	26.8	7.8	6.6-9.3	20.4	12.2	9.9-14.9	6.4	3.6	2.5-5.2	19.2	5.8	4.7-7.2	12.6	8.1	6.2-10.4	6.6	3.9	2.6-5.5
	2011-2015	33.4	8.8	7.5-10.3	25.0	13.7	11.4-16.5	8.4	4.4	3.1-6.0	18.4	4.9	4.0-6.1	13.2	7.7	5.9-9.9	5.2	2.6	1.7-3.8
2016-2020	32.0	7.6	6.4-8.9	23.6	11.7	9.7-14.2	8.4	3.9	2.8-5.4	19.8	4.7	3.8-5.8	13.2	6.7	5.1-8.6	6.6	3.0	2.0-4.4	
Pancreas	1976-1980	24.2	12.0	9.9-14.4	10.2	11.4	8.4-15.1	14.0	12.3	9.6-15.6	18.8	9.2	7.4-11.3	8.4	9.2	6.5-12.6	10.4	9.1	6.8-11.9
	1981-1985	29.0	13.4	11.3-15.7	14.4	16.0	12.4-20.3	14.6	11.2	8.8-14.1	27.4	12.6	10.6-15.0	12.2	13.7	10.4-17.7	15.2	11.7	9.2-14.7
	1986-1990	30.4	13.2	11.2-15.5	17.2	17.7	14.0-22.0	13.2	10.0	7.8-12.8	28.0	12.2	10.3-14.5	16.0	16.9	13.2-21.2	12.0	8.9	6.8-11.4
	1991-1995	31.2	12.8	10.9-15.0	16.2	16.4	12.9-20.6	15.0	10.7	8.4-13.5	29.0	12.0	10.1-14.1	15.4	15.9	12.4-20.0	13.6	9.6	7.5-12.2
	1996-2000	31.4	11.9	10.1-14.0	15.8	13.6	10.7-17.0	15.6	10.4	8.2-13.0	29.0	11.1	9.3-13.0	14.4	13.1	10.2-16.6	14.6	9.7	7.6-12.2
	2001-2005	36.6	12.5	10.8-14.5	18.6	14.4	11.6-17.7	18.0	10.9	8.7-13.4	29.6	10.1	8.6-11.9	15.0	11.8	9.2-14.9	14.6	8.7	6.8-11.0
	2006-2010	45.6	13.7	12.0-15.7	23.2	15.5	12.8-18.7	22.4	12.6	10.3-15.2	38.8	11.8	10.2-13.6	19.8	13.2	10.7-16.2	19.0	10.6	8.6-13.0
	2011-2015	53.0	14.4	12.7-16.3	25.0	14.8	12.2-17.7	28.0	13.8	11.6-16.4	44.4	12.0	10.5-13.8	20.8	12.6	10.3-15.4	23.6	11.4	9.4-13.8
2016-2020	55.0	13.8	12.1-15.5	31.8	17.2	14.6-20.2	23.2	10.9	8.9-13.2	41.6	10.2	8.8-11.7	24.0	12.8	10.6-15.4	17.6	8.0	6.3-9.9	
Larynx	1976-1980	11.4	5.2	3.9-6.7	9.4	9.9	7.2-13.2	2.0	1.6	0.8-3.0	5.4	2.4	1.6-3.5	4.4	4.8	2.9-7.3	1.0	0.8	0.2-1.8
	1981-1985	13.4	6.0	4.6-7.6	11.4	11.8	8.8-15.4	2.0	1.6	0.8-3.0	3.8	1.6	1.0-2.5	3.2	3.2	1.8-5.2	0.6	0.4	0.1-1.3
	1986-1990	11.2	4.8	3.6-6.2	9.8	9.5	7.0-12.6	1.4	1.1	0.5-2.3	2.6	1.2	0.6-2.0	2.2	2.3	1.1-4.1	0.4	0.3	0.0-1.1
	1991-1995	13.4	5.4	4.2-6.8	9.8	8.8	6.5-11.7	3.6	2.6	1.6-4.2	4.4	1.8	1.1-2.7	2.6	2.4	1.3-4.2	1.8	1.3	0.6-2.5
	1996-2000	11.0	4.1	3.1-5.3	7.0	5.9	4.1-8.3	4.0	2.9	1.7-4.4	4.2	1.6	1.0-2.4	2.8	2.4	1.3-4.1	1.4	1.0	0.4-2.1
	2001-2005	10.4	3.4	2.5-4.5	7.8	5.3	3.7-7.2	2.6	1.7	0.9-2.8	2.8	0.9	0.5-1.6	2.2	1.7	0.8-3.1	0.6	0.4	0.1-1.2
	2006-2010	11.8	3.4	2.6-4.4	10.8	6.8	5.0-8.9	1.0	0.5	0.2-1.2	2.8	0.9	0.5-1.5	2.2	1.6	0.8-2.8	0.6	0.3	0.1-0.9
	2011-2015	9.0	2.3	1.6-3.1	7.6	4.1	2.9-5.8	1.4	0.7	0.3-1.5	2.4	0.6	0.3-1.1	2.2	1.2	0.6-2.3	0.2	0.1	0.0-0.7
2016-2020	13.0	3.1	2.4-4.0	11.6	6.0	4.5-7.8	1.4	0.7	0.3-1.5	2.0	0.5	0.2-1.0	2.0	1.1	0.5-2.2	0.0	0.0	0.0-0.5	
Lung and bronchus	1976-1980	125.4	58.0	53.5-62.8	91.0	97.3	88.2-107.0	34.4	29.5	25.2-34.3	98.8	45.7	41.7-50.0	73.6	78.9	70.8-87.8	25.2	21.1	17.6-25.2
	1981-1985	143.4	63.0	58.4-67.8	97.8	102.2	92.9-112.0	45.6	35.7	31.1-40.7	113.8	50.3	46.2-54.7	77.0	82.2	73.9-91.2	36.8	28.6	24.5-33.0
	1986-1990	165.2	70.4	65.6-75.4	101.6	102.2	93.2-111.8	63.6	49.1	43.8-54.8	134.2	57.2	52.9-61.7	84.6	87.0	78.6-96.0	49.6	37.8	33.2-42.9
	1991-1995	182.2	74.3	69.5-79.3	108.2	102.8	94.1-112.1	74.0	54.2	48.8-60.1	148.4	60.8	56.5-65.3	84.6	81.9	74.0-90.3	63.8	46.3	41.3-51.7
	1996-2000	194.6	73.6	69.1-78.4	107.8	93.8	85.9-102.2	86.8	59.3	53.8-65.2	155.6	59.1	55.0-63.4	88.8	79.2	71.8-87.1	66.8	45.5	40.7-50.7
	2001-2005	213.4	73.2	68.8-77.7	116.8	90.6	83.2-98.4	96.6	60.3	55.0-66.0	151.4	52.3	48.6-56.1	79.6	63.9	57.6-70.6	71.8	44.3	39.8-49.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Dutchess**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	218.0	67.0	63.1-71.2	112.2	77.9	71.4-84.7	105.8	59.7	54.7-65.1	149.2	46.1	42.8-49.6	80.0	56.3	50.8-62.2	69.2	38.8	34.8-43.2
	2011-2015	226.8	62.2	58.6-66.0	110.4	67.2	61.6-73.2	116.4	58.7	54.0-63.8	142.0	39.2	36.3-42.2	74.6	46.2	41.5-51.3	67.4	33.7	30.2-37.6
	2016-2020	242.8	59.1	55.7-62.6	117.0	60.9	56.0-66.3	125.8	57.9	53.3-62.7	128.8	31.2	28.8-33.8	65.6	34.6	30.9-38.7	63.2	28.7	25.6-32.2
Melanoma of the skin	1976-1980	11.6	5.4	4.0-6.9	6.6	6.4	4.4-9.0	5.0	4.4	2.8-6.5	6.8	3.2	2.2-4.4	4.4	4.5	2.8-6.9	2.4	2.1	1.1-3.7
	1981-1985	19.4	8.6	7.0-10.5	9.2	8.7	6.4-11.7	10.2	8.5	6.3-11.2	5.6	2.5	1.6-3.6	2.0	2.0	1.0-3.7	3.6	2.9	1.7-4.5
	1986-1990	20.0	8.6	7.0-10.4	9.4	8.4	6.2-11.3	10.6	8.8	6.5-11.5	6.8	3.0	2.1-4.2	4.0	3.7	2.3-5.8	2.8	2.3	1.2-3.8
	1991-1995	15.4	6.1	4.8-7.7	8.2	7.3	5.2-9.9	7.2	5.2	3.7-7.3	8.2	3.4	2.4-4.6	4.0	3.7	2.2-5.7	4.2	3.1	1.9-4.7
	1996-2000	26.6	9.8	8.2-11.6	17.4	14.1	11.2-17.5	9.2	6.3	4.6-8.4	7.8	3.0	2.1-4.0	4.2	3.7	2.2-5.7	3.6	2.5	1.5-3.9
	2001-2005	34.6	11.5	9.9-13.4	19.2	13.5	10.9-16.5	15.4	9.8	7.7-12.2	8.2	2.8	2.0-3.8	5.2	3.8	2.4-5.5	3.0	1.9	1.0-3.1
	2006-2010	52.2	16.4	14.5-18.6	29.6	20.2	17.0-23.8	22.6	13.8	11.4-16.7	5.6	1.7	1.1-2.5	2.8	1.9	1.0-3.2	2.8	1.6	0.9-2.7
	2011-2015	76.2	21.3	19.2-23.7	44.4	27.3	23.7-31.2	31.8	17.1	14.5-20.1	9.4	2.6	1.9-3.5	6.0	3.7	2.5-5.4	3.4	1.8	1.0-3.0
2016-2020	64.8	16.6	14.8-18.6	41.8	22.4	19.4-25.7	23.0	11.9	9.7-14.5	9.2	2.3	1.7-3.1	5.2	2.8	1.8-4.2	4.0	1.9	1.1-3.0	
Female breast	1976-1980							114.6	101.8	93.5-110.6							43.4	38.3	33.4-43.9
	1981-1985							131.6	109.9	101.6-118.8							40.8	33.6	29.1-38.7
	1986-1990							158.8	125.6	116.9-134.7							47.6	36.4	31.9-41.4
	1991-1995							168.0	126.7	118.2-135.6							45.2	32.7	28.6-37.3
	1996-2000							203.2	140.7	132.2-149.7							46.8	31.2	27.3-35.5
	2001-2005							213.4	133.2	125.3-141.5							41.6	25.5	22.1-29.2
	2006-2010							248.4	143.0	135.1-151.3							42.4	23.1	20.0-26.5
	2011-2015							264.0	139.3	131.7-147.3							37.4	18.4	15.8-21.4
2016-2020							260.0	133.1	125.5-141.0							44.2	20.4	17.7-23.4	
Cervix uteri	1976-1980							13.6	12.3	9.5-15.7							5.4	4.8	3.2-7.1
	1981-1985							10.6	9.1	6.8-12.0							4.2	3.5	2.1-5.3
	1986-1990							13.2	10.3	7.9-13.1							3.6	2.7	1.6-4.3
	1991-1995							13.4	9.9	7.7-12.6							4.4	3.2	2.0-4.9
	1996-2000							14.6	10.2	8.0-12.9							3.2	2.3	1.3-3.7
	2001-2005							11.2	7.3	5.5-9.5							3.6	2.2	1.3-3.5
	2006-2010							12.4	7.9	6.0-10.2							2.8	1.6	0.9-2.8
	2011-2015							10.2	6.2	4.5-8.3							3.0	1.8	1.0-3.0
	2016-2020							9.6	5.8	4.2-7.8							2.6	1.2	0.6-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Dutchess**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							26.4	22.5	18.8-26.8							6.6	5.3	3.6-7.5
	1981-1985							29.4	23.7	19.9-27.9							6.6	5.0	3.5-7.1
	1986-1990							31.6	24.4	20.7-28.5							5.0	3.6	2.4-5.4
	1991-1995							30.8	23.0	19.5-27.0							6.0	4.2	2.8-6.0
	1996-2000							39.6	27.5	23.8-31.6							6.0	3.9	2.7-5.7
	2001-2005							52.6	32.8	28.9-37.0							7.6	4.6	3.3-6.4
	2006-2010							59.8	33.6	29.8-37.7							8.2	4.5	3.2-6.1
	2011-2015							62.2	31.9	28.3-35.7							8.8	4.2	3.0-5.7
	2016-2020							66.4	31.0	27.7-34.7							8.8	4.0	2.9-5.4
Ovary	1976-1980							17.4	15.2	12.1-18.8							11.0	9.5	7.1-12.4
	1981-1985							24.8	20.3	16.8-24.2							15.2	11.8	9.3-14.8
	1986-1990							21.2	16.4	13.4-19.9							14.4	11.0	8.6-13.8
	1991-1995							20.8	15.6	12.7-18.9							12.6	9.2	7.1-11.9
	1996-2000							22.2	15.4	12.6-18.5							13.0	8.8	6.8-11.2
	2001-2005							19.8	12.5	10.1-15.2							14.4	8.9	7.0-11.2
	2006-2010							22.8	13.1	10.8-15.8							14.4	8.1	6.3-10.3
	2011-2015							23.2	12.5	10.3-15.1							11.8	5.9	4.5-7.7
	2016-2020							20.2	9.9	7.9-12.1							11.6	5.3	4.0-7.0
Prostate	1976-1980				59.8	78.2	69.1-88.0							17.8	25.2	20.0-31.3			
	1981-1985				77.6	94.3	84.7-104.6							27.4	35.8	29.8-42.6			
	1986-1990				97.8	111.3	101.3-122.0							26.8	34.3	28.5-40.8			
	1991-1995				172.6	170.1	158.6-182.2							28.2	34.3	28.7-40.6			
	1996-2000				215.8	182.9	172.0-194.4							32.8	36.2	30.7-42.3			
	2001-2005				213.6	159.2	149.6-169.2							30.4	29.2	24.6-34.2			
	2006-2010				220.8	143.2	134.7-152.1							27.6	22.2	18.6-26.3			
	2011-2015				223.0	123.2	115.9-130.8							31.4	21.5	18.2-25.1			
	2016-2020				250.0	124.7	117.7-132.0							27.8	16.2	13.6-19.2			
Testis	1976-1980				4.2	3.5	2.1-5.4							0.2	0.2	0.0-1.0			
	1981-1985				5.2	3.9	2.5-5.8							0.4	0.3	0.0-1.3			
	1986-1990				6.0	3.9	2.6-5.7							0.2	0.2	0.0-1.0			
	1991-1995				7.2	4.8	3.4-6.8							0.6	0.4	0.1-1.3			
	1996-2000				8.8	6.1	4.4-8.3							0.2	0.1	0.0-0.8			
	2001-2005				7.6	5.2	3.6-7.1							0.0	0.0	0.0-0.6			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Dutchess**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				9.2	6.6	4.8-8.9							0.0	0.0	0.0-0.5			
	2011-2015				10.6	7.5	5.6-9.9							0.2	0.2	0.0-0.8			
	2016-2020				8.6	6.2	4.4-8.3							0.8	0.5	0.1-1.3			
Urinary bladder (incl. in situ)	1976-1980	50.2	24.4	21.5-27.7	38.6	45.6	39.0-52.9	11.6	10.3	7.8-13.4	11.8	6.1	4.6-7.8	9.6	12.4	8.9-16.6	2.2	2.1	1.0-3.7
	1981-1985	54.8	24.7	21.8-27.8	41.4	44.5	38.4-51.3	13.4	11.0	8.5-13.9	13.0	6.1	4.7-7.8	8.4	10.1	7.2-13.8	4.6	3.7	2.3-5.5
	1986-1990	54.6	23.9	21.1-26.9	38.6	41.5	35.6-48.0	16.0	12.1	9.6-15.1	8.2	3.6	2.6-4.9	6.4	7.6	5.1-10.8	1.8	1.3	0.6-2.5
	1991-1995	61.0	25.3	22.5-28.3	42.2	43.0	37.2-49.5	18.8	13.7	11.1-16.8	13.8	5.9	4.6-7.4	8.4	9.8	7.0-13.4	5.4	3.8	2.5-5.5
	1996-2000	67.0	25.3	22.7-28.2	47.8	43.0	37.6-49.0	19.2	12.9	10.4-15.8	16.6	6.4	5.1-8.0	9.8	9.9	7.2-13.1	6.8	4.3	3.0-6.1
	2001-2005	80.4	27.5	24.9-30.3	60.2	47.9	42.5-53.8	20.2	12.3	10.0-15.0	16.2	5.7	4.5-7.1	12.2	10.6	8.1-13.7	4.0	2.4	1.5-3.8
	2006-2010	88.0	27.3	24.8-30.0	66.6	48.2	43.1-53.8	21.4	11.9	9.7-14.4	13.8	4.4	3.4-5.5	11.0	8.8	6.6-11.4	2.8	1.5	0.8-2.6
	2011-2015	99.2	27.5	25.1-30.0	72.4	44.6	40.0-49.6	26.8	14.0	11.7-16.7	18.0	5.0	4.0-6.1	13.8	9.3	7.2-11.8	4.2	2.0	1.2-3.2
2016-2020	94.8	23.3	21.2-25.5	72.8	39.7	35.7-44.2	22.0	9.8	8.0-11.9	17.8	4.3	3.5-5.4	12.8	7.7	5.9-9.9	5.0	2.0	1.3-3.0	
Kidney and renal pelvis	1976-1980	17.4	8.2	6.6-10.2	10.0	10.7	7.8-14.2	7.4	6.3	4.4-8.8	8.0	3.9	2.8-5.3	4.0	4.7	2.8-7.3	4.0	3.5	2.1-5.4
	1981-1985	19.4	8.4	6.8-10.2	11.6	11.7	8.8-15.3	7.8	6.1	4.3-8.4	9.4	4.1	3.0-5.4	6.4	6.5	4.3-9.3	3.0	2.3	1.3-3.8
	1986-1990	21.4	9.4	7.7-11.3	14.0	14.0	10.8-17.8	7.4	5.7	4.0-7.9	10.2	4.4	3.3-5.8	6.6	7.0	4.8-9.9	3.6	2.7	1.6-4.3
	1991-1995	30.6	12.2	10.4-14.3	16.6	14.9	11.8-18.6	14.0	10.2	8.0-12.9	13.0	5.4	4.1-6.8	7.6	7.5	5.3-10.4	5.4	3.8	2.5-5.5
	1996-2000	31.4	11.7	9.9-13.6	19.8	16.5	13.3-20.1	11.6	8.0	6.1-10.4	7.6	2.9	2.1-4.0	4.6	4.3	2.7-6.5	3.0	2.0	1.1-3.3
	2001-2005	43.2	14.6	12.7-16.7	25.8	18.6	15.5-22.1	17.4	11.0	8.8-13.6	14.2	4.8	3.8-6.1	8.8	6.5	4.7-8.8	5.4	3.2	2.1-4.8
	2006-2010	50.4	14.9	13.1-16.9	32.4	20.2	17.1-23.6	18.0	10.1	8.1-12.5	10.8	3.3	2.5-4.3	6.6	4.7	3.2-6.7	4.2	2.3	1.4-3.5
	2011-2015	61.8	16.8	15.0-18.9	40.2	23.0	19.9-26.5	21.6	11.7	9.5-14.2	9.6	2.6	1.9-3.5	7.0	4.3	3.0-6.1	2.6	1.3	0.7-2.3
2016-2020	70.6	18.3	16.4-20.4	45.8	24.6	21.4-28.2	24.8	12.6	10.3-15.2	9.8	2.4	1.8-3.2	6.0	3.3	2.2-4.8	3.8	1.6	1.0-2.6	
Brain and other central nervous system	1976-1980	9.8	4.3	3.1-5.7	5.8	5.5	3.6-8.0	4.0	3.4	2.0-5.3	6.0	2.6	1.7-3.7	3.0	2.7	1.5-4.6	3.0	2.5	1.4-4.2
	1981-1985	15.0	6.4	5.0-8.0	8.2	7.5	5.4-10.3	6.8	5.4	3.7-7.6	7.8	3.3	2.4-4.6	4.0	3.7	2.2-5.7	3.8	3.0	1.8-4.7
	1986-1990	16.8	6.7	5.4-8.4	8.4	7.2	5.2-9.9	8.4	6.4	4.6-8.6	11.6	4.7	3.5-6.0	6.0	5.2	3.5-7.5	5.6	4.3	2.8-6.2
	1991-1995	19.0	7.4	6.0-9.0	10.0	8.5	6.3-11.3	9.0	6.5	4.8-8.7	11.0	4.4	3.3-5.7	6.6	5.5	3.8-7.8	4.4	3.1	2.0-4.8
	1996-2000	15.8	5.8	4.6-7.2	8.6	6.5	4.7-8.7	7.2	5.0	3.5-7.0	10.4	3.8	2.9-5.0	5.4	4.3	2.8-6.2	5.0	3.5	2.3-5.2
	2001-2005	20.6	7.1	5.8-8.6	11.6	8.5	6.5-11.1	9.0	5.9	4.3-8.0	11.8	4.0	3.1-5.2	6.4	4.8	3.3-6.8	5.4	3.5	2.3-5.1
	2006-2010	22.6	7.1	5.8-8.6	11.8	7.9	6.0-10.3	10.8	6.4	4.8-8.4	13.2	4.1	3.2-5.2	7.0	4.6	3.2-6.5	6.2	3.6	2.5-5.2
	2011-2015	24.6	7.4	6.1-8.9	14.0	8.7	6.8-11.1	10.6	6.2	4.6-8.3	13.8	3.7	2.9-4.7	8.6	4.9	3.5-6.7	5.2	2.7	1.7-4.0
2016-2020	24.6	6.9	5.7-8.4	13.2	7.5	5.7-9.6	11.4	6.5	4.8-8.5	18.0	4.8	3.8-5.9	9.2	5.2	3.7-7.0	8.8	4.5	3.2-6.2	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Dutchess**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	4.2	1.8	1.1-2.7	0.8	0.7	0.2-1.8	3.4	2.8	1.6-4.5	0.4	0.2	0.0-0.7	0.0	0.0	0.0-0.8	0.4	0.3	0.0-1.2
	1981-1985	8.0	3.4	2.4-4.6	1.8	1.6	0.7-3.1	6.2	5.2	3.5-7.4	0.6	0.3	0.1-0.9	0.2	0.3	0.0-1.4	0.4	0.3	0.0-1.2
	1986-1990	9.6	3.8	2.8-5.0	4.0	3.1	1.9-4.9	5.6	4.4	2.9-6.4	1.2	0.5	0.2-1.1	0.4	0.3	0.0-1.3	0.8	0.6	0.2-1.5
	1991-1995	14.8	5.7	4.5-7.2	5.0	4.4	2.8-6.6	9.8	7.6	5.6-10.0	0.4	0.2	0.0-0.6	0.4	0.6	0.1-1.9	0.0	0.0	0.0-0.6
	1996-2000	16.0	5.8	4.6-7.2	3.8	2.7	1.6-4.2	12.2	8.7	6.7-11.2	1.6	0.6	0.3-1.2	0.4	0.4	0.0-1.3	1.2	0.8	0.3-1.8
	2001-2005	36.8	12.3	10.6-14.3	10.2	6.8	5.0-8.9	26.6	18.1	15.2-21.5	1.2	0.4	0.1-0.9	0.4	0.3	0.0-1.0	0.8	0.5	0.1-1.3
	2006-2010	50.8	16.5	14.5-18.7	10.0	6.8	5.0-9.0	40.8	26.2	22.6-30.1	0.8	0.2	0.1-0.6	0.0	0.0	0.0-0.5	0.8	0.4	0.1-1.1
	2011-2015	61.8	19.2	17.1-21.6	17.0	10.4	8.2-13.0	44.8	28.3	24.5-32.4	2.2	0.6	0.3-1.2	0.8	0.5	0.1-1.4	1.4	0.7	0.3-1.6
	2016-2020	49.2	15.2	13.3-17.4	13.6	8.1	6.2-10.4	35.6	22.5	19.1-26.2	2.2	0.5	0.3-1.0	0.2	0.1	0.0-0.8	2.0	0.9	0.4-1.7
Hodgkin lymphoma	1976-1980	8.0	3.0	2.1-4.2	4.4	3.4	2.1-5.2	3.6	2.6	1.5-4.1	2.4	1.0	0.5-1.9	1.0	0.8	0.2-2.0	1.4	1.2	0.5-2.6
	1981-1985	9.8	3.7	2.7-5.0	4.2	3.2	1.9-4.9	5.6	4.4	2.9-6.4	2.2	0.9	0.4-1.7	0.6	0.5	0.1-1.6	1.6	1.2	0.5-2.4
	1986-1990	9.4	3.5	2.5-4.6	5.6	4.3	2.8-6.3	3.8	2.7	1.6-4.3	2.8	1.1	0.6-1.9	1.8	1.8	0.8-3.5	1.0	0.7	0.2-1.8
	1991-1995	9.2	3.4	2.5-4.5	3.6	2.6	1.5-4.2	5.6	4.1	2.7-5.9	2.0	0.8	0.4-1.4	1.2	0.9	0.3-2.1	0.8	0.6	0.2-1.5
	1996-2000	8.4	3.0	2.2-4.1	4.4	3.2	2.0-4.9	4.0	2.9	1.8-4.5	1.8	0.6	0.3-1.2	0.8	0.7	0.2-1.8	1.0	0.7	0.2-1.7
	2001-2005	10.0	3.4	2.5-4.5	5.4	3.8	2.5-5.6	4.6	3.2	2.0-4.9	1.2	0.4	0.2-0.9	0.6	0.4	0.1-1.2	0.6	0.4	0.1-1.3
	2006-2010	8.6	2.9	2.1-3.9	6.0	4.1	2.8-5.9	2.6	1.8	0.9-3.0	2.4	0.7	0.4-1.3	1.2	0.8	0.3-1.8	1.2	0.6	0.2-1.4
	2011-2015	10.8	3.6	2.7-4.7	5.6	3.7	2.4-5.4	5.2	3.6	2.3-5.3	0.8	0.3	0.1-0.7	0.6	0.4	0.1-1.2	0.2	0.1	0.0-0.8
	2016-2020	10.6	3.2	2.4-4.3	4.6	2.6	1.6-4.1	6.0	3.9	2.6-5.7	1.0	0.2	0.1-0.6	0.6	0.3	0.1-1.0	0.4	0.2	0.0-0.8
Non-Hodgkin lymphomas	1976-1980	17.2	7.8	6.2-9.7	10.0	9.8	7.2-13.1	7.2	6.2	4.3-8.7	9.0	4.2	3.1-5.7	4.6	5.0	3.1-7.6	4.4	3.7	2.3-5.7
	1981-1985	24.0	10.4	8.6-12.4	11.8	10.9	8.2-14.2	12.2	9.6	7.3-12.3	10.2	4.4	3.3-5.9	5.2	4.7	3.0-7.0	5.0	3.9	2.5-5.7
	1986-1990	40.6	17.3	15.0-19.9	21.4	21.7	17.6-26.4	19.2	14.7	11.9-17.9	18.8	8.3	6.7-10.2	9.6	10.8	7.8-14.4	9.2	6.8	5.0-9.1
	1991-1995	39.0	15.8	13.7-18.2	22.4	20.1	16.4-24.3	16.6	11.9	9.5-14.8	20.4	8.4	6.8-10.2	11.6	11.5	8.6-15.0	8.8	6.3	4.5-8.4
	1996-2000	56.4	21.2	18.8-23.9	28.6	24.6	20.6-29.1	27.8	18.7	15.7-22.1	22.8	8.7	7.2-10.5	10.0	9.3	6.8-12.3	12.8	8.2	6.3-10.5
	2001-2005	56.4	19.0	16.9-21.4	32.4	23.5	19.9-27.4	24.0	15.1	12.5-18.1	16.0	5.5	4.3-6.8	9.0	6.9	5.0-9.3	7.0	4.1	2.9-5.8
	2006-2010	67.6	20.9	18.7-23.3	38.2	26.1	22.4-30.1	29.4	16.7	14.1-19.7	19.2	6.0	4.9-7.4	9.0	6.6	4.8-8.9	10.2	5.6	4.2-7.4
	2011-2015	62.4	17.6	15.6-19.7	34.4	21.5	18.3-25.1	28.0	14.4	12.0-17.0	15.8	4.6	3.6-5.7	8.6	5.8	4.2-7.9	7.2	3.7	2.6-5.3
	2016-2020	78.2	20.3	18.3-22.5	46.8	26.3	22.9-30.0	31.4	15.3	12.9-18.0	18.2	4.6	3.7-5.7	12.0	6.9	5.3-9.0	6.2	2.8	1.8-4.0
Myeloma	1976-1980	6.8	3.3	2.3-4.7	3.4	4.3	2.4-6.9	3.4	2.9	1.7-4.7	2.8	1.4	0.7-2.3	1.6	1.9	0.8-3.7	1.2	1.0	0.4-2.2
	1981-1985	10.6	4.8	3.6-6.3	5.4	5.6	3.6-8.3	5.2	4.0	2.6-5.9	7.0	3.2	2.2-4.5	3.2	3.8	2.1-6.2	3.8	3.0	1.8-4.7
	1986-1990	13.0	5.7	4.4-7.3	5.6	6.1	4.0-8.9	7.4	5.5	3.9-7.7	10.2	4.5	3.4-6.0	4.8	4.9	3.1-7.4	5.4	4.1	2.7-6.0
	1991-1995	14.0	5.8	4.5-7.3	8.0	7.8	5.5-10.7	6.0	4.3	2.9-6.1	7.6	3.2	2.3-4.4	4.6	5.0	3.1-7.6	3.0	2.2	1.2-3.6
	1996-2000	13.2	5.0	3.9-6.3	7.4	6.4	4.5-8.9	5.8	3.9	2.6-5.7	10.0	3.8	2.8-5.1	5.0	4.5	2.9-6.7	5.0	3.3	2.1-4.9
	2001-2005	18.2	6.2	5.0-7.6	9.6	7.4	5.4-9.8	8.6	5.3	3.8-7.2	9.6	3.3	2.5-4.4	5.4	4.4	2.9-6.5	4.2	2.6	1.6-4.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Dutchess**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	23.4	7.2	6.0-8.7	15.8	10.9	8.6-13.7	7.6	4.1	2.9-5.7	11.6	3.7	2.8-4.8	6.4	4.7	3.2-6.7	5.2	2.8	1.8-4.2
	2011-2015	29.4	7.8	6.6-9.3	16.8	9.7	7.7-12.1	12.6	6.3	4.8-8.1	12.6	3.4	2.6-4.4	7.4	4.4	3.1-6.1	5.2	2.6	1.7-3.8
	2016-2020	25.6	6.3	5.2-7.5	14.0	7.0	5.4-9.0	11.6	5.5	4.1-7.2	11.6	2.8	2.1-3.7	5.0	2.6	1.7-4.0	6.6	2.9	2.0-4.3
Leukemias	1976-1980	26.0	12.4	10.3-14.8	14.4	15.7	12.1-20.0	11.6	10.1	7.6-13.1	13.8	6.6	5.1-8.4	6.4	7.3	4.8-10.5	7.4	6.4	4.5-8.9
	1981-1985	27.2	12.5	10.4-14.7	17.0	17.9	14.1-22.3	10.2	8.0	6.0-10.6	17.0	7.9	6.3-9.7	9.6	10.0	7.3-13.4	7.4	5.9	4.1-8.1
	1986-1990	24.6	10.3	8.5-12.3	14.0	14.1	10.8-18.0	10.6	7.9	5.9-10.4	16.0	6.9	5.4-8.6	10.4	11.0	8.1-14.6	5.6	4.1	2.7-5.9
	1991-1995	31.0	12.6	10.7-14.8	18.4	17.3	13.8-21.3	12.6	9.2	7.1-11.8	19.4	8.2	6.6-10.0	10.0	10.9	8.0-14.5	9.4	6.5	4.8-8.7
	1996-2000	35.2	13.3	11.4-15.4	21.2	18.2	14.8-22.0	14.0	9.6	7.5-12.2	23.8	9.1	7.5-10.9	14.2	12.8	10.0-16.2	9.6	6.5	4.8-8.6
	2001-2005	45.0	15.6	13.6-17.8	24.6	19.6	16.2-23.5	20.4	12.9	10.5-15.7	24.8	8.7	7.2-10.4	15.2	12.7	10.0-16.0	9.6	5.8	4.2-7.7
	2006-2010	48.6	15.3	13.4-17.3	30.0	20.6	17.4-24.3	18.6	11.0	8.8-13.6	24.6	7.6	6.3-9.1	13.8	9.9	7.7-12.7	10.8	5.9	4.4-7.8
	2011-2015	59.2	16.9	15.0-19.0	33.2	20.7	17.6-24.3	26.0	14.3	11.8-17.1	23.0	6.3	5.2-7.7	14.2	9.2	7.2-11.7	8.8	4.3	3.1-5.9
2016-2020	57.8	15.2	13.4-17.1	33.8	19.0	16.2-22.3	24.0	11.6	9.5-14.0	21.4	5.5	4.5-6.7	13.4	7.9	6.1-10.1	8.0	3.5	2.5-4.9	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Erie**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	4,136.6	415.1	409.3-421.0	2,091.0	513.5	503.0-524.1	2,045.6	360.3	353.2-367.6	2,285.6	230.6	226.3-235.0	1,220.6	307.9	299.7-316.3	1,065.0	184.0	179.0-189.1
	1981-1985	4,416.6	435.1	429.3-441.1	2,217.0	536.4	525.8-547.1	2,199.6	382.3	375.0-389.8	2,332.4	229.3	225.1-233.6	1,219.2	303.4	295.4-311.7	1,113.2	187.3	182.3-192.4
	1986-1990	4,743.8	454.6	448.7-460.6	2,345.0	547.1	536.7-557.7	2,398.8	406.5	399.0-414.1	2,414.2	230.5	226.3-234.8	1,258.6	304.2	296.3-312.3	1,155.6	187.9	183.0-193.0
	1991-1995	5,100.6	469.7	463.9-475.6	2,564.4	569.7	559.6-580.0	2,536.2	413.2	405.9-420.7	2,488.8	226.9	222.9-231.0	1,259.6	290.9	283.4-298.4	1,229.2	189.8	185.0-194.8
	1996-2000	5,535.2	502.9	496.9-508.9	2,775.2	600.4	590.3-610.6	2,760.0	444.4	436.9-452.1	2,399.0	212.1	208.3-216.0	1,187.8	264.9	258.1-271.8	1,211.2	180.9	176.3-185.6
	2001-2005	5,676.0	511.4	505.4-517.5	2,893.4	615.4	605.3-625.5	2,782.6	446.6	439.0-454.2	2,294.2	199.1	195.4-202.8	1,131.4	245.6	239.2-252.1	1,162.8	171.8	167.3-176.4
	2006-2010	6,040.2	539.3	533.1-545.5	3,087.8	634.0	623.9-644.2	2,952.4	477.4	469.5-485.4	2,209.6	188.5	184.9-192.1	1,082.8	225.7	219.7-231.8	1,126.8	165.9	161.5-170.5
	2011-2015	6,278.6	542.4	536.3-548.7	3,165.0	610.0	600.3-619.8	3,113.6	499.1	491.0-507.4	2,183.0	180.2	176.8-183.7	1,091.0	215.3	209.5-221.2	1,092.0	156.6	152.3-161.0
	2016-2020	6,334.4	520.4	514.4-526.3	3,183.0	571.6	562.5-580.9	3,151.4	486.9	478.9-495.0	2,056.4	161.0	157.8-164.2	1,024.4	189.0	183.7-194.4	1,032.0	142.3	138.3-146.4
Oral cavity and pharynx	1976-1980	122.0	11.9	10.9-12.9	82.8	18.9	17.0-20.9	39.2	6.8	5.8-7.8	48.6	4.7	4.1-5.3	36.8	8.3	7.1-9.7	11.8	2.0	1.5-2.6
	1981-1985	136.4	13.2	12.2-14.3	93.0	21.3	19.3-23.5	43.4	7.3	6.4-8.4	51.8	4.9	4.3-5.6	37.0	8.4	7.1-9.8	14.8	2.4	1.9-3.1
	1986-1990	121.4	11.9	10.9-12.9	81.2	18.2	16.4-20.1	40.2	7.0	6.1-8.1	44.2	4.3	3.7-4.9	29.4	6.8	5.7-8.0	14.8	2.5	1.9-3.2
	1991-1995	113.8	10.9	10.0-11.8	72.0	15.9	14.2-17.6	41.8	6.9	5.9-7.9	39.2	3.7	3.2-4.2	26.0	5.6	4.7-6.7	13.2	2.1	1.6-2.7
	1996-2000	117.8	11.1	10.2-12.0	76.2	16.4	14.8-18.1	41.6	6.6	5.7-7.6	36.8	3.4	2.9-3.9	22.4	4.8	4.0-5.8	14.4	2.2	1.7-2.7
	2001-2005	114.0	10.6	9.7-11.5	73.2	15.3	13.7-16.9	40.8	6.8	5.8-7.8	30.2	2.7	2.3-3.2	20.2	4.2	3.4-5.2	10.0	1.4	1.1-1.9
	2006-2010	130.6	11.7	10.8-12.6	87.2	17.4	15.8-19.2	43.4	7.0	6.1-8.1	27.4	2.4	2.0-2.8	18.2	3.7	2.9-4.5	9.2	1.4	1.0-1.8
	2011-2015	142.2	12.1	11.2-13.1	101.4	19.0	17.3-20.7	40.8	6.3	5.5-7.3	38.0	3.2	2.7-3.7	29.0	5.6	4.7-6.6	9.0	1.2	0.9-1.6
	2016-2020	170.2	13.6	12.7-14.6	118.4	20.5	18.9-22.3	51.8	7.8	6.8-8.9	38.8	2.9	2.5-3.4	28.6	4.9	4.1-5.8	10.2	1.4	1.0-1.8
Esophagus	1976-1980	55.8	5.5	4.8-6.2	41.8	9.8	8.4-11.3	14.0	2.4	1.9-3.0	49.6	4.9	4.2-5.5	37.2	8.6	7.4-10.0	12.4	2.1	1.6-2.7
	1981-1985	48.8	4.6	4.0-5.2	36.8	8.2	7.0-9.6	12.0	2.0	1.5-2.6	44.0	4.3	3.7-4.9	31.6	7.4	6.2-8.7	12.4	2.1	1.6-2.7
	1986-1990	58.0	5.4	4.8-6.1	43.0	9.6	8.3-11.1	15.0	2.6	2.0-3.3	50.6	4.7	4.2-5.4	38.0	8.6	7.4-10.0	12.6	2.1	1.6-2.7
	1991-1995	60.8	5.5	4.9-6.2	42.6	9.0	7.8-10.4	18.2	2.8	2.2-3.5	60.0	5.4	4.8-6.0	45.0	9.7	8.5-11.1	15.0	2.2	1.7-2.8
	1996-2000	62.2	5.5	4.9-6.2	45.4	9.6	8.4-11.0	16.8	2.5	2.0-3.1	52.4	4.6	4.1-5.3	36.4	7.9	6.7-9.1	16.0	2.4	1.9-3.0
	2001-2005	65.8	5.8	5.2-6.5	48.4	10.2	9.0-11.6	17.4	2.5	2.0-3.2	59.8	5.3	4.7-5.9	44.6	9.5	8.3-10.8	15.2	2.1	1.7-2.7
	2006-2010	77.4	6.8	6.1-7.5	59.4	12.0	10.7-13.5	18.0	2.6	2.1-3.2	61.8	5.2	4.7-5.9	45.4	9.2	8.0-10.4	16.4	2.3	1.8-2.9
	2011-2015	74.0	6.0	5.4-6.7	57.0	10.5	9.3-11.8	17.0	2.4	1.9-3.0	64.2	5.3	4.8-6.0	51.2	9.8	8.6-11.1	13.0	1.9	1.4-2.4
	2016-2020	70.8	5.6	5.0-6.2	54.8	9.6	8.5-10.9	16.0	2.3	1.8-2.9	56.6	4.3	3.8-4.9	46.2	8.0	7.0-9.2	10.4	1.4	1.0-1.9
Stomach	1976-1980	125.0	12.8	11.8-13.9	70.8	18.6	16.5-20.8	54.2	9.3	8.2-10.5	93.8	9.8	8.9-10.7	56.4	15.0	13.2-17.0	37.4	6.5	5.6-7.5
	1981-1985	103.6	10.4	9.5-11.3	61.8	15.4	13.6-17.3	41.8	7.0	6.1-8.0	79.0	7.9	7.1-8.7	45.4	11.3	9.8-13.0	33.6	5.6	4.8-6.6
	1986-1990	125.6	11.9	11.0-12.9	77.0	18.3	16.4-20.3	48.6	7.7	6.7-8.7	88.6	8.4	7.6-9.3	51.6	12.5	10.9-14.2	37.0	5.7	4.9-6.6
	1991-1995	106.8	9.7	8.9-10.5	65.4	14.8	13.1-16.5	41.4	6.1	5.3-7.1	73.8	6.7	6.0-7.4	44.4	10.2	8.9-11.7	29.4	4.3	3.6-5.1
	1996-2000	94.6	8.3	7.6-9.1	55.0	12.2	10.8-13.8	39.6	5.6	4.8-6.5	57.6	5.0	4.5-5.6	31.0	7.0	5.9-8.2	26.6	3.6	3.0-4.3
	2001-2005	85.8	7.5	6.8-8.3	50.6	10.8	9.5-12.3	35.2	5.2	4.4-6.0	54.4	4.7	4.1-5.2	32.8	7.1	6.1-8.3	21.6	3.0	2.4-3.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Erie**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	84.0	7.3	6.6-8.0	50.8	10.4	9.2-11.8	33.2	4.9	4.1-5.7	43.8	3.7	3.3-4.3	25.8	5.4	4.5-6.4	18.0	2.5	2.0-3.1
	2011-2015	96.6	8.2	7.4-9.0	58.2	11.3	10.0-12.7	38.4	5.8	5.0-6.8	43.4	3.5	3.1-4.1	28.4	5.6	4.7-6.6	15.0	2.1	1.6-2.6
	2016-2020	97.2	8.0	7.3-8.8	59.0	11.0	9.8-12.4	38.2	5.6	4.8-6.5	34.2	2.7	2.3-3.2	20.0	3.7	3.0-4.5	14.2	1.9	1.4-2.4
Colorectal	1976-1980	649.4	66.1	63.8-68.5	327.8	82.7	78.4-87.1	321.6	55.5	52.8-58.4	343.0	35.6	33.9-37.4	167.6	44.3	41.1-47.6	175.4	30.5	28.4-32.6
	1981-1985	682.8	67.6	65.3-69.9	349.6	86.6	82.3-91.0	333.2	55.5	52.8-58.3	331.0	33.3	31.6-34.9	163.4	42.9	39.8-46.2	167.6	27.7	25.9-29.7
	1986-1990	667.4	62.9	60.8-65.1	331.4	78.2	74.3-82.3	336.0	53.2	50.6-55.8	319.0	30.8	29.3-32.4	161.8	40.3	37.4-43.3	157.2	24.9	23.2-26.7
	1991-1995	636.6	57.3	55.3-59.3	322.4	73.1	69.4-76.9	314.2	47.1	44.7-49.5	282.6	25.5	24.1-26.8	135.4	32.0	29.5-34.6	147.2	21.6	20.1-23.3
	1996-2000	660.2	58.5	56.5-60.5	326.2	71.5	68.0-75.1	334.0	49.7	47.3-52.2	258.8	22.7	21.5-24.0	122.6	28.0	25.8-30.4	136.2	19.6	18.1-21.1
	2001-2005	614.0	53.8	51.9-55.7	306.2	65.8	62.5-69.2	307.8	45.5	43.2-47.9	224.2	19.1	18.0-20.2	110.6	24.3	22.3-26.4	113.6	15.9	14.6-17.3
	2006-2010	513.0	44.8	43.0-46.6	255.0	52.6	49.7-55.6	258.0	39.1	36.9-41.3	183.6	15.3	14.3-16.3	91.6	19.2	17.4-21.0	92.0	12.7	11.5-14.0
	2011-2015	476.4	40.8	39.2-42.5	235.4	46.1	43.5-48.9	241.0	36.5	34.4-38.8	166.2	13.4	12.5-14.4	81.0	16.0	14.5-17.7	85.2	11.4	10.3-12.7
2016-2020	452.4	37.7	36.1-39.3	222.6	41.6	39.1-44.2	229.8	34.5	32.4-36.6	159.4	12.5	11.7-13.5	81.0	15.0	13.6-16.6	78.4	10.6	9.5-11.7	
Colon excluding rectum	1976-1980	450.8	46.3	44.4-48.3	213.4	55.0	51.5-58.7	237.4	41.1	38.8-43.5	272.2	28.2	26.7-29.8	128.8	34.1	31.3-37.0	143.4	24.9	23.1-26.8
	1981-1985	486.0	48.5	46.5-50.5	233.0	58.8	55.2-62.5	253.0	42.2	39.9-44.6	272.0	27.4	25.9-28.9	130.2	34.8	32.0-37.8	141.8	23.4	21.7-25.3
	1986-1990	463.4	43.9	42.1-45.7	218.4	52.3	49.1-55.7	245.0	38.8	36.6-41.1	263.2	25.4	24.0-26.8	130.4	32.3	29.7-35.0	132.8	21.1	19.5-22.8
	1991-1995	447.8	40.3	38.7-42.1	213.2	49.0	46.0-52.1	234.6	34.9	32.9-37.1	235.0	21.2	20.0-22.5	111.2	26.6	24.3-29.0	123.8	18.1	16.7-19.7
	1996-2000	464.6	40.8	39.2-42.5	216.8	47.8	45.0-50.8	247.8	36.3	34.3-38.5	222.2	19.5	18.3-20.7	102.8	23.4	21.4-25.6	119.4	17.1	15.8-18.6
	2001-2005	444.8	38.6	37.0-40.3	215.0	46.4	43.6-49.3	229.8	33.5	31.6-35.6	197.8	16.8	15.7-17.9	96.2	21.1	19.2-23.0	101.6	14.1	12.9-15.5
	2006-2010	364.8	31.5	30.1-33.0	175.2	36.2	33.8-38.7	189.6	28.2	26.4-30.1	154.6	12.8	11.9-13.8	75.4	15.8	14.2-17.5	79.2	10.9	9.8-12.0
	2011-2015	326.8	27.7	26.4-29.1	152.6	30.0	27.9-32.3	174.2	25.8	24.1-27.7	129.6	10.5	9.7-11.3	60.6	12.0	10.7-13.4	69.0	9.3	8.3-10.4
2016-2020	318.6	26.4	25.1-27.8	148.4	27.9	25.8-30.0	170.2	25.1	23.4-27.0	115.8	9.0	8.3-9.8	57.6	10.6	9.4-12.0	58.2	7.7	6.8-8.7	
Rectum & rectosigmoid	1976-1980	198.6	19.7	18.5-21.0	114.4	27.7	25.3-30.2	84.2	14.4	13.1-15.9	70.8	7.4	6.6-8.2	38.8	10.2	8.7-11.8	32.0	5.6	4.7-6.5
	1981-1985	196.8	19.1	17.9-20.4	116.6	27.8	25.4-30.3	80.2	13.3	12.0-14.7	59.0	5.9	5.2-6.6	33.2	8.1	6.8-9.5	25.8	4.3	3.6-5.1
	1986-1990	204.0	19.0	17.9-20.3	113.0	25.9	23.7-28.2	91.0	14.4	13.1-15.8	55.8	5.4	4.8-6.1	31.4	8.0	6.7-9.4	24.4	3.8	3.2-4.6
	1991-1995	188.8	16.9	15.9-18.1	109.2	24.1	22.0-26.2	79.6	12.1	11.0-13.4	47.6	4.2	3.7-4.8	24.2	5.4	4.5-6.5	23.4	3.5	2.9-4.2
	1996-2000	195.6	17.6	16.6-18.8	109.4	23.7	21.7-25.7	86.2	13.4	12.2-14.8	36.6	3.2	2.8-3.7	19.8	4.6	3.7-5.6	16.8	2.4	1.9-3.0
	2001-2005	169.2	15.1	14.1-16.2	91.2	19.4	17.7-21.3	78.0	11.9	10.8-13.2	26.4	2.3	1.9-2.7	14.4	3.2	2.5-4.1	12.0	1.7	1.3-2.2
	2006-2010	148.2	13.2	12.3-14.2	79.8	16.4	14.8-18.1	68.4	10.9	9.7-12.2	29.0	2.5	2.1-2.9	16.2	3.4	2.7-4.2	12.8	1.8	1.4-2.4
	2011-2015	149.6	13.1	12.2-14.1	82.8	16.1	14.5-17.8	66.8	10.7	9.6-12.0	36.6	3.0	2.5-3.4	20.4	4.0	3.3-4.9	16.2	2.1	1.7-2.7
2016-2020	133.8	11.3	10.4-12.2	74.2	13.7	12.3-15.3	59.6	9.3	8.2-10.5	43.6	3.5	3.1-4.0	23.4	4.4	3.6-5.3	20.2	2.9	2.3-3.5	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Erie**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	32.4	3.2	2.7-3.7	21.6	5.1	4.1-6.2	10.8	1.8	1.4-2.4	26.6	2.6	2.2-3.1	16.4	3.8	3.0-4.8	10.2	1.8	1.3-2.3
	1981-1985	33.8	3.3	2.8-3.8	23.0	5.5	4.5-6.6	10.8	1.8	1.4-2.4	33.4	3.3	2.8-3.8	22.4	5.4	4.4-6.5	11.0	1.8	1.4-2.4
	1986-1990	35.6	3.3	2.8-3.8	25.2	5.5	4.6-6.7	10.4	1.7	1.3-2.2	33.4	3.1	2.6-3.6	21.6	4.8	3.9-5.8	11.8	1.9	1.4-2.5
	1991-1995	42.8	3.9	3.4-4.5	27.6	6.3	5.2-7.4	15.2	2.3	1.8-2.9	41.2	3.8	3.3-4.3	25.2	5.9	4.9-7.0	16.0	2.4	1.9-3.1
	1996-2000	44.2	3.9	3.4-4.5	26.8	5.8	4.9-6.9	17.4	2.5	2.0-3.1	41.6	3.7	3.2-4.2	22.0	4.8	4.0-5.9	19.6	2.9	2.3-3.5
	2001-2005	55.6	5.0	4.5-5.7	38.8	8.0	6.9-9.3	16.8	2.7	2.1-3.3	44.0	3.9	3.4-4.4	29.2	6.1	5.1-7.1	14.8	2.3	1.8-2.9
	2006-2010	68.6	6.0	5.4-6.7	49.0	9.7	8.5-11.1	19.6	3.0	2.4-3.6	60.8	5.3	4.7-5.9	39.2	7.9	6.9-9.2	21.6	3.2	2.6-3.9
	2011-2015	81.4	6.7	6.0-7.4	61.2	11.1	9.9-12.4	20.2	3.0	2.4-3.7	66.2	5.4	4.8-6.0	47.8	8.8	7.7-10.0	18.4	2.7	2.1-3.3
	2016-2020	112.2	8.5	7.8-9.3	82.0	13.9	12.6-15.4	30.2	4.2	3.5-4.9	74.4	5.7	5.1-6.3	51.8	8.8	7.7-10.0	22.6	3.2	2.6-3.9
Pancreas	1976-1980	135.4	13.6	12.5-14.7	63.6	15.1	13.4-16.9	71.8	12.3	11.1-13.7	127.8	12.8	11.8-13.9	61.8	14.6	12.9-16.5	66.0	11.3	10.1-12.6
	1981-1985	132.6	12.9	12.0-14.0	67.8	16.7	14.9-18.8	64.8	10.6	9.5-11.9	127.0	12.3	11.3-13.3	64.6	16.0	14.2-17.9	62.4	10.1	9.0-11.3
	1986-1990	127.0	12.1	11.1-13.1	61.6	14.5	12.8-16.3	65.4	10.3	9.2-11.5	122.0	11.5	10.6-12.5	61.0	14.5	12.8-16.3	61.0	9.6	8.5-10.7
	1991-1995	137.8	12.6	11.7-13.6	70.2	15.9	14.2-17.7	67.6	10.0	8.9-11.1	125.2	11.4	10.5-12.3	64.4	14.7	13.1-16.4	60.8	8.9	7.9-10.0
	1996-2000	144.4	12.8	11.9-13.8	70.4	15.5	13.9-17.2	74.0	10.8	9.7-12.0	134.4	11.8	10.9-12.7	66.4	14.7	13.1-16.4	68.0	9.8	8.8-11.0
	2001-2005	153.4	13.4	12.4-14.4	71.8	15.3	13.7-16.9	81.6	11.9	10.7-13.1	137.4	11.9	11.0-12.8	65.2	13.9	12.4-15.5	72.2	10.2	9.2-11.4
	2006-2010	154.0	13.2	12.3-14.2	72.0	15.0	13.5-16.7	82.0	12.0	10.8-13.2	132.4	11.2	10.4-12.1	60.8	12.6	11.2-14.1	71.6	10.3	9.3-11.5
	2011-2015	182.2	15.0	14.1-16.1	88.4	17.1	15.5-18.8	93.8	13.4	12.2-14.7	150.8	12.4	11.5-13.3	70.6	13.7	12.3-15.3	80.2	11.3	10.2-12.5
	2016-2020	200.0	16.0	15.0-17.1	101.4	18.4	16.8-20.2	98.6	14.0	12.7-15.3	150.0	11.8	10.9-12.7	76.0	13.9	12.5-15.4	74.0	10.1	9.1-11.3
Larynx	1976-1980	69.8	6.4	5.7-7.1	57.6	12.5	11.0-14.1	12.2	2.0	1.5-2.6	24.8	2.3	1.9-2.7	20.0	4.5	3.6-5.5	4.8	0.8	0.5-1.2
	1981-1985	66.4	6.2	5.5-6.9	53.6	11.6	10.2-13.1	12.8	2.3	1.7-3.0	22.8	2.1	1.7-2.5	18.8	4.3	3.4-5.3	4.0	0.6	0.4-0.9
	1986-1990	64.0	6.1	5.5-6.9	50.0	11.2	9.8-12.8	14.0	2.4	1.9-3.1	18.0	1.7	1.3-2.1	13.4	3.2	2.4-4.1	4.6	0.8	0.5-1.2
	1991-1995	65.6	6.1	5.4-6.8	50.8	11.1	9.8-12.6	14.8	2.4	1.9-3.1	20.4	1.9	1.5-2.3	15.2	3.4	2.7-4.3	5.2	0.8	0.5-1.2
	1996-2000	54.4	5.1	4.5-5.7	40.4	8.8	7.6-10.1	14.0	2.3	1.8-2.9	16.4	1.5	1.2-1.8	12.4	2.8	2.1-3.6	4.0	0.6	0.4-0.9
	2001-2005	53.2	4.8	4.2-5.4	41.4	8.7	7.5-9.9	11.8	2.0	1.5-2.6	15.8	1.4	1.1-1.7	12.6	2.7	2.0-3.4	3.2	0.5	0.3-0.8
	2006-2010	50.2	4.4	3.9-5.0	37.4	7.4	6.4-8.6	12.8	2.1	1.6-2.7	18.6	1.6	1.3-2.0	15.2	3.1	2.5-3.9	3.4	0.5	0.3-0.9
	2011-2015	44.6	3.8	3.3-4.3	36.6	7.0	6.0-8.1	8.0	1.3	0.9-1.8	15.0	1.2	1.0-1.5	12.6	2.4	1.8-3.1	2.4	0.4	0.2-0.7
	2016-2020	39.4	3.1	2.7-3.6	29.4	5.0	4.2-6.0	10.0	1.6	1.2-2.1	12.8	1.0	0.7-1.3	9.2	1.6	1.2-2.2	3.6	0.5	0.3-0.9
Lung and bronchus	1976-1980	663.4	63.4	61.2-65.7	493.4	115.3	110.5-120.2	170.0	28.5	26.5-30.5	542.2	52.5	50.5-54.5	406.8	97.0	92.6-101.6	135.4	22.8	21.1-24.7
	1981-1985	741.6	70.2	67.9-72.5	506.4	117.2	112.4-122.1	235.2	39.2	37.0-41.6	613.0	57.9	55.9-60.1	427.2	100.6	96.2-105.3	185.8	30.6	28.6-32.7
	1986-1990	793.0	73.5	71.2-75.9	513.0	115.4	110.8-120.2	280.0	45.5	43.1-48.0	665.0	61.5	59.4-63.7	438.4	100.0	95.6-104.5	226.6	36.5	34.3-38.7
	1991-1995	833.2	75.3	73.0-77.7	476.0	104.3	100.1-108.7	357.2	56.6	53.9-59.4	703.6	63.2	61.1-65.3	412.4	91.4	87.4-95.6	291.2	45.2	42.8-47.6
	1996-2000	881.6	78.5	76.2-80.9	480.2	103.0	98.9-107.3	401.4	62.1	59.3-64.9	697.4	61.6	59.6-63.7	389.0	84.1	80.4-88.0	308.4	46.7	44.4-49.2
	2001-2005	847.8	74.9	72.7-77.2	440.6	93.3	89.4-97.3	407.2	63.1	60.4-66.0	674.2	59.0	57.0-61.0	355.6	75.9	72.4-79.5	318.6	48.2	45.8-50.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Erie**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	868.8	75.7	73.5-78.0	429.0	88.1	84.3-91.9	439.8	67.6	64.8-70.6	629.2	54.0	52.1-55.9	323.4	66.7	63.4-70.0	305.8	45.5	43.2-47.9
	2011-2015	892.2	74.7	72.5-76.9	437.8	85.3	81.7-89.1	454.4	67.9	65.0-70.8	613.4	51.1	49.3-53.0	314.4	61.8	58.7-65.0	299.0	43.9	41.6-46.2
	2016-2020	871.4	68.1	66.0-70.2	400.0	71.8	68.6-75.1	471.4	65.9	63.2-68.7	498.0	38.5	37.0-40.1	240.6	43.7	41.2-46.4	257.4	34.8	32.9-36.9
Melanoma of the skin	1976-1980	48.6	4.9	4.3-5.6	23.2	5.2	4.3-6.3	25.4	4.8	4.0-5.8	18.4	1.8	1.5-2.3	10.0	2.2	1.6-3.0	8.4	1.6	1.1-2.1
	1981-1985	61.0	6.2	5.5-7.0	30.6	7.0	5.9-8.3	30.4	5.9	4.9-6.9	19.6	2.0	1.6-2.4	11.8	2.7	2.1-3.6	7.8	1.5	1.0-2.0
	1986-1990	67.8	6.7	6.0-7.4	39.8	8.7	7.5-10.0	28.0	5.1	4.3-6.1	24.4	2.4	1.9-2.8	14.8	3.3	2.5-4.2	9.6	1.7	1.2-2.3
	1991-1995	86.2	8.3	7.6-9.2	50.4	11.1	9.7-12.6	35.8	6.5	5.6-7.6	24.8	2.4	2.0-2.8	13.8	3.1	2.4-4.0	11.0	1.9	1.4-2.5
	1996-2000	114.6	11.0	10.2-12.0	61.8	13.6	12.1-15.2	52.8	9.4	8.3-10.6	22.6	2.1	1.7-2.5	13.6	3.0	2.3-3.8	9.0	1.5	1.1-2.0
	2001-2005	154.0	14.5	13.5-15.6	87.0	18.3	16.6-20.1	67.0	11.8	10.5-13.2	25.4	2.3	1.9-2.7	13.6	2.9	2.2-3.7	11.8	1.8	1.3-2.3
	2006-2010	226.0	21.1	19.9-22.4	132.2	27.6	25.5-29.8	93.8	16.7	15.1-18.3	32.0	2.9	2.5-3.4	20.6	4.3	3.5-5.2	11.4	1.8	1.3-2.3
	2011-2015	304.0	27.4	26.0-28.8	173.0	34.2	31.9-36.6	131.0	23.0	21.1-24.9	27.8	2.4	2.0-2.9	18.2	3.7	3.0-4.6	9.6	1.4	1.0-2.0
	2016-2020	354.4	30.0	28.6-31.5	200.8	36.7	34.4-39.1	153.6	25.6	23.7-27.6	28.8	2.3	1.9-2.7	18.2	3.4	2.7-4.2	10.6	1.4	1.0-1.9
Female breast	1976-1980							569.0	103.7	99.8-107.7							225.4	39.7	37.4-42.2
	1981-1985							608.0	110.1	106.0-114.2							231.2	40.0	37.6-42.4
	1986-1990							711.6	126.7	122.4-131.1							238.8	40.8	38.4-43.3
	1991-1995							732.0	124.5	120.4-128.7							233.4	38.0	35.8-40.3
	1996-2000							815.2	137.6	133.3-142.0							199.6	31.1	29.2-33.1
	2001-2005							800.0	132.5	128.4-136.8							183.6	27.8	26.0-29.7
	2006-2010							844.6	140.2	135.9-144.6							176.8	26.9	25.1-28.8
	2011-2015							903.8	148.6	144.1-153.2							166.2	24.4	22.7-26.2
	2016-2020							885.8	141.6	137.2-146.1							152.4	21.7	20.1-23.4
Cervix uteri	1976-1980							56.4	10.4	9.2-11.7							24.8	4.3	3.6-5.2
	1981-1985							51.4	9.6	8.4-10.9							17.8	3.2	2.6-4.0
	1986-1990							43.8	8.3	7.2-9.5							19.8	3.4	2.8-4.2
	1991-1995							43.8	7.8	6.8-9.0							13.8	2.3	1.8-3.0
	1996-2000							36.8	7.0	6.0-8.1							13.2	2.3	1.8-2.9
	2001-2005							34.4	6.7	5.7-7.8							10.8	2.0	1.5-2.6
	2006-2010							35.2	7.0	5.9-8.1							9.0	1.6	1.2-2.2
	2011-2015							35.2	7.0	5.9-8.2							11.2	2.0	1.5-2.6
	2016-2020							34.6	7.1	6.0-8.3							12.0	2.0	1.5-2.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Erie**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							143.8	24.4	22.6-26.3							30.6	5.2	4.4-6.1
	1981-1985							143.8	24.8	23.0-26.7							31.2	5.3	4.5-6.2
	1986-1990							145.6	24.1	22.3-26.0							33.6	5.3	4.5-6.1
	1991-1995							160.6	26.8	24.9-28.8							31.4	4.6	3.9-5.5
	1996-2000							164.6	27.4	25.6-29.4							27.8	4.2	3.5-5.0
	2001-2005							169.8	28.1	26.3-30.1							29.6	4.5	3.8-5.3
	2006-2010							176.2	28.6	26.7-30.6							30.6	4.5	3.8-5.3
	2011-2015							201.2	31.5	29.5-33.6							34.0	5.0	4.2-5.8
	2016-2020							220.6	33.3	31.2-35.4							38.4	5.4	4.7-6.3
Ovary	1976-1980							97.6	17.3	15.7-18.9							69.0	11.7	10.5-13.0
	1981-1985							104.6	18.6	17.0-20.4							69.4	11.7	10.4-13.0
	1986-1990							103.8	18.4	16.8-20.2							68.0	10.9	9.8-12.2
	1991-1995							113.8	19.5	17.9-21.2							67.8	10.7	9.5-11.9
	1996-2000							98.6	16.5	15.1-18.1							63.4	9.7	8.7-10.9
	2001-2005							87.4	14.4	13.0-15.9							60.8	9.1	8.1-10.2
	2006-2010							80.6	13.2	11.9-14.6							55.4	8.5	7.5-9.6
	2011-2015							81.0	13.3	11.9-14.7							49.4	7.4	6.4-8.4
	2016-2020							71.4	11.1	9.9-12.4							55.4	8.1	7.2-9.2
Prostate	1976-1980				297.4	81.8	77.4-86.4							106.4	32.7	29.8-35.7			
	1981-1985				350.2	92.9	88.3-97.6							104.2	30.3	27.6-33.2			
	1986-1990				439.0	107.1	102.4-112.0							126.8	36.4	33.4-39.5			
	1991-1995				654.8	144.4	139.3-149.6							149.8	39.0	36.1-42.0			
	1996-2000				805.0	171.2	165.9-176.7							137.0	32.6	30.2-35.2			
	2001-2005				882.2	185.9	180.5-191.5							117.8	27.3	25.1-29.6			
	2006-2010				920.0	184.7	179.4-190.2							100.6	21.9	20.0-23.9			
	2011-2015				890.0	163.1	158.2-168.0							93.4	19.1	17.4-20.9			
	2016-2020				854.8	141.8	137.5-146.2							98.0	18.7	17.0-20.4			
Testis	1976-1980				22.0	4.5	3.6-5.4							2.8	0.5	0.3-0.9			
	1981-1985				22.2	4.6	3.7-5.6							1.8	0.4	0.2-0.9			
	1986-1990				29.2	5.8	4.9-6.9							2.0	0.4	0.2-0.7			
	1991-1995				27.6	5.8	4.9-6.9							2.4	0.5	0.3-0.9			
	1996-2000				21.6	4.7	3.9-5.7							1.2	0.3	0.1-0.6			
	2001-2005				22.8	5.3	4.4-6.4							1.2	0.3	0.1-0.6			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Erie**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				28.6	6.8	5.7-8.1							0.8	0.2	0.0-0.5			
	2011-2015				27.4	6.4	5.3-7.6							1.2	0.3	0.1-0.6			
	2016-2020				28.2	6.6	5.5-7.8							0.6	0.1	0.0-0.4			
Urinary bladder (incl. in situ)	1976-1980	226.4	23.3	21.9-24.7	165.4	42.9	39.8-46.2	61.0	10.7	9.5-12.0	58.6	6.4	5.7-7.2	40.8	12.4	10.6-14.3	17.8	3.1	2.5-3.9
	1981-1985	235.8	23.1	21.7-24.5	170.6	42.1	39.1-45.2	65.2	11.1	9.9-12.4	56.2	5.8	5.1-6.5	39.4	11.5	9.8-13.3	16.8	2.8	2.2-3.4
	1986-1990	249.2	23.5	22.2-24.8	176.6	41.7	38.9-44.8	72.6	11.9	10.6-13.2	59.2	5.9	5.2-6.6	39.4	11.0	9.4-12.7	19.8	3.1	2.5-3.8
	1991-1995	265.4	24.0	22.7-25.4	190.2	42.9	40.1-45.8	75.2	11.7	10.5-12.9	59.0	5.3	4.7-5.9	40.4	9.8	8.4-11.3	18.6	2.7	2.2-3.4
	1996-2000	301.0	26.6	25.3-28.0	217.2	47.8	45.0-50.8	83.8	12.3	11.2-13.6	64.2	5.5	4.9-6.2	44.0	10.3	8.9-11.7	20.2	2.7	2.2-3.3
	2001-2005	288.2	25.1	23.8-26.5	203.0	43.7	41.1-46.5	85.2	12.6	11.4-13.9	59.4	4.9	4.3-5.5	36.0	8.2	7.0-9.5	23.4	3.1	2.5-3.7
	2006-2010	369.2	31.9	30.4-33.4	271.8	56.8	53.8-59.9	97.4	14.4	13.1-15.8	62.0	5.0	4.5-5.6	41.6	8.9	7.7-10.2	20.4	2.7	2.2-3.3
	2011-2015	357.8	29.9	28.5-31.4	256.0	51.1	48.3-54.0	101.8	14.8	13.5-16.2	66.6	5.3	4.7-5.9	44.8	9.0	7.9-10.3	21.8	2.9	2.3-3.5
2016-2020	343.4	27.0	25.7-28.3	242.0	44.4	41.8-47.0	101.4	13.9	12.7-15.3	60.8	4.7	4.2-5.3	40.2	7.8	6.7-9.0	20.6	2.7	2.2-3.3	
Kidney and renal pelvis	1976-1980	77.6	7.8	7.0-8.7	46.4	10.8	9.4-12.4	31.2	5.5	4.6-6.4	41.0	4.1	3.5-4.7	26.2	6.3	5.2-7.5	14.8	2.5	2.0-3.2
	1981-1985	97.6	9.6	8.7-10.5	61.8	14.4	12.7-16.2	35.8	6.3	5.4-7.3	45.2	4.5	3.9-5.1	26.2	6.3	5.2-7.5	19.0	3.2	2.6-4.0
	1986-1990	92.2	8.9	8.1-9.8	53.8	12.3	10.8-14.0	38.4	6.5	5.6-7.6	42.2	4.0	3.5-4.6	25.4	5.8	4.8-7.0	16.8	2.7	2.2-3.4
	1991-1995	113.0	10.6	9.8-11.6	65.4	14.5	12.9-16.2	47.6	7.7	6.7-8.8	46.2	4.3	3.7-4.9	25.0	5.6	4.6-6.7	21.2	3.2	2.6-3.9
	1996-2000	135.0	12.4	11.4-13.3	81.4	17.4	15.7-19.2	53.6	8.4	7.4-9.5	44.4	4.0	3.5-4.5	27.8	6.1	5.2-7.3	16.6	2.4	1.9-3.0
	2001-2005	163.6	14.8	13.8-15.8	99.6	21.1	19.3-23.1	64.0	10.1	9.0-11.3	51.8	4.5	4.0-5.1	31.6	6.8	5.8-8.0	20.2	3.0	2.4-3.6
	2006-2010	188.2	17.0	15.9-18.1	121.0	24.8	22.8-26.8	67.2	10.8	9.7-12.1	48.6	4.1	3.6-4.7	32.2	6.7	5.7-7.8	16.4	2.4	1.9-3.0
	2011-2015	215.0	18.6	17.4-19.7	134.2	26.0	24.0-28.1	80.8	12.6	11.3-13.9	45.6	3.7	3.2-4.2	28.4	5.5	4.6-6.5	17.2	2.4	1.9-3.0
2016-2020	222.8	18.6	17.4-19.7	139.8	25.7	23.8-27.8	83.0	12.8	11.5-14.1	39.2	3.1	2.7-3.6	22.8	4.1	3.4-5.0	16.4	2.3	1.8-2.9	
Brain and other central nervous system	1976-1980	62.2	5.9	5.3-6.6	35.4	7.4	6.3-8.7	26.8	4.8	4.0-5.7	44.2	4.1	3.6-4.7	24.6	5.3	4.4-6.4	19.6	3.3	2.7-4.1
	1981-1985	60.0	5.8	5.2-6.5	32.2	6.9	5.9-8.1	27.8	4.9	4.1-5.8	38.6	3.7	3.2-4.3	20.6	4.5	3.6-5.5	18.0	3.1	2.5-3.8
	1986-1990	70.4	6.9	6.2-7.7	35.2	7.7	6.6-9.0	35.2	6.3	5.4-7.4	42.0	4.0	3.5-4.6	21.6	4.6	3.8-5.6	20.4	3.5	2.8-4.2
	1991-1995	79.4	7.7	6.9-8.5	44.6	9.9	8.6-11.3	34.8	5.9	5.0-6.9	50.4	4.8	4.2-5.4	28.2	6.4	5.3-7.5	22.2	3.6	3.0-4.4
	1996-2000	76.6	7.4	6.7-8.2	43.2	9.3	8.1-10.7	33.4	6.0	5.1-7.0	43.4	4.0	3.5-4.6	26.4	5.7	4.7-6.8	17.0	2.8	2.2-3.5
	2001-2005	85.2	8.3	7.5-9.1	44.4	9.6	8.4-10.9	40.8	7.2	6.3-8.4	50.8	4.7	4.1-5.3	27.8	5.9	5.0-7.0	23.0	3.7	3.0-4.5
	2006-2010	80.0	7.8	7.1-8.6	43.0	9.2	8.0-10.5	37.0	6.8	5.8-7.9	47.8	4.4	3.8-5.0	26.0	5.4	4.5-6.4	21.8	3.6	2.9-4.4
	2011-2015	75.8	7.4	6.7-8.2	40.4	8.2	7.1-9.5	35.4	6.8	5.8-8.0	48.6	4.3	3.8-4.9	26.0	5.1	4.2-6.0	22.6	3.6	3.0-4.4
2016-2020	76.0	7.3	6.5-8.1	43.8	8.9	7.7-10.2	32.2	6.0	5.0-7.1	50.0	4.2	3.7-4.8	28.4	5.4	4.5-6.4	21.6	3.3	2.7-4.0	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Erie**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	41.8	4.4	3.8-5.0	13.8	3.2	2.5-4.1	28.0	5.4	4.5-6.4	7.8	0.8	0.6-1.2	3.4	0.9	0.5-1.5	4.4	0.8	0.5-1.2
	1981-1985	42.6	4.4	3.8-5.1	11.4	2.6	1.9-3.4	31.2	6.1	5.1-7.1	7.2	0.7	0.5-1.0	2.2	0.5	0.2-1.0	5.0	0.9	0.6-1.3
	1986-1990	45.4	4.7	4.1-5.4	11.4	2.5	1.9-3.3	34.0	6.6	5.6-7.7	4.4	0.4	0.3-0.7	1.4	0.3	0.1-0.7	3.0	0.5	0.3-0.8
	1991-1995	61.8	6.2	5.5-6.9	15.2	3.4	2.7-4.3	46.6	8.7	7.6-9.9	5.8	0.5	0.4-0.8	1.6	0.4	0.2-0.8	4.2	0.6	0.4-0.9
	1996-2000	77.6	7.9	7.1-8.7	18.6	4.0	3.2-4.9	59.0	11.5	10.2-12.9	5.0	0.4	0.3-0.6	1.4	0.3	0.1-0.6	3.6	0.5	0.3-0.8
	2001-2005	97.6	9.9	9.1-10.9	22.0	4.7	3.9-5.7	75.6	14.9	13.4-16.5	3.8	0.3	0.2-0.5	1.8	0.4	0.2-0.7	2.0	0.3	0.1-0.6
	2006-2010	154.4	15.9	14.7-17.1	34.4	7.4	6.3-8.6	120.0	23.9	22.0-26.0	6.6	0.5	0.4-0.8	2.4	0.5	0.2-0.8	4.2	0.6	0.4-0.9
	2011-2015	195.2	20.0	18.8-21.4	46.6	9.6	8.3-10.9	148.6	30.0	27.8-32.3	5.8	0.5	0.3-0.7	2.4	0.4	0.2-0.8	3.4	0.5	0.3-0.8
	2016-2020	146.2	15.3	14.2-16.5	42.8	8.9	7.7-10.3	103.4	21.5	19.6-23.5	4.4	0.4	0.2-0.6	2.2	0.5	0.2-0.9	2.2	0.3	0.1-0.5
Hodgkin lymphoma	1976-1980	36.8	3.4	2.9-3.9	20.4	4.1	3.3-5.0	16.4	2.8	2.2-3.5	12.8	1.2	0.9-1.6	7.0	1.5	1.0-2.1	5.8	1.0	0.7-1.5
	1981-1985	38.0	3.6	3.1-4.2	21.6	4.5	3.7-5.5	16.4	2.8	2.2-3.5	9.6	0.9	0.7-1.2	6.0	1.3	0.9-1.9	3.6	0.6	0.3-0.9
	1986-1990	33.2	3.2	2.7-3.8	18.4	3.8	3.1-4.7	14.8	2.7	2.1-3.4	10.8	1.1	0.8-1.5	7.2	1.6	1.1-2.2	3.6	0.7	0.4-1.1
	1991-1995	35.6	3.6	3.1-4.1	19.8	4.2	3.4-5.1	15.8	3.0	2.4-3.8	10.6	1.0	0.8-1.3	6.2	1.3	0.9-1.9	4.4	0.7	0.5-1.1
	1996-2000	30.2	3.2	2.7-3.7	16.8	3.7	2.9-4.5	13.4	2.7	2.1-3.4	5.6	0.6	0.4-0.8	3.0	0.6	0.4-1.1	2.6	0.4	0.2-0.8
	2001-2005	29.6	3.1	2.7-3.7	14.8	3.3	2.6-4.1	14.8	3.0	2.4-3.8	5.2	0.5	0.3-0.8	2.8	0.6	0.3-1.0	2.4	0.4	0.2-0.8
	2006-2010	31.0	3.2	2.7-3.7	16.4	3.6	2.8-4.5	14.6	2.9	2.2-3.6	4.2	0.4	0.2-0.6	1.8	0.4	0.2-0.7	2.4	0.4	0.2-0.7
	2011-2015	33.4	3.6	3.0-4.2	17.2	3.7	3.0-4.6	16.2	3.5	2.8-4.4	5.0	0.5	0.3-0.7	2.4	0.5	0.3-0.9	2.6	0.5	0.2-0.8
	2016-2020	29.6	3.0	2.5-3.6	17.4	3.8	3.0-4.7	12.2	2.4	1.8-3.1	2.4	0.2	0.1-0.3	1.2	0.2	0.1-0.5	1.2	0.2	0.1-0.4
Non-Hodgkin Lymphomas	1976-1980	114.6	11.3	10.4-12.3	59.8	13.7	12.1-15.5	54.8	9.4	8.3-10.7	66.6	6.6	5.9-7.4	38.4	9.3	7.9-10.8	28.2	4.7	4.0-5.6
	1981-1985	140.6	14.1	13.0-15.2	70.4	17.2	15.3-19.2	70.2	11.9	10.7-13.3	73.4	7.3	6.5-8.1	34.4	8.6	7.3-10.0	39.0	6.4	5.5-7.4
	1986-1990	164.2	16.1	15.0-17.2	80.0	18.7	16.8-20.7	84.2	14.2	12.8-15.7	87.8	8.6	7.8-9.5	46.8	11.4	9.9-13.1	41.0	6.7	5.8-7.7
	1991-1995	182.2	17.2	16.1-18.4	91.6	20.7	18.8-22.7	90.6	14.6	13.3-16.1	96.6	9.0	8.2-9.8	47.8	11.1	9.7-12.7	48.8	7.4	6.5-8.4
	1996-2000	200.8	18.4	17.3-19.6	97.2	21.3	19.4-23.3	103.6	16.3	14.9-17.8	102.4	9.0	8.3-9.9	47.0	10.5	9.2-12.0	55.4	8.1	7.2-9.2
	2001-2005	230.0	20.9	19.7-22.2	120.2	25.7	23.7-27.9	109.8	17.4	15.9-18.9	84.8	7.4	6.7-8.1	41.6	9.0	7.8-10.3	43.2	6.2	5.4-7.2
	2006-2010	244.4	22.0	20.8-23.3	125.6	26.3	24.2-28.4	118.8	18.7	17.2-20.4	82.4	7.0	6.3-7.7	42.0	8.9	7.7-10.2	40.4	5.8	5.0-6.7
	2011-2015	255.8	22.4	21.1-23.7	137.4	27.1	25.0-29.2	118.4	18.9	17.3-20.6	75.0	6.3	5.6-6.9	38.6	8.0	6.9-9.2	36.4	5.0	4.3-5.9
	2016-2020	277.6	23.0	21.8-24.3	148.6	27.9	25.9-30.1	129.0	19.4	17.8-21.0	72.6	5.7	5.1-6.3	45.2	8.6	7.5-9.8	27.4	3.8	3.2-4.5
Myeloma	1976-1980	46.4	4.7	4.1-5.4	25.4	6.5	5.3-7.8	21.0	3.6	2.9-4.4	33.4	3.4	2.9-4.0	18.2	4.7	3.7-5.9	15.2	2.6	2.0-3.2
	1981-1985	57.4	5.6	5.0-6.3	28.4	6.8	5.7-8.1	29.0	4.8	4.0-5.7	40.2	4.0	3.4-4.6	20.8	5.0	4.0-6.1	19.4	3.2	2.6-4.0
	1986-1990	60.8	5.8	5.1-6.5	28.8	6.8	5.7-8.1	32.0	5.0	4.3-5.9	34.4	3.3	2.8-3.8	18.6	4.5	3.6-5.5	15.8	2.4	1.9-3.0
	1991-1995	70.6	6.3	5.6-7.0	35.0	7.8	6.6-9.0	35.6	5.2	4.5-6.1	49.2	4.4	3.8-5.0	25.4	5.7	4.7-6.8	23.8	3.5	2.9-4.2
	1996-2000	75.2	6.6	5.9-7.3	37.0	7.9	6.8-9.2	38.2	5.7	4.9-6.6	47.6	4.1	3.6-4.7	22.0	4.9	4.0-5.9	25.6	3.6	3.0-4.3
	2001-2005	81.4	7.1	6.5-7.9	44.6	9.6	8.4-10.9	36.8	5.5	4.7-6.4	42.2	3.6	3.2-4.2	21.8	4.7	3.9-5.7	20.4	3.0	2.5-3.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Erie**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	86.8	7.5	6.8-8.3	43.8	9.0	7.8-10.2	43.0	6.5	5.6-7.5	43.8	3.6	3.2-4.2	20.0	4.1	3.4-5.0	23.8	3.3	2.7-4.0
	2011-2015	89.4	7.5	6.8-8.2	48.2	9.6	8.4-10.9	41.2	6.0	5.2-6.9	45.0	3.6	3.2-4.1	23.6	4.6	3.8-5.6	21.4	3.0	2.4-3.7
	2016-2020	101.2	8.2	7.5-9.0	51.0	9.5	8.3-10.8	50.2	7.4	6.4-8.4	35.6	2.8	2.4-3.3	19.4	3.8	3.0-4.6	16.2	2.2	1.7-2.7
Leukemias	1976-1980	130.2	13.4	12.4-14.5	74.8	19.1	17.0-21.3	55.4	9.9	8.7-11.2	86.8	9.0	8.1-9.9	48.4	12.5	10.9-14.3	38.4	6.8	5.8-7.8
	1981-1985	136.0	13.6	12.6-14.7	71.4	17.3	15.5-19.4	64.6	11.3	10.1-12.6	86.0	8.6	7.8-9.5	45.2	11.2	9.7-12.9	40.8	7.1	6.1-8.1
	1986-1990	136.0	13.4	12.4-14.5	77.6	19.3	17.3-21.5	58.4	9.8	8.7-11.0	76.2	7.4	6.7-8.2	43.0	11.0	9.5-12.6	33.2	5.4	4.6-6.3
	1991-1995	139.0	13.1	12.1-14.1	72.8	16.8	15.0-18.7	66.2	10.6	9.5-11.8	81.8	7.6	6.9-8.4	45.6	10.8	9.4-12.3	36.2	5.7	4.9-6.6
	1996-2000	160.0	14.5	13.5-15.5	90.0	19.8	18.0-21.8	70.0	10.8	9.6-12.0	85.2	7.5	6.8-8.2	47.4	10.5	9.2-12.0	37.8	5.5	4.7-6.4
	2001-2005	183.6	16.3	15.3-17.5	98.6	21.3	19.5-23.3	85.0	12.8	11.6-14.2	87.6	7.5	6.8-8.3	46.2	10.2	9.0-11.7	41.4	5.8	5.0-6.7
	2006-2010	191.0	17.3	16.2-18.4	105.8	22.3	20.4-24.3	85.2	13.5	12.2-15.0	89.8	7.7	7.0-8.5	51.2	10.8	9.5-12.3	38.6	5.5	4.7-6.4
	2011-2015	197.8	17.4	16.3-18.5	119.4	24.0	22.1-26.1	78.4	12.4	11.1-13.7	86.0	7.2	6.5-7.9	47.0	9.7	8.5-11.1	39.0	5.4	4.6-6.3
	2016-2020	214.8	18.1	17.0-19.2	130.4	24.7	22.8-26.7	84.4	12.8	11.6-14.2	83.0	6.6	5.9-7.3	49.8	9.7	8.5-11.0	33.2	4.3	3.7-5.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Essex**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	151.6	404.3	375.2-434.9	83.8	516.0	465.6-570.2	67.8	329.6	294.2-368.1	83.6	223.1	201.7-246.1	43.6	270.3	234.0-310.4	40.0	191.4	165.1-220.9
	1981-1985	184.4	467.4	437.1-499.4	98.8	591.1	538.3-647.7	85.6	390.2	352.8-430.7	104.4	261.7	239.3-285.7	58.4	343.7	303.9-387.1	46.0	203.5	177.3-232.7
	1986-1990	189.0	472.7	442.6-504.3	99.2	584.1	532.4-639.3	89.8	405.7	367.9-446.4	96.8	241.2	219.9-264.0	51.2	309.1	271.4-350.5	45.6	198.7	173.1-227.2
	1991-1995	228.8	536.9	505.9-569.3	117.6	646.3	594.0-701.9	111.2	475.5	435.8-518.1	102.0	237.1	216.8-258.8	49.6	284.4	249.4-322.9	52.4	213.1	187.5-241.4
	1996-2000	227.6	499.8	471.0-530.0	120.0	592.2	545.0-642.4	107.6	443.3	405.7-483.6	100.6	217.2	198.6-237.2	53.0	275.1	242.4-311.0	47.6	182.2	159.3-207.7
	2001-2005	237.8	484.5	457.2-513.1	126.6	568.1	524.2-614.9	111.2	432.7	396.5-471.6	95.8	190.6	173.9-208.7	52.2	247.4	217.8-280.0	43.6	155.0	134.7-177.9
	2006-2010	264.0	497.4	470.5-525.7	144.6	575.3	533.4-619.8	119.4	438.9	402.9-477.5	95.6	174.7	159.2-191.5	50.6	207.2	182.1-235.0	45.0	151.2	131.5-173.4
	2011-2015	265.2	465.2	439.6-492.2	136.6	502.3	464.2-543.1	128.6	444.8	409.0-483.4	95.6	159.4	145.1-174.9	49.8	184.9	162.2-210.3	45.8	142.2	123.8-163.1
	2016-2020	251.2	409.8	386.0-434.9	138.8	456.7	421.8-494.2	112.4	373.5	340.5-409.4	100.6	155.4	141.6-170.4	54.6	183.2	161.7-207.4	46.0	137.5	119.0-158.7
Oral cavity and pharynx	1976-1980	2.2	5.7	2.8-10.3	1.4	7.8	3.0-16.5	0.8	3.8	1.0-10.3	1.4	3.7	1.5-7.9	1.0	7.2	2.2-16.9	0.4	1.7	0.2-6.9
	1981-1985	4.0	9.7	5.9-15.2	3.0	17.6	9.6-29.4	1.0	4.1	1.3-10.1	1.6	3.7	1.6-7.4	1.0	5.9	1.8-14.0	0.6	2.3	0.5-7.5
	1986-1990	3.8	9.5	5.7-15.0	2.2	12.9	6.4-23.2	1.6	6.5	2.8-13.4	1.0	2.8	0.9-6.6	0.6	4.5	0.9-12.5	0.4	1.8	0.2-7.1
	1991-1995	6.2	14.9	10.1-21.3	4.0	20.8	12.6-32.4	2.2	9.6	4.7-17.7	1.0	2.4	0.8-5.7	0.4	2.3	0.3-8.1	0.6	2.2	0.4-7.3
	1996-2000	4.4	9.8	6.1-14.9	3.2	15.6	8.8-25.6	1.2	4.2	1.5-10.1	0.6	1.3	0.3-3.9	0.2	1.1	0.0-6.0	0.4	1.3	0.2-5.9
	2001-2005	6.6	13.4	9.2-19.0	3.8	16.2	9.7-25.6	2.8	10.4	5.6-18.1	1.2	2.4	0.9-5.5	0.6	2.6	0.5-8.0	0.6	2.3	0.4-7.5
	2006-2010	6.0	11.5	7.7-16.8	3.6	15.7	9.2-25.2	2.4	7.5	3.8-14.1	0.8	1.3	0.4-3.8	0.2	0.6	0.0-4.8	0.6	1.8	0.4-6.6
	2011-2015	7.6	13.4	9.4-18.8	5.2	18.4	11.8-27.8	2.4	8.1	4.0-15.4	2.8	4.8	2.6-8.5	2.0	7.2	3.4-14.0	0.8	2.7	0.7-8.3
	2016-2020	4.8	7.6	4.8-12.0	3.2	11.0	6.1-18.9	1.6	4.3	1.9-10.2	0.6	1.0	0.2-3.6	0.6	2.1	0.4-7.4	0.0	0.0	0.0-4.9
Esophagus	1976-1980	0.8	2.1	0.6-5.6	0.6	3.9	0.8-11.6	0.2	0.9	0.0-5.7	0.4	1.1	0.1-4.0	0.2	1.7	0.0-8.5	0.2	0.9	0.0-5.7
	1981-1985	2.6	5.8	3.1-10.2	2.0	10.8	5.0-20.3	0.6	2.3	0.5-7.5	2.6	5.8	3.1-10.1	2.0	9.6	4.6-18.2	0.6	2.5	0.5-7.9
	1986-1990	1.6	4.0	1.7-8.0	1.4	8.1	3.2-17.0	0.2	0.7	0.0-5.1	1.8	4.6	2.1-8.8	1.4	8.3	3.3-17.3	0.4	1.4	0.2-6.1
	1991-1995	1.8	4.6	2.1-8.7	1.6	9.7	4.1-19.0	0.2	0.6	0.0-4.8	1.8	4.5	2.1-8.7	1.2	6.9	2.5-15.0	0.6	1.9	0.4-6.7
	1996-2000	3.4	7.3	4.2-11.8	2.6	12.1	6.4-21.0	0.8	2.7	0.7-7.9	3.8	8.2	4.9-12.9	2.8	13.3	7.2-22.6	1.0	3.9	1.2-9.8
	2001-2005	3.2	6.4	3.7-10.6	2.6	11.6	6.2-20.1	0.6	2.0	0.4-6.9	3.4	6.9	4.0-11.2	2.6	10.9	5.8-19.0	0.8	3.1	0.8-8.7
	2006-2010	7.0	12.5	8.7-17.7	6.6	25.5	17.4-36.3	0.4	1.3	0.2-6.0	5.6	9.9	6.5-14.6	5.2	20.2	13.1-30.1	0.4	1.1	0.1-5.7
	2011-2015	2.2	3.6	1.8-6.9	1.6	5.3	2.3-11.4	0.6	2.1	0.4-7.4	2.6	4.2	2.2-7.7	2.0	7.1	3.4-13.9	0.6	1.9	0.4-7.0
	2016-2020	5.2	7.8	5.1-12.0	3.8	11.5	6.9-19.0	1.4	4.3	1.7-10.4	4.0	6.0	3.7-9.9	3.0	10.2	5.7-17.8	1.0	2.5	0.8-7.9
Stomach	1976-1980	2.2	5.9	2.9-10.7	1.6	9.9	4.1-19.9	0.6	2.5	0.5-8.1	1.4	3.5	1.4-7.4	1.0	6.4	1.9-15.4	0.4	1.5	0.2-6.5
	1981-1985	3.2	8.2	4.6-13.5	1.8	11.7	5.2-22.3	1.4	5.7	2.3-12.2	2.0	5.2	2.4-9.8	1.4	9.0	3.5-18.7	0.6	2.4	0.5-7.7
	1986-1990	3.2	8.0	4.5-13.1	1.8	11.8	5.3-22.3	1.4	5.3	2.1-11.7	2.0	4.6	2.2-8.6	0.6	3.2	0.6-9.6	1.4	5.3	2.1-11.5
	1991-1995	2.8	6.6	3.6-11.2	1.8	10.4	4.7-19.7	1.0	4.0	1.3-10.1	2.2	5.2	2.6-9.4	1.0	6.2	2.0-14.3	1.2	4.6	1.6-10.7
	1996-2000	3.8	8.2	4.9-12.9	2.2	10.5	5.2-19.0	1.6	5.4	2.2-11.5	1.2	2.7	1.0-6.1	0.8	4.2	1.1-10.8	0.4	1.5	0.2-6.3
	2001-2005	4.4	8.7	5.5-13.4	2.8	12.1	6.5-20.7	1.6	5.7	2.4-12.0	1.6	3.2	1.4-6.5	0.8	4.0	1.0-10.4	0.8	2.5	0.7-7.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Essex**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	3.0	5.6	3.1-9.5	1.8	7.4	3.3-14.4	1.2	3.9	1.4-9.6	1.6	2.9	1.2-6.0	1.0	4.0	1.3-9.9	0.6	1.4	0.3-5.9
	2011-2015	2.6	4.4	2.3-8.0	1.4	5.1	2.0-11.3	1.2	3.7	1.3-9.5	0.4	0.8	0.1-3.3	0.2	0.8	0.0-5.3	0.2	0.8	0.0-5.6
	2016-2020	3.8	5.7	3.4-9.5	3.4	10.7	6.1-18.2	0.4	1.2	0.2-6.4	0.6	1.0	0.2-3.7	0.4	1.4	0.1-6.5	0.2	0.6	0.0-5.7
Colorectal	1976-1980	20.6	55.2	44.9-67.2	8.8	55.6	39.8-75.5	11.8	53.5	40.6-69.4	12.2	32.8	24.9-42.3	5.0	29.3	18.6-43.8	7.2	33.5	23.3-46.8
	1981-1985	31.2	78.1	66.1-91.7	15.6	96.6	75.6-121.5	15.6	65.1	51.3-81.8	14.8	37.4	29.3-47.2	5.6	33.6	21.9-49.0	9.2	37.9	27.7-51.0
	1986-1990	28.0	68.6	57.6-81.1	13.6	79.5	61.2-101.3	14.4	58.6	45.7-74.4	12.2	30.5	23.3-39.3	5.4	35.0	22.7-51.1	6.8	28.1	19.4-39.8
	1991-1995	29.8	68.8	58.1-80.9	14.6	85.1	66.3-107.4	15.2	60.1	47.1-76.0	12.6	29.8	22.9-38.3	6.4	39.1	26.5-55.3	6.2	25.2	16.9-36.3
	1996-2000	31.0	66.6	56.5-78.2	18.2	89.5	71.8-110.3	12.8	49.5	37.8-64.0	13.0	27.9	21.5-35.8	6.8	35.2	24.2-49.5	6.2	24.0	16.1-34.8
	2001-2005	31.8	63.3	53.8-74.1	14.8	69.0	54.0-87.0	17.0	60.3	47.9-75.3	13.8	27.2	21.1-34.6	6.6	31.4	21.5-44.3	7.2	24.4	17.0-34.5
	2006-2010	25.4	46.1	38.3-55.1	14.6	56.4	44.0-71.5	10.8	35.3	26.3-47.0	9.2	16.2	11.9-21.9	5.8	24.1	16.0-35.0	3.4	10.4	6.0-17.6
	2011-2015	23.2	39.9	32.8-48.4	13.4	50.9	39.3-65.3	9.8	32.3	23.2-44.2	8.0	13.7	9.7-19.1	4.2	16.2	9.9-25.4	3.8	12.4	7.1-20.8
2016-2020	19.8	35.9	28.6-44.7	9.4	36.1	25.8-49.5	10.4	35.2	25.5-47.9	8.4	14.2	10.0-19.9	4.4	16.3	10.0-25.6	4.0	12.1	7.0-20.4	
Colon excluding rectum	1976-1980	16.6	44.4	35.2-55.3	6.8	43.4	29.5-61.3	9.8	43.6	32.2-58.0	9.4	25.4	18.5-34.0	4.0	24.4	14.6-38.2	5.4	24.8	16.2-36.5
	1981-1985	22.0	55.4	45.4-67.0	10.0	64.2	47.0-85.2	12.0	50.3	38.2-65.3	12.4	31.5	24.0-40.5	5.0	30.6	19.4-45.6	7.4	30.6	21.5-42.6
	1986-1990	21.4	52.3	42.8-63.4	9.4	55.4	40.4-74.1	12.0	47.7	36.3-61.9	11.0	27.6	20.7-36.0	5.0	33.2	21.2-49.2	6.0	24.6	16.5-35.7
	1991-1995	20.6	47.6	38.8-57.8	9.2	54.1	39.4-72.4	11.4	44.9	33.8-58.8	10.6	25.0	18.7-32.8	4.8	29.7	18.9-44.3	5.8	23.2	15.4-33.9
	1996-2000	23.6	50.4	41.6-60.5	13.6	67.0	51.8-85.4	10.0	38.4	28.2-51.4	11.8	25.2	19.2-32.7	6.2	32.3	21.8-46.2	5.6	21.3	14.0-31.4
	2001-2005	24.2	48.1	39.9-57.6	10.0	46.2	34.2-61.2	14.2	49.7	38.6-63.4	10.6	20.7	15.5-27.3	4.6	21.9	13.8-33.2	6.0	19.8	13.3-29.1
	2006-2010	17.8	32.3	25.8-40.0	9.4	37.1	27.1-49.9	8.4	27.4	19.5-38.0	7.4	13.0	9.1-18.2	4.6	19.7	12.4-29.9	2.8	8.6	4.6-15.4
	2011-2015	15.8	26.9	21.2-34.1	7.8	29.4	20.8-40.7	8.0	26.8	18.5-38.1	6.4	10.9	7.3-15.8	3.4	13.3	7.6-21.8	3.0	10.0	5.2-18.0
2016-2020	14.8	25.9	19.9-33.4	5.6	21.3	13.6-32.3	9.2	30.3	21.6-42.0	6.4	11.3	7.5-16.6	3.4	13.2	7.5-22.1	3.0	9.4	4.9-17.3	
Rectum & rectosigmoid	1976-1980	4.0	10.8	6.5-16.8	2.0	12.3	5.6-23.1	2.0	9.8	4.6-18.5	2.8	7.4	4.0-12.6	1.0	4.9	1.6-12.1	1.8	8.7	3.8-16.9
	1981-1985	9.2	22.7	16.5-30.5	5.6	32.5	21.2-47.6	3.6	14.8	8.8-23.9	2.4	6.0	3.1-10.5	0.6	3.0	0.6-9.3	1.8	7.3	3.3-14.4
	1986-1990	6.6	16.3	11.1-23.1	4.2	24.0	14.7-37.1	2.4	10.9	5.5-19.6	1.2	2.9	1.1-6.5	0.4	1.8	0.2-7.2	0.8	3.5	0.9-9.5
	1991-1995	9.2	21.2	15.5-28.4	5.4	31.0	20.2-45.3	3.8	15.2	9.0-24.4	2.0	4.9	2.3-9.0	1.6	9.4	4.0-18.5	0.4	2.0	0.2-7.5
	1996-2000	7.4	16.3	11.4-22.6	4.6	22.5	14.2-34.0	2.8	11.1	5.9-19.3	1.2	2.7	1.0-6.1	0.6	2.9	0.6-8.7	0.6	2.8	0.5-8.6
	2001-2005	7.6	15.2	10.8-21.1	4.8	22.8	14.5-34.3	2.8	10.6	5.7-18.5	3.2	6.5	3.7-10.7	2.0	9.4	4.5-17.6	1.2	4.6	1.6-10.7
	2006-2010	7.6	13.8	9.7-19.2	5.2	19.3	12.5-28.9	2.4	7.9	4.0-14.8	1.8	3.2	1.5-6.5	1.2	4.4	1.6-10.2	0.6	1.8	0.4-6.6
	2011-2015	7.4	13.0	9.0-18.3	5.6	21.6	14.2-31.8	1.8	5.4	2.4-11.6	1.6	2.8	1.2-6.0	0.8	2.9	0.8-8.4	0.8	2.4	0.6-7.6
2016-2020	5.0	10.0	6.2-15.4	3.8	14.8	8.6-24.2	1.2	4.9	1.5-12.2	2.0	2.9	1.4-6.1	1.0	3.0	1.0-8.4	1.0	2.7	0.9-8.2	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Essex**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	0.4	0.9	0.1-3.7	0.0	0.0	0.0-4.8	0.4	1.6	0.2-6.7	0.4	0.9	0.1-3.5	0.2	0.9	0.0-6.1	0.2	0.8	0.0-5.5
	1981-1985	1.6	4.3	1.8-8.7	1.4	8.6	3.3-17.9	0.2	0.9	0.0-5.7	1.2	2.8	1.0-6.4	1.2	6.2	2.2-13.9	0.0	0.0	0.0-4.1
	1986-1990	2.2	5.1	2.6-9.4	1.8	10.5	4.6-20.2	0.4	2.0	0.2-7.6	1.2	3.2	1.2-7.0	1.0	6.6	2.1-15.3	0.2	0.9	0.0-5.6
	1991-1995	1.8	4.3	2.0-8.4	1.4	7.3	2.9-15.3	0.4	1.7	0.2-6.7	2.4	5.6	2.9-10.0	2.0	10.6	5.0-19.6	0.4	1.5	0.2-6.2
	1996-2000	0.6	1.3	0.3-4.0	0.6	3.5	0.7-10.1	0.0	0.0	0.0-4.0	0.4	0.8	0.1-3.2	0.4	2.0	0.2-7.4	0.0	0.0	0.0-4.0
	2001-2005	2.4	5.0	2.6-8.9	1.4	6.3	2.5-13.2	1.0	3.8	1.2-9.6	2.6	5.1	2.7-9.0	1.0	4.5	1.4-10.7	1.6	5.9	2.5-12.3
	2006-2010	4.2	8.3	5.0-13.0	3.6	14.3	8.3-23.3	0.6	3.0	0.5-9.2	2.6	5.1	2.6-9.1	2.2	9.2	4.4-17.0	0.4	2.2	0.2-8.2
	2011-2015	4.0	6.3	3.9-10.2	3.0	9.8	5.4-16.9	1.0	3.0	1.0-8.4	3.4	5.2	3.0-8.8	3.0	9.5	5.3-16.5	0.4	1.3	0.2-6.2
	2016-2020	5.6	8.6	5.6-13.1	3.4	10.9	6.3-18.5	2.2	7.0	3.1-14.5	3.4	5.3	2.9-9.1	2.0	6.2	2.9-12.5	1.4	5.0	1.7-12.1
Pancreas	1976-1980	4.4	10.7	6.6-16.5	3.0	17.4	9.4-29.5	1.4	6.2	2.4-13.3	3.0	7.4	4.1-12.3	1.4	8.9	3.3-18.9	1.6	6.6	2.8-13.6
	1981-1985	7.6	19.6	13.8-27.1	4.4	25.9	15.9-39.7	3.2	13.5	7.6-22.4	6.6	17.1	11.7-24.2	3.8	23.2	13.7-36.6	2.8	12.0	6.4-20.7
	1986-1990	5.2	13.6	8.9-20.1	3.0	19.5	10.8-32.2	2.2	9.0	4.5-16.7	4.4	11.2	7.0-17.1	2.6	16.4	8.5-28.1	1.8	7.1	3.3-14.1
	1991-1995	4.8	10.5	6.7-15.8	3.0	15.8	8.7-26.4	1.8	6.4	2.9-12.9	6.2	13.8	9.4-19.8	3.0	15.4	8.5-25.7	3.2	12.2	6.9-20.4
	1996-2000	3.4	7.3	4.2-11.8	1.8	9.7	4.4-18.6	1.6	5.3	2.2-11.3	3.6	7.8	4.6-12.5	2.0	10.9	5.1-20.1	1.6	5.5	2.3-11.7
	2001-2005	6.2	12.2	8.3-17.5	3.6	16.4	9.6-26.3	2.6	9.4	5.0-16.8	4.6	9.1	5.8-13.9	2.4	10.7	5.5-19.1	2.2	7.8	3.8-14.7
	2006-2010	7.8	13.9	9.9-19.3	4.8	19.6	12.5-29.6	3.0	9.8	5.4-17.0	7.8	14.1	10.0-19.5	5.0	20.4	13.2-30.6	2.8	9.1	4.9-16.2
	2011-2015	8.0	13.3	9.5-18.6	4.0	14.5	8.7-23.0	4.0	12.6	7.7-20.4	6.8	11.2	7.7-16.1	3.6	12.8	7.5-20.9	3.2	9.8	5.6-17.0
	2016-2020	7.8	13.1	9.0-18.6	4.2	14.6	8.6-23.6	3.6	11.2	6.4-19.1	7.0	11.4	7.8-16.4	3.6	12.2	6.9-20.4	3.4	10.5	6.1-18.2
Larynx	1976-1980	3.8	9.9	5.9-15.7	2.8	17.9	9.5-30.5	1.0	4.3	1.4-10.6	1.6	4.3	1.8-8.8	1.2	7.2	2.5-16.0	0.4	1.8	0.2-7.1
	1981-1985	3.6	9.4	5.5-15.1	3.2	19.5	10.9-32.0	0.4	1.7	0.2-6.8	1.4	4.3	1.7-8.9	1.4	9.6	3.8-19.7	0.0	0.0	0.0-4.1
	1986-1990	2.2	5.2	2.6-9.5	2.0	10.7	5.1-20.0	0.2	0.7	0.0-5.2	1.0	2.5	0.8-5.8	1.0	5.5	1.8-13.1	0.0	0.0	0.0-4.1
	1991-1995	3.6	9.0	5.3-14.3	3.0	15.1	8.4-25.2	0.6	3.3	0.7-9.7	1.8	4.0	1.8-7.7	1.6	8.2	3.5-16.5	0.2	0.9	0.0-5.4
	1996-2000	2.8	6.4	3.5-10.8	2.2	10.3	5.1-18.6	0.6	2.7	0.6-8.4	2.0	4.3	2.1-8.1	1.4	6.5	2.6-13.7	0.6	2.3	0.5-7.5
	2001-2005	1.4	2.8	1.1-6.1	1.4	6.2	2.4-13.2	0.0	0.0	0.0-4.1	0.2	0.4	0.0-2.6	0.2	0.8	0.0-5.1	0.0	0.0	0.0-4.1
	2006-2010	1.8	3.4	1.5-6.8	1.6	6.2	2.6-12.8	0.2	0.7	0.0-5.1	1.2	2.2	0.8-5.1	1.2	4.4	1.6-10.4	0.0	0.0	0.0-4.2
	2011-2015	2.0	4.0	1.8-7.7	2.0	7.9	3.6-15.2	0.0	0.0	0.0-4.5	0.4	1.0	0.1-3.8	0.4	1.8	0.2-7.2	0.0	0.0	0.0-4.5
	2016-2020	2.0	3.1	1.5-6.4	1.4	4.4	1.7-10.3	0.6	1.9	0.4-7.3	0.8	1.4	0.4-4.2	0.6	2.0	0.4-7.3	0.2	0.7	0.0-5.7
Lung and bronchus	1976-1980	30.0	76.1	64.1-89.8	21.2	121.2	98.3-147.8	8.8	42.1	30.1-57.3	24.6	60.6	50.2-72.7	17.4	97.5	77.3-121.5	7.2	32.7	22.7-45.9
	1981-1985	36.8	90.5	77.6-105.0	24.8	137.3	113.6-164.6	12.0	56.3	42.4-73.4	27.8	67.5	56.5-80.1	20.2	113.1	91.4-138.4	7.6	35.1	24.4-49.0
	1986-1990	36.2	90.1	77.3-104.4	23.0	135.5	111.2-163.4	13.2	61.4	47.2-78.8	29.0	71.4	60.1-84.3	19.8	114.3	92.4-139.8	9.2	42.2	30.6-56.8
	1991-1995	37.8	87.2	75.1-100.7	20.4	110.2	89.5-134.3	17.4	71.0	56.6-88.1	29.0	65.8	55.4-77.5	15.2	83.9	65.7-105.4	13.8	54.5	42.3-69.5
	1996-2000	37.0	80.5	69.3-93.1	20.6	102.4	83.3-124.6	16.4	67.0	53.1-83.6	26.6	57.3	48.0-68.1	15.4	75.5	59.4-94.7	11.2	43.4	32.7-57.0
	2001-2005	36.6	73.4	63.2-85.1	20.4	91.8	74.6-111.9	16.2	60.5	47.8-75.9	29.0	58.1	49.0-68.5	15.6	72.2	56.8-90.5	13.4	49.7	38.3-63.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Essex**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	41.0	75.9	65.7-87.3	21.6	87.2	71.3-105.8	19.4	67.3	54.3-83.0	27.4	50.5	42.3-59.9	12.4	51.1	39.1-65.9	15.0	52.0	40.7-66.0
	2011-2015	44.4	73.2	63.8-83.9	22.2	76.9	63.0-93.3	22.2	71.4	58.5-86.8	28.6	46.8	39.3-55.5	14.4	51.9	40.4-66.0	14.2	43.3	33.7-55.5
	2016-2020	40.4	61.2	52.8-71.0	19.0	61.8	49.6-76.6	21.4	63.0	51.2-77.5	27.4	42.0	35.1-50.2	13.6	45.3	35.0-58.2	13.8	41.6	31.9-54.2
Melanoma of the skin	1976-1980	2.0	6.1	2.8-11.2	1.4	9.7	3.8-20.0	0.6	2.6	0.5-8.3	1.2	3.6	1.3-7.7	0.6	5.0	1.0-13.8	0.6	3.0	0.6-9.0
	1981-1985	1.6	5.0	2.1-9.9	1.0	7.3	2.3-16.7	0.6	3.4	0.6-10.1	1.2	3.1	1.1-7.0	0.4	2.5	0.3-9.1	0.8	3.4	0.9-9.3
	1986-1990	2.4	6.2	3.1-10.9	1.4	8.0	3.1-16.7	1.0	4.8	1.5-11.8	1.2	3.0	1.1-6.7	1.0	5.5	1.7-13.1	0.2	1.4	0.0-7.0
	1991-1995	5.0	12.4	8.0-18.4	2.6	14.2	7.5-24.4	2.4	11.6	5.9-20.6	0.6	1.4	0.3-4.2	0.4	2.1	0.3-7.8	0.2	1.1	0.0-6.0
	1996-2000	8.2	19.0	13.6-26.0	5.8	27.1	18.1-39.2	2.4	13.1	6.7-23.1	1.8	4.0	1.8-7.8	0.8	4.0	1.1-10.4	1.0	4.2	1.3-10.5
	2001-2005	8.8	18.8	13.6-25.5	5.0	23.4	15.0-34.8	3.8	16.6	9.7-26.6	2.0	4.1	1.9-7.7	1.4	5.9	2.4-12.6	0.6	2.1	0.4-7.2
	2006-2010	11.4	23.0	17.3-30.2	6.8	27.6	19.0-39.1	4.6	19.9	12.3-30.8	2.2	4.0	2.0-7.4	1.8	7.1	3.2-14.1	0.4	1.5	0.2-6.3
	2011-2015	13.2	24.1	18.4-31.2	6.0	22.8	15.2-33.3	7.2	27.7	18.8-39.7	2.2	3.9	1.9-7.3	1.6	6.7	2.9-13.6	0.6	1.5	0.3-6.4
	2016-2020	13.0	22.2	16.8-29.0	9.4	33.6	24.2-45.9	3.6	11.9	6.9-20.2	2.2	3.2	1.6-6.4	1.8	6.3	2.8-12.8	0.4	0.9	0.1-5.9
Female breast	1976-1980							18.8	96.9	77.5-119.6							7.4	38.2	26.5-53.2
	1981-1985							17.8	82.5	65.6-102.5							8.2	36.0	25.6-49.5
	1986-1990							24.2	109.7	90.5-132.0							9.0	39.1	28.3-52.9
	1991-1995							30.6	134.0	113.1-157.8							8.0	33.5	23.7-46.1
	1996-2000							28.4	119.8	100.6-142.0							5.4	20.9	13.6-31.0
	2001-2005							26.8	107.5	89.7-128.2							4.2	14.8	9.1-23.4
	2006-2010							32.6	121.2	102.7-142.3							3.6	12.1	7.0-19.9
	2011-2015							34.8	122.6	104.0-144.1							5.6	18.0	11.9-27.1
	2016-2020							29.0	99.4	82.4-119.3							5.0	14.0	8.9-22.2
Cervix uteri	1976-1980							2.2	15.0	7.3-26.9							1.2	7.7	2.6-16.8
	1981-1985							3.2	18.4	10.2-30.4							1.0	6.7	2.1-15.5
	1986-1990							2.2	12.8	6.2-23.0							0.8	3.8	1.0-10.2
	1991-1995							2.0	10.9	5.1-20.2							1.0	5.5	1.7-12.9
	1996-2000							1.8	8.3	3.7-16.1							0.8	3.7	1.0-9.9
	2001-2005							2.0	10.1	4.7-18.9							0.8	2.8	0.7-8.2
	2006-2010							1.8	6.5	2.8-13.2							0.6	1.9	0.4-6.7
	2011-2015							1.6	7.1	2.7-15.3							0.4	1.2	0.1-6.1
	2016-2020							1.6	8.8	3.4-18.5							0.6	2.7	0.4-9.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Essex**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							2.8	13.4	7.1-23.2							0.8	3.9	1.0-10.4
	1981-1985							7.0	29.8	20.5-42.2							1.8	7.9	3.6-15.4
	1986-1990							4.2	19.2	11.7-30.1							1.0	3.6	1.2-9.2
	1991-1995							7.4	31.6	22.1-44.1							1.0	3.6	1.2-9.3
	1996-2000							5.8	25.7	17.1-37.3							1.2	4.4	1.6-10.4
	2001-2005							6.0	23.5	15.8-34.1							0.6	1.8	0.4-6.7
	2006-2010							8.8	32.2	23.1-44.0							2.4	7.8	4.0-14.5
	2011-2015							7.4	22.8	15.9-32.4							1.8	5.2	2.3-11.1
	2016-2020							7.6	25.6	17.8-36.5							1.4	3.7	1.5-9.4
Ovary	1976-1980							2.6	12.9	6.7-22.4							1.8	8.9	4.0-17.2
	1981-1985							4.2	21.4	12.9-33.4							2.2	10.5	5.1-19.3
	1986-1990							4.0	20.4	12.2-32.0							2.4	11.4	5.8-20.3
	1991-1995							6.0	26.2	17.4-38.0							3.4	14.5	8.3-23.9
	1996-2000							3.8	16.0	9.5-25.6							3.8	15.4	9.2-24.7
	2001-2005							4.6	18.0	11.3-27.7							2.2	8.1	4.0-15.2
	2006-2010							3.0	11.1	6.2-19.0							2.4	8.6	4.4-15.8
	2011-2015							3.0	10.1	5.4-18.0							1.0	2.7	0.9-8.0
	2016-2020							2.2	7.0	3.3-14.1							2.6	9.4	4.5-17.8
Prostate	1976-1980				14.2	93.5	72.2-118.9							4.4	32.0	19.6-48.8			
	1981-1985				16.8	110.2	87.2-137.1							6.0	40.9	27.2-58.7			
	1986-1990				18.6	112.4	90.2-138.3							5.8	38.0	25.1-54.9			
	1991-1995				34.8	190.5	162.6-221.7							6.4	40.8	27.7-57.7			
	1996-2000				28.4	140.3	117.8-166.0							7.0	41.0	28.3-57.2			
	2001-2005				33.8	148.6	126.8-173.2							7.0	38.4	26.5-53.5			
	2006-2010				33.2	125.2	106.6-146.4							3.0	12.3	6.8-20.7			
	2011-2015				28.6	96.1	80.7-114.0							3.6	14.2	8.4-23.0			
	2016-2020				31.8	91.2	77.3-107.6							4.4	15.4	9.6-24.1			
Testis	1976-1980				0.6	3.9	0.7-11.7							0.0	0.0	0.0-4.8			
	1981-1985				0.8	5.3	1.4-13.5							0.0	0.0	0.0-4.5			
	1986-1990				1.2	5.3	1.9-12.2							0.2	1.1	0.0-6.3			
	1991-1995				1.4	6.5	2.6-13.9							0.2	1.2	0.0-6.4			
	1996-2000				1.8	8.9	4.0-17.2							0.0	0.0	0.0-3.9			
	2001-2005				0.8	3.6	1.0-9.5							0.2	0.8	0.0-5.2			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Essex**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				1.6	8.3	3.6-16.3							0.2	1.0	0.0-5.6			
	2011-2015				1.4	7.8	3.1-16.2							0.0	0.0	0.0-4.1			
	2016-2020				1.2	6.0	2.1-13.7							0.2	1.1	0.0-6.4			
Urinary bladder (incl. in situ)	1976-1980	10.8	27.8	20.7-36.5	9.4	58.3	42.2-78.5	1.4	6.8	2.7-14.4	2.4	6.2	3.2-11.0	1.6	9.5	4.0-19.1	0.8	4.1	1.1-10.7
	1981-1985	10.2	24.9	18.4-33.1	7.0	40.0	27.4-56.5	3.2	13.9	7.8-23.2	2.0	5.0	2.4-9.4	1.4	9.9	3.8-20.5	0.6	2.5	0.5-7.9
	1986-1990	9.0	21.8	15.9-29.4	6.8	41.2	28.2-58.0	2.2	9.6	4.7-17.7	2.4	5.8	3.0-10.2	1.6	10.4	4.3-20.6	0.8	3.0	0.8-8.6
	1991-1995	11.6	27.0	20.5-35.0	8.4	47.4	33.9-64.4	3.2	13.5	7.6-22.4	3.0	7.0	3.9-11.6	1.8	10.7	4.8-20.3	1.2	4.6	1.7-10.8
	1996-2000	13.0	27.8	21.4-35.6	8.8	44.0	31.7-59.3	4.2	16.1	9.9-25.2	1.6	3.5	1.5-7.1	1.0	5.5	1.8-12.9	0.6	2.0	0.4-6.8
	2001-2005	13.4	26.8	20.8-34.3	10.0	45.2	33.5-59.8	3.4	13.2	7.6-21.8	1.2	2.4	0.9-5.5	1.0	4.6	1.5-11.0	0.2	0.8	0.0-5.3
	2006-2010	14.6	27.1	21.2-34.3	12.0	48.1	36.6-62.4	2.6	8.4	4.5-15.3	1.4	2.6	1.0-5.7	0.8	3.3	0.9-8.9	0.6	1.9	0.4-6.7
	2011-2015	15.2	25.4	19.9-32.2	12.2	45.3	34.5-58.9	3.0	8.6	4.8-15.4	2.6	4.6	2.4-8.3	1.8	7.5	3.4-14.7	0.8	1.9	0.5-6.8
2016-2020	16.8	26.4	20.9-33.2	13.8	46.9	36.3-60.4	3.0	8.4	4.7-15.3	2.8	4.0	2.2-7.3	2.0	6.9	3.3-13.6	0.8	1.8	0.5-7.0	
Kidney and renal pelvis	1976-1980	3.0	7.3	4.0-12.2	2.6	15.1	7.8-26.4	0.4	1.5	0.2-6.5	1.4	3.6	1.4-7.6	1.2	7.7	2.6-17.1	0.2	0.7	0.0-5.3
	1981-1985	4.4	9.9	6.1-15.1	3.2	16.4	9.2-27.4	1.2	4.9	1.8-11.3	2.2	4.7	2.3-8.6	1.8	8.4	3.8-16.4	0.4	1.7	0.2-6.8
	1986-1990	4.0	11.1	6.7-17.4	2.8	16.7	9.0-28.3	1.2	6.8	2.4-15.0	2.6	6.6	3.5-11.5	1.8	10.1	4.5-19.4	0.8	3.7	0.9-10.0
	1991-1995	4.4	10.3	6.4-15.7	2.4	13.2	6.7-23.1	2.0	7.7	3.7-14.7	1.2	2.9	1.1-6.5	1.2	7.3	2.6-15.7	0.0	0.0	0.0-4.0
	1996-2000	5.6	12.2	8.1-17.8	3.0	14.6	8.1-24.3	2.6	10.5	5.5-18.5	2.6	5.8	3.1-10.1	1.2	6.4	2.3-14.0	1.4	5.5	2.2-11.9
	2001-2005	6.6	13.3	9.2-18.9	4.8	20.3	13.0-30.6	1.8	6.8	3.1-13.6	0.8	1.6	0.4-4.3	0.4	1.6	0.2-6.4	0.4	1.4	0.2-6.1
	2006-2010	7.0	13.0	9.0-18.3	4.4	18.0	11.1-27.7	2.6	10.1	5.2-18.1	1.4	2.7	1.1-5.8	0.8	3.3	0.9-8.9	0.6	2.2	0.4-7.3
	2011-2015	9.2	16.4	11.8-22.4	4.6	16.4	10.3-25.4	4.6	16.0	9.8-25.3	1.8	2.8	1.3-5.9	1.2	4.1	1.5-9.9	0.6	1.8	0.4-6.8
2016-2020	7.4	11.5	8.0-16.4	5.4	17.7	11.5-26.8	2.0	5.3	2.5-11.4	2.0	2.9	1.4-6.1	1.6	4.9	2.1-10.8	0.4	1.1	0.1-6.2	
Brain and other central nervous system	1976-1980	1.4	4.1	1.6-8.7	1.2	8.3	2.9-18.1	0.2	0.7	0.0-5.3	0.8	2.0	0.5-5.4	0.6	3.9	0.8-11.6	0.2	0.7	0.0-5.3
	1981-1985	3.2	8.5	4.8-13.9	1.8	10.9	4.8-20.9	1.4	6.7	2.6-14.2	1.6	4.1	1.7-8.3	1.0	5.6	1.7-13.4	0.6	2.5	0.5-7.9
	1986-1990	2.2	5.7	2.8-10.4	1.4	7.1	2.8-15.0	0.8	4.5	1.2-11.7	2.2	5.8	2.9-10.6	1.0	5.4	1.7-12.9	1.2	6.3	2.2-14.0
	1991-1995	3.4	9.0	5.2-14.5	1.6	8.9	3.8-17.6	1.8	9.0	4.0-17.3	1.6	3.8	1.6-7.7	0.6	2.7	0.6-8.3	1.0	4.8	1.5-11.6
	1996-2000	5.6	13.6	9.0-19.7	2.2	10.8	5.4-19.6	3.4	16.4	9.4-26.8	3.2	7.2	4.1-11.8	1.6	7.4	3.2-14.9	1.6	6.5	2.8-13.4
	2001-2005	2.4	5.3	2.7-9.5	1.0	4.6	1.5-11.0	1.4	6.4	2.4-13.8	2.0	4.0	1.9-7.6	1.0	4.6	1.5-11.0	1.0	3.8	1.2-9.6
	2006-2010	4.6	9.5	5.9-14.5	3.0	13.2	7.3-22.2	1.6	6.0	2.5-12.6	1.8	3.5	1.6-6.9	1.2	4.8	1.7-11.0	0.6	2.7	0.5-8.6
	2011-2015	4.8	10.3	6.3-15.8	2.6	11.8	6.0-21.0	2.2	8.8	4.0-17.1	4.0	7.1	4.3-11.5	2.0	8.3	3.8-15.9	2.0	5.9	2.8-12.1
2016-2020	2.4	3.5	1.8-6.7	1.4	4.1	1.6-9.8	1.0	2.9	0.9-8.5	2.6	4.3	2.2-8.0	1.6	5.3	2.2-11.5	1.0	3.5	0.9-10.1	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Essex**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	0.8	1.8	0.5-5.0	0.2	0.9	0.0-6.1	0.6	2.8	0.6-8.7	0.0	0.0	0.0-2.2	0.0	0.0	0.0-4.8	0.0	0.0	0.0-4.2
	1981-1985	0.6	1.9	0.4-5.6	0.2	0.8	0.0-5.8	0.4	3.0	0.4-10.1	0.2	0.4	0.0-2.7	0.2	0.8	0.0-5.8	0.0	0.0	0.0-4.1
	1986-1990	1.8	4.7	2.1-9.0	0.6	3.6	0.7-10.5	1.2	5.4	2.0-12.3	0.4	1.1	0.1-3.9	0.2	1.3	0.0-6.7	0.2	0.8	0.0-5.4
	1991-1995	1.0	2.4	0.8-5.8	0.2	0.8	0.0-5.3	0.8	4.3	1.2-11.1	0.2	0.4	0.0-2.6	0.0	0.0	0.0-4.0	0.2	0.7	0.0-5.1
	1996-2000	3.2	7.8	4.5-12.8	0.4	2.0	0.2-7.5	2.8	14.4	7.8-24.5	0.2	0.4	0.0-2.6	0.2	1.1	0.0-6.1	0.0	0.0	0.0-4.0
	2001-2005	2.8	6.3	3.4-10.7	0.8	3.3	0.9-9.0	2.0	10.0	4.7-18.9	0.0	0.0	0.0-1.9	0.0	0.0	0.0-3.9	0.0	0.0	0.0-4.1
	2006-2010	5.8	12.8	8.4-18.6	1.4	5.5	2.2-11.8	4.4	21.1	12.9-32.8	0.4	0.7	0.1-3.1	0.4	1.6	0.2-6.3	0.0	0.0	0.0-4.2
	2011-2015	5.8	13.3	8.7-19.5	1.0	3.9	1.2-9.9	4.8	24.3	15.1-37.2	0.6	1.1	0.2-3.6	0.0	0.0	0.0-4.1	0.6	2.0	0.4-7.3
	2016-2020	4.4	10.4	6.2-16.4	1.2	5.8	1.9-13.6	3.2	15.3	8.2-26.2	0.0	0.0	0.0-2.3	0.0	0.0	0.0-4.4	0.0	0.0	0.0-4.9
Hodgkin lymphoma	1976-1980	1.2	3.8	1.3-8.3	0.6	3.9	0.7-11.6	0.6	3.6	0.7-10.7	0.4	1.4	0.1-4.8	0.0	0.0	0.0-4.8	0.4	2.6	0.3-9.3
	1981-1985	1.0	2.7	0.8-6.3	0.4	2.7	0.3-9.4	0.6	2.5	0.5-7.9	0.6	1.4	0.3-4.4	0.2	1.1	0.0-6.4	0.4	1.7	0.2-6.8
	1986-1990	1.2	2.8	1.0-6.2	0.8	3.9	1.0-10.5	0.4	1.4	0.2-6.2	0.6	1.4	0.3-4.3	0.4	3.0	0.3-10.2	0.2	0.7	0.0-5.2
	1991-1995	2.0	4.9	2.3-9.1	1.2	5.1	1.9-11.7	0.8	4.8	1.3-12.1	1.0	2.5	0.8-6.0	0.6	3.2	0.7-9.5	0.4	1.8	0.2-7.2
	1996-2000	1.4	3.5	1.4-7.3	1.2	5.5	2.0-12.3	0.2	0.9	0.0-5.5	0.0	0.0	0.0-1.9	0.0	0.0	0.0-3.9	0.0	0.0	0.0-4.0
	2001-2005	1.2	2.7	1.0-6.1	0.4	1.9	0.2-7.1	0.8	3.4	0.8-9.5	0.0	0.0	0.0-1.9	0.0	0.0	0.0-3.9	0.0	0.0	0.0-4.1
	2006-2010	0.6	1.4	0.2-4.5	0.4	2.4	0.3-8.3	0.2	0.7	0.0-5.1	0.0	0.0	0.0-2.0	0.0	0.0	0.0-3.8	0.0	0.0	0.0-4.2
	2011-2015	1.2	3.3	1.2-7.3	1.0	5.2	1.6-12.4	0.2	1.4	0.0-7.2	0.0	0.0	0.0-2.1	0.0	0.0	0.0-4.1	0.0	0.0	0.0-4.5
	2016-2020	1.0	3.6	1.2-8.3	0.2	1.0	0.0-6.1	0.8	6.6	1.8-16.2	0.2	0.3	0.0-2.7	0.2	0.6	0.0-5.1	0.0	0.0	0.0-4.9
Non-Hodgkin lymphomas	1976-1980	5.0	12.9	8.2-19.4	3.0	17.8	9.7-29.8	2.0	8.4	4.0-16.0	2.6	6.7	3.5-11.6	1.2	7.1	2.5-15.8	1.4	5.9	2.3-12.7
	1981-1985	5.2	13.7	8.8-20.3	2.6	15.3	7.9-26.6	2.6	12.9	6.6-22.6	3.4	9.0	5.1-14.7	1.6	9.0	3.7-18.1	1.8	8.8	3.8-17.3
	1986-1990	8.2	21.4	15.2-29.1	4.6	25.5	16.0-38.6	3.6	16.9	9.7-27.3	3.4	8.6	5.0-13.9	1.2	6.9	2.4-15.2	2.2	9.8	4.8-18.1
	1991-1995	7.8	18.9	13.4-25.9	3.8	20.9	12.4-32.9	4.0	17.6	10.6-27.6	4.6	10.4	6.6-15.8	2.0	10.5	5.0-19.5	2.6	9.6	5.1-17.0
	1996-2000	9.0	20.2	14.7-27.2	4.6	22.3	14.1-33.6	4.4	16.9	10.4-26.3	3.8	8.3	5.0-13.1	1.2	6.4	2.3-13.9	2.6	9.5	5.0-17.0
	2001-2005	9.4	19.6	14.4-26.3	4.4	19.2	12.0-29.4	5.0	19.3	12.2-29.3	3.8	7.6	4.6-12.1	2.4	10.8	5.5-19.2	1.4	5.2	2.0-11.5
	2006-2010	11.2	21.4	16.1-28.1	7.0	28.1	19.4-39.5	4.2	14.7	9.0-23.3	3.8	6.9	4.1-11.0	1.0	4.6	1.5-10.9	2.8	8.8	4.7-15.7
	2011-2015	11.0	19.8	14.6-26.4	7.4	28.3	19.5-39.9	3.6	12.5	7.1-20.9	3.2	4.9	2.8-8.4	2.2	7.9	3.9-14.9	1.0	2.6	0.8-7.8
	2016-2020	12.8	20.5	15.6-26.9	7.2	23.2	16.0-33.1	5.6	18.8	11.9-28.8	3.4	4.9	2.8-8.4	2.2	6.9	3.4-13.4	1.2	3.5	1.3-9.3
Myeloma	1976-1980	2.8	7.7	4.1-13.1	2.0	12.6	5.7-23.8	0.8	3.6	1.0-9.6	2.0	5.2	2.5-9.7	1.2	7.6	2.7-16.9	0.8	3.4	0.9-9.4
	1981-1985	3.2	8.0	4.5-13.1	1.6	10.2	4.2-20.3	1.6	7.0	3.0-14.4	2.2	5.1	2.5-9.3	1.0	4.7	1.5-11.5	1.2	5.0	1.8-11.4
	1986-1990	4.2	11.3	7.0-17.4	2.6	17.1	9.0-29.2	1.6	7.0	3.0-14.2	2.2	5.7	2.8-10.3	1.2	8.4	3.0-17.9	1.0	4.0	1.3-10.1
	1991-1995	2.8	6.5	3.5-11.0	1.4	8.3	3.3-17.1	1.4	5.4	2.1-11.6	2.4	5.4	2.8-9.6	1.4	8.3	3.2-17.1	1.0	3.6	1.2-9.2
	1996-2000	2.8	5.8	3.2-10.0	1.2	6.4	2.3-14.0	1.6	5.8	2.5-12.1	1.4	2.9	1.2-6.3	0.4	2.4	0.3-8.4	1.0	3.7	1.2-9.3
	2001-2005	4.0	8.0	4.9-12.6	2.2	10.2	5.1-18.6	1.8	6.3	2.8-12.8	2.0	3.8	1.8-7.3	1.2	6.0	2.1-13.4	0.8	2.6	0.7-7.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Essex**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	5.8	10.8	7.2-15.9	3.2	12.6	7.1-21.0	2.6	9.0	4.7-16.2	3.0	5.3	3.0-9.0	1.8	7.5	3.4-14.7	1.2	3.6	1.3-9.0
	2011-2015	4.8	8.2	5.2-12.7	2.0	7.5	3.5-14.6	2.8	9.4	5.1-16.9	2.2	3.9	1.9-7.5	0.8	3.2	0.9-8.9	1.4	4.6	1.8-10.8
	2016-2020	3.6	4.9	2.9-8.4	2.6	7.7	4.0-14.3	1.0	2.5	0.8-7.8	1.6	2.1	0.9-4.9	0.6	1.7	0.3-6.7	1.0	2.6	0.8-8.0
Leukemias	1976-1980	7.4	20.1	14.1-28.0	4.4	29.3	18.0-44.8	3.0	14.5	8.0-24.4	4.4	12.8	8.0-19.5	2.2	16.2	7.9-29.0	2.2	11.1	5.5-20.3
	1981-1985	4.8	12.1	7.7-18.2	2.6	16.9	8.8-29.0	2.2	9.1	4.5-16.8	4.6	11.8	7.5-17.9	3.0	20.4	11.1-33.9	1.6	7.1	3.0-14.5
	1986-1990	5.2	12.3	8.0-18.2	2.6	14.1	7.3-24.6	2.6	10.9	5.8-19.1	2.6	6.3	3.3-10.9	1.6	8.5	3.7-17.0	1.0	4.1	1.3-10.2
	1991-1995	6.2	15.1	10.2-21.6	3.4	19.2	11.1-31.0	2.8	11.9	6.3-20.5	3.8	8.9	5.3-14.0	2.2	13.3	6.5-23.8	1.6	6.0	2.6-12.5
	1996-2000	9.0	19.8	14.4-26.7	4.0	19.9	12.1-31.0	5.0	19.1	12.2-28.8	4.0	8.8	5.4-13.7	2.0	10.3	4.9-19.1	2.0	7.0	3.3-13.7
	2001-2005	7.4	15.5	10.8-21.6	4.2	19.6	12.0-30.3	3.2	11.2	6.1-19.3	3.6	7.2	4.2-11.6	1.8	8.3	3.7-16.0	1.8	5.7	2.6-11.8
	2006-2010	8.6	16.9	12.1-23.1	5.8	22.8	15.2-33.3	2.8	10.8	5.6-19.2	3.4	6.2	3.6-10.3	1.6	6.4	2.7-13.2	1.8	5.4	2.4-11.4
	2011-2015	11.2	20.6	15.3-27.4	7.4	29.0	20.1-40.8	3.8	13.1	7.4-21.9	3.4	5.7	3.2-9.6	2.0	7.4	3.5-14.3	1.4	4.9	1.7-11.6
	2016-2020	7.8	13.0	9.0-18.5	5.0	18.1	11.3-28.1	2.8	7.8	4.2-14.5	4.0	5.8	3.5-9.5	2.2	6.8	3.4-13.4	1.8	4.8	2.2-10.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Franklin**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	160.8	382.3	355.8-410.3	82.8	464.3	418.5-513.6	78.0	334.8	301.5-370.7	93.2	221.2	201.2-242.7	48.2	276.1	240.6-315.1	45.0	187.4	163.3-214.2
	1981-1985	205.6	479.1	449.7-509.9	106.6	597.0	545.1-652.2	99.0	410.0	373.7-448.9	105.2	246.8	225.8-269.2	54.6	314.1	276.3-355.4	50.6	205.3	180.2-232.9
	1986-1990	211.0	483.5	454.3-514.0	111.0	602.3	551.5-656.5	100.0	404.5	369.1-442.5	111.2	255.9	234.8-278.3	57.6	318.5	281.5-358.9	53.6	212.8	187.6-240.5
	1991-1995	231.4	489.1	461.1-518.3	122.6	609.0	560.1-660.9	108.8	416.8	381.9-454.1	108.0	228.7	209.7-249.0	55.0	289.2	254.8-326.6	53.0	193.0	170.1-218.1
	1996-2000	265.6	532.7	504.4-562.2	133.2	612.2	565.6-661.6	132.4	490.3	453.2-529.8	104.8	210.7	193.0-229.5	54.8	266.1	234.7-300.3	50.0	174.8	153.6-198.4
	2001-2005	285.0	540.4	512.6-569.2	157.2	659.6	613.2-708.5	127.8	462.0	426.5-499.8	100.4	193.0	176.5-210.7	52.8	240.5	211.7-272.1	47.6	162.6	142.4-185.0
	2006-2010	279.6	491.0	465.3-517.7	150.6	563.7	522.8-606.8	129.0	441.0	407.2-477.2	105.0	188.9	173.0-205.9	54.4	220.0	193.8-248.7	50.6	167.4	147.3-189.8
	2011-2015	283.0	466.4	442.0-491.9	160.2	542.4	504.4-582.7	122.8	407.7	375.2-442.5	102.2	170.4	155.8-186.1	54.4	201.5	177.6-227.8	47.8	149.6	130.9-170.5
	2016-2020	304.2	466.6	442.8-491.5	174.2	532.8	497.0-570.7	130.0	414.4	381.5-449.5	100.4	153.6	140.2-168.0	51.6	169.4	148.8-192.1	48.8	141.3	123.7-161.2
Oral cavity and pharynx	1976-1980	4.8	10.7	6.8-16.0	4.4	24.0	14.5-37.3	0.4	1.5	0.2-5.8	2.0	4.2	2.0-7.9	1.6	7.8	3.3-15.7	0.4	1.5	0.2-5.8
	1981-1985	5.6	13.4	8.8-19.5	4.4	23.1	14.3-35.3	1.2	6.1	2.1-13.4	2.0	5.0	2.3-9.2	1.6	8.6	3.7-17.1	0.4	1.6	0.2-6.1
	1986-1990	5.8	14.4	9.6-20.7	3.4	19.9	11.5-31.8	2.4	10.1	5.1-17.9	2.0	5.2	2.5-9.5	1.0	6.8	2.1-15.6	1.0	5.2	1.6-12.0
	1991-1995	5.2	11.7	7.6-17.1	4.2	21.1	12.9-32.5	1.0	4.1	1.3-9.8	1.0	2.2	0.7-5.1	0.6	3.3	0.7-9.4	0.4	1.6	0.2-5.9
	1996-2000	3.2	6.5	3.7-10.6	1.8	7.7	3.5-14.7	1.4	5.1	2.0-10.7	1.0	2.1	0.7-4.8	0.6	2.4	0.5-7.2	0.4	1.3	0.2-5.4
	2001-2005	8.0	15.1	10.8-20.7	5.4	22.6	14.7-33.1	2.6	9.4	4.9-16.5	2.2	4.2	2.1-7.6	1.2	6.0	2.1-12.9	1.0	3.4	1.1-8.4
	2006-2010	7.0	11.6	8.1-16.3	5.6	19.1	12.5-27.9	1.4	4.4	1.8-9.6	1.0	1.7	0.6-4.2	1.0	3.3	1.1-7.9	0.0	0.0	0.0-3.2
	2011-2015	8.2	13.2	9.4-18.0	5.0	15.6	10.0-23.5	3.2	10.4	5.9-17.2	1.2	1.8	0.7-4.2	0.4	1.0	0.1-4.3	0.8	2.2	0.6-6.4
	2016-2020	6.2	10.0	6.7-14.4	4.2	13.5	8.2-21.1	2.0	6.4	3.0-12.4	0.8	1.3	0.3-3.5	0.8	2.9	0.8-7.5	0.0	0.0	0.0-3.4
Esophagus	1976-1980	1.0	2.6	0.8-6.1	0.8	5.4	1.3-14.1	0.2	0.9	0.0-5.0	1.6	3.9	1.7-7.8	1.4	8.3	3.0-17.6	0.2	0.9	0.0-5.0
	1981-1985	2.2	5.2	2.6-9.4	1.4	8.9	3.4-18.5	0.8	3.0	0.8-8.0	1.8	4.3	2.0-8.3	1.2	7.9	2.7-17.1	0.6	2.2	0.5-6.9
	1986-1990	3.0	7.1	3.9-11.7	2.0	12.1	5.4-22.6	1.0	4.6	1.4-10.8	1.6	4.1	1.8-8.1	1.2	9.4	3.2-20.2	0.4	1.9	0.2-6.9
	1991-1995	1.6	3.6	1.6-7.1	0.4	2.1	0.3-7.4	1.2	4.9	1.8-10.9	2.6	5.7	3.0-9.7	1.8	8.9	4.0-16.9	0.8	3.1	0.8-8.3
	1996-2000	4.2	8.5	5.3-13.0	3.6	16.2	9.5-25.7	0.6	2.2	0.5-6.8	3.8	7.7	4.6-12.0	2.6	11.3	6.0-19.4	1.2	4.4	1.6-9.8
	2001-2005	3.2	6.0	3.4-9.9	2.2	9.4	4.6-17.0	1.0	3.7	1.2-8.8	2.2	4.2	2.1-7.5	1.8	8.7	3.9-16.5	0.4	1.5	0.2-5.8
	2006-2010	3.6	6.0	3.5-9.5	3.0	10.1	5.5-17.0	0.6	2.2	0.4-6.6	2.0	3.4	1.6-6.4	1.4	4.9	1.9-10.6	0.6	2.1	0.4-6.5
	2011-2015	3.6	5.8	3.4-9.4	3.2	10.5	5.9-17.4	0.4	1.6	0.2-6.1	2.8	4.6	2.5-8.0	2.4	8.0	4.1-14.4	0.4	1.4	0.2-5.6
	2016-2020	5.0	7.4	4.7-11.2	4.4	13.0	8.0-20.2	0.6	1.6	0.3-5.7	4.0	6.0	3.6-9.5	3.4	10.3	5.9-16.9	0.6	2.1	0.4-6.9
Stomach	1976-1980	2.4	6.5	3.3-11.3	0.4	2.0	0.2-7.5	2.0	9.4	4.4-17.5	2.2	5.5	2.7-9.8	0.8	4.9	1.3-12.5	1.4	6.3	2.5-13.0
	1981-1985	4.2	9.5	5.8-14.6	2.2	11.8	5.8-21.3	2.0	7.3	3.5-13.8	3.0	6.5	3.6-10.8	1.4	7.0	2.7-14.7	1.6	6.0	2.6-12.1
	1986-1990	3.8	8.9	5.4-14.0	1.8	11.3	5.0-21.5	2.0	7.3	3.5-13.8	2.8	6.6	3.6-11.1	1.4	8.3	3.3-16.9	1.4	5.1	2.0-10.8
	1991-1995	3.2	6.8	3.9-11.1	1.8	8.7	3.9-16.6	1.4	5.2	2.0-10.9	2.2	4.7	2.3-8.4	1.4	6.8	2.7-14.2	0.8	2.7	0.7-7.2
	1996-2000	3.0	6.0	3.3-9.9	2.0	8.6	4.1-16.0	1.0	3.2	1.0-7.9	1.2	2.4	0.9-5.2	0.8	3.7	1.0-9.3	0.4	1.6	0.2-6.0
	2001-2005	3.2	6.2	3.5-10.0	1.4	5.9	2.3-12.4	1.8	6.2	2.8-12.2	1.6	3.1	1.3-6.2	0.8	3.5	0.9-9.1	0.8	2.5	0.7-7.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Franklin**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	2.4	4.5	2.3-7.9	1.2	4.8	1.7-10.6	1.2	4.3	1.5-9.8	0.8	1.4	0.4-3.7	0.4	1.8	0.2-6.2	0.4	1.3	0.2-5.3
	2011-2015	2.8	4.8	2.6-8.1	2.0	6.8	3.2-12.8	0.8	2.8	0.7-7.6	0.4	0.7	0.1-2.8	0.2	0.7	0.0-4.3	0.2	0.9	0.0-4.9
	2016-2020	4.2	6.2	3.8-9.8	2.2	6.7	3.3-12.5	2.0	5.9	2.8-11.6	1.6	2.5	1.1-5.1	0.8	2.3	0.6-6.5	0.8	2.6	0.7-7.3
Colorectal	1976-1980	27.4	66.1	55.3-78.4	13.6	77.4	59.3-99.2	13.8	58.0	44.9-73.8	13.2	32.7	25.1-41.7	5.6	36.5	23.6-53.6	7.6	31.2	22.0-43.2
	1981-1985	35.6	84.0	71.9-97.5	19.0	105.1	84.4-129.3	16.6	67.6	53.6-84.3	16.4	39.9	31.6-49.7	7.8	45.1	31.7-62.1	8.6	35.5	25.6-48.2
	1986-1990	28.0	65.0	54.6-76.9	14.2	76.3	59.1-96.8	13.8	53.8	41.7-68.4	15.0	35.0	27.5-44.0	8.0	46.4	32.6-63.9	7.0	26.5	18.4-37.1
	1991-1995	31.0	64.6	54.8-75.7	16.2	80.9	63.7-101.1	14.8	53.5	41.9-67.5	12.2	25.8	19.7-33.2	5.4	26.1	17.0-38.0	6.8	24.3	16.7-34.2
	1996-2000	35.4	70.9	60.9-82.2	15.8	73.2	57.7-91.4	19.6	66.9	54.2-81.9	14.0	28.2	22.0-35.6	7.2	36.1	25.0-50.1	6.8	21.7	15.0-30.8
	2001-2005	37.2	70.9	61.1-81.9	19.2	82.4	66.4-101.0	18.0	61.4	49.3-75.9	11.6	22.2	16.9-28.8	5.2	24.7	15.9-36.3	6.4	21.4	14.6-30.6
	2006-2010	30.0	53.2	45.0-62.6	16.6	65.2	51.4-81.4	13.4	44.8	34.6-57.3	11.8	21.6	16.4-28.0	6.2	25.6	17.1-36.6	5.6	18.0	11.9-26.4
	2011-2015	25.2	43.0	35.7-51.3	14.2	51.6	40.0-65.5	11.0	37.0	27.6-48.7	9.8	17.0	12.5-22.7	5.2	19.6	12.6-28.9	4.6	16.1	10.0-24.7
2016-2020	25.8	40.6	33.7-48.5	12.2	37.7	28.6-48.9	13.6	42.4	32.5-54.7	9.8	15.2	11.2-20.4	4.0	12.5	7.6-19.8	5.8	16.7	11.0-24.7	
Colon excluding rectum	1976-1980	18.8	45.1	36.3-55.4	8.8	51.4	36.7-69.9	10.0	41.1	30.4-54.4	9.6	24.1	17.6-32.0	4.0	26.4	15.5-41.3	5.6	23.2	15.3-33.8
	1981-1985	24.4	57.9	47.9-69.3	12.0	66.8	50.5-86.5	12.4	50.5	38.5-65.1	13.6	33.6	26.0-42.7	6.0	35.0	23.4-50.2	7.6	31.7	22.3-43.8
	1986-1990	19.2	44.5	35.9-54.4	9.2	48.5	35.2-65.2	10.0	39.3	29.0-52.2	13.4	30.9	23.8-39.3	7.0	38.8	26.6-54.4	6.4	24.3	16.6-34.5
	1991-1995	23.6	49.1	40.6-58.8	11.4	56.4	42.3-73.5	12.2	43.7	33.3-56.4	10.6	22.5	16.8-29.5	4.6	22.0	13.8-33.2	6.0	21.5	14.4-31.1
	1996-2000	27.4	54.9	46.1-64.9	11.2	53.0	39.8-68.9	16.2	55.3	43.8-69.1	12.4	25.0	19.2-32.0	5.8	29.1	19.4-41.8	6.6	21.0	14.4-30.0
	2001-2005	27.6	52.7	44.3-62.3	12.6	55.7	42.5-71.6	15.0	51.0	40.0-64.3	9.4	18.1	13.3-24.1	3.6	17.7	10.3-28.0	5.8	19.3	12.9-28.1
	2006-2010	21.4	38.8	31.8-47.0	11.4	46.0	34.4-60.1	10.0	33.5	24.7-44.5	9.4	17.3	12.7-23.1	5.0	20.8	13.3-31.0	4.4	14.3	8.9-22.0
	2011-2015	17.6	30.2	24.1-37.3	9.8	36.1	26.5-48.1	7.8	26.0	18.4-36.0	8.0	13.9	9.9-19.0	4.2	16.0	9.8-24.7	3.8	13.1	7.7-21.0
2016-2020	21.6	33.4	27.2-40.6	9.8	29.9	21.9-40.0	11.8	35.7	26.9-46.9	7.6	11.9	8.3-16.5	2.6	7.9	4.1-14.0	5.0	14.5	9.2-22.1	
Rectum & rectosigmoid	1976-1980	8.6	21.0	15.0-28.4	4.8	26.0	16.3-39.3	3.8	17.0	10.0-26.8	3.6	8.6	5.1-13.7	1.6	10.2	4.1-20.3	2.0	8.0	3.8-15.0
	1981-1985	11.2	26.1	19.6-34.1	7.0	38.3	26.2-53.9	4.2	17.1	10.5-26.5	2.8	6.3	3.4-10.7	1.8	10.1	4.4-19.6	1.0	3.9	1.2-9.3
	1986-1990	8.8	20.6	14.9-27.7	5.0	27.8	17.7-41.3	3.8	14.5	8.7-22.9	1.6	4.2	1.8-8.1	1.0	7.7	2.3-17.4	0.6	2.2	0.5-6.8
	1991-1995	7.4	15.5	10.9-21.5	4.8	24.5	15.4-36.8	2.6	9.8	5.2-17.1	1.6	3.3	1.4-6.5	0.8	4.0	1.1-10.3	0.8	2.7	0.7-7.4
	1996-2000	8.0	16.1	11.5-21.9	4.6	20.2	12.8-30.4	3.4	11.6	6.8-18.9	1.6	3.2	1.4-6.4	1.4	7.0	2.6-14.5	0.2	0.7	0.0-4.4
	2001-2005	9.6	18.2	13.4-24.2	6.6	26.7	18.2-37.7	3.0	10.5	5.8-17.6	2.2	4.1	2.1-7.5	1.6	7.0	2.9-13.9	0.6	2.1	0.4-6.6
	2006-2010	8.6	14.4	10.4-19.5	5.2	19.2	12.3-28.5	3.4	11.3	6.5-18.5	2.4	4.3	2.2-7.6	1.2	4.8	1.7-10.5	1.2	3.7	1.4-8.6
	2011-2015	7.6	12.8	9.0-17.7	4.4	15.5	9.6-23.8	3.2	10.9	6.1-18.3	1.8	3.2	1.4-6.1	1.0	3.6	1.1-8.6	0.8	3.0	0.8-8.1
2016-2020	4.2	7.2	4.4-11.3	2.4	7.8	4.0-14.0	1.8	6.7	2.8-13.5	2.2	3.4	1.7-6.3	1.4	4.6	1.8-9.8	0.8	2.2	0.6-6.6	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Franklin**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	1.0	2.2	0.7-5.3	0.8	4.4	1.1-11.5	0.2	0.8	0.0-4.7	0.6	1.2	0.2-3.7	0.6	2.7	0.5-8.7	0.0	0.0	0.0-3.4
	1981-1985	1.4	3.5	1.4-7.3	0.8	4.9	1.1-12.8	0.6	3.2	0.6-9.3	1.0	2.5	0.8-5.8	0.8	4.8	1.1-12.7	0.2	1.4	0.0-6.6
	1986-1990	1.4	3.7	1.5-7.5	0.8	4.7	1.3-11.8	0.6	2.2	0.5-6.9	0.6	1.5	0.3-4.2	0.0	0.0	0.0-3.9	0.6	2.2	0.5-6.9
	1991-1995	2.4	5.3	2.7-9.3	1.0	5.6	1.7-13.0	1.4	5.5	2.1-11.5	1.2	2.6	0.9-5.6	0.4	1.9	0.2-6.8	0.8	2.8	0.8-7.4
	1996-2000	3.0	6.0	3.4-10.0	2.0	9.0	4.3-16.5	1.0	3.2	1.0-7.9	3.0	6.1	3.4-10.0	2.2	10.5	5.1-18.8	0.8	2.7	0.7-7.5
	2001-2005	3.4	6.4	3.7-10.4	2.2	9.2	4.5-16.7	1.2	4.1	1.5-9.2	2.4	4.6	2.4-8.1	1.6	7.4	3.1-14.5	0.8	2.8	0.7-7.7
	2006-2010	4.2	7.3	4.5-11.2	3.0	10.8	5.9-18.1	1.2	4.0	1.5-9.2	2.2	4.1	2.0-7.4	1.8	8.0	3.5-15.3	0.4	1.5	0.2-5.6
	2011-2015	3.4	5.2	3.0-8.6	2.2	6.5	3.2-12.1	1.2	3.7	1.3-8.6	3.0	4.4	2.4-7.4	2.2	6.5	3.2-12.2	0.8	2.2	0.6-6.4
	2016-2020	6.0	9.0	6.0-13.1	3.6	10.5	6.1-17.1	2.4	7.5	3.7-13.9	3.2	4.8	2.7-8.1	2.0	5.9	2.8-11.3	1.2	3.4	1.2-8.2
Pancreas	1976-1980	4.6	10.9	6.9-16.5	2.2	12.2	6.0-21.9	2.4	10.2	5.2-18.1	5.2	12.5	8.1-18.5	2.4	13.6	6.9-24.1	2.8	11.3	6.1-19.3
	1981-1985	5.4	13.4	8.8-19.5	3.0	20.6	11.2-34.0	2.4	9.9	5.1-17.6	6.0	15.0	10.0-21.5	2.8	18.9	10.1-31.9	3.2	12.8	7.2-21.1
	1986-1990	6.0	14.0	9.4-20.1	3.2	18.6	10.4-30.3	2.8	10.5	5.7-17.8	6.0	14.0	9.4-20.0	3.0	17.6	9.7-29.2	3.0	11.3	6.3-18.9
	1991-1995	5.4	11.5	7.6-16.8	2.4	12.4	6.3-21.8	3.0	10.3	5.8-17.3	5.6	11.6	7.7-16.8	2.2	10.3	5.1-18.5	3.4	11.8	6.9-19.2
	1996-2000	5.4	10.7	7.1-15.6	3.0	14.6	8.0-24.2	2.4	7.8	4.0-14.0	5.0	10.0	6.5-14.8	3.0	16.5	9.0-27.2	2.0	6.5	3.1-12.4
	2001-2005	5.8	10.9	7.3-15.7	3.6	14.2	8.3-22.6	2.2	7.9	3.9-14.4	5.2	9.8	6.4-14.5	3.8	16.1	9.5-25.5	1.4	4.8	1.9-10.3
	2006-2010	8.8	15.3	11.1-20.7	4.8	17.4	10.9-26.3	4.0	13.3	8.1-20.9	7.2	12.9	9.0-17.9	2.4	10.1	5.1-17.9	4.8	15.8	10.1-23.8
	2011-2015	6.8	11.2	7.7-15.8	3.4	12.4	7.1-20.1	3.4	10.2	5.9-16.8	6.0	10.1	6.8-14.6	3.0	10.8	6.0-18.1	3.0	9.1	5.1-15.5
	2016-2020	10.6	15.9	11.9-21.1	6.4	21.5	14.5-30.6	4.2	12.0	7.4-19.1	7.6	11.1	7.8-15.4	4.8	15.9	10.1-24.1	2.8	7.5	4.1-13.3
Larynx	1976-1980	2.4	6.2	3.1-10.9	1.8	12.1	5.2-23.1	0.6	2.3	0.5-7.2	1.4	3.2	1.3-6.6	1.2	6.4	2.3-14.2	0.2	0.8	0.0-4.7
	1981-1985	3.0	6.9	3.8-11.5	2.2	10.6	5.2-19.3	0.8	3.7	1.0-9.7	1.2	2.8	1.0-6.1	0.8	4.4	1.1-11.4	0.4	1.5	0.2-5.9
	1986-1990	3.8	8.9	5.3-13.9	2.4	12.4	6.2-22.2	1.4	6.2	2.4-13.0	1.2	3.1	1.1-6.6	0.6	4.2	0.7-11.9	0.6	2.9	0.6-8.4
	1991-1995	1.2	2.4	0.9-5.3	0.8	3.5	0.9-9.1	0.4	1.4	0.2-5.5	0.6	1.1	0.2-3.3	0.4	1.6	0.2-6.1	0.2	0.7	0.0-4.2
	1996-2000	5.4	11.0	7.3-16.0	4.0	17.9	10.9-27.7	1.4	5.5	2.2-11.5	0.6	1.2	0.2-3.5	0.0	0.0	0.0-3.2	0.6	2.3	0.5-6.9
	2001-2005	4.4	8.1	5.1-12.3	3.8	14.0	8.4-22.1	0.6	2.3	0.5-6.9	2.0	3.9	1.9-7.2	1.4	5.9	2.3-12.2	0.6	1.9	0.4-6.2
	2006-2010	3.6	6.1	3.6-9.7	3.4	12.3	7.0-20.1	0.2	0.7	0.0-4.3	1.2	2.0	0.7-4.5	1.0	3.6	1.1-8.6	0.2	0.6	0.0-4.2
	2011-2015	3.8	5.6	3.4-9.0	2.6	7.6	4.0-13.5	1.2	3.8	1.4-8.8	1.6	2.5	1.1-5.1	1.4	4.7	1.8-9.9	0.2	0.5	0.0-4.0
	2016-2020	2.6	3.4	1.8-6.2	2.2	5.9	2.9-11.0	0.4	1.0	0.1-4.9	1.0	1.4	0.4-3.5	0.8	2.3	0.6-6.4	0.2	0.5	0.0-4.2
Lung and bronchus	1976-1980	28.4	62.8	52.8-74.3	20.0	100.6	81.6-122.9	8.4	33.5	24.1-45.6	23.0	52.5	43.2-63.2	17.2	92.7	73.5-115.3	5.8	24.1	16.0-35.1
	1981-1985	36.8	81.6	70.1-94.5	22.6	118.0	96.6-142.7	14.2	56.0	43.5-71.1	28.8	64.6	54.4-76.3	18.4	100.9	80.7-124.5	10.4	40.4	30.0-53.3
	1986-1990	39.4	86.9	75.0-100.1	25.6	133.3	110.5-159.4	13.8	54.9	42.4-69.9	35.8	79.7	68.3-92.5	22.8	118.6	97.3-143.1	13.0	51.6	39.6-66.2
	1991-1995	39.4	82.8	71.5-95.2	22.6	113.7	93.1-137.4	16.8	62.6	49.8-77.9	33.0	69.5	59.2-81.0	20.8	107.5	87.2-130.9	12.2	45.4	34.6-58.6
	1996-2000	40.8	82.1	71.2-94.2	24.6	116.0	96.0-138.7	16.2	59.0	46.7-73.7	30.8	62.1	52.7-72.7	18.4	88.4	70.9-108.6	12.4	45.3	34.6-58.4
	2001-2005	43.6	82.8	72.2-94.6	25.0	106.9	88.7-127.7	18.6	65.9	53.1-81.0	31.8	61.0	51.8-71.2	18.6	83.6	67.2-102.7	13.2	46.2	35.7-59.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Franklin**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	47.4	83.1	72.7-94.5	26.4	102.0	84.9-121.5	21.0	69.4	56.7-84.3	30.2	53.5	45.2-62.8	17.6	69.3	55.2-85.8	12.6	42.0	32.2-54.0
	2011-2015	47.2	78.6	68.8-89.6	25.4	89.6	74.3-107.1	21.8	70.4	57.6-85.5	34.6	57.1	48.8-66.5	18.2	65.8	52.6-81.3	16.4	51.2	40.5-64.0
	2016-2020	53.0	77.9	68.6-88.2	29.0	86.0	72.2-101.9	24.0	70.8	58.4-85.4	26.6	39.4	32.9-47.0	13.6	42.7	32.9-54.7	13.0	36.9	28.3-47.6
Melanoma of the skin	1976-1980	2.2	5.2	2.6-9.5	1.0	5.2	1.6-12.4	1.2	4.8	1.7-10.6	0.8	1.8	0.5-4.8	0.4	1.7	0.2-6.8	0.4	1.6	0.2-6.2
	1981-1985	2.8	7.4	4.0-12.4	1.8	10.5	4.7-19.9	1.0	4.3	1.4-10.3	0.8	2.0	0.5-5.0	0.2	1.3	0.0-6.7	0.6	2.7	0.5-8.1
	1986-1990	3.6	8.6	5.1-13.6	2.0	11.0	5.0-20.5	1.6	7.1	3.0-14.2	0.6	1.3	0.3-3.9	0.0	0.0	0.0-3.9	0.6	2.3	0.5-7.0
	1991-1995	3.4	7.0	4.1-11.3	1.6	7.5	3.2-14.8	1.8	7.0	3.1-13.4	1.2	2.5	0.9-5.4	1.0	4.7	1.5-11.0	0.2	0.7	0.0-4.4
	1996-2000	6.4	12.6	8.6-17.8	3.2	12.7	7.2-20.8	3.2	12.9	7.3-21.1	0.4	0.8	0.1-2.9	0.4	2.2	0.2-7.7	0.0	0.0	0.0-3.2
	2001-2005	11.2	21.2	16.0-27.6	6.0	23.7	15.9-34.0	5.2	20.2	13.1-29.9	1.2	2.3	0.9-5.1	1.0	4.7	1.5-11.0	0.2	0.6	0.0-4.2
	2006-2010	10.8	19.2	14.4-25.2	6.6	25.0	17.0-35.6	4.2	15.9	9.7-24.7	1.4	2.4	1.0-5.1	1.0	4.2	1.3-9.8	0.4	1.3	0.2-5.2
	2011-2015	10.8	17.7	13.2-23.3	6.6	21.7	14.8-30.9	4.2	14.1	8.6-22.1	1.2	2.3	0.8-5.0	0.8	3.4	0.9-8.7	0.4	1.3	0.2-5.4
2016-2020	12.2	19.2	14.6-25.0	7.2	22.4	15.5-31.5	5.0	18.1	11.3-27.6	0.6	0.9	0.2-2.9	0.4	1.6	0.2-5.7	0.2	0.5	0.0-4.0	
Female breast	1976-1980							17.2	77.3	61.3-96.1							8.8	38.0	27.4-51.5
	1981-1985							20.2	83.5	67.7-102.1							8.2	34.5	24.5-47.2
	1986-1990							27.0	114.5	95.5-136.1							9.6	40.6	29.7-54.3
	1991-1995							29.2	116.8	98.3-137.8							9.0	33.9	24.6-45.7
	1996-2000							37.2	141.6	121.7-163.8							7.0	25.5	17.7-35.8
	2001-2005							34.4	126.8	108.4-147.5							6.0	20.8	14.0-30.1
	2006-2010							32.4	109.9	93.4-128.6							6.4	22.4	15.2-32.0
	2011-2015							32.4	107.3	90.9-126.0							4.6	14.7	9.2-22.6
2016-2020							33.0	109.7	92.7-129.1							6.6	20.0	13.5-28.8	
Cervix uteri	1976-1980							4.6	24.4	15.1-36.8							2.0	9.2	4.3-17.1
	1981-1985							3.2	14.8	8.3-24.3							0.8	3.7	0.9-9.6
	1986-1990							1.8	9.3	4.2-17.7							0.8	4.0	1.0-10.1
	1991-1995							1.6	6.8	2.9-13.5							1.0	4.8	1.5-11.1
	1996-2000							3.4	13.6	7.9-22.0							0.8	3.1	0.8-8.1
	2001-2005							0.8	3.3	0.9-8.6							0.8	2.7	0.7-7.3
	2006-2010							1.8	6.8	3.1-13.2							0.6	1.8	0.4-5.9
	2011-2015							0.6	2.1	0.4-6.5							0.0	0.0	0.0-3.3
	2016-2020							1.6	5.8	2.4-11.9							0.8	2.9	0.7-8.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Franklin**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							5.2	21.0	13.6-31.0							1.0	4.4	1.3-10.6
	1981-1985							9.6	41.4	30.1-55.5							2.2	8.3	4.1-15.1
	1986-1990							7.4	30.8	21.5-42.9							0.4	1.5	0.2-5.8
	1991-1995							5.2	20.5	13.3-30.4							1.2	4.7	1.7-10.4
	1996-2000							7.0	27.8	19.3-38.9							1.2	4.0	1.5-9.1
	2001-2005							5.0	18.9	12.2-28.1							0.6	2.0	0.4-6.3
	2006-2010							9.4	33.3	24.3-44.7							1.6	5.1	2.2-10.5
	2011-2015							7.2	25.0	17.3-35.2							1.0	3.2	1.0-8.0
	2016-2020							8.4	23.5	16.8-32.5							1.2	3.3	1.2-8.0
Ovary	1976-1980							2.4	9.3	4.8-16.5							1.8	6.9	3.2-13.5
	1981-1985							2.6	10.4	5.4-18.2							2.4	10.8	5.4-19.2
	1986-1990							4.2	17.5	10.7-27.1							2.2	9.1	4.5-16.5
	1991-1995							3.6	15.1	8.8-24.0							2.8	9.8	5.3-16.8
	1996-2000							6.0	22.8	15.3-32.8							3.2	11.0	6.3-18.3
	2001-2005							4.8	18.1	11.6-27.2							3.0	11.1	6.2-18.6
	2006-2010							3.0	10.2	5.6-17.2							3.2	10.3	5.8-17.0
	2011-2015							1.6	5.3	2.3-11.0							1.0	3.0	1.0-7.7
	2016-2020							2.4	7.9	3.8-14.6							1.6	4.9	1.9-10.6
Prostate	1976-1980				14.8	88.9	68.7-112.9							6.2	40.7	27.1-58.2			
	1981-1985				18.2	110.8	88.0-137.3							5.4	35.2	22.7-51.7			
	1986-1990				20.0	116.9	94.2-143.1							4.2	25.6	15.4-39.6			
	1991-1995				34.6	178.7	152.3-208.2							6.0	37.4	24.8-53.6			
	1996-2000				29.2	138.0	116.1-162.6							5.6	29.5	19.4-42.6			
	2001-2005				41.0	173.0	149.7-199.0							2.2	11.9	5.9-21.0			
	2006-2010				39.4	141.2	121.7-163.0							4.4	21.2	13.1-32.2			
	2011-2015				47.8	152.0	132.8-173.3							4.8	21.0	13.4-31.1			
	2016-2020				48.8	141.9	124.2-161.6							4.4	16.3	10.2-24.9			
Testis	1976-1980				0.2	1.0	0.0-6.0							0.0	0.0	0.0-4.2			
	1981-1985				1.2	6.6	2.4-14.5							0.2	1.4	0.0-7.1			
	1986-1990				0.4	1.5	0.2-6.3							0.0	0.0	0.0-3.9			
	1991-1995				1.4	4.4	1.7-9.7							0.2	0.9	0.0-5.2			
	1996-2000				0.4	1.5	0.2-5.6							0.0	0.0	0.0-3.2			
	2001-2005				1.6	5.0	2.2-10.3							0.2	0.6	0.0-4.1			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Franklin**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				0.8	2.5	0.7-6.8							0.0	0.0	0.0-2.9			
	2011-2015				2.2	6.9	3.4-12.7							0.0	0.0	0.0-2.9			
	2016-2020				1.4	4.8	1.9-10.2							0.2	0.7	0.0-4.0			
Urinary bladder (incl. in situ)	1976-1980	11.0	26.0	19.5-34.0	8.0	46.4	32.8-63.7	3.0	12.3	6.8-20.5	3.8	9.6	5.7-15.0	2.4	15.0	7.5-26.6	1.4	5.6	2.3-11.8
	1981-1985	11.6	27.0	20.4-35.0	9.6	55.8	40.6-74.6	2.0	8.3	3.9-15.5	2.0	4.6	2.2-8.5	1.4	7.6	3.0-15.9	0.6	2.2	0.5-7.0
	1986-1990	11.2	25.1	18.9-32.7	8.0	43.1	30.4-59.1	3.2	12.0	6.9-19.8	2.6	5.8	3.0-9.9	1.8	10.5	4.7-19.9	0.8	2.8	0.8-7.6
	1991-1995	11.2	23.8	17.9-30.9	8.4	44.4	31.5-60.5	2.8	10.1	5.5-17.2	2.2	4.6	2.3-8.3	1.6	7.9	3.4-15.6	0.6	2.0	0.4-6.2
	1996-2000	18.8	38.1	30.8-46.7	13.2	66.4	51.0-84.7	5.6	20.9	13.7-30.6	3.0	6.1	3.4-10.1	2.2	13.8	6.8-24.2	0.8	3.0	0.8-7.9
	2001-2005	15.4	29.4	23.2-36.8	10.6	45.7	34.0-60.1	4.8	16.6	10.6-25.1	3.4	6.6	3.8-10.6	1.8	9.6	4.3-18.0	1.6	5.4	2.3-11.1
	2006-2010	16.8	30.7	24.5-38.1	12.2	50.0	38.0-64.5	4.6	15.1	9.5-23.0	4.2	7.8	4.8-12.0	2.8	11.7	6.3-19.8	1.4	4.6	1.8-9.9
	2011-2015	14.8	24.6	19.3-31.1	10.8	38.4	28.5-50.5	4.0	12.8	7.7-20.3	3.8	6.4	3.9-10.2	2.6	10.9	5.7-18.7	1.2	3.4	1.2-8.1
	2016-2020	17.4	26.8	21.4-33.4	13.6	43.9	33.8-56.2	3.8	12.3	7.1-20.0	2.8	4.3	2.4-7.5	1.4	5.1	2.0-10.8	1.4	3.6	1.4-8.2
Kidney and renal pelvis	1976-1980	1.6	3.6	1.5-7.3	1.2	6.0	2.1-13.4	0.4	1.3	0.2-5.5	1.0	2.6	0.8-6.1	0.8	4.6	1.2-11.8	0.2	0.8	0.0-4.7
	1981-1985	5.2	12.4	8.0-18.3	2.6	15.0	7.6-26.3	2.6	11.3	5.9-19.5	2.0	5.1	2.4-9.3	1.2	8.2	2.7-18.1	0.8	3.5	0.9-9.2
	1986-1990	5.8	13.6	9.1-19.7	3.8	20.3	12.1-31.9	2.0	7.7	3.7-14.5	4.2	9.5	5.8-14.6	2.4	12.7	6.5-22.4	1.8	6.6	3.0-12.8
	1991-1995	6.0	13.0	8.7-18.6	3.0	14.9	8.2-24.6	3.0	11.6	6.4-19.3	2.8	6.0	3.3-10.1	1.6	8.5	3.6-16.6	1.2	4.2	1.5-9.4
	1996-2000	6.2	12.5	8.5-17.7	3.4	16.4	9.4-26.3	2.8	10.5	5.7-17.9	2.6	5.3	2.8-9.0	0.8	4.1	1.1-10.3	1.8	6.0	2.7-11.9
	2001-2005	9.4	17.6	12.9-23.5	5.8	23.2	15.4-33.7	3.6	12.9	7.6-20.7	2.4	4.6	2.4-8.0	1.8	8.3	3.7-15.9	0.6	2.3	0.5-6.9
	2006-2010	9.6	17.2	12.6-22.8	5.4	20.5	13.3-30.1	4.2	14.8	9.1-23.0	2.4	4.2	2.1-7.4	1.8	6.4	2.9-12.3	0.6	1.8	0.4-5.8
	2011-2015	11.2	18.6	14.0-24.4	6.2	20.5	13.7-29.5	5.0	17.5	11.1-26.5	2.6	4.1	2.2-7.1	1.6	5.3	2.2-10.8	1.0	3.1	1.0-7.7
	2016-2020	11.0	16.1	12.0-21.2	7.4	20.7	14.4-29.0	3.6	10.5	6.1-17.3	2.8	4.6	2.5-7.9	1.8	6.0	2.7-11.7	1.0	3.0	0.9-7.9
Brain and other central nervous system	1976-1980	2.0	4.3	2.0-8.0	1.4	5.9	2.3-12.9	0.6	2.7	0.5-8.2	1.2	2.3	0.8-5.2	0.6	2.3	0.4-7.6	0.6	2.2	0.4-6.8
	1981-1985	3.6	8.1	4.7-12.9	2.0	9.5	4.4-18.0	1.6	6.2	2.7-12.6	2.2	5.3	2.6-9.6	1.6	8.1	3.4-16.3	0.6	2.3	0.5-7.2
	1986-1990	3.4	7.6	4.3-12.2	2.2	11.9	5.7-21.8	1.2	4.2	1.5-9.5	1.6	3.8	1.6-7.6	1.2	6.4	2.3-14.1	0.4	1.3	0.2-5.4
	1991-1995	4.2	8.8	5.4-13.5	2.0	8.9	4.2-16.7	2.2	8.7	4.3-15.7	2.4	4.8	2.5-8.5	1.0	4.6	1.5-10.8	1.4	5.0	2.0-10.6
	1996-2000	3.0	5.8	3.2-9.6	2.0	7.1	3.3-13.4	1.0	4.5	1.4-10.4	1.6	3.3	1.4-6.4	1.0	4.1	1.3-9.8	0.6	2.2	0.5-6.9
	2001-2005	4.6	8.8	5.6-13.3	2.6	9.7	5.1-16.8	2.0	6.8	3.2-12.8	1.6	3.0	1.3-6.0	1.2	4.6	1.7-10.2	0.4	1.5	0.2-5.8
	2006-2010	4.0	7.5	4.6-11.7	1.6	6.1	2.5-12.2	2.4	8.9	4.5-16.0	2.4	4.5	2.3-7.9	1.4	5.3	2.0-11.2	1.0	3.6	1.1-8.8
	2011-2015	4.0	6.9	4.2-10.8	2.0	6.7	3.2-12.7	2.0	6.6	3.1-12.7	3.0	4.9	2.7-8.2	1.0	3.6	1.1-8.6	2.0	6.1	2.9-11.7
	2016-2020	2.6	4.5	2.3-8.0	1.0	3.6	1.1-8.6	1.6	5.6	2.3-11.8	2.0	3.2	1.5-6.1	1.0	3.0	0.9-7.3	1.0	3.4	1.0-8.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Franklin**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	1.2	3.1	1.1-6.7	0.2	1.4	0.0-7.0	1.0	4.0	1.3-9.6	0.8	1.8	0.5-4.7	0.0	0.0	0.0-4.2	0.8	3.0	0.8-8.1
	1981-1985	1.0	2.5	0.8-5.9	0.4	1.7	0.2-6.9	0.6	3.3	0.6-9.5	0.4	0.9	0.1-3.4	0.0	0.0	0.0-4.1	0.4	1.5	0.2-5.9
	1986-1990	1.0	2.2	0.7-5.1	0.2	1.3	0.0-6.5	0.8	3.0	0.8-7.9	0.4	0.9	0.1-3.2	0.2	1.3	0.0-6.5	0.2	0.7	0.0-4.6
	1991-1995	0.8	1.5	0.4-4.0	0.4	1.7	0.2-6.4	0.4	1.5	0.2-5.6	0.0	0.0	0.0-1.6	0.0	0.0	0.0-3.4	0.0	0.0	0.0-3.2
	1996-2000	2.8	5.4	3.0-9.1	0.8	3.2	0.9-8.3	2.0	8.5	4.1-15.7	0.4	0.8	0.1-2.9	0.2	0.8	0.0-4.6	0.2	0.7	0.0-4.4
	2001-2005	3.0	5.6	3.1-9.3	0.6	2.3	0.5-6.9	2.4	9.6	4.9-17.0	0.4	0.8	0.1-2.8	0.0	0.0	0.0-3.1	0.4	1.2	0.1-5.0
	2006-2010	5.2	9.5	6.2-14.1	1.8	5.8	2.6-11.2	3.4	13.4	7.7-21.8	0.0	0.0	0.0-1.5	0.0	0.0	0.0-2.9	0.0	0.0	0.0-3.2
	2011-2015	5.6	9.9	6.5-14.4	2.2	7.4	3.6-13.7	3.4	14.2	8.1-23.1	0.8	1.4	0.4-3.6	0.8	2.9	0.8-7.6	0.0	0.0	0.0-3.3
	2016-2020	3.4	6.0	3.4-9.8	1.4	3.9	1.6-8.6	2.0	8.9	4.1-16.8	0.2	0.2	0.0-1.9	0.2	0.5	0.0-3.6	0.0	0.0	0.0-3.4
Hodgkin lymphoma	1976-1980	1.4	3.4	1.3-7.2	0.8	3.0	0.8-8.6	0.6	3.7	0.7-10.4	0.4	0.8	0.1-3.1	0.2	0.8	0.0-5.5	0.2	0.8	0.0-4.8
	1981-1985	1.6	3.9	1.6-7.8	1.2	6.5	2.3-14.4	0.4	1.5	0.2-6.0	1.0	2.5	0.8-5.9	0.8	4.2	1.1-11.0	0.2	0.8	0.0-4.8
	1986-1990	0.8	1.7	0.5-4.4	0.4	1.4	0.2-6.0	0.4	1.5	0.2-5.8	0.6	1.3	0.3-3.8	0.0	0.0	0.0-3.9	0.6	2.3	0.5-7.1
	1991-1995	1.6	3.1	1.3-6.2	1.0	3.4	1.1-8.5	0.6	2.8	0.6-8.0	0.0	0.0	0.0-1.6	0.0	0.0	0.0-3.4	0.0	0.0	0.0-3.2
	1996-2000	1.6	3.2	1.4-6.3	1.0	3.1	1.0-7.8	0.6	2.6	0.5-7.6	0.2	0.4	0.0-2.2	0.0	0.0	0.0-3.2	0.2	0.9	0.0-4.9
	2001-2005	2.6	4.8	2.5-8.2	2.2	7.3	3.6-13.4	0.4	2.0	0.2-6.9	0.4	0.8	0.1-2.8	0.4	1.7	0.2-6.3	0.0	0.0	0.0-3.3
	2006-2010	1.4	2.5	1.0-5.2	1.0	2.9	0.9-7.3	0.4	1.4	0.2-5.4	0.0	0.0	0.0-1.5	0.0	0.0	0.0-2.9	0.0	0.0	0.0-3.2
	2011-2015	1.4	3.0	1.2-6.1	0.8	3.6	1.0-8.9	0.6	2.9	0.6-8.2	0.0	0.0	0.0-1.5	0.0	0.0	0.0-2.9	0.0	0.0	0.0-3.3
	2016-2020	1.8	3.8	1.7-7.2	1.2	3.8	1.4-8.7	0.6	3.5	0.7-9.7	0.4	0.5	0.1-2.3	0.4	1.0	0.1-4.5	0.0	0.0	0.0-3.4
Non-Hodgkin lymphomas	1976-1980	3.6	8.1	4.7-12.9	1.8	10.9	4.7-21.1	1.8	6.5	3.0-12.7	2.4	4.9	2.5-8.7	1.2	5.0	1.8-11.5	1.2	4.4	1.6-9.9
	1981-1985	4.6	10.9	6.8-16.4	1.8	9.4	4.2-18.1	2.8	11.8	6.3-20.3	1.8	4.0	1.8-7.7	1.0	5.5	1.6-13.0	0.8	3.0	0.8-8.1
	1986-1990	7.2	15.3	10.7-21.2	4.8	21.8	13.9-32.8	2.4	9.4	4.8-16.7	4.2	9.2	5.7-14.1	2.4	11.4	5.8-20.2	1.8	7.5	3.4-14.5
	1991-1995	8.0	17.6	12.5-24.0	4.8	22.2	14.0-33.4	3.2	12.5	7.1-20.6	2.6	5.4	2.9-9.3	1.8	9.3	4.1-17.8	0.8	3.2	0.8-8.2
	1996-2000	13.8	27.7	21.5-35.0	7.8	32.1	22.7-44.1	6.0	22.2	14.9-32.0	5.0	10.1	6.5-14.9	3.6	15.4	9.1-24.4	1.4	4.8	1.9-10.3
	2001-2005	10.2	19.6	14.6-25.8	7.2	30.7	21.3-42.7	3.0	10.6	5.9-17.8	3.2	6.1	3.5-10.0	1.6	6.9	2.9-13.6	1.6	5.6	2.4-11.3
	2006-2010	11.6	20.5	15.6-26.6	5.4	20.4	13.2-30.0	6.2	20.4	13.8-29.3	2.6	4.9	2.6-8.4	1.0	4.7	1.5-10.8	1.6	5.1	2.2-10.6
	2011-2015	11.6	19.5	14.8-25.4	6.6	23.9	16.2-33.9	5.0	15.7	10.1-23.7	2.8	4.9	2.7-8.4	1.8	8.2	3.7-15.3	1.0	2.8	0.9-7.3
	2016-2020	11.4	18.4	13.8-24.1	7.0	23.6	16.3-33.2	4.4	14.4	8.7-22.5	3.0	4.8	2.7-8.1	1.8	6.4	2.9-12.3	1.2	3.3	1.2-8.0
Myeloma	1976-1980	1.2	2.8	1.0-6.1	0.2	1.3	0.0-6.7	1.0	3.9	1.3-9.4	0.6	1.2	0.3-3.8	0.0	0.0	0.0-4.2	0.6	2.2	0.5-7.0
	1981-1985	2.6	6.2	3.3-10.7	1.2	7.3	2.5-16.1	1.4	5.3	2.1-11.3	1.0	2.5	0.8-5.8	0.2	0.9	0.0-5.5	0.8	3.1	0.8-8.3
	1986-1990	3.0	7.3	4.1-12.1	2.2	12.5	6.1-22.2	0.8	3.0	0.8-8.0	2.4	5.3	2.7-9.2	1.8	9.5	4.2-18.1	0.6	2.2	0.4-6.7
	1991-1995	4.4	9.0	5.6-13.7	2.6	12.7	6.6-21.9	1.8	6.1	2.8-11.8	3.0	6.4	3.6-10.5	1.4	7.5	2.9-15.5	1.6	5.3	2.3-10.8
	1996-2000	3.6	7.2	4.3-11.4	2.4	11.1	5.7-19.3	1.2	4.2	1.5-9.5	1.6	3.2	1.4-6.3	1.0	5.0	1.6-11.4	0.6	2.1	0.4-6.6
	2001-2005	2.6	5.0	2.6-8.5	1.6	7.0	2.9-13.9	1.0	3.3	1.0-8.1	1.0	1.9	0.6-4.5	1.0	4.0	1.3-9.4	0.0	0.0	0.0-3.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Franklin**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	5.2	8.8	5.7-13.0	2.8	10.6	5.6-18.1	2.4	7.8	4.0-14.0	2.4	4.1	2.1-7.3	1.0	3.9	1.2-9.3	1.4	4.5	1.8-9.7
	2011-2015	3.6	5.5	3.2-9.0	1.8	6.0	2.6-11.7	1.8	5.3	2.4-10.7	2.8	4.6	2.5-7.8	1.2	4.2	1.5-9.4	1.6	4.9	2.1-10.2
	2016-2020	4.4	6.8	4.2-10.5	2.8	9.1	4.9-15.6	1.6	4.7	2.0-9.9	1.8	2.6	1.2-5.1	0.8	2.4	0.6-6.5	1.0	2.8	0.9-7.3
Leukemias	1976-1980	5.0	11.6	7.5-17.3	3.0	15.9	8.8-26.5	2.0	8.3	3.9-15.5	2.0	4.9	2.3-9.0	1.4	7.6	3.0-15.9	0.6	2.6	0.5-7.7
	1981-1985	7.0	16.2	11.2-22.7	3.4	17.2	9.8-28.0	3.6	15.3	8.9-24.6	4.2	9.6	5.9-14.8	2.0	10.6	4.9-19.9	2.2	9.6	4.6-17.5
	1986-1990	3.4	8.2	4.7-13.1	2.6	13.8	7.2-23.8	0.8	3.0	0.8-8.0	2.0	4.9	2.3-9.0	1.2	6.7	2.4-14.5	0.8	3.0	0.8-7.9
	1991-1995	8.2	17.2	12.3-23.4	4.0	19.5	11.6-30.3	4.2	15.8	9.7-24.3	5.0	10.6	6.8-15.7	1.8	9.9	4.3-18.9	3.2	11.0	6.3-18.1
	1996-2000	9.4	19.1	14.0-25.4	5.8	25.7	17.1-37.1	3.6	12.2	7.1-19.6	3.2	6.1	3.5-10.0	2.0	8.1	3.7-15.3	1.2	3.7	1.4-8.6
	2001-2005	10.8	21.0	15.8-27.5	6.0	27.7	18.4-39.7	4.8	17.0	10.8-25.7	4.6	9.2	5.8-13.8	2.8	13.7	7.4-22.8	1.8	5.7	2.6-11.2
	2006-2010	8.4	16.0	11.5-21.7	4.0	15.5	9.3-24.1	4.4	16.3	10.1-25.0	6.4	12.0	8.2-17.0	3.2	12.5	7.0-20.5	3.2	10.7	6.1-17.8
	2011-2015	9.6	16.7	12.2-22.3	6.4	23.4	15.8-33.3	3.2	10.9	6.1-18.1	3.4	5.8	3.4-9.4	2.0	7.8	3.6-14.5	1.4	4.7	1.8-10.3
	2016-2020	9.6	15.5	11.3-20.9	6.2	21.0	14.1-30.3	3.4	10.9	6.2-18.2	3.8	6.2	3.7-9.8	2.2	8.6	4.3-15.5	1.6	4.2	1.8-9.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Fulton**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	220.6	365.5	343.6-388.4	110.2	445.4	407.4-486.0	110.4	320.3	293.2-349.3	127.4	211.5	195.0-229.1	67.0	275.8	245.7-308.5	60.4	172.7	153.2-194.1
	1981-1985	243.0	395.3	372.8-418.9	125.6	480.9	442.5-521.8	117.4	348.0	319.1-379.0	138.4	220.2	203.7-237.7	76.8	302.3	271.5-335.6	61.6	173.6	154.0-195.2
	1986-1990	258.2	413.8	391.1-437.6	134.2	517.5	477.8-559.6	124.0	354.5	326.0-384.9	133.8	216.1	199.7-233.6	70.4	279.7	250.3-311.6	63.4	174.4	155.0-195.7
	1991-1995	291.8	457.9	434.4-482.5	153.6	565.4	525.4-607.7	138.2	391.5	361.8-423.0	146.6	226.4	210.0-243.6	72.4	270.4	242.7-300.3	74.2	196.3	176.1-218.2
	1996-2000	338.0	522.4	497.6-548.2	172.0	616.6	575.8-659.6	166.0	472.1	439.7-506.3	137.6	206.2	190.9-222.3	71.6	260.3	233.8-289.0	66.0	173.0	154.4-193.5
	2001-2005	342.4	517.5	493.1-542.9	175.0	594.2	555.3-635.2	167.4	469.7	437.6-503.7	133.4	193.5	179.0-209.0	70.0	242.1	217.3-269.1	63.4	162.3	144.5-181.8
	2006-2010	363.8	530.9	506.4-556.4	189.0	603.1	564.7-643.5	174.8	482.7	450.1-517.1	128.4	181.4	167.5-196.3	61.0	196.7	174.9-220.5	67.4	174.1	155.4-194.6
	2011-2015	377.2	527.2	503.0-552.4	186.0	560.3	523.9-598.7	191.2	507.6	474.5-542.5	132.0	176.9	163.4-191.4	67.4	207.1	185.1-231.1	64.6	155.3	138.3-174.2
	2016-2020	385.6	504.7	481.5-528.8	199.2	549.8	515.0-586.6	186.4	474.8	443.1-508.3	126.2	161.1	148.4-174.7	65.0	184.6	164.5-206.7	61.2	144.9	128.5-163.2
Oral cavity and pharynx	1976-1980	4.8	8.6	5.4-12.9	3.4	16.2	9.2-26.1	1.4	4.2	1.7-9.0	2.6	4.5	2.4-7.9	2.0	8.6	3.9-16.0	0.6	1.8	0.4-5.6
	1981-1985	5.6	9.5	6.2-14.0	4.0	16.8	10.1-26.3	1.6	4.4	1.8-9.1	0.8	1.1	0.3-3.1	0.6	2.5	0.4-7.6	0.2	0.5	0.0-3.3
	1986-1990	4.8	7.5	4.7-11.3	3.4	12.5	7.2-20.2	1.4	3.8	1.5-8.2	2.0	3.1	1.5-5.8	1.6	6.4	2.6-12.8	0.4	1.2	0.1-4.7
	1991-1995	5.6	10.3	6.8-14.9	3.0	11.8	6.6-19.5	2.6	8.9	4.6-15.4	2.4	4.1	2.1-7.2	1.4	5.1	2.0-10.6	1.0	3.4	1.0-8.1
	1996-2000	5.8	9.5	6.3-13.7	4.0	14.3	8.7-22.2	1.8	5.9	2.7-11.3	2.0	3.2	1.5-6.0	1.2	4.1	1.5-9.1	0.8	2.5	0.7-6.6
	2001-2005	9.4	14.1	10.3-18.9	6.6	22.5	15.4-31.7	2.8	7.9	4.3-13.6	1.6	2.5	1.1-5.0	1.2	4.2	1.5-9.3	0.4	1.3	0.2-4.9
	2006-2010	9.2	13.4	9.7-18.0	6.4	19.1	13.0-27.3	2.8	8.6	4.6-14.8	2.0	2.8	1.3-5.3	1.2	3.6	1.3-8.3	0.8	2.1	0.5-5.9
	2011-2015	9.6	12.7	9.3-17.1	6.6	17.9	12.2-25.7	3.0	8.2	4.4-14.0	2.4	3.1	1.6-5.6	1.6	4.6	1.9-9.5	0.8	1.9	0.5-5.7
	2016-2020	8.2	10.2	7.3-14.2	6.0	15.4	10.3-22.6	2.2	5.3	2.6-10.1	2.0	2.5	1.2-4.8	1.2	2.9	1.0-7.1	0.8	1.9	0.5-5.8
Esophagus	1976-1980	1.6	2.5	1.1-5.1	0.8	3.8	1.0-9.7	0.8	2.0	0.5-5.6	1.4	2.5	1.0-5.3	1.2	5.6	2.0-12.2	0.2	0.5	0.0-3.6
	1981-1985	1.4	2.1	0.8-4.5	1.0	3.9	1.2-9.5	0.4	1.0	0.1-4.2	1.8	2.9	1.3-5.5	1.2	4.9	1.7-10.8	0.6	1.5	0.3-4.9
	1986-1990	2.8	4.4	2.4-7.5	2.4	9.1	4.5-16.2	0.4	0.9	0.1-4.0	2.8	4.5	2.4-7.8	2.2	8.2	4.0-14.9	0.6	1.4	0.3-4.7
	1991-1995	3.4	5.5	3.2-8.9	2.6	10.5	5.5-18.2	0.8	1.9	0.5-5.5	3.0	4.6	2.5-7.7	2.4	9.3	4.6-16.4	0.6	1.4	0.3-4.7
	1996-2000	3.8	5.7	3.4-9.0	2.4	8.3	4.3-14.5	1.4	3.3	1.3-7.4	3.2	4.6	2.6-7.6	1.4	4.8	1.9-10.0	1.8	4.4	2.0-8.8
	2001-2005	3.4	5.1	2.9-8.2	3.2	10.5	6.0-17.2	0.2	0.6	0.0-3.7	4.2	6.1	3.7-9.4	3.8	13.2	7.9-20.7	0.4	1.0	0.1-4.2
	2006-2010	5.6	8.0	5.3-11.7	4.2	12.6	7.7-19.5	1.4	3.8	1.5-8.2	3.0	4.2	2.4-7.2	1.6	4.5	1.9-9.3	1.4	3.7	1.4-8.1
	2011-2015	4.8	6.6	4.1-10.0	3.6	10.5	6.1-17.1	1.2	2.8	1.0-6.7	5.2	7.2	4.6-10.8	4.8	14.7	9.3-22.3	0.4	1.0	0.1-4.2
	2016-2020	7.4	9.7	6.7-13.8	6.6	18.1	12.2-26.0	0.8	2.5	0.5-7.0	5.6	7.2	4.7-10.7	5.2	13.9	9.0-20.9	0.4	1.4	0.1-5.7
Stomach	1976-1980	6.2	10.3	6.9-14.8	2.8	11.0	5.8-18.8	3.4	9.5	5.4-15.5	5.4	8.9	5.8-13.2	3.2	13.0	7.2-21.6	2.2	6.0	3.0-11.1
	1981-1985	4.6	7.2	4.5-11.0	2.6	8.9	4.6-15.5	2.0	5.9	2.8-11.2	2.8	4.5	2.4-7.7	1.2	4.3	1.5-9.7	1.6	4.6	2.0-9.4
	1986-1990	5.8	9.4	6.3-13.7	3.4	12.5	7.2-20.2	2.4	6.8	3.4-12.3	2.6	4.2	2.2-7.3	1.2	4.7	1.7-10.3	1.4	3.5	1.4-7.7
	1991-1995	5.0	7.6	4.9-11.4	3.8	15.4	9.0-24.3	1.2	2.9	1.1-6.8	3.4	5.0	2.9-8.1	2.4	9.4	4.7-16.6	1.0	2.3	0.8-5.9
	1996-2000	4.2	6.2	3.9-9.7	2.2	8.2	4.0-14.7	2.0	5.0	2.4-9.7	2.0	3.0	1.4-5.6	0.6	2.0	0.4-6.1	1.4	3.6	1.4-7.9
	2001-2005	3.8	5.4	3.2-8.5	2.8	10.2	5.5-17.2	1.0	2.5	0.8-6.3	1.2	1.9	0.7-4.2	0.6	2.0	0.4-6.1	0.6	1.8	0.4-5.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Fulton**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	3.6	5.2	3.1-8.4	2.6	8.3	4.4-14.4	1.0	2.6	0.8-6.7	0.8	1.1	0.3-3.0	0.4	1.3	0.2-4.9	0.4	0.9	0.1-3.9
	2011-2015	4.4	6.2	3.8-9.7	3.4	11.0	6.3-18.0	1.0	2.1	0.7-5.8	0.6	0.8	0.2-2.6	0.2	0.7	0.0-4.0	0.4	0.9	0.1-4.1
	2016-2020	6.0	7.6	5.1-11.2	4.2	11.1	6.8-17.5	1.8	4.9	2.1-9.9	0.6	0.7	0.1-2.5	0.4	0.8	0.1-4.0	0.2	0.5	0.0-3.8
Colorectal	1976-1980	39.4	65.7	56.6-75.7	14.8	63.4	49.3-80.2	24.6	68.2	56.4-81.8	18.6	30.5	24.5-37.5	7.6	31.9	22.2-44.2	11.0	30.3	22.7-39.9
	1981-1985	41.2	64.5	55.8-74.2	20.2	79.0	63.8-96.8	21.0	55.4	45.1-67.5	25.4	39.7	33.0-47.5	13.0	52.2	39.7-67.2	12.4	32.4	24.7-41.8
	1986-1990	39.2	61.8	53.3-71.3	17.8	72.0	57.3-89.2	21.4	55.8	45.4-67.9	18.6	29.3	23.5-36.0	7.8	31.8	22.2-44.0	10.8	27.7	20.6-36.6
	1991-1995	41.4	61.7	53.5-70.9	19.2	69.6	56.2-85.3	22.2	54.9	45.0-66.5	17.6	26.9	21.5-33.3	6.8	24.6	16.9-34.5	10.8	26.8	19.9-35.4
	1996-2000	42.6	62.9	54.7-72.0	22.0	79.9	65.6-96.5	20.6	50.3	40.9-61.5	18.8	27.2	22.0-33.5	8.8	32.1	23.2-43.2	10.0	23.9	17.6-32.0
	2001-2005	43.2	62.8	54.7-71.9	20.2	69.1	56.2-84.1	23.0	58.2	47.8-70.5	16.2	23.3	18.5-29.1	6.6	22.9	15.7-32.2	9.6	23.4	17.1-31.5
	2006-2010	37.6	53.3	45.8-61.7	18.2	59.3	47.6-73.1	19.4	48.3	38.8-59.7	15.8	21.4	16.9-26.9	7.6	24.4	17.2-33.8	8.2	18.6	13.2-25.9
	2011-2015	30.4	42.2	35.6-49.8	14.2	43.3	33.6-55.1	16.2	42.2	33.3-53.1	12.4	17.1	13.0-22.2	6.2	19.2	12.9-27.6	6.2	15.7	10.4-22.9
2016-2020	29.2	38.8	32.5-46.1	13.6	39.4	30.2-50.6	15.6	38.6	30.2-49.0	11.0	14.7	10.8-19.5	5.8	16.9	11.1-25.0	5.2	12.2	7.7-18.9	
Colon excluding rectum	1976-1980	28.0	46.6	39.0-55.2	9.8	42.1	30.8-56.2	18.2	50.3	40.2-62.2	15.2	24.7	19.4-31.2	6.0	24.2	16.1-35.0	9.2	25.4	18.4-34.3
	1981-1985	32.0	50.0	42.4-58.6	14.6	56.0	43.3-71.1	17.4	46.2	36.8-57.4	22.0	34.4	28.2-41.6	11.2	44.5	33.1-58.5	10.8	28.2	21.1-37.1
	1986-1990	30.8	48.6	41.1-57.1	14.0	57.4	44.3-73.1	16.8	43.8	34.7-54.8	16.8	26.5	21.1-33.0	6.8	27.7	18.8-39.2	10.0	25.8	19.0-34.4
	1991-1995	30.2	44.9	38.0-52.8	13.4	49.3	38.1-62.8	16.8	40.4	32.2-50.4	15.2	23.2	18.2-29.2	5.6	20.2	13.3-29.4	9.6	23.7	17.3-31.9
	1996-2000	30.2	44.1	37.3-51.9	13.6	50.4	39.0-64.1	16.6	40.1	31.8-50.1	16.0	23.1	18.3-28.9	7.4	27.3	19.1-37.8	8.6	20.5	14.7-28.1
	2001-2005	31.0	44.4	37.7-52.1	12.6	43.8	33.6-56.2	18.4	45.4	36.4-56.2	14.4	20.6	16.1-26.1	5.8	20.2	13.5-29.1	8.6	20.8	14.9-28.4
	2006-2010	27.0	38.2	31.9-45.5	13.4	44.5	34.3-56.8	13.6	33.0	25.3-42.5	13.4	18.0	13.9-23.1	6.8	21.8	15.1-30.7	6.6	14.5	9.8-21.1
	2011-2015	21.6	30.3	24.7-36.8	9.8	30.8	22.6-41.2	11.8	30.6	23.1-40.0	10.0	13.6	10.0-18.2	5.2	16.4	10.6-24.4	4.8	11.5	7.2-17.7
2016-2020	19.8	26.5	21.4-32.6	7.6	22.7	15.8-31.7	12.2	29.9	22.6-39.2	8.6	11.8	8.4-16.4	4.2	13.1	7.9-20.6	4.4	10.5	6.3-16.9	
Rectum & rectosigmoid	1976-1980	11.4	19.1	14.4-24.9	5.0	21.3	13.5-31.7	6.4	17.9	12.2-25.5	3.4	5.8	3.3-9.3	1.6	7.6	3.1-15.1	1.8	5.0	2.3-9.7
	1981-1985	9.2	14.5	10.6-19.5	5.6	23.1	15.1-33.7	3.6	9.2	5.4-14.9	3.4	5.3	3.1-8.7	1.8	7.7	3.4-14.8	1.6	4.2	1.8-8.7
	1986-1990	8.4	13.2	9.5-18.0	3.8	14.6	8.6-23.0	4.6	11.9	7.4-18.3	1.8	2.7	1.2-5.3	1.0	4.1	1.2-9.8	0.8	1.9	0.5-5.4
	1991-1995	11.2	16.8	12.7-22.0	5.8	20.3	13.5-29.5	5.4	14.4	9.4-21.5	2.4	3.7	1.9-6.6	1.2	4.4	1.6-9.7	1.2	3.1	1.1-7.2
	1996-2000	12.4	18.7	14.3-24.1	8.4	29.5	21.2-40.0	4.0	10.3	6.2-16.3	2.8	4.1	2.2-7.0	1.4	4.8	1.9-10.0	1.4	3.4	1.4-7.5
	2001-2005	12.2	18.4	14.0-23.7	7.6	25.3	17.9-34.8	4.6	12.9	8.1-19.6	1.8	2.7	1.2-5.3	0.8	2.7	0.7-7.0	1.0	2.6	0.8-6.6
	2006-2010	10.6	15.1	11.2-19.9	4.8	14.8	9.4-22.3	5.8	15.4	10.1-22.6	2.4	3.4	1.7-6.1	0.8	2.6	0.7-6.9	1.6	4.1	1.7-8.6
	2011-2015	8.8	11.9	8.6-16.2	4.4	12.5	7.7-19.3	4.4	11.7	7.2-18.2	2.4	3.5	1.7-6.3	1.0	2.7	0.9-6.9	1.4	4.2	1.5-9.2
2016-2020	9.4	12.3	8.9-16.7	6.0	16.7	11.0-24.5	3.4	8.7	5.0-14.6	2.4	2.8	1.4-5.2	1.6	3.9	1.6-8.3	0.8	1.7	0.5-5.3	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Fulton**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	1.4	2.4	0.9-5.0	0.8	2.9	0.7-7.8	0.6	1.7	0.4-5.4	2.2	3.5	1.7-6.4	1.6	5.6	2.3-11.4	0.6	1.7	0.3-5.3
	1981-1985	1.2	1.9	0.7-4.3	1.2	5.1	1.8-11.2	0.0	0.0	0.0-2.6	1.6	2.5	1.1-5.1	1.4	5.6	2.2-11.5	0.2	0.5	0.0-3.4
	1986-1990	1.4	2.4	0.9-5.1	1.2	5.1	1.7-11.3	0.2	0.7	0.0-4.0	1.8	3.2	1.4-6.2	1.8	8.2	3.6-15.8	0.0	0.0	0.0-2.6
	1991-1995	2.2	3.3	1.7-6.1	0.8	2.8	0.7-7.2	1.4	3.3	1.3-7.2	1.6	2.3	1.0-4.6	1.2	3.9	1.4-8.8	0.4	0.8	0.1-3.7
	1996-2000	1.8	2.7	1.2-5.2	1.0	3.5	1.1-8.2	0.8	1.8	0.5-5.3	2.4	3.6	1.9-6.4	1.0	3.4	1.1-8.2	1.4	3.5	1.3-7.7
	2001-2005	3.4	5.1	3.0-8.3	3.2	10.7	6.1-17.4	0.2	0.5	0.0-3.4	1.8	2.7	1.2-5.2	1.2	4.1	1.5-9.1	0.6	1.5	0.3-4.9
	2006-2010	3.6	4.7	2.8-7.5	3.0	8.6	4.8-14.5	0.6	1.2	0.2-4.3	3.6	4.7	2.8-7.6	1.8	5.2	2.4-10.3	1.8	3.6	1.6-7.6
	2011-2015	5.0	6.4	4.1-9.7	3.8	10.6	6.3-17.1	1.2	2.6	0.9-6.3	3.4	4.2	2.4-7.0	1.8	4.9	2.2-9.8	1.6	3.3	1.4-7.3
	2016-2020	5.6	6.8	4.5-10.2	3.6	9.1	5.3-15.0	2.0	4.7	2.1-9.5	3.2	3.9	2.2-6.6	1.8	5.1	2.3-10.2	1.4	2.9	1.1-6.8
Pancreas	1976-1980	4.8	7.1	4.5-10.8	2.8	9.5	5.1-16.5	2.0	5.3	2.5-10.1	5.4	8.4	5.5-12.4	3.4	12.8	7.2-20.9	2.0	5.2	2.5-9.9
	1981-1985	6.8	10.4	7.2-14.8	4.0	14.3	8.5-22.7	2.8	7.4	4.0-12.8	8.2	12.4	8.8-17.0	5.0	18.7	11.7-28.2	3.2	8.3	4.7-13.9
	1986-1990	6.8	10.8	7.4-15.3	3.4	12.8	7.4-20.7	3.4	9.4	5.3-15.6	7.4	12.0	8.4-16.7	3.8	14.9	8.9-23.5	3.6	9.9	5.7-16.2
	1991-1995	9.0	13.5	9.8-18.2	4.6	16.5	10.3-24.9	4.4	10.4	6.5-16.2	9.4	14.3	10.5-19.2	5.0	18.4	11.8-27.4	4.4	10.0	6.3-15.6
	1996-2000	8.4	12.8	9.2-17.5	4.2	14.8	9.1-22.7	4.2	10.7	6.5-16.8	8.2	12.2	8.7-16.6	3.8	13.4	8.1-21.1	4.4	10.7	6.6-16.7
	2001-2005	7.2	10.6	7.4-14.8	3.2	10.4	5.9-17.1	4.0	10.6	6.4-16.8	7.4	10.8	7.6-15.0	3.6	11.9	7.0-19.0	3.8	9.8	5.9-15.7
	2006-2010	10.6	14.9	11.1-19.6	5.0	15.3	9.8-22.8	5.6	14.2	9.3-21.1	10.2	14.3	10.6-19.0	4.2	13.1	8.1-20.3	6.0	15.2	10.1-22.2
	2011-2015	13.8	19.1	14.8-24.4	6.2	18.3	12.3-26.4	7.6	19.5	13.6-27.4	10.4	13.8	10.3-18.3	5.4	16.1	10.6-23.9	5.0	11.9	7.6-18.1
	2016-2020	11.2	13.9	10.4-18.4	6.2	15.5	10.4-22.6	5.0	12.6	7.9-19.3	10.2	13.0	9.7-17.4	5.4	15.0	9.8-22.3	4.8	11.7	7.4-18.0
Larynx	1976-1980	3.4	5.5	3.2-8.9	3.4	12.2	7.0-19.9	0.0	0.0	0.0-2.7	2.2	3.5	1.7-6.4	1.8	7.4	3.3-14.2	0.4	0.9	0.1-4.0
	1981-1985	4.2	7.1	4.3-11.1	3.4	12.8	7.2-20.8	0.8	2.4	0.6-6.6	0.8	1.1	0.3-3.0	0.8	2.5	0.7-7.0	0.0	0.0	0.0-2.6
	1986-1990	3.8	5.7	3.4-9.1	3.0	10.3	5.7-17.3	0.8	1.9	0.5-5.4	0.6	1.0	0.2-3.1	0.4	1.5	0.2-5.5	0.2	0.7	0.0-4.0
	1991-1995	4.2	7.2	4.4-11.0	3.4	13.5	7.8-21.7	0.8	2.4	0.6-6.4	1.2	1.9	0.7-4.2	0.8	2.8	0.8-7.4	0.4	1.0	0.1-4.0
	1996-2000	2.8	4.5	2.5-7.7	1.8	6.6	3.0-12.7	1.0	3.0	1.0-7.2	1.2	1.9	0.7-4.2	0.8	3.1	0.8-8.0	0.4	1.0	0.1-4.1
	2001-2005	3.8	5.8	3.5-9.2	3.0	9.5	5.3-15.9	0.8	2.2	0.6-6.1	2.2	3.3	1.6-6.0	1.6	5.2	2.2-10.5	0.6	1.6	0.3-5.1
	2006-2010	2.8	4.0	2.2-6.9	2.0	6.2	2.9-11.6	0.8	2.0	0.5-5.6	0.6	0.8	0.2-2.7	0.4	1.2	0.1-4.7	0.2	0.6	0.0-3.6
	2011-2015	3.4	4.6	2.6-7.6	3.0	8.4	4.6-14.3	0.4	1.0	0.1-4.4	1.0	1.2	0.4-3.1	1.0	2.5	0.8-6.4	0.0	0.0	0.0-2.8
	2016-2020	2.8	3.6	2.0-6.4	2.6	7.2	3.8-12.8	0.2	0.5	0.0-3.6	0.4	0.5	0.1-2.1	0.2	0.5	0.0-3.7	0.2	0.5	0.0-3.6
Lung and bronchus	1976-1980	37.0	59.7	51.2-69.2	28.0	108.9	90.9-129.4	9.0	25.2	18.2-34.1	28.4	45.7	38.3-54.1	21.8	87.9	71.5-106.9	6.6	17.0	11.6-24.2
	1981-1985	42.6	65.2	56.6-74.9	30.8	110.5	93.1-130.1	11.8	32.6	24.5-42.6	35.8	55.0	47.1-64.0	26.0	95.7	79.4-114.5	9.8	27.9	20.2-37.5
	1986-1990	42.6	66.5	57.7-76.3	28.0	103.9	87.0-123.3	14.6	41.0	31.8-52.1	35.0	56.1	47.9-65.3	23.6	91.2	75.0-109.8	11.4	31.7	23.7-41.5
	1991-1995	50.2	77.6	68.2-88.0	31.6	112.9	95.8-132.3	18.6	53.5	42.9-66.1	42.6	64.4	55.9-73.8	26.6	96.0	80.1-114.0	16.0	42.7	33.6-53.6
	1996-2000	55.4	84.5	74.8-95.2	33.0	118.2	100.7-137.9	22.4	61.8	50.7-74.8	39.4	60.2	52.0-69.3	24.2	87.5	72.4-104.7	15.2	41.8	32.7-52.7
	2001-2005	59.4	88.2	78.4-98.9	30.0	102.3	86.5-120.1	29.4	78.5	66.1-92.7	40.8	59.5	51.5-68.3	22.4	77.1	63.4-92.9	18.4	48.0	38.5-59.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Fulton**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	53.0	76.6	67.5-86.6	28.0	90.9	76.3-107.7	25.0	67.6	56.1-80.9	36.6	53.0	45.5-61.4	18.8	60.1	48.4-73.8	17.8	49.5	39.6-61.4
	2011-2015	64.4	86.4	77.1-96.7	31.2	92.6	78.4-108.9	33.2	81.7	69.5-95.7	36.2	49.1	42.1-57.0	18.2	55.7	44.6-68.8	18.0	44.5	35.7-55.3
	2016-2020	64.4	80.4	71.7-90.1	30.8	82.4	69.6-97.2	33.6	79.9	67.9-93.6	36.0	45.6	39.0-53.1	17.6	48.3	38.5-60.1	18.4	45.1	36.0-56.0
Melanoma of the skin	1976-1980	1.8	4.0	1.8-7.6	1.2	5.7	2.0-12.4	0.6	2.6	0.5-7.5	1.2	2.6	0.9-5.5	1.0	5.2	1.6-11.9	0.2	0.8	0.0-4.2
	1981-1985	4.8	9.2	5.8-13.9	2.8	10.9	5.8-18.6	2.0	8.2	3.8-15.2	1.2	2.2	0.8-4.9	0.8	2.7	0.7-7.4	0.4	1.8	0.2-6.0
	1986-1990	3.4	6.2	3.6-10.1	1.8	7.3	3.2-14.2	1.6	6.1	2.6-12.1	2.0	4.2	2.0-7.8	1.0	4.8	1.5-11.0	1.0	3.6	1.1-8.6
	1991-1995	4.2	7.0	4.3-10.7	3.2	12.3	7.0-20.2	1.0	2.9	0.9-7.0	1.4	2.4	0.9-5.0	0.8	3.5	0.9-8.9	0.6	1.8	0.3-5.5
	1996-2000	7.6	12.8	9.0-17.6	4.0	14.2	8.6-22.0	3.6	12.3	7.2-19.6	2.0	2.9	1.4-5.5	1.8	6.3	2.9-12.1	0.2	0.7	0.0-3.8
	2001-2005	7.6	12.1	8.5-16.8	4.2	14.9	9.2-22.8	3.4	10.4	6.0-17.0	1.0	1.5	0.5-3.7	0.8	2.6	0.7-6.9	0.2	0.6	0.0-3.7
	2006-2010	10.2	15.2	11.2-20.1	5.2	16.3	10.6-24.2	5.0	14.1	8.9-21.4	2.4	3.6	1.8-6.4	1.6	5.3	2.3-10.7	0.8	2.2	0.6-6.0
	2011-2015	13.4	20.6	15.8-26.5	8.6	27.4	19.7-37.3	4.8	16.1	10.0-24.5	1.2	1.4	0.5-3.3	0.8	2.5	0.7-6.8	0.4	0.7	0.1-3.6
2016-2020	14.6	20.5	15.9-26.2	8.8	26.1	18.8-35.5	5.8	18.2	11.8-26.9	1.2	1.6	0.6-3.7	1.2	3.8	1.4-8.6	0.0	0.0	0.0-3.0	
Female breast	1976-1980							27.6	83.1	69.1-99.1							14.0	40.4	31.3-51.5
	1981-1985							34.6	109.1	92.5-127.8							13.0	38.7	29.4-50.0
	1986-1990							33.4	101.2	85.7-118.7							11.6	34.1	25.5-44.8
	1991-1995							35.8	107.3	91.6-125.0							13.2	37.7	28.9-48.6
	1996-2000							52.8	156.6	137.9-177.2							9.2	24.2	17.5-32.9
	2001-2005							41.4	121.7	105.4-139.9							11.4	30.7	23.1-40.2
	2006-2010							44.6	124.9	108.6-143.2							8.4	21.7	15.4-29.9
	2011-2015							50.2	137.2	120.0-156.2							7.2	18.4	12.7-26.0
2016-2020							48.8	122.6	106.9-140.2							8.8	20.9	14.8-28.9	
Cervix uteri	1976-1980							4.0	13.3	8.0-20.8							2.8	9.0	4.7-15.5
	1981-1985							3.6	11.9	6.9-19.1							1.4	4.0	1.5-8.5
	1986-1990							3.0	11.2	6.2-18.7							1.8	6.5	2.9-12.4
	1991-1995							3.6	11.9	6.9-19.0							1.2	4.3	1.6-9.4
	1996-2000							3.6	11.9	7.0-19.0							0.4	1.3	0.2-5.0
	2001-2005							3.2	11.4	6.4-18.7							0.8	2.6	0.7-6.9
	2006-2010							2.2	6.5	3.1-12.0							0.8	2.9	0.7-7.7
	2011-2015							1.6	5.4	2.3-11.0							0.6	1.9	0.4-6.0
	2016-2020							3.2	11.1	6.1-18.6							0.4	1.0	0.1-4.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Fulton**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							8.8	25.1	18.0-34.3							2.2	6.3	3.1-11.5
	1981-1985							10.4	31.5	23.1-42.0							1.6	4.0	1.7-8.3
	1986-1990							10.6	30.0	22.2-39.6							2.2	5.6	2.8-10.5
	1991-1995							9.6	28.1	20.5-37.8							2.2	5.5	2.6-10.3
	1996-2000							11.6	34.4	26.0-44.9							1.0	2.6	0.8-6.6
	2001-2005							12.0	34.5	26.2-44.7							2.4	5.9	3.0-10.8
	2006-2010							10.8	29.5	22.0-38.8							1.4	3.9	1.6-8.5
	2011-2015							13.4	35.0	26.8-45.0							1.8	4.5	2.0-9.1
	2016-2020							13.0	31.7	24.2-41.2							1.2	3.3	1.2-7.8
Ovary	1976-1980							6.0	17.7	11.8-25.6							3.6	10.9	6.3-17.6
	1981-1985							4.0	12.5	7.4-19.8							4.0	11.3	6.7-17.8
	1986-1990							4.2	12.2	7.4-19.1							3.6	10.2	5.9-16.5
	1991-1995							7.8	23.2	16.3-32.2							6.6	18.0	12.1-25.7
	1996-2000							6.6	18.6	12.6-26.5							5.2	13.7	8.8-20.5
	2001-2005							4.2	11.9	7.3-18.6							0.8	1.9	0.5-5.4
	2006-2010							4.6	12.3	7.7-18.8							3.2	8.8	5.0-14.6
	2011-2015							5.0	12.4	7.8-19.0							3.8	8.7	5.1-14.2
	2016-2020							4.6	11.7	7.3-18.3							2.0	4.2	2.0-8.5
Prostate	1976-1980				15.6	68.4	53.4-86.0							5.6	26.7	17.4-39.0			
	1981-1985				22.6	88.0	71.8-106.7							7.4	34.3	23.7-47.8			
	1986-1990				32.2	127.9	108.1-150.2							6.8	25.9	17.8-36.6			
	1991-1995				43.6	159.2	138.2-182.4							7.8	31.5	22.1-43.4			
	1996-2000				48.6	171.8	150.7-195.0							8.2	32.1	22.9-43.6			
	2001-2005				48.4	162.3	142.4-184.3							7.4	27.0	19.0-37.3			
	2006-2010				53.8	166.2	146.6-187.7							6.0	21.3	14.4-30.5			
	2011-2015				40.2	112.9	97.4-130.3							5.0	16.4	10.6-24.5			
	2016-2020				46.8	118.3	103.2-135.1							5.2	16.6	10.8-24.6			
Testis	1976-1980				0.4	1.8	0.2-6.4							0.0	0.0	0.0-3.1			
	1981-1985				1.6	6.0	2.5-11.9							0.2	0.7	0.0-4.3			
	1986-1990				0.8	3.2	0.8-8.2							0.0	0.0	0.0-3.0			
	1991-1995				1.4	5.4	2.1-11.1							0.0	0.0	0.0-2.9			
	1996-2000				1.2	4.3	1.6-9.4							0.0	0.0	0.0-2.8			
	2001-2005				1.6	6.2	2.7-12.2							0.0	0.0	0.0-2.8			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Fulton**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				1.6	6.2	2.7-12.3							0.0	0.0	0.0-2.8			
	2011-2015				1.8	7.3	3.3-13.9							0.2	0.5	0.0-3.6			
	2016-2020				1.6	6.9	3.0-13.5							0.0	0.0	0.0-2.9			
Urinary bladder (incl. in situ)	1976-1980	16.6	25.9	20.5-32.2	13.8	52.5	40.4-67.1	2.8	7.3	3.9-12.6	5.4	8.8	5.7-13.0	4.0	16.9	10.0-26.6	1.4	4.0	1.5-8.8
	1981-1985	12.2	19.2	14.6-24.9	9.2	35.2	25.4-47.5	3.0	8.1	4.5-13.8	4.0	6.4	3.9-10.1	3.0	12.9	7.1-21.4	1.0	2.5	0.8-6.3
	1986-1990	14.0	21.6	16.8-27.5	9.8	35.7	26.2-47.7	4.2	11.7	7.1-18.3	3.6	5.5	3.3-8.9	2.8	12.0	6.4-20.3	0.8	2.0	0.5-5.5
	1991-1995	15.0	22.9	18.0-28.9	11.6	42.3	32.0-55.0	3.4	8.8	5.0-14.5	3.8	5.9	3.5-9.4	2.4	9.3	4.7-16.3	1.4	3.2	1.3-7.1
	1996-2000	24.0	36.5	30.2-43.8	17.4	63.3	50.6-78.2	6.6	18.2	12.4-26.0	2.8	3.9	2.1-6.7	2.0	7.1	3.4-13.2	0.8	1.8	0.5-5.3
	2001-2005	23.0	33.9	28.0-40.8	15.4	52.3	41.2-65.5	7.6	20.8	14.6-29.0	4.0	5.6	3.4-8.7	2.6	9.1	4.8-15.6	1.4	3.1	1.2-7.0
	2006-2010	20.8	30.0	24.4-36.5	16.2	52.2	41.4-65.2	4.6	12.0	7.5-18.5	3.6	5.5	3.2-8.8	3.0	10.1	5.6-16.8	0.6	1.8	0.3-5.7
	2011-2015	26.2	36.6	30.5-43.7	19.4	62.1	50.1-76.2	6.8	16.6	11.3-23.7	3.4	4.4	2.5-7.3	2.6	8.7	4.6-15.1	0.8	1.6	0.4-5.1
2016-2020	25.8	32.7	27.2-39.2	19.2	54.6	44.0-67.3	6.6	16.2	10.9-23.4	4.0	4.7	2.9-7.6	3.4	9.9	5.7-16.3	0.6	1.3	0.2-4.7	
Kidney and renal pelvis	1976-1980	3.6	6.2	3.6-10.0	2.6	10.1	5.2-17.6	1.0	3.7	1.1-8.8	1.8	2.9	1.3-5.7	1.4	4.9	1.9-10.4	0.4	1.2	0.1-4.7
	1981-1985	4.8	8.8	5.6-13.3	3.0	14.6	8.0-24.1	1.8	4.6	2.1-9.1	3.4	6.1	3.5-9.9	2.2	9.5	4.6-17.1	1.2	3.5	1.3-8.0
	1986-1990	5.6	8.6	5.7-12.6	4.0	14.4	8.8-22.5	1.6	4.6	1.9-9.6	1.6	2.5	1.1-5.1	1.0	3.7	1.2-8.8	0.6	1.6	0.3-5.2
	1991-1995	8.2	12.7	9.1-17.4	3.6	12.4	7.3-19.8	4.6	12.4	7.8-19.1	3.6	5.4	3.2-8.7	1.4	5.0	2.0-10.5	2.2	5.7	2.8-10.7
	1996-2000	7.2	11.4	7.9-15.8	5.2	18.7	12.2-27.6	2.0	5.9	2.8-11.1	4.2	6.5	4.0-10.0	3.4	11.8	6.8-18.9	0.8	2.2	0.6-6.1
	2001-2005	10.4	15.6	11.6-20.5	6.2	20.7	14.0-29.5	4.2	11.6	7.1-18.2	3.8	5.4	3.2-8.5	2.6	9.1	4.8-15.6	1.2	2.9	1.0-6.9
	2006-2010	14.2	21.2	16.5-27.0	8.6	26.6	19.1-36.1	5.6	17.1	11.1-25.1	2.4	3.3	1.7-5.9	1.6	5.3	2.3-10.6	0.8	2.3	0.6-6.2
	2011-2015	13.6	20.0	15.4-25.6	8.2	24.5	17.4-33.8	5.4	15.3	9.9-22.8	2.8	3.6	1.9-6.4	1.6	5.0	2.0-10.3	1.2	2.4	0.9-6.0
2016-2020	14.4	19.0	14.7-24.3	8.8	24.5	17.6-33.5	5.6	14.2	9.2-21.3	3.0	3.8	2.1-6.5	1.6	4.2	1.7-8.8	1.4	3.5	1.4-7.8	
Brain and other central nervous system	1976-1980	2.6	3.9	2.1-6.9	2.0	7.1	3.3-13.5	0.6	1.6	0.3-5.2	2.6	4.5	2.3-7.9	2.4	8.8	4.4-15.7	0.2	1.3	0.0-5.7
	1981-1985	2.2	3.6	1.7-6.7	0.6	2.2	0.4-6.8	1.6	5.1	2.1-10.5	2.0	3.3	1.5-6.3	0.6	2.3	0.4-6.9	1.4	4.5	1.7-9.7
	1986-1990	4.2	7.3	4.5-11.4	2.2	9.0	4.4-16.2	2.0	5.9	2.8-11.1	3.6	6.2	3.6-9.9	1.4	6.1	2.4-12.7	2.2	5.8	2.9-10.7
	1991-1995	4.0	6.0	3.6-9.3	2.2	8.0	3.9-14.6	1.8	4.9	2.2-9.6	3.2	5.0	2.9-8.3	1.8	7.2	3.3-13.8	1.4	3.7	1.5-7.9
	1996-2000	2.8	4.3	2.4-7.4	2.6	9.2	4.9-15.9	0.2	0.5	0.0-3.4	2.2	3.4	1.7-6.1	1.6	5.8	2.5-11.6	0.6	1.8	0.4-5.6
	2001-2005	4.8	8.1	5.2-12.2	2.8	9.6	5.2-16.2	2.0	6.6	3.1-12.4	1.8	2.6	1.2-5.2	1.4	4.6	1.8-9.7	0.4	0.8	0.1-3.9
	2006-2010	4.8	7.3	4.6-11.1	3.0	10.3	5.7-17.2	1.8	4.5	2.0-9.1	2.8	3.9	2.1-6.8	1.4	4.3	1.7-9.2	1.4	3.6	1.4-8.0
	2011-2015	4.0	6.0	3.6-9.5	1.4	4.1	1.6-8.9	2.6	7.7	3.9-13.7	2.0	2.7	1.2-5.2	1.0	3.0	1.0-7.4	1.0	2.7	0.8-6.9
2016-2020	5.0	7.8	4.9-11.9	3.0	9.9	5.3-16.8	2.0	5.8	2.6-11.4	2.8	3.2	1.7-5.7	1.0	2.3	0.7-6.2	1.8	4.1	1.8-8.5	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Fulton**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	0.6	0.9	0.2-2.9	0.2	0.5	0.0-3.9	0.4	1.1	0.1-4.4	0.0	0.0	0.0-1.4	0.0	0.0	0.0-3.1	0.0	0.0	0.0-2.7
	1981-1985	2.0	3.2	1.5-6.0	0.6	2.1	0.4-6.6	1.4	4.4	1.7-9.2	0.2	0.3	0.0-2.0	0.2	1.0	0.0-5.2	0.0	0.0	0.0-2.6
	1986-1990	1.2	2.4	0.9-5.2	0.2	1.0	0.0-5.2	1.0	3.6	1.1-8.6	0.4	0.6	0.1-2.3	0.0	0.0	0.0-3.0	0.4	1.0	0.1-4.1
	1991-1995	1.2	2.0	0.7-4.5	0.6	2.1	0.4-6.3	0.6	2.1	0.4-6.1	0.2	0.4	0.0-2.1	0.2	0.8	0.0-4.5	0.0	0.0	0.0-2.6
	1996-2000	4.0	7.2	4.4-11.1	1.0	3.5	1.1-8.3	3.0	10.6	5.9-17.6	0.4	0.5	0.1-2.2	0.2	0.7	0.0-4.1	0.2	0.4	0.0-3.2
	2001-2005	4.8	8.3	5.2-12.4	0.8	2.9	0.8-7.6	4.0	13.3	8.0-20.8	0.2	0.3	0.0-1.7	0.0	0.0	0.0-2.8	0.2	0.4	0.0-3.2
	2006-2010	9.4	16.8	12.2-22.4	2.0	6.6	3.1-12.4	7.4	27.2	19.0-37.7	0.0	0.0	0.0-1.4	0.0	0.0	0.0-2.8	0.0	0.0	0.0-2.7
	2011-2015	9.2	16.4	11.8-22.1	2.6	8.5	4.4-14.9	6.6	24.1	16.2-34.4	0.0	0.0	0.0-1.4	0.0	0.0	0.0-2.9	0.0	0.0	0.0-2.8
	2016-2020	7.2	11.4	7.8-16.1	1.8	5.1	2.2-10.3	5.4	18.1	11.5-27.0	0.2	0.2	0.0-1.8	0.0	0.0	0.0-2.9	0.2	0.5	0.0-3.6
Hodgkin lymphoma	1976-1980	1.8	3.3	1.5-6.4	1.0	3.2	1.0-7.9	0.8	3.3	0.8-8.4	0.8	1.6	0.4-4.1	0.4	1.3	0.1-5.2	0.4	1.9	0.2-6.5
	1981-1985	2.2	3.7	1.8-6.8	0.8	2.9	0.8-7.7	1.4	4.7	1.8-10.1	0.6	0.9	0.2-2.8	0.4	1.2	0.1-5.0	0.2	0.6	0.0-3.7
	1986-1990	1.8	3.3	1.5-6.3	1.2	5.5	1.9-12.0	0.6	1.9	0.4-5.7	0.8	1.3	0.3-3.4	0.6	2.2	0.4-6.7	0.2	0.6	0.0-3.7
	1991-1995	1.8	3.4	1.5-6.4	0.6	2.3	0.5-6.7	1.2	4.1	1.5-9.1	0.0	0.0	0.0-1.3	0.0	0.0	0.0-2.9	0.0	0.0	0.0-2.6
	1996-2000	2.8	5.3	2.9-9.0	1.0	4.0	1.3-9.2	1.8	6.7	3.0-12.7	0.2	0.3	0.0-1.9	0.0	0.0	0.0-2.8	0.2	0.6	0.0-3.6
	2001-2005	3.0	5.3	2.9-8.8	1.8	6.3	2.8-12.0	1.2	4.3	1.5-9.5	0.4	0.7	0.1-2.7	0.0	0.0	0.0-2.8	0.4	1.4	0.1-5.2
	2006-2010	2.8	5.8	3.1-9.6	1.4	5.7	2.3-11.7	1.4	5.7	2.3-11.7	0.0	0.0	0.0-1.4	0.0	0.0	0.0-2.8	0.0	0.0	0.0-2.7
	2011-2015	2.6	4.2	2.2-7.4	1.6	5.0	2.1-10.3	1.0	3.3	1.0-8.2	0.2	0.3	0.0-1.9	0.0	0.0	0.0-2.9	0.2	0.5	0.0-3.5
	2016-2020	1.6	2.9	1.2-5.9	0.8	2.7	0.7-7.3	0.8	3.1	0.8-8.1	0.0	0.0	0.0-1.5	0.0	0.0	0.0-2.9	0.0	0.0	0.0-3.0
Non-Hodgkin lymphomas	1976-1980	7.2	11.4	7.9-16.0	4.4	15.3	9.4-23.7	2.8	7.9	4.3-13.5	3.4	5.6	3.2-9.1	2.4	8.6	4.3-15.4	1.0	2.9	0.9-7.1
	1981-1985	8.4	13.9	9.9-19.1	5.6	20.9	13.7-30.6	2.8	8.8	4.7-15.2	5.2	8.5	5.5-12.6	3.6	13.9	8.1-22.2	1.6	4.6	1.9-9.4
	1986-1990	6.8	11.3	7.7-16.0	3.2	12.2	6.9-20.1	3.6	10.0	5.7-16.3	5.6	9.7	6.3-14.1	3.6	14.0	8.2-22.4	2.0	5.8	2.7-11.2
	1991-1995	9.6	15.6	11.4-20.8	4.2	15.2	9.4-23.4	5.4	15.8	10.2-23.4	4.4	6.7	4.2-10.3	1.6	6.4	2.7-12.6	2.8	7.1	3.8-12.3
	1996-2000	10.2	16.0	11.9-21.2	4.4	15.9	10.0-24.2	5.8	16.3	10.8-23.7	6.2	9.6	6.5-13.7	2.2	7.7	3.8-13.8	4.0	10.8	6.5-17.1
	2001-2005	12.6	19.4	14.9-24.9	5.2	17.4	11.4-25.7	7.4	21.1	14.7-29.5	3.6	5.6	3.3-9.0	1.6	5.5	2.4-10.9	2.0	5.6	2.6-10.8
	2006-2010	15.2	22.8	17.9-28.8	9.2	30.5	22.2-40.9	6.0	16.3	10.8-23.9	3.4	4.7	2.7-7.7	1.2	3.9	1.4-8.7	2.2	5.4	2.6-10.2
	2011-2015	12.8	17.9	13.7-23.1	7.0	22.6	15.6-31.8	5.8	13.3	8.8-19.7	5.4	7.8	5.0-11.5	3.2	10.5	5.9-17.4	2.2	5.2	2.5-10.1
	2016-2020	15.4	21.5	16.7-27.3	9.4	28.5	20.6-38.7	6.0	14.5	9.6-21.5	2.4	3.3	1.6-6.0	1.4	4.4	1.7-9.6	1.0	2.4	0.8-6.2
Myeloma	1976-1980	2.4	4.2	2.1-7.5	1.0	5.3	1.6-12.2	1.4	3.7	1.5-8.1	1.4	2.2	0.9-4.7	0.6	2.4	0.4-7.1	0.8	2.2	0.6-6.0
	1981-1985	1.4	2.1	0.8-4.5	0.8	3.0	0.8-8.0	0.6	1.5	0.3-4.9	1.4	2.3	0.9-4.9	1.0	4.0	1.3-9.5	0.4	1.4	0.1-5.2
	1986-1990	2.6	4.2	2.2-7.3	1.2	5.5	1.8-12.1	1.4	3.9	1.5-8.5	3.0	4.7	2.6-7.9	1.8	7.7	3.3-14.9	1.2	2.8	1.0-6.6
	1991-1995	3.4	6.1	3.5-9.8	1.8	7.1	3.2-13.6	1.6	4.8	2.0-9.9	2.4	4.0	2.1-7.1	1.2	4.4	1.6-9.8	1.2	3.7	1.3-8.3
	1996-2000	2.0	3.0	1.5-5.7	1.6	5.5	2.4-11.0	0.4	1.2	0.1-4.7	1.4	2.1	0.9-4.5	0.8	2.8	0.8-7.3	0.6	1.4	0.3-4.8
	2001-2005	3.8	5.7	3.4-9.1	2.0	6.8	3.2-12.6	1.8	4.9	2.2-9.6	1.4	2.0	0.8-4.3	1.2	4.3	1.6-9.4	0.2	0.6	0.0-3.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Fulton**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	3.0	4.1	2.3-6.9	1.2	3.5	1.3-8.1	1.8	4.8	2.2-9.5	1.4	1.8	0.7-3.9	1.0	3.0	0.9-7.3	0.4	1.0	0.1-4.1
	2011-2015	6.8	8.9	6.1-12.7	3.8	10.9	6.5-17.5	3.0	6.6	3.6-11.6	1.8	2.4	1.1-4.7	0.4	1.2	0.1-4.8	1.4	2.9	1.1-6.8
	2016-2020	6.8	9.2	6.3-13.1	5.0	14.5	9.3-21.9	1.8	4.7	2.0-9.7	2.0	2.5	1.2-4.9	1.4	4.1	1.6-8.9	0.6	1.2	0.2-4.6
Leukemias	1976-1980	8.0	14.3	10.2-19.6	4.0	17.8	10.6-27.8	4.0	11.8	7.2-18.5	5.6	10.2	6.7-14.8	2.0	9.8	4.5-18.0	3.6	10.7	6.3-17.2
	1981-1985	6.4	10.6	7.1-15.1	3.8	16.0	9.3-25.4	2.6	7.3	3.7-13.1	5.4	9.3	6.0-13.6	3.2	13.8	7.7-22.7	2.2	6.3	3.0-11.8
	1986-1990	7.2	11.9	8.3-16.6	4.8	18.1	11.5-27.2	2.4	6.8	3.4-12.4	5.2	8.1	5.3-12.0	3.4	13.7	7.8-22.3	1.8	4.5	2.0-8.9
	1991-1995	5.0	8.5	5.5-12.7	2.8	10.8	5.9-18.2	2.2	5.8	2.8-10.8	3.0	4.8	2.7-8.0	1.6	5.9	2.5-11.8	1.4	3.6	1.4-7.8
	1996-2000	9.4	14.3	10.5-19.2	5.2	18.9	12.3-27.8	4.2	12.0	7.3-18.8	4.0	6.0	3.6-9.3	1.8	7.2	3.2-13.6	2.2	5.7	2.8-10.7
	2001-2005	8.4	12.6	9.0-17.1	5.2	18.0	11.7-26.5	3.2	8.5	4.7-14.4	4.6	6.5	4.1-9.9	3.0	10.2	5.7-17.0	1.6	3.6	1.5-7.7
	2006-2010	10.8	16.1	12.1-21.2	5.2	16.9	11.0-25.0	5.6	16.0	10.5-23.5	4.8	7.0	4.5-10.6	3.0	10.0	5.5-16.7	1.8	4.9	2.2-9.8
	2011-2015	13.6	19.2	14.8-24.7	7.0	21.8	15.0-30.8	6.6	18.1	12.1-26.1	4.4	5.6	3.5-8.7	2.6	7.9	4.1-13.9	1.8	3.7	1.6-7.8
2016-2020	13.8	19.3	14.8-24.7	6.6	20.0	13.6-28.6	7.2	18.8	12.8-26.9	3.6	5.3	3.1-8.6	2.0	6.9	3.3-12.9	1.6	4.1	1.7-8.8	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Genesee**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	215.4	393.7	370.2-418.4	111.8	476.1	436.1-518.7	103.6	343.3	313.8-374.8	111.2	204.7	187.8-222.7	58.2	256.1	226.6-288.2	53.0	170.4	150.3-192.5
	1981-1985	232.0	405.2	381.9-429.6	111.8	460.0	421.4-500.9	120.2	374.7	344.7-406.6	118.2	206.6	190.0-224.1	59.2	250.2	221.6-281.4	59.0	179.6	159.4-201.8
	1986-1990	267.0	448.5	424.5-473.6	136.8	543.5	502.2-587.1	130.2	399.0	368.2-431.7	119.6	200.4	184.5-217.3	60.8	250.9	222.7-281.7	58.8	170.5	151.3-191.6
	1991-1995	297.6	472.9	449.0-497.7	149.8	559.4	519.1-601.8	147.8	424.0	393.4-456.4	137.6	214.8	198.9-231.6	72.8	280.3	251.5-311.4	64.8	173.6	155.0-194.0
	1996-2000	334.6	510.5	486.2-535.6	178.6	625.6	584.8-668.5	156.0	434.7	404.2-466.9	135.8	202.2	187.3-218.1	75.2	271.0	243.9-300.3	60.6	157.7	140.2-176.9
	2001-2005	355.2	523.8	499.6-548.8	178.6	589.1	550.9-629.4	176.6	481.8	450.1-515.3	137.0	196.8	182.3-212.2	71.2	244.6	219.6-271.7	65.8	166.8	148.9-186.3
	2006-2010	363.2	499.9	476.9-523.8	194.0	588.6	551.7-627.3	169.2	433.1	403.6-464.3	133.8	178.1	164.7-192.4	69.6	218.4	195.8-243.0	64.2	152.9	136.2-171.2
	2011-2015	391.4	511.9	488.9-535.8	194.2	541.7	507.4-577.8	197.2	493.7	462.2-526.9	136.2	172.0	159.1-185.8	70.4	202.3	181.4-225.1	65.8	151.3	134.8-169.4
	2016-2020	397.2	496.1	473.6-519.5	208.4	543.3	509.8-578.7	188.8	461.5	430.8-494.0	138.6	164.8	152.5-178.0	79.4	209.3	188.8-231.7	59.2	132.3	117.0-149.2
Oral cavity and pharynx	1976-1980	3.4	5.7	3.3-9.2	2.2	8.2	4.0-15.0	1.2	3.6	1.3-8.0	2.0	3.7	1.8-6.9	1.0	4.7	1.5-10.7	1.0	2.9	0.9-7.0
	1981-1985	6.0	10.1	6.8-14.5	4.0	16.4	9.8-25.5	2.0	5.6	2.6-10.5	2.0	3.3	1.6-6.1	1.0	4.2	1.3-9.7	1.0	2.8	0.9-6.7
	1986-1990	6.0	9.8	6.5-14.0	4.6	17.0	10.6-25.6	1.4	4.8	1.9-10.0	2.2	3.6	1.8-6.5	1.2	5.0	1.8-10.8	1.0	2.6	0.9-6.4
	1991-1995	5.4	8.6	5.6-12.6	3.8	13.3	8.0-20.8	1.6	4.6	2.0-9.2	1.4	2.1	0.8-4.3	1.2	4.1	1.5-9.0	0.2	0.5	0.0-3.2
	1996-2000	6.4	10.0	6.8-14.2	3.8	12.7	7.6-19.9	2.6	7.2	3.8-12.6	1.6	2.6	1.1-5.1	1.6	5.3	2.3-10.5	0.0	0.0	0.0-2.4
	2001-2005	7.6	11.4	8.1-15.7	5.2	16.5	10.8-24.4	2.4	6.7	3.4-12.0	2.0	2.9	1.4-5.4	1.4	4.3	1.7-9.0	0.6	1.6	0.3-5.0
	2006-2010	8.2	10.8	7.8-14.8	5.8	16.1	10.8-23.4	2.4	5.6	2.9-10.1	1.0	1.3	0.4-3.2	0.8	2.3	0.6-6.1	0.2	0.3	0.0-2.9
	2011-2015	10.2	13.2	9.8-17.6	8.2	22.1	15.7-30.4	2.0	5.4	2.5-10.2	1.8	2.3	1.0-4.5	1.6	4.5	1.9-9.3	0.2	0.4	0.0-3.2
	2016-2020	11.6	14.6	11.0-19.2	8.2	21.5	15.2-29.7	3.4	8.5	4.8-14.3	2.8	3.1	1.7-5.4	2.4	5.9	3.0-10.7	0.4	0.8	0.1-3.8
Esophagus	1976-1980	1.4	2.4	1.0-5.1	1.2	5.2	1.8-11.4	0.2	0.5	0.0-3.4	1.8	3.2	1.4-6.1	1.2	5.5	1.9-11.9	0.6	1.7	0.3-5.3
	1981-1985	1.4	2.5	1.0-5.2	1.0	4.1	1.3-9.7	0.4	1.1	0.1-4.3	1.0	1.8	0.6-4.1	0.6	2.6	0.5-7.5	0.4	1.2	0.1-4.4
	1986-1990	2.0	3.2	1.5-6.0	1.4	4.9	1.9-10.2	0.6	1.6	0.3-5.0	1.6	2.7	1.1-5.3	1.0	3.4	1.1-8.1	0.6	1.6	0.3-5.0
	1991-1995	2.6	3.9	2.1-6.8	1.6	5.5	2.4-10.9	1.0	2.6	0.9-6.4	1.8	2.9	1.3-5.5	1.4	4.9	1.9-10.1	0.4	1.1	0.1-4.2
	1996-2000	1.8	2.7	1.2-5.2	1.6	5.2	2.3-10.4	0.2	0.4	0.0-3.0	2.2	3.3	1.6-6.0	2.0	6.5	3.1-12.1	0.2	0.5	0.0-3.2
	2001-2005	4.2	5.9	3.6-9.1	3.0	10.1	5.6-16.7	1.2	3.1	1.1-7.0	4.0	5.6	3.4-8.8	3.2	11.0	6.2-18.0	0.8	1.9	0.5-5.3
	2006-2010	4.0	5.3	3.2-8.3	3.6	10.2	6.0-16.4	0.4	0.9	0.1-3.8	4.4	5.7	3.6-8.8	3.8	10.9	6.5-17.2	0.6	1.4	0.3-4.5
	2011-2015	2.4	3.2	1.6-5.7	1.6	4.4	1.9-9.0	0.8	1.9	0.5-5.4	2.2	3.0	1.5-5.6	1.6	4.6	1.9-9.4	0.6	1.6	0.3-5.0
	2016-2020	5.0	5.9	3.8-9.0	4.2	10.5	6.4-16.5	0.8	1.7	0.5-5.2	5.0	5.8	3.7-8.8	4.0	9.6	5.8-15.2	1.0	2.1	0.7-5.7
Stomach	1976-1980	6.0	11.0	7.4-15.8	2.8	11.9	6.4-20.0	3.2	10.4	5.9-17.1	4.2	7.5	4.6-11.5	2.0	7.9	3.7-14.7	2.2	6.9	3.4-12.4
	1981-1985	3.0	5.3	2.9-8.7	1.4	5.6	2.2-11.6	1.6	4.9	2.0-9.9	2.2	4.1	2.0-7.3	0.8	3.8	1.0-9.4	1.4	4.4	1.7-9.2
	1986-1990	5.2	8.9	5.8-13.1	4.0	15.2	9.2-23.6	1.2	3.2	1.2-7.2	3.0	5.2	2.9-8.7	2.2	8.6	4.2-15.5	0.8	2.5	0.6-6.5
	1991-1995	5.2	7.8	5.1-11.5	3.0	10.3	5.7-17.1	2.2	5.8	2.9-10.5	3.4	5.2	3.0-8.5	2.4	8.8	4.5-15.4	1.0	2.5	0.8-6.1
	1996-2000	4.0	5.9	3.6-9.2	3.0	10.5	5.8-17.5	1.0	2.7	0.9-6.5	2.0	2.9	1.4-5.4	1.2	4.0	1.5-8.9	0.8	2.1	0.6-5.7
	2001-2005	5.6	8.0	5.3-11.6	4.0	13.2	8.0-20.5	1.6	3.8	1.6-7.9	3.0	4.1	2.3-6.9	2.2	7.4	3.7-13.3	0.8	1.5	0.4-4.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Genesee**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	5.4	7.3	4.8-10.7	4.6	14.6	9.2-22.0	0.8	2.0	0.5-5.6	1.8	2.2	1.0-4.4	1.6	5.5	2.3-10.8	0.2	0.5	0.0-3.3
	2011-2015	5.0	6.5	4.1-9.7	3.4	9.4	5.5-15.4	1.6	4.1	1.6-8.6	2.4	2.8	1.4-5.1	2.2	6.1	3.0-11.3	0.2	0.3	0.0-2.9
	2016-2020	5.2	6.8	4.3-10.3	2.8	7.3	3.9-12.8	2.4	6.6	3.3-12.1	1.6	1.9	0.8-4.0	1.2	3.0	1.1-7.0	0.4	0.9	0.1-4.1
Colorectal	1976-1980	38.0	70.6	60.8-81.5	20.8	90.4	73.4-110.0	17.2	56.2	44.8-69.5	17.8	34.1	27.3-42.0	9.0	42.7	30.8-57.4	8.8	28.7	20.7-38.6
	1981-1985	37.6	65.6	56.5-75.8	17.4	71.8	57.0-89.1	20.2	60.8	49.4-74.1	17.8	31.7	25.4-39.1	9.0	41.1	29.6-55.3	8.8	25.7	18.7-34.7
	1986-1990	33.8	57.7	49.3-67.2	17.2	72.7	57.8-90.2	16.6	47.6	37.8-59.2	17.2	28.7	22.9-35.4	9.4	38.6	28.2-51.6	7.8	22.1	15.6-30.4
	1991-1995	38.0	59.8	51.6-69.0	17.8	68.6	54.8-84.7	20.2	52.2	42.4-63.7	15.6	24.5	19.3-30.6	7.2	28.1	19.4-39.1	8.4	21.1	15.1-28.7
	1996-2000	45.6	68.3	59.7-77.9	23.6	83.7	69.1-100.4	22.0	56.4	46.2-68.3	19.8	29.2	23.7-35.6	10.2	36.4	27.0-47.9	9.6	23.1	16.9-30.9
	2001-2005	35.6	51.1	43.9-59.3	16.0	52.8	41.8-65.8	19.6	49.6	40.1-60.9	13.6	19.4	15.1-24.7	6.8	22.8	15.7-32.0	6.8	16.0	10.9-22.7
	2006-2010	35.0	46.9	40.2-54.6	17.0	52.3	41.7-65.0	18.0	44.0	35.1-54.6	11.8	15.1	11.4-19.6	5.8	19.0	12.7-27.4	6.0	13.3	8.9-19.5
	2011-2015	29.4	38.8	32.6-45.8	16.6	48.3	38.2-60.2	12.8	30.7	23.5-39.7	12.4	15.7	11.9-20.4	6.6	19.2	13.2-27.3	5.8	13.6	8.9-20.2
2016-2020	31.8	40.4	34.1-47.6	17.8	46.6	37.1-57.9	14.0	35.8	27.3-46.2	10.0	12.4	9.1-16.6	6.2	17.2	11.6-24.9	3.8	8.9	5.2-14.6	
Colon excluding rectum	1976-1980	26.6	49.6	41.4-58.9	14.2	62.7	48.6-79.5	12.4	40.3	30.8-51.8	13.6	25.8	20.0-32.8	6.0	29.1	19.3-41.7	7.6	24.1	17.0-33.2
	1981-1985	25.2	44.9	37.3-53.5	10.8	44.3	32.9-58.1	14.4	43.7	34.1-55.2	15.2	27.1	21.3-34.0	7.4	33.0	23.0-45.8	7.8	22.9	16.3-31.5
	1986-1990	22.0	37.9	31.1-45.7	10.4	44.7	33.0-59.0	11.6	33.7	25.5-43.8	14.8	24.8	19.4-31.2	7.8	32.2	22.7-44.2	7.0	19.9	13.8-27.9
	1991-1995	28.2	44.5	37.4-52.5	13.0	50.3	38.6-64.3	15.2	39.4	30.9-49.6	13.2	20.8	16.1-26.5	6.6	26.2	17.8-37.0	6.6	16.7	11.4-23.7
	1996-2000	31.4	47.0	39.9-55.0	17.2	60.7	48.4-75.1	14.2	36.3	28.2-46.1	17.0	25.2	20.1-31.2	8.2	28.9	20.7-39.3	8.8	21.1	15.3-28.7
	2001-2005	25.2	35.9	29.8-42.8	9.8	33.0	24.3-43.7	15.4	38.2	30.0-48.2	11.2	15.9	12.0-20.7	5.4	18.1	11.9-26.5	5.8	13.5	8.9-19.7
	2006-2010	23.0	30.4	25.0-36.6	9.6	29.7	21.8-39.6	13.4	31.4	24.2-40.4	8.8	11.1	8.0-15.0	4.0	13.2	8.0-20.5	4.8	10.2	6.4-15.7
	2011-2015	20.8	27.5	22.4-33.6	10.4	31.5	23.3-41.6	10.4	24.5	18.2-32.6	9.8	12.6	9.2-16.9	5.2	15.4	10.0-22.8	4.6	10.5	6.4-16.4
2016-2020	19.0	23.3	18.7-28.9	10.6	28.0	20.8-37.2	8.4	19.9	14.1-27.6	7.0	8.7	5.9-12.3	4.4	12.3	7.6-19.1	2.6	6.0	3.0-11.0	
Rectum & rectosigmoid	1976-1980	11.4	21.0	15.8-27.3	6.6	27.6	18.8-39.0	4.8	15.9	10.1-23.8	4.2	8.3	5.1-12.6	3.0	13.6	7.5-22.5	1.2	4.6	1.6-9.9
	1981-1985	12.4	20.8	15.9-26.7	6.6	27.5	18.5-39.1	5.8	17.2	11.4-24.9	2.6	4.6	2.4-7.9	1.6	8.1	3.3-15.8	1.0	2.8	0.9-6.8
	1986-1990	11.8	19.9	15.1-25.7	6.8	28.0	19.3-39.2	5.0	13.9	8.9-20.7	2.4	3.9	2.0-6.8	1.6	6.5	2.7-12.8	0.8	2.1	0.6-5.7
	1991-1995	9.8	15.4	11.3-20.4	4.8	18.3	11.6-27.4	5.0	12.8	8.3-19.2	2.4	3.7	1.9-6.4	0.6	1.9	0.4-5.7	1.8	4.4	2.0-8.6
	1996-2000	14.2	21.3	16.7-27.0	6.4	23.0	15.6-32.6	7.8	20.1	14.2-27.7	2.8	4.0	2.2-6.8	2.0	7.5	3.5-13.7	0.8	1.9	0.5-5.3
	2001-2005	10.4	15.3	11.4-20.1	6.2	19.8	13.4-28.2	4.2	11.4	7.0-17.7	2.4	3.5	1.8-6.2	1.4	4.6	1.8-9.7	1.0	2.5	0.8-6.2
	2006-2010	12.0	16.5	12.6-21.4	7.4	22.7	15.9-31.4	4.6	12.5	7.8-19.2	3.0	4.0	2.2-6.7	1.8	5.8	2.6-11.1	1.2	3.1	1.1-7.1
	2011-2015	8.6	11.2	8.1-15.3	6.2	16.8	11.3-24.3	2.4	6.2	3.1-11.2	2.6	3.1	1.6-5.6	1.4	3.8	1.5-8.3	1.2	3.1	1.1-7.2
2016-2020	12.8	17.0	12.9-22.2	7.2	18.6	12.8-26.2	5.6	15.9	10.1-23.8	3.0	3.7	2.0-6.4	1.8	4.9	2.2-9.7	1.2	2.9	1.0-7.0	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Genesee**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	1.4	2.3	0.9-4.9	0.6	2.2	0.4-6.7	0.8	2.3	0.6-6.2	2.2	3.7	1.8-6.7	1.2	5.1	1.8-11.1	1.0	2.9	0.9-6.9
	1981-1985	1.8	3.2	1.4-6.1	1.4	5.9	2.3-12.2	0.4	1.6	0.2-5.6	1.4	2.6	1.0-5.3	0.8	3.6	1.0-9.1	0.6	2.1	0.4-6.3
	1986-1990	1.4	2.4	1.0-5.0	0.4	1.8	0.2-6.1	1.0	2.7	0.9-6.4	1.4	2.5	1.0-5.1	0.6	3.0	0.5-8.4	0.8	2.1	0.6-5.7
	1991-1995	2.6	4.0	2.1-7.0	1.4	5.1	2.0-10.6	1.2	3.0	1.1-6.9	2.4	3.7	1.9-6.5	1.0	3.8	1.2-8.8	1.4	3.5	1.4-7.4
	1996-2000	2.2	3.5	1.7-6.2	1.4	4.6	1.8-9.5	0.8	2.3	0.6-6.1	3.4	5.3	3.1-8.4	1.4	4.7	1.9-9.7	2.0	5.8	2.7-10.7
	2001-2005	2.2	3.2	1.6-5.8	1.4	4.2	1.7-8.9	0.8	2.0	0.5-5.4	3.0	4.3	2.4-7.1	1.6	5.4	2.3-10.8	1.4	3.3	1.3-7.2
	2006-2010	3.0	4.0	2.2-6.8	1.6	4.3	1.8-8.8	1.4	3.7	1.4-7.9	3.4	4.4	2.6-7.2	2.0	6.0	2.8-11.2	1.4	3.3	1.3-7.2
	2011-2015	5.4	6.9	4.5-10.2	3.6	9.0	5.3-14.6	1.8	4.7	2.1-9.3	3.2	4.1	2.3-6.8	2.0	5.6	2.6-10.6	1.2	2.6	1.0-6.3
	2016-2020	4.6	5.1	3.2-7.9	3.4	8.1	4.7-13.4	1.2	2.3	0.8-5.8	4.4	5.1	3.1-7.9	3.2	8.0	4.5-13.5	1.2	2.5	0.9-6.2
Pancreas	1976-1980	6.2	12.2	8.3-17.4	3.6	17.0	9.8-27.1	2.6	8.6	4.6-14.7	5.8	10.9	7.2-15.7	3.4	14.7	8.3-23.9	2.4	7.7	4.0-13.6
	1981-1985	6.8	11.7	8.0-16.4	2.8	11.1	6.0-18.8	4.0	11.7	7.1-18.2	6.8	11.5	7.9-16.2	3.0	11.7	6.4-19.4	3.8	11.0	6.6-17.3
	1986-1990	5.2	8.6	5.6-12.7	1.8	6.9	3.1-13.1	3.4	9.2	5.3-14.9	3.8	6.4	3.8-10.0	1.2	4.7	1.7-10.4	2.6	7.0	3.7-12.1
	1991-1995	8.4	13.3	9.6-18.1	3.4	12.1	7.0-19.4	5.0	13.4	8.6-20.0	7.6	11.8	8.3-16.2	3.4	11.7	6.8-18.9	4.2	11.0	6.8-17.0
	1996-2000	9.4	14.1	10.4-18.8	5.8	19.9	13.3-28.7	3.6	9.2	5.4-14.8	10.0	14.8	11.0-19.6	5.2	18.1	11.8-26.5	4.8	12.3	7.8-18.5
	2001-2005	12.8	18.4	14.2-23.6	5.4	18.8	12.3-27.4	7.4	17.9	12.5-25.1	10.2	14.7	11.0-19.5	4.0	13.9	8.4-21.5	6.2	15.3	10.3-22.1
	2006-2010	9.0	12.3	8.9-16.6	4.0	12.3	7.5-19.3	5.0	11.5	7.3-17.4	7.4	10.3	7.2-14.4	3.4	10.8	6.2-17.4	4.0	9.1	5.5-14.6
	2011-2015	9.6	11.8	8.7-15.9	4.8	12.6	8.0-19.2	4.8	10.4	6.6-16.1	9.0	11.4	8.3-15.4	4.2	11.9	7.3-18.5	4.8	10.6	6.7-16.3
	2016-2020	10.4	12.9	9.5-17.2	4.8	12.3	7.8-18.8	5.6	13.5	8.6-20.4	11.2	12.8	9.6-16.8	5.8	14.7	9.7-21.5	5.4	11.0	7.1-16.7
Larynx	1976-1980	3.0	5.7	3.1-9.4	2.4	10.1	5.1-17.8	0.6	2.6	0.5-7.3	1.4	2.9	1.1-5.9	1.0	5.3	1.6-12.0	0.4	1.6	0.2-5.7
	1981-1985	2.4	4.2	2.2-7.4	2.2	8.3	4.1-15.0	0.2	0.8	0.0-4.2	1.4	2.3	0.9-4.8	1.4	5.8	2.2-12.1	0.0	0.0	0.0-2.5
	1986-1990	2.2	3.6	1.8-6.4	2.0	7.2	3.4-13.5	0.2	0.7	0.0-3.8	0.4	0.7	0.1-2.5	0.2	0.7	0.0-4.2	0.2	0.7	0.0-3.8
	1991-1995	3.0	5.1	2.9-8.5	2.0	8.2	3.9-15.0	1.0	2.5	0.8-6.2	1.4	2.1	0.9-4.4	1.0	3.4	1.1-8.0	0.4	1.0	0.1-4.0
	1996-2000	2.8	4.6	2.5-7.8	2.6	8.8	4.7-15.1	0.2	0.6	0.0-3.4	1.2	1.7	0.6-3.8	1.0	3.3	1.1-7.8	0.2	0.4	0.0-3.0
	2001-2005	2.2	3.2	1.6-5.8	1.8	5.8	2.6-11.1	0.4	1.1	0.1-4.3	0.4	0.5	0.1-2.1	0.2	0.7	0.0-3.8	0.2	0.5	0.0-3.3
	2006-2010	3.0	4.0	2.3-6.8	2.2	6.5	3.2-11.8	0.8	2.0	0.5-5.5	1.4	1.9	0.7-4.0	1.0	3.1	1.0-7.4	0.4	1.0	0.1-4.1
	2011-2015	4.8	5.9	3.8-9.0	3.6	9.1	5.3-14.7	1.2	3.1	1.1-7.2	1.0	1.1	0.4-2.9	1.0	2.6	0.8-6.6	0.0	0.0	0.0-2.6
	2016-2020	3.8	4.6	2.7-7.5	3.4	8.5	4.8-14.2	0.4	1.0	0.1-4.2	0.8	1.0	0.3-2.9	0.6	1.6	0.3-5.1	0.2	0.5	0.0-3.4
Lung and bronchus	1976-1980	31.8	55.4	47.0-64.9	22.6	91.5	74.9-110.6	9.2	29.6	21.6-39.8	24.4	42.8	35.4-51.2	17.4	72.3	57.5-89.6	7.0	21.7	15.1-30.4
	1981-1985	32.6	54.3	46.2-63.5	23.6	91.0	74.9-109.6	9.0	27.4	19.8-36.9	28.8	48.5	40.8-57.2	21.2	83.8	68.1-101.9	7.6	23.6	16.6-32.6
	1986-1990	42.0	68.4	59.4-78.5	27.2	103.1	86.0-122.4	14.8	42.7	33.4-53.8	31.2	50.7	43.0-59.4	20.0	76.9	62.1-94.0	11.2	32.1	24.2-42.0
	1991-1995	46.6	72.7	63.6-82.8	28.0	103.3	86.6-122.3	18.6	51.4	41.4-63.3	41.2	63.8	55.3-73.2	25.2	92.9	77.0-111.0	16.0	43.9	34.6-54.9
	1996-2000	53.2	80.6	71.2-91.0	34.2	119.1	101.8-138.6	19.0	52.3	42.2-64.2	39.4	58.6	50.6-67.4	25.6	91.3	76.0-108.8	13.8	37.2	28.8-47.3
	2001-2005	52.0	76.6	67.5-86.6	29.0	96.1	81.0-113.2	23.0	63.2	52.1-76.0	43.0	62.8	54.6-71.8	25.2	84.7	70.5-101.0	17.8	47.4	38.0-58.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Genesee**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	57.2	78.2	69.3-88.0	32.0	96.9	82.3-113.4	25.2	63.3	52.5-75.8	41.8	56.1	48.7-64.4	23.8	73.4	60.7-88.0	18.0	43.6	34.9-54.0
	2011-2015	60.2	76.6	68.1-86.1	27.4	75.7	63.4-89.9	32.8	78.9	67.0-92.5	44.6	56.3	49.0-64.4	23.2	65.8	54.2-79.3	21.4	50.1	40.9-61.0
	2016-2020	59.8	69.9	62.0-78.6	31.2	77.7	65.7-91.5	28.6	64.0	53.7-76.0	37.0	44.2	38.0-51.4	21.6	55.2	45.1-67.1	15.4	35.8	28.0-45.3
Melanoma of the skin	1976-1980	3.2	5.5	3.1-9.1	1.2	3.7	1.4-8.5	2.0	7.0	3.3-13.0	1.4	2.5	1.0-5.2	1.0	3.7	1.2-8.8	0.4	1.3	0.2-4.7
	1981-1985	2.0	3.3	1.6-6.2	0.6	2.0	0.4-6.2	1.4	4.1	1.6-8.7	0.6	1.2	0.2-3.4	0.0	0.0	0.0-2.9	0.6	2.2	0.4-6.4
	1986-1990	6.0	10.0	6.7-14.4	3.2	11.4	6.4-18.7	2.8	9.0	4.8-15.2	2.0	3.3	1.6-6.2	1.0	3.4	1.1-8.1	1.0	3.1	1.0-7.3
	1991-1995	4.8	7.8	5.0-11.6	3.0	10.3	5.7-17.1	1.8	5.9	2.7-11.1	1.0	1.6	0.5-3.8	1.0	3.7	1.2-8.6	0.0	0.0	0.0-2.3
	1996-2000	7.8	12.1	8.6-16.5	4.8	16.2	10.3-24.2	3.0	8.7	4.9-14.6	1.4	2.1	0.9-4.4	1.0	3.4	1.1-8.0	0.4	1.2	0.1-4.5
	2001-2005	9.8	15.0	11.1-19.8	5.2	17.0	11.0-24.9	4.6	13.5	8.5-20.4	1.4	2.0	0.8-4.3	0.8	2.7	0.7-7.1	0.6	1.6	0.3-5.0
	2006-2010	14.4	20.8	16.2-26.4	8.8	27.5	19.9-37.0	5.6	16.4	10.8-24.0	1.4	2.0	0.8-4.2	0.8	2.5	0.7-6.6	0.6	1.6	0.3-5.1
	2011-2015	19.8	25.9	20.9-31.8	13.6	37.2	28.7-47.6	6.2	16.5	11.0-23.8	2.6	3.4	1.8-6.0	2.2	5.8	2.8-10.9	0.4	1.0	0.1-4.1
2016-2020	23.2	29.8	24.4-36.3	13.2	35.3	26.9-45.5	10.0	25.7	18.6-34.8	2.0	2.1	1.0-4.2	1.8	4.0	1.8-8.1	0.2	0.3	0.0-3.1	
Female breast	1976-1980							25.8	89.4	74.3-106.6							10.4	33.8	25.2-44.4
	1981-1985							31.2	100.4	84.9-117.9							14.0	44.4	34.4-56.4
	1986-1990							39.2	123.3	106.2-142.4							12.6	38.3	29.2-49.3
	1991-1995							43.8	132.9	115.5-152.1							11.6	31.8	24.1-41.4
	1996-2000							47.2	135.4	118.5-154.1							10.2	27.6	20.5-36.6
	2001-2005							54.6	152.7	134.9-172.3							8.0	20.9	14.8-28.7
	2006-2010							47.4	123.6	108.1-140.9							7.8	19.3	13.6-26.8
	2011-2015							56.6	142.3	125.7-160.7							8.2	18.4	13.0-25.6
2016-2020							55.0	134.7	118.4-152.7							8.0	18.1	12.6-25.5	
Cervix uteri	1976-1980							3.4	11.5	6.6-18.7							1.2	4.1	1.5-9.0
	1981-1985							3.2	10.9	6.1-17.9							1.8	5.1	2.3-9.8
	1986-1990							3.6	13.0	7.6-20.7							0.6	1.6	0.3-5.0
	1991-1995							3.0	9.2	5.0-15.3							1.4	3.9	1.5-8.3
	1996-2000							3.8	12.1	7.2-18.9							1.0	2.6	0.8-6.4
	2001-2005							2.2	6.6	3.3-11.9							0.6	1.6	0.3-4.9
	2006-2010							1.6	5.4	2.3-10.8							0.6	1.6	0.3-5.1
	2011-2015							2.0	5.3	2.4-10.3							0.8	2.8	0.7-7.3
	2016-2020							1.0	3.7	1.1-9.0							0.4	0.9	0.1-4.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Genesee**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							10.0	32.5	24.0-43.2							2.0	6.4	3.1-11.9
	1981-1985							9.6	31.2	22.8-41.6							0.8	2.7	0.7-7.1
	1986-1990							9.2	28.1	20.4-37.8							2.2	5.8	2.9-10.6
	1991-1995							8.6	25.1	18.0-34.0							1.6	4.5	1.9-9.1
	1996-2000							9.4	27.5	20.1-36.8							0.8	1.9	0.5-5.3
	2001-2005							12.6	34.5	26.4-44.4							2.2	5.7	2.8-10.6
	2006-2010							11.6	29.4	22.2-38.4							3.8	9.3	5.6-15.0
	2011-2015							15.2	35.8	28.0-45.3							2.0	4.5	2.0-9.0
	2016-2020							13.0	30.5	23.1-39.7							2.6	5.6	3.0-10.1
Ovary	1976-1980							5.6	18.2	12.0-26.5							4.0	13.3	8.0-20.8
	1981-1985							8.0	25.6	18.1-35.1							5.2	15.9	10.3-23.5
	1986-1990							4.4	14.8	9.1-22.6							2.4	7.3	3.7-12.9
	1991-1995							4.8	14.9	9.4-22.3							2.6	7.0	3.7-12.1
	1996-2000							3.4	10.1	5.8-16.3							1.8	5.2	2.3-10.0
	2001-2005							6.2	17.7	11.9-25.3							3.2	8.3	4.7-13.8
	2006-2010							6.0	14.4	9.6-21.0							4.6	10.9	6.8-16.8
	2011-2015							7.2	18.9	12.9-26.7							3.4	8.2	4.8-13.6
	2016-2020							5.4	13.9	9.0-20.8							3.4	7.6	4.4-12.7
Prostate	1976-1980				20.6	93.5	75.8-114.0							5.8	28.9	19.2-41.5			
	1981-1985				24.2	106.9	88.2-128.2							5.0	23.9	15.1-35.5			
	1986-1990				32.6	135.3	114.7-158.5							7.6	34.4	24.1-47.4			
	1991-1995				37.8	139.2	119.6-161.1							9.4	40.8	29.7-54.6			
	1996-2000				50.2	174.3	153.2-197.5							8.6	34.5	24.7-46.6			
	2001-2005				57.6	187.9	166.8-211.1							6.8	24.8	17.1-34.8			
	2006-2010				53.8	159.7	141.0-180.3							6.0	20.2	13.6-29.0			
	2011-2015				43.6	116.4	101.3-133.4							4.4	13.3	8.3-20.4			
	2016-2020				52.6	126.9	111.7-143.8							9.0	25.6	18.6-34.5			
Testis	1976-1980				1.0	2.8	0.9-7.1							0.6	1.8	0.4-5.7			
	1981-1985				1.8	6.8	3.1-13.1							0.2	0.6	0.0-3.8			
	1986-1990				0.8	2.3	0.6-6.3							0.4	1.7	0.2-5.9			
	1991-1995				0.8	2.9	0.8-7.4							0.2	0.9	0.0-4.5			
	1996-2000				2.4	8.0	4.1-14.0							0.0	0.0	0.0-2.6			
	2001-2005				2.0	6.6	3.1-12.2							0.2	0.6	0.0-3.7			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Genesee**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				1.4	5.6	2.3-11.4							0.2	0.8	0.0-4.3			
	2011-2015				2.2	7.8	3.8-14.0							0.0	0.0	0.0-2.6			
	2016-2020				1.8	6.9	3.1-13.1							0.0	0.0	0.0-2.7			
Urinary bladder (incl. in situ)	1976-1980	11.6	20.7	15.6-26.8	8.6	36.3	25.9-49.2	3.0	9.2	5.1-15.3	4.0	7.4	4.5-11.5	3.0	13.9	7.6-22.9	1.0	3.1	1.0-7.4
	1981-1985	14.4	24.8	19.3-31.2	10.6	43.1	32.0-56.8	3.8	11.0	6.6-17.4	4.0	7.0	4.2-10.8	2.8	11.5	6.2-19.4	1.2	3.5	1.3-7.7
	1986-1990	17.8	29.3	23.5-36.1	12.8	51.2	39.1-65.8	5.0	14.0	9.0-20.8	2.4	4.1	2.1-7.2	1.8	8.5	3.8-16.1	0.6	1.6	0.3-5.0
	1991-1995	17.8	27.9	22.3-34.3	13.4	51.9	40.0-66.2	4.4	11.7	7.3-17.9	3.8	5.8	3.5-9.0	2.2	9.5	4.7-17.1	1.6	4.0	1.7-8.2
	1996-2000	16.2	24.3	19.3-30.3	11.8	42.1	31.9-54.4	4.4	12.0	7.5-18.4	2.8	4.1	2.3-7.0	2.0	7.2	3.4-13.3	0.8	2.0	0.5-5.4
	2001-2005	16.2	23.6	18.7-29.4	12.2	41.3	31.6-53.2	4.0	10.4	6.3-16.3	3.2	4.5	2.5-7.3	3.0	10.5	5.9-17.4	0.2	0.4	0.0-2.9
	2006-2010	17.6	23.2	18.6-28.7	14.8	44.9	35.2-56.6	2.8	6.3	3.4-11.0	5.2	6.6	4.3-9.8	3.4	10.9	6.3-17.6	1.8	4.1	1.8-8.3
	2011-2015	25.6	32.7	27.2-39.1	19.0	53.9	43.5-66.3	6.6	14.8	10.1-21.3	4.0	4.8	2.9-7.6	2.8	8.5	4.6-14.4	1.2	2.4	0.8-5.9
2016-2020	20.6	25.6	20.7-31.3	15.6	43.4	34.1-54.6	5.0	10.9	6.8-16.8	4.6	4.9	3.1-7.7	2.8	7.8	4.2-13.5	1.8	3.7	1.7-7.6	
Kidney and renal pelvis	1976-1980	6.6	12.2	8.3-17.2	3.8	15.0	8.9-23.6	2.8	9.1	4.9-15.3	3.0	5.2	2.9-8.7	1.0	3.6	1.2-8.7	2.0	6.1	2.9-11.4
	1981-1985	6.4	11.8	8.0-16.8	3.6	15.5	9.1-24.6	2.8	8.7	4.7-14.9	3.4	6.0	3.5-9.7	1.8	8.2	3.6-15.5	1.6	4.5	1.9-9.1
	1986-1990	5.6	10.1	6.7-14.6	3.8	15.6	9.3-24.5	1.8	6.5	2.9-12.3	2.8	5.0	2.7-8.3	1.4	6.0	2.4-12.2	1.4	4.5	1.7-9.4
	1991-1995	9.0	14.7	10.7-19.8	4.2	16.2	10.0-24.8	4.8	13.9	8.8-20.8	2.4	3.7	1.9-6.4	1.0	3.5	1.1-8.3	1.4	3.8	1.5-7.9
	1996-2000	8.6	13.4	9.7-18.1	4.8	16.6	10.6-24.8	3.8	10.7	6.4-17.0	2.2	3.4	1.7-6.1	1.4	5.2	2.0-10.7	0.8	2.2	0.6-5.9
	2001-2005	10.0	14.9	11.1-19.7	5.8	18.5	12.4-26.7	4.2	11.9	7.3-18.4	2.8	4.0	2.2-6.8	1.6	5.5	2.4-10.9	1.2	3.0	1.1-6.8
	2006-2010	11.6	15.8	11.9-20.5	7.2	21.0	14.6-29.3	4.4	11.2	6.9-17.2	1.8	2.3	1.1-4.6	1.6	4.9	2.1-9.9	0.2	0.5	0.0-3.3
	2011-2015	14.0	18.8	14.5-24.1	9.0	26.5	19.1-35.8	5.0	12.8	8.1-19.4	3.2	4.2	2.4-7.0	2.2	6.6	3.2-12.0	1.0	2.4	0.8-6.1
2016-2020	17.0	20.2	16.0-25.2	10.6	26.7	19.9-35.4	6.4	15.0	10.1-21.7	4.2	4.7	2.9-7.5	3.0	7.8	4.3-13.2	1.2	2.6	0.9-6.3	
Brain and other central nervous system	1976-1980	3.2	5.6	3.1-9.2	1.0	3.4	1.1-8.4	2.2	7.4	3.6-13.4	1.8	3.0	1.4-5.8	1.0	3.6	1.1-8.6	0.8	2.3	0.6-6.2
	1981-1985	4.6	8.3	5.2-12.6	3.2	13.4	7.5-21.8	1.4	3.7	1.5-7.9	1.6	2.8	1.2-5.7	1.0	3.9	1.2-9.2	0.6	1.9	0.4-5.6
	1986-1990	5.0	8.5	5.5-12.6	3.4	12.3	7.1-19.9	1.6	5.6	2.3-11.1	3.4	5.6	3.2-9.1	2.6	10.0	5.2-17.3	0.8	2.1	0.6-5.6
	1991-1995	6.0	9.8	6.6-14.0	3.4	12.8	7.3-20.5	2.6	7.8	4.1-13.5	2.6	4.3	2.3-7.4	2.0	6.9	3.3-12.9	0.6	1.9	0.4-5.5
	1996-2000	5.4	8.7	5.7-12.6	3.6	11.8	7.0-18.8	1.8	5.9	2.7-11.1	2.6	4.1	2.2-7.1	1.4	4.7	1.9-9.7	1.2	3.4	1.2-7.5
	2001-2005	5.0	7.9	5.1-11.7	2.0	6.6	3.1-12.2	3.0	9.0	5.0-15.1	3.8	5.9	3.5-9.2	1.8	5.7	2.6-10.9	2.0	5.9	2.8-11.0
	2006-2010	4.6	6.8	4.3-10.4	3.2	10.5	6.0-17.3	1.4	3.5	1.3-7.6	2.6	3.5	1.8-6.1	2.0	5.8	2.7-11.0	0.6	1.4	0.2-4.6
	2011-2015	5.8	8.5	5.6-12.4	3.6	11.0	6.3-17.7	2.2	6.3	3.0-11.7	3.4	4.4	2.5-7.2	2.2	6.4	3.1-11.8	1.2	2.7	1.0-6.4
2016-2020	5.0	6.9	4.3-10.6	2.6	7.4	3.8-13.1	2.4	6.8	3.3-12.5	3.8	5.0	2.9-8.2	2.2	6.4	3.0-12.2	1.6	3.4	1.4-7.4	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Genesee**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	1.6	3.3	1.4-6.5	0.8	3.7	1.0-9.3	0.8	2.8	0.8-7.3	0.6	1.0	0.2-3.0	0.4	1.3	0.2-5.0	0.2	0.6	0.0-3.7
	1981-1985	3.6	6.5	3.8-10.4	0.8	2.8	0.7-7.4	2.8	9.7	5.2-16.4	0.4	0.7	0.1-2.6	0.0	0.0	0.0-2.9	0.4	1.2	0.1-4.5
	1986-1990	3.2	6.2	3.5-10.0	0.4	1.6	0.2-5.7	2.8	10.4	5.6-17.6	0.0	0.0	0.0-1.3	0.0	0.0	0.0-2.8	0.0	0.0	0.0-2.4
	1991-1995	3.2	5.3	3.0-8.6	1.0	3.7	1.2-8.6	2.2	7.2	3.6-13.0	0.8	1.2	0.3-3.2	0.2	0.9	0.0-4.5	0.6	1.5	0.3-4.7
	1996-2000	3.2	5.2	3.0-8.4	0.4	1.3	0.2-4.9	2.8	9.0	4.9-15.1	0.2	0.3	0.0-1.7	0.0	0.0	0.0-2.6	0.2	0.5	0.0-3.2
	2001-2005	3.8	6.0	3.6-9.5	1.2	3.9	1.4-8.6	2.6	8.4	4.4-14.5	0.2	0.3	0.0-1.7	0.2	0.9	0.0-4.5	0.0	0.0	0.0-2.4
	2006-2010	6.0	9.0	6.0-12.9	1.6	4.6	2.0-9.4	4.4	13.3	8.2-20.4	0.8	1.2	0.3-3.1	0.4	1.2	0.1-4.6	0.4	1.1	0.1-4.4
	2011-2015	8.6	13.7	9.8-18.6	3.0	8.8	4.8-14.9	5.6	19.0	12.4-27.8	0.4	0.6	0.1-2.2	0.0	0.0	0.0-2.6	0.4	1.0	0.1-4.2
2016-2020	7.6	13.6	9.5-18.9	3.2	11.7	6.5-19.2	4.4	15.5	9.4-23.8	0.6	0.6	0.1-2.3	0.2	0.6	0.0-3.7	0.4	0.9	0.1-4.0	
Hodgkin lymphoma	1976-1980	1.8	3.0	1.3-5.8	1.4	5.2	2.0-10.9	0.4	1.1	0.1-4.3	0.6	0.9	0.2-2.8	0.2	0.6	0.0-4.0	0.4	1.1	0.1-4.4
	1981-1985	1.6	2.8	1.2-5.6	0.8	2.5	0.7-6.8	0.8	3.0	0.8-7.6	0.4	0.8	0.1-2.7	0.2	1.0	0.0-4.9	0.2	0.6	0.0-3.6
	1986-1990	2.0	3.4	1.6-6.3	0.6	1.9	0.4-5.9	1.4	4.9	1.9-10.0	0.2	0.4	0.0-2.0	0.2	0.7	0.0-4.1	0.0	0.0	0.0-2.4
	1991-1995	1.6	2.8	1.2-5.5	0.4	1.5	0.2-5.3	1.2	4.1	1.5-8.9	0.8	1.4	0.4-3.6	0.2	0.8	0.0-4.3	0.6	1.9	0.4-5.6
	1996-2000	0.6	1.2	0.2-3.4	0.2	0.9	0.0-4.4	0.4	1.6	0.2-5.5	0.0	0.0	0.0-1.2	0.0	0.0	0.0-2.6	0.0	0.0	0.0-2.4
	2001-2005	1.6	2.5	1.1-5.0	1.0	3.4	1.1-8.0	0.6	1.8	0.4-5.5	0.2	0.3	0.0-1.7	0.0	0.0	0.0-2.6	0.2	0.6	0.0-3.4
	2006-2010	2.4	4.1	2.1-7.2	1.6	5.4	2.3-10.8	0.8	2.9	0.8-7.4	0.0	0.0	0.0-1.3	0.0	0.0	0.0-2.6	0.0	0.0	0.0-2.5
	2011-2015	2.4	4.1	2.1-7.2	1.0	3.8	1.2-8.7	1.4	4.4	1.7-9.3	0.0	0.0	0.0-1.3	0.0	0.0	0.0-2.6	0.0	0.0	0.0-2.6
2016-2020	1.0	1.9	0.6-4.6	0.4	2.1	0.3-6.9	0.6	1.7	0.3-5.5	0.4	0.6	0.1-2.3	0.2	0.5	0.0-3.5	0.2	0.8	0.0-4.2	
Non-Hodgkin lymphomas	1976-1980	5.2	9.0	5.9-13.3	2.0	8.0	3.8-14.8	3.2	9.6	5.5-15.7	3.0	5.8	3.2-9.6	1.0	4.8	1.5-10.9	2.0	6.4	3.1-11.9
	1981-1985	6.0	10.9	7.3-15.6	2.4	10.5	5.3-18.5	3.6	11.0	6.5-17.6	4.6	8.1	5.1-12.3	1.8	6.8	3.0-13.0	2.8	8.5	4.6-14.5
	1986-1990	9.6	16.0	11.7-21.2	5.2	20.2	12.9-30.0	4.4	13.0	8.0-19.9	3.2	5.5	3.1-9.0	2.0	8.3	3.9-15.5	1.2	3.3	1.2-7.5
	1991-1995	10.2	16.3	12.1-21.4	5.2	19.2	12.3-28.5	5.0	15.0	9.6-22.4	5.6	8.9	5.9-12.9	3.4	14.3	8.1-23.0	2.2	6.2	3.0-11.2
	1996-2000	12.2	18.2	13.9-23.5	6.0	22.1	14.8-31.7	6.2	16.5	11.1-23.7	3.6	5.3	3.1-8.4	2.0	7.6	3.6-14.1	1.6	4.0	1.7-8.1
	2001-2005	15.8	23.7	18.7-29.6	7.2	23.7	16.6-33.0	8.6	23.2	16.7-31.5	4.8	7.1	4.5-10.6	2.2	7.7	3.8-13.9	2.6	7.1	3.8-12.4
	2006-2010	17.2	23.6	18.8-29.3	9.8	29.8	21.9-39.6	7.4	18.0	12.5-25.2	5.8	8.0	5.3-11.6	3.6	11.7	6.9-18.6	2.2	5.3	2.6-10.0
	2011-2015	14.6	18.8	14.6-23.9	7.4	21.5	15.0-29.9	7.2	17.1	11.8-24.2	3.8	4.9	2.9-7.9	2.4	7.4	3.8-13.2	1.4	3.5	1.3-7.7
2016-2020	15.4	19.3	15.1-24.5	7.6	19.7	13.8-27.5	7.8	19.3	13.3-27.1	3.2	3.7	2.1-6.3	1.6	4.6	2.0-9.4	1.6	3.0	1.3-6.7	
Myeloma	1976-1980	2.2	4.1	2.0-7.3	1.2	6.2	2.2-13.2	1.0	2.9	0.9-7.0	0.6	1.1	0.2-3.3	0.2	0.7	0.0-4.1	0.4	1.3	0.2-4.8
	1981-1985	2.0	3.5	1.7-6.5	1.0	4.4	1.4-10.1	1.0	3.0	1.0-7.2	1.6	2.8	1.2-5.5	0.6	2.4	0.5-7.0	1.0	2.9	0.9-7.0
	1986-1990	3.2	5.3	3.0-8.6	1.6	6.4	2.6-12.8	1.6	4.2	1.8-8.5	2.8	4.7	2.5-7.9	1.0	4.4	1.3-10.2	1.8	5.1	2.3-9.9
	1991-1995	4.8	7.4	4.7-11.0	2.8	10.4	5.5-17.7	2.0	5.3	2.5-10.0	2.0	3.1	1.5-5.7	1.2	4.7	1.6-10.4	0.8	2.0	0.5-5.3
	1996-2000	4.6	6.9	4.4-10.4	2.2	8.3	4.1-15.0	2.4	6.3	3.2-11.3	2.6	3.9	2.1-6.7	1.2	4.1	1.5-8.9	1.4	3.8	1.5-8.0
	2001-2005	2.6	3.7	2.0-6.4	1.4	4.7	1.9-9.8	1.2	3.2	1.1-7.2	1.8	2.5	1.1-4.8	0.8	2.7	0.7-6.9	1.0	2.3	0.7-5.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Genesee**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	3.2	4.3	2.4-7.1	1.6	5.3	2.3-10.6	1.6	3.5	1.5-7.5	1.0	1.3	0.4-3.2	0.8	2.6	0.7-6.7	0.2	0.6	0.0-3.4
	2011-2015	5.4	7.1	4.6-10.5	1.8	5.0	2.3-9.9	3.6	8.9	5.2-14.5	2.2	2.5	1.2-4.7	1.0	2.7	0.9-6.8	1.2	2.2	0.8-5.6
	2016-2020	5.6	6.8	4.4-10.0	3.0	8.4	4.6-14.3	2.6	5.8	3.1-10.4	3.0	3.6	2.0-6.1	1.4	4.0	1.6-8.5	1.6	3.7	1.6-7.8
Leukemias	1976-1980	8.2	15.7	11.2-21.3	5.8	26.6	17.6-38.4	2.4	7.8	4.0-13.7	5.4	10.3	6.7-15.0	3.8	17.3	10.3-27.2	1.6	5.3	2.2-10.5
	1981-1985	8.6	14.9	10.7-20.1	4.4	18.8	11.6-28.7	4.2	12.1	7.5-18.7	5.2	9.3	6.0-13.6	3.2	15.1	8.5-24.6	2.0	5.7	2.7-10.7
	1986-1990	9.4	15.8	11.6-21.1	5.4	21.6	14.0-31.7	4.0	12.1	7.3-18.9	4.6	7.8	4.9-11.7	2.2	9.9	4.8-17.7	2.4	7.0	3.6-12.5
	1991-1995	8.6	13.5	9.7-18.2	4.8	17.5	11.1-26.2	3.8	11.0	6.5-17.3	6.0	9.1	6.2-13.1	3.8	14.2	8.5-22.3	2.2	5.7	2.8-10.4
	1996-2000	12.4	19.0	14.5-24.3	7.0	25.8	17.8-36.0	5.4	13.9	9.1-20.6	4.4	6.6	4.1-10.0	2.2	8.2	4.0-14.7	2.2	5.3	2.6-9.9
	2001-2005	12.0	17.8	13.5-22.9	7.0	23.6	16.4-33.0	5.0	13.3	8.5-19.9	6.6	9.2	6.3-13.1	3.6	13.7	8.1-21.7	3.0	7.4	4.1-12.6
	2006-2010	12.4	17.3	13.2-22.4	7.0	21.7	15.0-30.3	5.4	14.0	9.0-20.8	5.4	7.1	4.6-10.4	3.0	9.7	5.4-16.1	2.4	5.0	2.5-9.2
	2011-2015	16.4	21.5	17.0-27.0	9.6	27.2	19.9-36.3	6.8	16.6	11.2-24.0	6.4	8.4	5.7-12.0	3.6	11.0	6.5-17.5	2.8	6.3	3.2-11.3
2016-2020	14.2	19.8	15.2-25.4	8.4	25.5	18.0-35.2	5.8	14.2	9.2-21.2	4.4	5.4	3.3-8.5	2.4	6.1	3.1-11.2	2.0	5.2	2.3-10.2	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Greene**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	189.8	404.3	378.2-431.9	100.0	502.2	457.1-550.4	89.8	346.2	313.4-381.5	100.8	208.8	190.5-228.4	54.2	261.4	230.0-295.9	46.6	174.0	151.6-199.0
	1981-1985	226.6	466.6	439.0-495.4	117.6	550.6	505.9-598.4	109.0	420.1	383.7-459.1	112.4	227.5	208.6-247.7	58.4	283.2	250.8-318.8	54.0	194.5	171.1-220.5
	1986-1990	238.6	471.0	444.1-499.1	125.4	561.9	518.0-608.6	113.2	413.2	378.5-450.3	111.4	221.2	202.8-240.8	59.2	269.1	238.7-302.4	52.2	191.6	168.0-217.6
	1991-1995	252.4	474.6	448.5-501.9	135.2	572.4	529.4-618.0	117.2	414.0	380.3-450.0	128.2	235.5	217.4-254.6	67.4	291.8	261.0-325.2	60.8	200.7	178.3-225.4
	1996-2000	293.8	528.1	501.3-556.0	161.4	640.1	596.2-686.5	132.4	450.9	416.6-487.5	136.2	240.5	222.7-259.4	75.8	310.0	279.1-343.3	60.4	189.5	168.4-212.7
	2001-2005	322.0	540.9	514.6-568.3	170.8	605.6	565.1-648.3	151.2	489.1	454.1-526.2	134.2	223.5	206.8-241.3	69.6	258.7	231.8-287.9	64.6	195.3	174.3-218.4
	2006-2010	345.0	530.4	505.2-556.6	191.6	612.8	573.8-653.8	153.4	464.3	431.0-499.6	138.0	210.5	194.9-227.1	76.6	250.8	225.8-277.9	61.4	178.4	158.6-200.2
	2011-2015	351.6	515.8	491.1-541.6	181.0	539.2	503.6-576.8	170.6	504.6	469.4-542.1	132.8	188.7	174.2-204.1	73.2	224.0	201.0-249.0	59.6	159.9	141.6-180.4
	2016-2020	358.2	494.6	470.6-519.6	186.4	509.3	475.8-544.9	171.8	490.3	455.2-527.7	125.4	164.2	151.2-178.3	71.0	192.5	172.5-214.5	54.4	139.2	122.3-158.3
Oral cavity and pharynx	1976-1980	6.2	13.5	9.0-19.5	4.4	20.9	12.9-32.1	1.8	7.4	3.2-14.6	2.0	5.1	2.3-9.4	1.4	7.8	3.1-16.2	0.6	3.0	0.5-9.0
	1981-1985	6.0	12.6	8.4-18.3	3.2	15.5	8.6-25.7	2.8	11.1	5.8-19.3	2.0	3.9	1.9-7.5	0.8	3.7	1.0-9.8	1.2	4.3	1.5-10.1
	1986-1990	5.2	10.4	6.7-15.4	3.4	15.3	8.7-24.7	1.8	6.0	2.7-11.9	2.6	4.8	2.6-8.4	1.4	6.1	2.4-12.8	1.2	4.1	1.4-9.4
	1991-1995	4.8	9.2	5.8-13.7	2.6	11.5	6.0-19.8	2.2	7.2	3.5-13.4	0.8	1.4	0.4-3.7	0.2	0.7	0.0-4.4	0.6	1.9	0.4-6.1
	1996-2000	7.4	13.6	9.6-18.9	4.6	17.9	11.3-27.0	2.8	10.9	5.8-18.7	2.2	3.9	1.9-7.1	2.0	7.2	3.4-13.5	0.2	0.5	0.0-4.0
	2001-2005	6.8	11.3	7.8-16.0	3.8	13.8	8.2-21.9	3.0	9.4	5.2-16.0	2.6	4.2	2.2-7.3	1.8	6.4	2.9-12.5	0.8	2.4	0.7-6.8
	2006-2010	9.0	13.6	9.8-18.4	7.4	23.2	16.2-32.4	1.6	5.2	2.1-10.9	1.0	1.5	0.5-3.8	0.4	1.2	0.1-5.0	0.6	1.7	0.4-5.9
	2011-2015	9.6	14.0	10.2-18.9	7.6	22.3	15.6-31.2	2.0	5.9	2.8-11.6	2.0	2.8	1.3-5.5	1.6	4.7	2.0-9.9	0.4	1.0	0.1-4.9
	2016-2020	7.2	9.3	6.4-13.4	5.8	15.0	9.8-22.3	1.4	3.7	1.4-8.8	2.2	2.7	1.3-5.2	1.8	4.4	2.0-9.1	0.4	1.0	0.1-5.1
Esophagus	1976-1980	2.2	5.0	2.4-9.1	1.4	7.3	2.8-15.4	0.8	2.7	0.7-7.6	2.2	4.6	2.2-8.4	1.8	8.3	3.7-16.2	0.4	1.4	0.2-5.9
	1981-1985	2.6	5.1	2.6-9.0	2.0	9.5	4.4-17.9	0.6	1.8	0.4-6.2	2.2	4.9	2.4-9.0	2.0	10.9	5.1-20.2	0.2	0.6	0.0-4.5
	1986-1990	3.8	7.2	4.3-11.5	2.4	10.9	5.6-19.3	1.4	4.3	1.7-9.4	2.2	4.2	2.1-7.7	1.4	6.1	2.4-12.8	0.8	2.4	0.7-6.9
	1991-1995	3.2	5.9	3.3-9.6	2.0	9.1	4.3-16.8	1.2	3.5	1.3-8.2	3.6	6.5	3.8-10.4	2.4	10.5	5.3-18.5	1.2	3.4	1.3-8.1
	1996-2000	5.0	9.1	5.9-13.6	3.4	13.4	7.8-21.7	1.6	5.3	2.2-10.9	3.2	5.7	3.2-9.3	2.2	9.0	4.5-16.2	1.0	2.7	0.9-7.0
	2001-2005	4.8	7.9	5.0-11.9	3.2	11.3	6.4-18.6	1.6	4.8	2.1-10.0	4.4	7.4	4.6-11.4	3.0	10.8	6.0-18.0	1.4	4.5	1.8-9.9
	2006-2010	5.2	7.7	5.0-11.6	3.0	9.5	5.2-16.1	2.2	6.3	3.1-11.8	4.0	5.9	3.6-9.4	3.0	9.5	5.2-16.1	1.0	3.0	1.0-7.7
	2011-2015	4.4	6.0	3.7-9.3	3.2	9.1	5.1-15.3	1.2	2.9	1.1-7.5	3.0	4.2	2.3-7.2	1.8	6.0	2.7-11.7	1.2	2.7	1.0-7.0
	2016-2020	5.2	7.2	4.6-11.1	3.8	10.7	6.2-17.6	1.4	3.6	1.4-8.5	3.4	4.4	2.5-7.5	2.6	6.9	3.6-12.6	0.8	1.6	0.4-5.7
Stomach	1976-1980	5.6	12.5	8.2-18.2	2.2	12.2	5.8-22.3	3.4	12.8	7.3-21.0	4.8	11.2	7.0-16.9	2.0	11.0	5.1-20.4	2.8	10.8	5.8-18.6
	1981-1985	3.6	7.1	4.2-11.4	2.6	12.7	6.6-22.2	1.0	3.3	1.1-8.3	2.0	4.0	1.9-7.5	0.8	4.1	1.1-10.7	1.2	3.9	1.4-9.1
	1986-1990	4.8	9.4	6.0-14.1	2.6	11.4	6.0-19.7	2.2	7.9	3.8-14.7	3.8	7.2	4.3-11.5	2.4	10.5	5.4-18.7	1.4	4.9	1.9-10.7
	1991-1995	4.2	7.6	4.7-11.8	2.0	8.7	4.1-16.0	2.2	6.5	3.2-12.1	3.0	5.5	3.1-9.1	1.2	5.6	2.0-12.2	1.8	5.3	2.4-10.6
	1996-2000	3.2	5.6	3.2-9.2	2.2	9.0	4.4-16.2	1.0	3.4	1.1-8.4	2.4	4.1	2.1-7.4	1.4	5.4	2.2-11.3	1.0	3.3	1.0-8.2
	2001-2005	4.8	7.9	5.1-12.0	3.8	13.6	8.1-21.6	1.0	2.7	0.9-7.0	1.6	2.7	1.2-5.5	1.4	5.5	2.2-11.4	0.2	0.5	0.0-4.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Greene**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	4.0	6.2	3.8-9.9	3.6	12.3	7.2-19.7	0.4	0.9	0.1-4.6	2.6	4.0	2.1-7.0	2.2	7.4	3.6-13.5	0.4	1.0	0.1-4.7
	2011-2015	5.2	7.5	4.8-11.2	3.6	11.7	6.9-19.0	1.6	4.2	1.7-9.4	3.6	5.2	3.0-8.6	2.8	9.1	4.9-15.8	0.8	1.7	0.5-5.7
	2016-2020	4.4	6.3	3.8-10.1	2.6	8.3	4.2-14.9	1.8	4.4	2.0-9.4	1.0	1.2	0.4-3.4	0.8	2.1	0.6-6.3	0.2	0.4	0.0-4.2
Colorectal	1976-1980	30.8	63.8	53.9-75.0	14.6	74.8	57.7-95.3	16.2	59.4	46.8-74.5	12.0	24.7	18.7-32.1	6.8	34.2	23.3-48.5	5.2	18.7	12.0-28.0
	1981-1985	38.0	75.2	64.7-86.9	21.0	96.7	78.7-117.6	17.0	58.5	46.4-73.0	15.8	32.0	25.2-40.1	7.4	35.0	24.4-48.7	8.4	30.6	21.8-42.0
	1986-1990	35.2	67.6	57.8-78.6	17.4	78.8	62.8-97.6	17.8	60.8	48.5-75.5	17.8	33.0	26.5-40.8	9.6	43.1	31.6-57.5	8.2	26.4	18.9-36.3
	1991-1995	33.6	61.7	52.6-71.9	17.6	76.0	60.7-93.9	16.0	52.1	41.1-65.3	16.2	29.7	23.6-37.1	7.8	35.6	25.2-48.8	8.4	26.3	18.8-36.0
	1996-2000	38.2	67.0	57.8-77.3	21.2	88.5	72.2-107.3	17.0	53.1	42.3-66.1	15.6	27.2	21.5-34.0	8.8	36.8	26.6-49.7	6.8	20.4	14.1-29.1
	2001-2005	37.8	63.1	54.4-73.0	21.4	77.2	63.0-93.6	16.4	49.6	39.3-62.1	14.2	23.7	18.5-30.1	7.2	27.1	18.9-37.8	7.0	20.0	13.9-28.4
	2006-2010	35.2	55.2	47.2-64.2	18.6	62.0	49.9-76.4	16.6	50.1	39.5-62.9	12.4	19.2	14.7-24.8	7.2	24.9	17.3-34.8	5.2	14.8	9.5-22.2
	2011-2015	29.0	42.5	35.7-50.4	12.4	38.4	29.2-49.9	16.6	45.5	35.9-57.3	12.8	18.4	14.1-23.9	6.2	18.9	12.7-27.4	6.6	17.5	11.8-25.5
	2016-2020	30.0	44.1	36.9-52.5	16.8	49.7	39.1-62.5	13.2	39.6	29.8-51.8	8.4	12.0	8.5-16.7	4.6	13.7	8.5-21.2	3.8	10.7	6.1-18.0
Colon excluding rectum	1976-1980	22.6	47.0	38.5-56.8	10.8	55.6	41.0-73.6	11.8	43.0	32.5-56.1	10.2	21.1	15.6-28.0	5.6	28.2	18.4-41.4	4.6	16.5	10.3-25.4
	1981-1985	27.4	54.8	45.8-65.0	14.2	66.6	51.7-84.5	13.2	45.3	34.8-58.3	12.6	25.3	19.3-32.7	5.6	25.6	16.8-37.4	7.0	25.0	17.1-35.4
	1986-1990	24.4	45.5	37.7-54.5	11.4	52.5	39.5-68.3	13.0	42.0	32.3-54.0	16.0	29.6	23.5-37.0	8.6	39.0	28.1-52.9	7.4	23.4	16.5-32.7
	1991-1995	23.6	42.8	35.3-51.3	11.6	50.2	37.9-65.2	12.0	38.0	28.8-49.4	15.4	28.3	22.3-35.5	7.2	32.7	22.7-45.4	8.2	25.7	18.3-35.4
	1996-2000	27.8	48.9	41.1-57.9	14.4	60.1	46.8-75.9	13.4	42.2	32.6-54.0	12.0	20.9	15.9-27.0	6.2	26.3	17.7-37.5	5.8	17.3	11.5-25.3
	2001-2005	28.4	47.4	39.9-56.0	14.4	51.5	40.1-65.3	14.0	42.1	32.7-53.8	11.2	18.5	14.0-24.2	5.2	19.7	12.8-29.1	6.0	17.2	11.5-25.0
	2006-2010	25.4	39.4	32.7-47.0	13.2	44.2	34.0-56.6	12.2	36.2	27.4-47.2	9.8	15.1	11.2-20.2	5.4	19.0	12.4-28.0	4.4	12.3	7.6-19.3
	2011-2015	20.4	29.0	23.5-35.5	8.2	24.9	17.7-34.3	12.2	32.2	24.4-42.2	10.0	14.3	10.5-19.2	4.6	14.1	8.9-21.7	5.4	14.3	9.2-21.7
	2016-2020	21.6	31.6	25.5-38.8	11.6	35.0	26.2-46.2	10.0	28.7	20.6-39.3	6.6	9.5	6.4-13.9	3.4	10.2	5.8-17.1	3.2	9.1	4.9-16.1
Rectum & rectosigmoid	1976-1980	8.2	16.8	12.0-23.1	3.8	19.2	11.1-30.8	4.4	16.3	10.1-25.3	1.8	3.6	1.6-7.0	1.2	6.0	2.0-13.6	0.6	2.2	0.4-7.0
	1981-1985	10.6	20.4	15.2-26.9	6.8	30.0	20.6-42.5	3.8	13.2	7.8-21.1	3.2	6.6	3.8-10.9	1.8	9.4	4.2-18.0	1.4	5.6	2.2-12.0
	1986-1990	10.8	22.1	16.5-29.1	6.0	26.3	17.6-37.9	4.8	18.8	11.7-28.6	1.8	3.4	1.5-6.7	1.0	4.1	1.3-9.9	0.8	3.0	0.8-8.2
	1991-1995	10.0	18.9	14.0-25.1	6.0	25.8	17.3-37.0	4.0	14.1	8.5-22.2	0.8	1.4	0.4-3.8	0.6	2.9	0.6-8.3	0.2	0.5	0.0-4.0
	1996-2000	10.4	18.1	13.5-23.8	6.8	28.4	19.5-39.9	3.6	10.9	6.5-17.7	3.6	6.3	3.7-10.1	2.6	10.5	5.5-18.2	1.0	3.2	1.0-8.0
	2001-2005	9.4	15.8	11.6-21.1	7.0	25.6	17.7-35.9	2.4	7.5	3.8-13.5	3.0	5.2	2.9-8.8	2.0	7.4	3.5-13.8	1.0	2.9	0.9-7.4
	2006-2010	9.8	15.8	11.6-21.1	5.4	17.8	11.7-26.3	4.4	13.9	8.5-21.8	2.6	4.0	2.1-7.1	1.8	5.9	2.6-11.5	0.8	2.4	0.6-6.9
	2011-2015	8.6	13.5	9.6-18.6	4.2	13.5	8.1-21.4	4.4	13.3	8.1-20.9	2.8	4.1	2.2-7.3	1.6	4.8	2.0-10.2	1.2	3.2	1.1-8.0
	2016-2020	8.4	12.6	8.9-17.4	5.2	14.7	9.4-22.2	3.2	10.9	6.0-18.6	1.8	2.5	1.1-5.1	1.2	3.5	1.3-8.2	0.6	1.6	0.3-5.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Greene**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	0.8	1.9	0.5-5.0	0.0	0.0	0.0-4.1	0.8	2.8	0.8-7.9	1.0	2.1	0.7-5.1	0.2	0.7	0.0-5.1	0.8	2.8	0.8-7.7
	1981-1985	4.2	8.5	5.2-13.1	1.8	7.6	3.4-14.7	2.4	8.8	4.4-16.0	2.2	4.1	2.0-7.5	1.0	4.2	1.3-10.3	1.2	3.9	1.4-9.1
	1986-1990	2.0	3.7	1.8-6.9	0.8	4.1	1.1-10.4	1.2	3.8	1.4-8.8	1.0	2.0	0.6-4.9	0.4	1.6	0.2-6.3	0.6	2.5	0.5-7.7
	1991-1995	3.2	6.0	3.4-9.9	2.4	10.1	5.2-17.8	0.8	2.2	0.6-6.4	2.4	4.3	2.2-7.7	1.6	6.3	2.7-12.6	0.8	2.1	0.6-6.2
	1996-2000	2.4	4.4	2.3-7.7	1.8	7.0	3.2-13.5	0.6	1.8	0.4-6.0	1.4	2.5	1.0-5.3	1.4	5.5	2.2-11.5	0.0	0.0	0.0-3.2
	2001-2005	4.8	8.2	5.2-12.3	4.6	16.1	10.1-24.4	0.2	0.6	0.0-4.3	3.4	6.0	3.5-9.7	3.2	11.2	6.4-18.4	0.2	0.6	0.0-4.3
	2006-2010	6.2	9.2	6.2-13.3	4.8	14.1	9.0-21.5	1.4	4.4	1.7-9.5	6.2	9.1	6.1-13.1	5.2	15.2	9.9-22.6	1.0	3.2	1.0-8.0
	2011-2015	3.2	4.6	2.6-7.9	3.0	9.0	4.9-15.4	0.2	0.4	0.0-4.1	3.6	4.9	2.8-8.1	2.8	7.8	4.2-13.7	0.8	2.1	0.6-6.4
	2016-2020	6.8	8.2	5.6-11.8	6.2	15.7	10.6-23.0	0.6	1.4	0.3-5.5	5.0	6.0	3.9-9.2	4.4	11.3	7.0-17.7	0.6	1.3	0.3-5.3
Pancreas	1976-1980	7.4	14.8	10.4-20.6	3.6	16.7	9.7-26.9	3.8	13.5	8.1-21.6	6.4	12.8	8.7-18.3	3.2	14.9	8.3-24.8	3.2	11.1	6.3-18.5
	1981-1985	6.6	12.8	8.8-18.2	3.4	16.1	9.2-26.0	3.2	11.0	6.2-18.4	6.0	11.5	7.7-16.7	2.8	12.8	6.9-21.9	3.2	10.9	6.1-18.3
	1986-1990	5.8	11.1	7.4-16.1	3.0	13.9	7.6-23.1	2.8	9.5	5.1-16.4	5.6	10.5	6.9-15.4	3.0	13.6	7.5-22.7	2.6	8.1	4.3-14.4
	1991-1995	5.8	10.4	6.9-15.0	2.2	8.8	4.3-15.9	3.6	11.6	6.8-18.8	5.8	10.5	7.0-15.2	2.2	9.4	4.6-16.9	3.6	11.8	6.9-19.2
	1996-2000	6.2	10.7	7.3-15.3	4.2	16.3	10.0-25.1	2.0	5.8	2.7-11.2	7.0	12.2	8.5-17.1	4.6	18.2	11.5-27.6	2.4	7.1	3.6-12.9
	2001-2005	9.2	14.8	10.8-19.9	3.8	13.6	8.1-21.6	5.4	15.6	10.2-23.1	7.4	12.0	8.4-16.7	3.0	10.8	6.0-18.2	4.4	13.1	8.2-20.3
	2006-2010	10.4	16.1	12.0-21.3	5.4	17.3	11.3-25.5	5.0	14.9	9.5-22.6	9.4	15.3	11.2-20.5	5.8	19.9	13.2-29.0	3.6	10.8	6.4-17.6
	2011-2015	12.8	17.4	13.4-22.6	7.0	20.1	13.9-28.6	5.8	15.6	10.4-23.1	12.8	17.9	13.7-23.1	7.0	21.1	14.6-29.8	5.8	15.6	10.3-23.2
	2016-2020	12.2	16.9	12.8-22.3	7.2	21.0	14.4-30.0	5.0	12.6	8.0-19.5	10.2	14.0	10.2-18.9	6.0	16.6	10.9-24.7	4.2	11.3	6.9-18.2
Larynx	1976-1980	3.8	7.8	4.6-12.5	3.0	14.2	7.8-23.8	0.8	2.6	0.7-7.5	0.6	1.2	0.3-3.8	0.4	2.0	0.2-7.4	0.2	0.8	0.0-5.0
	1981-1985	4.8	10.2	6.4-15.4	4.0	18.9	11.4-29.7	0.8	2.7	0.7-7.5	1.2	2.2	0.8-5.1	1.2	5.7	2.0-12.7	0.0	0.0	0.0-3.6
	1986-1990	3.4	6.2	3.6-10.1	2.4	9.8	5.0-17.5	1.0	3.2	1.0-8.1	1.4	3.2	1.2-6.7	1.0	5.1	1.6-11.9	0.4	1.8	0.2-6.8
	1991-1995	3.6	7.1	4.1-11.3	2.8	12.0	6.5-20.4	0.8	3.0	0.8-8.1	0.4	0.7	0.1-2.7	0.4	1.7	0.2-6.2	0.0	0.0	0.0-3.2
	1996-2000	3.8	6.9	4.1-10.8	3.2	12.3	7.0-20.1	0.6	1.9	0.4-6.1	1.8	3.4	1.5-6.5	1.6	6.8	2.9-13.4	0.2	0.5	0.0-4.0
	2001-2005	2.2	3.7	1.8-6.8	1.6	5.8	2.5-11.7	0.6	1.6	0.3-5.5	0.8	1.2	0.3-3.4	0.8	2.8	0.8-7.6	0.0	0.0	0.0-3.3
	2006-2010	2.4	3.6	1.9-6.5	2.0	6.3	3.0-12.0	0.4	1.1	0.1-5.0	1.2	1.7	0.6-3.9	0.6	1.9	0.4-6.1	0.6	1.5	0.3-5.4
	2011-2015	1.8	3.0	1.3-6.0	1.4	4.8	1.8-10.4	0.4	1.5	0.1-6.3	0.6	0.7	0.1-2.6	0.4	1.2	0.1-5.1	0.2	0.4	0.0-4.1
	2016-2020	3.0	3.7	2.0-6.5	2.6	6.5	3.4-11.8	0.4	1.0	0.1-5.0	0.6	0.8	0.2-2.8	0.4	1.1	0.1-5.0	0.2	0.6	0.0-4.5
Lung and bronchus	1976-1980	30.2	60.3	50.8-71.1	22.6	104.6	85.5-126.7	7.6	27.8	19.4-38.9	27.2	53.7	44.9-63.9	20.4	91.7	74.3-112.1	6.8	25.9	17.6-36.9
	1981-1985	37.0	74.2	63.6-86.2	24.8	113.4	93.7-136.0	12.2	46.8	35.2-61.1	27.8	54.9	45.9-65.2	18.0	82.5	65.9-102.1	9.8	35.4	25.8-47.6
	1986-1990	40.2	79.4	68.6-91.5	25.2	110.1	91.5-131.5	15.0	55.9	43.5-70.9	29.0	57.9	48.6-68.4	18.2	80.3	64.4-98.9	10.8	41.0	30.3-54.3
	1991-1995	51.0	93.8	82.6-106.3	30.4	126.2	106.6-148.3	20.6	71.6	58.1-87.4	40.8	74.4	64.4-85.5	25.2	105.2	87.3-125.6	15.6	53.2	41.8-66.9
	1996-2000	48.4	84.7	74.3-96.2	29.6	114.9	96.9-135.3	18.8	62.1	50.1-76.4	41.0	71.7	62.2-82.3	26.2	105.2	87.7-125.2	14.8	47.8	37.4-60.4
	2001-2005	49.8	82.2	72.3-93.2	24.0	84.0	69.5-100.9	25.8	81.7	68.1-97.5	35.4	58.5	50.2-68.0	19.2	68.0	54.9-83.3	16.2	48.8	38.7-61.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Greene**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	49.6	74.1	65.1-84.2	28.0	87.4	73.3-103.6	21.6	62.5	51.2-76.0	41.0	62.3	54.0-71.6	23.8	75.0	61.9-90.2	17.2	50.8	40.5-63.2
	2011-2015	61.2	84.5	75.1-94.9	33.4	98.4	83.8-115.2	27.8	72.3	60.6-86.2	39.6	54.5	47.0-62.9	21.8	65.3	53.4-79.4	17.8	45.7	36.6-57.0
	2016-2020	62.2	78.0	69.4-87.7	32.2	83.3	70.7-98.0	30.0	74.0	62.3-87.8	32.8	41.3	35.0-48.5	19.4	51.0	41.1-62.9	13.4	32.1	24.7-41.7
Melanoma of the skin	1976-1980	4.0	9.5	5.7-14.9	2.8	15.5	8.3-26.4	1.2	5.0	1.7-11.5	1.6	3.6	1.5-7.2	0.6	3.5	0.7-10.2	1.0	3.7	1.1-9.2
	1981-1985	3.0	7.0	3.8-11.7	1.6	8.5	3.6-16.8	1.4	6.3	2.4-13.3	1.2	3.1	1.1-6.8	1.0	6.1	1.9-14.0	0.2	0.8	0.0-5.1
	1986-1990	3.0	5.7	3.2-9.6	1.6	6.4	2.7-13.0	1.4	5.5	2.1-11.7	0.8	1.5	0.4-3.9	0.6	2.7	0.5-8.1	0.2	0.6	0.0-4.3
	1991-1995	3.8	7.3	4.4-11.5	1.6	7.5	3.2-14.7	2.2	7.4	3.7-13.6	1.2	2.2	0.8-4.9	0.8	3.2	0.9-8.4	0.4	1.2	0.1-5.1
	1996-2000	5.0	9.3	6.0-13.8	3.4	13.6	7.9-21.9	1.6	6.5	2.8-13.0	2.2	4.1	2.1-7.5	1.6	6.2	2.7-12.4	0.6	2.4	0.5-7.2
	2001-2005	10.6	18.7	14.0-24.7	6.2	22.8	15.4-32.5	4.4	15.1	9.3-23.4	1.4	2.3	0.9-4.9	1.4	5.1	2.0-10.7	0.0	0.0	0.0-3.3
	2006-2010	13.4	20.2	15.6-25.8	8.6	27.1	19.5-36.9	4.8	13.9	8.8-21.3	3.0	4.8	2.6-8.0	2.2	7.5	3.7-13.7	0.8	2.5	0.7-7.0
	2011-2015	12.6	19.3	14.6-25.1	7.4	22.4	15.6-31.6	5.2	17.2	10.8-26.2	2.2	2.9	1.4-5.6	1.8	5.0	2.1-10.2	0.4	0.9	0.1-4.6
	2016-2020	13.8	19.6	15.0-25.4	10.8	31.0	23.0-41.3	3.0	9.8	5.1-17.4	1.4	1.8	0.7-4.2	0.8	2.4	0.6-6.9	0.6	1.4	0.3-5.6
Female breast	1976-1980							21.4	84.0	67.9-102.8							9.2	35.3	25.4-47.9
	1981-1985							29.2	122.1	101.9-145.1							8.4	28.9	20.6-39.8
	1986-1990							29.2	107.9	90.6-127.8							9.6	39.8	28.9-53.5
	1991-1995							27.2	102.4	85.5-121.8							8.4	29.2	20.8-39.9
	1996-2000							37.8	134.1	115.4-155.2							10.2	31.9	23.6-42.4
	2001-2005							39.8	131.7	113.7-151.9							10.0	29.5	21.8-39.4
	2006-2010							41.6	126.0	109.0-145.1							9.0	25.0	18.1-34.0
	2011-2015							47.0	144.8	125.8-166.1							8.2	24.2	17.0-33.7
	2016-2020							50.4	144.4	125.6-165.4							9.0	25.6	18.0-35.7
Cervix uteri	1976-1980							3.2	15.1	8.3-25.1							1.0	3.6	1.1-9.1
	1981-1985							3.2	15.2	8.4-25.2							1.4	5.9	2.2-12.8
	1986-1990							4.4	20.4	12.5-31.2							1.8	8.0	3.5-15.4
	1991-1995							3.6	15.8	9.3-25.0							1.0	4.3	1.4-10.1
	1996-2000							3.6	13.4	7.9-21.5							1.0	3.8	1.2-9.2
	2001-2005							2.4	10.0	5.0-17.9							1.0	3.0	1.0-7.6
	2006-2010							3.0	11.2	6.1-19.0							0.6	2.4	0.4-7.3
	2011-2015							2.4	8.9	4.4-16.4							1.6	6.7	2.7-13.7
	2016-2020							1.6	5.7	2.1-12.4							0.4	1.1	0.1-5.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Greene**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							3.2	12.1	6.6-20.6							1.4	4.4	1.8-9.8
	1981-1985							6.4	25.9	17.4-37.3							1.4	5.7	2.1-12.4
	1986-1990							6.2	23.2	15.5-33.5							0.8	2.5	0.7-7.1
	1991-1995							6.4	23.8	16.1-33.9							1.0	3.5	1.1-8.6
	1996-2000							8.4	29.4	21.1-40.1							1.0	3.2	1.0-8.0
	2001-2005							8.6	28.1	20.3-38.3							1.8	5.2	2.4-10.5
	2006-2010							11.0	32.5	24.2-42.9							1.0	2.8	0.9-7.3
	2011-2015							11.8	32.7	24.6-42.9							2.2	5.7	2.8-11.1
	2016-2020							12.2	32.2	24.2-42.4							1.8	4.0	1.8-8.7
Ovary	1976-1980							3.8	16.8	9.8-26.9							2.4	9.0	4.5-16.3
	1981-1985							4.8	21.4	13.4-32.6							3.4	14.8	8.3-24.3
	1986-1990							3.6	13.8	7.9-22.4							3.2	12.6	7.0-20.8
	1991-1995							5.6	21.7	14.2-31.7							4.2	14.7	9.0-22.9
	1996-2000							5.0	17.3	11.1-25.9							3.6	11.4	6.7-18.5
	2001-2005							5.8	20.4	13.5-29.7							4.4	13.5	8.4-20.9
	2006-2010							4.2	13.9	8.3-22.0							3.6	12.2	7.0-20.0
	2011-2015							5.2	15.2	9.8-23.1							3.4	9.2	5.3-15.6
	2016-2020							4.0	12.1	7.0-19.8							3.2	8.5	4.7-14.9
Prostate	1976-1980				15.0	79.3	61.5-100.5							5.6	29.9	19.4-43.9			
	1981-1985				23.0	111.6	91.7-134.5							7.8	41.4	29.1-57.0			
	1986-1990				26.2	117.5	97.9-139.9							5.2	25.0	16.1-36.8			
	1991-1995				30.0	124.6	105.2-146.7							7.0	32.5	22.5-45.4			
	1996-2000				38.8	149.6	129.0-172.5							5.8	24.8	16.5-35.9			
	2001-2005				39.0	134.0	115.6-154.6							5.6	22.8	15.1-33.2			
	2006-2010				51.0	155.6	136.8-176.6							6.8	25.2	17.4-35.5			
	2011-2015				36.0	100.4	86.0-116.9							4.8	15.6	9.9-23.7			
	2016-2020				38.0	93.4	80.3-108.4							5.2	14.0	9.1-21.1			
Testis	1976-1980				1.2	6.4	2.3-14.3							0.0	0.0	0.0-4.1			
	1981-1985				0.8	3.7	1.0-9.8							0.2	0.8	0.0-5.1			
	1986-1990				2.0	8.8	4.2-16.5							0.0	0.0	0.0-3.6			
	1991-1995				2.0	7.8	3.7-14.5							0.6	2.6	0.5-7.7			
	1996-2000				1.0	4.0	1.3-9.5							0.0	0.0	0.0-3.1			
	2001-2005				1.2	4.2	1.5-9.4							0.2	1.0	0.0-5.2			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Greene**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				2.6	11.6	6.1-19.8							0.0	0.0	0.0-3.1			
	2011-2015				1.8	6.9	3.0-13.4							0.2	0.7	0.0-4.6			
	2016-2020				1.0	4.5	1.4-10.5							0.0	0.0	0.0-3.4			
Urinary bladder (incl. in situ)	1976-1980	11.2	23.3	17.5-30.5	7.8	41.1	28.6-57.0	3.4	12.0	6.9-19.8	2.0	4.5	2.1-8.4	1.2	7.3	2.5-15.9	0.8	3.2	0.8-8.6
	1981-1985	11.0	21.6	16.2-28.3	7.4	34.5	24.1-48.0	3.6	12.2	7.1-19.8	2.4	4.7	2.4-8.4	1.2	6.3	2.2-13.8	1.2	3.7	1.4-8.8
	1986-1990	14.0	27.1	21.1-34.5	11.2	52.4	39.3-68.4	2.8	9.5	5.1-16.5	2.4	4.8	2.5-8.6	2.4	13.6	6.9-23.6	0.0	0.0	0.0-3.4
	1991-1995	13.0	24.2	18.6-31.0	10.2	43.1	32.0-56.9	2.8	8.6	4.6-14.9	3.0	5.5	3.1-9.2	1.6	6.9	2.9-13.7	1.4	4.1	1.6-9.0
	1996-2000	16.8	29.5	23.5-36.6	12.4	49.2	37.6-63.3	4.4	12.8	8.0-20.0	4.6	7.8	5.0-11.9	2.6	11.6	6.1-19.9	2.0	5.7	2.7-11.1
	2001-2005	24.0	39.3	32.6-47.2	19.0	68.0	54.9-83.5	5.0	15.3	9.9-23.1	2.8	4.7	2.6-8.0	2.0	8.0	3.8-14.9	0.8	2.4	0.6-6.8
	2006-2010	22.2	33.5	27.5-40.5	18.0	58.4	46.8-72.3	4.2	11.4	7.0-18.0	3.8	5.6	3.4-9.0	2.8	8.9	4.8-15.4	1.0	2.7	0.9-7.1
	2011-2015	26.2	37.3	31.1-44.6	19.4	56.9	46.0-70.0	6.8	19.6	13.4-28.1	4.0	5.7	3.4-9.1	3.2	10.0	5.6-16.7	0.8	2.1	0.6-6.4
	2016-2020	21.0	28.7	23.3-35.2	16.2	45.1	35.6-56.8	4.8	13.2	8.2-20.8	5.0	6.5	4.2-10.0	4.0	11.0	6.7-17.6	1.0	2.5	0.8-7.0
Kidney and renal pelvis	1976-1980	3.8	7.8	4.7-12.5	2.4	10.8	5.5-19.3	1.4	5.8	2.2-12.3	2.2	4.2	2.1-7.8	1.4	6.2	2.5-13.4	0.8	2.9	0.8-8.1
	1981-1985	3.2	6.4	3.6-10.6	1.4	6.1	2.4-13.0	1.8	6.2	2.7-12.6	1.4	2.5	1.0-5.5	0.8	3.4	0.9-9.2	0.6	1.9	0.4-6.3
	1986-1990	6.4	12.8	8.7-18.2	3.8	17.2	10.2-27.1	2.6	8.5	4.5-15.1	1.0	1.8	0.6-4.5	0.6	2.4	0.5-7.5	0.4	1.2	0.1-5.2
	1991-1995	7.2	13.4	9.3-18.6	3.2	13.9	7.9-22.8	4.0	13.2	8.0-20.9	3.0	5.3	2.9-8.8	1.6	7.1	3.0-14.0	1.4	4.4	1.7-9.5
	1996-2000	11.0	19.5	14.7-25.5	7.0	27.5	19.1-38.5	4.0	12.9	7.8-20.4	2.8	5.2	2.8-8.9	2.0	8.3	4.0-15.4	0.8	2.2	0.6-6.3
	2001-2005	8.2	13.8	9.9-18.9	5.8	20.6	13.7-29.8	2.4	8.3	4.2-14.9	4.6	7.7	4.9-11.7	3.0	10.6	5.9-17.8	1.6	5.0	2.1-10.3
	2006-2010	8.2	12.7	9.1-17.4	6.0	19.1	12.8-27.7	2.2	7.0	3.4-13.1	2.8	4.2	2.3-7.3	2.4	7.8	3.9-13.9	0.4	1.3	0.1-5.3
	2011-2015	12.6	18.3	13.9-23.9	8.6	24.8	17.7-34.2	4.0	12.0	7.0-19.6	3.0	4.1	2.3-7.1	1.6	4.4	1.9-9.4	1.4	3.2	1.3-7.7
	2016-2020	11.2	15.4	11.5-20.6	7.2	19.8	13.6-28.3	4.0	11.2	6.7-18.4	2.2	3.0	1.5-5.8	2.0	5.5	2.6-10.9	0.2	0.7	0.0-5.0
Brain and other central nervous system	1976-1980	2.4	5.4	2.7-9.6	1.4	6.9	2.7-14.6	1.0	4.2	1.3-10.2	1.8	3.5	1.6-6.9	1.2	5.1	1.8-11.7	0.6	2.2	0.4-7.2
	1981-1985	2.6	5.1	2.7-8.9	1.8	7.9	3.6-15.3	0.8	3.0	0.8-8.1	1.2	2.2	0.8-5.0	0.4	1.7	0.2-6.7	0.8	2.6	0.7-7.5
	1986-1990	3.2	6.5	3.7-10.7	2.6	11.5	6.0-20.0	0.6	1.9	0.4-6.2	1.6	3.3	1.4-6.7	1.2	5.6	2.0-12.4	0.4	1.3	0.2-5.3
	1991-1995	3.4	6.7	3.9-10.9	2.2	9.7	4.8-17.4	1.2	4.3	1.5-9.8	1.4	3.0	1.2-6.2	1.0	4.2	1.4-10.0	0.4	1.5	0.2-5.8
	1996-2000	3.2	6.2	3.5-10.1	1.6	6.3	2.7-12.5	1.6	6.1	2.6-12.3	1.6	3.1	1.3-6.1	0.6	2.4	0.5-7.1	1.0	3.6	1.1-8.8
	2001-2005	3.0	5.3	2.9-8.8	1.4	4.8	1.9-10.1	1.6	5.6	2.3-11.5	2.4	4.4	2.2-7.8	1.4	5.2	2.1-11.0	1.0	3.5	1.0-8.7
	2006-2010	4.6	8.1	5.0-12.4	3.6	12.7	7.4-20.6	1.0	2.9	0.9-7.5	2.4	3.7	1.9-6.7	1.8	5.7	2.6-11.2	0.6	1.9	0.4-6.1
	2011-2015	3.2	5.9	3.2-10.0	1.8	5.7	2.5-11.4	1.4	6.8	2.6-14.2	2.8	4.0	2.1-7.2	2.2	6.6	3.1-12.5	0.6	1.4	0.3-5.4
	2016-2020	4.0	5.8	3.4-9.5	1.8	6.1	2.6-12.3	2.2	5.4	2.6-10.7	3.6	4.6	2.7-7.6	1.8	4.7	2.1-9.7	1.8	4.4	2.0-9.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Greene**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	2.2	5.5	2.7-9.9	1.2	7.9	2.8-17.0	1.0	4.8	1.5-11.5	0.0	0.0	0.0-1.9	0.0	0.0	0.0-4.1	0.0	0.0	0.0-3.6
	1981-1985	1.0	1.8	0.6-4.5	0.8	3.2	0.8-8.7	0.2	0.6	0.0-4.5	1.0	1.9	0.6-4.7	0.8	3.7	1.0-9.8	0.2	0.6	0.0-4.5
	1986-1990	1.6	3.3	1.4-6.7	0.4	1.8	0.2-6.7	1.2	4.8	1.7-10.8	0.2	0.3	0.0-2.3	0.0	0.0	0.0-3.6	0.2	0.6	0.0-4.3
	1991-1995	1.8	3.8	1.7-7.2	0.6	2.3	0.5-7.1	1.2	5.4	2.0-11.7	0.2	0.4	0.0-2.2	0.2	0.7	0.0-4.5	0.0	0.0	0.0-3.2
	1996-2000	1.6	3.3	1.4-6.5	0.8	3.3	0.9-8.5	0.8	3.4	0.9-8.8	0.0	0.0	0.0-1.5	0.0	0.0	0.0-3.1	0.0	0.0	0.0-3.2
	2001-2005	5.0	9.6	6.1-14.3	2.0	8.3	3.9-15.3	3.0	11.4	6.3-19.2	0.4	0.7	0.1-2.7	0.0	0.0	0.0-3.1	0.4	1.3	0.1-5.4
	2006-2010	8.2	14.1	10.0-19.4	2.2	8.0	3.8-14.7	6.0	20.7	13.7-30.2	0.2	0.2	0.0-1.9	0.0	0.0	0.0-3.1	0.2	0.5	0.0-4.0
	2011-2015	10.2	19.4	14.2-25.9	3.2	12.3	6.9-20.5	7.0	27.3	18.6-38.9	0.8	1.0	0.3-3.1	0.8	2.3	0.6-6.7	0.0	0.0	0.0-3.6
	2016-2020	11.2	20.9	15.4-27.8	2.6	7.9	3.9-14.4	8.6	35.7	25.0-49.5	1.0	1.2	0.4-3.4	0.4	1.1	0.1-5.0	0.6	1.2	0.3-5.2
Hodgkin lymphoma	1976-1980	1.6	4.1	1.7-8.1	0.8	4.1	1.1-10.7	0.8	3.9	1.0-10.1	0.4	1.1	0.1-3.9	0.2	0.8	0.0-5.4	0.2	1.4	0.0-6.8
	1981-1985	2.2	4.5	2.2-8.2	1.0	4.3	1.4-10.5	1.2	4.9	1.8-11.1	0.8	1.9	0.5-4.9	0.6	2.8	0.6-8.6	0.2	0.9	0.0-5.3
	1986-1990	1.2	2.7	1.0-6.0	1.0	3.9	1.2-9.6	0.2	1.2	0.0-6.0	0.2	0.3	0.0-2.3	0.2	0.8	0.0-4.9	0.0	0.0	0.0-3.4
	1991-1995	1.2	2.3	0.8-5.1	0.2	0.9	0.0-5.1	1.0	3.9	1.2-9.5	0.2	0.4	0.0-2.2	0.0	0.0	0.0-3.3	0.2	0.7	0.0-4.3
	1996-2000	1.8	3.7	1.7-7.0	0.8	3.2	0.9-8.3	1.0	4.3	1.3-10.3	0.2	0.3	0.0-2.1	0.0	0.0	0.0-3.1	0.2	0.5	0.0-4.0
	2001-2005	3.6	7.6	4.5-12.1	2.6	9.7	5.1-16.8	1.0	4.8	1.5-11.3	0.2	0.4	0.0-2.2	0.0	0.0	0.0-3.1	0.2	0.6	0.0-4.3
	2006-2010	1.4	2.7	1.0-5.6	0.4	1.1	0.1-4.7	1.0	4.9	1.6-11.4	0.6	1.1	0.2-3.3	0.6	2.0	0.4-6.1	0.0	0.0	0.0-3.3
	2011-2015	2.2	4.6	2.2-8.5	1.4	5.0	1.9-11.0	0.8	4.2	1.0-10.8	0.2	0.3	0.0-2.1	0.2	0.6	0.0-4.1	0.0	0.0	0.0-3.6
	2016-2020	2.4	4.2	2.1-7.6	1.6	5.7	2.4-11.7	0.8	2.5	0.6-7.4	0.2	0.6	0.0-3.1	0.2	1.2	0.0-5.9	0.0	0.0	0.0-3.8
Non-Hodgkin lymphomas	1976-1980	6.2	13.5	9.0-19.4	2.8	12.7	6.9-21.8	3.4	13.8	7.8-22.8	4.0	8.6	5.2-13.5	2.0	9.7	4.5-18.4	2.0	8.1	3.7-15.4
	1981-1985	9.4	18.1	13.2-24.3	5.2	23.0	14.9-34.2	4.2	13.7	8.4-21.5	4.6	8.9	5.6-13.6	2.6	11.2	5.9-19.6	2.0	7.5	3.4-14.4
	1986-1990	9.4	19.5	14.2-26.1	4.8	21.0	13.3-31.5	4.6	17.7	10.9-27.3	5.6	10.5	6.9-15.4	3.4	14.8	8.5-24.0	2.2	6.8	3.4-12.6
	1991-1995	9.0	17.8	12.9-24.0	5.6	25.6	16.9-37.1	3.4	12.0	6.9-19.6	4.4	8.3	5.2-12.7	2.4	11.1	5.6-19.4	2.0	6.7	3.2-12.8
	1996-2000	13.2	25.1	19.4-32.1	8.2	33.2	23.8-45.2	5.0	16.6	10.6-24.9	7.0	12.5	8.7-17.5	2.8	11.9	6.4-20.0	4.2	13.5	8.3-21.0
	2001-2005	13.6	22.7	17.6-28.9	8.0	27.6	19.6-37.9	5.6	16.7	11.1-24.6	4.2	7.0	4.3-10.8	2.8	10.0	5.4-17.0	1.4	4.2	1.6-9.3
	2006-2010	15.4	23.7	18.6-29.9	8.4	27.4	19.6-37.4	7.0	20.1	13.9-28.6	4.2	6.3	3.9-9.9	2.6	9.3	4.8-16.2	1.6	4.2	1.8-9.1
	2011-2015	15.8	24.0	18.8-30.2	8.6	26.4	19.0-36.1	7.2	22.2	15.2-31.8	4.8	7.0	4.4-10.7	3.6	11.1	6.5-18.0	1.2	3.2	1.1-7.9
	2016-2020	16.0	23.2	18.1-29.5	7.6	22.5	15.6-31.6	8.4	24.9	17.3-35.2	4.4	6.1	3.8-9.5	2.6	7.7	4.1-13.7	1.8	4.7	2.1-9.9
Myeloma	1976-1980	2.0	4.0	1.9-7.7	0.8	4.1	1.1-10.7	1.2	3.9	1.4-9.3	1.4	2.7	1.1-5.9	1.0	4.5	1.4-11.0	0.4	1.3	0.2-5.6
	1981-1985	2.8	5.5	3.0-9.4	1.4	7.0	2.8-14.7	1.4	4.4	1.8-9.8	2.4	4.7	2.4-8.4	1.4	7.1	2.8-14.8	1.0	3.2	1.0-8.1
	1986-1990	2.4	4.7	2.4-8.4	1.2	5.5	2.0-12.0	1.2	3.7	1.3-8.6	1.6	3.4	1.4-6.7	0.2	0.9	0.0-5.2	1.4	5.0	1.9-10.8
	1991-1995	1.8	3.2	1.5-6.2	1.6	7.2	3.1-14.3	0.2	0.5	0.0-4.0	2.4	4.2	2.2-7.5	1.8	8.2	3.7-15.6	0.6	1.8	0.4-5.8
	1996-2000	2.4	4.2	2.2-7.5	0.2	0.7	0.0-4.3	2.2	7.3	3.6-13.4	2.2	3.9	2.0-7.1	0.8	3.1	0.8-8.1	1.4	4.5	1.8-9.7
	2001-2005	3.2	5.2	3.0-8.7	1.6	5.6	2.4-11.3	1.6	4.7	2.0-9.8	1.4	2.3	0.9-4.8	0.8	2.6	0.7-7.2	0.6	1.8	0.4-5.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Greene**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	4.8	7.4	4.8-11.3	2.2	7.1	3.5-13.1	2.6	7.8	4.1-13.8	2.0	3.1	1.5-5.9	0.6	1.9	0.4-6.0	1.4	3.8	1.5-8.5
	2011-2015	4.2	5.9	3.6-9.4	2.6	8.1	4.2-14.5	1.6	3.9	1.7-8.7	2.6	3.8	2.0-6.8	1.8	5.7	2.6-11.3	0.8	2.2	0.6-6.6
	2016-2020	4.8	6.0	3.8-9.4	3.2	8.3	4.7-14.2	1.6	3.8	1.6-8.6	2.0	2.6	1.3-5.2	1.0	2.9	0.9-7.5	1.0	2.3	0.7-6.8
Leukemias	1976-1980	7.2	16.2	11.2-22.8	4.0	22.3	13.3-35.0	3.2	11.6	6.6-19.4	4.6	10.0	6.3-15.2	1.8	9.6	4.2-18.7	2.8	10.1	5.4-17.4
	1981-1985	7.0	15.1	10.4-21.2	3.6	19.0	11.0-30.3	3.4	12.5	7.2-20.5	4.8	9.7	6.2-14.7	2.2	11.9	5.8-21.5	2.6	8.5	4.5-15.1
	1986-1990	6.2	13.0	8.7-18.5	3.8	18.3	10.9-28.6	2.4	8.8	4.4-16.0	4.0	8.7	5.3-13.6	2.6	12.5	6.6-21.4	1.4	5.8	2.2-12.3
	1991-1995	6.4	12.5	8.5-17.8	3.8	16.9	10.1-26.5	2.6	7.9	4.2-14.0	4.4	8.4	5.3-12.9	2.4	10.8	5.5-18.9	2.0	6.4	3.0-12.1
	1996-2000	9.6	17.4	12.8-23.2	6.2	25.5	17.2-36.4	3.4	11.9	6.8-19.6	3.6	6.5	3.9-10.4	2.4	10.5	5.3-18.4	1.2	3.8	1.4-8.8
	2001-2005	9.8	16.5	12.1-21.9	5.4	19.9	13.0-29.3	4.4	14.4	8.9-22.4	6.0	10.3	6.9-14.8	2.8	11.7	6.3-19.7	3.2	10.2	5.7-17.2
	2006-2010	9.4	14.2	10.4-19.1	5.8	18.4	12.2-26.8	3.6	10.5	6.2-17.2	3.8	5.5	3.3-8.8	2.0	6.4	3.0-12.3	1.8	4.7	2.1-9.7
	2011-2015	10.8	15.8	11.7-21.1	8.0	24.0	16.8-33.4	2.8	8.5	4.5-15.0	3.6	5.7	3.2-9.4	2.4	7.8	3.9-14.3	1.2	3.9	1.3-9.6
2016-2020	13.6	19.4	14.8-25.2	7.6	21.4	14.9-30.1	6.0	17.7	11.5-26.5	4.8	6.2	3.9-9.6	2.6	7.0	3.7-12.7	2.2	5.4	2.6-10.8	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Hamilton**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	25.2	431.9	355.1-520.7	12.0	455.5	337.2-601.3	13.2	423.6	322.5-547.6	14.8	273.5	210.5-349.2	7.2	300.7	200.9-430.6	7.6	258.1	179.0-361.1
	1981-1985	27.8	478.3	398.0-570.5	14.2	563.3	428.5-726.1	13.6	443.6	339.6-570.7	16.2	283.4	222.6-356.1	9.6	417.0	299.5-563.7	6.6	199.3	135.4-285.5
	1986-1990	28.0	425.4	355.5-505.9	15.4	529.8	409.8-674.5	12.6	355.5	271.5-460.1	17.4	264.1	209.7-329.3	8.2	280.9	195.5-391.4	9.2	256.9	186.6-348.0
	1991-1995	31.6	452.7	383.1-532.4	16.0	514.0	402.4-648.1	15.6	430.1	338.0-542.2	18.8	266.0	213.8-328.1	10.8	356.8	263.7-472.8	8.0	207.8	147.5-287.9
	1996-2000	43.2	551.8	479.1-634.0	25.0	668.6	552.3-804.9	18.2	446.6	358.9-553.6	17.4	222.1	177.3-276.5	9.2	249.3	181.4-337.5	8.2	197.8	141.4-274.2
	2001-2005	48.2	599.3	523.7-684.8	28.6	758.7	633.8-904.4	19.6	460.7	370.9-570.4	18.8	232.0	186.5-287.5	10.4	282.1	207.9-378.0	8.4	198.7	141.8-276.6
	2006-2010	40.0	480.5	413.2-558.3	22.0	534.1	433.5-656.5	18.0	437.6	347.2-550.7	16.6	197.9	156.4-250.1	10.4	265.0	195.2-357.5	6.2	139.2	94.3-207.5
	2011-2015	47.0	543.3	469.3-628.6	22.4	480.2	391.2-591.1	24.6	608.9	492.5-749.9	16.8	187.2	147.4-238.4	9.8	222.6	161.2-307.7	7.0	152.2	104.7-224.1
	2016-2020	43.4	490.9	416.7-578.2	25.8	550.6	445.9-681.4	17.6	429.4	327.7-559.8	15.0	158.0	121.0-208.1	8.6	173.0	123.9-248.9	6.4	142.7	90.3-225.0
Oral cavity and pharynx	1976-1980	1.2	29.9	10.0-65.9	1.0	54.6	15.0-130.8	0.2	10.6	0.3-52.7	0.6	15.1	2.6-44.0	0.6	37.0	6.2-106.4	0.0	0.0	0.0-30.6
	1981-1985	0.4	6.4	0.6-26.6	0.0	0.0	0.0-33.9	0.4	12.5	1.3-51.5	0.2	4.2	0.1-23.7	0.2	10.8	0.3-55.7	0.0	0.0	0.0-30.9
	1986-1990	1.4	20.7	8.1-45.1	1.0	29.2	8.7-75.5	0.4	11.6	1.4-47.9	0.6	12.1	2.5-35.5	0.2	9.2	0.2-49.6	0.4	14.0	1.6-53.5
	1991-1995	0.8	12.4	3.1-34.1	0.4	13.2	1.3-52.8	0.4	11.3	1.2-47.4	0.4	6.5	0.7-25.7	0.2	9.2	0.2-48.0	0.2	4.2	0.1-35.2
	1996-2000	2.0	28.4	13.2-55.2	1.4	40.6	15.6-89.6	0.6	15.2	3.1-53.7	0.6	7.9	1.6-26.8	0.6	15.6	3.1-53.3	0.0	0.0	0.0-32.4
	2001-2005	1.2	19.9	6.4-47.4	1.0	33.9	10.1-85.4	0.2	4.4	0.1-40.0	0.6	7.6	1.5-27.1	0.4	11.3	1.4-49.7	0.2	4.4	0.1-39.9
	2006-2010	0.8	8.7	2.3-29.8	0.4	8.5	1.0-47.8	0.4	8.8	1.0-50.5	0.2	1.9	0.0-21.3	0.0	0.0	0.0-37.6	0.2	4.1	0.1-44.6
	2011-2015	1.0	10.4	3.2-33.2	1.0	20.7	6.3-66.2	0.0	0.0	0.0-42.3	0.0	0.0	0.0-20.8	0.0	0.0	0.0-41.4	0.0	0.0	0.0-42.3
	2016-2020	1.0	9.5	3.0-34.5	0.6	10.6	2.1-59.7	0.4	8.0	0.9-57.2	0.4	5.4	0.5-31.3	0.2	7.4	0.2-59.3	0.2	3.1	0.1-51.7
Esophagus	1976-1980	0.4	6.0	0.7-25.0	0.2	6.8	0.2-44.8	0.2	5.3	0.1-38.4	0.6	9.2	1.9-29.7	0.4	13.6	1.6-55.1	0.2	5.3	0.1-38.4
	1981-1985	0.6	13.1	2.6-38.5	0.6	33.7	6.3-96.1	0.0	0.0	0.0-30.9	0.6	11.6	2.1-35.6	0.6	28.3	4.6-86.1	0.0	0.0	0.0-30.9
	1986-1990	0.6	10.9	2.1-33.6	0.6	24.3	4.5-73.7	0.0	0.0	0.0-30.0	0.4	7.0	0.7-27.6	0.4	14.6	1.3-57.9	0.0	0.0	0.0-30.0
	1991-1995	0.2	4.3	0.1-23.0	0.2	9.2	0.2-48.0	0.0	0.0	0.0-29.6	0.2	3.6	0.1-21.1	0.2	11.1	0.3-53.7	0.0	0.0	0.0-29.6
	1996-2000	0.6	7.7	1.5-26.4	0.4	9.1	1.1-43.1	0.2	5.1	0.1-39.5	0.6	8.6	1.6-28.6	0.6	17.9	3.3-58.7	0.0	0.0	0.0-32.4
	2001-2005	0.6	6.3	1.3-24.7	0.6	11.9	2.5-48.2	0.0	0.0	0.0-34.4	0.6	7.0	1.4-26.0	0.6	13.5	2.7-51.4	0.0	0.0	0.0-34.4
	2006-2010	0.8	7.6	2.0-27.9	0.8	14.9	4.0-55.0	0.0	0.0	0.0-40.1	0.8	7.7	2.1-28.2	0.6	11.4	2.3-50.8	0.2	4.1	0.1-44.6
	2011-2015	0.4	6.4	0.6-30.1	0.2	4.1	0.1-45.7	0.2	9.3	0.2-57.7	0.8	10.1	2.4-34.3	0.6	12.0	2.4-55.0	0.2	9.3	0.2-57.7
	2016-2020	1.2	11.8	4.2-37.5	1.2	23.4	8.3-74.8	0.0	0.0	0.0-49.0	0.4	4.3	0.5-29.2	0.4	8.7	1.0-58.6	0.0	0.0	0.0-49.0
Stomach	1981-1985	0.4	8.3	0.8-31.2	0.4	20.9	1.8-76.5	0.0	0.0	0.0-30.9	0.4	9.8	1.2-34.2	0.4	26.3	3.0-86.8	0.0	0.0	0.0-30.9
	1986-1990	0.6	7.7	1.5-26.4	0.2	6.2	0.2-41.8	0.4	9.2	1.1-43.0	0.4	5.5	0.6-23.7	0.0	0.0	0.0-31.9	0.4	9.8	1.1-44.4
	1991-1995	0.4	5.2	0.6-22.4	0.4	11.3	1.2-47.7	0.0	0.0	0.0-29.6	0.4	5.3	0.6-22.7	0.4	11.8	1.3-48.5	0.0	0.0	0.0-29.6
	1996-2000	1.2	14.5	5.3-35.3	1.0	25.0	8.0-65.7	0.2	4.6	0.1-38.5	0.8	10.2	2.8-29.8	0.6	17.0	3.5-55.8	0.2	4.7	0.1-38.6
	2001-2005	0.6	7.6	1.5-27.3	0.6	18.6	3.5-61.8	0.0	0.0	0.0-34.4	0.8	9.7	2.6-29.8	0.6	18.6	3.5-61.8	0.2	3.6	0.1-38.5
	2006-2010	0.6	6.4	1.3-27.0	0.2	3.6	0.1-41.4	0.4	8.4	1.0-49.7	0.6	6.7	1.4-27.4	0.2	3.6	0.1-41.4	0.4	8.9	1.1-50.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Hamilton**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2011-2015	1.6	16.5	7.0-40.6	1.4	28.8	11.2-75.9	0.2	4.2	0.1-46.8	0.2	3.2	0.1-25.1	0.2	6.5	0.2-50.3	0.0	0.0	0.0-42.3
Colorectal	1976-1980	3.8	57.9	33.5-94.3	1.6	46.8	18.3-101.8	2.2	66.1	31.8-124.2	1.6	29.3	11.4-61.3	0.6	17.0	3.3-58.9	1.0	36.0	10.5-88.9
	1981-1985	4.8	80.8	50.2-123.8	2.2	75.7	35.1-143.9	2.6	88.8	45.3-157.4	2.4	37.2	19.0-67.4	1.0	41.8	13.0-100.4	1.4	37.9	15.1-84.0
	1986-1990	5.2	74.1	47.9-111.1	3.0	84.4	45.9-146.3	2.2	62.1	30.9-115.6	2.8	42.0	22.7-72.7	1.2	42.2	14.8-96.2	1.6	43.5	18.7-91.0
	1991-1995	6.0	85.2	57.0-123.9	3.0	100.1	54.0-170.4	3.0	80.3	44.3-137.6	2.4	36.8	18.7-66.1	1.4	54.1	20.7-114.2	1.0	26.2	8.3-67.5
	1996-2000	5.4	64.5	42.4-96.5	3.0	73.8	41.1-127.5	2.4	56.5	29.2-105.8	2.0	24.0	11.5-47.4	1.6	39.1	16.7-83.9	0.4	9.3	1.1-44.8
	2001-2005	5.0	60.7	39.1-92.9	2.6	65.4	34.5-119.2	2.4	52.9	27.2-101.4	1.8	21.2	9.6-44.5	0.8	18.0	4.8-57.1	1.0	23.2	7.5-64.6
	2006-2010	3.4	37.0	21.4-64.8	2.4	54.4	27.4-106.5	1.0	21.9	7.0-66.6	1.0	11.6	3.7-33.8	0.8	17.7	4.8-59.7	0.2	5.3	0.1-46.7
	2011-2015	3.4	47.2	24.6-84.8	1.2	27.2	9.5-75.0	2.2	68.5	28.5-141.4	1.8	19.6	8.9-44.6	1.2	25.8	9.3-72.3	0.6	13.3	2.7-57.8
2016-2020	4.2	48.7	28.0-85.0	2.0	36.3	16.9-89.4	2.2	60.9	26.9-129.3	1.4	14.7	5.8-41.2	0.6	10.0	2.0-59.0	0.8	18.9	5.0-70.5	
Colon excluding rectum	1976-1980	2.6	40.3	20.8-72.0	1.0	33.1	9.3-85.7	1.6	46.9	19.9-97.5	0.8	15.2	3.5-41.6	0.2	7.7	0.2-47.0	0.6	19.5	3.3-63.4
	1981-1985	2.8	50.0	26.2-86.8	0.8	26.0	6.9-72.6	2.0	70.9	32.4-135.5	1.8	28.8	13.0-57.1	0.6	29.0	5.8-84.1	1.2	33.2	12.1-78.2
	1986-1990	3.4	48.2	27.6-79.9	1.8	50.4	21.9-103.3	1.6	44.0	18.8-91.8	2.4	36.0	18.4-65.0	1.0	37.7	11.8-91.0	1.4	37.4	14.9-82.7
	1991-1995	5.0	70.6	45.2-106.4	2.2	76.4	36.4-141.4	2.8	73.0	39.4-127.8	1.8	26.4	11.8-52.3	1.0	35.8	10.8-87.1	0.8	19.1	5.1-56.4
	1996-2000	3.8	44.7	26.9-72.7	2.0	48.4	23.1-95.6	1.8	42.6	19.4-88.4	2.0	24.0	11.5-47.4	1.6	39.1	16.7-83.9	0.4	9.3	1.1-44.8
	2001-2005	3.2	38.2	21.7-65.7	1.2	28.8	10.4-71.9	2.0	43.5	20.8-89.6	1.0	12.3	3.9-33.3	0.2	4.0	0.1-38.2	0.8	18.8	5.1-59.0
	2006-2010	3.0	31.9	17.7-58.4	2.0	43.8	20.3-93.0	1.0	21.9	7.0-66.6	1.0	11.6	3.7-33.8	0.8	17.7	4.8-59.7	0.2	5.3	0.1-46.7
	2011-2015	2.4	36.0	15.8-72.2	0.6	15.2	2.9-60.7	1.8	58.1	21.1-129.6	1.0	11.6	3.7-34.9	0.6	13.8	2.8-57.9	0.4	9.4	1.1-53.3
2016-2020	3.2	37.3	19.2-71.5	1.4	25.6	9.9-77.1	1.8	48.2	18.6-113.3	1.2	13.3	4.8-39.7	0.4	7.3	0.9-56.3	0.8	18.9	5.0-70.5	
Rectum & rectosigmoid	1976-1980	1.2	17.5	5.4-43.0	0.6	13.7	2.8-51.9	0.6	19.3	3.2-63.0	0.8	14.1	3.2-39.6	0.4	9.3	1.1-46.3	0.4	16.5	1.8-60.7
	1981-1985	2.0	30.8	14.0-60.1	1.4	49.7	17.4-111.6	0.6	17.9	3.4-58.8	0.6	8.3	1.7-28.1	0.4	12.8	1.5-53.6	0.2	4.6	0.1-37.3
	1986-1990	1.8	25.9	11.7-51.6	1.2	34.0	12.0-80.9	0.6	18.2	3.7-57.8	0.4	6.0	0.7-24.8	0.2	4.5	0.1-37.8	0.2	6.2	0.2-40.0
	1991-1995	1.0	14.7	4.6-36.7	0.8	23.7	6.0-66.4	0.2	7.3	0.2-42.7	0.6	10.4	2.1-31.5	0.4	18.3	2.1-64.2	0.2	7.1	0.2-42.1
	1996-2000	1.6	19.8	8.5-42.3	1.0	25.3	8.2-66.1	0.6	13.9	2.9-51.0	0.0	0.0	0.0-15.4	0.0	0.0	0.0-30.6	0.0	0.0	0.0-32.4
	2001-2005	1.8	22.5	10.2-46.6	1.4	36.6	14.5-83.3	0.4	9.4	1.1-46.6	0.8	8.9	2.4-28.3	0.6	14.0	2.8-52.2	0.2	4.4	0.1-39.9
	2006-2010	0.4	5.1	0.6-25.7	0.4	10.6	1.3-51.6	0.0	0.0	0.0-40.1	0.0	0.0	0.0-19.2	0.0	0.0	0.0-37.6	0.0	0.0	0.0-40.1
	2011-2015	1.0	11.2	3.5-34.3	0.6	12.0	2.4-55.0	0.4	10.3	1.3-54.9	0.8	8.0	2.2-30.0	0.6	12.0	2.4-55.0	0.2	4.0	0.1-46.4
2016-2020	1.0	11.4	3.4-38.1	0.6	10.7	2.1-60.0	0.4	12.7	1.4-65.6	0.2	1.4	0.0-25.7	0.2	2.7	0.1-51.6	0.0	0.0	0.0-49.0	
Liver / intrahepatic bile duct	1981-1985	0.2	5.6	0.1-27.8	0.2	15.5	0.4-70.3	0.0	0.0	0.0-30.9	0.2	5.6	0.1-27.8	0.2	15.5	0.4-70.3	0.0	0.0	0.0-30.9
	1986-1990	0.4	5.1	0.6-22.5	0.2	5.0	0.1-38.9	0.2	5.2	0.1-37.7	0.4	4.6	0.6-21.6	0.2	5.0	0.1-38.9	0.2	4.0	0.1-35.3
	1991-1995	0.4	4.4	0.5-20.8	0.4	8.6	1.0-41.4	0.0	0.0	0.0-29.6	0.4	4.6	0.6-21.0	0.4	9.0	1.1-42.3	0.0	0.0	0.0-29.6
	2001-2005	0.2	2.1	0.1-19.3	0.2	4.0	0.1-38.2	0.0	0.0	0.0-34.4	0.4	5.3	0.6-24.1	0.2	5.3	0.1-40.9	0.2	5.6	0.1-42.2
	2006-2010	0.8	9.2	2.4-30.7	0.6	13.0	2.5-53.9	0.2	5.2	0.1-46.6	0.4	4.7	0.5-25.0	0.4	9.5	1.0-49.9	0.0	0.0	0.0-40.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Hamilton**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	2011-2015	0.4	4.7	0.6-26.4	0.4	9.5	1.1-52.8	0.0	0.0	0.0-42.3	0.2	1.6	0.0-22.3	0.2	3.1	0.1-44.2	0.0	0.0	0.0-42.3
	2016-2020	0.6	5.2	1.0-29.6	0.6	10.0	2.0-59.0	0.0	0.0	0.0-49.0	0.2	1.6	0.0-25.9	0.0	0.0	0.0-49.3	0.2	3.1	0.1-51.7
Pancreas	1976-1980	0.8	16.3	4.1-43.0	0.0	0.0	0.0-33.9	0.8	30.8	7.8-81.6	0.6	14.2	2.9-40.6	0.0	0.0	0.0-33.9	0.6	26.6	5.4-76.8
	1981-1985	1.2	19.2	6.8-44.6	0.6	24.5	4.7-74.5	0.6	15.8	3.1-54.1	1.0	13.9	4.2-36.2	0.6	21.2	3.7-68.8	0.4	8.9	1.1-42.9
	1986-1990	1.2	15.1	5.4-36.4	0.4	9.5	1.1-44.9	0.8	19.0	5.0-56.9	1.2	15.1	5.4-36.4	0.4	9.5	1.1-44.9	0.8	19.0	5.0-56.9
	1991-1995	1.2	18.1	6.5-41.3	1.0	37.7	11.5-90.6	0.2	5.2	0.1-37.4	1.0	15.0	4.8-37.1	0.8	30.2	7.5-80.1	0.2	5.2	0.1-37.4
	1996-2000	1.4	20.3	8.0-44.7	0.8	25.3	6.5-69.9	0.6	15.0	3.0-53.3	1.0	13.1	4.2-34.0	0.4	12.5	1.5-50.2	0.6	14.3	2.9-51.7
	2001-2005	0.6	6.9	1.4-26.0	0.0	0.0	0.0-33.5	0.6	12.1	2.5-49.4	1.0	14.3	4.3-37.6	0.8	25.0	6.3-71.1	0.2	4.1	0.1-39.2
	2006-2010	0.6	8.0	1.6-29.7	0.2	5.9	0.2-45.9	0.4	10.0	1.2-52.4	0.4	5.4	0.7-26.3	0.4	12.5	1.5-55.5	0.0	0.0	0.0-40.1
	2011-2015	0.8	8.5	2.3-30.8	0.2	3.2	0.1-44.4	0.6	13.3	2.7-57.8	1.0	11.1	3.6-34.2	0.4	8.9	1.1-51.6	0.6	13.3	2.7-57.8
2016-2020	2.4	23.3	11.9-50.9	2.0	38.2	18.0-91.8	0.4	7.8	0.9-57.0	0.8	8.0	2.2-33.0	0.6	12.0	2.4-61.8	0.2	4.7	0.1-54.0	
Larynx	1976-1980	0.6	11.1	2.1-34.2	0.6	24.7	4.5-76.0	0.0	0.0	0.0-30.6	0.0	0.0	0.0-15.9	0.0	0.0	0.0-33.9	0.0	0.0	0.0-30.6
	1981-1985	0.4	5.1	0.6-23.2	0.4	10.4	1.3-48.3	0.0	0.0	0.0-30.9	0.2	2.1	0.1-18.6	0.2	4.2	0.1-39.0	0.0	0.0	0.0-30.9
	1991-1995	0.0	0.0	0.0-14.6	0.0	0.0	0.0-29.7	0.0	0.0	0.0-29.6	0.2	3.0	0.1-19.6	0.2	7.3	0.2-42.6	0.0	0.0	0.0-29.6
	1996-2000	0.6	6.8	1.4-24.7	0.6	12.8	2.6-47.6	0.0	0.0	0.0-32.4	0.2	2.5	0.1-19.0	0.2	4.8	0.1-37.3	0.0	0.0	0.0-32.4
	2001-2005	0.2	2.5	0.1-20.1	0.2	5.3	0.1-40.9	0.0	0.0	0.0-34.4	0.4	4.1	0.5-21.8	0.4	8.5	1.0-44.2	0.0	0.0	0.0-34.4
	2006-2010	0.6	8.3	1.6-30.6	0.4	9.5	1.0-49.8	0.2	7.5	0.2-51.4	0.2	2.8	0.1-22.9	0.2	5.9	0.2-45.9	0.0	0.0	0.0-40.1
Lung and bronchus	1976-1980	6.0	96.9	63.3-142.6	4.0	144.0	83.0-232.6	2.0	56.8	25.9-111.2	3.8	60.4	34.7-98.6	2.0	67.8	29.2-134.7	1.8	53.6	23.2-108.2
	1981-1985	4.4	64.4	39.4-101.0	2.6	81.6	41.6-147.2	1.8	50.9	22.4-103.1	5.2	82.2	52.5-123.7	3.6	124.5	71.5-203.7	1.6	45.9	19.0-96.9
	1986-1990	3.8	53.1	31.2-86.0	2.6	80.5	39.9-146.7	1.2	34.2	12.0-80.3	4.6	69.1	42.9-106.7	3.2	99.4	53.6-170.2	1.4	45.4	17.6-97.9
	1991-1995	3.8	56.0	33.4-89.3	1.6	55.1	22.8-112.5	2.2	64.8	32.0-120.0	3.6	49.3	28.9-80.3	2.4	69.7	35.1-127.1	1.2	33.5	12.0-78.1
	1996-2000	6.4	79.5	54.2-114.9	3.4	94.3	53.9-156.8	3.0	70.8	39.5-123.6	4.4	54.1	33.8-84.7	1.8	52.3	23.3-104.7	2.6	60.0	31.9-109.8
	2001-2005	9.2	115.2	84.1-156.7	5.2	147.4	94.4-223.5	4.0	94.4	57.4-153.4	5.8	74.0	49.3-109.4	3.2	101.3	56.4-170.8	2.6	61.3	32.4-113.3
	2006-2010	5.6	68.5	45.2-103.5	3.0	71.3	39.3-128.0	2.6	65.2	34.4-122.4	4.6	54.1	34.1-85.9	2.6	61.7	32.2-116.1	2.0	47.2	22.5-98.8
	2011-2015	7.8	83.1	58.0-120.1	4.0	76.0	46.0-131.1	3.8	90.0	52.0-154.9	5.0	53.4	34.2-85.0	2.8	58.8	31.8-112.3	2.2	48.0	23.5-100.7
2016-2020	6.6	65.6	44.5-100.5	3.8	71.7	42.6-130.9	2.8	58.6	31.0-117.8	4.0	43.5	25.4-76.7	3.0	61.0	33.8-119.3	1.0	24.4	5.8-81.5	
Melanoma of the skin	1976-1980	0.6	15.5	3.2-43.4	0.0	0.0	0.0-33.9	0.6	29.7	6.1-83.4	0.2	5.3	0.1-26.7	0.0	0.0	0.0-33.9	0.2	10.6	0.3-52.8
	1981-1985	0.4	10.2	1.2-35.1	0.2	9.2	0.2-51.0	0.2	11.0	0.3-54.4	0.2	4.3	0.1-24.1	0.0	0.0	0.0-33.9	0.2	8.3	0.2-46.3
	1986-1990	0.6	10.2	1.9-31.9	0.4	12.7	1.3-52.6	0.2	6.2	0.2-40.0	0.6	9.4	1.9-29.8	0.2	5.0	0.1-38.9	0.4	12.3	1.5-49.5
	1991-1995	1.0	11.9	3.8-31.2	0.6	14.9	3.0-51.3	0.4	8.8	1.1-41.7	0.4	4.3	0.5-20.5	0.2	4.1	0.1-35.1	0.2	4.6	0.1-36.1
	1996-2000	2.4	32.8	16.6-60.1	2.0	57.8	26.8-112.2	0.4	10.6	1.2-47.6	0.6	6.7	1.4-24.4	0.4	9.1	1.1-43.1	0.2	4.7	0.1-38.8
	2001-2005	1.8	23.4	10.7-48.1	1.2	35.6	12.8-84.2	0.6	14.4	2.9-53.6	0.6	6.3	1.3-24.7	0.6	12.7	2.6-49.7	0.0	0.0	0.0-34.4
	2006-2010	1.6	23.7	8.7-53.8	1.0	36.1	8.4-98.1	0.6	13.1	2.7-55.5	0.2	2.8	0.1-22.9	0.0	0.0	0.0-37.6	0.2	5.3	0.1-46.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Hamilton**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Melanoma of the skin	2011-2015	2.6	29.8	14.9-58.5	1.8	39.6	17.1-90.8	0.8	19.8	4.3-68.9	0.6	5.3	1.1-26.5	0.6	10.4	2.1-52.4	0.0	0.0	0.0-42.3
	2016-2020	1.6	18.4	6.7-47.9	1.4	35.4	11.9-95.1	0.2	2.8	0.1-51.4	0.0	0.0	0.0-24.5	0.0	0.0	0.0-49.3	0.0	0.0	0.0-49.0
Female breast	1976-1980							2.4	73.0	36.0-133.9							1.4	43.3	17.0-94.0
	1981-1985							4.2	130.9	79.1-206.1							1.0	30.2	9.4-76.1
	1986-1990							2.8	84.2	45.4-145.7							1.4	40.1	16.0-87.6
	1991-1995							4.2	121.8	74.4-190.9							1.0	28.4	9.1-71.3
	1996-2000							5.2	133.7	86.9-201.8							1.8	43.6	19.8-90.2
	2001-2005							6.0	139.2	93.5-206.1							1.0	24.8	7.9-67.7
	2006-2010							5.2	127.2	79.5-200.2							0.8	16.2	4.4-59.0
	2011-2015							7.8	181.1	122.3-265.8							1.0	19.4	6.3-64.1
2016-2020							4.8	132.7	74.7-224.4							0.8	29.9	4.9-98.4	
Cervix uteri	1976-1980							0.2	4.2	0.1-36.2							0.2	7.5	0.2-44.0
	1981-1985							0.0	0.0	0.0-30.9							0.2	6.9	0.2-42.7
	1986-1990							0.2	6.2	0.2-40.0							0.0	0.0	0.0-30.0
	1991-1995							0.2	8.2	0.2-45.0							0.4	13.4	1.5-52.1
	2001-2005							0.2	4.1	0.1-39.2							0.2	4.1	0.1-39.2
	2006-2010							0.6	11.3	2.3-52.6							0.0	0.0	0.0-40.1
	2011-2015							0.2	5.2	0.1-48.5							0.0	0.0	0.0-42.3
	2016-2020							0.2	2.8	0.1-51.4							0.0	0.0	0.0-49.0
Corpus uteris and uterus, not otherwise specified	1976-1980							0.8	23.2	5.9-66.0							0.0	0.0	0.0-30.6
	1981-1985							1.6	65.0	27.0-130.6							0.0	0.0	0.0-30.9
	1986-1990							0.6	17.2	3.5-55.9							0.4	10.6	1.3-45.8
	1991-1995							1.4	42.9	16.8-92.6							0.6	14.0	2.9-49.3
	1996-2000							1.4	33.2	13.3-76.6							0.2	5.2	0.1-39.8
	2001-2005							1.0	22.4	7.3-63.1							0.0	0.0	0.0-34.4
	2006-2010							0.8	18.6	5.0-63.0							0.2	3.6	0.1-43.8
	2011-2015							1.6	43.0	16.0-101.3							0.4	9.1	1.1-52.9
2016-2020							2.0	49.5	17.9-117.8							0.4	6.3	0.7-54.7	
Ovary	1976-1980							0.2	4.5	0.1-36.7							0.4	12.0	1.3-49.7
	1981-1985							0.4	12.9	1.4-52.2							0.4	11.8	1.4-49.0
	1986-1990							0.8	19.8	5.3-58.3							0.6	12.1	2.5-46.0
	1991-1995							0.8	19.0	5.1-56.2							0.8	19.3	5.2-56.7
	1996-2000							0.4	9.4	1.1-45.0							0.4	9.7	1.2-45.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Hamilton**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Ovary	2001-2005							1.0	22.2	7.1-62.9							0.8	18.5	5.0-58.6
	2006-2010							0.6	13.2	2.7-55.8							0.2	4.3	0.1-45.0
	2011-2015							0.6	12.6	2.5-56.8							0.0	0.0	0.0-42.3
	2016-2020							0.2	3.5	0.1-52.2							0.2	4.7	0.1-54.0
Prostate	1976-1980				1.8	78.1	33.6-152.9							1.4	70.9	26.5-148.3			
	1981-1985				2.8	117.1	60.5-203.3							0.8	37.9	9.8-97.5			
	1986-1990				3.2	122.8	66.7-206.4							0.6	24.0	4.6-72.5			
	1991-1995				3.6	104.8	60.1-172.0							2.4	85.6	42.7-153.5			
	1996-2000				6.6	169.8	115.6-244.4							0.6	16.0	3.3-53.9			
	2001-2005				9.8	234.5	172.6-316.9							0.6	18.3	3.5-61.2			
	2006-2010				5.4	119.1	77.7-183.2							1.8	48.0	21.3-101.4			
	2011-2015				3.4	68.5	38.8-124.3							0.4	8.1	0.9-50.4			
2016-2020				5.8	114.6	75.1-182.4							1.0	22.3	7.1-74.3				
Testis	1976-1980				0.2	9.3	0.2-51.3							0.0	0.0	0.0-33.9			
	1981-1985				0.2	7.8	0.2-47.3							0.0	0.0	0.0-33.9			
	1986-1990				0.2	8.5	0.2-47.7							0.0	0.0	0.0-31.9			
	1996-2000				0.6	24.6	3.9-76.0							0.0	0.0	0.0-30.6			
	2001-2005				0.2	16.2	0.4-72.1							0.0	0.0	0.0-33.5			
	2011-2015				0.2	4.0	0.1-45.6							0.0	0.0	0.0-41.4			
Urinary bladder (incl. in situ)	1976-1980	0.8	14.6	3.7-39.5	0.8	30.1	7.7-81.6	0.0	0.0	0.0-30.6	1.0	20.1	6.1-48.3	0.8	34.7	8.3-92.8	0.2	8.5	0.2-46.8
	1981-1985	1.8	27.9	11.7-57.2	1.8	65.4	25.2-136.4	0.0	0.0	0.0-30.9	0.8	15.6	4.0-41.5	0.8	38.0	8.9-100.5	0.0	0.0	0.0-30.9
	1986-1990	1.6	28.3	11.8-57.4	1.2	52.0	17.7-116.1	0.4	13.0	1.4-51.5	0.4	7.5	0.9-27.8	0.4	19.0	2.3-66.4	0.0	0.0	0.0-30.0
	1991-1995	1.0	12.9	4.1-33.1	0.8	23.6	6.0-66.1	0.2	4.6	0.1-36.1	0.2	2.2	0.1-17.5	0.0	0.0	0.0-29.7	0.2	4.6	0.1-36.1
	1996-2000	1.8	22.2	10.1-45.6	1.8	48.7	21.4-99.2	0.0	0.0	0.0-32.4	0.6	7.7	1.6-26.3	0.6	17.0	3.5-55.8	0.0	0.0	0.0-32.4
	2001-2005	2.6	30.7	16.2-56.5	2.2	55.5	27.0-107.5	0.4	8.9	1.0-45.8	0.2	2.0	0.1-19.2	0.0	0.0	0.0-33.5	0.2	3.6	0.1-38.5
	2006-2010	2.0	26.3	12.3-53.6	1.8	51.9	23.2-107.6	0.2	4.0	0.1-44.5	0.8	9.7	2.6-31.6	0.8	23.4	6.1-70.2	0.0	0.0	0.0-40.1
	2011-2015	2.6	26.4	13.9-52.5	1.8	40.0	17.9-90.4	0.8	14.0	3.8-57.3	0.4	7.0	0.8-31.2	0.4	13.8	1.6-61.2	0.0	0.0	0.0-42.3
2016-2020	4.0	42.2	25.2-74.1	3.6	76.9	44.5-139.7	0.4	8.0	0.9-57.2	0.4	4.2	0.5-29.0	0.2	5.4	0.1-55.4	0.2	3.1	0.1-51.7	
Kidney and renal pelvis	1976-1980	0.4	6.4	0.8-25.8	0.2	6.8	0.2-44.8	0.2	6.0	0.2-40.1	0.4	6.0	0.7-25.0	0.2	6.8	0.2-44.8	0.2	5.3	0.1-38.4
	1981-1985	0.4	8.7	0.9-31.9	0.2	15.5	0.4-70.3	0.2	6.1	0.2-40.6	0.0	0.0	0.0-15.9	0.0	0.0	0.0-33.9	0.0	0.0	0.0-30.9
	1986-1990	0.8	12.7	3.3-34.9	0.4	12.8	1.3-52.9	0.4	12.0	1.4-48.7	0.6	12.4	2.5-36.2	0.2	13.7	0.3-63.4	0.4	12.4	1.5-49.7
	1991-1995	0.0	0.0	0.0-14.6	0.0	0.0	0.0-29.7	0.0	0.0	0.0-29.6	0.6	9.3	1.8-29.5	0.6	17.9	3.4-57.9	0.0	0.0	0.0-29.6
	1996-2000	1.2	14.3	5.2-35.0	0.8	19.2	5.1-57.3	0.4	9.1	1.1-44.4	0.4	4.8	0.6-22.2	0.2	5.7	0.1-39.5	0.2	4.4	0.1-38.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Hamilton**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Kidney and renal pelvis	2001-2005	1.8	23.8	10.1-50.0	1.4	39.0	14.0-90.3	0.4	9.1	1.1-46.1	0.4	4.2	0.5-22.0	0.4	7.9	1.0-43.2	0.0	0.0	0.0-34.4
	2006-2010	1.4	16.3	6.4-39.7	1.0	23.1	7.3-67.2	0.4	9.3	1.1-51.4	0.2	2.2	0.1-21.8	0.2	4.3	0.1-42.7	0.0	0.0	0.0-40.1
	2011-2015	1.4	15.6	6.1-39.9	0.6	14.8	2.9-59.9	0.8	16.4	4.4-61.1	0.2	2.7	0.1-24.1	0.0	0.0	0.0-41.4	0.2	5.2	0.1-48.5
	2016-2020	1.4	26.7	7.7-66.7	0.8	29.3	5.1-96.1	0.6	24.0	2.3-91.9	0.6	4.8	1.0-29.1	0.2	2.7	0.1-51.6	0.4	6.4	0.8-54.9
Brain and other central nervous system	1976-1980	0.8	12.8	3.3-35.5	0.4	12.7	1.5-53.3	0.4	13.3	1.4-53.1	0.4	5.5	0.6-24.0	0.2	6.8	0.2-44.8	0.2	4.5	0.1-36.7
	1986-1990	0.2	2.3	0.1-18.4	0.0	0.0	0.0-31.9	0.2	4.8	0.1-36.9	0.0	0.0	0.0-15.2	0.0	0.0	0.0-31.9	0.0	0.0	0.0-30.0
	1991-1995	1.0	16.9	5.0-41.8	0.2	10.5	0.3-52.0	0.8	21.6	5.7-61.4	0.6	9.0	1.8-28.7	0.0	0.0	0.0-29.7	0.6	17.0	3.4-55.4
	1996-2000	1.2	14.0	5.1-34.3	0.4	9.7	1.1-44.5	0.8	18.3	5.0-56.6	0.6	7.9	1.6-26.8	0.2	6.0	0.2-40.1	0.4	9.4	1.1-45.0
	2001-2005	0.6	7.7	1.6-27.3	0.2	5.6	0.1-41.6	0.4	10.0	1.2-47.9	0.2	2.2	0.1-19.5	0.2	4.3	0.1-38.8	0.0	0.0	0.0-34.4
	2006-2010	0.6	7.3	1.4-28.5	0.2	5.9	0.2-45.9	0.4	8.8	1.0-50.5	0.4	4.7	0.5-25.1	0.2	5.9	0.2-45.9	0.2	4.1	0.1-44.6
	2011-2015	1.0	15.9	3.9-45.1	0.6	13.6	2.7-57.6	0.4	18.8	1.4-77.2	0.6	5.7	1.2-27.0	0.2	4.1	0.1-45.7	0.4	7.5	0.9-50.2
	2016-2020	0.0	0.0	0.0-24.5	0.0	0.0	0.0-49.3	0.0	0.0	0.0-49.0	0.4	3.8	0.4-28.3	0.2	4.1	0.1-53.3	0.2	3.1	0.1-51.7
Thyroid	1976-1980	0.2	3.9	0.1-22.8	0.0	0.0	0.0-33.9	0.2	7.5	0.2-44.0	0.0	0.0	0.0-15.9	0.0	0.0	0.0-33.9	0.0	0.0	0.0-30.6
	1981-1985	0.4	8.8	1.1-31.6	0.0	0.0	0.0-33.9	0.4	16.2	1.9-59.5	0.2	4.2	0.1-23.7	0.0	0.0	0.0-33.9	0.2	6.9	0.2-42.7
	1996-2000	0.6	8.3	1.7-27.6	0.4	10.6	1.3-46.0	0.2	6.1	0.2-41.9	0.0	0.0	0.0-15.4	0.0	0.0	0.0-30.6	0.0	0.0	0.0-32.4
	2001-2005	0.6	7.4	1.5-26.7	0.2	5.6	0.1-41.6	0.4	9.1	1.1-46.2	0.2	2.5	0.1-20.1	0.0	0.0	0.0-33.5	0.2	4.7	0.1-40.5
	2006-2010	0.6	14.0	2.6-42.7	0.2	11.4	0.3-60.0	0.4	17.0	1.6-68.8	0.0	0.0	0.0-19.2	0.0	0.0	0.0-37.6	0.0	0.0	0.0-40.1
	2011-2015	1.4	16.4	5.1-43.7	0.6	10.3	2.1-52.2	0.8	22.7	4.0-77.1	0.0	0.0	0.0-20.8	0.0	0.0	0.0-41.4	0.0	0.0	0.0-42.3
	2016-2020	0.4	6.2	0.7-32.4	0.0	0.0	0.0-49.3	0.4	12.6	1.4-65.3	0.0	0.0	0.0-24.5	0.0	0.0	0.0-49.3	0.0	0.0	0.0-49.0
Hodgkin lymphoma	1976-1980	0.2	2.8	0.1-20.1	0.2	5.9	0.2-42.8	0.0	0.0	0.0-30.6	0.0	0.0	0.0-15.9	0.0	0.0	0.0-33.9	0.0	0.0	0.0-30.6
	1991-1995	0.2	2.3	0.1-17.8	0.0	0.0	0.0-29.7	0.2	4.6	0.1-36.1	0.2	2.3	0.1-17.8	0.0	0.0	0.0-29.7	0.2	4.6	0.1-36.1
	1996-2000	0.2	4.8	0.1-25.1	0.2	9.2	0.2-48.7	0.0	0.0	0.0-32.4	0.0	0.0	0.0-15.4	0.0	0.0	0.0-30.6	0.0	0.0	0.0-32.4
	2001-2005	0.4	11.1	1.3-38.0	0.2	9.1	0.2-50.3	0.2	15.8	0.4-71.5	0.0	0.0	0.0-16.7	0.0	0.0	0.0-33.5	0.0	0.0	0.0-34.4
	2006-2010	0.2	2.8	0.1-22.9	0.2	5.9	0.2-45.9	0.0	0.0	0.0-40.1	0.4	10.8	1.3-38.3	0.4	20.0	2.4-72.2	0.0	0.0	0.0-40.1
	2016-2020	0.2	1.6	0.0-25.9	0.2	3.2	0.1-52.1	0.0	0.0	0.0-49.0	0.2	1.6	0.0-25.9	0.2	3.2	0.1-52.1	0.0	0.0	0.0-49.0
Non-Hodgkin lymphomas	1976-1980	1.6	24.2	9.9-51.1	0.2	4.3	0.1-39.4	1.4	42.5	16.0-93.9	0.4	4.9	0.6-22.8	0.2	4.3	0.1-39.4	0.2	5.3	0.1-38.4
	1981-1985	0.4	8.5	0.9-31.3	0.4	17.7	2.1-65.0	0.0	0.0	0.0-30.9	0.2	5.8	0.1-28.4	0.0	0.0	0.0-33.9	0.2	11.0	0.3-54.4
	1986-1990	0.8	9.0	2.4-27.5	0.2	5.0	0.1-38.9	0.6	12.9	2.6-47.4	0.2	2.5	0.1-18.9	0.0	0.0	0.0-31.9	0.2	5.2	0.1-37.7
	1991-1995	2.4	36.0	18.4-64.9	1.8	61.9	27.9-120.2	0.6	14.0	2.9-49.3	0.6	8.3	1.7-27.1	0.4	11.3	1.2-47.7	0.2	5.2	0.1-37.4
	1996-2000	1.0	12.7	4.1-33.3	0.4	9.1	1.1-43.1	0.6	16.5	3.3-56.2	0.8	12.1	3.0-34.2	0.4	10.6	1.3-46.0	0.4	13.4	1.4-54.4
	2001-2005	1.2	13.8	5.0-34.9	0.8	19.4	5.2-59.8	0.4	8.5	1.0-44.9	0.0	0.0	0.0-16.7	0.0	0.0	0.0-33.5	0.0	0.0	0.0-34.4
	2006-2010	2.2	26.3	13.0-52.5	1.6	38.5	16.4-87.2	0.6	13.7	2.8-56.7	0.6	6.8	1.3-27.6	0.4	7.9	0.9-46.7	0.2	5.3	0.1-46.7
	2011-2015	2.2	22.6	11.2-47.9	1.2	24.0	8.6-69.4	1.0	21.6	6.9-67.8	0.6	6.4	1.3-28.2	0.2	4.9	0.1-47.2	0.4	7.9	1.0-50.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Hamilton**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Non-Hodgkin lymphomas	2016-2020	2.6	25.9	13.4-54.5	0.8	15.3	4.1-65.3	1.8	37.8	16.3-93.1	0.8	7.8	2.0-32.8	0.4	7.5	0.8-56.8	0.4	8.7	1.0-58.4
Myeloma	1981-1985	0.4	8.3	0.8-31.2	0.2	15.5	0.4-70.3	0.2	5.4	0.1-39.0	0.4	8.3	0.8-31.2	0.2	15.5	0.4-70.3	0.2	5.4	0.1-39.0
	1991-1995	0.4	4.3	0.5-20.4	0.4	8.8	1.1-41.7	0.0	0.0	0.0-29.6	0.2	2.1	0.1-17.5	0.2	4.4	0.1-35.7	0.0	0.0	0.0-29.6
	1996-2000	1.0	12.7	4.1-33.2	0.2	4.0	0.1-35.7	0.8	18.8	5.1-57.6	0.4	5.4	0.6-23.5	0.4	10.6	1.3-46.3	0.0	0.0	0.0-32.4
	2001-2005	0.6	7.1	1.5-26.3	0.4	10.6	1.3-48.3	0.2	4.4	0.1-39.9	0.2	2.0	0.1-19.2	0.0	0.0	0.0-33.5	0.2	3.6	0.1-38.5
	2006-2010	1.2	15.1	5.5-38.7	0.6	15.9	3.2-58.9	0.6	15.2	3.1-59.2	0.4	4.8	0.6-25.2	0.2	4.3	0.1-42.7	0.2	4.3	0.1-45.0
	2011-2015	1.0	13.0	3.8-37.9	0.2	4.3	0.1-46.1	0.8	21.7	5.0-71.7	0.6	6.9	1.4-29.0	0.4	9.8	1.2-53.3	0.2	4.0	0.1-46.4
	2016-2020	0.6	5.1	1.0-29.5	0.2	2.7	0.1-51.6	0.4	7.4	0.8-56.3	0.0	0.0	0.0-24.5	0.0	0.0	0.0-49.3	0.0	0.0	0.0-49.0
Leukemias	1976-1980	0.6	11.5	1.8-36.2	0.4	20.8	1.6-77.4	0.2	5.3	0.1-38.4	0.6	11.0	1.7-35.4	0.6	25.5	3.4-82.5	0.0	0.0	0.0-30.6
	1981-1985	0.4	5.9	0.7-24.9	0.4	12.8	1.5-53.6	0.0	0.0	0.0-30.9	0.8	18.3	4.9-46.4	0.4	26.3	3.0-86.8	0.4	13.8	1.7-53.6
	1986-1990	1.2	20.1	7.0-46.1	0.8	29.3	7.4-79.4	0.4	13.9	1.4-54.0	0.8	10.8	2.6-31.4	0.6	16.7	3.1-57.2	0.2	4.8	0.1-36.9
	1991-1995	0.8	11.0	2.9-31.1	0.6	22.7	4.0-69.4	0.2	4.2	0.1-35.2	0.4	5.2	0.6-22.4	0.2	7.3	0.2-42.6	0.2	4.2	0.1-35.2
	1996-2000	0.6	8.8	1.8-28.7	0.2	6.0	0.2-40.1	0.4	9.7	1.2-45.6	0.2	3.2	0.1-20.5	0.0	0.0	0.0-30.6	0.2	5.1	0.1-39.5
	2001-2005	0.6	7.4	1.5-26.9	0.6	15.8	3.2-55.4	0.0	0.0	0.0-34.4	0.4	4.7	0.6-23.1	0.4	10.1	1.2-47.4	0.0	0.0	0.0-34.4
	2006-2010	0.4	5.2	0.6-25.9	0.2	4.9	0.1-43.9	0.2	5.3	0.1-46.7	0.4	5.2	0.6-25.9	0.2	5.9	0.2-45.9	0.2	4.7	0.1-45.6
	2011-2015	2.8	37.6	19.0-70.5	2.2	64.0	29.3-128.2	0.6	12.8	2.6-57.0	1.0	15.9	3.9-45.1	0.8	28.0	5.8-85.3	0.2	4.4	0.1-47.2
	2016-2020	1.0	17.1	3.5-51.8	0.8	30.3	5.1-98.9	0.2	3.1	0.1-51.7	0.6	6.1	1.2-30.9	0.4	7.4	0.9-56.5	0.2	4.7	0.1-54.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Herkimer**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	274.4	396.9	375.6-419.2	133.4	457.6	422.0-495.3	141.0	365.6	338.2-394.7	140.6	203.4	188.3-219.4	71.4	253.5	226.7-282.4	69.2	172.9	154.7-192.6
	1981-1985	303.0	413.3	392.2-435.3	147.6	473.8	438.7-511.0	155.4	386.9	359.0-416.4	160.2	213.5	198.7-229.3	86.8	282.0	254.8-311.2	73.4	169.5	152.1-188.3
	1986-1990	307.4	406.1	385.5-427.5	153.8	478.9	444.4-515.3	153.6	367.3	340.6-395.6	164.4	212.1	197.5-227.5	84.4	264.7	239.1-292.3	80.0	179.5	161.8-198.8
	1991-1995	347.8	440.7	419.8-462.3	181.4	533.7	498.7-570.5	166.4	384.6	357.9-412.8	169.8	210.6	196.5-225.6	91.8	282.4	256.4-310.3	78.0	170.5	153.4-189.0
	1996-2000	400.6	506.4	484.3-529.4	197.4	573.3	537.8-610.6	203.2	465.2	436.2-495.7	175.6	214.8	200.7-229.6	88.4	259.8	235.9-285.5	87.2	185.9	168.4-204.8
	2001-2005	391.6	490.0	468.2-512.5	206.2	581.7	546.5-618.6	185.4	427.6	399.6-457.1	155.2	184.6	171.7-198.3	76.0	215.2	193.9-238.2	79.2	167.2	150.7-185.3
	2006-2010	408.6	497.6	475.8-520.2	208.4	546.3	513.2-581.2	200.2	465.3	435.9-496.3	136.8	159.5	147.6-172.2	68.2	184.3	165.0-205.2	68.6	143.9	128.7-160.7
	2011-2015	426.0	499.8	477.9-522.4	218.8	542.8	510.2-577.2	207.2	467.6	438.0-498.8	147.6	165.0	153.0-177.7	76.6	194.3	175.0-215.4	71.0	144.9	129.6-161.8
	2016-2020	426.6	479.5	458.3-501.6	228.0	526.1	494.6-559.2	198.6	440.7	411.8-471.3	134.4	143.1	132.2-154.9	73.8	172.4	154.7-191.7	60.6	118.7	105.2-133.7
Oral cavity and pharynx	1976-1980	8.2	11.7	8.3-16.0	5.6	19.2	12.5-28.2	2.6	6.1	3.2-10.6	3.2	4.4	2.5-7.3	2.0	6.8	3.1-12.8	1.2	2.7	1.0-6.2
	1981-1985	9.0	11.4	8.3-15.5	6.6	19.7	13.3-28.0	2.4	5.1	2.6-9.2	4.2	5.4	3.3-8.4	2.8	7.9	4.2-13.5	1.4	3.1	1.3-6.7
	1986-1990	7.4	9.9	6.9-13.8	4.4	13.9	8.6-21.3	3.0	6.9	3.8-11.8	3.2	4.1	2.3-6.8	1.4	4.5	1.7-9.6	1.8	3.7	1.7-7.4
	1991-1995	5.8	7.7	5.1-11.2	4.8	14.3	9.0-21.6	1.0	2.5	0.8-6.1	2.2	2.9	1.4-5.3	1.6	5.0	2.1-10.0	0.6	1.7	0.3-5.1
	1996-2000	6.4	8.5	5.8-12.1	4.2	12.1	7.5-18.6	2.2	5.7	2.8-10.4	2.2	2.9	1.4-5.3	1.8	5.4	2.5-10.4	0.4	0.8	0.1-3.4
	2001-2005	7.0	9.2	6.4-13.0	4.2	12.2	7.5-18.8	2.8	7.1	3.8-12.3	1.6	2.0	0.8-4.0	1.2	3.7	1.3-8.1	0.4	1.1	0.1-4.0
	2006-2010	10.6	13.0	9.7-17.2	7.8	20.5	14.5-28.3	2.8	6.7	3.5-11.7	1.8	2.0	0.9-4.0	1.6	4.0	1.7-8.2	0.2	0.5	0.0-3.2
	2011-2015	12.8	15.3	11.7-19.8	8.6	22.0	15.7-30.1	4.2	9.4	5.8-14.8	1.6	1.9	0.8-4.0	1.2	3.1	1.1-7.1	0.4	1.0	0.1-4.1
	2016-2020	12.0	13.1	9.8-17.2	7.6	17.0	11.8-23.8	4.4	10.1	6.1-15.9	3.2	3.2	1.8-5.4	2.0	4.3	2.0-8.5	1.2	2.5	0.9-6.0
Esophagus	1976-1980	3.2	4.6	2.6-7.6	3.0	10.6	5.8-17.7	0.2	0.4	0.0-2.9	2.0	2.9	1.4-5.4	2.0	7.3	3.3-13.5	0.0	0.0	0.0-2.2
	1981-1985	2.6	3.6	1.9-6.2	1.8	6.2	2.7-11.9	0.8	1.7	0.5-4.8	2.8	4.0	2.2-6.8	2.2	8.3	4.1-15.0	0.6	1.2	0.2-4.0
	1986-1990	1.8	2.3	1.0-4.5	1.6	4.6	2.0-9.3	0.2	0.4	0.0-2.7	1.8	2.2	1.0-4.3	1.4	4.2	1.6-8.8	0.4	0.8	0.1-3.3
	1991-1995	2.4	3.0	1.5-5.3	1.4	4.2	1.6-8.9	1.0	2.4	0.8-5.9	3.8	4.5	2.7-7.2	2.8	8.1	4.3-13.9	1.0	2.3	0.7-5.6
	1996-2000	4.0	5.4	3.3-8.4	2.4	6.9	3.6-12.2	1.6	4.0	1.7-8.1	3.4	4.4	2.6-7.2	2.6	7.4	3.9-12.8	0.8	2.0	0.5-5.5
	2001-2005	4.4	5.7	3.5-8.7	3.6	10.4	6.1-16.5	0.8	1.8	0.4-5.2	3.6	4.4	2.6-7.0	2.6	7.1	3.8-12.3	1.0	2.1	0.6-5.3
	2006-2010	7.2	8.6	6.0-12.0	5.2	13.4	8.7-19.9	2.0	4.0	1.9-7.9	4.0	4.5	2.7-7.1	2.8	7.4	4.0-12.7	1.2	2.4	0.9-5.7
	2011-2015	4.4	5.1	3.1-7.8	4.0	10.2	6.2-16.1	0.4	1.0	0.1-4.0	5.4	6.2	4.0-9.2	4.8	11.9	7.5-18.1	0.6	1.1	0.2-4.0
	2016-2020	4.0	3.6	2.2-5.9	2.8	5.4	2.9-9.6	1.2	1.9	0.7-4.9	5.0	4.8	3.1-7.3	4.0	8.3	5.0-13.4	1.0	1.6	0.5-4.6
Stomach	1976-1980	8.6	12.0	8.6-16.3	5.2	17.3	11.1-25.6	3.4	8.2	4.7-13.3	6.6	9.5	6.4-13.4	4.2	14.4	8.7-22.4	2.4	5.9	3.0-10.7
	1981-1985	7.6	9.8	6.9-13.5	3.8	12.5	7.3-19.9	3.8	8.3	5.0-13.3	5.8	7.2	4.8-10.5	3.0	9.5	5.1-16.2	2.8	5.9	3.2-10.2
	1986-1990	3.4	4.6	2.6-7.5	2.4	7.3	3.7-13.0	1.0	2.4	0.7-6.0	2.8	3.9	2.1-6.6	1.8	6.2	2.8-12.0	1.0	2.4	0.7-6.0
	1991-1995	7.2	8.9	6.2-12.4	4.6	14.3	8.9-21.6	2.6	5.3	2.8-9.4	5.0	6.1	3.9-9.1	3.6	10.8	6.3-17.3	1.4	2.7	1.1-5.9
	1996-2000	6.2	7.5	5.1-10.8	5.2	15.2	9.9-22.4	1.0	1.8	0.6-4.7	3.4	4.1	2.4-6.7	2.0	5.6	2.7-10.4	1.4	2.5	1.0-5.7
	2001-2005	5.2	6.5	4.2-9.6	3.6	10.3	6.1-16.4	1.6	3.7	1.5-7.7	3.0	3.4	1.9-5.8	1.8	5.4	2.5-10.4	1.2	2.1	0.8-5.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Herkimer**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	4.8	5.5	3.5-8.4	3.2	8.7	4.9-14.5	1.6	2.9	1.2-6.4	1.4	1.7	0.6-3.6	1.0	3.0	1.0-7.2	0.4	0.6	0.1-3.1
	2011-2015	7.8	8.5	6.0-11.9	5.2	12.3	7.9-18.4	2.6	5.1	2.7-9.2	2.8	3.0	1.6-5.3	1.8	4.5	2.0-8.9	1.0	1.8	0.6-4.9
	2016-2020	7.0	7.6	5.2-10.8	5.4	12.3	8.0-18.4	1.6	3.4	1.4-7.3	1.4	1.4	0.5-3.1	0.8	1.9	0.5-5.3	0.6	1.1	0.2-4.0
Colorectal	1976-1980	56.2	81.9	72.4-92.3	25.2	88.8	73.4-106.5	31.0	77.1	65.2-90.6	27.4	40.9	34.2-48.5	11.6	43.8	32.9-57.0	15.8	39.1	30.9-48.9
	1981-1985	57.2	76.7	67.9-86.4	28.2	91.1	76.0-108.3	29.0	68.0	57.1-80.5	29.2	39.0	32.9-46.1	14.4	48.1	37.2-61.2	14.8	32.7	25.6-41.2
	1986-1990	53.4	69.1	60.9-78.1	27.4	87.9	73.3-104.6	26.0	56.1	46.6-67.0	29.8	38.7	32.6-45.6	16.4	50.8	40.1-63.6	13.4	29.9	23.0-38.5
	1991-1995	50.6	62.1	54.6-70.4	24.0	69.6	57.5-83.6	26.6	56.8	47.2-67.8	20.0	24.3	19.7-29.7	8.6	27.5	19.6-37.3	11.4	23.2	17.5-30.5
	1996-2000	59.4	72.0	63.9-80.8	26.8	78.2	65.4-92.8	32.6	66.7	56.6-78.2	25.0	29.4	24.5-35.2	12.4	36.7	28.1-47.3	12.6	24.2	18.5-31.4
	2001-2005	48.4	58.4	51.2-66.4	23.2	65.9	54.4-79.2	25.2	52.6	43.5-63.2	13.8	16.1	12.5-20.5	6.8	19.0	13.1-26.7	7.0	13.6	9.3-19.4
	2006-2010	41.8	49.6	43.0-56.9	20.6	54.8	44.6-66.7	21.2	46.4	37.8-56.5	13.0	14.9	11.5-19.2	6.4	17.3	11.8-24.7	6.6	13.1	8.9-18.9
	2011-2015	38.6	44.8	38.5-52.0	21.0	56.2	45.6-68.6	17.6	34.4	27.4-42.9	15.2	17.3	13.5-22.0	7.0	18.9	13.0-26.6	8.2	15.7	11.1-21.9
	2016-2020	35.0	39.6	33.7-46.3	17.2	43.2	34.1-54.0	17.8	35.4	28.2-44.2	13.2	15.1	11.5-19.5	7.8	19.7	13.8-27.5	5.4	10.6	6.9-16.1
Colon excluding rectum	1976-1980	39.6	58.6	50.5-67.6	16.4	60.1	47.2-75.2	23.2	57.6	47.4-69.4	22.0	32.7	26.8-39.6	9.0	35.0	25.2-47.2	13.0	31.8	24.5-40.8
	1981-1985	42.2	56.2	48.7-64.5	20.0	66.9	53.8-82.0	22.2	50.6	41.5-61.3	25.4	33.6	27.9-40.1	12.0	39.2	29.5-51.0	13.4	29.2	22.6-37.3
	1986-1990	38.4	49.8	42.9-57.5	19.2	61.4	49.3-75.5	19.2	41.2	33.2-50.7	24.4	31.7	26.2-38.0	12.8	39.6	30.1-51.0	11.6	26.2	19.6-34.3
	1991-1995	36.6	45.3	38.9-52.5	16.0	47.5	37.4-59.5	20.6	44.3	35.9-54.2	15.8	19.0	15.0-23.8	6.8	21.5	14.7-30.3	9.0	18.6	13.4-25.2
	1996-2000	43.8	52.5	45.7-60.1	17.4	50.7	40.5-62.7	26.4	52.8	43.9-63.0	19.8	23.1	18.8-28.3	8.6	25.1	18.1-33.9	11.2	21.3	16.0-28.1
	2001-2005	37.2	44.6	38.4-51.7	17.4	49.6	39.6-61.3	19.8	40.7	32.9-50.1	11.0	12.9	9.7-17.0	5.8	16.2	10.8-23.5	5.2	10.3	6.6-15.5
	2006-2010	30.4	35.7	30.1-42.0	15.0	39.8	31.3-50.1	15.4	33.7	26.5-42.6	11.2	12.9	9.7-16.9	5.8	15.9	10.6-23.1	5.4	10.7	6.9-16.0
	2011-2015	25.4	28.8	23.9-34.6	12.4	33.1	25.1-42.9	13.0	24.8	19.0-32.1	13.0	14.3	11.0-18.4	6.0	15.4	10.3-22.3	7.0	13.0	9.0-18.6
2016-2020	25.4	28.0	23.2-33.6	11.0	27.0	20.1-35.7	14.4	28.2	21.9-36.1	10.2	11.1	8.2-14.8	5.4	12.7	8.3-18.9	4.8	9.5	6.0-14.8	
Rectum & rectosigmoid	1976-1980	16.6	23.3	18.5-29.1	8.8	28.7	20.6-39.0	7.8	19.5	13.7-26.9	5.4	8.2	5.4-12.0	2.6	8.8	4.6-15.3	2.8	7.3	3.9-12.4
	1981-1985	15.0	20.5	16.0-25.9	8.2	24.3	17.1-33.4	6.8	17.4	11.8-24.7	3.8	5.4	3.2-8.5	2.4	8.9	4.4-15.7	1.4	3.5	1.4-7.4
	1986-1990	15.0	19.3	15.1-24.4	8.2	26.5	18.8-36.3	6.8	14.8	10.1-21.1	5.4	7.1	4.6-10.4	3.6	11.3	6.6-18.1	1.8	3.8	1.7-7.5
	1991-1995	14.0	16.8	13.1-21.4	8.0	22.1	15.7-30.3	6.0	12.5	8.3-18.1	4.2	5.3	3.2-8.1	1.8	6.0	2.7-11.5	2.4	4.6	2.4-8.5
	1996-2000	15.6	19.5	15.3-24.4	9.4	27.5	20.1-36.7	6.2	13.9	9.4-20.1	5.2	6.3	4.1-9.3	3.8	11.7	7.0-18.3	1.4	2.9	1.2-6.3
	2001-2005	11.2	13.8	10.4-18.0	5.8	16.3	10.9-23.5	5.4	11.8	7.6-17.7	2.8	3.1	1.7-5.4	1.0	2.8	0.9-6.7	1.8	3.3	1.5-6.8
	2006-2010	11.4	13.9	10.5-18.2	5.6	14.9	9.8-21.9	5.8	12.7	8.4-18.5	1.8	2.0	0.9-4.0	0.6	1.4	0.3-4.6	1.2	2.5	0.9-5.8
	2011-2015	13.2	16.0	12.3-20.7	8.6	23.1	16.5-31.6	4.6	9.7	6.1-14.9	2.2	3.0	1.4-5.7	1.0	3.5	1.1-8.2	1.2	2.7	0.9-6.5
2016-2020	9.6	11.6	8.4-15.8	6.2	16.1	10.6-23.6	3.4	7.2	4.1-12.2	3.0	4.0	2.2-6.8	2.4	7.0	3.5-12.6	0.6	1.1	0.2-4.1	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Herkimer**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	1.2	1.4	0.5-3.2	0.8	1.9	0.5-5.4	0.4	0.9	0.1-3.5	0.4	0.6	0.1-2.2	0.2	0.5	0.0-3.3	0.2	0.5	0.0-3.1
	1981-1985	1.0	1.2	0.4-3.0	1.0	3.0	0.9-7.2	0.0	0.0	0.0-2.2	1.4	1.7	0.7-3.6	1.4	4.0	1.6-8.6	0.0	0.0	0.0-2.2
	1986-1990	2.6	3.4	1.8-5.9	1.6	5.3	2.2-10.7	1.0	2.0	0.7-5.1	2.6	3.5	1.8-6.1	2.2	7.2	3.5-13.1	0.4	0.9	0.1-3.5
	1991-1995	3.4	4.0	2.3-6.5	1.8	4.8	2.2-9.3	1.6	3.4	1.4-7.0	2.8	3.2	1.8-5.5	1.4	3.7	1.5-7.8	1.4	2.7	1.1-6.0
	1996-2000	3.2	3.9	2.2-6.4	1.6	4.7	2.0-9.4	1.6	3.4	1.4-7.1	3.2	3.8	2.2-6.3	1.8	4.9	2.3-9.6	1.4	2.7	1.1-6.0
	2001-2005	5.8	7.1	4.7-10.3	4.4	12.7	7.9-19.3	1.4	2.9	1.1-6.5	5.0	5.9	3.8-8.9	3.4	9.8	5.6-15.8	1.6	3.0	1.3-6.5
	2006-2010	3.2	3.7	2.1-6.1	2.4	6.2	3.2-11.1	0.8	1.9	0.5-5.3	3.8	4.3	2.5-6.8	2.2	5.9	2.9-10.8	1.6	3.2	1.3-6.7
	2011-2015	7.4	7.9	5.5-11.1	7.0	16.6	11.5-23.5	0.4	0.8	0.1-3.7	5.0	6.0	3.8-9.1	4.0	10.3	6.2-16.3	1.0	2.6	0.8-6.4
2016-2020	5.0	4.8	3.1-7.4	3.6	7.5	4.3-12.4	1.4	2.4	1.0-5.8	3.0	3.2	1.7-5.6	2.2	5.1	2.4-9.8	0.8	1.4	0.4-4.4	
Pancreas	1976-1980	7.0	10.0	6.9-14.1	3.6	12.0	6.9-19.4	3.4	8.2	4.8-13.4	7.6	10.7	7.5-14.8	3.4	11.6	6.5-18.9	4.2	9.9	6.1-15.4
	1981-1985	9.2	11.9	8.6-16.1	5.4	16.4	10.5-24.4	3.8	8.6	5.1-13.8	9.8	12.5	9.2-16.8	4.6	14.4	8.9-22.1	5.2	11.0	7.2-16.4
	1986-1990	7.6	9.2	6.4-12.7	3.4	11.1	6.2-18.1	4.2	8.4	5.2-13.1	7.6	9.4	6.6-13.1	3.4	11.3	6.4-18.4	4.2	8.3	5.2-13.0
	1991-1995	11.4	13.9	10.5-18.1	5.4	14.8	9.7-21.7	6.0	12.9	8.6-18.8	8.6	10.1	7.3-13.7	4.6	12.8	8.0-19.4	4.0	8.2	4.9-12.9
	1996-2000	11.2	13.6	10.2-17.7	5.8	16.7	11.2-24.2	5.4	10.8	7.1-16.1	10.4	12.6	9.4-16.6	5.4	15.8	10.4-23.2	5.0	10.6	6.8-16.1
	2001-2005	11.0	12.7	9.6-16.7	4.8	13.6	8.7-20.3	6.2	12.5	8.4-18.1	10.6	12.2	9.1-16.1	5.2	15.2	9.9-22.4	5.4	10.5	6.8-15.7
	2006-2010	10.8	12.4	9.3-16.3	4.8	12.9	8.2-19.4	6.0	11.9	7.9-17.4	9.8	11.2	8.3-15.0	4.0	10.9	6.7-17.1	5.8	11.6	7.7-17.0
	2011-2015	13.4	15.3	11.8-19.7	6.4	15.9	10.7-22.8	7.0	14.5	9.9-20.7	10.0	11.1	8.2-14.9	4.6	11.8	7.4-17.9	5.4	10.5	6.8-15.8
2016-2020	13.8	14.2	11.0-18.2	6.8	15.1	10.4-21.7	7.0	13.5	9.3-19.3	10.0	10.3	7.6-13.8	5.4	11.7	7.6-17.6	4.6	8.9	5.6-14.0	
Larynx	1976-1980	3.4	4.3	2.5-7.0	2.6	7.7	4.0-13.4	0.8	1.7	0.5-4.8	1.4	1.9	0.8-4.0	1.2	3.7	1.3-8.4	0.2	0.4	0.0-2.9
	1981-1985	4.6	5.9	3.7-8.9	2.6	7.6	3.9-13.2	2.0	4.4	2.1-8.5	1.8	2.3	1.0-4.4	1.2	4.4	1.5-9.7	0.6	1.2	0.2-4.0
	1986-1990	4.2	5.5	3.4-8.6	3.4	10.3	5.9-16.7	0.8	2.2	0.5-5.7	1.4	1.7	0.7-3.7	0.8	2.4	0.6-6.3	0.6	1.5	0.3-4.5
	1991-1995	2.6	3.5	1.8-6.1	1.8	5.6	2.5-10.7	0.8	2.1	0.5-5.6	1.0	1.4	0.4-3.3	0.8	2.4	0.6-6.2	0.2	0.4	0.0-2.7
	1996-2000	2.8	3.5	1.9-6.1	2.6	7.3	3.9-12.6	0.2	0.4	0.0-2.8	1.2	1.5	0.6-3.4	1.0	2.8	0.9-6.7	0.2	0.3	0.0-2.6
	2001-2005	4.0	5.5	3.3-8.5	3.6	10.2	6.0-16.3	0.4	1.1	0.1-4.1	0.8	1.2	0.3-3.1	0.6	2.0	0.4-5.8	0.2	0.5	0.0-3.0
	2006-2010	1.8	2.3	1.0-4.5	1.4	3.9	1.6-8.3	0.4	0.8	0.1-3.5	1.2	1.6	0.6-3.7	1.0	2.8	0.9-6.9	0.2	0.5	0.0-3.2
	2011-2015	2.8	3.1	1.7-5.4	2.0	4.9	2.3-9.4	0.8	1.6	0.4-4.7	0.2	0.2	0.0-1.5	0.2	0.4	0.0-3.0	0.0	0.0	0.0-2.4
2016-2020	2.6	3.4	1.7-6.2	1.6	4.4	1.8-9.2	1.0	2.4	0.6-6.4	1.0	0.9	0.3-2.6	0.8	1.7	0.4-4.9	0.2	0.4	0.0-3.1	
Lung and bronchus	1976-1980	37.2	52.1	44.7-60.4	27.4	91.5	76.1-109.0	9.8	25.6	18.7-34.2	29.6	41.8	35.2-49.3	22.4	78.1	63.7-94.7	7.2	18.1	12.5-25.4
	1981-1985	38.0	49.7	42.7-57.5	25.6	78.4	64.9-93.8	12.4	29.3	22.2-38.1	33.8	43.1	36.7-50.3	26.2	78.8	65.4-94.2	7.6	17.2	12.1-24.0
	1986-1990	47.4	60.5	52.9-68.9	31.2	93.2	78.7-109.7	16.2	38.7	30.4-48.6	42.8	54.7	47.5-62.7	26.2	79.3	65.8-94.7	16.6	38.2	30.1-47.8
	1991-1995	57.0	71.5	63.3-80.6	37.0	108.1	92.8-125.2	20.0	45.9	37.1-56.3	49.4	60.3	52.9-68.4	33.6	99.3	84.5-115.9	15.8	33.6	26.4-42.3
	1996-2000	63.2	78.8	70.3-88.1	36.0	103.0	88.4-119.4	27.2	61.2	51.2-72.8	45.4	56.5	49.3-64.5	25.6	74.1	61.7-88.2	19.8	44.1	35.6-54.0
	2001-2005	60.2	73.5	65.3-82.4	31.6	87.8	74.6-102.8	28.6	63.9	53.6-75.8	47.2	57.1	50.0-65.1	23.4	64.8	53.6-77.9	23.8	51.5	42.4-62.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Herkimer**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	61.0	72.0	64.0-80.7	31.2	81.8	69.4-96.0	29.8	66.3	55.8-78.3	40.8	47.1	40.8-54.2	22.6	59.0	48.5-71.2	18.2	39.0	31.2-48.2
	2011-2015	70.8	79.9	71.6-89.0	37.6	92.9	79.8-107.6	33.2	70.2	59.6-82.2	42.2	47.6	41.3-54.8	24.0	60.9	50.3-73.2	18.2	37.9	30.3-47.0
	2016-2020	63.2	66.9	59.6-75.1	31.6	71.0	60.0-83.6	31.6	62.8	53.2-73.9	36.2	37.9	32.5-44.2	19.6	44.0	35.5-54.2	16.6	32.7	25.9-41.2
Melanoma of the skin	1976-1980	3.8	6.2	3.6-9.9	2.0	5.9	2.7-11.2	1.8	6.7	2.9-12.6	2.0	3.0	1.4-5.5	1.0	2.8	0.9-6.8	1.0	3.1	0.9-7.4
	1981-1985	6.2	9.4	6.2-13.5	3.6	10.9	6.3-17.6	2.6	8.1	4.1-14.1	1.4	2.0	0.8-4.2	0.8	2.4	0.6-6.5	0.6	1.8	0.4-5.3
	1986-1990	3.8	5.2	3.1-8.2	2.8	8.3	4.5-14.1	1.0	2.7	0.8-6.5	3.4	4.2	2.4-6.8	2.2	6.1	3.0-11.2	1.2	2.5	0.9-5.8
	1991-1995	7.6	10.3	7.2-14.3	4.8	14.0	8.9-21.1	2.8	7.5	4.0-12.9	3.6	4.4	2.6-7.0	2.4	6.9	3.5-12.2	1.2	2.7	0.9-6.1
	1996-2000	6.6	9.2	6.3-12.9	3.4	10.1	5.8-16.2	3.2	8.9	5.1-14.7	2.2	3.0	1.5-5.5	2.0	6.0	2.9-11.1	0.2	0.3	0.0-2.7
	2001-2005	8.8	11.4	8.3-15.5	5.2	15.0	9.8-22.1	3.6	9.4	5.5-15.1	0.8	1.0	0.3-2.7	0.4	1.1	0.1-4.1	0.4	0.9	0.1-3.7
	2006-2010	14.2	16.9	13.2-21.5	9.0	23.6	17.1-31.9	5.2	11.3	7.3-17.0	1.4	1.6	0.6-3.4	0.8	2.1	0.6-5.7	0.6	1.2	0.2-4.1
	2011-2015	14.6	19.9	15.4-25.4	8.0	21.5	15.2-29.7	6.6	19.6	13.1-28.1	1.8	1.9	0.9-3.9	1.6	4.0	1.7-8.3	0.2	0.3	0.0-2.9
2016-2020	19.6	24.1	19.2-29.8	10.4	26.4	19.4-35.3	9.2	23.0	16.3-31.7	2.0	2.5	1.1-5.0	1.2	3.3	1.1-7.8	0.8	2.0	0.4-5.8	
Female breast	1976-1980							32.8	87.5	74.1-102.6							11.6	28.7	21.6-37.4
	1981-1985							36.2	93.9	80.1-109.4							11.0	27.1	20.2-35.7
	1986-1990							45.2	112.9	98.0-129.5							12.6	29.4	22.3-38.1
	1991-1995							42.6	104.9	90.8-120.7							15.8	37.7	29.5-47.4
	1996-2000							57.8	140.1	124.0-157.7							14.2	31.9	24.7-40.7
	2001-2005							47.8	115.1	100.7-131.2							8.0	18.6	13.2-25.7
	2006-2010							51.4	123.3	108.2-140.0							9.6	20.0	14.6-26.9
	2011-2015							56.0	130.3	114.7-147.5							9.0	18.0	13.0-24.7
2016-2020							50.6	113.6	99.1-129.8							7.0	14.2	9.7-20.6	
Cervix uteri	1976-1980							6.2	18.0	12.0-25.9							3.2	9.4	5.2-15.7
	1981-1985							7.6	22.6	15.7-31.4							2.8	7.4	3.9-12.8
	1986-1990							4.0	12.7	7.6-19.6							1.8	5.3	2.4-10.2
	1991-1995							3.4	9.3	5.3-15.0							0.2	0.4	0.0-2.7
	1996-2000							4.2	10.9	6.7-16.8							1.6	4.3	1.8-8.6
	2001-2005							1.8	4.8	2.2-9.4							0.6	1.3	0.3-4.2
	2006-2010							2.8	7.2	3.8-12.6							0.8	1.7	0.5-4.8
	2011-2015							2.6	8.6	4.4-15.0							0.4	1.5	0.2-5.2
	2016-2020							1.6	4.8	2.0-9.8							0.4	1.0	0.1-4.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Herkimer**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							10.8	27.0	19.9-35.7							2.4	5.5	2.8-9.8
	1981-1985							15.4	39.2	30.4-49.6							2.2	5.4	2.6-10.0
	1986-1990							11.6	27.7	20.7-36.2							3.2	7.1	4.0-12.0
	1991-1995							10.2	23.9	17.6-31.9							3.2	7.5	4.2-12.6
	1996-2000							9.6	22.5	16.5-30.2							1.6	3.2	1.4-6.7
	2001-2005							9.8	24.6	18.1-32.7							1.6	3.5	1.5-7.3
	2006-2010							15.2	35.9	28.1-45.3							1.2	2.7	1.0-6.3
	2011-2015							15.0	33.8	26.3-43.0							3.0	5.8	3.2-10.1
	2016-2020							14.2	28.9	22.2-37.3							3.4	6.3	3.6-10.6
Ovary	1976-1980							8.6	22.3	15.9-30.3							3.8	9.3	5.5-14.7
	1981-1985							5.0	13.0	8.3-19.5							5.0	12.0	7.6-18.2
	1986-1990							5.2	13.5	8.6-20.2							4.2	9.7	6.0-15.2
	1991-1995							5.2	12.8	8.2-19.1							3.2	7.0	3.9-11.7
	1996-2000							9.2	21.9	15.9-29.6							5.8	12.6	8.4-18.4
	2001-2005							5.4	13.1	8.5-19.3							5.0	11.0	7.0-16.5
	2006-2010							4.6	9.9	6.2-15.2							3.0	6.5	3.6-11.0
	2011-2015							6.2	12.2	8.1-17.9							4.2	8.7	5.3-13.8
	2016-2020							5.4	12.6	8.0-19.2							2.2	3.8	1.8-7.5
Prostate	1976-1980				20.4	77.5	62.7-94.6							6.0	23.3	15.5-33.4			
	1981-1985				25.0	86.1	71.1-103.3							10.0	35.2	25.7-46.9			
	1986-1990				30.8	97.0	81.7-114.3							7.2	22.6	15.5-31.6			
	1991-1995				46.4	136.9	119.3-156.3							11.0	38.3	28.6-50.2			
	1996-2000				54.2	155.4	137.3-175.2							9.4	29.0	21.2-38.7			
	2001-2005				56.8	156.7	138.9-176.2							9.0	26.3	19.1-35.3			
	2006-2010				59.2	147.5	131.0-165.7							4.8	12.9	8.3-19.4			
	2011-2015				51.2	115.9	101.9-131.5							5.8	14.6	9.7-21.2			
	2016-2020				66.6	137.0	122.4-153.3							6.6	15.8	10.8-22.6			
Testis	1976-1980				1.0	3.4	1.1-8.0							0.2	1.0	0.0-4.8			
	1981-1985				0.6	2.0	0.4-5.9							0.0	0.0	0.0-2.5			
	1986-1990				2.2	6.9	3.4-12.5							0.0	0.0	0.0-2.5			
	1991-1995				2.8	9.1	4.9-15.2							0.2	0.6	0.0-3.5			
	1996-2000				1.2	4.1	1.5-8.9							0.0	0.0	0.0-2.4			
	2001-2005				2.6	8.5	4.5-14.6							0.2	0.8	0.0-4.1			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Herkimer**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				1.0	3.7	1.2-8.5							0.2	0.6	0.0-3.5			
	2011-2015				2.6	8.9	4.6-15.4							0.4	1.3	0.1-4.8			
	2016-2020				1.4	4.4	1.7-9.3							0.4	1.2	0.1-4.6			
Urinary bladder (incl. in situ)	1976-1980	16.4	23.4	18.5-29.3	11.8	41.9	31.5-54.5	4.6	10.7	6.8-16.3	4.4	6.4	4.0-9.7	3.0	10.7	5.8-17.9	1.4	3.4	1.3-7.2
	1981-1985	20.6	27.9	22.6-34.1	15.0	49.0	38.1-62.1	5.6	14.1	9.2-20.7	4.6	6.4	4.0-9.6	3.2	10.6	6.0-17.4	1.4	3.5	1.3-7.6
	1986-1990	16.6	21.3	16.9-26.6	10.6	32.1	23.8-42.4	6.0	14.6	9.6-21.3	4.8	6.1	3.9-9.2	3.0	9.1	5.0-15.2	1.8	4.0	1.8-8.0
	1991-1995	15.6	19.6	15.4-24.6	11.4	32.6	24.6-42.4	4.2	9.1	5.5-14.2	4.2	5.2	3.2-8.1	3.6	11.5	6.7-18.4	0.6	1.1	0.2-3.8
	1996-2000	19.6	24.5	19.8-29.9	14.4	41.3	32.2-52.2	5.2	11.7	7.5-17.4	3.8	4.6	2.7-7.3	2.6	7.4	3.9-12.8	1.2	2.5	0.9-5.9
	2001-2005	26.0	31.5	26.2-37.5	20.4	57.9	47.1-70.4	5.6	11.3	7.4-16.7	3.2	3.8	2.2-6.3	1.8	5.5	2.5-10.5	1.4	3.0	1.2-6.7
	2006-2010	25.4	28.9	24.1-34.6	18.2	47.1	37.8-58.1	7.2	15.8	11.0-22.3	6.0	6.6	4.4-9.6	4.0	10.9	6.6-17.0	2.0	3.7	1.7-7.4
	2011-2015	23.0	26.3	21.6-31.8	17.6	44.6	35.5-55.4	5.4	11.2	7.3-16.8	3.4	3.7	2.1-6.1	2.8	7.1	3.9-12.3	0.6	1.4	0.3-4.5
	2016-2020	25.6	27.7	23.0-33.3	19.6	46.1	37.1-56.8	6.0	12.4	8.3-18.2	3.8	3.8	2.3-6.2	3.0	7.0	3.9-12.0	0.8	1.3	0.3-4.3
Kidney and renal pelvis	1976-1980	4.8	7.2	4.5-10.9	2.6	7.8	4.0-13.7	2.2	6.2	3.0-11.4	3.0	4.4	2.5-7.4	2.0	6.7	3.1-12.5	1.0	2.6	0.8-6.2
	1981-1985	6.4	8.8	5.9-12.7	3.4	10.7	6.0-17.5	3.0	6.4	3.6-10.8	3.6	4.7	2.7-7.7	2.0	6.0	2.8-11.4	1.6	3.3	1.4-6.9
	1986-1990	6.0	7.7	5.1-11.1	4.8	13.9	8.8-20.9	1.2	3.0	1.1-6.9	2.8	3.4	1.8-5.8	1.8	5.2	2.3-10.1	1.0	2.0	0.6-5.0
	1991-1995	5.2	6.1	4.0-9.1	3.0	8.5	4.7-14.2	2.2	4.5	2.2-8.3	1.8	2.0	0.9-3.9	0.6	1.6	0.3-4.8	1.2	2.2	0.8-5.2
	1996-2000	8.0	9.9	7.1-13.6	5.0	14.6	9.4-21.8	3.0	6.3	3.4-10.8	2.2	2.7	1.3-4.9	2.0	5.5	2.6-10.3	0.2	0.4	0.0-2.9
	2001-2005	14.0	17.5	13.6-22.2	7.6	21.1	14.9-29.2	6.4	14.5	9.8-20.9	5.4	6.0	3.9-8.9	2.4	6.7	3.4-11.9	3.0	5.1	2.8-8.9
	2006-2010	16.0	19.5	15.4-24.5	8.2	21.3	15.2-29.2	7.8	18.5	13.0-25.7	3.6	4.6	2.7-7.4	2.4	7.0	3.6-12.3	1.2	2.5	0.9-5.8
	2011-2015	16.2	19.8	15.5-24.9	9.8	25.1	18.3-33.7	6.4	15.3	10.2-22.3	2.6	2.8	1.4-4.9	1.6	3.8	1.6-7.9	1.0	2.3	0.7-5.8
	2016-2020	16.4	19.8	15.5-25.0	11.2	27.6	20.5-36.6	5.2	12.3	7.7-18.8	2.4	2.7	1.4-5.0	1.6	4.0	1.7-8.4	0.8	1.5	0.4-4.5
Brain and other central nervous system	1976-1980	2.8	3.9	2.1-6.6	1.6	4.3	1.8-8.8	1.2	3.6	1.3-7.8	2.0	2.6	1.2-4.9	1.6	4.9	2.0-10.0	0.4	1.0	0.1-3.8
	1981-1985	3.4	4.7	2.7-7.6	1.0	3.0	1.0-7.3	2.4	5.9	2.9-10.6	2.4	3.0	1.5-5.4	0.4	1.2	0.1-4.7	2.0	4.6	2.2-8.7
	1986-1990	4.8	6.5	4.1-9.9	2.8	8.9	4.8-15.1	2.0	4.4	2.1-8.5	3.2	4.1	2.3-6.8	1.8	5.5	2.5-10.7	1.4	3.0	1.2-6.5
	1991-1995	4.8	6.8	4.3-10.2	2.2	7.1	3.5-12.7	2.6	6.9	3.6-11.9	3.2	4.4	2.5-7.3	1.4	4.7	1.8-9.7	1.8	4.5	2.0-8.8
	1996-2000	6.2	8.5	5.7-12.1	2.6	7.6	4.1-13.2	3.6	8.8	5.1-14.4	4.6	6.2	3.9-9.4	2.8	8.3	4.5-14.1	1.8	4.2	1.8-8.4
	2001-2005	5.4	7.3	4.8-10.8	2.4	7.5	3.8-13.1	3.0	7.0	3.8-12.0	3.4	4.3	2.5-7.1	0.8	2.3	0.6-6.0	2.6	6.2	3.2-11.1
	2006-2010	5.2	7.5	4.8-11.1	2.8	7.8	4.2-13.4	2.4	7.4	3.7-13.0	2.8	4.0	2.1-6.8	2.4	7.1	3.6-12.7	0.4	1.0	0.1-4.0
	2011-2015	6.6	8.1	5.5-11.7	3.6	10.1	5.8-16.3	3.0	6.6	3.6-11.4	4.2	4.6	2.8-7.2	1.8	4.2	1.9-8.3	2.4	5.2	2.6-9.6
	2016-2020	5.8	7.6	4.9-11.2	2.6	6.6	3.4-11.6	3.2	8.8	4.7-15.0	4.2	4.7	2.8-7.5	2.4	5.8	3.0-10.6	1.8	3.7	1.5-7.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Herkimmer**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	2.2	3.7	1.8-6.7	0.4	1.1	0.1-4.4	1.8	6.0	2.6-11.4	0.4	0.6	0.1-2.1	0.0	0.0	0.0-2.6	0.4	0.9	0.1-3.7
	1981-1985	1.8	2.4	1.1-4.6	0.8	2.5	0.7-6.7	1.0	2.1	0.7-5.4	0.0	0.0	0.0-1.2	0.0	0.0	0.0-2.5	0.0	0.0	0.0-2.2
	1986-1990	1.8	2.8	1.3-5.4	1.0	3.1	1.0-7.4	0.8	2.8	0.7-6.9	0.0	0.0	0.0-1.1	0.0	0.0	0.0-2.5	0.0	0.0	0.0-2.2
	1991-1995	2.8	4.0	2.2-6.8	0.4	1.2	0.1-4.5	2.4	6.3	3.2-11.3	0.2	0.2	0.0-1.5	0.2	0.6	0.0-3.4	0.0	0.0	0.0-2.1
	1996-2000	5.6	8.2	5.4-11.8	2.4	7.5	3.9-13.2	3.2	8.8	5.0-14.5	0.8	1.0	0.3-2.6	0.2	0.5	0.0-3.3	0.6	1.2	0.2-4.1
	2001-2005	5.0	7.9	5.1-11.7	1.2	3.8	1.4-8.3	3.8	11.9	7.1-18.7	0.4	0.5	0.1-1.9	0.2	0.5	0.0-3.2	0.2	0.4	0.0-2.9
	2006-2010	11.8	18.1	13.6-23.5	3.2	9.8	5.4-16.3	8.6	25.7	18.3-35.0	0.8	0.9	0.3-2.6	0.4	1.1	0.1-4.2	0.4	0.7	0.1-3.4
	2011-2015	10.6	16.6	12.2-22.0	2.2	6.5	3.0-12.2	8.4	26.5	18.8-36.4	0.2	0.2	0.0-1.6	0.0	0.0	0.0-2.5	0.2	0.4	0.0-3.0
2016-2020	11.0	17.6	13.0-23.3	3.8	10.8	6.2-17.5	7.2	24.4	16.7-34.3	1.2	1.3	0.5-3.2	0.8	1.9	0.5-5.4	0.4	0.8	0.1-3.8	
Hodgkin lymphoma	1976-1980	1.8	2.2	1.0-4.3	0.8	1.9	0.5-5.4	1.0	2.6	0.8-6.2	0.4	0.5	0.1-2.0	0.2	0.5	0.0-3.4	0.2	0.5	0.0-3.1
	1981-1985	1.4	2.1	0.8-4.4	0.6	2.0	0.4-6.0	0.8	2.2	0.6-5.9	0.6	1.0	0.2-3.0	0.6	2.1	0.4-6.2	0.0	0.0	0.0-2.2
	1986-1990	3.0	4.7	2.6-7.8	2.2	7.4	3.6-13.3	0.8	2.3	0.6-5.9	0.6	0.7	0.2-2.3	0.2	0.6	0.0-3.5	0.4	0.8	0.1-3.4
	1991-1995	2.4	3.6	1.8-6.3	1.2	3.8	1.4-8.3	1.2	3.5	1.3-7.7	0.6	0.9	0.2-2.7	0.4	1.4	0.2-4.8	0.2	0.7	0.0-3.5
	1996-2000	2.0	3.1	1.5-5.7	1.6	5.0	2.2-9.9	0.4	1.2	0.1-4.3	0.8	1.0	0.3-2.7	0.2	0.8	0.0-4.0	0.6	1.4	0.3-4.5
	2001-2005	2.8	4.4	2.4-7.5	1.8	5.8	2.6-11.1	1.0	3.2	1.0-7.4	0.0	0.0	0.0-1.2	0.0	0.0	0.0-2.4	0.0	0.0	0.0-2.3
	2006-2010	3.6	5.6	3.3-9.0	2.4	7.3	3.7-12.9	1.2	4.1	1.5-8.9	0.2	0.2	0.0-1.5	0.2	0.5	0.0-3.3	0.0	0.0	0.0-2.3
	2011-2015	2.8	4.0	2.1-6.9	0.8	2.3	0.6-6.1	2.0	5.8	2.6-11.0	0.4	0.6	0.1-2.2	0.2	0.7	0.0-3.7	0.2	0.4	0.0-3.0
2016-2020	3.0	5.0	2.7-8.4	2.0	5.9	2.7-11.3	1.0	4.0	1.2-9.2	0.2	0.2	0.0-1.6	0.2	0.4	0.0-3.1	0.0	0.0	0.0-2.6	
Non-Hodgkin lymphomas	1976-1980	6.4	9.0	6.1-12.8	3.4	10.8	6.1-17.6	3.0	7.2	4.0-12.1	4.2	6.5	4.0-10.0	1.8	6.7	3.0-12.7	2.4	6.0	3.1-10.7
	1981-1985	8.4	10.9	7.8-14.8	4.4	13.3	8.1-20.6	4.0	9.6	5.8-15.0	4.4	5.7	3.6-8.7	2.6	9.2	4.7-16.1	1.8	4.0	1.8-7.8
	1986-1990	10.6	14.6	10.8-19.3	5.2	16.7	10.7-24.9	5.4	12.7	8.2-19.0	7.2	8.9	6.2-12.5	3.4	11.2	6.2-18.3	3.8	7.9	4.7-12.6
	1991-1995	15.6	20.0	15.8-25.2	8.0	23.6	16.7-32.3	7.6	16.6	11.6-23.2	5.4	6.8	4.4-10.0	2.8	9.1	4.9-15.4	2.6	5.6	2.9-9.9
	1996-2000	14.6	18.6	14.5-23.5	6.4	19.1	13.0-27.0	8.2	17.7	12.5-24.3	8.4	9.7	7.0-13.2	3.6	10.7	6.3-17.0	4.8	9.3	5.9-14.1
	2001-2005	16.0	19.6	15.5-24.5	5.8	16.8	11.2-24.3	10.2	21.8	16.1-29.2	5.6	6.6	4.3-9.6	2.2	6.5	3.2-11.8	3.4	6.3	3.6-10.5
	2006-2010	17.2	21.1	16.8-26.2	8.4	22.9	16.4-31.2	8.8	19.2	13.8-26.2	6.4	7.3	5.0-10.5	2.8	7.4	4.0-12.7	3.6	7.0	4.1-11.6
	2011-2015	15.0	17.0	13.2-21.6	8.0	19.7	13.9-27.2	7.0	14.2	9.7-20.4	5.4	6.3	4.1-9.3	1.6	4.1	1.7-8.5	3.8	7.8	4.6-12.6
2016-2020	17.2	19.3	15.2-24.1	9.0	22.4	16.1-30.4	8.2	16.8	11.8-23.6	3.0	3.4	1.9-5.9	1.2	3.2	1.1-7.6	1.8	3.4	1.5-7.0	
Myeloma	1976-1980	1.4	2.1	0.8-4.3	0.4	1.4	0.2-5.1	1.0	2.5	0.8-6.0	1.6	2.1	0.9-4.2	0.4	1.3	0.1-5.0	1.2	2.8	1.0-6.3
	1981-1985	2.8	3.9	2.1-6.6	1.8	6.0	2.6-11.6	1.0	2.3	0.8-5.7	2.0	3.0	1.5-5.6	1.0	4.0	1.2-9.1	1.0	2.2	0.7-5.5
	1986-1990	2.8	3.6	1.9-6.2	1.2	3.6	1.3-8.1	1.6	3.5	1.5-7.2	1.2	1.7	0.6-3.9	0.8	2.4	0.6-6.4	0.4	1.2	0.1-4.4
	1991-1995	4.0	4.9	2.9-7.6	2.8	8.4	4.5-14.4	1.2	2.6	1.0-6.0	2.8	3.4	1.8-5.8	1.6	4.8	2.0-9.7	1.2	2.4	0.9-5.6
	1996-2000	5.6	6.9	4.6-10.1	2.4	7.2	3.7-12.8	3.2	7.2	4.1-12.0	2.4	2.9	1.5-5.2	0.6	2.0	0.4-5.9	1.8	3.8	1.7-7.6
	2001-2005	4.6	5.3	3.4-8.1	3.2	9.0	5.1-14.8	1.4	2.7	1.1-6.1	3.6	4.2	2.5-6.8	2.4	7.0	3.6-12.4	1.2	2.5	0.9-5.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Herkimer**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	5.4	6.2	4.1-9.2	2.6	6.7	3.5-11.7	2.8	6.0	3.3-10.5	1.8	2.1	1.0-4.2	1.0	2.7	0.9-6.6	0.8	1.8	0.5-4.9
	2011-2015	5.0	6.0	3.8-9.0	3.2	8.3	4.7-13.8	1.8	4.1	1.9-8.2	2.4	2.6	1.4-4.8	1.2	2.9	1.0-6.7	1.2	2.6	1.0-6.2
	2016-2020	4.2	4.7	2.8-7.4	1.8	4.4	2.0-8.7	2.4	5.3	2.6-9.9	1.0	1.1	0.3-2.8	0.2	0.5	0.0-3.4	0.8	1.6	0.4-4.9
Leukemias	1976-1980	8.2	12.3	8.8-16.8	5.2	17.8	11.4-26.5	3.0	8.3	4.6-13.8	6.6	9.9	6.8-14.0	4.0	15.4	9.2-24.0	2.6	6.9	3.6-12.0
	1981-1985	9.4	13.0	9.5-17.5	5.4	17.3	11.1-25.7	4.0	10.5	6.3-16.6	4.4	5.9	3.6-9.0	2.8	9.9	5.2-17.0	1.6	3.7	1.6-7.6
	1986-1990	8.6	11.4	8.2-15.5	3.4	10.2	5.8-16.6	5.2	11.3	7.2-16.9	5.8	7.9	5.3-11.5	2.8	9.5	5.0-16.2	3.0	6.3	3.5-10.8
	1991-1995	11.0	14.0	10.5-18.3	6.8	20.6	14.1-29.0	4.2	9.1	5.5-14.3	5.2	6.9	4.5-10.2	2.8	9.0	4.8-15.3	2.4	5.8	2.9-10.5
	1996-2000	13.0	16.7	12.8-21.4	7.8	23.2	16.5-31.9	5.2	12.2	7.9-18.3	6.0	7.1	4.8-10.3	3.8	10.7	6.4-16.8	2.2	4.2	2.1-8.0
	2001-2005	12.8	16.6	12.7-21.3	8.6	24.3	17.6-32.9	4.2	10.4	6.3-16.2	6.2	7.7	5.2-11.1	4.0	11.0	6.7-17.2	2.2	5.8	2.8-10.6
	2006-2010	11.2	15.1	11.3-19.7	6.4	18.2	12.3-25.9	4.8	12.4	7.8-18.8	4.2	5.3	3.2-8.2	2.2	6.7	3.3-12.1	2.0	4.5	2.2-8.7
	2011-2015	13.4	15.1	11.6-19.5	8.4	20.6	14.6-28.3	5.0	10.2	6.3-15.8	4.6	5.5	3.4-8.6	2.4	6.4	3.3-11.4	2.2	5.3	2.4-10.2
2016-2020	15.6	18.5	14.4-23.6	9.4	23.2	16.7-31.6	6.2	14.3	9.4-21.0	6.8	7.6	5.2-10.8	4.4	11.0	6.8-17.0	2.4	4.8	2.3-9.1	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Jefferson**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	371.0	426.5	407.0-446.7	192.4	538.2	503.4-574.6	178.6	361.1	337.1-386.3	199.4	227.8	213.7-242.6	107.6	305.3	279.0-333.3	91.8	179.9	163.5-197.6
	1981-1985	381.4	463.0	442.1-484.6	191.6	571.8	534.8-610.6	189.8	407.2	380.9-434.7	201.8	244.3	229.2-260.0	106.2	326.0	297.7-356.0	95.6	196.4	178.8-215.3
	1986-1990	415.6	477.4	456.8-498.7	205.4	577.8	541.7-615.5	210.2	429.9	403.6-457.4	207.4	238.4	223.9-253.5	108.2	310.1	283.6-338.2	99.2	195.7	178.4-214.2
	1991-1995	422.4	439.8	421.1-459.1	204.2	513.8	481.9-547.1	218.2	404.4	380.3-429.5	201.8	212.3	199.3-225.9	97.0	263.3	239.6-288.5	104.8	188.0	172.0-205.1
	1996-2000	468.0	476.6	457.5-496.4	235.2	564.4	531.9-598.3	232.8	428.8	404.2-454.4	222.2	227.2	214.0-240.9	110.6	285.1	261.2-310.4	111.6	196.0	179.9-213.2
	2001-2005	547.0	534.2	514.3-554.6	298.2	667.1	633.0-702.5	248.8	444.3	419.7-469.9	210.8	206.0	193.7-218.8	106.8	255.4	233.7-278.5	104.0	177.2	162.1-193.3
	2006-2010	562.2	519.6	500.4-539.4	297.0	602.9	571.9-635.1	265.2	460.0	435.3-485.8	200.8	188.0	176.5-200.1	103.2	224.2	204.8-244.7	97.6	163.8	149.3-179.2
	2011-2015	582.2	519.2	500.2-538.8	296.8	569.7	540.3-600.2	285.4	484.7	459.3-511.0	199.2	179.1	168.0-190.8	101.8	211.0	192.6-230.6	97.4	159.5	145.4-174.7
	2016-2020	641.2	547.3	528.0-567.0	338.4	610.7	581.2-641.4	302.8	503.2	477.3-530.2	206.0	175.7	164.9-187.0	105.4	202.9	185.5-221.4	100.6	157.4	143.6-172.2
Oral cavity and pharynx	1976-1980	13.6	15.2	11.8-19.4	8.8	24.1	17.3-32.7	4.8	9.4	5.9-14.1	2.6	3.0	1.6-5.2	1.6	4.6	1.9-9.2	1.0	2.3	0.7-5.5
	1981-1985	10.0	12.2	9.0-16.2	5.8	16.7	11.0-24.2	4.2	9.4	5.7-14.5	3.2	3.9	2.2-6.3	2.0	6.1	2.9-11.3	1.2	2.4	0.9-5.3
	1986-1990	8.2	9.6	6.8-13.1	6.2	16.1	10.8-22.9	2.0	4.1	1.9-7.6	2.6	3.0	1.6-5.2	1.2	3.6	1.3-7.9	1.4	2.8	1.1-5.8
	1991-1995	8.2	8.7	6.2-11.7	5.4	12.9	8.4-18.8	2.8	5.4	2.9-9.1	2.2	2.3	1.2-4.1	1.2	3.1	1.1-6.6	1.0	1.8	0.6-4.2
	1996-2000	7.6	7.8	5.5-10.7	4.6	11.1	7.0-16.7	3.0	5.8	3.2-9.5	2.8	3.0	1.6-4.9	2.0	4.8	2.3-8.7	0.8	1.6	0.4-4.0
	2001-2005	9.6	9.2	6.8-12.2	7.6	15.0	10.6-20.7	2.0	3.7	1.8-6.9	2.6	2.5	1.3-4.3	1.8	3.6	1.6-6.8	0.8	1.4	0.4-3.6
	2006-2010	14.2	12.8	10.0-16.2	10.4	19.1	14.2-25.2	3.8	7.0	4.2-10.9	3.0	2.9	1.6-4.7	2.0	4.2	2.0-7.7	1.0	1.6	0.5-3.8
	2011-2015	17.2	14.9	11.9-18.4	12.2	23.0	17.4-29.6	5.0	8.0	5.1-11.8	4.2	3.7	2.3-5.7	3.0	6.1	3.4-10.2	1.2	2.0	0.7-4.5
	2016-2020	19.4	15.7	12.7-19.3	14.4	24.4	18.9-30.9	5.0	7.9	5.0-11.9	2.8	2.4	1.3-4.1	2.6	4.8	2.5-8.4	0.2	0.2	0.0-1.8
Esophagus	1976-1980	5.6	6.2	4.1-9.0	3.8	9.4	5.6-14.9	1.8	3.3	1.5-6.5	4.4	4.8	3.0-7.3	2.8	7.2	3.9-12.2	1.6	2.9	1.2-5.8
	1981-1985	3.8	4.3	2.6-6.7	1.8	4.3	2.0-8.3	2.0	3.9	1.9-7.3	3.8	4.4	2.6-6.8	2.6	6.5	3.4-11.2	1.2	2.3	0.8-5.2
	1986-1990	3.0	3.4	1.9-5.7	2.0	5.6	2.6-10.3	1.0	1.7	0.6-4.2	2.4	2.5	1.3-4.3	1.6	4.3	1.7-8.6	0.8	1.4	0.4-3.7
	1991-1995	5.8	6.2	4.2-8.9	3.8	10.5	6.2-16.4	2.0	3.7	1.7-6.8	5.0	5.3	3.4-7.8	4.0	10.9	6.5-16.8	1.0	1.7	0.5-4.0
	1996-2000	7.2	7.5	5.2-10.3	5.0	11.2	7.2-16.5	2.2	3.8	1.9-6.8	5.0	5.1	3.3-7.6	3.8	8.9	5.3-13.9	1.2	2.0	0.7-4.5
	2001-2005	7.6	7.5	5.3-10.3	6.0	12.9	8.7-18.5	1.6	2.9	1.2-5.7	6.0	5.8	3.9-8.3	4.4	9.6	5.9-14.6	1.6	2.8	1.2-5.5
	2006-2010	7.4	6.8	4.8-9.4	5.4	11.5	7.5-16.8	2.0	3.6	1.7-6.6	7.0	6.3	4.4-8.8	5.0	10.2	6.5-15.2	2.0	3.6	1.7-6.6
	2011-2015	7.2	6.6	4.6-9.2	6.2	12.2	8.2-17.4	1.0	1.5	0.5-3.5	6.4	5.7	3.9-8.0	5.4	10.5	6.9-15.4	1.0	1.5	0.5-3.5
	2016-2020	9.2	7.7	5.6-10.3	7.2	12.7	8.8-17.8	2.0	3.3	1.5-6.1	7.6	6.7	4.7-9.3	6.6	12.8	8.8-18.1	1.0	1.5	0.5-3.7
Stomach	1976-1980	7.0	7.7	5.4-10.8	4.0	11.2	6.6-17.6	3.0	5.5	3.1-9.2	6.4	7.1	4.9-10.1	3.8	10.6	6.2-16.9	2.6	5.0	2.6-8.6
	1981-1985	6.6	7.8	5.4-11.0	3.4	9.6	5.4-15.6	3.2	6.5	3.6-10.7	4.8	5.6	3.6-8.4	2.6	7.2	3.7-12.6	2.2	4.1	2.1-7.6
	1986-1990	8.8	9.8	7.1-13.2	5.0	14.7	9.3-21.9	3.8	7.2	4.3-11.4	7.6	9.0	6.3-12.3	4.8	15.1	9.5-22.6	2.8	5.5	3.0-9.4
	1991-1995	5.0	5.2	3.3-7.6	2.2	5.4	2.6-9.6	2.8	4.8	2.6-8.1	3.4	3.6	2.1-5.8	1.8	4.9	2.1-9.4	1.6	2.7	1.1-5.3
	1996-2000	5.0	5.2	3.3-7.6	3.8	9.7	5.7-15.0	1.2	2.0	0.7-4.5	3.6	3.6	2.1-5.7	2.0	4.8	2.3-8.9	1.6	2.7	1.1-5.4
	2001-2005	4.8	4.7	3.0-7.0	3.0	7.1	3.9-11.8	1.8	2.8	1.3-5.5	3.4	3.3	1.9-5.3	2.0	5.0	2.3-9.2	1.4	2.3	0.9-4.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Jefferson**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	6.8	6.4	4.4-8.9	5.0	10.1	6.5-14.9	1.8	3.2	1.4-6.0	2.4	2.2	1.1-3.9	1.8	3.7	1.7-7.0	0.6	0.9	0.2-2.8
	2011-2015	6.4	5.5	3.8-7.8	3.8	6.9	4.1-10.9	2.6	3.9	2.0-6.9	3.0	2.6	1.4-4.3	1.8	3.5	1.6-6.7	1.2	1.9	0.7-4.3
	2016-2020	7.6	6.6	4.6-9.1	4.8	9.3	5.9-13.9	2.8	4.7	2.5-8.0	1.4	1.3	0.5-2.6	0.2	0.3	0.0-2.0	1.2	2.0	0.7-4.5
Colorectal	1976-1980	67.6	77.3	69.1-86.1	31.8	88.2	74.5-103.6	35.8	69.9	59.9-81.2	32.2	37.1	31.5-43.3	14.6	42.1	32.6-53.3	17.6	33.7	26.9-41.7
	1981-1985	63.6	75.7	67.5-84.6	27.6	78.5	65.5-93.2	36.0	72.8	62.4-84.4	31.6	37.1	31.5-43.4	14.0	41.2	31.8-52.3	17.6	34.4	27.5-42.5
	1986-1990	53.6	61.6	54.4-69.5	23.0	66.6	54.7-80.2	30.6	59.3	50.1-69.8	27.8	31.9	26.7-37.7	13.4	38.4	29.5-48.9	14.4	27.0	21.0-34.2
	1991-1995	53.4	55.9	49.4-63.0	22.2	57.4	47.0-69.3	31.2	54.4	46.1-63.8	23.4	24.6	20.3-29.5	9.2	26.4	19.1-35.2	14.2	24.2	18.9-30.7
	1996-2000	62.2	63.5	56.7-71.0	32.8	80.0	68.0-93.5	29.4	50.7	42.8-59.7	23.2	23.6	19.5-28.3	10.8	28.3	21.1-37.0	12.4	20.2	15.4-26.0
	2001-2005	59.4	58.0	51.6-65.0	27.8	64.8	54.3-76.7	31.6	53.9	45.8-63.2	20.2	19.8	16.1-24.1	9.6	22.9	16.7-30.4	10.6	17.3	12.9-22.7
	2006-2010	53.0	49.1	43.3-55.4	27.8	58.6	49.0-69.3	25.2	41.3	34.3-49.4	20.6	19.3	15.7-23.4	9.8	21.6	15.8-28.6	10.8	17.7	13.2-23.2
	2011-2015	57.4	52.0	46.1-58.5	28.2	56.2	47.1-66.5	29.2	48.9	41.1-57.7	19.4	18.0	14.5-21.9	13.0	29.1	22.3-37.1	6.4	10.1	6.8-14.4
2016-2020	55.2	48.0	42.4-54.2	27.2	52.2	43.6-62.0	28.0	45.2	37.8-53.6	15.4	13.0	10.2-16.3	7.2	14.3	9.9-20.0	8.2	12.8	9.1-17.5	
Colon excluding rectum	1976-1980	48.0	55.2	48.3-62.7	21.0	57.5	46.6-70.1	27.0	53.7	44.8-63.8	24.8	28.2	23.4-33.8	11.0	31.0	23.0-40.8	13.8	26.5	20.5-33.7
	1981-1985	43.2	51.9	45.1-59.4	15.6	46.4	36.4-58.3	27.6	55.9	46.8-66.2	26.2	30.9	25.8-36.7	10.6	31.6	23.4-41.6	15.6	30.4	24.0-38.1
	1986-1990	38.8	44.8	38.7-51.6	15.2	44.8	35.0-56.2	23.6	45.4	37.4-54.6	25.4	29.1	24.3-34.7	11.8	33.6	25.3-43.5	13.6	25.6	19.8-32.7
	1991-1995	38.8	40.3	34.8-46.3	14.6	38.1	29.7-48.0	24.2	42.5	35.1-50.9	20.2	21.2	17.3-25.8	7.2	20.8	14.4-28.8	13.0	22.3	17.1-28.5
	1996-2000	45.2	46.1	40.3-52.5	21.4	53.2	43.4-64.4	23.8	40.7	33.7-48.8	22.0	22.4	18.4-27.0	10.4	27.2	20.2-35.7	11.6	18.9	14.3-24.6
	2001-2005	43.8	42.8	37.3-48.8	19.4	45.5	36.8-55.7	24.4	41.5	34.4-49.6	16.8	16.5	13.1-20.4	7.2	17.2	11.9-23.8	9.6	15.8	11.6-21.1
	2006-2010	38.8	36.2	31.2-41.7	18.0	39.3	31.5-48.5	20.8	33.7	27.4-40.9	18.4	17.3	13.9-21.2	8.4	18.9	13.5-25.6	10.0	16.3	12.0-21.6
	2011-2015	40.6	37.2	32.2-42.7	19.6	40.6	32.8-49.6	21.0	34.9	28.4-42.4	16.8	15.5	12.3-19.2	10.6	24.0	17.9-31.4	6.2	9.8	6.6-14.1
2016-2020	40.6	35.2	30.5-40.6	18.2	35.6	28.5-43.9	22.4	36.1	29.6-43.8	12.4	10.6	8.1-13.7	5.6	11.4	7.5-16.5	6.8	10.8	7.4-15.2	
Rectum & rectosigmoid	1976-1980	19.6	22.1	17.9-27.0	10.8	30.7	22.8-40.3	8.8	16.2	11.7-21.9	7.4	8.8	6.2-12.2	3.6	11.1	6.4-17.6	3.8	7.2	4.3-11.4
	1981-1985	20.4	23.8	19.3-28.9	12.0	32.0	24.2-41.5	8.4	16.9	12.1-23.0	5.4	6.3	4.1-9.1	3.4	9.6	5.4-15.5	2.0	4.0	1.9-7.5
	1986-1990	14.8	16.8	13.1-21.1	7.8	21.8	15.4-30.0	7.0	13.9	9.6-19.5	2.4	2.7	1.4-4.8	1.6	4.8	2.0-9.3	0.8	1.4	0.4-3.6
	1991-1995	14.6	15.6	12.2-19.7	7.6	19.3	13.6-26.5	7.0	12.0	8.3-16.8	3.2	3.4	1.9-5.4	2.0	5.6	2.6-10.1	1.2	1.9	0.7-4.3
	1996-2000	17.0	17.5	13.9-21.6	11.4	26.9	20.2-34.9	5.6	10.0	6.6-14.5	1.2	1.2	0.4-2.6	0.4	1.1	0.1-3.7	0.8	1.2	0.3-3.3
	2001-2005	15.6	15.2	12.0-19.0	8.4	19.3	13.8-26.1	7.2	12.5	8.7-17.3	3.4	3.3	1.9-5.3	2.4	5.7	2.9-10.0	1.0	1.4	0.5-3.5
	2006-2010	14.2	12.9	10.1-16.3	9.8	19.2	14.1-25.5	4.4	7.7	4.8-11.7	2.2	2.0	1.0-3.7	1.4	2.6	1.1-5.5	0.8	1.4	0.4-3.6
	2011-2015	16.8	14.8	11.8-18.4	8.6	15.6	11.2-21.2	8.2	14.0	10.0-19.1	2.6	2.5	1.3-4.2	2.4	5.1	2.6-8.9	0.2	0.3	0.0-1.8
2016-2020	14.6	12.8	9.9-16.1	9.0	16.6	12.0-22.4	5.6	9.0	5.9-13.3	3.0	2.4	1.3-4.0	1.6	3.0	1.2-5.9	1.4	2.0	0.8-4.3	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Jefferson**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	4.2	4.6	2.8-7.0	2.8	7.1	3.9-12.1	1.4	2.6	1.0-5.4	3.2	3.6	2.0-5.9	2.0	5.3	2.5-9.9	1.2	2.2	0.8-4.9
	1981-1985	2.4	3.1	1.6-5.4	1.4	5.0	2.0-10.2	1.0	2.1	0.7-4.9	3.2	3.8	2.2-6.2	2.2	7.4	3.6-13.3	1.0	2.0	0.6-4.7
	1986-1990	4.2	5.1	3.1-7.7	2.2	5.6	2.8-10.1	2.0	4.2	2.0-7.8	3.0	3.6	2.0-6.0	1.8	4.8	2.1-9.0	1.2	2.5	0.9-5.5
	1991-1995	3.2	3.4	1.9-5.5	2.2	5.8	2.8-10.4	1.0	1.7	0.5-4.1	2.8	2.9	1.6-4.8	2.0	4.9	2.3-8.9	0.8	1.4	0.4-3.5
	1996-2000	6.0	6.1	4.1-8.7	3.4	8.5	4.9-13.6	2.6	4.7	2.5-8.0	4.4	4.5	2.8-6.8	2.6	7.2	3.7-12.2	1.8	2.9	1.3-5.6
	2001-2005	6.2	6.2	4.2-8.7	4.2	10.3	6.3-15.6	2.0	3.4	1.6-6.4	4.6	4.6	2.9-6.8	3.0	7.4	4.1-12.1	1.6	2.5	1.1-5.1
	2006-2010	5.2	4.7	3.1-6.9	2.0	4.1	1.9-7.6	3.2	5.3	3.0-8.6	4.4	4.1	2.6-6.2	2.0	4.4	2.1-8.0	2.4	4.0	2.0-7.1
	2011-2015	5.8	4.9	3.2-7.0	3.8	7.0	4.1-11.0	2.0	3.3	1.6-6.1	3.6	3.2	1.9-5.0	2.4	5.2	2.6-9.1	1.2	1.9	0.7-4.3
	2016-2020	7.6	6.2	4.4-8.6	5.0	8.5	5.5-12.8	2.6	3.8	2.0-6.8	5.0	4.1	2.6-6.1	2.6	4.5	2.4-7.9	2.4	3.5	1.8-6.2
Pancreas	1976-1980	12.6	14.7	11.3-18.9	6.2	18.3	12.2-26.1	6.4	12.4	8.4-17.7	11.6	13.5	10.2-17.5	6.6	19.5	13.3-27.6	5.0	9.6	6.2-14.3
	1981-1985	12.4	14.5	11.1-18.7	6.4	16.9	11.4-24.1	6.0	12.0	8.0-17.3	12.6	14.9	11.4-19.1	6.0	16.0	10.7-23.0	6.6	13.2	9.0-18.7
	1986-1990	11.8	13.8	10.5-17.8	6.4	18.9	12.7-26.9	5.4	10.5	6.9-15.5	9.8	11.2	8.2-14.8	5.4	15.7	10.2-22.9	4.4	8.1	5.1-12.3
	1991-1995	9.2	9.7	7.1-12.9	4.4	11.5	7.1-17.4	4.8	7.9	5.1-11.9	8.8	9.3	6.8-12.5	4.8	12.3	7.8-18.4	4.0	6.5	3.9-10.1
	1996-2000	8.8	9.0	6.5-12.1	3.4	8.3	4.8-13.2	5.4	9.4	6.1-13.7	9.4	9.7	7.1-12.9	4.2	10.3	6.3-15.7	5.2	9.1	5.9-13.5
	2001-2005	9.2	9.0	6.6-12.0	4.8	10.3	6.5-15.3	4.4	7.0	4.4-10.8	9.2	9.0	6.6-12.0	4.6	10.0	6.3-14.9	4.6	7.2	4.5-10.9
	2006-2010	15.4	14.6	11.5-18.2	6.4	14.3	9.7-20.2	9.0	14.0	10.2-18.9	12.4	11.5	8.8-14.8	5.4	12.0	7.8-17.4	7.0	11.0	7.6-15.4
	2011-2015	14.6	13.2	10.3-16.6	7.2	14.1	9.8-19.6	7.4	12.3	8.6-17.1	12.6	11.4	8.7-14.6	5.8	11.4	7.5-16.5	6.8	11.1	7.6-15.7
	2016-2020	17.2	14.4	11.5-17.9	9.4	16.5	12.1-22.1	7.8	12.7	9.0-17.5	14.4	12.6	9.8-15.9	6.8	12.5	8.6-17.6	7.6	12.7	8.9-17.5
Larynx	1976-1980	7.0	7.7	5.3-10.8	6.4	16.8	11.3-23.9	0.6	1.6	0.3-4.6	2.4	2.5	1.3-4.4	2.2	5.7	2.7-10.4	0.2	0.4	0.0-2.3
	1981-1985	6.2	7.7	5.2-11.0	3.4	9.9	5.6-16.0	2.8	7.2	3.9-12.2	2.0	2.4	1.2-4.5	1.8	7.0	3.1-13.2	0.2	0.4	0.0-2.3
	1986-1990	9.2	10.7	7.8-14.3	7.0	19.0	13.2-26.5	2.2	4.8	2.3-8.6	3.2	3.6	2.0-5.8	2.2	6.1	3.0-10.9	1.0	1.8	0.6-4.3
	1991-1995	6.2	6.7	4.5-9.4	4.0	10.4	6.2-16.1	2.2	4.1	2.0-7.4	2.2	2.2	1.1-3.9	1.2	3.5	1.2-7.5	1.0	1.7	0.5-4.0
	1996-2000	5.8	6.0	4.0-8.5	5.0	11.3	7.3-16.6	0.8	1.5	0.4-3.9	2.0	2.1	1.0-3.8	1.6	3.7	1.6-7.2	0.4	0.7	0.1-2.6
	2001-2005	5.8	5.6	3.7-8.0	4.4	9.0	5.6-13.7	1.4	2.5	1.0-5.1	1.4	1.4	0.6-2.8	1.2	2.7	1.0-5.8	0.2	0.4	0.0-2.1
	2006-2010	4.4	4.0	2.5-6.1	3.2	6.0	3.4-9.8	1.2	2.3	0.8-4.9	2.0	1.8	0.9-3.4	1.4	3.0	1.2-6.1	0.6	1.2	0.3-3.5
	2011-2015	5.0	4.4	2.8-6.5	4.6	8.7	5.5-13.2	0.4	0.7	0.1-2.6	1.6	1.4	0.6-2.7	0.8	1.5	0.4-3.9	0.8	1.3	0.3-3.4
	2016-2020	5.2	4.2	2.7-6.2	3.4	5.4	3.1-8.8	1.8	3.2	1.4-6.2	1.6	1.2	0.5-2.4	1.2	2.0	0.7-4.4	0.4	0.5	0.1-2.2
Lung and bronchus	1976-1980	55.6	62.1	54.9-69.9	41.0	107.8	93.2-124.0	14.6	29.4	22.9-37.2	46.8	52.0	45.5-59.2	35.6	95.4	81.5-111.0	11.2	21.9	16.4-28.7
	1981-1985	70.8	84.7	76.0-94.2	47.0	136.0	118.7-155.1	23.8	50.6	41.7-60.8	57.8	69.5	61.6-78.2	39.8	116.4	100.3-134.3	18.0	39.2	31.3-48.5
	1986-1990	73.0	85.0	76.4-94.3	46.0	129.0	112.4-147.3	27.0	56.0	46.7-66.6	58.6	67.3	59.7-75.6	38.2	106.7	91.7-123.4	20.4	41.9	33.9-51.1
	1991-1995	77.4	81.8	73.8-90.4	43.2	108.7	94.4-124.5	34.2	64.5	55.1-75.1	61.0	64.0	57.0-71.6	34.2	88.6	75.4-103.2	26.8	49.2	41.1-58.4
	1996-2000	85.2	87.9	79.7-96.6	48.4	118.6	103.9-134.7	36.8	68.7	59.1-79.5	64.6	66.5	59.4-74.1	36.4	90.4	77.5-104.6	28.2	51.3	43.1-60.6
	2001-2005	89.8	88.6	80.6-97.2	49.0	109.8	96.2-124.6	40.8	73.1	63.3-83.9	68.4	67.4	60.4-74.9	37.6	85.4	73.4-98.6	30.8	54.6	46.3-64.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Jefferson**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	88.4	83.4	75.7-91.6	45.6	98.2	85.6-112.0	42.8	74.9	65.1-85.8	59.2	55.9	49.6-62.7	31.8	67.8	57.5-79.3	27.4	47.6	39.9-56.4
	2011-2015	98.8	88.6	80.9-96.9	49.6	96.6	84.7-109.7	49.2	82.1	72.0-93.3	59.0	52.8	46.8-59.2	28.4	56.5	47.4-66.9	30.6	50.3	42.5-59.1
	2016-2020	104.4	87.1	79.6-95.1	50.4	91.8	80.6-104.2	54.0	83.9	74.0-94.9	60.8	51.1	45.4-57.3	30.0	56.9	47.9-67.0	30.8	46.9	39.7-55.2
Melanoma of the skin	1976-1980	7.2	8.9	6.2-12.4	3.6	10.3	6.0-16.4	3.6	7.8	4.6-12.5	2.2	2.7	1.3-4.8	1.8	4.9	2.2-9.5	0.4	1.0	0.1-3.5
	1981-1985	7.2	9.6	6.7-13.4	3.0	9.3	5.1-15.5	4.2	10.7	6.4-16.5	2.0	2.9	1.4-5.4	1.8	6.1	2.7-11.5	0.2	0.4	0.0-2.4
	1986-1990	7.4	8.3	5.8-11.6	3.2	8.2	4.6-13.4	4.2	8.6	5.2-13.3	1.8	2.0	0.9-3.7	1.4	3.2	1.3-6.8	0.4	0.9	0.1-3.2
	1991-1995	3.6	3.6	2.1-5.7	2.8	6.6	3.5-11.2	0.8	1.5	0.4-3.7	0.8	0.8	0.2-2.1	0.2	0.6	0.0-3.0	0.6	1.1	0.2-3.2
	1996-2000	9.4	9.1	6.7-12.1	4.4	9.7	5.9-14.8	5.0	9.6	6.2-14.2	2.4	2.4	1.2-4.2	1.0	2.3	0.7-5.3	1.4	2.8	1.1-5.7
	2001-2005	13.4	12.9	10.0-16.4	8.6	19.6	14.0-26.5	4.8	8.6	5.5-12.9	1.6	1.6	0.7-3.1	1.4	3.3	1.3-6.8	0.2	0.4	0.0-2.1
	2006-2010	16.2	15.1	11.9-18.7	10.4	21.8	16.2-28.6	5.8	10.1	6.7-14.6	5.4	5.2	3.4-7.5	4.0	8.7	5.3-13.5	1.4	2.5	1.0-5.2
	2011-2015	27.0	24.0	20.0-28.5	13.2	25.8	19.8-33.0	13.8	23.5	18.1-29.9	1.8	1.4	0.6-2.7	1.0	1.6	0.5-3.8	0.8	1.1	0.3-2.9
	2016-2020	35.0	30.9	26.4-36.0	21.8	40.7	33.2-49.3	13.2	23.2	17.8-29.9	3.4	3.0	1.7-4.8	2.2	4.6	2.3-8.2	1.2	1.8	0.6-4.1
Female breast	1976-1980							42.6	87.6	75.8-100.6							17.4	35.4	28.1-43.9
	1981-1985							41.2	90.5	78.2-104.2							15.6	34.2	26.8-43.1
	1986-1990							57.6	122.1	108.0-137.4							17.0	33.6	26.7-41.8
	1991-1995							56.0	108.2	95.7-121.9							17.2	31.1	24.8-38.5
	1996-2000							67.2	128.4	114.9-143.0							17.0	30.5	24.3-37.8
	2001-2005							66.0	120.8	108.0-134.6							14.4	24.4	19.0-30.9
	2006-2010							70.4	124.5	111.7-138.4							11.2	19.4	14.6-25.3
	2011-2015							71.0	122.2	109.6-135.8							11.6	20.1	15.2-26.1
	2016-2020							77.8	132.3	119.0-146.7							11.0	17.6	13.1-23.2
Cervix uteri	1976-1980							5.4	12.3	8.0-18.0							2.4	5.2	2.6-9.1
	1981-1985							5.0	12.0	7.6-18.0							2.8	6.0	3.2-10.3
	1986-1990							5.2	12.1	7.8-17.7							2.2	5.2	2.5-9.4
	1991-1995							5.6	11.4	7.5-16.6							1.8	3.7	1.7-7.0
	1996-2000							4.8	8.8	5.6-13.1							1.8	3.6	1.6-6.8
	2001-2005							5.2	10.0	6.5-14.6							1.2	2.2	0.8-4.9
	2006-2010							5.0	9.7	6.3-14.3							1.2	2.3	0.8-4.9
	2011-2015							3.0	5.6	3.1-9.4							2.0	3.4	1.6-6.4
	2016-2020							3.6	7.4	4.3-11.7							1.2	2.3	0.8-5.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Jefferson**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							11.8	24.4	18.5-31.7							1.8	3.4	1.5-6.5
	1981-1985							9.0	20.7	14.9-28.0							2.2	4.2	2.1-7.7
	1986-1990							12.0	24.3	18.3-31.5							3.0	5.7	3.2-9.5
	1991-1995							10.0	19.6	14.5-26.0							2.0	3.5	1.7-6.6
	1996-2000							10.2	19.7	14.7-26.0							2.0	3.7	1.7-6.8
	2001-2005							13.6	24.8	19.3-31.5							2.0	3.4	1.6-6.3
	2006-2010							17.0	29.9	23.8-37.0							3.4	5.7	3.3-9.2
	2011-2015							19.4	32.8	26.4-40.1							2.6	4.3	2.3-7.5
	2016-2020							19.6	32.9	26.5-40.4							3.0	4.7	2.6-8.0
Ovary	1976-1980							8.8	19.0	13.6-25.8							6.0	12.7	8.4-18.4
	1981-1985							7.8	17.2	12.1-23.8							6.4	12.9	8.8-18.3
	1986-1990							7.6	16.9	11.8-23.4							3.2	6.8	3.8-11.1
	1991-1995							7.6	14.7	10.3-20.3							4.8	9.8	6.2-14.6
	1996-2000							8.6	16.2	11.7-21.8							5.2	9.1	5.9-13.5
	2001-2005							6.2	11.0	7.4-15.7							5.2	8.9	5.8-13.1
	2006-2010							8.8	15.2	11.0-20.4							4.8	8.1	5.2-12.2
	2011-2015							7.2	12.6	8.7-17.5							4.6	7.5	4.7-11.3
	2016-2020							7.0	12.0	8.3-16.9							4.8	8.0	5.0-12.1
Prostate	1976-1980				31.6	96.4	81.4-113.2							13.4	42.5	32.5-54.4			
	1981-1985				37.6	124.3	106.6-143.9							8.4	31.5	22.5-42.6			
	1986-1990				41.6	124.5	107.7-143.1							11.6	37.2	27.9-48.3			
	1991-1995				57.6	146.0	129.2-164.1							12.6	38.3	29.2-49.1			
	1996-2000				55.4	134.3	118.7-151.3							13.4	39.1	30.1-49.8			
	2001-2005				102.6	231.8	211.8-253.0							12.0	34.2	25.9-44.1			
	2006-2010				90.2	178.0	161.6-195.5							6.8	16.1	11.1-22.5			
	2011-2015				70.2	128.8	115.4-143.4							9.0	20.8	15.1-27.8			
	2016-2020				77.4	131.0	118.0-145.0							8.4	16.5	11.8-22.4			
Testis	1976-1980				0.6	1.5	0.3-4.7							0.2	0.5	0.0-3.0			
	1981-1985				3.0	6.7	3.6-11.4							0.0	0.0	0.0-2.1			
	1986-1990				2.4	5.2	2.6-9.4							0.2	0.4	0.0-2.6			
	1991-1995				1.4	1.9	0.7-4.2							0.2	0.2	0.0-1.9			
	1996-2000				3.2	5.3	3.0-8.8							0.0	0.0	0.0-1.5			
	2001-2005				5.0	8.3	5.3-12.4							0.2	0.3	0.0-2.1			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Jefferson**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				4.6	7.6	4.7-11.5							0.2	0.4	0.0-2.1			
	2011-2015				3.8	5.9	3.4-9.5							0.0	0.0	0.0-1.4			
	2016-2020				6.8	9.9	6.7-14.3							0.2	0.3	0.0-1.9			
Urinary bladder (incl. in situ)	1976-1980	21.2	24.4	19.9-29.6	16.8	49.1	38.9-61.2	4.4	8.4	5.2-12.9	7.0	8.3	5.8-11.6	4.2	14.5	8.8-22.1	2.8	5.5	3.0-9.5
	1981-1985	20.2	24.1	19.6-29.3	14.2	44.8	34.6-56.8	6.0	11.6	7.8-16.7	4.8	5.8	3.7-8.6	2.8	10.7	5.6-18.0	2.0	3.7	1.8-6.9
	1986-1990	24.4	28.0	23.2-33.5	16.8	49.3	38.9-61.3	7.6	14.9	10.4-20.6	4.8	5.4	3.5-8.1	3.0	8.3	4.6-13.6	1.8	3.3	1.5-6.4
	1991-1995	22.6	23.6	19.4-28.4	15.2	39.5	30.9-49.6	7.4	13.4	9.4-18.5	5.0	5.3	3.4-7.8	2.4	7.9	4.0-13.7	2.6	4.3	2.3-7.5
	1996-2000	26.8	27.6	23.1-32.6	17.6	43.6	34.8-53.9	9.2	16.1	11.7-21.5	8.4	8.6	6.2-11.6	4.2	12.0	7.3-18.3	4.2	6.8	4.2-10.6
	2001-2005	29.0	28.5	24.1-33.6	21.2	49.1	40.0-59.6	7.8	13.5	9.6-18.6	5.0	4.8	3.1-7.1	3.4	9.0	5.1-14.4	1.6	2.5	1.1-5.1
	2006-2010	30.6	28.3	24.0-33.2	22.4	48.5	39.8-58.6	8.2	13.5	9.7-18.5	6.2	5.9	4.0-8.4	4.4	9.9	6.2-15.0	1.8	2.8	1.3-5.4
	2011-2015	31.6	28.7	24.3-33.6	22.2	47.0	38.5-56.7	9.4	14.6	10.6-19.6	6.6	6.1	4.2-8.6	4.2	9.9	6.1-15.1	2.4	3.5	1.8-6.3
	2016-2020	34.6	29.8	25.4-34.7	25.4	48.9	40.6-58.4	9.2	14.2	10.3-19.1	5.4	4.5	3.0-6.6	3.8	8.1	4.8-12.7	1.6	2.1	0.9-4.4
Kidney and renal pelvis	1976-1980	6.2	7.1	4.8-10.2	3.8	11.2	6.5-17.7	2.4	4.7	2.4-8.4	3.4	4.0	2.3-6.5	2.6	7.8	4.0-13.5	0.8	1.8	0.5-4.6
	1981-1985	8.4	10.1	7.2-13.8	4.6	13.7	8.5-20.7	3.8	7.9	4.7-12.5	3.6	4.3	2.5-6.8	2.2	7.0	3.3-12.6	1.4	2.7	1.1-5.8
	1986-1990	9.6	11.4	8.4-15.2	5.8	17.0	11.2-24.4	3.8	8.0	4.8-12.6	4.0	4.6	2.8-7.2	2.2	6.8	3.3-12.2	1.8	3.5	1.6-6.8
	1991-1995	9.0	9.4	6.8-12.6	5.0	12.0	7.7-17.7	4.0	7.4	4.5-11.5	4.2	4.4	2.7-6.7	1.8	4.2	1.9-8.0	2.4	4.3	2.2-7.6
	1996-2000	10.6	10.8	8.1-14.1	6.0	13.9	9.3-20.0	4.6	8.7	5.5-13.0	5.0	5.1	3.3-7.5	3.0	8.3	4.6-13.7	2.0	3.5	1.7-6.5
	2001-2005	13.0	12.6	9.7-16.1	7.6	16.4	11.5-22.5	5.4	9.6	6.3-13.9	3.0	2.9	1.6-4.8	1.6	3.7	1.6-7.2	1.4	2.4	1.0-5.1
	2006-2010	16.0	14.4	11.4-17.9	9.8	18.0	13.3-23.9	6.2	10.8	7.3-15.4	5.0	4.5	2.9-6.7	4.2	8.5	5.2-13.0	0.8	1.5	0.4-3.7
	2011-2015	24.8	22.1	18.3-26.4	16.0	30.7	24.2-38.4	8.8	14.8	10.7-20.1	4.4	3.9	2.4-5.9	1.6	2.8	1.2-5.5	2.8	4.5	2.4-7.7
	2016-2020	21.4	18.3	15.0-22.3	13.4	24.9	19.2-31.8	8.0	12.9	9.2-17.8	4.8	4.1	2.6-6.1	3.2	6.3	3.6-10.3	1.6	2.4	1.0-4.9
Brain and other central nervous system	1976-1980	4.8	5.7	3.6-8.5	2.6	6.5	3.4-11.4	2.2	4.5	2.2-8.2	3.8	4.4	2.6-7.0	2.0	5.4	2.5-10.1	1.8	3.2	1.5-6.3
	1981-1985	6.2	7.6	5.1-10.8	3.2	8.8	4.9-14.4	3.0	6.6	3.6-10.9	4.2	5.6	3.4-8.5	2.4	8.0	4.1-14.0	1.8	3.8	1.7-7.2
	1986-1990	7.6	8.0	5.6-11.1	3.8	9.0	5.3-14.2	3.8	7.5	4.4-11.8	4.0	4.4	2.7-6.9	1.8	4.7	2.0-9.1	2.2	4.5	2.2-8.2
	1991-1995	7.4	7.1	4.9-9.9	3.0	5.9	3.1-10.2	4.4	8.7	5.4-13.2	3.0	3.1	1.7-5.0	1.4	3.1	1.2-6.4	1.6	3.2	1.4-6.3
	1996-2000	6.8	6.6	4.5-9.2	3.4	7.4	4.2-11.9	3.4	6.2	3.6-10.0	4.8	4.7	3.0-7.0	2.4	4.7	2.4-8.4	2.4	4.6	2.4-8.1
	2001-2005	7.4	7.1	5.0-9.8	5.2	10.3	6.7-15.1	2.2	4.1	2.1-7.4	2.8	2.8	1.5-4.7	1.6	3.5	1.5-6.8	1.2	2.3	0.9-5.1
	2006-2010	8.4	7.7	5.5-10.4	4.8	8.5	5.4-12.8	3.6	6.5	3.8-10.4	4.8	4.4	2.8-6.5	3.0	5.6	3.1-9.3	1.8	3.0	1.3-5.8
	2011-2015	9.2	8.0	5.8-10.7	5.2	8.7	5.6-12.8	4.0	7.3	4.4-11.3	4.6	4.0	2.5-6.0	2.6	4.8	2.5-8.2	2.0	3.2	1.5-6.0
	2016-2020	8.4	7.7	5.5-10.4	4.4	8.4	5.2-12.9	4.0	7.4	4.5-11.5	5.0	4.4	2.8-6.6	3.0	5.8	3.2-9.7	2.0	3.3	1.5-6.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Jefferson**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	3.2	4.0	2.2-6.6	1.2	2.9	1.0-6.5	2.0	5.1	2.4-9.4	0.6	0.6	0.1-1.8	0.4	0.8	0.1-3.4	0.2	0.4	0.0-2.2
	1981-1985	2.8	3.3	1.8-5.7	0.2	0.5	0.0-3.0	2.6	6.0	3.1-10.4	0.2	0.2	0.0-1.3	0.0	0.0	0.0-2.1	0.2	0.4	0.0-2.4
	1986-1990	2.0	2.1	1.0-3.9	1.0	2.1	0.6-5.2	1.0	2.3	0.7-5.2	0.4	0.5	0.1-1.7	0.2	0.7	0.0-3.3	0.2	0.5	0.0-2.5
	1991-1995	3.8	3.5	2.1-5.5	1.0	2.3	0.7-5.3	2.8	5.1	2.7-8.6	0.4	0.4	0.0-1.4	0.0	0.0	0.0-1.6	0.4	0.7	0.1-2.6
	1996-2000	3.8	3.6	2.1-5.6	1.2	2.4	0.9-5.3	2.6	4.7	2.5-8.0	0.2	0.2	0.0-1.1	0.2	0.5	0.0-2.4	0.0	0.0	0.0-1.4
	2001-2005	9.0	8.4	6.1-11.2	1.8	3.5	1.6-6.7	7.2	13.9	9.7-19.2	0.2	0.2	0.0-1.0	0.0	0.0	0.0-1.5	0.2	0.4	0.0-2.0
	2006-2010	10.6	9.5	7.1-12.4	2.2	3.6	1.7-6.5	8.4	15.9	11.4-21.5	0.0	0.0	0.0-0.7	0.0	0.0	0.0-1.4	0.0	0.0	0.0-1.3
	2011-2015	12.8	11.9	9.1-15.2	1.8	3.1	1.4-6.0	11.0	21.3	15.9-27.8	0.0	0.0	0.0-0.7	0.0	0.0	0.0-1.4	0.0	0.0	0.0-1.3
2016-2020	14.4	13.2	10.2-16.7	5.2	9.4	6.0-14.0	9.2	17.6	12.7-23.7	0.8	0.7	0.2-1.9	0.4	0.8	0.1-2.8	0.4	0.6	0.1-2.4	
Hodgkin lymphoma	1976-1980	3.0	3.7	2.0-6.1	1.6	4.0	1.7-8.1	1.4	3.4	1.3-7.0	1.8	2.2	1.0-4.2	0.8	2.2	0.6-5.9	1.0	1.9	0.6-4.6
	1981-1985	4.2	4.8	2.9-7.4	3.0	7.2	3.8-12.4	1.2	2.7	1.0-6.1	1.2	1.6	0.6-3.5	1.2	4.2	1.4-9.0	0.0	0.0	0.0-1.7
	1986-1990	3.8	3.9	2.3-6.2	2.4	4.8	2.3-8.9	1.4	3.1	1.2-6.4	1.0	1.1	0.3-2.7	0.8	2.0	0.5-5.2	0.2	0.3	0.0-2.0
	1991-1995	4.6	3.7	2.3-5.7	1.8	2.4	1.0-5.0	2.8	4.8	2.6-8.2	0.6	0.6	0.1-1.8	0.4	1.0	0.1-3.5	0.2	0.5	0.0-2.4
	1996-2000	2.8	2.4	1.3-4.1	1.8	3.0	1.3-6.0	1.0	1.9	0.6-4.4	0.4	0.4	0.0-1.4	0.4	0.9	0.1-3.2	0.0	0.0	0.0-1.4
	2001-2005	3.2	2.8	1.6-4.6	1.6	2.9	1.2-5.8	1.6	3.1	1.3-6.0	0.8	0.8	0.2-1.9	0.4	0.7	0.1-2.7	0.4	0.8	0.1-2.8
	2006-2010	3.6	3.0	1.8-4.8	2.8	4.5	2.4-7.8	0.8	1.3	0.4-3.5	0.0	0.0	0.0-0.7	0.0	0.0	0.0-1.4	0.0	0.0	0.0-1.3
	2011-2015	2.2	1.9	0.9-3.4	1.2	2.1	0.7-4.5	1.0	1.8	0.6-4.3	0.6	0.5	0.1-1.6	0.2	0.2	0.0-1.6	0.4	0.7	0.1-2.7
2016-2020	3.0	3.0	1.6-4.9	2.0	3.9	1.8-7.2	1.0	2.0	0.6-4.7	0.6	0.6	0.1-1.6	0.4	0.8	0.1-2.8	0.2	0.3	0.0-2.0	
Non-Hodgkin lymphomas	1976-1980	9.2	10.9	7.9-14.6	4.8	13.2	8.3-19.9	4.4	9.0	5.5-13.9	5.0	5.7	3.6-8.4	1.8	4.5	2.0-8.7	3.2	5.9	3.3-9.7
	1981-1985	10.0	12.4	9.2-16.5	5.8	17.2	11.4-24.9	4.2	8.9	5.4-13.7	6.0	7.2	4.8-10.3	3.6	10.2	5.9-16.4	2.4	5.1	2.6-9.0
	1986-1990	14.4	16.4	12.8-20.7	7.6	20.3	14.2-27.9	6.8	14.0	9.6-19.7	8.0	9.2	6.5-12.5	4.6	12.6	7.9-18.9	3.4	6.6	3.8-10.6
	1991-1995	14.0	14.2	11.1-18.0	6.2	15.5	10.4-22.2	7.8	13.6	9.6-18.7	6.8	7.1	4.9-9.9	3.0	8.2	4.5-13.4	3.8	6.4	3.8-10.1
	1996-2000	13.6	13.5	10.4-17.1	7.6	16.5	11.5-22.9	6.0	10.9	7.3-15.6	9.2	9.3	6.8-12.4	3.8	9.7	5.7-15.1	5.4	9.3	6.1-13.7
	2001-2005	19.2	18.9	15.3-23.1	10.4	23.8	17.7-31.3	8.8	15.5	11.3-20.9	7.0	6.8	4.7-9.4	2.6	6.8	3.6-11.6	4.4	7.1	4.4-10.8
	2006-2010	21.2	19.6	16.0-23.7	10.6	20.4	15.1-26.8	10.6	18.1	13.5-23.7	5.2	4.8	3.1-7.1	3.4	7.7	4.4-12.3	1.8	2.9	1.3-5.6
	2011-2015	22.4	20.8	17.0-25.1	14.0	27.7	21.5-35.2	8.4	14.6	10.4-19.8	3.4	3.2	1.8-5.1	2.0	4.4	2.1-8.1	1.4	2.1	0.8-4.4
2016-2020	28.2	24.8	20.7-29.3	15.6	30.2	23.7-37.8	12.6	21.1	16.1-27.3	5.0	4.4	2.9-6.6	2.8	6.2	3.4-10.3	2.2	3.5	1.7-6.5	
Myeloma	1976-1980	3.4	4.0	2.3-6.4	2.6	7.7	4.0-13.3	0.8	1.5	0.4-4.0	3.0	3.3	1.8-5.5	1.6	4.0	1.7-8.0	1.4	2.6	1.0-5.5
	1981-1985	3.4	4.2	2.4-6.7	1.6	5.5	2.3-10.7	1.8	3.5	1.6-6.7	3.4	4.2	2.4-6.7	2.0	6.4	3.0-11.7	1.4	2.6	1.1-5.6
	1986-1990	4.6	5.1	3.2-7.7	1.8	5.3	2.4-10.2	2.8	5.1	2.8-8.6	2.6	2.9	1.5-5.0	1.2	4.1	1.4-8.6	1.4	2.4	1.0-5.1
	1991-1995	4.0	4.3	2.6-6.6	1.6	4.5	1.9-8.8	2.4	4.6	2.4-8.1	3.8	4.2	2.5-6.5	1.6	5.0	2.1-9.6	2.2	4.4	2.2-7.9
	1996-2000	6.0	6.2	4.2-8.8	3.2	8.3	4.7-13.4	2.8	5.2	2.8-8.7	2.2	2.3	1.1-4.0	1.4	3.6	1.4-7.3	0.8	1.4	0.4-3.7
	2001-2005	6.2	6.2	4.2-8.8	3.6	8.4	5.0-13.3	2.6	4.5	2.4-7.7	4.2	4.2	2.6-6.3	2.8	6.7	3.6-11.2	1.4	2.3	0.9-4.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Jefferson**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	6.2	5.8	3.9-8.3	3.4	7.0	4.0-11.3	2.8	5.0	2.7-8.5	4.2	4.1	2.5-6.3	2.6	6.1	3.2-10.3	1.6	2.7	1.1-5.3
	2011-2015	7.0	6.4	4.4-9.0	3.2	6.1	3.4-10.0	3.8	6.4	3.8-10.2	3.4	2.9	1.7-4.7	2.2	4.4	2.1-7.8	1.2	1.8	0.7-4.1
	2016-2020	7.6	6.5	4.6-9.0	4.0	6.9	4.1-10.7	3.6	6.1	3.5-9.8	3.8	3.3	2.0-5.3	1.4	2.3	0.9-4.8	2.4	4.1	2.0-7.3
Leukemias	1976-1980	11.8	13.1	9.9-16.9	5.8	15.2	10.0-22.2	6.0	11.8	7.9-17.0	7.6	8.7	6.1-12.0	3.6	9.9	5.7-15.8	4.0	8.0	4.8-12.5
	1981-1985	10.6	12.6	9.4-16.6	6.2	18.1	12.0-26.1	4.4	8.6	5.3-13.2	5.4	6.4	4.2-9.4	3.6	10.7	6.1-17.1	1.8	3.5	1.6-6.7
	1986-1990	12.8	13.5	10.3-17.2	8.0	20.3	14.2-28.0	4.8	8.6	5.5-12.9	8.0	8.6	6.1-11.8	4.0	10.2	6.1-16.0	4.0	7.4	4.5-11.6
	1991-1995	14.2	14.6	11.4-18.4	6.6	16.1	11.0-22.7	7.6	13.1	9.2-18.1	7.6	7.9	5.6-10.9	3.6	9.2	5.4-14.5	4.0	7.0	4.2-11.0
	1996-2000	14.0	13.9	10.8-17.6	7.4	17.7	12.3-24.5	6.6	11.5	7.9-16.3	7.6	7.8	5.5-10.6	4.0	10.0	6.0-15.5	3.6	6.1	3.6-9.8
	2001-2005	16.6	15.9	12.7-19.7	8.2	18.2	12.9-24.8	8.4	14.0	10.1-19.0	7.4	7.1	5.0-9.7	3.8	9.2	5.5-14.4	3.6	5.8	3.4-9.3
	2006-2010	18.4	16.4	13.2-20.2	9.6	18.9	13.8-25.2	8.8	14.9	10.8-20.0	7.6	6.8	4.8-9.3	3.2	6.8	3.8-11.1	4.4	7.1	4.4-10.8
	2011-2015	18.8	16.7	13.4-20.5	10.4	19.6	14.5-25.8	8.4	14.2	10.1-19.4	7.0	6.1	4.2-8.6	3.4	6.5	3.7-10.6	3.6	5.8	3.3-9.3
2016-2020	21.8	18.8	15.4-22.7	13.4	25.4	19.5-32.4	8.4	14.3	10.2-19.5	7.0	6.2	4.3-8.6	4.4	8.7	5.4-13.2	2.6	4.1	2.2-7.1	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Lewis**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	78.2	352.5	317.9-389.8	41.8	425.7	367.9-489.6	36.4	304.5	261.1-353.1	38.6	174.5	150.5-201.1	21.8	224.1	182.5-271.8	16.8	138.2	109.9-171.5
	1981-1985	103.2	451.9	413.2-493.1	53.8	532.8	468.9-602.7	49.4	391.5	343.5-444.4	52.6	231.4	204.0-261.4	26.8	281.8	234.3-335.6	25.8	198.8	165.7-236.6
	1986-1990	105.0	442.4	405.0-482.4	52.4	497.4	437.3-563.1	52.6	404.3	356.1-457.2	54.4	231.5	204.5-260.9	29.4	283.8	238.4-334.9	25.0	188.2	156.1-224.8
	1991-1995	125.8	481.9	444.7-521.3	68.2	584.4	522.5-651.3	57.6	408.8	362.3-459.6	62.6	238.5	212.7-266.6	33.6	300.4	255.6-350.4	29.0	196.2	165.3-231.3
	1996-2000	135.4	486.4	450.3-524.6	71.8	567.3	509.0-630.3	63.6	431.6	384.9-482.4	61.4	221.5	197.3-247.8	35.2	287.4	245.7-334.0	26.2	173.9	145.2-206.9
	2001-2005	146.2	494.5	459.2-531.9	86.0	630.1	570.9-693.9	60.2	386.4	343.5-433.4	55.0	185.0	163.8-208.3	27.4	216.6	180.9-257.2	27.6	166.3	139.4-197.1
	2006-2010	160.6	506.6	471.8-543.3	85.6	558.7	506.3-615.2	75.0	463.8	417.1-514.4	61.0	189.0	168.3-211.7	36.2	243.5	208.9-282.3	24.8	139.3	115.6-166.9
	2011-2015	166.2	488.1	454.7-523.5	88.2	527.6	478.5-580.6	78.0	455.8	410.0-505.6	59.0	168.5	149.5-189.3	32.4	199.2	169.3-233.1	26.6	142.3	118.5-170.0
2016-2020	166.2	461.7	429.3-495.9	86.8	479.8	433.9-529.7	79.4	448.0	402.6-497.5	75.2	201.1	180.8-223.3	37.4	212.4	182.5-246.1	37.8	196.6	168.4-228.6	
Oral cavity and pharynx	1976-1980	0.4	2.3	0.3-7.8	0.2	2.7	0.1-13.0	0.2	1.6	0.0-9.0	0.0	0.0	0.0-3.2	0.0	0.0	0.0-7.2	0.0	0.0	0.0-6.2
	1981-1985	3.0	13.5	7.5-22.3	1.4	13.9	5.4-28.8	1.6	13.1	5.6-26.0	1.0	5.1	1.6-11.7	0.6	6.2	1.2-18.0	0.4	3.8	0.5-13.2
	1986-1990	1.4	6.7	2.7-13.7	1.2	11.0	4.0-24.0	0.2	2.2	0.1-10.7	0.4	1.8	0.2-6.3	0.2	1.4	0.0-8.9	0.2	2.2	0.1-10.7
	1991-1995	2.0	8.1	3.8-14.8	1.6	14.4	6.2-28.2	0.4	3.0	0.4-10.8	1.2	4.5	1.7-9.9	0.6	5.6	1.1-15.9	0.6	4.4	0.9-13.0
	1996-2000	3.4	12.4	7.2-19.9	2.6	19.5	10.4-33.7	0.8	5.3	1.4-13.9	1.4	5.0	2.0-10.4	1.2	8.8	3.2-19.4	0.2	1.2	0.0-7.6
	2001-2005	1.8	6.0	2.7-11.5	1.2	9.3	3.3-20.5	0.6	4.2	0.9-12.4	0.8	2.6	0.7-6.9	0.4	3.9	0.4-13.5	0.4	2.4	0.3-9.5
	2006-2010	3.8	11.6	6.9-18.3	2.8	17.0	9.2-29.1	1.0	5.7	1.8-14.0	1.4	4.2	1.7-9.0	1.2	7.8	2.8-17.5	0.2	0.9	0.0-6.8
	2011-2015	4.6	14.2	8.8-21.8	3.4	21.2	12.0-34.9	1.2	7.3	2.7-16.6	1.0	2.8	0.9-6.9	0.8	4.6	1.2-12.8	0.2	0.8	0.0-6.7
2016-2020	3.4	9.2	5.2-15.4	1.6	9.1	3.6-19.3	1.8	9.0	4.1-18.2	1.8	4.6	2.1-9.2	1.2	6.4	2.3-15.0	0.6	2.7	0.5-9.6	
Esophagus	1976-1980	0.4	1.7	0.2-6.3	0.4	3.7	0.4-13.5	0.0	0.0	0.0-6.2	0.4	1.7	0.2-6.1	0.4	3.6	0.4-13.4	0.0	0.0	0.0-6.2
	1981-1985	0.6	2.8	0.6-7.9	0.4	5.6	0.6-18.5	0.2	1.7	0.0-9.4	1.0	4.8	1.5-11.0	0.8	9.6	2.4-24.0	0.2	1.7	0.0-9.4
	1986-1990	1.2	5.3	1.9-11.5	0.6	5.5	1.1-16.1	0.6	4.2	0.9-12.7	1.4	5.8	2.3-12.0	0.4	3.4	0.4-12.5	1.0	7.0	2.3-16.6
	1991-1995	1.6	6.0	2.6-11.8	0.8	6.5	1.7-16.8	0.8	5.7	1.5-14.6	0.8	3.0	0.8-7.8	0.4	4.5	0.5-15.0	0.4	2.7	0.3-10.0
	1996-2000	1.8	6.3	2.9-12.1	1.6	11.6	5.0-23.2	0.2	1.2	0.0-7.6	1.4	4.9	2.0-10.2	1.2	8.8	3.2-19.4	0.2	1.2	0.0-7.6
	2001-2005	1.2	4.0	1.5-8.8	1.2	7.8	2.9-17.6	0.0	0.0	0.0-5.5	0.8	2.7	0.7-7.0	0.8	5.4	1.5-14.4	0.0	0.0	0.0-5.5
	2006-2010	1.0	3.2	1.0-7.6	1.0	6.6	2.1-15.8	0.0	0.0	0.0-5.5	1.6	5.2	2.2-10.4	1.6	11.4	4.9-22.7	0.0	0.0	0.0-5.5
	2011-2015	1.4	3.8	1.5-8.3	1.4	8.2	3.2-17.6	0.0	0.0	0.0-5.6	0.8	3.0	0.8-7.7	0.6	3.9	0.8-11.9	0.2	2.1	0.1-10.0
2016-2020	1.0	2.9	0.9-7.3	0.6	3.7	0.7-11.8	0.4	2.2	0.3-9.2	1.0	2.9	0.9-7.2	1.0	5.8	1.8-14.4	0.0	0.0	0.0-5.9	
Stomach	1976-1980	1.8	7.9	3.6-15.1	1.0	11.4	3.4-26.6	0.8	6.3	1.7-16.3	1.0	4.5	1.5-10.6	0.4	4.2	0.5-14.9	0.6	4.7	1.0-13.9
	1981-1985	2.0	8.7	4.1-16.0	0.8	6.7	1.8-17.6	1.2	9.1	3.3-20.2	1.6	7.0	3.0-13.8	0.6	6.5	1.1-19.0	1.0	7.5	2.4-17.9
	1986-1990	1.4	6.4	2.6-13.1	0.6	6.2	1.3-17.4	0.8	5.5	1.5-14.5	1.0	4.8	1.6-11.0	0.6	6.2	1.3-17.4	0.4	2.8	0.3-10.5
	1991-1995	1.8	7.0	3.2-13.3	0.4	3.0	0.4-11.2	1.4	9.6	3.8-20.0	1.4	5.3	2.1-11.0	0.8	7.0	1.8-17.9	0.6	4.1	0.8-12.3
	1996-2000	2.2	7.9	3.9-14.2	1.6	12.9	5.4-25.6	0.6	3.7	0.7-11.3	1.6	5.8	2.5-11.4	1.0	8.7	2.7-20.2	0.6	4.1	0.8-12.3
	2001-2005	1.0	3.3	1.1-7.9	0.4	2.8	0.3-10.5	0.6	3.7	0.7-11.5	1.0	3.3	1.1-7.9	0.6	4.2	0.9-12.7	0.4	2.3	0.3-9.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Lewis**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	2.6	8.2	4.3-14.1	2.2	14.4	7.2-26.1	0.4	1.8	0.2-8.2	1.0	3.0	1.0-7.2	0.6	3.6	0.7-11.4	0.4	2.1	0.2-8.7
	2011-2015	1.8	5.5	2.4-10.7	1.0	5.7	1.8-14.0	0.8	5.1	1.2-13.9	1.2	3.3	1.2-7.6	0.6	3.7	0.8-11.5	0.6	3.0	0.6-10.1
	2016-2020	2.4	6.3	3.2-11.5	1.8	9.6	4.3-19.2	0.6	2.7	0.5-9.6	0.8	2.2	0.6-6.1	0.8	4.4	1.2-12.4	0.0	0.0	0.0-5.9
Colorectal	1976-1980	15.6	69.6	54.9-87.0	7.0	73.2	49.9-103.1	8.6	69.7	50.2-94.2	8.4	37.9	27.2-51.3	4.2	44.4	26.8-68.7	4.2	33.8	20.8-51.9
	1981-1985	17.8	77.8	62.4-95.9	9.0	90.3	64.9-121.9	8.8	66.8	48.5-90.0	8.2	35.3	25.2-47.9	3.2	35.3	19.4-58.1	5.0	37.3	24.1-55.3
	1986-1990	15.6	64.6	50.9-80.8	6.4	63.9	43.1-90.7	9.2	66.7	48.7-89.3	9.6	40.9	30.1-54.2	4.8	50.0	31.6-74.7	4.8	34.1	21.8-50.9
	1991-1995	15.4	59.1	46.6-73.9	7.0	61.5	42.3-86.0	8.4	56.8	40.7-77.2	7.4	28.1	19.7-38.7	4.2	40.4	24.6-61.8	3.2	20.0	11.4-32.9
	1996-2000	21.8	77.8	63.9-94.0	10.8	86.4	64.4-113.3	11.0	70.2	52.7-91.8	8.8	32.0	23.3-43.1	3.6	29.1	17.0-46.2	5.2	33.3	21.6-49.1
	2001-2005	20.2	68.3	55.6-83.1	11.6	87.7	66.1-114.1	8.6	52.3	37.7-71.0	6.8	22.7	15.7-31.9	3.0	25.4	13.9-42.1	3.8	21.4	12.8-34.1
	2006-2010	17.0	53.2	42.4-66.0	8.8	57.5	41.5-77.7	8.2	49.2	35.1-67.4	5.6	17.5	11.6-25.5	4.0	27.8	16.9-43.3	1.6	8.4	3.6-17.4
	2011-2015	20.2	63.0	51.0-77.1	10.6	68.9	51.2-90.8	9.6	57.8	42.0-77.9	8.2	24.9	17.7-34.2	5.0	32.0	20.6-47.8	3.2	17.4	9.6-29.4
2016-2020	15.8	47.8	37.3-60.4	7.2	45.6	31.4-64.2	8.6	49.8	35.2-68.7	8.6	23.6	16.9-32.4	3.8	23.0	13.7-36.7	4.8	24.2	15.1-37.3	
Colon excluding rectum	1976-1980	9.8	43.6	32.2-57.7	3.8	37.5	22.1-59.1	6.0	47.8	32.2-68.5	6.2	27.4	18.6-39.0	3.2	31.6	17.7-51.8	3.0	23.7	13.2-39.4
	1981-1985	12.8	56.6	43.5-72.3	6.2	64.2	42.7-92.1	6.6	50.9	35.0-71.7	7.4	32.3	22.7-44.5	3.0	33.9	18.2-56.5	4.4	32.9	20.6-50.1
	1986-1990	10.2	42.2	31.4-55.6	4.0	39.2	23.4-61.1	6.2	44.1	29.9-62.8	8.2	34.8	24.9-47.3	4.0	42.3	25.3-65.5	4.2	29.6	18.3-45.5
	1991-1995	12.4	47.4	36.3-60.9	5.4	47.2	30.6-69.2	7.0	47.0	32.6-65.7	6.8	25.5	17.7-35.7	4.0	38.6	23.2-59.7	2.8	17.7	9.7-30.1
	1996-2000	16.8	60.0	47.8-74.4	7.6	62.3	43.7-86.1	9.2	59.0	43.0-79.2	8.2	29.7	21.3-40.4	3.2	25.8	14.5-42.2	5.0	32.2	20.8-47.9
	2001-2005	14.2	47.8	37.3-60.5	6.8	53.2	36.4-74.9	7.4	44.0	30.9-61.3	5.4	18.1	11.9-26.4	2.4	20.1	10.1-35.3	3.0	16.9	9.4-28.5
	2006-2010	12.2	38.3	29.3-49.5	6.4	42.2	28.7-60.1	5.8	34.9	23.3-50.8	4.2	13.1	8.1-20.2	2.8	20.1	10.9-33.9	1.4	7.2	2.8-15.7
	2011-2015	14.0	43.1	33.4-54.9	6.8	44.8	30.8-63.2	7.2	41.0	28.2-57.8	7.4	22.7	15.9-31.7	4.4	28.6	17.8-43.8	3.0	16.6	9.0-28.5
2016-2020	9.8	29.3	21.3-39.4	4.0	25.0	15.0-39.5	5.8	33.4	21.7-49.4	5.6	15.3	10.0-22.7	2.0	11.9	5.6-22.7	3.6	18.4	10.6-30.5	
Rectum & rectosigmoid	1976-1980	5.8	26.0	17.3-37.5	3.2	35.7	19.5-58.9	2.6	21.8	11.5-37.6	2.2	10.5	5.2-18.6	1.0	12.8	3.8-29.5	1.2	10.0	3.6-22.0
	1981-1985	5.0	21.2	13.6-31.4	2.8	26.1	14.0-44.1	2.2	15.9	7.9-28.9	0.8	3.0	0.8-7.9	0.2	1.4	0.0-9.3	0.6	4.4	0.9-13.2
	1986-1990	5.4	22.4	14.7-32.6	2.4	24.6	12.5-42.9	3.0	22.6	12.6-37.6	1.4	6.0	2.4-12.4	0.8	7.8	2.1-19.5	0.6	4.5	0.9-13.3
	1991-1995	3.0	11.7	6.5-19.3	1.6	14.3	6.1-28.1	1.4	9.8	3.9-20.5	0.6	2.5	0.5-7.2	0.2	1.8	0.0-9.5	0.4	2.3	0.3-9.1
	1996-2000	5.0	17.8	11.5-26.4	3.2	24.1	13.6-39.4	1.8	11.2	5.1-21.7	0.6	2.3	0.5-6.8	0.4	3.3	0.4-11.8	0.2	1.0	0.0-7.1
	2001-2005	6.0	20.5	13.8-29.4	4.8	34.5	22.0-51.8	1.2	8.2	3.0-18.2	1.4	4.6	1.9-9.7	0.6	5.3	1.0-15.3	0.8	4.6	1.2-12.5
	2006-2010	4.8	14.9	9.5-22.4	2.4	15.3	7.8-27.2	2.4	14.2	7.2-25.6	1.4	4.4	1.8-9.3	1.2	7.8	2.8-17.4	0.2	1.2	0.0-7.6
	2011-2015	6.2	20.0	13.4-28.8	3.8	24.0	14.3-38.2	2.4	16.9	8.4-30.2	0.8	2.2	0.6-6.1	0.6	3.4	0.7-10.9	0.2	0.8	0.0-6.7
2016-2020	6.0	18.5	12.1-27.1	3.2	20.5	11.3-34.3	2.8	16.4	8.5-29.0	3.0	8.3	4.6-14.2	1.8	11.2	5.0-21.9	1.2	5.7	2.1-13.9	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Lewis**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	0.4	2.0	0.2-6.9	0.2	3.6	0.1-16.0	0.2	1.4	0.0-8.5	0.6	2.9	0.6-8.3	0.4	4.4	0.5-15.3	0.2	1.4	0.0-8.5
	1981-1985	0.6	2.7	0.6-7.9	0.4	4.2	0.5-14.8	0.2	1.3	0.0-8.4	0.8	3.7	1.0-9.3	0.4	4.2	0.5-14.8	0.4	2.7	0.3-10.5
	1986-1990	1.4	6.0	2.4-12.4	0.8	7.3	2.0-18.5	0.6	5.5	1.1-15.6	0.6	2.5	0.5-7.2	0.6	5.2	1.1-15.2	0.0	0.0	0.0-5.8
	1991-1995	2.0	7.5	3.6-13.9	1.6	13.2	5.6-26.2	0.4	2.6	0.3-9.7	2.0	7.3	3.5-13.6	1.2	9.8	3.5-21.4	0.8	5.4	1.5-14.1
	1996-2000	1.6	5.8	2.5-11.5	0.6	4.7	1.0-13.8	1.0	6.4	2.0-15.3	1.6	6.0	2.6-11.8	0.8	6.1	1.6-15.8	0.8	5.2	1.4-13.8
	2001-2005	0.6	1.9	0.4-5.9	0.4	2.7	0.3-10.3	0.2	1.2	0.0-7.6	0.2	0.6	0.0-3.9	0.2	1.3	0.0-8.2	0.0	0.0	0.0-5.5
	2006-2010	1.8	5.4	2.5-10.5	1.4	9.4	3.7-19.9	0.4	2.5	0.3-9.7	1.0	3.0	1.0-7.3	0.6	4.3	0.8-12.7	0.4	2.1	0.2-8.7
	2011-2015	1.4	4.1	1.6-8.8	1.0	5.8	1.9-14.3	0.4	2.4	0.3-9.6	1.2	3.7	1.3-8.3	0.8	4.9	1.3-13.2	0.4	2.3	0.3-9.3
2016-2020	2.2	6.0	3.0-11.3	1.6	9.1	3.9-18.7	0.6	3.1	0.6-10.4	2.6	6.9	3.6-12.2	1.4	7.4	2.9-16.3	1.2	6.3	2.3-14.9	
Pancreas	1976-1980	1.0	3.9	1.2-9.2	0.2	1.5	0.0-9.6	0.8	5.7	1.6-15.2	2.0	8.5	4.0-15.8	0.8	9.1	2.2-23.3	1.2	8.7	3.2-19.4
	1981-1985	2.4	10.7	5.4-18.7	1.6	15.3	6.4-30.4	0.8	6.3	1.7-16.4	2.6	12.1	6.4-20.7	1.4	15.7	6.0-32.3	1.2	9.5	3.5-20.9
	1986-1990	2.6	11.0	5.8-18.8	1.4	13.7	5.2-28.3	1.2	9.3	3.3-20.5	2.8	12.5	6.8-20.9	1.4	16.0	6.2-32.5	1.4	10.5	4.1-22.0
	1991-1995	2.2	8.3	4.1-14.9	1.4	11.1	4.4-22.9	0.8	5.5	1.5-14.4	2.0	7.6	3.6-13.9	1.6	13.1	5.6-25.9	0.4	2.6	0.3-9.7
	1996-2000	3.4	11.8	6.9-19.0	2.4	18.4	9.3-32.6	1.0	7.1	2.3-16.8	3.0	10.7	6.0-17.7	2.2	17.6	8.7-31.8	0.8	5.9	1.6-15.1
	2001-2005	5.4	18.3	12.1-26.7	2.0	14.1	6.7-26.2	3.4	19.5	11.3-31.9	4.4	15.0	9.4-22.8	1.4	10.7	4.2-22.4	3.0	17.2	9.6-29.0
	2006-2010	6.0	18.6	12.5-26.7	3.6	24.5	14.4-39.1	2.4	14.1	7.2-25.1	5.6	17.1	11.4-24.9	3.6	24.4	14.3-38.9	2.0	11.2	5.3-21.2
	2011-2015	4.0	11.5	7.0-18.0	1.6	9.4	4.0-19.1	2.4	12.6	6.4-22.9	3.4	9.7	5.6-15.8	2.0	12.4	5.9-23.3	1.4	7.0	2.7-15.6
2016-2020	5.4	14.9	9.6-22.3	2.0	10.9	5.0-21.1	3.4	18.3	10.3-30.7	7.4	19.0	13.3-26.6	3.4	17.9	10.3-29.6	4.0	19.7	11.9-31.5	
Larynx	1976-1980	0.2	1.2	0.0-5.7	0.2	2.4	0.1-12.0	0.0	0.0	0.0-6.2	0.2	0.7	0.0-4.5	0.2	1.5	0.0-9.7	0.0	0.0	0.0-6.2
	1981-1985	1.4	6.5	2.5-13.4	1.2	13.0	4.5-28.3	0.2	1.4	0.0-8.7	0.8	3.4	0.9-8.8	0.6	6.2	1.2-17.8	0.2	1.3	0.0-8.4
	1986-1990	1.4	5.9	2.3-12.2	0.8	6.2	1.6-16.5	0.6	5.4	1.1-15.4	0.6	2.2	0.5-6.6	0.4	3.3	0.4-12.2	0.2	1.4	0.0-8.2
	1991-1995	1.2	4.4	1.6-9.6	1.2	8.9	3.2-19.8	0.0	0.0	0.0-5.4	0.6	2.3	0.5-6.7	0.6	5.0	1.0-14.5	0.0	0.0	0.0-5.4
	1996-2000	1.8	6.3	2.9-12.1	1.4	11.2	4.4-23.2	0.4	2.7	0.3-10.1	1.2	4.0	1.5-9.0	1.0	8.2	2.5-19.3	0.2	1.2	0.0-7.5
	2001-2005	0.8	2.6	0.7-6.9	0.4	2.6	0.3-10.1	0.4	2.6	0.3-9.8	0.8	2.7	0.7-7.2	0.4	3.1	0.4-11.4	0.4	2.6	0.3-9.9
	2006-2010	2.2	6.5	3.2-11.9	1.2	7.0	2.6-15.9	1.0	6.1	2.0-14.8	0.4	1.3	0.2-4.9	0.4	2.6	0.3-9.9	0.0	0.0	0.0-5.5
	2011-2015	1.0	3.2	1.0-7.8	0.8	4.3	1.1-12.1	0.2	2.1	0.1-10.0	0.8	2.4	0.6-6.6	0.6	2.7	0.6-9.3	0.2	2.1	0.1-10.0
2016-2020	0.8	2.4	0.6-6.8	0.4	1.6	0.2-7.8	0.4	3.2	0.3-11.7	0.4	1.0	0.1-4.3	0.2	0.8	0.0-6.7	0.2	1.1	0.0-7.6	
Lung and bronchus	1976-1980	13.8	61.3	47.5-77.8	11.2	108.4	81.2-141.6	2.6	22.5	11.9-38.5	9.2	41.1	30.0-54.9	7.6	74.3	52.0-102.6	1.6	13.6	5.8-26.8
	1981-1985	16.8	75.7	60.2-93.8	10.8	107.2	79.9-140.4	6.0	51.8	34.6-74.3	14.4	62.9	49.1-79.4	9.0	88.0	63.5-118.6	5.4	44.3	29.0-64.7
	1986-1990	17.8	74.0	59.3-91.3	13.0	118.2	90.5-151.4	4.8	36.7	23.4-54.8	14.0	60.1	46.7-76.1	10.8	100.8	75.1-132.3	3.2	27.2	15.4-44.2
	1991-1995	19.0	72.5	58.6-88.7	12.2	105.0	80.0-135.2	6.8	46.9	32.3-65.8	17.2	65.3	52.1-80.7	10.4	88.3	65.6-116.2	6.8	46.3	32.0-65.0
	1996-2000	17.8	63.2	50.7-77.9	10.2	80.9	59.9-106.8	7.6	51.4	36.2-70.8	14.2	51.7	40.3-65.2	9.8	81.0	59.6-107.5	4.4	30.1	18.8-45.8
	2001-2005	18.4	61.1	49.3-75.1	10.0	72.9	53.8-96.7	8.4	53.1	38.2-72.1	14.4	47.7	37.3-60.2	7.6	56.3	39.6-77.8	6.8	41.5	28.7-58.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Lewis**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	20.6	64.3	52.4-78.1	11.0	72.3	54.2-94.5	9.6	57.5	42.3-76.8	15.4	48.1	37.9-60.3	9.0	59.4	43.2-80.0	6.4	37.8	25.7-54.0
	2011-2015	21.8	61.6	50.5-74.7	15.2	91.2	71.5-114.8	6.6	35.1	23.9-50.1	16.4	46.3	36.7-57.8	9.6	59.1	43.4-79.0	6.8	35.8	24.6-50.9
	2016-2020	22.2	58.5	47.9-71.0	11.0	60.1	44.8-79.4	11.2	57.1	42.9-75.1	17.4	46.9	37.4-58.3	8.4	47.2	33.9-64.6	9.0	46.8	33.8-63.8
Melanoma of the skin	1976-1980	1.0	4.0	1.3-9.6	0.4	2.8	0.3-11.5	0.6	4.1	0.8-12.7	0.0	0.0	0.0-3.2	0.0	0.0	0.0-7.2	0.0	0.0	0.0-6.2
	1981-1985	2.2	10.2	5.0-18.2	1.2	11.8	4.3-25.6	1.0	9.4	2.9-21.6	0.8	3.5	0.9-9.0	0.4	3.9	0.4-14.0	0.4	3.4	0.4-12.2
	1986-1990	1.0	4.2	1.4-9.7	0.6	5.8	1.1-16.7	0.4	3.1	0.4-11.4	0.6	2.6	0.5-7.4	0.4	3.0	0.4-11.5	0.2	2.2	0.1-10.7
	1991-1995	2.8	10.6	5.8-17.8	1.2	8.3	3.0-18.5	1.6	11.3	4.8-22.5	1.0	4.1	1.3-9.5	0.4	3.1	0.4-11.5	0.6	4.1	0.8-12.3
	1996-2000	4.6	16.1	10.2-24.2	1.8	12.9	5.9-24.8	2.8	19.5	10.6-32.9	0.8	2.7	0.7-7.1	0.8	5.7	1.5-14.9	0.0	0.0	0.0-5.4
	2001-2005	3.4	11.9	6.9-19.2	2.2	16.2	7.9-29.4	1.2	8.9	3.2-19.5	1.0	3.5	1.1-8.3	0.6	3.9	0.8-12.0	0.4	3.0	0.3-10.8
	2006-2010	7.2	23.1	16.1-32.3	4.4	29.0	18.0-44.5	2.8	18.5	9.9-31.5	0.6	1.8	0.4-5.5	0.4	2.5	0.3-9.7	0.2	1.2	0.0-7.4
	2011-2015	8.6	27.2	19.4-37.1	4.2	25.7	15.8-39.9	4.4	29.3	17.8-45.5	1.4	4.1	1.6-8.8	0.8	5.5	1.5-14.4	0.6	2.8	0.6-9.6
2016-2020	6.8	19.9	13.4-28.7	3.8	20.7	12.1-33.5	3.0	19.7	10.4-33.8	0.8	2.2	0.6-6.1	0.6	3.3	0.7-10.7	0.2	1.1	0.0-7.6	
Female breast	1976-1980							8.4	73.1	52.3-99.3							2.6	23.8	12.5-40.8
	1981-1985							10.8	89.0	66.4-116.6							2.8	19.8	10.8-33.7
	1986-1990							14.2	113.6	88.2-144.0							3.8	29.6	17.6-46.7
	1991-1995							13.8	103.2	80.0-131.1							4.6	31.3	19.7-47.3
	1996-2000							14.8	102.1	79.9-128.7							4.2	27.4	16.8-42.2
	2001-2005							16.2	104.0	82.5-129.6							3.4	20.0	11.6-32.6
	2006-2010							18.0	114.2	91.5-141.0							2.4	14.4	7.4-25.7
	2011-2015							22.2	129.2	105.6-156.9							4.4	23.7	14.6-36.8
2016-2020							21.0	117.7	95.1-144.5							4.2	23.7	14.2-37.6	
Cervix uteri	1976-1980							1.8	15.6	6.9-29.8							0.8	6.7	1.8-17.3
	1981-1985							0.8	6.2	1.7-16.1							0.2	1.7	0.0-9.4
	1986-1990							1.6	12.4	5.3-24.7							0.8	6.2	1.7-16.0
	1991-1995							1.2	8.7	3.2-19.0							1.2	8.6	3.1-18.9
	1996-2000							1.2	9.2	3.4-20.0							0.4	3.5	0.4-12.0
	2001-2005							0.4	3.3	0.4-11.8							0.4	3.2	0.4-11.4
	2006-2010							1.8	14.1	6.3-26.9							0.4	2.6	0.3-9.8
	2011-2015							1.0	6.8	2.1-16.3							0.2	1.6	0.0-8.7
	2016-2020							1.0	6.9	2.0-17.0							0.8	5.7	1.4-15.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Lewis**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							3.2	26.9	15.2-43.9							0.6	5.0	1.0-14.6
	1981-1985							3.6	28.8	16.9-45.9							1.2	8.8	3.2-19.6
	1986-1990							4.0	30.9	18.7-48.0							1.0	7.4	2.4-17.4
	1991-1995							3.8	29.0	17.3-45.5							0.8	5.2	1.4-13.7
	1996-2000							4.2	30.0	18.5-46.0							1.0	7.4	2.4-17.3
	2001-2005							2.8	18.5	10.1-31.4							0.4	2.6	0.3-9.8
	2006-2010							6.4	39.5	26.8-56.2							1.8	10.3	4.7-20.2
	2011-2015							5.6	32.9	21.3-48.8							1.2	6.9	2.5-15.8
	2016-2020							4.6	23.5	14.6-36.6							2.2	11.0	5.2-21.0
Ovary	1976-1980							2.0	16.2	7.7-30.1							1.0	8.7	2.8-20.2
	1981-1985							1.8	14.6	6.5-28.1							1.4	11.3	4.5-23.5
	1986-1990							1.2	8.9	3.3-19.6							0.2	1.3	0.0-8.1
	1991-1995							2.2	15.8	7.8-28.6							1.8	12.2	5.5-23.5
	1996-2000							2.2	16.2	8.0-29.1							1.0	7.2	2.3-17.0
	2001-2005							2.6	16.8	8.9-29.1							2.0	13.3	6.4-24.7
	2006-2010							2.2	12.7	6.3-23.3							1.4	7.8	3.1-16.8
	2011-2015							2.6	14.2	7.5-25.1							1.4	7.7	3.1-16.7
	2016-2020							2.0	9.8	4.7-19.2							1.6	7.9	3.4-16.8
Prostate	1976-1980				9.4	100.0	72.7-133.8							2.2	24.6	11.9-44.1			
	1981-1985				10.8	109.5	81.1-144.0							4.2	51.8	31.4-79.3			
	1986-1990				11.2	116.4	87.3-151.4							3.4	32.9	18.7-53.1			
	1991-1995				20.4	179.8	145.6-219.3							4.0	39.1	23.4-60.6			
	1996-2000				18.6	148.3	119.1-182.3							5.0	48.3	30.9-71.3			
	2001-2005				31.8	227.8	193.2-266.8							4.0	35.6	21.3-55.2			
	2006-2010				24.0	152.1	125.9-182.5							2.4	16.8	8.6-29.6			
	2011-2015				19.2	107.8	87.0-132.4							3.0	18.2	10.1-30.6			
	2016-2020				25.6	123.0	102.2-147.5							4.4	26.4	16.5-40.5			
Testis	1976-1980				0.4	3.8	0.5-13.7							0.0	0.0	0.0-7.2			
	1981-1985				1.4	10.6	4.1-22.9							0.0	0.0	0.0-7.0			
	1986-1990				1.0	7.0	2.2-17.2							0.0	0.0	0.0-6.6			
	1991-1995				0.4	3.1	0.4-11.4							0.0	0.0	0.0-6.0			
	1996-2000				0.4	3.0	0.4-11.2							0.0	0.0	0.0-5.9			
	2001-2005				0.4	3.2	0.4-11.5							0.0	0.0	0.0-5.8			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Lewis**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				0.8	7.6	2.1-18.7							0.0	0.0	0.0-5.5			
	2011-2015				1.0	7.8	2.4-18.3							0.0	0.0	0.0-5.5			
	2016-2020				0.8	8.0	2.2-19.7							0.0	0.0	0.0-5.7			
Urinary bladder (incl. in situ)	1976-1980	3.8	17.8	10.6-27.8	3.8	38.7	23.1-60.6	0.0	0.0	0.0-6.2	1.0	4.6	1.5-10.7	1.0	11.2	3.5-25.6	0.0	0.0	0.0-6.2
	1981-1985	4.4	19.3	12.0-29.3	3.2	29.4	16.6-48.2	1.2	9.0	3.3-20.0	0.8	3.8	1.0-9.5	0.4	4.3	0.5-14.9	0.4	3.0	0.4-11.4
	1986-1990	2.8	12.0	6.5-20.1	2.0	20.0	9.3-36.8	0.8	6.0	1.6-15.5	1.0	4.4	1.4-10.2	0.2	2.0	0.1-10.5	0.8	5.8	1.6-15.1
	1991-1995	5.8	22.2	14.8-31.9	4.4	37.8	23.5-57.5	1.4	9.6	3.8-20.1	0.8	3.1	0.9-8.0	0.6	6.8	1.4-18.7	0.2	1.3	0.0-7.8
	1996-2000	6.4	23.3	15.9-33.0	5.2	43.0	27.8-63.3	1.2	8.9	3.2-19.5	0.6	2.2	0.5-6.4	0.6	5.8	1.1-16.6	0.0	0.0	0.0-5.4
	2001-2005	6.2	20.7	14.0-29.5	4.8	35.3	22.4-53.0	1.4	7.7	3.1-16.6	0.2	0.7	0.0-4.0	0.0	0.0	0.0-5.8	0.2	1.0	0.0-7.0
	2006-2010	6.6	20.5	14.1-29.1	5.2	34.8	22.6-51.4	1.4	8.5	3.3-18.1	1.0	3.1	1.0-7.4	0.4	2.3	0.3-9.3	0.6	3.1	0.6-10.3
	2011-2015	8.8	24.2	17.5-32.9	7.2	43.0	29.9-60.3	1.6	8.4	3.6-17.4	1.4	4.2	1.7-9.0	1.0	6.6	2.1-15.9	0.4	1.9	0.2-8.3
	2016-2020	9.2	24.4	17.7-33.0	6.8	37.2	25.6-52.8	2.4	13.1	6.5-24.1	1.6	4.3	1.8-8.9	1.0	6.2	2.0-15.0	0.6	3.1	0.6-10.4
Kidney and renal pelvis	1976-1980	1.2	5.3	1.9-11.6	0.6	5.1	1.0-15.5	0.6	5.3	1.1-15.2	0.2	1.0	0.0-5.2	0.0	0.0	0.0-7.2	0.2	1.6	0.0-9.0
	1981-1985	2.8	12.8	6.9-21.6	2.0	19.3	9.0-35.9	0.8	6.0	1.6-15.7	1.8	8.6	3.9-16.3	0.8	8.3	2.2-21.2	1.0	8.8	2.8-20.5
	1986-1990	2.4	10.1	5.2-17.7	1.2	12.4	4.3-27.0	1.2	9.4	3.3-20.6	1.0	3.6	1.2-8.7	0.4	2.9	0.3-11.4	0.6	3.9	0.8-12.1
	1991-1995	2.6	9.9	5.3-17.0	2.2	18.5	9.1-33.1	0.4	2.7	0.3-9.9	1.4	5.4	2.2-11.2	0.6	5.7	1.1-16.3	0.8	5.7	1.5-14.7
	1996-2000	3.4	12.3	7.1-19.7	1.8	13.7	6.2-26.2	1.6	10.6	4.5-21.2	1.0	3.6	1.2-8.5	1.0	7.6	2.4-17.8	0.0	0.0	0.0-5.4
	2001-2005	3.4	11.4	6.7-18.4	2.8	20.8	11.1-35.4	0.6	4.2	0.9-12.4	1.4	4.7	1.9-9.9	0.8	5.5	1.5-14.5	0.6	3.9	0.8-11.9
	2006-2010	7.2	22.0	15.3-30.6	3.2	21.4	12.1-35.1	4.0	23.6	14.3-37.2	2.8	8.9	4.8-15.1	1.6	11.7	5.0-23.3	1.2	6.9	2.4-16.1
	2011-2015	6.2	18.3	12.3-26.4	3.8	22.5	13.4-35.8	2.4	13.8	6.8-25.3	1.4	3.8	1.5-8.3	1.0	5.8	1.9-14.3	0.4	1.6	0.2-7.8
	2016-2020	4.6	12.8	7.9-19.9	2.6	12.8	6.7-23.1	2.0	13.1	6.0-25.2	1.0	2.6	0.8-6.6	0.8	4.4	1.2-12.3	0.2	0.9	0.0-7.1
Brain and other central nervous system	1976-1980	1.4	5.4	2.1-11.5	0.8	7.1	1.9-18.7	0.6	3.9	0.8-12.3	1.0	4.5	1.4-10.6	0.6	5.6	1.0-17.0	0.4	3.1	0.4-11.4
	1981-1985	1.4	5.5	2.2-11.5	0.6	4.5	0.9-14.0	0.8	6.3	1.7-16.4	1.0	4.1	1.3-9.6	0.6	4.7	1.0-14.5	0.4	3.4	0.4-12.3
	1986-1990	1.8	7.6	3.4-14.4	0.8	6.4	1.7-16.8	1.0	8.3	2.6-19.4	1.4	5.9	2.3-12.2	0.6	5.2	1.0-15.3	0.8	6.4	1.7-16.6
	1991-1995	0.6	2.0	0.4-6.1	0.2	1.4	0.0-8.5	0.4	2.5	0.3-9.7	0.4	1.4	0.2-5.1	0.4	2.9	0.3-10.9	0.0	0.0	0.0-5.4
	1996-2000	1.6	5.9	2.5-11.7	0.8	5.9	1.6-15.5	0.8	5.7	1.5-14.7	0.4	1.4	0.2-5.1	0.2	1.3	0.0-8.1	0.2	1.2	0.0-7.4
	2001-2005	1.6	5.5	2.4-11.0	0.8	5.5	1.5-14.4	0.8	5.3	1.4-13.9	1.2	4.3	1.6-9.4	0.6	4.6	0.9-13.7	0.6	4.0	0.8-12.0
	2006-2010	1.8	5.7	2.6-11.1	1.0	6.6	2.1-15.8	0.8	4.6	1.2-12.5	1.4	4.4	1.8-9.2	0.8	5.1	1.4-13.5	0.6	3.4	0.7-10.8
	2011-2015	2.2	7.1	3.4-13.0	0.8	4.2	1.1-11.8	1.4	9.7	3.7-20.5	1.8	4.8	2.2-9.5	0.8	4.2	1.1-11.8	1.0	5.1	1.6-13.0
	2016-2020	1.6	4.4	1.8-9.2	0.6	3.4	0.6-11.2	1.0	5.0	1.6-13.1	1.4	4.3	1.6-9.4	0.4	2.0	0.2-8.7	1.0	6.5	1.9-16.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Lewis**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	1.2	6.5	2.3-13.9	0.2	2.7	0.1-13.0	1.0	10.4	3.3-23.7	0.0	0.0	0.0-3.2	0.0	0.0	0.0-7.2	0.0	0.0	0.0-6.2
	1981-1985	0.8	3.5	0.9-9.0	0.2	2.1	0.1-11.1	0.6	4.8	1.0-14.4	0.0	0.0	0.0-3.2	0.0	0.0	0.0-7.0	0.0	0.0	0.0-6.2
	1986-1990	0.6	2.8	0.6-7.9	0.2	1.7	0.0-9.6	0.4	3.6	0.4-12.7	0.0	0.0	0.0-3.0	0.0	0.0	0.0-6.6	0.0	0.0	0.0-5.8
	1991-1995	0.4	1.5	0.2-5.3	0.2	1.6	0.0-8.9	0.2	1.5	0.0-8.3	0.0	0.0	0.0-2.8	0.0	0.0	0.0-6.0	0.0	0.0	0.0-5.4
	1996-2000	1.8	6.6	3.0-12.6	0.6	4.3	0.9-13.0	1.2	9.0	3.3-19.5	0.0	0.0	0.0-2.8	0.0	0.0	0.0-5.9	0.0	0.0	0.0-5.4
	2001-2005	3.6	13.5	7.9-21.5	0.4	3.7	0.4-13.0	3.2	24.7	14.0-40.3	0.2	0.6	0.0-3.9	0.0	0.0	0.0-5.8	0.2	1.2	0.0-7.6
	2006-2010	4.0	15.0	9.0-23.3	0.6	4.1	0.8-12.3	3.4	26.3	15.1-42.4	0.2	0.6	0.0-3.7	0.2	1.4	0.0-7.9	0.0	0.0	0.0-5.5
	2011-2015	4.2	14.6	8.9-22.7	1.6	11.0	4.5-22.4	2.6	18.3	9.5-31.7	0.0	0.0	0.0-2.8	0.0	0.0	0.0-5.5	0.0	0.0	0.0-5.6
	2016-2020	3.4	12.8	7.2-21.0	1.0	6.4	1.9-15.9	2.4	19.6	9.7-34.8	0.0	0.0	0.0-2.9	0.0	0.0	0.0-5.7	0.0	0.0	0.0-5.9
Hodgkin lymphoma	1976-1980	0.2	0.5	0.0-4.0	0.0	0.0	0.0-7.2	0.2	1.1	0.0-7.8	0.2	0.7	0.0-4.5	0.0	0.0	0.0-7.2	0.2	1.4	0.0-8.6
	1981-1985	0.6	2.4	0.5-7.2	0.4	3.5	0.4-13.1	0.2	1.3	0.0-8.4	0.2	0.7	0.0-4.4	0.2	1.4	0.0-9.3	0.0	0.0	0.0-6.2
	1986-1990	0.4	1.8	0.2-6.5	0.2	2.3	0.1-11.5	0.2	1.4	0.0-8.3	0.2	1.1	0.0-5.5	0.2	2.3	0.1-11.5	0.0	0.0	0.0-5.8
	1991-1995	0.6	2.1	0.4-6.3	0.2	1.4	0.0-8.5	0.4	2.8	0.3-10.2	0.2	0.7	0.0-4.1	0.0	0.0	0.0-6.0	0.2	1.3	0.0-7.8
	1996-2000	0.6	2.4	0.5-7.0	0.4	3.4	0.4-12.1	0.2	1.5	0.0-8.2	0.2	0.7	0.0-4.0	0.2	1.4	0.0-8.5	0.0	0.0	0.0-5.4
	2001-2005	1.2	4.7	1.7-10.2	0.8	6.2	1.7-16.1	0.4	3.1	0.4-11.1	0.0	0.0	0.0-2.8	0.0	0.0	0.0-5.8	0.0	0.0	0.0-5.5
	2006-2010	0.6	2.1	0.4-6.3	0.2	1.1	0.0-7.4	0.4	3.1	0.4-11.2	0.2	0.6	0.0-3.7	0.0	0.0	0.0-5.5	0.2	1.0	0.0-7.1
	2011-2015	0.2	0.8	0.0-4.3	0.2	1.6	0.0-8.6	0.0	0.0	0.0-5.6	0.0	0.0	0.0-2.8	0.0	0.0	0.0-5.5	0.0	0.0	0.0-5.6
	2016-2020	1.4	6.0	2.4-12.3	0.4	3.6	0.4-12.3	1.0	8.6	2.7-20.0	0.0	0.0	0.0-2.9	0.0	0.0	0.0-5.7	0.0	0.0	0.0-5.9
Non-Hodgkin lymphomas	1976-1980	2.2	11.2	5.5-20.0	1.4	15.2	5.7-31.5	0.8	8.6	2.2-21.4	0.4	2.0	0.2-6.9	0.2	2.3	0.1-11.9	0.2	1.6	0.0-9.0
	1981-1985	2.8	10.8	5.9-18.4	1.2	10.3	3.6-23.1	1.6	11.3	4.9-22.8	1.2	5.1	1.8-11.1	0.8	8.1	2.1-20.8	0.4	2.7	0.3-10.5
	1986-1990	4.6	20.0	12.6-30.0	2.8	28.3	15.2-47.5	1.8	14.6	6.5-27.8	1.2	5.3	1.9-11.5	1.0	11.7	3.7-26.5	0.2	1.4	0.0-8.2
	1991-1995	4.6	17.0	10.7-25.5	2.2	17.1	8.3-31.1	2.4	17.0	8.7-30.0	2.2	8.3	4.2-15.0	1.2	9.2	3.4-20.3	1.0	6.5	2.1-15.6
	1996-2000	5.4	19.1	12.5-27.9	2.4	17.8	9.2-31.3	3.0	19.0	10.5-31.8	2.6	9.3	4.9-16.0	1.2	9.1	3.3-20.1	1.4	8.8	3.5-18.5
	2001-2005	6.6	22.9	15.7-32.3	4.6	33.8	21.3-51.0	2.0	12.4	5.9-23.2	1.6	5.5	2.4-11.0	1.0	7.3	2.3-17.3	0.6	3.4	0.7-10.8
	2006-2010	6.6	20.4	14.0-28.9	3.8	24.6	14.6-38.9	2.8	15.7	8.5-27.0	2.8	8.8	4.8-15.1	2.0	13.8	6.6-25.8	0.8	4.3	1.2-12.0
	2011-2015	9.2	25.3	18.4-34.1	5.2	29.5	19.1-43.9	4.0	20.5	12.3-32.7	3.0	8.0	4.4-13.5	1.0	5.0	1.6-12.8	2.0	9.8	4.7-19.1
	2016-2020	8.4	25.2	17.7-34.8	6.0	37.5	24.7-54.8	2.4	13.3	6.5-24.7	3.2	7.8	4.4-13.1	2.0	11.1	5.2-21.3	1.2	5.3	1.9-13.1
Myeloma	1976-1980	0.8	3.4	0.9-8.8	0.6	5.0	1.0-15.1	0.2	1.6	0.0-9.0	0.6	2.7	0.5-7.8	0.4	3.4	0.4-12.9	0.2	1.4	0.0-8.5
	1981-1985	1.4	6.4	2.6-13.1	1.0	11.2	3.4-25.9	0.4	3.1	0.4-11.4	0.4	1.9	0.2-6.6	0.4	4.3	0.5-14.9	0.0	0.0	0.0-6.2
	1986-1990	1.2	4.9	1.8-10.6	0.2	2.4	0.1-11.9	1.0	7.4	2.4-17.4	0.8	3.3	0.9-8.4	0.2	1.9	0.0-10.1	0.6	4.4	0.9-13.0
	1991-1995	1.2	4.6	1.7-10.1	0.8	6.4	1.7-16.6	0.4	3.5	0.4-12.1	1.0	4.0	1.3-9.2	0.8	7.4	2.0-18.5	0.2	1.3	0.0-7.8
	1996-2000	2.8	10.1	5.5-17.0	1.4	11.6	4.5-24.0	1.4	8.9	3.5-18.8	1.0	3.5	1.1-8.2	0.2	1.4	0.0-8.5	0.8	5.3	1.4-13.9
	2001-2005	1.4	5.0	2.0-10.4	1.2	10.0	3.6-21.8	0.2	1.3	0.0-7.9	0.8	2.7	0.7-7.1	0.4	3.1	0.4-11.4	0.4	2.7	0.3-10.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Lewis**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	0.2	0.7	0.0-3.9	0.2	1.4	0.0-8.0	0.0	0.0	0.0-5.5	0.8	2.4	0.7-6.5	0.6	4.0	0.8-12.0	0.2	1.1	0.0-7.2
	2011-2015	2.2	6.3	3.1-11.5	1.6	9.1	3.9-18.6	0.6	3.6	0.7-11.3	0.8	2.2	0.6-6.2	0.8	4.7	1.2-12.7	0.0	0.0	0.0-5.6
	2016-2020	4.0	11.5	6.9-18.2	2.8	16.4	8.8-28.3	1.2	7.0	2.4-16.3	1.4	3.5	1.4-7.9	0.8	4.6	1.2-12.6	0.6	3.1	0.6-10.4
Leukemias	1976-1980	2.6	11.6	6.1-19.9	1.4	15.1	5.8-31.3	1.2	10.3	3.7-22.6	1.8	7.9	3.6-15.1	0.8	8.1	2.1-20.7	1.0	8.8	2.7-20.6
	1981-1985	2.8	12.4	6.7-20.8	1.6	16.3	6.7-32.3	1.2	8.7	3.2-19.4	1.0	4.9	1.5-11.2	0.2	2.6	0.1-12.8	0.8	5.8	1.6-15.2
	1986-1990	4.4	17.9	11.1-27.2	2.6	22.3	11.6-38.5	1.8	13.3	6.0-25.5	2.8	11.5	6.2-19.3	1.4	13.6	5.2-28.3	1.4	9.5	3.8-20.0
	1991-1995	5.2	19.9	13.0-29.2	2.4	18.7	9.6-32.9	2.8	19.9	10.8-33.6	2.2	9.1	4.5-16.2	0.8	6.8	1.9-17.3	1.4	10.1	4.0-21.0
	1996-2000	6.0	21.7	14.6-31.1	2.8	20.6	11.2-34.8	3.2	22.1	12.5-36.1	3.6	12.8	7.6-20.4	1.6	12.4	5.3-24.7	2.0	13.1	6.2-24.4
	2001-2005	7.6	26.7	18.8-36.7	4.0	30.0	18.2-46.7	3.6	23.7	13.9-38.0	3.2	11.1	6.3-18.1	1.6	12.0	5.1-23.8	1.6	8.9	3.8-18.3
	2006-2010	6.8	21.2	14.7-29.9	5.0	33.4	21.5-49.6	1.8	10.3	4.7-20.1	2.4	7.3	3.8-13.0	1.0	7.0	2.2-16.6	1.4	7.0	2.8-15.3
	2011-2015	6.0	17.6	11.8-25.5	3.4	21.4	12.3-34.8	2.6	14.3	7.5-25.4	2.2	6.5	3.2-11.9	1.6	10.6	4.6-21.3	0.6	3.5	0.7-11.2
	2016-2020	7.4	21.0	14.5-29.5	4.2	24.2	14.7-38.0	3.2	18.5	10.2-31.3	3.0	8.2	4.5-14.0	1.0	5.8	1.8-14.4	2.0	11.1	5.2-21.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Livingston**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	186.8	400.8	375.0-427.9	94.0	465.0	421.8-511.2	92.8	359.7	327.2-394.6	93.0	203.7	185.3-223.4	52.6	269.8	236.6-306.1	40.4	157.4	136.1-180.9
	1981-1985	213.0	432.9	406.9-460.1	101.2	493.2	449.3-540.0	111.8	410.6	376.6-446.7	106.8	220.5	201.9-240.2	53.6	266.1	233.9-301.2	53.2	195.5	172.4-220.8
	1986-1990	250.8	476.6	450.3-504.0	127.6	579.2	533.3-627.7	123.2	421.4	388.3-456.6	113.2	215.8	198.2-234.5	57.4	267.7	236.3-301.8	55.8	188.9	167.1-212.7
	1991-1995	298.4	529.0	502.4-556.7	164.2	680.6	633.3-730.4	134.2	433.2	400.8-467.6	118.8	214.3	197.4-232.3	64.2	288.9	256.8-323.6	54.6	172.4	152.4-194.3
	1996-2000	301.6	513.5	487.8-540.1	158.2	610.7	567.7-655.9	143.4	450.8	418.2-485.2	138.6	239.1	221.6-257.6	71.6	294.3	263.7-327.4	67.0	205.6	184.0-229.0
	2001-2005	340.2	529.8	504.8-555.7	182.2	633.4	592.0-676.9	158.0	461.4	429.5-495.0	132.6	208.5	192.9-225.0	68.4	256.2	229.0-285.6	64.2	179.2	159.9-200.2
	2006-2010	351.8	486.6	463.8-510.3	178.4	531.9	496.7-569.0	173.4	460.3	429.4-492.9	123.4	170.2	156.9-184.4	66.8	214.6	191.7-239.5	56.6	141.3	125.1-159.3
	2011-2015	375.8	479.9	457.8-502.9	199.4	533.7	500.1-569.1	176.4	440.8	410.9-472.5	135.4	170.2	157.4-183.9	70.0	194.9	174.5-217.1	65.4	151.0	134.5-169.1
	2016-2020	415.0	489.1	467.4-511.7	228.4	551.1	518.5-585.5	186.6	442.2	412.5-473.7	138.2	160.5	148.4-173.4	74.8	191.6	172.2-212.8	63.4	137.8	122.4-154.9
Oral cavity and pharynx	1976-1980	4.2	9.3	5.7-14.2	2.6	12.8	6.7-22.0	1.6	6.2	2.6-12.2	1.8	3.8	1.7-7.2	1.6	7.1	3.0-14.1	0.2	0.8	0.0-4.3
	1981-1985	4.6	9.1	5.7-13.7	2.4	11.7	5.8-20.7	2.2	7.6	3.8-13.7	2.0	4.1	1.9-7.6	1.2	5.4	1.9-11.9	0.8	2.7	0.7-7.1
	1986-1990	4.4	8.1	5.0-12.3	4.0	16.0	9.6-25.1	0.4	1.9	0.2-6.3	1.6	3.2	1.4-6.3	1.0	4.5	1.3-10.7	0.6	2.6	0.5-7.4
	1991-1995	6.0	10.5	7.1-15.0	5.4	22.0	14.2-32.4	0.6	2.0	0.4-5.7	2.4	4.3	2.2-7.5	2.2	9.2	4.4-16.6	0.2	0.6	0.0-3.5
	1996-2000	7.0	11.6	8.1-16.2	4.4	15.0	9.2-22.9	2.6	7.9	4.2-13.6	1.8	3.1	1.4-5.8	1.2	5.3	1.8-11.7	0.6	1.8	0.4-5.3
	2001-2005	7.0	10.6	7.4-14.8	4.6	15.8	9.8-23.9	2.4	7.2	3.7-12.7	1.6	2.5	1.1-5.0	1.0	3.3	1.1-7.9	0.6	1.7	0.3-5.2
	2006-2010	7.6	10.5	7.4-14.5	4.0	11.1	6.7-17.3	3.6	9.6	5.6-15.4	1.4	1.8	0.7-3.9	0.8	1.9	0.5-5.4	0.6	1.5	0.3-4.7
	2011-2015	8.6	11.3	8.1-15.5	6.0	15.9	10.6-23.1	2.6	6.9	3.5-12.3	2.8	3.7	2.0-6.4	2.0	5.6	2.6-10.6	0.8	1.8	0.5-5.2
	2016-2020	11.2	13.5	10.1-18.0	8.2	20.0	14.0-27.8	3.0	6.8	3.7-11.7	1.8	1.8	0.8-3.8	1.4	3.1	1.2-7.0	0.4	0.8	0.1-3.8
Esophagus	1976-1980	1.0	2.1	0.7-4.9	0.8	4.3	1.0-11.1	0.2	0.8	0.0-4.2	0.6	1.5	0.3-4.2	0.4	2.8	0.3-9.2	0.2	0.9	0.0-4.6
	1981-1985	1.4	2.5	1.0-5.2	1.0	3.9	1.3-9.4	0.4	1.3	0.2-5.0	1.4	2.6	1.0-5.4	1.2	5.1	1.8-11.3	0.2	0.7	0.0-4.0
	1986-1990	1.8	3.3	1.5-6.2	1.6	6.5	2.8-12.9	0.2	0.6	0.0-3.5	1.8	3.4	1.5-6.4	1.6	7.2	2.9-14.4	0.2	0.6	0.0-3.5
	1991-1995	2.4	4.3	2.2-7.5	2.2	9.4	4.7-16.8	0.2	0.7	0.0-3.6	3.2	5.9	3.4-9.5	2.8	12.5	6.8-20.9	0.4	1.2	0.1-4.4
	1996-2000	1.8	3.1	1.4-5.9	1.2	4.4	1.6-9.6	0.6	1.9	0.4-5.6	2.6	4.5	2.4-7.7	1.8	6.6	3.0-12.5	0.8	2.5	0.7-6.4
	2001-2005	3.4	5.6	3.2-8.9	2.6	9.8	5.1-16.8	0.8	2.5	0.7-6.5	2.4	3.8	1.9-6.6	2.0	7.8	3.6-14.3	0.4	1.1	0.1-4.3
	2006-2010	4.6	6.2	3.9-9.5	3.8	11.1	6.6-17.6	0.8	1.9	0.5-5.4	5.4	7.8	5.1-11.4	4.0	12.7	7.7-19.8	1.4	3.7	1.5-7.9
	2011-2015	5.4	6.8	4.4-10.0	4.8	13.4	8.5-20.3	0.6	1.2	0.2-4.3	3.4	3.9	2.3-6.5	2.4	6.0	3.0-11.0	1.0	2.0	0.6-5.3
	2016-2020	5.8	6.2	4.1-9.2	4.0	9.1	5.5-14.6	1.8	4.0	1.8-8.3	5.4	5.9	3.8-8.8	4.4	10.8	6.7-16.8	1.0	1.9	0.6-5.3
Stomach	1976-1980	3.0	6.5	3.6-10.8	1.8	8.3	3.7-15.9	1.2	4.7	1.7-10.2	2.8	6.2	3.3-10.4	1.6	7.6	3.2-15.0	1.2	4.7	1.7-10.2
	1981-1985	3.0	5.8	3.2-9.6	2.2	10.7	5.2-19.4	0.8	2.6	0.7-6.8	1.8	3.6	1.7-6.9	0.8	3.6	0.9-9.2	1.0	3.7	1.2-8.7
	1986-1990	3.8	7.4	4.4-11.5	2.2	10.0	4.8-18.0	1.6	5.3	2.3-10.6	2.6	4.9	2.6-8.4	1.4	6.6	2.5-13.6	1.2	4.1	1.5-9.1
	1991-1995	3.6	6.7	4.0-10.6	2.4	10.5	5.4-18.3	1.2	4.0	1.5-8.7	1.8	3.3	1.5-6.2	0.8	4.7	1.2-11.4	1.0	3.2	1.0-7.5
	1996-2000	3.4	5.8	3.3-9.2	2.2	8.3	4.0-15.0	1.2	3.9	1.4-8.5	1.2	2.1	0.8-4.5	0.8	3.1	0.8-7.9	0.4	1.2	0.1-4.4
	2001-2005	5.8	9.1	6.1-13.1	4.0	13.4	8.1-20.9	1.8	5.1	2.3-9.8	4.4	6.8	4.2-10.3	2.8	9.0	4.8-15.3	1.6	4.4	1.9-8.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Livingston**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	4.4	6.0	3.7-9.2	2.2	6.3	3.1-11.6	2.2	5.5	2.7-10.2	2.4	3.2	1.7-5.8	1.0	2.4	0.8-6.1	1.4	3.7	1.5-7.9
	2011-2015	4.6	5.6	3.5-8.5	2.4	5.8	3.0-10.6	2.2	5.2	2.5-9.7	2.2	2.7	1.3-5.1	0.8	2.3	0.6-6.2	1.4	3.0	1.2-6.8
	2016-2020	5.4	6.9	4.4-10.3	3.6	9.7	5.6-15.7	1.8	4.9	2.1-9.8	2.8	3.6	1.9-6.3	2.2	6.0	2.9-11.2	0.6	1.5	0.3-4.9
Colorectal	1976-1980	28.6	63.3	53.2-74.7	11.2	57.4	42.8-75.2	17.4	68.0	54.3-84.0	13.2	30.7	23.6-39.1	5.2	29.3	18.8-43.1	8.0	31.3	22.3-42.7
	1981-1985	32.6	66.0	56.2-77.1	14.0	69.5	53.5-88.4	18.6	65.9	53.1-80.9	18.4	38.3	30.8-47.0	9.2	49.2	35.5-66.1	9.2	32.3	23.6-43.1
	1986-1990	38.6	74.0	63.8-85.2	19.4	90.7	72.8-111.4	19.2	63.0	50.9-77.1	17.4	33.4	26.7-41.3	7.4	35.3	24.4-49.1	10.0	33.6	24.9-44.5
	1991-1995	40.2	71.3	61.7-81.8	21.0	90.5	73.4-110.3	19.2	58.5	47.3-71.5	12.2	22.1	16.9-28.3	6.6	30.7	20.7-43.5	5.6	16.8	11.1-24.4
	1996-2000	38.6	66.9	57.8-77.0	19.8	80.7	65.1-98.6	18.8	56.6	45.7-69.4	18.8	32.6	26.4-39.9	11.2	45.9	34.3-59.9	7.6	22.3	15.7-30.7
	2001-2005	43.2	67.3	58.6-77.0	21.6	76.3	62.2-92.5	21.6	60.3	49.4-73.1	12.2	19.1	14.6-24.6	6.4	24.3	16.4-34.5	5.8	15.4	10.3-22.3
	2006-2010	34.6	46.7	39.9-54.3	18.0	54.7	43.7-67.7	16.6	40.5	32.1-50.6	10.6	14.3	10.7-18.8	5.2	17.0	11.0-25.1	5.4	12.2	8.0-18.2
	2011-2015	33.0	41.9	35.5-49.1	18.0	48.8	38.9-60.6	15.0	36.3	28.3-46.2	13.0	16.3	12.5-21.0	7.4	20.4	14.2-28.5	5.6	13.2	8.6-19.6
2016-2020	28.8	35.2	29.4-41.7	17.2	42.9	34.1-53.5	11.6	29.8	22.2-39.2	11.4	14.1	10.6-18.7	5.6	15.6	10.2-22.9	5.8	13.2	8.6-19.8	
Colon excluding rectum	1976-1980	18.4	40.6	32.6-49.9	7.6	39.6	27.5-54.8	10.8	42.0	31.4-55.0	9.0	20.6	14.9-27.6	3.2	18.3	10.2-29.7	5.8	22.4	14.9-32.3
	1981-1985	22.0	45.0	36.9-54.4	7.8	40.1	28.0-55.3	14.2	50.9	39.6-64.3	14.0	29.7	23.1-37.6	6.4	35.5	23.8-50.3	7.6	26.8	18.9-36.9
	1986-1990	27.2	52.4	43.9-62.1	12.0	57.2	43.1-74.1	15.2	49.9	39.2-62.6	14.2	27.3	21.3-34.4	5.4	24.8	16.1-36.3	8.8	29.9	21.6-40.3
	1991-1995	27.6	49.2	41.3-58.1	13.2	58.0	44.3-74.3	14.4	44.0	34.4-55.5	10.8	19.4	14.6-25.4	5.4	25.1	16.1-36.8	5.4	16.1	10.6-23.5
	1996-2000	27.4	47.5	39.9-56.2	13.0	52.5	40.2-67.1	14.4	43.5	33.9-54.9	14.8	25.7	20.2-32.3	8.4	34.5	24.5-46.9	6.4	18.7	12.7-26.5
	2001-2005	30.4	47.9	40.6-56.2	14.4	52.0	40.4-65.7	16.0	44.4	35.1-55.5	9.6	15.0	11.1-20.0	4.8	18.1	11.4-27.1	4.8	12.7	8.1-19.1
	2006-2010	26.0	35.1	29.2-41.7	12.6	39.2	29.9-50.5	13.4	32.4	25.0-41.6	8.4	11.5	8.3-15.7	4.0	14.0	8.4-21.7	4.4	9.8	6.1-15.3
	2011-2015	22.6	29.5	24.2-35.8	12.8	35.9	27.4-46.3	9.8	23.9	17.5-32.0	10.8	13.7	10.2-18.0	6.4	17.7	11.9-25.3	4.4	10.3	6.3-16.1
2016-2020	18.2	22.4	17.9-27.8	10.2	26.2	19.3-34.8	8.0	20.3	14.2-28.3	10.2	12.6	9.3-16.9	5.0	14.2	9.1-21.3	5.2	11.5	7.3-17.6	
Rectum & rectosigmoid	1976-1980	10.2	22.7	16.8-29.9	3.6	17.9	10.3-28.5	6.6	25.9	17.8-36.5	4.2	10.1	6.2-15.4	2.0	11.0	5.1-20.3	2.2	8.9	4.4-15.9
	1981-1985	10.6	21.0	15.7-27.5	6.2	29.4	19.7-42.1	4.4	15.1	9.4-22.9	4.4	8.6	5.3-13.0	2.8	13.8	7.3-23.3	1.6	5.4	2.3-10.7
	1986-1990	11.4	21.5	16.3-27.9	7.4	33.6	23.2-46.8	4.0	13.0	7.9-20.2	3.2	6.1	3.5-10.0	2.0	10.5	4.7-19.5	1.2	3.7	1.4-8.2
	1991-1995	12.6	22.1	16.9-28.2	7.8	32.5	22.8-44.7	4.8	14.4	9.2-21.6	1.4	2.6	1.0-5.3	1.2	5.7	2.0-12.2	0.2	0.8	0.0-3.9
	1996-2000	11.2	19.3	14.6-25.1	6.8	28.2	19.2-39.7	4.4	13.2	8.2-20.0	4.0	6.9	4.2-10.6	2.8	11.4	6.0-19.2	1.2	3.6	1.3-7.9
	2001-2005	12.8	19.4	14.9-24.9	7.2	24.3	16.8-33.9	5.6	15.9	10.6-23.2	2.6	4.1	2.2-7.0	1.6	6.2	2.6-12.3	1.0	2.7	0.9-6.6
	2006-2010	8.6	11.6	8.3-15.8	5.4	15.5	10.1-22.9	3.2	8.0	4.5-13.5	2.2	2.7	1.4-5.0	1.2	3.0	1.1-6.9	1.0	2.4	0.8-6.0
	2011-2015	10.4	12.3	9.1-16.4	5.2	13.0	8.3-19.4	5.2	12.5	7.9-19.0	2.2	2.7	1.3-5.0	1.0	2.7	0.8-6.7	1.2	2.9	1.0-6.9
2016-2020	10.6	12.8	9.5-17.0	7.0	16.8	11.5-23.8	3.6	9.5	5.5-15.5	1.2	1.5	0.5-3.6	0.6	1.4	0.3-4.6	0.6	1.7	0.3-5.6	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Livingston**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	1.8	3.5	1.5-6.7	1.2	5.6	1.8-12.6	0.6	2.0	0.4-6.0	1.0	1.9	0.6-4.6	0.8	3.0	0.8-8.1	0.2	0.8	0.0-4.4
	1981-1985	1.4	3.0	1.2-6.1	1.0	5.2	1.5-12.2	0.4	1.7	0.2-5.9	1.2	2.8	1.0-5.9	0.6	4.6	0.9-12.2	0.6	2.4	0.5-6.9
	1986-1990	0.2	0.3	0.0-1.9	0.2	0.7	0.0-4.2	0.0	0.0	0.0-2.5	0.4	0.6	0.1-2.4	0.2	0.7	0.0-4.2	0.2	0.6	0.0-3.5
	1991-1995	2.0	3.5	1.7-6.3	1.6	6.9	2.8-13.7	0.4	1.3	0.2-4.7	2.6	4.6	2.5-7.9	1.0	4.7	1.4-10.9	1.6	5.0	2.2-10.0
	1996-2000	2.8	4.8	2.6-8.0	1.8	7.0	3.1-13.4	1.0	2.9	0.9-6.9	2.8	4.8	2.6-8.1	1.4	5.2	2.1-10.7	1.4	3.7	1.5-8.0
	2001-2005	2.2	3.4	1.7-6.1	1.6	4.9	2.1-9.8	0.6	2.1	0.4-6.0	1.8	2.7	1.2-5.1	1.0	3.3	1.0-7.8	0.8	2.2	0.6-5.9
	2006-2010	4.8	6.8	4.3-10.2	2.0	5.6	2.7-10.6	2.8	7.5	4.1-12.9	3.4	4.7	2.7-7.6	2.2	7.0	3.5-12.8	1.2	3.0	1.1-6.9
	2011-2015	4.6	5.9	3.7-9.1	2.6	6.8	3.6-12.0	2.0	5.1	2.3-10.0	4.8	6.0	3.8-9.1	2.8	7.4	4.0-12.8	2.0	4.6	2.1-8.9
2016-2020	7.0	7.6	5.2-10.8	5.6	12.7	8.3-18.8	1.4	3.0	1.2-6.9	5.6	6.4	4.2-9.4	4.2	10.4	6.3-16.3	1.4	3.1	1.2-7.0	
Pancreas	1976-1980	5.4	11.2	7.3-16.3	2.8	14.3	7.5-24.3	2.6	9.0	4.8-15.5	4.0	8.6	5.2-13.4	2.0	11.5	5.2-21.2	2.0	7.2	3.4-13.3
	1981-1985	8.4	17.2	12.4-23.3	4.2	21.5	13.0-33.1	4.2	15.2	9.3-23.3	7.6	15.6	11.0-21.5	4.0	20.6	12.3-32.1	3.6	12.6	7.4-20.0
	1986-1990	8.0	15.6	11.1-21.3	4.2	17.9	10.9-27.6	3.8	12.8	7.6-20.1	5.6	11.1	7.4-16.1	3.0	13.4	7.3-22.3	2.6	8.7	4.6-15.0
	1991-1995	6.2	11.1	7.5-15.7	3.0	12.9	7.0-21.5	3.2	10.0	5.7-16.3	6.0	10.7	7.2-15.3	2.6	11.2	5.7-19.4	3.4	10.6	6.1-17.0
	1996-2000	5.8	10.0	6.7-14.4	3.2	13.0	7.3-21.2	2.6	7.9	4.2-13.6	6.0	10.4	7.0-14.8	2.6	10.7	5.6-18.4	3.4	9.9	5.7-15.9
	2001-2005	8.8	13.7	9.9-18.4	4.8	17.1	10.9-25.6	4.0	10.8	6.6-16.9	8.2	12.8	9.2-17.4	4.6	17.5	10.9-26.4	3.6	9.6	5.7-15.4
	2006-2010	5.2	7.0	4.6-10.4	2.6	8.7	4.5-15.0	2.6	6.5	3.5-11.5	4.8	6.6	4.2-10.0	2.2	7.1	3.5-12.9	2.6	6.8	3.6-11.9
	2011-2015	8.0	9.8	7.0-13.5	4.0	10.6	6.4-16.7	4.0	9.0	5.4-14.5	7.8	9.0	6.4-12.5	3.4	8.6	4.9-14.2	4.4	9.1	5.6-14.2
2016-2020	9.8	11.1	8.2-14.9	5.4	13.1	8.5-19.5	4.4	9.1	5.7-14.4	8.6	9.7	7.0-13.3	4.6	11.6	7.2-17.8	4.0	8.0	4.8-12.9	
Larynx	1976-1980	2.8	5.8	3.1-9.8	2.0	9.1	4.3-16.8	0.8	3.0	0.8-7.8	1.0	1.9	0.6-4.5	0.6	2.7	0.5-8.2	0.4	1.3	0.2-5.1
	1981-1985	2.4	5.0	2.6-8.8	2.0	10.3	4.7-19.1	0.4	1.8	0.2-6.1	0.2	0.4	0.0-2.1	0.2	0.8	0.0-4.6	0.0	0.0	0.0-2.7
	1986-1990	3.8	7.3	4.4-11.5	3.2	14.1	8.0-22.9	0.6	1.9	0.4-5.7	0.4	0.7	0.1-2.5	0.4	1.4	0.2-5.3	0.0	0.0	0.0-2.5
	1991-1995	3.6	6.4	3.8-10.2	3.0	11.0	6.1-18.3	0.6	2.1	0.4-6.0	1.0	1.8	0.6-4.2	1.0	3.8	1.2-8.8	0.0	0.0	0.0-2.4
	1996-2000	2.8	4.8	2.6-8.0	2.4	8.9	4.5-15.5	0.4	1.3	0.2-4.6	0.4	0.7	0.1-2.5	0.4	1.4	0.2-5.1	0.0	0.0	0.0-2.4
	2001-2005	4.0	6.2	3.8-9.7	3.2	10.8	6.1-17.6	0.8	2.4	0.7-6.3	1.4	2.1	0.9-4.4	1.0	3.0	1.0-7.3	0.4	1.2	0.1-4.5
	2006-2010	2.2	3.4	1.7-6.1	1.8	6.0	2.7-11.5	0.4	1.0	0.1-4.1	1.2	1.7	0.6-3.8	0.8	2.6	0.7-6.8	0.4	0.8	0.1-3.6
	2011-2015	2.6	3.1	1.6-5.5	2.0	4.8	2.3-9.3	0.6	1.6	0.3-5.1	1.4	1.8	0.7-4.0	1.2	3.6	1.3-8.1	0.2	0.5	0.0-3.3
2016-2020	2.2	2.4	1.2-4.5	1.6	3.7	1.6-7.8	0.6	1.2	0.2-4.4	0.6	0.7	0.1-2.3	0.2	0.4	0.0-3.1	0.4	0.9	0.1-4.1	
Lung and bronchus	1976-1980	32.0	67.2	57.0-78.6	23.6	113.6	93.4-136.9	8.4	32.2	23.1-43.6	24.0	50.7	41.9-60.8	18.6	88.4	70.7-109.1	5.4	22.0	14.4-32.0
	1981-1985	33.8	68.1	58.1-79.3	21.8	100.1	81.5-121.5	12.0	45.7	34.6-59.0	27.6	56.1	47.0-66.4	18.2	85.0	67.8-105.0	9.4	36.4	26.5-48.5
	1986-1990	37.6	70.6	60.8-81.5	23.2	102.0	83.8-122.8	14.4	49.7	38.7-62.7	31.8	59.0	50.1-69.0	20.0	88.8	71.7-108.6	11.8	40.5	30.7-52.4
	1991-1995	43.2	77.5	67.4-88.5	27.2	110.9	92.7-131.6	16.0	53.7	42.5-66.9	37.2	66.9	57.6-77.2	23.0	97.2	79.8-117.2	14.2	46.6	36.4-58.9
	1996-2000	42.2	72.4	63.0-82.9	25.2	97.5	80.9-116.4	17.0	55.0	43.9-68.0	38.0	65.8	56.8-75.9	23.8	94.1	77.7-112.9	14.2	45.4	35.4-57.3
	2001-2005	47.2	74.1	64.9-84.2	27.4	95.4	79.8-113.1	19.8	57.7	46.9-70.4	40.0	63.1	54.6-72.5	22.2	80.8	66.2-97.6	17.8	51.5	41.3-63.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Livingston**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	55.6	76.4	67.6-86.0	28.8	87.1	73.1-102.9	26.8	67.5	56.4-80.3	37.0	50.8	43.7-58.8	20.4	62.9	51.0-76.7	16.6	42.2	33.5-52.6
	2011-2015	56.0	70.5	62.3-79.5	32.0	87.2	73.9-102.4	24.0	57.4	47.5-69.1	38.0	46.8	40.3-54.2	21.8	57.9	47.3-70.4	16.2	37.4	29.6-46.9
	2016-2020	59.0	65.8	58.4-74.1	26.4	62.0	51.6-74.1	32.6	69.4	58.9-81.5	39.0	44.0	37.9-50.8	19.2	46.5	37.5-57.3	19.8	41.9	33.9-51.5
Melanoma of the skin	1976-1980	2.6	5.5	2.9-9.4	1.6	7.3	2.9-14.8	1.0	4.0	1.3-9.4	1.6	3.5	1.5-6.9	1.2	5.9	2.1-12.9	0.4	1.4	0.2-5.3
	1981-1985	4.2	8.4	5.2-12.9	2.4	10.1	5.2-17.8	1.8	6.5	2.9-12.3	1.4	3.1	1.2-6.3	0.6	2.9	0.6-8.2	0.8	3.0	0.8-7.8
	1986-1990	6.4	11.4	7.8-16.2	3.6	15.6	9.0-25.0	2.8	8.5	4.6-14.4	2.0	3.5	1.7-6.5	1.0	4.4	1.3-10.6	1.0	3.2	1.0-7.6
	1991-1995	5.0	8.6	5.5-12.7	3.8	13.7	8.2-21.5	1.2	4.0	1.5-8.7	1.8	3.2	1.5-6.1	1.4	5.6	2.2-11.5	0.4	1.4	0.2-4.8
	1996-2000	7.8	13.0	9.2-17.7	4.6	17.1	10.7-25.8	3.2	10.2	5.8-16.7	2.4	4.0	2.1-6.9	1.6	5.9	2.4-11.9	0.8	2.5	0.7-6.5
	2001-2005	9.8	15.1	11.1-20.0	6.0	20.0	13.4-28.7	3.8	12.1	7.2-19.0	1.4	2.1	0.9-4.5	0.8	3.0	0.7-7.7	0.6	1.7	0.3-5.1
	2006-2010	11.4	16.9	12.7-22.1	6.4	18.7	12.7-26.7	5.0	15.7	10.0-23.5	1.2	1.6	0.6-3.6	1.2	3.2	1.2-7.3	0.0	0.0	0.0-2.4
	2011-2015	20.0	27.0	21.8-33.2	12.8	35.9	27.4-46.3	7.2	19.7	13.3-28.0	2.4	3.3	1.7-5.9	1.8	5.6	2.5-10.7	0.6	1.5	0.3-4.9
	2016-2020	22.0	26.7	21.7-32.6	14.0	35.9	27.7-46.0	8.0	19.1	13.3-26.8	2.8	3.2	1.7-5.6	1.8	4.6	2.0-9.2	1.0	1.8	0.6-5.1
Female breast	1976-1980							23.2	92.2	75.9-110.9							8.8	35.6	25.7-47.8
	1981-1985							32.6	126.5	107.4-147.8							10.4	38.4	28.5-50.6
	1986-1990							35.2	124.7	106.7-144.9							10.6	36.8	27.5-48.3
	1991-1995							41.6	139.2	120.7-159.5							7.4	24.0	16.8-33.1
	1996-2000							42.6	136.5	118.7-156.2							11.6	36.2	27.4-46.9
	2001-2005							45.4	133.8	116.9-152.6							8.2	22.8	16.3-31.2
	2006-2010							50.6	134.9	118.5-153.1							7.6	19.8	13.9-27.5
	2011-2015							54.6	138.0	121.4-156.4							9.2	20.2	14.5-27.6
	2016-2020							58.2	144.2	127.0-163.3							7.8	18.3	12.7-25.7
Cervix uteri	1976-1980							3.0	12.2	6.8-20.0							0.8	3.4	0.9-8.5
	1981-1985							2.2	7.3	3.6-13.3							0.6	2.4	0.5-7.0
	1986-1990							2.8	9.7	5.2-16.3							0.8	2.5	0.7-6.5
	1991-1995							2.4	7.7	4.0-13.6							0.6	2.1	0.4-6.1
	1996-2000							2.0	6.1	2.9-11.3							1.0	3.0	1.0-7.1
	2001-2005							1.8	5.5	2.5-10.6							1.2	3.4	1.3-7.6
	2006-2010							3.0	10.1	5.5-16.9							0.0	0.0	0.0-2.4
	2011-2015							2.0	7.8	3.6-14.4							1.4	4.0	1.5-8.8
	2016-2020							2.0	7.9	3.6-14.9							0.8	3.5	0.8-8.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Livingston**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							8.4	32.3	23.1-43.8							2.4	9.2	4.7-16.2
	1981-1985							8.2	29.2	20.8-39.8							2.6	8.5	4.5-14.7
	1986-1990							9.4	32.2	23.6-43.0							1.4	4.6	1.8-9.5
	1991-1995							9.4	31.1	22.8-41.4							2.0	6.1	2.9-11.3
	1996-2000							11.2	36.1	27.3-47.0							2.8	8.6	4.7-14.5
	2001-2005							9.4	28.0	20.5-37.3							1.2	3.0	1.1-6.9
	2006-2010							9.8	25.4	18.7-33.9							1.0	2.8	0.9-6.7
	2011-2015							11.8	28.2	21.2-36.9							0.8	1.7	0.4-5.0
	2016-2020							16.8	38.7	30.4-48.7							2.4	4.9	2.5-9.2
Ovary	1976-1980							3.4	13.9	8.0-22.3							3.2	12.1	6.8-19.7
	1981-1985							3.0	10.9	6.0-18.2							1.4	5.3	2.1-10.9
	1986-1990							5.2	18.0	11.7-26.5							3.2	10.7	6.1-17.4
	1991-1995							4.8	15.7	10.0-23.5							2.6	8.5	4.5-14.5
	1996-2000							4.6	14.7	9.3-22.1							2.8	8.7	4.7-14.7
	2001-2005							6.4	18.6	12.7-26.4							3.4	9.6	5.6-15.6
	2006-2010							6.2	15.9	10.7-22.8							3.6	8.5	5.0-13.8
	2011-2015							6.4	16.4	11.0-23.8							5.2	11.7	7.6-17.5
	2016-2020							3.0	6.6	3.6-11.5							3.4	7.5	4.3-12.7
Prostate	1976-1980				16.8	93.9	73.9-117.2							5.4	36.1	23.3-52.5			
	1981-1985				20.2	106.9	86.2-130.7							5.8	33.9	22.3-48.7			
	1986-1990				32.0	157.9	133.4-185.1							7.2	39.3	27.0-54.6			
	1991-1995				50.6	214.9	188.5-243.8							7.0	37.0	25.4-51.6			
	1996-2000				43.0	170.2	147.6-195.0							6.2	31.1	20.9-44.1			
	2001-2005				52.0	178.5	157.1-201.9							6.8	30.4	20.9-42.4			
	2006-2010				54.0	155.7	137.3-176.0							6.4	23.6	16.1-33.3			
	2011-2015				53.6	128.7	113.4-145.7							4.2	12.8	7.9-19.8			
	2016-2020				62.6	137.1	121.9-153.8							7.0	18.1	12.5-25.5			
Testis	1976-1980				1.4	5.2	2.0-11.3							0.6	1.5	0.3-5.5			
	1981-1985				1.4	4.9	1.9-10.7							0.0	0.0	0.0-3.3			
	1986-1990				1.2	4.2	1.5-9.5							0.0	0.0	0.0-3.0			
	1991-1995				1.8	5.4	2.5-10.6							0.0	0.0	0.0-2.8			
	1996-2000				2.0	5.6	2.7-10.6							0.4	1.1	0.1-4.4			
	2001-2005				2.2	6.4	3.2-11.7							0.0	0.0	0.0-2.6			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Livingston**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				1.4	4.7	1.8-9.8							0.0	0.0	0.0-2.5			
	2011-2015				1.4	5.0	2.0-10.4							0.0	0.0	0.0-2.5			
	2016-2020				2.6	7.8	4.0-13.8							0.0	0.0	0.0-2.6			
Urinary bladder (incl. in situ)	1976-1980	8.6	19.0	13.7-25.7	5.4	28.2	18.1-41.7	3.2	12.5	7.1-20.3	2.8	6.8	3.7-11.3	1.4	7.0	2.8-14.5	1.4	5.7	2.3-11.6
	1981-1985	11.4	23.7	17.9-30.8	8.4	42.8	30.3-58.4	3.0	11.0	6.1-18.2	2.4	5.3	2.7-9.2	1.6	9.0	3.8-17.6	0.8	2.9	0.8-7.5
	1986-1990	11.4	22.1	16.7-28.7	8.6	43.6	31.0-59.1	2.8	8.7	4.7-14.7	2.6	5.2	2.8-8.9	2.4	12.9	6.5-22.4	0.2	0.7	0.0-3.7
	1991-1995	15.2	27.3	21.5-34.2	12.0	52.2	39.4-67.6	3.2	9.7	5.6-15.9	3.0	5.5	3.1-9.1	2.2	13.2	6.4-23.3	0.8	2.3	0.6-6.1
	1996-2000	16.2	27.9	22.1-34.6	13.0	50.8	38.9-65.0	3.2	9.9	5.6-16.2	2.6	4.5	2.4-7.7	1.8	7.6	3.4-14.4	0.8	2.4	0.6-6.2
	2001-2005	21.8	33.9	27.8-40.9	17.0	61.8	49.0-76.8	4.8	13.7	8.8-20.6	3.0	4.9	2.7-8.1	1.8	7.2	3.2-13.5	1.2	3.2	1.2-7.2
	2006-2010	18.4	24.9	20.0-30.6	13.2	41.2	31.6-52.8	5.2	13.8	8.9-20.6	4.2	5.8	3.6-9.0	3.0	11.3	6.3-18.6	1.2	2.6	0.9-6.2
	2011-2015	19.6	24.8	20.1-30.5	15.0	42.8	33.5-54.0	4.6	10.8	6.7-16.7	5.2	6.4	4.2-9.6	4.2	12.4	7.6-19.2	1.0	2.3	0.7-5.9
	2016-2020	21.0	23.9	19.4-29.3	18.0	43.7	34.9-54.3	3.0	5.8	3.2-10.2	4.0	4.4	2.7-7.1	3.2	8.2	4.6-13.7	0.8	1.6	0.4-4.9
Kidney and renal pelvis	1976-1980	3.4	7.4	4.3-11.9	1.8	8.8	3.9-16.7	1.6	6.3	2.7-12.5	1.8	4.2	1.9-7.8	1.0	6.0	1.9-13.8	0.8	3.1	0.8-8.0
	1981-1985	3.4	6.4	3.7-10.3	1.6	6.3	2.7-12.6	1.8	6.1	2.8-11.7	2.0	4.0	1.9-7.4	0.8	3.3	0.9-8.7	1.2	4.5	1.6-9.8
	1986-1990	4.0	7.6	4.6-11.7	3.0	13.4	7.3-22.2	1.0	3.2	1.0-7.5	1.0	1.8	0.6-4.2	0.6	2.8	0.6-8.0	0.4	1.2	0.1-4.6
	1991-1995	5.8	10.1	6.8-14.6	3.6	13.5	8.0-21.4	2.2	7.0	3.5-12.6	2.0	3.5	1.7-6.5	1.2	4.5	1.7-9.9	0.8	2.4	0.6-6.2
	1996-2000	8.0	13.4	9.5-18.2	5.4	19.1	12.5-28.0	2.6	8.4	4.5-14.4	2.0	3.4	1.6-6.2	2.0	7.7	3.6-14.3	0.0	0.0	0.0-2.4
	2001-2005	11.6	18.0	13.7-23.3	6.6	22.4	15.3-31.7	5.0	14.4	9.3-21.4	4.4	6.9	4.3-10.5	2.4	8.9	4.5-15.7	2.0	5.5	2.6-10.3
	2006-2010	11.4	16.2	12.2-21.2	5.0	14.2	9.1-21.3	6.4	18.9	12.7-27.1	3.0	4.0	2.2-6.7	1.6	4.6	1.9-9.4	1.4	3.8	1.5-8.0
	2011-2015	12.6	16.3	12.4-21.1	9.0	24.2	17.4-32.8	3.6	9.0	5.2-14.8	2.0	2.9	1.4-5.5	1.2	3.5	1.2-8.0	0.8	2.3	0.6-6.3
	2016-2020	18.6	22.3	17.8-27.7	12.4	30.7	23.3-39.9	6.2	14.9	9.7-22.1	2.2	2.6	1.3-5.0	1.6	4.1	1.7-8.5	0.6	1.6	0.3-5.2
Brain and other central nervous system	1976-1980	2.6	5.3	2.8-9.2	1.6	7.3	2.9-14.8	1.0	4.1	1.3-9.7	1.8	3.7	1.7-7.0	1.2	5.5	2.0-12.1	0.6	2.0	0.4-6.0
	1981-1985	3.8	7.2	4.3-11.4	1.8	7.3	3.1-14.3	2.0	7.1	3.3-13.1	2.4	4.6	2.3-8.0	1.2	4.5	1.6-10.1	1.2	4.2	1.5-9.2
	1986-1990	3.6	6.5	3.8-10.3	1.6	6.3	2.7-12.6	2.0	6.4	3.0-12.0	2.2	4.0	2.0-7.2	1.0	4.0	1.3-9.3	1.2	3.8	1.4-8.4
	1991-1995	4.2	7.3	4.5-11.1	2.4	8.4	4.3-14.8	1.8	6.0	2.7-11.3	2.4	4.2	2.1-7.3	1.2	4.2	1.5-9.2	1.2	4.0	1.5-8.7
	1996-2000	5.8	9.4	6.2-13.5	3.2	10.2	5.8-16.7	2.6	7.8	4.1-13.5	2.6	4.4	2.4-7.6	1.2	3.9	1.4-8.6	1.4	4.6	1.8-9.4
	2001-2005	2.8	4.4	2.4-7.4	1.2	3.6	1.3-8.0	1.6	4.7	2.0-9.4	3.0	4.6	2.6-7.7	1.8	5.4	2.5-10.5	1.2	3.3	1.2-7.3
	2006-2010	4.6	6.3	3.9-9.5	2.6	7.4	3.9-13.0	2.0	5.6	2.6-10.6	4.4	5.9	3.7-9.1	2.8	8.6	4.6-14.6	1.6	3.8	1.6-7.8
	2011-2015	6.6	9.1	6.2-13.0	3.2	9.2	5.1-15.3	3.4	9.2	5.2-15.1	4.4	5.7	3.5-8.9	1.8	4.6	2.0-9.0	2.6	6.7	3.4-12.0
	2016-2020	6.6	9.9	6.6-14.1	3.8	11.4	6.7-18.2	2.8	7.9	4.1-14.0	4.0	5.2	3.1-8.4	1.8	5.0	2.3-9.8	2.2	5.4	2.5-10.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Livingston**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	1.4	2.7	1.0-5.7	0.2	0.7	0.0-4.7	1.2	4.2	1.5-9.3	0.2	0.6	0.0-2.8	0.2	1.2	0.0-5.8	0.0	0.0	0.0-2.8
	1981-1985	2.2	3.9	1.9-7.1	0.0	0.0	0.0-3.3	2.2	7.3	3.5-13.4	0.4	1.1	0.1-3.5	0.2	1.1	0.0-5.5	0.2	1.0	0.0-5.0
	1986-1990	2.2	3.5	1.7-6.4	0.8	2.4	0.6-6.6	1.4	4.5	1.8-9.4	0.2	0.3	0.0-1.9	0.2	0.6	0.0-4.1	0.0	0.0	0.0-2.5
	1991-1995	3.8	6.3	3.8-9.9	1.4	4.7	1.9-9.8	2.4	7.8	4.0-13.7	0.4	0.8	0.1-2.7	0.0	0.0	0.0-2.8	0.4	1.3	0.2-4.6
	1996-2000	2.0	3.0	1.4-5.6	0.2	1.3	0.0-5.9	1.8	5.4	2.5-10.4	0.2	0.4	0.0-1.9	0.2	1.3	0.0-5.9	0.0	0.0	0.0-2.4
	2001-2005	4.2	6.3	3.9-9.7	0.6	2.1	0.4-6.1	3.6	10.8	6.3-17.2	0.4	0.7	0.1-2.6	0.0	0.0	0.0-2.6	0.4	1.6	0.2-5.4
	2006-2010	8.2	12.8	9.1-17.5	3.2	10.2	5.8-16.7	5.0	16.4	10.4-24.4	0.2	0.3	0.0-1.8	0.2	0.7	0.0-3.7	0.0	0.0	0.0-2.4
	2011-2015	7.8	11.6	8.1-16.2	2.4	6.9	3.5-12.4	5.4	16.7	10.6-24.9	0.0	0.0	0.0-1.3	0.0	0.0	0.0-2.5	0.0	0.0	0.0-2.6
2016-2020	7.4	10.1	6.9-14.3	3.0	8.5	4.6-14.4	4.4	12.1	7.3-19.1	0.2	0.2	0.0-1.6	0.0	0.0	0.0-2.6	0.2	0.4	0.0-3.2	
Hodgkin lymphoma	1976-1980	2.4	4.6	2.2-8.3	1.4	5.8	2.1-12.5	1.0	3.4	1.0-8.3	0.8	1.6	0.4-4.2	0.8	3.8	0.9-10.0	0.0	0.0	0.0-2.8
	1981-1985	1.4	2.0	0.8-4.5	0.4	1.3	0.2-5.2	1.0	2.6	0.7-6.9	0.2	0.3	0.0-1.9	0.2	0.6	0.0-4.1	0.0	0.0	0.0-2.7
	1986-1990	1.6	2.3	1.0-4.6	1.0	2.8	0.9-7.2	0.6	1.8	0.4-5.5	0.6	0.9	0.2-2.8	0.4	1.1	0.1-4.7	0.2	0.7	0.0-3.9
	1991-1995	2.0	3.1	1.5-5.8	0.6	1.9	0.4-5.9	1.4	4.1	1.6-8.5	0.2	0.2	0.0-1.6	0.0	0.0	0.0-2.8	0.2	0.4	0.0-3.0
	1996-2000	2.0	3.2	1.5-5.9	1.0	3.2	1.0-7.6	1.0	3.0	1.0-7.1	0.8	1.3	0.4-3.4	0.4	1.2	0.1-4.7	0.4	1.2	0.1-4.4
	2001-2005	1.6	2.4	1.0-4.9	1.2	3.3	1.2-7.4	0.4	1.7	0.2-5.6	0.4	0.7	0.1-2.4	0.4	1.8	0.2-6.2	0.0	0.0	0.0-2.4
	2006-2010	1.2	1.5	0.5-3.5	0.6	1.5	0.3-4.8	0.6	1.5	0.3-4.9	0.0	0.0	0.0-1.2	0.0	0.0	0.0-2.5	0.0	0.0	0.0-2.4
	2011-2015	1.2	1.6	0.6-3.7	1.2	3.4	1.2-7.7	0.0	0.0	0.0-2.6	0.0	0.0	0.0-1.3	0.0	0.0	0.0-2.5	0.0	0.0	0.0-2.6
2016-2020	1.8	2.6	1.1-5.2	1.0	3.0	0.9-7.4	0.8	2.4	0.6-6.5	0.2	0.2	0.0-1.6	0.2	0.4	0.0-3.1	0.0	0.0	0.0-2.7	
Non-Hodgkin lymphomas	1976-1980	8.2	17.6	12.5-23.9	4.4	20.4	12.6-31.3	3.8	14.3	8.6-22.5	2.8	5.8	3.1-9.7	2.2	10.1	4.9-18.4	0.6	2.3	0.5-6.8
	1981-1985	6.4	12.9	8.8-18.2	3.2	15.9	8.8-26.0	3.2	11.2	6.4-18.3	2.8	5.9	3.2-9.8	1.0	4.4	1.4-10.3	1.8	7.3	3.3-13.8
	1986-1990	8.6	16.3	11.7-21.9	3.0	11.9	6.5-19.8	5.6	20.3	13.4-29.3	3.4	6.7	3.9-10.8	1.6	8.4	3.4-16.5	1.8	6.5	2.9-12.2
	1991-1995	12.0	20.5	15.6-26.5	6.8	26.5	18.0-37.4	5.2	16.4	10.7-24.1	4.4	7.7	4.8-11.7	2.0	9.0	4.1-16.8	2.4	7.5	3.9-13.2
	1996-2000	11.0	19.0	14.3-24.7	6.0	23.3	15.6-33.3	5.0	15.2	9.8-22.6	7.4	12.6	8.9-17.4	3.6	14.7	8.6-23.2	3.8	11.2	6.7-17.6
	2001-2005	13.8	21.2	16.5-26.9	6.8	22.4	15.4-31.6	7.0	19.8	13.8-27.7	4.2	6.6	4.1-10.2	1.6	5.7	2.4-11.2	2.6	7.2	3.8-12.5
	2006-2010	13.8	19.7	15.2-25.0	8.2	25.4	18.0-34.7	5.6	14.7	9.7-21.5	5.4	7.1	4.7-10.5	3.8	11.5	6.8-18.2	1.6	3.6	1.5-7.4
	2011-2015	15.0	18.5	14.5-23.4	8.4	22.7	16.2-31.0	6.6	15.0	10.2-21.5	4.0	5.1	3.1-8.0	2.8	8.6	4.7-14.7	1.2	2.7	1.0-6.5
2016-2020	14.8	17.4	13.5-22.1	8.6	21.9	15.6-30.0	6.2	13.8	9.1-20.2	3.6	3.9	2.3-6.4	2.4	6.5	3.3-11.7	1.2	2.1	0.8-5.4	
Myeloma	1976-1980	2.2	4.8	2.3-8.5	1.4	6.0	2.3-12.6	0.8	3.1	0.8-7.9	1.0	2.1	0.7-4.9	0.8	3.3	0.9-8.7	0.2	0.8	0.0-4.4
	1981-1985	2.0	3.8	1.8-7.1	1.0	3.7	1.2-9.0	1.0	3.8	1.2-8.8	1.2	2.5	0.9-5.3	0.6	3.3	0.6-9.3	0.6	2.0	0.4-6.0
	1986-1990	2.6	4.7	2.5-8.1	1.8	6.9	3.1-13.2	0.8	2.9	0.8-7.5	2.0	3.7	1.7-6.7	1.2	4.7	1.7-10.3	0.8	2.5	0.7-6.5
	1991-1995	2.4	4.3	2.2-7.5	0.6	2.5	0.5-7.1	1.8	5.2	2.4-10.1	1.8	3.3	1.5-6.1	0.6	2.0	0.4-5.9	1.2	3.3	1.2-7.4
	1996-2000	3.6	6.2	3.7-9.8	2.4	9.0	4.6-15.8	1.2	3.7	1.4-8.2	3.0	5.2	2.9-8.5	1.6	6.1	2.6-12.0	1.4	4.3	1.7-8.9
	2001-2005	4.2	6.7	4.1-10.3	3.0	11.3	6.2-18.6	1.2	3.2	1.2-7.3	2.4	3.9	2.0-6.8	1.6	6.8	2.9-13.2	0.8	2.2	0.6-5.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Livingston**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	5.2	6.9	4.5-10.3	2.8	8.3	4.5-14.2	2.4	5.8	3.0-10.5	3.0	4.4	2.5-7.3	2.2	7.8	3.9-14.0	0.8	2.1	0.6-5.8
	2011-2015	3.4	4.3	2.4-7.0	1.0	2.6	0.8-6.4	2.4	6.0	3.0-10.9	1.4	1.6	0.6-3.6	0.6	1.5	0.3-4.9	0.8	1.7	0.5-5.1
	2016-2020	5.4	6.5	4.2-9.7	4.0	10.7	6.4-16.9	1.4	3.2	1.2-7.2	1.6	1.9	0.8-3.9	0.8	2.1	0.6-5.9	0.8	1.8	0.5-5.2
Leukemias	1976-1980	7.2	14.4	10.0-20.1	3.6	15.6	9.0-25.3	3.6	12.9	7.5-20.6	4.0	8.5	5.1-13.3	2.2	12.0	5.7-21.5	1.8	6.1	2.7-11.8
	1981-1985	6.6	14.0	9.6-19.7	3.8	19.1	11.2-30.1	2.8	10.4	5.7-17.6	3.4	7.2	4.1-11.6	2.0	9.1	4.2-17.1	1.4	5.2	2.1-10.8
	1986-1990	9.0	16.8	12.2-22.5	4.8	21.0	13.2-31.4	4.2	14.2	8.7-21.8	5.4	10.5	6.9-15.2	2.6	13.4	7.0-22.7	2.8	9.2	5.0-15.6
	1991-1995	7.2	12.7	8.9-17.6	3.8	16.1	9.4-25.4	3.4	10.7	6.2-17.2	3.4	6.1	3.6-9.8	1.6	8.0	3.3-15.8	1.8	5.6	2.6-10.7
	1996-2000	10.8	18.1	13.6-23.7	6.6	23.7	16.1-33.5	4.2	12.9	8.0-19.8	5.8	9.8	6.6-14.1	2.8	10.7	5.8-17.9	3.0	8.9	4.9-14.7
	2001-2005	10.8	17.5	13.1-22.9	5.4	19.2	12.5-28.2	5.4	16.7	10.9-24.5	4.8	7.6	4.9-11.3	2.6	9.6	5.0-16.5	2.2	6.0	3.0-11.0
	2006-2010	9.6	13.6	9.9-18.1	5.0	14.6	9.4-21.9	4.6	11.7	7.3-17.9	4.0	5.7	3.4-8.8	1.2	3.9	1.4-8.5	2.8	6.7	3.6-11.6
	2011-2015	13.8	18.1	14.0-23.2	7.2	20.6	14.3-28.9	6.6	15.5	10.4-22.5	6.8	9.0	6.2-12.7	3.2	9.4	5.3-15.6	3.6	9.1	5.2-14.8
2016-2020	15.4	18.3	14.3-23.2	10.6	26.7	19.8-35.3	4.8	11.1	6.8-17.3	4.8	5.4	3.4-8.3	2.4	6.1	3.1-11.1	2.4	4.8	2.4-9.0	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Madison**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	218.8	422.9	397.6-449.2	109.6	502.5	458.8-548.9	109.2	373.8	342.7-406.9	115.2	224.4	206.0-243.9	62.2	287.4	254.5-323.0	53.0	182.9	161.3-206.6
	1981-1985	237.0	425.2	400.9-450.5	121.0	520.8	477.8-566.4	116.0	370.9	340.9-402.8	119.4	216.8	199.5-235.2	63.8	280.0	248.4-314.3	55.6	175.5	155.2-197.6
	1986-1990	282.6	474.5	449.8-500.2	147.6	589.6	545.8-635.8	135.0	405.5	375.1-437.6	137.2	233.7	216.4-252.0	74.6	305.7	273.9-339.9	62.6	185.2	165.1-207.2
	1991-1995	304.8	478.0	454.2-502.8	159.6	584.7	543.5-628.2	145.2	413.7	383.9-445.2	130.2	206.0	190.4-222.5	72.0	277.9	248.8-309.4	58.2	160.2	142.2-179.8
	1996-2000	351.0	516.4	492.5-541.1	179.4	592.6	553.7-633.5	171.6	464.8	434.1-497.1	149.4	221.8	206.2-238.3	77.8	272.6	245.5-301.8	71.6	188.5	169.4-209.2
	2001-2005	374.0	507.8	484.9-531.5	197.2	595.4	558.0-634.6	176.8	447.4	418.1-478.1	146.8	201.3	187.0-216.5	75.6	248.6	223.5-275.7	71.2	173.2	155.6-192.4
	2006-2010	433.6	533.8	511.2-557.1	243.2	646.5	609.8-684.9	190.4	443.0	414.8-472.7	143.6	178.0	165.1-191.6	75.6	218.0	196.1-241.8	68.0	151.9	136.0-169.3
	2011-2015	430.8	485.7	464.7-507.3	225.4	533.9	502.2-567.3	205.4	448.0	420.1-477.5	143.6	160.4	148.7-172.9	73.0	183.9	165.1-204.5	70.6	142.9	128.0-159.2
	2016-2020	448.2	464.4	444.4-485.0	244.6	521.0	491.0-552.5	203.6	418.3	391.6-446.6	141.4	143.9	133.2-155.4	78.6	174.5	157.1-193.4	62.8	119.2	105.9-133.9
Oral cavity and pharynx	1976-1980	7.2	12.9	9.0-18.0	4.2	17.5	10.5-27.3	3.0	10.0	5.5-16.5	2.8	5.2	2.8-8.8	2.2	9.4	4.5-17.0	0.6	2.0	0.4-5.9
	1981-1985	5.4	9.0	5.9-13.1	4.2	16.7	10.1-25.9	1.2	3.6	1.3-8.0	2.0	3.4	1.6-6.3	1.4	5.4	2.0-11.6	0.6	2.0	0.4-5.8
	1986-1990	6.8	11.1	7.7-15.6	4.8	19.3	11.9-29.3	2.0	5.7	2.7-10.6	2.6	4.1	2.2-7.0	1.6	5.7	2.4-11.5	1.0	2.9	0.9-6.9
	1991-1995	6.2	9.6	6.5-13.6	4.6	16.6	10.3-25.0	1.6	4.3	1.9-8.6	1.6	2.3	1.0-4.5	1.0	3.4	1.0-8.0	0.6	1.6	0.3-4.7
	1996-2000	7.4	10.8	7.6-14.9	4.2	13.6	8.3-21.0	3.2	8.8	5.0-14.3	1.6	2.4	1.0-4.7	1.2	4.7	1.6-10.2	0.4	1.1	0.1-4.0
	2001-2005	6.8	9.2	6.3-12.8	5.0	15.0	9.6-22.4	1.8	4.2	1.9-8.1	1.0	1.4	0.4-3.2	0.6	1.8	0.4-5.4	0.4	0.9	0.1-3.5
	2006-2010	8.2	9.7	6.9-13.3	5.6	14.3	9.4-21.1	2.6	5.5	2.9-9.6	2.0	2.4	1.1-4.5	1.4	3.6	1.4-7.6	0.6	1.2	0.2-3.9
	2011-2015	12.0	12.9	9.8-16.9	9.6	21.8	15.9-29.4	2.4	5.4	2.7-9.9	0.4	0.7	0.1-2.4	0.0	0.0	0.0-2.3	0.4	1.3	0.1-4.7
	2016-2020	12.6	13.5	10.2-17.6	10.6	24.1	17.7-32.2	2.0	3.8	1.8-7.6	1.8	1.8	0.8-3.6	1.2	2.6	0.9-6.1	0.6	1.1	0.2-3.8
Esophagus	1976-1980	1.6	2.9	1.2-5.7	1.2	5.8	1.9-12.9	0.4	1.2	0.1-4.4	1.0	2.0	0.6-4.7	1.0	5.4	1.6-12.4	0.0	0.0	0.0-2.5
	1981-1985	2.2	4.1	2.0-7.4	1.4	6.3	2.4-13.2	0.8	2.4	0.7-6.3	2.8	4.8	2.6-8.1	1.8	7.3	3.1-14.2	1.0	3.0	1.0-7.1
	1986-1990	3.6	6.1	3.6-9.7	3.0	12.4	6.7-20.8	0.6	1.7	0.3-5.0	3.0	5.1	2.8-8.4	2.4	10.0	5.0-17.6	0.6	1.6	0.3-4.9
	1991-1995	3.0	4.7	2.6-7.7	2.2	7.6	3.8-13.7	0.8	2.1	0.6-5.4	2.6	4.1	2.2-7.0	2.2	8.2	3.9-14.8	0.4	1.0	0.1-3.9
	1996-2000	2.4	3.6	1.9-6.3	2.0	7.2	3.3-13.3	0.4	1.0	0.1-3.8	1.8	2.7	1.2-5.1	1.2	4.2	1.5-9.1	0.6	1.5	0.3-4.6
	2001-2005	4.4	6.0	3.7-9.1	3.8	11.2	6.7-17.6	0.6	1.5	0.3-4.5	4.0	5.5	3.4-8.5	3.6	10.8	6.4-17.2	0.4	0.9	0.1-3.6
	2006-2010	4.4	5.4	3.4-8.2	3.6	9.4	5.5-15.1	0.8	1.8	0.5-5.0	5.2	6.7	4.3-9.8	3.4	9.5	5.5-15.4	1.8	4.1	1.9-8.1
	2011-2015	5.8	6.3	4.1-9.2	4.8	11.1	7.0-16.9	1.0	2.5	0.7-6.3	5.6	6.1	4.0-8.9	5.0	11.8	7.5-17.8	0.6	1.2	0.2-4.0
	2016-2020	6.2	5.9	4.0-8.5	5.0	9.9	6.3-15.1	1.2	2.1	0.8-5.1	4.4	4.0	2.5-6.3	3.8	7.3	4.3-11.9	0.6	1.0	0.2-3.5
Stomach	1976-1980	4.2	7.7	4.7-11.8	2.6	10.4	5.3-18.2	1.6	5.5	2.3-10.8	2.8	5.2	2.8-8.7	1.2	5.1	1.8-11.3	1.6	5.3	2.3-10.6
	1981-1985	3.8	7.1	4.2-11.0	2.4	11.2	5.5-19.7	1.4	4.4	1.7-9.1	2.6	4.7	2.5-8.1	1.2	5.1	1.8-11.1	1.4	4.2	1.7-8.7
	1986-1990	3.4	5.7	3.3-9.2	2.6	10.6	5.6-18.2	0.8	2.2	0.6-5.7	2.6	4.4	2.3-7.5	2.0	7.7	3.6-14.1	0.6	1.6	0.3-4.9
	1991-1995	2.8	4.5	2.4-7.5	2.2	8.3	4.1-14.9	0.6	1.6	0.3-4.7	2.4	3.8	2.0-6.7	2.4	9.6	4.9-16.6	0.0	0.0	0.0-2.1
	1996-2000	2.6	3.9	2.1-6.6	1.8	6.6	2.9-12.6	0.8	2.1	0.6-5.5	2.2	3.3	1.6-5.9	1.4	5.4	2.1-11.1	0.8	2.0	0.5-5.3
	2001-2005	3.8	5.3	3.2-8.3	2.6	8.6	4.5-14.8	1.2	3.1	1.1-7.0	3.2	4.5	2.6-7.4	1.6	5.7	2.4-11.2	1.6	4.1	1.7-8.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Madison**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	4.8	6.4	4.1-9.6	3.0	8.9	4.9-14.7	1.8	3.8	1.7-7.5	1.0	1.3	0.4-3.1	0.6	1.5	0.3-4.7	0.4	1.0	0.1-3.7
	2011-2015	4.8	5.4	3.4-8.2	3.0	7.1	3.9-12.1	1.8	3.8	1.7-7.6	0.6	0.7	0.1-2.2	0.2	0.6	0.0-3.4	0.4	0.7	0.1-3.1
	2016-2020	5.8	5.6	3.7-8.3	4.2	8.2	5.0-13.0	1.6	3.1	1.3-6.6	1.4	1.6	0.6-3.6	0.6	1.2	0.2-4.1	0.8	2.1	0.5-5.7
Colorectal	1976-1980	34.0	67.9	57.9-79.1	15.6	74.5	58.1-93.7	18.4	62.7	50.4-76.9	16.4	33.8	26.8-42.1	7.2	36.7	25.1-51.3	9.2	32.3	23.6-43.2
	1981-1985	45.4	84.3	73.5-96.1	22.6	100.0	81.5-121.3	22.8	72.8	60.0-87.6	19.0	35.0	28.2-42.8	9.4	42.2	30.3-56.7	9.6	29.5	21.7-39.2
	1986-1990	41.6	71.7	62.2-82.2	21.0	87.7	70.9-107.1	20.6	60.0	48.9-72.9	20.6	36.3	29.6-44.1	8.4	33.9	23.9-46.3	12.2	35.9	27.4-46.2
	1991-1995	33.8	54.2	46.3-63.1	15.2	57.4	44.9-72.3	18.6	51.2	41.3-62.8	16.2	26.3	20.9-32.7	8.6	34.7	24.7-47.2	7.6	20.6	14.5-28.3
	1996-2000	44.4	65.8	57.4-75.1	20.4	70.0	56.8-85.3	24.0	62.4	51.7-74.7	15.8	23.6	18.7-29.4	7.4	26.2	18.3-36.4	8.4	21.0	15.1-28.5
	2001-2005	47.2	65.0	56.9-73.9	23.2	75.4	61.9-90.8	24.0	58.4	48.3-69.9	18.6	25.7	20.7-31.5	9.0	31.0	22.4-41.7	9.6	22.2	16.4-29.7
	2006-2010	40.0	49.0	42.4-56.4	20.6	56.7	46.0-69.1	19.4	43.5	35.2-53.3	13.6	16.9	13.1-21.5	6.8	20.5	14.0-28.8	6.8	14.8	10.2-20.9
	2011-2015	35.0	40.4	34.5-47.2	17.2	42.4	33.6-52.9	17.8	38.0	30.3-47.2	14.2	16.6	12.9-21.1	6.8	18.3	12.5-25.9	7.4	15.4	10.7-21.6
2016-2020	34.0	36.8	31.3-43.1	19.8	47.3	38.0-58.2	14.2	28.0	21.7-35.9	12.0	13.1	9.9-17.2	6.4	15.7	10.5-22.8	5.6	10.7	7.1-16.0	
Colon excluding rectum	1976-1980	26.6	53.1	44.3-63.1	11.2	54.5	40.5-71.4	15.4	51.9	40.8-64.9	13.6	27.6	21.3-35.1	6.0	30.8	20.3-44.3	7.6	26.3	18.5-36.2
	1981-1985	32.4	59.9	50.9-70.0	15.8	67.3	52.6-84.7	16.6	53.2	42.2-66.0	15.6	28.6	22.5-35.7	7.4	31.5	21.7-43.9	8.2	25.2	18.1-34.3
	1986-1990	29.2	51.0	43.0-60.0	14.6	62.8	48.5-79.7	14.6	42.8	33.5-53.9	17.6	31.1	24.9-38.4	6.4	26.0	17.4-37.1	11.2	33.1	24.9-43.0
	1991-1995	27.0	43.3	36.3-51.3	10.6	40.0	29.6-52.6	16.4	45.1	35.8-56.0	13.2	21.5	16.6-27.3	6.8	28.5	19.3-40.2	6.4	16.9	11.6-24.0
	1996-2000	32.0	47.5	40.4-55.4	14.0	48.0	37.2-60.8	18.0	46.7	37.5-57.5	13.8	20.5	16.0-26.0	6.6	23.2	15.8-32.7	7.2	18.0	12.6-25.1
	2001-2005	35.0	48.4	41.5-56.2	15.0	49.4	38.6-62.2	20.0	48.2	39.1-58.8	14.4	20.0	15.6-25.2	7.2	25.3	17.5-35.2	7.2	16.4	11.4-22.9
	2006-2010	28.0	34.7	29.2-41.0	13.4	38.2	29.4-48.7	14.6	32.6	25.5-41.2	11.0	13.7	10.3-17.9	5.2	16.0	10.4-23.6	5.8	12.5	8.4-18.2
	2011-2015	24.0	27.6	22.8-33.2	11.2	27.9	20.8-36.6	12.8	27.0	20.7-34.9	10.8	12.4	9.3-16.3	4.6	12.4	7.8-18.8	6.2	12.9	8.6-18.7
2016-2020	23.0	24.4	20.0-29.6	12.2	28.7	21.7-37.5	10.8	20.7	15.4-27.4	9.2	10.3	7.4-14.0	4.8	12.3	7.6-18.8	4.4	8.4	5.2-13.2	
Rectum & rectosigmoid	1976-1980	7.4	14.8	10.4-20.5	4.4	20.0	12.2-30.5	3.0	10.8	6.0-17.8	2.8	6.2	3.4-10.4	1.2	5.9	2.0-13.0	1.6	6.0	2.6-11.7
	1981-1985	13.0	24.3	18.7-31.1	6.8	32.7	22.1-46.2	6.2	19.7	13.3-28.0	3.4	6.4	3.7-10.2	2.0	10.7	4.9-19.7	1.4	4.3	1.7-8.9
	1986-1990	12.4	20.8	15.9-26.6	6.4	24.9	16.7-35.6	6.0	17.2	11.6-24.6	3.0	5.2	2.9-8.5	2.0	7.9	3.6-14.8	1.0	2.8	0.9-6.7
	1991-1995	6.8	10.9	7.5-15.2	4.6	17.5	10.9-26.4	2.2	6.1	3.0-11.0	3.0	4.8	2.7-7.9	1.8	6.2	2.8-11.8	1.2	3.6	1.3-7.9
	1996-2000	12.4	18.4	14.1-23.5	6.4	22.1	14.9-31.3	6.0	15.7	10.5-22.4	2.0	3.1	1.5-5.6	0.8	3.1	0.8-7.7	1.2	3.0	1.1-6.6
	2001-2005	12.2	16.6	12.7-21.3	8.2	26.0	18.5-35.5	4.0	10.2	6.2-15.8	4.2	5.7	3.5-8.8	1.8	5.7	2.5-11.0	2.4	5.9	3.0-10.4
	2006-2010	12.0	14.3	10.9-18.5	7.2	18.5	12.8-25.9	4.8	10.9	6.9-16.4	2.6	3.2	1.7-5.5	1.6	4.5	1.8-9.0	1.0	2.3	0.7-5.5
	2011-2015	11.0	12.8	9.5-16.9	6.0	14.5	9.6-21.3	5.0	11.0	7.0-16.6	3.4	4.2	2.4-6.8	2.2	5.9	2.8-10.9	1.2	2.5	0.9-5.8
2016-2020	11.0	12.4	9.2-16.5	7.6	18.5	12.8-26.0	3.4	7.3	4.2-12.2	2.8	2.8	1.5-5.0	1.6	3.5	1.5-7.2	1.2	2.3	0.8-5.5	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Madison**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	2.6	4.7	2.5-8.1	1.4	5.5	2.1-11.6	1.2	3.8	1.4-8.4	1.0	1.6	0.5-3.7	0.6	1.9	0.4-5.9	0.4	1.2	0.1-4.5
	1981-1985	1.8	3.1	1.4-5.9	1.0	3.9	1.2-9.1	0.8	2.3	0.6-6.1	2.2	3.8	1.9-6.9	1.2	5.3	1.8-11.5	1.0	2.9	0.9-6.9
	1986-1990	1.4	2.5	1.0-5.1	0.6	3.0	0.5-8.6	0.8	2.4	0.7-6.2	1.6	3.0	1.3-5.8	1.0	5.3	1.6-12.2	0.6	1.8	0.4-5.4
	1991-1995	2.2	3.3	1.6-5.9	1.4	4.8	1.9-9.9	0.8	2.3	0.6-5.9	2.2	3.3	1.6-5.9	1.4	4.5	1.8-9.4	0.8	2.3	0.6-6.0
	1996-2000	3.2	4.8	2.7-7.8	2.0	6.6	3.1-12.1	1.2	3.4	1.2-7.5	2.0	3.0	1.4-5.5	1.0	3.2	1.0-7.4	1.0	2.5	0.8-5.9
	2001-2005	2.4	3.3	1.7-5.9	1.2	3.6	1.3-8.0	1.2	3.0	1.1-6.6	2.0	2.8	1.4-5.2	0.6	1.8	0.4-5.4	1.4	3.2	1.3-6.9
	2006-2010	3.8	4.7	2.8-7.4	2.6	7.3	3.8-12.6	1.2	2.6	0.9-5.9	2.8	3.6	2.0-6.1	2.0	6.0	2.8-11.1	0.8	1.7	0.4-4.6
	2011-2015	3.4	3.9	2.2-6.4	2.4	5.9	3.0-10.6	1.0	2.0	0.6-5.1	4.2	4.6	2.8-7.2	2.6	6.1	3.2-10.7	1.6	3.1	1.3-6.5
	2016-2020	4.8	4.6	2.9-7.0	3.4	6.7	3.8-11.1	1.4	2.9	1.1-6.3	2.4	2.4	1.2-4.4	1.6	3.4	1.4-7.1	0.8	1.5	0.4-4.5
Pancreas	1976-1980	4.8	8.8	5.6-13.2	3.4	14.7	8.1-24.1	1.4	4.7	1.9-9.7	6.0	11.3	7.5-16.2	4.2	18.0	10.7-28.1	1.8	6.2	2.8-11.7
	1981-1985	4.8	8.8	5.6-13.1	0.8	3.3	0.8-8.7	4.0	13.1	7.9-20.3	5.6	10.4	6.8-15.0	1.6	7.2	3.0-14.2	4.0	12.8	7.7-19.8
	1986-1990	7.4	12.9	9.0-17.8	4.2	18.5	11.0-28.6	3.2	9.6	5.5-15.7	5.6	9.6	6.4-13.9	3.2	14.5	7.9-23.8	2.4	7.1	3.7-12.5
	1991-1995	6.4	10.1	6.9-14.3	3.4	11.8	6.8-19.0	3.0	8.3	4.6-13.7	6.6	10.5	7.2-14.7	3.4	12.8	7.3-20.7	3.2	8.7	4.9-14.2
	1996-2000	9.6	14.3	10.5-18.9	5.8	19.6	13.0-28.4	3.8	9.9	5.9-15.5	7.6	11.3	8.0-15.6	4.2	14.7	8.9-22.6	3.4	8.9	5.2-14.3
	2001-2005	7.4	10.2	7.2-14.2	4.6	14.8	9.3-22.3	2.8	6.7	3.6-11.4	5.6	7.7	5.1-11.2	2.8	9.0	4.9-15.2	2.8	6.5	3.6-11.2
	2006-2010	10.6	12.8	9.6-16.9	4.4	12.0	7.4-18.4	6.2	13.8	9.3-19.8	8.6	10.5	7.6-14.2	4.0	11.3	6.8-17.7	4.6	10.1	6.4-15.4
	2011-2015	9.2	9.9	7.2-13.4	6.0	14.1	9.4-20.5	3.2	6.6	3.7-11.0	8.6	9.2	6.7-12.6	5.4	12.7	8.2-18.8	3.2	6.6	3.7-11.1
	2016-2020	13.2	13.6	10.4-17.6	7.2	15.1	10.5-21.2	6.0	12.1	8.0-17.9	10.0	10.0	7.4-13.4	5.0	10.8	6.9-16.3	5.0	9.1	5.9-13.9
Larynx	1976-1980	4.4	7.3	4.6-11.1	3.4	12.0	6.9-19.6	1.0	3.2	1.0-7.6	1.0	1.8	0.6-4.2	0.8	3.3	0.8-8.7	0.2	0.8	0.0-4.0
	1981-1985	2.2	3.7	1.8-6.7	2.0	7.3	3.4-13.6	0.2	0.7	0.0-3.8	1.2	1.8	0.7-4.0	1.2	3.9	1.4-8.9	0.0	0.0	0.0-2.4
	1986-1990	3.6	6.0	3.5-9.5	2.4	9.5	4.7-16.8	1.2	3.4	1.3-7.6	1.8	3.1	1.4-5.8	1.0	4.3	1.3-10.3	0.8	2.3	0.6-6.0
	1991-1995	4.0	6.2	3.8-9.5	3.0	9.9	5.5-16.4	1.0	3.0	1.0-6.9	0.6	0.8	0.2-2.5	0.4	1.2	0.1-4.5	0.2	0.5	0.0-3.0
	1996-2000	3.4	5.0	2.9-7.9	2.8	9.1	4.9-15.3	0.6	1.7	0.4-5.0	0.6	0.9	0.2-2.6	0.6	2.3	0.4-6.7	0.0	0.0	0.0-2.1
	2001-2005	3.0	4.1	2.3-6.8	2.4	7.4	3.8-13.1	0.6	1.7	0.3-5.0	1.4	2.0	0.8-4.1	1.0	3.5	1.1-8.1	0.4	0.9	0.1-3.4
	2006-2010	3.4	4.0	2.3-6.5	3.0	7.4	4.1-12.5	0.4	0.8	0.1-3.4	1.0	1.3	0.4-3.1	0.6	1.6	0.3-4.9	0.4	1.0	0.1-3.7
	2011-2015	2.8	3.1	1.7-5.4	2.0	4.8	2.3-9.1	0.8	1.7	0.4-4.7	0.4	0.4	0.1-1.8	0.4	0.9	0.1-3.7	0.0	0.0	0.0-2.2
	2016-2020	2.2	2.6	1.2-4.8	1.6	3.9	1.5-8.2	0.6	1.2	0.3-4.1	0.8	0.8	0.2-2.4	0.6	1.0	0.2-3.7	0.2	0.7	0.0-3.6
Lung and bronchus	1976-1980	28.0	52.5	44.0-62.1	20.4	87.9	71.1-107.4	7.6	25.1	17.7-34.5	26.0	48.3	40.2-57.5	19.8	84.9	68.4-104.1	6.2	20.4	13.7-29.0
	1981-1985	36.2	62.8	53.8-72.8	25.2	103.5	85.4-124.2	11.0	34.5	25.9-45.1	29.6	51.8	43.6-61.0	20.8	85.4	69.0-104.4	8.8	27.9	20.2-37.6
	1986-1990	48.8	80.8	70.9-91.7	32.8	125.6	106.4-147.0	16.0	47.5	37.6-59.3	40.8	67.6	58.6-77.7	27.4	105.5	88.0-125.3	13.4	39.6	30.6-50.4
	1991-1995	48.8	76.5	67.1-86.7	29.6	106.3	89.5-125.3	19.2	54.1	43.8-66.2	34.4	53.6	45.8-62.3	21.8	78.0	63.7-94.5	12.6	34.9	26.8-44.7
	1996-2000	62.2	91.3	81.5-102.1	36.2	117.8	101.1-136.5	26.0	70.5	58.8-83.7	49.0	72.4	63.6-82.0	29.6	99.6	83.9-117.3	19.4	52.2	42.3-63.7
	2001-2005	57.8	77.9	69.1-87.4	31.8	97.2	82.3-114.0	26.0	64.5	53.9-76.7	41.2	55.8	48.4-64.0	23.0	69.6	57.2-83.9	18.2	44.8	36.0-55.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Madison**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	70.0	86.2	77.3-95.9	36.6	101.3	86.9-117.5	33.4	75.3	64.2-87.9	46.0	56.7	49.5-64.6	25.2	69.4	57.5-82.9	20.8	47.2	38.4-57.4
	2011-2015	70.0	76.0	68.1-84.6	34.4	80.2	68.4-93.6	35.6	72.6	62.1-84.4	46.4	50.8	44.4-58.0	21.2	51.7	42.1-62.9	25.2	50.6	42.0-60.6
	2016-2020	63.8	61.7	55.0-69.2	34.6	69.2	59.1-80.9	29.2	55.4	46.4-65.7	42.4	40.7	35.3-46.8	22.6	46.8	38.3-56.8	19.8	35.5	28.8-43.6
Melanoma of the skin	1976-1980	3.2	6.4	3.6-10.4	1.8	8.5	3.5-16.8	1.4	5.3	2.1-10.7	1.6	3.1	1.3-6.2	1.4	7.2	2.7-15.0	0.2	0.6	0.0-3.6
	1981-1985	5.8	10.3	6.8-14.8	3.6	14.1	8.0-22.8	2.2	7.9	3.9-14.0	2.4	4.3	2.2-7.6	2.0	8.3	3.8-15.5	0.4	1.3	0.2-4.6
	1986-1990	4.6	7.3	4.6-11.0	2.4	8.3	4.2-14.7	2.2	6.7	3.3-12.1	1.6	2.7	1.2-5.3	1.0	3.5	1.1-8.3	0.6	2.0	0.4-5.9
	1991-1995	4.0	6.1	3.7-9.5	1.6	5.0	2.1-9.9	2.4	7.2	3.7-12.6	1.2	1.9	0.7-4.0	0.8	2.6	0.7-6.7	0.4	1.2	0.1-4.4
	1996-2000	6.0	8.8	5.9-12.6	4.0	13.0	7.8-20.1	2.0	5.6	2.7-10.4	1.2	1.8	0.7-3.9	0.8	3.1	0.8-7.8	0.4	1.1	0.1-4.1
	2001-2005	13.6	18.5	14.3-23.5	8.6	25.0	18.0-33.8	5.0	13.6	8.7-20.1	2.0	2.7	1.3-5.0	1.2	4.5	1.6-9.7	0.8	2.0	0.5-5.3
	2006-2010	15.2	19.5	15.3-24.6	7.8	21.1	14.8-29.2	7.4	18.9	13.1-26.3	2.0	2.3	1.1-4.3	1.4	3.5	1.4-7.5	0.6	1.3	0.3-4.0
	2011-2015	21.2	24.7	20.1-30.1	11.0	26.0	19.4-34.3	10.2	24.2	17.8-32.2	1.6	1.8	0.8-3.7	1.6	3.8	1.6-7.8	0.0	0.0	0.0-2.2
2016-2020	20.2	22.7	18.3-28.0	9.8	22.4	16.3-30.1	10.4	23.8	17.4-31.9	2.0	2.1	1.0-4.0	1.4	3.3	1.3-7.1	0.6	0.9	0.2-3.5	
Female breast	1976-1980							30.2	106.0	89.5-124.5							11.6	41.1	31.0-53.2
	1981-1985							33.2	108.9	92.7-127.1							10.6	34.3	25.6-45.0
	1986-1990							36.8	112.5	96.6-130.2							11.0	33.3	25.0-43.5
	1991-1995							47.0	136.7	119.6-155.5							9.2	26.0	19.0-34.7
	1996-2000							48.8	133.7	117.4-151.6							12.2	32.3	24.6-41.5
	2001-2005							49.2	124.8	109.6-141.6							11.8	28.5	21.7-36.9
	2006-2010							53.0	123.6	108.9-139.8							6.6	14.1	9.7-20.0
	2011-2015							57.2	125.1	110.5-141.2							6.8	13.0	8.9-18.6
2016-2020							60.2	126.8	112.0-143.1							6.8	12.9	8.8-18.5	
Cervix uteri	1976-1980							4.4	14.0	8.7-21.3							2.2	7.8	3.8-14.0
	1981-1985							2.6	7.4	3.9-12.8							1.2	3.5	1.3-7.7
	1986-1990							3.2	9.9	5.6-16.1							1.2	3.5	1.3-7.7
	1991-1995							3.4	9.8	5.7-15.7							0.8	2.2	0.6-5.8
	1996-2000							2.6	7.4	3.9-12.6							0.8	2.2	0.6-5.8
	2001-2005							1.8	4.5	2.1-8.7							0.8	1.9	0.5-5.1
	2006-2010							1.8	4.8	2.1-9.3							0.6	1.4	0.3-4.3
	2011-2015							1.6	4.4	1.8-8.9							0.8	1.9	0.5-5.3
	2016-2020							1.6	5.4	2.2-10.8							1.0	2.8	0.8-6.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Madison**

		Incidence									Mortality									
		Males and Females			Males			Females			Males and Females			Males			Females			
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	
Corpus uteri and uterus, not otherwise specified	1976-1980							11.0	36.5	27.4-47.7							2.6	8.8	4.6-15.0	
	1981-1985							7.6	23.7	16.6-32.7							1.2	3.8	1.4-8.3	
	1986-1990							8.6	26.4	19.0-35.6							1.8	5.1	2.3-9.7	
	1991-1995							8.6	25.6	18.4-34.5							1.6	4.3	1.9-8.6	
	1996-2000							11.0	29.9	22.5-39.0							1.2	3.1	1.1-6.8	
	2001-2005							11.6	29.4	22.3-38.1							2.0	4.9	2.3-9.1	
	2006-2010							10.0	23.7	17.5-31.5							1.4	3.1	1.2-6.6	
	2011-2015								15.0	30.5	23.8-38.8							2.0	4.0	1.9-7.7
	2016-2020								15.6	31.0	24.1-39.5							1.2	2.1	0.7-5.1
Ovary	1976-1980							4.6	16.0	10.1-24.1							2.8	10.2	5.5-17.1	
	1981-1985							5.4	17.5	11.5-25.6							3.4	11.2	6.4-18.0	
	1986-1990							5.8	17.9	11.9-25.8							2.4	7.3	3.7-12.8	
	1991-1995							6.4	19.0	13.0-26.9							3.2	9.1	5.2-14.9	
	1996-2000							6.0	16.4	11.1-23.5							4.0	10.7	6.5-16.5	
	2001-2005							6.0	15.9	10.7-22.9							4.6	11.8	7.4-17.8	
	2006-2010							4.8	11.1	7.1-16.8							4.0	9.2	5.6-14.5	
	2011-2015							4.4	10.1	6.1-15.7							2.8	5.4	2.9-9.4	
	2016-2020							5.0	10.3	6.5-15.8							2.2	5.0	2.4-9.4	
Prostate	1976-1980				19.2	102.3	81.8-125.9							6.8	38.6	26.2-54.2				
	1981-1985				22.2	104.0	84.6-126.3							6.8	37.4	25.4-52.4				
	1986-1990				35.2	149.4	127.1-174.2							7.6	36.7	25.3-50.8				
	1991-1995				51.4	189.5	166.2-215.0							8.6	38.9	27.8-52.7				
	1996-2000				48.4	160.3	140.3-182.2							7.6	30.1	21.1-41.5				
	2001-2005				58.0	166.1	147.3-186.8							7.0	27.8	19.2-38.8				
	2006-2010				87.8	217.2	196.9-239.0							7.0	22.7	15.7-31.8				
	2011-2015				60.2	133.2	118.2-149.7							5.8	15.4	10.3-22.4				
	2016-2020				69.8	133.4	119.4-148.8							7.6	18.8	13.2-26.0				
Testis	1976-1980				1.4	4.7	1.7-10.3							0.2	0.3	0.0-3.5				
	1981-1985				1.0	2.6	0.8-6.7							0.0	0.0	0.0-3.0				
	1986-1990				3.0	8.5	4.7-14.3							0.2	0.8	0.0-4.2				
	1991-1995				1.6	4.6	2.0-9.3							0.0	0.0	0.0-2.5				
	1996-2000				2.8	8.5	4.6-14.4							0.6	1.8	0.4-5.3				
	2001-2005				2.0	6.3	3.0-11.7							0.0	0.0	0.0-2.3				

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Madison**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				3.2	8.7	4.9-14.4							0.0	0.0	0.0-2.3			
	2011-2015				2.6	7.9	4.1-13.7							0.0	0.0	0.0-2.3			
	2016-2020				2.0	5.3	2.5-10.0							0.2	0.5	0.0-3.1			
Urinary bladder (incl. in situ)	1976-1980	13.8	28.0	21.7-35.5	10.6	51.1	37.6-67.4	3.2	11.6	6.6-18.7	4.4	8.9	5.5-13.5	2.8	13.1	6.9-22.2	1.6	5.7	2.4-11.1
	1981-1985	14.4	27.3	21.3-34.4	10.4	47.5	34.9-62.8	4.0	13.0	7.9-20.1	2.6	5.5	2.9-9.3	1.0	6.6	2.1-14.6	1.6	5.5	2.3-10.7
	1986-1990	15.0	25.6	20.1-32.1	10.6	43.9	32.5-57.8	4.4	12.6	7.9-19.1	4.2	7.8	4.8-11.8	2.8	14.3	7.6-23.9	1.4	4.0	1.6-8.3
	1991-1995	18.8	29.9	24.1-36.6	12.4	46.7	35.5-60.1	6.4	17.3	11.8-24.6	5.4	8.8	5.8-12.8	3.4	14.3	8.1-23.0	2.0	5.3	2.5-9.9
	1996-2000	16.4	24.0	19.1-29.8	12.4	42.2	32.1-54.3	4.0	10.9	6.6-16.8	3.2	4.6	2.7-7.6	1.8	6.2	2.8-11.9	1.4	3.7	1.5-7.7
	2001-2005	21.4	29.5	24.2-35.7	17.2	54.4	43.3-67.5	4.2	10.5	6.5-16.1	4.8	6.7	4.3-10.0	4.0	14.2	8.5-21.9	0.8	1.7	0.5-4.7
	2006-2010	25.2	32.1	26.7-38.3	19.8	57.0	46.1-69.7	5.4	12.4	8.1-18.2	4.2	5.4	3.3-8.4	3.4	10.6	6.1-17.0	0.8	1.6	0.4-4.6
	2011-2015	27.2	30.0	25.1-35.7	18.4	45.2	36.2-55.8	8.8	17.6	12.7-24.0	5.4	6.1	4.0-9.0	3.2	8.7	4.9-14.3	2.2	4.4	2.2-8.2
	2016-2020	26.0	27.0	22.4-32.3	20.4	46.8	37.9-57.4	5.6	10.6	6.9-15.8	6.4	6.7	4.5-9.6	4.6	11.4	7.1-17.3	1.8	3.2	1.5-6.6
Kidney and renal pelvis	1976-1980	6.0	11.5	7.6-16.4	4.0	18.0	10.6-28.3	2.0	6.4	3.1-11.9	2.6	4.9	2.6-8.4	1.8	7.2	3.2-13.9	0.8	2.7	0.7-7.0
	1981-1985	4.2	7.4	4.6-11.4	3.0	14.0	7.3-23.6	1.2	3.6	1.3-8.0	1.6	2.8	1.2-5.6	1.2	5.7	1.9-12.6	0.4	1.1	0.1-4.2
	1986-1990	7.8	13.3	9.4-18.2	3.6	13.0	7.6-20.8	4.2	13.4	8.3-20.6	2.8	4.6	2.5-7.8	1.8	6.0	2.7-11.6	1.0	3.4	1.1-7.9
	1991-1995	7.2	11.0	7.7-15.2	5.0	16.7	10.7-24.9	2.2	6.4	3.2-11.4	2.4	3.7	1.9-6.5	1.8	6.9	3.0-13.2	0.6	1.6	0.3-4.6
	1996-2000	10.0	14.6	10.8-19.2	7.0	21.6	15.0-30.1	3.0	8.1	4.5-13.4	3.0	4.4	2.5-7.3	2.0	6.9	3.3-12.8	1.0	2.5	0.8-6.0
	2001-2005	10.2	13.4	10.0-17.7	7.0	20.1	13.9-28.2	3.2	8.1	4.6-13.3	3.4	4.5	2.6-7.3	2.6	8.2	4.3-14.3	0.8	1.9	0.5-5.1
	2006-2010	11.6	14.5	11.0-18.9	8.2	22.7	16.2-31.0	3.4	8.0	4.6-13.1	2.8	3.4	1.9-5.8	2.6	7.3	3.8-12.6	0.2	0.4	0.0-2.8
	2011-2015	16.0	17.2	13.5-21.6	12.2	28.2	21.4-36.6	3.8	7.5	4.5-12.0	4.2	4.3	2.7-6.8	4.0	9.3	5.6-14.6	0.2	0.3	0.0-2.6
	2016-2020	18.0	18.4	14.7-23.0	11.6	24.4	18.3-32.1	6.4	13.4	8.9-19.5	1.8	1.8	0.8-3.6	1.2	2.7	1.0-6.3	0.6	1.0	0.2-3.7
Brain and other central nervous system	1976-1980	1.0	1.7	0.5-4.0	0.6	2.3	0.5-6.9	0.4	1.2	0.1-4.5	1.2	2.2	0.8-4.7	0.6	2.6	0.5-7.6	0.6	1.7	0.4-5.2
	1981-1985	3.8	6.6	3.9-10.3	1.6	5.6	2.4-11.3	2.2	7.0	3.4-12.7	1.6	3.1	1.3-6.1	0.6	2.3	0.5-6.9	1.0	3.4	1.1-7.9
	1986-1990	4.4	7.0	4.4-10.7	2.0	6.6	3.1-12.4	2.4	7.2	3.7-12.6	2.0	3.3	1.6-6.0	1.0	3.7	1.2-8.6	1.0	2.9	0.9-6.8
	1991-1995	3.2	5.1	2.9-8.2	1.4	5.3	2.0-11.0	1.8	5.1	2.3-9.7	3.0	4.6	2.6-7.6	1.6	5.5	2.2-11.1	1.4	3.8	1.5-7.8
	1996-2000	4.4	6.4	4.0-9.6	2.8	8.4	4.6-14.1	1.6	4.5	2.0-8.9	2.4	3.5	1.8-6.1	1.4	4.2	1.7-8.8	1.0	2.7	0.9-6.3
	2001-2005	5.8	8.0	5.3-11.5	3.4	10.1	5.9-16.3	2.4	6.4	3.2-11.2	3.0	4.2	2.3-6.9	1.6	4.6	2.0-9.2	1.4	3.5	1.4-7.3
	2006-2010	4.2	5.9	3.6-9.0	1.8	5.3	2.3-10.2	2.4	6.3	3.2-11.1	2.0	2.7	1.3-5.1	1.0	2.7	0.8-6.4	1.0	2.9	0.9-6.9
	2011-2015	4.6	5.7	3.6-8.8	3.2	8.4	4.7-14.0	1.4	3.2	1.2-7.0	2.8	3.4	1.8-5.8	2.0	5.1	2.4-9.6	0.8	1.8	0.5-4.9
	2016-2020	6.6	8.3	5.6-12.0	3.4	9.3	5.3-15.3	3.2	7.4	4.0-12.6	4.0	4.2	2.5-6.7	2.6	5.8	3.0-10.3	1.4	2.8	1.1-6.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Madison**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	1.4	2.8	1.1-5.7	0.8	3.1	0.8-8.1	0.6	2.3	0.5-6.6	0.4	0.7	0.1-2.5	0.2	0.9	0.0-4.7	0.2	0.6	0.0-3.6
	1981-1985	2.4	4.2	2.1-7.4	0.6	2.2	0.4-6.7	1.8	6.1	2.7-11.6	0.2	0.5	0.0-2.3	0.2	1.0	0.0-5.0	0.0	0.0	0.0-2.4
	1986-1990	2.2	3.8	1.9-6.8	0.4	1.3	0.1-5.0	1.8	6.0	2.6-11.3	0.2	0.4	0.0-1.9	0.0	0.0	0.0-2.8	0.2	0.6	0.0-3.5
	1991-1995	3.4	5.0	2.9-8.1	0.6	1.8	0.4-5.4	2.8	8.2	4.5-13.8	0.8	1.3	0.3-3.2	0.4	1.3	0.2-4.8	0.4	1.1	0.1-4.0
	1996-2000	5.6	8.2	5.4-11.8	0.4	1.2	0.1-4.4	5.2	14.6	9.5-21.5	0.2	0.3	0.0-1.6	0.0	0.0	0.0-2.4	0.2	0.5	0.0-2.9
	2001-2005	9.0	12.7	9.2-17.1	3.0	8.6	4.7-14.4	6.0	17.2	11.5-24.7	0.6	0.8	0.2-2.5	0.2	1.0	0.0-4.6	0.4	1.0	0.1-3.8
	2006-2010	9.6	13.4	9.8-18.0	2.4	6.5	3.3-11.6	7.2	20.5	14.2-28.7	0.2	0.2	0.0-1.4	0.0	0.0	0.0-2.3	0.2	0.4	0.0-2.7
	2011-2015	13.4	17.5	13.4-22.5	4.6	10.2	6.3-15.7	8.8	24.7	17.6-33.6	0.2	0.3	0.0-1.5	0.2	0.6	0.0-3.3	0.0	0.0	0.0-2.2
2016-2020	10.8	13.1	9.7-17.5	2.6	6.3	3.2-11.1	8.2	19.9	13.9-27.6	0.0	0.0	0.0-1.1	0.0	0.0	0.0-2.3	0.0	0.0	0.0-2.3	
Hodgkin lymphoma	1976-1980	1.0	1.6	0.5-3.8	0.2	0.4	0.0-3.6	0.8	2.7	0.7-7.0	0.2	0.3	0.0-1.9	0.2	0.7	0.0-4.2	0.0	0.0	0.0-2.5
	1981-1985	1.8	3.3	1.5-6.3	1.2	4.8	1.7-10.4	0.6	2.0	0.4-5.9	1.2	2.0	0.7-4.4	0.8	2.8	0.7-7.5	0.4	1.4	0.2-4.9
	1986-1990	1.0	1.3	0.4-3.2	0.6	1.4	0.3-4.7	0.4	1.2	0.1-4.3	0.4	0.8	0.1-2.7	0.2	0.8	0.0-4.2	0.2	0.8	0.0-4.0
	1991-1995	2.4	3.3	1.7-5.9	1.6	4.5	1.9-9.1	0.8	2.2	0.6-5.7	0.4	0.6	0.1-2.2	0.2	0.6	0.0-3.5	0.2	0.5	0.0-3.0
	1996-2000	3.2	4.4	2.5-7.2	1.4	3.9	1.5-8.3	1.8	5.0	2.3-9.5	1.0	1.5	0.5-3.5	0.8	2.4	0.7-6.3	0.2	0.6	0.0-3.2
	2001-2005	2.2	3.2	1.6-5.7	1.0	3.0	0.9-7.2	1.2	3.3	1.2-7.3	0.6	1.0	0.2-2.8	0.2	0.7	0.0-3.7	0.4	1.4	0.2-4.9
	2006-2010	1.8	2.2	1.0-4.3	0.8	1.8	0.5-5.1	1.0	2.4	0.7-5.8	0.2	0.3	0.0-1.5	0.0	0.0	0.0-2.3	0.2	0.4	0.0-2.8
	2011-2015	1.6	2.6	1.1-5.2	1.2	4.1	1.4-8.9	0.4	1.2	0.1-4.3	0.0	0.0	0.0-1.1	0.0	0.0	0.0-2.3	0.0	0.0	0.0-2.2
2016-2020	1.8	2.0	0.9-4.1	1.0	2.7	0.8-6.6	0.8	1.4	0.4-4.2	0.4	0.6	0.1-2.2	0.4	1.2	0.1-4.5	0.0	0.0	0.0-2.3	
Non-Hodgkin lymphomas	1976-1980	6.8	13.1	9.0-18.4	4.2	19.0	11.4-29.5	2.6	8.9	4.7-15.3	5.2	10.4	6.7-15.3	2.6	13.4	6.9-23.0	2.6	8.9	4.7-15.2
	1981-1985	6.8	11.3	7.8-15.9	5.0	18.6	11.8-27.8	1.8	5.9	2.6-11.2	3.2	5.9	3.4-9.7	2.6	11.5	5.8-20.0	0.6	1.9	0.4-5.5
	1986-1990	9.0	15.0	10.9-20.1	4.4	16.7	10.2-25.6	4.6	13.7	8.6-20.7	3.8	6.0	3.6-9.4	2.0	8.5	3.8-16.0	1.8	4.9	2.2-9.5
	1991-1995	9.0	14.1	10.3-18.9	4.6	16.8	10.4-25.5	4.4	12.2	7.6-18.5	5.6	8.8	5.8-12.7	2.4	8.6	4.3-15.3	3.2	9.0	5.1-14.6
	1996-2000	12.6	18.6	14.3-23.7	6.2	19.9	13.4-28.4	6.4	17.3	11.8-24.5	4.2	6.2	3.8-9.5	2.4	8.8	4.4-15.4	1.8	4.4	2.0-8.6
	2001-2005	13.4	18.2	14.1-23.2	5.2	15.2	9.9-22.5	8.2	21.4	15.3-29.1	5.0	7.0	4.5-10.3	3.4	10.2	5.9-16.6	1.6	4.0	1.7-8.1
	2006-2010	17.6	21.9	17.5-27.1	9.0	24.6	17.8-33.2	8.6	19.6	14.1-26.6	6.0	7.7	5.2-11.0	2.8	7.8	4.2-13.2	3.2	7.8	4.4-12.8
	2011-2015	18.6	21.7	17.4-26.9	9.6	24.2	17.6-32.7	9.0	20.2	14.5-27.4	5.4	6.8	4.4-10.1	3.2	9.0	5.0-14.9	2.2	4.9	2.4-9.0
2016-2020	20.0	21.2	17.1-26.1	10.6	24.0	17.7-31.9	9.4	18.5	13.5-25.2	5.8	6.4	4.3-9.4	3.2	8.1	4.6-13.4	2.6	4.8	2.6-8.7	
Myeloma	1976-1980	1.4	2.7	1.1-5.6	0.8	3.7	0.9-9.3	0.6	2.2	0.4-6.2	1.6	3.6	1.5-6.9	0.8	4.9	1.2-12.0	0.8	3.1	0.8-7.7
	1981-1985	2.4	4.3	2.2-7.5	1.8	7.7	3.4-14.9	0.6	1.7	0.4-5.2	1.6	3.0	1.3-6.0	1.2	5.6	1.9-12.2	0.4	1.2	0.1-4.4
	1986-1990	2.4	4.2	2.2-7.3	1.6	6.7	2.8-13.1	0.8	2.2	0.6-5.8	1.6	2.8	1.2-5.4	1.2	4.5	1.6-9.7	0.4	1.1	0.1-4.2
	1991-1995	3.6	5.9	3.5-9.2	2.0	9.4	4.3-17.2	1.6	4.4	1.9-8.8	3.0	4.8	2.7-7.9	1.6	7.0	2.8-13.8	1.4	3.8	1.5-8.0
	1996-2000	3.8	5.6	3.4-8.7	2.6	8.8	4.6-15.2	1.2	3.0	1.1-6.8	2.4	3.6	1.8-6.2	1.4	5.4	2.1-11.1	1.0	2.6	0.8-6.1
	2001-2005	5.4	7.3	4.8-10.6	2.4	7.0	3.6-12.3	3.0	7.3	4.1-12.2	3.2	4.4	2.5-7.1	2.0	6.3	2.9-11.6	1.2	2.8	1.0-6.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Madison**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	4.8	6.1	3.9-9.2	3.2	9.4	5.2-15.4	1.6	3.7	1.6-7.6	2.4	2.9	1.5-5.2	1.4	4.1	1.6-8.6	1.0	2.1	0.7-5.3
	2011-2015	5.8	6.6	4.4-9.7	3.0	7.8	4.3-13.2	2.8	5.8	3.1-10.1	2.4	2.8	1.4-5.0	1.6	4.5	1.9-9.0	0.8	1.5	0.4-4.4
	2016-2020	4.4	4.5	2.8-7.0	3.0	6.5	3.5-11.2	1.4	2.9	1.1-6.4	1.8	1.8	0.8-3.7	1.0	2.0	0.6-5.3	0.8	1.7	0.4-4.9
Leukemias	1976-1980	7.8	14.0	9.8-19.3	4.2	18.0	10.6-28.2	3.6	11.9	6.9-19.1	4.2	7.0	4.2-10.9	2.2	7.9	3.6-14.9	2.0	6.7	3.1-12.6
	1981-1985	4.8	8.4	5.3-12.5	2.4	10.2	5.0-18.1	2.4	7.3	3.8-12.9	4.8	8.5	5.4-12.8	2.8	12.3	6.4-21.0	2.0	6.0	2.9-11.1
	1986-1990	7.6	12.0	8.4-16.6	4.2	15.0	9.1-23.2	3.4	9.5	5.5-15.4	5.2	8.6	5.6-12.7	3.2	12.1	6.8-19.9	2.0	5.7	2.7-10.7
	1991-1995	7.2	11.4	8.0-15.8	4.8	18.8	11.8-28.2	2.4	6.8	3.5-12.0	4.6	7.4	4.7-11.0	2.4	10.1	5.1-17.7	2.2	6.0	3.0-10.8
	1996-2000	11.4	17.0	12.8-22.0	6.0	20.0	13.3-28.6	5.4	14.7	9.6-21.4	6.2	9.3	6.3-13.2	3.6	12.7	7.4-20.1	2.6	7.0	3.7-12.0
	2001-2005	8.6	11.9	8.6-16.1	4.8	15.2	9.6-22.8	3.8	10.0	6.0-15.7	6.4	9.0	6.1-12.7	3.4	11.6	6.7-18.6	3.0	7.3	4.1-12.2
	2006-2010	14.2	17.8	13.9-22.5	8.0	21.9	15.5-30.1	6.2	14.0	9.4-20.1	6.6	8.7	6.0-12.3	4.0	12.2	7.4-18.9	2.6	5.9	3.1-10.4
	2011-2015	13.8	16.9	13.1-21.6	7.4	20.0	13.8-27.9	6.4	13.7	9.3-19.7	5.4	5.8	3.8-8.6	3.0	7.6	4.2-12.8	2.4	4.4	2.3-8.1
	2016-2020	13.6	14.5	11.1-18.6	7.8	17.6	12.3-24.6	5.8	11.6	7.6-17.2	5.6	6.1	4.0-9.0	4.0	9.4	5.6-14.9	1.6	3.3	1.3-7.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Monroe**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	2,648.0	429.2	421.8-436.7	1,271.4	512.1	499.0-525.5	1,376.6	388.3	379.0-397.7	1,332.2	218.6	213.3-224.0	675.6	281.3	271.4-291.3	656.6	182.7	176.4-189.1
	1981-1985	2,766.6	430.1	422.9-437.5	1,350.2	525.8	512.8-539.0	1,416.4	383.6	374.5-392.8	1,360.8	212.4	207.4-217.6	706.2	283.6	273.9-293.5	654.6	172.5	166.6-178.7
	1986-1990	3,102.4	467.8	460.3-475.2	1,530.4	574.6	561.3-588.1	1,572.0	414.2	405.0-423.6	1,387.2	210.6	205.6-215.6	685.4	267.7	258.4-277.1	701.8	179.4	173.4-185.5
	1991-1995	3,595.2	510.2	502.7-517.7	1,887.6	650.9	637.6-664.5	1,707.6	426.2	417.1-435.5	1,484.2	210.3	205.5-215.1	744.8	270.2	261.4-279.3	739.4	176.4	170.7-182.3
	1996-2000	3,873.4	524.7	517.3-532.1	1,920.4	618.9	606.5-631.5	1,953.0	467.8	458.4-477.2	1,524.0	203.2	198.6-207.8	756.6	253.2	245.1-261.5	767.4	173.7	168.1-179.3
	2001-2005	3,872.4	495.9	488.9-503.0	1,955.0	585.3	573.7-597.1	1,917.4	435.3	426.5-444.3	1,512.8	188.9	184.6-193.2	743.0	229.5	222.1-237.1	769.8	163.2	157.9-168.5
	2006-2010	4,226.6	510.3	503.4-517.4	2,140.2	590.3	579.0-601.7	2,086.4	455.9	447.0-465.0	1,503.0	176.9	172.9-181.0	749.0	212.0	205.2-219.0	754.0	154.5	149.5-159.7
	2011-2015	4,344.8	490.2	483.5-496.9	2,125.0	532.9	522.6-543.4	2,219.8	463.5	454.6-472.5	1,508.4	165.1	161.3-168.9	757.2	196.5	190.2-203.0	751.2	143.9	139.2-148.7
	2016-2020	4,727.8	496.2	489.7-502.8	2,403.6	549.8	539.7-560.0	2,324.2	459.6	450.9-468.4	1,483.6	150.6	147.2-154.2	746.0	176.3	170.6-182.2	737.6	133.7	129.3-138.3
Oral cavity and pharynx	1976-1980	63.0	10.0	8.9-11.2	38.8	14.1	12.1-16.3	24.2	6.9	5.7-8.3	26.6	4.3	3.6-5.1	18.2	7.0	5.6-8.6	8.4	2.3	1.7-3.2
	1981-1985	59.8	9.2	8.2-10.3	37.8	13.5	11.6-15.6	22.0	6.2	5.0-7.4	22.0	3.5	2.8-4.2	13.4	5.2	4.0-6.6	8.6	2.4	1.7-3.3
	1986-1990	67.4	10.2	9.1-11.4	38.6	13.6	11.7-15.7	28.8	7.4	6.3-8.8	25.0	3.9	3.2-4.6	14.6	5.6	4.3-7.0	10.4	2.7	2.0-3.6
	1991-1995	70.2	10.2	9.1-11.3	44.4	15.1	13.1-17.2	25.8	6.3	5.3-7.5	21.8	3.1	2.6-3.8	13.0	4.6	3.5-5.8	8.8	2.0	1.5-2.7
	1996-2000	78.6	10.7	9.7-11.8	51.2	16.1	14.2-18.2	27.4	6.6	5.6-7.8	18.8	2.5	2.1-3.1	14.0	4.6	3.6-5.8	4.8	1.2	0.7-1.7
	2001-2005	76.4	9.8	8.8-10.9	46.8	13.4	11.7-15.2	29.6	6.8	5.7-8.0	22.8	2.9	2.4-3.5	13.4	3.9	3.0-4.9	9.4	2.0	1.4-2.7
	2006-2010	89.2	10.5	9.6-11.6	54.6	14.4	12.7-16.3	34.6	7.5	6.4-8.7	16.2	1.9	1.5-2.4	11.4	3.1	2.3-4.1	4.8	1.0	0.7-1.6
	2011-2015	105.0	11.7	10.7-12.8	70.0	17.1	15.3-19.0	35.0	7.1	6.0-8.3	23.2	2.5	2.1-3.0	12.0	3.0	2.3-3.9	11.2	2.1	1.5-2.7
	2016-2020	118.2	12.2	11.2-13.3	78.8	17.9	16.1-19.8	39.4	7.3	6.3-8.5	23.6	2.3	1.9-2.8	16.6	3.7	2.9-4.6	7.0	1.2	0.8-1.7
Esophagus	1976-1980	25.8	4.3	3.6-5.1	17.6	7.1	5.6-8.8	8.2	2.4	1.7-3.2	25.4	4.3	3.5-5.1	18.8	7.7	6.2-9.5	6.6	1.9	1.3-2.7
	1981-1985	25.6	3.8	3.2-4.6	20.4	7.3	5.9-8.9	5.2	1.4	0.9-2.0	28.4	4.4	3.7-5.2	22.2	8.5	6.9-10.2	6.2	1.6	1.1-2.3
	1986-1990	30.2	4.5	3.8-5.3	20.0	7.1	5.7-8.7	10.2	2.5	1.9-3.4	31.2	4.8	4.1-5.6	23.8	8.5	7.0-10.2	7.4	1.9	1.3-2.6
	1991-1995	38.0	5.4	4.7-6.3	25.6	8.8	7.3-10.5	12.4	3.0	2.3-3.9	39.2	5.5	4.8-6.4	28.8	9.8	8.2-11.6	10.4	2.5	1.9-3.3
	1996-2000	37.0	5.0	4.3-5.8	26.0	8.4	7.0-9.9	11.0	2.6	1.9-3.4	41.2	5.6	4.9-6.4	31.0	10.0	8.5-11.8	10.2	2.4	1.8-3.1
	2001-2005	34.2	4.4	3.7-5.1	24.8	7.4	6.1-8.8	9.4	2.0	1.5-2.7	37.4	4.7	4.1-5.5	26.0	7.8	6.5-9.3	11.4	2.4	1.8-3.2
	2006-2010	46.0	5.5	4.8-6.3	35.6	9.7	8.3-11.3	10.4	2.2	1.6-2.9	40.6	4.9	4.3-5.7	31.6	8.8	7.5-10.3	9.0	2.0	1.4-2.7
	2011-2015	42.4	4.6	4.0-5.3	30.8	7.7	6.5-9.1	11.6	2.1	1.6-2.8	41.8	4.6	4.0-5.3	32.2	8.1	6.9-9.5	9.6	1.8	1.3-2.4
	2016-2020	44.8	4.5	3.9-5.2	33.4	7.6	6.5-8.9	11.4	2.0	1.5-2.7	45.2	4.5	3.9-5.2	33.4	7.7	6.5-9.0	11.8	2.1	1.6-2.7
Stomach	1976-1980	72.6	12.0	10.8-13.3	43.0	17.4	15.1-20.0	29.6	8.1	6.9-9.5	46.6	7.8	6.8-8.9	24.2	10.2	8.4-12.3	22.4	6.1	5.0-7.4
	1981-1985	73.8	11.6	10.5-12.9	46.4	18.8	16.3-21.4	27.4	7.1	6.0-8.4	45.2	7.1	6.2-8.1	26.8	11.0	9.1-13.1	18.4	4.8	3.8-5.8
	1986-1990	59.8	9.0	8.0-10.1	39.2	14.4	12.4-16.7	20.6	5.2	4.2-6.3	38.6	5.9	5.1-6.8	22.4	8.6	7.0-10.4	16.2	4.0	3.1-4.9
	1991-1995	68.8	9.7	8.7-10.8	45.0	15.9	13.8-18.1	23.8	5.5	4.6-6.6	37.2	5.2	4.5-6.0	25.2	9.3	7.7-11.1	12.0	2.7	2.0-3.4
	1996-2000	65.4	8.8	7.8-9.8	42.4	13.9	12.1-15.9	23.0	5.1	4.2-6.2	35.6	4.7	4.0-5.5	21.6	7.2	5.9-8.8	14.0	3.0	2.4-3.9
	2001-2005	61.2	7.7	6.8-8.6	39.2	11.8	10.2-13.6	22.0	4.6	3.7-5.5	32.4	3.9	3.3-4.6	19.0	5.8	4.7-7.1	13.4	2.5	2.0-3.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Monroe**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	62.4	7.5	6.6-8.4	40.0	11.2	9.7-12.9	22.4	4.7	3.8-5.6	33.0	3.9	3.3-4.6	21.4	6.1	5.0-7.4	11.6	2.4	1.8-3.1
	2011-2015	63.0	7.0	6.2-7.8	41.2	10.5	9.1-12.1	21.8	4.4	3.6-5.4	26.6	3.0	2.5-3.6	15.8	4.1	3.2-5.1	10.8	2.2	1.6-2.9
	2016-2020	61.2	6.5	5.7-7.3	38.2	9.0	7.7-10.4	23.0	4.5	3.7-5.5	25.0	2.5	2.1-3.0	14.8	3.4	2.7-4.3	10.2	1.8	1.4-2.5
Colorectal	1976-1980	441.4	72.6	69.5-75.7	216.2	89.4	83.9-95.1	225.2	61.7	58.2-65.5	197.4	33.2	31.1-35.3	91.8	40.5	36.7-44.6	105.6	29.0	26.5-31.6
	1981-1985	436.6	68.5	65.6-71.4	216.4	86.9	81.6-92.5	220.2	57.0	53.6-60.5	196.8	31.0	29.1-33.0	98.0	40.3	36.7-44.2	98.8	25.3	23.1-27.7
	1986-1990	429.2	64.9	62.2-67.8	209.6	81.7	76.7-87.0	219.6	54.5	51.3-57.9	174.6	26.5	24.8-28.3	85.0	35.0	31.6-38.6	89.6	21.8	19.8-23.9
	1991-1995	427.8	60.8	58.3-63.5	205.4	73.5	69.0-78.3	222.4	52.2	49.1-55.4	174.2	24.7	23.1-26.4	79.4	30.6	27.6-33.8	94.8	21.5	19.6-23.6
	1996-2000	461.2	61.5	59.0-64.1	228.4	76.0	71.6-80.6	232.8	52.4	49.4-55.5	158.6	20.8	19.4-22.4	77.4	26.4	23.8-29.2	81.2	17.2	15.6-19.0
	2001-2005	407.4	51.3	49.1-53.6	199.2	60.1	56.4-63.9	208.2	44.8	42.0-47.7	154.0	18.8	17.5-20.2	74.4	23.3	21.0-25.8	79.6	16.0	14.4-17.7
	2006-2010	370.6	43.7	41.7-45.8	175.6	48.3	45.2-51.7	195.0	39.8	37.3-42.5	130.2	15.0	13.8-16.2	62.6	17.5	15.6-19.6	67.6	13.1	11.7-14.6
	2011-2015	339.0	37.9	36.1-39.8	167.8	42.1	39.2-45.1	171.2	34.6	32.2-37.1	120.0	12.9	11.9-14.1	58.6	15.0	13.3-16.9	61.4	11.2	9.9-12.6
2016-2020	338.2	36.1	34.3-37.9	173.8	41.1	38.3-44.0	164.4	32.1	29.8-34.5	122.6	12.4	11.4-13.5	63.0	14.7	13.1-16.5	59.6	10.6	9.4-12.0	
Colon excluding rectum	1976-1980	306.2	50.5	48.0-53.2	144.8	60.3	55.8-65.0	161.4	44.3	41.3-47.5	155.2	26.1	24.2-28.0	67.2	29.7	26.4-33.2	88.0	24.1	21.9-26.5
	1981-1985	306.0	48.2	45.8-50.7	143.0	58.1	53.7-62.6	163.0	42.3	39.4-45.4	165.0	26.0	24.2-27.8	78.8	32.5	29.2-36.0	86.2	22.1	20.1-24.3
	1986-1990	296.8	44.8	42.6-47.2	138.6	55.0	50.8-59.4	158.2	39.1	36.4-41.9	150.2	22.9	21.3-24.6	73.2	30.0	26.9-33.3	77.0	18.7	16.9-20.7
	1991-1995	295.8	42.0	39.9-44.2	137.0	49.8	46.0-53.7	158.8	36.8	34.3-39.5	146.4	20.7	19.2-22.3	65.8	25.4	22.6-28.4	80.6	18.1	16.4-20.0
	1996-2000	319.8	42.4	40.4-44.6	151.0	50.9	47.3-54.7	168.8	37.4	34.9-40.1	128.6	16.8	15.6-18.2	61.6	21.2	18.9-23.7	67.0	14.2	12.7-15.8
	2001-2005	294.8	37.0	35.1-39.0	139.0	42.3	39.2-45.6	155.8	33.1	30.8-35.6	127.2	15.5	14.3-16.8	62.2	19.6	17.4-21.9	65.0	13.0	11.6-14.5
	2006-2010	263.4	30.9	29.3-32.7	120.0	33.3	30.7-36.2	143.4	28.8	26.7-31.1	108.4	12.4	11.4-13.5	51.6	14.6	12.8-16.5	56.8	10.9	9.6-12.3
	2011-2015	234.6	26.0	24.5-27.6	105.6	26.8	24.5-29.3	129.0	25.5	23.5-27.6	96.6	10.3	9.4-11.3	44.2	11.4	9.9-13.0	52.4	9.4	8.2-10.6
2016-2020	241.6	25.7	24.2-27.3	118.8	28.3	26.0-30.8	122.8	23.7	21.8-25.8	94.8	9.6	8.8-10.6	46.6	11.0	9.6-12.5	48.2	8.6	7.5-9.8	
Rectum & rectosigmoid	1976-1980	135.2	22.0	20.4-23.8	71.4	29.1	26.1-32.4	63.8	17.5	15.6-19.5	42.2	7.1	6.2-8.1	24.6	10.8	8.9-13.0	17.6	4.8	3.9-6.0
	1981-1985	130.6	20.3	18.7-21.9	73.4	28.9	25.9-32.1	57.2	14.7	13.0-16.5	31.8	5.0	4.3-5.9	19.2	7.8	6.3-9.6	12.6	3.2	2.5-4.1
	1986-1990	132.4	20.1	18.6-21.7	71.0	26.8	24.0-29.8	61.4	15.5	13.8-17.3	24.4	3.6	3.0-4.3	11.8	5.0	3.8-6.6	12.6	3.0	2.3-3.9
	1991-1995	132.0	18.8	17.4-20.3	68.4	23.8	21.3-26.5	63.6	15.3	13.7-17.1	27.8	4.0	3.4-4.7	13.6	5.2	4.0-6.6	14.2	3.4	2.6-4.3
	1996-2000	141.4	19.1	17.7-20.5	77.4	25.1	22.6-27.7	64.0	15.0	13.3-16.7	30.0	4.0	3.4-4.7	15.8	5.2	4.1-6.5	14.2	3.1	2.4-3.9
	2001-2005	112.6	14.3	13.1-15.5	60.2	17.7	15.8-19.9	52.4	11.7	10.3-13.2	26.8	3.3	2.7-3.9	12.2	3.8	2.9-4.8	14.6	3.0	2.4-3.8
	2006-2010	107.2	12.8	11.7-13.9	55.6	15.0	13.3-16.9	51.6	11.0	9.7-12.5	21.8	2.6	2.1-3.1	11.0	3.0	2.2-3.9	10.8	2.2	1.6-2.9
	2011-2015	104.4	11.9	10.9-13.0	62.2	15.3	13.6-17.1	42.2	9.1	7.9-10.5	23.4	2.6	2.1-3.1	14.4	3.6	2.8-4.6	9.0	1.8	1.3-2.5
2016-2020	96.6	10.4	9.4-11.4	55.0	12.7	11.2-14.4	41.6	8.4	7.2-9.6	27.8	2.8	2.3-3.3	16.4	3.7	2.9-4.6	11.4	2.0	1.5-2.7	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Monroe**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	17.4	2.8	2.3-3.5	9.6	3.8	2.8-5.1	7.8	2.1	1.5-2.9	16.0	2.6	2.1-3.3	8.4	3.4	2.4-4.6	7.6	2.1	1.5-2.9
	1981-1985	19.2	3.0	2.4-3.6	12.0	4.4	3.4-5.7	7.2	1.9	1.3-2.7	14.4	2.3	1.8-2.9	9.2	3.6	2.6-4.8	5.2	1.4	0.9-2.0
	1986-1990	18.2	2.7	2.2-3.3	11.6	4.1	3.1-5.3	6.6	1.7	1.1-2.4	15.4	2.3	1.8-2.9	8.0	3.1	2.2-4.2	7.4	1.9	1.3-2.6
	1991-1995	26.8	3.8	3.2-4.5	17.0	5.8	4.6-7.2	9.8	2.3	1.7-3.1	24.2	3.4	2.8-4.1	16.0	5.6	4.4-7.0	8.2	1.9	1.4-2.6
	1996-2000	32.6	4.4	3.8-5.1	21.0	6.6	5.4-8.0	11.6	2.6	2.0-3.4	26.4	3.5	2.9-4.2	15.8	5.2	4.1-6.5	10.6	2.3	1.7-3.0
	2001-2005	39.0	5.0	4.3-5.8	26.2	7.6	6.4-9.0	12.8	2.9	2.2-3.7	35.8	4.5	3.9-5.2	21.6	6.3	5.1-7.6	14.2	3.0	2.4-3.9
	2006-2010	58.0	6.9	6.1-7.8	43.4	11.7	10.2-13.4	14.6	3.1	2.5-4.0	44.8	5.3	4.6-6.1	28.0	7.8	6.6-9.2	16.8	3.3	2.6-4.2
	2011-2015	63.6	6.8	6.1-7.6	43.0	10.2	8.8-11.7	20.6	4.0	3.3-4.9	51.8	5.6	4.9-6.3	34.2	8.3	7.1-9.7	17.6	3.4	2.7-4.2
	2016-2020	72.0	7.2	6.4-8.0	46.8	10.2	8.9-11.6	25.2	4.7	3.9-5.6	57.6	5.8	5.2-6.6	35.8	8.1	6.9-9.4	21.8	4.2	3.4-5.1
Pancreas	1976-1980	68.0	11.1	10.0-12.4	34.8	14.2	12.1-16.6	33.2	9.1	7.7-10.6	65.0	10.6	9.5-11.9	30.8	12.7	10.7-14.9	34.2	9.3	7.9-10.8
	1981-1985	71.8	11.1	10.0-12.3	36.2	13.3	11.4-15.5	35.6	9.1	7.8-10.6	70.4	11.0	9.9-12.2	35.8	13.5	11.5-15.7	34.6	8.9	7.6-10.3
	1986-1990	77.4	11.9	10.7-13.1	35.4	14.0	11.9-16.3	42.0	10.5	9.1-12.1	73.2	11.2	10.1-12.4	33.8	13.1	11.2-15.3	39.4	9.9	8.6-11.5
	1991-1995	90.0	12.8	11.6-14.0	41.0	14.6	12.7-16.8	49.0	11.6	10.1-13.1	86.8	12.3	11.2-13.5	41.4	14.9	12.9-17.1	45.4	10.6	9.2-12.1
	1996-2000	96.6	12.9	11.8-14.1	46.2	15.0	13.1-17.1	50.4	11.1	9.7-12.5	88.4	11.7	10.7-12.9	42.8	14.1	12.3-16.2	45.6	9.8	8.5-11.2
	2001-2005	102.4	12.8	11.8-14.0	51.4	15.5	13.7-17.5	51.0	10.8	9.5-12.2	90.8	11.3	10.3-12.4	41.2	12.6	10.9-14.4	49.6	10.4	9.1-11.8
	2006-2010	115.0	13.6	12.5-14.8	57.0	15.7	13.9-17.7	58.0	11.9	10.5-13.4	105.0	12.2	11.1-13.3	51.8	14.3	12.6-16.2	53.2	10.5	9.3-11.9
	2011-2015	127.2	13.9	12.8-15.0	64.4	16.1	14.3-18.0	62.8	12.2	10.8-13.7	115.4	12.5	11.5-13.6	57.0	14.6	12.9-16.5	58.4	10.8	9.6-12.2
	2016-2020	147.0	15.1	14.0-16.3	75.4	17.7	15.9-19.6	71.6	13.0	11.6-14.5	123.0	12.4	11.4-13.5	63.8	14.8	13.2-16.6	59.2	10.5	9.3-11.8
Larynx	1976-1980	39.8	6.2	5.4-7.1	34.0	12.3	10.5-14.3	5.8	1.6	1.1-2.3	8.4	1.3	1.0-1.8	6.6	2.6	1.8-3.7	1.8	0.5	0.2-1.0
	1981-1985	37.0	5.7	4.9-6.6	28.8	10.6	8.9-12.5	8.2	2.2	1.6-3.0	12.4	1.9	1.5-2.4	9.8	3.8	2.8-5.1	2.6	0.7	0.4-1.2
	1986-1990	38.0	5.8	5.0-6.7	29.4	10.5	8.8-12.3	8.6	2.4	1.7-3.3	10.6	1.7	1.2-2.2	8.4	3.2	2.3-4.3	2.2	0.6	0.3-1.0
	1991-1995	39.0	5.6	4.8-6.4	30.4	10.0	8.4-11.7	8.6	2.2	1.6-3.0	11.0	1.5	1.2-2.0	9.2	3.2	2.3-4.2	1.8	0.4	0.2-0.8
	1996-2000	35.4	4.9	4.2-5.7	27.4	8.6	7.2-10.1	8.0	2.0	1.4-2.7	10.8	1.5	1.1-1.9	8.2	2.6	1.9-3.6	2.6	0.6	0.3-1.1
	2001-2005	35.4	4.6	3.9-5.3	27.0	7.8	6.5-9.2	8.4	2.0	1.4-2.7	10.4	1.3	1.0-1.7	7.2	2.1	1.5-2.9	3.2	0.7	0.4-1.2
	2006-2010	31.2	3.7	3.1-4.3	24.6	6.6	5.5-7.9	6.6	1.4	1.0-2.0	9.8	1.1	0.8-1.5	7.0	1.9	1.3-2.7	2.8	0.6	0.3-1.0
	2011-2015	30.0	3.2	2.7-3.8	25.0	6.1	5.0-7.3	5.0	1.0	0.6-1.5	12.4	1.4	1.0-1.8	9.4	2.4	1.7-3.2	3.0	0.6	0.3-1.0
	2016-2020	27.2	2.7	2.3-3.2	20.4	4.5	3.6-5.5	6.8	1.3	0.9-1.9	9.0	0.9	0.6-1.2	6.2	1.3	0.9-1.9	2.8	0.5	0.3-0.9
Lung and bronchus	1976-1980	357.0	56.8	54.1-59.5	249.2	96.6	91.1-102.3	107.8	30.2	27.6-32.9	283.8	45.4	43.0-47.9	201.4	79.3	74.3-84.6	82.4	22.8	20.7-25.2
	1981-1985	399.2	61.1	58.4-63.8	263.6	98.9	93.4-104.5	135.6	36.3	33.6-39.2	319.8	49.1	46.7-51.6	218.6	83.1	78.1-88.3	101.2	27.2	24.8-29.7
	1986-1990	436.6	65.6	62.8-68.4	264.6	96.6	91.3-102.1	172.0	45.5	42.4-48.7	339.8	51.0	48.6-53.5	207.0	77.0	72.2-82.0	132.8	34.5	31.9-37.3
	1991-1995	481.4	68.3	65.6-71.1	277.0	95.2	90.2-100.4	204.4	50.9	47.8-54.2	383.8	54.3	51.9-56.8	227.0	78.7	74.1-83.5	156.8	38.4	35.7-41.2
	1996-2000	500.6	68.0	65.4-70.7	266.4	86.1	81.5-90.9	234.2	55.7	52.5-59.0	406.8	54.9	52.6-57.4	219.4	71.8	67.6-76.2	187.4	43.8	41.0-46.7
	2001-2005	506.4	65.1	62.6-67.7	265.6	81.0	76.7-85.5	240.8	53.8	50.8-57.0	396.2	50.6	48.4-52.9	205.0	63.1	59.3-67.1	191.2	41.7	39.1-44.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Monroe**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	563.2	68.0	65.5-70.6	284.4	79.2	75.1-83.5	278.8	60.0	56.8-63.3	413.4	49.5	47.4-51.7	214.8	60.5	56.9-64.3	198.6	41.8	39.2-44.5
	2011-2015	568.4	63.0	60.7-65.4	280.2	71.7	67.9-75.6	288.2	56.8	53.9-59.9	402.0	44.5	42.5-46.5	213.8	55.0	51.7-58.5	188.2	36.8	34.5-39.4
	2016-2020	614.6	61.5	59.3-63.7	297.8	67.6	64.1-71.2	316.8	57.5	54.6-60.4	346.0	34.8	33.1-36.5	180.8	42.0	39.3-44.9	165.2	29.7	27.6-31.8
Melanoma of the skin	1976-1980	52.8	8.5	7.5-9.6	31.8	11.3	9.6-13.3	21.0	6.2	5.0-7.5	15.2	2.5	2.0-3.2	10.6	4.0	3.0-5.3	4.6	1.4	0.9-2.1
	1981-1985	62.0	9.8	8.7-11.0	34.0	12.1	10.3-14.2	28.0	8.1	6.8-9.6	13.4	2.1	1.6-2.6	8.2	2.9	2.1-4.0	5.2	1.4	0.9-2.1
	1986-1990	65.6	9.9	8.9-11.1	38.0	13.5	11.5-15.6	27.6	7.7	6.4-9.1	17.2	2.7	2.1-3.3	9.8	3.7	2.7-5.0	7.4	2.0	1.4-2.8
	1991-1995	62.4	8.9	8.0-10.0	35.4	11.6	9.9-13.4	27.0	6.9	5.8-8.2	17.2	2.5	2.0-3.1	10.4	3.5	2.6-4.6	6.8	1.7	1.2-2.4
	1996-2000	81.6	11.1	10.1-12.3	44.2	13.7	12.0-15.7	37.4	9.4	8.1-10.8	16.2	2.2	1.8-2.7	9.4	3.0	2.2-3.9	6.8	1.6	1.1-2.3
	2001-2005	100.8	13.0	11.9-14.2	59.8	17.4	15.5-19.5	41.0	9.8	8.5-11.3	15.6	2.0	1.6-2.5	9.2	2.7	2.0-3.6	6.4	1.5	1.0-2.1
	2006-2010	157.2	19.6	18.3-21.1	86.0	24.0	21.8-26.4	71.2	16.9	15.1-18.8	20.6	2.4	2.0-2.9	13.4	3.6	2.8-4.6	7.2	1.5	1.1-2.2
	2011-2015	213.0	25.3	23.8-27.0	117.2	30.6	28.1-33.2	95.8	22.1	20.0-24.2	21.2	2.3	1.9-2.8	11.4	3.1	2.3-4.0	9.8	1.8	1.3-2.4
2016-2020	245.8	27.3	25.8-29.0	134.2	32.6	30.1-35.2	111.6	24.0	22.0-26.2	17.8	1.8	1.5-2.3	11.0	2.7	2.0-3.5	6.8	1.2	0.8-1.8	
Female breast	1976-1980							390.4	113.4	108.4-118.7							137.2	39.2	36.2-42.3
	1981-1985							413.0	115.8	110.7-121.0							139.2	37.5	34.7-40.5
	1986-1990							509.0	139.0	133.6-144.7							146.4	38.9	36.1-41.9
	1991-1995							534.0	137.9	132.6-143.3							133.8	33.0	30.5-35.6
	1996-2000							623.8	154.0	148.6-159.5							118.0	27.7	25.5-30.1
	2001-2005							590.6	137.3	132.3-142.4							108.4	23.6	21.6-25.8
	2006-2010							630.8	140.1	135.1-145.2							109.4	22.8	20.9-24.9
	2011-2015							690.4	147.4	142.3-152.5							97.0	19.0	17.3-20.9
2016-2020							726.4	148.1	143.1-153.3							98.2	18.5	16.8-20.3	
Cervix uteri	1976-1980							32.2	9.9	8.4-11.6							12.8	3.8	2.9-4.9
	1981-1985							30.0	8.7	7.3-10.3							12.6	3.7	2.8-4.7
	1986-1990							28.8	8.0	6.7-9.4							7.2	2.0	1.4-2.7
	1991-1995							34.4	9.0	7.7-10.4							6.4	1.6	1.1-2.2
	1996-2000							32.0	8.1	6.9-9.4							9.4	2.4	1.7-3.2
	2001-2005							23.2	5.8	4.8-6.9							6.0	1.4	0.9-2.0
	2006-2010							24.2	6.0	5.0-7.3							6.8	1.5	1.0-2.1
	2011-2015							27.8	6.6	5.5-7.8							11.4	2.5	1.9-3.3
	2016-2020							23.0	5.5	4.4-6.6							8.8	1.9	1.3-2.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Monroe**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							112.8	31.5	28.9-34.3							20.0	5.5	4.5-6.7
	1981-1985							90.0	24.8	22.5-27.3							16.2	4.3	3.4-5.3
	1986-1990							80.8	21.3	19.3-23.6							18.6	4.7	3.8-5.8
	1991-1995							100.2	25.7	23.4-28.1							20.6	4.9	4.0-5.9
	1996-2000							120.8	30.0	27.6-32.5							18.4	4.2	3.4-5.2
	2001-2005							120.0	27.8	25.6-30.1							18.0	3.8	3.0-4.7
	2006-2010							143.4	31.1	28.8-33.5							23.8	4.7	3.8-5.6
	2011-2015							156.8	31.4	29.2-33.7							28.4	5.5	4.6-6.5
	2016-2020							171.4	32.5	30.3-34.8							38.6	6.8	5.8-7.8
Ovary	1976-1980							62.8	17.5	15.6-19.6							44.6	12.5	10.9-14.2
	1981-1985							62.2	17.2	15.3-19.3							40.0	10.6	9.2-12.2
	1986-1990							58.2	16.0	14.1-17.9							39.4	10.3	8.9-11.8
	1991-1995							64.4	16.5	14.8-18.5							42.4	10.6	9.2-12.1
	1996-2000							70.8	17.3	15.5-19.2							42.8	10.0	8.7-11.4
	2001-2005							67.8	15.5	13.9-17.3							47.8	10.2	8.9-11.6
	2006-2010							54.8	11.9	10.5-13.5							44.4	9.4	8.1-10.7
	2011-2015							60.4	12.7	11.2-14.3							40.8	7.9	6.8-9.1
	2016-2020							47.0	9.6	8.3-11.0							34.0	6.1	5.2-7.1
Prostate	1976-1980				197.0	91.2	85.4-97.3							68.2	34.0	30.3-37.9			
	1981-1985				239.0	102.6	96.6-108.8							71.8	34.3	30.7-38.2			
	1986-1990				354.6	143.8	136.9-150.9							75.0	33.5	30.1-37.2			
	1991-1995				643.8	224.4	216.5-232.4							84.6	34.5	31.2-38.0			
	1996-2000				587.6	190.8	183.9-197.9							84.4	30.6	27.7-33.7			
	2001-2005				596.4	178.8	172.4-185.4							82.4	26.9	24.4-29.7			
	2006-2010				650.2	175.9	169.8-182.1							66.8	19.9	17.8-22.2			
	2011-2015				554.4	131.8	126.8-136.9							68.8	18.7	16.7-20.8			
	2016-2020				715.2	152.5	147.5-157.8							77.8	19.2	17.4-21.3			
Testis	1976-1980				16.8	4.9	3.9-6.2							2.6	0.7	0.4-1.2			
	1981-1985				17.2	4.6	3.7-5.8							1.0	0.3	0.1-0.8			
	1986-1990				20.6	5.6	4.5-6.8							0.6	0.2	0.0-0.6			
	1991-1995				19.0	5.1	4.1-6.3							0.6	0.2	0.0-0.6			
	1996-2000				27.6	7.7	6.4-9.1							1.0	0.3	0.1-0.7			
	2001-2005				22.2	6.3	5.2-7.6							0.6	0.2	0.0-0.5			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Monroe**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				21.0	6.2	5.1-7.5							0.4	0.1	0.0-0.4			
	2011-2015				19.6	5.8	4.7-7.1							0.4	0.1	0.0-0.4			
	2016-2020				26.4	7.8	6.5-9.2							0.6	0.2	0.0-0.5			
Urinary bladder (incl. in situ)	1976-1980	143.8	23.6	21.9-25.4	103.4	42.7	38.9-46.7	40.4	11.4	9.9-13.1	38.2	6.6	5.7-7.6	25.2	12.0	9.9-14.3	13.0	3.6	2.7-4.6
	1981-1985	141.2	22.1	20.5-23.8	100.8	41.6	37.9-45.6	40.4	10.2	8.9-11.8	40.2	6.5	5.6-7.4	28.4	13.1	11.0-15.5	11.8	3.0	2.3-3.9
	1986-1990	155.2	23.2	21.6-24.9	111.4	41.6	38.1-45.3	43.8	11.2	9.7-12.8	36.6	5.6	4.8-6.4	24.0	10.3	8.5-12.4	12.6	3.0	2.3-3.9
	1991-1995	173.2	24.6	23.0-26.3	128.2	45.5	41.9-49.2	45.0	10.7	9.4-12.2	34.2	4.8	4.1-5.6	22.8	9.2	7.5-11.1	11.4	2.5	1.9-3.3
	1996-2000	205.2	27.6	26.0-29.4	149.0	49.2	45.7-52.9	56.2	13.0	11.5-14.6	45.8	5.9	5.2-6.8	28.4	10.2	8.5-12.0	17.4	3.6	2.8-4.4
	2001-2005	217.2	27.5	25.9-29.2	158.6	48.7	45.3-52.2	58.6	12.4	11.0-14.0	44.8	5.5	4.8-6.2	30.6	9.7	8.2-11.4	14.2	2.8	2.2-3.6
	2006-2010	211.8	25.4	23.9-27.0	162.8	46.9	43.7-50.3	49.0	10.2	8.9-11.6	44.6	5.0	4.4-5.7	32.6	9.4	8.0-11.0	12.0	2.2	1.6-2.8
	2011-2015	222.0	24.7	23.2-26.3	166.6	43.7	40.8-46.9	55.4	10.7	9.4-12.1	42.8	4.5	3.9-5.1	32.8	8.8	7.5-10.3	10.0	1.6	1.2-2.2
2016-2020	234.0	23.6	22.2-25.0	175.4	41.1	38.4-44.0	58.6	10.5	9.3-11.8	40.2	4.0	3.4-4.6	27.8	6.8	5.7-8.0	12.4	2.0	1.5-2.6	
Kidney and renal pelvis	1976-1980	51.0	8.3	7.3-9.4	31.6	12.2	10.3-14.3	19.4	5.5	4.4-6.7	25.4	4.1	3.4-4.9	16.0	6.1	4.8-7.7	9.4	2.6	1.9-3.5
	1981-1985	61.6	9.6	8.5-10.7	37.2	14.0	12.0-16.3	24.4	6.7	5.5-8.0	24.2	3.7	3.1-4.4	15.4	5.8	4.6-7.4	8.8	2.2	1.6-3.0
	1986-1990	70.2	10.6	9.5-11.8	45.4	16.4	14.3-18.8	24.8	6.4	5.3-7.7	27.6	4.1	3.5-4.9	16.4	6.2	4.9-7.8	11.2	2.8	2.1-3.7
	1991-1995	74.2	10.7	9.6-11.8	46.8	15.9	13.9-18.1	27.4	6.9	5.8-8.2	32.4	4.7	4.0-5.5	20.8	7.5	6.1-9.1	11.6	2.8	2.1-3.6
	1996-2000	112.2	15.3	14.1-16.6	67.0	21.1	18.9-23.5	45.2	10.8	9.5-12.4	34.0	4.5	3.9-5.2	19.6	6.5	5.3-8.0	14.4	3.1	2.4-3.9
	2001-2005	116.4	14.9	13.7-16.2	68.4	19.9	17.9-22.2	48.0	10.9	9.5-12.4	34.6	4.3	3.7-5.0	20.6	6.2	5.0-7.5	14.0	2.8	2.2-3.6
	2006-2010	160.0	19.2	17.9-20.6	101.2	27.4	25.0-29.9	58.8	12.8	11.4-14.4	32.0	3.7	3.2-4.4	21.6	6.0	4.9-7.3	10.4	2.1	1.5-2.8
	2011-2015	155.4	17.8	16.5-19.1	98.8	24.9	22.7-27.2	56.6	11.7	10.3-13.2	33.6	3.7	3.1-4.3	21.2	5.5	4.5-6.7	12.4	2.3	1.8-3.0
2016-2020	183.8	19.8	18.5-21.2	123.4	28.8	26.5-31.3	60.4	12.2	10.8-13.8	29.8	3.0	2.5-3.5	19.0	4.5	3.6-5.5	10.8	1.9	1.4-2.5	
Brain and other central nervous system	1976-1980	43.2	6.4	5.6-7.4	24.2	7.9	6.5-9.5	19.0	5.2	4.2-6.4	30.4	4.6	3.9-5.4	17.2	5.9	4.7-7.3	13.2	3.6	2.8-4.6
	1981-1985	43.0	6.6	5.7-7.5	24.2	8.3	6.8-10.0	18.8	5.3	4.3-6.6	27.2	4.2	3.5-5.0	15.6	5.7	4.4-7.1	11.6	3.1	2.4-4.1
	1986-1990	46.4	6.8	5.9-7.8	26.2	8.7	7.2-10.3	20.2	5.3	4.3-6.4	29.4	4.4	3.7-5.1	16.4	5.5	4.4-6.9	13.0	3.3	2.6-4.2
	1991-1995	51.4	7.1	6.3-8.1	26.6	8.1	6.8-9.6	24.8	6.3	5.2-7.5	33.6	4.8	4.1-5.6	17.8	5.7	4.5-7.0	15.8	4.0	3.2-5.0
	1996-2000	50.2	6.9	6.0-7.8	27.6	8.2	6.9-9.7	22.6	5.7	4.7-6.8	33.6	4.6	3.9-5.3	17.6	5.5	4.4-6.7	16.0	3.9	3.1-4.8
	2001-2005	55.8	7.4	6.5-8.3	29.8	8.6	7.3-10.1	26.0	6.4	5.3-7.6	33.6	4.3	3.7-5.1	17.8	5.2	4.1-6.4	15.8	3.7	2.9-4.6
	2006-2010	50.4	6.5	5.7-7.3	24.4	7.0	5.8-8.4	26.0	6.2	5.1-7.4	36.0	4.4	3.8-5.2	20.0	5.6	4.6-6.8	16.0	3.6	2.8-4.5
	2011-2015	50.6	6.2	5.4-7.0	24.6	6.3	5.2-7.5	26.0	6.1	5.1-7.3	35.6	4.1	3.5-4.8	18.0	4.5	3.6-5.6	17.6	3.8	3.1-4.8
2016-2020	53.8	6.3	5.5-7.1	28.6	7.1	6.0-8.5	25.2	5.6	4.6-6.7	41.0	4.6	3.9-5.3	21.8	5.3	4.3-6.5	19.2	4.0	3.2-5.0	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Monroe**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	25.0	3.8	3.2-4.6	7.4	2.6	1.8-3.6	17.6	4.9	3.9-6.1	3.8	0.6	0.4-1.0	1.6	0.6	0.3-1.3	2.2	0.6	0.3-1.1
	1981-1985	26.4	4.0	3.3-4.8	6.4	2.2	1.5-3.2	20.0	5.6	4.6-6.9	1.6	0.3	0.1-0.5	0.2	0.1	0.0-0.6	1.4	0.4	0.2-0.8
	1986-1990	32.2	4.8	4.0-5.6	9.6	3.3	2.4-4.4	22.6	6.2	5.1-7.5	2.8	0.4	0.2-0.7	1.4	0.5	0.2-1.0	1.4	0.3	0.1-0.7
	1991-1995	37.4	5.2	4.5-6.0	10.0	3.1	2.3-4.2	27.4	7.2	6.0-8.5	3.2	0.5	0.3-0.7	1.6	0.5	0.2-1.0	1.6	0.4	0.2-0.8
	1996-2000	50.2	6.9	6.0-7.8	11.4	3.3	2.5-4.3	38.8	10.1	8.7-11.6	3.2	0.4	0.2-0.7	0.8	0.2	0.1-0.6	2.4	0.5	0.3-1.0
	2001-2005	61.6	8.2	7.3-9.2	17.0	4.8	3.8-5.9	44.6	11.5	10.0-13.1	4.8	0.6	0.4-0.9	2.8	0.8	0.4-1.3	2.0	0.4	0.2-0.8
	2006-2010	98.8	12.8	11.7-14.0	27.8	7.6	6.3-9.0	71.0	17.8	15.9-19.8	3.8	0.4	0.3-0.7	1.8	0.5	0.2-1.0	2.0	0.4	0.2-0.7
	2011-2015	121.6	15.5	14.2-16.8	35.8	9.2	7.9-10.8	85.8	21.3	19.3-23.5	5.4	0.5	0.4-0.8	1.8	0.4	0.2-0.8	3.6	0.6	0.4-1.0
2016-2020	117.0	14.5	13.3-15.8	37.8	9.6	8.2-11.1	79.2	19.1	17.1-21.1	4.8	0.5	0.3-0.8	1.6	0.4	0.2-0.8	3.2	0.6	0.3-1.0	
Hodgkin lymphoma	1976-1980	29.6	4.0	3.4-4.8	16.0	4.7	3.7-6.0	13.6	3.4	2.6-4.3	9.0	1.4	1.0-1.9	4.6	1.7	1.1-2.6	4.4	1.2	0.8-1.9
	1981-1985	22.2	3.1	2.5-3.7	12.4	3.7	2.8-4.8	9.8	2.5	1.8-3.3	5.6	0.8	0.6-1.2	3.8	1.3	0.8-2.1	1.8	0.4	0.2-0.8
	1986-1990	28.2	3.8	3.2-4.5	15.4	4.4	3.4-5.5	12.8	3.3	2.5-4.2	4.4	0.6	0.4-0.9	1.8	0.6	0.2-1.1	2.6	0.7	0.3-1.1
	1991-1995	29.6	3.9	3.3-4.6	14.8	4.0	3.2-5.1	14.8	3.7	2.9-4.7	4.4	0.6	0.4-0.9	2.0	0.5	0.3-1.0	2.4	0.6	0.3-1.1
	1996-2000	25.0	3.4	2.8-4.0	12.2	3.5	2.6-4.4	12.8	3.3	2.6-4.3	4.0	0.5	0.3-0.8	2.6	0.8	0.4-1.4	1.4	0.3	0.1-0.7
	2001-2005	26.4	3.6	3.0-4.3	14.0	4.0	3.1-5.1	12.4	3.3	2.5-4.2	4.2	0.5	0.3-0.8	2.2	0.6	0.3-1.2	2.0	0.5	0.2-0.9
	2006-2010	24.8	3.4	2.8-4.0	11.8	3.4	2.6-4.4	13.0	3.4	2.6-4.3	3.6	0.5	0.3-0.8	1.6	0.5	0.2-1.0	2.0	0.5	0.2-0.9
	2011-2015	26.8	3.4	2.8-4.0	12.0	3.1	2.4-4.0	14.8	3.7	2.8-4.6	1.8	0.2	0.1-0.4	0.6	0.2	0.0-0.5	1.2	0.3	0.1-0.6
2016-2020	26.8	3.6	3.0-4.2	16.8	4.7	3.7-5.9	10.0	2.6	1.9-3.4	2.8	0.3	0.2-0.5	2.0	0.5	0.2-0.9	0.8	0.1	0.0-0.4	
Non-Hodgkin lymphomas	1976-1980	86.0	13.8	12.5-15.2	48.0	18.3	15.9-20.9	38.0	10.6	9.1-12.2	45.6	7.3	6.4-8.3	22.8	9.1	7.5-11.1	22.8	6.1	5.0-7.4
	1981-1985	102.0	15.7	14.4-17.2	51.0	19.2	16.8-21.8	51.0	13.5	11.8-15.3	47.0	7.3	6.4-8.3	24.2	9.4	7.7-11.3	22.8	5.9	4.8-7.1
	1986-1990	128.4	19.2	17.7-20.8	70.8	25.3	22.6-28.2	57.6	14.8	13.1-16.6	58.2	8.8	7.8-9.8	28.6	11.0	9.2-13.0	29.6	7.4	6.2-8.7
	1991-1995	136.8	19.3	17.9-20.9	73.2	24.1	21.6-26.7	63.6	15.6	13.9-17.5	65.0	9.1	8.1-10.1	34.8	12.0	10.3-14.0	30.2	6.9	5.8-8.1
	1996-2000	162.4	21.9	20.5-23.5	77.8	24.5	22.1-27.1	84.6	20.0	18.1-22.0	62.8	8.4	7.5-9.4	32.0	10.4	8.8-12.1	30.8	6.9	5.8-8.1
	2001-2005	160.2	20.5	19.1-22.0	83.2	24.6	22.3-27.1	77.0	17.3	15.6-19.2	59.0	7.4	6.5-8.2	29.6	9.1	7.7-10.7	29.4	6.0	5.0-7.1
	2006-2010	191.2	23.2	21.7-24.7	99.4	27.6	25.2-30.2	91.8	19.7	17.9-21.7	59.2	6.9	6.1-7.7	29.0	8.1	6.8-9.5	30.2	5.8	4.9-6.9
	2011-2015	192.2	21.8	20.4-23.2	105.0	27.1	24.8-29.6	87.2	17.6	15.9-19.4	50.6	5.6	4.9-6.4	27.2	7.4	6.2-8.8	23.4	4.3	3.5-5.2
2016-2020	205.2	21.8	20.5-23.3	111.0	26.2	24.0-28.6	94.2	18.5	16.8-20.3	55.4	5.6	5.0-6.3	28.6	6.9	5.8-8.1	26.8	4.7	3.9-5.6	
Myeloma	1976-1980	33.8	5.6	4.8-6.6	14.2	6.2	4.8-7.9	19.6	5.4	4.4-6.6	24.0	4.0	3.3-4.8	11.0	4.6	3.4-6.1	13.0	3.6	2.8-4.6
	1981-1985	29.6	4.7	4.0-5.5	16.0	6.3	5.0-7.9	13.6	3.5	2.7-4.5	21.4	3.3	2.7-4.1	10.8	4.5	3.3-5.9	10.6	2.7	2.0-3.5
	1986-1990	44.2	6.9	6.0-7.8	22.2	8.5	6.9-10.2	22.0	5.6	4.6-6.8	25.8	4.0	3.3-4.8	12.6	5.1	3.8-6.5	13.2	3.3	2.6-4.2
	1991-1995	41.4	5.9	5.1-6.8	22.2	8.1	6.7-9.8	19.2	4.6	3.8-5.7	31.4	4.4	3.8-5.2	15.2	5.7	4.4-7.1	16.2	3.7	3.0-4.7
	1996-2000	48.4	6.5	5.7-7.4	24.4	7.9	6.5-9.4	24.0	5.6	4.6-6.7	31.0	4.1	3.5-4.8	15.6	5.3	4.2-6.6	15.4	3.5	2.7-4.4
	2001-2005	44.8	5.7	4.9-6.5	24.6	7.5	6.2-9.0	20.2	4.4	3.6-5.4	26.2	3.3	2.7-3.9	14.2	4.5	3.5-5.7	12.0	2.4	1.8-3.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Monroe**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	57.0	6.8	6.0-7.6	32.2	9.0	7.6-10.5	24.8	5.2	4.3-6.2	28.8	3.4	2.8-4.0	16.2	4.5	3.6-5.6	12.6	2.6	2.0-3.4
	2011-2015	63.2	6.9	6.2-7.7	35.2	8.8	7.5-10.2	28.0	5.5	4.6-6.5	34.0	3.6	3.1-4.3	19.6	5.0	4.0-6.1	14.4	2.6	2.0-3.3
	2016-2020	70.2	7.1	6.4-7.9	41.4	9.5	8.3-11.0	28.8	5.3	4.4-6.3	28.8	2.9	2.4-3.4	15.0	3.6	2.8-4.5	13.8	2.4	1.8-3.0
Leukemias	1976-1980	86.6	14.2	12.9-15.6	46.2	19.0	16.5-21.7	40.4	11.4	9.9-13.1	58.6	9.7	8.6-10.8	32.0	13.1	11.0-15.4	26.6	7.5	6.3-9.0
	1981-1985	81.6	12.7	11.5-14.0	42.6	16.4	14.2-18.9	39.0	10.5	9.0-12.1	48.6	7.7	6.7-8.7	25.6	10.7	8.8-12.8	23.0	5.9	4.9-7.2
	1986-1990	93.8	14.1	12.8-15.4	54.4	20.1	17.7-22.7	39.4	10.1	8.7-11.6	54.0	8.2	7.2-9.2	31.2	12.0	10.1-14.2	22.8	5.8	4.8-7.0
	1991-1995	104.4	14.7	13.5-16.1	61.0	21.1	18.7-23.7	43.4	10.7	9.3-12.3	51.2	7.2	6.4-8.2	28.0	10.4	8.7-12.3	23.2	5.4	4.4-6.5
	1996-2000	113.2	15.2	13.9-16.5	63.6	20.4	18.2-22.8	49.6	11.5	10.1-13.1	65.0	8.6	7.7-9.6	36.0	12.1	10.3-14.0	29.0	6.3	5.3-7.5
	2001-2005	126.6	16.0	14.8-17.3	70.2	20.9	18.7-23.2	56.4	12.4	10.9-14.0	63.4	7.8	7.0-8.7	35.6	11.0	9.4-12.8	27.8	5.7	4.8-6.8
	2006-2010	128.6	15.8	14.6-17.1	70.6	19.9	17.9-22.1	58.0	12.9	11.4-14.5	63.4	7.6	6.8-8.5	35.0	10.2	8.7-11.8	28.4	6.0	5.0-7.1
	2011-2015	157.6	18.0	16.7-19.3	88.4	23.2	21.0-25.5	69.2	14.2	12.6-15.8	69.2	7.6	6.8-8.5	36.8	9.7	8.4-11.3	32.4	6.1	5.1-7.1
2016-2020	135.2	14.3	13.2-15.5	78.6	18.6	16.8-20.6	56.6	10.9	9.6-12.3	58.6	6.1	5.4-6.9	34.4	8.5	7.2-9.9	24.2	4.4	3.6-5.4	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Montgomery**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	268.2	414.4	391.7-438.2	139.6	510.3	471.6-551.4	128.6	358.6	329.8-389.3	141.8	214.9	198.9-231.9	78.2	289.4	260.2-321.0	63.6	167.4	148.8-187.8
	1981-1985	284.2	432.4	409.3-456.5	143.8	524.3	485.2-565.8	140.4	383.7	354.0-415.3	139.0	207.1	191.5-223.7	75.2	280.5	251.8-311.6	63.8	162.9	144.5-183.1
	1986-1990	294.4	448.6	425.0-473.3	155.8	573.4	532.5-616.7	138.6	376.6	346.9-408.2	142.0	208.4	192.8-225.0	74.6	275.0	246.8-305.5	67.4	168.2	149.6-188.7
	1991-1995	352.4	523.0	498.0-549.1	190.4	680.1	636.6-725.8	162.0	428.8	397.7-461.8	156.8	224.8	208.8-241.7	84.2	305.6	276.5-337.0	72.6	178.2	159.2-199.1
	1996-2000	329.6	500.4	475.8-526.0	170.0	606.8	566.3-649.5	159.6	435.6	403.9-469.2	130.4	183.9	169.7-199.1	64.6	225.7	201.6-252.1	65.8	157.0	139.5-176.4
	2001-2005	336.8	526.7	501.1-553.3	167.0	599.2	559.0-641.7	169.8	477.7	444.2-513.1	123.0	176.5	162.4-191.5	59.6	207.2	184.2-232.4	63.4	157.4	139.5-177.2
	2006-2010	328.4	512.7	487.4-539.0	169.8	597.1	556.9-639.5	158.6	452.7	420.0-487.4	120.0	174.3	160.2-189.4	59.8	205.9	182.9-231.1	60.2	152.7	135.0-172.3
	2011-2015	323.0	497.5	472.7-523.5	156.0	528.2	490.7-567.9	167.0	480.4	446.7-516.1	110.8	162.5	148.8-177.1	58.0	195.8	173.4-220.4	52.8	138.4	121.5-157.4
	2016-2020	340.6	508.0	483.2-533.9	177.6	557.2	519.9-596.7	163.0	472.2	438.7-507.8	115.2	164.0	150.4-178.6	57.4	184.4	163.0-207.9	57.8	149.4	131.8-168.9
Oral cavity and pharynx	1976-1980	7.0	10.9	7.4-15.4	5.4	19.8	12.7-29.4	1.6	4.5	1.8-9.3	1.8	2.7	1.2-5.2	1.4	5.5	2.1-11.6	0.4	1.0	0.1-4.2
	1981-1985	6.0	8.6	5.7-12.5	4.2	13.7	8.3-21.6	1.8	5.5	2.3-11.0	1.6	2.6	1.1-5.4	1.0	3.8	1.2-9.2	0.6	2.4	0.4-7.1
	1986-1990	6.4	10.0	6.7-14.4	5.0	19.6	12.4-29.5	1.4	3.3	1.3-7.3	1.8	3.1	1.3-6.1	1.0	4.4	1.3-10.4	0.8	2.1	0.5-5.9
	1991-1995	8.4	12.4	8.8-17.1	4.6	15.6	9.8-23.8	3.8	9.7	5.6-15.9	2.0	2.5	1.2-4.9	1.0	3.2	1.0-7.9	1.0	2.0	0.7-5.5
	1996-2000	5.8	9.2	6.1-13.4	3.6	13.1	7.7-21.0	2.2	6.0	2.8-11.4	1.4	2.1	0.8-4.5	1.2	4.2	1.5-9.4	0.2	0.3	0.0-3.2
	2001-2005	8.2	13.3	9.4-18.3	5.6	20.8	13.7-30.3	2.6	6.6	3.4-12.0	1.6	2.4	1.0-5.0	0.8	2.7	0.7-7.3	0.8	2.6	0.7-7.0
	2006-2010	6.8	11.4	7.8-16.1	5.0	17.7	11.4-26.4	1.8	5.9	2.6-11.6	1.2	1.8	0.6-4.0	0.8	2.6	0.7-7.0	0.4	1.0	0.1-4.3
	2011-2015	7.8	13.0	9.1-18.0	5.8	20.5	13.5-29.9	2.0	6.2	2.8-12.0	2.6	4.2	2.2-7.4	2.2	7.6	3.7-14.0	0.4	1.2	0.1-4.9
	2016-2020	10.4	16.1	11.8-21.5	7.2	23.5	16.2-33.1	3.2	9.5	5.1-16.2	2.8	4.3	2.2-7.6	2.0	7.1	3.2-13.6	0.8	1.6	0.4-5.3
Esophagus	1976-1980	2.0	3.0	1.4-5.8	2.0	7.3	3.3-13.9	0.0	0.0	0.0-2.8	1.6	2.3	1.0-4.7	1.2	4.6	1.6-10.4	0.4	0.9	0.1-4.1
	1981-1985	2.2	3.2	1.6-5.9	1.4	4.8	1.9-10.4	0.8	2.2	0.5-6.3	2.2	2.9	1.4-5.5	1.6	5.1	2.1-10.5	0.6	1.3	0.3-4.6
	1986-1990	2.8	3.8	2.1-6.6	2.6	8.8	4.6-15.4	0.2	0.4	0.0-3.4	2.2	3.4	1.7-6.3	1.8	6.7	3.0-12.9	0.4	1.5	0.2-5.3
	1991-1995	5.0	7.3	4.6-11.0	2.8	9.9	5.4-16.9	2.2	5.0	2.4-9.6	3.4	5.3	3.0-8.7	2.4	9.1	4.6-16.1	1.0	2.5	0.8-6.5
	1996-2000	2.8	3.8	2.1-6.7	2.0	6.6	3.2-12.5	0.8	2.1	0.5-6.0	2.6	4.0	2.1-7.1	2.2	7.8	3.9-14.2	0.4	1.3	0.1-5.1
	2001-2005	2.6	4.1	2.1-7.1	1.6	5.6	2.4-11.3	1.0	3.2	1.0-7.9	3.2	4.7	2.7-7.9	2.2	7.7	3.8-14.0	1.0	2.6	0.9-6.7
	2006-2010	3.6	5.3	3.1-8.7	2.6	9.0	4.8-15.7	1.0	2.6	0.8-6.6	3.2	4.9	2.8-8.2	2.6	9.6	5.1-16.7	0.6	1.1	0.2-4.3
	2011-2015	3.2	5.1	2.8-8.5	2.4	8.3	4.2-14.8	0.8	2.3	0.5-6.7	3.4	5.7	3.2-9.3	2.8	9.9	5.4-16.9	0.6	2.2	0.4-6.7
	2016-2020	5.6	7.7	5.0-11.4	4.0	11.2	6.7-17.8	1.6	4.8	2.0-9.9	3.0	4.8	2.6-8.2	2.2	7.3	3.5-13.7	0.8	2.6	0.7-7.0
Stomach	1976-1980	7.4	10.8	7.5-15.2	3.6	13.5	7.8-21.9	3.8	8.8	5.3-14.1	5.8	8.3	5.5-12.3	3.2	11.0	6.0-18.6	2.6	6.1	3.2-10.9
	1981-1985	7.0	9.7	6.7-13.7	4.0	14.2	8.5-22.4	3.0	6.9	3.9-11.9	5.0	6.9	4.4-10.3	3.2	11.4	6.3-19.1	1.8	4.0	1.8-8.1
	1986-1990	4.6	7.0	4.3-10.8	3.6	13.2	7.6-21.3	1.0	3.0	0.8-7.6	4.4	6.0	3.7-9.4	3.2	11.6	6.4-19.3	1.2	2.5	0.9-6.3
	1991-1995	5.4	7.2	4.7-10.7	3.4	11.9	6.9-19.4	2.0	4.4	2.0-8.7	2.8	3.7	2.0-6.4	1.6	5.5	2.3-11.1	1.2	2.4	0.9-6.0
	1996-2000	5.4	8.2	5.3-12.1	3.6	13.3	7.8-21.2	1.8	4.4	1.9-9.1	2.6	3.6	1.9-6.3	1.6	5.3	2.3-10.7	1.0	2.3	0.7-6.2
	2001-2005	5.0	7.5	4.8-11.3	2.8	10.2	5.6-17.4	2.2	5.3	2.5-10.4	2.6	3.6	1.9-6.5	1.2	4.0	1.5-9.1	1.4	3.5	1.3-8.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Montgomery**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	4.8	7.2	4.6-10.9	3.0	10.1	5.6-17.0	1.8	4.2	1.8-8.8	1.8	2.3	1.0-4.8	1.0	3.4	1.1-8.2	0.8	1.2	0.3-4.4
	2011-2015	4.0	6.3	3.8-9.9	2.6	8.9	4.7-15.6	1.4	4.0	1.5-8.8	2.0	2.8	1.3-5.4	0.6	1.9	0.4-6.2	1.4	3.2	1.2-7.5
	2016-2020	4.0	6.1	3.6-9.6	2.6	7.6	4.0-13.6	1.4	4.8	1.9-10.3	1.0	1.5	0.5-3.8	0.6	1.5	0.3-5.2	0.4	1.5	0.2-5.7
Colorectal	1976-1980	51.6	76.2	67.0-86.3	28.6	104.3	87.3-123.7	23.0	57.1	47.0-69.0	25.0	37.2	30.8-44.6	13.8	52.6	40.4-67.3	11.2	27.0	20.4-35.5
	1981-1985	49.8	74.1	64.9-84.4	24.4	90.4	74.4-108.8	25.4	62.4	51.6-75.1	21.8	31.9	26.0-38.8	11.0	41.5	30.8-54.6	10.8	24.9	18.6-32.9
	1986-1990	48.4	70.2	61.3-80.0	23.4	87.1	71.5-105.1	25.0	57.4	47.4-69.2	21.2	30.3	24.7-36.9	11.4	43.0	32.2-56.3	9.8	21.4	15.8-28.8
	1991-1995	60.8	88.1	78.1-99.0	32.4	118.1	100.2-138.3	28.4	70.3	58.4-84.1	24.0	34.2	28.2-41.2	13.8	51.2	39.6-65.2	10.2	24.4	17.6-33.0
	1996-2000	47.8	68.5	59.9-78.2	22.0	80.0	65.6-96.7	25.8	61.4	50.6-74.0	18.6	25.1	20.1-31.0	5.8	21.9	14.6-31.6	12.8	28.0	21.1-36.7
	2001-2005	48.4	68.9	60.2-78.5	20.6	72.1	58.8-87.7	27.8	66.0	54.9-79.1	15.8	20.6	16.2-26.0	6.6	22.9	15.8-32.4	9.2	19.2	13.7-26.6
	2006-2010	40.2	58.6	50.5-67.7	19.6	68.4	55.3-83.7	20.6	50.9	41.0-62.7	12.4	17.0	12.9-22.2	4.8	16.3	10.4-24.6	7.6	17.8	12.3-25.4
	2011-2015	31.6	49.5	41.8-58.3	14.4	49.9	38.8-63.2	17.2	49.4	38.8-62.1	10.6	15.5	11.5-20.6	5.0	17.5	11.2-26.1	5.6	13.6	8.8-20.5
2016-2020	29.0	44.7	37.4-53.0	16.0	53.4	41.9-67.1	13.0	37.8	28.7-49.1	8.0	10.9	7.7-15.2	4.0	13.2	8.0-20.8	4.0	9.5	5.5-15.6	
Colon excluding rectum	1976-1980	38.8	57.2	49.3-66.1	21.6	79.3	64.5-96.5	17.2	42.3	33.8-52.7	19.2	27.7	22.4-34.1	10.2	38.0	27.9-50.6	9.0	21.7	15.8-29.4
	1981-1985	35.6	52.9	45.1-61.6	17.6	65.6	52.2-81.6	18.0	43.9	35.0-54.8	18.2	26.4	21.1-32.6	9.4	35.3	25.6-47.6	8.8	20.1	14.6-27.4
	1986-1990	35.2	51.0	43.5-59.5	17.8	65.9	52.4-81.8	17.4	40.3	31.9-50.5	17.2	24.2	19.2-30.1	10.2	37.7	27.7-50.0	7.0	15.3	10.6-21.7
	1991-1995	44.8	64.8	56.3-74.3	23.4	84.8	69.8-102.1	21.4	52.5	42.3-64.7	19.8	28.1	22.7-34.6	12.0	45.1	34.2-58.4	7.8	18.6	12.8-26.4
	1996-2000	35.2	49.7	42.4-58.0	15.4	56.4	44.4-70.8	19.8	46.0	36.9-57.0	15.0	19.8	15.4-25.1	4.4	16.9	10.6-25.7	10.6	22.6	16.6-30.5
	2001-2005	35.6	49.8	42.5-58.0	13.2	45.2	34.9-57.8	22.4	53.6	43.6-65.6	14.0	18.1	14.0-23.3	5.4	18.8	12.4-27.6	8.6	17.7	12.5-24.8
	2006-2010	31.2	44.6	37.7-52.6	14.0	48.9	38.0-62.1	17.2	41.5	32.7-52.2	10.2	13.8	10.2-18.5	3.0	9.8	5.4-16.4	7.2	17.1	11.7-24.6
	2011-2015	22.8	35.3	28.9-42.8	8.8	31.0	22.3-41.9	14.0	38.9	29.7-50.3	8.4	12.0	8.6-16.6	4.0	13.6	8.3-21.4	4.4	10.3	6.2-16.4
2016-2020	21.6	32.3	26.3-39.4	12.4	41.4	31.5-53.6	9.2	25.1	18.1-34.3	7.0	9.3	6.4-13.3	3.4	11.2	6.5-18.3	3.6	8.3	4.7-14.2	
Rectum & rectosigmoid	1976-1980	12.8	19.0	14.5-24.4	7.0	25.0	17.2-35.3	5.8	14.8	9.8-21.7	5.8	9.5	6.2-13.9	3.6	14.6	8.4-23.5	2.2	5.3	2.7-10.0
	1981-1985	14.2	21.3	16.4-27.2	6.8	24.7	16.7-35.2	7.4	18.5	12.8-26.1	3.6	5.5	3.2-9.0	1.6	6.1	2.5-12.5	2.0	4.8	2.2-9.5
	1986-1990	13.2	19.1	14.7-24.6	5.6	21.2	13.9-31.0	7.6	17.1	12.0-24.1	4.0	6.1	3.7-9.7	1.2	5.4	1.9-11.8	2.8	6.1	3.3-10.9
	1991-1995	16.0	23.3	18.3-29.3	9.0	33.3	24.1-44.9	7.0	17.8	12.0-25.5	4.2	6.1	3.7-9.5	1.8	6.2	2.8-12.0	2.4	5.7	2.8-10.7
	1996-2000	12.6	18.8	14.3-24.3	6.6	23.5	16.1-33.3	6.0	15.4	10.1-22.8	3.6	5.3	3.1-8.6	1.4	5.0	2.0-10.6	2.2	5.4	2.6-10.4
	2001-2005	12.8	19.1	14.6-24.6	7.4	26.9	18.9-37.2	5.4	12.4	7.9-18.9	1.8	2.4	1.1-4.9	1.2	4.1	1.5-9.2	0.6	1.5	0.3-5.2
	2006-2010	9.0	14.0	10.1-18.9	5.6	19.5	12.8-28.4	3.4	9.4	5.3-15.6	2.2	3.2	1.6-6.0	1.8	6.6	3.0-12.6	0.4	0.7	0.1-3.7
	2011-2015	8.8	14.2	10.2-19.3	5.6	18.9	12.4-27.7	3.2	10.4	5.8-17.4	2.2	3.5	1.7-6.4	1.0	3.8	1.2-9.1	1.2	3.3	1.2-7.9
2016-2020	7.4	12.4	8.5-17.4	3.6	11.9	6.8-19.5	3.8	12.7	7.4-20.5	1.0	1.6	0.5-4.0	0.6	2.0	0.4-6.4	0.4	1.1	0.1-4.7	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Montgomery**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	1.6	2.3	1.0-4.8	1.0	3.9	1.2-9.5	0.6	1.5	0.3-4.9	1.6	2.4	1.0-5.0	1.0	4.3	1.3-10.1	0.6	1.5	0.3-4.9
	1981-1985	1.6	2.6	1.1-5.3	0.6	2.3	0.4-6.9	1.0	3.0	0.9-7.6	0.6	1.0	0.2-3.2	0.4	1.6	0.2-5.9	0.2	1.0	0.0-5.0
	1986-1990	1.6	2.7	1.1-5.6	1.4	5.3	2.0-11.2	0.2	0.4	0.0-3.4	1.2	1.8	0.6-4.2	0.8	2.8	0.7-7.7	0.4	0.9	0.1-4.0
	1991-1995	3.4	5.1	2.9-8.4	2.4	9.0	4.6-16.0	1.0	2.4	0.7-6.3	2.6	3.6	1.9-6.3	2.0	7.3	3.4-13.6	0.6	1.1	0.2-4.1
	1996-2000	1.8	2.8	1.3-5.5	1.2	4.5	1.6-9.9	0.6	1.2	0.2-4.6	1.6	2.3	1.0-4.9	0.6	2.4	0.5-7.1	1.0	1.8	0.6-5.3
	2001-2005	2.2	3.3	1.6-6.2	1.8	6.2	2.8-12.0	0.4	1.0	0.1-4.5	2.0	3.0	1.4-5.7	2.0	6.8	3.2-12.8	0.0	0.0	0.0-2.9
	2006-2010	3.8	5.4	3.2-8.7	3.0	9.5	5.3-16.0	0.8	1.9	0.4-6.0	2.2	2.9	1.4-5.5	1.8	5.4	2.4-10.6	0.4	0.9	0.1-4.3
	2011-2015	3.0	3.9	2.2-6.7	1.4	3.6	1.4-8.1	1.6	4.2	1.8-9.0	1.8	2.3	1.0-4.6	1.0	2.8	0.9-7.1	0.8	2.0	0.5-5.9
	2016-2020	6.2	8.1	5.5-11.8	4.6	12.2	7.7-18.8	1.6	4.2	1.8-8.9	3.2	4.3	2.4-7.3	2.2	6.3	3.1-11.8	1.0	2.5	0.8-6.7
Pancreas	1976-1980	9.6	13.6	10.0-18.2	6.0	19.5	13.0-28.5	3.6	9.1	5.4-14.7	7.4	10.7	7.5-15.0	4.2	14.5	8.7-22.8	3.2	7.7	4.4-12.9
	1981-1985	11.0	16.6	12.4-21.9	6.4	23.9	16.1-34.2	4.6	12.2	7.6-18.8	8.6	12.7	9.1-17.3	4.8	18.7	11.8-28.1	3.8	9.4	5.6-15.1
	1986-1990	10.4	15.6	11.5-20.7	4.8	18.9	11.9-28.5	5.6	13.9	9.0-20.9	8.6	12.8	9.1-17.5	3.2	11.7	6.6-19.3	5.4	13.9	8.8-21.0
	1991-1995	9.4	12.9	9.4-17.4	4.0	13.8	8.3-21.7	5.4	11.9	7.7-17.8	8.6	11.8	8.5-16.1	4.2	14.6	8.9-22.6	4.4	9.4	5.8-14.8
	1996-2000	10.2	14.3	10.5-19.1	4.4	15.9	9.9-24.2	5.8	13.2	8.5-19.8	10.2	14.2	10.5-19.0	5.2	18.5	12.0-27.3	5.0	10.9	6.8-16.9
	2001-2005	9.6	14.0	10.3-18.8	5.6	19.4	12.8-28.2	4.0	10.0	5.9-16.1	5.8	9.0	6.0-13.2	2.8	9.9	5.4-16.8	3.0	8.4	4.5-14.5
	2006-2010	7.8	11.6	8.2-16.1	4.2	14.8	9.1-22.8	3.6	9.3	5.4-15.4	6.0	9.2	6.1-13.3	4.0	14.2	8.6-22.2	2.0	5.2	2.4-10.2
	2011-2015	11.6	16.5	12.4-21.6	5.8	19.7	13.1-28.7	5.8	15.1	10.0-22.3	9.0	13.1	9.5-17.8	4.8	16.7	10.6-25.2	4.2	10.7	6.5-17.0
	2016-2020	15.2	20.8	16.3-26.4	7.4	21.8	15.2-30.6	7.8	19.7	13.8-27.6	10.6	14.7	10.9-19.5	4.4	14.0	8.6-21.8	6.2	14.6	9.8-21.5
Larynx	1976-1980	3.8	5.5	3.2-8.9	3.2	10.9	6.0-18.2	0.6	1.3	0.3-4.5	1.4	1.9	0.7-4.1	1.2	3.8	1.3-8.7	0.2	0.5	0.0-3.5
	1981-1985	3.4	5.4	3.0-9.0	3.0	10.3	5.6-17.5	0.4	1.6	0.1-5.9	1.4	1.8	0.7-4.0	1.2	4.1	1.4-9.4	0.2	0.4	0.0-3.4
	1986-1990	5.0	7.6	4.8-11.5	4.2	14.2	8.7-22.2	0.8	2.7	0.7-7.2	1.4	1.8	0.7-4.0	1.4	4.7	1.8-10.0	0.0	0.0	0.0-2.8
	1991-1995	2.0	3.4	1.6-6.5	1.6	5.9	2.5-11.9	0.4	1.5	0.1-5.5	0.4	0.7	0.1-2.6	0.4	1.4	0.2-5.4	0.0	0.0	0.0-2.8
	1996-2000	2.6	4.2	2.2-7.4	2.2	8.2	4.0-14.8	0.4	1.0	0.1-4.3	0.2	0.2	0.0-1.8	0.2	0.8	0.0-4.5	0.0	0.0	0.0-2.8
	2001-2005	2.6	4.1	2.1-7.2	2.2	7.8	3.9-14.2	0.4	1.1	0.1-4.7	0.8	1.3	0.3-3.4	0.4	1.6	0.2-5.8	0.4	1.0	0.1-4.5
	2006-2010	3.2	5.2	2.9-8.6	2.0	7.5	3.6-14.0	1.2	3.0	1.0-7.3	0.8	1.2	0.3-3.2	0.4	1.4	0.2-5.5	0.4	0.8	0.1-4.1
	2011-2015	2.0	2.9	1.4-5.6	1.6	5.0	2.1-10.4	0.4	1.0	0.1-4.5	1.0	1.3	0.4-3.4	0.6	1.9	0.4-6.0	0.4	0.8	0.1-4.0
	2016-2020	1.0	1.7	0.5-4.1	0.4	1.4	0.2-5.4	0.6	1.9	0.4-5.9	0.2	0.4	0.0-2.2	0.2	0.8	0.0-4.6	0.0	0.0	0.0-3.0
Lung and bronchus	1976-1980	36.6	54.0	46.2-62.8	28.4	99.4	83.1-118.1	8.2	21.8	15.5-30.2	30.0	45.0	37.8-53.1	24.4	87.4	72.0-105.2	5.6	15.2	9.9-22.4
	1981-1985	42.6	62.1	53.7-71.6	29.6	99.1	83.2-117.3	13.0	34.9	26.4-45.4	29.0	42.0	35.2-49.8	20.6	71.1	57.6-87.0	8.4	22.0	15.5-30.6
	1986-1990	49.0	73.2	63.9-83.5	31.4	111.9	94.5-131.7	17.6	46.7	36.8-58.5	40.4	58.2	50.2-67.3	26.8	93.5	77.9-111.5	13.6	35.6	27.0-46.2
	1991-1995	54.0	80.6	71.0-91.3	34.0	123.4	105.1-144.1	20.0	51.6	41.5-63.6	41.8	61.2	52.9-70.5	27.6	99.3	83.1-117.8	14.2	36.8	28.3-47.1
	1996-2000	49.8	73.9	64.8-84.0	29.2	102.9	86.8-121.3	20.6	53.4	43.2-65.6	35.8	52.3	44.8-60.9	19.8	68.1	55.3-83.2	16.0	42.2	33.1-53.3
	2001-2005	49.0	75.1	65.8-85.4	27.6	99.6	83.6-117.9	21.4	56.7	46.1-69.4	30.6	45.5	38.5-53.6	16.6	58.6	46.6-72.9	14.0	35.9	27.6-46.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Montgomery**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	47.6	72.5	63.4-82.7	23.6	81.8	67.5-98.3	24.0	67.4	55.4-81.3	34.0	51.5	43.9-60.2	19.2	67.6	54.5-82.9	14.8	40.1	31.2-51.1
	2011-2015	53.4	79.8	70.2-90.3	28.4	93.3	78.2-110.5	25.0	70.0	57.9-84.2	31.8	45.8	38.8-53.8	16.8	54.0	42.9-67.4	15.0	39.8	31.1-50.5
	2016-2020	58.0	81.3	72.0-91.6	27.2	82.3	68.6-98.0	30.8	81.8	69.0-96.5	35.0	48.2	41.1-56.2	18.4	55.6	44.6-68.8	16.6	42.2	33.3-53.1
Melanoma of the skin	1976-1980	5.0	8.5	5.3-12.9	2.8	11.5	6.0-19.7	2.2	6.5	3.0-12.2	0.8	1.5	0.3-4.1	0.4	1.1	0.1-4.7	0.4	1.9	0.2-6.6
	1981-1985	3.8	6.8	4.0-10.9	2.2	7.0	3.4-13.0	1.6	6.9	2.8-13.6	2.2	3.6	1.7-6.6	2.0	7.8	3.6-14.6	0.2	0.4	0.0-3.4
	1986-1990	3.2	5.9	3.2-9.8	1.6	7.0	2.9-13.9	1.6	5.4	2.2-11.2	0.8	1.0	0.3-2.8	0.4	1.1	0.1-4.8	0.4	0.8	0.1-4.0
	1991-1995	3.6	6.6	3.8-10.6	2.4	10.1	5.1-17.7	1.2	3.2	1.1-7.6	0.8	1.6	0.4-4.1	0.4	1.7	0.2-6.2	0.4	1.3	0.1-4.9
	1996-2000	5.8	9.6	6.2-14.0	3.0	10.4	5.8-17.4	2.8	10.1	5.3-17.5	1.0	1.3	0.4-3.2	0.4	1.4	0.2-5.3	0.6	1.1	0.2-4.3
	2001-2005	9.0	16.2	11.7-21.9	5.2	20.1	13.1-29.7	3.8	12.9	7.5-20.8	2.0	2.9	1.3-5.5	1.2	4.1	1.5-9.2	0.8	2.2	0.5-6.3
	2006-2010	11.4	18.9	14.2-24.9	7.6	28.2	19.8-39.0	3.8	12.5	7.3-20.0	1.2	2.2	0.8-5.0	0.8	3.3	0.9-8.5	0.4	1.5	0.2-5.4
	2011-2015	11.6	19.7	14.8-25.9	5.2	18.6	11.9-27.8	6.4	21.3	14.2-30.6	2.2	3.9	1.9-7.2	1.6	6.3	2.6-12.6	0.6	2.0	0.4-6.2
2016-2020	10.6	16.8	12.3-22.4	7.8	25.1	17.5-35.0	2.8	10.2	5.3-17.7	1.0	1.3	0.4-3.3	0.6	1.6	0.3-5.4	0.4	1.1	0.1-4.7	
Female breast	1976-1980							29.6	84.8	70.9-100.7							12.8	34.5	26.2-44.8
	1981-1985							35.6	99.0	84.1-115.9							12.6	34.1	25.7-44.6
	1986-1990							36.4	111.6	94.7-130.7							10.4	27.9	20.3-37.5
	1991-1995							39.2	116.5	99.7-135.4							15.6	42.6	33.0-54.2
	1996-2000							39.8	116.0	99.5-134.4							8.2	21.2	14.9-29.5
	2001-2005							44.4	138.3	119.9-158.7							9.2	23.7	16.9-32.4
	2006-2010							37.2	111.1	95.0-129.3							8.2	20.9	14.7-29.3
	2011-2015							42.4	122.2	105.7-140.8							5.0	13.0	8.2-20.0
2016-2020							43.0	130.3	112.5-150.2							8.0	21.9	15.1-30.8	
Cervix uteri	1976-1980							5.2	20.3	12.8-30.4							1.6	4.8	1.9-10.1
	1981-1985							4.0	13.1	7.6-20.9							1.4	4.2	1.5-9.2
	1986-1990							3.0	12.2	6.6-20.4							1.4	5.2	1.9-11.1
	1991-1995							3.0	9.1	4.9-15.4							1.2	3.3	1.1-7.9
	1996-2000							1.4	5.5	2.2-11.3							0.2	0.8	0.0-4.5
	2001-2005							2.0	8.4	4.0-15.5							0.2	0.8	0.0-4.4
	2006-2010							3.0	10.0	5.5-16.9							1.0	3.7	1.1-8.8
	2011-2015							1.6	5.7	2.3-11.7							1.0	3.2	1.0-7.9
	2016-2020							1.6	5.7	2.4-11.6							0.8	2.8	0.7-7.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Montgomery**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							9.2	26.9	19.2-36.8							1.4	3.4	1.4-7.6
	1981-1985							11.4	32.5	24.1-42.9							2.6	6.0	3.2-10.9
	1986-1990							11.0	28.2	20.9-37.5							1.2	2.5	0.9-6.2
	1991-1995							10.4	27.1	19.8-36.4							2.2	5.2	2.5-10.0
	1996-2000							13.0	37.4	28.5-48.5							2.8	5.4	2.9-9.8
	2001-2005							11.0	32.8	24.4-43.3							1.6	3.9	1.6-8.4
	2006-2010							12.4	35.7	26.9-46.5							3.6	9.4	5.5-15.5
	2011-2015							11.2	30.8	22.9-40.6							2.2	5.7	2.7-10.9
	2016-2020							9.6	25.8	18.8-35.0							2.0	5.1	2.4-10.0
Ovary	1976-1980							8.4	25.9	18.2-35.7							6.0	17.4	11.5-25.4
	1981-1985							6.6	18.9	12.6-27.3							4.4	11.1	6.7-17.4
	1986-1990							5.4	14.3	9.1-21.6							3.6	9.3	5.2-15.5
	1991-1995							6.0	16.7	10.9-24.7							4.8	10.0	6.4-15.5
	1996-2000							5.2	16.9	10.8-25.2							2.2	5.4	2.6-10.4
	2001-2005							5.0	14.4	9.1-21.8							3.0	7.8	4.2-13.4
	2006-2010							3.8	9.9	5.8-16.1							3.0	7.3	3.9-12.8
	2011-2015							3.6	11.7	6.7-19.1							2.0	5.5	2.6-10.7
	2016-2020							2.4	7.3	3.6-13.3							2.0	5.2	2.5-10.2
Prostate	1976-1980				19.0	70.2	56.2-86.5							7.4	28.7	19.9-40.0			
	1981-1985				24.2	91.9	75.8-110.5							8.2	32.4	23.0-44.4			
	1986-1990				34.6	122.7	104.5-143.2							6.2	24.0	16.1-34.5			
	1991-1995				54.0	182.2	160.7-205.8							9.0	33.6	24.3-45.3			
	1996-2000				45.6	158.9	138.8-181.2							8.0	26.8	19.1-36.7			
	2001-2005				44.4	155.0	135.2-177.0							8.2	26.6	19.1-36.4			
	2006-2010				41.0	143.1	123.9-164.5							5.0	16.2	10.4-24.2			
	2011-2015				27.0	84.2	70.3-100.3							7.2	24.9	17.3-34.8			
	2016-2020				40.0	116.7	100.7-134.9							3.6	11.7	6.8-18.9			
Testis	1976-1980				0.6	2.8	0.5-8.3							0.0	0.0	0.0-3.1			
	1981-1985				0.8	3.7	0.9-9.4							0.0	0.0	0.0-3.1			
	1986-1990				0.6	2.9	0.6-8.3							0.0	0.0	0.0-3.1			
	1991-1995				1.6	7.3	3.1-14.4							0.2	0.9	0.0-4.9			
	1996-2000				2.0	9.1	4.3-16.7							0.0	0.0	0.0-3.0			
	2001-2005				2.2	9.1	4.5-16.4							0.4	1.4	0.2-5.3			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Montgomery**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				1.6	6.8	2.8-13.4							0.0	0.0	0.0-3.0			
	2011-2015				1.4	5.8	2.3-12.0							0.0	0.0	0.0-3.1			
	2016-2020				2.0	9.1	4.3-16.8							0.0	0.0	0.0-3.1			
Urinary bladder (incl. in situ)	1976-1980	14.8	23.9	18.6-30.3	8.8	34.0	24.4-46.2	6.0	17.4	11.4-25.7	5.8	8.8	5.9-12.9	3.8	14.3	8.4-22.7	2.0	5.5	2.5-10.7
	1981-1985	18.2	26.6	21.3-33.0	14.0	51.7	40.0-66.0	4.2	10.1	6.1-16.2	5.2	7.9	5.1-11.7	2.2	8.7	4.3-15.8	3.0	7.2	3.9-12.4
	1986-1990	17.0	24.5	19.5-30.6	13.4	49.2	37.8-62.9	3.6	8.0	4.7-13.3	4.4	6.2	3.9-9.6	2.6	11.5	6.0-19.8	1.8	3.8	1.7-7.9
	1991-1995	19.6	27.6	22.2-33.9	14.6	51.2	40.0-64.8	5.0	10.8	6.9-16.7	4.6	6.1	3.8-9.3	3.4	11.5	6.7-18.8	1.2	2.4	0.9-5.9
	1996-2000	22.0	32.3	26.4-39.2	17.0	60.1	47.9-74.5	5.0	11.8	7.4-18.2	5.0	6.7	4.3-10.2	2.6	9.2	4.9-15.9	2.4	5.3	2.7-10.0
	2001-2005	19.4	29.2	23.6-35.9	13.4	48.1	37.2-61.2	6.0	14.3	9.5-21.2	3.8	5.1	3.0-8.2	2.2	7.7	3.8-14.0	1.6	3.1	1.3-7.1
	2006-2010	19.4	29.2	23.5-35.9	15.6	54.3	42.7-68.1	3.8	10.6	6.1-17.3	4.0	5.7	3.4-9.1	2.8	9.6	5.2-16.4	1.2	3.5	1.2-8.1
	2011-2015	25.4	37.3	30.9-44.8	18.6	63.0	50.5-77.7	6.8	16.9	11.5-24.4	3.4	4.8	2.7-8.0	2.6	9.0	4.7-15.8	0.8	1.7	0.4-5.5
2016-2020	20.0	29.2	23.7-35.8	13.8	44.8	34.7-57.3	6.2	16.7	11.2-24.3	3.6	4.7	2.7-7.7	2.6	8.7	4.6-15.2	1.0	2.2	0.7-6.0	
Kidney and renal pelvis	1976-1980	5.8	8.5	5.6-12.4	3.6	11.9	6.9-19.4	2.2	5.4	2.7-10.1	2.6	3.6	1.9-6.3	1.4	4.2	1.6-9.2	1.2	2.9	1.1-6.9
	1981-1985	6.2	10.1	6.7-14.6	3.8	15.7	9.2-24.8	2.4	6.3	3.1-11.7	2.8	4.1	2.2-7.2	2.0	7.7	3.6-14.4	0.8	1.7	0.5-5.2
	1986-1990	7.2	11.2	7.7-15.8	4.4	16.7	10.3-25.7	2.8	7.8	4.0-13.8	3.2	5.0	2.7-8.4	1.6	6.5	2.7-13.0	1.6	4.3	1.7-9.1
	1991-1995	9.8	15.4	11.2-20.6	5.8	21.1	14.0-30.6	4.0	11.3	6.6-18.1	3.8	5.4	3.2-8.6	3.0	9.8	5.4-16.4	0.8	2.7	0.7-7.3
	1996-2000	11.6	17.5	13.1-22.8	5.8	21.1	14.1-30.5	5.8	14.5	9.5-21.6	3.2	4.2	2.4-7.1	2.0	7.0	3.4-13.2	1.2	2.2	0.8-5.7
	2001-2005	9.6	16.1	11.8-21.5	4.4	16.7	10.4-25.4	5.2	15.3	9.7-23.1	3.4	5.3	3.0-8.7	1.8	6.5	2.9-12.6	1.6	4.6	1.9-9.6
	2006-2010	11.8	18.6	14.0-24.2	7.4	24.9	17.4-34.6	4.4	12.4	7.4-19.6	3.6	5.4	3.1-8.9	1.6	6.0	2.5-12.0	2.0	4.5	1.9-9.2
	2011-2015	12.8	20.8	15.8-26.9	8.0	27.6	19.5-38.1	4.8	15.2	9.5-23.3	1.8	2.4	1.1-4.9	1.0	3.0	0.9-7.5	0.8	2.1	0.6-6.0
2016-2020	14.4	23.3	18.0-29.8	8.2	28.3	19.9-39.0	6.2	19.0	12.6-27.6	2.8	3.8	2.0-6.6	1.8	5.5	2.5-11.0	1.0	2.4	0.8-6.5	
Brain and other central nervous system	1976-1980	4.8	8.2	5.1-12.6	1.6	5.3	2.2-10.8	3.2	11.0	5.9-18.4	3.4	6.0	3.4-9.9	0.8	2.9	0.8-7.6	2.6	8.8	4.5-15.5
	1981-1985	3.6	6.4	3.7-10.3	1.2	5.0	1.8-11.1	2.4	7.3	3.6-13.1	2.0	3.1	1.4-5.9	0.6	2.0	0.4-6.2	1.4	4.1	1.6-8.9
	1986-1990	4.0	6.5	3.8-10.4	2.6	9.5	4.9-16.8	1.4	4.1	1.5-9.0	3.2	5.7	3.1-9.4	1.4	5.8	2.2-12.1	1.8	5.4	2.3-10.8
	1991-1995	4.0	5.9	3.5-9.3	2.0	7.5	3.5-14.1	2.0	4.9	2.3-9.5	3.2	4.3	2.4-7.2	2.4	8.4	4.3-14.9	0.8	1.6	0.4-4.9
	1996-2000	5.6	9.8	6.4-14.4	3.0	10.7	5.9-17.9	2.6	10.3	5.4-17.6	2.2	3.2	1.6-6.0	2.0	7.1	3.4-13.3	0.2	0.3	0.0-3.3
	2001-2005	3.6	7.1	4.2-11.3	1.8	7.5	3.4-14.2	1.8	7.0	3.1-13.4	2.4	4.4	2.3-7.9	1.0	3.7	1.2-8.9	1.4	5.3	2.1-11.0
	2006-2010	4.8	9.0	5.7-13.6	2.4	9.7	4.9-17.2	2.4	8.5	4.3-15.2	1.6	2.6	1.1-5.4	0.4	1.2	0.1-4.9	1.2	4.1	1.5-9.3
	2011-2015	3.8	6.6	3.9-10.6	1.8	6.2	2.7-12.2	2.0	7.2	3.3-13.6	1.8	3.2	1.4-6.3	1.2	4.3	1.5-9.7	0.6	2.4	0.5-7.2
2016-2020	4.6	8.7	5.4-13.3	3.0	11.8	6.5-19.8	1.6	5.7	2.3-11.6	3.8	6.4	3.7-10.2	2.2	8.7	4.2-15.9	1.6	4.1	1.7-8.8	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Montgomery**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	2.8	5.0	2.7-8.5	1.4	5.3	2.1-11.3	1.4	5.0	2.0-10.5	0.0	0.0	0.0-1.4	0.0	0.0	0.0-3.1	0.0	0.0	0.0-2.8
	1981-1985	2.0	3.8	1.7-7.0	0.8	3.4	0.9-8.6	1.2	4.2	1.4-9.5	0.2	0.3	0.0-1.8	0.2	0.6	0.0-4.1	0.0	0.0	0.0-2.8
	1986-1990	2.6	4.4	2.3-7.7	0.8	3.6	0.9-9.1	1.8	5.7	2.5-11.2	0.6	0.8	0.2-2.7	0.2	1.1	0.0-5.5	0.4	0.9	0.1-4.2
	1991-1995	2.4	5.1	2.6-8.8	0.4	2.0	0.2-6.8	2.0	7.9	3.6-14.7	0.4	0.5	0.1-2.2	0.2	1.0	0.0-5.0	0.2	0.4	0.0-3.3
	1996-2000	3.6	6.5	3.8-10.5	0.8	3.3	0.9-8.4	2.8	10.1	5.3-17.3	0.6	0.7	0.1-2.4	0.2	0.8	0.0-4.5	0.4	0.6	0.1-3.6
	2001-2005	5.6	10.9	7.2-15.9	1.6	5.7	2.4-11.4	4.0	16.3	9.9-25.1	0.2	0.2	0.0-1.8	0.2	0.7	0.0-4.3	0.0	0.0	0.0-2.9
	2006-2010	8.6	16.3	11.7-22.1	2.4	9.4	4.8-16.6	6.2	23.0	15.5-33.0	0.4	0.6	0.1-2.4	0.2	0.6	0.0-4.0	0.2	0.7	0.0-4.0
	2011-2015	8.0	14.2	10.0-19.7	2.0	7.9	3.7-14.9	6.0	20.6	13.6-30.0	0.6	1.0	0.2-3.1	0.4	1.6	0.2-6.0	0.2	0.5	0.0-3.8
2016-2020	7.2	12.7	8.7-17.8	2.0	5.4	2.5-10.5	5.2	19.8	12.7-29.4	0.4	0.5	0.1-2.3	0.2	0.6	0.0-4.0	0.2	0.4	0.0-3.6	
Hodgkin lymphoma	1976-1980	0.8	1.2	0.3-3.2	0.2	0.5	0.0-3.9	0.6	1.8	0.4-5.7	1.4	2.3	0.9-4.9	0.6	2.6	0.5-7.5	0.8	2.2	0.6-6.1
	1981-1985	2.0	4.2	2.0-7.7	1.0	3.9	1.3-9.3	1.0	4.4	1.4-10.1	0.0	0.0	0.0-1.4	0.0	0.0	0.0-3.1	0.0	0.0	0.0-2.8
	1986-1990	1.8	3.7	1.7-7.1	1.2	4.7	1.7-10.4	0.6	2.8	0.6-7.9	0.2	0.6	0.0-2.7	0.0	0.0	0.0-3.1	0.2	1.1	0.0-5.3
	1991-1995	2.2	4.4	2.1-7.9	1.2	5.3	1.9-11.5	1.0	3.4	1.0-8.3	0.6	1.1	0.2-3.3	0.2	0.8	0.0-4.5	0.4	1.2	0.1-4.9
	1996-2000	1.0	2.0	0.6-4.8	0.6	2.5	0.5-7.4	0.4	1.7	0.2-6.0	0.6	1.1	0.2-3.2	0.4	1.5	0.2-5.5	0.2	0.8	0.0-4.4
	2001-2005	1.6	3.6	1.5-7.0	1.0	4.2	1.3-9.8	0.6	2.9	0.6-8.1	0.2	0.5	0.0-2.4	0.2	0.9	0.0-4.9	0.0	0.0	0.0-2.9
	2006-2010	2.2	4.1	2.0-7.4	1.0	3.5	1.1-8.5	1.2	4.7	1.6-10.3	0.2	0.3	0.0-2.1	0.2	0.7	0.0-4.2	0.0	0.0	0.0-2.9
	2011-2015	1.2	2.5	0.9-5.4	0.8	3.5	1.0-8.9	0.4	1.4	0.1-5.5	0.4	0.7	0.1-2.6	0.2	0.9	0.0-4.9	0.2	0.5	0.0-3.7
2016-2020	1.0	1.8	0.5-4.3	0.4	1.5	0.2-5.7	0.6	2.0	0.4-6.3	0.0	0.0	0.0-1.5	0.0	0.0	0.0-3.1	0.0	0.0	0.0-3.0	
Non-Hodgkin lymphomas	1976-1980	7.2	11.3	7.8-16.0	4.4	15.4	9.4-23.8	2.8	8.8	4.5-15.5	2.4	3.6	1.8-6.4	1.2	4.4	1.6-9.9	1.2	3.1	1.1-7.3
	1981-1985	7.4	11.2	7.8-15.6	4.4	15.2	9.4-23.5	3.0	8.1	4.4-14.0	4.4	6.6	4.1-10.2	2.8	10.3	5.5-17.6	1.6	4.2	1.7-8.8
	1986-1990	7.6	12.4	8.6-17.4	4.4	17.7	10.9-27.1	3.2	8.2	4.5-14.1	3.6	5.6	3.2-9.0	1.8	6.8	3.0-13.2	1.8	4.0	1.8-8.3
	1991-1995	11.8	17.0	12.9-22.2	6.4	23.5	16.0-33.4	5.4	12.6	8.2-18.9	6.0	8.7	5.7-12.7	3.6	14.1	8.2-22.6	2.4	5.5	2.7-10.2
	1996-2000	14.2	22.9	17.7-29.1	6.8	25.2	17.4-35.4	7.4	20.5	14.0-29.1	4.2	6.4	3.9-9.9	2.6	9.5	5.0-16.4	1.6	3.9	1.6-8.4
	2001-2005	15.0	23.4	18.2-29.6	6.4	23.0	15.7-32.8	8.6	22.9	16.2-31.7	4.6	6.5	4.1-10.1	2.0	6.9	3.3-13.1	2.6	6.0	3.0-11.2
	2006-2010	13.4	20.8	16.0-26.7	5.0	17.4	11.1-26.0	8.4	23.8	16.9-32.8	4.6	6.2	3.9-9.6	2.6	8.6	4.6-15.0	2.0	4.5	2.1-8.9
	2011-2015	12.8	20.5	15.6-26.4	6.0	21.6	14.4-31.2	6.8	19.8	13.5-28.2	2.8	4.3	2.3-7.6	1.2	4.3	1.6-9.8	1.6	4.2	1.6-9.3
2016-2020	12.8	19.1	14.5-24.7	7.0	22.9	15.6-32.5	5.8	15.7	10.4-23.0	3.2	4.3	2.4-7.3	1.6	5.1	2.2-10.5	1.6	3.8	1.6-8.2	
Myeloma	1976-1980	2.4	3.7	1.9-6.6	0.8	3.2	0.8-8.4	1.6	4.1	1.8-8.5	2.2	3.2	1.6-5.9	1.2	4.3	1.5-9.8	1.0	2.5	0.8-6.3
	1981-1985	3.6	5.5	3.2-8.9	1.4	5.3	2.0-11.4	2.2	5.6	2.7-10.5	2.2	3.2	1.6-5.9	0.2	0.6	0.0-4.1	2.0	4.9	2.3-9.5
	1986-1990	4.4	6.8	4.2-10.6	2.4	9.1	4.6-16.1	2.0	5.7	2.5-11.2	3.0	4.0	2.2-6.8	1.8	6.5	2.9-12.7	1.2	2.6	1.0-6.4
	1991-1995	5.0	7.2	4.6-10.8	2.0	7.3	3.4-13.8	3.0	7.1	3.9-12.3	2.8	4.1	2.2-7.2	1.2	4.4	1.6-9.8	1.6	4.0	1.7-8.4
	1996-2000	4.2	5.9	3.6-9.3	2.4	8.5	4.4-15.1	1.8	4.7	2.0-9.6	2.6	4.1	2.2-7.2	1.0	3.8	1.2-9.0	1.6	4.4	1.8-9.2
	2001-2005	4.2	6.3	3.8-9.8	2.4	8.4	4.3-15.0	1.8	4.7	2.1-9.6	2.8	3.8	2.0-6.6	1.0	3.2	1.0-7.9	1.8	4.2	1.8-8.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Montgomery**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	4.6	6.9	4.3-10.6	2.8	9.8	5.3-16.7	1.8	4.2	1.9-8.7	2.2	3.5	1.7-6.4	1.2	4.2	1.5-9.4	1.0	2.9	0.9-7.2
	2011-2015	5.0	7.4	4.7-11.2	2.4	8.9	4.5-15.9	2.6	6.7	3.5-12.1	2.6	4.0	2.1-7.1	1.2	4.2	1.5-9.5	1.4	3.6	1.4-8.0
	2016-2020	6.6	10.1	6.8-14.4	3.8	13.4	7.9-21.3	2.8	7.0	3.7-12.5	1.4	1.8	0.7-4.1	0.6	2.3	0.5-6.9	0.8	1.4	0.4-4.8
Leukemias	1976-1980	10.4	15.8	11.7-20.9	7.0	28.3	19.5-39.8	3.4	8.0	4.7-13.3	6.8	10.1	7.0-14.3	4.2	16.7	10.1-25.9	2.6	6.3	3.4-11.3
	1981-1985	8.2	12.7	9.0-17.5	4.4	17.3	10.6-26.7	3.8	9.6	5.7-15.6	5.4	9.2	5.9-13.7	3.6	15.8	9.1-25.2	1.8	5.6	2.4-11.3
	1986-1990	5.8	8.7	5.8-12.8	3.2	12.0	6.7-19.8	2.6	6.2	3.2-11.1	5.4	7.4	4.9-11.1	2.8	9.5	5.0-16.4	2.6	5.5	2.9-10.0
	1991-1995	9.0	13.5	9.8-18.4	4.8	17.6	11.1-26.5	4.2	10.9	6.5-17.3	4.2	6.3	3.8-9.8	2.8	10.8	5.8-18.3	1.4	3.0	1.2-6.9
	1996-2000	7.6	12.1	8.5-16.8	4.8	17.3	11.0-25.9	2.8	8.6	4.5-14.9	3.6	5.1	3.0-8.3	2.2	7.5	3.7-13.7	1.4	3.5	1.3-8.0
	2001-2005	8.6	13.5	9.7-18.5	4.6	17.2	10.9-26.0	4.0	9.7	5.8-15.7	3.8	6.0	3.6-9.6	2.2	7.9	3.9-14.3	1.6	4.4	1.7-9.3
	2006-2010	12.2	19.3	14.6-25.2	7.0	25.8	17.7-36.2	5.2	14.0	8.8-21.4	4.6	6.3	3.9-9.8	2.2	7.5	3.7-13.9	2.4	5.1	2.5-9.7
	2011-2015	16.0	25.0	19.6-31.5	10.4	37.6	27.8-49.8	5.6	13.5	8.7-20.3	4.2	5.8	3.5-9.2	1.6	6.1	2.5-12.2	2.6	5.6	2.9-10.4
2016-2020	12.0	18.0	13.6-23.5	7.8	24.6	17.3-34.2	4.2	12.2	7.3-19.5	4.8	7.9	5.0-12.0	3.4	12.2	7.0-19.8	1.4	4.2	1.6-9.3	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Nassau**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	5,327.4	431.9	426.5-437.3	2,539.0	509.1	499.2-519.0	2,788.4	395.5	388.8-402.2	2,793.2	232.9	228.9-236.9	1,414.6	295.1	287.4-302.8	1,378.6	197.8	193.1-202.6
	1981-1985	5,871.8	441.9	436.7-447.2	2,836.6	515.7	506.3-525.1	3,035.2	405.6	399.0-412.2	2,924.0	225.6	221.9-229.5	1,462.2	282.4	275.3-289.7	1,461.8	194.7	190.2-199.3
	1986-1990	6,421.4	464.2	459.0-469.5	3,062.0	524.6	515.7-533.7	3,359.4	435.5	428.8-442.3	3,013.0	221.3	217.7-225.0	1,515.4	273.6	267.0-280.4	1,497.6	191.2	186.8-195.6
	1991-1995	7,159.4	495.8	490.6-501.0	3,586.0	578.1	569.3-587.0	3,573.4	447.5	440.9-454.2	2,959.0	207.3	203.9-210.7	1,443.6	248.8	242.8-255.0	1,515.4	184.8	180.6-189.0
	1996-2000	7,687.8	515.2	510.0-520.4	3,844.0	593.1	584.7-601.7	3,843.8	465.4	458.8-472.1	2,842.4	189.1	186.0-192.2	1,394.2	224.2	218.8-229.6	1,448.2	167.5	163.6-171.4
	2001-2005	8,039.4	517.7	512.6-522.8	3,999.0	591.5	583.2-599.8	4,040.4	469.6	463.1-476.2	2,728.0	171.5	168.6-174.4	1,319.4	200.5	195.7-205.5	1,408.6	153.6	150.0-157.3
	2006-2010	8,448.0	526.8	521.7-532.0	4,238.4	598.3	590.2-606.5	4,209.6	478.3	471.7-485.0	2,538.2	151.6	149.0-154.3	1,226.4	175.5	171.1-180.0	1,311.8	136.6	133.2-140.0
	2011-2015	8,580.0	504.3	499.4-509.2	4,205.8	548.3	540.7-555.9	4,374.2	477.2	470.6-483.8	2,496.2	139.2	136.7-141.8	1,197.2	157.9	153.8-162.0	1,299.0	126.7	123.5-130.0
	2016-2020	8,707.4	486.7	482.0-491.5	4,302.6	520.6	513.5-527.8	4,404.8	466.7	460.2-473.2	2,332.8	122.6	120.3-124.9	1,131.2	139.5	135.8-143.3	1,201.6	111.3	108.4-114.3
Oral cavity and pharynx	1976-1980	150.4	11.3	10.5-12.2	96.8	16.8	15.2-18.6	53.6	7.3	6.4-8.2	58.4	4.5	4.0-5.1	37.4	6.7	5.7-7.8	21.0	2.9	2.4-3.5
	1981-1985	146.6	10.5	9.7-11.3	95.8	15.8	14.3-17.4	50.8	6.5	5.7-7.3	58.0	4.2	3.7-4.7	39.2	6.5	5.5-7.5	18.8	2.4	1.9-3.0
	1986-1990	147.6	10.5	9.7-11.3	92.8	15.2	13.7-16.7	54.8	7.0	6.2-7.9	54.2	3.9	3.4-4.4	34.8	6.0	5.0-7.0	19.4	2.4	2.0-3.0
	1991-1995	132.8	9.2	8.5-10.0	81.2	12.9	11.7-14.3	51.6	6.5	5.7-7.4	41.6	2.9	2.5-3.3	27.2	4.3	3.6-5.2	14.4	1.8	1.4-2.2
	1996-2000	132.2	8.9	8.3-9.6	83.6	12.8	11.6-14.1	48.6	5.9	5.2-6.7	29.8	2.0	1.7-2.4	19.8	3.0	2.4-3.7	10.0	1.1	0.9-1.5
	2001-2005	138.8	9.0	8.4-9.8	86.6	12.6	11.4-13.9	52.2	6.1	5.4-6.9	29.6	1.9	1.6-2.2	19.2	2.8	2.3-3.5	10.4	1.1	0.8-1.5
	2006-2010	154.2	9.5	8.8-10.2	97.0	13.3	12.1-14.5	57.2	6.5	5.8-7.4	30.4	1.8	1.5-2.1	20.0	2.8	2.3-3.4	10.4	1.1	0.8-1.4
	2011-2015	175.0	10.1	9.4-10.8	111.8	14.2	13.0-15.4	63.2	6.7	6.0-7.5	26.0	1.4	1.2-1.7	14.6	1.8	1.4-2.3	11.4	1.1	0.8-1.4
	2016-2020	195.4	10.9	10.2-11.7	128.8	15.7	14.4-17.0	66.6	7.0	6.2-7.8	30.6	1.6	1.4-1.9	16.8	2.1	1.6-2.6	13.8	1.2	0.9-1.5
Esophagus	1976-1980	62.2	5.0	4.5-5.6	46.0	8.8	7.6-10.1	16.2	2.4	1.9-3.0	53.2	4.3	3.8-4.9	39.8	7.9	6.7-9.2	13.4	1.9	1.5-2.5
	1981-1985	56.2	4.1	3.6-4.6	41.0	6.9	5.9-8.0	15.2	2.0	1.6-2.5	51.0	3.8	3.4-4.4	36.8	6.6	5.6-7.7	14.2	1.9	1.5-2.4
	1986-1990	50.0	3.4	3.0-3.9	32.4	5.0	4.2-5.9	17.6	2.2	1.8-2.7	50.2	3.5	3.1-4.0	36.0	5.6	4.8-6.6	14.2	1.8	1.4-2.3
	1991-1995	63.0	4.3	3.8-4.8	46.0	7.2	6.3-8.2	17.0	2.0	1.6-2.5	54.0	3.7	3.2-4.1	39.0	6.0	5.2-7.0	15.0	1.7	1.4-2.2
	1996-2000	69.8	4.6	4.2-5.2	47.2	7.2	6.3-8.3	22.6	2.6	2.1-3.1	60.8	4.0	3.6-4.5	43.6	6.8	5.9-7.7	17.2	2.0	1.6-2.5
	2001-2005	68.6	4.3	3.9-4.8	50.4	7.4	6.5-8.4	18.2	1.9	1.5-2.3	60.2	3.8	3.4-4.2	44.4	6.5	5.7-7.4	15.8	1.6	1.3-2.1
	2006-2010	67.8	4.1	3.7-4.6	52.0	7.4	6.5-8.4	15.8	1.6	1.3-2.1	53.6	3.2	2.8-3.6	40.6	5.8	5.0-6.6	13.0	1.3	1.0-1.6
	2011-2015	75.6	4.3	3.9-4.8	55.2	7.1	6.3-8.0	20.4	2.0	1.7-2.5	54.2	3.0	2.7-3.4	40.6	5.3	4.6-6.1	13.6	1.2	1.0-1.6
	2016-2020	71.0	3.8	3.4-4.2	51.8	6.3	5.5-7.1	19.2	1.7	1.4-2.1	53.6	2.9	2.5-3.2	42.0	5.2	4.5-5.9	11.6	1.0	0.8-1.3
Stomach	1976-1980	139.4	12.1	11.2-13.1	74.2	15.8	14.1-17.7	65.2	9.7	8.7-10.8	105.6	9.5	8.7-10.4	54.4	12.2	10.6-13.9	51.2	7.7	6.8-8.7
	1981-1985	147.2	11.4	10.6-12.3	87.2	16.0	14.4-17.7	60.0	8.2	7.3-9.2	109.4	8.8	8.0-9.6	64.0	12.4	11.0-14.0	45.4	6.2	5.4-7.1
	1986-1990	141.8	10.6	9.8-11.4	85.8	14.8	13.4-16.4	56.0	7.4	6.5-8.3	95.4	7.2	6.6-7.9	55.0	9.8	8.6-11.2	40.4	5.3	4.6-6.1
	1991-1995	159.2	11.2	10.5-12.0	95.8	16.3	14.8-17.9	63.4	7.7	6.9-8.7	100.8	7.2	6.6-7.9	60.0	10.6	9.4-12.0	40.8	5.0	4.3-5.7
	1996-2000	149.8	10.0	9.3-10.8	89.4	14.3	13.0-15.8	60.4	6.9	6.2-7.8	78.8	5.3	4.8-5.8	43.2	7.0	6.1-8.0	35.6	4.1	3.5-4.8
	2001-2005	135.6	8.6	8.0-9.3	80.2	12.1	10.9-13.3	55.4	6.2	5.5-7.0	69.0	4.4	3.9-4.8	43.6	6.7	5.9-7.7	25.4	2.8	2.3-3.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Nassau**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	121.2	7.5	6.9-8.1	71.8	10.2	9.2-11.4	49.4	5.3	4.6-6.0	55.6	3.3	2.9-3.7	28.0	3.9	3.3-4.7	27.6	2.8	2.4-3.4
	2011-2015	143.6	8.5	7.9-9.1	86.2	11.5	10.4-12.6	57.4	6.1	5.4-6.9	58.6	3.4	3.0-3.8	33.4	4.4	3.8-5.2	25.2	2.5	2.1-3.1
	2016-2020	150.4	8.3	7.7-8.9	86.6	10.6	9.6-11.7	63.8	6.4	5.7-7.2	48.8	2.6	2.3-3.0	26.2	3.2	2.7-3.9	22.6	2.1	1.7-2.6
Colorectal	1976-1980	875.8	74.2	71.9-76.5	432.4	89.4	85.2-93.7	443.4	64.6	61.9-67.4	420.4	37.4	35.8-39.1	206.2	46.0	42.9-49.2	214.2	32.3	30.3-34.3
	1981-1985	953.0	73.9	71.8-76.1	494.4	92.5	88.5-96.6	458.6	61.5	59.0-64.2	447.0	36.3	34.8-37.9	217.4	44.8	41.9-47.9	229.6	31.4	29.6-33.3
	1986-1990	973.8	71.6	69.6-73.7	514.2	90.6	86.9-94.5	459.6	58.5	56.1-61.0	419.8	31.9	30.6-33.4	222.2	42.7	40.0-45.5	197.6	25.4	23.8-27.0
	1991-1995	924.2	64.3	62.4-66.2	462.6	76.7	73.5-80.1	461.6	55.6	53.3-57.9	371.4	26.4	25.2-27.6	189.6	33.5	31.3-35.9	181.8	21.8	20.4-23.3
	1996-2000	934.6	62.0	60.3-63.8	479.4	75.5	72.5-78.7	455.2	52.2	50.0-54.4	307.6	20.5	19.4-21.5	158.4	25.6	23.8-27.5	149.2	16.7	15.5-18.0
	2001-2005	885.4	55.8	54.1-57.4	436.2	64.8	62.1-67.6	449.2	48.8	46.8-50.9	286.2	17.8	16.9-18.8	139.0	21.1	19.6-22.8	147.2	15.4	14.3-16.6
	2006-2010	718.6	43.5	42.0-44.9	353.4	50.3	47.9-52.7	365.2	38.1	36.4-40.0	241.2	14.3	13.5-15.2	119.4	17.2	15.8-18.6	121.8	12.2	11.2-13.2
	2011-2015	676.8	38.7	37.4-40.1	335.8	44.2	42.0-46.4	341.0	34.3	32.7-36.1	222.4	12.1	11.4-12.8	111.2	14.5	13.3-15.8	111.2	10.1	9.2-11.0
	2016-2020	660.4	37.0	35.7-38.3	328.6	40.9	38.9-43.0	331.8	33.8	32.1-35.5	203.0	10.7	10.1-11.4	99.8	12.5	11.4-13.7	103.2	9.3	8.5-10.2
Colon excluding rectum	1976-1980	594.0	51.2	49.3-53.1	287.8	61.4	57.9-65.0	306.2	45.0	42.8-47.4	350.0	31.1	29.6-32.7	168.8	37.5	34.7-40.4	181.2	27.3	25.5-29.1
	1981-1985	664.4	52.1	50.3-54.0	334.4	63.9	60.5-67.3	330.0	44.6	42.4-46.8	384.4	31.2	29.7-32.6	185.2	38.0	35.3-40.8	199.2	27.2	25.5-28.9
	1986-1990	677.4	50.5	48.8-52.3	338.0	61.3	58.1-64.5	339.4	43.3	41.3-45.5	367.2	27.9	26.6-29.3	191.8	37.1	34.5-39.7	175.4	22.5	21.0-24.1
	1991-1995	658.0	45.9	44.3-47.5	319.8	53.5	50.7-56.3	338.2	40.7	38.7-42.7	318.6	22.6	21.5-23.8	160.8	28.7	26.6-30.8	157.8	18.9	17.6-20.3
	1996-2000	669.0	44.4	42.9-46.0	332.0	52.8	50.3-55.5	337.0	38.5	36.7-40.4	264.4	17.6	16.6-18.6	136.2	22.0	20.4-23.8	128.2	14.3	13.2-15.5
	2001-2005	631.6	39.6	38.2-41.0	299.2	44.7	42.5-47.1	332.4	35.9	34.1-37.6	240.6	15.0	14.1-15.9	113.6	17.3	15.9-18.8	127.0	13.2	12.2-14.3
	2006-2010	512.0	30.8	29.6-32.0	241.8	34.7	32.7-36.7	270.2	27.7	26.2-29.3	205.0	12.1	11.4-12.9	100.6	14.5	13.2-15.8	104.4	10.4	9.5-11.4
	2011-2015	481.6	27.3	26.2-28.4	230.0	30.4	28.6-32.2	251.6	24.8	23.4-26.2	189.0	10.2	9.6-10.9	90.0	11.8	10.7-12.9	99.0	8.9	8.1-9.8
	2016-2020	480.8	26.7	25.6-27.8	229.2	28.5	26.8-30.3	251.6	25.3	23.8-26.8	167.0	8.8	8.2-9.4	77.0	9.7	8.7-10.7	90.0	8.0	7.3-8.9
Rectum & rectosigmoid	1976-1980	281.8	23.0	21.8-24.3	144.6	28.0	25.8-30.3	137.2	19.6	18.1-21.2	70.4	6.3	5.6-7.0	37.4	8.5	7.2-9.9	33.0	5.0	4.3-5.8
	1981-1985	288.6	21.8	20.7-23.0	160.0	28.6	26.5-30.9	128.6	17.0	15.7-18.4	62.6	5.2	4.6-5.8	32.2	6.8	5.7-8.1	30.4	4.2	3.5-4.9
	1986-1990	296.4	21.1	20.0-22.2	176.2	29.4	27.3-31.5	120.2	15.2	14.0-16.5	52.6	4.0	3.5-4.5	30.4	5.7	4.7-6.7	22.2	2.9	2.3-3.5
	1991-1995	266.2	18.4	17.4-19.4	142.8	23.2	21.5-25.1	123.4	14.9	13.8-16.2	52.8	3.7	3.3-4.2	28.8	4.9	4.1-5.8	24.0	2.9	2.4-3.5
	1996-2000	265.6	17.6	16.7-18.6	147.4	22.7	21.1-24.4	118.2	13.7	12.6-14.8	43.2	2.9	2.5-3.3	22.2	3.5	2.9-4.3	21.0	2.4	2.0-2.9
	2001-2005	253.8	16.2	15.3-17.1	137.0	20.1	18.6-21.7	116.8	13.0	11.9-14.1	45.6	2.9	2.5-3.3	25.4	3.8	3.1-4.5	20.2	2.2	1.8-2.6
	2006-2010	206.6	12.7	11.9-13.5	111.6	15.6	14.3-17.0	95.0	10.4	9.5-11.4	36.2	2.2	1.9-2.5	18.8	2.7	2.2-3.3	17.4	1.7	1.4-2.2
	2011-2015	195.2	11.4	10.7-12.2	105.8	13.8	12.6-15.1	89.4	9.6	8.7-10.5	33.4	1.8	1.6-2.2	21.2	2.7	2.2-3.3	12.2	1.2	0.9-1.5
	2016-2020	179.6	10.3	9.6-11.1	99.4	12.4	11.3-13.6	80.2	8.5	7.6-9.4	36.0	2.0	1.7-2.3	22.8	2.8	2.3-3.4	13.2	1.3	1.0-1.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Nassau**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	36.6	3.1	2.7-3.6	23.0	4.8	3.9-5.8	13.6	2.0	1.6-2.6	34.0	2.9	2.4-3.4	19.0	3.9	3.1-4.8	15.0	2.2	1.7-2.8
	1981-1985	33.8	2.6	2.2-3.1	23.0	4.5	3.6-5.5	10.8	1.4	1.1-1.9	33.0	2.6	2.2-3.0	20.8	4.0	3.2-5.0	12.2	1.6	1.2-2.1
	1986-1990	48.6	3.6	3.1-4.1	30.0	5.2	4.4-6.2	18.6	2.4	1.9-2.9	42.8	3.2	2.8-3.7	26.6	4.9	4.1-5.9	16.2	2.1	1.7-2.7
	1991-1995	52.4	3.7	3.2-4.1	32.8	5.3	4.5-6.3	19.6	2.4	1.9-2.9	46.6	3.3	2.9-3.7	25.6	4.3	3.6-5.2	21.0	2.6	2.1-3.2
	1996-2000	65.4	4.3	3.9-4.8	43.8	6.8	5.9-7.8	21.6	2.5	2.0-3.0	52.2	3.5	3.1-3.9	32.8	5.0	4.3-5.9	19.4	2.2	1.8-2.7
	2001-2005	86.4	5.5	5.0-6.1	60.8	8.8	7.8-9.9	25.6	2.9	2.4-3.4	63.4	4.0	3.6-4.5	42.8	6.3	5.5-7.2	20.6	2.2	1.8-2.7
	2006-2010	107.8	6.6	6.0-7.1	77.8	10.7	9.6-11.8	30.0	3.2	2.7-3.8	80.0	4.7	4.3-5.2	53.4	7.5	6.6-8.4	26.6	2.7	2.2-3.2
	2011-2015	122.6	6.9	6.3-7.4	89.0	10.9	9.9-12.0	33.6	3.5	3.0-4.1	84.4	4.7	4.2-5.2	56.4	7.0	6.2-7.9	28.0	2.8	2.3-3.3
2016-2020	121.4	6.4	5.9-7.0	82.8	9.5	8.6-10.4	38.6	3.8	3.3-4.4	84.2	4.4	4.0-4.9	54.8	6.5	5.7-7.3	29.4	2.8	2.3-3.3	
Pancreas	1976-1980	174.2	14.8	13.8-15.9	93.4	19.9	17.9-22.0	80.8	11.8	10.7-13.0	163.6	13.9	12.9-15.0	86.0	18.4	16.5-20.5	77.6	11.3	10.2-12.5
	1981-1985	158.8	12.1	11.3-13.1	75.6	13.9	12.4-15.6	83.2	11.0	10.0-12.2	153.6	11.8	11.0-12.7	77.0	14.4	12.9-16.1	76.6	10.2	9.2-11.3
	1986-1990	187.2	13.6	12.7-14.5	91.6	15.2	13.7-16.8	95.6	12.2	11.1-13.3	167.6	12.3	11.4-13.2	83.4	14.2	12.7-15.7	84.2	10.7	9.7-11.8
	1991-1995	182.4	12.7	11.9-13.5	87.8	14.4	13.0-15.9	94.6	11.3	10.3-12.4	165.2	11.6	10.8-12.4	80.0	13.3	12.0-14.8	85.2	10.2	9.2-11.2
	1996-2000	212.4	14.1	13.3-15.0	102.6	16.1	14.7-17.6	109.8	12.5	11.5-13.6	183.6	12.2	11.4-13.0	90.0	14.4	13.0-15.8	93.6	10.6	9.7-11.6
	2001-2005	223.8	14.0	13.2-14.9	106.8	15.7	14.4-17.1	117.0	12.6	11.6-13.7	188.8	11.8	11.1-12.6	92.8	13.8	12.6-15.1	96.0	10.2	9.3-11.2
	2006-2010	243.6	14.6	13.8-15.5	123.4	17.5	16.2-19.0	120.2	12.3	11.3-13.4	192.4	11.4	10.7-12.1	96.8	13.8	12.6-15.1	95.6	9.6	8.8-10.6
	2011-2015	273.2	15.4	14.6-16.2	136.6	17.7	16.4-19.1	136.6	13.4	12.4-14.5	211.8	11.9	11.2-12.7	105.4	13.9	12.7-15.1	106.4	10.2	9.3-11.2
2016-2020	299.4	15.9	15.1-16.7	152.8	18.6	17.2-20.0	146.6	13.8	12.8-14.9	221.2	11.5	10.8-12.2	111.8	13.5	12.4-14.7	109.4	9.9	9.1-10.8	
Larynx	1976-1980	73.2	5.3	4.7-5.9	59.6	10.2	9.0-11.6	13.6	1.7	1.3-2.2	24.2	1.9	1.5-2.3	19.2	3.7	2.9-4.6	5.0	0.7	0.4-1.0
	1981-1985	79.4	5.4	4.8-5.9	62.4	9.7	8.6-10.9	17.0	2.1	1.6-2.6	22.4	1.6	1.3-1.9	17.8	3.0	2.3-3.7	4.6	0.6	0.4-0.9
	1986-1990	70.2	4.7	4.2-5.2	54.0	8.3	7.3-9.5	16.2	1.9	1.5-2.4	16.4	1.1	0.9-1.4	13.0	2.2	1.7-2.9	3.4	0.4	0.2-0.7
	1991-1995	70.2	4.7	4.2-5.3	53.6	8.2	7.2-9.3	16.6	2.0	1.6-2.5	17.4	1.2	0.9-1.4	11.8	1.9	1.4-2.4	5.6	0.7	0.4-1.0
	1996-2000	54.4	3.6	3.2-4.1	42.8	6.6	5.7-7.5	11.6	1.4	1.0-1.8	15.2	1.0	0.8-1.3	10.6	1.6	1.2-2.1	4.6	0.5	0.3-0.8
	2001-2005	52.8	3.4	3.0-3.8	41.0	6.0	5.2-6.9	11.8	1.4	1.1-1.8	16.4	1.0	0.8-1.3	12.8	1.9	1.5-2.5	3.6	0.4	0.2-0.7
	2006-2010	50.4	3.1	2.7-3.5	39.8	5.5	4.8-6.4	10.6	1.2	0.9-1.6	12.2	0.7	0.6-0.9	10.0	1.4	1.0-1.8	2.2	0.3	0.1-0.5
	2011-2015	43.4	2.4	2.1-2.8	34.2	4.3	3.6-5.0	9.2	1.0	0.7-1.3	13.4	0.7	0.6-0.9	8.6	1.1	0.8-1.5	4.8	0.5	0.3-0.7
2016-2020	45.4	2.4	2.1-2.8	36.4	4.3	3.7-5.0	9.0	0.9	0.6-1.2	11.2	0.6	0.4-0.8	10.0	1.2	0.9-1.6	1.2	0.1	0.0-0.3	
Lung and bronchus	1976-1980	745.8	58.2	56.3-60.2	501.8	96.2	92.1-100.5	244.0	33.3	31.4-35.3	608.8	48.0	46.3-49.8	420.8	82.2	78.4-86.2	188.0	25.8	24.2-27.6
	1981-1985	833.6	60.8	58.9-62.7	529.6	92.4	88.6-96.4	304.0	39.6	37.6-41.7	670.6	49.6	47.9-51.4	437.0	79.3	75.7-83.0	233.6	30.4	28.6-32.2
	1986-1990	912.8	64.3	62.4-66.2	529.0	89.0	85.4-92.8	383.8	48.2	46.0-50.4	727.4	51.6	49.9-53.3	433.2	73.9	70.6-77.4	294.2	36.9	35.0-38.9
	1991-1995	966.6	65.9	64.0-67.8	522.6	84.0	80.7-87.4	444.0	53.6	51.4-55.9	753.4	51.5	49.8-53.1	422.4	69.2	66.2-72.4	331.0	39.7	37.7-41.6
	1996-2000	1,000.4	65.8	64.0-67.7	515.0	79.5	76.4-82.6	485.4	56.4	54.2-58.7	718.0	47.2	45.6-48.7	382.6	59.8	57.1-62.6	335.4	38.5	36.6-40.4
	2001-2005	1,013.4	63.9	62.2-65.7	508.6	75.4	72.5-78.4	504.8	56.1	53.9-58.4	700.6	43.9	42.4-45.4	356.8	53.6	51.1-56.2	343.8	37.5	35.7-39.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Nassau**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	990.0	60.6	58.9-62.3	458.4	65.5	62.8-68.3	531.6	57.5	55.3-59.8	608.8	36.8	35.4-38.1	298.4	42.6	40.5-44.9	310.4	32.7	31.1-34.4
	2011-2015	982.4	56.5	54.9-58.2	450.0	60.0	57.5-62.6	532.4	54.5	52.4-56.7	573.0	32.3	31.1-33.5	276.2	36.5	34.6-38.6	296.8	29.4	27.9-31.0
	2016-2020	950.8	50.7	49.3-52.2	414.8	50.7	48.5-53.0	536.0	51.4	49.4-53.5	451.6	23.5	22.5-24.5	217.0	26.5	24.9-28.1	234.6	21.4	20.2-22.7
Melanoma of the skin	1976-1980	128.8	9.9	9.1-10.7	66.6	11.4	10.2-12.9	62.2	8.8	7.8-9.9	42.0	3.3	2.9-3.8	25.0	4.3	3.5-5.1	17.0	2.5	2.0-3.1
	1981-1985	164.6	12.1	11.2-12.9	86.8	14.5	13.1-16.1	77.8	10.5	9.5-11.6	41.4	3.1	2.7-3.6	23.8	4.3	3.5-5.2	17.6	2.4	1.9-2.9
	1986-1990	167.6	12.0	11.2-12.9	91.8	14.8	13.4-16.3	75.8	10.1	9.1-11.2	45.2	3.3	2.9-3.8	28.2	4.8	4.0-5.8	17.0	2.2	1.8-2.8
	1991-1995	154.6	10.9	10.1-11.7	89.2	14.2	12.9-15.7	65.4	8.5	7.6-9.5	47.4	3.4	3.0-3.8	27.8	4.7	3.9-5.6	19.6	2.5	2.0-3.1
	1996-2000	187.6	12.8	12.0-13.7	114.8	17.5	16.1-19.0	72.8	9.2	8.3-10.2	45.2	3.0	2.7-3.5	28.2	4.3	3.6-5.1	17.0	2.1	1.6-2.6
	2001-2005	316.4	20.8	19.7-21.8	180.2	26.6	24.9-28.5	136.2	16.5	15.3-17.9	46.4	3.0	2.6-3.4	27.8	4.2	3.5-4.9	18.6	2.1	1.7-2.6
	2006-2010	402.8	25.7	24.6-26.9	237.8	33.8	31.9-35.9	165.0	19.9	18.6-21.4	43.4	2.6	2.3-3.0	26.2	3.8	3.1-4.5	17.2	1.9	1.5-2.4
	2011-2015	390.2	23.1	22.1-24.2	223.2	29.5	27.8-31.4	167.0	18.6	17.3-20.0	43.6	2.5	2.1-2.8	28.0	3.7	3.1-4.4	15.6	1.6	1.2-2.0
2016-2020	389.6	21.9	20.9-22.9	234.2	28.9	27.2-30.7	155.4	16.8	15.6-18.1	32.8	1.7	1.5-2.1	21.0	2.7	2.2-3.2	11.8	1.1	0.8-1.4	
Female breast	1976-1980							822.8	115.8	112.2-119.5							302.6	42.0	39.9-44.2
	1981-1985							919.0	123.9	120.2-127.6							321.2	42.2	40.1-44.4
	1986-1990							1,040.6	136.7	132.9-140.6							323.6	41.3	39.2-43.4
	1991-1995							1,116.0	143.1	139.3-147.0							295.2	36.9	35.0-38.9
	1996-2000							1,204.0	149.4	145.6-153.2							271.4	32.2	30.5-34.0
	2001-2005							1,205.0	143.6	140.0-147.4							231.0	26.1	24.6-27.7
	2006-2010							1,260.0	147.0	143.3-150.8							203.2	21.7	20.4-23.1
	2011-2015							1,300.8	147.1	143.4-150.9							191.0	19.3	18.0-20.6
2016-2020							1,324.8	145.9	142.2-149.6							178.8	17.3	16.1-18.5	
Cervix uteri	1976-1980							61.2	8.7	7.7-9.7							20.6	2.9	2.4-3.5
	1981-1985							51.6	7.3	6.4-8.3							20.4	2.7	2.2-3.3
	1986-1990							63.2	9.1	8.1-10.1							18.6	2.5	2.0-3.1
	1991-1995							60.4	8.2	7.3-9.2							20.0	2.7	2.2-3.3
	1996-2000							66.6	8.8	7.9-9.8							19.4	2.4	2.0-3.0
	2001-2005							54.6	7.1	6.2-8.0							14.0	1.6	1.2-2.0
	2006-2010							52.2	7.0	6.2-8.0							13.8	1.7	1.3-2.1
	2011-2015							47.8	6.1	5.3-7.0							13.2	1.5	1.1-1.9
	2016-2020							52.0	6.6	5.8-7.5							13.8	1.5	1.1-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Nassau**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							200.6	27.0	25.3-28.7							46.0	6.6	5.8-7.5
	1981-1985							178.0	22.5	21.0-24.1							40.0	5.3	4.6-6.1
	1986-1990							194.4	24.2	22.7-25.8							38.8	4.8	4.2-5.6
	1991-1995							224.8	27.8	26.2-29.5							39.4	4.7	4.0-5.4
	1996-2000							242.4	29.5	27.8-31.2							38.6	4.4	3.8-5.0
	2001-2005							268.2	31.3	29.6-33.0							45.4	5.0	4.4-5.7
	2006-2010							289.4	32.6	30.9-34.3							45.2	4.8	4.2-5.5
	2011-2015							315.0	33.3	31.6-35.0							53.4	5.2	4.6-5.9
	2016-2020							321.4	32.0	30.4-33.7							62.4	6.0	5.3-6.7
Ovary	1976-1980							129.0	17.8	16.4-19.3							88.8	12.2	11.0-13.4
	1981-1985							137.2	18.1	16.7-19.5							84.6	11.0	10.0-12.1
	1986-1990							144.6	18.8	17.4-20.2							88.6	11.1	10.1-12.2
	1991-1995							142.8	18.1	16.8-19.5							86.8	10.7	9.7-11.8
	1996-2000							150.6	18.5	17.2-19.9							85.0	9.9	9.0-10.9
	2001-2005							139.4	16.5	15.3-17.8							87.4	9.7	8.8-10.6
	2006-2010							130.6	14.9	13.7-16.1							83.0	8.9	8.0-9.8
	2011-2015							112.4	12.3	11.3-13.4							74.6	7.6	6.9-8.5
	2016-2020							110.4	12.1	11.0-13.2							61.2	5.8	5.2-6.6
Prostate	1976-1980				337.4	81.9	77.7-86.3							118.4	33.1	30.3-36.1			
	1981-1985				428.0	90.6	86.4-94.9							135.6	33.3	30.6-36.1			
	1986-1990				545.6	99.9	95.8-104.1							163.6	35.5	32.9-38.2			
	1991-1995				1,043.6	164.0	159.3-168.7							165.4	33.4	31.0-35.9			
	1996-2000				1,173.4	176.7	172.1-181.3							156.0	27.6	25.6-29.7			
	2001-2005				1,178.8	171.9	167.5-176.3							141.2	22.7	21.0-24.5			
	2006-2010				1,315.8	181.2	176.8-185.7							126.0	18.3	16.9-19.8			
	2011-2015				1,123.8	138.7	135.1-142.5							105.8	14.2	13.0-15.5			
	2016-2020				1,258.4	141.7	138.2-145.4							110.8	13.7	12.6-14.9			
Testis	1976-1980				28.0	4.6	3.9-5.5							5.6	0.8	0.5-1.2			
	1981-1985				37.6	5.7	4.9-6.6							4.0	0.6	0.4-1.0			
	1986-1990				34.0	5.1	4.3-5.9							2.2	0.3	0.2-0.6			
	1991-1995				37.0	5.6	4.8-6.5							2.0	0.3	0.1-0.5			
	1996-2000				39.0	6.1	5.3-7.0							1.8	0.3	0.1-0.5			
	2001-2005				35.0	5.7	4.9-6.6							1.4	0.2	0.1-0.5			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Nassau**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				39.0	6.5	5.6-7.4							1.4	0.2	0.1-0.4			
	2011-2015				38.6	6.3	5.5-7.3							1.2	0.2	0.1-0.4			
	2016-2020				35.2	5.8	5.0-6.8							1.6	0.2	0.1-0.5			
Urinary bladder (incl. in situ)	1976-1980	280.0	23.3	22.1-24.6	206.2	43.1	40.2-46.1	73.8	10.7	9.6-11.8	74.8	7.0	6.3-7.8	53.4	13.3	11.7-15.2	21.4	3.3	2.7-4.0
	1981-1985	315.8	23.7	22.5-24.9	236.4	43.6	40.9-46.4	79.4	10.5	9.5-11.6	75.0	6.2	5.6-6.9	55.6	12.4	10.8-14.1	19.4	2.7	2.2-3.3
	1986-1990	348.6	25.1	23.9-26.4	255.2	43.9	41.3-46.6	93.4	11.9	10.8-13.0	71.0	5.6	5.0-6.2	49.4	10.4	9.0-11.9	21.6	2.8	2.3-3.4
	1991-1995	354.4	24.4	23.3-25.6	257.0	42.9	40.5-45.5	97.4	11.8	10.8-12.9	69.4	5.0	4.5-5.6	48.0	9.3	8.1-10.6	21.4	2.6	2.1-3.1
	1996-2000	410.6	27.1	25.9-28.3	305.8	48.2	45.8-50.7	104.8	12.1	11.0-13.1	67.2	4.4	4.0-4.9	47.2	8.2	7.1-9.3	20.0	2.2	1.8-2.7
	2001-2005	423.6	26.5	25.4-27.7	311.8	46.4	44.1-48.8	111.8	12.3	11.2-13.3	68.0	4.2	3.7-4.6	46.2	7.2	6.3-8.2	21.8	2.3	1.9-2.8
	2006-2010	447.8	26.9	25.8-28.1	328.0	46.7	44.4-49.0	119.8	12.6	11.5-13.6	72.6	4.2	3.8-4.6	51.4	7.5	6.6-8.5	21.2	2.0	1.6-2.4
	2011-2015	462.0	26.3	25.2-27.4	344.8	46.0	43.8-48.3	117.2	11.8	10.8-12.8	82.8	4.4	4.0-4.8	55.4	7.3	6.5-8.2	27.4	2.4	2.0-2.8
2016-2020	441.8	23.3	22.3-24.3	332.4	40.6	38.6-42.7	109.4	10.1	9.3-11.1	79.4	4.0	3.6-4.4	56.8	7.1	6.3-8.0	22.6	2.0	1.6-2.4	
Kidney and renal pelvis	1976-1980	110.2	8.6	7.9-9.4	73.6	13.1	11.6-14.7	36.6	5.3	4.5-6.1	64.4	5.2	4.6-5.8	43.4	8.2	7.0-9.5	21.0	3.0	2.4-3.6
	1981-1985	129.8	9.3	8.6-10.0	83.4	13.7	12.3-15.2	46.4	6.1	5.3-7.0	56.0	4.1	3.6-4.6	35.8	6.3	5.3-7.4	20.2	2.7	2.2-3.3
	1986-1990	153.6	10.8	10.0-11.6	92.0	14.8	13.4-16.4	61.6	7.8	6.9-8.7	61.4	4.3	3.8-4.8	37.0	6.0	5.1-7.1	24.4	3.0	2.5-3.6
	1991-1995	181.0	12.5	11.7-13.3	116.8	18.3	16.8-19.9	64.2	8.0	7.1-8.9	53.6	3.7	3.2-4.2	33.2	5.5	4.6-6.4	20.4	2.4	2.0-2.9
	1996-2000	196.8	13.2	12.4-14.0	121.4	18.5	17.1-20.1	75.4	9.1	8.2-10.1	54.4	3.6	3.2-4.1	32.6	5.1	4.3-6.0	21.8	2.5	2.0-3.0
	2001-2005	257.8	16.6	15.7-17.6	165.4	24.2	22.6-25.9	92.4	10.7	9.8-11.8	53.8	3.4	3.0-3.8	32.0	4.8	4.1-5.6	21.8	2.4	1.9-2.9
	2006-2010	292.4	18.3	17.4-19.3	188.8	26.5	24.8-28.3	103.6	11.6	10.6-12.7	54.0	3.2	2.8-3.6	33.6	4.8	4.1-5.5	20.4	2.0	1.6-2.5
	2011-2015	310.4	18.2	17.3-19.2	206.8	26.8	25.2-28.5	103.6	11.1	10.1-12.1	56.6	3.1	2.7-3.5	34.8	4.5	3.9-5.3	21.8	2.1	1.7-2.5
2016-2020	312.4	17.6	16.7-18.5	204.0	24.8	23.2-26.4	108.4	11.3	10.3-12.4	44.2	2.3	2.0-2.7	28.0	3.4	2.8-4.0	16.2	1.4	1.1-1.8	
Brain and other central nervous system	1976-1980	91.6	6.9	6.2-7.6	50.8	8.1	7.1-9.2	40.8	5.8	5.1-6.7	66.0	4.9	4.3-5.5	37.8	6.1	5.2-7.1	28.2	3.9	3.2-4.6
	1981-1985	94.0	6.9	6.2-7.5	52.8	8.6	7.5-9.8	41.2	5.6	4.8-6.4	63.4	4.5	4.0-5.0	32.2	5.1	4.3-6.0	31.2	4.1	3.5-4.9
	1986-1990	100.2	7.2	6.6-7.9	57.0	9.2	8.1-10.4	43.2	5.8	5.0-6.6	62.4	4.3	3.8-4.8	37.2	5.6	4.8-6.5	25.2	3.2	2.6-3.8
	1991-1995	107.6	7.5	6.9-8.2	60.6	9.4	8.3-10.5	47.0	6.0	5.3-6.9	65.6	4.5	4.0-5.0	35.2	5.3	4.6-6.2	30.4	3.8	3.2-4.5
	1996-2000	105.6	7.4	6.8-8.1	59.6	9.1	8.1-10.2	46.0	5.9	5.2-6.8	59.2	4.0	3.6-4.5	34.4	5.2	4.4-6.0	24.8	3.1	2.5-3.7
	2001-2005	104.4	7.2	6.6-7.9	58.0	8.7	7.7-9.8	46.4	6.0	5.2-6.9	61.8	4.1	3.7-4.6	36.0	5.3	4.6-6.2	25.8	3.1	2.6-3.7
	2006-2010	100.0	6.7	6.1-7.3	52.8	7.9	6.9-8.9	47.2	5.7	5.0-6.5	61.2	3.8	3.4-4.3	28.8	4.1	3.4-4.8	32.4	3.7	3.1-4.3
	2011-2015	105.2	6.7	6.2-7.4	58.2	8.0	7.1-9.0	47.0	5.7	4.9-6.5	66.6	3.9	3.5-4.4	37.0	5.0	4.3-5.8	29.6	3.1	2.6-3.6
2016-2020	113.6	7.2	6.6-7.8	61.2	8.2	7.3-9.2	52.4	6.3	5.5-7.2	69.0	3.9	3.5-4.4	39.8	4.9	4.2-5.7	29.2	3.1	2.6-3.7	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Nassau**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	58.2	4.6	4.1-5.2	19.8	3.4	2.7-4.2	38.4	5.6	4.8-6.5	8.0	0.7	0.5-1.0	2.6	0.6	0.3-1.1	5.4	0.8	0.5-1.2
	1981-1985	60.8	4.5	4.0-5.1	16.4	2.6	2.1-3.3	44.4	6.3	5.5-7.2	8.6	0.7	0.5-0.9	3.8	0.8	0.4-1.2	4.8	0.6	0.4-1.0
	1986-1990	69.2	5.1	4.6-5.7	19.6	3.1	2.5-3.8	49.6	7.0	6.1-7.9	9.2	0.7	0.5-0.9	3.2	0.5	0.3-0.8	6.0	0.8	0.5-1.1
	1991-1995	89.8	6.5	5.9-7.2	25.2	3.8	3.2-4.6	64.6	9.0	8.0-10.0	6.8	0.5	0.3-0.7	2.2	0.3	0.2-0.6	4.6	0.6	0.4-0.8
	1996-2000	133.6	9.6	8.9-10.4	34.6	5.2	4.5-6.1	99.0	13.8	12.6-15.0	8.6	0.6	0.4-0.8	4.0	0.6	0.4-0.9	4.6	0.5	0.3-0.8
	2001-2005	182.2	12.9	12.1-13.8	43.0	6.3	5.5-7.2	139.2	19.0	17.6-20.5	6.8	0.4	0.3-0.6	2.2	0.3	0.2-0.6	4.6	0.5	0.3-0.8
	2006-2010	275.4	19.2	18.2-20.3	77.0	11.2	10.1-12.4	198.4	26.7	25.0-28.5	8.6	0.5	0.4-0.7	4.2	0.6	0.4-0.9	4.4	0.5	0.3-0.7
	2011-2015	344.4	23.5	22.3-24.7	97.6	13.5	12.3-14.8	246.8	32.9	31.1-34.9	11.4	0.6	0.5-0.8	4.0	0.5	0.3-0.9	7.4	0.7	0.5-0.9
	2016-2020	333.6	22.9	21.8-24.1	97.6	13.3	12.1-14.6	236.0	32.2	30.3-34.2	11.6	0.6	0.5-0.8	5.6	0.7	0.5-1.1	6.0	0.5	0.4-0.8
Hodgkin lymphoma	1976-1980	59.4	4.4	3.9-4.9	33.4	5.2	4.4-6.1	26.0	3.6	3.0-4.3	17.4	1.3	1.1-1.7	9.4	1.6	1.1-2.1	8.0	1.1	0.8-1.5
	1981-1985	57.8	4.1	3.6-4.6	31.6	4.7	4.0-5.5	26.2	3.6	3.0-4.3	17.4	1.3	1.0-1.6	9.2	1.4	1.0-1.9	8.2	1.1	0.8-1.6
	1986-1990	59.6	4.3	3.8-4.9	27.2	4.2	3.5-5.0	32.4	4.5	3.8-5.2	12.4	0.9	0.7-1.2	7.4	1.2	0.9-1.7	5.0	0.7	0.4-1.0
	1991-1995	54.2	4.1	3.6-4.6	28.4	4.4	3.7-5.2	25.8	3.7	3.1-4.4	9.6	0.7	0.5-0.9	4.0	0.6	0.4-1.0	5.6	0.8	0.5-1.1
	1996-2000	49.6	3.8	3.3-4.3	23.4	3.7	3.1-4.5	26.2	4.0	3.3-4.7	8.6	0.6	0.4-0.8	4.8	0.7	0.5-1.1	3.8	0.5	0.3-0.8
	2001-2005	51.2	3.9	3.4-4.4	27.2	4.2	3.5-5.0	24.0	3.5	2.9-4.3	8.0	0.5	0.4-0.7	3.6	0.5	0.3-0.8	4.4	0.6	0.3-0.9
	2006-2010	49.4	3.7	3.2-4.2	29.6	4.5	3.8-5.3	19.8	2.9	2.3-3.6	5.6	0.4	0.2-0.5	3.6	0.5	0.3-0.8	2.0	0.2	0.1-0.4
	2011-2015	48.8	3.6	3.2-4.1	25.8	4.0	3.3-4.8	23.0	3.3	2.7-4.0	4.8	0.3	0.2-0.5	2.4	0.3	0.2-0.6	2.4	0.3	0.1-0.5
	2016-2020	50.0	3.5	3.1-4.0	25.0	3.6	3.0-4.3	25.0	3.5	2.9-4.2	5.2	0.3	0.2-0.4	2.8	0.3	0.2-0.6	2.4	0.2	0.1-0.4
Non-Hodgkin Lymphomas	1976-1980	172.2	13.5	12.6-14.5	91.8	16.0	14.5-17.7	80.4	11.4	10.3-12.6	89.6	7.1	6.5-7.9	43.4	7.5	6.4-8.7	46.2	6.6	5.8-7.6
	1981-1985	213.8	15.7	14.8-16.7	114.2	19.2	17.5-20.9	99.6	13.1	11.9-14.3	93.0	7.2	6.5-7.9	51.2	9.3	8.1-10.6	41.8	5.6	4.9-6.5
	1986-1990	255.2	18.1	17.1-19.2	136.8	21.7	20.0-23.5	118.4	15.2	14.0-16.5	114.6	8.4	7.7-9.1	63.2	10.5	9.3-11.8	51.4	6.6	5.8-7.4
	1991-1995	278.8	19.7	18.7-20.8	155.2	25.4	23.6-27.3	123.6	15.3	14.1-16.6	124.4	8.7	8.0-9.4	69.0	11.6	10.4-13.0	55.4	6.7	5.9-7.5
	1996-2000	324.0	21.9	20.8-23.0	167.6	26.2	24.4-28.0	156.4	18.8	17.5-20.2	130.8	8.7	8.1-9.4	68.2	11.0	9.9-12.3	62.6	7.2	6.4-8.1
	2001-2005	351.2	22.8	21.8-23.9	186.4	27.9	26.1-29.8	164.8	18.9	17.6-20.3	110.0	6.9	6.4-7.5	56.6	8.7	7.7-9.7	53.4	5.7	5.0-6.4
	2006-2010	393.4	24.7	23.6-25.9	204.2	29.2	27.4-31.1	189.2	21.3	19.9-22.8	99.6	5.9	5.4-6.5	55.4	8.0	7.1-9.0	44.2	4.4	3.8-5.1
	2011-2015	420.8	24.9	23.8-26.0	226.8	30.4	28.6-32.2	194.0	20.7	19.3-22.1	95.6	5.3	4.8-5.8	51.4	6.7	5.9-7.6	44.2	4.2	3.6-4.8
	2016-2020	401.2	22.7	21.6-23.7	212.6	26.7	25.1-28.4	188.6	19.5	18.2-20.8	95.4	5.1	4.6-5.6	50.0	6.3	5.5-7.1	45.4	4.2	3.7-4.8
Myeloma	1976-1980	55.4	4.6	4.1-5.2	28.6	5.9	4.9-7.1	26.8	3.9	3.2-4.6	34.8	3.0	2.5-3.5	18.6	3.9	3.1-4.8	16.2	2.4	1.9-3.0
	1981-1985	70.2	5.3	4.8-5.9	33.6	6.0	5.1-7.1	36.6	4.8	4.1-5.6	44.8	3.5	3.1-4.0	20.6	3.9	3.1-4.8	24.2	3.3	2.7-4.0
	1986-1990	72.0	5.3	4.8-5.9	38.8	6.9	5.9-8.1	33.2	4.3	3.7-5.0	47.8	3.6	3.1-4.1	25.4	4.9	4.0-5.9	22.4	2.8	2.3-3.4
	1991-1995	82.6	5.7	5.2-6.3	40.2	6.7	5.8-7.8	42.4	5.1	4.4-5.8	54.4	3.7	3.3-4.2	24.4	4.0	3.3-4.8	30.0	3.6	3.0-4.2
	1996-2000	94.0	6.2	5.7-6.8	45.4	7.1	6.2-8.1	48.6	5.7	5.0-6.4	51.2	3.4	3.0-3.8	27.6	4.5	3.7-5.3	23.6	2.6	2.2-3.2
	2001-2005	109.2	6.9	6.3-7.5	57.4	8.4	7.5-9.5	51.8	5.6	5.0-6.4	52.8	3.3	2.9-3.7	25.4	3.8	3.2-4.6	27.4	2.9	2.4-3.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Nassau**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	133.0	8.1	7.5-8.7	72.2	10.2	9.1-11.3	60.8	6.5	5.7-7.3	56.4	3.3	2.9-3.7	32.0	4.6	3.9-5.4	24.4	2.4	2.0-2.9
	2011-2015	155.6	8.9	8.3-9.6	85.6	11.3	10.2-12.4	70.0	7.0	6.3-7.8	57.6	3.2	2.8-3.6	29.4	3.8	3.2-4.5	28.2	2.7	2.3-3.2
	2016-2020	152.4	8.2	7.6-8.9	83.4	10.1	9.2-11.2	69.0	6.7	6.0-7.5	48.2	2.5	2.2-2.8	25.0	3.1	2.6-3.7	23.2	2.1	1.7-2.5
Leukemias	1976-1980	157.4	13.0	12.0-14.0	86.2	16.2	14.6-18.0	71.2	10.6	9.5-11.8	105.8	8.7	8.0-9.6	57.2	10.5	9.2-12.0	48.6	7.3	6.4-8.3
	1981-1985	168.4	13.1	12.2-14.0	91.2	15.7	14.2-17.3	77.2	10.9	9.8-12.1	107.6	8.5	7.8-9.3	55.6	10.5	9.1-11.9	52.0	7.2	6.4-8.2
	1986-1990	160.6	12.0	11.2-12.9	88.2	15.1	13.6-16.6	72.4	9.7	8.7-10.7	109.4	8.2	7.6-9.0	60.6	10.3	9.1-11.6	48.8	6.6	5.8-7.5
	1991-1995	181.0	13.1	12.2-14.0	100.2	16.9	15.3-18.5	80.8	10.5	9.5-11.6	106.2	7.7	7.0-8.3	57.0	10.1	8.9-11.4	49.2	6.1	5.4-7.0
	1996-2000	208.6	14.3	13.4-15.2	117.2	18.5	17.0-20.1	91.4	11.2	10.2-12.3	115.0	7.8	7.1-8.4	66.2	10.6	9.5-11.8	48.8	5.8	5.1-6.6
	2001-2005	227.6	14.9	14.1-15.8	130.0	19.6	18.1-21.2	97.6	11.4	10.3-12.4	117.2	7.5	6.9-8.1	65.4	10.0	8.9-11.1	51.8	5.7	5.0-6.5
	2006-2010	262.8	16.5	15.6-17.4	151.2	21.6	20.1-23.3	111.6	12.7	11.6-13.8	115.4	6.9	6.4-7.5	68.4	9.9	8.8-11.0	47.0	4.8	4.2-5.5
	2011-2015	324.6	19.3	18.3-20.2	187.6	25.3	23.6-27.0	137.0	14.7	13.6-15.9	112.6	6.4	5.9-7.0	65.8	9.0	8.0-10.0	46.8	4.5	3.9-5.1
2016-2020	288.2	16.5	15.6-17.4	173.0	22.0	20.5-23.5	115.2	12.3	11.2-13.4	108.0	5.7	5.2-6.2	65.0	8.3	7.4-9.2	43.0	3.8	3.3-4.4	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Niagara**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	876.0	407.7	395.4-420.3	446.2	486.6	465.4-508.6	429.8	359.1	343.7-375.0	485.6	228.2	219.0-237.8	260.6	291.3	274.6-308.6	225.0	186.6	175.7-198.1
	1981-1985	972.4	435.2	422.7-447.9	510.8	541.3	519.4-563.9	461.6	372.1	356.6-388.1	502.2	224.9	215.9-234.1	272.4	292.6	276.4-309.5	229.8	181.7	171.1-192.8
	1986-1990	1,039.4	442.9	430.7-455.5	521.8	526.3	505.3-547.9	517.6	398.2	382.6-414.4	517.0	219.6	211.0-228.4	275.2	286.9	271.1-303.3	241.8	178.8	168.6-189.5
	1991-1995	1,252.8	508.8	496.1-521.7	645.8	612.7	591.1-634.8	607.0	448.3	432.1-465.0	546.2	220.3	212.0-228.9	290.0	286.8	271.7-302.4	256.2	179.3	169.4-189.7
	1996-2000	1,306.2	519.8	507.2-532.6	689.6	641.1	619.6-663.2	616.6	441.7	426.0-457.9	540.6	211.1	203.1-219.2	278.4	266.9	252.8-281.6	262.2	177.2	167.6-187.3
	2001-2005	1,372.0	536.8	524.1-549.7	706.0	633.9	613.0-655.4	666.0	472.2	456.0-488.9	522.4	197.9	190.4-205.7	268.4	246.9	233.8-260.7	254.0	167.1	157.9-176.7
	2006-2010	1,396.8	522.9	510.5-535.5	718.4	602.2	582.4-622.5	678.4	469.6	453.5-486.2	500.8	181.6	174.4-188.9	250.0	213.5	201.7-225.9	250.8	161.0	152.0-170.5
	2011-2015	1,492.4	530.4	518.0-543.0	787.0	609.3	589.9-629.3	705.4	475.9	459.6-492.6	511.4	175.7	168.8-182.9	275.2	219.6	208.0-231.9	236.2	146.5	138.0-155.4
	2016-2020	1,594.4	541.7	529.3-554.3	842.4	608.5	589.5-628.0	752.0	493.7	477.0-510.8	520.8	170.4	163.7-177.3	277.8	206.7	195.6-218.3	243.0	145.0	136.6-153.9
Oral cavity and pharynx	1976-1980	28.2	13.0	10.9-15.4	19.0	19.7	15.8-24.3	9.2	7.9	5.7-10.6	10.0	4.7	3.5-6.3	7.0	7.7	5.2-10.9	3.0	2.4	1.3-4.0
	1981-1985	26.6	11.9	9.9-14.2	18.0	17.8	14.2-22.1	8.6	7.1	5.1-9.7	7.8	3.5	2.5-4.8	5.6	6.1	3.9-8.9	2.2	1.7	0.8-3.1
	1986-1990	23.8	10.4	8.5-12.5	16.4	16.1	12.7-20.1	7.4	5.8	4.1-8.2	8.2	3.5	2.5-4.8	5.6	5.8	3.7-8.5	2.6	2.0	1.0-3.5
	1991-1995	24.8	10.2	8.5-12.2	16.8	15.7	12.5-19.5	8.0	5.7	4.0-7.8	8.2	3.5	2.5-4.8	5.8	6.2	4.1-8.9	2.4	1.7	0.9-3.1
	1996-2000	22.0	9.1	7.4-10.9	15.4	14.6	11.5-18.3	6.6	5.0	3.4-7.1	7.6	3.1	2.2-4.2	5.2	4.7	3.1-7.0	2.4	1.9	1.0-3.3
	2001-2005	27.8	11.0	9.3-13.0	19.4	16.6	13.5-20.3	8.4	6.0	4.3-8.2	7.0	2.7	1.9-3.8	5.0	4.3	2.8-6.3	2.0	1.2	0.6-2.4
	2006-2010	34.6	12.6	10.7-14.6	24.0	19.1	15.8-22.9	10.6	7.2	5.3-9.5	7.2	2.5	1.8-3.6	5.4	4.4	2.9-6.5	1.8	1.1	0.5-2.2
	2011-2015	41.4	15.0	12.9-17.3	29.4	22.6	18.9-26.7	12.0	8.4	6.4-11.0	8.2	2.8	2.0-3.8	6.2	4.6	3.1-6.7	2.0	1.1	0.5-2.3
	2016-2020	41.2	13.5	11.6-15.7	26.8	18.2	15.1-21.8	14.4	9.4	7.2-12.1	8.4	2.7	1.9-3.8	6.0	4.1	2.7-6.1	2.4	1.6	0.8-3.0
Esophagus	1976-1980	10.6	4.8	3.6-6.3	7.2	7.3	5.0-10.3	3.4	2.8	1.6-4.6	9.2	4.4	3.2-5.9	6.2	6.6	4.4-9.5	3.0	2.6	1.4-4.4
	1981-1985	10.4	4.4	3.3-5.8	7.6	7.3	5.1-10.2	2.8	2.2	1.2-3.7	9.8	4.1	3.0-5.5	7.0	6.8	4.6-9.6	2.8	2.2	1.2-3.8
	1986-1990	11.6	4.7	3.6-6.2	7.6	6.9	4.8-9.5	4.0	2.8	1.7-4.4	9.0	3.6	2.6-4.9	6.8	6.2	4.3-8.8	2.2	1.5	0.8-2.8
	1991-1995	13.0	5.1	3.9-6.5	8.8	8.0	5.7-10.8	4.2	2.8	1.7-4.3	13.2	5.2	4.0-6.7	9.0	8.4	6.1-11.4	4.2	2.9	1.8-4.5
	1996-2000	16.6	6.6	5.3-8.2	14.2	13.0	10.1-16.5	2.4	1.7	0.9-3.0	11.2	4.4	3.3-5.7	10.2	9.6	7.1-12.7	1.0	0.7	0.2-1.7
	2001-2005	18.6	7.3	5.9-9.0	14.8	13.2	10.4-16.6	3.8	2.7	1.6-4.3	15.4	6.0	4.7-7.5	12.4	11.1	8.5-14.2	3.0	2.0	1.1-3.3
	2006-2010	16.8	6.1	4.8-7.6	12.4	10.0	7.6-12.9	4.4	2.9	1.8-4.5	17.4	6.2	4.9-7.7	13.8	11.4	8.8-14.5	3.6	2.1	1.3-3.5
	2011-2015	20.4	6.8	5.6-8.4	17.4	12.4	9.9-15.5	3.0	2.0	1.1-3.4	14.0	4.7	3.7-6.0	11.8	8.9	6.7-11.6	2.2	1.3	0.6-2.5
	2016-2020	20.2	6.3	5.1-7.7	16.4	11.1	8.8-14.0	3.8	2.3	1.3-3.7	18.2	5.6	4.5-7.0	14.6	9.7	7.5-12.4	3.6	2.2	1.3-3.7
Stomach	1976-1980	21.4	10.3	8.4-12.6	11.2	13.0	9.6-17.2	10.2	8.5	6.3-11.2	15.2	7.4	5.8-9.3	7.6	8.8	6.1-12.3	7.6	6.4	4.5-8.8
	1981-1985	18.8	7.8	6.3-9.7	12.8	12.0	9.1-15.5	6.0	4.7	3.2-6.9	15.4	6.8	5.3-8.5	9.6	10.3	7.5-13.8	5.8	4.5	3.0-6.6
	1986-1990	22.8	9.7	7.9-11.7	14.0	14.0	10.8-17.9	8.8	6.3	4.5-8.5	17.4	7.3	5.8-9.0	10.6	10.4	7.7-13.8	6.8	4.8	3.3-6.7
	1991-1995	19.2	7.9	6.4-9.6	12.2	12.0	9.0-15.5	7.0	4.9	3.4-6.9	11.8	4.9	3.7-6.3	8.4	9.0	6.4-12.3	3.4	2.3	1.3-3.7
	1996-2000	18.2	7.1	5.7-8.8	11.2	10.2	7.7-13.3	7.0	4.6	3.2-6.4	9.8	3.8	2.8-5.1	6.0	5.9	3.9-8.4	3.8	2.5	1.5-4.0
	2001-2005	19.6	7.3	6.0-9.0	10.2	9.5	7.0-12.5	9.4	6.0	4.4-8.1	12.4	4.6	3.6-6.0	6.4	5.9	4.0-8.4	6.0	3.8	2.5-5.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Niagara**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	19.8	7.3	5.9-8.9	12.4	10.4	8.0-13.4	7.4	4.5	3.1-6.4	9.8	3.5	2.6-4.7	5.8	5.0	3.4-7.3	4.0	2.4	1.4-3.9
	2011-2015	19.8	6.7	5.4-8.3	12.2	9.3	7.1-12.1	7.6	4.8	3.3-6.7	7.0	2.2	1.6-3.2	5.0	3.9	2.5-5.9	2.0	1.0	0.5-2.0
	2016-2020	20.4	6.8	5.5-8.4	12.6	9.4	7.2-12.2	7.8	4.7	3.3-6.6	6.8	2.2	1.5-3.2	4.0	3.0	1.8-4.9	2.8	1.6	0.9-2.8
Colorectal	1976-1980	135.0	63.7	58.9-68.9	69.6	77.5	69.1-86.6	65.4	53.5	47.8-59.7	71.0	34.0	30.5-37.8	39.0	43.9	37.6-50.9	32.0	26.7	22.7-31.2
	1981-1985	151.6	68.5	63.6-73.7	74.4	82.4	73.8-91.8	77.2	60.0	54.1-66.4	69.2	31.8	28.5-35.4	35.6	40.1	34.1-46.8	33.6	26.2	22.3-30.5
	1986-1990	141.8	59.6	55.2-64.2	68.8	69.3	61.8-77.4	73.0	52.8	47.4-58.7	73.2	31.8	28.5-35.2	36.6	39.8	34.0-46.4	36.6	26.7	22.9-30.9
	1991-1995	155.2	62.0	57.7-66.6	79.2	75.9	68.4-84.0	76.0	52.3	47.1-57.9	65.0	25.9	23.2-29.0	33.4	32.3	27.5-37.8	31.6	21.0	17.8-24.6
	1996-2000	163.4	63.5	59.2-68.1	83.2	79.4	71.8-87.6	80.2	53.0	47.9-58.5	58.2	22.4	19.9-25.1	29.2	29.5	24.8-34.8	29.0	18.1	15.3-21.4
	2001-2005	140.6	53.7	49.7-57.8	70.4	64.0	57.4-71.1	70.2	45.7	40.9-50.9	54.2	20.3	18.0-22.9	27.0	25.2	21.1-29.9	27.2	16.9	14.1-20.1
	2006-2010	119.0	43.5	40.0-47.2	60.2	50.9	45.3-57.1	58.8	38.0	33.7-42.8	43.4	15.5	13.5-17.7	20.4	17.7	14.4-21.6	23.0	14.0	11.5-17.0
	2011-2015	117.0	41.6	38.2-45.2	63.8	50.7	45.1-56.8	53.2	34.9	30.6-39.7	33.4	11.3	9.6-13.2	17.2	13.7	10.9-17.1	16.2	9.6	7.5-12.1
2016-2020	122.4	43.5	40.0-47.3	67.8	53.3	47.6-59.6	54.6	35.2	30.9-39.9	35.4	12.0	10.2-14.0	17.6	13.1	10.5-16.4	17.8	10.9	8.6-13.6	
Colon excluding rectum	1976-1980	90.4	43.6	39.6-47.9	44.0	51.7	44.7-59.4	46.4	38.1	33.3-43.4	50.2	24.2	21.2-27.5	26.8	30.5	25.2-36.5	23.4	19.5	16.1-23.4
	1981-1985	102.8	46.9	42.8-51.2	47.6	54.4	47.3-62.2	55.2	42.8	37.8-48.2	52.2	24.2	21.3-27.3	26.2	29.9	24.7-35.8	26.0	20.3	16.9-24.2
	1986-1990	105.4	44.3	40.6-48.4	50.8	51.9	45.5-59.0	54.6	39.1	34.6-44.2	60.0	25.7	22.8-28.8	30.0	32.4	27.1-38.3	30.0	21.4	18.0-25.1
	1991-1995	113.2	45.0	41.3-48.9	54.4	52.5	46.2-59.4	58.8	39.9	35.4-44.9	52.0	20.7	18.2-23.4	26.4	25.4	21.1-30.3	25.6	17.0	14.2-20.3
	1996-2000	120.4	46.5	42.8-50.4	57.2	55.4	49.0-62.3	63.2	41.3	36.8-46.2	50.6	19.3	17.0-21.9	24.4	24.9	20.6-29.9	26.2	16.3	13.6-19.4
	2001-2005	100.2	38.0	34.7-41.5	48.8	44.6	39.1-50.6	51.4	32.8	28.8-37.2	46.0	17.2	15.0-19.6	23.0	21.5	17.7-25.9	23.0	14.2	11.7-17.1
	2006-2010	84.8	31.2	28.3-34.4	40.4	34.6	29.9-39.8	44.4	28.7	25.0-32.9	36.0	12.8	11.0-14.9	15.6	13.6	10.8-17.1	20.4	12.3	10.0-15.1
	2011-2015	79.2	27.9	25.1-30.9	40.8	32.5	28.1-37.5	38.4	24.4	20.9-28.4	26.6	9.1	7.5-10.8	12.8	10.4	7.9-13.3	13.8	8.1	6.2-10.5
2016-2020	85.2	30.2	27.3-33.4	44.4	34.7	30.1-39.8	40.8	26.5	22.8-30.6	29.8	10.2	8.6-12.1	14.6	11.1	8.6-14.1	15.2	9.4	7.2-12.0	
Rectum & rectosigmoid	1976-1980	44.6	20.1	17.5-23.0	25.6	25.8	21.3-30.9	19.0	15.4	12.4-18.9	20.8	9.8	8.0-11.9	12.2	13.4	10.1-17.4	8.6	7.2	5.2-9.8
	1981-1985	48.8	21.6	18.9-24.6	26.8	28.0	23.2-33.5	22.0	17.2	14.1-20.8	17.0	7.6	6.1-9.5	9.4	10.2	7.4-13.7	7.6	5.8	4.1-8.1
	1986-1990	36.4	15.2	13.1-17.7	18.0	17.4	13.8-21.6	18.4	13.7	11.0-16.9	13.2	6.1	4.7-7.7	6.6	7.4	5.0-10.5	6.6	5.3	3.6-7.5
	1991-1995	42.0	17.0	14.8-19.5	24.8	23.4	19.4-28.1	17.2	12.4	9.8-15.4	13.0	5.2	4.0-6.7	7.0	6.9	4.7-9.7	6.0	4.0	2.7-5.7
	1996-2000	43.0	17.0	14.8-19.5	26.0	24.0	20.0-28.6	17.0	11.7	9.3-14.6	7.6	3.0	2.1-4.2	4.8	4.5	2.9-6.8	2.8	1.8	1.0-3.2
	2001-2005	40.4	15.7	13.6-18.0	21.6	19.4	15.9-23.5	18.8	12.9	10.4-15.9	8.2	3.1	2.2-4.3	4.0	3.7	2.2-5.8	4.2	2.7	1.6-4.2
	2006-2010	34.2	12.3	10.5-14.4	19.8	16.4	13.3-20.0	14.4	9.3	7.2-11.9	7.4	2.7	1.9-3.7	4.8	4.1	2.6-6.1	2.6	1.7	0.9-3.1
	2011-2015	37.8	13.7	11.8-15.9	23.0	18.2	14.9-22.0	14.8	10.5	8.1-13.4	6.8	2.2	1.5-3.2	4.4	3.4	2.1-5.2	2.4	1.5	0.7-2.7
2016-2020	37.2	13.3	11.3-15.5	23.4	18.6	15.3-22.5	13.8	8.7	6.7-11.2	5.6	1.8	1.2-2.6	3.0	2.1	1.1-3.5	2.6	1.5	0.8-2.7	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Niagara**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	6.6	3.2	2.2-4.6	2.4	2.4	1.2-4.3	4.2	3.7	2.3-5.7	6.4	3.0	2.0-4.3	3.0	3.0	1.6-5.1	3.4	3.0	1.7-4.8
	1981-1985	5.6	2.6	1.7-3.8	3.4	4.0	2.3-6.4	2.2	1.6	0.8-3.0	3.6	1.6	0.9-2.5	2.4	2.7	1.3-4.7	1.2	0.9	0.3-2.0
	1986-1990	7.8	3.2	2.3-4.5	5.0	4.9	3.1-7.4	2.8	1.9	1.1-3.3	6.2	2.7	1.8-3.9	4.0	4.7	2.8-7.3	2.2	1.5	0.8-2.8
	1991-1995	8.6	3.5	2.6-4.8	4.8	4.4	2.8-6.7	3.8	2.7	1.6-4.3	8.2	3.3	2.4-4.5	5.0	4.5	2.9-6.8	3.2	2.1	1.2-3.5
	1996-2000	10.4	4.1	3.0-5.4	7.6	7.1	5.0-9.8	2.8	1.7	0.9-3.0	10.8	4.1	3.1-5.4	6.6	6.2	4.3-8.8	4.2	2.6	1.6-4.1
	2001-2005	11.4	4.4	3.3-5.7	7.6	6.8	4.8-9.4	3.8	2.5	1.5-4.0	9.4	3.5	2.6-4.7	5.4	4.9	3.2-7.2	4.0	2.6	1.6-4.1
	2006-2010	15.4	5.7	4.5-7.2	10.2	8.4	6.2-11.1	5.2	3.4	2.2-5.1	13.2	4.9	3.7-6.2	8.0	6.9	4.9-9.5	5.2	3.0	2.0-4.6
	2011-2015	21.2	7.1	5.8-8.7	14.8	10.4	8.1-13.3	6.4	4.2	2.8-6.1	17.8	6.0	4.8-7.5	10.2	7.5	5.6-10.0	7.6	4.6	3.2-6.5
	2016-2020	26.2	8.2	6.9-9.9	18.6	12.5	10.0-15.5	7.6	4.5	3.2-6.4	18.2	5.7	4.6-7.1	11.4	7.7	5.8-10.1	6.8	4.0	2.8-5.8
Pancreas	1976-1980	31.2	14.9	12.6-17.5	16.8	18.5	14.6-23.2	14.4	12.1	9.5-15.4	27.8	13.3	11.1-15.7	14.8	16.7	12.9-21.2	13.0	10.7	8.3-13.8
	1981-1985	28.2	12.7	10.7-15.1	16.4	18.4	14.4-23.1	11.8	9.0	6.8-11.7	28.2	13.1	11.0-15.5	15.2	17.8	13.8-22.5	13.0	10.1	7.8-12.9
	1986-1990	27.4	11.5	9.6-13.6	14.2	14.5	11.1-18.5	13.2	9.3	7.2-11.9	25.8	10.7	8.9-12.7	14.2	13.9	10.7-17.8	11.6	8.2	6.2-10.7
	1991-1995	34.2	13.8	11.8-16.1	15.8	15.5	12.2-19.5	18.4	12.3	9.9-15.2	29.0	11.6	9.8-13.7	12.4	11.8	9.0-15.1	16.6	11.1	8.8-13.8
	1996-2000	33.8	13.1	11.2-15.2	13.6	12.9	9.9-16.4	20.2	13.1	10.6-16.0	28.8	11.1	9.3-13.1	13.0	12.1	9.3-15.5	15.8	10.1	7.9-12.6
	2001-2005	30.6	11.7	9.9-13.7	15.0	13.4	10.5-16.9	15.6	10.2	8.0-12.8	30.4	11.6	9.8-13.6	14.8	13.2	10.3-16.6	15.6	10.2	8.0-12.8
	2006-2010	28.4	10.3	8.6-12.2	16.0	13.3	10.5-16.6	12.4	8.0	6.1-10.4	29.2	10.3	8.7-12.2	16.2	13.6	10.8-17.0	13.0	7.9	6.1-10.2
	2011-2015	42.2	14.2	12.3-16.3	21.6	16.1	13.1-19.5	20.6	12.5	10.2-15.3	30.2	10.0	8.5-11.8	17.0	12.5	10.0-15.6	13.2	8.0	6.1-10.3
	2016-2020	47.2	15.4	13.4-17.6	25.8	19.0	15.7-22.7	21.4	12.5	10.2-15.4	37.0	12.2	10.4-14.1	19.0	14.1	11.3-17.4	18.0	10.6	8.5-13.2
Larynx	1976-1980	14.4	6.1	4.7-7.7	11.6	11.0	8.3-14.4	2.8	2.2	1.2-3.8	4.2	2.0	1.2-3.0	3.4	3.6	2.0-5.9	0.8	0.8	0.2-2.0
	1981-1985	14.8	6.6	5.1-8.3	12.6	12.9	9.8-16.7	2.2	1.8	0.9-3.3	4.4	1.8	1.1-2.8	3.6	3.9	2.2-6.3	0.8	0.6	0.2-1.5
	1986-1990	10.2	4.1	3.0-5.4	8.0	7.7	5.4-10.6	2.2	1.6	0.8-3.0	5.8	2.4	1.6-3.5	5.2	5.3	3.3-7.9	0.6	0.6	0.1-1.7
	1991-1995	13.2	5.6	4.3-7.1	9.4	8.9	6.5-11.8	3.8	3.1	1.8-4.9	4.2	1.6	1.0-2.5	3.4	3.0	1.7-4.8	0.8	0.6	0.2-1.6
	1996-2000	12.8	5.2	4.0-6.6	10.6	9.7	7.2-12.7	2.2	1.6	0.8-2.9	3.8	1.5	0.9-2.3	3.4	3.1	1.8-5.0	0.4	0.3	0.0-1.1
	2001-2005	12.8	5.0	3.9-6.5	10.0	8.6	6.4-11.3	2.8	2.0	1.1-3.4	2.4	0.9	0.5-1.6	1.4	1.3	0.5-2.7	1.0	0.7	0.2-1.6
	2006-2010	10.0	3.6	2.7-4.8	6.8	5.6	3.8-7.9	3.2	2.1	1.2-3.5	2.8	1.0	0.6-1.8	1.6	1.4	0.6-2.8	1.2	0.7	0.3-1.7
	2011-2015	11.8	3.8	2.9-4.9	9.2	6.5	4.7-8.7	2.6	1.6	0.8-2.9	3.4	1.1	0.6-1.8	2.6	1.8	1.0-3.2	0.8	0.5	0.1-1.4
	2016-2020	12.6	4.1	3.1-5.3	11.0	7.6	5.6-10.0	1.6	1.0	0.4-2.1	3.4	1.1	0.6-1.8	2.4	1.8	0.9-3.2	1.0	0.5	0.2-1.4
Lung and bronchus	1976-1980	140.6	62.9	58.2-67.9	101.2	105.6	96.0-115.8	39.4	32.7	28.2-37.8	119.2	53.0	48.7-57.6	87.8	93.3	84.3-103.1	31.4	24.9	21.1-29.3
	1981-1985	169.4	72.8	67.9-78.1	115.0	116.2	106.5-126.7	54.4	41.9	37.0-47.4	139.2	59.9	55.4-64.6	95.4	96.9	88.0-106.5	43.8	33.4	29.0-38.2
	1986-1990	182.4	75.8	70.8-81.0	113.8	110.5	101.2-120.3	68.6	51.5	46.0-57.4	146.4	60.0	55.6-64.6	93.8	90.0	81.7-98.9	52.6	38.9	34.3-44.1
	1991-1995	198.8	78.9	74.1-84.1	115.6	108.9	100.0-118.4	83.2	58.9	53.3-65.0	164.6	65.6	61.2-70.3	99.2	95.5	87.1-104.6	65.4	45.6	40.7-51.0
	1996-2000	202.8	79.4	74.6-84.5	113.2	104.1	95.6-113.1	89.6	62.2	56.5-68.3	162.2	63.1	58.9-67.7	91.6	84.1	76.5-92.3	70.6	48.5	43.5-54.0
	2001-2005	238.0	91.5	86.3-96.8	122.2	109.4	100.8-118.5	115.8	79.0	72.7-85.9	174.4	66.2	61.9-70.8	95.4	86.7	79.0-94.9	79.0	53.1	47.9-58.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Niagara**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	217.6	80.3	75.5-85.2	109.4	92.7	85.1-101.0	108.2	71.8	65.8-78.3	154.0	56.4	52.5-60.6	79.8	67.6	61.1-74.7	74.2	48.4	43.5-53.8
	2011-2015	240.8	82.7	78.0-87.6	127.0	98.4	90.7-106.6	113.8	72.1	66.2-78.5	164.8	56.6	52.7-60.7	91.6	72.5	65.9-79.7	73.2	45.7	41.0-50.8
	2016-2020	228.4	72.8	68.5-77.3	109.0	76.8	70.3-83.8	119.4	71.1	65.4-77.3	139.4	44.9	41.5-48.5	75.2	55.2	49.6-61.3	64.2	37.9	33.7-42.4
Melanoma of the skin	1976-1980	9.6	4.4	3.2-5.9	6.2	6.2	4.1-8.9	3.4	2.9	1.7-4.8	3.4	1.5	0.9-2.5	2.6	2.5	1.3-4.3	0.8	0.7	0.2-1.7
	1981-1985	12.0	5.4	4.1-7.1	6.8	6.8	4.6-9.6	5.2	4.6	2.9-6.8	2.4	1.0	0.5-1.8	1.6	1.3	0.6-2.8	0.8	0.6	0.2-1.6
	1986-1990	20.4	9.2	7.5-11.2	11.2	11.2	8.4-14.6	9.2	7.7	5.6-10.4	5.4	2.6	1.7-3.8	3.6	3.9	2.3-6.3	1.8	1.7	0.8-3.3
	1991-1995	21.6	9.2	7.5-11.2	13.2	12.5	9.6-16.0	8.4	6.7	4.8-9.1	4.6	2.1	1.3-3.1	2.8	2.7	1.5-4.6	1.8	1.6	0.7-3.0
	1996-2000	30.8	12.7	10.7-14.9	19.2	17.3	14.0-21.2	11.6	9.3	7.1-12.1	5.8	2.3	1.6-3.4	3.2	3.2	1.8-5.3	2.6	1.9	1.0-3.4
	2001-2005	41.2	16.9	14.6-19.4	25.8	23.2	19.4-27.6	15.4	12.5	9.8-15.6	5.6	2.1	1.4-3.1	4.0	3.6	2.2-5.6	1.6	1.0	0.4-2.2
	2006-2010	51.6	20.1	17.7-22.7	31.2	26.5	22.5-31.1	20.4	15.8	12.8-19.3	5.6	2.0	1.3-2.9	2.8	2.3	1.2-3.9	2.8	1.9	1.0-3.2
	2011-2015	70.4	26.0	23.2-29.0	38.4	29.7	25.6-34.4	32.0	24.0	20.2-28.3	6.0	2.1	1.4-3.1	4.8	3.8	2.4-5.7	1.2	0.9	0.3-2.1
2016-2020	77.0	28.3	25.4-31.5	40.8	30.6	26.4-35.4	36.2	27.4	23.2-32.1	5.2	1.7	1.1-2.6	2.8	2.2	1.2-3.8	2.4	1.3	0.6-2.4	
Female breast	1976-1980							112.2	96.0	88.0-104.5							48.2	40.5	35.4-46.0
	1981-1985							123.2	104.3	95.9-113.2							45.0	36.9	32.1-42.3
	1986-1990							156.2	124.7	115.8-134.1							43.4	33.3	28.9-38.2
	1991-1995							172.6	133.4	124.4-142.9							40.2	29.4	25.4-33.9
	1996-2000							178.2	131.9	123.3-141.0							44.8	31.7	27.6-36.3
	2001-2005							185.6	135.7	127.0-144.9							37.0	25.3	21.7-29.4
	2006-2010							197.0	138.6	129.9-147.7							35.4	23.6	20.2-27.6
	2011-2015							194.8	134.0	125.4-143.1							33.4	20.4	17.3-23.9
2016-2020							210.6	139.6	130.8-148.9							31.0	20.0	16.7-23.7	
Cervix uteri	1976-1980							17.0	15.2	12.0-18.9							6.2	5.2	3.5-7.4
	1981-1985							9.4	7.8	5.7-10.5							4.2	3.6	2.2-5.5
	1986-1990							9.8	8.2	6.0-10.9							3.6	2.7	1.6-4.4
	1991-1995							11.2	9.1	6.8-11.9							3.4	2.6	1.5-4.2
	1996-2000							10.6	8.6	6.4-11.2							3.6	2.6	1.5-4.2
	2001-2005							8.4	6.8	4.9-9.2							2.2	1.8	0.9-3.3
	2006-2010							7.2	6.1	4.2-8.5							1.8	1.4	0.6-2.7
	2011-2015							8.8	7.0	5.0-9.5							2.0	1.4	0.6-2.6
	2016-2020							10.6	10.4	7.7-13.8							2.2	1.3	0.6-2.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Niagara**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							30.6	24.1	20.3-28.4							6.0	4.8	3.2-6.9
	1981-1985							28.4	21.9	18.3-26.0							5.2	3.8	2.5-5.6
	1986-1990							29.0	23.3	19.5-27.6							6.0	4.3	2.9-6.2
	1991-1995							36.8	27.2	23.3-31.6							7.4	5.2	3.6-7.3
	1996-2000							36.4	27.8	23.9-32.2							5.2	3.4	2.2-5.1
	2001-2005							39.8	29.7	25.7-34.2							3.2	2.1	1.2-3.4
	2006-2010							44.0	31.2	27.1-35.7							5.4	3.7	2.4-5.4
	2011-2015							47.8	30.4	26.6-34.7							7.2	4.9	3.4-6.9
	2016-2020							49.4	32.5	28.3-37.1							9.2	5.2	3.8-7.1
Ovary	1976-1980							19.4	16.5	13.3-20.2							15.0	12.4	9.7-15.7
	1981-1985							19.0	15.7	12.6-19.3							12.0	9.9	7.5-12.8
	1986-1990							21.4	16.7	13.5-20.3							14.2	10.4	8.1-13.3
	1991-1995							27.2	21.1	17.6-25.1							15.2	10.6	8.3-13.3
	1996-2000							20.4	15.3	12.4-18.6							11.0	7.6	5.7-9.9
	2001-2005							22.2	16.3	13.3-19.7							12.6	8.5	6.5-11.0
	2006-2010							17.8	12.6	10.0-15.6							12.8	8.6	6.6-11.1
	2011-2015							17.6	11.7	9.3-14.6							11.0	7.4	5.5-9.8
	2016-2020							17.8	11.4	9.0-14.3							13.0	7.5	5.7-9.7
Prostate	1976-1980				70.8	87.2	77.8-97.3							27.2	38.2	31.7-45.4			
	1981-1985				94.4	108.3	98.3-119.1							27.2	34.5	28.6-41.1			
	1986-1990				102.4	107.7	98.1-118.1							31.0	38.1	32.0-44.9			
	1991-1995				188.0	174.9	163.5-186.8							31.4	34.5	29.1-40.6			
	1996-2000				198.6	182.4	171.1-194.3							28.0	29.5	24.7-34.9			
	2001-2005				198.2	175.4	164.6-186.7							23.4	23.1	19.1-27.8			
	2006-2010				207.8	168.6	158.4-179.3							19.2	16.9	13.7-20.7			
	2011-2015				193.6	140.6	131.7-150.1							20.4	17.2	14.0-20.9			
	2016-2020				226.0	148.1	139.4-157.4							24.0	18.9	15.6-22.7			
Testis	1976-1980				3.6	3.6	2.1-5.7							0.4	0.5	0.1-1.8			
	1981-1985				7.2	6.0	4.2-8.4							1.0	0.9	0.3-2.1			
	1986-1990				5.2	4.5	2.9-6.6							0.4	0.3	0.0-1.3			
	1991-1995				6.4	5.7	3.9-8.2							0.0	0.0	0.0-0.7			
	1996-2000				8.6	8.2	5.9-11.0							0.2	0.2	0.0-1.1			
	2001-2005				5.8	5.8	3.9-8.4							0.4	0.3	0.0-1.3			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Niagara**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				6.0	5.9	3.9-8.4							0.4	0.4	0.0-1.3			
	2011-2015				7.2	7.1	4.9-10.0							0.4	0.5	0.1-1.8			
	2016-2020				8.6	9.5	6.8-12.9							0.4	0.3	0.0-1.2			
Urinary bladder (incl. in situ)	1976-1980	49.8	23.3	20.4-26.5	36.6	40.2	34.2-46.9	13.2	11.0	8.5-14.1	13.2	6.2	4.8-8.0	10.2	11.7	8.5-15.6	3.0	2.5	1.4-4.2
	1981-1985	55.2	24.6	21.7-27.8	42.2	45.5	39.2-52.4	13.0	10.2	7.9-13.1	10.2	4.5	3.3-6.0	7.2	7.6	5.2-10.8	3.0	2.5	1.4-4.1
	1986-1990	58.0	24.4	21.6-27.4	44.4	44.4	38.5-51.0	13.6	9.9	7.6-12.6	10.2	4.5	3.4-6.0	7.4	8.4	5.8-11.7	2.8	2.1	1.1-3.5
	1991-1995	67.6	27.6	24.7-30.7	47.0	45.8	40.0-52.2	20.6	14.6	11.9-17.9	10.0	4.1	3.0-5.4	6.8	7.0	4.8-9.9	3.2	2.1	1.2-3.5
	1996-2000	82.8	32.4	29.4-35.7	59.2	56.6	50.2-63.6	23.6	15.7	13.0-18.9	15.0	5.8	4.6-7.3	11.0	11.6	8.6-15.1	4.0	2.7	1.6-4.2
	2001-2005	80.4	30.8	27.9-34.0	57.6	52.6	46.6-59.1	22.8	15.3	12.6-18.5	12.0	4.4	3.4-5.7	7.4	7.1	5.0-9.8	4.6	2.7	1.7-4.2
	2006-2010	86.4	31.6	28.6-34.7	62.6	53.7	47.8-60.0	23.8	15.5	12.8-18.7	15.4	5.3	4.2-6.7	9.4	8.2	6.0-11.0	6.0	3.7	2.4-5.4
	2011-2015	96.4	33.3	30.4-36.5	72.8	58.2	52.3-64.7	23.6	14.5	12.0-17.6	15.8	5.2	4.1-6.6	11.6	9.7	7.3-12.6	4.2	2.4	1.5-3.8
2016-2020	103.4	33.2	30.3-36.3	79.0	57.8	52.1-64.1	24.4	14.1	11.7-17.0	14.2	4.6	3.6-5.9	11.0	8.5	6.3-11.2	3.2	2.0	1.1-3.5	
Kidney and renal pelvis	1976-1980	15.8	7.1	5.6-8.9	9.0	9.8	7.0-13.3	6.8	5.4	3.7-7.6	8.2	3.8	2.7-5.2	4.6	5.0	3.1-7.6	3.6	3.0	1.7-4.7
	1981-1985	17.2	8.2	6.5-10.1	8.8	9.5	6.8-12.9	8.4	7.2	5.1-9.8	7.4	3.5	2.5-4.9	3.6	4.0	2.3-6.4	3.8	3.2	1.9-5.0
	1986-1990	16.4	7.0	5.5-8.7	10.0	10.0	7.3-13.3	6.4	4.7	3.2-6.7	7.6	3.1	2.2-4.3	3.4	3.8	2.1-6.1	4.2	2.8	1.7-4.4
	1991-1995	32.2	12.8	10.9-15.0	19.0	16.9	13.6-20.8	13.2	9.7	7.4-12.4	12.4	4.9	3.7-6.3	8.4	8.0	5.7-10.9	4.0	2.8	1.7-4.5
	1996-2000	35.6	14.1	12.1-16.4	21.0	19.2	15.7-23.3	14.6	10.2	7.9-12.9	11.0	4.3	3.2-5.6	6.4	6.2	4.2-8.8	4.6	3.0	1.9-4.6
	2001-2005	37.0	14.4	12.4-16.6	20.6	17.8	14.6-21.7	16.4	11.4	9.0-14.2	10.0	3.7	2.8-4.9	5.2	4.7	3.1-6.9	4.8	2.8	1.8-4.3
	2006-2010	43.2	16.1	14.0-18.4	26.4	21.6	18.1-25.7	16.8	11.7	9.3-14.6	10.4	3.8	2.8-5.0	6.0	4.8	3.2-7.0	4.4	2.9	1.8-4.5
	2011-2015	51.4	17.9	15.7-20.3	31.6	24.0	20.3-28.2	19.8	12.6	10.2-15.5	10.8	3.6	2.7-4.8	7.2	5.6	3.9-7.8	3.6	2.1	1.3-3.5
2016-2020	57.2	19.9	17.5-22.5	39.0	29.0	24.9-33.7	18.2	11.9	9.4-14.8	11.0	3.7	2.8-4.9	6.4	4.9	3.3-7.1	4.6	2.7	1.7-4.2	
Brain and other central nervous system	1976-1980	11.8	5.0	3.8-6.5	7.0	6.2	4.3-8.8	4.8	3.8	2.4-5.8	9.8	4.1	3.0-5.5	5.0	4.3	2.7-6.5	4.8	3.9	2.4-5.8
	1981-1985	14.0	6.3	4.9-8.0	6.8	6.7	4.6-9.5	7.2	6.0	4.1-8.4	11.2	5.0	3.7-6.5	5.6	5.5	3.6-8.1	5.6	4.6	3.0-6.8
	1986-1990	14.4	6.4	5.0-8.1	7.4	7.4	5.1-10.2	7.0	5.5	3.8-7.7	10.2	4.4	3.3-5.9	4.6	4.6	2.9-7.0	5.6	4.1	2.7-6.0
	1991-1995	16.6	7.2	5.7-8.9	8.8	8.1	5.9-11.0	7.8	6.3	4.5-8.7	9.6	4.2	3.0-5.5	4.8	4.6	2.9-6.9	4.8	3.9	2.5-5.9
	1996-2000	20.0	8.4	6.8-10.3	11.2	10.2	7.7-13.3	8.8	6.5	4.7-8.9	11.0	4.5	3.4-5.9	5.6	5.0	3.3-7.2	5.4	4.1	2.6-6.0
	2001-2005	19.0	7.9	6.4-9.7	10.8	9.9	7.4-13.0	8.2	6.0	4.3-8.3	11.4	4.5	3.4-5.9	6.2	5.5	3.7-7.8	5.2	3.7	2.4-5.4
	2006-2010	16.8	6.9	5.4-8.6	8.8	7.7	5.5-10.4	8.0	6.3	4.4-8.7	9.4	3.3	2.4-4.4	6.2	5.0	3.4-7.1	3.2	1.9	1.1-3.2
	2011-2015	21.0	8.5	6.9-10.4	12.4	10.5	8.0-13.6	8.6	6.8	4.9-9.4	10.6	3.7	2.8-4.9	6.0	4.5	3.0-6.5	4.6	3.2	2.0-4.9
2016-2020	17.0	6.5	5.1-8.2	10.2	8.3	6.1-11.2	6.8	4.8	3.2-6.9	14.0	4.8	3.7-6.2	9.0	6.5	4.7-8.9	5.0	3.2	2.0-5.0	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Niagara**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	10.2	4.9	3.6-6.5	3.8	4.0	2.3-6.3	6.4	5.7	3.8-8.1	2.0	0.9	0.4-1.7	0.6	0.5	0.1-1.5	1.4	1.2	0.5-2.4
	1981-1985	7.8	3.8	2.7-5.2	2.2	2.2	1.1-4.1	5.6	5.3	3.4-7.7	0.8	0.4	0.1-1.1	0.6	0.8	0.2-2.2	0.2	0.2	0.0-1.0
	1986-1990	8.8	4.2	3.0-5.6	3.0	3.0	1.6-5.0	5.8	5.3	3.5-7.6	1.0	0.4	0.1-1.0	0.8	0.8	0.2-2.0	0.2	0.1	0.0-0.9
	1991-1995	15.2	6.8	5.4-8.6	2.4	2.3	1.2-4.0	12.8	11.3	8.6-14.4	1.6	0.7	0.3-1.3	0.6	0.6	0.1-1.8	1.0	0.7	0.2-1.7
	1996-2000	21.2	9.3	7.6-11.3	4.8	4.3	2.7-6.4	16.4	14.1	11.2-17.5	1.2	0.5	0.2-1.0	0.6	0.5	0.1-1.6	0.6	0.4	0.1-1.3
	2001-2005	25.2	10.9	9.1-13.0	5.4	4.7	3.1-6.9	19.8	16.9	13.7-20.6	0.2	0.1	0.0-0.5	0.2	0.2	0.0-1.0	0.0	0.0	0.0-0.7
	2006-2010	26.6	11.5	9.6-13.7	7.0	6.1	4.2-8.5	19.6	16.7	13.4-20.4	0.6	0.2	0.0-0.7	0.2	0.1	0.0-0.9	0.4	0.3	0.0-1.1
	2011-2015	41.8	18.4	15.8-21.2	12.4	10.8	8.1-14.0	29.4	26.0	21.7-30.8	1.6	0.5	0.2-1.1	1.0	0.8	0.2-1.9	0.6	0.4	0.1-1.2
	2016-2020	37.0	16.5	14.1-19.3	10.2	9.1	6.6-12.2	26.8	23.8	19.7-28.5	1.8	0.7	0.3-1.3	1.0	0.9	0.3-2.1	0.8	0.6	0.1-1.7
Hodgkin lymphoma	1976-1980	7.8	3.2	2.2-4.4	4.8	4.4	2.7-6.7	3.0	2.1	1.2-3.6	2.6	1.3	0.7-2.3	1.8	2.2	1.0-4.2	0.8	0.8	0.2-1.9
	1981-1985	7.4	3.4	2.4-4.8	4.6	4.7	2.9-7.2	2.8	2.4	1.3-4.1	3.6	1.7	1.0-2.7	1.8	2.0	0.9-3.9	1.8	1.5	0.6-2.8
	1986-1990	7.6	3.3	2.3-4.6	3.4	3.3	1.9-5.5	4.2	3.5	2.1-5.3	1.4	0.5	0.2-1.2	0.8	0.6	0.2-1.8	0.6	0.5	0.1-1.4
	1991-1995	6.8	3.0	2.1-4.2	4.0	3.8	2.3-5.9	2.8	2.2	1.2-3.7	1.8	0.7	0.3-1.4	0.8	0.8	0.2-2.0	1.0	0.7	0.2-1.7
	1996-2000	6.6	3.1	2.1-4.3	3.4	3.3	1.9-5.2	3.2	2.8	1.6-4.6	0.8	0.3	0.1-0.8	0.8	0.7	0.2-1.9	0.0	0.0	0.0-0.6
	2001-2005	6.4	3.0	2.1-4.3	4.0	3.9	2.4-6.0	2.4	2.2	1.1-3.8	1.2	0.5	0.2-1.1	0.8	0.7	0.2-1.8	0.4	0.3	0.0-1.1
	2006-2010	5.8	2.6	1.7-3.7	3.2	3.0	1.7-4.9	2.6	2.1	1.1-3.7	0.6	0.3	0.1-0.8	0.4	0.4	0.0-1.3	0.2	0.1	0.0-0.9
	2011-2015	7.8	3.7	2.6-5.1	3.8	3.9	2.3-6.1	4.0	3.4	2.0-5.3	0.4	0.1	0.0-0.6	0.4	0.3	0.0-1.3	0.0	0.0	0.0-0.7
	2016-2020	7.4	3.3	2.3-4.7	4.2	3.9	2.4-6.1	3.2	2.9	1.6-4.7	0.8	0.3	0.1-0.8	0.6	0.5	0.1-1.7	0.2	0.1	0.0-0.9
Non-Hodgkin lymphomas	1976-1980	26.6	12.2	10.2-14.6	13.6	14.0	10.7-18.0	13.0	10.8	8.3-13.9	14.8	7.0	5.4-8.8	7.6	7.8	5.4-10.9	7.2	6.1	4.2-8.5
	1981-1985	31.8	14.4	12.2-16.9	15.4	16.6	13.0-21.0	16.4	12.8	10.1-16.0	18.0	7.8	6.2-9.6	8.8	9.1	6.5-12.4	9.2	6.9	5.0-9.3
	1986-1990	33.0	14.9	12.7-17.4	18.2	19.9	15.8-24.6	14.8	11.6	9.0-14.7	14.8	6.9	5.4-8.6	8.6	9.7	6.9-13.1	6.2	4.6	3.1-6.6
	1991-1995	44.8	18.6	16.2-21.3	23.4	23.0	18.9-27.6	21.4	15.5	12.6-18.8	25.2	10.3	8.5-12.3	14.8	15.1	11.7-19.0	10.4	7.2	5.4-9.5
	1996-2000	47.2	19.0	16.6-21.6	24.8	22.9	19.0-27.3	22.4	16.0	13.1-19.3	24.4	9.7	8.0-11.6	13.2	12.6	9.7-16.1	11.2	7.5	5.6-9.8
	2001-2005	52.4	20.8	18.4-23.5	29.8	26.9	22.7-31.6	22.6	16.3	13.4-19.7	18.0	6.8	5.5-8.4	9.4	8.6	6.3-11.5	8.6	5.6	4.0-7.7
	2006-2010	62.4	23.5	21.0-26.4	33.4	28.7	24.5-33.5	29.0	19.6	16.5-23.3	19.2	7.1	5.7-8.7	10.6	9.1	6.8-12.0	8.6	5.4	3.9-7.4
	2011-2015	61.4	21.5	19.1-24.2	33.0	26.4	22.4-30.9	28.4	18.0	15.0-21.3	19.6	7.0	5.6-8.6	11.2	9.6	7.3-12.6	8.4	5.1	3.6-7.1
	2016-2020	62.6	21.7	19.3-24.4	33.4	25.3	21.4-29.7	29.2	18.8	15.7-22.4	17.0	5.5	4.4-6.9	9.8	7.1	5.2-9.5	7.2	4.2	2.9-6.1
Myeloma	1976-1980	9.2	4.2	3.1-5.7	3.8	4.0	2.3-6.3	5.4	4.3	2.8-6.3	7.2	3.4	2.4-4.8	3.4	3.7	2.1-6.0	3.8	3.1	1.9-4.9
	1981-1985	11.2	5.1	3.8-6.6	5.4	5.4	3.5-8.0	5.8	4.6	3.1-6.6	7.4	3.2	2.3-4.5	4.0	3.8	2.3-6.0	3.4	2.6	1.5-4.3
	1986-1990	13.0	5.5	4.2-7.1	6.4	6.8	4.5-9.9	6.6	5.0	3.4-7.1	7.6	3.1	2.2-4.3	3.8	3.9	2.3-6.1	3.8	2.7	1.6-4.4
	1991-1995	16.6	6.9	5.5-8.6	7.6	8.8	6.2-12.2	9.0	6.2	4.5-8.3	9.8	4.0	3.0-5.4	4.8	5.3	3.3-7.9	5.0	3.4	2.2-5.1
	1996-2000	13.2	5.2	4.0-6.6	5.6	5.5	3.6-8.0	7.6	5.1	3.6-7.1	8.8	3.5	2.5-4.7	3.2	3.2	1.8-5.3	5.6	3.6	2.4-5.3
	2001-2005	15.4	5.9	4.7-7.4	7.8	6.9	4.9-9.5	7.6	5.0	3.5-7.0	7.2	2.8	1.9-3.8	3.2	2.9	1.7-4.8	4.0	2.6	1.6-4.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Niagara**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	17.8	6.8	5.4-8.4	8.0	6.9	4.9-9.4	9.8	6.8	5.0-9.1	8.8	3.2	2.3-4.3	4.4	3.8	2.4-5.8	4.4	2.8	1.7-4.3
	2011-2015	16.6	5.9	4.6-7.3	10.6	8.3	6.2-11.0	6.0	3.7	2.5-5.5	7.0	2.5	1.7-3.5	3.4	2.7	1.5-4.4	3.6	2.3	1.3-3.8
	2016-2020	18.8	6.2	5.0-7.7	10.6	8.0	5.9-10.7	8.2	4.6	3.3-6.4	9.0	3.0	2.2-4.1	5.8	4.5	3.0-6.6	3.2	1.8	1.0-3.1
Leukemias	1976-1980	30.0	14.2	11.9-16.7	15.8	17.0	13.2-21.5	14.2	11.8	9.2-14.9	19.4	9.1	7.3-11.2	10.6	11.2	8.2-14.9	8.8	7.4	5.4-10.0
	1981-1985	32.2	14.4	12.2-16.9	17.8	18.5	14.6-23.0	14.4	11.4	8.9-14.5	21.4	9.8	8.0-11.9	13.2	14.0	10.6-18.0	8.2	6.9	4.9-9.4
	1986-1990	31.6	13.8	11.7-16.2	17.6	17.6	13.9-21.8	14.0	11.0	8.5-14.0	20.6	9.0	7.3-11.0	11.0	11.6	8.6-15.3	9.6	7.2	5.3-9.7
	1991-1995	34.6	14.1	12.1-16.5	18.2	17.7	14.1-21.8	16.4	11.7	9.2-14.5	18.8	7.6	6.1-9.3	10.0	10.1	7.4-13.5	8.8	6.2	4.5-8.4
	1996-2000	36.8	14.8	12.8-17.2	21.6	20.1	16.4-24.3	15.2	10.4	8.1-13.1	21.6	8.5	7.0-10.3	11.2	10.2	7.7-13.3	10.4	6.9	5.1-9.1
	2001-2005	40.4	16.2	14.0-18.6	23.4	21.5	17.8-25.9	17.0	12.2	9.7-15.2	15.6	5.9	4.7-7.4	10.2	9.4	6.9-12.3	5.4	3.5	2.3-5.2
	2006-2010	40.2	15.3	13.2-17.6	22.0	19.4	15.9-23.4	18.2	12.5	10.0-15.5	18.0	6.8	5.4-8.3	9.2	8.1	5.9-10.8	8.8	5.8	4.2-8.0
	2011-2015	41.0	15.5	13.4-17.9	24.2	20.3	16.7-24.4	16.8	11.9	9.4-14.9	19.4	6.7	5.4-8.3	11.6	10.1	7.6-13.1	7.8	4.4	3.1-6.2
2016-2020	53.4	18.9	16.6-21.4	30.6	24.1	20.3-28.5	22.8	14.6	11.9-17.9	23.4	7.6	6.3-9.2	13.0	10.2	7.8-13.1	10.4	5.9	4.3-7.9	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Oneida**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	1,038.8	404.0	392.8-415.4	493.6	461.9	443.1-481.2	545.2	377.1	362.6-392.0	526.2	204.1	196.2-212.2	268.8	256.9	242.9-271.6	257.4	173.1	163.5-183.1
	1981-1985	1,174.2	441.9	430.4-453.7	564.8	518.0	498.3-538.2	609.4	408.8	393.8-424.3	577.6	214.7	206.8-222.8	303.4	282.2	267.7-297.4	274.2	174.0	164.7-183.8
	1986-1990	1,182.2	435.5	424.2-447.0	576.2	511.0	491.8-530.7	606.0	395.8	381.2-410.7	602.4	219.5	211.6-227.6	299.6	274.1	259.8-288.9	302.8	189.0	179.3-199.2
	1991-1995	1,284.0	460.3	449.0-471.9	658.2	557.4	538.1-577.2	625.8	403.6	389.0-418.5	620.4	217.0	209.4-224.9	307.6	270.1	256.5-284.3	312.8	185.2	175.9-195.0
	1996-2000	1,390.4	500.5	488.6-512.5	696.8	581.0	561.7-600.7	693.6	451.7	436.2-467.5	581.0	199.2	192.0-206.7	304.6	257.4	244.5-270.8	276.4	161.9	153.1-171.0
	2001-2005	1,482.0	526.7	514.6-538.9	768.8	621.4	601.9-641.5	713.2	463.3	447.6-479.4	555.8	186.7	179.8-193.9	275.8	225.6	213.8-237.9	280.0	163.0	154.2-172.1
	2006-2010	1,447.0	501.7	489.9-513.6	769.0	585.6	567.0-604.6	678.0	441.4	426.0-457.1	546.2	179.2	172.4-186.2	281.2	214.5	203.3-226.1	265.0	154.2	145.6-163.1
	2011-2015	1,479.8	493.5	481.9-505.2	760.2	541.6	524.1-559.5	719.6	461.0	445.3-477.2	535.0	169.4	162.9-176.1	287.2	208.8	198.0-220.1	247.8	141.0	132.9-149.5
	2016-2020	1,521.0	494.2	482.7-505.9	809.0	554.0	536.6-571.9	712.0	451.2	435.6-467.2	520.6	159.9	153.7-166.4	272.0	188.2	178.1-198.7	248.6	138.9	130.9-147.3
Oral cavity and pharynx	1976-1980	27.4	10.5	8.8-12.5	19.2	17.1	13.7-21.0	8.2	5.7	4.0-7.8	10.6	4.0	3.0-5.3	7.4	6.6	4.6-9.2	3.2	2.1	1.2-3.5
	1981-1985	34.4	12.4	10.5-14.4	23.4	19.7	16.2-23.8	11.0	7.2	5.3-9.4	13.2	4.7	3.6-6.0	8.4	7.2	5.1-9.8	4.8	2.9	1.9-4.4
	1986-1990	23.8	9.0	7.4-10.8	15.4	13.5	10.6-17.0	8.4	5.3	3.8-7.2	9.2	3.3	2.4-4.4	6.0	5.1	3.4-7.4	3.2	2.0	1.1-3.2
	1991-1995	28.0	10.5	8.8-12.4	18.0	15.5	12.4-19.1	10.0	6.3	4.6-8.4	10.4	3.7	2.7-4.9	5.6	5.0	3.3-7.3	4.8	2.7	1.7-4.1
	1996-2000	23.8	8.8	7.3-10.6	15.4	12.9	10.1-16.1	8.4	5.6	4.0-7.6	9.4	3.3	2.4-4.4	6.0	4.9	3.3-7.0	3.4	1.9	1.1-3.1
	2001-2005	30.6	11.4	9.6-13.4	19.4	15.6	12.6-19.0	11.2	7.8	5.8-10.2	5.4	1.9	1.3-2.8	4.4	3.6	2.2-5.4	1.0	0.6	0.2-1.5
	2006-2010	31.6	11.3	9.6-13.3	23.2	17.7	14.6-21.3	8.4	5.5	3.9-7.6	4.8	1.6	1.0-2.5	3.6	2.7	1.6-4.4	1.2	0.7	0.2-1.6
	2011-2015	39.2	13.0	11.2-15.0	28.2	19.4	16.3-23.0	11.0	7.1	5.3-9.4	7.6	2.3	1.6-3.3	6.2	4.2	2.8-6.0	1.4	0.6	0.2-1.5
	2016-2020	37.0	11.9	10.2-13.8	25.8	17.4	14.4-20.8	11.2	6.8	5.1-9.1	10.0	3.1	2.2-4.1	7.4	4.9	3.4-6.9	2.6	1.5	0.8-2.7
Esophagus	1976-1980	10.0	3.5	2.6-4.7	6.0	5.1	3.3-7.4	4.0	2.5	1.5-3.9	8.8	3.0	2.2-4.1	6.2	5.0	3.3-7.2	2.6	1.6	0.8-2.7
	1981-1985	13.8	5.0	3.8-6.3	8.4	7.9	5.6-10.7	5.4	3.2	2.1-4.7	12.0	4.4	3.4-5.7	8.8	8.7	6.2-11.7	3.2	1.9	1.1-3.2
	1986-1990	13.2	4.8	3.7-6.2	9.6	7.9	5.8-10.6	3.6	2.5	1.4-4.0	11.0	4.1	3.0-5.3	8.6	7.5	5.4-10.2	2.4	1.5	0.7-2.6
	1991-1995	15.4	5.4	4.3-6.8	11.2	8.9	6.7-11.6	4.2	2.4	1.5-3.8	15.4	5.4	4.3-6.8	10.8	8.8	6.6-11.6	4.6	2.8	1.8-4.3
	1996-2000	13.4	4.8	3.7-6.1	10.0	8.3	6.2-11.0	3.4	2.0	1.1-3.3	14.2	5.0	3.9-6.4	11.0	9.2	6.9-12.0	3.2	1.7	1.0-2.9
	2001-2005	12.8	4.6	3.5-5.9	9.8	8.1	6.0-10.7	3.0	1.8	1.0-3.1	13.0	4.5	3.4-5.7	10.6	8.8	6.6-11.5	2.4	1.3	0.7-2.5
	2006-2010	15.4	5.3	4.2-6.7	12.4	9.4	7.2-12.1	3.0	1.9	1.0-3.2	12.4	4.2	3.2-5.4	10.2	7.5	5.6-9.9	2.2	1.5	0.7-2.8
	2011-2015	16.4	5.3	4.2-6.7	12.4	9.0	6.9-11.6	4.0	2.4	1.4-3.9	14.4	4.7	3.6-5.9	11.6	8.3	6.2-10.8	2.8	1.6	0.9-2.9
	2016-2020	16.8	5.3	4.2-6.6	11.8	8.0	6.0-10.4	5.0	3.1	1.9-4.7	17.8	5.4	4.3-6.7	13.0	8.9	6.8-11.4	4.8	2.7	1.7-4.1
Stomach	1976-1980	29.0	11.6	9.8-13.8	19.8	19.3	15.5-23.6	9.2	6.5	4.7-8.8	20.8	8.3	6.8-10.1	12.0	11.5	8.7-14.9	8.8	6.1	4.4-8.3
	1981-1985	25.8	9.4	7.8-11.2	15.4	13.8	10.7-17.3	10.4	6.4	4.7-8.5	18.6	6.8	5.4-8.3	12.0	10.4	7.9-13.5	6.6	4.1	2.8-5.9
	1986-1990	27.6	10.3	8.6-12.2	17.8	17.2	13.7-21.3	9.8	5.8	4.3-7.8	17.6	6.5	5.2-8.1	9.8	10.1	7.4-13.5	7.8	4.5	3.1-6.2
	1991-1995	24.0	8.3	6.9-10.0	16.0	13.8	10.9-17.3	8.0	4.6	3.3-6.4	13.6	4.7	3.6-6.0	8.6	7.3	5.3-9.9	5.0	2.8	1.8-4.2
	1996-2000	28.8	9.7	8.2-11.4	18.2	14.9	12.0-18.4	10.6	5.5	4.1-7.3	15.0	4.9	3.9-6.2	9.2	7.6	5.5-10.1	5.8	2.9	1.9-4.3
	2001-2005	23.0	7.8	6.5-9.5	13.4	11.1	8.6-14.1	9.6	5.6	4.1-7.5	11.2	3.7	2.8-4.8	4.6	3.8	2.4-5.7	6.6	3.7	2.5-5.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Oneida**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	21.6	7.4	6.1-9.0	15.0	11.3	8.9-14.2	6.6	4.3	2.9-6.1	9.4	3.2	2.3-4.2	5.4	4.2	2.7-6.1	4.0	2.6	1.6-4.1
	2011-2015	24.6	8.1	6.7-9.7	15.2	10.7	8.4-13.5	9.4	5.9	4.3-8.1	7.2	2.4	1.7-3.4	4.6	3.5	2.2-5.3	2.6	1.6	0.8-2.9
	2016-2020	20.8	6.8	5.5-8.3	11.8	8.4	6.3-10.9	9.0	5.8	4.1-7.9	5.0	1.6	1.0-2.4	1.8	1.3	0.6-2.5	3.2	1.9	1.0-3.2
Colorectal	1976-1980	190.0	72.7	68.1-77.6	90.0	84.8	76.8-93.3	100.0	64.8	59.2-70.9	88.0	34.3	31.1-37.7	41.4	41.3	35.6-47.5	46.6	30.1	26.3-34.3
	1981-1985	203.2	75.0	70.4-79.9	99.0	91.4	83.3-100.2	104.2	64.2	58.7-70.2	91.0	33.6	30.6-36.9	42.6	40.1	34.7-46.0	48.4	29.2	25.6-33.2
	1986-1990	191.0	68.8	64.4-73.4	90.2	80.6	73.1-88.7	100.8	60.7	55.4-66.4	91.8	33.6	30.5-36.8	40.4	37.9	32.7-43.7	51.4	30.7	27.0-34.8
	1991-1995	175.0	60.8	56.8-65.0	82.4	71.8	64.9-79.3	92.6	54.3	49.4-59.7	82.4	28.1	25.5-31.0	39.0	35.5	30.6-41.0	43.4	23.8	20.7-27.3
	1996-2000	166.4	57.1	53.3-61.2	80.4	67.9	61.4-74.9	86.0	48.9	44.2-54.0	65.0	21.8	19.5-24.4	31.2	26.6	22.6-31.2	33.8	18.1	15.4-21.2
	2001-2005	164.6	56.0	52.2-60.0	85.6	69.9	63.4-76.8	79.0	45.6	41.0-50.6	54.4	17.8	15.7-20.1	28.0	22.8	19.2-27.0	26.4	14.4	12.0-17.3
	2006-2010	129.4	43.6	40.2-47.2	68.4	52.8	47.3-58.8	61.0	35.9	31.8-40.5	51.0	16.3	14.3-18.5	25.6	19.8	16.5-23.5	25.4	13.3	11.0-16.1
	2011-2015	105.4	35.0	32.0-38.3	51.2	37.7	33.1-42.8	54.2	33.0	29.0-37.5	42.6	13.5	11.7-15.5	23.4	17.3	14.2-20.8	19.2	10.8	8.6-13.4
2016-2020	106.6	35.4	32.4-38.7	52.4	37.9	33.3-42.9	54.2	34.0	29.9-38.7	38.4	11.4	9.8-13.3	20.4	13.9	11.3-17.0	18.0	9.5	7.5-11.9	
Colon excluding rectum	1976-1980	139.0	53.4	49.4-57.6	61.6	58.8	52.1-66.0	77.4	50.2	45.2-55.6	71.6	27.9	25.0-31.0	31.4	31.4	26.5-36.9	40.2	25.9	22.4-29.8
	1981-1985	147.8	54.7	50.8-58.9	70.0	64.9	58.0-72.3	77.8	47.8	43.1-52.9	75.8	28.2	25.4-31.2	35.0	33.0	28.1-38.5	40.8	24.8	21.4-28.6
	1986-1990	139.6	50.5	46.7-54.4	62.0	56.5	50.2-63.4	77.6	46.8	42.1-51.9	79.8	29.1	26.2-32.1	35.2	33.0	28.1-38.5	44.6	26.5	23.0-30.3
	1991-1995	121.8	42.3	39.0-45.9	53.8	48.4	42.6-54.7	68.0	40.0	35.7-44.6	70.8	24.1	21.7-26.8	32.4	29.7	25.2-34.8	38.4	21.1	18.1-24.4
	1996-2000	121.4	41.3	38.0-44.7	55.6	47.1	41.6-53.0	65.8	36.9	32.9-41.3	55.2	18.5	16.3-20.8	25.6	21.8	18.1-26.0	29.6	15.7	13.2-18.6
	2001-2005	116.2	38.9	35.8-42.3	58.8	48.2	42.8-54.1	57.4	31.9	28.1-36.0	45.8	15.0	13.1-17.1	24.4	19.9	16.5-23.8	21.4	11.7	9.5-14.4
	2006-2010	92.4	31.1	28.2-34.1	45.0	34.9	30.5-39.8	47.4	28.1	24.4-32.1	43.8	13.9	12.1-16.0	22.8	17.6	14.5-21.2	21.0	10.8	8.7-13.3
	2011-2015	76.4	25.5	22.9-28.3	33.4	25.0	21.2-29.2	43.0	26.1	22.5-30.1	35.6	11.1	9.5-12.9	19.2	14.1	11.4-17.3	16.4	9.0	7.0-11.3
2016-2020	72.2	23.6	21.1-26.3	33.4	24.1	20.5-28.2	38.8	23.6	20.2-27.5	31.6	9.3	7.8-10.9	15.6	10.7	8.4-13.5	16.0	8.1	6.4-10.3	
Rectum & rectosigmoid	1976-1980	51.0	19.4	17.0-22.0	28.4	26.0	21.8-30.8	22.6	14.6	12.0-17.7	16.4	6.4	5.1-8.0	10.0	9.9	7.3-13.1	6.4	4.3	2.9-6.1
	1981-1985	55.4	20.3	17.9-22.9	29.0	26.5	22.2-31.4	26.4	16.4	13.7-19.6	15.2	5.5	4.3-6.9	7.6	7.1	4.9-9.8	7.6	4.4	3.1-6.1
	1986-1990	51.4	18.3	16.1-20.8	28.2	24.1	20.2-28.5	23.2	13.9	11.4-16.8	12.0	4.5	3.4-5.8	5.2	4.9	3.1-7.2	6.8	4.2	2.9-6.0
	1991-1995	53.2	18.5	16.3-20.9	28.6	23.4	19.7-27.7	24.6	14.4	11.9-17.3	11.6	4.0	3.0-5.2	6.6	5.8	3.9-8.2	5.0	2.7	1.7-4.1
	1996-2000	45.0	15.8	13.8-18.1	24.8	20.8	17.3-24.9	20.2	12.0	9.7-14.7	9.8	3.4	2.5-4.5	5.6	4.8	3.2-7.0	4.2	2.4	1.5-3.8
	2001-2005	48.4	17.1	14.9-19.4	26.8	21.6	18.1-25.7	21.6	13.7	11.2-16.7	8.6	2.8	2.0-3.8	3.6	2.9	1.7-4.6	5.0	2.7	1.7-4.1
	2006-2010	37.0	12.5	10.7-14.5	23.4	17.9	14.8-21.5	13.6	7.9	6.0-10.2	7.2	2.4	1.7-3.4	2.8	2.2	1.2-3.7	4.4	2.5	1.5-4.0
	2011-2015	29.0	9.5	8.0-11.3	17.8	12.8	10.2-15.8	11.2	6.9	5.1-9.2	7.0	2.4	1.7-3.4	4.2	3.2	1.9-4.9	2.8	1.8	1.0-3.2
2016-2020	34.4	11.9	10.1-13.9	19.0	13.7	11.0-16.9	15.4	10.4	8.1-13.3	6.8	2.2	1.5-3.1	4.8	3.2	2.0-4.8	2.0	1.4	0.6-2.7	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Oneida**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	9.4	3.5	2.6-4.7	4.8	4.2	2.6-6.4	4.6	2.9	1.9-4.5	4.2	1.6	1.0-2.5	2.0	1.9	0.9-3.5	2.2	1.4	0.7-2.6
	1981-1985	10.6	4.1	3.1-5.4	4.4	4.1	2.5-6.2	6.2	4.2	2.8-6.1	6.2	2.5	1.7-3.5	3.0	2.8	1.5-4.6	3.2	2.3	1.3-3.8
	1986-1990	9.4	3.6	2.6-4.8	5.8	5.0	3.3-7.3	3.6	2.6	1.5-4.2	5.6	2.1	1.3-3.0	4.2	3.7	2.3-5.8	1.4	1.1	0.4-2.3
	1991-1995	9.2	3.3	2.4-4.4	5.6	4.8	3.1-6.9	3.6	2.2	1.3-3.6	8.2	2.8	2.0-3.8	5.2	4.4	2.8-6.4	3.0	1.6	0.9-2.7
	1996-2000	13.0	4.6	3.6-5.9	9.2	7.5	5.5-10.0	3.8	2.3	1.4-3.7	10.4	3.6	2.7-4.8	7.6	6.4	4.5-8.8	2.8	1.4	0.8-2.5
	2001-2005	18.8	6.7	5.4-8.3	11.6	9.2	7.0-11.9	7.2	4.4	3.0-6.2	10.6	3.6	2.7-4.7	5.6	4.4	2.9-6.4	5.0	2.6	1.7-4.0
	2006-2010	16.2	5.6	4.4-7.0	11.8	8.9	6.8-11.6	4.4	2.7	1.6-4.1	14.0	4.6	3.6-5.8	8.8	6.6	4.8-8.9	5.2	2.8	1.8-4.3
	2011-2015	19.8	6.4	5.2-7.8	13.4	9.4	7.2-12.0	6.4	3.9	2.6-5.6	16.2	5.0	4.0-6.3	10.4	7.3	5.4-9.7	5.8	3.2	2.1-4.8
2016-2020	25.4	7.9	6.6-9.5	16.2	10.9	8.6-13.7	9.2	5.3	3.8-7.3	17.4	5.3	4.2-6.6	11.0	7.4	5.5-9.7	6.4	3.7	2.4-5.4	
Pancreas	1976-1980	29.0	11.5	9.7-13.6	15.8	15.7	12.3-19.6	13.2	8.8	6.8-11.2	30.8	12.0	10.1-14.1	18.4	17.3	13.9-21.4	12.4	8.3	6.4-10.8
	1981-1985	28.8	10.9	9.1-12.8	14.0	14.3	11.1-18.2	14.8	9.0	7.1-11.4	29.2	11.1	9.3-13.0	16.2	15.9	12.5-19.9	13.0	8.0	6.2-10.3
	1986-1990	29.4	10.8	9.1-12.8	11.8	10.7	8.0-13.9	17.6	10.9	8.7-13.6	27.2	10.0	8.4-11.9	10.8	9.9	7.4-13.1	16.4	10.1	7.9-12.6
	1991-1995	35.2	12.3	10.5-14.3	16.0	13.9	10.9-17.3	19.2	10.8	8.7-13.3	31.6	10.9	9.2-12.7	14.4	12.6	9.8-15.9	17.2	9.2	7.3-11.4
	1996-2000	31.4	10.9	9.3-12.8	14.8	12.5	9.8-15.7	16.6	9.5	7.6-12.0	30.6	10.4	8.8-12.2	14.4	11.9	9.3-15.0	16.2	8.8	7.0-11.1
	2001-2005	36.8	12.5	10.8-14.5	15.8	12.6	10.0-15.7	21.0	12.8	10.4-15.6	31.8	10.7	9.1-12.6	14.0	11.2	8.7-14.2	17.8	10.6	8.4-13.2
	2006-2010	42.0	14.0	12.2-16.1	19.0	14.5	11.7-17.7	23.0	13.3	10.9-16.2	33.8	11.1	9.5-13.0	15.6	11.8	9.3-14.8	18.2	10.3	8.2-12.8
	2011-2015	44.8	14.4	12.6-16.5	21.8	15.6	12.7-18.9	23.0	13.5	11.0-16.4	35.6	11.4	9.7-13.3	17.2	12.3	9.8-15.3	18.4	10.8	8.6-13.4
2016-2020	52.0	16.0	14.1-18.2	30.0	20.6	17.3-24.3	22.0	12.3	10.0-15.0	36.4	10.9	9.3-12.7	21.6	14.9	12.2-18.2	14.8	7.7	6.0-9.7	
Larynx	1976-1980	13.0	4.8	3.7-6.2	10.2	8.6	6.3-11.4	2.8	1.9	1.0-3.3	5.4	2.0	1.3-2.9	4.4	3.9	2.4-6.0	1.0	0.7	0.2-1.7
	1981-1985	16.8	6.1	4.9-7.6	13.4	11.1	8.5-14.2	3.4	2.2	1.2-3.6	4.8	1.7	1.1-2.6	3.6	3.0	1.7-4.8	1.2	0.8	0.3-1.8
	1986-1990	15.4	5.7	4.5-7.2	10.8	9.0	6.7-11.8	4.6	3.2	2.0-4.8	5.0	1.7	1.1-2.5	2.8	2.2	1.2-3.8	2.2	1.3	0.6-2.4
	1991-1995	15.0	5.6	4.4-7.0	11.0	9.4	7.0-12.2	4.0	2.9	1.7-4.5	6.0	2.1	1.4-3.1	4.6	4.2	2.6-6.4	1.4	1.0	0.4-2.1
	1996-2000	14.4	5.5	4.3-6.9	11.4	9.6	7.2-12.4	3.0	2.0	1.1-3.4	5.0	1.8	1.2-2.7	4.0	3.2	1.9-4.9	1.0	0.7	0.2-1.7
	2001-2005	16.2	5.8	4.6-7.3	11.4	9.1	6.9-11.8	4.8	3.3	2.1-5.0	4.6	1.6	1.0-2.5	3.0	2.3	1.3-3.9	1.6	1.0	0.4-2.0
	2006-2010	14.4	5.0	3.9-6.4	11.0	8.4	6.3-11.0	3.4	2.2	1.3-3.6	3.4	1.1	0.6-1.8	2.4	1.8	0.9-3.1	1.0	0.6	0.2-1.4
	2011-2015	13.6	4.4	3.4-5.7	8.4	5.6	4.0-7.7	5.2	3.4	2.2-5.1	3.8	1.2	0.7-1.9	1.8	1.3	0.6-2.5	2.0	1.2	0.6-2.3
2016-2020	16.0	5.2	4.1-6.6	12.2	8.3	6.3-10.8	3.8	2.6	1.5-4.2	5.0	1.6	1.0-2.4	4.0	2.7	1.6-4.2	1.0	0.6	0.2-1.5	
Lung and bronchus	1976-1980	135.4	52.0	48.1-56.2	97.2	87.7	79.8-96.2	38.2	27.0	23.1-31.2	110.2	41.8	38.3-45.5	78.2	70.9	63.8-78.5	32.0	21.7	18.4-25.4
	1981-1985	172.8	63.1	58.9-67.6	116.0	100.9	92.5-109.8	56.8	37.8	33.4-42.7	134.8	48.9	45.2-52.9	95.2	83.4	75.8-91.6	39.6	25.2	21.7-29.2
	1986-1990	184.4	66.6	62.3-71.2	116.6	100.0	91.8-108.7	67.8	43.4	38.7-48.5	152.4	54.5	50.6-58.6	96.8	83.6	76.1-91.6	55.6	34.9	30.8-39.5
	1991-1995	208.8	74.6	70.0-79.3	120.8	102.6	94.4-111.2	88.0	55.6	50.4-61.3	163.0	57.3	53.4-61.5	97.6	82.4	75.1-90.2	65.4	39.9	35.6-44.7
	1996-2000	226.0	80.3	75.7-85.2	134.4	111.0	102.7-119.8	91.6	59.4	53.9-65.3	160.6	55.6	51.8-59.7	93.8	78.2	71.2-85.7	66.8	40.6	36.3-45.4
	2001-2005	225.4	79.1	74.5-83.9	121.2	97.6	90.0-105.7	104.2	65.7	60.0-71.7	159.4	55.2	51.4-59.2	91.0	73.6	67.0-80.8	68.4	42.1	37.7-47.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Oneida**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	241.0	81.4	76.8-86.2	126.8	95.7	88.3-103.6	114.2	71.6	65.7-77.9	163.2	54.5	50.8-58.5	89.0	67.4	61.2-74.0	74.2	45.2	40.6-50.3
	2011-2015	242.6	78.4	73.9-83.0	123.0	87.6	80.6-94.9	119.6	71.4	65.7-77.6	154.4	49.3	45.8-53.0	86.0	61.7	55.9-67.9	68.4	39.8	35.6-44.5
	2016-2020	221.8	68.4	64.4-72.7	112.6	76.4	70.1-83.2	109.2	62.5	57.2-68.2	133.0	40.7	37.6-44.1	67.8	45.5	40.7-50.7	65.2	36.8	32.8-41.3
Melanoma of the skin	1976-1980	11.2	4.9	3.7-6.5	5.8	5.5	3.6-7.9	5.4	4.7	3.0-6.9	5.2	2.2	1.4-3.2	3.8	3.7	2.2-5.8	1.4	1.1	0.4-2.3
	1981-1985	14.8	6.0	4.7-7.6	8.0	6.7	4.7-9.2	6.8	5.3	3.6-7.5	4.4	1.7	1.0-2.6	2.4	2.1	1.0-3.7	2.0	1.3	0.6-2.5
	1986-1990	12.0	4.8	3.7-6.3	7.4	6.9	4.8-9.6	4.6	3.4	2.1-5.2	5.0	2.1	1.3-3.1	2.8	2.5	1.4-4.3	2.2	1.7	0.8-3.1
	1991-1995	16.2	6.0	4.8-7.5	9.0	7.6	5.5-10.2	7.2	5.1	3.5-7.1	7.0	2.5	1.7-3.5	3.8	3.1	1.9-4.9	3.2	1.8	1.0-3.1
	1996-2000	25.4	9.7	8.1-11.6	14.2	11.9	9.3-15.0	11.2	8.6	6.4-11.2	5.2	1.9	1.3-2.8	3.6	3.0	1.8-4.8	1.6	1.0	0.4-2.1
	2001-2005	40.4	15.0	13.0-17.3	22.6	18.2	15.0-21.9	17.8	13.3	10.6-16.5	8.8	3.1	2.2-4.2	5.8	4.8	3.2-6.9	3.0	1.8	1.0-3.2
	2006-2010	44.2	16.0	13.9-18.3	23.4	18.3	15.1-22.0	20.8	14.9	12.1-18.3	5.8	2.0	1.3-2.9	3.0	2.3	1.3-3.9	2.8	1.7	0.9-3.0
	2011-2015	49.2	17.6	15.4-20.1	28.8	21.7	18.2-25.6	20.4	14.8	11.9-18.2	9.0	3.0	2.2-4.1	6.2	4.9	3.3-7.0	2.8	1.6	0.8-2.8
2016-2020	58.2	19.7	17.4-22.2	34.0	24.1	20.5-28.1	24.2	17.3	14.1-20.9	7.8	2.4	1.7-3.3	5.4	3.6	2.4-5.4	2.4	1.3	0.6-2.5	
Female breast	1976-1980							152.8	109.1	101.2-117.4							60.0	40.8	36.2-45.8
	1981-1985							161.0	111.8	103.8-120.2							55.0	36.2	31.9-41.0
	1986-1990							175.8	120.9	112.7-129.6							58.0	37.5	33.1-42.3
	1991-1995							168.6	115.1	107.1-123.4							56.8	35.3	31.2-39.9
	1996-2000							208.8	142.3	133.5-151.5							46.6	28.5	24.8-32.7
	2001-2005							178.6	120.8	112.7-129.2							39.8	23.3	20.0-26.9
	2006-2010							175.0	117.6	109.6-126.0							34.4	19.9	16.9-23.3
	2011-2015							192.0	126.9	118.6-135.7							29.2	16.3	13.6-19.4
2016-2020							189.0	122.0	113.8-130.6							30.2	18.0	15.1-21.4	
Cervix uteri	1976-1980							18.6	13.4	10.7-16.5							8.8	6.2	4.5-8.4
	1981-1985							18.4	14.1	11.2-17.4							6.2	4.2	2.8-6.1
	1986-1990							13.6	10.2	7.8-13.0							6.0	4.1	2.7-5.9
	1991-1995							10.2	7.7	5.7-10.2							3.4	2.5	1.4-4.1
	1996-2000							12.2	10.1	7.6-13.0							3.0	2.3	1.3-3.9
	2001-2005							10.6	8.5	6.3-11.2							2.8	2.1	1.2-3.6
	2006-2010							9.4	7.5	5.5-10.0							3.8	2.7	1.6-4.3
	2011-2015							7.2	5.4	3.7-7.6							4.2	2.8	1.7-4.4
2016-2020							7.0	5.2	3.6-7.4							1.6	0.9	0.4-1.9	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Oneida**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							46.8	31.9	27.8-36.4							5.4	3.5	2.3-5.1
	1981-1985							48.4	33.9	29.5-38.7							8.6	5.1	3.7-7.0
	1986-1990							35.6	23.0	19.6-26.8							8.8	5.2	3.8-7.1
	1991-1995							37.8	26.7	22.9-30.9							9.2	5.6	4.1-7.6
	1996-2000							40.0	27.2	23.4-31.4							7.0	4.3	2.9-6.0
	2001-2005							40.4	27.0	23.4-31.2							5.8	3.4	2.2-4.9
	2006-2010							40.8	26.9	23.3-31.0							5.8	3.5	2.3-5.2
	2011-2015							50.8	32.4	28.4-36.8							6.6	3.9	2.7-5.6
	2016-2020							53.2	33.0	29.0-37.5							11.2	6.5	4.9-8.5
Ovary	1976-1980							25.6	17.9	14.8-21.4							16.6	11.3	9.0-14.2
	1981-1985							23.6	17.5	14.4-21.1							15.8	10.6	8.3-13.4
	1986-1990							26.6	18.6	15.5-22.2							16.0	10.2	8.0-12.8
	1991-1995							27.8	18.5	15.4-21.9							17.2	10.8	8.5-13.4
	1996-2000							24.8	16.8	13.8-20.1							13.2	8.1	6.2-10.4
	2001-2005							24.6	16.6	13.7-19.9							18.4	11.4	9.1-14.1
	2006-2010							20.6	13.7	11.1-16.7							13.2	7.9	6.1-10.2
	2011-2015							24.4	15.7	12.9-18.9							12.6	7.3	5.5-9.5
	2016-2020							17.2	11.0	8.6-13.8							12.4	7.6	5.8-9.9
Prostate	1976-1980				72.4	73.2	65.6-81.4							29.2	32.7	27.4-38.5			
	1981-1985				95.6	94.4	85.8-103.5							32.0	32.7	27.7-38.4			
	1986-1990				109.8	101.3	92.7-110.5							34.8	35.3	30.1-41.2			
	1991-1995				180.4	150.2	140.3-160.5							40.8	39.1	33.7-45.0			
	1996-2000				173.4	142.8	133.4-152.7							37.2	33.1	28.4-38.2			
	2001-2005				218.8	174.8	164.6-185.6							28.8	24.7	20.8-29.1			
	2006-2010				210.8	155.6	146.2-165.4							23.8	18.3	15.2-22.0			
	2011-2015				190.6	127.8	119.7-136.4							21.8	15.8	12.9-19.1			
	2016-2020				231.4	146.2	137.7-155.1							22.8	16.2	13.3-19.6			
Testis	1976-1980				2.8	2.0	1.1-3.6							0.8	0.8	0.2-2.1			
	1981-1985				4.0	3.0	1.8-4.7							0.6	0.4	0.1-1.4			
	1986-1990				6.4	4.8	3.3-6.9							0.0	0.0	0.0-0.7			
	1991-1995				8.0	5.8	4.2-8.0							0.6	0.5	0.1-1.5			
	1996-2000				5.6	4.6	3.0-6.6							0.6	0.5	0.1-1.5			
	2001-2005				7.8	6.9	4.9-9.4							0.4	0.4	0.0-1.3			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Oneida**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				7.2	6.7	4.7-9.3							0.2	0.2	0.0-1.0			
	2011-2015				7.0	6.1	4.2-8.6							0.0	0.0	0.0-0.7			
	2016-2020				7.8	7.0	4.9-9.6							0.4	0.3	0.0-1.1			
Urinary bladder (incl. in situ)	1976-1980	57.4	22.3	19.8-25.1	42.8	41.6	36.0-47.8	14.6	9.9	7.7-12.5	12.8	5.0	3.8-6.4	9.4	9.5	6.9-12.7	3.4	2.2	1.3-3.6
	1981-1985	65.0	24.1	21.5-27.0	46.4	44.0	38.3-50.3	18.6	11.6	9.3-14.4	19.2	7.1	5.7-8.6	13.8	13.6	10.5-17.4	5.4	3.1	2.1-4.7
	1986-1990	62.2	23.0	20.5-25.8	46.8	43.2	37.6-49.3	15.4	9.5	7.4-12.0	14.0	5.2	4.0-6.6	10.4	11.0	8.1-14.5	3.6	2.0	1.2-3.2
	1991-1995	60.8	21.3	18.9-23.9	46.2	40.3	35.2-46.0	14.6	8.6	6.7-11.0	13.8	4.6	3.5-5.8	9.0	8.3	6.0-11.2	4.8	2.5	1.6-3.8
	1996-2000	84.2	29.4	26.7-32.4	61.6	52.2	46.5-58.4	22.6	13.9	11.4-16.9	15.8	5.2	4.1-6.5	10.8	9.8	7.3-12.8	5.0	2.6	1.7-4.0
	2001-2005	95.8	32.4	29.6-35.5	71.2	57.9	52.0-64.3	24.6	14.6	12.1-17.6	19.6	6.2	5.0-7.6	12.8	10.7	8.2-13.7	6.8	3.5	2.4-5.0
	2006-2010	86.0	28.2	25.6-31.1	64.8	49.4	44.1-55.2	21.2	12.2	9.9-14.9	18.6	5.7	4.6-7.0	11.6	8.9	6.7-11.5	7.0	3.8	2.6-5.5
	2011-2015	88.0	27.9	25.3-30.7	67.0	48.8	43.7-54.5	21.0	11.5	9.3-14.1	18.8	5.8	4.7-7.2	12.2	9.4	7.1-12.1	6.6	3.3	2.2-4.8
2016-2020	92.4	28.5	25.9-31.3	70.8	49.4	44.3-54.9	21.6	12.0	9.8-14.7	16.8	4.9	3.9-6.2	13.4	9.4	7.3-12.1	3.4	1.7	1.0-2.9	
Kidney and renal pelvis	1976-1980	22.0	8.3	6.8-10.0	14.2	12.4	9.6-15.8	7.8	5.2	3.7-7.3	10.0	3.7	2.8-5.0	5.0	4.4	2.8-6.6	5.0	3.2	2.1-4.8
	1981-1985	26.8	10.0	8.3-11.9	15.2	13.5	10.6-17.1	11.6	7.3	5.5-9.5	11.8	4.4	3.3-5.7	7.0	5.9	4.1-8.3	4.8	3.2	2.0-4.8
	1986-1990	26.4	9.6	8.0-11.4	16.2	13.8	10.9-17.3	10.2	6.6	4.8-8.8	13.0	4.7	3.6-6.1	7.2	6.3	4.3-8.7	5.8	3.8	2.5-5.6
	1991-1995	25.0	9.1	7.6-10.9	14.6	12.2	9.6-15.4	10.4	6.8	5.0-9.0	11.4	4.0	3.0-5.2	7.2	6.4	4.4-8.9	4.2	2.4	1.5-3.8
	1996-2000	37.2	13.8	11.8-15.9	23.6	19.8	16.4-23.8	13.6	8.7	6.7-11.2	10.6	3.6	2.7-4.8	7.2	6.0	4.2-8.4	3.4	1.8	1.0-3.0
	2001-2005	45.2	16.1	14.1-18.4	28.0	22.6	19.0-26.7	17.2	10.9	8.6-13.6	12.8	4.4	3.3-5.6	7.2	5.9	4.1-8.1	5.6	3.5	2.3-5.2
	2006-2010	51.4	18.7	16.4-21.2	30.2	23.9	20.2-28.1	21.2	14.3	11.6-17.5	11.4	3.8	2.8-4.9	7.0	5.3	3.6-7.4	4.4	2.6	1.6-4.1
	2011-2015	51.2	17.0	14.9-19.3	32.8	23.2	19.7-27.2	18.4	11.9	9.5-14.8	12.4	3.7	2.8-4.8	9.2	6.5	4.7-8.7	3.2	1.8	1.0-3.1
2016-2020	61.6	20.4	18.1-23.0	39.8	27.9	24.1-32.3	21.8	13.7	11.1-16.7	9.0	2.6	1.9-3.6	6.2	4.2	2.8-6.0	2.8	1.5	0.8-2.7	
Brain and other central nervous system	1976-1980	13.2	5.1	3.9-6.6	6.8	5.9	4.0-8.4	6.4	4.4	3.0-6.3	10.2	3.9	2.9-5.2	5.4	4.6	3.0-6.8	4.8	3.4	2.1-5.1
	1981-1985	18.6	7.3	5.8-9.0	8.2	7.3	5.2-10.0	10.4	7.4	5.5-9.8	9.2	3.5	2.6-4.8	4.6	4.4	2.7-6.7	4.6	3.2	2.0-5.0
	1986-1990	18.2	6.8	5.4-8.4	9.8	8.0	5.9-10.7	8.4	5.5	3.9-7.5	11.4	4.1	3.1-5.3	6.2	5.0	3.4-7.2	5.2	3.3	2.1-4.9
	1991-1995	24.8	9.3	7.7-11.1	13.8	11.6	9.0-14.8	11.0	7.5	5.6-9.9	14.8	5.4	4.2-6.8	8.4	7.1	5.1-9.6	6.4	4.1	2.8-5.9
	1996-2000	20.2	7.8	6.4-9.5	12.4	10.5	8.0-13.5	7.8	5.6	3.9-7.7	14.6	5.5	4.3-6.9	8.6	7.1	5.1-9.5	6.0	4.3	2.8-6.2
	2001-2005	18.8	7.0	5.7-8.7	10.6	8.8	6.6-11.5	8.2	5.3	3.8-7.4	10.6	4.0	3.0-5.3	5.8	4.7	3.2-6.8	4.8	3.3	2.1-5.1
	2006-2010	18.8	7.0	5.6-8.6	10.2	8.3	6.1-10.9	8.6	5.9	4.2-8.1	13.2	4.6	3.5-5.9	8.0	6.3	4.5-8.7	5.2	3.0	2.0-4.6
	2011-2015	23.2	8.8	7.2-10.6	13.6	11.0	8.5-14.0	9.6	6.8	4.9-9.2	11.2	3.7	2.8-4.8	6.6	4.8	3.3-6.9	4.6	2.8	1.7-4.3
2016-2020	17.0	6.6	5.2-8.3	9.8	8.2	6.0-11.0	7.2	5.1	3.5-7.2	13.2	4.5	3.4-5.8	7.2	4.9	3.4-6.9	6.0	4.1	2.7-6.1	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Oneida**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	7.6	3.5	2.4-4.8	2.2	2.2	1.1-4.0	5.4	4.5	2.9-6.6	1.2	0.5	0.2-1.1	0.6	0.6	0.1-1.7	0.6	0.4	0.1-1.3
	1981-1985	8.6	3.5	2.5-4.8	2.2	1.9	0.9-3.5	6.4	5.0	3.4-7.2	2.0	0.8	0.4-1.5	1.2	1.3	0.5-2.7	0.8	0.5	0.1-1.3
	1986-1990	7.0	2.8	1.9-3.9	1.6	1.3	0.5-2.5	5.4	4.1	2.6-6.0	1.2	0.4	0.2-1.0	0.2	0.2	0.0-0.9	1.0	0.6	0.2-1.6
	1991-1995	14.8	5.7	4.5-7.2	4.6	3.8	2.4-5.8	10.2	8.0	5.9-10.6	1.6	0.6	0.2-1.1	1.0	0.9	0.3-2.1	0.6	0.4	0.1-1.3
	1996-2000	21.4	8.8	7.2-10.7	4.8	4.0	2.6-6.0	16.6	13.8	11.0-17.2	1.2	0.4	0.1-0.9	0.0	0.0	0.0-0.6	1.2	0.7	0.2-1.6
	2001-2005	31.8	13.0	11.0-15.2	6.2	5.1	3.5-7.2	25.6	21.2	17.6-25.3	1.2	0.4	0.1-0.9	0.6	0.5	0.1-1.5	0.6	0.3	0.1-1.0
	2006-2010	43.2	17.2	15.0-19.7	8.4	6.6	4.7-8.9	34.8	28.0	23.9-32.7	0.8	0.2	0.1-0.6	0.2	0.1	0.0-0.9	0.6	0.2	0.0-0.9
	2011-2015	45.4	18.1	15.8-20.8	10.8	8.3	6.2-11.0	34.6	28.1	23.9-32.9	1.2	0.4	0.1-0.9	0.6	0.5	0.1-1.4	0.6	0.4	0.1-1.3
2016-2020	47.6	19.5	17.0-22.3	14.4	11.2	8.6-14.2	33.2	28.4	24.1-33.3	1.4	0.4	0.2-0.9	0.6	0.5	0.1-1.5	0.8	0.4	0.1-1.2	
Hodgkin lymphoma	1976-1980	7.8	3.1	2.1-4.2	3.6	3.2	1.9-5.1	4.2	3.0	1.8-4.6	2.4	0.9	0.5-1.7	1.4	1.3	0.5-2.8	1.0	0.6	0.2-1.6
	1981-1985	11.6	4.4	3.3-5.7	6.0	5.1	3.4-7.4	5.6	3.9	2.6-5.7	3.6	1.6	0.9-2.5	2.2	2.4	1.2-4.4	1.4	0.9	0.4-1.9
	1986-1990	8.4	3.3	2.3-4.5	4.8	3.9	2.4-5.8	3.6	2.6	1.5-4.2	2.0	0.8	0.4-1.5	1.0	0.8	0.2-1.9	1.0	0.8	0.3-1.9
	1991-1995	10.6	4.0	3.0-5.3	5.4	4.4	2.9-6.4	5.2	3.8	2.4-5.6	1.8	0.7	0.3-1.3	0.8	0.7	0.2-1.8	1.0	0.7	0.2-1.6
	1996-2000	7.2	3.0	2.1-4.1	3.0	2.5	1.4-4.1	4.2	3.5	2.1-5.4	1.6	0.6	0.3-1.2	0.2	0.2	0.0-1.0	1.4	1.0	0.4-2.1
	2001-2005	9.2	3.9	2.8-5.2	4.2	3.5	2.1-5.3	5.0	4.4	2.8-6.5	0.8	0.3	0.1-0.8	0.8	0.7	0.2-1.7	0.0	0.0	0.0-0.6
	2006-2010	7.6	3.2	2.3-4.4	4.0	3.5	2.1-5.3	3.6	3.1	1.8-4.9	2.4	0.9	0.5-1.6	1.6	1.3	0.5-2.5	0.8	0.7	0.2-1.8
	2011-2015	6.8	2.9	2.0-4.1	3.4	2.8	1.6-4.5	3.4	3.0	1.7-4.9	1.2	0.3	0.1-0.8	1.2	0.8	0.3-1.8	0.0	0.0	0.0-0.7
2016-2020	8.4	3.6	2.6-4.9	5.0	4.0	2.6-6.0	3.4	3.2	1.8-5.1	0.8	0.3	0.1-0.9	0.6	0.5	0.1-1.5	0.2	0.1	0.0-0.8	
Non-Hodgkin lymphomas	1976-1980	29.2	11.2	9.4-13.2	13.6	11.9	9.2-15.3	15.6	10.3	8.1-12.9	16.0	6.3	5.0-7.9	7.4	6.6	4.6-9.2	8.6	5.8	4.2-7.9
	1981-1985	30.6	11.7	9.9-13.8	15.4	13.9	10.8-17.5	15.2	10.5	8.2-13.3	16.6	6.2	4.9-7.7	8.6	8.0	5.7-11.0	8.0	4.9	3.5-6.7
	1986-1990	38.4	14.2	12.2-16.4	19.4	16.3	13.1-20.0	19.0	12.2	9.8-15.0	19.6	6.9	5.6-8.5	10.0	8.7	6.4-11.6	9.6	5.7	4.2-7.6
	1991-1995	49.0	17.9	15.7-20.3	26.4	22.0	18.4-26.2	22.6	14.1	11.6-17.1	27.0	9.6	8.0-11.4	12.2	10.6	8.1-13.7	14.8	8.9	6.9-11.3
	1996-2000	58.0	21.0	18.7-23.6	29.2	24.7	20.8-29.1	28.8	17.6	14.8-20.9	23.2	7.8	6.4-9.4	12.2	10.4	7.9-13.4	11.0	5.9	4.4-7.8
	2001-2005	63.0	22.4	20.0-25.1	31.6	25.6	21.8-30.0	31.4	19.4	16.4-22.9	19.8	6.3	5.1-7.7	8.8	7.2	5.2-9.6	11.0	5.6	4.2-7.5
	2006-2010	62.6	21.8	19.4-24.4	34.6	26.9	23.0-31.3	28.0	17.5	14.6-20.8	23.0	7.6	6.3-9.2	11.6	8.9	6.8-11.6	11.4	6.6	4.9-8.7
	2011-2015	67.6	22.8	20.3-25.5	39.2	29.2	25.2-33.8	28.4	17.3	14.4-20.7	19.4	6.0	4.8-7.4	10.6	7.9	5.9-10.4	8.8	4.5	3.2-6.2
2016-2020	67.2	22.2	19.8-24.9	34.6	24.9	21.2-29.0	32.6	20.4	17.2-24.0	21.8	6.9	5.7-8.5	12.0	9.0	6.8-11.6	9.8	5.2	3.8-7.1	
Myeloma	1976-1980	9.0	3.4	2.5-4.6	3.4	2.9	1.7-4.8	5.6	3.7	2.5-5.4	6.6	2.5	1.7-3.6	3.4	3.1	1.7-5.0	3.2	2.2	1.2-3.6
	1981-1985	13.0	4.9	3.7-6.2	7.2	6.8	4.7-9.6	5.8	3.7	2.5-5.4	6.8	2.6	1.8-3.6	3.0	3.0	1.6-4.9	3.8	2.4	1.4-3.8
	1986-1990	15.0	5.3	4.1-6.7	6.6	5.3	3.6-7.6	8.4	5.3	3.7-7.2	12.2	4.5	3.4-5.8	6.4	5.9	4.0-8.5	5.8	3.8	2.5-5.6
	1991-1995	13.0	4.6	3.5-5.8	6.6	5.4	3.7-7.7	6.4	3.7	2.5-5.3	8.8	3.1	2.2-4.1	3.8	3.1	1.9-4.9	5.0	2.8	1.8-4.2
	1996-2000	16.2	5.6	4.5-7.1	8.4	7.0	5.0-9.5	7.8	4.8	3.3-6.6	9.4	3.1	2.3-4.2	5.8	4.8	3.2-6.9	3.6	1.9	1.1-3.2
	2001-2005	18.8	6.2	5.0-7.7	8.4	6.8	4.9-9.2	10.4	6.0	4.4-7.9	10.2	3.3	2.5-4.4	6.4	5.2	3.6-7.4	3.8	2.1	1.2-3.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Oneida**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	20.0	6.8	5.5-8.4	11.0	8.4	6.3-11.0	9.0	5.6	4.0-7.6	8.8	2.7	2.0-3.7	4.2	3.1	1.9-4.9	4.6	2.4	1.5-3.7
	2011-2015	21.6	6.8	5.5-8.2	12.0	8.3	6.3-10.8	9.6	5.4	3.9-7.3	8.8	2.7	2.0-3.7	4.8	3.4	2.2-5.2	4.0	2.0	1.2-3.3
	2016-2020	17.4	5.7	4.5-7.1	11.2	8.2	6.1-10.8	6.2	3.8	2.5-5.4	9.0	2.8	2.0-3.8	5.4	3.9	2.6-5.8	3.6	2.0	1.1-3.3
Leukemias	1976-1980	31.0	12.0	10.1-14.1	17.6	16.6	13.2-20.6	13.4	9.2	7.1-11.8	19.2	7.3	5.9-9.0	10.6	10.4	7.7-13.8	8.6	5.8	4.1-7.8
	1981-1985	33.4	12.9	10.9-15.0	18.4	17.3	13.8-21.3	15.0	9.9	7.7-12.5	22.0	8.4	6.8-10.1	11.8	11.3	8.5-14.7	10.2	6.6	4.9-8.8
	1986-1990	37.8	14.1	12.1-16.3	23.0	19.7	16.2-23.8	14.8	9.4	7.3-11.9	20.4	7.7	6.2-9.4	11.6	10.2	7.7-13.3	8.8	5.7	4.1-7.7
	1991-1995	38.6	13.8	11.9-15.9	20.4	17.5	14.2-21.3	18.2	10.7	8.5-13.2	24.8	8.7	7.2-10.4	12.8	11.1	8.5-14.2	12.0	6.9	5.3-9.0
	1996-2000	46.8	16.7	14.6-19.0	25.2	21.5	17.9-25.7	21.6	13.3	10.8-16.2	21.4	7.4	6.1-9.0	12.0	10.0	7.6-12.9	9.4	5.5	4.0-7.4
	2001-2005	43.2	15.7	13.6-18.0	25.0	20.7	17.2-24.6	18.2	11.7	9.3-14.5	20.8	6.9	5.6-8.3	11.0	9.0	6.8-11.7	9.8	5.4	4.0-7.3
	2006-2010	41.0	14.4	12.5-16.6	25.4	19.9	16.5-23.7	15.6	10.1	7.8-12.7	21.6	7.2	5.9-8.8	12.4	9.7	7.5-12.5	9.2	5.2	3.8-7.2
	2011-2015	54.4	18.5	16.3-21.0	34.6	25.4	21.7-29.6	19.8	12.7	10.2-15.6	25.0	8.0	6.6-9.6	15.8	12.0	9.5-15.1	9.2	4.9	3.5-6.7
2016-2020	48.6	16.2	14.2-18.5	27.4	19.7	16.5-23.5	21.2	13.5	10.8-16.5	22.2	6.9	5.6-8.4	11.0	7.8	5.9-10.3	11.2	6.2	4.6-8.3	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Onondaga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	1,714.4	421.4	412.4-430.6	826.2	498.2	482.3-514.5	888.2	381.9	370.6-393.6	910.6	227.8	221.1-234.6	477.0	297.6	285.1-310.5	433.6	185.7	177.9-193.8
	1981-1985	1,898.8	443.0	434.0-452.2	934.8	531.3	515.4-547.5	964.0	393.5	382.3-405.1	940.8	221.0	214.6-227.5	489.0	289.0	277.0-301.3	451.8	180.4	173.0-188.2
	1986-1990	1,914.6	428.2	419.5-437.0	932.8	508.9	493.7-524.3	981.8	386.8	375.9-398.1	1,002.6	225.2	218.9-231.6	514.8	294.6	282.8-306.7	487.8	184.2	176.8-191.7
	1991-1995	2,199.6	469.8	461.0-478.7	1,092.2	562.1	546.8-577.6	1,107.4	418.0	406.9-429.4	1,018.0	217.4	211.4-223.5	505.6	276.0	264.9-287.3	512.4	184.1	177.0-191.5
	1996-2000	2,332.4	491.3	482.4-500.3	1,152.6	574.0	559.0-589.2	1,179.8	440.2	428.9-451.7	1,008.2	209.4	203.6-215.2	501.8	260.4	250.2-271.0	506.4	179.1	172.1-186.3
	2001-2005	2,569.6	522.7	513.6-531.8	1,324.2	625.4	610.3-640.7	1,245.4	451.4	440.1-462.9	983.8	195.9	190.4-201.4	492.4	241.4	231.8-251.2	491.4	167.1	160.5-174.0
	2006-2010	2,830.0	543.8	534.8-553.0	1,494.6	649.6	634.8-664.7	1,335.4	465.9	454.6-477.5	984.8	183.8	178.6-189.1	497.8	222.7	214.0-231.8	487.0	157.5	151.2-164.1
	2011-2015	2,881.2	516.3	507.6-525.0	1,458.8	575.7	562.3-589.5	1,422.4	473.1	461.8-484.6	1,003.6	175.3	170.4-180.4	507.6	207.8	199.6-216.2	496.0	153.0	146.9-159.4
	2016-2020	2,946.2	498.4	490.1-506.8	1,527.4	556.2	543.4-569.3	1,418.8	455.6	444.5-466.8	895.2	147.4	143.0-151.9	450.8	171.8	164.6-179.3	444.4	130.6	125.0-136.3
Oral cavity and pharynx	1976-1980	46.4	10.9	9.5-12.5	30.2	15.9	13.4-18.8	16.2	6.8	5.4-8.5	18.6	4.4	3.6-5.5	14.6	8.0	6.3-10.2	4.0	1.6	1.0-2.5
	1981-1985	51.6	11.4	10.0-12.9	35.6	17.9	15.3-20.9	16.0	6.2	4.9-7.7	15.0	3.4	2.7-4.3	10.0	5.1	3.7-6.8	5.0	2.0	1.3-2.9
	1986-1990	45.4	10.4	9.1-11.9	31.8	16.7	14.2-19.6	13.6	5.6	4.3-7.1	17.6	3.9	3.1-4.8	11.0	5.8	4.3-7.6	6.6	2.5	1.7-3.6
	1991-1995	41.2	9.0	7.8-10.4	27.4	13.7	11.5-16.3	13.8	5.5	4.3-7.0	11.2	2.5	1.9-3.2	8.4	4.3	3.1-5.9	2.8	1.1	0.6-1.8
	1996-2000	49.6	10.5	9.3-11.9	29.8	14.5	12.3-17.1	19.8	7.4	6.0-9.0	13.0	2.7	2.1-3.5	8.4	4.3	3.1-5.9	4.6	1.7	1.1-2.5
	2001-2005	52.6	10.7	9.4-12.1	34.0	15.5	13.2-18.0	18.6	6.6	5.3-8.2	11.2	2.2	1.7-2.9	7.0	3.2	2.2-4.5	4.2	1.4	0.9-2.2
	2006-2010	63.8	12.2	10.9-13.6	41.2	17.1	14.8-19.7	22.6	7.9	6.5-9.5	12.4	2.2	1.7-2.9	8.0	3.3	2.4-4.6	4.4	1.3	0.8-2.0
	2011-2015	68.6	12.3	11.0-13.7	45.4	17.9	15.6-20.5	23.2	7.4	6.1-8.9	11.0	1.9	1.4-2.5	5.8	2.3	1.5-3.4	5.2	1.5	1.0-2.3
	2016-2020	77.8	13.1	11.8-14.5	51.4	18.5	16.2-21.0	26.4	8.5	7.0-10.2	12.6	2.0	1.5-2.5	7.4	2.6	1.8-3.6	5.2	1.5	0.9-2.2
Esophagus	1976-1980	16.8	4.1	3.2-5.1	11.0	6.6	4.9-8.7	5.8	2.4	1.6-3.4	16.0	3.9	3.1-4.9	10.6	6.3	4.7-8.4	5.4	2.3	1.5-3.4
	1981-1985	17.0	3.9	3.1-4.8	11.0	6.2	4.6-8.1	6.0	2.3	1.5-3.3	14.8	3.4	2.6-4.3	9.6	5.6	4.0-7.4	5.2	1.9	1.3-2.8
	1986-1990	21.2	4.8	3.9-5.8	15.2	8.1	6.3-10.1	6.0	2.2	1.5-3.2	19.0	4.1	3.3-5.1	13.8	7.2	5.6-9.2	5.2	1.9	1.2-2.7
	1991-1995	26.2	5.6	4.7-6.7	18.0	9.1	7.3-11.3	8.2	3.0	2.1-4.1	23.0	4.9	4.0-5.9	16.8	8.7	6.9-10.8	6.2	2.2	1.5-3.1
	1996-2000	28.8	6.1	5.1-7.1	20.4	10.0	8.1-12.1	8.4	3.0	2.1-4.0	25.4	5.3	4.4-6.3	18.4	9.1	7.3-11.1	7.0	2.5	1.7-3.4
	2001-2005	29.0	5.8	4.9-6.9	23.2	10.8	8.9-13.0	5.8	1.8	1.2-2.7	28.0	5.6	4.7-6.6	20.6	9.7	7.9-11.8	7.4	2.3	1.6-3.2
	2006-2010	28.6	5.4	4.5-6.4	22.0	9.6	7.8-11.5	6.6	2.2	1.5-3.1	27.0	5.0	4.2-6.0	22.8	9.9	8.2-12.0	4.2	1.3	0.8-2.0
	2011-2015	25.6	4.4	3.7-5.3	20.0	7.9	6.4-9.6	5.6	1.8	1.1-2.6	30.6	5.3	4.5-6.3	23.4	9.3	7.7-11.2	7.2	2.2	1.5-3.1
	2016-2020	26.4	4.3	3.6-5.2	20.8	7.6	6.2-9.3	5.6	1.6	1.1-2.4	25.0	4.0	3.3-4.8	20.4	7.4	6.0-9.1	4.6	1.3	0.8-2.0
Stomach	1976-1980	39.4	10.0	8.6-11.5	23.4	14.2	11.6-17.2	16.0	7.1	5.6-8.8	30.0	7.7	6.5-9.1	18.6	11.9	9.5-14.8	11.4	5.1	3.8-6.6
	1981-1985	45.4	10.9	9.5-12.5	27.4	16.7	13.9-19.9	18.0	7.1	5.7-8.8	30.2	7.4	6.3-8.7	18.4	11.4	9.1-14.1	11.8	4.8	3.7-6.2
	1986-1990	35.4	8.0	6.8-9.2	23.0	12.6	10.4-15.3	12.4	4.6	3.5-6.0	24.8	5.6	4.6-6.7	16.6	9.5	7.5-11.8	8.2	3.0	2.2-4.2
	1991-1995	36.4	7.8	6.7-9.1	22.8	12.3	10.1-14.8	13.6	4.6	3.6-5.9	28.6	6.2	5.2-7.3	19.0	10.1	8.1-12.4	9.6	3.2	2.4-4.3
	1996-2000	35.8	7.5	6.4-8.7	19.6	10.0	8.1-12.2	16.2	5.7	4.5-7.1	20.0	4.1	3.4-5.0	10.6	5.5	4.1-7.2	9.4	3.2	2.4-4.3
	2001-2005	33.6	6.7	5.8-7.9	18.6	8.9	7.1-10.9	15.0	5.3	4.1-6.6	18.2	3.6	2.9-4.4	9.8	4.8	3.6-6.4	8.4	2.8	2.0-3.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Onondaga**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	33.4	6.3	5.4-7.4	22.6	9.9	8.2-12.0	10.8	3.3	2.5-4.4	13.0	2.5	1.9-3.2	8.8	3.9	2.8-5.3	4.2	1.3	0.8-2.0
	2011-2015	43.8	7.8	6.8-9.0	27.4	11.1	9.3-13.2	16.4	5.2	4.1-6.5	14.8	2.7	2.1-3.4	7.4	3.0	2.1-4.2	7.4	2.4	1.7-3.3
	2016-2020	46.4	8.0	7.0-9.1	30.6	11.6	9.7-13.6	15.8	5.2	4.1-6.6	14.6	2.5	1.9-3.2	8.2	3.2	2.3-4.4	6.4	2.0	1.3-2.9
Colorectal	1976-1980	264.4	66.6	63.0-70.4	128.6	80.8	74.3-87.6	135.8	57.5	53.3-62.1	126.4	32.9	30.3-35.6	60.2	40.4	35.7-45.5	66.2	28.5	25.5-31.8
	1981-1985	285.8	67.5	64.0-71.1	141.6	82.8	76.5-89.4	144.2	56.2	52.2-60.5	127.0	30.4	28.0-32.9	61.8	37.1	32.9-41.7	65.2	25.5	22.8-28.4
	1986-1990	261.0	58.4	55.2-61.7	123.2	69.5	63.9-75.5	137.8	50.9	47.1-54.9	123.8	28.2	26.0-30.5	60.4	35.8	31.7-40.3	63.4	23.2	20.7-25.9
	1991-1995	263.6	55.9	52.9-59.1	128.2	68.6	63.2-74.2	135.4	47.4	43.9-51.2	120.6	25.7	23.6-27.8	53.4	30.1	26.5-34.1	67.2	22.7	20.4-25.4
	1996-2000	261.6	54.3	51.4-57.4	124.2	63.4	58.4-68.7	137.4	47.8	44.2-51.6	110.0	22.7	20.8-24.7	51.6	27.1	23.8-30.7	58.4	19.6	17.4-22.1
	2001-2005	263.2	52.6	49.7-55.5	128.6	61.6	56.9-66.6	134.6	46.2	42.8-49.9	96.6	19.0	17.3-20.7	45.2	22.5	19.6-25.7	51.4	16.9	14.9-19.1
	2006-2010	229.0	43.3	40.8-45.9	107.4	47.2	43.2-51.4	121.6	39.8	36.7-43.3	81.2	15.0	13.6-16.6	36.4	16.5	14.2-19.1	44.8	13.9	12.1-15.9
	2011-2015	224.0	40.2	37.8-42.7	111.8	45.9	42.0-49.9	112.2	35.5	32.5-38.7	80.6	14.0	12.6-15.5	42.4	17.9	15.6-20.6	38.2	11.0	9.4-12.7
2016-2020	216.4	37.4	35.1-39.8	107.4	41.2	37.6-44.9	109.0	34.2	31.2-37.4	70.4	11.8	10.6-13.2	35.2	13.9	11.9-16.2	35.2	10.3	8.8-12.0	
Colon excluding rectum	1976-1980	185.8	47.2	44.1-50.3	85.4	54.2	48.9-59.9	100.4	42.8	39.1-46.7	103.4	27.0	24.7-29.4	48.0	32.6	28.4-37.3	55.4	23.9	21.2-26.9
	1981-1985	204.8	48.7	45.7-51.8	96.4	57.7	52.4-63.3	108.4	42.1	38.6-45.8	103.0	24.7	22.6-27.0	49.2	29.7	26.0-33.8	53.8	21.1	18.6-23.8
	1986-1990	192.4	43.3	40.5-46.1	84.2	48.9	44.1-54.1	108.2	39.9	36.6-43.4	99.0	22.4	20.5-24.5	46.2	27.3	23.8-31.2	52.8	19.3	17.0-21.8
	1991-1995	190.4	40.4	37.8-43.0	91.6	49.5	44.9-54.4	98.8	34.2	31.2-37.3	99.8	21.2	19.4-23.2	43.4	24.6	21.3-28.2	56.4	19.0	16.9-21.4
	1996-2000	194.4	40.2	37.7-42.8	90.8	46.7	42.5-51.3	103.6	35.7	32.6-38.9	88.8	18.3	16.6-20.1	40.4	21.2	18.3-24.4	48.4	16.2	14.2-18.4
	2001-2005	192.2	38.2	35.9-40.7	89.6	43.3	39.4-47.5	102.6	34.8	31.8-38.0	75.2	14.7	13.3-16.3	34.6	17.5	15.0-20.3	40.6	13.3	11.5-15.3
	2006-2010	170.0	32.0	29.8-34.2	75.0	33.2	29.9-36.8	95.0	30.8	28.0-33.8	63.4	11.8	10.5-13.2	27.2	12.5	10.5-14.8	36.2	11.2	9.6-13.0
	2011-2015	153.8	27.5	25.5-29.5	72.0	30.2	27.1-33.6	81.8	25.3	22.8-27.9	63.8	11.0	9.8-12.4	31.8	13.6	11.5-15.9	32.0	9.1	7.7-10.7
2016-2020	148.2	25.7	23.8-27.7	70.2	27.2	24.3-30.3	78.0	24.4	21.9-27.1	52.6	8.6	7.6-9.8	25.6	10.2	8.4-12.1	27.0	7.5	6.3-9.0	
Rectum & rectosigmoid	1976-1980	78.6	19.5	17.6-21.5	43.2	26.6	23.0-30.6	35.4	14.8	12.7-17.1	23.0	5.9	4.9-7.1	12.2	7.8	5.9-10.1	10.8	4.6	3.4-6.0
	1981-1985	81.0	18.8	17.0-20.7	45.2	25.1	21.8-28.7	35.8	14.1	12.1-16.4	24.0	5.7	4.7-6.8	12.6	7.4	5.6-9.5	11.4	4.4	3.3-5.7
	1986-1990	68.6	15.1	13.5-16.8	39.0	20.6	17.7-23.8	29.6	11.0	9.3-13.0	24.8	5.7	4.7-6.8	14.2	8.5	6.6-10.8	10.6	3.9	2.9-5.1
	1991-1995	73.2	15.6	14.0-17.3	36.6	19.1	16.4-22.1	36.6	13.3	11.4-15.4	20.8	4.4	3.6-5.4	10.0	5.5	4.0-7.3	10.8	3.7	2.8-4.9
	1996-2000	67.2	14.1	12.7-15.7	33.4	16.7	14.2-19.4	33.8	12.1	10.3-14.1	21.2	4.4	3.6-5.3	11.2	5.9	4.4-7.7	10.0	3.5	2.6-4.6
	2001-2005	71.0	14.3	12.9-15.9	39.0	18.3	15.8-21.1	32.0	11.5	9.7-13.4	21.4	4.2	3.5-5.1	10.6	5.0	3.7-6.6	10.8	3.6	2.7-4.7
	2006-2010	59.0	11.3	10.0-12.7	32.4	13.9	11.9-16.3	26.6	9.1	7.6-10.8	17.8	3.3	2.6-4.1	9.2	4.0	2.9-5.4	8.6	2.7	1.9-3.6
	2011-2015	70.2	12.7	11.4-14.2	39.8	15.7	13.5-18.1	30.4	10.3	8.6-12.1	16.8	2.9	2.3-3.7	10.6	4.4	3.2-5.7	6.2	1.9	1.2-2.7
2016-2020	68.2	11.7	10.4-13.1	37.2	14.0	12.0-16.2	31.0	9.8	8.2-11.6	17.8	3.2	2.5-3.9	9.6	3.7	2.7-5.0	8.2	2.8	1.9-3.8	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Onondaga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	8.6	2.1	1.5-2.8	4.4	2.6	1.6-4.1	4.2	1.8	1.1-2.7	7.8	1.8	1.3-2.5	4.8	2.7	1.7-4.0	3.0	1.3	0.7-2.1
	1981-1985	9.6	2.3	1.7-3.0	6.0	3.6	2.4-5.1	3.6	1.4	0.8-2.3	8.0	1.9	1.3-2.6	5.6	3.3	2.2-4.9	2.4	0.9	0.5-1.6
	1986-1990	13.8	3.0	2.3-3.8	9.4	4.9	3.6-6.6	4.4	1.6	1.0-2.5	12.6	2.7	2.1-3.5	8.6	4.5	3.2-6.1	4.0	1.4	0.9-2.3
	1991-1995	17.2	3.7	2.9-4.5	9.2	4.6	3.3-6.1	8.0	2.8	2.0-3.9	14.0	3.0	2.3-3.8	8.6	4.4	3.1-5.9	5.4	1.9	1.2-2.8
	1996-2000	22.4	4.7	3.9-5.6	14.4	7.2	5.6-9.0	8.0	2.8	2.0-3.9	16.2	3.4	2.7-4.2	11.0	5.5	4.1-7.2	5.2	1.8	1.2-2.6
	2001-2005	22.0	4.4	3.6-5.3	16.0	7.3	5.8-9.1	6.0	2.1	1.4-3.0	17.2	3.5	2.8-4.3	11.8	5.5	4.1-7.0	5.4	1.9	1.2-2.7
	2006-2010	35.2	6.6	5.6-7.7	25.4	10.6	8.8-12.6	9.8	3.5	2.6-4.6	26.6	5.0	4.2-6.0	17.6	7.6	6.1-9.4	9.0	3.1	2.2-4.1
	2011-2015	48.0	8.1	7.1-9.2	36.0	13.0	11.1-15.1	12.0	3.8	2.9-5.0	38.8	6.7	5.8-7.8	28.0	10.8	9.0-12.8	10.8	3.4	2.6-4.5
	2016-2020	55.8	8.8	7.8-9.9	41.0	14.2	12.2-16.3	14.8	4.4	3.4-5.6	31.6	5.1	4.3-6.0	21.6	7.5	6.1-9.2	10.0	3.0	2.2-4.0
Pancreas	1976-1980	44.6	11.2	9.7-12.7	22.6	13.9	11.3-16.8	22.0	9.2	7.6-11.1	42.8	10.8	9.4-12.3	21.0	13.2	10.6-16.1	21.8	9.1	7.5-11.0
	1981-1985	51.4	11.8	10.4-13.4	26.6	14.7	12.2-17.6	24.8	9.8	8.1-11.7	48.0	11.0	9.6-12.5	26.2	14.7	12.2-17.7	21.8	8.6	7.1-10.5
	1986-1990	50.0	11.2	9.8-12.7	23.0	12.6	10.3-15.2	27.0	9.9	8.3-11.7	50.6	11.2	9.9-12.7	22.6	12.4	10.1-15.0	28.0	10.1	8.5-12.0
	1991-1995	56.6	12.0	10.7-13.5	24.6	12.9	10.6-15.4	32.0	11.0	9.4-12.9	57.0	12.1	10.7-13.6	23.8	12.5	10.3-15.1	33.2	11.5	9.8-13.4
	1996-2000	64.8	13.5	12.1-15.0	28.6	14.8	12.5-17.5	36.2	12.7	10.9-14.8	57.8	12.0	10.6-13.4	23.4	12.4	10.2-14.9	34.4	11.9	10.2-13.9
	2001-2005	66.8	13.3	11.9-14.8	30.8	14.9	12.6-17.5	36.0	12.2	10.4-14.1	56.8	11.3	10.0-12.7	26.4	12.9	10.7-15.3	30.4	10.1	8.6-11.9
	2006-2010	77.2	14.6	13.2-16.2	37.4	16.7	14.3-19.3	39.8	12.8	11.0-14.8	68.4	12.6	11.3-14.0	32.4	14.4	12.3-16.8	36.0	11.1	9.5-12.9
	2011-2015	87.4	15.4	13.9-16.9	43.0	17.4	15.1-19.9	44.4	13.6	11.9-15.6	68.6	11.9	10.7-13.3	34.8	14.1	12.0-16.4	33.8	10.1	8.6-11.8
	2016-2020	88.0	14.5	13.2-16.0	44.4	16.3	14.2-18.7	43.6	13.0	11.3-15.0	72.2	11.7	10.5-13.0	34.6	12.9	11.0-15.0	37.6	10.7	9.2-12.4
Larynx	1976-1980	33.0	7.9	6.7-9.2	26.8	15.2	12.6-18.1	6.2	2.7	1.8-3.9	10.8	2.6	1.9-3.4	8.4	4.7	3.3-6.4	2.4	1.1	0.6-1.9
	1981-1985	30.4	6.9	5.8-8.1	22.8	12.0	9.8-14.5	7.6	3.0	2.1-4.1	7.2	1.6	1.1-2.2	5.2	2.8	1.8-4.1	2.0	0.7	0.4-1.4
	1986-1990	26.6	6.1	5.1-7.3	21.4	11.2	9.1-13.6	5.2	2.3	1.5-3.4	6.6	1.5	1.0-2.1	5.0	2.6	1.7-3.9	1.6	0.6	0.3-1.2
	1991-1995	25.4	5.5	4.6-6.6	18.8	9.6	7.7-11.7	6.6	2.5	1.7-3.6	6.4	1.4	0.9-1.9	5.4	2.9	1.9-4.3	1.0	0.4	0.1-1.0
	1996-2000	24.4	5.2	4.3-6.2	18.8	9.1	7.4-11.2	5.6	2.2	1.5-3.2	8.0	1.7	1.2-2.4	6.0	3.0	2.0-4.3	2.0	0.8	0.4-1.5
	2001-2005	21.0	4.3	3.5-5.2	17.8	8.2	6.6-10.1	3.2	1.2	0.7-2.0	5.6	1.1	0.8-1.7	4.8	2.4	1.5-3.6	0.8	0.3	0.1-0.8
	2006-2010	19.6	3.8	3.1-4.6	15.2	6.7	5.2-8.4	4.4	1.5	0.9-2.3	4.4	0.8	0.5-1.3	3.6	1.6	0.9-2.5	0.8	0.2	0.1-0.7
	2011-2015	24.6	4.3	3.5-5.1	19.6	7.5	6.1-9.2	5.0	1.5	1.0-2.3	6.4	1.1	0.7-1.6	5.8	2.2	1.5-3.2	0.6	0.2	0.0-0.6
	2016-2020	22.0	3.6	2.9-4.3	18.2	6.5	5.2-8.0	3.8	1.2	0.7-1.9	6.0	0.9	0.6-1.3	5.0	1.8	1.1-2.6	1.0	0.3	0.1-0.7
Lung and bronchus	1976-1980	265.4	63.9	60.4-67.5	186.8	108.5	101.3-116.0	78.6	33.7	30.4-37.2	219.2	52.9	49.7-56.2	157.6	92.3	85.7-99.3	61.6	26.2	23.3-29.4
	1981-1985	305.8	70.2	66.6-73.8	204.6	114.9	107.6-122.4	101.2	41.0	37.5-44.9	247.0	56.4	53.3-59.7	164.4	92.0	85.5-98.7	82.6	33.0	29.8-36.4
	1986-1990	323.0	71.8	68.3-75.4	204.8	111.6	104.6-118.9	118.2	45.4	41.8-49.3	273.6	60.6	57.3-63.9	176.4	97.7	91.1-104.6	97.2	36.9	33.7-40.4
	1991-1995	360.0	76.7	73.1-80.3	199.4	103.3	96.8-110.0	160.6	59.4	55.3-63.7	290.8	61.8	58.6-65.1	169.8	89.6	83.5-96.0	121.0	44.0	40.5-47.7
	1996-2000	387.6	81.0	77.4-84.7	207.0	103.3	97.0-109.9	180.6	66.4	62.1-70.9	287.8	59.8	56.8-63.0	160.8	80.7	75.2-86.6	127.0	45.5	42.0-49.3
	2001-2005	415.6	83.8	80.2-87.5	213.0	101.3	95.2-107.6	202.6	71.9	67.5-76.5	304.0	61.1	58.0-64.2	162.2	78.2	72.9-83.8	141.8	49.5	45.8-53.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Onondaga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	435.4	82.9	79.4-86.5	217.4	95.5	89.8-101.4	218.0	74.4	70.0-79.1	289.4	54.9	52.1-57.8	151.4	67.2	62.4-72.2	138.0	46.2	42.7-49.8
	2011-2015	436.2	76.7	73.5-80.1	216.8	86.5	81.3-91.9	219.4	69.7	65.6-74.1	277.6	48.7	46.1-51.4	144.2	58.1	53.9-62.6	133.4	42.1	38.8-45.5
	2016-2020	440.8	71.7	68.6-74.8	210.4	78.0	73.2-83.0	230.4	67.5	63.6-71.6	234.6	38.3	36.1-40.6	117.2	44.3	40.7-48.1	117.4	34.1	31.4-37.1
Melanoma of the skin	1976-1980	24.4	5.9	4.9-7.1	13.2	7.0	5.3-8.9	11.2	5.2	3.9-6.8	8.0	1.9	1.4-2.6	4.0	2.1	1.3-3.3	4.0	1.8	1.1-2.8
	1981-1985	38.2	9.0	7.8-10.5	17.6	9.4	7.4-11.7	20.6	8.8	7.2-10.8	7.4	1.8	1.3-2.5	3.6	2.0	1.2-3.2	3.8	1.7	1.0-2.6
	1986-1990	29.0	6.4	5.4-7.5	15.2	7.7	6.0-9.7	13.8	5.6	4.3-7.1	12.0	2.7	2.0-3.4	7.8	4.2	2.9-5.8	4.2	1.6	1.0-2.4
	1991-1995	31.4	6.8	5.8-7.9	18.8	9.1	7.3-11.1	12.6	5.0	3.8-6.4	9.8	2.0	1.5-2.7	6.2	3.1	2.1-4.4	3.6	1.3	0.8-2.1
	1996-2000	42.0	9.0	7.8-10.3	24.4	11.9	9.9-14.2	17.6	6.9	5.6-8.6	12.6	2.6	2.0-3.4	8.8	4.2	3.0-5.6	3.8	1.4	0.9-2.2
	2001-2005	63.2	13.1	11.7-14.6	36.4	17.4	14.9-20.1	26.8	10.4	8.7-12.4	14.8	3.0	2.4-3.8	9.0	4.4	3.2-5.8	5.8	2.1	1.4-3.0
	2006-2010	92.2	18.6	16.9-20.4	47.6	21.1	18.5-24.0	44.6	17.1	14.9-19.5	11.8	2.3	1.7-3.0	7.0	3.2	2.2-4.5	4.8	1.6	1.0-2.5
	2011-2015	104.2	19.4	17.7-21.2	59.0	24.2	21.4-27.2	45.2	16.2	14.0-18.5	16.0	2.8	2.2-3.5	10.4	4.2	3.1-5.6	5.6	1.8	1.2-2.7
	2016-2020	109.0	19.6	17.9-21.4	60.6	23.6	20.9-26.5	48.4	16.9	14.7-19.3	9.4	1.6	1.2-2.2	5.8	2.4	1.6-3.4	3.6	1.2	0.7-1.9
Female breast	1976-1980							241.8	106.4	100.4-112.7							97.0	41.5	37.8-45.4
	1981-1985							280.8	118.4	112.1-124.9							93.2	38.0	34.6-41.7
	1986-1990							287.0	116.7	110.6-123.1							96.0	37.8	34.4-41.5
	1991-1995							310.0	122.5	116.4-128.9							88.4	32.9	29.8-36.2
	1996-2000							354.8	137.5	131.1-144.1							72.8	26.4	23.7-29.3
	2001-2005							358.0	133.5	127.3-139.9							73.8	26.1	23.4-28.9
	2006-2010							368.6	131.2	125.2-137.5							69.8	22.6	20.3-25.2
	2011-2015							418.2	141.5	135.3-147.9							59.8	19.0	16.8-21.4
	2016-2020							412.8	137.0	130.9-143.4							59.0	18.0	15.9-20.4
Cervix uteri	1976-1980							31.2	13.7	11.6-16.1							10.4	4.7	3.5-6.2
	1981-1985							20.6	9.1	7.3-11.0							9.0	3.7	2.7-4.9
	1986-1990							31.0	13.1	11.1-15.4							8.4	3.3	2.3-4.5
	1991-1995							31.0	12.3	10.4-14.4							7.0	3.0	2.1-4.2
	1996-2000							17.4	7.0	5.6-8.7							5.8	2.2	1.5-3.2
	2001-2005							15.8	6.3	5.0-7.9							4.8	1.7	1.1-2.6
	2006-2010							17.0	6.9	5.5-8.6							4.6	1.7	1.0-2.5
	2011-2015							15.2	5.9	4.6-7.5							6.4	2.2	1.5-3.2
	2016-2020							16.0	6.5	5.0-8.1							4.4	1.5	0.9-2.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Onondaga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							85.0	35.5	32.2-39.1							14.4	6.1	4.8-7.7
	1981-1985							69.0	28.5	25.5-31.8							10.6	4.1	3.1-5.4
	1986-1990							60.4	24.9	22.1-27.9							11.0	4.0	3.0-5.2
	1991-1995							66.2	26.1	23.3-29.1							11.4	4.1	3.1-5.3
	1996-2000							61.2	23.4	20.8-26.2							13.6	4.8	3.8-6.2
	2001-2005							77.8	28.7	25.9-31.7							9.4	3.1	2.3-4.2
	2006-2010							82.8	28.9	26.2-31.9							14.6	4.7	3.6-5.9
	2011-2015							93.4	30.6	27.8-33.6							15.0	4.5	3.5-5.8
	2016-2020							93.8	29.2	26.5-32.1							15.2	4.4	3.5-5.6
Ovary	1976-1980							34.8	15.0	12.8-17.5							21.0	9.1	7.4-11.0
	1981-1985							37.4	15.6	13.4-18.0							25.2	9.8	8.2-11.7
	1986-1990							36.0	14.8	12.7-17.2							26.6	10.2	8.5-12.1
	1991-1995							45.2	17.4	15.2-19.9							26.0	9.5	7.9-11.4
	1996-2000							38.6	14.8	12.8-17.1							26.2	9.7	8.1-11.5
	2001-2005							38.2	14.1	12.2-16.3							21.4	7.3	6.0-8.8
	2006-2010							42.6	14.8	12.9-17.0							27.0	8.8	7.4-10.5
	2011-2015							33.4	11.6	9.8-13.5							29.0	9.1	7.6-10.8
	2016-2020							30.6	9.8	8.3-11.6							17.8	5.4	4.3-6.7
Prostate	1976-1980				127.0	86.8	79.8-94.2							48.2	36.1	31.5-41.2			
	1981-1985				149.2	91.9	85.1-99.0							50.4	36.8	32.2-41.8			
	1986-1990				191.2	109.5	102.3-117.0							51.0	34.2	30.0-38.9			
	1991-1995				311.6	159.8	151.7-168.2							58.8	36.3	32.1-40.8			
	1996-2000				326.4	161.4	153.6-169.6							56.6	33.0	29.1-37.1			
	2001-2005				421.8	197.0	188.7-205.7							42.0	22.5	19.5-25.8			
	2006-2010				502.2	213.6	205.2-222.2							45.6	21.3	18.6-24.3			
	2011-2015				417.0	154.6	147.9-161.6							43.2	18.6	16.2-21.3			
	2016-2020				468.8	155.9	149.4-162.5							39.0	15.3	13.2-17.7			
Testis	1976-1980				10.8	4.8	3.6-6.4							2.0	0.8	0.4-1.5			
	1981-1985				12.4	5.1	3.9-6.7							1.0	0.5	0.1-1.3			
	1986-1990				13.0	5.4	4.2-7.0							1.4	0.6	0.2-1.3			
	1991-1995				16.8	7.0	5.5-8.7							0.6	0.3	0.1-0.8			
	1996-2000				12.8	5.7	4.4-7.3							0.6	0.3	0.1-0.8			
	2001-2005				16.6	7.7	6.1-9.6							0.4	0.2	0.0-0.7			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Onondaga**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				13.6	6.4	4.9-8.1							0.8	0.4	0.1-0.9			
	2011-2015				16.2	7.7	6.1-9.6							0.6	0.3	0.1-0.8			
	2016-2020				14.4	6.8	5.3-8.6							0.4	0.2	0.0-0.6			
Urinary bladder (incl. in situ)	1976-1980	88.4	22.2	20.2-24.5	66.2	41.5	36.9-46.4	22.2	9.5	7.8-11.4	29.0	7.8	6.5-9.1	20.8	14.7	11.9-17.9	8.2	3.6	2.6-5.0
	1981-1985	101.8	23.7	21.7-25.9	73.6	42.8	38.3-47.6	28.2	11.2	9.4-13.2	28.4	6.9	5.8-8.2	19.0	13.3	10.6-16.3	9.4	3.7	2.7-4.9
	1986-1990	91.2	20.1	18.2-22.0	64.2	35.1	31.2-39.4	27.0	10.3	8.6-12.3	24.2	5.5	4.5-6.6	17.8	11.3	9.0-14.0	6.4	2.3	1.6-3.2
	1991-1995	104.2	22.1	20.2-24.1	69.8	37.1	33.2-41.3	34.4	12.3	10.5-14.3	25.8	5.5	4.6-6.5	15.0	9.0	7.0-11.4	10.8	3.6	2.7-4.7
	1996-2000	105.8	21.9	20.1-23.9	74.2	38.1	34.3-42.2	31.6	11.0	9.3-12.9	24.2	4.9	4.1-5.9	15.8	8.8	6.9-10.9	8.4	2.7	1.9-3.7
	2001-2005	130.2	26.1	24.2-28.2	95.4	45.9	41.9-50.3	34.8	11.9	10.2-13.9	25.4	5.0	4.1-5.9	16.6	8.5	6.8-10.6	8.8	2.8	2.0-3.8
	2006-2010	151.4	28.4	26.4-30.6	115.0	51.6	47.4-56.0	36.4	12.0	10.2-13.9	28.8	5.2	4.4-6.2	22.0	10.1	8.3-12.2	6.8	2.0	1.4-2.9
	2011-2015	150.2	26.6	24.7-28.6	109.6	45.0	41.2-49.0	40.6	13.1	11.3-15.1	26.0	4.5	3.7-5.3	17.8	7.5	6.0-9.2	8.2	2.5	1.8-3.4
	2016-2020	157.4	26.1	24.3-28.0	119.8	46.1	42.4-50.1	37.6	10.9	9.3-12.6	22.0	3.5	2.9-4.3	14.8	5.8	4.5-7.3	7.2	2.0	1.4-2.8
Kidney and renal pelvis	1976-1980	32.6	7.8	6.6-9.1	19.4	11.0	8.9-13.6	13.2	5.6	4.3-7.1	17.6	4.3	3.4-5.3	10.4	6.4	4.7-8.5	7.2	3.0	2.1-4.2
	1981-1985	39.2	9.1	7.8-10.5	23.2	12.3	10.1-14.8	16.0	6.5	5.2-8.2	16.2	3.7	3.0-4.7	9.0	4.8	3.4-6.6	7.2	3.0	2.1-4.1
	1986-1990	40.8	9.3	8.1-10.7	24.0	12.7	10.5-15.3	16.8	6.8	5.4-8.5	21.6	4.9	4.0-5.9	11.6	6.7	5.0-8.7	10.0	3.8	2.8-5.1
	1991-1995	51.4	10.9	9.6-12.4	26.8	13.6	11.4-16.2	24.6	9.2	7.6-11.0	18.4	3.9	3.1-4.8	9.4	5.1	3.7-6.8	9.0	3.2	2.3-4.3
	1996-2000	62.8	13.4	11.9-14.9	39.4	19.2	16.6-22.1	23.4	8.8	7.3-10.6	18.6	3.9	3.2-4.8	10.4	5.1	3.8-6.8	8.2	2.9	2.1-4.0
	2001-2005	76.4	15.6	14.1-17.3	47.0	22.1	19.3-25.1	29.4	10.8	9.1-12.7	17.4	3.5	2.8-4.3	11.4	5.6	4.2-7.3	6.0	2.0	1.4-3.0
	2006-2010	87.2	16.8	15.3-18.5	54.8	23.3	20.5-26.2	32.4	11.6	9.9-13.6	24.4	4.6	3.8-5.5	14.8	6.4	5.0-8.0	9.6	3.3	2.4-4.4
	2011-2015	90.0	16.1	14.6-17.7	55.6	21.6	19.1-24.4	34.4	11.5	9.8-13.4	18.8	3.3	2.6-4.0	12.0	4.9	3.8-6.4	6.8	2.0	1.4-2.9
	2016-2020	115.2	20.1	18.4-21.9	74.0	27.9	25.0-31.0	41.2	13.3	11.5-15.4	20.4	3.3	2.7-4.0	11.8	4.1	3.1-5.4	8.6	2.5	1.8-3.5
Brain and other central nervous system	1976-1980	30.0	7.2	6.0-8.5	16.2	8.5	6.6-10.6	13.8	6.1	4.7-7.7	24.2	5.8	4.8-6.9	11.8	6.1	4.6-8.0	12.4	5.4	4.1-6.9
	1981-1985	32.6	7.4	6.3-8.7	20.0	10.0	8.0-12.2	12.6	5.3	4.1-6.9	28.0	6.5	5.4-7.7	16.0	8.4	6.6-10.6	12.0	5.0	3.8-6.5
	1986-1990	30.4	6.6	5.6-7.8	16.4	8.0	6.3-10.0	14.0	5.4	4.2-6.9	20.8	4.7	3.8-5.7	11.8	6.2	4.7-8.1	9.0	3.4	2.5-4.6
	1991-1995	37.4	8.2	7.0-9.4	20.4	9.8	8.0-12.0	17.0	6.8	5.4-8.4	22.2	4.9	4.0-5.9	12.6	6.3	4.8-8.0	9.6	3.8	2.8-5.1
	1996-2000	37.0	7.9	6.8-9.1	22.0	10.6	8.7-12.8	15.0	5.7	4.4-7.1	23.0	4.9	4.0-5.9	12.2	5.8	4.4-7.4	10.8	4.0	3.0-5.2
	2001-2005	38.0	8.1	7.0-9.3	20.6	9.8	7.9-11.8	17.4	6.7	5.4-8.3	23.6	4.9	4.0-5.8	13.0	6.2	4.8-7.9	10.6	3.9	2.9-5.1
	2006-2010	36.8	7.5	6.5-8.7	18.6	8.3	6.7-10.2	18.2	6.9	5.5-8.5	21.4	4.2	3.4-5.1	11.8	5.3	4.0-6.9	9.6	3.3	2.4-4.4
	2011-2015	42.6	8.4	7.3-9.7	23.2	9.8	8.0-11.8	19.4	7.3	5.8-9.0	26.8	4.9	4.1-5.9	15.8	6.4	5.0-8.0	11.0	3.7	2.8-4.9
	2016-2020	35.4	6.4	5.5-7.5	21.2	8.3	6.7-10.1	14.2	4.8	3.7-6.1	31.8	5.8	4.9-6.8	18.8	7.4	5.9-9.1	13.0	4.4	3.4-5.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Onondaga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	15.0	3.6	2.8-4.5	5.0	2.7	1.7-4.1	10.0	4.5	3.3-5.9	2.8	0.8	0.4-1.3	1.4	0.9	0.4-2.0	1.4	0.6	0.3-1.3
	1981-1985	18.6	4.2	3.4-5.2	6.0	3.0	2.0-4.3	12.6	5.4	4.1-6.9	2.8	0.7	0.4-1.2	0.8	0.4	0.1-1.0	2.0	0.8	0.4-1.5
	1986-1990	17.6	3.9	3.1-4.8	4.6	2.2	1.4-3.4	13.0	5.5	4.2-7.1	2.2	0.5	0.2-0.9	1.4	0.8	0.3-1.7	0.8	0.3	0.1-0.8
	1991-1995	22.0	4.8	3.9-5.8	5.8	2.7	1.8-3.9	16.2	6.5	5.2-8.1	2.8	0.6	0.3-1.0	0.8	0.5	0.1-1.3	2.0	0.7	0.3-1.3
	1996-2000	35.0	7.6	6.5-8.8	9.2	4.4	3.2-5.8	25.8	10.7	8.9-12.7	2.0	0.4	0.2-0.8	0.4	0.2	0.0-0.7	1.6	0.6	0.2-1.1
	2001-2005	48.0	10.5	9.2-11.9	12.6	5.8	4.5-7.4	35.4	14.9	12.7-17.2	2.0	0.4	0.2-0.7	0.6	0.3	0.1-0.8	1.4	0.5	0.2-1.0
	2006-2010	73.2	15.3	13.7-17.0	18.4	8.1	6.5-9.9	54.8	22.0	19.4-24.8	3.6	0.7	0.4-1.1	2.0	0.9	0.4-1.6	1.6	0.5	0.2-1.0
	2011-2015	86.8	17.6	15.9-19.4	20.4	8.5	6.9-10.4	66.4	26.1	23.3-29.2	4.0	0.7	0.4-1.1	1.8	0.8	0.3-1.5	2.2	0.7	0.3-1.3
2016-2020	78.0	16.1	14.4-17.8	20.4	8.1	6.6-9.9	57.6	23.7	20.9-26.7	4.2	0.7	0.4-1.1	1.8	0.7	0.3-1.5	2.4	0.7	0.4-1.3	
Hodgkin lymphoma	1976-1980	16.0	3.3	2.6-4.1	9.0	4.0	2.9-5.4	7.0	2.5	1.8-3.6	4.6	1.0	0.6-1.5	3.2	1.5	0.8-2.5	1.4	0.5	0.2-1.1
	1981-1985	19.8	4.1	3.3-5.1	9.6	4.5	3.3-6.1	10.2	3.8	2.8-5.0	4.4	1.0	0.6-1.5	3.0	1.3	0.7-2.3	1.4	0.6	0.2-1.3
	1986-1990	16.6	3.5	2.8-4.4	8.0	3.6	2.6-5.0	8.6	3.3	2.4-4.5	6.2	1.4	0.9-2.0	3.6	1.8	1.1-2.9	2.6	1.0	0.5-1.7
	1991-1995	15.6	3.3	2.6-4.1	8.4	3.7	2.7-5.0	7.2	2.9	2.0-4.0	3.8	0.8	0.5-1.3	2.4	1.2	0.6-2.1	1.4	0.5	0.2-1.0
	1996-2000	17.4	3.8	3.0-4.7	9.8	4.6	3.4-6.1	7.6	3.1	2.2-4.3	2.8	0.6	0.3-1.0	1.0	0.5	0.2-1.2	1.8	0.7	0.3-1.4
	2001-2005	14.4	3.1	2.4-3.9	8.4	3.9	2.8-5.2	6.0	2.4	1.6-3.4	2.2	0.4	0.2-0.8	1.2	0.6	0.2-1.2	1.0	0.4	0.1-0.9
	2006-2010	15.4	3.4	2.6-4.2	9.6	4.4	3.2-5.9	5.8	2.4	1.6-3.5	2.8	0.6	0.3-1.0	2.2	1.0	0.5-1.8	0.6	0.2	0.0-0.6
	2011-2015	14.4	2.9	2.3-3.7	8.4	3.5	2.5-4.8	6.0	2.3	1.5-3.3	1.0	0.2	0.1-0.5	0.4	0.2	0.0-0.7	0.6	0.2	0.0-0.6
2016-2020	14.0	2.9	2.3-3.7	6.8	3.0	2.0-4.2	7.2	2.9	2.0-4.1	1.2	0.2	0.1-0.5	0.6	0.2	0.0-0.7	0.6	0.2	0.0-0.6	
Non-Hodgkin lymphomas	1976-1980	54.0	13.0	11.4-14.6	26.8	15.5	12.9-18.5	27.2	11.3	9.5-13.4	28.4	7.1	5.9-8.3	15.0	9.3	7.2-11.8	13.4	5.6	4.3-7.1
	1981-1985	61.4	14.4	12.8-16.2	34.4	19.6	16.6-22.8	27.0	10.7	9.0-12.7	30.2	7.1	6.0-8.3	15.8	9.4	7.4-11.8	14.4	5.7	4.4-7.2
	1986-1990	65.4	14.7	13.1-16.4	36.0	19.4	16.5-22.5	29.4	11.4	9.6-13.4	42.2	9.6	8.3-11.0	21.4	12.1	9.8-14.7	20.8	7.8	6.4-9.5
	1991-1995	79.2	17.0	15.4-18.8	37.6	18.5	15.9-21.5	41.6	15.8	13.7-18.1	37.2	8.0	6.9-9.3	16.4	8.7	6.9-10.9	20.8	7.3	6.0-8.9
	1996-2000	100.8	21.3	19.5-23.2	53.8	26.2	23.2-29.6	47.0	16.9	14.8-19.3	42.4	8.8	7.6-10.0	20.4	10.4	8.5-12.7	22.0	7.4	6.1-9.0
	2001-2005	104.8	21.3	19.5-23.2	53.4	25.2	22.2-28.4	51.4	18.0	15.8-20.4	35.8	7.1	6.1-8.3	19.8	9.7	7.9-11.8	16.0	5.2	4.1-6.5
	2006-2010	129.2	24.6	22.7-26.6	71.4	31.0	27.8-34.4	57.8	19.3	17.1-21.7	33.0	6.0	5.1-7.0	17.8	8.0	6.4-9.9	15.2	4.5	3.5-5.7
	2011-2015	123.8	22.1	20.4-24.0	68.2	27.4	24.5-30.6	55.6	18.0	15.9-20.3	38.4	6.6	5.7-7.6	20.4	8.5	6.9-10.4	18.0	5.1	4.1-6.4
2016-2020	115.2	19.9	18.2-21.7	56.8	21.7	19.2-24.5	58.4	18.3	16.2-20.7	25.8	4.3	3.5-5.1	15.0	6.0	4.7-7.6	10.8	2.8	2.1-3.8	
Myeloma	1976-1980	16.0	3.9	3.1-4.9	7.0	4.1	2.8-5.8	9.0	3.9	2.8-5.3	10.8	2.6	2.0-3.5	5.2	3.0	1.9-4.4	5.6	2.3	1.6-3.4
	1981-1985	18.6	4.4	3.6-5.4	9.4	5.6	4.1-7.5	9.2	3.7	2.7-5.0	14.2	3.3	2.5-4.1	7.4	4.4	3.0-6.1	6.8	2.7	1.8-3.8
	1986-1990	21.6	4.8	4.0-5.9	12.8	7.2	5.5-9.3	8.8	3.4	2.4-4.6	16.2	3.7	2.9-4.6	9.6	5.7	4.1-7.6	6.6	2.4	1.6-3.4
	1991-1995	25.0	5.3	4.4-6.3	14.2	7.5	5.8-9.5	10.8	3.9	2.9-5.1	15.6	3.2	2.6-4.1	9.0	4.8	3.4-6.4	6.6	2.3	1.6-3.2
	1996-2000	26.0	5.4	4.5-6.4	12.0	6.2	4.7-8.0	14.0	5.0	3.9-6.4	19.0	4.0	3.2-4.8	9.0	5.0	3.6-6.7	10.0	3.5	2.6-4.6
	2001-2005	36.0	7.2	6.2-8.4	17.2	8.4	6.7-10.3	18.8	6.5	5.2-8.0	19.0	3.7	3.0-4.6	9.2	4.4	3.2-5.9	9.8	3.2	2.3-4.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Onondaga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	36.2	6.9	5.9-8.0	17.4	7.7	6.1-9.5	18.8	6.4	5.1-7.9	15.0	2.8	2.2-3.5	7.0	3.2	2.2-4.5	8.0	2.5	1.8-3.5
	2011-2015	33.8	5.9	5.0-6.9	17.0	6.7	5.3-8.3	16.8	5.3	4.2-6.6	19.6	3.5	2.8-4.3	9.4	4.0	2.9-5.3	10.2	3.2	2.3-4.2
	2016-2020	48.4	8.5	7.4-9.7	24.8	9.6	7.9-11.5	23.6	7.7	6.3-9.2	14.2	2.3	1.8-3.0	6.4	2.5	1.7-3.5	7.8	2.2	1.6-3.1
Leukemias	1976-1980	50.4	12.5	10.9-14.1	27.2	15.9	13.2-19.0	23.2	10.0	8.3-12.1	36.6	9.1	7.8-10.5	19.6	12.0	9.6-14.8	17.0	7.1	5.7-8.8
	1981-1985	52.6	12.3	10.9-13.9	31.2	17.5	14.7-20.7	21.4	8.6	7.0-10.4	32.4	7.7	6.5-9.0	17.4	10.2	8.1-12.8	15.0	6.1	4.8-7.7
	1986-1990	51.0	11.3	9.9-12.8	24.6	12.6	10.4-15.2	26.4	10.2	8.5-12.2	33.4	7.4	6.3-8.7	15.0	8.0	6.2-10.1	18.4	6.9	5.5-8.5
	1991-1995	63.6	13.5	12.1-15.1	36.0	19.2	16.4-22.3	27.6	9.8	8.2-11.7	36.0	7.7	6.6-9.0	19.6	11.3	9.1-13.8	16.4	5.9	4.7-7.4
	1996-2000	71.6	14.9	13.4-16.6	38.4	19.5	16.8-22.5	33.2	11.9	10.2-13.9	39.2	8.1	7.0-9.3	21.8	11.2	9.1-13.5	17.4	6.1	4.8-7.5
	2001-2005	76.2	15.3	13.8-17.0	43.2	20.5	17.8-23.4	33.0	11.5	9.8-13.4	39.4	7.8	6.7-9.0	21.4	10.6	8.6-12.8	18.0	5.9	4.7-7.3
	2006-2010	89.8	17.5	15.9-19.2	54.8	24.6	21.8-27.8	35.0	11.8	10.1-13.7	42.6	8.1	7.0-9.3	27.4	12.6	10.6-14.9	15.2	4.8	3.8-6.1
	2011-2015	105.0	19.2	17.5-21.0	59.6	24.5	21.8-27.6	45.4	15.0	13.0-17.2	44.0	7.8	6.8-9.0	23.4	9.9	8.2-11.9	20.6	6.4	5.2-7.9
2016-2020	91.2	15.6	14.1-17.1	55.2	21.0	18.5-23.7	36.0	11.2	9.5-13.1	34.6	5.9	5.0-6.8	18.4	7.4	5.9-9.1	16.2	4.8	3.7-6.0	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**Ontario**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	321.6	407.5	387.5-428.2	159.4	459.3	426.8-493.5	162.2	378.9	352.8-406.4	170.6	217.6	203.0-232.9	88.4	263.1	238.2-289.8	82.2	187.2	169.3-206.5
	1981-1985	353.4	409.7	390.5-429.5	168.6	438.1	408.0-469.7	184.8	398.9	373.1-426.0	177.8	207.8	194.1-222.1	91.6	245.5	222.7-269.8	86.2	181.3	164.4-199.5
	1986-1990	408.4	445.2	425.9-465.2	198.4	492.5	461.3-525.1	210.0	421.6	396.1-448.3	185.8	201.9	189.0-215.4	97.0	243.6	221.7-267.0	88.8	173.3	157.4-190.5
	1991-1995	505.2	509.0	489.3-529.4	262.8	597.5	565.0-631.4	242.4	452.1	426.7-478.6	218.0	220.1	207.2-233.7	117.8	280.2	257.4-304.3	100.2	180.9	165.2-197.6
	1996-2000	542.6	521.2	501.7-541.2	279.4	612.6	580.5-646.1	263.2	463.7	438.8-489.7	225.6	215.7	203.3-228.7	114.8	260.8	239.6-283.3	110.8	184.1	169.0-200.3
	2001-2005	552.2	484.9	466.9-503.5	285.4	565.7	536.3-596.3	266.8	429.4	406.4-453.5	224.2	195.4	184.1-207.3	113.8	237.0	217.6-257.6	110.4	169.1	155.1-184.2
	2006-2010	647.8	506.0	488.4-524.2	341.8	582.1	554.2-611.1	306.0	446.8	424.1-470.6	230.6	177.8	167.5-188.6	120.8	215.0	197.9-233.2	109.8	153.1	140.2-167.0
	2011-2015	688.6	481.1	464.6-498.1	344.8	519.2	494.1-545.4	343.8	456.5	434.2-479.7	235.0	160.8	151.4-170.5	122.0	192.8	177.4-209.3	113.0	138.4	126.8-150.8
	2016-2020	802.2	508.1	491.7-524.9	427.6	570.8	545.9-596.6	374.6	464.7	442.4-487.8	252.0	152.0	143.5-160.9	133.4	186.0	171.7-201.2	118.6	129.0	118.6-140.4
Oral cavity and pharynx	1976-1980	8.2	10.4	7.4-14.1	4.2	11.5	7.0-17.8	4.0	9.5	5.7-14.8	2.2	2.8	1.4-5.0	1.6	4.6	1.9-9.0	0.6	1.2	0.2-3.7
	1981-1985	8.0	9.3	6.6-12.8	4.2	10.8	6.6-16.7	3.8	8.4	5.0-13.2	3.8	4.3	2.6-6.8	2.0	5.0	2.4-9.2	1.8	3.5	1.6-6.8
	1986-1990	7.6	8.5	6.0-11.7	4.2	10.9	6.7-16.8	3.4	7.1	4.1-11.5	2.4	2.6	1.3-4.5	1.6	3.6	1.5-7.1	0.8	1.7	0.4-4.4
	1991-1995	7.4	7.6	5.4-10.5	4.4	9.5	6.0-14.5	3.0	5.7	3.2-9.5	3.8	3.8	2.3-6.0	2.2	5.0	2.5-9.1	1.6	2.8	1.2-5.6
	1996-2000	12.0	11.5	8.8-14.8	8.2	16.6	11.9-22.6	3.8	6.7	4.0-10.5	1.8	1.7	0.8-3.3	1.4	3.0	1.2-6.3	0.4	0.6	0.1-2.3
	2001-2005	12.8	11.1	8.5-14.2	8.8	16.7	12.1-22.6	4.0	6.1	3.7-9.7	4.0	3.4	2.1-5.3	2.8	5.3	2.9-9.0	1.2	1.8	0.6-4.1
	2006-2010	15.8	12.1	9.6-15.2	10.6	16.8	12.5-22.2	5.2	7.8	5.1-11.6	2.8	2.2	1.2-3.8	1.8	2.9	1.3-5.7	1.0	1.5	0.5-3.7
	2011-2015	15.2	10.4	8.1-13.1	11.0	16.3	12.1-21.5	4.2	5.2	3.2-8.3	3.2	2.3	1.3-3.8	2.4	4.0	2.0-7.1	0.8	1.0	0.3-2.9
	2016-2020	22.2	13.3	10.8-16.2	15.4	19.0	14.8-24.1	6.8	8.6	5.8-12.4	4.2	2.5	1.5-3.9	3.4	4.6	2.6-7.6	0.8	1.0	0.3-2.9
Esophagus	1976-1980	2.4	2.7	1.4-4.7	2.2	5.3	2.6-9.7	0.2	0.4	0.0-2.5	1.2	1.3	0.5-2.9	1.2	2.8	1.0-6.3	0.0	0.0	0.0-1.7
	1981-1985	2.8	3.3	1.8-5.5	2.2	5.8	2.8-10.5	0.6	1.4	0.3-4.1	4.0	4.5	2.7-7.0	3.4	8.6	4.9-13.9	0.6	1.4	0.3-4.1
	1986-1990	3.4	3.6	2.1-5.8	2.4	5.5	2.8-9.7	1.0	1.9	0.6-4.5	4.0	4.2	2.6-6.5	3.0	7.0	3.8-11.8	1.0	2.1	0.7-4.9
	1991-1995	2.4	2.5	1.3-4.4	2.4	5.6	2.9-9.7	0.0	0.0	0.0-1.5	3.8	4.0	2.4-6.3	3.0	6.9	3.9-11.4	0.8	1.5	0.4-4.0
	1996-2000	4.2	4.0	2.5-6.2	3.6	7.5	4.4-11.9	0.6	1.2	0.2-3.4	4.2	4.0	2.5-6.2	3.4	7.0	4.1-11.3	0.8	1.4	0.4-3.7
	2001-2005	6.6	6.0	4.1-8.4	5.4	10.5	6.9-15.3	1.2	2.0	0.7-4.4	8.0	7.1	5.1-9.7	6.8	13.2	9.1-18.6	1.2	2.0	0.7-4.4
	2006-2010	8.2	6.4	4.6-8.7	5.6	9.3	6.1-13.6	2.6	3.9	2.1-6.9	6.0	4.6	3.1-6.7	4.8	8.2	5.2-12.3	1.2	1.7	0.6-3.9
	2011-2015	7.6	5.4	3.8-7.6	5.2	8.0	5.1-11.9	2.4	3.1	1.5-5.7	7.0	5.1	3.5-7.2	5.0	7.6	4.8-11.6	2.0	2.7	1.2-5.2
	2016-2020	10.2	5.8	4.3-7.7	9.4	12.0	8.7-16.2	0.8	0.8	0.2-2.5	7.4	4.4	3.1-6.2	6.6	8.8	6.0-12.7	0.8	0.8	0.2-2.5
Stomach	1976-1980	5.6	7.3	4.8-10.6	2.8	8.1	4.4-13.7	2.8	6.3	3.5-10.6	5.4	7.1	4.7-10.4	3.0	9.4	5.1-15.6	2.4	5.4	2.8-9.5
	1981-1985	7.8	9.1	6.5-12.5	5.4	15.3	9.9-22.3	2.4	4.7	2.4-8.3	5.2	6.2	4.0-9.1	3.4	10.2	5.8-16.5	1.8	3.7	1.7-7.0
	1986-1990	10.0	10.6	7.9-14.1	5.8	13.8	9.1-20.0	4.2	8.0	4.9-12.3	5.0	5.4	3.5-8.0	3.0	7.3	4.0-12.1	2.0	3.6	1.7-6.7
	1991-1995	9.4	9.5	6.9-12.6	6.4	14.4	9.8-20.4	3.0	5.1	2.8-8.5	5.8	5.9	4.0-8.5	4.0	9.7	5.8-15.0	1.8	3.1	1.4-6.0
	1996-2000	7.2	6.9	4.8-9.5	4.4	9.8	6.1-14.9	2.8	4.6	2.5-7.9	4.2	4.0	2.5-6.1	2.4	5.7	2.9-9.9	1.8	2.9	1.3-5.6
	2001-2005	7.4	6.5	4.6-9.0	4.8	9.1	5.8-13.7	2.6	4.3	2.3-7.5	4.4	3.8	2.4-5.8	2.2	4.8	2.4-8.5	2.2	3.2	1.6-6.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020  
Ontario**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	7.6	6.0	4.2-8.3	5.0	8.8	5.7-13.2	2.6	3.9	2.0-6.9	3.2	2.5	1.4-4.1	2.0	3.6	1.7-6.8	1.2	1.7	0.6-4.0
	2011-2015	6.2	4.4	3.0-6.4	3.8	6.1	3.6-9.7	2.4	3.1	1.6-5.7	3.2	2.4	1.3-4.1	1.4	2.7	1.0-5.6	1.8	2.3	1.1-4.7
	2016-2020	8.4	5.4	3.8-7.6	6.0	8.5	5.6-12.5	2.4	2.9	1.4-5.5	3.2	2.2	1.2-3.8	2.2	3.6	1.7-6.8	1.0	1.0	0.3-2.7
Colorectal	1976-1980	54.2	68.8	60.7-77.6	27.6	79.9	66.7-94.9	26.6	60.4	50.5-71.7	27.8	35.5	29.8-42.0	14.4	43.7	33.9-55.4	13.4	29.8	23.1-38.0
	1981-1985	61.8	71.6	63.7-80.1	27.8	72.0	60.1-85.4	34.0	70.1	59.9-81.6	27.4	31.7	26.6-37.6	14.2	37.1	28.8-47.2	13.2	26.4	20.4-33.6
	1986-1990	61.8	67.6	60.2-75.7	29.0	74.5	62.5-88.1	32.8	63.6	54.2-74.3	26.8	28.9	24.2-34.3	13.0	31.7	24.2-40.6	13.8	26.6	20.6-33.8
	1991-1995	66.8	67.4	60.3-75.0	33.0	79.6	67.5-93.0	33.8	60.1	51.3-70.0	28.0	28.5	23.9-33.6	13.6	34.0	26.2-43.4	14.4	25.5	19.9-32.2
	1996-2000	59.6	57.0	50.7-63.9	29.6	65.8	55.5-77.4	30.0	50.5	42.7-59.4	25.0	23.8	19.8-28.4	11.6	27.2	20.6-35.3	13.4	21.5	16.6-27.4
	2001-2005	58.2	50.2	44.6-56.4	31.2	63.7	54.0-74.7	27.0	40.0	33.4-47.6	19.6	16.7	13.6-20.4	9.4	20.7	15.1-27.6	10.2	13.9	10.3-18.6
	2006-2010	47.2	37.2	32.5-42.4	23.0	41.1	33.8-49.5	24.2	33.7	27.8-40.5	17.4	13.4	10.7-16.7	8.6	16.2	11.6-21.9	8.8	11.6	8.3-15.8
	2011-2015	46.2	31.9	27.8-36.5	19.8	29.3	23.6-36.0	26.4	33.8	28.0-40.5	14.8	9.8	7.6-12.4	5.2	8.2	5.3-12.3	9.6	10.9	7.9-14.7
2016-2020	52.2	34.7	30.4-39.5	23.4	33.7	27.6-40.8	28.8	35.7	29.7-42.6	19.2	11.9	9.6-14.7	10.0	14.3	10.5-19.2	9.2	10.1	7.3-13.8	
Colon excluding rectum	1976-1980	37.6	48.1	41.4-55.6	17.4	51.6	41.0-64.0	20.2	45.8	37.2-55.8	23.0	29.6	24.4-35.6	11.4	34.5	25.8-45.0	11.6	26.0	19.7-33.6
	1981-1985	45.4	52.9	46.2-60.3	19.2	50.4	40.5-61.9	26.2	53.7	44.9-63.9	23.0	26.4	21.7-31.7	11.4	29.4	22.0-38.3	11.6	23.1	17.5-29.9
	1986-1990	48.0	52.3	45.8-59.4	22.2	57.6	47.0-69.8	25.8	49.3	41.1-58.7	23.0	24.8	20.5-29.9	10.4	25.5	18.8-33.7	12.6	24.3	18.6-31.2
	1991-1995	52.2	52.7	46.5-59.5	26.0	63.6	52.8-75.8	26.2	46.2	38.6-54.9	24.4	24.8	20.6-29.7	11.4	29.0	21.7-37.7	13.0	23.0	17.7-29.5
	1996-2000	42.8	41.0	35.7-46.8	19.2	43.7	35.2-53.5	23.6	39.7	32.8-47.7	21.2	20.2	16.5-24.5	10.4	24.6	18.3-32.4	10.8	17.3	13.0-22.7
	2001-2005	42.2	36.4	31.6-41.7	23.0	47.4	39.0-57.1	19.2	28.0	22.6-34.5	17.0	14.5	11.5-18.0	8.2	18.3	13.1-24.9	8.8	12.0	8.7-16.4
	2006-2010	34.8	27.8	23.8-32.3	17.0	30.5	24.3-37.9	17.8	25.3	20.2-31.4	14.0	10.8	8.4-13.8	7.4	14.3	10.0-19.8	6.6	8.7	5.9-12.5
	2011-2015	31.6	21.6	18.3-25.4	12.0	17.3	13.1-22.6	19.6	24.9	20.0-30.6	12.0	7.9	6.0-10.3	3.6	5.7	3.3-9.2	8.4	9.5	6.8-13.2
2016-2020	38.2	25.5	21.8-29.6	15.0	22.0	17.1-28.0	23.2	28.4	23.2-34.6	15.6	9.6	7.6-12.2	7.6	11.3	7.9-15.8	8.0	8.5	6.0-11.8	
Rectum & rectosigmoid	1976-1980	16.6	20.6	16.4-25.6	10.2	28.3	20.9-37.5	6.4	14.6	9.9-20.7	4.8	5.9	3.7-8.8	3.0	9.2	5.0-15.4	1.8	3.9	1.8-7.5
	1981-1985	16.4	18.7	14.8-23.2	8.6	21.6	15.5-29.3	7.8	16.4	11.6-22.5	4.4	5.3	3.3-8.1	2.8	7.8	4.2-13.1	1.6	3.3	1.4-6.6
	1986-1990	13.8	15.4	11.9-19.5	6.8	16.9	11.5-23.8	7.0	14.4	10.0-20.1	3.8	4.1	2.5-6.4	2.6	6.2	3.2-10.7	1.2	2.3	0.8-5.1
	1991-1995	14.6	14.7	11.5-18.5	7.0	16.0	11.1-22.3	7.6	14.0	9.8-19.2	3.6	3.6	2.1-5.7	2.2	5.1	2.5-9.1	1.4	2.4	1.0-5.1
	1996-2000	16.8	16.1	12.8-19.9	10.4	22.1	16.5-29.1	6.4	10.8	7.4-15.3	3.8	3.6	2.2-5.7	1.2	2.6	1.0-5.7	2.6	4.1	2.2-7.2
	2001-2005	16.0	13.8	10.9-17.2	8.2	16.3	11.6-22.2	7.8	11.9	8.5-16.5	2.6	2.3	1.2-3.9	1.2	2.4	0.9-5.3	1.4	1.9	0.8-4.2
	2006-2010	12.4	9.4	7.2-12.1	6.0	10.6	7.1-15.2	6.4	8.4	5.7-12.0	3.4	2.6	1.5-4.2	1.2	1.9	0.7-4.3	2.2	2.9	1.4-5.5
	2011-2015	14.6	10.3	8.0-13.1	7.8	12.0	8.4-16.6	6.8	8.9	6.0-12.9	2.8	1.8	1.0-3.2	1.6	2.5	1.1-5.2	1.2	1.4	0.5-3.3
2016-2020	14.0	9.2	7.1-11.9	8.4	11.7	8.2-16.2	5.6	7.3	4.7-11.0	3.6	2.3	1.3-3.7	2.4	3.0	1.5-5.6	1.2	1.6	0.6-3.8	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**Ontario**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	2.4	3.0	1.5-5.3	1.8	5.0	2.2-9.5	0.6	1.4	0.3-4.1	1.6	2.0	0.9-4.0	1.0	2.7	0.8-6.4	0.6	1.6	0.3-4.7
	1981-1985	4.0	4.5	2.7-7.0	1.4	3.2	1.2-6.7	2.6	5.6	2.9-9.6	2.8	3.2	1.7-5.4	1.2	2.7	1.0-6.2	1.6	3.4	1.5-6.8
	1986-1990	3.0	3.2	1.8-5.4	1.2	3.0	1.0-6.6	1.8	3.5	1.6-6.7	2.4	2.6	1.4-4.6	0.4	0.9	0.1-3.4	2.0	3.8	1.8-7.2
	1991-1995	4.8	4.9	3.1-7.3	2.0	4.3	2.1-8.0	2.8	5.0	2.7-8.5	4.0	4.0	2.5-6.3	1.8	4.4	2.0-8.4	2.2	3.9	1.9-7.1
	1996-2000	5.8	5.6	3.7-8.0	3.2	7.2	4.1-11.7	2.6	4.6	2.4-8.0	4.4	4.2	2.6-6.4	2.8	5.9	3.2-10.0	1.6	2.7	1.2-5.5
	2001-2005	5.0	4.5	2.9-6.7	2.4	4.5	2.3-8.0	2.6	4.4	2.3-7.7	3.8	3.3	2.0-5.2	2.0	4.2	2.0-7.8	1.8	2.6	1.2-5.2
	2006-2010	6.2	4.9	3.3-7.0	4.8	8.3	5.3-12.5	1.4	1.9	0.7-4.1	6.4	4.7	3.2-6.7	4.2	6.7	4.1-10.5	2.2	2.7	1.3-5.1
	2011-2015	8.2	5.5	3.9-7.6	5.0	7.0	4.5-10.6	3.2	4.0	2.2-6.8	6.6	4.7	3.2-6.8	3.2	5.2	2.9-8.6	3.4	4.1	2.3-7.0
2016-2020	8.8	5.1	3.7-7.0	6.2	8.1	5.4-11.8	2.6	2.7	1.4-5.0	8.2	4.9	3.5-6.8	6.2	8.2	5.5-12.0	2.0	2.3	1.1-4.5	
Pancreas	1976-1980	6.4	8.2	5.6-11.6	3.4	9.1	5.2-14.8	3.0	6.9	3.9-11.4	8.2	10.7	7.6-14.6	3.6	10.2	5.9-16.3	4.6	10.8	6.8-16.2
	1981-1985	8.2	9.7	6.9-13.2	3.6	9.3	5.5-14.8	4.6	9.5	6.0-14.4	9.2	10.9	8.0-14.6	3.6	9.2	5.4-14.7	5.6	11.8	7.8-17.2
	1986-1990	9.8	11.0	8.1-14.5	6.0	15.2	10.0-21.8	3.8	7.5	4.5-11.8	10.4	11.4	8.5-15.0	6.6	16.0	10.9-22.7	3.8	7.4	4.4-11.6
	1991-1995	9.4	9.6	7.1-12.8	4.0	9.3	5.6-14.5	5.4	9.5	6.2-13.9	9.4	9.5	6.9-12.6	4.4	10.2	6.3-15.5	5.0	8.8	5.7-13.1
	1996-2000	11.8	11.2	8.5-14.5	5.2	11.8	7.7-17.4	6.6	10.9	7.5-15.5	10.2	9.7	7.2-12.8	4.0	9.6	5.8-14.9	6.2	10.1	6.8-14.4
	2001-2005	14.6	12.7	10.0-16.0	5.8	11.6	7.7-16.8	8.8	13.3	9.6-18.1	12.8	11.0	8.4-14.1	5.4	11.3	7.4-16.5	7.4	11.0	7.7-15.4
	2006-2010	15.6	11.5	9.1-14.5	7.2	12.4	8.6-17.3	8.4	11.6	8.3-15.9	15.8	11.9	9.4-15.0	8.0	13.3	9.4-18.3	7.8	11.2	7.9-15.6
	2011-2015	20.0	13.2	10.7-16.2	8.6	12.4	8.8-17.0	11.4	13.7	10.3-18.1	15.0	10.1	7.9-12.8	7.0	10.7	7.3-15.2	8.0	9.6	6.8-13.3
2016-2020	23.6	14.6	12.0-17.6	12.8	18.1	13.8-23.4	10.8	11.8	8.8-15.7	19.2	11.4	9.2-14.0	10.0	13.7	10.0-18.3	9.2	9.6	7.0-13.1	
Larynx	1976-1980	4.4	5.4	3.4-8.2	4.0	10.5	6.3-16.3	0.4	0.9	0.1-3.4	1.4	1.7	0.7-3.5	1.4	4.0	1.5-8.4	0.0	0.0	0.0-1.7
	1981-1985	5.0	6.0	3.9-8.9	4.2	11.2	6.8-17.2	0.8	1.7	0.4-4.4	1.0	1.2	0.4-2.8	0.8	2.4	0.6-6.1	0.2	0.4	0.0-2.3
	1986-1990	5.8	6.3	4.2-9.1	4.6	10.9	6.8-16.5	1.2	2.7	1.0-5.9	1.2	1.3	0.5-2.9	0.8	1.9	0.5-4.8	0.4	0.7	0.1-2.8
	1991-1995	5.4	5.6	3.7-8.2	4.0	9.0	5.4-13.9	1.4	2.6	1.0-5.4	1.6	1.6	0.7-3.2	0.8	1.9	0.5-4.8	0.8	1.5	0.4-3.9
	1996-2000	4.6	4.4	2.8-6.6	3.2	7.0	4.0-11.5	1.4	2.4	0.9-5.0	1.6	1.5	0.7-3.0	1.4	3.1	1.2-6.3	0.2	0.3	0.0-1.9
	2001-2005	3.8	3.3	2.0-5.2	3.4	6.5	3.8-10.5	0.4	0.7	0.1-2.6	1.6	1.3	0.6-2.7	1.4	3.0	1.2-6.2	0.2	0.3	0.0-2.0
	2006-2010	5.0	3.9	2.5-5.9	4.2	7.2	4.4-11.2	0.8	1.0	0.3-2.9	0.8	0.5	0.1-1.5	0.8	1.2	0.3-3.3	0.0	0.0	0.0-1.4
	2011-2015	5.2	3.5	2.3-5.3	3.6	5.5	3.2-8.9	1.6	1.8	0.8-3.9	3.0	2.1	1.2-3.6	1.8	3.2	1.4-6.1	1.2	1.4	0.5-3.4
2016-2020	5.4	3.2	2.1-4.8	4.6	5.8	3.6-9.0	0.8	0.9	0.2-2.8	1.2	0.8	0.3-1.9	1.2	1.8	0.6-4.1	0.0	0.0	0.0-1.4	
Lung and bronchus	1976-1980	47.8	59.9	52.4-68.1	35.6	100.3	85.7-116.7	12.2	27.8	21.2-35.9	39.4	49.4	42.6-56.9	28.6	80.2	67.2-94.8	10.8	24.6	18.4-32.3
	1981-1985	54.4	61.1	54.0-69.0	37.8	92.5	79.4-107.0	16.6	36.1	28.6-44.9	41.6	47.6	41.3-54.6	30.4	78.1	65.9-92.0	11.2	23.5	17.7-30.7
	1986-1990	66.0	69.6	62.2-77.6	39.6	93.7	80.8-108.1	26.4	51.5	43.0-61.3	48.8	51.8	45.4-58.8	30.0	72.5	61.1-85.4	18.8	36.1	29.1-44.3
	1991-1995	74.8	74.4	67.0-82.4	46.0	101.1	88.3-115.2	28.8	52.9	44.5-62.4	62.6	62.4	55.7-69.8	40.6	92.3	79.8-106.1	22.0	40.2	33.0-48.6
	1996-2000	79.8	76.4	69.1-84.3	46.4	101.4	88.6-115.4	33.4	58.0	49.5-67.6	69.8	66.7	59.9-74.1	41.0	91.0	78.8-104.5	28.8	48.7	41.0-57.5
	2001-2005	85.6	76.0	69.0-83.6	49.2	97.3	85.3-110.4	36.4	59.1	50.8-68.5	65.4	57.6	51.5-64.3	37.4	75.4	64.8-87.2	28.0	44.6	37.4-52.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**Ontario**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	94.4	73.6	67.0-80.7	51.0	87.5	76.9-99.3	43.4	62.9	54.7-72.2	72.4	56.3	50.6-62.5	40.0	69.4	60.0-80.0	32.4	47.0	39.9-55.1
	2011-2015	110.8	75.1	68.8-81.8	54.2	80.5	70.9-91.1	56.6	71.9	63.6-81.2	68.2	46.3	41.4-51.6	37.0	56.2	48.2-65.3	31.2	39.4	33.3-46.4
	2016-2020	98.4	58.8	53.6-64.4	48.0	63.0	55.1-71.9	50.4	56.3	49.3-64.1	61.6	36.6	32.5-41.1	30.0	39.9	33.6-47.2	31.6	34.5	29.3-40.7
Melanoma of the skin	1976-1980	5.2	6.8	4.4-10.0	3.0	9.0	4.9-14.9	2.2	5.4	2.6-9.8	1.4	2.0	0.8-4.1	0.8	3.1	0.8-7.5	0.6	1.4	0.3-4.2
	1981-1985	7.4	8.5	5.9-11.7	4.2	9.5	5.8-14.7	3.2	7.2	4.0-11.7	2.0	2.1	1.0-3.9	1.2	2.4	0.9-5.6	0.8	1.6	0.4-4.1
	1986-1990	6.8	7.2	5.0-10.1	4.2	9.3	5.7-14.3	2.6	5.2	2.7-8.9	2.8	3.1	1.7-5.2	2.0	4.6	2.2-8.6	0.8	1.6	0.4-4.3
	1991-1995	10.8	10.9	8.2-14.2	6.2	13.4	9.1-19.2	4.6	8.7	5.5-13.1	2.4	2.5	1.3-4.4	1.2	2.6	1.0-5.7	1.2	2.1	0.7-4.6
	1996-2000	12.6	12.3	9.5-15.8	7.6	16.2	11.5-22.4	5.0	9.1	5.8-13.5	4.0	3.9	2.4-6.1	2.4	5.2	2.7-9.1	1.6	2.6	1.1-5.2
	2001-2005	18.2	16.0	12.9-19.7	10.2	20.0	14.8-26.4	8.0	13.4	9.5-18.4	3.4	2.9	1.7-4.7	2.4	4.5	2.3-7.9	1.0	1.6	0.5-4.0
	2006-2010	35.0	27.7	23.7-32.3	20.6	35.8	29.0-43.6	14.4	21.7	16.8-27.6	3.8	2.9	1.7-4.6	2.6	4.2	2.2-7.4	1.2	1.6	0.6-3.8
	2011-2015	40.0	29.8	25.5-34.5	22.2	32.7	26.6-39.9	17.8	27.3	21.6-34.1	4.6	3.1	1.9-4.7	3.4	5.0	2.8-8.2	1.2	1.5	0.5-3.6
2016-2020	47.4	31.7	27.5-36.3	28.8	40.1	33.6-47.6	18.6	26.2	20.7-32.7	5.0	3.2	2.0-4.9	3.2	4.8	2.7-8.1	1.8	2.0	0.8-4.1	
Female breast	1976-1980							44.2	111.8	97.2-127.9							18.0	41.8	33.5-51.6
	1981-1985							47.4	108.2	94.6-123.2							18.8	42.3	34.0-51.9
	1986-1990							63.0	130.0	115.8-145.5							17.8	36.5	29.2-45.0
	1991-1995							75.2	142.8	128.6-158.1							16.2	29.4	23.3-36.6
	1996-2000							82.0	147.0	133.1-162.1							18.2	30.6	24.6-37.7
	2001-2005							84.2	137.8	124.8-151.8							14.6	23.1	18.0-29.2
	2006-2010							88.8	130.3	118.1-143.4							12.6	17.5	13.3-22.7
	2011-2015							94.8	126.6	115.0-139.2							13.8	17.4	13.3-22.4
2016-2020							122.0	151.3	138.7-164.8							15.6	17.4	13.6-22.1	
Cervix uteri	1976-1980							5.0	11.0	7.1-16.3							0.8	1.9	0.5-4.8
	1981-1985							4.2	8.8	5.4-13.5							1.4	2.9	1.2-6.0
	1986-1990							6.0	12.7	8.5-18.2							1.6	3.5	1.5-6.9
	1991-1995							5.6	11.1	7.3-16.1							1.0	2.0	0.6-4.7
	1996-2000							4.0	7.5	4.6-11.6							1.4	2.6	1.0-5.3
	2001-2005							3.2	5.8	3.2-9.5							1.2	1.9	0.7-4.3
	2006-2010							3.8	5.9	3.5-9.4							0.8	1.2	0.3-3.2
	2011-2015							4.2	7.7	4.6-12.1							1.4	1.9	0.7-4.2
	2016-2020							4.2	7.0	4.1-11.2							1.4	1.3	0.5-3.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Ontario**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							16.4	37.1	29.4-46.2							3.8	8.5	5.1-13.4
	1981-1985							14.8	31.6	24.7-39.9							3.2	6.2	3.5-10.2
	1986-1990							11.6	23.4	17.7-30.4							1.8	3.2	1.5-6.3
	1991-1995							17.4	33.8	27.0-41.8							3.4	5.9	3.4-9.5
	1996-2000							14.0	25.4	19.8-32.1							2.8	4.8	2.6-8.1
	2001-2005							17.8	29.3	23.4-36.1							2.8	4.2	2.3-7.2
	2006-2010							20.6	28.7	23.3-35.0							3.8	5.2	3.1-8.3
	2011-2015							27.8	33.7	28.2-40.1							2.2	2.2	1.1-4.3
	2016-2020							31.2	35.1	29.6-41.4							4.8	4.9	3.1-7.7
Ovary	1976-1980							8.6	20.2	14.5-27.3							8.0	18.0	12.8-24.6
	1981-1985							9.8	21.2	15.6-28.2							6.2	13.5	9.1-19.3
	1986-1990							9.2	18.7	13.6-25.1							7.2	13.9	9.7-19.4
	1991-1995							9.2	17.8	13.0-23.8							5.8	10.9	7.2-15.7
	1996-2000							11.6	21.0	15.9-27.2							4.2	7.2	4.4-11.1
	2001-2005							7.2	11.8	8.2-16.5							5.4	8.7	5.7-12.8
	2006-2010							11.2	16.2	12.2-21.3							7.2	9.8	6.8-13.8
	2011-2015							11.6	15.6	11.7-20.5							7.2	8.5	5.9-12.1
	2016-2020							5.4	7.2	4.6-10.9							6.2	6.8	4.6-10.0
Prostate	1976-1980				28.2	90.7	75.9-107.4							8.2	29.0	20.6-39.5			
	1981-1985				29.8	84.8	71.3-100.1							8.4	26.1	18.6-35.4			
	1986-1990				38.8	104.2	89.5-120.5							12.0	36.4	27.5-47.0			
	1991-1995				79.2	178.6	161.1-197.5							14.4	37.8	29.3-47.8			
	1996-2000				89.6	196.8	178.7-216.1							11.4	28.7	21.6-37.3			
	2001-2005				79.8	155.3	140.2-171.6							11.8	27.3	20.7-35.3			
	2006-2010				104.0	170.1	155.5-185.8							13.0	25.6	19.7-32.8			
	2011-2015				83.8	116.3	105.1-128.5							10.6	17.7	13.2-23.3			
	2016-2020				135.4	166.0	153.4-179.5							10.6	15.7	11.7-20.8			
Testis	1976-1980				3.2	7.4	4.1-12.3							0.4	0.8	0.1-3.3			
	1981-1985				1.6	3.6	1.5-7.2							0.2	0.4	0.0-2.5			
	1986-1990				3.2	6.6	3.7-11.0							0.6	1.8	0.3-5.1			
	1991-1995				3.4	6.9	4.0-11.2							0.2	0.4	0.0-2.3			
	1996-2000				2.6	5.4	2.9-9.3							0.6	1.3	0.3-3.8			
	2001-2005				3.2	6.3	3.6-10.3							0.0	0.0	0.0-1.6			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Ontario**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				4.2	9.6	5.9-14.7							0.2	0.5	0.0-2.5			
	2011-2015				3.6	8.1	4.7-12.9							0.4	1.0	0.1-3.5			
	2016-2020				3.8	8.8	5.2-13.7							0.0	0.0	0.0-1.5			
Urinary bladder (incl. in situ)	1976-1980	15.0	18.9	14.8-23.8	11.2	31.8	23.7-41.6	3.8	8.6	5.1-13.4	4.6	6.2	3.9-9.3	3.4	11.5	6.5-18.4	1.2	2.8	1.0-6.0
	1981-1985	20.6	23.6	19.2-28.7	15.2	40.8	31.9-51.4	5.4	10.8	7.1-15.9	5.4	6.4	4.2-9.3	4.4	12.2	7.5-18.6	1.0	2.1	0.7-4.9
	1986-1990	21.4	23.4	19.2-28.3	16.6	41.1	32.6-51.2	4.8	9.2	5.9-13.8	3.2	3.5	2.0-5.7	2.4	6.2	3.1-10.9	0.8	1.5	0.4-4.0
	1991-1995	25.2	25.4	21.1-30.3	18.6	45.1	36.1-55.5	6.6	11.8	8.1-16.7	6.0	6.0	4.1-8.6	4.4	11.4	7.0-17.4	1.6	2.9	1.2-5.8
	1996-2000	29.6	28.3	24.0-33.3	21.2	47.9	39.1-58.1	8.4	13.9	10.0-18.9	4.8	4.6	2.9-6.8	3.2	7.6	4.3-12.3	1.6	2.3	1.0-4.7
	2001-2005	26.2	22.6	18.9-26.9	20.4	42.4	34.4-51.6	5.8	8.5	5.7-12.4	6.6	5.7	3.9-8.1	5.0	11.0	7.1-16.2	1.6	2.2	0.9-4.6
	2006-2010	34.2	26.1	22.3-30.5	25.4	44.0	36.5-52.5	8.8	12.1	8.7-16.5	7.6	5.5	3.9-7.7	5.0	9.3	6.0-13.8	2.6	3.2	1.7-5.7
	2011-2015	39.4	27.4	23.6-31.7	29.0	46.5	39.1-55.0	10.4	13.3	9.9-17.8	6.8	4.6	3.2-6.6	4.2	6.9	4.3-10.8	2.6	3.1	1.6-5.7
	2016-2020	43.0	26.1	22.7-30.0	35.8	48.4	41.4-56.5	7.2	8.5	5.9-12.1	8.4	4.9	3.5-6.7	7.0	10.3	7.1-14.5	1.4	1.4	0.5-3.2
Kidney and renal pelvis	1976-1980	6.0	7.8	5.2-11.2	3.4	10.1	5.7-16.4	2.6	5.9	3.1-10.2	2.8	3.7	2.0-6.3	1.8	5.5	2.4-10.6	1.0	2.5	0.8-5.7
	1981-1985	6.0	6.9	4.6-9.9	3.4	9.3	5.3-15.0	2.6	5.1	2.7-8.9	2.4	2.9	1.5-5.0	1.0	3.4	1.1-7.6	1.4	2.7	1.1-5.7
	1986-1990	9.8	10.9	8.0-14.4	6.2	15.4	10.4-22.0	3.6	6.8	4.0-10.8	5.2	5.5	3.6-8.1	3.4	8.0	4.6-13.0	1.8	3.2	1.5-6.2
	1991-1995	15.2	15.4	12.1-19.3	9.6	20.8	15.3-27.7	5.6	11.1	7.4-16.1	5.8	6.0	4.0-8.6	4.0	9.7	5.9-15.1	1.8	3.4	1.5-6.4
	1996-2000	13.4	12.8	9.9-16.3	9.0	19.0	13.8-25.5	4.4	7.9	4.9-12.0	6.2	5.9	4.0-8.4	4.4	9.7	6.1-14.8	1.8	2.9	1.3-5.7
	2001-2005	20.6	18.4	15.0-22.3	11.6	22.3	16.9-29.0	9.0	14.5	10.5-19.6	3.6	3.2	1.9-5.1	1.6	3.3	1.4-6.6	2.0	3.0	1.4-5.7
	2006-2010	21.8	16.6	13.5-20.1	15.0	24.7	19.3-31.2	6.8	9.5	6.5-13.5	5.6	4.4	2.9-6.5	3.4	6.5	3.7-10.4	2.2	3.1	1.5-5.8
	2011-2015	25.4	17.7	14.6-21.2	18.2	27.3	21.8-33.9	7.2	9.0	6.2-12.8	6.2	4.1	2.8-6.0	4.4	6.9	4.2-10.6	1.8	1.7	0.8-3.6
	2016-2020	28.8	19.3	16.1-23.0	18.8	25.7	20.5-32.0	10.0	14.0	10.1-18.9	4.4	2.6	1.6-4.0	3.8	5.0	3.0-8.1	0.6	0.6	0.1-2.1
Brain and other central nervous system	1976-1980	3.6	4.1	2.4-6.5	1.6	3.9	1.7-7.8	2.0	4.3	2.0-8.0	2.6	3.1	1.6-5.3	1.4	3.3	1.3-7.0	1.2	2.9	1.0-6.3
	1981-1985	5.4	6.1	4.0-8.9	2.6	6.3	3.3-10.9	2.8	5.8	3.2-9.8	2.6	3.0	1.6-5.2	1.4	3.3	1.3-6.9	1.2	2.4	0.9-5.3
	1986-1990	5.0	5.2	3.4-7.7	3.0	6.4	3.6-10.7	2.0	4.1	1.9-7.6	3.4	3.4	2.0-5.5	2.0	4.1	2.0-7.7	1.4	2.8	1.1-5.9
	1991-1995	8.6	8.7	6.3-11.8	5.0	10.6	6.8-15.6	3.6	7.3	4.3-11.5	5.2	5.4	3.5-7.9	3.6	8.2	4.8-13.1	1.6	3.2	1.4-6.3
	1996-2000	7.6	7.4	5.2-10.1	3.4	7.1	4.1-11.4	4.2	7.7	4.7-11.8	5.4	5.2	3.4-7.5	2.8	5.7	3.1-9.6	2.6	4.3	2.3-7.5
	2001-2005	8.4	7.5	5.4-10.2	5.8	11.3	7.5-16.4	2.6	4.0	2.1-7.0	5.0	4.3	2.8-6.4	2.6	4.8	2.5-8.3	2.4	3.7	1.9-6.6
	2006-2010	9.8	8.4	6.1-11.2	5.8	10.8	7.1-15.7	4.0	6.1	3.5-9.7	5.4	4.3	2.8-6.3	3.2	5.6	3.2-9.3	2.2	3.0	1.5-5.6
	2011-2015	9.4	7.9	5.7-10.6	5.6	9.3	6.1-13.7	3.8	6.7	3.9-10.7	7.6	5.7	4.0-8.0	5.0	7.6	4.8-11.6	2.6	4.1	2.1-7.2
	2016-2020	9.2	6.5	4.7-8.9	5.4	8.0	5.2-12.0	3.8	5.5	3.2-9.0	6.8	4.4	3.0-6.4	4.4	6.0	3.6-9.4	2.4	3.1	1.5-5.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Ontario**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	3.4	4.0	2.3-6.5	0.6	1.6	0.3-4.7	2.8	6.0	3.3-10.2	1.0	1.3	0.4-3.0	0.2	0.5	0.0-3.0	0.8	1.7	0.5-4.5
	1981-1985	3.0	3.3	1.8-5.5	0.8	1.7	0.4-4.5	2.2	4.8	2.3-8.7	0.4	0.4	0.1-1.6	0.4	0.9	0.1-3.4	0.0	0.0	0.0-1.7
	1986-1990	4.2	4.5	2.7-6.8	1.2	2.5	0.9-5.7	3.0	6.4	3.5-10.6	0.4	0.4	0.1-1.6	0.4	1.0	0.1-3.6	0.0	0.0	0.0-1.6
	1991-1995	2.2	2.2	1.1-3.9	0.4	0.8	0.1-2.9	1.8	3.5	1.6-6.7	0.0	0.0	0.0-0.8	0.0	0.0	0.0-1.7	0.0	0.0	0.0-1.5
	1996-2000	4.0	3.9	2.4-6.0	1.6	3.1	1.3-6.1	2.4	4.7	2.4-8.2	0.2	0.2	0.0-1.1	0.2	0.5	0.0-2.5	0.0	0.0	0.0-1.4
	2001-2005	7.4	7.1	5.0-9.8	2.6	5.0	2.7-8.7	4.8	9.0	5.8-13.5	0.2	0.2	0.0-1.1	0.2	0.4	0.0-2.3	0.0	0.0	0.0-1.4
	2006-2010	14.4	12.7	9.9-16.2	3.6	6.9	4.0-11.1	10.8	18.3	13.6-24.1	0.4	0.3	0.0-1.2	0.4	0.6	0.1-2.6	0.0	0.0	0.0-1.4
	2011-2015	14.0	11.9	9.2-15.3	4.2	7.1	4.3-11.2	9.8	16.8	12.2-22.5	0.4	0.3	0.0-1.1	0.0	0.0	0.0-1.5	0.4	0.4	0.0-2.1
	2016-2020	14.4	13.2	10.2-16.8	4.0	7.2	4.3-11.4	10.4	18.9	13.9-25.2	1.2	0.7	0.3-1.8	0.4	0.7	0.1-2.6	0.8	0.8	0.2-2.5
Hodgkin lymphoma	1976-1980	3.0	3.0	1.7-5.1	1.0	2.3	0.7-5.6	2.0	3.7	1.8-7.0	1.2	1.4	0.5-3.1	0.8	1.8	0.5-4.9	0.4	0.9	0.1-3.4
	1981-1985	2.8	3.3	1.8-5.5	1.0	2.2	0.7-5.5	1.8	4.0	1.8-7.7	0.6	0.7	0.1-2.0	0.4	1.0	0.1-3.6	0.2	0.3	0.0-2.2
	1986-1990	3.6	3.7	2.2-5.9	1.8	3.9	1.8-7.6	1.8	3.3	1.5-6.3	0.6	0.6	0.1-1.9	0.2	0.7	0.0-3.2	0.4	0.8	0.1-2.9
	1991-1995	3.8	3.9	2.3-6.1	2.6	5.5	2.9-9.5	1.2	2.4	0.9-5.2	1.4	1.4	0.6-2.9	0.8	1.6	0.4-4.3	0.6	1.2	0.2-3.6
	1996-2000	3.4	3.5	2.0-5.6	1.8	3.8	1.7-7.2	1.6	3.4	1.5-6.7	0.6	0.6	0.1-1.7	0.6	1.2	0.3-3.7	0.0	0.0	0.0-1.4
	2001-2005	2.6	2.6	1.4-4.5	1.2	2.7	1.0-5.8	1.4	2.7	1.1-5.7	0.2	0.2	0.0-1.2	0.0	0.0	0.0-1.6	0.2	0.5	0.0-2.4
	2006-2010	3.4	3.1	1.8-5.1	1.8	3.4	1.5-6.6	1.6	2.9	1.2-5.8	0.8	0.5	0.1-1.5	0.8	1.1	0.3-3.1	0.0	0.0	0.0-1.4
	2011-2015	2.2	2.0	1.0-3.7	1.2	2.5	0.9-5.5	1.0	1.4	0.4-3.6	0.4	0.3	0.0-1.3	0.0	0.0	0.0-1.5	0.4	0.6	0.1-2.4
	2016-2020	5.2	4.6	2.9-6.8	2.4	4.3	2.2-7.7	2.8	5.0	2.6-8.6	0.8	0.5	0.1-1.4	0.2	0.3	0.0-2.1	0.6	0.6	0.1-2.2
Non-Hodgkin lymphomas	1976-1980	10.8	13.4	10.0-17.6	6.4	18.5	12.5-26.3	4.4	9.4	5.9-14.3	6.4	8.3	5.7-11.8	3.0	9.9	5.4-16.2	3.4	7.4	4.3-12.0
	1981-1985	10.4	12.0	8.9-15.8	4.8	11.7	7.4-17.5	5.6	12.3	8.1-17.9	5.8	6.9	4.6-9.9	3.2	8.8	4.9-14.5	2.6	5.3	2.8-9.1
	1986-1990	11.8	12.9	9.8-16.7	6.6	15.4	10.5-21.8	5.2	10.6	6.9-15.6	5.8	6.1	4.1-8.8	3.4	8.0	4.6-13.0	2.4	4.6	2.4-8.1
	1991-1995	15.4	15.5	12.3-19.5	7.6	16.9	11.9-23.3	7.8	14.1	10.0-19.4	8.8	9.0	6.6-12.2	4.8	11.6	7.3-17.3	4.0	7.1	4.3-11.0
	1996-2000	26.8	25.7	21.5-30.4	12.6	27.4	21.0-35.1	14.2	24.8	19.3-31.3	9.2	8.8	6.4-11.7	4.0	9.5	5.7-14.7	5.2	8.3	5.4-12.3
	2001-2005	20.0	17.9	14.5-21.8	10.8	21.0	15.7-27.5	9.2	14.6	10.7-19.7	8.6	7.6	5.5-10.3	3.4	7.0	4.1-11.2	5.2	7.7	5.0-11.5
	2006-2010	29.8	23.8	20.0-28.1	17.2	29.7	23.6-36.9	12.6	18.6	14.1-24.1	7.0	5.6	3.9-7.8	3.6	6.5	3.8-10.4	3.4	4.7	2.7-7.8
	2011-2015	27.0	18.6	15.5-22.2	15.6	23.9	18.7-30.1	11.4	14.2	10.6-18.8	8.4	5.7	4.1-7.8	5.4	8.7	5.7-12.8	3.0	3.3	1.8-5.8
	2016-2020	33.6	21.1	17.9-24.7	17.6	23.8	19.0-29.6	16.0	19.0	14.9-24.1	10.6	6.7	5.0-8.9	5.2	7.4	4.8-11.1	5.4	5.9	3.8-9.0
Myeloma	1976-1980	3.8	4.8	2.9-7.5	1.6	4.4	1.9-8.8	2.2	4.9	2.4-8.8	2.0	2.9	1.4-5.2	1.2	4.2	1.5-8.8	0.8	1.8	0.5-4.6
	1981-1985	2.6	2.9	1.5-4.9	1.4	3.5	1.4-7.2	1.2	2.2	0.8-5.0	2.0	2.2	1.0-4.1	0.8	1.9	0.5-5.0	1.2	2.3	0.9-5.2
	1986-1990	3.6	3.9	2.3-6.2	1.8	4.4	2.0-8.5	1.8	3.5	1.6-6.8	2.4	2.5	1.3-4.4	1.4	3.1	1.2-6.5	1.0	1.8	0.6-4.3
	1991-1995	5.2	5.3	3.4-7.7	3.4	8.6	4.9-13.8	1.8	3.2	1.4-6.1	3.2	3.2	1.8-5.2	2.0	5.2	2.4-9.5	1.2	2.0	0.7-4.5
	1996-2000	6.4	6.1	4.2-8.7	2.8	6.5	3.5-10.9	3.6	6.1	3.6-9.8	4.0	3.8	2.3-5.9	2.6	5.8	3.1-9.9	1.4	2.3	0.9-4.9
	2001-2005	6.6	5.6	3.8-7.9	4.4	8.7	5.4-13.3	2.2	3.3	1.6-6.0	3.0	2.5	1.4-4.2	1.4	3.3	1.3-6.7	1.6	2.3	1.0-4.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Ontario**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	6.4	5.0	3.4-7.2	3.2	5.6	3.2-9.3	3.2	4.3	2.4-7.2	4.0	3.1	1.9-4.9	2.0	3.7	1.8-6.9	2.0	2.6	1.2-5.0
	2011-2015	10.8	7.8	5.8-10.3	5.8	9.1	6.0-13.3	5.0	6.9	4.4-10.4	3.8	2.4	1.5-3.9	1.8	2.5	1.1-5.1	2.0	2.5	1.2-4.9
	2016-2020	10.2	6.2	4.6-8.3	6.8	9.0	6.1-12.8	3.4	3.8	2.2-6.4	5.6	3.4	2.2-5.1	3.4	4.7	2.7-7.7	2.2	2.4	1.2-4.7
Leukemias	1976-1980	10.2	13.1	9.7-17.3	5.6	16.7	10.9-24.4	4.6	10.5	6.6-15.8	8.0	10.2	7.2-13.9	5.0	15.7	9.9-23.3	3.0	6.6	3.7-11.0
	1981-1985	11.6	13.4	10.1-17.3	6.4	16.7	11.2-23.8	5.2	11.1	7.2-16.4	7.6	9.2	6.4-12.6	4.0	11.2	6.7-17.4	3.6	7.4	4.3-11.7
	1986-1990	14.4	16.1	12.6-20.3	9.4	23.5	17.1-31.4	5.0	10.3	6.6-15.2	7.8	8.8	6.2-12.0	5.4	14.5	9.4-21.2	2.4	4.7	2.4-8.3
	1991-1995	16.6	16.8	13.4-20.9	10.6	24.3	18.1-31.9	6.0	11.2	7.5-16.1	8.2	8.1	5.8-11.0	5.2	11.5	7.4-16.9	3.0	5.6	3.1-9.4
	1996-2000	18.8	18.0	14.6-22.1	10.8	24.8	18.5-32.4	8.0	13.6	9.7-18.6	8.0	7.7	5.5-10.5	4.2	9.7	6.0-14.8	3.8	6.3	3.8-9.9
	2001-2005	18.0	15.8	12.7-19.5	9.0	18.2	13.2-24.4	9.0	14.0	10.2-19.0	11.2	9.8	7.4-12.8	6.4	13.9	9.5-19.7	4.8	7.4	4.7-11.2
	2006-2010	19.4	14.9	12.0-18.3	11.2	18.8	14.1-24.7	8.2	11.9	8.4-16.4	8.4	6.8	4.8-9.2	5.2	9.7	6.3-14.3	3.2	4.5	2.5-7.6
	2011-2015	27.2	19.2	16.0-22.9	18.6	29.3	23.5-36.3	8.6	10.5	7.5-14.5	11.8	8.3	6.3-10.8	8.0	13.2	9.4-18.2	3.8	4.6	2.8-7.5
	2016-2020	29.0	18.7	15.7-22.3	17.6	24.7	19.6-30.7	11.4	14.2	10.5-18.9	12.4	7.4	5.6-9.6	6.4	9.0	6.1-12.9	6.0	6.4	4.3-9.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Orange**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	826.4	381.7	369.9-393.7	423.6	462.1	441.5-483.3	402.8	333.8	319.1-348.9	498.0	233.3	224.0-242.8	266.6	299.5	282.7-316.9	231.4	191.3	180.3-202.8
	1981-1985	1,000.0	430.4	418.4-442.7	507.0	518.2	497.3-539.8	493.0	378.4	363.4-393.8	551.6	241.5	232.4-250.8	289.2	308.6	292.0-325.7	262.4	199.4	188.6-210.6
	1986-1990	1,076.8	432.1	420.5-443.9	528.4	512.6	492.3-533.5	548.4	391.9	377.2-407.0	580.2	237.7	229.0-246.5	299.6	304.6	288.7-321.2	280.6	198.7	188.4-209.5
	1991-1995	1,341.0	499.4	487.5-511.6	667.4	598.2	577.3-619.6	673.6	446.3	431.3-461.7	614.8	234.7	226.4-243.1	317.4	304.2	288.8-320.2	297.4	195.5	185.6-205.7
	1996-2000	1,523.6	524.9	513.1-536.9	752.8	606.6	586.9-626.8	770.8	475.4	460.5-490.7	606.6	213.9	206.4-221.7	295.4	257.5	244.2-271.4	311.2	188.6	179.3-198.2
	2001-2005	1,693.4	524.9	513.6-536.3	849.0	603.4	584.8-622.5	844.4	473.4	459.1-488.0	620.8	199.3	192.3-206.5	299.0	234.2	222.1-246.7	321.8	178.5	169.8-187.5
	2006-2010	1,853.2	522.1	511.3-533.0	943.4	592.1	574.7-609.9	909.8	473.8	459.9-487.9	612.4	178.7	172.4-185.3	297.4	204.5	193.9-215.6	315.0	161.9	153.8-170.2
	2011-2015	1,960.2	501.8	491.6-512.0	969.8	540.4	524.7-556.5	990.4	478.9	465.3-492.7	638.8	168.7	162.8-174.8	319.8	195.4	185.5-205.5	319.0	150.3	142.8-158.0
	2016-2020	1,987.0	466.0	456.7-475.5	999.6	500.1	485.8-514.6	987.4	446.2	433.5-459.2	768.0	181.9	176.1-187.9	373.2	203.8	194.4-213.6	394.8	169.8	162.2-177.6
Oral cavity and pharynx	1976-1980	24.8	10.9	9.1-13.1	18.2	17.9	14.3-22.2	6.6	5.2	3.6-7.4	11.8	5.3	4.0-6.9	8.6	9.3	6.6-12.6	3.2	2.5	1.4-4.1
	1981-1985	29.0	12.4	10.4-14.6	22.2	21.7	17.7-26.2	6.8	5.1	3.5-7.2	12.4	5.3	4.0-6.8	9.8	10.1	7.3-13.5	2.6	1.8	1.0-3.2
	1986-1990	26.2	10.3	8.6-12.3	16.2	14.3	11.2-17.8	10.0	7.2	5.3-9.5	9.4	3.8	2.8-5.0	7.0	6.1	4.2-8.6	2.4	1.7	0.9-3.0
	1991-1995	29.4	10.9	9.2-12.8	19.6	16.8	13.6-20.6	9.8	6.7	4.9-8.8	8.8	3.3	2.4-4.5	6.0	5.4	3.6-7.7	2.8	1.9	1.0-3.2
	1996-2000	28.8	9.7	8.2-11.5	17.6	12.9	10.3-16.0	11.2	7.0	5.2-9.0	7.2	2.5	1.8-3.5	4.4	3.5	2.2-5.3	2.8	1.7	0.9-2.9
	2001-2005	29.4	8.8	7.4-10.3	21.4	13.4	10.9-16.3	8.0	4.6	3.3-6.3	8.2	2.5	1.8-3.5	5.2	3.2	2.1-4.8	3.0	1.7	1.0-2.8
	2006-2010	37.6	10.5	9.0-12.1	28.0	17.2	14.4-20.4	9.6	5.1	3.7-6.8	6.0	1.7	1.1-2.4	4.2	2.7	1.6-4.1	1.8	0.9	0.4-1.8
	2011-2015	42.2	10.2	8.8-11.7	32.4	17.0	14.4-19.9	9.8	4.4	3.2-5.8	7.4	1.9	1.3-2.6	5.6	3.1	2.0-4.6	1.8	0.8	0.4-1.6
	2016-2020	49.2	11.4	10.0-13.0	34.8	17.1	14.5-19.9	14.4	6.5	5.0-8.2	10.2	2.3	1.7-3.0	7.0	3.6	2.5-5.1	3.2	1.2	0.7-2.1
Esophagus	1976-1980	9.4	4.3	3.1-5.7	6.6	6.9	4.7-9.9	2.8	2.3	1.2-3.8	9.4	4.2	3.0-5.6	7.4	7.3	5.0-10.2	2.0	1.7	0.8-3.1
	1981-1985	10.6	4.3	3.2-5.7	7.8	7.5	5.3-10.5	2.8	2.0	1.1-3.4	10.8	4.5	3.3-5.8	8.0	8.1	5.6-11.2	2.8	2.0	1.1-3.4
	1986-1990	13.4	5.5	4.2-6.9	9.4	8.9	6.4-11.9	4.0	2.8	1.7-4.3	11.4	4.6	3.5-5.9	8.6	8.2	5.9-11.1	2.8	1.9	1.0-3.2
	1991-1995	14.4	5.5	4.3-6.9	9.6	8.2	6.0-10.9	4.8	3.0	1.9-4.5	12.2	4.6	3.5-5.9	9.4	8.0	5.9-10.7	2.8	1.8	1.0-3.1
	1996-2000	14.8	5.2	4.1-6.5	10.8	8.8	6.5-11.5	4.0	2.4	1.5-3.7	12.8	4.5	3.4-5.7	9.8	7.9	5.8-10.4	3.0	1.8	1.0-2.9
	2001-2005	15.4	4.9	3.8-6.1	12.6	9.1	7.0-11.7	2.8	1.5	0.8-2.5	15.0	4.8	3.8-6.0	12.8	9.8	7.5-12.6	2.2	1.2	0.6-2.1
	2006-2010	22.2	6.4	5.2-7.7	16.4	11.0	8.7-13.8	5.8	2.9	1.9-4.1	13.0	3.7	2.8-4.7	10.6	6.7	5.0-8.9	2.4	1.2	0.6-2.2
	2011-2015	19.2	4.9	3.9-6.0	15.0	8.4	6.5-10.6	4.2	1.9	1.2-3.0	15.8	4.1	3.2-5.1	11.6	6.5	4.9-8.6	4.2	1.9	1.2-3.0
	2016-2020	21.4	5.0	4.1-6.1	17.8	9.2	7.4-11.5	3.6	1.5	0.9-2.3	19.0	4.4	3.5-5.4	14.6	7.6	5.9-9.6	4.4	1.9	1.2-2.9
Stomach	1976-1980	25.2	11.8	9.8-14.1	16.4	18.3	14.4-23.0	8.8	7.1	5.2-9.6	21.4	10.2	8.3-12.4	13.2	15.3	11.7-19.7	8.2	6.6	4.7-9.0
	1981-1985	23.0	10.2	8.4-12.2	16.0	17.3	13.5-21.6	7.0	5.4	3.7-7.5	18.0	8.1	6.5-10.0	12.0	13.3	10.0-17.3	6.0	4.6	3.1-6.6
	1986-1990	31.2	12.9	10.9-15.1	18.6	17.9	14.4-22.1	12.6	8.9	6.9-11.5	21.0	8.8	7.2-10.7	11.6	12.0	9.0-15.6	9.4	6.8	5.0-9.0
	1991-1995	27.0	10.2	8.6-12.1	18.0	16.4	13.1-20.2	9.0	6.0	4.3-8.0	17.2	6.6	5.3-8.1	10.4	10.2	7.5-13.4	6.8	4.4	3.0-6.1
	1996-2000	31.8	11.1	9.4-13.0	20.8	17.0	13.8-20.7	11.0	6.5	4.9-8.5	16.4	5.7	4.6-7.1	10.2	8.8	6.5-11.6	6.2	3.7	2.5-5.2
	2001-2005	27.6	8.9	7.5-10.6	16.6	13.2	10.5-16.5	11.0	6.1	4.6-8.0	16.2	5.3	4.2-6.6	9.4	8.0	5.8-10.6	6.8	3.7	2.5-5.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Orange**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	24.6	7.4	6.1-8.8	14.6	10.3	8.0-13.0	10.0	5.5	4.1-7.3	10.8	3.2	2.4-4.2	6.4	4.7	3.2-6.7	4.4	2.2	1.4-3.3
	2011-2015	29.0	7.4	6.3-8.8	17.2	10.0	7.9-12.4	11.8	5.4	4.1-7.0	10.6	2.8	2.1-3.7	5.4	3.4	2.2-4.9	5.2	2.4	1.6-3.6
	2016-2020	35.4	8.4	7.2-9.8	21.0	10.6	8.6-12.9	14.4	6.6	5.2-8.4	14.6	3.5	2.7-4.4	7.8	4.0	2.8-5.6	6.8	3.1	2.1-4.4
Colorectal	1976-1980	135.8	64.5	59.7-69.7	71.4	83.3	74.4-92.9	64.4	52.7	47.0-58.9	77.4	37.4	33.7-41.4	39.0	46.1	39.5-53.4	38.4	31.7	27.3-36.5
	1981-1985	161.2	71.5	66.6-76.7	73.4	77.7	69.6-86.4	87.8	66.6	60.5-73.2	87.6	39.4	35.8-43.3	40.4	44.7	38.4-51.6	47.2	35.6	31.2-40.5
	1986-1990	148.2	61.0	56.6-65.6	75.6	76.4	68.5-84.8	72.6	50.6	45.5-56.1	79.0	32.9	29.7-36.4	36.8	38.4	32.9-44.6	42.2	29.3	25.5-33.5
	1991-1995	163.6	62.8	58.6-67.3	81.8	77.5	69.9-85.6	81.8	53.6	48.5-59.0	77.0	29.7	26.8-32.9	39.4	38.7	33.2-44.6	37.6	24.3	21.0-28.1
	1996-2000	195.8	69.1	64.8-73.6	93.8	80.0	72.7-87.8	102.0	61.7	56.4-67.3	65.8	23.4	21.0-26.1	32.4	30.1	25.5-35.2	33.4	19.7	16.8-22.9
	2001-2005	184.6	58.6	54.9-62.6	90.8	66.5	60.3-73.1	93.8	52.2	47.5-57.2	72.4	23.4	21.1-26.0	33.8	27.1	23.1-31.7	38.6	20.9	18.1-24.1
	2006-2010	168.8	48.6	45.3-52.1	85.0	54.4	49.2-60.1	83.8	43.7	39.5-48.1	50.4	14.9	13.1-16.9	22.4	15.8	12.9-19.1	28.0	14.2	11.9-16.8
	2011-2015	158.4	41.3	38.4-44.4	79.2	46.0	41.4-50.9	79.2	37.9	34.2-42.0	54.6	14.4	12.7-16.2	27.2	16.6	13.8-19.7	27.4	12.6	10.6-15.0
2016-2020	159.4	37.9	35.2-40.7	87.2	44.3	40.1-48.9	72.2	32.5	29.1-36.1	65.6	15.9	14.2-17.8	33.2	18.2	15.5-21.3	32.4	14.2	12.1-16.7	
Colon excluding rectum	1976-1980	98.8	47.4	43.3-51.9	50.2	58.8	51.3-66.9	48.6	39.9	35.0-45.3	62.6	30.1	26.8-33.7	30.4	35.7	29.9-42.1	32.2	26.3	22.4-30.8
	1981-1985	118.6	52.9	48.7-57.4	52.4	56.7	49.7-64.2	66.2	50.3	45.0-56.1	74.0	33.2	29.8-36.8	33.8	37.1	31.4-43.3	40.2	30.3	26.2-34.8
	1986-1990	104.0	43.1	39.4-47.0	50.8	53.2	46.6-60.5	53.2	36.8	32.5-41.6	68.2	28.5	25.5-31.7	31.4	33.4	28.2-39.2	36.8	25.5	21.9-29.5
	1991-1995	122.0	47.0	43.3-50.9	59.2	56.7	50.2-63.8	62.8	41.0	36.5-45.8	66.6	25.8	23.1-28.8	33.0	33.0	28.0-38.6	33.6	21.7	18.5-25.2
	1996-2000	144.6	51.2	47.5-55.1	67.8	58.9	52.6-65.6	76.8	46.1	41.5-50.9	54.2	19.3	17.1-21.7	26.6	24.9	20.8-29.6	27.6	16.2	13.6-19.2
	2001-2005	138.6	44.3	41.0-47.7	65.4	49.0	43.6-54.7	73.2	40.7	36.6-45.1	61.6	20.0	17.8-22.4	28.0	22.5	18.8-26.6	33.6	18.2	15.5-21.2
	2006-2010	119.2	34.5	31.7-37.4	57.6	37.5	33.1-42.3	61.6	31.9	28.4-35.7	39.2	11.5	9.9-13.3	15.8	11.1	8.7-13.9	23.4	11.9	9.8-14.3
	2011-2015	113.6	29.9	27.4-32.5	54.4	32.5	28.6-36.8	59.2	28.0	24.9-31.5	43.4	11.6	10.1-13.3	21.2	13.4	10.9-16.3	22.2	10.2	8.4-12.4
2016-2020	114.4	27.2	25.0-29.6	59.6	30.7	27.2-34.6	54.8	24.5	21.6-27.6	51.2	12.4	10.9-14.0	26.4	14.4	12.0-17.1	24.8	11.0	9.1-13.1	
Rectum & rectosigmoid	1976-1980	37.0	17.1	14.7-19.8	21.2	24.5	19.8-29.9	15.8	12.8	10.1-16.0	14.8	7.3	5.7-9.2	8.6	10.4	7.3-14.2	6.2	5.4	3.6-7.6
	1981-1985	42.6	18.6	16.1-21.3	21.0	21.0	17.0-25.6	21.6	16.4	13.4-19.8	13.6	6.3	4.8-7.9	6.6	7.6	5.1-10.8	7.0	5.3	3.7-7.4
	1986-1990	44.2	17.9	15.6-20.4	24.8	23.1	19.1-27.7	19.4	13.7	11.1-16.7	10.8	4.4	3.3-5.8	5.4	5.0	3.2-7.4	5.4	3.8	2.5-5.5
	1991-1995	41.6	15.8	13.7-18.1	22.6	20.8	17.0-25.1	19.0	12.6	10.2-15.4	10.4	3.9	2.9-5.1	6.4	5.6	3.8-8.0	4.0	2.6	1.6-4.1
	1996-2000	51.2	17.9	15.7-20.2	26.0	21.2	17.6-25.2	25.2	15.6	13.0-18.6	11.6	4.1	3.1-5.3	5.8	5.1	3.4-7.4	5.8	3.5	2.3-5.0
	2001-2005	46.0	14.4	12.6-16.4	25.4	17.6	14.6-21.0	20.6	11.5	9.4-13.9	10.8	3.4	2.6-4.5	5.8	4.7	3.1-6.8	5.0	2.7	1.8-4.0
	2006-2010	49.6	14.1	12.4-16.0	27.4	16.9	14.1-20.1	22.2	11.8	9.7-14.3	11.2	3.4	2.5-4.4	6.6	4.7	3.2-6.6	4.6	2.4	1.5-3.6
	2011-2015	44.8	11.4	9.9-13.1	24.8	13.4	11.1-16.1	20.0	9.9	8.0-12.1	11.2	2.7	2.1-3.6	6.0	3.2	2.1-4.7	5.2	2.4	1.5-3.5
2016-2020	45.0	10.6	9.3-12.2	27.6	13.6	11.4-16.2	17.4	8.0	6.4-9.9	14.4	3.5	2.7-4.4	6.8	3.8	2.6-5.4	7.6	3.3	2.3-4.6	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Orange**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	7.4	3.2	2.2-4.4	3.6	3.9	2.2-6.2	3.8	2.9	1.7-4.5	4.0	1.8	1.1-2.8	2.2	2.3	1.1-4.1	1.8	1.5	0.7-2.8
	1981-1985	8.2	3.4	2.4-4.6	5.2	4.9	3.1-7.4	3.0	2.3	1.3-3.7	6.4	2.8	1.9-3.9	3.2	3.3	1.8-5.5	3.2	2.4	1.4-3.9
	1986-1990	8.2	3.2	2.3-4.4	5.0	4.4	2.8-6.5	3.2	2.3	1.3-3.8	7.0	2.8	1.9-3.8	3.6	3.2	1.9-5.1	3.4	2.4	1.4-3.8
	1991-1995	10.0	3.8	2.8-5.0	6.6	6.2	4.2-8.8	3.4	2.2	1.3-3.5	10.2	3.9	2.9-5.2	6.6	6.5	4.4-9.2	3.6	2.3	1.4-3.6
	1996-2000	14.0	5.0	3.9-6.3	9.0	7.8	5.6-10.5	5.0	3.0	1.9-4.4	14.2	5.0	3.9-6.3	7.0	6.0	4.2-8.4	7.2	4.2	3.0-5.9
	2001-2005	20.8	6.5	5.3-7.9	16.0	11.7	9.2-14.7	4.8	2.6	1.7-3.9	14.0	4.5	3.5-5.7	8.8	6.9	5.0-9.3	5.2	2.8	1.8-4.1
	2006-2010	29.4	7.9	6.6-9.3	23.6	13.6	11.2-16.4	5.8	2.9	2.0-4.2	17.6	4.9	3.9-6.1	12.8	7.8	5.9-10.0	4.8	2.4	1.5-3.6
	2011-2015	40.0	9.7	8.4-11.2	28.0	14.5	12.1-17.3	12.0	5.7	4.3-7.4	26.0	6.6	5.5-7.8	16.4	8.6	6.8-10.8	9.6	4.7	3.4-6.2
	2016-2020	40.8	9.0	7.8-10.4	29.0	13.4	11.3-15.9	11.8	5.2	3.9-6.8	34.8	8.0	6.8-9.3	23.4	11.6	9.6-14.1	11.4	4.9	3.7-6.3
Pancreas	1976-1980	31.8	14.7	12.5-17.2	15.8	17.0	13.3-21.4	16.0	13.0	10.3-16.2	30.2	14.0	11.8-16.5	13.8	15.3	11.7-19.4	16.4	13.1	10.4-16.3
	1981-1985	31.8	13.6	11.5-15.9	15.2	15.2	11.9-19.1	16.6	12.3	9.7-15.2	28.6	12.1	10.2-14.3	12.6	12.1	9.2-15.5	16.0	11.9	9.4-14.8
	1986-1990	28.6	11.8	10.0-13.9	15.4	16.0	12.5-20.2	13.2	9.2	7.1-11.8	28.8	12.0	10.1-14.1	14.8	15.1	11.7-19.1	14.0	9.8	7.6-12.4
	1991-1995	35.6	13.8	11.8-15.9	17.2	16.7	13.2-20.7	18.4	11.9	9.6-14.6	33.6	13.0	11.1-15.1	16.4	16.2	12.8-20.2	17.2	11.0	8.8-13.6
	1996-2000	36.8	13.1	11.2-15.1	17.4	14.6	11.6-18.0	19.4	11.8	9.5-14.4	33.8	12.0	10.3-14.0	14.2	11.8	9.1-14.9	19.6	11.7	9.5-14.3
	2001-2005	38.0	12.1	10.4-13.9	17.4	13.2	10.5-16.3	20.6	11.3	9.2-13.7	32.2	10.3	8.8-12.1	14.4	11.1	8.6-14.1	17.8	9.6	7.7-11.9
	2006-2010	47.4	13.7	12.0-15.6	21.0	13.3	10.8-16.2	26.4	13.9	11.6-16.6	35.8	10.2	8.8-11.9	17.6	11.2	8.9-13.9	18.2	9.4	7.5-11.5
	2011-2015	56.4	14.2	12.6-16.1	27.0	14.6	12.1-17.4	29.4	13.8	11.6-16.2	46.0	11.9	10.4-13.6	20.0	11.6	9.3-14.2	26.0	12.0	10.0-14.3
	2016-2020	64.8	14.8	13.2-16.5	33.6	16.6	14.1-19.4	31.2	13.2	11.2-15.5	62.8	14.7	13.0-16.4	31.4	16.1	13.6-19.0	31.4	13.3	11.3-15.6
Larynx	1976-1980	9.6	4.2	3.1-5.6	8.2	8.2	5.8-11.2	1.4	1.1	0.4-2.3	4.6	2.0	1.3-3.1	4.0	4.2	2.4-6.6	0.6	0.5	0.1-1.5
	1981-1985	15.4	6.5	5.1-8.1	12.8	12.1	9.3-15.6	2.6	1.9	1.0-3.3	6.2	2.6	1.7-3.6	4.8	4.7	3.0-7.1	1.4	1.0	0.4-2.1
	1986-1990	16.4	6.5	5.1-8.0	13.2	12.1	9.2-15.4	3.2	2.5	1.4-4.1	4.6	1.9	1.2-2.9	3.8	3.8	2.2-6.0	0.8	0.6	0.2-1.4
	1991-1995	13.8	5.2	4.0-6.5	10.0	8.4	6.2-11.1	3.8	2.5	1.5-4.0	6.8	2.5	1.7-3.5	5.2	4.3	2.8-6.4	1.6	1.1	0.5-2.1
	1996-2000	13.0	4.5	3.4-5.7	9.8	7.5	5.5-9.9	3.2	2.1	1.2-3.4	3.6	1.3	0.7-2.0	2.4	2.0	1.0-3.5	1.2	0.8	0.3-1.7
	2001-2005	16.6	5.1	4.1-6.4	13.0	8.9	6.9-11.5	3.6	2.0	1.2-3.2	4.2	1.3	0.8-2.0	3.0	2.4	1.3-3.9	1.2	0.6	0.2-1.4
	2006-2010	15.6	4.4	3.4-5.5	11.0	6.6	4.9-8.7	4.6	2.4	1.5-3.7	4.2	1.3	0.8-1.9	3.4	2.4	1.4-3.8	0.8	0.4	0.1-1.0
	2011-2015	16.8	4.3	3.4-5.4	13.6	8.0	6.1-10.2	3.2	1.5	0.9-2.5	6.2	1.6	1.1-2.3	5.4	3.4	2.2-5.0	0.8	0.4	0.1-1.0
	2016-2020	10.0	2.2	1.6-3.0	7.0	3.3	2.3-4.7	3.0	1.3	0.7-2.2	3.0	0.7	0.4-1.2	2.0	1.1	0.5-2.1	1.0	0.4	0.1-1.0
Lung and bronchus	1976-1980	148.8	67.5	62.6-72.6	111.2	118.0	108.0-128.8	37.6	31.2	26.8-36.1	118.8	53.8	49.5-58.4	87.6	93.2	84.2-102.8	31.2	26.1	22.1-30.6
	1981-1985	175.0	74.5	69.6-79.7	116.2	116.1	106.5-126.3	58.8	44.5	39.5-49.9	140.4	59.6	55.2-64.2	96.4	95.6	87.0-104.9	44.0	33.4	29.0-38.1
	1986-1990	191.8	77.3	72.5-82.4	115.0	109.1	100.1-118.7	76.8	55.2	49.8-61.1	161.4	65.4	60.9-70.1	99.4	97.0	88.3-106.2	62.0	44.1	39.3-49.3
	1991-1995	219.6	83.1	78.2-88.2	132.8	120.9	111.6-130.7	86.8	58.0	52.6-63.7	176.4	67.2	62.8-71.8	106.0	98.4	89.9-107.4	70.4	47.1	42.3-52.3
	1996-2000	215.4	75.7	71.2-80.3	118.2	96.9	89.1-105.2	97.2	61.1	55.8-66.8	168.6	59.6	55.6-63.8	95.0	78.8	71.7-86.3	73.6	45.7	41.1-50.6
	2001-2005	234.0	74.6	70.3-79.0	126.0	92.9	85.5-100.6	108.0	62.0	56.9-67.5	168.6	54.4	50.7-58.2	90.8	69.0	62.6-75.8	77.8	44.3	40.0-49.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Orange**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	242.8	71.2	67.2-75.4	118.4	78.5	72.1-85.3	124.4	66.6	61.4-72.1	170.6	50.6	47.2-54.1	90.4	62.4	56.6-68.6	80.2	42.2	38.1-46.6
	2011-2015	253.2	66.7	63.0-70.5	126.6	74.6	68.6-80.8	126.6	60.8	56.1-65.9	168.6	45.2	42.2-48.5	91.4	55.3	50.2-60.8	77.2	37.5	33.8-41.5
	2016-2020	264.2	61.6	58.3-65.1	127.2	65.9	60.7-71.4	137.0	59.2	54.8-63.9	171.8	40.3	37.6-43.2	83.0	44.7	40.4-49.4	88.8	37.8	34.3-41.5
Melanoma of the skin	1976-1980	10.2	4.7	3.5-6.2	4.2	4.2	2.6-6.5	6.0	5.0	3.3-7.2	4.2	2.0	1.2-3.1	2.0	2.0	0.9-3.7	2.2	1.8	0.9-3.3
	1981-1985	16.6	7.1	5.6-8.8	9.6	8.7	6.4-11.5	7.0	5.5	3.8-7.7	5.6	2.4	1.6-3.5	5.0	4.6	2.9-6.8	0.6	0.4	0.1-1.3
	1986-1990	19.0	7.3	5.9-8.9	8.4	7.3	5.2-10.0	10.6	7.6	5.7-10.0	8.8	3.5	2.5-4.7	6.0	5.3	3.5-7.7	2.8	2.1	1.1-3.6
	1991-1995	25.0	9.0	7.5-10.7	12.8	11.1	8.4-14.3	12.2	8.0	6.1-10.3	7.0	2.6	1.8-3.6	3.6	3.0	1.7-4.8	3.4	2.2	1.3-3.5
	1996-2000	40.0	13.0	11.2-14.9	21.8	15.5	12.7-18.8	18.2	11.1	8.9-13.6	7.0	2.4	1.7-3.4	4.6	3.8	2.4-5.7	2.4	1.4	0.7-2.5
	2001-2005	36.8	11.1	9.5-12.8	20.4	13.7	11.1-16.8	16.4	9.3	7.4-11.5	9.2	2.7	2.0-3.7	5.8	4.0	2.6-5.8	3.4	1.9	1.1-3.1
	2006-2010	60.6	16.7	14.8-18.7	36.8	23.1	19.7-26.8	23.8	12.3	10.2-14.8	8.2	2.3	1.7-3.2	5.0	3.2	2.0-4.8	3.2	1.7	1.0-2.8
	2011-2015	80.6	20.3	18.4-22.5	47.6	26.2	22.8-29.8	33.0	15.8	13.4-18.5	10.0	2.7	2.0-3.6	7.0	4.1	2.8-5.7	3.0	1.5	0.8-2.5
	2016-2020	74.8	17.6	15.8-19.5	43.2	22.1	19.1-25.4	31.6	14.4	12.2-16.9	10.8	2.4	1.8-3.2	7.2	3.8	2.6-5.4	3.6	1.5	0.9-2.4
Female breast	1976-1980							104.6	89.4	81.7-97.5							44.8	38.1	33.2-43.5
	1981-1985							125.2	98.5	90.8-106.6							45.6	35.9	31.3-40.9
	1986-1990							150.4	109.4	101.6-117.5							51.8	37.5	33.0-42.4
	1991-1995							184.8	123.5	115.7-131.8							50.8	33.7	29.7-38.1
	1996-2000							222.0	137.8	129.8-146.2							54.0	32.8	29.0-37.0
	2001-2005							243.0	135.2	127.7-143.1							48.2	26.5	23.2-30.1
	2006-2010							253.8	129.9	122.7-137.3							49.0	24.9	21.9-28.3
	2011-2015							274.2	132.6	125.6-140.0							49.0	22.6	19.8-25.7
	2016-2020							291.8	133.6	126.6-140.9							58.0	25.1	22.2-28.2
Cervix uteri	1976-1980							19.0	15.5	12.5-19.0							7.8	6.7	4.7-9.2
	1981-1985							15.4	11.7	9.2-14.6							5.2	4.1	2.7-6.1
	1986-1990							13.4	9.7	7.4-12.3							4.0	2.7	1.7-4.2
	1991-1995							16.4	10.7	8.5-13.3							6.2	4.3	2.9-6.0
	1996-2000							14.6	9.0	7.0-11.3							6.4	3.9	2.7-5.5
	2001-2005							18.6	10.5	8.5-12.9							5.0	2.9	1.9-4.3
	2006-2010							15.6	8.4	6.6-10.5							6.2	3.1	2.1-4.4
	2011-2015							15.8	8.7	6.8-10.9							4.4	2.2	1.3-3.3
	2016-2020							18.0	9.3	7.4-11.5							7.0	3.3	2.3-4.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Orange**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							24.4	20.0	16.5-23.9							6.0	4.8	3.3-6.9
	1981-1985							27.0	21.0	17.5-24.9							10.0	7.4	5.5-9.7
	1986-1990							32.4	23.4	19.9-27.3							6.4	4.6	3.1-6.4
	1991-1995							38.4	26.5	22.9-30.6							6.4	4.2	2.9-6.0
	1996-2000							42.8	27.0	23.5-30.9							7.8	4.6	3.2-6.3
	2001-2005							50.8	28.8	25.3-32.5							10.6	5.9	4.4-7.7
	2006-2010							58.6	29.4	26.1-33.0							10.8	5.5	4.1-7.2
	2011-2015							76.0	34.9	31.4-38.7							11.6	5.4	4.1-7.1
	2016-2020							68.6	29.2	26.2-32.6							20.2	8.6	7.0-10.5
Ovary	1976-1980							20.6	16.8	13.7-20.5							15.2	12.4	9.7-15.6
	1981-1985							20.4	16.1	13.1-19.6							13.8	10.3	8.0-13.0
	1986-1990							24.6	17.8	14.7-21.2							15.4	11.2	8.8-14.0
	1991-1995							28.4	19.0	16.0-22.4							17.4	11.7	9.3-14.4
	1996-2000							27.8	17.5	14.7-20.6							13.4	8.3	6.4-10.5
	2001-2005							26.6	14.8	12.4-17.6							19.4	10.8	8.7-13.2
	2006-2010							29.0	14.9	12.6-17.6							20.2	10.4	8.5-12.7
	2011-2015							26.0	12.2	10.1-14.5							18.8	8.9	7.2-11.0
	2016-2020							25.8	11.6	9.6-13.8							17.2	7.5	6.0-9.4
Prostate	1976-1980				51.4	61.0	53.4-69.4							22.2	31.2	25.4-37.8			
	1981-1985				74.2	87.0	77.9-96.7							24.6	34.3	28.2-41.1			
	1986-1990				84.8	92.0	83.1-101.6							28.0	33.4	27.9-39.6			
	1991-1995				154.0	143.0	132.8-153.7							30.6	34.9	29.4-41.0			
	1996-2000				210.8	172.1	161.6-183.0							31.2	31.5	26.7-36.9			
	2001-2005				245.2	175.3	165.4-185.7							29.8	27.2	22.9-31.9			
	2006-2010				273.6	167.0	158.0-176.5							26.6	21.3	17.7-25.3			
	2011-2015				236.4	123.2	116.0-130.7							28.0	19.5	16.4-23.1			
	2016-2020				251.4	116.9	110.3-123.8							34.2	20.5	17.5-23.9			
Testis	1976-1980				4.4	3.5	2.1-5.3							0.6	0.4	0.1-1.3			
	1981-1985				6.4	4.9	3.3-7.0							1.2	1.2	0.4-2.6			
	1986-1990				6.6	4.0	2.8-5.8							1.4	0.9	0.4-2.0			
	1991-1995				6.8	4.0	2.7-5.6							0.2	0.1	0.0-0.7			
	1996-2000				8.4	5.0	3.6-6.8							0.0	0.0	0.0-0.5			
	2001-2005				13.0	7.5	5.7-9.5							0.0	0.0	0.0-0.5			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Orange**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				13.8	7.9	6.1-10.0							0.4	0.2	0.0-0.9			
	2011-2015				13.0	7.4	5.7-9.5							0.2	0.1	0.0-0.7			
	2016-2020				12.2	7.1	5.4-9.1							0.6	0.3	0.1-0.9			
Urinary bladder (incl. in situ)	1976-1980	42.6	20.1	17.5-23.1	32.0	36.4	30.7-42.8	10.6	8.7	6.5-11.5	14.4	7.1	5.5-9.0	10.2	12.7	9.3-16.9	4.2	3.5	2.2-5.4
	1981-1985	53.2	22.8	20.1-25.8	41.6	42.5	36.6-48.9	11.6	9.0	6.8-11.6	14.6	6.9	5.4-8.6	11.0	13.9	10.3-18.2	3.6	2.7	1.6-4.3
	1986-1990	51.2	21.1	18.6-23.9	36.6	38.5	32.8-44.8	14.6	10.2	8.0-12.9	11.8	5.1	3.9-6.6	7.4	9.0	6.2-12.5	4.4	3.0	1.9-4.6
	1991-1995	58.8	22.2	19.7-24.9	41.8	38.0	32.9-43.7	17.0	11.2	8.9-13.8	14.8	5.8	4.6-7.3	11.4	12.2	9.1-15.9	3.4	2.1	1.2-3.4
	1996-2000	73.6	25.8	23.2-28.6	54.4	45.7	40.3-51.6	19.2	11.8	9.6-14.4	14.8	5.3	4.2-6.6	10.2	9.9	7.3-13.1	4.6	2.7	1.7-4.1
	2001-2005	88.2	28.2	25.6-31.0	66.8	50.4	45.0-56.3	21.4	12.0	9.8-14.5	13.4	4.5	3.5-5.7	7.6	7.0	4.9-9.6	5.8	3.2	2.1-4.6
	2006-2010	95.8	28.1	25.6-30.7	69.2	47.2	42.2-52.6	26.6	14.0	11.7-16.7	20.0	5.9	4.8-7.2	13.0	10.1	7.7-12.9	7.0	3.4	2.4-4.8
	2011-2015	101.2	26.4	24.1-28.9	75.6	45.7	41.0-50.7	25.6	11.9	9.9-14.2	18.8	5.2	4.2-6.4	14.0	9.6	7.5-12.2	4.8	2.3	1.4-3.4
2016-2020	102.2	24.0	21.9-26.2	80.0	42.6	38.4-47.2	22.2	9.5	7.8-11.5	24.4	6.1	5.0-7.3	17.8	11.2	8.9-13.7	6.6	2.7	1.9-3.9	
Kidney and renal pelvis	1976-1980	14.2	6.4	5.0-8.1	8.8	8.7	6.2-11.7	5.4	4.5	2.9-6.5	9.2	4.3	3.1-5.7	4.8	5.0	3.1-7.5	4.4	3.6	2.3-5.5
	1981-1985	22.0	9.3	7.6-11.2	14.2	14.4	11.1-18.3	7.8	5.6	4.0-7.7	11.0	4.9	3.7-6.4	6.4	7.1	4.7-10.1	4.6	3.4	2.2-5.1
	1986-1990	23.0	9.2	7.6-11.0	13.8	12.4	9.5-15.8	9.2	6.7	4.9-8.9	12.2	5.0	3.8-6.4	9.0	9.2	6.5-12.4	3.2	2.2	1.3-3.6
	1991-1995	27.2	10.1	8.5-12.0	17.2	14.2	11.3-17.6	10.0	6.5	4.8-8.6	9.6	3.7	2.7-4.9	6.8	6.1	4.2-8.5	2.8	1.9	1.0-3.1
	1996-2000	41.8	14.2	12.4-16.3	23.6	18.2	15.0-21.9	18.2	11.2	9.0-13.8	11.6	4.1	3.1-5.2	5.8	4.8	3.2-6.8	5.8	3.5	2.4-5.1
	2001-2005	50.2	15.2	13.4-17.2	29.6	19.3	16.2-22.7	20.6	11.6	9.4-14.0	12.8	4.2	3.2-5.3	6.0	4.5	3.0-6.5	6.8	3.8	2.6-5.4
	2006-2010	64.8	18.0	16.0-20.1	43.8	26.3	22.8-30.2	21.0	11.2	9.2-13.6	12.8	3.7	2.8-4.7	8.4	5.3	3.8-7.2	4.4	2.3	1.4-3.5
	2011-2015	74.0	18.7	16.8-20.7	47.4	24.9	21.7-28.4	26.6	13.2	11.0-15.7	14.2	3.8	3.0-4.8	9.0	5.5	3.9-7.4	5.2	2.5	1.6-3.7
2016-2020	75.8	17.4	15.6-19.3	53.2	25.6	22.6-29.0	22.6	9.9	8.2-12.0	13.6	3.2	2.4-4.0	8.4	4.3	3.1-5.9	5.2	2.3	1.5-3.4	
Brain and other central nervous system	1976-1980	13.0	5.5	4.2-7.1	6.4	6.0	4.0-8.6	6.6	5.2	3.6-7.4	8.0	3.3	2.3-4.5	4.2	3.9	2.4-6.0	3.8	2.8	1.7-4.4
	1981-1985	17.2	6.9	5.5-8.6	9.0	7.8	5.6-10.6	8.2	6.2	4.4-8.5	9.2	3.8	2.8-5.0	4.2	3.8	2.3-5.9	5.0	3.8	2.4-5.6
	1986-1990	14.4	5.2	4.1-6.6	8.0	6.0	4.3-8.3	6.4	4.4	3.0-6.3	8.8	3.3	2.4-4.5	4.8	3.7	2.4-5.6	4.0	2.8	1.7-4.3
	1991-1995	20.4	7.1	5.7-8.6	10.6	7.7	5.7-10.1	9.8	6.6	4.9-8.8	10.8	3.9	2.9-5.1	6.2	4.9	3.3-6.9	4.6	3.1	2.0-4.7
	1996-2000	26.8	8.7	7.3-10.4	15.0	10.6	8.3-13.4	11.8	7.2	5.5-9.4	15.4	5.2	4.1-6.5	10.0	7.6	5.6-10.1	5.4	3.3	2.2-4.9
	2001-2005	26.6	7.9	6.6-9.4	14.0	8.7	6.7-11.0	12.6	7.1	5.5-9.2	13.4	4.2	3.2-5.3	6.4	4.3	2.9-6.2	7.0	4.1	2.8-5.7
	2006-2010	26.6	7.2	6.0-8.5	14.6	8.1	6.3-10.3	12.0	6.4	4.8-8.2	16.8	4.5	3.6-5.6	11.0	6.3	4.7-8.2	5.8	2.9	1.9-4.2
	2011-2015	24.2	6.3	5.2-7.6	14.6	8.0	6.2-10.1	9.6	4.9	3.6-6.5	15.2	3.9	3.1-5.0	8.2	4.7	3.3-6.4	7.0	3.3	2.3-4.7
2016-2020	23.0	5.6	4.6-6.8	15.6	8.0	6.3-10.0	7.4	3.7	2.6-5.1	20.6	4.9	4.0-6.0	11.6	5.8	4.3-7.5	9.0	4.1	3.0-5.6	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Orange**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	4.2	1.8	1.1-2.8	1.2	1.4	0.5-3.2	3.0	2.4	1.3-4.0	1.0	0.6	0.2-1.3	0.4	0.7	0.1-2.4	0.6	0.5	0.1-1.5
	1981-1985	9.8	4.0	2.9-5.3	2.4	2.3	1.2-4.1	7.4	5.6	3.9-7.7	2.0	0.9	0.4-1.6	1.0	1.0	0.3-2.3	1.0	0.7	0.2-1.7
	1986-1990	8.8	3.3	2.4-4.4	3.6	2.8	1.6-4.5	5.2	3.8	2.4-5.5	1.2	0.5	0.2-1.0	0.6	0.4	0.1-1.3	0.6	0.4	0.1-1.3
	1991-1995	14.0	4.7	3.7-6.0	3.6	2.6	1.5-4.1	10.4	6.7	5.0-8.8	1.6	0.6	0.3-1.2	0.6	0.4	0.1-1.3	1.0	0.6	0.2-1.4
	1996-2000	35.0	11.0	9.4-12.7	5.8	3.8	2.5-5.4	29.2	17.9	15.1-21.1	1.2	0.4	0.2-0.9	0.6	0.4	0.1-1.2	0.6	0.3	0.1-1.0
	2001-2005	60.4	17.2	15.3-19.3	12.2	7.2	5.5-9.3	48.2	27.2	23.8-30.8	1.4	0.5	0.2-0.9	0.4	0.4	0.0-1.2	1.0	0.6	0.2-1.3
	2006-2010	79.4	21.5	19.4-23.8	19.2	10.3	8.3-12.6	60.2	32.3	28.7-36.3	2.2	0.7	0.3-1.3	0.4	0.3	0.0-0.9	1.8	1.0	0.4-1.9
	2011-2015	91.0	24.6	22.3-27.0	21.8	11.9	9.7-14.4	69.2	37.2	33.3-41.5	1.8	0.5	0.2-1.0	1.2	0.8	0.3-1.7	0.6	0.3	0.1-0.9
	2016-2020	61.4	16.6	14.7-18.6	15.4	7.9	6.2-10.0	46.0	25.2	22.0-28.8	1.8	0.4	0.2-0.9	1.0	0.6	0.2-1.4	0.8	0.3	0.1-0.9
Hodgkin lymphoma	1976-1980	8.2	3.5	2.5-4.7	5.6	4.8	3.1-7.0	2.6	2.3	1.2-3.9	3.4	1.4	0.8-2.3	2.6	2.3	1.2-3.9	0.8	0.6	0.2-1.6
	1981-1985	8.4	3.0	2.2-4.1	5.0	3.4	2.2-5.1	3.4	2.6	1.5-4.3	2.8	1.1	0.6-1.8	1.8	1.4	0.6-2.7	1.0	0.7	0.2-1.7
	1986-1990	13.8	4.6	3.5-5.8	9.2	6.2	4.4-8.4	4.6	3.0	1.9-4.6	3.2	1.1	0.6-1.8	2.4	1.7	0.9-3.0	0.8	0.5	0.1-1.4
	1991-1995	10.2	3.2	2.4-4.2	5.4	3.2	2.1-4.7	4.8	3.0	1.9-4.5	2.4	0.8	0.4-1.4	1.4	0.8	0.3-1.8	1.0	0.7	0.2-1.5
	1996-2000	11.4	3.5	2.6-4.5	6.0	3.5	2.4-5.1	5.4	3.4	2.2-4.9	1.0	0.3	0.1-0.7	0.2	0.1	0.0-0.7	0.8	0.5	0.1-1.2
	2001-2005	14.2	4.2	3.2-5.3	8.8	5.3	3.8-7.2	5.4	3.2	2.1-4.6	1.4	0.4	0.2-0.9	0.6	0.3	0.1-1.1	0.8	0.4	0.1-1.2
	2006-2010	13.2	3.6	2.8-4.6	7.8	4.2	2.9-5.7	5.4	3.1	2.0-4.5	1.6	0.5	0.2-0.9	1.2	0.8	0.3-1.7	0.4	0.2	0.0-0.8
	2011-2015	11.8	3.2	2.4-4.1	7.2	4.0	2.8-5.6	4.6	2.4	1.5-3.7	1.4	0.4	0.1-0.8	1.2	0.7	0.2-1.6	0.2	0.1	0.0-0.5
	2016-2020	10.4	2.7	2.0-3.6	5.2	2.8	1.8-4.2	5.2	2.7	1.7-4.0	0.6	0.1	0.0-0.4	0.4	0.2	0.0-0.6	0.2	0.1	0.0-0.7
Non-Hodgkin lymphomas	1976-1980	24.8	11.2	9.3-13.4	9.6	9.2	6.7-12.3	15.2	12.5	9.8-15.7	16.4	7.5	5.9-9.3	7.6	7.6	5.3-10.5	8.8	7.1	5.2-9.6
	1981-1985	32.0	13.7	11.6-16.0	14.8	13.7	10.6-17.3	17.2	13.3	10.6-16.4	18.2	7.9	6.3-9.7	7.6	7.3	5.0-10.1	10.6	8.2	6.1-10.7
	1986-1990	35.6	14.2	12.2-16.4	18.4	16.8	13.4-20.7	17.2	12.1	9.6-14.9	19.8	8.2	6.6-10.0	10.2	10.5	7.7-13.9	9.6	6.7	4.9-8.8
	1991-1995	51.4	18.8	16.6-21.3	27.2	23.1	19.2-27.5	24.2	15.8	13.1-18.9	26.0	9.8	8.2-11.7	14.6	13.1	10.2-16.6	11.4	7.4	5.6-9.7
	1996-2000	66.0	22.5	20.1-25.1	35.2	27.8	23.8-32.4	30.8	18.9	16.1-22.2	25.6	9.0	7.5-10.7	10.6	9.4	7.0-12.4	15.0	9.2	7.2-11.5
	2001-2005	70.6	22.0	19.7-24.4	37.4	25.9	22.2-30.0	33.2	18.8	16.0-21.9	23.0	7.4	6.1-8.8	11.6	9.3	7.0-12.1	11.4	6.2	4.7-8.0
	2006-2010	76.8	22.0	19.8-24.4	41.4	26.7	23.0-30.7	35.4	18.3	15.7-21.3	23.2	6.9	5.7-8.3	12.4	8.7	6.6-11.2	10.8	5.7	4.3-7.5
	2011-2015	88.8	22.8	20.6-25.0	50.8	28.1	24.6-31.9	38.0	18.4	15.8-21.2	20.4	5.5	4.5-6.7	11.2	7.2	5.4-9.4	9.2	4.3	3.1-5.8
	2016-2020	94.4	22.2	20.2-24.4	50.0	25.9	22.7-29.4	44.4	19.6	17.0-22.4	28.4	6.7	5.7-8.0	16.0	8.9	7.0-11.2	12.4	5.1	3.9-6.6
Myeloma	1976-1980	8.8	4.2	3.0-5.7	4.6	5.2	3.2-7.8	4.2	3.5	2.2-5.4	5.2	2.5	1.6-3.6	3.2	3.6	2.0-5.8	2.0	1.6	0.8-3.0
	1981-1985	11.4	5.1	3.8-6.6	5.8	6.7	4.4-9.8	5.6	4.3	2.9-6.3	8.2	3.7	2.7-5.1	4.4	5.0	3.0-7.6	3.8	2.9	1.8-4.6
	1986-1990	11.6	4.8	3.6-6.1	5.6	5.5	3.6-8.1	6.0	4.3	2.9-6.1	8.6	3.6	2.6-4.9	4.2	4.7	2.8-7.2	4.4	3.1	2.0-4.8
	1991-1995	16.4	6.2	5.0-7.7	7.4	6.6	4.6-9.1	9.0	5.9	4.3-7.8	8.4	3.2	2.3-4.3	3.8	3.6	2.1-5.7	4.6	2.9	1.9-4.4
	1996-2000	17.8	6.3	5.0-7.7	10.0	8.6	6.3-11.3	7.8	4.7	3.3-6.4	11.4	4.1	3.1-5.3	5.4	4.9	3.2-7.1	6.0	3.5	2.4-5.1
	2001-2005	17.0	5.4	4.3-6.7	9.6	7.0	5.1-9.3	7.4	4.0	2.8-5.6	9.2	3.1	2.2-4.1	5.6	4.6	3.0-6.6	3.6	2.0	1.2-3.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Orange**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	24.2	6.9	5.7-8.3	12.8	7.9	6.0-10.1	11.4	6.0	4.6-7.9	10.0	2.9	2.2-3.9	4.6	3.0	1.8-4.5	5.4	2.7	1.8-4.0
	2011-2015	34.0	8.7	7.4-10.1	18.2	10.2	8.2-12.7	15.8	7.5	5.9-9.4	10.6	2.9	2.2-3.8	6.0	3.9	2.6-5.6	4.6	2.2	1.4-3.3
	2016-2020	34.0	8.0	6.8-9.3	17.6	8.9	7.1-11.1	16.4	7.0	5.5-8.7	17.4	4.1	3.3-5.1	8.4	4.6	3.3-6.2	9.0	3.7	2.7-5.0
Leukemias	1976-1980	20.8	9.7	7.9-11.8	11.6	12.6	9.4-16.4	9.2	7.7	5.6-10.3	16.0	7.3	5.7-9.1	8.4	9.3	6.6-12.8	7.6	6.1	4.3-8.4
	1981-1985	27.0	11.3	9.4-13.4	15.4	14.7	11.4-18.5	11.6	9.0	6.8-11.7	17.8	7.7	6.2-9.5	9.2	9.3	6.7-12.7	8.6	6.7	4.8-9.0
	1986-1990	31.4	12.1	10.2-14.1	17.8	16.0	12.7-19.9	13.6	9.2	7.2-11.7	19.0	7.7	6.2-9.4	10.4	10.4	7.6-13.8	8.6	5.9	4.3-8.0
	1991-1995	38.6	14.0	12.0-16.1	19.2	17.1	13.7-21.0	19.4	12.4	10.1-15.2	20.2	7.7	6.2-9.3	10.6	9.9	7.3-13.1	9.6	6.3	4.7-8.4
	1996-2000	39.2	13.3	11.5-15.3	22.2	17.1	14.0-20.7	17.0	10.2	8.2-12.7	21.8	7.6	6.2-9.1	12.6	11.0	8.4-14.1	9.2	5.5	4.0-7.3
	2001-2005	44.4	13.6	11.9-15.6	23.6	16.0	13.1-19.2	20.8	11.6	9.5-14.1	26.4	8.2	6.9-9.7	15.4	11.2	8.7-14.0	11.0	6.1	4.5-7.9
	2006-2010	55.8	15.9	14.1-17.9	32.8	20.8	17.6-24.3	23.0	11.8	9.7-14.2	25.2	7.5	6.3-9.0	13.6	9.2	7.1-11.8	11.6	6.0	4.5-7.8
	2011-2015	61.8	16.3	14.5-18.3	37.8	22.3	19.1-25.8	24.0	11.8	9.8-14.2	23.4	6.3	5.2-7.6	13.6	8.2	6.3-10.5	9.8	4.7	3.5-6.3
	2016-2020	53.0	12.7	11.2-14.4	30.4	15.9	13.4-18.7	22.6	10.4	8.5-12.5	30.2	7.3	6.2-8.6	16.0	9.1	7.1-11.4	14.2	6.2	4.8-7.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Orleans**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	136.0	386.7	357.7-417.5	70.4	457.0	408.8-509.0	65.6	344.7	307.5-385.0	66.0	183.8	164.2-205.0	36.4	230.9	197.3-268.3	29.6	149.1	125.7-175.6
	1981-1985	146.2	399.3	370.5-429.8	76.6	496.4	446.0-550.7	69.6	340.7	305.2-379.2	80.2	221.2	199.8-244.3	43.6	294.8	255.5-338.1	36.6	177.1	152.0-205.2
	1986-1990	164.2	420.0	391.5-450.1	83.4	493.2	445.4-544.5	80.8	378.2	341.6-417.7	81.0	210.1	189.9-231.8	43.4	265.3	230.0-304.2	37.6	174.6	150.1-202.0
	1991-1995	198.0	479.8	450.2-510.8	100.6	565.0	515.5-617.7	97.4	428.6	391.0-468.9	91.4	224.3	204.1-245.9	42.2	251.2	217.7-288.2	49.2	212.5	186.5-241.1
	1996-2000	229.8	539.6	508.8-571.7	119.4	641.6	590.5-695.9	110.4	472.0	433.2-513.5	93.4	219.6	200.1-240.4	50.4	282.8	248.4-320.4	43.0	177.5	154.3-203.2
	2001-2005	227.4	499.3	470.6-529.3	118.0	575.0	529.0-624.0	109.4	442.2	405.6-481.3	85.4	186.9	169.5-205.6	45.0	230.0	200.5-262.5	40.4	157.8	136.6-181.6
	2006-2010	237.4	474.9	447.9-503.1	121.4	529.1	487.0-573.9	116.0	439.1	403.5-477.1	87.0	173.5	157.4-190.9	44.2	201.6	175.4-230.6	42.8	152.7	132.7-175.2
	2011-2015	269.4	514.7	486.9-543.8	138.4	565.2	522.6-610.4	131.0	478.4	441.5-517.8	96.8	184.6	168.3-202.3	49.8	215.7	189.0-245.1	47.0	164.8	144.0-188.0
	2016-2020	292.8	535.1	506.9-564.5	149.4	560.2	519.2-603.8	143.4	523.5	484.1-565.5	101.2	182.5	166.5-199.7	53.8	211.0	185.7-239.0	47.4	162.2	141.6-185.3
Oral cavity and pharynx	1976-1980	4.0	11.5	6.9-17.8	2.8	17.7	9.5-30.0	1.2	6.8	2.4-14.8	1.8	5.1	2.3-9.7	1.4	8.4	3.4-17.4	0.4	2.0	0.2-7.4
	1981-1985	3.8	10.7	6.4-16.7	2.4	15.1	7.6-26.5	1.4	6.8	2.7-14.3	2.2	5.9	2.9-10.6	1.2	8.1	2.9-17.6	1.0	4.6	1.5-10.9
	1986-1990	2.6	7.0	3.7-11.9	1.6	9.8	4.1-19.4	1.0	4.8	1.5-11.3	0.8	2.1	0.6-5.3	0.6	3.2	0.6-9.6	0.2	0.9	0.0-5.3
	1991-1995	3.2	7.9	4.5-12.8	2.0	10.4	4.9-19.2	1.2	5.0	1.8-11.0	1.4	3.4	1.4-6.9	0.8	4.2	1.1-10.8	0.6	2.6	0.5-7.6
	1996-2000	4.6	10.9	6.9-16.3	3.0	15.9	8.8-26.3	1.6	6.5	2.8-12.9	1.2	2.7	1.0-6.0	1.0	4.6	1.5-10.9	0.2	0.7	0.0-4.5
	2001-2005	5.0	10.7	6.9-15.8	3.2	14.3	8.1-23.5	1.8	6.7	3.0-13.1	1.2	2.5	0.9-5.5	0.6	3.0	0.6-8.8	0.6	2.0	0.4-6.5
	2006-2010	2.8	4.9	2.7-8.4	1.8	6.3	2.9-12.5	1.0	3.5	1.1-8.6	1.2	2.3	0.8-5.3	1.0	4.3	1.3-10.4	0.2	0.7	0.0-4.7
	2011-2015	6.8	12.5	8.6-17.8	5.0	19.6	12.5-29.7	1.8	6.3	2.9-12.6	1.4	3.0	1.1-6.4	0.4	2.6	0.3-8.7	1.0	3.5	1.1-8.8
	2016-2020	6.6	11.2	7.6-16.1	3.6	13.0	7.6-21.3	3.0	9.9	5.4-17.0	1.8	3.1	1.4-6.2	1.2	4.0	1.4-9.6	0.6	2.0	0.4-6.9
Esophagus	1976-1980	1.2	3.1	1.1-6.8	1.0	5.5	1.7-13.1	0.2	0.9	0.0-5.4	0.8	2.0	0.5-5.2	0.4	2.0	0.2-7.7	0.4	1.9	0.2-7.2
	1981-1985	1.8	4.9	2.2-9.4	1.4	8.4	3.3-17.4	0.4	1.8	0.2-6.8	1.6	4.7	2.0-9.3	1.6	10.5	4.4-20.7	0.0	0.0	0.0-3.9
	1986-1990	1.2	2.8	1.0-6.3	1.0	5.0	1.6-11.9	0.2	0.9	0.0-5.2	1.2	2.8	1.0-6.2	0.8	3.9	1.1-10.4	0.4	1.7	0.2-6.4
	1991-1995	1.8	4.6	2.1-8.6	1.2	6.6	2.4-14.4	0.6	2.5	0.5-7.6	1.2	3.0	1.1-6.5	0.8	4.5	1.2-11.4	0.4	1.7	0.2-6.3
	1996-2000	2.8	6.6	3.6-11.1	1.8	10.2	4.6-19.3	1.0	4.5	1.5-10.5	2.2	5.1	2.6-9.2	1.6	9.0	3.8-17.7	0.6	2.5	0.5-7.5
	2001-2005	1.8	4.0	1.8-7.6	1.4	6.5	2.6-13.6	0.4	1.7	0.2-6.2	2.6	5.6	3.0-9.7	2.0	9.4	4.4-17.5	0.6	2.5	0.5-7.5
	2006-2010	2.4	5.0	2.6-9.0	1.8	8.5	3.8-16.3	0.6	2.5	0.5-7.6	2.4	4.5	2.3-8.0	2.0	8.1	3.8-15.3	0.4	1.4	0.2-5.6
	2011-2015	2.8	4.7	2.5-8.1	2.0	6.9	3.3-13.3	0.8	2.6	0.7-7.4	3.0	5.9	3.3-10.0	2.2	9.6	4.7-17.6	0.8	3.0	0.8-8.1
	2016-2020	4.6	8.2	5.1-12.7	3.2	11.2	6.2-19.1	1.4	5.2	2.0-11.3	4.0	7.1	4.3-11.3	2.8	9.7	5.2-17.1	1.2	4.5	1.6-10.5
Stomach	1976-1980	2.6	7.8	4.1-13.4	1.2	8.4	3.0-18.1	1.4	7.0	2.7-14.8	2.4	6.7	3.4-11.7	0.8	5.2	1.4-13.3	1.6	7.3	3.1-14.6
	1981-1985	3.4	9.7	5.6-15.5	2.4	17.6	8.9-30.7	1.0	4.3	1.4-10.5	3.2	9.2	5.2-14.9	2.0	14.7	6.9-26.9	1.2	5.3	1.9-11.9
	1986-1990	2.0	4.9	2.3-9.1	1.8	10.1	4.5-19.5	0.2	0.9	0.0-5.2	0.8	1.9	0.5-4.9	0.6	3.2	0.6-9.4	0.2	0.9	0.0-5.2
	1991-1995	4.2	10.2	6.3-15.6	2.2	11.9	5.9-21.4	2.0	8.9	4.2-16.4	2.8	6.9	3.7-11.5	1.0	5.0	1.6-11.8	1.8	8.5	3.8-16.1
	1996-2000	3.0	7.1	4.0-11.7	1.8	9.4	4.3-17.8	1.2	5.0	1.8-11.0	1.8	4.3	2.0-8.1	1.6	8.4	3.6-16.5	0.2	1.0	0.0-5.2
	2001-2005	2.2	4.7	2.4-8.6	1.4	6.4	2.5-13.3	0.8	3.2	0.9-8.5	1.0	2.1	0.7-5.1	0.6	3.0	0.6-8.8	0.4	1.5	0.2-5.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Orleans**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	3.8	7.5	4.5-11.9	3.0	12.9	7.2-21.7	0.8	3.0	0.8-8.1	1.4	2.9	1.2-6.1	0.8	3.4	0.9-9.0	0.6	2.4	0.5-7.3
	2011-2015	5.0	9.1	5.9-13.7	3.6	14.7	8.5-23.7	1.4	4.9	1.9-10.7	1.4	2.5	1.0-5.5	1.2	5.2	1.8-11.8	0.2	0.6	0.0-4.5
	2016-2020	2.4	4.6	2.4-8.4	1.6	6.3	2.7-12.9	0.8	3.0	0.8-8.3	1.0	1.5	0.5-4.0	1.0	3.1	1.0-8.4	0.0	0.0	0.0-3.9
Colorectal	1976-1980	19.6	56.7	45.9-69.3	8.8	57.8	41.6-78.1	10.8	55.4	41.4-72.6	10.2	29.2	21.6-38.4	3.2	22.3	12.4-36.4	7.0	34.9	24.2-48.8
	1981-1985	18.8	51.1	41.2-62.7	8.8	54.8	39.4-74.2	10.0	46.1	34.2-61.1	12.0	33.3	25.3-43.0	5.6	36.1	23.5-52.7	6.4	31.7	21.5-45.1
	1986-1990	24.2	62.9	52.1-75.3	9.8	58.2	42.6-77.4	14.4	64.8	50.5-81.9	9.8	25.7	19.0-34.1	5.4	33.0	21.4-48.3	4.4	19.4	12.1-29.7
	1991-1995	22.2	54.3	44.6-65.4	10.4	61.6	45.6-81.0	11.8	49.7	37.7-64.3	11.6	28.9	21.9-37.3	6.0	37.5	25.1-53.6	5.6	22.8	15.1-33.1
	1996-2000	28.4	66.9	56.4-78.9	14.6	83.0	64.7-104.7	13.8	55.2	42.9-70.1	11.0	25.9	19.5-33.7	6.0	34.2	22.9-48.8	5.0	19.3	12.4-28.9
	2001-2005	30.6	67.1	56.9-78.7	16.0	79.4	62.8-99.1	14.6	55.0	43.0-69.6	8.8	19.5	14.1-26.2	4.6	23.8	15.0-35.7	4.2	14.9	9.2-23.2
	2006-2010	26.6	53.1	44.4-63.1	12.6	54.3	41.5-70.0	14.0	51.1	39.7-65.0	7.2	15.0	10.4-20.8	2.8	13.9	7.6-23.5	4.4	15.6	9.7-24.0
	2011-2015	24.2	45.9	37.9-55.1	12.2	50.0	37.8-64.9	12.0	41.5	31.5-54.0	9.8	18.6	13.6-24.9	5.8	24.0	15.9-35.1	4.0	13.9	8.4-22.1
	2016-2020	24.6	45.5	37.5-54.9	12.6	47.2	35.8-61.5	12.0	43.6	32.7-57.2	8.0	14.7	10.4-20.4	4.6	18.7	11.6-28.8	3.4	11.6	6.7-19.3
Colon excluding rectum	1976-1980	13.6	39.8	30.8-50.5	5.4	37.8	24.6-55.2	8.2	41.9	29.9-57.1	7.0	20.0	13.9-27.9	1.6	11.5	4.7-22.6	5.4	27.3	17.8-39.9
	1981-1985	13.6	37.4	28.9-47.5	4.8	31.0	19.5-46.4	8.8	41.0	29.7-55.2	9.4	26.5	19.4-35.3	3.8	26.7	15.7-42.0	5.6	27.1	17.9-39.5
	1986-1990	14.6	38.3	30.0-48.3	5.6	33.6	22.0-48.9	9.0	41.1	29.8-55.3	8.4	22.5	16.2-30.5	4.6	29.1	18.1-43.8	3.8	16.9	10.1-26.7
	1991-1995	14.2	34.6	27.0-43.7	6.6	41.2	28.1-58.0	7.6	31.4	22.2-43.3	10.4	26.0	19.4-34.1	5.6	35.0	23.0-50.5	4.8	19.3	12.4-29.0
	1996-2000	19.0	44.7	36.2-54.6	8.2	48.0	34.2-65.2	10.8	43.3	32.4-56.7	9.8	23.1	17.1-30.5	5.4	30.9	20.2-45.0	4.4	17.3	10.7-26.4
	2001-2005	20.0	44.3	36.0-53.9	10.0	50.5	37.3-66.7	10.0	37.2	27.5-49.4	7.8	17.2	12.2-23.6	3.8	19.4	11.6-30.3	4.0	14.1	8.5-22.2
	2006-2010	20.6	40.8	33.2-49.6	9.2	40.0	29.1-53.8	11.4	41.1	31.0-53.7	6.8	14.1	9.8-19.9	2.6	13.0	6.9-22.3	4.2	15.0	9.2-23.4
	2011-2015	14.8	28.1	21.9-35.5	6.6	27.7	18.7-39.6	8.2	28.2	20.1-38.8	8.6	16.5	11.8-22.5	5.2	21.9	14.1-32.6	3.4	11.9	6.8-19.6
	2016-2020	16.4	31.0	24.3-39.0	8.4	32.3	22.8-44.6	8.0	29.1	20.4-40.6	5.6	10.1	6.6-15.0	3.4	13.4	7.5-22.2	2.2	7.3	3.6-13.8
Rectum & rectosigmoid	1976-1980	6.0	16.9	11.4-24.3	3.4	20.1	11.5-32.5	2.6	13.5	7.1-23.3	3.2	9.1	5.2-14.9	1.6	10.8	4.5-21.4	1.6	7.6	3.3-15.2
	1981-1985	5.2	13.8	8.9-20.3	4.0	23.8	14.3-37.3	1.2	5.2	1.9-11.6	2.6	6.9	3.6-11.8	1.8	9.3	4.2-18.1	0.8	4.6	1.2-11.7
	1986-1990	9.6	24.6	18.1-32.6	4.2	24.6	15.0-38.0	5.4	23.7	15.6-34.7	1.4	3.2	1.3-6.7	0.8	3.9	1.1-10.3	0.6	2.5	0.5-7.7
	1991-1995	8.0	19.7	14.0-26.8	3.8	20.3	12.2-31.8	4.2	18.3	11.3-28.1	1.2	2.9	1.1-6.3	0.4	2.5	0.3-8.7	0.8	3.4	0.9-8.9
	1996-2000	9.4	22.2	16.3-29.6	6.4	35.1	23.8-49.6	3.0	11.9	6.6-19.9	1.2	2.7	1.0-6.0	0.6	3.3	0.7-9.4	0.6	2.1	0.4-6.6
	2001-2005	10.6	22.9	17.1-30.0	6.0	28.9	19.4-41.5	4.6	17.8	11.3-27.0	1.0	2.3	0.7-5.3	0.8	4.4	1.2-11.0	0.2	0.8	0.0-4.9
	2006-2010	6.0	12.3	8.2-17.7	3.4	14.4	8.2-23.4	2.6	10.0	5.2-17.5	0.4	0.8	0.1-3.1	0.2	0.9	0.0-5.4	0.2	0.6	0.0-4.2
	2011-2015	9.4	17.8	13.0-23.9	5.6	22.3	14.7-32.8	3.8	13.3	8.0-21.3	1.2	2.1	0.8-4.9	0.6	2.1	0.4-7.0	0.6	2.1	0.4-6.8
	2016-2020	8.2	14.5	10.3-20.2	4.2	14.9	9.1-23.6	4.0	14.5	8.6-23.3	2.4	4.6	2.3-8.3	1.2	5.3	1.9-12.0	1.2	4.3	1.5-10.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Orleans**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	0.8	2.1	0.6-5.5	0.6	3.2	0.6-9.8	0.2	1.0	0.0-5.7	0.2	0.4	0.0-2.7	0.2	0.9	0.0-5.9	0.0	0.0	0.0-3.9
	1981-1985	0.4	1.1	0.1-4.1	0.2	0.9	0.0-6.0	0.2	1.3	0.0-6.6	0.8	2.2	0.6-5.8	0.4	2.3	0.3-8.6	0.4	2.1	0.2-7.8
	1986-1990	0.8	1.9	0.5-4.9	0.6	3.0	0.6-9.2	0.2	1.0	0.0-5.6	1.4	3.7	1.5-7.6	0.8	5.4	1.4-13.6	0.6	2.8	0.6-8.3
	1991-1995	1.4	3.5	1.4-7.2	0.6	3.6	0.7-10.2	0.8	3.6	1.0-9.2	0.8	2.0	0.5-5.1	0.2	1.5	0.0-7.2	0.6	2.5	0.5-7.4
	1996-2000	1.8	4.2	1.9-7.9	0.8	4.7	1.2-11.9	1.0	4.0	1.3-9.6	1.8	4.2	1.9-8.0	1.0	5.1	1.7-11.9	0.8	3.0	0.8-8.1
	2001-2005	1.4	3.2	1.3-6.6	1.4	7.6	3.0-15.4	0.0	0.0	0.0-3.4	1.4	3.1	1.2-6.4	1.2	6.8	2.5-14.6	0.2	0.8	0.0-4.9
	2006-2010	1.2	2.3	0.8-5.3	1.0	3.7	1.2-9.3	0.2	0.9	0.0-5.1	1.6	3.0	1.3-6.1	0.8	3.0	0.8-8.2	0.8	2.8	0.8-7.8
	2011-2015	4.0	6.8	4.1-10.9	2.2	8.0	3.9-15.0	1.8	5.7	2.6-11.5	3.6	6.6	3.9-10.7	1.6	6.6	2.8-13.5	2.0	6.9	3.3-13.3
	2016-2020	4.8	8.4	5.3-12.9	3.2	11.4	6.4-19.2	1.6	5.7	2.2-12.1	4.0	7.0	4.3-11.2	2.6	9.5	4.9-16.9	1.4	4.7	1.8-10.4
Pancreas	1976-1980	4.0	11.0	6.7-17.1	2.0	12.1	5.5-22.8	2.0	10.0	4.8-18.5	4.0	10.9	6.6-16.9	2.0	11.7	5.3-22.2	2.0	10.0	4.8-18.5
	1981-1985	4.0	10.9	6.6-16.9	1.6	10.0	4.2-19.8	2.4	11.0	5.6-19.5	4.2	12.0	7.4-18.3	2.0	14.2	6.6-25.9	2.2	10.0	4.9-18.2
	1986-1990	4.0	10.6	6.4-16.3	1.4	8.6	3.4-17.8	2.6	11.4	6.0-19.7	4.4	11.4	7.1-17.4	1.4	8.5	3.3-17.6	3.0	13.0	7.2-21.7
	1991-1995	3.2	7.8	4.4-12.6	1.0	5.2	1.6-12.3	2.2	9.4	4.7-17.1	3.4	8.3	4.8-13.3	1.0	5.2	1.6-12.3	2.4	10.3	5.3-18.2
	1996-2000	5.0	11.7	7.6-17.3	1.8	9.6	4.4-18.1	3.2	13.6	7.8-22.3	5.0	11.8	7.7-17.5	2.0	10.8	5.1-19.7	3.0	12.6	7.0-21.0
	2001-2005	5.4	11.7	7.7-17.1	2.6	12.6	6.6-21.6	2.8	11.2	6.1-19.0	5.8	12.5	8.3-17.9	2.8	13.0	7.1-21.9	3.0	11.2	6.2-18.9
	2006-2010	4.6	9.2	5.8-14.0	2.8	12.6	6.8-21.4	1.8	7.0	3.1-13.5	4.4	8.9	5.6-13.7	2.8	12.1	6.5-20.7	1.6	6.6	2.8-13.2
	2011-2015	9.0	17.2	12.5-23.3	5.2	20.4	13.2-30.4	3.8	13.7	8.1-21.9	6.4	12.0	8.2-17.3	3.6	14.3	8.3-23.1	2.8	10.0	5.4-17.4
	2016-2020	10.2	18.2	13.5-24.3	5.0	19.5	12.5-29.3	5.2	16.8	10.8-25.3	9.6	17.3	12.7-23.2	4.4	17.3	10.7-26.7	5.2	17.6	11.3-26.5
Larynx	1976-1980	1.2	3.2	1.2-7.0	1.0	5.6	1.8-13.3	0.2	1.1	0.0-6.0	0.2	0.5	0.0-2.8	0.2	1.0	0.0-6.2	0.0	0.0	0.0-3.9
	1981-1985	2.4	6.7	3.4-11.7	2.2	14.7	7.0-26.6	0.2	1.0	0.0-5.6	0.8	2.4	0.6-6.1	0.8	6.3	1.6-15.6	0.0	0.0	0.0-3.9
	1986-1990	2.2	5.9	2.9-10.7	1.8	9.9	4.5-18.8	0.4	2.5	0.3-8.4	0.6	1.6	0.3-4.7	0.4	2.9	0.3-10.0	0.2	0.8	0.0-4.9
	1991-1995	2.2	5.5	2.7-9.8	2.2	11.9	5.8-21.5	0.0	0.0	0.0-3.4	1.4	3.4	1.4-7.1	1.2	7.7	2.7-16.5	0.2	0.9	0.0-5.0
	1996-2000	2.8	6.6	3.6-11.0	2.4	12.1	6.2-21.1	0.4	1.8	0.2-6.5	0.0	0.0	0.0-1.7	0.0	0.0	0.0-3.7	0.0	0.0	0.0-3.3
	2001-2005	2.2	4.8	2.4-8.6	1.8	8.3	3.8-15.9	0.4	1.7	0.2-6.4	0.2	0.4	0.0-2.5	0.2	0.8	0.0-5.1	0.0	0.0	0.0-3.4
	2006-2010	2.8	5.3	2.9-9.1	2.2	9.5	4.7-17.3	0.6	2.0	0.4-6.4	0.4	0.7	0.1-2.9	0.2	0.8	0.0-4.9	0.2	0.6	0.0-4.4
	2011-2015	2.2	3.8	1.9-7.0	1.6	6.1	2.5-12.6	0.6	1.9	0.4-6.4	1.0	1.7	0.5-4.2	0.6	2.2	0.4-7.2	0.4	1.2	0.2-5.5
	2016-2020	2.0	4.0	1.8-7.7	1.2	5.3	1.8-12.3	0.8	2.7	0.7-7.8	0.4	0.8	0.1-3.3	0.4	1.6	0.2-6.7	0.0	0.0	0.0-3.9
Lung and bronchus	1976-1980	24.2	66.0	54.5-79.2	19.0	116.5	93.5-143.3	5.2	25.0	16.1-37.0	16.0	44.1	34.8-55.1	13.2	80.2	61.5-102.7	2.8	14.7	7.9-24.9
	1981-1985	27.6	74.0	62.0-87.6	18.6	115.0	92.0-141.9	9.0	45.7	33.1-61.5	20.6	54.6	44.4-66.3	13.8	85.6	65.8-109.3	6.8	33.6	23.1-47.3
	1986-1990	23.8	58.5	48.4-70.2	16.4	91.0	71.8-113.5	7.4	33.1	23.2-46.0	23.6	59.7	49.3-71.6	16.8	98.9	78.3-123.1	6.8	32.9	22.5-46.2
	1991-1995	29.6	71.8	60.7-84.4	18.6	106.0	85.1-130.2	11.0	49.0	36.8-63.9	24.4	59.5	49.3-71.0	13.6	79.0	61.0-100.5	10.8	48.1	36.0-62.9
	1996-2000	31.8	75.3	64.0-87.9	18.6	99.6	80.3-122.2	13.2	56.7	43.8-72.3	26.4	62.5	52.3-74.1	15.4	83.4	65.7-104.3	11.0	46.7	35.1-60.9
	2001-2005	34.8	76.5	65.5-88.9	18.4	90.5	72.7-111.2	16.4	65.6	52.1-81.7	25.8	56.5	47.1-67.2	14.4	72.8	56.7-91.9	11.4	45.5	34.4-59.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Orleans**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	42.6	84.1	73.1-96.4	24.2	104.6	86.5-125.5	18.4	68.9	55.4-84.9	30.0	59.8	50.5-70.4	17.0	78.1	62.1-96.9	13.0	47.0	36.2-60.3
	2011-2015	45.2	85.4	74.4-97.7	22.2	91.6	74.9-111.2	23.0	81.2	66.8-98.0	31.6	60.0	50.9-70.4	13.8	57.2	44.1-73.0	17.8	63.1	50.5-78.2
	2016-2020	52.8	92.7	81.6-105.1	26.8	101.4	84.4-121.2	26.0	87.0	72.4-104.2	30.2	53.6	45.2-63.3	16.4	63.2	49.7-79.4	13.8	46.1	35.7-58.9
Melanoma of the skin	1976-1980	2.6	7.4	3.9-12.7	1.6	9.7	4.0-19.4	1.0	6.0	1.9-13.8	1.2	3.3	1.2-7.2	1.0	5.4	1.7-12.9	0.2	1.5	0.0-7.2
	1981-1985	2.0	5.6	2.7-10.4	1.0	5.7	1.8-13.5	1.0	5.5	1.8-12.8	0.4	0.9	0.1-3.4	0.4	1.8	0.2-7.3	0.0	0.0	0.0-3.9
	1986-1990	3.0	7.4	4.1-12.3	1.0	4.7	1.5-11.5	2.0	9.6	4.5-17.9	0.8	2.0	0.5-5.3	0.8	4.2	1.1-10.9	0.0	0.0	0.0-3.6
	1991-1995	4.2	10.0	6.1-15.2	2.8	14.8	8.0-25.1	1.4	6.7	2.7-13.8	1.4	3.5	1.4-7.1	0.8	4.9	1.3-12.2	0.6	2.8	0.5-8.2
	1996-2000	6.4	14.9	10.2-21.1	3.4	16.7	9.7-26.9	3.0	13.3	7.4-22.0	1.4	3.4	1.3-6.9	1.2	6.2	2.3-13.5	0.2	0.8	0.0-4.8
	2001-2005	6.2	13.5	9.2-19.3	4.8	22.2	14.1-33.2	1.4	6.2	2.5-12.9	0.8	1.7	0.5-4.3	0.6	3.0	0.6-8.8	0.2	0.8	0.0-4.9
	2006-2010	7.6	15.1	10.7-20.9	4.2	18.9	11.5-29.1	3.4	13.0	7.5-21.2	1.2	2.7	1.0-6.0	1.0	5.0	1.6-11.8	0.2	1.1	0.0-5.7
	2011-2015	14.2	30.2	23.4-38.4	7.2	31.8	22.0-44.5	7.0	29.2	19.9-41.3	1.4	2.9	1.1-6.2	0.6	2.7	0.6-8.3	0.8	3.2	0.7-8.9
2016-2020	16.2	29.6	23.2-37.4	9.2	35.7	25.6-48.7	7.0	25.2	17.1-36.1	1.2	2.0	0.7-4.8	0.8	2.9	0.8-8.3	0.4	1.2	0.1-5.6	
Female breast	1976-1980							18.2	97.7	78.1-120.5							5.2	26.4	17.1-39.0
	1981-1985							15.8	79.8	62.7-100.0							6.8	32.3	22.2-45.5
	1986-1990							26.6	130.1	108.5-154.6							8.0	38.8	27.5-53.1
	1991-1995							27.8	124.2	104.1-146.9							7.4	33.0	23.1-45.7
	1996-2000							32.0	140.0	119.0-163.6							6.8	27.8	19.2-39.0
	2001-2005							30.2	124.2	105.1-145.9							5.6	22.7	15.0-33.0
	2006-2010							32.6	123.1	104.7-144.0							5.6	20.4	13.5-29.9
	2011-2015							31.0	114.2	96.5-134.3							3.6	12.7	7.5-20.6
2016-2020							35.0	131.6	111.9-154.1							5.0	16.9	10.8-25.6	
Cervix uteri	1976-1980							3.0	18.3	10.0-30.2							0.8	4.4	1.2-11.4
	1981-1985							2.0	9.9	4.7-18.4							0.8	3.8	1.0-9.9
	1986-1990							2.4	10.8	5.5-19.2							0.4	1.7	0.2-6.4
	1991-1995							3.2	15.0	8.5-24.5							0.8	3.1	0.8-8.3
	1996-2000							2.6	11.6	6.2-20.0							0.6	2.9	0.6-8.3
	2001-2005							3.6	16.0	9.4-25.3							0.2	0.9	0.0-5.0
	2006-2010							1.6	6.7	2.8-13.4							0.6	2.2	0.4-6.8
	2011-2015							1.6	6.1	2.5-12.7							0.4	1.3	0.1-5.6
2016-2020							2.0	9.5	4.3-18.1							1.4	6.6	2.4-14.4	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Orleans**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							5.4	28.7	18.8-41.9							1.0	5.0	1.6-11.7
	1981-1985							5.4	25.9	17.0-38.0							0.8	3.9	1.1-10.1
	1986-1990							3.8	16.6	9.9-26.2							1.0	4.8	1.5-11.4
	1991-1995							6.0	27.2	18.3-39.0							1.2	4.8	1.8-10.7
	1996-2000							7.6	32.8	23.2-45.2							1.4	5.9	2.4-12.4
	2001-2005							7.0	29.2	20.3-40.7							1.6	6.3	2.7-12.7
	2006-2010							6.6	25.6	17.5-36.3							1.2	3.9	1.4-9.1
	2011-2015							9.4	33.6	24.5-45.2							1.2	4.4	1.6-10.2
	2016-2020							10.8	40.1	29.6-53.4							1.0	3.0	1.0-8.1
Ovary	1976-1980							2.8	15.9	8.6-26.8							0.8	4.3	1.1-11.1
	1981-1985							2.8	13.5	7.3-23.0							1.8	9.2	4.1-17.5
	1986-1990							4.6	22.0	13.7-33.3							3.6	16.4	9.6-26.2
	1991-1995							4.4	19.6	12.2-29.8							4.0	17.8	10.8-27.6
	1996-2000							4.4	19.5	12.2-29.5							2.6	11.5	6.1-19.6
	2001-2005							2.8	11.1	6.1-18.9							1.2	5.1	1.8-11.2
	2006-2010							4.8	16.5	10.5-25.0							2.6	8.7	4.6-15.3
	2011-2015							3.4	11.8	6.8-19.4							2.0	6.7	3.2-12.9
	2016-2020							4.0	15.0	8.9-24.0							1.6	5.3	2.3-11.3
Prostate	1976-1980				13.2	98.8	75.9-126.0							4.8	35.8	22.6-53.4			
	1981-1985				14.8	102.3	79.4-129.2							5.0	40.6	25.9-59.8			
	1986-1990				20.8	135.1	109.4-164.5							3.8	27.1	16.0-42.3			
	1991-1995				29.4	170.7	143.7-201.1							6.8	42.9	29.4-60.1			
	1996-2000				36.0	194.9	167.1-225.7							5.0	30.8	19.7-45.4			
	2001-2005				26.4	128.5	107.3-152.8							4.2	24.0	14.8-36.6			
	2006-2010				30.4	130.4	110.2-153.5							2.0	9.6	4.5-17.7			
	2011-2015				30.6	115.3	97.4-135.7							3.2	15.9	9.0-26.0			
	2016-2020				36.0	125.5	107.3-146.2							3.6	16.2	9.6-26.0			
Testis	1976-1980				0.0	0.0	0.0-4.5							0.2	0.9	0.0-6.0			
	1981-1985				0.8	4.2	1.1-11.1							0.0	0.0	0.0-4.5			
	1986-1990				1.0	4.6	1.4-11.3							0.2	1.2	0.0-6.5			
	1991-1995				2.0	8.2	3.9-15.5							0.0	0.0	0.0-3.8			
	1996-2000				1.8	8.6	3.9-16.4							0.0	0.0	0.0-3.7			
	2001-2005				1.4	6.4	2.6-13.3							0.2	1.1	0.0-5.8			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Orleans**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				1.0	5.2	1.7-12.1							0.0	0.0	0.0-3.6			
	2011-2015				2.0	9.6	4.5-18.0							0.0	0.0	0.0-3.8			
	2016-2020				1.2	5.8	2.0-13.0							0.0	0.0	0.0-3.9			
Urinary bladder (incl. in situ)	1976-1980	6.6	18.4	12.5-25.9	4.8	29.9	18.9-44.9	1.8	8.3	3.8-16.1	0.8	1.9	0.5-5.2	0.6	3.8	0.7-11.3	0.2	0.8	0.0-5.3
	1981-1985	6.4	17.3	11.8-24.5	4.4	29.9	18.4-45.6	2.0	9.1	4.4-17.0	2.6	7.4	3.9-12.7	1.6	12.5	5.3-24.2	1.0	4.3	1.4-10.5
	1986-1990	9.2	23.3	17.0-31.1	7.4	44.5	30.9-61.8	1.8	8.1	3.6-15.5	2.6	6.6	3.5-11.3	2.0	12.5	5.8-23.1	0.6	2.3	0.5-7.3
	1991-1995	10.0	24.0	17.8-31.7	7.4	39.2	27.5-54.2	2.6	10.4	5.5-18.0	2.0	5.0	2.4-9.1	1.0	6.1	1.9-14.0	1.0	4.1	1.3-9.7
	1996-2000	12.8	30.2	23.3-38.5	9.4	52.5	38.4-69.9	3.4	14.0	8.1-22.6	3.8	8.9	5.4-13.9	2.6	16.2	8.5-27.5	1.2	4.6	1.7-10.3
	2001-2005	15.4	34.0	26.8-42.6	10.4	50.9	37.9-67.0	5.0	19.6	12.6-29.2	3.2	7.1	4.0-11.5	2.0	10.2	4.8-18.8	1.2	4.8	1.7-10.8
	2006-2010	11.8	23.7	18.0-30.8	9.8	44.3	32.5-58.9	2.0	7.7	3.6-14.5	2.4	4.9	2.5-8.7	0.8	3.7	1.0-9.7	1.6	5.8	2.5-11.9
	2011-2015	16.0	30.0	23.7-37.5	13.0	53.8	41.2-69.2	3.0	10.1	5.6-17.2	3.4	6.9	4.0-11.2	2.8	13.8	7.4-23.3	0.6	1.8	0.4-6.2
	2016-2020	16.6	29.2	23.1-36.5	13.0	48.7	37.2-62.9	3.6	12.7	7.4-20.8	4.0	7.2	4.4-11.5	2.8	11.6	6.2-20.0	1.2	4.2	1.5-10.0
Kidney and renal pelvis	1976-1980	3.2	8.7	4.9-14.2	1.8	11.3	5.0-21.8	1.4	6.5	2.6-13.8	1.2	2.9	1.1-6.5	0.8	4.7	1.2-12.6	0.4	1.8	0.2-6.8
	1981-1985	2.6	7.9	4.2-13.4	1.4	10.7	4.1-21.8	1.2	7.2	2.6-15.6	1.2	3.7	1.3-7.9	1.0	8.7	2.8-19.4	0.2	1.2	0.0-6.3
	1986-1990	3.8	9.5	5.7-15.0	2.4	13.3	6.8-23.4	1.4	5.9	2.4-12.4	2.0	5.3	2.5-9.8	1.2	7.6	2.7-16.6	0.8	3.6	1.0-9.4
	1991-1995	4.4	11.0	6.9-16.7	2.8	16.0	8.6-26.8	1.6	7.3	3.1-14.5	2.6	6.5	3.5-11.1	1.8	10.7	4.8-20.3	0.8	3.4	0.9-9.0
	1996-2000	5.4	12.5	8.3-18.2	3.6	19.3	11.4-30.6	1.8	7.2	3.3-13.8	2.0	4.7	2.3-8.6	1.2	6.4	2.3-13.8	0.8	3.4	0.9-8.8
	2001-2005	7.8	17.0	12.1-23.3	6.0	28.8	19.3-41.3	1.8	7.2	3.3-13.9	1.6	3.4	1.5-6.9	1.0	4.6	1.5-11.0	0.6	2.4	0.5-7.3
	2006-2010	6.8	13.2	9.1-18.6	4.0	17.6	10.6-27.5	2.8	10.0	5.4-17.3	2.4	4.8	2.4-8.5	2.0	9.1	4.3-17.0	0.4	1.3	0.2-5.5
	2011-2015	9.8	20.0	14.6-26.7	6.2	27.2	18.2-39.2	3.6	13.6	7.9-22.1	1.0	1.7	0.5-4.4	0.8	2.8	0.7-8.0	0.2	0.7	0.0-4.8
	2016-2020	8.6	16.2	11.6-22.3	5.4	20.7	13.3-30.9	3.2	13.0	7.2-21.8	2.6	4.6	2.4-8.2	1.8	6.5	2.9-13.2	0.8	2.7	0.7-7.9
Brain and other central nervous system	1976-1980	3.0	8.7	4.8-14.5	2.0	11.9	5.6-22.3	1.0	5.4	1.7-12.8	2.0	5.7	2.7-10.6	1.2	6.6	2.4-14.8	0.8	4.7	1.2-12.0
	1981-1985	2.8	8.3	4.5-13.9	1.4	9.2	3.7-18.8	1.4	7.5	2.9-15.6	2.2	6.3	3.1-11.3	1.2	7.7	2.8-16.6	1.0	4.5	1.5-10.8
	1986-1990	3.0	7.9	4.4-13.1	1.6	8.3	3.5-16.6	1.4	7.6	3.0-15.6	2.2	5.7	2.8-10.2	1.2	6.6	2.3-14.8	1.0	5.3	1.7-12.5
	1991-1995	3.0	7.2	4.0-11.8	1.0	5.7	1.8-13.3	2.0	8.6	4.1-16.1	2.0	4.8	2.3-8.7	0.4	2.1	0.3-7.7	1.6	7.0	3.0-13.9
	1996-2000	4.4	10.4	6.5-15.7	2.6	13.2	7.0-22.6	1.8	7.5	3.4-14.4	3.2	7.6	4.3-12.3	2.0	10.5	5.0-19.2	1.2	4.6	1.7-10.3
	2001-2005	3.0	6.6	3.7-10.9	1.2	5.3	1.9-11.8	1.8	7.2	3.2-14.0	2.4	5.0	2.6-8.9	1.0	4.4	1.4-10.4	1.4	4.9	1.9-10.6
	2006-2010	3.2	6.5	3.7-10.7	1.8	7.3	3.3-14.2	1.4	5.7	2.3-12.1	1.6	3.0	1.3-6.1	0.8	3.0	0.8-8.1	0.8	2.8	0.7-7.7
	2011-2015	2.8	5.8	3.0-10.0	1.2	5.1	1.8-11.6	1.6	6.5	2.6-13.5	3.6	7.0	4.1-11.4	2.0	7.7	3.6-14.9	1.6	6.2	2.5-12.8
	2016-2020	2.8	6.0	3.1-10.3	1.0	4.2	1.3-10.4	1.8	7.8	3.4-15.5	2.2	4.0	2.0-7.5	0.8	3.6	1.0-9.6	1.4	4.6	1.8-10.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Orleans**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	1.0	3.7	1.2-8.4	0.2	1.4	0.0-7.2	0.8	6.2	1.7-15.0	0.2	0.6	0.0-3.2	0.2	1.4	0.0-7.2	0.0	0.0	0.0-3.9
	1981-1985	2.0	5.6	2.6-10.4	0.6	3.3	0.6-10.0	1.4	7.5	2.9-15.5	0.2	0.5	0.0-2.9	0.2	1.0	0.0-6.2	0.0	0.0	0.0-3.9
	1986-1990	1.2	3.2	1.2-7.0	0.4	2.4	0.3-8.5	0.8	3.4	0.9-9.0	0.4	1.0	0.1-3.6	0.0	0.0	0.0-4.2	0.4	1.6	0.2-6.2
	1991-1995	2.0	4.8	2.3-8.9	0.6	3.2	0.7-9.3	1.4	6.6	2.6-13.6	0.0	0.0	0.0-1.7	0.0	0.0	0.0-3.8	0.0	0.0	0.0-3.4
	1996-2000	2.8	6.3	3.4-10.6	1.6	7.6	3.2-15.1	1.2	5.6	2.0-12.2	0.6	1.4	0.3-4.0	0.0	0.0	0.0-3.7	0.6	2.5	0.5-7.5
	2001-2005	3.6	7.8	4.6-12.4	1.4	6.6	2.6-13.8	2.2	9.2	4.5-16.6	0.2	0.5	0.0-2.6	0.2	1.1	0.0-5.8	0.0	0.0	0.0-3.4
	2006-2010	5.2	11.6	7.5-17.2	0.8	2.8	0.8-7.9	4.4	20.0	12.4-30.5	0.4	0.9	0.1-3.3	0.4	2.2	0.3-7.6	0.0	0.0	0.0-3.4
	2011-2015	7.6	14.6	10.2-20.5	1.6	5.6	2.4-11.8	6.0	23.8	15.8-34.7	0.4	0.7	0.1-2.9	0.4	2.2	0.3-7.8	0.0	0.0	0.0-3.6
2016-2020	5.8	12.4	8.2-18.3	1.0	4.2	1.3-10.4	4.8	20.9	13.0-31.8	0.0	0.0	0.0-1.9	0.0	0.0	0.0-3.9	0.0	0.0	0.0-3.9	
Hodgkin lymphoma	1976-1980	1.4	3.7	1.5-7.8	0.6	3.5	0.7-10.3	0.8	3.8	1.0-10.0	0.6	1.6	0.3-4.7	0.2	1.4	0.0-7.2	0.4	2.0	0.2-7.3
	1981-1985	1.4	3.3	1.3-7.0	0.6	3.4	0.7-10.1	0.8	3.6	1.0-9.5	0.0	0.0	0.0-2.0	0.0	0.0	0.0-4.5	0.0	0.0	0.0-3.9
	1986-1990	1.6	4.0	1.7-8.0	1.4	7.1	2.8-14.9	0.2	0.9	0.0-5.3	0.6	1.5	0.3-4.4	0.2	0.9	0.0-5.8	0.4	1.8	0.2-6.6
	1991-1995	1.4	3.1	1.3-6.5	0.8	3.2	0.9-8.8	0.6	2.7	0.5-8.0	0.4	0.9	0.1-3.3	0.2	0.7	0.0-5.0	0.2	0.7	0.0-4.5
	1996-2000	1.8	4.2	1.9-8.0	0.4	1.9	0.2-7.0	1.4	6.2	2.5-12.8	0.0	0.0	0.0-1.7	0.0	0.0	0.0-3.7	0.0	0.0	0.0-3.3
	2001-2005	1.2	2.7	1.0-6.0	0.8	3.9	1.1-10.1	0.4	1.4	0.2-5.7	0.2	0.6	0.0-2.9	0.0	0.0	0.0-3.6	0.2	1.1	0.0-5.8
	2006-2010	1.4	3.3	1.3-6.8	0.4	2.2	0.3-7.6	1.0	4.5	1.4-10.7	0.0	0.0	0.0-1.7	0.0	0.0	0.0-3.6	0.0	0.0	0.0-3.4
	2011-2015	1.2	2.9	1.0-6.4	0.4	2.2	0.3-7.8	0.8	3.5	0.9-9.3	0.2	0.5	0.0-2.8	0.2	1.0	0.0-5.6	0.0	0.0	0.0-3.6
2016-2020	2.2	4.7	2.3-8.6	1.0	4.2	1.3-10.3	1.2	5.3	1.9-12.1	0.2	0.4	0.0-2.7	0.2	0.9	0.0-5.5	0.0	0.0	0.0-3.9	
Non-Hodgkin lymphomas	1976-1980	2.2	6.0	3.0-10.8	1.6	9.2	3.9-18.4	0.6	3.3	0.6-9.6	1.6	4.3	1.8-8.6	0.8	4.7	1.2-12.6	0.8	3.9	1.0-10.1
	1981-1985	4.4	12.1	7.5-18.5	2.6	16.3	8.5-28.0	1.8	8.9	4.0-17.1	2.2	5.9	2.9-10.7	1.6	10.0	4.1-20.1	0.6	3.2	0.6-9.3
	1986-1990	4.0	10.1	6.1-15.6	1.6	8.8	3.7-17.4	2.4	11.0	5.6-19.5	2.6	7.0	3.7-11.9	1.6	9.1	3.9-18.1	1.0	4.7	1.5-11.1
	1991-1995	9.6	23.1	17.0-30.6	4.4	22.9	14.2-34.9	5.2	23.5	15.3-34.6	4.0	9.8	6.0-15.2	2.0	10.8	5.0-20.0	2.0	8.8	4.2-16.3
	1996-2000	8.6	20.1	14.5-27.1	3.4	17.7	10.2-28.5	5.2	23.0	15.0-33.9	3.6	8.4	5.0-13.3	1.8	9.4	4.3-17.7	1.8	7.3	3.3-14.0
	2001-2005	11.0	24.4	18.4-31.8	6.2	30.1	20.4-42.9	4.8	19.4	12.4-29.1	4.2	9.1	5.6-14.0	2.6	14.2	7.5-24.2	1.6	5.8	2.5-11.8
	2006-2010	9.8	20.1	14.8-26.7	5.2	23.4	15.1-34.6	4.6	16.5	10.4-25.2	3.0	6.0	3.3-10.0	2.0	9.8	4.7-18.1	1.0	3.0	1.0-7.7
	2011-2015	10.6	20.6	15.3-27.2	5.0	21.7	13.9-32.5	5.6	20.7	13.6-30.5	3.8	7.9	4.8-12.5	1.8	9.1	4.1-17.3	2.0	7.2	3.4-13.8
2016-2020	13.0	23.0	17.6-29.8	7.4	27.6	19.2-38.9	5.6	18.8	12.3-28.1	4.0	6.8	4.1-10.9	2.2	8.1	3.9-15.2	1.8	5.7	2.6-11.7	
Myeloma	1976-1980	1.6	4.1	1.7-8.1	0.8	5.3	1.3-13.5	0.8	3.6	1.0-9.6	1.4	3.8	1.5-7.9	0.8	4.9	1.3-12.5	0.6	2.8	0.6-8.4
	1981-1985	1.0	2.6	0.8-6.2	0.6	4.0	0.8-11.3	0.4	1.6	0.2-6.5	1.8	4.6	2.1-8.8	1.0	6.1	1.9-14.3	0.8	3.4	0.9-9.1
	1986-1990	1.0	2.6	0.8-6.2	0.6	3.3	0.7-9.8	0.4	2.2	0.3-7.8	1.0	2.5	0.8-5.9	0.6	3.8	0.8-10.7	0.4	1.7	0.2-6.4
	1991-1995	2.6	6.3	3.3-10.7	1.8	10.5	4.7-19.8	0.8	3.2	0.9-8.4	1.6	4.1	1.8-8.0	0.8	4.5	1.2-11.3	0.8	3.8	1.0-9.8
	1996-2000	2.2	5.1	2.6-9.2	1.0	5.5	1.8-12.7	1.2	5.3	1.9-11.6	1.2	2.8	1.0-6.1	0.8	5.1	1.3-12.6	0.4	1.7	0.2-6.3
	2001-2005	2.2	5.0	2.5-8.9	1.4	6.8	2.7-14.0	0.8	3.3	0.9-8.6	1.6	3.5	1.5-7.0	0.8	3.7	1.0-9.6	0.8	3.2	0.9-8.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Orleans**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	2.8	5.5	3.0-9.4	1.6	6.5	2.8-13.2	1.2	4.9	1.8-10.9	1.2	2.5	0.9-5.5	0.4	1.4	0.2-5.7	0.8	3.2	0.9-8.5
	2011-2015	2.2	4.2	2.1-7.9	1.0	4.5	1.4-10.9	1.2	4.3	1.5-10.1	0.8	1.7	0.4-4.4	0.6	3.1	0.6-9.1	0.2	0.8	0.0-5.0
	2016-2020	1.8	2.9	1.3-5.8	1.4	4.4	1.7-9.9	0.4	1.2	0.1-5.5	1.0	1.7	0.5-4.3	0.8	2.5	0.7-7.5	0.2	0.8	0.0-5.2
Leukemias	1976-1980	5.4	15.3	9.9-22.3	2.6	17.2	8.7-29.9	2.8	14.5	7.8-24.5	3.0	8.0	4.4-13.3	1.8	10.1	4.3-19.9	1.2	5.9	2.2-13.1
	1981-1985	5.4	15.5	10.1-22.6	3.6	24.2	14.0-38.5	1.8	9.2	4.1-17.7	3.4	9.7	5.6-15.5	1.4	9.6	3.7-19.9	2.0	9.9	4.6-18.4
	1986-1990	4.4	12.3	7.7-18.6	2.8	18.1	9.7-30.3	1.6	7.7	3.3-15.4	3.0	8.3	4.6-13.8	1.8	11.8	5.3-22.4	1.2	5.7	2.0-12.7
	1991-1995	6.6	16.1	11.1-22.6	3.8	22.4	13.3-35.1	2.8	12.2	6.6-20.5	3.4	8.4	4.9-13.5	1.4	8.4	3.3-17.1	2.0	8.4	4.0-15.6
	1996-2000	4.8	11.1	7.1-16.5	3.2	17.3	9.7-28.3	1.6	6.8	2.9-13.5	3.2	7.4	4.2-12.0	2.0	12.4	5.8-22.7	1.2	4.7	1.7-10.5
	2001-2005	8.4	18.5	13.3-25.0	4.6	23.0	14.5-34.6	3.8	15.4	9.2-24.2	3.2	6.9	3.9-11.3	2.0	10.5	4.9-19.3	1.2	4.4	1.6-10.0
	2006-2010	6.4	14.1	9.5-20.0	4.0	18.4	11.2-28.7	2.4	10.0	5.0-17.8	3.6	7.2	4.2-11.5	2.4	11.4	5.8-20.1	1.2	3.6	1.3-8.6
	2011-2015	9.6	19.5	14.2-26.1	6.4	27.0	18.3-38.6	3.2	13.2	7.3-22.0	3.4	6.4	3.7-10.5	2.4	10.7	5.4-19.0	1.0	2.9	0.9-7.7
	2016-2020	8.4	16.4	11.7-22.6	5.4	23.4	15.2-34.6	3.0	10.5	5.6-18.4	3.4	6.8	3.9-11.3	1.6	7.0	3.0-14.3	1.8	7.3	3.1-14.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Oswego**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	351.8	392.2	373.7-411.4	181.6	468.2	436.4-501.5	170.2	345.1	321.9-369.5	195.2	221.7	207.7-236.4	104.6	277.7	253.0-304.1	90.6	183.3	166.5-201.2
	1981-1985	389.4	414.9	396.4-434.2	192.2	477.0	445.6-509.8	197.2	381.4	357.5-406.5	198.8	214.4	201.0-228.4	106.4	274.7	250.6-300.4	92.4	177.4	161.3-194.6
	1986-1990	462.6	464.8	445.7-484.4	230.6	551.9	518.9-586.2	232.0	415.3	391.4-440.2	233.0	239.5	225.8-253.8	126.0	310.9	285.9-337.5	107.0	192.1	176.0-209.3
	1991-1995	517.6	486.0	467.4-505.3	258.6	572.0	539.9-605.4	259.0	438.4	414.7-463.2	234.2	224.4	211.7-237.7	121.4	285.3	261.9-310.0	112.8	187.1	171.9-203.3
	1996-2000	569.2	510.5	491.9-529.6	289.8	606.2	574.4-639.3	279.4	456.1	432.4-480.7	241.2	220.0	207.7-232.8	129.8	286.0	263.6-309.8	111.4	178.3	163.8-193.8
	2001-2005	659.4	551.9	533.2-571.2	350.2	662.6	631.1-695.2	309.2	476.8	453.2-501.3	245.4	208.3	196.8-220.4	132.6	266.9	246.4-288.7	112.8	172.4	158.4-187.3
	2006-2010	696.0	543.3	525.1-561.9	372.4	633.6	604.2-664.0	323.6	476.9	453.6-501.1	261.4	206.9	195.7-218.6	137.6	252.1	233.0-272.2	123.8	177.9	163.9-192.7
	2011-2015	733.0	525.5	508.1-543.3	382.2	571.4	545.0-598.7	350.8	488.6	465.3-512.7	251.4	184.3	174.0-195.0	141.4	230.1	212.8-248.4	110.0	148.6	136.2-161.9
	2016-2020	792.0	531.2	514.1-548.7	413.6	563.4	538.3-589.4	378.4	505.9	482.5-530.4	244.0	164.2	154.8-174.0	130.0	188.5	173.7-204.2	114.0	144.7	132.8-157.5
Oral cavity and pharynx	1976-1980	12.2	13.9	10.6-17.9	7.4	19.3	13.1-27.1	4.8	10.3	6.5-15.3	3.4	3.7	2.1-6.0	2.2	5.8	2.6-10.8	1.2	2.6	0.9-5.6
	1981-1985	8.0	8.3	5.9-11.3	6.2	13.2	8.9-18.8	1.8	3.6	1.6-6.7	3.0	3.0	1.7-5.0	2.2	4.5	2.2-8.2	0.8	1.9	0.5-4.7
	1986-1990	11.8	11.8	8.9-15.2	8.2	20.2	14.2-27.7	3.6	6.2	3.7-9.9	3.8	4.1	2.5-6.4	2.6	7.1	3.7-12.2	1.2	2.2	0.8-4.9
	1991-1995	9.6	9.0	6.6-11.9	6.8	15.1	10.3-21.2	2.8	4.8	2.6-8.0	2.4	2.3	1.2-4.0	1.4	3.5	1.3-7.2	1.0	1.7	0.6-4.0
	1996-2000	11.8	10.4	7.9-13.4	8.6	16.6	11.9-22.4	3.2	5.1	2.9-8.3	2.8	2.5	1.4-4.2	1.8	3.3	1.5-6.3	1.0	1.6	0.5-3.7
	2001-2005	11.4	9.2	6.9-11.9	9.2	15.6	11.3-20.9	2.2	3.5	1.7-6.3	2.2	1.7	0.9-3.1	1.8	3.2	1.4-6.2	0.4	0.6	0.1-2.3
	2006-2010	18.2	13.8	11.1-17.0	13.0	20.9	16.0-26.9	5.2	7.9	5.1-11.7	3.0	2.3	1.3-3.8	2.2	3.7	1.8-6.7	0.8	1.2	0.3-3.2
	2011-2015	19.6	13.3	10.7-16.3	15.6	22.2	17.4-28.0	4.0	5.3	3.2-8.4	3.4	2.3	1.3-3.8	3.2	4.7	2.6-7.8	0.2	0.4	0.0-1.9
	2016-2020	23.2	14.9	12.2-18.1	17.6	22.7	18.0-28.3	5.6	7.5	4.8-11.1	4.2	3.0	1.8-4.7	2.4	3.4	1.7-6.2	1.8	2.8	1.2-5.5
Esophagus	1976-1980	4.2	4.9	3.0-7.4	2.4	5.9	3.0-10.5	1.8	3.8	1.7-7.2	5.0	6.1	3.9-9.0	2.8	7.7	4.0-13.2	2.2	4.9	2.4-8.7
	1981-1985	3.6	3.5	2.1-5.6	2.4	5.5	2.8-9.7	1.2	2.0	0.7-4.5	2.6	2.5	1.3-4.2	1.6	3.5	1.5-7.1	1.0	1.7	0.5-4.0
	1986-1990	5.2	5.4	3.5-7.9	4.2	9.3	5.7-14.3	1.0	1.9	0.6-4.4	5.2	5.4	3.5-7.9	4.0	10.0	6.0-15.5	1.2	2.1	0.8-4.6
	1991-1995	4.8	4.7	3.0-7.0	4.0	9.8	5.8-15.2	0.8	1.3	0.4-3.4	4.2	4.2	2.6-6.3	3.6	9.1	5.2-14.6	0.6	1.0	0.2-2.8
	1996-2000	8.2	7.4	5.3-10.0	6.8	13.9	9.5-19.6	1.4	2.3	0.9-4.7	7.0	6.4	4.4-8.9	6.0	13.5	8.9-19.5	1.0	1.6	0.5-3.8
	2001-2005	7.8	6.7	4.7-9.1	6.8	12.5	8.6-17.5	1.0	1.6	0.5-3.7	7.2	6.0	4.2-8.3	6.0	10.8	7.2-15.5	1.2	1.8	0.7-4.0
	2006-2010	9.6	7.2	5.3-9.6	8.2	13.3	9.4-18.2	1.4	2.0	0.8-4.2	6.6	5.1	3.5-7.2	5.8	9.9	6.5-14.3	0.8	1.2	0.3-3.2
	2011-2015	7.8	5.1	3.6-7.1	6.4	8.5	5.7-12.3	1.4	1.9	0.7-4.1	8.6	5.7	4.1-7.8	7.8	11.0	7.7-15.3	0.8	1.0	0.3-2.7
	2016-2020	7.6	4.5	3.2-6.3	6.8	8.2	5.6-11.8	0.8	0.9	0.2-2.7	8.6	5.5	3.9-7.5	7.6	10.1	7.1-14.2	1.0	1.3	0.4-3.3
Stomach	1976-1980	9.2	10.5	7.6-14.1	6.8	18.8	12.6-26.6	2.4	4.8	2.5-8.4	8.0	9.2	6.5-12.6	5.6	15.9	10.3-23.3	2.4	4.8	2.5-8.4
	1981-1985	6.4	7.2	4.9-10.1	4.0	11.1	6.6-17.3	2.4	4.5	2.3-7.8	3.8	4.4	2.6-6.9	2.4	7.2	3.6-12.6	1.4	2.6	1.0-5.3
	1986-1990	9.6	9.9	7.3-13.2	6.6	16.8	11.3-23.8	3.0	5.1	2.8-8.4	7.0	7.4	5.1-10.3	4.6	12.0	7.4-18.1	2.4	4.1	2.1-7.2
	1991-1995	8.4	8.0	5.7-10.8	5.8	12.5	8.3-18.1	2.6	4.2	2.3-7.2	6.0	5.6	3.8-8.0	3.8	8.5	5.0-13.3	2.2	3.5	1.7-6.3
	1996-2000	8.6	7.8	5.6-10.5	6.2	13.2	8.8-18.8	2.4	3.8	2.0-6.7	3.4	3.1	1.8-4.9	2.6	5.8	3.0-9.9	0.8	1.3	0.4-3.3
	2001-2005	7.4	6.2	4.4-8.6	4.8	9.3	5.8-14.1	2.6	3.7	2.0-6.4	3.8	3.2	1.9-5.0	2.4	4.7	2.3-8.3	1.4	1.9	0.8-4.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Oswego**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	8.4	6.7	4.8-9.1	5.8	10.2	6.7-14.7	2.6	3.8	2.0-6.5	3.4	2.8	1.6-4.6	2.6	5.0	2.6-8.6	0.8	1.1	0.3-3.0
	2011-2015	11.4	7.8	5.9-10.3	8.6	12.0	8.6-16.5	2.8	4.2	2.3-7.2	2.2	1.6	0.8-2.9	2.0	3.2	1.5-6.1	0.2	0.3	0.0-1.7
	2016-2020	11.2	7.5	5.6-9.9	8.0	11.1	7.7-15.4	3.2	4.3	2.4-7.2	2.6	1.8	0.9-3.1	1.6	2.2	0.9-4.6	1.0	1.5	0.5-3.6
Colorectal	1976-1980	52.4	59.9	52.7-67.8	22.6	59.0	48.0-71.5	29.8	59.7	50.4-70.2	24.8	28.8	23.9-34.5	9.4	24.0	17.3-32.4	15.4	31.0	24.4-38.8
	1981-1985	60.4	66.4	59.0-74.4	26.2	66.3	54.9-79.2	34.2	65.6	56.0-76.3	29.2	32.3	27.2-38.1	13.2	35.8	27.3-46.1	16.0	30.3	24.0-37.8
	1986-1990	60.4	62.5	55.6-70.0	26.4	68.0	56.3-81.2	34.0	59.8	51.1-69.6	30.2	32.3	27.3-38.0	11.2	30.1	22.3-39.4	19.0	33.6	27.2-41.2
	1991-1995	60.6	58.2	51.8-65.1	31.8	72.8	61.5-85.5	28.8	47.3	39.9-55.7	26.0	25.9	21.6-30.7	13.2	33.0	25.2-42.3	12.8	21.0	16.1-26.8
	1996-2000	71.4	65.3	58.7-72.4	39.2	84.0	72.3-97.0	32.2	51.2	43.6-59.8	27.2	25.1	21.0-29.7	16.0	36.3	28.4-45.4	11.2	17.8	13.4-23.1
	2001-2005	71.4	60.6	54.5-67.3	36.6	71.7	61.4-83.1	34.8	52.5	45.0-61.0	25.4	22.1	18.4-26.3	14.6	31.4	24.4-39.6	10.8	16.1	12.1-21.1
	2006-2010	66.6	52.6	47.0-58.6	33.2	57.5	48.8-67.2	33.4	48.7	41.5-56.8	19.8	15.7	12.7-19.1	9.4	17.7	12.8-23.7	10.4	14.7	10.9-19.4
	2011-2015	53.8	39.6	34.9-44.8	27.0	42.2	35.1-50.3	26.8	36.7	30.6-43.8	22.4	16.6	13.6-20.0	12.6	20.8	15.8-26.9	9.8	12.9	9.5-17.2
2016-2020	55.8	38.4	33.9-43.4	28.2	39.8	33.2-47.4	27.6	36.9	30.8-44.0	17.6	11.9	9.5-14.8	9.8	14.3	10.4-19.1	7.8	9.5	6.8-13.2	
Colon excluding rectum	1976-1980	40.4	46.1	39.8-53.0	16.0	42.4	33.2-53.3	24.4	48.2	39.9-57.6	20.0	23.2	18.8-28.3	6.8	16.8	11.4-23.8	13.2	26.5	20.5-33.8
	1981-1985	44.0	47.5	41.3-54.3	17.8	43.5	34.5-54.0	26.2	49.4	41.2-58.7	25.8	28.7	23.9-34.1	11.8	32.6	24.4-42.5	14.0	26.4	20.5-33.5
	1986-1990	42.4	44.2	38.4-50.7	17.2	45.2	35.6-56.3	25.2	44.3	36.9-52.8	25.6	27.3	22.8-32.5	9.0	23.3	16.7-31.6	16.6	29.5	23.5-36.6
	1991-1995	43.8	42.2	36.8-48.2	20.6	47.0	38.1-57.4	23.2	38.4	31.7-46.1	20.6	20.5	16.7-24.9	9.8	24.3	17.7-32.4	10.8	17.8	13.3-23.2
	1996-2000	50.0	46.0	40.4-52.0	24.6	54.4	44.9-65.2	25.4	40.3	33.5-47.9	20.8	19.1	15.6-23.2	12.0	25.9	19.6-33.5	8.8	13.8	10.0-18.6
	2001-2005	51.4	44.1	38.9-49.9	25.8	51.6	42.9-61.5	25.6	38.7	32.2-46.1	20.6	18.0	14.6-21.8	11.8	25.4	19.1-32.9	8.8	13.0	9.4-17.6
	2006-2010	44.6	35.6	31.0-40.7	20.8	37.2	30.2-45.4	23.8	34.8	28.7-41.8	13.8	10.9	8.5-13.8	7.2	13.7	9.4-19.1	6.6	9.2	6.3-13.0
	2011-2015	38.4	28.5	24.5-32.9	18.4	29.2	23.3-36.1	20.0	27.4	22.2-33.6	18.2	13.6	10.9-16.8	10.8	18.2	13.5-23.9	7.4	9.8	6.9-13.6
2016-2020	40.2	28.0	24.1-32.3	18.2	25.4	20.2-31.5	22.0	29.9	24.4-36.3	12.8	8.6	6.6-11.1	6.8	9.5	6.5-13.6	6.0	7.6	5.1-11.0	
Rectum & rectosigmoid	1976-1980	12.0	13.8	10.5-17.9	6.6	16.5	11.1-23.5	5.4	11.5	7.5-16.8	4.8	5.6	3.6-8.4	2.6	7.2	3.7-12.5	2.2	4.5	2.2-8.0
	1981-1985	16.4	18.8	14.9-23.4	8.4	22.8	16.1-31.1	8.0	16.2	11.5-22.1	3.4	3.6	2.1-5.9	1.4	3.2	1.2-6.7	2.0	3.9	1.8-7.2
	1986-1990	18.0	18.2	14.6-22.5	9.2	22.8	16.4-30.7	8.8	15.5	11.2-20.8	4.6	5.0	3.1-7.5	2.2	6.8	3.3-12.0	2.4	4.1	2.1-7.2
	1991-1995	16.8	15.9	12.7-19.7	11.2	25.8	19.2-33.7	5.6	8.9	5.9-12.9	5.4	5.4	3.5-7.8	3.4	8.7	4.9-14.1	2.0	3.2	1.5-5.9
	1996-2000	21.4	19.3	15.8-23.3	14.6	29.6	23.1-37.4	6.8	11.0	7.6-15.3	6.4	5.9	4.1-8.4	4.0	10.3	6.1-16.1	2.4	3.9	2.0-6.9
	2001-2005	20.0	16.5	13.4-20.1	10.8	20.1	15.0-26.4	9.2	13.9	10.1-18.5	4.8	4.2	2.7-6.2	2.8	6.0	3.2-10.1	2.0	3.1	1.5-5.8
	2006-2010	22.0	17.0	13.9-20.5	12.4	20.2	15.4-26.1	9.6	13.9	10.2-18.5	6.0	4.8	3.2-6.9	2.2	4.0	1.9-7.3	3.8	5.5	3.3-8.7
	2011-2015	15.4	11.1	8.7-14.0	8.6	13.0	9.3-17.8	6.8	9.3	6.4-13.2	4.2	3.0	1.8-4.6	1.8	2.6	1.1-5.2	2.4	3.1	1.6-5.6
2016-2020	15.6	10.4	8.2-13.2	10.0	14.5	10.6-19.4	5.6	7.1	4.7-10.4	4.8	3.2	2.0-4.9	3.0	4.8	2.6-8.0	1.8	2.0	0.9-4.0	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Oswego**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	2.2	2.2	1.1-3.9	1.6	3.1	1.3-6.4	0.6	1.2	0.2-3.6	1.8	2.0	0.9-3.7	0.8	1.5	0.4-4.1	1.0	2.3	0.7-5.3
	1981-1985	2.8	3.2	1.7-5.3	1.6	4.1	1.6-8.2	1.2	2.6	0.9-5.5	1.8	1.9	0.8-3.6	1.4	3.1	1.2-6.6	0.4	0.8	0.1-2.9
	1986-1990	3.2	3.3	1.9-5.4	1.8	4.0	1.8-7.6	1.4	2.6	1.0-5.3	3.0	3.1	1.7-5.1	1.8	4.2	1.8-8.1	1.2	2.2	0.8-4.9
	1991-1995	4.0	4.0	2.4-6.1	2.8	6.5	3.5-11.1	1.2	2.2	0.8-4.7	2.6	2.5	1.3-4.3	2.0	4.6	2.1-8.6	0.6	1.0	0.2-2.8
	1996-2000	4.6	4.2	2.6-6.2	2.8	6.2	3.3-10.6	1.8	3.0	1.4-5.6	4.6	4.2	2.6-6.2	2.2	4.5	2.2-8.1	2.4	3.8	2.0-6.7
	2001-2005	5.4	4.6	3.1-6.8	4.0	7.3	4.4-11.3	1.4	2.3	0.9-4.8	6.2	5.5	3.7-7.8	4.4	8.5	5.3-12.9	1.8	3.0	1.4-5.7
	2006-2010	4.2	3.3	2.0-5.0	2.8	5.0	2.6-8.4	1.4	2.0	0.8-4.2	6.4	4.9	3.4-7.0	3.4	6.2	3.5-10.0	3.0	4.0	2.2-6.7
	2011-2015	11.0	7.7	5.7-10.1	6.8	9.6	6.5-13.6	4.2	5.6	3.4-8.7	7.6	5.6	4.0-7.8	5.0	8.4	5.3-12.6	2.6	3.5	1.8-6.1
	2016-2020	13.2	9.0	6.9-11.6	10.0	14.5	10.6-19.4	3.2	4.4	2.4-7.3	8.8	6.1	4.4-8.3	4.8	7.4	4.6-11.2	4.0	5.0	3.0-8.0
Pancreas	1976-1980	11.6	12.4	9.4-16.2	5.6	13.1	8.6-19.1	6.0	11.4	7.7-16.3	11.2	11.7	8.8-15.3	6.0	13.9	9.1-20.2	5.2	9.8	6.4-14.4
	1981-1985	10.2	11.5	8.5-15.1	5.6	17.1	11.1-25.0	4.6	8.7	5.5-13.2	9.2	9.8	7.2-13.2	4.4	12.2	7.4-18.8	4.8	8.7	5.5-12.9
	1986-1990	11.0	11.3	8.5-14.7	5.0	12.3	7.7-18.3	6.0	10.4	7.0-14.9	12.4	12.7	9.7-16.3	6.2	15.4	10.3-22.1	6.2	10.7	7.2-15.2
	1991-1995	13.0	12.6	9.7-16.0	6.4	14.4	9.7-20.4	6.6	11.1	7.6-15.6	9.4	9.3	6.8-12.3	4.8	11.4	7.2-17.1	4.6	7.7	4.9-11.6
	1996-2000	13.0	11.9	9.2-15.2	5.4	11.8	7.6-17.3	7.6	12.1	8.5-16.6	14.8	13.6	10.7-17.1	5.6	12.7	8.3-18.5	9.2	14.6	10.7-19.5
	2001-2005	14.6	12.4	9.7-15.6	8.6	16.5	11.8-22.4	6.0	9.1	6.1-13.0	11.2	9.5	7.2-12.3	5.6	10.9	7.1-15.8	5.6	8.5	5.6-12.3
	2006-2010	19.0	15.0	12.1-18.4	9.8	17.2	12.6-23.0	9.2	13.4	9.7-17.9	15.8	12.7	10.0-15.9	7.6	13.5	9.4-18.6	8.2	11.9	8.5-16.2
	2011-2015	18.0	13.0	10.4-16.1	9.0	14.5	10.4-19.6	9.0	12.1	8.8-16.4	14.8	11.1	8.7-14.0	6.6	11.1	7.5-15.8	8.2	11.1	7.9-15.2
	2016-2020	27.0	17.4	14.6-20.8	15.6	21.3	16.6-26.9	11.4	13.9	10.5-18.2	19.2	12.3	9.9-15.1	9.4	13.5	9.8-18.2	9.8	11.6	8.6-15.6
Larynx	1976-1980	6.4	6.8	4.6-9.6	4.2	9.8	5.9-15.1	2.2	4.7	2.3-8.3	3.2	3.3	1.9-5.4	2.0	5.0	2.3-9.3	1.2	2.2	0.8-4.8
	1981-1985	8.0	8.4	5.9-11.5	6.0	14.1	9.3-20.5	2.0	3.9	1.8-7.1	3.0	2.9	1.6-4.8	2.4	5.4	2.7-9.6	0.6	1.0	0.2-3.0
	1986-1990	7.0	7.3	5.0-10.1	5.4	12.6	8.2-18.5	1.6	3.1	1.3-6.1	2.4	2.3	1.2-4.0	2.0	4.1	2.0-7.7	0.4	0.7	0.1-2.5
	1991-1995	7.6	7.1	5.0-9.8	5.4	10.9	7.1-15.9	2.2	4.0	2.0-7.1	1.8	1.8	0.8-3.3	1.0	2.1	0.7-4.8	0.8	1.3	0.4-3.4
	1996-2000	6.0	5.3	3.6-7.6	4.4	8.6	5.4-13.1	1.6	2.7	1.2-5.3	2.2	2.0	1.0-3.5	1.8	3.4	1.5-6.5	0.4	0.6	0.1-2.3
	2001-2005	6.6	5.3	3.6-7.5	5.0	8.6	5.5-12.8	1.6	2.4	1.0-4.8	1.4	1.2	0.5-2.5	1.0	1.8	0.6-4.3	0.4	0.6	0.1-2.2
	2006-2010	8.4	6.5	4.7-8.9	6.8	11.5	7.9-16.3	1.6	2.4	1.0-4.9	2.0	1.5	0.7-2.9	1.6	2.9	1.2-5.9	0.4	0.5	0.1-2.0
	2011-2015	7.8	5.1	3.6-7.0	5.8	7.9	5.2-11.5	2.0	2.5	1.2-4.8	1.8	1.2	0.5-2.4	1.2	1.6	0.6-3.8	0.6	0.8	0.2-2.5
	2016-2020	7.6	5.0	3.5-7.0	5.6	7.4	4.8-10.9	2.0	2.8	1.3-5.3	1.6	0.9	0.4-2.0	1.2	1.4	0.5-3.3	0.4	0.5	0.1-2.1
Lung and bronchus	1976-1980	61.0	67.4	59.9-75.6	47.6	123.5	107.5-141.1	13.4	27.3	21.0-34.8	49.6	55.8	48.9-63.3	37.0	99.4	84.9-115.5	12.6	25.2	19.2-32.3
	1981-1985	65.2	66.7	59.5-74.5	46.0	107.0	93.1-122.4	19.2	37.2	29.9-45.5	51.2	52.7	46.3-59.7	36.0	87.4	74.4-101.8	15.2	28.7	22.5-36.1
	1986-1990	86.4	86.0	77.9-94.6	60.0	136.7	121.1-153.6	26.4	47.7	39.8-56.7	67.6	67.1	60.1-74.8	49.2	112.2	98.1-127.7	18.4	33.6	27.0-41.3
	1991-1995	86.0	80.5	73.1-88.6	47.8	102.8	89.8-117.1	38.2	65.1	56.1-75.0	71.4	66.9	60.1-74.2	43.6	96.0	83.2-110.1	27.8	46.5	39.0-54.9
	1996-2000	102.6	92.2	84.4-100.5	61.6	127.4	113.2-142.8	41.0	67.5	58.6-77.4	75.0	67.6	60.9-74.8	45.2	93.2	81.2-106.5	29.8	48.5	41.0-56.9
	2001-2005	112.4	94.8	87.1-103.0	59.8	114.4	101.5-128.5	52.6	82.0	72.3-92.6	78.4	66.2	59.8-73.2	43.6	84.7	73.5-97.0	34.8	53.8	46.1-62.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Oswego**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	129.2	101.8	94.0-110.1	64.4	112.8	100.4-126.2	64.8	95.1	84.9-106.2	89.8	71.1	64.6-78.1	48.4	86.0	75.2-97.9	41.4	60.8	52.7-69.8
	2011-2015	136.2	95.8	88.6-103.5	71.0	105.6	94.4-117.7	65.2	87.9	78.5-98.3	86.0	61.9	56.1-68.2	47.4	74.3	64.8-84.8	38.6	52.1	44.9-60.2
	2016-2020	151.0	97.4	90.4-104.9	76.8	105.2	94.5-116.8	74.2	90.7	81.5-100.7	74.8	48.5	43.6-53.9	43.0	60.3	52.2-69.3	31.8	38.5	32.7-45.2
Melanoma of the skin	1976-1980	6.2	7.0	4.7-9.9	3.6	9.7	5.7-15.4	2.6	4.8	2.5-8.3	2.4	2.6	1.3-4.5	1.8	4.1	1.8-7.8	0.6	1.2	0.2-3.4
	1981-1985	6.4	7.0	4.7-9.9	2.8	6.3	3.4-10.7	3.6	7.3	4.3-11.6	1.2	1.4	0.5-2.9	0.8	1.8	0.5-4.7	0.4	0.9	0.1-3.2
	1986-1990	6.8	6.4	4.4-9.0	3.0	5.9	3.3-9.9	3.8	6.4	3.8-10.1	3.6	3.6	2.1-5.6	2.2	4.4	2.2-7.9	1.4	2.7	1.1-5.5
	1991-1995	7.6	6.9	4.8-9.4	4.6	9.3	5.8-14.0	3.0	4.8	2.6-7.9	3.0	3.0	1.6-4.8	2.2	5.2	2.5-9.3	0.8	1.3	0.3-3.3
	1996-2000	11.0	9.7	7.3-12.6	6.4	13.2	8.9-18.9	4.6	7.6	4.8-11.5	1.4	1.3	0.5-2.6	1.0	2.5	0.7-5.7	0.4	0.7	0.1-2.4
	2001-2005	16.6	13.6	10.8-16.9	8.4	14.3	10.2-19.4	8.2	12.9	9.2-17.6	3.4	2.7	1.6-4.4	1.8	3.0	1.3-5.8	1.6	2.4	1.0-4.7
	2006-2010	20.2	16.3	13.2-19.9	11.2	19.1	14.3-24.9	9.0	14.3	10.4-19.3	5.2	3.9	2.6-5.8	3.2	4.9	2.8-8.1	2.0	2.7	1.3-5.2
	2011-2015	28.0	21.6	18.0-25.6	15.2	24.8	19.4-31.4	12.8	19.7	15.0-25.4	3.4	2.7	1.5-4.4	2.8	5.1	2.7-8.7	0.6	0.8	0.2-2.4
2016-2020	26.6	19.0	15.7-22.7	12.2	16.9	12.7-22.0	14.4	21.4	16.5-27.3	2.2	1.8	0.9-3.3	1.0	1.7	0.5-4.2	1.2	1.8	0.6-4.1	
Female breast	1976-1980							43.4	91.2	79.3-104.4							16.8	35.5	28.2-44.0
	1981-1985							54.8	110.4	97.4-124.5							17.4	34.9	27.8-43.2
	1986-1990							61.8	113.3	100.8-126.8							18.6	34.5	27.8-42.4
	1991-1995							74.0	128.4	115.5-142.2							20.8	35.2	28.7-42.7
	1996-2000							80.2	132.7	120.0-146.3							15.2	24.5	19.3-30.6
	2001-2005							85.4	131.7	119.4-144.9							14.8	22.6	17.7-28.4
	2006-2010							72.2	105.8	95.0-117.5							14.4	20.4	15.9-25.8
	2011-2015							89.6	123.4	112.0-135.8							13.2	17.6	13.5-22.6
2016-2020							103.2	141.5	129.0-155.0							13.8	18.0	13.9-23.1	
Cervix uteri	1976-1980							7.4	14.8	10.3-20.5							2.4	4.6	2.4-8.1
	1981-1985							7.6	15.3	10.7-21.2							2.0	4.4	2.0-8.0
	1986-1990							10.4	18.5	13.7-24.3							3.2	5.9	3.3-9.5
	1991-1995							10.4	17.4	12.9-22.8							3.2	5.7	3.2-9.2
	1996-2000							8.4	13.7	9.9-18.6							3.2	5.4	3.1-8.7
	2001-2005							5.8	8.9	5.9-12.8							3.0	4.7	2.6-7.8
	2006-2010							5.2	8.2	5.3-12.1							3.0	4.2	2.3-7.1
	2011-2015							3.6	5.4	3.1-8.7							1.4	2.1	0.8-4.4
	2016-2020							3.6	6.5	3.8-10.4							2.0	3.2	1.4-6.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Oswego**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							12.2	23.2	17.6-29.8							2.0	4.1	1.9-7.6
	1981-1985							15.4	29.1	22.9-36.6							3.2	5.8	3.3-9.5
	1986-1990							13.8	25.4	19.7-32.3							2.6	4.9	2.6-8.4
	1991-1995							17.0	29.4	23.5-36.4							3.2	5.3	3.0-8.5
	1996-2000							17.0	28.0	22.3-34.6							1.6	2.5	1.1-5.0
	2001-2005							19.8	30.6	24.9-37.3							2.0	3.1	1.5-5.7
	2006-2010							22.4	32.2	26.4-38.9							2.0	2.9	1.4-5.5
	2011-2015							27.0	36.6	30.5-43.7							1.8	2.4	1.1-4.7
	2016-2020							29.2	37.5	31.4-44.5							4.8	6.0	3.8-9.1
Ovary	1976-1980							7.4	14.7	10.3-20.4							7.2	14.0	9.8-19.5
	1981-1985							5.4	10.2	6.6-14.9							4.0	7.9	4.8-12.2
	1986-1990							8.6	16.3	11.8-22.1							3.2	5.8	3.3-9.5
	1991-1995							7.8	13.5	9.6-18.5							5.6	9.5	6.3-13.7
	1996-2000							7.6	12.6	8.9-17.3							5.8	9.4	6.3-13.5
	2001-2005							9.6	14.8	10.9-19.7							4.6	7.1	4.5-10.7
	2006-2010							7.2	10.3	7.1-14.3							5.0	7.0	4.5-10.5
	2011-2015							9.2	12.6	9.1-17.0							4.8	6.7	4.3-10.1
	2016-2020							8.0	11.6	8.1-16.1							5.4	6.6	4.3-9.8
Prostate	1976-1980				28.2	81.1	67.5-96.4							8.8	28.8	20.5-38.9			
	1981-1985				33.2	91.2	77.0-107.0							12.2	36.3	27.3-47.0			
	1986-1990				42.6	109.7	94.8-126.2							9.6	29.6	21.4-39.6			
	1991-1995				68.2	158.2	141.1-176.7							12.4	37.9	28.7-48.8			
	1996-2000				71.4	152.5	136.6-169.7							11.2	29.7	22.1-38.9			
	2001-2005				97.8	184.5	168.1-202.1							11.4	27.9	21.0-36.3			
	2006-2010				101.8	164.6	150.2-180.0							10.8	22.6	16.9-29.6			
	2011-2015				88.4	121.3	109.9-133.7							9.6	18.9	13.9-25.1			
	2016-2020				97.2	120.0	109.2-131.6							5.6	9.1	6.0-13.3			
Testis	1976-1980				1.0	1.9	0.6-4.6							0.4	0.9	0.1-3.4			
	1981-1985				2.8	4.9	2.4-8.9							0.2	0.3	0.0-2.2			
	1986-1990				2.4	3.9	2.0-7.0							0.0	0.0	0.0-1.6			
	1991-1995				3.4	4.9	2.8-8.0							0.0	0.0	0.0-1.5			
	1996-2000				2.2	3.5	1.7-6.4							0.2	0.3	0.0-2.0			
	2001-2005				5.6	9.2	6.1-13.4							0.2	0.4	0.0-2.0			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Oswego**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				3.2	5.9	3.3-9.6							0.0	0.0	0.0-1.3			
	2011-2015				4.6	9.0	5.6-13.6							0.2	0.3	0.0-1.7			
	2016-2020				5.4	10.5	6.8-15.4							0.2	0.5	0.0-2.4			
Urinary bladder (incl. in situ)	1976-1980	16.2	18.0	14.2-22.5	11.6	29.7	22.2-38.9	4.6	9.1	5.7-13.7	5.2	6.5	4.2-9.5	3.4	10.2	5.8-16.3	1.8	3.7	1.7-7.0
	1981-1985	22.4	24.6	20.2-29.7	16.4	43.4	34.0-54.4	6.0	10.9	7.4-15.6	6.6	7.7	5.3-10.8	5.0	16.1	10.2-23.9	1.6	3.0	1.3-5.9
	1986-1990	23.8	24.5	20.3-29.4	16.8	41.6	32.9-51.9	7.0	12.3	8.5-17.1	6.4	7.0	4.7-9.8	3.8	10.8	6.3-17.0	2.6	4.4	2.4-7.6
	1991-1995	24.8	23.9	19.8-28.5	16.6	40.0	31.4-50.0	8.2	13.1	9.4-17.8	6.6	6.6	4.5-9.2	3.6	9.6	5.5-15.3	3.0	4.7	2.6-7.8
	1996-2000	27.8	25.2	21.2-29.8	20.4	43.7	35.5-53.3	7.4	12.0	8.4-16.5	8.2	7.7	5.5-10.4	6.6	16.5	11.1-23.3	1.6	2.4	1.1-4.9
	2001-2005	40.8	34.6	30.0-39.7	31.0	62.7	53.0-73.7	9.8	14.7	10.8-19.4	5.8	5.0	3.3-7.2	4.8	10.6	6.7-15.9	1.0	1.4	0.5-3.5
	2006-2010	38.8	31.3	27.0-36.1	28.6	53.1	44.5-62.7	10.2	14.8	11.0-19.6	9.8	8.1	5.9-10.7	7.2	14.6	10.1-20.3	2.6	3.6	1.9-6.2
	2011-2015	37.4	27.0	23.2-31.3	27.6	44.1	36.8-52.4	9.8	13.2	9.7-17.6	7.2	5.4	3.8-7.5	4.8	8.3	5.2-12.5	2.4	3.2	1.7-5.8
2016-2020	44.2	28.7	25.0-32.9	36.8	51.6	44.2-60.0	7.4	9.0	6.3-12.7	4.8	3.3	2.1-5.0	3.6	5.4	3.1-8.8	1.2	1.6	0.6-3.7	
Kidney and renal pelvis	1976-1980	7.2	7.6	5.2-10.5	5.2	11.6	7.5-17.2	2.0	3.9	1.8-7.2	3.6	3.7	2.1-5.9	2.4	5.1	2.6-9.2	1.2	2.2	0.8-4.8
	1981-1985	10.4	11.2	8.3-14.7	6.0	14.2	9.4-20.4	4.4	8.9	5.5-13.5	4.8	5.3	3.4-7.8	3.2	7.9	4.4-12.9	1.6	3.3	1.4-6.4
	1986-1990	12.0	11.9	9.1-15.4	7.4	16.7	11.6-23.2	4.6	8.5	5.3-12.8	6.2	6.3	4.2-8.9	3.8	8.9	5.2-14.1	2.4	4.4	2.3-7.8
	1991-1995	14.8	13.7	10.7-17.2	8.6	17.5	12.5-23.7	6.2	10.8	7.3-15.3	6.2	6.0	4.1-8.5	3.8	8.6	5.1-13.5	2.4	4.0	2.1-7.0
	1996-2000	13.0	11.7	9.0-14.9	6.8	14.3	9.7-20.2	6.2	10.1	6.8-14.3	4.6	4.2	2.7-6.3	2.4	5.8	2.8-10.2	2.2	3.6	1.8-6.4
	2001-2005	17.0	14.1	11.3-17.5	10.2	18.3	13.5-24.1	6.8	10.8	7.4-15.1	4.8	4.0	2.5-5.9	3.0	5.3	2.9-8.7	1.8	2.8	1.3-5.4
	2006-2010	27.8	21.8	18.2-25.8	17.2	29.0	23.0-36.0	10.6	16.0	12.0-21.1	5.4	4.3	2.8-6.3	3.0	5.2	2.8-8.7	2.4	3.6	1.9-6.4
	2011-2015	27.2	18.7	15.6-22.2	16.2	22.4	17.6-28.1	11.0	15.1	11.3-19.9	4.6	3.2	2.0-4.9	3.2	4.8	2.7-7.9	1.4	1.8	0.7-4.0
2016-2020	36.2	24.9	21.2-29.0	23.6	32.1	26.3-38.9	12.6	18.0	13.6-23.3	5.6	3.6	2.4-5.3	4.0	5.4	3.2-8.6	1.6	2.0	0.9-4.3	
Brain and other central nervous system	1976-1980	5.2	5.5	3.6-8.1	3.2	6.9	3.8-11.3	2.0	4.2	2.0-7.7	4.4	5.1	3.1-7.7	3.0	7.2	4.0-12.1	1.4	2.9	1.2-6.0
	1981-1985	8.2	7.4	5.3-10.1	4.4	8.8	5.5-13.6	3.8	6.3	3.7-9.9	4.6	4.5	2.8-6.8	2.4	4.9	2.5-8.7	2.2	4.3	2.1-7.8
	1986-1990	7.0	6.1	4.2-8.6	3.2	6.4	3.6-10.5	3.8	5.8	3.5-9.2	5.8	5.5	3.7-8.0	4.0	8.6	5.1-13.6	1.8	2.8	1.3-5.3
	1991-1995	8.8	7.6	5.5-10.2	6.2	11.2	7.5-16.1	2.6	4.3	2.3-7.3	4.8	4.2	2.7-6.3	3.2	6.2	3.5-10.2	1.6	2.5	1.1-5.0
	1996-2000	9.6	8.3	6.1-11.0	5.0	9.1	5.9-13.5	4.6	7.5	4.7-11.2	5.2	4.6	3.0-6.8	2.8	5.0	2.7-8.5	2.4	3.8	2.0-6.7
	2001-2005	9.2	7.7	5.6-10.3	5.4	9.5	6.2-13.9	3.8	5.9	3.6-9.3	4.6	3.8	2.4-5.7	2.6	5.0	2.6-8.6	2.0	3.0	1.5-5.6
	2006-2010	10.2	8.1	6.0-10.7	6.4	11.4	7.7-16.2	3.8	5.6	3.3-8.8	6.4	5.0	3.4-7.1	4.2	7.4	4.5-11.4	2.2	3.1	1.5-5.7
	2011-2015	13.2	10.4	8.0-13.3	7.2	10.7	7.4-15.1	6.0	10.0	6.7-14.5	7.4	5.5	3.8-7.7	4.2	6.3	3.8-9.8	3.2	4.8	2.7-7.9
2016-2020	9.6	7.5	5.4-10.0	5.6	8.5	5.5-12.5	4.0	6.5	3.9-10.1	6.6	4.8	3.3-6.9	3.8	5.2	3.1-8.4	2.8	4.4	2.3-7.5	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Oswego**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	2.2	2.7	1.3-4.9	0.6	1.6	0.3-4.5	1.6	3.7	1.6-7.3	0.8	1.0	0.3-2.6	0.4	1.1	0.1-3.7	0.4	1.1	0.1-3.6
	1981-1985	2.8	3.1	1.7-5.2	0.8	1.9	0.5-4.8	2.0	4.4	2.1-8.0	1.2	1.3	0.5-2.7	0.4	1.0	0.1-3.5	0.8	1.5	0.4-4.0
	1986-1990	2.6	2.5	1.3-4.2	0.8	1.6	0.4-4.2	1.8	3.4	1.5-6.4	0.0	0.0	0.0-0.7	0.0	0.0	0.0-1.6	0.0	0.0	0.0-1.3
	1991-1995	5.0	4.4	2.9-6.6	1.2	2.0	0.7-4.5	3.8	6.4	3.9-10.1	0.4	0.4	0.0-1.4	0.2	0.4	0.0-2.1	0.2	0.3	0.0-1.8
	1996-2000	7.0	5.9	4.1-8.2	1.6	2.7	1.2-5.5	5.4	8.8	5.8-12.8	0.8	0.8	0.2-1.9	0.0	0.0	0.0-1.4	0.8	1.2	0.3-3.2
	2001-2005	12.2	10.1	7.7-13.0	3.4	5.9	3.4-9.5	8.8	14.6	10.6-19.6	0.0	0.0	0.0-0.6	0.0	0.0	0.0-1.4	0.0	0.0	0.0-1.2
	2006-2010	19.0	15.1	12.1-18.5	4.4	7.0	4.3-10.7	14.6	23.1	18.0-29.1	1.0	0.7	0.2-1.7	0.0	0.0	0.0-1.3	1.0	1.3	0.4-3.2
	2011-2015	20.8	17.2	13.9-20.9	5.2	8.5	5.4-12.7	15.6	25.7	20.1-32.3	0.2	0.1	0.0-0.8	0.2	0.3	0.0-1.7	0.0	0.0	0.0-1.3
	2016-2020	15.0	11.7	9.1-14.8	4.0	6.3	3.7-9.9	11.0	17.3	12.9-22.9	0.8	0.5	0.1-1.4	0.6	1.0	0.2-3.0	0.2	0.2	0.0-1.6
Hodgkin lymphoma	1976-1980	3.2	2.9	1.6-4.9	2.2	4.4	2.0-8.2	1.0	1.8	0.5-4.3	1.2	1.1	0.4-2.5	1.0	1.8	0.6-4.5	0.2	0.4	0.0-2.3
	1981-1985	1.8	1.5	0.7-3.0	0.6	1.5	0.3-4.3	1.2	1.7	0.6-3.9	1.2	1.2	0.4-2.6	0.8	1.8	0.4-4.8	0.4	0.6	0.1-2.4
	1986-1990	3.2	2.6	1.5-4.4	1.4	2.7	1.1-5.6	1.8	2.6	1.1-5.1	0.4	0.4	0.0-1.4	0.2	0.9	0.0-4.0	0.2	0.2	0.0-1.6
	1991-1995	4.6	3.6	2.3-5.5	1.8	3.1	1.4-6.0	2.8	4.2	2.3-7.1	1.0	0.8	0.3-1.9	0.2	0.4	0.0-2.1	0.8	1.2	0.3-3.0
	1996-2000	3.0	2.4	1.4-4.0	1.4	2.3	0.9-4.8	1.6	2.5	1.1-5.0	0.8	0.7	0.2-1.8	0.6	1.2	0.2-3.6	0.2	0.3	0.0-1.8
	2001-2005	3.4	2.8	1.6-4.5	1.4	2.4	0.9-5.0	2.0	3.2	1.5-6.0	0.4	0.3	0.0-1.2	0.2	0.3	0.0-1.8	0.2	0.3	0.0-1.7
	2006-2010	2.8	2.1	1.1-3.5	1.8	2.8	1.2-5.5	1.0	1.6	0.5-3.7	1.0	0.7	0.2-1.6	0.8	1.3	0.3-3.4	0.2	0.3	0.0-1.7
	2011-2015	5.4	4.6	3.0-6.7	3.8	6.6	3.9-10.5	1.6	2.6	1.1-5.2	0.6	0.3	0.1-1.1	0.6	0.7	0.1-2.3	0.0	0.0	0.0-1.3
	2016-2020	4.0	3.4	2.0-5.4	3.0	5.1	2.8-8.6	1.0	1.7	0.5-4.1	0.6	0.4	0.1-1.4	0.6	0.9	0.1-2.8	0.0	0.0	0.0-1.3
Non-Hodgkin lymphomas	1976-1980	11.0	11.8	8.8-15.5	5.8	12.6	8.3-18.3	5.2	10.3	6.7-15.1	6.2	6.8	4.6-9.7	2.6	5.5	2.8-9.7	3.6	7.3	4.3-11.5
	1981-1985	11.8	12.4	9.4-16.1	5.6	13.9	9.0-20.4	6.2	11.2	7.6-15.9	7.2	7.6	5.3-10.5	3.8	9.1	5.3-14.4	3.4	6.0	3.5-9.7
	1986-1990	16.2	16.5	13.0-20.5	7.8	18.7	13.1-25.7	8.4	14.7	10.6-20.0	6.4	6.8	4.6-9.6	3.2	8.2	4.6-13.3	3.2	5.6	3.2-9.2
	1991-1995	18.2	16.6	13.3-20.4	8.0	16.5	11.6-22.6	10.2	17.2	12.8-22.7	10.0	9.2	6.8-12.2	4.8	10.2	6.5-15.4	5.2	8.3	5.4-12.2
	1996-2000	24.8	22.0	18.3-26.3	11.2	22.5	16.8-29.4	13.6	22.2	17.2-28.2	8.8	8.0	5.8-10.7	3.4	7.3	4.2-11.8	5.4	8.5	5.6-12.4
	2001-2005	31.0	25.9	22.0-30.3	17.4	32.4	25.7-40.2	13.6	20.6	16.0-26.2	10.6	9.0	6.8-11.8	5.0	10.4	6.7-15.5	5.6	8.5	5.7-12.4
	2006-2010	28.6	22.5	18.9-26.6	15.0	25.4	19.8-32.1	13.6	20.5	15.8-26.1	9.0	6.9	5.0-9.2	5.0	8.7	5.5-13.1	4.0	5.3	3.2-8.3
	2011-2015	34.0	24.9	21.2-29.1	19.2	29.5	23.7-36.4	14.8	20.2	15.8-25.6	9.0	6.7	4.9-9.1	5.2	8.6	5.5-12.7	3.8	5.1	3.0-8.1
	2016-2020	34.8	23.5	20.0-27.4	18.2	25.8	20.6-32.1	16.6	21.6	17.1-27.0	7.0	5.0	3.4-7.0	3.2	5.0	2.8-8.2	3.8	4.9	2.9-7.8
Myeloma	1976-1980	2.2	2.6	1.3-4.6	1.6	4.7	1.9-9.3	0.6	1.1	0.2-3.3	1.8	2.3	1.0-4.3	1.0	3.0	0.9-7.1	0.8	1.7	0.5-4.2
	1981-1985	4.0	4.5	2.7-6.9	2.0	4.9	2.2-9.3	2.0	4.2	2.0-7.6	2.4	2.6	1.3-4.6	0.8	2.1	0.5-5.6	1.6	3.1	1.3-6.1
	1986-1990	4.0	4.1	2.5-6.4	1.8	5.0	2.1-9.6	2.2	3.7	1.8-6.6	3.0	2.9	1.6-4.8	1.8	4.5	1.9-8.6	1.2	2.1	0.8-4.5
	1991-1995	3.8	3.6	2.2-5.6	2.4	5.5	2.7-9.6	1.4	2.5	1.0-5.1	2.8	2.7	1.5-4.5	1.6	3.9	1.6-7.7	1.2	1.9	0.7-4.2
	1996-2000	8.0	7.2	5.1-9.8	4.2	9.2	5.6-14.2	3.8	6.2	3.7-9.7	4.8	4.4	2.8-6.5	2.8	6.2	3.3-10.5	2.0	3.2	1.5-5.9
	2001-2005	8.2	7.0	5.0-9.5	4.6	8.9	5.5-13.4	3.6	5.5	3.3-8.7	4.8	4.1	2.6-6.1	2.6	5.3	2.8-9.2	2.2	3.4	1.7-6.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Oswego**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	5.6	4.1	2.7-6.0	3.0	5.2	2.8-8.7	2.6	3.3	1.8-5.9	4.2	3.4	2.1-5.2	1.8	3.3	1.5-6.3	2.4	3.2	1.6-5.8
	2011-2015	8.0	5.7	4.0-7.9	4.8	7.6	4.8-11.6	3.2	4.1	2.3-6.8	4.4	3.3	2.1-5.1	3.0	5.3	2.9-8.9	1.4	1.9	0.8-4.1
	2016-2020	10.8	7.1	5.3-9.4	6.2	8.2	5.5-11.9	4.6	5.9	3.7-9.2	3.6	2.5	1.5-4.1	1.4	2.3	0.9-4.8	2.2	2.7	1.4-5.1
Leukemias	1976-1980	12.0	13.3	10.0-17.2	7.0	17.6	11.9-24.9	5.0	10.3	6.6-15.2	7.0	8.0	5.5-11.1	4.6	12.3	7.5-18.8	2.4	4.9	2.5-8.6
	1981-1985	11.6	12.1	9.1-15.7	6.6	16.2	10.8-23.1	5.0	9.1	5.9-13.5	7.4	8.1	5.7-11.2	3.8	10.4	6.0-16.4	3.6	6.6	3.9-10.5
	1986-1990	14.0	13.7	10.6-17.4	6.6	14.8	9.9-21.1	7.4	12.8	9.0-17.7	9.2	9.2	6.7-12.4	3.8	8.4	4.9-13.4	5.4	9.3	6.1-13.6
	1991-1995	15.0	13.8	10.8-17.3	9.4	19.9	14.4-26.8	5.6	9.1	6.0-13.2	8.4	7.7	5.6-10.5	5.4	11.0	7.1-16.1	3.0	5.0	2.8-8.2
	1996-2000	16.6	14.9	11.9-18.5	8.6	18.3	13.0-24.8	8.0	12.7	9.1-17.3	7.8	7.1	5.0-9.7	4.6	9.7	6.1-14.7	3.2	5.0	2.9-8.2
	2001-2005	21.4	18.2	14.9-22.0	13.6	26.1	20.1-33.4	7.8	12.0	8.5-16.4	9.2	7.9	5.7-10.5	5.8	11.4	7.6-16.5	3.4	5.3	3.1-8.5
	2006-2010	22.6	17.4	14.3-20.9	14.8	24.5	19.1-31.0	7.8	11.4	8.0-15.7	9.4	7.7	5.6-10.2	6.4	11.7	7.9-16.7	3.0	4.5	2.5-7.6
	2011-2015	24.2	18.4	15.2-22.1	15.8	26.1	20.5-32.7	8.4	12.0	8.5-16.4	8.0	6.0	4.2-8.2	4.4	7.6	4.7-11.7	3.6	5.0	2.9-8.1
2016-2020	22.4	15.7	12.9-19.1	12.6	19.4	14.8-25.1	9.8	13.0	9.5-17.5	9.6	6.8	5.0-9.1	6.0	10.0	6.7-14.5	3.6	4.3	2.5-6.9	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Otsego**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	249.4	417.2	393.7-441.7	127.6	496.9	457.7-538.4	121.8	371.3	341.2-403.5	115.4	191.3	175.7-207.9	61.0	242.0	214.7-271.7	54.4	157.1	138.5-177.7
	1981-1985	278.8	447.6	423.9-472.4	148.0	552.5	512.3-595.1	130.8	384.1	354.0-416.2	121.4	192.3	176.9-208.6	67.0	254.1	226.8-283.8	54.4	153.7	135.2-174.0
	1986-1990	305.6	475.5	451.5-500.5	150.6	538.0	499.4-578.6	155.0	440.0	408.3-473.4	130.0	199.2	184.0-215.4	66.4	245.5	219.2-274.0	63.6	169.8	151.0-190.3
	1991-1995	325.6	489.4	465.6-514.1	171.8	586.6	547.5-627.8	153.8	421.6	391.5-453.4	140.8	207.1	191.9-223.1	74.6	260.5	234.4-288.8	66.2	171.8	153.3-191.9
	1996-2000	340.0	494.0	470.6-518.2	173.0	563.3	526.1-602.5	167.0	446.9	416.5-479.1	135.6	190.4	176.3-205.4	73.6	245.8	221.0-272.5	62.0	152.3	135.5-170.8
	2001-2005	357.8	498.5	475.3-522.5	190.8	589.2	551.9-628.3	167.0	434.7	405.0-466.2	150.2	200.8	186.6-215.9	83.0	263.7	238.7-290.7	67.2	158.4	141.6-176.9
	2006-2010	366.6	477.0	454.8-500.0	190.8	534.1	500.1-570.0	175.8	434.2	404.6-465.6	142.8	179.2	166.1-193.2	79.8	230.9	208.5-255.2	63.0	142.8	126.9-160.3
	2011-2015	385.8	474.0	452.0-496.9	187.8	489.0	456.9-522.9	198.0	466.2	435.7-498.5	133.2	154.7	142.9-167.4	73.2	190.2	170.8-211.5	60.0	126.9	112.3-143.1
	2016-2020	400.8	470.9	448.9-493.8	203.6	495.8	464.2-529.3	197.2	457.8	426.8-490.7	136.8	150.8	139.1-163.3	69.6	169.0	151.1-188.7	67.2	136.8	121.4-153.8
Oral cavity and pharynx	1976-1980	5.2	9.1	5.8-13.4	4.0	15.9	9.6-24.8	1.2	4.1	1.4-9.2	2.4	4.1	2.1-7.3	1.4	5.5	2.2-11.6	1.0	2.9	0.9-7.1
	1981-1985	6.8	12.1	8.3-17.1	5.0	20.0	12.8-29.8	1.8	5.8	2.5-11.5	2.4	3.9	2.0-7.0	1.6	6.1	2.6-12.3	0.8	2.6	0.6-7.0
	1986-1990	6.4	10.5	7.1-14.9	3.4	12.4	7.2-20.0	3.0	8.4	4.6-14.1	1.6	2.3	1.0-4.7	1.0	3.2	1.0-7.7	0.6	1.4	0.3-4.6
	1991-1995	6.8	10.3	7.1-14.5	3.4	11.6	6.7-18.8	3.4	9.3	5.3-15.1	2.2	3.2	1.6-5.9	0.6	2.1	0.4-6.1	1.6	4.1	1.7-8.3
	1996-2000	8.8	13.2	9.6-17.8	6.0	19.9	13.4-28.5	2.8	7.2	3.9-12.4	2.4	3.4	1.8-6.1	2.0	6.3	3.0-11.8	0.4	0.7	0.1-3.4
	2001-2005	9.8	13.2	9.8-17.6	7.0	20.9	14.5-29.3	2.8	6.8	3.7-11.8	2.2	3.0	1.5-5.5	1.6	5.3	2.3-10.6	0.6	1.1	0.2-4.1
	2006-2010	10.8	14.2	10.6-18.8	7.2	19.5	13.6-27.3	3.6	9.7	5.6-15.7	3.6	4.5	2.6-7.3	2.0	5.1	2.4-9.8	1.6	3.9	1.7-8.2
	2011-2015	10.8	13.2	9.8-17.6	8.4	21.5	15.3-29.6	2.4	6.0	2.9-11.1	0.8	0.9	0.2-2.7	0.6	1.6	0.3-5.1	0.2	0.4	0.0-3.2
	2016-2020	12.6	14.0	10.6-18.4	8.0	19.3	13.6-26.9	4.6	9.6	5.9-15.4	2.8	2.9	1.6-5.2	2.0	4.4	2.1-8.8	0.8	1.5	0.4-4.8
Esophagus	1976-1980	2.8	4.4	2.4-7.4	1.6	6.0	2.5-12.1	1.2	3.2	1.2-7.4	3.0	4.8	2.6-7.9	1.8	6.4	2.9-12.4	1.2	3.2	1.2-7.4
	1981-1985	1.0	1.6	0.5-3.7	0.6	2.0	0.4-6.2	0.4	1.2	0.1-4.5	2.0	3.0	1.4-5.6	1.8	6.5	2.9-12.7	0.2	0.5	0.0-3.5
	1986-1990	2.6	3.9	2.1-6.8	2.0	7.3	3.5-13.5	0.6	1.4	0.3-4.6	2.0	3.1	1.5-5.8	1.6	6.3	2.7-12.6	0.4	1.0	0.1-4.1
	1991-1995	3.2	4.9	2.8-8.0	2.8	9.7	5.3-16.3	0.4	1.0	0.1-3.9	2.2	3.5	1.7-6.3	1.6	5.6	2.4-11.0	0.6	1.8	0.3-5.5
	1996-2000	3.6	5.2	3.1-8.2	2.6	8.4	4.4-14.5	1.0	2.2	0.7-5.7	2.2	3.1	1.6-5.7	1.8	6.1	2.8-11.8	0.4	0.7	0.1-3.4
	2001-2005	3.8	5.2	3.1-8.2	3.2	9.6	5.5-15.8	0.6	1.6	0.3-4.9	4.4	6.0	3.7-9.1	4.0	12.1	7.4-19.0	0.4	1.1	0.1-4.3
	2006-2010	7.0	8.5	5.9-12.0	6.8	17.8	12.2-25.2	0.2	0.3	0.0-3.0	4.2	5.0	3.1-7.9	3.8	9.7	5.8-15.6	0.4	0.7	0.1-3.5
	2011-2015	4.6	5.0	3.1-7.7	3.8	8.5	5.1-13.9	0.8	1.4	0.4-4.6	3.6	4.6	2.6-7.6	2.6	6.1	3.2-11.0	1.0	3.2	0.9-7.9
	2016-2020	6.2	6.8	4.5-10.1	5.2	12.4	7.9-19.0	1.0	1.7	0.6-5.1	5.0	5.5	3.4-8.5	4.2	10.2	6.1-16.3	0.8	1.3	0.4-4.6
Stomach	1976-1980	4.8	8.0	5.1-12.0	3.0	12.5	6.8-20.7	1.8	4.8	2.2-9.4	3.2	5.3	3.0-8.6	1.6	7.1	3.0-14.1	1.6	4.2	1.8-8.6
	1981-1985	5.4	9.0	5.9-13.3	3.8	15.9	9.4-25.0	1.6	3.8	1.6-7.9	2.6	4.0	2.1-7.0	2.0	7.9	3.7-14.7	0.6	1.6	0.3-5.0
	1986-1990	5.2	7.6	4.9-11.2	3.8	12.7	7.6-20.1	1.4	3.3	1.3-7.2	3.8	5.8	3.5-9.2	2.6	9.1	4.8-15.8	1.2	2.9	1.1-6.8
	1991-1995	3.6	5.6	3.3-8.9	2.2	7.7	3.8-13.9	1.4	3.6	1.4-7.8	3.0	4.5	2.5-7.4	2.0	6.8	3.2-12.6	1.0	2.3	0.7-5.8
	1996-2000	3.6	5.1	3.0-8.2	2.6	9.0	4.8-15.6	1.0	2.6	0.8-6.4	1.8	2.5	1.1-4.9	1.4	5.0	2.0-10.3	0.4	0.9	0.1-3.7
	2001-2005	6.8	9.1	6.3-12.8	5.2	15.4	10.0-22.8	1.6	3.7	1.5-7.7	4.4	5.8	3.6-9.0	3.4	10.5	6.1-17.1	1.0	2.0	0.6-5.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Otsego**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	4.0	5.0	3.0-8.0	2.4	6.8	3.5-12.2	1.6	3.9	1.6-8.3	1.0	1.3	0.4-3.2	0.6	1.9	0.4-5.8	0.4	0.8	0.1-3.6
	2011-2015	5.0	6.2	3.9-9.5	3.4	9.2	5.3-15.2	1.6	4.0	1.6-8.6	1.8	2.0	0.9-4.1	1.4	3.5	1.4-7.8	0.4	0.8	0.1-3.8
	2016-2020	2.8	3.2	1.7-5.8	1.6	4.1	1.6-8.9	1.2	2.3	0.8-5.9	0.8	1.1	0.2-3.3	0.8	2.3	0.5-6.7	0.0	0.0	0.0-3.0
Colorectal	1976-1980	42.0	68.0	58.9-78.1	19.8	75.2	60.7-92.2	22.2	62.0	50.8-75.1	18.2	29.3	23.5-36.1	9.4	36.2	26.3-48.5	8.8	23.7	17.1-32.1
	1981-1985	47.4	72.7	63.6-82.7	20.8	72.8	59.2-88.7	26.6	72.7	60.5-86.7	13.0	19.4	14.9-24.8	5.2	17.9	11.6-26.6	7.8	20.1	14.3-27.8
	1986-1990	42.8	63.9	55.6-73.3	19.2	66.7	53.7-81.8	23.6	62.0	51.1-74.8	18.0	27.7	22.2-34.2	8.4	32.0	22.8-43.5	9.6	25.1	18.4-33.7
	1991-1995	42.0	61.0	53.0-70.0	22.0	75.3	61.7-90.9	20.0	48.6	39.4-59.4	18.2	25.8	20.8-31.8	10.4	35.4	26.4-46.7	7.8	18.2	12.9-25.1
	1996-2000	39.2	55.2	47.7-63.6	18.4	59.6	47.9-73.2	20.8	50.7	41.3-61.8	17.6	24.1	19.4-29.8	7.2	23.4	16.4-32.5	10.4	23.1	17.2-30.7
	2001-2005	39.0	52.9	45.7-61.0	20.2	64.0	52.0-78.0	18.8	45.0	36.1-55.5	13.4	17.5	13.5-22.4	7.4	23.5	16.5-32.6	6.0	12.1	8.1-17.7
	2006-2010	33.4	42.7	36.3-49.9	16.4	47.0	37.2-58.8	17.0	37.8	29.9-47.4	13.6	17.1	13.2-21.9	6.6	19.1	13.1-27.1	7.0	14.8	10.1-21.1
	2011-2015	34.0	41.2	35.0-48.3	18.0	47.3	37.7-58.8	16.0	35.5	27.6-45.2	10.2	11.5	8.5-15.4	5.2	13.7	8.9-20.5	5.0	9.1	5.8-14.1
2016-2020	34.4	42.8	36.2-50.4	19.8	53.7	43.0-66.4	14.6	32.7	25.1-42.3	11.8	13.4	10.1-17.7	5.6	13.2	8.7-19.7	6.2	13.4	8.7-20.1	
Colon excluding rectum	1976-1980	29.4	48.2	40.6-56.9	14.0	54.6	42.2-69.5	15.4	43.1	33.8-54.3	13.6	22.0	17.0-27.9	7.4	28.3	19.7-39.3	6.2	16.5	11.2-23.6
	1981-1985	33.8	52.4	44.7-61.1	14.2	51.1	39.7-64.9	19.6	53.0	42.8-65.1	9.4	14.2	10.4-19.0	3.8	13.4	8.0-21.3	5.6	14.6	9.6-21.4
	1986-1990	30.0	44.6	37.6-52.4	12.0	42.2	32.0-54.7	18.0	47.5	37.9-58.8	15.0	23.0	18.0-28.9	6.6	25.4	17.3-36.0	8.4	22.0	15.6-30.1
	1991-1995	29.8	43.0	36.3-50.6	14.2	48.7	37.9-61.6	15.6	38.3	30.1-48.1	14.2	20.0	15.6-25.3	8.0	27.2	19.3-37.2	6.2	14.5	9.8-20.8
	1996-2000	27.4	38.2	32.0-45.2	12.6	40.1	30.7-51.4	14.8	35.5	27.8-44.9	14.8	20.2	15.8-25.4	5.8	18.9	12.6-27.2	9.0	20.2	14.7-27.4
	2001-2005	29.0	39.2	33.0-46.3	15.0	47.9	37.6-60.2	14.0	33.4	25.8-42.7	10.2	13.3	9.9-17.6	5.4	17.1	11.2-25.1	4.8	9.8	6.2-15.0
	2006-2010	24.4	30.8	25.5-37.0	10.6	29.9	22.3-39.5	13.8	30.9	23.8-39.8	10.4	13.1	9.8-17.4	4.4	12.9	8.0-19.7	6.0	12.8	8.5-19.0
	2011-2015	24.6	30.1	24.7-36.3	12.4	33.5	25.5-43.5	12.2	27.0	20.2-35.7	8.8	10.0	7.2-13.7	4.4	12.0	7.5-18.6	4.4	7.9	4.9-12.7
2016-2020	24.8	30.3	24.9-36.7	13.2	34.9	26.5-45.3	11.6	25.8	19.1-34.3	8.2	9.3	6.6-13.1	3.6	8.5	5.0-14.0	4.6	9.8	5.9-15.6	
Rectum & rectosigmoid	1976-1980	12.6	19.8	15.1-25.4	5.8	20.6	13.6-29.9	6.8	18.9	13.0-26.7	4.6	7.4	4.6-11.2	2.0	7.9	3.6-14.9	2.6	7.2	3.7-12.8
	1981-1985	13.6	20.3	15.7-25.9	6.6	21.7	14.8-30.8	7.0	19.7	13.5-27.8	3.6	5.2	3.1-8.4	1.4	4.5	1.8-9.6	2.2	5.5	2.8-10.3
	1986-1990	12.8	19.4	14.8-24.9	7.2	24.5	17.0-34.1	5.6	14.6	9.6-21.4	3.0	4.7	2.6-7.9	1.8	6.5	2.9-12.5	1.2	3.2	1.2-7.2
	1991-1995	12.2	18.0	13.7-23.2	7.8	26.6	18.9-36.5	4.4	10.3	6.4-15.9	4.0	5.9	3.6-9.1	2.4	8.2	4.2-14.5	1.6	3.7	1.6-7.6
	1996-2000	11.8	17.0	12.9-22.1	5.8	19.5	13.0-28.1	6.0	15.2	10.2-22.0	2.8	4.0	2.2-6.7	1.4	4.5	1.8-9.5	1.4	2.9	1.1-6.5
	2001-2005	10.0	13.7	10.1-18.1	5.2	16.2	10.5-23.9	4.8	11.6	7.4-17.6	3.2	4.2	2.4-7.0	2.0	6.4	3.0-11.9	1.2	2.3	0.8-5.7
	2006-2010	9.0	11.9	8.6-16.1	5.8	17.1	11.3-25.0	3.2	6.9	3.8-11.8	3.2	4.0	2.3-6.7	2.2	6.3	3.1-11.5	1.0	1.9	0.6-5.2
	2011-2015	9.4	11.1	8.1-15.2	5.6	13.8	8.9-20.6	3.8	8.5	4.9-13.9	1.4	1.5	0.6-3.4	0.8	1.7	0.4-5.1	0.6	1.2	0.2-4.4
2016-2020	9.6	12.5	9.0-17.1	6.6	18.8	12.6-27.2	3.0	6.9	3.6-12.4	3.6	4.1	2.3-6.8	2.0	4.7	2.2-9.3	1.6	3.6	1.4-8.2	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Otsego**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	1.0	1.6	0.5-3.8	0.6	1.8	0.4-5.8	0.4	1.2	0.1-4.6	0.6	1.0	0.2-3.0	0.2	0.6	0.0-3.9	0.4	1.2	0.1-4.6
	1981-1985	1.6	2.4	1.0-4.8	1.0	3.7	1.1-8.9	0.6	1.6	0.3-5.0	1.2	1.9	0.7-4.3	0.6	2.3	0.4-7.0	0.6	1.9	0.4-5.8
	1986-1990	1.4	2.2	0.9-4.7	0.8	3.0	0.8-7.7	0.6	1.4	0.3-4.7	1.2	2.1	0.7-4.7	0.8	3.3	0.9-8.3	0.4	0.9	0.1-3.9
	1991-1995	2.2	3.4	1.7-6.1	1.8	6.7	3.0-12.7	0.4	1.0	0.1-3.9	2.8	4.1	2.2-7.0	1.8	5.9	2.7-11.3	1.0	2.4	0.8-5.9
	1996-2000	3.2	4.6	2.6-7.6	1.6	5.1	2.2-10.1	1.6	4.0	1.7-8.3	3.2	4.6	2.6-7.5	1.6	5.5	2.3-10.9	1.6	3.9	1.7-8.1
	2001-2005	2.6	3.8	2.0-6.7	2.0	6.3	3.0-11.9	0.6	1.3	0.2-4.4	2.4	3.4	1.7-6.1	1.6	4.9	2.1-9.9	0.8	1.6	0.4-4.8
	2006-2010	3.0	3.6	2.0-6.2	2.8	7.1	3.8-12.2	0.2	0.4	0.0-3.2	1.2	1.7	0.6-3.8	1.2	3.4	1.3-7.8	0.0	0.0	0.0-2.6
	2011-2015	6.4	7.5	5.1-10.8	4.8	11.5	7.3-17.6	1.6	3.7	1.6-7.9	4.0	4.5	2.7-7.2	2.8	6.3	3.4-11.2	1.2	2.7	1.0-6.6
	2016-2020	4.4	5.1	3.1-8.1	2.4	5.5	2.8-10.2	2.0	4.8	2.2-9.7	3.2	3.9	2.1-6.7	1.8	4.6	2.1-9.3	1.4	3.5	1.2-8.0
Pancreas	1976-1980	10.4	17.1	12.7-22.5	5.4	20.6	13.4-30.3	5.0	14.1	9.1-21.2	8.6	14.0	10.1-19.0	4.6	17.5	11.0-26.6	4.0	10.5	6.4-16.5
	1981-1985	6.6	10.8	7.4-15.2	3.6	14.2	8.2-22.7	3.0	8.1	4.5-13.8	6.2	10.0	6.7-14.3	3.2	12.7	7.1-20.9	3.0	8.2	4.5-13.9
	1986-1990	7.6	11.9	8.3-16.4	3.6	13.0	7.6-20.8	4.0	9.6	5.9-15.1	6.6	10.5	7.1-14.8	3.6	13.2	7.7-21.1	3.0	7.2	4.0-12.2
	1991-1995	8.2	11.7	8.4-15.9	4.0	13.4	8.1-20.8	4.2	9.7	5.9-15.1	6.2	9.1	6.1-13.0	2.8	9.2	5.0-15.6	3.4	7.8	4.5-12.9
	1996-2000	8.2	11.4	8.2-15.5	3.6	11.4	6.7-18.2	4.6	11.4	7.2-17.4	7.4	10.4	7.3-14.4	3.6	11.6	6.8-18.4	3.8	9.3	5.5-14.9
	2001-2005	10.2	14.0	10.4-18.5	5.8	17.6	11.8-25.5	4.4	11.5	7.1-17.7	9.0	12.2	8.9-16.4	5.6	17.2	11.4-25.0	3.4	8.4	4.8-13.9
	2006-2010	9.6	12.1	8.9-16.2	5.4	15.5	10.2-22.9	4.2	8.8	5.4-14.0	9.2	11.5	8.4-15.5	5.0	14.0	9.0-20.9	4.2	9.3	5.7-14.6
	2011-2015	10.8	12.0	9.0-15.9	5.2	12.4	8.0-18.6	5.6	11.3	7.5-17.0	9.0	9.9	7.2-13.5	5.0	12.4	8.0-18.8	4.0	7.9	4.8-12.8
	2016-2020	12.2	12.9	9.8-16.9	6.0	14.2	9.5-20.8	6.2	12.0	8.0-17.7	10.8	11.2	8.3-14.9	4.8	11.3	7.1-17.5	6.0	11.0	7.4-16.5
Larynx	1976-1980	3.4	5.6	3.2-9.0	2.4	8.2	4.2-14.6	1.0	3.3	1.0-8.0	0.2	0.5	0.0-2.6	0.2	1.1	0.0-5.4	0.0	0.0	0.0-2.7
	1981-1985	2.2	3.5	1.7-6.3	2.0	7.1	3.4-13.4	0.2	0.6	0.0-3.7	0.6	1.1	0.2-3.4	0.4	1.3	0.2-5.1	0.2	1.1	0.0-5.2
	1986-1990	3.2	4.9	2.8-8.1	2.4	8.6	4.4-15.1	0.8	2.0	0.5-5.4	0.6	1.0	0.2-3.1	0.4	1.6	0.2-5.7	0.2	0.5	0.0-3.2
	1991-1995	3.8	5.7	3.4-9.0	2.6	8.8	4.7-15.2	1.2	3.0	1.1-6.9	0.6	0.9	0.2-2.7	0.6	2.2	0.5-6.4	0.0	0.0	0.0-2.4
	1996-2000	4.2	6.2	3.8-9.6	4.2	13.4	8.3-20.6	0.0	0.0	0.0-2.4	1.0	1.4	0.4-3.3	1.0	3.4	1.1-8.1	0.0	0.0	0.0-2.4
	2001-2005	3.0	4.2	2.3-7.0	1.8	5.6	2.6-10.9	1.2	3.1	1.1-7.1	0.4	0.5	0.1-2.1	0.4	1.0	0.1-4.2	0.0	0.0	0.0-2.5
	2006-2010	2.4	3.3	1.7-5.9	2.2	6.5	3.2-11.9	0.2	0.3	0.0-3.0	1.0	1.2	0.4-3.0	1.0	2.7	0.8-6.7	0.0	0.0	0.0-2.6
	2011-2015	3.8	4.1	2.5-6.7	3.2	7.7	4.4-13.0	0.6	1.2	0.3-4.5	1.0	1.1	0.3-2.8	0.8	2.1	0.6-5.9	0.2	0.4	0.0-3.4
	2016-2020	2.4	2.3	1.2-4.4	2.0	4.2	2.0-8.4	0.4	0.8	0.1-4.0	0.8	0.7	0.2-2.4	0.6	1.4	0.3-5.0	0.2	0.3	0.0-3.3
Lung and bronchus	1976-1980	32.0	53.9	45.6-63.3	24.2	91.8	75.6-110.3	7.8	25.5	17.8-35.3	21.8	36.6	29.9-44.5	16.2	62.9	49.5-78.7	5.6	16.9	11.0-24.9
	1981-1985	38.8	62.1	53.4-71.8	28.8	106.9	89.7-126.5	10.0	29.0	21.2-38.8	30.4	48.1	40.6-56.7	22.0	81.3	66.4-98.6	8.4	23.4	16.6-32.1
	1986-1990	44.6	68.7	59.8-78.5	25.6	89.4	74.3-106.7	19.0	54.5	43.7-67.2	31.4	47.6	40.4-55.9	20.4	72.8	59.2-88.7	11.0	30.8	22.9-40.5
	1991-1995	45.8	68.3	59.7-77.9	26.2	87.7	73.2-104.3	19.6	54.1	43.7-66.4	40.0	58.9	50.9-67.7	24.4	82.7	68.5-99.0	15.6	41.6	32.7-52.2
	1996-2000	49.6	69.6	61.2-78.9	31.2	100.9	85.6-118.3	18.4	47.1	37.9-58.0	37.4	52.1	44.9-60.2	24.8	81.2	67.4-97.1	12.6	32.0	24.5-41.2
	2001-2005	53.4	72.7	64.2-82.1	31.6	96.5	81.9-113.1	21.8	55.6	45.5-67.5	44.4	59.8	52.2-68.4	24.6	77.2	64.0-92.4	19.8	49.4	40.0-60.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Otsego**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	51.4	65.2	57.4-73.9	29.8	84.5	71.3-99.5	21.6	50.7	41.4-61.7	37.2	47.1	40.5-54.6	20.8	58.6	47.8-71.3	16.4	38.8	30.7-48.7
	2011-2015	52.8	61.6	54.2-69.9	28.4	70.8	59.4-84.0	24.4	54.4	44.8-65.7	37.8	44.1	37.8-51.2	22.4	56.5	46.3-68.5	15.4	34.3	26.7-43.6
	2016-2020	58.2	62.0	54.8-70.0	28.8	66.4	55.8-78.9	29.4	58.8	49.3-70.0	31.0	33.3	28.1-39.3	17.4	40.2	32.0-50.1	13.6	27.2	20.9-35.2
Melanoma of the skin	1976-1980	6.2	11.0	7.3-15.9	3.2	12.7	7.0-21.0	3.0	10.3	5.6-17.2	2.2	4.0	2.0-7.3	1.4	5.7	2.2-11.8	0.8	2.9	0.7-7.5
	1981-1985	7.2	12.2	8.5-17.1	4.4	17.1	10.5-26.1	2.8	9.2	4.9-15.6	1.2	1.8	0.7-4.0	0.6	2.9	0.6-8.3	0.6	1.5	0.3-4.9
	1986-1990	9.8	17.9	13.2-23.7	4.0	15.6	9.5-24.2	5.8	20.0	13.2-28.9	1.4	2.3	0.9-4.8	0.8	2.7	0.7-7.1	0.6	2.0	0.4-6.0
	1991-1995	10.2	17.0	12.6-22.5	5.6	20.4	13.5-29.5	4.6	14.0	8.8-21.2	2.4	3.9	2.0-6.9	0.8	3.0	0.8-7.5	1.6	4.7	2.0-9.5
	1996-2000	10.8	16.4	12.3-21.5	6.6	21.7	14.9-30.6	4.2	11.9	7.3-18.4	1.4	2.1	0.8-4.3	0.8	2.6	0.7-6.7	0.6	1.4	0.3-4.5
	2001-2005	12.0	17.7	13.4-22.9	6.4	20.3	13.8-28.8	5.6	16.3	10.7-23.9	2.2	2.9	1.5-5.4	1.0	3.2	1.0-7.6	1.2	2.9	1.1-6.8
	2006-2010	13.4	18.2	14.0-23.4	7.4	20.8	14.6-29.1	6.0	15.9	10.5-23.2	1.6	2.1	0.9-4.3	1.2	3.3	1.2-7.6	0.4	0.9	0.1-4.0
	2011-2015	20.4	26.6	21.4-32.8	13.4	37.0	28.2-47.8	7.0	18.0	12.1-25.9	1.6	1.8	0.8-3.8	1.2	3.0	1.1-7.0	0.4	0.8	0.1-3.8
2016-2020	22.0	27.6	22.3-34.0	14.0	35.0	27.0-45.0	8.0	22.6	15.3-32.2	2.4	2.5	1.2-4.8	1.2	2.9	1.0-7.2	1.2	2.1	0.7-5.6	
Female breast	1976-1980							36.0	114.7	97.8-133.7							10.2	31.9	23.4-42.4
	1981-1985							36.6	114.4	97.6-133.1							11.0	34.4	25.5-45.5
	1986-1990							38.8	114.4	98.2-132.6							11.6	32.1	24.1-41.9
	1991-1995							37.2	103.3	88.6-119.8							10.4	29.2	21.6-38.7
	1996-2000							52.2	148.0	130.2-167.5							11.6	29.1	22.0-38.1
	2001-2005							49.6	130.6	114.6-148.4							8.4	21.0	15.0-28.7
	2006-2010							47.0	119.5	104.0-136.8							8.2	19.8	14.0-27.4
	2011-2015							58.4	141.1	124.5-159.6							9.4	19.8	14.3-27.0
2016-2020							54.6	136.4	119.0-155.7							6.8	15.2	10.1-22.3	
Cervix uteri	1976-1980							2.8	9.4	4.9-16.1							1.2	3.7	1.3-8.3
	1981-1985							3.8	11.9	7.0-18.9							1.8	4.9	2.2-9.6
	1986-1990							2.6	8.2	4.2-14.4							1.0	2.6	0.8-6.4
	1991-1995							4.8	15.7	10.0-23.4							0.6	1.5	0.3-4.7
	1996-2000							2.8	9.0	4.8-15.2							0.2	0.7	0.0-3.8
	2001-2005							2.0	6.5	3.0-12.1							1.2	3.2	1.2-7.4
	2006-2010							3.4	10.1	5.7-16.5							0.8	2.1	0.6-5.8
	2011-2015							2.2	7.3	3.5-13.6							0.8	1.9	0.5-5.6
	2016-2020							2.0	6.2	2.6-12.5							0.8	2.6	0.5-7.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Otsego**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							7.2	22.1	15.2-31.1							1.6	4.4	1.9-9.1
	1981-1985							8.6	26.8	19.1-36.6							1.4	3.3	1.3-7.2
	1986-1990							9.0	27.6	19.8-37.4							1.2	3.9	1.4-8.8
	1991-1995							10.0	30.0	22.1-39.9							1.0	2.4	0.8-5.8
	1996-2000							11.4	31.4	23.7-40.9							0.6	1.4	0.3-4.5
	2001-2005							12.6	33.5	25.6-43.1							2.0	4.9	2.3-9.4
	2006-2010							16.2	40.7	32.1-51.2							1.2	2.6	0.9-6.3
	2011-2015							16.2	37.6	29.4-47.6							2.8	5.9	3.2-10.6
	2016-2020							16.6	39.3	30.4-50.2							2.6	4.6	2.4-8.7
Ovary	1976-1980							6.2	19.6	13.0-28.3							3.4	10.1	5.7-16.7
	1981-1985							6.0	17.7	11.7-25.8							4.6	12.9	8.1-19.8
	1986-1990							5.8	16.9	11.1-24.6							2.4	6.8	3.3-12.2
	1991-1995							6.6	19.2	13.1-27.3							3.4	8.6	5.0-14.1
	1996-2000							3.6	9.6	5.6-15.4							3.6	10.2	6.0-16.3
	2001-2005							4.8	12.8	8.1-19.4							2.8	6.5	3.5-11.2
	2006-2010							5.0	12.0	7.6-18.3							2.8	5.8	3.1-10.2
	2011-2015							6.0	14.2	9.3-21.0							4.2	9.1	5.5-14.6
	2016-2020							4.0	10.2	5.7-16.8							3.6	7.3	4.2-12.3
Prostate	1976-1980				27.2	111.3	92.8-132.3							6.8	28.6	19.6-40.2			
	1981-1985				36.0	135.1	115.6-157.0							9.6	38.8	28.3-51.7			
	1986-1990				40.4	143.3	123.8-164.9							10.6	41.0	30.5-53.9			
	1991-1995				53.8	179.1	158.0-202.3							10.6	42.0	31.2-55.1			
	1996-2000				44.0	140.8	122.7-160.9							10.8	37.0	27.7-48.4			
	2001-2005				46.4	139.0	121.5-158.3							9.6	31.8	23.4-42.3			
	2006-2010				47.0	123.9	108.3-141.2							9.6	29.3	21.6-39.1			
	2011-2015				34.8	84.1	71.7-98.2							7.6	20.1	14.2-28.0			
	2016-2020				41.2	88.6	76.7-102.2							8.2	19.8	14.2-27.3			
Testis	1976-1980				1.2	5.7	2.0-12.3							0.0	0.0	0.0-3.0			
	1981-1985				1.2	5.0	1.7-10.9							0.0	0.0	0.0-3.0			
	1986-1990				1.6	5.9	2.5-11.8							0.0	0.0	0.0-2.8			
	1991-1995				1.8	6.7	3.0-12.6							0.2	0.5	0.0-3.3			
	1996-2000				1.2	3.7	1.3-8.2							0.0	0.0	0.0-2.6			
	2001-2005				3.2	12.2	6.7-20.0							0.2	0.6	0.0-3.6			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Otsego**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				1.6	5.5	2.3-11.2							0.0	0.0	0.0-2.7			
	2011-2015				1.6	7.0	2.9-13.9							0.0	0.0	0.0-2.8			
	2016-2020				1.6	7.5	3.2-14.7							0.0	0.0	0.0-2.9			
Urinary bladder (incl. in situ)	1976-1980	13.2	21.6	16.6-27.6	10.2	38.6	28.6-51.1	3.0	8.4	4.6-14.3	3.0	5.3	2.9-8.8	2.2	9.2	4.5-16.5	0.8	2.0	0.5-5.6
	1981-1985	13.6	21.4	16.5-27.3	9.4	35.1	25.5-47.1	4.2	11.9	7.2-18.6	3.8	5.7	3.4-8.9	3.0	11.8	6.5-19.7	0.8	2.0	0.5-5.5
	1986-1990	19.0	28.8	23.2-35.3	12.8	47.1	36.0-60.4	6.2	17.1	11.4-24.7	3.8	5.8	3.5-9.1	2.4	9.8	4.9-17.3	1.4	3.8	1.5-8.2
	1991-1995	16.6	23.9	19.0-29.8	10.4	36.2	26.8-47.7	6.2	15.3	10.4-22.0	4.2	6.0	3.7-9.3	2.6	9.7	5.1-16.6	1.6	3.7	1.6-7.6
	1996-2000	20.8	29.4	24.0-35.7	15.0	49.6	38.9-62.4	5.8	14.2	9.5-20.7	2.6	3.6	1.9-6.3	1.8	6.5	2.9-12.4	0.8	1.7	0.5-4.9
	2001-2005	20.4	27.6	22.5-33.6	15.6	48.2	38.0-60.3	4.8	12.3	7.8-18.6	5.2	6.6	4.3-9.9	3.2	10.8	6.2-17.7	2.0	4.1	2.0-8.1
	2006-2010	19.6	24.0	19.4-29.4	15.2	41.6	32.6-52.4	4.4	9.9	6.2-15.4	6.0	7.4	5.0-10.7	4.8	14.5	9.3-21.8	1.2	2.6	0.9-6.3
	2011-2015	19.4	21.6	17.4-26.5	13.8	34.9	27.1-44.6	5.6	11.6	7.6-17.3	4.4	4.7	3.0-7.4	2.4	6.8	3.5-12.2	2.0	3.9	1.9-8.0
2016-2020	23.8	25.3	20.8-30.6	17.4	40.4	32.2-50.4	6.4	13.4	9.0-19.8	4.8	4.8	3.0-7.4	3.4	8.0	4.6-13.3	1.4	2.3	0.9-5.8	
Kidney and renal pelvis	1976-1980	2.8	4.8	2.6-8.3	1.8	7.2	3.2-13.9	1.0	2.8	0.9-6.9	2.2	3.7	1.8-6.8	1.4	5.4	2.1-11.3	0.8	2.4	0.6-6.4
	1981-1985	5.0	7.8	5.0-11.6	2.8	10.7	5.7-18.3	2.2	5.8	2.9-10.7	2.2	3.5	1.7-6.4	1.6	5.6	2.3-11.4	0.6	1.6	0.3-5.0
	1986-1990	4.8	7.7	4.9-11.5	2.8	10.8	5.8-18.2	2.0	5.7	2.7-10.7	2.4	3.8	1.9-6.8	1.8	7.0	3.1-13.4	0.6	1.5	0.3-4.9
	1991-1995	7.2	11.0	7.7-15.3	4.4	14.6	9.1-22.3	2.8	8.4	4.5-14.3	3.2	4.7	2.6-7.6	2.2	7.7	3.8-13.9	1.0	2.6	0.8-6.4
	1996-2000	8.0	11.5	8.2-15.7	4.2	13.9	8.6-21.4	3.8	9.8	5.8-15.6	2.2	3.2	1.6-5.8	0.8	2.8	0.8-7.3	1.4	3.9	1.5-8.2
	2001-2005	7.6	10.4	7.3-14.3	4.6	13.3	8.4-20.1	3.0	7.5	4.1-12.8	4.0	5.3	3.3-8.4	2.6	7.7	4.1-13.4	1.4	3.0	1.2-6.8
	2006-2010	11.8	15.0	11.4-19.6	8.4	23.9	17.1-32.7	3.4	7.6	4.4-12.6	3.0	3.8	2.1-6.4	2.2	6.1	3.0-11.3	0.8	1.6	0.4-4.8
	2011-2015	14.8	17.1	13.3-21.8	8.4	21.6	15.3-29.8	6.4	13.0	8.8-18.9	4.4	4.9	3.1-7.7	3.2	8.0	4.5-13.4	1.2	2.4	0.9-6.1
2016-2020	14.8	17.3	13.4-22.2	10.6	26.1	19.2-34.8	4.2	10.0	5.9-16.1	3.2	3.4	1.9-5.9	2.2	5.3	2.6-10.0	1.0	2.0	0.7-5.6	
Brain and other central nervous system	1976-1980	4.2	7.4	4.5-11.5	2.6	10.3	5.3-17.9	1.6	5.1	2.1-10.4	3.6	6.6	3.8-10.6	1.8	8.0	3.6-15.3	1.8	5.5	2.4-10.8
	1981-1985	4.2	7.5	4.6-11.6	3.0	11.5	6.3-19.1	1.2	4.5	1.6-9.8	2.0	3.6	1.7-6.8	1.8	7.0	3.1-13.6	0.2	1.1	0.0-5.2
	1986-1990	5.8	9.4	6.2-13.7	2.8	10.1	5.4-17.3	3.0	9.1	4.9-15.3	4.0	6.5	3.9-10.1	2.2	7.9	3.9-14.2	1.8	5.6	2.5-10.9
	1991-1995	3.4	5.6	3.3-9.1	1.6	5.7	2.5-11.3	1.8	5.7	2.5-10.9	2.8	4.5	2.4-7.6	1.4	4.6	1.8-9.6	1.4	4.6	1.8-9.6
	1996-2000	4.0	6.2	3.8-9.6	1.2	4.1	1.5-8.9	2.8	7.7	4.1-13.3	3.2	4.6	2.6-7.5	1.0	3.2	1.0-7.5	2.2	5.6	2.8-10.4
	2001-2005	5.0	7.8	5.0-11.7	2.8	9.1	4.9-15.5	2.2	7.1	3.4-13.0	2.8	3.7	2.0-6.3	1.2	3.7	1.4-8.3	1.6	3.8	1.6-7.9
	2006-2010	4.8	7.2	4.5-11.0	2.8	9.6	5.1-16.4	2.0	5.3	2.3-10.4	3.8	5.1	3.0-8.2	2.4	7.3	3.7-13.1	1.4	3.5	1.2-7.9
	2011-2015	6.8	10.9	7.2-15.7	3.6	11.9	6.6-19.6	3.2	9.6	5.0-16.5	3.2	3.8	2.1-6.4	1.8	4.3	2.0-8.8	1.4	3.2	1.2-7.3
2016-2020	6.8	8.9	5.9-13.0	3.6	10.4	5.8-17.2	3.2	7.8	4.1-13.7	4.4	5.5	3.3-8.8	2.0	5.6	2.5-10.9	2.4	5.6	2.6-10.7	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Otsego**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	0.8	1.7	0.4-4.3	0.2	0.7	0.0-4.2	0.6	2.7	0.5-7.6	0.0	0.0	0.0-1.4	0.0	0.0	0.0-3.0	0.0	0.0	0.0-2.7
	1981-1985	2.2	4.0	1.9-7.2	0.4	1.8	0.2-6.2	1.8	5.9	2.6-11.4	0.0	0.0	0.0-1.4	0.0	0.0	0.0-3.0	0.0	0.0	0.0-2.6
	1986-1990	3.8	6.0	3.6-9.6	1.6	5.2	2.2-10.5	2.2	6.8	3.3-12.5	0.2	0.3	0.0-1.8	0.0	0.0	0.0-2.8	0.2	0.5	0.0-3.3
	1991-1995	3.0	5.0	2.8-8.2	1.0	3.6	1.1-8.3	2.0	6.2	2.9-11.5	0.2	0.3	0.0-1.7	0.0	0.0	0.0-2.6	0.2	0.5	0.0-3.2
	1996-2000	4.0	6.7	4.1-10.4	1.0	3.1	1.0-7.3	3.0	9.9	5.5-16.5	0.4	0.6	0.1-2.1	0.2	0.6	0.0-3.8	0.2	0.5	0.0-3.2
	2001-2005	6.2	10.2	6.8-14.6	1.4	4.2	1.6-8.9	4.8	15.9	10.0-23.9	0.4	0.5	0.1-2.1	0.0	0.0	0.0-2.7	0.4	0.8	0.1-3.7
	2006-2010	8.4	14.1	10.0-19.3	2.0	6.1	2.8-11.5	6.4	22.0	14.7-31.6	0.4	0.5	0.1-2.0	0.0	0.0	0.0-2.7	0.4	0.9	0.1-3.9
	2011-2015	10.2	16.6	12.0-22.3	3.0	9.8	5.1-17.0	7.2	23.3	15.8-33.0	0.2	0.4	0.0-2.1	0.0	0.0	0.0-2.8	0.2	0.7	0.0-4.1
	2016-2020	8.8	15.2	10.6-21.0	3.4	11.9	6.5-19.9	5.4	18.5	11.6-27.9	0.2	0.4	0.0-2.3	0.0	0.0	0.0-2.9	0.2	0.9	0.0-4.6
Hodgkin lymphoma	1976-1980	2.4	3.3	1.6-6.1	1.6	4.1	1.7-8.8	0.8	2.6	0.6-7.0	1.0	1.5	0.5-3.7	0.4	1.3	0.1-5.2	0.6	1.6	0.3-5.2
	1981-1985	3.0	4.3	2.3-7.4	1.4	4.2	1.6-9.1	1.6	4.6	1.8-9.6	0.6	0.8	0.2-2.6	0.4	1.1	0.1-4.7	0.2	0.5	0.0-3.4
	1986-1990	1.6	2.3	0.9-4.7	0.4	1.7	0.2-5.9	1.2	2.6	1.0-6.2	0.6	0.9	0.2-2.8	0.2	0.7	0.0-4.1	0.4	1.0	0.1-4.0
	1991-1995	2.2	3.5	1.7-6.3	0.8	2.6	0.7-6.9	1.4	4.0	1.6-8.5	0.2	0.3	0.0-1.7	0.0	0.0	0.0-2.6	0.2	0.5	0.0-3.2
	1996-2000	2.6	4.1	2.2-7.1	2.0	6.9	3.3-12.7	0.6	1.4	0.3-4.6	0.4	0.6	0.1-2.2	0.2	0.7	0.0-3.9	0.2	0.4	0.0-2.9
	2001-2005	1.6	2.3	1.0-4.8	1.2	3.3	1.2-7.6	0.4	1.4	0.1-5.2	0.2	0.2	0.0-1.6	0.0	0.0	0.0-2.7	0.2	0.3	0.0-2.9
	2006-2010	2.4	4.4	2.2-7.8	1.2	4.4	1.5-9.8	1.2	4.4	1.5-9.7	0.2	0.2	0.0-1.6	0.2	0.4	0.0-3.3	0.0	0.0	0.0-2.6
	2011-2015	2.2	4.4	2.1-7.9	1.6	6.0	2.4-12.3	0.6	2.8	0.6-7.8	0.6	0.9	0.2-2.9	0.4	1.3	0.1-5.1	0.2	0.5	0.0-3.6
	2016-2020	1.2	1.9	0.6-4.5	0.6	1.8	0.3-6.1	0.6	1.9	0.3-6.3	0.0	0.0	0.0-1.5	0.0	0.0	0.0-2.9	0.0	0.0	0.0-3.0
Non-Hodgkin lymphomas	1976-1980	9.2	15.8	11.5-21.3	4.0	16.2	9.7-25.4	5.2	15.6	10.0-23.3	3.6	5.8	3.4-9.3	1.6	5.7	2.4-11.5	2.0	5.8	2.7-11.0
	1981-1985	9.4	16.0	11.6-21.5	6.0	24.6	16.3-35.4	3.4	9.7	5.5-15.9	6.0	9.8	6.5-14.1	4.2	17.0	10.3-26.3	1.8	5.4	2.3-10.6
	1986-1990	10.0	15.8	11.7-21.0	4.2	15.0	9.2-23.2	5.8	16.2	10.7-23.6	4.2	6.4	3.9-9.9	1.6	5.3	2.2-10.7	2.6	6.9	3.6-12.1
	1991-1995	13.8	21.5	16.7-27.3	5.8	21.9	14.6-31.5	8.0	21.7	15.3-29.9	6.2	9.3	6.3-13.2	2.6	9.5	5.0-16.3	3.6	9.1	5.3-14.7
	1996-2000	13.2	19.0	14.7-24.3	6.4	21.0	14.3-29.7	6.8	17.6	12.1-24.9	5.6	8.2	5.4-11.9	2.8	9.7	5.3-16.4	2.8	7.4	4.0-12.7
	2001-2005	15.0	21.7	16.9-27.3	8.2	26.2	18.7-35.7	6.8	18.2	12.4-25.9	6.0	8.2	5.5-11.8	3.0	9.7	5.4-16.1	3.0	6.9	3.8-11.9
	2006-2010	16.2	20.1	15.9-25.2	7.6	21.0	14.8-29.2	8.6	20.1	14.3-27.7	5.2	6.3	4.1-9.4	2.4	7.2	3.7-12.7	2.8	5.9	3.2-10.4
	2011-2015	17.0	21.6	17.1-27.2	9.2	26.2	18.9-35.7	7.8	17.9	12.6-25.0	4.2	5.1	3.1-8.1	3.0	8.2	4.5-14.0	1.2	2.7	0.9-6.7
	2016-2020	19.0	22.4	17.8-28.0	11.2	27.3	20.1-36.4	7.8	17.8	12.2-25.4	6.2	6.9	4.5-10.4	3.2	8.3	4.6-14.2	3.0	6.0	3.0-11.1
Myeloma	1976-1980	3.4	5.3	3.1-8.7	1.2	4.4	1.5-10.0	2.2	6.3	3.1-11.6	2.4	4.0	2.0-7.0	0.8	2.6	0.7-7.1	1.6	5.0	2.1-10.1
	1981-1985	3.2	4.7	2.7-7.8	2.0	7.3	3.4-13.6	1.2	2.9	1.1-6.8	2.0	3.0	1.4-5.6	0.8	3.3	0.9-8.5	1.2	2.9	1.1-6.8
	1986-1990	5.0	7.6	4.9-11.4	3.2	10.8	6.1-17.8	1.8	4.2	1.9-8.4	2.4	3.4	1.7-6.0	1.0	3.4	1.1-8.1	1.4	3.3	1.3-7.3
	1991-1995	4.0	6.1	3.7-9.5	2.8	10.1	5.5-17.1	1.2	3.3	1.2-7.4	1.8	2.6	1.2-5.0	1.2	4.2	1.5-9.2	0.6	1.3	0.3-4.2
	1996-2000	5.2	7.6	4.9-11.2	3.4	11.4	6.6-18.3	1.8	4.7	2.1-9.3	2.0	2.8	1.4-5.3	1.6	5.0	2.1-10.0	0.4	0.9	0.1-3.7
	2001-2005	3.4	4.4	2.6-7.2	1.6	4.6	2.0-9.3	1.8	4.3	1.9-8.5	2.2	2.9	1.4-5.3	0.8	2.6	0.7-6.9	1.4	3.0	1.2-6.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Otsego**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	4.2	5.3	3.2-8.3	3.0	7.8	4.3-13.3	1.2	3.1	1.0-7.3	3.4	4.3	2.5-7.1	2.2	6.3	3.1-11.6	1.2	3.1	1.0-7.4
	2011-2015	5.0	6.2	3.9-9.5	2.4	6.4	3.2-11.6	2.6	6.1	3.0-11.3	2.2	2.5	1.2-4.8	0.8	2.1	0.6-5.9	1.4	2.5	1.0-6.1
	2016-2020	5.6	6.7	4.3-10.1	4.0	10.3	6.0-16.7	1.6	3.2	1.4-7.1	2.6	2.9	1.5-5.3	1.2	3.1	1.1-7.5	1.4	2.6	1.0-6.2
Leukemias	1976-1980	5.6	8.6	5.7-12.5	3.6	13.9	8.0-22.4	2.0	5.4	2.6-10.3	5.4	8.7	5.7-12.7	2.4	10.2	5.1-18.1	3.0	8.3	4.6-14.1
	1981-1985	10.8	17.1	12.7-22.5	6.4	23.4	15.8-33.4	4.4	11.9	7.4-18.4	5.6	9.0	5.9-13.2	2.6	9.6	5.1-16.7	3.0	8.8	4.8-15.0
	1986-1990	12.6	19.3	14.8-24.8	6.2	22.4	15.1-32.0	6.4	16.9	11.4-24.2	5.8	8.5	5.7-12.3	2.4	8.8	4.5-15.4	3.4	8.0	4.6-13.1
	1991-1995	13.4	20.1	15.5-25.6	8.0	27.3	19.4-37.3	5.4	14.6	9.5-21.6	5.8	8.5	5.7-12.3	3.2	10.6	6.1-17.4	2.6	6.8	3.6-12.0
	1996-2000	10.6	15.6	11.7-20.5	6.8	22.1	15.3-31.0	3.8	9.4	5.6-15.0	4.2	5.9	3.6-9.1	2.4	8.0	4.1-14.1	1.8	4.7	2.1-9.3
	2001-2005	13.2	18.3	14.1-23.5	8.8	27.9	20.1-37.7	4.4	10.1	6.2-15.9	6.2	8.5	5.8-12.3	4.0	13.2	8.0-20.5	2.2	4.7	2.3-9.0
	2006-2010	19.0	25.3	20.3-31.2	10.2	30.6	22.6-40.7	8.8	20.4	14.5-28.0	7.2	9.5	6.6-13.4	4.6	14.1	8.8-21.4	2.6	5.7	2.9-10.4
	2011-2015	14.0	17.1	13.1-22.0	7.8	22.2	15.5-31.0	6.2	12.0	8.1-17.6	5.4	7.1	4.5-10.6	4.0	12.2	7.3-19.2	1.4	2.8	1.1-6.5
	2016-2020	14.0	15.6	11.9-20.1	8.0	20.0	14.0-28.1	6.0	11.6	7.7-17.4	4.4	5.6	3.4-9.0	2.0	6.6	3.0-12.7	2.4	4.5	2.3-8.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Putnam**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	227.8	391.9	368.8-416.1	120.6	470.4	431.2-512.0	107.2	340.7	311.9-371.3	124.2	225.7	207.7-244.7	69.0	294.0	261.6-329.1	55.2	178.3	157.5-201.0
	1981-1985	286.8	462.3	438.1-487.4	144.2	554.3	512.0-599.0	142.6	413.4	383.3-445.2	133.2	222.8	205.8-240.8	68.8	272.0	242.1-304.4	64.4	191.8	171.2-214.1
	1986-1990	270.6	408.7	386.8-431.4	130.8	464.2	427.2-503.4	139.8	381.8	353.9-411.4	142.2	224.1	207.6-241.4	78.0	286.7	257.2-318.4	64.2	180.4	161.1-201.3
	1991-1995	316.0	437.5	415.9-460.0	162.0	520.6	483.3-559.9	154.0	384.6	357.8-412.9	152.8	224.4	208.5-241.1	79.4	286.0	256.8-317.3	73.4	188.2	169.3-208.5
	1996-2000	435.0	532.1	509.6-555.4	222.4	611.2	573.8-650.3	212.6	477.0	448.5-506.9	166.4	219.1	204.2-234.7	80.8	253.3	227.8-280.7	85.6	200.4	181.7-220.4
	2001-2005	525.6	563.6	541.7-586.1	253.8	620.0	584.3-657.2	271.8	533.8	505.3-563.4	168.4	194.8	181.6-208.7	82.8	231.1	208.1-255.8	85.6	174.7	158.3-192.3
	2006-2010	560.0	530.7	510.6-551.4	277.2	576.4	544.7-609.4	282.8	500.9	474.6-528.4	157.4	158.4	147.2-170.1	80.4	188.0	169.1-208.4	77.0	138.3	124.6-153.1
	2011-2015	626.0	518.6	500.0-537.8	318.0	558.4	529.9-588.1	308.0	494.4	469.1-520.8	165.4	142.7	132.9-153.1	85.2	164.2	148.2-181.5	80.2	126.8	114.4-140.3
	2016-2020	622.8	468.2	451.3-485.7	305.4	477.4	452.8-503.1	317.4	470.0	446.0-495.0	163.4	123.7	115.2-132.8	83.8	141.0	127.3-155.8	79.6	112.6	101.5-124.7
Oral cavity and pharynx	1976-1980	7.4	12.3	8.6-17.0	6.0	20.7	13.7-30.0	1.4	4.7	1.8-9.5	2.4	3.6	1.9-6.4	1.2	3.8	1.4-8.5	1.2	3.4	1.2-7.6
	1981-1985	8.6	13.4	9.6-18.1	4.8	16.9	10.6-25.3	3.8	10.6	6.4-16.6	2.4	3.8	2.0-6.7	1.0	3.2	1.1-7.6	1.4	4.1	1.6-8.4
	1986-1990	5.6	8.6	5.7-12.4	3.0	11.1	6.0-18.6	2.6	7.1	3.8-12.0	2.2	3.4	1.7-6.0	1.4	4.7	1.8-9.7	0.8	2.3	0.6-5.8
	1991-1995	7.6	10.1	7.1-13.9	4.6	13.2	8.1-20.1	3.0	7.5	4.2-12.4	2.4	3.1	1.6-5.5	1.4	3.9	1.5-8.2	1.0	2.5	0.8-5.8
	1996-2000	9.6	11.2	8.2-14.9	5.8	12.4	8.2-18.1	3.8	9.2	5.5-14.3	1.2	1.6	0.6-3.4	0.6	1.6	0.3-4.7	0.6	1.4	0.3-4.1
	2001-2005	8.8	8.6	6.2-11.7	6.6	13.2	8.9-18.9	2.2	4.3	2.1-7.9	3.0	3.1	1.7-5.2	1.8	3.9	1.7-7.6	1.2	2.2	0.8-5.0
	2006-2010	10.2	9.0	6.6-11.9	8.0	15.2	10.7-21.0	2.2	3.3	1.7-6.2	2.2	2.1	1.0-3.9	1.8	4.0	1.7-7.7	0.4	0.6	0.1-2.5
	2011-2015	14.4	11.2	8.7-14.2	11.6	18.3	13.7-24.1	2.8	4.5	2.5-7.8	1.6	1.2	0.5-2.6	1.2	1.7	0.6-4.1	0.4	0.6	0.1-2.6
	2016-2020	16.6	11.6	9.2-14.5	12.8	18.3	14.0-23.8	3.8	5.2	3.1-8.4	3.0	2.1	1.2-3.7	2.2	2.9	1.4-5.6	0.8	1.4	0.3-3.8
Esophagus	1976-1980	2.4	4.5	2.3-7.9	2.2	9.4	4.4-17.2	0.2	0.7	0.0-3.6	2.0	3.8	1.7-6.9	1.8	8.0	3.4-15.4	0.2	0.7	0.0-3.6
	1981-1985	3.0	5.0	2.7-8.2	1.6	5.4	2.2-10.7	1.4	4.3	1.7-8.8	2.0	3.3	1.5-6.1	1.0	2.9	0.9-6.9	1.0	3.2	1.0-7.4
	1986-1990	2.8	4.5	2.4-7.4	1.8	6.7	2.8-12.9	1.0	2.8	0.9-6.5	2.6	4.2	2.2-7.1	2.2	8.2	3.9-14.9	0.4	1.1	0.1-4.0
	1991-1995	3.2	5.0	2.8-8.1	2.0	8.2	3.6-15.2	1.2	3.2	1.2-6.9	3.6	5.5	3.2-8.6	2.2	8.4	3.8-15.3	1.4	3.7	1.5-7.6
	1996-2000	6.2	8.3	5.6-11.8	4.4	13.0	7.9-19.9	1.8	4.4	2.0-8.4	4.0	5.3	3.2-8.2	2.8	8.1	4.2-13.8	1.2	3.0	1.1-6.4
	2001-2005	5.0	5.4	3.5-8.1	4.0	9.0	5.3-14.2	1.0	2.1	0.7-5.0	4.0	4.4	2.7-6.9	2.6	6.2	3.2-10.7	1.4	2.7	1.1-5.6
	2006-2010	6.0	5.5	3.6-8.0	5.0	9.8	6.1-14.9	1.0	1.9	0.6-4.6	4.8	4.9	3.1-7.4	3.8	8.7	5.1-13.9	1.0	1.9	0.6-4.5
	2011-2015	8.0	6.1	4.3-8.5	6.4	10.5	7.0-15.2	1.6	2.3	1.0-4.8	5.4	4.3	2.8-6.5	4.2	7.3	4.4-11.5	1.2	1.8	0.7-4.3
	2016-2020	4.6	3.3	2.1-5.1	4.0	5.8	3.5-9.3	0.6	1.0	0.2-3.3	4.2	2.9	1.8-4.6	3.4	5.1	2.9-8.4	0.8	1.1	0.3-3.3
Stomach	1976-1980	4.0	8.1	4.8-12.5	2.0	9.0	4.2-16.5	2.0	7.3	3.4-13.4	3.8	8.0	4.7-12.5	2.0	10.4	4.8-19.0	1.8	6.5	2.9-12.3
	1981-1985	9.0	15.6	11.3-20.9	6.4	26.1	17.3-37.3	2.6	7.9	4.2-13.5	5.2	9.6	6.2-14.1	4.2	18.0	10.7-27.9	1.0	3.4	1.1-7.8
	1986-1990	6.8	10.8	7.4-15.1	4.2	15.9	9.4-24.8	2.6	7.2	3.8-12.3	5.6	9.0	5.9-13.0	3.8	14.4	8.2-23.0	1.8	5.1	2.3-9.7
	1991-1995	4.4	6.6	4.1-10.0	2.8	10.3	5.3-17.4	1.6	4.2	1.8-8.2	4.4	7.0	4.4-10.6	2.6	10.8	5.5-18.6	1.8	4.7	2.1-8.8
	1996-2000	6.6	8.4	5.8-11.9	4.0	10.9	6.5-17.0	2.6	6.2	3.3-10.6	5.0	6.5	4.2-9.5	3.0	7.9	4.4-13.1	2.0	4.6	2.2-8.5
	2001-2005	6.6	6.9	4.7-9.8	4.0	9.3	5.4-14.8	2.6	5.0	2.6-8.7	3.2	3.6	2.1-5.9	1.8	5.2	2.2-10.2	1.4	2.7	1.1-5.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Putnam**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	9.6	9.2	6.7-12.3	6.8	13.5	9.1-19.2	2.8	5.4	2.9-9.2	3.8	3.8	2.2-6.0	2.2	4.5	2.1-8.3	1.6	3.2	1.4-6.4
	2011-2015	9.0	7.8	5.6-10.6	5.8	10.5	6.8-15.4	3.2	5.7	3.2-9.5	3.8	3.1	1.8-5.0	2.4	4.1	2.0-7.5	1.4	2.3	0.9-5.0
	2016-2020	9.6	7.1	5.2-9.5	4.2	7.4	4.5-11.6	5.4	7.2	4.7-10.7	3.4	3.0	1.7-4.8	1.6	2.9	1.2-5.8	1.8	3.3	1.4-6.4
Colorectal	1976-1980	36.6	65.5	56.1-75.9	19.4	80.3	64.2-98.9	17.2	55.5	44.2-68.8	15.8	28.7	22.6-36.0	8.2	33.7	23.7-46.3	7.6	24.7	17.3-34.0
	1981-1985	43.0	73.3	63.6-83.9	22.6	91.4	74.3-110.8	20.4	60.8	49.5-73.9	20.2	36.2	29.3-44.0	9.4	42.0	30.2-56.3	10.8	32.7	24.5-42.7
	1986-1990	33.8	53.5	45.7-62.3	16.6	61.8	48.7-77.2	17.2	48.1	38.4-59.4	13.4	21.5	16.6-27.4	7.6	28.7	19.9-39.8	5.8	16.4	11.0-23.5
	1991-1995	36.2	51.9	44.5-60.1	21.0	69.4	56.0-84.9	15.2	39.1	30.8-49.0	14.8	22.4	17.5-28.2	8.8	32.3	23.0-43.9	6.0	15.6	10.5-22.2
	1996-2000	54.0	70.2	62.0-79.2	27.0	79.0	65.5-94.3	27.0	63.2	53.0-74.9	17.2	22.8	18.2-28.3	7.0	21.8	14.8-30.7	10.2	24.1	17.9-31.7
	2001-2005	50.6	56.9	50.0-64.6	23.8	59.9	49.0-72.4	26.8	54.7	45.7-65.0	17.4	20.3	16.2-25.2	9.0	25.0	17.8-34.0	8.4	17.1	12.3-23.2
	2006-2010	42.2	40.9	35.4-47.0	19.2	44.7	35.7-55.2	23.0	39.8	32.7-48.0	14.0	14.7	11.4-18.6	7.4	19.8	13.7-27.5	6.6	11.8	8.1-16.8
	2011-2015	49.0	41.5	36.3-47.3	24.6	45.4	37.4-54.7	24.4	38.3	31.6-46.0	12.4	10.5	8.0-13.6	5.8	10.5	6.9-15.5	6.6	10.2	7.0-14.6
2016-2020	41.4	31.5	27.2-36.3	21.0	32.5	26.3-39.7	20.4	29.4	23.7-36.1	11.4	8.6	6.4-11.3	6.0	9.2	6.1-13.5	5.4	7.5	4.9-11.3	
Colon excluding rectum	1976-1980	27.4	49.0	40.9-58.2	13.6	58.2	44.3-74.6	13.8	43.7	33.9-55.5	12.8	22.7	17.3-29.1	6.4	26.4	17.5-37.8	6.4	20.0	13.6-28.3
	1981-1985	31.4	54.0	45.7-63.3	15.8	64.9	50.6-81.5	15.6	46.6	36.8-58.2	16.8	30.3	24.0-37.6	8.0	36.4	25.4-50.0	8.8	26.8	19.4-35.9
	1986-1990	23.0	36.7	30.2-44.1	11.0	42.6	31.6-55.9	12.0	33.6	25.6-43.2	11.4	18.6	14.0-24.1	6.4	24.1	16.1-34.4	5.0	14.3	9.3-21.1
	1991-1995	22.8	32.8	27.0-39.5	12.2	40.4	30.3-52.5	10.6	27.2	20.3-35.6	11.8	17.6	13.4-22.7	6.6	23.7	15.9-33.7	5.2	13.4	8.7-19.6
	1996-2000	39.8	51.9	44.9-59.8	19.0	56.8	45.3-70.0	20.8	48.3	39.4-58.6	15.0	20.3	15.9-25.4	6.0	19.5	12.8-28.1	9.0	21.3	15.5-28.5
	2001-2005	37.0	42.1	36.1-48.7	17.0	43.7	34.3-54.6	20.0	41.0	33.3-50.1	14.8	17.2	13.4-21.7	7.4	20.6	14.1-28.9	7.4	15.1	10.6-20.9
	2006-2010	30.6	30.1	25.4-35.5	13.8	33.2	25.4-42.5	16.8	29.2	23.1-36.3	11.6	12.3	9.3-16.0	6.0	16.8	11.1-24.1	5.6	10.2	6.7-14.9
	2011-2015	35.0	30.3	25.9-35.3	14.8	28.6	22.2-36.4	20.2	31.7	25.7-38.8	11.2	9.6	7.2-12.5	5.2	9.5	6.0-14.2	6.0	9.3	6.2-13.5
2016-2020	27.8	21.0	17.6-25.0	12.8	19.9	15.2-25.8	15.0	21.4	16.7-27.3	8.2	6.3	4.5-8.6	3.4	5.5	3.1-9.0	4.8	6.7	4.2-10.3	
Rectum & rectosigmoid	1976-1980	9.2	16.5	12.0-22.0	5.8	22.1	14.6-32.0	3.4	11.8	6.8-18.9	3.0	6.0	3.3-10.0	1.8	7.4	3.3-14.0	1.2	4.7	1.7-10.0
	1981-1985	11.6	19.3	14.5-25.0	6.8	26.5	17.7-37.7	4.8	14.2	9.1-21.1	3.4	5.9	3.4-9.4	1.4	5.6	2.2-11.4	2.0	6.0	2.8-11.0
	1986-1990	10.8	16.8	12.6-21.9	5.6	19.2	12.6-27.9	5.2	14.5	9.4-21.2	2.0	3.0	1.4-5.5	1.2	4.6	1.5-10.1	0.8	2.1	0.6-5.4
	1991-1995	13.4	19.1	14.8-24.3	8.8	29.0	20.6-39.5	4.6	11.9	7.6-17.9	3.0	4.8	2.7-7.9	2.2	8.6	4.1-15.5	0.8	2.2	0.6-5.4
	1996-2000	14.2	18.3	14.2-23.1	8.0	22.2	15.5-30.7	6.2	14.9	10.1-21.2	2.2	2.6	1.3-4.7	1.0	2.3	0.7-5.6	1.2	2.8	1.0-6.0
	2001-2005	13.6	14.8	11.5-18.9	6.8	16.2	11.0-23.0	6.8	13.7	9.4-19.3	2.6	3.1	1.6-5.4	1.6	4.4	1.8-8.7	1.0	2.0	0.6-4.8
	2006-2010	11.6	10.8	8.1-14.1	5.4	11.5	7.3-17.1	6.2	10.6	7.1-15.2	2.4	2.3	1.2-4.2	1.4	3.0	1.2-6.4	1.0	1.7	0.5-4.1
	2011-2015	14.0	11.2	8.7-14.3	9.8	16.8	12.2-22.6	4.2	6.6	4.0-10.4	1.2	1.0	0.3-2.2	0.6	1.1	0.2-3.5	0.6	0.9	0.2-3.0
2016-2020	13.6	10.5	8.0-13.4	8.2	12.6	8.9-17.4	5.4	7.9	5.1-11.9	3.2	2.3	1.3-3.9	2.6	3.8	2.0-6.8	0.6	0.8	0.2-2.8	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Putnam**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	1.8	3.1	1.3-5.9	1.4	5.6	2.0-12.1	0.4	1.2	0.1-4.4	1.4	2.2	0.9-4.6	1.2	4.0	1.5-8.8	0.2	0.7	0.0-3.6
	1981-1985	2.4	4.0	2.0-7.0	2.2	9.2	4.3-16.6	0.2	0.5	0.0-3.0	2.4	3.7	1.9-6.5	2.0	7.0	3.3-13.0	0.4	1.1	0.1-3.9
	1986-1990	1.0	1.7	0.5-3.9	0.4	1.1	0.1-4.2	0.6	1.8	0.4-5.1	1.4	2.4	0.9-4.8	0.6	1.9	0.4-5.5	0.8	2.4	0.6-6.0
	1991-1995	2.6	3.7	2.0-6.4	1.8	6.1	2.6-11.7	0.8	2.2	0.6-5.5	3.4	5.1	3.0-8.2	2.2	7.5	3.5-13.5	1.2	3.3	1.2-7.0
	1996-2000	5.0	6.2	4.0-9.1	2.6	6.5	3.4-11.3	2.4	5.7	2.9-10.0	3.6	4.5	2.6-7.1	2.2	5.7	2.8-10.3	1.4	3.3	1.3-6.9
	2001-2005	7.2	7.8	5.4-10.9	4.4	9.9	6.0-15.3	2.8	5.9	3.2-10.0	4.0	4.4	2.7-6.9	2.6	6.2	3.2-10.8	1.4	2.9	1.2-6.1
	2006-2010	6.2	5.8	3.9-8.3	3.6	6.2	3.6-10.2	2.6	5.2	2.8-9.0	4.0	3.9	2.4-6.1	2.0	3.6	1.7-7.0	2.0	3.9	1.9-7.3
	2011-2015	8.8	6.7	4.8-9.1	5.6	8.9	5.8-13.2	3.2	4.8	2.7-8.2	6.6	5.6	3.8-8.0	4.2	8.2	4.9-12.8	2.4	3.8	1.9-6.9
2016-2020	9.6	6.9	5.0-9.3	6.2	8.9	6.0-12.9	3.4	4.9	2.9-8.2	5.4	3.9	2.5-5.8	2.4	3.4	1.7-6.3	3.0	4.4	2.4-7.5	
Pancreas	1976-1980	8.8	16.3	11.7-21.9	5.6	21.7	14.2-31.7	3.2	11.1	6.3-18.1	6.4	12.0	8.1-17.0	3.4	14.4	8.0-23.4	3.0	9.9	5.4-16.4
	1981-1985	6.6	11.6	7.9-16.3	4.0	17.2	10.1-26.9	2.6	7.8	4.1-13.3	6.0	10.2	6.8-14.6	3.6	13.0	7.5-20.8	2.4	7.5	3.9-13.0
	1986-1990	7.0	11.1	7.7-15.5	2.8	10.1	5.5-17.0	4.2	12.0	7.4-18.3	6.2	10.1	6.8-14.3	2.8	10.0	5.3-16.8	3.4	9.9	5.8-15.7
	1991-1995	7.8	11.9	8.5-16.3	3.8	12.4	7.3-19.4	4.0	10.7	6.5-16.5	7.2	11.0	7.7-15.3	3.8	13.0	7.6-20.5	3.4	9.1	5.3-14.5
	1996-2000	11.8	15.0	11.4-19.4	6.4	18.0	11.9-25.8	5.4	12.5	8.2-18.2	11.2	14.8	11.1-19.3	5.6	17.0	11.0-24.9	5.6	12.9	8.6-18.7
	2001-2005	11.4	12.9	9.7-16.9	5.4	13.1	8.4-19.4	6.0	12.7	8.5-18.2	8.8	10.2	7.3-13.7	3.8	9.3	5.4-14.7	5.0	10.5	6.7-15.5
	2006-2010	14.6	14.4	11.2-18.3	6.6	14.3	9.6-20.6	8.0	14.8	10.5-20.3	10.2	9.8	7.3-13.1	4.8	9.4	5.8-14.3	5.4	9.7	6.4-14.3
	2011-2015	17.6	14.7	11.7-18.3	10.0	17.4	12.7-23.3	7.6	12.3	8.6-17.1	12.2	10.6	8.1-13.8	7.6	14.6	10.1-20.3	4.6	7.4	4.7-11.4
2016-2020	19.6	14.8	12.0-18.2	13.0	22.5	17.2-28.9	6.6	9.1	6.2-13.1	14.2	10.5	8.1-13.3	9.0	14.6	10.5-19.9	5.2	7.2	4.7-10.8	
Larynx	1976-1980	3.4	5.6	3.2-9.0	3.0	10.8	5.9-18.0	0.4	1.2	0.1-4.4	1.2	1.8	0.7-4.0	0.8	2.6	0.7-6.9	0.4	1.2	0.1-4.2
	1981-1985	6.2	10.0	6.7-14.2	4.6	16.2	10.0-24.6	1.6	5.0	2.1-9.7	1.8	3.4	1.5-6.5	1.2	6.2	2.0-13.2	0.6	2.0	0.4-5.6
	1986-1990	6.0	9.1	6.1-13.1	4.6	16.4	10.1-24.8	1.4	3.9	1.6-7.9	1.2	1.8	0.7-3.9	1.0	3.2	1.0-7.6	0.2	0.6	0.0-3.1
	1991-1995	4.0	5.1	3.1-7.9	3.6	9.4	5.5-15.1	0.4	1.0	0.1-3.6	2.4	3.7	1.9-6.3	2.0	7.2	3.4-13.2	0.4	1.0	0.1-3.7
	1996-2000	2.2	2.6	1.3-4.7	1.2	3.3	1.2-7.1	1.0	2.2	0.7-5.1	0.6	0.7	0.1-2.0	0.2	0.4	0.0-2.6	0.4	0.9	0.1-3.2
	2001-2005	4.0	3.8	2.3-6.0	3.0	6.1	3.4-10.4	1.0	1.7	0.6-4.2	0.8	1.0	0.3-2.5	0.8	2.5	0.6-6.2	0.0	0.0	0.0-1.6
	2006-2010	3.6	3.0	1.8-4.9	3.0	5.5	2.9-9.4	0.6	1.0	0.2-3.1	1.4	1.5	0.6-3.1	1.0	2.6	0.8-6.2	0.4	0.7	0.1-2.8
	2011-2015	3.6	2.7	1.6-4.5	3.2	5.0	2.8-8.5	0.4	0.6	0.1-2.6	1.0	0.7	0.2-1.9	1.0	1.5	0.5-4.0	0.0	0.0	0.0-1.6
2016-2020	2.2	1.5	0.7-2.8	2.0	2.9	1.3-5.6	0.2	0.2	0.0-1.9	0.8	0.6	0.1-1.6	0.8	1.1	0.3-3.3	0.0	0.0	0.0-1.6	
Lung and bronchus	1976-1980	35.4	61.2	52.3-71.1	26.2	103.4	85.6-123.7	9.2	28.7	20.9-38.3	31.2	56.2	47.5-65.9	23.8	99.8	81.6-120.5	7.4	23.5	16.5-32.5
	1981-1985	40.8	64.5	55.9-74.1	24.0	87.9	72.1-105.9	16.8	48.4	38.5-59.9	31.4	49.2	41.7-57.6	20.0	70.2	56.6-85.9	11.4	32.9	24.9-42.7
	1986-1990	49.6	75.0	65.8-85.0	30.6	106.6	89.5-125.9	19.0	52.5	42.4-64.1	42.8	65.9	57.2-75.4	27.4	98.2	81.6-116.9	15.4	42.9	33.8-53.6
	1991-1995	48.4	68.8	60.3-78.1	28.0	91.2	76.0-108.4	20.4	52.7	43.0-64.0	42.4	61.4	53.3-70.3	24.4	84.2	69.1-101.3	18.0	46.8	37.6-57.5
	1996-2000	66.6	86.0	76.9-95.8	34.6	103.2	87.7-120.6	32.0	75.2	63.9-87.9	48.2	63.6	55.7-72.2	27.8	84.7	70.5-100.7	20.4	49.3	40.2-59.9
	2001-2005	72.6	82.0	73.6-91.1	32.6	84.5	71.4-99.3	40.0	82.6	71.4-95.1	51.4	59.5	52.3-67.4	26.0	71.2	58.9-85.2	25.4	52.7	43.9-62.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Putnam**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	75.0	73.5	66.0-81.6	35.2	77.3	65.7-90.3	39.8	72.2	62.3-83.2	45.2	45.4	39.5-51.9	23.4	54.0	44.2-65.3	21.8	39.5	32.3-47.9
	2011-2015	76.2	63.1	56.7-70.0	36.6	69.0	58.9-80.3	39.6	60.1	51.9-69.5	45.0	38.6	33.6-44.1	22.8	44.2	36.1-53.5	22.2	34.9	28.6-42.3
	2016-2020	85.0	62.0	56.1-68.3	39.6	62.8	54.1-72.6	45.4	62.0	54.0-70.9	42.0	31.5	27.3-36.2	21.0	35.1	28.5-42.9	21.0	29.1	23.7-35.6
Melanoma of the skin	1976-1980	6.0	9.0	6.0-12.9	2.6	8.4	4.4-14.6	3.4	9.6	5.5-15.6	1.6	2.5	1.1-5.0	1.0	3.1	1.0-7.5	0.6	1.9	0.4-5.6
	1981-1985	8.4	12.1	8.6-16.4	3.8	13.4	7.7-21.4	4.6	11.7	7.4-17.8	3.0	4.4	2.4-7.3	2.0	6.5	3.0-12.0	1.0	2.5	0.8-6.0
	1986-1990	4.0	5.1	3.1-7.9	2.6	6.7	3.5-11.7	1.4	3.5	1.4-7.2	2.8	4.4	2.4-7.4	1.6	6.4	2.4-12.9	1.2	3.3	1.2-7.1
	1991-1995	5.6	7.4	4.9-10.8	4.0	12.7	7.5-19.9	1.6	3.7	1.6-7.4	2.8	4.0	2.2-6.8	1.8	7.0	3.0-13.5	1.0	2.5	0.8-5.8
	1996-2000	10.4	11.8	8.8-15.6	6.6	15.6	10.6-22.2	3.8	8.1	4.9-12.8	0.8	1.0	0.3-2.6	0.6	1.9	0.4-5.3	0.2	0.3	0.0-2.2
	2001-2005	20.0	21.3	17.2-26.0	11.0	26.9	19.8-35.5	9.0	17.5	12.7-23.6	2.4	2.5	1.3-4.4	1.6	3.5	1.4-7.1	0.8	1.5	0.4-4.1
	2006-2010	23.4	22.3	18.3-26.8	14.2	30.2	23.2-38.5	9.2	16.2	11.8-21.9	2.6	2.4	1.3-4.3	2.6	5.4	2.7-9.5	0.0	0.0	0.0-1.6
	2011-2015	26.6	22.1	18.4-26.4	14.4	25.6	19.7-32.7	12.2	20.0	15.1-26.1	3.0	2.6	1.4-4.4	1.8	3.2	1.4-6.4	1.2	1.8	0.6-4.2
	2016-2020	28.2	21.6	18.0-25.7	15.0	24.7	19.2-31.4	13.2	19.8	15.1-25.7	1.6	1.2	0.5-2.5	0.8	1.3	0.3-3.6	0.8	1.1	0.3-3.2
Female breast	1976-1980							30.2	97.5	82.4-114.5							11.0	36.6	27.5-47.7
	1981-1985							42.0	121.9	105.8-139.7							13.8	40.7	31.6-51.5
	1986-1990							42.6	114.4	99.4-130.9							14.4	40.1	31.3-50.4
	1991-1995							43.8	106.2	92.5-121.3							13.2	32.9	25.4-41.9
	1996-2000							58.2	125.0	110.9-140.4							14.2	32.2	25.1-40.7
	2001-2005							82.2	153.4	138.7-169.3							12.8	25.2	19.4-32.4
	2006-2010							87.2	151.1	136.9-166.4							13.4	23.0	17.7-29.4
	2011-2015							90.2	142.0	128.8-156.2							13.6	20.9	16.1-26.8
	2016-2020							100.2	150.4	136.8-165.0							9.4	13.2	9.6-17.9
Cervix uteri	1976-1980							3.4	10.0	5.7-16.1							1.2	4.0	1.4-8.7
	1981-1985							5.4	15.7	10.3-22.9							1.8	5.3	2.4-10.0
	1986-1990							3.4	8.8	5.1-14.2							1.2	3.4	1.2-7.3
	1991-1995							5.0	11.6	7.5-17.2							1.4	3.4	1.4-7.0
	1996-2000							4.6	8.8	5.6-13.3							2.0	4.0	1.9-7.5
	2001-2005							3.8	7.7	4.6-12.1							0.8	1.8	0.5-4.6
	2006-2010							2.8	4.9	2.6-8.4							0.2	0.4	0.0-2.4
	2011-2015							3.2	5.4	2.9-9.0							1.0	1.7	0.5-4.3
	2016-2020							2.8	5.6	2.9-9.7							0.6	1.0	0.2-3.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Putnam**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							7.2	23.7	16.5-32.8							2.0	6.3	3.0-11.7
	1981-1985							8.6	24.0	17.3-32.4							1.4	3.9	1.6-8.0
	1986-1990							7.8	21.5	15.3-29.4							0.2	0.6	0.0-3.2
	1991-1995							10.6	26.6	19.9-34.8							1.2	2.8	1.0-6.2
	1996-2000							11.4	25.2	19.0-32.7							1.6	3.6	1.6-7.2
	2001-2005							14.6	28.4	22.2-35.9							1.6	3.3	1.4-6.5
	2006-2010							18.0	30.6	24.5-37.8							2.0	3.5	1.7-6.7
	2011-2015							20.0	29.1	23.5-35.7							1.8	3.0	1.4-5.9
	2016-2020							21.2	29.3	23.8-35.9							3.2	4.3	2.5-7.4
Ovary	1976-1980							4.8	15.9	10.1-23.6							3.8	12.0	7.2-18.8
	1981-1985							5.6	16.6	11.0-24.0							4.4	13.3	8.3-20.1
	1986-1990							4.8	13.0	8.3-19.3							3.4	9.8	5.7-15.6
	1991-1995							6.4	15.7	10.7-22.2							5.2	13.1	8.5-19.2
	1996-2000							7.8	16.8	11.9-23.1							5.4	12.4	8.2-18.1
	2001-2005							9.4	17.7	13.0-23.7							4.4	8.9	5.5-13.5
	2006-2010							7.8	12.9	9.1-17.8							4.0	7.1	4.3-11.0
	2011-2015							10.6	17.6	13.0-23.4							3.8	6.4	3.8-10.2
	2016-2020							9.2	14.1	10.1-19.2							6.2	8.5	5.7-12.4
Prostate	1976-1980				15.4	69.0	53.4-87.2							5.6	30.4	19.6-44.3			
	1981-1985				21.8	97.7	79.2-118.8							6.0	30.2	19.8-43.4			
	1986-1990				22.4	89.1	72.7-107.8							8.8	38.8	27.6-52.4			
	1991-1995				41.8	136.1	117.4-156.7							7.4	32.2	22.2-44.5			
	1996-2000				57.4	153.9	135.7-173.7							7.8	31.1	21.8-42.6			
	2001-2005				71.0	173.2	154.6-193.3							6.8	23.7	16.2-33.1			
	2006-2010				83.0	161.1	145.2-178.4							6.6	19.2	13.1-27.0			
	2011-2015				80.0	128.6	115.7-142.6							6.4	15.1	10.2-21.4			
	2016-2020				75.8	105.3	94.7-117.0							5.0	8.9	5.7-13.4			
Testis	1976-1980				1.4	3.9	1.5-8.3							0.6	1.3	0.3-4.6			
	1981-1985				3.0	8.3	4.4-14.3							0.2	0.4	0.0-3.2			
	1986-1990				2.0	4.5	2.1-8.6							0.0	0.0	0.0-2.4			
	1991-1995				1.8	4.0	1.8-8.0							0.2	0.4	0.0-2.8			
	1996-2000				3.2	6.2	3.5-10.4							0.0	0.0	0.0-2.0			
	2001-2005				3.8	8.2	4.9-13.0							0.2	0.4	0.0-2.4			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Putnam**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				3.0	6.5	3.6-10.9							0.2	0.3	0.0-2.1			
	2011-2015				5.0	10.7	6.8-16.0							0.0	0.0	0.0-1.7			
	2016-2020				4.6	9.5	5.9-14.4							0.2	0.2	0.0-2.0			
Urinary bladder (incl. in situ)	1976-1980	14.0	24.7	19.1-31.4	10.4	40.4	29.5-53.7	3.6	12.2	7.1-19.4	4.6	9.5	5.9-14.3	3.6	18.1	10.3-28.9	1.0	3.5	1.1-8.1
	1981-1985	17.4	29.6	23.6-36.6	14.0	57.6	44.1-73.6	3.4	10.0	5.8-16.0	3.8	7.3	4.3-11.4	2.2	12.4	5.9-22.2	1.6	4.9	2.1-9.6
	1986-1990	15.6	24.1	19.0-30.1	10.4	35.5	26.1-46.9	5.2	14.8	9.6-21.6	3.6	6.2	3.7-9.8	2.4	9.3	4.8-16.2	1.2	3.5	1.3-7.6
	1991-1995	17.0	24.3	19.4-30.1	11.8	37.5	28.1-48.9	5.2	13.8	9.0-20.1	3.2	5.1	2.9-8.3	1.8	6.6	2.9-12.3	1.4	3.8	1.5-7.7
	1996-2000	22.4	28.6	23.4-34.4	15.0	43.3	33.6-54.8	7.4	17.1	12.0-23.6	3.8	5.2	3.1-8.0	2.6	7.9	4.1-13.7	1.2	2.9	1.1-6.2
	2001-2005	28.8	32.2	27.0-38.0	23.0	60.5	49.3-73.3	5.8	11.4	7.6-16.5	4.2	5.5	3.4-8.3	3.4	12.5	7.2-19.9	0.8	1.8	0.5-4.5
	2006-2010	28.2	28.6	23.9-33.9	20.2	46.5	37.4-57.0	8.0	15.1	10.7-20.7	3.4	3.6	2.1-5.8	2.6	6.3	3.2-10.9	0.8	1.5	0.4-4.0
	2011-2015	33.4	28.2	24.0-33.0	25.0	48.7	40.1-58.5	8.4	13.2	9.4-18.1	4.6	3.9	2.5-6.0	3.8	8.1	4.8-13.0	0.8	1.1	0.3-3.2
2016-2020	33.8	24.9	21.3-29.2	25.2	41.9	34.7-50.3	8.6	11.7	8.4-16.0	5.8	4.6	3.1-6.7	4.8	9.1	5.8-13.8	1.0	1.4	0.4-3.6	
Kidney and renal pelvis	1976-1980	3.6	5.9	3.5-9.3	2.6	8.6	4.5-14.9	1.0	3.2	1.0-7.4	2.8	5.3	2.8-8.9	1.2	5.2	1.7-11.6	1.6	5.2	2.2-10.3
	1981-1985	6.4	10.0	6.8-14.2	4.6	15.6	9.8-23.5	1.8	5.3	2.4-10.0	2.0	3.3	1.6-6.0	1.0	3.5	1.1-8.1	1.0	3.2	1.0-7.2
	1986-1990	6.8	10.2	7.0-14.3	5.2	17.6	11.2-26.2	1.6	4.5	2.0-8.9	3.0	4.8	2.6-7.9	2.4	8.5	4.1-15.2	0.6	1.8	0.4-5.0
	1991-1995	7.2	10.2	7.1-14.2	4.4	13.8	8.4-21.1	2.8	7.4	4.1-12.4	2.4	3.6	1.9-6.2	1.8	6.9	3.0-13.0	0.6	1.6	0.3-4.7
	1996-2000	15.6	18.3	14.4-22.9	9.6	23.5	17.0-31.5	6.0	13.7	9.2-19.6	2.8	3.7	2.0-6.2	2.0	6.3	2.8-11.8	0.8	1.9	0.5-4.9
	2001-2005	17.8	18.2	14.5-22.5	10.6	22.7	16.8-30.1	7.2	14.2	9.8-19.7	3.4	3.9	2.2-6.3	1.6	4.3	1.7-8.5	1.8	3.7	1.7-7.2
	2006-2010	23.8	23.0	18.9-27.6	14.2	29.6	22.8-37.7	9.6	17.0	12.5-22.8	3.4	3.3	1.9-5.4	2.6	5.8	2.9-10.1	0.8	1.4	0.4-3.7
	2011-2015	26.6	22.0	18.3-26.3	19.4	32.3	26.0-39.9	7.2	11.9	8.2-16.7	3.8	3.5	2.1-5.5	3.0	6.3	3.4-10.5	0.8	1.3	0.3-3.5
2016-2020	22.6	17.1	13.9-20.7	14.4	22.3	17.2-28.4	8.2	12.2	8.6-17.0	2.8	2.2	1.1-3.8	1.8	3.1	1.3-6.1	1.0	1.4	0.4-3.8	
Brain and other central nervous system	1976-1980	4.2	6.0	3.7-9.3	2.2	6.6	3.2-12.1	2.0	5.4	2.6-10.2	2.4	3.4	1.7-6.1	1.6	4.9	2.1-10.0	0.8	2.1	0.6-5.5
	1981-1985	4.2	5.8	3.6-9.0	3.2	9.4	5.2-15.6	1.0	2.7	0.8-6.3	2.6	3.6	1.9-6.2	1.4	4.1	1.6-8.6	1.2	3.3	1.2-7.2
	1986-1990	5.0	7.0	4.5-10.4	2.4	7.9	3.8-14.2	2.6	6.8	3.6-11.6	2.6	4.1	2.1-7.0	1.8	6.8	2.8-13.4	0.8	2.4	0.6-5.9
	1991-1995	6.0	7.5	5.0-10.8	3.0	7.9	4.3-13.2	3.0	6.9	3.9-11.5	4.6	6.2	3.9-9.4	2.6	7.6	4.0-13.2	2.0	5.0	2.4-9.1
	1996-2000	7.4	8.4	5.9-11.6	4.2	9.6	5.9-15.0	3.2	7.0	4.0-11.5	3.4	4.1	2.4-6.6	1.6	3.9	1.6-7.9	1.8	4.2	1.9-7.9
	2001-2005	6.6	7.1	4.9-10.1	3.8	8.7	5.1-13.8	2.8	6.2	3.3-10.5	4.0	4.2	2.5-6.6	2.0	4.5	2.0-8.4	2.0	4.0	1.9-7.4
	2006-2010	7.4	7.8	5.5-10.9	4.0	9.1	5.5-14.3	3.4	6.9	4.0-11.1	4.6	4.7	2.9-7.1	2.4	5.2	2.6-9.3	2.2	4.2	2.1-7.7
	2011-2015	8.0	7.2	5.1-10.0	4.8	8.6	5.4-13.2	3.2	5.8	3.2-9.7	4.0	3.2	1.9-5.0	2.6	4.0	2.1-7.3	1.4	2.2	0.9-4.8
2016-2020	8.6	6.7	4.8-9.2	5.6	9.4	6.1-13.9	3.0	4.3	2.3-7.5	4.6	3.5	2.2-5.4	2.6	3.8	2.0-6.9	2.0	3.2	1.4-6.2	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Putnam**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	3.0	4.9	2.7-8.2	0.4	1.1	0.1-4.6	2.6	8.2	4.3-14.1	0.8	1.5	0.4-3.9	0.2	0.7	0.0-4.1	0.6	2.1	0.4-6.0
	1981-1985	1.8	2.5	1.1-4.7	0.6	1.7	0.3-5.2	1.2	3.3	1.2-7.2	0.0	0.0	0.0-1.1	0.0	0.0	0.0-2.5	0.0	0.0	0.0-2.0
	1986-1990	2.2	2.8	1.4-5.0	0.4	1.0	0.1-4.0	1.8	4.5	2.0-8.6	0.4	0.6	0.1-2.0	0.4	1.1	0.1-4.2	0.0	0.0	0.0-1.9
	1991-1995	3.2	3.8	2.1-6.2	1.0	3.4	0.9-8.2	2.2	4.7	2.3-8.6	0.2	0.2	0.0-1.4	0.0	0.0	0.0-2.2	0.2	0.5	0.0-2.7
	1996-2000	8.2	8.7	6.2-11.9	2.8	5.5	3.0-9.5	5.4	11.7	7.6-17.3	0.4	0.6	0.1-2.1	0.0	0.0	0.0-2.0	0.4	1.0	0.1-3.6
	2001-2005	18.4	17.8	14.3-22.0	4.8	10.7	6.6-16.3	13.6	25.5	19.7-32.6	0.2	0.3	0.0-1.4	0.2	0.9	0.0-4.1	0.0	0.0	0.0-1.6
	2006-2010	21.0	18.9	15.3-23.0	4.8	8.7	5.4-13.3	16.2	29.0	22.9-36.3	0.6	0.5	0.1-1.6	0.4	0.6	0.1-2.6	0.2	0.3	0.0-2.1
	2011-2015	30.0	28.1	23.6-33.3	8.0	14.5	10.1-20.1	22.0	42.1	34.2-51.3	0.4	0.4	0.1-1.6	0.2	0.5	0.0-2.7	0.2	0.4	0.0-2.4
2016-2020	22.0	19.5	15.9-23.9	6.4	10.7	7.1-15.6	15.6	28.6	22.2-36.2	0.4	0.3	0.0-1.3	0.2	0.3	0.0-2.1	0.2	0.3	0.0-2.2	
Hodgkin lymphoma	1976-1980	3.4	4.4	2.5-7.2	2.0	5.2	2.4-10.1	1.4	3.5	1.3-7.4	1.0	1.6	0.5-3.8	0.8	2.6	0.7-6.8	0.2	0.7	0.0-3.7
	1981-1985	3.4	4.3	2.4-6.9	2.0	5.6	2.6-10.7	1.4	3.2	1.3-6.8	0.4	0.6	0.1-2.3	0.4	1.5	0.2-5.4	0.0	0.0	0.0-2.0
	1986-1990	3.8	4.5	2.7-7.2	1.8	4.1	1.9-8.2	2.0	4.9	2.3-9.1	1.0	1.2	0.4-2.9	0.8	1.7	0.5-4.9	0.2	0.6	0.0-3.1
	1991-1995	2.2	2.6	1.3-4.7	0.8	1.8	0.5-4.9	1.4	3.5	1.4-7.2	0.0	0.0	0.0-0.9	0.0	0.0	0.0-2.2	0.0	0.0	0.0-1.8
	1996-2000	4.0	4.3	2.6-6.7	3.0	6.3	3.5-10.6	1.0	2.2	0.7-5.2	0.2	0.3	0.0-1.4	0.0	0.0	0.0-2.0	0.2	0.5	0.0-2.7
	2001-2005	2.6	2.9	1.5-5.0	1.4	3.4	1.3-7.3	1.2	2.7	0.9-5.9	0.6	0.6	0.1-1.7	0.4	0.8	0.1-3.1	0.2	0.3	0.0-2.2
	2006-2010	4.2	4.4	2.7-6.8	2.0	4.2	2.0-7.8	2.2	4.4	2.1-8.0	0.4	0.4	0.0-1.5	0.2	0.5	0.0-2.6	0.2	0.3	0.0-2.1
	2011-2015	4.4	4.4	2.7-6.8	2.4	4.1	2.1-7.5	2.0	4.6	2.1-8.6	0.4	0.3	0.0-1.3	0.2	0.2	0.0-2.0	0.2	0.3	0.0-2.0
2016-2020	5.2	5.4	3.5-8.1	2.6	5.4	2.8-9.3	2.6	5.4	2.8-9.5	0.6	0.4	0.1-1.5	0.4	0.5	0.1-2.4	0.2	0.3	0.0-2.2	
Non-Hodgkin lymphomas	1976-1980	8.0	13.5	9.5-18.6	3.8	13.5	7.7-21.7	4.2	13.2	8.1-20.3	4.4	8.2	5.0-12.5	1.8	8.6	3.5-16.9	2.6	8.3	4.3-14.2
	1981-1985	9.6	14.7	10.7-19.5	5.0	16.0	10.2-23.9	4.6	13.1	8.3-19.7	4.6	7.2	4.5-10.8	2.4	7.1	3.7-12.6	2.2	6.6	3.2-11.7
	1986-1990	7.6	11.3	7.9-15.5	3.4	11.2	6.1-18.6	4.2	11.5	7.1-17.6	6.0	9.1	6.1-13.0	2.2	7.8	3.7-14.1	3.8	10.0	6.0-15.6
	1991-1995	12.8	17.4	13.3-22.2	7.0	22.3	15.0-31.6	5.8	14.6	9.7-21.0	7.4	10.8	7.5-14.8	3.8	13.7	7.9-21.5	3.6	9.1	5.4-14.4
	1996-2000	15.6	18.8	14.7-23.5	9.0	26.0	18.5-35.3	6.6	14.4	9.9-20.4	5.8	7.7	5.1-11.1	2.2	8.5	4.1-15.1	3.6	8.3	4.9-13.2
	2001-2005	22.4	24.3	19.9-29.4	12.0	27.9	20.9-36.4	10.4	21.1	15.7-27.8	6.8	8.0	5.5-11.2	4.6	12.9	7.9-19.7	2.2	4.6	2.3-8.3
	2006-2010	23.6	22.8	18.7-27.4	12.0	24.9	18.7-32.5	11.6	21.2	16.0-27.7	5.8	6.0	4.0-8.7	3.0	7.4	4.1-12.3	2.8	4.8	2.6-8.2
	2011-2015	27.4	22.3	18.6-26.6	17.2	28.9	22.9-36.2	10.2	16.3	12.0-21.7	5.6	5.2	3.4-7.6	2.0	3.6	1.7-6.9	3.6	6.2	3.6-10.1
2016-2020	31.8	23.7	20.0-28.0	16.0	24.8	19.5-31.3	15.8	23.0	18.0-29.1	7.2	5.5	3.8-7.8	4.2	7.3	4.4-11.4	3.0	4.1	2.2-7.0	
Myeloma	1976-1980	2.2	3.7	1.8-6.6	1.0	4.0	1.3-9.3	1.2	3.6	1.3-7.9	1.4	2.4	0.9-5.0	0.4	1.7	0.2-5.9	1.0	2.9	0.9-6.9
	1981-1985	3.2	5.4	3.0-8.7	1.8	7.3	3.1-14.1	1.4	4.1	1.7-8.5	2.0	3.4	1.6-6.2	1.0	4.7	1.4-10.9	1.0	2.8	0.9-6.4
	1986-1990	3.2	4.9	2.8-7.9	2.0	6.8	3.0-12.7	1.2	3.4	1.2-7.2	2.2	3.5	1.7-6.2	0.6	2.3	0.4-6.6	1.6	4.5	2.0-8.8
	1991-1995	4.6	6.8	4.3-10.2	2.2	7.9	3.8-14.3	2.4	5.9	3.1-10.4	3.4	5.0	2.9-8.0	2.0	6.2	2.9-11.4	1.4	3.5	1.4-7.2
	1996-2000	4.4	5.2	3.2-7.9	1.6	3.6	1.5-7.3	2.8	6.4	3.5-10.8	3.4	4.6	2.6-7.3	1.2	3.6	1.3-7.8	2.2	5.2	2.6-9.2
	2001-2005	8.2	8.7	6.2-11.9	4.6	9.9	6.2-15.2	3.6	7.1	4.2-11.4	3.6	4.0	2.3-6.4	2.6	6.1	3.1-10.6	1.0	2.0	0.6-4.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Putnam**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	6.8	6.4	4.4-9.1	3.2	6.4	3.5-10.7	3.6	6.2	3.6-10.0	3.2	3.1	1.7-5.1	1.8	3.8	1.7-7.4	1.4	2.3	0.9-5.0
	2011-2015	10.4	8.6	6.4-11.5	5.8	10.4	6.8-15.3	4.6	7.7	4.8-11.7	2.2	2.0	1.0-3.6	1.6	3.2	1.3-6.4	0.6	0.9	0.2-3.0
	2016-2020	8.6	6.6	4.7-9.1	6.0	10.3	6.8-14.9	2.6	3.6	1.9-6.5	2.8	2.1	1.1-3.6	1.8	3.1	1.4-6.2	1.0	1.4	0.5-3.7
Leukemias	1976-1980	7.4	12.5	8.6-17.5	4.8	18.3	11.1-28.2	2.6	8.0	4.1-13.8	6.0	10.1	6.7-14.6	4.0	15.2	8.8-24.1	2.0	6.2	2.9-11.5
	1981-1985	7.6	12.1	8.5-16.8	4.4	15.0	9.2-23.1	3.2	9.5	5.4-15.4	5.2	8.7	5.6-12.9	3.6	12.6	7.2-20.4	1.6	5.2	2.2-10.0
	1986-1990	7.2	10.5	7.3-14.6	4.4	14.0	8.4-21.7	2.8	7.6	4.1-12.7	4.8	7.3	4.6-10.8	3.0	9.9	5.3-16.5	1.8	5.2	2.4-9.8
	1991-1995	8.6	12.2	8.8-16.5	5.8	20.5	13.3-29.9	2.8	6.9	3.8-11.7	6.4	9.3	6.3-13.2	4.6	17.5	10.7-26.5	1.8	4.4	2.0-8.4
	1996-2000	13.4	16.8	13.0-21.4	8.4	24.0	16.9-32.8	5.0	11.3	7.3-16.7	6.4	8.4	5.7-11.8	4.4	12.5	7.6-19.1	2.0	4.8	2.3-8.8
	2001-2005	16.2	18.2	14.4-22.7	8.2	20.2	14.2-27.8	8.0	16.3	11.6-22.3	5.4	6.7	4.4-9.8	2.8	7.9	4.2-13.3	2.6	5.5	2.9-9.6
	2006-2010	15.2	15.2	11.9-19.1	10.2	21.7	15.9-29.0	5.0	9.3	6.0-13.9	4.8	5.0	3.1-7.5	3.4	8.6	4.9-14.0	1.4	2.2	0.9-4.8
	2011-2015	25.6	22.5	18.7-27.0	12.8	23.8	18.1-30.8	12.8	21.7	16.6-28.1	8.4	7.9	5.7-10.8	4.6	10.1	6.3-15.4	3.8	6.1	3.7-9.8
2016-2020	19.8	14.7	11.8-18.1	12.2	18.8	14.2-24.6	7.6	11.3	7.8-16.0	7.2	5.6	3.9-7.9	4.2	7.5	4.6-11.7	3.0	4.2	2.3-7.3	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Rensselaer**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	586.4	405.0	390.1-420.2	289.6	484.6	458.8-511.4	296.8	361.8	343.1-381.2	328.4	227.4	216.4-238.9	176.0	298.6	278.3-319.9	152.4	182.8	169.8-196.5
	1981-1985	654.4	445.2	429.7-461.0	325.0	550.0	522.4-578.5	329.4	390.4	371.2-410.3	337.6	229.2	218.2-240.6	174.4	300.9	280.4-322.5	163.2	187.9	174.9-201.6
	1986-1990	654.8	435.8	420.8-451.2	328.6	539.9	513.2-567.5	326.2	379.7	361.0-399.1	357.2	237.7	226.6-249.1	181.6	307.8	287.3-329.2	175.6	199.5	186.3-213.5
	1991-1995	655.4	421.8	407.4-436.6	331.8	513.2	488.2-539.1	323.6	369.5	351.4-388.3	357.6	230.3	219.7-241.3	182.8	296.7	277.3-317.1	174.8	191.2	178.6-204.6
	1996-2000	854.6	537.3	521.3-553.7	431.4	635.3	608.4-662.9	423.2	476.9	456.6-497.9	360.4	224.3	214.0-234.9	177.4	271.5	253.6-290.2	183.0	195.8	183.2-209.1
	2001-2005	894.0	541.0	525.2-557.2	457.4	638.3	612.1-665.4	436.6	477.2	457.2-497.9	353.4	212.3	202.5-222.4	179.8	262.5	245.4-280.5	173.6	181.7	169.7-194.4
	2006-2010	994.4	560.6	544.9-576.6	516.8	647.3	622.0-673.3	477.6	501.0	480.7-521.9	351.4	196.6	187.4-206.1	184.6	241.8	226.2-258.3	166.8	165.7	154.4-177.6
	2011-2015	985.6	516.3	501.6-531.3	493.0	565.1	542.3-588.7	492.6	483.1	463.6-503.3	347.8	180.3	171.7-189.2	180.6	216.9	202.6-232.0	167.2	155.8	145.1-167.1
2016-2020	1,032.8	504.2	490.1-518.7	537.8	562.6	540.8-585.2	495.0	461.7	443.0-481.1	316.6	152.6	145.0-160.6	161.6	178.6	166.1-191.8	155.0	135.0	125.3-145.2	
Oral cavity and pharynx	1976-1980	18.6	12.3	9.9-15.1	13.0	20.4	15.6-26.2	5.6	6.5	4.3-9.5	9.2	6.6	4.8-8.8	6.6	11.0	7.5-15.6	2.6	3.1	1.6-5.3
	1981-1985	20.0	13.6	11.0-16.6	14.4	22.4	17.4-28.4	5.6	6.8	4.5-9.9	8.6	5.6	4.0-7.6	5.6	8.7	5.7-12.7	3.0	3.2	1.8-5.4
	1986-1990	16.4	10.9	8.7-13.6	11.4	17.7	13.3-23.1	5.0	6.4	4.1-9.6	7.4	4.8	3.4-6.7	5.0	7.7	4.9-11.5	2.4	3.1	1.6-5.5
	1991-1995	15.0	9.9	7.8-12.4	10.4	15.8	11.7-20.7	4.6	5.9	3.7-8.9	4.0	2.6	1.6-4.0	2.8	4.4	2.4-7.5	1.2	1.4	0.5-3.1
	1996-2000	20.0	12.7	10.3-15.5	14.0	20.4	15.9-25.8	6.0	6.9	4.6-9.8	5.4	3.4	2.2-4.9	3.8	5.9	3.5-9.2	1.6	1.7	0.7-3.4
	2001-2005	20.2	12.1	9.8-14.7	13.4	16.9	13.1-21.6	6.8	7.4	5.1-10.4	4.6	2.7	1.7-4.1	2.4	2.9	1.5-5.1	2.2	2.2	1.1-4.0
	2006-2010	21.8	12.0	9.8-14.5	14.6	17.4	13.6-22.1	7.2	7.3	5.1-10.3	5.0	2.8	1.8-4.1	3.4	4.5	2.6-7.3	1.6	1.4	0.6-2.9
	2011-2015	22.8	11.7	9.6-14.2	16.8	18.9	15.0-23.6	6.0	5.3	3.5-7.7	3.2	1.6	0.9-2.7	2.6	2.9	1.5-5.1	0.6	0.5	0.1-1.7
2016-2020	31.8	14.5	12.3-17.1	24.4	24.1	19.9-29.0	7.4	6.1	4.3-8.5	7.0	3.2	2.2-4.6	4.8	4.9	3.1-7.6	2.2	1.8	0.9-3.4	
Esophagus	1976-1980	6.4	4.5	3.0-6.4	4.4	7.0	4.3-10.6	2.0	2.3	1.1-4.4	6.8	4.8	3.3-6.8	5.2	9.3	5.9-13.7	1.6	1.8	0.8-3.7
	1981-1985	8.4	5.4	3.9-7.4	6.4	9.9	6.7-14.1	2.0	2.1	1.0-3.9	5.4	3.6	2.4-5.3	5.0	8.5	5.4-12.7	0.4	0.4	0.1-1.7
	1986-1990	7.4	5.1	3.5-7.0	4.8	8.2	5.1-12.3	2.6	2.8	1.5-4.9	8.0	5.4	3.8-7.3	5.8	9.7	6.4-14.0	2.2	2.4	1.2-4.4
	1991-1995	10.8	6.9	5.1-9.0	7.6	11.2	7.9-15.5	3.2	3.4	1.9-5.6	8.6	5.4	3.9-7.3	5.6	8.2	5.4-11.9	3.0	3.3	1.8-5.5
	1996-2000	11.2	7.1	5.4-9.2	8.2	11.9	8.5-16.2	3.0	3.2	1.8-5.4	10.4	6.5	4.9-8.6	8.0	11.8	8.4-16.1	2.4	2.6	1.3-4.6
	2001-2005	13.4	7.9	6.1-10.1	8.8	11.8	8.5-15.9	4.6	4.8	3.0-7.3	12.6	7.5	5.7-9.6	8.6	11.8	8.5-15.9	4.0	3.8	2.3-6.0
	2006-2010	12.8	7.2	5.6-9.3	10.0	12.7	9.4-16.9	2.8	2.8	1.5-4.9	13.6	7.5	5.8-9.5	9.8	12.2	8.9-16.2	3.8	3.8	2.3-6.0
	2011-2015	11.0	5.6	4.2-7.4	8.0	8.9	6.3-12.2	3.0	2.9	1.6-4.9	8.8	4.8	3.5-6.5	6.8	8.1	5.5-11.4	2.0	2.2	1.0-4.1
2016-2020	16.2	7.6	6.0-9.6	13.2	14.0	10.7-18.1	3.0	2.5	1.4-4.3	11.8	5.6	4.2-7.3	9.8	10.4	7.6-13.9	2.0	1.7	0.8-3.4	
Stomach	1976-1980	14.6	10.7	8.4-13.5	8.4	15.8	11.1-21.6	6.2	7.8	5.2-11.1	12.4	9.0	6.9-11.6	7.2	13.3	9.1-18.7	5.2	6.3	4.1-9.3
	1981-1985	14.8	10.0	7.9-12.6	8.6	15.0	10.7-20.3	6.2	7.0	4.7-10.0	10.0	6.7	5.0-8.9	5.4	8.5	5.5-12.5	4.6	5.0	3.2-7.7
	1986-1990	14.6	9.8	7.7-12.4	9.2	15.6	11.3-20.9	5.4	5.8	3.8-8.4	10.0	6.9	5.1-9.1	5.8	10.5	6.9-15.1	4.2	4.6	2.8-7.0
	1991-1995	12.0	7.7	5.9-9.9	8.0	12.3	8.8-16.8	4.0	4.2	2.5-6.6	9.6	6.3	4.6-8.4	7.4	12.2	8.5-16.9	2.2	2.1	1.1-3.9
	1996-2000	12.0	7.5	5.7-9.7	6.8	9.8	6.8-13.7	5.2	5.6	3.6-8.2	7.8	4.9	3.5-6.6	4.0	6.0	3.7-9.4	3.8	3.9	2.3-6.2
	2001-2005	11.4	6.8	5.2-8.8	6.4	9.6	6.5-13.6	5.0	5.1	3.3-7.6	6.4	3.8	2.6-5.4	4.4	6.5	4.1-9.9	2.0	2.1	1.0-3.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Rensselaer**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	11.2	6.3	4.7-8.2	6.6	8.1	5.5-11.5	4.6	4.7	3.0-7.2	3.2	1.7	1.0-2.8	1.6	1.9	0.8-3.8	1.6	1.4	0.6-3.0
	2011-2015	12.0	6.7	5.1-8.7	7.4	9.0	6.2-12.5	4.6	4.5	2.8-7.0	6.2	3.3	2.2-4.8	3.2	4.0	2.2-6.5	3.0	2.7	1.5-4.7
	2016-2020	14.0	6.9	5.3-8.9	9.6	10.5	7.6-14.1	4.4	4.1	2.5-6.4	4.8	2.2	1.4-3.3	3.4	3.4	2.0-5.7	1.4	1.2	0.5-2.6
Colorectal	1976-1980	107.6	75.1	68.8-81.9	49.2	83.0	72.5-94.5	58.4	68.4	60.6-76.8	56.4	39.3	34.8-44.3	27.8	47.3	39.4-56.2	28.6	33.3	28.0-39.3
	1981-1985	103.6	70.7	64.6-77.1	53.0	94.1	82.7-106.7	50.6	56.1	49.3-63.6	58.6	40.6	36.1-45.6	29.2	53.9	45.1-63.7	29.4	33.3	28.1-39.3
	1986-1990	94.6	62.6	57.1-68.6	49.2	84.0	73.5-95.5	45.4	49.2	42.9-56.1	47.0	31.2	27.3-35.5	21.6	37.7	30.7-45.8	25.4	27.7	23.0-33.0
	1991-1995	83.4	53.3	48.3-58.7	38.4	61.6	53.0-71.0	45.0	46.9	40.9-53.5	43.2	27.6	24.0-31.6	21.4	34.7	28.3-42.0	21.8	22.9	18.8-27.8
	1996-2000	108.2	67.3	61.8-73.3	56.6	84.9	75.1-95.5	51.6	54.1	47.6-61.3	41.4	25.5	22.2-29.3	19.4	29.6	23.9-36.2	22.0	22.2	18.2-26.9
	2001-2005	101.6	61.1	55.9-66.7	49.0	69.7	61.1-79.2	52.6	53.3	47.0-60.3	34.2	20.6	17.6-23.9	16.6	24.3	19.3-30.2	17.6	17.1	13.7-21.2
	2006-2010	90.0	50.8	46.2-55.8	44.0	56.7	49.3-64.9	46.0	45.3	39.5-51.7	28.0	15.4	12.9-18.2	13.0	17.3	13.3-22.1	15.0	13.7	10.7-17.4
	2011-2015	80.0	42.3	38.1-46.7	39.0	46.8	40.3-54.1	41.0	39.3	33.9-45.3	29.8	15.5	13.1-18.3	15.6	19.6	15.4-24.6	14.2	12.7	9.8-16.3
2016-2020	72.4	37.3	33.4-41.4	37.4	42.5	36.4-49.3	35.0	32.5	27.7-37.9	19.8	9.8	7.9-12.0	11.0	12.6	9.4-16.5	8.8	7.5	5.4-10.3	
Colon excluding rectum	1976-1980	79.2	55.4	50.0-61.3	35.8	61.2	52.2-71.3	43.4	51.0	44.4-58.5	45.0	31.7	27.7-36.2	20.4	35.4	28.5-43.3	24.6	28.8	23.8-34.4
	1981-1985	75.4	51.9	46.7-57.4	35.8	64.7	55.1-75.2	39.6	44.0	38.0-50.7	47.0	32.7	28.6-37.2	22.8	42.0	34.3-50.8	24.2	27.6	22.8-33.1
	1986-1990	66.4	44.0	39.3-49.0	32.0	56.3	47.6-66.0	34.4	37.1	31.7-43.2	41.0	27.3	23.7-31.4	18.8	33.3	26.7-41.0	22.2	24.0	19.7-29.1
	1991-1995	65.4	42.0	37.5-46.8	27.6	44.5	37.2-52.6	37.8	39.4	33.9-45.5	37.0	23.6	20.3-27.3	17.2	27.6	21.9-34.2	19.8	20.8	16.8-25.4
	1996-2000	79.0	49.1	44.4-54.2	38.0	57.7	49.7-66.7	41.0	42.8	37.0-49.1	34.6	21.3	18.2-24.7	15.6	24.1	19.0-30.2	19.0	19.2	15.5-23.5
	2001-2005	75.4	45.2	40.8-50.0	35.0	50.3	43.0-58.4	40.4	40.4	34.9-46.5	27.8	16.7	14.0-19.8	12.8	18.8	14.4-24.0	15.0	14.5	11.4-18.3
	2006-2010	67.6	38.2	34.2-42.5	32.6	42.7	36.3-49.9	35.0	34.2	29.2-39.9	22.2	12.1	9.9-14.6	9.8	13.2	9.7-17.5	12.4	11.2	8.5-14.5
	2011-2015	56.8	30.4	26.9-34.2	25.8	31.5	26.2-37.7	31.0	29.5	24.9-34.8	23.6	12.3	10.1-14.8	11.0	13.8	10.3-18.2	12.6	11.4	8.6-14.8
2016-2020	51.2	25.8	22.7-29.3	23.6	26.6	21.9-32.1	27.6	24.9	20.8-29.7	14.8	7.3	5.7-9.3	7.6	8.7	6.1-12.1	7.2	6.0	4.2-8.6	
Rectum & rectosigmoid	1976-1980	28.4	19.7	16.5-23.3	13.4	21.7	16.7-27.8	15.0	17.3	13.6-21.8	11.4	7.6	5.7-9.9	7.4	11.9	8.3-16.6	4.0	4.5	2.8-7.0
	1981-1985	28.2	18.8	15.8-22.3	17.2	29.5	23.3-36.7	11.0	12.1	9.1-15.9	11.6	8.0	6.1-10.4	6.4	11.9	8.0-16.9	5.2	5.7	3.7-8.5
	1986-1990	28.2	18.7	15.7-22.1	17.2	27.8	22.0-34.4	11.0	12.0	9.0-15.8	6.0	3.9	2.6-5.6	2.8	4.4	2.3-7.5	3.2	3.6	2.0-6.0
	1991-1995	18.0	11.4	9.1-14.0	10.8	17.1	12.8-22.4	7.2	7.5	5.2-10.4	6.2	4.0	2.7-5.7	4.2	7.1	4.3-10.8	2.0	2.2	1.0-4.1
	1996-2000	29.2	18.2	15.4-21.4	18.6	27.1	21.9-33.3	10.6	11.4	8.5-15.0	6.8	4.2	2.9-5.9	3.8	5.5	3.3-8.5	3.0	3.1	1.7-5.1
	2001-2005	26.2	15.9	13.3-18.9	14.0	19.5	15.1-24.6	12.2	13.0	9.9-16.8	6.4	3.9	2.6-5.5	3.8	5.6	3.3-8.7	2.6	2.6	1.3-4.5
	2006-2010	22.4	12.6	10.4-15.2	11.4	14.0	10.6-18.3	11.0	11.0	8.3-14.5	5.8	3.3	2.2-4.7	3.2	4.1	2.3-6.8	2.6	2.5	1.3-4.4
	2011-2015	23.2	11.9	9.8-14.4	13.2	15.3	11.8-19.6	10.0	9.8	7.2-13.1	6.2	3.2	2.2-4.6	4.6	5.8	3.6-8.8	1.6	1.3	0.6-2.8
2016-2020	21.2	11.4	9.3-13.9	13.8	15.9	12.2-20.3	7.4	7.5	5.2-10.5	5.0	2.5	1.6-3.7	3.4	3.9	2.2-6.4	1.6	1.5	0.6-3.1	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Rensselaer**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	6.0	3.9	2.6-5.7	4.4	7.3	4.5-11.1	1.6	1.8	0.8-3.6	4.2	2.8	1.7-4.3	3.6	6.4	3.7-10.2	0.6	0.6	0.1-2.0
	1981-1985	3.4	2.2	1.3-3.5	1.8	2.9	1.3-5.6	1.6	1.8	0.8-3.6	2.6	1.7	0.9-2.9	1.4	2.3	0.9-4.9	1.2	1.4	0.5-3.1
	1986-1990	4.6	2.9	1.8-4.4	2.8	4.3	2.3-7.3	1.8	1.9	0.8-3.6	4.4	2.9	1.8-4.5	2.4	3.7	1.9-6.6	2.0	2.2	1.0-4.1
	1991-1995	8.2	5.2	3.7-7.0	6.0	10.3	6.8-14.7	2.2	2.3	1.2-4.3	9.0	5.8	4.2-7.8	6.2	10.3	6.9-14.7	2.8	3.1	1.7-5.4
	1996-2000	7.2	4.5	3.1-6.2	3.8	5.9	3.5-9.3	3.4	3.7	2.1-6.0	6.8	4.2	2.9-5.8	3.0	5.2	2.9-8.6	3.8	3.8	2.3-6.1
	2001-2005	7.2	4.3	3.0-5.9	4.8	6.5	4.1-9.8	2.4	2.6	1.3-4.7	7.8	4.6	3.3-6.3	5.8	7.9	5.3-11.4	2.0	2.1	1.0-4.0
	2006-2010	14.4	8.1	6.3-10.2	10.2	12.2	9.0-16.2	4.2	4.5	2.7-6.9	12.8	7.0	5.4-9.0	9.0	10.9	7.9-14.7	3.8	3.9	2.3-6.2
	2011-2015	17.6	8.5	6.8-10.6	13.8	14.3	11.0-18.3	3.8	3.5	2.1-5.6	13.4	6.7	5.2-8.6	9.6	10.3	7.5-13.9	3.8	3.6	2.2-5.8
	2016-2020	18.6	8.8	7.0-10.8	14.2	14.8	11.4-18.9	4.4	3.9	2.4-6.1	11.0	5.3	3.9-7.0	9.2	9.6	6.9-13.0	1.8	1.7	0.7-3.3
Pancreas	1976-1980	17.0	11.9	9.4-14.7	9.6	15.7	11.4-20.9	7.4	9.0	6.3-12.6	17.8	12.3	9.9-15.2	9.4	15.0	10.9-20.0	8.4	10.3	7.4-14.0
	1981-1985	18.4	12.7	10.2-15.6	8.6	15.6	11.1-21.2	9.8	10.9	8.0-14.5	16.0	11.1	8.8-13.9	7.8	14.3	9.9-19.7	8.2	9.2	6.6-12.6
	1986-1990	17.0	11.2	8.9-13.9	9.2	14.3	10.4-19.2	7.8	8.8	6.2-12.1	16.0	10.6	8.4-13.3	8.8	14.2	10.2-19.2	7.2	8.0	5.5-11.1
	1991-1995	22.6	14.9	12.2-17.9	9.4	14.6	10.7-19.5	13.2	14.9	11.5-19.1	21.0	13.7	11.2-16.5	8.8	14.2	10.2-19.1	12.2	13.3	10.1-17.2
	1996-2000	19.0	11.8	9.5-14.4	9.2	13.4	9.8-17.9	9.8	10.5	7.7-14.0	16.8	10.4	8.3-12.9	7.6	11.3	7.9-15.5	9.2	9.7	7.1-13.1
	2001-2005	21.2	12.8	10.5-15.5	10.6	14.5	10.9-19.1	10.6	11.3	8.4-14.8	19.6	11.8	9.6-14.4	8.8	11.9	8.6-16.1	10.8	11.4	8.6-15.0
	2006-2010	22.8	12.5	10.3-15.1	13.8	17.1	13.2-21.8	9.0	8.6	6.2-11.6	17.6	9.6	7.7-11.9	9.4	11.7	8.6-15.7	8.2	7.7	5.5-10.5
	2011-2015	25.6	13.4	11.1-16.0	14.0	16.3	12.6-20.7	11.6	10.9	8.2-14.2	21.2	10.7	8.7-13.0	12.8	14.1	10.8-18.2	8.4	7.6	5.4-10.4
	2016-2020	33.0	16.3	13.9-19.1	18.0	19.8	15.8-24.5	15.0	13.6	10.6-17.2	21.2	10.5	8.6-12.8	11.4	12.8	9.6-16.7	9.8	9.0	6.6-12.1
Larynx	1976-1980	10.4	7.1	5.3-9.4	7.8	12.8	8.9-17.6	2.6	3.2	1.6-5.5	3.6	2.5	1.5-3.9	2.8	4.8	2.5-8.1	0.8	1.1	0.3-2.8
	1981-1985	13.6	9.2	7.1-11.7	10.4	15.9	11.8-21.0	3.2	4.1	2.3-6.8	2.8	1.8	1.0-3.0	2.2	3.5	1.7-6.4	0.6	0.6	0.1-1.9
	1986-1990	11.2	7.3	5.5-9.5	9.0	13.7	9.9-18.4	2.2	2.7	1.3-4.9	3.4	2.1	1.2-3.4	3.0	4.6	2.6-7.7	0.4	0.4	0.1-1.7
	1991-1995	9.6	6.3	4.6-8.3	7.0	10.7	7.4-14.9	2.6	3.0	1.6-5.2	3.4	2.2	1.3-3.5	2.8	4.4	2.3-7.3	0.6	0.6	0.1-1.9
	1996-2000	8.4	5.4	3.9-7.3	7.6	10.6	7.5-14.6	0.8	0.9	0.2-2.4	3.0	1.9	1.0-3.1	2.2	3.1	1.6-5.6	0.8	0.8	0.2-2.1
	2001-2005	8.8	5.3	3.8-7.1	7.0	8.9	6.2-12.4	1.8	2.1	1.0-4.0	2.4	1.4	0.7-2.5	1.6	2.1	0.9-4.2	0.8	0.9	0.2-2.4
	2006-2010	12.0	6.5	5.0-8.5	9.6	11.1	8.1-14.8	2.4	2.6	1.3-4.6	3.4	1.8	1.0-2.9	3.0	3.8	2.1-6.4	0.4	0.4	0.0-1.5
	2011-2015	7.6	3.6	2.5-5.0	6.0	6.4	4.2-9.2	1.6	1.3	0.5-2.7	4.2	1.9	1.2-3.0	3.6	3.6	2.1-5.9	0.6	0.5	0.1-1.8
	2016-2020	8.6	4.1	2.9-5.5	7.4	7.7	5.4-10.8	1.2	1.0	0.3-2.3	1.2	0.6	0.2-1.3	1.0	1.2	0.4-2.8	0.2	0.1	0.0-1.2
Lung and bronchus	1976-1980	92.0	61.2	55.6-67.1	68.2	108.3	96.7-120.8	23.8	29.3	24.1-35.2	75.6	49.9	44.9-55.2	58.0	91.6	81.0-103.1	17.6	21.1	16.8-26.1
	1981-1985	110.0	73.5	67.4-80.1	71.6	116.2	104.1-129.4	38.4	45.3	39.0-52.4	83.8	56.3	50.9-62.1	56.0	91.3	80.6-103.0	27.8	33.0	27.6-39.2
	1986-1990	120.4	79.8	73.5-86.5	73.8	118.0	105.9-131.0	46.6	54.3	47.4-62.0	103.2	68.2	62.4-74.5	64.6	105.6	94.0-118.1	38.6	44.1	38.0-51.0
	1991-1995	128.0	82.2	75.9-88.9	72.0	110.6	99.2-122.9	56.0	63.9	56.5-72.0	106.8	68.7	62.9-74.8	61.6	96.6	85.9-108.2	45.2	50.3	43.8-57.4
	1996-2000	145.8	91.2	84.7-98.1	78.8	115.2	104.0-127.2	67.0	73.9	66.2-82.4	106.6	66.5	60.9-72.4	62.6	92.4	82.4-103.4	44.0	47.9	41.7-54.7
	2001-2005	139.4	84.5	78.3-91.0	72.2	101.4	91.1-112.5	67.2	73.4	65.7-81.7	105.2	63.4	58.1-69.1	55.0	79.4	70.2-89.5	50.2	53.7	47.2-60.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Rensselaer**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	157.4	88.6	82.5-95.1	82.0	103.9	93.9-114.6	75.4	78.5	70.7-87.0	109.4	61.7	56.6-67.2	59.6	77.0	68.4-86.4	49.8	51.2	45.0-58.2
	2011-2015	166.0	86.1	80.2-92.3	79.8	92.1	83.0-102.0	86.2	82.0	74.2-90.3	98.6	51.3	46.8-56.2	49.0	57.9	50.6-65.9	49.6	47.1	41.3-53.6
	2016-2020	167.0	78.0	72.7-83.6	80.0	82.9	74.7-91.8	87.0	75.2	68.1-82.9	90.2	42.6	38.6-46.8	43.8	46.5	40.3-53.3	46.4	40.6	35.4-46.4
Melanoma of the skin	1976-1980	7.8	5.2	3.6-7.1	4.2	6.5	3.9-10.2	3.6	4.4	2.6-7.1	3.2	2.1	1.2-3.5	1.2	1.7	0.6-3.9	2.0	2.5	1.2-4.6
	1981-1985	10.6	7.6	5.6-10.0	5.2	8.7	5.5-13.0	5.4	7.3	4.8-10.8	2.8	2.0	1.1-3.4	1.6	2.4	1.0-4.9	1.2	1.7	0.6-3.8
	1986-1990	11.6	8.0	6.1-10.4	7.0	11.2	7.7-15.7	4.6	5.7	3.5-8.6	4.4	3.1	2.0-4.8	2.2	3.6	1.8-6.4	2.2	2.7	1.3-4.9
	1991-1995	9.8	6.1	4.5-8.1	6.4	9.0	6.1-12.8	3.4	3.8	2.2-6.2	5.8	3.8	2.5-5.5	3.6	5.1	3.0-8.1	2.2	2.3	1.1-4.2
	1996-2000	16.4	10.5	8.3-13.0	9.4	13.1	9.6-17.5	7.0	8.2	5.7-11.4	3.0	1.9	1.1-3.1	2.4	3.4	1.8-6.0	0.6	0.6	0.1-1.9
	2001-2005	27.4	16.7	14.0-19.8	15.0	20.6	16.1-25.9	12.4	14.5	11.1-18.7	7.4	4.5	3.1-6.2	5.4	7.4	4.9-10.8	2.0	2.3	1.1-4.4
	2006-2010	33.6	19.3	16.4-22.5	20.8	25.9	21.1-31.5	12.8	13.8	10.6-17.8	6.2	3.4	2.3-4.9	4.6	5.8	3.6-8.8	1.6	1.5	0.6-3.0
	2011-2015	44.4	23.6	20.5-27.0	27.2	31.9	26.6-38.0	17.2	17.9	14.2-22.3	5.8	3.1	2.1-4.5	4.2	5.3	3.3-8.2	1.6	1.7	0.7-3.5
	2016-2020	44.6	22.2	19.2-25.4	25.2	28.3	23.4-33.9	19.4	18.0	14.5-22.3	7.0	3.2	2.2-4.6	4.2	4.7	2.9-7.4	2.8	2.2	1.2-3.8
Female breast	1976-1980							70.2	88.3	79.0-98.4							28.2	34.7	29.0-41.0
	1981-1985							86.6	108.5	98.2-119.6							32.0	37.8	32.0-44.3
	1986-1990							90.2	109.0	98.9-119.9							35.2	41.3	35.3-48.1
	1991-1995							86.4	103.1	93.4-113.5							33.6	37.9	32.2-44.2
	1996-2000							124.4	144.8	133.6-156.8							30.2	32.9	27.8-38.6
	2001-2005							118.8	131.6	121.1-142.7							25.4	26.8	22.3-32.0
	2006-2010							124.2	131.4	121.1-142.4							20.2	19.7	16.0-24.1
	2011-2015							132.2	132.4	122.2-143.3							22.8	21.2	17.3-25.6
	2016-2020							133.8	127.8	117.9-138.4							22.6	20.5	16.7-24.9
Cervix uteri	1976-1980							11.2	14.7	11.0-19.2							5.0	6.2	3.9-9.3
	1981-1985							9.8	12.1	8.9-16.2							3.4	3.8	2.2-6.1
	1986-1990							6.4	8.2	5.5-11.6							2.8	3.4	1.8-5.8
	1991-1995							5.2	6.2	4.0-9.1							1.6	1.7	0.7-3.4
	1996-2000							6.0	7.4	5.0-10.6							0.8	1.0	0.3-2.6
	2001-2005							5.2	6.4	4.1-9.3							1.8	2.2	1.0-4.2
	2006-2010							6.4	7.6	5.1-10.8							2.0	2.0	0.9-3.7
	2011-2015							6.0	6.4	4.2-9.3							2.4	2.2	1.1-4.0
	2016-2020							5.0	6.4	4.1-9.6							1.0	1.0	0.3-2.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Rensselaer**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							18.8	22.2	17.9-27.3							5.2	6.1	3.9-9.0
	1981-1985							20.0	24.4	19.7-29.9							4.4	4.8	3.0-7.4
	1986-1990							19.4	23.1	18.6-28.4							3.4	3.5	2.0-5.7
	1991-1995							15.2	18.1	14.2-22.7							3.8	4.0	2.4-6.4
	1996-2000							23.0	26.8	22.1-32.2							3.6	3.5	2.1-5.7
	2001-2005							30.0	33.6	28.4-39.5							4.2	4.3	2.7-6.7
	2006-2010							33.0	34.5	29.4-40.4							5.4	5.4	3.5-8.0
	2011-2015							30.8	29.0	24.5-34.1							5.4	5.3	3.4-7.8
	2016-2020							35.2	31.1	26.5-36.3							4.0	3.4	2.0-5.4
Ovary	1976-1980							13.2	17.0	13.1-21.8							8.6	10.3	7.4-14.0
	1981-1985							13.2	15.0	11.5-19.3							7.8	9.3	6.5-12.8
	1986-1990							14.0	16.4	12.7-20.8							11.2	12.9	9.7-16.9
	1991-1995							12.0	14.4	11.0-18.7							8.0	9.2	6.5-12.6
	1996-2000							16.6	19.1	15.1-23.7							10.8	12.1	9.1-15.9
	2001-2005							15.4	16.7	13.2-21.0							7.8	7.9	5.6-10.9
	2006-2010							13.8	14.0	10.9-17.9							8.8	8.8	6.4-12.0
	2011-2015							11.6	11.3	8.5-14.8							7.2	6.4	4.4-9.1
	2016-2020							13.0	13.0	9.9-16.8							9.0	7.6	5.4-10.3
Prostate	1976-1980				38.2	70.7	60.6-81.9							13.6	27.6	21.2-35.2			
	1981-1985				55.2	98.8	87.1-111.6							20.6	40.3	32.5-49.1			
	1986-1990				63.4	111.8	99.4-125.2							21.8	42.8	34.9-51.8			
	1991-1995				75.0	117.2	105.3-130.0							19.0	36.9	29.6-45.2			
	1996-2000				117.2	172.4	158.6-187.0							17.0	29.9	23.8-37.1			
	2001-2005				133.0	185.3	171.4-200.2							19.0	31.7	25.6-38.8			
	2006-2010				140.6	167.9	155.5-181.1							14.2	20.3	15.8-25.7			
	2011-2015				109.8	116.4	106.6-127.0							15.0	20.2	15.8-25.4			
	2016-2020				124.0	118.2	108.8-128.2							10.2	12.3	9.1-16.3			
Testis	1976-1980				2.4	3.9	2.0-7.0							0.4	0.4	0.0-1.9			
	1981-1985				2.4	3.0	1.5-5.5							0.4	0.6	0.1-2.3			
	1986-1990				3.8	4.8	2.9-7.7							1.0	1.1	0.3-2.8			
	1991-1995				2.8	3.4	1.9-5.9							0.2	0.2	0.0-1.4			
	1996-2000				3.2	4.1	2.4-6.8							0.0	0.0	0.0-1.0			
	2001-2005				4.4	6.1	3.8-9.3							0.0	0.0	0.0-1.0			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Rensselaer**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				5.2	6.9	4.5-10.2							0.0	0.0	0.0-1.0			
	2011-2015				6.2	8.1	5.4-11.5							0.0	0.0	0.0-1.0			
	2016-2020				5.4	7.2	4.7-10.6							0.2	0.2	0.0-1.2			
Urinary bladder (incl. in situ)	1976-1980	27.4	19.1	16.0-22.6	19.4	34.5	27.6-42.4	8.0	9.5	6.8-13.0	8.6	6.5	4.7-8.7	5.2	10.1	6.5-15.0	3.4	4.1	2.4-6.5
	1981-1985	34.2	23.1	19.7-26.9	25.2	43.8	36.2-52.5	9.0	9.8	7.1-13.1	9.6	6.5	4.8-8.7	6.2	11.8	7.9-16.8	3.4	3.7	2.1-5.9
	1986-1990	36.2	23.8	20.4-27.6	26.4	43.7	36.3-52.1	9.8	10.8	7.9-14.3	10.0	6.6	4.9-8.7	6.8	12.6	8.6-17.6	3.2	3.3	1.9-5.5
	1991-1995	32.8	20.9	17.8-24.4	23.4	37.0	30.4-44.5	9.4	10.3	7.5-13.8	9.0	5.8	4.3-7.8	6.2	11.0	7.4-15.6	2.8	2.7	1.5-4.7
	1996-2000	51.0	31.6	27.8-35.7	37.2	56.4	48.5-65.3	13.8	14.5	11.3-18.5	9.4	5.8	4.2-7.7	6.4	10.4	7.1-14.7	3.0	2.9	1.6-4.9
	2001-2005	46.4	28.0	24.5-31.9	33.6	48.2	41.1-56.1	12.8	13.6	10.4-17.4	8.4	5.0	3.6-6.8	6.0	9.3	6.2-13.3	2.4	2.5	1.3-4.5
	2006-2010	54.8	30.7	27.2-34.7	38.0	50.0	43.0-57.8	16.8	17.5	13.9-21.7	9.2	5.2	3.8-7.0	7.0	9.7	6.7-13.5	2.2	2.2	1.1-4.1
	2011-2015	57.8	30.6	27.1-34.4	39.8	49.2	42.5-56.8	18.0	16.2	12.9-20.1	8.8	4.6	3.3-6.3	5.4	6.8	4.5-10.1	3.4	3.1	1.8-5.1
	2016-2020	60.0	28.3	25.1-31.7	42.6	45.2	39.2-52.0	17.4	14.7	11.7-18.3	8.4	3.9	2.8-5.3	6.0	7.0	4.7-10.1	2.4	1.8	0.9-3.4
Kidney and renal pelvis	1976-1980	10.8	7.6	5.6-9.9	6.6	11.2	7.6-15.9	4.2	5.3	3.2-8.2	6.2	4.4	3.0-6.3	3.8	6.8	4.0-10.7	2.4	3.0	1.5-5.2
	1981-1985	15.2	10.2	8.0-12.8	8.6	13.7	9.8-18.7	6.6	7.5	5.1-10.6	6.8	4.4	3.0-6.2	3.6	6.1	3.5-9.7	3.2	3.3	1.9-5.4
	1986-1990	15.4	10.5	8.3-13.2	8.4	14.5	10.3-19.7	7.0	8.3	5.7-11.6	8.4	5.5	3.9-7.4	4.4	7.0	4.3-10.6	4.0	4.7	2.8-7.3
	1991-1995	15.8	10.7	8.4-13.3	10.2	16.3	12.1-21.5	5.6	6.8	4.5-10.0	7.2	4.9	3.4-6.8	4.4	7.2	4.4-10.9	2.8	3.6	2.0-6.1
	1996-2000	17.6	11.2	9.0-13.8	10.0	14.1	10.5-18.6	7.6	8.9	6.3-12.2	7.6	4.8	3.4-6.6	4.4	6.3	3.9-9.5	3.2	3.4	2.0-5.7
	2001-2005	26.2	15.9	13.3-18.9	17.6	24.2	19.3-29.8	8.6	9.7	7.0-13.2	8.0	4.9	3.5-6.7	6.2	8.9	6.0-12.7	1.8	2.1	0.9-4.0
	2006-2010	39.8	22.2	19.2-25.6	23.2	28.9	23.8-34.8	16.6	17.1	13.6-21.4	9.0	5.2	3.8-7.1	5.2	7.2	4.7-10.6	3.8	4.0	2.4-6.4
	2011-2015	39.4	20.8	17.9-24.1	23.6	27.7	22.8-33.4	15.8	15.0	11.8-19.0	8.8	4.5	3.2-6.1	4.4	5.0	3.1-7.7	4.4	4.1	2.5-6.4
	2016-2020	43.6	21.2	18.4-24.4	29.8	31.1	26.2-36.8	13.8	13.0	10.0-16.7	9.4	4.5	3.3-6.0	6.6	7.5	5.1-10.6	2.8	2.2	1.2-3.9
Brain and other central nervous system	1976-1980	8.2	5.4	3.9-7.4	4.8	7.2	4.6-10.8	3.4	3.8	2.2-6.2	4.8	3.2	2.0-4.8	2.6	4.1	2.2-7.1	2.2	2.5	1.2-4.5
	1981-1985	9.0	6.2	4.5-8.3	3.8	5.7	3.4-8.9	5.2	6.2	4.0-9.2	5.6	3.6	2.4-5.2	3.2	4.6	2.6-7.6	2.4	2.7	1.4-4.9
	1986-1990	10.2	7.0	5.2-9.3	4.0	5.9	3.6-9.1	6.2	7.8	5.3-11.2	6.8	4.6	3.1-6.4	2.6	4.0	2.1-7.0	4.2	5.1	3.1-7.8
	1991-1995	8.8	5.6	4.0-7.5	4.2	6.0	3.6-9.2	4.6	5.6	3.5-8.5	4.6	2.9	1.9-4.4	2.6	3.6	1.9-6.2	2.0	2.2	1.1-4.1
	1996-2000	13.0	8.3	6.4-10.6	6.0	8.7	5.9-12.5	7.0	8.1	5.6-11.3	6.4	4.1	2.8-5.8	3.2	5.0	2.8-8.1	3.2	3.7	2.1-6.0
	2001-2005	14.4	9.1	7.1-11.4	7.0	9.5	6.6-13.3	7.4	8.8	6.2-12.2	6.6	4.2	2.9-5.9	4.0	5.5	3.3-8.5	2.6	3.1	1.6-5.3
	2006-2010	13.4	8.2	6.3-10.4	7.8	9.7	6.9-13.4	5.6	6.8	4.5-9.9	7.2	4.1	2.9-5.8	3.8	4.3	2.5-6.8	3.4	3.9	2.2-6.3
	2011-2015	13.0	7.2	5.5-9.2	7.0	7.7	5.3-10.9	6.0	6.5	4.3-9.5	7.8	4.4	3.1-6.0	3.8	4.7	2.7-7.4	4.0	4.1	2.4-6.4
	2016-2020	12.2	6.8	5.1-8.8	6.8	7.3	5.0-10.4	5.4	6.4	4.1-9.5	9.0	4.5	3.2-6.1	6.0	6.5	4.3-9.5	3.0	2.8	1.5-4.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Rensselaer**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	5.8	3.8	2.5-5.6	1.8	2.9	1.3-5.6	4.0	4.6	2.8-7.2	1.0	0.7	0.2-1.6	0.2	0.2	0.0-1.5	0.8	0.9	0.2-2.4
	1981-1985	5.4	3.6	2.4-5.3	2.0	3.3	1.5-6.1	3.4	4.1	2.4-6.7	0.6	0.4	0.1-1.2	0.4	0.6	0.1-2.2	0.2	0.2	0.0-1.4
	1986-1990	5.0	3.3	2.1-4.9	1.4	2.2	0.9-4.6	3.6	4.2	2.4-6.7	0.4	0.3	0.0-1.0	0.0	0.0	0.0-1.1	0.4	0.5	0.1-1.8
	1991-1995	6.6	4.3	3.0-6.1	2.8	3.8	2.1-6.5	3.8	4.7	2.8-7.4	1.2	0.8	0.3-1.7	0.4	0.6	0.1-2.2	0.8	0.9	0.2-2.4
	1996-2000	11.2	7.2	5.5-9.4	2.6	3.6	1.9-6.2	8.6	10.8	7.8-14.5	1.6	1.0	0.4-2.0	0.6	0.9	0.2-2.5	1.0	1.1	0.3-2.6
	2001-2005	13.4	8.3	6.4-10.5	3.0	3.8	2.1-6.4	10.4	12.5	9.3-16.4	0.4	0.2	0.0-0.9	0.0	0.0	0.0-1.0	0.4	0.3	0.0-1.4
	2006-2010	26.8	16.1	13.5-19.2	7.0	8.4	5.8-11.9	19.8	23.5	19.0-28.8	0.8	0.5	0.1-1.2	0.6	0.7	0.1-2.2	0.2	0.2	0.0-1.4
	2011-2015	23.6	13.4	11.0-16.2	6.4	7.0	4.7-10.0	17.2	19.9	15.8-24.8	1.0	0.5	0.2-1.3	0.6	0.9	0.2-2.5	0.4	0.4	0.0-1.5
2016-2020	25.2	15.0	12.4-17.9	8.0	9.2	6.4-12.7	17.2	20.6	16.4-25.7	0.4	0.2	0.0-0.8	0.0	0.0	0.0-1.0	0.4	0.4	0.0-1.5	
Hodgkin lymphoma	1976-1980	5.4	3.2	2.1-4.8	3.0	3.7	2.0-6.4	2.4	2.7	1.4-4.9	2.4	1.7	0.9-3.0	0.8	1.1	0.3-3.0	1.6	2.1	0.9-4.2
	1981-1985	7.8	4.7	3.3-6.5	3.8	4.8	2.8-7.8	4.0	4.7	2.8-7.3	1.6	0.9	0.4-1.9	0.8	1.1	0.3-3.0	0.8	0.8	0.2-2.2
	1986-1990	4.8	2.8	1.8-4.2	2.2	2.5	1.2-4.6	2.6	2.9	1.6-5.1	0.8	0.5	0.1-1.3	0.2	0.3	0.0-1.8	0.6	0.6	0.1-1.9
	1991-1995	4.0	2.4	1.5-3.7	1.4	1.8	0.7-3.8	2.6	3.0	1.6-5.2	0.8	0.5	0.1-1.3	0.6	0.9	0.2-2.6	0.2	0.3	0.0-1.5
	1996-2000	4.6	3.0	1.9-4.5	2.0	2.8	1.3-5.3	2.6	3.4	1.8-5.8	0.4	0.3	0.0-0.9	0.2	0.3	0.0-1.5	0.2	0.3	0.0-1.4
	2001-2005	5.2	3.4	2.2-5.0	2.6	3.6	1.9-6.1	2.6	3.5	1.8-5.9	1.4	0.8	0.3-1.7	0.6	0.8	0.2-2.3	0.8	0.8	0.2-2.1
	2006-2010	6.8	4.1	2.8-5.7	3.2	3.9	2.2-6.4	3.6	4.2	2.5-6.7	2.2	1.2	0.6-2.2	1.0	1.2	0.4-2.9	1.2	1.2	0.4-2.8
	2011-2015	6.0	3.9	2.6-5.7	4.0	5.0	3.0-7.8	2.0	2.9	1.4-5.3	0.4	0.3	0.0-0.9	0.4	0.6	0.1-2.1	0.0	0.0	0.0-0.9
2016-2020	5.4	3.1	2.0-4.6	3.4	3.9	2.2-6.5	2.0	2.3	1.1-4.4	0.4	0.2	0.0-0.8	0.2	0.3	0.0-1.5	0.2	0.2	0.0-1.2	
Non-Hodgkin lymphomas	1976-1980	16.6	11.4	9.1-14.2	7.2	11.6	8.0-16.4	9.4	11.3	8.3-15.2	7.4	5.2	3.6-7.2	4.0	6.7	4.0-10.4	3.4	4.1	2.4-6.7
	1981-1985	17.4	12.1	9.7-15.0	8.4	14.1	10.1-19.2	9.0	10.5	7.6-14.2	9.0	5.9	4.3-7.9	4.0	5.9	3.5-9.2	5.0	5.5	3.6-8.3
	1986-1990	18.2	12.1	9.7-14.9	9.8	14.7	10.8-19.6	8.4	9.5	6.8-12.9	9.6	6.3	4.6-8.3	5.8	9.3	6.1-13.4	3.8	4.0	2.4-6.3
	1991-1995	19.4	12.6	10.2-15.4	11.4	17.0	12.8-22.2	8.0	9.0	6.4-12.4	10.2	6.6	4.9-8.7	5.8	9.0	6.0-13.0	4.4	4.8	3.0-7.3
	1996-2000	33.0	20.8	17.7-24.2	17.2	25.4	20.3-31.5	15.8	17.7	14.0-22.1	14.6	9.1	7.2-11.5	6.8	10.5	7.2-14.7	7.8	8.5	6.0-11.6
	2001-2005	31.2	19.0	16.1-22.2	15.6	21.9	17.2-27.4	15.6	17.7	13.9-22.1	10.4	6.2	4.7-8.2	4.8	7.1	4.5-10.5	5.6	5.8	3.8-8.4
	2006-2010	46.4	26.3	23.0-30.0	25.2	33.3	27.6-39.8	21.2	22.1	18.0-26.8	12.8	7.1	5.5-9.2	7.2	10.4	7.3-14.4	5.6	5.4	3.5-8.0
	2011-2015	42.4	22.3	19.3-25.6	23.0	26.2	21.5-31.7	19.4	19.0	15.3-23.4	11.4	5.9	4.4-7.7	6.8	8.5	5.8-12.0	4.6	4.2	2.6-6.4
2016-2020	44.0	22.2	19.2-25.4	23.4	25.8	21.1-31.1	20.6	18.9	15.3-23.2	11.4	6.0	4.5-7.8	5.6	7.0	4.6-10.2	5.8	5.3	3.5-7.8	
Myeloma	1976-1980	5.2	3.6	2.3-5.3	1.8	3.0	1.3-5.9	3.4	3.8	2.2-6.2	3.6	2.6	1.5-4.1	0.8	1.8	0.4-4.5	2.8	3.1	1.7-5.3
	1981-1985	8.0	5.3	3.8-7.3	3.4	5.7	3.3-9.3	4.6	5.0	3.1-7.5	4.2	2.8	1.7-4.2	1.4	2.2	0.8-4.7	2.8	3.0	1.6-5.1
	1986-1990	7.0	4.6	3.2-6.4	2.6	4.0	2.1-7.0	4.4	4.7	3.0-7.2	6.0	3.8	2.6-5.5	2.6	4.0	2.1-6.8	3.4	3.7	2.1-5.9
	1991-1995	6.6	4.3	3.0-6.1	2.2	3.7	1.8-6.6	4.4	5.0	3.1-7.7	5.6	3.6	2.4-5.2	1.4	2.4	0.9-4.9	4.2	4.5	2.8-7.0
	1996-2000	8.6	5.4	3.9-7.2	4.8	7.1	4.5-10.5	3.8	4.0	2.4-6.4	5.2	3.2	2.1-4.7	2.4	3.8	1.9-6.7	2.8	3.0	1.6-5.1
	2001-2005	10.2	6.0	4.5-8.0	5.4	7.7	5.1-11.3	4.8	4.9	3.1-7.3	5.6	3.3	2.2-4.9	3.0	4.4	2.4-7.2	2.6	2.6	1.4-4.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Rensselaer**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	13.0	7.1	5.5-9.2	7.0	8.6	6.0-12.1	6.0	6.0	4.0-8.7	5.2	2.8	1.8-4.2	3.0	3.7	2.1-6.2	2.2	2.0	1.0-3.7
	2011-2015	13.2	7.1	5.5-9.1	7.0	8.3	5.7-11.7	6.2	6.0	4.0-8.6	6.6	3.3	2.2-4.6	3.2	3.5	2.0-5.8	3.4	3.1	1.8-5.1
	2016-2020	14.2	6.6	5.1-8.4	8.4	8.4	5.9-11.5	5.8	5.0	3.3-7.3	5.4	2.5	1.6-3.7	1.8	1.8	0.8-3.6	3.6	2.9	1.7-4.7
Leukemias	1976-1980	22.6	15.7	12.9-18.9	13.4	22.1	16.8-28.6	9.2	11.6	8.4-15.6	12.2	8.4	6.4-10.9	7.0	12.4	8.4-17.6	5.2	6.1	4.0-9.0
	1981-1985	17.0	11.8	9.4-14.6	7.4	12.9	8.9-18.1	9.6	10.7	7.9-14.3	11.6	8.2	6.2-10.7	5.0	9.3	5.8-13.9	6.6	8.0	5.4-11.4
	1986-1990	16.6	10.9	8.6-13.5	10.2	16.5	12.1-22.0	6.4	7.4	5.0-10.5	11.0	7.4	5.5-9.6	6.4	11.2	7.5-15.9	4.6	5.3	3.3-8.0
	1991-1995	16.4	10.6	8.4-13.1	8.8	13.0	9.4-17.5	7.6	8.8	6.2-12.1	9.4	6.1	4.5-8.1	5.4	8.0	5.2-11.6	4.0	4.3	2.6-6.8
	1996-2000	20.2	12.7	10.3-15.4	11.0	16.0	12.0-20.9	9.2	10.1	7.4-13.6	10.4	6.5	4.9-8.6	4.8	7.2	4.6-10.7	5.6	6.1	4.0-8.9
	2001-2005	26.6	16.2	13.6-19.2	14.8	20.8	16.2-26.1	11.8	13.1	9.9-16.9	13.4	8.1	6.3-10.3	6.8	10.4	7.1-14.5	6.6	7.0	4.8-9.9
	2006-2010	30.2	17.3	14.6-20.3	17.4	22.7	18.1-28.1	12.8	13.5	10.3-17.3	14.2	8.1	6.3-10.3	9.2	12.3	9.0-16.5	5.0	5.1	3.2-7.7
	2011-2015	31.4	16.6	14.0-19.5	20.2	23.0	18.6-28.2	11.2	11.5	8.6-15.1	13.6	7.1	5.5-9.1	8.6	10.5	7.5-14.3	5.0	4.7	3.0-7.1
	2016-2020	36.4	18.2	15.6-21.2	23.0	24.9	20.4-30.2	13.4	12.9	9.8-16.6	10.0	4.8	3.6-6.4	8.0	9.0	6.4-12.4	2.0	1.7	0.8-3.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Rockland**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	780.8	406.8	393.8-420.1	370.4	482.8	459.3-507.1	410.4	371.6	355.6-388.1	398.0	215.3	205.7-225.2	203.2	280.7	262.4-299.9	194.8	179.8	168.6-191.5
	1981-1985	883.4	413.6	401.3-426.2	421.4	493.2	470.6-516.5	462.0	377.2	361.9-392.9	460.0	222.7	213.6-232.2	228.6	280.8	263.5-298.9	231.4	191.3	180.4-202.6
	1986-1990	995.2	430.9	418.7-443.2	453.0	489.5	467.9-511.7	542.2	409.3	394.0-425.1	486.2	218.5	209.8-227.5	240.0	276.1	259.5-293.4	246.2	187.5	177.2-198.4
	1991-1995	1,286.6	510.5	498.0-523.3	634.0	604.3	582.1-627.1	652.6	459.7	444.0-475.8	520.0	215.6	207.3-224.2	253.6	270.0	254.3-286.3	266.4	187.3	177.3-197.7
	1996-2000	1,453.8	529.6	517.4-542.0	739.8	624.1	603.3-645.4	714.0	465.6	450.4-481.2	504.6	189.2	181.8-196.8	247.0	229.8	216.5-243.6	257.6	164.9	155.9-174.2
	2001-2005	1,569.8	520.6	509.1-532.3	807.2	611.8	592.6-631.5	762.6	458.3	443.7-473.2	489.4	166.2	159.6-172.9	237.2	198.4	186.9-210.4	252.2	147.8	139.7-156.3
	2006-2010	1,683.8	512.4	501.4-523.5	867.6	594.0	576.1-612.3	816.2	455.1	441.0-469.5	461.2	140.9	135.2-146.9	219.0	159.7	150.2-169.6	242.2	128.9	121.6-136.5
	2011-2015	1,758.4	487.6	477.3-498.1	881.4	540.3	524.2-556.8	877.0	453.4	439.7-467.5	468.6	127.6	122.4-132.9	220.2	141.4	133.1-150.1	248.4	119.0	112.3-126.0
2016-2020	1,725.8	457.5	447.6-467.5	875.2	503.2	488.1-518.6	850.6	426.8	413.6-440.4	429.4	108.0	103.4-112.8	210.6	122.7	115.3-130.5	218.8	98.0	92.1-104.2	
Oral cavity and pharynx	1976-1980	17.4	8.6	6.9-10.6	10.0	11.8	8.5-15.8	7.4	6.5	4.6-9.0	7.2	3.6	2.5-5.0	4.4	5.5	3.3-8.6	2.8	2.5	1.4-4.2
	1981-1985	20.2	8.9	7.2-10.8	12.8	13.5	10.1-17.5	7.4	5.8	4.1-8.0	7.2	3.3	2.3-4.6	4.4	4.9	3.0-7.6	2.8	2.2	1.2-3.8
	1986-1990	17.2	7.1	5.7-8.8	10.4	9.5	6.9-12.6	6.8	5.1	3.5-7.1	5.6	2.4	1.6-3.4	4.6	4.6	2.8-7.0	1.0	0.7	0.2-1.7
	1991-1995	25.8	10.1	8.4-12.0	16.8	15.9	12.5-19.9	9.0	6.3	4.6-8.4	6.0	2.3	1.6-3.4	3.6	3.3	1.9-5.4	2.4	1.7	0.9-3.0
	1996-2000	22.8	8.3	6.8-9.9	15.4	12.7	9.9-16.0	7.4	4.9	3.5-6.8	6.6	2.5	1.7-3.5	3.8	3.4	2.0-5.3	2.8	1.8	1.0-3.1
	2001-2005	25.6	8.5	7.1-10.1	16.6	12.4	9.8-15.5	9.0	5.3	3.8-7.1	4.6	1.5	0.9-2.3	3.0	2.2	1.2-3.6	1.6	0.9	0.4-1.8
	2006-2010	24.8	7.4	6.1-8.8	16.4	10.4	8.2-13.0	8.4	4.8	3.5-6.6	4.4	1.3	0.8-2.0	3.0	1.8	1.0-3.1	1.4	0.8	0.3-1.7
	2011-2015	38.8	10.6	9.2-12.3	24.2	14.3	11.8-17.1	14.6	7.5	5.8-9.5	4.6	1.2	0.8-1.9	3.2	1.8	1.0-3.0	1.4	0.6	0.2-1.4
2016-2020	38.8	10.1	8.7-11.7	27.6	15.5	12.9-18.3	11.2	5.6	4.2-7.4	7.4	1.8	1.2-2.5	4.2	2.2	1.4-3.5	3.2	1.4	0.8-2.3	
Esophagus	1976-1980	7.2	3.9	2.7-5.4	3.8	5.3	3.1-8.4	3.4	3.1	1.8-5.0	6.2	3.3	2.2-4.7	4.2	5.3	3.2-8.2	2.0	1.9	0.9-3.4
	1981-1985	9.6	4.6	3.4-6.2	6.4	7.9	5.2-11.3	3.2	2.7	1.5-4.3	9.6	4.7	3.4-6.2	7.4	9.0	6.1-12.6	2.2	1.9	0.9-3.3
	1986-1990	11.0	4.8	3.6-6.2	7.8	7.7	5.4-10.8	3.2	2.5	1.4-4.1	9.8	4.5	3.3-5.9	6.8	7.5	5.1-10.7	3.0	2.4	1.3-4.0
	1991-1995	14.4	5.8	4.5-7.3	9.6	8.9	6.5-12.0	4.8	3.4	2.2-5.1	10.8	4.4	3.3-5.8	6.2	6.0	4.0-8.8	4.6	3.3	2.1-4.9
	1996-2000	12.4	4.5	3.4-5.8	9.6	8.2	5.9-10.9	2.8	1.8	1.0-3.1	13.0	4.9	3.7-6.2	10.2	8.9	6.5-11.7	2.8	1.9	1.0-3.1
	2001-2005	16.2	5.3	4.2-6.6	13.0	9.7	7.4-12.4	3.2	1.9	1.1-3.1	12.4	4.2	3.2-5.4	8.8	7.1	5.1-9.6	3.6	2.1	1.2-3.3
	2006-2010	16.8	5.0	4.0-6.2	12.0	8.3	6.3-10.8	4.8	2.4	1.6-3.7	10.6	3.2	2.4-4.2	8.6	6.0	4.3-8.1	2.0	1.1	0.5-2.0
	2011-2015	15.2	4.2	3.3-5.3	11.0	6.9	5.2-9.1	4.2	2.1	1.3-3.2	12.0	3.3	2.5-4.3	10.0	6.5	4.8-8.6	2.0	0.9	0.4-1.8
2016-2020	16.8	4.2	3.4-5.3	12.6	7.1	5.4-9.1	4.2	1.9	1.2-3.0	6.2	1.5	1.0-2.1	5.4	3.0	1.9-4.3	0.8	0.4	0.1-1.1	
Stomach	1976-1980	23.8	13.6	11.3-16.4	11.8	16.2	12.1-21.1	12.0	11.6	8.8-14.9	16.6	9.8	7.7-12.1	8.4	11.6	8.2-15.8	8.2	8.1	5.8-10.9
	1981-1985	27.0	13.6	11.4-16.2	16.4	22.2	17.3-27.9	10.6	8.8	6.6-11.5	18.8	9.5	7.6-11.6	10.2	13.7	9.9-18.3	8.6	7.2	5.2-9.7
	1986-1990	25.0	11.6	9.6-13.8	15.2	18.3	14.1-23.1	9.8	7.5	5.5-9.9	16.4	7.7	6.1-9.6	11.0	13.9	10.2-18.3	5.4	4.2	2.8-6.1
	1991-1995	29.8	12.7	10.7-14.9	18.0	18.8	14.9-23.3	11.8	8.4	6.4-10.8	21.6	9.3	7.6-11.3	12.4	13.5	10.1-17.5	9.2	6.6	4.8-8.8
	1996-2000	28.8	10.9	9.2-12.8	17.0	15.3	12.1-19.1	11.8	7.4	5.6-9.6	15.6	5.9	4.6-7.3	9.8	8.8	6.4-11.8	5.8	3.6	2.4-5.1
	2001-2005	32.4	11.0	9.4-12.8	19.2	15.5	12.5-19.1	13.2	7.8	6.0-9.9	16.6	5.6	4.5-7.0	10.6	9.7	7.2-12.8	6.0	3.3	2.2-4.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Rockland**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	28.0	8.5	7.1-10.0	17.6	12.4	9.9-15.4	10.4	5.6	4.2-7.4	12.8	3.8	2.9-4.9	7.2	5.1	3.5-7.1	5.6	2.9	1.9-4.3
	2011-2015	28.4	7.9	6.6-9.3	16.6	10.5	8.4-13.1	11.8	5.9	4.5-7.7	10.2	2.8	2.1-3.7	5.4	3.4	2.3-5.0	4.8	2.3	1.5-3.5
	2016-2020	26.8	7.2	6.0-8.5	14.0	8.0	6.2-10.2	12.8	6.5	4.9-8.4	11.0	3.0	2.2-3.9	5.4	3.4	2.2-4.9	5.6	2.7	1.7-4.0
Colorectal	1976-1980	127.6	70.5	65.1-76.3	64.2	88.4	78.3-99.4	63.4	59.6	53.2-66.5	60.6	33.7	29.9-37.7	29.2	42.1	35.0-50.0	31.4	29.4	25.0-34.4
	1981-1985	127.4	63.8	58.9-69.1	65.8	83.2	73.8-93.4	61.6	51.7	46.1-57.8	63.0	32.0	28.5-35.8	33.4	42.4	35.8-49.8	29.6	24.9	21.1-29.3
	1986-1990	138.6	63.5	58.8-68.5	69.2	79.2	70.4-88.6	69.4	53.3	47.8-59.2	65.6	30.9	27.6-34.5	33.0	39.6	33.4-46.6	32.6	25.2	21.5-29.4
	1991-1995	152.6	62.7	58.2-67.3	75.4	75.8	67.9-84.4	77.2	54.1	48.8-59.8	63.4	26.8	23.9-30.0	29.8	33.2	27.7-39.3	33.6	23.7	20.2-27.6
	1996-2000	173.0	65.4	61.1-69.9	83.4	76.2	68.7-84.2	89.6	57.7	52.5-63.4	62.2	23.8	21.2-26.7	31.8	30.7	25.9-36.0	30.4	19.2	16.3-22.6
	2001-2005	161.0	54.3	50.6-58.2	81.6	65.6	59.2-72.5	79.4	46.6	42.1-51.5	49.2	16.8	14.7-19.0	26.0	22.2	18.4-26.5	23.2	13.1	10.8-15.8
	2006-2010	138.4	42.4	39.3-45.8	67.6	49.0	43.9-54.6	70.8	38.2	34.3-42.5	39.4	12.1	10.5-14.0	20.8	15.3	12.5-18.6	18.6	9.6	7.7-11.8
	2011-2015	129.8	36.2	33.4-39.1	70.2	44.4	39.8-49.4	59.6	29.5	26.1-33.1	43.0	11.8	10.3-13.6	22.6	14.6	12.0-17.6	20.4	9.8	8.0-12.0
2016-2020	141.4	37.3	34.5-40.2	76.0	44.1	39.7-48.9	65.4	31.8	28.3-35.6	39.8	9.8	8.5-11.3	19.6	11.1	9.0-13.6	20.2	8.7	7.0-10.6	
Colon excluding rectum	1976-1980	90.8	50.6	46.0-55.6	45.0	63.9	55.2-73.5	45.8	43.0	37.6-48.9	51.4	28.6	25.2-32.4	24.0	35.3	28.8-42.6	27.4	25.6	21.5-30.2
	1981-1985	92.6	47.2	42.9-51.8	45.6	60.0	51.9-68.8	47.0	39.6	34.7-45.0	56.4	28.7	25.4-32.3	30.2	38.7	32.3-45.8	26.2	22.1	18.5-26.2
	1986-1990	102.4	47.1	43.1-51.5	50.6	58.7	51.1-66.9	51.8	39.8	35.1-45.0	53.6	25.2	22.2-28.4	25.8	30.4	25.0-36.5	27.8	21.6	18.1-25.5
	1991-1995	113.4	47.0	43.2-51.1	52.8	54.2	47.5-61.6	60.6	42.6	37.9-47.7	56.8	23.9	21.2-26.9	26.8	29.5	24.4-35.3	30.0	21.1	17.9-24.8
	1996-2000	132.4	50.1	46.3-54.1	59.8	55.9	49.4-62.9	72.6	46.4	41.7-51.5	55.6	21.4	18.9-24.0	29.0	28.2	23.6-33.4	26.6	16.8	14.1-20.0
	2001-2005	119.2	40.4	37.2-43.8	60.0	49.0	43.4-55.0	59.2	35.0	31.1-39.3	42.6	14.5	12.6-16.6	22.2	19.0	15.5-23.0	20.4	11.4	9.3-13.9
	2006-2010	103.8	31.8	29.1-34.7	48.8	35.3	31.0-40.1	55.0	29.5	26.1-33.3	33.2	10.2	8.7-11.9	17.0	12.5	10.0-15.5	16.2	8.3	6.6-10.4
	2011-2015	96.2	26.4	24.1-29.0	48.0	30.4	26.6-34.6	48.2	23.4	20.5-26.6	36.4	9.9	8.5-11.5	18.2	11.6	9.3-14.3	18.2	8.7	7.0-10.8
2016-2020	105.6	27.8	25.4-30.3	55.0	32.0	28.3-36.1	50.6	24.5	21.5-27.9	35.0	8.6	7.3-10.0	17.2	9.7	7.7-12.0	17.8	7.6	6.1-9.4	
Rectum & rectosigmoid	1976-1980	36.8	19.9	17.1-23.1	19.2	24.5	19.5-30.3	17.6	16.6	13.3-20.4	9.2	5.0	3.7-6.7	5.2	6.8	4.3-10.2	4.0	3.9	2.4-5.9
	1981-1985	34.8	16.6	14.2-19.3	20.2	23.2	18.6-28.6	14.6	12.1	9.5-15.2	6.6	3.4	2.3-4.7	3.2	3.7	2.1-6.1	3.4	2.8	1.6-4.5
	1986-1990	36.2	16.3	14.0-18.9	18.6	20.5	16.3-25.4	17.6	13.5	10.8-16.6	12.0	5.7	4.3-7.4	7.2	9.2	6.3-12.9	4.8	3.6	2.3-5.4
	1991-1995	39.2	15.7	13.5-18.0	22.6	21.6	17.6-26.2	16.6	11.5	9.2-14.3	6.6	2.9	2.0-4.1	3.0	3.7	1.9-6.1	3.6	2.6	1.5-4.1
	1996-2000	40.6	15.3	13.2-17.6	23.6	20.3	16.6-24.4	17.0	11.3	9.0-14.0	6.6	2.5	1.7-3.5	2.8	2.4	1.3-4.1	3.8	2.4	1.4-3.8
	2001-2005	41.8	13.9	12.1-15.9	21.6	16.6	13.6-20.2	20.2	11.6	9.5-14.2	6.6	2.3	1.6-3.2	3.8	3.2	1.9-4.9	2.8	1.7	0.9-2.9
	2006-2010	34.6	10.6	9.1-12.4	18.8	13.7	11.0-16.8	15.8	8.7	6.8-10.9	6.2	1.9	1.3-2.8	3.8	2.8	1.7-4.4	2.4	1.3	0.7-2.3
	2011-2015	33.6	9.7	8.3-11.3	22.2	14.0	11.5-16.9	11.4	6.1	4.6-8.0	6.6	1.9	1.3-2.8	4.4	3.0	1.8-4.5	2.2	1.1	0.6-2.1
2016-2020	35.8	9.5	8.1-11.0	21.0	12.1	9.8-14.7	14.8	7.3	5.6-9.2	4.8	1.2	0.8-1.9	2.4	1.4	0.7-2.5	2.4	1.1	0.5-1.9	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Rockland**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	5.4	3.1	2.0-4.5	3.0	4.0	2.1-6.7	2.4	2.3	1.2-4.0	3.6	2.0	1.2-3.2	2.0	2.6	1.2-4.8	1.6	1.5	0.7-3.0
	1981-1985	4.0	1.9	1.2-3.0	2.4	3.0	1.4-5.4	1.6	1.3	0.6-2.6	5.2	2.6	1.7-3.8	3.4	4.5	2.5-7.3	1.8	1.5	0.7-2.9
	1986-1990	7.4	3.0	2.1-4.1	5.2	4.5	2.9-6.8	2.2	1.6	0.8-2.9	7.4	3.0	2.1-4.2	5.6	5.6	3.6-8.2	1.8	1.3	0.6-2.5
	1991-1995	9.0	3.6	2.6-4.9	6.2	6.3	4.2-9.1	2.8	2.0	1.1-3.3	9.4	3.8	2.7-5.0	6.0	6.3	4.1-9.1	3.4	2.3	1.3-3.7
	1996-2000	17.0	6.4	5.1-8.0	11.6	10.2	7.7-13.3	5.4	3.5	2.3-5.1	10.6	3.9	2.9-5.2	7.8	6.6	4.6-9.1	2.8	1.9	1.0-3.1
	2001-2005	18.0	6.0	4.8-7.3	12.4	9.4	7.1-12.1	5.6	3.4	2.2-4.9	11.6	3.9	3.0-5.1	7.4	6.1	4.3-8.5	4.2	2.5	1.5-3.8
	2006-2010	20.0	6.0	4.9-7.3	13.4	9.1	7.0-11.6	6.6	3.7	2.5-5.3	12.8	3.9	3.0-4.9	7.6	5.4	3.8-7.4	5.2	2.7	1.7-4.0
	2011-2015	27.2	7.3	6.1-8.7	19.0	11.3	9.1-13.9	8.2	3.9	2.8-5.3	16.2	4.4	3.5-5.5	9.2	5.7	4.2-7.7	7.0	3.4	2.4-4.8
2016-2020	26.0	6.6	5.5-7.8	17.6	9.8	7.8-12.1	8.4	3.9	2.8-5.3	17.6	4.4	3.6-5.5	11.2	6.3	4.7-8.2	6.4	3.0	2.0-4.3	
Pancreas	1976-1980	26.0	14.4	12.0-17.1	13.6	18.2	13.9-23.5	12.4	11.7	8.9-14.9	25.0	14.0	11.6-16.7	12.6	17.3	13.0-22.5	12.4	11.7	9.0-15.0
	1981-1985	22.4	10.7	8.8-12.9	10.2	12.1	8.8-16.2	12.2	10.0	7.6-12.8	22.2	10.8	8.9-13.0	11.0	13.6	10.0-18.1	11.2	9.3	7.0-12.1
	1986-1990	29.2	13.4	11.3-15.8	14.0	15.5	11.8-19.8	15.2	11.7	9.2-14.6	27.8	12.9	10.8-15.2	12.8	14.6	11.0-18.9	15.0	11.6	9.1-14.5
	1991-1995	32.8	13.8	11.8-16.1	16.4	17.2	13.5-21.6	16.4	11.5	9.1-14.3	28.4	12.1	10.2-14.3	14.8	15.8	12.2-20.1	13.6	9.5	7.4-12.1
	1996-2000	35.6	13.4	11.5-15.6	16.8	15.3	12.1-19.1	18.8	12.3	9.9-15.0	30.2	11.6	9.8-13.7	14.4	14.3	11.0-18.2	15.8	10.2	8.1-12.7
	2001-2005	40.6	13.7	11.9-15.7	21.4	16.6	13.5-20.2	19.2	11.5	9.3-14.1	34.8	11.8	10.1-13.7	17.2	13.5	10.7-16.7	17.6	10.4	8.4-12.9
	2006-2010	43.2	13.4	11.7-15.3	20.8	15.0	12.2-18.2	22.4	12.0	9.8-14.4	36.6	11.1	9.6-12.9	17.4	12.4	9.9-15.3	19.2	10.1	8.1-12.3
	2011-2015	50.4	13.6	12.0-15.5	24.4	15.3	12.6-18.3	26.0	12.5	10.4-14.9	38.4	10.5	9.0-12.1	18.4	12.0	9.6-14.7	20.0	9.3	7.6-11.4
2016-2020	57.0	14.5	12.9-16.3	27.0	15.7	13.2-18.7	30.0	13.7	11.6-16.2	37.2	9.3	8.0-10.7	18.4	10.7	8.6-13.2	18.8	8.2	6.6-10.1	
Larynx	1976-1980	8.2	4.2	3.0-5.7	6.6	8.6	5.7-12.2	1.6	1.4	0.6-2.8	3.2	1.7	1.0-2.8	3.2	4.4	2.4-7.3	0.0	0.0	0.0-0.6
	1981-1985	12.2	5.2	4.0-6.7	10.6	9.9	7.3-13.1	1.6	1.3	0.6-2.5	2.4	1.1	0.5-1.9	1.8	1.8	0.7-3.6	0.6	0.5	0.1-1.5
	1986-1990	12.0	4.9	3.7-6.3	9.2	8.2	5.9-11.1	2.8	2.1	1.2-3.6	3.8	1.5	0.9-2.4	2.8	2.5	1.3-4.4	1.0	0.8	0.2-1.8
	1991-1995	13.4	5.3	4.1-6.7	10.2	9.9	7.2-13.2	3.2	2.3	1.3-3.7	3.6	1.5	0.9-2.4	3.0	3.3	1.8-5.6	0.6	0.4	0.1-1.2
	1996-2000	11.2	4.1	3.1-5.3	9.6	8.2	5.9-11.0	1.6	1.1	0.5-2.1	3.0	1.1	0.6-1.9	2.8	2.6	1.4-4.3	0.2	0.1	0.0-0.7
	2001-2005	10.6	3.4	2.6-4.5	8.4	6.3	4.5-8.5	2.2	1.3	0.7-2.4	2.4	0.8	0.4-1.4	2.0	1.5	0.7-2.8	0.4	0.2	0.0-0.9
	2006-2010	9.8	3.0	2.2-4.0	8.8	6.1	4.4-8.2	1.0	0.5	0.2-1.3	3.2	0.9	0.5-1.5	2.6	1.7	0.9-3.0	0.6	0.4	0.1-1.0
	2011-2015	10.0	2.8	2.1-3.7	8.6	5.3	3.8-7.2	1.4	0.8	0.3-1.6	2.8	0.8	0.4-1.3	2.8	1.8	1.0-3.1	0.0	0.0	0.0-0.5
2016-2020	8.6	2.3	1.6-3.1	7.4	4.3	3.0-6.0	1.2	0.6	0.2-1.3	1.6	0.4	0.2-0.8	1.4	0.8	0.3-1.7	0.2	0.1	0.0-0.6	
Lung and bronchus	1976-1980	103.8	53.7	49.1-58.6	70.6	90.7	80.8-101.2	33.2	30.0	25.6-35.0	85.0	44.8	40.6-49.4	60.2	81.1	71.7-91.4	24.8	22.4	18.6-26.7
	1981-1985	136.0	63.2	58.4-68.2	84.0	93.8	84.4-103.8	52.0	43.0	38.0-48.6	115.0	54.7	50.2-59.4	73.0	85.1	76.0-95.0	42.0	35.0	30.4-40.1
	1986-1990	138.2	59.8	55.3-64.5	78.2	82.3	73.8-91.6	60.0	45.6	40.6-51.1	109.2	47.8	43.8-52.0	65.8	71.4	63.3-80.1	43.4	33.2	28.9-38.0
	1991-1995	159.8	64.2	59.8-68.9	88.4	85.2	77.0-94.0	71.4	50.5	45.4-56.1	122.2	50.2	46.2-54.3	66.6	68.8	61.2-77.1	55.6	39.3	34.8-44.2
	1996-2000	167.4	61.2	57.1-65.6	84.0	73.0	65.9-80.7	83.4	54.1	49.0-59.6	118.2	44.0	40.5-47.8	63.8	56.9	50.6-63.7	54.4	35.0	31.0-39.5
	2001-2005	176.0	58.7	54.9-62.8	91.0	72.5	65.8-79.7	85.0	50.0	45.3-55.0	118.0	39.9	36.7-43.3	62.0	50.8	45.1-56.9	56.0	32.9	29.2-37.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Rockland**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	188.2	57.3	53.6-61.1	91.8	65.3	59.3-71.6	96.4	51.9	47.3-56.8	117.2	35.9	33.0-39.0	59.0	42.7	37.9-47.9	58.2	31.3	27.8-35.2
	2011-2015	183.0	50.0	46.8-53.4	81.8	51.7	46.7-57.0	101.2	49.4	45.1-54.0	102.2	27.7	25.3-30.3	47.4	30.2	26.5-34.4	54.8	26.4	23.3-29.8
	2016-2020	189.4	48.2	45.1-51.4	87.2	50.8	46.0-55.8	102.2	46.2	42.2-50.5	88.2	22.1	20.1-24.3	44.0	25.9	22.5-29.6	44.2	19.4	16.9-22.2
Melanoma of the skin	1976-1980	22.8	10.4	8.6-12.6	11.2	11.5	8.5-15.2	11.6	9.8	7.4-12.6	6.2	3.0	2.0-4.2	3.2	3.7	2.0-6.1	3.0	2.6	1.4-4.2
	1981-1985	18.6	7.8	6.3-9.6	7.8	7.3	5.1-10.2	10.8	8.6	6.4-11.2	5.2	2.3	1.5-3.5	2.4	2.9	1.3-5.2	2.8	2.3	1.2-3.8
	1986-1990	23.2	9.3	7.7-11.2	13.6	12.0	9.2-15.5	9.6	7.1	5.2-9.4	7.6	3.1	2.2-4.3	4.8	4.5	2.8-6.8	2.8	2.1	1.1-3.5
	1991-1995	29.0	11.1	9.4-13.1	16.2	13.9	11.0-17.4	12.8	9.0	6.9-11.5	8.0	3.1	2.2-4.2	5.6	4.8	3.1-7.0	2.4	1.6	0.8-2.8
	1996-2000	37.2	13.3	11.5-15.4	20.0	16.2	13.1-19.9	17.2	11.4	9.1-14.1	7.8	2.9	2.1-4.0	3.4	2.8	1.6-4.6	4.4	2.8	1.8-4.3
	2001-2005	50.2	16.8	14.8-19.0	32.2	23.4	19.8-27.4	18.0	11.4	9.1-14.0	8.0	2.7	1.9-3.6	4.6	3.6	2.2-5.4	3.4	2.0	1.1-3.2
	2006-2010	63.4	19.7	17.6-22.0	34.4	23.9	20.4-27.9	29.0	16.9	14.2-20.0	8.0	2.5	1.8-3.4	6.2	4.2	2.9-6.1	1.8	1.1	0.5-2.1
	2011-2015	61.6	17.3	15.4-19.4	37.6	23.0	19.8-26.6	24.0	12.9	10.6-15.6	9.0	2.4	1.7-3.2	5.0	3.1	2.0-4.6	4.0	1.7	1.0-2.8
	2016-2020	70.8	19.1	17.1-21.2	43.2	26.2	22.7-30.0	27.6	13.7	11.5-16.4	5.6	1.5	1.0-2.2	3.0	1.7	1.0-2.9	2.6	1.2	0.6-2.1
Female breast	1976-1980							120.6	108.6	100.0-117.6							43.6	40.1	34.9-45.8
	1981-1985							135.8	110.4	102.2-119.1							49.4	40.3	35.5-45.7
	1986-1990							177.8	133.6	124.9-142.7							57.0	43.0	38.1-48.3
	1991-1995							219.0	153.9	144.9-163.3							52.0	36.5	32.1-41.2
	1996-2000							229.2	149.5	140.9-158.4							49.0	31.4	27.6-35.6
	2001-2005							224.0	135.1	127.3-143.3							38.8	22.8	19.7-26.3
	2006-2010							242.4	136.0	128.3-144.0							35.8	19.0	16.3-22.1
	2011-2015							270.2	143.2	135.4-151.3							37.6	18.3	15.7-21.3
	2016-2020							250.6	132.0	124.5-139.9							36.2	17.4	14.9-20.3
Cervix uteri	1976-1980							11.0	9.7	7.3-12.6							4.0	3.6	2.2-5.5
	1981-1985							9.4	7.5	5.5-10.0							3.6	2.9	1.7-4.5
	1986-1990							10.8	7.9	5.9-10.3							3.8	2.9	1.7-4.5
	1991-1995							16.0	11.3	8.9-14.0							3.8	2.6	1.6-4.1
	1996-2000							13.0	8.7	6.7-11.1							4.6	3.0	1.9-4.5
	2001-2005							11.2	7.4	5.6-9.6							2.2	1.3	0.6-2.4
	2006-2010							12.4	7.6	5.8-9.8							3.2	1.8	1.0-3.0
	2011-2015							15.8	9.2	7.2-11.5							3.8	2.0	1.2-3.2
	2016-2020							10.0	6.0	4.3-8.0							2.2	1.1	0.5-2.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Rockland**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							22.6	20.4	16.8-24.6							3.8	3.6	2.2-5.6
	1981-1985							28.4	23.1	19.4-27.2							7.0	5.8	4.1-8.1
	1986-1990							30.2	22.5	19.0-26.4							5.2	3.9	2.5-5.7
	1991-1995							33.4	23.6	20.1-27.4							6.8	4.8	3.4-6.8
	1996-2000							41.0	26.5	23.0-30.5							7.6	4.9	3.4-6.7
	2001-2005							48.4	27.9	24.5-31.7							7.4	4.2	3.0-5.9
	2006-2010							57.6	31.2	27.6-35.0							9.0	4.8	3.5-6.5
	2011-2015							62.2	30.9	27.5-34.6							10.6	5.3	4.0-7.0
	2016-2020							61.8	29.2	25.9-32.7							8.0	3.5	2.5-4.9
Ovary	1976-1980							17.2	15.1	12.1-18.6							11.8	10.8	8.2-13.9
	1981-1985							17.6	14.1	11.3-17.4							10.8	8.7	6.6-11.4
	1986-1990							25.4	19.4	16.2-23.1							17.4	13.2	10.5-16.3
	1991-1995							24.6	17.3	14.4-20.7							15.0	10.5	8.2-13.2
	1996-2000							25.8	16.7	14.0-19.9							14.4	9.3	7.3-11.7
	2001-2005							24.0	14.4	11.9-17.3							16.8	9.8	7.8-12.2
	2006-2010							24.6	13.7	11.4-16.5							16.0	8.5	6.7-10.6
	2011-2015							21.6	11.1	9.1-13.5							12.2	5.9	4.5-7.6
	2016-2020							25.0	12.9	10.6-15.4							12.2	5.7	4.3-7.4
Prostate	1976-1980				50.0	79.5	69.4-90.4							16.0	28.1	22.0-35.2			
	1981-1985				62.2	87.4	77.4-98.2							16.8	26.2	20.6-32.7			
	1986-1990				75.4	92.6	82.8-103.1							26.0	36.6	30.3-43.7			
	1991-1995				166.6	156.6	145.5-168.3							26.8	33.9	28.1-40.4			
	1996-2000				250.2	205.0	193.4-217.1							23.6	25.9	21.3-31.3			
	2001-2005				258.0	187.4	177.1-198.1							22.2	21.9	17.9-26.4			
	2006-2010				312.6	201.9	191.9-212.4							18.6	14.7	11.9-18.1			
	2011-2015				262.8	153.1	144.8-161.8							16.4	11.1	8.8-13.8			
	2016-2020				248.2	133.5	126.1-141.3							21.8	12.9	10.5-15.6			
Testis	1976-1980				8.0	6.7	4.8-9.3							0.8	0.6	0.2-1.8			
	1981-1985				6.4	4.7	3.2-6.7							0.2	0.1	0.0-1.0			
	1986-1990				5.6	4.2	2.8-6.1							0.4	0.3	0.0-1.2			
	1991-1995				7.6	5.8	4.1-8.1							0.8	0.6	0.2-1.5			
	1996-2000				9.2	6.9	5.0-9.2							0.4	0.3	0.0-1.1			
	2001-2005				9.0	6.7	4.9-9.0							1.0	0.8	0.3-1.8			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Rockland**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				10.2	7.6	5.6-10.0							0.0	0.0	0.0-0.5			
	2011-2015				8.8	6.2	4.4-8.3							0.2	0.1	0.0-0.7			
	2016-2020				10.6	7.5	5.6-9.8							0.2	0.1	0.0-0.7			
Urinary bladder (incl. in situ)	1976-1980	37.8	20.4	17.6-23.6	26.4	35.2	29.1-42.2	11.4	10.7	8.1-13.8	10.8	6.6	5.0-8.6	9.0	14.8	10.6-19.9	1.8	1.9	0.9-3.5
	1981-1985	41.8	19.9	17.2-22.8	29.6	35.1	29.2-41.7	12.2	10.0	7.7-12.9	10.4	5.4	4.0-7.1	6.0	7.9	5.2-11.5	4.4	3.7	2.3-5.6
	1986-1990	50.6	22.7	20.0-25.8	36.8	42.1	35.8-49.2	13.8	10.7	8.3-13.5	13.0	6.5	5.0-8.3	7.6	10.8	7.5-14.9	5.4	4.2	2.8-6.2
	1991-1995	56.6	22.8	20.2-25.7	40.8	40.4	34.7-46.7	15.8	11.1	8.8-13.8	8.8	3.7	2.7-5.0	4.6	5.3	3.2-8.1	4.2	2.8	1.7-4.3
	1996-2000	71.8	26.4	23.8-29.4	52.4	46.0	40.4-52.2	19.4	12.5	10.2-15.3	8.2	3.2	2.3-4.4	5.6	6.0	3.9-8.7	2.6	1.7	0.9-3.0
	2001-2005	88.2	29.6	26.9-32.5	63.0	50.9	45.2-57.0	25.2	14.6	12.2-17.4	10.4	3.6	2.7-4.8	7.8	7.4	5.2-10.2	2.6	1.5	0.8-2.6
	2006-2010	75.8	23.0	20.7-25.5	56.4	41.2	36.4-46.3	19.4	10.1	8.2-12.4	10.6	3.2	2.4-4.2	7.6	6.0	4.2-8.3	3.0	1.5	0.8-2.5
	2011-2015	94.6	25.4	23.1-27.8	71.0	45.2	40.6-50.2	23.6	11.1	9.1-13.3	13.6	3.6	2.8-4.6	8.4	5.6	4.0-7.6	5.2	2.3	1.5-3.4
2016-2020	84.6	21.5	19.5-23.7	63.4	36.9	32.9-41.2	21.2	9.5	7.7-11.5	11.4	2.7	2.0-3.5	8.0	4.7	3.3-6.4	3.4	1.4	0.8-2.2	
Kidney and renal pelvis	1976-1980	13.0	6.5	5.0-8.3	8.6	9.2	6.6-12.6	4.4	4.0	2.5-6.1	8.6	4.4	3.2-6.0	6.0	6.6	4.3-9.6	2.6	2.4	1.3-4.2
	1981-1985	23.6	10.5	8.7-12.6	17.4	17.6	13.8-22.0	6.2	5.1	3.5-7.2	9.6	4.6	3.3-6.1	6.4	7.0	4.6-10.1	3.2	2.7	1.5-4.3
	1986-1990	23.4	10.1	8.3-12.1	13.4	13.2	10.1-17.0	10.0	7.5	5.6-9.9	9.4	4.3	3.1-5.7	6.0	6.6	4.3-9.5	3.4	2.5	1.5-4.1
	1991-1995	33.0	13.2	11.2-15.4	20.6	18.3	14.8-22.4	12.4	8.9	6.8-11.4	11.0	4.6	3.5-6.1	6.6	6.5	4.4-9.2	4.4	3.0	1.9-4.6
	1996-2000	41.4	14.8	12.9-17.0	25.2	20.2	16.7-24.2	16.2	10.6	8.4-13.2	8.0	2.9	2.1-4.0	5.4	4.5	2.9-6.7	2.6	1.7	0.9-2.9
	2001-2005	49.4	16.0	14.1-18.2	32.8	23.3	19.8-27.3	16.6	9.8	7.8-12.1	10.6	3.6	2.7-4.7	5.6	4.2	2.8-6.2	5.0	2.9	1.9-4.4
	2006-2010	54.6	16.6	14.7-18.7	34.8	23.6	20.2-27.5	19.8	11.3	9.1-13.8	10.8	3.3	2.4-4.3	7.0	5.1	3.5-7.1	3.8	2.0	1.2-3.2
	2011-2015	69.2	19.1	17.1-21.2	47.2	28.2	24.6-32.1	22.0	11.4	9.4-13.9	9.6	2.7	2.0-3.6	5.4	3.5	2.3-5.1	4.2	2.0	1.2-3.2
2016-2020	65.4	17.3	15.4-19.4	43.8	25.4	22.1-29.1	21.6	10.6	8.7-13.0	10.0	2.5	1.8-3.3	7.2	4.1	2.9-5.8	2.8	1.3	0.7-2.2	
Brain and other central nervous system	1976-1980	15.0	6.6	5.1-8.3	7.4	7.0	4.8-9.8	7.6	6.3	4.4-8.6	10.4	4.9	3.6-6.4	4.8	5.3	3.3-8.0	5.6	4.7	3.1-6.9
	1981-1985	15.8	6.5	5.1-8.1	8.2	7.2	5.0-9.9	7.6	5.9	4.2-8.1	9.0	4.0	2.9-5.3	4.4	4.2	2.5-6.5	4.6	3.7	2.4-5.6
	1986-1990	19.4	7.7	6.2-9.4	11.8	9.8	7.4-12.8	7.6	5.7	4.0-7.8	10.0	4.1	3.1-5.5	6.8	6.0	4.1-8.6	3.2	2.5	1.4-4.0
	1991-1995	17.8	6.7	5.4-8.3	10.0	8.2	6.0-10.9	7.8	5.5	3.9-7.5	11.4	4.5	3.4-5.8	6.0	5.4	3.5-7.8	5.4	3.8	2.5-5.5
	1996-2000	23.6	8.5	7.0-10.2	11.6	8.8	6.7-11.5	12.0	7.9	6.0-10.2	12.4	4.4	3.4-5.6	6.6	5.1	3.4-7.2	5.8	3.8	2.5-5.5
	2001-2005	22.4	7.5	6.2-9.0	13.2	10.0	7.7-12.8	9.2	5.7	4.1-7.6	12.2	4.1	3.2-5.3	7.4	5.5	3.9-7.6	4.8	2.9	1.9-4.4
	2006-2010	19.8	6.1	4.9-7.4	10.2	6.8	5.1-9.0	9.6	5.6	4.1-7.4	11.2	3.6	2.7-4.7	6.0	4.3	2.9-6.2	5.2	3.0	2.0-4.5
	2011-2015	20.2	5.9	4.8-7.2	11.6	7.0	5.3-9.1	8.6	4.8	3.5-6.6	9.8	2.8	2.1-3.8	5.4	3.2	2.1-4.7	4.4	2.4	1.5-3.7
2016-2020	24.4	6.7	5.6-8.1	14.2	8.3	6.4-10.5	10.2	5.2	3.9-7.0	12.8	3.3	2.5-4.3	7.0	4.0	2.8-5.6	5.8	2.7	1.8-3.9	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Rockland**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	9.6	4.3	3.1-5.7	3.0	3.2	1.7-5.5	6.6	5.3	3.6-7.5	1.2	0.7	0.3-1.6	0.4	0.8	0.1-2.7	0.8	0.8	0.2-1.9
	1981-1985	13.4	5.6	4.3-7.1	3.4	3.5	1.9-5.8	10.0	7.7	5.7-10.2	1.4	0.7	0.3-1.4	0.8	0.9	0.2-2.3	0.6	0.5	0.1-1.5
	1986-1990	18.0	6.8	5.5-8.4	5.0	4.0	2.6-5.9	13.0	9.4	7.2-11.9	1.2	0.5	0.2-1.2	0.4	0.5	0.1-1.7	0.8	0.6	0.2-1.5
	1991-1995	23.0	8.6	7.1-10.3	4.8	4.0	2.5-6.0	18.2	13.0	10.4-15.9	2.6	1.0	0.6-1.8	1.0	1.0	0.3-2.4	1.6	1.1	0.5-2.1
	1996-2000	30.8	10.9	9.3-12.8	8.6	6.3	4.5-8.5	22.2	15.1	12.4-18.2	1.4	0.5	0.2-1.1	0.6	0.5	0.1-1.5	0.8	0.5	0.1-1.3
	2001-2005	54.2	18.4	16.3-20.8	13.4	9.6	7.4-12.2	40.8	26.7	23.1-30.7	1.8	0.6	0.3-1.2	0.8	0.6	0.2-1.5	1.0	0.6	0.2-1.4
	2006-2010	64.6	21.4	19.1-23.9	15.8	11.1	8.7-13.9	48.8	31.1	27.2-35.4	3.0	0.9	0.5-1.5	1.2	0.9	0.3-1.9	1.8	0.9	0.4-1.7
	2011-2015	71.4	22.0	19.7-24.4	20.8	13.3	10.8-16.3	50.6	30.2	26.4-34.3	2.6	0.7	0.4-1.2	1.4	0.8	0.3-1.7	1.2	0.5	0.2-1.2
2016-2020	65.0	20.9	18.6-23.4	18.2	11.8	9.5-14.6	46.8	29.7	25.9-34.0	2.0	0.5	0.2-1.0	0.6	0.3	0.1-1.0	1.4	0.7	0.3-1.5	
Hodgkin lymphoma	1976-1980	10.4	4.4	3.3-5.8	5.8	5.0	3.2-7.3	4.6	3.7	2.3-5.7	3.8	1.8	1.1-2.9	1.6	1.5	0.6-3.1	2.2	2.0	1.0-3.6
	1981-1985	10.4	4.0	3.0-5.3	6.2	5.3	3.5-7.6	4.2	3.0	1.9-4.7	2.6	1.2	0.6-2.0	1.6	1.7	0.7-3.5	1.0	0.8	0.3-1.9
	1986-1990	8.0	3.0	2.1-4.1	4.4	3.6	2.2-5.7	3.6	2.6	1.5-4.1	2.4	1.0	0.5-1.7	1.8	1.8	0.7-3.5	0.6	0.4	0.1-1.2
	1991-1995	11.0	4.2	3.2-5.5	6.8	5.7	3.8-8.0	4.2	3.0	1.9-4.6	1.8	0.8	0.3-1.4	1.2	1.2	0.4-2.7	0.6	0.4	0.1-1.3
	1996-2000	11.8	4.3	3.3-5.6	6.0	4.6	3.1-6.6	5.8	4.1	2.8-5.9	0.2	0.1	0.0-0.4	0.0	0.0	0.0-0.6	0.2	0.1	0.0-0.7
	2001-2005	13.4	4.9	3.8-6.2	5.6	4.2	2.8-6.0	7.8	5.6	3.9-7.6	2.4	0.8	0.4-1.5	0.8	0.6	0.1-1.5	1.6	1.1	0.5-2.1
	2006-2010	9.0	3.0	2.2-4.1	5.6	3.9	2.6-5.7	3.4	2.2	1.3-3.5	1.6	0.5	0.2-1.0	0.6	0.4	0.1-1.2	1.0	0.6	0.2-1.5
	2011-2015	9.8	3.0	2.2-4.0	5.2	3.3	2.2-4.9	4.6	3.0	1.9-4.5	0.6	0.2	0.0-0.5	0.2	0.1	0.0-0.7	0.4	0.2	0.0-0.8
2016-2020	9.4	2.9	2.1-3.9	5.4	3.4	2.2-5.0	4.0	2.5	1.5-3.9	2.0	0.6	0.3-1.1	1.8	1.1	0.5-2.2	0.2	0.1	0.0-0.8	
Non-Hodgkin lymphomas	1976-1980	27.4	13.9	11.6-16.5	14.6	17.5	13.4-22.4	12.8	11.5	8.8-14.6	14.0	7.5	5.8-9.5	7.2	9.4	6.4-13.3	6.8	6.2	4.3-8.7
	1981-1985	31.2	14.3	12.1-16.8	16.4	17.0	13.3-21.4	14.8	12.1	9.5-15.2	18.0	8.7	7.0-10.8	10.2	12.3	8.9-16.5	7.8	6.5	4.6-8.9
	1986-1990	37.4	15.9	13.7-18.4	20.0	19.2	15.4-23.7	17.4	13.2	10.6-16.3	17.2	7.6	6.0-9.4	7.8	7.6	5.3-10.6	9.4	7.2	5.3-9.6
	1991-1995	63.6	25.2	22.5-28.2	37.2	34.3	29.3-39.9	26.4	18.4	15.4-21.9	26.8	11.2	9.3-13.3	15.2	15.5	12.0-19.6	11.6	8.2	6.2-10.6
	1996-2000	61.4	22.5	20.0-25.1	36.8	29.9	25.6-34.7	24.6	16.3	13.6-19.5	26.8	10.0	8.4-11.9	15.6	14.1	11.0-17.7	11.2	7.0	5.3-9.1
	2001-2005	68.2	22.8	20.5-25.4	36.4	27.9	23.9-32.4	31.8	18.8	16.0-22.0	21.4	7.2	5.9-8.8	12.0	10.4	7.8-13.5	9.4	5.5	4.0-7.3
	2006-2010	83.0	25.7	23.2-28.3	43.4	30.1	26.2-34.5	39.6	22.0	19.0-25.4	19.0	5.9	4.8-7.2	10.0	7.1	5.3-9.5	9.0	4.8	3.5-6.4
	2011-2015	86.2	24.3	22.0-26.8	45.6	29.3	25.6-33.5	40.6	20.4	17.6-23.6	19.4	5.2	4.2-6.4	10.6	6.7	5.0-8.8	8.8	4.0	2.9-5.4
2016-2020	87.4	23.9	21.6-26.3	46.8	28.7	25.1-32.8	40.6	20.1	17.3-23.2	14.0	3.5	2.7-4.5	6.8	4.1	2.8-5.7	7.2	3.1	2.2-4.4	
Myeloma	1976-1980	9.6	5.1	3.8-6.8	5.2	7.0	4.4-10.4	4.4	4.0	2.5-6.1	5.0	2.9	1.8-4.2	2.4	3.8	1.9-6.7	2.6	2.4	1.3-4.1
	1981-1985	9.0	4.3	3.1-5.7	5.2	6.5	4.1-9.7	3.8	3.1	1.9-4.9	5.4	2.6	1.7-3.8	3.8	5.0	2.8-7.9	1.6	1.3	0.6-2.6
	1986-1990	10.2	4.5	3.3-5.9	6.0	6.4	4.2-9.3	4.2	3.2	1.9-4.8	8.2	3.7	2.7-5.1	4.4	5.2	3.1-8.0	3.8	2.9	1.7-4.5
	1991-1995	17.6	7.4	5.9-9.1	9.8	10.6	7.6-14.2	7.8	5.7	4.1-7.8	12.6	5.3	4.0-6.8	7.4	7.6	5.2-10.7	5.2	3.8	2.5-5.6
	1996-2000	18.2	6.8	5.5-8.4	6.8	6.0	4.1-8.5	11.4	7.2	5.4-9.4	12.8	4.9	3.8-6.3	6.0	6.4	4.2-9.2	6.8	4.4	3.0-6.1
	2001-2005	20.6	6.9	5.6-8.4	10.6	8.1	6.0-10.6	10.0	6.0	4.4-7.9	8.4	2.8	2.0-3.8	3.0	2.5	1.4-4.2	5.4	3.1	2.0-4.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Rockland**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	23.4	7.1	5.8-8.5	11.6	7.9	5.9-10.2	11.8	6.5	4.9-8.4	7.8	2.5	1.8-3.4	4.6	3.4	2.1-5.0	3.2	1.9	1.1-3.0
	2011-2015	35.2	9.3	8.0-10.8	18.4	11.0	8.8-13.5	16.8	8.0	6.4-10.0	10.8	2.9	2.2-3.8	4.6	3.0	1.9-4.5	6.2	2.9	1.9-4.1
	2016-2020	26.6	6.7	5.6-8.0	14.8	8.3	6.5-10.5	11.8	5.6	4.2-7.3	8.8	2.1	1.5-2.9	4.4	2.4	1.5-3.7	4.4	1.9	1.1-2.9
Leukemias	1976-1980	21.6	10.8	8.8-13.1	14.0	17.4	13.2-22.4	7.6	6.7	4.7-9.2	14.8	7.5	5.9-9.5	9.2	11.4	8.1-15.5	5.6	5.0	3.3-7.2
	1981-1985	27.4	13.0	10.9-15.4	15.4	17.8	13.7-22.7	12.0	9.9	7.5-12.7	18.2	8.7	7.0-10.7	9.0	11.0	7.8-15.0	9.2	7.5	5.5-10.0
	1986-1990	29.2	12.5	10.5-14.8	15.6	17.5	13.6-22.2	13.6	10.1	7.8-12.8	22.2	9.9	8.1-12.0	13.2	15.8	12.0-20.4	9.0	6.7	4.9-9.0
	1991-1995	30.2	12.0	10.2-14.1	16.6	15.7	12.3-19.7	13.6	9.6	7.4-12.2	17.2	7.1	5.6-8.7	9.8	10.4	7.5-13.9	7.4	5.3	3.7-7.3
	1996-2000	36.8	13.5	11.6-15.6	22.0	18.3	14.9-22.2	14.8	9.8	7.7-12.3	18.6	7.0	5.6-8.6	9.8	8.9	6.5-11.9	8.8	5.5	4.0-7.5
	2001-2005	45.8	15.2	13.3-17.3	25.8	18.7	15.6-22.3	20.0	12.2	9.9-14.9	23.6	8.2	6.8-9.8	12.4	10.1	7.7-13.0	11.2	6.7	5.0-8.7
	2006-2010	59.0	18.3	16.2-20.5	34.2	24.3	20.8-28.3	24.8	13.6	11.3-16.3	19.8	6.2	5.1-7.6	10.0	7.6	5.6-10.0	9.8	5.3	3.9-7.0
	2011-2015	66.4	18.4	16.5-20.6	38.8	24.1	20.8-27.8	27.6	14.0	11.7-16.6	24.8	6.7	5.5-8.0	13.2	8.3	6.4-10.6	11.6	5.6	4.2-7.3
	2016-2020	63.6	17.0	15.2-19.1	38.8	23.2	20.0-26.8	24.8	11.9	9.9-14.3	21.4	5.5	4.5-6.7	13.8	8.4	6.5-10.7	7.6	3.4	2.4-4.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**St. Lawrence**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	347.4	363.7	346.5-381.5	177.6	439.9	410.0-471.3	169.8	319.1	297.7-341.8	190.4	200.4	187.7-213.8	99.4	251.5	228.8-275.8	91.0	167.3	152.0-183.6
	1981-1985	389.4	394.9	377.3-413.2	196.0	468.5	438.3-500.1	193.4	350.1	327.9-373.4	210.4	213.8	200.9-227.4	111.4	277.5	253.7-302.7	99.0	176.1	160.7-192.6
	1986-1990	430.8	424.9	406.9-443.5	224.6	527.2	495.8-560.1	206.2	363.1	340.8-386.4	227.4	223.6	210.6-237.1	123.4	290.9	267.5-315.6	104.0	177.7	162.6-194.0
	1991-1995	493.8	460.6	442.5-479.3	248.4	541.7	511.1-573.7	245.4	413.9	390.7-438.0	252.0	234.6	221.8-248.1	131.8	295.9	273.0-320.2	120.2	194.1	178.7-210.4
	1996-2000	574.8	514.7	496.0-533.9	282.6	568.2	538.5-599.2	292.2	480.3	455.8-505.9	242.6	217.2	205.1-229.8	130.8	278.5	257.0-301.3	111.8	174.9	160.6-190.2
	2001-2005	643.6	547.2	528.4-566.5	343.8	660.7	629.2-693.3	299.8	472.8	448.9-497.6	248.6	211.2	199.6-223.3	130.0	263.0	242.7-284.5	118.6	178.8	164.6-194.0
	2006-2010	633.0	511.2	493.4-529.6	337.6	591.9	563.4-621.5	295.4	453.2	429.9-477.4	241.0	196.3	185.2-207.8	128.0	235.3	217.0-254.6	113.0	166.1	152.5-180.6
	2011-2015	644.6	485.3	468.4-502.8	354.2	557.1	530.8-584.5	290.4	427.5	405.1-450.9	227.6	170.4	160.5-180.8	115.6	191.4	175.7-208.1	112.0	155.3	142.4-169.0
	2016-2020	655.4	465.0	448.6-481.8	353.2	516.7	492.2-542.3	302.2	425.3	403.0-448.6	223.0	154.6	145.5-164.2	119.2	180.6	166.0-196.2	103.8	133.4	121.8-145.9
Oral cavity and pharynx	1976-1980	7.0	7.0	4.9-9.8	5.0	10.8	6.9-16.1	2.0	3.7	1.8-6.9	3.0	3.3	1.8-5.4	2.0	5.6	2.5-10.5	1.0	1.8	0.6-4.3
	1981-1985	10.2	10.4	7.7-13.7	7.2	16.8	11.6-23.6	3.0	5.5	3.0-9.2	2.2	2.3	1.1-4.0	1.4	3.2	1.3-6.7	0.8	1.4	0.4-3.7
	1986-1990	7.8	7.9	5.5-10.8	5.8	13.8	9.1-19.9	2.0	3.4	1.6-6.2	3.2	3.2	1.8-5.2	1.8	4.0	1.8-7.7	1.4	2.5	1.0-5.1
	1991-1995	10.4	9.8	7.3-12.9	7.4	15.7	11.0-21.7	3.0	5.1	2.8-8.4	2.8	2.5	1.4-4.2	1.8	3.8	1.7-7.2	1.0	1.5	0.5-3.6
	1996-2000	11.6	10.6	8.0-13.7	6.8	14.3	9.8-20.1	4.8	7.9	5.0-11.8	2.4	2.1	1.1-3.7	1.8	3.5	1.6-6.7	0.6	1.0	0.2-3.0
	2001-2005	10.2	8.6	6.4-11.3	7.8	13.8	9.8-18.9	2.4	3.9	2.0-6.9	2.6	2.2	1.2-3.7	1.6	2.8	1.2-5.6	1.0	1.5	0.5-3.6
	2006-2010	15.2	12.0	9.4-15.1	11.6	19.0	14.4-24.8	3.6	5.4	3.2-8.6	4.4	3.4	2.1-5.1	3.4	5.7	3.3-9.3	1.0	1.4	0.5-3.5
	2011-2015	13.6	9.6	7.4-12.2	10.0	14.5	10.7-19.4	3.6	4.7	2.8-7.7	1.6	1.2	0.5-2.4	1.2	2.0	0.7-4.4	0.4	0.5	0.1-2.1
	2016-2020	21.2	14.1	11.5-17.1	15.2	21.6	16.9-27.3	6.0	7.2	4.8-10.5	3.6	2.3	1.4-3.8	2.6	3.7	1.9-6.5	1.0	1.0	0.3-2.8
Esophagus	1976-1980	3.8	3.8	2.3-6.0	2.8	6.2	3.3-10.5	1.0	1.8	0.6-4.2	4.0	4.1	2.5-6.3	3.2	7.2	4.1-11.9	0.8	1.5	0.4-3.7
	1981-1985	3.0	3.1	1.7-5.2	2.0	5.2	2.4-9.5	1.0	1.9	0.6-4.5	2.8	3.1	1.7-5.1	1.6	4.8	2.0-9.3	1.2	2.2	0.8-4.9
	1986-1990	4.8	4.8	3.1-7.2	3.8	10.0	5.9-15.7	1.0	1.7	0.5-4.0	3.6	3.6	2.1-5.7	3.4	8.6	4.9-14.0	0.2	0.3	0.0-1.9
	1991-1995	6.8	6.3	4.4-8.9	5.6	12.0	7.9-17.5	1.2	1.9	0.7-4.2	6.0	5.6	3.7-8.0	4.2	8.3	5.1-12.7	1.8	2.9	1.3-5.6
	1996-2000	7.6	6.8	4.8-9.3	6.2	13.0	8.7-18.6	1.4	2.1	0.8-4.4	6.2	5.6	3.8-7.9	5.0	10.9	7.0-16.3	1.2	1.8	0.7-4.0
	2001-2005	10.4	8.7	6.5-11.5	8.0	14.9	10.6-20.4	2.4	3.6	1.8-6.3	7.6	6.5	4.6-8.9	5.8	11.3	7.5-16.3	1.8	2.6	1.2-5.1
	2006-2010	8.4	6.6	4.8-9.0	5.8	10.2	6.7-14.7	2.6	3.8	2.0-6.6	8.2	6.6	4.7-9.0	6.0	10.0	6.7-14.3	2.2	3.4	1.7-6.2
	2011-2015	9.4	6.6	4.9-8.9	7.2	10.6	7.4-14.9	2.2	2.9	1.4-5.4	7.4	5.4	3.8-7.5	6.2	9.9	6.7-14.3	1.2	1.6	0.6-3.8
	2016-2020	8.0	5.7	4.0-7.9	6.8	10.2	6.9-14.4	1.2	1.5	0.6-3.7	9.2	6.5	4.7-8.8	8.4	12.4	8.8-16.9	0.8	1.2	0.3-3.2
Stomach	1976-1980	9.8	10.4	7.7-13.8	6.4	16.7	11.1-23.8	3.4	6.2	3.6-10.0	8.4	9.1	6.5-12.3	4.6	11.8	7.3-18.0	3.8	6.9	4.2-10.8
	1981-1985	7.0	7.3	5.1-10.2	2.8	7.8	4.1-13.2	4.2	7.1	4.4-10.9	5.2	5.4	3.5-7.9	2.2	6.0	2.8-10.9	3.0	5.0	2.8-8.3
	1986-1990	5.4	5.1	3.4-7.5	3.6	8.0	4.7-12.7	1.8	2.9	1.3-5.7	4.4	4.4	2.7-6.6	3.0	6.8	3.8-11.3	1.4	2.3	0.9-4.9
	1991-1995	7.6	7.4	5.2-10.1	4.6	10.7	6.7-16.2	3.0	4.6	2.5-7.6	5.2	5.0	3.2-7.3	2.8	7.6	4.1-12.8	2.4	3.8	2.0-6.7
	1996-2000	7.4	6.6	4.6-9.1	4.2	8.4	5.2-12.9	3.2	5.0	2.9-8.2	4.0	3.6	2.2-5.5	1.6	3.4	1.4-6.6	2.4	3.7	1.9-6.5
	2001-2005	6.4	5.3	3.6-7.5	4.2	7.5	4.6-11.6	2.2	3.6	1.8-6.6	3.0	2.5	1.4-4.1	2.2	4.4	2.2-7.9	0.8	1.3	0.3-3.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**St. Lawrence**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	6.8	5.4	3.7-7.6	5.4	9.4	6.1-13.7	1.4	2.0	0.8-4.2	3.0	2.5	1.4-4.2	2.0	3.5	1.7-6.6	1.0	1.7	0.5-4.0
	2011-2015	9.0	6.5	4.7-8.8	7.0	11.2	7.7-15.7	2.0	2.5	1.2-4.9	2.8	2.0	1.1-3.5	1.8	3.2	1.4-6.1	1.0	1.2	0.4-3.2
	2016-2020	9.8	7.1	5.2-9.5	7.0	10.4	7.1-14.6	2.8	4.4	2.3-7.7	3.2	2.3	1.3-3.8	2.8	4.4	2.4-7.6	0.4	0.6	0.1-2.4
Colorectal	1976-1980	54.8	58.8	51.9-66.3	25.4	64.6	53.3-77.5	29.4	54.9	46.3-64.7	29.6	31.6	26.7-37.2	12.8	32.8	24.8-42.3	16.8	30.6	24.4-38.0
	1981-1985	67.8	69.6	62.3-77.5	31.0	76.0	64.0-89.5	36.8	64.4	55.3-74.5	33.6	34.4	29.3-40.1	16.6	41.0	32.2-51.3	17.0	29.9	23.8-37.1
	1986-1990	67.0	65.6	58.7-73.1	34.0	78.5	66.7-91.6	33.0	55.8	47.5-65.1	33.6	33.4	28.5-38.9	16.0	37.5	29.5-47.0	17.6	29.2	23.3-36.1
	1991-1995	71.0	66.5	59.7-73.8	33.8	75.1	63.9-87.7	37.2	60.6	52.1-70.1	37.2	34.9	30.0-40.3	16.4	37.4	29.5-46.8	20.8	32.7	26.6-39.7
	1996-2000	79.6	71.4	64.6-78.8	36.8	76.5	65.6-88.6	42.8	66.8	58.1-76.5	30.2	26.9	22.8-31.6	17.0	37.2	29.5-46.2	13.2	20.1	15.5-25.7
	2001-2005	79.8	67.6	61.2-74.6	40.4	78.8	68.1-90.7	39.4	58.6	50.6-67.5	30.2	25.6	21.7-30.0	16.4	33.2	26.3-41.4	13.8	20.2	15.7-25.7
	2006-2010	66.0	53.4	47.7-59.5	34.6	62.9	53.7-73.3	31.4	46.0	39.0-53.9	23.6	19.2	15.9-23.1	11.2	20.8	15.6-27.2	12.4	17.0	13.0-22.0
	2011-2015	55.4	42.2	37.3-47.7	29.4	47.8	40.1-56.4	26.0	37.0	30.7-44.2	16.2	12.5	9.9-15.7	8.8	15.4	11.1-20.8	7.4	10.5	7.3-14.7
2016-2020	51.2	37.3	32.8-42.4	27.0	42.1	35.1-50.1	24.2	32.4	26.7-39.2	21.2	15.3	12.4-18.6	10.8	16.8	12.5-22.2	10.4	13.8	10.2-18.5	
Colon excluding rectum	1976-1980	40.0	43.0	37.2-49.5	16.8	44.2	34.8-55.3	23.2	43.0	35.4-51.6	22.4	24.1	19.8-29.1	9.4	24.5	17.6-33.1	13.0	23.8	18.3-30.4
	1981-1985	47.8	49.3	43.1-56.0	20.6	52.2	42.1-63.8	27.2	47.1	39.4-55.8	26.6	27.3	22.8-32.4	12.8	31.8	24.1-41.1	13.8	24.6	19.1-31.2
	1986-1990	47.8	46.7	40.9-53.1	22.4	51.9	42.4-62.8	25.4	42.6	35.5-50.9	28.0	27.8	23.4-32.9	13.0	30.6	23.3-39.2	15.0	25.0	19.6-31.5
	1991-1995	51.8	48.5	42.8-54.9	22.8	51.3	42.0-61.9	29.0	47.0	39.6-55.4	32.2	30.3	25.8-35.4	13.8	32.5	25.0-41.5	18.4	28.8	23.2-35.4
	1996-2000	60.8	54.4	48.4-60.9	26.4	55.2	46.0-65.7	34.4	53.1	45.4-61.7	23.8	21.2	17.6-25.4	12.8	28.7	21.9-36.9	11.0	16.6	12.5-21.8
	2001-2005	58.4	49.5	44.0-55.5	27.6	54.0	45.2-64.0	30.8	45.6	38.6-53.5	23.6	20.0	16.6-24.0	11.8	24.0	18.2-31.1	11.8	17.1	13.0-22.2
	2006-2010	48.0	38.3	33.6-43.6	23.0	41.8	34.4-50.4	25.0	36.2	30.1-43.3	19.4	15.6	12.6-19.1	8.8	16.0	11.5-21.6	10.6	14.5	10.8-19.1
	2011-2015	35.4	27.0	23.1-31.4	18.0	29.3	23.4-36.2	17.4	24.8	19.7-30.8	10.8	8.3	6.2-10.9	5.6	9.7	6.4-14.1	5.2	7.3	4.7-10.9
2016-2020	35.4	26.0	22.2-30.3	18.2	28.6	22.9-35.4	17.2	23.0	18.2-28.8	15.6	11.3	8.9-14.3	8.0	12.3	8.7-17.0	7.6	10.3	7.2-14.5	
Rectum & rectosigmoid	1976-1980	14.8	15.8	12.3-19.8	8.6	20.4	14.6-27.7	6.2	12.0	8.1-17.0	7.2	7.5	5.2-10.4	3.4	8.2	4.7-13.3	3.8	6.9	4.1-10.8
	1981-1985	20.0	20.3	16.5-24.8	10.4	23.9	17.6-31.5	9.6	17.3	12.7-23.0	7.0	7.1	4.9-9.9	3.8	9.2	5.4-14.5	3.2	5.3	3.0-8.7
	1986-1990	19.2	18.8	15.2-23.0	11.6	26.6	20.0-34.6	7.6	13.1	9.2-18.1	5.6	5.6	3.7-8.1	3.0	7.0	3.8-11.5	2.6	4.2	2.2-7.2
	1991-1995	19.2	17.9	14.5-21.9	11.0	23.8	17.8-31.1	8.2	13.6	9.7-18.5	5.0	4.6	3.0-6.8	2.6	4.9	2.6-8.4	2.4	3.9	2.0-6.8
	1996-2000	18.8	17.0	13.8-20.9	10.4	21.3	15.9-28.0	8.4	13.7	9.9-18.6	6.4	5.7	3.9-8.0	4.2	8.4	5.2-13.0	2.2	3.5	1.7-6.3
	2001-2005	21.4	18.1	14.9-21.9	12.8	24.8	19.0-31.8	8.6	13.0	9.4-17.6	6.6	5.6	3.8-7.9	4.6	9.2	5.8-13.8	2.0	3.1	1.5-5.8
	2006-2010	18.0	15.0	12.0-18.5	11.6	21.1	16.0-27.4	6.4	9.7	6.6-13.9	4.2	3.6	2.2-5.5	2.4	4.8	2.5-8.4	1.8	2.6	1.2-5.1
	2011-2015	20.0	15.2	12.3-18.6	11.4	18.5	13.9-24.2	8.6	12.2	8.7-16.7	5.4	4.3	2.8-6.3	3.2	5.7	3.2-9.4	2.2	3.2	1.6-6.0
2016-2020	15.8	11.3	8.9-14.3	8.8	13.5	9.7-18.4	7.0	9.4	6.4-13.4	5.6	3.9	2.6-5.8	2.8	4.5	2.4-7.8	2.8	3.5	1.9-6.2	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**St. Lawrence**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	3.4	3.6	2.1-5.7	1.6	4.2	1.7-8.3	1.8	3.4	1.5-6.4	2.0	2.1	1.0-3.8	1.4	4.0	1.5-8.2	0.6	1.1	0.2-3.1
	1981-1985	3.4	3.3	1.9-5.3	2.2	4.5	2.2-8.2	1.2	2.0	0.7-4.4	3.8	3.7	2.2-5.9	2.2	5.0	2.5-9.1	1.6	2.7	1.2-5.4
	1986-1990	5.0	5.2	3.3-7.7	3.2	7.9	4.4-12.9	1.8	3.1	1.4-6.0	4.8	4.8	3.0-7.1	3.2	7.1	4.0-11.5	1.6	2.6	1.1-5.2
	1991-1995	6.0	5.6	3.8-8.0	3.6	8.0	4.7-12.7	2.4	3.9	2.0-6.9	2.6	2.4	1.3-4.1	1.6	3.4	1.5-6.8	1.0	1.5	0.5-3.7
	1996-2000	7.2	6.5	4.6-9.1	4.8	9.8	6.3-14.6	2.4	3.8	2.0-6.8	4.0	3.6	2.2-5.5	2.8	5.5	3.0-9.3	1.2	1.9	0.7-4.3
	2001-2005	4.4	3.8	2.4-5.8	3.2	5.9	3.3-9.7	1.2	2.0	0.7-4.4	5.6	4.8	3.2-6.9	3.6	7.1	4.1-11.3	2.0	3.2	1.5-6.0
	2006-2010	4.2	3.2	2.0-5.0	2.2	3.6	1.8-6.5	2.0	2.8	1.4-5.4	5.4	4.5	2.9-6.6	2.8	5.1	2.7-8.5	2.6	3.8	2.0-6.6
	2011-2015	8.2	6.0	4.3-8.3	5.4	7.8	5.1-11.5	2.8	4.0	2.2-7.0	5.6	4.2	2.8-6.2	2.8	4.2	2.3-7.3	2.8	3.9	2.1-6.8
2016-2020	8.6	5.7	4.1-7.8	5.4	7.7	5.0-11.4	3.2	3.9	2.2-6.7	7.0	4.6	3.2-6.4	3.4	4.9	2.8-8.1	3.6	4.1	2.4-6.8	
Pancreas	1976-1980	9.4	10.6	7.7-14.1	3.8	9.3	5.5-14.6	5.6	10.4	6.9-15.1	9.6	10.7	7.8-14.2	4.2	10.4	6.3-16.1	5.4	10.2	6.7-14.8
	1981-1985	10.0	10.5	7.8-13.9	5.6	14.6	9.5-21.3	4.4	7.6	4.7-11.5	9.2	9.5	6.9-12.8	4.6	11.2	6.9-16.9	4.6	7.8	5.0-11.8
	1986-1990	10.6	10.2	7.6-13.3	5.2	10.9	7.1-16.1	5.4	8.9	5.8-13.0	9.4	9.2	6.8-12.3	4.8	10.8	6.8-16.3	4.6	7.6	4.8-11.6
	1991-1995	14.4	13.2	10.3-16.6	7.2	14.9	10.4-20.7	7.2	11.6	8.1-16.1	13.4	12.3	9.5-15.7	6.4	13.2	9.0-18.7	7.0	11.3	7.8-15.8
	1996-2000	12.2	10.9	8.3-14.0	6.0	12.0	8.0-17.2	6.2	9.5	6.4-13.6	12.0	10.8	8.2-13.9	6.8	14.3	9.8-20.1	5.2	7.8	5.1-11.6
	2001-2005	12.6	10.6	8.1-13.6	6.4	12.1	8.2-17.1	6.2	9.4	6.3-13.4	12.6	10.6	8.1-13.6	6.4	12.2	8.3-17.3	6.2	9.3	6.3-13.3
	2006-2010	14.6	11.5	9.0-14.5	7.0	12.0	8.3-16.8	7.6	10.9	7.7-15.1	12.8	10.2	7.8-13.0	7.0	11.9	8.2-16.7	5.8	8.2	5.5-12.0
	2011-2015	18.2	13.5	10.8-16.7	9.6	15.7	11.5-21.0	8.6	11.5	8.3-15.8	15.6	11.4	9.0-14.3	7.4	11.8	8.2-16.5	8.2	10.9	7.7-15.0
2016-2020	20.2	13.3	10.8-16.3	8.6	11.5	8.3-15.8	11.6	14.7	11.1-19.4	17.4	11.7	9.3-14.5	9.2	13.2	9.6-17.8	8.2	10.0	7.1-13.8	
Larynx	1976-1980	6.2	5.9	4.0-8.4	4.6	9.8	6.1-14.8	1.6	2.8	1.2-5.5	2.8	2.9	1.5-4.8	2.6	6.0	3.1-10.4	0.2	0.3	0.0-2.0
	1981-1985	5.0	5.0	3.2-7.4	3.4	7.3	4.2-11.8	1.6	3.1	1.3-6.1	2.6	2.3	1.2-4.1	1.8	3.8	1.7-7.3	0.8	1.4	0.4-3.6
	1986-1990	11.4	11.4	8.6-14.9	9.0	20.1	14.5-27.0	2.4	4.5	2.3-7.9	2.8	2.9	1.6-4.9	2.4	5.5	2.8-9.7	0.4	0.7	0.1-2.5
	1991-1995	4.2	3.8	2.4-5.8	3.8	7.9	4.7-12.4	0.4	0.7	0.1-2.5	2.4	2.2	1.1-3.9	1.8	3.6	1.6-6.9	0.6	1.0	0.2-3.1
	1996-2000	6.8	6.2	4.3-8.6	5.0	9.8	6.3-14.5	1.8	3.3	1.5-6.2	0.6	0.5	0.1-1.5	0.6	1.1	0.2-3.2	0.0	0.0	0.0-1.3
	2001-2005	7.2	6.0	4.2-8.4	5.0	9.1	5.9-13.5	2.2	3.6	1.8-6.5	1.6	1.3	0.6-2.7	1.4	2.6	1.0-5.4	0.2	0.3	0.0-1.9
	2006-2010	6.4	5.0	3.4-7.1	5.0	8.2	5.3-12.3	1.4	2.0	0.8-4.3	2.0	1.5	0.7-2.8	1.6	2.6	1.1-5.2	0.4	0.5	0.1-2.2
	2011-2015	5.8	4.1	2.7-6.0	4.4	6.4	4.0-10.0	1.4	2.1	0.8-4.5	1.4	1.1	0.4-2.3	0.8	1.4	0.4-3.7	0.6	0.8	0.2-2.6
2016-2020	8.2	5.7	4.0-7.8	5.6	8.0	5.3-11.8	2.6	3.4	1.8-6.1	0.8	0.5	0.1-1.4	0.8	1.1	0.3-3.1	0.0	0.0	0.0-1.5	
Lung and bronchus	1976-1980	51.2	50.6	44.5-57.3	39.8	92.1	79.3-106.4	11.4	20.7	15.6-26.9	41.4	41.4	35.9-47.5	31.8	73.8	62.4-86.7	9.6	17.7	13.0-23.5
	1981-1985	76.4	75.2	67.7-83.3	52.0	119.1	104.6-134.9	24.4	43.4	35.9-52.1	59.8	58.7	52.1-65.8	40.8	94.7	81.6-109.1	19.0	33.5	27.0-41.1
	1986-1990	77.0	75.2	67.8-83.3	49.4	112.5	98.5-127.9	27.6	48.7	40.7-57.7	70.0	68.1	61.1-75.7	45.4	105.1	91.4-120.1	24.6	42.9	35.5-51.3
	1991-1995	90.2	83.0	75.4-91.0	51.8	109.5	96.2-124.0	38.4	63.0	54.3-72.7	71.8	66.3	59.6-73.6	45.2	96.9	84.4-110.7	26.6	43.6	36.4-51.8
	1996-2000	99.4	88.2	80.6-96.3	56.6	113.3	100.3-127.5	42.8	69.4	60.3-79.4	77.8	69.3	62.5-76.5	44.6	90.8	79.1-103.7	33.2	53.4	45.6-62.3
	2001-2005	109.8	92.7	85.1-100.9	60.8	117.5	104.4-131.8	49.0	75.5	66.3-85.7	72.4	61.1	55.0-67.8	40.0	79.6	68.7-91.7	32.4	49.4	42.1-57.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020  
St. Lawrence**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	116.0	93.2	85.7-101.2	62.4	111.8	99.5-125.2	53.6	79.9	70.5-90.2	81.0	65.5	59.2-72.3	43.4	77.6	67.4-88.9	37.6	55.6	47.9-64.4
	2011-2015	99.8	73.2	66.8-80.1	51.8	81.6	71.7-92.5	48.0	66.5	58.2-75.7	76.0	56.1	50.5-62.2	42.2	67.0	58.0-77.0	33.8	47.4	40.4-55.3
	2016-2020	108.0	73.5	67.3-80.1	54.0	77.2	68.0-87.3	54.0	70.6	62.2-79.9	60.2	40.4	35.9-45.4	30.4	44.3	37.3-52.2	29.8	38.1	32.1-45.0
Melanoma of the skin	1976-1980	3.0	3.5	1.9-5.8	2.0	4.6	2.1-8.5	1.0	2.0	0.7-4.7	1.2	1.3	0.5-2.8	0.8	1.8	0.5-4.7	0.4	0.8	0.1-3.0
	1981-1985	5.8	5.9	3.9-8.5	3.0	6.7	3.6-11.1	2.8	5.2	2.8-8.7	1.0	1.0	0.3-2.4	0.6	1.3	0.3-3.8	0.4	0.8	0.1-2.8
	1986-1990	4.6	4.6	2.9-7.0	3.4	7.6	4.4-12.3	1.2	2.0	0.7-4.5	2.4	2.3	1.2-4.1	1.6	3.3	1.4-6.6	0.8	1.3	0.4-3.4
	1991-1995	3.6	3.4	2.0-5.4	1.8	4.2	1.9-8.0	1.8	3.0	1.4-5.8	1.4	1.4	0.5-2.8	1.0	2.1	0.7-4.9	0.4	0.8	0.1-2.8
	1996-2000	11.4	10.5	7.9-13.6	6.2	12.2	8.3-17.5	5.2	9.1	5.9-13.4	2.2	1.9	0.9-3.4	1.4	2.6	1.0-5.4	0.8	1.2	0.3-3.3
	2001-2005	21.8	18.7	15.4-22.6	13.0	23.9	18.4-30.6	8.8	15.0	10.9-20.3	2.8	2.4	1.3-4.0	1.2	2.2	0.8-4.9	1.6	2.6	1.1-5.2
	2006-2010	30.2	25.5	21.5-30.0	15.6	27.6	21.7-34.6	14.6	25.1	19.5-31.8	2.2	1.8	0.9-3.3	1.0	2.0	0.6-4.6	1.2	1.9	0.7-4.3
	2011-2015	29.2	23.9	20.1-28.3	16.0	26.0	20.4-32.6	13.2	23.3	17.8-29.9	3.0	2.3	1.3-3.9	2.0	3.3	1.5-6.1	1.0	1.5	0.4-3.7
2016-2020	28.2	21.2	17.7-25.3	16.4	24.9	19.6-31.2	11.8	18.6	13.9-24.5	2.2	1.6	0.8-3.0	2.0	3.6	1.7-6.7	0.2	0.2	0.0-1.7	
Female breast	1976-1980							42.4	82.5	71.5-94.6							19.6	36.0	29.1-44.0
	1981-1985							50.0	93.6	82.1-106.3							17.4	32.2	25.7-40.0
	1986-1990							56.8	103.0	91.1-116.0							19.0	33.9	27.3-41.6
	1991-1995							63.4	111.1	99.0-124.2							15.2	25.0	19.6-31.4
	1996-2000							82.2	138.1	125.0-152.2							14.0	21.9	17.0-27.7
	2001-2005							79.6	127.8	115.5-141.2							16.2	24.6	19.5-30.7
	2006-2010							71.0	110.4	99.0-122.8							11.8	17.7	13.4-23.0
	2011-2015							71.8	106.8	95.7-119.0							14.8	20.4	15.9-25.8
2016-2020							76.2	110.2	98.7-122.6							10.0	13.3	9.8-17.9	
Cervix uteri	1976-1980							8.0	15.6	11.0-21.4							2.6	4.6	2.4-8.0
	1981-1985							6.0	11.6	7.7-16.7							2.6	5.3	2.8-9.2
	1986-1990							6.6	12.7	8.7-18.0							2.4	4.6	2.3-8.0
	1991-1995							8.0	15.0	10.6-20.4							2.6	4.8	2.6-8.3
	1996-2000							7.4	12.9	9.0-17.8							2.2	3.6	1.8-6.5
	2001-2005							6.4	11.2	7.6-16.0							2.8	4.5	2.4-7.7
	2006-2010							3.4	6.0	3.4-9.8							1.2	1.9	0.7-4.4
	2011-2015							3.6	6.2	3.6-10.1							1.4	2.4	0.9-5.2
	2016-2020							3.2	6.3	3.5-10.4							1.4	2.2	0.8-4.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**St. Lawrence**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							14.0	26.5	20.5-33.6							3.4	6.2	3.6-10.0
	1981-1985							11.6	21.8	16.4-28.3							1.8	3.7	1.6-7.0
	1986-1990							9.2	17.1	12.4-23.0							1.4	2.2	0.9-4.7
	1991-1995							11.0	19.6	14.7-25.5							2.8	4.7	2.6-8.0
	1996-2000							13.4	22.8	17.6-29.1							2.2	3.3	1.6-6.1
	2001-2005							20.6	33.0	26.9-40.2							2.6	4.0	2.1-7.0
	2006-2010							20.0	30.7	24.9-37.6							3.2	4.7	2.7-7.7
	2011-2015							21.6	31.3	25.5-38.1							5.4	7.6	5.0-11.2
	2016-2020							23.0	31.3	25.6-38.1							3.2	3.8	2.2-6.5
Ovary	1976-1980							7.8	14.2	10.0-19.6							4.4	7.9	4.9-12.1
	1981-1985							4.6	8.4	5.3-12.8							4.0	7.0	4.3-10.9
	1986-1990							6.0	10.5	7.0-15.1							3.8	6.3	3.8-9.9
	1991-1995							8.6	14.1	10.2-19.1							7.8	12.6	8.9-17.3
	1996-2000							10.6	17.5	13.1-23.0							6.6	10.4	7.1-14.6
	2001-2005							8.6	13.6	9.8-18.3							6.0	9.3	6.3-13.4
	2006-2010							5.8	9.0	6.0-13.1							3.8	5.6	3.4-8.9
	2011-2015							8.6	13.1	9.4-18.0							4.6	5.9	3.7-9.1
	2016-2020							4.6	7.4	4.5-11.6							3.8	5.4	3.1-8.7
Prostate	1976-1980				34.0	91.6	77.7-107.1							10.4	31.6	23.2-41.7			
	1981-1985				28.4	76.8	64.1-91.0							9.8	30.6	22.3-40.7			
	1986-1990				44.8	114.9	99.8-131.4							12.4	32.7	24.8-42.1			
	1991-1995				57.8	130.7	115.6-147.2							16.6	43.2	34.1-53.7			
	1996-2000				66.8	132.4	118.4-147.7							13.6	34.8	26.8-44.3			
	2001-2005				98.0	187.5	171.0-205.2							13.4	30.5	23.5-38.9			
	2006-2010				89.2	151.1	137.1-166.1							10.2	20.9	15.5-27.5			
	2011-2015				96.8	144.9	132.0-158.9							10.0	19.8	14.6-26.1			
	2016-2020				91.2	125.9	114.4-138.5							8.2	13.4	9.6-18.3			
Testis	1976-1980				1.4	2.5	0.9-5.6							0.2	0.2	0.0-2.0			
	1981-1985				2.8	5.2	2.7-9.2							0.4	0.8	0.1-3.1			
	1986-1990				1.6	2.5	1.0-5.2							0.2	0.4	0.0-2.4			
	1991-1995				2.0	3.3	1.6-6.3							0.2	0.3	0.0-2.0			
	1996-2000				2.0	3.4	1.6-6.4							0.0	0.0	0.0-1.5			
	2001-2005				2.6	5.0	2.7-8.6							0.2	0.3	0.0-2.0			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**St. Lawrence**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				3.6	6.4	3.7-10.3							0.0	0.0	0.0-1.4			
	2011-2015				2.2	3.9	1.9-7.1							0.0	0.0	0.0-1.4			
	2016-2020				3.6	7.3	4.2-11.6							0.0	0.0	0.0-1.4			
Urinary bladder (incl. in situ)	1976-1980	24.8	26.0	21.5-31.0	17.2	44.1	34.8-54.9	7.6	13.7	9.6-18.8	8.0	9.3	6.6-12.6	5.4	17.0	11.0-24.7	2.6	4.8	2.5-8.2
	1981-1985	22.0	22.7	18.6-27.4	17.2	43.1	34.1-53.7	4.8	9.1	5.8-13.7	7.2	7.6	5.3-10.6	5.0	14.4	9.1-21.4	2.2	3.7	1.9-6.7
	1986-1990	26.4	26.2	21.9-31.1	19.2	47.9	38.5-58.8	7.2	12.1	8.4-16.8	5.6	5.7	3.8-8.2	4.0	10.6	6.3-16.4	1.6	2.8	1.2-5.6
	1991-1995	24.2	22.5	18.7-27.0	16.8	38.9	30.8-48.4	7.4	12.0	8.4-16.6	5.8	5.3	3.6-7.6	3.6	8.5	4.9-13.6	2.2	3.4	1.7-6.1
	1996-2000	30.6	27.4	23.2-32.1	20.6	42.7	34.7-51.9	10.0	15.9	11.8-21.1	4.8	4.3	2.8-6.4	2.0	4.8	2.2-8.8	2.8	4.3	2.3-7.2
	2001-2005	38.2	32.4	28.0-37.4	27.0	54.1	45.2-64.2	11.2	17.0	12.8-22.2	7.0	6.0	4.2-8.4	3.8	8.2	4.9-12.9	3.2	4.4	2.5-7.3
	2006-2010	38.0	30.3	26.1-35.0	26.8	48.1	40.1-57.2	11.2	16.2	12.2-21.1	6.6	5.4	3.7-7.6	3.2	6.7	3.8-10.9	3.4	4.6	2.7-7.5
	2011-2015	44.2	33.0	28.7-37.7	34.4	57.9	49.4-67.5	9.8	13.7	10.1-18.4	4.2	3.4	2.1-5.2	2.6	4.8	2.5-8.2	1.6	2.2	0.9-4.6
2016-2020	40.6	27.6	23.9-31.8	28.4	42.7	35.8-50.5	12.2	15.6	11.9-20.4	7.2	4.9	3.4-6.9	4.8	7.4	4.7-11.2	2.4	2.9	1.5-5.4	
Kidney and renal pelvis	1976-1980	5.8	5.7	3.8-8.2	4.0	9.7	5.8-15.1	1.8	2.9	1.3-5.7	2.8	2.8	1.5-4.8	2.6	6.3	3.2-11.0	0.2	0.3	0.0-1.9
	1981-1985	6.6	6.4	4.4-9.1	4.0	8.4	5.0-13.2	2.6	4.7	2.5-8.2	4.6	4.5	2.8-6.8	2.2	5.1	2.4-9.3	2.4	4.2	2.1-7.4
	1986-1990	11.2	10.7	8.1-14.0	6.6	13.6	9.3-19.3	4.6	8.2	5.2-12.4	5.0	4.9	3.2-7.3	3.2	7.5	4.2-12.3	1.8	3.1	1.4-5.9
	1991-1995	10.8	9.9	7.5-13.0	6.6	14.1	9.6-20.0	4.2	7.2	4.4-11.1	6.0	5.6	3.8-8.1	4.4	10.2	6.3-15.5	1.6	2.8	1.2-5.5
	1996-2000	11.8	10.7	8.1-13.8	5.8	11.4	7.6-16.4	6.0	9.9	6.7-14.2	3.4	3.0	1.8-4.9	1.8	4.0	1.8-7.6	1.6	2.4	1.1-4.9
	2001-2005	15.6	13.3	10.5-16.6	8.4	15.7	11.3-21.3	7.2	10.9	7.6-15.3	5.8	4.9	3.3-7.1	3.8	7.0	4.2-11.0	2.0	2.8	1.3-5.3
	2006-2010	15.0	12.0	9.4-15.1	7.6	13.2	9.2-18.2	7.4	11.4	8.0-15.8	3.6	3.0	1.7-4.7	2.2	3.7	1.8-6.8	1.4	2.4	1.0-5.1
	2011-2015	21.4	16.4	13.4-19.9	14.2	22.4	17.4-28.4	7.2	10.8	7.5-15.1	4.2	3.1	1.9-4.8	2.2	3.7	1.8-6.7	2.0	2.8	1.3-5.3
2016-2020	26.0	19.7	16.3-23.6	17.0	26.2	20.7-32.8	9.0	13.3	9.6-18.2	5.0	3.3	2.1-5.0	2.8	3.8	2.0-6.7	2.2	2.7	1.3-5.1	
Brain and other central nervous system	1976-1980	4.8	5.0	3.1-7.5	2.6	5.8	3.0-10.0	2.2	4.4	2.1-7.9	3.6	3.8	2.2-6.0	2.2	4.7	2.3-8.6	1.4	3.1	1.2-6.3
	1981-1985	4.2	4.2	2.6-6.4	1.8	3.7	1.6-7.2	2.4	4.4	2.3-7.8	3.4	3.4	2.0-5.5	2.6	6.0	3.1-10.5	0.8	1.4	0.4-3.6
	1986-1990	6.6	6.4	4.4-9.1	3.4	6.7	3.9-10.8	3.2	6.1	3.4-10.0	3.6	3.5	2.0-5.5	1.6	3.4	1.4-6.7	2.0	3.5	1.7-6.6
	1991-1995	9.8	8.7	6.4-11.6	5.0	9.5	6.1-14.1	4.8	8.0	5.1-12.0	6.4	5.8	3.9-8.2	3.0	6.2	3.4-10.3	3.4	5.5	3.2-8.8
	1996-2000	9.2	8.2	6.0-11.0	5.4	10.3	6.7-15.0	3.8	6.4	3.8-10.0	7.0	6.3	4.4-8.8	4.4	8.5	5.3-12.9	2.6	4.3	2.3-7.4
	2001-2005	8.2	7.2	5.1-9.8	4.2	7.7	4.7-11.8	4.0	6.9	4.1-10.7	4.8	4.1	2.6-6.1	2.8	5.0	2.7-8.5	2.0	3.2	1.5-6.0
	2006-2010	5.2	4.7	3.0-6.9	3.6	6.6	3.9-10.6	1.6	2.8	1.2-5.6	4.0	3.6	2.2-5.6	2.6	4.7	2.5-8.1	1.4	2.5	1.0-5.2
	2011-2015	9.8	7.2	5.3-9.6	5.2	7.9	5.1-11.7	4.6	6.6	4.1-10.1	5.8	4.1	2.7-6.0	3.0	4.6	2.5-7.8	2.8	3.6	2.0-6.3
2016-2020	8.2	7.0	4.9-9.7	4.4	7.8	4.8-12.1	3.8	6.4	3.7-10.4	4.8	3.7	2.3-5.7	3.0	5.2	2.8-8.7	1.8	2.4	1.1-4.8	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**St. Lawrence**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	2.4	2.5	1.3-4.4	0.8	1.7	0.5-4.5	1.6	3.0	1.3-6.0	1.0	1.0	0.3-2.4	0.4	0.9	0.1-3.3	0.6	1.0	0.2-3.1
	1981-1985	2.6	2.3	1.2-4.0	1.0	2.2	0.7-5.1	1.6	2.5	1.0-5.0	0.6	0.6	0.1-1.8	0.2	0.4	0.0-2.3	0.4	0.8	0.1-2.8
	1986-1990	2.0	1.9	0.9-3.5	0.2	0.3	0.0-2.2	1.8	3.3	1.5-6.3	0.0	0.0	0.0-0.7	0.0	0.0	0.0-1.6	0.0	0.0	0.0-1.4
	1991-1995	3.6	3.3	2.0-5.3	0.6	1.2	0.2-3.6	3.0	5.4	3.0-8.9	0.2	0.2	0.0-1.1	0.2	0.5	0.0-2.7	0.0	0.0	0.0-1.3
	1996-2000	7.6	7.0	4.9-9.6	2.0	3.9	1.8-7.2	5.6	10.4	6.9-15.0	1.2	1.1	0.4-2.3	0.2	0.4	0.0-2.1	1.0	1.5	0.5-3.7
	2001-2005	6.4	5.5	3.8-7.8	2.4	4.4	2.2-7.7	4.0	7.0	4.2-10.9	0.4	0.3	0.0-1.2	0.2	0.4	0.0-2.1	0.2	0.3	0.0-1.9
	2006-2010	9.8	8.9	6.6-11.9	1.8	3.0	1.3-5.7	8.0	15.1	10.6-20.7	1.0	0.8	0.3-1.9	0.2	0.3	0.0-2.0	0.8	1.2	0.3-3.3
	2011-2015	14.8	13.2	10.3-16.8	4.6	7.6	4.7-11.6	10.2	19.3	14.2-25.6	0.4	0.3	0.0-1.2	0.2	0.3	0.0-2.0	0.2	0.3	0.0-1.9
	2016-2020	9.0	8.0	5.7-10.8	1.4	2.5	1.0-5.3	7.6	13.9	9.6-19.4	0.8	0.6	0.2-1.6	0.4	0.7	0.1-2.7	0.4	0.6	0.1-2.4
Hodgkin lymphoma	1976-1980	3.2	2.8	1.6-4.7	1.4	2.2	0.9-5.0	1.8	3.3	1.4-6.3	1.8	1.9	0.8-3.6	0.4	0.8	0.1-3.2	1.4	2.7	1.0-5.6
	1981-1985	3.6	3.5	2.0-5.5	2.2	4.4	2.1-8.0	1.4	2.4	1.0-5.2	1.0	0.9	0.3-2.1	0.4	0.9	0.1-3.3	0.6	0.8	0.2-2.7
	1986-1990	1.6	1.3	0.6-2.7	0.8	1.2	0.3-3.5	0.8	1.4	0.4-3.7	0.6	0.4	0.1-1.4	0.4	0.4	0.1-2.2	0.2	0.3	0.0-2.0
	1991-1995	2.6	2.3	1.2-4.0	0.8	2.0	0.5-5.0	1.8	3.2	1.4-6.1	0.4	0.3	0.0-1.2	0.2	0.3	0.0-2.1	0.2	0.3	0.0-1.9
	1996-2000	3.0	2.6	1.4-4.2	1.6	2.6	1.1-5.3	1.4	2.5	1.0-5.1	1.0	0.8	0.3-1.9	0.4	0.5	0.1-2.2	0.6	1.0	0.2-3.0
	2001-2005	2.4	2.1	1.1-3.8	1.8	3.2	1.4-6.1	0.6	1.1	0.2-3.3	0.0	0.0	0.0-0.7	0.0	0.0	0.0-1.4	0.0	0.0	0.0-1.4
	2006-2010	3.6	2.9	1.7-4.7	2.6	3.9	2.0-6.9	1.0	1.9	0.6-4.5	0.8	0.8	0.2-2.1	0.4	0.8	0.1-2.8	0.4	1.0	0.1-3.3
	2011-2015	3.2	2.7	1.5-4.5	2.0	3.3	1.6-6.2	1.2	2.2	0.8-4.9	0.2	0.2	0.0-1.0	0.0	0.0	0.0-1.4	0.2	0.3	0.0-1.9
	2016-2020	1.6	1.6	0.7-3.1	0.6	1.0	0.2-3.0	1.0	2.2	0.7-5.2	0.4	0.3	0.0-1.1	0.4	0.6	0.1-2.3	0.0	0.0	0.0-1.5
Non-Hodgkin lymphomas	1976-1980	9.6	9.9	7.3-13.3	4.6	11.2	6.9-17.1	5.0	9.2	5.9-13.7	5.4	5.3	3.4-7.7	3.2	7.8	4.2-12.9	2.2	3.9	1.9-7.1
	1981-1985	10.4	10.8	8.0-14.2	6.4	14.7	10.0-20.9	4.0	7.6	4.6-11.8	7.6	7.8	5.4-10.7	4.8	10.9	6.9-16.4	2.8	5.1	2.8-8.7
	1986-1990	13.0	12.8	9.8-16.3	5.4	12.2	7.9-18.0	7.6	13.2	9.3-18.3	7.4	7.4	5.2-10.2	3.2	8.5	4.8-13.8	4.2	7.0	4.3-10.7
	1991-1995	16.6	15.3	12.2-19.0	7.4	14.8	10.3-20.6	9.2	15.3	11.2-20.5	9.4	8.8	6.4-11.7	3.8	8.0	4.7-12.6	5.6	9.0	6.0-13.1
	1996-2000	23.2	20.8	17.2-24.9	13.4	26.4	20.3-33.6	9.8	16.2	11.9-21.5	9.0	8.1	5.9-10.8	5.6	11.2	7.4-16.3	3.4	5.4	3.1-8.7
	2001-2005	25.6	22.2	18.5-26.5	14.2	27.6	21.4-34.9	11.4	18.3	13.8-23.8	8.8	7.6	5.5-10.3	4.8	10.1	6.4-15.1	4.0	5.9	3.6-9.3
	2006-2010	27.6	22.1	18.5-26.2	13.8	24.2	18.7-30.8	13.8	20.3	15.8-25.9	8.2	6.6	4.7-9.0	4.6	8.8	5.5-13.3	3.6	5.1	3.0-8.3
	2011-2015	27.8	21.3	17.8-25.2	15.8	25.4	20.0-31.9	12.0	17.2	13.0-22.4	7.6	5.6	4.0-7.8	4.4	7.3	4.5-11.2	3.2	4.0	2.3-6.7
	2016-2020	24.2	17.3	14.3-20.9	14.6	21.0	16.3-26.8	9.6	13.5	9.8-18.2	5.6	3.9	2.6-5.7	2.6	3.7	2.0-6.6	3.0	3.9	2.2-6.7
Myeloma	1976-1980	4.4	4.4	2.7-6.7	2.6	5.5	2.9-9.4	1.8	3.4	1.5-6.4	3.4	3.7	2.1-5.9	2.0	4.5	2.1-8.4	1.4	2.6	1.1-5.5
	1981-1985	3.6	3.4	2.0-5.5	1.6	3.4	1.5-6.9	2.0	3.3	1.6-6.1	3.4	3.5	2.0-5.6	1.6	4.4	1.8-8.7	1.8	3.0	1.3-5.7
	1986-1990	5.6	5.7	3.7-8.2	2.6	6.5	3.4-11.0	3.0	5.2	2.9-8.6	4.0	3.8	2.3-5.8	2.2	4.7	2.3-8.6	1.8	2.9	1.3-5.6
	1991-1995	7.4	7.1	5.0-9.8	3.4	8.1	4.6-13.0	4.0	6.9	4.2-10.7	3.0	2.9	1.6-4.8	1.6	3.8	1.6-7.5	1.4	2.4	0.9-5.0
	1996-2000	7.0	6.3	4.4-8.7	3.8	7.7	4.6-12.1	3.2	4.9	2.8-8.0	5.0	4.5	2.9-6.7	2.4	5.1	2.6-8.9	2.6	3.8	2.0-6.7
	2001-2005	8.2	6.9	5.0-9.4	4.4	8.8	5.5-13.5	3.8	5.5	3.3-8.7	6.2	5.3	3.6-7.6	3.2	7.0	3.9-11.4	3.0	4.3	2.4-7.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**St. Lawrence**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	6.6	5.2	3.6-7.4	3.4	5.9	3.4-9.6	3.2	4.7	2.7-7.7	5.2	4.2	2.7-6.2	3.2	6.3	3.6-10.2	2.0	2.8	1.3-5.3
	2011-2015	8.4	6.2	4.4-8.5	5.8	9.5	6.3-13.8	2.6	3.4	1.8-6.0	3.2	2.3	1.3-3.8	1.4	2.1	0.8-4.6	1.8	2.5	1.1-5.0
	2016-2020	9.6	6.8	5.0-9.2	5.8	9.1	6.0-13.2	3.8	4.9	2.8-8.0	3.8	2.8	1.7-4.5	2.8	4.7	2.5-7.9	1.0	1.6	0.5-4.1
Leukemias	1976-1980	9.6	9.9	7.2-13.2	4.6	10.7	6.7-16.2	5.0	9.3	5.9-13.8	7.2	7.3	5.0-10.1	3.0	7.7	4.2-12.8	4.2	7.5	4.6-11.6
	1981-1985	13.0	12.7	9.7-16.2	8.2	19.4	13.6-26.8	4.8	8.1	5.1-12.1	9.8	9.9	7.3-13.1	6.2	16.4	10.8-23.7	3.6	6.4	3.7-10.1
	1986-1990	11.6	11.4	8.7-14.8	6.8	15.3	10.4-21.6	4.8	8.4	5.3-12.5	6.8	6.5	4.5-9.0	4.4	9.6	5.8-14.8	2.4	3.9	2.0-6.8
	1991-1995	19.2	17.9	14.5-21.9	11.2	23.6	17.7-30.9	8.0	13.3	9.4-18.1	10.8	10.0	7.5-13.0	6.6	14.4	9.8-20.4	4.2	6.7	4.1-10.3
	1996-2000	20.4	18.5	15.1-22.4	10.8	21.3	15.9-27.8	9.6	16.3	12.0-21.7	9.4	8.5	6.2-11.3	6.0	13.0	8.7-18.7	3.4	5.5	3.2-8.8
	2001-2005	20.2	17.2	14.0-20.9	12.4	24.9	18.9-32.1	7.8	12.1	8.6-16.7	9.4	8.0	5.9-10.7	5.8	11.8	7.8-17.0	3.6	5.4	3.2-8.7
	2006-2010	19.8	16.6	13.4-20.3	12.6	23.5	17.9-30.1	7.2	11.1	7.7-15.5	12.2	10.3	7.8-13.2	9.0	18.1	13.1-24.3	3.2	4.5	2.6-7.5
	2011-2015	20.8	16.4	13.3-20.0	12.4	20.9	15.9-27.0	8.4	12.0	8.6-16.5	8.0	6.3	4.5-8.6	4.6	8.3	5.2-12.5	3.4	5.0	2.9-8.2
	2016-2020	24.4	17.7	14.6-21.4	15.4	22.8	17.8-28.9	9.0	13.0	9.2-17.8	7.8	5.9	4.2-8.2	5.0	8.4	5.4-12.6	2.8	3.5	1.9-6.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Saratoga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	458.0	397.1	380.5-414.2	235.0	491.0	461.0-522.3	223.0	344.6	324.3-365.7	250.4	224.1	211.5-237.2	136.6	296.8	273.1-321.9	113.8	178.5	164.0-194.0
	1981-1985	546.6	427.5	411.2-444.2	278.4	507.6	479.4-537.0	268.2	375.9	355.8-396.7	282.8	228.9	216.9-241.4	152.0	296.4	274.1-319.9	130.8	186.2	172.1-201.1
	1986-1990	628.0	431.6	416.3-447.2	316.0	509.8	483.4-537.3	312.0	385.7	366.6-405.4	317.2	224.7	213.6-236.2	167.4	285.6	265.3-307.0	149.8	185.4	172.3-199.3
	1991-1995	750.8	453.2	438.6-468.0	393.2	549.4	524.2-575.4	357.6	390.3	372.4-408.9	342.0	213.9	203.8-224.4	184.6	278.5	259.8-298.1	157.4	173.0	161.1-185.5
	1996-2000	946.0	509.0	494.6-523.8	466.6	573.3	549.5-597.8	479.4	467.9	449.3-487.1	368.4	204.8	195.5-214.4	183.0	243.0	226.9-260.0	185.4	180.8	169.3-192.9
	2001-2005	1,113.8	521.9	508.2-535.9	568.8	603.4	580.9-626.6	545.0	466.0	448.6-484.0	394.2	189.3	181.0-197.9	196.6	223.6	209.4-238.4	197.6	167.2	156.9-178.1
	2006-2010	1,276.6	524.4	511.4-537.7	663.4	602.3	581.4-623.8	613.2	468.5	451.8-485.7	422.0	178.0	170.4-185.9	216.0	212.6	199.8-226.0	206.0	154.6	145.1-164.5
	2011-2015	1,406.0	515.5	503.2-528.0	717.8	571.5	552.4-591.2	688.2	476.1	459.9-492.8	450.4	166.4	159.5-173.6	241.4	204.5	192.8-216.7	209.0	139.3	130.8-148.3
2016-2020	1,531.0	493.0	481.7-504.6	784.8	535.8	518.7-553.4	746.2	464.8	449.4-480.6	461.8	146.9	140.8-153.2	241.8	174.1	164.2-184.5	220.0	127.3	119.7-135.3	
Oral cavity and pharynx	1976-1980	14.2	12.8	9.9-16.2	11.6	24.6	18.3-32.3	2.6	4.3	2.3-7.3	7.0	6.4	4.4-8.9	5.4	12.4	7.9-18.3	1.6	2.6	1.1-5.1
	1981-1985	16.6	12.5	9.9-15.5	11.4	17.9	13.4-23.4	5.2	7.5	4.9-11.0	5.4	4.0	2.6-5.8	3.0	5.1	2.7-8.6	2.4	3.2	1.7-5.7
	1986-1990	14.6	9.8	7.6-12.3	10.2	16.0	11.6-21.3	4.4	5.5	3.4-8.3	5.8	4.0	2.6-5.7	4.4	7.4	4.5-11.5	1.4	1.6	0.7-3.4
	1991-1995	21.0	12.2	9.9-14.8	14.8	19.7	15.2-24.9	6.2	6.7	4.5-9.5	5.8	3.4	2.3-5.0	4.8	6.5	4.1-9.7	1.0	1.1	0.3-2.5
	1996-2000	18.4	9.6	7.7-11.7	11.6	12.5	9.5-16.2	6.8	6.6	4.6-9.2	2.6	1.4	0.8-2.4	1.6	1.9	0.8-3.7	1.0	1.0	0.3-2.3
	2001-2005	24.8	11.4	9.4-13.6	16.0	15.9	12.6-19.9	8.8	7.6	5.5-10.2	5.8	2.7	1.8-3.9	4.0	4.3	2.6-6.7	1.8	1.5	0.7-2.9
	2006-2010	31.0	12.4	10.5-14.5	21.4	18.3	14.9-22.2	9.6	7.1	5.2-9.5	4.8	1.9	1.2-2.9	2.8	2.5	1.3-4.2	2.0	1.3	0.6-2.5
	2011-2015	33.0	11.8	10.0-13.8	23.2	16.8	13.8-20.3	9.8	7.1	5.2-9.5	6.0	2.1	1.4-3.0	3.8	2.8	1.7-4.5	2.2	1.5	0.7-2.7
2016-2020	45.8	14.2	12.4-16.3	32.0	20.7	17.5-24.4	13.8	8.3	6.4-10.7	8.6	2.7	1.9-3.7	6.0	3.9	2.6-5.8	2.6	1.5	0.8-2.8	
Esophagus	1976-1980	4.0	3.8	2.3-5.9	2.8	6.8	3.5-11.8	1.2	1.9	0.7-4.2	3.6	3.3	1.9-5.2	2.6	5.3	2.7-9.4	1.0	1.6	0.5-3.7
	1981-1985	4.2	3.3	2.0-5.0	2.6	4.6	2.4-8.0	1.6	2.3	1.0-4.5	3.6	2.9	1.7-4.6	3.0	6.0	3.2-10.0	0.6	0.8	0.2-2.4
	1986-1990	6.8	4.8	3.3-6.7	4.8	8.0	5.0-12.0	2.0	2.3	1.1-4.3	4.6	3.2	2.0-4.7	3.2	5.2	2.9-8.6	1.4	1.7	0.7-3.5
	1991-1995	9.8	5.9	4.4-7.8	8.0	10.8	7.6-14.9	1.8	2.0	0.9-3.8	8.4	5.2	3.7-7.0	6.8	10.2	6.9-14.5	1.6	1.8	0.8-3.5
	1996-2000	8.4	4.6	3.3-6.2	6.4	8.1	5.5-11.6	2.0	2.0	0.9-3.6	9.6	5.2	3.9-6.9	7.8	10.0	7.0-13.8	1.8	1.7	0.8-3.3
	2001-2005	12.8	6.1	4.7-7.8	10.2	11.2	8.3-14.7	2.6	2.2	1.2-3.8	11.6	5.5	4.2-7.1	9.2	9.7	7.1-13.0	2.4	2.1	1.1-3.6
	2006-2010	12.8	5.2	4.0-6.7	10.4	9.2	6.8-12.2	2.4	1.7	0.9-3.1	10.2	4.1	3.1-5.5	8.8	8.1	5.8-10.9	1.4	1.1	0.4-2.3
	2011-2015	17.8	6.4	5.1-7.9	14.8	11.6	9.1-14.8	3.0	2.0	1.1-3.4	13.8	5.0	3.8-6.3	10.0	8.1	5.9-10.8	3.8	2.4	1.4-3.8
2016-2020	19.8	6.1	5.0-7.5	15.4	10.3	8.1-13.0	4.4	2.5	1.6-3.9	12.2	3.8	2.9-4.9	9.8	6.7	4.9-8.9	2.4	1.3	0.7-2.4	
Stomach	1976-1980	11.8	11.1	8.3-14.3	6.4	13.6	9.0-19.5	5.4	9.0	5.9-13.0	6.8	6.5	4.4-9.1	3.2	7.1	3.8-11.9	3.6	5.9	3.5-9.3
	1981-1985	12.6	10.5	8.0-13.5	7.8	15.1	10.5-20.9	4.8	7.0	4.5-10.4	10.4	8.7	6.5-11.5	6.2	11.9	7.9-17.2	4.2	6.2	3.8-9.4
	1986-1990	10.4	7.4	5.5-9.7	7.6	14.2	9.7-19.8	2.8	3.4	1.9-5.8	9.8	7.1	5.2-9.4	6.6	11.2	7.5-16.0	3.2	4.0	2.3-6.6
	1991-1995	10.0	6.3	4.6-8.3	7.0	9.6	6.6-13.4	3.0	3.3	1.9-5.5	6.4	4.0	2.7-5.6	4.2	5.9	3.6-9.2	2.2	2.4	1.2-4.3
	1996-2000	14.2	7.9	6.1-9.9	10.4	13.1	9.7-17.3	3.8	3.7	2.2-5.8	8.2	4.6	3.3-6.3	5.2	6.6	4.2-9.8	3.0	2.9	1.6-4.8
	2001-2005	13.8	6.5	5.1-8.3	9.2	9.4	6.9-12.6	4.6	3.8	2.4-5.8	8.6	4.2	3.0-5.6	5.2	5.5	3.5-8.1	3.4	2.8	1.6-4.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Saratoga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	16.0	6.7	5.3-8.4	10.2	10.0	7.3-13.2	5.8	4.3	2.8-6.2	8.0	3.3	2.4-4.6	5.8	5.7	3.8-8.3	2.2	1.5	0.8-2.8
	2011-2015	13.8	4.7	3.6-6.0	9.6	7.4	5.4-9.9	4.2	2.8	1.7-4.4	7.0	2.4	1.6-3.3	4.4	3.4	2.1-5.3	2.6	1.7	0.9-2.9
	2016-2020	17.2	5.4	4.3-6.7	11.0	7.4	5.5-9.7	6.2	3.6	2.4-5.2	6.6	2.1	1.4-3.0	4.0	2.9	1.7-4.5	2.6	1.5	0.8-2.7
Colorectal	1976-1980	72.2	65.8	59.0-73.2	35.6	78.4	66.4-91.6	36.6	57.6	49.5-66.7	35.2	33.3	28.5-38.7	16.8	38.7	30.3-48.4	18.4	29.6	23.8-36.4
	1981-1985	96.6	78.6	71.6-86.1	48.4	95.7	83.1-109.4	48.2	67.7	59.4-76.9	42.0	36.2	31.4-41.6	20.2	45.5	36.4-55.9	21.8	31.4	25.7-37.9
	1986-1990	87.4	63.3	57.4-69.6	44.0	74.5	64.4-85.6	43.4	55.3	48.1-63.2	42.2	31.4	27.2-35.9	21.6	38.7	31.3-47.2	20.6	26.4	21.5-32.0
	1991-1995	93.0	58.0	52.8-63.6	52.2	76.9	67.4-87.2	40.8	44.4	38.5-50.9	41.2	26.5	23.0-30.4	22.0	35.5	28.8-43.2	19.2	21.3	17.2-26.0
	1996-2000	112.2	62.8	57.7-68.2	54.2	71.6	63.0-81.0	58.0	56.3	50.0-63.2	41.6	23.6	20.5-27.0	22.8	31.6	25.8-38.2	18.8	18.1	14.6-22.1
	2001-2005	126.2	60.1	55.5-65.0	59.6	65.1	57.8-73.2	66.6	56.8	50.8-63.3	38.4	18.7	16.1-21.5	17.6	20.4	16.2-25.2	20.8	17.2	14.0-20.8
	2006-2010	107.8	45.2	41.4-49.2	52.8	49.9	44.0-56.5	55.0	41.8	36.9-47.1	33.6	14.1	12.0-16.4	16.0	15.9	12.5-19.9	17.6	12.7	10.1-15.7
	2011-2015	111.8	41.2	37.7-44.8	57.4	46.7	41.3-52.6	54.4	36.9	32.5-41.7	33.8	12.5	10.7-14.6	17.0	14.7	11.7-18.3	16.8	10.7	8.5-13.4
2016-2020	113.4	37.8	34.7-41.2	58.6	43.0	38.0-48.4	54.8	33.3	29.3-37.7	40.6	13.0	11.2-15.0	21.2	15.4	12.5-18.7	19.4	10.8	8.7-13.4	
Colon excluding rectum	1976-1980	49.6	45.8	40.1-52.0	22.4	51.0	41.3-62.1	27.2	42.9	35.9-50.8	29.4	27.9	23.5-32.9	13.2	30.3	23.0-39.1	16.2	26.2	20.8-32.6
	1981-1985	65.0	53.9	48.1-60.2	30.4	61.6	51.6-72.9	34.6	49.2	42.1-57.2	35.6	31.0	26.5-36.0	17.4	40.1	31.5-50.1	18.2	26.2	21.1-32.2
	1986-1990	66.2	49.0	43.7-54.6	32.4	58.2	49.1-68.5	33.8	43.0	36.8-50.1	37.0	27.8	23.9-32.2	19.4	35.8	28.6-44.2	17.6	22.6	18.1-27.9
	1991-1995	65.4	41.1	36.8-45.9	34.6	51.2	43.5-59.7	30.8	33.5	28.4-39.2	34.2	22.2	18.9-25.8	17.8	29.4	23.3-36.6	16.4	18.2	14.4-22.6
	1996-2000	82.4	46.4	42.0-51.2	36.0	48.1	41.1-55.9	46.4	45.0	39.4-51.2	35.6	20.3	17.4-23.5	19.2	27.0	21.6-33.1	16.4	15.8	12.5-19.6
	2001-2005	90.0	43.3	39.4-47.5	41.0	45.3	39.2-52.1	49.0	41.9	36.8-47.5	31.0	15.1	12.8-17.7	12.6	14.6	11.1-18.8	18.4	15.2	12.3-18.8
	2006-2010	79.4	33.7	30.4-37.2	36.6	35.5	30.4-41.2	42.8	32.7	28.4-37.5	27.2	11.6	9.7-13.7	11.8	12.0	9.1-15.5	15.4	11.3	8.8-14.2
	2011-2015	76.4	28.3	25.4-31.3	38.2	31.2	26.8-36.1	38.2	25.8	22.2-29.9	27.0	10.0	8.4-11.9	13.2	11.6	8.9-14.8	13.8	8.7	6.7-11.1
2016-2020	81.4	27.0	24.4-29.8	40.4	29.7	25.7-34.3	41.0	24.4	21.1-28.2	29.2	9.2	7.8-10.9	15.2	10.9	8.5-13.7	14.0	7.9	6.1-10.1	
Rectum & rectosigmoid	1976-1980	22.6	20.0	16.4-24.2	13.2	27.4	20.8-35.3	9.4	14.7	10.8-19.6	5.8	5.4	3.6-7.8	3.6	8.4	4.7-13.4	2.2	3.4	1.7-6.1
	1981-1985	31.6	24.7	20.9-29.0	18.0	34.0	26.9-42.4	13.6	18.5	14.3-23.5	6.4	5.2	3.5-7.4	2.8	5.3	2.7-9.2	3.6	5.1	3.0-8.1
	1986-1990	21.2	14.4	11.7-17.4	11.6	16.2	12.3-21.1	9.6	12.2	9.0-16.2	5.2	3.5	2.3-5.2	2.2	2.9	1.4-5.2	3.0	3.8	2.1-6.2
	1991-1995	27.6	16.9	14.2-20.0	17.6	25.7	20.4-31.9	10.0	10.9	8.1-14.4	7.0	4.3	3.0-6.0	4.2	6.1	3.6-9.5	2.8	3.1	1.7-5.2
	1996-2000	29.8	16.3	13.8-19.2	18.2	23.5	18.8-29.1	11.6	11.4	8.6-14.7	6.0	3.3	2.2-4.7	3.6	4.6	2.7-7.4	2.4	2.3	1.2-4.0
	2001-2005	36.2	16.8	14.4-19.5	18.6	19.8	15.9-24.4	17.6	14.9	12.0-18.4	7.4	3.5	2.5-4.9	5.0	5.8	3.7-8.6	2.4	1.9	1.0-3.4
	2006-2010	28.4	11.5	9.7-13.6	16.2	14.4	11.4-18.1	12.2	9.0	6.9-11.7	6.4	2.5	1.7-3.6	4.2	3.9	2.4-6.1	2.2	1.5	0.7-2.7
	2011-2015	35.4	12.9	11.0-15.0	19.2	15.5	12.5-19.1	16.2	11.0	8.7-13.8	6.8	2.5	1.7-3.5	3.8	3.1	1.9-5.0	3.0	2.0	1.1-3.4
2016-2020	32.0	10.8	9.1-12.7	18.2	13.2	10.6-16.4	13.8	8.9	6.8-11.4	11.4	3.7	2.8-4.9	6.0	4.5	3.0-6.5	5.4	3.0	1.9-4.5	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Saratoga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	3.0	2.4	1.3-3.9	1.6	2.8	1.2-5.5	1.4	2.0	0.8-4.1	2.4	1.8	0.9-3.2	1.6	2.5	1.1-5.1	0.8	1.1	0.3-2.9
	1981-1985	3.4	2.5	1.4-4.1	2.4	3.8	1.9-6.9	1.0	1.4	0.4-3.2	4.0	3.2	1.9-5.0	2.2	4.0	1.9-7.4	1.8	2.6	1.2-4.9
	1986-1990	6.2	4.3	2.9-6.1	3.6	5.5	3.2-8.8	2.6	3.3	1.7-5.6	6.6	4.6	3.2-6.5	4.6	7.5	4.7-11.4	2.0	2.4	1.2-4.5
	1991-1995	6.0	3.7	2.5-5.3	4.0	5.7	3.4-8.9	2.0	2.2	1.0-4.0	4.4	2.8	1.7-4.2	3.2	4.8	2.7-7.9	1.2	1.4	0.5-3.0
	1996-2000	10.0	5.5	4.0-7.2	7.8	9.6	6.7-13.2	2.2	2.1	1.0-3.8	8.4	4.8	3.4-6.4	6.0	8.0	5.3-11.5	2.4	2.3	1.2-4.0
	2001-2005	9.6	4.5	3.3-6.0	5.8	6.2	4.1-9.0	3.8	3.1	1.8-4.9	5.8	2.8	1.9-4.0	3.2	3.6	2.0-5.9	2.6	2.0	1.1-3.5
	2006-2010	12.2	5.2	3.9-6.7	9.2	8.4	6.1-11.3	3.0	2.3	1.3-3.9	11.6	4.8	3.6-6.2	7.8	7.7	5.4-10.5	3.8	2.7	1.6-4.3
	2011-2015	18.6	6.6	5.3-8.2	14.6	10.7	8.3-13.6	4.0	3.2	1.9-5.0	13.2	4.8	3.7-6.1	9.4	7.1	5.2-9.6	3.8	2.7	1.6-4.3
2016-2020	26.8	8.2	6.8-9.7	20.2	13.1	10.6-16.1	6.6	3.8	2.6-5.5	15.0	4.5	3.6-5.8	11.8	7.7	5.8-10.0	3.2	1.9	1.1-3.2	
Pancreas	1976-1980	16.2	14.2	11.2-17.7	9.2	19.5	13.8-26.4	7.0	10.8	7.5-15.0	14.4	13.1	10.2-16.6	7.6	16.7	11.4-23.5	6.8	10.9	7.5-15.2
	1981-1985	16.2	13.2	10.4-16.4	9.4	17.5	12.5-23.6	6.8	10.0	6.9-14.0	15.6	12.6	9.9-15.8	8.8	16.5	11.7-22.5	6.8	9.9	6.8-13.8
	1986-1990	16.6	11.9	9.4-14.7	7.6	12.4	8.6-17.2	9.0	11.2	8.1-14.9	17.6	12.5	10.0-15.4	8.4	13.5	9.6-18.4	9.2	11.3	8.2-15.0
	1991-1995	21.8	13.9	11.4-16.8	10.0	14.6	10.7-19.4	11.8	13.1	10.0-16.9	19.0	12.4	10.1-15.2	8.2	12.7	8.9-17.3	10.8	11.9	9.0-15.6
	1996-2000	23.0	12.8	10.5-15.3	11.6	14.3	10.8-18.6	11.4	11.2	8.4-14.5	22.0	12.3	10.1-14.8	10.4	13.0	9.6-17.1	11.6	11.3	8.6-14.6
	2001-2005	29.6	14.2	12.0-16.7	16.0	17.8	14.0-22.2	13.6	11.3	8.8-14.4	24.0	11.7	9.7-14.0	13.6	15.5	12.0-19.8	10.4	8.5	6.4-11.3
	2006-2010	32.8	13.6	11.5-15.9	16.6	15.3	12.1-19.1	16.2	12.1	9.6-15.2	27.2	11.2	9.4-13.3	13.0	11.8	9.0-15.2	14.2	10.6	8.2-13.4
	2011-2015	40.4	14.9	12.8-17.1	21.6	17.2	14.1-21.0	18.8	12.6	10.2-15.6	30.8	11.6	9.8-13.6	16.6	13.6	10.8-17.0	14.2	9.8	7.6-12.5
2016-2020	41.8	13.0	11.2-14.9	24.6	16.6	13.7-20.0	17.2	9.9	7.9-12.4	31.6	9.7	8.3-11.5	16.4	10.9	8.6-13.7	15.2	8.7	6.8-11.1	
Larynx	1976-1980	7.2	5.9	4.1-8.3	6.6	12.3	8.3-17.4	0.6	0.8	0.2-2.5	2.8	2.5	1.3-4.2	2.4	5.3	2.5-9.4	0.4	0.7	0.1-2.3
	1981-1985	8.2	6.1	4.3-8.3	6.4	10.5	7.0-15.1	1.8	2.4	1.1-4.6	2.6	1.9	1.0-3.3	2.0	3.0	1.4-5.6	0.6	0.9	0.2-2.5
	1986-1990	9.2	6.3	4.6-8.4	7.6	11.4	8.0-15.9	1.6	2.1	0.9-4.0	2.4	1.8	0.9-3.1	2.0	3.5	1.5-6.6	0.4	0.5	0.1-1.9
	1991-1995	7.4	4.4	3.1-6.1	5.2	7.3	4.7-10.9	2.2	2.4	1.2-4.3	2.8	1.7	0.9-2.8	2.2	3.2	1.5-5.8	0.6	0.6	0.1-1.8
	1996-2000	7.2	3.8	2.7-5.3	5.8	6.6	4.4-9.6	1.4	1.4	0.6-2.8	2.4	1.4	0.7-2.4	1.4	1.9	0.7-3.9	1.0	1.0	0.3-2.3
	2001-2005	9.0	4.2	3.0-5.6	6.4	6.6	4.4-9.4	2.6	2.3	1.2-3.9	2.2	1.0	0.5-1.8	1.6	1.7	0.7-3.5	0.6	0.5	0.1-1.4
	2006-2010	11.0	4.5	3.4-5.9	8.6	7.8	5.6-10.5	2.4	1.8	0.9-3.2	3.8	1.6	0.9-2.5	3.2	2.8	1.6-4.7	0.6	0.5	0.1-1.4
	2011-2015	10.6	3.6	2.7-4.8	7.8	5.7	4.0-7.9	2.8	1.8	0.9-3.1	2.6	1.0	0.5-1.7	2.6	2.1	1.1-3.7	0.0	0.0	0.0-0.7
2016-2020	9.2	2.9	2.1-3.9	7.4	4.9	3.4-6.9	1.8	1.2	0.5-2.4	2.0	0.6	0.3-1.1	1.0	0.6	0.2-1.5	1.0	0.5	0.2-1.3	
Lung and bronchus	1976-1980	71.6	60.1	53.9-66.8	52.0	102.1	89.3-116.1	19.6	29.6	24.0-36.2	61.6	52.5	46.7-58.8	45.2	90.2	78.1-103.6	16.4	25.3	20.1-31.4
	1981-1985	90.2	69.1	62.8-75.9	61.0	107.6	95.2-121.1	29.2	40.6	34.2-47.8	78.4	60.6	54.6-67.0	54.4	99.4	87.2-112.7	24.0	33.2	27.5-39.8
	1986-1990	101.6	68.7	62.8-75.1	67.4	105.3	93.8-117.7	34.2	41.8	35.7-48.6	84.0	57.7	52.2-63.5	56.4	90.6	79.8-102.4	27.6	33.7	28.3-39.9
	1991-1995	124.6	75.7	69.8-81.9	76.2	105.0	94.3-116.5	48.4	53.1	46.6-60.3	99.6	60.8	55.6-66.5	63.8	90.4	80.4-101.3	35.8	39.3	33.8-45.6
	1996-2000	140.6	76.4	70.9-82.3	75.2	94.5	84.9-104.8	65.4	64.8	57.9-72.2	110.8	61.1	56.1-66.4	58.4	75.7	67.0-85.2	52.4	51.7	45.6-58.4
	2001-2005	159.6	75.8	70.6-81.3	84.6	90.8	82.1-100.1	75.0	64.6	58.1-71.5	116.0	55.4	50.9-60.1	61.6	66.7	59.2-74.7	54.4	46.9	41.5-52.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Saratoga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	187.8	78.5	73.5-83.8	96.4	91.1	83.0-99.9	91.4	69.9	63.5-76.7	127.2	54.1	49.9-58.6	67.6	66.2	59.2-73.8	59.6	45.5	40.4-51.1
	2011-2015	196.2	70.8	66.3-75.5	98.0	79.0	72.0-86.6	98.2	64.8	59.1-71.0	134.4	49.4	45.6-53.3	71.2	58.3	52.2-64.9	63.2	42.6	37.9-47.7
	2016-2020	213.2	65.5	61.6-69.7	102.6	69.6	63.6-76.1	110.6	63.2	58.0-68.9	124.4	38.9	35.9-42.2	60.0	42.3	37.5-47.5	64.4	37.0	33.0-41.4
Melanoma of the skin	1976-1980	9.0	6.9	4.9-9.2	4.8	7.9	5.0-12.0	4.2	5.8	3.5-8.9	3.2	2.7	1.5-4.5	2.2	4.5	2.2-8.1	1.0	1.5	0.5-3.4
	1981-1985	12.2	8.2	6.3-10.6	6.4	9.1	6.2-13.0	5.8	7.4	4.9-10.7	4.0	2.6	1.6-4.1	3.2	4.3	2.5-7.2	0.8	0.9	0.2-2.4
	1986-1990	13.0	8.4	6.4-10.7	6.8	9.4	6.3-13.3	6.2	7.4	5.0-10.5	3.6	2.3	1.4-3.7	1.4	1.9	0.7-3.9	2.2	2.6	1.3-4.8
	1991-1995	15.8	8.9	7.0-11.1	9.4	11.0	8.0-14.7	6.4	7.0	4.8-9.8	4.2	2.4	1.5-3.7	2.4	2.9	1.5-5.2	1.8	2.0	0.9-3.7
	1996-2000	23.6	12.1	10.0-14.5	13.0	14.1	10.8-18.1	10.6	10.3	7.7-13.5	5.0	2.6	1.7-3.8	2.2	2.3	1.1-4.2	2.8	2.7	1.5-4.6
	2001-2005	33.0	15.2	12.9-17.7	18.6	19.1	15.4-23.5	14.4	12.6	9.8-15.9	4.6	2.1	1.3-3.1	3.2	3.1	1.8-5.2	1.4	1.2	0.5-2.6
	2006-2010	52.2	21.2	18.7-24.0	29.2	26.0	21.8-30.7	23.0	18.1	14.9-21.8	6.6	2.8	1.9-4.0	4.8	4.8	3.0-7.2	1.8	1.3	0.6-2.6
	2011-2015	74.0	27.6	24.8-30.7	47.8	38.8	33.9-44.2	26.2	18.6	15.4-22.2	10.2	3.8	2.8-5.0	6.8	5.4	3.7-7.6	3.4	2.5	1.4-4.1
2016-2020	76.8	25.7	23.1-28.5	43.0	30.4	26.3-34.9	33.8	22.3	18.9-26.1	8.4	2.8	2.0-3.8	6.2	4.5	3.0-6.5	2.2	1.3	0.6-2.5	
Female breast	1976-1980							58.6	92.1	81.7-103.5							20.8	31.6	25.8-38.4
	1981-1985							70.8	100.6	90.2-111.7							23.4	33.5	27.6-40.1
	1986-1990							95.0	117.8	107.4-129.0							31.6	38.6	32.8-45.2
	1991-1995							116.4	127.6	117.4-138.5							30.2	33.4	28.3-39.2
	1996-2000							157.4	153.2	142.7-164.4							29.4	28.6	24.2-33.7
	2001-2005							164.8	140.2	130.8-150.2							27.2	22.8	19.1-27.0
	2006-2010							179.2	135.8	126.9-145.2							32.8	24.4	20.8-28.6
	2011-2015							196.8	135.3	126.8-144.3							28.4	18.1	15.2-21.5
2016-2020							230.0	145.9	137.3-155.1							29.4	17.3	14.5-20.4	
Cervix uteri	1976-1980							9.2	14.2	10.3-18.9							3.0	5.0	2.8-8.2
	1981-1985							7.4	10.0	7.0-13.9							2.6	3.6	1.9-6.2
	1986-1990							6.6	8.0	5.5-11.2							2.4	3.2	1.6-5.6
	1991-1995							6.4	6.7	4.6-9.4							1.4	1.5	0.6-3.2
	1996-2000							10.2	9.6	7.1-12.6							1.8	1.8	0.8-3.4
	2001-2005							9.0	7.8	5.7-10.4							2.6	2.2	1.2-3.9
	2006-2010							8.8	7.5	5.4-10.1							2.2	1.8	0.9-3.3
	2011-2015							6.4	5.3	3.5-7.5							1.2	0.9	0.3-2.0
2016-2020							6.2	5.0	3.3-7.2							1.6	1.1	0.5-2.3	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Saratoga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							13.8	21.2	16.5-26.9							3.8	6.3	3.8-9.9
	1981-1985							15.4	21.5	16.9-26.9							4.4	6.5	4.1-9.8
	1986-1990							17.6	22.0	17.6-27.2							3.6	4.6	2.7-7.3
	1991-1995							16.6	18.2	14.4-22.5							3.4	3.7	2.2-5.9
	1996-2000							29.2	28.6	24.2-33.7							6.0	5.8	3.9-8.3
	2001-2005							31.6	26.4	22.4-30.9							5.8	4.9	3.3-7.1
	2006-2010							32.0	23.2	19.7-27.2							3.0	2.4	1.3-4.0
	2011-2015							49.4	31.5	27.6-35.8							5.4	3.3	2.2-4.9
	2016-2020							45.6	25.9	22.6-29.7							9.4	5.3	3.8-7.1
Ovary	1976-1980							9.4	14.6	10.6-19.4							7.0	11.2	7.8-15.6
	1981-1985							10.4	14.3	10.7-18.9							6.2	8.6	5.8-12.2
	1986-1990							12.6	15.9	12.2-20.4							8.6	10.9	7.9-14.7
	1991-1995							14.4	15.6	12.2-19.7							6.8	7.5	5.2-10.5
	1996-2000							18.2	17.8	14.3-21.9							10.6	10.4	7.8-13.6
	2001-2005							14.8	12.6	9.9-15.9							12.6	10.6	8.1-13.5
	2006-2010							18.2	13.7	11.0-16.9							12.2	9.1	6.9-11.8
	2011-2015							16.4	10.9	8.7-13.7							8.6	5.8	4.2-7.9
	2016-2020							16.6	10.0	7.9-12.6							9.6	5.2	3.8-7.1
Prostate	1976-1980				30.0	79.2	66.2-93.8							12.8	37.9	28.8-48.7			
	1981-1985				38.4	81.7	69.7-94.9							12.4	30.8	23.1-39.8			
	1986-1990				57.8	103.2	90.9-116.5							16.6	35.7	27.9-44.8			
	1991-1995				94.2	138.2	125.4-151.8							21.2	38.5	31.1-46.9			
	1996-2000				128.0	158.9	146.5-172.1							19.0	30.0	24.0-36.8			
	2001-2005				159.8	165.1	153.6-177.3							19.8	26.5	21.4-32.3			
	2006-2010				180.2	153.6	143.4-164.3							16.4	18.0	14.2-22.3			
	2011-2015				161.6	118.0	109.8-126.8							19.4	18.6	15.0-22.7			
	2016-2020				198.0	123.6	115.8-131.8							22.6	18.5	15.2-22.3			
Testis	1976-1980				2.6	3.3	1.7-5.8							0.6	0.7	0.1-2.4			
	1981-1985				5.0	6.2	3.9-9.4							0.4	0.4	0.1-1.9			
	1986-1990				4.8	4.6	2.9-7.0							0.2	0.2	0.0-1.4			
	1991-1995				6.4	6.2	4.2-8.9							0.6	0.6	0.1-1.9			
	1996-2000				6.4	6.2	4.2-8.8							0.0	0.0	0.0-0.9			
	2001-2005				6.2	6.0	4.1-8.6							0.2	0.2	0.0-1.1			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Saratoga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				10.4	10.5	7.8-13.8							0.0	0.0	0.0-0.7			
	2011-2015				10.4	9.6	7.2-12.7							0.2	0.1	0.0-0.9			
	2016-2020				8.4	7.7	5.5-10.5							0.2	0.1	0.0-0.8			
Urinary bladder (incl. in situ)	1976-1980	25.8	23.8	19.8-28.4	20.0	47.3	37.9-58.2	5.8	9.0	6.0-13.0	7.2	7.2	5.0-10.0	5.6	15.7	10.1-22.9	1.6	2.4	1.0-4.8
	1981-1985	27.8	22.2	18.6-26.3	21.6	40.6	32.9-49.5	6.2	8.9	6.0-12.6	6.6	6.0	4.1-8.4	5.0	12.2	7.8-18.1	1.6	2.4	1.0-4.7
	1986-1990	37.6	26.2	22.5-30.3	26.8	44.3	36.7-53.0	10.8	13.1	9.8-17.1	7.6	5.8	4.1-7.9	5.4	10.2	6.5-15.0	2.2	2.8	1.4-4.9
	1991-1995	36.2	22.4	19.2-25.9	25.8	38.5	31.8-46.0	10.4	11.5	8.6-15.1	8.2	5.3	3.8-7.1	5.0	8.2	5.2-12.2	3.2	3.5	2.0-5.7
	1996-2000	46.2	25.4	22.2-28.9	35.0	44.3	37.8-51.6	11.2	10.8	8.2-14.1	11.6	6.6	5.0-8.6	8.0	11.4	8.0-15.6	3.6	3.4	2.0-5.5
	2001-2005	70.8	33.6	30.2-37.3	52.8	58.3	51.3-66.0	18.0	15.5	12.4-19.1	10.6	5.2	3.9-6.9	7.8	10.0	7.1-13.7	2.8	2.3	1.3-4.0
	2006-2010	72.0	29.9	26.8-33.2	54.6	51.9	45.8-58.6	17.4	12.9	10.3-16.0	12.2	5.2	3.9-6.6	8.0	8.5	6.0-11.6	4.2	2.9	1.8-4.6
	2011-2015	87.4	32.3	29.3-35.6	64.8	54.5	48.5-60.9	22.6	15.0	12.3-18.1	12.2	4.6	3.5-5.9	8.2	7.5	5.3-10.2	4.0	2.7	1.6-4.2
2016-2020	85.6	27.1	24.6-29.9	67.8	48.0	42.9-53.5	17.8	10.0	8.0-12.4	12.0	3.8	2.9-4.9	8.4	6.2	4.5-8.5	3.6	2.0	1.2-3.4	
Kidney and renal pelvis	1976-1980	8.4	6.9	4.9-9.3	5.0	9.3	5.8-13.9	3.4	5.3	3.0-8.4	5.6	4.7	3.1-6.9	3.4	6.3	3.5-10.2	2.2	3.4	1.7-6.0
	1981-1985	7.6	5.9	4.2-8.2	5.0	9.1	5.7-13.7	2.6	3.6	1.9-6.1	3.4	2.7	1.6-4.3	1.8	3.2	1.4-6.2	1.6	2.3	1.0-4.5
	1986-1990	17.2	11.7	9.3-14.5	11.4	16.8	12.6-21.9	5.8	7.3	4.9-10.4	4.8	3.4	2.2-5.1	3.6	5.9	3.4-9.5	1.2	1.6	0.6-3.4
	1991-1995	17.6	10.5	8.4-13.0	11.4	15.0	11.2-19.6	6.2	6.9	4.7-9.8	6.6	4.2	2.9-5.9	3.4	5.1	2.8-8.3	3.2	3.5	2.0-5.7
	1996-2000	25.4	13.5	11.2-16.1	16.6	19.8	15.7-24.6	8.8	8.6	6.3-11.6	7.8	4.3	3.0-5.9	4.4	5.7	3.5-8.7	3.4	3.4	2.0-5.4
	2001-2005	30.6	14.4	12.2-16.9	18.2	18.7	15.0-23.1	12.4	10.5	8.0-13.5	9.2	4.4	3.2-5.8	5.8	6.0	4.0-8.7	3.4	2.9	1.7-4.7
	2006-2010	46.0	18.4	16.1-21.0	29.6	25.7	21.6-30.4	16.4	12.5	9.9-15.6	10.6	4.5	3.4-5.9	7.2	7.2	5.0-10.0	3.4	2.5	1.4-4.0
	2011-2015	57.0	21.1	18.6-23.7	36.6	28.9	24.7-33.5	20.4	14.2	11.5-17.4	12.4	4.4	3.4-5.7	7.4	5.8	4.0-8.1	5.0	3.2	2.1-4.8
2016-2020	54.6	17.9	15.8-20.3	36.4	25.3	21.6-29.5	18.2	11.5	9.1-14.3	9.2	3.0	2.2-4.1	6.0	4.3	2.9-6.3	3.2	2.0	1.1-3.3	
Brain and other central nervous system	1976-1980	9.2	6.8	4.9-9.2	5.6	8.6	5.5-12.7	3.6	5.2	3.1-8.4	9.8	7.4	5.4-9.9	5.8	9.0	5.9-13.3	4.0	5.9	3.6-9.2
	1981-1985	11.8	8.2	6.2-10.6	5.6	7.6	5.0-11.2	6.2	8.4	5.7-11.9	6.4	4.7	3.2-6.6	3.4	4.6	2.7-7.6	3.0	4.2	2.3-6.9
	1986-1990	11.4	7.1	5.3-9.2	7.2	9.7	6.7-13.6	4.2	4.8	2.9-7.3	9.0	5.9	4.3-7.9	5.0	7.2	4.6-10.8	4.0	4.6	2.8-7.1
	1991-1995	14.2	8.0	6.2-10.1	8.0	9.4	6.7-12.9	6.2	6.5	4.4-9.3	8.0	4.6	3.3-6.3	4.8	5.9	3.8-8.9	3.2	3.4	1.9-5.5
	1996-2000	17.4	9.1	7.3-11.2	8.8	9.6	6.9-13.0	8.6	8.4	6.1-11.4	9.4	4.9	3.6-6.6	6.2	6.8	4.6-9.8	3.2	3.1	1.8-5.1
	2001-2005	13.2	6.1	4.7-7.7	6.4	6.2	4.2-8.8	6.8	5.9	4.0-8.2	9.0	4.2	3.0-5.6	2.8	2.8	1.5-4.7	6.2	5.5	3.7-7.8
	2006-2010	18.6	8.2	6.6-10.0	10.4	9.5	7.0-12.5	8.2	6.9	4.9-9.5	9.0	3.9	2.8-5.3	5.4	5.1	3.3-7.4	3.6	2.9	1.7-4.7
	2011-2015	19.4	7.7	6.2-9.5	12.0	10.1	7.7-13.1	7.4	5.4	3.7-7.6	13.2	4.8	3.7-6.1	8.6	6.5	4.7-8.9	4.6	3.1	1.9-4.8
2016-2020	19.0	7.0	5.6-8.6	12.6	9.1	6.9-11.8	6.4	5.1	3.4-7.3	13.2	4.4	3.4-5.7	8.6	5.9	4.2-8.1	4.6	3.0	1.9-4.7	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Saratoga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	2.4	1.6	0.8-3.0	0.0	0.0	0.0-1.4	2.4	3.1	1.5-5.6	0.6	0.7	0.1-1.9	0.0	0.0	0.0-1.4	0.6	1.1	0.2-3.0
	1981-1985	4.6	3.2	2.0-4.9	2.0	2.9	1.4-5.5	2.6	3.4	1.8-5.8	0.4	0.4	0.1-1.4	0.0	0.0	0.0-1.3	0.4	0.6	0.1-2.2
	1986-1990	7.2	4.3	3.0-6.0	3.4	4.5	2.6-7.3	3.8	4.3	2.6-6.8	0.4	0.3	0.0-0.9	0.0	0.0	0.0-1.1	0.4	0.5	0.1-1.7
	1991-1995	6.0	3.2	2.2-4.7	1.6	2.0	0.9-4.1	4.4	4.4	2.7-6.7	0.8	0.5	0.1-1.3	0.4	0.5	0.1-1.8	0.4	0.5	0.1-1.6
	1996-2000	18.0	9.0	7.2-11.0	4.4	4.6	2.8-7.0	13.6	13.3	10.3-16.9	0.4	0.2	0.0-0.7	0.2	0.2	0.0-1.3	0.2	0.2	0.0-1.1
	2001-2005	25.4	11.7	9.7-13.9	5.8	5.4	3.6-7.9	19.6	17.7	14.3-21.6	1.0	0.5	0.2-1.2	0.4	0.4	0.0-1.5	0.6	0.6	0.1-1.7
	2006-2010	41.8	17.4	15.1-20.0	10.8	9.5	7.1-12.4	31.0	25.3	21.4-29.7	0.2	0.1	0.0-0.5	0.2	0.3	0.0-1.3	0.0	0.0	0.0-0.7
	2011-2015	52.4	21.6	19.0-24.5	13.2	10.9	8.3-13.9	39.2	32.2	27.7-37.3	0.6	0.2	0.0-0.7	0.4	0.3	0.0-1.3	0.2	0.1	0.0-0.9
	2016-2020	50.6	19.9	17.5-22.7	14.4	11.0	8.5-14.0	36.2	29.0	24.8-33.8	1.8	0.6	0.3-1.1	1.2	0.9	0.3-2.0	0.6	0.3	0.1-1.2
Hodgkin lymphoma	1976-1980	6.4	4.6	3.1-6.6	3.0	4.3	2.3-7.2	3.4	4.6	2.6-7.5	1.2	1.0	0.3-2.1	0.2	0.4	0.0-2.1	1.0	1.4	0.4-3.4
	1981-1985	6.4	4.0	2.7-5.7	3.4	4.6	2.6-7.5	3.0	3.5	1.9-5.8	2.4	1.8	0.9-3.2	1.8	2.6	1.2-5.0	0.6	0.9	0.2-2.6
	1986-1990	4.4	2.4	1.5-3.7	2.2	2.3	1.2-4.4	2.2	2.4	1.2-4.4	2.0	1.4	0.6-2.5	0.8	1.3	0.3-3.3	1.2	1.5	0.5-3.2
	1991-1995	6.2	3.2	2.1-4.5	3.0	3.1	1.7-5.3	3.2	3.2	1.8-5.2	1.2	0.7	0.3-1.6	0.4	0.4	0.1-1.7	0.8	1.0	0.3-2.4
	1996-2000	5.4	2.6	1.7-3.9	3.2	3.3	1.9-5.4	2.2	2.1	1.0-3.7	0.2	0.1	0.0-0.5	0.0	0.0	0.0-0.9	0.2	0.2	0.0-1.1
	2001-2005	4.4	2.2	1.4-3.4	3.0	3.1	1.7-5.2	1.4	1.3	0.5-2.8	0.4	0.2	0.0-0.7	0.2	0.2	0.0-1.2	0.2	0.2	0.0-1.0
	2006-2010	6.4	2.9	2.0-4.1	3.4	3.3	1.9-5.3	3.0	2.7	1.5-4.5	0.2	0.1	0.0-0.6	0.0	0.0	0.0-0.7	0.2	0.2	0.0-1.2
	2011-2015	5.6	2.3	1.5-3.4	2.6	2.2	1.1-3.8	3.0	2.4	1.3-4.0	1.2	0.4	0.2-1.0	0.2	0.2	0.0-1.0	1.0	0.7	0.2-1.7
	2016-2020	7.2	3.1	2.1-4.3	3.4	3.1	1.8-5.1	3.8	3.1	1.8-5.0	0.6	0.2	0.0-0.7	0.4	0.3	0.0-1.1	0.2	0.2	0.0-1.2
Non-Hodgkin lymphomas	1976-1980	14.2	11.5	8.9-14.6	7.6	13.9	9.5-19.4	6.6	10.0	6.9-14.1	6.4	5.0	3.4-7.1	2.6	4.5	2.2-8.1	3.8	5.8	3.5-9.1
	1981-1985	18.2	14.3	11.5-17.6	9.4	15.2	11.0-20.6	8.8	12.6	9.1-16.9	10.2	8.3	6.1-11.0	5.0	9.3	5.8-13.9	5.2	7.5	4.9-10.9
	1986-1990	22.4	15.2	12.4-18.3	9.2	13.5	9.7-18.2	13.2	16.4	12.6-20.9	12.8	8.9	6.8-11.4	5.6	8.3	5.4-12.2	7.2	8.9	6.2-12.3
	1991-1995	25.6	15.1	12.5-17.9	14.2	17.4	13.4-22.1	11.4	12.6	9.5-16.3	15.2	9.2	7.3-11.6	7.2	9.6	6.5-13.5	8.0	8.7	6.2-11.8
	1996-2000	37.0	19.8	17.1-22.9	20.6	24.0	19.5-29.2	16.4	16.0	12.7-19.8	13.6	7.5	5.8-9.5	7.0	8.6	5.9-12.1	6.6	6.3	4.3-8.9
	2001-2005	49.8	23.4	20.6-26.5	25.0	25.8	21.4-30.9	24.8	21.6	17.9-25.7	13.2	6.4	4.9-8.2	6.4	7.2	4.9-10.3	6.8	5.6	3.9-7.9
	2006-2010	58.6	24.4	21.7-27.5	33.2	30.7	26.1-35.9	25.4	19.8	16.4-23.6	13.8	6.0	4.6-7.6	7.6	7.8	5.5-10.8	6.2	4.7	3.2-6.8
	2011-2015	65.6	25.3	22.5-28.2	34.4	29.3	24.9-34.2	31.2	22.4	18.9-26.3	16.4	6.4	5.0-8.0	9.2	8.2	6.0-11.1	7.2	5.0	3.5-7.1
	2016-2020	72.0	23.2	20.8-25.8	39.8	27.7	23.9-32.0	32.2	19.5	16.5-22.9	18.6	6.1	4.9-7.5	10.6	7.7	5.7-10.2	8.0	4.8	3.4-6.6
Myeloma	1976-1980	4.4	3.8	2.3-5.7	3.0	6.3	3.4-10.6	1.4	2.0	0.8-4.2	4.0	3.2	1.9-5.0	2.4	4.8	2.3-8.5	1.6	2.2	0.9-4.3
	1981-1985	5.2	4.2	2.8-6.2	3.4	6.5	3.6-10.5	1.8	2.6	1.2-4.9	3.4	2.9	1.7-4.7	2.0	4.3	2.0-8.0	1.4	2.0	0.8-4.1
	1986-1990	5.2	3.8	2.5-5.6	3.4	5.7	3.2-9.2	1.8	2.4	1.1-4.5	5.2	3.9	2.5-5.6	3.0	5.3	2.9-8.8	2.2	2.8	1.4-5.0
	1991-1995	6.4	4.0	2.7-5.6	2.8	3.5	1.9-5.9	3.6	3.9	2.3-6.2	4.4	2.8	1.7-4.2	2.0	2.7	1.3-5.1	2.4	2.7	1.4-4.6
	1996-2000	10.2	5.6	4.1-7.3	5.0	5.9	3.8-8.7	5.2	5.1	3.3-7.5	5.2	2.9	1.9-4.3	3.0	3.9	2.1-6.4	2.2	2.1	1.1-3.8
	2001-2005	12.2	6.0	4.6-7.8	8.2	9.3	6.6-12.6	4.0	3.5	2.1-5.4	7.4	3.6	2.5-5.0	4.2	4.6	2.8-7.0	3.2	2.6	1.5-4.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Saratoga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	14.8	6.1	4.7-7.6	8.2	7.8	5.5-10.7	6.6	4.9	3.3-6.9	8.0	3.4	2.4-4.6	4.4	4.1	2.5-6.3	3.6	2.8	1.7-4.5
	2011-2015	19.8	7.2	5.8-8.8	11.6	9.2	6.9-12.1	8.2	5.7	4.0-7.8	8.2	3.1	2.2-4.3	5.2	4.9	3.1-7.2	3.0	2.0	1.1-3.5
	2016-2020	24.8	7.9	6.5-9.4	15.6	10.3	8.1-12.9	9.2	5.8	4.2-7.9	8.0	2.5	1.8-3.4	5.6	4.1	2.7-6.0	2.4	1.4	0.7-2.5
Leukemias	1976-1980	15.6	12.8	10.0-16.1	8.4	16.2	11.2-22.4	7.2	10.5	7.3-14.6	9.2	7.8	5.6-10.5	4.4	8.6	5.1-13.4	4.8	7.4	4.7-11.1
	1981-1985	18.2	13.8	11.1-17.1	10.6	19.1	13.9-25.5	7.6	10.4	7.3-14.2	12.6	10.1	7.7-13.0	7.4	15.9	10.8-22.3	5.2	7.0	4.6-10.3
	1986-1990	12.6	8.5	6.5-10.9	5.4	8.9	5.6-13.3	7.2	8.6	6.0-12.0	10.8	7.2	5.4-9.4	5.6	9.1	5.8-13.4	5.2	6.1	4.0-8.9
	1991-1995	20.8	12.7	10.4-15.4	10.0	14.3	10.4-19.1	10.8	11.6	8.7-15.1	13.8	8.6	6.6-10.9	7.2	10.8	7.4-15.1	6.6	7.1	4.9-10.0
	1996-2000	21.2	11.5	9.4-13.9	12.0	14.3	10.8-18.5	9.2	9.0	6.6-12.1	11.0	6.2	4.7-8.1	3.8	5.1	3.0-8.0	7.2	7.1	4.9-9.8
	2001-2005	32.2	15.4	13.1-18.0	18.4	20.0	16.0-24.6	13.8	11.7	9.1-14.9	15.0	7.4	5.8-9.3	8.2	10.0	7.1-13.6	6.8	5.8	4.0-8.1
	2006-2010	38.4	15.9	13.7-18.4	22.8	20.4	16.7-24.6	15.6	12.1	9.5-15.2	15.0	6.6	5.2-8.3	8.8	8.4	6.1-11.4	6.2	4.9	3.3-7.0
	2011-2015	48.4	18.9	16.5-21.5	28.0	23.5	19.6-27.9	20.4	15.2	12.3-18.6	20.0	7.7	6.3-9.4	12.6	11.2	8.6-14.5	7.4	5.1	3.6-7.2
	2016-2020	45.2	14.9	13.0-17.1	26.6	19.0	15.8-22.7	18.6	11.4	9.1-14.2	16.4	5.6	4.4-7.0	9.6	7.6	5.5-10.2	6.8	3.9	2.7-5.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schenectady**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	697.6	431.3	416.7-446.2	348.8	531.8	505.8-558.7	348.8	380.2	362.0-399.2	359.4	221.8	211.4-232.6	187.4	292.0	272.6-312.4	172.0	182.0	169.7-194.9
	1981-1985	755.0	449.1	434.6-464.1	368.2	534.7	509.6-560.7	386.8	404.3	385.7-423.5	382.4	225.2	215.0-235.7	194.2	291.8	272.9-311.5	188.2	188.9	176.7-201.8
	1986-1990	765.6	446.0	431.6-460.7	365.6	524.5	499.9-549.9	400.0	413.3	394.6-432.8	382.4	218.1	208.3-228.4	189.6	276.8	258.9-295.6	192.8	187.2	175.1-199.9
	1991-1995	863.2	489.9	475.1-505.0	445.4	605.8	580.5-631.9	417.8	421.7	403.1-440.9	379.8	210.1	200.6-219.9	186.4	260.8	244.0-278.4	193.4	180.7	169.1-192.9
	1996-2000	854.4	485.5	470.8-500.5	429.6	573.1	549.1-598.0	424.8	428.5	409.8-447.8	389.8	209.7	200.3-219.3	189.4	255.5	239.4-272.4	200.4	181.8	170.3-194.0
	2001-2005	871.0	492.0	477.3-507.0	434.6	568.5	544.7-593.0	436.4	445.8	426.8-465.6	353.4	187.7	178.9-196.9	171.6	223.5	208.8-239.1	181.8	167.1	156.0-178.8
	2006-2010	908.0	497.0	482.3-512.0	456.2	569.6	546.1-593.8	451.8	450.5	431.5-470.1	350.8	180.1	171.6-189.0	187.0	233.4	218.5-249.0	163.8	145.3	135.1-156.1
	2011-2015	907.0	477.3	463.1-491.9	437.6	509.3	487.6-531.8	469.4	459.7	440.5-479.5	329.0	165.2	157.1-173.7	166.6	198.9	185.2-213.2	162.4	142.5	132.4-153.2
	2016-2020	960.0	481.5	467.6-495.9	477.2	522.3	500.9-544.4	482.8	457.7	438.8-477.2	316.2	150.0	142.5-157.8	153.2	169.8	157.7-182.7	163.0	137.7	128.0-147.9
Oral cavity and pharynx	1976-1980	20.8	12.1	9.9-14.8	12.0	16.7	12.6-21.8	8.8	8.9	6.4-12.0	9.4	5.8	4.2-7.8	5.4	8.5	5.4-12.5	4.0	4.1	2.5-6.4
	1981-1985	24.8	15.3	12.7-18.4	15.8	22.4	17.6-28.1	9.0	10.6	7.6-14.4	9.0	5.1	3.7-6.8	5.0	6.7	4.2-10.1	4.0	3.8	2.3-6.1
	1986-1990	20.0	11.3	9.2-13.9	13.0	17.8	13.6-23.0	7.0	6.9	4.8-9.8	7.2	4.2	2.9-5.8	4.6	6.7	4.2-10.2	2.6	2.6	1.3-4.6
	1991-1995	18.0	10.8	8.7-13.4	12.6	17.7	13.6-22.8	5.4	5.5	3.5-8.1	5.8	3.2	2.1-4.7	3.2	4.2	2.4-6.9	2.6	2.5	1.3-4.4
	1996-2000	18.2	10.8	8.6-13.3	13.0	17.4	13.4-22.2	5.2	5.4	3.5-8.1	4.8	2.6	1.7-3.9	3.4	4.6	2.7-7.3	1.4	1.3	0.5-2.8
	2001-2005	14.6	8.3	6.5-10.5	9.4	12.0	8.8-16.0	5.2	5.3	3.4-7.9	3.6	1.8	1.1-3.0	2.2	2.7	1.4-4.9	1.4	1.2	0.5-2.7
	2006-2010	19.4	10.5	8.5-12.9	15.0	17.6	13.8-22.2	4.4	4.6	2.8-7.1	2.2	1.1	0.6-2.1	2.0	2.3	1.1-4.3	0.2	0.1	0.0-1.1
	2011-2015	23.6	12.2	10.0-14.7	16.4	18.6	14.7-23.3	7.2	6.5	4.5-9.2	4.8	2.4	1.6-3.7	3.8	4.4	2.6-7.0	1.0	1.0	0.3-2.4
	2016-2020	25.4	12.8	10.6-15.3	18.2	19.9	15.9-24.6	7.2	6.6	4.5-9.3	6.4	3.2	2.1-4.6	4.0	4.2	2.5-6.7	2.4	2.1	1.1-3.9
Esophagus	1976-1980	6.8	4.0	2.7-5.7	6.0	8.9	5.8-12.9	0.8	0.8	0.2-2.1	6.2	3.6	2.4-5.2	4.6	6.7	4.1-10.3	1.6	1.6	0.7-3.3
	1981-1985	7.2	4.1	2.9-5.7	5.4	7.6	5.0-11.3	1.8	1.7	0.8-3.3	7.6	4.4	3.1-6.1	6.0	8.9	5.9-12.9	1.6	1.5	0.6-3.0
	1986-1990	5.6	3.3	2.1-4.8	4.0	5.3	3.2-8.3	1.6	1.5	0.6-3.2	5.8	3.1	2.1-4.6	4.4	5.5	3.4-8.5	1.4	1.4	0.5-3.0
	1991-1995	7.4	4.1	2.9-5.8	5.0	6.6	4.2-9.8	2.4	2.3	1.1-4.1	4.0	2.2	1.3-3.4	2.8	3.6	2.0-6.1	1.2	1.0	0.4-2.4
	1996-2000	11.8	6.8	5.1-8.8	9.0	12.1	8.9-16.3	2.8	2.4	1.3-4.3	11.0	6.2	4.6-8.1	8.2	11.1	7.9-15.0	2.8	2.2	1.2-3.9
	2001-2005	8.8	4.8	3.5-6.5	7.2	9.4	6.5-13.0	1.6	1.5	0.6-3.1	9.2	5.0	3.6-6.7	7.4	9.6	6.7-13.2	1.8	1.7	0.7-3.3
	2006-2010	11.0	6.0	4.5-7.9	8.6	10.8	7.8-14.7	2.4	2.2	1.1-4.1	9.4	5.2	3.8-7.0	7.6	9.8	6.9-13.6	1.8	1.7	0.8-3.5
	2011-2015	10.8	5.5	4.1-7.3	8.4	9.3	6.6-12.7	2.4	2.2	1.1-4.1	10.8	5.6	4.2-7.4	7.2	8.3	5.8-11.7	3.6	3.4	2.0-5.5
	2016-2020	10.2	5.2	3.8-6.9	7.0	8.3	5.7-11.6	3.2	3.0	1.7-5.1	8.4	4.2	3.0-5.8	6.4	7.2	4.9-10.3	2.0	1.8	0.9-3.5
Stomach	1976-1980	18.8	11.8	9.5-14.5	11.4	17.7	13.1-23.3	7.4	8.1	5.6-11.2	15.0	9.5	7.4-12.0	9.6	15.2	11.0-20.4	5.4	5.7	3.7-8.4
	1981-1985	17.8	10.4	8.3-12.8	11.8	18.0	13.5-23.4	6.0	5.9	3.9-8.5	12.6	7.3	5.6-9.4	8.0	12.3	8.6-17.0	4.6	4.4	2.8-6.8
	1986-1990	14.8	8.4	6.6-10.7	9.8	14.4	10.6-19.2	5.0	4.6	2.9-7.0	9.8	5.5	4.1-7.4	6.4	9.0	6.1-12.8	3.4	3.2	1.8-5.2
	1991-1995	15.8	9.0	7.1-11.2	7.6	10.3	7.2-14.2	8.2	7.6	5.3-10.5	10.8	6.1	4.5-8.0	5.8	8.2	5.4-11.8	5.0	4.7	3.0-7.1
	1996-2000	14.2	8.0	6.2-10.1	9.4	12.8	9.4-17.1	4.8	4.6	2.9-7.0	6.2	3.3	2.3-4.8	3.8	5.2	3.1-8.1	2.4	2.0	1.0-3.7
	2001-2005	15.6	8.4	6.6-10.6	9.0	11.7	8.5-15.7	6.6	5.7	3.8-8.2	8.2	4.4	3.2-6.1	4.2	5.4	3.4-8.3	4.0	3.6	2.1-5.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schenectady**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	13.0	7.0	5.4-9.0	9.0	11.4	8.3-15.3	4.0	3.7	2.3-5.9	7.2	3.7	2.6-5.2	4.2	5.3	3.3-8.2	3.0	2.6	1.4-4.4
	2011-2015	16.0	8.0	6.3-10.1	9.4	10.9	7.9-14.6	6.6	5.8	4.0-8.4	5.2	2.5	1.6-3.7	3.6	4.1	2.4-6.6	1.6	1.2	0.5-2.6
	2016-2020	15.2	7.4	5.8-9.4	9.4	10.4	7.6-14.0	5.8	5.0	3.3-7.4	3.8	1.7	1.0-2.8	2.4	2.6	1.3-4.6	1.4	1.1	0.4-2.5
Colorectal	1976-1980	121.0	73.6	67.7-79.8	58.4	88.7	78.3-100.0	62.6	64.8	57.7-72.6	58.4	36.0	31.9-40.5	27.2	43.5	36.1-51.9	31.2	32.0	27.1-37.5
	1981-1985	131.2	76.6	70.7-82.8	66.2	96.7	86.2-108.1	65.0	62.8	56.1-70.3	63.8	37.5	33.4-41.9	30.2	46.9	39.5-55.4	33.6	31.8	27.2-37.2
	1986-1990	108.8	60.4	55.4-65.8	52.2	72.4	63.6-82.1	56.6	53.6	47.4-60.6	51.6	28.4	25.0-32.2	25.0	36.3	30.0-43.5	26.6	24.1	20.1-28.7
	1991-1995	111.6	60.9	55.9-66.2	55.6	76.7	67.8-86.4	56.0	51.4	45.4-58.0	46.6	25.0	21.8-28.5	21.0	29.3	23.9-35.6	25.6	22.1	18.3-26.4
	1996-2000	101.8	54.7	50.0-59.8	46.4	61.9	54.2-70.5	55.4	49.1	43.4-55.6	44.4	22.5	19.6-25.7	20.0	26.6	21.6-32.4	24.4	19.7	16.3-23.8
	2001-2005	88.4	48.1	43.7-52.9	47.0	60.8	53.2-69.1	41.4	38.9	33.6-44.8	31.8	16.4	13.9-19.3	15.4	19.7	15.6-24.7	16.4	14.3	11.3-18.0
	2006-2010	91.8	49.1	44.6-53.9	43.4	54.6	47.5-62.5	48.4	44.8	39.1-51.2	35.0	17.5	14.9-20.4	17.4	21.9	17.5-27.1	17.6	13.7	10.8-17.1
	2011-2015	65.0	34.0	30.3-38.0	34.6	41.1	35.0-47.9	30.4	28.3	23.8-33.5	27.4	13.4	11.2-16.0	12.6	14.6	11.2-18.9	14.8	12.7	9.8-16.3
2016-2020	64.0	33.4	29.7-37.5	36.2	42.1	36.0-49.0	27.8	25.6	21.4-30.5	19.0	8.6	6.9-10.6	9.0	9.4	6.8-12.7	10.0	7.6	5.6-10.3	
Colon excluding rectum	1976-1980	86.8	52.9	48.0-58.2	41.0	63.1	54.3-72.8	45.8	47.2	41.2-53.9	49.2	30.3	26.5-34.4	23.2	36.6	29.9-44.3	26.0	26.7	22.2-31.8
	1981-1985	97.4	57.3	52.2-62.7	46.4	69.4	60.5-79.3	51.0	49.2	43.3-55.8	53.0	31.4	27.7-35.5	25.0	39.4	32.5-47.2	28.0	26.6	22.3-31.5
	1986-1990	79.6	44.5	40.1-49.1	37.6	52.0	44.6-60.2	42.0	40.0	34.6-46.0	41.6	23.0	19.9-26.4	19.8	28.6	23.1-35.1	21.8	19.8	16.2-24.0
	1991-1995	82.0	44.3	40.0-48.8	38.8	53.6	46.2-61.8	43.2	39.0	33.8-44.8	37.6	20.0	17.2-23.1	16.8	23.3	18.5-29.0	20.8	17.9	14.5-21.8
	1996-2000	75.8	40.5	36.5-44.9	35.4	47.4	40.6-54.9	40.4	35.6	30.7-41.1	35.8	18.1	15.5-21.0	16.8	22.4	17.9-27.8	19.0	15.3	12.3-18.9
	2001-2005	65.2	35.1	31.3-39.2	34.8	44.9	38.5-52.2	30.4	27.9	23.5-33.0	25.4	13.0	10.8-15.5	13.2	17.0	13.1-21.6	12.2	10.5	7.9-13.7
	2006-2010	64.8	34.5	30.7-38.6	30.0	38.1	32.1-44.8	34.8	31.8	27.1-37.2	28.8	14.2	11.9-16.9	13.8	17.4	13.5-22.1	15.0	11.4	8.9-14.6
	2011-2015	45.6	24.1	20.9-27.5	21.8	26.5	21.7-32.2	23.8	22.1	18.2-26.8	22.2	10.9	8.9-13.3	9.8	11.5	8.5-15.4	12.4	10.6	8.0-14.0
2016-2020	43.0	22.1	19.2-25.4	23.2	26.8	22.0-32.4	19.8	17.9	14.4-22.0	14.2	6.5	5.0-8.2	6.4	6.8	4.6-9.7	7.8	5.8	4.1-8.2	
Rectum & rectosigmoid	1976-1980	34.2	20.7	17.6-24.1	17.4	25.6	20.2-31.9	16.8	17.6	14.0-22.0	9.2	5.7	4.2-7.7	4.0	6.9	4.0-10.8	5.2	5.3	3.4-7.8
	1981-1985	33.8	19.3	16.4-22.5	19.8	27.3	22.0-33.5	14.0	13.6	10.5-17.3	10.8	6.1	4.5-8.0	5.2	7.5	4.8-11.2	5.6	5.2	3.5-7.7
	1986-1990	29.2	16.0	13.4-18.8	14.6	20.5	15.9-25.9	14.6	13.7	10.7-17.4	10.0	5.4	4.0-7.2	5.2	7.7	4.9-11.4	4.8	4.3	2.7-6.6
	1991-1995	29.6	16.6	14.0-19.6	16.8	23.1	18.3-28.7	12.8	12.4	9.4-16.0	9.0	5.0	3.6-6.8	4.2	6.0	3.7-9.2	4.8	4.2	2.7-6.5
	1996-2000	26.0	14.2	11.8-16.9	11.0	14.6	11.0-19.0	15.0	13.5	10.6-17.2	8.6	4.4	3.2-6.0	3.2	4.2	2.4-6.8	5.4	4.4	2.9-6.7
	2001-2005	23.2	13.1	10.8-15.7	12.2	15.8	12.1-20.4	11.0	11.0	8.2-14.4	6.4	3.4	2.3-4.9	2.2	2.8	1.4-5.0	4.2	3.8	2.3-6.0
	2006-2010	27.0	14.6	12.2-17.4	13.4	16.5	12.8-21.1	13.6	13.0	10.0-16.7	6.2	3.3	2.2-4.7	3.6	4.5	2.7-7.2	2.6	2.2	1.1-4.0
	2011-2015	19.4	9.9	8.0-12.2	12.8	14.5	11.1-18.7	6.6	6.2	4.2-8.8	5.2	2.5	1.6-3.8	2.8	3.1	1.7-5.3	2.4	2.1	1.0-3.8
2016-2020	21.0	11.3	9.2-13.8	13.0	15.3	11.7-19.7	8.0	7.7	5.4-10.7	4.8	2.2	1.4-3.3	2.6	2.6	1.4-4.7	2.2	1.8	0.9-3.4	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schenectady**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	4.8	3.0	1.9-4.5	3.2	4.9	2.7-8.1	1.6	1.6	0.7-3.3	4.6	2.7	1.7-4.2	3.0	3.9	2.2-6.6	1.6	1.9	0.8-3.8
	1981-1985	6.4	3.9	2.6-5.6	3.6	4.9	2.9-7.9	2.8	3.0	1.6-5.1	4.0	2.3	1.4-3.6	2.4	3.2	1.7-5.8	1.6	1.4	0.6-3.0
	1986-1990	5.8	3.7	2.4-5.3	4.0	6.0	3.6-9.4	1.8	2.3	1.0-4.5	4.6	2.9	1.8-4.3	3.4	5.4	3.1-8.8	1.2	1.2	0.4-2.8
	1991-1995	4.2	2.3	1.4-3.6	1.6	2.1	0.9-4.3	2.6	2.4	1.2-4.3	4.8	2.7	1.7-4.1	2.2	2.8	1.4-5.1	2.6	2.5	1.3-4.4
	1996-2000	7.8	4.4	3.1-6.1	5.6	7.4	4.9-10.7	2.2	2.1	1.0-3.9	5.2	2.8	1.8-4.1	3.8	5.0	3.0-7.8	1.4	1.2	0.5-2.7
	2001-2005	8.0	4.4	3.1-6.0	6.0	7.5	5.1-10.8	2.0	2.0	0.9-3.8	5.8	3.1	2.0-4.5	4.8	6.0	3.8-9.0	1.0	0.9	0.3-2.3
	2006-2010	12.0	6.5	4.9-8.4	7.6	9.2	6.5-12.7	4.4	4.3	2.6-6.6	8.2	4.5	3.2-6.1	5.2	6.5	4.2-9.5	3.0	3.0	1.6-5.1
	2011-2015	13.8	6.6	5.1-8.5	11.0	11.5	8.6-15.1	2.8	2.6	1.4-4.6	10.0	5.1	3.7-6.8	7.8	9.2	6.5-12.7	2.2	2.0	1.0-3.8
2016-2020	18.6	8.9	7.1-11.0	12.4	13.0	9.9-16.9	6.2	5.4	3.6-7.8	12.4	6.1	4.6-7.8	7.4	8.4	5.8-11.7	5.0	4.2	2.7-6.4	
Pancreas	1976-1980	19.6	12.0	9.7-14.7	11.0	17.1	12.7-22.5	8.6	8.8	6.3-12.0	17.6	10.9	8.7-13.5	9.6	15.3	11.1-20.5	8.0	8.2	5.8-11.2
	1981-1985	20.2	12.1	9.8-14.8	9.6	14.2	10.4-19.1	10.6	10.4	7.8-13.8	19.8	11.7	9.5-14.3	9.4	13.9	10.1-18.6	10.4	10.3	7.6-13.7
	1986-1990	20.8	12.0	9.7-14.6	10.6	15.9	11.8-21.0	10.2	10.0	7.4-13.3	20.0	11.6	9.4-14.2	10.2	15.2	11.2-20.1	9.8	9.3	6.8-12.5
	1991-1995	24.2	13.1	10.8-15.7	11.8	16.5	12.5-21.4	12.4	11.1	8.5-14.4	22.8	12.3	10.2-14.9	11.2	16.3	12.3-21.3	11.6	10.3	7.8-13.5
	1996-2000	19.2	10.5	8.5-12.9	7.6	10.1	7.1-13.8	11.6	10.5	7.9-13.7	22.0	12.0	9.8-14.5	9.8	13.2	9.8-17.5	12.2	10.9	8.3-14.2
	2001-2005	27.6	14.7	12.3-17.4	12.6	16.3	12.5-20.9	15.0	13.4	10.4-17.0	24.6	12.8	10.6-15.3	10.8	14.2	10.6-18.5	13.8	11.6	9.0-15.0
	2006-2010	25.2	13.3	11.1-16.0	13.8	17.1	13.3-21.7	11.4	10.8	8.2-14.2	20.8	10.6	8.6-12.9	11.4	14.0	10.5-18.2	9.4	8.4	6.1-11.4
	2011-2015	30.2	15.6	13.2-18.4	15.0	17.8	13.9-22.5	15.2	13.7	10.7-17.4	23.2	12.1	10.0-14.6	12.2	14.8	11.2-19.1	11.0	9.8	7.3-13.0
2016-2020	33.8	16.9	14.3-19.7	16.4	18.6	14.7-23.3	17.4	15.7	12.4-19.6	27.0	13.1	10.9-15.6	12.0	13.2	10.0-17.2	15.0	12.9	10.1-16.4	
Larynx	1976-1980	11.0	6.4	4.8-8.4	8.2	11.5	8.1-15.8	2.8	2.8	1.5-4.8	3.4	2.0	1.1-3.2	2.0	3.4	1.5-6.3	1.4	1.4	0.5-2.9
	1981-1985	11.6	6.9	5.2-9.0	10.0	13.9	10.2-18.5	1.6	1.7	0.7-3.6	2.8	1.6	0.9-2.7	2.2	2.8	1.4-5.1	0.6	0.8	0.1-2.3
	1986-1990	11.8	6.9	5.2-9.0	8.8	12.6	9.1-17.1	3.0	3.0	1.6-5.1	4.2	2.4	1.5-3.7	3.4	5.2	3.0-8.4	0.8	0.7	0.2-1.9
	1991-1995	12.2	7.1	5.4-9.2	9.4	13.1	9.6-17.6	2.8	3.0	1.6-5.2	2.6	1.3	0.7-2.3	1.8	2.5	1.1-4.7	0.8	0.7	0.2-2.0
	1996-2000	5.8	3.5	2.4-5.1	3.6	4.9	2.9-7.8	2.2	2.5	1.2-4.5	2.4	1.4	0.7-2.5	1.8	2.5	1.2-4.8	0.6	0.5	0.1-1.8
	2001-2005	6.0	3.4	2.3-4.9	4.6	6.1	3.9-9.2	1.4	1.5	0.6-3.2	3.0	1.5	0.9-2.6	2.0	2.7	1.3-4.9	1.0	0.9	0.3-2.3
	2006-2010	6.6	3.8	2.6-5.4	5.2	6.4	4.2-9.5	1.4	1.6	0.6-3.4	1.8	1.0	0.5-2.0	1.6	2.1	0.9-4.2	0.2	0.2	0.0-1.3
	2011-2015	6.0	3.0	2.0-4.3	5.2	5.7	3.7-8.5	0.8	0.8	0.2-2.3	1.2	0.6	0.2-1.5	1.2	1.5	0.6-3.4	0.0	0.0	0.0-0.9
2016-2020	8.0	3.8	2.7-5.2	6.2	6.5	4.3-9.3	1.8	1.5	0.7-3.1	1.2	0.5	0.2-1.2	1.0	0.9	0.3-2.3	0.2	0.1	0.0-1.1	
Lung and bronchus	1976-1980	105.8	62.7	57.4-68.5	74.0	108.0	96.8-120.2	31.8	32.9	27.9-38.6	85.8	50.9	46.1-56.1	61.2	90.0	79.8-101.2	24.6	25.2	20.9-30.2
	1981-1985	113.0	65.5	60.1-71.4	72.0	102.8	92.1-114.5	41.0	41.1	35.5-47.4	95.0	55.1	50.1-60.4	62.4	89.4	79.4-100.3	32.6	33.0	27.9-38.7
	1986-1990	124.6	72.0	66.3-78.1	76.0	108.3	97.4-120.1	48.6	49.5	43.2-56.4	100.2	56.5	51.5-61.8	61.0	85.7	76.1-96.2	39.2	38.3	32.9-44.3
	1991-1995	132.8	74.1	68.5-80.0	77.4	104.5	94.2-115.6	55.4	54.1	47.7-61.1	104.6	58.3	53.3-63.6	63.0	86.9	77.4-97.2	41.6	39.5	34.1-45.4
	1996-2000	128.8	71.8	66.3-77.6	68.2	90.7	81.3-100.9	60.6	58.5	51.9-65.8	108.8	59.4	54.4-64.7	58.4	77.8	69.1-87.3	50.4	47.1	41.3-53.6
	2001-2005	131.2	73.1	67.6-79.0	67.2	88.0	78.9-98.0	64.0	63.2	56.3-70.8	96.2	52.9	48.2-57.9	49.2	64.6	56.7-73.2	47.0	45.0	39.3-51.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schenectady**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	139.8	74.8	69.3-80.7	69.0	86.6	77.6-96.4	70.8	66.8	59.8-74.4	99.8	52.6	48.0-57.5	54.0	67.6	59.7-76.4	45.8	42.1	36.6-48.2
	2011-2015	136.2	69.1	63.9-74.6	65.8	77.5	69.1-86.6	70.4	63.6	56.9-70.8	84.4	42.8	38.7-47.3	43.2	51.1	44.3-58.6	41.2	36.7	31.7-42.3
	2016-2020	140.8	67.5	62.5-72.8	69.6	76.3	68.3-85.0	71.2	61.6	55.2-68.6	83.6	40.2	36.3-44.3	40.6	45.9	39.6-52.9	43.0	36.3	31.5-41.8
Melanoma of the skin	1976-1980	11.6	7.8	5.8-10.1	5.8	8.2	5.4-11.9	5.8	7.6	5.0-11.1	4.4	2.8	1.7-4.3	2.0	3.0	1.4-5.6	2.4	2.7	1.3-4.9
	1981-1985	19.4	12.4	10.0-15.3	11.4	15.8	11.8-20.7	8.0	9.7	6.8-13.4	4.6	3.0	1.8-4.5	2.4	3.3	1.7-6.0	2.2	2.8	1.4-5.2
	1986-1990	14.4	9.1	7.0-11.5	8.2	11.2	7.9-15.3	6.2	7.2	4.8-10.4	5.2	3.5	2.2-5.1	3.6	5.6	3.3-8.9	1.6	2.0	0.8-4.1
	1991-1995	14.8	9.0	7.0-11.3	8.2	11.7	8.3-15.9	6.6	7.3	4.9-10.4	4.2	2.4	1.5-3.8	2.0	3.0	1.4-5.5	2.2	2.1	1.0-3.9
	1996-2000	12.4	7.1	5.4-9.2	7.4	9.9	6.9-13.6	5.0	4.8	3.1-7.3	4.6	2.6	1.6-3.9	2.8	3.8	2.1-6.4	1.8	1.7	0.8-3.4
	2001-2005	22.0	12.9	10.6-15.6	13.4	17.7	13.7-22.6	8.6	9.8	7.0-13.3	6.4	3.5	2.4-5.0	3.2	4.2	2.4-6.9	3.2	3.3	1.8-5.4
	2006-2010	31.0	18.0	15.2-21.1	17.8	22.3	17.8-27.5	13.2	15.3	11.8-19.6	4.6	2.4	1.5-3.7	3.4	4.2	2.4-6.8	1.2	1.0	0.4-2.5
	2011-2015	33.6	17.5	14.9-20.5	18.8	22.2	17.8-27.3	14.8	14.4	11.1-18.3	5.0	2.6	1.7-4.0	3.6	4.6	2.7-7.3	1.4	1.2	0.4-2.6
2016-2020	32.8	16.6	14.0-19.4	18.2	19.6	15.7-24.3	14.6	14.3	11.1-18.2	4.2	2.0	1.2-3.1	2.2	2.5	1.2-4.6	2.0	1.5	0.7-3.0	
Female breast	1976-1980							87.2	99.7	90.1-109.9							31.0	34.6	29.2-40.8
	1981-1985							103.0	111.9	101.9-122.5							33.2	34.0	28.9-39.8
	1986-1990							123.2	130.3	119.8-141.6							40.6	42.2	36.4-48.8
	1991-1995							108.6	116.0	106.1-126.6							32.8	32.1	27.2-37.7
	1996-2000							134.8	144.5	133.5-156.3							28.8	27.4	22.9-32.5
	2001-2005							131.2	139.3	128.6-150.6							27.0	25.3	21.0-30.2
	2006-2010							129.0	131.0	120.8-141.8							23.2	21.9	17.9-26.4
	2011-2015							138.0	141.4	130.7-152.8							21.6	18.6	15.1-22.7
2016-2020							146.2	143.6	132.9-155.0							24.0	21.2	17.4-25.7	
Cervix uteri	1976-1980							8.8	10.7	7.6-14.5							4.6	5.3	3.3-8.1
	1981-1985							9.6	10.8	7.8-14.5							3.2	3.8	2.1-6.3
	1986-1990							11.6	13.7	10.3-17.8							2.8	2.8	1.5-4.9
	1991-1995							8.4	9.6	6.9-13.1							2.4	2.3	1.2-4.2
	1996-2000							5.4	6.8	4.4-9.9							1.2	1.3	0.4-2.9
	2001-2005							5.2	6.1	3.9-9.1							1.4	1.7	0.7-3.5
	2006-2010							4.2	4.7	2.9-7.3							2.6	2.7	1.4-4.7
	2011-2015							7.2	8.7	6.0-12.2							1.2	1.5	0.5-3.3
	2016-2020							5.4	6.6	4.3-9.8							1.4	1.3	0.5-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schenectady**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							27.6	30.3	25.3-36.1							4.6	4.6	2.9-7.1
	1981-1985							27.0	28.7	23.9-34.2							5.4	5.3	3.5-8.0
	1986-1990							22.4	24.0	19.6-29.2							5.4	5.0	3.3-7.4
	1991-1995							22.2	22.8	18.6-27.7							4.0	3.7	2.2-5.8
	1996-2000							23.4	25.1	20.7-30.3							3.4	2.9	1.7-4.8
	2001-2005							24.2	26.1	21.6-31.3							4.8	4.6	2.9-7.0
	2006-2010							30.0	30.3	25.6-35.7							3.8	3.3	1.9-5.3
	2011-2015							32.0	30.9	26.2-36.2							5.8	5.3	3.5-7.8
	2016-2020							33.8	30.8	26.2-36.0							5.8	4.6	3.0-6.7
Ovary	1976-1980							14.0	15.4	11.9-19.7							11.2	11.7	8.8-15.4
	1981-1985							18.8	21.0	16.8-26.0							11.2	11.6	8.7-15.3
	1986-1990							18.4	21.0	16.7-26.0							12.4	12.5	9.5-16.2
	1991-1995							18.2	19.0	15.2-23.5							10.6	10.7	7.9-14.2
	1996-2000							15.0	16.4	12.9-20.7							10.0	9.4	7.0-12.6
	2001-2005							14.2	15.4	12.0-19.6							10.0	9.7	7.1-12.9
	2006-2010							9.0	9.1	6.6-12.3							5.4	5.0	3.3-7.5
	2011-2015							13.2	12.9	9.8-16.6							8.8	7.8	5.6-10.7
	2016-2020							9.8	9.1	6.7-12.3							7.2	6.1	4.2-8.6
Prostate	1976-1980				53.6	90.5	79.5-102.6							18.0	33.2	26.4-41.0			
	1981-1985				53.6	79.6	70.0-90.1							17.4	28.5	22.6-35.4			
	1986-1990				68.2	100.8	90.1-112.5							23.6	37.8	31.1-45.6			
	1991-1995				125.2	167.0	153.9-180.8							20.8	30.9	25.2-37.6			
	1996-2000				132.0	173.9	160.8-187.7							20.6	28.6	23.3-34.7			
	2001-2005				113.6	148.5	136.5-161.2							21.4	27.6	22.6-33.4			
	2006-2010				111.0	137.6	126.2-149.7							17.8	22.3	17.9-27.5			
	2011-2015				86.6	93.3	84.5-102.8							15.0	17.9	14.0-22.6			
	2016-2020				120.0	123.0	113.1-133.6							16.8	18.7	14.9-23.3			
Testis	1976-1980				4.4	5.4	3.4-8.3							0.4	0.8	0.1-2.7			
	1981-1985				4.6	5.9	3.7-9.0							0.2	0.3	0.0-1.6			
	1986-1990				4.2	5.5	3.4-8.5							0.0	0.0	0.0-1.1			
	1991-1995				4.8	6.6	4.2-9.8							0.2	0.3	0.0-1.6			
	1996-2000				3.2	4.7	2.7-7.6							0.2	0.3	0.0-1.5			
	2001-2005				3.8	5.6	3.3-8.7							0.0	0.0	0.0-1.0			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schenectady**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				5.0	7.2	4.6-10.6							0.0	0.0	0.0-1.0			
	2011-2015				5.0	7.3	4.7-10.8							0.0	0.0	0.0-1.0			
	2016-2020				4.2	6.3	3.9-9.6							0.0	0.0	0.0-1.0			
Urinary bladder (incl. in situ)	1976-1980	41.6	26.0	22.5-29.8	29.4	46.6	39.0-55.2	12.2	13.2	10.0-17.1	14.0	9.0	7.0-11.3	8.8	14.8	10.6-20.1	5.2	5.5	3.6-8.2
	1981-1985	46.8	27.2	23.7-31.0	32.6	48.4	41.0-56.8	14.2	14.0	10.9-17.8	13.2	7.9	6.1-10.1	9.4	15.4	11.2-20.7	3.8	3.6	2.2-5.8
	1986-1990	41.0	23.4	20.2-26.9	28.4	40.7	34.1-48.2	12.6	12.2	9.2-15.8	10.2	5.6	4.2-7.4	4.8	6.8	4.3-10.2	5.4	4.6	3.0-6.8
	1991-1995	47.0	26.2	22.9-29.9	33.6	45.4	38.7-52.9	13.4	12.8	9.8-16.5	9.0	4.7	3.5-6.4	5.2	7.3	4.7-10.8	3.8	3.2	1.9-5.1
	1996-2000	44.4	24.7	21.5-28.3	32.6	43.2	36.8-50.4	11.8	11.4	8.6-14.9	12.8	6.5	5.0-8.4	8.4	11.3	8.1-15.4	4.4	3.8	2.3-5.9
	2001-2005	59.8	32.8	29.2-36.9	42.8	56.0	48.7-64.0	17.0	15.6	12.4-19.5	13.8	7.0	5.4-8.9	9.4	12.3	9.0-16.4	4.4	3.4	2.1-5.3
	2006-2010	52.0	26.7	23.5-30.3	36.8	45.3	38.9-52.4	15.2	13.8	10.7-17.5	10.8	4.9	3.7-6.5	6.6	8.0	5.5-11.2	4.2	3.4	2.0-5.4
	2011-2015	55.8	29.1	25.7-32.9	41.8	51.0	44.2-58.7	14.0	12.9	9.9-16.5	12.6	5.7	4.4-7.4	8.6	9.8	7.0-13.3	4.0	3.2	1.8-5.2
2016-2020	48.6	23.1	20.2-26.3	35.4	39.3	33.6-45.8	13.2	10.9	8.4-14.2	10.0	4.5	3.3-6.0	7.4	8.2	5.7-11.4	2.6	2.0	1.0-3.6	
Kidney and renal pelvis	1976-1980	12.4	7.8	5.9-10.0	7.2	11.4	7.8-16.0	5.2	5.3	3.4-7.8	4.2	2.7	1.7-4.2	2.2	3.5	1.7-6.3	2.0	2.1	1.0-3.9
	1981-1985	11.8	7.1	5.4-9.3	6.2	9.2	6.1-13.2	5.6	5.6	3.6-8.2	5.8	3.4	2.2-4.9	2.6	3.5	1.8-6.1	3.2	3.2	1.8-5.4
	1986-1990	16.6	9.8	7.8-12.3	10.4	14.7	10.9-19.4	6.2	6.4	4.2-9.2	5.8	3.6	2.4-5.2	3.4	5.2	3.0-8.4	2.4	2.3	1.2-4.2
	1991-1995	18.4	10.5	8.4-12.9	9.8	13.1	9.6-17.4	8.6	8.6	6.2-11.8	7.0	3.9	2.7-5.4	4.0	5.5	3.3-8.5	3.0	2.8	1.5-4.7
	1996-2000	20.8	12.1	9.9-14.7	14.4	19.2	15.0-24.2	6.4	7.0	4.7-10.0	9.8	5.5	4.1-7.4	5.2	6.8	4.5-10.1	4.6	4.6	2.9-7.1
	2001-2005	30.0	17.2	14.6-20.3	20.2	26.4	21.5-32.1	9.8	9.8	7.2-13.1	6.0	3.3	2.2-4.8	3.4	4.4	2.5-7.0	2.6	2.4	1.2-4.2
	2006-2010	30.6	16.8	14.2-19.8	21.4	26.8	21.9-32.5	9.2	8.8	6.4-11.9	7.6	3.9	2.7-5.4	5.2	6.5	4.3-9.6	2.4	2.0	1.0-3.8
	2011-2015	29.6	15.6	13.1-18.4	20.4	23.1	18.7-28.2	9.2	9.3	6.7-12.7	7.8	4.0	2.8-5.5	5.4	6.3	4.1-9.3	2.4	2.3	1.1-4.2
2016-2020	36.2	18.3	15.7-21.3	21.8	23.8	19.4-28.8	14.4	14.2	10.9-18.1	5.2	2.4	1.6-3.6	3.4	3.5	2.0-5.8	1.8	1.5	0.6-3.2	
Brain and other central nervous system	1976-1980	9.2	5.8	4.2-7.8	6.0	8.3	5.5-12.0	3.2	3.6	2.0-5.9	5.6	3.5	2.3-5.1	3.4	4.4	2.5-7.3	2.2	2.5	1.2-4.5
	1981-1985	9.2	6.0	4.3-8.1	6.0	8.4	5.6-12.1	3.2	4.1	2.3-6.8	5.2	3.1	2.0-4.5	3.4	4.8	2.7-7.7	1.8	1.9	0.8-3.7
	1986-1990	10.4	6.8	5.0-9.0	5.6	7.9	5.2-11.5	4.8	6.1	3.8-9.2	5.2	3.3	2.1-4.9	3.0	4.0	2.2-6.7	2.2	2.8	1.3-5.0
	1991-1995	12.6	8.0	6.1-10.2	6.0	8.4	5.6-12.0	6.6	7.6	5.1-10.8	7.8	4.7	3.3-6.5	4.2	5.6	3.4-8.6	3.6	4.0	2.3-6.4
	1996-2000	12.2	7.4	5.6-9.5	6.6	9.1	6.3-12.8	5.6	6.0	3.9-8.8	8.8	5.3	3.8-7.1	4.6	6.4	4.0-9.6	4.2	4.5	2.8-7.1
	2001-2005	13.4	8.3	6.4-10.5	7.0	9.6	6.6-13.3	6.4	7.2	4.9-10.3	6.8	4.0	2.8-5.7	2.6	3.5	1.9-6.0	4.2	4.4	2.7-6.9
	2006-2010	12.4	7.1	5.4-9.1	7.0	8.7	6.0-12.1	5.4	5.7	3.7-8.5	6.6	3.6	2.5-5.2	4.2	5.3	3.2-8.2	2.4	2.2	1.1-4.1
	2011-2015	14.8	8.6	6.7-10.9	7.8	9.6	6.8-13.2	7.0	7.6	5.2-10.7	9.2	5.1	3.7-6.8	3.2	3.7	2.1-6.1	6.0	6.2	4.1-9.0
2016-2020	11.4	6.6	5.0-8.7	7.0	8.6	5.9-12.0	4.4	5.0	3.0-7.7	8.2	4.2	3.0-5.8	5.0	5.6	3.6-8.4	3.2	3.0	1.6-5.0	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schenectady**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	5.4	3.8	2.5-5.7	1.2	1.9	0.7-4.2	4.2	5.6	3.4-8.6	0.8	0.4	0.1-1.1	0.4	0.4	0.1-1.8	0.4	0.4	0.0-1.5
	1981-1985	4.4	2.8	1.7-4.3	1.6	2.1	0.9-4.3	2.8	3.2	1.7-5.5	1.0	0.5	0.2-1.3	0.4	0.5	0.1-1.9	0.6	0.5	0.1-1.8
	1986-1990	6.4	4.1	2.8-5.9	1.2	1.4	0.5-3.1	5.2	6.6	4.2-9.8	0.2	0.1	0.0-0.7	0.2	0.2	0.0-1.4	0.0	0.0	0.0-1.0
	1991-1995	13.8	8.9	6.9-11.3	3.0	4.4	2.4-7.3	10.8	13.4	10.0-17.5	0.6	0.3	0.1-1.0	0.0	0.0	0.0-1.0	0.6	0.5	0.1-1.6
	1996-2000	9.8	6.5	4.8-8.6	2.2	3.1	1.5-5.5	7.6	9.5	6.7-13.1	0.6	0.3	0.1-1.1	0.4	0.6	0.1-2.1	0.2	0.2	0.0-1.3
	2001-2005	15.4	10.2	8.0-12.8	2.6	3.6	1.9-6.2	12.8	16.3	12.5-20.9	0.4	0.2	0.0-0.9	0.0	0.0	0.0-1.0	0.4	0.4	0.0-1.6
	2006-2010	19.6	12.5	10.1-15.3	4.4	5.7	3.5-8.7	15.2	18.8	14.7-23.6	0.0	0.0	0.0-0.5	0.0	0.0	0.0-1.0	0.0	0.0	0.0-0.9
	2011-2015	26.8	16.2	13.5-19.4	5.4	6.4	4.2-9.5	21.4	25.8	21.0-31.3	0.6	0.3	0.1-1.0	0.4	0.5	0.1-1.8	0.2	0.2	0.0-1.2
2016-2020	27.2	16.5	13.7-19.6	6.4	7.4	5.0-10.7	20.8	25.2	20.4-30.8	0.2	0.1	0.0-0.7	0.0	0.0	0.0-1.0	0.2	0.2	0.0-1.4	
Hodgkin lymphoma	1976-1980	5.8	3.6	2.3-5.2	3.4	4.7	2.7-7.8	2.4	2.5	1.3-4.5	1.4	0.9	0.3-1.9	0.8	1.4	0.3-3.7	0.6	0.6	0.1-1.8
	1981-1985	7.8	4.7	3.3-6.5	4.4	5.7	3.5-8.8	3.4	3.6	2.1-5.9	1.2	0.6	0.2-1.4	0.6	0.7	0.1-2.1	0.6	0.6	0.1-1.9
	1986-1990	7.0	4.5	3.1-6.3	3.8	5.3	3.2-8.4	3.2	3.9	2.2-6.3	1.6	0.9	0.4-1.9	1.0	1.3	0.4-3.2	0.6	0.7	0.1-2.1
	1991-1995	6.2	3.8	2.6-5.5	3.4	4.6	2.6-7.4	2.8	3.3	1.8-5.7	1.6	1.0	0.4-2.0	1.0	1.3	0.4-3.1	0.6	0.6	0.1-2.0
	1996-2000	3.2	2.1	1.2-3.5	1.8	2.6	1.2-5.0	1.4	1.8	0.7-3.7	1.6	1.0	0.4-2.0	0.6	0.9	0.2-2.5	1.0	1.2	0.4-2.9
	2001-2005	4.4	3.0	1.8-4.5	3.2	4.5	2.6-7.4	1.2	1.6	0.6-3.4	0.6	0.3	0.1-1.0	0.6	0.7	0.1-2.2	0.0	0.0	0.0-1.0
	2006-2010	6.0	4.0	2.7-5.8	3.4	4.7	2.7-7.5	2.6	3.3	1.7-5.7	0.4	0.2	0.0-0.9	0.2	0.3	0.0-1.6	0.2	0.1	0.0-1.1
	2011-2015	3.8	2.4	1.5-3.9	1.8	2.3	1.0-4.5	2.0	2.6	1.2-4.8	0.6	0.3	0.1-0.9	0.4	0.4	0.0-1.7	0.2	0.1	0.0-1.1
2016-2020	6.0	3.7	2.5-5.4	3.8	4.9	2.9-7.8	2.2	2.6	1.3-4.8	1.0	0.4	0.1-1.1	0.8	0.8	0.2-2.3	0.2	0.2	0.0-1.2	
Non-Hodgkin lymphomas	1976-1980	22.0	13.5	11.0-16.4	11.8	16.8	12.6-21.9	10.2	11.2	8.2-14.9	11.8	7.3	5.5-9.5	5.8	8.4	5.5-12.3	6.0	6.2	4.2-9.0
	1981-1985	23.0	14.1	11.5-17.0	10.4	15.7	11.6-20.8	12.6	13.6	10.3-17.6	12.2	7.1	5.4-9.3	5.6	8.5	5.5-12.6	6.6	6.5	4.4-9.3
	1986-1990	25.4	15.2	12.6-18.2	13.2	19.9	15.2-25.4	12.2	12.0	9.1-15.6	11.4	6.6	5.0-8.6	5.4	7.7	5.0-11.3	6.0	5.8	3.9-8.4
	1991-1995	36.8	21.4	18.3-24.7	21.0	28.4	23.2-34.5	15.8	15.8	12.4-19.9	17.0	9.4	7.5-11.7	9.4	13.5	9.8-18.0	7.6	6.8	4.7-9.5
	1996-2000	40.8	23.4	20.3-26.9	21.8	29.5	24.2-35.7	19.0	18.1	14.5-22.3	20.6	11.4	9.3-13.9	9.6	13.1	9.6-17.4	11.0	10.1	7.5-13.3
	2001-2005	36.4	20.6	17.7-23.9	17.0	22.4	17.9-27.7	19.4	19.7	15.9-24.3	12.6	6.5	5.0-8.4	7.6	10.0	7.0-13.7	5.0	4.1	2.6-6.3
	2006-2010	42.4	22.9	19.8-26.3	23.8	29.4	24.3-35.3	18.6	17.9	14.3-22.2	14.8	7.7	6.0-9.7	9.2	12.0	8.7-16.1	5.6	4.8	3.1-7.1
	2011-2015	40.6	21.0	18.1-24.2	24.4	28.4	23.5-34.2	16.2	14.9	11.7-18.8	12.0	5.9	4.4-7.7	5.6	6.9	4.5-10.0	6.4	5.0	3.4-7.4
2016-2020	47.8	23.8	20.8-27.2	24.6	26.9	22.2-32.3	23.2	20.9	17.2-25.4	10.2	4.6	3.4-6.1	5.8	6.5	4.3-9.4	4.4	3.1	1.9-4.9	
Myeloma	1976-1980	9.2	5.3	3.9-7.2	3.8	5.8	3.3-9.4	5.4	5.3	3.5-7.8	4.8	3.0	1.9-4.6	3.4	6.0	3.3-9.9	1.4	1.4	0.6-3.0
	1981-1985	6.4	3.7	2.5-5.3	2.6	4.1	2.1-7.2	3.8	3.6	2.1-5.7	6.6	3.8	2.6-5.4	2.2	3.4	1.6-6.2	4.4	4.2	2.6-6.5
	1986-1990	5.6	3.1	2.0-4.5	3.2	4.7	2.6-7.7	2.4	2.1	1.1-3.9	4.8	2.6	1.7-3.9	1.8	2.7	1.2-5.2	3.0	2.6	1.5-4.5
	1991-1995	6.6	3.6	2.5-5.1	3.2	4.2	2.4-6.8	3.4	3.4	1.9-5.5	5.8	3.0	2.0-4.3	3.4	4.5	2.6-7.2	2.4	2.0	1.0-3.6
	1996-2000	7.2	3.9	2.7-5.4	4.0	5.3	3.2-8.2	3.2	2.9	1.6-4.9	4.2	2.2	1.4-3.4	2.0	2.6	1.3-4.9	2.2	2.1	1.0-3.9
	2001-2005	9.6	5.0	3.7-6.7	5.4	6.9	4.6-10.1	4.2	3.8	2.3-6.0	4.4	2.3	1.4-3.5	2.4	3.1	1.6-5.5	2.0	1.7	0.8-3.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schenectady**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	11.2	5.8	4.4-7.6	7.4	9.0	6.3-12.4	3.8	3.6	2.2-5.8	6.6	3.2	2.2-4.6	2.8	3.5	1.9-5.9	3.8	3.2	1.9-5.2
	2011-2015	15.6	8.1	6.3-10.2	8.2	8.8	6.3-12.0	7.4	7.3	5.1-10.3	5.2	2.8	1.8-4.1	3.0	3.9	2.1-6.4	2.2	1.9	0.9-3.6
	2016-2020	15.0	7.0	5.5-8.9	8.8	9.6	7.0-13.1	6.2	5.1	3.4-7.4	7.2	3.2	2.2-4.5	3.6	3.9	2.3-6.4	3.6	2.7	1.6-4.6
Leukemias	1976-1980	25.8	16.6	13.8-19.8	14.6	23.2	17.9-29.5	11.2	12.4	9.3-16.2	13.0	8.3	6.3-10.6	6.2	9.2	6.2-13.3	6.8	7.6	5.2-10.7
	1981-1985	24.0	14.6	12.0-17.5	15.6	22.8	17.9-28.7	8.4	8.8	6.2-12.0	13.8	8.5	6.6-10.8	8.0	12.8	9.0-17.6	5.8	6.2	4.1-9.1
	1986-1990	21.4	12.5	10.2-15.2	12.2	18.0	13.6-23.3	9.2	9.3	6.7-12.7	15.0	8.7	6.8-10.9	8.8	13.4	9.6-18.1	6.2	6.0	4.0-8.7
	1991-1995	23.6	13.4	11.1-16.1	14.2	18.8	14.6-23.8	9.4	9.1	6.6-12.3	15.4	8.6	6.8-10.8	8.6	11.5	8.3-15.6	6.8	6.3	4.3-9.0
	1996-2000	22.2	12.3	10.1-14.9	10.4	13.9	10.3-18.2	11.8	10.6	8.0-13.9	14.8	7.8	6.1-9.9	6.4	8.4	5.8-12.0	8.4	7.1	5.1-9.8
	2001-2005	25.2	14.1	11.7-16.9	15.2	19.7	15.5-24.7	10.0	10.2	7.5-13.7	11.8	6.1	4.6-7.9	6.2	7.9	5.4-11.3	5.6	5.2	3.4-7.8
	2006-2010	30.8	17.6	14.9-20.7	18.4	23.9	19.2-29.4	12.4	13.6	10.3-17.6	12.2	6.4	4.9-8.3	8.0	10.1	7.2-13.9	4.2	4.1	2.5-6.5
	2011-2015	36.4	19.1	16.4-22.2	18.8	23.3	18.8-28.7	17.6	15.7	12.5-19.7	15.4	7.9	6.2-10.0	8.6	11.0	7.9-14.9	6.8	5.7	3.8-8.2
2016-2020	29.6	15.4	12.9-18.2	20.2	23.2	18.8-28.4	9.4	9.7	7.0-13.1	11.4	5.5	4.1-7.2	7.8	8.8	6.2-12.2	3.6	3.3	1.9-5.4	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schoharie**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	116.0	413.3	379.6-449.1	61.6	478.0	424.6-536.3	54.4	360.5	317.8-407.5	59.8	212.8	189.0-238.8	31.0	249.8	210.8-293.8	28.8	188.3	158.2-222.6
	1981-1985	120.6	400.7	368.7-434.7	64.2	489.2	435.1-548.0	56.4	352.0	310.9-397.1	65.6	214.2	191.3-239.2	35.6	271.2	231.4-315.7	30.0	174.6	147.3-205.8
	1986-1990	139.4	433.6	401.5-467.5	69.8	511.8	457.2-570.9	69.6	405.1	362.6-451.3	71.6	223.3	200.4-248.0	35.6	275.2	234.5-320.7	36.0	200.6	171.7-233.1
	1991-1995	151.8	443.6	412.3-476.6	81.8	534.0	482.1-590.0	70.0	383.4	343.6-426.6	73.4	215.4	193.8-238.9	38.0	269.3	230.9-312.0	35.4	192.0	164.3-223.2
	1996-2000	157.0	449.8	418.8-482.5	81.2	506.8	458.0-559.4	75.8	406.3	366.0-450.1	82.4	232.7	210.8-256.5	42.6	283.0	245.5-324.6	39.8	201.4	174.1-232.1
	2001-2005	183.0	491.0	459.5-524.2	89.4	511.6	464.5-562.3	93.6	479.2	436.1-525.7	75.8	199.5	179.8-220.8	39.0	228.6	197.1-263.7	36.8	174.0	149.6-201.7
	2006-2010	187.8	467.3	437.3-499.0	101.8	520.2	474.9-568.9	86.0	418.7	379.1-461.6	75.0	187.2	168.5-207.6	41.6	221.8	192.0-255.0	33.4	156.0	132.8-182.5
	2011-2015	200.4	457.6	428.7-488.1	108.0	503.3	460.2-549.7	92.4	418.3	379.5-460.4	71.0	160.9	144.2-179.1	39.6	194.9	168.0-225.2	31.4	131.9	111.6-155.3
	2016-2020	218.0	456.0	427.7-485.9	114.2	477.5	437.3-520.9	103.8	437.8	398.1-480.7	73.0	148.8	133.4-165.7	40.8	171.9	148.4-198.7	32.2	127.2	107.5-150.1
Oral cavity and pharynx	1976-1980	3.0	10.4	5.7-17.4	2.0	15.7	7.3-29.2	1.0	5.7	1.9-14.0	0.2	0.8	0.0-4.4	0.0	0.0	0.0-6.0	0.2	1.7	0.0-8.7
	1981-1985	3.8	11.5	6.9-18.2	2.8	19.9	10.6-34.0	1.0	5.2	1.7-12.9	1.4	4.6	1.8-9.7	1.2	8.8	3.1-19.6	0.2	1.1	0.0-6.8
	1986-1990	3.0	9.5	5.3-15.7	2.2	19.0	9.1-34.4	0.8	4.1	1.1-11.1	1.0	3.0	1.0-7.1	1.0	6.5	2.0-15.5	0.0	0.0	0.0-4.8
	1991-1995	3.2	9.2	5.2-15.0	1.4	8.4	3.3-17.6	1.8	9.7	4.4-18.8	0.4	1.3	0.2-4.7	0.2	1.4	0.0-7.6	0.2	1.1	0.0-6.4
	1996-2000	4.2	12.0	7.4-18.4	1.8	10.3	4.7-20.0	2.4	12.5	6.4-22.3	0.8	2.2	0.6-5.9	0.2	1.1	0.0-7.0	0.6	3.2	0.7-9.9
	2001-2005	2.8	7.4	4.1-12.7	1.8	9.7	4.4-19.1	1.0	5.2	1.7-12.7	1.2	3.4	1.2-7.5	0.6	3.8	0.8-11.4	0.6	2.5	0.5-8.5
	2006-2010	4.4	11.2	6.9-17.3	3.0	16.2	9.0-27.4	1.4	7.0	2.7-15.2	0.6	1.5	0.3-4.8	0.0	0.0	0.0-4.9	0.6	2.8	0.6-9.1
	2011-2015	4.8	12.1	7.5-18.5	3.8	19.3	11.2-31.3	1.0	5.2	1.5-13.2	0.6	1.8	0.3-5.5	0.4	2.6	0.3-9.8	0.2	0.8	0.0-6.2
	2016-2020	7.4	15.4	10.7-21.9	5.6	24.2	15.8-36.2	1.8	6.9	3.1-14.6	1.2	2.3	0.8-5.8	0.8	3.0	0.8-9.4	0.4	1.5	0.2-7.5
Esophagus	1976-1980	0.8	2.6	0.7-6.8	0.6	4.3	0.9-13.2	0.2	1.1	0.0-7.1	0.6	1.8	0.4-5.5	0.6	3.7	0.8-11.7	0.0	0.0	0.0-5.4
	1981-1985	0.8	2.4	0.7-6.5	0.6	4.2	0.8-12.6	0.2	1.1	0.0-6.8	1.6	5.8	2.5-11.5	1.2	10.9	3.9-23.5	0.4	2.3	0.3-8.8
	1986-1990	0.8	2.1	0.6-5.8	0.8	4.9	1.2-13.2	0.0	0.0	0.0-4.8	0.4	1.2	0.1-4.5	0.4	2.4	0.3-9.4	0.0	0.0	0.0-4.8
	1991-1995	2.2	5.8	2.9-10.6	2.0	12.0	5.5-22.7	0.2	1.0	0.0-6.3	2.6	7.2	3.8-12.4	2.2	15.0	7.1-27.4	0.4	1.8	0.2-7.4
	1996-2000	1.0	2.9	0.9-6.8	0.8	5.0	1.4-12.9	0.2	1.2	0.0-6.8	2.2	6.3	3.2-11.5	1.8	11.2	5.1-21.4	0.4	1.9	0.2-7.8
	2001-2005	1.8	4.8	2.2-9.4	1.4	8.0	3.2-17.0	0.4	1.8	0.2-7.5	1.2	3.1	1.1-7.0	0.8	4.6	1.2-12.1	0.4	1.7	0.2-7.4
	2006-2010	3.8	9.3	5.5-14.9	3.2	16.3	9.1-27.3	0.6	3.0	0.6-9.4	3.0	7.3	4.0-12.3	2.4	11.6	5.9-21.0	0.6	3.0	0.6-9.4
	2011-2015	3.2	7.0	4.0-11.8	2.6	11.6	6.1-20.7	0.6	2.3	0.5-8.3	3.4	8.1	4.7-13.3	2.4	12.1	6.2-21.8	1.0	4.1	1.3-10.9
	2016-2020	2.6	5.1	2.6-9.4	2.0	8.0	3.7-16.2	0.6	2.3	0.4-8.7	1.4	2.4	0.9-5.7	1.2	4.2	1.5-10.7	0.2	0.6	0.0-6.1
Stomach	1976-1980	3.2	11.3	6.4-18.5	1.6	11.0	4.7-22.2	1.6	10.4	4.4-20.7	2.4	8.4	4.3-14.8	1.2	8.6	3.1-19.2	1.2	7.8	2.8-17.4
	1981-1985	2.4	8.3	4.2-14.6	2.4	18.4	9.3-32.3	0.0	0.0	0.0-5.1	2.4	8.0	4.1-14.1	1.8	12.9	5.9-24.9	0.6	3.0	0.6-9.7
	1986-1990	2.6	8.3	4.4-14.3	1.6	11.9	4.8-23.9	1.0	5.2	1.7-12.7	1.6	5.2	2.2-10.3	1.0	8.0	2.4-18.9	0.6	3.0	0.6-9.5
	1991-1995	2.8	8.3	4.5-14.0	1.8	12.7	5.6-24.4	1.0	5.1	1.6-12.3	1.4	4.0	1.6-8.3	1.0	6.1	2.0-14.6	0.4	1.9	0.2-7.6
	1996-2000	3.2	9.0	5.2-14.8	2.6	16.9	8.9-29.0	0.6	3.1	0.6-9.6	1.8	5.1	2.3-9.8	1.2	9.0	3.2-19.5	0.6	3.3	0.7-10.0
	2001-2005	2.2	5.8	2.9-10.6	1.6	9.0	3.9-18.2	0.6	2.7	0.6-8.9	1.6	4.1	1.8-8.3	1.0	5.6	1.8-13.5	0.6	2.7	0.5-8.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schoharie**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	3.4	8.4	4.8-13.8	1.8	9.6	4.3-18.9	1.6	6.8	2.9-14.2	1.2	3.4	1.2-7.6	0.8	4.8	1.3-12.6	0.4	2.0	0.2-7.9
	2011-2015	3.4	8.4	4.8-13.8	1.8	10.1	4.5-19.7	1.6	6.7	2.9-14.4	0.6	1.3	0.3-4.5	0.4	1.7	0.2-7.7	0.2	0.9	0.0-6.4
	2016-2020	2.2	4.7	2.3-9.0	1.4	6.0	2.4-13.5	0.8	3.5	0.9-10.3	1.0	2.0	0.6-5.5	0.8	3.5	0.9-10.2	0.2	0.6	0.0-6.1
Colorectal	1976-1980	19.0	68.9	55.4-84.5	10.0	77.3	56.8-102.7	9.0	60.7	43.8-82.1	8.8	32.7	23.7-44.1	4.4	36.0	22.1-55.2	4.4	29.8	18.5-45.6
	1981-1985	20.0	64.5	52.4-78.8	11.0	83.3	62.0-109.4	9.0	52.1	37.7-70.3	9.2	28.7	20.9-38.5	5.2	37.7	24.1-56.3	4.0	22.5	13.6-35.2
	1986-1990	20.2	61.3	49.8-74.8	9.4	71.8	51.9-96.5	10.8	57.7	43.1-75.8	12.0	37.7	28.7-48.7	5.2	42.6	27.3-63.0	6.8	36.4	25.0-51.4
	1991-1995	22.0	63.7	52.3-76.9	11.4	79.7	59.9-103.9	10.6	53.3	39.7-70.2	8.6	25.6	18.5-34.5	3.8	29.5	17.4-46.3	4.8	25.7	16.4-38.7
	1996-2000	21.2	59.9	49.0-72.5	9.8	64.3	47.2-85.4	11.4	56.9	43.0-74.2	8.8	24.6	17.9-33.1	3.4	24.4	14.0-39.4	5.4	24.9	16.4-36.9
	2001-2005	18.0	47.2	37.9-58.2	9.4	55.1	40.2-73.7	8.6	40.4	29.2-55.0	9.6	25.6	18.8-34.1	4.2	25.4	15.6-39.3	5.4	24.0	15.8-35.6
	2006-2010	19.8	47.9	38.8-58.6	10.0	50.1	36.9-66.7	9.8	45.7	33.7-61.0	5.8	14.5	9.7-21.0	3.2	17.3	9.8-28.6	2.6	12.0	6.4-21.2
	2011-2015	17.0	40.3	31.9-50.4	9.6	46.8	34.1-63.0	7.4	34.5	23.8-48.7	6.2	13.9	9.3-20.1	2.6	11.9	6.1-21.4	3.6	15.2	9.0-25.0
2016-2020	15.4	33.3	26.0-42.4	8.0	36.7	25.7-51.3	7.4	29.4	20.4-42.0	5.4	11.6	7.5-17.6	2.6	11.8	6.0-21.6	2.8	11.3	6.0-20.4	
Colon excluding rectum	1976-1980	12.8	46.5	35.6-59.6	6.2	47.9	32.2-68.7	6.6	43.3	29.5-61.4	7.0	26.0	18.0-36.3	3.2	24.5	13.7-40.3	3.8	25.7	15.3-40.5
	1981-1985	12.8	40.1	30.8-51.5	6.8	49.3	33.7-69.8	6.0	33.5	22.5-48.3	7.2	22.4	15.6-31.2	4.0	28.2	16.8-44.5	3.2	18.3	10.3-30.2
	1986-1990	14.8	45.4	35.5-57.2	6.6	52.8	35.7-74.9	8.2	43.6	31.1-59.7	10.2	32.2	23.9-42.5	4.2	35.8	21.7-55.1	6.0	32.3	21.6-46.7
	1991-1995	15.6	45.3	35.7-56.6	7.4	51.6	35.9-71.7	8.2	41.9	29.9-57.4	7.4	21.9	15.4-30.2	2.6	21.6	11.2-37.0	4.8	25.7	16.4-38.7
	1996-2000	16.6	47.0	37.5-58.4	7.4	48.3	33.8-66.9	9.2	46.1	33.7-62.0	7.0	19.6	13.6-27.4	2.2	15.6	7.6-28.2	4.8	22.5	14.3-34.0
	2001-2005	11.6	30.5	23.1-39.6	5.6	33.4	22.0-48.9	6.0	28.3	19.0-40.9	7.8	20.8	14.8-28.6	3.4	21.0	12.1-33.9	4.4	19.6	12.2-30.3
	2006-2010	15.0	37.0	29.0-46.6	7.0	35.5	24.5-50.0	8.0	38.1	27.1-52.4	4.8	11.9	7.6-18.0	2.4	12.8	6.5-22.9	2.4	11.2	5.8-20.2
	2011-2015	13.6	32.3	24.8-41.5	7.0	34.8	23.9-49.5	6.6	29.9	20.2-43.2	4.6	10.4	6.5-16.1	1.6	7.4	3.0-15.8	3.0	12.9	7.2-22.2
2016-2020	10.4	22.4	16.4-30.2	5.0	23.3	14.6-35.9	5.4	21.1	13.6-32.2	4.8	10.2	6.3-15.9	2.4	10.9	5.3-20.5	2.4	9.3	4.6-17.8	
Rectum & rectosigmoid	1976-1980	6.2	22.4	15.1-32.0	3.8	29.4	17.4-46.4	2.4	17.4	8.7-31.0	1.8	6.7	3.1-12.8	1.2	11.6	4.0-25.0	0.6	4.2	0.8-12.4
	1981-1985	7.2	24.4	17.0-34.0	4.2	34.0	20.6-52.5	3.0	18.5	10.2-31.1	2.0	6.3	3.0-11.8	1.2	9.5	3.2-21.1	0.8	4.2	1.1-11.5
	1986-1990	5.4	15.9	10.5-23.4	2.8	19.0	10.1-32.5	2.6	14.0	7.4-24.4	1.8	5.6	2.5-10.6	1.0	6.8	2.2-16.3	0.8	4.1	1.1-11.1
	1991-1995	6.4	18.5	12.6-26.1	4.0	28.1	16.9-43.8	2.4	11.3	5.8-20.3	1.2	3.7	1.3-8.1	1.2	7.9	2.9-17.2	0.0	0.0	0.0-4.5
	1996-2000	4.6	12.8	8.1-19.4	2.4	16.0	8.1-28.2	2.2	10.8	5.4-19.8	1.8	5.0	2.3-9.7	1.2	8.8	3.1-19.2	0.6	2.5	0.5-8.3
	2001-2005	6.4	16.7	11.4-23.8	3.8	21.6	13.0-34.1	2.6	12.1	6.4-21.4	1.8	4.7	2.2-9.2	0.8	4.5	1.2-11.9	1.0	4.4	1.4-11.3
	2006-2010	4.8	10.9	7.0-16.6	3.0	14.6	8.1-24.8	1.8	7.7	3.5-15.4	1.0	2.6	0.8-6.3	0.8	4.5	1.2-12.0	0.2	0.8	0.0-5.9
	2011-2015	3.4	8.0	4.6-13.3	2.6	11.9	6.3-21.3	0.8	4.6	1.2-12.4	1.6	3.4	1.5-7.3	1.0	4.5	1.4-11.5	0.6	2.3	0.5-8.3
2016-2020	5.0	10.9	7.0-16.5	3.0	13.4	7.4-23.1	2.0	8.3	3.9-16.6	0.6	1.4	0.3-4.8	0.2	0.9	0.0-6.6	0.4	2.0	0.2-8.4	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schoharie**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	0.8	2.9	0.8-7.5	0.6	5.2	1.0-15.1	0.2	1.2	0.0-7.4	1.0	3.8	1.2-8.9	0.2	2.0	0.1-10.2	0.8	5.2	1.4-13.7
	1981-1985	0.2	0.6	0.0-3.6	0.2	1.3	0.0-8.1	0.0	0.0	0.0-5.1	0.6	2.0	0.4-6.0	0.4	3.0	0.4-11.0	0.2	1.0	0.0-6.7
	1986-1990	0.4	1.3	0.2-4.7	0.2	1.6	0.0-8.5	0.2	0.9	0.0-6.3	1.2	3.6	1.3-7.9	0.4	2.6	0.3-9.9	0.8	4.1	1.1-11.1
	1991-1995	0.6	1.7	0.4-5.1	0.4	2.6	0.3-9.6	0.2	1.0	0.0-6.3	1.4	4.2	1.7-8.7	0.6	4.2	0.9-12.2	0.8	4.4	1.2-11.4
	1996-2000	1.6	4.8	2.1-9.5	0.8	5.0	1.3-12.9	0.8	4.3	1.1-11.5	0.8	2.3	0.6-6.1	0.4	2.5	0.3-9.3	0.4	2.0	0.2-8.0
	2001-2005	1.2	3.2	1.2-7.1	1.0	6.0	1.9-14.4	0.2	1.0	0.0-6.4	1.0	2.7	0.9-6.6	0.6	4.0	0.8-11.8	0.4	2.0	0.2-8.0
	2006-2010	2.8	6.8	3.7-11.7	1.8	9.0	4.1-17.7	1.0	4.7	1.5-11.7	1.4	3.5	1.4-7.5	0.8	4.1	1.1-11.1	0.6	2.8	0.6-9.1
	2011-2015	3.8	8.3	5.0-13.5	2.6	11.2	5.8-20.1	1.2	5.2	1.9-12.4	2.8	6.6	3.6-11.4	2.0	9.6	4.5-18.5	0.8	3.6	1.0-10.3
2016-2020	3.8	7.0	4.2-11.6	3.2	11.6	6.6-20.1	0.6	2.5	0.5-9.0	2.0	3.7	1.8-7.6	1.6	6.1	2.6-13.3	0.4	1.5	0.2-7.3	
Pancreas	1976-1980	4.0	12.9	7.9-20.2	3.0	21.2	11.7-35.4	1.0	6.4	2.1-15.2	3.8	12.8	7.7-20.2	3.0	23.0	12.6-38.4	0.8	4.9	1.3-13.1
	1981-1985	2.8	8.8	4.8-14.9	1.2	9.3	3.2-20.8	1.6	8.9	3.8-18.0	2.2	6.8	3.4-12.4	1.0	7.7	2.3-18.5	1.2	7.1	2.6-15.8
	1986-1990	2.8	9.5	5.1-16.1	1.8	14.4	6.3-27.5	1.0	7.1	2.2-16.5	3.0	9.6	5.3-16.0	1.8	13.3	5.9-25.4	1.2	7.1	2.5-15.9
	1991-1995	4.8	14.1	9.0-21.0	2.6	17.8	9.1-31.0	2.2	11.7	5.8-21.3	4.4	12.7	7.9-19.4	2.4	16.6	8.2-29.6	2.0	10.2	4.8-19.3
	1996-2000	2.6	7.3	3.9-12.6	1.4	8.2	3.3-17.3	1.2	5.6	2.0-12.9	3.6	10.0	6.0-16.0	2.0	12.2	5.8-22.6	1.6	7.4	3.2-15.3
	2001-2005	4.8	12.8	8.2-19.2	1.8	10.4	4.7-20.3	3.0	14.8	8.3-25.0	3.4	9.1	5.3-14.7	1.8	10.4	4.7-20.2	1.6	7.9	3.3-16.2
	2006-2010	5.2	13.2	8.6-19.6	2.6	14.6	7.7-25.5	2.6	12.0	6.3-21.2	4.6	11.4	7.2-17.4	2.6	14.1	7.4-24.7	2.0	9.2	4.4-17.6
	2011-2015	5.0	12.2	7.8-18.4	2.6	12.6	6.7-22.4	2.4	11.1	5.5-20.7	3.8	9.0	5.4-14.4	2.2	11.5	5.7-21.2	1.6	6.6	2.8-14.2
2016-2020	7.2	14.0	9.7-19.9	2.8	11.9	6.5-21.1	4.4	16.1	10.0-25.7	5.0	10.5	6.6-16.1	2.2	9.9	4.9-18.7	2.8	11.6	6.0-21.1	
Larynx	1976-1980	1.2	3.6	1.3-8.0	1.2	7.7	2.8-17.4	0.0	0.0	0.0-5.4	0.4	1.2	0.1-4.7	0.4	2.6	0.3-10.1	0.0	0.0	0.0-5.4
	1981-1985	2.2	8.3	4.0-15.1	1.6	12.6	5.3-25.1	0.6	4.2	0.8-12.3	0.4	1.6	0.2-5.7	0.4	3.3	0.3-11.8	0.0	0.0	0.0-5.1
	1986-1990	1.8	5.5	2.5-10.6	1.6	11.4	4.8-22.7	0.2	1.3	0.0-7.4	0.4	1.0	0.1-4.1	0.4	2.0	0.2-8.6	0.0	0.0	0.0-4.8
	1991-1995	1.8	5.4	2.5-10.4	1.2	7.3	2.7-16.2	0.6	3.9	0.8-11.4	0.4	1.1	0.1-4.1	0.4	2.3	0.3-8.9	0.0	0.0	0.0-4.5
	1996-2000	1.8	5.1	2.3-9.8	1.6	9.6	4.1-19.3	0.2	1.0	0.0-6.4	0.6	1.7	0.4-5.2	0.6	3.8	0.8-11.3	0.0	0.0	0.0-4.7
	2001-2005	1.0	2.6	0.8-6.4	0.8	4.6	1.2-12.2	0.2	1.0	0.0-6.3	0.4	1.0	0.1-4.0	0.4	2.0	0.2-8.1	0.0	0.0	0.0-4.8
	2006-2010	1.4	2.8	1.1-6.4	1.4	5.7	2.3-12.8	0.0	0.0	0.0-4.8	0.0	0.0	0.0-2.4	0.0	0.0	0.0-4.9	0.0	0.0	0.0-4.8
	2011-2015	1.0	1.7	0.5-4.8	1.0	3.4	1.1-9.6	0.0	0.0	0.0-5.1	0.4	0.8	0.1-3.7	0.4	1.9	0.2-8.1	0.0	0.0	0.0-5.1
2016-2020	1.0	2.0	0.6-5.5	1.0	4.1	1.3-10.9	0.0	0.0	0.0-5.4	0.4	0.9	0.1-3.9	0.4	1.8	0.2-7.8	0.0	0.0	0.0-5.4	
Lung and bronchus	1976-1980	19.0	66.9	53.8-82.1	14.6	113.9	88.6-144.2	4.4	26.9	16.8-41.1	12.2	43.5	33.1-56.2	10.6	87.1	64.7-114.6	1.6	9.7	4.2-19.7
	1981-1985	20.0	65.0	52.6-79.4	12.8	93.3	71.2-120.2	7.2	44.7	30.9-62.6	18.4	60.0	48.1-73.9	11.6	86.0	64.5-112.3	6.8	39.9	27.4-56.3
	1986-1990	23.6	71.2	58.7-85.5	13.8	97.2	74.8-124.1	9.8	56.2	41.3-74.9	20.2	62.2	50.5-75.8	12.8	93.8	71.3-120.8	7.4	43.9	30.5-61.1
	1991-1995	27.0	78.0	65.3-92.5	15.4	96.1	75.4-120.7	11.6	62.6	47.4-81.4	24.4	71.0	58.8-84.9	14.0	93.1	71.9-118.5	10.4	57.5	42.8-75.8
	1996-2000	27.8	78.8	66.2-93.2	17.2	106.5	84.9-131.9	10.6	56.3	42.1-74.1	25.0	70.9	59.0-84.5	15.6	101.3	79.7-126.9	9.4	48.8	35.7-65.4
	2001-2005	32.6	85.2	72.6-99.6	17.2	97.2	77.5-120.6	15.4	77.3	60.9-97.1	26.0	67.4	56.3-80.2	14.8	84.8	66.3-107.0	11.2	55.2	41.7-72.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schoharie**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	30.4	74.5	63.0-87.7	17.4	87.9	70.0-109.1	13.0	61.5	47.2-79.1	23.0	55.9	46.0-67.4	13.6	69.5	53.7-88.8	9.4	42.6	31.2-57.3
	2011-2015	29.6	64.0	53.9-75.6	15.6	70.8	55.6-89.4	14.0	58.8	45.6-75.2	18.8	41.4	33.3-51.1	9.2	43.2	31.4-58.4	9.6	39.9	29.2-54.0
	2016-2020	32.0	62.4	52.8-73.6	16.6	66.5	52.5-83.8	15.4	58.6	45.8-74.8	21.6	44.9	36.4-55.0	11.4	49.1	36.7-64.9	10.2	41.0	29.9-55.5
Melanoma of the skin	1976-1980	2.8	11.8	6.3-19.8	2.2	19.9	9.8-35.5	0.6	3.5	0.7-10.8	0.8	3.3	0.9-8.4	0.6	4.9	1.0-14.5	0.2	2.3	0.1-10.8
	1981-1985	4.8	17.2	10.8-25.8	3.8	29.2	17.2-46.3	1.0	7.1	2.2-16.6	1.0	3.3	1.1-7.9	0.8	5.8	1.5-15.2	0.2	1.0	0.0-6.7
	1986-1990	3.6	11.8	6.9-18.8	2.0	14.6	6.9-27.1	1.6	9.7	4.0-19.5	1.0	3.6	1.1-8.3	1.0	8.5	2.6-19.9	0.0	0.0	0.0-4.8
	1991-1995	3.0	8.9	4.9-14.7	1.0	6.5	2.1-15.3	2.0	11.1	5.2-20.6	1.2	3.2	1.2-7.1	0.6	3.7	0.8-11.2	0.6	2.8	0.6-8.8
	1996-2000	4.2	12.7	7.8-19.5	2.8	17.1	9.3-28.9	1.4	8.0	3.2-16.9	1.0	3.0	1.0-7.1	1.0	6.1	2.0-14.6	0.0	0.0	0.0-4.7
	2001-2005	7.0	18.8	13.1-26.4	4.0	22.1	13.5-34.5	3.0	15.5	8.5-26.1	1.2	3.2	1.2-7.2	1.0	5.6	1.8-13.6	0.2	0.9	0.0-6.1
	2006-2010	5.4	14.5	9.4-21.4	2.4	11.1	5.7-20.1	3.0	17.9	9.7-30.1	1.6	3.6	1.5-7.4	1.2	6.2	2.2-14.1	0.4	1.5	0.2-6.8
	2011-2015	11.8	29.1	21.8-38.2	7.6	39.2	27.2-54.9	4.2	21.0	12.6-33.3	1.8	4.0	1.8-8.1	1.6	7.8	3.3-16.2	0.2	0.7	0.0-6.0
2016-2020	8.4	19.2	13.4-26.9	4.6	20.2	12.4-31.8	3.8	18.8	10.6-31.0	1.2	3.1	1.1-7.3	1.2	6.5	2.3-15.1	0.0	0.0	0.0-5.4	
Female breast	1976-1980							15.0	99.0	77.3-124.9							6.2	40.7	27.4-58.3
	1981-1985							16.4	101.0	79.8-126.2							8.2	49.0	34.8-67.2
	1986-1990							16.6	98.9	78.3-123.4							7.0	37.6	25.9-53.0
	1991-1995							16.2	91.2	72.1-113.9							5.8	33.0	21.9-47.7
	1996-2000							20.8	114.0	93.0-138.6							6.6	34.3	23.5-48.5
	2001-2005							28.2	143.6	120.6-170.0							4.4	20.4	12.8-31.6
	2006-2010							18.8	92.1	74.1-113.5							4.0	19.8	11.8-31.4
	2011-2015							26.0	119.4	99.1-143.1							2.6	11.0	5.8-19.7
2016-2020							31.0	132.7	110.8-158.0							3.6	14.2	8.2-23.7	
Cervix uteri	1976-1980							2.0	14.9	6.9-27.6							1.2	8.5	3.0-18.6
	1981-1985							1.8	12.8	5.7-24.4							0.6	3.3	0.7-10.3
	1986-1990							2.2	17.1	8.4-30.3							0.2	1.8	0.0-8.6
	1991-1995							1.8	11.9	5.4-22.4							0.6	4.4	0.9-12.4
	1996-2000							2.2	12.4	6.1-22.5							0.6	3.2	0.6-9.8
	2001-2005							2.0	12.9	6.0-24.0							0.4	2.1	0.3-8.1
	2006-2010							1.6	10.1	4.2-20.2							1.2	6.4	2.2-14.5
	2011-2015							0.4	2.7	0.2-10.3							0.4	1.4	0.2-6.9
	2016-2020							1.6	10.0	4.0-20.7							0.4	1.8	0.2-8.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schoharie**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							3.6	27.4	15.9-43.8							1.4	9.1	3.6-19.0
	1981-1985							4.4	30.7	18.9-46.9							0.4	2.1	0.3-8.6
	1986-1990							4.8	27.3	17.3-41.1							0.2	1.8	0.0-8.6
	1991-1995							4.0	22.5	13.6-35.1							1.4	7.3	2.9-15.5
	1996-2000							4.8	26.6	17.0-39.9							1.4	7.8	3.1-16.4
	2001-2005							4.8	24.3	15.5-36.6							1.0	5.0	1.6-12.2
	2006-2010							7.0	32.4	22.4-45.9							1.0	5.2	1.5-13.0
	2011-2015							10.2	40.8	30.2-54.6							2.2	10.4	4.9-19.8
	2016-2020							7.8	31.1	21.7-44.1							0.6	2.3	0.5-8.4
Ovary	1976-1980							2.2	15.8	7.7-28.5							2.2	15.4	7.4-28.0
	1981-1985							2.0	13.3	6.1-25.1							1.0	5.5	1.8-13.5
	1986-1990							2.6	15.2	8.0-26.4							1.8	9.3	4.2-18.1
	1991-1995							2.8	16.3	8.8-27.5							1.2	6.2	2.2-13.9
	1996-2000							2.6	14.2	7.5-24.6							2.4	13.3	6.8-23.5
	2001-2005							2.4	12.7	6.5-22.7							1.6	7.5	3.2-15.5
	2006-2010							3.6	17.6	10.2-28.7							2.6	12.5	6.5-22.1
	2011-2015							1.4	5.4	2.1-12.4							1.0	3.4	1.1-9.6
	2016-2020							1.6	7.3	2.8-16.1							1.4	5.1	2.0-12.0
Prostate	1976-1980				9.2	77.5	56.2-104.1							2.4	20.1	10.0-35.6			
	1981-1985				9.2	75.5	54.4-101.7							4.6	39.5	24.7-59.4			
	1986-1990				15.2	113.5	88.4-143.3							3.6	32.4	18.7-51.5			
	1991-1995				24.4	156.7	129.5-188.0							5.6	44.3	29.1-64.3			
	1996-2000				20.6	124.4	101.3-151.3							6.2	43.9	29.5-62.7			
	2001-2005				21.6	119.6	97.8-145.1							2.8	17.9	9.7-30.2			
	2006-2010				24.6	119.5	98.9-143.4							4.0	23.3	14.1-36.4			
	2011-2015				18.2	75.6	60.5-93.9							3.4	17.9	10.3-29.3			
	2016-2020				28.0	108.1	90.6-128.6							5.0	20.6	13.2-31.4			
Testis	1976-1980				0.8	7.1	1.9-17.8							0.0	0.0	0.0-6.0			
	1981-1985				1.0	6.8	2.1-16.5							0.2	1.4	0.0-8.2			
	1986-1990				0.8	5.1	1.4-13.5							0.0	0.0	0.0-5.5			
	1991-1995				0.6	3.9	0.8-11.6							0.0	0.0	0.0-5.0			
	1996-2000				1.0	7.0	2.3-16.3							0.0	0.0	0.0-5.0			
	2001-2005				1.0	7.3	2.3-17.0							0.0	0.0	0.0-5.0			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schoharie**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				1.6	10.7	4.4-21.3							0.0	0.0	0.0-4.9			
	2011-2015				1.2	8.4	3.0-18.4							0.0	0.0	0.0-5.1			
	2016-2020				1.0	7.1	2.2-17.0							0.0	0.0	0.0-5.3			
Urinary bladder (incl. in situ)	1976-1980	7.6	25.2	17.8-34.8	5.6	42.3	27.8-61.7	2.0	11.5	5.5-21.7	2.0	7.3	3.5-13.4	1.8	16.1	7.2-30.6	0.2	1.2	0.0-7.4
	1981-1985	6.6	21.1	14.5-29.9	5.4	42.8	27.7-62.9	1.2	6.3	2.3-14.3	1.8	5.8	2.6-11.2	0.8	7.2	1.8-18.2	1.0	5.6	1.8-13.7
	1986-1990	6.2	20.2	13.6-28.8	4.6	35.4	22.0-53.6	1.6	9.8	4.1-19.6	2.0	6.3	3.0-11.7	1.2	10.6	3.8-22.8	0.8	4.6	1.2-12.1
	1991-1995	8.6	25.2	18.2-34.1	6.0	40.7	27.0-58.8	2.6	14.1	7.4-24.5	2.0	5.9	2.8-10.9	1.0	8.3	2.5-19.2	1.0	4.4	1.4-11.0
	1996-2000	7.2	20.3	14.2-28.2	5.2	33.5	21.7-49.5	2.0	9.9	4.7-18.9	3.2	9.0	5.2-14.8	1.6	12.7	5.4-24.9	1.6	7.6	3.2-15.6
	2001-2005	11.2	29.9	22.5-38.9	8.4	49.8	35.7-67.7	2.8	14.0	7.6-24.1	2.2	5.8	2.9-10.7	1.8	10.6	4.8-20.5	0.4	1.9	0.2-7.7
	2006-2010	9.8	23.4	17.2-31.2	7.2	36.6	25.4-51.3	2.6	11.6	6.1-20.6	2.6	6.2	3.3-11.0	2.0	10.6	5.0-20.1	0.6	2.4	0.5-8.3
	2011-2015	13.6	29.9	23.1-38.3	10.8	49.1	36.7-64.9	2.8	12.0	6.5-21.1	1.6	3.5	1.5-7.4	1.4	6.7	2.6-14.8	0.2	0.8	0.0-6.2
	2016-2020	13.2	26.4	20.2-34.1	10.2	42.1	31.0-56.5	3.0	11.9	6.6-20.9	1.4	2.7	1.1-6.3	1.0	3.9	1.3-10.6	0.4	1.6	0.2-7.5
Kidney and renal pelvis	1976-1980	2.2	7.8	3.9-14.1	1.0	9.8	3.0-22.6	1.2	6.9	2.5-15.6	2.0	7.0	3.3-13.0	0.8	7.7	2.0-19.4	1.2	7.1	2.6-15.9
	1981-1985	2.2	7.3	3.6-13.3	1.6	10.9	4.6-21.9	0.6	4.5	0.8-13.1	1.2	4.0	1.4-8.9	0.8	5.6	1.5-14.9	0.4	3.1	0.4-10.9
	1986-1990	2.0	6.3	2.9-11.7	1.2	6.4	2.3-14.8	0.8	5.7	1.5-14.7	1.2	3.7	1.3-8.2	0.6	3.1	0.6-10.0	0.6	4.1	0.8-11.9
	1991-1995	4.4	13.0	8.1-19.8	2.8	17.5	9.5-29.7	1.6	9.1	3.9-18.2	1.8	5.2	2.4-10.0	1.2	8.0	2.9-17.6	0.6	3.2	0.7-9.7
	1996-2000	1.6	4.5	1.9-8.9	1.0	6.4	2.0-15.3	0.6	3.2	0.7-9.9	1.0	2.9	0.9-6.8	1.0	6.6	2.1-15.6	0.0	0.0	0.0-4.7
	2001-2005	5.2	13.7	8.9-20.2	3.2	17.6	10.0-29.2	2.0	10.0	4.8-18.8	0.8	2.1	0.6-5.6	0.8	4.7	1.2-12.6	0.0	0.0	0.0-4.8
	2006-2010	5.4	14.2	9.2-21.0	3.8	20.9	12.4-33.3	1.6	7.3	3.1-15.2	2.0	4.8	2.3-9.1	1.4	7.6	3.0-16.2	0.6	2.4	0.5-8.3
	2011-2015	9.4	22.3	16.1-30.3	7.0	32.9	22.5-46.8	2.4	12.2	6.0-22.6	2.8	6.5	3.5-11.2	1.8	9.3	4.2-18.4	1.0	4.0	1.3-10.6
	2016-2020	10.0	19.9	14.6-26.9	6.8	27.5	18.7-39.7	3.2	12.5	7.0-21.7	2.2	4.3	2.2-8.4	1.0	4.1	1.3-11.0	1.2	4.6	1.7-11.5
Brain and other central nervous system	1976-1980	1.2	3.5	1.3-8.1	1.0	5.9	1.9-14.7	0.2	1.5	0.0-8.2	1.8	5.8	2.7-11.3	1.0	6.5	2.1-15.9	0.8	5.2	1.4-13.7
	1981-1985	1.6	4.8	2.1-9.7	1.2	7.5	2.7-17.0	0.4	2.6	0.3-9.7	1.0	3.2	1.0-7.7	0.6	4.3	0.9-12.9	0.4	2.6	0.3-9.7
	1986-1990	1.8	5.8	2.6-11.2	1.2	8.4	3.0-18.5	0.6	3.4	0.7-10.3	1.2	3.6	1.3-8.1	0.4	2.6	0.3-10.0	0.8	4.5	1.2-12.0
	1991-1995	1.8	5.8	2.6-10.9	0.6	4.3	0.9-12.5	1.2	7.3	2.6-15.9	0.2	0.5	0.0-3.2	0.0	0.0	0.0-5.0	0.2	1.1	0.0-6.4
	1996-2000	2.2	6.2	3.1-11.2	1.8	11.0	5.0-21.1	0.4	2.2	0.3-8.5	1.4	4.1	1.7-8.6	0.8	4.7	1.3-12.4	0.6	3.6	0.7-10.6
	2001-2005	1.8	4.7	2.1-9.2	1.2	6.3	2.3-14.3	0.6	2.8	0.6-9.0	1.4	3.8	1.5-8.0	0.6	3.4	0.7-10.4	0.8	3.8	1.0-10.4
	2006-2010	2.6	7.4	3.8-13.0	1.4	8.7	3.2-18.5	1.2	6.1	2.2-13.8	1.6	3.8	1.6-7.8	1.2	5.7	2.1-13.1	0.4	1.8	0.2-7.5
	2011-2015	2.8	7.0	3.7-12.2	1.2	5.7	2.0-13.3	1.6	8.3	3.4-17.2	2.4	5.0	2.6-9.3	1.2	4.9	1.8-12.0	1.2	5.1	1.8-12.2
	2016-2020	3.4	7.5	4.1-12.9	3.0	12.3	6.5-21.8	0.4	3.1	0.3-11.3	2.6	5.1	2.6-9.4	2.4	9.3	4.6-17.8	0.2	0.9	0.0-6.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schoharie**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	0.6	2.0	0.4-6.1	0.0	0.0	0.0-6.0	0.6	3.9	0.8-11.8	0.2	0.6	0.0-3.7	0.0	0.0	0.0-6.0	0.2	1.1	0.0-7.3
	1981-1985	0.2	0.6	0.0-3.6	0.2	1.3	0.0-8.1	0.0	0.0	0.0-5.1	0.0	0.0	0.0-2.7	0.0	0.0	0.0-5.8	0.0	0.0	0.0-5.1
	1986-1990	1.6	4.6	1.9-9.4	0.4	2.0	0.2-8.6	1.2	6.6	2.3-15.0	0.4	1.3	0.2-4.8	0.0	0.0	0.0-5.5	0.4	2.0	0.2-8.1
	1991-1995	1.4	4.3	1.7-8.9	0.8	5.3	1.4-13.7	0.6	3.4	0.7-10.1	0.0	0.0	0.0-2.3	0.0	0.0	0.0-5.0	0.0	0.0	0.0-4.5
	1996-2000	4.0	12.7	7.7-19.6	1.4	9.8	3.9-20.2	2.6	16.4	8.6-28.1	0.2	0.6	0.0-3.3	0.2	2.0	0.0-9.3	0.0	0.0	0.0-4.7
	2001-2005	3.0	10.0	5.5-16.6	0.4	2.6	0.3-9.5	2.6	17.7	9.2-30.4	0.4	1.0	0.1-4.0	0.4	2.0	0.2-8.1	0.0	0.0	0.0-4.8
	2006-2010	5.0	14.3	9.1-21.5	1.0	5.2	1.6-12.9	4.0	23.4	14.0-36.7	0.0	0.0	0.0-2.4	0.0	0.0	0.0-4.9	0.0	0.0	0.0-4.8
	2011-2015	5.0	15.0	9.4-22.6	2.0	12.3	5.5-23.5	3.0	17.5	9.5-29.6	0.0	0.0	0.0-2.6	0.0	0.0	0.0-5.1	0.0	0.0	0.0-5.1
	2016-2020	6.4	18.1	12.0-26.4	1.2	6.8	2.1-16.2	5.2	29.7	18.7-44.8	0.2	0.4	0.0-3.3	0.0	0.0	0.0-5.3	0.2	0.9	0.0-6.6
Hodgkin lymphoma	1976-1980	1.0	3.7	1.1-9.0	0.6	4.5	0.8-13.7	0.4	3.0	0.2-11.5	0.0	0.0	0.0-2.8	0.0	0.0	0.0-6.0	0.0	0.0	0.0-5.4
	1981-1985	0.6	2.6	0.5-7.3	0.2	1.7	0.0-9.0	0.4	3.7	0.4-12.5	0.0	0.0	0.0-2.7	0.0	0.0	0.0-5.8	0.0	0.0	0.0-5.1
	1986-1990	0.4	1.2	0.1-4.5	0.2	1.0	0.0-7.1	0.2	1.3	0.0-7.4	0.2	0.6	0.0-3.6	0.2	1.8	0.0-9.2	0.0	0.0	0.0-4.8
	1991-1995	0.8	2.7	0.7-6.8	0.0	0.0	0.0-5.0	0.8	5.3	1.4-13.3	0.2	0.8	0.0-3.9	0.2	1.5	0.0-8.0	0.0	0.0	0.0-4.5
	1996-2000	1.2	3.7	1.4-8.2	0.4	2.2	0.3-8.5	0.8	5.2	1.4-13.3	0.2	0.6	0.0-3.4	0.2	1.3	0.0-7.4	0.0	0.0	0.0-4.7
	2001-2005	1.6	4.6	2.0-9.3	0.8	4.9	1.3-12.8	0.8	4.3	1.2-11.5	0.2	0.6	0.0-3.4	0.2	1.3	0.0-7.2	0.0	0.0	0.0-4.8
	2006-2010	1.0	2.4	0.8-6.0	0.8	3.8	1.0-10.6	0.2	1.0	0.0-6.5	0.2	0.6	0.0-3.4	0.0	0.0	0.0-4.9	0.2	1.1	0.0-6.6
	2011-2015	0.6	1.8	0.3-5.5	0.4	2.6	0.2-9.8	0.2	0.9	0.0-6.4	0.2	0.5	0.0-3.3	0.0	0.0	0.0-5.1	0.2	0.9	0.0-6.4
	2016-2020	1.2	3.8	1.3-8.6	0.8	4.8	1.3-12.9	0.4	2.7	0.2-10.5	0.0	0.0	0.0-2.7	0.0	0.0	0.0-5.3	0.0	0.0	0.0-5.4
Non-Hodgkin lymphomas	1976-1980	4.2	14.2	8.7-21.9	0.8	6.6	1.7-16.9	3.4	21.1	12.2-34.3	3.0	10.1	5.6-16.9	0.6	4.5	0.9-13.6	2.4	14.9	7.6-26.5
	1981-1985	4.4	15.4	9.5-23.5	2.6	23.1	11.9-39.7	1.8	11.6	5.2-22.6	1.4	4.3	1.7-9.1	0.6	3.8	0.8-11.7	0.8	4.4	1.2-12.0
	1986-1990	5.4	16.8	11.0-24.6	2.2	15.4	7.5-28.0	3.2	18.9	10.6-31.1	2.4	7.5	3.8-13.2	0.6	4.4	0.9-13.0	1.8	10.1	4.6-19.6
	1991-1995	4.8	14.0	8.9-20.9	2.2	14.8	7.2-26.9	2.6	13.6	7.1-23.6	2.2	6.8	3.4-12.2	0.6	5.2	1.0-14.6	1.6	8.3	3.5-16.9
	1996-2000	6.6	19.1	13.1-26.9	3.2	19.7	11.1-32.4	3.4	18.3	10.6-29.6	2.2	6.1	3.0-11.0	0.8	5.5	1.4-14.1	1.4	7.1	2.8-15.0
	2001-2005	8.2	22.2	15.9-30.3	4.2	24.9	15.3-38.5	4.0	20.5	12.5-32.2	2.6	7.0	3.7-12.2	1.6	10.4	4.4-20.7	1.0	5.0	1.6-12.4
	2006-2010	7.0	17.9	12.3-25.4	4.6	23.7	14.7-36.5	2.4	12.1	6.1-21.7	2.6	6.5	3.4-11.5	2.0	10.9	5.0-20.7	0.6	2.8	0.6-9.1
	2011-2015	8.4	19.2	13.8-26.4	5.6	26.9	17.7-39.7	2.8	12.3	6.6-21.6	3.2	7.7	4.3-13.0	2.8	14.3	7.6-25.1	0.4	1.7	0.2-7.5
	2016-2020	7.6	15.9	11.1-22.6	3.8	16.1	9.6-26.1	3.8	16.3	9.4-26.9	3.6	7.3	4.2-12.2	2.0	7.8	3.7-15.6	1.6	6.7	2.7-14.6
Myeloma	1976-1980	1.0	3.2	1.0-7.6	0.6	3.8	0.8-11.8	0.4	2.4	0.3-9.4	0.6	2.0	0.4-6.1	0.0	0.0	0.0-6.0	0.6	3.4	0.7-10.7
	1981-1985	1.6	5.1	2.2-10.2	1.0	8.1	2.4-19.3	0.6	3.3	0.7-10.3	0.6	1.8	0.4-5.5	0.6	4.2	0.8-12.8	0.0	0.0	0.0-5.1
	1986-1990	1.2	3.9	1.4-8.6	0.6	5.6	1.1-16.0	0.6	2.8	0.6-9.0	0.6	2.1	0.4-6.2	0.4	3.1	0.4-11.2	0.2	0.9	0.0-6.3
	1991-1995	1.4	4.0	1.6-8.4	1.0	6.1	2.0-14.6	0.4	1.8	0.2-7.3	0.6	1.6	0.3-4.9	0.6	4.5	0.8-13.2	0.0	0.0	0.0-4.5
	1996-2000	3.0	8.6	4.8-14.2	1.4	8.2	3.3-17.2	1.6	7.9	3.4-16.1	1.6	4.4	1.9-8.9	0.6	3.5	0.7-10.6	1.0	4.3	1.4-11.0
	2001-2005	2.8	7.4	4.1-12.7	1.0	5.6	1.8-13.6	1.8	8.8	4.0-17.3	2.0	5.2	2.5-9.7	1.2	6.2	2.3-14.1	0.8	3.7	1.0-10.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schoharie**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	2.2	5.3	2.6-9.8	1.4	7.8	3.1-16.5	0.8	3.2	0.9-9.4	1.2	2.8	1.0-6.5	0.6	2.8	0.6-9.2	0.6	2.7	0.6-8.9
	2011-2015	3.0	6.6	3.7-11.4	1.8	7.7	3.4-15.8	1.2	5.7	2.0-13.4	1.0	2.2	0.7-5.6	0.6	2.8	0.5-9.2	0.4	1.5	0.2-7.1
	2016-2020	3.0	5.7	3.1-10.3	2.2	7.7	3.8-15.3	0.8	3.8	0.8-11.5	1.0	2.2	0.7-5.7	0.8	3.5	0.9-10.2	0.2	1.1	0.0-7.2
Leukemias	1976-1980	3.0	11.7	6.4-19.5	1.2	8.2	2.9-18.4	1.8	14.4	6.3-27.7	1.6	5.4	2.2-10.9	0.4	2.1	0.3-9.1	1.2	8.2	2.8-18.4
	1981-1985	3.2	11.6	6.5-19.0	1.8	14.5	6.4-28.0	1.4	8.5	3.3-18.2	2.8	9.6	5.1-16.3	1.4	11.6	4.5-24.0	1.4	8.3	3.1-17.9
	1986-1990	5.8	17.7	11.8-25.7	3.6	24.9	14.4-40.2	2.2	12.8	6.2-23.4	2.8	8.6	4.7-14.5	1.6	11.3	4.6-23.0	1.2	6.3	2.2-14.3
	1991-1995	2.4	6.9	3.5-12.1	1.6	10.3	4.2-20.8	0.8	4.5	1.2-11.8	1.8	5.4	2.5-10.4	1.0	7.5	2.3-17.6	0.8	4.6	1.2-11.9
	1996-2000	3.0	8.6	4.8-14.3	1.6	9.6	4.1-19.2	1.4	7.3	2.9-15.5	1.6	4.4	1.9-8.9	0.6	3.3	0.7-10.2	1.0	4.8	1.5-12.0
	2001-2005	6.2	17.0	11.5-24.4	3.6	20.4	12.0-32.8	2.6	13.8	7.2-24.2	2.2	5.8	2.9-10.5	1.0	6.1	2.0-14.5	1.2	5.4	2.0-12.5
	2006-2010	5.4	14.1	9.1-20.9	3.6	20.3	11.8-32.7	1.8	8.8	3.8-17.6	3.0	8.0	4.4-13.5	2.4	13.9	7.0-24.8	0.6	3.4	0.6-10.6
	2011-2015	6.6	14.4	9.9-20.6	4.6	20.6	12.9-31.8	2.0	8.6	4.1-16.9	2.6	6.0	3.2-10.7	1.2	6.0	2.2-14.0	1.4	6.3	2.5-14.0
	2016-2020	7.8	16.8	11.6-23.8	5.6	25.5	16.3-38.4	2.2	8.2	4.0-15.9	3.2	6.9	3.9-11.7	2.0	9.3	4.4-18.0	1.2	4.8	1.7-12.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schuyler**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	59.6	354.7	314.9-398.1	31.2	426.2	359.5-501.2	28.4	310.5	260.4-367.3	28.4	169.4	142.2-200.2	15.6	211.6	165.8-265.7	12.8	137.5	105.1-176.6
	1981-1985	76.0	430.9	387.9-477.4	36.8	469.7	402.1-545.1	39.2	414.5	357.0-478.5	36.2	204.2	175.0-236.8	17.8	229.4	182.6-284.1	18.4	186.4	149.5-229.8
	1986-1990	81.4	435.2	393.5-480.1	40.8	502.8	434.6-578.5	40.6	398.5	344.6-458.6	44.0	235.8	205.3-269.5	25.6	329.9	273.7-393.7	18.4	177.8	142.7-219.1
	1991-1995	96.2	474.6	432.8-519.3	50.0	559.5	490.9-634.8	46.2	429.1	374.6-489.4	43.4	213.4	185.8-244.0	20.6	240.4	195.4-292.4	22.8	197.8	162.7-238.6
	1996-2000	107.6	505.8	463.9-550.7	54.2	571.0	503.9-644.5	53.4	457.6	403.6-517.2	44.0	203.9	177.8-232.9	23.8	267.8	220.8-321.7	20.2	162.9	132.2-199.0
	2001-2005	116.8	517.5	476.0-561.8	61.8	599.2	533.1-671.5	55.0	448.8	396.5-506.6	42.8	187.0	162.6-214.1	21.8	225.6	184.4-273.3	21.0	163.2	133.1-198.9
	2006-2010	118.2	498.4	458.3-541.4	59.8	551.5	488.8-620.4	58.4	468.3	414.9-527.3	42.8	178.5	155.1-204.8	24.0	234.8	193.5-282.6	18.8	144.9	116.6-178.8
	2011-2015	123.6	481.9	443.6-523.0	64.4	537.4	478.1-602.7	59.2	449.0	397.5-506.1	44.0	170.6	148.4-195.5	23.0	201.1	164.7-243.6	21.0	150.6	122.7-183.9
	2016-2020	135.6	504.0	464.5-546.3	67.2	506.2	450.3-567.9	68.4	513.6	457.1-575.9	49.8	182.9	160.0-208.6	26.4	208.8	173.4-250.0	23.4	163.5	133.7-199.0
Oral cavity and pharynx	1976-1980	2.2	12.8	6.3-23.1	1.6	18.9	8.0-38.0	0.6	6.4	1.3-19.2	0.2	1.3	0.0-7.1	0.2	2.7	0.1-14.8	0.0	0.0	0.0-8.5
	1981-1985	1.4	7.2	2.8-15.3	1.0	11.1	3.4-27.0	0.4	5.1	0.6-17.8	0.8	4.0	1.1-10.5	0.4	3.8	0.5-15.2	0.4	4.0	0.5-15.0
	1986-1990	1.2	6.2	2.2-13.7	0.8	8.8	2.3-23.0	0.4	3.7	0.4-14.1	0.4	2.1	0.2-7.6	0.4	4.4	0.5-16.4	0.0	0.0	0.0-7.9
	1991-1995	2.0	9.7	4.6-18.0	1.0	10.3	3.4-24.4	1.0	8.8	2.8-21.3	0.0	0.0	0.0-3.9	0.0	0.0	0.0-8.2	0.0	0.0	0.0-7.7
	1996-2000	0.8	3.9	1.1-10.1	0.6	6.2	1.3-18.4	0.2	1.9	0.0-11.1	0.2	1.0	0.0-5.6	0.0	0.0	0.0-8.1	0.2	1.9	0.0-11.1
	2001-2005	1.6	6.6	2.8-13.4	1.0	9.0	2.9-22.0	0.6	4.7	1.0-15.1	0.0	0.0	0.0-4.0	0.0	0.0	0.0-8.2	0.0	0.0	0.0-8.0
	2006-2010	4.0	16.5	9.9-26.2	2.8	23.8	12.7-41.5	1.2	8.8	3.2-21.0	0.8	3.1	0.8-8.8	0.4	2.6	0.3-12.3	0.4	3.3	0.4-13.7
	2011-2015	4.4	18.2	11.1-28.5	2.8	23.4	12.1-41.5	1.6	13.6	5.7-28.4	0.4	1.3	0.2-6.1	0.4	2.5	0.3-12.3	0.0	0.0	0.0-8.8
	2016-2020	3.6	13.5	7.7-22.4	2.2	16.3	7.6-31.6	1.4	10.7	4.1-24.4	0.8	2.5	0.6-7.9	0.8	4.9	1.2-15.8	0.0	0.0	0.0-9.1
Esophagus	1976-1980	0.8	4.5	1.2-11.7	0.8	9.4	2.5-24.6	0.0	0.0	0.0-8.5	0.2	1.2	0.0-6.8	0.2	2.5	0.1-14.3	0.0	0.0	0.0-8.5
	1981-1985	0.6	3.5	0.7-10.3	0.4	5.0	0.6-18.3	0.2	1.9	0.0-11.5	0.8	4.8	1.2-12.3	0.8	9.7	2.6-25.3	0.0	0.0	0.0-8.2
	1986-1990	0.6	2.9	0.6-8.8	0.6	7.6	1.4-22.3	0.0	0.0	0.0-7.9	0.4	2.2	0.3-7.9	0.4	5.7	0.6-19.9	0.0	0.0	0.0-7.9
	1991-1995	0.8	4.0	1.1-10.3	0.8	8.0	2.2-21.0	0.0	0.0	0.0-7.7	0.6	3.2	0.7-9.4	0.2	2.2	0.1-12.4	0.4	4.3	0.5-15.3
	1996-2000	1.6	7.2	3.1-14.5	1.0	9.3	3.0-22.3	0.6	4.5	0.9-14.7	1.2	5.6	2.1-12.4	1.0	10.0	3.2-23.7	0.2	1.3	0.0-9.8
	2001-2005	1.2	5.2	1.9-11.7	0.8	7.4	2.0-19.9	0.4	3.0	0.4-12.5	1.0	4.3	1.4-10.4	0.8	7.2	1.9-19.4	0.2	1.3	0.0-9.9
	2006-2010	1.6	7.7	3.3-15.5	1.4	14.6	5.8-30.7	0.2	1.1	0.0-9.9	0.6	2.2	0.5-7.5	0.4	3.0	0.4-13.2	0.2	1.5	0.0-10.9
	2011-2015	0.8	2.8	0.8-8.3	0.8	6.7	1.7-19.0	0.0	0.0	0.0-8.8	2.0	7.7	3.6-15.0	1.8	16.3	7.2-32.3	0.2	1.3	0.0-10.5
	2016-2020	1.0	4.0	1.2-10.4	1.0	8.5	2.6-21.9	0.0	0.0	0.0-9.1	0.8	3.3	0.8-9.4	0.8	7.2	1.8-20.1	0.0	0.0	0.0-9.1
Stomach	1976-1980	1.0	6.4	2.0-14.8	0.8	14.8	3.9-36.0	0.2	1.7	0.0-11.3	0.2	1.0	0.0-6.0	0.2	2.1	0.1-13.2	0.0	0.0	0.0-8.5
	1981-1985	2.0	10.8	5.1-20.0	1.6	18.0	7.7-36.0	0.4	3.7	0.4-14.4	1.0	5.9	1.9-13.8	0.8	10.8	2.8-27.8	0.2	1.6	0.0-10.8
	1986-1990	1.6	8.3	3.6-16.5	1.2	16.5	5.9-35.4	0.4	3.3	0.4-13.2	1.2	6.3	2.3-13.8	1.0	13.9	4.4-32.0	0.2	1.6	0.0-10.4
	1991-1995	2.0	9.5	4.5-17.6	0.8	8.1	2.2-21.3	1.2	11.4	4.1-25.1	1.0	4.8	1.6-11.4	0.2	1.9	0.0-11.5	0.8	7.2	1.9-19.0
	1996-2000	1.6	7.4	3.2-14.8	1.4	15.7	6.2-32.7	0.2	1.3	0.0-9.8	1.2	5.4	2.0-12.1	1.0	13.2	4.1-30.3	0.2	1.3	0.0-9.8
	2001-2005	1.4	6.0	2.4-12.7	1.0	9.1	2.9-22.2	0.4	3.3	0.4-13.2	1.0	4.2	1.4-10.4	0.4	3.9	0.4-14.8	0.6	4.8	1.0-15.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schuyler**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	1.6	6.9	2.9-14.4	1.2	11.3	3.9-25.9	0.4	3.3	0.4-13.7	0.8	3.7	1.0-9.9	0.4	3.6	0.4-14.5	0.4	3.5	0.4-14.2
	2011-2015	0.8	3.1	0.8-9.0	0.6	4.4	0.8-15.3	0.2	2.1	0.1-12.5	0.4	1.4	0.2-6.3	0.2	1.1	0.0-10.2	0.2	1.6	0.0-11.2
	2016-2020	1.8	6.7	3.0-13.8	1.4	10.1	4.0-22.8	0.4	3.6	0.3-15.0	0.8	2.5	0.7-7.9	0.4	3.0	0.4-13.4	0.4	2.1	0.3-11.7
Colorectal	1976-1980	9.8	58.5	43.1-77.6	4.6	64.9	40.3-98.3	5.2	54.7	35.5-80.7	4.6	28.2	17.7-42.4	1.6	20.8	8.9-41.2	3.0	30.9	17.1-51.4
	1981-1985	12.0	63.9	48.6-82.6	4.8	60.5	38.1-91.1	7.2	69.1	48.2-96.3	5.4	29.8	19.5-43.7	2.8	36.9	19.6-62.5	2.6	24.7	13.1-42.9
	1986-1990	10.2	55.0	40.8-72.5	5.0	63.8	40.8-94.7	5.2	48.9	31.8-72.3	4.2	22.0	13.6-33.8	1.8	21.6	9.6-41.5	2.4	21.3	11.0-37.9
	1991-1995	13.2	63.8	49.3-81.4	7.4	90.9	63.3-126.0	5.8	49.8	33.1-72.4	4.4	20.6	12.9-31.4	3.0	37.8	20.8-62.6	1.4	11.2	4.5-23.9
	1996-2000	15.4	71.7	56.6-89.8	7.0	76.5	52.8-107.0	8.4	67.1	48.1-91.7	5.4	24.6	16.2-36.0	3.0	35.7	19.6-59.1	2.4	19.0	9.7-34.3
	2001-2005	15.0	65.4	51.4-82.4	7.0	69.7	48.2-97.8	8.0	58.4	41.5-80.8	5.6	23.3	15.5-34.1	2.6	28.2	14.6-48.9	3.0	20.7	11.5-35.6
	2006-2010	12.8	53.1	40.7-68.3	5.0	46.7	29.8-70.4	7.8	56.4	39.9-78.6	5.2	20.4	13.3-30.5	2.4	21.8	11.0-39.3	2.8	19.6	10.7-34.5
	2011-2015	12.4	53.8	40.8-69.9	7.0	60.8	41.6-86.3	5.4	47.7	30.7-71.2	3.4	13.4	7.7-22.1	1.4	11.6	4.5-25.7	2.0	14.4	6.7-28.4
2016-2020	11.4	44.4	33.0-59.0	5.2	43.6	27.5-66.3	6.2	45.4	29.9-67.2	3.6	14.1	8.0-23.5	2.6	21.5	10.8-39.3	1.0	6.0	1.9-16.9	
Colon excluding rectum	1976-1980	7.4	44.3	31.0-61.3	3.4	44.0	25.2-71.2	4.0	43.0	26.1-66.9	4.0	24.4	14.8-37.9	1.4	17.7	7.0-36.8	2.6	26.9	14.2-46.6
	1981-1985	9.4	49.7	36.4-66.5	3.8	46.9	27.7-74.1	5.6	53.4	35.4-77.8	4.6	25.3	15.9-38.2	2.2	27.5	13.4-49.9	2.4	23.1	11.9-40.9
	1986-1990	7.8	41.5	29.4-56.8	3.4	42.9	24.6-69.0	4.4	40.6	25.3-62.1	3.4	17.3	10.0-27.9	1.6	18.3	7.7-36.5	1.8	15.7	7.1-30.5
	1991-1995	10.4	50.7	37.8-66.6	5.2	68.6	44.3-100.6	5.2	43.5	28.2-64.6	3.8	17.8	10.7-28.0	2.4	31.6	16.0-55.1	1.4	11.2	4.5-23.9
	1996-2000	11.2	51.9	39.2-67.6	5.6	62.2	40.9-90.5	5.6	44.4	29.3-65.2	4.6	20.8	13.2-31.5	2.8	33.2	17.8-56.0	1.8	13.4	6.1-26.8
	2001-2005	11.6	51.0	38.6-66.2	4.8	50.2	31.8-75.3	6.8	50.1	34.4-71.3	4.8	20.0	12.8-30.2	2.2	24.9	12.1-45.1	2.6	17.7	9.4-31.7
	2006-2010	9.4	39.5	28.9-53.0	3.4	33.6	19.2-54.9	6.0	43.2	28.9-63.2	4.6	18.0	11.4-27.7	2.0	18.4	8.6-35.1	2.6	18.3	9.7-32.9
	2011-2015	9.4	42.4	30.7-57.3	4.8	43.8	27.5-66.7	4.6	40.7	25.0-63.0	2.4	9.3	4.7-16.9	1.0	8.5	2.6-21.6	1.4	9.5	3.7-21.9
2016-2020	7.0	27.3	18.6-39.2	2.4	21.0	10.3-38.8	4.6	33.0	20.2-52.2	2.6	10.9	5.5-19.8	1.8	16.2	6.8-33.0	0.8	4.9	1.3-15.7	
Rectum & rectosigmoid	1976-1980	2.4	14.2	7.2-24.9	1.2	20.9	7.3-44.6	1.2	11.7	4.2-26.2	0.6	3.7	0.7-10.9	0.2	3.1	0.1-16.0	0.4	3.9	0.5-15.1
	1981-1985	2.6	14.2	7.5-24.5	1.0	13.6	4.1-32.0	1.6	15.7	6.6-31.7	0.8	4.5	1.2-11.7	0.6	9.3	1.8-26.5	0.2	1.6	0.0-10.8
	1986-1990	2.4	13.5	6.9-23.7	1.6	20.9	8.8-41.2	0.8	8.3	2.2-21.6	0.8	4.7	1.3-11.9	0.2	3.4	0.1-16.1	0.6	5.7	1.2-17.2
	1991-1995	2.8	13.1	7.1-22.3	2.2	22.3	10.9-40.7	0.6	6.3	1.3-18.5	0.6	2.8	0.6-8.5	0.6	6.3	1.3-18.7	0.0	0.0	0.0-7.7
	1996-2000	4.2	19.8	12.3-30.4	1.4	14.3	5.7-29.8	2.8	22.7	12.3-39.1	0.8	3.7	1.0-9.8	0.2	2.5	0.1-13.0	0.6	5.6	1.2-16.9
	2001-2005	3.4	14.5	8.4-23.6	2.2	19.6	9.7-35.8	1.2	8.3	3.0-19.8	0.8	3.3	0.9-9.0	0.4	3.3	0.4-13.3	0.4	3.1	0.4-12.7
	2006-2010	3.4	13.6	7.8-22.4	1.6	13.2	5.6-27.5	1.8	13.2	6.0-26.8	0.6	2.4	0.5-7.8	0.4	3.4	0.4-14.0	0.2	1.3	0.0-10.4
	2011-2015	3.0	11.4	6.3-19.6	2.2	17.0	8.2-32.3	0.8	7.0	1.9-19.2	1.0	4.1	1.3-10.4	0.4	3.1	0.4-13.5	0.6	4.9	1.0-16.1
2016-2020	4.4	17.1	10.3-27.1	2.8	22.6	11.7-40.2	1.6	12.4	4.9-26.9	1.0	3.2	1.0-8.8	0.8	5.4	1.4-16.5	0.2	1.0	0.0-10.4	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schuyler**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	0.2	1.4	0.0-7.2	0.2	3.9	0.1-18.4	0.0	0.0	0.0-8.5	0.6	3.4	0.7-10.1	0.2	3.9	0.1-18.4	0.4	3.7	0.4-14.5
	1981-1985	0.8	4.3	1.2-11.2	0.2	1.9	0.0-12.4	0.6	6.1	1.3-18.3	0.4	2.2	0.3-8.1	0.0	0.0	0.0-9.3	0.4	4.0	0.5-15.0
	1986-1990	1.0	5.2	1.7-12.4	1.0	14.5	4.5-33.3	0.0	0.0	0.0-7.9	0.6	3.3	0.7-9.7	0.6	9.1	1.8-25.5	0.0	0.0	0.0-7.9
	1991-1995	1.4	7.0	2.8-14.4	0.8	8.5	2.3-22.0	0.6	4.9	1.0-15.4	0.6	2.7	0.6-8.3	0.4	4.3	0.5-15.7	0.2	1.5	0.0-10.1
	1996-2000	0.8	3.8	1.0-9.9	0.6	5.9	1.2-17.8	0.2	1.3	0.0-9.6	0.2	0.8	0.0-5.2	0.2	1.7	0.0-10.8	0.0	0.0	0.0-7.8
	2001-2005	1.2	4.9	1.8-11.2	0.6	6.3	1.2-19.1	0.6	4.1	0.8-14.1	1.0	4.2	1.4-10.4	0.2	1.7	0.0-11.0	0.8	5.8	1.6-16.5
	2006-2010	1.8	7.0	3.2-13.9	1.6	12.1	5.2-25.4	0.2	1.6	0.0-11.0	1.2	5.0	1.8-11.5	0.8	6.6	1.8-18.5	0.4	3.1	0.4-13.3
	2011-2015	1.2	4.7	1.7-11.0	1.0	8.3	2.6-21.0	0.2	1.1	0.0-10.2	1.0	4.0	1.3-10.2	0.6	4.9	1.0-16.1	0.4	2.8	0.3-12.7
	2016-2020	2.2	8.1	3.9-15.7	1.2	8.4	3.0-20.6	1.0	7.9	2.2-21.0	1.4	6.0	2.2-13.4	0.8	6.3	1.7-18.2	0.6	5.8	1.0-18.8
Pancreas	1976-1980	1.0	6.8	2.2-15.6	0.8	12.6	3.4-31.3	0.2	2.5	0.1-13.5	1.4	8.7	3.4-17.9	1.0	14.5	4.5-33.6	0.4	4.3	0.5-16.0
	1981-1985	1.8	9.9	4.5-19.1	1.2	14.7	5.2-32.5	0.6	6.7	1.3-19.7	1.0	5.7	1.8-13.3	0.4	6.0	0.7-20.7	0.6	6.2	1.3-18.5
	1986-1990	1.8	9.6	4.3-18.3	0.6	6.3	1.3-19.1	1.2	12.1	4.3-26.8	2.2	11.6	5.7-20.9	0.8	8.7	2.3-22.9	1.4	13.3	5.2-28.0
	1991-1995	1.4	6.8	2.7-14.2	1.0	11.2	3.5-26.3	0.4	3.6	0.4-13.7	1.6	7.8	3.4-15.6	0.8	8.7	2.3-22.4	0.8	7.2	2.0-19.0
	1996-2000	0.8	3.7	1.0-9.8	0.4	3.7	0.5-14.4	0.4	3.5	0.4-13.5	0.6	2.8	0.6-8.5	0.4	5.3	0.6-18.3	0.2	1.6	0.0-10.5
	2001-2005	2.0	8.7	4.2-16.4	1.4	13.7	5.4-29.2	0.6	5.6	1.1-17.1	1.4	6.3	2.5-13.3	0.6	7.0	1.4-20.4	0.8	7.3	2.0-19.4
	2006-2010	2.6	11.7	6.2-20.5	1.4	14.1	5.6-29.8	1.2	9.8	3.5-23.0	2.0	8.4	4.0-16.0	1.6	16.3	6.9-33.0	0.4	2.8	0.3-12.7
	2011-2015	3.0	11.8	6.5-20.1	1.4	13.0	5.0-28.1	1.6	11.4	4.9-24.3	2.6	10.4	5.5-18.6	1.6	14.2	5.9-29.5	1.0	7.5	2.4-19.4
	2016-2020	3.6	14.1	8.1-23.3	1.4	12.4	4.8-27.3	2.2	15.6	7.5-30.3	2.8	11.5	6.1-20.3	0.8	8.8	2.3-23.2	2.0	14.2	6.5-28.5
Larynx	1976-1980	1.2	7.8	2.8-17.0	1.0	13.3	4.2-31.3	0.2	2.7	0.1-14.0	0.0	0.0	0.0-4.5	0.0	0.0	0.0-9.8	0.0	0.0	0.0-8.5
	1981-1985	1.8	9.2	4.1-18.0	1.8	20.2	8.9-39.5	0.0	0.0	0.0-8.2	0.6	3.2	0.6-9.5	0.6	7.6	1.4-22.8	0.0	0.0	0.0-8.2
	1986-1990	1.6	7.9	3.4-15.7	1.2	12.2	4.4-27.1	0.4	3.4	0.4-13.5	0.8	4.1	1.1-10.7	0.4	4.1	0.5-15.7	0.4	4.2	0.5-15.6
	1991-1995	1.0	5.0	1.6-11.9	0.2	1.7	0.0-11.0	0.8	8.3	2.2-21.4	0.8	3.8	1.0-9.9	0.6	6.4	1.3-19.0	0.2	1.4	0.0-9.8
	1996-2000	0.6	2.8	0.6-8.4	0.4	3.6	0.4-14.0	0.2	2.0	0.0-11.3	0.2	0.9	0.0-5.5	0.2	1.8	0.0-11.2	0.0	0.0	0.0-7.8
	2001-2005	1.0	4.5	1.4-11.0	1.0	9.8	3.1-23.5	0.0	0.0	0.0-8.0	0.0	0.0	0.0-4.0	0.0	0.0	0.0-8.2	0.0	0.0	0.0-8.0
	2006-2010	0.8	3.2	0.9-9.0	0.8	6.6	1.8-18.6	0.0	0.0	0.0-8.5	0.0	0.0	0.0-4.2	0.0	0.0	0.0-8.7	0.0	0.0	0.0-8.5
	2011-2015	2.0	7.5	3.6-14.6	1.4	10.6	4.2-23.7	0.6	4.6	1.0-15.5	0.4	1.3	0.2-6.2	0.2	1.1	0.0-10.2	0.2	1.6	0.0-11.1
	2016-2020	1.2	4.0	1.4-10.0	1.0	7.1	2.2-19.2	0.2	1.0	0.0-10.4	0.2	0.7	0.0-5.5	0.2	1.4	0.0-11.0	0.0	0.0	0.0-9.1
Lung and bronchus	1976-1980	9.0	53.4	38.7-71.8	6.6	85.8	58.3-121.6	2.4	25.4	12.9-44.9	7.8	44.3	31.3-60.9	5.6	71.6	46.9-104.6	2.2	21.5	10.6-39.1
	1981-1985	13.8	77.1	59.7-97.9	9.0	111.1	80.1-149.9	4.8	48.8	31.0-73.4	8.8	48.8	35.2-65.9	5.6	66.4	43.5-96.9	3.2	35.1	19.7-57.6
	1986-1990	16.8	89.3	71.0-110.9	12.0	144.6	109.6-187.0	4.8	47.7	30.3-71.6	14.2	75.1	58.5-95.1	10.2	124.1	91.6-164.1	4.0	39.6	23.9-61.8
	1991-1995	16.8	82.1	65.4-101.9	10.8	120.8	90.3-158.1	6.0	54.1	36.3-78.0	13.4	65.6	50.8-83.5	8.4	93.8	67.3-127.1	5.0	42.4	27.3-63.3
	1996-2000	19.0	89.3	72.2-109.3	11.2	120.1	90.4-156.5	7.8	68.2	48.3-93.9	13.8	64.5	50.2-81.8	9.6	104.4	76.5-139.2	4.2	35.1	21.6-54.6
	2001-2005	18.2	80.2	64.5-98.8	10.2	98.1	72.8-129.6	8.0	66.3	47.2-91.1	13.8	60.6	47.1-76.9	8.4	84.0	60.2-114.2	5.4	44.5	29.2-65.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schuyler**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	18.6	75.8	61.0-93.5	10.8	102.4	76.1-135.1	7.8	60.3	42.8-83.6	13.2	55.2	42.6-70.8	7.4	70.9	49.3-99.0	5.8	45.4	30.3-66.4
	2011-2015	20.8	77.7	63.3-94.9	10.6	87.4	64.7-116.3	10.2	71.9	53.3-96.0	14.6	55.0	43.0-69.9	7.0	60.0	41.1-85.3	7.6	55.1	38.8-77.1
	2016-2020	22.4	77.2	63.1-94.0	11.6	83.5	62.7-110.1	10.8	72.1	53.5-96.4	14.6	50.8	39.6-64.8	7.6	56.7	39.6-79.7	7.0	46.0	31.6-66.2
Melanoma of the skin	1976-1980	1.2	8.7	3.1-18.6	0.4	6.0	0.7-20.8	0.8	10.6	2.7-26.8	0.8	5.1	1.4-13.1	0.6	7.3	1.5-21.7	0.2	3.1	0.1-15.2
	1981-1985	2.0	10.9	5.2-20.3	1.2	14.3	5.1-31.6	0.8	8.0	2.2-21.0	0.4	2.5	0.3-9.0	0.4	5.1	0.6-18.7	0.0	0.0	0.0-8.2
	1986-1990	0.8	3.8	1.0-10.1	0.4	4.1	0.5-15.6	0.4	3.3	0.4-13.2	0.6	3.3	0.7-9.8	0.6	6.9	1.4-20.5	0.0	0.0	0.0-7.9
	1991-1995	1.8	9.8	4.5-18.6	0.8	8.3	2.2-21.7	1.0	11.8	3.8-27.1	0.2	1.3	0.0-6.6	0.0	0.0	0.0-8.2	0.2	2.9	0.1-13.9
	1996-2000	3.8	18.1	10.9-28.4	2.4	24.4	12.4-43.2	1.4	11.8	4.7-25.3	1.4	6.4	2.6-13.5	0.8	6.9	1.9-18.6	0.6	5.2	1.0-16.1
	2001-2005	4.0	17.1	10.4-26.7	2.6	23.8	12.5-41.7	1.4	11.3	4.5-24.3	0.4	1.8	0.2-7.0	0.4	4.9	0.6-17.4	0.0	0.0	0.0-8.0
	2006-2010	4.6	20.3	12.7-30.9	2.4	22.7	11.5-40.8	2.2	19.1	9.4-35.3	0.4	1.7	0.2-7.0	0.4	4.7	0.6-16.9	0.0	0.0	0.0-8.5
	2011-2015	8.0	32.1	22.7-44.5	4.2	35.8	21.8-56.2	3.8	28.2	16.6-45.9	1.0	3.9	1.3-10.0	0.6	5.0	1.0-16.3	0.4	2.7	0.3-12.6
2016-2020	5.8	20.9	13.8-31.1	4.4	31.9	19.6-50.4	1.4	12.2	4.7-27.1	1.4	5.1	2.0-11.6	1.2	10.1	3.6-23.8	0.2	1.3	0.0-10.8	
Female breast	1976-1980							7.8	86.8	61.1-119.6							2.0	25.0	11.7-46.1
	1981-1985							10.2	117.3	86.4-155.3							3.2	32.0	17.8-53.1
	1986-1990							10.2	101.1	74.7-133.8							4.4	41.7	25.9-63.9
	1991-1995							10.6	103.3	77.0-135.8							3.8	34.5	20.6-54.6
	1996-2000							14.2	129.4	100.7-163.9							2.4	20.0	10.2-35.8
	2001-2005							15.0	123.5	96.8-156.0							2.6	20.5	10.8-36.2
	2006-2010							14.2	112.7	87.4-143.7							1.8	13.4	6.1-27.1
	2011-2015							14.8	110.0	85.7-139.9							1.8	11.4	5.2-23.9
2016-2020							19.8	153.7	123.2-190.3							2.8	25.2	12.7-45.0	
Cervix uteri	1976-1980							0.8	10.0	2.6-25.5							0.8	9.8	2.5-25.3
	1981-1985							2.4	27.8	14.0-49.2							1.2	14.6	5.1-31.9
	1986-1990							0.4	3.8	0.5-14.4							0.2	2.7	0.1-13.5
	1996-2000							0.6	5.5	1.1-16.9							0.4	3.6	0.4-13.8
	2001-2005							1.0	8.2	2.6-20.3							0.6	4.8	1.0-15.3
	2006-2010							0.8	7.8	1.9-21.3							0.2	2.9	0.1-14.7
	2011-2015							0.4	3.4	0.4-14.1							0.4	2.8	0.3-12.6
	2016-2020							0.4	4.3	0.3-17.1							0.0	0.0	0.0-9.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schuyler**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							2.0	23.8	11.1-44.1							0.0	0.0	0.0-8.5
	1981-1985							3.4	34.1	19.6-55.3							1.0	9.4	3.0-22.6
	1986-1990							3.0	28.9	16.0-48.3							0.6	6.0	1.2-18.0
	1991-1995							3.4	32.8	18.9-53.0							0.4	3.0	0.4-12.3
	1996-2000							3.2	30.5	17.3-49.8							0.2	2.0	0.0-11.3
	2001-2005							4.8	39.0	24.9-59.0							1.2	9.0	3.3-21.0
	2006-2010							4.4	34.0	21.2-52.7							0.4	2.6	0.3-12.2
	2011-2015							4.2	28.8	17.7-45.8							1.2	9.0	3.3-21.4
	2016-2020							4.4	30.4	18.4-48.6							1.0	6.6	2.1-18.0
Ovary	1976-1980							1.0	11.4	3.6-26.9							1.6	16.1	6.9-32.4
	1981-1985							1.0	9.8	3.2-23.5							0.8	8.1	2.2-21.3
	1986-1990							1.8	17.4	7.9-33.6							0.6	6.5	1.3-19.1
	1991-1995							2.6	24.4	12.8-42.3							2.2	19.2	9.5-35.1
	1996-2000							1.8	14.0	6.4-27.7							2.6	20.6	10.8-36.2
	2001-2005							2.4	21.6	11.0-38.5							1.4	12.6	4.8-27.0
	2006-2010							2.6	22.7	11.8-40.3							0.6	6.2	1.1-19.1
	2011-2015							1.2	8.8	3.1-21.3							1.2	8.1	2.9-19.9
	2016-2020							1.6	11.1	4.5-24.5							0.8	5.4	1.4-16.6
Prostate	1976-1980				5.2	74.8	48.1-110.4							2.0	34.2	16.1-62.2			
	1981-1985				5.2	74.7	48.2-109.9							1.2	17.4	6.0-37.9			
	1986-1990				8.8	112.0	80.6-151.1							4.2	61.9	37.9-94.4			
	1991-1995				13.0	143.8	110.2-184.3							3.2	39.9	22.3-65.1			
	1996-2000				15.4	157.0	123.4-197.1							2.0	23.1	10.8-42.8			
	2001-2005				18.2	169.2	135.8-208.8							1.2	13.0	4.7-28.5			
	2006-2010				17.2	145.8	115.8-181.9							2.4	29.9	15.4-52.1			
	2011-2015				12.6	92.2	70.1-120.1							1.6	15.0	6.2-30.9			
	2016-2020				17.2	117.4	92.6-148.0							2.8	23.9	12.8-41.6			
Testis	1976-1980				0.2	2.3	0.1-13.8							0.0	0.0	0.0-9.8			
	1981-1985				0.2	2.7	0.1-14.5							0.0	0.0	0.0-9.3			
	1986-1990				0.2	2.6	0.1-13.7							0.0	0.0	0.0-8.8			
	1991-1995				0.2	2.2	0.1-12.3							0.2	2.2	0.1-12.3			
	1996-2000				1.0	11.3	3.7-26.4							0.0	0.0	0.0-8.1			
	2001-2005				0.6	8.6	1.8-23.9							0.0	0.0	0.0-8.2			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schuyler**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				0.8	9.4	2.4-24.6							0.2	1.3	0.0-10.5			
	2011-2015				0.2	2.3	0.1-13.0							0.0	0.0	0.0-8.8			
	2016-2020				0.8	11.2	2.9-28.4							0.0	0.0	0.0-9.1			
Urinary bladder (incl. in situ)	1976-1980	3.2	19.5	11.1-31.7	2.4	37.0	18.5-65.1	0.8	8.9	2.4-23.1	0.4	2.3	0.3-8.6	0.2	2.2	0.1-13.5	0.2	1.9	0.0-11.8
	1981-1985	3.2	17.3	9.8-28.3	2.4	30.2	15.1-53.7	0.8	7.8	2.1-20.6	0.6	3.3	0.7-9.8	0.4	6.0	0.7-20.7	0.2	2.0	0.1-11.9
	1986-1990	4.4	23.6	14.7-35.9	2.8	34.0	18.2-57.5	1.6	16.0	6.8-32.1	0.8	4.4	1.2-11.2	0.6	9.1	1.8-25.5	0.2	1.8	0.0-11.0
	1991-1995	6.2	30.5	20.6-43.4	4.0	41.2	25.0-64.1	2.2	20.3	10.0-36.9	1.2	5.9	2.2-13.0	0.2	2.8	0.1-14.2	1.0	7.6	2.5-19.0
	1996-2000	5.8	26.7	17.9-38.5	4.2	47.9	29.3-73.7	1.6	12.5	5.3-25.8	1.2	5.6	2.1-12.5	1.2	15.4	5.5-33.2	0.0	0.0	0.0-7.8
	2001-2005	5.0	22.2	14.3-33.1	4.6	45.2	28.4-68.4	0.4	3.3	0.4-13.2	0.8	3.3	0.9-9.0	0.6	6.2	1.2-18.5	0.2	1.2	0.0-9.6
	2006-2010	5.2	21.9	14.3-32.7	4.2	41.6	25.4-64.5	1.0	7.4	2.4-19.1	1.2	5.1	1.9-11.7	0.8	7.9	2.1-21.0	0.4	2.9	0.3-12.7
	2011-2015	8.4	32.8	23.5-45.0	7.2	65.3	45.3-91.8	1.2	8.1	2.9-19.9	1.2	5.1	1.9-11.8	0.8	7.5	2.0-20.4	0.4	2.8	0.3-12.7
	2016-2020	5.6	19.2	12.7-28.7	4.6	34.1	21.4-53.1	1.0	6.4	2.0-17.7	2.4	8.4	4.3-15.7	1.8	16.0	7.2-31.8	0.6	3.2	0.7-13.1
Kidney and renal pelvis	1976-1980	1.2	7.8	2.8-17.0	0.6	8.3	1.6-24.2	0.6	7.3	1.4-21.3	0.0	0.0	0.0-4.5	0.0	0.0	0.0-9.8	0.0	0.0	0.0-8.5
	1981-1985	1.4	7.6	3.0-15.8	1.4	17.6	6.9-36.5	0.0	0.0	0.0-8.2	1.0	5.4	1.7-12.7	0.8	10.6	2.7-27.2	0.2	2.0	0.1-11.9
	1986-1990	2.6	15.4	8.1-26.3	1.0	11.9	3.8-27.9	1.6	18.8	8.0-36.8	1.0	5.7	1.8-13.3	0.8	9.7	2.6-24.9	0.2	2.1	0.1-11.9
	1991-1995	2.8	13.8	7.5-23.3	1.4	14.9	5.9-31.1	1.4	12.5	4.9-26.5	1.0	4.9	1.6-11.7	0.2	2.0	0.1-11.9	0.8	6.9	1.8-18.7
	1996-2000	2.8	12.8	7.0-21.7	0.8	10.0	2.6-25.2	2.0	16.6	7.9-31.4	1.4	6.2	2.5-13.0	0.8	9.6	2.6-24.2	0.6	4.7	0.9-15.0
	2001-2005	2.6	12.2	6.5-21.2	1.4	13.1	5.2-27.8	1.2	10.7	3.8-24.2	1.2	5.3	1.9-11.9	0.6	6.9	1.4-20.2	0.6	4.2	0.9-14.2
	2006-2010	2.8	11.8	6.3-20.5	0.8	6.9	1.7-19.4	2.0	16.9	7.8-32.6	0.6	2.4	0.5-7.9	0.4	3.9	0.4-15.3	0.2	1.6	0.0-11.0
	2011-2015	4.2	15.7	9.6-24.7	2.6	22.9	12.0-40.6	1.6	10.4	4.4-22.6	1.0	3.6	1.2-9.4	0.8	6.9	1.8-19.5	0.2	1.3	0.0-10.4
	2016-2020	4.8	18.7	11.6-29.2	2.8	21.4	11.1-38.3	2.0	16.1	7.2-31.9	1.8	6.0	2.7-12.5	0.6	4.1	0.7-15.0	1.2	7.5	2.7-19.0
Brain and other central nervous system	1976-1980	1.6	7.5	3.2-15.2	1.2	12.1	4.4-27.5	0.4	3.4	0.4-13.9	1.0	4.9	1.6-11.9	0.8	8.6	2.3-23.0	0.2	1.9	0.0-11.6
	1981-1985	1.2	7.1	2.5-15.5	0.6	8.1	1.6-23.2	0.6	6.2	1.3-18.5	0.6	3.1	0.6-9.4	0.4	5.0	0.6-18.0	0.2	1.7	0.0-10.9
	1986-1990	1.4	7.6	3.0-15.7	0.8	8.8	2.4-22.8	0.6	6.3	1.2-18.6	0.8	4.9	1.3-12.4	0.4	5.0	0.6-17.8	0.4	4.7	0.5-16.7
	1991-1995	1.2	6.1	2.2-13.3	1.0	11.0	3.5-25.8	0.2	1.4	0.0-9.8	0.2	1.0	0.0-5.7	0.0	0.0	0.0-8.2	0.2	1.4	0.0-9.8
	1996-2000	1.6	7.6	3.2-15.2	0.4	4.0	0.5-14.9	1.2	10.2	3.7-23.2	1.2	5.6	2.0-12.4	0.6	6.1	1.3-18.1	0.6	4.9	1.0-15.4
	2001-2005	1.4	6.6	2.6-13.8	1.0	11.1	3.5-26.2	0.4	3.5	0.4-14.0	1.0	4.4	1.4-10.7	0.8	8.5	2.2-22.2	0.2	1.2	0.0-9.6
	2006-2010	1.6	6.8	2.9-14.1	1.0	8.2	2.5-20.9	0.6	5.2	1.1-16.6	1.2	4.6	1.6-10.9	0.8	6.5	1.7-18.6	0.4	2.7	0.3-12.2
	2011-2015	0.4	1.6	0.2-6.7	0.2	1.6	0.0-11.2	0.2	1.6	0.0-11.1	0.2	0.6	0.0-5.2	0.2	1.3	0.0-10.5	0.0	0.0	0.0-8.8
	2016-2020	2.6	12.4	6.2-22.2	1.8	17.3	7.3-34.8	0.8	8.3	2.0-22.7	1.6	6.0	2.5-12.9	0.6	4.1	0.8-14.8	1.0	7.4	2.2-19.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schuyler**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	0.2	0.9	0.0-5.9	0.0	0.0	0.0-9.8	0.2	1.7	0.0-11.3	0.0	0.0	0.0-4.5	0.0	0.0	0.0-9.8	0.0	0.0	0.0-8.5
	1981-1985	0.2	1.4	0.0-7.1	0.2	3.8	0.1-17.9	0.0	0.0	0.0-8.2	0.2	1.4	0.0-7.1	0.2	3.8	0.1-17.9	0.0	0.0	0.0-8.2
	1986-1990	0.4	2.3	0.3-8.2	0.0	0.0	0.0-8.8	0.4	4.7	0.6-16.6	0.0	0.0	0.0-4.1	0.0	0.0	0.0-8.8	0.0	0.0	0.0-7.9
	1991-1995	0.8	4.4	1.2-11.2	0.0	0.0	0.0-8.2	0.8	8.9	2.4-22.4	0.0	0.0	0.0-3.9	0.0	0.0	0.0-8.2	0.0	0.0	0.0-7.7
	1996-2000	1.2	6.0	2.2-13.1	0.2	1.8	0.0-11.2	1.0	9.4	3.0-22.5	0.2	0.9	0.0-5.5	0.0	0.0	0.0-8.1	0.2	1.8	0.0-11.0
	2001-2005	2.0	10.6	5.0-19.6	0.4	4.6	0.5-16.9	1.6	16.3	6.9-32.6	0.0	0.0	0.0-4.0	0.0	0.0	0.0-8.2	0.0	0.0	0.0-8.0
	2006-2010	2.6	11.9	6.1-21.0	0.4	3.1	0.4-13.3	2.2	20.6	9.9-38.2	0.0	0.0	0.0-4.2	0.0	0.0	0.0-8.7	0.0	0.0	0.0-8.5
	2011-2015	2.4	11.0	5.5-20.0	1.0	7.8	2.4-20.2	1.4	14.6	5.6-31.1	0.2	0.9	0.0-5.8	0.2	2.0	0.1-12.3	0.0	0.0	0.0-8.8
2016-2020	2.4	12.4	6.1-22.6	0.4	4.5	0.3-17.8	2.0	20.6	9.4-39.4	0.2	0.8	0.0-5.8	0.0	0.0	0.0-9.1	0.2	1.7	0.0-11.7	
Hodgkin lymphoma	1976-1980	0.2	1.3	0.0-7.1	0.2	2.7	0.1-14.8	0.0	0.0	0.0-8.5	0.4	2.3	0.3-8.6	0.2	2.7	0.1-14.8	0.2	1.9	0.0-11.9
	1981-1985	0.2	0.9	0.0-5.7	0.0	0.0	0.0-9.3	0.2	1.8	0.0-11.2	0.0	0.0	0.0-4.3	0.0	0.0	0.0-9.3	0.0	0.0	0.0-8.2
	1991-1995	0.8	4.3	1.2-11.0	0.4	4.0	0.5-15.1	0.4	4.8	0.6-16.5	0.0	0.0	0.0-3.9	0.0	0.0	0.0-8.2	0.0	0.0	0.0-7.7
	1996-2000	0.6	2.8	0.6-8.4	0.6	5.3	1.1-16.5	0.0	0.0	0.0-7.8	0.2	1.1	0.0-6.0	0.2	2.2	0.1-12.4	0.0	0.0	0.0-7.8
	2001-2005	0.6	3.0	0.6-8.9	0.6	5.7	1.2-17.6	0.0	0.0	0.0-8.0	0.0	0.0	0.0-4.0	0.0	0.0	0.0-8.2	0.0	0.0	0.0-8.0
	2006-2010	0.6	3.1	0.6-9.4	0.2	1.3	0.0-10.5	0.4	4.7	0.5-17.0	0.0	0.0	0.0-4.2	0.0	0.0	0.0-8.7	0.0	0.0	0.0-8.5
	2011-2015	0.8	4.9	1.3-12.5	0.6	7.0	1.4-20.5	0.2	2.6	0.1-13.9	0.0	0.0	0.0-4.4	0.0	0.0	0.0-8.8	0.0	0.0	0.0-8.8
	2016-2020	0.4	2.9	0.3-10.0	0.2	3.5	0.1-16.8	0.2	2.7	0.1-14.3	0.0	0.0	0.0-4.5	0.0	0.0	0.0-9.1	0.0	0.0	0.0-9.1
Non-Hodgkin lymphomas	1976-1980	2.8	15.8	8.5-26.7	1.4	18.0	6.9-38.0	1.4	15.2	6.0-31.7	1.2	6.4	2.3-14.3	0.8	8.8	2.3-23.4	0.4	4.4	0.5-16.4
	1981-1985	2.6	15.0	7.8-25.9	1.6	18.5	7.8-37.0	1.0	11.6	3.6-27.4	1.2	6.1	2.2-13.7	0.6	7.5	1.3-22.6	0.6	5.7	1.2-17.4
	1986-1990	3.2	16.9	9.5-27.6	0.8	9.4	2.5-24.2	2.4	23.3	11.8-41.3	1.6	8.2	3.5-16.4	1.2	13.4	4.8-29.6	0.4	3.6	0.4-13.8
	1991-1995	3.2	16.2	9.2-26.4	1.4	16.7	6.7-34.3	1.8	15.7	7.1-30.6	2.0	10.4	5.0-19.2	0.6	7.5	1.5-21.4	1.4	12.5	5.0-26.4
	1996-2000	4.4	20.7	13.0-31.6	1.2	12.0	4.4-26.6	3.2	26.2	14.9-43.5	1.6	7.4	3.2-14.8	0.2	1.8	0.0-11.2	1.4	10.7	4.2-23.3
	2001-2005	5.6	25.4	16.8-37.0	2.4	23.5	12.0-41.6	3.2	26.1	14.8-43.3	1.8	7.8	3.6-15.2	0.6	6.1	1.2-18.3	1.2	9.1	3.3-21.1
	2006-2010	4.0	17.2	10.5-27.1	1.6	15.9	6.7-32.3	2.4	18.8	9.6-34.1	0.8	3.1	0.8-8.9	0.4	4.1	0.4-15.7	0.4	3.1	0.4-13.2
	2011-2015	5.0	18.1	11.6-27.5	2.2	19.2	9.4-35.8	2.8	18.8	10.1-33.6	2.0	8.0	3.8-15.4	1.0	10.2	3.2-24.8	1.0	7.7	2.5-19.8
2016-2020	7.0	23.4	15.9-33.7	3.2	22.6	12.2-39.5	3.8	24.9	14.7-41.1	1.2	4.3	1.5-10.4	0.4	2.7	0.3-12.9	0.8	5.3	1.4-16.3	
Myeloma	1976-1980	0.4	2.2	0.3-8.2	0.4	4.6	0.5-17.4	0.0	0.0	0.0-8.5	0.4	2.2	0.3-8.2	0.4	4.6	0.5-17.4	0.0	0.0	0.0-8.5
	1981-1985	1.0	5.9	1.9-13.8	0.4	6.3	0.7-21.3	0.6	6.2	1.3-18.5	0.4	2.5	0.3-8.8	0.0	0.0	0.0-9.3	0.4	4.2	0.5-15.5
	1986-1990	1.4	7.5	3.0-15.5	0.6	6.3	1.3-19.1	0.8	7.5	2.0-19.8	0.4	2.1	0.2-7.6	0.2	2.2	0.1-12.8	0.2	1.6	0.0-10.4
	1991-1995	0.8	3.7	1.0-9.7	0.4	3.6	0.4-14.0	0.4	3.3	0.4-12.9	0.8	4.0	1.1-10.3	0.0	0.0	0.0-8.2	0.8	6.4	1.8-17.5
	1996-2000	1.4	6.5	2.6-13.6	0.2	1.7	0.0-10.8	1.2	10.9	4.0-24.3	0.6	2.9	0.6-8.7	0.0	0.0	0.0-8.1	0.6	5.7	1.2-17.1
	2001-2005	1.6	7.1	3.1-14.3	1.2	11.0	4.0-24.7	0.4	3.3	0.4-13.2	1.0	4.7	1.5-11.3	0.8	7.9	2.1-20.7	0.2	1.5	0.0-10.4
	2006-2010	1.6	6.6	2.8-13.6	0.8	7.2	1.8-19.9	0.8	6.4	1.7-18.0	1.0	4.4	1.4-10.9	0.6	6.5	1.3-19.5	0.4	3.1	0.4-13.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Schuyler**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2011-2015	3.0	11.6	6.5-19.9	1.6	14.0	5.9-29.2	1.4	10.5	4.2-23.4	0.4	1.6	0.2-6.8	0.4	3.5	0.4-14.3	0.0	0.0	0.0-8.8
	2016-2020	1.6	6.0	2.5-12.9	0.8	6.0	1.6-17.7	0.8	6.1	1.5-18.2	1.0	3.7	1.2-9.7	0.8	6.0	1.6-17.7	0.2	1.7	0.0-11.7
Leukemias	1976-1980	2.6	14.4	7.5-24.9	1.0	13.1	3.9-31.3	1.6	15.8	6.8-31.9	0.6	3.6	0.7-10.7	0.2	4.4	0.1-19.9	0.4	4.3	0.5-15.9
	1981-1985	2.6	15.6	8.2-26.8	1.6	22.0	9.1-43.7	1.0	10.2	3.1-24.7	2.4	13.8	7.0-24.3	1.2	16.0	5.6-35.0	1.2	10.8	3.9-24.5
	1986-1990	3.4	18.9	11.0-30.4	1.4	18.6	7.3-38.1	2.0	20.6	9.8-38.3	0.8	4.5	1.2-11.4	0.4	6.9	0.8-22.5	0.4	4.3	0.5-15.6
	1991-1995	3.4	17.4	10.1-28.0	1.0	12.8	4.1-29.2	2.4	21.2	10.8-37.8	1.8	9.0	4.1-17.2	0.6	7.0	1.4-20.2	1.2	11.1	4.0-24.8
	1996-2000	3.2	14.9	8.5-24.4	2.0	19.8	9.5-36.9	1.2	8.7	3.1-20.4	2.0	9.8	4.7-18.2	1.2	12.9	4.7-28.1	0.8	5.7	1.5-16.3
	2001-2005	2.8	12.2	6.7-20.9	1.8	17.0	7.7-33.1	1.0	7.2	2.3-18.4	2.0	8.6	4.1-16.3	1.2	10.5	3.8-23.9	0.8	6.5	1.7-17.8
	2006-2010	4.2	19.7	12.0-30.7	2.4	27.0	13.7-47.8	1.8	16.2	7.2-32.1	1.8	8.3	3.8-16.2	1.6	17.8	7.6-35.4	0.2	1.6	0.0-11.0
	2011-2015	4.2	18.3	10.9-28.9	2.6	21.8	11.1-39.3	1.6	15.3	6.1-32.1	2.2	9.2	4.5-17.1	1.2	10.9	3.9-25.1	1.0	6.1	2.0-17.0
2016-2020	4.6	17.4	10.7-27.3	1.8	14.5	6.2-29.8	2.8	20.5	10.7-36.9	2.0	6.8	3.2-13.6	1.2	10.2	3.6-23.9	0.8	5.1	1.4-16.0	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Seneca**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	121.4	354.8	326.7-384.8	60.2	411.1	363.6-462.8	61.2	327.2	290.7-367.0	63.6	186.1	165.9-208.0	35.6	252.7	215.0-294.8	28.0	142.9	119.8-169.2
	1981-1985	141.6	411.6	381.2-443.8	69.8	468.2	418.4-522.3	71.8	384.7	344.8-428.0	68.4	198.0	177.2-220.5	32.8	223.2	189.2-261.5	35.6	182.9	156.4-212.7
	1986-1990	158.0	446.2	415.1-479.0	75.2	497.0	446.5-551.6	82.8	421.4	380.6-465.4	73.4	207.7	186.7-230.4	37.2	249.0	213.5-288.5	36.2	181.5	155.4-210.9
	1991-1995	174.0	483.4	451.5-517.0	90.0	564.4	512.3-620.4	84.0	433.8	392.5-478.3	78.2	216.2	195.2-239.0	42.4	278.4	241.1-319.6	35.8	172.3	147.6-200.2
	1996-2000	189.8	509.8	477.8-543.5	103.4	606.1	554.4-661.3	86.4	436.5	395.8-480.4	83.0	220.3	199.6-242.7	41.0	247.7	214.5-284.5	42.0	201.3	174.7-231.1
	2001-2005	212.2	520.2	489.2-552.7	112.4	595.7	547.2-647.4	99.8	472.0	430.7-516.4	75.0	179.6	161.9-198.9	38.6	211.4	182.4-243.7	36.4	158.0	135.6-183.5
	2006-2010	220.6	512.3	482.1-544.0	115.8	562.3	516.9-610.8	104.8	476.2	434.9-520.5	85.6	195.0	176.8-214.6	45.4	231.2	201.8-263.8	40.2	166.7	144.0-192.4
	2011-2015	221.6	487.6	458.4-518.4	116.4	528.6	485.6-574.7	105.2	466.6	425.4-510.9	75.6	163.6	147.1-181.6	39.6	184.1	158.9-212.4	36.0	148.2	126.3-173.1
	2016-2020	238.4	510.9	480.8-542.4	128.4	554.4	510.9-600.9	110.0	484.9	442.1-531.0	77.8	158.3	142.5-175.5	41.4	180.8	156.4-208.2	36.4	143.3	122.5-167.2
Oral cavity and pharynx	1976-1980	1.2	3.9	1.4-8.5	0.6	4.6	0.9-13.1	0.6	3.1	0.6-9.5	0.8	2.2	0.6-5.8	0.4	2.9	0.3-10.5	0.4	1.8	0.2-7.2
	1981-1985	4.4	11.3	7.0-17.4	3.4	18.9	10.9-30.9	1.0	4.7	1.5-11.4	1.4	3.7	1.4-7.8	1.0	5.7	1.8-13.9	0.4	1.9	0.2-7.5
	1986-1990	3.4	10.1	5.8-16.3	2.6	16.8	8.9-28.9	0.8	3.9	1.0-10.5	1.4	3.9	1.6-8.2	1.2	7.2	2.6-15.9	0.2	0.8	0.0-5.6
	1991-1995	3.0	8.6	4.7-14.2	2.0	11.9	5.6-22.1	1.0	5.2	1.6-12.6	1.2	3.9	1.4-8.5	1.0	7.1	2.3-16.3	0.2	1.0	0.0-5.9
	1996-2000	4.6	12.6	7.9-18.9	2.8	16.5	9.0-27.8	1.8	9.1	4.1-17.8	1.2	3.2	1.2-7.0	0.6	3.3	0.7-10.1	0.6	2.7	0.5-8.6
	2001-2005	4.2	10.3	6.4-15.9	3.0	16.0	8.9-26.6	1.2	5.0	1.8-11.6	0.6	1.4	0.3-4.3	0.2	0.9	0.0-5.8	0.4	1.7	0.2-6.9
	2006-2010	5.2	11.5	7.4-17.1	3.8	16.9	10.1-26.9	1.4	6.2	2.4-13.7	1.2	2.7	1.0-6.2	0.4	2.0	0.2-7.5	0.8	2.9	0.8-8.6
	2011-2015	3.4	6.9	4.0-11.4	2.6	11.2	5.9-19.7	0.8	3.4	0.9-9.6	0.6	1.2	0.2-3.9	0.6	2.5	0.5-8.0	0.0	0.0	0.0-4.6
	2016-2020	8.0	18.0	12.6-25.0	5.6	24.5	15.8-36.4	2.4	11.1	5.5-20.3	1.4	2.6	1.0-5.9	1.4	5.6	2.2-12.3	0.0	0.0	0.0-4.9
Esophagus	1976-1980	0.4	1.3	0.2-4.7	0.2	1.3	0.0-7.6	0.2	1.2	0.0-6.6	0.6	2.0	0.4-5.8	0.4	3.0	0.4-10.6	0.2	1.2	0.0-6.6
	1981-1985	1.2	3.8	1.4-8.4	0.6	4.5	0.9-13.1	0.6	3.5	0.7-10.4	1.4	4.2	1.7-8.7	0.8	5.0	1.3-13.4	0.6	3.5	0.7-10.4
	1986-1990	1.2	3.6	1.3-7.8	1.0	6.0	1.9-14.4	0.2	1.0	0.0-6.0	1.2	3.5	1.3-7.7	1.0	6.5	2.0-15.4	0.2	0.8	0.0-5.5
	1991-1995	1.0	2.7	0.9-6.4	0.6	4.3	0.9-12.3	0.4	1.8	0.2-7.2	1.0	2.7	0.9-6.5	0.8	5.1	1.4-13.1	0.2	0.9	0.0-5.8
	1996-2000	1.8	4.8	2.2-9.2	1.2	7.1	2.6-15.6	0.6	2.8	0.6-8.7	1.6	4.2	1.8-8.3	1.0	6.6	2.1-15.3	0.6	2.7	0.5-8.5
	2001-2005	2.6	6.6	3.5-11.3	2.2	11.5	5.7-20.8	0.4	2.0	0.2-7.6	2.0	5.0	2.4-9.4	1.4	7.8	3.1-16.1	0.6	2.8	0.6-8.7
	2006-2010	1.6	4.0	1.7-8.0	1.4	7.0	2.8-14.6	0.2	1.1	0.0-6.3	2.0	4.9	2.3-9.2	1.6	8.2	3.5-16.3	0.4	2.0	0.2-7.8
	2011-2015	2.2	5.1	2.5-9.5	1.6	8.0	3.3-16.3	0.6	2.3	0.5-7.9	2.8	5.8	3.1-10.0	1.8	7.8	3.5-15.5	1.0	3.6	1.2-9.6
	2016-2020	3.8	7.4	4.4-12.0	3.6	14.9	8.7-24.2	0.2	0.9	0.0-6.1	3.6	7.0	4.1-11.4	3.4	13.9	8.0-23.0	0.2	0.9	0.0-6.1
Stomach	1976-1980	2.6	7.7	4.0-13.3	1.2	7.2	2.5-16.3	1.4	7.6	2.9-16.0	2.0	5.8	2.7-10.7	0.8	5.4	1.4-14.1	1.2	5.9	2.1-13.3
	1981-1985	4.2	12.6	7.7-19.4	2.6	19.9	10.3-34.1	1.6	7.2	3.1-14.7	2.2	6.3	3.1-11.4	1.2	8.4	3.0-18.5	1.0	4.5	1.4-11.0
	1986-1990	4.0	11.3	6.9-17.6	3.0	21.5	11.7-35.7	1.0	4.3	1.4-10.7	2.6	7.4	3.9-12.7	1.6	11.8	5.0-23.2	1.0	4.5	1.5-11.1
	1991-1995	1.8	4.9	2.2-9.5	1.2	7.8	2.7-17.3	0.6	2.3	0.5-7.7	1.0	2.8	0.9-6.7	0.4	2.3	0.3-8.8	0.6	2.3	0.5-7.7
	1996-2000	3.0	7.9	4.4-13.1	2.2	13.6	6.7-24.6	0.8	4.2	1.1-11.0	1.4	3.7	1.5-7.7	1.0	6.2	1.9-14.6	0.4	2.1	0.2-8.0
	2001-2005	1.6	3.8	1.6-7.7	0.6	3.2	0.7-9.6	1.0	4.2	1.3-10.5	1.2	2.8	1.0-6.3	0.6	3.3	0.7-9.7	0.6	2.1	0.4-7.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Seneca**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	3.4	7.8	4.5-12.7	1.6	7.7	3.3-15.6	1.8	7.1	3.0-14.5	1.8	4.4	2.0-8.5	0.6	2.9	0.6-8.9	1.2	5.4	1.8-12.7
	2011-2015	4.6	9.1	5.7-14.1	3.4	15.2	8.7-24.9	1.2	4.4	1.6-10.8	1.4	2.6	1.1-5.9	1.0	4.8	1.5-11.6	0.4	1.2	0.1-6.1
	2016-2020	3.4	7.3	4.1-12.2	2.6	10.0	5.3-18.0	0.8	5.4	1.4-13.8	1.0	1.8	0.6-4.7	0.8	3.0	0.8-8.8	0.2	0.6	0.0-5.6
Colorectal	1976-1980	19.6	56.3	45.5-68.9	9.0	63.9	45.8-86.4	10.6	52.8	39.4-69.6	10.8	32.9	24.6-43.1	5.4	44.7	28.8-65.7	5.4	27.1	17.8-39.7
	1981-1985	27.4	76.7	64.2-90.9	14.2	97.0	74.8-123.5	13.2	64.8	50.0-82.9	10.4	29.1	21.7-38.3	3.8	26.4	15.5-41.8	6.6	31.3	21.5-44.3
	1986-1990	25.8	71.4	59.4-85.0	12.4	78.9	59.9-101.9	13.4	62.6	48.3-80.2	10.6	29.2	21.8-38.4	5.6	34.2	22.4-50.0	5.0	24.6	15.7-36.9
	1991-1995	22.8	62.8	51.7-75.6	11.6	78.1	58.8-101.6	11.2	53.7	40.4-70.2	11.0	30.4	22.8-39.7	5.8	39.6	26.1-57.3	5.2	24.5	15.9-36.4
	1996-2000	23.4	62.2	51.5-74.7	11.8	70.4	53.4-91.1	11.6	55.4	42.0-72.2	10.4	27.5	20.5-36.1	3.8	23.0	13.7-36.2	6.6	30.1	20.6-42.8
	2001-2005	22.4	53.9	44.3-64.9	12.0	63.8	48.6-82.3	10.4	45.6	33.9-60.4	7.2	17.2	12.0-23.9	3.2	18.2	10.3-29.6	4.0	16.7	10.0-26.5
	2006-2010	20.2	45.8	37.2-55.9	11.6	55.6	42.1-72.3	8.6	37.6	26.9-51.5	5.8	13.0	8.7-18.9	3.8	19.7	11.8-30.9	2.0	7.9	3.7-15.3
	2011-2015	18.8	40.7	32.7-50.3	10.0	46.4	34.2-61.6	8.8	38.1	27.1-52.5	6.4	13.8	9.3-20.0	3.4	14.8	8.5-24.3	3.0	13.5	7.0-23.7
	2016-2020	17.2	37.7	29.7-47.2	10.0	43.4	31.8-58.2	7.2	33.4	22.7-47.7	5.2	11.1	7.1-16.7	3.0	13.6	7.4-23.1	2.2	8.9	4.3-16.9
Colon excluding rectum	1976-1980	13.2	38.8	29.8-49.5	5.8	44.2	29.1-64.0	7.4	36.4	25.4-50.6	8.2	25.0	17.8-34.0	4.4	35.3	21.6-53.9	3.8	18.7	11.2-29.5
	1981-1985	19.8	56.2	45.5-68.6	9.8	70.1	51.1-93.7	10.0	49.2	36.4-65.2	9.2	25.8	18.8-34.5	3.4	23.9	13.5-38.7	5.8	27.4	18.3-39.7
	1986-1990	18.6	51.0	41.1-62.6	9.0	58.2	41.9-78.5	9.6	43.9	32.2-58.7	8.6	23.3	16.8-31.5	4.6	26.0	16.3-39.4	4.0	20.5	12.3-32.2
	1991-1995	18.2	50.2	40.3-61.8	9.0	62.6	45.2-84.3	9.2	44.1	32.1-59.3	10.2	28.1	20.9-37.1	5.2	36.0	23.1-53.2	5.0	23.8	15.3-35.5
	1996-2000	17.4	46.3	37.0-57.2	7.4	44.2	31.0-61.2	10.0	47.5	35.1-63.1	8.2	21.5	15.4-29.3	2.8	16.8	9.1-28.3	5.4	24.4	16.0-36.0
	2001-2005	16.0	37.8	29.9-47.2	8.2	44.7	32.0-60.8	7.8	32.9	23.3-45.6	5.6	13.3	8.8-19.4	2.4	13.9	7.1-24.3	3.2	13.4	7.5-22.5
	2006-2010	14.8	34.0	26.6-42.9	7.6	37.1	26.2-51.3	7.2	31.5	21.7-44.5	4.0	8.8	5.3-13.8	2.4	12.4	6.4-21.9	1.6	6.2	2.6-13.2
	2011-2015	13.4	29.0	22.3-37.1	7.2	34.5	24.1-48.2	6.2	25.7	17.2-37.4	5.2	11.2	7.2-16.8	2.8	12.4	6.7-21.4	2.4	10.1	4.9-19.0
	2016-2020	11.8	25.0	18.7-32.8	6.6	28.3	19.2-40.7	5.2	22.7	14.3-34.5	3.8	7.9	4.7-12.8	2.2	9.2	4.5-17.2	1.6	6.9	2.9-14.5
Rectum & rectosigmoid	1976-1980	6.4	17.5	11.9-25.0	3.2	19.6	10.8-32.8	3.2	16.4	9.3-27.1	2.6	7.9	4.2-13.6	1.0	9.5	2.8-21.7	1.6	8.4	3.6-16.8
	1981-1985	7.6	20.5	14.5-28.4	4.4	26.9	16.5-41.4	3.2	15.7	8.9-25.9	1.2	3.3	1.2-7.4	0.4	2.6	0.3-9.6	0.8	3.9	1.0-10.4
	1986-1990	7.2	20.4	14.2-28.4	3.4	20.7	11.9-33.6	3.8	18.8	11.0-30.0	2.0	5.9	2.8-10.9	1.0	8.3	2.6-18.7	1.0	4.1	1.3-10.3
	1991-1995	4.6	12.6	7.9-19.0	2.6	15.5	8.2-26.8	2.0	9.6	4.6-18.1	0.8	2.3	0.6-5.9	0.6	3.6	0.7-10.8	0.2	0.8	0.0-5.5
	1996-2000	6.0	16.0	10.8-22.9	4.4	26.2	16.3-39.8	1.6	8.0	3.4-16.2	2.2	6.0	3.0-10.8	1.0	6.3	2.0-14.8	1.2	5.7	2.0-13.0
	2001-2005	6.4	16.1	11.0-22.8	3.8	19.1	11.5-30.1	2.6	12.7	6.7-22.2	1.6	3.8	1.6-7.7	0.8	4.3	1.2-11.1	0.8	3.3	0.9-9.2
	2006-2010	5.4	11.8	7.8-17.5	4.0	18.5	11.3-28.9	1.4	6.1	2.4-13.3	1.8	4.2	1.9-8.2	1.4	7.3	2.9-15.2	0.4	1.6	0.2-7.0
	2011-2015	5.4	11.8	7.6-17.6	2.8	11.8	6.4-20.4	2.6	12.5	6.1-22.7	1.2	2.7	0.9-6.4	0.6	2.4	0.5-7.8	0.6	3.4	0.5-10.7
	2016-2020	5.4	12.7	8.1-19.0	3.4	15.1	8.5-25.0	2.0	10.7	4.7-20.7	1.4	3.2	1.1-7.0	0.8	4.3	1.1-11.6	0.6	2.0	0.4-7.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Seneca**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	1.2	3.3	1.2-7.3	1.2	9.4	3.0-21.0	0.0	0.0	0.0-4.4	1.4	4.1	1.6-8.5	1.4	10.2	3.8-21.3	0.0	0.0	0.0-4.4
	1981-1985	0.8	2.5	0.6-6.6	0.0	0.0	0.0-5.1	0.8	5.0	1.2-12.9	0.8	2.4	0.6-6.4	0.0	0.0	0.0-5.1	0.8	4.4	1.0-11.7
	1986-1990	0.6	1.6	0.3-4.9	0.2	1.4	0.0-7.6	0.4	1.8	0.2-7.2	0.4	1.4	0.2-4.9	0.4	3.0	0.4-10.5	0.0	0.0	0.0-4.3
	1991-1995	2.4	6.6	3.4-11.7	1.8	10.8	4.9-20.8	0.6	3.2	0.6-9.6	2.0	5.5	2.6-10.2	1.8	11.4	5.2-21.8	0.2	0.9	0.0-5.9
	1996-2000	1.2	3.2	1.2-7.1	0.8	4.8	1.3-12.5	0.4	2.0	0.2-7.8	1.2	3.2	1.2-7.0	0.6	4.3	0.9-12.3	0.6	2.4	0.5-7.9
	2001-2005	3.0	7.2	4.0-12.1	2.2	12.2	6.1-21.9	0.8	3.4	0.9-9.5	1.8	4.2	1.9-8.2	1.4	8.1	3.2-16.7	0.4	1.5	0.2-6.5
	2006-2010	2.0	4.7	2.2-8.8	1.2	6.2	2.2-13.7	0.8	3.7	1.0-10.0	1.8	4.3	2.0-8.4	1.0	5.0	1.6-11.9	0.8	3.7	1.0-10.0
	2011-2015	3.6	8.2	4.8-13.3	2.2	9.4	4.6-17.4	1.4	7.5	2.8-16.1	1.4	3.7	1.4-7.9	1.0	4.7	1.5-11.4	0.4	3.2	0.4-10.9
	2016-2020	3.0	6.0	3.2-10.4	1.6	6.7	2.8-13.9	1.4	6.0	2.0-14.0	3.8	6.9	4.1-11.2	2.6	10.5	5.5-18.7	1.2	4.1	1.5-10.3
Pancreas	1976-1980	4.0	11.3	6.9-17.6	2.8	19.1	9.9-32.9	1.2	5.8	2.1-13.1	3.8	10.9	6.5-17.1	2.2	14.5	6.9-26.6	1.6	7.5	3.2-15.3
	1981-1985	2.4	6.6	3.4-11.7	0.6	3.2	0.7-10.2	1.8	9.2	4.1-17.8	2.4	7.0	3.6-12.3	0.6	4.4	0.9-12.9	1.8	9.6	4.3-18.4
	1986-1990	4.8	13.6	8.7-20.4	2.4	15.6	8.0-27.5	2.4	11.2	5.7-20.2	4.2	11.4	7.0-17.5	2.4	15.6	7.9-27.4	1.8	7.7	3.5-15.3
	1991-1995	5.6	15.7	10.4-22.8	4.2	27.5	16.9-42.2	1.4	6.8	2.7-14.4	4.4	12.5	7.8-19.0	3.0	19.8	10.9-32.8	1.4	7.1	2.8-15.0
	1996-2000	4.4	11.6	7.3-17.7	1.8	10.4	4.7-20.0	2.6	12.2	6.4-21.3	4.4	11.4	7.2-17.4	2.4	14.9	7.6-26.1	2.0	9.0	4.3-17.0
	2001-2005	5.2	12.1	7.9-17.9	2.6	14.0	7.4-24.1	2.6	10.1	5.3-18.0	4.6	10.8	6.9-16.4	2.2	11.6	5.8-20.9	2.4	9.5	4.8-17.3
	2006-2010	6.0	14.2	9.5-20.4	2.4	12.5	6.4-22.1	3.6	15.5	9.0-25.3	5.4	11.7	7.7-17.3	2.2	10.9	5.4-19.7	3.2	11.9	6.7-20.1
	2011-2015	6.4	14.1	9.5-20.2	4.0	19.7	11.9-30.8	2.4	8.6	4.4-16.0	4.6	10.4	6.5-15.9	2.6	12.7	6.7-22.0	2.0	8.5	3.9-16.7
	2016-2020	8.2	16.1	11.5-22.3	4.8	20.6	13.0-31.4	3.4	11.8	6.8-20.0	5.6	10.8	7.1-16.1	2.6	11.4	5.9-20.1	3.0	10.6	5.9-18.5
Larynx	1976-1980	2.2	5.8	2.9-10.6	1.8	10.5	4.7-20.5	0.4	2.0	0.2-7.7	1.0	2.7	0.9-6.5	1.0	6.4	2.0-15.4	0.0	0.0	0.0-4.4
	1981-1985	1.6	4.6	1.9-9.3	1.4	9.5	3.7-19.8	0.2	0.9	0.0-5.8	0.8	2.6	0.7-6.6	0.8	5.6	1.5-14.3	0.0	0.0	0.0-4.4
	1986-1990	3.0	8.2	4.5-13.8	3.0	17.6	9.7-29.4	0.0	0.0	0.0-4.3	0.8	2.2	0.6-5.7	0.8	4.5	1.2-12.1	0.0	0.0	0.0-4.3
	1991-1995	1.6	4.5	1.9-8.9	1.6	9.4	4.0-18.9	0.0	0.0	0.0-4.3	0.8	2.1	0.6-5.6	0.6	3.6	0.7-10.8	0.2	0.9	0.0-5.9
	1996-2000	1.8	4.9	2.2-9.4	1.2	7.1	2.6-15.5	0.6	2.7	0.6-8.6	0.6	1.6	0.3-4.8	0.6	3.4	0.7-10.3	0.0	0.0	0.0-4.4
	2001-2005	1.4	3.4	1.4-7.1	0.8	4.0	1.1-10.6	0.6	3.0	0.6-9.2	0.0	0.0	0.0-2.1	0.0	0.0	0.0-4.2	0.0	0.0	0.0-4.3
	2006-2010	1.8	4.1	1.8-8.0	1.2	5.4	2.0-12.3	0.6	2.7	0.6-8.6	0.0	0.0	0.0-2.1	0.0	0.0	0.0-4.1	0.0	0.0	0.0-4.5
	2011-2015	1.6	3.0	1.3-6.3	1.4	6.0	2.4-13.0	0.2	0.7	0.0-5.7	0.4	0.8	0.1-3.5	0.4	2.0	0.2-7.5	0.0	0.0	0.0-4.6
	2016-2020	1.4	3.1	1.2-6.9	1.0	4.9	1.6-12.0	0.4	1.5	0.2-6.9	0.8	1.5	0.4-4.4	0.4	1.4	0.2-6.3	0.4	1.6	0.2-7.2
Lung and bronchus	1976-1980	18.0	52.4	42.0-64.7	14.6	95.5	74.0-121.2	3.4	19.2	10.9-31.1	15.2	43.5	34.1-54.6	12.4	80.8	61.1-104.7	2.8	15.2	8.2-25.9
	1981-1985	21.8	62.6	51.1-75.9	13.0	82.9	63.4-106.5	8.8	46.4	33.3-62.9	18.8	54.1	43.5-66.5	11.8	75.9	57.3-98.6	7.0	36.6	25.2-51.6
	1986-1990	21.2	59.3	48.3-72.0	11.6	76.6	57.6-99.7	9.6	48.4	35.3-64.8	17.8	50.4	40.3-62.3	10.0	65.2	47.9-86.5	7.8	39.7	27.9-54.8
	1991-1995	25.8	69.9	58.2-83.2	14.8	91.3	71.3-115.1	11.0	52.9	39.7-69.3	19.4	53.0	42.9-64.8	11.6	74.4	56.1-96.7	7.8	37.2	26.3-51.4
	1996-2000	33.4	89.1	76.1-103.8	19.8	114.5	92.9-139.7	13.6	68.2	52.8-86.9	27.2	72.3	60.6-85.7	16.0	93.3	73.9-116.5	11.2	55.3	41.6-72.2
	2001-2005	33.0	79.8	68.1-93.1	18.8	98.5	79.5-120.7	14.2	65.6	51.1-83.2	25.2	60.2	50.2-71.8	15.0	78.7	61.8-98.8	10.2	44.9	33.3-59.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Seneca**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	35.0	80.0	68.5-93.1	19.0	92.3	74.5-113.2	16.0	69.7	55.0-87.4	22.2	50.9	41.8-61.5	13.0	64.4	49.5-82.4	9.2	39.8	29.0-53.7
	2011-2015	38.4	81.6	70.2-94.4	19.8	89.7	72.6-109.8	18.6	76.9	61.6-95.2	22.2	47.4	38.8-57.5	12.2	55.6	42.3-72.1	10.0	41.9	30.8-56.2
	2016-2020	35.2	70.8	60.4-82.6	17.2	72.4	57.6-90.2	18.0	71.2	56.7-88.8	22.2	44.2	36.2-53.7	10.4	43.9	32.5-58.3	11.8	45.1	34.1-59.2
Melanoma of the skin	1976-1980	2.6	7.5	3.9-13.0	1.6	8.9	3.7-18.1	1.0	6.3	1.9-14.7	0.4	1.3	0.1-4.7	0.2	1.0	0.0-6.8	0.2	1.6	0.0-7.9
	1981-1985	3.0	9.2	5.0-15.4	1.8	10.6	4.7-20.7	1.2	8.0	2.9-17.3	0.4	1.0	0.1-4.0	0.2	1.2	0.0-7.1	0.2	0.9	0.0-5.8
	1986-1990	2.4	7.7	3.9-13.5	1.6	10.8	4.6-21.4	0.8	4.6	1.2-11.8	1.0	3.0	0.9-7.0	1.0	6.4	2.0-15.1	0.0	0.0	0.0-4.3
	1991-1995	2.8	8.3	4.5-14.0	1.4	9.2	3.5-19.2	1.4	8.5	3.4-17.6	1.4	3.9	1.6-8.2	1.0	5.9	1.9-14.2	0.4	1.8	0.2-7.1
	1996-2000	4.8	13.4	8.6-20.1	2.8	16.3	8.9-27.5	2.0	11.0	5.2-20.4	1.2	3.3	1.2-7.3	0.8	4.8	1.3-12.4	0.4	2.1	0.2-7.9
	2001-2005	6.4	16.6	11.3-23.5	3.8	20.1	12.1-31.6	2.6	14.5	7.6-25.2	1.6	4.0	1.7-8.0	1.2	6.6	2.4-14.4	0.4	2.1	0.3-7.9
	2006-2010	5.2	12.9	8.4-19.1	3.0	15.4	8.6-25.6	2.2	11.5	5.5-21.1	1.2	2.8	1.0-6.3	0.8	3.8	1.0-10.0	0.4	1.7	0.2-7.0
	2011-2015	11.0	26.9	20.1-35.4	4.8	22.8	14.5-34.4	6.2	32.8	21.8-47.6	0.2	0.4	0.0-2.7	0.0	0.0	0.0-4.2	0.2	0.7	0.0-5.6
	2016-2020	12.6	28.4	21.4-37.1	8.0	35.9	25.1-49.9	4.6	22.2	13.4-34.7	1.2	2.8	1.0-6.6	0.2	0.7	0.0-5.4	1.0	4.9	1.4-12.6
Female breast	1976-1980							16.4	91.5	72.3-114.3							4.0	21.4	12.9-33.5
	1981-1985							17.4	100.4	79.7-124.8							6.8	37.2	25.3-52.6
	1986-1990							27.6	147.3	123.0-175.0							7.6	39.8	27.8-55.2
	1991-1995							24.4	130.1	107.6-155.9							4.8	23.9	15.2-35.9
	1996-2000							23.8	122.4	101.2-146.9							5.8	27.8	18.5-40.4
	2001-2005							30.4	145.5	123.0-171.2							4.8	21.3	13.6-32.3
	2006-2010							31.0	144.3	121.8-170.0							5.8	24.7	16.3-36.2
	2011-2015							26.6	124.1	102.7-148.7							5.6	23.7	15.3-35.4
	2016-2020							30.8	140.1	117.0-166.6							3.4	15.1	8.4-25.3
Cervix uteri	1976-1980							2.8	16.3	8.8-27.5							1.4	7.3	2.9-15.3
	1981-1985							2.0	13.9	6.5-25.6							0.8	4.1	1.1-10.8
	1986-1990							2.2	13.3	6.5-23.9							0.8	5.1	1.4-12.9
	1991-1995							3.2	19.2	10.9-31.3							0.6	3.3	0.6-9.9
	1996-2000							1.2	6.9	2.5-15.2							0.4	2.2	0.3-8.1
	2001-2005							1.4	8.0	3.2-16.6							0.2	1.0	0.0-6.1
	2006-2010							1.4	7.8	3.1-16.4							0.6	2.8	0.6-8.8
	2011-2015							1.4	7.4	2.7-16.0							0.4	1.6	0.2-7.0
	2016-2020							1.8	10.5	4.5-20.6							0.4	1.9	0.2-8.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Seneca**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							6.2	32.2	21.6-46.3							1.6	7.5	3.2-15.2
	1981-1985							6.0	30.9	20.7-44.7							1.2	5.6	2.0-12.7
	1986-1990							4.4	22.8	14.1-35.1							1.0	4.6	1.5-11.2
	1991-1995							5.8	30.9	20.6-44.8							1.0	5.3	1.7-12.6
	1996-2000							4.6	24.2	15.3-36.6							1.6	8.5	3.6-17.1
	2001-2005							8.6	40.6	29.3-55.3							2.2	9.4	4.7-17.5
	2006-2010							6.2	27.8	18.7-40.2							2.6	10.3	5.4-18.5
	2011-2015							8.2	34.6	24.5-47.9							1.0	3.6	1.1-9.6
	2016-2020							10.2	43.8	31.9-59.1							1.0	4.4	1.3-11.5
Ovary	1976-1980							2.6	14.5	7.5-25.2							1.8	9.9	4.5-19.2
	1981-1985							3.2	18.2	10.1-30.1							1.0	6.2	2.0-14.5
	1986-1990							3.6	19.2	11.2-30.8							2.0	10.6	5.0-19.9
	1991-1995							2.4	12.3	6.3-21.9							2.8	13.3	7.2-22.7
	1996-2000							3.0	15.9	8.9-26.6							1.8	8.5	3.9-16.7
	2001-2005							3.6	17.8	10.5-28.6							3.0	14.1	7.8-23.7
	2006-2010							3.2	13.7	7.6-23.2							1.6	6.0	2.5-12.8
	2011-2015							2.2	9.4	4.6-17.6							1.8	8.5	3.8-16.9
	2016-2020							3.6	15.0	8.6-24.8							2.4	8.9	4.5-16.7
Prostate	1976-1980				10.0	75.3	54.9-100.4							2.6	23.4	12.0-40.1			
	1981-1985				10.2	69.5	51.1-92.1							2.6	19.0	9.9-32.8			
	1986-1990				15.8	105.2	82.6-131.9							4.6	33.5	21.0-50.5			
	1991-1995				26.2	158.4	131.8-188.8							4.6	32.5	20.2-49.2			
	1996-2000				33.2	190.9	162.7-222.7							5.6	35.0	23.2-50.7			
	2001-2005				33.0	174.2	148.5-203.2							3.8	23.0	13.8-35.8			
	2006-2010				30.2	142.4	120.3-167.5							4.2	23.8	14.7-36.4			
	2011-2015				26.4	110.9	92.4-132.3							3.0	14.8	8.2-24.7			
	2016-2020				31.8	125.1	106.0-147.0							2.6	11.7	6.2-20.6			
Testis	1976-1980				0.4	2.5	0.3-9.6							0.0	0.0	0.0-5.1			
	1981-1985				0.2	0.9	0.0-6.6							0.0	0.0	0.0-5.1			
	1986-1990				1.0	5.7	1.8-13.8							0.0	0.0	0.0-4.8			
	1991-1995				1.6	9.0	3.9-18.0							0.0	0.0	0.0-4.8			
	1996-2000				0.6	3.6	0.7-10.6							0.0	0.0	0.0-4.6			
	2001-2005				1.0	5.6	1.8-13.1							0.0	0.0	0.0-4.2			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Seneca**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				1.0	5.1	1.6-12.1							0.2	0.9	0.0-5.6			
	2011-2015				0.8	3.8	1.0-10.3							0.0	0.0	0.0-4.2			
	2016-2020				2.0	11.2	5.2-20.9							0.0	0.0	0.0-4.3			
Urinary bladder (incl. in situ)	1976-1980	6.0	17.4	11.7-25.0	4.0	28.9	17.1-45.4	2.0	9.8	4.7-18.4	1.6	4.9	2.1-9.7	1.0	9.1	2.7-21.0	0.6	2.8	0.6-8.8
	1981-1985	5.8	17.0	11.3-24.6	4.4	31.7	19.4-48.6	1.4	7.4	2.9-15.5	1.0	3.2	1.0-7.3	1.0	9.6	3.1-21.5	0.0	0.0	0.0-4.4
	1986-1990	8.2	21.5	15.4-29.4	6.2	38.8	25.9-55.7	2.0	9.6	4.5-18.2	2.2	6.1	3.0-10.9	1.2	8.8	3.2-19.1	1.0	4.5	1.4-11.0
	1991-1995	7.4	20.1	14.1-27.9	4.6	30.3	19.0-45.8	2.8	13.3	7.2-22.8	3.0	8.1	4.5-13.5	2.4	16.5	8.3-29.0	0.6	2.6	0.5-8.3
	1996-2000	10.8	28.2	21.2-36.9	8.4	50.0	35.8-67.9	2.4	10.9	5.6-19.5	1.6	4.2	1.8-8.5	1.0	7.1	2.2-16.4	0.6	3.0	0.6-9.2
	2001-2005	12.6	30.4	23.3-39.0	10.4	55.4	41.3-72.9	2.2	10.1	4.9-18.7	1.8	4.2	1.9-8.1	1.4	8.0	3.2-16.4	0.4	1.3	0.2-6.2
	2006-2010	13.2	30.5	23.6-39.1	10.2	51.4	38.1-67.9	3.0	12.9	7.1-22.1	4.0	8.8	5.3-13.8	3.4	17.4	10.0-28.1	0.6	2.1	0.4-7.5
	2011-2015	11.8	25.1	19.0-32.8	7.2	33.4	23.2-46.9	4.6	17.3	10.9-26.9	0.8	1.7	0.4-4.6	0.4	2.0	0.2-7.6	0.4	1.2	0.1-6.1
2016-2020	9.6	18.5	13.5-25.0	7.4	30.8	21.5-43.3	2.2	7.3	3.6-14.2	1.2	2.2	0.8-5.3	0.6	3.0	0.6-9.1	0.6	1.9	0.4-7.4	
Kidney and renal pelvis	1976-1980	3.0	8.8	4.9-14.6	1.4	7.9	3.2-16.9	1.6	9.0	3.8-17.9	2.2	6.0	3.0-10.9	1.2	6.8	2.5-15.4	1.0	5.0	1.6-12.2
	1981-1985	3.0	8.7	4.8-14.4	1.8	11.1	4.9-21.6	1.2	5.8	2.1-13.1	0.8	2.5	0.7-6.4	0.4	2.4	0.3-9.1	0.4	2.6	0.3-9.0
	1986-1990	3.2	8.8	5.0-14.5	2.4	16.6	8.3-29.3	0.8	3.4	0.9-9.4	1.0	2.5	0.8-6.1	0.8	5.3	1.3-13.9	0.2	0.9	0.0-5.8
	1991-1995	3.4	9.8	5.6-15.7	1.8	11.2	5.1-21.5	1.6	9.1	3.8-18.0	1.4	3.7	1.5-7.7	1.0	6.0	1.9-14.3	0.4	1.8	0.2-7.1
	1996-2000	4.0	10.7	6.5-16.6	1.6	9.7	4.2-19.2	2.4	11.6	5.9-20.7	1.2	3.3	1.2-7.3	0.6	3.1	0.6-9.5	0.6	3.3	0.5-10.2
	2001-2005	6.0	14.6	9.8-20.9	3.8	20.1	12.1-31.6	2.2	10.5	5.2-19.2	1.6	3.9	1.7-7.9	0.6	3.2	0.7-9.6	1.0	4.8	1.5-11.7
	2006-2010	10.0	22.9	16.9-30.4	7.2	34.0	23.7-47.5	2.8	12.1	6.6-21.0	2.2	4.8	2.4-8.9	2.0	9.4	4.5-17.6	0.2	1.1	0.0-6.3
	2011-2015	7.6	17.6	12.3-24.6	4.4	18.9	11.6-29.4	3.2	16.0	9.0-26.7	2.4	4.8	2.5-8.8	1.6	6.2	2.6-13.0	0.8	3.2	0.9-9.2
2016-2020	9.4	21.5	15.5-29.1	7.2	32.2	22.3-45.3	2.2	11.3	5.1-21.6	1.2	2.5	0.9-5.9	0.8	3.0	0.8-8.8	0.4	2.1	0.2-8.2	
Brain and other central nervous system	1976-1980	1.4	4.7	1.8-9.7	1.2	8.3	3.0-18.2	0.2	1.2	0.0-6.6	0.4	1.7	0.2-5.7	0.4	3.5	0.4-11.7	0.0	0.0	0.0-4.4
	1981-1985	2.0	5.7	2.7-10.7	0.6	3.5	0.7-10.8	1.4	7.4	2.9-15.7	1.8	6.0	2.7-11.4	0.8	5.9	1.6-15.0	1.0	5.9	1.9-13.8
	1986-1990	1.6	4.9	2.1-9.8	0.6	4.7	0.9-13.4	1.0	5.6	1.8-13.4	0.8	2.5	0.7-6.4	0.2	1.5	0.0-7.8	0.6	2.9	0.6-9.0
	1991-1995	1.8	5.3	2.4-10.1	1.4	8.9	3.5-18.4	0.4	2.1	0.3-8.0	1.4	3.9	1.5-8.1	1.0	6.1	2.0-14.4	0.4	1.8	0.2-7.3
	1996-2000	2.6	7.5	4.0-12.8	1.4	8.3	3.3-17.3	1.2	6.8	2.4-15.0	0.8	2.2	0.6-5.8	0.6	3.4	0.7-10.3	0.2	1.0	0.0-6.1
	2001-2005	2.8	7.4	4.0-12.5	2.0	10.4	5.0-19.3	0.8	4.3	1.1-11.3	0.8	2.1	0.6-5.4	0.4	2.1	0.3-7.9	0.4	2.3	0.3-8.3
	2006-2010	3.8	9.7	5.8-15.4	1.2	5.4	1.9-12.4	2.6	14.5	7.5-25.2	3.2	7.6	4.3-12.6	1.8	8.6	3.9-16.6	1.4	7.1	2.7-15.1
	2011-2015	5.4	13.3	8.5-19.7	3.0	15.5	8.5-26.0	2.4	11.8	5.7-21.6	3.0	6.3	3.5-10.7	1.8	8.5	3.8-16.6	1.2	4.2	1.5-10.4
2016-2020	2.6	6.1	3.1-10.9	1.8	8.6	3.8-16.9	0.8	3.6	0.9-10.5	1.4	2.9	1.2-6.5	0.6	2.3	0.5-7.7	0.8	3.7	0.9-10.5	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Seneca**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	1.0	3.2	1.0-7.4	0.0	0.0	0.0-5.1	1.0	5.9	1.9-14.0	0.2	0.6	0.0-3.4	0.0	0.0	0.0-5.1	0.2	0.9	0.0-5.8
	1981-1985	1.0	2.6	0.8-6.3	0.6	4.7	0.8-13.7	0.4	1.9	0.2-7.4	0.2	0.5	0.0-3.2	0.0	0.0	0.0-5.1	0.2	1.0	0.0-6.1
	1986-1990	1.4	4.1	1.6-8.5	0.2	1.5	0.0-7.8	1.2	6.8	2.5-14.9	0.2	0.6	0.0-3.2	0.2	2.1	0.1-9.7	0.0	0.0	0.0-4.3
	1991-1995	1.2	3.4	1.2-7.4	0.2	1.0	0.0-6.4	1.0	5.2	1.6-12.6	0.2	0.6	0.0-3.2	0.0	0.0	0.0-4.8	0.2	0.8	0.0-5.5
	1996-2000	1.4	4.1	1.7-8.5	0.8	4.5	1.2-11.8	0.6	3.7	0.8-10.7	0.2	0.5	0.0-3.1	0.0	0.0	0.0-4.6	0.2	0.9	0.0-5.8
	2001-2005	2.8	8.1	4.4-13.6	0.6	3.2	0.7-9.6	2.2	14.5	7.2-25.7	0.2	0.5	0.0-3.1	0.2	1.0	0.0-6.0	0.0	0.0	0.0-4.3
	2006-2010	2.8	8.0	4.3-13.5	0.0	0.0	0.0-4.1	2.8	17.4	9.3-29.4	0.0	0.0	0.0-2.1	0.0	0.0	0.0-4.1	0.0	0.0	0.0-4.5
	2011-2015	4.8	14.5	9.1-21.9	2.0	10.7	5.0-20.1	2.8	19.1	10.1-32.6	0.4	1.1	0.1-4.0	0.2	1.2	0.0-6.6	0.2	0.7	0.0-5.7
2016-2020	4.8	13.9	8.6-21.0	1.4	6.9	2.6-15.0	3.4	21.8	12.2-35.7	0.0	0.0	0.0-2.3	0.0	0.0	0.0-4.3	0.0	0.0	0.0-4.9	
Hodgkin lymphoma	1976-1980	0.6	2.0	0.4-5.9	0.6	4.1	0.8-12.2	0.0	0.0	0.0-4.4	0.6	1.5	0.3-4.6	0.2	1.0	0.0-6.8	0.4	2.0	0.2-7.7
	1981-1985	1.2	3.3	1.2-7.4	0.4	2.1	0.2-8.3	0.8	4.6	1.2-11.9	0.6	1.9	0.4-5.5	0.4	2.5	0.3-9.4	0.2	1.0	0.0-6.0
	1986-1990	1.0	3.0	0.9-7.0	0.4	2.6	0.3-9.6	0.6	3.4	0.7-10.0	0.0	0.0	0.0-2.2	0.0	0.0	0.0-4.8	0.0	0.0	0.0-4.3
	1991-1995	1.8	5.5	2.5-10.5	0.8	4.8	1.3-12.6	1.0	6.1	1.9-14.3	0.0	0.0	0.0-2.2	0.0	0.0	0.0-4.8	0.0	0.0	0.0-4.3
	1996-2000	1.6	4.7	2.0-9.3	0.8	4.8	1.3-12.5	0.8	4.6	1.2-11.8	0.2	0.6	0.0-3.4	0.2	1.3	0.0-7.0	0.0	0.0	0.0-4.4
	2001-2005	1.8	4.8	2.2-9.2	1.2	6.2	2.3-13.7	0.6	3.0	0.6-9.3	0.4	1.0	0.1-3.9	0.2	1.1	0.0-6.4	0.2	0.7	0.0-5.4
	2006-2010	1.0	2.8	0.9-6.6	0.8	4.4	1.2-11.4	0.2	0.6	0.0-5.3	0.8	2.2	0.6-5.7	0.2	1.3	0.0-6.9	0.6	2.6	0.5-8.6
	2011-2015	1.2	3.1	1.1-7.1	0.6	3.2	0.6-9.6	0.6	2.9	0.5-9.2	0.2	0.6	0.0-3.3	0.0	0.0	0.0-4.2	0.2	1.3	0.0-7.1
2016-2020	2.2	5.5	2.6-10.2	1.2	6.5	2.3-14.5	1.0	4.6	1.4-11.8	0.2	0.4	0.0-2.8	0.0	0.0	0.0-4.3	0.2	0.7	0.0-5.8	
Non-Hodgkin lymphomas	1976-1980	4.2	12.0	7.4-18.5	1.8	11.6	4.9-22.9	2.4	12.4	6.4-22.0	2.6	7.5	3.9-12.9	0.8	5.1	1.3-13.4	1.8	9.1	4.1-17.7
	1981-1985	5.2	15.7	10.1-23.1	3.4	25.2	14.2-40.9	1.8	9.2	4.1-17.8	1.8	5.0	2.3-9.6	1.4	10.9	4.0-22.9	0.4	1.8	0.2-7.2
	1986-1990	5.4	15.0	9.9-22.0	2.4	14.6	7.4-25.9	3.0	15.1	8.3-25.3	3.0	8.2	4.6-13.7	1.2	8.7	3.1-18.8	1.8	8.4	3.8-16.5
	1991-1995	7.0	20.4	14.2-28.5	3.6	24.0	14.0-38.2	3.4	18.4	10.6-29.8	2.8	7.8	4.3-13.3	1.4	9.0	3.5-18.9	1.4	6.2	2.5-13.4
	1996-2000	7.0	19.1	13.3-26.6	4.0	22.8	13.9-35.4	3.0	15.7	8.6-26.4	3.2	8.7	5.0-14.2	1.8	11.0	4.9-21.0	1.4	6.9	2.7-14.7
	2001-2005	8.4	20.9	15.0-28.4	4.4	23.3	14.6-35.4	4.0	19.3	11.6-30.3	1.8	4.3	1.9-8.2	1.4	7.8	3.1-16.1	0.4	1.5	0.2-6.5
	2006-2010	11.4	26.3	19.9-34.4	6.6	31.5	21.6-44.6	4.8	20.5	13.0-31.2	3.2	7.3	4.1-12.0	1.8	8.7	4.0-16.9	1.4	6.0	2.3-13.2
	2011-2015	9.6	20.5	15.0-27.6	6.2	28.6	19.2-41.2	3.4	13.2	7.6-22.1	2.4	5.1	2.6-9.3	1.0	5.5	1.7-13.0	1.4	4.5	1.8-10.6
2016-2020	9.0	19.1	13.8-26.0	5.8	26.3	17.5-38.4	3.2	12.7	7.0-21.7	3.6	7.3	4.3-12.0	2.4	11.1	5.6-20.0	1.2	4.3	1.6-10.7	
Myeloma	1976-1980	1.4	3.9	1.6-8.2	0.4	1.9	0.2-8.1	1.0	5.5	1.8-13.0	0.8	2.2	0.6-5.8	0.4	2.4	0.3-9.1	0.4	1.8	0.2-7.2
	1981-1985	1.8	4.8	2.1-9.2	1.0	5.9	1.8-14.3	0.8	4.3	1.1-11.4	1.4	3.6	1.4-7.6	0.6	3.1	0.6-9.8	0.8	3.9	1.1-10.4
	1986-1990	1.4	4.4	1.8-9.2	0.6	4.7	1.0-13.2	0.8	3.7	1.0-10.0	1.0	2.8	0.9-6.6	0.4	3.5	0.4-11.8	0.6	2.6	0.5-8.3
	1991-1995	1.6	4.5	1.9-9.0	0.6	3.4	0.7-10.4	1.0	4.7	1.5-11.5	1.2	3.2	1.2-7.1	0.2	1.3	0.0-7.3	1.0	4.4	1.4-10.9
	1996-2000	2.2	6.0	3.0-10.8	1.0	6.8	2.2-15.7	1.2	5.8	2.1-13.2	1.2	3.1	1.1-6.9	0.2	1.3	0.0-7.2	1.0	4.4	1.4-10.9
	2001-2005	2.2	5.2	2.6-9.5	1.2	6.8	2.5-14.9	1.0	4.3	1.4-10.7	1.2	2.9	1.1-6.5	0.8	4.4	1.2-11.4	0.4	1.8	0.2-7.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Seneca**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	3.4	7.9	4.6-12.8	1.8	9.6	4.4-18.4	1.6	7.1	3.0-14.7	1.8	4.3	1.9-8.3	0.8	3.9	1.0-10.3	1.0	4.5	1.4-11.2
	2011-2015	2.8	5.8	3.2-10.1	2.2	10.2	5.1-18.7	0.6	2.2	0.4-7.7	1.6	3.1	1.3-6.5	0.8	3.8	1.0-10.1	0.8	2.9	0.8-8.6
	2016-2020	1.8	3.4	1.5-7.0	1.0	4.5	1.4-11.2	0.8	2.8	0.8-8.6	1.2	2.3	0.9-5.6	0.8	3.3	0.9-9.2	0.4	1.6	0.2-7.2
Leukemias	1976-1980	3.6	10.4	6.1-16.5	2.4	15.7	7.9-27.9	1.2	6.0	2.2-13.5	2.4	7.2	3.7-12.7	1.4	9.8	3.9-20.4	1.0	5.0	1.6-12.0
	1981-1985	4.0	11.7	7.1-18.2	2.4	16.0	8.0-28.6	1.6	8.0	3.4-16.1	2.4	6.9	3.6-12.2	1.4	9.7	3.7-20.4	1.0	5.1	1.6-12.3
	1986-1990	3.8	11.1	6.6-17.4	2.2	17.6	8.6-31.3	1.6	7.6	3.2-15.3	3.2	9.5	5.4-15.5	1.8	13.1	5.9-24.8	1.4	6.8	2.7-14.4
	1991-1995	4.4	12.5	7.8-19.0	2.6	17.2	8.9-29.7	1.8	10.4	4.7-19.8	3.0	8.2	4.6-13.7	1.8	12.0	5.2-23.2	1.2	6.7	2.4-14.9
	1996-2000	6.4	17.7	12.1-25.1	4.6	28.2	17.8-42.6	1.8	8.7	3.9-17.0	3.6	9.7	5.7-15.4	1.4	8.7	3.4-18.1	2.2	10.3	5.1-19.0
	2001-2005	8.6	21.3	15.3-28.8	4.0	21.0	12.8-32.6	4.6	20.4	12.7-31.3	3.0	7.5	4.2-12.6	1.4	7.5	3.0-15.6	1.6	6.8	2.8-14.2
	2006-2010	8.2	18.9	13.5-25.9	4.4	21.2	13.2-32.4	3.8	16.7	9.9-26.8	4.6	10.5	6.6-16.0	1.6	8.3	3.5-16.5	3.0	11.9	6.6-20.4
	2011-2015	8.0	18.4	13.0-25.5	5.2	24.8	16.0-36.9	2.8	11.7	6.3-20.5	3.6	7.9	4.6-12.8	2.8	12.9	7.0-22.1	0.8	2.9	0.8-8.7
	2016-2020	7.8	17.8	12.4-24.8	6.0	30.1	20.1-43.6	1.8	7.0	3.2-14.5	3.6	7.9	4.6-12.9	3.0	15.3	8.5-25.5	0.6	2.5	0.5-8.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Steuben**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	371.2	382.1	364.6-400.2	194.4	453.6	424.4-484.2	176.8	334.8	312.8-358.1	203.0	210.6	197.6-224.1	112.0	267.9	245.3-291.9	91.0	170.4	154.9-187.1
	1981-1985	429.0	430.2	411.8-449.2	211.8	486.7	456.8-518.1	217.2	403.5	379.2-428.9	218.4	217.5	204.6-231.0	115.0	266.7	244.5-290.3	103.4	186.4	170.3-203.6
	1986-1990	471.0	453.0	434.6-472.0	249.6	551.0	519.7-583.5	221.4	393.7	370.3-418.3	234.2	224.9	212.1-238.4	122.8	280.4	257.8-304.4	111.4	190.7	174.9-207.6
	1991-1995	533.0	486.5	468.0-505.5	284.0	582.1	551.6-614.0	249.0	425.2	401.5-450.0	247.2	223.6	211.2-236.6	138.6	296.9	274.6-320.5	108.6	175.8	161.1-191.6
	1996-2000	546.2	491.0	472.7-509.9	280.6	557.4	528.3-587.7	265.6	441.2	417.5-466.0	244.2	215.3	203.4-227.8	130.4	268.9	248.4-290.7	113.8	178.9	164.3-194.5
	2001-2005	596.0	515.5	497.1-534.5	311.4	590.4	561.2-620.7	284.6	458.4	434.5-483.3	240.2	203.8	192.4-215.7	121.0	235.7	217.1-255.4	119.2	179.8	165.5-195.1
	2006-2010	575.8	477.8	460.2-495.8	301.4	530.1	503.3-558.0	274.4	440.9	417.3-465.6	239.8	194.0	183.1-205.4	128.2	234.6	216.6-253.8	111.6	164.8	151.2-179.4
	2011-2015	611.6	471.3	454.3-488.8	328.0	529.1	503.2-556.0	283.6	427.2	404.4-451.1	231.2	174.0	163.9-184.5	123.2	205.1	188.9-222.3	108.0	151.5	138.6-165.3
	2016-2020	615.8	459.7	442.9-477.1	316.4	480.6	456.4-506.0	299.4	445.9	422.3-470.6	223.6	160.4	150.9-170.4	117.2	182.7	167.9-198.6	106.4	142.9	130.6-156.3
Oral cavity and pharynx	1976-1980	12.6	12.6	9.7-16.3	9.4	21.3	15.4-28.7	3.2	5.7	3.2-9.3	4.0	3.9	2.4-6.0	2.6	5.4	2.8-9.3	1.4	2.5	1.0-5.2
	1981-1985	14.4	13.8	10.8-17.5	10.0	21.0	15.4-27.9	4.4	8.1	5.0-12.4	5.0	4.8	3.1-7.1	4.4	9.6	5.9-14.7	0.6	1.0	0.2-3.1
	1986-1990	12.4	12.1	9.2-15.6	8.6	18.4	13.2-24.9	3.8	6.7	4.0-10.6	3.6	3.5	2.0-5.6	2.2	5.2	2.5-9.4	1.4	2.6	1.0-5.4
	1991-1995	13.4	12.2	9.4-15.5	9.6	19.0	13.9-25.3	3.8	6.9	4.1-10.9	4.2	3.7	2.3-5.8	3.0	6.3	3.4-10.5	1.2	2.1	0.8-4.7
	1996-2000	11.4	10.2	7.8-13.3	9.0	17.8	13.0-23.9	2.4	3.8	2.0-6.8	3.2	2.8	1.6-4.6	2.4	4.6	2.4-8.1	0.8	1.3	0.4-3.6
	2001-2005	13.6	11.7	9.1-14.9	9.4	17.2	12.6-23.0	4.2	6.6	4.1-10.2	4.0	3.3	2.0-5.2	3.2	5.9	3.4-9.7	0.8	1.1	0.3-3.1
	2006-2010	14.8	12.1	9.5-15.3	9.0	15.0	10.9-20.3	5.8	9.3	6.2-13.6	4.2	3.3	2.0-5.1	2.4	4.0	2.0-7.2	1.8	2.7	1.2-5.3
	2011-2015	17.6	13.0	10.3-16.1	12.0	17.9	13.6-23.3	5.6	8.2	5.3-12.1	3.8	2.9	1.8-4.7	3.0	5.2	2.9-8.8	0.8	1.1	0.3-3.2
	2016-2020	17.6	12.7	10.1-15.9	14.0	20.8	16.0-26.8	3.6	5.0	2.9-8.2	3.6	2.6	1.5-4.3	2.4	3.6	1.8-6.6	1.2	1.6	0.6-3.9
Esophagus	1976-1980	4.8	4.9	3.1-7.3	3.6	9.0	5.2-14.4	1.2	2.0	0.7-4.6	4.8	4.8	3.1-7.2	4.0	9.0	5.4-14.1	0.8	1.4	0.4-3.7
	1981-1985	3.4	3.3	1.9-5.3	2.2	5.2	2.5-9.4	1.2	2.1	0.8-4.6	3.0	3.0	1.7-5.0	1.6	4.3	1.8-8.4	1.4	2.5	1.0-5.4
	1986-1990	3.8	3.5	2.1-5.5	3.0	5.6	3.1-9.3	0.8	1.7	0.4-4.4	4.4	3.9	2.4-5.9	3.4	6.3	3.7-10.3	1.0	1.8	0.6-4.3
	1991-1995	6.6	6.1	4.2-8.6	4.8	9.9	6.3-14.9	1.8	2.9	1.3-5.6	6.8	6.3	4.3-8.8	5.4	10.9	7.1-16.0	1.4	2.3	0.9-4.8
	1996-2000	4.6	4.2	2.6-6.3	3.8	7.5	4.5-11.9	0.8	1.2	0.3-3.2	6.8	6.2	4.3-8.6	5.2	10.3	6.7-15.2	1.6	2.6	1.1-5.3
	2001-2005	4.8	4.0	2.6-6.0	3.8	6.8	4.1-10.8	1.0	1.4	0.4-3.4	5.4	4.6	3.0-6.7	4.0	7.2	4.4-11.2	1.4	2.0	0.8-4.4
	2006-2010	8.2	6.5	4.6-8.9	6.4	11.1	7.6-15.9	1.8	2.7	1.2-5.3	6.2	5.2	3.5-7.4	5.4	9.9	6.5-14.5	0.8	1.2	0.3-3.3
	2011-2015	5.6	4.2	2.7-6.2	4.6	7.3	4.6-11.3	1.0	1.3	0.4-3.4	6.6	5.0	3.4-7.1	5.8	9.6	6.4-14.0	0.8	0.9	0.2-2.8
	2016-2020	7.0	5.6	3.8-8.0	5.4	9.2	5.9-13.7	1.6	2.4	1.0-5.1	5.8	4.2	2.8-6.2	4.6	7.0	4.4-10.8	1.2	1.8	0.6-4.3
Stomach	1976-1980	7.6	8.0	5.7-11.1	5.2	12.1	7.8-17.9	2.4	4.8	2.4-8.4	5.8	6.2	4.1-8.9	3.8	9.8	5.8-15.4	2.0	3.8	1.8-7.0
	1981-1985	9.6	9.8	7.2-13.0	5.0	11.5	7.4-17.2	4.6	8.1	5.1-12.3	6.8	6.8	4.7-9.6	3.8	9.0	5.3-14.2	3.0	5.0	2.8-8.4
	1986-1990	7.0	6.7	4.6-9.3	4.4	9.4	5.8-14.4	2.6	4.5	2.3-7.8	5.8	5.6	3.8-8.1	2.8	6.9	3.7-11.7	3.0	4.8	2.7-8.1
	1991-1995	7.6	7.1	5.0-9.8	5.2	11.1	7.1-16.3	2.4	3.7	1.9-6.6	5.6	5.0	3.3-7.3	4.0	8.4	5.1-13.2	1.6	2.4	1.0-4.9
	1996-2000	9.8	8.7	6.4-11.5	6.0	11.6	7.8-16.6	3.8	5.7	3.4-9.0	4.6	4.1	2.6-6.1	2.4	4.7	2.4-8.2	2.2	3.3	1.6-6.0
	2001-2005	7.6	6.5	4.6-8.9	4.8	9.2	5.9-13.8	2.8	4.1	2.2-7.1	3.4	2.9	1.7-4.6	1.8	3.5	1.6-6.7	1.6	2.3	1.0-4.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Steuben**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	7.6	6.2	4.4-8.5	4.8	8.7	5.5-13.1	2.8	4.0	2.1-6.9	3.0	2.5	1.4-4.1	1.6	3.1	1.3-6.2	1.4	1.9	0.8-4.2
	2011-2015	6.8	5.4	3.7-7.7	4.8	8.4	5.3-12.7	2.0	3.0	1.4-5.9	2.6	1.8	1.0-3.3	1.8	2.8	1.3-5.6	0.8	1.2	0.3-3.3
	2016-2020	5.6	4.2	2.8-6.3	3.8	5.7	3.4-9.2	1.8	3.0	1.3-6.1	2.4	1.8	0.9-3.3	1.0	1.8	0.6-4.4	1.4	1.8	0.7-4.1
Colorectal	1976-1980	59.6	61.0	54.2-68.4	28.4	65.8	55.0-77.9	31.2	56.7	48.1-66.4	30.2	31.7	26.7-37.2	13.4	33.6	25.7-42.9	16.8	30.8	24.5-38.3
	1981-1985	64.2	64.7	57.7-72.4	30.8	71.4	60.2-84.1	33.4	59.0	50.3-68.9	29.8	30.4	25.6-35.7	14.4	35.0	27.1-44.5	15.4	26.9	21.1-33.8
	1986-1990	66.2	63.7	57.0-71.1	32.4	72.8	61.7-85.4	33.8	57.8	49.2-67.4	31.6	31.0	26.3-36.3	14.8	35.4	27.5-44.8	16.8	27.9	22.2-34.7
	1991-1995	65.4	59.3	53.0-66.2	34.4	72.7	61.9-84.7	31.0	48.8	41.3-57.3	27.2	24.4	20.5-29.0	15.2	33.0	25.8-41.4	12.0	18.2	13.9-23.6
	1996-2000	64.0	56.4	50.3-62.9	31.6	63.9	54.2-74.9	32.4	48.9	41.6-57.2	30.0	26.2	22.2-30.8	14.8	31.2	24.3-39.3	15.2	22.6	17.8-28.5
	2001-2005	70.0	59.1	53.0-65.6	30.6	58.5	49.5-68.6	39.4	58.4	50.5-67.4	24.6	20.8	17.3-24.9	13.4	26.4	20.4-33.7	11.2	15.9	12.0-20.9
	2006-2010	56.6	47.0	41.7-52.9	27.4	50.0	41.9-59.3	29.2	44.2	37.2-52.3	19.4	15.9	12.9-19.5	10.0	19.2	14.2-25.4	9.4	13.2	9.6-17.7
	2011-2015	54.8	41.4	36.5-46.7	31.4	51.3	43.5-60.2	23.4	33.7	27.6-40.7	20.2	14.7	12.0-18.0	11.0	18.4	13.8-24.2	9.2	12.0	8.7-16.3
2016-2020	55.6	41.3	36.4-46.8	28.6	45.0	37.6-53.4	27.0	38.1	31.6-45.6	20.2	14.8	12.0-18.2	10.8	17.8	13.3-23.5	9.4	12.7	9.2-17.3	
Colon excluding rectum	1976-1980	44.4	45.8	39.9-52.3	21.2	49.6	40.3-60.4	23.2	42.2	34.8-50.7	24.6	25.9	21.4-30.9	10.6	26.5	19.6-35.0	14.0	25.7	20.0-32.6
	1981-1985	45.6	46.0	40.2-52.5	20.0	46.2	37.2-56.5	25.6	45.6	37.9-54.5	23.4	24.0	19.8-28.8	11.2	27.1	20.2-35.6	12.2	21.2	16.2-27.4
	1986-1990	48.4	47.2	41.4-53.6	22.6	51.9	42.5-62.8	25.8	44.1	36.7-52.6	27.6	27.0	22.6-31.9	13.2	31.8	24.3-40.7	14.4	23.8	18.6-30.1
	1991-1995	50.0	45.6	40.1-51.7	24.6	52.0	43.0-62.3	25.4	40.3	33.5-48.2	24.4	22.0	18.2-26.3	13.8	29.9	23.1-38.0	10.6	16.2	12.1-21.4
	1996-2000	47.0	41.3	36.2-47.0	21.6	44.0	36.0-53.3	25.4	38.6	32.1-46.1	24.8	21.7	18.0-25.9	11.8	24.8	18.8-32.2	13.0	19.4	14.9-24.9
	2001-2005	51.0	42.9	37.8-48.5	21.0	40.5	33.1-49.2	30.0	44.1	37.3-51.9	19.0	16.1	13.0-19.8	9.8	19.6	14.5-26.0	9.2	13.4	9.7-18.0
	2006-2010	42.0	34.8	30.2-40.0	19.6	36.1	29.2-44.2	22.4	33.4	27.4-40.4	17.0	14.0	11.1-17.3	8.6	16.5	11.9-22.3	8.4	11.8	8.5-16.2
	2011-2015	41.6	31.2	27.1-35.9	21.8	35.3	28.9-42.8	19.8	28.2	22.7-34.7	14.6	10.7	8.4-13.5	7.0	12.0	8.3-16.9	7.6	9.8	6.9-13.7
2016-2020	40.4	30.1	26.0-34.8	20.6	33.2	26.9-40.6	19.8	27.3	22.0-33.7	15.6	11.4	9.0-14.4	8.6	14.4	10.3-19.6	7.0	9.2	6.3-13.2	
Rectum & rectosigmoid	1976-1980	15.2	15.2	11.9-19.1	7.2	16.1	11.1-22.6	8.0	14.5	10.3-19.9	5.6	5.8	3.8-8.4	2.8	7.0	3.8-11.9	2.8	5.1	2.7-8.6
	1981-1985	18.6	18.7	15.0-23.0	10.8	25.3	18.8-33.2	7.8	13.4	9.5-18.5	6.4	6.4	4.4-9.1	3.2	7.9	4.4-12.9	3.2	5.7	3.2-9.4
	1986-1990	17.8	16.6	13.3-20.5	9.8	20.9	15.3-27.9	8.0	13.7	9.7-18.8	4.0	4.0	2.4-6.2	1.6	3.6	1.5-7.3	2.4	4.1	2.1-7.4
	1991-1995	15.4	13.7	10.8-17.2	9.8	20.6	15.1-27.4	5.6	8.5	5.6-12.4	2.8	2.5	1.3-4.2	1.4	3.1	1.2-6.3	1.4	2.0	0.8-4.3
	1996-2000	17.0	15.0	12.0-18.6	10.0	19.9	14.7-26.4	7.0	10.3	7.1-14.5	5.2	4.5	2.9-6.7	3.0	6.3	3.5-10.5	2.2	3.2	1.6-5.9
	2001-2005	19.0	16.2	13.1-19.8	9.6	18.0	13.2-23.9	9.4	14.4	10.5-19.3	5.6	4.7	3.1-6.8	3.6	6.8	4.0-10.9	2.0	2.6	1.2-5.0
	2006-2010	14.6	12.2	9.5-15.4	7.8	13.9	9.8-19.2	6.8	10.9	7.5-15.4	2.4	2.0	1.0-3.5	1.4	2.7	1.1-5.7	1.0	1.4	0.4-3.5
	2011-2015	13.2	10.2	7.8-13.0	9.6	16.0	11.7-21.4	3.6	5.5	3.2-8.9	5.6	4.1	2.7-6.0	4.0	6.4	3.9-10.1	1.6	2.2	0.9-4.6
2016-2020	15.2	11.2	8.7-14.3	8.0	11.7	8.3-16.3	7.2	10.8	7.3-15.3	4.6	3.4	2.1-5.2	2.2	3.5	1.7-6.5	2.4	3.5	1.7-6.5	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Steuben**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	3.4	3.3	1.9-5.4	2.0	4.4	2.1-8.2	1.4	2.7	1.0-5.6	3.4	3.4	2.0-5.5	2.2	4.9	2.4-8.9	1.2	2.2	0.8-4.8
	1981-1985	3.2	3.0	1.7-4.9	1.6	3.8	1.6-7.6	1.6	2.7	1.1-5.4	2.8	2.7	1.5-4.6	1.8	4.0	1.8-7.8	1.0	1.7	0.5-4.0
	1986-1990	4.4	4.4	2.7-6.7	3.6	8.1	4.7-13.0	0.8	1.3	0.3-3.4	3.0	2.8	1.6-4.7	1.8	3.5	1.6-6.8	1.2	2.2	0.8-4.9
	1991-1995	4.8	4.1	2.6-6.2	3.6	7.5	4.4-12.1	1.2	1.8	0.7-4.1	4.8	4.1	2.6-6.2	3.6	7.8	4.5-12.5	1.2	1.8	0.7-4.2
	1996-2000	4.2	3.7	2.3-5.8	3.2	6.1	3.5-10.0	1.0	1.5	0.5-3.7	3.6	3.2	1.9-5.1	2.4	4.8	2.5-8.5	1.2	2.1	0.7-4.6
	2001-2005	6.0	5.0	3.4-7.3	4.0	7.6	4.6-11.9	2.0	2.9	1.4-5.5	4.6	3.9	2.4-5.8	2.6	5.0	2.6-8.6	2.0	2.9	1.4-5.5
	2006-2010	5.2	4.0	2.6-6.0	4.2	7.0	4.3-10.9	1.0	1.4	0.5-3.6	3.2	2.6	1.5-4.3	2.8	5.2	2.8-8.9	0.4	0.5	0.1-2.1
	2011-2015	8.6	6.4	4.6-8.7	6.2	9.5	6.4-13.8	2.4	3.5	1.8-6.4	8.6	6.3	4.6-8.7	5.6	8.4	5.5-12.4	3.0	4.4	2.4-7.5
	2016-2020	7.0	4.6	3.2-6.5	4.0	5.3	3.2-8.6	3.0	3.8	2.1-6.6	5.4	3.5	2.3-5.3	2.8	3.7	2.0-6.5	2.6	3.3	1.7-6.0
Pancreas	1976-1980	8.8	8.6	6.2-11.6	5.0	11.1	7.0-16.7	3.8	6.6	4.0-10.4	9.0	8.6	6.3-11.6	5.4	11.7	7.5-17.3	3.6	6.4	3.8-10.2
	1981-1985	12.4	12.5	9.5-16.1	7.0	16.1	11.1-22.6	5.4	9.8	6.4-14.3	10.8	10.8	8.1-14.2	6.2	14.7	9.9-21.0	4.6	8.3	5.2-12.6
	1986-1990	12.2	11.8	9.0-15.2	5.6	12.9	8.5-18.8	6.6	11.1	7.6-15.7	11.0	10.9	8.2-14.2	5.0	11.4	7.3-16.9	6.0	10.2	6.8-14.7
	1991-1995	12.0	11.0	8.4-14.2	6.2	13.4	9.0-19.2	5.8	8.9	5.9-12.9	11.6	10.5	8.0-13.6	6.0	12.7	8.5-18.3	5.6	8.3	5.5-12.1
	1996-2000	12.4	10.8	8.3-13.9	6.4	12.6	8.6-17.8	6.0	9.2	6.2-13.4	10.4	9.1	6.8-11.9	5.8	11.7	7.8-16.9	4.6	6.9	4.3-10.5
	2001-2005	14.8	12.5	9.8-15.7	7.2	14.3	10.0-19.9	7.6	11.1	7.8-15.3	14.4	12.0	9.4-15.2	6.4	12.5	8.5-17.8	8.0	11.8	8.4-16.2
	2006-2010	14.2	11.3	8.8-14.3	7.4	13.2	9.3-18.4	6.8	9.8	6.8-13.9	12.6	10.0	7.7-12.9	6.6	11.9	8.2-16.9	6.0	8.7	5.8-12.6
	2011-2015	17.2	12.6	10.0-15.7	8.6	13.3	9.6-18.1	8.6	12.2	8.7-16.7	13.2	9.8	7.5-12.5	7.4	12.2	8.6-17.0	5.8	7.9	5.3-11.6
	2016-2020	20.2	14.7	11.9-18.1	9.2	14.6	10.6-19.7	11.0	14.8	11.0-19.6	17.4	12.5	10.0-15.6	9.0	14.4	10.4-19.5	8.4	10.8	7.7-14.9
Larynx	1976-1980	5.4	5.8	3.8-8.5	4.6	11.2	6.9-17.1	0.8	1.4	0.4-3.7	2.2	2.4	1.2-4.3	2.2	5.9	2.8-10.5	0.0	0.0	0.0-1.5
	1981-1985	7.2	7.0	4.9-9.8	6.4	13.5	9.1-19.3	0.8	1.7	0.4-4.3	2.6	2.7	1.4-4.6	1.8	4.2	1.9-8.2	0.8	1.4	0.4-3.8
	1986-1990	7.6	7.3	5.1-10.1	5.0	10.4	6.7-15.5	2.6	4.7	2.5-8.2	2.2	2.0	1.0-3.6	2.2	4.3	2.1-7.8	0.0	0.0	0.0-1.5
	1991-1995	7.6	6.7	4.8-9.3	6.6	12.6	8.6-17.7	1.0	1.6	0.5-4.0	2.8	2.4	1.3-4.0	1.8	3.4	1.5-6.6	1.0	1.4	0.5-3.6
	1996-2000	6.8	6.3	4.3-8.8	5.2	10.0	6.5-14.7	1.6	2.8	1.2-5.5	2.0	1.8	0.8-3.3	1.8	3.4	1.6-6.6	0.2	0.3	0.0-1.8
	2001-2005	3.6	3.1	1.8-4.9	2.6	4.8	2.5-8.3	1.0	1.6	0.5-3.8	2.2	1.9	0.9-3.4	1.8	3.4	1.5-6.5	0.4	0.6	0.1-2.5
	2006-2010	5.2	4.1	2.7-6.1	5.0	8.3	5.3-12.4	0.2	0.3	0.0-1.9	2.2	1.7	0.9-3.2	2.0	3.3	1.5-6.2	0.2	0.3	0.0-2.1
	2011-2015	7.4	5.6	3.9-7.8	6.2	9.7	6.5-14.0	1.2	1.8	0.6-4.1	2.0	1.4	0.7-2.7	1.6	2.4	1.0-4.9	0.4	0.6	0.1-2.4
	2016-2020	7.6	5.7	4.0-8.0	5.4	8.2	5.3-12.2	2.2	3.4	1.7-6.3	2.8	2.1	1.1-3.7	2.4	3.7	1.9-6.8	0.4	0.6	0.1-2.5
Lung and bronchus	1976-1980	63.8	64.7	57.7-72.4	47.4	107.1	93.5-122.2	16.4	31.0	24.5-38.7	51.8	52.2	45.9-59.0	37.6	84.6	72.6-98.1	14.2	26.4	20.5-33.4
	1981-1985	78.0	75.4	67.9-83.4	51.0	110.1	96.5-125.0	27.0	50.3	42.0-59.9	61.0	59.0	52.4-66.2	41.4	90.3	78.0-104.0	19.6	36.1	29.1-44.2
	1986-1990	85.6	80.2	72.7-88.3	56.8	120.3	106.2-135.6	28.8	50.9	42.8-60.2	69.8	65.6	58.8-73.0	43.6	94.3	81.8-108.2	26.2	45.7	38.1-54.5
	1991-1995	91.2	82.5	75.0-90.5	55.4	113.2	100.0-127.6	35.8	60.1	51.5-69.8	74.6	67.3	60.5-74.5	46.8	97.5	85.1-111.1	27.8	46.0	38.6-54.6
	1996-2000	89.8	79.0	71.8-86.7	52.0	101.6	89.6-114.9	37.8	60.5	52.1-69.9	68.0	59.7	53.5-66.5	43.6	86.8	75.5-99.2	24.4	38.6	32.0-46.3
	2001-2005	93.6	79.7	72.6-87.3	51.8	98.7	86.9-111.6	41.8	65.9	57.2-75.6	72.4	61.3	55.2-68.0	40.2	76.4	66.2-87.9	32.2	49.5	42.1-57.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Steuben**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	106.8	85.5	78.3-93.1	63.2	109.4	97.5-122.4	43.6	66.1	57.5-75.7	82.2	66.0	59.7-72.8	49.4	88.0	77.2-99.9	32.8	49.0	41.7-57.4
	2011-2015	104.2	78.7	71.9-85.9	57.0	92.4	81.8-104.1	47.2	68.0	59.3-77.6	73.6	55.6	50.0-61.7	41.4	68.8	59.6-79.1	32.2	44.7	37.9-52.4
	2016-2020	93.2	64.5	58.6-70.8	50.0	73.0	64.0-83.0	43.2	56.9	49.4-65.3	65.0	45.2	40.4-50.6	35.0	51.8	44.2-60.4	30.0	39.3	33.2-46.4
Melanoma of the skin	1976-1980	6.2	6.9	4.7-9.8	2.8	6.7	3.6-11.4	3.4	6.8	4.0-11.0	1.8	1.9	0.9-3.7	1.2	2.5	0.9-5.7	0.6	1.4	0.3-3.9
	1981-1985	7.0	7.4	5.1-10.4	3.6	7.8	4.5-12.5	3.4	6.8	3.9-11.0	1.6	1.9	0.8-3.6	0.6	1.4	0.3-4.1	1.0	2.1	0.7-5.0
	1986-1990	8.2	8.6	6.1-11.7	5.8	13.1	8.7-18.9	2.4	4.7	2.4-8.3	2.2	2.1	1.1-3.9	1.4	3.1	1.2-6.6	0.8	1.5	0.4-3.9
	1991-1995	10.8	10.3	7.7-13.4	6.2	11.8	8.0-16.8	4.6	8.7	5.5-13.1	2.6	2.4	1.3-4.2	1.2	2.3	0.8-5.0	1.4	2.5	1.0-5.2
	1996-2000	15.0	14.0	11.0-17.5	9.2	18.3	13.4-24.5	5.8	9.9	6.6-14.4	2.0	1.8	0.9-3.4	1.4	3.0	1.2-6.3	0.6	1.1	0.2-3.3
	2001-2005	22.8	21.1	17.4-25.5	13.4	25.9	20.0-33.0	9.4	17.7	12.9-23.7	3.0	2.7	1.5-4.5	1.2	2.7	1.0-5.9	1.8	3.0	1.4-5.9
	2006-2010	26.4	23.4	19.5-27.9	14.4	26.3	20.5-33.3	12.0	22.8	17.3-29.6	4.0	3.4	2.1-5.4	2.2	4.2	2.1-7.6	1.8	2.8	1.2-5.5
	2011-2015	27.0	22.8	19.0-27.2	15.0	25.7	20.1-32.5	12.0	21.2	16.0-27.7	3.4	2.7	1.5-4.5	1.8	3.0	1.3-6.0	1.6	2.7	1.1-5.6
	2016-2020	31.6	26.0	21.9-30.7	18.2	29.9	23.8-37.1	13.4	23.6	17.9-30.5	4.4	3.5	2.2-5.5	3.0	5.3	2.8-9.0	1.4	2.0	0.8-4.4
Female breast	1976-1980							41.6	82.8	71.7-95.1							17.2	33.9	27.0-42.0
	1981-1985							52.4	100.7	88.6-114.1							19.4	36.0	29.0-44.1
	1986-1990							62.2	113.4	100.9-127.2							16.4	28.4	22.4-35.4
	1991-1995							71.8	128.0	114.8-142.2							15.0	25.7	20.1-32.4
	1996-2000							75.4	131.3	118.2-145.5							20.8	34.0	27.7-41.3
	2001-2005							81.2	134.4	121.5-148.4							17.2	26.5	21.1-32.9
	2006-2010							70.8	113.4	101.7-126.2							17.0	25.7	20.5-32.0
	2011-2015							72.4	107.8	96.5-120.0							11.8	17.0	12.8-22.3
	2016-2020							87.0	131.7	118.8-145.7							13.6	18.2	13.9-23.6
Cervix uteri	1976-1980							8.4	15.9	11.3-21.6							3.4	6.3	3.6-10.2
	1981-1985							10.0	20.7	15.2-27.5							1.6	3.0	1.3-6.0
	1986-1990							8.2	16.0	11.4-21.8							2.8	4.7	2.6-8.1
	1991-1995							6.6	12.5	8.5-17.6							1.8	3.3	1.5-6.4
	1996-2000							6.2	11.5	7.7-16.4							2.8	4.9	2.7-8.3
	2001-2005							5.4	10.8	7.1-15.8							1.4	2.2	0.9-4.8
	2006-2010							5.2	11.0	7.1-16.2							1.2	1.8	0.6-4.2
	2011-2015							4.0	6.3	3.7-10.1							1.4	1.8	0.7-4.0
	2016-2020							4.6	7.9	4.8-12.2							1.6	2.1	0.9-4.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Steuben**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							14.0	25.1	19.5-31.8							2.8	5.1	2.8-8.6
	1981-1985							14.6	26.2	20.4-33.2							4.0	7.0	4.2-10.9
	1986-1990							18.4	33.3	26.6-41.1							3.4	5.9	3.4-9.6
	1991-1995							16.8	28.3	22.4-35.2							3.4	5.2	3.0-8.5
	1996-2000							16.6	28.4	22.6-35.4							3.6	6.0	3.5-9.6
	2001-2005							16.8	27.4	21.8-34.1							2.8	3.9	2.1-6.8
	2006-2010							16.4	25.4	20.1-31.7							3.8	5.5	3.3-8.8
	2011-2015							19.8	28.3	22.8-34.8							3.2	4.8	2.7-8.1
	2016-2020							23.6	35.6	29.1-43.3							4.2	5.8	3.5-9.2
Ovary	1976-1980							8.6	17.3	12.4-23.5							6.4	12.4	8.4-17.7
	1981-1985							9.6	18.2	13.3-24.4							6.2	11.2	7.6-16.1
	1986-1990							8.4	15.4	11.0-21.1							5.4	9.3	6.1-13.7
	1991-1995							10.8	19.1	14.2-25.0							6.6	10.8	7.4-15.4
	1996-2000							11.0	18.5	13.9-24.2							6.6	10.3	7.0-14.6
	2001-2005							8.4	12.9	9.3-17.6							7.4	11.2	7.8-15.6
	2006-2010							7.6	12.3	8.6-17.2							5.6	8.3	5.5-12.2
	2011-2015							7.6	12.3	8.6-17.2							5.8	8.0	5.3-11.8
	2016-2020							4.8	7.3	4.5-11.3							4.4	6.5	4.0-10.3
Prostate	1976-1980				30.2	76.9	64.7-90.6							12.0	34.9	26.4-45.0			
	1981-1985				34.0	84.5	71.9-98.7							10.6	27.9	20.7-36.8			
	1986-1990				53.2	122.1	107.4-138.3							11.6	30.2	22.6-39.2			
	1991-1995				80.0	159.6	144.0-176.4							18.8	43.9	35.3-54.0			
	1996-2000				72.8	142.3	127.9-157.9							16.4	37.5	29.7-46.6			
	2001-2005				88.4	162.2	147.3-178.2							11.4	23.8	18.0-30.9			
	2006-2010				74.2	124.5	112.0-138.1							10.0	19.9	14.8-26.3			
	2011-2015				72.4	107.3	96.4-119.4							7.6	13.6	9.6-18.7			
	2016-2020				81.6	112.6	101.7-124.6							11.2	18.0	13.5-23.6			
Testis	1976-1980				2.6	5.5	2.9-9.6							0.6	1.2	0.2-3.6			
	1981-1985				1.2	2.9	1.0-6.4							0.0	0.0	0.0-1.7			
	1986-1990				2.4	5.1	2.6-8.9							0.4	0.9	0.1-3.2			
	1991-1995				2.0	4.1	2.0-7.6							0.2	0.4	0.0-2.2			
	1996-2000				1.8	3.9	1.8-7.3							0.4	0.9	0.1-3.1			
	2001-2005				3.4	7.7	4.5-12.3							0.0	0.0	0.0-1.6			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Steuben**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				3.4	7.5	4.3-12.1							0.0	0.0	0.0-1.5			
	2011-2015				3.8	8.6	5.2-13.5							0.0	0.0	0.0-1.6			
	2016-2020				1.4	3.3	1.3-6.8							0.4	0.5	0.1-2.3			
Urinary bladder (incl. in situ)	1976-1980	18.6	18.7	15.0-22.9	13.6	31.2	24.0-39.9	5.0	9.4	6.0-13.9	4.8	4.8	3.0-7.1	3.4	8.2	4.6-13.2	1.4	2.4	1.0-5.1
	1981-1985	25.0	25.3	21.0-30.2	19.4	47.1	37.9-57.8	5.6	9.7	6.4-14.1	8.6	8.7	6.3-11.8	5.6	13.8	9.0-20.1	3.0	5.2	2.9-8.7
	1986-1990	22.0	20.4	16.7-24.7	18.4	40.1	32.0-49.5	3.6	5.7	3.4-9.1	5.6	5.4	3.6-7.9	3.8	9.9	5.9-15.5	1.8	2.8	1.3-5.5
	1991-1995	26.4	23.7	19.8-28.2	20.6	43.1	34.9-52.5	5.8	9.3	6.2-13.5	7.6	6.7	4.8-9.3	5.2	11.6	7.5-17.2	2.4	3.3	1.7-6.0
	1996-2000	30.0	26.5	22.4-31.2	22.0	44.2	36.2-53.4	8.0	12.5	8.9-17.2	5.4	4.7	3.1-6.9	4.2	9.3	5.7-14.2	1.2	1.7	0.6-4.0
	2001-2005	30.2	25.6	21.7-30.1	23.8	46.1	38.2-55.3	6.4	9.9	6.7-14.1	5.6	4.8	3.2-6.9	3.0	5.7	3.2-9.5	2.6	3.9	2.0-6.8
	2006-2010	33.0	26.6	22.7-31.1	25.2	45.5	37.8-54.3	7.8	11.6	8.2-16.0	7.2	5.9	4.1-8.3	4.8	9.3	5.9-14.0	2.4	3.4	1.7-6.2
	2011-2015	37.4	28.4	24.4-32.9	29.2	48.2	40.6-56.9	8.2	11.7	8.3-16.2	5.2	3.7	2.4-5.5	3.6	5.6	3.3-9.1	1.6	2.1	0.9-4.5
2016-2020	31.6	22.5	19.0-26.5	23.6	36.8	30.2-44.4	8.0	10.9	7.7-15.2	6.2	4.3	2.9-6.3	5.8	9.4	6.2-13.6	0.4	0.6	0.1-2.5	
Kidney and renal pelvis	1976-1980	6.2	6.6	4.4-9.4	2.8	6.0	3.2-10.2	3.4	6.6	3.8-10.6	3.6	3.7	2.2-5.9	1.8	3.6	1.6-6.9	1.8	3.7	1.6-7.0
	1981-1985	11.2	11.8	8.8-15.4	6.4	14.9	10.1-21.2	4.8	9.5	6.0-14.2	3.8	3.8	2.3-5.9	2.8	6.1	3.3-10.3	1.0	2.0	0.6-4.7
	1986-1990	9.8	9.5	7.0-12.7	5.2	11.1	7.2-16.5	4.6	7.9	5.0-12.0	5.6	5.6	3.7-8.2	3.4	8.4	4.8-13.6	2.2	3.7	1.8-6.8
	1991-1995	11.8	11.0	8.3-14.2	7.0	14.6	10.1-20.4	4.8	7.8	4.9-11.7	5.0	4.7	3.0-6.9	2.0	4.4	2.1-8.2	3.0	5.0	2.8-8.4
	1996-2000	16.4	14.9	11.8-18.5	8.2	16.7	12.0-22.8	8.2	13.6	9.7-18.7	5.6	4.9	3.2-7.1	3.2	6.7	3.8-11.0	2.4	3.5	1.8-6.3
	2001-2005	17.0	15.0	12.0-18.6	10.6	19.8	14.8-26.0	6.4	10.5	7.2-15.0	4.2	3.6	2.2-5.5	2.6	5.0	2.7-8.7	1.6	2.7	1.1-5.4
	2006-2010	15.6	13.3	10.5-16.7	8.8	15.6	11.3-21.0	6.8	11.7	8.0-16.5	5.4	4.4	2.9-6.4	4.0	7.3	4.5-11.5	1.4	2.2	0.9-4.7
	2011-2015	19.0	14.9	12.0-18.4	12.6	20.6	15.7-26.7	6.4	9.7	6.5-14.0	2.8	2.1	1.1-3.7	1.8	3.1	1.4-6.0	1.0	1.4	0.4-3.6
2016-2020	22.2	18.2	14.8-22.2	12.4	22.2	16.8-28.9	9.8	14.2	10.4-19.2	3.8	2.7	1.6-4.4	1.8	2.5	1.1-5.2	2.0	2.9	1.4-5.6	
Brain and other central nervous system	1976-1980	5.2	5.5	3.5-8.1	3.6	8.0	4.7-12.8	1.6	3.1	1.3-6.3	3.2	3.6	2.0-6.0	2.4	5.9	3.0-10.4	0.8	1.4	0.4-3.8
	1981-1985	8.2	8.1	5.8-11.1	4.4	9.8	6.0-15.0	3.8	6.9	4.1-11.0	4.6	4.6	2.9-7.0	2.4	5.7	2.9-10.1	2.2	4.0	1.9-7.2
	1986-1990	6.6	6.4	4.4-9.0	4.6	9.1	5.7-13.8	2.0	3.5	1.6-6.5	6.6	6.5	4.5-9.2	3.6	7.3	4.3-11.7	3.0	5.5	3.0-9.1
	1991-1995	6.0	5.4	3.6-7.7	2.6	5.1	2.7-8.9	3.4	5.6	3.3-9.1	3.8	3.5	2.1-5.5	1.4	2.7	1.1-5.7	2.4	4.2	2.1-7.4
	1996-2000	8.4	7.9	5.7-10.7	4.6	8.9	5.7-13.5	3.8	6.9	4.2-10.9	4.0	3.6	2.2-5.6	1.8	3.6	1.6-6.8	2.2	3.7	1.8-6.7
	2001-2005	10.0	9.0	6.6-11.9	5.2	9.8	6.4-14.4	4.8	8.4	5.4-12.7	5.8	5.0	3.3-7.2	3.0	5.4	3.0-9.0	2.8	4.5	2.4-7.7
	2006-2010	7.4	6.7	4.7-9.4	4.2	7.8	4.8-12.1	3.2	6.0	3.4-9.9	4.6	3.8	2.4-5.8	2.0	3.4	1.6-6.5	2.6	4.3	2.3-7.6
	2011-2015	10.0	8.5	6.2-11.4	6.2	11.2	7.5-16.2	3.8	6.0	3.5-9.7	7.6	6.3	4.4-8.8	3.8	6.8	4.0-10.8	3.8	5.9	3.4-9.5
2016-2020	6.8	6.7	4.6-9.6	3.6	7.1	4.1-11.4	3.2	6.3	3.5-10.6	4.4	3.8	2.3-6.0	2.4	4.1	2.0-7.4	2.0	3.7	1.7-7.2	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Steuben**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	3.0	2.8	1.6-4.7	0.6	1.1	0.2-3.4	2.4	4.5	2.3-7.9	0.2	0.2	0.0-1.1	0.0	0.0	0.0-1.7	0.2	0.3	0.0-2.1
	1981-1985	2.8	2.7	1.5-4.5	0.8	1.5	0.4-4.1	2.0	3.8	1.8-7.0	0.0	0.0	0.0-0.8	0.0	0.0	0.0-1.7	0.0	0.0	0.0-1.5
	1986-1990	3.8	3.8	2.3-6.0	1.2	2.6	0.9-5.7	2.6	4.7	2.5-8.2	1.4	1.2	0.5-2.6	0.6	1.0	0.2-3.3	0.8	1.3	0.3-3.4
	1991-1995	4.8	4.8	3.1-7.2	0.6	1.6	0.3-4.5	4.2	8.1	5.0-12.4	0.2	0.2	0.0-1.0	0.0	0.0	0.0-1.6	0.2	0.3	0.0-1.9
	1996-2000	4.4	4.6	2.9-7.0	0.6	1.3	0.3-3.8	3.8	7.9	4.7-12.3	1.4	1.3	0.5-2.8	0.2	0.4	0.0-2.4	1.2	2.2	0.8-4.8
	2001-2005	4.6	4.6	2.9-6.9	0.8	1.5	0.4-4.0	3.8	7.4	4.4-11.6	0.4	0.3	0.0-1.3	0.2	0.4	0.0-2.2	0.2	0.3	0.0-1.9
	2006-2010	12.0	12.3	9.3-15.9	2.0	3.5	1.7-6.7	10.0	20.8	15.3-27.6	0.6	0.5	0.1-1.5	0.0	0.0	0.0-1.5	0.6	0.8	0.2-2.7
	2011-2015	14.6	14.3	11.1-18.2	3.6	6.4	3.7-10.4	11.0	22.3	16.6-29.3	1.4	1.0	0.4-2.1	0.4	0.5	0.1-2.2	1.0	1.4	0.4-3.6
2016-2020	12.8	12.0	9.1-15.6	4.0	7.4	4.4-11.7	8.8	16.7	11.8-22.9	0.4	0.3	0.0-1.2	0.4	0.6	0.1-2.6	0.0	0.0	0.0-1.6	
Hodgkin lymphoma	1976-1980	2.4	2.4	1.2-4.3	1.4	2.9	1.1-6.2	1.0	1.9	0.6-4.5	0.6	0.7	0.1-1.9	0.4	1.0	0.1-3.6	0.2	0.3	0.0-2.1
	1981-1985	3.2	3.3	1.8-5.4	1.2	2.9	1.0-6.4	2.0	3.5	1.7-6.6	0.0	0.0	0.0-0.8	0.0	0.0	0.0-1.7	0.0	0.0	0.0-1.5
	1986-1990	3.2	3.3	1.9-5.4	2.0	4.6	2.2-8.4	1.2	2.1	0.8-4.8	0.8	0.8	0.2-2.1	0.4	0.9	0.1-3.3	0.4	0.9	0.1-3.1
	1991-1995	2.6	2.7	1.4-4.6	1.4	2.9	1.1-5.9	1.2	2.4	0.9-5.3	0.2	0.2	0.0-1.1	0.2	0.4	0.0-2.2	0.0	0.0	0.0-1.4
	1996-2000	1.6	1.7	0.7-3.4	1.2	2.5	0.9-5.5	0.4	1.0	0.1-3.3	0.4	0.4	0.0-1.4	0.2	0.4	0.0-2.4	0.2	0.4	0.0-2.2
	2001-2005	4.0	4.1	2.5-6.3	2.2	4.4	2.2-7.9	1.8	3.8	1.7-7.2	0.8	0.7	0.2-1.8	0.4	0.7	0.1-2.8	0.4	0.6	0.1-2.4
	2006-2010	3.4	3.8	2.2-6.0	2.4	5.1	2.6-8.9	1.0	2.5	0.8-5.7	0.2	0.3	0.0-1.3	0.0	0.0	0.0-1.5	0.2	0.5	0.0-2.6
	2011-2015	3.4	3.4	1.9-5.5	1.8	3.5	1.6-6.8	1.6	3.4	1.4-6.8	0.2	0.2	0.0-1.2	0.0	0.0	0.0-1.6	0.2	0.4	0.0-2.3
2016-2020	1.6	1.7	0.7-3.4	1.0	2.0	0.6-4.9	0.6	1.3	0.2-3.9	0.4	0.4	0.0-1.6	0.2	0.3	0.0-2.0	0.2	0.6	0.0-3.0	
Non-Hodgkin lymphomas	1976-1980	12.4	12.9	9.8-16.6	7.4	17.4	12.0-24.1	5.0	9.3	6.0-13.8	6.4	6.8	4.6-9.6	4.2	10.6	6.5-16.3	2.2	3.9	1.9-7.1
	1981-1985	11.6	11.0	8.3-14.3	4.2	9.3	5.6-14.3	7.4	12.4	8.7-17.2	6.4	6.4	4.3-9.1	3.8	9.0	5.3-14.2	2.6	4.4	2.3-7.6
	1986-1990	17.2	16.5	13.1-20.4	10.2	23.2	17.0-30.9	7.0	12.0	8.3-16.9	8.8	8.5	6.2-11.5	4.4	10.8	6.6-16.5	4.4	7.6	4.7-11.6
	1991-1995	16.0	15.1	12.0-18.9	8.8	18.5	13.4-25.0	7.2	12.4	8.6-17.3	8.0	7.4	5.2-10.1	5.0	11.3	7.2-16.8	3.0	4.8	2.7-8.1
	1996-2000	23.0	20.8	17.1-25.0	11.8	23.6	17.9-30.5	11.2	18.0	13.6-23.6	9.4	8.2	6.0-11.0	5.0	9.7	6.3-14.4	4.4	6.6	4.1-10.2
	2001-2005	20.8	18.3	14.9-22.2	11.4	22.3	16.9-29.1	9.4	14.9	10.9-19.9	8.8	7.7	5.6-10.4	4.0	7.8	4.8-12.2	4.8	7.4	4.7-11.1
	2006-2010	22.4	18.3	15.0-22.1	11.4	20.1	15.1-26.2	11.0	16.5	12.4-21.7	6.6	5.1	3.5-7.3	3.4	5.8	3.4-9.5	3.2	4.6	2.6-7.7
	2011-2015	25.6	20.2	16.8-24.2	12.2	20.6	15.7-26.6	13.4	20.0	15.3-25.7	8.6	6.7	4.8-9.1	5.8	9.7	6.5-14.1	2.8	3.9	2.0-6.8
2016-2020	20.0	15.5	12.5-19.1	10.6	17.1	12.7-22.6	9.4	13.9	10.0-19.0	6.6	5.0	3.4-7.1	3.0	4.9	2.7-8.4	3.6	4.9	2.8-8.1	
Myeloma	1976-1980	5.8	6.1	4.1-8.8	1.6	4.0	1.7-7.9	4.2	7.9	4.8-12.1	3.4	3.6	2.1-5.8	0.8	2.2	0.6-5.5	2.6	4.6	2.4-7.9
	1981-1985	5.6	5.7	3.8-8.3	3.6	9.1	5.3-14.6	2.0	3.5	1.7-6.5	3.4	3.2	1.9-5.3	1.8	3.9	1.7-7.7	1.6	2.8	1.2-5.7
	1986-1990	5.2	5.0	3.2-7.3	2.4	5.6	2.8-9.8	2.8	4.7	2.5-8.0	5.4	5.1	3.3-7.4	2.8	6.9	3.7-11.6	2.6	4.2	2.3-7.4
	1991-1995	6.8	6.1	4.2-8.6	3.6	7.9	4.6-12.5	3.2	5.3	3.0-8.8	4.4	3.9	2.4-6.0	2.4	5.5	2.7-9.6	2.0	3.0	1.4-5.8
	1996-2000	8.0	6.9	4.9-9.4	4.2	8.4	5.1-12.9	3.8	5.8	3.5-9.2	4.8	4.2	2.7-6.3	2.8	5.6	3.0-9.5	2.0	3.2	1.5-6.0
	2001-2005	6.6	5.5	3.8-7.7	3.6	7.2	4.2-11.4	3.0	4.4	2.4-7.4	6.0	5.0	3.3-7.1	2.8	5.8	3.1-9.8	3.2	4.5	2.6-7.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Steuben**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	6.4	5.0	3.4-7.1	3.2	5.5	3.1-9.1	3.2	4.4	2.5-7.3	4.0	3.2	1.9-5.0	2.0	3.7	1.7-6.9	2.0	2.5	1.2-4.9
	2011-2015	8.6	6.7	4.8-9.2	5.2	8.4	5.4-12.6	3.4	5.1	2.9-8.5	4.0	3.0	1.8-4.7	2.0	3.3	1.5-6.2	2.0	2.7	1.3-5.2
	2016-2020	6.8	5.1	3.5-7.4	4.2	6.8	4.1-10.6	2.6	3.8	2.0-6.9	2.6	1.8	0.9-3.2	1.4	2.2	0.9-4.8	1.2	1.4	0.5-3.5
Leukemias	1976-1980	14.0	14.8	11.4-18.7	8.4	19.9	14.1-27.2	5.6	11.1	7.3-16.2	10.0	10.6	7.8-14.0	6.6	15.6	10.5-22.2	3.4	6.7	3.9-10.8
	1981-1985	10.6	10.8	8.0-14.1	5.2	13.4	8.6-19.7	5.4	9.6	6.3-14.1	8.4	8.7	6.2-11.8	3.8	9.6	5.7-15.2	4.6	8.3	5.2-12.6
	1986-1990	13.0	12.7	9.7-16.2	8.2	19.1	13.5-26.2	4.8	9.0	5.7-13.5	8.0	7.7	5.5-10.5	5.4	13.1	8.4-19.2	2.6	4.5	2.4-7.8
	1991-1995	17.6	16.3	13.1-20.2	9.8	20.6	15.1-27.5	7.8	13.2	9.3-18.3	11.6	10.5	7.9-13.6	6.2	13.0	8.7-18.6	5.4	9.1	5.9-13.3
	1996-2000	17.8	16.3	13.1-20.1	8.6	17.7	12.8-24.0	9.2	15.5	11.2-20.8	8.8	7.9	5.7-10.6	5.4	11.3	7.4-16.5	3.4	5.1	3.0-8.4
	2001-2005	17.6	15.3	12.2-18.9	12.0	23.1	17.6-29.8	5.6	9.0	5.9-13.1	8.4	7.2	5.2-9.8	5.0	9.9	6.4-14.7	3.4	5.4	3.1-8.8
	2006-2010	14.2	12.1	9.4-15.4	6.2	11.0	7.4-15.8	8.0	13.5	9.5-18.6	8.0	6.6	4.7-9.0	5.2	9.4	6.1-13.9	2.8	4.3	2.3-7.5
	2011-2015	16.8	12.9	10.2-16.1	10.2	16.7	12.3-22.2	6.6	9.7	6.5-13.9	8.6	6.7	4.8-9.1	4.2	7.2	4.4-11.2	4.4	6.3	3.8-9.8
2016-2020	21.0	15.2	12.3-18.6	13.0	19.0	14.6-24.6	8.0	11.6	8.2-16.2	7.8	5.7	4.0-7.9	3.8	6.0	3.6-9.5	4.0	5.4	3.2-8.7	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Suffolk**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	3,973.8	411.9	406.1-417.9	1,945.6	491.5	480.9-502.2	2,028.2	371.1	363.9-378.5	2,178.8	231.7	227.3-236.2	1,145.2	299.4	291.0-307.9	1,033.6	191.6	186.4-197.0
	1981-1985	4,662.6	437.6	431.9-443.4	2,248.8	514.7	504.5-525.0	2,413.8	400.3	393.2-407.6	2,469.0	238.3	234.0-242.6	1,257.0	303.5	295.5-311.7	1,212.0	202.3	197.2-207.5
	1986-1990	5,437.2	472.2	466.6-478.0	2,599.0	546.2	536.2-556.3	2,838.2	437.4	430.2-444.7	2,726.4	243.9	239.8-248.1	1,386.4	310.4	302.6-318.3	1,340.0	206.8	201.8-211.8
	1991-1995	6,071.0	491.2	485.7-496.8	3,007.2	575.2	565.6-584.9	3,063.8	443.4	436.4-450.5	2,824.4	235.8	231.9-239.8	1,425.2	297.3	290.1-304.7	1,399.2	201.4	196.7-206.2
	1996-2000	7,186.6	535.0	529.5-540.6	3,610.4	621.1	611.8-630.5	3,576.2	478.5	471.5-485.5	2,753.6	210.3	206.8-213.8	1,345.0	251.9	245.7-258.3	1,408.6	185.9	181.6-190.3
	2001-2005	8,085.4	544.2	538.9-549.6	4,079.0	630.6	621.8-639.5	4,006.4	487.2	480.4-494.0	2,840.2	194.6	191.4-197.9	1,396.4	232.2	226.7-237.9	1,443.8	172.0	168.0-176.0
	2006-2010	8,885.2	548.3	543.2-553.5	4,497.2	623.9	615.6-632.3	4,388.0	496.9	490.3-503.7	2,794.0	173.9	171.0-176.8	1,367.0	201.7	196.9-206.6	1,427.0	156.0	152.4-159.7
	2011-2015	9,311.8	525.1	520.2-530.0	4,679.4	581.0	573.4-588.6	4,632.4	488.7	482.3-495.2	2,777.6	154.9	152.3-157.6	1,394.8	181.5	177.2-185.9	1,382.8	137.2	133.9-140.5
	2016-2020	9,735.8	508.1	503.5-512.8	4,924.8	554.3	547.2-561.5	4,811.0	478.5	472.2-484.8	2,679.2	136.1	133.7-138.5	1,339.8	156.8	153.0-160.7	1,339.4	122.0	119.1-125.1
Oral cavity and pharynx	1976-1980	114.8	11.3	10.4-12.3	72.2	16.3	14.5-18.2	42.6	7.7	6.7-8.8	49.2	5.0	4.4-5.6	31.6	7.5	6.3-8.8	17.6	3.2	2.5-3.9
	1981-1985	121.4	11.0	10.1-11.9	83.4	17.2	15.5-19.1	38.0	6.3	5.4-7.2	44.2	4.2	3.6-4.8	27.4	6.3	5.2-7.5	16.8	2.8	2.2-3.5
	1986-1990	129.8	11.0	10.1-11.9	81.2	15.5	13.9-17.1	48.6	7.4	6.5-8.4	49.2	4.3	3.8-4.9	31.6	6.4	5.4-7.5	17.6	2.8	2.2-3.4
	1991-1995	129.0	10.3	9.5-11.1	85.4	15.3	13.9-16.9	43.6	6.3	5.5-7.2	36.6	3.0	2.6-3.5	23.2	4.6	3.7-5.6	13.4	1.9	1.5-2.5
	1996-2000	136.8	10.1	9.4-10.9	90.4	15.0	13.6-16.5	46.4	6.2	5.5-7.1	31.8	2.4	2.0-2.8	19.2	3.4	2.7-4.2	12.6	1.7	1.3-2.2
	2001-2005	139.2	9.2	8.6-10.0	97.0	14.3	13.0-15.6	42.2	5.0	4.4-5.8	31.0	2.1	1.8-2.5	21.2	3.3	2.7-4.0	9.8	1.1	0.8-1.5
	2006-2010	175.8	10.5	9.8-11.2	120.0	15.4	14.2-16.7	55.8	6.4	5.6-7.2	28.4	1.7	1.5-2.0	19.2	2.8	2.2-3.4	9.2	1.0	0.7-1.3
	2011-2015	210.2	11.5	10.8-12.2	149.4	17.5	16.3-18.9	60.8	6.3	5.6-7.1	33.4	1.8	1.6-2.1	22.2	2.8	2.3-3.4	11.2	1.1	0.8-1.4
	2016-2020	237.4	12.0	11.4-12.8	161.8	17.6	16.4-18.9	75.6	7.3	6.5-8.1	37.2	2.0	1.7-2.3	25.4	3.0	2.5-3.5	11.8	1.1	0.9-1.5
Esophagus	1976-1980	42.2	4.6	4.0-5.3	32.8	8.5	7.1-10.0	9.4	1.8	1.4-2.5	39.8	4.3	3.7-5.0	30.6	7.8	6.5-9.3	9.2	1.8	1.3-2.4
	1981-1985	49.4	4.6	4.1-5.3	34.2	7.5	6.4-8.8	15.2	2.5	2.0-3.2	45.6	4.3	3.8-4.9	32.0	7.1	6.0-8.4	13.6	2.3	1.8-2.9
	1986-1990	52.0	4.6	4.0-5.2	36.8	7.5	6.4-8.8	15.2	2.3	1.8-2.9	47.2	4.2	3.7-4.8	34.2	7.3	6.2-8.5	13.0	2.0	1.5-2.5
	1991-1995	58.2	4.7	4.2-5.3	40.2	7.5	6.5-8.7	18.0	2.6	2.1-3.2	57.2	4.7	4.1-5.2	42.2	7.9	6.8-9.1	15.0	2.2	1.7-2.7
	1996-2000	58.6	4.4	3.9-4.9	38.6	6.5	5.6-7.5	20.0	2.6	2.1-3.2	56.4	4.2	3.8-4.8	38.8	6.6	5.7-7.6	17.6	2.3	1.8-2.8
	2001-2005	80.2	5.4	4.9-6.0	62.6	9.6	8.6-10.8	17.6	2.1	1.7-2.6	70.2	4.8	4.3-5.3	56.0	8.8	7.8-9.9	14.2	1.6	1.3-2.1
	2006-2010	92.4	5.7	5.2-6.3	69.2	9.7	8.7-10.8	23.2	2.6	2.1-3.1	76.8	4.8	4.3-5.3	59.2	8.3	7.4-9.4	17.6	1.9	1.6-2.4
	2011-2015	94.2	5.1	4.7-5.6	72.2	8.7	7.8-9.6	22.0	2.2	1.8-2.7	71.4	3.9	3.5-4.3	55.6	6.9	6.1-7.7	15.8	1.6	1.2-1.9
	2016-2020	89.2	4.4	4.0-4.8	70.2	7.5	6.7-8.4	19.0	1.7	1.4-2.1	71.4	3.5	3.1-3.9	56.6	6.2	5.5-7.0	14.8	1.3	1.0-1.7
Stomach	1976-1980	110.2	12.3	11.2-13.4	69.6	18.9	16.8-21.2	40.6	7.8	6.7-8.9	82.0	9.4	8.4-10.3	49.8	13.8	12.0-15.8	32.2	6.3	5.4-7.4
	1981-1985	125.4	12.3	11.3-13.3	74.8	17.8	15.9-19.8	50.6	8.5	7.5-9.6	90.6	9.0	8.2-9.9	55.8	13.5	11.9-15.3	34.8	5.9	5.1-6.9
	1986-1990	136.6	12.4	11.5-13.4	85.2	19.3	17.4-21.4	51.4	7.9	7.0-9.0	94.2	8.7	7.9-9.5	59.0	13.9	12.3-15.7	35.2	5.4	4.6-6.3
	1991-1995	121.6	10.2	9.4-11.0	75.0	14.8	13.3-16.5	46.6	6.7	5.9-7.6	76.8	6.6	5.9-7.3	46.4	9.8	8.5-11.2	30.4	4.3	3.7-5.1
	1996-2000	133.8	10.2	9.4-11.0	87.2	15.8	14.3-17.5	46.6	6.0	5.3-6.9	72.4	5.6	5.0-6.2	42.8	8.0	6.9-9.2	29.6	3.8	3.2-4.4
	2001-2005	133.8	9.1	8.5-9.9	82.0	12.9	11.7-14.3	51.8	6.2	5.5-7.0	63.8	4.4	3.9-4.9	35.6	5.9	5.0-6.9	28.2	3.3	2.8-3.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Suffolk**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	119.0	7.4	6.8-8.0	76.8	10.9	9.8-12.1	42.2	4.7	4.0-5.3	51.4	3.2	2.8-3.6	32.4	4.8	4.0-5.6	19.0	2.1	1.7-2.5
	2011-2015	137.0	7.7	7.2-8.3	87.6	11.0	10.0-12.1	49.4	5.2	4.6-5.9	52.2	2.9	2.6-3.3	33.6	4.2	3.6-4.9	18.6	1.9	1.6-2.4
	2016-2020	146.6	7.6	7.0-8.2	85.2	9.6	8.7-10.6	61.4	6.0	5.3-6.7	54.0	2.7	2.4-3.1	29.0	3.3	2.8-3.9	25.0	2.3	1.9-2.8
Colorectal	1976-1980	628.8	69.2	66.7-71.7	314.0	84.3	79.9-88.9	314.8	59.7	56.7-62.7	330.8	37.0	35.2-38.9	164.0	45.2	41.9-48.7	166.8	31.8	29.7-34.1
	1981-1985	747.4	73.6	71.2-76.0	372.8	89.8	85.5-94.3	374.6	63.0	60.2-66.0	347.8	35.2	33.5-36.9	171.8	43.9	40.8-47.2	176.0	29.9	28.0-32.0
	1986-1990	815.6	73.6	71.3-75.9	412.2	89.7	85.6-94.0	403.4	62.5	59.8-65.3	383.6	35.3	33.8-37.0	197.8	46.0	43.0-49.2	185.8	28.7	26.9-30.6
	1991-1995	775.6	64.8	62.8-66.9	395.0	79.7	76.0-83.5	380.6	54.7	52.3-57.3	351.6	30.1	28.7-31.6	175.4	38.0	35.4-40.7	176.2	25.1	23.5-26.9
	1996-2000	869.8	66.3	64.3-68.3	448.6	80.2	76.8-83.7	421.2	55.6	53.2-58.0	317.8	24.6	23.4-25.9	157.8	30.1	27.9-32.3	160.0	20.7	19.3-22.2
	2001-2005	868.6	59.3	57.5-61.1	430.0	68.4	65.5-71.4	438.6	52.3	50.1-54.6	298.8	20.6	19.6-21.7	145.4	24.2	22.4-26.1	153.4	17.9	16.6-19.2
	2006-2010	790.0	49.1	47.5-50.6	390.0	55.4	53.0-58.0	400.0	43.9	42.0-45.9	271.6	16.9	16.0-17.9	135.6	20.1	18.6-21.7	136.0	14.3	13.3-15.5
	2011-2015	720.6	40.7	39.3-42.0	386.2	48.9	46.7-51.2	334.4	34.1	32.5-35.8	233.0	12.8	12.1-13.6	124.2	16.2	14.9-17.5	108.8	10.5	9.6-11.4
	2016-2020	755.2	39.5	38.2-40.8	398.0	45.6	43.5-47.7	357.2	34.5	32.9-36.2	227.2	11.6	11.0-12.4	117.4	13.9	12.8-15.1	109.8	9.9	9.1-10.8
Colon excluding rectum	1976-1980	452.4	50.3	48.2-52.4	214.0	58.5	54.7-62.3	238.4	45.3	42.8-48.0	272.8	30.5	28.9-32.2	129.4	35.7	32.8-38.8	143.4	27.3	25.3-29.4
	1981-1985	546.2	54.2	52.1-56.3	261.4	63.5	59.9-67.3	284.8	48.1	45.6-50.7	298.6	30.1	28.6-31.7	146.0	36.9	34.1-39.9	152.6	25.9	24.1-27.8
	1986-1990	580.0	52.8	50.9-54.8	286.4	63.3	59.8-66.9	293.6	45.5	43.2-47.9	337.8	31.1	29.6-32.6	172.6	40.1	37.3-43.0	165.2	25.5	23.8-27.3
	1991-1995	556.2	46.8	45.0-48.6	279.0	57.5	54.3-60.7	277.2	39.8	37.7-42.0	306.0	26.2	24.9-27.6	152.2	33.2	30.7-35.7	153.8	22.0	20.4-23.6
	1996-2000	636.6	48.8	47.1-50.5	316.0	57.6	54.7-60.6	320.6	42.2	40.2-44.3	282.4	21.9	20.8-23.1	138.0	26.5	24.5-28.7	144.4	18.7	17.3-20.1
	2001-2005	635.0	43.5	42.0-45.0	300.8	48.5	46.0-51.0	334.2	39.8	37.9-41.7	255.0	17.6	16.7-18.6	121.0	20.3	18.7-22.0	134.0	15.7	14.5-16.9
	2006-2010	575.8	35.9	34.6-37.3	276.4	39.9	37.8-42.1	299.4	32.7	31.0-34.4	227.0	14.1	13.3-15.0	110.6	16.5	15.1-17.9	116.4	12.2	11.2-13.3
	2011-2015	525.2	29.7	28.6-30.9	269.6	34.5	32.6-36.4	255.6	26.1	24.7-27.6	197.4	10.9	10.2-11.6	103.0	13.5	12.3-14.8	94.4	9.0	8.2-9.9
	2016-2020	535.4	28.0	26.9-29.1	266.6	30.8	29.1-32.5	268.8	25.7	24.3-27.2	181.4	9.3	8.7-9.9	91.4	10.8	9.8-11.9	90.0	8.1	7.4-9.0
Rectum & rectosigmoid	1976-1980	176.4	18.9	17.7-20.3	100.0	25.9	23.5-28.4	76.4	14.3	12.9-15.9	58.0	6.5	5.8-7.3	34.6	9.5	8.0-11.1	23.4	4.5	3.7-5.4
	1981-1985	201.2	19.4	18.2-20.6	111.4	26.3	24.0-28.8	89.8	14.9	13.6-16.4	49.2	5.1	4.5-5.8	25.8	7.0	5.7-8.4	23.4	4.0	3.3-4.8
	1986-1990	235.6	20.8	19.6-22.0	125.8	26.4	24.3-28.7	109.8	17.0	15.6-18.5	45.8	4.3	3.7-4.9	25.2	6.0	4.9-7.2	20.6	3.2	2.6-3.9
	1991-1995	219.4	18.1	17.0-19.2	116.0	22.2	20.4-24.2	103.4	14.9	13.7-16.3	45.6	3.9	3.4-4.4	23.2	4.8	3.9-5.9	22.4	3.2	2.6-3.8
	1996-2000	233.2	17.5	16.5-18.6	132.6	22.6	20.9-24.4	100.6	13.4	12.2-14.6	35.4	2.7	2.4-3.2	19.8	3.5	2.9-4.3	15.6	2.0	1.6-2.6
	2001-2005	233.6	15.8	14.9-16.7	129.2	19.9	18.4-21.6	104.4	12.5	11.5-13.7	43.8	3.0	2.6-3.4	24.4	3.9	3.2-4.7	19.4	2.2	1.8-2.7
	2006-2010	214.2	13.1	12.4-14.0	113.6	15.5	14.3-16.9	100.6	11.3	10.3-12.3	44.6	2.8	2.5-3.2	25.0	3.6	3.0-4.4	19.6	2.1	1.7-2.6
	2011-2015	195.4	10.9	10.3-11.7	116.6	14.4	13.2-15.7	78.8	8.0	7.2-8.9	35.6	2.0	1.7-2.3	21.2	2.7	2.2-3.2	14.4	1.4	1.1-1.8
	2016-2020	219.8	11.5	10.8-12.2	131.4	14.8	13.6-16.0	88.4	8.8	8.0-9.7	45.8	2.3	2.0-2.7	26.0	3.1	2.6-3.7	19.8	1.8	1.4-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Suffolk**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	30.0	3.0	2.6-3.6	19.0	4.4	3.5-5.5	11.0	2.0	1.5-2.6	27.0	2.8	2.3-3.3	16.2	3.8	3.0-4.8	10.8	2.0	1.5-2.6
	1981-1985	30.8	3.1	2.6-3.6	17.8	4.2	3.3-5.3	13.0	2.2	1.7-2.8	26.6	2.7	2.2-3.2	15.0	3.7	2.8-4.7	11.6	2.0	1.5-2.6
	1986-1990	36.6	3.2	2.8-3.7	21.4	4.5	3.6-5.4	15.2	2.3	1.8-2.9	36.0	3.2	2.7-3.7	21.2	4.6	3.7-5.6	14.8	2.3	1.8-2.9
	1991-1995	52.2	4.3	3.8-4.9	35.2	7.2	6.1-8.5	17.0	2.4	1.9-3.0	48.6	4.1	3.6-4.6	30.2	6.3	5.2-7.4	18.4	2.6	2.1-3.2
	1996-2000	56.8	4.3	3.8-4.8	38.0	6.4	5.5-7.4	18.8	2.5	2.0-3.0	50.6	3.9	3.4-4.4	30.6	5.4	4.6-6.3	20.0	2.6	2.1-3.2
	2001-2005	76.2	5.1	4.6-5.6	54.4	8.0	7.0-9.0	21.8	2.6	2.2-3.2	63.2	4.3	3.8-4.8	39.0	6.1	5.2-7.0	24.2	2.9	2.4-3.5
	2006-2010	113.6	6.8	6.2-7.4	83.6	10.9	9.8-12.0	30.0	3.3	2.8-3.9	77.6	4.7	4.3-5.2	49.8	6.7	5.8-7.6	27.8	3.1	2.6-3.6
	2011-2015	149.6	8.1	7.5-8.7	108.6	12.7	11.6-13.8	41.0	4.1	3.6-4.8	106.6	5.9	5.4-6.4	73.2	9.0	8.1-10.0	33.4	3.3	2.8-3.9
	2016-2020	155.6	7.7	7.1-8.3	109.8	11.8	10.8-12.9	45.8	4.2	3.7-4.9	98.2	4.9	4.5-5.4	64.6	7.2	6.4-8.0	33.6	3.0	2.6-3.5
Pancreas	1976-1980	123.2	13.5	12.4-14.6	61.0	16.1	14.2-18.1	62.2	11.7	10.4-13.1	119.4	12.9	11.9-14.1	61.6	16.2	14.3-18.2	57.8	10.8	9.6-12.1
	1981-1985	127.6	12.5	11.5-13.5	56.4	13.3	11.7-15.0	71.2	12.0	10.8-13.3	121.8	11.9	11.0-12.9	54.4	12.7	11.1-14.4	67.4	11.3	10.2-12.6
	1986-1990	145.6	13.2	12.2-14.2	69.6	15.4	13.7-17.2	76.0	11.7	10.5-12.9	141.4	12.9	11.9-13.9	67.8	14.9	13.3-16.7	73.6	11.4	10.2-12.6
	1991-1995	163.0	13.6	12.7-14.6	77.2	15.2	13.6-16.8	85.8	12.4	11.2-13.6	151.2	12.7	11.8-13.7	69.0	13.8	12.3-15.5	82.2	11.9	10.7-13.1
	1996-2000	168.6	12.8	12.0-13.7	79.2	14.3	12.9-15.8	89.4	11.8	10.7-13.0	155.0	11.9	11.1-12.8	72.6	13.4	12.0-14.9	82.4	10.9	9.8-12.0
	2001-2005	208.0	14.2	13.4-15.1	100.4	16.0	14.6-17.4	107.6	12.9	11.8-14.0	180.4	12.4	11.6-13.3	87.0	14.2	12.8-15.6	93.4	11.1	10.1-12.1
	2006-2010	242.8	15.1	14.3-16.0	114.6	16.3	14.9-17.7	128.2	14.0	12.9-15.2	193.8	12.1	11.3-12.8	93.6	13.3	12.1-14.6	100.2	10.8	9.9-11.8
	2011-2015	275.2	15.3	14.5-16.1	146.8	18.4	17.1-19.9	128.4	12.7	11.7-13.8	210.4	11.6	10.9-12.4	110.2	13.9	12.7-15.1	100.2	9.7	8.9-10.6
	2016-2020	315.2	16.0	15.2-16.9	159.2	18.0	16.7-19.3	156.0	14.4	13.4-15.5	221.8	11.3	10.6-12.0	110.6	12.6	11.6-13.8	111.2	10.1	9.3-11.0
Larynx	1976-1980	58.4	5.7	5.1-6.4	46.8	10.6	9.2-12.1	11.6	2.0	1.5-2.6	20.0	2.1	1.7-2.5	15.4	3.8	3.0-4.8	4.6	0.9	0.5-1.3
	1981-1985	66.6	6.0	5.4-6.7	54.2	11.2	9.8-12.7	12.4	2.1	1.6-2.6	24.4	2.2	1.9-2.7	19.0	4.2	3.3-5.2	5.4	0.9	0.6-1.3
	1986-1990	77.0	6.4	5.7-7.1	57.6	10.9	9.6-12.3	19.4	3.0	2.4-3.6	20.0	1.7	1.4-2.1	16.4	3.4	2.7-4.3	3.6	0.6	0.3-0.9
	1991-1995	67.0	5.3	4.8-5.9	52.0	9.3	8.2-10.5	15.0	2.2	1.7-2.7	22.4	1.9	1.5-2.2	18.0	3.7	3.0-4.6	4.4	0.6	0.4-1.0
	1996-2000	68.4	5.0	4.5-5.6	51.6	8.7	7.6-9.8	16.8	2.3	1.8-2.8	21.2	1.6	1.3-1.9	15.8	2.7	2.1-3.4	5.4	0.7	0.5-1.1
	2001-2005	58.6	3.9	3.5-4.4	47.4	7.2	6.3-8.3	11.2	1.4	1.0-1.8	20.8	1.4	1.2-1.7	17.2	2.8	2.2-3.5	3.6	0.4	0.2-0.7
	2006-2010	60.4	3.7	3.2-4.1	44.2	6.0	5.2-6.8	16.2	1.8	1.5-2.3	18.4	1.1	0.9-1.4	14.0	2.0	1.5-2.5	4.4	0.5	0.3-0.7
	2011-2015	62.4	3.4	3.0-3.8	50.0	6.0	5.3-6.9	12.4	1.3	1.0-1.6	16.0	0.9	0.7-1.1	12.8	1.6	1.2-2.0	3.2	0.3	0.2-0.5
	2016-2020	58.0	2.9	2.5-3.2	46.6	5.1	4.5-5.8	11.4	1.1	0.8-1.4	15.6	0.8	0.6-1.0	12.2	1.4	1.0-1.8	3.4	0.3	0.2-0.5
Lung and bronchus	1976-1980	616.8	62.8	60.6-65.1	434.4	106.1	101.4-111.0	182.4	32.9	30.8-35.2	525.0	53.9	51.8-56.1	375.0	93.5	89.0-98.1	150.0	27.1	25.2-29.2
	1981-1985	755.0	70.3	68.1-72.6	494.2	111.7	107.1-116.5	260.8	43.1	40.8-45.5	640.0	60.2	58.1-62.3	417.2	96.5	92.2-101.0	222.8	36.8	34.7-39.1
	1986-1990	866.8	74.8	72.6-77.1	521.8	108.7	104.3-113.2	345.0	52.9	50.5-55.5	724.8	63.3	61.2-65.4	446.0	94.8	90.7-99.1	278.8	43.0	40.8-45.3
	1991-1995	934.2	75.9	73.7-78.1	517.0	99.5	95.5-103.6	417.2	60.5	57.9-63.2	787.6	64.7	62.7-66.8	457.6	90.8	86.9-94.8	330.0	47.7	45.4-50.0
	1996-2000	987.2	74.0	72.0-76.1	520.8	91.7	88.1-95.4	466.4	62.4	59.9-65.0	734.2	55.7	53.9-57.5	395.8	71.7	68.4-75.0	338.4	45.1	43.0-47.4
	2001-2005	1,098.6	74.6	72.6-76.6	551.2	87.7	84.4-91.1	547.4	66.4	63.9-69.0	777.8	53.2	51.5-54.9	408.2	66.2	63.3-69.2	369.6	44.5	42.5-46.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Suffolk**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	1,199.4	74.7	72.8-76.6	572.2	82.4	79.3-85.5	627.2	70.0	67.5-72.5	761.4	47.5	46.0-49.0	369.6	53.9	51.5-56.5	391.8	43.4	41.5-45.4
	2011-2015	1,202.4	67.4	65.6-69.1	560.8	71.7	69.0-74.5	641.6	64.9	62.6-67.2	724.8	40.6	39.3-42.0	351.6	45.4	43.3-47.6	373.2	37.3	35.6-39.1
	2016-2020	1,249.6	62.9	61.4-64.5	573.6	65.6	63.2-68.1	676.0	61.6	59.5-63.8	594.0	29.9	28.8-31.0	291.2	33.8	32.1-35.7	302.8	27.2	25.8-28.6
Melanoma of the skin	1976-1980	85.0	8.0	7.3-8.8	43.2	9.2	8.0-10.6	41.8	7.3	6.3-8.3	33.2	3.3	2.8-3.9	18.8	4.3	3.4-5.3	14.4	2.6	2.0-3.3
	1981-1985	111.6	9.9	9.1-10.7	56.8	11.5	10.1-13.0	54.8	9.0	7.9-10.1	35.8	3.3	2.8-3.8	19.6	4.1	3.3-5.1	16.2	2.7	2.2-3.4
	1986-1990	150.4	12.5	11.6-13.4	83.2	16.0	14.4-17.7	67.2	10.2	9.2-11.4	42.4	3.6	3.2-4.2	25.4	5.0	4.1-6.0	17.0	2.6	2.1-3.3
	1991-1995	136.2	10.7	9.9-11.5	85.2	15.0	13.5-16.5	51.0	7.3	6.4-8.3	43.0	3.5	3.0-4.0	28.2	5.5	4.5-6.5	14.8	2.1	1.7-2.7
	1996-2000	191.6	14.1	13.2-15.0	109.2	18.3	16.7-19.9	82.4	11.1	10.0-12.2	41.4	3.1	2.7-3.6	26.2	4.6	3.9-5.5	15.2	2.0	1.6-2.5
	2001-2005	278.6	18.6	17.7-19.6	159.0	23.8	22.1-25.5	119.6	14.9	13.7-16.1	43.8	3.0	2.6-3.4	25.8	4.1	3.4-4.9	18.0	2.2	1.8-2.7
	2006-2010	402.4	25.1	24.0-26.3	237.6	33.0	31.1-35.0	164.8	19.4	18.1-20.8	53.8	3.3	3.0-3.8	34.6	4.9	4.2-5.7	19.2	2.1	1.7-2.6
	2011-2015	479.6	27.4	26.3-28.6	290.8	36.6	34.7-38.6	188.8	20.6	19.3-22.0	47.8	2.7	2.4-3.1	31.8	4.2	3.5-4.9	16.0	1.6	1.3-2.0
2016-2020	474.4	24.9	23.9-25.9	283.2	32.4	30.7-34.2	191.2	19.3	18.1-20.7	38.0	1.9	1.6-2.2	25.2	2.9	2.4-3.5	12.8	1.2	0.9-1.5	
Female breast	1976-1980							572.2	104.9	101.1-108.8							223.8	41.3	38.9-43.8
	1981-1985							663.8	110.8	107.0-114.7							243.4	40.7	38.4-43.0
	1986-1990							838.8	130.4	126.5-134.4							276.6	42.7	40.5-45.1
	1991-1995							932.2	135.8	131.9-139.7							260.0	37.6	35.6-39.7
	1996-2000							1,124.8	151.0	147.1-155.0							244.4	32.3	30.5-34.2
	2001-2005							1,175.8	142.8	139.2-146.5							237.6	28.2	26.6-29.8
	2006-2010							1,251.8	141.6	138.1-145.2							215.6	23.5	22.1-25.0
	2011-2015							1,322.0	140.4	136.9-143.9							201.8	20.2	19.0-21.5
2016-2020							1,371.2	139.9	136.4-143.4							191.2	17.8	16.6-19.0	
Cervix uteri	1976-1980							61.6	10.8	9.6-12.1							23.8	4.4	3.6-5.3
	1981-1985							55.6	9.1	8.1-10.3							19.4	3.2	2.6-3.9
	1986-1990							64.2	9.8	8.8-11.0							23.6	3.7	3.0-4.4
	1991-1995							68.2	9.8	8.8-10.9							18.2	2.6	2.1-3.2
	1996-2000							77.8	10.5	9.5-11.6							20.8	2.8	2.3-3.4
	2001-2005							61.8	7.7	6.9-8.7							21.8	2.7	2.2-3.2
	2006-2010							67.4	8.3	7.4-9.3							17.2	1.9	1.5-2.4
	2011-2015							57.8	7.1	6.3-8.0							19.8	2.2	1.8-2.7
2016-2020							46.0	5.5	4.8-6.4							13.4	1.5	1.1-1.9	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Suffolk**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							131.2	23.8	22.0-25.7							29.8	5.6	4.7-6.5
	1981-1985							161.6	26.4	24.6-28.3							33.8	5.6	4.8-6.5
	1986-1990							156.6	24.1	22.5-25.9							34.4	5.2	4.5-6.1
	1991-1995							185.8	27.1	25.4-28.9							37.6	5.4	4.6-6.2
	1996-2000							217.2	29.1	27.4-30.9							39.8	5.3	4.5-6.0
	2001-2005							267.8	32.2	30.5-34.0							40.4	4.8	4.2-5.6
	2006-2010							269.4	29.8	28.2-31.5							43.0	4.8	4.2-5.5
	2011-2015							322.0	32.4	30.8-34.1							55.6	5.4	4.8-6.1
	2016-2020							341.8	32.2	30.6-33.8							61.2	5.7	5.0-6.4
Ovary	1976-1980							85.4	15.4	14.0-16.9							56.2	10.3	9.1-11.6
	1981-1985							113.4	18.8	17.3-20.5							71.2	11.8	10.6-13.1
	1986-1990							109.8	17.0	15.6-18.5							66.8	10.3	9.2-11.5
	1991-1995							118.8	17.3	15.9-18.7							68.0	9.8	8.8-11.0
	1996-2000							119.2	16.1	14.8-17.4							68.8	9.2	8.2-10.2
	2001-2005							132.4	16.1	14.9-17.4							83.8	10.1	9.1-11.1
	2006-2010							131.2	14.8	13.7-16.0							72.6	8.1	7.3-9.0
	2011-2015							130.2	13.6	12.5-14.7							81.0	7.9	7.1-8.7
	2016-2020							121.2	12.2	11.2-13.2							71.8	6.5	5.8-7.2
Prostate	1976-1980				238.0	72.2	67.9-76.8							95.2	31.5	28.5-34.6			
	1981-1985				308.4	81.4	77.1-85.8							125.4	37.5	34.4-40.7			
	1986-1990				393.0	93.5	89.2-98.0							149.6	41.4	38.3-44.6			
	1991-1995				742.4	145.1	140.3-150.1							160.6	40.7	37.8-43.8			
	1996-2000				1,058.0	179.9	174.9-184.9							140.0	31.3	28.9-33.7			
	2001-2005				1,196.8	182.9	178.2-187.7							125.6	24.2	22.3-26.2			
	2006-2010				1,323.4	177.8	173.5-182.2							119.8	19.5	18.0-21.2			
	2011-2015				1,095.8	127.7	124.2-131.2							113.0	16.0	14.7-17.4			
	2016-2020				1,316.4	138.2	134.9-141.7							129.8	15.9	14.7-17.2			
Testis	1976-1980				24.4	3.9	3.2-4.7							4.0	0.7	0.4-1.1			
	1981-1985				30.8	4.5	3.8-5.3							2.4	0.4	0.2-0.7			
	1986-1990				39.2	5.6	4.8-6.4							4.0	0.6	0.3-0.9			
	1991-1995				41.6	6.0	5.2-6.9							2.4	0.4	0.2-0.7			
	1996-2000				51.6	7.3	6.5-8.3							1.2	0.2	0.1-0.4			
	2001-2005				49.6	7.0	6.1-7.9							2.4	0.3	0.2-0.6			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Suffolk**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				50.2	7.0	6.2-8.0							1.6	0.2	0.1-0.5			
	2011-2015				50.6	7.3	6.4-8.3							1.0	0.1	0.0-0.3			
	2016-2020				50.2	7.3	6.4-8.3							2.2	0.3	0.1-0.5			
Urinary bladder (incl. in situ)	1976-1980	215.4	23.4	22.0-24.8	163.0	43.5	40.4-46.9	52.4	9.9	8.7-11.1	57.2	6.6	5.9-7.4	40.8	12.3	10.5-14.2	16.4	3.2	2.5-3.9
	1981-1985	243.4	23.5	22.1-24.8	179.8	42.6	39.6-45.7	63.6	10.7	9.5-11.9	63.6	6.7	6.0-7.5	45.2	13.2	11.5-15.2	18.4	3.2	2.5-3.9
	1986-1990	266.8	23.4	22.2-24.7	197.0	41.9	39.2-44.8	69.8	10.7	9.6-11.9	58.6	5.6	5.0-6.3	41.4	10.8	9.3-12.4	17.2	2.7	2.1-3.3
	1991-1995	282.8	23.2	22.0-24.5	203.0	40.2	37.6-42.9	79.8	11.4	10.3-12.6	66.8	5.8	5.2-6.5	46.0	10.7	9.2-12.2	20.8	2.9	2.4-3.6
	1996-2000	377.2	28.5	27.2-29.8	279.2	50.2	47.5-53.0	98.0	13.0	11.9-14.2	68.8	5.4	4.8-6.0	50.2	10.3	9.0-11.7	18.6	2.4	1.9-2.9
	2001-2005	441.6	30.2	29.0-31.5	328.8	53.1	50.5-55.8	112.8	13.5	12.4-14.6	77.2	5.4	4.9-6.0	56.0	10.2	9.0-11.5	21.2	2.4	2.0-3.0
	2006-2010	458.2	28.6	27.4-29.8	347.8	50.7	48.3-53.2	110.4	12.2	11.2-13.3	81.2	5.1	4.6-5.6	58.2	9.2	8.1-10.3	23.0	2.4	2.0-2.9
	2011-2015	512.6	28.8	27.7-29.9	382.8	49.9	47.6-52.2	129.8	12.9	11.9-13.9	86.8	4.8	4.4-5.3	60.8	8.3	7.4-9.4	26.0	2.5	2.1-3.0
2016-2020	522.0	26.7	25.7-27.8	389.2	45.5	43.5-47.7	132.8	12.2	11.2-13.1	93.8	4.8	4.3-5.2	67.0	8.3	7.4-9.3	26.8	2.3	1.9-2.7	
Kidney and renal pelvis	1976-1980	74.4	7.7	6.9-8.5	44.0	10.8	9.3-12.5	30.4	5.6	4.7-6.5	41.4	4.3	3.7-4.9	25.8	6.5	5.3-7.8	15.6	2.8	2.2-3.5
	1981-1985	96.0	8.9	8.1-9.7	55.6	11.7	10.3-13.2	40.4	6.7	5.8-7.7	47.2	4.5	3.9-5.1	26.6	5.9	4.9-7.1	20.6	3.4	2.8-4.2
	1986-1990	134.8	11.6	10.7-12.5	82.6	16.1	14.5-17.8	52.2	8.1	7.1-9.1	55.0	4.8	4.2-5.4	32.6	6.4	5.4-7.5	22.4	3.4	2.8-4.1
	1991-1995	150.2	11.9	11.1-12.8	91.6	16.7	15.2-18.4	58.6	8.4	7.5-9.5	50.0	4.1	3.6-4.7	31.0	6.2	5.3-7.4	19.0	2.7	2.2-3.3
	1996-2000	184.4	13.6	12.7-14.5	116.2	19.2	17.7-20.9	68.2	9.1	8.2-10.2	56.2	4.2	3.7-4.7	35.6	6.3	5.4-7.4	20.6	2.7	2.2-3.3
	2001-2005	247.2	16.5	15.5-17.4	158.6	23.5	21.8-25.2	88.6	10.9	9.9-12.0	57.6	3.9	3.5-4.4	38.8	6.3	5.4-7.2	18.8	2.2	1.8-2.7
	2006-2010	310.8	19.1	18.2-20.1	193.0	25.9	24.3-27.7	117.8	13.5	12.4-14.6	57.0	3.5	3.1-4.0	33.0	4.8	4.1-5.6	24.0	2.6	2.1-3.1
	2011-2015	363.8	20.5	19.5-21.5	242.2	29.4	27.7-31.1	121.6	13.0	11.9-14.1	60.6	3.4	3.0-3.8	42.8	5.5	4.8-6.3	17.8	1.7	1.4-2.1
2016-2020	366.0	19.1	18.2-20.0	233.8	26.2	24.6-27.8	132.2	12.9	11.9-14.0	51.6	2.6	2.3-2.9	33.4	3.8	3.3-4.5	18.2	1.6	1.3-2.0	
Brain and other central nervous system	1976-1980	70.0	6.2	5.5-6.9	42.2	8.2	7.0-9.5	27.8	4.6	3.9-5.5	43.8	4.0	3.4-4.5	25.4	5.0	4.2-6.0	18.4	3.1	2.5-3.8
	1981-1985	78.4	6.6	5.9-7.3	40.4	7.3	6.3-8.5	38.0	6.0	5.1-6.9	51.6	4.4	3.9-5.0	26.2	4.8	4.0-5.8	25.4	4.1	3.4-4.8
	1986-1990	98.4	8.0	7.3-8.7	52.0	9.4	8.2-10.7	46.4	7.0	6.1-8.0	55.4	4.6	4.1-5.2	31.6	6.0	5.0-7.0	23.8	3.6	2.9-4.3
	1991-1995	101.4	7.9	7.2-8.6	55.8	9.5	8.4-10.8	45.6	6.6	5.8-7.5	59.2	4.7	4.2-5.3	33.8	6.2	5.2-7.2	25.4	3.7	3.0-4.3
	1996-2000	115.8	8.5	7.8-9.2	63.6	9.9	8.8-11.1	52.2	7.2	6.3-8.1	56.4	4.2	3.7-4.7	29.8	4.7	4.0-5.5	26.6	3.6	3.0-4.2
	2001-2005	114.6	7.8	7.1-8.4	65.0	9.3	8.3-10.4	49.6	6.3	5.5-7.1	63.2	4.2	3.8-4.7	35.2	5.1	4.4-6.0	28.0	3.4	2.9-4.1
	2006-2010	112.2	7.2	6.6-7.8	64.4	8.8	7.9-9.9	47.8	5.9	5.1-6.7	62.8	4.0	3.5-4.4	39.4	5.4	4.7-6.3	23.4	2.7	2.2-3.2
	2011-2015	123.4	7.4	6.8-8.1	69.8	8.8	7.9-9.8	53.6	6.2	5.4-7.0	73.0	4.1	3.7-4.5	40.4	4.9	4.2-5.6	32.6	3.4	2.9-4.0
2016-2020	124.6	7.3	6.7-7.9	65.2	8.0	7.1-8.9	59.4	6.6	5.9-7.5	81.4	4.2	3.8-4.6	45.2	5.0	4.4-5.7	36.2	3.5	3.0-4.0	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Suffolk**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	40.8	3.7	3.2-4.3	10.6	2.1	1.5-2.7	30.2	5.1	4.3-6.0	6.4	0.7	0.5-1.0	2.8	0.7	0.4-1.1	3.6	0.7	0.4-1.1
	1981-1985	41.4	3.5	3.0-4.0	10.8	1.9	1.4-2.5	30.6	4.8	4.1-5.7	4.6	0.5	0.3-0.7	1.6	0.4	0.2-0.8	3.0	0.5	0.3-0.8
	1986-1990	45.4	3.6	3.1-4.1	11.8	2.0	1.5-2.6	33.6	5.0	4.3-5.9	5.6	0.5	0.3-0.7	1.4	0.3	0.1-0.6	4.2	0.6	0.4-1.0
	1991-1995	68.6	5.1	4.6-5.7	21.6	3.4	2.8-4.1	47.0	6.7	5.9-7.6	6.8	0.6	0.4-0.8	2.8	0.5	0.3-0.9	4.0	0.6	0.4-0.9
	1996-2000	127.4	9.1	8.4-9.8	37.4	5.6	4.8-6.5	90.0	12.4	11.3-13.6	8.0	0.6	0.4-0.9	3.2	0.7	0.4-1.1	4.8	0.6	0.4-0.9
	2001-2005	199.8	13.2	12.4-14.0	53.4	7.4	6.5-8.4	146.4	18.7	17.4-20.2	8.4	0.6	0.4-0.8	3.8	0.6	0.4-1.0	4.6	0.5	0.3-0.8
	2006-2010	304.0	19.3	18.3-20.3	81.0	10.6	9.6-11.7	223.0	27.7	26.1-29.5	8.8	0.6	0.4-0.7	4.6	0.7	0.4-1.0	4.2	0.4	0.3-0.7
	2011-2015	384.4	23.8	22.7-25.0	101.6	12.6	11.5-13.8	282.8	34.9	33.0-36.8	11.8	0.7	0.5-0.9	6.2	0.8	0.5-1.1	5.6	0.5	0.3-0.8
2016-2020	342.8	21.4	20.4-22.5	103.8	12.5	11.4-13.7	239.0	30.3	28.5-32.1	11.4	0.6	0.4-0.8	4.0	0.5	0.3-0.7	7.4	0.7	0.5-1.0	
Hodgkin lymphoma	1976-1980	43.4	3.4	2.9-3.9	24.2	4.0	3.3-4.8	19.2	2.8	2.3-3.5	16.8	1.5	1.2-1.8	8.8	1.6	1.2-2.2	8.0	1.3	0.9-1.7
	1981-1985	52.0	3.9	3.4-4.4	25.6	3.9	3.3-4.7	26.4	3.8	3.2-4.6	14.2	1.1	0.9-1.4	6.8	1.2	0.8-1.7	7.4	1.1	0.8-1.5
	1986-1990	50.8	3.7	3.2-4.2	25.4	3.9	3.2-4.6	25.4	3.5	2.9-4.2	14.0	1.1	0.8-1.4	8.4	1.4	1.0-1.9	5.6	0.8	0.5-1.2
	1991-1995	53.0	4.0	3.5-4.5	27.8	4.3	3.6-5.1	25.2	3.6	3.0-4.3	10.6	0.8	0.6-1.1	5.8	1.0	0.6-1.4	4.8	0.7	0.4-1.0
	1996-2000	53.6	3.9	3.4-4.4	29.0	4.3	3.6-5.1	24.6	3.5	2.9-4.2	8.6	0.6	0.5-0.8	4.4	0.7	0.4-1.0	4.2	0.6	0.4-0.9
	2001-2005	56.4	4.0	3.5-4.5	28.6	4.1	3.4-4.8	27.8	3.9	3.3-4.6	5.2	0.4	0.2-0.5	2.4	0.4	0.2-0.6	2.8	0.4	0.2-0.6
	2006-2010	51.6	3.5	3.1-3.9	26.0	3.6	3.0-4.3	25.6	3.4	2.8-4.0	6.4	0.4	0.3-0.6	3.2	0.5	0.3-0.7	3.2	0.4	0.2-0.6
	2011-2015	53.4	3.5	3.1-4.0	31.4	4.2	3.5-4.9	22.0	2.9	2.4-3.5	4.4	0.3	0.2-0.4	2.2	0.3	0.1-0.5	2.2	0.2	0.1-0.4
2016-2020	53.8	3.4	3.0-3.9	27.8	3.5	3.0-4.2	26.0	3.4	2.8-4.0	4.6	0.3	0.2-0.4	2.2	0.3	0.1-0.5	2.4	0.3	0.1-0.5	
Non-Hodgkin Lymphomas	1976-1980	112.6	11.3	10.3-12.3	59.6	13.3	11.7-15.0	53.0	9.6	8.5-10.9	63.6	6.5	5.8-7.3	31.8	7.3	6.1-8.6	31.8	5.8	4.9-6.8
	1981-1985	137.2	12.3	11.4-13.2	70.8	13.9	12.4-15.5	66.4	10.8	9.6-12.0	77.2	7.1	6.4-7.9	40.6	8.6	7.4-10.0	36.6	6.1	5.2-7.0
	1986-1990	195.6	16.4	15.4-17.5	102.0	19.2	17.4-21.0	93.6	14.2	13.0-15.6	91.8	8.1	7.4-8.9	48.6	10.0	8.7-11.4	43.2	6.7	5.8-7.7
	1991-1995	219.8	17.4	16.4-18.5	120.2	20.9	19.2-22.8	99.6	14.4	13.1-15.7	104.4	8.6	7.8-9.3	48.6	9.0	7.9-10.3	55.8	8.0	7.1-9.0
	1996-2000	284.0	21.1	20.0-22.2	147.2	24.3	22.5-26.2	136.8	18.4	17.0-19.8	110.6	8.4	7.7-9.2	57.0	10.3	9.1-11.6	53.6	7.1	6.3-8.0
	2001-2005	337.2	22.8	21.7-23.9	180.4	27.5	25.7-29.4	156.8	19.2	17.8-20.6	105.4	7.3	6.7-7.9	59.0	9.6	8.5-10.8	46.4	5.5	4.8-6.3
	2006-2010	374.6	23.4	22.3-24.5	199.6	28.1	26.4-30.0	175.0	19.8	18.5-21.2	100.6	6.3	5.8-6.9	53.2	7.8	6.9-8.8	47.4	5.2	4.5-5.9
	2011-2015	439.2	25.2	24.1-26.3	246.0	31.5	29.7-33.3	193.2	20.2	18.9-21.5	96.4	5.5	5.0-6.0	57.2	7.7	6.8-8.6	39.2	3.9	3.3-4.5
2016-2020	427.8	22.6	21.7-23.7	232.6	27.2	25.6-28.8	195.2	19.1	17.9-20.4	90.8	4.6	4.2-5.1	51.4	6.1	5.4-6.9	39.4	3.6	3.1-4.2	
Myeloma	1976-1980	42.4	4.5	3.9-5.1	20.6	5.1	4.1-6.3	21.8	4.1	3.3-4.9	28.6	3.1	2.6-3.6	14.2	3.7	2.8-4.8	14.4	2.7	2.1-3.4
	1981-1985	44.0	4.2	3.6-4.8	25.0	5.8	4.7-7.0	19.0	3.1	2.5-3.8	32.8	3.2	2.7-3.7	17.2	4.1	3.2-5.2	15.6	2.6	2.0-3.2
	1986-1990	58.4	5.3	4.7-5.9	30.4	6.6	5.5-7.8	28.0	4.3	3.6-5.1	38.0	3.4	3.0-4.0	19.0	4.2	3.3-5.2	19.0	2.9	2.4-3.6
	1991-1995	59.2	4.9	4.3-5.5	31.2	6.1	5.2-7.2	28.0	4.0	3.4-4.7	42.6	3.6	3.1-4.1	20.2	4.2	3.4-5.2	22.4	3.2	2.6-3.8
	1996-2000	87.0	6.6	6.0-7.2	44.8	7.9	6.9-9.0	42.2	5.6	4.9-6.4	46.6	3.6	3.2-4.1	25.0	4.8	4.0-5.8	21.6	2.8	2.3-3.4
	2001-2005	101.8	6.9	6.3-7.5	57.0	8.9	7.8-10.0	44.8	5.4	4.7-6.1	48.8	3.4	3.0-3.9	26.6	4.5	3.8-5.4	22.2	2.7	2.2-3.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Suffolk**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	126.0	7.7	7.1-8.4	65.6	9.2	8.2-10.3	60.4	6.6	5.9-7.5	51.4	3.2	2.8-3.6	26.8	4.0	3.3-4.7	24.6	2.7	2.2-3.2
	2011-2015	161.0	8.9	8.3-9.6	89.6	11.1	10.1-12.3	71.4	7.2	6.4-8.0	53.0	2.9	2.6-3.3	28.2	3.6	3.0-4.3	24.8	2.4	2.0-2.9
	2016-2020	160.0	8.2	7.6-8.8	91.8	10.3	9.4-11.4	68.2	6.4	5.7-7.1	51.2	2.5	2.2-2.9	29.0	3.4	2.8-4.0	22.2	1.9	1.6-2.3
Leukemias	1976-1980	123.8	12.4	11.4-13.4	68.2	15.8	14.0-17.7	55.6	9.9	8.8-11.2	73.2	7.5	6.7-8.3	39.6	9.8	8.3-11.4	33.6	6.0	5.1-7.0
	1981-1985	124.0	11.5	10.6-12.5	75.2	16.3	14.6-18.2	48.8	8.0	7.1-9.1	83.2	7.8	7.1-8.6	48.4	10.8	9.4-12.4	34.8	5.7	4.9-6.7
	1986-1990	148.6	12.7	11.8-13.7	82.2	16.2	14.6-18.0	66.4	10.2	9.1-11.4	92.2	8.2	7.5-9.0	46.8	10.0	8.6-11.4	45.4	7.0	6.1-8.0
	1991-1995	166.0	13.5	12.5-14.4	92.8	17.7	16.0-19.4	73.2	10.6	9.6-11.8	94.6	7.9	7.2-8.6	53.4	11.0	9.6-12.4	41.2	5.9	5.1-6.8
	1996-2000	194.8	14.6	13.7-15.5	113.0	19.2	17.6-20.9	81.8	10.8	9.8-11.9	113.8	8.7	8.0-9.4	62.0	11.4	10.1-12.8	51.8	6.8	6.0-7.7
	2001-2005	247.4	16.9	16.0-17.9	139.4	21.8	20.2-23.6	108.0	13.3	12.2-14.5	106.2	7.4	6.8-8.0	61.2	10.3	9.2-11.6	45.0	5.4	4.7-6.1
	2006-2010	277.0	17.3	16.4-18.3	163.8	23.0	21.4-24.7	113.2	12.9	11.8-14.0	109.0	6.9	6.4-7.5	65.2	9.8	8.7-10.9	43.8	4.8	4.2-5.5
	2011-2015	363.0	21.1	20.2-22.2	211.6	27.5	25.8-29.3	151.4	16.1	14.9-17.3	108.8	6.3	5.7-6.8	64.2	8.7	7.7-9.7	44.6	4.6	4.0-5.3
	2016-2020	350.4	18.9	18.0-19.9	206.6	24.5	23.0-26.1	143.8	14.5	13.4-15.6	109.0	5.7	5.2-6.2	64.2	7.9	7.0-8.8	44.8	4.1	3.6-4.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Sullivan**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	274.2	391.0	369.8-413.1	150.8	471.1	435.7-508.5	123.4	337.3	310.2-366.1	153.8	220.7	204.8-237.5	85.8	274.4	247.1-303.7	68.0	182.0	162.6-203.1
	1981-1985	335.0	476.1	452.9-500.2	183.8	566.8	528.9-606.7	151.2	414.7	384.6-446.5	172.6	244.8	228.3-262.3	99.6	311.9	283.7-342.1	73.0	194.2	174.3-216.0
	1986-1990	358.2	488.5	465.7-512.2	192.0	573.1	536.1-611.8	166.2	430.4	400.8-461.6	183.6	250.5	234.2-267.6	99.2	296.8	270.4-325.1	84.4	216.3	195.6-238.6
	1991-1995	403.6	521.0	498.2-544.5	217.2	607.5	570.9-645.9	186.4	462.0	432.3-493.2	195.0	253.3	237.4-269.9	108.0	306.1	280.1-333.8	87.0	211.0	191.4-232.2
	1996-2000	430.8	532.2	509.9-555.3	232.0	611.1	575.6-648.4	198.8	472.0	442.9-502.6	179.8	224.4	209.9-239.7	100.4	280.2	255.4-306.9	79.4	183.5	165.8-202.6
	2001-2005	441.2	509.3	488.1-531.2	226.0	544.7	512.6-578.4	215.2	485.4	456.6-515.7	181.4	211.8	198.1-226.2	97.4	246.7	224.5-270.5	84.0	184.0	166.8-202.7
	2006-2010	469.4	508.4	487.7-529.9	248.0	546.3	515.4-578.7	221.4	479.4	451.0-509.3	172.0	191.1	178.4-204.6	90.6	220.0	199.5-242.1	81.4	170.2	153.8-187.9
	2011-2015	451.0	459.7	440.3-479.8	237.6	492.1	463.5-522.1	213.4	438.6	411.6-466.9	165.6	168.0	156.5-180.2	91.6	196.7	178.4-216.4	74.0	146.0	131.1-162.3
	2016-2020	457.2	440.2	421.6-459.5	239.6	452.4	426.2-479.8	217.6	435.4	408.4-463.8	163.8	155.7	144.9-167.1	86.4	169.5	153.4-187.0	77.4	144.6	130.1-160.4
Oral cavity and pharynx	1976-1980	6.6	9.4	6.4-13.5	4.8	12.8	8.1-19.4	1.8	6.1	2.7-11.7	3.2	4.4	2.5-7.3	2.8	8.1	4.3-13.9	0.4	1.0	0.1-4.1
	1981-1985	8.6	12.1	8.7-16.6	5.4	16.2	10.4-24.0	3.2	8.9	5.0-14.9	3.2	4.3	2.4-7.2	2.6	7.0	3.6-12.2	0.6	1.9	0.3-5.8
	1986-1990	10.4	15.4	11.4-20.4	7.0	22.0	15.1-30.9	3.4	10.0	5.7-16.3	4.4	6.5	4.0-9.9	2.6	8.5	4.4-14.8	1.8	5.2	2.3-10.1
	1991-1995	7.8	9.7	6.8-13.3	5.8	16.2	10.7-23.5	2.0	4.4	2.1-8.3	3.2	4.0	2.3-6.6	2.0	5.5	2.6-10.1	1.2	2.7	1.0-6.1
	1996-2000	9.0	10.8	7.9-14.5	7.0	17.3	12.0-24.3	2.0	4.7	2.2-8.8	1.4	1.7	0.7-3.6	0.8	1.9	0.5-5.1	0.6	1.3	0.3-4.1
	2001-2005	10.6	12.1	9.1-15.9	6.6	15.4	10.5-21.8	4.0	9.0	5.5-14.0	3.0	3.5	1.9-5.8	2.0	4.7	2.2-8.8	1.0	2.1	0.7-5.2
	2006-2010	11.0	11.2	8.4-14.7	8.4	17.0	12.2-23.2	2.6	5.3	2.8-9.3	3.0	3.3	1.8-5.5	2.2	4.7	2.3-8.6	0.8	1.8	0.5-4.8
	2011-2015	8.4	8.3	5.9-11.4	5.6	10.6	6.9-15.7	2.8	5.9	3.1-10.4	1.2	1.1	0.4-2.7	0.8	1.6	0.4-4.4	0.4	0.7	0.1-3.0
	2016-2020	15.0	14.1	11.0-17.9	12.6	23.4	17.8-30.4	2.4	4.8	2.4-8.8	2.2	1.8	0.9-3.4	2.0	3.4	1.6-6.7	0.2	0.3	0.0-2.5
Esophagus	1976-1980	4.2	6.4	3.9-10.0	3.2	11.1	6.0-18.5	1.0	2.8	0.9-6.8	3.4	5.3	3.0-8.6	2.8	10.0	5.2-17.3	0.6	1.7	0.3-5.2
	1981-1985	5.0	7.5	4.7-11.2	3.6	11.8	6.7-19.1	1.4	4.4	1.7-9.2	4.4	6.4	3.9-9.8	3.8	12.5	7.3-20.0	0.6	1.8	0.3-5.3
	1986-1990	4.6	6.8	4.3-10.3	3.6	12.3	7.2-19.5	1.0	2.3	0.7-5.7	3.4	4.5	2.6-7.3	2.2	6.6	3.2-12.1	1.2	2.9	1.0-6.5
	1991-1995	3.4	4.3	2.5-6.9	3.0	7.7	4.3-12.9	0.4	0.9	0.1-3.5	4.2	5.3	3.3-8.2	3.8	9.9	5.9-15.7	0.4	0.9	0.1-3.5
	1996-2000	4.8	5.9	3.8-8.9	3.6	8.9	5.3-14.2	1.2	2.6	1.0-5.9	4.2	5.1	3.2-7.9	3.6	9.1	5.3-14.7	0.6	1.3	0.3-4.0
	2001-2005	4.8	5.4	3.5-8.1	3.2	7.5	4.2-12.3	1.6	3.5	1.5-7.1	5.8	6.6	4.4-9.5	4.6	11.5	7.2-17.5	1.2	2.5	0.9-5.6
	2006-2010	5.4	6.0	4.0-8.9	4.4	10.4	6.4-16.0	1.0	2.2	0.7-5.4	2.8	3.2	1.7-5.4	2.0	4.7	2.2-8.9	0.8	1.7	0.4-4.6
	2011-2015	6.4	6.2	4.2-8.9	5.8	12.0	7.9-17.7	0.6	1.2	0.2-3.8	6.6	6.7	4.5-9.5	6.2	13.1	8.7-18.9	0.4	0.8	0.1-3.4
	2016-2020	5.2	4.8	3.1-7.3	3.8	7.0	4.2-11.3	1.4	2.5	1.0-5.6	3.2	2.9	1.6-4.9	2.4	4.2	2.1-7.7	0.8	1.6	0.4-4.4
Stomach	1976-1980	8.2	12.3	8.7-17.0	5.6	20.4	12.9-30.4	2.6	7.3	3.8-12.8	5.8	8.5	5.6-12.4	3.8	13.6	7.6-22.0	2.0	5.3	2.5-10.1
	1981-1985	11.6	16.4	12.4-21.4	8.2	24.1	17.1-33.1	3.4	9.9	5.6-16.0	7.0	10.5	7.2-14.7	5.2	17.3	11.1-25.8	1.8	5.3	2.4-10.2
	1986-1990	9.2	13.3	9.7-17.9	4.6	14.5	9.0-22.1	4.6	12.6	7.9-19.1	6.2	8.5	5.8-12.2	3.2	9.1	5.2-15.0	3.0	7.6	4.2-12.7
	1991-1995	8.6	11.2	8.1-15.2	6.8	20.0	13.7-28.2	1.8	4.0	1.8-7.8	5.2	7.1	4.7-10.5	3.6	11.7	6.8-18.5	1.6	3.8	1.6-7.7
	1996-2000	9.0	11.5	8.4-15.5	6.4	17.9	12.1-25.6	2.6	6.0	3.2-10.5	4.8	6.1	3.9-9.1	3.4	9.2	5.3-14.9	1.4	3.5	1.4-7.3
	2001-2005	7.6	9.1	6.4-12.6	4.4	10.3	6.4-15.7	3.2	7.1	4.0-11.7	3.6	4.1	2.4-6.6	2.2	4.8	2.4-8.8	1.4	3.0	1.2-6.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Sullivan**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	6.2	6.3	4.2-9.0	4.2	8.6	5.2-13.5	2.0	4.0	1.9-7.6	4.0	4.3	2.6-6.7	2.4	5.0	2.5-9.1	1.6	3.5	1.5-7.1
	2011-2015	6.4	6.8	4.6-9.8	5.2	11.6	7.4-17.3	1.2	2.4	0.8-5.5	3.4	3.5	2.0-5.7	2.6	5.8	2.9-10.2	0.8	1.4	0.4-4.1
	2016-2020	5.2	4.8	3.1-7.3	3.4	6.3	3.6-10.4	1.8	3.4	1.5-7.0	2.0	1.8	0.8-3.5	1.8	3.3	1.5-6.7	0.2	0.4	0.0-2.7
Colorectal	1976-1980	42.0	59.7	51.6-68.6	22.0	67.4	54.8-82.1	20.0	52.8	42.7-64.6	22.8	33.6	27.5-40.6	11.6	38.4	28.6-50.4	11.2	29.6	22.2-38.7
	1981-1985	48.2	68.7	60.0-78.2	24.0	74.7	61.3-90.1	24.2	63.5	52.4-76.3	20.6	29.2	23.7-35.6	10.8	35.0	25.7-46.5	9.8	25.2	18.6-33.6
	1986-1990	56.8	74.9	66.3-84.3	29.0	84.5	70.9-100.0	27.8	65.7	55.1-77.8	27.2	36.9	30.8-43.8	15.2	45.6	35.6-57.5	12.0	29.8	22.6-38.6
	1991-1995	55.2	70.8	62.6-79.8	28.6	81.9	68.6-97.0	26.6	62.1	51.9-73.9	26.2	33.7	28.1-40.1	14.2	41.0	31.6-52.1	12.0	27.6	21.0-35.7
	1996-2000	55.6	68.9	61.0-77.6	31.6	86.0	72.7-101.1	24.0	54.2	45.0-65.0	19.8	25.2	20.4-30.7	12.8	37.2	28.3-47.9	7.0	15.7	10.9-22.0
	2001-2005	46.6	53.7	47.0-61.2	23.4	56.7	46.7-68.3	23.2	50.9	42.0-61.2	17.8	21.1	16.9-26.1	9.0	22.0	15.9-29.8	8.8	19.1	13.8-25.8
	2006-2010	39.4	42.9	37.0-49.5	20.6	47.0	38.0-57.4	18.8	39.7	31.9-48.9	14.0	15.7	12.2-19.9	6.4	16.4	11.0-23.4	7.6	14.8	10.5-20.6
	2011-2015	38.8	39.7	34.2-46.0	21.4	45.6	37.1-55.6	17.4	34.3	27.3-42.7	14.2	14.2	11.0-18.2	7.6	15.7	10.9-22.0	6.6	13.3	9.0-19.0
2016-2020	40.2	39.7	34.2-45.9	25.0	49.2	40.6-59.1	15.2	30.3	23.5-38.5	14.8	14.2	11.1-18.0	8.4	16.5	11.8-22.7	6.4	11.7	8.0-16.9	
Colon excluding rectum	1976-1980	29.4	42.8	35.9-50.6	15.4	47.5	37.0-60.1	14.0	38.2	29.6-48.5	18.0	25.8	20.6-31.9	8.8	26.8	19.2-36.6	9.2	24.3	17.6-32.7
	1981-1985	36.2	51.6	44.2-60.0	17.4	53.3	42.2-66.5	18.8	49.8	40.0-61.4	17.2	24.5	19.5-30.5	9.0	29.6	21.0-40.4	8.2	21.2	15.2-29.1
	1986-1990	40.8	54.1	46.8-62.2	20.2	59.1	47.8-72.2	20.6	49.1	40.0-59.8	23.6	31.9	26.3-38.3	13.0	38.5	29.4-49.5	10.6	26.1	19.4-34.4
	1991-1995	40.4	52.4	45.4-60.3	19.6	58.6	47.2-71.9	20.8	49.0	39.9-59.6	21.6	27.9	22.8-33.7	11.4	33.4	25.0-43.6	10.2	23.6	17.5-31.2
	1996-2000	38.8	48.4	41.8-55.8	20.6	57.7	46.8-70.5	18.2	40.9	32.9-50.4	16.2	20.7	16.4-25.7	10.6	31.1	22.9-41.1	5.6	12.5	8.3-18.3
	2001-2005	33.8	39.3	33.6-45.8	17.0	41.8	33.2-52.1	16.8	37.3	29.7-46.3	14.2	16.8	13.1-21.2	6.8	16.3	11.2-23.1	7.4	16.0	11.2-22.2
	2006-2010	30.0	32.9	27.8-38.8	15.0	34.7	27.1-43.9	15.0	31.6	24.8-39.9	11.8	13.2	10.0-17.1	5.2	13.4	8.6-19.8	6.6	13.0	8.9-18.4
	2011-2015	27.2	28.0	23.4-33.3	15.0	32.2	25.1-40.8	12.2	24.0	18.2-31.2	11.0	11.5	8.6-15.1	5.8	12.5	8.2-18.3	5.2	11.0	7.1-16.4
2016-2020	27.4	26.0	21.7-31.0	16.2	30.9	24.3-38.8	11.2	21.2	15.8-28.0	11.6	11.1	8.4-14.6	6.2	12.3	8.3-17.8	5.4	9.8	6.4-14.6	
Rectum & rectosigmoid	1976-1980	12.6	16.9	12.8-21.8	6.6	19.9	13.4-28.5	6.0	14.6	9.8-21.2	4.8	7.8	4.9-11.8	2.8	11.6	6.0-19.7	2.0	5.3	2.5-10.0
	1981-1985	12.0	17.0	12.9-22.1	6.6	21.4	14.4-30.4	5.4	13.6	8.9-20.1	3.4	4.7	2.7-7.6	1.8	5.4	2.4-10.6	1.6	4.0	1.7-8.1
	1986-1990	16.0	20.8	16.4-26.1	8.8	25.4	18.3-34.5	7.2	16.5	11.6-23.1	3.6	5.0	2.9-8.0	2.2	7.1	3.4-12.8	1.4	3.7	1.4-7.8
	1991-1995	14.8	18.4	14.4-23.2	9.0	23.3	16.9-31.4	5.8	13.1	8.8-19.1	4.6	5.8	3.7-8.8	2.8	7.6	4.0-13.0	1.8	4.0	1.8-7.8
	1996-2000	16.8	20.5	16.3-25.4	11.0	28.3	21.2-37.1	5.8	13.3	8.9-19.3	3.6	4.5	2.7-7.2	2.2	6.1	2.9-11.0	1.4	3.2	1.3-6.8
	2001-2005	12.8	14.4	11.0-18.4	6.4	14.9	10.1-21.3	6.4	13.6	9.3-19.4	3.6	4.4	2.6-6.9	2.2	5.7	2.8-10.4	1.4	3.1	1.2-6.6
	2006-2010	9.4	9.9	7.3-13.4	5.6	12.2	8.0-18.0	3.8	8.1	4.8-12.9	2.2	2.5	1.2-4.5	1.2	3.0	1.1-6.8	1.0	1.8	0.6-4.7
	2011-2015	11.6	11.7	8.8-15.4	6.4	13.3	9.0-19.2	5.2	10.3	6.7-15.4	3.2	2.8	1.6-4.7	1.8	3.3	1.5-6.6	1.4	2.3	0.9-5.2
2016-2020	12.8	13.7	10.4-17.7	8.8	18.3	13.1-25.0	4.0	9.0	5.3-14.5	3.2	3.1	1.7-5.2	2.2	4.2	2.0-7.9	1.0	2.0	0.6-5.0	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Sullivan**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	3.6	5.0	2.9-8.1	2.2	6.4	3.1-11.8	1.4	4.1	1.6-8.6	3.6	5.2	3.0-8.3	1.8	5.4	2.4-10.6	1.8	4.9	2.2-9.5
	1981-1985	2.4	3.7	1.9-6.6	1.2	3.6	1.2-8.1	1.2	3.8	1.3-8.4	1.8	2.4	1.1-4.7	1.4	4.1	1.6-8.8	0.4	1.1	0.1-4.2
	1986-1990	4.8	7.1	4.5-10.6	2.4	8.1	4.0-14.3	2.4	6.5	3.3-11.5	3.0	4.5	2.5-7.4	1.4	5.5	2.1-11.1	1.6	4.1	1.7-8.3
	1991-1995	5.0	6.5	4.2-9.6	2.6	7.1	3.7-12.4	2.4	5.7	2.9-10.1	3.6	4.7	2.7-7.4	2.2	6.2	3.0-11.2	1.4	3.5	1.4-7.3
	1996-2000	6.6	8.2	5.7-11.6	5.2	13.6	8.8-20.0	1.4	3.1	1.2-6.5	5.4	6.8	4.5-10.0	4.4	11.8	7.3-18.1	1.0	2.2	0.7-5.4
	2001-2005	5.0	5.7	3.7-8.6	3.4	7.4	4.3-12.1	1.6	3.4	1.4-6.9	4.6	5.5	3.5-8.3	2.8	6.7	3.6-11.4	1.8	3.7	1.7-7.3
	2006-2010	8.8	9.2	6.6-12.4	7.4	14.7	10.3-20.6	1.4	3.3	1.3-6.9	5.0	5.4	3.5-8.1	4.2	9.1	5.6-14.2	0.8	1.9	0.5-5.0
	2011-2015	10.4	9.1	6.7-12.1	8.6	14.8	10.6-20.3	1.8	3.4	1.5-6.9	7.8	7.4	5.2-10.3	6.0	10.9	7.3-15.9	1.8	3.7	1.7-7.4
	2016-2020	10.0	8.6	6.3-11.5	8.4	14.6	10.4-20.1	1.6	2.7	1.1-5.8	5.8	4.8	3.2-7.2	4.8	8.2	5.2-12.6	1.0	1.6	0.5-4.3
Pancreas	1976-1980	8.4	12.6	9.0-17.2	4.6	14.2	8.7-21.8	3.8	10.8	6.4-17.1	8.8	13.4	9.6-18.1	4.6	15.1	9.2-23.2	4.2	11.8	7.3-18.2
	1981-1985	11.8	16.8	12.7-21.9	6.2	20.1	13.3-29.0	5.6	15.2	9.9-22.3	9.4	13.4	9.7-18.0	5.0	14.9	9.4-22.4	4.4	12.0	7.4-18.5
	1986-1990	11.2	15.0	11.3-19.6	5.4	15.3	10.0-22.5	5.8	14.5	9.6-21.1	9.8	13.1	9.6-17.5	5.0	14.6	9.3-21.8	4.8	12.3	7.7-18.6
	1991-1995	12.4	15.9	12.2-20.5	6.8	18.6	12.8-26.0	5.6	13.5	8.9-19.7	10.4	13.7	10.2-18.0	5.6	15.5	10.3-22.5	4.8	11.7	7.4-17.6
	1996-2000	10.4	12.9	9.6-17.0	4.4	12.2	7.4-18.7	6.0	13.5	9.1-19.5	9.6	11.8	8.7-15.7	4.0	10.7	6.4-16.8	5.6	12.8	8.5-18.6
	2001-2005	10.0	11.4	8.5-15.2	6.0	15.0	9.9-21.7	4.0	8.8	5.3-13.7	9.4	10.8	7.9-14.4	5.4	13.9	8.9-20.5	4.0	8.7	5.3-13.6
	2006-2010	12.6	13.7	10.5-17.6	7.6	16.2	11.4-22.6	5.0	10.2	6.5-15.2	11.6	12.5	9.4-16.3	6.2	13.4	9.0-19.4	5.4	10.5	6.9-15.5
	2011-2015	14.2	13.8	10.7-17.6	8.4	16.6	11.8-22.9	5.8	10.9	7.2-16.0	9.2	9.6	7.0-13.0	5.4	11.6	7.5-17.3	3.8	7.6	4.5-12.1
	2016-2020	17.4	16.1	12.8-20.1	8.6	16.0	11.5-21.9	8.8	16.4	11.7-22.5	15.6	14.9	11.7-18.8	7.8	14.9	10.5-20.8	7.8	14.8	10.4-20.8
Larynx	1976-1980	5.8	7.9	5.2-11.5	5.0	14.1	9.0-21.2	0.8	2.4	0.6-6.4	1.8	2.4	1.1-4.7	1.6	4.9	2.0-10.0	0.2	0.4	0.0-3.0
	1981-1985	5.2	7.3	4.6-10.8	4.8	14.1	8.9-21.4	0.4	0.8	0.1-3.6	1.8	2.5	1.1-4.8	1.6	4.7	2.0-9.6	0.2	0.4	0.0-3.0
	1986-1990	5.4	7.2	4.7-10.6	4.6	14.3	8.9-21.7	0.8	1.7	0.5-4.7	1.8	2.5	1.1-4.7	1.4	4.1	1.6-8.6	0.4	1.0	0.1-4.0
	1991-1995	6.6	8.5	5.8-12.0	4.6	12.6	7.9-19.2	2.0	4.7	2.2-8.8	2.6	3.2	1.7-5.6	2.2	6.2	3.0-11.2	0.4	0.8	0.1-3.4
	1996-2000	4.6	5.8	3.6-8.7	3.4	9.6	5.5-15.6	1.2	2.9	1.1-6.5	1.6	2.0	0.8-3.9	1.6	4.4	1.8-8.9	0.0	0.0	0.0-2.1
	2001-2005	5.0	5.5	3.5-8.2	3.4	7.2	4.2-11.7	1.6	3.8	1.6-7.7	1.0	1.1	0.4-2.7	0.8	1.8	0.5-5.0	0.2	0.5	0.0-2.9
	2006-2010	4.4	4.8	3.0-7.3	3.0	6.5	3.5-11.0	1.4	3.3	1.3-6.9	1.6	1.8	0.8-3.6	1.0	2.9	0.9-6.8	0.6	1.3	0.3-4.0
	2011-2015	4.2	3.9	2.4-6.2	3.6	6.7	3.9-10.9	0.6	1.1	0.2-3.8	1.2	1.2	0.4-2.8	1.2	2.4	0.9-5.5	0.0	0.0	0.0-2.1
	2016-2020	4.2	4.0	2.5-6.4	3.2	6.4	3.6-10.7	1.0	1.8	0.6-4.7	0.8	0.7	0.2-2.1	0.6	1.2	0.2-3.8	0.2	0.4	0.0-2.7
Lung and bronchus	1976-1980	49.0	67.2	58.8-76.5	33.8	99.6	84.4-116.9	15.2	40.5	31.6-51.1	39.6	53.1	45.7-61.3	29.0	84.8	70.7-100.8	10.6	27.5	20.4-36.3
	1981-1985	59.2	82.1	72.7-92.4	42.0	125.2	108.2-144.1	17.2	45.9	36.3-57.3	44.2	61.2	53.1-70.1	32.6	98.6	83.5-115.8	11.6	30.1	22.6-39.3
	1986-1990	64.2	83.5	74.5-93.4	42.0	119.8	103.7-137.8	22.2	53.4	43.8-64.7	48.0	63.3	55.4-72.0	31.6	89.4	75.6-104.9	16.4	40.9	32.3-51.1
	1991-1995	70.6	90.7	81.4-100.8	43.8	119.7	104.0-137.0	26.8	65.9	55.1-78.4	55.6	71.2	63.0-80.2	36.8	100.2	85.9-116.1	18.8	46.6	37.5-57.2
	1996-2000	63.4	76.9	68.6-85.9	37.4	95.5	82.1-110.6	26.0	60.0	50.1-71.3	49.8	60.7	53.3-68.7	30.2	78.1	65.9-91.9	19.6	44.7	36.3-54.6
	2001-2005	71.6	80.6	72.4-89.5	41.0	97.7	84.5-112.5	30.6	66.5	56.4-78.1	53.4	60.9	53.8-68.7	30.4	74.3	62.7-87.5	23.0	50.1	41.3-60.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Sullivan**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	76.4	81.8	73.7-90.6	36.6	81.4	69.7-94.6	39.8	84.1	72.7-97.0	51.6	55.8	49.1-63.2	27.2	62.8	52.4-74.7	24.4	51.5	42.7-61.8
	2011-2015	69.2	68.7	61.5-76.5	36.6	76.3	65.3-88.8	32.6	62.8	53.4-73.6	49.6	48.4	42.5-55.1	25.0	51.6	42.6-62.0	24.6	46.6	38.6-56.0
	2016-2020	71.0	64.6	57.9-71.9	34.2	64.5	54.9-75.4	36.8	65.2	55.9-75.8	45.6	41.0	35.8-46.9	25.2	47.8	39.6-57.3	20.4	34.9	28.4-42.8
Melanoma of the skin	1976-1980	4.0	6.3	3.8-9.9	2.8	9.8	5.2-16.8	1.2	3.6	1.2-8.0	1.0	1.6	0.5-3.9	0.6	2.4	0.5-6.7	0.4	0.8	0.1-3.6
	1981-1985	6.2	9.4	6.3-13.5	4.0	11.0	6.6-17.3	2.2	8.0	3.9-14.4	3.0	4.3	2.4-7.2	1.6	4.8	2.0-9.6	1.4	3.8	1.4-8.1
	1986-1990	5.6	7.4	4.9-10.8	3.2	8.7	4.9-14.3	2.4	6.0	3.1-10.7	2.2	3.2	1.6-5.8	1.0	2.8	0.9-6.7	1.2	3.6	1.3-7.9
	1991-1995	7.6	9.9	6.9-13.6	3.0	8.2	4.5-13.6	4.6	11.7	7.3-17.7	3.6	4.9	2.9-7.7	2.6	7.1	3.8-12.3	1.0	2.7	0.9-6.4
	1996-2000	9.4	11.6	8.5-15.5	4.0	10.3	6.2-16.1	5.4	12.6	8.3-18.5	1.4	1.9	0.7-3.9	0.6	1.9	0.3-5.6	0.8	1.9	0.5-5.0
	2001-2005	10.6	12.4	9.3-16.3	6.4	15.7	10.6-22.4	4.2	9.4	5.8-14.5	1.6	1.9	0.8-3.7	0.6	1.3	0.3-4.1	1.0	2.1	0.7-5.2
	2006-2010	14.4	15.2	11.8-19.3	8.6	19.1	13.7-26.1	5.8	11.7	7.8-17.0	1.8	1.9	0.9-3.8	1.0	2.2	0.7-5.3	0.8	1.8	0.5-4.9
	2011-2015	19.2	20.9	16.8-25.8	11.6	26.2	19.6-34.3	7.6	17.5	12.1-24.4	2.0	1.9	0.9-3.8	1.4	2.6	1.0-5.7	0.6	1.3	0.2-4.2
2016-2020	16.6	16.4	12.9-20.6	10.4	19.3	14.2-25.8	6.2	13.5	8.9-19.7	2.0	2.0	0.9-3.8	1.4	3.1	1.2-6.7	0.6	1.0	0.2-3.5	
Female breast	1976-1980							27.8	80.0	66.6-95.3							12.0	33.0	24.8-43.0
	1981-1985							37.6	103.9	89.0-120.5							14.4	39.0	30.2-49.6
	1986-1990							38.4	104.7	89.9-121.2							16.6	43.8	34.6-54.7
	1991-1995							47.6	121.8	106.5-138.6							14.0	35.6	27.6-45.3
	1996-2000							54.6	131.8	116.6-148.6							11.2	26.4	19.9-34.5
	2001-2005							58.8	133.3	118.4-149.6							9.2	20.6	15.1-27.7
	2006-2010							55.2	117.7	103.9-132.8							10.2	21.4	15.9-28.4
	2011-2015							55.2	113.5	100.0-128.5							9.0	17.3	12.5-23.5
2016-2020							56.0	115.7	101.7-131.2							13.2	26.9	20.5-34.6	
Cervix uteri	1976-1980							6.2	17.5	11.6-25.3							1.8	4.8	2.1-9.5
	1981-1985							4.2	12.4	7.5-19.3							2.6	6.9	3.6-12.0
	1986-1990							6.4	19.2	13.0-27.4							1.8	5.0	2.2-9.7
	1991-1995							4.6	12.6	8.0-19.1							1.8	5.0	2.2-9.6
	1996-2000							4.6	12.4	7.8-18.7							2.0	4.8	2.3-8.9
	2001-2005							4.8	11.3	7.2-17.0							1.4	3.5	1.4-7.4
	2006-2010							4.2	10.4	6.4-16.1							1.2	2.8	1.0-6.4
	2011-2015							4.6	11.5	7.1-17.6							1.6	3.6	1.5-7.5
	2016-2020							4.2	9.1	5.4-14.4							1.4	2.9	1.1-6.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Sullivan**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							9.6	25.4	18.5-34.1							2.4	5.7	2.9-10.2
	1981-1985							10.6	29.5	21.8-39.1							3.6	9.6	5.6-15.5
	1986-1990							9.2	24.4	17.7-33.0							2.2	5.6	2.8-10.2
	1991-1995							11.4	28.9	21.8-37.7							1.6	3.7	1.6-7.5
	1996-2000							15.0	35.4	27.8-44.6							2.8	6.3	3.4-10.7
	2001-2005							13.0	28.9	22.2-37.0							1.8	4.0	1.8-7.8
	2006-2010							16.0	33.5	26.4-41.9							2.4	5.2	2.6-9.2
	2011-2015							16.2	31.6	24.9-39.8							2.8	5.7	3.1-9.9
	2016-2020							17.8	34.7	27.4-43.5							2.4	5.0	2.5-9.2
Ovary	1976-1980							3.8	11.3	6.6-18.0							3.6	11.0	6.4-17.6
	1981-1985							5.4	15.2	9.9-22.4							2.6	6.8	3.5-12.0
	1986-1990							6.6	17.8	12.1-25.3							4.4	11.8	7.3-18.2
	1991-1995							7.2	17.9	12.4-25.0							6.2	14.4	9.7-20.6
	1996-2000							8.2	19.4	13.9-26.4							6.0	13.4	9.0-19.3
	2001-2005							9.8	22.0	16.3-29.3							4.8	10.5	6.7-15.8
	2006-2010							6.0	13.1	8.7-18.9							4.8	10.1	6.4-15.2
	2011-2015							6.8	14.5	9.9-20.7							3.6	7.2	4.2-11.7
	2016-2020							3.8	7.9	4.6-12.7							2.6	5.1	2.6-9.2
Prostate	1976-1980				26.8	91.9	75.8-110.2							8.8	29.6	20.7-40.8			
	1981-1985				30.2	97.0	81.3-114.9							9.8	32.4	23.5-43.4			
	1986-1990				35.8	112.8	96.2-131.4							10.8	32.8	24.4-43.2			
	1991-1995				52.6	148.9	130.8-168.7							8.6	27.9	19.9-37.8			
	1996-2000				64.0	164.5	146.5-184.0							12.4	38.9	29.4-50.2			
	2001-2005				60.8	140.2	124.5-157.3							10.8	32.3	23.9-42.6			
	2006-2010				65.0	136.8	121.9-153.1							9.8	29.1	21.3-38.7			
	2011-2015				46.8	89.3	77.8-102.1							7.8	20.3	14.3-27.9			
	2016-2020				53.0	90.7	79.9-102.8							5.6	11.8	7.7-17.4			
Testis	1976-1980				0.2	0.4	0.0-3.2							0.0	0.0	0.0-2.6			
	1981-1985				1.0	3.0	0.9-7.3							0.0	0.0	0.0-2.5			
	1986-1990				1.4	4.1	1.6-8.6							0.0	0.0	0.0-2.3			
	1991-1995				1.2	3.0	1.1-6.6							0.0	0.0	0.0-2.1			
	1996-2000				1.6	4.5	1.9-8.9							0.0	0.0	0.0-2.1			
	2001-2005				1.8	5.1	2.3-9.6							0.4	1.2	0.1-4.1			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Sullivan**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				1.0	2.5	0.8-5.9							0.4	1.1	0.1-3.9			
	2011-2015				1.8	4.6	2.0-9.0							0.2	0.4	0.0-2.6			
	2016-2020				1.8	5.1	2.3-9.7							0.2	0.3	0.0-2.5			
Urinary bladder (incl. in situ)	1976-1980	15.8	21.8	17.1-27.4	11.6	35.4	26.3-46.6	4.2	10.4	6.4-16.2	4.6	6.3	3.9-9.7	3.0	10.4	5.5-17.8	1.6	3.8	1.6-7.7
	1981-1985	17.0	24.4	19.4-30.4	14.0	47.4	36.2-60.7	3.0	7.2	4.0-12.2	6.2	9.2	6.2-13.2	4.4	16.4	9.8-25.3	1.8	4.7	2.1-9.2
	1986-1990	15.0	20.1	15.7-25.3	11.6	34.9	26.2-45.5	3.4	8.2	4.7-13.3	4.0	5.6	3.4-8.7	2.8	9.8	5.2-16.6	1.2	2.8	1.0-6.3
	1991-1995	16.6	21.5	17.1-26.8	12.2	34.7	26.3-44.9	4.4	10.3	6.4-15.7	3.6	4.8	2.9-7.7	2.0	6.1	2.9-11.3	1.6	3.8	1.6-7.6
	1996-2000	24.2	29.7	24.7-35.6	18.0	49.7	39.6-61.5	6.2	14.0	9.5-20.0	5.6	7.1	4.7-10.3	4.0	12.8	7.6-19.9	1.6	3.5	1.5-7.2
	2001-2005	24.0	27.4	22.7-32.8	16.8	41.9	33.1-52.3	7.2	15.9	11.1-22.1	5.0	5.9	3.8-8.8	3.4	8.9	5.1-14.4	1.6	3.5	1.5-7.2
	2006-2010	26.6	29.5	24.6-35.1	20.0	47.5	38.3-58.2	6.6	14.2	9.7-20.1	3.8	4.3	2.6-6.8	2.6	6.8	3.5-11.8	1.2	2.4	0.9-5.4
	2011-2015	25.6	26.5	22.0-31.7	20.2	45.4	36.7-55.7	5.4	10.3	6.7-15.3	5.4	5.4	3.6-8.0	3.6	7.6	4.4-12.4	1.8	3.3	1.5-6.6
	2016-2020	25.4	24.0	19.9-28.7	19.4	38.9	31.3-47.9	6.0	11.0	7.3-16.1	3.0	3.1	1.7-5.3	2.8	6.7	3.6-11.4	0.2	0.4	0.0-2.7
Kidney and renal pelvis	1976-1980	5.4	7.6	4.9-11.2	3.0	8.8	4.6-15.1	2.4	6.8	3.4-12.2	3.0	4.6	2.5-7.7	1.2	3.8	1.2-8.9	1.8	5.4	2.4-10.4
	1981-1985	5.4	7.7	5.0-11.3	3.6	10.3	6.0-16.6	1.8	5.3	2.3-10.2	2.6	3.4	1.8-6.0	2.0	5.2	2.5-9.8	0.6	1.9	0.3-5.7
	1986-1990	6.6	9.5	6.5-13.4	3.6	11.4	6.7-18.2	3.0	8.0	4.4-13.4	2.8	4.1	2.2-6.9	2.0	6.5	3.1-11.9	0.8	1.9	0.5-5.1
	1991-1995	9.2	11.7	8.5-15.7	5.0	14.2	9.1-21.1	4.2	10.0	6.1-15.5	4.2	5.2	3.2-8.1	2.8	7.5	4.1-12.8	1.4	3.3	1.3-6.9
	1996-2000	8.2	10.1	7.2-13.7	4.8	12.0	7.6-17.9	3.4	7.8	4.6-12.7	2.4	3.1	1.6-5.4	1.6	4.1	1.8-8.2	0.8	1.8	0.5-4.8
	2001-2005	11.2	13.2	9.9-17.2	6.4	16.3	11.0-23.2	4.8	10.8	6.9-16.3	2.8	3.4	1.8-5.7	2.2	6.0	2.9-11.0	0.6	1.4	0.3-4.3
	2006-2010	15.4	16.3	12.8-20.5	9.4	20.0	14.6-26.9	6.0	12.3	8.3-17.9	4.0	4.2	2.5-6.6	2.2	4.6	2.3-8.6	1.8	3.7	1.7-7.2
	2011-2015	20.0	20.8	16.8-25.5	14.4	29.8	23.1-38.0	5.6	11.9	7.8-17.5	4.8	5.1	3.2-7.7	3.8	8.1	4.8-13.0	1.0	2.4	0.8-5.8
	2016-2020	15.4	15.3	11.9-19.4	8.8	17.4	12.5-23.8	6.6	13.3	8.9-19.2	4.8	4.6	2.9-7.1	2.8	5.9	3.2-10.2	2.0	3.2	1.5-6.5
Brain and other central nervous system	1976-1980	4.2	5.9	3.6-9.1	2.2	6.8	3.3-12.4	2.0	5.2	2.4-9.9	2.0	2.7	1.2-5.1	1.0	3.1	1.0-7.4	1.0	2.2	0.7-5.6
	1981-1985	4.8	7.6	4.7-11.4	2.4	7.5	3.8-13.3	2.4	7.7	3.8-13.7	3.2	4.8	2.7-8.0	1.8	5.5	2.4-10.8	1.4	4.7	1.8-9.8
	1986-1990	4.4	6.3	3.9-9.6	2.8	8.3	4.5-14.1	1.6	4.4	1.8-8.8	2.4	3.3	1.7-5.9	1.4	4.3	1.7-8.9	1.0	2.4	0.8-5.9
	1991-1995	6.0	7.9	5.3-11.3	2.0	5.0	2.4-9.4	4.0	10.5	6.4-16.3	3.8	4.9	2.9-7.7	1.6	4.1	1.8-8.3	2.2	5.6	2.7-10.1
	1996-2000	6.8	8.8	6.1-12.3	3.8	9.9	5.9-15.7	3.0	7.9	4.4-13.1	1.4	1.8	0.7-3.8	0.6	1.9	0.4-5.7	0.8	2.0	0.5-5.2
	2001-2005	5.8	7.3	4.8-10.5	2.6	7.0	3.7-12.1	3.2	7.7	4.3-12.6	4.8	5.4	3.5-8.1	2.6	5.9	3.1-10.2	2.2	4.7	2.3-8.6
	2006-2010	6.0	6.8	4.5-9.8	4.2	9.3	5.7-14.5	1.8	4.3	1.9-8.4	3.0	3.4	1.9-5.7	2.4	5.5	2.8-9.7	0.6	1.2	0.2-3.8
	2011-2015	6.8	7.7	5.3-10.9	4.6	10.1	6.3-15.4	2.2	5.4	2.6-9.8	3.4	3.5	2.0-5.8	2.0	4.4	2.1-8.4	1.4	2.8	1.1-6.2
	2016-2020	5.2	5.6	3.6-8.4	3.4	7.8	4.5-12.8	1.8	3.6	1.6-7.3	3.0	3.3	1.8-5.6	1.8	4.4	1.9-8.5	1.2	2.1	0.8-5.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Sullivan**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	1.0	1.4	0.4-3.3	0.0	0.0	0.0-2.6	1.0	2.7	0.8-6.5	0.4	0.4	0.1-1.9	0.2	0.4	0.0-3.2	0.2	0.4	0.0-3.1
	1981-1985	2.8	4.8	2.6-8.1	0.6	2.3	0.5-6.5	2.2	7.5	3.7-13.5	0.4	0.6	0.1-2.3	0.0	0.0	0.0-2.5	0.4	1.0	0.1-4.0
	1986-1990	2.4	3.5	1.8-6.3	0.6	1.8	0.4-5.3	1.8	5.3	2.3-10.2	0.4	0.6	0.1-2.1	0.2	0.7	0.0-3.7	0.2	0.6	0.0-3.4
	1991-1995	3.4	4.5	2.6-7.3	1.8	4.7	2.1-9.0	1.6	4.4	1.9-8.8	0.8	0.9	0.3-2.5	0.4	1.0	0.1-3.9	0.4	0.9	0.1-3.5
	1996-2000	7.2	9.6	6.7-13.3	2.2	5.5	2.7-10.0	5.0	13.9	9.0-20.7	1.4	1.8	0.7-3.7	0.8	2.0	0.5-5.3	0.6	1.3	0.3-4.1
	2001-2005	10.2	12.9	9.6-17.1	2.2	4.9	2.4-9.1	8.0	21.3	15.1-29.1	1.2	1.4	0.5-3.2	0.2	0.6	0.0-3.2	1.0	2.1	0.7-5.1
	2006-2010	15.2	18.8	14.7-23.6	3.4	8.1	4.6-13.2	11.8	30.0	22.7-39.0	0.8	0.9	0.2-2.4	0.0	0.0	0.0-2.0	0.8	1.6	0.4-4.4
	2011-2015	14.0	15.9	12.2-20.3	4.0	9.0	5.4-14.3	10.0	23.0	16.8-30.7	0.2	0.1	0.0-1.2	0.2	0.3	0.0-2.4	0.0	0.0	0.0-2.1
2016-2020	12.0	13.9	10.4-18.2	2.8	5.7	3.0-9.9	9.2	22.9	16.4-31.0	0.0	0.0	0.0-1.0	0.0	0.0	0.0-2.0	0.0	0.0	0.0-2.1	
Hodgkin lymphoma	1976-1980	3.6	4.5	2.7-7.3	2.0	4.9	2.3-9.5	1.6	4.3	1.8-8.7	1.8	2.6	1.1-5.1	1.0	2.9	0.9-7.2	0.8	2.2	0.6-5.8
	1981-1985	2.6	4.1	2.1-7.1	1.6	5.3	2.2-10.5	1.0	2.6	0.8-6.4	1.0	1.4	0.4-3.4	0.6	1.8	0.4-5.5	0.4	1.0	0.1-3.9
	1986-1990	1.6	2.5	1.1-5.0	1.0	3.2	1.0-7.4	0.6	1.9	0.4-5.6	0.6	1.0	0.2-3.0	0.4	1.4	0.2-4.9	0.2	0.5	0.0-3.2
	1991-1995	1.6	2.4	1.0-4.6	1.0	2.9	0.9-6.7	0.6	1.8	0.4-5.1	0.4	0.7	0.1-2.3	0.0	0.0	0.0-2.1	0.4	1.4	0.2-4.6
	1996-2000	2.2	3.3	1.6-5.9	1.2	3.5	1.3-7.7	1.0	3.0	0.9-6.9	0.4	0.7	0.1-2.3	0.2	0.5	0.0-3.0	0.2	0.8	0.0-3.8
	2001-2005	2.4	3.2	1.6-5.6	1.2	3.0	1.1-6.8	1.2	3.3	1.2-7.2	1.0	1.1	0.3-2.6	0.4	0.9	0.1-3.4	0.6	1.3	0.3-4.0
	2006-2010	3.6	4.7	2.7-7.5	2.4	6.6	3.3-11.6	1.2	2.9	1.0-6.6	0.0	0.0	0.0-1.0	0.0	0.0	0.0-2.0	0.0	0.0	0.0-2.0
	2011-2015	1.6	2.1	0.9-4.2	1.2	3.0	1.1-6.7	0.4	1.1	0.1-4.1	0.0	0.0	0.0-1.0	0.0	0.0	0.0-2.0	0.0	0.0	0.0-2.1
2016-2020	2.4	2.9	1.4-5.2	1.6	3.9	1.6-7.8	0.8	1.9	0.4-5.2	0.2	0.2	0.0-1.3	0.2	0.4	0.0-2.7	0.0	0.0	0.0-2.1	
Non-Hodgkin lymphomas	1976-1980	6.2	7.9	5.3-11.5	3.0	8.8	4.7-14.9	3.2	7.2	4.0-12.0	2.8	3.9	2.1-6.6	1.8	6.2	2.7-11.9	1.0	2.2	0.7-5.5
	1981-1985	9.6	12.6	9.2-16.8	5.8	16.2	10.7-23.6	3.8	9.5	5.7-15.1	4.4	6.3	3.9-9.6	2.6	8.3	4.3-14.5	1.8	4.5	2.0-8.8
	1986-1990	16.8	23.0	18.2-28.6	11.0	31.2	23.3-40.9	5.8	14.5	9.6-21.1	6.8	9.6	6.6-13.6	4.4	13.4	8.3-20.5	2.4	6.1	3.1-10.9
	1991-1995	21.2	27.6	22.6-33.5	10.4	28.8	21.3-38.1	10.8	26.9	20.1-35.2	10.6	13.7	10.3-18.1	5.6	15.1	10.0-21.9	5.0	11.8	7.6-17.5
	1996-2000	15.4	19.5	15.3-24.4	7.4	19.6	13.7-27.2	8.0	19.6	13.9-26.8	7.6	9.5	6.7-13.2	3.4	9.3	5.3-15.1	4.2	9.9	6.1-15.2
	2001-2005	19.6	23.1	18.7-28.2	11.0	26.6	19.9-34.8	8.6	19.9	14.3-27.0	6.4	7.4	5.0-10.5	4.2	9.6	5.9-15.0	2.2	5.0	2.4-9.2
	2006-2010	17.8	19.5	15.6-24.1	8.8	19.1	13.7-26.0	9.0	19.9	14.4-26.8	5.2	6.2	4.0-9.1	3.8	10.1	6.0-16.0	1.4	3.1	1.2-6.5
	2011-2015	21.6	22.7	18.5-27.6	12.0	26.1	19.7-33.9	9.6	19.5	14.2-26.3	3.2	3.6	2.1-6.0	2.4	5.7	2.9-10.0	0.8	1.7	0.4-4.6
2016-2020	17.8	16.9	13.5-21.1	9.6	17.5	12.8-23.5	8.2	17.1	12.0-23.8	2.8	2.6	1.4-4.5	2.0	3.8	1.8-7.3	0.8	1.5	0.4-4.5	
Myeloma	1976-1980	3.6	4.5	2.6-7.3	1.2	3.0	1.1-7.0	2.4	5.9	3.0-10.7	3.0	4.0	2.2-6.8	1.2	3.0	1.1-7.0	1.8	5.0	2.2-9.8
	1981-1985	5.6	7.7	5.0-11.3	3.0	8.2	4.5-13.9	2.6	6.7	3.5-11.8	4.4	5.5	3.4-8.4	2.2	5.4	2.7-10.0	2.2	5.1	2.5-9.5
	1986-1990	3.4	4.6	2.7-7.5	1.8	5.4	2.5-10.4	1.6	3.9	1.7-7.9	1.6	2.3	1.0-4.7	0.8	2.7	0.7-6.8	0.8	1.9	0.5-5.2
	1991-1995	2.8	3.8	2.1-6.4	1.6	5.8	2.4-11.3	1.2	2.8	1.0-6.3	1.8	2.4	1.1-4.7	1.0	3.3	1.0-7.7	0.8	1.7	0.5-4.7
	1996-2000	5.6	6.8	4.5-9.9	3.6	9.8	5.7-15.8	2.0	4.5	2.2-8.5	3.6	4.5	2.7-7.2	2.4	7.7	3.8-13.6	1.2	2.6	1.0-6.0
	2001-2005	5.0	5.9	3.8-8.8	3.2	8.9	4.9-14.6	1.8	4.0	1.8-7.7	3.0	3.6	2.0-6.0	1.4	3.6	1.4-7.7	1.6	3.4	1.5-7.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Sullivan**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	6.6	7.1	4.9-10.1	4.6	9.9	6.2-15.1	2.0	4.5	2.2-8.5	2.8	3.2	1.8-5.5	2.4	6.7	3.3-11.8	0.4	0.9	0.1-3.4
	2011-2015	7.4	7.2	5.1-10.1	4.4	8.6	5.3-13.3	3.0	5.9	3.3-10.0	1.8	1.9	0.9-3.7	0.8	1.8	0.5-4.7	1.0	2.1	0.7-5.2
	2016-2020	7.4	7.0	4.9-9.8	4.8	9.3	5.9-14.2	2.6	4.8	2.5-8.6	2.0	2.1	1.0-4.0	1.4	3.4	1.3-7.1	0.6	1.2	0.3-4.0
Leukemias	1976-1980	8.8	12.6	9.0-17.1	6.0	18.3	12.0-26.6	2.8	7.4	4.0-12.6	5.2	8.4	5.4-12.4	2.8	9.9	5.2-16.9	2.4	7.1	3.6-12.5
	1981-1985	13.2	17.9	13.8-23.0	7.4	21.0	14.6-29.3	5.8	14.8	9.8-21.6	6.8	9.6	6.6-13.6	3.6	10.6	6.2-17.0	3.2	8.3	4.7-13.8
	1986-1990	9.8	13.7	10.1-18.2	4.4	13.0	8.0-19.8	5.4	14.1	9.2-20.8	5.8	8.0	5.3-11.7	3.0	9.3	5.0-15.5	2.8	7.8	4.2-13.3
	1991-1995	10.4	13.6	10.1-17.9	5.8	16.1	10.6-23.4	4.6	11.5	7.2-17.4	6.4	8.3	5.7-11.8	4.0	11.1	6.7-17.3	2.4	5.4	2.8-9.7
	1996-2000	13.2	16.7	12.9-21.3	7.4	20.6	14.3-28.7	5.8	13.7	9.1-19.9	8.6	11.1	8.0-15.0	4.6	13.3	8.2-20.2	4.0	10.1	6.1-15.8
	2001-2005	12.8	16.0	12.3-20.4	7.6	20.3	14.3-28.0	5.2	12.1	7.9-17.9	6.6	8.0	5.5-11.3	4.2	11.3	6.9-17.4	2.4	5.3	2.7-9.4
	2006-2010	14.4	16.7	13.0-21.2	8.0	18.2	12.9-25.1	6.4	15.4	10.4-21.9	5.4	6.2	4.1-9.1	2.8	6.7	3.6-11.4	2.6	5.7	3.0-10.0
	2011-2015	10.6	11.1	8.2-14.7	6.2	13.4	9.0-19.4	4.4	9.0	5.5-13.9	6.0	6.4	4.3-9.3	3.8	9.0	5.3-14.3	2.2	4.6	2.2-8.6
2016-2020	14.4	15.2	11.7-19.4	8.2	18.1	12.8-25.0	6.2	12.3	8.2-17.9	6.2	6.0	4.0-8.6	2.6	6.0	3.1-10.4	3.6	6.0	3.6-10.0	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Tioga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	163.4	423.9	394.6-454.6	85.4	521.8	470.4-576.9	78.0	363.5	328.0-401.7	80.8	212.2	191.5-234.3	43.8	265.8	229.6-305.7	37.0	175.1	150.6-202.4
	1981-1985	164.6	397.1	370.0-425.7	84.8	487.2	439.3-538.6	79.8	344.5	311.2-380.3	78.4	192.9	173.9-213.2	43.0	256.2	221.3-294.5	35.4	150.8	129.2-174.8
	1986-1990	203.0	450.3	422.7-479.3	101.4	538.7	489.8-590.8	101.6	405.9	371.2-443.0	92.4	207.8	189.0-227.9	48.2	262.5	228.3-300.2	44.2	175.5	153.0-200.3
	1991-1995	238.6	491.1	463.4-520.0	133.4	640.1	589.8-693.4	105.2	394.7	361.6-430.0	104.4	222.1	203.3-242.1	52.2	275.8	241.3-313.5	52.2	194.4	171.5-219.6
	1996-2000	249.6	473.0	447.0-500.1	136.8	587.8	542.8-635.4	112.8	390.8	359.1-424.6	103.6	200.8	183.8-219.0	60.2	284.1	251.3-319.8	43.4	147.2	128.2-168.3
	2001-2005	287.0	500.2	474.5-527.0	155.0	596.8	554.4-641.6	132.0	429.5	397.1-464.0	106.4	187.0	171.4-203.7	57.0	234.6	207.2-264.7	49.4	155.9	136.9-176.9
	2006-2010	322.4	518.6	493.2-545.0	172.0	593.6	553.7-635.9	150.4	463.4	430.2-498.6	107.0	172.4	158.0-187.8	57.8	208.6	184.7-235.0	49.2	147.2	129.1-167.2
	2011-2015	316.0	477.9	453.9-503.0	164.2	517.6	481.7-555.5	151.8	446.4	413.8-481.0	95.6	139.9	127.4-153.3	48.6	156.2	136.8-177.7	47.0	128.6	112.2-146.9
	2016-2020	337.4	479.9	456.0-504.8	178.0	515.4	480.7-552.2	159.4	453.3	420.2-488.6	94.4	127.8	116.2-140.5	52.2	153.1	134.6-173.8	42.2	106.4	92.0-122.8
Oral cavity and pharynx	1976-1980	4.6	11.0	6.9-16.6	2.8	14.7	7.8-25.1	1.8	8.1	3.7-15.4	2.0	4.6	2.2-8.5	1.4	6.7	2.7-14.1	0.6	2.7	0.5-7.9
	1981-1985	2.8	6.7	3.6-11.2	1.6	8.8	3.5-17.7	1.2	5.1	1.9-11.2	0.6	1.4	0.3-4.1	0.2	0.8	0.0-5.1	0.4	1.7	0.2-6.2
	1986-1990	4.0	8.3	5.0-12.8	2.8	13.2	6.9-22.7	1.2	4.5	1.7-9.9	1.2	2.6	0.9-5.6	0.6	2.9	0.6-8.7	0.6	2.3	0.5-6.8
	1991-1995	5.8	11.5	7.7-16.6	4.2	17.8	10.9-27.4	1.6	5.7	2.5-11.3	0.8	1.6	0.4-4.2	0.6	3.1	0.6-9.1	0.2	0.7	0.0-4.0
	1996-2000	4.6	8.6	5.4-13.0	2.2	9.6	4.6-17.4	2.4	8.1	4.2-14.3	1.6	3.1	1.3-6.1	1.0	5.2	1.6-12.2	0.6	2.1	0.4-6.2
	2001-2005	4.8	8.4	5.4-12.5	2.8	9.9	5.4-16.9	2.0	6.5	3.1-12.3	0.6	0.9	0.2-2.9	0.6	1.9	0.4-6.1	0.0	0.0	0.0-2.9
	2006-2010	6.0	9.4	6.3-13.6	3.8	12.8	7.6-20.4	2.2	6.5	3.2-11.9	1.2	2.0	0.7-4.4	0.4	1.9	0.2-6.6	0.8	2.4	0.7-6.7
	2011-2015	4.8	6.8	4.3-10.4	3.8	10.9	6.4-17.6	1.0	2.8	0.9-7.1	0.8	1.1	0.3-3.1	0.6	1.8	0.4-5.8	0.2	0.5	0.0-3.8
	2016-2020	8.6	12.0	8.5-16.6	7.2	21.4	14.6-30.4	1.4	3.0	1.2-7.2	1.2	1.7	0.6-4.2	1.0	3.1	0.9-7.9	0.2	0.4	0.0-3.8
Esophagus	1976-1980	1.6	4.2	1.8-8.2	0.4	2.1	0.2-7.6	1.2	5.7	2.1-12.2	1.0	2.4	0.8-5.7	0.4	1.9	0.2-7.3	0.6	2.8	0.6-8.0
	1981-1985	2.2	5.7	2.8-10.1	1.6	10.4	4.3-20.4	0.6	2.5	0.5-7.3	1.8	4.8	2.2-9.0	1.0	7.5	2.3-16.8	0.8	3.4	0.9-8.7
	1986-1990	1.8	3.7	1.7-7.0	1.8	8.0	3.6-15.4	0.0	0.0	0.0-2.9	1.6	3.3	1.4-6.6	1.6	7.2	3.0-14.5	0.0	0.0	0.0-2.9
	1991-1995	2.8	5.6	3.0-9.4	2.2	9.2	4.5-16.8	0.6	2.4	0.5-6.9	2.4	5.0	2.6-8.7	1.8	9.1	3.8-17.8	0.6	2.3	0.5-6.7
	1996-2000	2.4	4.5	2.3-7.8	1.6	6.2	2.6-12.4	0.8	2.6	0.7-7.0	2.6	4.7	2.5-8.1	1.8	6.6	3.0-12.8	0.8	2.7	0.7-7.0
	2001-2005	3.6	6.1	3.6-9.8	2.6	9.9	5.1-17.3	1.0	3.2	1.0-7.7	3.2	5.3	3.0-8.8	2.6	9.0	4.8-15.7	0.6	1.9	0.4-5.9
	2006-2010	2.4	3.6	1.9-6.5	1.6	5.1	2.2-10.4	0.8	2.3	0.6-6.3	2.2	3.5	1.7-6.4	1.8	6.4	2.8-12.4	0.4	1.1	0.1-4.6
	2011-2015	3.2	4.6	2.6-7.7	3.0	9.5	5.2-16.1	0.2	0.5	0.0-3.8	2.4	3.5	1.8-6.3	2.2	7.1	3.5-13.2	0.2	0.6	0.0-4.1
	2016-2020	4.4	5.5	3.4-8.7	3.6	8.7	5.1-14.4	0.8	2.5	0.6-7.2	3.2	4.5	2.5-7.7	2.8	8.4	4.4-14.8	0.4	1.0	0.1-4.6
Stomach	1976-1980	3.4	8.7	5.0-14.0	2.4	13.2	6.6-23.4	1.0	4.7	1.5-11.0	2.4	6.5	3.3-11.3	1.8	10.8	4.8-20.6	0.6	2.9	0.6-8.3
	1981-1985	3.2	8.4	4.7-13.5	2.2	12.5	6.1-22.5	1.0	4.4	1.4-10.2	2.2	5.6	2.8-10.0	1.4	6.9	2.8-14.4	0.8	3.7	1.0-9.2
	1986-1990	3.8	8.7	5.2-13.5	2.8	16.3	8.4-27.8	1.0	3.8	1.2-9.0	2.2	5.6	2.7-9.8	1.4	10.2	3.9-20.6	0.8	3.1	0.8-7.9
	1991-1995	3.4	7.5	4.3-11.9	2.4	13.3	6.5-23.4	1.0	3.7	1.2-8.7	2.0	4.2	2.0-7.7	1.4	6.7	2.6-13.8	0.6	2.3	0.5-6.7
	1996-2000	4.8	9.4	6.0-14.1	2.8	13.1	6.9-22.4	2.0	6.5	3.1-12.1	3.4	6.9	4.0-11.0	2.0	11.1	5.0-20.5	1.4	4.5	1.8-9.6
	2001-2005	5.2	8.9	5.8-13.2	3.4	12.4	7.2-20.1	1.8	5.9	2.7-11.4	2.2	3.8	1.9-6.8	1.2	4.2	1.5-9.5	1.0	3.2	1.0-7.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Tioga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	3.0	4.9	2.7-8.2	2.0	7.1	3.3-13.4	1.0	3.2	1.0-7.9	2.0	3.2	1.5-6.0	1.8	6.5	2.9-12.7	0.2	0.6	0.0-3.9
	2011-2015	4.8	7.1	4.5-10.8	3.2	9.8	5.5-16.3	1.6	5.1	2.1-10.6	1.2	1.7	0.6-4.0	0.6	1.7	0.3-5.8	0.6	1.7	0.3-5.6
	2016-2020	4.0	5.4	3.2-8.8	2.2	6.1	2.9-11.9	1.8	4.5	2.0-9.4	2.0	2.8	1.3-5.6	1.0	2.9	0.9-7.6	1.0	2.7	0.8-7.0
Colorectal	1976-1980	24.2	65.7	54.3-78.7	12.4	81.2	61.0-105.4	11.8	55.7	42.3-71.9	10.6	28.2	20.9-37.0	4.0	22.6	13.4-35.3	6.6	30.7	21.1-43.2
	1981-1985	25.2	63.4	52.7-75.7	12.8	76.9	58.2-99.2	12.4	54.4	41.7-69.8	10.4	27.0	20.0-35.4	5.4	35.7	22.9-52.5	5.0	21.5	13.9-31.8
	1986-1990	31.2	71.6	60.6-83.8	14.2	77.1	59.3-98.3	17.0	67.8	54.1-83.8	12.6	28.2	21.6-36.1	5.8	29.3	19.2-42.5	6.8	26.3	18.2-36.8
	1991-1995	29.0	62.0	52.2-73.0	13.6	68.1	51.9-87.4	15.4	57.5	45.3-71.9	14.6	31.3	24.5-39.4	5.8	29.6	19.2-43.2	8.8	32.4	23.5-43.5
	1996-2000	28.0	54.6	45.9-64.4	15.4	66.7	52.2-83.8	12.6	42.3	32.4-54.2	10.2	20.3	15.1-26.7	6.2	30.5	20.3-43.7	4.0	13.4	8.2-20.8
	2001-2005	32.2	55.8	47.5-65.3	16.8	65.7	52.0-81.9	15.4	48.4	38.1-60.7	12.0	20.9	16.0-27.0	6.6	26.4	18.0-37.4	5.4	16.6	10.9-24.4
	2006-2010	30.6	49.7	42.0-58.3	17.6	63.4	50.5-78.7	13.0	38.6	29.7-49.6	8.4	13.6	9.8-18.5	5.2	19.9	12.8-29.5	3.2	9.6	5.5-15.9
	2011-2015	27.8	41.2	34.5-49.0	15.2	48.2	37.7-60.9	12.6	35.7	27.1-46.3	8.0	11.6	8.2-15.9	3.4	11.3	6.5-18.5	4.6	11.9	7.5-18.4
2016-2020	29.6	43.8	36.6-52.1	16.2	49.9	39.1-63.1	13.4	37.8	28.7-49.1	8.4	11.1	7.9-15.3	4.4	12.0	7.5-18.8	4.0	10.5	6.3-17.0	
Colon excluding rectum	1976-1980	16.4	44.2	34.9-55.0	7.2	47.6	32.4-66.9	9.2	43.1	31.5-57.6	8.6	22.9	16.5-31.0	2.8	16.4	8.7-27.8	5.8	26.8	17.9-38.6
	1981-1985	19.4	49.4	39.9-60.3	9.6	57.3	41.5-76.7	9.8	43.3	32.0-57.2	8.6	22.3	16.1-30.1	4.2	28.3	16.9-43.7	4.4	18.9	11.8-28.7
	1986-1990	22.0	50.7	41.5-61.2	9.4	51.7	37.3-69.5	12.6	50.1	38.5-64.1	10.2	23.1	17.1-30.5	5.0	26.0	16.5-38.8	5.2	20.2	13.2-29.6
	1991-1995	21.6	46.3	37.9-55.9	10.0	49.4	35.8-66.1	11.6	43.5	33.0-56.2	11.4	24.6	18.6-31.9	4.6	22.6	13.8-34.4	6.8	25.2	17.4-35.3
	1996-2000	19.0	37.6	30.4-46.0	9.8	43.2	31.7-57.4	9.2	30.8	22.5-41.3	8.2	16.4	11.7-22.3	5.0	24.2	15.4-36.1	3.2	10.7	6.1-17.6
	2001-2005	23.8	41.4	34.3-49.7	11.6	45.3	34.1-59.0	12.2	38.3	29.3-49.4	9.4	16.5	12.1-22.0	4.8	20.0	12.6-30.1	4.6	14.2	9.0-21.5
	2006-2010	21.6	35.7	29.2-43.2	11.0	41.5	30.9-54.5	10.6	31.9	23.8-42.1	6.6	10.7	7.3-15.1	4.0	15.2	9.2-23.8	2.6	7.7	4.1-13.6
	2011-2015	20.8	31.2	25.3-38.0	11.2	35.3	26.5-46.3	9.6	28.3	20.6-38.1	6.4	9.4	6.4-13.4	2.8	9.1	4.9-15.7	3.6	9.5	5.6-15.5
2016-2020	20.2	29.7	23.9-36.7	10.8	33.7	24.9-44.8	9.4	26.0	18.7-35.6	6.2	8.2	5.5-12.0	3.2	8.7	4.9-14.7	3.0	8.0	4.4-14.1	
Rectum & rectosigmoid	1976-1980	7.8	21.5	15.2-29.5	5.2	33.6	21.2-49.9	2.6	12.6	6.6-21.4	2.0	5.2	2.5-9.7	1.2	6.2	2.1-13.8	0.8	3.9	1.1-9.8
	1981-1985	5.8	14.1	9.4-20.3	3.2	19.6	10.7-32.4	2.6	11.1	5.9-19.1	1.8	4.6	2.1-8.8	1.2	7.4	2.5-16.2	0.6	2.6	0.5-7.5
	1986-1990	9.2	20.9	15.2-27.9	4.8	25.4	15.7-38.5	4.4	17.7	11.0-26.7	2.4	5.1	2.6-8.9	0.8	3.2	0.9-8.6	1.6	6.1	2.6-12.0
	1991-1995	7.4	15.7	11.0-21.7	3.6	18.7	10.6-30.0	3.8	14.0	8.4-21.9	3.2	6.7	3.8-10.9	1.2	7.0	2.3-15.4	2.0	7.2	3.4-13.3
	1996-2000	9.0	17.0	12.4-22.8	5.6	23.5	15.3-34.4	3.4	11.4	6.6-18.4	2.0	3.9	1.9-7.2	1.2	6.2	2.1-13.8	0.8	2.7	0.7-7.1
	2001-2005	8.4	14.4	10.4-19.6	5.2	20.4	13.1-30.4	3.2	10.1	5.7-16.6	2.6	4.4	2.4-7.7	1.8	6.4	2.9-12.4	0.8	2.4	0.7-6.6
	2006-2010	9.0	14.0	10.2-18.9	6.6	21.9	15.0-31.2	2.4	6.7	3.5-12.2	1.8	2.9	1.3-5.7	1.2	4.7	1.6-10.5	0.6	1.9	0.4-5.9
	2011-2015	7.0	10.1	6.9-14.3	4.0	12.9	7.7-20.5	3.0	7.4	4.1-12.8	1.6	2.2	0.9-4.6	0.6	2.2	0.5-6.7	1.0	2.4	0.8-6.4
2016-2020	9.4	14.1	10.1-19.2	5.4	16.2	10.4-24.5	4.0	11.8	6.9-19.1	2.2	2.9	1.4-5.5	1.2	3.3	1.2-8.0	1.0	2.5	0.8-6.7	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Tioga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	1.0	2.7	0.9-6.2	0.8	5.2	1.4-13.1	0.2	1.0	0.0-5.3	0.6	1.9	0.4-5.2	0.2	1.7	0.0-8.0	0.4	2.2	0.3-7.3
	1981-1985	0.4	0.9	0.1-3.3	0.2	0.9	0.0-5.4	0.2	0.9	0.0-4.7	0.4	0.9	0.1-3.3	0.2	0.9	0.0-5.4	0.2	0.9	0.0-4.7
	1986-1990	0.8	1.7	0.5-4.4	0.4	2.1	0.3-7.3	0.4	1.6	0.2-5.7	0.4	0.8	0.1-3.0	0.2	1.0	0.0-5.6	0.2	0.7	0.0-4.1
	1991-1995	1.0	2.1	0.7-4.8	0.8	4.4	1.0-11.3	0.2	0.8	0.0-4.3	1.2	2.5	0.9-5.3	0.8	4.2	1.0-11.0	0.4	1.7	0.2-5.8
	1996-2000	2.0	3.7	1.8-6.8	1.8	7.5	3.2-14.7	0.2	0.7	0.0-4.1	1.6	3.1	1.3-6.0	1.2	5.4	1.8-12.1	0.4	1.3	0.2-5.0
	2001-2005	2.8	4.9	2.6-8.3	2.0	7.7	3.6-14.5	0.8	2.8	0.7-7.3	2.4	4.2	2.1-7.3	1.8	7.0	3.1-13.6	0.6	1.8	0.4-5.7
	2006-2010	2.6	4.0	2.1-7.0	2.6	8.7	4.5-15.2	0.0	0.0	0.0-3.0	3.4	5.1	3.0-8.4	2.6	8.2	4.3-14.4	0.8	2.2	0.6-6.1
	2011-2015	6.6	9.6	6.6-13.7	4.0	12.7	7.6-20.1	2.6	6.7	3.5-12.0	3.8	5.5	3.3-8.9	2.2	7.2	3.5-13.2	1.6	4.1	1.8-8.8
	2016-2020	5.4	6.9	4.5-10.3	4.2	11.2	6.9-17.8	1.2	3.1	1.1-7.6	3.0	4.1	2.2-7.2	2.4	7.5	3.7-13.7	0.6	1.3	0.3-5.0
Pancreas	1976-1980	5.8	15.1	10.0-21.8	3.6	21.1	12.1-33.8	2.2	10.1	5.0-18.1	5.8	15.6	10.3-22.4	3.6	22.3	12.7-35.8	2.2	10.2	5.0-18.2
	1981-1985	4.2	9.9	6.1-15.2	2.4	12.7	6.3-22.5	1.8	7.8	3.5-14.8	3.8	9.2	5.5-14.4	2.0	11.2	5.2-20.6	1.8	7.9	3.6-15.0
	1986-1990	2.8	7.1	3.9-11.8	1.8	12.8	5.6-24.0	1.0	4.4	1.4-10.0	2.6	6.4	3.4-10.8	1.8	12.2	5.2-23.1	0.8	3.3	0.9-8.4
	1991-1995	6.2	13.2	9.0-18.8	4.2	21.7	13.2-33.4	2.0	7.5	3.6-13.7	5.6	11.6	7.7-16.8	3.4	16.9	9.6-27.3	2.2	8.1	4.0-14.5
	1996-2000	4.0	7.7	4.7-11.9	2.0	8.1	3.8-15.0	2.0	6.9	3.3-12.8	3.4	6.7	3.9-10.7	1.6	7.1	2.9-14.3	1.8	6.1	2.8-11.7
	2001-2005	7.6	12.8	9.0-17.6	4.0	14.1	8.6-22.0	3.6	11.4	6.7-18.2	7.2	12.2	8.5-17.0	3.6	12.7	7.5-20.3	3.6	11.3	6.7-18.2
	2006-2010	8.4	12.9	9.3-17.6	3.4	11.3	6.5-18.4	5.0	13.8	8.9-20.8	5.6	8.8	5.8-12.8	1.6	5.2	2.2-10.6	4.0	11.1	6.7-17.5
	2011-2015	8.4	12.0	8.6-16.5	3.4	9.9	5.7-16.4	5.0	13.4	8.5-20.3	6.8	9.6	6.6-13.7	3.2	9.5	5.4-16.0	3.6	9.6	5.6-15.8
	2016-2020	7.0	9.4	6.5-13.5	3.6	10.8	6.3-17.8	3.4	8.0	4.7-13.5	6.6	9.1	6.2-13.3	3.6	11.2	6.4-18.5	3.0	7.0	3.9-12.4
Larynx	1976-1980	1.8	4.3	2.0-8.3	1.4	8.0	3.1-16.6	0.4	1.7	0.2-6.1	0.4	1.1	0.1-3.9	0.2	0.9	0.0-5.5	0.2	1.1	0.0-5.6
	1981-1985	1.6	3.9	1.7-7.6	1.6	8.7	3.6-17.3	0.0	0.0	0.0-3.0	0.8	1.7	0.5-4.5	0.8	3.9	1.0-10.3	0.0	0.0	0.0-3.0
	1986-1990	2.4	4.9	2.5-8.6	2.0	9.4	4.3-17.6	0.4	1.5	0.2-5.5	0.6	1.3	0.3-3.9	0.6	2.9	0.6-8.7	0.0	0.0	0.0-2.9
	1991-1995	2.2	4.4	2.2-8.0	2.0	9.3	4.3-17.3	0.2	0.8	0.0-4.4	0.6	1.2	0.2-3.6	0.6	3.0	0.5-8.8	0.0	0.0	0.0-2.8
	1996-2000	3.0	5.6	3.1-9.3	2.2	8.9	4.4-16.2	0.8	2.8	0.8-7.3	0.4	0.8	0.1-2.9	0.4	1.9	0.2-6.8	0.0	0.0	0.0-2.8
	2001-2005	2.2	3.7	1.9-6.8	1.6	6.3	2.6-12.8	0.6	1.8	0.4-5.7	0.2	0.4	0.0-2.1	0.2	0.9	0.0-5.0	0.0	0.0	0.0-2.9
	2006-2010	2.8	4.6	2.5-7.8	1.6	5.8	2.4-11.7	1.2	3.7	1.3-8.5	0.4	0.7	0.1-2.7	0.2	0.7	0.0-4.4	0.2	0.5	0.0-3.7
	2011-2015	2.6	4.1	2.2-7.2	2.2	7.0	3.5-13.0	0.4	1.4	0.2-5.4	0.2	0.4	0.0-2.3	0.2	0.8	0.0-4.6	0.0	0.0	0.0-3.1
	2016-2020	2.2	2.7	1.3-5.2	2.2	5.6	2.8-10.8	0.0	0.0	0.0-3.3	0.4	0.5	0.1-2.3	0.4	1.0	0.1-4.7	0.0	0.0	0.0-3.3
Lung and bronchus	1976-1980	27.4	69.3	58.0-82.1	21.4	126.5	102.7-153.8	6.0	27.1	18.2-38.7	21.0	52.7	43.0-64.0	17.2	98.8	78.1-122.9	3.8	17.9	10.7-27.8
	1981-1985	24.2	56.7	46.9-67.9	17.4	95.2	75.4-118.4	6.8	28.1	19.4-39.4	20.4	48.0	39.0-58.3	15.0	83.2	64.7-105.0	5.4	22.3	14.6-32.5
	1986-1990	38.0	82.7	71.2-95.5	24.8	129.1	106.0-155.4	13.2	51.6	39.9-65.7	26.8	58.7	49.0-69.6	16.8	87.5	68.6-109.6	10.0	39.2	29.0-51.7
	1991-1995	32.6	67.4	57.4-78.7	21.0	105.5	85.1-128.9	11.6	43.7	33.2-56.5	28.2	58.6	49.2-69.1	16.6	83.8	65.8-105.0	11.6	43.5	33.0-56.3
	1996-2000	36.4	68.0	58.4-78.7	22.0	90.6	74.1-109.8	14.4	49.0	38.3-61.8	29.2	55.4	46.7-65.2	18.0	77.6	61.8-96.2	11.2	38.4	29.0-49.9
	2001-2005	36.6	63.0	54.2-73.0	19.4	75.0	60.4-92.0	17.2	54.5	43.6-67.5	30.2	52.4	44.4-61.6	17.0	67.8	53.8-84.3	13.2	41.5	32.1-53.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Tioga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	46.6	73.0	63.9-83.2	26.4	90.6	75.5-108.0	20.2	61.1	49.6-74.6	31.4	49.5	42.0-58.0	19.2	66.4	53.5-81.6	12.2	37.8	28.8-48.9
	2011-2015	44.8	64.2	55.9-73.4	22.6	68.3	56.1-82.6	22.2	61.5	50.3-74.6	29.0	41.5	34.9-49.1	14.4	45.0	35.1-57.2	14.6	39.7	31.0-50.5
	2016-2020	44.2	58.3	50.7-66.8	23.6	65.8	54.3-79.4	20.6	51.5	41.8-63.3	25.2	33.1	27.5-39.7	13.4	38.3	29.6-49.2	11.8	28.5	21.6-37.5
Melanoma of the skin	1976-1980	2.6	6.0	3.2-10.2	1.2	5.5	2.0-12.3	1.4	6.4	2.5-13.1	0.4	0.9	0.1-3.4	0.2	0.9	0.0-5.5	0.2	1.0	0.0-5.2
	1981-1985	3.8	8.5	5.0-13.3	1.6	9.6	3.8-19.4	2.2	9.1	4.5-16.4	0.4	0.8	0.1-3.0	0.4	1.7	0.2-6.5	0.0	0.0	0.0-3.0
	1986-1990	6.0	12.3	8.3-17.7	2.8	12.9	6.8-22.0	3.2	12.3	7.0-20.0	1.2	2.4	0.8-5.2	1.0	4.1	1.3-10.1	0.2	0.9	0.0-4.6
	1991-1995	6.8	13.3	9.2-18.6	3.0	12.4	6.8-20.8	3.8	14.7	8.8-22.9	0.8	1.7	0.5-4.3	0.6	2.6	0.5-7.8	0.2	0.8	0.0-4.4
	1996-2000	7.6	14.3	10.1-19.7	5.2	19.5	12.7-28.7	2.4	8.6	4.4-15.2	2.2	4.2	2.1-7.5	1.8	7.1	3.2-13.7	0.4	1.3	0.2-5.1
	2001-2005	9.4	16.2	11.9-21.6	6.8	25.5	17.5-36.0	2.6	8.6	4.6-14.9	2.0	3.4	1.6-6.3	1.4	5.6	2.1-11.9	0.6	1.9	0.4-5.9
	2006-2010	15.2	26.0	20.3-32.7	8.0	28.5	20.2-39.2	7.2	25.7	17.8-36.0	2.0	3.0	1.4-5.7	1.6	4.9	2.1-10.1	0.4	1.1	0.1-4.7
	2011-2015	17.0	28.8	22.7-36.1	11.0	39.1	29.1-51.5	6.0	20.1	13.2-29.5	1.4	2.5	1.0-5.5	0.2	0.7	0.0-4.4	1.2	4.4	1.5-10.1
2016-2020	19.2	29.7	23.7-36.9	11.8	38.9	29.1-51.0	7.4	21.4	14.6-30.6	1.2	2.2	0.7-5.0	0.8	2.9	0.8-7.9	0.4	1.8	0.2-6.7	
Female breast	1976-1980							22.2	104.1	85.5-125.4							6.4	30.6	20.8-43.1
	1981-1985							22.4	98.9	81.3-119.1							7.0	29.9	20.7-41.6
	1986-1990							27.8	112.2	94.2-132.6							7.6	31.1	21.9-42.7
	1991-1995							33.4	126.5	108.0-147.3							10.6	39.7	29.7-51.9
	1996-2000							33.8	118.6	101.3-137.9							6.6	22.2	15.3-31.4
	2001-2005							37.0	120.6	103.7-139.5							7.2	23.2	16.2-32.4
	2006-2010							42.6	131.2	113.8-150.6							6.6	20.1	13.7-28.7
	2011-2015							43.4	126.0	109.1-145.0							4.4	12.3	7.4-19.3
2016-2020							51.6	151.6	132.4-173.0							6.8	18.7	12.6-27.1	
Cervix uteri	1976-1980							3.0	14.6	8.1-24.0							1.6	8.4	3.6-16.2
	1981-1985							2.8	11.8	6.4-19.9							0.8	3.5	0.9-8.8
	1986-1990							2.4	9.4	4.8-16.5							1.4	5.7	2.3-11.8
	1991-1995							1.8	6.7	3.1-12.8							0.4	1.4	0.2-5.3
	1996-2000							2.0	7.2	3.4-13.3							0.0	0.0	0.0-2.8
	2001-2005							2.0	7.3	3.4-13.6							0.4	1.4	0.2-5.4
	2006-2010							3.0	11.3	6.2-19.0							1.2	4.6	1.6-10.4
	2011-2015							1.4	4.8	1.7-10.5							0.2	0.6	0.0-3.9
	2016-2020							2.2	9.2	4.4-17.0							1.2	3.8	1.3-9.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Tioga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							5.8	26.4	17.6-37.9							1.0	4.4	1.4-10.3
	1981-1985							5.8	24.8	16.6-35.7							1.0	4.3	1.4-10.0
	1986-1990							6.6	27.7	19.0-38.9							1.4	5.4	2.1-11.2
	1991-1995							4.2	16.1	10.0-24.6							1.4	5.2	2.1-10.8
	1996-2000							7.2	25.2	17.6-34.9							0.6	2.0	0.4-6.1
	2001-2005							11.0	36.0	27.1-47.2							2.0	6.2	3.0-11.7
	2006-2010							8.8	26.3	19.0-35.8							1.4	4.1	1.6-8.8
	2011-2015							11.6	31.4	23.7-41.2							0.6	1.5	0.3-5.1
	2016-2020							13.0	34.9	26.4-45.5							1.4	3.7	1.4-8.5
Ovary	1976-1980							4.4	20.5	12.8-31.0							2.4	11.2	5.8-19.5
	1981-1985							3.8	15.6	9.3-24.4							2.6	10.7	5.7-18.4
	1986-1990							3.8	16.1	9.6-25.0							2.6	10.9	5.8-18.5
	1991-1995							5.2	19.6	12.8-28.8							4.0	15.0	9.1-23.2
	1996-2000							3.2	11.2	6.4-18.3							3.4	12.2	7.1-19.6
	2001-2005							3.8	12.7	7.6-20.1							1.8	6.0	2.7-11.7
	2006-2010							3.4	11.0	6.3-18.0							2.0	6.1	2.9-11.5
	2011-2015							3.2	10.1	5.6-17.0							2.2	5.8	2.9-11.0
	2016-2020							3.0	8.2	4.4-14.2							1.6	4.1	1.7-8.8
Prostate	1976-1980				13.0	99.2	75.3-127.5							3.6	31.5	18.1-49.7			
	1981-1985				14.8	100.3	77.6-126.8							4.2	30.8	18.6-47.1			
	1986-1990				14.4	84.1	64.8-107.0							4.8	29.4	18.2-44.4			
	1991-1995				40.6	199.8	171.9-230.6							5.4	35.0	22.6-51.1			
	1996-2000				41.0	175.1	151.1-201.9							8.2	43.8	30.8-59.9			
	2001-2005				49.8	187.4	164.3-212.9							5.4	27.2	17.6-39.8			
	2006-2010				49.2	160.1	140.4-182.0							5.0	20.2	12.9-30.1			
	2011-2015				39.8	113.8	98.3-131.4							4.8	16.5	10.5-24.8			
	2016-2020				42.2	108.6	94.2-124.9							4.6	13.1	8.3-20.2			
Testis	1976-1980				0.8	3.1	0.8-8.8							0.0	0.0	0.0-4.1			
	1981-1985				1.6	6.5	2.7-13.2							0.4	1.7	0.2-6.6			
	1986-1990				0.8	2.8	0.8-7.7							0.2	0.7	0.0-4.8			
	1991-1995				2.2	8.4	4.1-15.3							0.0	0.0	0.0-3.4			
	1996-2000				1.6	6.4	2.7-12.7							0.0	0.0	0.0-3.3			
	2001-2005				1.4	6.8	2.7-13.9							0.2	0.6	0.0-4.2			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Tioga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				2.6	11.9	6.2-20.3							0.0	0.0	0.0-3.1			
	2011-2015				3.2	15.1	8.5-24.5							0.0	0.0	0.0-3.1			
	2016-2020				1.6	8.0	3.4-15.7							0.0	0.0	0.0-3.3			
Urinary bladder (incl. in situ)	1976-1980	9.8	26.1	19.2-34.6	8.0	50.1	35.0-69.0	1.8	8.4	3.8-15.9	2.4	6.0	3.1-10.6	1.4	7.1	2.7-15.0	1.0	4.8	1.5-11.1
	1981-1985	9.2	22.5	16.4-30.1	6.6	36.7	24.6-52.2	2.6	11.8	6.3-20.1	2.8	7.8	4.2-13.0	1.6	11.1	4.5-21.8	1.2	5.7	2.1-12.2
	1986-1990	11.2	25.1	18.9-32.7	7.8	44.4	30.7-61.6	3.4	13.1	7.6-21.0	1.6	3.7	1.6-7.3	1.4	8.7	3.4-17.8	0.2	0.7	0.0-4.2
	1991-1995	14.0	28.9	22.5-36.6	10.6	51.7	38.0-68.4	3.4	12.8	7.5-20.5	2.8	6.4	3.5-10.7	1.8	12.2	5.3-23.0	1.0	3.8	1.2-8.9
	1996-2000	13.0	25.2	19.4-32.2	9.6	46.8	33.8-62.8	3.4	11.8	6.9-19.1	3.0	6.0	3.3-9.9	2.4	13.5	6.7-23.8	0.6	2.0	0.4-6.1
	2001-2005	17.8	31.1	24.9-38.3	14.2	56.4	43.7-71.6	3.6	11.3	6.7-18.1	3.6	6.7	4.0-10.6	2.4	11.7	5.8-20.7	1.2	3.5	1.3-8.0
	2006-2010	17.2	27.3	21.8-33.9	13.2	46.7	35.9-59.9	4.0	11.6	7.1-18.3	4.2	6.9	4.3-10.7	2.6	10.0	5.2-17.4	1.6	4.7	2.0-9.6
	2011-2015	18.6	26.3	21.1-32.4	14.0	43.4	33.7-55.4	4.6	11.8	7.5-18.3	2.8	3.9	2.1-6.8	2.0	6.4	3.0-12.2	0.8	1.8	0.5-5.6
	2016-2020	21.4	28.1	22.9-34.3	17.2	48.9	38.9-61.0	4.2	10.1	6.2-16.2	2.6	3.5	1.9-6.3	1.8	5.0	2.2-10.1	0.8	2.1	0.6-6.3
Kidney and renal pelvis	1976-1980	2.4	7.0	3.5-12.1	1.2	8.1	2.8-17.4	1.2	5.9	2.2-12.8	2.2	6.6	3.2-11.6	1.6	10.4	4.3-20.5	0.6	3.3	0.7-9.1
	1981-1985	2.6	6.3	3.3-10.8	1.4	7.6	2.8-16.1	1.2	5.2	1.9-11.2	2.0	5.3	2.5-9.7	1.2	7.3	2.4-16.0	0.8	3.7	1.0-9.2
	1986-1990	4.8	10.6	6.8-15.9	3.4	17.5	9.9-28.5	1.4	5.6	2.2-11.6	2.4	5.5	2.8-9.6	1.8	10.1	4.3-19.7	0.6	2.4	0.5-6.9
	1991-1995	6.0	12.0	8.0-17.1	4.6	20.5	12.7-31.1	1.4	5.3	2.1-10.9	2.6	5.1	2.7-8.8	1.4	6.8	2.5-14.4	1.2	4.6	1.7-10.1
	1996-2000	7.8	14.9	10.6-20.4	3.4	14.2	8.1-23.1	4.4	15.3	9.5-23.2	1.0	2.1	0.7-4.9	0.6	3.2	0.6-8.9	0.4	1.5	0.2-5.3
	2001-2005	9.0	15.7	11.4-21.1	5.2	19.5	12.6-29.1	3.8	13.0	7.7-20.6	2.8	4.8	2.6-8.2	1.6	6.3	2.6-12.8	1.2	3.7	1.3-8.3
	2006-2010	12.8	20.2	15.5-26.0	7.6	25.0	17.6-34.7	5.2	15.3	9.9-22.8	2.8	4.6	2.5-7.9	1.8	6.0	2.7-11.7	1.0	2.6	0.8-6.7
	2011-2015	8.6	12.8	9.2-17.5	4.2	13.2	8.0-20.7	4.4	12.8	7.9-19.9	3.0	4.3	2.4-7.3	1.0	3.4	1.1-8.3	2.0	5.5	2.6-10.6
	2016-2020	12.0	16.8	12.7-22.1	8.8	25.1	18.0-34.4	3.2	9.0	4.9-15.5	1.8	2.6	1.1-5.3	1.2	3.8	1.3-8.9	0.6	1.5	0.3-5.3
Brain and other central nervous system	1976-1980	2.8	5.8	3.1-9.8	1.6	6.5	2.8-13.4	1.2	5.1	1.8-11.2	1.6	3.4	1.5-6.9	0.6	2.8	0.6-8.5	1.0	4.1	1.3-9.9
	1981-1985	2.2	4.9	2.4-8.8	1.0	4.3	1.4-10.5	1.2	5.1	1.8-11.1	2.6	6.0	3.2-10.3	1.2	5.4	1.9-12.1	1.4	6.2	2.5-12.7
	1986-1990	3.0	5.9	3.3-9.9	1.6	6.4	2.7-13.0	1.4	5.6	2.2-11.5	1.4	2.6	1.1-5.5	0.8	3.0	0.8-8.2	0.6	2.3	0.5-6.8
	1991-1995	3.2	6.2	3.5-10.1	2.2	8.1	4.1-14.9	1.0	3.7	1.2-8.7	1.8	3.5	1.6-6.6	1.0	3.7	1.2-8.9	0.8	3.1	0.8-7.9
	1996-2000	3.4	6.7	3.9-10.7	2.0	8.2	3.8-15.2	1.4	5.2	2.1-10.7	2.4	4.4	2.3-7.8	1.6	6.4	2.7-12.8	0.8	2.8	0.8-7.3
	2001-2005	4.6	8.4	5.3-12.7	3.4	13.7	7.8-22.1	1.2	4.2	1.5-9.4	3.6	6.4	3.8-10.3	2.2	8.8	4.3-16.0	1.4	4.6	1.8-9.7
	2006-2010	5.0	8.9	5.7-13.3	3.4	12.1	6.9-19.8	1.6	5.6	2.3-11.5	2.4	4.0	2.0-7.2	1.4	4.9	1.9-10.5	1.0	3.0	0.9-7.4
	2011-2015	3.0	5.4	3.0-9.1	2.4	9.3	4.8-16.5	0.6	2.1	0.4-6.6	3.4	5.4	3.1-8.9	2.6	9.3	4.8-16.3	0.8	2.1	0.6-6.1
	2016-2020	3.6	5.0	2.9-8.2	2.4	7.7	3.9-14.1	1.2	2.6	1.0-6.7	3.0	4.0	2.2-7.0	2.0	6.0	2.8-11.6	1.0	2.3	0.7-6.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Tioga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	0.6	1.3	0.3-3.8	0.2	1.0	0.0-5.8	0.4	1.6	0.2-6.1	0.0	0.0	0.0-1.8	0.0	0.0	0.0-4.1	0.0	0.0	0.0-3.2
	1981-1985	2.0	4.4	2.1-8.1	0.8	3.3	0.9-9.0	1.2	5.4	2.0-11.8	0.0	0.0	0.0-1.6	0.0	0.0	0.0-3.9	0.0	0.0	0.0-3.0
	1986-1990	1.8	3.7	1.7-7.0	0.4	1.7	0.2-6.5	1.4	5.3	2.1-11.0	0.4	0.9	0.1-3.3	0.2	0.7	0.0-4.8	0.2	0.8	0.0-4.4
	1991-1995	1.8	3.3	1.5-6.3	0.4	1.6	0.2-6.0	1.4	5.0	2.0-10.3	0.0	0.0	0.0-1.5	0.0	0.0	0.0-3.4	0.0	0.0	0.0-2.8
	1996-2000	3.6	6.7	4.0-10.7	1.2	4.5	1.6-10.0	2.4	8.9	4.6-15.6	0.0	0.0	0.0-1.5	0.0	0.0	0.0-3.3	0.0	0.0	0.0-2.8
	2001-2005	6.0	11.5	7.7-16.5	2.0	7.3	3.5-13.6	4.0	15.8	9.5-24.6	0.4	0.7	0.1-2.6	0.4	1.6	0.2-5.9	0.0	0.0	0.0-2.9
	2006-2010	8.4	15.3	10.9-20.9	2.0	7.6	3.5-14.3	6.4	22.7	15.3-32.5	0.4	0.7	0.1-2.7	0.0	0.0	0.0-3.1	0.4	1.2	0.1-4.8
	2011-2015	12.2	23.3	17.5-30.3	3.8	12.6	7.4-20.3	8.4	34.3	24.2-46.9	0.2	0.3	0.0-2.1	0.0	0.0	0.0-3.1	0.2	0.6	0.0-4.1
	2016-2020	10.4	20.6	15.1-27.5	3.2	13.0	7.2-21.6	7.2	28.4	19.3-40.3	0.2	0.3	0.0-2.0	0.2	0.5	0.0-4.1	0.0	0.0	0.0-3.3
Hodgkin lymphoma	1976-1980	1.2	2.2	0.8-5.0	1.0	3.9	1.2-9.9	0.2	0.6	0.0-4.1	0.4	0.9	0.1-3.4	0.4	1.9	0.2-7.2	0.0	0.0	0.0-3.2
	1981-1985	2.2	4.5	2.2-8.1	1.4	5.6	2.2-12.1	0.8	3.3	0.9-8.5	0.6	1.3	0.3-3.8	0.2	1.1	0.0-5.9	0.4	1.5	0.2-5.7
	1986-1990	2.4	4.9	2.5-8.6	1.2	5.2	1.9-11.5	1.2	4.7	1.7-10.3	1.2	2.7	1.0-5.9	0.8	3.3	0.9-8.8	0.4	1.6	0.2-5.8
	1991-1995	2.2	4.5	2.2-8.0	1.0	4.4	1.4-10.4	1.2	4.6	1.7-9.9	0.6	1.5	0.3-4.1	0.4	3.1	0.3-9.9	0.2	0.7	0.0-4.0
	1996-2000	2.2	4.4	2.2-8.0	0.6	3.1	0.5-9.1	1.6	6.0	2.5-11.9	0.6	1.2	0.2-3.6	0.4	2.4	0.2-8.2	0.2	0.7	0.0-4.1
	2001-2005	1.8	3.3	1.5-6.3	1.0	3.7	1.2-9.0	0.8	2.6	0.7-7.0	0.2	0.4	0.0-2.1	0.2	0.9	0.0-5.0	0.0	0.0	0.0-2.9
	2006-2010	1.0	1.6	0.5-3.8	0.6	2.0	0.4-6.2	0.4	1.2	0.1-4.8	0.0	0.0	0.0-1.5	0.0	0.0	0.0-3.1	0.0	0.0	0.0-3.0
	2011-2015	1.8	3.2	1.4-6.3	1.0	3.6	1.1-8.8	0.8	2.9	0.7-7.7	0.0	0.0	0.0-1.6	0.0	0.0	0.0-3.1	0.0	0.0	0.0-3.1
	2016-2020	0.2	0.5	0.0-2.7	0.2	1.0	0.0-5.3	0.0	0.0	0.0-3.3	0.0	0.0	0.0-1.6	0.0	0.0	0.0-3.3	0.0	0.0	0.0-3.3
Non-Hodgkin lymphomas	1976-1980	4.8	12.0	7.6-17.9	2.4	11.4	5.8-20.4	2.4	11.9	6.1-20.7	2.4	6.5	3.3-11.3	1.4	9.2	3.4-19.3	1.0	4.8	1.5-11.0
	1981-1985	4.6	11.1	7.0-16.6	2.8	15.6	8.2-26.5	1.8	7.6	3.5-14.4	2.2	5.6	2.8-10.0	0.8	5.5	1.3-13.8	1.4	6.2	2.5-12.8
	1986-1990	7.0	15.7	10.9-21.9	4.2	21.5	12.8-33.5	2.8	11.0	6.0-18.5	3.8	8.4	5.0-13.2	1.8	9.1	4.0-17.4	2.0	7.9	3.8-14.5
	1991-1995	6.8	13.8	9.5-19.3	3.8	17.9	10.5-28.4	3.0	11.1	6.2-18.3	4.0	8.6	5.2-13.3	2.6	14.2	7.2-24.6	1.4	5.0	2.0-10.5
	1996-2000	10.6	20.3	15.1-26.6	5.2	22.0	14.1-32.6	5.4	18.6	12.3-27.2	4.4	8.7	5.4-13.3	2.6	12.4	6.4-21.4	1.8	6.0	2.7-11.5
	2001-2005	11.4	20.1	15.2-26.1	4.8	17.9	11.4-26.9	6.6	22.0	15.1-31.1	3.0	5.3	2.9-8.8	1.2	4.5	1.6-10.0	1.8	5.6	2.5-10.9
	2006-2010	11.8	18.9	14.3-24.6	7.0	25.4	17.5-35.7	4.8	14.0	8.9-21.2	3.6	6.1	3.6-9.7	1.4	6.1	2.4-12.6	2.2	6.8	3.3-12.6
	2011-2015	12.2	19.4	14.7-25.2	7.2	24.9	17.2-35.0	5.0	14.1	9.1-21.4	5.2	7.9	5.1-11.7	3.0	9.5	5.3-16.0	2.2	6.2	3.1-11.6
	2016-2020	11.8	17.1	12.7-22.5	6.4	17.9	12.1-25.9	5.4	16.8	10.6-25.4	3.6	4.5	2.6-7.4	2.6	7.2	3.8-12.9	1.0	2.3	0.8-6.4
Myeloma	1976-1980	1.6	4.5	1.9-8.7	1.4	9.4	3.6-19.3	0.2	0.8	0.0-4.6	0.6	1.4	0.3-4.1	0.6	2.9	0.6-8.9	0.0	0.0	0.0-3.2
	1981-1985	2.0	5.2	2.4-9.4	1.6	10.6	4.3-20.8	0.4	1.6	0.2-5.9	1.4	3.8	1.5-7.7	1.4	10.0	3.8-20.4	0.0	0.0	0.0-3.0
	1986-1990	2.2	5.1	2.5-9.2	1.4	7.9	2.9-16.4	0.8	3.1	0.9-8.1	1.8	4.1	1.8-7.7	1.0	4.9	1.5-11.5	0.8	3.2	0.9-8.1
	1991-1995	2.8	5.6	3.1-9.5	1.6	7.6	3.0-15.4	1.2	4.4	1.6-9.6	1.0	2.2	0.7-5.1	0.4	2.6	0.2-9.0	0.6	2.1	0.4-6.4
	1996-2000	4.2	8.0	5.0-12.3	2.6	12.8	6.5-22.1	1.6	5.6	2.4-11.1	3.6	7.0	4.2-11.2	2.4	12.5	6.2-22.0	1.2	4.2	1.5-9.3
	2001-2005	3.6	6.5	3.8-10.3	2.2	10.2	4.9-18.4	1.4	4.3	1.7-9.3	2.4	4.3	2.2-7.5	2.0	8.8	4.1-16.5	0.4	1.3	0.2-5.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Tioga**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	3.6	5.8	3.5-9.4	1.4	5.1	2.0-10.8	2.2	6.7	3.4-12.4	2.6	4.1	2.2-7.2	0.8	3.0	0.8-8.0	1.8	5.0	2.3-9.9
	2011-2015	5.8	8.9	5.9-12.9	3.4	11.8	6.8-19.2	2.4	6.3	3.3-11.7	1.6	2.4	1.0-5.0	1.2	4.1	1.5-9.3	0.4	1.2	0.1-4.9
	2016-2020	4.8	6.3	4.0-9.7	1.8	5.1	2.3-10.3	3.0	7.8	4.3-13.6	1.6	2.1	0.9-4.5	1.4	4.1	1.6-9.0	0.2	0.5	0.0-4.0
Leukemias	1976-1980	4.6	11.8	7.3-17.8	1.8	10.2	4.4-19.8	2.8	12.8	7.0-21.5	2.8	6.6	3.5-11.3	1.0	5.3	1.5-13.2	1.8	8.0	3.7-15.3
	1981-1985	6.4	14.7	10.0-20.8	4.2	22.2	13.4-34.3	2.2	8.8	4.4-15.9	2.2	5.1	2.5-9.2	1.6	7.9	3.3-15.9	0.6	2.4	0.5-7.1
	1986-1990	7.0	15.4	10.6-21.4	5.4	27.4	17.5-40.6	1.6	6.4	2.8-12.6	3.2	7.4	4.2-12.0	1.8	10.5	4.4-20.4	1.4	5.5	2.2-11.3
	1991-1995	8.6	16.9	12.2-22.9	4.4	19.0	11.6-29.3	4.2	15.2	9.4-23.3	3.8	7.9	4.7-12.4	2.6	13.0	6.6-22.6	1.2	4.4	1.6-9.6
	1996-2000	8.8	16.5	12.0-22.2	6.8	29.2	19.8-41.4	2.0	6.7	3.2-12.6	5.2	10.0	6.5-14.7	3.4	16.6	9.4-26.9	1.8	6.2	2.8-11.8
	2001-2005	9.2	16.8	12.3-22.4	5.6	22.3	14.7-32.5	3.6	11.1	6.6-17.9	4.2	7.6	4.7-11.6	2.2	9.7	4.7-17.6	2.0	6.2	3.0-11.8
	2006-2010	10.6	17.4	13.0-23.0	6.6	22.9	15.6-32.7	4.0	12.3	7.4-19.4	4.0	7.0	4.2-10.9	2.4	9.2	4.7-16.2	1.6	4.6	2.0-9.5
	2011-2015	11.6	17.9	13.4-23.5	6.6	21.9	14.8-31.3	5.0	14.2	9.0-21.7	3.2	4.8	2.7-8.0	1.8	5.3	2.4-10.6	1.4	4.1	1.5-9.1
2016-2020	10.6	16.0	11.8-21.4	7.0	22.0	15.0-31.3	3.6	10.9	6.2-18.1	3.2	4.3	2.5-7.3	2.2	6.6	3.3-12.4	1.0	2.5	0.8-6.7	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Tompkins**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	217.2	375.4	352.9-398.9	102.6	425.3	387.3-465.7	114.6	349.3	320.7-379.7	105.4	187.5	171.6-204.5	53.4	238.2	209.2-269.7	52.0	157.4	138.6-177.9
	1981-1985	256.0	423.0	399.7-447.2	122.0	486.9	447.4-528.7	134.0	391.1	361.5-422.5	119.8	199.8	183.9-216.6	62.4	255.6	226.9-286.8	57.4	164.4	145.7-184.9
	1986-1990	293.8	451.3	428.2-475.3	142.6	539.7	499.2-582.4	151.2	408.8	379.6-439.6	134.0	209.7	194.0-226.3	66.4	259.3	231.2-289.7	67.6	178.6	159.8-198.9
	1991-1995	335.6	475.7	453.1-499.2	169.2	580.5	540.8-622.2	166.4	415.7	387.6-445.3	140.8	204.7	189.8-220.4	71.4	268.1	240.1-298.1	69.4	171.6	153.8-190.8
	1996-2000	370.2	493.2	470.9-516.3	187.0	582.7	545.2-622.0	183.2	431.5	403.7-460.7	143.4	192.1	178.3-206.7	68.0	228.2	204.0-254.3	75.4	171.9	154.8-190.4
	2001-2005	397.6	484.4	463.1-506.4	209.2	585.5	549.9-622.8	188.4	415.5	388.9-443.4	141.2	172.7	160.1-186.0	66.2	197.9	176.8-220.7	75.0	156.9	141.1-174.0
	2006-2010	463.2	509.6	488.6-531.4	238.0	581.8	548.2-616.9	225.2	455.6	428.6-483.9	144.6	159.2	147.6-171.5	69.4	183.9	164.7-204.7	75.2	144.0	129.3-159.8
	2011-2015	442.6	446.2	427.1-465.9	209.2	462.6	433.9-492.7	233.4	439.2	413.4-466.3	154.8	154.2	143.2-165.8	74.2	171.2	153.7-190.1	80.6	143.4	129.2-158.7
	2016-2020	480.8	440.9	422.8-459.6	245.6	479.9	452.5-508.7	235.2	412.3	388.1-437.8	148.2	135.0	125.2-145.4	74.8	155.8	139.9-173.0	73.4	120.4	108.0-133.9
Oral cavity and pharynx	1976-1980	4.2	7.2	4.4-11.0	2.8	11.4	6.0-19.3	1.4	4.5	1.8-9.3	2.0	3.4	1.6-6.3	1.4	6.3	2.4-12.9	0.6	1.8	0.4-5.2
	1981-1985	4.6	7.4	4.6-11.1	3.6	13.5	7.7-21.6	1.0	3.1	1.0-7.1	1.2	1.9	0.7-4.2	1.2	5.0	1.7-11.0	0.0	0.0	0.0-2.1
	1986-1990	6.0	9.0	6.0-12.8	3.0	10.6	5.8-17.7	3.0	7.9	4.4-13.1	1.6	2.6	1.1-5.0	0.4	1.4	0.2-4.9	1.2	3.1	1.2-6.8
	1991-1995	6.4	8.9	6.0-12.5	4.0	12.4	7.4-19.2	2.4	5.5	2.8-9.7	2.4	3.4	1.8-6.0	0.8	3.0	0.7-7.5	1.6	3.9	1.7-7.7
	1996-2000	7.0	9.3	6.5-13.0	5.6	16.5	10.9-23.9	1.4	3.5	1.4-7.1	1.6	2.1	0.9-4.1	0.8	2.3	0.6-6.0	0.8	1.8	0.5-4.6
	2001-2005	9.2	11.1	8.1-14.8	6.2	16.0	10.8-22.9	3.0	6.7	3.7-11.1	2.0	2.5	1.2-4.6	1.6	4.7	2.0-9.2	0.4	0.8	0.1-3.0
	2006-2010	8.4	8.9	6.4-12.2	5.6	12.4	8.1-18.1	2.8	5.7	3.0-9.7	0.8	0.8	0.2-2.1	0.8	1.6	0.4-4.3	0.0	0.0	0.0-1.8
	2011-2015	11.4	11.0	8.2-14.5	7.0	13.8	9.4-19.5	4.4	8.4	5.1-13.0	2.0	1.7	0.8-3.3	1.6	3.0	1.3-6.3	0.4	0.6	0.1-2.7
	2016-2020	12.2	11.2	8.5-14.5	8.4	16.5	11.7-22.6	3.8	6.3	3.7-10.3	1.4	1.2	0.5-2.6	1.0	1.9	0.6-4.8	0.4	0.7	0.1-2.8
Esophagus	1976-1980	1.6	2.7	1.1-5.2	1.2	4.6	1.6-10.0	0.4	1.1	0.1-4.2	1.6	2.5	1.1-5.0	1.2	4.3	1.6-9.4	0.4	1.2	0.1-4.2
	1981-1985	1.2	2.1	0.7-4.4	0.8	3.7	1.0-9.1	0.4	1.1	0.1-3.9	1.2	2.2	0.8-4.8	0.8	3.7	1.0-9.2	0.4	1.1	0.1-4.0
	1986-1990	4.2	6.7	4.1-10.2	2.4	8.9	4.5-15.6	1.8	4.8	2.2-9.2	2.4	3.6	1.9-6.3	1.2	4.5	1.5-9.8	1.2	3.1	1.1-6.8
	1991-1995	4.0	5.8	3.5-8.9	3.2	10.5	5.9-17.1	0.8	2.2	0.6-5.6	2.8	4.1	2.2-6.8	2.2	8.4	4.1-14.9	0.6	1.6	0.3-4.6
	1996-2000	4.2	5.7	3.5-8.7	3.6	11.1	6.5-17.5	0.6	1.4	0.3-4.1	3.6	4.9	2.9-7.8	3.0	9.9	5.5-16.3	0.6	1.3	0.3-4.0
	2001-2005	3.0	3.9	2.2-6.4	2.2	6.8	3.4-12.0	0.8	1.7	0.5-4.6	3.2	4.0	2.3-6.4	2.6	7.3	3.8-12.5	0.6	1.5	0.3-4.4
	2006-2010	5.6	6.4	4.2-9.3	4.6	11.5	7.2-17.4	1.0	2.3	0.7-5.5	4.6	5.3	3.4-8.1	4.0	10.8	6.5-16.7	0.6	1.3	0.3-4.1
	2011-2015	5.8	5.5	3.6-8.0	4.2	9.0	5.5-14.0	1.6	2.7	1.1-5.7	3.8	3.7	2.2-5.8	3.2	7.3	4.1-12.0	0.6	1.0	0.2-3.3
	2016-2020	7.6	6.5	4.6-9.1	6.8	12.5	8.5-17.8	0.8	1.4	0.4-3.8	4.8	4.1	2.6-6.3	3.0	5.8	3.1-9.8	1.8	2.7	1.2-5.4
Stomach	1976-1980	4.6	8.7	5.5-12.9	3.2	14.0	7.8-22.8	1.4	4.7	1.9-9.5	3.0	5.4	3.0-8.9	1.8	8.5	3.7-16.0	1.2	3.5	1.3-7.6
	1981-1985	6.6	11.2	7.7-15.7	3.4	14.1	8.0-22.7	3.2	9.0	5.1-14.6	5.4	9.3	6.1-13.5	2.2	9.2	4.4-16.6	3.2	9.1	5.2-14.8
	1986-1990	5.2	8.2	5.4-12.0	2.6	9.9	5.2-16.8	2.6	7.0	3.7-12.0	3.8	6.1	3.7-9.5	2.2	8.6	4.2-15.4	1.6	4.4	1.8-8.6
	1991-1995	5.4	7.9	5.2-11.4	4.4	15.9	9.8-24.2	1.0	2.6	0.8-6.0	3.2	4.6	2.6-7.5	2.0	8.0	3.7-14.7	1.2	3.0	1.1-6.5
	1996-2000	5.8	7.6	5.1-11.0	3.4	11.8	6.7-18.9	2.4	5.3	2.7-9.4	3.8	5.1	3.1-7.9	2.4	8.3	4.2-14.4	1.4	2.9	1.1-6.1
	2001-2005	4.6	5.7	3.6-8.6	3.4	9.9	5.7-15.9	1.2	2.6	0.9-5.8	2.6	3.1	1.6-5.3	1.6	4.8	2.0-9.4	1.0	1.8	0.6-4.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Tompkins**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	5.4	6.5	4.2-9.5	2.8	7.5	4.0-12.6	2.6	6.0	3.2-10.4	3.0	3.3	1.8-5.5	1.8	4.5	2.0-8.7	1.2	2.5	0.9-5.7
	2011-2015	4.8	5.0	3.2-7.6	2.4	5.7	2.9-10.1	2.4	4.4	2.2-7.9	1.6	1.8	0.8-3.6	0.6	1.5	0.3-4.3	1.0	2.0	0.6-4.8
	2016-2020	5.4	5.7	3.7-8.4	3.4	7.6	4.4-12.4	2.0	4.0	1.9-7.6	2.4	2.3	1.2-4.1	1.4	3.2	1.2-6.7	1.0	1.4	0.5-3.7
Colorectal	1976-1980	36.4	66.4	57.0-76.9	15.6	68.6	53.7-86.2	20.8	64.4	52.5-78.1	19.2	34.4	27.7-42.0	7.8	32.5	22.6-44.8	11.4	34.5	26.1-44.7
	1981-1985	34.8	57.9	49.5-67.2	14.2	57.2	44.2-72.6	20.6	57.4	46.8-69.7	17.8	30.3	24.3-37.4	7.4	30.9	21.4-42.8	10.4	29.0	21.6-38.1
	1986-1990	43.8	69.3	60.4-79.2	22.8	89.7	73.5-108.2	21.0	56.5	46.1-68.5	18.8	29.6	23.9-36.2	10.4	41.5	30.6-54.6	8.4	21.4	15.4-29.0
	1991-1995	38.2	56.0	48.3-64.5	20.6	75.2	61.0-91.4	17.6	42.6	34.1-52.6	18.4	27.1	21.8-33.2	9.0	33.5	24.2-44.9	9.4	22.1	16.2-29.5
	1996-2000	47.8	63.9	56.1-72.6	21.2	68.0	55.4-82.4	26.6	60.9	50.9-72.3	20.4	27.2	22.2-33.0	9.6	33.3	24.4-44.3	10.8	23.7	17.8-31.1
	2001-2005	41.2	50.2	43.5-57.6	19.4	55.2	44.5-67.5	21.8	46.2	37.7-56.0	17.2	20.6	16.4-25.4	7.0	20.4	14.2-28.4	10.2	19.6	14.4-26.0
	2006-2010	39.6	43.7	37.7-50.4	17.2	43.4	34.5-53.8	22.4	43.7	35.7-52.9	14.0	14.6	11.3-18.6	5.4	13.9	9.1-20.3	8.6	15.0	10.7-20.6
	2011-2015	31.4	32.1	27.1-37.8	14.2	32.1	24.8-40.8	17.2	32.0	25.4-40.0	12.4	12.6	9.6-16.3	7.2	16.5	11.4-23.0	5.2	10.0	6.4-15.0
2016-2020	26.6	25.1	20.8-29.9	13.8	27.6	21.3-35.3	12.8	22.4	17.1-29.0	9.0	8.4	6.1-11.5	4.6	9.7	6.0-14.8	4.4	7.6	4.7-11.9	
Colon excluding rectum	1976-1980	26.4	48.5	40.5-57.5	10.6	48.1	35.5-63.4	15.8	48.6	38.4-60.6	16.2	29.2	23.1-36.3	6.2	26.5	17.6-38.0	10.0	30.2	22.4-39.8
	1981-1985	23.6	40.0	33.0-47.9	9.0	37.0	26.7-49.7	14.6	41.2	32.2-51.9	14.8	25.3	19.8-31.8	6.2	26.9	18.0-38.3	8.6	23.9	17.3-32.3
	1986-1990	31.2	49.7	42.1-58.1	16.4	64.2	50.7-79.9	14.8	39.4	30.9-49.6	15.2	24.0	18.9-30.0	7.8	31.3	22.0-43.0	7.4	18.7	13.2-25.8
	1991-1995	28.2	41.1	34.5-48.4	14.4	53.9	41.9-68.1	13.8	32.7	25.4-41.4	16.8	24.8	19.8-30.7	7.6	28.6	20.0-39.3	9.2	21.7	15.9-29.1
	1996-2000	35.4	47.4	40.7-55.0	15.2	49.2	38.5-61.7	20.2	46.0	37.4-56.0	16.4	21.9	17.4-27.2	7.6	26.3	18.4-36.3	8.8	19.4	14.0-26.1
	2001-2005	30.6	37.1	31.4-43.6	13.2	38.5	29.6-49.2	17.4	36.2	28.8-44.9	14.8	17.6	13.8-22.2	6.4	18.6	12.7-26.3	8.4	16.0	11.4-21.9
	2006-2010	28.6	31.5	26.4-37.2	11.2	27.9	20.9-36.5	17.4	33.7	26.8-41.9	11.6	12.1	9.2-15.8	4.2	10.9	6.7-16.8	7.4	12.9	8.9-18.1
	2011-2015	23.8	24.5	20.1-29.5	10.2	23.2	17.1-30.8	13.6	25.1	19.3-32.2	10.8	11.1	8.2-14.6	6.4	14.7	10.0-21.0	4.4	8.7	5.3-13.4
2016-2020	19.4	18.7	15.1-23.0	8.8	17.9	12.9-24.4	10.6	18.9	14.0-25.1	7.8	7.3	5.1-10.1	3.8	7.8	4.6-12.5	4.0	7.0	4.2-11.1	
Rectum & rectosigmoid	1976-1980	10.0	17.9	13.3-23.7	5.0	20.5	13.1-30.5	5.0	15.8	10.2-23.3	3.0	5.2	2.9-8.6	1.6	6.0	2.5-11.9	1.4	4.3	1.7-8.8
	1981-1985	11.2	17.9	13.5-23.3	5.2	20.2	12.9-29.9	6.0	16.2	10.9-23.2	3.0	5.0	2.8-8.3	1.2	4.0	1.4-9.0	1.8	5.1	2.3-9.7
	1986-1990	12.6	19.6	15.1-25.1	6.4	25.5	17.1-36.3	6.2	17.1	11.6-24.3	3.6	5.6	3.3-8.9	2.6	10.1	5.3-17.4	1.0	2.7	0.9-6.2
	1991-1995	10.0	14.9	11.1-19.6	6.2	21.2	14.3-30.1	3.8	9.9	6.0-15.5	1.6	2.3	1.0-4.5	1.4	4.9	1.9-10.0	0.2	0.4	0.0-2.4
	1996-2000	12.4	16.5	12.6-21.1	6.0	18.8	12.6-26.8	6.4	14.9	10.2-21.1	4.0	5.3	3.2-8.1	2.0	7.0	3.3-12.8	2.0	4.4	2.1-8.1
	2001-2005	10.6	13.0	9.7-17.1	6.2	16.6	11.2-23.7	4.4	10.0	6.2-15.3	2.4	2.9	1.5-5.1	0.6	1.8	0.4-5.2	1.8	3.6	1.6-7.0
	2006-2010	11.0	12.2	9.1-16.0	6.0	15.4	10.3-22.2	5.0	10.0	6.4-15.0	2.4	2.5	1.3-4.4	1.2	3.0	1.0-6.6	1.2	2.1	0.8-4.9
	2011-2015	7.6	7.7	5.3-10.7	4.0	8.9	5.3-14.0	3.6	6.9	4.0-11.2	1.6	1.5	0.6-3.1	0.8	1.7	0.4-4.7	0.8	1.4	0.4-3.8
2016-2020	7.2	6.4	4.4-9.0	5.0	9.6	6.1-14.6	2.2	3.5	1.7-6.6	1.2	1.1	0.4-2.6	0.8	1.9	0.5-4.9	0.4	0.7	0.1-2.8	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Tompkins**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	1.4	2.2	0.9-4.7	0.8	3.5	0.9-8.7	0.6	1.4	0.3-4.5	1.4	2.1	0.8-4.4	0.6	2.9	0.5-8.1	0.8	1.9	0.5-5.3
	1981-1985	1.0	1.5	0.5-3.5	1.0	3.5	1.1-8.4	0.0	0.0	0.0-2.1	0.8	1.1	0.3-3.0	0.4	1.2	0.1-4.5	0.4	1.1	0.1-3.9
	1986-1990	0.6	0.8	0.1-2.4	0.4	1.7	0.2-5.6	0.2	0.2	0.0-2.3	1.2	2.0	0.7-4.2	0.8	3.5	0.9-8.5	0.4	1.0	0.1-3.7
	1991-1995	1.6	2.3	1.0-4.4	0.6	2.1	0.4-5.9	1.0	2.2	0.7-5.3	1.0	1.4	0.5-3.3	0.4	1.5	0.2-5.0	0.6	1.3	0.3-4.0
	1996-2000	2.8	3.8	2.0-6.3	2.0	6.4	3.0-11.8	0.8	2.0	0.5-5.0	1.0	1.3	0.4-3.1	0.8	2.7	0.7-6.9	0.2	0.5	0.0-2.7
	2001-2005	3.8	4.5	2.7-7.1	2.6	7.2	3.8-12.3	1.2	2.4	0.9-5.4	1.6	1.9	0.8-3.9	1.0	2.8	0.9-6.6	0.6	1.2	0.2-3.7
	2006-2010	4.6	5.5	3.4-8.3	2.6	7.2	3.8-12.3	2.0	4.0	1.8-7.6	3.0	3.6	2.0-5.9	1.8	5.2	2.3-9.8	1.2	2.5	0.9-5.6
	2011-2015	5.4	5.1	3.3-7.5	4.6	9.5	5.9-14.5	0.8	1.4	0.4-3.9	5.6	5.2	3.4-7.7	4.0	8.3	5.0-13.0	1.6	2.8	1.2-5.8
2016-2020	7.2	6.2	4.3-8.7	4.8	8.9	5.6-13.5	2.4	3.9	2.0-7.2	6.2	5.3	3.6-7.7	3.8	7.3	4.3-11.7	2.4	3.8	1.9-6.9	
Pancreas	1976-1980	5.2	9.5	6.2-13.9	2.6	11.8	6.2-20.1	2.6	7.6	4.0-13.0	5.2	9.9	6.4-14.4	3.0	14.9	8.2-24.4	2.2	6.5	3.2-11.7
	1981-1985	5.4	9.3	6.1-13.5	1.8	7.6	3.4-14.4	3.6	10.6	6.2-16.8	4.8	8.4	5.4-12.5	1.6	6.6	2.8-12.8	3.2	9.2	5.3-15.0
	1986-1990	9.2	14.6	10.7-19.4	4.8	18.3	11.5-27.3	4.4	11.8	7.4-17.9	8.2	12.9	9.2-17.4	3.8	14.2	8.4-22.2	4.4	11.6	7.2-17.6
	1991-1995	6.8	10.0	6.9-13.9	2.4	8.1	4.1-14.0	4.4	10.9	6.8-16.6	7.0	10.3	7.1-14.2	2.8	9.3	5.0-15.6	4.2	10.4	6.4-15.9
	1996-2000	8.0	10.5	7.5-14.3	3.4	10.7	6.1-17.2	4.6	10.7	6.7-16.1	7.6	10.2	7.2-13.9	3.2	10.7	6.0-17.4	4.4	10.3	6.4-15.7
	2001-2005	10.4	12.6	9.4-16.6	5.2	14.3	9.3-21.1	5.2	11.0	7.1-16.3	9.8	11.9	8.8-15.8	5.6	15.9	10.5-23.1	4.2	8.8	5.4-13.6
	2006-2010	11.6	12.3	9.3-16.0	4.8	12.3	7.8-18.4	6.8	12.0	8.2-17.1	10.4	11.1	8.2-14.7	4.4	11.6	7.2-17.7	6.0	10.2	6.8-14.9
	2011-2015	14.6	14.7	11.4-18.6	7.0	17.0	11.7-23.9	7.6	12.8	8.9-17.9	11.6	11.7	8.8-15.2	6.4	16.0	10.9-22.8	5.2	8.1	5.2-12.2
2016-2020	15.6	14.0	11.0-17.6	7.8	14.8	10.4-20.5	7.8	13.1	9.1-18.2	12.6	11.2	8.5-14.4	5.6	10.9	7.1-16.1	7.0	11.1	7.7-15.7	
Larynx	1976-1980	3.2	5.2	3.0-8.5	2.4	8.7	4.4-15.3	0.8	2.5	0.7-6.4	0.4	0.7	0.1-2.5	0.4	2.0	0.2-6.7	0.0	0.0	0.0-2.2
	1981-1985	3.8	6.4	3.8-10.0	2.2	8.1	4.0-14.4	1.6	5.2	2.2-10.1	0.2	0.3	0.0-1.7	0.0	0.0	0.0-2.6	0.2	0.5	0.0-3.0
	1986-1990	4.2	6.8	4.2-10.3	3.0	11.8	6.5-19.5	1.2	3.3	1.2-7.1	1.4	2.0	0.8-4.2	0.8	2.5	0.7-6.5	0.6	1.6	0.3-4.6
	1991-1995	3.4	4.9	2.8-7.8	2.4	8.2	4.2-14.2	1.0	2.5	0.8-5.8	1.4	2.1	0.8-4.2	1.2	4.3	1.6-9.2	0.2	0.5	0.0-2.8
	1996-2000	3.8	5.2	3.1-8.1	2.8	8.4	4.6-14.0	1.0	2.6	0.8-6.0	1.0	1.4	0.5-3.2	0.6	1.9	0.4-5.5	0.4	1.1	0.1-3.7
	2001-2005	3.8	4.4	2.6-6.9	2.8	7.6	4.1-12.8	1.0	2.0	0.6-4.9	1.8	2.1	1.0-4.1	1.0	2.8	0.9-6.5	0.8	1.7	0.4-4.5
	2006-2010	2.4	2.8	1.4-4.9	2.2	5.6	2.7-10.1	0.2	0.4	0.0-2.4	0.0	0.0	0.0-0.9	0.0	0.0	0.0-1.9	0.0	0.0	0.0-1.8
	2011-2015	2.6	2.6	1.4-4.6	2.2	4.5	2.2-8.4	0.4	1.0	0.1-3.7	1.6	1.8	0.7-3.5	1.4	3.7	1.4-7.6	0.2	0.3	0.0-2.1
2016-2020	3.2	2.8	1.6-4.8	3.0	5.6	3.0-9.6	0.2	0.5	0.0-2.6	1.2	1.1	0.4-2.6	1.2	2.5	0.9-5.7	0.0	0.0	0.0-1.8	
Lung and bronchus	1976-1980	29.6	51.6	43.5-60.6	19.2	80.2	64.4-98.4	10.4	32.5	24.1-42.7	22.4	40.0	32.8-48.1	15.4	68.4	53.4-85.8	7.0	21.5	14.9-29.9
	1981-1985	38.8	64.8	55.9-74.6	27.6	110.6	92.4-131.1	11.2	34.0	25.5-44.2	27.6	46.4	38.9-54.9	19.6	79.5	64.2-97.2	8.0	24.8	17.6-33.8
	1986-1990	41.6	64.4	55.9-73.8	24.4	93.6	77.3-112.1	17.2	46.5	37.1-57.5	34.2	53.3	45.6-62.0	21.0	80.1	65.1-97.3	13.2	35.9	27.7-45.8
	1991-1995	44.6	64.6	56.4-73.7	25.8	89.6	74.5-106.7	18.8	47.8	38.6-58.6	36.6	53.2	45.7-61.4	22.0	79.1	64.6-95.5	14.6	36.7	28.7-46.2
	1996-2000	47.8	65.0	57.0-73.8	23.2	73.0	60.2-87.6	24.6	59.0	49.0-70.5	37.6	51.2	44.2-59.1	18.8	60.0	48.3-73.4	18.8	44.6	36.0-54.7
	2001-2005	48.0	60.0	52.6-68.2	21.6	62.7	51.3-75.8	26.4	59.3	49.5-70.5	38.0	47.6	41.0-54.9	16.2	48.4	38.3-60.2	21.8	48.2	39.4-58.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Tompkins**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	51.2	58.8	51.6-66.6	23.6	60.4	49.7-72.6	27.6	58.1	48.6-68.9	37.8	43.4	37.3-50.2	18.0	46.6	37.3-57.5	19.8	41.0	33.1-50.2
	2011-2015	55.8	55.6	49.1-62.7	26.2	57.6	47.8-68.8	29.6	54.4	45.8-64.2	38.2	37.9	32.6-43.9	18.2	40.1	32.0-49.6	20.0	36.0	29.1-44.1
	2016-2020	59.2	52.1	46.2-58.6	29.8	59.0	49.7-69.7	29.4	47.5	39.9-56.1	35.0	31.5	26.9-36.7	20.6	41.9	34.0-51.1	14.4	23.7	18.4-30.1
Melanoma of the skin	1976-1980	4.4	6.7	4.0-10.3	2.0	7.1	3.1-13.7	2.4	6.9	3.4-12.4	0.8	0.9	0.2-2.6	0.4	0.9	0.1-4.4	0.4	0.9	0.1-3.6
	1981-1985	8.2	12.4	8.8-17.0	4.6	16.5	10.3-25.1	3.6	9.6	5.5-15.6	2.6	4.2	2.2-7.2	2.2	8.1	3.9-14.6	0.4	1.3	0.2-4.6
	1986-1990	8.4	11.0	7.8-15.1	4.0	14.9	8.8-23.3	4.4	9.6	5.9-14.9	2.2	3.2	1.6-5.8	1.0	3.3	1.1-7.7	1.2	2.9	1.1-6.5
	1991-1995	9.8	12.4	9.0-16.5	4.2	12.7	7.5-19.9	5.6	12.9	8.4-18.9	0.8	1.2	0.3-2.9	0.6	1.9	0.4-5.5	0.2	0.4	0.0-2.4
	1996-2000	12.2	15.8	12.0-20.3	6.6	18.7	12.8-26.4	5.6	13.4	8.9-19.5	1.0	1.3	0.4-3.0	0.4	1.7	0.2-5.7	0.6	1.5	0.3-4.2
	2001-2005	15.6	18.3	14.4-22.9	7.4	19.5	13.6-27.0	8.2	17.5	12.5-24.0	3.4	4.2	2.4-6.7	2.0	5.4	2.6-10.0	1.4	2.7	1.1-5.9
	2006-2010	25.2	27.4	22.7-32.8	12.8	31.8	24.3-40.8	12.4	24.8	18.8-32.0	1.8	1.9	0.8-3.6	1.2	3.3	1.2-7.1	0.6	1.0	0.2-3.2
	2011-2015	22.8	23.1	18.9-28.0	14.0	32.0	24.7-40.7	8.8	17.0	12.1-23.2	2.6	2.5	1.3-4.4	1.4	3.4	1.4-7.2	1.2	2.2	0.8-4.9
	2016-2020	26.4	24.7	20.5-29.5	14.6	30.2	23.5-38.3	11.8	20.1	15.1-26.3	2.0	1.8	0.8-3.6	1.4	3.1	1.2-6.5	0.6	1.1	0.2-3.5
Female breast	1976-1980							28.0	89.2	74.7-105.5							11.0	34.4	25.8-44.9
	1981-1985							40.0	120.0	103.6-138.1							9.2	26.5	19.3-35.5
	1986-1990							46.0	129.3	112.9-147.4							11.4	29.9	22.6-38.9
	1991-1995							51.2	131.5	115.7-148.7							12.8	32.8	25.2-41.9
	1996-2000							54.6	131.3	116.1-147.9							10.4	24.2	18.1-31.9
	2001-2005							57.4	128.2	113.7-144.1							9.2	19.0	13.8-25.5
	2006-2010							63.4	131.0	116.6-146.7							12.2	22.2	16.9-28.9
	2011-2015							64.2	128.0	113.9-143.5							10.8	20.0	14.8-26.4
	2016-2020							79.6	145.9	131.3-161.8							10.6	18.0	13.3-23.9
Cervix uteri	1976-1980							4.2	12.6	7.7-19.4							1.2	3.8	1.4-8.2
	1981-1985							2.8	8.2	4.4-13.8							0.8	2.1	0.6-5.5
	1986-1990							2.6	6.9	3.6-12.0							1.2	3.7	1.3-7.9
	1991-1995							2.2	5.4	2.7-9.7							1.0	2.7	0.9-6.2
	1996-2000							1.8	4.5	2.1-8.6							0.4	0.9	0.1-3.4
	2001-2005							2.8	6.5	3.5-10.9							1.0	2.3	0.7-5.4
	2006-2010							3.0	6.3	3.5-10.6							1.2	2.0	0.7-4.6
	2011-2015							4.0	8.5	5.1-13.4							0.8	1.7	0.4-4.5
	2016-2020							1.2	2.8	1.0-6.2							0.4	0.6	0.1-2.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Tompkins**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							9.6	29.2	21.4-38.7							1.6	4.8	2.1-9.4
	1981-1985							8.0	23.5	16.7-32.1							1.8	4.9	2.2-9.4
	1986-1990							8.0	22.9	16.2-31.2							2.0	5.0	2.4-9.3
	1991-1995							10.6	28.1	21.0-36.7							1.0	2.6	0.8-5.9
	1996-2000							11.0	26.2	19.7-34.2							2.0	4.6	2.2-8.6
	2001-2005							8.6	19.3	13.9-26.1							1.2	2.9	1.0-6.2
	2006-2010							19.2	38.3	30.9-47.1							2.6	4.9	2.6-8.6
	2011-2015							19.8	35.2	28.4-43.2							2.6	4.6	2.4-8.2
	2016-2020							13.8	22.1	17.0-28.5							2.8	4.2	2.3-7.4
Ovary	1976-1980							4.0	12.1	7.3-18.8							2.6	7.9	4.2-13.6
	1981-1985							7.6	23.7	16.6-32.6							5.4	15.5	10.1-22.5
	1986-1990							5.4	14.2	9.3-20.8							4.0	10.3	6.3-16.0
	1991-1995							7.8	20.1	14.2-27.5							4.0	9.8	6.0-15.2
	1996-2000							6.0	13.3	8.9-19.1							5.0	11.8	7.6-17.4
	2001-2005							4.4	10.3	6.4-15.6							4.2	9.2	5.7-14.2
	2006-2010							6.8	14.4	9.9-20.3							3.6	6.8	4.0-10.9
	2011-2015							7.4	13.5	9.3-19.0							6.0	10.3	6.8-14.9
	2016-2020							6.4	11.3	7.6-16.3							3.6	5.8	3.3-9.5
Prostate	1976-1980				15.2	71.2	55.5-89.6							5.8	30.3	20.0-43.5			
	1981-1985				19.8	89.4	72.1-109.2							6.8	32.4	22.1-45.4			
	1986-1990				28.6	117.6	98.6-138.9							7.0	31.5	21.6-44.0			
	1991-1995				51.6	188.0	165.3-212.8							10.8	46.1	34.3-60.2			
	1996-2000				60.8	191.5	170.3-214.5							9.8	38.2	28.0-50.5			
	2001-2005				80.8	225.5	203.8-249.0							6.0	20.5	13.8-29.0			
	2006-2010				84.2	198.7	179.7-219.3							9.2	26.6	19.5-35.5			
	2011-2015				47.6	98.1	85.5-112.0							6.8	16.2	11.1-22.7			
	2016-2020				68.2	121.6	108.6-135.8							8.2	17.8	12.7-24.4			
Testis	1976-1980				2.4	4.0	1.8-8.3							0.2	0.4	0.0-3.4			
	1981-1985				2.6	6.4	3.1-11.7							0.2	0.9	0.0-4.5			
	1986-1990				2.8	5.4	2.9-9.6							0.4	1.1	0.1-4.4			
	1991-1995				2.6	4.9	2.5-9.0							0.0	0.0	0.0-2.2			
	1996-2000				3.6	6.9	3.9-11.4							0.2	0.2	0.0-2.2			
	2001-2005				2.4	4.2	1.9-8.1							0.0	0.0	0.0-2.0			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Tompkins**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				3.8	6.5	3.7-10.8							0.2	0.4	0.0-2.6			
	2011-2015				3.8	9.0	5.3-14.2							0.2	0.1	0.0-2.0			
	2016-2020				3.8	8.9	5.2-14.2							0.2	0.3	0.0-2.4			
Urinary bladder (incl. in situ)	1976-1980	11.6	20.5	15.5-26.6	7.2	30.3	20.8-42.3	4.4	13.5	8.4-20.5	2.4	4.4	2.3-7.6	1.2	5.9	2.0-12.6	1.2	3.6	1.3-7.7
	1981-1985	14.0	23.3	18.1-29.5	10.2	41.1	30.3-54.3	3.8	10.8	6.5-16.8	2.6	4.1	2.2-7.1	1.8	7.5	3.3-14.3	0.8	2.1	0.6-5.4
	1986-1990	12.2	18.6	14.2-23.9	8.6	33.9	24.2-45.9	3.6	8.8	5.2-14.0	2.2	3.6	1.8-6.4	1.4	5.2	2.1-10.7	0.8	2.1	0.6-5.3
	1991-1995	18.2	26.5	21.3-32.5	10.4	36.8	27.2-48.5	7.8	19.3	13.7-26.4	3.8	5.5	3.3-8.6	2.2	9.6	4.7-17.1	1.6	3.6	1.5-7.1
	1996-2000	21.2	28.5	23.4-34.5	16.2	52.5	41.5-65.4	5.0	11.3	7.2-16.7	3.8	5.1	3.1-8.0	2.4	7.8	4.0-13.5	1.4	3.1	1.2-6.5
	2001-2005	23.8	29.5	24.4-35.4	17.8	52.2	41.8-64.3	6.0	13.3	8.9-19.1	4.2	5.3	3.3-8.2	2.8	8.6	4.7-14.4	1.4	2.7	1.1-5.9
	2006-2010	25.0	27.4	22.7-32.7	18.2	46.2	37.0-57.0	6.8	13.2	9.0-18.7	5.0	5.4	3.5-8.1	3.6	9.5	5.6-15.1	1.4	2.7	1.1-5.8
	2011-2015	25.0	26.0	21.5-31.1	18.4	44.6	35.7-55.1	6.6	11.8	8.0-16.8	4.2	4.2	2.6-6.6	2.8	6.8	3.7-11.6	1.4	2.4	0.9-5.3
2016-2020	24.2	22.1	18.3-26.6	17.4	36.5	29.0-45.3	6.8	11.6	7.9-16.4	2.8	2.7	1.4-4.6	1.8	4.0	1.8-7.8	1.0	1.7	0.5-4.3	
Kidney and renal pelvis	1976-1980	3.4	6.0	3.4-9.6	1.8	7.6	3.4-14.4	1.6	4.7	2.0-9.2	1.8	3.2	1.5-6.1	1.2	5.5	1.9-11.8	0.6	1.7	0.4-5.1
	1981-1985	4.6	7.7	4.8-11.5	2.8	10.9	5.9-18.3	1.8	5.0	2.3-9.6	2.0	3.3	1.6-6.0	1.4	5.0	2.0-10.3	0.6	1.6	0.3-4.8
	1986-1990	6.6	10.6	7.3-14.8	4.2	16.6	10.1-25.4	2.4	6.6	3.4-11.4	4.6	7.3	4.6-10.9	2.8	11.3	6.0-18.9	1.8	5.0	2.3-9.4
	1991-1995	6.4	9.1	6.2-12.8	4.2	14.6	8.9-22.4	2.2	5.4	2.7-9.8	2.4	3.6	1.8-6.2	1.6	5.9	2.5-11.5	0.8	2.0	0.5-5.2
	1996-2000	7.6	10.3	7.3-14.1	5.0	14.9	9.6-22.0	2.6	6.6	3.5-11.3	2.0	2.7	1.3-5.0	1.4	4.5	1.8-9.1	0.6	1.4	0.3-4.1
	2001-2005	9.4	11.7	8.6-15.6	5.0	14.7	9.5-21.7	4.4	10.1	6.3-15.4	2.6	3.2	1.7-5.5	1.4	4.7	1.9-9.4	1.2	2.4	0.9-5.5
	2006-2010	10.2	11.2	8.3-14.9	6.6	15.8	10.8-22.4	3.6	7.1	4.1-11.5	4.4	5.1	3.2-7.8	2.8	7.3	4.0-12.4	1.6	3.1	1.3-6.4
	2011-2015	12.4	12.3	9.3-16.0	8.2	18.5	13.1-25.5	4.2	7.4	4.5-11.6	3.4	3.7	2.1-6.0	2.0	4.8	2.2-9.0	1.4	2.7	1.1-5.8
2016-2020	14.2	13.5	10.4-17.1	9.8	20.0	14.6-26.8	4.4	7.9	4.9-12.2	3.6	3.5	2.0-5.7	2.4	5.6	2.8-9.9	1.2	2.1	0.7-4.9	
Brain and other central nervous system	1976-1980	4.2	6.9	4.2-10.6	2.4	9.3	4.7-16.5	1.8	5.1	2.3-9.9	2.6	4.4	2.3-7.5	1.6	6.0	2.5-12.0	1.0	3.0	1.0-6.9
	1981-1985	4.4	6.6	4.0-10.1	2.8	9.1	4.7-15.7	1.6	4.9	2.0-9.7	2.4	3.8	1.9-6.7	1.4	4.5	1.8-9.4	1.0	3.2	1.0-7.4
	1986-1990	5.0	6.6	4.2-9.9	3.6	10.4	6.0-16.7	1.4	3.4	1.3-7.1	2.6	3.5	1.8-6.1	1.6	5.5	2.3-11.0	1.0	2.0	0.6-5.0
	1991-1995	7.0	9.1	6.3-12.7	3.4	9.4	5.3-15.4	3.6	9.1	5.3-14.4	4.6	6.2	3.9-9.4	2.6	7.9	4.2-13.6	2.0	5.0	2.4-9.2
	1996-2000	5.0	6.3	4.1-9.4	3.4	9.2	5.3-14.8	1.6	3.6	1.5-7.1	3.2	4.2	2.4-6.8	1.6	4.4	1.9-8.6	1.6	3.7	1.6-7.4
	2001-2005	5.8	7.1	4.7-10.2	2.2	6.3	3.1-11.3	3.6	8.3	4.8-13.3	3.6	4.4	2.6-7.0	2.0	5.6	2.6-10.3	1.6	3.5	1.5-7.0
	2006-2010	8.2	9.9	7.0-13.5	6.0	14.7	9.8-21.3	2.2	5.1	2.5-9.3	4.2	5.2	3.2-8.0	2.2	5.9	2.9-10.5	2.0	4.6	2.2-8.5
	2011-2015	8.0	7.7	5.4-10.7	4.2	9.2	5.5-14.4	3.8	6.7	4.0-10.7	5.2	5.1	3.3-7.6	3.0	6.7	3.7-11.3	2.2	3.8	1.8-7.1
2016-2020	5.0	4.8	3.0-7.2	3.4	6.7	3.8-11.0	1.6	3.1	1.2-6.4	6.8	6.8	4.6-9.7	4.8	10.1	6.3-15.4	2.0	3.8	1.8-7.3	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Tompkins**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	2.4	2.5	1.1-4.8	0.4	1.1	0.1-4.6	2.0	3.7	1.5-7.7	0.2	0.4	0.0-2.1	0.0	0.0	0.0-2.8	0.2	0.6	0.0-3.4
	1981-1985	2.4	3.5	1.7-6.3	0.8	2.4	0.6-6.5	1.6	4.5	1.8-9.2	0.0	0.0	0.0-1.1	0.0	0.0	0.0-2.6	0.0	0.0	0.0-2.1
	1986-1990	4.2	5.3	3.2-8.2	1.2	3.2	1.1-7.2	3.0	7.1	3.8-12.1	0.6	0.9	0.2-2.7	0.2	0.6	0.0-3.6	0.4	1.0	0.1-3.7
	1991-1995	4.8	6.2	3.9-9.3	1.6	5.0	2.1-9.9	3.2	7.7	4.3-12.7	0.2	0.3	0.0-1.6	0.0	0.0	0.0-2.2	0.2	0.5	0.0-2.8
	1996-2000	4.2	4.9	3.0-7.6	1.2	3.3	1.2-7.3	3.0	6.6	3.7-11.0	0.2	0.3	0.0-1.4	0.2	0.7	0.0-3.5	0.0	0.0	0.0-1.8
	2001-2005	4.4	5.0	3.1-7.7	0.8	2.1	0.6-5.3	3.6	7.6	4.4-12.3	0.4	0.4	0.1-1.6	0.2	0.8	0.0-3.6	0.2	0.3	0.0-2.2
	2006-2010	13.6	14.5	11.2-18.6	3.8	9.8	5.8-15.4	9.8	19.4	14.1-26.0	0.2	0.2	0.0-1.3	0.2	0.5	0.0-2.7	0.0	0.0	0.0-1.8
	2011-2015	12.8	13.2	10.1-17.1	4.0	8.1	4.8-12.8	8.8	18.1	12.9-24.6	0.6	0.6	0.1-1.8	0.0	0.0	0.0-1.9	0.6	1.0	0.2-3.2
2016-2020	11.0	11.1	8.3-14.7	2.2	4.3	2.0-8.2	8.8	17.5	12.5-23.9	0.8	0.7	0.2-2.0	0.2	0.5	0.0-2.9	0.6	0.8	0.2-2.9	
Hodgkin lymphoma	1976-1980	3.6	3.7	2.0-6.3	2.4	5.1	2.2-10.1	1.2	2.8	0.9-6.5	0.4	0.7	0.1-2.5	0.4	2.0	0.2-6.7	0.0	0.0	0.0-2.2
	1981-1985	3.4	4.7	2.7-7.8	1.8	5.8	2.4-11.5	1.6	4.4	1.8-8.9	1.0	1.3	0.4-3.1	0.8	2.3	0.5-6.7	0.2	0.6	0.0-3.1
	1986-1990	2.8	2.4	1.2-4.5	1.6	2.8	1.0-6.6	1.2	1.8	0.6-4.6	0.8	0.8	0.2-2.4	0.4	0.7	0.1-3.6	0.4	0.8	0.1-3.3
	1991-1995	2.4	2.2	1.1-4.1	1.0	2.0	0.6-5.3	1.4	2.3	0.9-5.2	0.6	0.9	0.2-2.6	0.4	1.2	0.1-4.3	0.2	0.5	0.0-2.8
	1996-2000	3.4	2.8	1.5-4.8	1.2	2.4	0.8-5.7	2.2	3.4	1.5-6.6	0.2	0.3	0.0-1.5	0.0	0.0	0.0-2.1	0.2	0.6	0.0-2.9
	2001-2005	3.0	3.4	1.9-5.8	2.4	6.1	3.1-10.9	0.6	1.1	0.2-3.6	0.2	0.3	0.0-1.5	0.0	0.0	0.0-2.0	0.2	0.6	0.0-2.9
	2006-2010	3.6	3.7	2.1-6.0	2.8	5.6	2.9-9.7	0.8	1.8	0.4-4.7	0.6	0.7	0.1-2.0	0.2	0.7	0.0-3.3	0.4	0.7	0.1-2.8
	2011-2015	4.0	4.1	2.4-6.5	2.2	5.1	2.5-9.4	1.8	3.2	1.4-6.5	0.6	0.7	0.1-2.0	0.0	0.0	0.0-1.9	0.6	1.2	0.2-3.8
2016-2020	2.6	2.8	1.5-5.0	1.6	3.5	1.4-7.2	1.0	2.3	0.7-5.5	0.0	0.0	0.0-0.9	0.0	0.0	0.0-1.9	0.0	0.0	0.0-1.8	
Non-Hodgkin lymphomas	1976-1980	5.8	9.5	6.3-13.8	3.4	12.7	7.1-20.8	2.4	7.2	3.6-12.7	3.2	5.0	2.8-8.2	2.2	8.8	4.2-16.0	1.0	2.5	0.8-6.1
	1981-1985	11.4	18.9	14.3-24.6	6.4	26.5	17.7-37.7	5.0	14.4	9.2-21.2	6.2	10.2	6.9-14.5	4.2	16.7	10.2-25.7	2.0	5.7	2.7-10.6
	1986-1990	10.0	15.2	11.2-20.2	5.6	20.7	13.3-30.4	4.4	12.1	7.5-18.5	4.6	7.6	4.8-11.4	1.6	7.4	3.0-14.5	3.0	8.2	4.5-13.6
	1991-1995	12.0	16.2	12.3-20.9	7.2	22.4	15.4-31.4	4.8	11.7	7.4-17.6	4.6	6.4	4.0-9.6	3.2	11.4	6.3-18.6	1.4	3.4	1.4-7.1
	1996-2000	13.0	17.4	13.4-22.1	6.6	21.4	14.6-30.2	6.4	15.3	10.4-21.6	4.6	6.1	3.9-9.1	2.0	6.9	3.2-12.6	2.6	5.4	2.8-9.4
	2001-2005	18.4	22.2	17.8-27.3	9.4	26.0	18.9-34.8	9.0	19.4	14.0-26.1	5.6	6.6	4.4-9.6	2.4	7.7	3.9-13.3	3.2	6.4	3.6-10.5
	2006-2010	19.2	21.8	17.5-26.7	10.0	27.0	19.9-35.6	9.2	18.5	13.4-25.0	4.6	5.1	3.2-7.7	2.6	6.8	3.5-11.6	2.0	3.9	1.8-7.3
	2011-2015	21.0	21.4	17.4-26.2	10.6	23.6	17.4-31.3	10.4	19.7	14.5-26.2	4.8	5.3	3.4-8.0	2.8	7.5	4.1-12.6	2.0	3.6	1.7-6.9
2016-2020	21.8	20.5	16.8-24.9	12.2	26.0	19.7-33.7	9.6	16.0	11.7-21.5	5.0	4.7	3.0-7.0	2.0	4.1	1.9-7.7	3.0	5.0	2.7-8.5	
Myeloma	1976-1980	3.0	5.3	2.9-8.6	1.6	6.6	2.8-13.0	1.4	4.3	1.7-8.9	0.6	1.0	0.2-2.8	0.4	1.4	0.2-5.2	0.2	0.5	0.0-3.1
	1981-1985	3.0	5.2	2.9-8.6	1.4	5.1	2.0-10.4	1.6	4.6	2.0-9.1	2.0	3.4	1.6-6.2	1.0	4.3	1.3-9.8	1.0	2.6	0.9-6.2
	1986-1990	2.6	3.8	2.0-6.5	1.6	6.0	2.4-11.9	1.0	2.6	0.8-6.0	2.0	3.2	1.5-5.8	1.0	4.3	1.4-9.7	1.0	2.6	0.8-6.0
	1991-1995	3.6	5.1	3.0-8.1	2.0	6.9	3.2-12.8	1.6	3.9	1.7-7.7	2.0	2.9	1.4-5.3	0.8	3.7	0.9-9.0	1.2	2.9	1.0-6.3
	1996-2000	3.4	4.6	2.7-7.3	2.0	6.7	3.2-12.1	1.4	3.3	1.3-6.8	2.8	3.6	1.9-6.0	1.6	4.9	2.0-9.7	1.2	2.6	1.0-5.9
	2001-2005	5.0	6.4	4.1-9.5	3.4	10.0	5.8-16.0	1.6	3.5	1.5-7.0	4.2	5.2	3.2-7.9	2.6	7.8	4.1-13.2	1.6	3.1	1.3-6.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Tompkins**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	5.4	5.8	3.8-8.6	3.6	9.1	5.3-14.4	1.8	3.6	1.6-7.0	2.2	2.7	1.3-4.8	1.2	3.6	1.3-7.8	1.0	2.2	0.7-5.2
	2011-2015	8.0	7.7	5.4-10.6	4.8	10.5	6.6-15.9	3.2	5.4	3.0-9.1	3.4	3.4	1.9-5.5	2.0	4.5	2.1-8.4	1.4	2.6	1.0-5.7
	2016-2020	12.6	11.1	8.4-14.3	7.2	13.2	9.1-18.6	5.4	9.2	6.0-13.8	2.8	2.4	1.3-4.1	1.8	3.3	1.5-6.7	1.0	1.5	0.5-4.0
Leukemias	1976-1980	9.2	15.3	11.1-20.6	5.8	25.2	16.5-36.5	3.4	9.4	5.3-15.4	4.0	7.2	4.4-11.2	2.6	12.3	6.3-21.1	1.4	4.2	1.7-8.7
	1981-1985	8.0	12.3	8.7-16.9	4.4	16.7	10.1-25.8	3.6	9.2	5.4-14.7	5.4	8.4	5.4-12.3	2.6	11.1	5.6-19.3	2.8	7.4	3.9-12.6
	1986-1990	8.4	13.4	9.6-18.1	5.6	20.4	13.4-29.6	2.8	8.4	4.5-14.0	6.2	10.0	6.8-14.1	4.4	17.1	10.6-25.8	1.8	5.1	2.3-9.6
	1991-1995	11.4	16.1	12.1-20.8	5.4	17.8	11.5-26.1	6.0	15.0	10.1-21.4	4.8	7.2	4.6-10.6	2.2	8.5	4.2-15.0	2.6	6.4	3.4-10.9
	1996-2000	10.8	14.6	11.0-19.0	3.8	12.4	7.4-19.4	7.0	16.8	11.6-23.4	6.0	7.8	5.2-11.1	2.6	7.8	4.0-13.4	3.4	7.8	4.5-12.5
	2001-2005	12.0	14.7	11.2-19.0	6.4	17.5	11.8-24.8	5.6	12.8	8.4-18.6	4.0	4.8	2.9-7.4	2.2	6.8	3.3-12.0	1.8	3.5	1.6-6.9
	2006-2010	15.8	17.8	14.0-22.4	9.6	24.2	17.6-32.3	6.2	12.2	8.1-17.6	5.4	5.5	3.5-8.1	3.0	7.3	4.0-12.4	2.4	4.1	2.1-7.5
	2011-2015	18.8	19.3	15.4-23.8	9.4	21.9	15.9-29.4	9.4	17.3	12.5-23.4	8.2	8.3	5.9-11.4	3.4	8.2	4.7-13.2	4.8	8.6	5.3-13.1
	2016-2020	18.8	17.3	13.8-21.3	10.4	21.9	16.2-29.0	8.4	13.6	9.7-18.7	6.4	5.9	4.0-8.5	3.0	7.1	3.9-11.7	3.4	5.1	2.9-8.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Ulster**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	566.4	377.3	363.3-391.7	287.0	449.8	425.9-474.7	279.4	335.9	318.1-354.4	328.2	219.8	209.1-230.9	172.8	278.9	259.8-298.9	155.4	182.3	169.5-195.9
	1981-1985	647.2	415.0	400.6-429.8	315.4	480.7	456.5-505.9	331.8	383.6	365.0-403.0	344.2	223.4	212.9-234.4	179.2	281.9	263.1-301.6	165.0	187.2	174.4-200.7
	1986-1990	688.6	428.0	413.7-442.8	340.8	496.6	472.6-521.5	347.8	392.2	373.6-411.4	356.6	221.1	210.9-231.7	184.8	275.3	257.3-294.2	171.8	186.1	173.7-199.2
	1991-1995	798.8	462.5	448.2-477.1	401.6	541.5	517.4-566.3	397.2	419.3	400.8-438.3	390.8	227.7	217.7-238.1	203.4	286.3	268.4-304.9	187.4	190.4	178.3-203.2
	1996-2000	940.2	510.1	495.6-524.9	474.2	583.2	559.6-607.6	466.0	459.2	440.6-478.4	389.2	211.5	202.2-221.1	198.8	259.0	242.8-276.0	190.4	180.2	168.9-192.2
	2001-2005	997.6	500.2	486.3-514.3	512.6	575.4	552.9-598.5	485.0	449.3	431.4-467.7	411.6	207.0	198.1-216.1	202.0	238.8	223.9-254.3	209.6	185.6	174.4-197.3
	2006-2010	1,104.4	507.0	493.5-520.9	563.4	566.5	545.3-588.4	541.0	466.9	449.1-485.3	388.0	178.3	170.4-186.6	193.6	205.0	192.0-218.7	194.4	161.2	151.1-172.0
	2011-2015	1,098.4	463.5	450.9-476.3	560.8	500.9	482.0-520.5	537.6	440.2	423.0-458.0	388.4	160.9	153.7-168.4	196.8	183.2	171.6-195.3	191.6	144.4	135.1-154.3
2016-2020	1,147.6	449.5	437.4-461.8	583.4	474.3	456.6-492.5	564.2	437.1	420.1-454.7	390.8	148.1	141.4-155.0	203.6	169.6	159.1-180.7	187.2	130.9	122.3-140.0	
Oral cavity and pharynx	1976-1980	14.8	9.9	7.8-12.5	10.4	16.1	11.9-21.2	4.4	5.2	3.2-7.9	7.4	4.9	3.4-6.7	5.2	8.0	5.1-11.9	2.2	2.5	1.2-4.6
	1981-1985	18.0	11.2	9.0-13.9	13.2	18.9	14.5-24.2	4.8	5.4	3.4-8.1	6.6	4.1	2.8-5.7	4.4	6.5	4.0-10.0	2.2	2.3	1.1-4.2
	1986-1990	17.4	10.6	8.4-13.1	13.0	17.8	13.7-22.8	4.4	4.7	3.0-7.3	7.2	4.4	3.1-6.1	4.6	6.1	3.9-9.2	2.6	2.8	1.5-4.9
	1991-1995	21.4	12.4	10.2-15.0	14.2	17.9	14.0-22.7	7.2	7.5	5.2-10.4	8.2	4.7	3.3-6.3	5.8	7.3	4.9-10.5	2.4	2.4	1.2-4.3
	1996-2000	19.8	10.7	8.7-13.0	13.6	16.6	12.8-21.1	6.2	6.0	4.1-8.6	5.2	2.8	1.8-4.1	4.0	5.0	3.0-7.7	1.2	1.0	0.4-2.3
	2001-2005	19.6	9.7	7.9-11.8	13.8	14.2	11.0-18.0	5.8	5.2	3.5-7.6	4.6	2.3	1.4-3.5	2.8	3.3	1.8-5.6	1.8	1.5	0.7-3.0
	2006-2010	26.6	11.9	9.9-14.1	19.8	18.1	14.6-22.2	6.8	6.0	4.1-8.4	5.6	2.5	1.7-3.7	3.0	2.7	1.5-4.6	2.6	2.2	1.1-3.8
	2011-2015	28.8	11.6	9.7-13.7	20.8	17.6	14.3-21.5	8.0	6.1	4.4-8.5	3.8	1.6	0.9-2.5	3.0	2.9	1.6-4.8	0.8	0.6	0.2-1.7
2016-2020	39.6	14.9	12.8-17.3	27.4	21.6	18.0-25.7	12.2	8.8	6.7-11.6	4.4	1.7	1.1-2.7	3.8	3.3	2.0-5.3	0.6	0.5	0.1-1.6	
Esophagus	1976-1980	7.2	4.8	3.3-6.6	4.8	7.1	4.5-10.7	2.4	2.9	1.5-5.2	6.0	4.1	2.7-5.9	4.4	7.0	4.3-10.7	1.6	1.9	0.8-3.9
	1981-1985	6.0	3.8	2.5-5.4	4.4	6.3	3.9-9.7	1.6	1.7	0.7-3.3	3.6	2.4	1.4-3.8	2.6	4.0	2.1-6.9	1.0	1.1	0.3-2.5
	1986-1990	5.8	3.3	2.2-4.8	4.8	6.3	4.0-9.5	1.0	1.0	0.3-2.4	5.4	3.2	2.1-4.6	4.2	5.3	3.3-8.2	1.2	1.2	0.4-2.7
	1991-1995	8.4	4.8	3.4-6.5	6.6	8.5	5.8-12.1	1.8	1.8	0.8-3.4	7.0	4.0	2.8-5.6	5.2	6.8	4.4-10.1	1.8	1.8	0.8-3.4
	1996-2000	11.4	6.2	4.7-8.0	9.0	10.4	7.6-13.9	2.4	2.3	1.2-4.1	8.4	4.5	3.3-6.1	5.8	6.8	4.5-9.7	2.6	2.5	1.3-4.3
	2001-2005	11.6	5.8	4.4-7.5	9.0	10.3	7.4-13.9	2.6	2.3	1.2-4.1	14.2	7.1	5.6-9.0	12.6	14.2	10.8-18.2	1.6	1.4	0.6-2.8
	2006-2010	11.8	5.1	3.9-6.7	8.6	8.1	5.8-11.1	3.2	2.6	1.4-4.3	9.4	4.1	3.0-5.6	7.6	7.4	5.2-10.3	1.8	1.4	0.6-2.8
	2011-2015	12.2	5.1	3.9-6.6	9.6	8.9	6.5-12.0	2.6	1.8	0.9-3.2	11.8	5.0	3.7-6.5	8.8	8.2	5.9-11.2	3.0	2.0	1.1-3.5
2016-2020	17.2	6.5	5.2-8.1	12.6	10.1	7.7-13.1	4.6	3.2	2.0-5.1	13.0	4.8	3.7-6.2	9.2	7.2	5.2-9.8	3.8	2.5	1.5-4.2	
Stomach	1976-1980	14.2	9.4	7.3-11.8	9.6	15.2	11.0-20.4	4.6	5.2	3.3-8.0	12.4	8.1	6.1-10.4	8.2	12.7	9.0-17.4	4.2	4.6	2.9-7.2
	1981-1985	16.4	10.5	8.3-13.1	10.0	15.2	11.2-20.2	6.4	6.9	4.7-9.8	12.6	8.3	6.4-10.7	7.8	12.4	8.7-17.0	4.8	5.3	3.4-8.0
	1986-1990	16.2	10.0	7.9-12.4	9.2	13.6	9.9-18.3	7.0	7.3	5.1-10.2	11.4	7.0	5.3-9.1	7.0	10.2	7.0-14.2	4.4	4.6	2.9-7.1
	1991-1995	17.0	9.9	7.9-12.3	9.8	14.1	10.3-18.8	7.2	6.8	4.7-9.4	13.8	8.2	6.4-10.4	7.8	12.0	8.4-16.5	6.0	5.7	3.8-8.2
	1996-2000	12.8	6.9	5.3-8.9	8.2	9.8	7.0-13.4	4.6	4.3	2.7-6.5	7.8	4.2	3.0-5.8	4.2	5.3	3.2-8.1	3.6	3.2	1.9-5.2
	2001-2005	14.4	7.2	5.7-9.1	11.2	13.2	9.9-17.2	3.2	2.9	1.6-4.8	4.2	2.1	1.3-3.2	2.0	2.6	1.2-4.8	2.2	1.9	0.9-3.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Ulster**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	12.2	5.6	4.2-7.2	8.4	8.4	6.0-11.4	3.8	3.2	1.9-5.1	5.2	2.4	1.6-3.6	2.6	2.6	1.4-4.6	2.6	2.3	1.2-4.0
	2011-2015	15.6	6.5	5.1-8.2	11.6	10.2	7.7-13.4	4.0	3.3	1.9-5.2	6.2	2.5	1.7-3.6	4.2	3.8	2.3-5.9	2.0	1.4	0.7-2.8
	2016-2020	13.8	5.3	4.1-6.8	9.2	7.2	5.2-9.8	4.6	3.7	2.3-5.8	4.2	1.6	1.0-2.5	3.2	2.7	1.5-4.6	1.0	0.6	0.2-1.7
Colorectal	1976-1980	96.0	64.1	58.4-70.2	48.0	76.1	66.4-86.7	48.0	55.7	48.8-63.3	49.4	33.4	29.3-37.9	23.4	38.3	31.4-46.2	26.0	30.1	25.1-35.8
	1981-1985	114.2	72.4	66.5-78.7	54.8	82.6	72.8-93.3	59.4	65.5	58.2-73.5	52.4	34.1	30.0-38.5	23.6	36.9	30.3-44.3	28.8	32.0	26.9-37.8
	1986-1990	107.6	67.4	61.7-73.4	53.4	79.9	70.4-90.3	54.2	58.3	51.5-65.8	53.0	33.1	29.2-37.4	25.2	38.1	31.6-45.6	27.8	29.2	24.5-34.6
	1991-1995	99.4	57.7	52.7-63.0	50.2	69.6	61.0-78.9	49.2	49.4	43.3-56.0	48.2	28.1	24.7-31.9	23.6	32.9	27.1-39.5	24.6	24.1	20.0-28.8
	1996-2000	129.2	70.2	64.9-75.9	62.0	76.9	68.5-86.1	67.2	62.6	56.0-69.8	48.0	26.1	22.9-29.7	22.6	29.5	24.2-35.6	25.4	22.7	18.9-27.1
	2001-2005	108.4	54.3	49.8-59.1	57.8	66.1	58.6-74.4	50.6	44.7	39.3-50.7	43.6	22.0	19.2-25.1	22.4	26.7	21.9-32.2	21.2	17.9	14.6-21.7
	2006-2010	111.4	50.9	46.7-55.4	55.0	56.4	49.8-63.7	56.4	46.2	40.9-52.1	37.0	16.9	14.5-19.5	17.6	19.1	15.2-23.6	19.4	15.4	12.5-19.0
	2011-2015	91.2	38.9	35.3-42.8	46.0	41.1	35.8-47.0	45.2	36.8	32.0-42.3	34.2	14.1	12.1-16.5	16.6	14.8	11.7-18.5	17.6	13.2	10.5-16.5
2016-2020	98.0	39.6	36.0-43.5	52.6	46.1	40.5-52.3	45.4	34.3	29.7-39.4	31.2	12.4	10.5-14.7	16.0	14.2	11.2-17.9	15.2	10.8	8.4-13.8	
Colon excluding rectum	1976-1980	65.0	43.4	38.7-48.5	28.4	45.4	37.9-53.8	36.6	42.0	36.1-48.7	38.8	26.4	22.8-30.5	17.8	29.2	23.3-36.2	21.0	24.4	19.9-29.6
	1981-1985	85.4	54.4	49.3-59.9	37.4	57.0	48.9-66.1	48.0	52.8	46.3-60.0	45.0	29.2	25.5-33.3	20.4	31.8	25.8-38.8	24.6	27.3	22.7-32.7
	1986-1990	80.6	50.7	45.8-55.9	36.6	55.8	47.8-64.7	44.0	47.4	41.2-54.2	47.2	29.5	25.8-33.6	21.8	33.3	27.2-40.3	25.4	26.7	22.2-31.9
	1991-1995	70.8	41.2	37.0-45.8	33.0	46.5	39.5-54.3	37.8	37.6	32.4-43.4	43.0	25.2	21.9-28.8	20.4	28.5	23.1-34.8	22.6	22.1	18.2-26.6
	1996-2000	93.0	50.6	46.1-55.5	40.4	50.8	43.9-58.4	52.6	48.7	43.0-55.0	42.4	23.1	20.1-26.4	18.6	24.2	19.4-29.8	23.8	21.1	17.5-25.4
	2001-2005	75.8	38.3	34.5-42.3	39.8	46.6	40.2-53.7	36.0	31.4	27.0-36.5	37.8	19.1	16.5-22.1	19.8	23.8	19.2-29.1	18.0	15.1	12.1-18.7
	2006-2010	82.6	37.9	34.3-41.8	38.0	39.8	34.2-46.0	44.6	36.4	31.7-41.6	30.6	14.0	11.8-16.4	14.0	15.2	11.8-19.4	16.6	13.2	10.5-16.6
	2011-2015	66.8	28.4	25.4-31.8	33.0	29.8	25.3-35.0	33.8	27.1	23.0-31.7	27.4	11.3	9.5-13.5	13.0	11.7	9.0-15.1	14.4	10.5	8.2-13.4
2016-2020	66.0	26.8	23.9-30.0	33.2	29.9	25.4-35.1	32.8	24.4	20.6-28.8	22.4	8.8	7.2-10.7	11.6	10.3	7.8-13.5	10.8	7.6	5.6-10.1	
Rectum & rectosigmoid	1976-1980	31.0	20.7	17.6-24.3	19.6	30.7	24.7-37.6	11.4	13.7	10.3-17.8	10.6	7.0	5.2-9.2	5.6	9.1	5.9-13.3	5.0	5.7	3.7-8.5
	1981-1985	28.8	18.0	15.2-21.3	17.4	25.5	20.3-31.7	11.4	12.7	9.6-16.5	7.4	4.9	3.4-6.8	3.2	5.0	2.8-8.2	4.2	4.7	2.9-7.2
	1986-1990	27.0	16.7	14.0-19.8	16.8	24.1	19.1-29.9	10.2	11.0	8.1-14.5	5.8	3.6	2.4-5.1	3.4	4.8	2.8-7.7	2.4	2.5	1.3-4.4
	1991-1995	28.6	16.5	13.9-19.4	17.2	23.1	18.4-28.7	11.4	11.8	8.9-15.3	5.2	3.0	1.9-4.4	3.2	4.4	2.5-7.1	2.0	2.0	0.9-3.7
	1996-2000	36.2	19.6	16.9-22.7	21.6	26.2	21.4-31.7	14.6	13.9	10.9-17.5	5.6	3.0	2.0-4.4	4.0	5.3	3.2-8.2	1.6	1.6	0.7-3.2
	2001-2005	32.6	16.1	13.7-18.7	18.0	19.5	15.7-24.1	14.6	13.2	10.4-16.7	5.8	2.9	1.9-4.1	2.6	2.9	1.5-5.0	3.2	2.8	1.6-4.6
	2006-2010	28.8	13.0	10.9-15.3	17.0	16.6	13.2-20.7	11.8	9.9	7.5-12.8	6.4	2.9	2.0-4.1	3.6	3.8	2.2-6.2	2.8	2.2	1.2-3.8
	2011-2015	24.4	10.5	8.7-12.7	13.0	11.3	8.6-14.6	11.4	9.8	7.3-12.9	6.8	2.8	1.9-4.0	3.6	3.1	1.8-5.0	3.2	2.7	1.5-4.6
2016-2020	32.0	12.8	10.8-15.1	19.4	16.1	13.0-19.9	12.6	9.9	7.5-12.9	8.8	3.6	2.6-4.9	4.4	3.9	2.4-6.1	4.4	3.2	2.0-5.1	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Ulster**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	4.2	2.8	1.7-4.3	2.6	4.3	2.2-7.4	1.6	1.7	0.7-3.5	2.6	1.8	0.9-3.0	1.4	2.6	1.0-5.3	1.2	1.3	0.5-2.8
	1981-1985	6.4	4.0	2.7-5.7	3.4	5.2	3.0-8.4	3.0	3.2	1.8-5.3	5.0	3.2	2.0-4.7	3.2	5.2	2.9-8.4	1.8	2.0	0.9-3.9
	1986-1990	3.8	2.2	1.3-3.5	2.6	3.5	1.8-6.0	1.2	1.3	0.5-2.9	4.4	2.6	1.6-4.0	2.6	3.5	1.8-6.0	1.8	1.9	0.9-3.8
	1991-1995	7.6	4.4	3.1-6.1	5.4	7.1	4.6-10.4	2.2	2.2	1.1-4.0	5.2	3.0	1.9-4.4	3.8	5.1	3.0-8.0	1.4	1.5	0.6-3.1
	1996-2000	11.0	6.0	4.5-7.8	6.2	7.5	5.0-10.6	4.8	4.6	2.9-6.9	8.8	4.8	3.5-6.4	3.6	4.2	2.5-6.6	5.2	4.7	3.1-7.0
	2001-2005	14.8	7.4	5.8-9.3	11.4	12.1	9.1-15.8	3.4	3.3	1.9-5.3	12.8	6.4	4.9-8.2	9.6	10.5	7.7-14.0	3.2	2.8	1.6-4.6
	2006-2010	16.4	7.1	5.6-8.8	13.2	12.0	9.2-15.4	3.2	2.5	1.4-4.3	12.8	5.8	4.4-7.4	9.6	9.4	6.9-12.6	3.2	2.8	1.6-4.6
	2011-2015	24.4	9.6	7.9-11.5	18.8	15.5	12.4-19.2	5.6	4.3	2.8-6.4	16.8	6.7	5.3-8.3	12.6	10.7	8.1-13.9	4.2	3.1	1.9-4.9
2016-2020	26.6	9.6	8.0-11.5	20.8	15.4	12.5-18.9	5.8	4.4	2.9-6.6	17.4	6.4	5.1-8.0	12.2	9.3	7.1-12.2	5.2	3.9	2.5-5.9	
Pancreas	1976-1980	22.4	14.9	12.2-17.9	13.6	21.7	16.7-27.7	8.8	9.8	7.1-13.2	20.2	13.6	11.0-16.6	11.8	19.3	14.5-25.1	8.4	9.5	6.8-12.9
	1981-1985	14.6	9.4	7.4-11.9	6.4	10.1	6.8-14.4	8.2	9.1	6.5-12.4	14.2	9.5	7.4-12.0	7.8	12.3	8.6-16.9	6.4	7.5	5.1-10.6
	1986-1990	20.8	12.7	10.4-15.5	8.6	12.7	9.1-17.2	12.2	12.8	9.8-16.5	18.6	11.4	9.2-14.0	7.0	10.7	7.4-15.0	11.6	11.9	9.0-15.5
	1991-1995	21.2	12.3	10.1-14.9	12.0	16.8	12.7-21.8	9.2	8.9	6.5-12.0	20.4	11.9	9.7-14.5	10.6	15.2	11.2-19.9	9.8	9.6	7.1-12.8
	1996-2000	21.8	11.8	9.7-14.2	11.0	14.0	10.5-18.3	10.8	9.9	7.4-12.9	20.2	10.9	8.9-13.3	11.0	13.8	10.4-18.0	9.2	8.3	6.1-11.1
	2001-2005	29.8	14.8	12.5-17.4	14.0	15.7	12.2-20.0	15.8	14.2	11.2-17.7	26.6	13.4	11.2-15.9	12.6	14.8	11.3-19.1	14.0	12.3	9.6-15.6
	2006-2010	27.6	12.6	10.6-15.0	13.2	13.1	10.0-16.7	14.4	11.7	9.1-14.9	25.4	11.6	9.7-13.9	11.8	11.9	9.0-15.5	13.6	11.0	8.5-14.1
	2011-2015	30.2	12.3	10.3-14.5	17.0	15.1	12.0-18.8	13.2	10.0	7.7-12.9	22.0	8.9	7.3-10.8	11.6	10.4	7.8-13.6	10.4	7.8	5.8-10.4
2016-2020	34.6	12.9	11.0-15.1	19.2	15.3	12.3-18.8	15.4	10.8	8.4-13.7	26.2	9.6	8.0-11.5	14.6	11.4	8.9-14.5	11.6	8.0	6.0-10.5	
Larynx	1976-1980	5.8	3.8	2.5-5.4	5.2	7.8	5.0-11.5	0.6	0.7	0.2-2.2	2.0	1.3	0.6-2.4	1.6	2.5	1.0-5.0	0.4	0.5	0.1-1.8
	1981-1985	7.4	4.7	3.3-6.6	5.8	8.4	5.6-12.2	1.6	1.9	0.8-3.7	2.6	1.5	0.8-2.7	2.2	3.2	1.5-5.9	0.4	0.4	0.1-1.7
	1986-1990	8.6	5.5	3.9-7.4	6.4	9.4	6.4-13.4	2.2	2.5	1.2-4.5	4.0	2.4	1.5-3.7	2.8	4.0	2.1-6.7	1.2	1.2	0.4-2.7
	1991-1995	7.8	4.4	3.2-6.1	5.4	6.6	4.3-9.6	2.4	2.7	1.4-4.7	4.2	2.5	1.5-3.8	2.6	3.6	1.8-6.1	1.6	1.7	0.7-3.3
	1996-2000	8.0	4.3	3.1-5.9	6.2	7.5	5.1-10.7	1.8	1.8	0.8-3.5	2.6	1.4	0.8-2.4	2.0	2.6	1.2-4.8	0.6	0.6	0.1-1.9
	2001-2005	10.0	4.9	3.6-6.5	8.0	8.3	5.9-11.3	2.0	1.9	0.9-3.5	3.6	1.7	1.0-2.8	2.8	2.9	1.6-5.0	0.8	0.7	0.2-1.9
	2006-2010	10.0	4.7	3.5-6.3	8.2	8.4	5.9-11.4	1.8	1.6	0.7-3.1	2.2	1.0	0.5-1.8	1.4	1.4	0.5-2.9	0.8	0.6	0.2-1.7
	2011-2015	7.2	2.8	2.0-4.0	5.8	5.0	3.3-7.3	1.4	1.0	0.4-2.2	2.4	0.9	0.5-1.7	2.0	1.6	0.8-3.1	0.4	0.3	0.0-1.3
2016-2020	6.4	2.4	1.6-3.4	5.0	3.6	2.3-5.6	1.4	1.1	0.4-2.6	2.0	0.7	0.4-1.5	1.4	1.1	0.4-2.4	0.6	0.5	0.1-1.6	
Lung and bronchus	1976-1980	92.4	60.3	54.8-66.2	63.2	95.6	85.0-107.1	29.2	35.1	29.5-41.4	74.0	48.3	43.4-53.5	52.2	80.5	70.6-91.2	21.8	25.6	20.9-31.0
	1981-1985	99.6	62.6	57.2-68.5	66.2	98.5	87.8-110.0	33.4	38.1	32.4-44.5	83.0	52.5	47.5-57.9	57.6	86.4	76.4-97.3	25.4	29.6	24.5-35.3
	1986-1990	115.0	70.2	64.5-76.3	74.4	106.6	95.6-118.3	40.6	45.3	39.2-52.1	90.6	55.0	50.0-60.4	60.4	85.9	76.2-96.5	30.2	32.7	27.6-38.5
	1991-1995	122.0	70.2	64.7-76.0	73.4	97.0	87.2-107.7	48.6	50.3	44.1-57.1	102.8	59.3	54.3-64.7	64.6	86.8	77.4-97.0	38.2	39.6	34.1-45.7
	1996-2000	140.2	75.9	70.4-81.8	76.8	93.7	84.4-103.7	63.4	61.8	55.1-69.1	107.0	58.1	53.3-63.2	59.6	74.2	65.9-83.2	47.4	45.4	39.8-51.7
	2001-2005	148.6	74.1	68.9-79.7	73.0	81.5	73.2-90.5	75.6	68.6	61.9-76.0	111.2	55.8	51.2-60.7	55.4	63.4	56.0-71.5	55.8	50.3	44.5-56.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Ulster**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	157.2	71.9	66.9-77.2	76.2	78.5	70.7-87.0	81.0	68.2	61.6-75.3	108.8	49.9	45.7-54.4	55.4	57.0	50.4-64.4	53.4	44.9	39.6-50.7
	2011-2015	167.6	69.0	64.3-73.9	85.4	76.7	69.4-84.6	82.2	62.8	56.7-69.4	108.2	45.1	41.3-49.1	56.4	51.7	45.7-58.3	51.8	39.4	34.7-44.8
	2016-2020	166.8	61.3	57.1-65.8	81.0	64.3	58.0-71.2	85.8	58.9	53.4-65.0	95.8	35.7	32.5-39.1	51.2	41.8	36.7-47.4	44.6	30.5	26.6-35.0
Melanoma of the skin	1976-1980	10.4	7.3	5.4-9.6	5.6	8.5	5.6-12.3	4.8	6.3	4.0-9.5	4.6	3.1	1.9-4.7	3.6	5.3	3.1-8.4	1.0	1.1	0.4-2.7
	1981-1985	12.4	8.5	6.5-10.9	7.0	10.9	7.5-15.3	5.4	7.2	4.7-10.6	4.6	3.1	1.9-4.7	3.4	5.3	3.0-8.6	1.2	1.4	0.5-3.2
	1986-1990	9.8	6.2	4.5-8.2	6.0	8.6	5.7-12.4	3.8	4.3	2.5-6.8	5.2	3.4	2.2-5.0	3.4	5.4	3.1-8.6	1.8	2.2	1.0-4.3
	1991-1995	13.0	7.6	5.9-9.7	7.8	10.4	7.3-14.3	5.2	5.7	3.7-8.4	6.8	3.9	2.7-5.5	5.0	7.2	4.6-10.7	1.8	1.8	0.8-3.5
	1996-2000	21.6	11.7	9.6-14.2	12.0	14.5	11.0-18.7	9.6	9.9	7.3-13.1	6.2	3.3	2.3-4.8	4.0	4.6	2.8-7.2	2.2	2.2	1.1-4.0
	2001-2005	28.6	14.3	12.1-16.9	15.4	16.8	13.2-21.1	13.2	12.8	9.9-16.4	7.0	3.6	2.5-5.0	4.4	5.2	3.2-8.0	2.6	2.7	1.4-4.7
	2006-2010	43.4	20.3	17.6-23.2	27.6	29.3	24.5-34.8	15.8	13.9	10.9-17.4	7.0	3.2	2.2-4.5	5.0	5.2	3.3-7.8	2.0	1.6	0.8-3.1
	2011-2015	41.8	18.4	15.9-21.2	21.6	19.6	16.0-23.9	20.2	18.5	14.9-22.7	6.8	2.8	1.9-4.0	4.6	4.1	2.6-6.3	2.2	1.9	0.9-3.5
	2016-2020	42.4	17.0	14.7-19.6	23.6	21.1	17.3-25.5	18.8	14.1	11.2-17.5	5.0	2.2	1.4-3.3	1.6	1.5	0.6-3.0	3.4	2.9	1.6-5.0
Female breast	1976-1980							70.8	90.6	81.2-100.8							28.0	34.9	29.3-41.4
	1981-1985							88.4	107.3	97.2-118.0							31.2	36.2	30.6-42.4
	1986-1990							96.4	113.2	103.1-124.0							32.6	37.1	31.5-43.3
	1991-1995							119.8	131.3	120.9-142.4							38.0	40.3	34.7-46.5
	1996-2000							138.8	140.4	130.1-151.3							33.8	33.5	28.6-39.0
	2001-2005							134.0	125.2	115.9-135.2							32.2	29.0	24.7-33.9
	2006-2010							154.8	133.5	124.0-143.5							26.6	22.3	18.6-26.6
	2011-2015							153.4	124.7	115.6-134.3							26.2	19.9	16.5-23.8
	2016-2020							156.8	126.2	116.9-136.1							29.2	20.8	17.5-24.8
Cervix uteri	1976-1980							9.0	11.2	8.1-15.1							5.4	6.2	4.0-9.1
	1981-1985							11.2	14.2	10.7-18.6							3.6	4.2	2.5-6.7
	1986-1990							11.6	14.2	10.7-18.4							4.2	4.7	2.9-7.2
	1991-1995							12.8	14.3	11.0-18.3							2.8	3.3	1.8-5.5
	1996-2000							8.8	9.4	6.8-12.7							2.8	3.0	1.6-5.0
	2001-2005							9.6	9.5	7.0-12.7							3.6	3.3	1.9-5.3
	2006-2010							11.6	11.3	8.5-14.7							3.4	3.0	1.7-4.8
	2011-2015							8.0	8.4	5.9-11.6							2.8	2.2	1.2-3.8
	2016-2020							6.8	7.2	4.9-10.3							2.2	1.9	0.9-3.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Ulster**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							19.8	23.4	18.9-28.6							5.4	5.9	3.9-8.7
	1981-1985							22.2	25.4	20.8-30.7							5.2	5.7	3.7-8.5
	1986-1990							22.4	26.5	21.7-32.1							3.8	4.4	2.6-6.9
	1991-1995							26.0	28.1	23.4-33.5							4.2	4.1	2.5-6.3
	1996-2000							28.8	29.1	24.5-34.3							3.6	3.4	2.0-5.5
	2001-2005							33.0	30.6	26.1-35.7							8.2	7.2	5.2-9.9
	2006-2010							36.0	30.3	25.9-35.2							7.2	6.1	4.2-8.5
	2011-2015							36.8	28.7	24.5-33.4							6.8	5.3	3.6-7.5
	2016-2020							43.0	31.5	27.2-36.3							9.2	6.6	4.8-9.1
Ovary	1976-1980							12.6	14.7	11.2-18.9							9.4	11.1	8.1-14.8
	1981-1985							14.4	17.2	13.3-21.7							9.2	10.3	7.5-13.9
	1986-1990							16.4	18.5	14.6-23.1							9.4	10.0	7.3-13.4
	1991-1995							16.6	17.6	14.0-21.9							11.2	11.3	8.5-14.8
	1996-2000							18.4	18.5	14.9-22.7							8.2	8.0	5.7-10.9
	2001-2005							16.6	15.6	12.4-19.4							13.4	11.8	9.1-15.1
	2006-2010							15.8	13.9	11.0-17.5							11.6	9.7	7.3-12.7
	2011-2015							16.8	14.3	11.2-17.9							12.0	9.9	7.5-12.9
	2016-2020							16.4	12.8	10.0-16.2							7.6	5.2	3.6-7.4
Prostate	1976-1980				42.0	72.2	62.4-83.1							15.8	29.8	23.3-37.3			
	1981-1985				53.2	87.8	77.2-99.4							19.8	35.3	28.5-43.1			
	1986-1990				55.6	88.8	78.4-100.2							21.8	37.0	30.2-44.8			
	1991-1995				93.8	133.5	121.3-146.5							26.2	42.3	35.2-50.4			
	1996-2000				130.8	161.5	149.2-174.6							26.8	42.0	35.0-49.8			
	2001-2005				144.2	160.4	148.7-172.7							21.0	29.2	23.8-35.4			
	2006-2010				150.4	144.9	134.5-155.9							19.2	22.8	18.4-27.9			
	2011-2015				129.2	105.2	97.0-113.9							16.4	16.8	13.3-21.0			
	2016-2020				133.6	99.7	92.1-107.8							19.0	16.6	13.4-20.5			
Testis	1976-1980				3.4	4.1	2.3-6.7							0.4	0.6	0.1-2.3			
	1981-1985				3.4	4.2	2.4-6.8							0.4	0.5	0.1-2.1			
	1986-1990				3.2	3.4	1.9-5.6							0.0	0.0	0.0-1.0			
	1991-1995				3.8	3.9	2.4-6.2							0.2	0.2	0.0-1.3			
	1996-2000				6.0	6.5	4.4-9.3							0.2	0.2	0.0-1.3			
	2001-2005				5.6	6.3	4.2-9.1							0.4	0.4	0.0-1.5			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Ulster**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				5.6	6.6	4.3-9.5							0.2	0.2	0.0-1.3			
	2011-2015				6.2	7.5	5.0-10.6							0.0	0.0	0.0-0.9			
	2016-2020				4.6	5.3	3.3-8.1							0.0	0.0	0.0-0.9			
Urinary bladder (incl. in situ)	1976-1980	30.6	20.9	17.6-24.5	22.6	37.8	30.9-45.7	8.0	9.0	6.4-12.4	14.6	10.0	7.8-12.6	11.2	19.4	14.4-25.3	3.4	4.0	2.3-6.5
	1981-1985	31.8	20.4	17.3-23.8	23.4	36.2	29.7-43.5	8.4	9.3	6.7-12.7	10.8	7.3	5.4-9.5	6.6	10.8	7.3-15.3	4.2	4.7	2.9-7.2
	1986-1990	36.8	22.4	19.2-25.9	27.8	40.9	34.2-48.5	9.0	9.4	6.8-12.6	10.0	6.2	4.6-8.2	6.4	9.7	6.6-13.8	3.6	3.7	2.2-5.9
	1991-1995	35.2	20.3	17.4-23.6	25.6	35.3	29.3-42.1	9.6	9.3	6.8-12.4	11.0	6.5	4.9-8.5	6.6	10.7	7.3-15.0	4.4	4.2	2.6-6.4
	1996-2000	42.8	23.2	20.2-26.6	31.2	39.8	33.7-46.8	11.6	11.0	8.4-14.3	9.8	5.4	4.0-7.1	6.6	9.6	6.5-13.5	3.2	3.0	1.7-4.9
	2001-2005	54.2	27.5	24.3-31.0	39.0	46.2	39.8-53.3	15.2	13.7	10.8-17.2	9.8	5.0	3.7-6.6	6.4	7.7	5.2-10.9	3.4	2.9	1.7-4.7
	2006-2010	56.6	25.6	22.7-28.9	45.2	46.3	40.3-52.9	11.4	9.7	7.3-12.7	9.0	4.2	3.1-5.7	6.6	7.7	5.3-10.9	2.4	2.0	1.0-3.6
	2011-2015	61.0	25.7	22.8-28.8	47.0	44.3	38.7-50.6	14.0	10.8	8.3-13.8	12.8	5.1	3.9-6.5	10.0	9.6	7.1-12.8	2.8	1.8	1.0-3.3
2016-2020	62.8	23.1	20.6-25.9	46.4	37.9	33.0-43.3	16.4	11.1	8.8-14.0	13.8	5.1	4.0-6.5	10.0	8.8	6.5-11.8	3.8	2.4	1.4-4.0	
Kidney and renal pelvis	1976-1980	13.0	8.5	6.5-10.9	7.2	11.0	7.6-15.4	5.8	6.8	4.5-9.8	7.0	4.8	3.3-6.7	4.6	7.7	4.8-11.7	2.4	2.7	1.4-4.8
	1981-1985	15.2	9.6	7.6-12.1	9.4	13.8	10.0-18.5	5.8	6.5	4.3-9.3	8.2	5.1	3.6-6.9	4.6	6.9	4.3-10.4	3.6	3.8	2.2-6.0
	1986-1990	14.2	8.6	6.7-10.9	8.6	11.9	8.6-16.2	5.6	6.3	4.1-9.1	6.6	4.1	2.8-5.8	4.0	5.9	3.6-9.1	2.6	2.8	1.5-4.8
	1991-1995	17.8	10.3	8.3-12.7	10.8	13.7	10.2-17.9	7.0	7.5	5.2-10.5	6.2	3.6	2.4-5.1	3.4	4.8	2.7-7.6	2.8	2.8	1.5-4.7
	1996-2000	25.8	14.1	11.8-16.7	15.2	18.6	14.6-23.4	10.6	10.5	7.9-13.8	9.8	5.3	3.9-7.0	6.2	7.5	5.1-10.7	3.6	3.4	2.0-5.4
	2001-2005	25.4	12.8	10.7-15.3	15.2	16.1	12.6-20.2	10.2	9.5	7.1-12.6	8.0	4.0	2.8-5.4	5.4	6.4	4.2-9.4	2.6	2.3	1.2-4.0
	2006-2010	36.4	16.8	14.4-19.5	23.6	23.4	19.3-28.2	12.8	11.0	8.4-14.2	9.4	4.3	3.1-5.7	6.8	7.4	5.1-10.5	2.6	2.0	1.1-3.6
	2011-2015	34.0	14.2	12.1-16.6	21.6	19.5	15.9-23.8	12.4	9.6	7.3-12.5	7.0	2.9	2.0-4.1	3.8	3.6	2.2-5.8	3.2	2.2	1.2-3.8
2016-2020	38.2	15.8	13.5-18.4	26.0	21.4	17.7-25.6	12.2	10.9	8.1-14.3	8.0	3.1	2.2-4.3	4.2	3.2	2.0-5.1	3.8	2.8	1.6-4.6	
Brain and other central nervous system	1976-1980	10.0	5.9	4.3-7.8	6.4	8.0	5.5-11.5	3.6	3.8	2.2-6.1	7.0	4.4	3.0-6.1	3.6	4.8	2.8-7.7	3.4	3.9	2.2-6.3
	1981-1985	10.0	6.4	4.7-8.5	5.2	7.4	4.8-10.9	4.8	5.7	3.6-8.6	7.4	4.7	3.3-6.5	3.8	5.4	3.2-8.6	3.6	4.1	2.4-6.6
	1986-1990	10.0	6.1	4.5-8.0	5.8	7.2	4.8-10.5	4.2	4.9	3.0-7.5	4.8	2.9	1.9-4.4	3.2	4.3	2.4-7.0	1.6	1.7	0.7-3.5
	1991-1995	13.2	7.6	5.9-9.7	7.8	9.9	7.0-13.6	5.4	5.9	3.9-8.6	6.4	3.7	2.5-5.3	4.2	5.3	3.3-8.2	2.2	2.4	1.2-4.3
	1996-2000	12.4	6.9	5.3-8.9	7.8	9.2	6.5-12.5	4.6	4.9	3.1-7.4	6.6	3.6	2.5-5.1	4.0	4.8	2.9-7.5	2.6	2.6	1.4-4.5
	2001-2005	16.0	8.4	6.7-10.5	9.2	10.7	7.7-14.3	6.8	6.7	4.6-9.5	9.4	4.7	3.5-6.3	5.8	6.2	4.1-9.0	3.6	3.3	2.0-5.3
	2006-2010	14.0	6.8	5.3-8.7	6.0	6.2	4.2-9.0	8.0	7.7	5.4-10.7	9.8	4.7	3.4-6.2	5.0	5.2	3.3-7.7	4.8	4.3	2.7-6.5
	2011-2015	15.2	7.6	5.9-9.6	8.0	8.3	5.8-11.5	7.2	6.9	4.7-9.8	8.4	3.6	2.6-5.0	4.2	4.1	2.5-6.4	4.2	3.3	2.0-5.2
2016-2020	15.0	6.5	5.0-8.3	9.2	8.0	5.7-10.8	5.8	4.9	3.2-7.4	10.2	3.9	2.9-5.3	5.6	4.5	2.9-6.8	4.6	3.2	2.0-5.0	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Ulster**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	3.6	2.2	1.3-3.6	1.2	1.5	0.5-3.4	2.4	3.0	1.5-5.3	0.8	0.5	0.1-1.3	0.4	0.6	0.1-2.2	0.4	0.4	0.0-1.6
	1981-1985	3.2	1.9	1.1-3.1	0.8	1.2	0.3-3.0	2.4	2.5	1.3-4.5	1.2	0.8	0.3-1.8	0.4	0.7	0.1-2.3	0.8	1.0	0.2-2.5
	1986-1990	5.0	3.1	2.0-4.7	1.6	2.2	0.9-4.3	3.4	4.2	2.4-6.8	0.2	0.1	0.0-0.7	0.2	0.3	0.0-1.5	0.0	0.0	0.0-0.9
	1991-1995	7.2	4.1	2.8-5.6	1.4	1.6	0.7-3.4	5.8	6.5	4.3-9.4	0.4	0.2	0.0-0.8	0.2	0.2	0.0-1.3	0.2	0.2	0.0-1.1
	1996-2000	10.8	6.0	4.5-7.8	3.6	4.1	2.4-6.6	7.2	7.8	5.5-10.9	1.0	0.6	0.2-1.3	0.0	0.0	0.0-0.9	1.0	0.9	0.3-2.2
	2001-2005	17.2	8.9	7.1-11.1	3.2	3.3	1.9-5.4	14.0	14.5	11.3-18.4	0.6	0.3	0.1-0.9	0.4	0.5	0.1-1.9	0.2	0.2	0.0-1.1
	2006-2010	29.0	14.6	12.3-17.3	7.4	7.8	5.4-10.8	21.6	22.1	18.0-26.8	0.8	0.4	0.1-1.0	0.2	0.1	0.0-1.1	0.6	0.5	0.1-1.7
	2011-2015	32.6	15.9	13.4-18.6	7.8	6.9	4.8-9.6	24.8	25.1	20.7-30.2	1.2	0.5	0.2-1.2	0.4	0.4	0.0-1.5	0.8	0.6	0.2-1.8
2016-2020	30.4	15.4	12.9-18.2	7.0	6.4	4.3-9.2	23.4	24.6	20.1-29.8	1.4	0.5	0.2-1.1	0.8	0.7	0.2-1.9	0.6	0.3	0.1-1.3	
Hodgkin lymphoma	1976-1980	5.4	3.2	2.1-4.8	3.2	4.2	2.3-7.0	2.2	2.4	1.2-4.4	1.6	0.9	0.4-1.9	1.0	1.3	0.4-3.3	0.6	0.6	0.1-1.8
	1981-1985	4.8	2.7	1.7-4.1	1.8	2.2	1.0-4.4	3.0	3.3	1.8-5.5	1.4	1.0	0.4-2.0	0.8	1.1	0.3-2.9	0.6	0.7	0.1-2.2
	1986-1990	4.4	2.4	1.5-3.7	2.6	2.8	1.5-4.9	1.8	2.0	0.9-3.8	1.8	1.1	0.5-2.0	1.4	1.7	0.6-3.5	0.4	0.4	0.1-1.6
	1991-1995	7.6	4.4	3.1-6.0	5.6	6.5	4.3-9.5	2.0	2.3	1.1-4.2	0.8	0.5	0.1-1.2	0.6	0.7	0.1-2.1	0.2	0.2	0.0-1.3
	1996-2000	6.2	3.5	2.3-4.9	4.0	4.6	2.8-7.2	2.2	2.3	1.2-4.2	1.2	0.6	0.2-1.4	0.2	0.2	0.0-1.3	1.0	1.0	0.3-2.4
	2001-2005	8.0	4.4	3.2-6.1	4.6	5.4	3.4-8.1	3.4	3.9	2.3-6.3	1.2	0.7	0.2-1.4	0.8	1.0	0.3-2.7	0.4	0.4	0.0-1.4
	2006-2010	9.6	4.9	3.6-6.6	5.6	5.9	3.9-8.6	4.0	4.0	2.4-6.3	1.0	0.6	0.2-1.3	0.6	0.6	0.1-1.9	0.4	0.5	0.1-1.8
	2011-2015	6.6	3.4	2.3-4.9	4.0	4.5	2.7-7.0	2.6	2.6	1.3-4.6	1.0	0.5	0.1-1.2	0.8	0.9	0.2-2.3	0.2	0.1	0.0-1.1
2016-2020	6.6	3.6	2.4-5.1	3.6	3.8	2.2-6.1	3.0	3.5	1.9-5.8	1.2	0.5	0.2-1.1	1.2	1.0	0.4-2.4	0.0	0.0	0.0-0.9	
Non-Hodgkin Lymphomas	1976-1980	18.4	12.0	9.6-14.8	9.2	13.0	9.4-17.4	9.2	10.8	7.9-14.5	8.6	5.8	4.2-7.9	4.8	7.2	4.6-10.9	3.8	4.6	2.8-7.3
	1981-1985	18.8	11.9	9.6-14.6	9.4	13.8	10.0-18.5	9.4	10.2	7.5-13.6	11.0	7.0	5.3-9.2	5.4	8.3	5.4-12.1	5.6	6.0	4.0-8.8
	1986-1990	22.6	13.9	11.4-16.8	14.4	19.8	15.4-25.1	8.2	9.1	6.5-12.4	12.2	7.7	5.9-9.9	7.2	11.2	7.7-15.6	5.0	5.4	3.5-8.0
	1991-1995	32.0	18.5	15.7-21.6	18.0	23.2	18.5-28.6	14.0	14.6	11.4-18.5	15.2	8.9	7.0-11.1	8.6	12.1	8.7-16.3	6.6	6.7	4.6-9.5
	1996-2000	35.0	19.1	16.3-22.1	17.6	21.7	17.3-26.8	17.4	16.6	13.3-20.6	13.4	7.3	5.7-9.3	7.2	9.3	6.5-12.9	6.2	5.8	3.9-8.2
	2001-2005	39.2	19.6	17.0-22.6	20.2	22.0	17.9-26.9	19.0	17.8	14.4-21.8	14.4	7.3	5.7-9.2	6.6	7.3	5.0-10.3	7.8	6.9	4.9-9.6
	2006-2010	51.4	24.1	21.2-27.3	26.8	27.8	23.2-33.1	24.6	21.1	17.5-25.4	13.0	6.1	4.7-7.9	7.2	7.8	5.4-10.8	5.8	4.7	3.1-6.9
	2011-2015	50.0	22.0	19.3-25.1	28.0	26.6	22.3-31.6	22.0	18.7	15.2-22.9	15.2	6.2	4.8-7.8	8.8	8.4	6.1-11.4	6.4	4.6	3.1-6.8
2016-2020	52.4	20.9	18.3-23.7	28.6	24.0	20.1-28.5	23.8	18.5	15.2-22.4	13.6	5.2	4.0-6.7	8.2	7.2	5.1-9.9	5.4	3.6	2.3-5.4	
Myeloma	1976-1980	5.4	3.7	2.4-5.4	2.2	3.6	1.7-6.4	3.2	4.0	2.2-6.5	5.4	3.7	2.4-5.4	2.6	4.2	2.2-7.2	2.8	3.6	1.9-6.0
	1981-1985	9.0	6.1	4.4-8.1	3.4	5.8	3.3-9.3	5.6	6.4	4.2-9.3	5.4	3.6	2.3-5.2	2.6	4.3	2.2-7.4	2.8	3.1	1.7-5.2
	1986-1990	5.6	3.6	2.4-5.2	2.6	3.5	1.9-6.0	3.0	3.3	1.8-5.5	4.2	2.7	1.7-4.1	1.6	2.5	1.0-4.9	2.6	2.8	1.5-4.9
	1991-1995	9.4	5.5	4.0-7.3	4.8	6.4	4.0-9.7	4.6	4.7	3.0-7.2	6.0	3.5	2.4-5.1	2.4	3.5	1.8-6.2	3.6	3.6	2.1-5.7
	1996-2000	11.2	6.1	4.6-7.9	6.6	8.5	5.8-12.0	4.6	4.4	2.8-6.7	6.4	3.5	2.4-4.9	3.6	4.6	2.7-7.3	2.8	2.6	1.4-4.5
	2001-2005	9.4	4.7	3.5-6.3	5.2	6.0	3.9-8.8	4.2	3.6	2.2-5.6	5.8	2.9	1.9-4.1	3.8	4.5	2.7-7.2	2.0	1.8	0.8-3.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Ulster**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	13.2	6.0	4.7-7.7	8.0	8.4	5.9-11.5	5.2	4.2	2.7-6.3	4.4	2.1	1.3-3.2	2.4	2.8	1.4-5.0	2.0	1.6	0.7-3.0
	2011-2015	16.8	7.1	5.6-8.9	9.6	8.9	6.5-11.9	7.2	5.7	3.9-8.1	5.4	2.3	1.5-3.4	2.0	2.0	0.9-3.8	3.4	2.4	1.4-4.0
	2016-2020	17.2	6.6	5.2-8.3	11.6	9.5	7.1-12.5	5.6	4.3	2.8-6.5	6.6	2.4	1.7-3.5	4.2	3.4	2.1-5.4	2.4	1.6	0.8-3.0
Leukemias	1976-1980	15.4	10.2	8.0-12.8	9.2	13.6	9.8-18.3	6.2	7.6	5.1-10.8	9.4	6.3	4.6-8.4	4.6	7.0	4.4-10.7	4.8	6.0	3.8-9.0
	1981-1985	17.2	11.4	9.1-14.1	9.4	14.8	10.7-19.8	7.8	9.1	6.4-12.5	10.0	6.7	5.0-8.9	5.8	9.8	6.4-14.2	4.2	5.0	3.0-7.6
	1986-1990	22.6	14.2	11.7-17.1	10.6	16.3	12.1-21.4	12.0	13.1	10.0-17.0	14.0	8.9	6.9-11.2	7.8	12.6	8.8-17.3	6.2	6.8	4.6-9.7
	1991-1995	24.6	14.3	11.9-17.1	15.0	19.6	15.3-24.7	9.6	10.3	7.6-13.8	14.0	8.1	6.3-10.3	8.8	12.3	8.8-16.6	5.2	5.4	3.5-7.9
	1996-2000	29.6	16.2	13.7-19.1	17.0	21.2	16.9-26.3	12.6	12.4	9.5-15.9	14.2	7.7	6.0-9.7	8.4	10.9	7.8-14.8	5.8	5.4	3.6-7.8
	2001-2005	31.4	16.2	13.7-18.9	17.6	20.3	16.2-25.1	13.8	12.7	9.9-16.2	15.8	8.0	6.4-10.0	7.4	8.7	6.1-12.1	8.4	7.1	5.1-9.8
	2006-2010	30.0	14.0	11.8-16.5	18.4	19.4	15.6-23.9	11.6	10.3	7.7-13.5	14.2	6.7	5.2-8.5	8.2	9.3	6.6-12.7	6.0	5.0	3.3-7.2
	2011-2015	38.6	17.1	14.7-19.9	23.8	22.6	18.6-27.3	14.8	12.4	9.6-15.8	18.4	8.1	6.4-10.0	10.8	11.1	8.3-14.7	7.6	6.0	4.1-8.4
2016-2020	34.8	13.7	11.6-16.0	20.6	17.5	14.2-21.5	14.2	10.2	7.8-13.2	16.4	6.3	5.0-7.9	9.8	8.3	6.1-11.2	6.6	4.7	3.1-6.9	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Warren**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	219.0	415.3	390.6-441.2	106.4	492.0	449.1-537.8	112.6	377.8	346.5-411.2	116.0	219.9	202.0-238.9	59.8	283.7	250.8-319.6	56.2	184.4	163.0-207.8
	1981-1985	269.4	481.1	455.3-508.0	130.8	575.5	530.5-623.1	138.6	436.1	403.3-470.8	136.4	244.5	226.2-263.9	68.0	309.6	276.1-345.7	68.4	209.8	187.6-233.9
	1986-1990	284.2	474.2	449.5-499.9	139.0	570.1	527.0-615.5	145.2	423.6	392.5-456.6	134.8	223.3	206.6-241.1	67.6	286.9	256.1-320.2	67.2	186.9	167.0-208.6
	1991-1995	305.4	464.7	441.5-488.8	153.0	557.7	517.9-599.7	152.4	412.5	383.2-443.6	138.8	208.9	193.5-225.1	67.8	257.5	230.1-287.2	71.0	179.3	160.8-199.4
	1996-2000	354.0	500.5	477.4-524.5	177.6	578.0	540.1-617.9	176.4	448.8	419.3-480.0	152.6	212.8	197.9-228.5	76.4	259.8	233.9-287.7	76.2	182.9	164.8-202.6
	2001-2005	408.2	528.7	505.8-552.3	212.0	618.7	581.4-657.7	196.2	465.1	436.1-495.8	159.4	204.4	190.4-219.2	79.4	243.8	220.0-269.4	80.0	179.4	162.1-198.3
	2006-2010	462.4	538.8	516.6-561.7	238.4	606.0	571.4-642.4	224.0	490.9	461.8-521.6	166.8	191.9	179.0-205.7	86.4	229.8	208.2-253.0	80.4	164.9	148.9-182.5
	2011-2015	500.4	545.6	523.5-568.4	257.8	599.0	565.6-633.9	242.6	507.0	477.2-538.4	173.0	178.3	166.3-191.0	91.8	215.5	195.8-236.9	81.2	151.8	136.9-168.1
	2016-2020	501.2	502.5	481.9-523.9	260.8	543.6	513.1-575.6	240.4	470.2	442.0-499.9	170.2	163.6	152.4-175.6	92.0	196.0	177.8-215.7	78.2	136.1	122.4-151.3
Oral cavity and pharynx	1976-1980	6.8	12.5	8.6-17.6	4.4	18.9	11.6-29.2	2.4	7.9	4.0-13.9	3.8	7.0	4.2-11.1	2.6	11.9	6.1-20.9	1.2	4.2	1.4-9.4
	1981-1985	7.2	12.4	8.6-17.3	4.2	17.6	10.7-27.3	3.0	9.3	5.1-15.5	2.6	4.5	2.4-7.9	1.6	6.8	2.8-13.8	1.0	3.4	1.0-8.1
	1986-1990	6.6	11.1	7.6-15.7	3.8	14.9	8.8-23.6	2.8	8.5	4.6-14.5	2.4	3.8	2.0-6.7	1.0	3.8	1.2-9.0	1.4	3.7	1.5-8.0
	1991-1995	7.4	11.6	8.1-16.1	4.4	15.4	9.6-23.5	3.0	8.1	4.5-13.7	2.0	3.2	1.5-5.8	1.6	5.5	2.3-10.9	0.4	1.1	0.1-4.3
	1996-2000	9.4	13.4	9.8-17.8	6.0	19.7	13.2-28.3	3.4	8.6	5.0-14.0	1.6	2.3	1.0-4.6	0.8	2.6	0.7-6.7	0.8	2.0	0.5-5.5
	2001-2005	11.4	14.6	11.1-19.1	7.4	20.6	14.4-28.7	4.0	9.6	5.8-15.2	3.6	4.5	2.6-7.2	1.8	5.3	2.4-10.3	1.8	4.0	1.8-7.9
	2006-2010	12.2	14.4	10.9-18.7	9.2	22.4	16.3-30.2	3.0	6.9	3.7-11.8	3.2	3.5	2.0-5.9	2.2	5.3	2.6-9.9	1.0	1.7	0.6-4.7
	2011-2015	14.4	15.5	12.0-19.8	9.6	21.8	15.9-29.4	4.8	9.5	6.0-14.6	3.2	3.3	1.9-5.6	2.0	4.8	2.2-9.2	1.2	2.0	0.7-5.1
	2016-2020	14.8	15.1	11.7-19.3	10.2	22.0	16.2-29.6	4.6	8.8	5.4-13.9	3.2	3.3	1.8-5.6	2.4	4.8	2.4-9.0	0.8	1.8	0.4-5.2
Esophagus	1976-1980	1.6	2.8	1.2-5.7	1.4	6.2	2.4-13.0	0.2	0.6	0.0-3.7	1.2	2.2	0.8-4.9	0.8	3.9	1.0-9.9	0.4	1.4	0.2-5.1
	1981-1985	2.8	4.7	2.6-8.0	1.2	4.4	1.6-9.9	1.6	4.4	1.9-9.0	2.4	4.2	2.1-7.4	1.0	3.5	1.1-8.5	1.4	3.9	1.6-8.4
	1986-1990	4.0	6.4	3.9-10.0	2.2	7.9	3.9-14.3	1.8	4.4	2.0-8.7	2.8	4.4	2.4-7.4	1.8	6.6	3.0-12.7	1.0	2.6	0.8-6.3
	1991-1995	3.0	4.5	2.5-7.5	1.6	5.6	2.4-11.0	1.4	3.3	1.3-7.1	2.6	4.1	2.2-7.0	1.6	5.8	2.5-11.4	1.0	2.2	0.7-5.6
	1996-2000	2.6	3.6	1.9-6.2	2.0	6.0	2.9-11.2	0.6	1.3	0.3-4.3	2.6	3.6	1.9-6.3	1.4	4.4	1.8-9.2	1.2	2.7	1.0-6.3
	2001-2005	4.2	5.3	3.3-8.2	3.2	9.7	5.5-16.0	1.0	2.1	0.7-5.4	4.0	5.0	3.1-7.9	3.4	9.8	5.7-15.9	0.6	1.2	0.3-4.1
	2006-2010	5.6	6.5	4.3-9.5	4.8	12.0	7.6-18.2	0.8	1.7	0.4-4.8	4.4	5.2	3.2-8.0	4.0	10.4	6.3-16.3	0.4	0.9	0.1-3.7
	2011-2015	7.0	6.9	4.8-9.8	6.0	12.9	8.6-18.9	1.0	1.8	0.6-4.9	5.4	5.6	3.6-8.3	3.8	8.7	5.2-14.0	1.6	2.8	1.2-6.1
	2016-2020	8.4	7.9	5.7-11.0	6.8	13.6	9.3-19.5	1.6	3.2	1.4-7.0	5.8	5.4	3.6-8.1	4.2	8.3	5.1-13.3	1.6	3.3	1.4-7.0
Stomach	1976-1980	3.2	6.1	3.5-10.0	2.2	11.6	5.6-20.7	1.0	3.1	1.0-7.4	2.8	5.3	2.9-8.9	1.4	6.9	2.7-14.3	1.4	4.2	1.7-8.9
	1981-1985	4.8	8.8	5.6-13.1	2.6	11.6	6.0-20.1	2.2	6.6	3.2-12.1	2.6	4.5	2.4-7.8	0.8	3.5	0.9-9.0	1.8	5.2	2.3-10.2
	1986-1990	3.8	6.3	3.8-9.9	2.4	9.9	5.0-17.4	1.4	3.9	1.5-8.3	3.8	6.5	3.9-10.2	1.6	7.3	3.1-14.4	2.2	5.9	2.9-10.9
	1991-1995	4.4	6.5	4.1-9.9	2.2	7.6	3.7-13.8	2.2	5.3	2.6-9.8	3.2	4.7	2.7-7.7	1.6	5.5	2.3-11.0	1.6	3.7	1.6-7.6
	1996-2000	6.6	9.3	6.4-13.1	4.8	15.3	9.7-22.8	1.8	4.4	2.0-8.6	3.8	5.2	3.2-8.3	3.2	11.0	6.2-17.9	0.6	1.3	0.3-4.3
	2001-2005	5.4	6.8	4.5-10.0	3.8	10.9	6.6-17.2	1.6	3.7	1.6-7.6	2.2	2.7	1.4-5.0	1.6	4.7	2.0-9.4	0.6	1.3	0.3-4.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Warren**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	5.8	6.6	4.4-9.7	4.2	9.8	6.0-15.4	1.6	3.6	1.5-7.5	2.2	2.5	1.2-4.7	1.4	3.4	1.4-7.4	0.8	1.9	0.5-5.3
	2011-2015	7.0	7.1	4.9-10.1	4.8	11.1	7.0-16.9	2.2	3.8	1.9-7.5	2.6	2.6	1.4-4.8	2.2	5.0	2.4-9.3	0.4	0.9	0.1-3.8
	2016-2020	4.0	3.9	2.3-6.3	3.0	6.6	3.6-11.4	1.0	1.4	0.5-4.3	2.2	2.4	1.1-4.6	1.0	2.7	0.8-6.8	1.2	1.9	0.7-5.0
Colorectal	1976-1980	35.0	65.2	55.8-75.8	16.4	73.8	58.1-92.4	18.6	58.6	47.2-72.0	22.4	42.1	34.5-50.8	10.8	51.2	37.7-67.7	11.6	36.7	27.8-47.8
	1981-1985	49.0	86.0	75.4-97.6	20.6	90.8	73.6-110.7	28.4	83.8	70.4-99.2	18.2	32.8	26.3-40.4	6.8	29.6	20.2-41.7	11.4	34.7	26.1-45.4
	1986-1990	42.2	70.4	61.1-80.7	18.6	77.7	62.3-95.6	23.6	66.2	54.5-79.8	19.0	31.6	25.5-38.8	8.2	35.7	25.4-48.7	10.8	28.7	21.4-37.8
	1991-1995	42.4	63.4	55.1-72.6	22.4	83.9	68.7-101.4	20.0	50.8	41.2-62.2	16.4	24.5	19.5-30.4	6.6	25.9	17.6-36.6	9.8	23.1	17.0-30.8
	1996-2000	46.2	64.4	56.3-73.3	20.2	69.4	56.2-84.6	26.0	62.0	51.7-73.9	15.8	22.0	17.4-27.4	7.0	24.6	17.0-34.4	8.8	19.8	14.4-26.9
	2001-2005	42.6	54.4	47.3-62.3	21.0	62.1	50.6-75.4	21.6	47.5	38.9-57.6	14.6	18.8	14.7-23.7	7.2	22.3	15.5-31.1	7.4	15.7	11.0-21.9
	2006-2010	35.2	39.4	33.8-45.9	18.0	44.7	35.8-55.3	17.2	34.9	27.8-43.5	16.0	18.7	14.8-23.4	8.8	25.0	18.1-33.7	7.2	13.9	9.7-19.7
	2011-2015	38.2	43.8	37.5-50.9	17.2	42.9	33.9-53.6	21.0	43.9	35.4-54.0	12.4	12.4	9.5-16.2	5.8	13.6	9.1-19.9	6.6	11.5	7.8-16.7
2016-2020	32.4	34.5	29.0-40.8	16.8	39.4	31.0-49.6	15.6	30.1	23.3-38.5	13.0	13.1	10.0-17.1	7.0	16.4	11.2-23.6	6.0	9.6	6.4-14.4	
Colon excluding rectum	1976-1980	25.0	46.8	38.8-55.9	10.6	46.3	34.3-61.0	14.4	45.8	35.7-57.9	17.4	32.5	25.9-40.2	8.2	37.5	26.3-51.6	9.2	29.5	21.5-39.6
	1981-1985	34.4	60.9	52.0-70.9	12.6	55.0	41.9-70.9	21.8	64.8	53.0-78.6	16.2	29.3	23.2-36.5	5.6	24.6	16.1-35.8	10.6	32.0	23.8-42.1
	1986-1990	32.0	52.9	44.9-61.9	12.8	53.8	41.1-69.1	19.2	52.8	42.5-64.9	16.8	27.7	22.1-34.4	7.2	30.5	21.1-42.5	9.6	25.6	18.7-34.2
	1991-1995	27.8	41.4	34.7-48.9	14.0	53.3	41.2-67.7	13.8	34.1	26.4-43.4	14.4	21.5	16.8-27.1	5.6	22.2	14.6-32.3	8.8	20.6	14.9-28.0
	1996-2000	35.4	49.3	42.2-57.1	13.8	48.2	37.3-61.3	21.6	51.3	42.0-62.3	11.8	16.4	12.5-21.2	5.0	17.9	11.5-26.6	6.8	15.3	10.6-21.6
	2001-2005	31.8	40.7	34.6-47.7	15.2	45.5	35.8-57.2	16.6	36.3	28.9-45.3	10.8	13.9	10.5-18.3	5.2	16.8	10.9-24.8	5.6	11.6	7.7-17.1
	2006-2010	24.0	27.0	22.4-32.5	11.0	27.5	20.6-36.2	13.0	26.5	20.4-34.1	11.6	13.5	10.2-17.6	6.4	18.1	12.3-25.7	5.2	10.1	6.6-15.3
	2011-2015	25.4	28.5	23.6-34.3	10.0	25.2	18.4-33.7	15.4	31.1	24.2-39.6	9.2	9.1	6.6-12.3	4.4	10.3	6.4-16.0	4.8	8.1	5.1-12.6
2016-2020	23.8	24.8	20.2-30.1	12.4	28.8	21.7-37.7	11.4	21.1	15.6-28.2	8.8	9.1	6.5-12.6	5.0	12.1	7.7-18.5	3.8	6.1	3.6-10.3	
Rectum & rectosigmoid	1976-1980	10.0	18.4	13.6-24.4	5.8	27.6	18.0-40.1	4.2	12.8	7.9-19.7	5.0	9.6	6.2-14.3	2.6	13.7	7.0-23.7	2.4	7.2	3.7-12.9
	1981-1985	14.6	25.1	19.6-31.7	8.0	35.8	25.3-49.0	6.6	19.0	13.0-26.9	2.0	3.6	1.7-6.6	1.2	5.0	1.8-11.1	0.8	2.8	0.7-7.3
	1986-1990	10.2	17.5	13.0-23.1	5.8	23.9	15.8-34.5	4.4	13.4	8.3-20.6	2.2	3.9	1.9-7.0	1.0	5.2	1.7-11.9	1.2	3.1	1.1-7.2
	1991-1995	14.6	22.0	17.2-27.8	8.4	30.6	21.9-41.6	6.2	16.7	11.3-24.0	2.0	3.0	1.4-5.5	1.0	3.7	1.2-8.6	1.0	2.4	0.8-6.0
	1996-2000	10.8	15.2	11.4-19.8	6.4	21.2	14.4-30.1	4.4	10.6	6.6-16.4	4.0	5.6	3.4-8.7	2.0	6.7	3.2-12.3	2.0	4.5	2.2-8.7
	2001-2005	10.8	13.7	10.3-18.0	5.8	16.5	11.0-24.0	5.0	11.1	7.2-16.7	3.8	4.8	2.9-7.6	2.0	5.5	2.6-10.4	1.8	4.1	1.8-8.0
	2006-2010	11.2	12.4	9.3-16.3	7.0	17.1	11.8-24.2	4.2	8.4	5.1-13.3	4.4	5.2	3.2-8.0	2.4	6.9	3.6-12.2	2.0	3.8	1.8-7.5
	2011-2015	12.8	15.2	11.5-19.8	7.2	17.7	12.1-25.1	5.6	12.8	8.2-19.2	3.2	3.3	1.9-5.7	1.4	3.3	1.3-7.2	1.8	3.5	1.5-7.2
2016-2020	8.6	9.7	6.9-13.5	4.4	10.6	6.4-16.7	4.2	8.9	5.2-14.5	4.2	4.0	2.4-6.5	2.0	4.3	1.9-8.7	2.2	3.5	1.7-7.0	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Warren**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	1.4	2.6	1.0-5.4	1.2	5.5	2.0-11.9	0.2	0.5	0.0-3.6	0.8	1.7	0.5-4.2	0.8	4.0	1.1-10.0	0.0	0.0	0.0-2.7
	1981-1985	2.4	4.3	2.2-7.6	1.8	8.0	3.5-15.3	0.6	1.8	0.4-5.4	2.4	4.5	2.3-8.0	1.8	8.7	3.8-16.6	0.6	2.2	0.4-6.5
	1986-1990	1.4	2.3	0.9-4.8	1.2	5.6	2.0-12.1	0.2	0.5	0.0-3.2	1.6	2.8	1.2-5.5	1.4	7.0	2.7-14.3	0.2	0.5	0.0-3.2
	1991-1995	4.0	6.0	3.6-9.3	1.8	6.2	2.8-11.9	2.2	5.6	2.8-10.3	3.2	4.6	2.6-7.5	1.8	7.0	3.1-13.4	1.4	3.2	1.3-6.9
	1996-2000	3.4	4.8	2.8-7.7	2.4	8.4	4.3-14.8	1.0	2.3	0.7-5.7	2.4	3.4	1.7-6.0	1.4	4.8	1.9-10.0	1.0	2.4	0.8-5.9
	2001-2005	2.2	2.8	1.4-5.1	1.8	5.3	2.4-10.2	0.4	0.9	0.1-3.7	2.2	2.8	1.4-5.1	1.4	4.3	1.7-9.0	0.8	1.8	0.5-5.0
	2006-2010	4.8	5.5	3.5-8.4	3.4	7.8	4.5-12.9	1.4	3.4	1.3-7.5	3.8	4.4	2.6-7.0	2.4	6.2	3.1-11.0	1.4	2.9	1.2-6.5
	2011-2015	5.8	5.6	3.7-8.3	4.2	8.5	5.2-13.5	1.6	2.9	1.2-6.3	4.4	4.3	2.7-6.8	3.2	7.1	4.0-11.9	1.2	2.0	0.7-5.0
	2016-2020	8.0	7.5	5.3-10.5	5.6	11.1	7.2-16.7	2.4	4.2	2.1-7.9	6.6	5.9	4.0-8.6	4.6	8.8	5.5-13.8	2.0	3.5	1.7-7.1
Pancreas	1976-1980	5.8	11.1	7.4-16.0	2.4	11.7	5.7-20.8	3.4	10.9	6.3-17.6	7.0	13.1	9.1-18.3	3.0	15.2	8.1-25.6	4.0	12.3	7.5-19.2
	1981-1985	6.6	11.9	8.1-16.8	4.4	19.0	11.6-29.0	2.2	6.3	3.1-11.5	5.6	9.8	6.5-14.2	3.6	15.3	8.8-24.5	2.0	5.8	2.8-10.9
	1986-1990	8.6	13.9	10.1-18.9	4.2	17.4	10.5-27.0	4.4	11.9	7.4-18.3	7.4	12.0	8.4-16.6	3.4	13.6	7.7-22.1	4.0	10.8	6.5-17.0
	1991-1995	6.4	9.6	6.6-13.6	4.0	15.8	9.4-24.5	2.4	6.2	3.2-11.0	6.4	9.4	6.5-13.4	3.2	13.0	7.3-21.3	3.2	8.0	4.5-13.1
	1996-2000	9.4	13.2	9.7-17.6	3.6	11.8	6.9-18.8	5.8	14.4	9.6-20.9	6.0	8.3	5.6-11.9	2.6	8.7	4.6-14.9	3.4	8.0	4.6-13.1
	2001-2005	7.0	9.1	6.3-12.7	2.8	8.2	4.5-14.0	4.2	9.4	5.8-14.7	8.8	11.2	8.1-15.1	4.2	11.9	7.3-18.5	4.6	10.1	6.3-15.4
	2006-2010	9.8	10.6	7.8-14.2	5.8	14.5	9.6-21.2	4.0	8.0	4.8-12.8	9.4	10.4	7.6-14.0	5.0	12.6	8.1-19.0	4.4	8.9	5.5-13.8
	2011-2015	13.6	13.8	10.7-17.7	8.0	18.2	12.9-25.2	5.6	10.7	7.0-16.0	11.4	11.7	8.8-15.4	6.4	15.2	10.4-21.9	5.0	9.4	6.0-14.3
	2016-2020	12.6	11.3	8.6-14.8	6.0	11.3	7.5-16.7	6.6	11.3	7.7-16.5	10.4	9.5	7.0-12.7	4.6	8.9	5.6-13.9	5.8	9.6	6.4-14.4
Larynx	1976-1980	2.0	3.8	1.8-7.0	1.8	8.3	3.6-16.2	0.2	0.7	0.0-3.9	0.6	1.1	0.2-3.2	0.4	1.5	0.2-5.9	0.2	0.6	0.0-3.7
	1981-1985	2.2	3.6	1.8-6.6	1.8	6.8	3.0-13.2	0.4	1.4	0.2-5.2	1.2	2.0	0.7-4.5	1.0	4.0	1.3-9.6	0.2	0.6	0.0-3.8
	1986-1990	4.2	7.1	4.4-11.0	3.4	13.8	7.9-22.3	0.8	3.0	0.8-7.6	1.4	2.4	0.9-5.0	1.0	3.7	1.2-8.9	0.4	1.3	0.1-4.8
	1991-1995	4.4	6.8	4.2-10.3	3.4	11.6	6.7-18.8	1.0	3.1	1.0-7.3	0.6	0.8	0.2-2.5	0.4	1.7	0.2-6.1	0.2	0.5	0.0-3.2
	1996-2000	4.2	5.9	3.7-9.1	4.0	12.1	7.4-18.9	0.2	0.6	0.0-3.3	0.8	1.1	0.3-3.0	0.6	1.8	0.4-5.5	0.2	0.6	0.0-3.3
	2001-2005	2.2	2.7	1.4-5.0	1.6	4.1	1.8-8.3	0.6	1.4	0.3-4.5	1.8	2.1	1.0-4.2	1.4	3.6	1.4-7.7	0.4	0.9	0.1-3.7
	2006-2010	5.6	6.7	4.4-9.9	4.8	12.2	7.7-18.4	0.8	1.8	0.5-5.1	1.2	1.5	0.6-3.5	0.6	1.7	0.4-5.2	0.6	1.4	0.3-4.5
	2011-2015	6.0	6.5	4.3-9.5	4.6	10.9	6.8-16.7	1.4	2.6	1.0-6.0	1.2	1.2	0.4-3.0	0.6	1.4	0.3-4.6	0.6	1.0	0.2-3.8
	2016-2020	2.6	2.6	1.3-4.8	2.0	3.6	1.7-7.3	0.6	1.6	0.2-5.3	1.4	1.5	0.6-3.4	1.4	3.1	1.2-7.1	0.0	0.0	0.0-2.6
Lung and bronchus	1976-1980	42.2	77.4	67.1-88.9	28.6	126.0	105.4-149.5	13.6	44.7	34.5-57.0	30.2	55.7	47.0-65.6	19.8	88.4	71.2-108.5	10.4	34.7	25.7-45.9
	1981-1985	40.4	71.0	61.3-81.7	25.8	108.2	89.9-129.2	14.6	44.6	34.7-56.5	36.2	62.6	53.6-72.6	24.0	101.8	83.8-122.3	12.2	35.6	27.1-46.1
	1986-1990	45.0	75.3	65.6-86.0	28.4	113.9	95.3-134.9	16.6	48.1	38.1-60.1	34.4	57.0	48.7-66.3	20.6	82.9	67.1-101.1	13.8	39.5	30.6-50.3
	1991-1995	50.8	75.6	66.5-85.6	26.6	97.3	81.1-115.7	24.2	61.6	51.0-73.9	36.8	54.5	46.9-63.0	20.8	77.1	62.7-93.7	16.0	39.6	31.3-49.5
	1996-2000	56.2	78.3	69.4-88.1	31.6	100.3	85.1-117.4	24.6	61.4	51.0-73.5	45.0	62.5	54.6-71.3	26.0	84.5	70.4-100.6	19.0	47.1	38.0-57.8
	2001-2005	62.4	79.8	71.1-89.2	31.2	91.7	77.6-107.5	31.2	72.1	61.2-84.6	43.0	54.8	47.7-62.7	23.4	68.7	56.7-82.7	19.6	44.7	36.3-54.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Warren**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	67.8	77.5	69.3-86.4	33.8	87.3	74.4-102.0	34.0	70.8	60.4-82.7	45.2	51.9	45.3-59.3	23.4	61.2	50.4-73.7	21.8	45.2	37.0-54.9
	2011-2015	76.8	79.3	71.4-88.0	41.0	92.5	80.0-106.7	35.8	68.3	58.4-79.8	43.6	45.3	39.4-52.1	22.6	50.6	41.5-61.3	21.0	40.6	32.9-49.8
	2016-2020	76.4	70.0	63.0-77.7	39.2	76.3	65.7-88.3	37.2	64.5	55.4-75.1	45.8	42.8	37.3-49.0	24.6	50.1	41.4-60.3	21.2	36.6	29.8-44.8
Melanoma of the skin	1976-1980	3.2	6.5	3.6-10.6	1.8	7.5	3.3-14.6	1.4	5.8	2.2-11.8	1.2	3.1	1.1-6.5	0.8	4.2	1.1-10.7	0.4	2.0	0.2-6.7
	1981-1985	7.0	12.7	8.8-17.8	4.6	18.6	11.6-28.2	2.4	7.9	4.0-14.1	0.6	1.0	0.2-3.1	0.4	2.4	0.2-8.0	0.2	0.6	0.0-3.6
	1986-1990	6.4	10.9	7.4-15.5	3.6	13.5	7.9-21.5	2.8	8.6	4.6-14.7	1.8	2.9	1.3-5.5	1.4	5.3	2.0-11.1	0.4	1.2	0.1-4.6
	1991-1995	7.0	11.3	7.8-15.7	4.6	17.0	10.6-25.6	2.4	7.1	3.6-12.6	2.2	3.4	1.7-6.0	1.6	5.6	2.4-11.1	0.6	1.7	0.3-5.0
	1996-2000	10.8	15.8	11.9-20.7	5.6	18.2	12.0-26.4	5.2	14.4	9.3-21.2	2.2	3.2	1.6-5.7	1.4	4.4	1.8-9.2	0.8	2.1	0.6-5.6
	2001-2005	20.0	27.2	22.1-33.2	10.2	29.6	22.0-39.2	9.8	25.2	18.5-33.6	2.8	3.8	2.1-6.4	1.0	2.8	0.9-6.8	1.8	4.7	2.1-9.1
	2006-2010	23.4	28.7	23.6-34.7	13.8	35.7	27.6-45.6	9.6	23.8	17.3-32.1	2.8	3.0	1.7-5.3	1.8	4.5	2.0-8.9	1.0	2.0	0.6-5.1
	2011-2015	23.4	26.2	21.5-31.7	14.6	34.1	26.6-43.3	8.8	20.1	14.3-27.6	2.8	3.0	1.6-5.3	1.8	3.9	1.7-8.0	1.0	2.1	0.6-5.4
	2016-2020	28.0	32.9	27.2-39.4	17.4	41.0	32.4-51.3	10.6	27.5	20.0-36.9	3.8	3.8	2.3-6.2	3.2	7.3	4.1-12.3	0.6	1.1	0.2-4.0
Female breast	1976-1980							32.0	109.8	93.0-128.7							8.6	28.5	20.5-38.7
	1981-1985							34.4	114.3	97.3-133.5							15.4	50.5	39.5-63.5
	1986-1990							42.0	123.8	107.2-142.4							12.4	35.8	27.2-46.2
	1991-1995							46.4	132.6	115.7-151.2							14.6	40.0	31.1-50.7
	1996-2000							50.4	131.4	115.5-148.9							13.4	33.4	25.8-42.7
	2001-2005							53.0	127.8	112.7-144.5							13.8	32.2	24.9-41.1
	2006-2010							59.6	133.2	118.1-149.8							9.8	20.1	14.8-27.0
	2011-2015							68.2	142.4	126.9-159.4							10.6	19.3	14.4-25.7
	2016-2020							74.6	146.0	130.5-163.1							8.8	15.6	11.1-21.7
Cervix uteri	1976-1980							4.0	15.2	9.2-23.7							2.0	7.4	3.5-13.8
	1981-1985							4.6	15.9	10.0-24.0							2.4	9.2	4.7-16.2
	1986-1990							3.8	12.8	7.6-20.1							1.4	4.8	1.9-10.0
	1991-1995							2.6	7.9	4.1-13.6							1.0	3.1	1.0-7.3
	1996-2000							4.2	11.7	7.2-18.0							1.6	4.0	1.7-8.1
	2001-2005							3.0	7.6	4.2-12.8							1.2	2.7	1.0-6.2
	2006-2010							3.4	7.6	4.3-12.5							1.0	2.3	0.7-5.7
	2011-2015							1.8	6.0	2.6-11.7							1.2	2.9	0.9-6.9
	2016-2020							1.2	2.4	0.8-5.8							0.2	1.0	0.0-4.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Warren**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							7.6	25.4	17.8-35.2							1.2	3.6	1.3-8.0
	1981-1985							7.4	24.6	17.1-34.2							1.4	4.0	1.6-8.5
	1986-1990							9.4	30.2	22.0-40.6							2.0	5.8	2.7-11.0
	1991-1995							6.2	17.8	12.0-25.5							0.8	2.0	0.5-5.4
	1996-2000							8.2	20.8	14.9-28.5							2.0	4.5	2.2-8.7
	2001-2005							12.2	28.9	22.1-37.4							2.6	6.1	3.2-10.7
	2006-2010							18.8	39.0	31.4-48.1							3.6	7.6	4.4-12.4
	2011-2015							12.6	25.2	19.1-32.8							2.8	5.6	3.0-9.9
	2016-2020							15.4	30.0	23.2-38.6							2.6	3.8	2.0-7.3
Ovary	1976-1980							4.8	16.7	10.6-25.0							2.0	6.3	3.0-11.8
	1981-1985							5.8	18.6	12.3-27.0							3.4	10.0	5.8-16.3
	1986-1990							6.8	20.7	14.2-29.3							2.8	7.5	4.1-12.9
	1991-1995							4.6	12.2	7.7-18.5							4.6	12.4	7.8-18.9
	1996-2000							5.4	15.6	10.2-22.8							2.0	5.0	2.4-9.5
	2001-2005							5.6	13.9	9.2-20.3							4.2	9.6	5.9-14.9
	2006-2010							7.8	18.2	12.8-25.3							4.2	8.6	5.3-13.6
	2011-2015							4.8	11.0	6.8-17.1							3.0	5.8	3.2-10.0
	2016-2020							4.6	9.4	5.6-14.9							3.6	7.4	4.1-12.6
Prostate	1976-1980				17.6	91.6	72.5-113.9							5.2	29.2	18.5-43.3			
	1981-1985				22.8	109.5	89.5-132.3							7.4	40.6	28.1-56.4			
	1986-1990				27.8	122.1	101.8-145.0							10.0	48.7	35.7-64.6			
	1991-1995				38.4	136.2	117.2-157.4							7.0	30.4	21.0-42.5			
	1996-2000				45.2	142.8	124.6-163.0							8.2	29.7	21.1-40.5			
	2001-2005				63.0	178.5	159.1-199.8							9.0	30.0	21.8-40.3			
	2006-2010				68.2	167.5	149.9-186.7							6.4	18.5	12.6-26.3			
	2011-2015				59.6	127.8	113.4-143.7							9.2	23.3	17.0-31.4			
	2016-2020				71.4	134.6	120.7-150.1							8.2	17.5	12.5-24.2			
Testis	1976-1980				0.8	3.9	1.0-10.1							0.0	0.0	0.0-3.4			
	1981-1985				1.0	3.7	1.1-8.9							0.0	0.0	0.0-3.2			
	1986-1990				1.0	3.6	1.2-8.6							0.2	0.8	0.0-4.4			
	1991-1995				1.2	4.0	1.5-8.8							0.0	0.0	0.0-2.7			
	1996-2000				3.2	10.8	6.2-17.6							0.2	0.7	0.0-3.8			
	2001-2005				2.0	6.5	3.1-12.0							0.2	0.5	0.0-3.3			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Warren**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				1.4	5.1	2.0-10.5							0.2	0.5	0.0-3.2			
	2011-2015				2.4	7.9	4.0-14.1							0.6	1.4	0.3-4.6			
	2016-2020				2.0	7.0	3.2-13.1							0.0	0.0	0.0-2.6			
Urinary bladder (incl. in situ)	1976-1980	12.4	23.3	17.8-30.0	9.2	44.9	32.4-60.4	3.2	10.0	5.7-16.4	4.6	9.2	5.8-13.8	3.4	18.5	10.4-29.9	1.2	3.8	1.4-8.5
	1981-1985	13.6	24.2	18.7-30.8	9.2	42.7	30.8-57.5	4.4	13.2	8.2-20.3	3.2	5.8	3.3-9.5	2.0	10.3	4.8-19.1	1.2	3.4	1.2-7.6
	1986-1990	14.8	24.4	19.1-30.7	11.0	46.3	34.5-60.7	3.8	10.2	6.0-16.2	4.0	6.6	4.0-10.3	3.0	14.2	7.7-23.4	1.0	2.3	0.8-5.9
	1991-1995	18.2	27.1	21.8-33.4	12.4	46.6	35.4-60.2	5.8	14.9	9.9-21.6	2.8	4.1	2.2-6.9	1.4	5.0	1.9-10.5	1.4	3.1	1.2-6.7
	1996-2000	17.4	23.9	19.1-29.5	12.6	41.3	31.6-53.1	4.8	11.1	7.1-16.8	3.6	5.0	2.9-7.9	2.0	7.1	3.4-13.2	1.6	3.5	1.5-7.3
	2001-2005	25.0	32.0	26.6-38.2	19.8	59.6	48.3-72.9	5.2	11.8	7.6-17.5	3.4	4.4	2.5-7.1	1.6	4.9	2.1-9.7	1.8	3.8	1.7-7.6
	2006-2010	25.0	28.5	23.6-34.1	17.2	44.2	35.2-54.9	7.8	15.3	10.9-21.4	4.0	4.4	2.7-7.0	3.4	9.1	5.3-14.9	0.6	1.3	0.3-4.2
	2011-2015	28.0	28.4	23.8-33.7	22.0	49.3	40.4-59.9	6.0	11.6	7.6-17.1	6.8	7.3	5.0-10.4	4.4	10.5	6.5-16.2	2.4	5.2	2.5-9.7
2016-2020	26.8	24.6	20.5-29.4	19.4	39.9	32.2-49.3	7.4	11.6	8.1-16.6	5.4	5.1	3.3-7.7	4.2	9.3	5.7-14.7	1.2	1.7	0.6-4.6	
Kidney and renal pelvis	1976-1980	4.0	7.2	4.4-11.2	2.2	10.1	4.8-18.5	1.8	5.5	2.5-10.6	2.2	4.3	2.1-7.7	0.8	2.9	0.8-7.9	1.4	5.0	2.0-10.4
	1981-1985	6.6	11.8	8.0-16.6	3.6	15.9	9.2-25.4	3.0	9.2	5.1-15.5	3.4	5.9	3.4-9.5	1.6	8.0	3.3-15.7	1.8	4.9	2.2-9.6
	1986-1990	8.4	13.6	9.7-18.5	4.4	17.0	10.5-26.0	4.0	11.0	6.6-17.3	3.6	5.8	3.4-9.3	2.2	9.3	4.5-16.9	1.4	4.0	1.5-8.5
	1991-1995	6.6	9.8	6.7-13.8	3.0	10.3	5.7-17.2	3.6	9.1	5.3-14.6	3.4	5.3	3.1-8.5	1.4	5.1	2.0-10.5	2.0	5.2	2.4-9.8
	1996-2000	8.8	12.3	8.9-16.5	4.2	14.6	8.9-22.5	4.6	11.2	7.0-17.0	3.6	4.9	2.9-7.8	2.0	7.0	3.3-12.9	1.6	3.5	1.5-7.2
	2001-2005	12.2	15.9	12.1-20.5	7.6	21.5	15.2-29.7	4.6	11.0	6.9-16.9	3.0	3.8	2.1-6.4	1.4	4.4	1.7-9.2	1.6	3.6	1.6-7.5
	2006-2010	15.2	17.3	13.6-21.9	8.4	20.5	14.6-28.1	6.8	15.0	10.3-21.4	3.8	4.5	2.7-7.2	2.0	5.1	2.4-9.8	1.8	4.0	1.7-8.0
	2011-2015	18.6	20.6	16.4-25.6	11.0	26.2	19.4-34.7	7.6	14.8	10.2-21.0	6.2	6.3	4.2-9.2	3.6	8.4	4.9-13.7	2.6	4.7	2.5-8.6
2016-2020	16.2	16.4	12.8-20.9	10.0	21.7	15.7-29.5	6.2	11.6	7.7-17.3	3.8	3.6	2.1-5.9	2.2	4.5	2.2-8.7	1.6	2.8	1.2-6.2	
Brain and other central nervous system	1976-1980	1.6	3.0	1.3-5.9	0.6	2.8	0.5-8.3	1.0	3.0	1.0-7.3	1.8	3.4	1.5-6.5	0.8	3.6	0.9-9.3	1.0	3.4	1.1-8.1
	1981-1985	3.6	6.7	3.9-10.7	2.0	8.3	3.9-15.5	1.6	5.4	2.3-10.9	2.2	4.1	2.0-7.3	1.4	7.1	2.7-14.6	0.8	2.2	0.6-5.9
	1986-1990	5.4	9.3	6.1-13.6	3.0	11.1	6.1-18.6	2.4	7.6	3.9-13.6	3.0	5.3	2.9-8.8	1.8	7.8	3.5-14.7	1.2	3.2	1.2-7.3
	1991-1995	3.4	5.3	3.1-8.5	2.2	7.6	3.8-13.7	1.2	3.6	1.3-8.0	3.0	4.8	2.7-7.9	2.6	8.9	4.7-15.3	0.4	1.3	0.2-4.6
	1996-2000	4.8	7.0	4.5-10.5	2.8	8.7	4.8-14.7	2.0	5.3	2.5-10.0	3.2	4.8	2.7-7.8	1.4	4.6	1.8-9.5	1.8	4.7	2.1-9.1
	2001-2005	5.2	7.3	4.7-10.9	3.6	11.0	6.5-17.6	1.6	4.2	1.8-8.5	3.8	5.0	3.0-7.9	2.6	7.7	4.1-13.4	1.2	2.9	1.1-6.6
	2006-2010	5.6	7.8	5.1-11.5	3.0	8.9	4.9-14.9	2.6	6.7	3.5-11.7	4.0	5.3	3.2-8.4	2.2	6.3	3.1-11.5	1.8	4.4	2.0-8.7
	2011-2015	8.4	11.3	7.9-15.6	3.8	10.8	6.2-17.3	4.6	12.0	7.3-18.7	4.8	5.6	3.5-8.7	2.8	7.1	3.7-12.5	2.0	4.5	2.0-8.9
2016-2020	6.6	8.7	5.8-12.6	3.6	10.0	5.7-16.4	3.0	7.8	4.1-13.6	5.0	5.3	3.3-8.1	3.4	7.9	4.5-13.3	1.6	3.1	1.2-6.9	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Warren**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	0.8	1.5	0.4-4.0	0.2	0.9	0.0-5.1	0.6	1.9	0.4-5.7	0.2	0.4	0.0-2.2	0.0	0.0	0.0-3.4	0.2	0.7	0.0-3.9
	1981-1985	2.0	3.3	1.6-6.2	0.4	1.4	0.2-5.4	1.6	5.1	2.2-10.1	0.2	0.4	0.0-2.1	0.0	0.0	0.0-3.2	0.2	0.6	0.0-3.6
	1986-1990	1.2	2.2	0.8-4.8	0.6	2.3	0.5-6.9	0.6	1.9	0.3-5.7	0.2	0.3	0.0-1.7	0.0	0.0	0.0-3.0	0.2	0.5	0.0-3.2
	1991-1995	2.8	4.4	2.4-7.4	0.2	1.2	0.0-5.4	2.6	7.8	4.1-13.4	0.6	0.9	0.2-2.7	0.2	1.2	0.0-5.4	0.4	0.9	0.1-3.6
	1996-2000	3.8	6.0	3.6-9.4	0.8	2.5	0.7-6.6	3.0	9.4	5.2-15.5	0.4	0.6	0.1-2.2	0.2	0.6	0.0-3.7	0.2	0.5	0.0-3.1
	2001-2005	5.0	7.4	4.7-11.0	0.6	1.8	0.4-5.5	4.4	12.5	7.7-19.1	0.0	0.0	0.0-1.2	0.0	0.0	0.0-2.5	0.0	0.0	0.0-2.3
	2006-2010	12.2	16.8	12.7-21.8	3.0	7.9	4.3-13.3	9.2	25.5	18.4-34.5	0.8	0.9	0.3-2.6	0.4	0.9	0.1-3.8	0.4	0.8	0.1-3.5
	2011-2015	18.6	25.5	20.3-31.7	4.2	11.0	6.6-17.3	14.4	39.2	30.1-50.3	0.4	0.4	0.0-1.7	0.0	0.0	0.0-2.5	0.4	0.5	0.1-3.1
2016-2020	14.2	19.7	15.1-25.3	3.4	9.1	5.0-15.2	10.8	30.1	22.0-40.1	0.4	0.3	0.0-1.7	0.2	0.5	0.0-3.3	0.2	0.3	0.0-2.9	
Hodgkin lymphoma	1976-1980	0.8	1.4	0.4-3.8	0.6	2.4	0.5-7.3	0.2	0.6	0.0-3.8	0.4	0.8	0.1-2.8	0.2	0.8	0.0-4.7	0.2	0.7	0.0-3.9
	1981-1985	2.0	3.4	1.6-6.3	1.2	4.7	1.5-10.9	0.8	2.7	0.7-7.1	0.0	0.0	0.0-1.4	0.0	0.0	0.0-3.2	0.0	0.0	0.0-2.6
	1986-1990	2.6	4.1	2.1-7.0	1.4	4.6	1.8-9.8	1.2	3.6	1.3-8.1	0.0	0.0	0.0-1.3	0.0	0.0	0.0-3.0	0.0	0.0	0.0-2.5
	1991-1995	2.0	3.3	1.6-6.0	1.0	3.3	1.1-7.8	1.0	3.2	1.0-7.5	0.4	0.6	0.1-2.1	0.2	0.6	0.0-3.8	0.2	0.5	0.0-3.2
	1996-2000	1.2	2.0	0.7-4.4	0.6	1.9	0.4-5.8	0.6	2.4	0.5-6.6	0.0	0.0	0.0-1.2	0.0	0.0	0.0-2.6	0.0	0.0	0.0-2.3
	2001-2005	2.8	4.2	2.2-7.1	1.2	3.7	1.3-8.2	1.6	4.5	1.9-9.2	0.0	0.0	0.0-1.2	0.0	0.0	0.0-2.5	0.0	0.0	0.0-2.3
	2006-2010	1.2	1.8	0.6-3.9	0.6	1.8	0.3-5.4	0.6	1.7	0.3-5.1	1.4	1.9	0.7-4.1	1.0	2.6	0.8-6.5	0.4	1.2	0.1-4.6
	2011-2015	1.4	2.1	0.8-4.4	0.6	1.8	0.3-5.6	0.8	2.4	0.6-6.3	0.4	0.4	0.0-1.8	0.2	0.5	0.0-3.4	0.2	0.4	0.0-3.1
2016-2020	1.6	2.2	0.9-4.5	1.0	2.9	0.9-7.1	0.6	1.5	0.2-5.2	0.6	0.8	0.1-2.6	0.2	0.3	0.0-3.0	0.4	1.2	0.1-4.8	
Non-Hodgkin lymphomas	1976-1980	7.8	16.1	11.4-22.2	4.0	18.7	11.2-29.2	3.8	13.3	7.9-21.0	4.4	8.6	5.3-13.1	1.8	7.7	3.4-14.9	2.6	8.6	4.5-14.8
	1981-1985	9.8	17.6	12.9-23.5	6.6	27.7	18.8-39.3	3.2	10.0	5.6-16.6	4.8	9.1	5.8-13.6	3.6	16.1	9.4-25.7	1.2	4.1	1.4-9.1
	1986-1990	9.8	15.9	11.7-21.1	5.8	20.8	13.9-30.1	4.0	12.1	7.3-18.9	4.2	6.9	4.2-10.6	2.2	8.0	4.0-14.6	2.0	5.6	2.6-10.5
	1991-1995	10.6	16.2	12.1-21.3	6.2	21.4	14.5-30.5	4.4	11.5	7.2-17.6	5.2	7.7	5.0-11.4	3.2	10.9	6.2-17.8	2.0	5.0	2.4-9.5
	1996-2000	14.2	20.2	15.8-25.5	7.6	24.6	17.4-34.0	6.6	16.4	11.3-23.3	6.2	8.7	5.9-12.4	3.2	10.1	5.7-16.5	3.0	7.6	4.2-12.7
	2001-2005	16.2	20.6	16.3-25.7	8.2	24.1	17.2-33.0	8.0	18.5	13.2-25.5	4.8	6.2	3.9-9.2	2.4	8.5	4.3-14.9	2.4	5.0	2.6-9.1
	2006-2010	21.2	24.7	20.1-30.1	12.0	31.9	24.2-41.4	9.2	19.3	14.0-26.2	6.2	7.1	4.8-10.2	3.8	10.2	6.1-16.3	2.4	4.4	2.3-8.2
	2011-2015	23.6	24.9	20.5-30.0	11.6	26.8	20.2-35.2	12.0	23.8	18.0-31.1	6.6	6.6	4.5-9.4	3.8	8.5	5.1-13.7	2.8	5.1	2.7-9.1
2016-2020	22.2	22.9	18.7-28.0	11.0	23.6	17.5-31.4	11.2	22.4	16.6-29.8	5.6	5.6	3.6-8.4	2.0	4.9	2.2-9.5	3.6	6.2	3.7-10.5	
Myeloma	1976-1980	1.8	3.9	1.7-7.3	1.2	5.1	1.8-11.3	0.6	2.9	0.6-8.1	1.6	3.0	1.3-6.0	0.8	3.3	0.9-8.5	0.8	3.1	0.8-7.9
	1981-1985	3.4	6.1	3.5-9.8	2.2	9.6	4.7-17.2	1.2	3.5	1.3-7.9	2.2	4.1	2.0-7.3	1.6	7.2	3.0-14.2	0.6	1.8	0.4-5.4
	1986-1990	3.0	5.3	2.9-8.8	2.2	10.0	4.9-17.9	0.8	2.3	0.6-6.1	1.2	1.9	0.7-4.3	0.8	3.2	0.8-8.2	0.4	1.1	0.1-4.3
	1991-1995	3.6	5.5	3.2-8.7	3.0	11.3	6.2-18.7	0.6	1.8	0.4-5.4	3.4	5.1	3.0-8.2	2.6	9.9	5.2-17.0	0.8	1.8	0.5-5.0
	1996-2000	3.6	5.0	3.0-8.0	2.2	7.8	3.8-14.1	1.4	3.7	1.5-7.9	3.8	5.2	3.1-8.2	3.2	11.9	6.7-19.5	0.6	1.3	0.3-4.3
	2001-2005	4.0	5.2	3.2-8.1	2.4	6.9	3.6-12.2	1.6	3.7	1.6-7.6	3.0	3.8	2.1-6.3	2.0	6.0	2.9-11.3	1.0	2.3	0.7-5.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Warren**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	5.4	6.0	3.9-8.9	3.2	8.5	4.8-14.1	2.2	4.2	2.1-8.0	1.4	1.6	0.6-3.5	0.8	2.4	0.7-6.3	0.6	1.3	0.3-4.2
	2011-2015	8.8	9.1	6.5-12.4	5.2	12.3	7.9-18.5	3.6	6.7	3.9-11.1	2.6	2.6	1.4-4.7	2.0	5.1	2.4-9.7	0.6	1.0	0.2-3.9
	2016-2020	9.8	9.0	6.6-12.1	4.8	9.0	5.7-13.9	5.0	9.0	5.7-13.8	4.4	4.0	2.5-6.3	2.6	5.3	2.8-9.6	1.8	2.7	1.2-6.0
Leukemias	1976-1980	7.8	13.6	9.6-18.8	3.6	13.1	7.7-21.1	4.2	13.2	8.1-20.3	4.2	7.4	4.5-11.4	2.8	12.5	6.4-21.8	1.4	4.2	1.7-8.9
	1981-1985	6.8	11.8	8.1-16.5	4.6	19.4	12.0-29.5	2.2	6.4	3.2-11.7	4.6	8.0	5.1-12.1	2.4	10.7	5.4-18.9	2.2	6.2	3.1-11.4
	1986-1990	7.2	12.1	8.4-16.8	4.2	16.7	10.2-25.6	3.0	7.6	4.2-12.8	5.0	8.3	5.3-12.2	2.6	10.8	5.7-18.5	2.4	6.1	3.1-10.9
	1991-1995	9.2	13.9	10.2-18.6	5.6	21.5	14.1-31.2	3.6	10.0	5.8-16.0	5.4	7.9	5.2-11.6	3.4	12.9	7.3-20.9	2.0	5.2	2.5-9.8
	1996-2000	10.0	14.4	10.7-19.0	6.0	20.4	13.7-29.3	4.0	10.0	6.0-15.7	6.2	8.6	5.9-12.3	3.4	12.0	6.9-19.3	2.8	6.3	3.4-11.0
	2001-2005	13.2	17.6	13.6-22.6	7.6	23.4	16.5-32.3	5.6	12.9	8.4-19.0	8.2	10.5	7.6-14.4	5.4	17.0	11.1-25.0	2.8	5.8	3.2-10.1
	2006-2010	13.4	15.7	12.1-20.0	7.2	19.6	13.6-27.4	6.2	12.9	8.6-18.7	6.0	6.9	4.6-10.1	3.2	8.5	4.8-14.1	2.8	5.5	2.9-9.8
	2011-2015	18.4	21.7	17.2-26.9	10.0	26.5	19.4-35.4	8.4	17.8	12.5-24.8	9.4	9.9	7.3-13.4	5.6	14.0	9.2-20.5	3.8	7.0	4.2-11.4
	2016-2020	15.8	15.8	12.3-20.1	10.0	22.5	16.4-30.4	5.8	10.0	6.5-15.1	7.2	7.0	4.9-10.0	4.0	8.6	5.2-13.8	3.2	5.5	3.1-9.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Washington**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	182.2	356.5	333.3-380.9	95.8	446.6	405.4-490.6	86.4	304.5	275.8-335.3	114.4	224.6	206.2-244.1	63.2	303.0	268.7-340.2	51.2	177.6	156.1-201.2
	1981-1985	223.4	420.7	396.0-446.6	107.2	469.5	429.1-512.6	116.2	396.1	363.7-430.7	118.6	221.4	203.7-240.2	61.4	275.6	244.4-309.6	57.2	184.6	163.5-207.9
	1986-1990	254.0	447.6	423.0-473.2	135.2	561.3	518.2-606.9	118.8	378.6	348.1-411.1	125.2	222.2	205.0-240.5	67.4	295.4	263.4-329.9	57.8	176.9	156.7-199.0
	1991-1995	273.0	442.4	419.0-466.7	147.8	552.4	512.0-595.1	125.2	373.4	344.2-404.4	141.4	230.9	214.1-248.7	74.2	293.2	263.1-325.7	67.2	191.8	171.6-213.9
	1996-2000	339.8	521.1	496.6-546.5	174.2	597.6	557.9-639.3	165.6	471.0	439.1-504.7	144.4	221.2	205.3-237.9	75.4	273.1	245.6-302.7	69.0	186.4	167.1-207.5
	2001-2005	357.8	515.8	492.1-540.4	193.2	618.9	579.8-659.9	164.6	447.5	417.1-479.6	143.4	205.3	190.5-221.0	77.2	259.8	234.0-287.7	66.2	169.0	151.1-188.6
	2006-2010	413.2	544.3	520.8-568.7	217.4	612.8	576.2-651.2	195.8	499.8	468.3-533.1	145.8	188.4	174.9-202.8	72.8	214.1	192.2-237.9	73.0	173.9	156.3-193.2
	2011-2015	402.4	496.3	474.3-519.1	212.6	547.5	514.2-582.5	189.8	461.7	431.6-493.5	146.6	178.3	165.4-192.0	81.8	222.1	200.6-245.4	64.8	145.4	129.6-162.8
	2016-2020	432.2	493.7	472.2-515.9	222.4	508.0	477.6-540.0	209.8	492.4	461.1-525.4	150.4	166.6	154.6-179.4	75.0	176.9	159.0-196.6	75.4	158.5	142.4-176.3
Oral cavity and pharynx	1976-1980	3.8	7.3	4.4-11.4	3.4	16.7	9.4-27.0	0.4	1.3	0.2-5.0	1.4	2.9	1.2-5.9	1.0	5.2	1.6-12.2	0.4	1.4	0.2-5.1
	1981-1985	4.8	9.0	5.7-13.5	3.8	17.1	10.1-27.0	1.0	2.9	0.9-7.1	2.4	4.6	2.4-8.1	1.4	7.7	2.9-15.8	1.0	3.0	1.0-7.2
	1986-1990	4.4	7.8	4.9-11.9	2.6	9.8	5.1-16.8	1.8	6.6	2.9-12.6	2.0	3.8	1.8-7.0	1.6	6.3	2.7-12.5	0.4	1.1	0.1-4.4
	1991-1995	5.8	9.5	6.3-13.6	4.4	15.8	9.8-24.0	1.4	4.4	1.7-9.2	1.4	2.3	0.9-4.7	1.0	3.4	1.1-8.1	0.4	1.2	0.1-4.5
	1996-2000	6.4	9.8	6.7-13.9	4.2	15.0	9.2-23.1	2.2	5.8	2.9-10.7	2.8	4.2	2.3-7.2	2.0	7.2	3.4-13.3	0.8	2.1	0.6-5.6
	2001-2005	10.2	14.7	10.9-19.4	7.0	22.2	15.3-31.0	3.2	8.6	4.9-14.2	2.0	2.8	1.3-5.2	1.8	6.0	2.7-11.5	0.2	0.4	0.0-3.2
	2006-2010	9.0	11.3	8.2-15.2	6.2	16.5	11.1-23.7	2.8	6.7	3.6-11.6	1.2	1.5	0.5-3.4	1.2	3.3	1.2-7.5	0.0	0.0	0.0-2.5
	2011-2015	9.0	10.3	7.5-14.0	5.8	13.5	9.0-19.8	3.2	7.2	4.0-12.2	2.2	2.8	1.4-5.1	1.2	2.9	1.0-6.6	1.0	2.5	0.8-6.3
	2016-2020	12.8	13.6	10.4-17.6	10.0	21.5	15.8-28.8	2.8	5.9	3.2-10.6	2.0	2.2	1.0-4.4	1.0	2.3	0.7-6.0	1.0	1.9	0.6-5.3
Esophagus	1976-1980	2.0	4.0	1.9-7.4	1.0	4.5	1.4-10.7	1.0	3.3	1.1-7.9	1.6	3.2	1.3-6.2	0.4	1.3	0.2-5.4	1.2	4.0	1.5-8.8
	1981-1985	1.6	3.1	1.3-6.1	1.0	4.5	1.3-10.7	0.6	2.4	0.5-7.0	2.4	4.5	2.3-8.0	1.8	8.0	3.4-15.4	0.6	2.4	0.4-6.9
	1986-1990	2.0	3.4	1.6-6.3	1.2	4.6	1.7-10.2	0.8	2.4	0.6-6.4	2.0	3.5	1.7-6.5	0.8	3.1	0.8-8.1	1.2	3.5	1.3-7.9
	1991-1995	2.0	3.4	1.6-6.2	1.2	4.7	1.7-10.1	0.8	2.3	0.6-6.1	2.4	4.0	2.0-6.9	2.0	7.4	3.5-13.6	0.4	1.1	0.1-4.2
	1996-2000	4.8	7.4	4.7-11.0	4.2	15.0	9.2-23.0	0.6	1.7	0.3-5.2	2.8	4.2	2.3-7.1	2.2	8.0	3.9-14.5	0.6	1.7	0.3-5.2
	2001-2005	5.0	7.2	4.6-10.6	3.4	10.7	6.2-17.2	1.6	4.1	1.8-8.5	4.6	6.6	4.2-10.0	3.0	9.9	5.5-16.4	1.6	4.0	1.7-8.2
	2006-2010	6.4	8.0	5.5-11.5	5.0	13.6	8.7-20.4	1.4	3.4	1.4-7.4	3.2	4.1	2.3-6.8	2.6	7.3	3.8-12.7	0.6	1.3	0.3-4.5
	2011-2015	4.2	4.8	3.0-7.5	3.0	7.3	4.0-12.4	1.2	2.5	0.9-6.1	4.2	4.8	2.9-7.4	3.6	8.8	5.1-14.3	0.6	1.6	0.3-5.0
	2016-2020	6.4	7.1	4.8-10.3	4.6	10.2	6.4-15.8	1.8	4.5	2.0-9.0	3.8	4.3	2.5-7.0	2.4	5.6	2.8-10.2	1.4	3.4	1.2-7.7
Stomach	1976-1980	4.4	8.2	5.1-12.5	2.8	13.0	6.9-22.1	1.6	5.0	2.2-10.1	3.4	6.8	4.0-11.0	1.8	9.3	4.1-17.9	1.6	5.2	2.2-10.4
	1981-1985	4.4	8.4	5.2-12.8	2.0	7.8	3.7-14.5	2.4	8.1	4.1-14.4	1.8	3.4	1.6-6.6	0.8	3.7	1.0-9.4	1.0	3.0	1.0-7.2
	1986-1990	4.8	8.7	5.6-13.0	2.6	12.2	6.4-20.9	2.2	7.1	3.5-13.0	3.2	5.9	3.3-9.5	1.8	9.5	4.2-17.8	1.4	4.1	1.7-8.7
	1991-1995	5.4	8.7	5.7-12.6	3.6	13.7	8.0-21.9	1.8	5.0	2.2-9.6	3.0	5.0	2.8-8.2	2.2	9.2	4.5-16.5	0.8	2.4	0.6-6.3
	1996-2000	3.4	5.2	3.1-8.4	1.2	4.9	1.7-10.6	2.2	6.1	3.1-11.2	2.2	3.4	1.7-6.1	1.0	3.8	1.2-8.9	1.2	3.3	1.2-7.4
	2001-2005	6.2	8.7	5.9-12.4	3.8	12.5	7.4-19.8	2.4	6.2	3.2-11.0	2.6	3.7	2.0-6.4	1.8	6.6	2.9-12.5	0.8	2.0	0.5-5.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Washington**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	4.8	6.0	3.8-9.1	2.8	7.6	4.1-13.1	2.0	4.8	2.3-9.1	3.2	3.9	2.2-6.5	2.2	6.4	3.1-11.6	1.0	2.2	0.7-5.6
	2011-2015	6.2	7.5	5.0-10.9	4.0	9.8	5.8-15.5	2.2	5.0	2.4-9.5	2.0	2.4	1.1-4.6	1.0	2.6	0.8-6.5	1.0	2.0	0.6-5.3
	2016-2020	5.4	5.9	3.8-9.0	3.4	7.4	4.2-12.4	2.0	4.3	1.9-8.7	1.6	1.9	0.8-4.0	1.2	3.1	1.1-7.1	0.4	0.7	0.1-3.5
Colorectal	1976-1980	32.4	63.2	53.7-73.9	15.0	67.7	52.6-85.7	17.4	58.8	46.9-72.8	18.6	36.7	29.5-45.0	9.4	44.4	32.0-59.7	9.2	31.4	22.8-42.1
	1981-1985	40.6	76.4	66.1-87.9	16.8	75.9	60.0-94.5	23.8	76.5	63.2-91.9	18.0	33.7	27.1-41.6	8.0	37.4	26.2-51.6	10.0	30.9	22.8-41.0
	1986-1990	36.2	63.7	54.6-73.8	16.8	71.5	56.5-89.2	19.4	58.0	46.9-71.0	17.8	32.2	25.8-39.7	8.8	39.3	28.1-53.1	9.0	26.4	19.2-35.6
	1991-1995	36.8	60.3	51.8-69.7	15.6	62.2	48.7-78.0	21.2	58.7	48.0-71.3	18.8	30.8	24.8-37.7	7.6	30.0	21.1-41.4	11.2	30.4	22.9-39.7
	1996-2000	44.2	67.5	58.9-77.0	21.2	76.5	62.3-92.8	23.0	62.4	51.4-75.1	16.2	24.7	19.6-30.7	7.8	29.7	20.9-40.8	8.4	21.8	15.7-29.8
	2001-2005	36.2	51.9	44.6-60.1	16.2	52.8	41.8-65.9	20.0	51.6	41.9-63.0	15.2	21.7	17.1-27.2	6.8	23.6	16.1-33.1	8.4	20.7	14.9-28.3
	2006-2010	31.0	40.0	33.9-46.9	15.2	42.6	33.4-53.7	15.8	37.1	29.2-46.7	11.4	14.6	11.0-19.0	5.2	15.3	9.9-22.7	6.2	14.0	9.4-20.3
	2011-2015	33.2	40.7	34.7-47.7	17.0	42.8	34.0-53.3	16.2	36.7	28.9-46.3	12.2	14.3	10.9-18.5	5.2	12.6	8.2-18.9	7.0	14.3	9.9-20.5
2016-2020	32.2	38.4	32.5-45.2	16.2	39.3	31.0-49.4	16.0	38.0	29.6-48.3	13.2	14.6	11.1-18.8	7.0	15.8	10.8-22.5	6.2	12.8	8.5-18.9	
Colon excluding rectum	1976-1980	21.8	42.8	35.0-51.8	9.2	43.3	31.2-58.4	12.6	42.2	32.3-54.2	12.8	25.0	19.2-32.0	6.0	27.0	17.8-39.0	6.8	23.3	16.0-32.8
	1981-1985	28.2	53.2	44.6-62.8	11.0	50.7	37.7-66.4	17.2	55.2	44.0-68.5	14.0	26.6	20.7-33.7	6.6	31.2	21.0-44.4	7.4	23.3	16.3-32.4
	1986-1990	25.0	44.2	36.8-52.8	10.4	44.7	32.9-59.1	14.6	44.0	34.4-55.6	15.8	28.5	22.5-35.5	7.2	32.3	22.2-45.2	8.6	25.3	18.2-34.4
	1991-1995	26.8	43.6	36.5-51.7	9.8	39.3	28.8-52.3	17.0	46.7	37.2-58.0	16.6	27.3	21.7-33.9	6.2	25.1	16.8-35.8	10.4	28.3	21.1-37.4
	1996-2000	32.0	49.0	41.7-57.2	14.2	51.7	40.1-65.5	17.8	48.0	38.5-59.3	12.4	18.9	14.5-24.3	6.0	23.6	15.7-33.9	6.4	16.6	11.3-23.7
	2001-2005	26.6	38.2	32.0-45.3	11.2	37.0	27.7-48.3	15.4	39.6	31.2-49.7	12.0	17.1	13.0-22.0	5.0	17.0	10.9-25.3	7.0	17.3	12.0-24.4
	2006-2010	25.0	32.5	27.0-38.8	10.8	31.3	23.4-41.2	14.2	32.8	25.5-41.7	9.2	11.6	8.5-15.6	4.2	12.1	7.4-18.7	5.0	10.8	7.0-16.4
	2011-2015	25.2	31.2	25.9-37.4	12.8	32.9	25.2-42.4	12.4	28.0	21.2-36.5	10.0	11.5	8.5-15.4	4.6	11.3	7.1-17.3	5.4	10.4	6.8-15.7
2016-2020	21.6	26.6	21.7-32.5	9.4	23.9	17.4-32.2	12.2	29.0	21.7-38.1	9.2	10.7	7.7-14.6	4.4	10.6	6.5-16.6	4.8	10.4	6.5-16.1	
Rectum & rectosigmoid	1976-1980	10.6	20.4	15.2-26.7	5.8	24.4	16.0-35.6	4.8	16.6	10.5-24.9	5.8	11.7	7.8-16.8	3.4	17.4	9.8-28.2	2.4	8.0	4.1-14.2
	1981-1985	12.4	23.3	17.8-30.0	5.8	25.2	16.7-36.5	6.6	21.3	14.6-30.2	4.0	7.1	4.3-11.1	1.4	6.2	2.4-12.9	2.6	7.6	4.0-13.2
	1986-1990	11.2	19.4	14.6-25.3	6.4	26.8	18.0-38.2	4.8	14.0	9.0-21.0	2.0	3.8	1.8-6.9	1.6	6.9	2.9-13.6	0.4	1.1	0.1-4.4
	1991-1995	10.0	16.6	12.3-22.0	5.8	22.8	15.0-33.0	4.2	12.0	7.4-18.6	2.2	3.4	1.7-6.2	1.4	5.0	2.0-10.3	0.8	2.1	0.6-5.6
	1996-2000	12.2	18.5	14.2-23.8	7.0	24.8	17.1-34.6	5.2	14.4	9.4-21.3	3.8	5.7	3.5-9.0	1.8	6.0	2.7-11.5	2.0	5.2	2.5-9.9
	2001-2005	9.6	13.7	10.1-18.3	5.0	15.9	10.2-23.6	4.6	12.0	7.5-18.2	3.2	4.7	2.7-7.6	1.8	6.6	2.9-12.5	1.4	3.4	1.3-7.4
	2006-2010	6.0	7.5	5.0-10.9	4.4	11.3	7.0-17.4	1.6	4.3	1.8-8.9	2.2	2.9	1.5-5.4	1.0	3.3	1.0-7.7	1.2	3.2	1.1-7.3
	2011-2015	8.0	9.5	6.7-13.2	4.2	9.8	6.0-15.5	3.8	8.7	5.1-14.2	2.2	2.7	1.3-5.0	0.6	1.3	0.3-4.4	1.6	4.0	1.7-8.2
2016-2020	10.6	11.8	8.7-15.7	6.8	15.4	10.5-22.0	3.8	9.0	5.2-14.8	4.0	3.8	2.3-6.2	2.6	5.1	2.7-9.3	1.4	2.5	1.0-5.9	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Washington**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	2.0	3.8	1.8-7.0	1.0	4.7	1.4-11.1	1.0	3.3	1.1-7.9	1.4	2.8	1.1-5.7	0.8	4.6	1.2-11.4	0.6	2.0	0.4-5.9
	1981-1985	1.0	1.8	0.6-4.2	0.6	2.1	0.4-6.5	0.4	1.2	0.1-4.8	1.6	2.9	1.2-5.7	1.0	4.3	1.2-10.5	0.6	1.8	0.4-5.6
	1986-1990	2.8	4.7	2.6-8.0	2.2	7.8	3.9-14.2	0.6	1.7	0.4-5.4	2.2	3.6	1.8-6.5	1.2	4.5	1.6-10.1	1.0	2.8	0.9-6.8
	1991-1995	2.4	3.7	1.9-6.4	1.8	5.8	2.7-11.2	0.6	1.6	0.3-4.9	2.0	3.1	1.5-5.7	1.4	4.5	1.8-9.5	0.6	1.6	0.3-4.9
	1996-2000	2.8	4.2	2.3-7.1	1.4	4.4	1.8-9.2	1.4	3.8	1.5-8.0	2.8	4.3	2.3-7.2	1.6	5.4	2.3-10.7	1.2	3.2	1.2-7.3
	2001-2005	4.0	5.7	3.5-8.9	2.8	8.6	4.7-14.6	1.2	3.1	1.1-7.1	3.8	5.2	3.2-8.3	2.6	8.0	4.2-13.8	1.2	3.0	1.1-7.0
	2006-2010	3.8	4.9	3.0-7.8	2.6	7.0	3.7-12.2	1.2	2.9	1.0-6.6	4.6	5.8	3.7-8.9	2.6	6.9	3.6-12.0	2.0	4.8	2.3-9.1
	2011-2015	6.0	6.9	4.6-10.0	4.2	9.8	6.0-15.5	1.8	3.8	1.7-7.7	3.6	4.4	2.6-7.1	2.2	5.5	2.7-10.3	1.4	2.9	1.1-6.6
	2016-2020	8.6	9.5	6.8-13.0	6.4	14.1	9.6-20.4	2.2	5.4	2.6-10.2	5.4	6.3	4.1-9.4	4.0	9.4	5.7-15.0	1.4	3.4	1.2-7.7
Pancreas	1976-1980	9.0	17.8	12.9-23.9	5.8	29.4	19.3-42.5	3.2	11.3	6.3-18.5	8.4	16.5	11.8-22.4	5.4	25.8	16.6-38.0	3.0	10.8	5.9-18.0
	1981-1985	5.4	10.1	6.7-14.8	2.4	11.8	5.8-20.8	3.0	9.2	5.1-15.4	6.2	11.7	7.9-16.6	2.6	11.9	6.1-20.7	3.6	11.0	6.5-17.6
	1986-1990	4.2	7.2	4.5-11.1	2.2	9.6	4.7-17.1	2.0	5.6	2.7-10.5	4.0	6.8	4.1-10.5	2.2	9.5	4.7-16.9	1.8	5.1	2.3-9.9
	1991-1995	6.6	10.7	7.3-15.0	3.4	12.7	7.2-20.6	3.2	9.3	5.3-15.3	5.4	8.7	5.7-12.7	2.4	9.4	4.6-16.7	3.0	8.6	4.8-14.4
	1996-2000	10.6	16.4	12.3-21.5	5.2	18.9	12.2-27.8	5.4	14.6	9.6-21.5	9.2	14.1	10.3-18.8	4.6	16.4	10.3-24.7	4.6	12.3	7.8-18.7
	2001-2005	7.4	10.6	7.4-14.6	4.2	13.5	8.3-20.8	3.2	7.9	4.5-13.1	7.8	11.1	7.9-15.3	3.6	11.5	6.8-18.3	4.2	10.2	6.3-15.9
	2006-2010	12.0	15.4	11.7-20.0	6.2	18.1	12.1-25.9	5.8	13.9	9.3-20.4	10.4	13.0	9.7-17.2	5.0	14.0	8.9-21.0	5.4	12.9	8.4-19.2
	2011-2015	10.2	12.3	9.1-16.3	6.4	16.7	11.3-23.9	3.8	8.3	5.0-13.4	7.4	9.3	6.5-13.0	4.6	12.7	7.9-19.2	2.8	6.4	3.5-11.1
	2016-2020	14.4	15.9	12.4-20.3	8.0	17.8	12.6-24.8	6.4	13.6	9.2-19.8	12.0	13.4	10.1-17.4	7.4	17.3	12.0-24.3	4.6	9.4	5.9-14.7
Larynx	1976-1980	1.8	3.0	1.4-5.9	1.4	5.7	2.2-12.0	0.4	1.2	0.1-4.6	1.0	1.7	0.6-4.2	1.0	4.1	1.3-9.9	0.0	0.0	0.0-2.8
	1981-1985	2.6	5.0	2.6-8.5	2.0	9.4	4.3-17.4	0.6	1.9	0.4-5.8	1.4	2.5	1.0-5.2	1.0	4.0	1.2-9.6	0.4	1.2	0.1-4.8
	1986-1990	2.2	4.0	2.0-7.3	1.6	6.2	2.6-12.3	0.6	2.1	0.4-6.2	0.8	1.6	0.4-4.1	0.8	4.0	1.0-10.1	0.0	0.0	0.0-2.6
	1991-1995	2.2	3.5	1.7-6.3	1.6	5.6	2.4-11.0	0.6	1.8	0.3-5.4	0.6	0.9	0.2-2.6	0.6	1.9	0.4-5.7	0.0	0.0	0.0-2.4
	1996-2000	2.8	4.3	2.3-7.2	2.2	6.8	3.4-12.4	0.6	1.7	0.3-5.3	1.0	1.5	0.5-3.6	0.6	2.2	0.4-6.3	0.4	1.2	0.1-4.6
	2001-2005	4.2	6.1	3.8-9.4	3.2	10.3	5.8-16.8	1.0	2.7	0.9-6.7	1.8	2.6	1.2-5.0	1.4	5.2	2.1-10.6	0.4	1.1	0.1-4.4
	2006-2010	4.4	5.8	3.6-8.9	3.6	9.9	5.8-15.8	0.8	1.8	0.5-5.2	0.8	0.9	0.2-2.6	0.8	2.0	0.5-5.4	0.0	0.0	0.0-2.5
	2011-2015	4.2	5.0	3.0-7.8	3.0	7.5	4.1-12.7	1.2	3.1	1.1-7.3	2.0	2.4	1.1-4.6	2.0	5.4	2.5-10.1	0.0	0.0	0.0-2.6
	2016-2020	4.8	4.8	3.1-7.4	3.6	7.9	4.6-13.0	1.2	2.2	0.8-5.6	1.0	1.0	0.3-2.6	0.8	1.8	0.5-5.2	0.2	0.3	0.0-3.1
Lung and bronchus	1976-1980	31.2	59.8	50.6-70.1	22.4	100.0	81.5-121.3	8.8	31.9	23.0-43.0	25.2	48.3	40.1-57.7	20.0	91.5	73.6-112.2	5.2	18.4	11.9-27.3
	1981-1985	37.0	69.0	59.2-79.9	23.6	101.7	83.7-122.3	13.4	44.8	34.5-57.3	31.4	58.6	49.7-68.7	20.8	89.8	73.0-109.3	10.6	34.8	25.9-45.8
	1986-1990	47.8	83.1	72.8-94.5	31.4	127.5	107.8-149.6	16.4	50.8	40.2-63.4	38.6	66.4	57.3-76.6	25.0	101.5	84.0-121.5	13.6	41.7	32.2-53.1
	1991-1995	49.4	79.5	69.8-90.1	31.8	118.4	100.3-138.8	17.6	51.4	41.1-63.6	41.4	67.1	58.2-77.0	25.8	97.4	80.8-116.2	15.6	45.4	35.8-57.0
	1996-2000	61.4	93.2	83.0-104.2	36.2	122.0	104.7-141.4	25.2	70.0	58.2-83.5	45.6	69.4	60.7-79.1	26.6	92.7	77.3-110.1	19.0	51.8	41.8-63.6
	2001-2005	58.6	83.5	74.2-93.7	33.0	103.9	88.4-121.3	25.6	67.9	56.6-80.9	41.2	58.8	51.1-67.5	26.4	85.0	70.9-101.1	14.8	38.2	29.9-48.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Washington**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	72.4	95.4	85.7-105.9	38.0	109.5	94.3-126.6	34.4	84.8	72.5-98.8	45.6	59.2	51.7-67.5	22.6	65.5	53.8-79.0	23.0	55.5	45.8-67.0
	2011-2015	67.4	81.2	72.6-90.6	37.2	97.4	83.6-112.9	30.2	68.5	57.8-80.9	45.6	55.0	48.0-62.9	27.0	71.2	59.5-84.7	18.6	42.6	34.3-52.7
	2016-2020	69.4	75.0	67.2-83.7	33.0	74.5	63.2-87.3	36.4	76.5	65.6-89.0	40.0	43.2	37.3-49.9	18.8	43.4	34.9-53.7	21.2	43.0	35.1-52.5
Melanoma of the skin	1976-1980	1.8	3.4	1.6-6.6	0.8	3.4	0.9-9.0	1.0	3.7	1.2-8.6	1.2	2.5	0.9-5.5	0.6	2.7	0.5-8.0	0.6	2.9	0.6-8.0
	1981-1985	4.0	8.8	5.3-13.6	2.2	9.7	4.7-17.4	1.8	8.6	3.9-16.0	0.6	1.0	0.2-3.0	0.4	1.4	0.2-5.5	0.2	0.6	0.0-3.8
	1986-1990	3.2	6.0	3.4-9.9	2.4	9.9	5.0-17.5	0.8	3.2	0.8-8.1	1.6	3.1	1.3-6.2	1.4	6.1	2.4-12.6	0.2	0.6	0.0-3.6
	1991-1995	5.6	9.2	6.1-13.3	2.8	11.0	5.8-18.7	2.8	8.8	4.7-14.8	1.8	3.1	1.4-5.8	0.4	1.3	0.2-4.9	1.4	4.3	1.7-9.1
	1996-2000	7.8	12.3	8.7-16.8	4.8	16.2	10.3-24.1	3.0	9.1	5.0-15.2	1.6	2.4	1.1-4.9	1.2	4.1	1.5-9.0	0.4	1.0	0.1-4.0
	2001-2005	11.8	17.2	13.1-22.3	7.8	24.8	17.6-34.1	4.0	11.5	7.0-18.1	2.6	3.8	2.0-6.5	1.4	5.1	2.0-10.6	1.2	3.0	1.1-7.0
	2006-2010	15.8	21.2	16.7-26.6	9.4	27.2	19.9-36.5	6.4	17.1	11.6-24.5	2.2	2.9	1.4-5.3	1.4	4.2	1.7-8.8	0.8	1.9	0.5-5.3
	2011-2015	19.2	24.9	20.1-30.7	10.2	29.1	21.6-38.6	9.0	23.8	16.9-32.6	2.0	2.7	1.2-5.1	1.6	5.4	2.3-10.7	0.4	0.9	0.1-3.9
2016-2020	17.8	20.4	16.2-25.5	10.8	24.7	18.4-32.7	7.0	17.9	12.1-25.8	2.2	2.4	1.2-4.6	1.4	3.6	1.4-7.7	0.8	1.6	0.4-4.8	
Female breast	1976-1980							19.6	70.6	57.0-86.4							10.0	36.3	26.7-48.3
	1981-1985							26.4	97.2	80.8-115.9							9.4	31.8	23.1-42.7
	1986-1990							28.6	99.8	83.7-118.1							8.6	26.9	19.3-36.5
	1991-1995							28.4	87.1	73.1-103.0							9.0	26.6	19.3-35.9
	1996-2000							47.2	137.4	120.3-156.4							8.2	22.6	16.2-30.9
	2001-2005							47.8	131.1	114.9-149.1							8.8	23.0	16.7-31.1
	2006-2010							50.6	131.7	115.7-149.5							9.8	22.8	16.8-30.5
	2011-2015							45.8	114.1	99.3-130.7							7.6	17.1	11.9-24.0
2016-2020							60.6	146.3	129.1-165.2							8.8	18.8	13.4-26.0	
Cervix uteri	1976-1980							4.6	19.6	12.2-29.6							2.4	8.5	4.3-15.1
	1981-1985							4.6	18.3	11.5-27.6							1.6	5.3	2.3-10.7
	1986-1990							2.4	7.6	3.9-13.4							0.8	3.0	0.8-7.7
	1991-1995							4.2	14.6	8.9-22.3							1.0	3.1	1.0-7.3
	1996-2000							2.8	8.9	4.8-15.0							1.0	3.3	1.1-7.7
	2001-2005							3.2	9.9	5.6-16.3							1.2	3.1	1.1-7.0
	2006-2010							4.4	13.2	8.1-20.4							1.0	2.5	0.8-6.3
	2011-2015							2.4	6.6	3.3-12.0							0.6	1.2	0.2-4.3
	2016-2020							3.4	11.9	6.7-19.4							1.4	3.1	1.2-7.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Washington**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							5.6	19.0	12.6-27.7							1.2	3.7	1.4-8.4
	1981-1985							7.0	23.8	16.4-33.5							1.2	3.8	1.4-8.4
	1986-1990							6.0	20.3	13.6-29.2							1.0	3.0	1.0-7.3
	1991-1995							7.4	23.3	16.3-32.3							2.6	7.3	3.8-12.7
	1996-2000							9.8	28.4	21.0-37.7							2.4	6.3	3.2-11.3
	2001-2005							9.6	26.2	19.3-34.9							2.6	6.6	3.5-11.6
	2006-2010							15.0	37.6	29.5-47.5							2.6	6.2	3.3-11.1
	2011-2015							16.4	37.3	29.5-46.8							2.2	4.6	2.3-8.8
	2016-2020							12.4	26.7	20.2-34.9							3.2	7.6	4.1-13.1
Ovary	1976-1980							3.0	11.0	6.0-18.4							2.2	8.0	3.9-14.5
	1981-1985							6.2	21.9	14.7-31.4							4.8	17.0	10.7-25.6
	1986-1990							5.2	17.4	11.2-25.7							3.8	13.0	7.7-20.6
	1991-1995							5.0	15.0	9.6-22.3							3.8	11.0	6.6-17.3
	1996-2000							7.0	20.7	14.4-29.0							4.2	11.7	7.2-18.1
	2001-2005							6.0	16.3	11.0-23.5							3.6	9.1	5.4-14.7
	2006-2010							4.8	12.3	7.8-18.6							3.4	8.7	5.0-14.2
	2011-2015							5.0	12.8	8.1-19.4							3.8	9.2	5.4-14.7
	2016-2020							4.6	10.4	6.4-16.3							4.0	8.0	4.9-13.0
Prostate	1976-1980				14.2	74.6	57.5-94.8							6.6	37.9	25.6-53.5			
	1981-1985				17.2	81.1	64.1-100.9							6.6	35.6	24.1-50.2			
	1986-1990				28.2	126.9	105.9-150.5							7.6	37.9	26.3-52.3			
	1991-1995				41.2	151.9	131.3-174.9							9.2	41.6	30.1-55.7			
	1996-2000				45.2	154.4	134.7-176.3							7.0	29.2	20.2-40.6			
	2001-2005				50.6	159.4	140.1-180.7							5.4	21.3	13.9-31.0			
	2006-2010				58.8	159.3	141.2-179.1							8.0	28.3	20.1-38.5			
	2011-2015				46.4	111.3	97.1-127.1							7.4	21.8	15.2-30.2			
	2016-2020				47.6	100.6	88.0-114.8							6.4	15.2	10.3-21.8			
Testis	1976-1980				0.2	0.6	0.0-4.1							0.2	1.0	0.0-5.3			
	1981-1985				3.0	10.8	5.9-18.1							0.4	1.2	0.1-5.0			
	1986-1990				1.6	5.2	2.2-10.5							0.2	0.6	0.0-3.9			
	1991-1995				1.0	2.8	0.9-6.8							0.6	1.9	0.4-5.7			
	1996-2000				2.0	6.0	2.9-11.2							0.2	0.6	0.0-3.6			
	2001-2005				2.8	9.1	4.9-15.2							0.0	0.0	0.0-2.5			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Washington**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				3.0	9.5	5.3-15.8							0.0	0.0	0.0-2.4			
	2011-2015				2.2	7.5	3.7-13.4							0.2	0.8	0.0-4.1			
	2016-2020				1.0	3.7	1.2-8.5							0.0	0.0	0.0-2.5			
Urinary bladder (incl. in situ)	1976-1980	8.6	16.5	11.9-22.3	6.4	29.6	19.8-42.3	2.2	7.3	3.6-13.3	6.0	11.6	7.8-16.7	3.6	17.1	9.8-27.5	2.4	7.7	4.0-13.6
	1981-1985	11.0	20.2	15.1-26.4	7.8	35.9	25.1-49.5	3.2	9.8	5.5-16.3	3.2	6.0	3.4-9.9	2.2	10.4	5.0-18.8	1.0	2.9	1.0-7.2
	1986-1990	16.6	28.7	22.8-35.7	11.4	45.9	34.5-59.8	5.2	15.3	9.9-22.7	4.0	7.0	4.3-10.9	2.2	10.4	5.0-18.7	1.8	5.2	2.4-10.1
	1991-1995	13.0	21.3	16.4-27.2	9.0	35.4	25.5-47.7	4.0	11.2	6.8-17.6	4.8	7.9	5.0-11.8	3.2	14.4	8.0-23.4	1.6	4.0	1.7-8.2
	1996-2000	16.4	24.9	19.8-30.9	12.4	41.9	32.0-53.9	4.0	10.5	6.4-16.5	3.2	4.9	2.8-7.9	1.8	6.1	2.8-11.7	1.4	3.3	1.3-7.2
	2001-2005	18.8	27.2	22.0-33.4	14.8	52.1	40.7-65.6	4.0	10.5	6.4-16.4	4.8	6.9	4.4-10.3	3.4	12.1	6.9-19.4	1.4	3.3	1.3-7.2
	2006-2010	22.0	28.3	23.2-34.2	16.6	48.6	38.5-60.6	5.4	13.2	8.6-19.6	3.6	4.7	2.8-7.5	2.8	8.3	4.5-14.2	0.8	2.1	0.6-5.7
	2011-2015	24.6	30.1	25.0-36.1	19.0	51.0	41.1-62.7	5.6	13.5	8.8-19.9	5.2	6.2	4.0-9.3	4.2	11.6	7.1-18.0	1.0	2.9	0.9-7.2
	2016-2020	22.0	23.8	19.4-29.0	16.6	37.3	29.5-46.8	5.4	11.8	7.6-17.8	1.8	2.2	1.0-4.5	1.4	3.8	1.5-8.2	0.4	0.7	0.1-3.7
Kidney and renal pelvis	1976-1980	2.4	4.6	2.3-8.1	1.6	7.1	3.0-14.1	0.8	2.5	0.7-6.6	1.6	3.0	1.3-6.0	1.2	5.1	1.8-11.3	0.4	1.4	0.2-5.1
	1981-1985	5.0	8.8	5.7-13.1	3.0	11.5	6.3-19.3	2.0	6.1	2.9-11.4	2.0	3.9	1.9-7.2	0.8	3.1	0.8-8.2	1.2	3.6	1.3-8.2
	1986-1990	7.0	11.8	8.2-16.5	3.8	16.7	9.8-26.3	3.2	9.1	5.2-15.0	3.6	6.2	3.7-9.9	2.2	10.7	5.1-19.2	1.4	3.9	1.6-8.4
	1991-1995	5.4	8.4	5.5-12.2	3.4	11.4	6.6-18.4	2.0	5.8	2.7-10.9	2.8	4.6	2.5-7.7	1.6	6.7	2.8-13.2	1.2	3.4	1.2-7.7
	1996-2000	10.0	15.4	11.4-20.4	6.6	21.8	15.0-30.8	3.4	9.7	5.6-15.7	3.0	4.6	2.5-7.6	2.2	7.4	3.6-13.4	0.8	2.1	0.6-5.6
	2001-2005	9.6	14.0	10.3-18.7	6.2	19.9	13.4-28.4	3.4	9.1	5.3-14.9	2.2	3.1	1.6-5.7	1.2	3.9	1.4-8.7	1.0	2.6	0.8-6.4
	2006-2010	13.8	18.5	14.3-23.6	9.0	26.0	18.9-35.0	4.8	12.2	7.7-18.5	2.6	3.3	1.7-5.7	1.8	4.7	2.1-9.2	0.8	2.1	0.6-5.8
	2011-2015	14.2	16.7	13.0-21.4	9.0	22.8	16.4-30.9	5.2	11.1	7.2-16.7	3.6	4.5	2.6-7.2	2.2	5.9	2.9-10.8	1.4	2.9	1.1-6.5
	2016-2020	15.4	16.7	13.1-21.1	11.4	25.3	19.0-33.2	4.0	8.7	5.3-13.9	5.0	5.4	3.5-8.2	3.4	7.8	4.4-12.9	1.6	3.5	1.5-7.4
Brain and other central nervous system	1976-1980	1.6	2.9	1.2-5.8	0.8	3.5	0.9-9.2	0.8	2.5	0.7-6.7	0.6	1.2	0.2-3.5	0.2	1.3	0.0-6.1	0.4	1.3	0.2-4.9
	1981-1985	3.0	5.0	2.8-8.4	2.2	7.7	3.8-14.1	0.8	2.5	0.7-6.6	3.0	5.2	2.9-8.7	2.0	7.6	3.6-14.2	1.0	3.2	1.0-7.6
	1986-1990	5.6	10.2	6.7-14.8	3.8	15.0	8.9-23.7	1.8	6.5	2.9-12.5	3.6	6.8	4.0-10.7	2.4	10.5	5.3-18.4	1.2	4.5	1.6-9.8
	1991-1995	3.6	6.0	3.5-9.5	2.2	7.6	3.7-13.8	1.4	4.2	1.7-8.9	2.6	4.2	2.2-7.2	1.2	4.4	1.6-9.7	1.4	4.0	1.6-8.4
	1996-2000	5.0	7.8	5.0-11.5	2.4	7.5	3.9-13.2	2.6	7.9	4.2-13.6	2.2	3.3	1.7-6.0	1.2	3.7	1.3-8.2	1.0	3.0	1.0-7.2
	2001-2005	4.8	7.1	4.6-10.7	3.2	9.4	5.3-15.4	1.6	4.8	2.0-9.6	2.4	3.5	1.8-6.2	1.6	4.9	2.1-9.8	0.8	2.3	0.6-6.1
	2006-2010	5.8	8.0	5.3-11.6	3.8	10.2	6.1-16.2	2.0	5.9	2.8-11.2	4.0	5.0	3.1-7.9	2.8	7.1	3.8-12.3	1.2	3.0	1.1-7.0
	2011-2015	5.4	7.9	5.1-11.7	4.0	12.1	7.3-19.0	1.4	3.5	1.3-7.9	3.8	4.5	2.7-7.2	2.4	6.0	3.0-10.8	1.4	3.3	1.3-7.4
	2016-2020	6.6	7.8	5.2-11.2	4.4	10.7	6.6-16.7	2.2	4.7	2.3-9.2	6.0	6.9	4.6-10.2	3.6	7.9	4.6-13.0	2.4	6.4	3.1-11.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Washington**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	0.6	1.2	0.2-3.4	0.2	0.8	0.0-4.8	0.4	1.5	0.2-5.5	0.4	0.8	0.1-2.9	0.0	0.0	0.0-3.3	0.4	1.4	0.2-5.1
	1981-1985	1.0	1.6	0.5-3.9	0.2	0.7	0.0-4.3	0.8	2.8	0.8-7.2	0.0	0.0	0.0-1.4	0.0	0.0	0.0-3.2	0.0	0.0	0.0-2.7
	1986-1990	2.8	4.7	2.5-7.9	1.4	5.3	2.1-11.0	1.4	4.1	1.6-8.7	0.4	0.7	0.1-2.5	0.0	0.0	0.0-2.9	0.4	1.0	0.1-4.2
	1991-1995	1.0	1.7	0.6-4.0	0.2	0.7	0.0-4.1	0.8	2.7	0.7-6.9	0.4	0.6	0.1-2.2	0.0	0.0	0.0-2.7	0.4	1.1	0.1-4.2
	1996-2000	4.4	6.9	4.3-10.5	1.0	3.1	1.0-7.3	3.4	11.2	6.4-18.0	0.4	0.6	0.1-2.3	0.2	0.6	0.0-3.7	0.2	0.4	0.0-3.1
	2001-2005	7.2	11.0	7.7-15.2	1.8	5.5	2.5-10.5	5.4	17.3	11.3-25.4	0.6	0.8	0.2-2.5	0.2	0.7	0.0-3.9	0.4	1.0	0.1-4.2
	2006-2010	13.2	19.1	14.7-24.5	3.0	8.4	4.7-14.2	10.2	30.7	22.7-40.7	0.4	0.5	0.1-1.9	0.0	0.0	0.0-2.4	0.4	0.8	0.1-3.7
	2011-2015	16.6	23.6	18.6-29.5	4.2	10.1	6.2-15.9	12.4	38.9	29.4-50.4	0.4	0.6	0.1-2.1	0.4	1.2	0.1-4.4	0.0	0.0	0.0-2.6
	2016-2020	12.4	18.8	14.2-24.4	4.0	11.1	6.6-17.7	8.4	28.0	19.8-38.5	0.2	0.4	0.0-2.2	0.0	0.0	0.0-2.5	0.2	1.0	0.0-4.9
Hodgkin lymphoma	1976-1980	1.0	1.7	0.5-4.2	0.4	1.1	0.1-4.9	0.6	1.8	0.4-5.6	1.2	2.3	0.8-5.1	0.4	1.8	0.2-6.8	0.8	2.9	0.8-7.6
	1981-1985	2.4	4.0	2.0-7.1	1.0	3.3	1.0-8.1	1.4	4.4	1.8-9.2	1.2	2.1	0.8-4.7	0.8	2.7	0.7-7.3	0.4	1.2	0.1-4.6
	1986-1990	2.8	4.7	2.5-8.0	1.8	5.9	2.6-11.6	1.0	3.4	1.1-7.9	0.2	0.4	0.0-2.2	0.2	0.9	0.0-4.6	0.0	0.0	0.0-2.6
	1991-1995	2.8	4.5	2.5-7.6	1.6	4.8	2.1-9.7	1.2	3.9	1.4-8.6	0.6	1.0	0.2-2.9	0.4	1.4	0.2-5.2	0.2	0.5	0.0-3.2
	1996-2000	2.2	3.5	1.8-6.3	1.4	4.2	1.7-8.8	0.8	2.7	0.7-6.8	0.4	0.6	0.1-2.2	0.0	0.0	0.0-2.6	0.4	1.2	0.1-4.5
	2001-2005	1.6	2.6	1.1-5.1	0.8	2.4	0.7-6.4	0.8	2.7	0.7-6.8	0.2	0.4	0.0-2.0	0.2	0.7	0.0-3.9	0.0	0.0	0.0-2.5
	2006-2010	2.2	3.4	1.7-6.2	1.2	3.0	1.1-6.9	1.0	4.1	1.3-9.2	0.2	0.3	0.0-1.8	0.0	0.0	0.0-2.4	0.2	0.7	0.0-3.8
	2011-2015	1.8	2.6	1.2-5.1	1.4	4.0	1.5-8.4	0.4	1.0	0.1-4.1	0.6	0.7	0.1-2.3	0.2	0.4	0.0-3.0	0.4	0.9	0.1-3.9
	2016-2020	0.6	0.9	0.1-2.7	0.4	0.8	0.1-3.7	0.2	1.0	0.0-4.8	0.0	0.0	0.0-1.3	0.0	0.0	0.0-2.5	0.0	0.0	0.0-2.7
Non-Hodgkin lymphomas	1976-1980	5.8	11.6	7.7-16.8	4.0	19.5	11.6-30.5	1.8	6.5	2.9-12.5	3.8	7.2	4.3-11.3	2.4	10.9	5.3-19.6	1.4	4.7	1.9-9.9
	1981-1985	6.8	12.3	8.5-17.3	3.0	12.6	6.8-21.2	3.8	12.2	7.3-19.3	4.2	7.6	4.7-11.7	2.6	11.4	5.9-19.8	1.6	5.4	2.3-10.8
	1986-1990	10.8	19.1	14.3-25.0	6.2	25.7	17.1-36.9	4.6	13.5	8.5-20.4	4.6	8.2	5.2-12.3	2.0	9.7	4.4-17.9	2.6	7.4	3.9-12.9
	1991-1995	11.6	18.9	14.3-24.4	6.8	23.9	16.4-33.7	4.8	14.3	9.1-21.5	4.6	7.8	4.9-11.7	2.2	8.3	4.1-14.9	2.4	7.0	3.6-12.4
	1996-2000	14.0	21.4	16.6-27.0	5.6	18.1	12.0-26.2	8.4	22.9	16.5-31.2	7.0	11.0	7.7-15.3	3.4	11.9	6.9-19.1	3.6	9.6	5.7-15.5
	2001-2005	14.0	20.1	15.7-25.5	9.4	29.2	21.3-39.0	4.6	12.3	7.7-18.7	5.0	7.2	4.7-10.7	1.8	5.8	2.6-11.0	3.2	7.9	4.5-13.2
	2006-2010	16.6	22.0	17.5-27.5	9.6	28.5	20.9-38.0	7.0	16.8	11.7-23.7	5.4	7.0	4.6-10.3	3.6	10.6	6.2-17.0	1.8	4.1	1.9-8.2
	2011-2015	16.8	21.0	16.6-26.2	10.2	27.1	20.0-36.1	6.6	15.2	10.4-21.7	5.4	6.9	4.5-10.2	3.8	10.7	6.4-17.0	1.6	3.9	1.7-8.1
	2016-2020	18.2	21.2	16.9-26.3	11.8	28.0	21.1-36.6	6.4	14.1	9.4-20.6	4.4	5.3	3.3-8.3	2.2	5.9	2.9-11.0	2.2	4.7	2.3-9.1
Myeloma	1976-1980	2.4	4.4	2.3-7.8	1.0	3.7	1.2-8.9	1.4	4.7	1.9-9.8	1.0	1.6	0.5-4.0	0.8	2.7	0.7-7.3	0.2	0.6	0.0-3.8
	1981-1985	2.8	5.2	2.9-8.8	1.6	7.2	3.0-14.3	1.2	4.0	1.5-8.8	1.4	2.7	1.1-5.5	0.4	2.1	0.2-7.2	1.0	3.2	1.0-7.6
	1986-1990	3.6	6.5	3.8-10.4	1.8	8.4	3.7-15.8	1.8	4.9	2.2-9.7	1.6	2.9	1.2-5.7	1.2	5.6	1.9-12.1	0.4	1.2	0.1-4.5
	1991-1995	3.8	6.4	3.8-10.0	1.6	7.5	3.1-14.8	2.2	6.8	3.4-12.4	4.4	7.2	4.5-11.0	2.4	9.1	4.6-16.0	2.0	5.8	2.8-10.9
	1996-2000	3.6	5.5	3.2-8.7	2.2	7.7	3.8-13.8	1.4	3.3	1.3-7.3	2.0	3.1	1.5-5.7	1.4	6.1	2.4-12.3	0.6	1.5	0.3-4.7
	2001-2005	2.4	3.5	1.8-6.1	1.6	5.1	2.2-10.1	0.8	2.1	0.6-5.8	2.2	3.1	1.6-5.7	1.2	3.9	1.4-8.5	1.0	2.4	0.8-6.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Washington**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	3.4	4.4	2.5-7.1	2.4	6.6	3.4-11.8	1.0	2.2	0.7-5.7	2.0	2.6	1.3-5.0	1.4	4.3	1.7-8.9	0.6	1.3	0.3-4.3
	2011-2015	7.8	9.0	6.4-12.5	4.8	10.8	6.8-16.5	3.0	6.9	3.8-11.8	2.6	3.1	1.6-5.5	1.6	4.5	1.9-9.1	1.0	1.8	0.6-5.0
	2016-2020	9.8	10.6	7.8-14.3	5.8	13.1	8.7-19.4	4.0	8.5	5.2-13.7	3.6	3.8	2.2-6.2	2.0	4.5	2.1-8.7	1.6	3.2	1.3-6.9
Leukemias	1976-1980	8.8	16.2	11.7-21.9	5.2	24.8	15.6-37.1	3.6	11.5	6.8-18.4	5.8	10.8	7.2-15.6	3.2	16.2	8.8-26.9	2.6	8.4	4.5-14.6
	1981-1985	8.0	14.8	10.5-20.2	4.8	20.9	13.0-31.5	3.2	10.2	5.8-16.9	5.0	9.2	5.9-13.6	2.6	10.5	5.5-18.2	2.4	7.7	3.9-13.7
	1986-1990	6.8	12.1	8.3-16.9	3.8	15.5	9.2-24.4	3.0	8.9	4.9-14.9	4.4	7.5	4.7-11.5	2.0	7.7	3.6-14.4	2.4	7.2	3.6-12.8
	1991-1995	7.4	12.3	8.7-17.0	3.8	14.8	8.7-23.4	3.6	10.4	6.1-16.6	5.6	9.2	6.1-13.3	3.6	14.7	8.5-23.5	2.0	5.3	2.5-10.0
	1996-2000	8.0	12.3	8.8-16.8	5.4	19.7	12.8-28.8	2.6	7.8	4.1-13.7	5.0	7.6	4.9-11.3	3.6	13.5	7.9-21.5	1.4	3.4	1.4-7.4
	2001-2005	12.0	17.9	13.6-23.1	6.8	23.3	16.0-32.8	5.2	15.1	9.8-22.4	6.2	9.2	6.2-13.1	3.6	13.5	7.8-21.4	2.6	7.4	3.9-13.0
	2006-2010	9.6	13.5	9.9-18.1	6.2	18.0	12.1-25.8	3.4	9.4	5.3-15.4	3.0	4.2	2.3-7.0	1.4	4.3	1.7-8.9	1.6	4.1	1.7-8.5
	2011-2015	11.6	14.8	11.2-19.3	6.0	16.5	11.0-23.8	5.6	14.2	9.3-21.0	4.2	5.4	3.3-8.4	2.0	6.1	2.9-11.4	2.2	4.9	2.3-9.4
	2016-2020	12.8	16.1	12.2-20.8	7.4	19.6	13.6-27.5	5.4	12.5	8.0-18.9	5.2	6.0	3.9-9.0	2.8	7.1	3.9-12.3	2.4	4.9	2.5-9.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Wayne**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	293.4	409.1	388.0-431.0	149.2	482.2	446.5-519.8	144.2	366.9	340.2-395.2	144.8	208.8	193.6-224.8	82.0	276.8	249.4-306.3	62.8	161.1	143.6-180.1
	1981-1985	319.6	424.3	403.4-445.9	168.8	523.8	487.6-561.9	150.8	363.0	337.1-390.2	165.6	224.2	209.0-240.2	93.0	301.1	273.2-331.0	72.6	173.5	155.9-192.5
	1986-1990	376.8	474.9	453.5-497.0	194.4	584.4	546.8-623.7	182.4	416.4	389.5-444.6	175.4	225.9	211.1-241.4	95.4	302.7	275.0-332.2	80.0	179.5	162.2-198.1
	1991-1995	419.4	498.4	477.2-520.2	224.8	627.8	590.5-666.6	194.6	418.2	392.2-445.5	184.2	222.9	208.7-237.8	97.4	290.4	264.3-318.3	86.8	183.6	166.7-201.8
	1996-2000	463.0	511.2	490.6-532.5	244.2	616.0	581.3-652.3	218.8	439.7	413.9-466.7	181.6	203.0	190.0-216.6	93.4	245.8	223.5-269.6	88.2	174.0	158.1-191.2
	2001-2005	501.8	511.7	491.7-532.2	259.4	586.9	554.7-620.4	242.4	461.4	435.6-488.3	190.8	196.1	183.8-209.0	98.0	234.2	213.5-256.4	92.8	171.4	156.0-187.9
	2006-2010	563.4	524.5	505.0-544.6	305.0	613.9	582.7-646.3	258.4	455.3	430.4-481.4	194.4	181.8	170.5-193.8	104.4	221.9	202.8-242.3	90.0	152.9	139.0-168.0
	2011-2015	598.8	512.4	493.6-531.7	302.6	542.5	514.6-571.5	296.2	492.2	466.6-519.0	206.2	174.7	164.0-186.0	103.6	195.5	178.6-213.6	102.6	161.3	147.3-176.4
	2016-2020	639.4	507.1	488.9-525.9	335.0	548.5	521.4-576.7	304.4	474.6	449.8-500.5	214.8	167.3	157.2-178.0	112.2	190.9	175.0-208.1	102.6	149.2	136.1-163.3
Oral cavity and pharynx	1976-1980	6.0	8.0	5.4-11.5	4.4	13.1	8.1-20.0	1.6	4.0	1.7-7.9	3.0	4.4	2.4-7.2	1.6	5.9	2.4-11.7	1.4	3.4	1.4-7.1
	1981-1985	6.6	8.9	6.1-12.5	4.6	13.9	8.6-21.1	2.0	4.8	2.3-8.8	3.0	3.8	2.1-6.3	2.0	6.2	2.9-11.5	1.0	2.3	0.7-5.4
	1986-1990	9.6	12.0	8.9-16.0	6.4	17.7	12.0-25.2	3.2	7.5	4.2-12.2	2.8	3.6	1.9-6.0	2.4	6.9	3.5-12.2	0.4	0.9	0.1-3.2
	1991-1995	9.4	11.2	8.2-14.9	7.0	19.4	13.3-27.1	2.4	5.2	2.7-9.1	2.6	3.1	1.7-5.3	2.0	5.6	2.6-10.4	0.6	1.3	0.3-3.7
	1996-2000	9.8	10.6	7.9-14.1	6.8	15.6	10.8-21.9	3.0	6.2	3.5-10.2	2.8	3.1	1.7-5.3	2.0	5.5	2.6-10.1	0.8	1.6	0.4-4.1
	2001-2005	10.4	10.4	7.7-13.7	7.4	15.4	10.8-21.4	3.0	6.0	3.4-10.0	1.4	1.4	0.6-2.9	1.2	2.8	1.0-6.1	0.2	0.4	0.0-2.4
	2006-2010	10.2	8.7	6.5-11.6	7.4	13.4	9.3-18.7	2.8	4.6	2.5-7.9	3.4	3.0	1.7-4.8	2.8	5.2	2.8-9.0	0.6	0.8	0.2-2.8
	2011-2015	13.8	11.1	8.6-14.2	9.6	16.2	11.8-21.7	4.2	6.8	4.1-10.6	1.2	0.9	0.3-2.1	1.0	1.6	0.5-4.0	0.2	0.3	0.0-2.2
	2016-2020	16.4	12.0	9.5-15.1	12.2	18.7	14.2-24.4	4.2	5.9	3.6-9.4	2.8	2.0	1.1-3.5	2.2	3.4	1.7-6.5	0.6	0.8	0.2-2.9
Esophagus	1976-1980	2.6	3.9	2.0-6.6	2.2	7.5	3.7-13.5	0.4	1.2	0.1-4.0	1.6	2.3	1.0-4.4	1.4	4.6	1.8-9.4	0.2	0.6	0.0-3.0
	1981-1985	2.8	3.6	2.0-6.1	2.0	6.1	2.9-11.3	0.8	1.7	0.5-4.4	3.2	3.9	2.2-6.4	2.4	6.6	3.4-11.8	0.8	1.7	0.5-4.5
	1986-1990	3.4	4.5	2.6-7.2	2.2	7.3	3.5-13.1	1.2	2.6	0.9-5.6	3.6	4.7	2.8-7.5	3.0	9.2	5.1-15.3	0.6	1.3	0.3-3.8
	1991-1995	3.4	4.0	2.3-6.4	2.4	6.1	3.1-10.8	1.0	2.0	0.7-4.8	6.4	7.6	5.2-10.7	5.2	13.8	8.9-20.4	1.2	2.3	0.9-5.2
	1996-2000	3.2	3.5	2.0-5.7	2.2	5.3	2.6-9.6	1.0	1.9	0.6-4.5	2.8	3.1	1.7-5.2	2.2	5.1	2.5-9.2	0.6	1.1	0.2-3.3
	2001-2005	6.4	6.4	4.4-9.0	4.4	9.8	6.1-15.0	2.0	3.7	1.7-6.9	6.2	6.3	4.3-9.0	4.4	10.1	6.2-15.4	1.8	3.4	1.6-6.6
	2006-2010	6.8	6.2	4.3-8.8	6.2	12.1	8.1-17.3	0.6	0.8	0.2-2.8	5.4	5.2	3.4-7.7	5.2	11.0	7.1-16.2	0.2	0.4	0.0-2.3
	2011-2015	8.4	6.5	4.6-8.9	6.6	10.9	7.4-15.7	1.8	2.7	1.2-5.5	7.4	6.1	4.2-8.5	6.0	10.8	7.2-15.7	1.4	2.0	0.8-4.4
	2016-2020	6.6	5.1	3.4-7.3	6.2	10.3	6.9-15.0	0.4	0.5	0.1-2.3	6.2	4.3	2.9-6.3	5.2	7.6	4.9-11.5	1.0	1.3	0.4-3.5
Stomach	1976-1980	8.6	12.5	9.0-16.9	6.4	21.2	14.2-30.3	2.2	6.0	3.0-10.7	6.0	8.9	6.0-12.8	3.6	12.3	7.1-19.5	2.4	6.3	3.2-10.9
	1981-1985	8.8	12.0	8.6-16.1	5.6	18.9	12.3-27.4	3.2	7.1	4.1-11.6	5.2	7.1	4.6-10.4	3.4	11.4	6.5-18.4	1.8	4.0	1.8-7.7
	1986-1990	7.8	10.3	7.3-14.1	4.8	14.4	9.1-21.5	3.0	7.1	4.0-11.8	5.0	6.5	4.2-9.6	3.0	9.7	5.3-16.0	2.0	4.4	2.1-8.0
	1991-1995	8.4	10.1	7.3-13.7	6.0	17.6	11.6-25.3	2.4	4.9	2.5-8.6	3.8	4.7	2.8-7.3	1.8	6.3	2.8-11.9	2.0	4.2	2.0-7.8
	1996-2000	7.6	8.5	6.0-11.6	5.2	12.8	8.3-18.9	2.4	4.4	2.3-7.8	5.0	5.6	3.6-8.3	3.2	8.1	4.5-13.2	1.8	3.2	1.4-6.2
	2001-2005	6.6	6.8	4.6-9.5	4.2	9.8	6.0-15.1	2.4	4.5	2.3-7.9	3.2	3.2	1.9-5.3	1.6	3.9	1.7-7.8	1.6	2.6	1.1-5.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Wayne**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	7.8	7.4	5.2-10.2	5.2	10.7	6.9-15.8	2.6	4.8	2.5-8.5	4.0	3.8	2.3-5.9	2.6	5.2	2.7-9.1	1.4	2.5	1.0-5.4
	2011-2015	6.2	5.2	3.5-7.5	4.0	7.2	4.4-11.4	2.2	3.6	1.8-6.7	2.4	2.1	1.1-3.8	1.6	3.1	1.3-6.2	0.8	1.4	0.4-3.8
	2016-2020	6.8	5.2	3.5-7.5	3.8	6.2	3.6-10.1	3.0	4.3	2.3-7.6	4.2	3.4	2.0-5.4	2.8	5.1	2.7-8.8	1.4	1.8	0.7-4.2
Colorectal	1976-1980	46.6	65.0	56.7-74.0	22.6	74.7	61.0-90.5	24.0	59.0	48.8-70.7	24.2	35.2	29.1-42.2	13.0	45.5	34.6-58.4	11.2	28.1	21.2-36.5
	1981-1985	49.8	67.8	59.5-76.8	25.4	82.2	68.0-98.4	24.4	58.1	48.1-69.4	22.0	31.5	25.8-38.0	11.2	38.7	28.9-50.6	10.8	26.2	19.6-34.2
	1986-1990	46.4	59.1	51.7-67.3	25.2	77.5	64.0-92.7	21.2	46.6	38.1-56.4	24.4	31.6	26.2-37.7	13.8	44.8	34.5-57.1	10.6	23.4	17.5-30.6
	1991-1995	47.8	57.8	50.7-65.6	20.4	60.8	49.2-74.1	27.4	56.5	47.4-66.8	19.6	24.3	19.7-29.6	9.6	31.2	22.7-41.6	10.0	20.5	15.2-27.1
	1996-2000	66.0	74.2	66.4-82.6	35.0	90.7	77.5-105.4	31.0	60.8	51.6-71.2	23.4	26.5	21.9-31.8	12.6	35.0	26.7-44.9	10.8	21.0	15.8-27.5
	2001-2005	53.2	54.7	48.3-61.7	26.6	60.9	50.8-72.5	26.6	50.0	41.8-59.4	16.8	17.6	14.0-21.8	7.6	18.2	12.8-25.1	9.2	16.9	12.3-22.6
	2006-2010	52.0	48.2	42.4-54.5	24.0	48.6	40.1-58.4	28.0	47.9	40.2-56.7	16.6	15.8	12.6-19.7	8.2	17.3	12.3-23.6	8.4	14.2	10.2-19.4
	2011-2015	47.2	40.7	35.6-46.5	23.0	40.5	33.2-48.9	24.2	41.2	33.9-49.6	17.0	14.0	11.1-17.4	9.4	16.1	11.7-21.7	7.6	12.2	8.5-17.0
2016-2020	52.0	42.8	37.5-48.7	28.8	50.1	41.9-59.5	23.2	36.2	29.6-44.0	16.0	12.7	10.0-16.0	8.8	15.4	11.0-21.0	7.2	10.4	7.1-14.7	
Colon excluding rectum	1976-1980	32.2	44.7	37.9-52.3	14.2	45.9	35.4-58.4	18.0	44.2	35.5-54.4	18.6	26.8	21.5-32.9	10.2	35.0	25.7-46.4	8.4	21.0	15.1-28.4
	1981-1985	34.4	47.4	40.5-55.1	15.6	52.2	40.8-65.6	18.8	45.1	36.4-55.3	17.4	24.7	19.8-30.5	8.4	28.8	20.5-39.2	9.0	21.8	15.8-29.1
	1986-1990	33.0	42.2	35.9-49.2	17.6	55.3	43.9-68.5	15.4	33.7	26.6-42.2	21.4	28.0	22.9-33.8	12.0	39.6	29.8-51.4	9.4	20.9	15.3-27.8
	1991-1995	34.0	41.3	35.3-48.0	13.0	40.0	30.6-51.1	21.0	43.4	35.5-52.6	16.0	19.9	15.8-24.7	7.6	25.1	17.5-34.6	8.4	17.2	12.4-23.3
	1996-2000	47.4	53.4	46.8-60.6	24.0	62.9	51.9-75.4	23.4	45.7	37.7-54.8	16.8	19.1	15.3-23.7	8.4	24.2	17.3-32.7	8.4	16.3	11.7-22.1
	2001-2005	36.0	37.2	31.9-43.1	16.2	38.0	30.0-47.5	19.8	37.3	30.3-45.5	13.6	14.3	11.1-18.1	6.0	14.4	9.6-20.6	7.6	14.1	9.9-19.4
	2006-2010	34.2	31.7	27.1-36.9	14.8	30.9	24.1-39.0	19.4	32.8	26.5-40.1	11.2	10.8	8.2-14.1	5.0	10.9	7.0-16.2	6.2	10.5	7.1-15.1
	2011-2015	34.8	30.2	25.8-35.2	16.8	30.5	24.1-38.1	18.0	30.1	24.0-37.4	12.8	10.5	8.1-13.6	6.6	11.3	7.7-16.2	6.2	10.0	6.7-14.4
2016-2020	35.0	29.4	25.0-34.4	18.6	33.7	26.9-41.9	16.4	25.4	20.0-31.9	11.2	9.3	6.9-12.3	5.4	9.9	6.4-14.8	5.8	8.6	5.6-12.7	
Rectum & rectosigmoid	1976-1980	14.4	20.3	15.8-25.6	8.4	28.8	20.4-39.2	6.0	14.8	9.9-21.2	5.6	8.4	5.6-12.2	2.8	10.5	5.5-17.7	2.8	7.1	3.9-11.9
	1981-1985	15.4	20.4	16.0-25.5	9.8	30.0	21.9-39.9	5.6	13.0	8.6-18.8	4.6	6.7	4.3-10.1	2.8	9.9	5.3-16.6	1.8	4.4	2.0-8.4
	1986-1990	13.4	16.9	13.1-21.5	7.6	22.2	15.6-30.6	5.8	12.9	8.6-18.6	3.0	3.6	2.0-5.9	1.8	5.2	2.3-9.9	1.2	2.5	0.9-5.5
	1991-1995	13.8	16.5	12.8-20.8	7.4	20.8	14.5-28.9	6.4	13.1	8.9-18.5	3.6	4.4	2.6-6.9	2.0	6.1	2.8-11.2	1.6	3.3	1.4-6.6
	1996-2000	18.6	20.8	16.8-25.5	11.0	27.8	20.8-36.4	7.6	15.1	10.7-20.8	6.6	7.4	5.1-10.4	4.2	10.8	6.6-16.6	2.4	4.7	2.4-8.3
	2001-2005	17.2	17.5	14.0-21.6	10.4	22.9	17.0-30.2	6.8	12.7	8.8-17.9	3.2	3.3	1.9-5.3	1.6	3.9	1.6-7.7	1.6	2.8	1.2-5.7
	2006-2010	17.8	16.4	13.2-20.3	9.2	17.7	12.9-23.9	8.6	15.1	10.9-20.5	5.4	5.0	3.3-7.4	3.2	6.4	3.6-10.6	2.2	3.7	1.8-6.8
	2011-2015	12.4	10.5	8.0-13.7	6.2	10.0	6.7-14.5	6.2	11.0	7.3-16.0	4.2	3.4	2.1-5.3	2.8	4.8	2.6-8.3	1.4	2.2	0.9-4.8
2016-2020	17.0	13.4	10.6-16.8	10.2	16.3	12.0-21.8	6.8	10.9	7.3-15.7	4.8	3.4	2.2-5.3	3.4	5.4	3.1-9.0	1.4	1.8	0.7-4.1	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Wayne**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	1.8	2.6	1.2-4.9	1.2	3.7	1.3-8.0	0.6	1.5	0.3-4.5	1.6	2.2	1.0-4.4	1.0	3.1	1.0-7.2	0.6	1.5	0.3-4.4
	1981-1985	2.8	3.7	2.0-6.2	1.8	5.5	2.4-10.4	1.0	2.3	0.7-5.3	2.2	3.0	1.5-5.4	1.4	5.1	2.0-10.3	0.8	1.8	0.5-4.6
	1986-1990	2.4	3.1	1.6-5.5	2.0	5.9	2.8-10.8	0.4	0.9	0.1-3.2	1.6	2.1	0.9-4.1	1.4	4.9	1.9-10.0	0.2	0.4	0.0-2.4
	1991-1995	2.4	2.8	1.4-4.9	1.4	4.0	1.6-8.2	1.0	2.1	0.7-4.9	3.0	3.5	2.0-5.8	1.6	4.5	1.9-8.9	1.4	2.9	1.2-6.1
	1996-2000	3.2	3.5	2.0-5.7	2.2	5.5	2.7-10.0	1.0	2.0	0.7-4.8	3.8	4.2	2.6-6.6	1.8	4.6	2.1-8.8	2.0	4.0	1.9-7.4
	2001-2005	4.8	5.1	3.3-7.6	3.2	7.3	4.1-11.9	1.6	3.1	1.4-6.3	3.4	3.6	2.1-5.8	2.6	6.2	3.3-10.7	0.8	1.5	0.4-3.9
	2006-2010	6.2	5.7	3.8-8.2	4.2	8.0	4.9-12.5	2.0	3.5	1.6-6.5	5.4	4.9	3.2-7.3	3.2	6.2	3.5-10.2	2.2	3.7	1.8-6.8
	2011-2015	7.0	5.4	3.7-7.7	5.0	7.9	5.1-12.0	2.0	3.0	1.4-5.8	4.4	3.4	2.1-5.4	2.2	3.6	1.7-6.7	2.2	3.3	1.6-6.2
	2016-2020	9.2	6.8	4.9-9.2	6.8	10.3	7.0-14.8	2.4	3.5	1.8-6.5	4.8	3.5	2.2-5.4	3.8	5.9	3.5-9.6	1.0	1.3	0.4-3.5
Pancreas	1976-1980	9.0	12.6	9.1-16.9	5.2	17.1	10.9-25.3	3.8	9.2	5.5-14.4	8.6	12.1	8.7-16.4	5.4	18.1	11.6-26.6	3.2	7.7	4.4-12.6
	1981-1985	6.4	8.8	6.0-12.4	3.2	8.6	4.9-14.1	3.2	8.0	4.5-12.9	5.4	7.4	4.8-10.7	2.6	7.2	3.8-12.3	2.8	6.8	3.7-11.4
	1986-1990	7.8	10.3	7.3-14.1	2.4	7.5	3.8-13.2	5.4	12.0	7.9-17.5	8.4	11.5	8.3-15.6	2.4	7.4	3.7-13.1	6.0	13.5	9.1-19.3
	1991-1995	10.6	12.8	9.6-16.7	6.2	17.5	11.7-25.1	4.4	9.2	5.7-14.0	9.4	11.4	8.4-15.2	5.0	14.4	9.2-21.4	4.4	9.2	5.8-14.0
	1996-2000	9.0	10.1	7.3-13.5	4.6	12.2	7.6-18.3	4.4	8.6	5.4-13.1	9.8	11.0	8.1-14.5	4.6	12.1	7.6-18.1	5.2	10.1	6.6-14.9
	2001-2005	14.2	14.7	11.5-18.6	6.2	14.5	9.8-20.6	8.0	14.2	10.1-19.5	11.4	11.7	8.9-15.2	6.0	14.3	9.6-20.5	5.4	9.6	6.3-14.1
	2006-2010	16.0	14.9	11.8-18.7	7.8	15.5	10.9-21.4	8.2	14.1	10.1-19.3	12.6	11.7	9.0-15.1	6.4	12.7	8.6-18.1	6.2	10.5	7.1-15.1
	2011-2015	12.0	10.0	7.6-13.0	5.2	9.7	6.2-14.4	6.8	10.2	7.0-14.6	10.2	8.7	6.5-11.6	4.6	9.3	5.8-14.1	5.6	8.3	5.5-12.3
	2016-2020	17.6	14.0	11.2-17.5	8.6	14.8	10.6-20.4	9.0	13.4	9.7-18.2	15.4	12.4	9.7-15.6	7.4	12.4	8.6-17.5	8.0	12.2	8.6-16.9
Larynx	1976-1980	3.8	5.0	3.0-7.9	3.8	11.0	6.5-17.4	0.0	0.0	0.0-1.8	1.6	2.1	0.9-4.1	1.6	4.3	1.8-8.6	0.0	0.0	0.0-1.8
	1981-1985	5.6	6.9	4.6-10.1	4.2	12.1	7.4-18.6	1.4	3.2	1.3-6.7	1.6	2.1	0.9-4.0	1.4	4.4	1.7-9.0	0.2	0.4	0.0-2.4
	1986-1990	4.6	5.8	3.7-8.7	3.2	9.1	5.1-14.9	1.4	3.2	1.3-6.7	1.4	1.8	0.7-3.7	0.8	3.0	0.8-7.5	0.6	1.5	0.3-4.2
	1991-1995	5.2	6.0	3.9-8.8	4.2	10.3	6.4-15.8	1.0	2.1	0.7-4.9	1.8	2.2	1.0-4.2	1.6	4.7	2.0-9.3	0.2	0.5	0.0-2.7
	1996-2000	3.4	3.7	2.1-5.9	2.6	6.1	3.2-10.5	0.8	1.7	0.5-4.3	1.6	1.8	0.8-3.6	1.2	3.1	1.1-6.7	0.4	0.7	0.1-2.8
	2001-2005	6.2	6.2	4.2-8.8	4.8	9.9	6.3-14.8	1.4	2.7	1.1-5.7	2.8	2.9	1.6-4.9	1.8	3.9	1.8-7.5	1.0	1.9	0.6-4.6
	2006-2010	4.2	3.8	2.3-5.8	3.4	6.6	3.8-10.8	0.8	1.3	0.4-3.6	1.4	1.4	0.6-2.9	1.2	2.9	1.0-6.3	0.2	0.4	0.0-2.3
	2011-2015	3.4	2.7	1.6-4.5	2.6	4.4	2.3-7.8	0.8	1.2	0.3-3.4	1.8	1.4	0.6-2.8	1.0	1.6	0.5-4.1	0.8	1.0	0.3-3.1
	2016-2020	4.0	3.0	1.8-4.9	3.4	5.4	3.0-9.0	0.6	0.8	0.2-2.9	1.4	1.1	0.4-2.4	1.2	2.0	0.7-4.7	0.2	0.2	0.0-2.0
Lung and bronchus	1976-1980	37.4	50.5	43.4-58.5	28.8	90.3	75.6-107.0	8.6	20.3	14.6-27.4	31.8	43.4	36.8-50.9	25.6	80.4	66.5-96.3	6.2	15.1	10.2-21.4
	1981-1985	55.0	71.3	63.0-80.3	41.6	122.4	105.8-140.8	13.4	33.1	25.5-42.2	45.4	58.9	51.4-67.2	34.8	106.5	90.7-124.1	10.6	24.3	18.1-31.9
	1986-1990	58.8	72.3	64.2-81.1	40.8	116.6	100.7-134.1	18.0	39.8	32.0-49.1	49.8	62.6	55.0-70.9	34.4	102.9	87.6-120.0	15.4	34.3	27.0-43.0
	1991-1995	64.0	76.5	68.3-85.3	37.4	103.2	88.6-119.4	26.6	57.6	48.2-68.4	51.0	60.9	53.6-68.9	29.6	83.2	70.1-98.1	21.4	45.7	37.4-55.3
	1996-2000	60.8	67.2	59.9-75.3	38.2	94.1	81.0-108.7	22.6	46.0	37.8-55.3	45.4	50.5	44.1-57.5	26.2	65.0	54.1-77.3	19.2	39.0	31.6-47.7
	2001-2005	74.6	75.6	68.1-83.8	39.6	87.3	75.3-100.6	35.0	66.1	56.7-76.8	56.4	57.6	51.1-64.8	32.2	72.9	61.9-85.3	24.2	45.6	37.8-54.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Wayne**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	79.2	73.3	66.2-81.0	41.2	85.1	73.6-97.9	38.0	65.7	56.6-75.9	59.8	55.3	49.1-62.0	32.2	67.3	57.0-78.8	27.6	47.3	39.7-56.1
	2011-2015	88.8	74.3	67.4-81.7	40.8	74.2	64.1-85.5	48.0	75.8	66.4-86.4	62.2	52.4	46.7-58.8	30.2	56.0	47.2-66.0	32.0	51.1	43.4-59.9
	2016-2020	96.0	72.4	65.9-79.5	48.2	76.7	67.0-87.5	47.8	68.6	60.0-78.3	59.4	45.5	40.4-51.2	31.0	50.7	42.8-59.8	28.4	40.8	34.3-48.4
Melanoma of the skin	1976-1980	5.4	7.1	4.7-10.5	3.2	9.2	5.1-15.3	2.2	5.9	2.9-10.6	1.2	1.6	0.6-3.4	0.8	2.0	0.5-5.4	0.4	1.0	0.1-3.7
	1981-1985	4.4	5.8	3.6-8.8	2.2	6.0	3.0-10.8	2.2	5.5	2.7-9.9	3.0	3.7	2.1-6.2	2.0	5.1	2.4-9.5	1.0	2.2	0.7-5.2
	1986-1990	6.8	8.6	6.0-12.1	2.8	7.6	4.1-12.9	4.0	9.6	5.8-14.9	1.8	2.4	1.1-4.5	1.0	2.8	0.9-6.5	0.8	1.9	0.5-4.9
	1991-1995	7.8	9.2	6.5-12.6	3.8	10.1	6.0-16.0	4.0	8.8	5.3-13.6	1.6	1.9	0.8-3.7	0.4	0.8	0.1-3.2	1.2	2.7	1.0-5.8
	1996-2000	12.8	14.1	10.9-18.1	8.0	19.9	14.1-27.1	4.8	10.0	6.4-14.9	2.4	2.6	1.4-4.6	1.0	2.7	0.8-6.2	1.4	2.9	1.2-5.9
	2001-2005	11.6	11.7	8.8-15.2	5.4	12.4	8.0-18.1	6.2	12.2	8.2-17.4	3.6	3.8	2.2-6.0	2.2	5.3	2.6-9.5	1.4	2.7	1.1-5.7
	2006-2010	21.4	20.9	17.1-25.4	11.8	24.5	18.5-31.8	9.6	18.6	13.6-24.8	2.2	2.1	1.0-3.8	1.6	3.2	1.3-6.4	0.6	1.0	0.2-3.1
	2011-2015	25.0	22.7	18.8-27.2	13.0	24.3	18.6-31.2	12.0	22.1	16.7-28.8	4.6	3.6	2.3-5.6	2.6	4.5	2.4-8.0	2.0	2.9	1.4-5.8
2016-2020	34.0	30.1	25.5-35.3	19.0	33.1	26.4-41.0	15.0	27.8	21.5-35.4	3.2	2.4	1.4-4.1	1.6	3.1	1.3-6.3	1.6	2.1	0.9-4.6	
Female breast	1976-1980							35.8	92.8	79.5-107.6							10.6	27.5	20.5-36.0
	1981-1985							43.6	107.5	93.5-123.0							14.8	36.2	28.3-45.6
	1986-1990							58.0	136.5	121.0-153.3							15.2	35.6	28.0-44.7
	1991-1995							56.6	125.2	110.9-140.7							14.2	30.8	24.0-38.9
	1996-2000							71.0	143.9	129.3-159.7							16.4	32.5	25.8-40.4
	2001-2005							68.6	129.9	116.4-144.5							13.4	24.2	18.8-30.9
	2006-2010							72.2	125.6	112.8-139.5							12.6	21.7	16.6-27.9
	2011-2015							87.4	146.7	132.8-161.8							13.0	20.4	15.7-26.3
2016-2020							88.0	140.8	127.2-155.5							12.0	19.2	14.4-25.3	
Cervix uteri	1976-1980							7.0	18.6	12.8-25.9							1.6	4.4	1.9-8.7
	1981-1985							4.8	12.1	7.7-18.1							1.2	2.8	1.0-6.2
	1986-1990							5.2	11.8	7.7-17.4							1.6	3.8	1.6-7.5
	1991-1995							5.4	11.6	7.6-16.9							1.6	3.6	1.5-7.0
	1996-2000							4.0	8.4	5.1-13.1							1.2	2.4	0.9-5.3
	2001-2005							3.8	7.5	4.5-11.9							2.2	4.1	2.0-7.5
	2006-2010							3.4	7.1	4.0-11.5							1.6	2.9	1.2-6.0
	2011-2015							6.6	14.4	9.7-20.5							2.4	4.6	2.3-8.3
	2016-2020							2.8	5.6	2.9-9.8							1.2	2.9	1.0-6.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Wayne**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							15.4	38.8	30.5-48.7							1.4	3.7	1.5-7.7
	1981-1985							11.4	27.3	20.6-35.4							3.8	9.6	5.7-14.9
	1986-1990							12.8	29.3	22.5-37.5							1.8	4.0	1.8-7.7
	1991-1995							10.6	22.6	16.9-29.6							3.0	6.1	3.4-10.1
	1996-2000							14.0	28.8	22.4-36.4							2.2	4.5	2.2-8.0
	2001-2005							14.2	26.5	20.7-33.5							2.4	4.3	2.2-7.7
	2006-2010							15.0	25.8	20.2-32.6							3.0	4.9	2.7-8.2
	2011-2015							21.8	32.3	26.4-39.3							1.8	2.6	1.2-5.3
	2016-2020							24.2	35.9	29.6-43.3							4.6	6.2	3.9-9.6
Ovary	1976-1980							7.4	20.4	14.3-28.2							4.6	12.6	7.9-18.9
	1981-1985							7.4	18.0	12.6-24.9							4.2	9.8	6.1-15.1
	1986-1990							8.0	18.2	13.0-24.9							5.4	12.0	7.9-17.5
	1991-1995							7.8	16.8	11.9-23.0							4.4	9.6	6.0-14.5
	1996-2000							6.4	13.2	9.0-18.6							3.8	7.7	4.7-12.1
	2001-2005							10.2	19.5	14.4-25.7							3.8	7.0	4.2-11.1
	2006-2010							7.0	12.5	8.6-17.6							4.0	6.9	4.2-10.9
	2011-2015							8.4	13.3	9.4-18.3							5.6	8.3	5.4-12.2
	2016-2020							9.0	13.7	9.8-18.7							5.8	8.5	5.6-12.6
Prostate	1976-1980				23.8	85.8	70.4-103.4							7.8	31.2	21.8-42.7			
	1981-1985				25.6	88.6	73.3-105.9							7.6	31.8	22.1-43.7			
	1986-1990				42.6	140.4	121.5-161.2							12.0	44.7	33.7-57.8			
	1991-1995				70.4	203.3	182.1-226.2							13.0	45.6	34.8-58.3			
	1996-2000				65.6	170.3	152.1-190.2							10.4	31.9	23.7-41.9			
	2001-2005				79.8	181.3	163.6-200.4							10.2	29.2	21.6-38.3			
	2006-2010				97.4	186.9	170.3-204.8							9.4	22.7	16.6-30.3			
	2011-2015				88.0	146.1	132.4-160.9							11.0	22.7	17.0-29.7			
	2016-2020				90.0	133.9	121.5-147.4							9.6	17.4	12.7-23.3			
Testis	1976-1980				1.4	3.2	1.3-6.9							0.6	1.6	0.3-4.9			
	1981-1985				2.0	4.5	2.1-8.5							0.0	0.0	0.0-2.1			
	1986-1990				2.6	5.8	3.0-10.2							0.0	0.0	0.0-2.0			
	1991-1995				2.0	4.3	2.0-8.1							0.2	0.5	0.0-2.9			
	1996-2000				3.4	7.5	4.3-12.1							0.6	1.6	0.3-4.7			
	2001-2005				2.0	4.4	2.1-8.2							0.0	0.0	0.0-1.7			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Wayne**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				3.4	7.6	4.4-12.3							0.2	0.3	0.0-2.2			
	2011-2015				3.8	9.0	5.4-14.1							0.0	0.0	0.0-1.7			
	2016-2020				3.4	9.1	5.2-14.6							0.8	2.5	0.7-6.0			
Urinary bladder (incl. in situ)	1976-1980	19.8	27.2	22.0-33.2	16.0	49.7	38.9-62.4	3.8	9.9	5.9-15.4	5.4	8.2	5.4-12.0	3.6	12.8	7.4-20.4	1.8	5.0	2.3-9.3
	1981-1985	17.6	23.2	18.6-28.7	14.0	44.7	34.3-57.0	3.6	8.3	4.9-13.3	3.8	5.5	3.3-8.5	2.6	10.2	5.2-17.3	1.2	3.0	1.1-6.4
	1986-1990	21.0	26.8	21.9-32.5	16.4	51.6	40.6-64.4	4.6	10.2	6.5-15.4	5.2	7.0	4.5-10.2	4.0	14.6	8.8-22.6	1.2	2.5	0.9-5.6
	1991-1995	23.2	28.1	23.2-33.6	16.2	45.7	36.0-57.1	7.0	14.5	10.0-20.2	6.6	8.1	5.6-11.4	4.2	13.9	8.4-21.3	2.4	4.7	2.4-8.3
	1996-2000	23.6	26.4	21.8-31.6	17.0	45.9	36.5-57.0	6.6	12.8	8.8-18.0	4.8	5.4	3.4-8.0	3.4	9.4	5.4-15.1	1.4	2.6	1.0-5.5
	2001-2005	30.2	31.5	26.6-36.9	22.8	55.2	45.3-66.5	7.4	14.0	9.8-19.4	5.0	5.3	3.4-7.8	3.2	8.9	5.0-14.4	1.8	2.9	1.3-5.8
	2006-2010	32.2	30.4	25.9-35.6	23.0	48.4	39.8-58.4	9.2	16.2	11.8-21.8	3.6	3.6	2.1-5.8	3.2	7.3	4.2-12.0	0.4	0.6	0.1-2.4
	2011-2015	34.0	28.9	24.7-33.7	25.0	47.5	39.3-56.9	9.0	14.3	10.3-19.4	6.8	5.9	4.1-8.4	4.8	9.8	6.2-14.8	2.0	3.3	1.5-6.2
2016-2020	32.0	25.0	21.2-29.4	26.4	45.0	37.4-53.7	5.6	8.0	5.3-11.8	8.4	6.2	4.5-8.6	6.0	10.7	7.2-15.5	2.4	3.0	1.6-5.7	
Kidney and renal pelvis	1976-1980	6.8	9.8	6.8-13.8	3.2	10.9	6.1-17.9	3.6	9.2	5.4-14.5	2.8	4.2	2.3-7.1	1.4	4.9	1.9-10.0	1.4	3.9	1.5-7.9
	1981-1985	5.8	7.9	5.3-11.4	3.8	12.4	7.3-19.6	2.0	4.8	2.3-8.9	3.0	4.2	2.3-6.9	2.0	7.0	3.2-13.0	1.0	2.4	0.8-5.6
	1986-1990	10.0	12.7	9.4-16.7	6.2	17.9	12.0-25.5	3.8	8.7	5.2-13.7	3.6	4.6	2.7-7.2	2.2	6.2	3.1-11.1	1.4	3.0	1.2-6.3
	1991-1995	11.2	13.5	10.2-17.5	7.2	20.2	14.0-28.2	4.0	8.7	5.3-13.5	6.0	7.4	5.0-10.5	3.6	10.4	6.0-16.4	2.4	5.4	2.8-9.4
	1996-2000	11.8	12.9	9.8-16.6	9.0	20.6	15.0-27.7	2.8	5.5	3.0-9.3	3.6	4.0	2.4-6.4	2.4	5.8	3.0-10.2	1.2	2.4	0.9-5.3
	2001-2005	13.4	13.5	10.5-17.2	7.2	15.8	11.0-22.1	6.2	11.9	8.1-17.0	2.8	2.8	1.5-4.7	1.4	3.0	1.2-6.3	1.4	2.5	1.0-5.4
	2006-2010	20.0	19.1	15.5-23.3	12.8	26.7	20.4-34.3	7.2	12.8	8.9-17.8	3.2	3.0	1.7-4.9	2.2	4.4	2.1-8.0	1.0	1.9	0.6-4.6
	2011-2015	26.8	23.4	19.5-27.8	16.2	29.3	23.1-36.8	10.6	17.7	13.1-23.4	3.8	3.2	1.9-5.1	2.6	5.1	2.7-8.9	1.2	1.8	0.7-4.3
2016-2020	28.4	22.1	18.4-26.3	17.4	28.4	22.4-35.5	11.0	16.4	12.2-21.7	5.0	3.7	2.4-5.6	3.2	4.9	2.7-8.3	1.8	2.7	1.2-5.5	
Brain and other central nervous system	1976-1980	3.6	4.4	2.6-7.1	2.0	5.3	2.5-10.1	1.6	3.8	1.6-7.6	1.8	2.4	1.1-4.5	1.2	3.9	1.4-8.6	0.6	1.3	0.3-3.8
	1981-1985	4.6	5.2	3.3-7.9	2.0	4.8	2.3-9.1	2.6	5.6	2.9-9.7	3.0	3.8	2.1-6.3	1.4	3.7	1.4-7.7	1.6	3.8	1.6-7.5
	1986-1990	5.0	5.9	3.8-8.7	3.2	8.3	4.7-13.6	1.8	3.8	1.7-7.2	3.0	3.5	1.9-5.8	1.2	3.5	1.2-7.6	1.8	3.7	1.7-7.1
	1991-1995	6.6	7.5	5.1-10.5	3.8	8.9	5.3-14.0	2.8	5.9	3.2-9.9	3.8	4.4	2.6-6.9	2.0	4.8	2.3-9.0	1.8	3.7	1.7-7.1
	1996-2000	6.8	7.2	5.0-10.1	4.0	9.2	5.5-14.3	2.8	5.7	3.1-9.6	4.4	4.7	3.0-7.2	2.0	4.6	2.2-8.5	2.4	4.8	2.5-8.5
	2001-2005	6.4	6.6	4.5-9.4	2.4	5.0	2.5-8.9	4.0	7.9	4.8-12.2	3.6	3.6	2.1-5.7	1.8	3.6	1.6-6.9	1.8	3.5	1.6-6.8
	2006-2010	7.4	7.3	5.1-10.2	3.6	8.2	4.8-13.0	3.8	6.9	4.1-10.9	5.4	5.0	3.3-7.3	3.0	6.0	3.3-10.1	2.4	4.1	2.1-7.3
	2011-2015	8.8	8.5	6.1-11.6	6.2	12.4	8.2-17.9	2.6	4.8	2.5-8.5	8.0	7.8	5.5-10.8	4.4	9.1	5.6-14.1	3.6	6.6	3.8-10.7
2016-2020	7.2	5.9	4.1-8.4	3.8	6.1	3.5-9.9	3.4	5.6	3.2-9.3	5.8	4.5	3.0-6.7	3.8	5.8	3.4-9.4	2.0	3.5	1.6-6.7	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Wayne**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	1.8	2.2	1.0-4.2	0.4	0.9	0.1-3.6	1.4	3.3	1.3-6.9	0.2	0.3	0.0-1.5	0.0	0.0	0.0-2.2	0.2	0.5	0.0-2.8
	1981-1985	1.6	2.1	0.9-4.2	0.8	2.4	0.6-6.2	0.8	1.9	0.5-5.0	0.4	0.5	0.1-1.9	0.2	0.4	0.0-2.7	0.2	0.5	0.0-2.7
	1986-1990	2.6	3.0	1.6-5.1	1.0	2.5	0.8-5.9	1.6	3.6	1.5-7.1	0.6	0.7	0.1-2.0	0.4	1.1	0.1-3.9	0.2	0.4	0.0-2.4
	1991-1995	3.4	3.7	2.2-6.0	0.8	1.7	0.5-4.6	2.6	5.5	2.9-9.5	0.0	0.0	0.0-0.8	0.0	0.0	0.0-1.9	0.0	0.0	0.0-1.6
	1996-2000	4.6	5.0	3.2-7.6	2.0	4.7	2.2-8.8	2.6	5.6	3.0-9.6	0.4	0.5	0.1-1.6	0.2	0.6	0.0-3.1	0.2	0.4	0.0-2.2
	2001-2005	5.2	5.5	3.6-8.1	1.6	3.3	1.4-6.5	3.6	7.8	4.5-12.3	0.0	0.0	0.0-0.8	0.0	0.0	0.0-1.7	0.0	0.0	0.0-1.6
	2006-2010	9.2	9.4	6.8-12.6	2.6	5.3	2.8-9.3	6.6	13.5	9.2-19.2	0.4	0.4	0.0-1.4	0.4	0.8	0.1-3.0	0.0	0.0	0.0-1.6
	2011-2015	11.0	10.5	7.8-13.9	4.0	7.3	4.4-11.5	7.0	13.8	9.4-19.5	0.8	0.6	0.2-1.8	0.4	0.7	0.1-2.8	0.4	0.5	0.1-2.4
2016-2020	11.2	10.9	8.0-14.5	3.4	7.4	4.1-12.2	7.8	14.5	10.0-20.4	0.6	0.5	0.1-1.6	0.4	0.8	0.1-3.1	0.2	0.3	0.0-2.1	
Hodgkin lymphoma	1976-1980	2.4	2.8	1.4-5.0	1.4	4.0	1.5-8.5	1.0	1.9	0.6-4.7	0.4	0.7	0.1-2.2	0.4	1.5	0.2-5.2	0.0	0.0	0.0-1.8
	1981-1985	3.8	4.1	2.5-6.5	1.6	3.3	1.4-6.8	2.2	4.9	2.4-8.9	1.0	1.2	0.4-2.9	0.8	2.0	0.5-5.3	0.2	0.5	0.0-2.6
	1986-1990	3.0	3.6	2.0-5.9	1.2	2.6	1.0-5.9	1.8	4.6	2.1-8.7	0.4	0.5	0.1-1.9	0.2	0.5	0.0-2.8	0.2	0.6	0.0-3.1
	1991-1995	3.2	3.6	2.1-5.9	1.8	4.6	2.0-9.0	1.4	3.1	1.2-6.4	0.2	0.2	0.0-1.3	0.0	0.0	0.0-1.9	0.2	0.5	0.0-2.5
	1996-2000	2.2	2.7	1.3-4.8	1.2	3.0	1.1-6.5	1.0	2.2	0.7-5.2	0.6	0.7	0.1-1.9	0.4	0.9	0.1-3.2	0.2	0.3	0.0-2.1
	2001-2005	4.2	4.9	3.0-7.4	2.6	6.1	3.2-10.6	1.6	3.4	1.4-6.7	0.0	0.0	0.0-0.8	0.0	0.0	0.0-1.7	0.0	0.0	0.0-1.6
	2006-2010	4.4	5.1	3.2-7.7	2.0	4.6	2.2-8.5	2.4	5.5	2.8-9.7	0.2	0.2	0.0-1.2	0.0	0.0	0.0-1.7	0.2	0.3	0.0-2.2
	2011-2015	3.8	4.3	2.6-6.7	2.6	5.6	2.9-9.6	1.2	3.1	1.1-6.6	0.4	0.5	0.1-1.7	0.0	0.0	0.0-1.7	0.4	0.9	0.1-3.4
2016-2020	2.2	2.5	1.2-4.6	1.0	2.3	0.7-5.4	1.2	2.7	0.9-6.1	0.2	0.2	0.0-1.2	0.2	0.4	0.0-2.4	0.0	0.0	0.0-1.7	
Non-Hodgkin lymphomas	1976-1980	11.2	15.1	11.3-19.7	6.4	19.3	12.9-27.6	4.8	11.7	7.4-17.4	5.4	8.3	5.4-12.1	2.2	7.9	3.8-14.1	3.2	8.4	4.8-13.5
	1981-1985	10.8	13.7	10.2-17.9	5.8	16.6	10.9-24.2	5.0	11.4	7.3-16.9	6.2	8.0	5.4-11.3	4.2	13.3	8.0-20.6	2.0	4.5	2.1-8.3
	1986-1990	17.4	21.7	17.4-26.8	9.0	25.4	18.3-34.3	8.4	19.4	13.9-26.2	6.2	7.6	5.2-10.9	3.2	9.1	5.1-15.0	3.0	6.6	3.7-10.9
	1991-1995	18.0	20.9	16.8-25.6	9.4	23.2	16.9-30.9	8.6	18.9	13.7-25.5	9.0	10.8	7.9-14.5	4.0	11.0	6.6-17.2	5.0	10.7	6.9-15.8
	1996-2000	21.0	23.3	19.1-28.3	11.0	28.5	21.3-37.3	10.0	20.6	15.3-27.2	8.0	9.0	6.4-12.2	4.8	13.3	8.4-19.8	3.2	6.4	3.7-10.5
	2001-2005	22.8	23.3	19.2-28.0	11.8	25.0	18.9-32.4	11.0	21.4	16.1-27.9	8.6	9.0	6.5-12.1	4.6	10.6	6.7-16.0	4.0	7.5	4.5-11.6
	2006-2010	24.6	22.7	18.8-27.1	13.6	27.3	21.1-34.8	11.0	19.1	14.3-25.1	8.2	7.8	5.6-10.6	4.6	9.9	6.2-15.0	3.6	6.1	3.6-9.9
	2011-2015	23.0	19.8	16.2-23.9	12.0	21.7	16.4-28.3	11.0	17.5	13.1-23.2	7.2	5.6	3.9-7.9	3.4	5.6	3.2-9.2	3.8	5.2	3.1-8.4
2016-2020	30.0	24.7	20.7-29.2	17.0	29.0	22.9-36.3	13.0	20.9	15.9-27.2	7.0	5.7	3.9-8.0	3.0	5.2	2.9-8.9	4.0	6.0	3.6-9.6	
Myeloma	1976-1980	2.4	3.5	1.8-6.1	1.0	3.0	0.9-7.2	1.4	3.5	1.4-7.3	3.2	4.4	2.5-7.2	1.6	4.9	2.0-9.7	1.6	4.0	1.7-7.9
	1981-1985	3.4	4.8	2.8-7.6	2.4	8.1	4.0-14.1	1.0	2.8	0.9-6.4	1.4	2.0	0.8-4.0	0.8	2.6	0.6-6.7	0.6	1.7	0.3-4.8
	1986-1990	3.8	5.0	3.0-7.7	1.8	5.0	2.3-9.5	2.0	4.3	2.1-8.0	2.0	2.8	1.3-5.1	0.8	2.9	0.8-7.1	1.2	2.6	1.0-5.7
	1991-1995	3.6	4.2	2.5-6.7	2.6	7.4	3.8-12.7	1.0	2.3	0.7-5.3	2.8	3.3	1.8-5.5	2.2	5.9	2.9-10.6	0.6	1.2	0.3-3.6
	1996-2000	5.6	6.3	4.2-9.1	3.0	7.4	4.1-12.2	2.6	5.1	2.7-8.8	3.6	4.1	2.4-6.4	1.8	5.1	2.3-9.6	1.8	3.5	1.6-6.7
	2001-2005	6.4	6.7	4.6-9.5	4.0	9.5	5.8-14.7	2.4	4.4	2.3-7.8	4.0	4.2	2.6-6.5	2.6	6.1	3.2-10.5	1.4	2.7	1.1-5.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Wayne**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	5.6	4.9	3.2-7.2	2.2	4.2	2.0-7.7	3.4	5.4	3.1-8.8	2.2	2.0	1.0-3.7	1.2	2.6	0.9-5.7	1.0	1.6	0.5-4.0
	2011-2015	8.6	7.4	5.3-10.1	4.8	9.3	5.9-14.2	3.8	5.7	3.4-9.1	4.2	3.6	2.2-5.7	2.6	5.2	2.8-9.1	1.6	2.4	1.0-5.0
	2016-2020	9.0	7.2	5.1-9.8	5.6	9.6	6.2-14.3	3.4	4.7	2.7-7.9	3.4	2.7	1.6-4.5	1.6	2.7	1.1-5.6	1.8	2.7	1.2-5.4
Leukemias	1976-1980	12.0	16.7	12.7-21.6	5.8	18.9	12.3-27.6	6.2	15.6	10.5-22.2	5.4	7.9	5.1-11.5	2.8	9.6	4.9-16.4	2.6	6.8	3.6-11.6
	1981-1985	10.0	13.7	10.1-18.1	6.6	22.5	15.3-31.6	3.4	7.7	4.4-12.3	8.0	11.3	8.0-15.4	5.0	17.0	10.8-25.3	3.0	7.2	4.0-11.9
	1986-1990	10.6	12.8	9.6-16.8	6.6	18.1	12.2-25.8	4.0	8.6	5.2-13.3	5.2	6.6	4.3-9.7	3.6	11.0	6.3-17.5	1.6	3.4	1.5-6.8
	1991-1995	10.8	12.7	9.5-16.6	6.6	17.9	12.1-25.4	4.2	8.7	5.4-13.3	5.8	7.1	4.7-10.2	3.4	9.7	5.5-15.6	2.4	4.9	2.5-8.6
	1996-2000	17.6	19.3	15.4-23.7	10.2	25.0	18.5-33.0	7.4	14.6	10.3-20.2	8.6	9.7	7.0-13.1	4.6	12.0	7.5-18.0	4.0	7.7	4.7-12.0
	2001-2005	11.8	12.4	9.4-16.0	7.4	16.4	11.5-22.8	4.4	9.2	5.7-14.1	6.8	6.9	4.7-9.6	4.2	9.3	5.7-14.4	2.6	4.6	2.4-8.0
	2006-2010	17.2	16.6	13.2-20.6	10.6	22.1	16.5-29.1	6.6	11.8	8.1-16.8	6.8	6.4	4.4-9.0	3.6	7.9	4.6-12.7	3.2	5.4	3.1-9.0
	2011-2015	18.4	16.5	13.2-20.4	11.6	22.2	16.7-29.0	6.8	11.2	7.6-16.0	8.0	7.4	5.2-10.2	4.2	8.3	5.1-12.9	3.8	6.2	3.6-10.1
	2016-2020	20.6	17.0	13.8-20.9	11.4	20.3	15.2-26.7	9.2	13.8	9.9-19.0	8.4	6.5	4.7-9.0	4.4	8.3	5.2-12.8	4.0	5.1	3.1-8.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Westchester**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	3,569.4	403.2	397.1-409.2	1,685.0	470.3	459.7-481.0	1,884.4	371.2	363.6-378.9	1,920.2	219.3	214.9-223.9	977.0	282.0	273.7-290.6	943.2	183.2	177.9-188.5
	1981-1985	3,851.0	419.2	413.2-425.2	1,833.8	492.5	482.0-503.1	2,017.2	384.1	376.5-391.8	1,988.8	216.2	211.9-220.6	986.0	273.0	265.1-281.1	1,002.8	184.6	179.5-189.9
	1986-1990	3,911.0	415.3	409.5-421.3	1,839.4	480.0	469.9-490.3	2,071.6	385.1	377.6-392.8	2,003.2	211.9	207.8-216.2	987.4	266.0	258.3-273.8	1,015.8	181.8	176.7-186.9
	1991-1995	4,379.6	452.9	446.8-458.9	2,123.0	529.8	519.6-540.2	2,256.6	410.9	403.3-418.7	1,993.8	204.6	200.6-208.7	972.4	255.1	247.9-262.6	1,021.4	177.3	172.4-182.3
	1996-2000	4,980.8	503.8	497.6-510.1	2,496.0	601.0	590.4-611.7	2,484.8	443.9	436.1-451.9	1,916.0	191.2	187.4-195.1	958.4	240.8	234.0-247.8	957.6	162.2	157.6-166.9
	2001-2005	5,188.4	501.3	495.2-507.4	2,571.8	584.3	574.2-594.5	2,616.6	448.1	440.4-456.0	1,803.0	170.1	166.6-173.7	848.2	199.7	193.7-205.8	954.8	152.9	148.5-157.3
	2006-2010	5,483.2	504.1	498.1-510.2	2,708.6	573.8	564.0-583.6	2,774.6	459.1	451.3-467.0	1,732.6	154.0	150.8-157.4	830.2	181.1	175.6-186.8	902.4	137.7	133.6-141.8
	2011-2015	5,567.4	476.3	470.6-482.1	2,711.6	521.8	512.9-530.8	2,855.8	448.5	441.0-456.2	1,640.8	134.2	131.3-137.2	777.0	152.9	148.0-157.9	863.8	122.3	118.6-126.1
	2016-2020	5,607.6	450.8	445.4-456.3	2,740.2	486.3	478.0-494.7	2,867.4	430.3	423.0-437.7	1,485.4	112.2	109.6-114.8	728.6	131.0	126.7-135.4	756.8	99.8	96.5-103.1
Oral cavity and pharynx	1976-1980	99.8	10.8	9.9-11.9	63.8	16.7	14.8-18.8	36.0	6.9	5.9-8.0	45.6	4.9	4.3-5.6	29.4	7.5	6.3-9.0	16.2	3.1	2.4-3.8
	1981-1985	102.6	11.1	10.1-12.1	66.2	17.0	15.2-19.0	36.4	7.0	6.0-8.1	38.0	4.1	3.5-4.7	24.8	6.4	5.3-7.7	13.2	2.6	2.0-3.3
	1986-1990	103.6	11.1	10.1-12.1	70.4	17.5	15.6-19.4	33.2	6.1	5.2-7.1	41.6	4.4	3.9-5.1	27.4	7.2	6.0-8.6	14.2	2.5	2.0-3.2
	1991-1995	84.2	8.8	8.0-9.7	53.4	13.1	11.5-14.8	30.8	5.6	4.7-6.5	30.0	3.1	2.6-3.6	19.4	4.7	3.8-5.7	10.6	1.9	1.4-2.5
	1996-2000	90.6	9.2	8.4-10.1	58.0	13.5	11.9-15.1	32.6	5.9	5.0-6.8	25.0	2.5	2.1-3.0	15.4	3.8	3.0-4.7	9.6	1.6	1.2-2.1
	2001-2005	95.0	9.2	8.3-10.0	57.8	12.7	11.3-14.3	37.2	6.3	5.4-7.3	19.2	1.8	1.4-2.2	11.8	2.7	2.1-3.5	7.4	1.1	0.8-1.5
	2006-2010	109.4	10.0	9.1-10.8	73.2	15.0	13.5-16.6	36.2	5.9	5.1-6.9	20.0	1.8	1.4-2.2	13.2	2.7	2.1-3.5	6.8	1.0	0.7-1.4
	2011-2015	118.4	10.0	9.2-10.9	81.8	15.3	13.8-16.9	36.6	5.7	4.9-6.6	20.4	1.7	1.4-2.1	13.8	2.7	2.1-3.4	6.6	0.9	0.6-1.3
	2016-2020	117.8	9.2	8.5-10.0	80.2	13.7	12.3-15.1	37.6	5.5	4.7-6.4	20.4	1.5	1.2-1.9	13.8	2.4	1.8-3.0	6.6	0.9	0.6-1.3
Esophagus	1976-1980	44.8	4.9	4.3-5.6	29.4	7.8	6.5-9.2	15.4	2.9	2.3-3.7	40.4	4.3	3.8-5.0	27.4	7.1	5.9-8.5	13.0	2.4	1.9-3.1
	1981-1985	48.6	5.1	4.5-5.8	32.2	8.1	6.9-9.6	16.4	3.0	2.4-3.7	44.2	4.6	4.0-5.3	30.8	8.0	6.7-9.5	13.4	2.4	1.9-3.1
	1986-1990	43.6	4.6	4.0-5.3	28.6	7.3	6.1-8.6	15.0	2.7	2.1-3.4	44.8	4.7	4.1-5.4	29.6	7.5	6.3-8.8	15.2	2.7	2.1-3.4
	1991-1995	46.4	4.7	4.1-5.4	30.8	7.6	6.4-8.9	15.6	2.7	2.1-3.3	38.4	3.9	3.4-4.5	25.8	6.5	5.4-7.7	12.6	2.1	1.6-2.7
	1996-2000	47.2	4.7	4.2-5.4	32.4	7.9	6.7-9.2	14.8	2.6	2.0-3.2	39.0	3.9	3.4-4.5	27.8	6.8	5.7-8.1	11.2	1.9	1.4-2.5
	2001-2005	47.0	4.5	3.9-5.1	31.2	7.1	6.0-8.3	15.8	2.5	2.0-3.1	36.4	3.5	3.0-4.0	26.0	5.9	4.9-7.0	10.4	1.7	1.2-2.2
	2006-2010	49.4	4.4	3.9-5.0	34.6	7.2	6.2-8.4	14.8	2.3	1.8-2.9	43.6	3.9	3.4-4.4	31.4	6.6	5.6-7.8	12.2	1.8	1.4-2.4
	2011-2015	44.2	3.7	3.2-4.2	28.8	5.5	4.6-6.5	15.4	2.2	1.7-2.8	35.4	2.9	2.5-3.4	23.4	4.5	3.7-5.4	12.0	1.6	1.2-2.1
	2016-2020	43.6	3.4	3.0-3.9	31.6	5.7	4.8-6.7	12.0	1.6	1.2-2.1	33.8	2.6	2.2-3.1	25.4	4.6	3.8-5.5	8.4	1.1	0.8-1.5
Stomach	1976-1980	113.8	13.2	12.1-14.3	65.2	18.7	16.6-21.0	48.6	9.4	8.3-10.7	89.4	10.5	9.5-11.5	49.4	14.7	12.8-16.8	40.0	7.8	6.7-8.9
	1981-1985	107.4	11.8	10.8-12.9	63.2	17.3	15.4-19.4	44.2	8.1	7.0-9.2	80.8	8.9	8.0-9.8	47.2	13.2	11.5-15.1	33.6	6.0	5.1-7.0
	1986-1990	98.2	10.4	9.5-11.4	56.8	15.6	13.8-17.6	41.4	7.2	6.2-8.2	71.2	7.6	6.8-8.4	38.4	10.7	9.2-12.4	32.8	5.6	4.8-6.6
	1991-1995	111.0	11.4	10.5-12.4	61.6	16.1	14.3-18.0	49.4	8.4	7.3-9.5	71.0	7.3	6.6-8.1	41.2	11.1	9.6-12.7	29.8	5.0	4.2-5.9
	1996-2000	103.0	10.3	9.4-11.2	62.0	15.3	13.6-17.1	41.0	6.7	5.8-7.7	65.0	6.5	5.8-7.2	36.4	9.2	7.9-10.6	28.6	4.7	3.9-5.5
	2001-2005	99.2	9.5	8.6-10.3	57.6	13.3	11.8-14.9	41.6	6.7	5.8-7.7	54.4	5.1	4.5-5.8	29.4	6.8	5.8-8.0	25.0	3.8	3.2-4.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Westchester**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	89.2	8.0	7.3-8.8	49.4	10.7	9.4-12.2	39.8	6.2	5.3-7.1	48.4	4.2	3.6-4.7	28.0	6.0	5.1-7.1	20.4	3.0	2.4-3.6
	2011-2015	101.0	8.5	7.7-9.3	56.8	11.2	9.9-12.6	44.2	6.4	5.5-7.3	41.2	3.4	2.9-3.9	22.8	4.5	3.7-5.4	18.4	2.6	2.1-3.2
	2016-2020	100.0	8.1	7.4-8.8	56.8	10.4	9.2-11.7	43.2	6.4	5.5-7.3	40.2	3.1	2.7-3.6	24.8	4.4	3.7-5.3	15.4	2.2	1.7-2.7
Colorectal	1976-1980	599.2	68.3	65.8-70.9	288.8	81.8	77.4-86.4	310.4	59.7	56.7-62.7	297.0	34.9	33.1-36.8	142.0	43.0	39.7-46.5	155.0	30.1	28.0-32.3
	1981-1985	616.2	66.9	64.6-69.4	314.4	86.2	81.8-90.8	301.8	54.7	52.0-57.6	306.2	33.9	32.2-35.7	147.4	43.1	39.9-46.5	158.8	28.6	26.6-30.7
	1986-1990	553.4	58.4	56.2-60.6	275.4	73.9	69.9-78.0	278.0	48.8	46.3-51.5	267.0	28.2	26.7-29.8	134.2	36.7	33.9-39.7	132.8	22.9	21.2-24.8
	1991-1995	559.0	57.2	55.1-59.4	271.6	69.3	65.6-73.2	287.4	48.8	46.3-51.4	250.6	25.5	24.1-26.9	118.2	31.4	28.9-34.1	132.4	21.9	20.3-23.7
	1996-2000	600.8	59.9	57.8-62.1	299.6	74.0	70.3-77.9	301.2	50.7	48.1-53.3	212.0	20.9	19.7-22.2	108.0	27.5	25.2-30.0	104.0	16.6	15.2-18.2
	2001-2005	549.4	52.1	50.2-54.1	261.6	59.8	56.6-63.2	287.8	46.2	43.8-48.7	191.6	17.8	16.7-19.0	86.0	20.4	18.5-22.4	105.6	16.0	14.6-17.5
	2006-2010	482.0	43.5	41.8-45.3	234.0	49.9	47.1-52.9	248.0	38.7	36.6-41.0	164.2	14.4	13.4-15.4	75.6	16.5	14.9-18.3	88.6	13.0	11.8-14.3
	2011-2015	439.0	37.2	35.6-38.8	218.8	42.9	40.3-45.5	220.2	32.7	30.7-34.7	145.8	11.9	11.0-12.8	70.6	13.7	12.3-15.2	75.2	10.5	9.4-11.7
2016-2020	409.4	32.8	31.4-34.3	208.4	37.7	35.4-40.2	201.0	29.0	27.1-30.9	128.8	9.6	8.8-10.4	64.8	11.6	10.4-13.0	64.0	8.0	7.1-9.0	
Colon excluding rectum	1976-1980	425.6	48.9	46.8-51.1	194.6	56.3	52.6-60.2	231.0	44.6	42.0-47.3	240.2	28.3	26.7-30.0	111.2	33.9	30.9-37.0	129.0	25.1	23.2-27.1
	1981-1985	437.8	47.8	45.8-49.9	215.2	60.1	56.4-64.0	222.6	40.3	38.0-42.8	256.6	28.4	26.9-30.1	122.4	35.9	32.9-39.0	134.2	24.2	22.4-26.1
	1986-1990	405.2	42.7	40.9-44.6	198.8	53.4	50.0-56.9	206.4	36.1	33.9-38.4	230.6	24.3	23.0-25.8	116.8	32.1	29.5-34.9	113.8	19.6	18.0-21.3
	1991-1995	416.0	42.6	40.8-44.5	192.4	49.4	46.2-52.6	223.6	37.8	35.5-40.1	217.6	22.1	20.8-23.5	99.4	26.4	24.0-28.9	118.2	19.5	18.0-21.2
	1996-2000	441.8	43.9	42.1-45.8	213.8	53.3	50.1-56.6	228.0	38.1	35.9-40.4	187.2	18.5	17.3-19.7	94.4	24.1	22.0-26.4	92.8	14.8	13.5-16.2
	2001-2005	404.6	38.3	36.6-40.0	185.0	42.6	39.9-45.5	219.6	34.9	32.8-37.1	164.4	15.2	14.2-16.3	72.0	17.1	15.4-19.0	92.4	14.0	12.7-15.3
	2006-2010	357.8	32.3	30.8-33.8	166.6	35.8	33.4-38.4	191.2	29.6	27.8-31.6	135.6	11.9	11.0-12.8	61.2	13.4	11.9-15.0	74.4	10.8	9.7-12.0
	2011-2015	318.4	26.8	25.5-28.2	153.0	30.2	28.1-32.5	165.4	24.0	22.4-25.8	120.0	9.7	8.9-10.6	57.6	11.1	9.9-12.5	62.4	8.7	7.7-9.7
2016-2020	291.2	23.2	22.0-24.5	141.6	25.8	23.9-27.8	149.6	21.2	19.7-22.9	101.2	7.5	6.8-8.2	51.0	9.1	8.0-10.3	50.2	6.3	5.5-7.1	
Rectum & rectosigmoid	1976-1980	173.6	19.4	18.1-20.7	94.2	25.5	23.1-28.0	79.4	15.1	13.6-16.7	56.8	6.6	5.9-7.5	30.8	9.2	7.7-10.9	26.0	5.0	4.2-6.0
	1981-1985	178.4	19.1	17.9-20.5	99.2	26.1	23.8-28.6	79.2	14.4	13.0-15.9	49.6	5.5	4.8-6.2	25.0	7.2	6.0-8.7	24.6	4.4	3.7-5.3
	1986-1990	148.2	15.7	14.6-16.9	76.6	20.5	18.4-22.7	71.6	12.7	11.4-14.1	36.4	3.9	3.3-4.5	17.4	4.6	3.6-5.7	19.0	3.3	2.7-4.1
	1991-1995	143.0	14.6	13.5-15.7	79.2	19.9	18.0-22.0	63.8	11.0	9.9-12.4	33.0	3.4	2.9-3.9	18.8	5.0	4.0-6.2	14.2	2.4	1.9-3.0
	1996-2000	159.0	16.0	14.9-17.1	85.8	20.7	18.8-22.8	73.2	12.6	11.3-14.0	24.8	2.5	2.0-2.9	13.6	3.4	2.6-4.3	11.2	1.8	1.4-2.4
	2001-2005	144.8	13.8	12.8-14.9	76.6	17.2	15.5-19.0	68.2	11.3	10.1-12.6	27.2	2.6	2.2-3.0	14.0	3.2	2.5-4.1	13.2	2.0	1.6-2.6
	2006-2010	124.2	11.2	10.4-12.2	67.4	14.1	12.6-15.7	56.8	9.1	8.0-10.2	28.6	2.5	2.1-3.0	14.4	3.1	2.4-3.9	14.2	2.1	1.7-2.7
	2011-2015	120.6	10.4	9.5-11.3	65.8	12.6	11.3-14.1	54.8	8.6	7.6-9.8	25.8	2.1	1.8-2.5	13.0	2.5	1.9-3.2	12.8	1.8	1.4-2.4
2016-2020	118.2	9.6	8.8-10.4	66.8	11.9	10.7-13.3	51.4	7.7	6.8-8.8	27.6	2.1	1.8-2.5	13.8	2.5	1.9-3.2	13.8	1.7	1.3-2.2	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**Westchester**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	26.8	3.0	2.5-3.5	16.2	4.3	3.4-5.4	10.6	2.0	1.5-2.7	20.4	2.2	1.8-2.7	12.4	3.3	2.5-4.3	8.0	1.5	1.1-2.1
	1981-1985	21.8	2.3	1.9-2.8	14.4	3.6	2.8-4.6	7.4	1.4	1.0-1.9	22.2	2.4	1.9-2.9	13.4	3.5	2.7-4.5	8.8	1.6	1.2-2.2
	1986-1990	33.6	3.6	3.0-4.2	22.8	6.0	5.0-7.3	10.8	1.9	1.4-2.5	31.8	3.4	2.9-4.0	20.0	5.1	4.1-6.2	11.8	2.1	1.6-2.7
	1991-1995	46.0	4.7	4.1-5.4	25.6	6.4	5.3-7.6	20.4	3.5	2.9-4.3	38.4	3.9	3.4-4.5	21.8	5.5	4.5-6.7	16.6	2.8	2.2-3.5
	1996-2000	56.6	5.7	5.1-6.5	38.4	9.3	8.0-10.7	18.2	3.1	2.5-3.8	43.4	4.4	3.8-5.0	27.6	6.7	5.6-7.9	15.8	2.6	2.1-3.3
	2001-2005	62.8	6.0	5.4-6.7	42.4	9.5	8.2-10.8	20.4	3.4	2.7-4.1	47.8	4.5	4.0-5.1	31.0	7.1	6.0-8.3	16.8	2.6	2.1-3.2
	2006-2010	79.2	7.1	6.4-7.8	60.0	12.0	10.7-13.5	19.2	3.0	2.4-3.7	53.4	4.8	4.2-5.4	35.4	7.3	6.3-8.5	18.0	2.8	2.2-3.5
	2011-2015	86.8	7.1	6.5-7.8	57.8	10.4	9.2-11.7	29.0	4.5	3.7-5.3	65.6	5.4	4.8-6.0	38.6	7.2	6.2-8.4	27.0	3.9	3.2-4.6
	2016-2020	100.0	7.7	7.1-8.5	69.6	12.0	10.8-13.4	30.4	4.3	3.6-5.1	58.2	4.4	3.9-5.0	40.4	7.0	6.0-8.0	17.8	2.5	2.0-3.1
Pancreas	1976-1980	114.8	13.1	12.0-14.2	58.6	16.4	14.4-18.5	56.2	10.8	9.5-12.1	112.0	12.8	11.7-13.9	56.4	15.9	14.0-18.0	55.6	10.6	9.4-12.0
	1981-1985	124.2	13.4	12.3-14.5	60.0	15.6	13.8-17.6	64.2	11.6	10.4-13.0	119.2	13.0	12.0-14.1	56.0	15.2	13.4-17.2	63.2	11.4	10.2-12.8
	1986-1990	117.4	12.3	11.3-13.3	54.0	14.4	12.7-16.3	63.4	11.0	9.8-12.3	112.8	11.8	10.8-12.8	52.6	13.8	12.1-15.6	60.2	10.4	9.3-11.7
	1991-1995	126.8	12.9	11.9-14.0	55.6	14.3	12.6-16.1	71.2	12.0	10.7-13.3	114.4	11.6	10.7-12.6	49.8	12.8	11.3-14.6	64.6	10.7	9.5-11.9
	1996-2000	136.4	13.6	12.6-14.6	66.6	16.3	14.6-18.1	69.8	11.6	10.4-12.9	123.6	12.3	11.3-13.3	65.4	16.3	14.5-18.1	58.2	9.6	8.5-10.7
	2001-2005	133.2	12.6	11.6-13.6	59.6	13.7	12.2-15.4	73.6	11.5	10.4-12.8	119.4	11.2	10.3-12.1	50.8	11.9	10.5-13.5	68.6	10.7	9.6-11.9
	2006-2010	154.4	13.7	12.8-14.7	71.8	15.3	13.7-16.9	82.6	12.5	11.3-13.8	122.8	10.8	10.0-11.7	59.2	12.6	11.2-14.2	63.6	9.4	8.4-10.5
	2011-2015	166.8	13.8	12.8-14.7	79.6	15.4	13.9-17.1	87.2	12.6	11.4-13.9	127.6	10.4	9.6-11.2	60.4	11.8	10.5-13.3	67.2	9.3	8.3-10.4
	2016-2020	193.2	14.9	13.9-15.9	98.6	17.8	16.2-19.5	94.6	12.5	11.4-13.8	132.0	10.0	9.2-10.8	65.8	11.8	10.6-13.2	66.2	8.5	7.6-9.5
Larynx	1976-1980	50.2	5.3	4.6-6.0	42.6	10.7	9.3-12.3	7.6	1.4	1.0-1.9	19.8	2.2	1.8-2.7	18.0	4.8	3.8-6.0	1.8	0.4	0.2-0.7
	1981-1985	50.2	5.2	4.6-5.9	41.2	9.9	8.6-11.4	9.0	1.7	1.2-2.3	16.0	1.7	1.3-2.1	11.6	3.0	2.3-4.0	4.4	0.8	0.5-1.2
	1986-1990	49.6	5.2	4.6-5.9	40.2	9.8	8.5-11.3	9.4	1.7	1.3-2.3	20.0	2.1	1.7-2.5	16.0	4.0	3.1-5.0	4.0	0.7	0.4-1.1
	1991-1995	42.2	4.3	3.8-5.0	36.2	8.7	7.5-10.1	6.0	1.0	0.7-1.5	14.8	1.5	1.2-1.9	13.0	3.2	2.5-4.1	1.8	0.3	0.1-0.6
	1996-2000	43.2	4.4	3.8-5.0	33.8	7.9	6.7-9.2	9.4	1.7	1.3-2.3	12.4	1.2	1.0-1.6	9.2	2.2	1.6-3.0	3.2	0.6	0.3-0.9
	2001-2005	32.2	3.1	2.6-3.6	27.2	6.1	5.1-7.2	5.0	0.9	0.6-1.3	10.4	1.0	0.7-1.3	7.6	1.7	1.2-2.4	2.8	0.5	0.3-0.8
	2006-2010	35.8	3.3	2.8-3.8	27.8	5.9	4.9-7.0	8.0	1.3	0.9-1.8	8.2	0.7	0.5-1.0	6.2	1.3	0.9-1.8	2.0	0.3	0.1-0.5
	2011-2015	31.6	2.6	2.2-3.1	26.2	5.0	4.1-5.9	5.4	0.8	0.5-1.2	7.8	0.6	0.5-0.9	6.8	1.3	0.9-1.8	1.0	0.2	0.1-0.4
	2016-2020	30.2	2.3	1.9-2.7	24.4	4.1	3.4-5.0	5.8	0.8	0.5-1.1	8.2	0.6	0.4-0.8	7.2	1.3	0.9-1.8	1.0	0.1	0.0-0.4
Lung and bronchus	1976-1980	509.8	55.0	52.9-57.2	340.4	89.0	84.6-93.5	169.4	32.0	29.9-34.3	419.4	45.7	43.7-47.7	284.6	76.3	72.2-80.6	134.8	25.3	23.4-27.4
	1981-1985	557.4	58.7	56.5-61.0	350.0	90.1	85.7-94.6	207.4	38.3	35.9-40.7	449.6	47.4	45.5-49.5	287.0	74.9	70.9-79.0	162.6	29.8	27.7-31.9
	1986-1990	578.2	60.2	58.0-62.5	337.4	86.1	81.9-90.4	240.8	43.8	41.3-46.4	480.8	50.0	48.0-52.1	286.2	74.1	70.2-78.1	194.6	34.8	32.6-37.1
	1991-1995	597.4	61.1	59.0-63.4	325.8	80.9	77.0-85.0	271.6	48.5	45.9-51.1	487.2	49.7	47.8-51.7	279.2	70.4	66.7-74.2	208.0	36.4	34.2-38.7
	1996-2000	629.2	63.1	60.9-65.4	326.2	79.0	75.2-82.9	303.0	52.8	50.1-55.5	459.4	46.0	44.1-47.9	251.0	61.5	58.1-65.0	208.4	35.8	33.7-38.1
	2001-2005	604.4	58.0	55.9-60.1	296.2	68.4	64.9-71.9	308.2	51.3	48.8-54.0	426.6	40.6	38.9-42.4	221.8	51.6	48.6-54.7	204.8	33.3	31.3-35.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry**  
**Cancer Incidence and Mortality Trends, 1976-2020**  
**Westchester**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	630.4	57.5	55.5-59.6	294.0	63.7	60.4-67.0	336.4	53.6	51.0-56.3	413.8	37.3	35.7-39.0	204.8	44.9	42.2-47.8	209.0	32.2	30.2-34.3
	2011-2015	600.2	50.0	48.2-51.8	273.0	53.3	50.4-56.2	327.2	47.8	45.5-50.3	384.8	31.6	30.2-33.1	183.2	36.1	33.8-38.6	201.6	28.4	26.6-30.2
	2016-2020	583.8	44.8	43.1-46.5	263.0	46.9	44.3-49.6	320.8	43.6	41.4-45.8	277.4	20.8	19.7-21.9	132.8	23.8	22.0-25.7	144.6	18.5	17.2-20.0
Melanoma of the skin	1976-1980	70.8	8.2	7.3-9.1	35.8	9.3	7.9-10.8	35.0	7.6	6.5-8.8	20.0	2.3	1.9-2.9	10.4	2.7	2.0-3.6	9.6	2.0	1.5-2.7
	1981-1985	83.0	9.3	8.4-10.2	46.0	12.0	10.5-13.8	37.0	7.6	6.5-8.8	24.2	2.7	2.2-3.2	13.8	3.5	2.7-4.5	10.4	2.0	1.5-2.7
	1986-1990	81.4	8.9	8.1-9.8	41.6	10.7	9.3-12.3	39.8	7.9	6.8-9.0	26.0	2.8	2.4-3.4	16.4	4.3	3.4-5.3	9.6	1.9	1.4-2.5
	1991-1995	91.8	9.6	8.8-10.6	53.4	12.9	11.4-14.6	38.4	7.3	6.3-8.4	25.8	2.7	2.2-3.2	15.6	4.0	3.1-5.0	10.2	1.8	1.4-2.4
	1996-2000	118.0	12.0	11.1-13.1	69.8	16.7	15.0-18.6	48.2	8.9	7.8-10.1	29.0	2.9	2.5-3.4	19.0	4.6	3.7-5.7	10.0	1.8	1.3-2.4
	2001-2005	179.0	17.4	16.3-18.6	107.0	24.0	22.0-26.1	72.0	12.8	11.5-14.2	22.8	2.2	1.8-2.6	13.2	3.0	2.3-3.9	9.6	1.6	1.2-2.1
	2006-2010	208.2	19.3	18.1-20.5	119.8	25.6	23.5-27.7	88.4	15.1	13.7-16.6	25.8	2.3	1.9-2.7	15.6	3.4	2.7-4.3	10.2	1.5	1.1-2.0
	2011-2015	245.6	21.1	19.9-22.3	146.2	28.6	26.5-30.8	99.4	15.8	14.4-17.3	30.0	2.5	2.1-2.9	17.0	3.4	2.7-4.2	13.0	1.8	1.4-2.3
	2016-2020	222.2	17.7	16.6-18.8	127.2	23.0	21.2-24.9	95.0	13.8	12.6-15.2	18.2	1.3	1.1-1.7	11.6	2.1	1.6-2.7	6.6	0.8	0.5-1.1
Female breast	1976-1980							536.2	108.4	104.3-112.7							198.8	39.3	36.9-41.9
	1981-1985							587.2	115.8	111.5-120.1							211.8	39.8	37.4-42.3
	1986-1990							631.2	121.5	117.3-125.9							210.2	38.8	36.4-41.3
	1991-1995							679.6	128.6	124.2-133.0							190.0	34.2	32.1-36.5
	1996-2000							781.8	144.1	139.6-148.7							170.2	29.5	27.5-31.5
	2001-2005							772.6	135.4	131.1-139.8							159.2	26.1	24.3-28.0
	2006-2010							842.0	142.5	138.2-147.0							136.4	21.3	19.7-23.0
	2011-2015							885.0	143.2	138.9-147.6							126.6	18.6	17.1-20.1
	2016-2020							911.6	142.9	138.6-147.3							121.2	17.1	15.7-18.6
Cervix uteri	1976-1980							53.2	11.4	10.1-12.9							18.0	3.8	3.0-4.7
	1981-1985							41.2	8.7	7.5-10.0							15.0	2.9	2.3-3.7
	1986-1990							46.4	9.5	8.3-10.8							16.2	3.1	2.5-3.9
	1991-1995							50.8	10.1	8.9-11.5							14.2	2.8	2.2-3.5
	1996-2000							43.6	8.4	7.3-9.6							11.4	2.1	1.6-2.8
	2001-2005							43.2	8.1	7.1-9.3							12.8	2.3	1.7-2.9
	2006-2010							38.0	7.2	6.2-8.3							10.8	1.9	1.4-2.5
	2011-2015							36.2	6.4	5.5-7.5							10.2	1.6	1.2-2.2
	2016-2020							34.2	6.3	5.3-7.3							7.8	1.2	0.9-1.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**  
**Westchester**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uterus and uterus, not otherwise specified	1976-1980							117.8	22.2	20.4-24.1							26.6	5.0	4.1-5.9
	1981-1985							129.4	24.0	22.2-26.0							27.0	4.8	4.0-5.7
	1986-1990							113.4	20.9	19.2-22.8							26.0	4.5	3.8-5.4
	1991-1995							114.8	21.1	19.4-22.9							27.8	4.8	4.0-5.6
	1996-2000							149.0	27.0	25.1-29.1							26.8	4.4	3.7-5.3
	2001-2005							175.8	30.6	28.6-32.7							32.4	5.3	4.5-6.2
	2006-2010							176.2	29.2	27.2-31.2							39.4	6.2	5.3-7.1
	2011-2015							199.8	30.7	28.8-32.7							39.2	5.6	4.8-6.5
	2016-2020							201.6	29.0	27.2-30.9							39.2	5.2	4.5-6.1
Ovary	1976-1980							86.6	17.0	15.4-18.7							49.8	9.6	8.5-10.9
	1981-1985							85.2	16.6	15.1-18.3							56.2	10.4	9.2-11.7
	1986-1990							95.6	18.2	16.6-20.0							60.4	11.0	9.7-12.3
	1991-1995							91.0	17.3	15.7-19.0							58.4	10.4	9.3-11.7
	1996-2000							94.2	17.3	15.7-18.9							60.2	10.4	9.3-11.7
	2001-2005							88.6	15.4	14.0-17.0							54.2	9.0	8.0-10.2
	2006-2010							85.6	14.5	13.1-16.0							53.4	8.4	7.4-9.5
	2011-2015							80.6	12.9	11.6-14.2							52.2	7.6	6.7-8.6
	2016-2020							80.8	12.6	11.4-14.0							44.0	6.0	5.2-6.8
Prostate	1976-1980				247.6	78.9	74.3-83.7							108.0	38.2	34.8-41.7			
	1981-1985				297.6	87.0	82.5-91.8							102.8	32.7	29.8-35.8			
	1986-1990				337.8	92.3	87.8-97.0							117.6	35.5	32.5-38.5			
	1991-1995				592.8	146.8	141.5-152.3							130.6	37.9	35.0-41.0			
	1996-2000				769.4	182.3	176.6-188.2							113.2	31.3	28.7-34.0			
	2001-2005				809.2	181.5	176.0-187.2							100.4	25.0	22.9-27.3			
	2006-2010				852.8	176.7	171.4-182.1							98.0	22.2	20.3-24.2			
	2011-2015				758.6	139.8	135.3-144.4							86.4	17.4	15.8-19.2			
	2016-2020				797.0	134.1	129.9-138.4							80.6	14.7	13.3-16.2			
Testis	1976-1980				14.2	3.5	2.7-4.4							1.2	0.3	0.1-0.7			
	1981-1985				19.0	4.5	3.6-5.5							2.0	0.5	0.2-0.9			
	1986-1990				19.4	4.3	3.5-5.3							0.6	0.1	0.0-0.4			
	1991-1995				20.2	4.6	3.7-5.6							1.0	0.2	0.1-0.5			
	1996-2000				20.6	4.7	3.8-5.7							1.2	0.3	0.1-0.6			
	2001-2005				30.8	7.0	6.0-8.2							0.8	0.2	0.0-0.5			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Westchester**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				27.2	6.3	5.3-7.5							0.6	0.1	0.0-0.4			
	2011-2015				26.4	6.0	5.0-7.2							0.2	0.0	0.0-0.2			
	2016-2020				24.6	5.6	4.7-6.7							1.4	0.3	0.1-0.6			
Urinary bladder (incl. in situ)	1976-1980	188.8	21.6	20.3-23.1	134.6	39.1	36.0-42.3	54.2	10.5	9.3-11.9	53.8	6.5	5.8-7.4	35.8	11.5	9.8-13.4	18.0	3.6	2.9-4.4
	1981-1985	200.4	21.5	20.2-22.9	139.4	38.1	35.2-41.2	61.0	11.1	9.9-12.4	50.2	5.7	5.0-6.4	35.2	10.9	9.2-12.7	15.0	2.7	2.1-3.4
	1986-1990	199.0	20.7	19.5-22.1	138.4	36.2	33.5-39.1	60.6	10.7	9.5-12.0	48.8	5.2	4.5-5.9	31.6	9.4	7.9-11.0	17.2	2.9	2.4-3.7
	1991-1995	196.6	20.1	18.8-21.4	131.8	33.9	31.3-36.6	64.8	11.3	10.1-12.6	42.0	4.3	3.7-4.9	26.8	7.9	6.6-9.4	15.2	2.5	2.0-3.1
	1996-2000	259.6	25.9	24.5-27.3	180.6	44.8	41.9-47.8	79.0	13.4	12.1-14.8	49.0	4.8	4.2-5.4	33.6	8.7	7.5-10.2	15.4	2.4	1.9-3.0
	2001-2005	277.2	26.4	25.0-27.8	199.2	46.5	43.7-49.5	78.0	12.6	11.4-14.0	48.2	4.5	3.9-5.1	32.2	7.9	6.7-9.2	16.0	2.4	1.9-3.1
	2006-2010	252.2	22.6	21.4-23.9	183.2	39.9	37.3-42.6	69.0	10.6	9.5-11.8	47.6	4.2	3.6-4.7	33.2	7.4	6.3-8.6	14.4	2.0	1.6-2.6
	2011-2015	278.8	23.0	21.8-24.3	205.2	40.5	38.0-43.1	73.6	10.5	9.4-11.6	44.4	3.4	3.0-3.9	31.0	6.1	5.2-7.2	13.4	1.7	1.3-2.1
2016-2020	270.4	20.7	19.6-21.9	195.2	35.1	32.9-37.4	75.2	10.1	9.1-11.2	45.0	3.2	2.8-3.7	30.4	5.5	4.7-6.5	14.6	1.7	1.3-2.2	
Kidney and renal pelvis	1976-1980	58.0	6.4	5.7-7.2	36.2	9.4	8.0-11.0	21.8	4.2	3.4-5.0	30.2	3.3	2.8-3.9	17.6	4.7	3.7-5.8	12.6	2.3	1.8-3.0
	1981-1985	79.6	8.7	7.9-9.6	49.4	12.6	11.1-14.4	30.2	5.8	4.9-6.8	34.4	3.8	3.2-4.4	20.6	5.4	4.4-6.6	13.8	2.6	2.0-3.3
	1986-1990	84.6	9.0	8.1-9.9	53.2	13.7	12.0-15.5	31.4	5.9	5.0-7.0	36.0	3.8	3.3-4.4	20.2	5.4	4.3-6.6	15.8	2.8	2.2-3.6
	1991-1995	98.2	10.2	9.3-11.2	58.2	14.3	12.7-16.1	40.0	7.3	6.3-8.4	33.8	3.5	3.0-4.1	21.2	5.4	4.4-6.6	12.6	2.2	1.6-2.8
	1996-2000	127.0	12.9	11.9-14.0	83.4	19.6	17.7-21.6	43.6	7.7	6.7-8.8	33.6	3.4	2.9-3.9	19.2	4.8	3.9-5.9	14.4	2.5	1.9-3.1
	2001-2005	147.8	14.4	13.4-15.5	92.8	20.8	18.9-22.8	55.0	9.5	8.4-10.7	37.8	3.6	3.1-4.1	24.0	5.6	4.6-6.7	13.8	2.1	1.7-2.7
	2006-2010	167.6	15.4	14.3-16.4	105.0	22.1	20.2-24.1	62.6	10.0	8.9-11.3	30.4	2.6	2.2-3.1	18.0	3.8	3.0-4.7	12.4	1.8	1.4-2.3
	2011-2015	179.0	15.3	14.3-16.4	121.4	23.2	21.3-25.2	57.6	9.0	8.0-10.1	28.2	2.3	1.9-2.7	20.0	4.0	3.2-4.8	8.2	1.1	0.8-1.5
2016-2020	188.8	15.2	14.2-16.3	126.0	22.3	20.5-24.1	62.8	9.4	8.3-10.5	25.4	2.0	1.6-2.3	17.8	3.2	2.6-4.0	7.6	1.0	0.7-1.5	
Brain and other central nervous system	1976-1980	53.4	5.8	5.2-6.6	30.8	7.6	6.4-8.9	22.6	4.5	3.7-5.5	38.6	4.2	3.6-4.9	23.4	5.9	4.9-7.2	15.2	2.9	2.3-3.6
	1981-1985	56.8	6.2	5.5-7.0	28.6	7.1	6.0-8.4	28.2	5.5	4.6-6.5	38.0	4.1	3.6-4.8	21.2	5.5	4.5-6.7	16.8	3.1	2.5-3.9
	1986-1990	70.6	7.7	6.9-8.6	40.0	9.7	8.4-11.2	30.6	6.1	5.1-7.1	45.8	4.9	4.3-5.6	24.8	6.1	5.0-7.3	21.0	4.1	3.3-5.0
	1991-1995	66.6	7.1	6.4-7.9	33.2	8.1	6.9-9.4	33.4	6.5	5.6-7.6	38.6	4.1	3.5-4.7	21.4	5.3	4.3-6.4	17.2	3.2	2.6-4.0
	1996-2000	74.8	7.8	7.0-8.7	36.4	8.6	7.4-10.0	38.4	7.4	6.4-8.5	43.4	4.5	3.9-5.1	23.8	5.7	4.7-6.8	19.6	3.6	3.0-4.5
	2001-2005	78.0	7.8	7.0-8.6	43.6	9.8	8.5-11.1	34.4	6.3	5.4-7.3	40.8	4.0	3.4-4.6	21.8	4.9	4.0-5.9	19.0	3.3	2.7-4.1
	2006-2010	70.6	6.8	6.1-7.6	41.0	8.8	7.6-10.1	29.6	5.2	4.3-6.1	39.6	3.6	3.1-4.2	23.8	5.1	4.2-6.1	15.8	2.6	2.0-3.3
	2011-2015	70.8	6.4	5.8-7.2	37.8	7.5	6.5-8.7	33.0	5.5	4.6-6.4	43.6	3.7	3.2-4.3	22.2	4.3	3.5-5.2	21.4	3.2	2.6-3.9
2016-2020	71.6	6.5	5.8-7.2	38.8	7.6	6.6-8.8	32.8	5.5	4.6-6.5	44.8	3.7	3.2-4.3	26.0	4.8	4.0-5.7	18.8	2.8	2.3-3.5	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Westchester**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	31.2	3.6	3.1-4.3	10.0	2.5	1.8-3.3	21.2	4.6	3.7-5.5	7.0	0.8	0.6-1.2	2.0	0.6	0.3-1.1	5.0	1.0	0.6-1.4
	1981-1985	35.2	4.0	3.4-4.6	12.4	3.2	2.4-4.1	22.8	4.7	3.9-5.7	5.8	0.6	0.4-0.9	2.6	0.8	0.4-1.3	3.2	0.6	0.3-0.9
	1986-1990	40.6	4.5	3.9-5.1	11.8	2.9	2.2-3.7	28.8	5.9	5.0-6.9	6.0	0.6	0.4-0.9	1.8	0.5	0.2-0.9	4.2	0.7	0.5-1.1
	1991-1995	46.8	5.0	4.4-5.7	14.4	3.4	2.6-4.2	32.4	6.5	5.6-7.6	5.2	0.5	0.3-0.8	2.4	0.6	0.3-1.0	2.8	0.5	0.3-0.8
	1996-2000	84.0	8.9	8.0-9.8	21.8	5.0	4.1-6.0	62.2	12.4	11.0-13.9	7.8	0.8	0.5-1.1	3.2	0.8	0.5-1.3	4.6	0.7	0.5-1.1
	2001-2005	127.8	13.1	12.1-14.1	32.8	7.1	6.1-8.3	95.0	18.5	16.8-20.2	5.2	0.5	0.3-0.7	2.2	0.5	0.2-0.9	3.0	0.5	0.3-0.8
	2006-2010	190.6	19.2	18.0-20.5	49.4	10.4	9.1-11.8	141.2	27.2	25.2-29.4	5.4	0.5	0.3-0.7	2.8	0.6	0.3-1.0	2.6	0.4	0.2-0.6
	2011-2015	223.2	21.7	20.4-23.0	64.6	12.9	11.5-14.4	158.6	30.0	27.8-32.2	6.0	0.5	0.3-0.7	3.0	0.6	0.3-1.0	3.0	0.4	0.2-0.7
Hodgkin lymphoma	1976-1980	32.4	3.7	3.1-4.3	17.0	4.3	3.4-5.3	15.4	3.2	2.5-4.1	12.8	1.5	1.2-2.0	8.8	2.5	1.8-3.4	4.0	0.8	0.5-1.3
	1981-1985	34.6	3.8	3.2-4.4	17.2	4.2	3.3-5.2	17.4	3.4	2.7-4.2	9.8	1.0	0.8-1.4	5.8	1.3	0.9-2.0	4.0	0.7	0.5-1.2
	1986-1990	29.0	3.1	2.6-3.6	15.0	3.4	2.7-4.3	14.0	2.8	2.2-3.6	10.0	1.1	0.8-1.4	6.6	1.6	1.1-2.3	3.4	0.6	0.4-1.0
	1991-1995	31.2	3.4	2.9-4.0	17.6	4.1	3.3-5.0	13.6	2.7	2.1-3.5	7.0	0.7	0.5-1.0	3.6	0.9	0.5-1.5	3.4	0.6	0.4-1.0
	1996-2000	35.2	3.9	3.3-4.5	19.0	4.5	3.6-5.5	16.2	3.4	2.7-4.2	5.0	0.6	0.4-0.8	2.8	0.7	0.4-1.2	2.2	0.4	0.2-0.8
	2001-2005	37.2	4.1	3.6-4.8	18.6	4.3	3.5-5.3	18.6	4.0	3.2-4.9	6.4	0.6	0.4-0.9	2.6	0.6	0.3-1.0	3.8	0.7	0.4-1.1
	2006-2010	32.2	3.5	3.0-4.1	14.6	3.3	2.6-4.1	17.6	3.8	3.0-4.7	3.2	0.3	0.2-0.5	1.0	0.2	0.1-0.5	2.2	0.4	0.2-0.7
	2011-2015	35.8	3.7	3.2-4.3	20.0	4.3	3.5-5.3	15.8	3.2	2.5-4.0	2.8	0.2	0.1-0.4	1.6	0.3	0.1-0.6	1.2	0.2	0.1-0.4
2016-2020	31.8	3.2	2.7-3.7	16.4	3.3	2.6-4.1	15.4	3.1	2.4-3.9	3.2	0.2	0.1-0.4	1.4	0.2	0.1-0.5	1.8	0.2	0.1-0.5	
Non-Hodgkin Lymphomas	1976-1980	106.2	11.9	10.9-13.0	53.4	14.1	12.4-16.0	52.8	10.2	9.0-11.6	56.8	6.4	5.6-7.2	29.2	7.7	6.4-9.1	27.6	5.4	4.5-6.3
	1981-1985	130.2	14.3	13.2-15.5	66.0	17.3	15.5-19.4	64.2	12.1	10.8-13.5	62.2	6.8	6.1-7.6	34.2	9.3	7.9-10.9	28.0	5.1	4.3-6.1
	1986-1990	143.0	15.3	14.2-16.5	77.4	19.5	17.6-21.6	65.6	12.1	10.8-13.6	68.2	7.2	6.5-8.1	35.4	9.3	7.9-10.8	32.8	5.8	4.9-6.8
	1991-1995	175.2	18.3	17.1-19.5	87.4	21.8	19.8-24.0	87.8	15.7	14.3-17.3	81.4	8.4	7.6-9.2	42.2	11.1	9.6-12.8	39.2	6.7	5.7-7.7
	1996-2000	211.4	21.5	20.2-22.8	109.8	26.3	24.1-28.6	101.6	17.8	16.3-19.4	88.8	8.9	8.1-9.8	47.6	11.7	10.2-13.3	41.2	6.9	6.0-7.9
	2001-2005	232.6	22.5	21.3-23.9	120.6	27.5	25.3-29.8	112.0	19.1	17.5-20.8	77.4	7.3	6.6-8.1	39.6	9.3	8.1-10.7	37.8	5.8	5.0-6.7
	2006-2010	260.6	24.2	22.9-25.5	138.4	29.8	27.6-32.1	122.2	19.7	18.1-21.4	72.4	6.4	5.7-7.1	37.4	8.3	7.1-9.6	35.0	5.2	4.4-6.0
	2011-2015	277.2	23.8	22.6-25.1	145.4	28.4	26.3-30.6	131.8	20.4	18.9-22.1	61.8	5.0	4.5-5.7	28.6	5.7	4.8-6.7	33.2	4.5	3.9-5.3
2016-2020	277.8	22.4	21.2-23.7	153.6	28.2	26.2-30.4	124.2	17.8	16.4-19.3	54.2	4.0	3.5-4.5	28.4	5.0	4.2-6.0	25.8	3.2	2.7-3.9	
Myeloma	1976-1980	43.0	4.8	4.2-5.5	19.4	5.2	4.2-6.4	23.6	4.6	3.8-5.5	29.2	3.3	2.7-3.9	13.2	3.7	2.8-4.8	16.0	3.0	2.4-3.8
	1981-1985	45.2	4.9	4.3-5.6	20.8	5.7	4.6-7.0	24.4	4.4	3.7-5.3	31.2	3.4	2.9-3.9	15.0	4.1	3.2-5.2	16.2	2.9	2.3-3.6
	1986-1990	46.0	4.8	4.2-5.4	21.6	5.6	4.5-6.8	24.4	4.3	3.5-5.1	29.2	3.0	2.6-3.6	13.0	3.6	2.7-4.6	16.2	2.8	2.2-3.4
	1991-1995	56.2	5.8	5.1-6.5	26.4	6.6	5.5-7.8	29.8	5.1	4.3-6.0	38.6	3.9	3.4-4.5	15.2	3.9	3.0-4.9	23.4	3.9	3.2-4.7
	1996-2000	65.6	6.6	5.9-7.3	37.4	9.1	7.8-10.5	28.2	4.8	4.0-5.6	36.2	3.6	3.1-4.2	19.8	4.9	4.0-6.0	16.4	2.7	2.2-3.4
	2001-2005	70.8	6.8	6.1-7.5	39.0	8.9	7.7-10.2	31.8	5.2	4.5-6.1	33.4	3.1	2.7-3.7	19.2	4.5	3.6-5.5	14.2	2.3	1.8-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Westchester**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	96.6	8.7	7.9-9.5	51.4	10.8	9.5-12.3	45.2	7.0	6.1-8.0	39.0	3.4	3.0-4.0	20.0	4.3	3.5-5.2	19.0	2.8	2.3-3.5
	2011-2015	103.6	8.8	8.0-9.6	61.6	12.1	10.7-13.5	42.0	6.2	5.4-7.2	34.6	2.8	2.4-3.2	16.8	3.2	2.6-4.0	17.8	2.4	1.9-3.0
	2016-2020	111.2	8.7	8.0-9.5	59.8	10.7	9.5-12.0	51.4	7.3	6.4-8.2	37.2	2.8	2.4-3.2	19.0	3.4	2.8-4.2	18.2	2.3	1.8-2.9
Leukemias	1976-1980	106.6	12.3	11.2-13.4	56.0	15.7	13.8-17.8	50.6	10.2	8.9-11.5	69.4	8.2	7.3-9.1	37.4	11.0	9.4-12.8	32.0	6.5	5.5-7.6
	1981-1985	115.2	13.1	12.0-14.2	60.8	16.7	14.8-18.8	54.4	10.7	9.4-12.1	71.4	8.0	7.2-8.9	35.8	9.8	8.3-11.4	35.6	6.8	5.9-7.9
	1986-1990	107.8	11.8	10.8-12.8	56.2	14.8	13.1-16.7	51.6	9.6	8.4-10.9	70.6	7.6	6.8-8.4	39.0	10.4	9.0-12.0	31.6	5.7	4.8-6.6
	1991-1995	123.2	13.0	12.0-14.1	71.4	18.2	16.4-20.3	51.8	9.6	8.4-10.8	77.4	8.1	7.3-9.0	42.8	11.4	9.9-13.1	34.6	6.2	5.3-7.2
	1996-2000	140.8	14.4	13.3-15.5	82.2	20.2	18.3-22.3	58.6	10.3	9.1-11.5	79.2	8.0	7.2-8.8	45.2	11.2	9.8-12.8	34.0	5.7	4.9-6.7
	2001-2005	157.8	15.3	14.3-16.5	90.6	20.8	19.0-22.9	67.2	11.4	10.2-12.7	77.2	7.4	6.7-8.2	44.8	10.7	9.3-12.1	32.4	5.2	4.4-6.1
	2006-2010	189.4	17.6	16.5-18.8	106.8	23.2	21.3-25.3	82.6	13.4	12.1-14.8	78.4	7.0	6.4-7.8	43.4	9.7	8.4-11.1	35.0	5.2	4.4-6.1
	2011-2015	218.6	18.8	17.7-20.0	126.6	25.0	23.1-27.1	92.0	14.0	12.7-15.5	71.2	5.9	5.3-6.5	40.8	8.3	7.2-9.5	30.4	4.2	3.5-5.0
	2016-2020	220.6	18.0	16.9-19.1	126.4	23.2	21.4-25.1	94.2	13.8	12.5-15.2	68.0	5.1	4.6-5.7	38.8	7.1	6.1-8.1	29.2	3.8	3.2-4.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Wyoming**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	129.0	370.0	341.5-400.1	66.6	432.1	385.2-482.8	62.4	337.2	300.0-377.7	70.0	200.2	179.5-222.6	37.4	250.7	214.8-290.5	32.6	168.7	143.4-197.2
	1981-1985	154.0	420.9	391.3-452.1	80.8	508.8	458.9-562.3	73.2	367.9	330.4-408.6	78.8	218.7	197.4-241.6	42.8	282.4	244.7-323.9	36.0	178.7	153.0-207.4
	1986-1990	166.0	429.8	400.7-460.5	82.2	474.0	427.9-523.4	83.8	408.4	369.4-450.3	75.0	198.7	178.9-220.0	38.8	237.0	204.0-273.6	36.2	174.1	149.1-202.1
	1991-1995	185.4	461.1	431.8-492.0	101.0	575.6	525.0-629.5	84.4	392.7	355.5-432.8	83.8	207.6	188.1-228.5	46.8	272.1	237.4-310.2	37.0	164.1	141.0-190.0
	1996-2000	223.2	528.4	497.8-560.3	121.4	648.1	596.5-702.8	101.8	456.1	417.0-498.0	83.0	197.4	178.9-217.4	46.2	259.0	226.1-295.2	36.8	157.3	135.2-182.2
	2001-2005	238.2	534.0	503.9-565.4	127.4	634.2	584.6-686.8	110.8	473.5	434.4-515.2	88.4	198.8	180.6-218.3	45.6	237.9	207.2-271.7	42.8	174.8	151.9-200.4
	2006-2010	274.2	578.5	547.8-610.5	149.8	658.6	610.6-709.4	124.4	525.3	483.9-569.4	88.4	187.9	170.6-206.5	47.4	230.8	201.3-263.2	41.0	162.0	140.2-186.6
	2011-2015	277.2	553.5	524.1-584.3	145.8	596.2	551.8-643.4	131.4	540.2	498.1-585.0	85.0	174.2	157.8-192.0	44.0	198.8	172.4-228.2	41.0	158.1	136.8-182.2
	2016-2020	280.4	517.4	489.8-546.3	153.2	560.0	519.5-603.0	127.2	498.2	458.0-541.2	88.0	163.3	148.1-179.8	49.8	197.6	172.9-225.1	38.2	139.8	120.1-162.2
Oral cavity and pharynx	1976-1980	2.6	8.1	4.3-13.8	1.0	6.7	2.1-15.5	1.6	9.2	3.9-18.3	1.0	3.1	1.0-7.2	0.6	4.7	0.9-13.0	0.4	2.2	0.3-7.9
	1981-1985	5.0	14.1	9.1-20.9	3.8	22.7	13.6-35.5	1.2	6.7	2.4-14.8	1.8	4.8	2.2-9.1	1.6	9.2	3.9-18.1	0.2	0.9	0.0-5.3
	1986-1990	3.2	8.3	4.7-13.5	2.4	13.0	6.6-22.8	0.8	4.4	1.1-11.2	1.4	4.1	1.6-8.3	1.0	6.3	2.0-14.5	0.4	3.0	0.4-9.8
	1991-1995	2.8	6.7	3.6-11.2	1.6	7.9	3.3-15.9	1.2	5.1	1.9-11.4	1.2	2.9	1.1-6.3	0.6	2.9	0.6-8.6	0.6	2.4	0.5-7.4
	1996-2000	4.8	11.4	7.3-17.0	3.4	17.4	10.1-27.8	1.4	6.2	2.5-13.0	1.4	3.4	1.4-6.9	1.0	6.0	1.9-13.8	0.4	1.6	0.2-6.3
	2001-2005	4.0	8.8	5.4-13.7	3.2	14.8	8.4-24.2	0.8	3.7	1.0-9.7	1.4	3.1	1.3-6.6	1.2	5.5	2.0-12.2	0.2	0.6	0.0-4.7
	2006-2010	7.2	14.8	10.3-20.7	5.2	20.3	13.1-30.3	2.0	8.4	4.0-16.0	1.4	2.7	1.1-5.8	1.0	4.5	1.4-11.0	0.4	1.3	0.2-5.8
	2011-2015	8.2	15.6	11.1-21.5	6.0	22.7	15.0-33.1	2.2	7.8	3.9-14.7	1.4	2.6	1.0-5.7	1.2	4.6	1.5-10.8	0.2	0.8	0.0-5.3
	2016-2020	6.6	11.3	7.7-16.3	4.8	17.1	10.7-26.3	1.8	6.6	2.9-13.6	1.4	2.3	0.9-5.1	1.2	4.0	1.4-9.5	0.2	0.6	0.0-5.0
Esophagus	1976-1980	1.4	3.6	1.4-7.5	1.2	7.4	2.5-16.4	0.2	0.9	0.0-5.5	1.0	2.7	0.8-6.3	0.8	4.7	1.2-12.2	0.2	1.2	0.0-6.2
	1981-1985	1.8	5.1	2.3-9.6	1.6	11.4	4.8-22.1	0.2	1.0	0.0-5.9	1.0	2.8	0.9-6.5	0.8	6.0	1.5-14.8	0.2	1.0	0.0-5.9
	1986-1990	1.8	4.9	2.2-9.2	1.2	6.6	2.4-14.4	0.6	2.9	0.6-8.7	1.6	4.4	1.9-8.6	1.2	7.2	2.6-15.5	0.4	1.8	0.2-6.9
	1991-1995	1.2	3.1	1.1-6.7	0.8	4.2	1.1-10.6	0.4	1.7	0.2-6.4	1.8	4.4	2.0-8.3	1.4	7.4	2.9-15.4	0.4	1.6	0.2-6.2
	1996-2000	2.4	5.8	3.0-10.1	2.4	13.2	6.7-23.0	0.0	0.0	0.0-3.7	1.8	4.4	2.0-8.3	1.8	9.6	4.4-18.0	0.0	0.0	0.0-3.7
	2001-2005	2.6	5.7	3.0-9.8	1.8	7.7	3.5-15.0	0.8	3.5	0.9-9.4	2.4	5.4	2.8-9.5	1.6	7.3	3.1-14.5	0.8	3.2	0.8-8.7
	2006-2010	3.6	7.4	4.4-11.9	2.8	11.2	6.0-19.3	0.8	3.0	0.8-8.4	2.6	5.4	2.8-9.4	2.2	9.2	4.5-16.9	0.4	1.8	0.2-6.9
	2011-2015	5.4	10.2	6.7-15.1	4.2	15.9	9.6-25.1	1.2	4.5	1.6-10.6	3.8	7.4	4.4-11.9	3.0	13.1	7.1-22.2	0.8	2.7	0.7-7.9
	2016-2020	3.2	5.7	3.2-9.6	3.0	10.7	5.9-18.4	0.2	0.6	0.0-5.0	2.6	4.8	2.5-8.6	2.2	8.4	4.1-15.6	0.4	1.4	0.2-6.3
Stomach	1976-1980	3.2	8.9	5.1-14.5	1.6	9.8	4.0-19.4	1.6	8.0	3.4-15.9	1.8	5.0	2.2-9.5	0.8	4.1	1.1-10.8	1.0	5.0	1.6-11.8
	1981-1985	4.0	10.5	6.4-16.2	2.2	12.6	6.1-22.8	1.8	8.2	3.8-15.9	3.2	8.6	4.9-14.0	1.6	10.1	4.2-20.1	1.6	7.4	3.2-14.8
	1986-1990	2.2	6.1	3.0-10.9	1.6	10.1	4.3-19.7	0.6	3.2	0.6-9.4	1.6	4.4	1.9-8.6	1.2	6.9	2.5-15.0	0.4	2.4	0.3-8.3
	1991-1995	3.2	8.0	4.5-12.9	1.4	7.6	3.0-15.5	1.8	7.1	3.3-13.9	1.8	4.7	2.2-8.9	1.2	8.2	2.9-17.3	0.6	2.3	0.5-7.2
	1996-2000	4.0	9.6	5.8-14.8	2.4	13.9	7.1-24.3	1.6	6.1	2.6-12.5	2.2	5.2	2.6-9.2	1.0	5.7	1.9-13.1	1.2	4.9	1.8-11.0
	2001-2005	4.4	10.1	6.3-15.4	2.2	11.2	5.5-20.2	2.2	9.0	4.4-16.5	2.4	5.4	2.8-9.5	0.8	4.4	1.1-11.2	1.6	6.3	2.7-13.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Wyoming**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	3.2	6.6	3.7-10.9	2.8	11.3	6.0-19.5	0.4	2.0	0.2-7.3	0.8	2.0	0.5-5.0	0.6	2.9	0.6-8.6	0.2	1.0	0.0-5.7
	2011-2015	3.2	6.5	3.7-10.9	2.4	10.5	5.3-18.9	0.8	3.3	0.8-9.2	0.8	1.7	0.5-4.6	0.6	3.1	0.6-9.1	0.2	0.6	0.0-5.0
	2016-2020	3.6	6.3	3.7-10.3	3.0	10.3	5.7-17.7	0.6	2.0	0.4-7.2	1.4	2.4	0.9-5.4	0.8	2.4	0.6-7.1	0.6	2.1	0.4-7.3
Colorectal	1976-1980	19.4	56.4	45.6-68.9	9.4	68.3	49.6-91.1	10.0	50.8	37.5-67.4	9.6	27.4	20.2-36.4	5.2	35.7	22.9-52.6	4.4	21.4	13.4-32.6
	1981-1985	23.8	66.5	55.0-79.7	12.0	78.5	59.4-101.5	11.8	57.1	43.3-73.9	12.8	36.5	28.0-46.6	6.4	47.4	32.0-66.9	6.4	30.4	20.7-43.2
	1986-1990	25.0	64.8	53.8-77.3	11.4	67.5	50.8-87.8	13.6	61.3	47.4-78.1	9.4	24.9	18.3-33.1	4.2	25.9	15.9-39.6	5.2	22.1	14.4-32.7
	1991-1995	24.0	59.1	49.0-70.7	12.6	73.1	55.7-94.0	11.4	49.1	37.1-64.0	12.0	29.5	22.5-38.0	6.2	36.8	24.6-52.5	5.8	24.9	16.6-36.1
	1996-2000	29.8	71.0	60.0-83.4	14.4	81.0	63.1-102.2	15.4	65.4	51.5-82.2	12.2	29.0	22.2-37.3	5.4	32.1	21.0-46.6	6.8	28.3	19.5-39.9
	2001-2005	30.4	68.7	58.2-80.6	16.6	90.2	71.3-112.3	13.8	55.6	43.1-70.9	9.8	21.9	16.2-29.1	5.4	32.6	21.2-47.4	4.4	16.2	10.1-25.1
	2006-2010	29.4	64.3	54.2-75.8	14.6	69.7	54.1-88.4	14.8	60.1	47.0-76.1	9.0	19.1	13.9-25.7	4.8	22.6	14.2-34.0	4.2	14.9	9.2-23.5
	2011-2015	23.2	49.3	40.6-59.4	12.4	54.2	41.0-70.3	10.8	43.5	32.4-57.4	7.8	16.7	11.8-23.0	3.8	18.2	10.7-28.9	4.0	15.3	9.2-24.3
2016-2020	21.6	43.4	35.4-52.9	11.8	45.5	34.3-59.5	9.8	43.1	31.1-58.4	6.6	12.8	8.7-18.3	3.4	13.0	7.3-21.6	3.2	13.2	7.2-22.5	
Colon excluding rectum	1976-1980	13.4	39.7	30.6-50.4	6.6	49.1	33.4-69.1	6.8	35.3	24.2-49.7	7.8	22.8	16.1-31.1	4.0	29.2	17.5-45.2	3.8	18.5	11.1-29.1
	1981-1985	18.6	51.5	41.5-63.2	8.0	50.7	35.9-69.4	10.6	51.6	38.5-67.8	9.6	27.2	20.0-36.1	4.4	31.8	19.6-48.0	5.2	24.9	16.1-36.7
	1986-1990	16.6	43.2	34.4-53.7	6.2	38.1	25.6-54.3	10.4	46.3	34.4-61.0	9.0	23.8	17.3-31.8	4.0	24.5	14.8-37.9	5.0	21.2	13.7-31.6
	1991-1995	18.2	45.5	36.6-55.8	9.2	54.1	39.3-72.4	9.0	38.9	28.2-52.4	9.8	24.1	17.8-31.9	4.4	26.1	16.1-39.5	5.4	23.2	15.2-34.1
	1996-2000	24.0	57.2	47.4-68.4	10.6	60.9	45.3-79.8	13.4	56.4	43.6-72.0	11.4	27.2	20.6-35.2	5.2	30.9	20.0-45.2	6.2	26.2	17.7-37.5
	2001-2005	21.2	48.1	39.4-58.3	11.0	61.7	46.1-80.6	10.2	41.0	30.4-54.3	8.4	18.8	13.6-25.5	4.2	25.9	15.9-39.5	4.2	15.6	9.6-24.3
	2006-2010	21.0	46.3	37.7-56.2	10.0	46.7	34.2-62.1	11.0	44.8	33.5-58.9	8.8	18.7	13.6-25.3	4.6	21.9	13.6-33.3	4.2	14.9	9.2-23.5
	2011-2015	18.0	38.5	30.8-47.5	8.8	40.1	28.7-54.4	9.2	36.1	26.2-48.9	6.8	14.5	10.0-20.4	3.2	14.7	8.2-24.4	3.6	13.8	8.0-22.5
2016-2020	12.8	25.8	19.7-33.3	6.6	25.9	17.6-37.1	6.2	26.7	17.6-39.1	4.2	8.0	4.9-12.6	2.2	8.4	4.1-15.7	2.0	7.7	3.5-15.3	
Rectum & rectosigmoid	1976-1980	6.0	16.7	11.2-24.0	2.8	19.2	10.2-32.3	3.2	15.5	8.8-25.5	1.8	4.7	2.1-9.0	1.2	6.5	2.3-14.3	0.6	2.9	0.6-8.7
	1981-1985	5.2	15.0	9.7-22.0	4.0	27.8	16.7-42.9	1.2	5.5	2.0-12.2	3.2	9.3	5.3-15.1	2.0	15.6	7.3-28.4	1.2	5.5	2.0-12.3
	1986-1990	8.4	21.5	15.5-29.2	5.2	29.4	19.1-43.2	3.2	15.0	8.5-24.7	0.4	1.1	0.1-3.9	0.2	1.4	0.0-7.0	0.2	0.9	0.0-5.4
	1991-1995	5.8	13.6	9.1-19.6	3.4	19.0	10.7-30.8	2.4	10.2	5.3-18.2	2.2	5.4	2.7-9.7	1.8	10.7	4.6-20.5	0.4	1.7	0.2-6.4
	1996-2000	5.8	13.8	9.2-19.8	3.8	20.1	12.1-31.3	2.0	9.1	4.3-16.9	0.8	1.8	0.5-4.8	0.2	1.2	0.0-6.2	0.6	2.1	0.4-6.9
	2001-2005	9.2	20.6	15.1-27.5	5.6	28.5	18.7-41.5	3.6	14.7	8.6-23.6	1.4	3.1	1.2-6.5	1.2	6.6	2.4-14.4	0.2	0.6	0.0-4.7
	2006-2010	8.4	18.1	12.9-24.6	4.6	23.1	14.4-35.0	3.8	15.3	9.2-24.5	0.2	0.3	0.0-2.3	0.2	0.6	0.0-4.6	0.0	0.0	0.0-3.9
	2011-2015	5.2	10.8	7.0-16.1	3.6	14.1	8.1-23.0	1.6	7.4	3.2-14.9	1.0	2.2	0.7-5.3	0.6	3.5	0.7-9.9	0.4	1.5	0.2-6.3
2016-2020	8.8	17.7	12.6-24.1	5.2	19.6	12.6-29.3	3.6	16.4	9.2-27.0	2.4	4.8	2.4-8.7	1.2	4.6	1.5-10.7	1.2	5.5	1.9-12.7	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**  
**Wyoming**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	0.4	1.0	0.1-3.7	0.2	1.2	0.0-6.7	0.2	0.9	0.0-5.4	1.0	2.8	0.9-6.5	0.4	2.8	0.3-9.8	0.6	3.1	0.6-9.1
	1981-1985	1.0	2.6	0.8-6.1	0.8	5.0	1.3-13.0	0.2	1.0	0.0-5.8	2.0	5.8	2.8-10.5	0.8	6.3	1.6-15.5	1.2	5.7	2.1-12.7
	1986-1990	1.4	3.9	1.6-8.0	1.0	6.6	2.1-15.0	0.4	2.1	0.2-7.5	1.0	2.8	0.9-6.4	0.6	4.3	0.8-12.1	0.4	2.5	0.3-8.6
	1991-1995	1.2	3.1	1.1-6.7	0.8	4.8	1.3-12.0	0.4	1.7	0.2-6.5	1.8	4.5	2.0-8.5	0.8	4.7	1.3-11.7	1.0	4.5	1.4-10.6
	1996-2000	2.0	4.8	2.3-8.8	1.0	4.9	1.6-11.4	1.0	4.2	1.4-10.2	1.0	2.4	0.8-5.6	0.6	2.9	0.6-8.5	0.4	1.9	0.2-7.0
	2001-2005	2.4	5.3	2.7-9.4	1.4	5.9	2.3-12.4	1.0	4.4	1.4-10.7	1.6	3.7	1.6-7.3	0.8	3.8	1.0-9.9	0.8	3.4	0.9-9.2
	2006-2010	1.6	3.1	1.3-6.3	1.4	5.6	2.2-12.1	0.2	0.8	0.0-5.3	2.4	4.9	2.5-8.7	1.8	8.1	3.5-15.8	0.6	2.1	0.4-7.0
	2011-2015	3.8	7.4	4.3-11.8	3.0	12.6	6.8-21.5	0.8	2.6	0.7-7.8	2.8	5.1	2.8-8.9	2.0	7.3	3.4-14.1	0.8	2.7	0.7-7.9
2016-2020	4.2	8.0	4.8-12.5	3.4	12.2	7.0-20.2	0.8	2.9	0.8-8.4	3.4	6.3	3.6-10.4	2.8	10.3	5.5-18.0	0.6	1.9	0.4-6.8	
Pancreas	1976-1980	5.0	14.8	9.5-21.9	2.4	16.1	8.1-28.1	2.6	13.3	7.0-22.9	3.6	10.7	6.3-16.8	1.8	12.3	5.5-23.2	1.8	8.5	3.9-16.3
	1981-1985	4.2	11.8	7.3-18.1	1.6	13.2	5.6-25.3	2.6	12.0	6.3-20.7	3.2	8.8	5.0-14.3	1.6	12.0	5.0-23.4	1.6	7.3	3.1-14.7
	1986-1990	3.6	9.3	5.5-14.8	1.8	10.5	4.7-19.8	1.8	8.5	3.8-16.3	4.2	11.1	6.8-17.0	2.2	13.3	6.6-23.6	2.0	9.1	4.3-17.0
	1991-1995	4.6	11.3	7.2-17.0	2.8	15.8	8.5-26.4	1.8	7.3	3.3-14.2	3.4	8.5	4.9-13.5	2.0	12.0	5.7-21.8	1.4	5.7	2.3-12.0
	1996-2000	5.4	12.8	8.5-18.7	3.6	19.3	11.3-30.5	1.8	7.5	3.4-14.6	5.6	13.3	8.8-19.2	3.2	17.3	9.8-28.1	2.4	9.6	4.9-17.1
	2001-2005	4.4	9.6	6.0-14.7	1.4	6.1	2.4-12.8	3.0	12.9	7.2-21.6	3.2	7.3	4.2-11.9	1.2	5.5	2.0-12.2	2.0	8.9	4.2-16.6
	2006-2010	8.2	16.8	12.0-23.0	5.6	23.7	15.5-34.8	2.6	9.7	5.1-17.2	6.0	12.3	8.2-17.7	3.8	16.5	9.7-26.3	2.2	8.1	4.0-15.1
	2011-2015	6.8	14.3	9.9-20.2	3.8	17.2	10.2-27.3	3.0	11.7	6.5-19.8	6.4	13.4	9.1-19.1	4.2	19.0	11.6-29.6	2.2	8.8	4.4-16.4
2016-2020	9.6	17.6	12.9-23.6	5.2	18.5	11.9-27.8	4.4	16.3	10.1-25.3	8.2	15.0	10.7-20.6	4.4	16.2	9.9-25.3	3.8	14.3	8.5-23.1	
Larynx	1976-1980	1.6	4.5	1.9-8.9	1.4	9.0	3.5-18.5	0.2	1.1	0.0-6.0	0.2	0.5	0.0-3.0	0.0	0.0	0.0-4.3	0.2	1.0	0.0-5.8
	1981-1985	2.0	4.8	2.3-8.9	1.8	9.3	4.2-17.8	0.2	0.8	0.0-5.3	0.4	1.0	0.1-3.6	0.4	2.2	0.3-8.0	0.0	0.0	0.0-3.9
	1986-1990	2.0	5.1	2.4-9.4	2.0	10.6	5.0-19.6	0.0	0.0	0.0-3.8	0.6	1.4	0.3-4.1	0.6	2.9	0.6-8.7	0.0	0.0	0.0-3.8
	1991-1995	2.2	5.7	2.9-10.2	1.4	8.8	3.5-18.0	0.8	3.6	1.0-9.5	0.6	1.5	0.3-4.3	0.4	2.3	0.3-7.9	0.2	0.8	0.0-5.1
	1996-2000	1.2	2.9	1.1-6.3	1.0	4.7	1.5-11.0	0.2	0.9	0.0-5.3	0.4	0.9	0.1-3.4	0.4	1.9	0.2-7.0	0.0	0.0	0.0-3.7
	2001-2005	2.6	5.6	3.0-9.6	2.0	9.0	4.2-16.7	0.6	2.3	0.5-7.4	1.0	2.3	0.7-5.4	0.8	3.8	1.0-9.7	0.2	0.9	0.0-5.4
	2006-2010	1.0	2.1	0.7-5.0	1.0	4.4	1.4-10.8	0.0	0.0	0.0-3.9	0.6	1.1	0.2-3.6	0.6	2.8	0.5-8.4	0.0	0.0	0.0-3.9
	2011-2015	2.2	3.7	1.8-6.9	1.4	4.2	1.7-9.4	0.8	3.3	0.9-9.1	0.8	1.6	0.4-4.5	0.6	2.6	0.5-8.2	0.2	1.0	0.0-5.8
2016-2020	2.2	3.5	1.7-6.7	1.8	5.7	2.5-11.7	0.4	1.4	0.2-6.3	1.2	2.2	0.8-5.2	1.0	3.8	1.2-9.5	0.2	0.7	0.0-5.2	
Lung and bronchus	1976-1980	21.0	58.7	47.8-71.2	16.2	102.1	80.4-127.6	4.8	26.5	16.8-39.8	16.2	44.9	35.5-55.9	12.4	78.8	59.8-101.5	3.8	19.3	11.5-30.5
	1981-1985	23.8	62.3	51.5-74.7	17.6	104.0	82.9-128.7	6.2	30.7	20.6-44.1	19.2	52.4	42.3-64.1	15.4	95.0	74.5-119.1	3.8	20.4	12.0-32.1
	1986-1990	23.2	61.7	50.9-74.1	15.4	91.6	72.0-114.8	7.8	38.4	27.1-52.8	19.6	51.7	41.9-63.1	12.8	78.8	60.3-101.0	6.8	32.4	22.3-45.6
	1991-1995	25.8	64.2	53.5-76.3	17.6	96.8	77.4-119.5	8.2	36.8	26.2-50.2	22.8	56.7	46.8-68.2	15.0	83.3	65.3-104.7	7.8	35.1	24.8-48.4
	1996-2000	30.8	73.2	62.1-85.7	20.2	109.5	88.7-133.4	10.6	48.6	36.4-63.8	21.8	51.9	42.7-62.7	14.4	80.2	62.4-101.2	7.4	32.6	22.9-45.2
	2001-2005	32.0	72.3	61.5-84.5	18.6	95.2	76.4-117.0	13.4	56.1	43.4-71.7	24.4	55.1	45.7-65.9	14.6	74.2	57.8-93.7	9.8	40.1	29.6-53.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Wyoming**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	33.8	71.6	61.1-83.5	18.8	85.7	68.7-105.6	15.0	62.0	48.6-78.2	23.6	49.6	41.0-59.7	13.8	67.4	52.0-85.9	9.8	39.2	28.9-52.3
	2011-2015	37.4	74.2	63.7-85.9	18.2	75.7	60.3-93.8	19.2	74.0	59.7-91.1	25.0	51.3	42.6-61.4	11.6	51.1	38.5-66.7	13.4	51.3	39.6-65.8
	2016-2020	35.8	62.6	53.6-72.9	19.2	69.0	55.5-85.1	16.6	56.5	44.8-70.9	20.4	36.8	29.9-45.0	12.6	48.6	36.9-63.0	7.8	27.0	19.1-37.7
Melanoma of the skin	1976-1980	1.2	3.8	1.3-8.1	0.6	4.2	0.8-11.8	0.6	3.8	0.7-10.9	1.0	3.0	0.9-7.1	0.8	5.2	1.3-13.3	0.2	0.9	0.0-5.4
	1981-1985	4.0	11.1	6.6-17.2	1.8	10.1	4.5-19.5	2.2	11.8	5.8-21.3	0.8	1.9	0.5-5.1	0.2	1.2	0.0-6.5	0.6	2.7	0.6-8.3
	1986-1990	4.8	11.5	7.3-17.3	2.4	11.7	5.9-20.9	2.4	11.5	5.9-20.3	1.2	3.2	1.1-6.9	0.8	4.1	1.0-10.8	0.4	2.1	0.2-7.5
	1991-1995	4.4	10.6	6.6-16.1	2.2	11.8	5.7-21.4	2.2	10.0	4.9-18.0	1.6	4.3	1.8-8.3	1.0	5.3	1.7-12.4	0.6	2.9	0.6-8.7
	1996-2000	5.4	12.8	8.4-18.6	3.0	17.2	9.3-28.5	2.4	11.1	5.7-19.6	1.2	2.8	1.0-6.1	0.8	4.2	1.0-11.0	0.4	1.7	0.2-6.6
	2001-2005	6.6	14.6	10.0-20.5	4.4	21.5	13.2-33.0	2.2	9.5	4.7-17.4	0.8	1.9	0.5-4.8	0.4	3.1	0.4-9.9	0.4	1.8	0.2-6.9
	2006-2010	10.8	23.0	17.2-30.2	5.8	24.9	16.4-36.3	5.0	23.9	15.2-35.7	1.2	2.8	1.0-6.1	0.8	3.2	0.9-8.7	0.4	2.2	0.3-8.0
	2011-2015	11.2	22.8	17.1-29.8	6.6	25.8	17.5-36.8	4.6	22.0	13.6-33.7	1.2	2.5	0.9-5.7	0.8	3.1	0.8-8.6	0.4	1.5	0.2-6.3
	2016-2020	16.6	31.1	24.5-39.0	10.8	39.2	29.1-51.9	5.8	24.2	15.7-35.9	2.2	4.2	2.1-7.8	1.2	5.2	1.8-11.8	1.0	4.2	1.3-10.6
Female breast	1976-1980							13.2	73.2	56.2-93.6							8.6	48.4	34.7-65.5
	1981-1985							20.6	105.7	85.8-128.8							7.8	40.3	28.4-55.4
	1986-1990							26.0	133.5	111.1-159.2							6.6	32.5	22.2-46.0
	1991-1995							27.2	131.9	110.3-156.5							6.0	27.2	18.3-39.1
	1996-2000							29.8	135.5	114.4-159.4							5.2	22.3	14.5-33.0
	2001-2005							32.0	138.0	117.3-161.4							6.2	26.0	17.6-37.3
	2006-2010							34.8	145.9	124.6-169.9							6.2	24.7	16.6-35.7
	2011-2015							40.0	168.5	145.0-194.9							4.4	16.4	10.2-25.5
	2016-2020							39.6	156.7	134.3-182.1							5.2	18.3	11.9-27.5
Cervix uteri	1976-1980							3.2	18.6	10.5-30.3							0.6	3.4	0.7-10.0
	1981-1985							2.4	13.3	6.7-23.5							1.2	6.2	2.2-13.6
	1986-1990							1.2	5.7	2.0-12.6							0.8	3.9	1.0-10.1
	1991-1995							2.6	12.8	6.8-22.0							0.6	3.3	0.7-9.5
	1996-2000							2.6	12.7	6.8-21.8							0.2	1.0	0.0-5.5
	2001-2005							1.4	6.2	2.4-13.1							1.2	4.8	1.7-10.9
	2006-2010							2.0	9.7	4.5-18.2							0.2	0.9	0.0-5.5
	2011-2015							1.0	4.8	1.4-11.9							0.2	1.0	0.0-5.9
	2016-2020							2.0	9.3	4.2-18.1							0.6	2.1	0.4-7.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Wyoming**

		Incidence									Mortality									
		Males and Females			Males			Females			Males and Females			Males			Females			
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	
Corpus uteri and uterus, not otherwise specified	1976-1980							4.8	26.7	16.9-39.9							1.0	4.8	1.5-11.4	
	1981-1985							4.0	21.4	12.8-33.4							0.4	1.9	0.2-7.0	
	1986-1990							4.2	20.4	12.4-31.4							0.4	1.6	0.2-6.4	
	1991-1995							6.2	30.0	20.2-42.8							1.0	4.0	1.3-9.7	
	1996-2000							8.0	36.7	26.2-50.2							0.4	1.9	0.2-7.0	
	2001-2005							7.6	33.5	23.7-46.3							1.4	6.0	2.4-12.8	
	2006-2010							7.6	32.4	22.7-45.0							1.0	3.9	1.2-9.7	
	2011-2015								10.4	37.9	28.2-50.3							2.6	9.3	4.9-16.6
	2016-2020								10.6	38.7	28.7-51.6							1.2	4.1	1.5-10.0
Ovary	1976-1980							4.0	22.9	13.8-35.5							3.0	16.0	8.9-26.5	
	1981-1985							4.0	20.1	12.1-31.4							1.8	9.4	4.2-18.1	
	1986-1990							3.6	18.3	10.7-29.1							2.2	11.5	5.6-20.8	
	1991-1995							3.0	15.1	8.3-25.0							3.0	13.3	7.4-22.2	
	1996-2000							4.8	21.7	13.8-32.5							2.8	12.3	6.7-21.0	
	2001-2005							3.8	16.8	10.0-26.5							3.4	13.5	7.8-22.1	
	2006-2010							5.6	24.0	15.8-35.2							2.8	12.1	6.6-20.7	
	2011-2015							3.6	14.3	8.3-23.4							2.0	7.7	3.7-14.9	
	2016-2020								2.0	7.7	3.5-15.3							1.6	5.4	2.3-11.7
Prostate	1976-1980				11.6	80.0	60.1-103.9							2.8	21.7	11.6-36.3				
	1981-1985				15.2	104.4	81.8-131.1							5.4	37.0	24.0-54.1				
	1986-1990				16.0	97.5	76.7-121.9							4.4	29.6	18.3-44.8				
	1991-1995				27.0	166.3	138.8-197.5							5.6	36.8	24.1-53.4				
	1996-2000				32.4	179.8	152.9-210.0							5.6	34.9	23.1-50.3				
	2001-2005				34.8	172.4	147.2-200.6							3.8	22.8	13.6-35.6				
	2006-2010				44.2	188.3	163.5-215.9							4.2	25.8	15.9-39.2				
	2011-2015				40.0	153.4	131.9-177.6							3.8	20.2	12.0-31.7				
	2016-2020				42.0	141.8	122.6-163.4							4.6	21.1	13.2-32.1				
Testis	1976-1980				1.4	6.1	2.4-13.2							0.2	0.7	0.0-5.4				
	1981-1985				0.8	3.2	0.8-9.0							0.0	0.0	0.0-4.2				
	1986-1990				1.8	7.1	3.1-14.3							0.2	1.1	0.0-6.1				
	1991-1995				1.2	3.8	1.4-9.2							0.0	0.0	0.0-3.8				
	1996-2000				2.4	9.7	4.8-17.7							0.0	0.0	0.0-3.6				
	2001-2005				1.6	6.0	2.6-12.2							0.0	0.0	0.0-3.6				

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Wyoming**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2006-2010				1.4	5.9	2.3-12.5							0.0	0.0	0.0-3.6			
	2011-2015				0.2	0.8	0.0-5.0							0.0	0.0	0.0-3.7			
	2016-2020				1.4	5.9	2.3-12.7							0.2	1.1	0.0-5.8			
Urinary bladder (incl. in situ)	1976-1980	6.8	19.9	13.7-27.9	4.8	31.6	20.0-47.3	2.0	10.1	4.8-18.7	4.2	12.7	7.8-19.3	3.0	22.9	12.6-37.6	1.2	6.0	2.2-13.3
	1981-1985	8.6	24.5	17.6-33.0	6.0	39.3	26.2-56.2	2.6	13.7	7.1-23.6	2.2	6.5	3.2-11.5	1.4	8.9	3.6-18.2	0.8	3.8	1.0-10.0
	1986-1990	8.8	22.7	16.5-30.6	6.0	33.7	22.5-48.4	2.8	12.3	6.7-20.8	2.2	5.7	2.8-10.1	1.0	5.0	1.6-11.8	1.2	5.1	1.9-11.5
	1991-1995	9.0	22.2	16.2-29.7	7.6	45.8	32.0-63.1	1.4	6.4	2.5-13.4	2.0	4.9	2.4-9.0	1.6	11.2	4.6-21.7	0.4	1.6	0.2-6.2
	1996-2000	12.4	29.4	22.5-37.7	10.6	57.1	42.5-74.9	1.8	7.1	3.2-14.0	1.0	2.4	0.8-5.6	0.4	2.1	0.3-7.4	0.6	2.2	0.5-7.2
	2001-2005	14.4	32.5	25.4-41.1	10.8	59.1	44.0-77.3	3.6	14.9	8.8-23.9	2.0	4.6	2.2-8.5	1.6	8.4	3.5-16.5	0.4	1.6	0.2-6.4
	2006-2010	20.0	43.5	35.3-53.1	15.8	77.6	60.9-97.3	4.2	17.4	10.7-27.0	3.4	7.1	4.1-11.6	2.6	13.9	7.3-23.9	0.8	2.5	0.7-7.4
	2011-2015	14.0	28.2	21.8-35.9	12.0	51.5	38.8-67.2	2.0	9.0	4.2-17.1	1.8	3.6	1.6-7.0	1.4	6.4	2.4-13.6	0.4	2.0	0.2-7.7
2016-2020	14.4	27.5	21.4-34.9	11.4	46.3	34.7-60.7	3.0	11.1	6.1-19.0	3.4	7.1	4.1-11.6	2.6	12.7	6.7-21.9	0.8	3.3	0.9-9.2	
Kidney and renal pelvis	1976-1980	3.2	9.3	5.3-15.1	1.8	10.9	4.8-20.8	1.4	7.6	3.0-15.7	1.6	4.6	2.0-9.1	1.0	6.3	2.0-14.6	0.6	3.2	0.7-9.5
	1981-1985	2.8	7.1	3.8-12.0	1.2	5.8	2.1-13.0	1.6	8.7	3.6-17.3	1.6	4.3	1.8-8.6	1.2	7.7	2.8-16.7	0.4	1.9	0.2-7.1
	1986-1990	4.4	11.6	7.2-17.6	2.2	12.6	6.2-22.5	2.2	10.6	5.2-19.2	2.2	5.7	2.8-10.2	0.8	4.3	1.1-11.2	1.4	7.2	2.8-15.0
	1991-1995	5.8	14.7	9.8-21.1	4.0	23.5	14.1-36.3	1.8	8.6	3.9-16.6	2.8	7.2	3.9-12.1	1.8	11.0	4.8-20.9	1.0	4.6	1.5-10.9
	1996-2000	5.6	13.3	8.8-19.2	3.4	17.2	9.9-27.5	2.2	10.2	5.1-18.4	1.8	4.3	2.0-8.1	1.0	5.4	1.7-12.5	0.8	3.7	1.0-9.6
	2001-2005	6.6	14.1	9.7-19.8	4.8	20.5	13.0-30.9	1.8	7.3	3.3-14.3	2.4	5.2	2.7-9.2	1.2	4.8	1.7-10.8	1.2	5.0	1.8-11.4
	2006-2010	10.2	20.8	15.4-27.5	6.8	27.7	18.9-39.4	3.4	13.7	7.9-22.5	3.2	6.7	3.8-11.0	2.0	9.3	4.3-17.3	1.2	3.8	1.4-9.1
	2011-2015	11.2	20.6	15.5-27.1	7.0	27.2	18.6-38.5	4.2	15.3	9.3-24.2	1.8	3.6	1.6-7.0	1.6	7.1	2.9-14.6	0.2	1.0	0.0-5.8
2016-2020	7.8	14.9	10.4-20.7	6.2	24.5	16.4-35.5	1.6	5.5	2.4-11.8	1.6	2.8	1.2-6.0	1.2	4.0	1.4-9.6	0.4	1.8	0.2-7.4	
Brain and other central nervous system	1976-1980	2.6	6.9	3.6-11.9	1.2	6.7	2.3-14.9	1.4	7.8	3.0-16.3	2.4	6.1	3.1-10.8	1.2	6.7	2.3-14.9	1.2	5.8	2.1-12.8
	1981-1985	1.0	2.4	0.8-5.9	0.6	3.1	0.6-9.6	0.4	1.8	0.2-7.0	1.4	3.6	1.4-7.6	0.8	4.3	1.2-11.3	0.6	3.2	0.6-9.6
	1986-1990	3.0	7.4	4.1-12.3	1.6	9.0	3.8-17.9	1.4	6.6	2.7-13.8	1.2	3.1	1.1-6.9	0.8	4.7	1.2-12.1	0.4	1.8	0.2-6.7
	1991-1995	3.2	8.0	4.5-13.0	2.4	11.2	5.7-19.7	0.8	4.0	1.1-10.2	1.8	4.6	2.1-8.6	0.8	4.0	1.1-10.2	1.0	5.2	1.7-12.0
	1996-2000	3.2	7.4	4.2-12.1	1.8	8.0	3.6-15.5	1.4	7.0	2.8-14.4	1.8	4.0	1.8-7.7	0.8	3.4	0.9-8.9	1.0	5.0	1.6-11.6
	2001-2005	3.2	7.5	4.2-12.2	2.6	11.9	6.3-20.6	0.6	2.9	0.6-8.7	1.8	3.9	1.8-7.5	1.8	7.8	3.5-15.1	0.0	0.0	0.0-3.8
	2006-2010	5.8	13.4	9.0-19.4	3.4	14.9	8.6-24.1	2.4	11.2	5.7-19.9	2.0	4.5	2.1-8.4	1.0	4.3	1.3-10.4	1.0	4.1	1.3-10.2
	2011-2015	3.6	7.9	4.6-12.7	2.2	10.7	5.2-19.4	1.4	5.4	2.1-11.8	1.8	4.2	1.9-8.1	1.2	5.5	2.0-12.2	0.6	3.0	0.5-9.4
2016-2020	2.4	4.5	2.2-8.3	1.0	3.1	0.9-8.4	1.4	6.1	2.2-13.5	2.2	3.7	1.8-7.0	1.4	3.9	1.6-9.1	0.8	3.3	0.8-9.4	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Wyoming**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	1.6	4.5	1.9-8.9	0.2	0.8	0.0-5.6	1.4	7.9	3.0-16.5	0.4	1.1	0.1-4.0	0.2	1.3	0.0-6.7	0.2	0.9	0.0-5.5
	1981-1985	1.0	2.7	0.8-6.3	0.6	3.3	0.7-9.7	0.4	2.0	0.2-7.5	0.2	0.5	0.0-2.9	0.2	1.2	0.0-6.5	0.0	0.0	0.0-3.9
	1986-1990	0.6	1.4	0.3-4.2	0.0	0.0	0.0-4.0	0.6	3.1	0.6-9.0	0.2	0.5	0.0-2.8	0.0	0.0	0.0-4.0	0.2	0.9	0.0-5.3
	1991-1995	1.2	2.6	0.9-5.8	0.6	3.6	0.7-10.2	0.6	2.6	0.5-8.0	0.2	0.5	0.0-2.7	0.2	1.4	0.0-6.7	0.0	0.0	0.0-3.7
	1996-2000	3.4	7.7	4.5-12.4	1.0	5.2	1.7-12.1	2.4	11.5	5.9-20.2	0.0	0.0	0.0-1.8	0.0	0.0	0.0-3.6	0.0	0.0	0.0-3.7
	2001-2005	3.6	7.6	4.5-12.1	0.8	3.1	0.8-8.4	2.8	13.2	7.2-22.3	0.0	0.0	0.0-1.8	0.0	0.0	0.0-3.6	0.0	0.0	0.0-3.8
	2006-2010	6.2	13.7	9.3-19.6	1.4	6.0	2.3-12.8	4.8	24.2	15.2-36.4	0.4	1.0	0.1-3.5	0.2	1.3	0.0-6.5	0.2	1.0	0.0-5.7
	2011-2015	8.2	17.7	12.5-24.3	1.4	5.4	2.0-11.8	6.8	34.4	23.3-48.8	0.2	0.4	0.0-2.6	0.0	0.0	0.0-3.7	0.2	0.6	0.0-5.0
	2016-2020	6.2	12.8	8.5-18.5	1.2	4.7	1.6-10.9	5.0	22.8	14.3-34.7	0.2	0.3	0.0-2.3	0.0	0.0	0.0-3.8	0.2	0.6	0.0-5.0
Hodgkin lymphoma	1976-1980	1.6	3.6	1.5-7.2	1.0	4.7	1.3-11.8	0.6	2.5	0.5-7.9	0.6	1.5	0.3-4.6	0.4	2.7	0.3-9.5	0.2	0.8	0.0-5.4
	1981-1985	1.2	2.7	0.9-6.1	0.6	2.8	0.5-8.8	0.6	2.5	0.5-7.9	0.2	0.6	0.0-3.2	0.0	0.0	0.0-4.2	0.2	1.0	0.0-5.8
	1986-1990	1.6	4.1	1.7-8.2	1.2	5.6	1.9-12.8	0.4	2.4	0.3-8.2	0.4	1.1	0.1-4.0	0.2	1.3	0.0-6.7	0.2	0.8	0.0-5.2
	1991-1995	2.2	5.2	2.6-9.4	1.0	5.1	1.6-12.0	1.2	6.3	2.3-13.7	0.8	1.5	0.4-4.0	0.4	1.1	0.1-5.4	0.4	1.9	0.2-7.0
	1996-2000	2.0	4.5	2.1-8.3	1.4	5.3	2.1-11.5	0.6	2.9	0.6-8.5	0.2	0.5	0.0-2.7	0.2	1.2	0.0-6.0	0.0	0.0	0.0-3.7
	2001-2005	1.8	4.1	1.9-7.8	0.6	2.2	0.5-7.1	1.2	6.4	2.3-14.0	0.6	1.3	0.3-4.0	0.6	3.6	0.7-10.1	0.0	0.0	0.0-3.8
	2006-2010	1.6	3.3	1.4-6.7	0.8	2.7	0.7-7.5	0.8	4.1	1.1-10.6	0.0	0.0	0.0-1.8	0.0	0.0	0.0-3.6	0.0	0.0	0.0-3.9
	2011-2015	1.8	4.4	2.0-8.4	1.2	5.5	2.0-12.2	0.6	2.9	0.6-8.8	0.0	0.0	0.0-1.9	0.0	0.0	0.0-3.7	0.0	0.0	0.0-4.1
	2016-2020	1.2	2.4	0.8-5.6	0.6	2.2	0.4-7.2	0.6	2.7	0.5-8.8	0.2	0.4	0.0-2.7	0.0	0.0	0.0-3.8	0.2	0.8	0.0-5.6
Non-Hodgkin lymphomas	1976-1980	4.2	12.5	7.7-19.1	2.4	14.1	7.1-25.0	1.8	9.9	4.5-18.8	1.4	4.1	1.6-8.5	1.2	7.6	2.7-16.6	0.2	0.9	0.0-5.7
	1981-1985	5.2	14.0	9.1-20.6	3.0	17.2	9.5-28.7	2.2	10.1	5.0-18.3	1.4	3.9	1.5-8.0	0.6	3.5	0.7-10.2	0.8	3.9	1.1-10.3
	1986-1990	7.8	20.7	14.7-28.4	3.6	20.8	12.1-33.0	4.2	20.8	12.7-32.0	2.8	7.6	4.1-12.7	1.0	6.2	1.9-14.4	1.8	9.7	4.3-18.6
	1991-1995	9.0	22.8	16.6-30.5	5.8	30.5	20.0-44.4	3.2	15.2	8.6-25.0	2.8	6.9	3.8-11.6	2.2	11.6	5.8-20.8	0.6	2.8	0.6-8.3
	1996-2000	8.6	20.1	14.6-27.2	4.6	22.8	14.3-34.5	4.0	17.9	10.9-27.9	2.8	6.7	3.7-11.2	1.6	9.7	4.1-19.0	1.2	4.9	1.8-11.0
	2001-2005	9.2	20.7	15.1-27.7	4.8	23.5	14.9-35.3	4.4	19.6	12.1-30.0	3.0	6.7	3.7-11.1	1.8	10.1	4.5-19.1	1.2	4.4	1.6-10.3
	2006-2010	9.0	18.2	13.2-24.6	5.0	19.7	12.5-29.7	4.0	16.8	10.1-26.4	3.6	8.0	4.7-12.7	1.2	5.2	1.8-11.9	2.4	11.5	5.8-20.4
	2011-2015	11.6	24.2	18.2-31.5	6.0	25.7	17.1-37.3	5.6	22.8	14.9-33.7	2.0	4.0	1.9-7.5	1.0	4.0	1.2-9.9	1.0	3.7	1.2-9.4
	2016-2020	13.2	25.0	19.2-32.3	7.6	28.3	19.7-39.6	5.6	24.3	15.6-36.3	2.8	5.6	3.0-9.6	1.8	8.4	3.7-16.4	1.0	4.2	1.3-10.6
Myeloma	1976-1980	2.8	8.2	4.4-13.7	1.2	8.5	3.0-18.3	1.6	7.6	3.3-15.2	1.0	3.3	1.1-7.6	0.8	6.9	1.9-16.7	0.2	0.9	0.0-5.7
	1981-1985	2.0	5.6	2.7-10.4	0.8	4.6	1.1-12.0	1.2	6.8	2.4-14.9	1.2	3.3	1.2-7.1	0.4	2.7	0.3-9.3	0.8	4.1	1.1-10.7
	1986-1990	1.0	2.8	0.9-6.4	1.0	6.7	2.1-15.4	0.0	0.0	0.0-3.8	0.2	0.6	0.0-3.0	0.2	1.2	0.0-6.2	0.0	0.0	0.0-3.8
	1991-1995	2.2	5.6	2.8-10.0	1.8	10.8	4.8-20.5	0.4	1.8	0.2-6.7	1.0	2.5	0.8-5.9	0.8	6.3	1.7-15.2	0.2	0.8	0.0-5.1
	1996-2000	2.2	5.2	2.6-9.3	0.6	4.1	0.8-11.3	1.6	7.0	3.0-14.0	1.6	3.8	1.6-7.5	0.6	3.1	0.6-9.0	1.0	4.3	1.4-10.4
	2001-2005	4.4	10.0	6.2-15.2	2.6	12.6	6.6-21.8	1.8	6.9	3.1-13.7	2.4	5.3	2.8-9.4	1.2	6.3	2.2-13.7	1.2	4.4	1.6-10.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Wyoming**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	3.2	7.0	4.0-11.5	2.2	10.5	5.1-19.0	1.0	4.1	1.3-10.2	1.6	3.6	1.6-7.2	1.0	5.3	1.7-12.3	0.6	2.3	0.5-7.5
	2011-2015	3.8	8.0	4.8-12.7	2.6	11.6	6.1-20.1	1.2	5.4	1.9-12.3	2.0	4.5	2.1-8.4	1.0	4.7	1.5-11.3	1.0	4.0	1.3-10.0
	2016-2020	3.6	7.2	4.1-11.7	2.2	8.8	4.3-16.2	1.4	6.1	2.2-13.6	1.8	3.4	1.5-6.8	0.8	2.7	0.7-7.8	1.0	4.2	1.3-10.6
Leukemias	1976-1980	4.8	14.0	8.9-20.8	3.0	19.3	10.6-31.9	1.8	9.4	4.2-17.9	2.0	5.9	2.8-10.8	1.2	8.4	2.9-18.3	0.8	4.1	1.1-10.6
	1981-1985	4.8	12.9	8.2-19.2	2.8	17.9	9.6-30.1	2.0	9.0	4.3-16.9	2.4	7.1	3.6-12.3	1.0	7.8	2.5-17.6	1.4	6.2	2.5-13.1
	1986-1990	5.4	13.9	9.1-20.3	3.6	20.5	11.9-32.5	1.8	8.4	3.8-16.3	4.4	11.4	7.1-17.3	2.4	14.3	7.1-25.2	2.0	9.5	4.5-17.7
	1991-1995	5.0	12.4	8.0-18.4	3.2	16.8	9.4-27.7	1.8	8.5	3.8-16.4	3.2	7.9	4.5-12.8	2.0	10.2	4.8-18.8	1.2	5.8	2.1-12.7
	1996-2000	4.6	10.9	6.9-16.3	3.2	17.2	9.7-28.1	1.4	6.1	2.4-12.9	2.6	6.1	3.2-10.4	2.0	10.5	4.9-19.4	0.6	2.5	0.5-7.8
	2001-2005	8.8	20.6	14.9-27.7	4.6	22.6	14.2-34.2	4.2	18.6	11.4-28.8	3.4	7.8	4.6-12.6	1.4	7.0	2.7-14.6	2.0	8.8	4.2-16.6
	2006-2010	7.8	16.4	11.6-22.6	4.6	21.3	13.2-32.4	3.2	13.2	7.4-21.9	2.8	6.2	3.3-10.6	1.6	8.7	3.6-17.1	1.2	4.5	1.6-10.6
	2011-2015	9.6	19.6	14.3-26.3	5.4	21.6	14.0-32.2	4.2	17.5	10.6-27.5	4.0	8.7	5.3-13.6	2.0	8.9	4.2-16.8	2.0	8.4	3.9-16.0
	2016-2020	10.6	20.3	15.0-26.9	7.4	30.5	21.0-42.8	3.2	12.3	6.8-20.8	2.6	5.2	2.7-9.1	1.4	5.6	2.2-12.0	1.2	5.0	1.8-11.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Yates**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
All Invasive Malignant Tumors	1976-1980	87.0	380.2	344.5-418.8	49.6	517.7	452.4-589.5	37.4	292.4	250.6-339.4	45.6	201.9	175.9-230.8	27.2	298.7	248.1-356.2	18.4	141.6	113.2-175.1
	1981-1985	96.0	403.6	367.5-442.3	49.0	496.9	434.5-565.7	47.0	353.4	308.0-403.7	46.4	192.5	168.1-219.5	24.0	242.5	199.5-291.8	22.4	163.5	133.8-198.1
	1986-1990	130.4	526.5	486.2-569.3	71.4	673.2	603.3-749.0	59.0	437.0	387.0-491.8	63.6	252.3	225.0-282.0	34.2	330.3	281.3-385.3	29.4	201.7	169.8-238.2
	1991-1995	139.0	526.7	487.8-568.0	75.4	637.1	573.1-706.3	63.6	455.7	405.7-510.2	60.2	223.8	199.0-251.0	32.2	289.7	245.8-339.2	28.0	185.6	155.5-220.1
	1996-2000	141.2	499.5	463.1-538.2	73.4	570.4	512.8-632.8	67.8	452.7	405.0-504.8	61.0	209.8	186.8-235.0	31.6	254.4	215.8-298.1	29.4	179.3	151.1-211.6
	2001-2005	151.0	511.7	475.6-550.1	82.6	602.2	544.7-664.3	68.4	445.3	398.5-496.4	56.0	185.1	164.0-208.4	28.8	215.7	181.4-254.7	27.2	163.7	137.0-194.5
	2006-2010	152.4	479.9	445.7-516.2	80.2	537.9	485.2-595.0	72.2	428.9	384.6-477.2	55.2	170.7	150.9-192.6	28.8	195.3	164.2-230.9	26.4	152.3	126.8-181.9
	2011-2015	162.8	497.9	462.6-535.4	81.6	511.3	461.0-566.0	81.2	492.2	442.4-546.2	61.0	180.7	160.3-203.2	31.4	208.6	176.3-245.5	29.6	159.2	133.5-188.9
	2016-2020	176.6	498.6	464.1-535.2	96.6	553.5	502.9-608.3	80.0	457.6	410.0-509.5	56.2	150.7	133.0-170.4	28.2	164.2	137.4-195.2	28.0	142.1	118.5-169.6
Oral cavity and pharynx	1976-1980	3.4	16.2	9.2-26.3	2.6	29.0	15.1-50.2	0.8	5.6	1.5-15.5	1.6	7.8	3.2-15.7	1.4	16.9	6.4-35.3	0.2	1.4	0.0-9.3
	1981-1985	2.0	9.5	4.5-17.7	1.4	14.1	5.4-29.7	0.6	6.6	1.2-18.8	0.8	4.1	1.1-10.4	0.8	9.8	2.6-24.7	0.0	0.0	0.0-6.8
	1986-1990	3.0	12.2	6.8-20.4	1.8	18.8	8.4-36.0	1.2	8.1	2.9-18.3	1.2	4.9	1.8-10.8	1.0	11.3	3.5-26.3	0.2	1.3	0.0-8.6
	1991-1995	4.6	17.8	11.2-27.0	3.4	27.5	15.9-44.6	1.2	8.9	3.2-19.8	1.8	6.8	3.1-13.1	1.4	11.3	4.5-23.7	0.4	2.6	0.3-10.2
	1996-2000	2.8	10.0	5.4-16.9	1.2	10.3	3.7-22.5	1.6	10.0	4.3-20.3	0.8	2.6	0.7-7.1	0.2	1.3	0.0-8.5	0.6	3.1	0.6-10.5
	2001-2005	3.0	10.6	5.9-17.8	2.6	19.9	10.4-34.5	0.4	2.6	0.3-10.1	1.2	4.1	1.5-9.3	0.8	5.9	1.6-15.8	0.4	2.4	0.3-9.6
	2006-2010	2.0	5.8	2.7-11.0	1.4	8.5	3.4-18.5	0.6	3.1	0.6-10.4	0.6	1.7	0.4-5.6	0.2	1.5	0.0-8.9	0.4	2.1	0.3-9.0
	2011-2015	4.6	15.2	9.2-23.6	3.4	22.3	12.6-37.0	1.2	9.1	2.8-21.2	0.2	0.5	0.0-3.8	0.2	1.0	0.0-7.9	0.0	0.0	0.0-6.1
	2016-2020	5.8	19.2	12.3-28.7	4.2	26.3	15.8-41.7	1.6	13.8	5.4-28.2	1.0	2.4	0.8-6.5	0.8	4.1	1.1-12.3	0.2	0.8	0.0-7.2
Esophagus	1976-1980	1.6	7.3	3.0-14.8	0.8	7.0	1.9-18.9	0.8	8.2	2.0-21.2	1.2	5.8	2.0-13.0	0.6	6.0	1.1-18.4	0.6	6.9	1.3-19.6
	1981-1985	0.6	2.5	0.5-7.5	0.4	4.9	0.5-17.2	0.2	1.4	0.0-9.1	0.4	1.7	0.2-6.4	0.2	3.2	0.1-14.9	0.2	1.4	0.0-9.1
	1986-1990	1.8	7.5	3.4-14.5	1.4	13.0	5.1-27.2	0.4	3.0	0.4-11.6	1.4	5.6	2.2-11.8	1.0	7.7	2.5-18.8	0.4	3.8	0.4-13.5
	1991-1995	1.8	7.2	3.2-13.9	1.0	8.5	2.7-20.2	0.8	6.1	1.5-16.1	1.0	3.9	1.2-9.2	0.4	3.2	0.4-12.2	0.6	4.2	0.8-12.9
	1996-2000	0.4	1.3	0.2-5.2	0.2	1.3	0.0-8.5	0.2	1.2	0.0-7.7	1.0	3.5	1.1-8.3	0.6	4.3	0.9-13.3	0.4	2.2	0.3-9.2
	2001-2005	1.0	3.3	1.1-8.1	1.0	7.1	2.3-17.3	0.0	0.0	0.0-5.9	1.4	4.9	2.0-10.3	0.8	5.9	1.6-15.7	0.6	3.9	0.8-12.1
	2006-2010	1.8	5.5	2.5-10.8	1.2	7.8	2.8-17.8	0.6	3.3	0.7-10.8	1.4	4.4	1.8-9.4	0.8	5.3	1.4-14.5	0.6	3.5	0.7-11.2
	2011-2015	3.2	8.9	5.0-15.0	2.2	13.7	6.7-25.6	1.0	4.7	1.5-12.6	2.0	6.0	2.8-11.7	1.8	11.9	5.3-23.5	0.2	1.0	0.0-7.5
	2016-2020	3.0	8.4	4.6-14.7	2.2	13.5	6.4-25.6	0.8	4.1	1.1-12.0	3.6	9.4	5.4-15.7	2.8	16.2	8.5-28.7	0.8	3.3	0.9-10.6
Stomach	1976-1980	2.6	12.1	6.3-21.1	1.6	20.2	8.3-40.2	1.0	7.0	2.3-17.4	2.2	10.6	5.2-19.3	1.4	19.1	7.4-39.0	0.8	5.6	1.5-15.5
	1981-1985	1.8	8.2	3.7-15.7	1.2	14.3	5.0-31.2	0.6	5.0	0.9-15.2	1.0	4.3	1.4-10.4	0.4	4.4	0.5-15.8	0.6	3.9	0.8-12.7
	1986-1990	0.4	1.4	0.2-5.5	0.4	2.8	0.3-11.7	0.0	0.0	0.0-6.4	0.4	1.9	0.2-6.9	0.4	3.9	0.4-14.5	0.0	0.0	0.0-6.4
	1991-1995	1.8	6.7	3.0-12.9	1.6	15.4	6.5-30.4	0.2	1.3	0.0-8.2	1.2	4.4	1.6-9.7	1.0	9.7	3.1-22.5	0.2	1.3	0.0-8.2
	1996-2000	1.4	4.8	1.9-10.2	0.6	4.3	0.9-13.2	0.8	4.6	1.2-13.0	0.6	2.2	0.4-6.6	0.0	0.0	0.0-6.3	0.6	3.7	0.7-11.8
	2001-2005	2.8	9.6	5.2-16.3	1.4	10.0	4.0-21.2	1.4	8.8	3.5-18.9	0.8	2.6	0.7-7.0	0.4	2.7	0.3-10.8	0.4	2.2	0.3-9.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.



**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Yates**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Stomach	2006-2010	0.4	1.2	0.1-5.0	0.4	2.5	0.3-10.2	0.0	0.0	0.0-5.9	0.6	1.9	0.4-6.0	0.4	2.5	0.3-10.2	0.2	1.3	0.0-8.1
	2011-2015	1.4	4.2	1.7-9.1	1.2	8.4	3.1-19.0	0.2	1.0	0.0-7.5	0.2	0.7	0.0-4.3	0.2	1.5	0.0-8.9	0.0	0.0	0.0-6.1
	2016-2020	2.0	4.7	2.2-9.4	1.2	6.3	2.3-15.2	0.8	3.3	0.9-10.6	0.6	1.4	0.3-5.0	0.2	0.8	0.0-7.6	0.4	1.8	0.2-8.6
Colorectal	1976-1980	16.4	71.1	56.3-88.7	9.0	98.9	70.9-134.0	7.4	54.7	38.3-76.2	7.0	30.7	21.3-43.0	3.6	43.0	24.9-68.5	3.4	24.0	13.9-39.2
	1981-1985	16.8	67.8	53.9-84.3	8.4	86.8	61.7-118.5	8.4	56.5	40.6-77.1	6.2	25.7	17.4-36.8	2.8	29.4	15.6-50.1	3.4	23.8	13.7-39.0
	1986-1990	19.2	78.3	63.2-96.1	10.4	102.9	76.1-135.9	8.8	62.5	44.9-85.0	8.8	35.7	25.8-48.1	5.2	52.2	33.7-77.0	3.6	24.9	14.5-40.3
	1991-1995	19.2	71.4	57.7-87.5	9.6	81.0	59.4-108.0	9.6	64.1	46.8-85.9	6.6	25.2	17.2-35.6	2.8	24.9	13.5-41.9	3.8	26.2	15.5-41.5
	1996-2000	19.4	67.1	54.4-82.1	9.8	76.6	56.4-101.9	9.6	61.4	45.0-82.2	8.4	29.0	20.9-39.4	4.0	33.0	20.0-51.4	4.4	26.7	16.5-41.3
	2001-2005	18.0	60.1	48.2-74.2	11.6	88.5	66.8-115.2	6.4	38.6	26.1-55.3	6.6	22.0	15.1-31.1	3.0	23.6	13.0-39.4	3.6	21.4	12.5-34.8
	2006-2010	12.2	37.1	28.2-48.0	6.8	44.2	30.3-62.7	5.4	29.9	19.6-44.3	4.6	13.9	8.8-21.2	3.0	20.2	11.2-34.1	1.6	9.0	3.8-18.6
	2011-2015	13.2	42.5	32.4-54.9	5.6	37.5	24.4-55.4	7.6	46.3	31.8-65.3	5.0	16.1	10.0-24.5	2.6	17.5	9.0-31.1	2.4	14.1	6.7-26.5
	2016-2020	9.2	24.8	17.9-34.0	5.6	31.9	20.7-47.7	3.6	18.1	10.4-30.2	4.0	10.2	6.1-16.4	1.8	9.9	4.4-20.2	2.2	10.8	5.2-20.9
Colon excluding rectum	1976-1980	12.2	53.5	40.7-69.1	6.6	75.4	50.9-107.2	5.6	40.7	26.8-59.7	4.6	20.8	13.1-31.4	2.4	32.3	16.4-56.0	2.2	15.3	7.6-28.3
	1981-1985	13.8	55.7	43.2-70.8	7.8	80.0	56.1-110.5	6.0	40.6	27.3-58.9	6.0	25.0	16.8-36.0	2.6	28.0	14.5-48.5	3.4	23.8	13.7-39.0
	1986-1990	15.8	64.7	51.0-81.0	7.8	80.0	56.2-110.2	8.0	55.7	39.3-76.9	6.8	27.5	19.0-38.6	4.0	40.6	24.4-63.1	2.8	19.6	10.5-33.8
	1991-1995	16.0	59.9	47.4-74.9	7.8	65.2	46.0-89.7	8.2	55.4	39.3-76.0	6.0	23.1	15.5-33.2	2.4	21.6	11.1-37.8	3.6	25.2	14.8-40.5
	1996-2000	15.2	52.4	41.2-65.7	7.4	58.8	41.2-81.6	7.8	48.9	34.5-67.7	7.2	24.7	17.3-34.4	3.4	28.8	16.6-46.3	3.8	22.3	13.3-35.8
	2001-2005	14.4	47.9	37.4-60.6	9.4	72.5	52.9-97.1	5.0	29.3	18.8-44.1	5.8	18.8	12.6-27.2	2.6	21.0	11.0-36.4	3.2	17.7	10.1-29.5
	2006-2010	8.0	24.1	17.2-33.1	4.4	28.5	17.8-44.0	3.6	20.0	11.8-32.4	4.0	12.0	7.3-18.9	2.6	17.5	9.2-30.7	1.4	7.7	3.1-16.9
	2011-2015	10.0	32.5	23.6-43.7	4.6	31.8	19.7-48.9	5.4	32.7	20.7-49.5	4.2	14.1	8.4-22.3	2.2	15.7	7.6-29.0	2.0	12.6	5.5-24.7
	2016-2020	5.6	15.2	9.8-22.7	2.6	14.8	7.5-26.9	3.0	14.6	7.9-25.7	3.2	7.9	4.5-13.6	1.4	7.1	2.8-16.3	1.8	8.6	3.8-18.0
Rectum & rectosigmoid	1976-1980	4.2	17.6	10.8-27.3	2.4	23.5	11.6-42.2	1.8	14.0	6.3-27.3	2.4	9.9	5.1-17.7	1.2	10.7	3.9-24.3	1.2	8.6	3.2-19.7
	1981-1985	3.0	12.1	6.7-20.3	0.6	6.8	1.3-19.7	2.4	15.8	8.2-28.5	0.2	0.7	0.0-4.6	0.2	1.4	0.0-9.9	0.0	0.0	0.0-6.8
	1986-1990	3.4	13.6	7.8-22.1	2.6	22.9	12.1-39.7	0.8	6.8	1.7-17.8	2.0	8.2	3.9-15.3	1.2	11.6	4.2-25.4	0.8	5.4	1.4-14.7
	1991-1995	3.2	11.5	6.6-18.9	1.8	15.8	7.1-30.4	1.4	8.7	3.4-18.8	0.6	2.1	0.4-6.4	0.4	3.3	0.4-12.4	0.2	0.9	0.0-7.3
	1996-2000	4.2	14.8	9.1-22.8	2.4	17.8	9.1-31.6	1.8	12.5	5.6-24.1	1.2	4.3	1.6-9.6	0.6	4.3	0.9-13.2	0.6	4.4	0.9-13.2
	2001-2005	3.6	12.2	7.1-19.6	2.2	16.0	7.9-29.3	1.4	9.3	3.6-19.9	0.8	3.2	0.8-8.3	0.4	2.6	0.3-10.4	0.4	3.8	0.4-13.0
	2006-2010	4.2	13.0	7.9-20.3	2.4	15.7	7.8-28.5	1.8	9.9	4.5-19.8	0.6	1.9	0.4-6.1	0.4	2.7	0.3-10.6	0.2	1.3	0.0-8.0
	2011-2015	3.2	10.0	5.6-16.7	1.0	5.7	1.8-14.7	2.2	13.5	6.5-25.3	0.8	1.9	0.5-5.7	0.4	1.8	0.2-8.8	0.4	1.6	0.2-8.1
	2016-2020	3.6	9.7	5.5-16.0	3.0	17.1	9.3-29.6	0.6	3.5	0.6-11.6	0.8	2.2	0.6-6.4	0.4	2.8	0.3-11.0	0.4	2.2	0.3-9.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Yates**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Liver / intrahepatic bile duct	1976-1980	0.2	0.7	0.0-4.9	0.2	1.7	0.0-11.0	0.0	0.0	0.0-7.1	0.2	0.9	0.0-5.3	0.0	0.0	0.0-8.2	0.2	1.3	0.0-9.0
	1981-1985	0.8	3.9	1.0-10.0	0.2	2.0	0.0-11.4	0.6	5.4	0.9-16.5	0.6	3.0	0.5-8.8	0.2	2.0	0.0-11.4	0.4	4.1	0.4-14.8
	1986-1990	1.4	5.4	2.2-11.5	0.6	5.7	1.1-16.9	0.8	5.9	1.6-15.8	0.8	2.9	0.8-7.9	0.4	3.7	0.4-13.9	0.4	2.7	0.3-10.8
	1991-1995	0.4	1.4	0.2-5.3	0.2	1.4	0.0-9.0	0.2	1.3	0.0-8.2	0.8	3.1	0.8-8.1	0.6	5.3	1.1-15.6	0.2	1.2	0.0-7.9
	1996-2000	0.8	2.9	0.8-7.5	0.4	3.0	0.4-11.4	0.4	2.5	0.3-10.0	0.4	1.3	0.2-5.1	0.2	1.4	0.0-8.6	0.2	0.9	0.0-7.2
	2001-2005	0.8	2.6	0.7-7.0	0.8	5.6	1.5-15.1	0.0	0.0	0.0-5.9	1.2	3.9	1.4-8.8	0.8	5.6	1.5-15.1	0.4	2.2	0.3-9.2
	2006-2010	1.6	4.8	2.1-9.9	1.2	7.4	2.7-17.1	0.4	1.9	0.2-8.6	1.6	4.8	2.1-9.9	1.2	8.1	2.9-18.4	0.4	1.9	0.2-8.6
	2011-2015	2.0	6.1	2.9-11.8	0.8	4.6	1.2-13.3	1.2	7.6	2.7-17.4	1.0	2.5	0.8-6.6	0.4	2.1	0.2-9.5	0.6	2.5	0.5-9.5
	2016-2020	1.8	4.4	2.0-9.0	0.8	4.0	1.0-12.1	1.0	4.9	1.6-13.0	1.6	4.7	2.0-9.9	0.6	3.6	0.7-11.9	1.0	5.6	1.7-14.4
Pancreas	1976-1980	2.0	8.6	4.0-16.2	1.6	15.9	6.6-32.1	0.4	3.0	0.4-11.9	3.0	13.3	7.2-22.4	2.0	21.4	9.9-40.1	1.0	6.7	2.2-16.7
	1981-1985	3.6	14.3	8.3-23.0	1.8	15.6	7.0-30.7	1.8	13.9	6.1-27.3	4.0	16.4	9.9-25.7	1.8	17.2	7.6-33.4	2.2	16.8	8.1-31.1
	1986-1990	2.4	9.5	4.9-16.9	0.4	3.6	0.4-13.3	2.0	13.5	6.4-25.6	2.2	8.9	4.4-16.0	0.4	3.6	0.4-13.3	1.8	12.6	5.7-24.6
	1991-1995	2.6	9.5	5.0-16.4	1.6	12.9	5.5-26.0	1.0	5.8	1.9-14.6	3.2	12.1	6.9-19.9	1.8	16.1	7.2-30.9	1.4	8.0	3.2-17.4
	1996-2000	3.2	11.2	6.4-18.4	1.2	9.8	3.5-21.5	2.0	13.2	6.2-25.0	3.2	11.0	6.3-18.0	1.2	10.1	3.6-22.1	2.0	12.7	6.0-24.0
	2001-2005	3.4	11.0	6.4-17.9	1.2	9.7	3.5-21.6	2.2	12.6	6.3-23.4	3.6	11.8	7.0-18.9	1.4	10.9	4.3-23.1	2.2	12.7	6.3-23.7
	2006-2010	3.6	11.1	6.6-17.9	1.6	10.1	4.3-20.8	2.0	11.8	5.6-22.5	3.6	11.1	6.6-17.9	1.2	7.4	2.7-17.2	2.4	14.0	7.2-25.2
	2011-2015	3.0	7.9	4.4-13.6	1.8	9.8	4.4-19.9	1.2	5.8	2.1-14.1	3.2	8.9	5.1-15.0	1.6	9.8	4.1-20.3	1.6	8.3	3.6-17.6
	2016-2020	6.0	15.0	10.1-22.1	3.4	18.2	10.5-30.5	2.6	11.9	6.3-21.9	4.8	11.9	7.5-18.3	2.8	14.7	7.9-26.1	2.0	9.2	4.4-18.4
Larynx	1976-1980	0.4	1.6	0.2-6.3	0.4	4.4	0.5-16.3	0.0	0.0	0.0-7.1	0.4	1.9	0.2-7.0	0.2	1.5	0.0-10.5	0.2	2.2	0.1-11.5
	1981-1985	2.0	8.6	4.0-16.1	1.8	17.1	7.7-33.1	0.2	1.3	0.0-8.9	0.2	0.8	0.0-4.8	0.2	1.7	0.0-10.6	0.0	0.0	0.0-6.8
	1986-1990	2.2	8.5	4.2-15.6	1.6	14.1	6.0-28.3	0.6	4.0	0.8-12.7	0.2	0.7	0.0-4.4	0.2	1.4	0.0-9.5	0.0	0.0	0.0-6.4
	1991-1995	0.6	2.5	0.5-7.4	0.6	5.4	1.1-15.8	0.0	0.0	0.0-6.0	0.4	1.4	0.2-5.4	0.2	1.8	0.0-10.0	0.2	1.2	0.0-8.0
	1996-2000	1.0	3.7	1.2-8.7	1.0	7.6	2.4-18.2	0.0	0.0	0.0-5.9	0.6	2.3	0.5-6.8	0.4	3.4	0.4-12.3	0.2	1.3	0.0-8.0
	2001-2005	1.8	5.8	2.6-11.3	1.2	8.0	2.9-18.2	0.6	3.8	0.8-11.9	0.4	1.3	0.2-5.2	0.4	2.8	0.3-11.0	0.0	0.0	0.0-5.9
	2006-2010	1.4	4.3	1.7-9.3	1.2	8.0	2.9-18.2	0.2	1.1	0.0-7.6	0.0	0.0	0.0-3.0	0.0	0.0	0.0-6.3	0.0	0.0	0.0-5.9
	2011-2015	0.8	2.4	0.6-6.8	0.8	5.1	1.4-14.2	0.0	0.0	0.0-6.1	0.2	0.5	0.0-3.8	0.2	1.3	0.0-8.6	0.0	0.0	0.0-6.1
	2016-2020	1.4	3.6	1.4-8.2	1.2	7.0	2.5-16.6	0.2	1.1	0.0-7.9	0.8	2.2	0.6-6.3	0.2	1.4	0.0-8.8	0.6	3.2	0.7-10.9
Lung and bronchus	1976-1980	13.2	55.1	42.4-70.5	11.0	107.6	80.1-141.6	2.2	17.2	8.5-31.5	10.0	41.1	30.3-54.6	8.6	82.1	58.5-112.2	1.4	11.0	4.3-23.3
	1981-1985	13.8	57.2	44.2-72.9	8.6	86.8	62.0-118.1	5.2	37.0	24.0-55.0	13.0	52.4	40.2-67.2	8.4	81.4	58.0-111.2	4.6	30.7	19.4-46.8
	1986-1990	22.2	85.5	70.1-103.4	15.2	129.6	101.6-163.1	7.0	50.7	34.9-71.4	18.8	72.5	58.4-89.0	12.4	111.5	84.8-143.8	6.4	43.6	29.6-62.3
	1991-1995	23.4	85.2	70.3-102.4	14.2	119.0	92.4-150.8	9.2	60.3	43.9-81.1	18.2	67.4	54.1-83.0	11.2	97.9	73.4-127.9	7.0	47.7	32.9-67.0
	1996-2000	23.2	79.1	65.4-95.1	13.2	97.9	75.5-125.0	10.0	64.7	47.9-85.8	16.2	54.8	43.5-68.3	11.4	87.5	66.0-113.9	4.8	30.4	19.4-45.7
	2001-2005	23.0	76.1	62.8-91.6	11.6	83.6	63.3-108.7	11.4	70.7	53.5-92.2	17.4	57.3	45.8-70.8	8.8	63.7	46.1-86.2	8.6	52.7	38.0-71.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Yates**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Lung and bronchus	2006-2010	22.4	69.3	56.9-83.9	10.6	71.5	53.4-94.2	11.8	68.4	51.6-89.3	14.8	44.7	35.0-56.5	7.6	49.6	34.9-68.8	7.2	40.5	28.0-57.0
	2011-2015	24.8	69.5	57.6-83.4	13.4	81.5	62.9-104.5	11.4	61.9	46.7-81.1	17.8	50.6	40.5-62.9	8.6	53.7	38.6-73.4	9.2	49.3	35.7-66.9
	2016-2020	25.0	65.5	54.1-78.9	12.0	66.8	50.3-87.7	13.0	64.2	49.0-83.4	13.8	36.7	28.3-47.2	7.2	42.4	29.1-60.2	6.6	31.4	21.5-45.4
Melanoma of the skin	1976-1980	1.6	7.9	3.2-15.8	1.0	10.1	3.1-24.4	0.6	5.6	1.0-16.9	0.8	3.9	1.0-10.2	0.6	6.4	1.2-19.2	0.2	1.3	0.0-9.0
	1981-1985	3.2	14.7	8.3-24.1	1.6	15.9	6.7-31.6	1.6	12.8	5.3-25.9	1.0	4.8	1.5-11.5	0.6	5.7	1.1-17.2	0.4	4.2	0.4-14.9
	1986-1990	3.8	17.1	10.1-26.9	1.4	12.3	4.8-25.9	2.4	21.5	10.8-38.1	1.6	6.3	2.7-12.8	1.0	7.8	2.5-19.1	0.6	5.1	0.9-15.2
	1991-1995	1.8	7.3	3.3-14.0	1.2	10.9	4.0-23.7	0.6	3.2	0.6-10.6	0.4	1.6	0.2-6.0	0.4	3.3	0.4-12.4	0.0	0.0	0.0-6.0
	1996-2000	2.8	11.0	6.0-18.5	1.0	7.8	2.5-18.6	1.8	13.6	6.1-26.0	1.4	4.9	2.0-10.3	0.8	6.8	1.8-17.5	0.6	4.3	0.9-13.0
	2001-2005	5.6	19.6	12.9-28.6	3.4	23.0	13.3-37.4	2.2	16.4	8.1-29.7	0.6	1.9	0.4-6.1	0.4	2.7	0.3-10.7	0.2	1.1	0.0-7.6
	2006-2010	6.2	20.7	13.9-29.8	3.6	26.9	15.7-43.2	2.6	16.4	8.6-28.9	0.4	1.1	0.1-4.8	0.2	1.0	0.0-7.7	0.2	1.3	0.0-8.1
	2011-2015	10.2	36.6	26.5-49.2	6.0	41.7	27.5-60.9	4.2	34.3	20.3-53.8	1.6	4.5	1.9-9.5	1.2	7.8	2.8-17.9	0.4	2.1	0.2-9.3
	2016-2020	9.8	29.9	21.5-40.7	6.8	41.2	27.6-59.4	3.0	19.6	10.3-34.1	0.2	0.6	0.0-4.1	0.0	0.0	0.0-6.5	0.2	1.2	0.0-8.1
Female breast	1976-1980							9.6	81.1	58.6-109.2							2.6	21.0	10.8-37.0
	1981-1985							11.8	93.0	69.8-121.4							3.8	26.3	15.8-41.8
	1986-1990							17.2	129.6	102.7-161.5							3.6	27.0	15.7-43.5
	1991-1995							17.4	135.0	107.5-167.4							4.4	30.2	18.7-46.3
	1996-2000							19.6	136.5	110.4-167.1							4.6	28.5	17.9-43.5
	2001-2005							20.8	139.7	113.7-170.1							3.2	19.3	11.0-32.1
	2006-2010							20.6	123.9	100.6-151.2							4.0	24.4	14.5-38.8
	2011-2015							21.4	129.5	104.9-158.5							3.0	16.7	9.0-28.8
	2016-2020							21.8	128.3	103.4-157.7							2.8	17.0	8.9-29.7
Cervix uteri	1976-1980							1.0	8.0	2.6-19.3							1.2	12.1	4.1-26.7
	1981-1985							1.4	12.2	4.6-25.8							0.4	3.0	0.4-11.7
	1986-1990							1.8	14.4	6.4-27.8							1.2	7.4	2.7-17.2
	1991-1995							2.2	17.7	8.7-32.1							1.2	8.3	2.9-18.6
	1996-2000							0.8	6.6	1.8-16.8							0.4	2.4	0.3-9.8
	2001-2005							0.8	5.9	1.6-15.3							0.0	0.0	0.0-5.9
	2006-2010							0.6	3.7	0.7-11.6							0.2	1.3	0.0-8.1
	2011-2015							1.2	10.5	3.5-23.4							0.6	5.0	0.8-15.3
	2016-2020							1.0	8.3	2.4-20.1							0.2	2.6	0.1-12.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Yates**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Corpus uteri and uterus, not otherwise specified	1976-1980							1.8	12.3	5.6-24.4							0.2	1.4	0.0-9.2
	1981-1985							2.2	15.3	7.6-28.3							0.6	4.2	0.8-13.3
	1986-1990							3.4	26.4	15.1-43.0							0.6	3.9	0.8-12.3
	1991-1995							4.0	29.3	17.7-45.8							0.4	2.2	0.3-9.4
	1996-2000							4.0	27.5	16.7-43.0							0.6	3.1	0.6-10.3
	2001-2005							3.8	24.9	14.9-39.4							0.8	5.4	1.4-14.5
	2006-2010							6.0	34.6	23.1-50.3							1.4	7.0	2.8-15.7
	2011-2015							6.6	36.6	24.7-52.8							0.6	3.7	0.8-11.9
	2016-2020							6.2	29.3	19.4-43.3							2.0	9.1	4.2-18.5
Ovary	1976-1980							3.2	22.9	13.0-37.9							1.8	11.9	5.4-23.7
	1981-1985							3.0	24.9	13.3-42.3							2.0	16.4	7.5-31.1
	1986-1990							1.8	13.9	6.2-26.8							2.0	13.1	6.3-24.9
	1991-1995							1.8	11.9	5.4-23.3							0.8	5.1	1.4-13.7
	1996-2000							2.2	13.5	6.6-25.1							1.4	7.9	3.1-17.3
	2001-2005							2.8	19.1	10.3-32.8							1.4	8.5	3.4-18.2
	2006-2010							2.8	16.2	8.8-27.9							1.6	10.5	4.3-21.6
	2011-2015							2.6	13.5	7.1-24.2							2.4	13.1	6.7-24.0
	2016-2020							2.2	14.6	6.7-27.8							0.6	3.4	0.7-11.3
Prostate	1976-1980				6.8	73.6	50.0-104.3							2.4	28.6	14.1-50.9			
	1981-1985				11.0	117.0	87.1-153.7							3.2	35.4	19.6-58.4			
	1986-1990				14.8	139.4	108.5-176.2							2.8	33.4	18.0-55.8			
	1991-1995				23.4	194.9	160.4-234.6							3.8	37.9	22.5-59.2			
	1996-2000				23.4	177.1	146.1-212.9							4.4	36.8	22.8-56.0			
	2001-2005				25.0	178.9	148.6-213.9							3.2	25.7	14.6-42.0			
	2006-2010				22.2	138.8	113.8-168.1							2.2	16.4	8.1-29.8			
	2011-2015				17.6	98.1	78.4-122.0							3.6	25.1	14.9-40.2			
	2016-2020				28.8	152.2	127.7-180.7							3.4	20.8	12.0-34.3			
Testis	1976-1980				0.4	3.8	0.5-14.5							0.0	0.0	0.0-8.2			
	1986-1990				0.8	7.7	2.0-19.9							0.0	0.0	0.0-7.3			
	1991-1995				0.6	5.9	1.2-16.9							0.0	0.0	0.0-6.7			
	1996-2000				1.4	13.2	5.3-26.9							0.0	0.0	0.0-6.3			
	2001-2005				0.2	2.1	0.1-10.7							0.0	0.0	0.0-6.4			
	2006-2010				0.4	4.5	0.5-15.2							0.2	1.1	0.0-8.0			

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Yates**

Site of Cancer	Time Period	Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
		Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Testis	2011-2015				0.4	4.0	0.5-13.8							0.0	0.0	0.0-6.4			
	2016-2020				0.6	4.4	0.8-13.9							0.0	0.0	0.0-6.5			
Urinary bladder (incl. in situ)	1976-1980	4.8	19.9	12.6-30.0	4.4	43.3	26.6-66.7	0.4	2.9	0.4-11.7	0.4	1.9	0.2-7.1	0.0	0.0	0.0-8.2	0.4	2.9	0.4-11.7
	1981-1985	4.8	19.7	12.5-29.7	3.6	33.6	19.5-54.1	1.2	9.5	3.2-21.6	1.6	6.5	2.8-13.1	1.2	12.7	4.5-27.9	0.4	2.9	0.3-11.5
	1986-1990	7.6	29.7	20.9-41.1	6.6	62.7	42.6-88.9	1.0	7.1	2.2-17.2	2.0	7.6	3.7-14.3	1.2	11.8	4.2-26.0	0.8	4.7	1.2-13.3
	1991-1995	6.8	24.9	17.2-35.0	4.4	38.8	24.1-59.1	2.4	15.3	7.8-27.4	2.6	9.6	5.1-16.5	1.6	14.9	6.3-29.3	1.0	6.0	1.9-15.2
	1996-2000	6.0	20.6	13.8-29.6	5.0	40.5	26.0-60.3	1.0	6.1	2.0-15.1	1.8	6.0	2.7-11.6	1.4	12.4	4.9-25.7	0.4	2.4	0.3-9.7
	2001-2005	8.4	27.7	19.9-37.6	6.8	49.3	34.0-69.5	1.6	10.0	4.3-20.3	2.0	6.6	3.2-12.4	1.8	14.5	6.5-27.8	0.2	1.1	0.0-7.6
	2006-2010	8.0	24.7	17.6-34.1	6.4	43.0	29.2-61.5	1.6	7.9	3.4-16.8	1.8	5.7	2.6-11.1	1.2	8.8	3.2-19.6	0.6	3.1	0.6-10.4
	2011-2015	8.0	21.6	15.3-29.9	7.0	40.4	27.9-57.3	1.0	5.2	1.7-13.5	1.0	2.5	0.8-6.6	1.0	5.9	1.9-15.0	0.0	0.0	0.0-6.1
2016-2020	12.2	33.2	25.1-43.3	9.6	55.3	40.4-74.6	2.6	15.3	7.8-27.4	2.0	5.4	2.6-10.5	0.8	5.6	1.5-15.0	1.2	5.9	2.1-14.3	
Kidney and renal pelvis	1976-1980	1.0	4.9	1.6-11.7	0.4	4.8	0.5-17.1	0.6	4.4	0.9-13.8	0.8	3.4	0.9-9.0	0.4	3.4	0.4-13.6	0.4	2.7	0.3-11.1
	1981-1985	1.8	7.2	3.3-14.0	1.6	13.9	6.0-28.2	0.2	1.3	0.0-8.8	0.8	3.4	0.9-9.0	0.6	5.8	1.2-17.2	0.2	1.3	0.0-8.8
	1986-1990	1.4	5.7	2.2-12.0	0.8	6.5	1.7-17.6	0.6	5.0	1.0-14.9	0.2	0.8	0.0-4.8	0.0	0.0	0.0-7.3	0.2	1.6	0.0-9.2
	1991-1995	2.4	10.1	5.2-17.8	1.8	15.8	7.1-30.2	0.6	5.4	1.1-15.5	1.0	3.9	1.3-9.3	0.6	5.7	1.1-16.7	0.4	3.5	0.4-12.4
	1996-2000	1.8	6.1	2.8-11.8	1.4	9.8	3.9-20.7	0.4	2.4	0.3-9.7	0.2	0.8	0.0-4.4	0.0	0.0	0.0-6.3	0.2	1.5	0.0-8.6
	2001-2005	4.8	16.2	10.3-24.3	3.4	24.0	13.9-39.1	1.4	8.6	3.3-18.6	0.6	1.9	0.4-6.1	0.4	2.7	0.3-10.8	0.2	0.9	0.0-7.3
	2006-2010	6.4	20.7	14.0-29.6	4.8	31.1	19.6-47.3	1.6	10.4	4.3-21.3	1.2	3.4	1.2-7.9	1.0	6.6	2.1-16.3	0.2	0.9	0.0-7.2
	2011-2015	6.6	20.1	13.4-29.1	4.0	27.1	15.8-43.6	2.6	13.2	6.8-23.9	1.4	4.3	1.7-9.4	0.6	4.1	0.8-12.9	0.8	4.4	1.0-12.9
2016-2020	5.6	17.1	10.8-25.9	4.2	26.7	15.8-42.6	1.4	8.8	3.0-20.1	0.6	1.5	0.3-5.4	0.2	1.1	0.0-8.2	0.4	2.0	0.2-9.1	
Brain and other central nervous system	1976-1980	1.8	8.5	3.8-16.5	1.0	9.8	3.1-23.5	0.8	8.0	2.0-20.6	0.6	3.0	0.6-8.8	0.4	4.7	0.6-16.6	0.2	1.3	0.0-9.0
	1981-1985	1.6	6.8	2.9-13.7	0.6	5.8	1.1-17.5	1.0	7.7	2.4-18.6	0.4	1.9	0.2-7.0	0.4	4.0	0.4-14.8	0.0	0.0	0.0-6.8
	1986-1990	1.4	5.6	2.2-11.6	0.6	5.9	1.2-17.2	0.8	5.5	1.5-14.9	1.6	6.6	2.8-13.2	0.8	7.5	2.0-19.5	0.8	5.4	1.4-14.7
	1991-1995	2.0	7.6	3.6-14.2	0.8	6.3	1.7-16.7	1.2	8.9	3.2-19.7	1.4	5.0	2.0-10.6	0.8	6.6	1.8-17.3	0.6	3.9	0.8-12.1
	1996-2000	2.6	9.9	5.2-17.1	1.4	11.3	4.5-23.6	1.2	8.0	2.9-17.9	1.2	4.8	1.7-10.4	0.6	5.3	1.1-15.4	0.6	3.7	0.8-11.6
	2001-2005	2.8	9.4	5.1-16.1	1.2	8.3	3.0-18.8	1.6	10.1	4.3-20.5	2.6	8.6	4.6-15.0	1.2	8.3	3.0-18.7	1.4	8.7	3.5-18.7
	2006-2010	2.6	8.9	4.7-15.5	1.4	10.4	4.0-22.0	1.2	7.3	2.7-16.6	1.8	5.8	2.6-11.3	1.0	7.0	2.2-17.0	0.8	4.5	1.2-12.6
	2011-2015	2.0	6.4	3.0-12.3	0.8	4.9	1.3-13.8	1.2	8.4	2.9-19.0	1.8	6.1	2.7-12.1	1.2	9.1	3.1-20.6	0.6	3.7	0.8-11.9
2016-2020	2.2	5.7	2.8-10.8	1.2	6.2	2.2-15.2	1.0	5.3	1.7-13.7	1.2	3.1	1.1-7.5	0.4	2.1	0.2-9.5	0.8	4.1	1.1-12.0	

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Yates**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Thyroid	1976-1980	0.8	3.8	1.0-10.1	0.0	0.0	0.0-8.2	0.8	6.7	1.6-18.0	0.4	2.0	0.2-7.2	0.2	3.6	0.1-16.5	0.2	1.3	0.0-9.0
	1981-1985	0.6	2.6	0.5-7.7	0.2	3.2	0.1-14.9	0.4	3.1	0.4-11.9	0.0	0.0	0.0-3.5	0.0	0.0	0.0-7.8	0.0	0.0	0.0-6.8
	1986-1990	0.4	2.1	0.3-7.3	0.0	0.0	0.0-7.3	0.4	4.2	0.5-14.4	0.0	0.0	0.0-3.3	0.0	0.0	0.0-7.3	0.0	0.0	0.0-6.4
	1991-1995	2.0	8.6	4.1-15.8	0.6	4.7	0.9-14.3	1.4	12.1	4.8-24.7	0.4	1.7	0.2-6.0	0.2	1.9	0.0-10.3	0.2	1.3	0.0-8.1
	1996-2000	2.4	10.2	5.3-17.9	0.4	3.1	0.4-11.6	2.0	17.0	8.1-31.2	0.2	0.6	0.0-4.1	0.0	0.0	0.0-6.3	0.2	1.2	0.0-7.8
	2001-2005	1.8	6.6	3.0-12.8	0.8	5.9	1.6-15.7	1.0	7.3	2.3-17.4	0.0	0.0	0.0-3.0	0.0	0.0	0.0-6.4	0.0	0.0	0.0-5.9
	2006-2010	3.2	13.4	7.3-22.2	0.8	6.6	1.6-17.4	2.4	19.7	9.7-35.2	0.0	0.0	0.0-3.0	0.0	0.0	0.0-6.3	0.0	0.0	0.0-5.9
	2011-2015	4.6	17.3	10.6-26.6	1.2	7.9	2.9-18.1	3.4	26.5	14.8-43.5	0.6	1.8	0.4-5.8	0.2	1.2	0.0-8.3	0.4	2.1	0.2-9.2
	2016-2020	3.2	14.4	7.9-23.8	0.4	2.5	0.3-10.4	2.8	26.3	13.9-44.6	0.0	0.0	0.0-3.2	0.0	0.0	0.0-6.5	0.0	0.0	0.0-6.3
Hodgkin lymphoma	1976-1980	0.6	2.8	0.6-8.3	0.6	5.6	1.1-17.3	0.0	0.0	0.0-7.1	0.2	0.8	0.0-5.1	0.0	0.0	0.0-8.2	0.2	1.4	0.0-9.3
	1981-1985	0.6	2.7	0.5-8.1	0.4	3.8	0.5-14.2	0.2	2.2	0.1-11.4	0.0	0.0	0.0-3.5	0.0	0.0	0.0-7.8	0.0	0.0	0.0-6.8
	1986-1990	0.8	3.0	0.8-8.0	0.8	6.6	1.8-17.6	0.0	0.0	0.0-6.4	0.2	0.8	0.0-4.7	0.2	1.6	0.0-10.0	0.0	0.0	0.0-6.4
	1991-1995	1.0	4.3	1.4-10.1	0.2	1.4	0.0-9.1	0.8	7.0	1.9-17.7	0.6	2.4	0.5-7.2	0.4	3.1	0.4-12.0	0.2	1.7	0.0-9.3
	1996-2000	0.8	3.8	1.0-9.5	0.8	7.7	2.1-19.4	0.0	0.0	0.0-5.9	0.0	0.0	0.0-3.0	0.0	0.0	0.0-6.3	0.0	0.0	0.0-5.9
	2001-2005	0.2	1.3	0.0-6.0	0.0	0.0	0.0-6.4	0.2	2.5	0.1-11.5	0.0	0.0	0.0-3.0	0.0	0.0	0.0-6.4	0.0	0.0	0.0-5.9
	2006-2010	0.6	2.5	0.5-7.4	0.4	3.9	0.5-13.6	0.2	1.3	0.0-8.0	0.2	0.7	0.0-4.2	0.0	0.0	0.0-6.3	0.2	1.3	0.0-8.0
	2011-2015	0.8	2.7	0.7-7.5	0.6	4.1	0.7-13.1	0.2	1.4	0.0-8.5	0.4	1.8	0.2-6.5	0.4	3.9	0.4-13.9	0.0	0.0	0.0-6.1
	2016-2020	0.8	2.4	0.6-6.9	0.6	4.0	0.8-13.0	0.2	1.0	0.0-7.7	0.0	0.0	0.0-3.2	0.0	0.0	0.0-6.5	0.0	0.0	0.0-6.3
Non-Hodgkin lymphomas	1976-1980	1.6	7.8	3.3-15.5	1.4	17.1	6.4-35.6	0.2	1.4	0.0-9.3	1.8	8.8	3.9-16.9	1.6	21.3	8.7-42.2	0.2	1.6	0.0-9.7
	1981-1985	2.8	11.1	6.0-18.9	0.8	7.3	1.9-19.5	2.0	13.3	6.4-25.4	1.2	5.4	1.9-12.1	0.6	5.1	1.0-15.8	0.6	5.4	0.9-16.5
	1986-1990	4.2	17.0	10.5-26.2	1.8	17.3	7.7-33.2	2.4	16.9	8.6-30.2	3.4	14.1	8.2-22.8	1.6	17.0	7.1-33.6	1.8	12.1	5.4-23.8
	1991-1995	4.4	16.7	10.4-25.5	3.0	26.2	14.5-43.7	1.4	9.9	3.9-20.9	1.0	3.7	1.2-8.8	0.4	4.5	0.5-15.2	0.6	4.0	0.8-12.5
	1996-2000	6.0	21.4	14.4-30.8	3.0	25.5	14.2-42.3	3.0	18.2	10.1-30.9	2.6	9.0	4.8-15.5	1.2	9.2	3.4-20.4	1.4	7.5	3.0-16.6
	2001-2005	6.0	20.6	13.8-29.7	4.2	31.8	19.4-49.3	1.8	11.5	5.2-22.3	1.6	5.2	2.3-10.6	0.8	5.7	1.5-15.2	0.8	4.3	1.1-12.1
	2006-2010	9.6	31.3	22.8-42.0	5.2	37.2	23.8-55.7	4.4	25.6	15.9-39.4	2.8	8.5	4.6-14.6	2.0	13.0	6.1-24.8	0.8	4.3	1.2-12.2
	2011-2015	8.8	26.9	19.2-36.9	6.8	45.0	30.7-64.1	2.0	10.4	4.8-20.6	1.4	3.9	1.5-8.6	0.4	2.1	0.2-9.5	1.0	4.9	1.6-13.0
	2016-2020	6.6	20.7	13.8-30.1	3.6	24.8	14.1-40.7	3.0	17.2	9.0-30.2	1.4	3.4	1.4-7.8	0.8	4.6	1.2-13.3	0.6	2.5	0.5-9.6
Myeloma	1976-1980	2.0	8.0	3.8-15.1	1.2	11.3	4.0-25.5	0.8	5.8	1.6-15.9	1.2	4.7	1.7-10.7	0.8	7.0	1.9-19.1	0.4	3.1	0.4-12.1
	1981-1985	1.2	4.7	1.7-10.6	0.6	5.1	1.0-15.8	0.6	4.0	0.8-12.8	0.8	3.1	0.8-8.3	0.6	6.0	1.0-18.4	0.2	1.4	0.0-9.1
	1986-1990	1.6	6.6	2.8-13.3	1.2	11.7	4.2-25.7	0.4	2.3	0.3-10.0	1.6	6.1	2.6-12.3	1.2	11.1	4.1-24.5	0.4	2.3	0.3-9.9
	1991-1995	2.4	9.5	4.8-16.7	0.6	5.7	1.1-16.5	1.8	11.9	5.4-23.3	0.8	3.2	0.8-8.3	0.4	3.6	0.4-13.0	0.4	2.5	0.3-10.0
	1996-2000	2.0	7.0	3.3-13.0	1.2	9.6	3.5-21.2	0.8	5.6	1.5-14.8	0.8	2.9	0.8-7.7	0.4	3.4	0.4-12.3	0.4	2.9	0.3-11.0
	2001-2005	1.2	3.9	1.4-8.9	0.8	6.4	1.7-16.7	0.4	2.4	0.3-9.7	1.4	4.5	1.8-9.7	1.0	7.8	2.5-18.7	0.4	2.5	0.3-9.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.

**New York State Department of Health, New York State Cancer Registry  
Cancer Incidence and Mortality Trends, 1976-2020**

**Yates**

		Incidence									Mortality								
		Males and Females			Males			Females			Males and Females			Males			Females		
Site of Cancer	Time Period	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Cases	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI	Avg Ann Deaths	Rate per 100,000	95% CI
Myeloma	2006-2010	1.8	5.5	2.5-10.8	1.0	6.8	2.2-16.5	0.8	4.7	1.3-13.0	0.6	1.7	0.3-5.6	0.2	1.2	0.0-8.1	0.4	2.3	0.3-9.5
	2011-2015	1.4	4.4	1.7-9.5	0.4	2.5	0.3-10.3	1.0	6.0	1.8-15.2	0.8	2.5	0.7-6.8	0.4	2.9	0.4-11.2	0.4	2.1	0.2-9.2
	2016-2020	3.0	8.2	4.5-14.1	1.6	9.0	3.8-19.0	1.4	7.6	3.0-17.1	1.0	2.6	0.8-6.9	0.4	2.5	0.3-10.4	0.6	2.7	0.5-10.0
Leukemias	1976-1980	4.4	20.2	12.5-30.9	2.8	30.2	15.9-51.7	1.6	14.0	5.8-28.4	3.2	15.5	8.8-25.4	1.8	21.8	9.5-41.8	1.4	12.3	4.7-26.1
	1981-1985	2.4	11.3	5.7-20.0	1.4	16.0	6.2-33.1	1.0	8.6	2.5-20.8	2.0	8.3	4.0-15.6	1.0	9.9	3.0-23.9	1.0	7.6	2.4-18.6
	1986-1990	4.6	19.8	12.5-29.9	4.0	43.5	26.3-67.4	0.6	3.6	0.7-11.9	2.0	8.3	3.9-15.4	1.0	11.1	3.5-25.8	1.0	6.5	2.1-16.2
	1991-1995	3.4	13.2	7.6-21.2	1.6	13.0	5.5-26.1	1.8	12.7	5.7-24.7	1.6	6.1	2.6-12.2	0.6	5.1	1.0-15.1	1.0	5.9	1.9-14.6
	1996-2000	5.4	18.9	12.4-27.7	3.6	28.0	16.5-44.6	1.8	11.3	5.1-22.2	3.4	11.8	6.8-19.1	1.6	12.2	5.2-24.3	1.8	10.6	4.8-20.8
	2001-2005	3.0	10.6	5.9-17.7	0.8	5.9	1.5-15.8	2.2	15.1	7.4-27.7	1.0	3.3	1.1-8.0	0.6	5.6	1.1-16.2	0.4	2.4	0.3-9.6
	2006-2010	5.2	17.4	11.1-25.9	3.2	23.4	13.1-38.7	2.0	12.4	5.6-23.8	2.0	6.3	3.0-12.0	1.4	9.2	3.7-19.8	0.6	4.0	0.8-12.3
	2011-2015	4.6	13.8	8.6-21.3	2.0	13.6	6.3-26.0	2.6	14.4	7.5-25.9	2.0	6.2	2.9-11.9	1.2	9.1	3.3-20.3	0.8	4.1	1.1-11.9
	2016-2020	6.0	18.7	12.3-27.5	2.6	16.3	8.5-28.9	3.4	21.0	11.5-35.4	2.2	6.6	3.1-12.5	0.8	4.6	1.2-13.4	1.4	7.7	2.8-17.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2022.

Rates based on fewer than 4 cases or deaths per year are unstable and should be used with caution.

The data in this table are not identical to the data in the corresponding 2016-2020 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.