
New York State HIV/AIDS County Surveillance Report

(Excludes State Prison Inmates)



For Cases Diagnosed Through December 2011

Bureau of HIV/AIDS Epidemiology AIDS Institute New York State Department of Health August 2013

TABLE OF CONTENTS Excludes State Prison Inmates

T	Name Wards Charle HIV/AIDC Carries Carries III-man Damant	Page
	New York State HIV/AIDS County Surveillance Report	
Technical Notes on	the New York State HIV/AIDS County Surveillance Report	5-9
Definition of Term	s Used in this Report	.10-13
Guide to the Tables	S	.14-18
	NYS, EXCLUDING NYC, by County	10.01
Summary Table A	Average Annual Newly Diagnosed HIV Cases by County	19-21
Summary Table B	AIDS Cases by county of Residence at Diagnosis and Year of Diagnosis	22-23
Summary Table C	HIV Cases, Newly Diagnosed by County of Residence at Diagnosis and Year	of
-	Diagnosis	24-25
Summary Table D	Deaths Among AIDS Cases by County of Residence at Diagnosis and Year of	
-	Death	.26-27

Counties in New York State

The following tables have been generated for each of the counties in New York State:

- 1A Living HIV and AIDS Cases by Gender, Age, Race/Ethnicity and Risk
- 1B Living HIV and AIDS Cases by Gender and Risk
- **2A** Average Annual Newly Diagnosed HIV and AIDS Cases and Cumulative AIDS by Gender, Age, Race/Ethnicity and Risk
- 2B Newly Diagnosed HIV and AIDS Cases and Cumulative AIDS by Gender and Risk
- **3** Newly Diagnosed HIV and AIDS Cases by Year of Diagnosis and Deaths of AIDS Cases by Year

Introduction to the New York State HIV/AIDS County Surveillance Report

Overview

The County Surveillance Report presents recent data on newly diagnosed HIV cases, persons living with HIV/AIDS and deaths among AIDS cases in the counties of New York State. The County Report is a companion piece to the NYS HIV/AIDS Annual Surveillance Report (available at http://www.health.state.ny.us/diseases/aids/statistics/annual/index.htm). Like the Annual Surveillance Report, the County Report divides new HIV diagnoses into subcategories based on concurrent or subsequent AIDS diagnoses, and displays diagnosis and prevalence rates.

Format of the Report

The report covers the individual counties of New York State, except for the five boroughs of New York City, which are covered in the Annual Surveillance Report, and Hamilton County, which has too few HIV/AIDS cases to populate the tables. The report is issued in two parallel sections, one including and one excluding inmates in state correctional facilities. The sections contain four summary tables listing all counties, followed by sets of detailed tables for each individual county. The summary tables (A-D) are:

- (A) Average Annual Newly Diagnosed HIV Cases by Diagnosis Type,
- (B) AIDS Cases by Year of Diagnosis,
- (C) Newly Diagnosed HIV Cases by Year of Diagnosis, and
- (D) Deaths among AIDS Cases by Year of Death.

The county detail tables (1-3) are:

- (1) Living HIV and AIDS Cases
- (2) Average Annual Newly Diagnosed HIV and AIDS Cases, and
- (3) Newly Diagnosed HIV and AIDS Cases by Year of Diagnosis and Deaths of AIDS Cases by Year of Death

Living HIV (not AIDS), Living AIDS Cases and Living HIV/AIDS Cases

- The tables feature **prevalence rates** (living cases per 100,000 population) and show the population.
- Age is **current age** for the time point being reported, e.g. December 2012.

Newly Diagnosed HIV Cases

- All HIV cases newly diagnosed during the reporting period are counted, regardless of concurrent or subsequent AIDS diagnosis or vital status.
- Total new HIV diagnoses are subcategorized by AIDS diagnosis status. "Late" diagnoses are defined as HIV diagnoses followed by an AIDS diagnosis within 12 months. "Late" diagnoses are further subdivided into "concurrent" (AIDS diagnosis within one month of HIV diagnosis) or "other late" (AIDS diagnosis within 12 months). Remaining HIV diagnoses those who have not developed AIDS to date or developed AIDS more than 12 months after HIV diagnosis are reported as "other".
- Late and concurrent HIV diagnoses are also presented as a percent of total new HIV diagnoses.
- Case rates (cases per 100,000 population per year) are given for new HIV diagnoses.

Newly Diagnosed AIDS Cases

- Case rates are given.
- Cumulative AIDS cases are shown next to new AIDS diagnoses.

Special Note

Criteria for the inclusion of cases in this report have changed. In previous reports, persons living with diagnosed HIV infection (PLWDHI) whose initial HIV diagnosis occurred outside the United States were included in the tabulations only if they were subsequently diagnosed with AIDS in NYS. Foreign-diagnosed cases are now included regardless of AIDS status. In previous reports, persons diagnosed with HIV in other US states and diagnosed with AIDS in New York were included in the tabulations of PLWDHI regardless of the date of the AIDS diagnosis. Such cases are now included only if the AIDS diagnosis date was before the end of the calendar year of the report. The net result of these changes is an increase of 700 cases in the statewide tally of PLWDHI over what would have been shown.

Technical Notes on the New York State HIV/AIDS County Surveillance Report

Introduction

This HIV/AIDS County Surveillance Report is produced by the Bureau of HIV/AIDS Epidemiology (BHAE), AIDS Institute, New York State Department of Health (NYSDOH).

Printed copies can be obtained by contacting:

Division of Epidemiology, Evaluation and Research P.O. Box 2073, ESP Station
Albany NY 12220-0073
(518) 474-4284
bhae@health.state.ny.us

Information presented in this report is generated from data collected and maintained by BHAE and the HIV Epidemiology and Field Services (HEFS) Program of the NYC Department of Health and Mental Hygiene (NYCDOHMH).

Source Data

This report was produced using the March 2013 BHAE Statewide Analysis file. The Analysis file combines surveillance data from the New York State and New York City HIV/AIDS surveillance registries.

Reporting Procedures

On June 1, 2000, the NYSDOH implemented the HIV Reporting and Partner Notification Law (Chapter 163 of the Laws of 1998, PHL 21 Title III). This law enhanced the existing AIDS case reporting by requiring health care providers to report all initial determinations of HIV infection, HIV-related illness and AIDS to the NYS Commissioner of Health. Under the amended HIV

¹ HIV-infected patients diagnosed with tuberculosis (TB) are also reportable to the local TB control program. Questions regarding TB reporting should be directed to your local program or the NYS TB Control Program at (518) 474-4845.

testing law (Chapter 308 of the Laws 2010), providers are required to report such determinations within 14 days of receipt of positive laboratory result or after diagnosis, whichever is sooner. Until June 2005, laboratories were required to report all confirmatory HIV antibody tests, all detectable viral loads and all CD4<500 cells/mm³ or 29% of total lymphocytes. As of June 2005, all viral load and CD4 tests are reportable, as well as confirmatory HIV antibody tests and genotype results of antiretroviral drug resistance tests. In 2010, laboratory reporting requirements were broadened to include electronic reporting of any laboratory test, tests or series of tests approved for the diagnosis of HIV or for the periodic monitoring of HIV infection.

Laboratories conducting HIV-related testing must report electronically to the NYSDOH. For more information call 518-474-4284.

All NYS providers should report using the NYS Confidential Medical Provider HIV/AIDS and Partner/Contact Report Form (PRF) DOH-4189 Revised 3/09. Copies can be obtained by calling 518-474-4284. Additional information about provider reporting is available at http://www.health.state.ny.us/diseases/aids/.

NYS providers **outside** of NYC should mail the completed PRF to:

Division of Epidemiology, Evaluation and Research P.O. Box 2073, ESP Station Albany, New York 12220-0073

NYC providers should call (212) 442-3388 to arrange pickup of the completed PRF.

Record Updates:

HIV/AIDS surveillance files are never closed. When additional information is received on a confirmed case after the initial confirmation, data for that case may be updated, resulting in changes in the data reported to the public. For example, new information may indicate that two surveillance cases in fact refer to a single individual. Surveillance system records for the two cases will be combined, reducing the case count for the geographic area, demographic or risk group of the combined cases.

HIV/AIDS Surveillance Case Definition:

A uniform surveillance case definition for AIDS was established by the Centers for Disease Control and Prevention (CDC) in the early 1980s, and has since been modified three times to expand the range of AIDS-indicator diseases and conditions and to incorporate HIV diagnostic tests. Modifications in 1985 and in 1987 were followed by the 1993 expansion of the AIDS case definition for adults and adolescents, which includes HIV-infected individuals with CD4+ T-lymphocyte counts below 200 cells/mm³ or below 14 percent of total lymphocytes, and individuals diagnosed with pulmonary tuberculosis, recurrent pneumonia or invasive cervical cancer.

Particularly due to the new immunologic criterion of the expanded case definition, the number of cases reported to the NYSDOH and NYCDOHMH grew immensely following the 1993 modification. The immunologic criterion permitted a substantial number of AIDS diagnoses to be made before the occurrence of the first opportunistic condition. These cases were effectively diagnosed earlier in the spectrum of HIV disease than other AIDS cases diagnosed under the old case definition. The effect of this new criterion was to create a large and sudden increase in the number of AIDS cases confirmed around 1993, producing the false impression of a dramatic increase in AIDS cases during that period.

In 2008 CDC revised the HIV case definition², designating AIDS as "stage 3 HIV infection". This County Report continues to use the term AIDS to refer to this stage of disease. The 2008 case definition also requires laboratory-confirmed evidence of HIV infection. Since virtually all HIV cases entered into the NYS surveillance system meet this requirement it is not expected to affect case counts.

Deduplication:

The HIV/AIDS case data used in this County Report have undergone both inter- and intra-state deduplication.

² Revised Surveillance Case Definitions for HIV Infection Among Adults, Adolescents, and Children Aged <18 Months and for HIV Infection and AIDS Among Children Aged 18 Months to <13 Years — United States, 2008;MMWR, Dec. 5 2008, vol. 57, No. RR-10

CDC's process for interstate deduplication of HIV/AIDS cases, known as Routine Interstate Duplicate Review (RIDR), is conducted twice annually. If a case is reported by two or more states, the earliest date of diagnosis is used to determine which state the case is assigned to for national tabulations. Cases in the New York registry that are known through RIDR to have been initially diagnosed in another state are excluded from the County Report.

The County Report case data have also been deduplicated between NYC and the rest of NYS. In all of NYS excluding NYC, HIV/AIDS surveillance is conducted by the NYSDOH's BHAE. In NYC, HIV/AIDS surveillance is conducted by the NYC DOHMH HEFS Program. Often, the same case will be reported to both jurisdictions. The NYC and NYS departments of health report HIV/AIDS data separately to CDC, and the combined file can contain certain cases twice. At the national level, CDC deduplicates the data between the two jurisdictions for its own use. The goal of CDC's process is to ensure that each case is counted only once and is credited to the case's state of residence at the time of the earliest report. This deduplicated count provides the basis for Ryan White and other federal funding to NYS. The data used in this report have been deduplicated by BHAE following an algorithm similar to that used by CDC. Since May 2010, BHAE has performed further adjustments to the data using information from an intrastate extension of the RIDR process. This removes several hundred additional duplicate records, but importantly, returns to the analysis files some 3,000 unique cases that are excluded by the base procedure.

Ongoing refinement of the data may also contribute to geographical reassignment of cases at the state, regional, and/or county and local levels.

Data Completeness:

Newly diagnosed HIV and AIDS cases: Historical data indicate that counts of new HIV and AIDS diagnoses are approximately 95% complete by one calendar year after the close of the reporting period.

Living HIV and AIDS Cases: Because living cases include cases diagnosed in previous years as well as newly-diagnosed cases, delays in case reporting and ascertainment have little practical effect on the count. However, counts of living cases

may be affected by delays in death reporting and incompleteness of matching death records to surveillance data.

Definition of Terms Used in this Report

<u>Case</u>: "Case" refers to a report for which confirmatory clinical and demographic data required by the CDC have been obtained. Reports received without sufficient information to confirm the suspected case or a confirmed case reported without required demographic variables or risk are investigated by NYS or NYC public health representatives who examine medical records to confirm case status, obtain demographic information, clinical status and reported HIV exposure (risk).

HIV and AIDS Case Categories:

- *Living HIV Cases* All reported individuals living with HIV infection who have not developed AIDS. This category includes HIV cases reported by a provider indicating an HIV diagnosis or an HIV illness and/or by a laboratory reporting a confirmatory test result.
- Newly Diagnosed HIV Cases HIV cases that have been reported for the first time by a laboratory reporting a confirmatory test result or by a physician reporting an initial HIV diagnosis, regardless of concurrent or subsequent AIDS diagnosis. Although DOH staff seek to obtain an accurate diagnosis date, please note that HIV reporting began in June 2000, so cases diagnosed after then by a provider other than the provider reporting the case may be misclassified as newly diagnosed. (See **Diagnosis date**, below.)
- Living AIDS Cases All reported individuals living with AIDS.
- Newly Diagnosed AIDS Cases Cases reported with AIDS for the first time. This
 includes HIV cases that have never been reported before and cases initially reported with
 HIV infection that have subsequently developed AIDS.
- *Cumulative AIDS Cases* All AIDS cases reported since 1983, both living and deceased. There is no corresponding category for HIV cases.

Diagnosis date: The diagnosis date refers to the earliest date on which a clinical or laboratory diagnosis of HIV and AIDS is documented on a provider report form, laboratory report or in the patient's medical chart. This date may be months, or even years, earlier than the date the case was reported to the health department.

HIV Exposure Category: Information on possible mode(s) of HIV exposure is typically gathered by health care providers during routine medical history taking and is obtained from review of the patient's medical records, or may be obtained during routine follow-up for Partner Services. An individual could have been exposed to HIV through more than one route. However, for the purpose of surveillance, only one exposure mechanism is considered. (The category of "men who have sex with men and inject drugs" is the only exception to this.)

Persons identified with more than one potential exposure route are classified in accordance with the following CDC-defined hierarchy of exposure category:

Adults:

- Men who have sex with men and inject drugs
- Men who have sex with men (includes bisexual men)
- Injection drug users
- Hemophiliacs or persons with other clotting disorders
- Heterosexual contact with an injection drug user, a bisexual man, a hemophiliac, an HIV-positive blood product recipient, or a person with documented HIV or AIDS
- Transfusion or transplant recipient
- Confirmed other risk
- Female presumed heterosexual contact (FPHC)
- Adult with undetermined mode of exposure

Children:

CDC categorizes a case as "pediatric" if the diagnosis was made before age 13.

- Hemophiliacs or children with other clotting disorder
- Mother is an injection drug user
- Mother had sex with an injection drug user, a bisexual man, a hemophiliac, an HIV-positive blood transfusion recipient, an individual with documented HIV or AIDS
- Mother received a transfusion or blood products
- Mother had AIDS or is HIV-positive
- Transfusion or transplant recipient

- Pediatric other risk
- Child with undetermined mode of exposure

Based on this hierarchy, for example, a woman who may have been exposed to HIV through both injection drug use and through heterosexual contact with an HIV-infected partner would be classified only as an injection drug user, since that mode of exposure is ranked higher than heterosexual contact on the hierarchy.

Under the standard CDC risk hierarchy, an individual for whom all but heterosexual HIV risk has been ruled out can be classified as heterosexually exposed only if the partner's risk and/or HIV status is documented. In the absence of such information, the case is classified as having an undetermined risk category. In contrast, men who have sex with men and injection drug users need only be behaviorally identified in one of these groups, with no additional documentation required. As a result of the more rigorous standards for identifying heterosexually exposed individuals, it is likely that many heterosexually exposed cases fail to meet these standards. The Female Presumed Heterosexual Contact categorization addresses this issue in women whose surveillance record indicates heterosexual activity. There is no corresponding classification for males; thus a portion of male cases with an undetermined exposure category may be heterosexually exposed.

Individuals with an undetermined mode of exposure may fall into one of several categories: persons currently under investigation by public health representatives, persons for whom no mode of exposure was identified because the individual died or was lost to follow-up, persons with lack of exposure information in his/her medical chart, and persons for whom adequate follow-up information revealed no identifiable exposure mechanism.

Age:

Age at Diagnosis: The individual's age at the initial diagnosis of the condition in question (i.e., HIV or AIDS). As noted above, CDC categorizes a case as "pediatric" if the diagnosis was made before age 13.

Current Age: The individual's age at the end of the reporting period.

<u>Living Cases</u>: Vital status of cases is determined from updated case reports, medical record reviews, death certificate files maintained by the NYSDOH and NYCDOH Vital Records, and from Social Security death data and the National Death Index. Unless known to be deceased from these sources, cases are presumed to be alive.

<u>Prisoners</u>: Cases categorized as "Prisoners" have been diagnosed and/or reported while incarcerated in city or county jails or one of the NYS Department of Corrections and Community Supervision (DOCCS) facilities throughout NYS. Most prisoner reports come from a NYSDOCCS facility. For cases diagnosed while incarcerated, residence reflects the address of the correctional facility, rather than pre-incarceration home address. In some counties prisoners make up a significant but shifting proportion of new diagnoses and living cases, and the local HIV burden is better represented by removing prisoner cases from the tabulations. The County Reports present HIV surveillance data both including and excluding prisoners. Refer to table titles and footnotes to determine inclusion or exclusion of prisoner cases.

Residence: In this report, new HIV and AIDS diagnoses are presented by county of residence at the time of diagnosis, as reported in the surveillance record. Similarly, living AIDS cases are assigned to the county of AIDS diagnosis. Living HIV cases diagnosed with AIDS after the close of the report period are presented by county of HIV diagnosis. For example, a case diagnosed with HIV in Albany County in 2010 and diagnosed with AIDS in Monroe County in 2012, would be included among living HIV cases in Albany County in the County Report for December 2011.

Guide to the Tables

There are two types of tables in the County Reports. Summary tables contain one row for each of the 56 counties (all NYS counties, excluding the five boroughs of New York City and Hamilton County.) The summary tables present newly diagnosed HIV cases by type and year of diagnosis, and AIDS cases by year of diagnosis and year of death. The county detail tables present demographic and risk breakouts for new and living HIV and AIDS cases.

The tables of newly diagnosed cases present 3-year averages. Case counts in smaller counties may show great variation across years; the 3-year average is more stable and representative than any one year.

Guide to the Tables Summary Table A

New York State Excluding New York City

Summary Table A Average Annual Newly Diagnosed HIV Cases¹ (January 2009 - December 2011), by County Excludes State Prison Inmates

	Late Diagnoses (AII	DS within 1 yr of H	IV diagnosis)	Other HIV Diagnoses	All HIV Dia	gnoses ²	Percent of All	HIV Diagnoses
	Concurrent HIV&AIDS (A)	Other Late (B)	All Late (C=A+B)	(D)	(E=C+	D)	Concurrent (A÷E x 100)	All Late (C÷E x 100)
County	Annual Average	AnnualAverage	AnnualAverage	AnnualAverage	AnnualAverage	Case Rate ³	%	%
Albany	8.7	1.3	10.0	21.0	31.0	10.2	28.0	32.3
Allegany					1.3	2.7		
Broome					11.0	5.5		
Cattaraugus					2.0	2.5		

HIV Diagnoses: All newly diagnosed HIV cases, regardless of concurrent or subsequent AIDS diagnosis. Some of these cases may *also* be diagnosed with AIDS during the reporting period (in this case, January 2009-December 2011), and those AIDS diagnoses will be reported in the AIDS diagnosis tables (Summary Table B, County Tables 2A, 2B, 3).

Late Diagnoses: Late diagnoses are defined as those with an AIDS diagnosis within 12 months of the HIV diagnosis. For counties where the 3-year total of each of the subcategories (Concurrent HIV/AIDS, Other Late and Other) is at least 3 cases, the sub-category averages are displayed.

Concurrent HIV & AIDS: Newly diagnosed HIV cases with an AIDS diagnosis within a month of the HIV diagnosis date.

Other Late: Newly diagnosed HIV cases with an AIDS diagnosis 2 to 12 months after the HIV diagnosis date.

Other HIV Diagnoses: Newly diagnosed HIV cases without a subsequent AIDS diagnosis or with an AIDS diagnosis *more than 12 months after* the HIV diagnosis date.

Case (diagnosis) Rate: The number of newly diagnosed cases per 100,000 population (i.e. Case rate=new diagnoses divided by the county population x 100,000. County populations are shown on Table 1A of each of the individual counties.)

Percent Concurrent: The percent of newly diagnosed cases that are also diagnosed with AIDS *within one month* of the HIV diagnosis date.

Percent Late: The percent of newly diagnosed cases that are also diagnosed with AIDS *within twelve months* of the HIV diagnosis date. This includes concurrent diagnoses.

Guide to the Tables
Detail Table 1A

Albany County Table 1A

Living HIV and AIDS Cases as of December 2011

by Gender, Age, Race/Ethnicity and Risk Excludes State Prison Inmates

	Livin	g HIV (not	AIDS) Cases]	Living A	AIDS Cases	Livi	ng HIV and	1 AIDS Cases	
	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
	440	100.0	144.9	547	100.0	180.2	987	100.0	325.1	303,565
Male	288	65.5		367	67.1		655	66.4		
Female	152	34.5		180	32.9		332	33.6		
12 & under	6	1.4		0	0.0		6	0.6		
13-19	8	1.8		4	0.7		12	1.2		
20-24	19	4.3		6	1.1		25	2.5		
	Female 12 & under 13-19	Number 440 Male 288 Female 152 12 & under 6 13-19 8	Number Col % 440 100.0 Male 288 65.5 Female 152 34.5 12 & under 6 1.4 13-19 8 1.8	440 100.0 144.9 Male 288 65.5 . Female 152 34.5 . 12 & under 6 1.4 . 13-19 8 1.8 .	Number Col % Prevalence Rate 2 Number 440 100.0 144.9 547 Male 288 65.5 . 367 Female 152 34.5 . 180 12 & under 6 1.4 . 0 13-19 8 1.8 . 4	Number Col % Prevalence Rate ² Number Col % 440 100.0 144.9 547 100.0 Male 288 65.5 . 367 67.1 Female 152 34.5 . 180 32.9 12 & under 6 1.4 . 0 0.0 13-19 8 1.8 . 4 0.7	Number Col % Prevalence Rate ² Number Col % Prevalence Rate ² Male 288 65.5 . 367 67.1 . Female 152 34.5 . 180 32.9 . 12 & under 6 1.4 . 0 0.0 . 13-19 8 1.8 . 4 0.7 .	Number Col % Prevalence Rate ² Number Col % Prevalence Rate ² Number Male 288 65.5 . 367 67.1 . 655 Female 152 34.5 . 180 32.9 . 332 12 & under 6 1.4 . 0 0.0 . 6 13-19 8 1.8 . 4 0.7 . 12	Number Col % Prevalence Rate ² Number Col % Prevalence Rate ² Number Col % Male 288 65.5 . 367 67.1 . 655 66.4 Female 152 34.5 . 180 32.9 . 332 33.6 12 & under 6 1.4 . 0 0.0 . 6 0.6 13-19 8 1.8 . 4 0.7 . 12 1.2	Number Col % Prevalence Rate ² Number Col % Prevalence Rate ² Number Col % Prevalence Rate ² Male 288 65.5 . 367 67.1 . 655 66.4 . Female 152 34.5 . 180 32.9 . 332 33.6 . 12 & under 6 1.4 . 0 0.0 . 6 0.6 . 13-19 8 1.8 . 4 0.7 . 12 1.2 .

Living HIV/AIDS Cases: Persons living with HIV (not AIDS) and AIDS at the end of the report period (in this case December 2011). These are summed to produce a total count of people living with HIV and AIDS.

Prevalence Rate: The number of living cases per 100,000 population. Rates are not presented for population sub-groups because in most cases the numbers are too small to be statistically meaningful.

Area Population: Estimated population in 2010. Source: U.S. Bureau of the Census, Annual County Resident Population Estimates by Age, Sex, Race and Hispanic Origin, 2011.

Guide to the Tables
Detail Table 2A

Albany County Table 2A

Average Annual Newly Diagnosed HIV and AIDS Cases (2009-2011) and Cumulative AIDS (through 2011) by Gender, Age, Race/Ethnicity and Risk Excludes State Prison Inmates

		HIV Diagn	oses (Annua	l Average)	AIDS Diagr	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		31.0	100.0	10.2	21.3	100.0	7.0	1,196	100.0	303,565
Gender	Male	20.7	66.7		14.7	68.8		893	74.7	
	Female	10.3	33.3		6.7	31.3		303	25.3	
Age at Diagnosis	12 & under	0.3	1.1		0.0	0.0		14	1.2	
	13-19	1.0	3.2		0.3	1.6		11	0.9	
	20-24	3.0	9.7		0.7	3.1		32	2.7	

HIV Diagnoses: All newly diagnosed HIV cases, regardless of concurrent or subsequent AIDS diagnosis. Some of these cases may *also* be diagnosed with AIDS during the reporting period, and those AIDS diagnoses will be reported in the AIDS diagnosis column.

AIDS Diagnoses: AIDS diagnoses may occur among persons newly diagnosed with HIV (i.e., within the report period) as well as among those diagnosed earlier. HIV diagnoses and AIDS diagnoses **cannot** be added in a meaningful way.

Case Rate: The AIDS case (diagnosis) rate is calculated the same way as the HIV case (diagnosis) rate (see Guide to Summary Table A). Note: For population subgroups that did not have any HIV or AIDS diagnoses in the reporting period, no line appears in the table.

$Summary\ Table\ A$ Average Annual Newly Diagnosed HIV Cases 1 (January 2009 - December 2011), by County

	Late Diagnoses (AII	DS within 1 yr of H	IV diagnosis)	Other HIV Diagnoses	All HIV Dia	gnoses ²	Percent of All	HIV Diagnoses
	Concurrent HIV&AIDS (A)	Other Late (B)	All Late (C=A+B)	(D)	(E=C+	D)	Concurrent (A÷E x 100)	All Late (C÷E x 100)
County	Annual Average	AnnualAverage	AnnualAverage	AnnualAverage	AnnualAverage	Case Rate ³	%	%
Albany	8.7	1.3	10.0	21.0	31.0	10.2	28.0	32.3
Allegany					1.3	2.7		
Broome					11.0	5.5		
Cattaraugus					2.0	2.5		
Cayuga	1.0	1.0	2.0	3.7	5.7	7.1	17.6	35.3
Chautauqua					7.0	5.2		
Chemung					3.3	3.8		
Chenango					1.0	2.0		
Clinton					0.7	0.8		
Columbia					3.7	5.9		
Cortland					1.3	2.7		
Delaware					1.7	3.5		
Dutchess	4.7	2.3	7.0	10.3	17.3	5.8	26.9	40.4
Erie	26.3	6.0	32.3	66.0	98.3	10.7	26.8	32.9
Essex					1.0	2.6		
Franklin					2.0	3.9		
Fulton					2.3	4.2		
Genessee					2.3	3.9		
Greene					1.3	2.7		
Herkimer					3.3	5.2		
Jefferson					3.7	3.1		
Lewis					0.0	0.0		

¹ Data as of March 2013

²Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

³ Per 100,000 population

${\it Summary Table A} \\ {\it Average Annual Newly Diagnosed HIV Cases}^1 \ ({\it January 2009 - December 2011}), by County \\ {\it Summary Table A} \\ {\it Average Annual Newly Diagnosed HIV Cases}^1 \ ({\it January 2009 - December 2011}), by County \\ {\it Summary Table A} \\ {\it Average Annual Newly Diagnosed HIV Cases}^1 \ ({\it January 2009 - December 2011}), by County \\ {\it Summary Table A} \\ {\it Average Annual Newly Diagnosed HIV Cases}^1 \ ({\it January 2009 - December 2011}), by County \\ {\it Summary 2009 - December 2011}), by County \\ {\it Summary 2009 - December 2011}), by County \\ {\it Summary 2009 - December 2011}), by County \\ {\it Summary 2009 - December 2011}), by County \\ {\it Summary 2009 - December 2011}), by County \\ {\it Summary 2009 - December 2011}), by County \\ {\it Summary 2009 - December 2011}), by County \\ {\it Summary 2009 - December 2011}), by County \\ {\it Summary 2009 - December 2011}), by County \\ {\it Summary 2009 - December 2011}), by County \\ {\it Summary 2009 - December 2011}), by County \\ {\it Summary 2009 - December 2011}), by County \\ {\it Summary 2009 - December 2011}), by County \\ {\it Summary 2009 - December 2011}), by County \\ {\it Summary 2009 - December 2011}), by County \\ {\it Summary 2011}), by Coun$

	Late Diagnoses (AII	DS within 1 yr of H	IV diagnosis)	Other HIV Diagnoses	All HIV Dia	gnoses ²	Percent of All	HIV Diagnoses
	Concurrent HIV&AIDS (A)	Other Late (B)	All Late (C=A+B)	(D)	(E=C+	D)	Concurrent (A÷E x 100)	All Late (C÷E x 100)
County	AnnualAverage	AnnualAverage	AnnualAverage	AnnualAverage	AnnualAverage	Case Rate ³	%	%
Livingston					2.7	4.1		
Madison					1.3	1.8		
Monroe	18.7	6.0	24.7	62.0	86.7	11.6	21.5	28.5
Montgomery					1.3	2.7		
Nassau	39.0	7.0	46.0	57.0	103.0	7.7	37.9	44.7
Niagara	1.3	2.0	3.3	5.7	9.0	4.2	14.8	37.0
Oneida	2.7	1.3	4.0	7.7	11.7	5.0	22.9	34.3
Onondaga	9.7	3.0	12.7	26.0	38.7	8.3	25.0	32.8
Ontario					1.7	1.5		
Orange	8.0	3.3	11.3	14.3	25.7	6.8	31.2	44.2
Orleans					2.3	5.5		
Oswego					5.0	4.1		
Otsego					1.7	2.7		•
Putnam					6.3	6.3		
Rensselaer					9.0	5.6		
Rockland	6.0	1.0	7.0	15.0	22.0	7.0	27.3	31.8
St Lawrence					2.0	1.8		
Saratoga					5.3	2.4		
Schenectady	3.0	1.7	4.7	6.7	11.3	7.3	26.5	41.2
Schoharie					0.0	0.0		
Schuyler					0.3	1.8		
Seneca					1.0	2.8		

¹ Data as of March 2013

²Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

³ Per 100,000 population

$Summary\ Table\ A$ Average Annual Newly Diagnosed HIV Cases 1 (January 2009 - December 2011), by County

	Late Diagnoses (AII	DS within 1 yr of H	IV diagnosis)	Other HIV Diagnoses	All HIV Dia	gnoses ²	Percent of All	HIV Diagnoses
	Concurrent HIV&AIDS (A)	Other Late (B)	All Late (C=A+B)	(D)	(E=C+	D)	Concurrent (A÷E x 100)	All Late (C÷E x 100)
County	Annual Average	Annual Average	AnnualAverage	AnnualAverage	AnnualAverage	Case Rate ³	%	%
Steuben					2.7	2.7	-	
Suffolk	31.7	6.7	38.3	70.0	108.3	7.2	29.2	35.4
Sullivan					2.7	3.5		
Tioga					1.0	2.0	•	
Tompkins					4.0	3.9	•	
Ulster					10.7	5.8		
Warren					1.3	2.0		
Washington					2.7	4.2		
Wayne					4.7	5.0		
Westchester	28.0	7.0	35.0	72.7	107.7	11.3	26.0	32.5
Wyoming					0.3	0.8		
Yates					0.0	0.0		

¹ Data as of March 2013

²Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

³ Per 100,000 population

Summary Table B

AIDS Cases by County of Residence at Diagnosis and Year of Diagnosis 1

County	< 1986	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	Cumulative
Albany	17	16	26	32	48	53	77	88	94	69	73	63	54	43	44	40	34	48	42	55	29	40	21	26	20	28	16	1,196
Allegany	2	0	0	0	1	1	1	1	0	1	4	2	0	1	1	0	0	4	1	1	0	2	0	1	3	0	1	28
Broome	4	2	6	10	11	18	24	20	23	19	17	23	22	13	10	9	15	15	14	17	14	25	24	9	16	9	7	396
Cattaraugus	1	2	1	1	2	2	5	4	4	3	2	2	2	2	0	2	1	1	1	3	3	3	5	2	2	0	2	58
Cayuga	3	2	2	2	3	3	3	7	4	4	7	5	3	5	2	4	4	2	5	3	3	3	5	5	2	2	2	95
Chautauqua	2	0	1	2	4	2	4	9	9	8	12	6	3	9	5	6	8	12	4	7	9	5	8	7	7	4	3	156
Chemung	2	4	3	4	9	9	14	12	11	12	13	4	6	3	0	3	3	0	4	3	10	1	3	3	1	4	3	144
Chenango	1	0	0	2	0	1	1	4	2	0	6	3	2	0	1	3	5	5	2	1	3	0	1	2	2	1	2	50
Clinton	3	0	2	1	2	5	5	2	8	2	6	3	1	3	3	1	1	2	1	4	2	2	5	4	1	0	1	70
Columbia	2	3	1	2	4	5	9	14	5	4	10	2	4	2	3	4	2	1	3	5	5	2	4	1	4	2	3	106
Cortland	0	0	2	0	1	0	1	0	2	1	1	1	2	3	0	0	1	5	0	5	0	1	1	1	1	1	0	30
Delaware	0	0	0	1	2	4	1	3	2	3	6	4	2	0	1	0	4	4	4	2	3	1	2	3	0	3	4	59
Dutchess	13	15	29	32	41	47	63	67	105	95	63	67	31	26	36	41	29	26	22	36	23	21	14	16	17	11	13	999
Erie	20	23	36	55	82	78	103	122	174	161	161	144	78	86	102	127	93	66	88	69	68	59	84	55	61	54	72	2,321
Essex	3	0	1	4	3	2	1	2	3	2	7	1	1	0	0	1	0	1	1	0	1	1	0	1	0	1	0	37
Franklin	0	1	0	1	1	0	4	1	1	4	4	3	3	0	0	2	4	1	0	2	0	2	1	1	2	1	3	42
Fulton	3	1	0	3	0	2	4	4	1	2	3	1	1	4	3	1	1	1	2	4	2	3	2	4	1	1	3	57
Genessee	0	1	1	1	1	4	3	4	2	4	5	1	1	3	5	2	2	3	1	0	1	0	3	3	2	1	3	57
Greene	1	0	2	3	3	2	5	5	4	4	7	3	3	2	5	2	3	3	7	3	1	2	2	4	1	3	1	81
Herkimer	0	2	0	3	3	1	5	1	2	2	5	3	0	2	1	0	1	0	2	1	4	2	2	0	0	5	3	50
Jefferson	1	0	3	1	3	2	1	7	10	6	4	4	1	3	3	4	2	2	3	0	3	6	4	2	5	8	2	90
Lewis	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1
Livingston	1	1	2	6	3	2	5	2	6	7	3	0	2	3	2	1	4	2	2	0	1	2	4	2	1	1	2	67
Madison	0	2	0	1	1	5	4	0	6	0	2	2	3	3	0	4	1	0	0	1	2	1	0	1	1	2	1	43
Monroe	23	37	50	72	79	89	120	163	196	204	201	138	113	80	92	79	107	95	71	81	96	77	88	74	62	44	69	2,600
Montgomery	1	0	2	4	2	1	7	6	4	8	3	5	5	5	3	2	2	3	2	2	4	2	2	3	0	2	2	82
Nassau	134	119	143	190	226	224	246	293	304	251	294	235	164	150	151	166	154	149	123	118	94	93	94	93	87	80	72	4,447
Niagara	2	4	5	6	14	21	18	17	22	8	15	11	14	10	6	13	11	5	11	8	11	7	7	8	2	4	8	268
Oneida	5	6	6	10	12	24	21	34	39	20	23	21	17	14	19	15	5	10	15	14	13	7	11	13	12	4	5	395
Onondaga	34	12	24	33	42	49	58	119	110	77	80	84	54	32	44	75	38	39	41	31	38	27	36	35	26	20	31	1,289
Ontario	0	1	1	2	2	2	1	4	6	6	6	2	1	7	2	0	4	4	3	2	4	4	1	1	2	1	1	70

Summary Table B

AIDS Cases by County of Residence at Diagnosis and Year of Diagnosis¹

County	< 1986	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	Cumulative
Orange	15	16	29	35	52	71	87	89	93	110	77	53	59	36	47	47	41	43	57	43	43	27	28	30	28	18	20	1,294
Orleans	0	0	0	0	0	4	3	2	0	5	4	3	4	2	0	2	0	3	3	2	2	1	3	2	1	2	3	51
Oswego	2	2	0	4	8	2	4	9	5	3	7	4	6	2	6	2	1	8	1	4	7	4	1	0	3	7	3	105
Otsego	1	0	1	3	10	2	4	5	5	2	8	6	2	1	4	5	0	2	5	2	5	3	1	1	2	1	1	82
Putnam	7	2	8	10	7	9	12	16	11	12	17	11	6	10	10	5	5	2	5	2	1	5	8	3	3	6	2	195
Rensselaer	2	3	5	6	13	13	20	28	19	20	21	20	4	8	11	12	5	15	14	15	8	10	11	7	3	9	4	306
Rockland	31	20	28	29	44	30	51	90	79	64	56	55	35	41	30	45	38	27	25	30	43	25	18	18	18	14	14	998
St Lawrence	2	1	2	0	4	3	5	5	2	3	1	3	1	8	5	6	3	3	5	4	3	2	4	1	4	0	1	81
Saratoga	3	3	5	4	8	12	15	14	14	15	12	8	10	7	7	5	7	6	13	10	10	4	5	1	3	4	8	213
Schenectady	14	10	5	8	16	26	27	19	26	35	35	22	25	23	12	21	12	17	16	20	17	21	11	10	4	14	10	476
Schoharie	1	1	0	1	1	3	4	3	6	4	2	1	0	0	3	0	0	0	1	1	0	2	1	2	2	0	0	39
Schuyler	0	0	1	0	1	2	1	1	3	1	0	0	0	0	0	0	1	0	0	0	1	0	1	0	0	1	1	15
Seneca	0	0	1	1	0	1	0	2	1	2	2	2	0	1	0	1	0	0	2	0	2	0	0	2	0	1	0	21
Steuben	1	2	3	4	4	4	5	9	5	3	2	5	3	8	5	3	1	2	2	2	2	1	3	5	1	3	2	90
Suffolk	104	103	145	190	189	210	273	270	335	301	313	252	186	137	136	138	123	118	132	108	101	96	105	102	77	63	77	4,384
Sullivan	10	10	13	23	26	21	31	45	63	49	33	20	13	12	18	14	15	19	12	14	11	9	9	9	4	1	5	509
Tioga	1	0	1	2	1	3	3	1	2	1	4	1	0	0	0	1	2	0	4	1	1	0	1	5	1	1	0	37
Tompkins	2	2	4	5	5	8	7	12	5	10	7	13	7	7	5	5	4	8	3	3	4	4	9	4	4	3	1	151
Ulster	7	7	20	19	38	40	50	47	57	56	51	21	21	20	25	22	22	11	19	18	9	16	8	8	15	5	9	641
Warren	2	2	1	4	2	4	2	5	3	2	7	3	3	5	2	2	4	5	0	5	3	0	3	2	2	4	0	77
Washington	0	2	2	4	1	4	3	3	1	3	3	0	3	0	1	3	2	2	0	2	1	0	1	0	1	2	1	45
Wayne	4	3	1	2	1	5	4	8	7	5	5	6	6	7	2	4	6	3	2	3	2	7	2	2	4	2	4	107
Westchester	110	110	172	190	232	246	293	393	345	347	339	276	225	215	171	179	173	167	171	139	129	115	122	100	89	85	62	5,195
Wyoming	0	0	0	1	0	1	0	0	4	2	1	1	0	2	2	1	0	0	0	0	0	0	0	0	0	0	0	15
Yates	0	0	0	0	1	0	2	1	2	1	2	0	0	0	0	0	0	0	1	1	0	1	0	1	0	0	0	13

Summary Table CHIV Cases¹, Newly Diagnosed by County of Residence at Diagnosis and Year of Diagnosis²

County	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Albany	32	48	53	43	50	38	32	34	43	32	31	30
Allegany	1	1	3	0	1	2	2	0	1	2	0	2
Broome	14	20	13	14	13	16	16	13	8	14	7	12
Cattaraugus	0	1	4	3	1	1	3	5	3	2	0	4
Cayuga	9	5	2	5	2	5	4	4	2	9	3	5
Chautauqua	7	15	9	3	10	14	6	10	9	7	8	6
Chemung	2	5	4	6	5	7	6	2	7	1	3	6
Chenango	0	2	3	1	1	1	1	0	0	0	0	3
Clinton	5	3	3	1	4	2	2	6	4	0	0	2
Columbia	3	3	1	5	4	3	5	4	2	4	2	5
Cortland	0	1	1	0	0	1	1	3	2	1	2	1
Delaware	1	4	7	2	1	0	2	2	4	0	4	1
Dutchess	33	33	17	18	31	34	26	17	13	18	14	20
Erie	69	90	74	79	86	82	95	99	82	104	93	98
Essex	1	1	1	2	1	0	0	0	0	1	2	0
Franklin	2	4	0	1	2	0	0	0	0	4	1	1
Fulton	2	2	0	3	4	2	0	1	5	1	3	3
Genessee	1	5	4	3	0	1	2	5	1	1	3	3
Greene	0	6	2	1	4	1	2	5	3	4	0	0
Herkimer	1	2	1	2	0	1	2	1	2	0	6	4
Jefferson	4	1	3	2	2	3	3	3	3	1	8	2
Lewis	0	1	0	1	0	0	0	0	0	0	0	0
Livingston	2	0	0	3	1	1	1	2	1	4	2	2
Madison	1	1	0	2	1	3	4	0	1	2	1	1
Monroe	76	91	113	92	101	79	73	97	89	89	70	101
Montgomery	3	2	4	3	2	3	3	1	4	2	1	1
Nassau	149	156	130	112	113	124	110	100	133	89	100	120
Niagara	7	3	6	9	12	18	9	5	9	7	10	10
Oneida	9	9	14	14	16	11	9	12	9	13	13	9

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

² Data as of March 2013

 $^{^3}_{\it HIV}$ reporting began in June 2000. No earlier data are available

Summary Table C

HIV Cases¹, Newly Diagnosed by County of Residence at Diagnosis and Year of Diagnosis²

Excludes State Prison Inmates

County	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Onondaga	25	46	44	55	44	42	32	35	43	33	38	45
Ontario	1	4	6	3	2	3	2	5	3	1	3	1
Orange	38	39	32	42	34	43	39	35	28	32	24	21
Orleans	2	3	2	2	1	2	2	2	1	1	1	5
Oswego	1	5	11	3	3	3	3	5	4	7	5	3
Otsego	5	4	5	2	2	2	3	3	1	0	4	1
Putnam	7	3	3	5	1	2	5	5	2	9	6	4
Rensselaer	11	8	14	12	20	9	9	9	11	7	13	7
Rockland	27	29	19	25	32	35	33	14	24	21	20	25
St Lawrence	6	3	4	4	5	0	2	0	1	2	2	2
Saratoga	4	10	4	9	7	9	7	7	5	7	3	6
Schenectady	9	18	26	22	24	16	17	11	11	7	11	16
Schoharie	0	2	5	1	2	0	0	1	2	0	0	0
Schuyler	0	0	0	2	0	1	1	1	1	0	1	0
Seneca	2	1	2	2	2	0	0	1	2	0	2	1
Steuben	6	5	1	2	1	2	2	2	3	3	3	2
Suffolk	86	124	121	143	136	123	111	126	108	108	113	104
Sullivan	12	14	10	15	19	6	8	10	10	1	3	4
Tioga	1	4	0	2	2	3	1	0	3	1	1	1
Tompkins	4	2	10	4	6	5	7	5	6	4	5	3
Ulster	10	16	16	18	16	17	19	5	9	11	9	12
Warren	2	4	5	4	4	1	1	2	1	1	3	0
Washington	1	1	0	2	1	0	2	0	1	1	3	4
Wayne	5	7	9	3	4	6	9	3	4	5	5	4
Westchester	209	139	133	129	148	109	144	129	128	109	115	99
Wyoming	2	0	0	1	0	1	0	0	0	0	0	1
Yates	1	0	1	2	1	0	2	0	0	0	0	0

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^3}_{\it HIV}$ reporting began in June 2000. No earlier data are available

Summary Table D

Deaths Among AIDS Cases by County of Residence at Diagnosis and Year of Death 2

County	< 1985	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Albany	2	7	13	10	14	22	37	34	49	49	61	49	42	31	25	21	23	21	13	15	2004	12	20	16	13	12	17	7
Allegany	0	0	2	0	0	0	1	0	0	1	01	2	3	0	0	0	1	0	0	13	0	0	0	0	0	0	0	0
Broome	1	0	4	4	4	6	6	15	15	12	11	18	17	3	6	3	4	7	5	6	11	6	6	6	6	1	6	1
Cattaraugus	0	0	2	1	0	1	2	2	13	1	4	2	3	1	0	2	0	2	1	1	0	0	1	1	0	0	0	0
Cayuga	3	0	0	2	3	2	2	1	4	3	3	5	3	1	0	1	1	0	1	0	1	2	2	2	0	0	1	0
Chautauqua	0	0	2	0	1	4	0	1	1	7	6	7	7	3	2	5	1	1	2	1	6	3	2	2	2	4	4	4
Chemung	1	0	1	2	3	7	6	5	11	8	12	10	8	1	0	1	1	4	4	1	2	1	2	3	2	0	1	1
Chenango	0	0	0	0	2	1	0	0	2	0	1	4	2	0	0	0	1	1	1	0	1	1	2	0	0	0	1	2
Clinton	0	2	1	1	0	1	0	3	6	2	6	2	1	1	2	0	0	2	0	0	0	3	0	1	0	0	0	0
Columbia	0	0	3	1	1	4	2	7	9	5	2	3	5	2	1	1	3	1	2	0	1	0	1	0	0	1	1	0
Cortland	0	0	0	0	0	3	0	0	0	2	0	1	0	0	2	1	0	0	1	2	1	2	0	0	0	0	0	0
Delaware	0	0	0	0	0	1	2	1	2	3	1	1	3	0	1	1	0	2	1	1	0	2	0	0	0	0	1	1
Dutchess	3	3	8	16	24	33	18	35	33	50	59	68	47	23	16	28	15	17	23	16	18	10	16	10	8	11	13	8
Erie	5	11	12	19	19	49	62	70	80	71	114	154	81	49	40	41	46	51	30	50	39	37	36	37	29	34	30	24
Essex	2	1	0	0	1	2	1	1	4	2	1	1	2	2	0	0	0	0	2	0	0	1	1	0	0	0	0	0
Franklin	0	0	0	1	1	1	0	1	1	2	2	2	1	1	0	0	0	1	2	0	1	0	1	0	0	2	0	0
Fulton	0	1	2	0	1	1	0	3	2	1	4	1	0	1	0	1	0	1	0	1	0	1	0	1	1	0	0	0
Genessee	0	0	0	0	1	3	1	3	3	2	2	1	2	0	1	1	0	0	2	0	0	2	1	0	0	1	1	1
Greene	1	0	0	1	2	3	0	1	4	1	5	2	2	1	1	0	5	1	1	0	2	2	1	0	2	2	1	1
Herkimer	0	0	1	0	2	1	1	1	4	3	0	2	2	0	0	0	0	0	0	1	2	0	0	0	0	0	0	1
Jefferson	0	0	0	3	1	0	2	1	3	3	2	7	4	1	2	2	0	1	0	1	2	1	1	2	0	0	0	1
Lewis	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Livingston	0	0	2	2	1	0	3	2	4	4	1	2	3	1	1	1	1	2	1	0	0	1	0	0	1	2	0	1
Madison	0	0	1	0	1	1	1	2	1	2	2	3	1	1	0	1	0	1	1	0	1	3	0	0	1	0	1	1
Monroe	6	10	17	26	31	46	50	85	94	79	116	122	97	59	46	51	53	38	45	41	53	41	42	40	37	25	26	32
Montgomery	1	0	0	0	1	4	2	1	5	3	5	5	3	1	0	1	3	2	0	3	1	2	0	0	2	3	0	1
Nassau	41	38	70	101	114	149	150	188	191	201	238	224	137	80	68	74	60	67	56	79	54	55	53	59	41	39	43	28
Niagara	0	0	4	3	3	8	10	13	21	11	20	9	6	9	4	2	4	4	1	3	5	13	4	2	4	4	2	3
Oneida	1	2	3	4	5	9	9	11	14	28	20	18	21	10	6	7	9	5	3	6	5	8	1	9	5	4	4	1

¹ Death may be due to causes other than AIDS

²Data as of March 2013

Summary Table D

Deaths Among AIDS Cases by County of Residence at Diagnosis and Year of Death 2

County	< 1985	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011
Onondaga	5	13	16	12	24	24	21	34	53	56	80	65	57	19	20	20	24	17	16	14	13	12	19	23	25	18	21	15
Ontario	0	0	0	1	1	1	2	0	2	6	2	3	1	2	1	0	0	1	1	0	0	2	1	0	0	2	2	0
Orange	3	5	11	18	20	38	30	57	59	68	58	84	44	34	23	24	30	17	17	23	20	24	19	18	23	13	11	6
Orleans	0	0	0	0	0	0	0	2	4	2	0	3	2	1	1	2	0	0	1	0	2	3	1	0	0	1	1	0
Oswego	0	0	2	1	0	6	3	4	2	7	2	4	4	0	0	1	1	1	0	4	0	4	1	1	0	3	2	1
Otsego	1	0	0	0	2	3	4	4	3	0	4	6	4	0	0	2	2	0	1	0	2	1	5	1	3	0	0	2
Putnam	1	4	3	5	5	4	7	3	7	15	12	14	6	3	4	1	2	0	2	1	1	1	0	1	1	1	0	2
Rensselaer	1	0	1	4	4	3	8	10	11	17	11	14	12	8	2	5	5	7	4	5	5	7	5	3	4	4	5	7
Rockland	11	8	16	15	17	33	27	20	46	49	47	46	28	15	18	20	13	19	20	22	12	10	10	3	7	11	9	4
St Lawrence	0	1	1	0	1	4	2	3	3	1	4	1	1	0	0	2	1	1	1	0	2	3	1	0	1	1	3	0
Saratoga	0	0	3	5	2	5	3	7	12	7	16	12	4	5	4	1	4	7	3	3	4	3	4	1	2	2	1	0
Schenectady	4	2	10	5	6	9	17	14	21	11	21	16	21	10	11	8	8	6	11	4	8	8	8	7	7	6	5	6
Schoharie	0	1	1	0	0	1	1	3	1	5	2	4	3	0	0	2	1	0	0	0	0	0	0	0	0	0	0	0
Schuyler	0	0	0	1	0	0	0	2	1	4	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Seneca	0	0	0	0	1	0	1	0	1	2	0	1	2	1	0	0	0	0	2	0	0	0	0	1	0	0	0	0
Steuben	0	0	2	3	3	0	3	3	6	4	3	5	5	1	1	3	0	1	3	0	2	2	0	0	3	0	1	0
Suffolk	24	35	70	81	126	136	142	166	211	196	226	207	156	94	79	74	57	67	71	66	64	77	55	53	42	38	39	34
Sullivan	1	4	4	12	17	19	16	17	26	22	37	43	22	13	6	6	16	10	8	9	4	6	6	1	6	1	5	4
Tioga	0	1	0	0	0	2	2	0	2	4	1	0	2	1	0	0	0	1	1	0	0	0	0	0	0	0	1	0
Tompkins	1	0	2	2	5	2	4	6	5	5	7	4	5	4	3	2	1	2	0	2	2	1	2	0	1	1	2	0
Ulster	1	4	4	9	10	14	23	24	32	36	35	30	36	18	7	8	13	14	10	7	11	5	6	8	12	7	6	8
Warren	1	1	1	2	0	0	5	5	1	1	3	1	4	0	0	0	1	5	0	2	1	3	1	1	3	0	1	1
Washington	0	0	1	1	1	3	1	3	4	3	1	1	1	2	0	0	0	1	0	0	2	1	0	1	0	0	1	0
Wayne	1	2	1	1	2	3	2	4	2	8	5	5	1	0	2	3	0	2	1	1	1	2	1	1	1	1	3	0
Westchester	17	40	75	106	129	156	184	186	224	224	263	260	167	100	101	87	112	65	74	89	68	67	60	64	71	60	61	40
Wyoming	0	0	0	0	1	0	0	1	0	2	2	0	0	0	1	0	0	0	0	1	0	0	0	1	0	0	0	1
Yates	0	0	0	0	0	0	0	2	0	2	2	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0

¹ Death may be due to causes other than AIDS

²Data as of March 2013

Living HIV and AIDS Cases as of December 2011

by Gender, Age, Race/Ethnicity and Risk Excludes State Prison Inmates

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	1 AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		440	100.0	144.9	547	100.0	180.2	987	100.0	325.1	303,565
Gender	Male	288	65.5		367	67.1		655	66.4		
	Female	152	34.5		180	32.9		332	33.6		
Current Age	12 & under	6	1.4		0	0.0		6	0.6		
	13-19	8	1.8		4	0.7		12	1.2		
	20-24	19	4.3		6	1.1		25	2.5		
	25-29	23	5.2		9	1.6		32	3.2		
	30-39	79	18.0		64	11.7		143	14.5		
	40-49	154	35.0		208	38.0		362	36.7		
	50-59	102	23.2		188	34.4		290	29.4		
	60+	48	10.9		68	12.4		116	11.8		
	Unk	1	0.2		0	0.0		1	0.1		
Race/Ethnicity	White	189	43.0		177	32.4		366	37.1		
	Black	172	39.1		216	39.5		388	39.3		
	Hispanic	42	9.5		86	15.7		128	13.0		
	Multi Race	33	7.5		63	11.5		96	9.7		
	Others	4	0.9		5	0.9		9	0.9		
Risk	MSM	188	42.7		196	35.8		384	38.9		
	IDU	48	10.9		103	18.8		151	15.3		
	MSM/IDU	14	3.2		26	4.8		40	4.1		
	Heterosexual	88	20.0		120	21.9		208	21.1		
	Fem Pres Het Cont	50	11.4		50	9.1		100	10.1		
	Blood Prod	2	0.5		0	0.0		2	0.2		
	Pediatric Risk	16	3.6		8	1.5		24	2.4		
	Unk	34	7.7		44	8.0		78	7.9		

¹ Data as of March 2013

²Per 100,000 population

Living HIV and AIDS Cases as of December 2011

by Gender and Risk Excludes State Prison Inmates

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	/Iale	Fe	male	Т	otal	M	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		288	100.0	152	100.0	440	100.0	367	100.0	180	100.0	547	100.0	655	100.0	332	100.0	987	100.0
Risk	MSM	188	65.3	0	0.0	188	42.7	196	53.4	0	0.0	196	35.8	384	58.6	0	0.0	384	38.9
	IDU	27	9.4	21	13.8	48	10.9	65	17.7	38	21.1	103	18.8	92	14.0	59	17.8	151	15.3
	MSM/IDU	14	4.9	0	0.0	14	3.2	26	7.1	0	0.0	26	4.8	40	6.1	0	0.0	40	4.1
	Heterosexual	14	4.9	74	48.7	88	20.0	32	8.7	88	48.9	120	21.9	46	7.0	162	48.8	208	21.1
	Fem Pres Het Cont	0	0.0	50	32.9	50	11.4	0	0.0	50	27.8	50	9.1	0	0.0	100	30.1	100	10.1
	Blood Prod	2	0.7	0	0.0	2	0.5	0	0.0	0	0.0	0	0.0	2	0.3	0	0.0	2	0.2
	Pediatric Risk	9	3.1	7	4.6	16	3.6	4	1.1	4	2.2	8	1.5	13	2.0	11	3.3	24	2.4
	Unk	34	11.8	0	0.0	34	7.7	44	12.0	0	0.0	44	8.0	78	11.9	0	0.0	78	7.9

Average Annual Newly Diagnosed HIV and AIDS Cases (2009-2011) and Cumulative AIDS (through 2011)

by Gender, Age, Race/Ethnicity and Risk Excludes State Prison Inmates

		HIV Diagn	oses (Annua	ıl Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		31.0	100.0	10.2	21.3	100.0	7.0	1,196	100.0	303,565
Gender	Male	20.7	66.7		14.7	68.8		893	74.7	
	Female	10.3	33.3		6.7	31.3		303	25.3	
Age at Diagnosis	12 & under	0.3	1.1		0.0	0.0		14	1.2	
	13-19	1.0	3.2		0.3	1.6		11	0.9	
	20-24	3.0	9.7		0.7	3.1		32	2.7	
	25-29	3.7	11.8		1.3	6.3		165	13.8	
	30-39	8.3	26.9		5.0	23.4		455	38.0	
	40-49	7.7	24.7		8.0	37.5		374	31.3	
	50-59	4.7	15.1		5.0	23.4		123	10.3	
	60+	2.3	7.5		1.0	4.7		22	1.8	
	Unk	0.0	0.0		0.0	0.0		0	0.0	
Race/Ethnicity	White	12.0	38.7		4.3	20.3		464	38.8	
	Black	13.3	43.0		11.7	54.7		471	39.4	
	Hispanic	2.7	8.6		2.0	9.4		170	14.2	
	Multi Race	2.0	6.5		3.0	14.1		85	7.1	
	Others	1.0	3.2		0.3	1.6		6	0.5	
Risk	MSM	13.0	41.9		6.3	29.7		433	36.2	
	IDU	1.3	4.3		2.3	10.9		343	28.7	
	MSM/IDU	0.3	1.1		1.0	4.7		64	5.4	
	Heterosexual	7.7	24.7		6.7	31.3		177	14.8	
	Fem Pres Het Cont	4.0	12.9		1.3	6.3		68	5.7	
	Blood Prod	0.0	0.0		0.0	0.0		8	0.7	
	Pediatric Risk	0.3	1.1		0.3	1.6		18	1.5	
	Unk	4.3	14.0		3.3	15.6		85	7.1	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

Newly Diagnosed HIV and AIDS Cases (2009-2011) and Cumulative AIDS (through 2011)

by Gender and Risk Excludes State Prison Inmates

		I	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	M	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		20.7	100.0	10.3	100.0	31.0	100.0	14.7	100.0	6.7	100.0	21.3	100.0	893	100.0	303	100.0	1,196	100.0
Risk	MSM	13.0	62.9	0.0	0.0	13.0	41.9	6.3	43.2	0.0	0.0	6.3	29.7	433	48.5	0	0.0	433	36.2
	IDU	1.0	4.8	0.3	3.2	1.3	4.3	1.7	11.4	0.7	10.0	2.3	10.9	237	26.5	106	35.0	343	28.7
	MSM/IDU	0.3	1.6	0.0	0.0	0.3	1.1	1.0	6.8	0.0	0.0	1.0	4.7	64	7.2	0	0.0	64	5.4
	Heterosexual	1.7	8.1	6.0	58.1	7.7	24.7	2.0	13.6	4.7	70.0	6.7	31.3	55	6.2	122	40.3	177	14.8
	Fem Pres Het Cont	0.0	0.0	4.0	38.7	4.0	12.9	0.0	0.0	1.3	20.0	1.3	6.3	0	0.0	68	22.4	68	5.7
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	7	0.8	1	0.3	8	0.7
	Pediatric Risk	0.3	1.6	0.0	0.0	0.3	1.1	0.3	2.3	0.0	0.0	0.3	1.6	12	1.3	6	2.0	18	1.5
	Unk	4.3	21.0	0.0	0.0	4.3	14.0	3.3	22.7	0.0	0.0	3.3	15.6	85	9.5	0	0.0	85	7.1

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		17	9
1986		16	13
1987		26	10
1988		32	14
1989		48	22
1990		53	37
1991		77	34
1992		88	49
1993		94	49
1994	٠	69	61
1995	٠	73	49
1996	٠	63	42
1997	•	54	31
1998	٠	43	25
1999		44	21
2000*	32	40	23
2001	48	34	21
2002	53	48	13
2003	43	42	15
2004	50	55	20
2005	38	29	12
2006	32	40	20
2007	34	21	16
2008	43	26	13
2009	32	20	12
2010	31	28	17
2011	30	16	7

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

Living HIV and AIDS Cases as of December 2011

by Gender, Age, Race/Ethnicity and Risk Excludes State Prison Inmates

		Livin	g HIV (not	AIDS) Cases]	Living A	AIDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		10	100.0	20.5	17	100.0	34.9	27	100.0	55.4	48,778
Gender	Male	7	70.0		12	70.6		19	70.4		
	Female	3	30.0		5	29.4		8	29.6		
Current Age	20-24	1	10.0		0	0.0		1	3.7		
	25-29	0	0.0		0	0.0		0	0.0		
	30-39	3	30.0		2	11.8		5	18.5		
	40-49	1	10.0		10	58.8		11	40.7		
	50-59	3	30.0		5	29.4		8	29.6		
	60+	2	20.0		0	0.0		2	7.4		
Race/Ethnicity	White	8	80.0		10	58.8		18	66.7		
	Black	0	0.0		0	0.0		0	0.0		
	Hispanic	0	0.0		2	11.8		2	7.4		
	Multi Race	2	20.0		5	29.4		7	25.9		
Risk	MSM	5	50.0		8	47.1		13	48.1		
	IDU	3	30.0		1	5.9		4	14.8		
	MSM/IDU	0	0.0		1	5.9		1	3.7		
	Heterosexual	2	20.0		4	23.5		6	22.2		
	Fem Pres Het Cont	0	0.0		1	5.9		1	3.7		
	Blood Prod	0	0.0		0	0.0		0	0.0		
	Unk	0	0.0		2	11.8		2	7.4		

¹ Data as of March 2013

²Per 100,000 population

Living HIV and AIDS Cases as of December 2011

by Gender and Risk Excludes State Prison Inmates

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	/Iale	Fe	male	Т	otal	M	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		7	100.0	3	100.0	10	100.0	12	100.0	5	100.0	17	100.0	19	100.0	8	100.0	27	100.0
Risk	MSM	5	71.4	0	0.0	5	50.0	8	66.7	0	0.0	8	47.1	13	68.4	0	0.0	13	48.1
	IDU	2	28.6	1	33.3	3	30.0	1	8.3	0	0.0	1	5.9	3	15.8	1	12.5	4	14.8
	MSM/IDU	0	0.0	0	0.0	0	0.0	1	8.3	0	0.0	1	5.9	1	5.3	0	0.0	1	3.7
	Heterosexual	0	0.0	2	66.7	2	20.0	0	0.0	4	80.0	4	23.5	0	0.0	6	75.0	6	22.2
	Fem Pres Het Cont	0	0.0	0	0.0	0	0.0	0	0.0	1	20.0	1	5.9	0	0.0	1	12.5	1	3.7
	Blood Prod	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Unk	0	0.0	0	0.0	0	0.0	2	16.7	0	0.0	2	11.8	2	10.5	0	0.0	2	7.4

Average Annual Newly Diagnosed HIV and AIDS Cases (2009-2011) and Cumulative AIDS (through 2011)

by Gender, Age, Race/Ethnicity and Risk Excludes State Prison Inmates

		HIV Diagn	oses (Annua	l Average)	AIDS Diagr	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		1.3	100.0	2.7	1.3	100.0	2.7	28	100.0	48,778
Gender	Male	1.3	100.0		1.3	100.0		21	75.0	
	Female	0.0	0.0		0.0	0.0		7	25.0	
Age at Diagnosis	20-24	0.0	0.0		0.0	0.0		2	7.1	
	25-29	0.0	0.0		0.0	0.0		4	14.3	
	30-39	0.3	25.0		0.0	0.0		12	42.9	
	40-49	0.7	50.0		0.7	50.0		7	25.0	
	50-59	0.3	25.0		0.7	50.0		2	7.1	
	60+	0.0	0.0		0.0	0.0		1	3.6	
Race/Ethnicity	White	0.7	50.0		0.3	25.0		19	67.9	
	Black	0.0	0.0		0.0	0.0		1	3.6	
	Hispanic	0.0	0.0		0.0	0.0		3	10.7	
	Multi Race	0.7	50.0		1.0	75.0		5	17.9	
Risk	MSM	1.3	100.0		1.3	100.0		13	46.4	
	IDU	0.0	0.0		0.0	0.0		4	14.3	
	MSM/IDU	0.0	0.0		0.0	0.0		2	7.1	
	Heterosexual	0.0	0.0		0.0	0.0		4	14.3	
	Fem Pres Het Cont	0.0	0.0		0.0	0.0		1	3.6	
	Blood Prod	0.0	0.0		0.0	0.0		1	3.6	
	Unk	0.0	0.0		0.0	0.0		3	10.7	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

Newly Diagnosed HIV and AIDS Cases (2009-2011) and Cumulative AIDS (through 2011)

by Gender and Risk Excludes State Prison Inmates

]	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Averag	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fei	nale	To	otal	M	ale	Fer	nale	To	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		1.3	100.0	0.0	0.0	1.3	100.0	1.3	100.0	0.0	0.0	1.3	100.0	21	100.0	7	100.0	28	100.0
Risk	MSM	1.3	100.0	0.0	0.0	1.3	100.0	1.3	100.0	0.0	0.0	1.3	100.0	13	61.9	0	0.0	13	46.4
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3	14.3	1	14.3	4	14.3
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	9.5	0	0.0	2	7.1
	Heterosexual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	4	57.1	4	14.3
	Fem Pres Het Cont	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	1	14.3	1	3.6
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	1	14.3	1	3.6
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3	14.3	0	0.0	3	10.7

¹Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		2	
1986			2
1989		1	
1990		1	1
1991		1	
1992		1	
1993			1
1994		1	
1995		4	2
1996		2	3
1998		1	
1999		1	
2000*	1		1
2001	1	•	
2002	3	4	
2003		1	1
2004	1	1	
2005	2		
2006	2	2	
2008	1	1	
2009	2	3	
2011	2	1	

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		142	100.0	71.3	207	100.0	104.0	349	100.0	175.3	199,031
Gender	Male	85	59.9		144	69.6		229	65.6		
	Female	57	40.1		63	30.4		120	34.4		
Current Age	12 & under	0	0.0		0	0.0		0	0.0		
	13-19	3	2.1		0	0.0		3	0.9		
	20-24	7	4.9		4	1.9		11	3.2		
	25-29	11	7.7		7	3.4		18	5.2		
	30-39	28	19.7		30	14.5		58	16.6		
	40-49	46	32.4		67	32.4		113	32.4		
	50-59	33	23.2		75	36.2		108	30.9		
	60+	14	9.9		24	11.6		38	10.9		
Race/Ethnicity	White	72	50.7		106	51.2		178	51.0		
	Black	38	26.8		47	22.7		85	24.4		
	Hispanic	21	14.8		36	17.4		57	16.3		
	Multi Race	9	6.3		18	8.7		27	7.7		
	Others	2	1.4		0	0.0		2	0.6		
Risk	MSM	52	36.6		71	34.3		123	35.2		
	IDU	19	13.4		38	18.4		57	16.3		
	MSM/IDU	2	1.4		14	6.8		16	4.6		
	Heterosexual	31	21.8		34	16.4		65	18.6		
	Fem Pres Het Cont	24	16.9		16	7.7		40	11.5		
	Blood Prod	0	0.0		1	0.5		1	0.3		
	Pediatric Risk	3	2.1		4	1.9		7	2.0		
	Unk	11	7.7		29	14.0		40	11.5		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	/Iale	Fe	male	Т	otal	M	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		85	100.0	57	100.0	142	100.0	144	100.0	63	100.0	207	100.0	229	100.0	120	100.0	349	100.0
Risk	MSM	52	61.2	0	0.0	52	36.6	71	49.3	0	0.0	71	34.3	123	53.7	0	0.0	123	35.2
	IDU	11	12.9	8	14.0	19	13.4	21	14.6	17	27.0	38	18.4	32	14.0	25	20.8	57	16.3
	MSM/IDU	2	2.4	0	0.0	2	1.4	14	9.7	0	0.0	14	6.8	16	7.0	0	0.0	16	4.6
	Heterosexual	6	7.1	25	43.9	31	21.8	7	4.9	27	42.9	34	16.4	13	5.7	52	43.3	65	18.6
	Fem Pres Het Cont	0	0.0	24	42.1	24	16.9	0	0.0	16	25.4	16	7.7	0	0.0	40	33.3	40	11.5
	Blood Prod	0	0.0	0	0.0	0	0.0	1	0.7	0	0.0	1	0.5	1	0.4	0	0.0	1	0.3
	Pediatric Risk	3	3.5	0	0.0	3	2.1	2	1.4	2	3.2	4	1.9	5	2.2	2	1.7	7	2.0
	Unk	11	12.9	0	0.0	11	7.7	28	19.4	1	1.6	29	14.0	39	17.0	1	0.8	40	11.5

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		11.0	100.0	5.5	10.7	100.0	5.4	396	100.0	199,031
Gender	Male	7.7	69.7		7.3	68.8		288	72.7	
	Female	3.3	30.3		3.3	31.3		108	27.3	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		2	0.5	
	13-19	0.3	3.0		0.7	6.3		7	1.8	
	20-24	2.0	18.2		0.7	6.3		19	4.8	
	25-29	1.0	9.1		0.3	3.1		45	11.4	
	30-39	3.0	27.3		2.0	18.8		154	38.9	
	40-49	2.0	18.2		4.7	43.8		115	29.0	
	50-59	2.3	21.2		2.3	21.9		47	11.9	
	60+	0.3	3.0		0.0	0.0		7	1.8	
Race/Ethnicity	White	6.0	54.5		5.7	53.1		230	58.1	
	Black	2.7	24.2		2.7	25.0		82	20.7	
	Hispanic	1.7	15.2		2.0	18.8		59	14.9	
	Multi Race	0.3	3.0		0.3	3.1		25	6.3	
	Others	0.3	3.0		0.0	0.0		0	0.0	
Risk	MSM	5.0	45.5		3.0	28.1		149	37.6	
	IDU	0.7	6.1		2.3	21.9		90	22.7	
	MSM/IDU	0.3	3.0		1.0	9.4		28	7.1	
	Heterosexual	2.7	24.2		1.7	15.6		51	12.9	
	Fem Pres Het Cont	1.0	9.1		0.0	0.0		24	6.1	
	Blood Prod	0.0	0.0		0.0	0.0		2	0.5	
	Pediatric Risk	0.0	0.0		0.7	6.3		7	1.8	
	Unk	1.3	12.1		2.0	18.8		45	11.4	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Diag	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		7.7	100.0	3.3	100.0	11.0	100.0	7.3	100.0	3.3	100.0	10.7	100.0	288	100.0	108	100.0	396	100.0
Risk	MSM	5.0	65.2	0.0	0.0	5.0	45.5	3.0	40.9	0.0	0.0	3.0	28.1	149	51.7	0	0.0	149	37.6
	IDU	0.3	4.3	0.3	10.0	0.7	6.1	1.0	13.6	1.3	40.0	2.3	21.9	55	19.1	35	32.4	90	22.7
	MSM/IDU	0.3	4.3	0.0	0.0	0.3	3.0	1.0	13.6	0.0	0.0	1.0	9.4	28	9.7	0	0.0	28	7.1
	Heterosexual	1.0	13.0	1.7	50.0	2.7	24.2	0.7	9.1	1.0	30.0	1.7	15.6	9	3.1	42	38.9	51	12.9
	Fem Pres Het Cont	0.0	0.0	1.0	30.0	1.0	9.1	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	24	22.2	24	6.1
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	0.7	0	0.0	2	0.5
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.7	20.0	0.7	6.3	2	0.7	5	4.6	7	1.8
	Unk	1.0	13.0	0.3	10.0	1.3	12.1	1.7	22.7	0.3	10.0	2.0	18.8	43	14.9	2	1.9	45	11.4

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		4	1
1986		2	4
1987		6	4
1988		10	4
1989		11	6
1990		18	6
1991		24	15
1992		20	15
1993		23	12
1994		19	11
1995		17	18
1996		23	17
1997		22	3
1998		13	6
1999		10	3
2000*	14	9	4
2001	20	15	7
2002	13	15	5
2003	14	14	6
2004	13	17	11
2005	16	14	6
2006	16	25	6
2007	13	24	6
2008	8	9	6
2009	14	16	1
2010	7	9	6
2011	12	7	1

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		19	100.0	23.8	30	100.0	37.6	49	100.0	61.4	79,832
Gender	Male	10	52.6		24	80.0		34	69.4		
	Female	9	47.4		6	20.0		15	30.6		
Current Age	13-19	0	0.0		0	0.0		0	0.0		
	20-24	2	10.5		0	0.0		2	4.1		
	25-29	1	5.3		0	0.0		1	2.0		
	30-39	6	31.6		7	23.3		13	26.5		
	40-49	4	21.1		14	46.7		18	36.7		
	50-59	2	10.5		8	26.7		10	20.4		
	60+	4	21.1		1	3.3		5	10.2		
Race/Ethnicity	White	18	94.7		24	80.0		42	85.7		
	Black	1	5.3		2	6.7		3	6.1		
	Hispanic	0	0.0		1	3.3		1	2.0		
	Multi Race	0	0.0		2	6.7		2	4.1		
	Others	0	0.0		1	3.3		1	2.0		
Risk	MSM	8	42.1		14	46.7		22	44.9		
	IDU	0	0.0		3	10.0		3	6.1		
	MSM/IDU	1	5.3		3	10.0		4	8.2		
	Heterosexual	9	47.4		5	16.7		14	28.6		
	Fem Pres Het Cont	1	5.3		3	10.0		4	8.2		
	Blood Prod	0	0.0		1	3.3		1	2.0		
	Unk	0	0.0		1	3.3		1	2.0		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	/Iale	Fe	male	Т	otal	М	ale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		10	100.0	9	100.0	19	100.0	24	100.0	6	100.0	30	100.0	34	100.0	15	100.0	49	100.0
Risk	MSM	8	80.0	0	0.0	8	42.1	14	58.3	0	0.0	14	46.7	22	64.7	0	0.0	22	44.9
	IDU	0	0.0	0	0.0	0	0.0	3	12.5	0	0.0	3	10.0	3	8.8	0	0.0	3	6.1
	MSM/IDU	1	10.0	0	0.0	1	5.3	3	12.5	0	0.0	3	10.0	4	11.8	0	0.0	4	8.2
	Heterosexual	1	10.0	8	88.9	9	47.4	2	8.3	3	50.0	5	16.7	3	8.8	11	73.3	14	28.6
	Fem Pres Het Cont	0	0.0	1	11.1	1	5.3	0	0.0	3	50.0	3	10.0	0	0.0	4	26.7	4	8.2
	Blood Prod	0	0.0	0	0.0	0	0.0	1	4.2	0	0.0	1	3.3	1	2.9	0	0.0	1	2.0
	Unk	0	0.0	0	0.0	0	0.0	1	4.2	0	0.0	1	3.3	1	2.9	0	0.0	1	2.0

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		2.0	100.0	2.5	1.3	100.0	1.7	58	100.0	79,832
Gender	Male	2.0	100.0		1.0	75.0		43	74.1	
	Female	0.0	0.0		0.3	25.0		15	25.9	
Age at Diagnosis	13-19	0.0	0.0		0.0	0.0		1	1.7	
	20-24	0.3	16.7		0.0	0.0	•	3	5.2	
	25-29	0.3	16.7		0.0	0.0		9	15.5	
	30-39	0.3	16.7		0.7	50.0		29	50.0	
	40-49	0.7	33.3		0.7	50.0		14	24.1	
	50-59	0.3	16.7		0.0	0.0		1	1.7	
	60+	0.0	0.0		0.0	0.0		1	1.7	
Race/Ethnicity	White	2.0	100.0		1.3	100.0		47	81.0	
	Black	0.0	0.0		0.0	0.0		5	8.6	
	Hispanic	0.0	0.0		0.0	0.0		2	3.4	
	Multi Race	0.0	0.0		0.0	0.0	•	3	5.2	
	Others	0.0	0.0		0.0	0.0		1	1.7	
Risk	MSM	1.7	83.3		1.0	75.0		29	50.0	
	IDU	0.0	0.0		0.0	0.0	•	8	13.8	
	MSM/IDU	0.3	16.7		0.0	0.0		3	5.2	
	Heterosexual	0.0	0.0		0.0	0.0		10	17.2	
	Fem Pres Het Cont	0.0	0.0		0.3	25.0		4	6.9	
	Blood Prod	0.0	0.0		0.0	0.0		2	3.4	
	Unk	0.0	0.0		0.0	0.0		2	3.4	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		1	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	To	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		2.0	100.0	0.0	0.0	2.0	100.0	1.0	100.0	0.3	100.0	1.3	100.0	43	100.0	15	100.0	58	100.0
Risk	MSM	1.7	83.3	0.0	0.0	1.7	83.3	1.0	100.0	0.0	0.0	1.0	75.0	29	67.4	0	0.0	29	50.0
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	5	11.6	3	20.0	8	13.8
	MSM/IDU	0.3	16.7	0.0	0.0	0.3	16.7	0.0	0.0	0.0	0.0	0.0	0.0	3	7.0	0	0.0	3	5.2
	Heterosexual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	4.7	8	53.3	10	17.2
	Fem Pres Het Cont	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	100.0	0.3	25.0	0	0.0	4	26.7	4	6.9
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	4.7	0	0.0	2	3.4
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	4.7	0	0.0	2	3.4

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		1	
1986		2	2
1987	•	1	1
1988		1	
1989		2	1
1990	٠	2	2
1991		5	2
1992		4	1
1993	٠	4	1
1994		3	4
1995		2	2
1996		2	3
1997		2	1
1998		2	
1999			2
2000*	٠	2	
2001	1	1	2
2002	4	1	1
2003	3	1	1
2004	1	3	
2005	1	3	
2006	3	3	1
2007	5	5	1
2008	3	2	
2009	2	2	
2011	4	2	

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	1 AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		34	100.0	42.6	53	100.0	66.5	87	100.0	109.1	79,738
Gender	Male	22	64.7		40	75.5		62	71.3		
	Female	12	35.3		13	24.5		25	28.7		
Current Age	13-19	0	0.0		0	0.0		0	0.0		
	20-24	4	11.8		1	1.9		5	5.7		
	25-29	2	5.9		0	0.0		2	2.3		
	30-39	9	26.5		0	0.0		9	10.3		
	40-49	13	38.2		25	47.2		38	43.7		
	50-59	4	11.8		17	32.1		21	24.1		
	60+	2	5.9		10	18.9		12	13.8		
Race/Ethnicity	White	28	82.4		30	56.6		58	66.7		
	Black	4	11.8		17	32.1		21	24.1		
	Hispanic	2	5.9		3	5.7		5	5.7		
	Multi Race	0	0.0		3	5.7		3	3.4		
Risk	MSM	18	52.9		19	35.8		37	42.5		
	IDU	1	2.9		6	11.3		7	8.0		
	MSM/IDU	2	5.9		3	5.7		5	5.7		
	Heterosexual	7	20.6		13	24.5		20	23.0		
	Fem Pres Het Cont	4	11.8		2	3.8		6	6.9		
	Blood Prod	0	0.0		0	0.0		0	0.0		
	Pediatric Risk	0	0.0		0	0.0		0	0.0		
	Unk	2	5.9		10	18.9		12	13.8		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		M	ale	Fe	male	To	otal	N	Male	Fe	emale	7	otal	M	[ale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		22	100.0	12	100.0	34	100.0	40	100.0	13	100.0	53	100.0	62	100.0	25	100.0	87	100.0
Risk	MSM	18	81.8	0	0.0	18	52.9	19	47.5	0	0.0	19	35.8	37	59.7	0	0.0	37	42.5
	IDU	0	0.0	1	8.3	1	2.9	4	10.0	2	15.4	6	11.3	4	6.5	3	12.0	7	8.0
	MSM/IDU	2	9.1	0	0.0	2	5.9	3	7.5	0	0.0	3	5.7	5	8.1	0	0.0	5	5.7
	Heterosexual	0	0.0	7	58.3	7	20.6	4	10.0	9	69.2	13	24.5	4	6.5	16	64.0	20	23.0
	Fem Pres Het Cont	0	0.0	4	33.3	4	11.8	0	0.0	2	15.4	2	3.8	0	0.0	6	24.0	6	6.9
	Blood Prod	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Pediatric Risk	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Unk	2	9.1	0	0.0	2	5.9	10	25.0	0	0.0	10	18.9	12	19.4	0	0.0	12	13.8

		HIV Diagn	oses (Annua	l Average)	AIDS Diagr	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		5.7	100.0	7.1	2.0	100.0	2.5	95	100.0	79,738
Gender	Male	4.3	76.5		1.0	50.0		74	77.9	
	Female	1.3	23.5		1.0	50.0		21	22.1	
Age at Diagnosis	13-19	0.0	0.0		0.0	0.0		1	1.1	
	20-24	1.7	29.4		0.3	16.7	•	2	2.1	
	25-29	0.7	11.8		0.0	0.0		5	5.3	
	30-39	1.0	17.6		0.0	0.0		41	43.2	
	40-49	1.3	23.5		0.3	16.7		33	34.7	
	50-59	1.0	17.6		0.7	33.3		7	7.4	
	60+	0.0	0.0		0.7	33.3		6	6.3	
Race/Ethnicity	White	4.3	76.5		1.0	50.0		59	62.1	
	Black	1.0	17.6		0.7	33.3		24	25.3	
	Hispanic	0.0	0.0		0.0	0.0		8	8.4	
	Multi Race	0.3	5.9		0.3	16.7		4	4.2	
Risk	MSM	2.7	47.1		0.7	33.3		37	38.9	
	IDU	0.0	0.0		0.0	0.0		16	16.8	
	MSM/IDU	0.7	11.8		0.0	0.0		3	3.2	
	Heterosexual	1.3	23.5		1.0	50.0		18	18.9	
	Fem Pres Het Cont	0.0	0.0		0.0	0.0		5	5.3	
	Blood Prod	0.0	0.0		0.0	0.0		2	2.1	
	Pediatric Risk	0.0	0.0		0.0	0.0		1	1.1	
	Unk	1.0	17.6		0.3	16.7		13	13.7	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Diag	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	M	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		4.3	100.0	1.3	100.0	5.7	100.0	1.0	100.0	1.0	100.0	2.0	100.0	74	100.0	21	100.0	95	100.0
Risk	MSM	2.7	61.5	0.0	0.0	2.7	47.1	0.7	66.7	0.0	0.0	0.7	33.3	37	50.0	0	0.0	37	38.9
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	13	17.6	3	14.3	16	16.8
	MSM/IDU	0.7	15.4	0.0	0.0	0.7	11.8	0.0	0.0	0.0	0.0	0.0	0.0	3	4.1	0	0.0	3	3.2
	Heterosexual	0.0	0.0	1.3	100.0	1.3	23.5	0.0	0.0	1.0	100.0	1.0	50.0	6	8.1	12	57.1	18	18.9
	Fem Pres Het Cont	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	5	23.8	5	5.3
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	1.4	1	4.8	2	2.1
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	1.4	0	0.0	1	1.1
	Unk	1.0	23.1	0.0	0.0	1.0	17.6	0.3	33.3	0.0	0.0	0.3	16.7	13	17.6	0	0.0	13	13.7

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		3	3
1986		2	
1987		2	2
1988		2	3
1989		3	2
1990		3	2
1991		3	1
1992		7	4
1993		4	3
1994		4	3
1995		7	5
1996		5	3
1997		3	1
1998		5	
1999		2	1
2000*	9	4	1
2001	5	4	
2002	2	2	1
2003	5	5	
2004	2	3	1
2005	5	3	2
2006	4	3	2
2007	4	5	2
2008	2	5	
2009	9	2	
2010	3	2	1
2011	5	2	

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		90	100.0	67.0	79	100.0	58.8	169	100.0	125.8	134,368
Gender	Male	53	58.9		58	73.4		111	65.7		
	Female	37	41.1		21	26.6		58	34.3		
Current Age	12 & under	1	1.1		0	0.0		1	0.6		
	13-19	2	2.2		0	0.0		2	1.2		
	20-24	5	5.6		1	1.3		6	3.6		
	25-29	12	13.3		4	5.1		16	9.5		
	30-39	20	22.2		15	19.0		35	20.7		
	40-49	30	33.3		29	36.7		59	34.9		
	50-59	16	17.8		16	20.3		32	18.9		
	60+	4	4.4		14	17.7		18	10.7		
Race/Ethnicity	White	50	55.6		45	57.0		95	56.2		
	Black	7	7.8		9	11.4		16	9.5		
	Hispanic	28	31.1		18	22.8		46	27.2		
	Multi Race	5	5.6		7	8.9		12	7.1		
Risk	MSM	35	38.9		38	48.1		73	43.2		
	IDU	8	8.9		11	13.9		19	11.2		
	MSM/IDU	5	5.6		7	8.9		12	7.1		
	Heterosexual	32	35.6		17	21.5		49	29.0		
	Fem Pres Het Cont	5	5.6		2	2.5		7	4.1		
	Blood Prod	0	0.0		1	1.3		1	0.6		
	Pediatric Risk	3	3.3		0	0.0		3	1.8		
	Unk	2	2.2		3	3.8		5	3.0		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	Male	Fe	male	1	otal	M	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		53	100.0	37	100.0	90	100.0	58	100.0	21	100.0	79	100.0	111	100.0	58	100.0	169	100.0
Risk	MSM	35	66.0	0	0.0	35	38.9	38	65.5	0	0.0	38	48.1	73	65.8	0	0.0	73	43.2
	IDU	4	7.5	4	10.8	8	8.9	8	13.8	3	14.3	11	13.9	12	10.8	7	12.1	19	11.2
	MSM/IDU	5	9.4	0	0.0	5	5.6	7	12.1	0	0.0	7	8.9	12	10.8	0	0.0	12	7.1
	Heterosexual	5	9.4	27	73.0	32	35.6	1	1.7	16	76.2	17	21.5	6	5.4	43	74.1	49	29.0
	Fem Pres Het Cont	0	0.0	5	13.5	5	5.6	0	0.0	2	9.5	2	2.5	0	0.0	7	12.1	7	4.1
	Blood Prod	0	0.0	0	0.0	0	0.0	1	1.7	0	0.0	1	1.3	1	0.9	0	0.0	1	0.6
	Pediatric Risk	2	3.8	1	2.7	3	3.3	0	0.0	0	0.0	0	0.0	2	1.8	1	1.7	3	1.8
	Unk	2	3.8	0	0.0	2	2.2	3	5.2	0	0.0	3	3.8	5	4.5	0	0.0	5	3.0

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		7.0	100.0	5.2	4.7	100.0	3.5	156	100.0	134,368
Gender	Male	5.0	71.4		4.0	85.7		115	73.7	
	Female	2.0	28.6		0.7	14.3		41	26.3	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		0	0.0	
	13-19	0.0	0.0		0.3	7.1		4	2.6	
	20-24	1.0	14.3		0.0	0.0		9	5.8	
	25-29	1.0	14.3		0.3	7.1		20	12.8	
	30-39	1.0	14.3		0.3	7.1		55	35.3	
	40-49	3.3	47.6		2.3	50.0		41	26.3	
	50-59	0.7	9.5		1.3	28.6		20	12.8	
	60+	0.0	0.0		0.0	0.0		7	4.5	
Race/Ethnicity	White	4.3	61.9		3.0	64.3		91	58.3	
	Black	1.0	14.3		1.0	21.4		21	13.5	
	Hispanic	1.3	19.0		0.7	14.3		34	21.8	
	Multi Race	0.3	4.8		0.0	0.0		10	6.4	
Risk	MSM	3.3	47.6		3.0	64.3		69	44.2	
	IDU	1.3	19.0		0.7	14.3		30	19.2	
	MSM/IDU	0.7	9.5		0.3	7.1		11	7.1	
	Heterosexual	1.3	19.0		0.3	7.1		36	23.1	
	Fem Pres Het Cont	0.0	0.0		0.0	0.0		3	1.9	
_	Blood Prod	0.0	0.0		0.0	0.0		3	1.9	
	Pediatric Risk	0.0	0.0		0.0	0.0		0	0.0	
	Unk	0.3	4.8		0.3	7.1		4	2.6	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	M	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		5.0	100.0	2.0	100.0	7.0	100.0	4.0	100.0	0.7	100.0	4.7	100.0	115	100.0	41	100.0	156	100.0
Risk	MSM	3.3	66.7	0.0	0.0	3.3	47.6	3.0	75.0	0.0	0.0	3.0	64.3	69	60.0	0	0.0	69	44.2
	IDU	0.7	13.3	0.7	33.3	1.3	19.0	0.3	8.3	0.3	50.0	0.7	14.3	23	20.0	7	17.1	30	19.2
	MSM/IDU	0.7	13.3	0.0	0.0	0.7	9.5	0.3	8.3	0.0	0.0	0.3	7.1	11	9.6	0	0.0	11	7.1
	Heterosexual	0.0	0.0	1.3	66.7	1.3	19.0	0.0	0.0	0.3	50.0	0.3	7.1	5	4.3	31	75.6	36	23.1
	Fem Pres Het Cont	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	3	7.3	3	1.9
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3	2.6	0	0.0	3	1.9
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	0	0.0	0	0.0
	Unk	0.3	6.7	0.0	0.0	0.3	4.8	0.3	8.3	0.0	0.0	0.3	7.1	4	3.5	0	0.0	4	2.6

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		2	
1986			2
1987		1	
1988		2	1
1989		4	4
1990		2	
1991		4	1
1992		9	1
1993		9	7
1994		8	6
1995		12	7
1996		6	7
1997		3	3
1998		9	2
1999		5	5
2000*	7	6	1
2001	15	8	1
2002	9	12	2
2003	3	4	1
2004	10	7	6
2005	14	9	3
2006	6	5	2
2007	10	8	2
2008	9	7	2
2009	7	7	4
2010	8	4	4
2011	6	3	4

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		66	100.0	74.3	46	100.0	51.8	112	100.0	126.1	88,840
Gender	Male	44	66.7		35	76.1		79	70.5		
	Female	22	33.3		11	23.9		33	29.5		
Current Age	12 & under	0	0.0		0	0.0		0	0.0		
	13-19	1	1.5		0	0.0		1	0.9		
	20-24	5	7.6		0	0.0		5	4.5		
	25-29	3	4.5		3	6.5		6	5.4		
	30-39	13	19.7		7	15.2		20	17.9		
	40-49	26	39.4		16	34.8		42	37.5		
	50-59	11	16.7		13	28.3		24	21.4		
	60+	6	9.1		7	15.2		13	11.6		
	Unk	1	1.5		0	0.0		1	0.9		
Race/Ethnicity	White	45	68.2		28	60.9		73	65.2		
	Black	13	19.7		12	26.1		25	22.3		
	Hispanic	5	7.6		6	13.0		11	9.8		
	Multi Race	3	4.5		0	0.0		3	2.7		
Risk	MSM	24	36.4		17	37.0		41	36.6		
	IDU	6	9.1		13	28.3		19	17.0		
	MSM/IDU	5	7.6		2	4.3		7	6.3		
	Heterosexual	13	19.7		6	13.0		19	17.0		
	Fem Pres Het Cont	8	12.1		3	6.5		11	9.8		
	Pediatric Risk	0	0.0		0	0.0		0	0.0		
	Unk	10	15.2		5	10.9		15	13.4		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	Male	Fe	male	Т	otal	M	[ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		44	100.0	22	100.0	66	100.0	35	100.0	11	100.0	46	100.0	79	100.0	33	100.0	112	100.0
Risk	MSM	24	54.5	0	0.0	24	36.4	17	48.6	0	0.0	17	37.0	41	51.9	0	0.0	41	36.6
	IDU	3	6.8	3	13.6	6	9.1	9	25.7	4	36.4	13	28.3	12	15.2	7	21.2	19	17.0
	MSM/IDU	5	11.4	0	0.0	5	7.6	2	5.7	0	0.0	2	4.3	7	8.9	0	0.0	7	6.3
	Heterosexual	2	4.5	11	50.0	13	19.7	2	5.7	4	36.4	6	13.0	4	5.1	15	45.5	19	17.0
	Fem Pres Het Cont	0	0.0	8	36.4	8	12.1	0	0.0	3	27.3	3	6.5	0	0.0	11	33.3	11	9.8
	Pediatric Risk	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Unk	10	22.7	0	0.0	10	15.2	5	14.3	0	0.0	5	10.9	15	19.0	0	0.0	15	13.4

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		3.3	100.0	3.8	2.7	100.0	3.0	144	100.0	88,840
Gender	Male	3.0	90.0		2.0	75.0		110	76.4	
	Female	0.3	10.0		0.7	25.0		34	23.6	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		4	2.8	
	13-19	0.0	0.0		0.0	0.0		0	0.0	
	20-24	1.0	30.0		0.0	0.0		5	3.5	
	25-29	0.0	0.0		0.0	0.0		16	11.1	
	30-39	1.0	30.0		1.0	37.5		67	46.5	
	40-49	1.0	30.0		1.0	37.5		35	24.3	
	50-59	0.3	10.0		0.7	25.0		15	10.4	
	60+	0.0	0.0		0.0	0.0		2	1.4	
	Unk	0.0	0.0		0.0	0.0		0	0.0	
Race/Ethnicity	White	2.0	60.0		1.7	62.5		96	66.7	
	Black	1.0	30.0		1.0	37.5	•	31	21.5	
	Hispanic	0.3	10.0		0.0	0.0		16	11.1	
	Multi Race	0.0	0.0		0.0	0.0		1	0.7	
Risk	MSM	2.0	60.0		1.3	50.0		60	41.7	
	IDU	0.0	0.0		0.0	0.0		46	31.9	
	MSM/IDU	0.3	10.0		0.0	0.0		3	2.1	
	Heterosexual	0.3	10.0		0.0	0.0		17	11.8	
	Fem Pres Het Cont	0.0	0.0		0.7	25.0		5	3.5	
	Pediatric Risk	0.0	0.0		0.0	0.0		4	2.8	
	Unk	0.7	20.0		0.7	25.0		9	6.3	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		1	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fei	nale	To	otal	М	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		3.0	100.0	0.3	100.0	3.3	100.0	2.0	100.0	0.7	100.0	2.7	100.0	110	100.0	34	100.0	144	100.0
Risk	MSM	2.0	66.7	0.0	0.0	2.0	60.0	1.3	66.7	0.0	0.0	1.3	50.0	60	54.5	0	0.0	60	41.7
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	34	30.9	12	35.3	46	31.9
	MSM/IDU	0.3	11.1	0.0	0.0	0.3	10.0	0.0	0.0	0.0	0.0	0.0	0.0	3	2.7	0	0.0	3	2.1
	Heterosexual	0.0	0.0	0.3	100.0	0.3	10.0	0.0	0.0	0.0	0.0	0.0	0.0	3	2.7	14	41.2	17	11.8
	Fem Pres Het Cont	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.7	100.0	0.7	25.0	0	0.0	5	14.7	5	3.5
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	0.9	3	8.8	4	2.8
	Unk	0.7	22.2	0.0	0.0	0.7	20.0	0.7	33.3	0.0	0.0	0.7	25.0	9	8.2	0	0.0	9	6.3

¹Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		2	1
1986	•	4	1
1987		3	2
1988		4	3
1989		9	7
1990	•	9	6
1991	٠	14	5
1992		12	11
1993	٠	11	8
1994		12	12
1995		13	10
1996		4	8
1997		6	1
1998		3	
1999			1
2000*	2	3	1
2001	5	3	4
2002	4		4
2003	6	4	1
2004	5	3	2
2005	7	10	1
2006	6	1	2
2007	2	3	3
2008	7	3	2
2009	1	1	
2010	3	4	1
2011	6	3	1

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases		Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		16	100.0	31.9	30	100.0	59.9	46	100.0	91.8	50,118
Gender	Male	11	68.8		23	76.7		34	73.9		
	Female	5	31.3		7	23.3		12	26.1		
Current Age	13-19	1	6.3		0	0.0		1	2.2		
	20-24	1	6.3		1	3.3		2	4.3		
	25-29	0	0.0		0	0.0		0	0.0		
	30-39	1	6.3		3	10.0		4	8.7		
	40-49	5	31.3		11	36.7		16	34.8		
	50-59	7	43.8		9	30.0		16	34.8		
	60+	1	6.3		6	20.0		7	15.2		
Race/Ethnicity	White	15	93.8		22	73.3		37	80.4		
	Black	0	0.0		2	6.7		2	4.3		
	Hispanic	1	6.3		3	10.0		4	8.7		
	Multi Race	0	0.0		3	10.0		3	6.5		
Risk	MSM	6	37.5		17	56.7		23	50.0		
	IDU	6	37.5		1	3.3		7	15.2		
	MSM/IDU	0	0.0		1	3.3		1	2.2		
	Heterosexual	1	6.3		4	13.3		5	10.9		
	Fem Pres Het Cont	0	0.0		3	10.0		3	6.5		
	Blood Prod	1	6.3		0	0.0		1	2.2		
	Pediatric Risk	2	12.5		0	0.0		2	4.3		
_	Unk	0	0.0		4	13.3		4	8.7		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	Male	Fe	emale	1	otal	M	[ale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		11	100.0	5	100.0	16	100.0	23	100.0	7	100.0	30	100.0	34	100.0	12	100.0	46	100.0
Risk	MSM	6	54.5	0	0.0	6	37.5	17	73.9	0	0.0	17	56.7	23	67.6	0	0.0	23	50.0
	IDU	4	36.4	2	40.0	6	37.5	0	0.0	1	14.3	1	3.3	4	11.8	3	25.0	7	15.2
	MSM/IDU	0	0.0	0	0.0	0	0.0	1	4.3	0	0.0	1	3.3	1	2.9	0	0.0	1	2.2
	Heterosexual	0	0.0	1	20.0	1	6.3	1	4.3	3	42.9	4	13.3	1	2.9	4	33.3	5	10.9
	Fem Pres Het Cont	0	0.0	0	0.0	0	0.0	0	0.0	3	42.9	3	10.0	0	0.0	3	25.0	3	6.5
	Blood Prod	1	9.1	0	0.0	1	6.3	0	0.0	0	0.0	0	0.0	1	2.9	0	0.0	1	2.2
	Pediatric Risk	0	0.0	2	40.0	2	12.5	0	0.0	0	0.0	0	0.0	0	0.0	2	16.7	2	4.3
	Unk	0	0.0	0	0.0	0	0.0	4	17.4	0	0.0	4	13.3	4	11.8	0	0.0	4	8.7

		HIV Diagn	oses (Annua	ıl Average)	AIDS Diagr	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		1.0	100.0	2.0	1.7	100.0	3.3	50	100.0	50,118
Gender	Male	1.0	100.0		1.3	80.0		36	72.0	
	Female	0.0	0.0		0.3	20.0		14	28.0	
Age at Diagnosis	13-19	0.0	0.0		0.0	0.0		0	0.0	
	20-24	0.3	33.3		0.3	20.0		2	4.0	
	25-29	0.0	0.0		0.0	0.0		4	8.0	
	30-39	0.3	33.3		0.3	20.0		23	46.0	
	40-49	0.3	33.3		0.3	20.0		13	26.0	
	50-59	0.0	0.0		0.7	40.0		7	14.0	
	60+	0.0	0.0		0.0	0.0		1	2.0	
Race/Ethnicity	White	1.0	100.0		1.7	100.0		35	70.0	
	Black	0.0	0.0		0.0	0.0		6	12.0	
	Hispanic	0.0	0.0		0.0	0.0		6	12.0	
	Multi Race	0.0	0.0		0.0	0.0		3	6.0	
Risk	MSM	1.0	100.0		0.7	40.0		24	48.0	
	IDU	0.0	0.0		0.0	0.0		7	14.0	
	MSM/IDU	0.0	0.0		0.0	0.0		2	4.0	
	Heterosexual	0.0	0.0		0.7	40.0		8	16.0	
	Fem Pres Het Cont	0.0	0.0		0.0	0.0		5	10.0	
	Blood Prod	0.0	0.0		0.0	0.0		1	2.0	
	Pediatric Risk	0.0	0.0		0.0	0.0		0	0.0	
	Unk	0.0	0.0		0.3	20.0		3	6.0	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		1.0	100.0	0.0	0.0	1.0	100.0	1.3	100.0	0.3	100.0	1.7	100.0	36	100.0	14	100.0	50	100.0
Risk	MSM	1.0	100.0	0.0	0.0	1.0	100.0	0.7	50.0	0.0	0.0	0.7	40.0	24	66.7	0	0.0	24	48.0
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	4	11.1	3	21.4	7	14.0
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	5.6	0	0.0	2	4.0
	Heterosexual	0.0	0.0	0.0	0.0	0.0	0.0	0.3	25.0	0.3	100.0	0.7	40.0	3	8.3	5	35.7	8	16.0
	Fem Pres Het Cont	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	5	35.7	5	10.0
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	1	7.1	1	2.0
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	0	0.0	0	0.0
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.3	25.0	0.0	0.0	0.3	20.0	3	8.3	0	0.0	3	6.0

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		1	
1988		2	2
1989			1
1990		1	
1991		1	
1992		4	2
1993		2	
1994			1
1995		6	4
1996		3	2
1997		2	
1999		1	
2000*		3	1
2001	2	5	1
2002	3	5	1
2003	1	2	
2004	1	1	1
2005	1	3	1
2006	1		2
2007		1	
2008		2	
2009		2	
2010		1	1
2011	3	2	2

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		31	100.0	37.8	36	100.0	43.9	67	100.0	81.8	81,945
Gender	Male	27	87.1		29	80.6		56	83.6		
	Female	4	12.9		7	19.4		11	16.4		
Current Age	12 & under	0	0.0		0	0.0		0	0.0		
	13-19	1	3.2		0	0.0		1	1.5		
	20-24	0	0.0		0	0.0		0	0.0		
	25-29	3	9.7		0	0.0		3	4.5		
	30-39	3	9.7		7	19.4		10	14.9		
	40-49	6	19.4		14	38.9		20	29.9		
	50-59	15	48.4		11	30.6		26	38.8		
	60+	3	9.7		4	11.1		7	10.4		
Race/Ethnicity	White	23	74.2		27	75.0		50	74.6		
	Black	4	12.9		2	5.6		6	9.0		
	Hispanic	2	6.5		5	13.9		7	10.4		
	Multi Race	1	3.2		2	5.6		3	4.5		
	Others	1	3.2		0	0.0		1	1.5		
Risk	MSM	19	61.3		18	50.0		37	55.2		
	IDU	0	0.0		6	16.7		6	9.0		
	MSM/IDU	0	0.0		4	11.1		4	6.0		
	Heterosexual	3	9.7		2	5.6		5	7.5		
	Fem Pres Het Cont	1	3.2		3	8.3		4	6.0		
	Blood Prod	0	0.0		0	0.0		0	0.0		
	Pediatric Risk	1	3.2		0	0.0		1	1.5		
	Unk	7	22.6		3	8.3		10	14.9		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	IDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	Male	Fe	male	Т	otal	M	[ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		27	100.0	4	100.0	31	100.0	29	100.0	7	100.0	36	100.0	56	100.0	11	100.0	67	100.0
Risk	MSM	19	70.4	0	0.0	19	61.3	18	62.1	0	0.0	18	50.0	37	66.1	0	0.0	37	55.2
	IDU	0	0.0	0	0.0	0	0.0	4	13.8	2	28.6	6	16.7	4	7.1	2	18.2	6	9.0
	MSM/IDU	0	0.0	0	0.0	0	0.0	4	13.8	0	0.0	4	11.1	4	7.1	0	0.0	4	6.0
	Heterosexual	0	0.0	3	75.0	3	9.7	0	0.0	2	28.6	2	5.6	0	0.0	5	45.5	5	7.5
	Fem Pres Het Cont	0	0.0	1	25.0	1	3.2	0	0.0	3	42.9	3	8.3	0	0.0	4	36.4	4	6.0
	Blood Prod	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Pediatric Risk	1	3.7	0	0.0	1	3.2	0	0.0	0	0.0	0	0.0	1	1.8	0	0.0	1	1.5
	Unk	7	25.9	0	0.0	7	22.6	3	10.3	0	0.0	3	8.3	10	17.9	0	0.0	10	14.9

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		0.7	100.0	0.8	0.7	100.0	0.8	70	100.0	81,945
Gender	Male	0.3	50.0		0.3	50.0		56	80.0	
	Female	0.3	50.0		0.3	50.0		14	20.0	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		1	1.4	
	13-19	0.0	0.0		0.0	0.0		0	0.0	
	20-24	0.0	0.0		0.0	0.0		2	2.9	
	25-29	0.0	0.0		0.0	0.0		8	11.4	
	30-39	0.3	50.0		0.7	100.0		36	51.4	
	40-49	0.0	0.0		0.0	0.0		15	21.4	
	50-59	0.3	50.0		0.0	0.0		6	8.6	
	60+	0.0	0.0		0.0	0.0		2	2.9	
Race/Ethnicity	White	0.7	100.0		0.7	100.0		56	80.0	
	Black	0.0	0.0		0.0	0.0		6	8.6	
	Hispanic	0.0	0.0		0.0	0.0		6	8.6	
	Multi Race	0.0	0.0		0.0	0.0		2	2.9	
	Others	0.0	0.0		0.0	0.0		0	0.0	
Risk	MSM	0.3	50.0		0.3	50.0		32	45.7	
	IDU	0.0	0.0		0.3	50.0		11	15.7	
	MSM/IDU	0.0	0.0		0.0	0.0		6	8.6	
	Heterosexual	0.0	0.0		0.0	0.0		4	5.7	
	Fem Pres Het Cont	0.3	50.0		0.0	0.0		7	10.0	
	Blood Prod	0.0	0.0		0.0	0.0		1	1.4	
	Pediatric Risk	0.0	0.0		0.0	0.0		1	1.4	
	Unk	0.0	0.0		0.0	0.0		8	11.4	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	M	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		0.3	100.0	0.3	100.0	0.7	100.0	0.3	100.0	0.3	100.0	0.7	100.0	56	100.0	14	100.0	70	100.0
Risk	MSM	0.3	100.0	0.0	0.0	0.3	50.0	0.3	100.0	0.0	0.0	0.3	50.0	32	57.1	0	0.0	32	45.7
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	100.0	0.3	50.0	8	14.3	3	21.4	11	15.7
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	6	10.7	0	0.0	6	8.6
	Heterosexual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	1.8	3	21.4	4	5.7
	Fem Pres Het Cont	0.0	0.0	0.3	100.0	0.3	50.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	7	50.0	7	10.0
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	1.8	0	0.0	1	1.4
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	1	7.1	1	1.4
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	8	14.3	0	0.0	8	11.4

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		3	2
1986			1
1987		2	1
1988		1	
1989		2	1
1990		5	
1991		5	3
1992		2	6
1993		8	2
1994		2	6
1995		6	2
1996		3	1
1997		1	1
1998		3	2
1999		3	
2000*	5	1	
2001	3	1	2
2002	3	2	
2003	1	1	
2004	4	4	
2005	2	2	3
2006	2	2	
2007	6	5	1
2008	4	4	
2009		1	
2011	2	1	

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		55	100.0	87.9	56	100.0	89.5	111	100.0	177.5	62,550
Gender	Male	38	69.1		43	76.8		81	73.0		
	Female	17	30.9		13	23.2		30	27.0		
Current Age	12 & under	2	3.6		0	0.0		2	1.8		
	13-19	2	3.6		0	0.0		2	1.8		
	20-24	1	1.8		2	3.6		3	2.7		
	25-29	1	1.8		0	0.0		1	0.9		
	30-39	7	12.7		8	14.3		15	13.5		
	40-49	9	16.4		19	33.9		28	25.2		
	50-59	23	41.8		13	23.2		36	32.4		
	60+	10	18.2		14	25.0		24	21.6		
Race/Ethnicity	White	32	58.2		32	57.1		64	57.7		
	Black	9	16.4		7	12.5		16	14.4		
	Hispanic	7	12.7		15	26.8		22	19.8		
	Multi Race	6	10.9		2	3.6		8	7.2		
	Others	1	1.8		0	0.0		1	0.9		
Risk	MSM	33	60.0		23	41.1		56	50.5		
	IDU	1	1.8		10	17.9		11	9.9		
	MSM/IDU	0	0.0		1	1.8		1	0.9		
	Heterosexual	8	14.5		16	28.6		24	21.6		
	Fem Pres Het Cont	9	16.4		0	0.0		9	8.1		
	Blood Prod	0	0.0		0	0.0		0	0.0		
	Pediatric Risk	3	5.5		0	0.0		3	2.7		
	Unk	1	1.8		6	10.7		7	6.3		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	Male	Fe	male	1	otal	M	Iale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		38	100.0	17	100.0	55	100.0	43	100.0	13	100.0	56	100.0	81	100.0	30	100.0	111	100.0
Risk	MSM	33	86.8	0	0.0	33	60.0	23	53.5	0	0.0	23	41.1	56	69.1	0	0.0	56	50.5
	IDU	1	2.6	0	0.0	1	1.8	7	16.3	3	23.1	10	17.9	8	9.9	3	10.0	11	9.9
	MSM/IDU	0	0.0	0	0.0	0	0.0	1	2.3	0	0.0	1	1.8	1	1.2	0	0.0	1	0.9
	Heterosexual	2	5.3	6	35.3	8	14.5	6	14.0	10	76.9	16	28.6	8	9.9	16	53.3	24	21.6
	Fem Pres Het Cont	0	0.0	9	52.9	9	16.4	0	0.0	0	0.0	0	0.0	0	0.0	9	30.0	9	8.1
	Blood Prod	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Pediatric Risk	1	2.6	2	11.8	3	5.5	0	0.0	0	0.0	0	0.0	1	1.2	2	6.7	3	2.7
	Unk	1	2.6	0	0.0	1	1.8	6	14.0	0	0.0	6	10.7	7	8.6	0	0.0	7	6.3

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		3.7	100.0	5.9	3.0	100.0	4.8	106	100.0	62,550
Gender	Male	1.7	45.5		1.7	55.6		83	78.3	
	Female	2.0	54.5		1.3	44.4		23	21.7	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		1	0.9	
	13-19	0.3	9.1		0.0	0.0		0	0.0	
	20-24	1.0	27.3		0.7	22.2		4	3.8	
	25-29	0.3	9.1		0.0	0.0		4	3.8	
	30-39	0.7	18.2		1.0	33.3		42	39.6	
	40-49	0.3	9.1		0.3	11.1		34	32.1	
	50-59	0.7	18.2		0.7	22.2		13	12.3	
	60+	0.3	9.1		0.3	11.1		8	7.5	
Race/Ethnicity	White	2.3	63.6		1.7	55.6		71	67.0	
	Black	0.3	9.1		0.7	22.2		19	17.9	
	Hispanic	0.7	18.2		0.7	22.2		14	13.2	
	Multi Race	0.3	9.1		0.0	0.0		2	1.9	
	Others	0.0	0.0		0.0	0.0		0	0.0	
Risk	MSM	0.7	18.2		0.3	11.1		44	41.5	
	IDU	0.3	9.1		0.3	11.1		27	25.5	
	MSM/IDU	0.0	0.0		0.0	0.0		4	3.8	
	Heterosexual	2.0	54.5		2.0	66.7		18	17.0	
	Fem Pres Het Cont	0.7	18.2		0.0	0.0		0	0.0	
	Blood Prod	0.0	0.0		0.0	0.0		1	0.9	
	Pediatric Risk	0.0	0.0		0.0	0.0		1	0.9	
	Unk	0.0	0.0		0.3	11.1		11	10.4	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	M	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		1.7	100.0	2.0	100.0	3.7	100.0	1.7	100.0	1.3	100.0	3.0	100.0	83	100.0	23	100.0	106	100.0
Risk	MSM	0.7	40.0	0.0	0.0	0.7	18.2	0.3	20.0	0.0	0.0	0.3	11.1	44	53.0	0	0.0	44	41.5
	IDU	0.3	20.0	0.0	0.0	0.3	9.1	0.3	20.0	0.0	0.0	0.3	11.1	16	19.3	11	47.8	27	25.5
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	4	4.8	0	0.0	4	3.8
	Heterosexual	0.7	40.0	1.3	66.7	2.0	54.5	0.7	40.0	1.3	100.0	2.0	66.7	7	8.4	11	47.8	18	17.0
	Fem Pres Het Cont	0.0	0.0	0.7	33.3	0.7	18.2	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	0	0.0	0	0.0
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	1.2	0	0.0	1	0.9
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	1	4.3	1	0.9
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.3	20.0	0.0	0.0	0.3	11.1	11	13.3	0	0.0	11	10.4

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		2	
1986		3	3
1987		1	1
1988		2	1
1989		4	4
1990		5	2
1991	•	9	7
1992		14	9
1993		5	5
1994		4	2
1995		10	3
1996		2	5
1997		4	2
1998	•	2	1
1999		3	1
2000*	3	4	3
2001	3	2	1
2002	1	1	2
2003	5	3	
2004	4	5	1
2005	3	5	
2006	5	2	1
2007	4	4	
2008	2	1	
2009	4	4	1
2010	2	2	1
2011	5	3	

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		15	100.0	30.4	15	100.0	30.4	30	100.0	60.8	49,363
Gender	Male	12	80.0		12	80.0		24	80.0		
	Female	3	20.0		3	20.0		6	20.0		
Current Age	20-24	1	6.7		1	6.7		2	6.7		
	25-29	1	6.7		0	0.0		1	3.3		
	30-39	5	33.3		3	20.0		8	26.7		
	40-49	2	13.3		5	33.3		7	23.3		
	50-59	5	33.3		3	20.0		8	26.7		
	60+	1	6.7		3	20.0		4	13.3		
Race/Ethnicity	White	11	73.3		11	73.3		22	73.3		
	Black	1	6.7		1	6.7		2	6.7		
	Hispanic	1	6.7		3	20.0		4	13.3		
	Multi Race	2	13.3		0	0.0		2	6.7		
Risk	MSM	9	60.0		7	46.7		16	53.3		
	IDU	1	6.7		1	6.7		2	6.7		
	MSM/IDU	0	0.0		0	0.0		0	0.0		
	Heterosexual	2	13.3		3	20.0		5	16.7		
	Fem Pres Het Cont	2	13.3		0	0.0		2	6.7		
	Blood Prod	0	0.0		1	6.7		1	3.3		
	Unk	1	6.7		3	20.0		4	13.3		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		M	ale	Fe	male	To	otal	N	Male	Fe	male	Т	otal	M	[ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		12	100.0	3	100.0	15	100.0	12	100.0	3	100.0	15	100.0	24	100.0	6	100.0	30	100.0
Risk	MSM	9	75.0	0	0.0	9	60.0	7	58.3	0	0.0	7	46.7	16	66.7	0	0.0	16	53.3
	IDU	1	8.3	0	0.0	1	6.7	1	8.3	0	0.0	1	6.7	2	8.3	0	0.0	2	6.7
	MSM/IDU	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Heterosexual	1	8.3	1	33.3	2	13.3	1	8.3	2	66.7	3	20.0	2	8.3	3	50.0	5	16.7
	Fem Pres Het Cont	0	0.0	2	66.7	2	13.3	0	0.0	0	0.0	0	0.0	0	0.0	2	33.3	2	6.7
	Blood Prod	0	0.0	0	0.0	0	0.0	0	0.0	1	33.3	1	6.7	0	0.0	1	16.7	1	3.3
	Unk	1	8.3	0	0.0	1	6.7	3	25.0	0	0.0	3	20.0	4	16.7	0	0.0	4	13.3

		HIV Diagn	oses (Annua	l Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		1.3	100.0	2.7	0.7	100.0	1.4	30	100.0	49,363
Gender	Male	1.3	100.0		0.7	100.0		25	83.3	
	Female	0.0	0.0		0.0	0.0		5	16.7	
Age at Diagnosis	20-24	0.7	50.0		0.3	50.0		4	13.3	
	25-29	0.3	25.0		0.0	0.0		2	6.7	
	30-39	0.0	0.0		0.0	0.0		12	40.0	
	40-49	0.3	25.0		0.3	50.0		8	26.7	
	50-59	0.0	0.0		0.0	0.0		4	13.3	
	60+	0.0	0.0		0.0	0.0		0	0.0	
Race/Ethnicity	White	1.3	100.0		0.3	50.0		24	80.0	
	Black	0.0	0.0		0.3	50.0		2	6.7	
	Hispanic	0.0	0.0		0.0	0.0		3	10.0	
	Multi Race	0.0	0.0		0.0	0.0		1	3.3	
Risk	MSM	1.3	100.0		0.3	50.0		13	43.3	
	IDU	0.0	0.0		0.3	50.0		4	13.3	
	MSM/IDU	0.0	0.0		0.0	0.0		2	6.7	
	Heterosexual	0.0	0.0		0.0	0.0		4	13.3	
	Fem Pres Het Cont	0.0	0.0		0.0	0.0		0	0.0	
	Blood Prod	0.0	0.0		0.0	0.0		1	3.3	
	Unk	0.0	0.0		0.0	0.0		6	20.0	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		1	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	To	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		1.3	100.0	0.0	0.0	1.3	100.0	0.7	100.0	0.0	0.0	0.7	100.0	25	100.0	5	100.0	30	100.0
Risk	MSM	1.3	100.0	0.0	0.0	1.3	100.0	0.3	50.0	0.0	0.0	0.3	50.0	13	52.0	0	0.0	13	43.3
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.3	50.0	0.0	0.0	0.3	50.0	3	12.0	1	20.0	4	13.3
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	8.0	0	0.0	2	6.7
	Heterosexual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	4.0	3	60.0	4	13.3
	Fem Pres Het Cont	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	0	0.0	0	0.0
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	1	20.0	1	3.3
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	6	24.0	0	0.0	6	20.0

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
1987		2	
1989		1	3
1991		1	
1993		2	2
1994		1	
1995		1	1
1996		1	
1997		2	
1998		3	2
1999			1
2001	1	1	
2002	1	5	1
2003			2
2004		5	1
2005	1		2
2006	1	1	
2007	3	1	
2008	2	1	
2009	1	1	
2010	2	1	
2011	1		

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		18	100.0	37.8	36	100.0	75.7	54	100.0	113.5	47,559
Gender	Male	13	72.2		26	72.2		39	72.2		
	Female	5	27.8		10	27.8		15	27.8		
Current Age	13-19	0	0.0		0	0.0		0	0.0		
	20-24	1	5.6		1	2.8		2	3.7		
	25-29	1	5.6		0	0.0		1	1.9		
	30-39	0	0.0		4	11.1		4	7.4		
	40-49	9	50.0		10	27.8		19	35.2		
	50-59	3	16.7		14	38.9		17	31.5		
	60+	4	22.2		7	19.4		11	20.4		
Race/Ethnicity	White	14	77.8		22	61.1		36	66.7		
	Black	2	11.1		5	13.9		7	13.0		
	Hispanic	2	11.1		4	11.1		6	11.1		
	Multi Race	0	0.0		4	11.1		4	7.4		
	Others	0	0.0		1	2.8		1	1.9		
Risk	MSM	7	38.9		15	41.7		22	40.7		
	IDU	3	16.7		9	25.0		12	22.2		
	MSM/IDU	1	5.6		2	5.6		3	5.6		
	Heterosexual	4	22.2		4	11.1		8	14.8		
	Fem Pres Het Cont	0	0.0		2	5.6		2	3.7		
	Blood Prod	0	0.0		1	2.8		1	1.9		
	Unk	3	16.7		3	8.3		6	11.1		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	Т	otal	N	/Iale	Fe	male	Т	otal	М	ale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		13	100.0	5	100.0	18	100.0	26	100.0	10	100.0	36	100.0	39	100.0	15	100.0	54	100.0
Risk	MSM	7	53.8	0	0.0	7	38.9	15	57.7	0	0.0	15	41.7	22	56.4	0	0.0	22	40.7
	IDU	1	7.7	2	40.0	3	16.7	5	19.2	4	40.0	9	25.0	6	15.4	6	40.0	12	22.2
	MSM/IDU	1	7.7	0	0.0	1	5.6	2	7.7	0	0.0	2	5.6	3	7.7	0	0.0	3	5.6
	Heterosexual	1	7.7	3	60.0	4	22.2	0	0.0	4	40.0	4	11.1	1	2.6	7	46.7	8	14.8
	Fem Pres Het Cont	0	0.0	0	0.0	0	0.0	0	0.0	2	20.0	2	5.6	0	0.0	2	13.3	2	3.7
	Blood Prod	0	0.0	0	0.0	0	0.0	1	3.8	0	0.0	1	2.8	1	2.6	0	0.0	1	1.9
	Unk	3	23.1	0	0.0	3	16.7	3	11.5	0	0.0	3	8.3	6	15.4	0	0.0	6	11.1

		HIV Diagn	oses (Annua	ıl Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		1.7	100.0	3.5	2.3	100.0	4.9	59	100.0	47,559
Gender	Male	1.0	60.0		2.0	85.7		47	79.7	
	Female	0.7	40.0		0.3	14.3		12	20.3	
Age at Diagnosis	13-19	0.0	0.0		0.0	0.0		1	1.7	
	20-24	0.3	20.0		0.0	0.0		0	0.0	
	25-29	0.0	0.0		0.0	0.0		2	3.4	
	30-39	0.3	20.0		0.7	28.6		19	32.2	
	40-49	0.3	20.0		0.7	28.6		29	49.2	
	50-59	0.0	0.0		1.0	42.9		7	11.9	
	60+	0.7	40.0		0.0	0.0		1	1.7	
Race/Ethnicity	White	1.3	80.0		1.7	71.4		42	71.2	
	Black	0.0	0.0		0.0	0.0		7	11.9	
	Hispanic	0.3	20.0		0.3	14.3		4	6.8	
	Multi Race	0.0	0.0		0.0	0.0		5	8.5	
	Others	0.0	0.0		0.3	14.3		1	1.7	
Risk	MSM	0.7	40.0		1.3	57.1		26	44.1	
	IDU	0.3	20.0		0.3	14.3		14	23.7	
	MSM/IDU	0.0	0.0		0.0	0.0		2	3.4	
	Heterosexual	0.3	20.0		0.3	14.3		6	10.2	
	Fem Pres Het Cont	0.3	20.0		0.0	0.0		3	5.1	
	Blood Prod	0.0	0.0		0.0	0.0		1	1.7	
	Unk	0.0	0.0		0.3	14.3		7	11.9	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		1	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fei	nale	To	otal	М	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		1.0	100.0	0.7	100.0	1.7	100.0	2.0	100.0	0.3	100.0	2.3	100.0	47	100.0	12	100.0	59	100.0
Risk	MSM	0.7	66.7	0.0	0.0	0.7	40.0	1.3	66.7	0.0	0.0	1.3	57.1	26	55.3	0	0.0	26	44.1
	IDU	0.0	0.0	0.3	50.0	0.3	20.0	0.0	0.0	0.3	100.0	0.3	14.3	10	21.3	4	33.3	14	23.7
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	4.3	0	0.0	2	3.4
	Heterosexual	0.3	33.3	0.0	0.0	0.3	20.0	0.3	16.7	0.0	0.0	0.3	14.3	1	2.1	5	41.7	6	10.2
	Fem Pres Het Cont	0.0	0.0	0.3	50.0	0.3	20.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	3	25.0	3	5.1
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	2.1	0	0.0	1	1.7
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.3	16.7	0.0	0.0	0.3	14.3	7	14.9	0	0.0	7	11.9

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
1988		1	
1989		2	1
1990		4	2
1991		1	1
1992		3	2
1993		2	3
1994		3	1
1995		6	1
1996		4	3
1997		2	
1998			1
1999		1	1
2000*	1		
2001	4	4	2
2002	7	4	1
2003	2	4	1
2004	1	2	
2005		3	2
2006	2	1	
2007	2	2	
2008	4	3	
2010	4	3	1
2011	1	4	1

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	AIDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		246	100.0	82.6	371	100.0	124.5	617	100.0	207.0	297,999
Gender	Male	149	60.6		251	67.7		400	64.8		
	Female	97	39.4		120	32.3		217	35.2		
Current Age	12 & under	1	0.4		0	0.0		1	0.2		
	13-19	7	2.8		2	0.5		9	1.5		
	20-24	13	5.3		10	2.7		23	3.7		
	25-29	15	6.1		8	2.2		23	3.7		
	30-39	33	13.4		34	9.2		67	10.9		
	40-49	72	29.3		106	28.6		178	28.8		
	50-59	82	33.3		150	40.4		232	37.6		
	60+	23	9.3		61	16.4		84	13.6		
Race/Ethnicity	White	99	40.2		117	31.5		216	35.0		
	Black	83	33.7		140	37.7		223	36.1		
	Hispanic	37	15.0		83	22.4		120	19.4		
	Multi Race	22	8.9		29	7.8		51	178 28.8 . 232 37.6 . 84 13.6 . 216 35.0 . 223 36.1 . 120 19.4 .		
	Others	4	1.6		2	0.5		6	1.0		
	Unk	1	0.4		0	0.0		1	0.2		
Risk	MSM	76	30.9		92	24.8		168	27.2		
	IDU	42	17.1		101	27.2		143	23.2		
	MSM/IDU	8	3.3		16	4.3		24	3.9		
	Heterosexual	82	33.3		98	26.4		180	29.2		
	Fem Pres Het Cont	10	4.1		18	4.9		28	4.5		
	Blood Prod	0	0.0		3	0.8		3	0.5		
	Pediatric Risk	12	4.9		6	1.6		18	2.9		
	Unk	16	6.5		37	10.0		53	8.6		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		M	ale	Fe	male	To	otal	N	Male	Fe	emale	1	otal	M	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		149	100.0	97	100.0	246	100.0	251	100.0	120	100.0	371	100.0	400	100.0	217	100.0	617	100.0
Risk	MSM	76	51.0	0	0.0	76	30.9	92	36.7	0	0.0	92	24.8	168	42.0	0	0.0	168	27.2
	IDU	22	14.8	20	20.6	42	17.1	73	29.1	28	23.3	101	27.2	95	23.8	48	22.1	143	23.2
	MSM/IDU	8	5.4	0	0.0	8	3.3	16	6.4	0	0.0	16	4.3	24	6.0	0	0.0	24	3.9
	Heterosexual	21	14.1	61	62.9	82	33.3	30	12.0	68	56.7	98	26.4	51	12.8	129	59.4	180	29.2
	Fem Pres Het Cont	0	0.0	10	10.3	10	4.1	0	0.0	18	15.0	18	4.9	0	0.0	28	12.9	28	4.5
	Blood Prod	0	0.0	0	0.0	0	0.0	1	0.4	2	1.7	3	0.8	1	0.3	2	0.9	3	0.5
	Pediatric Risk	6	4.0	6	6.2	12	4.9	3	1.2	3	2.5	6	1.6	9	2.3	9	4.1	18	2.9
	Unk	16	10.7	0	0.0	16	6.5	36	14.3	1	0.8	37	10.0	52	13.0	1	0.5	53	8.6

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate 3	Number	Col %	Case Rate 3	Number	Col %	Population
Total		17.3	100.0	5.8	13.7	100.0	4.6	999	100.0	297,999
Gender	Male	12.0	69.2		9.0	65.9		751	75.2	
	Female	5.3	30.8		4.7	34.1		248	24.8	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		8	0.8	
	13-19	1.0	5.8		0.3	2.4		11	1.1	
	20-24	3.0	17.3		1.7	12.2		24	2.4	
	25-29	2.3	13.5		1.3	9.8		84	8.4	
	30-39	3.0	17.3		2.7	19.5		416	41.6	
	40-49	4.0	23.1		4.3	31.7		312	31.2	
	50-59	3.0	17.3		3.0	22.0		117	11.7	
	60+	1.0	5.8		0.3	2.4		27	2.7	
Race/Ethnicity	White	6.3	36.5		4.3	31.7		358	35.8	
	Black	4.7	26.9		5.3	39.0		376	37.6	
	Hispanic	4.0	23.1		2.7	19.5		217	21.7	
	Multi Race	1.7	9.6		1.0	7.3		41	4.1	
	Others	0.7	3.8		0.3	2.4		7	0.7	
	Unk	0.0	0.0		0.0	0.0		0	0.0	
Risk	MSM	7.3	42.3		4.0	29.3		242	24.2	
	IDU	2.7	15.4		2.0	14.6		448	44.8	
	MSM/IDU	0.7	3.8		1.0	7.3		38	3.8	
	Heterosexual	5.0	28.8		4.0	29.3		161	16.1	
	Fem Pres Het Cont	0.7	3.8		1.0	7.3		30	3.0	
	Blood Prod	0.0	0.0		0.0	0.0		11	1.1	
	Pediatric Risk	0.0	0.0		0.3	2.4		13	1.3	
	Unk	1.0	5.8		1.3	9.8		56	5.6	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Diag	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		12.0	100.0	5.3	100.0	17.3	100.0	9.0	100.0	4.7	100.0	13.7	100.0	751	100.0	248	100.0	999	100.0
Risk	MSM	7.3	61.1	0.0	0.0	7.3	42.3	4.0	44.4	0.0	0.0	4.0	29.3	242	32.2	0	0.0	242	24.2
	IDU	1.7	13.9	1.0	18.8	2.7	15.4	1.0	11.1	1.0	21.4	2.0	14.6	349	46.5	99	39.9	448	44.8
	MSM/IDU	0.7	5.6	0.0	0.0	0.7	3.8	1.0	11.1	0.0	0.0	1.0	7.3	38	5.1	0	0.0	38	3.8
	Heterosexual	1.3	11.1	3.7	68.8	5.0	28.8	1.7	18.5	2.3	50.0	4.0	29.3	54	7.2	107	43.1	161	16.1
	Fem Pres Het Cont	0.0	0.0	0.7	12.5	0.7	3.8	0.0	0.0	1.0	21.4	1.0	7.3	0	0.0	30	12.1	30	3.0
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	6	0.8	5	2.0	11	1.1
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	7.1	0.3	2.4	7	0.9	6	2.4	13	1.3
	Unk	1.0	8.3	0.0	0.0	1.0	5.8	1.3	14.8	0.0	0.0	1.3	9.8	55	7.3	1	0.4	56	5.6

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		13	6
1986		15	8
1987		29	16
1988	•	32	24
1989		41	33
1990		47	18
1991		63	35
1992		67	33
1993		105	50
1994		95	59
1995		63	68
1996		67	47
1997		31	23
1998		26	16
1999		36	28
2000*	33	41	15
2001	33	29	17
2002	17	26	23
2003	18	22	16
2004	31	36	18
2005	34	23	10
2006	26	21	16
2007	17	14	10
2008	13	16	8
2009	18	17	11
2010	14	11	13
2011	20	13	8

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	1 AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		858	100.0	93.5	1,013	100.0	110.3	1,871	100.0	203.8	918,028
Gender	Male	555	64.7		701	69.2		1,256	67.1		
	Female	303	35.3		312	30.8		615	32.9		
Current Age	12 & under	8	0.9		2	0.2		10	0.5		
	13-19	11	1.3		4	0.4		15	0.8		
	20-24	47	5.5		19	1.9		66	3.5		
	25-29	85	9.9		26	2.6		111	5.9		
	30-39	193	22.5		123	12.1		316	16.9		
	40-49	262	30.5		376	37.1		638	34.1		
	50-59	190	22.1		331	32.7		521	27.8		
	60+	62	7.2		132	13.0		194	10.4		
Race/Ethnicity	White	290	33.8		330	32.6		620	33.1		
	Black	353	41.1		442	43.6		795	42.5		
	Hispanic	117	13.6		146	14.4		263	14.1		
	Multi Race	80	9.3		81	8.0		161	8.6		
	Others	18	2.1		14	1.4		32	1.7		
Risk	MSM	370	43.1		428	42.3		798	42.7		
	IDU	106	12.4		165	16.3		271	14.5		
	MSM/IDU	32	3.7		59	5.8		91	4.9		
	Heterosexual	255	29.7		299	29.5		554	29.6		
	Fem Pres Het Cont	41	4.8		24	2.4		65	3.5		
	Blood Prod	2	0.2		5	0.5		7	0.4		
	Pediatric Risk	15	1.7		12	1.2		27	1.4		
	Unk	37	4.3		21	2.1		58	3.1		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Ca	ses			Living	HIV a	nd AIDS	Cases	
		M	ale	Fe	male	Т	otal	I	Male	Fe	male	Т	otal	M	[ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		555	100.0	303	100.0	858	100.0	701	100.0	312	100.0	1,013	100.0	1,256	100.0	615	100.0	1,871	100.0
Risk	MSM	370	66.7	0	0.0	370	43.1	428	61.1	0	0.0	428	42.3	798	63.5	0	0.0	798	42.7
	IDU	53	9.5	53	17.5	106	12.4	101	14.4	64	20.5	165	16.3	154	12.3	117	19.0	271	14.5
	MSM/IDU	32	5.8	0	0.0	32	3.7	59	8.4	0	0.0	59	5.8	91	7.2	0	0.0	91	4.9
	Heterosexual	54	9.7	201	66.3	255	29.7	82	11.7	217	69.6	299	29.5	136	10.8	418	68.0	554	29.6
	Fem Pres Het Cont	0	0.0	41	13.5	41	4.8	0	0.0	24	7.7	24	2.4	0	0.0	65	10.6	65	3.5
	Blood Prod	2	0.4	0	0.0	2	0.2	5	0.7	0	0.0	5	0.5	7	0.6	0	0.0	7	0.4
	Pediatric Risk	7	1.3	8	2.6	15	1.7	6	0.9	6	1.9	12	1.2	13	1.0	14	2.3	27	1.4
	Unk	37	6.7	0	0.0	37	4.3	20	2.9	1	0.3	21	2.1	57	4.5	1	0.2	58	3.1

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		98.3	100.0	10.7	62.3	100.0	6.8	2,321	100.0	918,028
Gender	Male	65.0	66.1		39.0	62.6		1,747	75.3	
	Female	33.3	33.9		23.3	37.4		574	24.7	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		17	0.7	
	13-19	4.0	4.1		0.7	1.1		23	1.0	
	20-24	14.7	14.9		3.7	5.9		95	4.1	
	25-29	11.7	11.9		5.3	8.6		237	10.2	
	30-39	24.0	24.4		16.7	26.7		846	36.4	
	40-49	27.3	27.8		22.0	35.3		748	32.2	
	50-59	13.7	13.9		10.7	17.1		274	11.8	
	60+	3.0	3.1		3.3	5.3		81	3.5	
Race/Ethnicity	White	32.3	32.9		12.7	20.3		850	36.6	
	Black	44.0	44.7		32.7	52.4		1,068	46.0	
	Hispanic	12.3	12.5		9.7	15.5		280	12.1	
	Multi Race	5.3	5.4		5.3	8.6		106	4.6	
	Others	4.3	4.4		2.0	3.2		17	0.7	
Risk	MSM	43.0	43.7		22.7	36.4		1,033	44.5	
	IDU	7.7	7.8		9.0	14.4		563	24.3	
	MSM/IDU	2.7	2.7		2.7	4.3		114	4.9	
	Heterosexual	32.3	32.9		21.7	34.8		474	20.4	
	Fem Pres Het Cont	5.3	5.4		3.3	5.3		36	1.6	
	Blood Prod	0.0	0.0		0.0	0.0		23	1.0	
	Pediatric Risk	0.3	0.3		0.7	1.1		27	1.2	
	Unk	7.0	7.1		2.3	3.7		51	2.2	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	То	otal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		65.0	100.0	33.3	100.0	98.3	100.0	39.0	100.0	23.3	100.0	62.3	100.0	1,747	100.0	574	100.0	2,321	100.0
Risk	MSM	43.0	66.2	0.0	0.0	43.0	43.7	22.7	58.1	0.0	0.0	22.7	36.4	1,033	59.1	0	0.0	1,033	44.5
	IDU	4.0	6.2	3.7	11.0	7.7	7.8	4.7	12.0	4.3	18.6	9.0	14.4	398	22.8	165	28.7	563	24.3
	MSM/IDU	2.7	4.1	0.0	0.0	2.7	2.7	2.7	6.8	0.0	0.0	2.7	4.3	114	6.5	0	0.0	114	4.9
	Heterosexual	8.0	12.3	24.3	73.0	32.3	32.9	6.3	16.2	15.3	65.7	21.7	34.8	117	6.7	357	62.2	474	20.4
	Fem Pres Het Cont	0.0	0.0	5.3	16.0	5.3	5.4	0.0	0.0	3.3	14.3	3.3	5.3	0	0.0	36	6.3	36	1.6
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	20	1.1	3	0.5	23	1.0
	Pediatric Risk	0.3	0.5	0.0	0.0	0.3	0.3	0.3	0.9	0.3	1.4	0.7	1.1	17	1.0	10	1.7	27	1.2
	Unk	7.0	10.8	0.0	0.0	7.0	7.1	2.3	6.0	0.0	0.0	2.3	3.7	48	2.7	3	0.5	51	2.2

¹Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		20	16
1986		23	12
1987		36	19
1988		55	19
1989		82	49
1990		78	62
1991		103	70
1992		122	80
1993		174	71
1994		161	114
1995		161	154
1996		144	81
1997		78	49
1998		86	40
1999		102	41
2000*	69	127	46
2001	90	93	51
2002	74	66	30
2003	79	88	50
2004	86	69	39
2005	82	68	37
2006	95	59	36
2007	99	84	37
2008	82	55	29
2009	104	61	34
2010	93	54	30
2011	98	72	24

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		11	100.0	28.1	14	100.0	35.7	25	100.0	63.8	39,181
Gender	Male	7	63.6		11	78.6		18	72.0		
	Female	4	36.4		3	21.4		7	28.0		
Current Age	20-24	1	9.1		0	0.0		1	4.0		
	25-29	0	0.0		0	0.0		0	0.0		
	30-39	1	9.1		0	0.0		1	4.0		
	40-49	0	0.0		8	57.1		8	32.0		
	50-59	5	45.5		5	35.7		10	40.0		
	60+	4	36.4		1	7.1		5	20.0		
Race/Ethnicity	White	6	54.5		6	42.9		12	48.0		
	Black	1	9.1		3	21.4		4	16.0		
	Hispanic	4	36.4		3	21.4		7	28.0		
	Multi Race	0	0.0		2	14.3		2	8.0		
	Others	0	0.0		0	0.0		0	0.0		
Risk	MSM	2	18.2		3	21.4		5	20.0		
	IDU	4	36.4		4	28.6		8	32.0		
	MSM/IDU	0	0.0		0	0.0		0	0.0		
	Heterosexual	4	36.4		3	21.4		7	28.0		
	Fem Pres Het Cont	1	9.1		1	7.1		2	8.0		
	Unk	0	0.0		3	21.4		3	12.0		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	/Iale	Fe	male	Т	otal	М	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		7	100.0	4	100.0	11	100.0	11	100.0	3	100.0	14	100.0	18	100.0	7	100.0	25	100.0
Risk	MSM	2	28.6	0	0.0	2	18.2	3	27.3	0	0.0	3	21.4	5	27.8	0	0.0	5	20.0
	IDU	4	57.1	0	0.0	4	36.4	3	27.3	1	33.3	4	28.6	7	38.9	1	14.3	8	32.0
	MSM/IDU	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Heterosexual	1	14.3	3	75.0	4	36.4	2	18.2	1	33.3	3	21.4	3	16.7	4	57.1	7	28.0
	Fem Pres Het Cont	0	0.0	1	25.0	1	9.1	0	0.0	1	33.3	1	7.1	0	0.0	2	28.6	2	8.0
	Unk	0	0.0	0	0.0	0	0.0	3	27.3	0	0.0	3	21.4	3	16.7	0	0.0	3	12.0

		HIV Diagn	oses (Annua	l Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		1.0	100.0	2.6	0.3	100.0	0.9	37	100.0	39,181
Gender	Male	0.7	66.7		0.3	100.0		33	89.2	
	Female	0.3	33.3		0.0	0.0		4	10.8	
Age at Diagnosis	20-24	0.7	66.7		0.3	100.0		4	10.8	
	25-29	0.0	0.0		0.0	0.0		6	16.2	
	30-39	0.0	0.0		0.0	0.0		11	29.7	
	40-49	0.0	0.0		0.0	0.0		10	27.0	
	50-59	0.3	33.3		0.0	0.0		5	13.5	
	60+	0.0	0.0		0.0	0.0		1	2.7	
Race/Ethnicity	White	0.3	33.3		0.0	0.0		27	73.0	
	Black	0.0	0.0		0.0	0.0		4	10.8	
	Hispanic	0.7	66.7		0.3	100.0		3	8.1	
	Multi Race	0.0	0.0		0.0	0.0		2	5.4	
	Others	0.0	0.0		0.0	0.0		1	2.7	
Risk	MSM	0.7	66.7		0.3	100.0		19	51.4	
	IDU	0.0	0.0		0.0	0.0		7	18.9	
	MSM/IDU	0.0	0.0		0.0	0.0		1	2.7	
	Heterosexual	0.0	0.0		0.0	0.0		5	13.5	
	Fem Pres Het Cont	0.3	33.3		0.0	0.0		1	2.7	
	Unk	0.0	0.0		0.0	0.0		4	10.8	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

]	HIV Diag	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS C	Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		0.7	100.0	0.3	100.0	1.0	100.0	0.3	100.0	0.0	0.0	0.3	100.0	33	100.0	4	100.0	37	100.0
Risk	MSM	0.7	100.0	0.0	0.0	0.7	66.7	0.3	100.0	0.0	0.0	0.3	100.0	19	57.6	0	0.0	19	51.4
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	6	18.2	1	25.0	7	18.9
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	3.0	0	0.0	1	2.7
	Heterosexual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3	9.1	2	50.0	5	13.5
	Fem Pres Het Cont	0.0	0.0	0.3	100.0	0.3	33.3	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	1	25.0	1	2.7
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	4	12.1	0	0.0	4	10.8

¹Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		3	3
1987		1	
1988		4	1
1989		3	2
1990		2	1
1991		1	1
1992		2	4
1993		3	2
1994		2	1
1995		7	1
1996		1	2
1997		1	2
2000*	1	1	
2001	1		
2002	1	1	2
2003	2	1	
2004	1		
2005		1	1
2006		1	1
2008		1	
2009	1		
2010	2	1	

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		11	100.0	21.3	23	100.0	44.6	34	100.0	66.0	51,551
Gender	Male	9	81.8		18	78.3		27	79.4		
	Female	2	18.2		5	21.7		7	20.6		
Current Age	20-24	0	0.0		0	0.0		0	0.0		
	25-29	0	0.0		2	8.7		2	5.9		
	30-39	1	9.1		3	13.0		4	11.8		
	40-49	5	45.5		10	43.5		15	44.1		
	50-59	3	27.3		5	21.7		8	23.5		
	60+	2	18.2		3	13.0		5	14.7		
Race/Ethnicity	White	6	54.5		17	73.9		23	67.6		
	Black	1	9.1		0	0.0		1	2.9		
	Hispanic	2	18.2		5	21.7		7	20.6		
	Multi Race	1	9.1		1	4.3		2	5.9		
	Others	1	9.1		0	0.0		1	2.9		
Risk	MSM	4	36.4		13	56.5		17	50.0		
	IDU	3	27.3		4	17.4		7	20.6		
	MSM/IDU	0	0.0		1	4.3		1	2.9		
	Heterosexual	2	18.2		4	17.4		6	17.6		
	Fem Pres Het Cont	0	0.0		1	4.3		1	2.9		
	Unk	2	18.2		0	0.0		2	5.9		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		M	ale	Fe	male	To	otal	N	Male	Fe	male	Т	otal	M	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		9	100.0	2	100.0	11	100.0	18	100.0	5	100.0	23	100.0	27	100.0	7	100.0	34	100.0
Risk	MSM	4	44.4	0	0.0	4	36.4	13	72.2	0	0.0	13	56.5	17	63.0	0	0.0	17	50.0
	IDU	3	33.3	0	0.0	3	27.3	2	11.1	2	40.0	4	17.4	5	18.5	2	28.6	7	20.6
	MSM/IDU	0	0.0	0	0.0	0	0.0	1	5.6	0	0.0	1	4.3	1	3.7	0	0.0	1	2.9
	Heterosexual	0	0.0	2	100.0	2	18.2	2	11.1	2	40.0	4	17.4	2	7.4	4	57.1	6	17.6
	Fem Pres Het Cont	0	0.0	0	0.0	0	0.0	0	0.0	1	20.0	1	4.3	0	0.0	1	14.3	1	2.9
	Unk	2	22.2	0	0.0	2	18.2	0	0.0	0	0.0	0	0.0	2	7.4	0	0.0	2	5.9

		HIV Diagn	oses (Annua	l Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		2.0	100.0	3.9	2.0	100.0	3.9	42	100.0	51,551
Gender	Male	1.7	83.3		1.7	83.3		32	76.2	
	Female	0.3	16.7		0.3	16.7		10	23.8	
Age at Diagnosis	20-24	0.0	0.0		0.0	0.0		1	2.4	
	25-29	0.7	33.3		0.7	33.3		5	11.9	
	30-39	0.3	16.7		0.7	33.3		18	42.9	
	40-49	0.7	33.3		0.3	16.7		11	26.2	
	50-59	0.3	16.7		0.3	16.7		5	11.9	
	60+	0.0	0.0		0.0	0.0		2	4.8	
Race/Ethnicity	White	1.3	66.7		1.7	83.3		34	81.0	
	Black	0.0	0.0		0.0	0.0		2	4.8	
	Hispanic	0.3	16.7		0.3	16.7		5	11.9	
	Multi Race	0.0	0.0		0.0	0.0		1	2.4	
	Others	0.3	16.7		0.0	0.0		0	0.0	
Risk	MSM	1.7	83.3		1.7	83.3		20	47.6	
	IDU	0.0	0.0		0.0	0.0		11	26.2	
	MSM/IDU	0.0	0.0		0.0	0.0		1	2.4	
	Heterosexual	0.3	16.7		0.3	16.7		5	11.9	
	Fem Pres Het Cont	0.0	0.0		0.0	0.0		2	4.8	
	Unk	0.0	0.0		0.0	0.0		3	7.1	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Diag	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Averag	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	M	ale	Fer	nale	To	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		1.7	100.0	0.3	100.0	2.0	100.0	1.7	100.0	0.3	100.0	2.0	100.0	32	100.0	10	100.0	42	100.0
Risk	MSM	1.7	100.0	0.0	0.0	1.7	83.3	1.7	100.0	0.0	0.0	1.7	83.3	20	62.5	0	0.0	20	47.6
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	6	18.8	5	50.0	11	26.2
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	3.1	0	0.0	1	2.4
	Heterosexual	0.0	0.0	0.3	100.0	0.3	16.7	0.0	0.0	0.3	100.0	0.3	16.7	2	6.3	3	30.0	5	11.9
	Fem Pres Het Cont	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	2	20.0	2	4.8
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3	9.4	0	0.0	3	7.1

¹Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
1986		1	
1987			1
1988		1	1
1989		1	1
1991		4	1
1992		1	1
1993		1	2
1994		4	2
1995		4	2
1996		3	1
1997		3	1
2000*	2	2	
2001	4	4	1
2002		1	2
2003	1		
2004	2	2	1
2006		2	1
2007		1	•
2008		1	
2009	4	2	2
2010	1	1	
2011	1	3	

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		34	100.0	61.6	34	100.0	61.6	68	100.0	123.2	55,180
Gender	Male	19	55.9		26	76.5		45	66.2		
	Female	15	44.1		8	23.5		23	33.8		
Current Age	12 & under	0	0.0		0	0.0		0	0.0		
	13-19	1	2.9		0	0.0		1	1.5		
	20-24	1	2.9		2	5.9		3	4.4		
	25-29	2	5.9		0	0.0		2	2.9		•
	30-39	6	17.6		6	17.6		12	17.6		
	40-49	13	38.2		8	23.5		21	30.9		
	50-59	10	29.4		12	35.3		22	32.4		
	60+	1	2.9		6	17.6		7	10.3		
Race/Ethnicity	White	23	67.6		24	70.6		47	69.1		•
	Black	0	0.0		3	8.8		3	4.4		
	Hispanic	7	20.6		4	11.8		11	16.2		
	Multi Race	4	11.8		1	2.9		5	7.4		
	Others	0	0.0		2	5.9		2	2.9		
Risk	MSM	12	35.3		14	41.2		26	38.2		•
	IDU	4	11.8		2	5.9		6	8.8		
	MSM/IDU	0	0.0		0	0.0		0	0.0		
	Heterosexual	11	32.4		5	14.7		16	23.5		
	Fem Pres Het Cont	5	14.7		3	8.8		8	11.8		
	Blood Prod	0	0.0		0	0.0		0	0.0		
	Pediatric Risk	0	0.0		2	5.9		2	2.9		
	Unk	2	5.9		8	23.5		10	14.7		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	Т	otal	N	Male	Fe	emale	7	otal	M	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		19	100.0	15	100.0	34	100.0	26	100.0	8	100.0	34	100.0	45	100.0	23	100.0	68	100.0
Risk	MSM	12	63.2	0	0.0	12	35.3	14	53.8	0	0.0	14	41.2	26	57.8	0	0.0	26	38.2
	IDU	1	5.3	3	20.0	4	11.8	2	7.7	0	0.0	2	5.9	3	6.7	3	13.0	6	8.8
	MSM/IDU	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Heterosexual	4	21.1	7	46.7	11	32.4	2	7.7	3	37.5	5	14.7	6	13.3	10	43.5	16	23.5
	Fem Pres Het Cont	0	0.0	5	33.3	5	14.7	0	0.0	3	37.5	3	8.8	0	0.0	8	34.8	8	11.8
	Blood Prod	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Pediatric Risk	0	0.0	0	0.0	0	0.0	0	0.0	2	25.0	2	5.9	0	0.0	2	8.7	2	2.9
	Unk	2	10.5	0	0.0	2	5.9	8	30.8	0	0.0	8	23.5	10	22.2	0	0.0	10	14.7

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		2.3	100.0	4.2	1.7	100.0	3.0	57	100.0	55,180
Gender	Male	1.3	57.1		1.7	100.0		44	77.2	
	Female	1.0	42.9		0.0	0.0		13	22.8	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		1	1.8	
	13-19	0.3	14.3		0.0	0.0		1	1.8	
	20-24	0.3	14.3		0.0	0.0		1	1.8	
	25-29	0.3	14.3		0.3	20.0		6	10.5	
	30-39	0.3	14.3		0.7	40.0		20	35.1	
	40-49	0.7	28.6		0.3	20.0		18	31.6	
	50-59	0.3	14.3		0.0	0.0		8	14.0	
	60+	0.0	0.0		0.3	20.0		2	3.5	
Race/Ethnicity	White	2.3	100.0		0.7	40.0		44	77.2	
	Black	0.0	0.0		0.7	40.0		5	8.8	
	Hispanic	0.0	0.0		0.0	0.0		5	8.8	
	Multi Race	0.0	0.0		0.3	20.0		1	1.8	
	Others	0.0	0.0		0.0	0.0		2	3.5	
Risk	MSM	1.0	42.9		1.0	60.0		26	45.6	
	IDU	0.0	0.0		0.0	0.0		5	8.8	
	MSM/IDU	0.0	0.0		0.0	0.0		2	3.5	
	Heterosexual	0.7	28.6		0.3	20.0		9	15.8	
	Fem Pres Het Cont	0.7	28.6		0.0	0.0		3	5.3	
	Blood Prod	0.0	0.0		0.0	0.0		2	3.5	
	Pediatric Risk	0.0	0.0		0.0	0.0		2	3.5	
	Unk	0.0	0.0		0.3	20.0		8	14.0	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Diag	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		1.3	100.0	1.0	100.0	2.3	100.0	1.7	100.0	0.0	0.0	1.7	100.0	44	100.0	13	100.0	57	100.0
Risk	MSM	1.0	75.0	0.0	0.0	1.0	42.9	1.0	60.0	0.0	0.0	1.0	60.0	26	59.1	0	0.0	26	45.6
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	5	11.4	0	0.0	5	8.8
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	4.5	0	0.0	2	3.5
	Heterosexual	0.3	25.0	0.3	33.3	0.7	28.6	0.3	20.0	0.0	0.0	0.3	20.0	2	4.5	7	53.8	9	15.8
	Fem Pres Het Cont	0.0	0.0	0.7	66.7	0.7	28.6	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	3	23.1	3	5.3
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	2.3	1	7.7	2	3.5
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	2	15.4	2	3.5
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.3	20.0	0.0	0.0	0.3	20.0	8	18.2	0	0.0	8	14.0

¹Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

² Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		3	1
1986		1	2
1988	•	3	1
1989			1
1990	٠	2	
1991	٠	4	3
1992		4	2
1993		1	1
1994	٠	2	4
1995		3	1
1996		1	
1997		1	1
1998		4	
1999		3	1
2000*	2	1	
2001	2	1	1
2002	•	1	
2003	3	2	1
2004	4	4	
2005	2	2	1
2006		3	
2007	1	2	1
2008	5	4	1
2009	1	1	
2010	3	1	
2011	3	3	

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	1 AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		22	100.0	36.7	29	100.0	48.3	51	100.0	85.0	59,993
Gender	Male	14	63.6		27	93.1		41	80.4		
	Female	8	36.4		2	6.9		10	19.6		
Current Age	20-24	2	9.1		0	0.0		2	3.9		
	25-29	0	0.0		1	3.4		1	2.0		
	30-39	6	27.3		5	17.2		11	21.6		
	40-49	10	45.5		13	44.8		23	45.1		
	50-59	3	13.6		8	27.6		11	21.6		
	60+	1	4.5		2	6.9		3	5.9		
Race/Ethnicity	White	12	54.5		15	51.7		27	52.9		
	Black	4	18.2		4	13.8		8	15.7		
	Hispanic	3	13.6		8	27.6		11	21.6		
	Multi Race	3	13.6		2	6.9		5	9.8		
	Others	0	0.0		0	0.0		0	0.0		
Risk	MSM	9	40.9		12	41.4		21	41.2		
	IDU	4	18.2		6	20.7		10	19.6		
	MSM/IDU	1	4.5		2	6.9		3	5.9		
	Heterosexual	6	27.3		4	13.8		10	19.6		
	Fem Pres Het Cont	2	9.1		0	0.0		2	3.9		
	Blood Prod	0	0.0		0	0.0		0	0.0		
	Unk	0	0.0		5	17.2		5	9.8		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	/Iale	Fe	male	Т	otal	M	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		14	100.0	8	100.0	22	100.0	27	100.0	2	100.0	29	100.0	41	100.0	10	100.0	51	100.0
Risk	MSM	9	64.3	0	0.0	9	40.9	12	44.4	0	0.0	12	41.4	21	51.2	0	0.0	21	41.2
	IDU	3	21.4	1	12.5	4	18.2	5	18.5	1	50.0	6	20.7	8	19.5	2	20.0	10	19.6
	MSM/IDU	1	7.1	0	0.0	1	4.5	2	7.4	0	0.0	2	6.9	3	7.3	0	0.0	3	5.9
	Heterosexual	1	7.1	5	62.5	6	27.3	3	11.1	1	50.0	4	13.8	4	9.8	6	60.0	10	19.6
	Fem Pres Het Cont	0	0.0	2	25.0	2	9.1	0	0.0	0	0.0	0	0.0	0	0.0	2	20.0	2	3.9
	Blood Prod	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Unk	0	0.0	0	0.0	0	0.0	5	18.5	0	0.0	5	17.2	5	12.2	0	0.0	5	9.8

		HIV Diagn	oses (Annua	ıl Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		2.3	100.0	3.9	2.0	100.0	3.3	57	100.0	59,993
Gender	Male	1.7	71.4		1.7	83.3		51	89.5	
	Female	0.7	28.6		0.3	16.7		6	10.5	
Age at Diagnosis	20-24	0.3	14.3		0.3	16.7		3	5.3	
	25-29	0.0	0.0		0.0	0.0		6	10.5	
	30-39	1.0	42.9		0.7	33.3		21	36.8	
	40-49	0.3	14.3		1.0	50.0		18	31.6	
	50-59	0.7	28.6		0.0	0.0		7	12.3	
	60+	0.0	0.0		0.0	0.0		2	3.5	
Race/Ethnicity	White	1.3	57.1		1.3	66.7		38	66.7	
	Black	0.0	0.0		0.0	0.0		6	10.5	
	Hispanic	1.0	42.9		0.7	33.3		9	15.8	
	Multi Race	0.0	0.0		0.0	0.0		3	5.3	
	Others	0.0	0.0		0.0	0.0		1	1.8	
Risk	MSM	0.7	28.6		0.7	33.3		30	52.6	
	IDU	0.3	14.3		0.0	0.0		8	14.0	
	MSM/IDU	0.0	0.0		0.0	0.0		2	3.5	
	Heterosexual	0.3	14.3		0.3	16.7		7	12.3	
	Fem Pres Het Cont	0.0	0.0		0.0	0.0		2	3.5	
	Blood Prod	0.0	0.0		0.0	0.0		1	1.8	
	Unk	1.0	42.9		1.0	50.0		7	12.3	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		1	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fei	nale	To	otal	М	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		1.7	100.0	0.7	100.0	2.3	100.0	1.7	100.0	0.3	100.0	2.0	100.0	51	100.0	6	100.0	57	100.0
Risk	MSM	0.7	40.0	0.0	0.0	0.7	28.6	0.7	40.0	0.0	0.0	0.7	33.3	30	58.8	0	0.0	30	52.6
	IDU	0.0	0.0	0.3	50.0	0.3	14.3	0.0	0.0	0.0	0.0	0.0	0.0	7	13.7	1	16.7	8	14.0
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	3.9	0	0.0	2	3.5
	Heterosexual	0.0	0.0	0.3	50.0	0.3	14.3	0.0	0.0	0.3	100.0	0.3	16.7	4	7.8	3	50.0	7	12.3
	Fem Pres Het Cont	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	2	33.3	2	3.5
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	2.0	0	0.0	1	1.8
	Unk	1.0	60.0	0.0	0.0	1.0	42.9	1.0	60.0	0.0	0.0	1.0	50.0	7	13.7	0	0.0	7	12.3

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
1986		1	
1987		1	
1988		1	1
1989		1	3
1990		4	1
1991		3	3
1992		4	3
1993		2	2
1994		4	2
1995		5	1
1996		1	2
1997		1	
1998		3	1
1999		5	1
2000*	1	2	
2001	5	2	
2002	4	3	2
2003	3	1	
2005	1	1	2
2006	2		1
2007	5	3	
2008	1	3	
2009	1	2	1
2010	3	1	1
2011	3	3	1

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		30	100.0	61.3	39	100.0	79.7	69	100.0	140.9	48,954
Gender	Male	20	66.7		29	74.4		49	71.0		
	Female	10	33.3		10	25.6		20	29.0		
Current Age	13-19	0	0.0		0	0.0		0	0.0		
	20-24	0	0.0		0	0.0		0	0.0		
	25-29	0	0.0		0	0.0		0	0.0		
	30-39	6	20.0		4	10.3		10	14.5		
	40-49	8	26.7		11	28.2		19	27.5		
	50-59	12	40.0		19	48.7		31	44.9		
	60+	4	13.3		5	12.8		9	13.0		
Race/Ethnicity	White	23	76.7		30	76.9		53	76.8		
	Black	3	10.0		1	2.6		4	5.8		
	Hispanic	4	13.3		3	7.7		7	10.1		
	Multi Race	0	0.0		5	12.8		5	7.2		
Risk	MSM	15	50.0		18	46.2		33	47.8		
	IDU	1	3.3		5	12.8		6	8.7		
	MSM/IDU	0	0.0		1	2.6		1	1.4		
	Heterosexual	8	26.7		9	23.1		17	24.6		
	Fem Pres Het Cont	4	13.3		3	7.7		7	10.1		
	Blood Prod	0	0.0		0	0.0		0	0.0		
	Pediatric Risk	0	0.0		1	2.6		1	1.4		
	Unk	2	6.7		2	5.1		4	5.8		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	Male	Fe	male	Т	otal	M	[ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		20	100.0	10	100.0	30	100.0	29	100.0	10	100.0	39	100.0	49	100.0	20	100.0	69	100.0
Risk	MSM	15	75.0	0	0.0	15	50.0	18	62.1	0	0.0	18	46.2	33	67.3	0	0.0	33	47.8
	IDU	1	5.0	0	0.0	1	3.3	5	17.2	0	0.0	5	12.8	6	12.2	0	0.0	6	8.7
	MSM/IDU	0	0.0	0	0.0	0	0.0	1	3.4	0	0.0	1	2.6	1	2.0	0	0.0	1	1.4
	Heterosexual	2	10.0	6	60.0	8	26.7	2	6.9	7	70.0	9	23.1	4	8.2	13	65.0	17	24.6
	Fem Pres Het Cont	0	0.0	4	40.0	4	13.3	0	0.0	3	30.0	3	7.7	0	0.0	7	35.0	7	10.1
	Blood Prod	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Pediatric Risk	0	0.0	0	0.0	0	0.0	1	3.4	0	0.0	1	2.6	1	2.0	0	0.0	1	1.4
	Unk	2	10.0	0	0.0	2	6.7	2	6.9	0	0.0	2	5.1	4	8.2	0	0.0	4	5.8

		HIV Diagn	oses (Annua	ıl Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		1.3	100.0	2.7	1.7	100.0	3.4	81	100.0	48,954
Gender	Male	1.0	75.0		0.7	40.0		66	81.5	
	Female	0.3	25.0		1.0	60.0		15	18.5	
Age at Diagnosis	13-19	0.0	0.0		0.0	0.0		3	3.7	
	20-24	0.0	0.0		0.0	0.0		1	1.2	
	25-29	0.0	0.0		0.0	0.0		3	3.7	
	30-39	0.0	0.0		0.3	20.0		27	33.3	
	40-49	0.3	25.0		1.0	60.0		36	44.4	
	50-59	0.7	50.0		0.3	20.0		8	9.9	
	60+	0.3	25.0		0.0	0.0		3	3.7	
Race/Ethnicity	White	1.3	100.0		1.7	100.0		57	70.4	
	Black	0.0	0.0		0.0	0.0		4	4.9	
	Hispanic	0.0	0.0		0.0	0.0		15	18.5	
	Multi Race	0.0	0.0		0.0	0.0		5	6.2	
Risk	MSM	0.3	25.0		0.3	20.0		33	40.7	
	IDU	0.0	0.0		0.0	0.0		18	22.2	
	MSM/IDU	0.0	0.0		0.0	0.0		2	2.5	
	Heterosexual	0.7	50.0		0.7	40.0		15	18.5	
	Fem Pres Het Cont	0.0	0.0		0.3	20.0		4	4.9	
	Blood Prod	0.0	0.0		0.0	0.0		1	1.2	
	Pediatric Risk	0.0	0.0		0.0	0.0		1	1.2	
	Unk	0.3	25.0		0.3	20.0		7	8.6	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Diag	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		1.0	100.0	0.3	100.0	1.3	100.0	0.7	100.0	1.0	100.0	1.7	100.0	66	100.0	15	100.0	81	100.0
Risk	MSM	0.3	33.3	0.0	0.0	0.3	25.0	0.3	50.0	0.0	0.0	0.3	20.0	33	50.0	0	0.0	33	40.7
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	16	24.2	2	13.3	18	22.2
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	3.0	0	0.0	2	2.5
	Heterosexual	0.3	33.3	0.3	100.0	0.7	50.0	0.0	0.0	0.7	66.7	0.7	40.0	6	9.1	9	60.0	15	18.5
	Fem Pres Het Cont	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	33.3	0.3	20.0	0	0.0	4	26.7	4	4.9
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	1.5	0	0.0	1	1.2
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	1.5	0	0.0	1	1.2
	Unk	0.3	33.3	0.0	0.0	0.3	25.0	0.3	50.0	0.0	0.0	0.3	20.0	7	10.6	0	0.0	7	8.6

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

² Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		1	1
1987		2	1
1988		3	2
1989		3	3
1990	٠	2	
1991		5	1
1992		5	4
1993		4	1
1994	٠	4	5
1995		7	2
1996		3	2
1997		3	1
1998		2	1
1999		5	
2000*		2	5
2001	6	3	1
2002	2	3	1
2003	1	7	
2004	4	3	2
2005	1	1	2
2006	2	2	1
2007	5	2	
2008	3	4	2
2009	4	1	2
2010		3	1
2011		1	1

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	d AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		21	100.0	32.7	29	100.0	45.2	50	100.0	77.9	64,160
Gender	Male	17	81.0		24	82.8		41	82.0		
	Female	4	19.0		5	17.2		9	18.0		
Current Age	13-19	0	0.0		0	0.0		0	0.0		
	20-24	0	0.0		2	6.9		2	4.0		
	25-29	3	14.3		1	3.4		4	8.0		
	30-39	2	9.5		0	0.0		2	4.0		
	40-49	9	42.9		12	41.4		21	42.0		
	50-59	4	19.0		8	27.6		12	24.0		
	60+	3	14.3		6	20.7		9	18.0		
Race/Ethnicity	White	17	81.0		17	58.6		34	68.0		
	Black	1	4.8		3	10.3		4	8.0		
	Hispanic	3	14.3		1	3.4		4	8.0		
	Multi Race	0	0.0		6	20.7		6	12.0		
	Others	0	0.0		2	6.9		2	4.0		
Risk	MSM	14	66.7		14	48.3		28	56.0		
	IDU	1	4.8		3	10.3		4	8.0		
	MSM/IDU	0	0.0		1	3.4		1	2.0		
	Heterosexual	3	14.3		5	17.2		8	16.0		
	Fem Pres Het Cont	1	4.8		3	10.3		4	8.0		
	Pediatric Risk	0	0.0		1	3.4		1	2.0		
	Unk	2	9.5		2	6.9		4	8.0		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	/Iale	Fe	male	Т	otal	M	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		17	100.0	4	100.0	21	100.0	24	100.0	5	100.0	29	100.0	41	100.0	9	100.0	50	100.0
Risk	MSM	14	82.4	0	0.0	14	66.7	14	58.3	0	0.0	14	48.3	28	68.3	0	0.0	28	56.0
	IDU	1	5.9	0	0.0	1	4.8	3	12.5	0	0.0	3	10.3	4	9.8	0	0.0	4	8.0
	MSM/IDU	0	0.0	0	0.0	0	0.0	1	4.2	0	0.0	1	3.4	1	2.4	0	0.0	1	2.0
	Heterosexual	0	0.0	3	75.0	3	14.3	3	12.5	2	40.0	5	17.2	3	7.3	5	55.6	8	16.0
	Fem Pres Het Cont	0	0.0	1	25.0	1	4.8	0	0.0	3	60.0	3	10.3	0	0.0	4	44.4	4	8.0
	Pediatric Risk	0	0.0	0	0.0	0	0.0	1	4.2	0	0.0	1	3.4	1	2.4	0	0.0	1	2.0
	Unk	2	11.8	0	0.0	2	9.5	2	8.3	0	0.0	2	6.9	4	9.8	0	0.0	4	8.0

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		3.3	100.0	5.2	2.7	100.0	4.2	50	100.0	64,160
Gender	Male	3.0	90.0		2.0	75.0		41	82.0	
	Female	0.3	10.0		0.7	25.0		9	18.0	
Age at Diagnosis	13-19	0.0	0.0		0.0	0.0		1	2.0	
	20-24	0.3	10.0		0.3	12.5		2	4.0	
	25-29	0.3	10.0		0.3	12.5		3	6.0	
	30-39	0.0	0.0		0.0	0.0		21	42.0	
	40-49	2.0	60.0		1.3	50.0		16	32.0	
	50-59	0.3	10.0		0.0	0.0		2	4.0	
	60+	0.3	10.0		0.7	25.0		5	10.0	
Race/Ethnicity	White	3.0	90.0		1.7	62.5		33	66.0	
	Black	0.0	0.0		0.3	12.5		8	16.0	
	Hispanic	0.0	0.0		0.0	0.0		1	2.0	
	Multi Race	0.3	10.0		0.7	25.0		6	12.0	
	Others	0.0	0.0		0.0	0.0		2	4.0	
Risk	MSM	2.3	70.0		1.0	37.5		28	56.0	
	IDU	0.0	0.0		0.0	0.0		4	8.0	
	MSM/IDU	0.3	10.0		0.3	12.5		1	2.0	
	Heterosexual	0.3	10.0		1.0	37.5		8	16.0	
	Fem Pres Het Cont	0.3	10.0		0.3	12.5		3	6.0	
	Pediatric Risk	0.0	0.0		0.0	0.0		1	2.0	
	Unk	0.0	0.0		0.0	0.0		5	10.0	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		1	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fei	nale	To	otal	М	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		3.0	100.0	0.3	100.0	3.3	100.0	2.0	100.0	0.7	100.0	2.7	100.0	41	100.0	9	100.0	50	100.0
Risk	MSM	2.3	77.8	0.0	0.0	2.3	70.0	1.0	50.0	0.0	0.0	1.0	37.5	28	68.3	0	0.0	28	56.0
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3	7.3	1	11.1	4	8.0
	MSM/IDU	0.3	11.1	0.0	0.0	0.3	10.0	0.3	16.7	0.0	0.0	0.3	12.5	1	2.4	0	0.0	1	2.0
	Heterosexual	0.3	11.1	0.0	0.0	0.3	10.0	0.7	33.3	0.3	50.0	1.0	37.5	3	7.3	5	55.6	8	16.0
	Fem Pres Het Cont	0.0	0.0	0.3	100.0	0.3	10.0	0.0	0.0	0.3	50.0	0.3	12.5	0	0.0	3	33.3	3	6.0
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	2.4	0	0.0	1	2.0
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	5	12.2	0	0.0	5	10.0

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
1986		2	1
1988		3	2
1989		3	1
1990		1	1
1991		5	1
1992		1	4
1993		2	3
1994		2	
1995		5	2
1996		3	2
1998		2	
1999		1	
2000*	1		
2001	2	1	
2002	1	٠	
2003	2	2	1
2004		1	2
2005	1	4	
2006	2	2	
2007	1	2	
2008	2		
2010	6	5	
2011	4	3	1

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	AIDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		26	100.0	22.1	50	100.0	42.4	76	100.0	64.5	117,910
Gender	Male	15	57.7		32	64.0		47	61.8		
	Female	11	42.3		18	36.0		29	38.2		
Current Age	12 & under	0	0.0		0	0.0		0	0.0		
	13-19	0	0.0		0	0.0		0	0.0		
	20-24	0	0.0		1	2.0		1	1.3		
	25-29	3	11.5		0	0.0		3	3.9		
	30-39	8	30.8		10	20.0		18	23.7		
	40-49	8	30.8		16	32.0		24	31.6		
	50-59	5	19.2		18	36.0		23	30.3		
	60+	2	7.7		5	10.0		7	9.2		
Race/Ethnicity	White	14	53.8		32	64.0		46	60.5		
	Black	6	23.1		8	16.0		14	18.4		
	Hispanic	3	11.5		6	12.0		9	11.8		
	Multi Race	2	7.7		3	6.0		5	6.6		
	Others	1	3.8		1	2.0		2	2.6		
Risk	MSM	11	42.3		15	30.0		26	34.2		
	IDU	1	3.8		7	14.0		8	10.5		
	MSM/IDU	1	3.8		1	2.0		2	2.6		
	Heterosexual	10	38.5		12	24.0		22	28.9		
	Fem Pres Het Cont	2	7.7		5	10.0		7	9.2		
	Blood Prod	0	0.0		1	2.0		1	1.3		
	Pediatric Risk	0	0.0		1	2.0		1	1.3		
	Unk	1	3.8		8	16.0		9	11.8		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	[ale	Fe	male	To	otal	N	Male	Fe	male	1	otal	M	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		15	100.0	11	100.0	26	100.0	32	100.0	18	100.0	50	100.0	47	100.0	29	100.0	76	100.0
Risk	MSM	11	73.3	0	0.0	11	42.3	15	46.9	0	0.0	15	30.0	26	55.3	0	0.0	26	34.2
	IDU	0	0.0	1	9.1	1	3.8	6	18.8	1	5.6	7	14.0	6	12.8	2	6.9	8	10.5
	MSM/IDU	1	6.7	0	0.0	1	3.8	1	3.1	0	0.0	1	2.0	2	4.3	0	0.0	2	2.6
	Heterosexual	2	13.3	8	72.7	10	38.5	1	3.1	11	61.1	12	24.0	3	6.4	19	65.5	22	28.9
	Fem Pres Het Cont	0	0.0	2	18.2	2	7.7	0	0.0	5	27.8	5	10.0	0	0.0	7	24.1	7	9.2
	Blood Prod	0	0.0	0	0.0	0	0.0	1	3.1	0	0.0	1	2.0	1	2.1	0	0.0	1	1.3
	Pediatric Risk	0	0.0	0	0.0	0	0.0	0	0.0	1	5.6	1	2.0	0	0.0	1	3.4	1	1.3
	Unk	1	6.7	0	0.0	1	3.8	8	25.0	0	0.0	8	16.0	9	19.1	0	0.0	9	11.8

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		3.7	100.0	3.1	5.0	100.0	4.2	90	100.0	117,910
Gender	Male	1.7	45.5		2.7	53.3		60	66.7	
	Female	2.0	54.5		2.3	46.7		30	33.3	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		1	1.1	
	13-19	0.0	0.0		0.3	6.7		2	2.2	
	20-24	0.0	0.0		0.0	0.0		2	2.2	
	25-29	0.0	0.0		0.7	13.3		16	17.8	
	30-39	1.0	27.3		1.3	26.7		33	36.7	
	40-49	1.7	45.5		1.3	26.7		19	21.1	
	50-59	1.0	27.3		1.3	26.7		12	13.3	
	60+	0.0	0.0		0.0	0.0		5	5.6	
Race/Ethnicity	White	2.0	54.5		2.3	46.7		61	67.8	
	Black	1.0	27.3		1.3	26.7		16	17.8	
	Hispanic	0.7	18.2		0.7	13.3		9	10.0	
	Multi Race	0.0	0.0		0.3	6.7		3	3.3	
	Others	0.0	0.0		0.3	6.7		1	1.1	
Risk	MSM	1.0	27.3		1.0	20.0		31	34.4	
	IDU	0.3	9.1		1.0	20.0		17	18.9	
	MSM/IDU	0.0	0.0		0.0	0.0		2	2.2	
	Heterosexual	0.7	18.2		1.3	26.7		19	21.1	
	Fem Pres Het Cont	1.0	27.3		0.7	13.3		8	8.9	
	Blood Prod	0.0	0.0		0.0	0.0		1	1.1	
	Pediatric Risk	0.0	0.0		0.3	6.7		2	2.2	
	Unk	0.7	18.2		0.7	13.3		10	11.1	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Diag	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		1.7	100.0	2.0	100.0	3.7	100.0	2.7	100.0	2.3	100.0	5.0	100.0	60	100.0	30	100.0	90	100.0
Risk	MSM	1.0	60.0	0.0	0.0	1.0	27.3	1.0	37.5	0.0	0.0	1.0	20.0	31	51.7	0	0.0	31	34.4
	IDU	0.0	0.0	0.3	16.7	0.3	9.1	0.7	25.0	0.3	14.3	1.0	20.0	12	20.0	5	16.7	17	18.9
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	3.3	0	0.0	2	2.2
	Heterosexual	0.0	0.0	0.7	33.3	0.7	18.2	0.3	12.5	1.0	42.9	1.3	26.7	4	6.7	15	50.0	19	21.1
	Fem Pres Het Cont	0.0	0.0	1.0	50.0	1.0	27.3	0.0	0.0	0.7	28.6	0.7	13.3	0	0.0	8	26.7	8	8.9
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	1.7	0	0.0	1	1.1
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	14.3	0.3	6.7	0	0.0	2	6.7	2	2.2
	Unk	0.7	40.0	0.0	0.0	0.7	18.2	0.7	25.0	0.0	0.0	0.7	13.3	10	16.7	0	0.0	10	11.1

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		1	
1987	٠	3	3
1988	•	1	1
1989		3	
1990	٠	2	2
1991	٠	1	1
1992		7	3
1993		10	3
1994		6	2
1995		4	7
1996		4	4
1997		1	1
1998		3	2
1999		3	2
2000*	4	4	
2001	1	2	1
2002	3	2	
2003	2	3	1
2004	2		2
2005	3	3	1
2006	3	6	1
2007	3	4	2
2008	3	2	
2009	1	5	
2010	8	8	
2011	2	2	1

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	1AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		5	100.0	18.5	1	100.0	3.7	6	100.0	22.2	27,072
Gender	Male	3	60.0		1	100.0		4	66.7		
	Female	2	40.0		0	0.0		2	33.3		
Current Age	30-39	0	0.0		0	0.0		0	0.0		
	40-49	1	20.0		1	100.0		2	33.3		
	50-59	4	80.0		0	0.0		4	66.7		
Race/Ethnicity	White	3	60.0		1	100.0		4	66.7		•
	Black	1	20.0		0	0.0		1	16.7		
	Hispanic	1	20.0		0	0.0		1	16.7		
Risk	MSM	3	60.0		0	0.0		3	50.0		
	Heterosexual	1	20.0		1	100.0		2	33.3		
	Fem Pres Het Cont	1	20.0		0	0.0		1	16.7		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		M	ale	Fe	male	Т	otal	N	Male	Fe	male	Т	otal	M	[ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		3	100.0	2	100.0	5	100.0	1	100.0	0	0.0	1	100.0	4	100.0	2	100.0	6	100.0
Risk	MSM	3	100.0	0	0.0	3	60.0	0	0.0	0	0.0	0	0.0	3	75.0	0	0.0	3	50.0
	Heterosexual	0	0.0	1	50.0	1	20.0	1	100.0	0	0.0	1	100.0	1	25.0	1	50.0	2	33.3
	Fem Pres Het Cont	0	0.0	1	50.0	1	20.0	0	0.0	0	0.0	0	0.0	0	0.0	1	50.0	1	16.7

		HIV Diagn	oses (Annua	l Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		0.0	0.0	0.0	0.0	0.0	0.0	1	100.0	27,072
Gender	Male	0.0	0.0	•	0.0	0.0		1	100.0	
	Female	0.0	0.0	•	0.0	0.0		0	0.0	
Age at Diagnosis	30-39	0.0	0.0	•	0.0	0.0		1	100.0	
	40-49	0.0	0.0	•	0.0	0.0		0	0.0	
	50-59	0.0	0.0		0.0	0.0		0	0.0	
Race/Ethnicity	White	0.0	0.0	•	0.0	0.0		1	100.0	
	Black	0.0	0.0		0.0	0.0		0	0.0	
	Hispanic	0.0	0.0		0.0	0.0		0	0.0	
Risk	MSM	0.0	0.0		0.0	0.0		0	0.0	
	Heterosexual	0.0	0.0	·	0.0	0.0		1	100.0	
	Fem Pres Het Cont	0.0	0.0		0.0	0.0		0	0.0	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS C	Cases	
		M	ale	Fer	nale	To	otal	M	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	100.0	0	0.0	1	100.0
Risk	MSM	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	0	0.0	0	0.0
	Heterosexual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	100.0	0	0.0	1	100.0
	Fem Pres Het Cont	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	0	0.0	0	0.0

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
2001	1		
2003	1	1	

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	1 AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		17	100.0	26.1	31	100.0	47.6	48	100.0	73.8	65,070
Gender	Male	12	70.6		23	74.2		35	72.9		
	Female	5	29.4		8	25.8		13	27.1		
Current Age	12 & under	0	0.0		0	0.0		0	0.0		
	13-19	0	0.0		0	0.0		0	0.0		
	20-24	0	0.0		1	3.2		1	2.1		
	25-29	0	0.0		2	6.5		2	4.2		
	30-39	3	17.6		2	6.5		5	10.4		
	40-49	8	47.1		12	38.7		20	41.7		
	50-59	2	11.8		10	32.3		12	25.0		
	60+	4	23.5		4	12.9		8	16.7		
Race/Ethnicity	White	15	88.2		21	67.7		36	75.0		
	Black	0	0.0		2	6.5		2	4.2		
	Hispanic	2	11.8		6	19.4		8	16.7		
	Multi Race	0	0.0		1	3.2		1	2.1		
	Others	0	0.0		1	3.2		1	2.1		
Risk	MSM	6	35.3		13	41.9		19	39.6		
	IDU	0	0.0		3	9.7		3	6.3		
	MSM/IDU	2	11.8		1	3.2		3	6.3		
	Heterosexual	2	11.8		7	22.6		9	18.8		
	Fem Pres Het Cont	3	17.6		2	6.5		5	10.4		
	Blood Prod	0	0.0		1	3.2		1	2.1		
	Pediatric Risk	0	0.0		1	3.2		1	2.1		
	Unk	4	23.5		3	9.7		7	14.6		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	Male	Fe	emale	1	otal	M	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		12	100.0	5	100.0	17	100.0	23	100.0	8	100.0	31	100.0	35	100.0	13	100.0	48	100.0
Risk	MSM	6	50.0	0	0.0	6	35.3	13	56.5	0	0.0	13	41.9	19	54.3	0	0.0	19	39.6
	IDU	0	0.0	0	0.0	0	0.0	3	13.0	0	0.0	3	9.7	3	8.6	0	0.0	3	6.3
	MSM/IDU	2	16.7	0	0.0	2	11.8	1	4.3	0	0.0	1	3.2	3	8.6	0	0.0	3	6.3
	Heterosexual	0	0.0	2	40.0	2	11.8	2	8.7	5	62.5	7	22.6	2	5.7	7	53.8	9	18.8
	Fem Pres Het Cont	0	0.0	3	60.0	3	17.6	0	0.0	2	25.0	2	6.5	0	0.0	5	38.5	5	10.4
	Blood Prod	0	0.0	0	0.0	0	0.0	1	4.3	0	0.0	1	3.2	1	2.9	0	0.0	1	2.1
	Pediatric Risk	0	0.0	0	0.0	0	0.0	0	0.0	1	12.5	1	3.2	0	0.0	1	7.7	1	2.1
	Unk	4	33.3	0	0.0	4	23.5	3	13.0	0	0.0	3	9.7	7	20.0	0	0.0	7	14.6

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		2.7	100.0	4.1	1.3	100.0	2.0	67	100.0	65,070
Gender	Male	1.7	62.5		0.7	50.0		53	79.1	
	Female	1.0	37.5		0.7	50.0		14	20.9	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		1	1.5	
	13-19	0.0	0.0		0.0	0.0		4	6.0	
	20-24	0.0	0.0		0.0	0.0		3	4.5	
	25-29	0.3	12.5		0.3	25.0		9	13.4	
	30-39	0.7	25.0		0.0	0.0		19	28.4	
	40-49	1.0	37.5		1.0	75.0		27	40.3	
	50-59	0.3	12.5		0.0	0.0		3	4.5	
	60+	0.3	12.5		0.0	0.0		1	1.5	
Race/Ethnicity	White	2.0	75.0		1.0	75.0		52	77.6	
	Black	0.0	0.0		0.0	0.0		6	9.0	
	Hispanic	0.7	25.0		0.3	25.0		6	9.0	
	Multi Race	0.0	0.0		0.0	0.0		2	3.0	
	Others	0.0	0.0		0.0	0.0		1	1.5	
Risk	MSM	0.7	25.0		0.7	50.0		24	35.8	
	IDU	0.0	0.0		0.0	0.0		11	16.4	
	MSM/IDU	0.3	12.5		0.0	0.0		2	3.0	
	Heterosexual	0.7	25.0		0.7	50.0		8	11.9	
	Fem Pres Het Cont	0.3	12.5		0.0	0.0		6	9.0	
	Blood Prod	0.0	0.0		0.0	0.0		6	9.0	
	Pediatric Risk	0.0	0.0		0.0	0.0		3	4.5	
	Unk	0.7	25.0		0.0	0.0		7	10.4	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

]	HIV Dia	gnoses ((Annual A	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS C	Cases	
		M	ale	Fer	nale	To	otal	M	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		1.7	100.0	1.0	100.0	2.7	100.0	0.7	100.0	0.7	100.0	1.3	100.0	53	100.0	14	100.0	67	100.0
Risk	MSM	0.7	40.0	0.0	0.0	0.7	25.0	0.7	100.0	0.0	0.0	0.7	50.0	24	45.3	0	0.0	24	35.8
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	10	18.9	1	7.1	11	16.4
	MSM/IDU	0.3	20.0	0.0	0.0	0.3	12.5	0.0	0.0	0.0	0.0	0.0	0.0	2	3.8	0	0.0	2	3.0
	Heterosexual	0.0	0.0	0.7	66.7	0.7	25.0	0.0	0.0	0.7	100.0	0.7	50.0	2	3.8	6	42.9	8	11.9
	Fem Pres Het Cont	0.0	0.0	0.3	33.3	0.3	12.5	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	6	42.9	6	9.0
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	6	11.3	0	0.0	6	9.0
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	3.8	1	7.1	3	4.5
	Unk	0.7	40.0	0.0	0.0	0.7	25.0	0.0	0.0	0.0	0.0	0.0	0.0	7	13.2	0	0.0	7	10.4

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		1	
1986		1	2
1987		2	2
1988		6	1
1989		3	
1990		2	3
1991	•	5	2
1992	•	2	4
1993		6	4
1994		7	1
1995		3	2
1996			3
1997		2	1
1998	•	3	1
1999		2	1
2000*	2	1	1
2001		4	2
2002	•	2	1
2003	3	2	
2004	1		
2005	1	1	1
2006	1	2	
2007	2	4	
2008	1	2	1
2009	4	1	2
2010	2	1	
2011	2	2	1

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	I AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		19	100.0	25.9	17	100.0	23.2	36	100.0	49.1	73,365
Gender	Male	16	84.2		10	58.8		26	72.2		
	Female	3	15.8		7	41.2		10	27.8		
Current Age	13-19	0	0.0		0	0.0		0	0.0		
	25-29	1	5.3		0	0.0		1	2.8		
	30-39	3	15.8		2	11.8		5	13.9		
	40-49	9	47.4		9	52.9		18	50.0		
	50-59	4	21.1		3	17.6		7	19.4		
	60+	2	10.5		3	17.6		5	13.9		
Race/Ethnicity	White	15	78.9		12	70.6		27	75.0		
	Black	3	15.8		0	0.0		3	8.3		
	Hispanic	1	5.3		2	11.8		3	8.3		
	Multi Race	0	0.0		2	11.8		2	5.6		
	Others	0	0.0		1	5.9		1	2.8		
Risk	MSM	11	57.9		5	29.4		16	44.4		
	IDU	0	0.0		4	23.5		4	11.1		
	MSM/IDU	0	0.0		1	5.9		1	2.8		
	Heterosexual	1	5.3		4	23.5		5	13.9		
	Fem Pres Het Cont	2	10.5		1	5.9		3	8.3		
	Blood Prod	0	0.0		1	5.9		1	2.8		
	Pediatric Risk	0	0.0		0	0.0		0	0.0		
	Unk	5	26.3		1	5.9		6	16.7		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		M	ale	Fe	male	To	otal	N	Male	Fe	emale	1	otal	M	[ale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		16	100.0	3	100.0	19	100.0	10	100.0	7	100.0	17	100.0	26	100.0	10	100.0	36	100.0
Risk	MSM	11	68.8	0	0.0	11	57.9	5	50.0	0	0.0	5	29.4	16	61.5	0	0.0	16	44.4
	IDU	0	0.0	0	0.0	0	0.0	2	20.0	2	28.6	4	23.5	2	7.7	2	20.0	4	11.1
	MSM/IDU	0	0.0	0	0.0	0	0.0	1	10.0	0	0.0	1	5.9	1	3.8	0	0.0	1	2.8
	Heterosexual	0	0.0	1	33.3	1	5.3	0	0.0	4	57.1	4	23.5	0	0.0	5	50.0	5	13.9
	Fem Pres Het Cont	0	0.0	2	66.7	2	10.5	0	0.0	1	14.3	1	5.9	0	0.0	3	30.0	3	8.3
	Blood Prod	0	0.0	0	0.0	0	0.0	1	10.0	0	0.0	1	5.9	1	3.8	0	0.0	1	2.8
	Pediatric Risk	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Unk	5	31.3	0	0.0	5	26.3	1	10.0	0	0.0	1	5.9	6	23.1	0	0.0	6	16.7

		HIV Diagn	oses (Annua	ıl Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		1.3	100.0	1.8	1.3	100.0	1.8	43	100.0	73,365
Gender	Male	1.3	100.0		1.0	75.0		32	74.4	
	Female	0.0	0.0		0.3	25.0		11	25.6	
Age at Diagnosis	13-19	0.0	0.0		0.0	0.0		2	4.7	
	25-29	0.0	0.0		0.3	25.0		4	9.3	
	30-39	0.7	50.0		0.7	50.0		24	55.8	
	40-49	0.0	0.0		0.0	0.0		6	14.0	
	50-59	0.7	50.0		0.3	25.0		6	14.0	
	60+	0.0	0.0		0.0	0.0		1	2.3	
Race/Ethnicity	White	1.3	100.0		1.0	75.0		35	81.4	
	Black	0.0	0.0		0.0	0.0		2	4.7	
	Hispanic	0.0	0.0		0.0	0.0		3	7.0	
	Multi Race	0.0	0.0		0.3	25.0		2	4.7	
	Others	0.0	0.0		0.0	0.0		1	2.3	
Risk	MSM	1.3	100.0		1.0	75.0		15	34.9	
	IDU	0.0	0.0		0.0	0.0		11	25.6	
	MSM/IDU	0.0	0.0		0.0	0.0		3	7.0	
	Heterosexual	0.0	0.0		0.3	25.0		8	18.6	
	Fem Pres Het Cont	0.0	0.0		0.0	0.0		1	2.3	
	Blood Prod	0.0	0.0		0.0	0.0		2	4.7	
	Pediatric Risk	0.0	0.0		0.0	0.0		1	2.3	
	Unk	0.0	0.0		0.0	0.0		2	4.7	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	То	otal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		1.3	100.0	0.0	0.0	1.3	100.0	1.0	100.0	0.3	100.0	1.3	100.0	32	100.0	11	100.0	43	100.0
Risk	MSM	1.3	100.0	0.0	0.0	1.3	100.0	1.0	100.0	0.0	0.0	1.0	75.0	15	46.9	0	0.0	15	34.9
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	7	21.9	4	36.4	11	25.6
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3	9.4	0	0.0	3	7.0
	Heterosexual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	100.0	0.3	25.0	2	6.3	6	54.5	8	18.6
	Fem Pres Het Cont	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	1	9.1	1	2.3
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	6.3	0	0.0	2	4.7
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	3.1	0	0.0	1	2.3
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	6.3	0	0.0	2	4.7

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
1986		2	1
1988		1	1
1989		1	1
1990		5	1
1991		4	2
1992			1
1993		6	2
1994			2
1995		2	3
1996		2	1
1997		3	1
1998		3	
1999			1
2000*	1	4	
2001	1	1	1
2002			1
2003	2		
2004	1	1	1
2005	3	2	3
2006	4	1	
2008	1	1	1
2009	2	1	
2010	1	2	1
2011	1	1	1

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		930 100.0 676 72.7 254 27.3 0 0.0 20 2.2 69 7.4 73 7.8 159 17.1 283 30.4 240 25.8 86 9.2 342 36.8 380 40.9 144 15.5 60 6.5 4 0.4 446 48.0 94 10.1 30 3.2 136 14.6 115 12.4		AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		930	100.0	124.7	1,198	100.0	160.7	2,128	100.0	285.4	745,625
Gender	Male	676	72.7		835	69.7		1,511	71.0		
	Female	254	27.3		363	30.3		617	29.0		
Current Age	12 & under	0	0.0		1	0.1		1	0.0		
	13-19	20	2.2		3	0.3		23	1.1		
	20-24	69	7.4		27	2.3		96	4.5		
	25-29	73	7.8		34	2.8		107	5.0		
	30-39	159	17.1		105	8.8		264	12.4		
	40-49	283	30.4		404	33.7		687	32.3		
	50-59	240	25.8		433	36.1		673	31.6		
	60+	86	9.2		191	15.9		277	13.0		
Race/Ethnicity	White	342	36.8		410	34.2		752	35.3		
	Black	380	40.9		509	42.5		889	41.8		
	Hispanic	144	15.5		187	15.6		331	15.6		
	Multi Race	60	6.5		89	7.4		149	7.0		
	Others	4	0.4		3	0.3		7	0.3		
Risk	MSM	446	48.0		441	36.8		887	41.7		
	IDU	94	10.1		245	20.5		339	15.9		
	MSM/IDU	30	3.2		49	4.1		79	3.7		
	Heterosexual	136	14.6		188	15.7		324	15.2		
	Fem Pres Het Cont	115	12.4		129	10.8		244	11.5		
	Blood Prod	0	0.0		6	0.5		6	0.3		
	Pediatric Risk	14	1.5		13	1.1		27	1.3		
	Unk	95	10.2		127	10.6		222	10.4		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Ca	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	I	Male	Fe	male	Т	otal	M	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		676	100.0	254	100.0	930	100.0	835	100.0	363	100.0	1,198	100.0	1,511	100.0	617	100.0	2,128	100.0
Risk	MSM	446	66.0	0	0.0	446	48.0	441	52.8	0	0.0	441	36.8	887	58.7	0	0.0	887	41.7
	IDU	60	8.9	34	13.4	94	10.1	150	18.0	95	26.2	245	20.5	210	13.9	129	20.9	339	15.9
	MSM/IDU	30	4.4	0	0.0	30	3.2	49	5.9	0	0.0	49	4.1	79	5.2	0	0.0	79	3.7
	Heterosexual	41	6.1	95	37.4	136	14.6	56	6.7	132	36.4	188	15.7	97	6.4	227	36.8	324	15.2
	Fem Pres Het Cont	0	0.0	115	45.3	115	12.4	0	0.0	129	35.5	129	10.8	0	0.0	244	39.5	244	11.5
	Blood Prod	0	0.0	0	0.0	0	0.0	3	0.4	3	0.8	6	0.5	3	0.2	3	0.5	6	0.3
	Pediatric Risk	7	1.0	7	2.8	14	1.5	9	1.1	4	1.1	13	1.1	16	1.1	11	1.8	27	1.3
	Unk	92	13.6	3	1.2	95	10.2	127	15.2	0	0.0	127	10.6	219	14.5	3	0.5	222	10.4

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		86.7	100.0	11.6	58.3	100.0	7.8	2,600	100.0	745,625
Gender	Male	69.0	79.6		41.0	70.3		1,902	73.2	
	Female	17.7	20.4		17.3	29.7		698	26.8	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		13	0.5	
	13-19	6.7	7.7		1.0	1.7		16	0.6	
	20-24	19.3	22.3		6.3	10.9		83	3.2	
	25-29	11.3	13.1		3.3	5.7		279	10.7	
	30-39	15.0	17.3		11.3	19.4		995	38.3	
	40-49	18.0	20.8		16.3	28.0		821	31.6	
	50-59	11.3	13.1		10.7	18.3		282	10.8	
	60+	5.0	5.8		9.3	16.0		111	4.3	
Race/Ethnicity	White	27.0	31.2		16.0	27.4		1,049	40.3	
	Black	43.3	50.0		28.0	48.0		1,080	41.5	
	Hispanic	12.3	14.2		9.3	16.0		356	13.7	
	Multi Race	3.3	3.8		4.7	8.0		108	4.2	
	Others	0.7	0.8		0.3	0.6		7	0.3	
Risk	MSM	51.3	59.2		24.3	41.7		971	37.3	
	IDU	4.7	5.4		7.3	12.6		752	28.9	
	MSM/IDU	2.7	3.1		2.3	4.0		113	4.3	
	Heterosexual	10.7	12.3		10.3	17.7		287	11.0	
	Fem Pres Het Cont	9.3	10.8		8.0	13.7		186	7.2	
	Blood Prod	0.0	0.0		0.0	0.0		22	0.8	
	Pediatric Risk	0.0	0.0		0.3	0.6		19	0.7	
	Unk	8.0	9.2		5.7	9.7		250	9.6	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	M	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		69.0	100.0	17.7	100.0	86.7	100.0	41.0	100.0	17.3	100.0	58.3	100.0	1,902	100.0	698	100.0	2,600	100.0
Risk	MSM	51.3	74.4	0.0	0.0	51.3	59.2	24.3	59.3	0.0	0.0	24.3	41.7	971	51.1	0	0.0	971	37.3
	IDU	3.0	4.3	1.7	9.4	4.7	5.4	3.7	8.9	3.7	21.2	7.3	12.6	464	24.4	288	41.3	752	28.9
	MSM/IDU	2.7	3.9	0.0	0.0	2.7	3.1	2.3	5.7	0.0	0.0	2.3	4.0	113	5.9	0	0.0	113	4.3
	Heterosexual	4.7	6.8	6.0	34.0	10.7	12.3	4.7	11.4	5.7	32.7	10.3	17.7	74	3.9	213	30.5	287	11.0
	Fem Pres Het Cont	0.0	0.0	9.3	52.8	9.3	10.8	0.0	0.0	8.0	46.2	8.0	13.7	0	0.0	186	26.6	186	7.2
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	16	0.8	6	0.9	22	0.8
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.8	0.0	0.0	0.3	0.6	14	0.7	5	0.7	19	0.7
	Unk	7.3	10.6	0.7	3.8	8.0	9.2	5.7	13.8	0.0	0.0	5.7	9.7	250	13.1	0	0.0	250	9.6

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		23	16
1986		37	17
1987		50	26
1988		72	31
1989		79	46
1990		89	50
1991		120	85
1992		163	94
1993		196	79
1994		204	116
1995		201	122
1996		138	97
1997		113	59
1998		80	46
1999		92	51
2000*	76	79	53
2001	91	107	38
2002	113	95	45
2003	92	71	41
2004	101	81	53
2005	79	96	41
2006	73	77	42
2007	97	88	40
2008	89	74	37
2009	89	62	25
2010	70	44	26
2011	101	69	32

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		33	100.0	66.1	34	100.0	68.1	67	100.0	134.2	49,916
Gender	Male	17	51.5		17	50.0		34	50.7		
	Female	16	48.5		17	50.0		33	49.3		
Current Age	20-24	3	9.1		0	0.0		3	4.5		
	25-29	0	0.0		0	0.0		0	0.0		
	30-39	5	15.2		5	14.7		10	14.9		
	40-49	9	27.3		14	41.2		23	34.3		
	50-59	10	30.3		12	35.3		22	32.8		
	60+	6	18.2		3	8.8		9	13.4		
Race/Ethnicity	White	11	33.3		12	35.3		23	34.3		
	Black	3	9.1		3	8.8		6	9.0		
	Hispanic	17	51.5		17	50.0		34	50.7		
	Multi Race	2	6.1		2	5.9		4	6.0		
Risk	MSM	7	21.2		5	14.7		12	17.9		
	IDU	6	18.2		8	23.5		14	20.9		
	MSM/IDU	1	3.0		2	5.9		3	4.5		
	Heterosexual	12	36.4		10	29.4		22	32.8		
	Fem Pres Het Cont	4	12.1		3	8.8		7	10.4		
	Blood Prod	0	0.0		0	0.0		0	0.0		
	Pediatric Risk	1	3.0		0	0.0		1	1.5		
	Unk	2	6.1		6	17.6		8	11.9		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		M	ale	Fe	male	Т	otal	N	Male	Fe	emale	1	otal	M	ale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		17	100.0	16	100.0	33	100.0	17	100.0	17	100.0	34	100.0	34	100.0	33	100.0	67	100.0
Risk	MSM	7	41.2	0	0.0	7	21.2	5	29.4	0	0.0	5	14.7	12	35.3	0	0.0	12	17.9
	IDU	3	17.6	3	18.8	6	18.2	3	17.6	5	29.4	8	23.5	6	17.6	8	24.2	14	20.9
	MSM/IDU	1	5.9	0	0.0	1	3.0	2	11.8	0	0.0	2	5.9	3	8.8	0	0.0	3	4.5
	Heterosexual	4	23.5	8	50.0	12	36.4	1	5.9	9	52.9	10	29.4	5	14.7	17	51.5	22	32.8
	Fem Pres Het Cont	0	0.0	4	25.0	4	12.1	0	0.0	3	17.6	3	8.8	0	0.0	7	21.2	7	10.4
	Blood Prod	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Pediatric Risk	0	0.0	1	6.3	1	3.0	0	0.0	0	0.0	0	0.0	0	0.0	1	3.0	1	1.5
	Unk	2	11.8	0	0.0	2	6.1	6	35.3	0	0.0	6	17.6	8	23.5	0	0.0	8	11.9

		HIV Diagn	oses (Annua	ıl Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		1.3	100.0	2.7	1.3	100.0	2.7	82	100.0	49,916
Gender	Male	0.7	50.0		1.0	75.0		56	68.3	
	Female	0.7	50.0		0.3	25.0		26	31.7	
Age at Diagnosis	20-24	0.0	0.0		0.0	0.0		3	3.7	
	25-29	0.7	50.0		0.0	0.0		13	15.9	
	30-39	0.3	25.0		0.0	0.0		30	36.6	
	40-49	0.0	0.0		0.7	50.0		19	23.2	
	50-59	0.3	25.0		0.7	50.0		14	17.1	
	60+	0.0	0.0		0.0	0.0		3	3.7	
Race/Ethnicity	White	0.7	50.0		0.3	25.0		37	45.1	
	Black	0.7	50.0		0.0	0.0		9	11.0	
	Hispanic	0.0	0.0		1.0	75.0		33	40.2	
	Multi Race	0.0	0.0		0.0	0.0		3	3.7	
Risk	MSM	0.3	25.0		0.3	25.0		16	19.5	
	IDU	0.3	25.0		0.3	25.0		28	34.1	
	MSM/IDU	0.0	0.0		0.0	0.0		3	3.7	
	Heterosexual	0.7	50.0		0.3	25.0		16	19.5	
	Fem Pres Het Cont	0.0	0.0		0.0	0.0		6	7.3	
	Blood Prod	0.0	0.0		0.0	0.0		3	3.7	
	Pediatric Risk	0.0	0.0		0.0	0.0		0	0.0	
	Unk	0.0	0.0		0.3	25.0		10	12.2	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Diag	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS C	Cases	
		M	ale	Fer	nale	To	otal	M	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		0.7	100.0	0.7	100.0	1.3	100.0	1.0	100.0	0.3	100.0	1.3	100.0	56	100.0	26	100.0	82	100.0
Risk	MSM	0.3	50.0	0.0	0.0	0.3	25.0	0.3	33.3	0.0	0.0	0.3	25.0	16	28.6	0	0.0	16	19.5
	IDU	0.0	0.0	0.3	50.0	0.3	25.0	0.3	33.3	0.0	0.0	0.3	25.0	18	32.1	10	38.5	28	34.1
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3	5.4	0	0.0	3	3.7
	Heterosexual	0.3	50.0	0.3	50.0	0.7	50.0	0.0	0.0	0.3	100.0	0.3	25.0	6	10.7	10	38.5	16	19.5
	Fem Pres Het Cont	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	6	23.1	6	7.3
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3	5.4	0	0.0	3	3.7
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	0	0.0	0	0.0
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.3	33.3	0.0	0.0	0.3	25.0	10	17.9	0	0.0	10	12.2

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		1	1
1987		2	
1988		4	1
1989		2	4
1990	٠	1	2
1991		7	1
1992		6	5
1993		4	3
1994	٠	8	5
1995		3	5
1996		5	3
1997		5	1
1998		5	
1999		3	1
2000*	3	2	3
2001	2	2	2
2002	4	3	
2003	3	2	3
2004	2	2	1
2005	3	4	2
2006	3	2	
2007	1	2	
2008	4	3	2
2009	2		3
2010	1	2	
2011	1	2	1

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	d AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		1,110	100.0	82.6	1,753	100.0	130.4	2,863	100.0	213.0	1,344,436
Gender	Male	713	64.2		1,199	68.4		1,912	66.8		
	Female	397	35.8		554	31.6		951	33.2		
Current Age	12 & under	6	0.5		0	0.0		6	0.2		
	13-19	36	3.2		7	0.4		43	1.5		
	20-24	65	5.9		38	2.2		103	3.6		
	25-29	79	7.1		55	3.1		134	4.7		
	30-39	212	19.1		169	9.6		381	13.3		
	40-49	313	28.2		556	31.7		869	30.4		
	50-59	284	25.6		629	35.9		913	31.9		
	60+	114	10.3		299	17.1		413	14.4		
	Unk	1	0.1		0	0.0		1	0.0		
Race/Ethnicity	White	369	33.2		523	29.8		892	31.2		
	Black	429	38.6		692	39.5		1,121	39.2		
	Hispanic	242	21.8		382	21.8		624	21.8		
	Multi Race	57	5.1		142	8.1		199	7.0		
	Others	13	1.2		14	0.8		27	0.9		
	Unk	0	0.0		0	0.0		0	0.0		
Risk	MSM	380	34.2		498	28.4		878	30.7		
	IDU	87	7.8		299	17.1		386	13.5		
	MSM/IDU	26	2.3		52	3.0		78	2.7		
	Heterosexual	198	17.8		357	20.4		555	19.4		
	Fem Pres Het Cont	183	16.5		198	11.3		381	13.3		
	Blood Prod	3	0.3		18	1.0		21	0.7		
	Pediatric Risk	42	3.8		29	1.7		71	2.5		
	Unk	191	17.2		302	17.2		493	17.2		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	Iale	Fe	male	Т	otal	М	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		713	100.0	397	100.0	1,110	100.0	1,199	100.0	554	100.0	1,753	100.0	1,912	100.0	951	100.0	2,863	100.0
Risk	MSM	380	53.3	0	0.0	380	34.2	498	41.5	0	0.0	498	28.4	878	45.9	0	0.0	878	30.7
	IDU	53	7.4	34	8.6	87	7.8	208	17.3	91	16.4	299	17.1	261	13.7	125	13.1	386	13.5
	MSM/IDU	26	3.6	0	0.0	26	2.3	52	4.3	0	0.0	52	3.0	78	4.1	0	0.0	78	2.7
	Heterosexual	44	6.2	154	38.8	198	17.8	109	9.1	248	44.8	357	20.4	153	8.0	402	42.3	555	19.4
	Fem Pres Het Cont	0	0.0	183	46.1	183	16.5	0	0.0	198	35.7	198	11.3	0	0.0	381	40.1	381	13.3
	Blood Prod	0	0.0	3	0.8	3	0.3	12	1.0	6	1.1	18	1.0	12	0.6	9	0.9	21	0.7
	Pediatric Risk	19	2.7	23	5.8	42	3.8	18	1.5	11	2.0	29	1.7	37	1.9	34	3.6	71	2.5
	Unk	191	26.8	0	0.0	191	17.2	302	25.2	0	0.0	302	17.2	493	25.8	0	0.0	493	17.2

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		103.0	100.0	7.7	79.7	100.0	5.9	4,447	100.0	1,344,436
Gender	Male	76.7	74.4		54.3	68.2		3,268	73.5	
	Female	26.3	25.6		25.3	31.8		1,179	26.5	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		45	1.0	
	13-19	7.0	6.8		1.7	2.1		40	0.9	
	20-24	19.3	18.8		8.0	10.0		140	3.1	
	25-29	12.7	12.3		8.7	10.9		435	9.8	
	30-39	22.0	21.4		16.3	20.5		1,752	39.4	
	40-49	21.0	20.4		20.7	25.9		1,325	29.8	
	50-59	16.7	16.2		19.0	23.8		499	11.2	
	60+	4.3	4.2		5.3	6.7		211	4.7	
	Unk	0.0	0.0		0.0	0.0		0	0.0	
Race/Ethnicity	White	26.0	25.2		13.0	16.3		1,788	40.2	
	Black	40.7	39.5		34.3	43.1		1,843	41.4	
	Hispanic	31.7	30.7		24.0	30.1		588	13.2	
	Multi Race	3.0	2.9		7.3	9.2		201	4.5	
	Others	1.7	1.6		1.0	1.3		26	0.6	
	Unk	0.0	0.0		0.0	0.0		1	0.0	
Risk	MSM	48.3	46.9		27.7	34.7		1,237	27.8	
	IDU	2.0	1.9		4.0	5.0		1,383	31.1	
	MSM/IDU	1.0	1.0		1.0	1.3		163	3.7	
	Heterosexual	14.3	13.9		16.7	20.9		630	14.2	
	Fem Pres Het Cont	16.7	16.2		13.3	16.7		313	7.0	
	Blood Prod	0.0	0.0		0.0	0.0		65	1.5	
	Pediatric Risk	0.0	0.0		1.3	1.7		70	1.6	
	Unk	20.7	20.1		15.7	19.7		586	13.2	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	M	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		76.7	100.0	26.3	100.0	103.0	100.0	54.3	100.0	25.3	100.0	79.7	100.0	3,268	100.0	1,179	100.0	4,447	100.0
Risk	MSM	48.3	63.0	0.0	0.0	48.3	46.9	27.7	50.9	0.0	0.0	27.7	34.7	1,237	37.9	0	0.0	1,237	27.8
	IDU	1.7	2.2	0.3	1.3	2.0	1.9	3.0	5.5	1.0	3.9	4.0	5.0	1,011	30.9	372	31.6	1,383	31.1
	MSM/IDU	1.0	1.3	0.0	0.0	1.0	1.0	1.0	1.8	0.0	0.0	1.0	1.3	163	5.0	0	0.0	163	3.7
	Heterosexual	5.0	6.5	9.3	35.4	14.3	13.9	6.7	12.3	10.0	39.5	16.7	20.9	185	5.7	445	37.7	630	14.2
	Fem Pres Het Cont	0.0	0.0	16.7	63.3	16.7	16.2	0.0	0.0	13.3	52.6	13.3	16.7	0	0.0	313	26.5	313	7.0
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	48	1.5	17	1.4	65	1.5
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.6	1.0	3.9	1.3	1.7	40	1.2	30	2.5	70	1.6
	Unk	20.7	27.0	0.0	0.0	20.7	20.1	15.7	28.8	0.0	0.0	15.7	19.7	584	17.9	2	0.2	586	13.2

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		134	79
1986		119	70
1987		143	101
1988		190	114
1989		226	149
1990		224	150
1991		246	188
1992		293	191
1993		304	201
1994		251	238
1995		294	224
1996		235	137
1997		164	80
1998		150	68
1999		151	74
2000*	149	166	60
2001	156	154	67
2002	130	149	56
2003	112	123	79
2004	113	118	54
2005	124	94	55
2006	110	93	53
2007	100	94	59
2008	133	93	41
2009	89	87	39
2010	100	80	43
2011	120	72	28

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		85	100.0	39.3	96	100.0	44.4	181	100.0	83.8	216,011
Gender	Male	60	70.6		63	65.6		123	68.0		
	Female	25	29.4		33	34.4		58	32.0		
Current Age	12 & under	0	0.0		0	0.0		0	0.0		
	13-19	2	2.4		1	1.0		3	1.7		
	20-24	4	4.7		3	3.1		7	3.9		
	25-29	10	11.8		5	5.2		15	8.3		
	30-39	17	20.0		10	10.4		27	14.9		
	40-49	26	30.6		40	41.7		66	36.5		
	50-59	20	23.5		27	28.1		47	26.0		
	60+	6	7.1		10	10.4		16	8.8		
Race/Ethnicity	White	45	52.9		47	49.0		92	50.8		
	Black	23	27.1		28	29.2		51	28.2		
	Hispanic	9	10.6		10	10.4		19	10.5		
	Multi Race	8	9.4		11	11.5		19	10.5		
	Others	0	0.0		0	0.0		0	0.0		
Risk	MSM	38	44.7		38	39.6		76	42.0		
	IDU	11	12.9		23	24.0		34	18.8		
	MSM/IDU	5	5.9		5	5.2		10	5.5		
	Heterosexual	17	20.0		23	24.0		40	22.1		
	Fem Pres Het Cont	8	9.4		2	2.1		10	5.5		
	Blood Prod	1	1.2		0	0.0		1	0.6		
	Pediatric Risk	0	0.0		3	3.1		3	1.7		
	Unk	5	5.9		2	2.1		7	3.9		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	Male	Fe	male	7	otal	M	Iale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		60	100.0	25	100.0	85	100.0	63	100.0	33	100.0	96	100.0	123	100.0	58	100.0	181	100.0
Risk	MSM	38	63.3	0	0.0	38	44.7	38	60.3	0	0.0	38	39.6	76	61.8	0	0.0	76	42.0
	IDU	9	15.0	2	8.0	11	12.9	14	22.2	9	27.3	23	24.0	23	18.7	11	19.0	34	18.8
	MSM/IDU	5	8.3	0	0.0	5	5.9	5	7.9	0	0.0	5	5.2	10	8.1	0	0.0	10	5.5
	Heterosexual	2	3.3	15	60.0	17	20.0	3	4.8	20	60.6	23	24.0	5	4.1	35	60.3	40	22.1
	Fem Pres Het Cont	0	0.0	8	32.0	8	9.4	0	0.0	2	6.1	2	2.1	0	0.0	10	17.2	10	5.5
	Blood Prod	1	1.7	0	0.0	1	1.2	0	0.0	0	0.0	0	0.0	1	0.8	0	0.0	1	0.6
	Pediatric Risk	0	0.0	0	0.0	0	0.0	1	1.6	2	6.1	3	3.1	1	0.8	2	3.4	3	1.7
	Unk	5	8.3	0	0.0	5	5.9	2	3.2	0	0.0	2	2.1	7	5.7	0	0.0	7	3.9

		HIV Diagn	oses (Annua	ıl Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		9.0	100.0	4.2	4.7	100.0	2.2	268	100.0	216,011
Gender	Male	7.3	81.5		3.7	78.6		186	69.4	
	Female	1.7	18.5		1.0	21.4		82	30.6	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		3	1.1	
	13-19	0.7	7.4		0.0	0.0		5	1.9	
	20-24	2.7	29.6		1.7	35.7		11	4.1	
	25-29	1.0	11.1		0.0	0.0		34	12.7	
	30-39	1.3	14.8		1.0	21.4		98	36.6	
	40-49	2.3	25.9		1.3	28.6		83	31.0	
	50-59	0.3	3.7		0.3	7.1		28	10.4	
	60+	0.7	7.4		0.3	7.1		6	2.2	
Race/Ethnicity	White	4.7	51.9		2.3	50.0		141	52.6	
	Black	3.7	40.7		2.0	42.9		91	34.0	
	Hispanic	0.0	0.0		0.0	0.0		19	7.1	
	Multi Race	0.7	7.4		0.3	7.1		15	5.6	
	Others	0.0	0.0		0.0	0.0		2	0.7	
Risk	MSM	5.7	63.0		2.7	57.1		98	36.6	
	IDU	0.7	7.4		0.3	7.1		86	32.1	
	MSM/IDU	0.3	3.7		0.3	7.1		16	6.0	
	Heterosexual	2.0	22.2		1.0	21.4		45	16.8	
	Fem Pres Het Cont	0.3	3.7		0.3	7.1		7	2.6	
	Blood Prod	0.0	0.0		0.0	0.0		3	1.1	
	Pediatric Risk	0.0	0.0		0.0	0.0		6	2.2	
	Unk	0.0	0.0		0.0	0.0		7	2.6	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

]	HIV Diag	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	M	ale	Fer	nale	To	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		7.3	100.0	1.7	100.0	9.0	100.0	3.7	100.0	1.0	100.0	4.7	100.0	186	100.0	82	100.0	268	100.0
Risk	MSM	5.7	77.3	0.0	0.0	5.7	63.0	2.7	72.7	0.0	0.0	2.7	57.1	98	52.7	0	0.0	98	36.6
	IDU	0.3	4.5	0.3	20.0	0.7	7.4	0.3	9.1	0.0	0.0	0.3	7.1	53	28.5	33	40.2	86	32.1
	MSM/IDU	0.3	4.5	0.0	0.0	0.3	3.7	0.3	9.1	0.0	0.0	0.3	7.1	16	8.6	0	0.0	16	6.0
	Heterosexual	1.0	13.6	1.0	60.0	2.0	22.2	0.3	9.1	0.7	66.7	1.0	21.4	7	3.8	38	46.3	45	16.8
	Fem Pres Het Cont	0.0	0.0	0.3	20.0	0.3	3.7	0.0	0.0	0.3	33.3	0.3	7.1	0	0.0	7	8.5	7	2.6
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	0.5	2	2.4	3	1.1
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	4	2.2	2	2.4	6	2.2
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	7	3.8	0	0.0	7	2.6

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		2	
1986		4	4
1987		5	3
1988		6	3
1989		14	8
1990		21	10
1991		18	13
1992		17	21
1993		22	11
1994		8	20
1995		15	9
1996		11	6
1997		14	9
1998		10	4
1999		6	2
2000*	7	13	4
2001	3	11	4
2002	6	5	1
2003	9	11	3
2004	12	8	5
2005	18	11	13
2006	9	7	4
2007	5	7	2
2008	9	8	4
2009	7	2	4
2010	10	4	2
2011	10	8	3

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	d AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		114	100.0	48.7	167	100.0	71.3	281	100.0	119.9	234,287
Gender	Male	65	57.0		106	63.5		171	60.9		
	Female	49	43.0		61	36.5		110	39.1		
Current Age	12 & under	1	0.9		0	0.0		1	0.4		
	13-19	2	1.8		2	1.2		4	1.4		
	20-24	8	7.0		3	1.8		11	3.9		
	25-29	12	10.5		5	3.0		17	6.0		
	30-39	25	21.9		29	17.4		54	19.2		
	40-49	34	29.8		63	37.7		97	34.5		
	50-59	20	17.5		44	26.3		64	22.8		
	60+	12	10.5		21	12.6		33	11.7		
Race/Ethnicity	White	53	46.5		84	50.3		137	48.8		
	Black	26	22.8		49	29.3		75	26.7		
	Hispanic	21	18.4		19	11.4		40	14.2		
	Multi Race	9	7.9		13	7.8		22	7.8		
	Others	5	4.4		2	1.2		7	2.5		
Risk	MSM	43	37.7		61	36.5		104	37.0		
	IDU	9	7.9		29	17.4		38	13.5		
	MSM/IDU	2	1.8		8	4.8		10	3.6		
	Heterosexual	27	23.7		27	16.2		54	19.2		
	Fem Pres Het Cont	23	20.2		24	14.4		47	16.7		
	Blood Prod	0	0.0		1	0.6		1	0.4		
	Pediatric Risk	2	1.8		5	3.0		7	2.5		
	Unk	8	7.0		12	7.2		20	7.1		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		M	ale	Fe	male	To	otal	N	Male	Fe	emale	7	otal	M	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		65	100.0	49	100.0	114	100.0	106	100.0	61	100.0	167	100.0	171	100.0	110	100.0	281	100.0
Risk	MSM	43	66.2	0	0.0	43	37.7	61	57.5	0	0.0	61	36.5	104	60.8	0	0.0	104	37.0
	IDU	5	7.7	4	8.2	9	7.9	18	17.0	11	18.0	29	17.4	23	13.5	15	13.6	38	13.5
	MSM/IDU	2	3.1	0	0.0	2	1.8	8	7.5	0	0.0	8	4.8	10	5.8	0	0.0	10	3.6
	Heterosexual	7	10.8	20	40.8	27	23.7	3	2.8	24	39.3	27	16.2	10	5.8	44	40.0	54	19.2
	Fem Pres Het Cont	0	0.0	23	46.9	23	20.2	0	0.0	24	39.3	24	14.4	0	0.0	47	42.7	47	16.7
	Blood Prod	0	0.0	0	0.0	0	0.0	1	0.9	0	0.0	1	0.6	1	0.6	0	0.0	1	0.4
	Pediatric Risk	0	0.0	2	4.1	2	1.8	3	2.8	2	3.3	5	3.0	3	1.8	4	3.6	7	2.5
	Unk	8	12.3	0	0.0	8	7.0	12	11.3	0	0.0	12	7.2	20	11.7	0	0.0	20	7.1

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		11.7	100.0	5.0	7.0	100.0	3.0	395	100.0	234,287
Gender	Male	6.0	51.4		3.3	47.6		276	69.9	
	Female	5.7	48.6		3.7	52.4		119	30.1	
Age at Diagnosis	12 & under	0.3	2.9		0.0	0.0		2	0.5	
	13-19	0.7	5.7		0.3	4.8		6	1.5	
	20-24	4.0	34.3		0.7	9.5		17	4.3	
	25-29	1.0	8.6		1.3	19.0		51	12.9	
	30-39	2.0	17.1		1.7	23.8		162	41.0	
	40-49	2.7	22.9		1.7	23.8		112	28.4	
	50-59	0.3	2.9		1.0	14.3		34	8.6	
	60+	0.7	5.7		0.3	4.8		11	2.8	
Race/Ethnicity	White	7.0	60.0		3.3	47.6		189	47.8	
	Black	2.3	20.0		1.7	23.8		129	32.7	
	Hispanic	1.3	11.4		1.0	14.3		53	13.4	
	Multi Race	1.0	8.6		0.7	9.5		21	5.3	
	Others	0.0	0.0		0.3	4.8		3	0.8	
Risk	MSM	3.7	31.4		2.0	28.6		129	32.7	
	IDU	1.0	8.6		0.3	4.8		110	27.8	
	MSM/IDU	0.3	2.9		0.0	0.0		25	6.3	
	Heterosexual	2.3	20.0		2.0	28.6		57	14.4	
	Fem Pres Het Cont	3.0	25.7		1.3	19.0		35	8.9	
	Blood Prod	0.0	0.0		0.0	0.0		5	1.3	
	Pediatric Risk	0.3	2.9		0.3	4.8		8	2.0	
	Unk	1.0	8.6		1.0	14.3		26	6.6	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS C	Cases	
		M	ale	Fer	nale	To	otal	M	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		6.0	100.0	5.7	100.0	11.7	100.0	3.3	100.0	3.7	100.0	7.0	100.0	276	100.0	119	100.0	395	100.0
Risk	MSM	3.7	61.1	0.0	0.0	3.7	31.4	2.0	60.0	0.0	0.0	2.0	28.6	129	46.7	0	0.0	129	32.7
	IDU	0.3	5.6	0.7	11.8	1.0	8.6	0.0	0.0	0.3	9.1	0.3	4.8	75	27.2	35	29.4	110	27.8
	MSM/IDU	0.3	5.6	0.0	0.0	0.3	2.9	0.0	0.0	0.0	0.0	0.0	0.0	25	9.1	0	0.0	25	6.3
	Heterosexual	0.7	11.1	1.7	29.4	2.3	20.0	0.3	10.0	1.7	45.5	2.0	28.6	12	4.3	45	37.8	57	14.4
	Fem Pres Het Cont	0.0	0.0	3.0	52.9	3.0	25.7	0.0	0.0	1.3	36.4	1.3	19.0	0	0.0	35	29.4	35	8.9
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	5	1.8	0	0.0	5	1.3
	Pediatric Risk	0.0	0.0	0.3	5.9	0.3	2.9	0.0	0.0	0.3	9.1	0.3	4.8	4	1.4	4	3.4	8	2.0
	Unk	1.0	16.7	0.0	0.0	1.0	8.6	1.0	30.0	0.0	0.0	1.0	14.3	26	9.4	0	0.0	26	6.6

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		5	3
1986		6	3
1987	•	6	4
1988		10	5
1989	٠	12	9
1990	٠	24	9
1991		21	11
1992	٠	34	14
1993		39	28
1994		20	20
1995	٠	23	18
1996	٠	21	21
1997	٠	17	10
1998		14	6
1999		19	7
2000*	9	15	9
2001	9	5	5
2002	14	10	3
2003	14	15	6
2004	16	14	5
2005	11	13	8
2006	9	7	1
2007	12	11	9
2008	9	13	5
2009	13	12	4
2010	13	4	4
2011	9	5	1

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		356	100.0	76.2	558	100.0	119.5	914	100.0	195.7	466,960
Gender	Male	225	63.2		384	68.8		609	66.6		
	Female	131	36.8		174	31.2		305	33.4		
Current Age	12 & under	2	0.6		0	0.0		2	0.2		
	13-19	13	3.7		0	0.0		13	1.4		
	20-24	28	7.9		7	1.3		35	3.8		
	25-29	35	9.8		15	2.7		50	5.5		
	30-39	70	19.7		63	11.3		133	14.6		
	40-49	106	29.8		202	36.2		308	33.7		
	50-59	77	21.6		206	36.9		283	31.0		
	60+	25	7.0		65	11.6		90	9.8		
Race/Ethnicity	White	154	43.3		252	45.2		406	44.4		
	Black	135	37.9		194	34.8		329	36.0		
	Hispanic	34	9.6		56	10.0		90	9.8		
	Multi Race	28	7.9		47	8.4		75	8.2		
	Others	5	1.4		9	1.6		14	1.5		
Risk	MSM	148	41.6		223	40.0		371	40.6		
	IDU	39	11.0		85	15.2		124	13.6		
	MSM/IDU	3	0.8		27	4.8		30	3.3		
	Heterosexual	105	29.5		150	26.9		255	27.9		
	Fem Pres Het Cont	23	6.5		20	3.6		43	4.7		
	Blood Prod	0	0.0		5	0.9		5	0.5		
	Pediatric Risk	10	2.8		5	0.9		15	1.6		
	Unk	28	7.9		43	7.7		71	7.8		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	Male	Fe	male	Т	otal	M	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		225	100.0	131	100.0	356	100.0	384	100.0	174	100.0	558	100.0	609	100.0	305	100.0	914	100.0
Risk	MSM	148	65.8	0	0.0	148	41.6	223	58.1	0	0.0	223	40.0	371	60.9	0	0.0	371	40.6
	IDU	22	9.8	17	13.0	39	11.0	49	12.8	36	20.7	85	15.2	71	11.7	53	17.4	124	13.6
	MSM/IDU	3	1.3	0	0.0	3	0.8	27	7.0	0	0.0	27	4.8	30	4.9	0	0.0	30	3.3
	Heterosexual	18	8.0	87	66.4	105	29.5	37	9.6	113	64.9	150	26.9	55	9.0	200	65.6	255	27.9
	Fem Pres Het Cont	0	0.0	23	17.6	23	6.5	0	0.0	20	11.5	20	3.6	0	0.0	43	14.1	43	4.7
	Blood Prod	0	0.0	0	0.0	0	0.0	5	1.3	0	0.0	5	0.9	5	0.8	0	0.0	5	0.5
	Pediatric Risk	6	2.7	4	3.1	10	2.8	2	0.5	3	1.7	5	0.9	8	1.3	7	2.3	15	1.6
	Unk	28	12.4	0	0.0	28	7.9	41	10.7	2	1.1	43	7.7	69	11.3	2	0.7	71	7.8

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		38.7	100.0	8.3	25.7	100.0	5.5	1,289	100.0	466,960
Gender	Male	28.7	74.1		18.3	71.4		980	76.0	
	Female	10.0	25.9		7.3	28.6		309	24.0	
Age at Diagnosis	12 & under	0.3	0.9		0.0	0.0		8	0.6	
	13-19	3.0	7.8		0.3	1.3		13	1.0	
	20-24	6.0	15.5		0.7	2.6		54	4.2	
	25-29	5.7	14.7		2.7	10.4		153	11.9	
	30-39	6.7	17.2		4.0	15.6		509	39.5	
	40-49	8.7	22.4		10.7	41.6		388	30.1	
	50-59	6.0	15.5		6.0	23.4		130	10.1	
	60+	2.3	6.0		1.3	5.2		34	2.6	
Race/Ethnicity	White	14.0	36.2		9.0	35.1		692	53.7	
	Black	16.0	41.4		11.0	42.9		426	33.0	
	Hispanic	5.0	12.9		2.7	10.4		100	7.8	
	Multi Race	2.3	6.0		2.0	7.8		56	4.3	
	Others	1.3	3.4		1.0	3.9		15	1.2	
Risk	MSM	18.7	48.3		10.0	39.0		556	43.1	
	IDU	1.3	3.4		2.0	7.8		271	21.0	
	MSM/IDU	0.7	1.7		0.7	2.6		77	6.0	
	Heterosexual	7.7	19.8		6.7	26.0		239	18.5	
	Fem Pres Het Cont	3.3	8.6		1.0	3.9		34	2.6	
	Blood Prod	0.0	0.0		0.0	0.0		26	2.0	
	Pediatric Risk	0.3	0.9		0.3	1.3		11	0.9	
	Unk	6.7	17.2		5.0	19.5		75	5.8	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Diaş	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	To	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		28.7	100.0	10.0	100.0	38.7	100.0	18.3	100.0	7.3	100.0	25.7	100.0	980	100.0	309	100.0	1,289	100.0
Risk	MSM	18.7	65.1	0.0	0.0	18.7	48.3	10.0	54.5	0.0	0.0	10.0	39.0	556	56.7	0	0.0	556	43.1
	IDU	1.3	4.7	0.0	0.0	1.3	3.4	1.7	9.1	0.3	4.5	2.0	7.8	180	18.4	91	29.4	271	21.0
	MSM/IDU	0.7	2.3	0.0	0.0	0.7	1.7	0.7	3.6	0.0	0.0	0.7	2.6	77	7.9	0	0.0	77	6.0
	Heterosexual	2.0	7.0	5.7	56.7	7.7	19.8	1.7	9.1	5.0	68.2	6.7	26.0	67	6.8	172	55.7	239	18.5
	Fem Pres Het Cont	0.0	0.0	3.3	33.3	3.3	8.6	0.0	0.0	1.0	13.6	1.0	3.9	0	0.0	34	11.0	34	2.6
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	21	2.1	5	1.6	26	2.0
	Pediatric Risk	0.0	0.0	0.3	3.3	0.3	0.9	0.0	0.0	0.3	4.5	0.3	1.3	6	0.6	5	1.6	11	0.9
	Unk	6.0	20.9	0.7	6.7	6.7	17.2	4.3	23.6	0.7	9.1	5.0	19.5	73	7.4	2	0.6	75	5.8

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		34	18
1986		12	16
1987		24	12
1988		33	24
1989		42	24
1990		49	21
1991		58	34
1992		119	53
1993		110	56
1994		77	80
1995		80	65
1996		84	57
1997		54	19
1998		32	20
1999		44	20
2000*	25	75	24
2001	46	38	17
2002	44	39	16
2003	55	41	14
2004	44	31	13
2005	42	38	12
2006	32	27	19
2007	35	36	23
2008	43	35	25
2009	33	26	18
2010	38	20	21
2011	45	31	15

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	1 AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		41	100.0	37.8	39	100.0	35.9	80	100.0	73.7	108,525
Gender	Male	24	58.5		26	66.7		50	62.5		
	Female	17	41.5		13	33.3		30	37.5		
Current Age	13-19	0	0.0		0	0.0		0	0.0		
	20-24	0	0.0		0	0.0		0	0.0		
	25-29	2	4.9		1	2.6		3	3.8		
	30-39	10	24.4		4	10.3		14	17.5		
	40-49	11	26.8		14	35.9		25	31.3		
	50-59	15	36.6		13	33.3		28	35.0		
	60+	3	7.3		7	17.9		10	12.5		
Race/Ethnicity	White	25	61.0		24	61.5		49	61.3		
	Black	6	14.6		5	12.8		11	13.8		
	Hispanic	9	22.0		9	23.1		18	22.5		
	Multi Race	1	2.4		1	2.6		2	2.5		
Risk	MSM	14	34.1		16	41.0		30	37.5		
	IDU	4	9.8		2	5.1		6	7.5		
	MSM/IDU	0	0.0		2	5.1		2	2.5		
	Heterosexual	9	22.0		9	23.1		18	22.5		
	Fem Pres Het Cont	7	17.1		7	17.9		14	17.5		
	Blood Prod	0	0.0		0	0.0		0	0.0		
	Unk	7	17.1		3	7.7		10	12.5		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		M	ale	Fe	male	To	otal	N	Aale	Fe	male	Т	otal	M	[ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		24	100.0	17	100.0	41	100.0	26	100.0	13	100.0	39	100.0	50	100.0	30	100.0	80	100.0
Risk	MSM	14	58.3	0	0.0	14	34.1	16	61.5	0	0.0	16	41.0	30	60.0	0	0.0	30	37.5
	IDU	2	8.3	2	11.8	4	9.8	2	7.7	0	0.0	2	5.1	4	8.0	2	6.7	6	7.5
	MSM/IDU	0	0.0	0	0.0	0	0.0	2	7.7	0	0.0	2	5.1	2	4.0	0	0.0	2	2.5
	Heterosexual	1	4.2	8	47.1	9	22.0	3	11.5	6	46.2	9	23.1	4	8.0	14	46.7	18	22.5
	Fem Pres Het Cont	0	0.0	7	41.2	7	17.1	0	0.0	7	53.8	7	17.9	0	0.0	14	46.7	14	17.5
	Blood Prod	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Unk	7	29.2	0	0.0	7	17.1	3	11.5	0	0.0	3	7.7	10	20.0	0	0.0	10	12.5

		HIV Diagn	oses (Annua	ıl Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		1.7	100.0	1.5	1.3	100.0	1.2	70	100.0	108,525
Gender	Male	1.7	100.0		1.0	75.0		53	75.7	
	Female	0.0	0.0		0.3	25.0		17	24.3	
Age at Diagnosis	13-19	0.0	0.0		0.0	0.0		1	1.4	
	20-24	0.0	0.0		0.0	0.0		1	1.4	
	25-29	0.0	0.0		0.0	0.0		6	8.6	
	30-39	0.7	40.0		0.7	50.0		33	47.1	
	40-49	0.3	20.0		0.0	0.0		21	30.0	
	50-59	0.7	40.0		0.3	25.0		4	5.7	
	60+	0.0	0.0		0.3	25.0		4	5.7	
Race/Ethnicity	White	1.3	80.0		0.7	50.0		46	65.7	
	Black	0.0	0.0		0.0	0.0		8	11.4	
	Hispanic	0.3	20.0		0.7	50.0		15	21.4	
	Multi Race	0.0	0.0		0.0	0.0		1	1.4	
Risk	MSM	1.7	100.0		1.0	75.0		30	42.9	
	IDU	0.0	0.0		0.0	0.0		12	17.1	
	MSM/IDU	0.0	0.0		0.0	0.0		2	2.9	
	Heterosexual	0.0	0.0		0.3	25.0		11	15.7	
	Fem Pres Het Cont	0.0	0.0		0.0	0.0		7	10.0	
	Blood Prod	0.0	0.0		0.0	0.0		2	2.9	
_	Unk	0.0	0.0		0.0	0.0		6	8.6	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		1	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS C	Cases	
		M	ale	Fei	nale	To	otal	М	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		1.7	100.0	0.0	0.0	1.7	100.0	1.0	100.0	0.3	100.0	1.3	100.0	53	100.0	17	100.0	70	100.0
Risk	MSM	1.7	100.0	0.0	0.0	1.7	100.0	1.0	100.0	0.0	0.0	1.0	75.0	30	56.6	0	0.0	30	42.9
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	10	18.9	2	11.8	12	17.1
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	3.8	0	0.0	2	2.9
	Heterosexual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	100.0	0.3	25.0	3	5.7	8	47.1	11	15.7
	Fem Pres Het Cont	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	7	41.2	7	10.0
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	3.8	0	0.0	2	2.9
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	6	11.3	0	0.0	6	8.6

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
1986		1	
1987		1	1
1988		2	1
1989		2	1
1990		2	2
1991		1	
1992		4	2
1993		6	6
1994		6	2
1995		6	3
1996		2	1
1997		1	2
1998		7	1
1999		2	
2000*	1		
2001	4	4	1
2002	6	4	1
2003	3	3	
2004	2	2	
2005	3	4	2
2006	2	4	1
2007	5	1	
2008	3	1	
2009	1	2	2
2010	3	1	2
2011	1	1	

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	d AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		312	100.0	83.2	503	100.0	134.2	815	100.0	217.4	374,872
Gender	Male	192	61.5		341	67.8		533	65.4		
	Female	120	38.5		162	32.2		282	34.6		
Current Age	12 & under	3	1.0		0	0.0		3	0.4		
	13-19	6	1.9		3	0.6		9	1.1		
	20-24	20	6.4		14	2.8		34	4.2		
	25-29	21	6.7		8	1.6		29	3.6		
	30-39	50	16.0		41	8.2		91	11.2		
	40-49	88	28.2		150	29.8		238	29.2		
	50-59	97	31.1		206	41.0		303	37.2		
	60+	26	8.3		81	16.1		107	13.1		
	Unk	1	0.3		0	0.0		1	0.1		
Race/Ethnicity	White	113	36.2		130	25.8		243	29.8		
	Black	83	26.6		167	33.2		250	30.7		
	Hispanic	86	27.6		152	30.2		238	29.2		
	Multi Race	27	8.7		52	10.3		79	9.7		
	Others	3	1.0		2	0.4		5	0.6		
Risk	MSM	96	30.8		98	19.5		194	23.8		
	IDU	35	11.2		142	28.2		177	21.7		
	MSM/IDU	8	2.6		24	4.8		32	3.9		
	Heterosexual	108	34.6		179	35.6		287	35.2		
	Fem Pres Het Cont	24	7.7		14	2.8		38	4.7		
	Blood Prod	1	0.3		4	0.8		5	0.6		
	Pediatric Risk	13	4.2		14	2.8		27	3.3		
	Unk	27	8.7		28	5.6		55	6.7		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	[ale	Fe	male	To	otal	N	Male	Fe	male	1	otal	M	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		192	100.0	120	100.0	312	100.0	341	100.0	162	100.0	503	100.0	533	100.0	282	100.0	815	100.0
Risk	MSM	96	50.0	0	0.0	96	30.8	98	28.7	0	0.0	98	19.5	194	36.4	0	0.0	194	23.8
	IDU	24	12.5	11	9.2	35	11.2	107	31.4	35	21.6	142	28.2	131	24.6	46	16.3	177	21.7
	MSM/IDU	8	4.2	0	0.0	8	2.6	24	7.0	0	0.0	24	4.8	32	6.0	0	0.0	32	3.9
	Heterosexual	28	14.6	80	66.7	108	34.6	75	22.0	104	64.2	179	35.6	103	19.3	184	65.2	287	35.2
	Fem Pres Het Cont	0	0.0	24	20.0	24	7.7	0	0.0	14	8.6	14	2.8	0	0.0	38	13.5	38	4.7
	Blood Prod	1	0.5	0	0.0	1	0.3	2	0.6	2	1.2	4	0.8	3	0.6	2	0.7	5	0.6
	Pediatric Risk	8	4.2	5	4.2	13	4.2	7	2.1	7	4.3	14	2.8	15	2.8	12	4.3	27	3.3
	Unk	27	14.1	0	0.0	27	8.7	28	8.2	0	0.0	28	5.6	55	10.3	0	0.0	55	6.7

		HIV Diagn	oses (Annua	al Average)	AIDS Diagr	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		25.7	100.0	6.8	22.0	100.0	5.9	1,294	100.0	374,872
Gender	Male	18.0	70.1		13.7	62.1		954	73.7	
	Female	7.7	29.9		8.3	37.9		340	26.3	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		16	1.2	
	13-19	1.0	3.9		0.7	3.0		11	0.9	
	20-24	3.7	14.3		1.0	4.5		30	2.3	
	25-29	3.0	11.7		2.0	9.1		95	7.3	
	30-39	5.0	19.5		6.3	28.8		481	37.2	
	40-49	6.7	26.0		6.7	30.3		431	33.3	
	50-59	4.0	15.6		4.3	19.7		174	13.4	
	60+	2.3	9.1		1.0	4.5		56	4.3	
	Unk	0.0	0.0		0.0	0.0		0	0.0	
Race/Ethnicity	White	4.7	18.2		4.7	21.2		415	32.1	
	Black	10.7	41.6		7.7	34.8		444	34.3	
	Hispanic	8.7	33.8		8.7	39.4		356	27.5	
	Multi Race	1.3	5.2		0.7	3.0		77	6.0	
	Others	0.3	1.3		0.3	1.5		2	0.2	
Risk	MSM	11.7	45.5		6.7	30.3		255	19.7	
	IDU	3.0	11.7		5.0	22.7		539	41.7	
	MSM/IDU	0.0	0.0		0.0	0.0		63	4.9	
	Heterosexual	9.7	37.7		8.7	39.4		278	21.5	
	Fem Pres Het Cont	0.3	1.3		0.0	0.0		31	2.4	
	Blood Prod	0.0	0.0		0.0	0.0		21	1.6	
	Pediatric Risk	0.0	0.0		0.7	3.0		22	1.7	
	Unk	1.0	3.9		1.0	4.5		85	6.6	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	То	otal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		18.0	100.0	7.7	100.0	25.7	100.0	13.7	100.0	8.3	100.0	22.0	100.0	954	100.0	340	100.0	1,294	100.0
Risk	MSM	11.7	64.8	0.0	0.0	11.7	45.5	6.7	48.8	0.0	0.0	6.7	30.3	255	26.7	0	0.0	255	19.7
	IDU	2.0	11.1	1.0	13.0	3.0	11.7	2.7	19.5	2.3	28.0	5.0	22.7	423	44.3	116	34.1	539	41.7
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	63	6.6	0	0.0	63	4.9
	Heterosexual	3.3	18.5	6.3	82.6	9.7	37.7	2.7	19.5	6.0	72.0	8.7	39.4	102	10.7	176	51.8	278	21.5
	Fem Pres Het Cont	0.0	0.0	0.3	4.3	0.3	1.3	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	31	9.1	31	2.4
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	15	1.6	6	1.8	21	1.6
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.7	4.9	0.0	0.0	0.7	3.0	11	1.2	11	3.2	22	1.7
	Unk	1.0	5.6	0.0	0.0	1.0	3.9	1.0	7.3	0.0	0.0	1.0	4.5	85	8.9	0	0.0	85	6.6

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		15	8
1986		16	11
1987		29	18
1988		35	20
1989		52	38
1990		71	30
1991	•	87	57
1992		89	59
1993		93	68
1994		110	58
1995	•	77	84
1996		53	44
1997		59	34
1998	•	36	23
1999		47	24
2000*	38	47	30
2001	39	41	17
2002	32	43	17
2003	42	57	23
2004	34	43	20
2005	43	43	24
2006	39	27	19
2007	35	28	18
2008	28	30	23
2009	32	28	13
2010	24	18	11
2011	21	20	6

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	1 AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		15	100.0	35.2	26	100.0	61.0	41	100.0	96.2	42,622
Gender	Male	9	60.0		16	61.5		25	61.0		
	Female	6	40.0		10	38.5		16	39.0		
Current Age	13-19	0	0.0		1	3.8		1	2.4		
	20-24	1	6.7		1	3.8		2	4.9		
	25-29	2	13.3		2	7.7		4	9.8		
	30-39	5	33.3		4	15.4		9	22.0		
	40-49	4	26.7		9	34.6		13	31.7		
	50-59	1	6.7		8	30.8		9	22.0		
	60+	2	13.3		1	3.8		3	7.3		
Race/Ethnicity	White	9	60.0		15	57.7		24	58.5		
	Black	1	6.7		8	30.8		9	22.0		
	Hispanic	5	33.3		3	11.5		8	19.5		
	Multi Race	0	0.0		0	0.0		0	0.0		
Risk	MSM	7	46.7		10	38.5		17	41.5		
	IDU	3	20.0		3	11.5		6	14.6		
	MSM/IDU	0	0.0		2	7.7		2	4.9		
	Heterosexual	1	6.7		2	7.7		3	7.3		
	Fem Pres Het Cont	4	26.7		7	26.9		11	26.8		
	Blood Prod	0	0.0		0	0.0		0	0.0		
	Unk	0	0.0		2	7.7		2	4.9		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	/Iale	Fe	male	Т	otal	M	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		9	100.0	6	100.0	15	100.0	16	100.0	10	100.0	26	100.0	25	100.0	16	100.0	41	100.0
Risk	MSM	7	77.8	0	0.0	7	46.7	10	62.5	0	0.0	10	38.5	17	68.0	0	0.0	17	41.5
	IDU	2	22.2	1	16.7	3	20.0	2	12.5	1	10.0	3	11.5	4	16.0	2	12.5	6	14.6
	MSM/IDU	0	0.0	0	0.0	0	0.0	2	12.5	0	0.0	2	7.7	2	8.0	0	0.0	2	4.9
	Heterosexual	0	0.0	1	16.7	1	6.7	0	0.0	2	20.0	2	7.7	0	0.0	3	18.8	3	7.3
	Fem Pres Het Cont	0	0.0	4	66.7	4	26.7	0	0.0	7	70.0	7	26.9	0	0.0	11	68.8	11	26.8
	Blood Prod	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Unk	0	0.0	0	0.0	0	0.0	2	12.5	0	0.0	2	7.7	2	8.0	0	0.0	2	4.9

		HIV Diagn	oses (Annua	ıl Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		2.3	100.0	5.5	2.0	100.0	4.7	51	100.0	42,622
Gender	Male	1.3	57.1		1.0	50.0		37	72.5	
	Female	1.0	42.9		1.0	50.0		14	27.5	
Age at Diagnosis	13-19	0.0	0.0		0.3	16.7		2	3.9	
	20-24	0.3	14.3		0.3	16.7		4	7.8	
	25-29	0.7	28.6		0.0	0.0		6	11.8	
	30-39	1.0	42.9		0.7	33.3		19	37.3	
	40-49	0.0	0.0		0.3	16.7		13	25.5	
	50-59	0.3	14.3		0.3	16.7		4	7.8	
	60+	0.0	0.0		0.0	0.0		3	5.9	
Race/Ethnicity	White	1.7	71.4		0.7	33.3		28	54.9	
	Black	0.7	28.6		1.0	50.0		17	33.3	
	Hispanic	0.0	0.0		0.3	16.7		5	9.8	
	Multi Race	0.0	0.0		0.0	0.0		1	2.0	
Risk	MSM	1.0	42.9		0.3	16.7		18	35.3	
	IDU	0.0	0.0		0.0	0.0		8	15.7	
	MSM/IDU	0.3	14.3		0.3	16.7		4	7.8	
	Heterosexual	0.0	0.0		0.3	16.7		3	5.9	
	Fem Pres Het Cont	1.0	42.9		0.7	33.3		9	17.6	
	Blood Prod	0.0	0.0		0.0	0.0		1	2.0	
	Unk	0.0	0.0		0.3	16.7		8	15.7	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		1	HIV Dia	gnoses (Annual A	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS C	Cases	
		M	ale	Fei	nale	To	otal	M	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		1.3	100.0	1.0	100.0	2.3	100.0	1.0	100.0	1.0	100.0	2.0	100.0	37	100.0	14	100.0	51	100.0
Risk	MSM	1.0	75.0	0.0	0.0	1.0	42.9	0.3	33.3	0.0	0.0	0.3	16.7	18	48.6	0	0.0	18	35.3
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	6	16.2	2	14.3	8	15.7
	MSM/IDU	0.3	25.0	0.0	0.0	0.3	14.3	0.3	33.3	0.0	0.0	0.3	16.7	4	10.8	0	0.0	4	7.8
	Heterosexual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	33.3	0.3	16.7	0	0.0	3	21.4	3	5.9
	Fem Pres Het Cont	0.0	0.0	1.0	100.0	1.0	42.9	0.0	0.0	0.7	66.7	0.7	33.3	0	0.0	9	64.3	9	17.6
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	2.7	0	0.0	1	2.0
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.3	33.3	0.0	0.0	0.3	16.7	8	21.6	0	0.0	8	15.7

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
1990		4	
1991		3	2
1992		2	4
1993			2
1994		5	
1995		4	3
1996		3	2
1997		4	1
1998		2	1
1999			2
2000*	2	2	
2001	3		
2002	2	3	1
2003	2	3	
2004	1	2	2
2005	2	2	3
2006	2	1	1
2007	2	3	
2008	1	2	
2009	1	1	1
2010	1	2	1
2011	5	3	

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		37	100.0	30.3	52	100.0	42.5	89	100.0	72.8	122,228
Gender	Male	28	75.7		46	88.5		74	83.1		
	Female	9	24.3		6	11.5		15	16.9		
Current Age	12 & under	0	0.0		0	0.0		0	0.0		
	13-19	0	0.0		1	1.9		1	1.1		
	20-24	1	2.7		2	3.8		3	3.4		
	25-29	5	13.5		0	0.0		5	5.6		
	30-39	9	24.3		12	23.1		21	23.6		
	40-49	11	29.7		21	40.4		32	36.0		
	50-59	8	21.6		11	21.2		19	21.3		
	60+	3	8.1		5	9.6		8	9.0		
Race/Ethnicity	White	33	89.2		41	78.8		74	83.1		
	Black	1	2.7		5	9.6		6	6.7		
	Hispanic	1	2.7		4	7.7		5	5.6		
	Multi Race	2	5.4		2	3.8		4	4.5		
Risk	MSM	22	59.5		31	59.6		53	59.6		
	IDU	0	0.0		4	7.7		4	4.5		
	MSM/IDU	2	5.4		2	3.8		4	4.5		
	Heterosexual	8	21.6		8	15.4		16	18.0		
	Fem Pres Het Cont	2	5.4		1	1.9		3	3.4		
	Blood Prod	0	0.0		0	0.0		0	0.0		
	Pediatric Risk	0	0.0		0	0.0		0	0.0		
	Unk	3	8.1		6	11.5		9	10.1		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	Male	Fe	emale	Т	otal	M	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		28	100.0	9	100.0	37	100.0	46	100.0	6	100.0	52	100.0	74	100.0	15	100.0	89	100.0
Risk	MSM	22	78.6	0	0.0	22	59.5	31	67.4	0	0.0	31	59.6	53	71.6	0	0.0	53	59.6
	IDU	0	0.0	0	0.0	0	0.0	4	8.7	0	0.0	4	7.7	4	5.4	0	0.0	4	4.5
	MSM/IDU	2	7.1	0	0.0	2	5.4	2	4.3	0	0.0	2	3.8	4	5.4	0	0.0	4	4.5
	Heterosexual	1	3.6	7	77.8	8	21.6	3	6.5	5	83.3	8	15.4	4	5.4	12	80.0	16	18.0
	Fem Pres Het Cont	0	0.0	2	22.2	2	5.4	0	0.0	1	16.7	1	1.9	0	0.0	3	20.0	3	3.4
	Blood Prod	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Pediatric Risk	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Unk	3	10.7	0	0.0	3	8.1	6	13.0	0	0.0	6	11.5	9	12.2	0	0.0	9	10.1

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		5.0	100.0	4.1	4.3	100.0	3.5	105	100.0	122,228
Gender	Male	4.7	93.3		3.7	84.6		90	85.7	
	Female	0.3	6.7		0.7	15.4		15	14.3	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		1	1.0	
	13-19	0.0	0.0		0.3	7.7		2	1.9	
	20-24	0.3	6.7		0.7	15.4		6	5.7	
	25-29	1.3	26.7		0.7	15.4		10	9.5	
	30-39	1.0	20.0		1.3	30.8		49	46.7	
	40-49	1.0	20.0		0.7	15.4		27	25.7	
	50-59	1.0	20.0		0.3	7.7		7	6.7	
	60+	0.3	6.7		0.3	7.7		3	2.9	
Race/Ethnicity	White	4.3	86.7		3.3	76.9		86	81.9	
	Black	0.0	0.0		0.0	0.0		11	10.5	
	Hispanic	0.3	6.7		0.7	15.4		4	3.8	
	Multi Race	0.3	6.7		0.3	7.7		4	3.8	
Risk	MSM	4.0	80.0		3.0	69.2		62	59.0	
	IDU	0.0	0.0		0.0	0.0		10	9.5	
	MSM/IDU	0.0	0.0		0.3	7.7		4	3.8	
	Heterosexual	0.3	6.7		0.3	7.7		17	16.2	
	Fem Pres Het Cont	0.0	0.0		0.3	7.7		1	1.0	
	Blood Prod	0.0	0.0		0.0	0.0		3	2.9	
	Pediatric Risk	0.0	0.0		0.0	0.0		2	1.9	
	Unk	0.7	13.3		0.3	7.7		6	5.7	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Diag	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	M	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		4.7	100.0	0.3	100.0	5.0	100.0	3.7	100.0	0.7	100.0	4.3	100.0	90	100.0	15	100.0	105	100.0
Risk	MSM	4.0	85.7	0.0	0.0	4.0	80.0	3.0	81.8	0.0	0.0	3.0	69.2	62	68.9	0	0.0	62	59.0
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	9	10.0	1	6.7	10	9.5
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.3	9.1	0.0	0.0	0.3	7.7	4	4.4	0	0.0	4	3.8
	Heterosexual	0.0	0.0	0.3	100.0	0.3	6.7	0.0	0.0	0.3	50.0	0.3	7.7	4	4.4	13	86.7	17	16.2
	Fem Pres Het Cont	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	50.0	0.3	7.7	0	0.0	1	6.7	1	1.0
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3	3.3	0	0.0	3	2.9
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	2.2	0	0.0	2	1.9
	Unk	0.7	14.3	0.0	0.0	0.7	13.3	0.3	9.1	0.0	0.0	0.3	7.7	6	6.7	0	0.0	6	5.7

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		2	
1986		2	2
1987			1
1988		4	
1989		8	6
1990		2	3
1991	•	4	4
1992		9	2
1993		5	7
1994		3	2
1995		7	4
1996		4	4
1997		6	
1998	•	2	
1999		6	1
2000*	1	2	1
2001	5	1	1
2002	11	8	
2003	3	1	4
2004	3	4	
2005	3	7	4
2006	3	4	1
2007	5	1	1
2008	4		
2009	7	3	3
2010	5	7	2
2011	3	3	1

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		23	100.0	37.1	34	100.0	54.9	57	100.0	92.1	61,917
Gender	Male	17	73.9		21	61.8		38	66.7		
	Female	6	26.1		13	38.2		19	33.3		
Current Age	13-19	1	4.3		0	0.0		1	1.8		
	20-24	1	4.3		0	0.0		1	1.8		
	25-29	2	8.7		1	2.9		3	5.3		
	30-39	4	17.4		6	17.6		10	17.5		
	40-49	6	26.1		7	20.6		13	22.8		
	50-59	7	30.4		11	32.4		18	31.6		
	60+	2	8.7		9	26.5		11	19.3		
Race/Ethnicity	White	13	56.5		23	67.6		36	63.2		
	Black	5	21.7		4	11.8		9	15.8		
	Hispanic	3	13.0		4	11.8		7	12.3		
	Multi Race	2	8.7		3	8.8		5	8.8		
Risk	MSM	9	39.1		13	38.2		22	38.6		
	IDU	2	8.7		3	8.8		5	8.8		
	MSM/IDU	0	0.0		2	5.9		2	3.5		
	Heterosexual	4	17.4		8	23.5		12	21.1		
	Fem Pres Het Cont	3	13.0		4	11.8		7	12.3		
	Blood Prod	0	0.0		0	0.0		0	0.0		
	Unk	5	21.7		4	11.8		9	15.8		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	Male	Fe	male	Т	otal	М	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		17	100.0	6	100.0	23	100.0	21	100.0	13	100.0	34	100.0	38	100.0	19	100.0	57	100.0
Risk	MSM	9	52.9	0	0.0	9	39.1	13	61.9	0	0.0	13	38.2	22	57.9	0	0.0	22	38.6
	IDU	1	5.9	1	16.7	2	8.7	1	4.8	2	15.4	3	8.8	2	5.3	3	15.8	5	8.8
	MSM/IDU	0	0.0	0	0.0	0	0.0	2	9.5	0	0.0	2	5.9	2	5.3	0	0.0	2	3.5
	Heterosexual	2	11.8	2	33.3	4	17.4	1	4.8	7	53.8	8	23.5	3	7.9	9	47.4	12	21.1
	Fem Pres Het Cont	0	0.0	3	50.0	3	13.0	0	0.0	4	30.8	4	11.8	0	0.0	7	36.8	7	12.3
	Blood Prod	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Unk	5	29.4	0	0.0	5	21.7	4	19.0	0	0.0	4	11.8	9	23.7	0	0.0	9	15.8

		HIV Diagn	oses (Annua	ıl Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		1.7	100.0	2.7	1.3	100.0	2.2	82	100.0	61,917
Gender	Male	1.3	80.0		1.0	75.0		60	73.2	
	Female	0.3	20.0		0.3	25.0		22	26.8	
Age at Diagnosis	13-19	0.3	20.0		0.0	0.0		2	2.4	
	20-24	0.3	20.0		0.0	0.0		1	1.2	
	25-29	0.0	0.0		0.0	0.0		12	14.6	
	30-39	0.0	0.0		0.0	0.0		29	35.4	
	40-49	0.7	40.0		0.3	25.0		25	30.5	
	50-59	0.3	20.0		0.7	50.0		10	12.2	
	60+	0.0	0.0		0.3	25.0		3	3.7	
Race/Ethnicity	White	1.0	60.0		1.3	100.0		66	80.5	
	Black	0.7	40.0		0.0	0.0		9	11.0	
	Hispanic	0.0	0.0		0.0	0.0		4	4.9	
	Multi Race	0.0	0.0		0.0	0.0		3	3.7	
Risk	MSM	0.3	20.0		1.0	75.0		34	41.5	
	IDU	0.0	0.0		0.0	0.0		10	12.2	
	MSM/IDU	0.0	0.0		0.0	0.0		5	6.1	
	Heterosexual	0.7	40.0		0.3	25.0		16	19.5	
	Fem Pres Het Cont	0.0	0.0		0.0	0.0		5	6.1	
	Blood Prod	0.0	0.0		0.0	0.0		3	3.7	
	Unk	0.7	40.0		0.0	0.0		9	11.0	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

]	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		М	ale	Fei	nale	To	otal	М	ale	Fer	nale	To	otal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		1.3	100.0	0.3	100.0	1.7	100.0	1.0	100.0	0.3	100.0	1.3	100.0	60	100.0	22	100.0	82	100.0
Risk	MSM	0.3	25.0	0.0	0.0	0.3	20.0	1.0	100.0	0.0	0.0	1.0	75.0	34	56.7	0	0.0	34	41.5
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	6	10.0	4	18.2	10	12.2
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	5	8.3	0	0.0	5	6.1
	Heterosexual	0.3	25.0	0.3	100.0	0.7	40.0	0.0	0.0	0.3	100.0	0.3	25.0	3	5.0	13	59.1	16	19.5
	Fem Pres Het Cont	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	5	22.7	5	6.1
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3	5.0	0	0.0	3	3.7
	Unk	0.7	50.0	0.0	0.0	0.7	40.0	0.0	0.0	0.0	0.0	0.0	0.0	9	15.0	0	0.0	9	11.0

¹Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

² Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		1	1
1987		1	
1988		3	2
1989		10	3
1990		2	4
1991		4	4
1992		5	3
1993		5	
1994		2	4
1995		8	6
1996		6	4
1997		2	
1998		1	
1999		4	2
2000*	5	5	2
2001	4		
2002	5	2	1
2003	2	5	
2004	2	2	2
2005	2	5	1
2006	3	3	5
2007	3	1	1
2008	1	1	3
2009		2	
2010	4	1	
2011	1	1	2

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		41	100.0	41.0	89	100.0	89.1	130	100.0	130.1	99,933
Gender	Male	28	68.3		60	67.4		88	67.7		
	Female	13	31.7		29	32.6		42	32.3		
Current Age	12 & under	0	0.0		0	0.0		0	0.0		
	13-19	0	0.0		1	1.1		1	0.8		
	20-24	1	2.4		1	1.1		2	1.5		
	25-29	1	2.4		0	0.0		1	0.8		
	30-39	9	22.0		11	12.4		20	15.4		
	40-49	17	41.5		25	28.1		42	32.3		
	50-59	10	24.4		37	41.6		47	36.2		
	60+	3	7.3		14	15.7		17	13.1		
Race/Ethnicity	White	19	46.3		49	55.1		68	52.3		
	Black	8	19.5		14	15.7		22	16.9		
	Hispanic	12	29.3		23	25.8		35	26.9		
	Multi Race	2	4.9		3	3.4		5	3.8		
Risk	MSM	15	36.6		35	39.3		50	38.5		
	IDU	9	22.0		19	21.3		28	21.5		
	MSM/IDU	2	4.9		2	2.2		4	3.1		
	Heterosexual	8	19.5		24	27.0		32	24.6		
	Fem Pres Het Cont	1	2.4		3	3.4		4	3.1		
	Blood Prod	0	0.0		0	0.0		0	0.0		
	Pediatric Risk	0	0.0		2	2.2		2	1.5		
	Unk	6	14.6		4	4.5		10	7.7		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	Male	Fe	male	7	otal	M	Iale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		28	100.0	13	100.0	41	100.0	60	100.0	29	100.0	89	100.0	88	100.0	42	100.0	130	100.0
Risk	MSM	15	53.6	0	0.0	15	36.6	35	58.3	0	0.0	35	39.3	50	56.8	0	0.0	50	38.5
	IDU	2	7.1	7	53.8	9	22.0	7	11.7	12	41.4	19	21.3	9	10.2	19	45.2	28	21.5
	MSM/IDU	2	7.1	0	0.0	2	4.9	2	3.3	0	0.0	2	2.2	4	4.5	0	0.0	4	3.1
	Heterosexual	3	10.7	5	38.5	8	19.5	11	18.3	13	44.8	24	27.0	14	15.9	18	42.9	32	24.6
	Fem Pres Het Cont	0	0.0	1	7.7	1	2.4	0	0.0	3	10.3	3	3.4	0	0.0	4	9.5	4	3.1
	Blood Prod	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Pediatric Risk	0	0.0	0	0.0	0	0.0	1	1.7	1	3.4	2	2.2	1	1.1	1	2.4	2	1.5
	Unk	6	21.4	0	0.0	6	14.6	4	6.7	0	0.0	4	4.5	10	11.4	0	0.0	10	7.7

		HIV Diagn	oses (Annua	ıl Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate 3	Number	Col %	Case Rate 3	Number	Col %	Population
Total		6.3	100.0	6.3	3.7	100.0	3.7	195	100.0	99,933
Gender	Male	4.3	68.4		1.7	45.5		143	73.3	
	Female	2.0	31.6		2.0	54.5		52	26.7	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		5	2.6	
	13-19	0.0	0.0		0.0	0.0	•	0	0.0	
	20-24	0.3	5.3		0.0	0.0	٠	3	1.5	
	25-29	0.7	10.5		0.0	0.0		17	8.7	
	30-39	1.7	26.3		1.0	27.3		67	34.4	
	40-49	2.7	42.1		2.0	54.5		85	43.6	
	50-59	1.0	15.8		0.7	18.2		11	5.6	
	60+	0.0	0.0		0.0	0.0		7	3.6	
Race/Ethnicity	White	3.3	52.6		2.0	54.5		124	63.6	
	Black	0.3	5.3		1.0	27.3	٠	27	13.8	
	Hispanic	2.7	42.1		0.7	18.2	•	41	21.0	
	Multi Race	0.0	0.0		0.0	0.0		3	1.5	
Risk	MSM	2.7	42.1		1.0	27.3		76	39.0	
	IDU	0.0	0.0		0.0	0.0		58	29.7	
	MSM/IDU	0.0	0.0		0.0	0.0		9	4.6	
	Heterosexual	2.3	36.8		2.3	63.6		36	18.5	
	Fem Pres Het Cont	0.3	5.3		0.3	9.1		3	1.5	
	Blood Prod	0.0	0.0		0.0	0.0		1	0.5	
	Pediatric Risk	0.0	0.0		0.0	0.0		5	2.6	
	Unk	1.0	15.8		0.0	0.0		7	3.6	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	M	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		4.3	100.0	2.0	100.0	6.3	100.0	1.7	100.0	2.0	100.0	3.7	100.0	143	100.0	52	100.0	195	100.0
Risk	MSM	2.7	61.5	0.0	0.0	2.7	42.1	1.0	60.0	0.0	0.0	1.0	27.3	76	53.1	0	0.0	76	39.0
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	35	24.5	23	44.2	58	29.7
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	9	6.3	0	0.0	9	4.6
	Heterosexual	0.7	15.4	1.7	83.3	2.3	36.8	0.7	40.0	1.7	83.3	2.3	63.6	14	9.8	22	42.3	36	18.5
	Fem Pres Het Cont	0.0	0.0	0.3	16.7	0.3	5.3	0.0	0.0	0.3	16.7	0.3	9.1	0	0.0	3	5.8	3	1.5
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	0.7	0	0.0	1	0.5
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	0.7	4	7.7	5	2.6
	Unk	1.0	23.1	0.0	0.0	1.0	15.8	0.0	0.0	0.0	0.0	0.0	0.0	7	4.9	0	0.0	7	3.6

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		7	5
1986		2	3
1987		8	5
1988		10	5
1989		7	4
1990		9	7
1991		12	3
1992		16	7
1993		11	15
1994		12	12
1995		17	14
1996		11	6
1997		6	3
1998		10	4
1999		10	1
2000*	7	5	2
2001	3	5	
2002	3	2	2
2003	5	5	1
2004	1	2	1
2005	2	1	1
2006	5	5	
2007	5	8	1
2008	2	3	1
2009	9	3	1
2010	6	6	
2011	4	2	2

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		93	100.0	58.3	138	100.0	86.6	231	100.0	144.9	159,395
Gender	Male	56	60.2		96	69.6		152	65.8		
	Female	37	39.8		42	30.4		79	34.2		
Current Age	12 & under	0	0.0		0	0.0		0	0.0		
	13-19	2	2.2		1	0.7		3	1.3		
	20-24	10	10.8		1	0.7		11	4.8		
	25-29	4	4.3		4	2.9		8	3.5		
	30-39	18	19.4		14	10.1		32	13.9		
	40-49	28	30.1		50	36.2		78	33.8		
	50-59	25	26.9		45	32.6		70	30.3		
	60+	6	6.5		23	16.7		29	12.6		
Race/Ethnicity	White	51	54.8		77	55.8		128	55.4		
	Black	18	19.4		31	22.5		49	21.2		
	Hispanic	12	12.9		15	10.9		27	11.7		
	Multi Race	11	11.8		14	10.1		25	10.8		
	Others	1	1.1		1	0.7		2	0.9		
Risk	MSM	37	39.8		49	35.5		86	37.2		
	IDU	6	6.5		20	14.5		26	11.3		
	MSM/IDU	7	7.5		8	5.8		15	6.5		
	Heterosexual	22	23.7		23	16.7		45	19.5		
	Fem Pres Het Cont	12	12.9		12	8.7		24	10.4		
	Blood Prod	0	0.0		2	1.4		2	0.9		
	Pediatric Risk	1	1.1		3	2.2		4	1.7		
	Unk	8	8.6		21	15.2		29	12.6		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	IDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	Male	Fe	male	Т	otal	M	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		56	100.0	37	100.0	93	100.0	96	100.0	42	100.0	138	100.0	152	100.0	79	100.0	231	100.0
Risk	MSM	37	66.1	0	0.0	37	39.8	49	51.0	0	0.0	49	35.5	86	56.6	0	0.0	86	37.2
	IDU	2	3.6	4	10.8	6	6.5	10	10.4	10	23.8	20	14.5	12	7.9	14	17.7	26	11.3
	MSM/IDU	7	12.5	0	0.0	7	7.5	8	8.3	0	0.0	8	5.8	15	9.9	0	0.0	15	6.5
	Heterosexual	1	1.8	21	56.8	22	23.7	6	6.3	17	40.5	23	16.7	7	4.6	38	48.1	45	19.5
	Fem Pres Het Cont	0	0.0	12	32.4	12	12.9	0	0.0	12	28.6	12	8.7	0	0.0	24	30.4	24	10.4
	Blood Prod	0	0.0	0	0.0	0	0.0	1	1.0	1	2.4	2	1.4	1	0.7	1	1.3	2	0.9
	Pediatric Risk	1	1.8	0	0.0	1	1.1	1	1.0	2	4.8	3	2.2	2	1.3	2	2.5	4	1.7
	Unk	8	14.3	0	0.0	8	8.6	21	21.9	0	0.0	21	15.2	29	19.1	0	0.0	29	12.6

		HIV Diagn	oses (Annua	al Average)	AIDS Diagr	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		9.0	100.0	5.6	5.3	100.0	3.3	306	100.0	159,395
Gender	Male	6.7	74.1		5.3	100.0		231	75.5	
	Female	2.3	25.9		0.0	0.0		75	24.5	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		3	1.0	
	13-19	0.7	7.4		0.0	0.0		3	1.0	
	20-24	1.7	18.5		0.0	0.0		14	4.6	
	25-29	1.0	11.1		0.3	6.3		37	12.1	
	30-39	0.3	3.7		0.7	12.5		115	37.6	
	40-49	3.3	37.0		1.7	31.3		86	28.1	
	50-59	1.7	18.5		1.3	25.0		35	11.4	
	60+	0.3	3.7		1.3	25.0		13	4.2	
Race/Ethnicity	White	4.0	44.4		3.3	62.5		186	60.8	
	Black	3.0	33.3		1.3	25.0		70	22.9	
	Hispanic	1.0	11.1		0.7	12.5		30	9.8	
	Multi Race	1.0	11.1		0.0	0.0		20	6.5	
	Others	0.0	0.0		0.0	0.0		0	0.0	
Risk	MSM	3.7	40.7		2.3	43.8		127	41.5	
	IDU	0.3	3.7		0.0	0.0		63	20.6	
	MSM/IDU	0.3	3.7		0.7	12.5		13	4.2	
	Heterosexual	2.0	22.2		0.3	6.3		44	14.4	
	Fem Pres Het Cont	0.7	7.4		0.0	0.0		16	5.2	
	Blood Prod	0.0	0.0		0.0	0.0		5	1.6	
	Pediatric Risk	0.0	0.0		0.0	0.0		4	1.3	
	Unk	2.0	22.2		2.0	37.5		34	11.1	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	M	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		6.7	100.0	2.3	100.0	9.0	100.0	5.3	100.0	0.0	0.0	5.3	100.0	231	100.0	75	100.0	306	100.0
Risk	MSM	3.7	55.0	0.0	0.0	3.7	40.7	2.3	43.8	0.0	0.0	2.3	43.8	127	55.0	0	0.0	127	41.5
	IDU	0.0	0.0	0.3	14.3	0.3	3.7	0.0	0.0	0.0	0.0	0.0	0.0	44	19.0	19	25.3	63	20.6
	MSM/IDU	0.3	5.0	0.0	0.0	0.3	3.7	0.7	12.5	0.0	0.0	0.7	12.5	13	5.6	0	0.0	13	4.2
	Heterosexual	0.7	10.0	1.3	57.1	2.0	22.2	0.3	6.3	0.0	0.0	0.3	6.3	8	3.5	36	48.0	44	14.4
	Fem Pres Het Cont	0.0	0.0	0.7	28.6	0.7	7.4	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	16	21.3	16	5.2
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	4	1.7	1	1.3	5	1.6
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	0.4	3	4.0	4	1.3
	Unk	2.0	30.0	0.0	0.0	2.0	22.2	2.0	37.5	0.0	0.0	2.0	37.5	34	14.7	0	0.0	34	11.1

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		2	1
1986		3	1
1987		5	4
1988		6	4
1989		13	3
1990		13	8
1991		20	10
1992		28	11
1993		19	17
1994		20	11
1995		21	14
1996		20	12
1997		4	8
1998		8	2
1999		11	5
2000*	11	12	5
2001	8	5	7
2002	14	15	4
2003	12	14	5
2004	20	15	5
2005	9	8	7
2006	9	10	5
2007	9	11	3
2008	11	7	4
2009	7	3	4
2010	13	9	5
2011	7	4	7

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		243	100.0	77.1	442	100.0	140.2	685	100.0	217.4	315,158
Gender	Male	146	60.1		296	67.0		442	64.5		
	Female	96	39.5		146	33.0		242	35.3		
	Unk	1	0.4		0	0.0		1	0.1		
Current Age	12 & under	3	1.2		0	0.0		3	0.4		
	13-19	12	4.9		5	1.1		17	2.5		
	20-24	8	3.3		10	2.3		18	2.6		
	25-29	11	4.5		4	0.9		15	2.2		
	30-39	43	17.7		34	7.7		77	11.2		
	40-49	75	30.9		138	31.2		213	31.1		
	50-59	64	26.3		161	36.4		225	32.8		
	60+	27	11.1		90	20.4		117	17.1		
Race/Ethnicity	White	55	22.6		96	21.7		151	22.0		
	Black	106	43.6		197	44.6		303	44.2		
	Hispanic	57	23.5		95	21.5		152	22.2		
	Multi Race	23	9.5		54	12.2		77	11.2		
	Others	2	0.8		0	0.0		2	0.3		
Risk	MSM	63	25.9		91	20.6		154	22.5		
	IDU	25	10.3		72	16.3		97	14.2		
	MSM/IDU	5	2.1		20	4.5		25	3.6		
	Heterosexual	100	41.2		175	39.6		275	40.1		
	Fem Pres Het Cont	13	5.3		17	3.8		30	4.4		
	Blood Prod	0	0.0		3	0.7		3	0.4		
	Pediatric Risk	15	6.2		12	2.7		27	3.9	44.2 . 22.2 . 11.2 . 0.3 . 22.5 . 14.2 . 3.6 . 40.1 . 4.4 . 0.4 . 3.9 .	
	Unk	22	9.1		52	11.8		74	10.8		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	Т	otal	N	Male	Fe	male	1	otal	M	Iale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		146	100.0	96	100.0	243	100.0	296	100.0	146	100.0	442	100.0	442	100.0	242	100.0	684	100.0
Risk	MSM	63	43.2	0	0.0	63	25.9	91	30.7	0	0.0	91	20.6	154	34.8	0	0.0	154	22.5
	IDU	14	9.6	10	10.4	25	10.3	47	15.9	25	17.1	72	16.3	61	13.8	35	14.5	96	14.0
	MSM/IDU	5	3.4	0	0.0	5	2.1	20	6.8	0	0.0	20	4.5	25	5.7	0	0.0	25	3.7
	Heterosexual	34	23.3	66	68.8	100	41.2	76	25.7	99	67.8	175	39.6	110	24.9	165	68.2	275	40.2
	Fem Pres Het Cont	0	0.0	13	13.5	13	5.3	0	0.0	17	11.6	17	3.8	0	0.0	30	12.4	30	4.4
	Blood Prod	0	0.0	0	0.0	0	0.0	2	0.7	1	0.7	3	0.7	2	0.5	1	0.4	3	0.4
	Pediatric Risk	8	5.5	7	7.3	15	6.2	8	2.7	4	2.7	12	2.7	16	3.6	11	4.5	27	3.9
	Unk	22	15.1	0	0.0	22	9.1	52	17.6	0	0.0	52	11.8	74	16.7	0	0.0	74	10.8

		HIV Diagn	oses (Annua	ıl Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		22.0	100.0	7.0	15.3	100.0	4.9	998	100.0	315,158
Gender	Male	13.7	62.1		9.7	63.0		714	71.5	
	Female	8.3	37.9		5.7	37.0		284	28.5	
	Unk	0.0	0.0		0.0	0.0		0	0.0	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		21	2.1	
	13-19	1.0	4.5		0.7	4.3		10	1.0	
	20-24	2.0	9.1		1.0	6.5		23	2.3	
	25-29	2.7	12.1		1.0	6.5		93	9.3	
	30-39	3.7	16.7		2.0	13.0		348	34.9	
	40-49	5.7	25.8		5.3	34.8		329	33.0	
	50-59	2.7	12.1		2.7	17.4		108	10.8	
	60+	4.3	19.7		2.7	17.4		66	6.6	
Race/Ethnicity	White	5.0	22.7		3.3	21.7		323	32.4	
	Black	10.3	47.0		6.7	43.5		435	43.6	
	Hispanic	5.7	25.8		4.7	30.4		176	17.6	
	Multi Race	1.0	4.5		0.7	4.3		61	6.1	
	Others	0.0	0.0		0.0	0.0		3	0.3	
Risk	MSM	6.0	27.3		2.3	15.2		262	26.3	
	IDU	2.0	9.1		2.3	15.2		227	22.7	
	MSM/IDU	1.0	4.5		0.0	0.0		40	4.0	
	Heterosexual	11.0	50.0		7.7	50.0		270	27.1	
	Fem Pres Het Cont	1.0	4.5		1.0	6.5		50	5.0	
		0.0	0.0		0.0	0.0		13	1.3	
	Pediatric Risk	0.3	1.5		0.7	4.3		25	2.5	
	Unk	0.7	3.0		1.3	8.7		111	11.1	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS C	Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		13.7	100.0	8.3	100.0	22.0	100.0	9.7	100.0	5.7	100.0	15.3	100.0	714	100.0	284	100.0	998	100.0
Risk	MSM	6.0	43.9	0.0	0.0	6.0	27.3	2.3	24.1	0.0	0.0	2.3	15.2	262	36.7	0	0.0	262	26.3
	IDU	0.7	4.9	1.3	16.0	2.0	9.1	1.3	13.8	1.0	17.6	2.3	15.2	156	21.8	71	25.0	227	22.7
	MSM/IDU	1.0	7.3	0.0	0.0	1.0	4.5	0.0	0.0	0.0	0.0	0.0	0.0	40	5.6	0	0.0	40	4.0
	Heterosexual	5.3	39.0	5.7	68.0	11.0	50.0	4.3	44.8	3.3	58.8	7.7	50.0	121	16.9	149	52.5	270	27.1
	Fem Pres Het Cont	0.0	0.0	1.0	12.0	1.0	4.5	0.0	0.0	1.0	17.6	1.0	6.5	0	0.0	50	17.6	50	5.0
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	9	1.3	4	1.4	13	1.3
	Pediatric Risk	0.0	0.0	0.3	4.0	0.3	1.5	0.3	3.4	0.3	5.9	0.7	4.3	15	2.1	10	3.5	25	2.5
	Unk	0.7	4.9	0.0	0.0	0.7	3.0	1.3	13.8	0.0	0.0	1.3	8.7	111	15.5	0	0.0	111	11.1

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		31	19
1986		20	16
1987		28	15
1988		29	17
1989		44	33
1990	•	30	27
1991	٠	51	20
1992		90	46
1993	٠	79	49
1994		64	47
1995		56	46
1996		55	28
1997		35	15
1998		41	18
1999		30	20
2000*	27	45	13
2001	29	38	19
2002	19	27	20
2003	25	25	22
2004	32	30	12
2005	35	43	10
2006	33	25	10
2007	14	18	3
2008	24	18	7
2009	21	18	11
2010	20	14	9
2011	25	14	4

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		34	100.0	30.4	43	100.0	38.5	77	100.0	68.9	111,690
Gender	Male	24	70.6		29	67.4		53	68.8		
	Female	10	29.4		14	32.6		24	31.2		
Current Age	13-19	1	2.9		0	0.0		1	1.3		
	20-24	0	0.0		0	0.0		0	0.0		
	25-29	3	8.8		1	2.3		4	5.2		
	30-39	6	17.6		6	14.0		12	15.6		
	40-49	13	38.2		17	39.5		30	39.0		
	50-59	6	17.6		13	30.2		19	24.7		
	60+	5	14.7		6	14.0		11	14.3		
Race/Ethnicity	White	25	73.5		32	74.4		57	74.0		
	Black	4	11.8		3	7.0		7	9.1		
	Hispanic	4	11.8		6	14.0		10	13.0		
	Multi Race	1	2.9		2	4.7		3	3.9		
Risk	MSM	13	38.2		16	37.2		29	37.7		
	IDU	3	8.8		2	4.7		5	6.5		
	MSM/IDU	2	5.9		2	4.7		4	5.2		
	Heterosexual	4	11.8		10	23.3		14	18.2		
	Fem Pres Het Cont	6	17.6		6	14.0		12	15.6		
	Blood Prod	1	2.9		0	0.0		1	1.3		
	Unk	5	14.7		7	16.3		12	15.6		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	/Iale	Fe	male	Т	otal	М	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		24	100.0	10	100.0	34	100.0	29	100.0	14	100.0	43	100.0	53	100.0	24	100.0	77	100.0
Risk	MSM	13	54.2	0	0.0	13	38.2	16	55.2	0	0.0	16	37.2	29	54.7	0	0.0	29	37.7
	IDU	3	12.5	0	0.0	3	8.8	2	6.9	0	0.0	2	4.7	5	9.4	0	0.0	5	6.5
	MSM/IDU	2	8.3	0	0.0	2	5.9	2	6.9	0	0.0	2	4.7	4	7.5	0	0.0	4	5.2
	Heterosexual	0	0.0	4	40.0	4	11.8	2	6.9	8	57.1	10	23.3	2	3.8	12	50.0	14	18.2
	Fem Pres Het Cont	0	0.0	6	60.0	6	17.6	0	0.0	6	42.9	6	14.0	0	0.0	12	50.0	12	15.6
	Blood Prod	1	4.2	0	0.0	1	2.9	0	0.0	0	0.0	0	0.0	1	1.9	0	0.0	1	1.3
	Unk	5	20.8	0	0.0	5	14.7	7	24.1	0	0.0	7	16.3	12	22.6	0	0.0	12	15.6

		HIV Diagn	oses (Annua	ıl Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		2.0	100.0	1.8	1.7	100.0	1.5	81	100.0	111,690
Gender	Male	1.7	83.3		1.0	60.0		63	77.8	
	Female	0.3	16.7		0.7	40.0		18	22.2	
Age at Diagnosis	13-19	0.0	0.0		0.0	0.0		2	2.5	
	20-24	0.0	0.0		0.0	0.0	•	3	3.7	
	25-29	0.3	16.7		0.3	20.0		7	8.6	
	30-39	0.3	16.7		0.0	0.0		36	44.4	
	40-49	0.3	16.7		0.7	40.0		20	24.7	
	50-59	0.7	33.3		0.7	40.0		11	13.6	
	60+	0.3	16.7		0.0	0.0		2	2.5	
Race/Ethnicity	White	1.7	83.3		1.0	60.0		58	71.6	
	Black	0.0	0.0		0.0	0.0		9	11.1	
	Hispanic	0.0	0.0		0.3	20.0		12	14.8	
	Multi Race	0.3	16.7		0.3	20.0		2	2.5	
Risk	MSM	1.7	83.3		0.7	40.0		29	35.8	
	IDU	0.0	0.0		0.3	20.0		9	11.1	
	MSM/IDU	0.0	0.0		0.0	0.0		8	9.9	
	Heterosexual	0.0	0.0		0.3	20.0		15	18.5	
	Fem Pres Het Cont	0.3	16.7		0.3	20.0		6	7.4	
	Blood Prod	0.0	0.0		0.0	0.0		4	4.9	
	Unk	0.0	0.0		0.0	0.0		10	12.3	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

Newly Diagnosed HIV and AIDS Cases¹ (2009-2011) and Cumulative AIDS² (through 2011)

]	HIV Diaș	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	To	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		1.7	100.0	0.3	100.0	2.0	100.0	1.0	100.0	0.7	100.0	1.7	100.0	63	100.0	18	100.0	81	100.0
Risk	MSM	1.7	100.0	0.0	0.0	1.7	83.3	0.7	66.7	0.0	0.0	0.7	40.0	29	46.0	0	0.0	29	35.8
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.3	33.3	0.0	0.0	0.3	20.0	9	14.3	0	0.0	9	11.1
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	8	12.7	0	0.0	8	9.9
	Heterosexual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	50.0	0.3	20.0	4	6.3	11	61.1	15	18.5
	Fem Pres Het Cont	0.0	0.0	0.3	100.0	0.3	16.7	0.0	0.0	0.3	50.0	0.3	20.0	0	0.0	6	33.3	6	7.4
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3	4.8	1	5.6	4	4.9
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	10	15.9	0	0.0	10	12.3

¹Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

² Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		2	1
1986		1	1
1987		2	
1988			1
1989		4	4
1990		3	2
1991		5	3
1992		5	3
1993		2	1
1994		3	4
1995		1	1
1996		3	1
1997		1	
1998		8	
1999		5	2
2000*	6	6	1
2001	3	3	1
2002	4	3	1
2003	4	5	
2004	5	4	2
2005		3	3
2006	2	2	1
2007		4	
2008	1	1	1
2009	2	4	1
2010	2		3
2011	2	1	

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		70	100.0	31.7	94	100.0	42.6	164	100.0	74.2	220,882
Gender	Male	59	84.3		71	75.5		130	79.3		
	Female	11	15.7		23	24.5		34	20.7		
Current Age	12 & under	0	0.0		0	0.0		0	0.0		
	13-19	0	0.0		2	2.1		2	1.2		
	20-24	4	5.7		0	0.0		4	2.4		
	25-29	5	7.1		3	3.2		8	4.9		
	30-39	7	10.0		8	8.5		15	9.1		
	40-49	31	44.3		26	27.7		57	34.8		
	50-59	15	21.4		43	45.7		58	35.4		
	60+	8	11.4		12	12.8		20	12.2		
Race/Ethnicity	White	52	74.3		61	64.9		113	68.9		
	Black	7	10.0		13	13.8		20	12.2		
	Hispanic	3	4.3		13	13.8		16	9.8		
	Multi Race	7	10.0		7	7.4		14	8.5		
	Others	1	1.4		0	0.0		1	0.6		
Risk	MSM	41	58.6		41	43.6		82	50.0		
	IDU	7	10.0		11	11.7		18	11.0		
	MSM/IDU	3	4.3		7	7.4		10	6.1		
	Heterosexual	6	8.6		15	16.0		21	12.8		
	Fem Pres Het Cont	3	4.3		6	6.4		9	5.5		
	Blood Prod	0	0.0		4	4.3		4	2.4		
	Pediatric Risk	2	2.9		4	4.3		6	3.7		
	Unk	8	11.4		6	6.4		14	8.5		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	Т	otal	N	Male	Fe	male	1	otal	M	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		59	100.0	11	100.0	70	100.0	71	100.0	23	100.0	94	100.0	130	100.0	34	100.0	164	100.0
Risk	MSM	41	69.5	0	0.0	41	58.6	41	57.7	0	0.0	41	43.6	82	63.1	0	0.0	82	50.0
	IDU	4	6.8	3	27.3	7	10.0	7	9.9	4	17.4	11	11.7	11	8.5	7	20.6	18	11.0
	MSM/IDU	3	5.1	0	0.0	3	4.3	7	9.9	0	0.0	7	7.4	10	7.7	0	0.0	10	6.1
	Heterosexual	1	1.7	5	45.5	6	8.6	4	5.6	11	47.8	15	16.0	5	3.8	16	47.1	21	12.8
	Fem Pres Het Cont	0	0.0	3	27.3	3	4.3	0	0.0	6	26.1	6	6.4	0	0.0	9	26.5	9	5.5
	Blood Prod	0	0.0	0	0.0	0	0.0	3	4.2	1	4.3	4	4.3	3	2.3	1	2.9	4	2.4
	Pediatric Risk	2	3.4	0	0.0	2	2.9	3	4.2	1	4.3	4	4.3	5	3.8	1	2.9	6	3.7
	Unk	8	13.6	0	0.0	8	11.4	6	8.5	0	0.0	6	6.4	14	10.8	0	0.0	14	8.5

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		5.3	100.0	2.4	5.0	100.0	2.3	213	100.0	220,882
Gender	Male	4.3	81.3		3.0	60.0		167	78.4	
	Female	1.0	18.8		2.0	40.0		46	21.6	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		3	1.4	
	13-19	0.3	6.3		0.7	13.3		4	1.9	
	20-24	0.3	6.3		0.0	0.0		5	2.3	
	25-29	0.3	6.3		0.3	6.7		20	9.4	
	30-39	1.0	18.8		1.7	33.3		79	37.1	
	40-49	2.3	43.8		1.7	33.3		74	34.7	
	50-59	1.0	18.8		0.7	13.3		24	11.3	
	60+	0.0	0.0		0.0	0.0		4	1.9	
Race/Ethnicity	White	3.7	68.8		2.3	46.7		152	71.4	
	Black	0.3	6.3		1.0	20.0		27	12.7	
	Hispanic	0.7	12.5		1.0	20.0		23	10.8	
	Multi Race	0.7	12.5		0.7	13.3		11	5.2	
	Others	0.0	0.0		0.0	0.0		0	0.0	
Risk	MSM	2.7	50.0		1.3	26.7		96	45.1	
	IDU	0.3	6.3		0.3	6.7		40	18.8	
	MSM/IDU	0.7	12.5		0.3	6.7		13	6.1	
	Heterosexual	0.7	12.5		1.7	33.3		23	10.8	
	Fem Pres Het Cont	0.0	0.0		0.0	0.0		12	5.6	
	Blood Prod	0.0	0.0		0.0	0.0		10	4.7	
	Pediatric Risk	0.0	0.0		0.7	13.3		8	3.8	
	Unk	1.0	18.8		0.7	13.3		11	5.2	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Diag	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS C	Cases	
		M	ale	Fer	nale	To	otal	M	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		4.3	100.0	1.0	100.0	5.3	100.0	3.0	100.0	2.0	100.0	5.0	100.0	167	100.0	46	100.0	213	100.0
Risk	MSM	2.7	61.5	0.0	0.0	2.7	50.0	1.3	44.4	0.0	0.0	1.3	26.7	96	57.5	0	0.0	96	45.1
	IDU	0.0	0.0	0.3	33.3	0.3	6.3	0.3	11.1	0.0	0.0	0.3	6.7	27	16.2	13	28.3	40	18.8
	MSM/IDU	0.7	15.4	0.0	0.0	0.7	12.5	0.3	11.1	0.0	0.0	0.3	6.7	13	7.8	0	0.0	13	6.1
	Heterosexual	0.0	0.0	0.7	66.7	0.7	12.5	0.0	0.0	1.7	83.3	1.7	33.3	6	3.6	17	37.0	23	10.8
	Fem Pres Het Cont	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	12	26.1	12	5.6
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	8	4.8	2	4.3	10	4.7
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.3	11.1	0.3	16.7	0.7	13.3	6	3.6	2	4.3	8	3.8
	Unk	1.0	23.1	0.0	0.0	1.0	18.8	0.7	22.2	0.0	0.0	0.7	13.3	11	6.6	0	0.0	11	5.2

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		3	
1986		3	3
1987		5	5
1988		4	2
1989		8	5
1990		12	3
1991		15	7
1992		14	12
1993		14	7
1994		15	16
1995		12	12
1996		8	4
1997		10	5
1998		7	4
1999		7	1
2000*	4	5	4
2001	10	7	7
2002	4	6	3
2003	9	13	3
2004	7	10	4
2005	9	10	3
2006	7	4	4
2007	7	5	1
2008	5	1	2
2009	7	3	2
2010	3	4	1
2011	6	8	

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		173	100.0	111.6	207	100.0	133.5	380	100.0	245.1	155,058
Gender	Male	85	49.1		113	54.6		198	52.1		
	Female	88	50.9		94	45.4		182	47.9		
Current Age	12 & under	3	1.7		0	0.0		3	0.8		
	13-19	4	2.3		1	0.5		5	1.3		
	20-24	5	2.9		2	1.0		7	1.8		
	25-29	16	9.2		7	3.4		23	6.1		
	30-39	28	16.2		18	8.7		46	12.1		
	40-49	57	32.9		89	43.0		146	38.4		
	50-59	46	26.6		61	29.5		107	28.2		
	60+	14	8.1		29	14.0		43	11.3		
Race/Ethnicity	White	61	35.3		67	32.4		128	33.7		
	Black	43	24.9		60	29.0		103	27.1		
	Hispanic	33	19.1		41	19.8		74	19.5		
	Multi Race	35	20.2		37	17.9		72	18.9		
	Others	1	0.6		2	1.0		3	0.8		
Risk	MSM	44	25.4		47	22.7		91	23.9		
	IDU	22	12.7		47	22.7		69	18.2		
	MSM/IDU	4	2.3		12	5.8		16	4.2		
	Heterosexual	54	31.2		65	31.4		119	31.3		
	Fem Pres Het Cont	31	17.9		18	8.7		49	12.9		
	Blood Prod	0	0.0		1	0.5		1	0.3		
	Pediatric Risk	8	4.6		3	1.4		11	2.9		
	Unk	10	5.8		14	6.8		24	6.3		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	IDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	Male	Fe	male	Т	otal	M	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		85	100.0	88	100.0	173	100.0	113	100.0	94	100.0	207	100.0	198	100.0	182	100.0	380	100.0
Risk	MSM	44	51.8	0	0.0	44	25.4	47	41.6	0	0.0	47	22.7	91	46.0	0	0.0	91	23.9
	IDU	9	10.6	13	14.8	22	12.7	25	22.1	22	23.4	47	22.7	34	17.2	35	19.2	69	18.2
	MSM/IDU	4	4.7	0	0.0	4	2.3	12	10.6	0	0.0	12	5.8	16	8.1	0	0.0	16	4.2
	Heterosexual	15	17.6	39	44.3	54	31.2	14	12.4	51	54.3	65	31.4	29	14.6	90	49.5	119	31.3
	Fem Pres Het Cont	0	0.0	31	35.2	31	17.9	0	0.0	18	19.1	18	8.7	0	0.0	49	26.9	49	12.9
	Blood Prod	0	0.0	0	0.0	0	0.0	1	0.9	0	0.0	1	0.5	1	0.5	0	0.0	1	0.3
	Pediatric Risk	3	3.5	5	5.7	8	4.6	0	0.0	3	3.2	3	1.4	3	1.5	8	4.4	11	2.9
	Unk	10	11.8	0	0.0	10	5.8	14	12.4	0	0.0	14	6.8	24	12.1	0	0.0	24	6.3

		HIV Diagn	oses (Annua	ıl Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		11.3	100.0	7.3	9.3	100.0	6.0	476	100.0	155,058
Gender	Male	6.0	52.9		5.0	53.6		294	61.8	
	Female	5.3	47.1		4.3	46.4		182	38.2	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		6	1.3	
	13-19	0.0	0.0		0.0	0.0		3	0.6	
	20-24	1.3	11.8		0.3	3.6		16	3.4	
	25-29	1.0	8.8		0.7	7.1		53	11.1	
	30-39	1.7	14.7		2.0	21.4		168	35.3	
	40-49	4.0	35.3		3.7	39.3		172	36.1	
	50-59	3.0	26.5		2.0	21.4		51	10.7	
	60+	0.3	2.9		0.7	7.1		7	1.5	
Race/Ethnicity	White	3.3	29.4		4.3	46.4		206	43.3	
	Black	4.0	35.3		3.0	32.1		152	31.9	
	Hispanic	2.3	20.6		1.3	14.3		72	15.1	
	Multi Race	1.7	14.7		0.7	7.1		44	9.2	
	Others	0.0	0.0		0.0	0.0		2	0.4	
Risk	MSM	1.3	11.8		2.0	21.4		120	25.2	
	IDU	1.7	14.7		2.3	25.0		146	30.7	
	MSM/IDU	1.3	11.8		1.0	10.7		26	5.5	
	Heterosexual	4.3	38.2		2.7	28.6		98	20.6	
	Fem Pres Het Cont	1.7	14.7		0.7	7.1		38	8.0	
_	Blood Prod	0.0	0.0		0.0	0.0		4	0.8	
	Pediatric Risk	0.0	0.0		0.3	3.6		8	1.7	
	Unk	1.0	8.8		0.3	3.6		36	7.6	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		1	HIV Diaș	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	To	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		6.0	100.0	5.3	100.0	11.3	100.0	5.0	100.0	4.3	100.0	9.3	100.0	294	100.0	182	100.0	476	100.0
Risk	MSM	1.3	22.2	0.0	0.0	1.3	11.8	2.0	40.0	0.0	0.0	2.0	21.4	120	40.8	0	0.0	120	25.2
	IDU	0.7	11.1	1.0	18.8	1.7	14.7	1.0	20.0	1.3	30.8	2.3	25.0	86	29.3	60	33.0	146	30.7
	MSM/IDU	1.3	22.2	0.0	0.0	1.3	11.8	1.0	20.0	0.0	0.0	1.0	10.7	26	8.8	0	0.0	26	5.5
	Heterosexual	1.7	27.8	2.7	50.0	4.3	38.2	0.7	13.3	2.0	46.2	2.7	28.6	22	7.5	76	41.8	98	20.6
	Fem Pres Het Cont	0.0	0.0	1.7	31.3	1.7	14.7	0.0	0.0	0.7	15.4	0.7	7.1	0	0.0	38	20.9	38	8.0
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3	1.0	1	0.5	4	0.8
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	7.7	0.3	3.6	1	0.3	7	3.8	8	1.7
	Unk	1.0	16.7	0.0	0.0	1.0	8.8	0.3	6.7	0.0	0.0	0.3	3.6	36	12.2	0	0.0	36	7.6

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		14	6
1986		10	10
1987		5	5
1988		8	6
1989		16	9
1990		26	17
1991		27	14
1992		19	21
1993		26	11
1994		35	21
1995		35	16
1996		22	21
1997		25	10
1998		23	11
1999		12	8
2000*	9	21	8
2001	18	12	6
2002	26	17	11
2003	22	16	4
2004	24	20	8
2005	16	17	8
2006	17	21	8
2007	11	11	7
2008	11	10	7
2009	7	4	6
2010	11	14	5
2011	16	10	6

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		11	100.0	33.8	14	100.0	43.0	25	100.0	76.7	32,578
Gender	Male	8	72.7		10	71.4		18	72.0		
	Female	3	27.3		4	28.6		7	28.0		
Current Age	20-24	0	0.0		0	0.0		0	0.0		
	25-29	0	0.0		0	0.0		0	0.0		
	30-39	1	9.1		3	21.4		4	16.0		
	40-49	5	45.5		3	21.4		8	32.0		
	50-59	4	36.4		7	50.0		11	44.0		
	60+	1	9.1		1	7.1		2	8.0		
Race/Ethnicity	White	5	45.5		9	64.3		14	56.0		
	Black	3	27.3		3	21.4		6	24.0		
	Hispanic	2	18.2		1	7.1		3	12.0		
	Multi Race	0	0.0		1	7.1		1	4.0		
	Others	1	9.1		0	0.0		1	4.0		
Risk	MSM	5	45.5		5	35.7		10	40.0		
	IDU	0	0.0		3	21.4		3	12.0		
	MSM/IDU	0	0.0		0	0.0		0	0.0		
	Heterosexual	2	18.2		3	21.4		5	20.0		
	Fem Pres Het Cont	2	18.2		0	0.0		2	8.0		
	Blood Prod	0	0.0		0	0.0		0	0.0		
	Unk	2	18.2		3	21.4		5	20.0		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		M	ale	Fe	male	Т	otal	N	Aale	Fe	male	Т	otal	M	[ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		8	100.0	3	100.0	11	100.0	10	100.0	4	100.0	14	100.0	18	100.0	7	100.0	25	100.0
Risk	MSM	5	62.5	0	0.0	5	45.5	5	50.0	0	0.0	5	35.7	10	55.6	0	0.0	10	40.0
	IDU	0	0.0	0	0.0	0	0.0	2	20.0	1	25.0	3	21.4	2	11.1	1	14.3	3	12.0
	MSM/IDU	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Heterosexual	1	12.5	1	33.3	2	18.2	0	0.0	3	75.0	3	21.4	1	5.6	4	57.1	5	20.0
	Fem Pres Het Cont	0	0.0	2	66.7	2	18.2	0	0.0	0	0.0	0	0.0	0	0.0	2	28.6	2	8.0
	Blood Prod	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Unk	2	25.0	0	0.0	2	18.2	3	30.0	0	0.0	3	21.4	5	27.8	0	0.0	5	20.0

		HIV Diagn	oses (Annua	ıl Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		0.0	0.0	0.0	0.7	100.0	2.0	39	100.0	32,578
Gender	Male	0.0	0.0		0.3	50.0		34	87.2	
	Female	0.0	0.0		0.3	50.0		5	12.8	
Age at Diagnosis	20-24	0.0	0.0		0.0	0.0		1	2.6	
	25-29	0.0	0.0		0.3	50.0		1	2.6	
	30-39	0.0	0.0		0.0	0.0		15	38.5	
	40-49	0.0	0.0		0.0	0.0		14	35.9	
	50-59	0.0	0.0		0.3	50.0		7	17.9	
	60+	0.0	0.0		0.0	0.0		1	2.6	
Race/Ethnicity	White	0.0	0.0		0.3	50.0		30	76.9	
	Black	0.0	0.0		0.0	0.0		4	10.3	
	Hispanic	0.0	0.0		0.3	50.0		4	10.3	
	Multi Race	0.0	0.0		0.0	0.0		1	2.6	
	Others	0.0	0.0		0.0	0.0		0	0.0	
Risk	MSM	0.0	0.0		0.3	50.0		15	38.5	
	IDU	0.0	0.0		0.0	0.0		11	28.2	
	MSM/IDU	0.0	0.0		0.0	0.0		1	2.6	
	Heterosexual	0.0	0.0		0.3	50.0		4	10.3	
	Fem Pres Het Cont	0.0	0.0		0.0	0.0		1	2.6	
	Blood Prod	0.0	0.0		0.0	0.0		1	2.6	
	Unk	0.0	0.0		0.0	0.0		6	15.4	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

Newly Diagnosed HIV and AIDS Cases¹ (2009-2011) and Cumulative AIDS² (through 2011)

		1	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fei	nale	To	otal	М	ale	Fer	nale	To	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		0.0	0.0	0.0	0.0	0.0	0.0	0.3	100.0	0.3	100.0	0.7	100.0	34	100.0	5	100.0	39	100.0
Risk	MSM	0.0	0.0	0.0	0.0	0.0	0.0	0.3	100.0	0.0	0.0	0.3	50.0	15	44.1	0	0.0	15	38.5
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	10	29.4	1	20.0	11	28.2
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	2.9	0	0.0	1	2.6
	Heterosexual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	100.0	0.3	50.0	1	2.9	3	60.0	4	10.3
	Fem Pres Het Cont	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	1	20.0	1	2.6
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	2.9	0	0.0	1	2.6
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	6	17.6	0	0.0	6	15.4

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		1	1
1986		1	1
1988	•	1	
1989		1	1
1990	٠	3	1
1991	•	4	3
1992		3	1
1993		6	5
1994		4	2
1995		2	4
1996		1	3
1999	٠	3	2
2000*			1
2001	2	•	
2002	5	٠	
2003	1	1	
2004	2	1	
2006		2	
2007	1	1	
2008	2	2	
2009		2	

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	I AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		7	100.0	38.1	6	100.0	32.7	13	100.0	70.8	18,361
Gender	Male	6	85.7		5	83.3		11	84.6		
	Female	1	14.3		1	16.7		2	15.4		
Current Age	25-29	1	14.3		1	16.7		2	15.4		
	40-49	4	57.1		1	16.7		5	38.5		
	50-59	1	14.3		3	50.0		4	30.8		
	60+	1	14.3		1	16.7		2	15.4		
Race/Ethnicity	White	5	71.4		3	50.0		8	61.5		
	Black	1	14.3		1	16.7		2	15.4		
	Hispanic	1	14.3		2	33.3		3	23.1		
Risk	MSM	5	71.4		4	66.7		9	69.2		
	IDU	0	0.0		2	33.3		2	15.4		
	MSM/IDU	0	0.0		0	0.0		0	0.0		
	Heterosexual	1	14.3		0	0.0		1	7.7		
	Unk	1	14.3		0	0.0		1	7.7		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		M	ale	Fe	male	Т	otal	N	Aale	Fe	male	1	otal	M	Iale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		6	100.0	1	100.0	7	100.0	5	100.0	1	100.0	6	100.0	11	100.0	2	100.0	13	100.0
Risk	MSM	5	83.3	0	0.0	5	71.4	4	80.0	0	0.0	4	66.7	9	81.8	0	0.0	9	69.2
	IDU	0	0.0	0	0.0	0	0.0	1	20.0	1	100.0	2	33.3	1	9.1	1	50.0	2	15.4
	MSM/IDU	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Heterosexual	0	0.0	1	100.0	1	14.3	0	0.0	0	0.0	0	0.0	0	0.0	1	50.0	1	7.7
	Unk	1	16.7	0	0.0	1	14.3	0	0.0	0	0.0	0	0.0	1	9.1	0	0.0	1	7.7

		HIV Diagn	oses (Annua	l Average)	AIDS Diagr	noses (Annua	ıl Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		0.3	100.0	1.8	0.7	100.0	3.6	15	100.0	18,36
Gender	Male	0.3	100.0		0.7	100.0		15	100.0	
	Female	0.0	0.0		0.0	0.0		0	0.0	
Age at Diagnosis	25-29	0.0	0.0		0.0	0.0		2	13.3	
	40-49	0.0	0.0		0.3	50.0		5	33.3	
	50-59	0.0	0.0		0.3	50.0		7	46.7	
	60+	0.3	100.0		0.0	0.0		1	6.7	
Race/Ethnicity	White	0.3	100.0		0.0	0.0		13	86.7	
	Black	0.0	0.0		0.3	50.0		1	6.7	
	Hispanic	0.0	0.0		0.3	50.0		1	6.7	
Risk	MSM	0.0	0.0		0.3	50.0		10	66.7	
	IDU	0.0	0.0		0.3	50.0		1	6.7	
	MSM/IDU	0.0	0.0		0.0	0.0		2	13.3	
	Heterosexual	0.0	0.0		0.0	0.0		1	6.7	
	Unk	0.3	100.0		0.0	0.0		1	6.7	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

Newly Diagnosed HIV and AIDS Cases¹ (2009-2011) and Cumulative AIDS² (through 2011)

		1	HIV Diag	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	M	ale	Fer	nale	To	otal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		0.3	100.0	0.0	0.0	0.3	100.0	0.7	100.0	0.0	0.0	0.7	100.0	15	100.0	0	0.0	15	100.0
Risk	MSM	0.0	0.0	0.0	0.0	0.0	0.0	0.3	50.0	0.0	0.0	0.3	50.0	10	66.7	0	0.0	10	66.7
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.3	50.0	0.0	0.0	0.3	50.0	1	6.7	0	0.0	1	6.7
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	13.3	0	0.0	2	13.3
	Heterosexual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	6.7	0	0.0	1	6.7
	Unk	0.3	100.0	0.0	0.0	0.3	100.0	0.0	0.0	0.0	0.0	0.0	0.0	1	6.7	0	0.0	1	6.7

¹Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
1987		1	1
1989		1	
1990		2	
1991		1	2
1992		1	1
1993		3	4
1994		1	
1995			1
1996			1
2001		1	
2003	2		
2005	1	1	
2006	1		
2007	1	1	
2008	1		
2010	1	1	
2011		1	

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	1 AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		16	100.0	45.5	9	100.0	25.6	25	100.0	71.0	35,198
Gender	Male	13	81.3		6	66.7		19	76.0		
	Female	3	18.8		3	33.3		6	24.0		
Current Age	12 & under	0	0.0		0	0.0		0	0.0		
	20-24	1	6.3		1	11.1		2	8.0		
	25-29	1	6.3		0	0.0		1	4.0		
	30-39	3	18.8		1	11.1		4	16.0		
	40-49	6	37.5		4	44.4		10	40.0		
	50-59	3	18.8		3	33.3		6	24.0		
	60+	2	12.5		0	0.0		2	8.0		
Race/Ethnicity	White	9	56.3		6	66.7		15	60.0		
	Black	3	18.8		0	0.0		3	12.0		
	Hispanic	4	25.0		0	0.0		4	16.0		
	Multi Race	0	0.0		3	33.3		3	12.0		
Risk	MSM	6	37.5		4	44.4		10	40.0		
	IDU	3	18.8		1	11.1		4	16.0		
	Heterosexual	2	12.5		2	22.2		4	16.0		
	Fem Pres Het Cont	1	6.3		0	0.0		1	4.0		
	Blood Prod	0	0.0		1	11.1		1	4.0		
	Pediatric Risk	0	0.0		1	11.1		1	4.0		
	Unk	4	25.0		0	0.0		4	16.0		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	Male	Fe	male	Т	otal	M	[ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		13	100.0	3	100.0	16	100.0	6	100.0	3	100.0	9	100.0	19	100.0	6	100.0	25	100.0
Risk	MSM	6	46.2	0	0.0	6	37.5	4	66.7	0	0.0	4	44.4	10	52.6	0	0.0	10	40.0
	IDU	1	7.7	2	66.7	3	18.8	0	0.0	1	33.3	1	11.1	1	5.3	3	50.0	4	16.0
	Heterosexual	2	15.4	0	0.0	2	12.5	1	16.7	1	33.3	2	22.2	3	15.8	1	16.7	4	16.0
	Fem Pres Het Cont	0	0.0	1	33.3	1	6.3	0	0.0	0	0.0	0	0.0	0	0.0	1	16.7	1	4.0
	Blood Prod	0	0.0	0	0.0	0	0.0	1	16.7	0	0.0	1	11.1	1	5.3	0	0.0	1	4.0
	Pediatric Risk	0	0.0	0	0.0	0	0.0	0	0.0	1	33.3	1	11.1	0	0.0	1	16.7	1	4.0
	Unk	4	30.8	0	0.0	4	25.0	0	0.0	0	0.0	0	0.0	4	21.1	0	0.0	4	16.0

		HIV Diagn	oses (Annua	ıl Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		1.0	100.0	2.8	0.3	100.0	0.9	21	100.0	35,198
Gender	Male	0.7	66.7		0.0	0.0		14	66.7	
	Female	0.3	33.3		0.3	100.0		7	33.3	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		1	4.8	
	20-24	0.3	33.3		0.0	0.0		1	4.8	
	25-29	0.0	0.0		0.0	0.0		2	9.5	
	30-39	0.0	0.0		0.3	100.0		7	33.3	
	40-49	0.3	33.3		0.0	0.0		8	38.1	
	50-59	0.0	0.0		0.0	0.0		2	9.5	
	60+	0.3	33.3		0.0	0.0		0	0.0	
Race/Ethnicity	White	1.0	100.0		0.3	100.0		18	85.7	
	Black	0.0	0.0		0.0	0.0		0	0.0	
	Hispanic	0.0	0.0		0.0	0.0		0	0.0	
	Multi Race	0.0	0.0		0.0	0.0		3	14.3	
Risk	MSM	0.3	33.3		0.0	0.0		10	47.6	
	IDU	0.0	0.0		0.0	0.0		5	23.8	
	Heterosexual	0.0	0.0		0.3	100.0		3	14.3	
	Fem Pres Het Cont	0.3	33.3		0.0	0.0		1	4.8	
	Blood Prod	0.0	0.0		0.0	0.0		1	4.8	
	Pediatric Risk	0.0	0.0		0.0	0.0		1	4.8	
	Unk	0.3	33.3		0.0	0.0		0	0.0	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

Newly Diagnosed HIV and AIDS Cases¹ (2009-2011) and Cumulative AIDS² (through 2011)

		1	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fei	nale	To	otal	M	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		0.7	100.0	0.3	100.0	1.0	100.0	0.0	0.0	0.3	100.0	0.3	100.0	14	100.0	7	100.0	21	100.0
Risk	MSM	0.3	50.0	0.0	0.0	0.3	33.3	0.0	0.0	0.0	0.0	0.0	0.0	10	71.4	0	0.0	10	47.6
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	14.3	3	42.9	5	23.8
	Heterosexual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	100.0	0.3	100.0	1	7.1	2	28.6	3	14.3
	Fem Pres Het Cont	0.0	0.0	0.3	100.0	0.3	33.3	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	1	14.3	1	4.8
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	7.1	0	0.0	1	4.8
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	1	14.3	1	4.8
	Unk	0.3	50.0	0.0	0.0	0.3	33.3	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	0	0.0	0	0.0

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
1987		1	
1988		1	1
1990		1	1
1992		2	1
1993		1	2
1994		2	
1995		2	1
1996		2	2
1997			1
1998		1	
2000*	2	1	
2001	1		
2002	2		2
2003	2	2	
2004	2		
2005		2	
2007	1		1
2008	2	2	
2010	2	1	
2011	1		

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		34	100.0	34.3	38	100.0	38.4	72	100.0	72.7	99,033
Gender	Male	26	76.5		27	71.1		53	73.6		
	Female	8	23.5		11	28.9		19	26.4		
Current Age	12 & under	0	0.0		0	0.0		0	0.0		
	20-24	1	2.9		0	0.0		1	1.4		
	25-29	5	14.7		2	5.3		7	9.7		
	30-39	4	11.8		6	15.8		10	13.9		
	40-49	11	32.4		11	28.9		22	30.6		
	50-59	8	23.5		12	31.6		20	27.8		
	60+	5	14.7		7	18.4		12	16.7		
Race/Ethnicity	White	20	58.8		25	65.8		45	62.5		
	Black	7	20.6		7	18.4		14	19.4		
	Hispanic	2	5.9		4	10.5		6	8.3		
	Multi Race	4	11.8		2	5.3		6	8.3		
	Others	1	2.9		0	0.0		1	1.4		
Risk	MSM	16	47.1		16	42.1		32	44.4		
	IDU	5	14.7		5	13.2		10	13.9		
	MSM/IDU	0	0.0		2	5.3		2	2.8		
	Heterosexual	9	26.5		11	28.9		20	27.8		
	Fem Pres Het Cont	2	5.9		2	5.3		4	5.6		
	Blood Prod	0	0.0		1	2.6		1	1.4		
	Pediatric Risk	0	0.0		0	0.0		0	0.0		
	Unk	2	5.9		1	2.6		3	4.2		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	Male	Fe	male	1	otal	M	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		26	100.0	8	100.0	34	100.0	27	100.0	11	100.0	38	100.0	53	100.0	19	100.0	72	100.0
Risk	MSM	16	61.5	0	0.0	16	47.1	16	59.3	0	0.0	16	42.1	32	60.4	0	0.0	32	44.4
	IDU	5	19.2	0	0.0	5	14.7	4	14.8	1	9.1	5	13.2	9	17.0	1	5.3	10	13.9
	MSM/IDU	0	0.0	0	0.0	0	0.0	2	7.4	0	0.0	2	5.3	2	3.8	0	0.0	2	2.8
	Heterosexual	3	11.5	6	75.0	9	26.5	4	14.8	7	63.6	11	28.9	7	13.2	13	68.4	20	27.8
	Fem Pres Het Cont	0	0.0	2	25.0	2	5.9	0	0.0	2	18.2	2	5.3	0	0.0	4	21.1	4	5.6
	Blood Prod	0	0.0	0	0.0	0	0.0	0	0.0	1	9.1	1	2.6	0	0.0	1	5.3	1	1.4
	Pediatric Risk	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Unk	2	7.7	0	0.0	2	5.9	1	3.7	0	0.0	1	2.6	3	5.7	0	0.0	3	4.2

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate 3	Number	Col %	Case Rate 3	Number	Col %	Population
Total		2.7	100.0	2.7	2.0	100.0	2.0	90	100.0	99,033
Gender	Male	2.3	87.5		1.7	83.3		71	78.9	
	Female	0.3	12.5		0.3	16.7		19	21.1	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		1	1.1	
	20-24	0.7	25.0		0.0	0.0		5	5.6	
	25-29	1.3	50.0		0.3	16.7		12	13.3	
	30-39	0.3	12.5		0.3	16.7		33	36.7	
	40-49	0.3	12.5		0.3	16.7		26	28.9	
	50-59	0.0	0.0		1.0	50.0		8	8.9	
	60+	0.0	0.0		0.0	0.0		5	5.6	
Race/Ethnicity	White	2.3	87.5		1.7	83.3		72	80.0	
	Black	0.0	0.0		0.0	0.0	•	13	14.4	
	Hispanic	0.0	0.0		0.0	0.0		3	3.3	
	Multi Race	0.3	12.5		0.3	16.7	•	2	2.2	
	Others	0.0	0.0		0.0	0.0		0	0.0	
Risk	MSM	2.3	87.5		1.0	50.0		40	44.4	
	IDU	0.0	0.0		0.0	0.0		17	18.9	
	MSM/IDU	0.0	0.0		0.3	16.7		4	4.4	
	Heterosexual	0.0	0.0		0.3	16.7		16	17.8	
	Fem Pres Het Cont	0.3	12.5		0.3	16.7		4	4.4	
	Blood Prod	0.0	0.0		0.0	0.0		3	3.3	
	Pediatric Risk	0.0	0.0		0.0	0.0		1	1.1	
	Unk	0.0	0.0		0.0	0.0		5	5.6	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

Newly Diagnosed HIV and AIDS Cases¹ (2009-2011) and Cumulative AIDS² (through 2011)

		I	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	M	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		2.3	100.0	0.3	100.0	2.7	100.0	1.7	100.0	0.3	100.0	2.0	100.0	71	100.0	19	100.0	90	100.0
Risk	MSM	2.3	100.0	0.0	0.0	2.3	87.5	1.0	60.0	0.0	0.0	1.0	50.0	40	56.3	0	0.0	40	44.4
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	13	18.3	4	21.1	17	18.9
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.3	20.0	0.0	0.0	0.3	16.7	4	5.6	0	0.0	4	4.4
	Heterosexual	0.0	0.0	0.0	0.0	0.0	0.0	0.3	20.0	0.0	0.0	0.3	16.7	6	8.5	10	52.6	16	17.8
	Fem Pres Het Cont	0.0	0.0	0.3	100.0	0.3	12.5	0.0	0.0	0.3	100.0	0.3	16.7	0	0.0	4	21.1	4	4.4
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	2.8	1	5.3	3	3.3
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	1.4	0	0.0	1	1.1
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	5	7.0	0	0.0	5	5.6

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

² Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		1	
1986		2	2
1987		3	3
1988	•	4	3
1989		4	
1990		4	3
1991		5	3
1992		9	6
1993		5	4
1994		3	3
1995		2	5
1996		5	5
1997		3	1
1998	•	8	1
1999		5	3
2000*	6	3	
2001	5	1	1
2002	1	2	3
2003	2	2	
2004	1	2	2
2005	2	2	2
2006	2	1	
2007	2	3	
2008	3	5	3
2009	3	1	
2010	3	3	1
2011	2	2	

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		1,142	100.0	76.2	1,708	100.0	114.0	2,850	100.0	190.2	1,498,816
Gender	Male	754	66.0		1,179	69.0		1,933	67.8		
	Female	388	34.0		529	31.0		917	32.2		
Current Age	12 & under	12	1.1		0	0.0		12	0.4		
	13-19	26	2.3		10	0.6		36	1.3		
	20-24	68	6.0		39	2.3		107	3.8		
	25-29	96	8.4		47	2.8		143	5.0		
	30-39	206	18.0		207	12.1		413	14.5		
	40-49	341	29.9		537	31.4		878	30.8		
	50-59	276	24.2		599	35.1		875	30.7		
	60+	114	10.0		269	15.7		383	13.4		
	Unk	3	0.3		0	0.0		3	0.1		
Race/Ethnicity	White	472	41.3		641	37.5		1,113	39.1		
	Black	293	25.7		468	27.4		761	26.7		
	Hispanic	285	25.0		448	26.2		733	25.7		
	Multi Race	76	6.7		141	8.3		217	7.6		
	Others	13	1.1		10	0.6		23	0.8		
	Unk	3	0.3		0	0.0		3	0.1		
Risk	MSM	484	42.4		669	39.2		1,153	40.5		
	IDU	96	8.4		307	18.0		403	14.1		
	MSM/IDU	34	3.0		76	4.4		110	3.9		
	Heterosexual	235	20.6		309	18.1		544	19.1		
	Fem Pres Het Cont	109	9.5		124	7.3		233	8.2		
	Blood Prod	6	0.5		8	0.5		14	0.5		
	Pediatric Risk	49	4.3		46	2.7		95	3.3		
	Unk	129	11.3		169	9.9		298	10.5		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	Т	otal	M	Iale	Fe	male	Т	otal	M	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		754	100.0	388	100.0	1,142	100.0	1,179	100.0	529	100.0	1,708	100.0	1,933	100.0	917	100.0	2,850	100.0
Risk	MSM	484	64.2	0	0.0	484	42.4	669	56.7	0	0.0	669	39.2	1,153	59.6	0	0.0	1,153	40.5
	IDU	49	6.5	47	12.1	96	8.4	182	15.4	125	23.6	307	18.0	231	12.0	172	18.8	403	14.1
	MSM/IDU	34	4.5	0	0.0	34	3.0	76	6.4	0	0.0	76	4.4	110	5.7	0	0.0	110	3.9
	Heterosexual	34	4.5	201	51.8	235	20.6	51	4.3	258	48.8	309	18.1	85	4.4	459	50.1	544	19.1
	Fem Pres Het Cont	0	0.0	109	28.1	109	9.5	0	0.0	124	23.4	124	7.3	0	0.0	233	25.4	233	8.2
	Blood Prod	4	0.5	2	0.5	6	0.5	7	0.6	1	0.2	8	0.5	11	0.6	3	0.3	14	0.5
	Pediatric Risk	22	2.9	27	7.0	49	4.3	26	2.2	20	3.8	46	2.7	48	2.5	47	5.1	95	3.3
	Unk	127	16.8	2	0.5	129	11.3	168	14.2	1	0.2	169	9.9	295	15.3	3	0.3	298	10.5

		HIV Diagn	oses (Annua	l Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		108.3	100.0	7.2	72.3	100.0	4.8	4,384	100.0	1,498,816
Gender	Male	85.0	78.5		51.3	71.0		3,208	73.2	
	Female	23.3	21.5		21.0	29.0		1,176	26.8	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		67	1.5	
	13-19	5.0	4.6		0.3	0.5		52	1.2	
	20-24	19.0	17.5		5.0	6.9		127	2.9	
	25-29	13.7	12.6		6.0	8.3		389	8.9	
	30-39	26.0	24.0		16.7	23.0		1,730	39.5	
	40-49	27.3	25.2		24.7	34.1		1,396	31.8	
	50-59	13.3	12.3		14.7	20.3		470	10.7	
	60+	4.0	3.7		5.0	6.9		153	3.5	
	Unk	0.0	0.0		0.0	0.0		0	0.0	
Race/Ethnicity	White	36.0	33.2		23.0	31.8		2,068	47.2	
	Black	27.7	25.5		20.7	28.6		1,268	28.9	
	Hispanic	33.3	30.8		20.7	28.6		851	19.4	
	Multi Race	8.7	8.0		7.3	10.1		177	4.0	
	Others	2.7	2.5		0.7	0.9		20	0.5	
	Unk	0.0	0.0		0.0	0.0		0	0.0	
Risk	MSM	61.7	56.9		35.3	48.8		1,516	34.6	
	IDU	2.3	2.2		6.0	8.3		1,361	31.0	
	MSM/IDU	3.0	2.8		2.7	3.7		251	5.7	
	Heterosexual	16.7	15.4		13.7	18.9		549	12.5	
	Fem Pres Het Cont	9.3	8.6		5.0	6.9		208	4.7	
	Blood Prod	0.0	0.0		0.0	0.0		62	1.4	
	Pediatric Risk	0.0	0.0		1.0	1.4		98	2.2	
	Unk	15.3	14.2		8.7	12.0		339	7.7	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		85.0	100.0	23.3	100.0	108.3	100.0	51.3	100.0	21.0	100.0	72.3	100.0	3,208	100.0	1,176	100.0	4,384	100.0
Risk	MSM	61.7	72.5	0.0	0.0	61.7	56.9	35.3	68.8	0.0	0.0	35.3	48.8	1,516	47.3	0	0.0	1,516	34.6
	IDU	1.0	1.2	1.3	5.7	2.3	2.2	2.0	3.9	4.0	19.0	6.0	8.3	915	28.5	446	37.9	1,361	31.0
	MSM/IDU	3.0	3.5	0.0	0.0	3.0	2.8	2.7	5.2	0.0	0.0	2.7	3.7	251	7.8	0	0.0	251	5.7
	Heterosexual	4.0	4.7	12.7	54.3	16.7	15.4	2.3	4.5	11.3	54.0	13.7	18.9	88	2.7	461	39.2	549	12.5
	Fem Pres Het Cont	0.0	0.0	9.3	40.0	9.3	8.6	0.0	0.0	5.0	23.8	5.0	6.9	0	0.0	208	17.7	208	4.7
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	47	1.5	15	1.3	62	1.4
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.7	1.3	0.3	1.6	1.0	1.4	54	1.7	44	3.7	98	2.2
	Unk	15.3	18.0	0.0	0.0	15.3	14.2	8.3	16.2	0.3	1.6	8.7	12.0	337	10.5	2	0.2	339	7.7

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		104	59
1986		103	70
1987		145	81
1988		190	126
1989		189	136
1990		210	142
1991		273	166
1992		270	211
1993		335	196
1994		301	226
1995		313	207
1996		252	156
1997		186	94
1998		137	79
1999		136	74
2000*	86	138	57
2001	124	123	67
2002	121	118	71
2003	143	132	66
2004	136	108	64
2005	123	101	77
2006	111	96	55
2007	126	105	53
2008	108	102	42
2009	108	77	38
2010	113	63	39
2011	104	77	34

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		106	100.0	137.8	168	100.0	218.5	274	100.0	356.3	76,900
Gender	Male	61	57.5		95	56.5		156	56.9		
	Female	45	42.5		73	43.5		118	43.1		
Current Age	12 & under	1	0.9		0	0.0		1	0.4		
	13-19	2	1.9		1	0.6		3	1.1		
	20-24	6	5.7		0	0.0		6	2.2		
	25-29	12	11.3		2	1.2		14	5.1		
	30-39	12	11.3		12	7.1		24	8.8		
	40-49	33	31.1		44	26.2		77	28.1		
	50-59	29	27.4		67	39.9		96	35.0		
	60+	11	10.4		42	25.0		53	19.3		
Race/Ethnicity	White	27	25.5		42	25.0		69	25.2		
	Black	34	32.1		50	29.8		84	30.7		
	Hispanic	29	27.4		47	28.0		76	27.7		
	Multi Race	13	12.3		29	17.3		42	15.3		
	Others	3	2.8		0	0.0		3	1.1		
Risk	MSM	22	20.8		31	18.5		53	19.3		
	IDU	31	29.2		63	37.5		94	34.3		
	MSM/IDU	8	7.5		6	3.6		14	5.1		
	Heterosexual	31	29.2		44	26.2		75	27.4		
	Fem Pres Het Cont	7	6.6		10	6.0		17	6.2		
	Blood Prod	0	0.0		1	0.6		1	0.4		
	Pediatric Risk	3	2.8		3	1.8		6	2.2		
	Unk	4	3.8		10	6.0		14	5.1		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	Т	otal	N	Male	Fe	emale	Т	otal	M	[ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		61	100.0	45	100.0	106	100.0	95	100.0	73	100.0	168	100.0	156	100.0	118	100.0	274	100.0
Risk	MSM	22	36.1	0	0.0	22	20.8	31	32.6	0	0.0	31	18.5	53	34.0	0	0.0	53	19.3
	IDU	19	31.1	12	26.7	31	29.2	44	46.3	19	26.0	63	37.5	63	40.4	31	26.3	94	34.3
	MSM/IDU	8	13.1	0	0.0	8	7.5	6	6.3	0	0.0	6	3.6	14	9.0	0	0.0	14	5.1
	Heterosexual	8	13.1	23	51.1	31	29.2	4	4.2	40	54.8	44	26.2	12	7.7	63	53.4	75	27.4
	Fem Pres Het Cont	0	0.0	7	15.6	7	6.6	0	0.0	10	13.7	10	6.0	0	0.0	17	14.4	17	6.2
	Blood Prod	0	0.0	0	0.0	0	0.0	0	0.0	1	1.4	1	0.6	0	0.0	1	0.8	1	0.4
	Pediatric Risk	0	0.0	3	6.7	3	2.8	0	0.0	3	4.1	3	1.8	0	0.0	6	5.1	6	2.2
	Unk	4	6.6	0	0.0	4	3.8	10	10.5	0	0.0	10	6.0	14	9.0	0	0.0	14	5.1

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		2.7	100.0	3.5	3.3	100.0	4.3	509	100.0	76,900
Gender	Male	1.7	62.5		2.0	60.0		360	70.7	
	Female	1.0	37.5		1.3	40.0		149	29.3	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		6	1.2	
	13-19	0.0	0.0		0.0	0.0		3	0.6	
	20-24	0.7	25.0		0.0	0.0		8	1.6	
	25-29	0.7	25.0		0.0	0.0		44	8.6	
	30-39	0.3	12.5		0.3	10.0		188	36.9	
	40-49	0.7	25.0		1.3	40.0		185	36.3	
	50-59	0.3	12.5		0.3	10.0		53	10.4	
	60+	0.0	0.0		1.3	40.0		22	4.3	
Race/Ethnicity	White	0.0	0.0		0.7	20.0		149	29.3	
	Black	1.3	50.0		1.3	40.0		189	37.1	
	Hispanic	0.7	25.0		1.3	40.0		133	26.1	
	Multi Race	0.0	0.0		0.0	0.0		37	7.3	
	Others	0.7	25.0		0.0	0.0		1	0.2	
Risk	MSM	1.0	37.5		0.0	0.0		87	17.1	
	IDU	0.7	25.0		2.7	80.0		257	50.5	
	MSM/IDU	0.3	12.5		0.0	0.0		18	3.5	
	Heterosexual	0.7	25.0		0.3	10.0		84	16.5	
	Fem Pres Het Cont	0.0	0.0		0.3	10.0		23	4.5	
	Blood Prod	0.0	0.0		0.0	0.0		5	1.0	
	Pediatric Risk	0.0	0.0		0.0	0.0		9	1.8	
	Unk	0.0	0.0		0.0	0.0		26	5.1	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		1	HIV Diag	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS C	Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		1.7	100.0	1.0	100.0	2.7	100.0	2.0	100.0	1.3	100.0	3.3	100.0	360	100.0	149	100.0	509	100.0
Risk	MSM	1.0	60.0	0.0	0.0	1.0	37.5	0.0	0.0	0.0	0.0	0.0	0.0	87	24.2	0	0.0	87	17.1
	IDU	0.3	20.0	0.3	33.3	0.7	25.0	2.0	100.0	0.7	50.0	2.7	80.0	212	58.9	45	30.2	257	50.5
	MSM/IDU	0.3	20.0	0.0	0.0	0.3	12.5	0.0	0.0	0.0	0.0	0.0	0.0	18	5.0	0	0.0	18	3.5
	Heterosexual	0.0	0.0	0.7	66.7	0.7	25.0	0.0	0.0	0.3	25.0	0.3	10.0	12	3.3	72	48.3	84	16.5
	Fem Pres Het Cont	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	25.0	0.3	10.0	0	0.0	23	15.4	23	4.5
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	0.6	3	2.0	5	1.0
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3	0.8	6	4.0	9	1.8
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	26	7.2	0	0.0	26	5.1

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		10	5
1986		10	4
1987		13	12
1988		23	17
1989		26	19
1990		21	16
1991		31	17
1992		45	26
1993		63	22
1994		49	37
1995		33	43
1996		20	22
1997		13	13
1998		12	6
1999		18	6
2000*	12	14	16
2001	14	15	10
2002	10	19	8
2003	15	12	9
2004	19	14	4
2005	6	11	6
2006	8	9	6
2007	10	9	1
2008	10	9	6
2009	1	4	1
2010	3	1	5
2011	4	5	4

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Living HIV (not A) Number Col % I 16 100.0 1 11 68.8 5 5 31.3 0 0.0 3 18.8 5 31.3 4 25.0 2 12.5 2 12.5 14 87.5 1 6.3 0 0.0 1 6.3 7 43.8 0 0.0 1 6.3		AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		16	100.0	31.3	20	100.0	39.2	36	100.0	70.5	51,043
Gender	Male	11	68.8		14	70.0		25	69.4		
	Female	5	31.3		6	30.0		11	30.6		
Current Age	13-19	0	0.0		0	0.0		0	0.0		
	20-24	0	0.0		1	5.0		1	2.8		
	25-29	3	18.8		0	0.0		3	8.3		
	30-39	5	31.3		2	10.0		7	19.4		
	40-49	4	25.0		8	40.0		12	33.3		
	50-59	2	12.5		7	35.0		9	25.0		
	60+	2	12.5		2	10.0		4	11.1		
Race/Ethnicity	White	14	87.5		14	70.0		28	77.8		
	Black	1	6.3		1	5.0		2	5.6		
	Hispanic	0	0.0		4	20.0		4	11.1		
	Multi Race	1	6.3		1	5.0		2	5.6		
Risk	MSM	7	43.8		8	40.0		15	41.7		
	IDU	0	0.0		4	20.0		4	11.1		
	MSM/IDU	1	6.3		1	5.0		2	5.6		
	Heterosexual	3	18.8		4	20.0		7	19.4		
	Fem Pres Het Cont	2	12.5		1	5.0		3	8.3		
	Pediatric Risk	0	0.0		1	5.0		1	2.8		
	Unk	3	18.8		1	5.0		4	11.1		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		M	ale	Fe	male	Т	otal	N	Aale	Fe	male	Т	otal	M	[ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		11	100.0	5	100.0	16	100.0	14	100.0	6	100.0	20	100.0	25	100.0	11	100.0	36	100.0
Risk	MSM	7	63.6	0	0.0	7	43.8	8	57.1	0	0.0	8	40.0	15	60.0	0	0.0	15	41.7
	IDU	0	0.0	0	0.0	0	0.0	3	21.4	1	16.7	4	20.0	3	12.0	1	9.1	4	11.1
	MSM/IDU	1	9.1	0	0.0	1	6.3	1	7.1	0	0.0	1	5.0	2	8.0	0	0.0	2	5.6
	Heterosexual	0	0.0	3	60.0	3	18.8	1	7.1	3	50.0	4	20.0	1	4.0	6	54.5	7	19.4
	Fem Pres Het Cont	0	0.0	2	40.0	2	12.5	0	0.0	1	16.7	1	5.0	0	0.0	3	27.3	3	8.3
	Pediatric Risk	0	0.0	0	0.0	0	0.0	0	0.0	1	16.7	1	5.0	0	0.0	1	9.1	1	2.8
	Unk	3	27.3	0	0.0	3	18.8	1	7.1	0	0.0	1	5.0	4	16.0	0	0.0	4	11.1

		HIV Diagn	oses (Annua	ıl Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		1.0	100.0	2.0	0.7	100.0	1.3	37	100.0	51,043
Gender	Male	0.7	66.7		0.3	50.0		31	83.8	
	Female	0.3	33.3		0.3	50.0		6	16.2	
Age at Diagnosis	13-19	0.0	0.0		0.0	0.0		1	2.7	
	20-24	0.0	0.0		0.0	0.0		2	5.4	
	25-29	0.0	0.0		0.0	0.0		2	5.4	
	30-39	0.3	33.3		0.0	0.0		14	37.8	
	40-49	0.3	33.3		0.3	50.0		15	40.5	
	50-59	0.0	0.0		0.0	0.0		2	5.4	
	60+	0.3	33.3		0.3	50.0		1	2.7	
Race/Ethnicity	White	0.7	66.7		0.3	50.0		31	83.8	
	Black	0.0	0.0		0.0	0.0		2	5.4	
	Hispanic	0.3	33.3		0.3	50.0		3	8.1	
	Multi Race	0.0	0.0		0.0	0.0		1	2.7	
Risk	MSM	0.7	66.7		0.3	50.0		20	54.1	
	IDU	0.0	0.0		0.0	0.0		6	16.2	
	MSM/IDU	0.0	0.0		0.0	0.0		4	10.8	
	Heterosexual	0.0	0.0		0.0	0.0		3	8.1	
	Fem Pres Het Cont	0.3	33.3		0.3	50.0		2	5.4	
	Pediatric Risk	0.0	0.0		0.0	0.0		1	2.7	
	Unk	0.0	0.0		0.0	0.0		1	2.7	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		1	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fei	nale	To	otal	М	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		0.7	100.0	0.3	100.0	1.0	100.0	0.3	100.0	0.3	100.0	0.7	100.0	31	100.0	6	100.0	37	100.0
Risk	MSM	0.7	100.0	0.0	0.0	0.7	66.7	0.3	100.0	0.0	0.0	0.3	50.0	20	64.5	0	0.0	20	54.1
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	5	16.1	1	16.7	6	16.2
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	4	12.9	0	0.0	4	10.8
	Heterosexual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	3.2	2	33.3	3	8.1
	Fem Pres Het Cont	0.0	0.0	0.3	100.0	0.3	33.3	0.0	0.0	0.3	100.0	0.3	50.0	0	0.0	2	33.3	2	5.4
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	1	16.7	1	2.7
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	3.2	0	0.0	1	2.7

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		1	1
1987		1	
1988		2	
1989		1	2
1990		3	2
1991		3	
1992		1	2
1993		2	4
1994		1	1
1995		4	
1996		1	2
1997			1
2000*	1	1	
2001	4	2	1
2002			1
2003	2	4	
2004	2	1	
2005	3	1	
2006	1	٠	
2007		1	
2008	3	5	
2009	1	1	
2010	1	1	1
2011	1		

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		50	100.0	49.2	83	100.0	81.6	133	100.0	130.7	101,723
Gender	Male	30	60.0		62	74.7		92	69.2		
	Female	20	40.0		21	25.3		41	30.8		
Current Age	12 & under	1	2.0		0	0.0		1	0.8		
	13-19	1	2.0		0	0.0		1	0.8		
	20-24	2	4.0		1	1.2		3	2.3		
	25-29	4	8.0		5	6.0		9	6.8		
	30-39	11	22.0		9	10.8		20	15.0		
	40-49	17	34.0		36	43.4		53	39.8		
	50-59	10	20.0		23	27.7		33	24.8		
	60+	4	8.0		9	10.8		13	9.8		
Race/Ethnicity	White	30	60.0		50	60.2		80	60.2		
	Black	10	20.0		13	15.7		23	17.3		
	Hispanic	5	10.0		12	14.5		17	12.8		
	Multi Race	1	2.0		7	8.4		8	6.0		
	Others	4	8.0		1	1.2		5	3.8		
Risk	MSM	21	42.0		39	47.0		60	45.1		
	IDU	1	2.0		6	7.2		7	5.3		
	MSM/IDU	2	4.0		9	10.8		11	8.3		
	Heterosexual	9	18.0		13	15.7		22	16.5		
	Fem Pres Het Cont	9	18.0		8	9.6		17	12.8		
	Blood Prod	0	0.0		1	1.2		1	0.8		
	Pediatric Risk	2	4.0		0	0.0		2	1.5		
	Unk	6	12.0		7	8.4		13	9.8		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	IDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	Male	Fe	male	Т	otal	M	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		30	100.0	20	100.0	50	100.0	62	100.0	21	100.0	83	100.0	92	100.0	41	100.0	133	100.0
Risk	MSM	21	70.0	0	0.0	21	42.0	39	62.9	0	0.0	39	47.0	60	65.2	0	0.0	60	45.1
	IDU	0	0.0	1	5.0	1	2.0	4	6.5	2	9.5	6	7.2	4	4.3	3	7.3	7	5.3
	MSM/IDU	2	6.7	0	0.0	2	4.0	9	14.5	0	0.0	9	10.8	11	12.0	0	0.0	11	8.3
	Heterosexual	1	3.3	8	40.0	9	18.0	2	3.2	11	52.4	13	15.7	3	3.3	19	46.3	22	16.5
	Fem Pres Het Cont	0	0.0	9	45.0	9	18.0	0	0.0	8	38.1	8	9.6	0	0.0	17	41.5	17	12.8
	Blood Prod	0	0.0	0	0.0	0	0.0	1	1.6	0	0.0	1	1.2	1	1.1	0	0.0	1	0.8
	Pediatric Risk	1	3.3	1	5.0	2	4.0	0	0.0	0	0.0	0	0.0	1	1.1	1	2.4	2	1.5
	Unk	5	16.7	1	5.0	6	12.0	7	11.3	0	0.0	7	8.4	12	13.0	1	2.4	13	9.8

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		4.0	100.0	3.9	2.7	100.0	2.6	151	100.0	101,723
Gender	Male	3.3	83.3		1.3	50.0		118	78.1	
	Female	0.7	16.7		1.3	50.0		33	21.9	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		0	0.0	
	13-19	0.0	0.0		0.0	0.0		2	1.3	
	20-24	0.3	8.3		0.0	0.0		10	6.6	
	25-29	0.3	8.3		0.3	12.5		24	15.9	
	30-39	0.7	16.7		0.3	12.5		55	36.4	
	40-49	1.3	33.3		1.0	37.5		46	30.5	
	50-59	1.0	25.0		0.7	25.0		10	6.6	
	60+	0.3	8.3		0.3	12.5		4	2.6	
Race/Ethnicity	White	2.7	66.7		1.7	62.5		94	62.3	
	Black	0.7	16.7		0.3	12.5		31	20.5	
	Hispanic	0.0	0.0		0.0	0.0		18	11.9	
	Multi Race	0.3	8.3		0.7	25.0		7	4.6	
	Others	0.3	8.3		0.0	0.0		1	0.7	
Risk	MSM	2.3	58.3		1.0	37.5		69	45.7	
	IDU	0.3	8.3		0.0	0.0		27	17.9	
	MSM/IDU	0.0	0.0		0.0	0.0		12	7.9	
	Heterosexual	0.3	8.3		0.7	25.0		16	10.6	
	Fem Pres Het Cont	0.3	8.3		0.7	25.0		11	7.3	
	Blood Prod	0.0	0.0		0.0	0.0		5	3.3	
	Pediatric Risk	0.0	0.0		0.0	0.0		0	0.0	
	Unk	0.7	16.7		0.3	12.5		11	7.3	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	То	otal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		3.3	100.0	0.7	100.0	4.0	100.0	1.3	100.0	1.3	100.0	2.7	100.0	118	100.0	33	100.0	151	100.0
Risk	MSM	2.3	70.0	0.0	0.0	2.3	58.3	1.0	75.0	0.0	0.0	1.0	37.5	69	58.5	0	0.0	69	45.7
	IDU	0.3	10.0	0.0	0.0	0.3	8.3	0.0	0.0	0.0	0.0	0.0	0.0	18	15.3	9	27.3	27	17.9
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	12	10.2	0	0.0	12	7.9
	Heterosexual	0.0	0.0	0.3	50.0	0.3	8.3	0.0	0.0	0.7	50.0	0.7	25.0	4	3.4	12	36.4	16	10.6
	Fem Pres Het Cont	0.0	0.0	0.3	50.0	0.3	8.3	0.0	0.0	0.7	50.0	0.7	25.0	0	0.0	11	33.3	11	7.3
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	4	3.4	1	3.0	5	3.3
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	0	0.0	0	0.0
	Unk	0.7	20.0	0.0	0.0	0.7	16.7	0.3	25.0	0.0	0.0	0.3	12.5	11	9.3	0	0.0	11	7.3

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		2	1
1986		2	2
1987		4	2
1988		5	5
1989		5	2
1990		8	4
1991		7	6
1992		12	5
1993		5	5
1994		10	7
1995		7	4
1996		13	5
1997		7	4
1998		7	3
1999		5	2
2000*	4	5	1
2001	2	4	2
2002	10	8	
2003	4	3	2
2004	6	3	2
2005	5	4	1
2006	7	4	2
2007	5	9	
2008	6	4	1
2009	4	4	1
2010	5	3	2
2011	3	1	

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		144	100.0	78.9	243	100.0	133.2	387	100.0	212.1	182,448
Gender	Male	87	60.4		159	65.4		246	63.6		
	Female	57	39.6		84	34.6		141	36.4		
Current Age	12 & under	0	0.0		0	0.0		0	0.0		
	13-19	2	1.4		1	0.4		3	0.8		
	20-24	3	2.1		4	1.6		7	1.8		
	25-29	5	3.5		5	2.1		10	2.6		
	30-39	25	17.4		18	7.4		43	11.1		
	40-49	47	32.6		83	34.2		130	33.6		
	50-59	41	28.5		85	35.0		126	32.6		
	60+	21	14.6		47	19.3		68	17.6		
Race/Ethnicity	White	97	67.4		112	46.1		209	54.0		
	Black	22	15.3		48	19.8		70	18.1		
	Hispanic	14	9.7		55	22.6		69	17.8		
	Multi Race	11	7.6		27	11.1		38	9.8		
	Others	0	0.0		1	0.4		1	0.3		
Risk	MSM	42	29.2		63	25.9		105	27.1		
	IDU	31	21.5		68	28.0		99	25.6		
	MSM/IDU	7	4.9		14	5.8		21	5.4		
	Heterosexual	43	29.9		64	26.3		107	27.6		
	Fem Pres Het Cont	9	6.3		13	5.3		22	5.7		
	Blood Prod	0	0.0		0	0.0		0	0.0		
	Pediatric Risk	1	0.7		6	2.5		7	1.8		
	Unk	11	7.6		15	6.2		26	6.7		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	/Iale	Fe	male	Т	otal	M	[ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		87	100.0	57	100.0	144	100.0	159	100.0	84	100.0	243	100.0	246	100.0	141	100.0	387	100.0
Risk	MSM	42	48.3	0	0.0	42	29.2	63	39.6	0	0.0	63	25.9	105	42.7	0	0.0	105	27.1
	IDU	19	21.8	12	21.1	31	21.5	48	30.2	20	23.8	68	28.0	67	27.2	32	22.7	99	25.6
	MSM/IDU	7	8.0	0	0.0	7	4.9	14	8.8	0	0.0	14	5.8	21	8.5	0	0.0	21	5.4
	Heterosexual	7	8.0	36	63.2	43	29.9	18	11.3	46	54.8	64	26.3	25	10.2	82	58.2	107	27.6
	Fem Pres Het Cont	0	0.0	9	15.8	9	6.3	0	0.0	13	15.5	13	5.3	0	0.0	22	15.6	22	5.7
	Blood Prod	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Pediatric Risk	1	1.1	0	0.0	1	0.7	1	0.6	5	6.0	6	2.5	2	0.8	5	3.5	7	1.8
	Unk	11	12.6	0	0.0	11	7.6	15	9.4	0	0.0	15	6.2	26	10.6	0	0.0	26	6.7

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		10.7	100.0	5.8	9.7	100.0	5.3	641	100.0	182,448
Gender	Male	8.0	75.0		7.3	75.9		484	75.5	
	Female	2.7	25.0		2.3	24.1		157	24.5	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		9	1.4	
	13-19	0.3	3.1		0.3	3.4		6	0.9	
	20-24	1.0	9.4		0.3	3.4		11	1.7	
	25-29	1.7	15.6		1.0	10.3		65	10.1	
	30-39	1.7	15.6		0.7	6.9		256	39.9	
	40-49	3.7	34.4		4.0	41.4		207	32.3	
	50-59	2.0	18.8		3.0	31.0		65	10.1	
	60+	0.3	3.1		0.3	3.4		22	3.4	
Race/Ethnicity	White	7.0	65.6		5.3	55.2		338	52.7	
	Black	1.3	12.5		2.3	24.1		127	19.8	
	Hispanic	2.0	18.8		1.7	17.2		137	21.4	
	Multi Race	0.3	3.1		0.3	3.4		38	5.9	
	Others	0.0	0.0		0.0	0.0		1	0.2	
Risk	MSM	4.0	37.5		3.3	34.5		181	28.2	
	IDU	1.0	9.4		1.3	13.8		251	39.2	
	MSM/IDU	0.0	0.0		0.0	0.0		38	5.9	
	Heterosexual	3.7	34.4		3.0	31.0		96	15.0	
	Fem Pres Het Cont	0.3	3.1		0.3	3.4		19	3.0	
	Blood Prod	0.0	0.0		0.0	0.0		2	0.3	
	Pediatric Risk	0.0	0.0		0.3	3.4		11	1.7	
	Unk	1.7	15.6		1.3	13.8		43	6.7	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		1	HIV Diag	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		8.0	100.0	2.7	100.0	10.7	100.0	7.3	100.0	2.3	100.0	9.7	100.0	484	100.0	157	100.0	641	100.0
Risk	MSM	4.0	50.0	0.0	0.0	4.0	37.5	3.3	45.5	0.0	0.0	3.3	34.5	181	37.4	0	0.0	181	28.2
	IDU	1.0	12.5	0.0	0.0	1.0	9.4	1.3	18.2	0.0	0.0	1.3	13.8	190	39.3	61	38.9	251	39.2
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	38	7.9	0	0.0	38	5.9
	Heterosexual	1.3	16.7	2.3	87.5	3.7	34.4	1.3	18.2	1.7	71.4	3.0	31.0	27	5.6	69	43.9	96	15.0
	Fem Pres Het Cont	0.0	0.0	0.3	12.5	0.3	3.1	0.0	0.0	0.3	14.3	0.3	3.4	0	0.0	19	12.1	19	3.0
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	0.4	0	0.0	2	0.3
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	14.3	0.3	3.4	3	0.6	8	5.1	11	1.7
	Unk	1.7	20.8	0.0	0.0	1.7	15.6	1.3	18.2	0.0	0.0	1.3	13.8	43	8.9	0	0.0	43	6.7

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		7	5
1986		7	4
1987		20	9
1988		19	10
1989		38	14
1990		40	23
1991		50	24
1992		47	32
1993		57	36
1994		56	35
1995		51	30
1996		21	36
1997		21	18
1998		20	7
1999	•	25	8
2000*	10	22	13
2001	16	22	14
2002	16	11	10
2003	18	19	7
2004	16	18	11
2005	17	9	5
2006	19	16	6
2007	5	8	8
2008	9	8	12
2009	11	15	7
2010	9	5	6
2011	12	9	8

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		29	100.0	44.1	37	100.0	56.2	66	100.0	100.3	65,831
Gender	Male	21	72.4		28	75.7		49	74.2		
	Female	8	27.6		9	24.3		17	25.8		
Current Age	12 & under	1	3.4		0	0.0		1	1.5		
	13-19	1	3.4		0	0.0		1	1.5		
	20-24	2	6.9		3	8.1		5	7.6		
	25-29	0	0.0		1	2.7		1	1.5		
	30-39	6	20.7		4	10.8		10	15.2		
	40-49	10	34.5		12	32.4		22	33.3		
	50-59	7	24.1		13	35.1		20	30.3		
	60+	2	6.9		4	10.8		6	9.1		
Race/Ethnicity	White	21	72.4		23	62.2		44	66.7		
	Black	0	0.0		2	5.4		2	3.0		
	Hispanic	6	20.7		8	21.6		14	21.2		
	Multi Race	1	3.4		4	10.8		5	7.6		
	Others	1	3.4		0	0.0		1	1.5		
Risk	MSM	15	51.7		15	40.5		30	45.5		
	IDU	2	6.9		6	16.2		8	12.1		
	MSM/IDU	0	0.0		0	0.0		0	0.0		
	Heterosexual	5	17.2		8	21.6		13	19.7		
	Fem Pres Het Cont	2	6.9		3	8.1		5	7.6		
	Blood Prod	0	0.0		2	5.4		2	3.0		
	Pediatric Risk	3	10.3		3	8.1		6	9.1		
	Unk	2	6.9		0	0.0		2	3.0		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	Male	Fe	male	Т	otal	M	[ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		21	100.0	8	100.0	29	100.0	28	100.0	9	100.0	37	100.0	49	100.0	17	100.0	66	100.0
Risk	MSM	15	71.4	0	0.0	15	51.7	15	53.6	0	0.0	15	40.5	30	61.2	0	0.0	30	45.5
	IDU	2	9.5	0	0.0	2	6.9	6	21.4	0	0.0	6	16.2	8	16.3	0	0.0	8	12.1
	MSM/IDU	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Heterosexual	0	0.0	5	62.5	5	17.2	3	10.7	5	55.6	8	21.6	3	6.1	10	58.8	13	19.7
	Fem Pres Het Cont	0	0.0	2	25.0	2	6.9	0	0.0	3	33.3	3	8.1	0	0.0	5	29.4	5	7.6
	Blood Prod	0	0.0	0	0.0	0	0.0	2	7.1	0	0.0	2	5.4	2	4.1	0	0.0	2	3.0
	Pediatric Risk	2	9.5	1	12.5	3	10.3	2	7.1	1	11.1	3	8.1	4	8.2	2	11.8	6	9.1
	Unk	2	9.5	0	0.0	2	6.9	0	0.0	0	0.0	0	0.0	2	4.1	0	0.0	2	3.0

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		1.3	100.0	2.0	2.0	100.0	3.0	77	100.0	65,831
Gender	Male	0.3	25.0		0.7	33.3		65	84.4	
	Female	1.0	75.0		1.3	66.7		12	15.6	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		2	2.6	
	13-19	0.0	0.0		0.0	0.0		1	1.3	
	20-24	0.0	0.0		0.3	16.7		3	3.9	
	25-29	0.3	25.0		0.3	16.7		6	7.8	
	30-39	0.3	25.0		0.3	16.7		32	41.6	
	40-49	0.7	50.0		0.7	33.3		25	32.5	
	50-59	0.0	0.0		0.3	16.7		6	7.8	
	60+	0.0	0.0		0.0	0.0		2	2.6	
Race/Ethnicity	White	1.3	100.0		1.7	83.3		57	74.0	
	Black	0.0	0.0		0.3	16.7		7	9.1	
	Hispanic	0.0	0.0		0.0	0.0		7	9.1	
	Multi Race	0.0	0.0		0.0	0.0		6	7.8	
	Others	0.0	0.0		0.0	0.0		0	0.0	
Risk	MSM	0.3	25.0		0.3	16.7		36	46.8	
	IDU	0.0	0.0		0.0	0.0		14	18.2	
	MSM/IDU	0.0	0.0		0.0	0.0		2	2.6	
	Heterosexual	0.7	50.0		1.0	50.0		9	11.7	
	Fem Pres Het Cont	0.3	25.0		0.3	16.7		6	7.8	
	Blood Prod	0.0	0.0		0.0	0.0		5	6.5	
	Pediatric Risk	0.0	0.0		0.3	16.7		3	3.9	
	Unk	0.0	0.0		0.0	0.0		2	2.6	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

]	HIV Diag	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS C	Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		0.3	100.0	1.0	100.0	1.3	100.0	0.7	100.0	1.3	100.0	2.0	100.0	65	100.0	12	100.0	77	100.0
Risk	MSM	0.3	100.0	0.0	0.0	0.3	25.0	0.3	50.0	0.0	0.0	0.3	16.7	36	55.4	0	0.0	36	46.8
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	13	20.0	1	8.3	14	18.2
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	3.1	0	0.0	2	2.6
	Heterosexual	0.0	0.0	0.7	66.7	0.7	50.0	0.3	50.0	0.7	50.0	1.0	50.0	5	7.7	4	33.3	9	11.7
	Fem Pres Het Cont	0.0	0.0	0.3	33.3	0.3	25.0	0.0	0.0	0.3	25.0	0.3	16.7	0	0.0	6	50.0	6	7.8
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	5	7.7	0	0.0	5	6.5
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	25.0	0.3	16.7	2	3.1	1	8.3	3	3.9
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	3.1	0	0.0	2	2.6

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		2	2
1986		2	1
1987		1	2
1988		4	
1989		2	
1990		4	5
1991		2	5
1992		5	1
1993		3	1
1994		2	3
1995		7	1
1996		3	4
1997		3	
1998		5	
1999		2	
2000*	2	2	1
2001	4	4	5
2002	5	5	
2003	4		2
2004	4	5	1
2005	1	3	3
2006	1		1
2007	2	3	1
2008	1	2	3
2009	1	2	
2010	3	4	1
2011			1

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		18	100.0	28.5	18	100.0	28.5	36	100.0	57.0	63,165
Gender	Male	12	66.7		11	61.1		23	63.9		
	Female	6	33.3		7	38.9		13	36.1		
Current Age	12 & under	1	5.6		0	0.0		1	2.8		
	13-19	2	11.1		0	0.0		2	5.6		
	20-24	1	5.6		0	0.0		1	2.8		
	25-29	1	5.6		0	0.0		1	2.8		
	30-39	2	11.1		1	5.6		3	8.3		
	40-49	4	22.2		10	55.6		14	38.9		
	50-59	6	33.3		5	27.8		11	30.6		
	60+	1	5.6		2	11.1		3	8.3		
Race/Ethnicity	White	14	77.8		16	88.9		30	83.3		
	Black	2	11.1		0	0.0		2	5.6		
	Hispanic	2	11.1		2	11.1		4	11.1		
Risk	MSM	8	44.4		8	44.4		16	44.4		
	IDU	2	11.1		0	0.0		2	5.6		
	MSM/IDU	0	0.0		0	0.0		0	0.0		
	Heterosexual	2	11.1		6	33.3		8	22.2		
	Fem Pres Het Cont	1	5.6		1	5.6		2	5.6		
	Blood Prod	0	0.0		1	5.6		1	2.8		
	Pediatric Risk	2	11.1		0	0.0		2	5.6		
	Unk	3	16.7		2	11.1		5	13.9		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	Male	Fe	male	1	otal	M	ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		12	100.0	6	100.0	18	100.0	11	100.0	7	100.0	18	100.0	23	100.0	13	100.0	36	100.0
Risk	MSM	8	66.7	0	0.0	8	44.4	8	72.7	0	0.0	8	44.4	16	69.6	0	0.0	16	44.4
	IDU	0	0.0	2	33.3	2	11.1	0	0.0	0	0.0	0	0.0	0	0.0	2	15.4	2	5.6
	MSM/IDU	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Heterosexual	1	8.3	1	16.7	2	11.1	0	0.0	6	85.7	6	33.3	1	4.3	7	53.8	8	22.2
	Fem Pres Het Cont	0	0.0	1	16.7	1	5.6	0	0.0	1	14.3	1	5.6	0	0.0	2	15.4	2	5.6
	Blood Prod	0	0.0	0	0.0	0	0.0	1	9.1	0	0.0	1	5.6	1	4.3	0	0.0	1	2.8
	Pediatric Risk	0	0.0	2	33.3	2	11.1	0	0.0	0	0.0	0	0.0	0	0.0	2	15.4	2	5.6
	Unk	3	25.0	0	0.0	3	16.7	2	18.2	0	0.0	2	11.1	5	21.7	0	0.0	5	13.9

		HIV Diagn	oses (Annua	l Average)	AIDS Diagr	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		2.7	100.0	4.2	1.3	100.0	2.1	45	100.0	63,165
Gender	Male	2.3	87.5		1.3	100.0		39	86.7	
	Female	0.3	12.5		0.0	0.0		6	13.3	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		0	0.0	
	13-19	0.3	12.5		0.0	0.0	•	0	0.0	
	20-24	0.3	12.5		0.0	0.0		2	4.4	
	25-29	0.0	0.0		0.0	0.0		8	17.8	
	30-39	0.3	12.5		0.3	25.0		21	46.7	
	40-49	0.7	25.0		0.7	50.0		10	22.2	
	50-59	0.7	25.0		0.0	0.0		3	6.7	
	60+	0.3	12.5		0.3	25.0		1	2.2	
Race/Ethnicity	White	2.3	87.5		1.3	100.0		37	82.2	
	Black	0.3	12.5		0.0	0.0		2	4.4	
	Hispanic	0.0	0.0		0.0	0.0		6	13.3	
Risk	MSM	1.7	62.5		0.7	50.0		22	48.9	
	IDU	0.0	0.0		0.0	0.0		9	20.0	
	MSM/IDU	0.0	0.0		0.0	0.0		2	4.4	
	Heterosexual	0.7	25.0		0.3	25.0		7	15.6	
	Fem Pres Het Cont	0.0	0.0		0.0	0.0		1	2.2	
	Blood Prod	0.0	0.0		0.0	0.0		1	2.2	
	Pediatric Risk	0.0	0.0		0.0	0.0		0	0.0	
	Unk	0.3	12.5		0.3	25.0		3	6.7	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		I	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		2.3	100.0	0.3	100.0	2.7	100.0	1.3	100.0	0.0	0.0	1.3	100.0	39	100.0	6	100.0	45	100.0
Risk	MSM	1.7	71.4	0.0	0.0	1.7	62.5	0.7	50.0	0.0	0.0	0.7	50.0	22	56.4	0	0.0	22	48.9
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	9	23.1	0	0.0	9	20.0
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	5.1	0	0.0	2	4.4
	Heterosexual	0.3	14.3	0.3	100.0	0.7	25.0	0.3	25.0	0.0	0.0	0.3	25.0	2	5.1	5	83.3	7	15.6
	Fem Pres Het Cont	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	1	16.7	1	2.2
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	2.6	0	0.0	1	2.2
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	0	0.0	0	0.0
	Unk	0.3	14.3	0.0	0.0	0.3	12.5	0.3	25.0	0.0	0.0	0.3	25.0	3	7.7	0	0.0	3	6.7

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
1986		2	1
1987		2	1
1988		4	1
1989		1	3
1990		4	1
1991		3	3
1992		3	4
1993		1	3
1994		3	1
1995		3	1
1996			1
1997		3	2
1999		1	
2000*	1	3	
2001	1	2	1
2002		2	
2003	2		
2004	1	2	2
2005		1	1
2006	2		
2007		1	1
2008	1		
2009	1	1	
2010	3	2	1
2011	4	1	

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		55	100.0	58.9	52	100.0	55.7	107	100.0	114.5	93,436
Gender	Male	37	67.3		37	71.2		74	69.2		
	Female	18	32.7		15	28.8		33	30.8		
Current Age	12 & under	0	0.0		0	0.0		0	0.0		
	13-19	1	1.8		0	0.0		1	0.9		
	20-24	0	0.0		0	0.0		0	0.0		
	25-29	2	3.6		1	1.9		3	2.8		
	30-39	11	20.0		7	13.5		18	16.8		
	40-49	19	34.5		16	30.8		35	32.7		
	50-59	15	27.3		18	34.6		33	30.8		
	60+	7	12.7		10	19.2		17	15.9		
Race/Ethnicity	White	28	50.9		30	57.7		58	54.2		
	Black	15	27.3		12	23.1		27	25.2		
	Hispanic	8	14.5		8	15.4		16	15.0		
	Multi Race	4	7.3		2	3.8		6	5.6		
Risk	MSM	19	34.5		11	21.2		30	28.0		
	IDU	3	5.5		8	15.4		11	10.3		
	MSM/IDU	2	3.6		2	3.8		4	3.7		
	Heterosexual	10	18.2		18	34.6		28	26.2		
	Fem Pres Het Cont	9	16.4		3	5.8		12	11.2		
	Pediatric Risk	1	1.8		0	0.0		1	0.9		
	Unk	11	20.0		10	19.2		21	19.6		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	Male	Fe	male	Т	otal	M	[ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		37	100.0	18	100.0	55	100.0	37	100.0	15	100.0	52	100.0	74	100.0	33	100.0	107	100.0
Risk	MSM	19	51.4	0	0.0	19	34.5	11	29.7	0	0.0	11	21.2	30	40.5	0	0.0	30	28.0
	IDU	3	8.1	0	0.0	3	5.5	6	16.2	2	13.3	8	15.4	9	12.2	2	6.1	11	10.3
	MSM/IDU	2	5.4	0	0.0	2	3.6	2	5.4	0	0.0	2	3.8	4	5.4	0	0.0	4	3.7
	Heterosexual	2	5.4	8	44.4	10	18.2	8	21.6	10	66.7	18	34.6	10	13.5	18	54.5	28	26.2
	Fem Pres Het Cont	0	0.0	9	50.0	9	16.4	0	0.0	3	20.0	3	5.8	0	0.0	12	36.4	12	11.2
	Pediatric Risk	0	0.0	1	5.6	1	1.8	0	0.0	0	0.0	0	0.0	0	0.0	1	3.0	1	0.9
	Unk	11	29.7	0	0.0	11	20.0	10	27.0	0	0.0	10	19.2	21	28.4	0	0.0	21	19.6

		HIV Diagn	oses (Annua	l Average)	AIDS Diagr	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		4.7	100.0	5.0	3.3	100.0	3.6	107	100.0	93,436
Gender	Male	3.3	71.4		2.3	70.0		85	79.4	
	Female	1.3	28.6		1.0	30.0		22	20.6	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		1	0.9	
	13-19	0.0	0.0		0.0	0.0	•	0	0.0	
	20-24	0.7	14.3		0.3	10.0		4	3.7	
	25-29	0.0	0.0		0.0	0.0		9	8.4	
	30-39	1.0	21.4		0.7	20.0		39	36.4	
	40-49	1.0	21.4		0.3	10.0		28	26.2	
	50-59	1.0	21.4		1.0	30.0		17	15.9	
	60+	1.0	21.4		1.0	30.0		9	8.4	
Race/Ethnicity	White	3.3	71.4		3.0	90.0		69	64.5	
	Black	1.0	21.4		0.0	0.0		22	20.6	
	Hispanic	0.3	7.1		0.3	10.0		13	12.1	
	Multi Race	0.0	0.0		0.0	0.0		3	2.8	
Risk	MSM	1.7	35.7		1.0	30.0		36	33.6	
	IDU	0.0	0.0		0.3	10.0		16	15.0	
	MSM/IDU	0.0	0.0		0.0	0.0		8	7.5	
	Heterosexual	1.0	21.4		1.0	30.0		21	19.6	
	Fem Pres Het Cont	1.0	21.4		0.3	10.0		3	2.8	
	Pediatric Risk	0.0	0.0		0.0	0.0		1	0.9	
	Unk	1.0	21.4		0.7	20.0		22	20.6	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

Newly Diagnosed HIV and AIDS Cases¹ (2009-2011) and Cumulative AIDS² (through 2011)

]	HIV Dia	gnoses (Annual A	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fei	nale	To	otal	М	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		3.3	100.0	1.3	100.0	4.7	100.0	2.3	100.0	1.0	100.0	3.3	100.0	85	100.0	22	100.0	107	100.0
Risk	MSM	1.7	50.0	0.0	0.0	1.7	35.7	1.0	42.9	0.0	0.0	1.0	30.0	36	42.4	0	0.0	36	33.6
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	33.3	0.3	10.0	10	11.8	6	27.3	16	15.0
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	8	9.4	0	0.0	8	7.5
	Heterosexual	0.7	20.0	0.3	25.0	1.0	21.4	0.7	28.6	0.3	33.3	1.0	30.0	9	10.6	12	54.5	21	19.6
	Fem Pres Het Cont	0.0	0.0	1.0	75.0	1.0	21.4	0.0	0.0	0.3	33.3	0.3	10.0	0	0.0	3	13.6	3	2.8
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	1	4.5	1	0.9
	Unk	1.0	30.0	0.0	0.0	1.0	21.4	0.7	28.6	0.0	0.0	0.7	20.0	22	25.9	0	0.0	22	20.6

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		4	3
1986		3	1
1987		1	1
1988		2	2
1989		1	3
1990		5	2
1991		4	4
1992		8	2
1993		7	8
1994		5	5
1995		5	5
1996		6	1
1997		6	
1998		7	2
1999	•	2	3
2000*	5	4	
2001	7	6	2
2002	9	3	1
2003	3	2	1
2004	4	3	1
2005	6	2	2
2006	9	7	1
2007	3	2	1
2008	4	2	1
2009	5	4	1
2010	5	2	3
2011	4	4	

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		1,172	100.0	122.6	2,050	100.0	214.5	3,222	100.0	337.1	955,899
Gender	Male	731	62.4		1,323	64.5		2,054	63.7		
	Female	441	37.6		727	35.5		1,168	36.3		
Current Age	12 & under	9	0.8		0	0.0		9	0.3		
	13-19	27	2.3		7	0.3		34	1.1		
	20-24	60	5.1		27	1.3		87	2.7		
	25-29	84	7.2		56	2.7		140	4.3		
	30-39	221	18.9		202	9.9		423	13.1		
	40-49	340	29.0		619	30.2		959	29.8		
	50-59	293	25.0		764	37.3		1,057	32.8		
	60+	137	11.7		373	18.2		510	15.8		
	Unk	1	0.1		2	0.1		3	0.1		
Race/Ethnicity	White	282	24.1		402	19.6		684	21.2		
	Black	501	42.7		891	43.5		1,392	43.2		
	Hispanic	312	26.6		597	29.1		909	28.2		
	Multi Race	68	5.8		150	7.3		218	6.8		
	Others	9	0.8		10	0.5		19	0.6		
Risk	MSM	404	34.5		536	26.1		940	29.2		
	IDU	110	9.4		554	27.0		664	20.6		
	MSM/IDU	20	1.7		97	4.7		117	3.6		
	Heterosexual	375	32.0		640	31.2		1,015	31.5		
	Fem Pres Het Cont	110	9.4		75	3.7		185	5.7		
	Blood Prod	1	0.1		7	0.3		8	0.2		
	Pediatric Risk	36	3.1		28	1.4		64	2.0		
	Unk	116	9.9		113	5.5		229	7.1		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	To	otal	N	Iale	Fe	male	Т	otal	M	[ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		731	100.0	441	100.0	1,172	100.0	1,323	100.0	727	100.0	2,050	100.0	2,054	100.0	1,168	100.0	3,222	100.0
Risk	MSM	404	55.3	0	0.0	404	34.5	536	40.5	0	0.0	536	26.1	940	45.8	0	0.0	940	29.2
	IDU	65	8.9	45	10.2	110	9.4	355	26.8	199	27.4	554	27.0	420	20.4	244	20.9	664	20.6
	MSM/IDU	20	2.7	0	0.0	20	1.7	97	7.3	0	0.0	97	4.7	117	5.7	0	0.0	117	3.6
	Heterosexual	104	14.2	271	61.5	375	32.0	206	15.6	434	59.7	640	31.2	310	15.1	705	60.4	1,015	31.5
	Fem Pres Het Cont	0	0.0	110	24.9	110	9.4	0	0.0	75	10.3	75	3.7	0	0.0	185	15.8	185	5.7
	Blood Prod	1	0.1	0	0.0	1	0.1	4	0.3	3	0.4	7	0.3	5	0.2	3	0.3	8	0.2
	Pediatric Risk	21	2.9	15	3.4	36	3.1	12	0.9	16	2.2	28	1.4	33	1.6	31	2.7	64	2.0
	Unk	116	15.9	0	0.0	116	9.9	113	8.5	0	0.0	113	5.5	229	11.1	0	0.0	229	7.1

		HIV Diagn	oses (Annua	al Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		107.7	100.0	11.3	78.7	100.0	8.2	5,195	100.0	955,899
Gender	Male	76.7	71.2		56.3	71.6		3,671	70.7	
	Female	31.0	28.8		22.3	28.4		1,524	29.3	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		58	1.1	
	13-19	3.0	2.8		1.3	1.7		46	0.9	
	20-24	13.3	12.4		5.0	6.4		140	2.7	
	25-29	14.3	13.3		5.7	7.2		418	8.0	
	30-39	22.7	21.1		19.7	25.0		1,960	37.7	
	40-49	28.0	26.0		20.7	26.3		1,696	32.6	
	50-59	18.0	16.7		18.0	22.9		640	12.3	
	60+	8.3	7.7		8.3	10.6		237	4.6	
	Unk	0.0	0.0		0.0	0.0		0	0.0	
Race/Ethnicity	White	15.7	14.6		11.3	14.4		1,449	27.9	
	Black	46.7	43.3		35.0	44.5		2,391	46.0	
	Hispanic	36.0	33.4		24.7	31.4		1,115	21.5	
	Multi Race	8.0	7.4		6.7	8.5		221	4.3	
	Others	1.3	1.2		1.0	1.3		19	0.4	
Risk	MSM	46.7	43.3		28.7	36.4		1,301	25.0	
	IDU	3.7	3.4		8.7	11.0		2,112	40.7	
	MSM/IDU	1.3	1.2		2.0	2.5		253	4.9	
	Heterosexual	37.7	35.0		27.7	35.2		1,065	20.5	
	Fem Pres Het Cont	8.0	7.4		3.3	4.2		124	2.4	
	Blood Prod	0.0	0.0		0.0	0.0		46	0.9	
	Pediatric Risk	0.0	0.0		0.7	0.8		77	1.5	
	Unk	10.3	9.6		7.7	9.7		217	4.2	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

Newly Diagnosed HIV and AIDS Cases¹ (2009-2011) and Cumulative AIDS² (through 2011)

		I	HIV Dia	gnoses (Annual	Average	e)	A	AIDS Dia	gnoses	(Annual	Average	e)		Cum	ulativ	e AIDS (Cases	
		M	ale	Fer	nale	To	otal	М	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		76.7	100.0	31.0	100.0	107.7	100.0	56.3	100.0	22.3	100.0	78.7	100.0	3,671	100.0	1,524	100.0	5,195	100.0
Risk	MSM	46.7	60.9	0.0	0.0	46.7	43.3	28.7	50.9	0.0	0.0	28.7	36.4	1,301	35.4	0	0.0	1,301	25.0
	IDU	3.0	3.9	0.7	2.2	3.7	3.4	4.7	8.3	4.0	17.9	8.7	11.0	1,509	41.1	603	39.6	2,112	40.7
	MSM/IDU	1.3	1.7	0.0	0.0	1.3	1.2	2.0	3.6	0.0	0.0	2.0	2.5	253	6.9	0	0.0	253	4.9
	Heterosexual	15.3	20.0	22.3	72.0	37.7	35.0	12.7	22.5	15.0	67.2	27.7	35.2	322	8.8	743	48.8	1,065	20.5
	Fem Pres Het Cont	0.0	0.0	8.0	25.8	8.0	7.4	0.0	0.0	3.3	14.9	3.3	4.2	0	0.0	124	8.1	124	2.4
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	33	0.9	13	0.9	46	0.9
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.7	1.2	0.0	0.0	0.7	0.8	36	1.0	41	2.7	77	1.5
	Unk	10.3	13.5	0.0	0.0	10.3	9.6	7.7	13.6	0.0	0.0	7.7	9.7	217	5.9	0	0.0	217	4.2

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
<1986		110	57
1986		110	75
1987		172	106
1988		190	129
1989		232	156
1990		246	184
1991		293	186
1992		393	224
1993		345	224
1994		347	263
1995		339	260
1996		276	167
1997		225	100
1998		215	101
1999		171	87
2000*	209	179	112
2001	139	173	65
2002	133	167	74
2003	129	171	89
2004	148	139	68
2005	109	129	67
2006	144	115	60
2007	129	122	64
2008	128	100	71
2009	109	89	60
2010	115	85	61
2011	99	62	40

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	l AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		6	100.0	14.3	5	100.0	11.9	11	100.0	26.2	41,944
Gender	Male	5	83.3		3	60.0		8	72.7		
	Female	1	16.7		2	40.0		3	27.3		
Current Age	12 & under	0	0.0		1	20.0		1	9.1		
	13-19	1	16.7		0	0.0		1	9.1		
	25-29	1	16.7		0	0.0		1	9.1		
	30-39	1	16.7		0	0.0		1	9.1		
	40-49	2	33.3		3	60.0		5	45.5		
	50-59	0	0.0		0	0.0		0	0.0		
	60+	1	16.7		1	20.0		2	18.2		
Race/Ethnicity	White	5	83.3		2	40.0		7	63.6		
	Black	0	0.0		1	20.0		1	9.1		
	Multi Race	1	16.7		1	20.0		2	18.2		
	Others	0	0.0		1	20.0		1	9.1		
Risk	MSM	4	66.7		1	20.0		5	45.5		
	IDU	0	0.0		0	0.0		0	0.0		
	MSM/IDU	0	0.0		0	0.0		0	0.0		
	Heterosexual	0	0.0		2	40.0		2	18.2		
	Fem Pres Het Cont	1	16.7		0	0.0		1	9.1		
	Blood Prod	0	0.0		1	20.0		1	9.1		
	Pediatric Risk	1	16.7		1	20.0		2	18.2		

¹ Data as of March 2013

²Per 100,000 population

			Living	HIV (n	ot AIDS) Cases			Li	ving A	AIDS Cas	ses			Living	HIV a	nd AIDS	Cases	
		М	ale	Fe	male	Т	otal	N	Male	Fe	emale	Т	otal	M	[ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		5	100.0	1	100.0	6	100.0	3	100.0	2	100.0	5	100.0	8	100.0	3	100.0	11	100.0
Risk	MSM	4	80.0	0	0.0	4	66.7	1	33.3	0	0.0	1	20.0	5	62.5	0	0.0	5	45.5
	IDU	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	MSM/IDU	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Heterosexual	0	0.0	0	0.0	0	0.0	0	0.0	2	100.0	2	40.0	0	0.0	2	66.7	2	18.2
	Fem Pres Het Cont	0	0.0	1	100.0	1	16.7	0	0.0	0	0.0	0	0.0	0	0.0	1	33.3	1	9.1
	Blood Prod	0	0.0	0	0.0	0	0.0	1	33.3	0	0.0	1	20.0	1	12.5	0	0.0	1	9.1
	Pediatric Risk	1	20.0	0	0.0	1	16.7	1	33.3	0	0.0	1	20.0	2	25.0	0	0.0	2	18.2

		HIV Diagn	oses (Annua	ıl Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		0.3	100.0	0.8	0.0	0.0	0.0	15	100.0	41,944
Gender	Male	0.3	100.0		0.0	0.0		12	80.0	
	Female	0.0	0.0		0.0	0.0		3	20.0	
Age at Diagnosis	12 & under	0.0	0.0		0.0	0.0		1	6.7	
	13-19	0.0	0.0		0.0	0.0		0	0.0	
	25-29	0.0	0.0		0.0	0.0		2	13.3	
	30-39	0.3	100.0		0.0	0.0		5	33.3	
	40-49	0.0	0.0		0.0	0.0		4	26.7	
	50-59	0.0	0.0		0.0	0.0		2	13.3	
	60+	0.0	0.0		0.0	0.0		1	6.7	
Race/Ethnicity	White	0.3	100.0		0.0	0.0		12	80.0	
	Black	0.0	0.0		0.0	0.0		1	6.7	
	Multi Race	0.0	0.0		0.0	0.0		1	6.7	
	Others	0.0	0.0		0.0	0.0		1	6.7	
Risk	MSM	0.3	100.0		0.0	0.0		8	53.3	
	IDU	0.0	0.0		0.0	0.0		1	6.7	
	MSM/IDU	0.0	0.0		0.0	0.0		1	6.7	
	Heterosexual	0.0	0.0		0.0	0.0		3	20.0	
	Fem Pres Het Cont	0.0	0.0		0.0	0.0		0	0.0	
	Blood Prod	0.0	0.0		0.0	0.0		1	6.7	
	Pediatric Risk	0.0	0.0		0.0	0.0		1	6.7	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

		1	HIV Diagnoses (Annual Average)							AIDS Diagnoses (Annual Average)							Cumulative AIDS Cases						
		M	ale	Fei	Female		otal	М	ale	Fer	nale	То	tal	N	Iale	Fe	male	Т	otal				
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %				
Total		0.3	100.0	0.0	0.0	0.3	100.0	0.0	0.0	0.0	0.0	0.0	0.0	12	100.0	3	100.0	15	100.0				
Risk	MSM	0.3	100.0	0.0	0.0	0.3	100.0	0.0	0.0	0.0	0.0	0.0	0.0	8	66.7	0	0.0	8	53.3				
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	8.3	0	0.0	1	6.7				
	MSM/IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	8.3	0	0.0	1	6.7				
	Heterosexual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	3	100.0	3	20.0				
	Fem Pres Het Cont	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	0	0.0	0	0.0				
	Blood Prod	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	8.3	0	0.0	1	6.7				
	Pediatric Risk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	8.3	0	0.0	1	6.7				

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
1988		1	1
1990		1	
1991			1
1993		4	2
1994		2	2
1995		1	
1996		1	
1998		2	1
1999		2	
2000*	2	1	
2003	1		1
2005	1		
2007			1
2011	1		1

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.

		Livin	g HIV (not	AIDS) Cases]	Living A	IDS Cases	Livi	ng HIV and	1 AIDS Cases	
		Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Number	Col %	Prevalence Rate ²	Population
Total		6	100.0	23.6	6	100.0	23.6	12	100.0	47.1	25,454
Gender	Male	4	66.7		2	33.3		6	50.0		
	Female	2	33.3		4	66.7		6	50.0		
Current Age	20-24	0	0.0		0	0.0		0	0.0		
	25-29	2	33.3		1	16.7		3	25.0		
	30-39	0	0.0		0	0.0		0	0.0		
	40-49	3	50.0		1	16.7		4	33.3		
	50-59	1	16.7		2	33.3		3	25.0		
	60+	0	0.0		2	33.3		2	16.7		
Race/Ethnicity	White	2	33.3		6	100.0		8	66.7		
	Black	4	66.7		0	0.0		4	33.3		
	Hispanic	0	0.0		0	0.0		0	0.0		
	Others	0	0.0		0	0.0		0	0.0		
Risk	MSM	2	33.3		0	0.0		2	16.7		
	IDU	0	0.0		4	66.7		4	33.3		
	Heterosexual	1	16.7		0	0.0		1	8.3		
	Fem Pres Het Cont	1	16.7		2	33.3		3	25.0		
	Unk	2	33.3		0	0.0		2	16.7		

¹ Data as of March 2013

²Per 100,000 population

	Living HIV (not AIDS								Living AIDS Cases Living HIV and AIDS Ca							Cases	Cases		
			Male		Female		otal	N	Male	Fe	male	1	otal	M	[ale	Fe	male	T	otal
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %
Total		4	100.0	2	100.0	6	100.0	2	100.0	4	100.0	6	100.0	6	100.0	6	100.0	12	100.0
Risk	MSM	2	50.0	0	0.0	2	33.3	0	0.0	0	0.0	0	0.0	2	33.3	0	0.0	2	16.7
	IDU	0	0.0	0	0.0	0	0.0	2	100.0	2	50.0	4	66.7	2	33.3	2	33.3	4	33.3
	Heterosexual	0	0.0	1	50.0	1	16.7	0	0.0	0	0.0	0	0.0	0	0.0	1	16.7	1	8.3
	Fem Pres Het Cont	0	0.0	1	50.0	1	16.7	0	0.0	2	50.0	2	33.3	0	0.0	3	50.0	3	25.0
	Unk	2	50.0	0	0.0	2	33.3	0	0.0	0	0.0	0	0.0	2	33.3	0	0.0	2	16.7

		HIV Diagn	oses (Annua	l Average)	AIDS Diag	noses (Annua	al Average)	Cumulative	AIDS Cases	
		Number	Col %	Case Rate ³	Number	Col %	Case Rate ³	Number	Col %	Population
Total		0.0	0.0	0.0	0.0	0.0	0.0	13	100.0	25,454
Gender	Male	0.0	0.0		0.0	0.0		8	61.5	
	Female	0.0	0.0		0.0	0.0		5	38.5	
Age at Diagnosis	20-24	0.0	0.0		0.0	0.0		1	7.7	
	25-29	0.0	0.0		0.0	0.0		0	0.0	
	30-39	0.0	0.0		0.0	0.0		5	38.5	
	40-49	0.0	0.0		0.0	0.0		6	46.2	
	50-59	0.0	0.0		0.0	0.0		1	7.7	
	60+	0.0	0.0		0.0	0.0		0	0.0	
Race/Ethnicity	White	0.0	0.0		0.0	0.0		11	84.6	
	Black	0.0	0.0		0.0	0.0		0	0.0	
	Hispanic	0.0	0.0		0.0	0.0		1	7.7	
	Others	0.0	0.0		0.0	0.0		1	7.7	
Risk	MSM	0.0	0.0		0.0	0.0		5	38.5	
	IDU	0.0	0.0		0.0	0.0		4	30.8	
	Heterosexual	0.0	0.0		0.0	0.0		2	15.4	
	Fem Pres Het Cont	0.0	0.0		0.0	0.0		2	15.4	
	Unk	0.0	0.0		0.0	0.0		0	0.0	

¹ Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

³Per 100,000 population

Newly Diagnosed HIV and AIDS Cases¹ (2009-2011) and Cumulative AIDS² (through 2011)

]	HIV Diagnoses (Annual Average)							AIDS Diagnoses (Annual Average)							Cumulative AIDS Cases						
			Male		Female		otal	М	ale	Fer	nale	To	otal	N	Iale	Fe	male	Т	otal				
		#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %	#	Col %				
Total		0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	8	100.0	5	100.0	13	100.0				
Risk	MSM	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	5	62.5	0	0.0	5	38.5				
	IDU	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2	25.0	2	40.0	4	30.8				
	Heterosexual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	12.5	1	20.0	2	15.4				
	Fem Pres Het Cont	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	2	40.0	2	15.4				
	Unk	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	0	0.0	0	0.0				

¹Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way

²Data as of March 2013

Newly Diagnosed HIV and AIDS Cases¹ by Year of Diagnosis² and Deaths Among Persons Living With AIDS (PLWA) by Year Excludes State Prison Inmates

Year	HIV Diagnoses	AIDS Diagnoses	Deaths Among PLWA ³
1989		1	
1991		2	2
1992		1	
1993		2	2
1994		1	2
1995		2	
1996			1
2000*	1		
2002	1		
2003	2	1	
2004	1	1	1
2006	2	1	
2008		1	

Persons diagnosed with HIV may also be diagnosed with AIDS in the same year or a later year and their AIDS diagnosis will be counted in the AIDS diagnosis tables. HIV and AIDS diagnoses cannot be added together in a meaningful way.

²Data as of March 2013

 $^{^{3}}$ Deaths among AIDS cases. However, death may be due to causes other than AIDS.

^{*}HIV reporting began in June 2000. No earlier data are available.