

HIV-Positive Adults in Care in New York State (Excluding NYC), Medical Monitoring Project, 2009-2014

The Medical Monitoring Project (MMP)

- MMP is a surveillance system funded by the Centers for Disease Control and Prevention and implemented by local health departments. It collects behavioral and medical data about HIV-positive adults receiving medical care in the United States.
- From 2009 to 2014, MMP interviewed 618 HIV-positive adults receiving care in New York State (excluding NYC). Their responses reflect their experiences during the 12 months before their interview, unless otherwise noted. All data presented are weighted.
- The information in this factsheet can guide policy decisions, resource allocation, and evaluation of treatment and prevention initiatives.

Characteristics of HIV-Positive Adults in Care in New York State, 2009-2014


- 66% were male and 33% were female
- 40% were men who have sex with men
- 31% were black/African-American, 15% were Hispanic or Latino, and 45% were white
- 16% had been diagnosed with HIV less than 5 years at the time of their interview
- 38% had private insurance, 54% public insurance only, 5% had Ryan White coverage only, and 1% were uninsured
- 46% had a household income at or below the poverty line
- 6% experienced homelessness

HIV Treatment and Prevention Measures among HIV-Positive Adults in Care in New York State, Medical Monitoring Project (MMP), 2009-2014*


Characteristic	Prescription of ART ^{1,2} (%)	ART Dose Adherence ^{3,4} (%)	Sustained Viral Suppression ^{2,5} (%)	Receipt of Condoms ^{3,6} (%)	HIV Prevention Counseling ^{3,7} (%)
Total	96	89	71	52	47
Age					
18-29 years	89	84	50	75	76
30-39 years	94	90	65	62	49
40-49 years	93	86	68	53	48
≥50 years	98	92	77	46	43
Gender					
Male	96	93	74	52	47
Female	95	83	65	50	49
Race/Ethnicity					
Black/African-American	97	85	59	64	62
Hispanic/Latino	93	88	68	56	48
White	96	93	79	40	40
Insurance					
Any Private Insurance	96	92	78	41	45
Public Insurance Only	97	88	68	58	48
Ryan White Coverage Only	94	83	66	59	63
Uninsured	33	§	13	77	45
Sexual Behavior					
MSM ⁸	94	94	74	46	42
MSW ⁹	98	91	72	60	50
WSM ¹⁰	95	81	62	50	50

* All data are from the past 12 months unless specified otherwise; §Estimates suppressed because coefficient of variation for the estimate was ≥30%.¹Antiretroviral therapy; ²Documented in the medical record; ³Self-reported; ⁴Past three days; ⁵All viral loads in past 12 months undetectable or <200 copies/ml; ⁶Received free condoms, not counting those given by a friend, relative, or sex partner; ⁷Given by a healthcare provider; ⁸Men who have sex with men; ⁹Men who have sex with women only; ¹⁰Women who have sex with men

Met and Unmet Needs for Services among HIV-Positive Adults in Care in New York State, Medical Monitoring Project, 2009 – 2014



Behavioral and Clinical Characteristics



Notes: any sex = any oral, anal, or vaginal sex from 2009 to 2013, but in 2014 oral sex was excluded; condomless sex = vaginal or anal sex without a condom; condomless sex with HIV-negative or unknown-status partner=engaged in condomless sex with an HIV-negative partner or a partner whose status was unknown; all sexual behavior information is self-reported.

Documentation of MMP methods can be found here: http://www.cdc.gov/hiv/pdf/HSSR_MMP_2010-PDF01.pdf

Questions? Contact us:

New York State Department of Health
 Bureau of HIV/AIDS Epidemiology
 Corning Tower, Empire State Plaza Station
 Albany, NY 12237-0001
 518-474-4284

Project Coordinator

Megan Swanson
 Megan.Swanson@health.ny.gov

<http://www.cdc.gov/hiv/statistics/systems/mmp/>
<https://www.health.ny.gov/diseases/aids/general/about/>

Substance Use¹

- 43% were current smokers
- 17% engaged in binge drinking²
- 2% used injection drugs³

¹Self-reported

²Binge drinking is defined for men as 5 or more drinks in a sitting in the past 30 days and for women as 4 or more drinks in a sitting in the past 30 days.

³Past 12 months

STD Testing Among Sexually Active Persons in the Past 12 Months

- 53% were tested for syphilis
- 36% were tested for gonorrhea
- 36% were tested for chlamydia

Influenza Vaccination in the Past 12 Months

- 87% received an influenza vaccine



Department of Health

Medical Monitoring Project

