

**Expanded BRFSS, 2003**

**Chemung, Schuyler, Tioga Locality Tables**

TABLE OF CONTENTS:

Core 1: Health Status / Healthy Days.....	1
Core 2: Health Care Access.....	4
Core 3: Exercise.....	8
Core 4: Diabetes.....	9
Core 5: Asthma.....	10
Core 6: Arthritis.....	12
Core 7: Tobacco Use.....	14
Core 8: Tobacco ETS.....	17
Core 9: Alcohol Consumption.....	19
Core 10: Weight Status.....	21
Core 11: Mammography.....	23
Core 12: Sexual Behavior.....	25
Core 13: Family Planning.....	28
Core 14: Cardiovascular Disease.....	29
Core 15: Prostate Cancer Screening.....	31
Core 16: Colorectal Cancer Screening.....	37

**New York State Department Of Health**

Expanded BRFSS, 2003 - New York State Department of Health

## Self-rated Health Status

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	Fair or Poor % <sup>2</sup>	n	Good to Excellent %	C.I. <sup>3</sup>
Total	111	15.5	521	84.5	3.0
Gender					
Male	36	12.4	219	87.6	4.1
Female	75	18.4	302	81.6	4.4
Age					
18-34	9	5.7	130	94.3	4.1
35-54	39	14.6	215	85.4	4.6
≥55	62	24.7	169	75.3	6.2
Education					
≤High School	74	21.5	236	78.5	5.0
>High School	36	9.5	284	90.5	3.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Reported Number of Physical Distress Days in the Past Month

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	≥14 days % <sup>2</sup>	n	< 14 days %	C.I. <sup>3</sup>
Total	77	11.1	542	88.9	2.6
Gender					
Male	27	9.5	224	90.5	3.6
Female	50	12.6	318	87.4	3.7
Age					
18-34	13	9.0	126	91.0	5.1
35-54	23	8.1	228	91.9	3.5
≥55	41	17.1	180	82.9	5.3
Education					
≤High School	46	13.8	252	86.2	4.2
>High School	30	8.2	289	91.8	3.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Reported Number of Mental Distress Days in the Past Month

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	≥14 days % <sup>2</sup>	n	< 14 days %	C.I. <sup>3</sup>
Total	73	10.3	549	89.7	2.5
Gender					
Male	21	8.3	230	91.7	3.6
Female	52	12.3	319	87.7	3.4
Age					
18-34	23	13.7	116	86.3	5.7
35-54	34	11.6	217	88.4	4.0
≥55	14	5.0	210	95.0	2.8
Education					
≤High School	38	11.3	264	88.7	3.7
>High School	35	9.5	283	90.5	3.3

<sup>1</sup> Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup> Weighted Percent

<sup>3</sup> 95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Health Insurance Coverage, 18-64 Years of Age

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	394	82.1	84	17.9	3.9
Gender					
Male	168	80.1	40	19.9	6.2
Female	226	84.2	44	15.8	4.7
Age					
18-34	106	75.7	34	24.3	7.8
35-54	210	82.7	44	17.3	5.3
55-64	78	94.1	6	5.9	4.7
Education					
≤High School	167	77.9	46	22.1	6.3
>High School	226	85.6	38	14.4	4.8

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Health Insurance Coverage, Aged 18 and Older

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	545	85.4	88	14.6	3.2
Gender					
Male	213	82.9	42	17.1	5.3
Female	332	87.8	46	12.2	3.6
Age					
18-34	106	75.7	34	24.3	7.8
35-54	210	82.7	44	17.3	5.3
≥55	222	96.8	9	3.2	2.2
Education					
≤High School	263	83.2	48	16.8	4.9
>High School	280	87.5	40	12.5	4.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Reported Having a Routine Physical Exam in Past Two Years

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	537	85.2	89	14.8	3.1
Gender					
Male	192	78.1	59	21.9	5.5
Female	345	91.8	30	8.2	3.0
Age					
18-34	111	84.4	22	15.6	6.6
35-54	208	79.9	47	20.1	5.6
≥55	213	93.0	17	7.0	3.3
Education					
≤High School	259	83.0	48	17.0	4.9
>High School	276	87.1	41	12.9	3.9

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Time in Past 12 Months when Needed Medical Care but Could Not Get It

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	38	5.5	597	94.5	1.9
Gender					
Male	12	3.5	245	96.5	2.1
Female	26	7.4	352	92.6	3.0
Age					
18-34	10	5.4	131	94.6	3.6
35-54	20	7.7	235	92.3	3.5
≥55	8	2.9	223	97.1	2.1
Education					
≤High School	23	7.7	289	92.3	3.2
>High School	15	3.4	306	96.6	1.9

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval



Expanded BRFSS, 2003 - New York State Department of Health

## Participated in Leisure-Time Physical Activity or Exercise During the Past 30 Days

*Chemung, Schuyler, Tioga*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	470	75.3	164	24.7	3.7
Gender					
Male	197	77.5	60	22.5	5.6
Female	273	73.1	104	26.9	5.0
Age					
18-34	111	80.9	30	19.1	6.7
35-54	203	78.6	52	21.4	5.6
≥55	148	65.3	82	34.7	7.1
Education					
≤High School	201	66.4	110	33.6	5.8
>High School	267	83.5	54	16.5	4.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Physician Diagnosed Diabetes

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	54	8.3	581	91.7	2.4
Gender					
Male	18	6.4	239	93.6	3.1
Female	36	10.0	342	90.0	3.6
Age					
18-34	2	0.6	139	99.4	0.9
35-54	15	6.4	240	93.6	3.5
≥55	37	17.4	194	82.6	5.8
Education					
≤High School	36	11.0	276	89.0	3.9
>High School	18	5.7	303	94.3	2.8

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Lifetime Asthma Diagnosed by a Medical Professional

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	79	12.9	556	87.1	3.0
Gender					
Male	23	9.4	234	90.6	4.0
Female	56	16.3	322	83.7	4.3
Age					
18-34	22	15.7	119	84.3	6.6
35-54	36	14.0	219	86.0	4.6
≥55	20	8.9	211	91.1	4.4
Education					
≤High School	37	12.9	275	87.1	4.4
>High School	42	13.0	279	87.0	4.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Current Asthma Diagnosed by a Medical Professional

*Chemung, Schuylers, Tioga*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	58	9.2	577	90.8	2.5
Gender					
Male	16	6.5	241	93.5	3.2
Female	42	11.8	336	88.2	3.8
Age					
18-34	15	9.1	126	90.9	4.8
35-54	25	10.4	230	89.6	4.2
≥55	18	8.1	213	91.9	4.3
Education					
≤High School	30	9.6	282	90.4	3.7
>High School	28	8.8	293	91.2	3.4

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Arthritis Diagnosed by a Medical Professional

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	203	28.1	431	71.9	3.8
Gender					
Male	65	23.9	191	76.1	5.6
Female	138	32.0	240	68.0	5.0
Age					
18-34	15	7.7	125	92.3	4.2
35-54	63	24.9	192	75.1	5.9
≥55	124	50.1	107	49.9	7.3
Education					
≤High School	113	31.3	199	68.7	5.6
>High School	89	24.9	231	75.1	5.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Adults with Doctor Diagnosed Arthritis or Possible Arthritis (Chronic Joint Symptoms Only)

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	Doctor Diagnosed % <sup>2</sup>	C.I. <sup>3</sup>	n	Possible Arthritis (CJS Only) %	C.I.	n	No Arthritis or CJS %	C.I.
Total	203	28.4	3.8	114	18.6	3.5	313	53.1	4.3
Gender									
Male	65	24.5	5.7	50	21.3	5.7	137	54.3	6.8
Female	138	32.0	5.0	64	16.0	4.0	176	52.0	5.5
Age									
18-34	15	7.9	4.3	28	18.5	6.8	95	73.6	7.6
35-54	63	25.2	5.9	54	22.6	5.9	136	52.2	6.8
≥55	124	50.1	7.3	28	12.3	5.1	79	37.7	7.2
Education									
≤High School	113	32.0	5.6	56	20.2	5.2	139	47.7	6.3
>High School	89	24.9	5.1	57	16.7	4.5	174	58.4	5.9

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Ever Smoked 100 Cigarettes in Lifetime

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	340	54.1	293	45.9	4.3
Gender					
Male	152	58.3	104	41.7	6.7
Female	188	50.2	189	49.8	5.6
Age					
18-34	73	51.3	68	48.7	9.0
35-54	133	51.0	122	49.0	6.8
≥55	129	60.4	100	39.6	7.1
Education					
≤High School	188	62.5	122	37.5	6.0
>High School	150	45.9	171	54.1	6.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Current Smoker

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	179	28.8	453	71.2	4.0
Gender					
Male	72	29.1	183	70.9	6.2
Female	107	28.6	270	71.4	5.0
Age					
18-34	55	38.5	86	61.5	8.8
35-54	83	30.7	172	69.3	6.2
≥55	37	17.2	191	82.8	5.8
Education					
≤High School	109	36.5	200	63.5	6.1
>High School	68	21.2	253	78.8	5.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval



Expanded BRFSS, 2003 - New York State Department of Health

## Status of Cigarette Smoking

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	Everyday Smoker % <sup>2</sup>	C.I. <sup>3</sup>	n	Someday Smoker %	C.I.	n	Former or Never Smoked %	C.I.
Total	147	23.8	3.8	32	5.0	1.9	453	71.2	4.0
Gender									
Male	62	25.7	6.1	10	3.5	2.3	183	70.9	6.2
Female	85	22.1	4.5	22	6.5	3.0	270	71.4	5.0
Age									
18-34	46	31.9	8.5	9	6.6	4.4	86	61.5	8.8
35-54	71	26.4	5.9	12	4.3	2.7	172	69.3	6.2
≥55	26	12.4	5.1	11	4.8	3.3	191	82.8	5.8
Education									
≤High School	96	32.6	5.9	13	3.9	2.4	200	63.5	6.1
>High School	50	15.4	4.5	18	5.9	2.9	253	78.8	5.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Home Smoking Policies

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	Not Allowed % <sup>2</sup>	C.I. <sup>3</sup>	n	Allowed Some Places %	C.I.	n	Allowed Anywhere %	C.I.
Total	374	59.7	4.3	98	15.3	3.1	161	25.0	3.8
Gender									
Male	152	60.2	6.6	30	12.3	4.5	73	27.6	6.1
Female	222	59.2	5.5	68	18.2	4.2	88	22.6	4.7
Age									
18-34	80	59.6	8.8	29	19.7	7.2	31	20.7	7.1
35-54	148	59.2	6.6	45	16.8	4.8	62	23.9	5.9
≥55	141	60.3	7.3	23	9.6	4.1	66	30.1	7.0
Education									
≤High School	157	49.7	6.2	58	18.3	4.7	96	32.0	6.0
>High School	217	69.6	5.5	39	12.2	4.0	64	18.2	4.5

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Home Smoking

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	192	30.1	442	69.9	4.0
Gender					
Male	77	30.0	179	70.0	6.3
Female	115	30.3	263	69.7	5.1
Age					
18-34	45	30.7	96	69.3	8.3
35-54	86	31.1	169	68.9	6.2
≥55	59	28.6	171	71.4	7.0
Education					
≤High School	115	37.8	196	62.2	6.1
>High School	77	23.0	244	77.0	5.1

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## At Risk for Binge Drinking (Drank in Past Month and had 5 or more Drinks per drink Occasion)

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	88	16.3	535	83.7	3.4
Gender					
Male	59	25.2	191	74.8	6.0
Female	29	8.0	344	92.0	3.0
Age					
18-34	38	29.8	100	70.2	8.6
35-54	35	14.5	215	85.5	4.8
≥55	14	6.9	213	93.1	3.7
Education					
≤High School	39	15.4	267	84.6	4.9
>High School	49	17.2	267	82.8	4.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**At Risk for Heavy Drinking (Males More Than 2 Drinks and Females More Than 1 Drink per Day in Past Month)**

*Chemung, Schuyler, Tioga*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	36	5.9	590	94.1	2.1
Gender					
Male	15	6.1	238	93.9	3.4
Female	21	5.6	352	94.4	2.5
Age					
18-34	11	8.6	129	91.4	5.4
35-54	15	5.1	236	94.9	2.7
≥55	9	4.1	218	95.9	2.7
Education					
≤High School	17	6.3	291	93.7	3.3
>High School	19	5.4	298	94.6	2.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Overweight or Obese, BMI(Body Mass Index) 25 or More**

*Chemung, Schuyler, Tioga*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	375	63.3	218	36.7	4.4
Gender					
Male	177	67.8	77	32.2	6.4
Female	198	58.4	141	41.6	5.8
Age					
18-34	70	51.4	60	48.6	9.4
35-54	150	64.8	88	35.2	6.7
≥55	152	71.4	68	28.6	6.6
Education					
≤High School	190	66.6	98	33.4	6.1
>High School	185	60.4	119	39.6	6.2

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Weight Status (Not Overweight BMI<25, Overweight BMI 25-29, Obese BMI 30+)**

*Chemung, Schuylers, Tioga*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Underweight\ Normal %<sup>2</sup></b>	<b>C.I.<sup>3</sup></b>	<b>n</b>	<b>Overweight %</b>	<b>C.I.</b>	<b>n</b>	<b>Obese %</b>	<b>C.I.</b>
Total	218	36.7	4.4	229	40.2	4.4	146	23.0	3.7
Gender									
Male	77	32.2	6.4	121	47.1	6.8	56	20.7	5.3
Female	141	41.6	5.8	108	32.8	5.5	90	25.6	5.1
Age									
18-34	60	48.6	9.4	42	33.0	8.9	28	18.4	6.7
35-54	88	35.2	6.7	92	41.2	7.0	58	23.6	5.9
≥55	68	28.6	6.6	92	44.6	7.5	60	26.7	6.7
Education									
≤High School	98	33.4	6.1	107	38.9	6.4	83	27.7	5.8
>High School	119	39.6	6.2	122	41.6	6.2	63	18.7	4.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Ever Had Mammography Screening, Women Aged 40 Years and Older**

*Chemung, Schuyler, Tioga*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Female	246	92.7	18	7.3	3.8
Age					
40-64	150	92.2	13	7.8	4.5
≥65	96	93.6	5	6.4	7.0
Education					
≤High School	128	91.9	9	8.1	6.0
>High School	117	93.4	9	6.6	4.5

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval



Expanded BRFSS, 2003 - New York State Department of Health

**Received Mammography Screening in Past Two Years, Women Aged 40 Years and Older**

*Chemung, Schuyler, Tioga*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Female	204	77.3	57	22.7	5.7
Age					
40-64	127	78.7	36	21.3	6.7
≥65	77	74.7	21	25.3	10.6
Education					
≤High School	102	74.1	32	25.9	8.6
>High School	101	80.7	25	19.3	7.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Ever Counseled by Medical Professional on Prevention of STD's Through Condom Use, Aged 64 or Younger**

*Chemung, Schuyler, Tioga*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	61	12.2	413	87.8	3.2
Gender					
Male	12	6.5	194	93.5	3.9
Female	49	18.2	219	81.8	5.1
Age					
18-34	37	24.3	103	75.7	7.5
35-54	24	7.8	227	92.2	3.5
55-64	0	0.0	83	100	0.0
Education					
≤High School	30	12.9	180	87.1	4.9
>High School	31	11.8	232	88.2	4.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Sexually Active, Age 18-64 Years Old**

*Chemung, Schuyler, Tioga*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	360	82.3	89	17.7	3.9
Gender					
Male	156	80.9	38	19.1	6.2
Female	204	83.7	51	16.3	4.7
Age					
18-34	117	86.0	18	14.0	6.5
35-54	195	83.8	45	16.2	5.1
55-64	48	68.2	26	31.8	12.2
Education					
≤High School	162	82.7	35	17.3	5.8
>High School	197	81.9	54	18.1	5.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Used Condom Last Time Had Sexual Intercourse, Aged 64 or Younger

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	61	16.6	307	83.4	4.3
Gender					
Male	27	19.1	132	80.9	7.1
Female	34	14.3	175	85.7	4.9
Age					
18-34	35	31.3	84	68.7	9.3
35-54	24	9.1	174	90.9	3.9
55-64	2	4.0	49	96.0	5.5
Education					
≤High School	27	13.6	140	86.4	5.7
>High School	34	19.3	166	80.7	6.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Birth Control(BC) Use To Prevent Pregnancy Among Reproductive Aged Men(18-59) and Women(18-44)

*Chemung, Schuyler, Tioga*

Demographic Groups	Use BC			No BC at risk			No BC		
	n <sup>1</sup>	% <sup>2</sup>	C.I. <sup>3</sup>	n	%	C.I.	n	%	C.I.
Total	216	65.9	5.7	44	13.6	4.0	65	20.6	5.0
Gender									
Male	105	57.9	7.8	29	15.0	5.5	50	27.2	7.2
Female	111	79.0	7.2	15	11.3	5.8	15	9.7	5.1
Age									
18-34	97	67.7	8.6	21	15.8	6.6	20	16.5	7.0
35-59	119	64.3	7.6	23	11.7	4.9	45	24.0	6.9
Education									
≤High School	104	68.4	8.3	17	11.3	5.7	28	20.3	7.2
>High School	111	63.5	7.9	27	15.6	5.8	37	20.9	6.9

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Physician Diagnosed Heart Attack, Angina or Stroke**

*Chemung, Schuylers, Tioga*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	57	7.6	571	92.4	2.1
Gender					
Male	25	8.9	227	91.1	3.4
Female	32	6.3	344	93.7	2.4
Age					
18-64	18	2.9	455	97.1	1.4
≥65	39	26.6	108	73.4	8.0
Education					
≤High School	35	9.4	274	90.6	3.2
>High School	22	5.9	295	94.1	2.6

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Physician Diagnosed Coronary Heart Disease (Heart Attack or Angina)**

*Chemung, Schuyler, Tioga*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	42	5.7	586	94.3	1.8
Gender					
Male	21	7.6	231	92.4	3.2
Female	21	3.9	355	96.1	1.8
Age					
18-64	12	2.0	461	98.0	1.2
≥65	30	20.7	117	79.3	7.3
Education					
≤High School	25	6.4	284	93.6	2.6
>High School	17	5.0	300	95.0	2.5

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Ever Had Prostate Specific Antigen Test, Men Aged 40 and Older**

*Chemung, Schuyler, Tioga*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Male	98	63.8	55	36.2	8.3
Age					
40-64	60	53.9	49	46.1	10.2
≥65	38	87.8	6	12.2	9.7
Education					
≤High School	43	53.6	34	46.4	12.2
>High School	54	74.3	21	25.7	10.2

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval



Expanded BRFSS, 2003 - New York State Department of Health

## Received Prostate Specific Antigen Test in the Past Two Years, Men Aged 40 and Older

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Male	84	55.7	69	44.3	8.5
Age					
40-64	50	45.5	59	54.5	10.2
≥65	34	80.5	10	19.5	11.5
Education					
≤High School	36	45.2	41	54.8	12.1
>High School	48	67.9	27	32.1	10.9

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Ever Had a Digital Rectal Exam, Men Aged 40 and Older

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Male	132	82.2	29	17.8	6.5
Age					
40-64	94	80.3	22	19.7	8.2
≥65	38	87.0	7	13.0	9.6
Education					
≤High School	57	70.1	24	29.9	11.1
>High School	74	94.4	5	5.6	4.8

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Ever Used a Blood Stool Test at Home, Aged 50 and Older

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	151	53.9	129	46.1	6.7
Gender					
Male	51	49.7	53	50.3	10.5
Female	100	57.7	76	42.3	8.3
Age					
50-64	63	48.7	73	51.3	9.4
≥65	88	59.4	56	40.6	9.4
Education					
≤High School	72	45.7	77	54.3	9.0
>High School	79	63.2	51	36.8	9.4

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Used a Blood Stool Test at Home in the Past Year, Aged 50 and Older

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	75	26.5	200	73.5	5.9
Gender					
Male	22	23.0	80	77.0	9.3
Female	53	29.7	120	70.3	7.6
Age					
50-64	30	24.0	105	76.0	8.5
≥65	45	29.2	95	70.8	8.3
Education					
≤High School	35	22.3	112	77.7	7.4
>High School	40	31.3	87	68.7	9.3

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Used a Blood Stool Test at Home in the Past 2 Years, Aged 50 and Older

*Chemung, Schuyler, Tioga*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	105	37.2	170	62.8	6.5
Gender					
Male	32	30.8	70	69.2	9.9
Female	73	42.9	100	57.1	8.3
Age					
50-64	46	35.9	89	64.1	9.2
≥65	59	38.6	81	61.4	9.1
Education					
≤High School	52	33.6	95	66.4	8.5
>High School	53	41.4	74	58.6	9.8

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

## Ever Had a Sigmoidoscopy or Colonoscopy, Aged 50 and Older

*Chemung, Schuyler, Tioga*

Demographic Groups	n <sup>1</sup>	Yes % <sup>2</sup>	n	No %	C.I. <sup>3</sup>
Total	157	59.7	124	40.3	6.5
Gender					
Male	65	67.2	39	32.8	9.9
Female	92	53.1	85	46.9	8.3
Age					
50-64	68	52.8	70	47.2	9.3
≥65	89	67.1	54	32.9	8.5
Education					
≤High School	82	60.8	67	39.2	8.5
>High School	74	58.3	57	41.7	9.7

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Had a Sigmoidoscopy or Colonoscopy in the Past 10 Years, Aged 50 and Older**

*Chemung, Schuyler, Tioga*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	148	56.4	133	43.6	6.5
Gender					
Male	60	62.5	44	37.5	10.2
Female	88	51.0	89	49.0	8.3
Age					
50-64	62	48.1	76	51.9	9.3
≥65	86	65.2	57	34.8	8.7
Education					
≤High School	79	58.3	70	41.7	8.7
>High School	69	54.7	62	45.3	9.8

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

**Had a Blood Stool Test Using a Home Kit in the Past Year or Had a Lower Endoscopy in the Past 10 Years, Aged 50 and Older**

*Chemung, Schuyler, Tioga*

<b>Demographic Groups</b>	<b>n<sup>1</sup></b>	<b>Yes %<sup>2</sup></b>	<b>n</b>	<b>No %</b>	<b>C.I.<sup>3</sup></b>
Total	173	65.1	111	34.9	6.1
Gender					
Male	65	67.3	40	32.7	9.6
Female	108	63.2	71	36.8	7.8
Age					
50-64	73	58.0	65	42.0	9.1
≥65	100	72.6	46	27.4	7.8
Education					
≤High School	91	63.6	61	36.4	8.3
>High School	82	67.2	49	32.8	9.0

<sup>1</sup>Percents based on row denominators of less than 50 are unstable and should be used with caution

<sup>2</sup>Weighted Percent

<sup>3</sup>95% Confidence Interval