

# **New York State Expanded Behavioral Risk Factor Surveillance System**

**Montgomery County**

**Final Report  
July 2008 - June 2009**

**New York State Department of Health  
December 2009**

## Contents

<b>Introduction</b>	12
Table 1. Montgomery County and New York State adult US Census Bureau population estimates, 2007	13
<b>General Health</b>	
Table 2a. Fair or poor general health among adults in Montgomery County and New York State	14
Table 2b. Fair or poor general health among adults in Montgomery County and New York State - Age-adjusted	15
Table 3a. Poor physical health 14 or more days within the past month among adults in Montgomery County and New York State	16
Table 3b. Poor physical health 14 or more days within the past month among adults in Montgomery County and New York State - Age-adjusted	17
Table 4a. Poor mental health 14 or more days within the past month among adults in Montgomery County and New York State	18
Table 4b. Poor mental health 14 or more days within the past month among adults in Montgomery County and New York State - Age-adjusted	19
Table 5a. Poor physical or mental health 14 or more days within the past month among adults in Montgomery County and New York State	20
Table 5b. Poor physical or mental health 14 or more days within the past month among adults in Montgomery County and New York State - Age-adjusted	21
<b>Quality Health Care</b>	
Table 6a. Health insurance among adults in Montgomery County and New York State	22
Table 6b. Health insurance among adults in Montgomery County and New York State - Age-adjusted	23
Table 7a. Health insurance among adults aged 18-64 in Montgomery County and New York State	24
Table 7b. Health insurance among adults aged 18-64 in Montgomery County and New York State - Age-adjusted	25

## Contents

Table 8a. No health insurance among adults in Montgomery County and New York State	26
Table 8b. No health insurance among adults in Montgomery County and New York State - Age-adjusted	27
Table 9a. No health insurance among adults aged 18-64 in Montgomery County and New York State	28
Table 9b. No health insurance among adults aged 18-64 in Montgomery County and New York State - Age-adjusted	29
Table 10a. Adults with regular health care provider in Montgomery County and New York State	30
Table 10b. Adults with regular health care provider in Montgomery County and New York State - Age-adjusted	31
Table 11a. Cost prevented visit to doctor within the past year among adults in Montgomery County and New York State	32
Table 11b. Cost prevented visit to doctor within the past year among adults in Montgomery County and New York State - Age-adjusted	33
Table 12a. Visited a doctor for routine checkup within the past year among adults in Montgomery County and New York State	34
Table 12b. Visited a doctor for routine checkup within the past year among adults in Montgomery County and New York State - Age-adjusted	35
Table 13a. Visited a doctor for routine checkup within the past 2 years among adults in Montgomery County and New York State	36
Table 13b. Visited a doctor for routine checkup within the past 2 years among adults in Montgomery County and New York State - Age-adjusted	37
Table 14a. Dentist visit within the past year among adults in Montgomery County and New York State	38
Table 14b. Dentist visit within the past year among adults in Montgomery County and New York State - Age-adjusted	39
Table 15a. Had teeth cleaned within the past year among adults in Montgomery County and New York State	40

## Contents

Table 15b. Had teeth cleaned within the past year among adults in Montgomery County and New York State - Age-adjusted	41
Table 16a. Permanent teeth extracted due to decay or gum disease among adults in Montgomery County and New York State	42
Table 16b. Permanent teeth extracted due to decay or gum disease among adults in Montgomery County and New York State - Age-adjusted	43
Table 17a. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in Montgomery County and New York State	44
Table 17b. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in Montgomery County and New York State - Age-adjusted	45
Table 18a. Flu shot within the past 12 months in Montgomery County and New York State	46
Table 18b. Flu shot within the past 12 months in Montgomery County and New York State - Age-adjusted	47
Table 19a. Flu vaccine that was sprayed in nose within the past 12 months in Montgomery County and New York State	48
Table 19b. Flu vaccine that was sprayed in nose within the past 12 months in Montgomery County and New York State - Age-adjusted	49
Table 20a. Flu shot or flu vaccine sprayed in nose within the past 12 months in Montgomery County and New York State	50
Table 20b. Flu shot or flu vaccine sprayed in nose within the past 12 months in Montgomery County and New York State - Age-adjusted	51
Table 21a. Flu shot among adults aged 65 and older in Montgomery County and New York State	52
Table 21b. Flu shot among adults aged 65 and older in Montgomery County and New York State - Age-adjusted	53
Table 22a. Pneumonia shot or pneumococcal vaccine in Montgomery County and New York State	54
Table 22b. Pneumonia shot or pneumococcal vaccine in Montgomery County and New York State - Age-adjusted	55

## Contents

Table 23a. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in Montgomery County and New York State	56
Table 23b. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in Montgomery County and New York State - Age-adjusted	57

### Cancer Screening

Table 24a. Ever had a mammogram among women aged 40 and older in Montgomery County and New York State	58
Table 24b. Ever had a mammogram among women aged 40 and older in Montgomery County and New York State - Age-adjusted	59
Table 25a. Had mammogram within the past 2 years among women aged 40 and older in Montgomery County and New York State	60
Table 25b. Had mammogram within the past 2 years among women aged 40 and older in Montgomery County and New York State - Age-adjusted	61
Table 26a. Had mammogram within the past 2 years among women aged 50 and older in Montgomery County and New York State	62
Table 26b. Had mammogram within the past 2 years among women aged 50 and older in Montgomery County and New York State - Age-adjusted	63
Table 27a. Ever had a Pap test among women in Montgomery County and New York State	64
Table 27b. Ever had a Pap test among women in Montgomery County and New York State - Age-adjusted	65
Table 28a. Had a Pap test within the past 3 years among women in Montgomery County and New York State	66
Table 28b. Had a Pap test within the past 3 years among women in Montgomery County and New York State - Age-adjusted	67
Table 29a. Ever had a digital rectal exam among men aged 40 and older in Montgomery County and New York State	68
Table 29b. Ever had a digital rectal exam among men aged 40 and older in Montgomery County and New York State - Age-adjusted	69

## Contents

Table 30a. Had digital rectal exam within the past 2 years among men aged 40 and older in Montgomery County and New York State	70
Table 30b. Had digital rectal exam within the past 2 years among men aged 40 and older in Montgomery County and New York State - Age-adjusted	71
Table 31a. Ever had a prostate specific antigen test among men aged 40 and older in Montgomery County and New York State	72
Table 31b. Ever had a prostate specific antigen test among men aged 40 and older in Montgomery County and New York State - Age-adjusted	73
Table 32a. Had prostate specific antigen test within the past 2 years among men aged 40 and older in Montgomery County and New York State	74
Table 32b. Had prostate specific antigen test within the past 2 years among men aged 40 and older in Montgomery County and New York State - Age-adjusted	75
Table 33a. Prostate cancer among men aged 40 and older in Montgomery County and New York State	76
Table 33b. Prostate cancer among men aged 40 and older in Montgomery County and New York State - Age-adjusted	77
Table 34a. Home blood stool test ever used among adults aged 50 and older in Montgomery County and New York State	78
Table 34b. Home blood stool test ever used among adults aged 50 and older in Montgomery County and New York State - Age-adjusted	79
Table 35a. Home blood stool test used within the past year among adults aged 50 and older in Montgomery County and New York State	80
Table 35b. Home blood stool test used within the past year among adults aged 50 and older in Montgomery County and New York State - Age-adjusted	81
Table 36a. Home blood stool test used within the past 2 years among adults aged 50 and older in Montgomery County and New York State	82
Table 36b. Home blood stool test used within the past 2 years among adults aged 50 and older in Montgomery County and New York State - Age-adjusted	83
Table 37a. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Montgomery County and New York State	84

## Contents

Table 37b. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Montgomery County and New York State - Age-adjusted	85
Table 38a. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Montgomery County and New York State	86
Table 38b. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Montgomery County and New York State - Age-adjusted	87
Table 39a. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Montgomery County and New York State	88
Table 39b. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Montgomery County and New York State - Age-adjusted	89

### Chronic Diseases

Table 40a. High blood pressure among adults in Montgomery County and New York State	90
Table 40b. High blood pressure among adults in Montgomery County and New York State - Age-adjusted	91
Table 41a. Blood pressure medication among adults with high blood pressure in Montgomery County and New York State	92
Table 41b. Blood pressure medication among adults with high blood pressure in Montgomery County and New York State - Age-adjusted	93
Table 42a. Ever had blood cholesterol checked among adults in Montgomery County and New York State	94
Table 42b. Ever had blood cholesterol checked among adults in Montgomery County and New York State - Age-adjusted	95
Table 43a. Blood cholesterol checked within the past 5 years among adults in Montgomery County and New York State	96
Table 43b. Blood cholesterol checked within the past 5 years among adults in Montgomery County and New York State - Age-adjusted	97
Table 44a. Cardiovascular disease among adults in Montgomery County and New York State	98

## Contents

Table 44b. Cardiovascular disease among adults in Montgomery County and New York State - Age-adjusted	99
Table 45a. Coronary heart disease among adults in Montgomery County and New York State	100
Table 45b. Coronary heart disease among adults in Montgomery County and New York State - Age-adjusted	101
Table 46a. Ever diagnosed with diabetes among adults in Montgomery County and New York State	102
Table 46b. Ever diagnosed with diabetes among adults in Montgomery County and New York State - Age-adjusted	103
Table 47a. Ever diagnosed with asthma (lifetime) among adults in Montgomery County and New York State	104
Table 47b. Ever diagnosed with asthma (lifetime) among adults in Montgomery County and New York State - Age-adjusted	105
Table 48a. Current asthma among adults in Montgomery County and New York State	106
Table 48b. Current asthma among adults in Montgomery County and New York State - Age-adjusted	107
Table 49a. Chronic joint symptoms among adults in Montgomery County and New York State	108
Table 49b. Chronic joint symptoms among adults in Montgomery County and New York State - Age-adjusted	109
Table 50a. Arthritis among adults in Montgomery County and New York State	110
Table 50b. Arthritis among adults in Montgomery County and New York State - Age-adjusted	111
Table 51a. Activity limitations because of physical, mental or emotional problems among adults in Montgomery County and New York State	112
Table 51b. Activity limitations because of physical, mental or emotional problems among adults in Montgomery County and New York State - Age-adjusted	113
Table 52a. Have health problem that requires the use of special equipment among adults in Montgomery County and New York State	114

## Contents

Table 52b. Have health problem that requires the use of special equipment among adults in Montgomery County and New York State - Age-adjusted	115
Table 53a. Disability among adults in Montgomery County and New York State	116
Table 53b. Disability among adults in Montgomery County and New York State - Age-adjusted	117
<b>Health Behaviors</b>	
Table 54a. Current smoking among adults in Montgomery County and New York State	118
Table 54b. Current smoking among adults in Montgomery County and New York State - Age-adjusted	119
Table 55a. Everyday smoking among adults in Montgomery County and New York State	120
Table 55b. Everyday smoking among adults in Montgomery County and New York State - Age-adjusted	121
Table 56a. Adults living in homes in which smoking is prohibited in Montgomery County and New York State	122
Table 56b. Adults living in homes in which smoking is prohibited in Montgomery County and New York State - Age-adjusted	123
Table 57a. Binge drinking within the past month among adults in Montgomery County and New York State	124
Table 57b. Binge drinking within the past month among adults in Montgomery County and New York State - Age-adjusted	125
Table 58a. Heavy drinking within the past month among adults in Montgomery County and New York State	126
Table 58b. Heavy drinking within the past month among adults in Montgomery County and New York State - Age-adjusted	127
Table 59a. Never or rarely been asked about sexual history during routine check-up among adults in Montgomery County and New York State	128
Table 59b. Never or rarely been asked about sexual history during routine check-up among adults in Montgomery County and New York State - Age-adjusted	129

## Contents

Table 60a. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults in Montgomery County and New York State	130
Table 60b. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults in Montgomery County and New York State - Age-adjusted	131
Table 61a. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults in Montgomery County and New York State	132
Table 61b. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults in Montgomery County and New York State - Age-adjusted	133

### Overweight / Obesity

Table 62a. Overweight among adults in Montgomery County and New York State	134
Table 62b. Overweight among adults in Montgomery County and New York State - Age-adjusted	135
Table 63a. Obesity among adults in Montgomery County and New York State	136
Table 63b. Obesity among adults in Montgomery County and New York State - Age-adjusted	137
Table 64a. Overweight or obesity among adults in Montgomery County and New York State	138
Table 64b. Overweight or obesity among adults in Montgomery County and New York State - Age-adjusted	139
Table 65a. Received advice about weight by a health professional among adults in Montgomery County and New York State	140
Table 65b. Received advice about weight by a health professional among adults in Montgomery County and New York State - Age-adjusted	141
Table 66a. Received advice to lose weight by a health professional among adults provided weight advice in Montgomery County and New York State	142
Table 66b. Received advice to lose weight by a health professional among adults provided weight advice in Montgomery County and New York State - Age-adjusted	143

## Contents

### Physical Activity / Nutrition

Table 67a. No leisure-time physical activity among adults in Montgomery County and New York State	144
Table 67b. No leisure-time physical activity among adults in Montgomery County and New York State - Age-adjusted	145
Table 68a. Consumption of 5 or more servings of fruits and vegetables per day among adults in Montgomery County and New York State	146
Table 68b. Consumption of 5 or more servings of fruits and vegetables per day among adults in Montgomery County and New York State - Age-adjusted	147

### Injury

Table 69a. Had a fall within the past 3 months among adults in Montgomery County and New York State	148
Table 69b. Had a fall within the past 3 months among adults in Montgomery County and New York State - Age-adjusted	149
Table 70a. Had a fall that caused injury within the past 3 months among adults in Montgomery County and New York State	150
Table 70b. Had a fall that caused injury within the past 3 months among adults in Montgomery County and New York State - Age-adjusted	151

## Introduction

The New York State Expanded Behavioral Risk Factor Surveillance System (Expanded BRFSS) is a random digit dialed telephone survey of adults (18 years of age and older) that is representative of the non-institutionalized civilian population with landline telephones living in New York State. The goal of the Expanded BRFSS is to collect county-specific data on preventive health practices, risk behaviors, injuries and preventable chronic and infectious diseases. Topics assessed by the Expanded BRFSS include tobacco use, physical inactivity, diet, use of cancer screening services and other factors linked to the leading causes of morbidity and mortality. For each topic, health indicators are based on answers to one or more questions in the survey. The survey results are weighted to more accurately represent the adult population living in each county. Weighting procedures adjust for the differential selection probabilities by race, age and gender. More detailed information on weighting procedures is available in the technical brief.

Survey data collection occurred in 58 geographic areas of New York State, including each of the 57 counties outside of New York City and New York City (5 counties) as a single area. A standard questionnaire was used in all areas. Data collection for the 2009 Expanded BRFSS took place during the period of July 2008 through June 2009. The goal was to complete about 650 interviews in each area. The final number of interviews varies because of the sampling design of the survey and differential response rates. More detailed information on the sampling design is available in the technical brief.

The New York State Department of Health contracted with RTI International, which is responsible for the data collection, sampling, weighting and creation of the statistical reports. RTI International subcontracted with Clearwater Research Inc. for data collection and additional expertise in sample design and data quality control.

This report presents both crude estimates and estimates that have been age-adjusted to the Year 2000 US Standard Population when they represent the entire adult population of an area. Estimates for specific age groups are not age-adjusted. Age adjustment is a statistical process applied to rates that allows communities with different age structures to be compared. Counties can compare their data with results from New York State as a whole and with other counties. Confidence intervals at the 95% level associated with each health indicator give an estimated range of values which is likely to include the actual prevalence of the indicator. Confidence intervals are similar to margins of error. When the confidence intervals of two estimates of the same indicator from different areas do not overlap, they may be said to be statistically significantly different, i.e., these differences are unlikely related to chance and are considered true differences. If there is any value that is included in both intervals, the two estimates are not statistically significantly different.

The survey results for each of the 69 health indicators included in the survey are typically provided by gender, age, race and ethnicity, education and income in the Expanded BRFSS county reports. The estimates for some of these groupings may be suppressed in geographic areas with low proportions of adult respondents in these groupings.

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 1. Montgomery County and New York State adult US Census Bureau population estimates,<sup>1</sup> 2007**

	Montgomery County		New York State	
	Adult Population <sup>1</sup>	Percent	Adult Population <sup>1</sup>	Percent
<b>Total</b>	37,649	100	14,884,315	100
<b>Gender</b>				
Male	17,784	47.2	7,103,791	47.7
Female	19,865	52.8	7,780,524	52.3
<b>Age (years)</b>				
18-34	10,196	27.1	4,497,479	30.2
35-44	6,141	16.3	2,844,115	19.1
45-54	7,180	19.1	2,854,316	19.2
55-64	5,719	15.2	2,142,000	14.4
≥ 65	8,413	22.3	2,546,405	17.1
<b>Race/Ethnicity</b>				
White Non-Hispanic	34,056	90.5	9,343,884	62.8
Black Non-Hispanic	521	1.4	2,156,941	14.5
Hispanic	2,714	7.2	2,258,717	15.2
Other Non-Hispanic	358	1.0	1,124,773	7.6

<sup>1</sup> Source: National Center for Health Statistics. Postcensal estimates of the resident population of the United States for July 1, 2000-July 1, 2007, by year, county, age, bridged race, Hispanic origin, and sex (Vintage 2007). Prepared under a collaborative arrangement with the U.S. Census Bureau; released August 7, 2008. Available from: [http://www.cdc.gov/nchs/nvss/bridged\\_race.htm](http://www.cdc.gov/nchs/nvss/bridged_race.htm) as of November 25, 2009.

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 2a. Fair or poor general health among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	640	6,500	17.3	(13.9 - 20.6)	38,124	2,531,400	17.1	(15.1 - 19.0)
<b>Gender</b>								
Male	224	3,000	17.2	(11.8 - 22.5)	14,599	1,069,700	15.1	(12.4 - 17.8)
Female	416	3,400	17.3	(13.2 - 21.5)	23,525	1,461,700	18.8	(16.1 - 21.6)
<b>Age (years)</b>								
18-34	*	*	*	*	3,940	535,800	12.0	(7.2 - 16.8)
35-44	*	*	*	*	4,985	289,200	10.0	(6.8 - 13.3)
45-54	138	1,000	13.5	(6.8 - 20.2)	7,819	507,800	17.9	(14.0 - 21.8)
55-64	126	1,100	20.1	(12.6 - 27.7)	8,757	424,700	19.9	(16.0 - 23.9)
≥ 65	231	2,700	31.9	(25.3 - 38.5)	12,623	773,900	30.5	(26.8 - 34.2)
<b>Race/Ethnicity</b>								
White Non-Hispanic	612	5,700	16.9	(13.6 - 20.2)	35,811	1,226,600	13.2	(11.7 - 14.6)
Black Non-Hispanic	*	*	*	*	739	457,300	21.3	(14.8 - 27.9)
Hispanic	*	*	*	*	861	684,100	30.4	(22.2 - 38.6)
Other Non-Hispanic	*	*	*	*	713	163,400	14.5	(5.1 - 24.0)
<b>Education</b>								
≤ High School	324	4,000	22.6	(17.4 - 27.8)	15,682	1,466,500	26.3	(22.6 - 30.0)
Some College	171	1,300	11.8	(6.2 - 17.4)	9,904	508,500	14.3	(10.9 - 17.8)
≥ College Degree	145	1,100	13.1	(6.9 - 19.4)	12,457	493,500	8.8	(6.5 - 11.1)
<b>Income</b>								
≤ \$24,999	192	2,200	24.6	(17.3 - 32.0)	9,633	1,269,200	36.3	(30.1 - 42.4)
\$25,000-\$49,999	184	2,500	23.2	(15.9 - 30.5)	9,649	447,600	15.4	(11.9 - 18.9)
\$50,000-\$74,999	*	*	*	*	5,769	237,800	12.1	(8.0 - 16.2)
≥ \$75,000	*	*	*	*	8,562	218,300	4.9	(3.6 - 6.1)

<sup>1</sup> Defined as reporting fair or poor general health

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have fair or poor health (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 2b. Fair or poor general health among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	640	15.7	(12.3 - 19.1)	38,124	16.7	(14.8 - 18.6)
<b>Gender</b>						
Male	224	16.3	(10.9 - 21.7)	14,599	15.3	(12.6 - 18.0)
Female	416	15.1	(10.9 - 19.4)	23,525	18.0	(15.2 - 20.7)
<b>Race/Ethnicity</b>						
White Non-Hispanic	612	15.4	(12.0 - 18.7)	35,811	12.9	(11.4 - 14.4)
Black Non-Hispanic	*	*	*	739	20.8	(14.2 - 27.3)
Hispanic	*	*	*	861	29.9	(21.8 - 37.9)
Other Non-Hispanic	*	*	*	713	13.8	(4.8 - 22.8)
<b>Education</b>						
≤ High School	324	21.2	(15.6 - 26.7)	15,682	25.8	(22.1 - 29.5)
Some College	171	10.7	(5.0 - 16.4)	9,904	14.1	(10.7 - 17.6)
≥ College Degree	145	11.5	(5.7 - 17.4)	12,457	8.5	(6.3 - 10.8)
<b>Income</b>						
≤ \$24,999	192	22.7	(14.9 - 30.5)	9,633	35.9	(29.7 - 42.0)
\$25,000-\$49,999	184	21.3	(13.7 - 28.9)	9,649	15.0	(11.6 - 18.5)
\$50,000-\$74,999	*	*	*	5,769	11.8	(7.7 - 16.0)
≥ \$75,000	*	*	*	8,562	4.7	(3.5 - 6.0)

<sup>1</sup> Defined as reporting fair or poor general health

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 3a. Poor physical health 14 or more days within the past month among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	624	4,100	11.0	(8.4 - 13.6)	37,327	1,472,900	10.1	(8.7 - 11.6)
<b>Gender</b>								
Male	226	1,800	10.0	(6.1 - 13.8)	14,325	604,200	8.7	(7.0 - 10.5)
Female	398	2,300	12.0	(8.5 - 15.4)	23,002	868,600	11.4	(9.2 - 13.6)
<b>Age (years)</b>								
18-34	*	*	*	*	3,891	268,100	6.1	(2.7 - 9.5)
35-44	*	*	*	*	4,948	232,100	8.1	(5.2 - 11.1)
45-54	136	500	6.5	(2.5 - 10.5)	7,764	350,900	12.5	(9.4 - 15.6)
55-64	123	900	16.7	(9.7 - 23.7)	8,621	271,200	13.0	(10.3 - 15.8)
≥ 65	220	1,700	20.7	(15.0 - 26.4)	12,103	350,600	14.5	(12.1 - 16.9)
<b>Race/Ethnicity</b>								
White Non-Hispanic	597	3,500	10.6	(8.1 - 13.1)	35,073	929,700	10.2	(8.8 - 11.6)
Black Non-Hispanic	*	*	*	*	724	208,200	9.9	(4.8 - 14.9)
Hispanic	*	*	*	*	835	250,300	11.5	(6.5 - 16.4)
Other Non-Hispanic	*	*	*	*	695	84,600	7.6	(1.3 - 13.9)
<b>Education</b>								
≤ High School	311	2,300	13.3	(9.4 - 17.3)	15,178	692,800	12.9	(10.6 - 15.3)
Some College	169	900	8.7	(4.3 - 13.0)	9,751	362,500	10.3	(7.4 - 13.2)
≥ College Degree	144	800	9.2	(3.9 - 14.5)	12,327	359,300	6.4	(4.8 - 8.1)
<b>Income</b>								
≤ \$24,999	184	1,500	17.3	(11.2 - 23.4)	9,294	625,600	18.4	(13.6 - 23.3)
\$25,000-\$49,999	180	1,100	10.5	(6.0 - 14.9)	9,495	279,100	9.8	(7.3 - 12.3)
\$50,000-\$74,999	*	*	*	*	5,733	186,400	9.5	(5.8 - 13.2)
≥ \$75,000	*	*	*	*	8,510	219,100	4.9	(3.6 - 6.2)

<sup>1</sup> Defined as 14 or more days within the past month when physical health was not good

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults with poor physical health 14 or more days within the past month (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 3b. Poor physical health 14 or more days within the past month among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	624	9.9	(7.3 - 12.5)	37,327	9.9	(8.4 - 11.4)
<b>Gender</b>						
Male	226	9.3	(5.4 - 13.2)	14,325	8.7	(6.9 - 10.5)
Female	398	10.5	(7.1 - 13.9)	23,002	11.0	(8.7 - 13.2)
<b>Race/Ethnicity</b>						
White Non-Hispanic	597	9.6	(7.0 - 12.1)	35,073	9.9	(8.5 - 11.3)
Black Non-Hispanic	*	*	*	724	9.7	(4.6 - 14.9)
Hispanic	*	*	*	835	11.3	(6.3 - 16.2)
Other Non-Hispanic	*	*	*	695	7.6	(1.2 - 14.0)
<b>Education</b>						
≤ High School	311	12.0	(8.1 - 16.0)	15,178	12.7	(10.2 - 15.1)
Some College	169	7.7	(3.3 - 12.0)	9,751	10.1	(7.2 - 12.9)
≥ College Degree	144	8.7	(3.4 - 14.1)	12,327	6.3	(4.7 - 8.0)
<b>Income</b>						
≤ \$24,999	184	15.7	(9.5 - 21.9)	9,294	18.4	(13.5 - 23.4)
\$25,000-\$49,999	180	9.1	(4.8 - 13.5)	9,495	9.4	(7.0 - 11.9)
\$50,000-\$74,999	*	*	*	5,733	9.0	(5.3 - 12.6)
≥ \$75,000	*	*	*	8,510	4.9	(3.5 - 6.2)

<sup>1</sup> Defined as 14 or more days within the past month when physical health was not good

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 4a. Poor mental health 14 or more days within the past month among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	630	3,800	10.1	(7.5 - 12.7)	37,528	1,473,200	10.1	(8.6 - 11.5)
<b>Gender</b>								
Male	224	1,100	6.3	(3.0 - 9.7)	14,407	555,200	7.9	(6.3 - 9.6)
Female	406	2,600	13.5	(9.7 - 17.4)	23,121	918,000	12.0	(9.7 - 14.3)
<b>Age (years)</b>								
18-34	71	1,100	10.9	(4.3 - 17.6)	3,889	496,700	11.2	(7.4 - 15.1)
35-44	*	*	*	*	4,947	254,100	8.9	(6.7 - 11.1)
45-54	134	500	6.7	(2.7 - 10.7)	7,749	347,200	12.3	(9.0 - 15.6)
55-64	125	900	16.7	(9.8 - 23.6)	8,623	211,300	10.1	(7.9 - 12.3)
≥ 65	227	800	9.2	(5.2 - 13.3)	12,320	163,900	6.7	(4.7 - 8.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	603	3,300	9.9	(7.3 - 12.4)	35,266	899,200	9.8	(8.4 - 11.2)
Black Non-Hispanic	*	*	*	*	722	227,800	10.9	(5.4 - 16.4)
Hispanic	*	*	*	*	845	263,700	11.8	(6.8 - 16.8)
Other Non-Hispanic	*	*	*	*	695	82,500	7.4	(1.4 - 13.5)
<b>Education</b>								
≤ High School	319	1,900	10.8	(7.0 - 14.6)	15,335	665,000	12.2	(10.0 - 14.5)
Some College	170	1,100	9.7	(5.1 - 14.3)	9,771	404,500	11.5	(8.3 - 14.8)
≥ College Degree	141	800	9.2	(3.7 - 14.8)	12,342	346,600	6.2	(4.5 - 7.9)
<b>Income</b>								
≤ \$24,999	188	1,100	12.3	(6.9 - 17.8)	9,398	611,500	17.9	(13.0 - 22.8)
\$25,000-\$49,999	183	900	8.6	(4.5 - 12.6)	9,547	260,300	9.0	(6.7 - 11.4)
\$50,000-\$74,999	94	600	9.1	(3.0 - 15.1)	5,726	187,300	9.6	(6.0 - 13.1)
≥ \$75,000	*	*	*	*	8,520	276,500	6.2	(4.6 - 7.7)

<sup>1</sup> Defined as 14 or more days within the past month when mental health was not good

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults with poor mental health 14 or more days within the past month (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 4b. Poor mental health 14 or more days within the past month among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	630	9.9	(7.1 - 12.6)	37,528	10.2	(8.7 - 11.7)
<b>Gender</b>						
Male	224	5.9	(2.6 - 9.3)	14,407	7.9	(6.3 - 9.6)
Female	406	13.4	(9.2 - 17.6)	23,121	12.2	(9.8 - 14.6)
<b>Race/Ethnicity</b>						
White Non-Hispanic	603	9.7	(6.9 - 12.5)	35,266	10.0	(8.5 - 11.4)
Black Non-Hispanic	*	*	*	722	11.1	(5.5 - 16.7)
Hispanic	*	*	*	845	11.5	(6.5 - 16.4)
Other Non-Hispanic	*	*	*	695	7.6	(1.5 - 13.7)
<b>Education</b>						
≤ High School	319	10.1	(6.1 - 14.0)	15,335	12.3	(10.0 - 14.6)
Some College	170	9.3	(4.4 - 14.2)	9,771	11.8	(8.5 - 15.1)
≥ College Degree	141	10.2	(4.0 - 16.4)	12,342	6.3	(4.6 - 8.0)
<b>Income</b>						
≤ \$24,999	188	12.1	(6.0 - 18.1)	9,398	18.2	(13.2 - 23.1)
\$25,000-\$49,999	183	7.9	(4.0 - 11.9)	9,547	9.4	(6.9 - 11.9)
\$50,000-\$74,999	94	9.4	(2.6 - 16.2)	5,726	9.5	(5.9 - 13.1)
≥ \$75,000	*	*	*	8,520	6.2	(4.6 - 7.7)

<sup>1</sup> Defined as 14 or more days within the past month when mental health was not good

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 5a. Poor physical or mental health 14 or more days within the past month among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	622	6,500	17.6	(14.3 - 20.8)	37,005	2,385,900	16.5	(14.8 - 18.2)
<b>Gender</b>								
Male	226	2,400	13.3	(8.8 - 17.8)	14,231	937,700	13.6	(11.4 - 15.8)
Female	396	4,200	21.5	(16.9 - 26.1)	22,774	1,448,200	19.1	(16.5 - 21.7)
<b>Age (years)</b>								
18-34	71	1,400	14.2	(6.6 - 21.8)	3,859	602,600	13.8	(9.7 - 17.9)
35-44	74	700	10.9	(3.9 - 18.0)	4,924	388,300	13.7	(10.4 - 17.0)
45-54	134	800	11.6	(6.3 - 16.8)	7,722	551,000	19.6	(15.8 - 23.4)
55-64	123	1,500	26.7	(18.2 - 35.2)	8,545	395,500	19.2	(16.1 - 22.3)
≥ 65	220	2,100	25.8	(19.6 - 31.9)	11,955	448,500	18.8	(15.9 - 21.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	595	5,900	17.7	(14.4 - 21.0)	34,778	1,485,700	16.3	(14.8 - 17.8)
Black Non-Hispanic	*	*	*	*	715	343,700	16.5	(10.3 - 22.7)
Hispanic	*	*	*	*	832	412,900	19.0	(12.6 - 25.3)
Other Non-Hispanic	*	*	*	*	680	143,500	13.0	(5.0 - 21.0)
<b>Education</b>								
≤ High School	311	3,400	19.6	(14.8 - 24.4)	15,018	1,077,200	20.3	(17.3 - 23.2)
Some College	168	1,600	14.9	(9.2 - 20.5)	9,672	615,500	17.6	(14.1 - 21.1)
≥ College Degree	143	1,500	16.9	(9.8 - 23.9)	12,244	634,200	11.4	(9.2 - 13.6)
<b>Income</b>								
≤ \$24,999	183	2,300	25.5	(17.6 - 33.3)	9,199	924,000	27.3	(21.9 - 32.8)
\$25,000-\$49,999	180	1,600	15.6	(10.1 - 21.1)	9,425	456,000	16.1	(13.0 - 19.2)
\$50,000-\$74,999	94	900	14.1	(6.5 - 21.7)	5,696	312,000	16.0	(11.4 - 20.5)
≥ \$75,000	*	*	*	*	8,475	447,600	10.0	(8.1 - 12.0)

<sup>1</sup> Defined as 14 or more days within the past month when physical or mental health prevented usual activities, such as self-care, work or recreation

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults with poor physical or mental health 14 or more days within the past month (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 5b. Poor physical or mental health 14 or more days within the past month among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	622	16.5	(13.1 - 19.9)	37,005	16.4	(14.7 - 18.2)
<b>Gender</b>						
Male	226	12.4	(7.9 - 16.9)	14,231	13.6	(11.4 - 15.8)
Female	396	20.2	(15.3 - 25.1)	22,774	19.0	(16.3 - 21.6)
<b>Race/Ethnicity</b>						
White Non-Hispanic	595	16.7	(13.2 - 20.2)	34,778	16.3	(14.7 - 17.8)
Black Non-Hispanic	*	*	*	715	16.7	(10.3 - 23.0)
Hispanic	*	*	*	832	18.5	(12.2 - 24.8)
Other Non-Hispanic	*	*	*	680	13.1	(5.0 - 21.3)
<b>Education</b>						
≤ High School	311	18.0	(13.0 - 23.0)	15,018	20.1	(17.1 - 23.1)
Some College	168	13.8	(8.0 - 19.6)	9,672	17.6	(14.1 - 21.1)
≥ College Degree	143	17.0	(9.5 - 24.5)	12,244	11.4	(9.2 - 13.7)
<b>Income</b>						
≤ \$24,999	183	24.0	(15.6 - 32.4)	9,199	27.6	(22.1 - 33.1)
\$25,000-\$49,999	180	14.2	(8.7 - 19.6)	9,425	16.1	(12.9 - 19.3)
\$50,000-\$74,999	94	14.8	(6.4 - 23.2)	5,696	15.6	(11.0 - 20.1)
≥ \$75,000	*	*	*	8,475	10.0	(8.0 - 12.0)

<sup>1</sup> Defined as 14 or more days within the past month when physical or mental health prevented usual activities, such as self-care, work or recreation

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 6a. Health insurance among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	641	33,000	87.8	(83.3 - 92.3)	38,119	12,854,200	86.6	(84.2 - 89.0)
<b>Gender</b>								
Male	227	14,800	83.0	(74.9 - 91.1)	14,597	5,881,100	83.0	(78.9 - 87.1)
Female	414	18,200	92.1	(87.9 - 96.2)	23,522	6,973,100	89.9	(87.2 - 92.6)
<b>Age (years)</b>								
18-34	*	*	*	*	3,910	3,302,100	74.3	(67.8 - 80.7)
35-44	74	5,300	85.8	(76.4 - 95.1)	4,986	2,560,300	88.9	(84.7 - 93.1)
45-54	138	6,200	86.8	(79.2 - 94.4)	7,831	2,530,100	88.9	(84.6 - 93.2)
55-64	127	5,300	92.8	(88.0 - 97.5)	8,764	1,986,700	92.8	(90.2 - 95.5)
≥ 65	231	8,300	98.6	(97.0 - 100)	12,628	2,475,000	97.8	(96.7 - 98.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	613	30,600	90.1	(86.7 - 93.4)	35,811	8,591,200	92.2	(90.6 - 93.8)
Black Non-Hispanic	*	*	*	*	736	1,781,600	82.7	(75.3 - 90.1)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	324	15,900	89.0	(84.0 - 94.1)	15,673	4,448,900	79.7	(74.9 - 84.5)
Some College	*	*	*	*	9,906	3,004,600	85.0	(80.1 - 89.8)
≥ College Degree	145	8,300	94.9	(89.4 - 100)	12,459	5,361,700	95.3	(93.3 - 97.2)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,640	2,437,200	69.7	(62.5 - 76.8)
\$25,000-\$49,999	184	9,600	90.1	(83.9 - 96.3)	9,662	2,384,000	82.2	(76.1 - 88.3)
\$50,000-\$74,999	95	5,900	91.6	(83.4 - 99.8)	5,772	1,855,700	94.0	(91.2 - 96.7)
≥ \$75,000	97	7,200	97.1	(93.1 - 100)	8,567	4,388,400	97.5	(96.6 - 98.5)

<sup>1</sup> Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have health insurance (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 6b. Health insurance among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	641	86.8	(82.0 - 91.7)	38,119	86.7	(84.4 - 89.1)
<b>Gender</b>						
Male	227	82.1	(73.6 - 90.6)	14,597	83.6	(79.7 - 87.5)
Female	414	91.0	(86.2 - 95.9)	23,522	89.6	(86.8 - 92.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	613	89.3	(85.5 - 93.0)	35,811	92.1	(90.5 - 93.7)
Black Non-Hispanic	*	*	*	736	82.5	(75.0 - 90.0)
Hispanic	*	*	*	860	68.7	(59.0 - 78.4)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	324	87.9	(82.1 - 93.7)	15,673	80.1	(75.4 - 84.7)
Some College	*	*	*	9,906	84.6	(79.7 - 89.4)
≥ College Degree	145	94.6	(88.5 - 100)	12,459	95.4	(93.6 - 97.2)
<b>Income</b>						
≤ \$24,999	*	*	*	9,640	70.2	(63.2 - 77.1)
\$25,000-\$49,999	184	88.2	(80.8 - 95.6)	9,662	81.9	(76.0 - 87.9)
\$50,000-\$74,999	95	90.4	(81.0 - 99.8)	5,772	94.0	(91.3 - 96.7)
≥ \$75,000	97	97.0	(92.9 - 100)	8,567	97.5	(96.6 - 98.5)

<sup>1</sup> Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 7a. Health insurance among adults aged 18-64<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	403	24,500	84.5	(78.8 - 90.2)	25,165	10,251,300	84.2	(81.3 - 87.1)
<b>Gender</b>								
Male	148	11,500	79.7	(70.0 - 89.4)	10,078	4,839,900	80.3	(75.5 - 85.0)
Female	255	13,000	89.3	(83.8 - 94.9)	15,087	5,411,400	88.0	(84.7 - 91.3)
<b>Age (years)</b>								
18-34	*	*	*	*	3,910	3,302,100	74.3	(67.8 - 80.7)
35-44	74	5,300	85.8	(76.4 - 95.1)	4,981	2,556,600	88.9	(84.7 - 93.1)
45-54	138	6,200	86.8	(79.2 - 94.4)	7,791	2,472,900	88.7	(84.3 - 93.0)
55-64	120	5,000	92.5	(87.5 - 97.4)	8,483	1,919,700	92.8	(90.1 - 95.5)
≥ 65	-	-	-	-	-	-	-	-
<b>Race/Ethnicity</b>								
White Non-Hispanic	381	22,500	87.3	(83.0 - 91.7)	23,354	6,582,200	90.3	(88.3 - 92.4)
Black Non-Hispanic	*	*	*	*	567	1,464,100	80.2	(71.6 - 88.7)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	164	10,300	84.8	(77.6 - 92.0)	9,037	3,191,500	74.6	(68.6 - 80.6)
Some College	*	*	*	*	7,031	2,540,400	82.9	(77.4 - 88.4)
≥ College Degree	110	6,800	93.9	(87.4 - 100)	9,068	4,502,400	94.5	(92.2 - 96.8)
<b>Income</b>								
≤ \$24,999	*	*	*	*	4,918	1,707,200	62.6	(54.1 - 71.2)
\$25,000-\$49,999	115	6,900	86.8	(78.6 - 95.0)	6,180	1,783,000	77.7	(70.3 - 85.1)
\$50,000-\$74,999	81	5,200	90.6	(81.4 - 99.7)	4,554	1,553,400	92.9	(89.7 - 96.1)
≥ \$75,000	91	6,900	96.9	(92.7 - 100)	7,324	3,987,600	97.3	(96.3 - 98.3)

<sup>1</sup> Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 18-64 who have health insurance (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 7b. Health insurance among adults aged 18-64<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	403	84.4	(78.6 - 90.1)	25,165	84.3	(81.5 - 87.2)
<b>Gender</b>						
Male	*	*	*	10,078	80.5	(75.9 - 85.1)
Female	255	89.2	(83.5 - 95.0)	15,087	87.9	(84.6 - 91.2)
<b>Race/Ethnicity</b>						
White Non-Hispanic	381	87.2	(82.6 - 91.8)	23,354	90.3	(88.3 - 92.3)
Black Non-Hispanic	*	*	*	567	80.0	(71.3 - 88.6)
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	164	84.7	(77.1 - 92.3)	9,037	75.1	(69.3 - 80.9)
Some College	*	*	*	7,031	82.6	(77.1 - 88.1)
≥ College Degree	110	93.8	(86.9 - 100)	9,068	94.6	(92.5 - 96.8)
<b>Income</b>						
≤ \$24,999	*	*	*	4,918	63.2	(54.9 - 71.6)
\$25,000-\$49,999	115	85.3	(76.3 - 94.4)	6,180	77.5	(70.2 - 84.7)
\$50,000-\$74,999	*	*	*	4,554	93.0	(89.8 - 96.2)
≥ \$75,000	91	96.9	(92.6 - 100)	7,324	97.3	(96.3 - 98.3)

<sup>1</sup> Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 8a. No health insurance among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	641	4,600	12.2	(7.7 - 16.7)	38,119	1,989,300	13.4	(11.0 - 15.8)
<b>Gender</b>								
Male	227	3,000	17.0	(8.9 - 25.1)	14,597	1,207,400	17.0	(12.9 - 21.1)
Female	414	1,600	7.9	(3.8 - 12.1)	23,522	781,900	10.1	(7.4 - 12.8)
<b>Age (years)</b>								
18-34	*	*	*	*	3,910	1,144,100	25.7	(19.3 - 32.2)
35-44	*	*	*	*	4,986	319,200	11.1	(6.9 - 15.3)
45-54	138	900	13.2	(5.6 - 20.8)	7,831	316,500	11.1	(6.8 - 15.4)
55-64	127	400	7.2	(2.5 - 12.0)	8,764	153,300	7.2	(4.5 - 9.8)
≥ 65	*	*	*	*	12,628	56,200	2.2	(1.2 - 3.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	613	3,400	9.9	(6.6 - 13.3)	35,811	727,100	7.8	(6.2 - 9.4)
Black Non-Hispanic	*	*	*	*	736	372,800	17.3	(9.9 - 24.7)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	324	2,000	11.0	(5.9 - 16.0)	15,673	1,134,400	20.3	(15.5 - 25.1)
Some College	*	*	*	*	9,906	531,400	15.0	(10.2 - 19.9)
≥ College Degree	*	*	*	*	12,459	266,300	4.7	(2.8 - 6.7)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,640	1,061,200	30.3	(23.2 - 37.5)
\$25,000-\$49,999	184	1,100	9.9	(3.7 - 16.1)	9,662	516,700	17.8	(11.7 - 23.9)
\$50,000-\$74,999	*	*	*	*	5,772	119,100	6.0	(3.3 - 8.8)
≥ \$75,000	*	*	*	*	8,567	111,100	2.5	(1.5 - 3.4)

<sup>1</sup> Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have no health insurance (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 8b. No health insurance among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	641	13.2	(8.3 - 18.0)	38,119	13.3	(10.9 - 15.6)
<b>Gender</b>						
Male	227	17.9	(9.4 - 26.4)	14,597	16.4	(12.5 - 20.3)
Female	414	9.0	(4.1 - 13.8)	23,522	10.4	(7.7 - 13.2)
<b>Race/Ethnicity</b>						
White Non-Hispanic	613	10.7	(7.0 - 14.5)	35,811	7.9	(6.3 - 9.5)
Black Non-Hispanic	*	*	*	736	17.5	(10.0 - 25.0)
Hispanic	*	*	*	860	31.3	(21.6 - 41.0)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	324	12.1	(6.3 - 17.9)	15,673	19.9	(15.3 - 24.6)
Some College	*	*	*	9,906	15.4	(10.6 - 20.3)
≥ College Degree	*	*	*	12,459	4.6	(2.8 - 6.4)
<b>Income</b>						
≤ \$24,999	*	*	*	9,640	29.8	(22.9 - 36.8)
\$25,000-\$49,999	184	11.8	(4.4 - 19.2)	9,662	18.1	(12.1 - 24.0)
\$50,000-\$74,999	*	*	*	5,772	6.0	(3.3 - 8.7)
≥ \$75,000	*	*	*	8,567	2.5	(1.5 - 3.4)

<sup>1</sup> Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 9a. No health insurance among adults aged 18-64<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	403	4,500	15.5	(9.8 - 21.2)	25,165	1,928,400	15.8	(12.9 - 18.7)
<b>Gender</b>								
Male	148	2,900	20.3	(10.6 - 30.0)	10,078	1,190,200	19.7	(15.0 - 24.5)
Female	255	1,500	10.7	(5.1 - 16.2)	15,087	738,200	12.0	(8.7 - 15.3)
<b>Age (years)</b>								
18-34	*	*	*	*	3,910	1,144,100	25.7	(19.3 - 32.2)
35-44	*	*	*	*	4,981	319,200	11.1	(6.9 - 15.3)
45-54	138	900	13.2	(5.6 - 20.8)	7,791	316,100	11.3	(7.0 - 15.7)
55-64	120	400	7.5	(2.6 - 12.5)	8,483	148,900	7.2	(4.5 - 9.9)
≥ 65	-	-	-	-	-	-	-	-
<b>Race/Ethnicity</b>								
White Non-Hispanic	381	3,300	12.7	(8.3 - 17.0)	23,354	703,100	9.7	(7.6 - 11.7)
Black Non-Hispanic	*	*	*	*	567	361,800	19.8	(11.3 - 28.4)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	164	1,800	15.2	(8.0 - 22.4)	9,037	1,087,700	25.4	(19.4 - 31.4)
Some College	*	*	*	*	7,031	522,800	17.1	(11.6 - 22.6)
≥ College Degree	*	*	*	*	9,068	261,200	5.5	(3.2 - 7.8)
<b>Income</b>								
≤ \$24,999	*	*	*	*	4,918	1,018,000	37.4	(28.8 - 45.9)
\$25,000-\$49,999	115	1,100	13.2	(5.0 - 21.4)	6,180	511,100	22.3	(14.9 - 29.7)
\$50,000-\$74,999	*	*	*	*	4,554	118,800	7.1	(3.9 - 10.3)
≥ \$75,000	*	*	*	*	7,324	110,700	2.7	(1.7 - 3.7)

<sup>1</sup> Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 18-64 who have no health insurance (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 9b. No health insurance among adults aged 18-64<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	403	15.6	(9.9 - 21.4)	25,165	15.7	(12.8 - 18.5)
<b>Gender</b>						
Male	*	*	*	10,078	19.5	(14.9 - 24.1)
Female	255	10.8	(5.0 - 16.5)	15,087	12.1	(8.8 - 15.4)
<b>Race/Ethnicity</b>						
White Non-Hispanic	381	12.8	(8.2 - 17.4)	23,354	9.7	(7.7 - 11.7)
Black Non-Hispanic	*	*	*	567	20.0	(11.4 - 28.7)
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	164	15.3	(7.7 - 22.9)	9,037	24.9	(19.1 - 30.7)
Some College	*	*	*	7,031	17.4	(11.9 - 22.9)
≥ College Degree	*	*	*	9,068	5.4	(3.2 - 7.5)
<b>Income</b>						
≤ \$24,999	*	*	*	4,918	36.8	(28.4 - 45.1)
\$25,000-\$49,999	115	14.7	(5.6 - 23.7)	6,180	22.5	(15.3 - 29.8)
\$50,000-\$74,999	*	*	*	4,554	7.0	(3.8 - 10.2)
≥ \$75,000	*	*	*	7,324	2.7	(1.7 - 3.7)

<sup>1</sup> Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 10a. Adults with regular health care provider<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	642	33,100	87.9	(83.4 - 92.3)	38,135	12,324,300	83.0	(80.4 - 85.6)
<b>Gender</b>								
Male	227	14,300	80.4	(72.1 - 88.7)	14,593	5,494,300	77.6	(73.2 - 82.0)
Female	415	18,800	94.6	(91.5 - 97.7)	23,542	6,830,000	87.8	(85.1 - 90.5)
<b>Age (years)</b>								
18-34	*	*	*	*	3,929	2,931,600	65.8	(59.2 - 72.5)
35-44	74	5,400	87.8	(78.9 - 96.8)	4,988	2,471,000	86.1	(81.9 - 90.3)
45-54	138	6,000	83.8	(75.9 - 91.6)	7,823	2,529,300	88.6	(84.6 - 92.7)
55-64	128	5,400	94.8	(90.9 - 98.7)	8,770	2,002,000	93.6	(91.7 - 95.4)
≥ 65	231	8,200	97.4	(95.3 - 99.6)	12,625	2,390,400	94.1	(92.2 - 96.0)
<b>Race/Ethnicity</b>								
White Non-Hispanic	614	30,900	90.8	(87.6 - 94.1)	35,821	8,227,700	88.2	(86.4 - 89.9)
Black Non-Hispanic	*	*	*	*	740	1,753,200	81.3	(72.6 - 89.9)
Hispanic	*	*	*	*	860	1,453,100	64.7	(54.7 - 74.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	324	15,700	87.9	(82.6 - 93.2)	15,692	4,249,200	75.9	(70.9 - 80.8)
Some College	*	*	*	*	9,904	2,935,000	82.8	(77.2 - 88.4)
≥ College Degree	146	8,300	94.4	(89.8 - 99.0)	12,456	5,046,100	89.8	(87.1 - 92.6)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,644	2,554,200	72.7	(65.9 - 79.6)
\$25,000-\$49,999	185	9,800	91.8	(86.1 - 97.4)	9,655	2,243,100	77.3	(70.0 - 84.6)
\$50,000-\$74,999	95	5,800	90.4	(83.5 - 97.4)	5,771	1,767,500	89.5	(85.2 - 93.9)
≥ \$75,000	97	6,900	92.9	(85.9 - 99.9)	8,560	4,122,800	91.9	(90.2 - 93.7)

<sup>1</sup> Defined as having one or more persons respondent thinks of as a personal doctor or health care provider

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have at least one person they think of as their personal doctor or health care provider (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 10b. Adults with regular health care provider<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	642	87.3	(82.6 - 92.0)	38,135	83.0	(80.4 - 85.5)
<b>Gender</b>						
Male	227	79.4	(70.7 - 88.2)	14,593	78.1	(73.9 - 82.3)
Female	415	94.4	(90.9 - 97.8)	23,542	87.4	(84.6 - 90.1)
<b>Race/Ethnicity</b>						
White Non-Hispanic	614	90.4	(86.8 - 93.9)	35,821	88.0	(86.2 - 89.7)
Black Non-Hispanic	*	*	*	740	81.4	(73.0 - 89.8)
Hispanic	*	*	*	860	65.2	(55.5 - 75.0)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	324	86.7	(80.6 - 92.8)	15,692	75.8	(71.0 - 80.6)
Some College	*	*	*	9,904	82.9	(77.5 - 88.3)
≥ College Degree	146	94.8	(90.5 - 99.0)	12,456	89.8	(87.1 - 92.4)
<b>Income</b>						
≤ \$24,999	*	*	*	9,644	72.9	(66.1 - 79.6)
\$25,000-\$49,999	185	91.1	(84.9 - 97.4)	9,655	77.4	(70.5 - 84.3)
\$50,000-\$74,999	95	91.1	(84.3 - 97.8)	5,771	89.6	(85.3 - 94.0)
≥ \$75,000	97	92.8	(85.4 - 100)	8,560	91.6	(89.8 - 93.4)

<sup>1</sup> Defined as having one or more persons respondent thinks of as a personal doctor or health care provider

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 11a. Cost prevented visit to doctor within the past year among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	642	4,300	11.6	(8.3 - 14.8)	38,143	2,034,800	13.7	(11.6 - 15.8)
<b>Gender</b>								
Male	227	2,300	12.9	(7.3 - 18.5)	14,607	903,300	12.7	(9.5 - 16.0)
Female	415	2,100	10.4	(6.9 - 13.9)	23,536	1,131,500	14.6	(11.8 - 17.4)
<b>Age (years)</b>								
18-34	71	1,500	15.1	(6.5 - 23.7)	3,939	947,300	21.2	(15.6 - 26.9)
35-44	74	800	13.6	(4.9 - 22.2)	4,987	347,900	12.1	(8.0 - 16.2)
45-54	138	1,100	15.0	(7.7 - 22.3)	7,825	382,200	13.5	(9.9 - 17.1)
55-64	128	400	7.2	(2.6 - 11.8)	8,766	236,800	11.1	(8.0 - 14.2)
≥ 65	231	500	5.8	(2.3 - 9.4)	12,626	120,600	4.8	(2.5 - 7.0)
<b>Race/Ethnicity</b>								
White Non-Hispanic	614	3,700	10.8	(7.6 - 13.9)	35,833	984,600	10.6	(8.9 - 12.2)
Black Non-Hispanic	*	*	*	*	738	280,000	13.1	(7.2 - 19.0)
Hispanic	*	*	*	*	859	597,900	26.5	(17.3 - 35.7)
Other Non-Hispanic	*	*	*	*	713	172,300	15.3	(5.7 - 24.9)
<b>Education</b>								
≤ High School	325	1,800	10.0	(5.7 - 14.3)	15,696	996,600	17.8	(13.8 - 21.9)
Some College	172	1,800	16.4	(8.8 - 23.9)	9,911	512,400	14.5	(10.4 - 18.6)
≥ College Degree	145	800	8.7	(3.6 - 13.9)	12,453	466,900	8.3	(6.0 - 10.6)
<b>Income</b>								
≤ \$24,999	193	1,200	13.7	(7.4 - 19.9)	9,639	854,500	24.3	(18.4 - 30.3)
\$25,000-\$49,999	185	1,700	15.8	(8.8 - 22.8)	9,656	532,900	18.4	(12.5 - 24.3)
\$50,000-\$74,999	*	*	*	*	5,776	202,700	10.3	(6.0 - 14.5)
≥ \$75,000	*	*	*	*	8,561	192,000	4.3	(2.9 - 5.6)

<sup>1</sup> Defined as needing to see a doctor within the past year but not being able to because of cost

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who did not see a doctor because of cost within the past year (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 11b. Cost prevented visit to doctor within the past year among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	642	12.1	(8.5 - 15.6)	38,143	13.8	(11.7 - 15.9)
<b>Gender</b>						
Male	227	13.6	(7.5 - 19.7)	14,607	12.5	(9.4 - 15.6)
Female	415	10.7	(6.8 - 14.6)	23,536	15.0	(12.1 - 17.9)
<b>Race/Ethnicity</b>						
White Non-Hispanic	614	11.3	(7.8 - 14.8)	35,833	10.8	(9.2 - 12.5)
Black Non-Hispanic	*	*	*	738	13.2	(7.2 - 19.3)
Hispanic	*	*	*	859	26.1	(17.2 - 35.1)
Other Non-Hispanic	*	*	*	713	14.8	(5.5 - 24.2)
<b>Education</b>						
≤ High School	325	10.1	(5.4 - 14.8)	15,696	17.8	(13.8 - 21.7)
Some College	172	17.2	(9.1 - 25.2)	9,911	15.3	(11.0 - 19.5)
≥ College Degree	145	9.3	(3.8 - 14.9)	12,453	8.2	(6.0 - 10.5)
<b>Income</b>						
≤ \$24,999	193	14.0	(7.1 - 20.9)	9,639	24.5	(18.6 - 30.4)
\$25,000-\$49,999	185	17.7	(9.7 - 25.7)	9,656	18.7	(12.9 - 24.6)
\$50,000-\$74,999	*	*	*	5,776	10.4	(6.1 - 14.7)
≥ \$75,000	*	*	*	8,561	4.4	(3.1 - 5.8)

<sup>1</sup> Defined as needing to see a doctor within the past year but not being able to because of cost

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 12a. Visited a doctor for routine checkup within the past year among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	637	28,200	75.2	(70.2 - 80.2)	38,026	10,809,300	72.9	(70.6 - 75.3)
<b>Gender</b>								
Male	225	12,400	69.8	(61.3 - 78.4)	14,592	4,738,400	66.9	(63.0 - 70.9)
Female	412	15,800	80.0	(74.9 - 85.2)	23,434	6,070,900	78.5	(75.9 - 81.0)
<b>Age (years)</b>								
18-34	*	*	*	*	3,921	2,747,400	61.7	(55.6 - 67.9)
35-44	*	*	*	*	4,973	1,883,100	65.5	(60.2 - 70.7)
45-54	137	4,800	67.3	(57.7 - 76.8)	7,808	2,174,100	76.6	(73.2 - 80.0)
55-64	128	4,800	83.7	(76.5 - 90.9)	8,751	1,728,600	81.4	(78.7 - 84.2)
≥ 65	227	7,700	93.3	(89.9 - 96.8)	12,573	2,276,000	90.0	(87.5 - 92.5)
<b>Race/Ethnicity</b>								
White Non-Hispanic	609	25,800	76.2	(71.7 - 80.7)	35,721	6,746,200	72.4	(70.6 - 74.2)
Black Non-Hispanic	*	*	*	*	736	1,688,900	78.8	(70.4 - 87.3)
Hispanic	*	*	*	*	857	1,550,900	69.3	(60.0 - 78.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	321	13,900	77.8	(71.3 - 84.3)	15,640	4,150,600	74.7	(70.7 - 78.7)
Some College	*	*	*	*	9,880	2,524,000	71.3	(66.1 - 76.5)
≥ College Degree	144	6,800	78.2	(70.1 - 86.2)	12,428	4,042,900	71.9	(68.5 - 75.3)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,616	2,469,200	70.4	(64.2 - 76.6)
\$25,000-\$49,999	185	8,300	76.9	(69.1 - 84.8)	9,631	2,068,400	71.3	(65.3 - 77.3)
\$50,000-\$74,999	*	*	*	*	5,765	1,466,000	74.8	(69.6 - 80.0)
≥ \$75,000	*	*	*	*	8,547	3,176,500	70.7	(67.3 - 74.2)

<sup>1</sup> Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past year

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have visited a doctor for a routine checkup within the past year (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 12b. Visited a doctor for routine checkup within the past year among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	637	73.5	(68.1 - 78.9)	38,026	72.7	(70.3 - 75.0)
<b>Gender</b>						
Male	225	68.7	(59.6 - 77.7)	14,592	67.1	(63.3 - 70.9)
Female	412	77.8	(71.9 - 83.8)	23,434	77.7	(75.1 - 80.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	609	74.4	(69.4 - 79.4)	35,721	71.9	(70.1 - 73.7)
Black Non-Hispanic	*	*	*	736	79.3	(71.3 - 87.4)
Hispanic	*	*	*	857	69.2	(60.1 - 78.2)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	321	75.9	(68.5 - 83.3)	15,640	74.4	(70.4 - 78.4)
Some College	*	*	*	9,880	70.9	(65.9 - 76.0)
≥ College Degree	144	76.8	(68.2 - 85.4)	12,428	71.7	(68.3 - 75.0)
<b>Income</b>						
≤ \$24,999	*	*	*	9,616	70.2	(64.1 - 76.2)
\$25,000-\$49,999	185	75.8	(67.1 - 84.4)	9,631	71.1	(65.3 - 76.9)
\$50,000-\$74,999	*	*	*	5,765	74.5	(69.2 - 79.8)
≥ \$75,000	*	*	*	8,547	70.3	(66.9 - 73.8)

<sup>1</sup> Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past year

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 13a. Visited a doctor for routine checkup within the past 2 years among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	637	31,500	84.1	(79.4 - 88.7)	38,026	12,726,600	85.9	(83.9 - 87.9)
<b>Gender</b>								
Male	225	14,000	78.8	(70.6 - 87.1)	14,592	5,791,900	81.8	(78.2 - 85.4)
Female	412	17,500	88.7	(84.4 - 93.1)	23,434	6,934,700	89.6	(87.9 - 91.3)
<b>Age (years)</b>								
18-34	*	*	*	*	3,921	3,535,400	79.4	(73.9 - 85.0)
35-44	*	*	*	*	4,973	2,355,600	81.9	(78.0 - 85.7)
45-54	137	5,600	78.4	(69.5 - 87.3)	7,808	2,507,900	88.4	(85.6 - 91.2)
55-64	128	5,200	91.0	(85.5 - 96.5)	8,751	1,924,800	90.7	(88.6 - 92.7)
≥ 65	227	7,800	94.4	(91.2 - 97.6)	12,573	2,402,900	95.0	(93.5 - 96.5)
<b>Race/Ethnicity</b>								
White Non-Hispanic	609	28,800	85.1	(81.2 - 89.0)	35,721	8,070,000	86.6	(85.4 - 87.9)
Black Non-Hispanic	*	*	*	*	736	1,914,500	89.4	(82.3 - 96.5)
Hispanic	*	*	*	*	857	1,832,800	81.9	(73.7 - 90.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	321	15,000	84.1	(78.2 - 90.1)	15,640	4,713,700	84.8	(81.2 - 88.4)
Some College	*	*	*	*	9,880	2,990,500	84.5	(79.6 - 89.4)
≥ College Degree	144	7,700	88.7	(82.5 - 94.9)	12,428	4,928,300	87.6	(85.3 - 89.9)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,616	2,911,700	83.0	(77.4 - 88.6)
\$25,000-\$49,999	185	9,100	84.6	(77.6 - 91.6)	9,631	2,375,500	81.9	(76.1 - 87.7)
\$50,000-\$74,999	94	5,500	86.4	(77.5 - 95.3)	5,765	1,733,500	88.4	(84.9 - 91.9)
≥ \$75,000	97	6,400	86.4	(78.2 - 94.6)	8,547	3,907,000	87.0	(84.8 - 89.2)

<sup>1</sup> Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past 2 years

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have visited a doctor for a routine checkup within the past 2 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 13b. Visited a doctor for routine checkup within the past 2 years among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	637	83.1	(78.1 - 88.1)	38,026	85.7	(83.7 - 87.6)
<b>Gender</b>						
Male	225	78.2	(69.4 - 86.9)	14,592	81.9	(78.4 - 85.3)
Female	412	87.5	(82.4 - 92.6)	23,434	89.1	(87.4 - 90.9)
<b>Race/Ethnicity</b>						
White Non-Hispanic	609	84.1	(79.8 - 88.5)	35,721	86.2	(85.0 - 87.5)
Black Non-Hispanic	*	*	*	736	89.6	(83.0 - 96.3)
Hispanic	*	*	*	857	81.8	(73.8 - 89.8)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	321	82.8	(75.9 - 89.6)	15,640	84.6	(81.1 - 88.1)
Some College	*	*	*	9,880	84.4	(79.8 - 89.0)
≥ College Degree	144	87.4	(80.5 - 94.3)	12,428	87.3	(85.0 - 89.6)
<b>Income</b>						
≤ \$24,999	*	*	*	9,616	83.2	(77.8 - 88.5)
\$25,000-\$49,999	185	84.1	(76.5 - 91.7)	9,631	81.9	(76.3 - 87.4)
\$50,000-\$74,999	94	86.0	(76.3 - 95.6)	5,765	88.1	(84.4 - 91.8)
≥ \$75,000	97	85.5	(76.6 - 94.4)	8,547	86.4	(84.2 - 88.7)

<sup>1</sup> Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past 2 years

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 14a. Dentist visit within the past year among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	638	24,900	66.4	(61.3 - 71.5)	38,026	10,540,400	71.1	(68.6 - 73.6)
<b>Gender</b>								
Male	225	11,000	62.4	(53.7 - 71.0)	14,573	4,845,800	68.4	(64.5 - 72.2)
Female	413	13,800	70.0	(64.5 - 75.5)	23,453	5,694,600	73.6	(70.4 - 76.8)
<b>Age (years)</b>								
18-34	*	*	*	*	3,924	2,789,700	62.6	(56.3 - 69.0)
35-44	*	*	*	*	4,979	2,181,000	75.8	(70.9 - 80.7)
45-54	137	5,100	72.4	(63.4 - 81.3)	7,814	2,149,300	75.8	(71.9 - 79.7)
55-64	127	4,100	71.5	(62.9 - 80.0)	8,742	1,690,500	79.2	(75.9 - 82.6)
≥ 65	229	4,700	55.9	(48.8 - 63.0)	12,567	1,729,800	68.5	(64.7 - 72.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	610	23,500	69.3	(64.8 - 73.9)	35,721	6,959,900	74.7	(72.7 - 76.6)
Black Non-Hispanic	*	*	*	*	735	1,405,800	66.2	(57.9 - 74.5)
Hispanic	*	*	*	*	862	1,425,600	63.1	(53.7 - 72.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	323	9,500	53.0	(45.9 - 60.2)	15,613	3,620,700	65.1	(60.7 - 69.5)
Some College	*	*	*	*	9,880	2,504,700	70.8	(66.1 - 75.5)
≥ College Degree	145	7,400	85.0	(78.6 - 91.3)	12,453	4,402,400	78.2	(74.6 - 81.8)
<b>Income</b>								
≤ \$24,999	193	3,200	35.5	(26.1 - 45.0)	9,594	1,972,200	56.2	(49.8 - 62.6)
\$25,000-\$49,999	185	7,200	67.6	(59.0 - 76.1)	9,649	1,896,000	65.8	(59.5 - 72.1)
\$50,000-\$74,999	95	5,300	81.8	(72.3 - 91.3)	5,763	1,482,400	75.2	(69.4 - 81.0)
≥ \$75,000	97	6,700	90.1	(83.7 - 96.4)	8,564	3,707,900	82.4	(79.5 - 85.3)

<sup>1</sup> Defined as having visited a dentist or a dental clinic for any reason within the past year

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have visited a dentist or a dental clinic for any reason within the past year (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 14b. Dentist visit within the past year among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	638	67.5	(62.1 - 73.0)	38,026	71.1	(68.6 - 73.5)
<b>Gender</b>						
Male	225	62.1	(53.0 - 71.3)	14,573	68.3	(64.6 - 72.1)
Female	413	72.4	(66.4 - 78.3)	23,453	73.6	(70.4 - 76.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	610	70.5	(65.6 - 75.5)	35,721	74.4	(72.4 - 76.3)
Black Non-Hispanic	*	*	*	735	66.6	(58.3 - 74.9)
Hispanic	*	*	*	862	63.6	(54.4 - 72.9)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	323	54.1	(46.1 - 62.2)	15,613	65.3	(60.9 - 69.6)
Some College	*	*	*	9,880	70.4	(65.7 - 75.1)
≥ College Degree	145	85.7	(79.4 - 91.9)	12,453	78.1	(74.6 - 81.7)
<b>Income</b>						
≤ \$24,999	*	*	*	9,594	56.4	(50.1 - 62.8)
\$25,000-\$49,999	185	68.2	(58.9 - 77.5)	9,649	65.8	(59.6 - 72.0)
\$50,000-\$74,999	*	*	*	5,763	75.0	(69.1 - 80.8)
≥ \$75,000	97	90.9	(84.8 - 97.0)	8,564	82.1	(79.1 - 85.0)

<sup>1</sup> Defined as having visited a dentist or a dental clinic for any reason within the past year

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 15a. Had teeth cleaned within the past year among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	566	23,400	68.3	(62.9 - 73.7)	34,070	9,985,900	71.7	(69.1 - 74.3)
<b>Gender</b>								
Male	197	10,200	62.3	(53.1 - 71.5)	13,173	4,616,400	69.0	(65.0 - 73.1)
Female	369	13,200	73.8	(68.3 - 79.3)	20,897	5,369,500	74.2	(70.8 - 77.5)
<b>Age (years)</b>								
18-34	*	*	*	*	3,860	2,746,400	63.0	(56.5 - 69.5)
35-44	*	*	*	*	4,875	2,112,100	75.5	(70.7 - 80.3)
45-54	125	4,500	71.1	(61.4 - 80.7)	7,438	2,022,600	73.3	(69.1 - 77.4)
55-64	109	3,800	77.5	(69.0 - 86.0)	7,906	1,560,900	77.7	(73.8 - 81.6)
≥ 65	189	4,400	62.4	(54.7 - 70.1)	9,991	1,543,900	77.1	(73.7 - 80.5)
<b>Race/Ethnicity</b>								
White Non-Hispanic	540	21,900	71.2	(66.4 - 76.0)	31,982	6,601,500	75.4	(73.4 - 77.4)
Black Non-Hispanic	*	*	*	*	680	1,227,200	61.6	(52.1 - 71.0)
Hispanic	*	*	*	*	789	1,421,300	66.6	(57.1 - 76.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	265	9,000	58.5	(50.7 - 66.4)	12,740	3,388,600	67.7	(63.1 - 72.3)
Some College	*	*	*	*	9,145	2,286,800	68.5	(62.9 - 74.1)
≥ College Degree	138	7,000	83.2	(76.1 - 90.4)	12,128	4,298,500	78.3	(74.8 - 81.7)
<b>Income</b>								
≤ \$24,999	*	*	*	*	7,525	1,861,800	60.3	(53.5 - 67.1)
\$25,000-\$49,999	173	7,000	69.0	(60.2 - 77.8)	8,709	1,681,700	62.6	(55.4 - 69.8)
\$50,000-\$74,999	*	*	*	*	5,534	1,447,800	76.3	(70.7 - 81.8)
≥ \$75,000	95	6,500	89.3	(82.8 - 95.8)	8,448	3,617,500	80.8	(77.8 - 83.8)

<sup>1</sup> Defined as having had teeth cleaned by a dentist or dental hygienist within the past year

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had their teeth cleaned by a dentist or dental hygienist within the past year (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 15b. Had teeth cleaned within the past year among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	566	68.8	(63.1 - 74.5)	34,070	71.7	(69.1 - 74.2)
<b>Gender</b>						
Male	197	62.0	(52.3 - 71.6)	13,173	69.2	(65.3 - 73.1)
Female	369	75.0	(69.1 - 81.0)	20,897	73.9	(70.5 - 77.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	540	71.8	(66.6 - 77.0)	31,982	75.1	(73.0 - 77.1)
Black Non-Hispanic	*	*	*	680	62.0	(52.7 - 71.4)
Hispanic	*	*	*	789	66.9	(57.5 - 76.2)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	265	58.4	(49.7 - 67.2)	12,740	67.7	(63.2 - 72.3)
Some College	*	*	*	9,145	68.1	(62.6 - 73.6)
≥ College Degree	138	83.6	(76.3 - 90.8)	12,128	78.3	(74.9 - 81.7)
<b>Income</b>						
≤ \$24,999	*	*	*	7,525	60.2	(53.4 - 67.0)
\$25,000-\$49,999	173	68.8	(59.2 - 78.4)	8,709	62.7	(55.7 - 69.7)
\$50,000-\$74,999	*	*	*	5,534	76.4	(70.9 - 81.9)
≥ \$75,000	95	90.4	(84.2 - 96.6)	8,448	80.4	(77.4 - 83.4)

<sup>1</sup> Defined as having had teeth cleaned by a dentist or dental hygienist within the past year

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 16a. Permanent teeth extracted due to decay or gum disease among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	631	19,700	52.9	(47.7 - 58.2)	37,652	7,426,500	50.4	(47.8 - 52.9)
<b>Gender</b>								
Male	224	8,800	49.8	(41.3 - 58.4)	14,457	3,367,000	47.8	(43.9 - 51.6)
Female	407	10,900	55.7	(49.6 - 61.9)	23,195	4,059,500	52.7	(49.5 - 56.0)
<b>Age (years)</b>								
18-34	*	*	*	*	3,921	1,121,600	25.4	(19.7 - 31.0)
35-44	*	*	*	*	4,964	1,250,000	43.6	(37.9 - 49.2)
45-54	137	4,200	58.5	(48.8 - 68.3)	7,780	1,595,300	56.1	(51.4 - 60.7)
55-64	126	3,800	68.1	(58.8 - 77.5)	8,648	1,382,800	65.5	(61.6 - 69.4)
≥ 65	224	6,700	82.3	(76.6 - 87.9)	12,339	2,076,900	83.0	(80.6 - 85.4)
<b>Race/Ethnicity</b>								
White Non-Hispanic	604	18,400	54.7	(49.6 - 59.7)	35,364	4,425,700	47.9	(46.0 - 49.9)
Black Non-Hispanic	*	*	*	*	727	1,244,700	58.2	(48.9 - 67.4)
Hispanic	*	*	*	*	853	1,220,500	54.2	(44.9 - 63.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	318	11,400	65.1	(57.6 - 72.6)	15,427	3,436,000	61.9	(57.2 - 66.6)
Some College	169	4,900	45.1	(35.2 - 54.9)	9,798	1,726,200	49.3	(44.3 - 54.2)
≥ College Degree	144	3,300	38.2	(28.7 - 47.7)	12,351	2,242,600	40.1	(36.5 - 43.6)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,483	2,171,900	62.3	(55.8 - 68.8)
\$25,000-\$49,999	182	5,400	50.4	(41.2 - 59.5)	9,552	1,577,400	54.7	(48.3 - 61.0)
\$50,000-\$74,999	*	*	*	*	5,744	1,000,700	51.5	(45.6 - 57.4)
≥ \$75,000	*	*	*	*	8,516	1,738,200	38.8	(35.3 - 42.2)

<sup>1</sup> Defined as having had at least one permanent tooth extracted because of tooth decay or gum disease

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had at least one permanent tooth extracted because of tooth decay or gum disease (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 16b. Permanent teeth extracted due to decay or gum disease among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	631	49.3	(43.7 - 54.8)	37,652	49.4	(46.9 - 51.9)
<b>Gender</b>						
Male	224	47.6	(38.7 - 56.5)	14,457	47.9	(44.1 - 51.7)
Female	407	50.9	(44.1 - 57.6)	23,195	50.7	(47.4 - 54.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	604	50.7	(45.3 - 56.2)	35,364	46.5	(44.5 - 48.5)
Black Non-Hispanic	*	*	*	727	57.7	(48.6 - 66.9)
Hispanic	*	*	*	853	53.8	(44.6 - 62.9)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	318	61.6	(53.1 - 70.0)	15,427	61.3	(56.7 - 66.0)
Some College	*	*	*	9,798	47.9	(43.0 - 52.9)
≥ College Degree	144	35.3	(25.7 - 44.9)	12,351	39.0	(35.5 - 42.5)
<b>Income</b>						
≤ \$24,999	*	*	*	9,483	61.6	(55.2 - 68.1)
\$25,000-\$49,999	182	45.2	(35.6 - 54.9)	9,552	54.2	(48.0 - 60.3)
\$50,000-\$74,999	*	*	*	5,744	49.9	(44.0 - 55.9)
≥ \$75,000	*	*	*	8,516	37.7	(34.2 - 41.1)

<sup>1</sup> Defined as having had at least one permanent tooth extracted because of tooth decay or gum disease

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 17a. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	224	1,400	16.7	(11.7 - 21.8)	12,339	491,200	19.6	(16.0 - 23.3)
<b>Gender</b>								
Male	75	500	16.3	(8.1 - 24.5)	4,352	167,200	16.4	(11.8 - 20.9)
Female	149	800	17.0	(10.5 - 23.4)	7,987	324,000	21.8	(16.6 - 27.0)
<b>Age (years)</b>								
≥ 65	224	1,400	16.7	(11.7 - 21.8)	12,339	491,200	19.6	(16.0 - 23.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	218	1,300	17.1	(11.8 - 22.3)	11,885	322,600	16.8	(14.3 - 19.2)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	152	1,100	20.5	(13.8 - 27.2)	6,358	314,100	25.3	(19.9 - 30.7)
Some College	*	*	*	*	2,736	90,600	20.2	(12.1 - 28.2)
≥ College Degree	*	*	*	*	3,214	80,500	10.1	(3.6 - 16.5)
<b>Income</b>								
≤ \$24,999	108	900	27.2	(17.9 - 36.6)	4,558	199,600	26.9	(19.3 - 34.4)
\$25,000-\$49,999	*	*	*	*	3,365	123,900	21.4	(13.1 - 29.6)
\$50,000-\$74,999	*	*	*	*	1,179	21,400	7.3	(3.8 - 10.8)
≥ \$75,000	*	*	*	*	1,187	9,600	2.5	(1.0 - 4.0)

<sup>1</sup> Defined as having all permanent teeth extracted because of tooth decay or gum disease among adults aged 65 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 65 and older who have had all permanent teeth extracted because of tooth decay or gum disease (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 17b. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	224	16.7	(11.5 - 21.9)	12,339	19.4	(15.7 - 23.2)
<b>Gender</b>						
Male	75	16.4	(8.2 - 24.6)	4,352	16.5	(11.5 - 21.4)
Female	149	17.0	(10.5 - 23.4)	7,987	22.1	(16.6 - 27.7)
<b>Race/Ethnicity</b>						
White Non-Hispanic	218	17.1	(11.8 - 22.4)	11,885	16.6	(13.9 - 19.3)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	152	20.4	(13.6 - 27.3)	6,358	24.9	(19.3 - 30.5)
Some College	*	*	*	2,736	21.0	(12.1 - 29.9)
≥ College Degree	*	*	*	3,214	10.0	(3.6 - 16.3)
<b>Income</b>						
≤ \$24,999	108	27.6	(17.9 - 37.3)	4,558	27.3	(19.1 - 35.5)
\$25,000-\$49,999	*	*	*	3,365	21.0	(12.9 - 29.1)
\$50,000-\$74,999	*	*	*	1,179	7.6	(3.6 - 11.5)
≥ \$75,000	*	*	*	1,187	2.2	(0.9 - 3.5)

<sup>1</sup> Defined as having all permanent teeth extracted because of tooth decay or gum disease among adults aged 65 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 18a. Flu shot within the past 12 months<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	641	14,400	38.4	(33.7 - 43.1)	38,079	6,255,900	42.3	(39.9 - 44.8)
<b>Gender</b>								
Male	226	5,900	33.7	(26.2 - 41.1)	14,579	2,798,900	39.6	(35.8 - 43.5)
Female	415	8,500	42.7	(36.9 - 48.4)	23,500	3,456,900	44.8	(41.6 - 47.9)
<b>Age (years)</b>								
18-49	211	4,200	21.2	(14.8 - 27.6)	12,505	2,685,300	31.1	(27.3 - 34.9)
50-64	198	4,300	47.0	(39.0 - 55.0)	12,972	1,662,900	46.0	(42.6 - 49.5)
≥ 65	232	5,900	70.1	(63.7 - 76.6)	12,602	1,907,700	75.2	(72.0 - 78.4)
<b>Race/Ethnicity</b>								
White Non-Hispanic	614	13,700	40.3	(35.6 - 45.0)	35,770	4,029,000	43.3	(41.4 - 45.3)
Black Non-Hispanic	*	*	*	*	738	907,000	42.1	(33.1 - 51.1)
Hispanic	*	*	*	*	860	882,900	39.6	(30.4 - 48.9)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	323	7,000	39.6	(32.9 - 46.4)	15,663	2,457,800	44.3	(39.8 - 48.8)
Some College	172	3,600	33.0	(24.4 - 41.5)	9,898	1,377,700	38.9	(34.0 - 43.9)
≥ College Degree	146	3,700	42.9	(33.0 - 52.7)	12,436	2,404,400	43.0	(39.5 - 46.4)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,635	1,656,800	47.6	(41.2 - 54.0)
\$25,000-\$49,999	184	4,700	44.4	(35.6 - 53.2)	9,647	1,200,600	41.4	(35.6 - 47.3)
\$50,000-\$74,999	*	*	*	*	5,763	792,300	40.2	(34.3 - 46.2)
≥ \$75,000	97	2,100	27.9	(18.2 - 37.6)	8,548	1,776,100	39.6	(36.3 - 42.8)

<sup>1</sup> Defined as having had a flu shot within the past 12 months

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had a flu shot within the past 12 months (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 18b. Flu shot within the past 12 months<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	641	34.7	(29.9 - 39.6)	38,079	41.7	(39.2 - 44.1)
<b>Gender</b>						
Male	226	31.6	(23.9 - 39.2)	14,579	40.0	(36.2 - 43.8)
Female	415	37.5	(31.4 - 43.6)	23,500	43.1	(40.0 - 46.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	614	36.6	(31.7 - 41.6)	35,770	42.4	(40.4 - 44.3)
Black Non-Hispanic	*	*	*	738	42.3	(33.3 - 51.3)
Hispanic	*	*	*	860	39.0	(29.9 - 48.1)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	323	35.4	(28.2 - 42.5)	15,663	43.7	(39.1 - 48.2)
Some College	172	30.1	(21.5 - 38.8)	9,898	37.8	(33.0 - 42.7)
≥ College Degree	146	39.2	(29.3 - 49.2)	12,436	42.5	(39.1 - 46.0)
<b>Income</b>						
≤ \$24,999	*	*	*	9,635	47.3	(41.0 - 53.7)
\$25,000-\$49,999	184	39.4	(30.2 - 48.6)	9,647	40.3	(34.5 - 46.0)
\$50,000-\$74,999	*	*	*	5,763	39.4	(33.4 - 45.4)
≥ \$75,000	97	25.5	(16.1 - 34.9)	8,548	39.1	(35.8 - 42.4)

<sup>1</sup> Defined as having had a flu shot within the past 12 months

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 19a. Flu vaccine that was sprayed in nose within the past 12 months<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	*	*	*	*	38,106	248,300	1.7	(0.6 - 2.8)
<b>Gender</b>								
Male	*	*	*	*	14,586	148,000	2.1	(0.0 - 4.2)
Female	*	*	*	*	23,520	100,300	1.3	(0.5 - 2.1)
<b>Age (years)</b>								
18-49	*	*	*	*	12,535	184,900	2.1	(0.3 - 3.9)
50-64	*	*	*	*	12,968	36,000	1.0	(0.2 - 1.8)
≥ 65	*	*	*	*	12,603	27,400	1.1	(0.4 - 1.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	*	*	*	*	35,796	66,000	0.7	(0.4 - 1.0)
Black Non-Hispanic	*	*	*	*	740	38,500	1.8	(0.0 - 4.1)
Hispanic	*	*	*	*	860	133,700	5.9	(0.0 - 12.3)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	15,679	172,500	3.1	(0.3 - 5.9)
Some College	*	*	*	*	9,897	43,500	1.2	(0.4 - 2.0)
≥ College Degree	*	*	*	*	12,449	31,600	0.6	(0.3 - 0.8)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,636	138,400	3.9	(0.0 - 8.1)
\$25,000-\$49,999	*	*	*	*	9,648	41,500	1.4	(0.0 - 3.1)
\$50,000-\$74,999	*	*	*	*	5,766	13,300	0.7	(0.2 - 1.2)
≥ \$75,000	*	*	*	*	8,561	35,300	0.8	(0.3 - 1.3)

<sup>1</sup> Defined as having had flu vaccine sprayed in nose within the past 12 months

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had flu vaccine sprayed in nose within the past 12 months (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 19b. Flu vaccine that was sprayed in nose within the past 12 months<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	*	*	*	38,106	1.7	(0.6 - 2.7)
<b>Gender</b>						
Male	*	*	*	14,586	2.0	(0.1 - 3.9)
Female	*	*	*	23,520	1.4	(0.5 - 2.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	*	*	*	35,796	0.7	(0.4 - 1.0)
Black Non-Hispanic	*	*	*	740	1.9	(0.0 - 4.4)
Hispanic	*	*	*	860	5.9	(0.0 - 11.9)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	15,679	3.1	(0.4 - 5.8)
Some College	*	*	*	9,897	1.2	(0.4 - 2.0)
≥ College Degree	*	*	*	12,449	0.6	(0.3 - 0.8)
<b>Income</b>						
≤ \$24,999	*	*	*	9,636	3.8	(0.0 - 7.7)
\$25,000-\$49,999	*	*	*	9,648	1.5	(0.0 - 3.4)
\$50,000-\$74,999	*	*	*	5,766	0.7	(0.1 - 1.2)
≥ \$75,000	*	*	*	8,561	0.8	(0.3 - 1.3)

<sup>1</sup> Defined as having had flu vaccine sprayed in nose within the past 12 months

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 20a. Flu shot or flu vaccine sprayed in nose within the past 12 months<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	641	14,800	39.5	(34.8 - 44.3)	38,059	6,302,300	42.7	(40.2 - 45.1)
<b>Gender</b>								
Male	226	5,900	33.7	(26.2 - 41.1)	14,568	2,822,700	40.0	(36.1 - 43.9)
Female	415	8,900	44.8	(38.9 - 50.6)	23,491	3,479,500	45.1	(42.0 - 48.2)
<b>Age (years)</b>								
18-49	211	4,700	23.3	(16.5 - 30.1)	12,501	2,726,400	31.6	(27.8 - 35.4)
50-64	198	4,300	47.0	(39.0 - 55.0)	12,964	1,664,200	46.1	(42.7 - 49.5)
≥ 65	232	5,900	70.1	(63.7 - 76.6)	12,594	1,911,700	75.4	(72.2 - 78.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	614	13,900	41.0	(36.3 - 45.8)	35,750	4,048,700	43.6	(41.7 - 45.5)
Black Non-Hispanic	*	*	*	*	738	910,000	42.2	(33.2 - 51.2)
Hispanic	*	*	*	*	860	903,300	40.6	(31.3 - 49.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	323	7,400	42.0	(35.1 - 48.9)	15,651	2,480,100	44.7	(40.2 - 49.3)
Some College	172	3,600	33.0	(24.4 - 41.5)	9,895	1,387,300	39.2	(34.3 - 44.1)
≥ College Degree	146	3,700	42.9	(33.0 - 52.7)	12,431	2,418,300	43.2	(39.8 - 46.7)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,626	1,675,100	48.2	(41.8 - 54.6)
\$25,000-\$49,999	184	5,100	48.3	(39.2 - 57.4)	9,643	1,208,300	41.7	(35.8 - 47.6)
\$50,000-\$74,999	*	*	*	*	5,761	795,600	40.4	(34.5 - 46.3)
≥ \$75,000	97	2,100	27.9	(18.2 - 37.6)	8,548	1,791,200	39.9	(36.6 - 43.2)

<sup>1</sup> Defined as having had a flu shot or flu vaccine sprayed in nose within the past 12 months

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had a flu shot or flu vaccine sprayed in nose within the past 12 months (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 20b. Flu shot or flu vaccine sprayed in nose within the past 12 months<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	641	36.0	(31.0 - 41.0)	38,059	42.0	(39.6 - 44.5)
<b>Gender</b>						
Male	226	31.6	(23.9 - 39.2)	14,568	40.4	(36.6 - 44.2)
Female	415	39.9	(33.6 - 46.3)	23,491	43.5	(40.3 - 46.7)
<b>Race/Ethnicity</b>						
White Non-Hispanic	614	37.5	(32.5 - 42.6)	35,750	42.6	(40.7 - 44.6)
Black Non-Hispanic	*	*	*	738	42.4	(33.4 - 51.4)
Hispanic	*	*	*	860	40.0	(30.9 - 49.1)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	323	38.2	(30.6 - 45.7)	15,651	44.1	(39.6 - 48.6)
Some College	172	30.1	(21.5 - 38.8)	9,895	38.1	(33.2 - 43.0)
≥ College Degree	146	39.2	(29.3 - 49.2)	12,431	42.8	(39.4 - 46.3)
<b>Income</b>						
≤ \$24,999	*	*	*	9,626	48.0	(41.6 - 54.3)
\$25,000-\$49,999	184	44.0	(34.3 - 53.7)	9,643	40.5	(34.8 - 46.3)
\$50,000-\$74,999	*	*	*	5,761	39.6	(33.6 - 45.6)
≥ \$75,000	97	25.5	(16.1 - 34.9)	8,548	39.4	(36.1 - 42.7)

<sup>1</sup> Defined as having had a flu shot or flu vaccine sprayed in nose within the past 12 months

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 21a. Flu shot among adults aged 65 and older<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	232	5,900	70.1	(63.7 - 76.6)	12,602	1,907,700	75.2	(72.0 - 78.4)
<b>Gender</b>								
Male	*	*	*	*	4,417	772,900	74.6	(69.3 - 79.8)
Female	155	3,600	69.7	(61.9 - 77.5)	8,185	1,134,800	75.6	(71.6 - 79.6)
<b>Age (years)</b>								
≥ 65	232	5,900	70.1	(63.7 - 76.6)	12,602	1,907,700	75.2	(72.0 - 78.4)
<b>Race/Ethnicity</b>								
White Non-Hispanic	226	5,600	69.4	(62.8 - 76.0)	12,137	1,487,100	76.0	(73.5 - 78.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	157	3,800	68.3	(60.2 - 76.3)	6,518	930,800	74.0	(68.9 - 79.2)
Some College	*	*	*	*	2,790	317,800	69.7	(62.7 - 76.7)
≥ College Degree	*	*	*	*	3,261	654,000	80.5	(76.4 - 84.6)
<b>Income</b>								
≤ \$24,999	110	2,200	64.3	(54.6 - 74.1)	4,665	545,300	72.4	(65.6 - 79.3)
\$25,000-\$49,999	*	*	*	*	3,422	430,600	72.8	(67.0 - 78.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	1,198	301,000	77.5	(70.9 - 84.2)

<sup>1</sup> Defined as having had a flu shot during the past 12 months among adults aged 65 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 65 and older who have had a flu shot within the past 12 months (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 21b. Flu shot among adults aged 65 and older<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	232	70.1	(63.4 - 76.8)	12,602	75.0	(71.5 - 78.5)
<b>Gender</b>						
Male	*	*	*	4,417	74.4	(68.7 - 80.1)
Female	155	69.7	(61.9 - 77.4)	8,185	75.5	(71.3 - 79.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	226	69.4	(62.5 - 76.3)	12,137	75.9	(73.0 - 78.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	157	67.7	(59.1 - 76.2)	6,518	73.3	(67.5 - 79.1)
Some College	*	*	*	2,790	70.0	(62.7 - 77.2)
≥ College Degree	*	*	*	3,261	80.7	(76.2 - 85.1)
<b>Income</b>						
≤ \$24,999	110	65.5	(55.7 - 75.4)	4,665	72.6	(65.1 - 80.1)
\$25,000-\$49,999	66	80.8	(70.9 - 90.7)	3,422	71.4	(65.0 - 77.9)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	1,198	78.0	(70.8 - 85.1)

<sup>1</sup> Defined as having had a flu shot during the past 12 months among adults aged 65 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 22a. Pneumonia shot or pneumococcal vaccine<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	602	10,100	29.8	(25.4 - 34.2)	35,303	3,456,800	26.1	(23.9 - 28.3)
<b>Gender</b>								
Male	199	4,100	27.7	(20.4 - 35.1)	13,052	1,542,200	25.2	(21.5 - 28.8)
Female	403	6,000	31.4	(26.1 - 36.6)	22,251	1,914,600	26.9	(24.2 - 29.6)
<b>Age (years)</b>								
18-49	191	2,800	16.3	(9.9 - 22.7)	10,943	1,097,000	14.6	(11.3 - 17.9)
50-64	185	1,700	20.6	(14.4 - 26.7)	12,179	758,000	23.0	(19.8 - 26.2)
≥ 65	226	5,500	68.4	(61.7 - 75.1)	12,181	1,601,700	65.7	(61.7 - 69.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	578	9,400	30.3	(25.9 - 34.7)	33,225	2,358,700	28.7	(26.9 - 30.5)
Black Non-Hispanic	*	*	*	*	685	584,400	28.4	(19.5 - 37.3)
Hispanic	*	*	*	*	769	408,000	19.5	(12.6 - 26.4)
Other Non-Hispanic	*	*	*	*	624	105,600	12.2	(2.3 - 22.2)
<b>Education</b>								
≤ High School	308	5,400	33.5	(27.2 - 39.7)	14,700	1,535,300	30.4	(26.4 - 34.3)
Some College	160	2,500	24.5	(16.1 - 32.9)	9,151	836,200	27.2	(21.9 - 32.5)
≥ College Degree	134	2,200	29.0	(20.2 - 37.9)	11,375	1,070,700	21.4	(18.6 - 24.1)
<b>Income</b>								
≤ \$24,999	188	3,200	36.1	(26.5 - 45.7)	9,082	1,050,900	33.1	(27.6 - 38.6)
\$25,000-\$49,999	173	3,100	32.0	(23.8 - 40.2)	9,033	797,500	30.3	(24.1 - 36.6)
\$50,000-\$74,999	*	*	*	*	5,320	378,800	21.8	(17.4 - 26.1)
≥ \$75,000	89	900	14.0	(5.8 - 22.2)	7,773	648,700	16.3	(14.1 - 18.5)

<sup>1</sup> Defined as having ever had a pneumonia shot or pneumococcal vaccine

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have ever had a pneumonia shot or pneumococcal vaccine (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 22b. Pneumonia shot or pneumococcal vaccine<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	602	26.5	(22.0 - 31.0)	35,303	25.4	(23.2 - 27.6)
<b>Gender</b>						
Male	199	26.8	(19.1 - 34.5)	13,052	25.8	(22.3 - 29.4)
Female	403	26.2	(20.8 - 31.6)	22,251	25.1	(22.4 - 27.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	578	27.0	(22.4 - 31.6)	33,225	27.7	(25.9 - 29.5)
Black Non-Hispanic	*	*	*	685	27.9	(19.1 - 36.7)
Hispanic	*	*	*	769	19.6	(12.8 - 26.5)
Other Non-Hispanic	*	*	*	624	11.3	(2.0 - 20.7)
<b>Education</b>						
≤ High School	308	29.3	(22.7 - 35.8)	14,700	29.5	(25.6 - 33.5)
Some College	160	23.2	(14.5 - 32.0)	9,151	26.2	(21.0 - 31.3)
≥ College Degree	134	25.3	(16.5 - 34.2)	11,375	21.0	(18.3 - 23.8)
<b>Income</b>						
≤ \$24,999	*	*	*	9,082	32.3	(26.9 - 37.8)
\$25,000-\$49,999	173	27.5	(19.1 - 35.9)	9,033	29.0	(23.0 - 35.0)
\$50,000-\$74,999	*	*	*	5,320	21.2	(16.8 - 25.7)
≥ \$75,000	89	13.2	(4.9 - 21.4)	7,773	16.2	(13.9 - 18.5)

<sup>1</sup> Defined as having ever had a pneumonia shot or pneumococcal vaccine

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 23a. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	226	5,500	68.4	(61.7 - 75.1)	12,181	1,601,700	65.7	(61.7 - 69.7)
<b>Gender</b>								
Male	*	*	*	*	4,180	606,400	62.1	(55.5 - 68.7)
Female	155	3,500	67.9	(59.8 - 76.0)	8,001	995,300	68.0	(63.1 - 73.0)
<b>Age (years)</b>								
≥ 65	226	5,500	68.4	(61.7 - 75.1)	12,181	1,601,700	65.7	(61.7 - 69.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	221	5,200	67.6	(60.8 - 74.4)	11,735	1,281,300	68.2	(64.9 - 71.5)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	154	3,700	68.5	(60.6 - 76.5)	6,345	806,200	65.6	(59.5 - 71.7)
Some College	*	*	*	*	2,694	289,500	66.8	(60.1 - 73.4)
≥ College Degree	*	*	*	*	3,111	501,100	65.6	(58.5 - 72.7)
<b>Income</b>								
≤ \$24,999	109	2,100	63.0	(53.0 - 73.0)	4,533	497,100	67.7	(59.9 - 75.5)
\$25,000-\$49,999	*	*	*	*	3,324	344,300	61.5	(53.3 - 69.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	1,138	236,300	64.5	(55.2 - 73.8)

<sup>1</sup> Defined as having ever had a pneumonia shot or pneumococcal vaccine among adults aged 65 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 65 and older who have ever had a pneumonia shot or pneumococcal vaccine (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 23b. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	226	68.4	(61.6 - 75.3)	12,181	64.7	(60.4 - 69.1)
<b>Gender</b>						
Male	*	*	*	4,180	61.5	(54.3 - 68.6)
Female	155	67.9	(59.8 - 76.0)	8,001	67.6	(62.4 - 72.9)
<b>Race/Ethnicity</b>						
White Non-Hispanic	221	67.7	(60.6 - 74.7)	11,735	67.1	(63.2 - 70.9)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	154	68.1	(59.8 - 76.4)	6,345	64.2	(57.4 - 71.0)
Some College	*	*	*	2,694	65.8	(58.4 - 73.3)
≥ College Degree	*	*	*	3,111	65.3	(57.7 - 72.9)
<b>Income</b>						
≤ \$24,999	*	*	*	4,533	66.0	(57.0 - 74.9)
\$25,000-\$49,999	*	*	*	3,324	60.3	(51.9 - 68.6)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	*	*

<sup>1</sup> Defined as having ever had a pneumonia shot or pneumococcal vaccine among adults aged 65 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 24a. Ever had a mammogram among women aged 40 and older<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	341	11,500	88.9	(84.9 - 93.0)	19,491	4,328,800	91.8	(89.7 - 94.0)
<b>Gender</b>								
Male	-	-	-	-	-	-	-	-
Female	341	11,500	88.9	(84.9 - 93.0)	19,491	4,328,800	91.8	(89.7 - 94.0)
<b>Age (years)</b>								
40-49	*	*	*	*	3,851	1,287,500	85.0	(79.7 - 90.3)
50-64	120	4,400	95.9	(92.4 - 99.4)	7,452	1,641,700	96.0	(94.1 - 97.9)
≥ 65	155	4,600	89.1	(84.2 - 93.9)	8,188	1,399,600	94.1	(90.5 - 97.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	330	10,900	88.4	(84.2 - 92.6)	18,548	2,976,100	93.9	(92.7 - 95.1)
Black Non-Hispanic	*	*	*	*	342	746,600	93.8	(87.3 - 100)
Hispanic	*	*	*	*	333	447,600	91.8	(84.6 - 99.0)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	180	5,700	87.9	(82.7 - 93.1)	8,266	1,698,700	91.0	(87.4 - 94.5)
Some College	92	3,300	88.7	(79.5 - 98.0)	5,322	1,055,500	93.6	(90.2 - 97.0)
≥ College Degree	69	2,500	91.7	(84.9 - 98.6)	5,872	1,560,700	91.6	(87.5 - 95.6)
<b>Income</b>								
≤ \$24,999	125	3,400	85.5	(78.5 - 92.6)	5,691	1,023,400	93.5	(89.7 - 97.4)
\$25,000-\$49,999	92	3,300	92.1	(86.2 - 97.9)	4,857	860,300	89.5	(83.6 - 95.5)
\$50,000-\$74,999	*	*	*	*	2,648	547,100	94.6	(92.2 - 96.9)
≥ \$75,000	*	*	*	*	3,582	1,185,100	92.2	(88.2 - 96.1)

<sup>1</sup> Defined as having ever had a mammogram among women aged 40 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of women aged 40 and older who have ever had a mammogram (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 24b. Ever had a mammogram among women aged 40 and older<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	341	87.0	(81.6 - 92.5)	19,491	91.1	(88.7 - 93.6)
<b>Gender</b>						
Male	-	-	-	-	-	-
Female	341	87.0	(81.6 - 92.5)	19,491	91.1	(88.7 - 93.6)
<b>Race/Ethnicity</b>						
White Non-Hispanic	330	86.5	(80.8 - 92.1)	18,548	93.3	(92.0 - 94.7)
Black Non-Hispanic	*	*	*	342	93.1	(85.7 - 100)
Hispanic	*	*	*	333	91.0	(82.8 - 99.2)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	180	86.6	(79.9 - 93.4)	8,266	90.2	(86.1 - 94.2)
Some College	*	*	*	5,322	92.9	(88.9 - 96.9)
≥ College Degree	69	91.4	(83.5 - 99.3)	5,872	90.9	(86.6 - 95.3)
<b>Income</b>						
≤ \$24,999	125	85.4	(76.7 - 94.1)	5,691	93.0	(88.5 - 97.4)
\$25,000-\$49,999	92	89.6	(81.3 - 98.0)	4,857	88.8	(82.5 - 95.1)
\$50,000-\$74,999	*	*	*	2,648	94.3	(91.7 - 96.8)
≥ \$75,000	*	*	*	3,582	91.2	(86.8 - 95.6)

<sup>1</sup> Defined as having ever had a mammogram among women aged 40 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 25a. Had mammogram within the past 2 years among women aged 40 and older<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	338	9,500	74.3	(69.0 - 79.6)	19,376	3,760,000	80.3	(77.8 - 82.9)
<b>Gender</b>								
Male	-	-	-	-	-	-	-	-
Female	338	9,500	74.3	(69.0 - 79.6)	19,376	3,760,000	80.3	(77.8 - 82.9)
<b>Age (years)</b>								
40-49	*	*	*	*	3,842	1,122,700	74.4	(68.5 - 80.2)
50-64	120	3,600	77.8	(69.7 - 85.9)	7,428	1,487,900	87.2	(84.5 - 89.8)
≥ 65	152	3,700	73.3	(65.7 - 80.9)	8,106	1,149,400	78.5	(74.2 - 82.9)
<b>Race/Ethnicity</b>								
White Non-Hispanic	327	9,000	73.8	(68.3 - 79.2)	18,438	2,531,500	80.2	(78.3 - 82.2)
Black Non-Hispanic	*	*	*	*	339	667,100	85.6	(77.5 - 93.8)
Hispanic	*	*	*	*	332	419,700	86.1	(77.5 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	177	4,700	72.9	(65.5 - 80.3)	8,204	1,486,900	80.5	(76.4 - 84.6)
Some College	*	*	*	*	5,297	929,400	83.0	(78.8 - 87.1)
≥ College Degree	*	*	*	*	5,845	1,334,800	78.6	(73.9 - 83.2)
<b>Income</b>								
≤ \$24,999	122	2,500	65.1	(55.4 - 74.9)	5,650	882,300	80.8	(75.9 - 85.7)
\$25,000-\$49,999	92	2,900	81.8	(73.3 - 90.3)	4,846	732,300	76.3	(69.7 - 82.9)
\$50,000-\$74,999	*	*	*	*	2,641	494,200	85.6	(81.9 - 89.4)
≥ \$75,000	*	*	*	*	3,572	1,025,900	80.0	(75.2 - 84.8)

<sup>1</sup> Defined as having had a mammogram within the past 2 years among women aged 40 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of women aged 40 and older who have had a mammogram within the past 2 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 25b. Had mammogram within the past 2 years among women aged 40 and older<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	338	73.0	(66.7 - 79.2)	19,376	79.7	(76.9 - 82.5)
<b>Gender</b>						
Male	-	-	-	-	-	-
Female	338	73.0	(66.7 - 79.2)	19,376	79.7	(76.9 - 82.5)
<b>Race/Ethnicity</b>						
White Non-Hispanic	327	72.4	(66.0 - 78.8)	18,438	79.7	(77.6 - 81.8)
Black Non-Hispanic	*	*	*	339	85.3	(76.5 - 94.2)
Hispanic	*	*	*	332	85.2	(75.6 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	177	71.0	(62.3 - 79.7)	8,204	80.1	(75.6 - 84.7)
Some College	*	*	*	5,297	82.3	(77.7 - 86.9)
≥ College Degree	*	*	*	5,845	77.8	(72.9 - 82.8)
<b>Income</b>						
≤ \$24,999	*	*	*	5,650	80.8	(75.4 - 86.2)
\$25,000-\$49,999	92	81.1	(71.4 - 90.7)	4,846	75.7	(68.7 - 82.6)
\$50,000-\$74,999	*	*	*	2,641	85.5	(81.6 - 89.4)
≥ \$75,000	*	*	*	3,572	79.0	(73.8 - 84.2)

<sup>1</sup> Defined as having had a mammogram within the past 2 years among women aged 40 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 26a. Had mammogram within the past 2 years among women aged 50 and older<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	272	7,300	75.4	(69.9 - 81.0)	15,534	2,637,300	83.2	(80.7 - 85.7)
<b>Gender</b>								
Male	-	-	-	-	-	-	-	-
Female	272	7,300	75.4	(69.9 - 81.0)	15,534	2,637,300	83.2	(80.7 - 85.7)
<b>Age (years)</b>								
50-64	120	3,600	77.8	(69.7 - 85.9)	7,428	1,487,900	87.2	(84.5 - 89.8)
≥ 65	152	3,700	73.3	(65.7 - 80.9)	8,106	1,149,400	78.5	(74.2 - 82.9)
<b>Race/Ethnicity</b>								
White Non-Hispanic	262	6,800	74.9	(69.2 - 80.6)	14,887	1,907,000	82.3	(80.3 - 84.4)
Black Non-Hispanic	*	*	*	*	249	385,600	89.9	(83.2 - 96.7)
Hispanic	*	*	*	*	220	286,700	93.9	(89.1 - 98.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	162	4,200	74.5	(67.2 - 81.8)	7,026	1,094,500	83.3	(80.0 - 86.6)
Some College	*	*	*	*	4,124	646,700	87.3	(84.2 - 90.3)
≥ College Degree	*	*	*	*	4,356	890,500	80.6	(75.1 - 86.1)
<b>Income</b>								
≤ \$24,999	110	2,200	66.7	(57.0 - 76.4)	4,991	649,100	81.0	(76.8 - 85.2)
\$25,000-\$49,999	74	2,300	82.3	(73.0 - 91.5)	3,913	533,300	78.6	(70.9 - 86.2)
\$50,000-\$74,999	*	*	*	*	1,943	359,300	89.6	(86.3 - 93.0)
≥ \$75,000	*	*	*	*	2,303	601,300	84.6	(79.3 - 89.9)

<sup>1</sup> Defined as having had a mammogram within the past 2 years among women aged 50 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of women aged 50 and older who have had a mammogram within the past 2 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 26b. Had mammogram within the past 2 years among women aged 50 and older<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	272	75.7	(70.0 - 81.3)	15,534	83.1	(80.3 - 85.8)
<b>Gender</b>						
Male	-	-	-	-	-	-
Female	272	75.7	(70.0 - 81.3)	15,534	83.1	(80.3 - 85.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	262	75.1	(69.3 - 80.9)	14,887	82.1	(79.9 - 84.3)
Black Non-Hispanic	*	*	*	249	90.4	(83.6 - 97.2)
Hispanic	*	*	*	220	94.1	(89.3 - 99.0)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	162	74.4	(66.7 - 82.0)	7,026	83.6	(80.1 - 87.0)
Some College	*	*	*	4,124	87.3	(84.1 - 90.5)
≥ College Degree	*	*	*	4,356	79.9	(73.9 - 86.0)
<b>Income</b>						
≤ \$24,999	110	66.9	(57.0 - 76.8)	4,991	81.3	(76.9 - 85.6)
\$25,000-\$49,999	74	83.1	(74.1 - 92.1)	3,913	78.1	(69.9 - 86.4)
\$50,000-\$74,999	*	*	*	1,943	90.2	(87.1 - 93.4)
≥ \$75,000	*	*	*	2,303	83.9	(77.9 - 89.9)

<sup>1</sup> Defined as having had a mammogram within the past 2 years among women aged 50 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 27a. Ever had a Pap test among women<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	412	18,600	94.5	(90.7 - 98.2)	23,381	7,045,300	92.1	(90.0 - 94.3)
<b>Gender</b>								
Male	-	-	-	-	-	-	-	-
Female	412	18,600	94.5	(90.7 - 98.2)	23,381	7,045,300	92.1	(90.0 - 94.3)
<b>Age (years)</b>								
18-34	*	*	*	*	2,377	1,896,600	85.2	(79.3 - 91.0)
35-44	*	*	*	*	3,023	1,301,500	92.6	(87.7 - 97.5)
45-54	93	3,600	100	(100 - 100)	4,657	1,420,300	98.3	(97.0 - 99.6)
55-64	75	2,800	98.6	(96.6 - 100)	5,187	1,071,200	97.2	(95.0 - 99.3)
≥ 65	153	4,800	95.5	(92.0 - 98.9)	8,137	1,355,700	92.3	(88.5 - 96.1)
<b>Race/Ethnicity</b>								
White Non-Hispanic	396	17,300	94.1	(90.1 - 98.1)	21,976	4,439,700	94.2	(92.1 - 96.3)
Black Non-Hispanic	*	*	*	*	481	1,231,600	91.2	(84.0 - 98.4)
Hispanic	*	*	*	*	533	984,500	90.1	(85.3 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	208	9,100	95.1	(89.8 - 100)	9,526	2,540,500	88.3	(83.8 - 92.7)
Some College	110	4,900	91.4	(81.9 - 100)	6,462	1,782,200	93.5	(90.4 - 96.6)
≥ College Degree	94	4,600	96.8	(93.6 - 100)	7,345	2,643,200	95.0	(92.2 - 97.9)
<b>Income</b>								
≤ \$24,999	138	4,800	88.2	(78.5 - 97.9)	6,526	1,717,200	93.7	(90.9 - 96.6)
\$25,000-\$49,999	117	5,700	99.6	(99.0 - 100)	5,787	1,378,400	91.0	(85.6 - 96.4)
\$50,000-\$74,999	*	*	*	*	3,310	933,900	96.0	(91.8 - 100)
≥ \$75,000	55	3,400	100	(100 - 100)	4,570	1,943,300	95.4	(92.6 - 98.2)

<sup>1</sup> Defined as having ever had a Pap test among women

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of women who have ever had a Pap test (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 27b. Ever had a Pap test among women<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	412	94.0	(89.5 - 98.4)	23,381	91.8	(89.6 - 94.0)
<b>Gender</b>						
Male	-	-	-	-	-	-
Female	412	94.0	(89.5 - 98.4)	23,381	91.8	(89.6 - 94.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	396	93.5	(88.8 - 98.3)	21,976	94.0	(91.8 - 96.2)
Black Non-Hispanic	*	*	*	481	90.8	(83.5 - 98.1)
Hispanic	*	*	*	533	89.7	(84.6 - 94.7)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	208	94.4	(87.8 - 100)	9,526	87.8	(83.2 - 92.4)
Some College	*	*	*	6,462	93.1	(89.7 - 96.4)
≥ College Degree	94	97.5	(94.9 - 100)	7,345	94.9	(92.0 - 97.7)
<b>Income</b>						
≤ \$24,999	*	*	*	6,526	93.3	(90.1 - 96.5)
\$25,000-\$49,999	117	99.7	(99.1 - 100)	5,787	91.1	(85.7 - 96.5)
\$50,000-\$74,999	*	*	*	3,310	96.0	(91.9 - 100)
≥ \$75,000	55	100	(100 - 100)	4,570	95.0	(91.9 - 98.1)

<sup>1</sup> Defined as having ever had a Pap test among women

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 28a. Had a Pap test within the past 3 years among women<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	408	14,700	75.0	(69.9 - 80.0)	23,109	6,205,600	82.2	(79.8 - 84.6)
<b>Gender</b>								
Male	-	-	-	-	-	-	-	-
Female	408	14,700	75.0	(69.9 - 80.0)	23,109	6,205,600	82.2	(79.8 - 84.6)
<b>Age (years)</b>								
18-34	*	*	*	*	2,371	1,818,400	81.8	(75.5 - 88.1)
35-44	*	*	*	*	3,012	1,220,300	86.9	(81.8 - 92.0)
45-54	93	3,100	85.4	(77.7 - 93.1)	4,640	1,311,500	92.0	(89.8 - 94.3)
55-64	*	*	*	*	5,152	924,300	84.5	(81.0 - 88.0)
≥ 65	150	2,600	53.0	(44.2 - 61.7)	7,934	931,100	66.4	(61.8 - 71.0)
<b>Race/Ethnicity</b>								
White Non-Hispanic	392	13,500	74.1	(68.8 - 79.3)	21,720	3,807,100	81.5	(79.2 - 83.8)
Black Non-Hispanic	*	*	*	*	473	1,142,800	86.7	(79.1 - 94.3)
Hispanic	*	*	*	*	527	894,900	83.6	(76.3 - 91.0)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	204	6,400	67.4	(59.5 - 75.3)	9,384	2,132,600	76.0	(71.1 - 81.0)
Some College	*	*	*	*	6,401	1,594,300	84.0	(80.4 - 87.5)
≥ College Degree	94	4,100	86.7	(80.4 - 93.0)	7,281	2,406,000	86.9	(83.7 - 90.1)
<b>Income</b>								
≤ \$24,999	*	*	*	*	6,441	1,453,900	82.0	(77.9 - 86.0)
\$25,000-\$49,999	116	4,800	84.3	(77.8 - 90.8)	5,740	1,182,000	78.3	(72.6 - 83.9)
\$50,000-\$74,999	*	*	*	*	3,296	825,000	85.0	(78.0 - 91.9)
≥ \$75,000	55	3,100	93.1	(86.1 - 100)	4,555	1,810,200	89.0	(85.7 - 92.2)

<sup>1</sup> Defined as having had a Pap test within the past 3 years among women

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of women who have had a Pap test within the past 3 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 28b. Had a Pap test within the past 3 years among women<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	408	78.0	(72.6 - 83.4)	23,109	82.7	(80.2 - 85.1)
<b>Gender</b>						
Male	-	-	-	-	-	-
Female	408	78.0	(72.6 - 83.4)	23,109	82.7	(80.2 - 85.1)
<b>Race/Ethnicity</b>						
White Non-Hispanic	392	77.5	(71.9 - 83.1)	21,720	82.3	(80.0 - 84.6)
Black Non-Hispanic	*	*	*	473	87.0	(79.3 - 94.6)
Hispanic	*	*	*	527	83.4	(75.9 - 90.9)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	204	71.2	(62.5 - 79.9)	9,384	76.4	(71.3 - 81.5)
Some College	*	*	*	6,401	84.2	(80.5 - 88.0)
≥ College Degree	94	89.2	(83.7 - 94.6)	7,281	87.5	(84.3 - 90.6)
<b>Income</b>						
≤ \$24,999	*	*	*	6,441	82.7	(78.5 - 86.9)
\$25,000-\$49,999	116	88.0	(82.7 - 93.4)	5,740	79.2	(73.5 - 84.8)
\$50,000-\$74,999	*	*	*	3,296	85.7	(78.7 - 92.8)
≥ \$75,000	55	93.6	(86.4 - 100)	4,555	88.9	(85.5 - 92.3)

<sup>1</sup> Defined as having had a Pap test within the past 3 years among women

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 29a. Ever had a digital rectal exam among men aged 40 and older<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	184	9,100	80.8	(73.9 - 87.7)	12,033	3,176,500	73.8	(69.7 - 77.8)
<b>Gender</b>								
Male	184	9,100	80.8	(73.9 - 87.7)	12,033	3,176,500	73.8	(69.7 - 77.8)
Female	-	-	-	-	-	-	-	-
<b>Age (years)</b>								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	3,098	922,800	66.8	(59.9 - 73.8)
55-64	53	2,500	87.8	(78.4 - 97.2)	3,504	865,600	87.1	(82.5 - 91.7)
≥ 65	73	2,800	88.9	(81.1 - 96.8)	4,348	953,900	92.9	(91.0 - 94.9)
<b>Race/Ethnicity</b>								
White Non-Hispanic	177	8,400	80.6	(73.4 - 87.8)	11,411	2,452,100	80.3	(77.7 - 82.9)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	98	4,300	77.2	(67.3 - 87.1)	5,013	931,600	64.1	(57.3 - 71.0)
Some College	*	*	*	*	2,695	674,200	75.5	(67.8 - 83.2)
≥ College Degree	*	*	*	*	4,302	1,568,000	80.5	(74.0 - 87.0)
<b>Income</b>								
≤ \$24,999	*	*	*	*	2,573	573,300	65.9	(56.0 - 75.9)
\$25,000-\$49,999	*	*	*	*	3,168	530,100	74.6	(67.2 - 82.1)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,309	1,351,100	78.0	(72.6 - 83.5)

<sup>1</sup> Defined as having ever had a digital rectal exam among men aged 40 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of men aged 40 and older who have ever had a digital rectal exam (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 29b. Ever had a digital rectal exam among men aged 40 and older<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	184	79.4	(71.5 - 87.2)	12,033	73.7	(69.6 - 77.8)
<b>Gender</b>						
Male	184	79.4	(71.5 - 87.2)	12,033	73.7	(69.6 - 77.8)
Female	-	-	-	-	-	-
<b>Race/Ethnicity</b>						
White Non-Hispanic	177	78.8	(70.5 - 87.0)	11,411	79.9	(77.2 - 82.5)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	5,013	64.7	(57.8 - 71.7)
Some College	*	*	*	2,695	75.0	(67.1 - 82.9)
≥ College Degree	*	*	*	4,302	80.3	(73.8 - 86.8)
<b>Income</b>						
≤ \$24,999	*	*	*	2,573	66.9	(56.9 - 76.9)
\$25,000-\$49,999	*	*	*	3,168	74.8	(67.3 - 82.2)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,309	77.5	(72.0 - 83.0)

<sup>1</sup> Defined as having ever had a digital rectal exam among men aged 40 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 30a. Had digital rectal exam within the past 2 years among men aged 40 and older<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	182	7,000	62.7	(54.3 - 71.1)	11,923	2,380,300	55.8	(51.8 - 59.8)
<b>Gender</b>								
Male	182	7,000	62.7	(54.3 - 71.1)	11,923	2,380,300	55.8	(51.8 - 59.8)
Female	-	-	-	-	-	-	-	-
<b>Age (years)</b>								
40-44	*	*	*	*	1,078	246,800	27.5	(18.5 - 36.5)
45-54	*	*	*	*	3,074	650,000	47.6	(40.8 - 54.4)
55-64	*	*	*	*	3,467	673,500	68.1	(62.4 - 73.8)
≥ 65	72	2,400	80.2	(70.4 - 90.0)	4,304	809,900	79.7	(75.0 - 84.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	175	6,500	62.9	(54.4 - 71.5)	11,305	1,875,600	62.2	(59.2 - 65.2)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	4,963	718,700	49.6	(43.2 - 56.1)
Some College	*	*	*	*	2,665	511,500	58.5	(50.8 - 66.3)
≥ College Degree	*	*	*	*	4,277	1,148,400	59.4	(52.8 - 66.0)
<b>Income</b>								
≤ \$24,999	*	*	*	*	2,553	388,000	45.0	(35.0 - 55.0)
\$25,000-\$49,999	*	*	*	*	3,139	422,400	60.5	(53.0 - 68.0)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,295	978,300	56.9	(51.1 - 62.7)

<sup>1</sup> Defined as having had a digital rectal exam within the past 2 years among men aged 40 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of men aged 40 and older who have had a digital rectal exam within the past 2 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 30b. Had digital rectal exam within the past 2 years among men aged 40 and older<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	182	61.7	(52.6 - 70.7)	11,923	55.9	(51.7 - 60.0)
<b>Gender</b>						
Male	182	61.7	(52.6 - 70.7)	11,923	55.9	(51.7 - 60.0)
Female	-	-	-	-	-	-
<b>Race/Ethnicity</b>						
White Non-Hispanic	175	62.0	(52.9 - 71.0)	11,305	62.0	(58.9 - 65.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	4,963	50.5	(43.7 - 57.3)
Some College	*	*	*	2,665	58.3	(50.4 - 66.2)
≥ College Degree	*	*	*	4,277	59.2	(52.4 - 65.9)
<b>Income</b>						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,139	60.8	(53.1 - 68.4)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,295	56.5	(50.5 - 62.4)

<sup>1</sup> Defined as having had a digital rectal exam within the past 2 years among men aged 40 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 31a. Ever had a prostate specific antigen test among men aged 40 and older<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	177	7,000	64.3	(55.7 - 72.9)	11,557	2,822,600	67.6	(63.9 - 71.2)
<b>Gender</b>								
Male	177	7,000	64.3	(55.7 - 72.9)	11,557	2,822,600	67.6	(63.9 - 71.2)
Female	-	-	-	-	-	-	-	-
<b>Age (years)</b>								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,938	760,600	57.4	(50.6 - 64.3)
55-64	*	*	*	*	3,373	811,400	83.4	(78.8 - 88.1)
≥ 65	71	2,900	96.1	(92.7 - 99.5)	4,215	905,400	90.4	(87.1 - 93.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	170	6,600	65.5	(56.8 - 74.1)	10,950	1,965,400	67.0	(63.9 - 70.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	4,775	829,200	59.0	(52.4 - 65.6)
Some College	*	*	*	*	2,584	535,000	61.9	(53.7 - 70.2)
≥ College Degree	*	*	*	*	4,176	1,455,700	76.8	(72.5 - 81.1)
<b>Income</b>								
≤ \$24,999	*	*	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	*	3,048	444,100	64.1	(56.0 - 72.3)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,188	1,211,100	72.3	(67.6 - 77.0)

<sup>1</sup> Defined as having ever had a prostate specific antigen test among men aged 40 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of men aged 40 and older who have ever had a prostate specific antigen test (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 31b. Ever had a prostate specific antigen test among men aged 40 and older<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	177	62.0	(52.7 - 71.4)	11,557	67.4	(63.7 - 71.1)
<b>Gender</b>						
Male	177	62.0	(52.7 - 71.4)	11,557	67.4	(63.7 - 71.1)
Female	-	-	-	-	-	-
<b>Race/Ethnicity</b>						
White Non-Hispanic	170	63.1	(53.7 - 72.4)	10,950	66.6	(63.4 - 69.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	4,775	59.7	(52.9 - 66.4)
Some College	*	*	*	2,584	61.4	(53.0 - 69.9)
≥ College Degree	*	*	*	4,176	76.3	(71.9 - 80.7)
<b>Income</b>						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,048	64.2	(56.1 - 72.4)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,188	71.5	(66.6 - 76.3)

<sup>1</sup> Defined as having ever had a prostate specific antigen test among men aged 40 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 32a. Had prostate specific antigen test within the past 2 years among men aged 40 and older<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	174	6,300	58.4	(49.6 - 67.2)	11,437	2,493,800	59.9	(56.0 - 63.9)
<b>Gender</b>								
Male	174	6,300	58.4	(49.6 - 67.2)	11,437	2,493,800	59.9	(56.0 - 63.9)
Female	-	-	-	-	-	-	-	-
<b>Age (years)</b>								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,917	659,400	49.9	(42.8 - 57.0)
55-64	*	*	*	*	3,326	719,700	74.5	(69.4 - 79.7)
≥ 65	70	2,700	89.0	(80.6 - 97.4)	4,166	830,000	83.3	(78.4 - 88.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	167	6,000	60.0	(51.1 - 68.9)	10,836	1,762,200	60.4	(57.3 - 63.5)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	4,727	734,200	52.4	(45.7 - 59.1)
Some College	*	*	*	*	2,558	483,800	56.3	(48.2 - 64.4)
≥ College Degree	*	*	*	*	4,133	1,273,200	67.4	(61.6 - 73.3)
<b>Income</b>								
≤ \$24,999	*	*	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	*	3,021	410,800	59.6	(51.5 - 67.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,170	1,080,100	64.6	(59.1 - 70.0)

<sup>1</sup> Defined as having had a prostate specific antigen test within the past 2 years among men aged 40 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of men aged 40 and older who have had a prostate specific antigen test within the past 2 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 32b. Had prostate specific antigen test within the past 2 years among men aged 40 and older<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	174	56.4	(47.0 - 65.8)	11,437	59.8	(55.8 - 63.9)
<b>Gender</b>						
Male	174	56.4	(47.0 - 65.8)	11,437	59.8	(55.8 - 63.9)
Female	-	-	-	-	-	-
<b>Race/Ethnicity</b>						
White Non-Hispanic	167	57.8	(48.4 - 67.3)	10,836	60.1	(56.9 - 63.4)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	4,727	52.9	(45.9 - 59.9)
Some College	*	*	*	2,558	56.0	(47.7 - 64.2)
≥ College Degree	*	*	*	4,133	67.1	(61.2 - 73.1)
<b>Income</b>						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,021	59.8	(51.7 - 67.9)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,170	63.9	(58.3 - 69.6)

<sup>1</sup> Defined as having had a prostate specific antigen test within the past 2 years among men aged 40 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 33a. Prostate cancer among men aged 40 and older<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	185	600	5.4	(2.2 - 8.6)	12,133	178,200	4.1	(3.0 - 5.2)
<b>Gender</b>								
Male	185	600	5.4	(2.2 - 8.6)	12,133	178,200	4.1	(3.0 - 5.2)
Female	-	-	-	-	-	-	-	-
<b>Age (years)</b>								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	3,120	6,100	0.4	(0.0 - 0.9)
55-64	*	*	*	*	3,532	31,600	3.1	(1.5 - 4.8)
≥ 65	74	500	14.2	(5.9 - 22.6)	4,388	140,400	13.6	(9.7 - 17.4)
<b>Race/Ethnicity</b>								
White Non-Hispanic	178	600	5.8	(2.4 - 9.2)	11,502	143,700	4.7	(3.5 - 5.8)
Black Non-Hispanic	*	*	*	*	184	23,500	5.1	(0.0 - 11.0)
Hispanic	*	*	*	*	197	7,700	1.6	(0.0 - 3.7)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	*	*	*	*	5,065	46,300	3.1	(2.2 - 4.1)
Some College	*	*	*	*	2,709	39,200	4.4	(1.3 - 7.5)
≥ College Degree	*	*	*	*	4,336	92,500	4.7	(3.0 - 6.5)
<b>Income</b>								
≤ \$24,999	*	*	*	*	2,597	35,800	4.1	(2.2 - 6.0)
\$25,000-\$49,999	*	*	*	*	3,196	29,600	4.1	(2.9 - 5.4)
\$50,000-\$74,999	*	*	*	*	2,007	20,300	2.8	(1.2 - 4.4)
≥ \$75,000	*	*	*	*	3,322	56,500	3.2	(1.7 - 4.8)

<sup>1</sup> Defined as having ever been told by a health professional that respondent had prostate cancer among men aged 40 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of men aged 40 and older who have ever been told they had prostate cancer (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 33b. Prostate cancer among men aged 40 and older<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	185	5.7	(2.1 - 9.3)	12,133	4.5	(3.2 - 5.7)
<b>Gender</b>						
Male	185	5.7	(2.1 - 9.3)	12,133	4.5	(3.2 - 5.7)
Female	-	-	-	-	-	-
<b>Race/Ethnicity</b>						
White Non-Hispanic	178	6.1	(2.2 - 10.1)	11,502	5.1	(3.8 - 6.4)
Black Non-Hispanic	*	*	*	184	5.7	(0.0 - 12.8)
Hispanic	*	*	*	197	1.8	(0.0 - 4.4)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	5,065	3.3	(2.3 - 4.3)
Some College	*	*	*	2,709	5.0	(1.1 - 9.0)
≥ College Degree	*	*	*	4,336	5.2	(3.1 - 7.2)
<b>Income</b>						
≤ \$24,999	*	*	*	2,597	4.5	(2.2 - 6.7)
\$25,000-\$49,999	*	*	*	3,196	4.3	(3.0 - 5.6)
\$50,000-\$74,999	*	*	*	2,007	3.0	(1.1 - 5.0)
≥ \$75,000	*	*	*	3,322	3.5	(1.6 - 5.4)

<sup>1</sup> Defined as having ever been told by a health professional that respondent had prostate cancer among men aged 40 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 34a. Home blood stool test ever used among adults aged 50 and older<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	423	6,000	34.8	(29.5 - 40.1)	25,322	2,288,500	38.0	(35.5 - 40.4)
<b>Gender</b>								
Male	147	3,000	40.1	(31.0 - 49.2)	9,571	1,081,900	38.9	(35.1 - 42.6)
Female	276	3,000	30.6	(24.6 - 36.7)	15,751	1,206,600	37.2	(34.0 - 40.4)
<b>Age (years)</b>								
50-64	198	2,800	30.3	(22.5 - 38.2)	12,834	1,192,000	34.0	(30.8 - 37.2)
≥ 65	225	3,200	39.8	(32.7 - 46.8)	12,472	1,093,400	43.6	(39.8 - 47.4)
<b>Race/Ethnicity</b>								
White Non-Hispanic	409	5,900	36.1	(30.7 - 41.5)	24,213	1,754,500	39.1	(37.1 - 41.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	243	3,100	33.2	(26.3 - 40.0)	11,146	823,500	34.9	(30.5 - 39.2)
Some College	*	*	*	*	6,261	480,300	37.0	(32.6 - 41.4)
≥ College Degree	*	*	*	*	7,851	973,700	41.6	(37.8 - 45.3)
<b>Income</b>								
≤ \$24,999	153	1,600	32.6	(24.2 - 41.0)	7,222	505,000	35.4	(29.7 - 41.2)
\$25,000-\$49,999	*	*	*	*	6,609	467,600	40.3	(35.2 - 45.4)
\$50,000-\$74,999	*	*	*	*	3,500	313,000	38.2	(32.6 - 43.9)
≥ \$75,000	*	*	*	*	4,647	657,800	37.3	(33.0 - 41.6)

<sup>1</sup> Defined as having ever used a home blood stool test among adults aged 50 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 50 and older who have ever used a home blood stool test (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 34b. Home blood stool test ever used among adults aged 50 and older<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	423	34.3	(28.9 - 39.6)	25,322	38.1	(35.5 - 40.8)
<b>Gender</b>						
Male	147	40.5	(31.3 - 49.6)	9,571	39.4	(35.3 - 43.5)
Female	276	29.0	(23.0 - 35.0)	15,751	36.9	(33.5 - 40.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	409	35.6	(30.0 - 41.1)	24,213	38.9	(36.7 - 41.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	243	32.4	(25.4 - 39.5)	11,146	35.1	(30.3 - 39.9)
Some College	*	*	*	6,261	36.6	(32.0 - 41.3)
≥ College Degree	*	*	*	7,851	41.9	(37.9 - 46.0)
<b>Income</b>						
≤ \$24,999	153	32.2	(23.5 - 40.8)	7,222	35.9	(29.5 - 42.4)
\$25,000-\$49,999	*	*	*	6,609	40.2	(34.8 - 45.6)
\$50,000-\$74,999	*	*	*	3,500	38.1	(31.9 - 44.3)
≥ \$75,000	*	*	*	4,647	37.5	(32.9 - 42.2)

<sup>1</sup> Defined as having ever used a home blood stool test among adults aged 50 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 35a. Home blood stool test used within the past year among adults aged 50 and older<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	418	2,500	14.5	(10.5 - 18.4)	25,074	863,300	14.4	(12.6 - 16.2)
<b>Gender</b>								
Male	144	1,500	20.4	(13.2 - 27.6)	9,508	444,400	16.0	(13.2 - 18.9)
Female	274	1,000	9.9	(5.7 - 14.1)	15,566	418,900	13.1	(10.8 - 15.4)
<b>Age (years)</b>								
50-64	196	1,300	14.6	(8.7 - 20.6)	12,759	454,400	13.0	(10.6 - 15.5)
≥ 65	222	1,100	14.3	(9.3 - 19.2)	12,299	406,400	16.4	(13.7 - 19.0)
<b>Race/Ethnicity</b>								
White Non-Hispanic	404	2,400	15.0	(10.9 - 19.1)	23,971	634,300	14.3	(12.8 - 15.7)
Black Non-Hispanic	*	*	*	*	396	136,000	18.6	(9.4 - 27.8)
Hispanic	*	*	*	*	349	53,400	9.9	(2.5 - 17.2)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	239	1,400	15.4	(10.0 - 20.8)	11,026	288,600	12.3	(9.4 - 15.3)
Some College	104	600	13.1	(5.8 - 20.5)	6,208	212,500	16.4	(12.5 - 20.4)
≥ College Degree	75	500	13.6	(4.6 - 22.6)	7,778	352,400	15.2	(12.5 - 17.9)
<b>Income</b>								
≤ \$24,999	152	800	15.1	(8.4 - 21.9)	7,157	196,300	13.9	(10.1 - 17.6)
\$25,000-\$49,999	120	1,000	18.4	(10.3 - 26.5)	6,562	221,500	19.2	(14.1 - 24.3)
\$50,000-\$74,999	*	*	*	*	3,469	111,900	13.7	(10.1 - 17.4)
≥ \$75,000	*	*	*	*	4,623	205,100	11.7	(8.8 - 14.6)

<sup>1</sup> Defined as having used a home blood stool test within the past year among adults aged 50 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 50 and older who have used a home blood stool test within the past year (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 35b. Home blood stool test used within the past year among adults aged 50 and older<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	418	14.6	(10.5 - 18.6)	25,074	14.6	(12.6 - 16.7)
<b>Gender</b>						
Male	144	20.2	(13.1 - 27.4)	9,508	16.4	(13.2 - 19.6)
Female	274	9.8	(5.5 - 14.1)	15,566	12.9	(10.4 - 15.4)
<b>Race/Ethnicity</b>						
White Non-Hispanic	404	15.1	(10.9 - 19.3)	23,971	14.2	(12.7 - 15.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	349	10.2	(1.7 - 18.7)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	239	15.4	(9.8 - 21.0)	11,026	12.7	(9.2 - 16.1)
Some College	104	13.6	(5.9 - 21.4)	6,208	16.6	(12.3 - 20.9)
≥ College Degree	75	13.4	(4.7 - 22.2)	7,778	15.3	(12.4 - 18.3)
<b>Income</b>						
≤ \$24,999	152	15.9	(8.7 - 23.1)	7,157	14.3	(9.8 - 18.7)
\$25,000-\$49,999	120	17.8	(9.8 - 25.7)	6,562	19.3	(13.8 - 24.8)
\$50,000-\$74,999	*	*	*	3,469	13.8	(9.8 - 17.7)
≥ \$75,000	*	*	*	4,623	11.8	(8.6 - 15.0)

<sup>1</sup> Defined as having used a home blood stool test within the past year among adults aged 50 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 36a. Home blood stool test used within the past 2 years among adults aged 50 and older<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	418	3,600	21.3	(16.5 - 26.0)	25,074	1,218,000	20.4	(18.3 - 22.5)
<b>Gender</b>								
Male	144	2,200	30.0	(21.4 - 38.7)	9,508	593,100	21.4	(18.3 - 24.5)
Female	274	1,400	14.5	(9.7 - 19.3)	15,566	624,900	19.5	(16.6 - 22.4)
<b>Age (years)</b>								
50-64	196	2,000	22.2	(14.9 - 29.5)	12,759	629,900	18.1	(15.4 - 20.7)
≥ 65	222	1,600	20.2	(14.4 - 26.0)	12,299	585,500	23.6	(20.2 - 27.0)
<b>Race/Ethnicity</b>								
White Non-Hispanic	404	3,500	21.8	(16.9 - 26.7)	23,971	877,500	19.7	(18.2 - 21.3)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	239	1,800	19.2	(13.4 - 25.0)	11,026	454,600	19.4	(15.6 - 23.3)
Some College	104	900	20.8	(11.8 - 29.8)	6,208	262,200	20.3	(16.3 - 24.2)
≥ College Degree	*	*	*	*	7,778	491,400	21.2	(18.1 - 24.2)
<b>Income</b>								
≤ \$24,999	152	1,000	20.0	(12.4 - 27.6)	7,157	263,900	18.6	(14.6 - 22.7)
\$25,000-\$49,999	*	*	*	*	6,562	285,900	24.8	(19.6 - 30.0)
\$50,000-\$74,999	*	*	*	*	3,469	167,500	20.6	(15.6 - 25.5)
≥ \$75,000	*	*	*	*	4,623	294,900	16.8	(13.7 - 20.0)

<sup>1</sup> Defined as having used a home blood stool test within the past 2 years among adults aged 50 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 50 and older who have used a home blood stool test within the past 2 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 36b. Home blood stool test used within the past 2 years among adults aged 50 and older<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	418	21.5	(16.6 - 26.4)	25,074	20.5	(18.2 - 22.8)
<b>Gender</b>						
Male	144	30.1	(21.4 - 38.8)	9,508	21.7	(18.3 - 25.1)
Female	274	14.3	(9.4 - 19.1)	15,566	19.4	(16.3 - 22.5)
<b>Race/Ethnicity</b>						
White Non-Hispanic	404	22.1	(17.1 - 27.2)	23,971	19.7	(18.0 - 21.3)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	239	19.2	(13.2 - 25.2)	11,026	19.5	(15.4 - 23.7)
Some College	104	21.7	(12.3 - 31.1)	6,208	20.4	(16.1 - 24.7)
≥ College Degree	*	*	*	7,778	21.4	(18.1 - 24.7)
<b>Income</b>						
≤ \$24,999	152	20.6	(12.6 - 28.7)	7,157	18.9	(14.2 - 23.6)
\$25,000-\$49,999	*	*	*	6,562	24.8	(19.3 - 30.3)
\$50,000-\$74,999	*	*	*	3,469	20.6	(15.3 - 25.9)
≥ \$75,000	*	*	*	4,623	16.9	(13.4 - 20.4)

<sup>1</sup> Defined as having used a home blood stool test within the past 2 years among adults aged 50 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 37a. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	423	11,500	66.9	(61.9 - 72.0)	25,357	4,019,100	66.6	(64.1 - 69.1)
<b>Gender</b>								
Male	145	5,200	69.1	(60.6 - 77.5)	9,582	1,894,100	68.0	(64.2 - 71.7)
Female	278	6,400	65.3	(59.1 - 71.5)	15,775	2,125,000	65.5	(62.1 - 68.8)
<b>Age (years)</b>								
50-54	*	*	*	*	4,179	718,900	50.6	(44.6 - 56.5)
55-64	128	4,400	76.9	(69.4 - 84.4)	8,695	1,493,500	71.0	(67.3 - 74.8)
≥ 65	225	5,800	71.8	(65.5 - 78.2)	12,466	1,802,000	72.1	(68.6 - 75.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	408	10,900	67.2	(62.1 - 72.3)	24,248	3,093,900	68.8	(66.8 - 70.8)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	243	6,300	66.4	(59.6 - 73.1)	11,121	1,445,100	61.2	(56.7 - 65.8)
Some College	*	*	*	*	6,287	835,300	64.3	(59.7 - 69.0)
≥ College Degree	*	*	*	*	7,884	1,732,400	73.7	(70.0 - 77.4)
<b>Income</b>								
≤ \$24,999	154	2,900	57.3	(48.5 - 66.0)	7,220	892,600	62.7	(56.8 - 68.6)
\$25,000-\$49,999	122	3,700	70.9	(61.6 - 80.2)	6,619	697,300	60.1	(54.6 - 65.7)
\$50,000-\$74,999	*	*	*	*	3,514	613,700	74.2	(68.9 - 79.6)
≥ \$75,000	*	*	*	*	4,665	1,294,400	73.3	(69.2 - 77.5)

<sup>1</sup> Defined as having ever had a sigmoidoscopy or colonoscopy among adults aged 50 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 50 and older who have ever had a sigmoidoscopy or colonoscopy (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 37b. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	423	65.5	(60.2 - 70.8)	25,357	66.6	(63.9 - 69.3)
<b>Gender</b>						
Male	145	68.2	(59.6 - 76.8)	9,582	68.5	(64.5 - 72.5)
Female	278	63.3	(56.8 - 69.8)	15,775	64.8	(61.2 - 68.4)
<b>Race/Ethnicity</b>						
White Non-Hispanic	408	66.0	(60.7 - 71.3)	24,248	68.7	(66.5 - 70.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	243	64.8	(57.6 - 72.0)	11,121	61.3	(56.4 - 66.2)
Some College	*	*	*	6,287	63.8	(58.8 - 68.8)
≥ College Degree	*	*	*	7,884	74.0	(70.2 - 77.9)
<b>Income</b>						
≤ \$24,999	154	55.8	(46.7 - 64.9)	7,220	63.0	(56.7 - 69.3)
\$25,000-\$49,999	122	70.2	(60.4 - 79.9)	6,619	59.7	(53.8 - 65.6)
\$50,000-\$74,999	*	*	*	3,514	74.4	(68.3 - 80.5)
≥ \$75,000	*	*	*	4,665	73.6	(69.3 - 78.0)

<sup>1</sup> Defined as having ever had a sigmoidoscopy or colonoscopy among adults aged 50 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 38a. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	421	10,900	63.2	(58.1 - 68.4)	25,169	3,855,800	64.6	(62.1 - 67.1)
<b>Gender</b>								
Male	145	5,000	66.5	(57.9 - 75.0)	9,521	1,835,400	66.4	(62.6 - 70.1)
Female	276	5,900	60.8	(54.4 - 67.1)	15,648	2,020,400	63.1	(59.8 - 66.4)
<b>Age (years)</b>								
50-54	*	*	*	*	4,169	700,500	49.3	(43.3 - 55.3)
55-64	126	4,100	72.9	(64.8 - 80.9)	8,652	1,455,100	69.4	(65.7 - 73.2)
≥ 65	225	5,500	67.3	(60.7 - 73.9)	12,331	1,695,700	69.5	(65.9 - 73.1)
<b>Race/Ethnicity</b>								
White Non-Hispanic	406	10,300	63.3	(58.1 - 68.5)	24,069	2,981,500	66.6	(64.6 - 68.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	241	5,900	62.1	(55.1 - 69.0)	11,017	1,360,400	59.0	(54.5 - 63.5)
Some College	*	*	*	*	6,251	807,100	62.4	(57.8 - 67.1)
≥ College Degree	*	*	*	*	7,841	1,682,500	71.7	(68.0 - 75.4)
<b>Income</b>								
≤ \$24,999	154	2,600	50.7	(41.9 - 59.6)	7,163	858,900	61.0	(55.1 - 67.0)
\$25,000-\$49,999	122	3,500	66.7	(57.1 - 76.3)	6,584	664,900	57.7	(52.2 - 63.2)
\$50,000-\$74,999	*	*	*	*	3,501	602,600	73.1	(67.7 - 78.5)
≥ \$75,000	*	*	*	*	4,655	1,260,400	71.5	(67.3 - 75.7)

<sup>1</sup> Defined as having had a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 50 and older who have had a sigmoidoscopy or colonoscopy within the past 10 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 38b. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	421	61.9	(56.5 - 67.3)	25,169	64.6	(61.9 - 67.3)
<b>Gender</b>						
Male	145	65.6	(56.8 - 74.3)	9,521	66.9	(62.9 - 70.9)
Female	276	58.9	(52.3 - 65.5)	15,648	62.4	(58.9 - 66.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	406	62.2	(56.8 - 67.7)	24,069	66.5	(64.3 - 68.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	241	60.5	(53.2 - 67.8)	11,017	59.0	(54.1 - 63.9)
Some College	*	*	*	6,251	61.9	(56.9 - 66.9)
≥ College Degree	*	*	*	7,841	72.1	(68.2 - 75.9)
<b>Income</b>						
≤ \$24,999	154	49.5	(40.3 - 58.6)	7,163	61.3	(54.9 - 67.6)
\$25,000-\$49,999	122	66.2	(56.3 - 76.2)	6,584	57.2	(51.4 - 63.0)
\$50,000-\$74,999	*	*	*	3,501	73.3	(67.2 - 79.4)
≥ \$75,000	*	*	*	4,655	71.8	(67.4 - 76.2)

<sup>1</sup> Defined as having had a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 39a. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	411	11,200	66.3	(61.2 - 71.4)	24,792	4,049,200	69.1	(66.7 - 71.6)
<b>Gender</b>								
Male	140	5,100	70.5	(62.0 - 79.0)	9,418	1,945,200	70.7	(67.0 - 74.3)
Female	271	6,000	63.2	(56.9 - 69.5)	15,374	2,104,100	67.8	(64.6 - 71.1)
<b>Age (years)</b>								
50-54	*	*	*	*	4,132	751,200	54.1	(48.0 - 60.1)
55-64	119	4,200	76.5	(68.8 - 84.2)	8,366	1,487,600	73.7	(70.0 - 77.4)
≥ 65	222	5,600	70.0	(63.5 - 76.5)	12,294	1,810,500	74.0	(70.6 - 77.4)
<b>Race/Ethnicity</b>								
White Non-Hispanic	396	10,600	66.5	(61.4 - 71.7)	23,726	3,091,400	70.4	(68.4 - 72.4)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	236	5,900	64.5	(57.5 - 71.4)	10,872	1,461,600	64.2	(59.8 - 68.7)
Some College	102	2,900	68.7	(58.8 - 78.6)	6,170	873,300	68.4	(63.9 - 72.8)
≥ College Degree	*	*	*	*	7,708	1,709,000	74.6	(70.9 - 78.3)
<b>Income</b>								
≤ \$24,999	153	2,700	53.1	(44.2 - 62.0)	7,092	923,300	66.0	(60.1 - 71.9)
\$25,000-\$49,999	116	3,600	72.4	(63.0 - 81.7)	6,526	730,000	64.2	(58.9 - 69.6)
\$50,000-\$74,999	*	*	*	*	3,467	621,500	76.6	(71.4 - 81.9)
≥ \$75,000	*	*	*	*	4,615	1,292,500	73.8	(69.8 - 77.9)

<sup>1</sup> Defined as having used a home blood stool test within the past year or a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 years and older

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults aged 50 and older who have used a home blood stool test in the past year or sigmoidoscopy or colonoscopy within the past 10 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 39b. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	411	65.1	(59.7 - 70.4)	24,792	69.3	(66.7 - 71.9)
<b>Gender</b>						
Male	140	69.4	(60.7 - 78.1)	9,418	71.5	(67.6 - 75.4)
Female	271	61.5	(54.9 - 68.1)	15,374	67.2	(63.7 - 70.7)
<b>Race/Ethnicity</b>						
White Non-Hispanic	396	65.5	(60.1 - 70.9)	23,726	70.3	(68.1 - 72.4)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	236	62.9	(55.6 - 70.3)	10,872	64.5	(59.8 - 69.3)
Some College	*	*	*	6,170	68.2	(63.4 - 72.9)
≥ College Degree	*	*	*	7,708	75.0	(71.2 - 78.8)
<b>Income</b>						
≤ \$24,999	153	51.8	(42.6 - 61.0)	7,092	66.4	(60.2 - 72.6)
\$25,000-\$49,999	116	71.9	(62.2 - 81.7)	6,526	63.8	(58.2 - 69.5)
\$50,000-\$74,999	*	*	*	3,467	76.7	(70.6 - 82.7)
≥ \$75,000	*	*	*	4,615	74.4	(70.2 - 78.6)

<sup>1</sup> Defined as having used a home blood stool test within the past year or a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 years and older

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 40a. High blood pressure among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	640	12,000	31.9	(27.6 - 36.2)	38,125	3,974,800	26.8	(24.9 - 28.7)
<b>Gender</b>								
Male	227	5,900	33.3	(26.3 - 40.4)	14,611	1,960,500	27.7	(24.7 - 30.7)
Female	413	6,000	30.6	(25.5 - 35.7)	23,514	2,014,300	26.0	(23.6 - 28.3)
<b>Age (years)</b>								
18-34	*	*	*	*	3,935	312,200	7.0	(4.2 - 9.8)
35-44	74	900	14.7	(5.4 - 24.1)	4,984	426,200	14.8	(11.6 - 18.1)
45-54	138	2,100	29.3	(20.3 - 38.2)	7,825	862,400	30.3	(25.9 - 34.8)
55-64	127	3,100	55.4	(45.7 - 65.1)	8,765	890,800	41.6	(37.6 - 45.7)
≥ 65	230	5,000	59.9	(52.9 - 66.9)	12,616	1,483,200	58.5	(54.7 - 62.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	612	11,500	33.9	(29.5 - 38.2)	35,814	2,650,900	28.4	(26.9 - 30.0)
Black Non-Hispanic	*	*	*	*	736	558,500	26.2	(19.6 - 32.8)
Hispanic	*	*	*	*	863	538,100	23.8	(17.0 - 30.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	323	6,700	37.8	(31.3 - 44.2)	15,688	1,851,200	33.1	(29.4 - 36.9)
Some College	171	3,200	28.8	(20.8 - 36.8)	9,906	851,100	24.1	(20.9 - 27.3)
≥ College Degree	146	2,100	23.8	(15.9 - 31.6)	12,452	1,253,400	22.3	(19.7 - 24.9)
<b>Income</b>								
≤ \$24,999	194	3,600	39.6	(29.9 - 49.4)	9,640	1,153,000	32.8	(27.6 - 38.1)
\$25,000-\$49,999	184	3,800	35.2	(27.0 - 43.3)	9,657	807,800	27.9	(23.7 - 32.1)
\$50,000-\$74,999	*	*	*	*	5,773	507,600	25.8	(21.4 - 30.1)
≥ \$75,000	97	1,900	25.1	(15.8 - 34.4)	8,560	942,500	21.0	(18.4 - 23.6)

<sup>1</sup> Defined as having ever been told by a health professional that respondent has high blood pressure

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have ever been told by a health professional that they have high blood pressure (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 40b. High blood pressure among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	640	27.5	(23.3 - 31.8)	38,125	25.7	(23.8 - 27.5)
<b>Gender</b>						
Male	227	30.4	(23.3 - 37.5)	14,611	27.7	(24.7 - 30.7)
Female	413	24.9	(19.9 - 30.0)	23,514	23.8	(21.5 - 26.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	612	29.4	(24.9 - 33.8)	35,814	27.2	(25.6 - 28.7)
Black Non-Hispanic	*	*	*	736	24.9	(18.4 - 31.3)
Hispanic	*	*	*	863	23.3	(16.6 - 30.0)
Other Non-Hispanic	*	*	*	712	19.3	(9.5 - 29.1)
<b>Education</b>						
≤ High School	323	33.7	(26.9 - 40.6)	15,688	32.1	(28.4 - 35.8)
Some College	171	24.5	(16.9 - 32.0)	9,906	22.8	(19.7 - 25.9)
≥ College Degree	146	19.6	(12.4 - 26.8)	12,452	21.2	(18.6 - 23.8)
<b>Income</b>						
≤ \$24,999	194	34.6	(24.8 - 44.4)	9,640	31.8	(26.7 - 37.0)
\$25,000-\$49,999	184	29.9	(21.7 - 38.1)	9,657	26.9	(22.8 - 30.9)
\$50,000-\$74,999	*	*	*	5,773	24.8	(20.4 - 29.2)
≥ \$75,000	97	22.2	(13.1 - 31.3)	8,560	19.9	(17.3 - 22.5)

<sup>1</sup> Defined as having ever been told by a health professional that respondent has high blood pressure

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 41a. Blood pressure medication among adults with high blood pressure<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	263	10,100	84.7	(78.3 - 91.1)	15,288	3,145,800	79.2	(76.0 - 82.4)
<b>Gender</b>								
Male	101	5,200	89.0	(80.7 - 97.4)	6,080	1,432,100	73.1	(67.6 - 78.5)
Female	162	4,900	80.5	(71.0 - 90.0)	9,208	1,713,700	85.2	(82.1 - 88.2)
<b>Age (years)</b>								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,315	645,500	74.9	(66.8 - 82.9)
55-64	69	3,000	95.2	(90.2 - 100)	4,045	782,300	87.8	(84.5 - 91.1)
≥ 65	139	4,800	96.2	(92.8 - 99.6)	7,675	1,383,400	93.4	(91.1 - 95.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	256	9,600	84.1	(77.5 - 90.7)	14,442	2,117,300	79.9	(77.5 - 82.4)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	160	5,500	81.0	(71.6 - 90.4)	7,354	1,482,800	80.2	(74.7 - 85.6)
Some College	61	2,800	89.1	(80.4 - 97.8)	3,868	632,000	74.3	(67.8 - 80.7)
≥ College Degree	*	*	*	*	4,035	1,013,200	80.9	(76.8 - 84.9)
<b>Income</b>								
≤ \$24,999	98	3,200	88.9	(81.5 - 96.3)	4,898	948,200	82.3	(75.7 - 88.9)
\$25,000-\$49,999	*	*	*	*	4,088	668,300	82.8	(77.6 - 88.0)
\$50,000-\$74,999	*	*	*	*	1,971	391,500	77.2	(67.7 - 86.6)
≥ \$75,000	*	*	*	*	2,397	683,700	72.5	(66.1 - 78.9)

<sup>1</sup> Defined as currently taking medicine for high blood pressure among adults with high blood pressure

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults with high blood pressure who are taking medicine for high blood pressure (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 41b. Blood pressure medication among adults with high blood pressure<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	263	80.6	(72.2 - 89.0)	15,288	78.5	(75.2 - 81.7)
<b>Gender</b>						
Male	*	*	*	6,080	73.3	(67.9 - 78.7)
Female	*	*	*	9,208	84.0	(80.7 - 87.2)
<b>Race/Ethnicity</b>						
White Non-Hispanic	256	79.9	(71.3 - 88.5)	14,442	79.0	(76.5 - 81.6)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	*	*	*	7,354	79.7	(74.3 - 85.2)
Some College	*	*	*	3,868	72.5	(65.7 - 79.3)
≥ College Degree	*	*	*	4,035	80.4	(76.1 - 84.6)
<b>Income</b>						
≤ \$24,999	98	86.9	(77.5 - 96.2)	4,898	81.6	(75.0 - 88.3)
\$25,000-\$49,999	*	*	*	4,088	82.1	(76.7 - 87.5)
\$50,000-\$74,999	*	*	*	1,971	76.4	(66.7 - 86.1)
≥ \$75,000	*	*	*	2,397	71.6	(64.8 - 78.3)

<sup>1</sup> Defined as currently taking medicine for high blood pressure among adults with high blood pressure

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 42a. Ever had blood cholesterol checked among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	634	29,700	80.1	(75.5 - 84.7)	37,450	11,496,600	80.3	(77.7 - 83.0)
<b>Gender</b>								
Male	222	13,900	80.1	(72.8 - 87.4)	14,389	5,288,400	77.7	(73.3 - 82.0)
Female	412	15,700	80.1	(74.4 - 85.8)	23,061	6,208,200	82.7	(79.6 - 85.8)
<b>Age (years)</b>								
18-34	*	*	*	*	3,684	2,296,400	55.3	(48.7 - 61.9)
35-44	*	*	*	*	4,883	2,264,200	81.6	(76.6 - 86.5)
45-54	137	6,500	91.4	(85.9 - 96.8)	7,731	2,539,400	91.0	(87.0 - 94.9)
55-64	128	5,600	97.5	(94.7 - 100)	8,718	2,014,900	95.4	(93.5 - 97.2)
≥ 65	229	8,100	96.7	(93.8 - 99.6)	12,434	2,381,700	95.9	(94.3 - 97.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	606	26,900	80.4	(75.7 - 85.1)	35,198	7,832,600	86.3	(84.8 - 87.9)
Black Non-Hispanic	*	*	*	*	720	1,588,900	77.8	(68.4 - 87.2)
Hispanic	*	*	*	*	842	1,224,600	56.8	(46.8 - 66.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	322	13,800	77.8	(70.8 - 84.7)	15,342	3,795,700	70.6	(65.6 - 75.6)
Some College	171	8,800	80.3	(71.9 - 88.8)	9,732	2,599,800	77.4	(71.7 - 83.2)
≥ College Degree	141	7,100	84.8	(76.9 - 92.7)	12,296	5,008,300	91.3	(88.9 - 93.6)
<b>Income</b>								
≤ \$24,999	193	7,200	78.6	(69.2 - 88.0)	9,437	2,277,500	68.4	(61.5 - 75.3)
\$25,000-\$49,999	183	8,400	80.2	(71.8 - 88.6)	9,492	2,053,000	71.9	(64.5 - 79.3)
\$50,000-\$74,999	*	*	*	*	5,721	1,736,100	90.0	(87.7 - 92.3)
≥ \$75,000	95	6,400	87.7	(79.0 - 96.3)	8,460	4,019,300	92.2	(90.8 - 93.6)

<sup>1</sup> Defined as ever having blood cholesterol checked

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have ever had blood cholesterol checked (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 42b. Ever had blood cholesterol checked among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	634	76.9	(71.7 - 82.1)	37,450	79.9	(77.3 - 82.5)
<b>Gender</b>						
Male	222	77.9	(69.8 - 86.0)	14,389	78.0	(73.9 - 82.1)
Female	412	76.1	(69.4 - 82.8)	23,061	81.6	(78.4 - 84.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	606	76.9	(71.5 - 82.3)	35,198	85.6	(84.0 - 87.2)
Black Non-Hispanic	*	*	*	720	77.5	(68.2 - 86.7)
Hispanic	*	*	*	842	57.2	(47.4 - 67.0)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	322	73.5	(65.4 - 81.6)	15,342	70.2	(65.2 - 75.1)
Some College	171	77.0	(67.3 - 86.6)	9,732	76.7	(71.1 - 82.3)
≥ College Degree	141	83.5	(74.9 - 92.0)	12,296	90.9	(88.6 - 93.2)
<b>Income</b>						
≤ \$24,999	*	*	*	9,437	68.1	(61.3 - 74.9)
\$25,000-\$49,999	183	76.3	(66.6 - 86.0)	9,492	71.6	(64.4 - 78.7)
\$50,000-\$74,999	*	*	*	5,721	89.6	(87.2 - 92.0)
≥ \$75,000	95	86.5	(77.1 - 95.9)	8,460	91.7	(90.1 - 93.2)

<sup>1</sup> Defined as ever having blood cholesterol checked

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 43a. Blood cholesterol checked within the past 5 years among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	630	28,600	77.7	(73.0 - 82.4)	37,201	11,035,200	77.8	(75.1 - 80.4)
<b>Gender</b>								
Male	220	13,400	77.8	(70.3 - 85.2)	14,307	5,020,700	74.5	(70.2 - 78.9)
Female	410	15,200	77.6	(71.8 - 83.4)	22,894	6,014,600	80.7	(77.6 - 83.8)
<b>Age (years)</b>								
18-34	*	*	*	*	3,647	2,179,800	53.3	(46.7 - 59.9)
35-44	*	*	*	*	4,862	2,161,200	78.0	(72.8 - 83.2)
45-54	136	6,200	87.2	(80.7 - 93.8)	7,697	2,408,600	86.6	(82.4 - 90.8)
55-64	127	5,200	92.2	(86.9 - 97.4)	8,677	1,957,900	93.3	(91.4 - 95.3)
≥ 65	228	8,000	96.4	(93.5 - 99.4)	12,318	2,327,800	94.8	(93.1 - 96.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	602	25,900	77.8	(73.0 - 82.6)	34,960	7,486,600	83.4	(81.8 - 85.0)
Black Non-Hispanic	*	*	*	*	719	1,561,700	76.5	(67.1 - 85.9)
Hispanic	*	*	*	*	838	1,184,000	55.6	(45.6 - 65.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	320	13,300	75.6	(68.5 - 82.7)	15,215	3,612,500	68.3	(63.2 - 73.3)
Some College	170	8,400	77.3	(68.6 - 86.0)	9,673	2,524,200	75.4	(69.8 - 81.1)
≥ College Degree	140	6,900	82.6	(74.5 - 90.8)	12,236	4,806,500	88.1	(85.5 - 90.6)
<b>Income</b>								
≤ \$24,999	193	6,900	75.9	(66.4 - 85.5)	9,350	2,214,800	66.9	(60.1 - 73.8)
\$25,000-\$49,999	182	8,100	78.0	(69.5 - 86.6)	9,454	1,936,600	68.8	(61.5 - 76.1)
\$50,000-\$74,999	*	*	*	*	5,701	1,640,400	85.9	(82.4 - 89.5)
≥ \$75,000	95	6,200	84.7	(75.7 - 93.8)	8,435	3,872,400	89.2	(87.5 - 90.9)

<sup>1</sup> Defined as having had blood cholesterol checked within the past 5 years

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had blood cholesterol checked within the past 5 years (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 43b. Blood cholesterol checked within the past 5 years among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	630	74.5	(69.3 - 79.8)	37,201	77.3	(74.7 - 79.9)
<b>Gender</b>						
Male	220	76.0	(67.8 - 84.2)	14,307	74.8	(70.7 - 79.0)
Female	410	73.3	(66.5 - 80.0)	22,894	79.5	(76.3 - 82.7)
<b>Race/Ethnicity</b>						
White Non-Hispanic	602	74.3	(68.8 - 79.7)	34,960	82.6	(81.0 - 84.3)
Black Non-Hispanic	*	*	*	719	76.2	(66.9 - 85.5)
Hispanic	*	*	*	838	55.9	(46.1 - 65.8)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	320	71.6	(63.5 - 79.8)	15,215	67.8	(62.9 - 72.8)
Some College	170	74.0	(64.3 - 83.8)	9,673	74.7	(69.1 - 80.2)
≥ College Degree	140	80.9	(71.9 - 89.8)	12,236	87.5	(85.0 - 90.1)
<b>Income</b>						
≤ \$24,999	*	*	*	9,350	66.6	(59.8 - 73.4)
\$25,000-\$49,999	182	74.4	(64.6 - 84.1)	9,454	68.5	(61.4 - 75.6)
\$50,000-\$74,999	*	*	*	5,701	85.4	(81.7 - 89.0)
≥ \$75,000	95	83.4	(73.6 - 93.3)	8,435	88.5	(86.7 - 90.3)

<sup>1</sup> Defined as having had blood cholesterol checked within the past 5 years

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 44a. Cardiovascular disease among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	638	3,500	9.4	(7.3 - 11.6)	37,810	1,135,000	7.7	(6.5 - 8.8)
<b>Gender</b>								
Male	224	1,900	10.8	(7.1 - 14.5)	14,477	664,600	9.4	(7.5 - 11.4)
Female	414	1,600	8.2	(5.8 - 10.7)	23,333	470,400	6.1	(4.8 - 7.4)
<b>Age (years)</b>								
18-34	*	*	*	*	3,930	100,800	2.3	(0.0 - 4.8)
35-44	*	*	*	*	4,971	103,400	3.6	(1.5 - 5.7)
45-54	*	*	*	*	7,768	131,300	4.6	(2.9 - 6.4)
55-64	126	600	9.8	(4.3 - 15.3)	8,699	224,200	10.6	(8.3 - 12.9)
≥ 65	229	2,600	30.9	(24.3 - 37.5)	12,442	575,300	23.0	(19.9 - 26.1)
<b>Race/Ethnicity</b>								
White Non-Hispanic	610	3,300	9.7	(7.5 - 12.0)	35,522	772,200	8.3	(7.4 - 9.2)
Black Non-Hispanic	*	*	*	*	728	140,700	6.6	(3.3 - 9.9)
Hispanic	*	*	*	*	854	163,300	7.3	(2.0 - 12.5)
Other Non-Hispanic	*	*	*	*	706	58,800	5.2	(0.9 - 9.5)
<b>Education</b>								
≤ High School	321	2,000	11.5	(8.1 - 14.8)	15,486	590,800	10.7	(8.2 - 13.2)
Some College	172	700	5.9	(2.8 - 9.1)	9,849	227,100	6.4	(4.7 - 8.1)
≥ College Degree	145	900	9.8	(5.0 - 14.6)	12,393	314,600	5.6	(4.3 - 6.9)
<b>Income</b>								
≤ \$24,999	192	1,500	16.0	(10.3 - 21.7)	9,505	448,100	12.9	(9.1 - 16.7)
\$25,000-\$49,999	184	1,100	10.7	(6.5 - 14.8)	9,571	179,300	6.2	(4.8 - 7.6)
\$50,000-\$74,999	*	*	*	*	5,751	166,900	8.5	(5.5 - 11.4)
≥ \$75,000	*	*	*	*	8,538	183,500	4.1	(2.9 - 5.3)

<sup>1</sup> Defined as having ever been told by a health professional that respondent had a heart attack or angina or a stroke

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults with cardiovascular disease (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 44b. Cardiovascular disease among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	638	7.7	(5.8 - 9.6)	37,810	7.6	(6.4 - 8.7)
<b>Gender</b>						
Male	224	9.8	(6.3 - 13.3)	14,477	9.7	(7.8 - 11.7)
Female	414	5.9	(4.0 - 7.7)	23,333	5.6	(4.3 - 6.9)
<b>Race/Ethnicity</b>						
White Non-Hispanic	610	8.0	(6.0 - 10.0)	35,522	8.2	(7.2 - 9.1)
Black Non-Hispanic	*	*	*	728	6.8	(3.2 - 10.4)
Hispanic	*	*	*	854	7.0	(2.0 - 12.0)
Other Non-Hispanic	*	*	*	706	5.5	(0.9 - 10.0)
<b>Education</b>						
≤ High School	321	9.5	(6.5 - 12.4)	15,486	10.5	(8.0 - 13.0)
Some College	172	5.0	(1.9 - 8.1)	9,849	6.4	(4.5 - 8.2)
≥ College Degree	145	7.8	(3.8 - 11.9)	12,393	5.5	(4.2 - 6.9)
<b>Income</b>						
≤ \$24,999	192	13.5	(8.3 - 18.7)	9,505	12.8	(9.0 - 16.6)
\$25,000-\$49,999	184	8.6	(5.1 - 12.1)	9,571	6.0	(4.7 - 7.4)
\$50,000-\$74,999	*	*	*	5,751	8.5	(5.4 - 11.6)
≥ \$75,000	*	*	*	8,538	4.0	(2.7 - 5.2)

<sup>1</sup> Defined as having ever been told by a health professional that respondent had a heart attack or angina or a stroke

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 45a. Coronary heart disease among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	638	3,000	8.1	(6.2 - 10.1)	37,838	929,400	6.3	(5.3 - 7.3)
<b>Gender</b>								
Male	223	1,600	9.1	(5.8 - 12.4)	14,479	598,600	8.5	(6.6 - 10.4)
Female	415	1,400	7.3	(5.0 - 9.6)	23,359	330,800	4.3	(3.4 - 5.2)
<b>Age (years)</b>								
18-34	*	*	*	*	3,932	63,400	1.4	(0.0 - 3.6)
35-44	*	*	*	*	4,973	95,300	3.3	(1.2 - 5.4)
45-54	*	*	*	*	7,778	108,300	3.8	(2.1 - 5.6)
55-64	126	500	8.9	(3.6 - 14.1)	8,703	188,600	8.9	(6.7 - 11.1)
≥ 65	229	2,300	27.3	(21.0 - 33.7)	12,452	473,800	18.9	(16.1 - 21.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	611	2,900	8.5	(6.4 - 10.5)	35,552	636,700	6.9	(6.0 - 7.7)
Black Non-Hispanic	*	*	*	*	728	113,100	5.3	(2.3 - 8.4)
Hispanic	*	*	*	*	853	128,000	5.7	(1.1 - 10.3)
Other Non-Hispanic	*	*	*	*	705	51,500	4.6	(0.5 - 8.7)
<b>Education</b>								
≤ High School	321	1,800	10.4	(7.2 - 13.6)	15,485	474,500	8.6	(6.3 - 10.8)
Some College	172	500	4.6	(2.1 - 7.1)	9,860	193,000	5.5	(3.9 - 7.1)
≥ College Degree	145	700	7.9	(3.6 - 12.2)	12,411	259,700	4.6	(3.4 - 5.8)
<b>Income</b>								
≤ \$24,999	191	1,300	14.3	(8.9 - 19.7)	9,507	372,300	10.7	(7.3 - 14.1)
\$25,000-\$49,999	184	1,000	9.7	(5.7 - 13.8)	9,585	136,300	4.7	(3.8 - 5.7)
\$50,000-\$74,999	*	*	*	*	5,755	154,600	7.8	(4.9 - 10.7)
≥ \$75,000	*	*	*	*	8,541	151,800	3.4	(2.3 - 4.5)

<sup>1</sup> Defined as having ever been told by a health professional that respondent had a heart attack or angina

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults with coronary heart disease (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 45b. Coronary heart disease among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	638	6.5	(4.9 - 8.2)	37,838	6.3	(5.2 - 7.3)
<b>Gender</b>						
Male	223	8.0	(5.0 - 11.0)	14,479	8.8	(6.9 - 10.8)
Female	415	5.2	(3.5 - 7.0)	23,359	3.9	(3.0 - 4.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	611	6.8	(5.1 - 8.6)	35,552	6.8	(5.9 - 7.7)
Black Non-Hispanic	*	*	*	728	5.6	(2.2 - 9.0)
Hispanic	*	*	*	853	5.4	(1.1 - 9.6)
Other Non-Hispanic	*	*	*	705	4.8	(0.4 - 9.2)
<b>Education</b>						
≤ High School	321	8.6	(5.8 - 11.4)	15,485	8.5	(6.2 - 10.7)
Some College	172	3.6	(1.6 - 5.5)	9,860	5.5	(3.7 - 7.2)
≥ College Degree	145	6.4	(2.7 - 10.0)	12,411	4.7	(3.4 - 5.9)
<b>Income</b>						
≤ \$24,999	191	12.1	(7.2 - 17.0)	9,507	10.7	(7.2 - 14.1)
\$25,000-\$49,999	184	7.9	(4.5 - 11.3)	9,585	4.7	(3.6 - 5.7)
\$50,000-\$74,999	*	*	*	5,755	8.0	(4.8 - 11.1)
≥ \$75,000	*	*	*	8,541	3.3	(2.1 - 4.5)

<sup>1</sup> Defined as having ever been told by a health professional that respondent had a heart attack or angina

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 46a. Ever diagnosed with diabetes among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	643	3,400	9.0	(6.7 - 11.2)	38,174	1,382,700	9.3	(8.0 - 10.5)
<b>Gender</b>								
Male	227	1,500	8.3	(4.9 - 11.8)	14,626	530,400	7.5	(6.2 - 8.8)
Female	416	1,900	9.5	(6.6 - 12.5)	23,548	852,300	11.0	(8.9 - 13.0)
<b>Age (years)</b>								
18-34	*	*	*	*	3,941	150,200	3.4	(1.0 - 5.8)
35-44	*	*	*	*	4,987	165,100	5.7	(3.6 - 7.9)
45-54	138	500	6.7	(2.3 - 11.1)	7,830	279,000	9.8	(6.7 - 12.8)
55-64	128	800	14.1	(7.4 - 20.8)	8,771	332,900	15.5	(12.3 - 18.8)
≥ 65	232	1,600	19.3	(13.7 - 24.9)	12,645	455,600	17.9	(15.0 - 20.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	615	3,100	9.2	(6.9 - 11.5)	35,862	809,300	8.7	(7.8 - 9.5)
Black Non-Hispanic	*	*	*	*	739	261,000	12.1	(6.6 - 17.6)
Hispanic	*	*	*	*	862	227,300	10.1	(5.4 - 14.8)
Other Non-Hispanic	*	*	*	*	711	85,100	7.6	(2.3 - 12.8)
<b>Education</b>								
≤ High School	325	2,300	12.9	(9.0 - 16.9)	15,709	730,000	13.0	(10.5 - 15.6)
Some College	172	600	5.3	(2.3 - 8.2)	9,918	303,500	8.6	(5.9 - 11.2)
≥ College Degree	146	500	5.5	(1.8 - 9.2)	12,464	337,500	6.0	(4.8 - 7.2)
<b>Income</b>								
≤ \$24,999	194	1,400	15.5	(9.9 - 21.1)	9,652	543,400	15.5	(11.4 - 19.5)
\$25,000-\$49,999	185	1,300	12.1	(7.0 - 17.3)	9,667	291,000	10.0	(7.5 - 12.6)
\$50,000-\$74,999	*	*	*	*	5,777	151,000	7.6	(5.0 - 10.3)
≥ \$75,000	*	*	*	*	8,565	220,800	4.9	(3.8 - 6.1)

<sup>1</sup> Defined as having ever been told by a doctor that respondent had diabetes, excluding pre-diabetes and women with diabetes only when pregnant

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have diabetes (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 46b. Ever diagnosed with diabetes among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	643	7.7	(5.6 - 9.9)	38,174	9.0	(7.8 - 10.3)
<b>Gender</b>						
Male	227	7.3	(4.1 - 10.4)	14,626	7.5	(6.1 - 8.9)
Female	416	8.1	(5.2 - 11.0)	23,548	10.4	(8.4 - 12.4)
<b>Race/Ethnicity</b>						
White Non-Hispanic	615	8.0	(5.8 - 10.3)	35,862	8.4	(7.5 - 9.2)
Black Non-Hispanic	*	*	*	739	11.4	(6.0 - 16.7)
Hispanic	*	*	*	862	10.1	(5.4 - 14.9)
Other Non-Hispanic	*	*	*	711	7.7	(2.3 - 13.0)
<b>Education</b>						
≤ High School	325	11.9	(7.8 - 16.0)	15,709	12.6	(10.0 - 15.2)
Some College	172	3.9	(1.7 - 6.2)	9,918	8.3	(5.6 - 10.9)
≥ College Degree	146	4.7	(1.6 - 7.8)	12,464	5.9	(4.7 - 7.2)
<b>Income</b>						
≤ \$24,999	194	13.4	(8.1 - 18.7)	9,652	15.2	(11.1 - 19.2)
\$25,000-\$49,999	185	11.5	(6.0 - 17.0)	9,667	9.7	(7.1 - 12.2)
\$50,000-\$74,999	*	*	*	5,777	7.6	(4.9 - 10.2)
≥ \$75,000	*	*	*	8,565	4.7	(3.6 - 5.9)

<sup>1</sup> Defined as having ever been told by a doctor that respondent had diabetes, excluding pre-diabetes and women with diabetes only when pregnant

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 47a. Ever diagnosed with asthma (lifetime) among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	640	5,700	15.1	(10.9 - 19.3)	38,096	2,225,100	15.0	(13.2 - 16.8)
<b>Gender</b>								
Male	225	1,800	10.3	(3.1 - 17.6)	14,601	875,800	12.4	(10.1 - 14.7)
Female	415	3,800	19.3	(14.7 - 24.0)	23,495	1,349,300	17.4	(14.6 - 20.1)
<b>Age (years)</b>								
18-34	*	*	*	*	3,931	825,100	18.5	(13.8 - 23.3)
35-44	74	800	13.6	(6.0 - 21.2)	4,972	431,700	15.1	(11.2 - 19.1)
45-54	137	500	6.5	(2.6 - 10.3)	7,819	365,900	12.8	(9.7 - 16.0)
55-64	126	600	10.5	(5.4 - 15.7)	8,756	316,700	14.8	(11.9 - 17.7)
≥ 65	232	1,200	13.9	(9.0 - 18.7)	12,618	285,700	11.2	(8.6 - 13.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	612	4,600	13.6	(10.5 - 16.7)	35,783	1,365,700	14.6	(13.0 - 16.3)
Black Non-Hispanic	*	*	*	*	737	367,000	17.2	(10.2 - 24.2)
Hispanic	*	*	*	*	863	317,100	14.1	(9.0 - 19.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	324	2,400	13.4	(9.0 - 17.8)	15,675	831,600	14.9	(12.0 - 17.8)
Some College	*	*	*	*	9,893	587,500	16.6	(13.1 - 20.1)
≥ College Degree	145	1,600	18.8	(11.3 - 26.2)	12,444	739,000	13.2	(10.4 - 15.9)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,643	708,900	20.2	(14.9 - 25.5)
\$25,000-\$49,999	184	1,700	16.0	(9.7 - 22.3)	9,641	389,800	13.4	(10.0 - 16.9)
\$50,000-\$74,999	95	700	11.0	(4.1 - 18.0)	5,762	257,800	13.2	(10.0 - 16.5)
≥ \$75,000	96	800	11.3	(4.3 - 18.3)	8,562	577,900	12.8	(10.4 - 15.3)

<sup>1</sup> Defined as adults having ever been told by a health professional that they had asthma

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have lifetime asthma (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 47b. Ever diagnosed with asthma (lifetime) among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	640	15.6	(11.2 - 20.1)	38,096	15.2	(13.3 - 17.0)
<b>Gender</b>						
Male	225	10.8	(3.2 - 18.4)	14,601	12.6	(10.3 - 14.9)
Female	415	19.9	(14.7 - 25.2)	23,495	17.5	(14.8 - 20.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	612	14.2	(10.6 - 17.7)	35,783	14.9	(13.2 - 16.6)
Black Non-Hispanic	*	*	*	737	17.1	(10.1 - 24.1)
Hispanic	*	*	*	863	14.5	(9.3 - 19.7)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	324	13.0	(8.2 - 17.8)	15,675	15.1	(12.1 - 18.0)
Some College	*	*	*	9,893	16.9	(13.4 - 20.4)
≥ College Degree	145	20.0	(11.9 - 28.1)	12,444	13.3	(10.6 - 16.0)
<b>Income</b>						
≤ \$24,999	*	*	*	9,643	20.3	(15.0 - 25.6)
\$25,000-\$49,999	184	16.6	(9.5 - 23.7)	9,641	13.7	(10.2 - 17.2)
\$50,000-\$74,999	95	12.9	(4.7 - 21.0)	5,762	13.5	(10.0 - 17.0)
≥ \$75,000	96	11.7	(4.2 - 19.3)	8,562	13.0	(10.6 - 15.4)

<sup>1</sup> Defined as adults having ever been told by a health professional that they had asthma

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 48a. Current asthma among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	639	4,400	11.7	(8.8 - 14.7)	37,933	1,415,000	9.6	(8.1 - 11.0)
<b>Gender</b>								
Male	225	1,100	6.1	(2.7 - 9.5)	14,546	440,700	6.3	(4.8 - 7.7)
Female	414	3,300	16.7	(12.2 - 21.2)	23,387	974,300	12.6	(10.2 - 15.0)
<b>Age (years)</b>								
18-34	71	1,900	18.6	(9.8 - 27.3)	3,909	467,200	10.5	(6.8 - 14.3)
35-44	73	700	10.9	(4.2 - 17.7)	4,947	246,800	8.7	(5.8 - 11.6)
45-54	137	400	6.0	(2.3 - 9.7)	7,787	240,000	8.4	(6.1 - 10.8)
55-64	126	400	7.7	(3.2 - 12.2)	8,726	231,100	10.8	(8.1 - 13.5)
≥ 65	232	1,000	11.5	(7.1 - 16.0)	12,564	229,900	9.1	(6.5 - 11.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	611	4,000	11.7	(8.7 - 14.7)	35,629	888,400	9.6	(8.2 - 11.0)
Black Non-Hispanic	*	*	*	*	736	246,900	11.6	(5.7 - 17.5)
Hispanic	*	*	*	*	859	186,800	8.3	(4.4 - 12.2)
Other Non-Hispanic	*	*	*	*	709	93,000	8.3	(1.4 - 15.2)
<b>Education</b>								
≤ High School	323	2,000	11.0	(6.9 - 15.1)	15,594	595,100	10.7	(8.2 - 13.2)
Some College	171	800	7.2	(2.7 - 11.8)	9,861	322,000	9.1	(7.0 - 11.3)
≥ College Degree	145	1,600	18.8	(11.3 - 26.2)	12,394	430,900	7.7	(5.8 - 9.7)
<b>Income</b>								
≤ \$24,999	193	1,000	10.9	(6.1 - 15.7)	9,600	489,200	14.0	(9.7 - 18.3)
\$25,000-\$49,999	184	1,400	13.5	(7.5 - 19.6)	9,590	268,400	9.3	(6.2 - 12.4)
\$50,000-\$74,999	*	*	*	*	5,741	143,400	7.4	(5.1 - 9.6)
≥ \$75,000	96	800	10.7	(3.8 - 17.6)	8,541	339,200	7.6	(5.8 - 9.4)

<sup>1</sup> Defined as adults having ever been told by a health professional that they had asthma and still have asthma

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who currently have asthma (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 48b. Current asthma among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	639	12.3	(9.0 - 15.6)	37,933	9.7	(8.2 - 11.2)
<b>Gender</b>						
Male	225	6.3	(2.6 - 10.0)	14,546	6.4	(4.9 - 7.9)
Female	414	17.6	(12.5 - 22.7)	23,387	12.7	(10.2 - 15.1)
<b>Race/Ethnicity</b>						
White Non-Hispanic	611	12.4	(9.0 - 15.7)	35,629	9.7	(8.3 - 11.2)
Black Non-Hispanic	*	*	*	736	11.5	(5.6 - 17.3)
Hispanic	*	*	*	859	8.6	(4.5 - 12.6)
Other Non-Hispanic	*	*	*	709	8.3	(1.3 - 15.3)
<b>Education</b>						
≤ High School	323	10.8	(6.3 - 15.4)	15,594	10.8	(8.3 - 13.3)
Some College	171	8.2	(3.0 - 13.4)	9,861	9.4	(7.2 - 11.6)
≥ College Degree	145	20.0	(11.9 - 28.1)	12,394	7.8	(5.8 - 9.7)
<b>Income</b>						
≤ \$24,999	193	10.8	(5.5 - 16.2)	9,600	14.1	(9.7 - 18.4)
\$25,000-\$49,999	184	14.4	(7.5 - 21.2)	9,590	9.5	(6.3 - 12.7)
\$50,000-\$74,999	*	*	*	5,741	7.5	(5.0 - 10.1)
≥ \$75,000	96	11.3	(3.8 - 18.8)	8,541	7.7	(5.9 - 9.5)

<sup>1</sup> Defined as adults having ever been told by a health professional that they had asthma and still have asthma

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 49a. Chronic joint symptoms among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	642	16,900	45.0	(40.0 - 49.9)	38,099	6,377,100	42.9	(40.5 - 45.3)
<b>Gender</b>								
Male	226	7,500	42.2	(34.1 - 50.4)	14,595	3,033,600	42.8	(39.0 - 46.6)
Female	416	9,400	47.4	(41.5 - 53.3)	23,504	3,343,500	43.0	(40.0 - 46.0)
<b>Age (years)</b>								
18-34	*	*	*	*	3,933	1,261,300	28.3	(22.6 - 34.0)
35-44	*	*	*	*	4,984	1,110,900	38.6	(33.1 - 44.0)
45-54	138	3,800	53.6	(43.8 - 63.3)	7,817	1,456,500	51.2	(46.4 - 55.9)
55-64	128	3,400	60.0	(50.5 - 69.6)	8,751	1,167,400	54.6	(50.4 - 58.8)
≥ 65	231	4,900	58.7	(51.6 - 65.7)	12,614	1,381,000	54.3	(50.5 - 58.1)
<b>Race/Ethnicity</b>								
White Non-Hispanic	614	15,700	46.2	(41.4 - 51.1)	35,785	4,267,600	45.8	(43.8 - 47.7)
Black Non-Hispanic	*	*	*	*	740	890,500	41.3	(32.5 - 50.0)
Hispanic	*	*	*	*	860	827,500	36.7	(27.7 - 45.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	324	9,100	51.0	(43.8 - 58.1)	15,676	2,450,000	43.8	(39.4 - 48.1)
Some College	172	4,300	39.5	(30.2 - 48.8)	9,900	1,598,300	45.1	(40.1 - 50.1)
≥ College Degree	146	3,500	39.6	(30.0 - 49.2)	12,440	2,313,100	41.1	(37.8 - 44.5)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,641	1,483,000	42.2	(36.3 - 48.1)
\$25,000-\$49,999	184	4,300	39.8	(31.2 - 48.4)	9,647	1,332,100	45.9	(39.8 - 52.1)
\$50,000-\$74,999	*	*	*	*	5,768	894,200	45.3	(39.5 - 51.1)
≥ \$75,000	*	*	*	*	8,549	1,874,200	41.7	(38.4 - 45.1)

<sup>1</sup> Defined as adults having symptoms of pain, aching or stiffness in or around a joint during the past 30 days and began more than 3 months ago

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have chronic joint symptoms (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 49b. Chronic joint symptoms among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	642	41.8	(36.5 - 47.1)	38,099	42.1	(39.7 - 44.5)
<b>Gender</b>						
Male	226	40.4	(31.9 - 48.9)	14,595	42.6	(38.8 - 46.3)
Female	416	43.0	(36.5 - 49.5)	23,504	41.6	(38.6 - 44.7)
<b>Race/Ethnicity</b>						
White Non-Hispanic	614	43.0	(37.7 - 48.2)	35,785	44.7	(42.7 - 46.6)
Black Non-Hispanic	*	*	*	740	40.2	(31.5 - 48.9)
Hispanic	*	*	*	860	36.6	(27.8 - 45.4)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	324	47.6	(39.7 - 55.6)	15,676	43.1	(38.8 - 47.5)
Some College	172	36.5	(26.9 - 46.0)	9,900	43.9	(39.0 - 48.8)
≥ College Degree	146	37.5	(27.7 - 47.3)	12,440	40.3	(37.0 - 43.7)
<b>Income</b>						
≤ \$24,999	*	*	*	9,641	41.6	(35.7 - 47.4)
\$25,000-\$49,999	184	35.3	(26.5 - 44.1)	9,647	45.0	(39.0 - 51.0)
\$50,000-\$74,999	*	*	*	5,768	44.3	(38.4 - 50.2)
≥ \$75,000	*	*	*	8,549	40.8	(37.4 - 44.2)

<sup>1</sup> Defined as adults having symptoms of pain, aching or stiffness in or around a joint during the past 30 days and began more than 3 months ago

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 50a. Arthritis among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	643	13,600	36.1	(31.6 - 40.6)	38,013	4,102,800	27.7	(25.9 - 29.6)
<b>Gender</b>								
Male	227	5,500	31.0	(24.0 - 38.0)	14,555	1,642,700	23.3	(20.7 - 25.8)
Female	416	8,100	40.7	(35.1 - 46.3)	23,458	2,460,100	31.8	(29.2 - 34.4)
<b>Age (years)</b>								
18-34	*	*	*	*	3,927	312,100	7.0	(4.9 - 9.1)
35-44	*	*	*	*	4,972	412,500	14.5	(11.3 - 17.6)
45-54	138	2,600	35.6	(26.3 - 44.9)	7,789	869,100	30.7	(26.6 - 34.8)
55-64	128	3,400	60.3	(50.7 - 69.9)	8,729	999,300	47.0	(42.9 - 51.1)
≥ 65	232	5,600	66.8	(60.0 - 73.7)	12,596	1,509,800	59.6	(55.8 - 63.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	615	13,000	38.1	(33.6 - 42.7)	35,710	2,967,700	31.9	(30.3 - 33.6)
Black Non-Hispanic	*	*	*	*	737	588,100	27.3	(20.2 - 34.4)
Hispanic	*	*	*	*	856	380,700	17.1	(11.8 - 22.5)
Other Non-Hispanic	*	*	*	*	710	166,300	14.8	(7.8 - 21.8)
<b>Education</b>								
≤ High School	325	7,800	43.6	(36.7 - 50.4)	15,652	1,742,700	31.3	(27.8 - 34.8)
Some College	172	3,600	32.5	(24.2 - 40.8)	9,874	902,000	25.5	(22.3 - 28.6)
≥ College Degree	146	2,200	25.5	(17.9 - 33.0)	12,407	1,437,800	25.6	(22.9 - 28.4)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,618	1,112,500	32.0	(27.1 - 36.9)
\$25,000-\$49,999	185	4,000	37.2	(28.9 - 45.6)	9,624	821,900	28.4	(24.2 - 32.6)
\$50,000-\$74,999	*	*	*	*	5,753	542,500	27.5	(23.2 - 31.9)
≥ \$75,000	97	2,100	28.4	(18.5 - 38.3)	8,536	1,012,800	22.6	(19.8 - 25.3)

<sup>1</sup> Defined as adults having ever been told by a health professional that they have some form of arthritis, rheumatoid arthritis, gout, lupus or fibromyalgia (excluding back or neck)

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have ever been told by a health professional they have some form of arthritis (excluding back or neck) (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 50b. Arthritis among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	643	31.3	(26.8 - 35.8)	38,013	26.6	(24.8 - 28.5)
<b>Gender</b>						
Male	227	28.4	(21.4 - 35.5)	14,555	23.5	(20.9 - 26.1)
Female	416	33.8	(28.2 - 39.5)	23,458	29.5	(27.0 - 32.1)
<b>Race/Ethnicity</b>						
White Non-Hispanic	615	33.2	(28.6 - 37.9)	35,710	30.4	(28.7 - 32.0)
Black Non-Hispanic	*	*	*	737	26.5	(19.3 - 33.6)
Hispanic	*	*	*	856	17.2	(11.8 - 22.6)
Other Non-Hispanic	*	*	*	710	14.9	(7.7 - 22.0)
<b>Education</b>						
≤ High School	325	38.9	(31.5 - 46.2)	15,652	30.4	(26.9 - 33.8)
Some College	172	27.4	(19.6 - 35.1)	9,874	24.1	(21.0 - 27.1)
≥ College Degree	146	21.9	(14.8 - 29.0)	12,407	24.7	(21.9 - 27.5)
<b>Income</b>						
≤ \$24,999	194	36.2	(26.3 - 46.2)	9,618	31.2	(26.3 - 36.1)
\$25,000-\$49,999	185	32.2	(23.7 - 40.7)	9,624	27.3	(23.2 - 31.3)
\$50,000-\$74,999	*	*	*	5,753	26.3	(21.9 - 30.7)
≥ \$75,000	97	24.7	(15.2 - 34.2)	8,536	21.6	(18.8 - 24.3)

<sup>1</sup> Defined as adults having ever been told by a health professional that they have some form of arthritis, rheumatoid arthritis, gout, lupus or fibromyalgia (excluding back or neck)

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 51a. Activity limitations because of physical, mental or emotional problems among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	638	8,300	22.3	(18.5 - 26.1)	37,998	3,053,000	20.6	(18.7 - 22.5)
<b>Gender</b>								
Male	223	3,700	21.3	(15.2 - 27.5)	14,566	1,318,400	18.6	(15.7 - 21.5)
Female	415	4,600	23.1	(18.5 - 27.7)	23,432	1,734,600	22.4	(19.8 - 25.0)
<b>Age (years)</b>								
18-34	71	1,400	13.7	(5.4 - 22.0)	3,929	580,000	13.0	(8.2 - 17.9)
35-44	74	1,000	15.9	(6.9 - 25.0)	4,973	418,600	14.5	(11.0 - 18.1)
45-54	136	1,600	22.0	(14.0 - 30.1)	7,812	620,200	21.7	(18.3 - 25.2)
55-64	127	2,000	35.0	(25.9 - 44.1)	8,739	651,600	30.5	(26.5 - 34.6)
≥ 65	230	2,400	29.1	(22.9 - 35.3)	12,545	782,700	31.1	(27.7 - 34.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	610	7,600	22.4	(18.6 - 26.2)	35,688	2,037,200	21.9	(20.3 - 23.5)
Black Non-Hispanic	*	*	*	*	735	429,700	20.1	(13.5 - 26.7)
Hispanic	*	*	*	*	863	448,200	19.8	(11.9 - 27.8)
Other Non-Hispanic	*	*	*	*	712	137,900	12.3	(5.2 - 19.3)
<b>Education</b>								
≤ High School	323	4,300	24.1	(18.4 - 29.7)	15,626	1,359,600	24.4	(20.6 - 28.1)
Some College	170	2,600	24.3	(16.7 - 32.0)	9,875	700,800	19.8	(16.5 - 23.2)
≥ College Degree	145	1,400	16.1	(9.7 - 22.4)	12,415	924,800	16.5	(14.2 - 18.7)
<b>Income</b>								
≤ \$24,999	192	2,600	28.4	(20.1 - 36.7)	9,603	1,055,800	30.1	(24.7 - 35.5)
\$25,000-\$49,999	184	2,100	20.1	(13.9 - 26.3)	9,628	630,900	21.8	(16.6 - 27.0)
\$50,000-\$74,999	95	1,100	16.9	(9.1 - 24.6)	5,760	408,100	20.7	(16.1 - 25.2)
≥ \$75,000	97	1,400	18.8	(10.1 - 27.5)	8,544	583,300	13.0	(11.1 - 14.9)

<sup>1</sup> Defined as activity limitations because of physical, mental or emotional problems

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who are limited in any activities because of physical, mental or emotional problems (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 51b. Activity limitations because of physical, mental or emotional problems among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	638	20.7	(16.7 - 24.6)	37,998	20.1	(18.2 - 22.0)
<b>Gender</b>						
Male	223	20.7	(14.2 - 27.1)	14,566	18.6	(15.8 - 21.4)
Female	415	20.7	(15.9 - 25.4)	23,432	21.5	(18.9 - 24.1)
<b>Race/Ethnicity</b>						
White Non-Hispanic	610	20.7	(16.7 - 24.7)	35,688	21.3	(19.7 - 22.9)
Black Non-Hispanic	*	*	*	735	19.9	(13.2 - 26.5)
Hispanic	*	*	*	863	19.1	(11.5 - 26.8)
Other Non-Hispanic	*	*	*	712	12.4	(5.3 - 19.6)
<b>Education</b>						
≤ High School	323	23.0	(16.8 - 29.1)	15,626	23.8	(20.1 - 27.4)
Some College	170	22.1	(14.4 - 29.7)	9,875	19.3	(16.0 - 22.6)
≥ College Degree	145	14.5	(8.4 - 20.7)	12,415	16.1	(13.9 - 18.3)
<b>Income</b>						
≤ \$24,999	192	27.2	(18.1 - 36.3)	9,603	29.7	(24.3 - 35.2)
\$25,000-\$49,999	184	17.1	(11.0 - 23.2)	9,628	21.2	(16.1 - 26.3)
\$50,000-\$74,999	95	15.8	(8.0 - 23.6)	5,760	20.1	(15.6 - 24.6)
≥ \$75,000	97	18.2	(9.4 - 27.1)	8,544	12.7	(10.8 - 14.6)

<sup>1</sup> Defined as activity limitations because of physical, mental or emotional problems

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 52a. Have health problem that requires the use of special equipment among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	641	3,000	8.0	(6.0 - 10.0)	38,171	1,169,400	7.9	(6.9 - 8.9)
<b>Gender</b>								
Male	226	1,200	6.6	(3.6 - 9.5)	14,617	460,700	6.5	(5.3 - 7.7)
Female	415	1,800	9.2	(6.5 - 12.0)	23,554	708,700	9.1	(7.6 - 10.7)
<b>Age (years)</b>								
18-34	*	*	*	*	3,940	68,800	1.5	(0.7 - 2.4)
35-44	*	*	*	*	4,989	122,500	4.3	(2.4 - 6.1)
45-54	*	*	*	*	7,836	182,500	6.4	(4.7 - 8.1)
55-64	127	600	10.8	(5.4 - 16.3)	8,773	219,600	10.3	(7.8 - 12.8)
≥ 65	231	1,700	20.6	(15.0 - 26.1)	12,633	576,100	22.7	(19.0 - 26.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	613	2,600	7.7	(5.7 - 9.6)	35,853	802,000	8.6	(7.6 - 9.6)
Black Non-Hispanic	*	*	*	*	739	235,800	10.9	(6.4 - 15.5)
Hispanic	*	*	*	*	864	104,100	4.6	(2.1 - 7.1)
Other Non-Hispanic	*	*	*	*	715	27,600	2.5	(0.8 - 4.1)
<b>Education</b>								
≤ High School	324	1,900	10.7	(7.4 - 14.0)	15,711	604,200	10.8	(8.7 - 12.8)
Some College	171	400	3.9	(1.3 - 6.5)	9,914	260,000	7.3	(5.5 - 9.1)
≥ College Degree	146	700	7.5	(3.2 - 11.9)	12,463	298,900	5.3	(4.1 - 6.5)
<b>Income</b>								
≤ \$24,999	194	1,400	14.9	(9.3 - 20.5)	9,656	430,500	12.2	(9.6 - 14.8)
\$25,000-\$49,999	184	600	5.8	(2.9 - 8.7)	9,663	238,900	8.2	(5.9 - 10.5)
\$50,000-\$74,999	*	*	*	*	5,775	103,000	5.2	(3.6 - 6.8)
≥ \$75,000	*	*	*	*	8,566	136,000	3.0	(2.1 - 4.0)

<sup>1</sup> Defined as having any health problem that requires the use of special equipment such as a cane, a wheelchair, a special bed, or a special telephone

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have any health problem that requires the use of special equipment (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 52b. Have health problem that requires the use of special equipment among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	641	6.8	(4.8 - 8.7)	38,171	7.6	(6.6 - 8.6)
<b>Gender</b>						
Male	226	6.2	(3.2 - 9.1)	14,617	6.8	(5.5 - 8.0)
Female	415	7.3	(4.8 - 9.9)	23,554	8.4	(6.9 - 9.9)
<b>Race/Ethnicity</b>						
White Non-Hispanic	613	6.4	(4.6 - 8.2)	35,853	8.3	(7.3 - 9.3)
Black Non-Hispanic	*	*	*	739	10.5	(6.1 - 15.0)
Hispanic	*	*	*	864	4.5	(2.1 - 6.9)
Other Non-Hispanic	*	*	*	715	2.2	(0.7 - 3.8)
<b>Education</b>						
≤ High School	324	9.0	(5.9 - 12.1)	15,711	10.3	(8.3 - 12.3)
Some College	171	3.4	(0.7 - 6.2)	9,914	7.1	(5.2 - 8.9)
≥ College Degree	146	6.8	(2.5 - 11.0)	12,463	5.3	(4.1 - 6.6)
<b>Income</b>						
≤ \$24,999	194	13.1	(7.5 - 18.7)	9,656	11.9	(9.4 - 14.5)
\$25,000-\$49,999	184	4.5	(2.2 - 6.9)	9,663	7.7	(5.5 - 10.0)
\$50,000-\$74,999	*	*	*	5,775	5.1	(3.4 - 6.8)
≥ \$75,000	*	*	*	8,566	2.9	(2.0 - 3.9)

<sup>1</sup> Defined as having any health problem that requires the use of special equipment such as a cane, a wheelchair, a special bed, or a special telephone

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 53a. Disability among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	638	9,300	24.8	(20.9 - 28.7)	38,016	3,394,200	22.9	(20.9 - 24.9)
<b>Gender</b>								
Male	223	4,200	24.1	(17.7 - 30.6)	14,568	1,469,700	20.7	(17.8 - 23.7)
Female	415	5,000	25.4	(20.7 - 30.1)	23,448	1,924,600	24.8	(22.1 - 27.5)
<b>Age (years)</b>								
18-34	71	1,400	13.7	(5.4 - 22.0)	3,931	590,300	13.3	(8.4 - 18.1)
35-44	74	1,000	15.9	(6.9 - 25.0)	4,974	447,900	15.6	(11.9 - 19.3)
45-54	136	1,700	24.2	(16.0 - 32.5)	7,811	643,900	22.6	(19.1 - 26.0)
55-64	127	2,100	37.6	(28.3 - 47.0)	8,740	702,100	33.0	(28.9 - 37.0)
≥ 65	230	3,100	36.8	(30.1 - 43.5)	12,560	1,010,000	39.9	(36.1 - 43.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	610	8,500	25.1	(21.1 - 29.0)	35,703	2,253,400	24.2	(22.6 - 25.9)
Black Non-Hispanic	*	*	*	*	736	511,700	23.8	(16.8 - 30.8)
Hispanic	*	*	*	*	864	484,300	21.4	(13.4 - 29.5)
Other Non-Hispanic	*	*	*	*	713	144,800	12.9	(5.8 - 19.9)
<b>Education</b>								
≤ High School	323	5,100	28.4	(22.5 - 34.3)	15,638	1,544,200	27.6	(23.7 - 31.4)
Some College	170	2,700	24.8	(17.1 - 32.4)	9,876	755,200	21.4	(17.9 - 24.8)
≥ College Degree	145	1,500	17.4	(10.8 - 24.0)	12,419	1,021,300	18.2	(15.9 - 20.5)
<b>Income</b>								
≤ \$24,999	192	3,100	34.1	(25.0 - 43.3)	9,613	1,179,100	33.6	(28.0 - 39.2)
\$25,000-\$49,999	184	2,300	21.1	(14.8 - 27.5)	9,629	699,000	24.2	(18.9 - 29.4)
\$50,000-\$74,999	95	1,300	19.6	(11.3 - 27.8)	5,757	430,100	21.8	(17.2 - 26.4)
≥ \$75,000	97	1,400	18.8	(10.1 - 27.5)	8,546	632,700	14.1	(12.1 - 16.1)

<sup>1</sup> Defined as having activity limitations due to physical, mental or emotional problems or have health problems that require the use of special equipment

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults with a disability (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 53b. Disability among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	638	22.7	(18.6 - 26.7)	38,016	22.3	(20.4 - 24.3)
<b>Gender</b>						
Male	223	23.1	(16.4 - 29.7)	14,568	20.9	(18.0 - 23.8)
Female	415	22.3	(17.5 - 27.2)	23,448	23.6	(21.0 - 26.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	610	22.9	(18.8 - 27.0)	35,703	23.6	(21.9 - 25.3)
Black Non-Hispanic	*	*	*	736	23.5	(16.5 - 30.5)
Hispanic	*	*	*	864	20.6	(12.9 - 28.3)
Other Non-Hispanic	*	*	*	713	12.9	(5.8 - 20.1)
<b>Education</b>						
≤ High School	323	26.7	(20.3 - 33.1)	15,638	26.8	(23.0 - 30.6)
Some College	170	22.4	(14.7 - 30.0)	9,876	20.9	(17.4 - 24.3)
≥ College Degree	145	15.6	(9.2 - 22.0)	12,419	17.9	(15.5 - 20.3)
<b>Income</b>						
≤ \$24,999	192	32.0	(22.3 - 41.8)	9,613	33.2	(27.6 - 38.8)
\$25,000-\$49,999	184	17.9	(11.7 - 24.2)	9,629	23.5	(18.3 - 28.6)
\$50,000-\$74,999	95	18.1	(9.9 - 26.3)	5,757	21.2	(16.6 - 25.8)
≥ \$75,000	97	18.2	(9.4 - 27.1)	8,546	13.8	(11.7 - 15.8)

<sup>1</sup> Defined as having activity limitations due to physical, mental or emotional problems or have health problems that require the use of special equipment

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 54a. Current smoking among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	640	8,200	21.9	(17.6 - 26.2)	38,029	2,483,500	16.8	(15.1 - 18.6)
<b>Gender</b>								
Male	225	3,500	19.7	(13.1 - 26.3)	14,559	1,356,400	19.3	(16.3 - 22.2)
Female	415	4,800	23.9	(18.4 - 29.5)	23,470	1,127,100	14.6	(12.7 - 16.6)
<b>Age (years)</b>								
18-34	*	*	*	*	3,929	993,500	22.3	(17.5 - 27.1)
35-44	74	1,100	17.4	(7.9 - 27.0)	4,977	421,100	14.8	(11.6 - 17.9)
45-54	138	2,200	30.7	(21.6 - 39.7)	7,807	508,800	18.3	(15.4 - 21.3)
55-64	128	1,000	18.0	(10.9 - 25.1)	8,744	314,500	14.7	(12.1 - 17.3)
≥ 65	230	800	9.7	(5.6 - 13.9)	12,572	245,600	9.7	(7.2 - 12.3)
<b>Race/Ethnicity</b>								
White Non-Hispanic	612	7,200	21.2	(17.0 - 25.4)	35,729	1,615,500	17.3	(15.9 - 18.8)
Black Non-Hispanic	*	*	*	*	735	284,100	13.6	(8.9 - 18.2)
Hispanic	*	*	*	*	858	350,900	15.8	(9.6 - 21.9)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	322	5,000	28.4	(21.6 - 35.2)	15,654	1,237,300	22.4	(19.1 - 25.6)
Some College	172	2,000	18.6	(11.3 - 25.9)	9,875	670,600	19.1	(15.5 - 22.8)
≥ College Degree	146	1,100	13.0	(6.0 - 20.1)	12,421	563,300	10.0	(7.5 - 12.5)
<b>Income</b>								
≤ \$24,999	192	2,600	28.6	(19.2 - 37.9)	9,626	835,100	24.1	(19.1 - 29.1)
\$25,000-\$49,999	185	2,800	26.0	(17.5 - 34.5)	9,629	558,100	19.4	(15.6 - 23.2)
\$50,000-\$74,999	95	1,200	18.6	(8.6 - 28.6)	5,756	338,100	17.3	(13.0 - 21.7)
≥ \$75,000	*	*	*	*	8,549	468,700	10.4	(8.6 - 12.3)

<sup>1</sup> Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day or some days

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adult current smokers (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 54b. Current smoking among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	640	22.9	(18.1 - 27.6)	38,029	17.0	(15.3 - 18.8)
<b>Gender</b>						
Male	225	19.2	(12.2 - 26.1)	14,559	19.2	(16.3 - 22.1)
Female	415	26.1	(19.8 - 32.5)	23,470	15.0	(13.0 - 17.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	612	22.0	(17.3 - 26.7)	35,729	17.7	(16.2 - 19.2)
Black Non-Hispanic	*	*	*	735	13.5	(8.8 - 18.2)
Hispanic	*	*	*	858	15.9	(9.8 - 22.0)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	322	30.4	(22.7 - 38.1)	15,654	22.8	(19.5 - 26.0)
Some College	172	18.5	(10.9 - 26.2)	9,875	19.4	(15.8 - 22.9)
≥ College Degree	146	14.2	(6.3 - 22.2)	12,421	10.0	(7.6 - 12.5)
<b>Income</b>						
≤ \$24,999	*	*	*	9,626	24.1	(19.2 - 29.1)
\$25,000-\$49,999	185	27.8	(18.3 - 37.2)	9,629	19.9	(16.0 - 23.7)
\$50,000-\$74,999	*	*	*	5,756	17.8	(13.3 - 22.3)
≥ \$75,000	*	*	*	8,549	10.6	(8.6 - 12.5)

<sup>1</sup> Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day or some days

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 55a. Everyday smoking among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	640	6,200	16.7	(12.9 - 20.5)	38,029	1,626,800	11.0	(9.8 - 12.2)
<b>Gender</b>								
Male	225	2,700	15.5	(9.6 - 21.3)	14,559	848,400	12.0	(10.0 - 14.1)
Female	415	3,500	17.8	(12.9 - 22.7)	23,470	778,400	10.1	(8.7 - 11.5)
<b>Age (years)</b>								
18-34	*	*	*	*	3,929	559,800	12.6	(9.6 - 15.6)
35-44	74	1,000	15.9	(6.6 - 25.3)	4,977	315,200	11.0	(8.4 - 13.7)
45-54	138	1,900	26.5	(17.7 - 35.2)	7,807	377,000	13.6	(11.1 - 16.0)
55-64	128	800	13.9	(7.8 - 20.1)	8,744	203,000	9.5	(7.5 - 11.5)
≥ 65	230	600	7.6	(3.8 - 11.5)	12,572	171,700	6.8	(4.8 - 8.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	612	5,700	16.7	(12.9 - 20.5)	35,729	1,158,100	12.4	(11.2 - 13.7)
Black Non-Hispanic	*	*	*	*	735	197,700	9.4	(5.6 - 13.3)
Hispanic	*	*	*	*	858	176,200	7.9	(4.2 - 11.6)
Other Non-Hispanic	*	*	*	*	707	94,900	8.5	(3.3 - 13.7)
<b>Education</b>								
≤ High School	322	3,700	21.1	(15.0 - 27.3)	15,654	881,900	15.9	(13.4 - 18.5)
Some College	172	2,000	18.5	(11.2 - 25.7)	9,875	410,200	11.7	(9.6 - 13.8)
≥ College Degree	*	*	*	*	12,421	323,500	5.8	(4.3 - 7.2)
<b>Income</b>								
≤ \$24,999	192	2,100	23.0	(14.6 - 31.4)	9,626	452,800	13.1	(10.3 - 15.8)
\$25,000-\$49,999	185	2,100	19.9	(12.2 - 27.5)	9,629	425,900	14.8	(11.9 - 17.7)
\$50,000-\$74,999	95	800	12.8	(4.7 - 20.9)	5,756	222,700	11.4	(8.5 - 14.4)
≥ \$75,000	*	*	*	*	8,549	320,200	7.1	(5.5 - 8.8)

<sup>1</sup> Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have smoked at least 100 cigarettes in lifetime and currently smoking everyday (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 55b. Everyday smoking among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	640	17.4	(13.2 - 21.6)	38,029	11.3	(10.0 - 12.5)
<b>Gender</b>						
Male	225	15.2	(9.0 - 21.3)	14,559	12.1	(10.1 - 14.1)
Female	415	19.4	(13.7 - 25.0)	23,470	10.5	(9.0 - 12.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	612	17.2	(13.1 - 21.4)	35,729	12.8	(11.5 - 14.1)
Black Non-Hispanic	*	*	*	735	9.4	(5.5 - 13.4)
Hispanic	*	*	*	858	8.0	(4.3 - 11.7)
Other Non-Hispanic	*	*	*	707	8.5	(3.2 - 13.8)
<b>Education</b>						
≤ High School	322	22.9	(15.9 - 29.9)	15,654	16.4	(13.8 - 19.0)
Some College	172	18.4	(10.7 - 26.1)	9,875	12.1	(9.9 - 14.3)
≥ College Degree	*	*	*	12,421	5.8	(4.3 - 7.3)
<b>Income</b>						
≤ \$24,999	192	25.1	(15.2 - 34.9)	9,626	13.4	(10.6 - 16.2)
\$25,000-\$49,999	185	21.6	(13.0 - 30.2)	9,629	15.3	(12.3 - 18.3)
\$50,000-\$74,999	95	12.6	(4.1 - 21.1)	5,756	11.7	(8.7 - 14.8)
≥ \$75,000	*	*	*	8,549	7.2	(5.5 - 8.9)

<sup>1</sup> Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 56a. Adults living in homes in which smoking is prohibited<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	636	27,400	73.5	(68.6 - 78.3)	38,045	11,997,100	80.8	(79.0 - 82.6)
<b>Gender</b>								
Male	224	12,700	72.5	(64.1 - 80.9)	14,577	5,634,400	79.4	(76.6 - 82.2)
Female	412	14,700	74.3	(69.2 - 79.5)	23,468	6,362,800	82.1	(79.9 - 84.3)
<b>Age (years)</b>								
18-34	*	*	*	*	3,934	3,655,200	82.1	(77.8 - 86.4)
35-44	*	*	*	*	4,985	2,479,700	86.1	(82.5 - 89.7)
45-54	135	5,100	72.5	(63.9 - 81.0)	7,808	2,194,400	77.2	(73.5 - 80.8)
55-64	128	4,500	78.7	(71.2 - 86.2)	8,753	1,732,500	81.0	(78.3 - 83.7)
≥ 65	228	6,200	74.4	(68.3 - 80.5)	12,565	1,935,400	76.5	(73.4 - 79.5)
<b>Race/Ethnicity</b>								
White Non-Hispanic	608	25,200	74.8	(70.6 - 79.1)	35,737	7,426,900	79.8	(78.3 - 81.2)
Black Non-Hispanic	*	*	*	*	738	1,692,100	78.6	(71.9 - 85.3)
Hispanic	*	*	*	*	863	2,003,900	88.7	(84.1 - 93.3)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	321	11,600	65.2	(58.3 - 72.0)	15,647	4,307,200	77.1	(74.0 - 80.2)
Some College	*	*	*	*	9,883	2,795,400	78.9	(75.0 - 82.8)
≥ College Degree	143	7,800	91.3	(87.1 - 95.5)	12,437	4,817,300	85.7	(83.3 - 88.2)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,608	2,627,700	74.9	(70.0 - 79.7)
\$25,000-\$49,999	184	7,900	74.9	(67.1 - 82.6)	9,640	2,314,500	79.8	(76.0 - 83.5)
\$50,000-\$74,999	95	5,200	81.4	(71.8 - 91.1)	5,766	1,551,300	78.7	(74.0 - 83.4)
≥ \$75,000	96	6,900	92.9	(87.7 - 98.0)	8,556	3,883,200	86.5	(84.3 - 88.7)

<sup>1</sup> Defined as having a rule that prohibits smoking anywhere inside the home

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who live in a home with a rule that prohibits smoking anywhere inside the home (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 56b. Adults living in homes in which smoking is prohibited<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	636	73.1	(67.9 - 78.3)	38,045	80.9	(79.2 - 82.7)
<b>Gender</b>						
Male	224	72.3	(63.4 - 81.2)	14,577	79.4	(76.7 - 82.2)
Female	412	73.8	(67.9 - 79.7)	23,468	82.3	(80.0 - 84.5)
<b>Race/Ethnicity</b>						
White Non-Hispanic	608	74.7	(70.0 - 79.4)	35,737	79.9	(78.4 - 81.4)
Black Non-Hispanic	*	*	*	738	78.6	(71.8 - 85.4)
Hispanic	*	*	*	863	88.9	(84.4 - 93.4)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	321	63.2	(55.4 - 71.0)	15,647	76.9	(73.8 - 80.0)
Some College	*	*	*	9,883	79.1	(75.2 - 83.0)
≥ College Degree	143	91.3	(86.9 - 95.7)	12,437	86.0	(83.5 - 88.5)
<b>Income</b>						
≤ \$24,999	*	*	*	9,608	74.9	(70.1 - 79.7)
\$25,000-\$49,999	184	74.0	(65.2 - 82.7)	9,640	79.8	(76.1 - 83.6)
\$50,000-\$74,999	*	*	*	5,766	78.6	(73.8 - 83.3)
≥ \$75,000	96	93.6	(88.5 - 98.6)	8,556	86.6	(84.3 - 88.9)

<sup>1</sup> Defined as having a rule that prohibits smoking anywhere inside the home

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 57a. Binge drinking within the past month among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	638	4,800	12.8	(9.3 - 16.4)	37,745	2,602,000	17.7	(15.7 - 19.8)
<b>Gender</b>								
Male	224	3,200	18.4	(12.0 - 24.7)	14,420	1,712,400	24.5	(20.9 - 28.1)
Female	414	1,600	7.9	(4.3 - 11.6)	23,325	889,600	11.6	(9.5 - 13.7)
<b>Age (years)</b>								
18-34	71	1,700	17.1	(7.7 - 26.5)	3,895	1,254,300	28.5	(22.8 - 34.2)
35-44	*	*	*	*	4,934	584,800	20.6	(16.5 - 24.7)
45-54	136	700	10.3	(4.8 - 15.8)	7,738	467,600	16.6	(13.3 - 19.9)
55-64	*	*	*	*	8,663	207,200	9.8	(7.9 - 11.8)
≥ 65	231	500	6.2	(2.8 - 9.6)	12,515	88,100	3.5	(2.5 - 4.5)
<b>Race/Ethnicity</b>								
White Non-Hispanic	610	4,400	13.0	(9.4 - 16.5)	35,453	1,761,900	19.2	(17.5 - 20.8)
Black Non-Hispanic	*	*	*	*	728	247,000	11.6	(6.3 - 17.0)
Hispanic	*	*	*	*	854	460,600	20.5	(11.3 - 29.8)
Other Non-Hispanic	*	*	*	*	710	132,500	12.1	(3.4 - 20.7)
<b>Education</b>								
≤ High School	322	2,200	12.1	(6.9 - 17.4)	15,491	945,800	17.4	(13.2 - 21.6)
Some College	170	1,800	16.4	(9.0 - 23.7)	9,819	636,600	18.1	(14.6 - 21.5)
≥ College Degree	146	900	9.9	(3.9 - 15.8)	12,358	1,018,700	18.2	(15.3 - 21.1)
<b>Income</b>								
≤ \$24,999	194	1,000	10.7	(3.5 - 17.9)	9,557	469,100	13.5	(8.5 - 18.5)
\$25,000-\$49,999	184	1,400	12.8	(6.4 - 19.1)	9,575	534,300	18.5	(13.0 - 24.0)
\$50,000-\$74,999	*	*	*	*	5,726	378,200	19.2	(14.1 - 24.3)
≥ \$75,000	97	1,500	20.7	(11.1 - 30.2)	8,501	1,011,400	22.6	(19.7 - 25.6)

<sup>1</sup> Defined as men having 5 or more drinks or women having 4 or more drinks on 1 or more occasion within the past month

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adult binge drinkers within the past month (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 57b. Binge drinking within the past month among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	638	14.3	(10.2 - 18.4)	37,745	18.1	(16.1 - 20.2)
<b>Gender</b>						
Male	224	20.3	(13.1 - 27.5)	14,420	24.5	(21.0 - 27.9)
Female	414	9.0	(4.8 - 13.2)	23,325	12.4	(10.2 - 14.5)
<b>Race/Ethnicity</b>						
White Non-Hispanic	610	14.4	(10.3 - 18.5)	35,453	19.8	(18.1 - 21.6)
Black Non-Hispanic	*	*	*	728	11.8	(6.4 - 17.1)
Hispanic	*	*	*	854	20.2	(11.3 - 29.1)
Other Non-Hispanic	*	*	*	710	12.1	(3.4 - 20.8)
<b>Education</b>						
≤ High School	322	13.9	(7.7 - 20.1)	15,491	17.5	(13.5 - 21.5)
Some College	170	17.8	(9.6 - 26.0)	9,819	18.6	(15.1 - 22.0)
≥ College Degree	146	10.7	(4.2 - 17.2)	12,358	18.8	(15.9 - 21.7)
<b>Income</b>						
≤ \$24,999	194	12.3	(3.8 - 20.8)	9,557	13.4	(8.6 - 18.2)
\$25,000-\$49,999	184	14.5	(7.0 - 21.9)	9,575	18.9	(13.5 - 24.2)
\$50,000-\$74,999	*	*	*	5,726	19.9	(14.8 - 25.1)
≥ \$75,000	*	*	*	8,501	23.2	(20.2 - 26.1)

<sup>1</sup> Defined as men having 5 or more drinks or women having 4 or more drinks on 1 or more occasion within the past month

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 58a. Heavy drinking within the past month among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	631	1,400	3.8	(1.9 - 5.6)	37,553	718,700	4.9	(4.2 - 5.6)
<b>Gender</b>								
Male	218	600	3.7	(0.4 - 7.0)	14,330	333,300	4.8	(3.9 - 5.7)
Female	413	800	3.8	(1.9 - 5.8)	23,223	385,400	5.0	(4.0 - 6.1)
<b>Age (years)</b>								
18-34	*	*	*	*	3,866	224,900	5.1	(3.5 - 6.8)
35-44	*	*	*	*	4,924	137,200	4.8	(3.0 - 6.7)
45-54	*	*	*	*	7,715	144,600	5.2	(4.1 - 6.2)
55-64	*	*	*	*	8,624	117,400	5.6	(3.9 - 7.3)
≥ 65	229	400	4.8	(2.1 - 7.5)	12,424	94,600	3.8	(2.7 - 4.9)
<b>Race/Ethnicity</b>								
White Non-Hispanic	603	1,300	4.0	(2.0 - 6.0)	35,279	572,400	6.3	(5.4 - 7.1)
Black Non-Hispanic	*	*	*	*	723	58,100	2.8	(0.6 - 4.9)
Hispanic	*	*	*	*	848	60,900	2.7	(0.5 - 4.9)
Other Non-Hispanic	*	*	*	*	703	27,200	2.5	(0.8 - 4.2)
<b>Education</b>								
≤ High School	320	800	4.8	(1.6 - 8.0)	15,411	228,100	4.2	(3.2 - 5.2)
Some College	*	*	*	*	9,759	163,700	4.7	(3.6 - 5.8)
≥ College Degree	*	*	*	*	12,306	326,800	5.9	(4.5 - 7.3)
<b>Income</b>								
≤ \$24,999	194	300	3.4	(1.3 - 5.4)	9,512	104,200	3.0	(1.8 - 4.1)
\$25,000-\$49,999	*	*	*	*	9,537	158,700	5.5	(3.9 - 7.2)
\$50,000-\$74,999	*	*	*	*	5,708	95,400	4.9	(3.1 - 6.7)
≥ \$75,000	*	*	*	*	8,481	313,900	7.0	(5.4 - 8.6)

<sup>1</sup> Defined as adult men averaging more than 2 alcoholic drinks per day and adult women averaging more than 1 alcoholic drink per day within the past month

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adult heavy drinkers within the past month (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 58b. Heavy drinking within the past month among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	631	3.8	(1.8 - 5.9)	37,553	5.0	(4.3 - 5.8)
<b>Gender</b>						
Male	218	3.9	(0.2 - 7.5)	14,330	4.9	(3.9 - 5.9)
Female	413	3.8	(1.6 - 5.9)	23,223	5.1	(4.0 - 6.2)
<b>Race/Ethnicity</b>						
White Non-Hispanic	603	4.1	(1.9 - 6.4)	35,279	6.4	(5.5 - 7.2)
Black Non-Hispanic	*	*	*	723	2.9	(0.6 - 5.1)
Hispanic	*	*	*	848	2.8	(0.5 - 5.1)
Other Non-Hispanic	*	*	*	703	2.5	(0.8 - 4.2)
<b>Education</b>						
≤ High School	320	5.0	(1.4 - 8.7)	15,411	4.3	(3.2 - 5.3)
Some College	*	*	*	9,759	4.8	(3.6 - 5.9)
≥ College Degree	*	*	*	12,306	6.0	(4.5 - 7.4)
<b>Income</b>						
≤ \$24,999	194	3.1	(1.2 - 5.1)	9,512	3.0	(1.9 - 4.1)
\$25,000-\$49,999	*	*	*	9,537	5.8	(4.0 - 7.5)
\$50,000-\$74,999	*	*	*	5,708	5.0	(3.1 - 7.0)
≥ \$75,000	*	*	*	8,481	7.0	(5.4 - 8.7)

<sup>1</sup> Defined as adult men averaging more than 2 alcoholic drinks per day and adult women averaging more than 1 alcoholic drink per day within the past month

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 59a. Never or rarely been asked about sexual history during routine check-up among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	596	24,000	69.1	(64.1 - 74.1)	35,193	8,480,600	61.9	(59.2 - 64.5)
<b>Gender</b>								
Male	205	13,000	81.3	(74.0 - 88.7)	13,299	4,357,400	67.7	(63.6 - 71.7)
Female	391	11,000	58.7	(52.4 - 65.0)	21,894	4,123,300	56.8	(53.3 - 60.2)
<b>Age (years)</b>								
18-34	*	*	*	*	3,507	1,804,600	44.3	(37.8 - 50.8)
35-44	*	*	*	*	4,531	1,635,300	62.2	(56.6 - 67.8)
45-54	129	4,800	70.1	(61.3 - 78.9)	7,215	1,715,600	65.4	(60.5 - 70.3)
55-64	123	4,000	73.9	(65.3 - 82.5)	8,168	1,416,200	70.7	(66.6 - 74.8)
≥ 65	212	6,300	83.1	(77.7 - 88.6)	11,772	1,908,900	80.3	(76.9 - 83.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	568	21,600	69.1	(64.3 - 74.0)	33,090	5,940,300	68.5	(66.3 - 70.7)
Black Non-Hispanic	*	*	*	*	683	831,800	40.8	(31.8 - 49.9)
Hispanic	*	*	*	*	784	1,071,700	53.4	(43.6 - 63.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	297	11,300	68.3	(61.1 - 75.4)	14,264	3,001,200	60.1	(55.4 - 64.9)
Some College	157	7,200	73.4	(64.7 - 82.0)	9,156	1,972,800	60.3	(54.9 - 65.7)
≥ College Degree	*	*	*	*	11,706	3,476,000	65.0	(61.2 - 68.7)
<b>Income</b>								
≤ \$24,999	175	5,900	71.2	(61.2 - 81.1)	8,772	1,529,400	49.3	(42.6 - 56.0)
\$25,000-\$49,999	176	7,100	69.7	(60.8 - 78.6)	8,932	1,705,000	64.4	(58.0 - 70.7)
\$50,000-\$74,999	*	*	*	*	5,433	1,217,000	64.5	(58.5 - 70.5)
≥ \$75,000	*	*	*	*	8,108	2,934,300	68.2	(64.7 - 71.7)

<sup>1</sup> Defined as respondents who never or rarely (hardly ever) were asked about their sexual history, such as about sexual partners and sexual practices, during a routine check-up at a doctor's office or clinic

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who were never or rarely asked about their sexual history during a routine check-up at a doctor's office or clinic (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 59b. Never or rarely been asked about sexual history during routine check-up among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	596	67.2	(61.7 - 72.7)	35,193	61.4	(58.7 - 64.0)
<b>Gender</b>						
Male	205	81.1	(73.3 - 89.0)	13,299	67.9	(64.0 - 71.9)
Female	391	55.4	(48.4 - 62.4)	21,894	55.5	(52.0 - 59.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	568	67.0	(61.6 - 72.4)	33,090	67.8	(65.6 - 70.1)
Black Non-Hispanic	*	*	*	683	41.1	(32.0 - 50.2)
Hispanic	*	*	*	784	53.0	(43.3 - 62.6)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	297	67.1	(59.0 - 75.1)	14,264	59.5	(54.7 - 64.3)
Some College	157	70.9	(61.1 - 80.6)	9,156	59.7	(54.4 - 65.0)
≥ College Degree	*	*	*	11,706	64.5	(60.8 - 68.2)
<b>Income</b>						
≤ \$24,999	*	*	*	8,772	48.7	(42.0 - 55.4)
\$25,000-\$49,999	176	67.8	(57.9 - 77.8)	8,932	63.7	(57.5 - 70.0)
\$50,000-\$74,999	*	*	*	5,433	64.2	(58.2 - 70.2)
≥ \$75,000	*	*	*	8,108	67.5	(64.0 - 71.0)

<sup>1</sup> Defined as respondents who never or rarely (hardly ever) were asked about their sexual history, such as about sexual partners and sexual practices, during a routine check-up at a doctor's office or clinic

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 60a. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	372	18,800	74.5	(68.5 - 80.6)	22,788	7,011,100	77.9	(75.2 - 80.5)
<b>Gender</b>								
Male	146	10,200	76.5	(67.2 - 85.8)	9,034	3,618,500	81.0	(77.2 - 84.8)
Female	226	8,700	72.4	(64.9 - 79.8)	13,754	3,392,600	74.8	(71.2 - 78.4)
<b>Age (years)</b>								
18-34	*	*	*	*	3,213	2,362,500	70.8	(65.1 - 76.6)
35-44	*	*	*	*	3,513	1,313,400	71.7	(65.6 - 77.7)
45-54	94	4,100	77.5	(67.7 - 87.3)	4,975	1,290,600	80.3	(75.9 - 84.8)
55-64	*	*	*	*	4,955	932,200	87.2	(84.1 - 90.3)
≥ 65	93	3,200	95.5	(91.0 - 100)	6,132	1,112,400	95.9	(94.4 - 97.5)
<b>Race/Ethnicity</b>								
White Non-Hispanic	357	17,400	77.0	(71.5 - 82.6)	21,540	4,875,200	78.3	(75.6 - 80.9)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	491	1,002,400	84.8	(77.8 - 91.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	174	8,800	77.8	(69.3 - 86.4)	8,879	2,581,700	82.1	(77.9 - 86.3)
Some College	*	*	*	*	5,953	1,599,500	77.5	(72.6 - 82.3)
≥ College Degree	*	*	*	*	7,928	2,824,800	75.6	(71.8 - 79.5)
<b>Income</b>								
≤ \$24,999	97	4,600	82.6	(72.8 - 92.4)	5,491	1,602,900	78.7	(71.9 - 85.6)
\$25,000-\$49,999	*	*	*	*	5,912	1,335,900	78.8	(73.8 - 83.9)
\$50,000-\$74,999	*	*	*	*	3,717	987,500	76.6	(70.4 - 82.8)
≥ \$75,000	*	*	*	*	5,771	2,321,600	77.6	(74.0 - 81.2)

<sup>1</sup> Defined as respondent estimating there were hardly any (0-1 out of 10) or a few (2-3 out of 10) persons their age in their community who have had a sexually transmitted disease such as gonorrhea, chlamydia, herpes or syphilis

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who believed hardly any or a few people their age in their community have had a sexually transmitted disease (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 60b. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	372	72.8	(66.4 - 79.3)	22,788	77.3	(74.7 - 79.9)
<b>Gender</b>						
Male	146	75.6	(65.8 - 85.4)	9,034	81.1	(77.4 - 84.8)
Female	226	69.9	(61.7 - 78.1)	13,754	73.6	(70.0 - 77.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	357	75.3	(69.3 - 81.4)	21,540	77.5	(74.9 - 80.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	491	84.6	(77.6 - 91.5)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	174	75.2	(65.5 - 84.9)	8,879	81.6	(77.4 - 85.7)
Some College	*	*	*	5,953	76.8	(72.0 - 81.6)
≥ College Degree	*	*	*	7,928	75.1	(71.3 - 79.0)
<b>Income</b>						
≤ \$24,999	*	*	*	5,491	77.9	(71.0 - 84.7)
\$25,000-\$49,999	*	*	*	5,912	77.9	(72.8 - 83.0)
\$50,000-\$74,999	*	*	*	3,717	76.1	(70.0 - 82.3)
≥ \$75,000	*	*	*	5,771	77.0	(73.4 - 80.6)

<sup>1</sup> Defined as respondent estimating there were hardly any (0-1 out of 10) or a few (2-3 out of 10) persons their age in their community who have had a sexually transmitted disease such as gonorrhea, chlamydia, herpes or syphilis

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 61a. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	601	30,300	84.8	(81.2 - 88.4)	36,287	12,738,300	89.7	(88.2 - 91.3)
<b>Gender</b>								
Male	206	14,000	83.8	(77.7 - 89.9)	13,893	6,104,600	89.4	(87.2 - 91.5)
Female	395	16,300	85.6	(81.5 - 89.8)	22,394	6,633,700	90.1	(87.9 - 92.2)
<b>Age (years)</b>								
18-34	69	8,600	86.9	(77.6 - 96.1)	3,850	4,007,400	91.3	(87.5 - 95.1)
35-44	72	5,600	93.2	(86.5 - 100)	4,875	2,578,600	93.2	(90.4 - 96.0)
45-54	133	6,000	86.2	(78.9 - 93.5)	7,607	2,416,800	90.5	(87.9 - 93.2)
55-64	123	4,800	88.0	(82.1 - 93.8)	8,411	1,876,500	92.2	(90.0 - 94.5)
≥ 65	204	5,300	71.4	(64.6 - 78.1)	11,544	1,858,900	79.7	(76.6 - 82.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	575	26,900	83.7	(79.8 - 87.5)	34,102	8,050,700	89.4	(87.7 - 91.0)
Black Non-Hispanic	*	*	*	*	702	1,822,200	91.9	(88.2 - 95.6)
Hispanic	*	*	*	*	820	1,967,200	89.9	(85.2 - 94.7)
Other Non-Hispanic	*	*	*	*	663	898,200	88.6	(79.5 - 97.6)
<b>Education</b>								
≤ High School	295	12,500	75.3	(68.9 - 81.7)	14,507	4,385,300	84.6	(81.9 - 87.2)
Some College	162	9,700	92.1	(87.1 - 97.2)	9,571	3,094,300	90.0	(86.7 - 93.4)
≥ College Degree	144	8,100	93.9	(89.6 - 98.3)	12,142	5,228,800	95.4	(94.0 - 96.7)
<b>Income</b>								
≤ \$24,999	181	6,600	76.1	(67.4 - 84.8)	9,021	2,833,500	86.5	(82.6 - 90.3)
\$25,000-\$49,999	178	9,100	87.6	(82.1 - 93.1)	9,261	2,529,900	89.5	(86.5 - 92.5)
\$50,000-\$74,999	92	5,300	86.4	(76.4 - 96.4)	5,644	1,740,000	91.2	(87.4 - 95.1)
≥ \$75,000	94	6,700	92.7	(86.8 - 98.7)	8,397	4,184,300	94.4	(92.5 - 96.3)

<sup>1</sup> Defined as believing it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who believed it is at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 61b. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	601	85.8	(82.0 - 89.6)	36,287	89.8	(88.3 - 91.4)
<b>Gender</b>						
Male	206	84.2	(77.8 - 90.6)	13,893	89.1	(86.9 - 91.3)
Female	395	87.2	(82.8 - 91.7)	22,394	90.5	(88.3 - 92.7)
<b>Race/Ethnicity</b>						
White Non-Hispanic	575	84.7	(80.6 - 88.8)	34,102	89.5	(87.8 - 91.2)
Black Non-Hispanic	*	*	*	702	91.7	(87.8 - 95.6)
Hispanic	*	*	*	820	89.6	(84.7 - 94.5)
Other Non-Hispanic	*	*	*	663	89.0	(80.5 - 97.6)
<b>Education</b>						
≤ High School	295	75.5	(68.3 - 82.7)	14,507	84.6	(81.8 - 87.4)
Some College	162	93.1	(87.9 - 98.2)	9,571	90.2	(86.9 - 93.5)
≥ College Degree	144	95.2	(91.5 - 98.9)	12,142	95.4	(94.0 - 96.7)
<b>Income</b>						
≤ \$24,999	181	76.3	(66.4 - 86.1)	9,021	86.5	(82.6 - 90.4)
\$25,000-\$49,999	178	89.4	(84.2 - 94.6)	9,261	89.6	(86.5 - 92.6)
\$50,000-\$74,999	*	*	*	5,644	91.1	(86.9 - 95.3)
≥ \$75,000	94	93.4	(87.5 - 99.3)	8,397	94.5	(92.6 - 96.4)

<sup>1</sup> Defined as believing it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 62a. Overweight among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	613	12,900	35.5	(30.5 - 40.4)	36,710	5,213,000	36.4	(34.0 - 38.8)
<b>Gender</b>								
Male	224	6,600	37.4	(29.4 - 45.5)	14,502	3,073,200	43.7	(39.9 - 47.6)
Female	389	6,300	33.6	(27.7 - 39.6)	22,208	2,139,800	29.3	(26.4 - 32.2)
<b>Age (years)</b>								
18-34	*	*	*	*	3,789	1,295,200	30.1	(24.7 - 35.6)
35-44	*	*	*	*	4,785	1,052,000	37.5	(31.7 - 43.3)
45-54	130	2,300	33.5	(24.1 - 42.9)	7,495	1,053,300	38.8	(34.1 - 43.4)
55-64	119	2,200	40.1	(30.1 - 50.1)	8,356	798,100	38.8	(34.8 - 42.7)
≥ 65	223	3,000	37.3	(30.4 - 44.2)	12,285	1,014,400	41.5	(37.6 - 45.4)
<b>Race/Ethnicity</b>								
White Non-Hispanic	586	11,100	33.8	(29.1 - 38.5)	34,501	3,282,200	36.4	(34.4 - 38.3)
Black Non-Hispanic	*	*	*	*	708	740,000	36.7	(28.3 - 45.1)
Hispanic	*	*	*	*	813	781,500	35.6	(27.1 - 44.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	306	5,600	32.9	(26.1 - 39.7)	15,079	1,889,700	35.4	(31.2 - 39.5)
Some College	167	4,100	37.9	(28.3 - 47.4)	9,515	1,314,500	38.7	(33.8 - 43.7)
≥ College Degree	*	*	*	*	12,062	1,996,900	36.3	(32.8 - 39.7)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,346	1,170,900	34.0	(28.2 - 39.8)
\$25,000-\$49,999	183	3,400	31.7	(23.5 - 39.9)	9,368	1,136,000	40.4	(34.6 - 46.2)
\$50,000-\$74,999	*	*	*	*	5,615	738,700	38.3	(32.3 - 44.3)
≥ \$75,000	*	*	*	*	8,337	1,659,100	37.5	(34.1 - 40.9)

<sup>1</sup> Defined as having a body mass index (BMI) more than 24.9 but less than 30.0. BMI is calculated as weight in kilograms divided by the square of height in meters.

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who are overweight (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 62b. Overweight among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	613	35.0	(29.7 - 40.4)	36,710	36.2	(33.8 - 38.6)
<b>Gender</b>						
Male	224	37.6	(29.0 - 46.2)	14,502	43.9	(40.1 - 47.6)
Female	389	32.6	(26.0 - 39.3)	22,208	28.9	(25.9 - 31.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	586	33.2	(28.1 - 38.4)	34,501	36.0	(34.0 - 38.0)
Black Non-Hispanic	*	*	*	708	37.0	(28.5 - 45.5)
Hispanic	*	*	*	813	36.0	(27.5 - 44.4)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	306	32.3	(24.6 - 39.9)	15,079	35.2	(31.1 - 39.4)
Some College	*	*	*	9,515	38.3	(33.3 - 43.2)
≥ College Degree	*	*	*	12,062	36.2	(32.7 - 39.8)
<b>Income</b>						
≤ \$24,999	*	*	*	9,346	33.8	(28.0 - 39.5)
\$25,000-\$49,999	183	30.1	(21.4 - 38.8)	9,368	40.6	(34.8 - 46.4)
\$50,000-\$74,999	*	*	*	5,615	38.6	(32.5 - 44.7)
≥ \$75,000	*	*	*	8,337	37.2	(33.7 - 40.6)

<sup>1</sup> Defined as having a body mass index (BMI) more than 24.9 but less than 30.0. BMI is calculated as weight in kilograms divided by the square of height in meters.

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 63a. Obesity among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	613	11,600	31.8	(26.7 - 36.8)	36,710	3,327,400	23.2	(21.2 - 25.3)
<b>Gender</b>								
Male	224	6,600	37.5	(28.9 - 46.0)	14,502	1,505,800	21.4	(19.0 - 23.9)
Female	389	5,000	26.4	(21.1 - 31.7)	22,208	1,821,600	25.0	(21.8 - 28.1)
<b>Age (years)</b>								
18-34	*	*	*	*	3,789	730,600	17.0	(12.4 - 21.6)
35-44	*	*	*	*	4,785	662,200	23.6	(19.4 - 27.9)
45-54	130	2,100	30.8	(21.4 - 40.3)	7,495	816,800	30.1	(25.3 - 34.8)
55-64	*	*	*	*	8,356	563,900	27.4	(23.7 - 31.1)
≥ 65	223	2,400	29.0	(22.3 - 35.8)	12,285	554,000	22.7	(19.2 - 26.1)
<b>Race/Ethnicity</b>								
White Non-Hispanic	586	10,000	30.6	(25.9 - 35.2)	34,501	2,062,700	22.9	(21.2 - 24.6)
Black Non-Hispanic	*	*	*	*	708	703,100	34.9	(26.1 - 43.7)
Hispanic	*	*	*	*	813	456,400	20.8	(14.3 - 27.2)
Other Non-Hispanic	*	*	*	*	688	105,200	9.6	(4.0 - 15.3)
<b>Education</b>								
≤ High School	306	5,900	34.8	(27.8 - 41.9)	15,079	1,561,600	29.2	(25.3 - 33.2)
Some College	*	*	*	*	9,515	792,800	23.4	(19.4 - 27.3)
≥ College Degree	140	2,100	25.1	(16.4 - 33.9)	12,062	915,000	16.6	(14.4 - 18.8)
<b>Income</b>								
≤ \$24,999	*	*	*	*	9,346	974,800	28.3	(22.6 - 34.0)
\$25,000-\$49,999	183	3,000	27.8	(19.4 - 36.2)	9,368	584,800	20.8	(17.1 - 24.5)
\$50,000-\$74,999	*	*	*	*	5,615	475,300	24.6	(20.0 - 29.3)
≥ \$75,000	93	2,000	27.7	(17.8 - 37.6)	8,337	867,700	19.6	(17.0 - 22.3)

<sup>1</sup> Defined as having a body mass index (BMI) of 30.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who are obese (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 63b. Obesity among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	613	31.0	(25.6 - 36.4)	36,710	23.1	(21.0 - 25.1)
<b>Gender</b>						
Male	224	36.7	(27.7 - 45.7)	14,502	21.5	(19.0 - 24.0)
Female	389	25.7	(19.9 - 31.5)	22,208	24.6	(21.5 - 27.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	586	29.6	(24.6 - 34.5)	34,501	22.8	(21.0 - 24.5)
Black Non-Hispanic	*	*	*	708	34.8	(25.9 - 43.6)
Hispanic	*	*	*	813	20.5	(14.1 - 26.9)
Other Non-Hispanic	*	*	*	688	9.5	(3.8 - 15.2)
<b>Education</b>						
≤ High School	306	34.2	(26.4 - 42.0)	15,079	29.4	(25.4 - 33.3)
Some College	*	*	*	9,515	23.1	(19.2 - 27.1)
≥ College Degree	140	25.6	(16.3 - 34.8)	12,062	16.4	(14.2 - 18.5)
<b>Income</b>						
≤ \$24,999	*	*	*	9,346	28.3	(22.6 - 34.1)
\$25,000-\$49,999	183	27.5	(18.4 - 36.7)	9,368	21.0	(17.2 - 24.8)
\$50,000-\$74,999	*	*	*	5,615	24.0	(19.4 - 28.7)
≥ \$75,000	93	23.8	(14.6 - 33.1)	8,337	19.3	(16.7 - 21.9)

<sup>1</sup> Defined as having a body mass index (BMI) of 30.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 64a. Overweight or obesity among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	613	24,500	67.3	(62.4 - 72.1)	36,710	8,540,500	59.6	(57.0 - 62.2)
<b>Gender</b>								
Male	224	13,200	74.9	(67.6 - 82.1)	14,502	4,579,100	65.2	(61.0 - 69.4)
Female	389	11,300	60.1	(54.0 - 66.2)	22,208	3,961,400	54.3	(51.0 - 57.6)
<b>Age (years)</b>								
18-34	*	*	*	*	3,789	2,025,800	47.1	(40.8 - 53.5)
35-44	*	*	*	*	4,785	1,714,200	61.1	(55.6 - 66.6)
45-54	130	4,400	64.3	(54.6 - 74.0)	7,495	1,870,100	68.8	(64.6 - 73.1)
55-64	119	4,600	85.4	(78.9 - 92.0)	8,356	1,362,000	66.2	(61.9 - 70.4)
≥ 65	223	5,400	66.3	(59.5 - 73.2)	12,285	1,568,400	64.1	(60.6 - 67.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	586	21,100	64.3	(59.4 - 69.3)	34,501	5,344,900	59.3	(57.2 - 61.3)
Black Non-Hispanic	*	*	*	*	708	1,443,100	71.6	(62.5 - 80.7)
Hispanic	*	*	*	*	813	1,237,900	56.4	(46.5 - 66.2)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	306	11,600	67.8	(60.7 - 74.8)	15,079	3,451,400	64.6	(59.8 - 69.4)
Some College	167	7,600	70.0	(61.1 - 78.9)	9,515	2,107,300	62.1	(57.0 - 67.2)
≥ College Degree	140	5,300	62.7	(53.0 - 72.4)	12,062	2,911,900	52.9	(49.2 - 56.6)
<b>Income</b>								
≤ \$24,999	185	6,400	73.3	(64.2 - 82.5)	9,346	2,145,700	62.3	(55.7 - 68.9)
\$25,000-\$49,999	183	6,300	59.5	(50.5 - 68.5)	9,368	1,720,700	61.2	(54.5 - 67.9)
\$50,000-\$74,999	92	4,900	79.1	(69.7 - 88.6)	5,615	1,214,000	62.9	(57.1 - 68.8)
≥ \$75,000	*	*	*	*	8,337	2,526,700	57.1	(53.6 - 60.6)

<sup>1</sup> Defined as having a body mass index (BMI) of 25.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who are overweight or obese (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 64b. Overweight or obesity among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	613	66.0	(60.8 - 71.3)	36,710	59.3	(56.8 - 61.9)
<b>Gender</b>						
Male	224	74.3	(66.6 - 82.0)	14,502	65.4	(61.3 - 69.4)
Female	389	58.3	(51.4 - 65.2)	22,208	53.5	(50.1 - 56.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	586	62.8	(57.4 - 68.2)	34,501	58.8	(56.7 - 60.8)
Black Non-Hispanic	*	*	*	708	71.8	(63.0 - 80.6)
Hispanic	*	*	*	813	56.5	(46.8 - 66.1)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	306	66.5	(58.5 - 74.5)	15,079	64.6	(59.9 - 69.3)
Some College	167	68.7	(59.1 - 78.3)	9,515	61.4	(56.4 - 66.5)
≥ College Degree	*	*	*	12,062	52.6	(48.9 - 56.3)
<b>Income</b>						
≤ \$24,999	*	*	*	9,346	62.1	(55.6 - 68.6)
\$25,000-\$49,999	183	57.6	(47.7 - 67.5)	9,368	61.6	(55.1 - 68.1)
\$50,000-\$74,999	*	*	*	5,615	62.6	(56.7 - 68.6)
≥ \$75,000	*	*	*	8,337	56.5	(52.9 - 60.0)

<sup>1</sup> Defined as having a body mass index (BMI) of 25.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 65a. Received advice about weight by a health professional among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	612	9,700	26.7	(22.5 - 30.9)	36,729	3,929,700	27.4	(25.1 - 29.7)
<b>Gender</b>								
Male	223	4,900	27.6	(20.8 - 34.4)	14,499	1,654,800	23.6	(20.6 - 26.6)
Female	389	4,900	25.9	(20.9 - 31.0)	22,230	2,274,900	31.0	(27.7 - 34.3)
<b>Age (years)</b>								
18-34	71	1,000	9.8	(3.2 - 16.3)	3,796	895,500	20.7	(15.3 - 26.2)
35-44	*	*	*	*	4,779	722,700	25.8	(20.9 - 30.6)
45-54	130	2,100	30.6	(21.6 - 39.5)	7,490	910,900	33.5	(29.0 - 38.1)
55-64	*	*	*	*	8,351	741,100	36.1	(32.0 - 40.3)
≥ 65	222	2,500	31.3	(24.5 - 38.1)	12,313	659,500	26.9	(23.0 - 30.7)
<b>Race/Ethnicity</b>								
White Non-Hispanic	585	8,700	26.6	(22.5 - 30.8)	34,502	2,386,200	26.5	(24.6 - 28.4)
Black Non-Hispanic	*	*	*	*	709	615,700	30.4	(22.0 - 38.8)
Hispanic	*	*	*	*	830	555,400	25.1	(17.8 - 32.4)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	307	5,000	29.5	(23.1 - 35.8)	15,116	1,469,700	27.5	(23.6 - 31.4)
Some College	165	2,100	19.7	(13.0 - 26.5)	9,515	946,800	27.8	(23.2 - 32.4)
≥ College Degree	140	2,600	30.1	(21.1 - 39.1)	12,041	1,445,500	26.2	(23.0 - 29.4)
<b>Income</b>								
≤ \$24,999	186	2,100	24.2	(16.4 - 32.0)	9,360	1,079,400	31.4	(25.3 - 37.5)
\$25,000-\$49,999	182	2,700	26.0	(18.8 - 33.1)	9,365	696,900	24.6	(19.8 - 29.4)
\$50,000-\$74,999	*	*	*	*	5,603	507,100	26.3	(21.6 - 31.0)
≥ \$75,000	*	*	*	*	8,334	1,207,100	27.3	(24.2 - 30.5)

<sup>1</sup> Defined as being advised by a doctor, nurse or other health professional about their weight (lose, gain or maintain)

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults provided advice by a health professional about their weight (lose, gain or maintain) (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 65b. Received advice about weight by a health professional among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	612	25.7	(21.3 - 30.1)	36,729	27.1	(24.8 - 29.4)
<b>Gender</b>						
Male	223	26.4	(19.3 - 33.5)	14,499	23.4	(20.5 - 26.3)
Female	389	25.1	(19.6 - 30.5)	22,230	30.6	(27.3 - 34.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	585	25.5	(21.1 - 29.9)	34,502	26.1	(24.2 - 28.0)
Black Non-Hispanic	*	*	*	709	30.4	(21.9 - 38.9)
Hispanic	*	*	*	830	24.9	(17.7 - 32.2)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	307	28.4	(21.5 - 35.3)	15,116	27.3	(23.4 - 31.2)
Some College	165	18.0	(11.3 - 24.6)	9,515	27.4	(22.9 - 32.0)
≥ College Degree	140	30.5	(21.1 - 39.9)	12,041	25.9	(22.7 - 29.0)
<b>Income</b>						
≤ \$24,999	186	22.5	(14.4 - 30.5)	9,360	31.0	(24.9 - 37.0)
\$25,000-\$49,999	182	24.6	(17.2 - 32.0)	9,365	24.9	(20.0 - 29.8)
\$50,000-\$74,999	*	*	*	5,603	25.9	(21.2 - 30.6)
≥ \$75,000	*	*	*	8,334	26.9	(23.7 - 30.0)

<sup>1</sup> Defined as being advised by a doctor, nurse or other health professional about their weight (lose, gain or maintain)

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 66a. Received advice to lose weight by a health professional among adults provided weight advice<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	184	8,200	84.3	(78.0 - 90.5)	9,724	3,412,900	86.8	(83.9 - 89.8)
<b>Gender</b>								
Male	73	4,200	87.4	(78.3 - 96.5)	3,807	1,468,700	88.8	(85.7 - 91.8)
Female	111	3,900	81.1	(72.5 - 89.7)	5,917	1,944,200	85.5	(80.8 - 90.1)
<b>Age (years)</b>								
18-34	*	*	*	*	681	779,700	87.1	(80.6 - 93.6)
35-44	*	*	*	*	1,215	622,100	86.1	(76.9 - 95.2)
45-54	*	*	*	*	2,238	813,300	89.3	(84.4 - 94.2)
55-64	*	*	*	*	2,697	645,500	87.1	(81.2 - 93.0)
≥ 65	67	2,300	90.0	(82.2 - 97.9)	2,893	552,300	83.7	(76.9 - 90.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	175	7,200	83.0	(76.2 - 89.8)	9,082	2,076,700	87.0	(84.5 - 89.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	251	483,600	87.1	(78.5 - 95.7)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	98	4,500	89.8	(82.6 - 97.0)	3,962	1,239,700	84.3	(78.6 - 90.1)
Some College	*	*	*	*	2,633	814,800	86.1	(80.1 - 92.0)
≥ College Degree	*	*	*	*	3,115	1,290,900	89.3	(85.3 - 93.3)
<b>Income</b>								
≤ \$24,999	55	1,900	90.2	(81.2 - 99.1)	2,539	922,100	85.4	(78.7 - 92.1)
\$25,000-\$49,999	56	2,400	86.4	(76.5 - 96.4)	2,482	594,100	85.2	(78.4 - 92.1)
\$50,000-\$74,999	*	*	*	*	1,521	446,100	88.0	(83.2 - 92.7)
≥ \$75,000	*	*	*	*	2,270	1,073,200	88.9	(83.7 - 94.1)

<sup>1</sup> Defined as provided advice by a health professional about their weight and advised to lose weight

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults provided advice by a health professional about their weight and advised to lose weight (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 66b. Received advice to lose weight by a health professional among adults provided weight advice<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	184	82.9	(75.7 - 90.1)	9,724	86.9	(83.8 - 89.9)
<b>Gender</b>						
Male	*	*	*	3,807	88.7	(85.7 - 91.8)
Female	111	79.6	(70.0 - 89.2)	5,917	85.5	(80.7 - 90.3)
<b>Race/Ethnicity</b>						
White Non-Hispanic	175	81.3	(73.5 - 89.2)	9,082	87.0	(84.4 - 89.6)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	251	87.6	(79.3 - 95.8)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	98	89.2	(81.2 - 97.3)	3,962	84.6	(78.7 - 90.5)
Some College	*	*	*	2,633	85.6	(79.5 - 91.7)
≥ College Degree	*	*	*	3,115	89.4	(85.4 - 93.3)
<b>Income</b>						
≤ \$24,999	*	*	*	2,539	85.7	(79.0 - 92.3)
\$25,000-\$49,999	*	*	*	2,482	85.3	(78.4 - 92.3)
\$50,000-\$74,999	*	*	*	1,521	88.0	(83.1 - 92.8)
≥ \$75,000	*	*	*	2,270	88.7	(83.1 - 94.2)

<sup>1</sup> Defined as provided advice by a health professional about their weight and advised to lose weight

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 67a. No leisure-time physical activity among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	642	9,800	26.1	(22.0 - 30.1)	38,128	3,541,700	23.9	(21.6 - 26.1)
<b>Gender</b>								
Male	226	4,000	22.5	(16.5 - 28.6)	14,602	1,482,500	20.9	(17.8 - 24.0)
Female	416	5,800	29.3	(24.0 - 34.5)	23,526	2,059,200	26.6	(23.4 - 29.7)
<b>Age (years)</b>								
18-34	*	*	*	*	3,938	982,900	22.0	(16.6 - 27.5)
35-44	74	1,300	20.8	(10.9 - 30.7)	4,988	556,000	19.3	(15.0 - 23.6)
45-54	137	2,100	29.5	(20.4 - 38.6)	7,827	759,000	26.8	(21.7 - 31.9)
55-64	128	1,800	31.9	(22.9 - 40.9)	8,765	464,900	21.7	(18.2 - 25.2)
≥ 65	232	3,500	41.2	(34.2 - 48.1)	12,610	778,800	30.7	(27.3 - 34.2)
<b>Race/Ethnicity</b>								
White Non-Hispanic	614	8,900	26.2	(22.2 - 30.3)	35,816	1,892,800	20.3	(18.6 - 22.0)
Black Non-Hispanic	*	*	*	*	739	690,000	32.0	(23.4 - 40.6)
Hispanic	*	*	*	*	860	600,800	26.7	(19.1 - 34.4)
Other Non-Hispanic	*	*	*	*	*	*	*	*
<b>Education</b>								
≤ High School	325	6,200	34.7	(28.3 - 41.2)	15,683	1,810,200	32.5	(28.4 - 36.5)
Some College	171	2,200	19.7	(12.8 - 26.6)	9,906	814,200	23.0	(18.6 - 27.3)
≥ College Degree	146	1,400	16.3	(9.8 - 22.8)	12,458	841,000	14.9	(11.9 - 18.0)
<b>Income</b>								
≤ \$24,999	194	3,100	34.0	(24.8 - 43.3)	9,640	1,332,200	38.0	(31.7 - 44.3)
\$25,000-\$49,999	185	3,200	29.5	(21.7 - 37.3)	9,658	648,000	22.3	(18.3 - 26.3)
\$50,000-\$74,999	95	1,400	21.7	(12.1 - 31.4)	5,773	430,700	21.8	(16.6 - 27.0)
≥ \$75,000	97	1,100	15.0	(7.5 - 22.6)	8,565	560,800	12.5	(10.3 - 14.6)

<sup>1</sup> Defined as having not participated in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have not participated in any physical activities or exercises (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 67b. No leisure-time physical activity among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	642	24.1	(19.9 - 28.3)	38,128	23.7	(21.5 - 25.9)
<b>Gender</b>						
Male	226	20.9	(14.9 - 27.0)	14,602	20.9	(17.8 - 23.9)
Female	416	26.9	(21.2 - 32.5)	23,526	26.3	(23.1 - 29.5)
<b>Race/Ethnicity</b>						
White Non-Hispanic	614	24.1	(19.9 - 28.3)	35,816	20.0	(18.3 - 21.7)
Black Non-Hispanic	*	*	*	739	32.0	(23.3 - 40.6)
Hispanic	*	*	*	860	26.9	(19.2 - 34.5)
Other Non-Hispanic	*	*	*	*	*	*
<b>Education</b>						
≤ High School	325	32.2	(25.4 - 39.1)	15,683	32.3	(28.3 - 36.4)
Some College	171	18.8	(11.6 - 26.1)	9,906	22.8	(18.5 - 27.2)
≥ College Degree	146	15.3	(8.7 - 21.9)	12,458	14.9	(11.8 - 17.9)
<b>Income</b>						
≤ \$24,999	*	*	*	9,640	37.9	(31.6 - 44.2)
\$25,000-\$49,999	185	28.0	(19.7 - 36.3)	9,658	22.0	(18.0 - 26.0)
\$50,000-\$74,999	*	*	*	5,773	21.6	(16.3 - 26.8)
≥ \$75,000	97	13.3	(6.4 - 20.3)	8,565	12.6	(10.3 - 14.9)

<sup>1</sup> Defined as having not participated in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 68a. Consumption of 5 or more servings of fruits and vegetables per day among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	643	9,200	24.5	(20.2 - 28.8)	38,203	4,023,400	27.0	(24.9 - 29.2)
<b>Gender</b>								
Male	227	3,300	18.4	(11.6 - 25.2)	14,634	1,603,900	22.6	(19.4 - 25.8)
Female	416	5,900	30.0	(24.5 - 35.4)	23,569	2,419,500	31.1	(28.2 - 34.0)
<b>Age (years)</b>								
18-34	*	*	*	*	3,940	1,261,700	28.3	(22.7 - 33.9)
35-44	74	1,300	20.5	(10.6 - 30.4)	4,989	688,000	23.9	(19.5 - 28.3)
45-54	138	1,800	25.3	(16.8 - 33.8)	7,837	746,800	26.2	(22.3 - 30.1)
55-64	128	1,600	27.9	(18.8 - 37.0)	8,777	579,100	27.0	(23.6 - 30.5)
≥ 65	232	2,200	26.6	(20.3 - 33.0)	12,660	747,800	29.4	(26.0 - 32.8)
<b>Race/Ethnicity</b>								
White Non-Hispanic	615	8,500	24.8	(20.7 - 29.0)	35,884	2,538,000	27.2	(25.4 - 28.9)
Black Non-Hispanic	*	*	*	*	740	572,400	26.5	(19.2 - 33.9)
Hispanic	*	*	*	*	864	672,200	29.8	(21.0 - 38.5)
Other Non-Hispanic	*	*	*	*	715	240,800	21.4	(12.0 - 30.8)
<b>Education</b>								
≤ High School	325	3,200	17.6	(12.5 - 22.7)	15,726	1,309,800	23.4	(19.3 - 27.4)
Some College	172	2,500	22.5	(15.1 - 29.9)	9,924	964,300	27.2	(23.0 - 31.3)
≥ College Degree	*	*	*	*	12,471	1,747,500	31.0	(27.9 - 34.1)
<b>Income</b>								
≤ \$24,999	194	2,000	22.1	(14.2 - 30.0)	9,667	831,500	23.7	(18.6 - 28.7)
\$25,000-\$49,999	185	2,700	25.6	(18.0 - 33.2)	9,669	917,200	31.6	(25.6 - 37.6)
\$50,000-\$74,999	95	1,300	20.5	(11.2 - 29.9)	5,777	501,000	25.3	(21.4 - 29.3)
≥ \$75,000	*	*	*	*	8,569	1,260,100	28.0	(25.0 - 31.0)

<sup>1</sup> Defined as having consumed 5 or more servings of fruits and vegetables in an average day

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who consume 5 or more servings of fruits and vegetables per day (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 68b. Consumption of 5 or more servings of fruits and vegetables per day among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	643	24.0	(19.4 - 28.6)	38,203	27.1	(24.9 - 29.2)
<b>Gender</b>						
Male	227	18.0	(11.0 - 25.1)	14,634	22.7	(19.6 - 25.9)
Female	416	29.4	(23.4 - 35.4)	23,569	31.0	(28.1 - 34.0)
<b>Race/Ethnicity</b>						
White Non-Hispanic	615	24.6	(20.0 - 29.1)	35,884	27.1	(25.4 - 28.9)
Black Non-Hispanic	*	*	*	740	26.8	(19.4 - 34.2)
Hispanic	*	*	*	864	29.5	(20.9 - 38.1)
Other Non-Hispanic	*	*	*	715	21.8	(12.2 - 31.5)
<b>Education</b>						
≤ High School	325	17.1	(11.4 - 22.8)	15,726	23.3	(19.3 - 27.3)
Some College	172	20.8	(13.3 - 28.3)	9,924	27.1	(22.9 - 31.2)
≥ College Degree	*	*	*	12,471	31.2	(28.0 - 34.3)
<b>Income</b>						
≤ \$24,999	194	22.0	(12.9 - 31.1)	9,667	23.9	(18.8 - 28.9)
\$25,000-\$49,999	185	25.2	(17.1 - 33.3)	9,669	32.0	(26.0 - 38.0)
\$50,000-\$74,999	95	18.9	(9.4 - 28.5)	5,777	25.0	(21.0 - 28.9)
≥ \$75,000	*	*	*	8,569	28.0	(24.9 - 31.1)

<sup>1</sup> Defined as having consumed 5 or more servings of fruits and vegetables in an average day

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 69a. Had a fall within the past 3 months among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	488	3,800	18.1	(14.1 - 22.0)	28,783	1,244,800	17.0	(14.9 - 19.0)
<b>Gender</b>								
Male	171	1,700	17.4	(11.0 - 23.8)	10,966	489,400	14.4	(11.9 - 16.8)
Female	317	2,100	18.6	(13.8 - 23.3)	17,817	755,300	19.2	(16.1 - 22.3)
<b>Age (years)</b>								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	137	1,300	18.2	(11.1 - 25.4)	7,741	445,500	16.0	(12.0 - 20.0)
55-64	121	900	16.7	(8.7 - 24.8)	8,454	336,000	16.5	(13.4 - 19.6)
≥ 65	230	1,600	18.7	(13.2 - 24.3)	12,588	463,300	18.4	(15.2 - 21.6)
<b>Race/Ethnicity</b>								
White Non-Hispanic	472	3,500	17.5	(13.6 - 21.4)	27,456	869,000	16.6	(15.1 - 18.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	424	76,100	11.0	(4.5 - 17.4)
Other Non-Hispanic	*	*	*	*	445	44,300	10.0	(3.1 - 16.8)
<b>Education</b>								
≤ High School	270	2,300	20.6	(15.2 - 26.1)	12,417	493,000	17.3	(13.3 - 21.3)
Some College	123	700	12.7	(5.9 - 19.5)	7,232	299,400	18.7	(14.6 - 22.7)
≥ College Degree	95	800	18.4	(9.2 - 27.6)	9,086	441,500	15.4	(12.9 - 17.9)
<b>Income</b>								
≤ \$24,999	166	1,000	17.3	(11.2 - 23.5)	7,824	423,900	25.0	(18.5 - 31.6)
\$25,000-\$49,999	136	1,400	22.8	(15.0 - 30.7)	7,445	223,600	16.0	(12.8 - 19.2)
\$50,000-\$74,999	*	*	*	*	4,141	151,200	14.6	(10.8 - 18.4)
≥ \$75,000	*	*	*	*	5,874	287,100	12.5	(10.0 - 15.1)

<sup>1</sup> Defined as respondent unintentionally coming to rest on the ground or another lower level one or more times within the past 3 months

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had a fall within the past 3 months (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009  
Montgomery County**

**Table 69b. Had a fall within the past 3 months among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	488	18.1	(14.1 - 22.1)	28,783	17.3	(15.0 - 19.5)
<b>Gender</b>						
Male	171	17.4	(11.0 - 23.8)	10,966	15.0	(12.1 - 17.8)
Female	317	18.7	(13.9 - 23.6)	17,817	19.4	(16.0 - 22.8)
<b>Race/Ethnicity</b>						
White Non-Hispanic	472	17.7	(13.7 - 21.7)	27,456	16.9	(15.2 - 18.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	424	10.7	(4.2 - 17.1)
Other Non-Hispanic	*	*	*	445	10.0	(2.9 - 17.0)
<b>Education</b>						
≤ High School	270	20.3	(14.8 - 25.8)	12,417	17.4	(13.0 - 21.7)
Some College	123	13.0	(5.9 - 20.0)	7,232	19.0	(14.7 - 23.2)
≥ College Degree	95	19.2	(9.9 - 28.4)	9,086	16.0	(13.1 - 18.8)
<b>Income</b>						
≤ \$24,999	166	18.0	(11.5 - 24.5)	7,824	25.3	(18.3 - 32.4)
\$25,000-\$49,999	136	22.7	(14.8 - 30.5)	7,445	15.9	(12.6 - 19.2)
\$50,000-\$74,999	*	*	*	4,141	14.7	(10.6 - 18.8)
≥ \$75,000	*	*	*	5,874	12.9	(9.9 - 15.8)

<sup>1</sup> Defined as respondent unintentionally coming to rest on the ground or another lower level one or more times within the past 3 months

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 70a. Had a fall that caused injury within the past 3 months among adults<sup>1</sup> in Montgomery County and New York State**

	Montgomery County				New York State			
	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>2</sup>	Est. # of Adults <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	488	1,200	5.7	(3.6 - 7.8)	28,760	335,700	4.6	(3.7 - 5.5)
<b>Gender</b>								
Male	*	*	*	*	10,957	120,000	3.5	(2.5 - 4.6)
Female	317	1,000	8.4	(5.0 - 11.8)	17,803	215,600	5.5	(4.1 - 6.9)
<b>Age (years)</b>								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	7,737	115,100	4.1	(2.9 - 5.4)
55-64	*	*	*	*	8,449	90,300	4.4	(2.7 - 6.1)
≥ 65	230	600	7.6	(3.8 - 11.4)	12,574	130,300	5.2	(3.4 - 6.9)
<b>Race/Ethnicity</b>								
White Non-Hispanic	472	1,000	5.1	(3.2 - 7.1)	27,433	270,800	5.2	(4.2 - 6.1)
Black Non-Hispanic	*	*	*	*	458	47,900	4.9	(0.9 - 9.0)
Hispanic	*	*	*	*	424	12,400	1.8	(0.2 - 3.3)
Other Non-Hispanic	*	*	*	*	445	4,600	1.0	(0.4 - 1.6)
<b>Education</b>								
≤ High School	270	900	8.2	(4.8 - 11.7)	12,404	148,900	5.2	(3.6 - 6.8)
Some College	*	*	*	*	7,228	85,400	5.3	(3.1 - 7.5)
≥ College Degree	*	*	*	*	9,081	101,300	3.5	(2.4 - 4.6)
<b>Income</b>								
≤ \$24,999	166	400	7.0	(2.8 - 11.1)	7,812	90,200	5.3	(3.6 - 7.1)
\$25,000-\$49,999	136	500	8.3	(3.7 - 12.8)	7,442	93,200	6.7	(3.9 - 9.4)
\$50,000-\$74,999	*	*	*	*	4,139	37,000	3.6	(1.8 - 5.3)
≥ \$75,000	*	*	*	*	5,872	54,700	2.4	(1.5 - 3.3)

<sup>1</sup> Defined as as respondent unintentionally coming to rest on the ground or another lower level and caused respondent to limit regular activities for at least a day or to go see a doctor one or more times within the past 3 months

<sup>2</sup> Sample size from survey

<sup>3</sup> Estimated number of adults who have had a fall that caused an injury within the past 3 months (rounded to the nearest hundred)

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009**  
**Montgomery County**

**Table 70b. Had a fall that caused injury within the past 3 months among adults<sup>1</sup> in Montgomery County and New York State - Age-Adjusted<sup>2</sup>**

	Montgomery County			New York State		
	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>	n <sup>3</sup>	Percent	(C.I.) <sup>4</sup>
<b>Total</b>	488	5.5	(3.5 - 7.5)	28,760	4.6	(3.6 - 5.5)
<b>Gender</b>						
Male	*	*	*	10,957	3.6	(2.4 - 4.7)
Female	317	8.3	(5.0 - 11.6)	17,803	5.5	(4.1 - 6.9)
<b>Race/Ethnicity</b>						
White Non-Hispanic	472	5.0	(3.1 - 6.9)	27,433	5.2	(4.2 - 6.2)
Black Non-Hispanic	*	*	*	458	4.7	(0.7 - 8.7)
Hispanic	*	*	*	424	1.8	(0.1 - 3.4)
Other Non-Hispanic	*	*	*	445	1.0	(0.4 - 1.6)
<b>Education</b>						
≤ High School	270	7.8	(4.6 - 11.0)	12,404	5.2	(3.5 - 6.8)
Some College	*	*	*	7,228	5.3	(3.1 - 7.5)
≥ College Degree	*	*	*	9,081	3.6	(2.4 - 4.7)
<b>Income</b>						
≤ \$24,999	166	7.4	(2.9 - 11.9)	7,812	5.1	(3.3 - 7.0)
\$25,000-\$49,999	136	7.7	(3.4 - 12.0)	7,442	6.6	(3.9 - 9.2)
\$50,000-\$74,999	*	*	*	4,139	3.7	(1.7 - 5.8)
≥ \$75,000	*	*	*	5,872	2.4	(1.4 - 3.3)

<sup>1</sup> Defined as as respondent unintentionally coming to rest on the ground or another lower level and caused respondent to limit regular activities for at least a day or to go see a doctor one or more times within the past 3 months

<sup>2</sup> Prevalence estimates are age-adjusted to the 2000 US Standard population

<sup>3</sup> Sample size from survey

<sup>4</sup> 95% confidence interval

\* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)