

New York State Expanded Behavioral Risk Factor Surveillance System

Schoharie County

**Final Report
July 2008 - June 2009**

**New York State Department of Health
December 2009**

Contents

Introduction	12
Table 1. Schoharie County and New York State adult US Census Bureau population estimates, 2007	13
General Health	
Table 2a. Fair or poor general health among adults in Schoharie County and New York State	14
Table 2b. Fair or poor general health among adults in Schoharie County and New York State - Age-adjusted	15
Table 3a. Poor physical health 14 or more days within the past month among adults in Schoharie County and New York State	16
Table 3b. Poor physical health 14 or more days within the past month among adults in Schoharie County and New York State - Age-adjusted	17
Table 4a. Poor mental health 14 or more days within the past month among adults in Schoharie County and New York State	18
Table 4b. Poor mental health 14 or more days within the past month among adults in Schoharie County and New York State - Age-adjusted	19
Table 5a. Poor physical or mental health 14 or more days within the past month among adults in Schoharie County and New York State	20
Table 5b. Poor physical or mental health 14 or more days within the past month among adults in Schoharie County and New York State - Age-adjusted	21
Quality Health Care	
Table 6a. Health insurance among adults in Schoharie County and New York State	22
Table 6b. Health insurance among adults in Schoharie County and New York State - Age-adjusted	23
Table 7a. Health insurance among adults aged 18-64 in Schoharie County and New York State	24
Table 7b. Health insurance among adults aged 18-64 in Schoharie County and New York State - Age-adjusted	25

Contents

Table 8a. No health insurance among adults in Schoharie County and New York State	26
Table 8b. No health insurance among adults in Schoharie County and New York State - Age-adjusted	27
Table 9a. No health insurance among adults aged 18-64 in Schoharie County and New York State	28
Table 9b. No health insurance among adults aged 18-64 in Schoharie County and New York State - Age-adjusted	29
Table 10a. Adults with regular health care provider in Schoharie County and New York State	30
Table 10b. Adults with regular health care provider in Schoharie County and New York State - Age-adjusted	31
Table 11a. Cost prevented visit to doctor within the past year among adults in Schoharie County and New York State	32
Table 11b. Cost prevented visit to doctor within the past year among adults in Schoharie County and New York State - Age-adjusted	33
Table 12a. Visited a doctor for routine checkup within the past year among adults in Schoharie County and New York State	34
Table 12b. Visited a doctor for routine checkup within the past year among adults in Schoharie County and New York State - Age-adjusted	35
Table 13a. Visited a doctor for routine checkup within the past 2 years among adults in Schoharie County and New York State	36
Table 13b. Visited a doctor for routine checkup within the past 2 years among adults in Schoharie County and New York State - Age-adjusted	37
Table 14a. Dentist visit within the past year among adults in Schoharie County and New York State	38
Table 14b. Dentist visit within the past year among adults in Schoharie County and New York State - Age-adjusted	39
Table 15a. Had teeth cleaned within the past year among adults in Schoharie County and New York State	40
Table 15b. Had teeth cleaned within the past year among adults in Schoharie County and New York State - Age-adjusted	41

Contents

Table 16a. Permanent teeth extracted due to decay or gum disease among adults in Schoharie County and New York State	42
Table 16b. Permanent teeth extracted due to decay or gum disease among adults in Schoharie County and New York State - Age-adjusted	43
Table 17a. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in Schoharie County and New York State	44
Table 17b. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in Schoharie County and New York State - Age-adjusted	45
Table 18a. Flu shot within the past 12 months in Schoharie County and New York State	46
Table 18b. Flu shot within the past 12 months in Schoharie County and New York State - Age-adjusted	47
Table 19a. Flu vaccine that was sprayed in nose within the past 12 months in Schoharie County and New York State	48
Table 19b. Flu vaccine that was sprayed in nose within the past 12 months in Schoharie County and New York State - Age-adjusted	49
Table 20a. Flu shot or flu vaccine sprayed in nose within the past 12 months in Schoharie County and New York State	50
Table 20b. Flu shot or flu vaccine sprayed in nose within the past 12 months in Schoharie County and New York State - Age-adjusted	51
Table 21a. Flu shot among adults aged 65 and older in Schoharie County and New York State	52
Table 21b. Flu shot among adults aged 65 and older in Schoharie County and New York State - Age-adjusted	53
Table 22a. Pneumonia shot or pneumococcal vaccine in Schoharie County and New York State	54
Table 22b. Pneumonia shot or pneumococcal vaccine in Schoharie County and New York State - Age-adjusted	55
Table 23a. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in Schoharie County and New York State	56
Table 23b. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in Schoharie County and New York State - Age-adjusted	57

Contents

Cancer Screening

Table 24a. Ever had a mammogram among women aged 40 and older in Schoharie County and New York State	58
Table 24b. Ever had a mammogram among women aged 40 and older in Schoharie County and New York State - Age-adjusted	59
Table 25a. Had mammogram within the past 2 years among women aged 40 and older in Schoharie County and New York State	60
Table 25b. Had mammogram within the past 2 years among women aged 40 and older in Schoharie County and New York State - Age-adjusted	61
Table 26a. Had mammogram within the past 2 years among women aged 50 and older in Schoharie County and New York State	62
Table 26b. Had mammogram within the past 2 years among women aged 50 and older in Schoharie County and New York State - Age-adjusted	63
Table 27a. Ever had a Pap test among women in Schoharie County and New York State	64
Table 27b. Ever had a Pap test among women in Schoharie County and New York State - Age-adjusted	65
Table 28a. Had a Pap test within the past 3 years among women in Schoharie County and New York State	66
Table 28b. Had a Pap test within the past 3 years among women in Schoharie County and New York State - Age-adjusted	67
Table 29a. Ever had a digital rectal exam among men aged 40 and older in Schoharie County and New York State	68
Table 29b. Ever had a digital rectal exam among men aged 40 and older in Schoharie County and New York State - Age-adjusted	69
Table 30a. Had digital rectal exam within the past 2 years among men aged 40 and older in Schoharie County and New York State	70
Table 30b. Had digital rectal exam within the past 2 years among men aged 40 and older in Schoharie County and New York State - Age-adjusted	71

Contents

Table 31a. Ever had a prostate specific antigen test among men aged 40 and older in Schoharie County and New York State	72
Table 31b. Ever had a prostate specific antigen test among men aged 40 and older in Schoharie County and New York State - Age-adjusted	73
Table 32a. Had prostate specific antigen test within the past 2 years among men aged 40 and older in Schoharie County and New York State	74
Table 32b. Had prostate specific antigen test within the past 2 years among men aged 40 and older in Schoharie County and New York State - Age-adjusted	75
Table 33a. Prostate cancer among men aged 40 and older in Schoharie County and New York State	76
Table 33b. Prostate cancer among men aged 40 and older in Schoharie County and New York State - Age-adjusted	77
Table 34a. Home blood stool test ever used among adults aged 50 and older in Schoharie County and New York State	78
Table 34b. Home blood stool test ever used among adults aged 50 and older in Schoharie County and New York State - Age-adjusted	79
Table 35a. Home blood stool test used within the past year among adults aged 50 and older in Schoharie County and New York State	80
Table 35b. Home blood stool test used within the past year among adults aged 50 and older in Schoharie County and New York State - Age-adjusted	81
Table 36a. Home blood stool test used within the past 2 years among adults aged 50 and older in Schoharie County and New York State	82
Table 36b. Home blood stool test used within the past 2 years among adults aged 50 and older in Schoharie County and New York State - Age-adjusted	83
Table 37a. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Schoharie County and New York State	84
Table 37b. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Schoharie County and New York State - Age-adjusted	85
Table 38a. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Schoharie County and New York State	86

Contents

Table 38b. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Schoharie County and New York State - Age-adjusted	87
Table 39a. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Schoharie County and New York State	88
Table 39b. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Schoharie County and New York State - Age-adjusted	89
Chronic Diseases	
Table 40a. High blood pressure among adults in Schoharie County and New York State	90
Table 40b. High blood pressure among adults in Schoharie County and New York State - Age-adjusted	91
Table 41a. Blood pressure medication among adults with high blood pressure in Schoharie County and New York State	92
Table 41b. Blood pressure medication among adults with high blood pressure in Schoharie County and New York State - Age-adjusted	93
Table 42a. Ever had blood cholesterol checked among adults in Schoharie County and New York State	94
Table 42b. Ever had blood cholesterol checked among adults in Schoharie County and New York State - Age-adjusted	95
Table 43a. Blood cholesterol checked within the past 5 years among adults in Schoharie County and New York State	96
Table 43b. Blood cholesterol checked within the past 5 years among adults in Schoharie County and New York State - Age-adjusted	97
Table 44a. Cardiovascular disease among adults in Schoharie County and New York State	98
Table 44b. Cardiovascular disease among adults in Schoharie County and New York State - Age-adjusted	99
Table 45a. Coronary heart disease among adults in Schoharie County and New York State	100

Contents

Table 45b. Coronary heart disease among adults in Schoharie County and New York State - Age-adjusted	101
Table 46a. Ever diagnosed with diabetes among adults in Schoharie County and New York State	102
Table 46b. Ever diagnosed with diabetes among adults in Schoharie County and New York State - Age-adjusted	103
Table 47a. Ever diagnosed with asthma (lifetime) among adults in Schoharie County and New York State	104
Table 47b. Ever diagnosed with asthma (lifetime) among adults in Schoharie County and New York State - Age-adjusted	105
Table 48a. Current asthma among adults in Schoharie County and New York State	106
Table 48b. Current asthma among adults in Schoharie County and New York State - Age-adjusted	107
Table 49a. Chronic joint symptoms among adults in Schoharie County and New York State	108
Table 49b. Chronic joint symptoms among adults in Schoharie County and New York State - Age-adjusted	109
Table 50a. Arthritis among adults in Schoharie County and New York State	110
Table 50b. Arthritis among adults in Schoharie County and New York State - Age-adjusted	111
Table 51a. Activity limitations because of physical, mental or emotional problems among adults in Schoharie County and New York State	112
Table 51b. Activity limitations because of physical, mental or emotional problems among adults in Schoharie County and New York State - Age-adjusted	113
Table 52a. Have health problem that requires the use of special equipment among adults in Schoharie County and New York State	114
Table 52b. Have health problem that requires the use of special equipment among adults in Schoharie County and New York State - Age-adjusted	115
Table 53a. Disability among adults in Schoharie County and New York State	116
Table 53b. Disability among adults in Schoharie County and New York State - Age-adjusted	117

Contents

Health Behaviors

Table 54a. Current smoking among adults in Schoharie County and New York State	118
Table 54b. Current smoking among adults in Schoharie County and New York State - Age-adjusted	119
Table 55a. Everyday smoking among adults in Schoharie County and New York State	120
Table 55b. Everyday smoking among adults in Schoharie County and New York State - Age-adjusted	121
Table 56a. Adults living in homes in which smoking is prohibited in Schoharie County and New York State	122
Table 56b. Adults living in homes in which smoking is prohibited in Schoharie County and New York State - Age-adjusted	123
Table 57a. Binge drinking within the past month among adults in Schoharie County and New York State	124
Table 57b. Binge drinking within the past month among adults in Schoharie County and New York State - Age-adjusted	125
Table 58a. Heavy drinking within the past month among adults in Schoharie County and New York State	126
Table 58b. Heavy drinking within the past month among adults in Schoharie County and New York State - Age-adjusted	127
Table 59a. Never or rarely been asked about sexual history during routine check-up among adults in Schoharie County and New York State	128
Table 59b. Never or rarely been asked about sexual history during routine check-up among adults in Schoharie County and New York State - Age-adjusted	129
Table 60a. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults in Schoharie County and New York State	130
Table 60b. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults in Schoharie County and New York State - Age-adjusted	131

Contents

Table 61a. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults in Schoharie County and New York State	132
Table 61b. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults in Schoharie County and New York State - Age-adjusted	133

Overweight / Obesity

Table 62a. Overweight among adults in Schoharie County and New York State	134
Table 62b. Overweight among adults in Schoharie County and New York State - Age-adjusted	135
Table 63a. Obesity among adults in Schoharie County and New York State	136
Table 63b. Obesity among adults in Schoharie County and New York State - Age-adjusted	137
Table 64a. Overweight or obesity among adults in Schoharie County and New York State	138
Table 64b. Overweight or obesity among adults in Schoharie County and New York State - Age-adjusted	139
Table 65a. Received advice about weight by a health professional among adults in Schoharie County and New York State	140
Table 65b. Received advice about weight by a health professional among adults in Schoharie County and New York State - Age-adjusted	141
Table 66a. Received advice to lose weight by a health professional among adults provided weight advice in Schoharie County and New York State	142
Table 66b. Received advice to lose weight by a health professional among adults provided weight advice in Schoharie County and New York State - Age-adjusted	143

Physical Activity / Nutrition

Table 67a. No leisure-time physical activity among adults in Schoharie County and New York State	144
Table 67b. No leisure-time physical activity among adults in Schoharie County and New York State - Age-adjusted	145

Contents

Table 68a. Consumption of 5 or more servings of fruits and vegetables per day among adults in Schoharie County and New York State	146
Table 68b. Consumption of 5 or more servings of fruits and vegetables per day among adults in Schoharie County and New York State - Age-adjusted	147

Injury

Table 69a. Had a fall within the past 3 months among adults in Schoharie County and New York State	148
Table 69b. Had a fall within the past 3 months among adults in Schoharie County and New York State - Age-adjusted	149
Table 70a. Had a fall that caused injury within the past 3 months among adults in Schoharie County and New York State	150
Table 70b. Had a fall that caused injury within the past 3 months among adults in Schoharie County and New York State - Age-adjusted	151

Introduction

The New York State Expanded Behavioral Risk Factor Surveillance System (Expanded BRFSS) is a random digit dialed telephone survey of adults (18 years of age and older) that is representative of the non-institutionalized civilian population with landline telephones living in New York State. The goal of the Expanded BRFSS is to collect county-specific data on preventive health practices, risk behaviors, injuries and preventable chronic and infectious diseases. Topics assessed by the Expanded BRFSS include tobacco use, physical inactivity, diet, use of cancer screening services and other factors linked to the leading causes of morbidity and mortality. For each topic, health indicators are based on answers to one or more questions in the survey. The survey results are weighted to more accurately represent the adult population living in each county. Weighting procedures adjust for the differential selection probabilities by race, age and gender. More detailed information on weighting procedures is available in the technical brief.

Survey data collection occurred in 58 geographic areas of New York State, including each of the 57 counties outside of New York City and New York City (5 counties) as a single area. A standard questionnaire was used in all areas. Data collection for the 2009 Expanded BRFSS took place during the period of July 2008 through June 2009. The goal was to complete about 650 interviews in each area. The final number of interviews varies because of the sampling design of the survey and differential response rates. More detailed information on the sampling design is available in the technical brief.

The New York State Department of Health contracted with RTI International, which is responsible for the data collection, sampling, weighting and creation of the statistical reports. RTI International subcontracted with Clearwater Research Inc. for data collection and additional expertise in sample design and data quality control.

This report presents both crude estimates and estimates that have been age-adjusted to the Year 2000 US Standard Population when they represent the entire adult population of an area. Estimates for specific age groups are not age-adjusted. Age adjustment is a statistical process applied to rates that allows communities with different age structures to be compared. Counties can compare their data with results from New York State as a whole and with other counties. Confidence intervals at the 95% level associated with each health indicator give an estimated range of values which is likely to include the actual prevalence of the indicator. Confidence intervals are similar to margins of error. When the confidence intervals of two estimates of the same indicator from different areas do not overlap, they may be said to be statistically significantly different, i.e., these differences are unlikely related to chance and are considered true differences. If there is any value that is included in both intervals, the two estimates are not statistically significantly different.

The survey results for each of the 69 health indicators included in the survey are typically provided by gender, age, race and ethnicity, education and income in the Expanded BRFSS county reports. The estimates for some of these groupings may be suppressed in geographic areas with low proportions of adult respondents in these groupings.

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 1. Schoharie County and New York State adult US Census Bureau population estimates,¹ 2007

	Schoharie County		New York State	
	Adult Population ¹	Percent	Adult Population ¹	Percent
Total	25,658	100	14,884,315	100
Gender				
Male	12,703	49.5	7,103,791	47.7
Female	12,955	50.5	7,780,524	52.3
Age (years)				
18-34	7,886	30.7	4,497,479	30.2
35-44	4,107	16.0	2,844,115	19.1
45-54	4,784	18.6	2,854,316	19.2
55-64	4,010	15.6	2,142,000	14.4
≥ 65	4,871	19.0	2,546,405	17.1
Race/Ethnicity				
White Non-Hispanic	24,469	95.4	9,343,884	62.8
Black Non-Hispanic	397	1.5	2,156,941	14.5
Hispanic	562	2.2	2,258,717	15.2
Other Non-Hispanic	230	0.9	1,124,773	7.6

¹ Source: National Center for Health Statistics. Postcensal estimates of the resident population of the United States for July 1, 2000-July 1, 2007, by year, county, age, bridged race, Hispanic origin, and sex (Vintage 2007). Prepared under a collaborative arrangement with the U.S. Census Bureau; released August 7, 2008. Available from: http://www.cdc.gov/nchs/nvss/bridged_race.htm as of November 25, 2009.

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 2a. Fair or poor general health among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	643	3,400	13.5	(10.3 - 16.8)	38,124	2,531,400	17.1	(15.1 - 19.0)
Gender								
Male	242	1,500	11.9	(7.0 - 16.8)	14,599	1,069,700	15.1	(12.4 - 17.8)
Female	401	2,000	15.1	(10.9 - 19.3)	23,525	1,461,700	18.8	(16.1 - 21.6)
Age (years)								
18-34	*	*	*	*	3,940	535,800	12.0	(7.2 - 16.8)
35-44	*	*	*	*	4,985	289,200	10.0	(6.8 - 13.3)
45-54	141	500	9.7	(4.6 - 14.8)	7,819	507,800	17.9	(14.0 - 21.8)
55-64	163	800	20.1	(13.3 - 27.0)	8,757	424,700	19.9	(16.0 - 23.9)
≥ 65	217	1,100	22.6	(16.7 - 28.5)	12,623	773,900	30.5	(26.8 - 34.2)
Race/Ethnicity								
White Non-Hispanic	628	3,400	13.7	(10.4 - 17.0)	35,811	1,226,600	13.2	(11.7 - 14.6)
Black Non-Hispanic	*	*	*	*	739	457,300	21.3	(14.8 - 27.9)
Hispanic	*	*	*	*	861	684,100	30.4	(22.2 - 38.6)
Other Non-Hispanic	*	*	*	*	713	163,400	14.5	(5.1 - 24.0)
Education								
≤ High School	302	2,100	17.4	(11.9 - 23.0)	15,682	1,466,500	26.3	(22.6 - 30.0)
Some College	173	800	10.8	(6.0 - 15.6)	9,904	508,500	14.3	(10.9 - 17.8)
≥ College Degree	167	500	9.0	(4.0 - 14.1)	12,457	493,500	8.8	(6.5 - 11.1)
Income								
≤ \$24,999	171	1,300	23.8	(15.5 - 32.0)	9,633	1,269,200	36.3	(30.1 - 42.4)
\$25,000-\$49,999	198	1,200	13.9	(8.0 - 19.8)	9,649	447,600	15.4	(11.9 - 18.9)
\$50,000-\$74,999	*	*	*	*	5,769	237,800	12.1	(8.0 - 16.2)
≥ \$75,000	*	*	*	*	8,562	218,300	4.9	(3.6 - 6.1)

¹ Defined as reporting fair or poor general health

² Sample size from survey

³ Estimated number of adults who have fair or poor health (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 2b. Fair or poor general health among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	643	12.8	(9.5 - 16.2)	38,124	16.7	(14.8 - 18.6)
Gender						
Male	242	11.4	(6.3 - 16.5)	14,599	15.3	(12.6 - 18.0)
Female	401	14.2	(9.9 - 18.6)	23,525	18.0	(15.2 - 20.7)
Race/Ethnicity						
White Non-Hispanic	628	13.0	(9.6 - 16.4)	35,811	12.9	(11.4 - 14.4)
Black Non-Hispanic	*	*	*	739	20.8	(14.2 - 27.3)
Hispanic	*	*	*	861	29.9	(21.8 - 37.9)
Other Non-Hispanic	*	*	*	713	13.8	(4.8 - 22.8)
Education						
≤ High School	302	16.6	(10.8 - 22.4)	15,682	25.8	(22.1 - 29.5)
Some College	173	10.0	(5.1 - 14.8)	9,904	14.1	(10.7 - 17.6)
≥ College Degree	167	8.6	(3.6 - 13.7)	12,457	8.5	(6.3 - 10.8)
Income						
≤ \$24,999	171	21.9	(13.4 - 30.3)	9,633	35.9	(29.7 - 42.0)
\$25,000-\$49,999	198	13.6	(7.4 - 19.8)	9,649	15.0	(11.6 - 18.5)
\$50,000-\$74,999	*	*	*	5,769	11.8	(7.7 - 16.0)
≥ \$75,000	*	*	*	8,562	4.7	(3.5 - 6.0)

¹ Defined as reporting fair or poor general health

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 3a. Poor physical health 14 or more days within the past month among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	625	2,700	10.9	(7.4 - 14.5)	37,327	1,472,900	10.1	(8.7 - 11.6)
Gender								
Male	235	1,600	13.0	(6.6 - 19.4)	14,325	604,200	8.7	(7.0 - 10.5)
Female	390	1,100	8.9	(5.9 - 12.0)	23,002	868,600	11.4	(9.2 - 13.6)
Age (years)								
18-34	*	*	*	*	3,891	268,100	6.1	(2.7 - 9.5)
35-44	*	*	*	*	4,948	232,100	8.1	(5.2 - 11.1)
45-54	*	*	*	*	7,764	350,900	12.5	(9.4 - 15.6)
55-64	156	600	15.8	(9.6 - 22.0)	8,621	271,200	13.0	(10.3 - 15.8)
≥ 65	206	700	14.2	(8.8 - 19.5)	12,103	350,600	14.5	(12.1 - 16.9)
Race/Ethnicity								
White Non-Hispanic	611	2,600	10.8	(7.2 - 14.3)	35,073	929,700	10.2	(8.8 - 11.6)
Black Non-Hispanic	*	*	*	*	724	208,200	9.9	(4.8 - 14.9)
Hispanic	*	*	*	*	835	250,300	11.5	(6.5 - 16.4)
Other Non-Hispanic	*	*	*	*	695	84,600	7.6	(1.3 - 13.9)
Education								
≤ High School	288	1,900	15.9	(9.6 - 22.3)	15,178	692,800	12.9	(10.6 - 15.3)
Some College	171	600	8.6	(3.2 - 14.1)	9,751	362,500	10.3	(7.4 - 13.2)
≥ College Degree	165	200	4.0	(1.2 - 6.8)	12,327	359,300	6.4	(4.8 - 8.1)
Income								
≤ \$24,999	162	1,000	18.4	(8.9 - 27.9)	9,294	625,600	18.4	(13.6 - 23.3)
\$25,000-\$49,999	198	1,200	13.5	(7.1 - 20.0)	9,495	279,100	9.8	(7.3 - 12.3)
\$50,000-\$74,999	*	*	*	*	5,733	186,400	9.5	(5.8 - 13.2)
≥ \$75,000	*	*	*	*	8,510	219,100	4.9	(3.6 - 6.2)

¹ Defined as 14 or more days within the past month when physical health was not good

² Sample size from survey

³ Estimated number of adults with poor physical health 14 or more days within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 3b. Poor physical health 14 or more days within the past month among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	625	10.8	(7.0 - 14.6)	37,327	9.9	(8.4 - 11.4)
Gender						
Male	235	13.4	(6.5 - 20.3)	14,325	8.7	(6.9 - 10.5)
Female	390	8.2	(5.0 - 11.5)	23,002	11.0	(8.7 - 13.2)
Race/Ethnicity						
White Non-Hispanic	611	10.5	(6.7 - 14.4)	35,073	9.9	(8.5 - 11.3)
Black Non-Hispanic	*	*	*	724	9.7	(4.6 - 14.9)
Hispanic	*	*	*	835	11.3	(6.3 - 16.2)
Other Non-Hispanic	*	*	*	695	7.6	(1.2 - 14.0)
Education						
≤ High School	288	15.9	(9.0 - 22.8)	15,178	12.7	(10.2 - 15.1)
Some College	171	8.4	(2.6 - 14.1)	9,751	10.1	(7.2 - 12.9)
≥ College Degree	165	3.6	(1.0 - 6.2)	12,327	6.3	(4.7 - 8.0)
Income						
≤ \$24,999	*	*	*	9,294	18.4	(13.5 - 23.4)
\$25,000-\$49,999	198	13.9	(6.9 - 20.9)	9,495	9.4	(7.0 - 11.9)
\$50,000-\$74,999	*	*	*	5,733	9.0	(5.3 - 12.6)
≥ \$75,000	*	*	*	8,510	4.9	(3.5 - 6.2)

¹ Defined as 14 or more days within the past month when physical health was not good

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 4a. Poor mental health 14 or more days within the past month among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	641	2,500	10.1	(6.9 - 13.2)	37,528	1,473,200	10.1	(8.6 - 11.5)
Gender								
Male	241	800	6.1	(1.9 - 10.3)	14,407	555,200	7.9	(6.3 - 9.6)
Female	400	1,800	14.0	(9.2 - 18.7)	23,121	918,000	12.0	(9.7 - 14.3)
Age (years)								
18-34	*	*	*	*	3,889	496,700	11.2	(7.4 - 15.1)
35-44	74	900	15.6	(6.2 - 25.0)	4,947	254,100	8.9	(6.7 - 11.1)
45-54	139	400	8.1	(3.5 - 12.7)	7,749	347,200	12.3	(9.0 - 15.6)
55-64	164	400	9.8	(5.2 - 14.3)	8,623	211,300	10.1	(7.9 - 12.3)
≥ 65	217	300	6.6	(2.9 - 10.3)	12,320	163,900	6.7	(4.7 - 8.7)
Race/Ethnicity								
White Non-Hispanic	626	2,500	10.1	(6.8 - 13.4)	35,266	899,200	9.8	(8.4 - 11.2)
Black Non-Hispanic	*	*	*	*	722	227,800	10.9	(5.4 - 16.4)
Hispanic	*	*	*	*	845	263,700	11.8	(6.8 - 16.8)
Other Non-Hispanic	*	*	*	*	695	82,500	7.4	(1.4 - 13.5)
Education								
≤ High School	301	1,400	11.5	(6.5 - 16.5)	15,335	665,000	12.2	(10.0 - 14.5)
Some College	174	600	8.7	(3.7 - 13.6)	9,771	404,500	11.5	(8.3 - 14.8)
≥ College Degree	165	500	8.8	(2.1 - 15.6)	12,342	346,600	6.2	(4.5 - 7.9)
Income								
≤ \$24,999	169	800	14.0	(6.5 - 21.4)	9,398	611,500	17.9	(13.0 - 22.8)
\$25,000-\$49,999	200	1,000	11.6	(5.0 - 18.1)	9,547	260,300	9.0	(6.7 - 11.4)
\$50,000-\$74,999	*	*	*	*	5,726	187,300	9.6	(6.0 - 13.1)
≥ \$75,000	*	*	*	*	8,520	276,500	6.2	(4.6 - 7.7)

¹ Defined as 14 or more days within the past month when mental health was not good

² Sample size from survey

³ Estimated number of adults with poor mental health 14 or more days within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 4b. Poor mental health 14 or more days within the past month among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	641	10.4	(7.0 - 13.9)	37,528	10.2	(8.7 - 11.7)
Gender						
Male	241	6.2	(1.7 - 10.7)	14,407	7.9	(6.3 - 9.6)
Female	400	14.6	(9.4 - 19.8)	23,121	12.2	(9.8 - 14.6)
Race/Ethnicity						
White Non-Hispanic	626	10.5	(7.0 - 14.1)	35,266	10.0	(8.5 - 11.4)
Black Non-Hispanic	*	*	*	722	11.1	(5.5 - 16.7)
Hispanic	*	*	*	845	11.5	(6.5 - 16.4)
Other Non-Hispanic	*	*	*	695	7.6	(1.5 - 13.7)
Education						
≤ High School	301	12.0	(6.6 - 17.4)	15,335	12.3	(10.0 - 14.6)
Some College	174	9.3	(3.8 - 14.7)	9,771	11.8	(8.5 - 15.1)
≥ College Degree	165	8.8	(1.4 - 16.2)	12,342	6.3	(4.6 - 8.0)
Income						
≤ \$24,999	169	14.2	(6.1 - 22.3)	9,398	18.2	(13.2 - 23.1)
\$25,000-\$49,999	200	11.8	(4.8 - 18.7)	9,547	9.4	(6.9 - 11.9)
\$50,000-\$74,999	*	*	*	5,726	9.5	(5.9 - 13.1)
≥ \$75,000	*	*	*	8,520	6.2	(4.6 - 7.7)

¹ Defined as 14 or more days within the past month when mental health was not good

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 5a. Poor physical or mental health 14 or more days within the past month among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	623	4,600	18.4	(14.0 - 22.8)	37,005	2,385,900	16.5	(14.8 - 18.2)
Gender								
Male	233	2,000	16.7	(9.6 - 23.8)	14,231	937,700	13.6	(11.4 - 15.8)
Female	390	2,500	20.0	(14.7 - 25.3)	22,774	1,448,200	19.1	(16.5 - 21.7)
Age (years)								
18-34	*	*	*	*	3,859	602,600	13.8	(9.7 - 17.9)
35-44	*	*	*	*	4,924	388,300	13.7	(10.4 - 17.0)
45-54	139	500	10.0	(5.0 - 15.0)	7,722	551,000	19.6	(15.8 - 23.4)
55-64	156	700	19.4	(12.9 - 26.0)	8,545	395,500	19.2	(16.1 - 22.3)
≥ 65	208	800	16.5	(10.9 - 22.1)	11,955	448,500	18.8	(15.9 - 21.7)
Race/Ethnicity								
White Non-Hispanic	609	4,500	18.4	(13.9 - 22.9)	34,778	1,485,700	16.3	(14.8 - 17.8)
Black Non-Hispanic	*	*	*	*	715	343,700	16.5	(10.3 - 22.7)
Hispanic	*	*	*	*	832	412,900	19.0	(12.6 - 25.3)
Other Non-Hispanic	*	*	*	*	680	143,500	13.0	(5.0 - 21.0)
Education								
≤ High School	287	2,800	23.9	(16.6 - 31.2)	15,018	1,077,200	20.3	(17.3 - 23.2)
Some College	172	1,200	15.7	(8.6 - 22.7)	9,672	615,500	17.6	(14.1 - 21.1)
≥ College Degree	163	600	11.0	(3.9 - 18.1)	12,244	634,200	11.4	(9.2 - 13.6)
Income								
≤ \$24,999	*	*	*	*	9,199	924,000	27.3	(21.9 - 32.8)
\$25,000-\$49,999	198	2,000	23.6	(15.1 - 32.1)	9,425	456,000	16.1	(13.0 - 19.2)
\$50,000-\$74,999	76	400	11.5	(4.0 - 19.0)	5,696	312,000	16.0	(11.4 - 20.5)
≥ \$75,000	*	*	*	*	8,475	447,600	10.0	(8.1 - 12.0)

¹ Defined as 14 or more days within the past month when physical or mental health prevented usual activities, such as self-care, work or recreation

² Sample size from survey

³ Estimated number of adults with poor physical or mental health 14 or more days within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 5b. Poor physical or mental health 14 or more days within the past month among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	623	18.9	(14.1 - 23.7)	37,005	16.4	(14.7 - 18.2)
Gender						
Male	233	17.3	(9.6 - 25.0)	14,231	13.6	(11.4 - 15.8)
Female	390	20.5	(14.7 - 26.3)	22,774	19.0	(16.3 - 21.6)
Race/Ethnicity						
White Non-Hispanic	609	18.9	(14.0 - 23.8)	34,778	16.3	(14.7 - 17.8)
Black Non-Hispanic	*	*	*	715	16.7	(10.3 - 23.0)
Hispanic	*	*	*	832	18.5	(12.2 - 24.8)
Other Non-Hispanic	*	*	*	680	13.1	(5.0 - 21.3)
Education						
≤ High School	287	24.5	(16.6 - 32.5)	15,018	20.1	(17.1 - 23.1)
Some College	172	16.3	(8.7 - 23.9)	9,672	17.6	(14.1 - 21.1)
≥ College Degree	163	10.9	(3.2 - 18.6)	12,244	11.4	(9.2 - 13.7)
Income						
≤ \$24,999	*	*	*	9,199	27.6	(22.1 - 33.1)
\$25,000-\$49,999	198	24.5	(15.4 - 33.6)	9,425	16.1	(12.9 - 19.3)
\$50,000-\$74,999	76	11.4	(3.5 - 19.2)	5,696	15.6	(11.0 - 20.1)
≥ \$75,000	*	*	*	8,475	10.0	(8.0 - 12.0)

¹ Defined as 14 or more days within the past month when physical or mental health prevented usual activities, such as self-care, work or recreation

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 6a. Health insurance among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	644	20,900	82.7	(77.6 - 87.8)	38,119	12,854,200	86.6	(84.2 - 89.0)
Gender								
Male	243	9,900	79.0	(70.5 - 87.4)	14,597	5,881,100	83.0	(78.9 - 87.1)
Female	401	11,000	86.4	(80.7 - 92.1)	23,522	6,973,100	89.9	(87.2 - 92.6)
Age (years)								
18-34	*	*	*	*	3,910	3,302,100	74.3	(67.8 - 80.7)
35-44	*	*	*	*	4,986	2,560,300	88.9	(84.7 - 93.1)
45-54	142	4,000	84.6	(77.2 - 92.1)	7,831	2,530,100	88.9	(84.6 - 93.2)
55-64	164	3,600	90.2	(84.6 - 95.8)	8,764	1,986,700	92.8	(90.2 - 95.5)
≥ 65	217	4,700	97.7	(95.4 - 99.9)	12,628	2,475,000	97.8	(96.7 - 98.8)
Race/Ethnicity								
White Non-Hispanic	629	20,500	83.5	(78.4 - 88.6)	35,811	8,591,200	92.2	(90.6 - 93.8)
Black Non-Hispanic	*	*	*	*	736	1,781,600	82.7	(75.3 - 90.1)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	302	9,600	80.3	(72.8 - 87.9)	15,673	4,448,900	79.7	(74.9 - 84.5)
Some College	*	*	*	*	9,906	3,004,600	85.0	(80.1 - 89.8)
≥ College Degree	167	5,700	95.2	(89.0 - 100)	12,459	5,361,700	95.3	(93.3 - 97.2)
Income								
≤ \$24,999	*	*	*	*	9,640	2,437,200	69.7	(62.5 - 76.8)
\$25,000-\$49,999	200	6,300	74.8	(64.8 - 84.7)	9,662	2,384,000	82.2	(76.1 - 88.3)
\$50,000-\$74,999	*	*	*	*	5,772	1,855,700	94.0	(91.2 - 96.7)
≥ \$75,000	119	5,500	99.6	(98.8 - 100)	8,567	4,388,400	97.5	(96.6 - 98.5)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Sample size from survey

³ Estimated number of adults who have health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 6b. Health insurance among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	644	81.9	(76.3 - 87.4)	38,119	86.7	(84.4 - 89.1)
Gender						
Male	243	78.1	(69.0 - 87.2)	14,597	83.6	(79.7 - 87.5)
Female	401	85.6	(79.4 - 91.8)	23,522	89.6	(86.8 - 92.3)
Race/Ethnicity						
White Non-Hispanic	629	82.6	(77.1 - 88.1)	35,811	92.1	(90.5 - 93.7)
Black Non-Hispanic	*	*	*	736	82.5	(75.0 - 90.0)
Hispanic	*	*	*	860	68.7	(59.0 - 78.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	302	79.8	(71.7 - 87.9)	15,673	80.1	(75.4 - 84.7)
Some College	*	*	*	9,906	84.6	(79.7 - 89.4)
≥ College Degree	167	95.0	(88.2 - 100)	12,459	95.4	(93.6 - 97.2)
Income						
≤ \$24,999	*	*	*	9,640	70.2	(63.2 - 77.1)
\$25,000-\$49,999	*	*	*	9,662	81.9	(76.0 - 87.9)
\$50,000-\$74,999	*	*	*	5,772	94.0	(91.3 - 96.7)
≥ \$75,000	119	99.6	(98.9 - 100)	8,567	97.5	(96.6 - 98.5)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 7a. Health insurance among adults aged 18-64¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	423	16,200	79.2	(73.0 - 85.4)	25,165	10,251,300	84.2	(81.3 - 87.1)
Gender								
Male	164	7,800	75.2	(65.2 - 85.1)	10,078	4,839,900	80.3	(75.5 - 85.0)
Female	259	8,400	83.4	(76.3 - 90.5)	15,087	5,411,400	88.0	(84.7 - 91.3)
Age (years)								
18-34	*	*	*	*	3,910	3,302,100	74.3	(67.8 - 80.7)
35-44	*	*	*	*	4,981	2,556,600	88.9	(84.7 - 93.1)
45-54	142	4,000	84.6	(77.2 - 92.1)	7,791	2,472,900	88.7	(84.3 - 93.0)
55-64	160	3,600	90.4	(84.8 - 96.0)	8,483	1,919,700	92.8	(90.1 - 95.5)
≥ 65	-	-	-	-	-	-	-	-
Race/Ethnicity								
White Non-Hispanic	413	15,900	80.1	(73.9 - 86.3)	23,354	6,582,200	90.3	(88.3 - 92.4)
Black Non-Hispanic	*	*	*	*	567	1,464,100	80.2	(71.6 - 88.7)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	181	7,200	75.7	(66.4 - 85.0)	9,037	3,191,500	74.6	(68.6 - 80.6)
Some College	*	*	*	*	7,031	2,540,400	82.9	(77.4 - 88.4)
≥ College Degree	123	4,500	94.0	(86.3 - 100)	9,068	4,502,400	94.5	(92.2 - 96.8)
Income								
≤ \$24,999	*	*	*	*	4,918	1,707,200	62.6	(54.1 - 71.2)
\$25,000-\$49,999	*	*	*	*	6,180	1,783,000	77.7	(70.3 - 85.1)
\$50,000-\$74,999	*	*	*	*	4,554	1,553,400	92.9	(89.7 - 96.1)
≥ \$75,000	101	4,900	99.6	(98.7 - 100)	7,324	3,987,600	97.3	(96.3 - 98.3)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Sample size from survey

³ Estimated number of adults aged 18-64 who have health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 7b. Health insurance among adults aged 18-64¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	423	78.6	(72.1 - 85.2)	25,165	84.3	(81.5 - 87.2)
Gender						
Male	*	*	*	10,078	80.5	(75.9 - 85.1)
Female	259	83.0	(75.6 - 90.4)	15,087	87.9	(84.6 - 91.2)
Race/Ethnicity						
White Non-Hispanic	413	79.5	(72.9 - 86.0)	23,354	90.3	(88.3 - 92.3)
Black Non-Hispanic	*	*	*	567	80.0	(71.3 - 88.6)
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	181	75.7	(65.9 - 85.4)	9,037	75.1	(69.3 - 80.9)
Some College	*	*	*	7,031	82.6	(77.1 - 88.1)
≥ College Degree	123	93.9	(85.6 - 100)	9,068	94.6	(92.5 - 96.8)
Income						
≤ \$24,999	*	*	*	4,918	63.2	(54.9 - 71.6)
\$25,000-\$49,999	*	*	*	6,180	77.5	(70.2 - 84.7)
\$50,000-\$74,999	*	*	*	4,554	93.0	(89.8 - 96.2)
≥ \$75,000	101	99.6	(98.8 - 100)	7,324	97.3	(96.3 - 98.3)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 8a. No health insurance among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	644	4,400	17.3	(12.2 - 22.4)	38,119	1,989,300	13.4	(11.0 - 15.8)
Gender								
Male	243	2,600	21.0	(12.6 - 29.5)	14,597	1,207,400	17.0	(12.9 - 21.1)
Female	401	1,700	13.6	(7.9 - 19.3)	23,522	781,900	10.1	(7.4 - 12.8)
Age (years)								
18-34	*	*	*	*	3,910	1,144,100	25.7	(19.3 - 32.2)
35-44	*	*	*	*	4,986	319,200	11.1	(6.9 - 15.3)
45-54	142	700	15.4	(7.9 - 22.8)	7,831	316,500	11.1	(6.8 - 15.4)
55-64	164	400	9.8	(4.2 - 15.4)	8,764	153,300	7.2	(4.5 - 9.8)
≥ 65	*	*	*	*	12,628	56,200	2.2	(1.2 - 3.3)
Race/Ethnicity								
White Non-Hispanic	629	4,100	16.5	(11.4 - 21.6)	35,811	727,100	7.8	(6.2 - 9.4)
Black Non-Hispanic	*	*	*	*	736	372,800	17.3	(9.9 - 24.7)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	302	2,400	19.7	(12.1 - 27.2)	15,673	1,134,400	20.3	(15.5 - 25.1)
Some College	*	*	*	*	9,906	531,400	15.0	(10.2 - 19.9)
≥ College Degree	*	*	*	*	12,459	266,300	4.7	(2.8 - 6.7)
Income								
≤ \$24,999	*	*	*	*	9,640	1,061,200	30.3	(23.2 - 37.5)
\$25,000-\$49,999	200	2,100	25.2	(15.3 - 35.2)	9,662	516,700	17.8	(11.7 - 23.9)
\$50,000-\$74,999	*	*	*	*	5,772	119,100	6.0	(3.3 - 8.8)
≥ \$75,000	*	*	*	*	8,567	111,100	2.5	(1.5 - 3.4)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Sample size from survey

³ Estimated number of adults who have no health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 8b. No health insurance among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	644	18.1	(12.6 - 23.7)	38,119	13.3	(10.9 - 15.6)
Gender						
Male	243	21.9	(12.8 - 31.0)	14,597	16.4	(12.5 - 20.3)
Female	401	14.4	(8.2 - 20.6)	23,522	10.4	(7.7 - 13.2)
Race/Ethnicity						
White Non-Hispanic	629	17.4	(11.9 - 22.9)	35,811	7.9	(6.3 - 9.5)
Black Non-Hispanic	*	*	*	736	17.5	(10.0 - 25.0)
Hispanic	*	*	*	860	31.3	(21.6 - 41.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	302	20.2	(12.1 - 28.3)	15,673	19.9	(15.3 - 24.6)
Some College	*	*	*	9,906	15.4	(10.6 - 20.3)
≥ College Degree	*	*	*	12,459	4.6	(2.8 - 6.4)
Income						
≤ \$24,999	*	*	*	9,640	29.8	(22.9 - 36.8)
\$25,000-\$49,999	*	*	*	9,662	18.1	(12.1 - 24.0)
\$50,000-\$74,999	*	*	*	5,772	6.0	(3.3 - 8.7)
≥ \$75,000	*	*	*	8,567	2.5	(1.5 - 3.4)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 9a. No health insurance among adults aged 18-64¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	423	4,200	20.8	(14.6 - 27.0)	25,165	1,928,400	15.8	(12.9 - 18.7)
Gender								
Male	164	2,600	24.8	(14.9 - 34.8)	10,078	1,190,200	19.7	(15.0 - 24.5)
Female	259	1,700	16.6	(9.5 - 23.7)	15,087	738,200	12.0	(8.7 - 15.3)
Age (years)								
18-34	*	*	*	*	3,910	1,144,100	25.7	(19.3 - 32.2)
35-44	*	*	*	*	4,981	319,200	11.1	(6.9 - 15.3)
45-54	142	700	15.4	(7.9 - 22.8)	7,791	316,100	11.3	(7.0 - 15.7)
55-64	160	400	9.6	(4.0 - 15.2)	8,483	148,900	7.2	(4.5 - 9.9)
≥ 65	-	-	-	-	-	-	-	-
Race/Ethnicity								
White Non-Hispanic	413	3,900	19.9	(13.7 - 26.1)	23,354	703,100	9.7	(7.6 - 11.7)
Black Non-Hispanic	*	*	*	*	567	361,800	19.8	(11.3 - 28.4)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	181	2,300	24.3	(15.0 - 33.6)	9,037	1,087,700	25.4	(19.4 - 31.4)
Some College	*	*	*	*	7,031	522,800	17.1	(11.6 - 22.6)
≥ College Degree	*	*	*	*	9,068	261,200	5.5	(3.2 - 7.8)
Income								
≤ \$24,999	*	*	*	*	4,918	1,018,000	37.4	(28.8 - 45.9)
\$25,000-\$49,999	*	*	*	*	6,180	511,100	22.3	(14.9 - 29.7)
\$50,000-\$74,999	*	*	*	*	4,554	118,800	7.1	(3.9 - 10.3)
≥ \$75,000	*	*	*	*	7,324	110,700	2.7	(1.7 - 3.7)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Sample size from survey

³ Estimated number of adults aged 18-64 who have no health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 9b. No health insurance among adults aged 18-64¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	423	21.4	(14.8 - 27.9)	25,165	15.7	(12.8 - 18.5)
Gender						
Male	*	*	*	10,078	19.5	(14.9 - 24.1)
Female	259	17.0	(9.6 - 24.4)	15,087	12.1	(8.8 - 15.4)
Race/Ethnicity						
White Non-Hispanic	413	20.5	(14.0 - 27.1)	23,354	9.7	(7.7 - 11.7)
Black Non-Hispanic	*	*	*	567	20.0	(11.4 - 28.7)
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	181	24.3	(14.6 - 34.1)	9,037	24.9	(19.1 - 30.7)
Some College	*	*	*	7,031	17.4	(11.9 - 22.9)
≥ College Degree	*	*	*	9,068	5.4	(3.2 - 7.5)
Income						
≤ \$24,999	*	*	*	4,918	36.8	(28.4 - 45.1)
\$25,000-\$49,999	*	*	*	6,180	22.5	(15.3 - 29.8)
\$50,000-\$74,999	*	*	*	4,554	7.0	(3.8 - 10.2)
≥ \$75,000	*	*	*	7,324	2.7	(1.7 - 3.7)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 10a. Adults with regular health care provider¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	644	22,000	86.0	(81.7 - 90.4)	38,135	12,324,300	83.0	(80.4 - 85.6)
Gender								
Male	243	10,400	82.5	(75.3 - 89.7)	14,593	5,494,300	77.6	(73.2 - 82.0)
Female	401	11,600	89.6	(84.8 - 94.3)	23,542	6,830,000	87.8	(85.1 - 90.5)
Age (years)								
18-34	*	*	*	*	3,929	2,931,600	65.8	(59.2 - 72.5)
35-44	*	*	*	*	4,988	2,471,000	86.1	(81.9 - 90.3)
45-54	142	4,300	90.2	(84.8 - 95.5)	7,823	2,529,300	88.6	(84.6 - 92.7)
55-64	164	3,600	89.9	(84.5 - 95.3)	8,770	2,002,000	93.6	(91.7 - 95.4)
≥ 65	215	4,500	94.9	(91.5 - 98.3)	12,625	2,390,400	94.1	(92.2 - 96.0)
Race/Ethnicity								
White Non-Hispanic	629	21,500	86.5	(82.1 - 90.8)	35,821	8,227,700	88.2	(86.4 - 89.9)
Black Non-Hispanic	*	*	*	*	740	1,753,200	81.3	(72.6 - 89.9)
Hispanic	*	*	*	*	860	1,453,100	64.7	(54.7 - 74.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	303	10,500	86.7	(81.1 - 92.2)	15,692	4,249,200	75.9	(70.9 - 80.8)
Some College	*	*	*	*	9,904	2,935,000	82.8	(77.2 - 88.4)
≥ College Degree	166	5,300	90.3	(83.3 - 97.3)	12,456	5,046,100	89.8	(87.1 - 92.6)
Income								
≤ \$24,999	171	4,600	82.8	(73.9 - 91.8)	9,644	2,554,200	72.7	(65.9 - 79.6)
\$25,000-\$49,999	200	7,100	81.7	(73.3 - 90.2)	9,655	2,243,100	77.3	(70.0 - 84.6)
\$50,000-\$74,999	75	2,800	91.7	(82.6 - 100)	5,771	1,767,500	89.5	(85.2 - 93.9)
≥ \$75,000	119	4,900	90.0	(80.2 - 99.9)	8,560	4,122,800	91.9	(90.2 - 93.7)

¹ Defined as having one or more persons respondent thinks of as a personal doctor or health care provider

² Sample size from survey

³ Estimated number of adults who have at least one person they think of as their personal doctor or health care provider (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 10b. Adults with regular health care provider¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	644	85.3	(80.6 - 90.0)	38,135	83.0	(80.4 - 85.5)
Gender						
Male	243	82.0	(74.3 - 89.8)	14,593	78.1	(73.9 - 82.3)
Female	401	88.6	(83.3 - 93.8)	23,542	87.4	(84.6 - 90.1)
Race/Ethnicity						
White Non-Hispanic	629	85.7	(81.1 - 90.4)	35,821	88.0	(86.2 - 89.7)
Black Non-Hispanic	*	*	*	740	81.4	(73.0 - 89.8)
Hispanic	*	*	*	860	65.2	(55.5 - 75.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	303	86.2	(80.2 - 92.1)	15,692	75.8	(71.0 - 80.6)
Some College	*	*	*	9,904	82.9	(77.5 - 88.3)
≥ College Degree	166	90.0	(82.4 - 97.6)	12,456	89.8	(87.1 - 92.4)
Income						
≤ \$24,999	171	82.1	(72.2 - 91.9)	9,644	72.9	(66.1 - 79.6)
\$25,000-\$49,999	200	80.6	(71.6 - 89.7)	9,655	77.4	(70.5 - 84.3)
\$50,000-\$74,999	75	91.3	(81.7 - 100)	5,771	89.6	(85.3 - 94.0)
≥ \$75,000	*	*	*	8,560	91.6	(89.8 - 93.4)

¹ Defined as having one or more persons respondent thinks of as a personal doctor or health care provider

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 11a. Cost prevented visit to doctor within the past year among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	647	2,900	11.5	(8.0 - 15.0)	38,143	2,034,800	13.7	(11.6 - 15.8)
Gender								
Male	244	1,400	10.8	(5.5 - 16.0)	14,607	903,300	12.7	(9.5 - 16.0)
Female	403	1,600	12.2	(7.5 - 16.9)	23,536	1,131,500	14.6	(11.8 - 17.4)
Age (years)								
18-34	*	*	*	*	3,939	947,300	21.2	(15.6 - 26.9)
35-44	*	*	*	*	4,987	347,900	12.1	(8.0 - 16.2)
45-54	142	700	15.2	(7.5 - 23.0)	7,825	382,200	13.5	(9.9 - 17.1)
55-64	164	300	7.6	(3.4 - 11.7)	8,766	236,800	11.1	(8.0 - 14.2)
≥ 65	*	*	*	*	12,626	120,600	4.8	(2.5 - 7.0)
Race/Ethnicity								
White Non-Hispanic	632	2,800	11.3	(7.8 - 14.7)	35,833	984,600	10.6	(8.9 - 12.2)
Black Non-Hispanic	*	*	*	*	738	280,000	13.1	(7.2 - 19.0)
Hispanic	*	*	*	*	859	597,900	26.5	(17.3 - 35.7)
Other Non-Hispanic	*	*	*	*	713	172,300	15.3	(5.7 - 24.9)
Education								
≤ High School	304	1,600	12.9	(7.7 - 18.1)	15,696	996,600	17.8	(13.8 - 21.9)
Some College	175	1,200	16.5	(8.3 - 24.6)	9,911	512,400	14.5	(10.4 - 18.6)
≥ College Degree	*	*	*	*	12,453	466,900	8.3	(6.0 - 10.6)
Income								
≤ \$24,999	171	900	16.4	(8.9 - 24.0)	9,639	854,500	24.3	(18.4 - 30.3)
\$25,000-\$49,999	201	1,200	13.7	(6.6 - 20.9)	9,656	532,900	18.4	(12.5 - 24.3)
\$50,000-\$74,999	*	*	*	*	5,776	202,700	10.3	(6.0 - 14.5)
≥ \$75,000	*	*	*	*	8,561	192,000	4.3	(2.9 - 5.6)

¹ Defined as needing to see a doctor within the past year but not being able to because of cost

² Sample size from survey

³ Estimated number of adults who did not see a doctor because of cost within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 11b. Cost prevented visit to doctor within the past year among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	647	12.1	(8.3 - 15.9)	38,143	13.8	(11.7 - 15.9)
Gender						
Male	244	11.0	(5.4 - 16.5)	14,607	12.5	(9.4 - 15.6)
Female	403	13.2	(8.0 - 18.3)	23,536	15.0	(12.1 - 17.9)
Race/Ethnicity						
White Non-Hispanic	632	11.9	(8.1 - 15.7)	35,833	10.8	(9.2 - 12.5)
Black Non-Hispanic	*	*	*	738	13.2	(7.2 - 19.3)
Hispanic	*	*	*	859	26.1	(17.2 - 35.1)
Other Non-Hispanic	*	*	*	713	14.8	(5.5 - 24.2)
Education						
≤ High School	304	13.2	(7.7 - 18.8)	15,696	17.8	(13.8 - 21.7)
Some College	175	17.7	(8.9 - 26.4)	9,911	15.3	(11.0 - 19.5)
≥ College Degree	*	*	*	12,453	8.2	(6.0 - 10.5)
Income						
≤ \$24,999	171	16.1	(8.2 - 24.0)	9,639	24.5	(18.6 - 30.4)
\$25,000-\$49,999	201	14.7	(7.1 - 22.3)	9,656	18.7	(12.9 - 24.6)
\$50,000-\$74,999	*	*	*	5,776	10.4	(6.1 - 14.7)
≥ \$75,000	*	*	*	8,561	4.4	(3.1 - 5.8)

¹ Defined as needing to see a doctor within the past year but not being able to because of cost

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 12a. Visited a doctor for routine checkup within the past year among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	646	17,400	68.0	(62.4 - 73.6)	38,026	10,809,300	72.9	(70.6 - 75.3)
Gender								
Male	245	8,300	65.4	(56.7 - 74.2)	14,592	4,738,400	66.9	(63.0 - 70.9)
Female	401	9,100	70.6	(63.6 - 77.6)	23,434	6,070,900	78.5	(75.9 - 81.0)
Age (years)								
18-34	*	*	*	*	3,921	2,747,400	61.7	(55.6 - 67.9)
35-44	*	*	*	*	4,973	1,883,100	65.5	(60.2 - 70.7)
45-54	142	3,400	71.4	(63.2 - 79.6)	7,808	2,174,100	76.6	(73.2 - 80.0)
55-64	164	3,200	79.6	(72.7 - 86.5)	8,751	1,728,600	81.4	(78.7 - 84.2)
≥ 65	218	4,200	87.0	(81.9 - 92.1)	12,573	2,276,000	90.0	(87.5 - 92.5)
Race/Ethnicity								
White Non-Hispanic	631	16,900	67.9	(62.2 - 73.6)	35,721	6,746,200	72.4	(70.6 - 74.2)
Black Non-Hispanic	*	*	*	*	736	1,688,900	78.8	(70.4 - 87.3)
Hispanic	*	*	*	*	857	1,550,900	69.3	(60.0 - 78.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	305	8,200	67.3	(59.5 - 75.0)	15,640	4,150,600	74.7	(70.7 - 78.7)
Some College	*	*	*	*	9,880	2,524,000	71.3	(66.1 - 76.5)
≥ College Degree	*	*	*	*	12,428	4,042,900	71.9	(68.5 - 75.3)
Income								
≤ \$24,999	*	*	*	*	9,616	2,469,200	70.4	(64.2 - 76.6)
\$25,000-\$49,999	*	*	*	*	9,631	2,068,400	71.3	(65.3 - 77.3)
\$50,000-\$74,999	*	*	*	*	5,765	1,466,000	74.8	(69.6 - 80.0)
≥ \$75,000	*	*	*	*	8,547	3,176,500	70.7	(67.3 - 74.2)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past year

² Sample size from survey

³ Estimated number of adults who have visited a doctor for a routine checkup within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 12b. Visited a doctor for routine checkup within the past year among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	646	66.7	(60.7 - 72.7)	38,026	72.7	(70.3 - 75.0)
Gender						
Male	245	64.2	(54.8 - 73.6)	14,592	67.1	(63.3 - 70.9)
Female	401	69.1	(61.6 - 76.6)	23,434	77.7	(75.1 - 80.3)
Race/Ethnicity						
White Non-Hispanic	631	66.6	(60.5 - 72.6)	35,721	71.9	(70.1 - 73.7)
Black Non-Hispanic	*	*	*	736	79.3	(71.3 - 87.4)
Hispanic	*	*	*	857	69.2	(60.1 - 78.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	305	66.3	(58.0 - 74.6)	15,640	74.4	(70.4 - 78.4)
Some College	*	*	*	9,880	70.9	(65.9 - 76.0)
≥ College Degree	*	*	*	12,428	71.7	(68.3 - 75.0)
Income						
≤ \$24,999	*	*	*	9,616	70.2	(64.1 - 76.2)
\$25,000-\$49,999	*	*	*	9,631	71.1	(65.3 - 76.9)
\$50,000-\$74,999	*	*	*	5,765	74.5	(69.2 - 79.8)
≥ \$75,000	*	*	*	8,547	70.3	(66.9 - 73.8)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past year

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 13a. Visited a doctor for routine checkup within the past 2 years among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	646	20,100	78.5	(73.1 - 83.8)	38,026	12,726,600	85.9	(83.9 - 87.9)
Gender								
Male	245	9,700	76.0	(67.7 - 84.3)	14,592	5,791,900	81.8	(78.2 - 85.4)
Female	401	10,400	80.9	(74.2 - 87.5)	23,434	6,934,700	89.6	(87.9 - 91.3)
Age (years)								
18-34	*	*	*	*	3,921	3,535,400	79.4	(73.9 - 85.0)
35-44	*	*	*	*	4,973	2,355,600	81.9	(78.0 - 85.7)
45-54	142	4,000	83.4	(76.8 - 90.0)	7,808	2,507,900	88.4	(85.6 - 91.2)
55-64	164	3,500	86.2	(80.2 - 92.2)	8,751	1,924,800	90.7	(88.6 - 92.7)
≥ 65	218	4,400	91.3	(86.8 - 95.7)	12,573	2,402,900	95.0	(93.5 - 96.5)
Race/Ethnicity								
White Non-Hispanic	631	19,600	78.6	(73.3 - 84.0)	35,721	8,070,000	86.6	(85.4 - 87.9)
Black Non-Hispanic	*	*	*	*	736	1,914,500	89.4	(82.3 - 96.5)
Hispanic	*	*	*	*	857	1,832,800	81.9	(73.7 - 90.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	305	9,700	79.4	(72.5 - 86.3)	15,640	4,713,700	84.8	(81.2 - 88.4)
Some College	*	*	*	*	9,880	2,990,500	84.5	(79.6 - 89.4)
≥ College Degree	*	*	*	*	12,428	4,928,300	87.6	(85.3 - 89.9)
Income								
≤ \$24,999	*	*	*	*	9,616	2,911,700	83.0	(77.4 - 88.6)
\$25,000-\$49,999	201	6,600	76.2	(66.5 - 85.9)	9,631	2,375,500	81.9	(76.1 - 87.7)
\$50,000-\$74,999	76	2,800	89.2	(79.7 - 98.6)	5,765	1,733,500	88.4	(84.9 - 91.9)
≥ \$75,000	*	*	*	*	8,547	3,907,000	87.0	(84.8 - 89.2)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past 2 years

² Sample size from survey

³ Estimated number of adults who have visited a doctor for a routine checkup within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 13b. Visited a doctor for routine checkup within the past 2 years among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	646	77.6	(71.9 - 83.3)	38,026	85.7	(83.7 - 87.6)
Gender						
Male	245	75.4	(66.5 - 84.3)	14,592	81.9	(78.4 - 85.3)
Female	401	79.8	(72.6 - 86.9)	23,434	89.1	(87.4 - 90.9)
Race/Ethnicity						
White Non-Hispanic	631	77.8	(72.0 - 83.6)	35,721	86.2	(85.0 - 87.5)
Black Non-Hispanic	*	*	*	736	89.6	(83.0 - 96.3)
Hispanic	*	*	*	857	81.8	(73.8 - 89.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	305	79.0	(71.6 - 86.5)	15,640	84.6	(81.1 - 88.1)
Some College	*	*	*	9,880	84.4	(79.8 - 89.0)
≥ College Degree	*	*	*	12,428	87.3	(85.0 - 89.6)
Income						
≤ \$24,999	*	*	*	9,616	83.2	(77.8 - 88.5)
\$25,000-\$49,999	*	*	*	9,631	81.9	(76.3 - 87.4)
\$50,000-\$74,999	76	89.6	(80.0 - 99.2)	5,765	88.1	(84.4 - 91.8)
≥ \$75,000	*	*	*	8,547	86.4	(84.2 - 88.7)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past 2 years

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 14a. Dentist visit within the past year among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	644	16,200	63.4	(57.9 - 68.8)	38,026	10,540,400	71.1	(68.6 - 73.6)
Gender								
Male	244	7,400	58.3	(49.7 - 67.0)	14,573	4,845,800	68.4	(64.5 - 72.2)
Female	400	8,800	68.3	(61.8 - 74.9)	23,453	5,694,600	73.6	(70.4 - 76.8)
Age (years)								
18-34	*	*	*	*	3,924	2,789,700	62.6	(56.3 - 69.0)
35-44	*	*	*	*	4,979	2,181,000	75.8	(70.9 - 80.7)
45-54	142	3,300	68.3	(59.4 - 77.3)	7,814	2,149,300	75.8	(71.9 - 79.7)
55-64	163	2,700	68.5	(60.4 - 76.5)	8,742	1,690,500	79.2	(75.9 - 82.6)
≥ 65	217	3,300	67.9	(60.9 - 75.0)	12,567	1,729,800	68.5	(64.7 - 72.3)
Race/Ethnicity								
White Non-Hispanic	629	16,100	64.7	(59.2 - 70.2)	35,721	6,959,900	74.7	(72.7 - 76.6)
Black Non-Hispanic	*	*	*	*	735	1,405,800	66.2	(57.9 - 74.5)
Hispanic	*	*	*	*	862	1,425,600	63.1	(53.7 - 72.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	302	6,300	51.9	(44.0 - 59.9)	15,613	3,620,700	65.1	(60.7 - 69.5)
Some College	*	*	*	*	9,880	2,504,700	70.8	(66.1 - 75.5)
≥ College Degree	167	4,900	82.1	(74.2 - 90.0)	12,453	4,402,400	78.2	(74.6 - 81.8)
Income								
≤ \$24,999	*	*	*	*	9,594	1,972,200	56.2	(49.8 - 62.6)
\$25,000-\$49,999	201	5,000	58.0	(48.3 - 67.8)	9,649	1,896,000	65.8	(59.5 - 72.1)
\$50,000-\$74,999	76	2,600	83.0	(74.4 - 91.6)	5,763	1,482,400	75.2	(69.4 - 81.0)
≥ \$75,000	119	4,500	82.3	(73.8 - 90.8)	8,564	3,707,900	82.4	(79.5 - 85.3)

¹ Defined as having visited a dentist or a dental clinic for any reason within the past year

² Sample size from survey

³ Estimated number of adults who have visited a dentist or a dental clinic for any reason within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 14b. Dentist visit within the past year among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	644	63.0	(57.2 - 68.8)	38,026	71.1	(68.6 - 73.5)
Gender						
Male	244	57.7	(48.4 - 66.9)	14,573	68.3	(64.6 - 72.1)
Female	400	68.3	(61.3 - 75.3)	23,453	73.6	(70.4 - 76.8)
Race/Ethnicity						
White Non-Hispanic	629	64.3	(58.5 - 70.2)	35,721	74.4	(72.4 - 76.3)
Black Non-Hispanic	*	*	*	735	66.6	(58.3 - 74.9)
Hispanic	*	*	*	862	63.6	(54.4 - 72.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	302	51.8	(43.3 - 60.3)	15,613	65.3	(60.9 - 69.6)
Some College	*	*	*	9,880	70.4	(65.7 - 75.1)
≥ College Degree	167	81.2	(72.6 - 89.8)	12,453	78.1	(74.6 - 81.7)
Income						
≤ \$24,999	*	*	*	9,594	56.4	(50.1 - 62.8)
\$25,000-\$49,999	*	*	*	9,649	65.8	(59.6 - 72.0)
\$50,000-\$74,999	76	84.6	(76.3 - 92.8)	5,763	75.0	(69.1 - 80.8)
≥ \$75,000	119	82.4	(73.3 - 91.5)	8,564	82.1	(79.1 - 85.0)

¹ Defined as having visited a dentist or a dental clinic for any reason within the past year

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 15a. Had teeth cleaned within the past year among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	587	14,700	62.3	(56.7 - 68.0)	34,070	9,985,900	71.7	(69.1 - 74.3)
Gender								
Male	222	6,500	56.8	(47.8 - 65.7)	13,173	4,616,400	69.0	(65.0 - 73.1)
Female	365	8,200	67.6	(60.7 - 74.5)	20,897	5,369,500	74.2	(70.8 - 77.5)
Age (years)								
18-34	*	*	*	*	3,860	2,746,400	63.0	(56.5 - 69.5)
35-44	*	*	*	*	4,875	2,112,100	75.5	(70.7 - 80.3)
45-54	133	3,000	67.3	(58.0 - 76.7)	7,438	2,022,600	73.3	(69.1 - 77.4)
55-64	151	2,500	67.4	(59.0 - 75.8)	7,906	1,560,900	77.7	(73.8 - 81.6)
≥ 65	185	3,000	71.3	(63.8 - 78.8)	9,991	1,543,900	77.1	(73.7 - 80.5)
Race/Ethnicity								
White Non-Hispanic	573	14,600	63.6	(57.8 - 69.3)	31,982	6,601,500	75.4	(73.4 - 77.4)
Black Non-Hispanic	*	*	*	*	680	1,227,200	61.6	(52.1 - 71.0)
Hispanic	*	*	*	*	789	1,421,300	66.6	(57.1 - 76.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	260	5,500	51.8	(43.4 - 60.1)	12,740	3,388,600	67.7	(63.1 - 72.3)
Some College	*	*	*	*	9,145	2,286,800	68.5	(62.9 - 74.1)
≥ College Degree	161	4,700	80.8	(72.5 - 89.1)	12,128	4,298,500	78.3	(74.8 - 81.7)
Income								
≤ \$24,999	*	*	*	*	7,525	1,861,800	60.3	(53.5 - 67.1)
\$25,000-\$49,999	191	4,600	55.0	(45.0 - 64.9)	8,709	1,681,700	62.6	(55.4 - 69.8)
\$50,000-\$74,999	73	2,600	84.2	(75.6 - 92.7)	5,534	1,447,800	76.3	(70.7 - 81.8)
≥ \$75,000	114	4,200	82.8	(73.9 - 91.7)	8,448	3,617,500	80.8	(77.8 - 83.8)

¹ Defined as having had teeth cleaned by a dentist or dental hygienist within the past year

² Sample size from survey

³ Estimated number of adults who have had their teeth cleaned by a dentist or dental hygienist within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 15b. Had teeth cleaned within the past year among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	587	61.9	(55.9 - 68.0)	34,070	71.7	(69.1 - 74.2)
Gender						
Male	222	55.7	(46.1 - 65.3)	13,173	69.2	(65.3 - 73.1)
Female	365	67.8	(60.4 - 75.1)	20,897	73.9	(70.5 - 77.3)
Race/Ethnicity						
White Non-Hispanic	573	63.0	(56.9 - 69.1)	31,982	75.1	(73.0 - 77.1)
Black Non-Hispanic	*	*	*	680	62.0	(52.7 - 71.4)
Hispanic	*	*	*	789	66.9	(57.5 - 76.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	260	51.7	(42.8 - 60.6)	12,740	67.7	(63.2 - 72.3)
Some College	*	*	*	9,145	68.1	(62.6 - 73.6)
≥ College Degree	161	80.0	(71.0 - 88.9)	12,128	78.3	(74.9 - 81.7)
Income						
≤ \$24,999	*	*	*	7,525	60.2	(53.4 - 67.0)
\$25,000-\$49,999	*	*	*	8,709	62.7	(55.7 - 69.7)
\$50,000-\$74,999	73	85.4	(77.1 - 93.7)	5,534	76.4	(70.9 - 81.9)
≥ \$75,000	114	82.5	(72.9 - 92.1)	8,448	80.4	(77.4 - 83.4)

¹ Defined as having had teeth cleaned by a dentist or dental hygienist within the past year

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 16a. Permanent teeth extracted due to decay or gum disease among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	637	13,300	52.6	(47.0 - 58.3)	37,652	7,426,500	50.4	(47.8 - 52.9)
Gender								
Male	240	6,800	54.8	(46.0 - 63.6)	14,457	3,367,000	47.8	(43.9 - 51.6)
Female	397	6,500	50.6	(43.6 - 57.6)	23,195	4,059,500	52.7	(49.5 - 56.0)
Age (years)								
18-34	*	*	*	*	3,921	1,121,600	25.4	(19.7 - 31.0)
35-44	*	*	*	*	4,964	1,250,000	43.6	(37.9 - 49.2)
45-54	141	2,800	58.6	(49.2 - 68.0)	7,780	1,595,300	56.1	(51.4 - 60.7)
55-64	162	2,600	64.9	(56.7 - 73.2)	8,648	1,382,800	65.5	(61.6 - 69.4)
≥ 65	212	4,000	84.8	(79.0 - 90.7)	12,339	2,076,900	83.0	(80.6 - 85.4)
Race/Ethnicity								
White Non-Hispanic	622	13,100	53.2	(47.5 - 58.9)	35,364	4,425,700	47.9	(46.0 - 49.9)
Black Non-Hispanic	*	*	*	*	727	1,244,700	58.2	(48.9 - 67.4)
Hispanic	*	*	*	*	853	1,220,500	54.2	(44.9 - 63.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	295	7,600	64.6	(56.5 - 72.6)	15,427	3,436,000	61.9	(57.2 - 66.6)
Some College	*	*	*	*	9,798	1,726,200	49.3	(44.3 - 54.2)
≥ College Degree	167	2,400	39.7	(30.4 - 49.1)	12,351	2,242,600	40.1	(36.5 - 43.6)
Income								
≤ \$24,999	*	*	*	*	9,483	2,171,900	62.3	(55.8 - 68.8)
\$25,000-\$49,999	200	4,300	50.2	(40.4 - 60.0)	9,552	1,577,400	54.7	(48.3 - 61.0)
\$50,000-\$74,999	*	*	*	*	5,744	1,000,700	51.5	(45.6 - 57.4)
≥ \$75,000	*	*	*	*	8,516	1,738,200	38.8	(35.3 - 42.2)

¹ Defined as having had at least one permanent tooth extracted because of tooth decay or gum disease

² Sample size from survey

³ Estimated number of adults who have had at least one permanent tooth extracted because of tooth decay or gum disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 16b. Permanent teeth extracted due to decay or gum disease among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	637	50.5	(44.5 - 56.4)	37,652	49.4	(46.9 - 51.9)
Gender						
Male	240	53.5	(44.1 - 62.9)	14,457	47.9	(44.1 - 51.7)
Female	397	47.6	(40.3 - 54.9)	23,195	50.7	(47.4 - 54.0)
Race/Ethnicity						
White Non-Hispanic	622	51.0	(44.9 - 57.0)	35,364	46.5	(44.5 - 48.5)
Black Non-Hispanic	*	*	*	727	57.7	(48.6 - 66.9)
Hispanic	*	*	*	853	53.8	(44.6 - 62.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	295	62.3	(53.7 - 70.9)	15,427	61.3	(56.7 - 66.0)
Some College	*	*	*	9,798	47.9	(43.0 - 52.9)
≥ College Degree	167	38.0	(28.3 - 47.7)	12,351	39.0	(35.5 - 42.5)
Income						
≤ \$24,999	*	*	*	9,483	61.6	(55.2 - 68.1)
\$25,000-\$49,999	*	*	*	9,552	54.2	(48.0 - 60.3)
\$50,000-\$74,999	*	*	*	5,744	49.9	(44.0 - 55.9)
≥ \$75,000	*	*	*	8,516	37.7	(34.2 - 41.1)

¹ Defined as having had at least one permanent tooth extracted because of tooth decay or gum disease

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 17a. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	212	600	12.9	(8.2 - 17.5)	12,339	491,200	19.6	(16.0 - 23.3)
Gender								
Male	76	200	11.1	(4.0 - 18.2)	4,352	167,200	16.4	(11.8 - 20.9)
Female	136	400	14.3	(8.2 - 20.4)	7,987	324,000	21.8	(16.6 - 27.0)
Age (years)								
≥ 65	212	600	12.9	(8.2 - 17.5)	12,339	491,200	19.6	(16.0 - 23.3)
Race/Ethnicity								
White Non-Hispanic	208	600	12.5	(7.9 - 17.2)	11,885	322,600	16.8	(14.3 - 19.2)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	113	500	21.4	(13.3 - 29.4)	6,358	314,100	25.3	(19.9 - 30.7)
Some College	*	*	*	*	2,736	90,600	20.2	(12.1 - 28.2)
≥ College Degree	*	*	*	*	3,214	80,500	10.1	(3.6 - 16.5)
Income								
≤ \$24,999	90	400	25.2	(15.2 - 35.1)	4,558	199,600	26.9	(19.3 - 34.4)
\$25,000-\$49,999	*	*	*	*	3,365	123,900	21.4	(13.1 - 29.6)
\$50,000-\$74,999	*	*	*	*	1,179	21,400	7.3	(3.8 - 10.8)
≥ \$75,000	*	*	*	*	1,187	9,600	2.5	(1.0 - 4.0)

¹ Defined as having all permanent teeth extracted because of tooth decay or gum disease among adults aged 65 and older

² Sample size from survey

³ Estimated number of adults aged 65 and older who have had all permanent teeth extracted because of tooth decay or gum disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 17b. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	212	12.7	(8.0 - 17.4)	12,339	19.4	(15.7 - 23.2)
Gender						
Male	76	11.0	(3.9 - 18.1)	4,352	16.5	(11.5 - 21.4)
Female	136	14.3	(8.2 - 20.5)	7,987	22.1	(16.6 - 27.7)
Race/Ethnicity						
White Non-Hispanic	208	12.3	(7.7 - 16.9)	11,885	16.6	(13.9 - 19.3)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	113	21.0	(12.8 - 29.1)	6,358	24.9	(19.3 - 30.5)
Some College	*	*	*	2,736	21.0	(12.1 - 29.9)
≥ College Degree	*	*	*	3,214	10.0	(3.6 - 16.3)
Income						
≤ \$24,999	*	*	*	4,558	27.3	(19.1 - 35.5)
\$25,000-\$49,999	*	*	*	3,365	21.0	(12.9 - 29.1)
\$50,000-\$74,999	*	*	*	1,179	7.6	(3.6 - 11.5)
≥ \$75,000	*	*	*	1,187	2.2	(0.9 - 3.5)

¹ Defined as having all permanent teeth extracted because of tooth decay or gum disease among adults aged 65 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 18a. Flu shot within the past 12 months¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	646	9,300	36.4	(31.4 - 41.3)	38,079	6,255,900	42.3	(39.9 - 44.8)
Gender								
Male	245	3,300	26.1	(19.9 - 32.3)	14,579	2,798,900	39.6	(35.8 - 43.5)
Female	401	6,000	46.5	(39.6 - 53.4)	23,500	3,456,900	44.8	(41.6 - 47.9)
Age (years)								
18-49	183	2,800	20.5	(13.3 - 27.7)	12,505	2,685,300	31.1	(27.3 - 34.9)
50-64	246	3,100	44.5	(37.3 - 51.6)	12,972	1,662,900	46.0	(42.6 - 49.5)
≥ 65	217	3,400	70.0	(63.1 - 77.0)	12,602	1,907,700	75.2	(72.0 - 78.4)
Race/Ethnicity								
White Non-Hispanic	631	8,900	35.8	(30.9 - 40.8)	35,770	4,029,000	43.3	(41.4 - 45.3)
Black Non-Hispanic	*	*	*	*	738	907,000	42.1	(33.1 - 51.1)
Hispanic	*	*	*	*	860	882,900	39.6	(30.4 - 48.9)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	304	4,100	33.5	(26.7 - 40.3)	15,663	2,457,800	44.3	(39.8 - 48.8)
Some College	175	2,700	35.9	(26.2 - 45.6)	9,898	1,377,700	38.9	(34.0 - 43.9)
≥ College Degree	*	*	*	*	12,436	2,404,400	43.0	(39.5 - 46.4)
Income								
≤ \$24,999	172	2,100	37.1	(27.3 - 47.0)	9,635	1,656,800	47.6	(41.2 - 54.0)
\$25,000-\$49,999	201	2,900	33.6	(25.0 - 42.2)	9,647	1,200,600	41.4	(35.6 - 47.3)
\$50,000-\$74,999	*	*	*	*	5,763	792,300	40.2	(34.3 - 46.2)
≥ \$75,000	*	*	*	*	8,548	1,776,100	39.6	(36.3 - 42.8)

¹ Defined as having had a flu shot within the past 12 months

² Sample size from survey

³ Estimated number of adults who have had a flu shot within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 18b. Flu shot within the past 12 months¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	646	34.2	(29.1 - 39.4)	38,079	41.7	(39.2 - 44.1)
Gender						
Male	245	24.5	(18.3 - 30.8)	14,579	40.0	(36.2 - 43.8)
Female	401	43.8	(36.5 - 51.1)	23,500	43.1	(40.0 - 46.3)
Race/Ethnicity						
White Non-Hispanic	631	33.7	(28.5 - 38.9)	35,770	42.4	(40.4 - 44.3)
Black Non-Hispanic	*	*	*	738	42.3	(33.3 - 51.3)
Hispanic	*	*	*	860	39.0	(29.9 - 48.1)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	304	31.5	(24.5 - 38.5)	15,663	43.7	(39.1 - 48.2)
Some College	175	32.9	(23.1 - 42.7)	9,898	37.8	(33.0 - 42.7)
≥ College Degree	*	*	*	12,436	42.5	(39.1 - 46.0)
Income						
≤ \$24,999	*	*	*	9,635	47.3	(41.0 - 53.7)
\$25,000-\$49,999	201	31.7	(22.9 - 40.5)	9,647	40.3	(34.5 - 46.0)
\$50,000-\$74,999	*	*	*	5,763	39.4	(33.4 - 45.4)
≥ \$75,000	*	*	*	8,548	39.1	(35.8 - 42.4)

¹ Defined as having had a flu shot within the past 12 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 19a. Flu vaccine that was sprayed in nose within the past 12 months¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	*	*	*	*	38,106	248,300	1.7	(0.6 - 2.8)
Gender								
Male	*	*	*	*	14,586	148,000	2.1	(0.0 - 4.2)
Female	*	*	*	*	23,520	100,300	1.3	(0.5 - 2.1)
Age (years)								
18-49	*	*	*	*	12,535	184,900	2.1	(0.3 - 3.9)
50-64	*	*	*	*	12,968	36,000	1.0	(0.2 - 1.8)
≥ 65	*	*	*	*	12,603	27,400	1.1	(0.4 - 1.8)
Race/Ethnicity								
White Non-Hispanic	*	*	*	*	35,796	66,000	0.7	(0.4 - 1.0)
Black Non-Hispanic	*	*	*	*	740	38,500	1.8	(0.0 - 4.1)
Hispanic	*	*	*	*	860	133,700	5.9	(0.0 - 12.3)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,679	172,500	3.1	(0.3 - 5.9)
Some College	*	*	*	*	9,897	43,500	1.2	(0.4 - 2.0)
≥ College Degree	*	*	*	*	12,449	31,600	0.6	(0.3 - 0.8)
Income								
≤ \$24,999	*	*	*	*	9,636	138,400	3.9	(0.0 - 8.1)
\$25,000-\$49,999	*	*	*	*	9,648	41,500	1.4	(0.0 - 3.1)
\$50,000-\$74,999	*	*	*	*	5,766	13,300	0.7	(0.2 - 1.2)
≥ \$75,000	*	*	*	*	8,561	35,300	0.8	(0.3 - 1.3)

¹ Defined as having had flu vaccine sprayed in nose within the past 12 months

² Sample size from survey

³ Estimated number of adults who have had flu vaccine sprayed in nose within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 19b. Flu vaccine that was sprayed in nose within the past 12 months¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	*	*	*	38,106	1.7	(0.6 - 2.7)
Gender						
Male	*	*	*	14,586	2.0	(0.1 - 3.9)
Female	*	*	*	23,520	1.4	(0.5 - 2.3)
Race/Ethnicity						
White Non-Hispanic	*	*	*	35,796	0.7	(0.4 - 1.0)
Black Non-Hispanic	*	*	*	740	1.9	(0.0 - 4.4)
Hispanic	*	*	*	860	5.9	(0.0 - 11.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	15,679	3.1	(0.4 - 5.8)
Some College	*	*	*	9,897	1.2	(0.4 - 2.0)
≥ College Degree	*	*	*	12,449	0.6	(0.3 - 0.8)
Income						
≤ \$24,999	*	*	*	9,636	3.8	(0.0 - 7.7)
\$25,000-\$49,999	*	*	*	9,648	1.5	(0.0 - 3.4)
\$50,000-\$74,999	*	*	*	5,766	0.7	(0.1 - 1.2)
≥ \$75,000	*	*	*	8,561	0.8	(0.3 - 1.3)

¹ Defined as having had flu vaccine sprayed in nose within the past 12 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 20a. Flu shot or flu vaccine sprayed in nose within the past 12 months¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	646	9,300	36.4	(31.4 - 41.3)	38,059	6,302,300	42.7	(40.2 - 45.1)
Gender								
Male	245	3,300	26.1	(19.9 - 32.3)	14,568	2,822,700	40.0	(36.1 - 43.9)
Female	401	6,000	46.5	(39.6 - 53.4)	23,491	3,479,500	45.1	(42.0 - 48.2)
Age (years)								
18-49	183	2,800	20.5	(13.3 - 27.7)	12,501	2,726,400	31.6	(27.8 - 35.4)
50-64	246	3,100	44.5	(37.3 - 51.6)	12,964	1,664,200	46.1	(42.7 - 49.5)
≥ 65	217	3,400	70.0	(63.1 - 77.0)	12,594	1,911,700	75.4	(72.2 - 78.6)
Race/Ethnicity								
White Non-Hispanic	631	8,900	35.8	(30.9 - 40.8)	35,750	4,048,700	43.6	(41.7 - 45.5)
Black Non-Hispanic	*	*	*	*	738	910,000	42.2	(33.2 - 51.2)
Hispanic	*	*	*	*	860	903,300	40.6	(31.3 - 49.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	304	4,100	33.5	(26.7 - 40.3)	15,651	2,480,100	44.7	(40.2 - 49.3)
Some College	175	2,700	35.9	(26.2 - 45.6)	9,895	1,387,300	39.2	(34.3 - 44.1)
≥ College Degree	*	*	*	*	12,431	2,418,300	43.2	(39.8 - 46.7)
Income								
≤ \$24,999	172	2,100	37.1	(27.3 - 47.0)	9,626	1,675,100	48.2	(41.8 - 54.6)
\$25,000-\$49,999	201	2,900	33.6	(25.0 - 42.2)	9,643	1,208,300	41.7	(35.8 - 47.6)
\$50,000-\$74,999	*	*	*	*	5,761	795,600	40.4	(34.5 - 46.3)
≥ \$75,000	*	*	*	*	8,548	1,791,200	39.9	(36.6 - 43.2)

¹ Defined as having had a flu shot or flu vaccine sprayed in nose within the past 12 months

² Sample size from survey

³ Estimated number of adults who have had a flu shot or flu vaccine sprayed in nose within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 20b. Flu shot or flu vaccine sprayed in nose within the past 12 months¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	646	34.2	(29.1 - 39.4)	38,059	42.0	(39.6 - 44.5)
Gender						
Male	245	24.5	(18.3 - 30.8)	14,568	40.4	(36.6 - 44.2)
Female	401	43.8	(36.5 - 51.1)	23,491	43.5	(40.3 - 46.7)
Race/Ethnicity						
White Non-Hispanic	631	33.7	(28.5 - 38.9)	35,750	42.6	(40.7 - 44.6)
Black Non-Hispanic	*	*	*	738	42.4	(33.4 - 51.4)
Hispanic	*	*	*	860	40.0	(30.9 - 49.1)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	304	31.5	(24.5 - 38.5)	15,651	44.1	(39.6 - 48.6)
Some College	175	32.9	(23.1 - 42.7)	9,895	38.1	(33.2 - 43.0)
≥ College Degree	*	*	*	12,431	42.8	(39.4 - 46.3)
Income						
≤ \$24,999	*	*	*	9,626	48.0	(41.6 - 54.3)
\$25,000-\$49,999	201	31.7	(22.9 - 40.5)	9,643	40.5	(34.8 - 46.3)
\$50,000-\$74,999	*	*	*	5,761	39.6	(33.6 - 45.6)
≥ \$75,000	*	*	*	8,548	39.4	(36.1 - 42.7)

¹ Defined as having had a flu shot or flu vaccine sprayed in nose within the past 12 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 21a. Flu shot among adults aged 65 and older¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	217	3,400	70.0	(63.1 - 77.0)	12,602	1,907,700	75.2	(72.0 - 78.4)
Gender								
Male	*	*	*	*	4,417	772,900	74.6	(69.3 - 79.8)
Female	138	1,800	68.3	(59.2 - 77.3)	8,185	1,134,800	75.6	(71.6 - 79.6)
Age (years)								
≥ 65	217	3,400	70.0	(63.1 - 77.0)	12,602	1,907,700	75.2	(72.0 - 78.4)
Race/Ethnicity								
White Non-Hispanic	213	3,300	69.3	(62.3 - 76.4)	12,137	1,487,100	76.0	(73.5 - 78.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	118	1,700	69.9	(60.6 - 79.2)	6,518	930,800	74.0	(68.9 - 79.2)
Some College	*	*	*	*	2,790	317,800	69.7	(62.7 - 76.7)
≥ College Degree	*	*	*	*	3,261	654,000	80.5	(76.4 - 84.6)
Income								
≤ \$24,999	*	*	*	*	4,665	545,300	72.4	(65.6 - 79.3)
\$25,000-\$49,999	*	*	*	*	3,422	430,600	72.8	(67.0 - 78.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	1,198	301,000	77.5	(70.9 - 84.2)

¹ Defined as having had a flu shot during the past 12 months among adults aged 65 and older

² Sample size from survey

³ Estimated number of adults aged 65 and older who have had a flu shot within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 21b. Flu shot among adults aged 65 and older¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	217	70.2	(63.2 - 77.2)	12,602	75.0	(71.5 - 78.5)
Gender						
Male	*	*	*	4,417	74.4	(68.7 - 80.1)
Female	138	68.3	(59.2 - 77.3)	8,185	75.5	(71.3 - 79.8)
Race/Ethnicity						
White Non-Hispanic	213	69.5	(62.4 - 76.6)	12,137	75.9	(73.0 - 78.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	118	70.3	(61.0 - 79.6)	6,518	73.3	(67.5 - 79.1)
Some College	*	*	*	2,790	70.0	(62.7 - 77.2)
≥ College Degree	*	*	*	3,261	80.7	(76.2 - 85.1)
Income						
≤ \$24,999	*	*	*	4,665	72.6	(65.1 - 80.1)
\$25,000-\$49,999	*	*	*	3,422	71.4	(65.0 - 77.9)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	1,198	78.0	(70.8 - 85.1)

¹ Defined as having had a flu shot during the past 12 months among adults aged 65 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 22a. Pneumonia shot or pneumococcal vaccine¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	600	5,500	25.1	(20.8 - 29.4)	35,303	3,456,800	26.1	(23.9 - 28.3)
Gender								
Male	220	2,200	21.5	(14.9 - 28.1)	13,052	1,542,200	25.2	(21.5 - 28.8)
Female	380	3,200	28.4	(22.9 - 33.9)	22,251	1,914,600	26.9	(24.2 - 29.6)
Age (years)								
18-49	154	1,200	10.9	(4.4 - 17.4)	10,943	1,097,000	14.6	(11.3 - 17.9)
50-64	233	1,300	19.8	(14.4 - 25.2)	12,179	758,000	23.0	(19.8 - 26.2)
≥ 65	213	3,100	64.6	(57.4 - 71.9)	12,181	1,601,700	65.7	(61.7 - 69.7)
Race/Ethnicity								
White Non-Hispanic	589	5,300	24.6	(20.3 - 29.0)	33,225	2,358,700	28.7	(26.9 - 30.5)
Black Non-Hispanic	*	*	*	*	685	584,400	28.4	(19.5 - 37.3)
Hispanic	*	*	*	*	769	408,000	19.5	(12.6 - 26.4)
Other Non-Hispanic	*	*	*	*	624	105,600	12.2	(2.3 - 22.2)
Education								
≤ High School	286	3,000	28.1	(21.2 - 35.0)	14,700	1,535,300	30.4	(26.4 - 34.3)
Some College	158	1,200	19.7	(13.1 - 26.2)	9,151	836,200	27.2	(21.9 - 32.5)
≥ College Degree	155	1,200	25.3	(17.9 - 32.7)	11,375	1,070,700	21.4	(18.6 - 24.1)
Income								
≤ \$24,999	*	*	*	*	9,082	1,050,900	33.1	(27.6 - 38.6)
\$25,000-\$49,999	188	1,500	20.1	(13.5 - 26.7)	9,033	797,500	30.3	(24.1 - 36.6)
\$50,000-\$74,999	73	500	19.7	(10.2 - 29.1)	5,320	378,800	21.8	(17.4 - 26.1)
≥ \$75,000	107	700	14.3	(7.7 - 20.8)	7,773	648,700	16.3	(14.1 - 18.5)

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine

² Sample size from survey

³ Estimated number of adults who have ever had a pneumonia shot or pneumococcal vaccine (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 22b. Pneumonia shot or pneumococcal vaccine¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	600	23.5	(19.1 - 28.0)	35,303	25.4	(23.2 - 27.6)
Gender						
Male	220	21.0	(14.1 - 27.9)	13,052	25.8	(22.3 - 29.4)
Female	380	25.8	(20.2 - 31.5)	22,251	25.1	(22.4 - 27.8)
Race/Ethnicity						
White Non-Hispanic	589	23.0	(18.6 - 27.5)	33,225	27.7	(25.9 - 29.5)
Black Non-Hispanic	*	*	*	685	27.9	(19.1 - 36.7)
Hispanic	*	*	*	769	19.6	(12.8 - 26.5)
Other Non-Hispanic	*	*	*	624	11.3	(2.0 - 20.7)
Education						
≤ High School	286	27.0	(19.7 - 34.3)	14,700	29.5	(25.6 - 33.5)
Some College	158	17.1	(10.9 - 23.4)	9,151	26.2	(21.0 - 31.3)
≥ College Degree	155	24.0	(16.6 - 31.4)	11,375	21.0	(18.3 - 23.8)
Income						
≤ \$24,999	*	*	*	9,082	32.3	(26.9 - 37.8)
\$25,000-\$49,999	188	19.3	(12.4 - 26.2)	9,033	29.0	(23.0 - 35.0)
\$50,000-\$74,999	73	17.9	(8.8 - 27.0)	5,320	21.2	(16.8 - 25.7)
≥ \$75,000	107	13.4	(7.0 - 19.9)	7,773	16.2	(13.9 - 18.5)

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 23a. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	213	3,100	64.6	(57.4 - 71.9)	12,181	1,601,700	65.7	(61.7 - 69.7)
Gender								
Male	*	*	*	*	4,180	606,400	62.1	(55.5 - 68.7)
Female	136	1,700	64.6	(55.4 - 73.8)	8,001	995,300	68.0	(63.1 - 73.0)
Age (years)								
≥ 65	213	3,100	64.6	(57.4 - 71.9)	12,181	1,601,700	65.7	(61.7 - 69.7)
Race/Ethnicity								
White Non-Hispanic	209	2,900	63.8	(56.4 - 71.2)	11,735	1,281,300	68.2	(64.9 - 71.5)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	115	1,600	65.7	(56.1 - 75.3)	6,345	806,200	65.6	(59.5 - 71.7)
Some College	*	*	*	*	2,694	289,500	66.8	(60.1 - 73.4)
≥ College Degree	*	*	*	*	3,111	501,100	65.6	(58.5 - 72.7)
Income								
≤ \$24,999	89	1,400	80.4	(71.3 - 89.5)	4,533	497,100	67.7	(59.9 - 75.5)
\$25,000-\$49,999	*	*	*	*	3,324	344,300	61.5	(53.3 - 69.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	1,138	236,300	64.5	(55.2 - 73.8)

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine among adults aged 65 and older

² Sample size from survey

³ Estimated number of adults aged 65 and older who have ever had a pneumonia shot or pneumococcal vaccine (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 23b. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	213	64.6	(57.3 - 72.0)	12,181	64.7	(60.4 - 69.1)
Gender						
Male	*	*	*	4,180	61.5	(54.3 - 68.6)
Female	136	64.6	(55.4 - 73.8)	8,001	67.6	(62.4 - 72.9)
Race/Ethnicity						
White Non-Hispanic	209	63.8	(56.3 - 71.3)	11,735	67.1	(63.2 - 70.9)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	115	66.1	(56.4 - 75.8)	6,345	64.2	(57.4 - 71.0)
Some College	*	*	*	2,694	65.8	(58.4 - 73.3)
≥ College Degree	*	*	*	3,111	65.3	(57.7 - 72.9)
Income						
≤ \$24,999	89	79.9	(70.6 - 89.2)	4,533	66.0	(57.0 - 74.9)
\$25,000-\$49,999	*	*	*	3,324	60.3	(51.9 - 68.6)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	*	*

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine among adults aged 65 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 24a. Ever had a mammogram among women aged 40 and older¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	347	7,400	89.8	(85.7 - 93.9)	19,491	4,328,800	91.8	(89.7 - 94.0)
Gender								
Male	-	-	-	-	-	-	-	-
Female	347	7,400	89.8	(85.7 - 93.9)	19,491	4,328,800	91.8	(89.7 - 94.0)
Age (years)								
40-49	*	*	*	*	3,851	1,287,500	85.0	(79.7 - 90.3)
50-64	143	3,200	97.0	(94.3 - 99.6)	7,452	1,641,700	96.0	(94.1 - 97.9)
≥ 65	138	2,500	93.1	(88.8 - 97.5)	8,188	1,399,600	94.1	(90.5 - 97.6)
Race/Ethnicity								
White Non-Hispanic	339	7,300	89.8	(85.6 - 93.9)	18,548	2,976,100	93.9	(92.7 - 95.1)
Black Non-Hispanic	*	*	*	*	342	746,600	93.8	(87.3 - 100)
Hispanic	*	*	*	*	333	447,600	91.8	(84.6 - 99.0)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	160	3,200	84.6	(77.1 - 92.2)	8,266	1,698,700	91.0	(87.4 - 94.5)
Some College	94	2,100	93.2	(87.3 - 99.1)	5,322	1,055,500	93.6	(90.2 - 97.0)
≥ College Degree	92	2,100	95.1	(90.5 - 99.7)	5,872	1,560,700	91.6	(87.5 - 95.6)
Income								
≤ \$24,999	100	1,600	86.7	(77.9 - 95.6)	5,691	1,023,400	93.5	(89.7 - 97.4)
\$25,000-\$49,999	97	2,000	87.2	(77.8 - 96.6)	4,857	860,300	89.5	(83.6 - 95.5)
\$50,000-\$74,999	*	*	*	*	2,648	547,100	94.6	(92.2 - 96.9)
≥ \$75,000	57	1,600	91.1	(82.9 - 99.2)	3,582	1,185,100	92.2	(88.2 - 96.1)

¹ Defined as having ever had a mammogram among women aged 40 and older

² Sample size from survey

³ Estimated number of women aged 40 and older who have ever had a mammogram (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 24b. Ever had a mammogram among women aged 40 and older¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	347	88.0	(82.8 - 93.3)	19,491	91.1	(88.7 - 93.6)
Gender						
Male	-	-	-	-	-	-
Female	347	88.0	(82.8 - 93.3)	19,491	91.1	(88.7 - 93.6)
Race/Ethnicity						
White Non-Hispanic	339	88.0	(82.6 - 93.3)	18,548	93.3	(92.0 - 94.7)
Black Non-Hispanic	*	*	*	342	93.1	(85.7 - 100)
Hispanic	*	*	*	333	91.0	(82.8 - 99.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	160	82.3	(72.9 - 91.8)	8,266	90.2	(86.1 - 94.2)
Some College	94	91.2	(83.2 - 99.3)	5,322	92.9	(88.9 - 96.9)
≥ College Degree	92	94.4	(88.7 - 100)	5,872	90.9	(86.6 - 95.3)
Income						
≤ \$24,999	*	*	*	5,691	93.0	(88.5 - 97.4)
\$25,000-\$49,999	*	*	*	4,857	88.8	(82.5 - 95.1)
\$50,000-\$74,999	*	*	*	2,648	94.3	(91.7 - 96.8)
≥ \$75,000	*	*	*	3,582	91.2	(86.8 - 95.6)

¹ Defined as having ever had a mammogram among women aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 25a. Had mammogram within the past 2 years among women aged 40 and older¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	344	6,500	78.7	(73.5 - 83.8)	19,376	3,760,000	80.3	(77.8 - 82.9)
Gender								
Male	-	-	-	-	-	-	-	-
Female	344	6,500	78.7	(73.5 - 83.8)	19,376	3,760,000	80.3	(77.8 - 82.9)
Age (years)								
40-49	*	*	*	*	3,842	1,122,700	74.4	(68.5 - 80.2)
50-64	143	2,900	88.2	(82.5 - 93.9)	7,428	1,487,900	87.2	(84.5 - 89.8)
≥ 65	135	2,000	76.6	(68.5 - 84.6)	8,106	1,149,400	78.5	(74.2 - 82.9)
Race/Ethnicity								
White Non-Hispanic	336	6,300	78.7	(73.6 - 83.9)	18,438	2,531,500	80.2	(78.3 - 82.2)
Black Non-Hispanic	*	*	*	*	339	667,100	85.6	(77.5 - 93.8)
Hispanic	*	*	*	*	332	419,700	86.1	(77.5 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	159	2,900	75.9	(67.7 - 84.1)	8,204	1,486,900	80.5	(76.4 - 84.6)
Some College	93	1,700	76.1	(66.1 - 86.1)	5,297	929,400	83.0	(78.8 - 87.1)
≥ College Degree	91	1,900	85.7	(77.9 - 93.6)	5,845	1,334,800	78.6	(73.9 - 83.2)
Income								
≤ \$24,999	*	*	*	*	5,650	882,300	80.8	(75.9 - 85.7)
\$25,000-\$49,999	*	*	*	*	4,846	732,300	76.3	(69.7 - 82.9)
\$50,000-\$74,999	*	*	*	*	2,641	494,200	85.6	(81.9 - 89.4)
≥ \$75,000	*	*	*	*	3,572	1,025,900	80.0	(75.2 - 84.8)

¹ Defined as having had a mammogram within the past 2 years among women aged 40 and older

² Sample size from survey

³ Estimated number of women aged 40 and older who have had a mammogram within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 25b. Had mammogram within the past 2 years among women aged 40 and older¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	344	77.2	(71.3 - 83.2)	19,376	79.7	(76.9 - 82.5)
Gender						
Male	-	-	-	-	-	-
Female	344	77.2	(71.3 - 83.2)	19,376	79.7	(76.9 - 82.5)
Race/Ethnicity						
White Non-Hispanic	336	77.2	(71.2 - 83.2)	18,438	79.7	(77.6 - 81.8)
Black Non-Hispanic	*	*	*	339	85.3	(76.5 - 94.2)
Hispanic	*	*	*	332	85.2	(75.6 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	159	74.3	(64.6 - 84.0)	8,204	80.1	(75.6 - 84.7)
Some College	*	*	*	5,297	82.3	(77.7 - 86.9)
≥ College Degree	91	83.8	(74.3 - 93.3)	5,845	77.8	(72.9 - 82.8)
Income						
≤ \$24,999	*	*	*	5,650	80.8	(75.4 - 86.2)
\$25,000-\$49,999	*	*	*	4,846	75.7	(68.7 - 82.6)
\$50,000-\$74,999	*	*	*	2,641	85.5	(81.6 - 89.4)
≥ \$75,000	*	*	*	3,572	79.0	(73.8 - 84.2)

¹ Defined as having had a mammogram within the past 2 years among women aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 26a. Had mammogram within the past 2 years among women aged 50 and older¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	278	4,900	83.0	(78.2 - 87.9)	15,534	2,637,300	83.2	(80.7 - 85.7)
Gender								
Male	-	-	-	-	-	-	-	-
Female	278	4,900	83.0	(78.2 - 87.9)	15,534	2,637,300	83.2	(80.7 - 85.7)
Age (years)								
50-64	143	2,900	88.2	(82.5 - 93.9)	7,428	1,487,900	87.2	(84.5 - 89.8)
≥ 65	135	2,000	76.6	(68.5 - 84.6)	8,106	1,149,400	78.5	(74.2 - 82.9)
Race/Ethnicity								
White Non-Hispanic	272	4,800	83.4	(78.6 - 88.3)	14,887	1,907,000	82.3	(80.3 - 84.4)
Black Non-Hispanic	*	*	*	*	249	385,600	89.9	(83.2 - 96.7)
Hispanic	*	*	*	*	220	286,700	93.9	(89.1 - 98.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	134	2,300	82.2	(75.3 - 89.1)	7,026	1,094,500	83.3	(80.0 - 86.6)
Some College	*	*	*	*	4,124	646,700	87.3	(84.2 - 90.3)
≥ College Degree	68	1,300	90.4	(83.4 - 97.4)	4,356	890,500	80.6	(75.1 - 86.1)
Income								
≤ \$24,999	*	*	*	*	4,991	649,100	81.0	(76.8 - 85.2)
\$25,000-\$49,999	73	1,300	87.5	(78.8 - 96.1)	3,913	533,300	78.6	(70.9 - 86.2)
\$50,000-\$74,999	*	*	*	*	1,943	359,300	89.6	(86.3 - 93.0)
≥ \$75,000	*	*	*	*	2,303	601,300	84.6	(79.3 - 89.9)

¹ Defined as having had a mammogram within the past 2 years among women aged 50 and older

² Sample size from survey

³ Estimated number of women aged 50 and older who have had a mammogram within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 26b. Had mammogram within the past 2 years among women aged 50 and older¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	278	83.3	(78.5 - 88.1)	15,534	83.1	(80.3 - 85.8)
Gender						
Male	-	-	-	-	-	-
Female	278	83.3	(78.5 - 88.1)	15,534	83.1	(80.3 - 85.8)
Race/Ethnicity						
White Non-Hispanic	272	83.7	(78.9 - 88.5)	14,887	82.1	(79.9 - 84.3)
Black Non-Hispanic	*	*	*	249	90.4	(83.6 - 97.2)
Hispanic	*	*	*	220	94.1	(89.3 - 99.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	134	82.7	(75.9 - 89.5)	7,026	83.6	(80.1 - 87.0)
Some College	*	*	*	4,124	87.3	(84.1 - 90.5)
≥ College Degree	68	90.1	(82.8 - 97.3)	4,356	79.9	(73.9 - 86.0)
Income						
≤ \$24,999	*	*	*	4,991	81.3	(76.9 - 85.6)
\$25,000-\$49,999	73	87.7	(79.2 - 96.2)	3,913	78.1	(69.9 - 86.4)
\$50,000-\$74,999	*	*	*	1,943	90.2	(87.1 - 93.4)
≥ \$75,000	*	*	*	2,303	83.9	(77.9 - 89.9)

¹ Defined as having had a mammogram within the past 2 years among women aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 27a. Ever had a Pap test among women¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	399	12,300	95.6	(91.3 - 99.9)	23,381	7,045,300	92.1	(90.0 - 94.3)
Gender								
Male	-	-	-	-	-	-	-	-
Female	399	12,300	95.6	(91.3 - 99.9)	23,381	7,045,300	92.1	(90.0 - 94.3)
Age (years)								
18-34	*	*	*	*	2,377	1,896,600	85.2	(79.3 - 91.0)
35-44	*	*	*	*	3,023	1,301,500	92.6	(87.7 - 97.5)
45-54	85	2,400	100	(100 - 100)	4,657	1,420,300	98.3	(97.0 - 99.6)
55-64	98	2,000	99.5	(98.5 - 100)	5,187	1,071,200	97.2	(95.0 - 99.3)
≥ 65	138	2,500	94.6	(91.4 - 97.8)	8,137	1,355,700	92.3	(88.5 - 96.1)
Race/Ethnicity								
White Non-Hispanic	388	12,000	95.7	(91.3 - 100)	21,976	4,439,700	94.2	(92.1 - 96.3)
Black Non-Hispanic	*	*	*	*	481	1,231,600	91.2	(84.0 - 98.4)
Hispanic	*	*	*	*	533	984,500	90.1	(85.3 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	186	5,800	98.4	(97.2 - 99.6)	9,526	2,540,500	88.3	(83.8 - 92.7)
Some College	*	*	*	*	6,462	1,782,200	93.5	(90.4 - 96.6)
≥ College Degree	102	2,900	99.6	(98.7 - 100)	7,345	2,643,200	95.0	(92.2 - 97.9)
Income								
≤ \$24,999	108	2,500	97.7	(95.5 - 99.8)	6,526	1,717,200	93.7	(90.9 - 96.6)
\$25,000-\$49,999	120	3,900	94.5	(85.2 - 100)	5,787	1,378,400	91.0	(85.6 - 96.4)
\$50,000-\$74,999	*	*	*	*	3,310	933,900	96.0	(91.8 - 100)
≥ \$75,000	63	2,400	100	(100 - 100)	4,570	1,943,300	95.4	(92.6 - 98.2)

¹ Defined as having ever had a Pap test among women

² Sample size from survey

³ Estimated number of women who have ever had a Pap test (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 27b. Ever had a Pap test among women¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	399	95.6	(91.1 - 100)	23,381	91.8	(89.6 - 94.0)
Gender						
Male	-	-	-	-	-	-
Female	399	95.6	(91.1 - 100)	23,381	91.8	(89.6 - 94.0)
Race/Ethnicity						
White Non-Hispanic	388	95.6	(91.0 - 100)	21,976	94.0	(91.8 - 96.2)
Black Non-Hispanic	*	*	*	481	90.8	(83.5 - 98.1)
Hispanic	*	*	*	533	89.7	(84.6 - 94.7)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	186	98.7	(97.7 - 99.7)	9,526	87.8	(83.2 - 92.4)
Some College	*	*	*	6,462	93.1	(89.7 - 96.4)
≥ College Degree	102	99.6	(98.9 - 100)	7,345	94.9	(92.0 - 97.7)
Income						
≤ \$24,999	108	98.0	(96.1 - 99.8)	6,526	93.3	(90.1 - 96.5)
\$25,000-\$49,999	120	94.4	(84.9 - 100)	5,787	91.1	(85.7 - 96.5)
\$50,000-\$74,999	*	*	*	3,310	96.0	(91.9 - 100)
≥ \$75,000	63	100	(100 - 100)	4,570	95.0	(91.9 - 98.1)

¹ Defined as having ever had a Pap test among women

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 28a. Had a Pap test within the past 3 years among women¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	391	9,300	73.4	(67.2 - 79.6)	23,109	6,205,600	82.2	(79.8 - 84.6)
Gender								
Male	-	-	-	-	-	-	-	-
Female	391	9,300	73.4	(67.2 - 79.6)	23,109	6,205,600	82.2	(79.8 - 84.6)
Age (years)								
18-34	*	*	*	*	2,371	1,818,400	81.8	(75.5 - 88.1)
35-44	*	*	*	*	3,012	1,220,300	86.9	(81.8 - 92.0)
45-54	85	2,000	86.7	(79.4 - 94.0)	4,640	1,311,500	92.0	(89.8 - 94.3)
55-64	97	1,500	75.2	(66.0 - 84.3)	5,152	924,300	84.5	(81.0 - 88.0)
≥ 65	131	1,400	53.6	(44.1 - 63.0)	7,934	931,100	66.4	(61.8 - 71.0)
Race/Ethnicity								
White Non-Hispanic	380	9,100	73.3	(67.0 - 79.7)	21,720	3,807,100	81.5	(79.2 - 83.8)
Black Non-Hispanic	*	*	*	*	473	1,142,800	86.7	(79.1 - 94.3)
Hispanic	*	*	*	*	527	894,900	83.6	(76.3 - 91.0)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	183	4,200	71.0	(62.4 - 79.6)	9,384	2,132,600	76.0	(71.1 - 81.0)
Some College	*	*	*	*	6,401	1,594,300	84.0	(80.4 - 87.5)
≥ College Degree	98	2,400	83.2	(75.5 - 90.8)	7,281	2,406,000	86.9	(83.7 - 90.1)
Income								
≤ \$24,999	*	*	*	*	6,441	1,453,900	82.0	(77.9 - 86.0)
\$25,000-\$49,999	*	*	*	*	5,740	1,182,000	78.3	(72.6 - 83.9)
\$50,000-\$74,999	*	*	*	*	3,296	825,000	85.0	(78.0 - 91.9)
≥ \$75,000	62	2,000	85.4	(76.7 - 94.2)	4,555	1,810,200	89.0	(85.7 - 92.2)

¹ Defined as having had a Pap test within the past 3 years among women

² Sample size from survey

³ Estimated number of women who have had a Pap test within the past 3 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 28b. Had a Pap test within the past 3 years among women¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	391	74.4	(67.8 - 81.0)	23,109	82.7	(80.2 - 85.1)
Gender						
Male	-	-	-	-	-	-
Female	391	74.4	(67.8 - 81.0)	23,109	82.7	(80.2 - 85.1)
Race/Ethnicity						
White Non-Hispanic	380	74.2	(67.5 - 81.0)	21,720	82.3	(80.0 - 84.6)
Black Non-Hispanic	*	*	*	473	87.0	(79.3 - 94.6)
Hispanic	*	*	*	527	83.4	(75.9 - 90.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	183	72.5	(63.4 - 81.6)	9,384	76.4	(71.3 - 81.5)
Some College	*	*	*	6,401	84.2	(80.5 - 88.0)
≥ College Degree	98	84.2	(76.3 - 92.1)	7,281	87.5	(84.3 - 90.6)
Income						
≤ \$24,999	*	*	*	6,441	82.7	(78.5 - 86.9)
\$25,000-\$49,999	*	*	*	5,740	79.2	(73.5 - 84.8)
\$50,000-\$74,999	*	*	*	3,296	85.7	(78.7 - 92.8)
≥ \$75,000	62	85.9	(76.6 - 95.1)	4,555	88.9	(85.5 - 92.3)

¹ Defined as having had a Pap test within the past 3 years among women

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 29a. Ever had a digital rectal exam among men aged 40 and older¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	218	6,800	75.8	(67.6 - 84.1)	12,033	3,176,500	73.8	(69.7 - 77.8)
Gender								
Male	218	6,800	75.8	(67.6 - 84.1)	12,033	3,176,500	73.8	(69.7 - 77.8)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	3,098	922,800	66.8	(59.9 - 73.8)
55-64	66	1,700	82.4	(73.0 - 91.8)	3,504	865,600	87.1	(82.5 - 91.7)
≥ 65	79	1,900	87.7	(80.0 - 95.4)	4,348	953,900	92.9	(91.0 - 94.9)
Race/Ethnicity								
White Non-Hispanic	214	6,500	76.6	(68.5 - 84.7)	11,411	2,452,100	80.3	(77.7 - 82.9)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	5,013	931,600	64.1	(57.3 - 71.0)
Some College	*	*	*	*	2,695	674,200	75.5	(67.8 - 83.2)
≥ College Degree	*	*	*	*	4,302	1,568,000	80.5	(74.0 - 87.0)
Income								
≤ \$24,999	*	*	*	*	2,573	573,300	65.9	(56.0 - 75.9)
\$25,000-\$49,999	*	*	*	*	3,168	530,100	74.6	(67.2 - 82.1)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,309	1,351,100	78.0	(72.6 - 83.5)

¹ Defined as having ever had a digital rectal exam among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have ever had a digital rectal exam (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 29b. Ever had a digital rectal exam among men aged 40 and older¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	218	74.9	(65.9 - 84.0)	12,033	73.7	(69.6 - 77.8)
Gender						
Male	218	74.9	(65.9 - 84.0)	12,033	73.7	(69.6 - 77.8)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	214	75.9	(66.9 - 84.8)	11,411	79.9	(77.2 - 82.5)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	5,013	64.7	(57.8 - 71.7)
Some College	*	*	*	2,695	75.0	(67.1 - 82.9)
≥ College Degree	*	*	*	4,302	80.3	(73.8 - 86.8)
Income						
≤ \$24,999	*	*	*	2,573	66.9	(56.9 - 76.9)
\$25,000-\$49,999	*	*	*	3,168	74.8	(67.3 - 82.2)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,309	77.5	(72.0 - 83.0)

¹ Defined as having ever had a digital rectal exam among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 30a. Had digital rectal exam within the past 2 years among men aged 40 and older¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	217	5,100	57.2	(48.5 - 65.9)	11,923	2,380,300	55.8	(51.8 - 59.8)
Gender								
Male	217	5,100	57.2	(48.5 - 65.9)	11,923	2,380,300	55.8	(51.8 - 59.8)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	1,078	246,800	27.5	(18.5 - 36.5)
45-54	*	*	*	*	3,074	650,000	47.6	(40.8 - 54.4)
55-64	*	*	*	*	3,467	673,500	68.1	(62.4 - 73.8)
≥ 65	*	*	*	*	4,304	809,900	79.7	(75.0 - 84.3)
Race/Ethnicity								
White Non-Hispanic	213	4,900	57.1	(48.4 - 65.8)	11,305	1,875,600	62.2	(59.2 - 65.2)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	4,963	718,700	49.6	(43.2 - 56.1)
Some College	*	*	*	*	2,665	511,500	58.5	(50.8 - 66.3)
≥ College Degree	*	*	*	*	4,277	1,148,400	59.4	(52.8 - 66.0)
Income								
≤ \$24,999	*	*	*	*	2,553	388,000	45.0	(35.0 - 55.0)
\$25,000-\$49,999	*	*	*	*	3,139	422,400	60.5	(53.0 - 68.0)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,295	978,300	56.9	(51.1 - 62.7)

¹ Defined as having had a digital rectal exam within the past 2 years among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have had a digital rectal exam within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 30b. Had digital rectal exam within the past 2 years among men aged 40 and older¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	217	56.7	(47.3 - 66.1)	11,923	55.9	(51.7 - 60.0)
Gender						
Male	217	56.7	(47.3 - 66.1)	11,923	55.9	(51.7 - 60.0)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	213	56.8	(47.4 - 66.2)	11,305	62.0	(58.9 - 65.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	4,963	50.5	(43.7 - 57.3)
Some College	*	*	*	2,665	58.3	(50.4 - 66.2)
≥ College Degree	*	*	*	4,277	59.2	(52.4 - 65.9)
Income						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,139	60.8	(53.1 - 68.4)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,295	56.5	(50.5 - 62.4)

¹ Defined as having had a digital rectal exam within the past 2 years among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 31a. Ever had a prostate specific antigen test among men aged 40 and older¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	201	4,800	58.2	(49.1 - 67.3)	11,557	2,822,600	67.6	(63.9 - 71.2)
Gender								
Male	201	4,800	58.2	(49.1 - 67.3)	11,557	2,822,600	67.6	(63.9 - 71.2)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,938	760,600	57.4	(50.6 - 64.3)
55-64	*	*	*	*	3,373	811,400	83.4	(78.8 - 88.1)
≥ 65	72	1,700	82.5	(73.3 - 91.7)	4,215	905,400	90.4	(87.1 - 93.6)
Race/Ethnicity								
White Non-Hispanic	197	4,600	58.2	(49.0 - 67.3)	10,950	1,965,400	67.0	(63.9 - 70.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	4,775	829,200	59.0	(52.4 - 65.6)
Some College	*	*	*	*	2,584	535,000	61.9	(53.7 - 70.2)
≥ College Degree	*	*	*	*	4,176	1,455,700	76.8	(72.5 - 81.1)
Income								
≤ \$24,999	*	*	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	*	3,048	444,100	64.1	(56.0 - 72.3)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,188	1,211,100	72.3	(67.6 - 77.0)

¹ Defined as having ever had a prostate specific antigen test among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have ever had a prostate specific antigen test (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 31b. Ever had a prostate specific antigen test among men aged 40 and older¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	201	57.7	(47.8 - 67.5)	11,557	67.4	(63.7 - 71.1)
Gender						
Male	201	57.7	(47.8 - 67.5)	11,557	67.4	(63.7 - 71.1)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	197	57.8	(47.9 - 67.7)	10,950	66.6	(63.4 - 69.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	4,775	59.7	(52.9 - 66.4)
Some College	*	*	*	2,584	61.4	(53.0 - 69.9)
≥ College Degree	*	*	*	4,176	76.3	(71.9 - 80.7)
Income						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,048	64.2	(56.1 - 72.4)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,188	71.5	(66.6 - 76.3)

¹ Defined as having ever had a prostate specific antigen test among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 32a. Had prostate specific antigen test within the past 2 years among men aged 40 and older¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	201	4,200	50.8	(41.8 - 59.9)	11,437	2,493,800	59.9	(56.0 - 63.9)
Gender								
Male	201	4,200	50.8	(41.8 - 59.9)	11,437	2,493,800	59.9	(56.0 - 63.9)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,917	659,400	49.9	(42.8 - 57.0)
55-64	*	*	*	*	3,326	719,700	74.5	(69.4 - 79.7)
≥ 65	*	*	*	*	4,166	830,000	83.3	(78.4 - 88.3)
Race/Ethnicity								
White Non-Hispanic	197	4,000	50.4	(41.4 - 59.5)	10,836	1,762,200	60.4	(57.3 - 63.5)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	4,727	734,200	52.4	(45.7 - 59.1)
Some College	*	*	*	*	2,558	483,800	56.3	(48.2 - 64.4)
≥ College Degree	*	*	*	*	4,133	1,273,200	67.4	(61.6 - 73.3)
Income								
≤ \$24,999	*	*	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	*	3,021	410,800	59.6	(51.5 - 67.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,170	1,080,100	64.6	(59.1 - 70.0)

¹ Defined as having had a prostate specific antigen test within the past 2 years among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have had a prostate specific antigen test within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 32b. Had prostate specific antigen test within the past 2 years among men aged 40 and older¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	201	50.0	(40.3 - 59.7)	11,437	59.8	(55.8 - 63.9)
Gender						
Male	201	50.0	(40.3 - 59.7)	11,437	59.8	(55.8 - 63.9)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	197	49.8	(40.0 - 59.5)	10,836	60.1	(56.9 - 63.4)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	4,727	52.9	(45.9 - 59.9)
Some College	*	*	*	2,558	56.0	(47.7 - 64.2)
≥ College Degree	*	*	*	4,133	67.1	(61.2 - 73.1)
Income						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,021	59.8	(51.7 - 67.9)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,170	63.9	(58.3 - 69.6)

¹ Defined as having had a prostate specific antigen test within the past 2 years among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 33a. Prostate cancer among men aged 40 and older¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	217	400	4.0	(1.4 - 6.6)	12,133	178,200	4.1	(3.0 - 5.2)
Gender								
Male	217	400	4.0	(1.4 - 6.6)	12,133	178,200	4.1	(3.0 - 5.2)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	3,120	6,100	0.4	(0.0 - 0.9)
55-64	*	*	*	*	3,532	31,600	3.1	(1.5 - 4.8)
≥ 65	78	300	15.0	(5.6 - 24.5)	4,388	140,400	13.6	(9.7 - 17.4)
Race/Ethnicity								
White Non-Hispanic	213	400	4.2	(1.5 - 6.9)	11,502	143,700	4.7	(3.5 - 5.8)
Black Non-Hispanic	*	*	*	*	184	23,500	5.1	(0.0 - 11.0)
Hispanic	*	*	*	*	197	7,700	1.6	(0.0 - 3.7)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	5,065	46,300	3.1	(2.2 - 4.1)
Some College	*	*	*	*	2,709	39,200	4.4	(1.3 - 7.5)
≥ College Degree	*	*	*	*	4,336	92,500	4.7	(3.0 - 6.5)
Income								
≤ \$24,999	*	*	*	*	2,597	35,800	4.1	(2.2 - 6.0)
\$25,000-\$49,999	*	*	*	*	3,196	29,600	4.1	(2.9 - 5.4)
\$50,000-\$74,999	*	*	*	*	2,007	20,300	2.8	(1.2 - 4.4)
≥ \$75,000	*	*	*	*	3,322	56,500	3.2	(1.7 - 4.8)

¹ Defined as having ever been told by a health professional that respondent had prostate cancer among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have ever been told they had prostate cancer (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 33b. Prostate cancer among men aged 40 and older¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	217	4.0	(1.4 - 6.7)	12,133	4.5	(3.2 - 5.7)
Gender						
Male	217	4.0	(1.4 - 6.7)	12,133	4.5	(3.2 - 5.7)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	213	4.2	(1.4 - 7.0)	11,502	5.1	(3.8 - 6.4)
Black Non-Hispanic	*	*	*	184	5.7	(0.0 - 12.8)
Hispanic	*	*	*	197	1.8	(0.0 - 4.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	5,065	3.3	(2.3 - 4.3)
Some College	*	*	*	2,709	5.0	(1.1 - 9.0)
≥ College Degree	*	*	*	4,336	5.2	(3.1 - 7.2)
Income						
≤ \$24,999	*	*	*	2,597	4.5	(2.2 - 6.7)
\$25,000-\$49,999	*	*	*	3,196	4.3	(3.0 - 5.6)
\$50,000-\$74,999	*	*	*	2,007	3.0	(1.1 - 5.0)
≥ \$75,000	*	*	*	3,322	3.5	(1.6 - 5.4)

¹ Defined as having ever been told by a health professional that respondent had prostate cancer among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 34a. Home blood stool test ever used among adults aged 50 and older¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	457	5,200	44.7	(39.5 - 49.9)	25,322	2,288,500	38.0	(35.5 - 40.4)
Gender								
Male	177	2,300	40.1	(32.0 - 48.2)	9,571	1,081,900	38.9	(35.1 - 42.6)
Female	280	2,900	49.2	(42.7 - 55.7)	15,751	1,206,600	37.2	(34.0 - 40.4)
Age (years)								
50-64	243	2,600	37.2	(30.2 - 44.2)	12,834	1,192,000	34.0	(30.8 - 37.2)
≥ 65	214	2,700	55.5	(48.0 - 63.0)	12,472	1,093,400	43.6	(39.8 - 47.4)
Race/Ethnicity								
White Non-Hispanic	448	4,900	43.4	(38.3 - 48.6)	24,213	1,754,500	39.1	(37.1 - 41.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	220	2,200	39.4	(32.3 - 46.6)	11,146	823,500	34.9	(30.5 - 39.2)
Some College	*	*	*	*	6,261	480,300	37.0	(32.6 - 41.4)
≥ College Degree	*	*	*	*	7,851	973,700	41.6	(37.8 - 45.3)
Income								
≤ \$24,999	141	1,300	45.3	(36.0 - 54.7)	7,222	505,000	35.4	(29.7 - 41.2)
\$25,000-\$49,999	*	*	*	*	6,609	467,600	40.3	(35.2 - 45.4)
\$50,000-\$74,999	*	*	*	*	3,500	313,000	38.2	(32.6 - 43.9)
≥ \$75,000	*	*	*	*	4,647	657,800	37.3	(33.0 - 41.6)

¹ Defined as having ever used a home blood stool test among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have ever used a home blood stool test (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 34b. Home blood stool test ever used among adults aged 50 and older¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	457	44.6	(39.2 - 49.9)	25,322	38.1	(35.5 - 40.8)
Gender						
Male	177	40.4	(32.2 - 48.6)	9,571	39.4	(35.3 - 43.5)
Female	280	48.9	(42.3 - 55.5)	15,751	36.9	(33.5 - 40.3)
Race/Ethnicity						
White Non-Hispanic	448	43.3	(38.0 - 48.5)	24,213	38.9	(36.7 - 41.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	220	39.1	(31.8 - 46.4)	11,146	35.1	(30.3 - 39.9)
Some College	*	*	*	6,261	36.6	(32.0 - 41.3)
≥ College Degree	*	*	*	7,851	41.9	(37.9 - 46.0)
Income						
≤ \$24,999	141	45.6	(36.2 - 55.1)	7,222	35.9	(29.5 - 42.4)
\$25,000-\$49,999	*	*	*	6,609	40.2	(34.8 - 45.6)
\$50,000-\$74,999	*	*	*	3,500	38.1	(31.9 - 44.3)
≥ \$75,000	*	*	*	4,647	37.5	(32.9 - 42.2)

¹ Defined as having ever used a home blood stool test among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 35a. Home blood stool test used within the past year among adults aged 50 and older¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	453	1,800	15.9	(12.2 - 19.6)	25,074	863,300	14.4	(12.6 - 16.2)
Gender								
Male	176	1,000	16.4	(10.6 - 22.2)	9,508	444,400	16.0	(13.2 - 18.9)
Female	277	900	15.4	(10.9 - 19.9)	15,566	418,900	13.1	(10.8 - 15.4)
Age (years)								
50-64	241	900	13.7	(9.0 - 18.3)	12,759	454,400	13.0	(10.6 - 15.5)
≥ 65	212	900	19.2	(13.2 - 25.1)	12,299	406,400	16.4	(13.7 - 19.0)
Race/Ethnicity								
White Non-Hispanic	444	1,800	15.8	(12.1 - 19.5)	23,971	634,300	14.3	(12.8 - 15.7)
Black Non-Hispanic	*	*	*	*	396	136,000	18.6	(9.4 - 27.8)
Hispanic	*	*	*	*	349	53,400	9.9	(2.5 - 17.2)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	218	900	15.5	(10.4 - 20.6)	11,026	288,600	12.3	(9.4 - 15.3)
Some College	120	400	14.1	(7.1 - 21.1)	6,208	212,500	16.4	(12.5 - 20.4)
≥ College Degree	114	500	17.8	(10.0 - 25.6)	7,778	352,400	15.2	(12.5 - 17.9)
Income								
≤ \$24,999	139	500	16.4	(9.5 - 23.3)	7,157	196,300	13.9	(10.1 - 17.6)
\$25,000-\$49,999	127	500	15.4	(8.5 - 22.3)	6,562	221,500	19.2	(14.1 - 24.3)
\$50,000-\$74,999	*	*	*	*	3,469	111,900	13.7	(10.1 - 17.4)
≥ \$75,000	80	400	15.6	(7.3 - 23.9)	4,623	205,100	11.7	(8.8 - 14.6)

¹ Defined as having used a home blood stool test within the past year among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have used a home blood stool test within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 35b. Home blood stool test used within the past year among adults aged 50 and older¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	453	16.0	(12.2 - 19.7)	25,074	14.6	(12.6 - 16.7)
Gender						
Male	176	16.4	(10.5 - 22.2)	9,508	16.4	(13.2 - 19.6)
Female	277	15.6	(11.0 - 20.2)	15,566	12.9	(10.4 - 15.4)
Race/Ethnicity						
White Non-Hispanic	444	15.9	(12.2 - 19.7)	23,971	14.2	(12.7 - 15.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	349	10.2	(1.7 - 18.7)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	218	15.5	(10.4 - 20.6)	11,026	12.7	(9.2 - 16.1)
Some College	120	14.5	(7.2 - 21.8)	6,208	16.6	(12.3 - 20.9)
≥ College Degree	114	17.5	(9.7 - 25.4)	7,778	15.3	(12.4 - 18.3)
Income						
≤ \$24,999	139	16.9	(9.8 - 24.1)	7,157	14.3	(9.8 - 18.7)
\$25,000-\$49,999	127	15.6	(8.5 - 22.7)	6,562	19.3	(13.8 - 24.8)
\$50,000-\$74,999	*	*	*	3,469	13.8	(9.8 - 17.7)
≥ \$75,000	80	15.6	(7.1 - 24.1)	4,623	11.8	(8.6 - 15.0)

¹ Defined as having used a home blood stool test within the past year among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 36a. Home blood stool test used within the past 2 years among adults aged 50 and older¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	453	2,800	24.3	(19.8 - 28.9)	25,074	1,218,000	20.4	(18.3 - 22.5)
Gender								
Male	176	1,500	26.0	(18.6 - 33.5)	9,508	593,100	21.4	(18.3 - 24.5)
Female	277	1,300	22.7	(17.4 - 28.0)	15,566	624,900	19.5	(16.6 - 22.4)
Age (years)								
50-64	241	1,400	20.8	(14.7 - 26.9)	12,759	629,900	18.1	(15.4 - 20.7)
≥ 65	212	1,400	29.5	(22.6 - 36.4)	12,299	585,500	23.6	(20.2 - 27.0)
Race/Ethnicity								
White Non-Hispanic	444	2,600	23.3	(19.0 - 27.6)	23,971	877,500	19.7	(18.2 - 21.3)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	218	1,100	20.0	(14.4 - 25.6)	11,026	454,600	19.4	(15.6 - 23.3)
Some College	*	*	*	*	6,208	262,200	20.3	(16.3 - 24.2)
≥ College Degree	114	900	28.1	(18.9 - 37.3)	7,778	491,400	21.2	(18.1 - 24.2)
Income								
≤ \$24,999	139	700	25.0	(17.0 - 33.1)	7,157	263,900	18.6	(14.6 - 22.7)
\$25,000-\$49,999	127	900	25.8	(15.9 - 35.6)	6,562	285,900	24.8	(19.6 - 30.0)
\$50,000-\$74,999	*	*	*	*	3,469	167,500	20.6	(15.6 - 25.5)
≥ \$75,000	80	600	21.5	(12.1 - 30.8)	4,623	294,900	16.8	(13.7 - 20.0)

¹ Defined as having used a home blood stool test within the past 2 years among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have used a home blood stool test within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 36b. Home blood stool test used within the past 2 years among adults aged 50 and older¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	453	24.5	(19.8 - 29.2)	25,074	20.5	(18.2 - 22.8)
Gender						
Male	176	26.2	(18.6 - 33.8)	9,508	21.7	(18.3 - 25.1)
Female	277	22.7	(17.3 - 28.0)	15,566	19.4	(16.3 - 22.5)
Race/Ethnicity						
White Non-Hispanic	444	23.4	(19.0 - 27.8)	23,971	19.7	(18.0 - 21.3)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	218	19.9	(14.3 - 25.5)	11,026	19.5	(15.4 - 23.7)
Some College	*	*	*	6,208	20.4	(16.1 - 24.7)
≥ College Degree	114	28.5	(19.1 - 38.0)	7,778	21.4	(18.1 - 24.7)
Income						
≤ \$24,999	139	25.8	(17.4 - 34.1)	7,157	18.9	(14.2 - 23.6)
\$25,000-\$49,999	*	*	*	6,562	24.8	(19.3 - 30.3)
\$50,000-\$74,999	*	*	*	3,469	20.6	(15.3 - 25.9)
≥ \$75,000	80	21.7	(12.1 - 31.3)	4,623	16.9	(13.4 - 20.4)

¹ Defined as having used a home blood stool test within the past 2 years among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 37a. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	460	7,000	59.2	(54.0 - 64.4)	25,357	4,019,100	66.6	(64.1 - 69.1)
Gender								
Male	176	3,100	54.1	(45.8 - 62.3)	9,582	1,894,100	68.0	(64.2 - 71.7)
Female	284	3,800	64.1	(58.0 - 70.3)	15,775	2,125,000	65.5	(62.1 - 68.8)
Age (years)								
50-54	*	*	*	*	4,179	718,900	50.6	(44.6 - 56.5)
55-64	164	2,300	57.2	(48.8 - 65.7)	8,695	1,493,500	71.0	(67.3 - 74.8)
≥ 65	214	3,300	68.8	(61.9 - 75.7)	12,466	1,802,000	72.1	(68.6 - 75.7)
Race/Ethnicity								
White Non-Hispanic	450	6,800	59.5	(54.4 - 64.7)	24,248	3,093,900	68.8	(66.8 - 70.8)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	223	3,100	55.2	(47.8 - 62.7)	11,121	1,445,100	61.2	(56.7 - 65.8)
Some College	*	*	*	*	6,287	835,300	64.3	(59.7 - 69.0)
≥ College Degree	116	2,200	72.1	(63.0 - 81.2)	7,884	1,732,400	73.7	(70.0 - 77.4)
Income								
≤ \$24,999	140	1,500	52.1	(42.6 - 61.6)	7,220	892,600	62.7	(56.8 - 68.6)
\$25,000-\$49,999	*	*	*	*	6,619	697,300	60.1	(54.6 - 65.7)
\$50,000-\$74,999	*	*	*	*	3,514	613,700	74.2	(68.9 - 79.6)
≥ \$75,000	*	*	*	*	4,665	1,294,400	73.3	(69.2 - 77.5)

¹ Defined as having ever had a sigmoidoscopy or colonoscopy among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have ever had a sigmoidoscopy or colonoscopy (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 37b. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	460	59.1	(53.7 - 64.4)	25,357	66.6	(63.9 - 69.3)
Gender						
Male	176	54.4	(46.0 - 62.8)	9,582	68.5	(64.5 - 72.5)
Female	284	63.9	(57.6 - 70.1)	15,775	64.8	(61.2 - 68.4)
Race/Ethnicity						
White Non-Hispanic	450	59.6	(54.3 - 64.8)	24,248	68.7	(66.5 - 70.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	223	55.5	(47.9 - 63.1)	11,121	61.3	(56.4 - 66.2)
Some College	*	*	*	6,287	63.8	(58.8 - 68.8)
≥ College Degree	116	72.4	(63.2 - 81.6)	7,884	74.0	(70.2 - 77.9)
Income						
≤ \$24,999	140	52.0	(42.4 - 61.6)	7,220	63.0	(56.7 - 69.3)
\$25,000-\$49,999	*	*	*	6,619	59.7	(53.8 - 65.6)
\$50,000-\$74,999	*	*	*	3,514	74.4	(68.3 - 80.5)
≥ \$75,000	*	*	*	4,665	73.6	(69.3 - 78.0)

¹ Defined as having ever had a sigmoidoscopy or colonoscopy among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 38a. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	459	6,800	58.1	(52.9 - 63.3)	25,169	3,855,800	64.6	(62.1 - 67.1)
Gender								
Male	176	3,100	53.5	(45.2 - 61.8)	9,521	1,835,400	66.4	(62.6 - 70.1)
Female	283	3,700	62.5	(56.3 - 68.7)	15,648	2,020,400	63.1	(59.8 - 66.4)
Age (years)								
50-54	*	*	*	*	4,169	700,500	49.3	(43.3 - 55.3)
55-64	164	2,300	57.0	(48.5 - 65.4)	8,652	1,455,100	69.4	(65.7 - 73.2)
≥ 65	213	3,200	66.3	(59.3 - 73.4)	12,331	1,695,700	69.5	(65.9 - 73.1)
Race/Ethnicity								
White Non-Hispanic	449	6,600	58.4	(53.3 - 63.6)	24,069	2,981,500	66.6	(64.6 - 68.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	223	3,000	53.7	(46.2 - 61.1)	11,017	1,360,400	59.0	(54.5 - 63.5)
Some College	*	*	*	*	6,251	807,100	62.4	(57.8 - 67.1)
≥ College Degree	116	2,200	71.7	(62.6 - 80.8)	7,841	1,682,500	71.7	(68.0 - 75.4)
Income								
≤ \$24,999	140	1,500	50.5	(41.0 - 59.9)	7,163	858,900	61.0	(55.1 - 67.0)
\$25,000-\$49,999	*	*	*	*	6,584	664,900	57.7	(52.2 - 63.2)
\$50,000-\$74,999	*	*	*	*	3,501	602,600	73.1	(67.7 - 78.5)
≥ \$75,000	*	*	*	*	4,655	1,260,400	71.5	(67.3 - 75.7)

¹ Defined as having had a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have had a sigmoidoscopy or colonoscopy within the past 10 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 38b. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	459	58.0	(52.6 - 63.3)	25,169	64.6	(61.9 - 67.3)
Gender						
Male	176	53.8	(45.4 - 62.2)	9,521	66.9	(62.9 - 70.9)
Female	283	62.3	(56.0 - 68.6)	15,648	62.4	(58.9 - 66.0)
Race/Ethnicity						
White Non-Hispanic	449	58.4	(53.2 - 63.7)	24,069	66.5	(64.3 - 68.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	223	54.0	(46.3 - 61.6)	11,017	59.0	(54.1 - 63.9)
Some College	*	*	*	6,251	61.9	(56.9 - 66.9)
≥ College Degree	116	72.0	(62.8 - 81.2)	7,841	72.1	(68.2 - 75.9)
Income						
≤ \$24,999	140	50.5	(40.9 - 60.1)	7,163	61.3	(54.9 - 67.6)
\$25,000-\$49,999	*	*	*	6,584	57.2	(51.4 - 63.0)
\$50,000-\$74,999	*	*	*	3,501	73.3	(67.2 - 79.4)
≥ \$75,000	*	*	*	4,655	71.8	(67.4 - 76.2)

¹ Defined as having had a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 39a. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	454	7,300	62.7	(57.5 - 67.9)	24,792	4,049,200	69.1	(66.7 - 71.6)
Gender								
Male	175	3,300	56.9	(48.6 - 65.1)	9,418	1,945,200	70.7	(67.0 - 74.3)
Female	279	4,000	68.4	(62.4 - 74.4)	15,374	2,104,100	67.8	(64.6 - 71.1)
Age (years)								
50-54	*	*	*	*	4,132	751,200	54.1	(48.0 - 60.1)
55-64	160	2,400	61.3	(52.9 - 69.8)	8,366	1,487,600	73.7	(70.0 - 77.4)
≥ 65	212	3,400	71.4	(64.6 - 78.1)	12,294	1,810,500	74.0	(70.6 - 77.4)
Race/Ethnicity								
White Non-Hispanic	445	7,100	63.2	(58.1 - 68.3)	23,726	3,091,400	70.4	(68.4 - 72.4)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	218	3,300	59.8	(52.3 - 67.2)	10,872	1,461,600	64.2	(59.8 - 68.7)
Some College	*	*	*	*	6,170	873,300	68.4	(63.9 - 72.8)
≥ College Degree	116	2,200	72.6	(63.5 - 81.6)	7,708	1,709,000	74.6	(70.9 - 78.3)
Income								
≤ \$24,999	138	1,700	57.4	(47.9 - 67.0)	7,092	923,300	66.0	(60.1 - 71.9)
\$25,000-\$49,999	*	*	*	*	6,526	730,000	64.2	(58.9 - 69.6)
\$50,000-\$74,999	*	*	*	*	3,467	621,500	76.6	(71.4 - 81.9)
≥ \$75,000	*	*	*	*	4,615	1,292,500	73.8	(69.8 - 77.9)

¹ Defined as having used a home blood stool test within the past year or a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 years and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have used a home blood stool test in the past year or sigmoidoscopy or colonoscopy within the past 10 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 39b. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	454	62.6	(57.3 - 67.9)	24,792	69.3	(66.7 - 71.9)
Gender						
Male	175	57.1	(48.6 - 65.5)	9,418	71.5	(67.6 - 75.4)
Female	279	68.3	(62.3 - 74.4)	15,374	67.2	(63.7 - 70.7)
Race/Ethnicity						
White Non-Hispanic	445	63.2	(58.1 - 68.4)	23,726	70.3	(68.1 - 72.4)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	218	60.1	(52.5 - 67.7)	10,872	64.5	(59.8 - 69.3)
Some College	*	*	*	6,170	68.2	(63.4 - 72.9)
≥ College Degree	116	72.8	(63.7 - 82.0)	7,708	75.0	(71.2 - 78.8)
Income						
≤ \$24,999	138	57.6	(47.9 - 67.3)	7,092	66.4	(60.2 - 72.6)
\$25,000-\$49,999	*	*	*	6,526	63.8	(58.2 - 69.5)
\$50,000-\$74,999	*	*	*	3,467	76.7	(70.6 - 82.7)
≥ \$75,000	*	*	*	4,615	74.4	(70.2 - 78.6)

¹ Defined as having used a home blood stool test within the past year or a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 years and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 40a. High blood pressure among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	646	7,900	30.9	(26.3 - 35.5)	38,125	3,974,800	26.8	(24.9 - 28.7)
Gender								
Male	245	4,000	31.8	(24.3 - 39.2)	14,611	1,960,500	27.7	(24.7 - 30.7)
Female	401	3,900	30.0	(24.6 - 35.5)	23,514	2,014,300	26.0	(23.6 - 28.3)
Age (years)								
18-34	*	*	*	*	3,935	312,200	7.0	(4.2 - 9.8)
35-44	74	900	15.2	(6.2 - 24.3)	4,984	426,200	14.8	(11.6 - 18.1)
45-54	142	1,200	25.6	(17.0 - 34.1)	7,825	862,400	30.3	(25.9 - 34.8)
55-64	163	2,000	50.4	(41.9 - 58.9)	8,765	890,800	41.6	(37.6 - 45.7)
≥ 65	218	2,900	60.1	(52.9 - 67.4)	12,616	1,483,200	58.5	(54.7 - 62.3)
Race/Ethnicity								
White Non-Hispanic	631	7,600	30.6	(26.0 - 35.2)	35,814	2,650,900	28.4	(26.9 - 30.0)
Black Non-Hispanic	*	*	*	*	736	558,500	26.2	(19.6 - 32.8)
Hispanic	*	*	*	*	863	538,100	23.8	(17.0 - 30.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	303	3,600	29.9	(23.3 - 36.4)	15,688	1,851,200	33.1	(29.4 - 36.9)
Some College	175	2,500	32.6	(23.4 - 41.8)	9,906	851,100	24.1	(20.9 - 27.3)
≥ College Degree	167	1,800	30.9	(22.0 - 39.8)	12,452	1,253,400	22.3	(19.7 - 24.9)
Income								
≤ \$24,999	*	*	*	*	9,640	1,153,000	32.8	(27.6 - 38.1)
\$25,000-\$49,999	201	2,600	29.9	(22.1 - 37.7)	9,657	807,800	27.9	(23.7 - 32.1)
\$50,000-\$74,999	*	*	*	*	5,773	507,600	25.8	(21.4 - 30.1)
≥ \$75,000	119	1,300	23.3	(14.7 - 32.0)	8,560	942,500	21.0	(18.4 - 23.6)

¹ Defined as having ever been told by a health professional that respondent has high blood pressure

² Sample size from survey

³ Estimated number of adults who have ever been told by a health professional that they have high blood pressure (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 40b. High blood pressure among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	646	28.7	(24.0 - 33.4)	38,125	25.7	(23.8 - 27.5)
Gender						
Male	245	30.4	(22.6 - 38.1)	14,611	27.7	(24.7 - 30.7)
Female	401	27.0	(21.6 - 32.4)	23,514	23.8	(21.5 - 26.0)
Race/Ethnicity						
White Non-Hispanic	631	28.5	(23.7 - 33.2)	35,814	27.2	(25.6 - 28.7)
Black Non-Hispanic	*	*	*	736	24.9	(18.4 - 31.3)
Hispanic	*	*	*	863	23.3	(16.6 - 30.0)
Other Non-Hispanic	*	*	*	712	19.3	(9.5 - 29.1)
Education						
≤ High School	303	27.7	(21.0 - 34.5)	15,688	32.1	(28.4 - 35.8)
Some College	175	29.6	(20.4 - 38.9)	9,906	22.8	(19.7 - 25.9)
≥ College Degree	167	29.6	(20.3 - 38.9)	12,452	21.2	(18.6 - 23.8)
Income						
≤ \$24,999	*	*	*	9,640	31.8	(26.7 - 37.0)
\$25,000-\$49,999	201	27.7	(19.8 - 35.5)	9,657	26.9	(22.8 - 30.9)
\$50,000-\$74,999	*	*	*	5,773	24.8	(20.4 - 29.2)
≥ \$75,000	119	22.3	(13.4 - 31.2)	8,560	19.9	(17.3 - 22.5)

¹ Defined as having ever been told by a health professional that respondent has high blood pressure

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 41a. Blood pressure medication among adults with high blood pressure¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	264	5,800	72.9	(64.5 - 81.4)	15,288	3,145,800	79.2	(76.0 - 82.4)
Gender								
Male	*	*	*	*	6,080	1,432,100	73.1	(67.6 - 78.5)
Female	165	3,200	81.9	(74.0 - 89.8)	9,208	1,713,700	85.2	(82.1 - 88.2)
Age (years)								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,315	645,500	74.9	(66.8 - 82.9)
55-64	81	1,700	82.5	(73.5 - 91.6)	4,045	782,300	87.8	(84.5 - 91.1)
≥ 65	130	2,700	91.3	(85.4 - 97.3)	7,675	1,383,400	93.4	(91.1 - 95.7)
Race/Ethnicity								
White Non-Hispanic	258	5,600	73.9	(65.5 - 82.3)	14,442	2,117,300	79.9	(77.5 - 82.4)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	7,354	1,482,800	80.2	(74.7 - 85.6)
Some College	*	*	*	*	3,868	632,000	74.3	(67.8 - 80.7)
≥ College Degree	*	*	*	*	4,035	1,013,200	80.9	(76.8 - 84.9)
Income								
≤ \$24,999	*	*	*	*	4,898	948,200	82.3	(75.7 - 88.9)
\$25,000-\$49,999	*	*	*	*	4,088	668,300	82.8	(77.6 - 88.0)
\$50,000-\$74,999	*	*	*	*	1,971	391,500	77.2	(67.7 - 86.6)
≥ \$75,000	*	*	*	*	2,397	683,700	72.5	(66.1 - 78.9)

¹ Defined as currently taking medicine for high blood pressure among adults with high blood pressure

² Sample size from survey

³ Estimated number of adults with high blood pressure who are taking medicine for high blood pressure (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 41b. Blood pressure medication among adults with high blood pressure¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	264	70.3	(60.8 - 79.7)	15,288	78.5	(75.2 - 81.7)
Gender						
Male	*	*	*	6,080	73.3	(67.9 - 78.7)
Female	165	78.8	(69.1 - 88.5)	9,208	84.0	(80.7 - 87.2)
Race/Ethnicity						
White Non-Hispanic	258	71.3	(61.8 - 80.8)	14,442	79.0	(76.5 - 81.6)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	7,354	79.7	(74.3 - 85.2)
Some College	*	*	*	3,868	72.5	(65.7 - 79.3)
≥ College Degree	*	*	*	4,035	80.4	(76.1 - 84.6)
Income						
≤ \$24,999	*	*	*	4,898	81.6	(75.0 - 88.3)
\$25,000-\$49,999	*	*	*	4,088	82.1	(76.7 - 87.5)
\$50,000-\$74,999	*	*	*	1,971	76.4	(66.7 - 86.1)
≥ \$75,000	*	*	*	2,397	71.6	(64.8 - 78.3)

¹ Defined as currently taking medicine for high blood pressure among adults with high blood pressure

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 42a. Ever had blood cholesterol checked among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	634	20,100	80.4	(75.0 - 85.8)	37,450	11,496,600	80.3	(77.7 - 83.0)
Gender								
Male	241	10,200	81.9	(73.8 - 90.0)	14,389	5,288,400	77.7	(73.3 - 82.0)
Female	393	9,800	78.9	(71.8 - 86.0)	23,061	6,208,200	82.7	(79.6 - 85.8)
Age (years)								
18-34	*	*	*	*	3,684	2,296,400	55.3	(48.7 - 61.9)
35-44	*	*	*	*	4,883	2,264,200	81.6	(76.6 - 86.5)
45-54	136	4,300	91.7	(87.3 - 96.0)	7,731	2,539,400	91.0	(87.0 - 94.9)
55-64	164	3,900	97.7	(95.5 - 99.9)	8,718	2,014,900	95.4	(93.5 - 97.2)
≥ 65	216	4,500	92.7	(88.8 - 96.7)	12,434	2,381,700	95.9	(94.3 - 97.6)
Race/Ethnicity								
White Non-Hispanic	620	19,600	80.5	(75.1 - 85.9)	35,198	7,832,600	86.3	(84.8 - 87.9)
Black Non-Hispanic	*	*	*	*	720	1,588,900	77.8	(68.4 - 87.2)
Hispanic	*	*	*	*	842	1,224,600	56.8	(46.8 - 66.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	295	9,600	81.6	(75.1 - 88.2)	15,342	3,795,700	70.6	(65.6 - 75.6)
Some College	*	*	*	*	9,732	2,599,800	77.4	(71.7 - 83.2)
≥ College Degree	164	5,300	91.1	(83.6 - 98.7)	12,296	5,008,300	91.3	(88.9 - 93.6)
Income								
≤ \$24,999	*	*	*	*	9,437	2,277,500	68.4	(61.5 - 75.3)
\$25,000-\$49,999	*	*	*	*	9,492	2,053,000	71.9	(64.5 - 79.3)
\$50,000-\$74,999	*	*	*	*	5,721	1,736,100	90.0	(87.7 - 92.3)
≥ \$75,000	*	*	*	*	8,460	4,019,300	92.2	(90.8 - 93.6)

¹ Defined as ever having blood cholesterol checked

² Sample size from survey

³ Estimated number of adults who have ever had blood cholesterol checked (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 42b. Ever had blood cholesterol checked among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	634	78.6	(72.7 - 84.4)	37,450	79.9	(77.3 - 82.5)
Gender						
Male	241	80.4	(71.6 - 89.2)	14,389	78.0	(73.9 - 82.1)
Female	393	76.7	(69.1 - 84.4)	23,061	81.6	(78.4 - 84.8)
Race/Ethnicity						
White Non-Hispanic	620	78.7	(72.8 - 84.6)	35,198	85.6	(84.0 - 87.2)
Black Non-Hispanic	*	*	*	720	77.5	(68.2 - 86.7)
Hispanic	*	*	*	842	57.2	(47.4 - 67.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	295	79.9	(72.7 - 87.2)	15,342	70.2	(65.2 - 75.1)
Some College	*	*	*	9,732	76.7	(71.1 - 82.3)
≥ College Degree	164	89.9	(81.4 - 98.4)	12,296	90.9	(88.6 - 93.2)
Income						
≤ \$24,999	*	*	*	9,437	68.1	(61.3 - 74.9)
\$25,000-\$49,999	*	*	*	9,492	71.6	(64.4 - 78.7)
\$50,000-\$74,999	*	*	*	5,721	89.6	(87.2 - 92.0)
≥ \$75,000	*	*	*	8,460	91.7	(90.1 - 93.2)

¹ Defined as ever having blood cholesterol checked

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 43a. Blood cholesterol checked within the past 5 years among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	630	18,800	76.3	(70.8 - 81.7)	37,201	11,035,200	77.8	(75.1 - 80.4)
Gender								
Male	238	9,200	75.1	(66.8 - 83.4)	14,307	5,020,700	74.5	(70.2 - 78.9)
Female	392	9,600	77.4	(70.4 - 84.5)	22,894	6,014,600	80.7	(77.6 - 83.8)
Age (years)								
18-34	*	*	*	*	3,647	2,179,800	53.3	(46.7 - 59.9)
35-44	*	*	*	*	4,862	2,161,200	78.0	(72.8 - 83.2)
45-54	135	3,700	80.5	(73.2 - 87.9)	7,697	2,408,600	86.6	(82.4 - 90.8)
55-64	164	3,600	90.3	(85.5 - 95.2)	8,677	1,957,900	93.3	(91.4 - 95.3)
≥ 65	214	4,400	90.9	(86.6 - 95.2)	12,318	2,327,800	94.8	(93.1 - 96.6)
Race/Ethnicity								
White Non-Hispanic	616	18,300	76.5	(71.0 - 82.0)	34,960	7,486,600	83.4	(81.8 - 85.0)
Black Non-Hispanic	*	*	*	*	719	1,561,700	76.5	(67.1 - 85.9)
Hispanic	*	*	*	*	838	1,184,000	55.6	(45.6 - 65.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	293	8,800	76.9	(70.1 - 83.8)	15,215	3,612,500	68.3	(63.2 - 73.3)
Some College	*	*	*	*	9,673	2,524,200	75.4	(69.8 - 81.1)
≥ College Degree	164	5,000	86.1	(78.0 - 94.2)	12,236	4,806,500	88.1	(85.5 - 90.6)
Income								
≤ \$24,999	*	*	*	*	9,350	2,214,800	66.9	(60.1 - 73.8)
\$25,000-\$49,999	*	*	*	*	9,454	1,936,600	68.8	(61.5 - 76.1)
\$50,000-\$74,999	*	*	*	*	5,701	1,640,400	85.9	(82.4 - 89.5)
≥ \$75,000	*	*	*	*	8,435	3,872,400	89.2	(87.5 - 90.9)

¹ Defined as having had blood cholesterol checked within the past 5 years

² Sample size from survey

³ Estimated number of adults who have had blood cholesterol checked within the past 5 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 43b. Blood cholesterol checked within the past 5 years among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	630	74.7	(68.8 - 80.6)	37,201	77.3	(74.7 - 79.9)
Gender						
Male	238	74.1	(65.2 - 83.1)	14,307	74.8	(70.7 - 79.0)
Female	392	75.2	(67.6 - 82.9)	22,894	79.5	(76.3 - 82.7)
Race/Ethnicity						
White Non-Hispanic	616	74.9	(69.0 - 80.9)	34,960	82.6	(81.0 - 84.3)
Black Non-Hispanic	*	*	*	719	76.2	(66.9 - 85.5)
Hispanic	*	*	*	838	55.9	(46.1 - 65.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	293	75.5	(68.0 - 83.0)	15,215	67.8	(62.9 - 72.8)
Some College	*	*	*	9,673	74.7	(69.1 - 80.2)
≥ College Degree	164	85.3	(76.4 - 94.2)	12,236	87.5	(85.0 - 90.1)
Income						
≤ \$24,999	*	*	*	9,350	66.6	(59.8 - 73.4)
\$25,000-\$49,999	*	*	*	9,454	68.5	(61.4 - 75.6)
\$50,000-\$74,999	*	*	*	5,701	85.4	(81.7 - 89.0)
≥ \$75,000	*	*	*	8,435	88.5	(86.7 - 90.3)

¹ Defined as having had blood cholesterol checked within the past 5 years

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 44a. Cardiovascular disease among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	642	2,400	9.5	(6.9 - 12.0)	37,810	1,135,000	7.7	(6.5 - 8.8)
Gender								
Male	243	1,500	12.1	(7.5 - 16.7)	14,477	664,600	9.4	(7.5 - 11.4)
Female	399	900	6.9	(4.6 - 9.2)	23,333	470,400	6.1	(4.8 - 7.4)
Age (years)								
18-34	*	*	*	*	3,930	100,800	2.3	(0.0 - 4.8)
35-44	*	*	*	*	4,971	103,400	3.6	(1.5 - 5.7)
45-54	*	*	*	*	7,768	131,300	4.6	(2.9 - 6.4)
55-64	164	400	10.4	(5.5 - 15.4)	8,699	224,200	10.6	(8.3 - 12.9)
≥ 65	216	1,400	28.8	(22.0 - 35.5)	12,442	575,300	23.0	(19.9 - 26.1)
Race/Ethnicity								
White Non-Hispanic	628	2,400	9.6	(7.0 - 12.2)	35,522	772,200	8.3	(7.4 - 9.2)
Black Non-Hispanic	*	*	*	*	728	140,700	6.6	(3.3 - 9.9)
Hispanic	*	*	*	*	854	163,300	7.3	(2.0 - 12.5)
Other Non-Hispanic	*	*	*	*	706	58,800	5.2	(0.9 - 9.5)
Education								
≤ High School	302	1,100	8.9	(5.5 - 12.2)	15,486	590,800	10.7	(8.2 - 13.2)
Some College	173	600	8.2	(4.6 - 11.8)	9,849	227,100	6.4	(4.7 - 8.1)
≥ College Degree	166	700	12.3	(5.2 - 19.4)	12,393	314,600	5.6	(4.3 - 6.9)
Income								
≤ \$24,999	168	1,100	20.3	(12.1 - 28.6)	9,505	448,100	12.9	(9.1 - 16.7)
\$25,000-\$49,999	200	500	5.7	(2.8 - 8.6)	9,571	179,300	6.2	(4.8 - 7.6)
\$50,000-\$74,999	*	*	*	*	5,751	166,900	8.5	(5.5 - 11.4)
≥ \$75,000	118	300	5.9	(2.0 - 9.9)	8,538	183,500	4.1	(2.9 - 5.3)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina or a stroke

² Sample size from survey

³ Estimated number of adults with cardiovascular disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 44b. Cardiovascular disease among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	642	8.8	(6.2 - 11.4)	37,810	7.6	(6.4 - 8.7)
Gender						
Male	243	11.7	(7.0 - 16.5)	14,477	9.7	(7.8 - 11.7)
Female	399	5.9	(3.8 - 8.0)	23,333	5.6	(4.3 - 6.9)
Race/Ethnicity						
White Non-Hispanic	628	8.9	(6.3 - 11.6)	35,522	8.2	(7.2 - 9.1)
Black Non-Hispanic	*	*	*	728	6.8	(3.2 - 10.4)
Hispanic	*	*	*	854	7.0	(2.0 - 12.0)
Other Non-Hispanic	*	*	*	706	5.5	(0.9 - 10.0)
Education						
≤ High School	302	8.4	(5.0 - 11.7)	15,486	10.5	(8.0 - 13.0)
Some College	173	7.0	(3.9 - 10.2)	9,849	6.4	(4.5 - 8.2)
≥ College Degree	166	11.9	(4.4 - 19.5)	12,393	5.5	(4.2 - 6.9)
Income						
≤ \$24,999	168	19.4	(10.6 - 28.1)	9,505	12.8	(9.0 - 16.6)
\$25,000-\$49,999	200	5.0	(2.4 - 7.6)	9,571	6.0	(4.7 - 7.4)
\$50,000-\$74,999	*	*	*	5,751	8.5	(5.4 - 11.6)
≥ \$75,000	118	5.8	(1.8 - 9.7)	8,538	4.0	(2.7 - 5.2)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina or a stroke

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 45a. Coronary heart disease among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	641	2,000	8.0	(5.6 - 10.4)	37,838	929,400	6.3	(5.3 - 7.3)
Gender								
Male	242	1,300	10.6	(6.2 - 15.0)	14,479	598,600	8.5	(6.6 - 10.4)
Female	399	700	5.5	(3.5 - 7.5)	23,359	330,800	4.3	(3.4 - 5.2)
Age (years)								
18-34	*	*	*	*	3,932	63,400	1.4	(0.0 - 3.6)
35-44	*	*	*	*	4,973	95,300	3.3	(1.2 - 5.4)
45-54	*	*	*	*	7,778	108,300	3.8	(2.1 - 5.6)
55-64	164	400	9.4	(4.7 - 14.2)	8,703	188,600	8.9	(6.7 - 11.1)
≥ 65	215	1,100	23.4	(17.1 - 29.8)	12,452	473,800	18.9	(16.1 - 21.7)
Race/Ethnicity								
White Non-Hispanic	627	2,000	8.2	(5.7 - 10.6)	35,552	636,700	6.9	(6.0 - 7.7)
Black Non-Hispanic	*	*	*	*	728	113,100	5.3	(2.3 - 8.4)
Hispanic	*	*	*	*	853	128,000	5.7	(1.1 - 10.3)
Other Non-Hispanic	*	*	*	*	705	51,500	4.6	(0.5 - 8.7)
Education								
≤ High School	301	900	7.4	(4.3 - 10.5)	15,485	474,500	8.6	(6.3 - 10.8)
Some College	173	500	6.5	(3.3 - 9.6)	9,860	193,000	5.5	(3.9 - 7.1)
≥ College Degree	166	700	11.3	(4.4 - 18.2)	12,411	259,700	4.6	(3.4 - 5.8)
Income								
≤ \$24,999	167	1,000	18.4	(10.3 - 26.4)	9,507	372,300	10.7	(7.3 - 14.1)
\$25,000-\$49,999	200	400	4.4	(1.9 - 6.9)	9,585	136,300	4.7	(3.8 - 5.7)
\$50,000-\$74,999	*	*	*	*	5,755	154,600	7.8	(4.9 - 10.7)
≥ \$75,000	*	*	*	*	8,541	151,800	3.4	(2.3 - 4.5)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina

² Sample size from survey

³ Estimated number of adults with coronary heart disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 45b. Coronary heart disease among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	641	7.5	(5.0 - 10.0)	37,838	6.3	(5.2 - 7.3)
Gender						
Male	242	10.4	(5.8 - 15.0)	14,479	8.8	(6.9 - 10.8)
Female	399	4.7	(2.9 - 6.5)	23,359	3.9	(3.0 - 4.8)
Race/Ethnicity						
White Non-Hispanic	627	7.6	(5.1 - 10.1)	35,552	6.8	(5.9 - 7.7)
Black Non-Hispanic	*	*	*	728	5.6	(2.2 - 9.0)
Hispanic	*	*	*	853	5.4	(1.1 - 9.6)
Other Non-Hispanic	*	*	*	705	4.8	(0.4 - 9.2)
Education						
≤ High School	301	7.1	(3.9 - 10.3)	15,485	8.5	(6.2 - 10.7)
Some College	173	5.5	(2.7 - 8.2)	9,860	5.5	(3.7 - 7.2)
≥ College Degree	166	11.0	(3.6 - 18.4)	12,411	4.7	(3.4 - 5.9)
Income						
≤ \$24,999	167	17.4	(8.9 - 26.0)	9,507	10.7	(7.2 - 14.1)
\$25,000-\$49,999	200	3.9	(1.6 - 6.2)	9,585	4.7	(3.6 - 5.7)
\$50,000-\$74,999	*	*	*	5,755	8.0	(4.8 - 11.1)
≥ \$75,000	*	*	*	8,541	3.3	(2.1 - 4.5)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 46a. Ever diagnosed with diabetes among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	648	2,200	8.4	(6.2 - 10.7)	38,174	1,382,700	9.3	(8.0 - 10.5)
Gender								
Male	245	1,100	9.0	(5.3 - 12.6)	14,626	530,400	7.5	(6.2 - 8.8)
Female	403	1,000	7.9	(5.3 - 10.5)	23,548	852,300	11.0	(8.9 - 13.0)
Age (years)								
18-34	*	*	*	*	3,941	150,200	3.4	(1.0 - 5.8)
35-44	*	*	*	*	4,987	165,100	5.7	(3.6 - 7.9)
45-54	142	400	8.0	(3.4 - 12.7)	7,830	279,000	9.8	(6.7 - 12.8)
55-64	164	600	13.8	(7.8 - 19.8)	8,771	332,900	15.5	(12.3 - 18.8)
≥ 65	219	1,100	21.8	(15.2 - 28.3)	12,645	455,600	17.9	(15.0 - 20.8)
Race/Ethnicity								
White Non-Hispanic	633	2,100	8.6	(6.3 - 10.9)	35,862	809,300	8.7	(7.8 - 9.5)
Black Non-Hispanic	*	*	*	*	739	261,000	12.1	(6.6 - 17.6)
Hispanic	*	*	*	*	862	227,300	10.1	(5.4 - 14.8)
Other Non-Hispanic	*	*	*	*	711	85,100	7.6	(2.3 - 12.8)
Education								
≤ High School	305	1,000	8.5	(5.0 - 11.9)	15,709	730,000	13.0	(10.5 - 15.6)
Some College	175	600	7.6	(4.0 - 11.1)	9,918	303,500	8.6	(5.9 - 11.2)
≥ College Degree	167	600	9.5	(4.9 - 14.1)	12,464	337,500	6.0	(4.8 - 7.2)
Income								
≤ \$24,999	172	800	14.6	(8.0 - 21.3)	9,652	543,400	15.5	(11.4 - 19.5)
\$25,000-\$49,999	201	500	6.1	(2.9 - 9.3)	9,667	291,000	10.0	(7.5 - 12.6)
\$50,000-\$74,999	*	*	*	*	5,777	151,000	7.6	(5.0 - 10.3)
≥ \$75,000	119	300	5.8	(1.9 - 9.7)	8,565	220,800	4.9	(3.8 - 6.1)

¹ Defined as having ever been told by a doctor that respondent had diabetes, excluding pre-diabetes and women with diabetes only when pregnant

² Sample size from survey

³ Estimated number of adults who have diabetes (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 46b. Ever diagnosed with diabetes among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	648	7.6	(5.5 - 9.7)	38,174	9.0	(7.8 - 10.3)
Gender						
Male	245	8.4	(4.8 - 12.0)	14,626	7.5	(6.1 - 8.9)
Female	403	6.8	(4.5 - 9.2)	23,548	10.4	(8.4 - 12.4)
Race/Ethnicity						
White Non-Hispanic	633	7.8	(5.6 - 10.0)	35,862	8.4	(7.5 - 9.2)
Black Non-Hispanic	*	*	*	739	11.4	(6.0 - 16.7)
Hispanic	*	*	*	862	10.1	(5.4 - 14.9)
Other Non-Hispanic	*	*	*	711	7.7	(2.3 - 13.0)
Education						
≤ High School	305	7.5	(4.2 - 10.8)	15,709	12.6	(10.0 - 15.2)
Some College	175	6.7	(3.4 - 10.0)	9,918	8.3	(5.6 - 10.9)
≥ College Degree	167	9.1	(4.6 - 13.6)	12,464	5.9	(4.7 - 7.2)
Income						
≤ \$24,999	172	13.6	(6.8 - 20.4)	9,652	15.2	(11.1 - 19.2)
\$25,000-\$49,999	201	5.6	(2.5 - 8.8)	9,667	9.7	(7.1 - 12.2)
\$50,000-\$74,999	*	*	*	5,777	7.6	(4.9 - 10.2)
≥ \$75,000	119	5.4	(1.6 - 9.1)	8,565	4.7	(3.6 - 5.9)

¹ Defined as having ever been told by a doctor that respondent had diabetes, excluding pre-diabetes and women with diabetes only when pregnant

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 47a. Ever diagnosed with asthma (lifetime) among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	642	2,700	10.8	(7.7 - 13.9)	38,096	2,225,100	15.0	(13.2 - 16.8)
Gender								
Male	244	1,100	8.4	(3.8 - 13.1)	14,601	875,800	12.4	(10.1 - 14.7)
Female	398	1,700	13.2	(9.0 - 17.3)	23,495	1,349,300	17.4	(14.6 - 20.1)
Age (years)								
18-34	*	*	*	*	3,931	825,100	18.5	(13.8 - 23.3)
35-44	74	600	11.1	(3.5 - 18.6)	4,972	431,700	15.1	(11.2 - 19.1)
45-54	142	400	9.1	(4.5 - 13.8)	7,819	365,900	12.8	(9.7 - 16.0)
55-64	162	600	14.5	(8.5 - 20.4)	8,756	316,700	14.8	(11.9 - 17.7)
≥ 65	217	500	10.3	(5.9 - 14.7)	12,618	285,700	11.2	(8.6 - 13.8)
Race/Ethnicity								
White Non-Hispanic	627	2,700	10.9	(7.7 - 14.1)	35,783	1,365,700	14.6	(13.0 - 16.3)
Black Non-Hispanic	*	*	*	*	737	367,000	17.2	(10.2 - 24.2)
Hispanic	*	*	*	*	863	317,100	14.1	(9.0 - 19.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	300	1,600	13.7	(8.2 - 19.3)	15,675	831,600	14.9	(12.0 - 17.8)
Some College	174	500	7.0	(2.7 - 11.3)	9,893	587,500	16.6	(13.1 - 20.1)
≥ College Degree	167	600	9.7	(5.1 - 14.3)	12,444	739,000	13.2	(10.4 - 15.9)
Income								
≤ \$24,999	171	600	10.8	(5.9 - 15.8)	9,643	708,900	20.2	(14.9 - 25.5)
\$25,000-\$49,999	199	800	9.0	(3.8 - 14.3)	9,641	389,800	13.4	(10.0 - 16.9)
\$50,000-\$74,999	76	300	9.9	(2.9 - 17.0)	5,762	257,800	13.2	(10.0 - 16.5)
≥ \$75,000	*	*	*	*	8,562	577,900	12.8	(10.4 - 15.3)

¹ Defined as adults having ever been told by a health professional that they had asthma

² Sample size from survey

³ Estimated number of adults who have lifetime asthma (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 47b. Ever diagnosed with asthma (lifetime) among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	642	10.8	(7.5 - 14.2)	38,096	15.2	(13.3 - 17.0)
Gender						
Male	244	8.3	(3.4 - 13.3)	14,601	12.6	(10.3 - 14.9)
Female	398	13.3	(8.8 - 17.9)	23,495	17.5	(14.8 - 20.3)
Race/Ethnicity						
White Non-Hispanic	627	10.9	(7.4 - 14.3)	35,783	14.9	(13.2 - 16.6)
Black Non-Hispanic	*	*	*	737	17.1	(10.1 - 24.1)
Hispanic	*	*	*	863	14.5	(9.3 - 19.7)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	300	14.3	(8.2 - 20.3)	15,675	15.1	(12.1 - 18.0)
Some College	174	6.7	(2.2 - 11.1)	9,893	16.9	(13.4 - 20.4)
≥ College Degree	167	9.3	(4.6 - 14.0)	12,444	13.3	(10.6 - 16.0)
Income						
≤ \$24,999	171	9.8	(5.2 - 14.4)	9,643	20.3	(15.0 - 25.6)
\$25,000-\$49,999	199	9.5	(3.7 - 15.3)	9,641	13.7	(10.2 - 17.2)
\$50,000-\$74,999	76	9.9	(2.3 - 17.6)	5,762	13.5	(10.0 - 17.0)
≥ \$75,000	*	*	*	8,562	13.0	(10.6 - 15.4)

¹ Defined as adults having ever been told by a health professional that they had asthma

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 48a. Current asthma among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	639	2,100	8.2	(5.4 - 11.1)	37,933	1,415,000	9.6	(8.1 - 11.0)
Gender								
Male	242	700	5.9	(1.7 - 10.1)	14,546	440,700	6.3	(4.8 - 7.7)
Female	397	1,300	10.5	(6.7 - 14.4)	23,387	974,300	12.6	(10.2 - 15.0)
Age (years)								
18-34	*	*	*	*	3,909	467,200	10.5	(6.8 - 14.3)
35-44	74	600	11.1	(3.5 - 18.6)	4,947	246,800	8.7	(5.8 - 11.6)
45-54	*	*	*	*	7,787	240,000	8.4	(6.1 - 10.8)
55-64	161	500	12.8	(7.0 - 18.6)	8,726	231,100	10.8	(8.1 - 13.5)
≥ 65	216	300	6.0	(2.6 - 9.4)	12,564	229,900	9.1	(6.5 - 11.6)
Race/Ethnicity								
White Non-Hispanic	624	2,000	8.3	(5.4 - 11.2)	35,629	888,400	9.6	(8.2 - 11.0)
Black Non-Hispanic	*	*	*	*	736	246,900	11.6	(5.7 - 17.5)
Hispanic	*	*	*	*	859	186,800	8.3	(4.4 - 12.2)
Other Non-Hispanic	*	*	*	*	709	93,000	8.3	(1.4 - 15.2)
Education								
≤ High School	298	1,300	10.6	(5.5 - 15.8)	15,594	595,100	10.7	(8.2 - 13.2)
Some College	173	400	5.6	(1.6 - 9.7)	9,861	322,000	9.1	(7.0 - 11.3)
≥ College Degree	167	400	6.7	(3.0 - 10.4)	12,394	430,900	7.7	(5.8 - 9.7)
Income								
≤ \$24,999	170	400	6.9	(2.9 - 10.8)	9,600	489,200	14.0	(9.7 - 18.3)
\$25,000-\$49,999	199	700	8.3	(3.1 - 13.4)	9,590	268,400	9.3	(6.2 - 12.4)
\$50,000-\$74,999	*	*	*	*	5,741	143,400	7.4	(5.1 - 9.6)
≥ \$75,000	*	*	*	*	8,541	339,200	7.6	(5.8 - 9.4)

¹ Defined as adults having ever been told by a health professional that they had asthma and still have asthma

² Sample size from survey

³ Estimated number of adults who currently have asthma (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 48b. Current asthma among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	639	8.4	(5.3 - 11.5)	37,933	9.7	(8.2 - 11.2)
Gender						
Male	242	5.9	(1.4 - 10.3)	14,546	6.4	(4.9 - 7.9)
Female	397	10.9	(6.6 - 15.2)	23,387	12.7	(10.2 - 15.1)
Race/Ethnicity						
White Non-Hispanic	624	8.4	(5.2 - 11.5)	35,629	9.7	(8.3 - 11.2)
Black Non-Hispanic	*	*	*	736	11.5	(5.6 - 17.3)
Hispanic	*	*	*	859	8.6	(4.5 - 12.6)
Other Non-Hispanic	*	*	*	709	8.3	(1.3 - 15.3)
Education						
≤ High School	298	11.3	(5.6 - 17.0)	15,594	10.8	(8.3 - 13.3)
Some College	173	5.3	(1.2 - 9.5)	9,861	9.4	(7.2 - 11.6)
≥ College Degree	167	6.3	(2.6 - 10.1)	12,394	7.8	(5.8 - 9.7)
Income						
≤ \$24,999	170	6.0	(2.5 - 9.5)	9,600	14.1	(9.7 - 18.4)
\$25,000-\$49,999	199	8.8	(3.1 - 14.6)	9,590	9.5	(6.3 - 12.7)
\$50,000-\$74,999	*	*	*	5,741	7.5	(5.0 - 10.1)
≥ \$75,000	*	*	*	8,541	7.7	(5.9 - 9.5)

¹ Defined as adults having ever been told by a health professional that they had asthma and still have asthma

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 49a. Chronic joint symptoms among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	646	13,300	52.0	(46.4 - 57.5)	38,099	6,377,100	42.9	(40.5 - 45.3)
Gender								
Male	245	6,800	53.5	(44.9 - 62.1)	14,595	3,033,600	42.8	(39.0 - 46.6)
Female	401	6,500	50.4	(43.5 - 57.4)	23,504	3,343,500	43.0	(40.0 - 46.0)
Age (years)								
18-34	*	*	*	*	3,933	1,261,300	28.3	(22.6 - 34.0)
35-44	*	*	*	*	4,984	1,110,900	38.6	(33.1 - 44.0)
45-54	141	2,300	48.6	(39.3 - 57.9)	7,817	1,456,500	51.2	(46.4 - 55.9)
55-64	164	2,400	60.5	(52.1 - 68.8)	8,751	1,167,400	54.6	(50.4 - 58.8)
≥ 65	218	3,000	61.3	(54.0 - 68.5)	12,614	1,381,000	54.3	(50.5 - 58.1)
Race/Ethnicity								
White Non-Hispanic	631	12,700	51.0	(45.4 - 56.6)	35,785	4,267,600	45.8	(43.8 - 47.7)
Black Non-Hispanic	*	*	*	*	740	890,500	41.3	(32.5 - 50.0)
Hispanic	*	*	*	*	860	827,500	36.7	(27.7 - 45.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	305	6,000	49.2	(41.3 - 57.1)	15,676	2,450,000	43.8	(39.4 - 48.1)
Some College	*	*	*	*	9,900	1,598,300	45.1	(40.1 - 50.1)
≥ College Degree	*	*	*	*	12,440	2,313,100	41.1	(37.8 - 44.5)
Income								
≤ \$24,999	*	*	*	*	9,641	1,483,000	42.2	(36.3 - 48.1)
\$25,000-\$49,999	200	4,900	56.7	(46.9 - 66.5)	9,647	1,332,100	45.9	(39.8 - 52.1)
\$50,000-\$74,999	*	*	*	*	5,768	894,200	45.3	(39.5 - 51.1)
≥ \$75,000	*	*	*	*	8,549	1,874,200	41.7	(38.4 - 45.1)

¹ Defined as adults having symptoms of pain, aching or stiffness in or around a joint during the past 30 days and began more than 3 months ago

² Sample size from survey

³ Estimated number of adults who have chronic joint symptoms (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 49b. Chronic joint symptoms among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	646	51.1	(45.2 - 57.0)	38,099	42.1	(39.7 - 44.5)
Gender						
Male	245	53.5	(44.3 - 62.7)	14,595	42.6	(38.8 - 46.3)
Female	401	48.6	(41.3 - 56.0)	23,504	41.6	(38.6 - 44.7)
Race/Ethnicity						
White Non-Hispanic	631	50.1	(44.1 - 56.1)	35,785	44.7	(42.7 - 46.6)
Black Non-Hispanic	*	*	*	740	40.2	(31.5 - 48.9)
Hispanic	*	*	*	860	36.6	(27.8 - 45.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	305	48.1	(39.7 - 56.5)	15,676	43.1	(38.8 - 47.5)
Some College	*	*	*	9,900	43.9	(39.0 - 48.8)
≥ College Degree	*	*	*	12,440	40.3	(37.0 - 43.7)
Income						
≤ \$24,999	*	*	*	9,641	41.6	(35.7 - 47.4)
\$25,000-\$49,999	*	*	*	9,647	45.0	(39.0 - 51.0)
\$50,000-\$74,999	*	*	*	5,768	44.3	(38.4 - 50.2)
≥ \$75,000	*	*	*	8,549	40.8	(37.4 - 44.2)

¹ Defined as adults having symptoms of pain, aching or stiffness in or around a joint during the past 30 days and began more than 3 months ago

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 50a. Arthritis among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	644	9,200	35.9	(31.1 - 40.8)	38,013	4,102,800	27.7	(25.9 - 29.6)
Gender								
Male	244	4,200	33.4	(25.8 - 41.0)	14,555	1,642,700	23.3	(20.7 - 25.8)
Female	400	4,900	38.5	(32.3 - 44.6)	23,458	2,460,100	31.8	(29.2 - 34.4)
Age (years)								
18-34	*	*	*	*	3,927	312,100	7.0	(4.9 - 9.1)
35-44	*	*	*	*	4,972	412,500	14.5	(11.3 - 17.6)
45-54	142	1,300	27.5	(19.4 - 35.7)	7,789	869,100	30.7	(26.6 - 34.8)
55-64	161	2,300	58.5	(50.1 - 67.0)	8,729	999,300	47.0	(42.9 - 51.1)
≥ 65	219	2,900	59.9	(52.6 - 67.2)	12,596	1,509,800	59.6	(55.8 - 63.3)
Race/Ethnicity								
White Non-Hispanic	629	8,900	36.1	(31.2 - 41.1)	35,710	2,967,700	31.9	(30.3 - 33.6)
Black Non-Hispanic	*	*	*	*	737	588,100	27.3	(20.2 - 34.4)
Hispanic	*	*	*	*	856	380,700	17.1	(11.8 - 22.5)
Other Non-Hispanic	*	*	*	*	710	166,300	14.8	(7.8 - 21.8)
Education								
≤ High School	302	4,700	38.8	(31.4 - 46.2)	15,652	1,742,700	31.3	(27.8 - 34.8)
Some College	175	2,400	31.6	(22.6 - 40.6)	9,874	902,000	25.5	(22.3 - 28.6)
≥ College Degree	166	2,100	35.4	(26.6 - 44.2)	12,407	1,437,800	25.6	(22.9 - 28.4)
Income								
≤ \$24,999	*	*	*	*	9,618	1,112,500	32.0	(27.1 - 36.9)
\$25,000-\$49,999	199	2,700	31.7	(23.7 - 39.7)	9,624	821,900	28.4	(24.2 - 32.6)
\$50,000-\$74,999	*	*	*	*	5,753	542,500	27.5	(23.2 - 31.9)
≥ \$75,000	*	*	*	*	8,536	1,012,800	22.6	(19.8 - 25.3)

¹ Defined as adults having ever been told by a health professional that they have some form of arthritis, rheumatoid arthritis, gout, lupus or fibromyalgia (excluding back or neck)

² Sample size from survey

³ Estimated number of adults who have ever been told by a health professional they have some form of arthritis (excluding back or neck) (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 50b. Arthritis among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	644	33.9	(28.8 - 38.9)	38,013	26.6	(24.8 - 28.5)
Gender						
Male	244	32.4	(24.4 - 40.3)	14,555	23.5	(20.9 - 26.1)
Female	400	35.3	(29.0 - 41.6)	23,458	29.5	(27.0 - 32.1)
Race/Ethnicity						
White Non-Hispanic	629	34.0	(28.8 - 39.2)	35,710	30.4	(28.7 - 32.0)
Black Non-Hispanic	*	*	*	737	26.5	(19.3 - 33.6)
Hispanic	*	*	*	856	17.2	(11.8 - 22.6)
Other Non-Hispanic	*	*	*	710	14.9	(7.7 - 22.0)
Education						
≤ High School	302	36.9	(29.1 - 44.7)	15,652	30.4	(26.9 - 33.8)
Some College	175	29.6	(20.3 - 38.8)	9,874	24.1	(21.0 - 27.1)
≥ College Degree	166	32.8	(23.9 - 41.8)	12,407	24.7	(21.9 - 27.5)
Income						
≤ \$24,999	*	*	*	9,618	31.2	(26.3 - 36.1)
\$25,000-\$49,999	199	29.7	(21.5 - 37.9)	9,624	27.3	(23.2 - 31.3)
\$50,000-\$74,999	*	*	*	5,753	26.3	(21.9 - 30.7)
≥ \$75,000	*	*	*	8,536	21.6	(18.8 - 24.3)

¹ Defined as adults having ever been told by a health professional that they have some form of arthritis, rheumatoid arthritis, gout, lupus or fibromyalgia (excluding back or neck)

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 51a. Activity limitations because of physical, mental or emotional problems among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	644	5,400	21.3	(17.1 - 25.5)	37,998	3,053,000	20.6	(18.7 - 22.5)
Gender								
Male	243	2,700	21.7	(14.7 - 28.6)	14,566	1,318,400	18.6	(15.7 - 21.5)
Female	401	2,700	21.0	(16.3 - 25.6)	23,432	1,734,600	22.4	(19.8 - 25.0)
Age (years)								
18-34	*	*	*	*	3,929	580,000	13.0	(8.2 - 17.9)
35-44	*	*	*	*	4,973	418,600	14.5	(11.0 - 18.1)
45-54	141	600	13.1	(6.1 - 20.1)	7,812	620,200	21.7	(18.3 - 25.2)
55-64	164	1,200	29.5	(22.0 - 37.0)	8,739	651,600	30.5	(26.5 - 34.6)
≥ 65	216	1,600	34.1	(27.0 - 41.2)	12,545	782,700	31.1	(27.7 - 34.6)
Race/Ethnicity								
White Non-Hispanic	629	5,200	21.0	(16.8 - 25.2)	35,688	2,037,200	21.9	(20.3 - 23.5)
Black Non-Hispanic	*	*	*	*	735	429,700	20.1	(13.5 - 26.7)
Hispanic	*	*	*	*	863	448,200	19.8	(11.9 - 27.8)
Other Non-Hispanic	*	*	*	*	712	137,900	12.3	(5.2 - 19.3)
Education								
≤ High School	303	3,000	25.0	(18.1 - 31.9)	15,626	1,359,600	24.4	(20.6 - 28.1)
Some College	174	1,500	19.7	(12.4 - 26.9)	9,875	700,800	19.8	(16.5 - 23.2)
≥ College Degree	166	900	15.8	(10.2 - 21.5)	12,415	924,800	16.5	(14.2 - 18.7)
Income								
≤ \$24,999	*	*	*	*	9,603	1,055,800	30.1	(24.7 - 35.5)
\$25,000-\$49,999	200	2,000	23.5	(16.1 - 31.0)	9,628	630,900	21.8	(16.6 - 27.0)
\$50,000-\$74,999	76	500	15.6	(7.8 - 23.4)	5,760	408,100	20.7	(16.1 - 25.2)
≥ \$75,000	119	500	8.4	(3.4 - 13.4)	8,544	583,300	13.0	(11.1 - 14.9)

¹ Defined as activity limitations because of physical, mental or emotional problems

² Sample size from survey

³ Estimated number of adults who are limited in any activities because of physical, mental or emotional problems (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 51b. Activity limitations because of physical, mental or emotional problems among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	644	20.6	(16.2 - 25.1)	37,998	20.1	(18.2 - 22.0)
Gender						
Male	243	21.5	(14.1 - 28.8)	14,566	18.6	(15.8 - 21.4)
Female	401	19.9	(14.9 - 24.8)	23,432	21.5	(18.9 - 24.1)
Race/Ethnicity						
White Non-Hispanic	629	20.4	(16.0 - 24.8)	35,688	21.3	(19.7 - 22.9)
Black Non-Hispanic	*	*	*	735	19.9	(13.2 - 26.5)
Hispanic	*	*	*	863	19.1	(11.5 - 26.8)
Other Non-Hispanic	*	*	*	712	12.4	(5.3 - 19.6)
Education						
≤ High School	303	24.8	(17.4 - 32.2)	15,626	23.8	(20.1 - 27.4)
Some College	174	18.5	(11.1 - 25.9)	9,875	19.3	(16.0 - 22.6)
≥ College Degree	166	14.9	(9.3 - 20.6)	12,415	16.1	(13.9 - 18.3)
Income						
≤ \$24,999	*	*	*	9,603	29.7	(24.3 - 35.2)
\$25,000-\$49,999	200	22.6	(14.9 - 30.3)	9,628	21.2	(16.1 - 26.3)
\$50,000-\$74,999	76	14.1	(6.6 - 21.5)	5,760	20.1	(15.6 - 24.6)
≥ \$75,000	119	8.6	(3.2 - 14.1)	8,544	12.7	(10.8 - 14.6)

¹ Defined as activity limitations because of physical, mental or emotional problems

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 52a. Have health problem that requires the use of special equipment among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	648	1,600	6.4	(4.4 - 8.3)	38,171	1,169,400	7.9	(6.9 - 8.9)
Gender								
Male	245	800	6.0	(3.0 - 9.0)	14,617	460,700	6.5	(5.3 - 7.7)
Female	403	900	6.7	(4.1 - 9.2)	23,554	708,700	9.1	(7.6 - 10.7)
Age (years)								
18-34	*	*	*	*	3,940	68,800	1.5	(0.7 - 2.4)
35-44	*	*	*	*	4,989	122,500	4.3	(2.4 - 6.1)
45-54	*	*	*	*	7,836	182,500	6.4	(4.7 - 8.1)
55-64	164	300	6.8	(2.8 - 10.7)	8,773	219,600	10.3	(7.8 - 12.8)
≥ 65	219	900	19.0	(13.0 - 24.9)	12,633	576,100	22.7	(19.0 - 26.3)
Race/Ethnicity								
White Non-Hispanic	633	1,600	6.5	(4.4 - 8.5)	35,853	802,000	8.6	(7.6 - 9.6)
Black Non-Hispanic	*	*	*	*	739	235,800	10.9	(6.4 - 15.5)
Hispanic	*	*	*	*	864	104,100	4.6	(2.1 - 7.1)
Other Non-Hispanic	*	*	*	*	715	27,600	2.5	(0.8 - 4.1)
Education								
≤ High School	305	900	7.3	(4.0 - 10.6)	15,711	604,200	10.8	(8.7 - 12.8)
Some College	175	400	5.3	(2.2 - 8.4)	9,914	260,000	7.3	(5.5 - 9.1)
≥ College Degree	167	300	5.7	(2.3 - 9.1)	12,463	298,900	5.3	(4.1 - 6.5)
Income								
≤ \$24,999	172	600	10.7	(6.1 - 15.3)	9,656	430,500	12.2	(9.6 - 14.8)
\$25,000-\$49,999	201	500	6.1	(2.5 - 9.8)	9,663	238,900	8.2	(5.9 - 10.5)
\$50,000-\$74,999	*	*	*	*	5,775	103,000	5.2	(3.6 - 6.8)
≥ \$75,000	*	*	*	*	8,566	136,000	3.0	(2.1 - 4.0)

¹ Defined as having any health problem that requires the use of special equipment such as a cane, a wheelchair, a special bed, or a special telephone

² Sample size from survey

³ Estimated number of adults who have any health problem that requires the use of special equipment (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 52b. Have health problem that requires the use of special equipment among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	648	5.9	(3.9 - 7.9)	38,171	7.6	(6.6 - 8.6)
Gender						
Male	245	5.8	(2.7 - 8.9)	14,617	6.8	(5.5 - 8.0)
Female	403	6.1	(3.4 - 8.7)	23,554	8.4	(6.9 - 9.9)
Race/Ethnicity						
White Non-Hispanic	633	6.0	(4.0 - 8.1)	35,853	8.3	(7.3 - 9.3)
Black Non-Hispanic	*	*	*	739	10.5	(6.1 - 15.0)
Hispanic	*	*	*	864	4.5	(2.1 - 6.9)
Other Non-Hispanic	*	*	*	715	2.2	(0.7 - 3.8)
Education						
≤ High School	305	6.7	(3.4 - 10.1)	15,711	10.3	(8.3 - 12.3)
Some College	175	5.0	(1.8 - 8.2)	9,914	7.1	(5.2 - 8.9)
≥ College Degree	167	5.5	(2.0 - 8.9)	12,463	5.3	(4.1 - 6.6)
Income						
≤ \$24,999	172	9.9	(5.4 - 14.3)	9,656	11.9	(9.4 - 14.5)
\$25,000-\$49,999	201	5.7	(2.0 - 9.4)	9,663	7.7	(5.5 - 10.0)
\$50,000-\$74,999	*	*	*	5,775	5.1	(3.4 - 6.8)
≥ \$75,000	*	*	*	8,566	2.9	(2.0 - 3.9)

¹ Defined as having any health problem that requires the use of special equipment such as a cane, a wheelchair, a special bed, or a special telephone

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 53a. Disability among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	645	5,600	22.0	(17.8 - 26.2)	38,016	3,394,200	22.9	(20.9 - 24.9)
Gender								
Male	244	2,800	21.9	(14.9 - 28.8)	14,568	1,469,700	20.7	(17.8 - 23.7)
Female	401	2,800	22.1	(17.3 - 26.8)	23,448	1,924,600	24.8	(22.1 - 27.5)
Age (years)								
18-34	*	*	*	*	3,931	590,300	13.3	(8.4 - 18.1)
35-44	*	*	*	*	4,974	447,900	15.6	(11.9 - 19.3)
45-54	141	600	13.1	(6.1 - 20.1)	7,811	643,900	22.6	(19.1 - 26.0)
55-64	164	1,200	30.6	(23.0 - 38.2)	8,740	702,100	33.0	(28.9 - 37.0)
≥ 65	217	1,800	36.7	(29.5 - 43.8)	12,560	1,010,000	39.9	(36.1 - 43.7)
Race/Ethnicity								
White Non-Hispanic	630	5,400	21.7	(17.5 - 25.9)	35,703	2,253,400	24.2	(22.6 - 25.9)
Black Non-Hispanic	*	*	*	*	736	511,700	23.8	(16.8 - 30.8)
Hispanic	*	*	*	*	864	484,300	21.4	(13.4 - 29.5)
Other Non-Hispanic	*	*	*	*	713	144,800	12.9	(5.8 - 19.9)
Education								
≤ High School	303	3,100	25.5	(18.6 - 32.4)	15,638	1,544,200	27.6	(23.7 - 31.4)
Some College	174	1,600	20.8	(13.4 - 28.1)	9,876	755,200	21.4	(17.9 - 24.8)
≥ College Degree	167	1,000	16.3	(10.6 - 22.0)	12,419	1,021,300	18.2	(15.9 - 20.5)
Income								
≤ \$24,999	*	*	*	*	9,613	1,179,100	33.6	(28.0 - 39.2)
\$25,000-\$49,999	200	2,000	23.7	(16.2 - 31.1)	9,629	699,000	24.2	(18.9 - 29.4)
\$50,000-\$74,999	76	500	16.6	(8.5 - 24.6)	5,757	430,100	21.8	(17.2 - 26.4)
≥ \$75,000	119	500	8.7	(3.6 - 13.7)	8,546	632,700	14.1	(12.1 - 16.1)

¹ Defined as having activity limitations due to physical, mental or emotional problems or have health problems that require the use of special equipment

² Sample size from survey

³ Estimated number of adults with a disability (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 53b. Disability among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	645	21.2	(16.8 - 25.6)	38,016	22.3	(20.4 - 24.3)
Gender						
Male	244	21.6	(14.3 - 29.0)	14,568	20.9	(18.0 - 23.8)
Female	401	20.8	(15.8 - 25.7)	23,448	23.6	(21.0 - 26.3)
Race/Ethnicity						
White Non-Hispanic	630	21.0	(16.5 - 25.4)	35,703	23.6	(21.9 - 25.3)
Black Non-Hispanic	*	*	*	736	23.5	(16.5 - 30.5)
Hispanic	*	*	*	864	20.6	(12.9 - 28.3)
Other Non-Hispanic	*	*	*	713	12.9	(5.8 - 20.1)
Education						
≤ High School	303	25.2	(17.8 - 32.6)	15,638	26.8	(23.0 - 30.6)
Some College	174	19.4	(11.9 - 26.8)	9,876	20.9	(17.4 - 24.3)
≥ College Degree	167	15.3	(9.6 - 21.0)	12,419	17.9	(15.5 - 20.3)
Income						
≤ \$24,999	*	*	*	9,613	33.2	(27.6 - 38.8)
\$25,000-\$49,999	200	22.7	(15.0 - 30.5)	9,629	23.5	(18.3 - 28.6)
\$50,000-\$74,999	76	14.8	(7.2 - 22.4)	5,757	21.2	(16.6 - 25.8)
≥ \$75,000	119	8.8	(3.4 - 14.3)	8,546	13.8	(11.7 - 15.8)

¹ Defined as having activity limitations due to physical, mental or emotional problems or have health problems that require the use of special equipment

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 54a. Current smoking among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	645	6,100	23.9	(18.8 - 29.0)	38,029	2,483,500	16.8	(15.1 - 18.6)
Gender								
Male	244	2,800	21.9	(14.1 - 29.6)	14,559	1,356,400	19.3	(16.3 - 22.2)
Female	401	3,300	25.8	(19.1 - 32.5)	23,470	1,127,100	14.6	(12.7 - 16.6)
Age (years)								
18-34	*	*	*	*	3,929	993,500	22.3	(17.5 - 27.1)
35-44	*	*	*	*	4,977	421,100	14.8	(11.6 - 17.9)
45-54	142	1,300	27.6	(19.0 - 36.1)	7,807	508,800	18.3	(15.4 - 21.3)
55-64	161	800	20.3	(13.4 - 27.1)	8,744	314,500	14.7	(12.1 - 17.3)
≥ 65	219	400	8.4	(4.5 - 12.3)	12,572	245,600	9.7	(7.2 - 12.3)
Race/Ethnicity								
White Non-Hispanic	630	5,900	23.7	(18.5 - 28.8)	35,729	1,615,500	17.3	(15.9 - 18.8)
Black Non-Hispanic	*	*	*	*	735	284,100	13.6	(8.9 - 18.2)
Hispanic	*	*	*	*	858	350,900	15.8	(9.6 - 21.9)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	304	3,900	32.1	(24.1 - 40.2)	15,654	1,237,300	22.4	(19.1 - 25.6)
Some College	174	1,800	24.2	(14.4 - 34.1)	9,875	670,600	19.1	(15.5 - 22.8)
≥ College Degree	166	400	6.1	(2.2 - 10.0)	12,421	563,300	10.0	(7.5 - 12.5)
Income								
≤ \$24,999	*	*	*	*	9,626	835,100	24.1	(19.1 - 29.1)
\$25,000-\$49,999	201	2,000	22.6	(14.6 - 30.6)	9,629	558,100	19.4	(15.6 - 23.2)
\$50,000-\$74,999	75	400	14.3	(6.5 - 22.1)	5,756	338,100	17.3	(13.0 - 21.7)
≥ \$75,000	*	*	*	*	8,549	468,700	10.4	(8.6 - 12.3)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day or some days

² Sample size from survey

³ Estimated number of adult current smokers (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 54b. Current smoking among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	645	24.3	(18.9 - 29.8)	38,029	17.0	(15.3 - 18.8)
Gender						
Male	244	22.0	(13.7 - 30.3)	14,559	19.2	(16.3 - 22.1)
Female	401	26.6	(19.5 - 33.8)	23,470	15.0	(13.0 - 17.0)
Race/Ethnicity						
White Non-Hispanic	630	24.1	(18.6 - 29.6)	35,729	17.7	(16.2 - 19.2)
Black Non-Hispanic	*	*	*	735	13.5	(8.8 - 18.2)
Hispanic	*	*	*	858	15.9	(9.8 - 22.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	304	32.9	(24.3 - 41.5)	15,654	22.8	(19.5 - 26.0)
Some College	*	*	*	9,875	19.4	(15.8 - 22.9)
≥ College Degree	166	6.5	(2.2 - 10.8)	12,421	10.0	(7.6 - 12.5)
Income						
≤ \$24,999	*	*	*	9,626	24.1	(19.2 - 29.1)
\$25,000-\$49,999	201	22.4	(14.1 - 30.8)	9,629	19.9	(16.0 - 23.7)
\$50,000-\$74,999	75	12.5	(5.2 - 19.8)	5,756	17.8	(13.3 - 22.3)
≥ \$75,000	*	*	*	8,549	10.6	(8.6 - 12.5)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day or some days

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 55a. Everyday smoking among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	645	5,100	19.8	(14.9 - 24.8)	38,029	1,626,800	11.0	(9.8 - 12.2)
Gender								
Male	244	2,500	20.0	(12.4 - 27.6)	14,559	848,400	12.0	(10.0 - 14.1)
Female	401	2,500	19.6	(13.3 - 26.0)	23,470	778,400	10.1	(8.7 - 11.5)
Age (years)								
18-34	*	*	*	*	3,929	559,800	12.6	(9.6 - 15.6)
35-44	*	*	*	*	4,977	315,200	11.0	(8.4 - 13.7)
45-54	142	1,000	20.5	(12.5 - 28.4)	7,807	377,000	13.6	(11.1 - 16.0)
55-64	161	700	16.7	(10.3 - 23.1)	8,744	203,000	9.5	(7.5 - 11.5)
≥ 65	219	300	6.8	(3.2 - 10.4)	12,572	171,700	6.8	(4.8 - 8.8)
Race/Ethnicity								
White Non-Hispanic	630	4,900	19.8	(14.8 - 24.8)	35,729	1,158,100	12.4	(11.2 - 13.7)
Black Non-Hispanic	*	*	*	*	735	197,700	9.4	(5.6 - 13.3)
Hispanic	*	*	*	*	858	176,200	7.9	(4.2 - 11.6)
Other Non-Hispanic	*	*	*	*	707	94,900	8.5	(3.3 - 13.7)
Education								
≤ High School	304	3,400	28.0	(20.0 - 36.0)	15,654	881,900	15.9	(13.4 - 18.5)
Some College	174	1,400	18.9	(9.7 - 28.1)	9,875	410,200	11.7	(9.6 - 13.8)
≥ College Degree	*	*	*	*	12,421	323,500	5.8	(4.3 - 7.2)
Income								
≤ \$24,999	*	*	*	*	9,626	452,800	13.1	(10.3 - 15.8)
\$25,000-\$49,999	201	1,700	19.3	(11.8 - 26.8)	9,629	425,900	14.8	(11.9 - 17.7)
\$50,000-\$74,999	*	*	*	*	5,756	222,700	11.4	(8.5 - 14.4)
≥ \$75,000	*	*	*	*	8,549	320,200	7.1	(5.5 - 8.8)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day

² Sample size from survey

³ Estimated number of adults who have smoked at least 100 cigarettes in lifetime and currently smoking everyday (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 55b. Everyday smoking among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	645	20.3	(15.0 - 25.6)	38,029	11.3	(10.0 - 12.5)
Gender						
Male	244	20.1	(12.0 - 28.3)	14,559	12.1	(10.1 - 14.1)
Female	401	20.4	(13.6 - 27.3)	23,470	10.5	(9.0 - 12.0)
Race/Ethnicity						
White Non-Hispanic	630	20.3	(15.0 - 25.7)	35,729	12.8	(11.5 - 14.1)
Black Non-Hispanic	*	*	*	735	9.4	(5.5 - 13.4)
Hispanic	*	*	*	858	8.0	(4.3 - 11.7)
Other Non-Hispanic	*	*	*	707	8.5	(3.2 - 13.8)
Education						
≤ High School	304	28.6	(20.0 - 37.1)	15,654	16.4	(13.8 - 19.0)
Some College	174	18.9	(9.3 - 28.6)	9,875	12.1	(9.9 - 14.3)
≥ College Degree	*	*	*	12,421	5.8	(4.3 - 7.3)
Income						
≤ \$24,999	*	*	*	9,626	13.4	(10.6 - 16.2)
\$25,000-\$49,999	201	18.9	(11.2 - 26.7)	9,629	15.3	(12.3 - 18.3)
\$50,000-\$74,999	*	*	*	5,756	11.7	(8.7 - 14.8)
≥ \$75,000	*	*	*	8,549	7.2	(5.5 - 8.9)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 56a. Adults living in homes in which smoking is prohibited¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	644	19,100	74.5	(69.6 - 79.4)	38,045	11,997,100	80.8	(79.0 - 82.6)
Gender								
Male	245	9,000	71.1	(63.2 - 78.9)	14,577	5,634,400	79.4	(76.6 - 82.2)
Female	399	10,000	77.9	(72.1 - 83.6)	23,468	6,362,800	82.1	(79.9 - 84.3)
Age (years)								
18-34	*	*	*	*	3,934	3,655,200	82.1	(77.8 - 86.4)
35-44	*	*	*	*	4,985	2,479,700	86.1	(82.5 - 89.7)
45-54	142	3,300	69.4	(60.8 - 78.1)	7,808	2,194,400	77.2	(73.5 - 80.8)
55-64	161	2,800	71.0	(63.1 - 78.8)	8,753	1,732,500	81.0	(78.3 - 83.7)
≥ 65	218	3,800	79.2	(73.5 - 84.8)	12,565	1,935,400	76.5	(73.4 - 79.5)
Race/Ethnicity								
White Non-Hispanic	629	18,500	74.4	(69.5 - 79.3)	35,737	7,426,900	79.8	(78.3 - 81.2)
Black Non-Hispanic	*	*	*	*	738	1,692,100	78.6	(71.9 - 85.3)
Hispanic	*	*	*	*	863	2,003,900	88.7	(84.1 - 93.3)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	301	7,700	64.0	(56.0 - 72.0)	15,647	4,307,200	77.1	(74.0 - 80.2)
Some College	175	6,100	80.6	(73.0 - 88.2)	9,883	2,795,400	78.9	(75.0 - 82.8)
≥ College Degree	167	5,300	88.4	(83.1 - 93.7)	12,437	4,817,300	85.7	(83.3 - 88.2)
Income								
≤ \$24,999	*	*	*	*	9,608	2,627,700	74.9	(70.0 - 79.7)
\$25,000-\$49,999	200	6,600	76.0	(67.8 - 84.3)	9,640	2,314,500	79.8	(76.0 - 83.5)
\$50,000-\$74,999	76	2,600	83.9	(75.7 - 92.2)	5,766	1,551,300	78.7	(74.0 - 83.4)
≥ \$75,000	119	4,800	87.7	(81.8 - 93.7)	8,556	3,883,200	86.5	(84.3 - 88.7)

¹ Defined as having a rule that prohibits smoking anywhere inside the home

² Sample size from survey

³ Estimated number of adults who live in a home with a rule that prohibits smoking anywhere inside the home (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 56b. Adults living in homes in which smoking is prohibited¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	644	74.9	(69.7 - 80.0)	38,045	80.9	(79.2 - 82.7)
Gender						
Male	245	71.3	(63.0 - 79.7)	14,577	79.4	(76.7 - 82.2)
Female	399	78.4	(72.3 - 84.4)	23,468	82.3	(80.0 - 84.5)
Race/Ethnicity						
White Non-Hispanic	629	74.7	(69.5 - 80.0)	35,737	79.9	(78.4 - 81.4)
Black Non-Hispanic	*	*	*	738	78.6	(71.8 - 85.4)
Hispanic	*	*	*	863	88.9	(84.4 - 93.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	301	64.1	(55.6 - 72.7)	15,647	76.9	(73.8 - 80.0)
Some College	175	81.5	(73.8 - 89.2)	9,883	79.1	(75.2 - 83.0)
≥ College Degree	167	88.8	(83.4 - 94.2)	12,437	86.0	(83.5 - 88.5)
Income						
≤ \$24,999	*	*	*	9,608	74.9	(70.1 - 79.7)
\$25,000-\$49,999	200	76.7	(68.1 - 85.3)	9,640	79.8	(76.1 - 83.6)
\$50,000-\$74,999	76	85.8	(78.1 - 93.5)	5,766	78.6	(73.8 - 83.3)
≥ \$75,000	119	87.8	(81.7 - 94.0)	8,556	86.6	(84.3 - 88.9)

¹ Defined as having a rule that prohibits smoking anywhere inside the home

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 57a. Binge drinking within the past month among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	641	4,000	15.9	(11.4 - 20.4)	37,745	2,602,000	17.7	(15.7 - 19.8)
Gender								
Male	243	2,900	22.7	(14.9 - 30.5)	14,420	1,712,400	24.5	(20.9 - 28.1)
Female	398	1,200	9.2	(5.1 - 13.2)	23,325	889,600	11.6	(9.5 - 13.7)
Age (years)								
18-34	*	*	*	*	3,895	1,254,300	28.5	(22.8 - 34.2)
35-44	*	*	*	*	4,934	584,800	20.6	(16.5 - 24.7)
45-54	141	700	15.3	(8.9 - 21.7)	7,738	467,600	16.6	(13.3 - 19.9)
55-64	163	400	9.2	(4.8 - 13.7)	8,663	207,200	9.8	(7.9 - 11.8)
≥ 65	*	*	*	*	12,515	88,100	3.5	(2.5 - 4.5)
Race/Ethnicity								
White Non-Hispanic	626	4,000	16.4	(11.8 - 21.0)	35,453	1,761,900	19.2	(17.5 - 20.8)
Black Non-Hispanic	*	*	*	*	728	247,000	11.6	(6.3 - 17.0)
Hispanic	*	*	*	*	854	460,600	20.5	(11.3 - 29.8)
Other Non-Hispanic	*	*	*	*	710	132,500	12.1	(3.4 - 20.7)
Education								
≤ High School	301	1,700	14.4	(8.4 - 20.3)	15,491	945,800	17.4	(13.2 - 21.6)
Some College	174	1,400	18.8	(8.8 - 28.8)	9,819	636,600	18.1	(14.6 - 21.5)
≥ College Degree	165	900	15.6	(7.3 - 23.8)	12,358	1,018,700	18.2	(15.3 - 21.1)
Income								
≤ \$24,999	172	400	7.8	(1.6 - 13.9)	9,557	469,100	13.5	(8.5 - 18.5)
\$25,000-\$49,999	199	1,400	16.3	(8.4 - 24.1)	9,575	534,300	18.5	(13.0 - 24.0)
\$50,000-\$74,999	*	*	*	*	5,726	378,200	19.2	(14.1 - 24.3)
≥ \$75,000	*	*	*	*	8,501	1,011,400	22.6	(19.7 - 25.6)

¹ Defined as men having 5 or more drinks or women having 4 or more drinks on 1 or more occasion within the past month

² Sample size from survey

³ Estimated number of adult binge drinkers within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 57b. Binge drinking within the past month among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	641	16.7	(11.8 - 21.6)	37,745	18.1	(16.1 - 20.2)
Gender						
Male	243	23.7	(15.2 - 32.1)	14,420	24.5	(21.0 - 27.9)
Female	398	9.8	(5.4 - 14.3)	23,325	12.4	(10.2 - 14.5)
Race/Ethnicity						
White Non-Hispanic	626	17.2	(12.2 - 22.2)	35,453	19.8	(18.1 - 21.6)
Black Non-Hispanic	*	*	*	728	11.8	(6.4 - 17.1)
Hispanic	*	*	*	854	20.2	(11.3 - 29.1)
Other Non-Hispanic	*	*	*	710	12.1	(3.4 - 20.8)
Education						
≤ High School	301	14.9	(8.5 - 21.4)	15,491	17.5	(13.5 - 21.5)
Some College	*	*	*	9,819	18.6	(15.1 - 22.0)
≥ College Degree	165	16.8	(7.7 - 25.9)	12,358	18.8	(15.9 - 21.7)
Income						
≤ \$24,999	172	8.0	(1.3 - 14.7)	9,557	13.4	(8.6 - 18.2)
\$25,000-\$49,999	199	16.9	(8.6 - 25.3)	9,575	18.9	(13.5 - 24.2)
\$50,000-\$74,999	*	*	*	5,726	19.9	(14.8 - 25.1)
≥ \$75,000	*	*	*	8,501	23.2	(20.2 - 26.1)

¹ Defined as men having 5 or more drinks or women having 4 or more drinks on 1 or more occasion within the past month

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 58a. Heavy drinking within the past month among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	635	1,400	5.6	(3.2 - 8.1)	37,553	718,700	4.9	(4.2 - 5.6)
Gender								
Male	237	800	6.6	(2.6 - 10.5)	14,330	333,300	4.8	(3.9 - 5.7)
Female	398	600	4.7	(1.7 - 7.8)	23,223	385,400	5.0	(4.0 - 6.1)
Age (years)								
18-34	*	*	*	*	3,866	224,900	5.1	(3.5 - 6.8)
35-44	*	*	*	*	4,924	137,200	4.8	(3.0 - 6.7)
45-54	*	*	*	*	7,715	144,600	5.2	(4.1 - 6.2)
55-64	161	200	6.3	(2.3 - 10.3)	8,624	117,400	5.6	(3.9 - 7.3)
≥ 65	*	*	*	*	12,424	94,600	3.8	(2.7 - 4.9)
Race/Ethnicity								
White Non-Hispanic	620	1,400	5.8	(3.2 - 8.3)	35,279	572,400	6.3	(5.4 - 7.1)
Black Non-Hispanic	*	*	*	*	723	58,100	2.8	(0.6 - 4.9)
Hispanic	*	*	*	*	848	60,900	2.7	(0.5 - 4.9)
Other Non-Hispanic	*	*	*	*	703	27,200	2.5	(0.8 - 4.2)
Education								
≤ High School	298	400	3.8	(1.0 - 6.6)	15,411	228,100	4.2	(3.2 - 5.2)
Some College	174	700	8.9	(2.5 - 15.3)	9,759	163,700	4.7	(3.6 - 5.8)
≥ College Degree	162	300	5.2	(1.7 - 8.7)	12,306	326,800	5.9	(4.5 - 7.3)
Income								
≤ \$24,999	*	*	*	*	9,512	104,200	3.0	(1.8 - 4.1)
\$25,000-\$49,999	200	400	5.0	(1.2 - 8.8)	9,537	158,700	5.5	(3.9 - 7.2)
\$50,000-\$74,999	*	*	*	*	5,708	95,400	4.9	(3.1 - 6.7)
≥ \$75,000	*	*	*	*	8,481	313,900	7.0	(5.4 - 8.6)

¹ Defined as adult men averaging more than 2 alcoholic drinks per day and adult women averaging more than 1 alcoholic drink per day within the past month

² Sample size from survey

³ Estimated number of adult heavy drinkers within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 58b. Heavy drinking within the past month among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	635	5.8	(3.1 - 8.5)	37,553	5.0	(4.3 - 5.8)
Gender						
Male	237	6.5	(2.3 - 10.7)	14,330	4.9	(3.9 - 5.9)
Female	398	5.1	(1.8 - 8.4)	23,223	5.1	(4.0 - 6.2)
Race/Ethnicity						
White Non-Hispanic	620	6.0	(3.2 - 8.7)	35,279	6.4	(5.5 - 7.2)
Black Non-Hispanic	*	*	*	723	2.9	(0.6 - 5.1)
Hispanic	*	*	*	848	2.8	(0.5 - 5.1)
Other Non-Hispanic	*	*	*	703	2.5	(0.8 - 4.2)
Education						
≤ High School	298	3.8	(0.8 - 6.8)	15,411	4.3	(3.2 - 5.3)
Some College	174	9.4	(2.5 - 16.3)	9,759	4.8	(3.6 - 5.9)
≥ College Degree	162	5.2	(1.5 - 8.9)	12,306	6.0	(4.5 - 7.4)
Income						
≤ \$24,999	*	*	*	9,512	3.0	(1.9 - 4.1)
\$25,000-\$49,999	200	5.3	(1.2 - 9.5)	9,537	5.8	(4.0 - 7.5)
\$50,000-\$74,999	*	*	*	5,708	5.0	(3.1 - 7.0)
≥ \$75,000	*	*	*	8,481	7.0	(5.4 - 8.7)

¹ Defined as adult men averaging more than 2 alcoholic drinks per day and adult women averaging more than 1 alcoholic drink per day within the past month

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 59a. Never or rarely been asked about sexual history during routine check-up among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	594	15,500	67.7	(62.3 - 73.2)	35,193	8,480,600	61.9	(59.2 - 64.5)
Gender								
Male	223	9,000	79.8	(72.4 - 87.2)	13,299	4,357,400	67.7	(63.6 - 71.7)
Female	371	6,500	56.0	(48.7 - 63.3)	21,894	4,123,300	56.8	(53.3 - 60.2)
Age (years)								
18-34	*	*	*	*	3,507	1,804,600	44.3	(37.8 - 50.8)
35-44	*	*	*	*	4,531	1,635,300	62.2	(56.6 - 67.8)
45-54	132	2,600	59.6	(50.0 - 69.2)	7,215	1,715,600	65.4	(60.5 - 70.3)
55-64	151	2,900	77.9	(70.7 - 85.2)	8,168	1,416,200	70.7	(66.6 - 74.8)
≥ 65	205	3,500	77.2	(70.6 - 83.7)	11,772	1,908,900	80.3	(76.9 - 83.8)
Race/Ethnicity								
White Non-Hispanic	580	15,200	68.0	(62.5 - 73.6)	33,090	5,940,300	68.5	(66.3 - 70.7)
Black Non-Hispanic	*	*	*	*	683	831,800	40.8	(31.8 - 49.9)
Hispanic	*	*	*	*	784	1,071,700	53.4	(43.6 - 63.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	275	7,800	71.4	(64.1 - 78.8)	14,264	3,001,200	60.1	(55.4 - 64.9)
Some College	*	*	*	*	9,156	1,972,800	60.3	(54.9 - 65.7)
≥ College Degree	*	*	*	*	11,706	3,476,000	65.0	(61.2 - 68.7)
Income								
≤ \$24,999	*	*	*	*	8,772	1,529,400	49.3	(42.6 - 56.0)
\$25,000-\$49,999	184	4,800	63.0	(53.2 - 72.8)	8,932	1,705,000	64.4	(58.0 - 70.7)
\$50,000-\$74,999	*	*	*	*	5,433	1,217,000	64.5	(58.5 - 70.5)
≥ \$75,000	*	*	*	*	8,108	2,934,300	68.2	(64.7 - 71.7)

¹ Defined as respondents who never or rarely (hardly ever) were asked about their sexual history, such as about sexual partners and sexual practices, during a routine check-up at a doctor's office or clinic

² Sample size from survey

³ Estimated number of adults who were never or rarely asked about their sexual history during a routine check-up at a doctor's office or clinic (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 59b. Never or rarely been asked about sexual history during routine check-up among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	594	66.4	(60.5 - 72.3)	35,193	61.4	(58.7 - 64.0)
Gender						
Male	223	79.7	(71.8 - 87.6)	13,299	67.9	(64.0 - 71.9)
Female	371	53.5	(45.7 - 61.3)	21,894	55.5	(52.0 - 59.0)
Race/Ethnicity						
White Non-Hispanic	580	66.8	(60.8 - 72.7)	33,090	67.8	(65.6 - 70.1)
Black Non-Hispanic	*	*	*	683	41.1	(32.0 - 50.2)
Hispanic	*	*	*	784	53.0	(43.3 - 62.6)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	275	70.7	(62.8 - 78.7)	14,264	59.5	(54.7 - 64.3)
Some College	*	*	*	9,156	59.7	(54.4 - 65.0)
≥ College Degree	*	*	*	11,706	64.5	(60.8 - 68.2)
Income						
≤ \$24,999	*	*	*	8,772	48.7	(42.0 - 55.4)
\$25,000-\$49,999	*	*	*	8,932	63.7	(57.5 - 70.0)
\$50,000-\$74,999	*	*	*	5,433	64.2	(58.2 - 70.2)
≥ \$75,000	*	*	*	8,108	67.5	(64.0 - 71.0)

¹ Defined as respondents who never or rarely (hardly ever) were asked about their sexual history, such as about sexual partners and sexual practices, during a routine check-up at a doctor's office or clinic

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 60a. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	368	14,000	83.2	(77.4 - 89.0)	22,788	7,011,100	77.9	(75.2 - 80.5)
Gender								
Male	135	7,300	91.6	(83.8 - 99.5)	9,034	3,618,500	81.0	(77.2 - 84.8)
Female	233	6,700	75.6	(67.4 - 83.9)	13,754	3,392,600	74.8	(71.2 - 78.4)
Age (years)								
18-34	*	*	*	*	3,213	2,362,500	70.8	(65.1 - 76.6)
35-44	50	3,500	86.8	(77.6 - 96.1)	3,513	1,313,400	71.7	(65.6 - 77.7)
45-54	89	2,600	86.5	(79.0 - 93.9)	4,975	1,290,600	80.3	(75.9 - 84.8)
55-64	88	2,000	93.4	(89.1 - 97.7)	4,955	932,200	87.2	(84.1 - 90.3)
≥ 65	98	2,100	96.5	(92.3 - 100)	6,132	1,112,400	95.9	(94.4 - 97.5)
Race/Ethnicity								
White Non-Hispanic	361	13,500	82.8	(76.9 - 88.7)	21,540	4,875,200	78.3	(75.6 - 80.9)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	491	1,002,400	84.8	(77.8 - 91.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	172	6,700	87.2	(80.0 - 94.4)	8,879	2,581,700	82.1	(77.9 - 86.3)
Some College	*	*	*	*	5,953	1,599,500	77.5	(72.6 - 82.3)
≥ College Degree	95	3,100	84.1	(75.5 - 92.7)	7,928	2,824,800	75.6	(71.8 - 79.5)
Income								
≤ \$24,999	*	*	*	*	5,491	1,602,900	78.7	(71.9 - 85.6)
\$25,000-\$49,999	125	5,400	86.4	(78.8 - 93.9)	5,912	1,335,900	78.8	(73.8 - 83.9)
\$50,000-\$74,999	*	*	*	*	3,717	987,500	76.6	(70.4 - 82.8)
≥ \$75,000	*	*	*	*	5,771	2,321,600	77.6	(74.0 - 81.2)

¹ Defined as respondent estimating there were hardly any (0-1 out of 10) or a few (2-3 out of 10) persons their age in their community who have had a sexually transmitted disease such as gonorrhea, chlamydia, herpes or syphilis

² Sample size from survey

³ Estimated number of adults who believed hardly any or a few people their age in their community have had a sexually transmitted disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 60b. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	368	82.3	(76.1 - 88.5)	22,788	77.3	(74.7 - 79.9)
Gender						
Male	135	91.0	(82.6 - 99.5)	9,034	81.1	(77.4 - 84.8)
Female	233	74.5	(65.9 - 83.2)	13,754	73.6	(70.0 - 77.3)
Race/Ethnicity						
White Non-Hispanic	361	81.8	(75.5 - 88.2)	21,540	77.5	(74.9 - 80.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	491	84.6	(77.6 - 91.5)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	172	86.4	(78.6 - 94.2)	8,879	81.6	(77.4 - 85.7)
Some College	*	*	*	5,953	76.8	(72.0 - 81.6)
≥ College Degree	95	83.0	(73.6 - 92.4)	7,928	75.1	(71.3 - 79.0)
Income						
≤ \$24,999	*	*	*	5,491	77.9	(71.0 - 84.7)
\$25,000-\$49,999	125	85.4	(77.3 - 93.5)	5,912	77.9	(72.8 - 83.0)
\$50,000-\$74,999	*	*	*	3,717	76.1	(70.0 - 82.3)
≥ \$75,000	*	*	*	5,771	77.0	(73.4 - 80.6)

¹ Defined as respondent estimating there were hardly any (0-1 out of 10) or a few (2-3 out of 10) persons their age in their community who have had a sexually transmitted disease such as gonorrhea, chlamydia, herpes or syphilis

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 61a. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	607	22,300	90.9	(87.8 - 94.0)	36,287	12,738,300	89.7	(88.2 - 91.3)
Gender								
Male	225	10,800	90.0	(84.6 - 95.3)	13,893	6,104,600	89.4	(87.2 - 91.5)
Female	382	11,500	91.8	(88.6 - 95.0)	22,394	6,633,700	90.1	(87.9 - 92.2)
Age (years)								
18-34	*	*	*	*	3,850	4,007,400	91.3	(87.5 - 95.1)
35-44	72	5,000	89.5	(80.5 - 98.6)	4,875	2,578,600	93.2	(90.4 - 96.0)
45-54	139	4,500	97.0	(94.6 - 99.5)	7,607	2,416,800	90.5	(87.9 - 93.2)
55-64	155	3,500	91.6	(87.2 - 95.9)	8,411	1,876,500	92.2	(90.0 - 94.5)
≥ 65	192	3,400	79.5	(73.1 - 86.0)	11,544	1,858,900	79.7	(76.6 - 82.8)
Race/Ethnicity								
White Non-Hispanic	593	21,700	91.1	(87.9 - 94.2)	34,102	8,050,700	89.4	(87.7 - 91.0)
Black Non-Hispanic	*	*	*	*	702	1,822,200	91.9	(88.2 - 95.6)
Hispanic	*	*	*	*	820	1,967,200	89.9	(85.2 - 94.7)
Other Non-Hispanic	*	*	*	*	663	898,200	88.6	(79.5 - 97.6)
Education								
≤ High School	276	9,800	85.7	(79.9 - 91.4)	14,507	4,385,300	84.6	(81.9 - 87.2)
Some College	166	7,000	95.3	(92.0 - 98.6)	9,571	3,094,300	90.0	(86.7 - 93.4)
≥ College Degree	164	5,600	95.5	(91.0 - 99.9)	12,142	5,228,800	95.4	(94.0 - 96.7)
Income								
≤ \$24,999	157	4,700	89.1	(84.3 - 93.9)	9,021	2,833,500	86.5	(82.6 - 90.3)
\$25,000-\$49,999	195	7,700	90.5	(84.7 - 96.4)	9,261	2,529,900	89.5	(86.5 - 92.5)
\$50,000-\$74,999	74	2,800	92.3	(83.3 - 100)	5,644	1,740,000	91.2	(87.4 - 95.1)
≥ \$75,000	114	4,900	92.8	(84.9 - 100)	8,397	4,184,300	94.4	(92.5 - 96.3)

¹ Defined as believing it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community

² Sample size from survey

³ Estimated number of adults who believed it is at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 61b. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	607	91.1	(87.7 - 94.4)	36,287	89.8	(88.3 - 91.4)
Gender						
Male	225	89.8	(84.0 - 95.5)	13,893	89.1	(86.9 - 91.3)
Female	382	92.3	(88.9 - 95.7)	22,394	90.5	(88.3 - 92.7)
Race/Ethnicity						
White Non-Hispanic	593	91.2	(87.8 - 94.6)	34,102	89.5	(87.8 - 91.2)
Black Non-Hispanic	*	*	*	702	91.7	(87.8 - 95.6)
Hispanic	*	*	*	820	89.6	(84.7 - 94.5)
Other Non-Hispanic	*	*	*	663	89.0	(80.5 - 97.6)
Education						
≤ High School	276	86.3	(80.2 - 92.4)	14,507	84.6	(81.8 - 87.4)
Some College	166	95.0	(91.0 - 98.9)	9,571	90.2	(86.9 - 93.5)
≥ College Degree	164	95.5	(90.7 - 100)	12,142	95.4	(94.0 - 96.7)
Income						
≤ \$24,999	157	90.3	(85.8 - 94.8)	9,021	86.5	(82.6 - 90.4)
\$25,000-\$49,999	195	90.2	(83.8 - 96.6)	9,261	89.6	(86.5 - 92.6)
\$50,000-\$74,999	74	91.9	(82.4 - 100)	5,644	91.1	(86.9 - 95.3)
≥ \$75,000	114	92.6	(84.0 - 100)	8,397	94.5	(92.6 - 96.4)

¹ Defined as believing it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 62a. Overweight among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	625	8,900	36.2	(30.8 - 41.6)	36,710	5,213,000	36.4	(34.0 - 38.8)
Gender								
Male	243	5,100	40.6	(32.0 - 49.1)	14,502	3,073,200	43.7	(39.9 - 47.6)
Female	382	3,800	31.6	(25.2 - 38.1)	22,208	2,139,800	29.3	(26.4 - 32.2)
Age (years)								
18-34	*	*	*	*	3,789	1,295,200	30.1	(24.7 - 35.6)
35-44	*	*	*	*	4,785	1,052,000	37.5	(31.7 - 43.3)
45-54	137	1,600	34.9	(26.0 - 43.8)	7,495	1,053,300	38.8	(34.1 - 43.4)
55-64	159	1,200	30.2	(22.2 - 38.1)	8,356	798,100	38.8	(34.8 - 42.7)
≥ 65	214	2,200	45.6	(38.1 - 53.2)	12,285	1,014,400	41.5	(37.6 - 45.4)
Race/Ethnicity								
White Non-Hispanic	610	8,500	35.7	(30.2 - 41.2)	34,501	3,282,200	36.4	(34.4 - 38.3)
Black Non-Hispanic	*	*	*	*	708	740,000	36.7	(28.3 - 45.1)
Hispanic	*	*	*	*	813	781,500	35.6	(27.1 - 44.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	294	4,000	34.9	(27.0 - 42.8)	15,079	1,889,700	35.4	(31.2 - 39.5)
Some College	*	*	*	*	9,515	1,314,500	38.7	(33.8 - 43.7)
≥ College Degree	*	*	*	*	12,062	1,996,900	36.3	(32.8 - 39.7)
Income								
≤ \$24,999	*	*	*	*	9,346	1,170,900	34.0	(28.2 - 39.8)
\$25,000-\$49,999	195	2,900	34.7	(25.7 - 43.7)	9,368	1,136,000	40.4	(34.6 - 46.2)
\$50,000-\$74,999	*	*	*	*	5,615	738,700	38.3	(32.3 - 44.3)
≥ \$75,000	*	*	*	*	8,337	1,659,100	37.5	(34.1 - 40.9)

¹ Defined as having a body mass index (BMI) more than 24.9 but less than 30.0. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Sample size from survey

³ Estimated number of adults who are overweight (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 62b. Overweight among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	625	36.2	(30.4 - 42.0)	36,710	36.2	(33.8 - 38.6)
Gender						
Male	243	40.9	(31.8 - 50.0)	14,502	43.9	(40.1 - 47.6)
Female	382	31.4	(24.5 - 38.2)	22,208	28.9	(25.9 - 31.8)
Race/Ethnicity						
White Non-Hispanic	610	35.7	(29.8 - 41.5)	34,501	36.0	(34.0 - 38.0)
Black Non-Hispanic	*	*	*	708	37.0	(28.5 - 45.5)
Hispanic	*	*	*	813	36.0	(27.5 - 44.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	294	35.1	(26.6 - 43.6)	15,079	35.2	(31.1 - 39.4)
Some College	*	*	*	9,515	38.3	(33.3 - 43.2)
≥ College Degree	*	*	*	12,062	36.2	(32.7 - 39.8)
Income						
≤ \$24,999	*	*	*	9,346	33.8	(28.0 - 39.5)
\$25,000-\$49,999	195	34.1	(24.6 - 43.5)	9,368	40.6	(34.8 - 46.4)
\$50,000-\$74,999	*	*	*	5,615	38.6	(32.5 - 44.7)
≥ \$75,000	*	*	*	8,337	37.2	(33.7 - 40.6)

¹ Defined as having a body mass index (BMI) more than 24.9 but less than 30.0. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 63a. Obesity among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	625	7,400	29.9	(24.7 - 35.1)	36,710	3,327,400	23.2	(21.2 - 25.3)
Gender								
Male	243	3,900	30.7	(22.8 - 38.6)	14,502	1,505,800	21.4	(19.0 - 23.9)
Female	382	3,500	29.1	(22.4 - 35.8)	22,208	1,821,600	25.0	(21.8 - 28.1)
Age (years)								
18-34	*	*	*	*	3,789	730,600	17.0	(12.4 - 21.6)
35-44	*	*	*	*	4,785	662,200	23.6	(19.4 - 27.9)
45-54	137	1,400	29.9	(20.8 - 39.1)	7,495	816,800	30.1	(25.3 - 34.8)
55-64	159	1,300	34.1	(26.1 - 42.2)	8,356	563,900	27.4	(23.7 - 31.1)
≥ 65	214	1,200	25.9	(19.1 - 32.6)	12,285	554,000	22.7	(19.2 - 26.1)
Race/Ethnicity								
White Non-Hispanic	610	7,100	29.7	(24.4 - 35.0)	34,501	2,062,700	22.9	(21.2 - 24.6)
Black Non-Hispanic	*	*	*	*	708	703,100	34.9	(26.1 - 43.7)
Hispanic	*	*	*	*	813	456,400	20.8	(14.3 - 27.2)
Other Non-Hispanic	*	*	*	*	688	105,200	9.6	(4.0 - 15.3)
Education								
≤ High School	294	4,100	35.2	(27.4 - 43.0)	15,079	1,561,600	29.2	(25.3 - 33.2)
Some College	168	1,800	24.2	(15.2 - 33.2)	9,515	792,800	23.4	(19.4 - 27.3)
≥ College Degree	*	*	*	*	12,062	915,000	16.6	(14.4 - 18.8)
Income								
≤ \$24,999	171	1,700	30.1	(20.3 - 39.9)	9,346	974,800	28.3	(22.6 - 34.0)
\$25,000-\$49,999	195	2,800	33.0	(23.5 - 42.4)	9,368	584,800	20.8	(17.1 - 24.5)
\$50,000-\$74,999	74	500	17.6	(8.8 - 26.5)	5,615	475,300	24.6	(20.0 - 29.3)
≥ \$75,000	*	*	*	*	8,337	867,700	19.6	(17.0 - 22.3)

¹ Defined as having a body mass index (BMI) of 30.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Sample size from survey

³ Estimated number of adults who are obese (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 63b. Obesity among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	625	29.8	(24.2 - 35.3)	36,710	23.1	(21.0 - 25.1)
Gender						
Male	243	30.7	(22.2 - 39.1)	14,502	21.5	(19.0 - 24.0)
Female	382	28.8	(21.6 - 36.0)	22,208	24.6	(21.5 - 27.8)
Race/Ethnicity						
White Non-Hispanic	610	29.6	(23.9 - 35.2)	34,501	22.8	(21.0 - 24.5)
Black Non-Hispanic	*	*	*	708	34.8	(25.9 - 43.6)
Hispanic	*	*	*	813	20.5	(14.1 - 26.9)
Other Non-Hispanic	*	*	*	688	9.5	(3.8 - 15.2)
Education						
≤ High School	294	35.7	(27.3 - 44.0)	15,079	29.4	(25.4 - 33.3)
Some College	168	23.2	(13.8 - 32.6)	9,515	23.1	(19.2 - 27.1)
≥ College Degree	*	*	*	12,062	16.4	(14.2 - 18.5)
Income						
≤ \$24,999	*	*	*	9,346	28.3	(22.6 - 34.1)
\$25,000-\$49,999	*	*	*	9,368	21.0	(17.2 - 24.8)
\$50,000-\$74,999	74	16.5	(7.7 - 25.3)	5,615	24.0	(19.4 - 28.7)
≥ \$75,000	*	*	*	8,337	19.3	(16.7 - 21.9)

¹ Defined as having a body mass index (BMI) of 30.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 64a. Overweight or obesity among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	625	16,300	66.1	(60.8 - 71.4)	36,710	8,540,500	59.6	(57.0 - 62.2)
Gender								
Male	243	9,000	71.3	(63.5 - 79.1)	14,502	4,579,100	65.2	(61.0 - 69.4)
Female	382	7,300	60.7	(53.7 - 67.7)	22,208	3,961,400	54.3	(51.0 - 57.6)
Age (years)								
18-34	*	*	*	*	3,789	2,025,800	47.1	(40.8 - 53.5)
35-44	*	*	*	*	4,785	1,714,200	61.1	(55.6 - 66.6)
45-54	137	3,000	64.8	(56.0 - 73.7)	7,495	1,870,100	68.8	(64.6 - 73.1)
55-64	159	2,500	64.3	(56.0 - 72.5)	8,356	1,362,000	66.2	(61.9 - 70.4)
≥ 65	214	3,400	71.5	(65.0 - 78.1)	12,285	1,568,400	64.1	(60.6 - 67.7)
Race/Ethnicity								
White Non-Hispanic	610	15,600	65.4	(60.0 - 70.8)	34,501	5,344,900	59.3	(57.2 - 61.3)
Black Non-Hispanic	*	*	*	*	708	1,443,100	71.6	(62.5 - 80.7)
Hispanic	*	*	*	*	813	1,237,900	56.4	(46.5 - 66.2)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	294	8,100	70.1	(63.2 - 77.0)	15,079	3,451,400	64.6	(59.8 - 69.4)
Some College	*	*	*	*	9,515	2,107,300	62.1	(57.0 - 67.2)
≥ College Degree	163	3,900	67.2	(58.2 - 76.1)	12,062	2,911,900	52.9	(49.2 - 56.6)
Income								
≤ \$24,999	*	*	*	*	9,346	2,145,700	62.3	(55.7 - 68.9)
\$25,000-\$49,999	195	5,700	67.7	(58.2 - 77.2)	9,368	1,720,700	61.2	(54.5 - 67.9)
\$50,000-\$74,999	*	*	*	*	5,615	1,214,000	62.9	(57.1 - 68.8)
≥ \$75,000	*	*	*	*	8,337	2,526,700	57.1	(53.6 - 60.6)

¹ Defined as having a body mass index (BMI) of 25.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Sample size from survey

³ Estimated number of adults who are overweight or obese (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 64b. Overweight or obesity among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	625	66.0	(60.3 - 71.7)	36,710	59.3	(56.8 - 61.9)
Gender						
Male	243	71.6	(63.2 - 79.9)	14,502	65.4	(61.3 - 69.4)
Female	382	60.2	(52.7 - 67.7)	22,208	53.5	(50.1 - 56.8)
Race/Ethnicity						
White Non-Hispanic	610	65.2	(59.5 - 71.0)	34,501	58.8	(56.7 - 60.8)
Black Non-Hispanic	*	*	*	708	71.8	(63.0 - 80.6)
Hispanic	*	*	*	813	56.5	(46.8 - 66.1)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	294	70.8	(63.5 - 78.0)	15,079	64.6	(59.9 - 69.3)
Some College	*	*	*	9,515	61.4	(56.4 - 66.5)
≥ College Degree	163	67.8	(58.4 - 77.2)	12,062	52.6	(48.9 - 56.3)
Income						
≤ \$24,999	*	*	*	9,346	62.1	(55.6 - 68.6)
\$25,000-\$49,999	*	*	*	9,368	61.6	(55.1 - 68.1)
\$50,000-\$74,999	*	*	*	5,615	62.6	(56.7 - 68.6)
≥ \$75,000	*	*	*	8,337	56.5	(52.9 - 60.0)

¹ Defined as having a body mass index (BMI) of 25.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 65a. Received advice about weight by a health professional among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	627	5,800	23.5	(19.1 - 27.9)	36,729	3,929,700	27.4	(25.1 - 29.7)
Gender								
Male	243	2,700	21.6	(15.3 - 27.9)	14,499	1,654,800	23.6	(20.6 - 26.6)
Female	384	3,100	25.5	(19.5 - 31.5)	22,230	2,274,900	31.0	(27.7 - 34.3)
Age (years)								
18-34	*	*	*	*	3,796	895,500	20.7	(15.3 - 26.2)
35-44	*	*	*	*	4,779	722,700	25.8	(20.9 - 30.6)
45-54	136	1,200	25.5	(16.5 - 34.4)	7,490	910,900	33.5	(29.0 - 38.1)
55-64	159	1,400	35.4	(27.3 - 43.6)	8,351	741,100	36.1	(32.0 - 40.3)
≥ 65	217	1,500	31.1	(23.9 - 38.3)	12,313	659,500	26.9	(23.0 - 30.7)
Race/Ethnicity								
White Non-Hispanic	612	5,500	23.2	(18.8 - 27.6)	34,502	2,386,200	26.5	(24.6 - 28.4)
Black Non-Hispanic	*	*	*	*	709	615,700	30.4	(22.0 - 38.8)
Hispanic	*	*	*	*	830	555,400	25.1	(17.8 - 32.4)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	297	2,500	21.4	(16.0 - 26.9)	15,116	1,469,700	27.5	(23.6 - 31.4)
Some College	168	1,500	20.2	(12.9 - 27.5)	9,515	946,800	27.8	(23.2 - 32.4)
≥ College Degree	*	*	*	*	12,041	1,445,500	26.2	(23.0 - 29.4)
Income								
≤ \$24,999	171	1,100	20.0	(13.1 - 26.9)	9,360	1,079,400	31.4	(25.3 - 37.5)
\$25,000-\$49,999	195	2,000	23.4	(15.6 - 31.2)	9,365	696,900	24.6	(19.8 - 29.4)
\$50,000-\$74,999	*	*	*	*	5,603	507,100	26.3	(21.6 - 31.0)
≥ \$75,000	*	*	*	*	8,334	1,207,100	27.3	(24.2 - 30.5)

¹ Defined as being advised by a doctor, nurse or other health professional about their weight (lose, gain or maintain)

² Sample size from survey

³ Estimated number of adults provided advice by a health professional about their weight (lose, gain or maintain) (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 65b. Received advice about weight by a health professional among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	627	22.8	(18.2 - 27.4)	36,729	27.1	(24.8 - 29.4)
Gender						
Male	243	20.6	(14.1 - 27.1)	14,499	23.4	(20.5 - 26.3)
Female	384	25.1	(18.6 - 31.6)	22,230	30.6	(27.3 - 34.0)
Race/Ethnicity						
White Non-Hispanic	612	22.6	(17.9 - 27.2)	34,502	26.1	(24.2 - 28.0)
Black Non-Hispanic	*	*	*	709	30.4	(21.9 - 38.9)
Hispanic	*	*	*	830	24.9	(17.7 - 32.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	297	20.6	(14.9 - 26.3)	15,116	27.3	(23.4 - 31.2)
Some College	168	18.9	(11.6 - 26.2)	9,515	27.4	(22.9 - 32.0)
≥ College Degree	*	*	*	12,041	25.9	(22.7 - 29.0)
Income						
≤ \$24,999	171	18.6	(11.6 - 25.7)	9,360	31.0	(24.9 - 37.0)
\$25,000-\$49,999	195	22.8	(14.7 - 30.9)	9,365	24.9	(20.0 - 29.8)
\$50,000-\$74,999	*	*	*	5,603	25.9	(21.2 - 30.6)
≥ \$75,000	*	*	*	8,334	26.9	(23.7 - 30.0)

¹ Defined as being advised by a doctor, nurse or other health professional about their weight (lose, gain or maintain)

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 66a. Received advice to lose weight by a health professional among adults provided weight advice¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	169	4,800	82.1	(72.3 - 91.9)	9,724	3,412,900	86.8	(83.9 - 89.8)
Gender								
Male	64	2,400	88.8	(78.9 - 98.7)	3,807	1,468,700	88.8	(85.7 - 91.8)
Female	*	*	*	*	5,917	1,944,200	85.5	(80.8 - 90.1)
Age (years)								
18-34	*	*	*	*	681	779,700	87.1	(80.6 - 93.6)
35-44	*	*	*	*	1,215	622,100	86.1	(76.9 - 95.2)
45-54	*	*	*	*	2,238	813,300	89.3	(84.4 - 94.2)
55-64	*	*	*	*	2,697	645,500	87.1	(81.2 - 93.0)
≥ 65	59	1,400	93.0	(87.2 - 98.8)	2,893	552,300	83.7	(76.9 - 90.6)
Race/Ethnicity								
White Non-Hispanic	*	*	*	*	9,082	2,076,700	87.0	(84.5 - 89.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	251	483,600	87.1	(78.5 - 95.7)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	82	2,200	87.5	(79.8 - 95.2)	3,962	1,239,700	84.3	(78.6 - 90.1)
Some College	*	*	*	*	2,633	814,800	86.1	(80.1 - 92.0)
≥ College Degree	*	*	*	*	3,115	1,290,900	89.3	(85.3 - 93.3)
Income								
≤ \$24,999	*	*	*	*	2,539	922,100	85.4	(78.7 - 92.1)
\$25,000-\$49,999	*	*	*	*	2,482	594,100	85.2	(78.4 - 92.1)
\$50,000-\$74,999	*	*	*	*	1,521	446,100	88.0	(83.2 - 92.7)
≥ \$75,000	*	*	*	*	2,270	1,073,200	88.9	(83.7 - 94.1)

¹ Defined as provided advice by a health professional about their weight and advised to lose weight

² Sample size from survey

³ Estimated number of adults provided advice by a health professional about their weight and advised to lose weight (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 66b. Received advice to lose weight by a health professional among adults provided weight advice¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	*	*	*	9,724	86.9	(83.8 - 89.9)
Gender						
Male	*	*	*	3,807	88.7	(85.7 - 91.8)
Female	*	*	*	5,917	85.5	(80.7 - 90.3)
Race/Ethnicity						
White Non-Hispanic	*	*	*	9,082	87.0	(84.4 - 89.6)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	251	87.6	(79.3 - 95.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	82	88.3	(80.0 - 96.5)	3,962	84.6	(78.7 - 90.5)
Some College	*	*	*	2,633	85.6	(79.5 - 91.7)
≥ College Degree	*	*	*	3,115	89.4	(85.4 - 93.3)
Income						
≤ \$24,999	*	*	*	2,539	85.7	(79.0 - 92.3)
\$25,000-\$49,999	*	*	*	2,482	85.3	(78.4 - 92.3)
\$50,000-\$74,999	*	*	*	1,521	88.0	(83.1 - 92.8)
≥ \$75,000	*	*	*	2,270	88.7	(83.1 - 94.2)

¹ Defined as provided advice by a health professional about their weight and advised to lose weight

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 67a. No leisure-time physical activity among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	647	5,000	19.6	(15.5 - 23.6)	38,128	3,541,700	23.9	(21.6 - 26.1)
Gender								
Male	244	2,600	20.5	(13.7 - 27.3)	14,602	1,482,500	20.9	(17.8 - 24.0)
Female	403	2,400	18.6	(14.1 - 23.2)	23,526	2,059,200	26.6	(23.4 - 29.7)
Age (years)								
18-34	*	*	*	*	3,938	982,900	22.0	(16.6 - 27.5)
35-44	*	*	*	*	4,988	556,000	19.3	(15.0 - 23.6)
45-54	141	500	11.5	(5.9 - 17.0)	7,827	759,000	26.8	(21.7 - 31.9)
55-64	164	1,100	27.5	(19.8 - 35.3)	8,765	464,900	21.7	(18.2 - 25.2)
≥ 65	219	1,600	32.3	(25.4 - 39.2)	12,610	778,800	30.7	(27.3 - 34.2)
Race/Ethnicity								
White Non-Hispanic	632	4,900	19.7	(15.6 - 23.9)	35,816	1,892,800	20.3	(18.6 - 22.0)
Black Non-Hispanic	*	*	*	*	739	690,000	32.0	(23.4 - 40.6)
Hispanic	*	*	*	*	860	600,800	26.7	(19.1 - 34.4)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	305	3,600	29.9	(22.7 - 37.1)	15,683	1,810,200	32.5	(28.4 - 36.5)
Some College	175	900	11.8	(6.9 - 16.6)	9,906	814,200	23.0	(18.6 - 27.3)
≥ College Degree	166	500	8.2	(4.3 - 12.1)	12,458	841,000	14.9	(11.9 - 18.0)
Income								
≤ \$24,999	*	*	*	*	9,640	1,332,200	38.0	(31.7 - 44.3)
\$25,000-\$49,999	201	1,400	16.2	(10.4 - 22.0)	9,658	648,000	22.3	(18.3 - 26.3)
\$50,000-\$74,999	76	500	15.3	(7.3 - 23.2)	5,773	430,700	21.8	(16.6 - 27.0)
≥ \$75,000	118	800	14.2	(6.8 - 21.6)	8,565	560,800	12.5	(10.3 - 14.6)

¹ Defined as having not participated in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise

² Sample size from survey

³ Estimated number of adults who have not participated in any physical activities or exercises (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 67b. No leisure-time physical activity among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	647	19.0	(14.7 - 23.4)	38,128	23.7	(21.5 - 25.9)
Gender						
Male	244	20.2	(13.0 - 27.4)	14,602	20.9	(17.8 - 23.9)
Female	403	17.9	(13.0 - 22.8)	23,526	26.3	(23.1 - 29.5)
Race/Ethnicity						
White Non-Hispanic	632	19.1	(14.7 - 23.6)	35,816	20.0	(18.3 - 21.7)
Black Non-Hispanic	*	*	*	739	32.0	(23.3 - 40.6)
Hispanic	*	*	*	860	26.9	(19.2 - 34.5)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	305	29.5	(21.7 - 37.3)	15,683	32.3	(28.3 - 36.4)
Some College	175	11.2	(6.3 - 16.1)	9,906	22.8	(18.5 - 27.2)
≥ College Degree	166	7.3	(3.6 - 11.1)	12,458	14.9	(11.8 - 17.9)
Income						
≤ \$24,999	*	*	*	9,640	37.9	(31.6 - 44.2)
\$25,000-\$49,999	201	15.8	(9.6 - 22.0)	9,658	22.0	(18.0 - 26.0)
\$50,000-\$74,999	76	14.0	(6.3 - 21.8)	5,773	21.6	(16.3 - 26.8)
≥ \$75,000	118	14.0	(6.1 - 21.9)	8,565	12.6	(10.3 - 14.9)

¹ Defined as having not participated in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 68a. Consumption of 5 or more servings of fruits and vegetables per day among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	648	6,500	25.4	(20.9 - 29.9)	38,203	4,023,400	27.0	(24.9 - 29.2)
Gender								
Male	245	2,500	20.0	(13.2 - 26.9)	14,634	1,603,900	22.6	(19.4 - 25.8)
Female	403	4,000	30.7	(24.8 - 36.6)	23,569	2,419,500	31.1	(28.2 - 34.0)
Age (years)								
18-34	*	*	*	*	3,940	1,261,700	28.3	(22.7 - 33.9)
35-44	*	*	*	*	4,989	688,000	23.9	(19.5 - 28.3)
45-54	142	1,200	25.1	(17.2 - 33.1)	7,837	746,800	26.2	(22.3 - 30.1)
55-64	164	1,000	26.2	(18.7 - 33.6)	8,777	579,100	27.0	(23.6 - 30.5)
≥ 65	219	1,600	33.7	(26.7 - 40.7)	12,660	747,800	29.4	(26.0 - 32.8)
Race/Ethnicity								
White Non-Hispanic	633	6,200	24.8	(20.3 - 29.3)	35,884	2,538,000	27.2	(25.4 - 28.9)
Black Non-Hispanic	*	*	*	*	740	572,400	26.5	(19.2 - 33.9)
Hispanic	*	*	*	*	864	672,200	29.8	(21.0 - 38.5)
Other Non-Hispanic	*	*	*	*	715	240,800	21.4	(12.0 - 30.8)
Education								
≤ High School	305	2,800	23.3	(16.7 - 29.8)	15,726	1,309,800	23.4	(19.3 - 27.4)
Some College	175	2,000	26.7	(17.4 - 36.0)	9,924	964,300	27.2	(23.0 - 31.3)
≥ College Degree	167	1,700	28.4	(20.4 - 36.4)	12,471	1,747,500	31.0	(27.9 - 34.1)
Income								
≤ \$24,999	172	1,400	25.2	(15.3 - 35.2)	9,667	831,500	23.7	(18.6 - 28.7)
\$25,000-\$49,999	201	2,400	27.7	(19.3 - 36.1)	9,669	917,200	31.6	(25.6 - 37.6)
\$50,000-\$74,999	76	400	13.5	(5.7 - 21.3)	5,777	501,000	25.3	(21.4 - 29.3)
≥ \$75,000	119	1,500	27.8	(17.8 - 37.8)	8,569	1,260,100	28.0	(25.0 - 31.0)

¹ Defined as having consumed 5 or more servings of fruits and vegetables in an average day

² Sample size from survey

³ Estimated number of adults who consume 5 or more servings of fruits and vegetables per day (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 68b. Consumption of 5 or more servings of fruits and vegetables per day among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	648	25.2	(20.4 - 30.0)	38,203	27.1	(24.9 - 29.2)
Gender						
Male	245	20.1	(12.8 - 27.5)	14,634	22.7	(19.6 - 25.9)
Female	403	30.1	(23.8 - 36.4)	23,569	31.0	(28.1 - 34.0)
Race/Ethnicity						
White Non-Hispanic	633	24.5	(19.7 - 29.3)	35,884	27.1	(25.4 - 28.9)
Black Non-Hispanic	*	*	*	740	26.8	(19.4 - 34.2)
Hispanic	*	*	*	864	29.5	(20.9 - 38.1)
Other Non-Hispanic	*	*	*	715	21.8	(12.2 - 31.5)
Education						
≤ High School	305	23.3	(16.3 - 30.3)	15,726	23.3	(19.3 - 27.3)
Some College	175	26.6	(16.7 - 36.5)	9,924	27.1	(22.9 - 31.2)
≥ College Degree	167	27.4	(19.1 - 35.7)	12,471	31.2	(28.0 - 34.3)
Income						
≤ \$24,999	*	*	*	9,667	23.9	(18.8 - 28.9)
\$25,000-\$49,999	201	28.0	(19.0 - 36.9)	9,669	32.0	(26.0 - 38.0)
\$50,000-\$74,999	76	13.7	(5.3 - 22.1)	5,777	25.0	(21.0 - 28.9)
≥ \$75,000	*	*	*	8,569	28.0	(24.9 - 31.1)

¹ Defined as having consumed 5 or more servings of fruits and vegetables in an average day

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 69a. Had a fall within the past 3 months among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	517	2,500	18.6	(14.8 - 22.4)	28,783	1,244,800	17.0	(14.9 - 19.0)
Gender								
Male	199	1,100	17.5	(11.7 - 23.4)	10,966	489,400	14.4	(11.9 - 16.8)
Female	318	1,400	19.6	(14.7 - 24.5)	17,817	755,300	19.2	(16.1 - 22.3)
Age (years)								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	139	900	19.9	(12.7 - 27.2)	7,741	445,500	16.0	(12.0 - 20.0)
55-64	160	500	13.2	(7.9 - 18.6)	8,454	336,000	16.5	(13.4 - 19.6)
≥ 65	218	1,100	21.7	(15.3 - 28.1)	12,588	463,300	18.4	(15.2 - 21.6)
Race/Ethnicity								
White Non-Hispanic	507	2,400	18.2	(14.4 - 22.0)	27,456	869,000	16.6	(15.1 - 18.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	424	76,100	11.0	(4.5 - 17.4)
Other Non-Hispanic	*	*	*	*	445	44,300	10.0	(3.1 - 16.8)
Education								
≤ High School	244	1,300	20.6	(14.7 - 26.5)	12,417	493,000	17.3	(13.3 - 21.3)
Some College	135	500	13.9	(7.4 - 20.3)	7,232	299,400	18.7	(14.6 - 22.7)
≥ College Degree	137	700	19.8	(12.7 - 27.0)	9,086	441,500	15.4	(12.9 - 17.9)
Income								
≤ \$24,999	149	600	19.7	(12.5 - 27.0)	7,824	423,900	25.0	(18.5 - 31.6)
\$25,000-\$49,999	153	700	17.4	(10.5 - 24.3)	7,445	223,600	16.0	(12.8 - 19.2)
\$50,000-\$74,999	*	*	*	*	4,141	151,200	14.6	(10.8 - 18.4)
≥ \$75,000	92	500	18.1	(9.0 - 27.1)	5,874	287,100	12.5	(10.0 - 15.1)

¹ Defined as respondent unintentionally coming to rest on the ground or another lower level one or more times within the past 3 months

² Sample size from survey

³ Estimated number of adults who have had a fall within the past 3 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 69b. Had a fall within the past 3 months among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	517	19.1	(15.1 - 23.0)	28,783	17.3	(15.0 - 19.5)
Gender						
Male	199	18.1	(12.0 - 24.1)	10,966	15.0	(12.1 - 17.8)
Female	318	20.1	(15.0 - 25.1)	17,817	19.4	(16.0 - 22.8)
Race/Ethnicity						
White Non-Hispanic	507	18.7	(14.7 - 22.7)	27,456	16.9	(15.2 - 18.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	424	10.7	(4.2 - 17.1)
Other Non-Hispanic	*	*	*	445	10.0	(2.9 - 17.0)
Education						
≤ High School	244	21.0	(14.8 - 27.1)	12,417	17.4	(13.0 - 21.7)
Some College	135	14.4	(7.6 - 21.2)	7,232	19.0	(14.7 - 23.2)
≥ College Degree	137	20.3	(12.9 - 27.8)	9,086	16.0	(13.1 - 18.8)
Income						
≤ \$24,999	149	20.1	(12.6 - 27.6)	7,824	25.3	(18.3 - 32.4)
\$25,000-\$49,999	153	17.9	(10.7 - 25.1)	7,445	15.9	(12.6 - 19.2)
\$50,000-\$74,999	*	*	*	4,141	14.7	(10.6 - 18.8)
≥ \$75,000	92	19.0	(9.4 - 28.6)	5,874	12.9	(9.9 - 15.8)

¹ Defined as respondent unintentionally coming to rest on the ground or another lower level one or more times within the past 3 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Schoharie County**

Table 70a. Had a fall that caused injury within the past 3 months among adults¹ in Schoharie County and New York State

	Schoharie County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	517	500	3.4	(1.8 - 4.9)	28,760	335,700	4.6	(3.7 - 5.5)
Gender								
Male	*	*	*	*	10,957	120,000	3.5	(2.5 - 4.6)
Female	318	300	4.8	(2.4 - 7.3)	17,803	215,600	5.5	(4.1 - 6.9)
Age (years)								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	7,737	115,100	4.1	(2.9 - 5.4)
55-64	*	*	*	*	8,449	90,300	4.4	(2.7 - 6.1)
≥ 65	*	*	*	*	12,574	130,300	5.2	(3.4 - 6.9)
Race/Ethnicity								
White Non-Hispanic	507	400	2.9	(1.4 - 4.3)	27,433	270,800	5.2	(4.2 - 6.1)
Black Non-Hispanic	*	*	*	*	458	47,900	4.9	(0.9 - 9.0)
Hispanic	*	*	*	*	424	12,400	1.8	(0.2 - 3.3)
Other Non-Hispanic	*	*	*	*	445	4,600	1.0	(0.4 - 1.6)
Education								
≤ High School	244	200	3.4	(1.2 - 5.7)	12,404	148,900	5.2	(3.6 - 6.8)
Some College	*	*	*	*	7,228	85,400	5.3	(3.1 - 7.5)
≥ College Degree	*	*	*	*	9,081	101,300	3.5	(2.4 - 4.6)
Income								
≤ \$24,999	*	*	*	*	7,812	90,200	5.3	(3.6 - 7.1)
\$25,000-\$49,999	*	*	*	*	7,442	93,200	6.7	(3.9 - 9.4)
\$50,000-\$74,999	*	*	*	*	4,139	37,000	3.6	(1.8 - 5.3)
≥ \$75,000	*	*	*	*	5,872	54,700	2.4	(1.5 - 3.3)

¹ Defined as as respondent unintentionally coming to rest on the ground or another lower level and caused respondent to limit regular activities for at least a day or to go see a doctor one or more times within the past 3 months

² Sample size from survey

³ Estimated number of adults who have had a fall that caused an injury within the past 3 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Schoharie County

Table 70b. Had a fall that caused injury within the past 3 months among adults¹ in Schoharie County and New York State - Age-Adjusted²

	Schoharie County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	517	3.2	(1.7 - 4.7)	28,760	4.6	(3.6 - 5.5)
Gender						
Male	*	*	*	10,957	3.6	(2.4 - 4.7)
Female	318	4.7	(2.3 - 7.1)	17,803	5.5	(4.1 - 6.9)
Race/Ethnicity						
White Non-Hispanic	507	2.7	(1.4 - 4.1)	27,433	5.2	(4.2 - 6.2)
Black Non-Hispanic	*	*	*	458	4.7	(0.7 - 8.7)
Hispanic	*	*	*	424	1.8	(0.1 - 3.4)
Other Non-Hispanic	*	*	*	445	1.0	(0.4 - 1.6)
Education						
≤ High School	244	3.1	(1.1 - 5.2)	12,404	5.2	(3.5 - 6.8)
Some College	*	*	*	7,228	5.3	(3.1 - 7.5)
≥ College Degree	*	*	*	9,081	3.6	(2.4 - 4.7)
Income						
≤ \$24,999	*	*	*	7,812	5.1	(3.3 - 7.0)
\$25,000-\$49,999	*	*	*	7,442	6.6	(3.9 - 9.2)
\$50,000-\$74,999	*	*	*	4,139	3.7	(1.7 - 5.8)
≥ \$75,000	*	*	*	5,872	2.4	(1.4 - 3.3)

¹ Defined as as respondent unintentionally coming to rest on the ground or another lower level and caused respondent to limit regular activities for at least a day or to go see a doctor one or more times within the past 3 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)