

New York State Expanded Behavioral Risk Factor Surveillance System

Wayne County

**Final Report
July 2008 - June 2009**

**New York State Department of Health
December 2009**

Contents

Introduction	12
Table 1. Wayne County and New York State adult US Census Bureau population estimates, 2007	13
General Health	
Table 2a. Fair or poor general health among adults in Wayne County and New York State	14
Table 2b. Fair or poor general health among adults in Wayne County and New York State - Age-adjusted	15
Table 3a. Poor physical health 14 or more days within the past month among adults in Wayne County and New York State	16
Table 3b. Poor physical health 14 or more days within the past month among adults in Wayne County and New York State - Age-adjusted	17
Table 4a. Poor mental health 14 or more days within the past month among adults in Wayne County and New York State	18
Table 4b. Poor mental health 14 or more days within the past month among adults in Wayne County and New York State - Age-adjusted	19
Table 5a. Poor physical or mental health 14 or more days within the past month among adults in Wayne County and New York State	20
Table 5b. Poor physical or mental health 14 or more days within the past month among adults in Wayne County and New York State - Age-adjusted	21
Quality Health Care	
Table 6a. Health insurance among adults in Wayne County and New York State	22
Table 6b. Health insurance among adults in Wayne County and New York State - Age-adjusted	23
Table 7a. Health insurance among adults aged 18-64 in Wayne County and New York State	24
Table 7b. Health insurance among adults aged 18-64 in Wayne County and New York State - Age-adjusted	25

Contents

Table 8a. No health insurance among adults in Wayne County and New York State	26
Table 8b. No health insurance among adults in Wayne County and New York State - Age-adjusted	27
Table 9a. No health insurance among adults aged 18-64 in Wayne County and New York State	28
Table 9b. No health insurance among adults aged 18-64 in Wayne County and New York State - Age-adjusted	29
Table 10a. Adults with regular health care provider in Wayne County and New York State	30
Table 10b. Adults with regular health care provider in Wayne County and New York State - Age-adjusted	31
Table 11a. Cost prevented visit to doctor within the past year among adults in Wayne County and New York State	32
Table 11b. Cost prevented visit to doctor within the past year among adults in Wayne County and New York State - Age-adjusted	33
Table 12a. Visited a doctor for routine checkup within the past year among adults in Wayne County and New York State	34
Table 12b. Visited a doctor for routine checkup within the past year among adults in Wayne County and New York State - Age-adjusted	35
Table 13a. Visited a doctor for routine checkup within the past 2 years among adults in Wayne County and New York State	36
Table 13b. Visited a doctor for routine checkup within the past 2 years among adults in Wayne County and New York State - Age-adjusted	37
Table 14a. Dentist visit within the past year among adults in Wayne County and New York State	38
Table 14b. Dentist visit within the past year among adults in Wayne County and New York State - Age-adjusted	39
Table 15a. Had teeth cleaned within the past year among adults in Wayne County and New York State	40
Table 15b. Had teeth cleaned within the past year among adults in Wayne County and New York State - Age-adjusted	41

Contents

Table 16a. Permanent teeth extracted due to decay or gum disease among adults in Wayne County and New York State	42
Table 16b. Permanent teeth extracted due to decay or gum disease among adults in Wayne County and New York State - Age-adjusted	43
Table 17a. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in Wayne County and New York State	44
Table 17b. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older in Wayne County and New York State - Age-adjusted	45
Table 18a. Flu shot within the past 12 months in Wayne County and New York State	46
Table 18b. Flu shot within the past 12 months in Wayne County and New York State - Age-adjusted	47
Table 19a. Flu vaccine that was sprayed in nose within the past 12 months in Wayne County and New York State	48
Table 19b. Flu vaccine that was sprayed in nose within the past 12 months in Wayne County and New York State - Age-adjusted	49
Table 20a. Flu shot or flu vaccine sprayed in nose within the past 12 months in Wayne County and New York State	50
Table 20b. Flu shot or flu vaccine sprayed in nose within the past 12 months in Wayne County and New York State - Age-adjusted	51
Table 21a. Flu shot among adults aged 65 and older in Wayne County and New York State	52
Table 21b. Flu shot among adults aged 65 and older in Wayne County and New York State - Age-adjusted	53
Table 22a. Pneumonia shot or pneumococcal vaccine in Wayne County and New York State	54
Table 22b. Pneumonia shot or pneumococcal vaccine in Wayne County and New York State - Age-adjusted	55
Table 23a. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in Wayne County and New York State	56
Table 23b. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older in Wayne County and New York State - Age-adjusted	57

Contents

Cancer Screening

Table 24a. Ever had a mammogram among women aged 40 and older in Wayne County and New York State	58
Table 24b. Ever had a mammogram among women aged 40 and older in Wayne County and New York State - Age-adjusted	59
Table 25a. Had mammogram within the past 2 years among women aged 40 and older in Wayne County and New York State	60
Table 25b. Had mammogram within the past 2 years among women aged 40 and older in Wayne County and New York State - Age-adjusted	61
Table 26a. Had mammogram within the past 2 years among women aged 50 and older in Wayne County and New York State	62
Table 26b. Had mammogram within the past 2 years among women aged 50 and older in Wayne County and New York State - Age-adjusted	63
Table 27a. Ever had a Pap test among women in Wayne County and New York State	64
Table 27b. Ever had a Pap test among women in Wayne County and New York State - Age-adjusted	65
Table 28a. Had a Pap test within the past 3 years among women in Wayne County and New York State	66
Table 28b. Had a Pap test within the past 3 years among women in Wayne County and New York State - Age-adjusted	67
Table 29a. Ever had a digital rectal exam among men aged 40 and older in Wayne County and New York State	68
Table 29b. Ever had a digital rectal exam among men aged 40 and older in Wayne County and New York State - Age-adjusted	69
Table 30a. Had digital rectal exam within the past 2 years among men aged 40 and older in Wayne County and New York State	70
Table 30b. Had digital rectal exam within the past 2 years among men aged 40 and older in Wayne County and New York State - Age-adjusted	71

Contents

Table 31a. Ever had a prostate specific antigen test among men aged 40 and older in Wayne County and New York State	72
Table 31b. Ever had a prostate specific antigen test among men aged 40 and older in Wayne County and New York State - Age-adjusted	73
Table 32a. Had prostate specific antigen test within the past 2 years among men aged 40 and older in Wayne County and New York State	74
Table 32b. Had prostate specific antigen test within the past 2 years among men aged 40 and older in Wayne County and New York State - Age-adjusted	75
Table 33a. Prostate cancer among men aged 40 and older in Wayne County and New York State	76
Table 33b. Prostate cancer among men aged 40 and older in Wayne County and New York State - Age-adjusted	77
Table 34a. Home blood stool test ever used among adults aged 50 and older in Wayne County and New York State	78
Table 34b. Home blood stool test ever used among adults aged 50 and older in Wayne County and New York State - Age-adjusted	79
Table 35a. Home blood stool test used within the past year among adults aged 50 and older in Wayne County and New York State	80
Table 35b. Home blood stool test used within the past year among adults aged 50 and older in Wayne County and New York State - Age-adjusted	81
Table 36a. Home blood stool test used within the past 2 years among adults aged 50 and older in Wayne County and New York State	82
Table 36b. Home blood stool test used within the past 2 years among adults aged 50 and older in Wayne County and New York State - Age-adjusted	83
Table 37a. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Wayne County and New York State	84
Table 37b. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older in Wayne County and New York State - Age-adjusted	85
Table 38a. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Wayne County and New York State	86

Contents

Table 38b. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Wayne County and New York State - Age-adjusted	87
Table 39a. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Wayne County and New York State	88
Table 39b. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older in Wayne County and New York State - Age-adjusted	89
Chronic Diseases	
Table 40a. High blood pressure among adults in Wayne County and New York State	90
Table 40b. High blood pressure among adults in Wayne County and New York State - Age-adjusted	91
Table 41a. Blood pressure medication among adults with high blood pressure in Wayne County and New York State	92
Table 41b. Blood pressure medication among adults with high blood pressure in Wayne County and New York State - Age-adjusted	93
Table 42a. Ever had blood cholesterol checked among adults in Wayne County and New York State	94
Table 42b. Ever had blood cholesterol checked among adults in Wayne County and New York State - Age-adjusted	95
Table 43a. Blood cholesterol checked within the past 5 years among adults in Wayne County and New York State	96
Table 43b. Blood cholesterol checked within the past 5 years among adults in Wayne County and New York State - Age-adjusted	97
Table 44a. Cardiovascular disease among adults in Wayne County and New York State	98
Table 44b. Cardiovascular disease among adults in Wayne County and New York State - Age-adjusted	99
Table 45a. Coronary heart disease among adults in Wayne County and New York State	100

Contents

Table 45b. Coronary heart disease among adults in Wayne County and New York State - Age-adjusted	101
Table 46a. Ever diagnosed with diabetes among adults in Wayne County and New York State	102
Table 46b. Ever diagnosed with diabetes among adults in Wayne County and New York State - Age-adjusted	103
Table 47a. Ever diagnosed with asthma (lifetime) among adults in Wayne County and New York State	104
Table 47b. Ever diagnosed with asthma (lifetime) among adults in Wayne County and New York State - Age-adjusted	105
Table 48a. Current asthma among adults in Wayne County and New York State	106
Table 48b. Current asthma among adults in Wayne County and New York State - Age-adjusted	107
Table 49a. Chronic joint symptoms among adults in Wayne County and New York State	108
Table 49b. Chronic joint symptoms among adults in Wayne County and New York State - Age-adjusted	109
Table 50a. Arthritis among adults in Wayne County and New York State	110
Table 50b. Arthritis among adults in Wayne County and New York State - Age-adjusted	111
Table 51a. Activity limitations because of physical, mental or emotional problems among adults in Wayne County and New York State	112
Table 51b. Activity limitations because of physical, mental or emotional problems among adults in Wayne County and New York State - Age-adjusted	113
Table 52a. Have health problem that requires the use of special equipment among adults in Wayne County and New York State	114
Table 52b. Have health problem that requires the use of special equipment among adults in Wayne County and New York State - Age-adjusted	115
Table 53a. Disability among adults in Wayne County and New York State	116
Table 53b. Disability among adults in Wayne County and New York State - Age-adjusted	117

Contents

Health Behaviors

Table 54a. Current smoking among adults in Wayne County and New York State	118
Table 54b. Current smoking among adults in Wayne County and New York State - Age-adjusted	119
Table 55a. Everyday smoking among adults in Wayne County and New York State	120
Table 55b. Everyday smoking among adults in Wayne County and New York State - Age-adjusted	121
Table 56a. Adults living in homes in which smoking is prohibited in Wayne County and New York State	122
Table 56b. Adults living in homes in which smoking is prohibited in Wayne County and New York State - Age-adjusted	123
Table 57a. Binge drinking within the past month among adults in Wayne County and New York State	124
Table 57b. Binge drinking within the past month among adults in Wayne County and New York State - Age-adjusted	125
Table 58a. Heavy drinking within the past month among adults in Wayne County and New York State	126
Table 58b. Heavy drinking within the past month among adults in Wayne County and New York State - Age-adjusted	127
Table 59a. Never or rarely been asked about sexual history during routine check-up among adults in Wayne County and New York State	128
Table 59b. Never or rarely been asked about sexual history during routine check-up among adults in Wayne County and New York State - Age-adjusted	129
Table 60a. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults in Wayne County and New York State	130
Table 60b. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults in Wayne County and New York State - Age-adjusted	131

Contents

Table 61a. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults in Wayne County and New York State	132
Table 61b. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults in Wayne County and New York State - Age-adjusted	133

Overweight / Obesity

Table 62a. Overweight among adults in Wayne County and New York State	134
Table 62b. Overweight among adults in Wayne County and New York State - Age-adjusted	135
Table 63a. Obesity among adults in Wayne County and New York State	136
Table 63b. Obesity among adults in Wayne County and New York State - Age-adjusted	137
Table 64a. Overweight or obesity among adults in Wayne County and New York State	138
Table 64b. Overweight or obesity among adults in Wayne County and New York State - Age-adjusted	139
Table 65a. Received advice about weight by a health professional among adults in Wayne County and New York State	140
Table 65b. Received advice about weight by a health professional among adults in Wayne County and New York State - Age-adjusted	141
Table 66a. Received advice to lose weight by a health professional among adults provided weight advice in Wayne County and New York State	142
Table 66b. Received advice to lose weight by a health professional among adults provided weight advice in Wayne County and New York State - Age-adjusted	143

Physical Activity / Nutrition

Table 67a. No leisure-time physical activity among adults in Wayne County and New York State	144
Table 67b. No leisure-time physical activity among adults in Wayne County and New York State - Age-adjusted	145

Contents

Table 68a. Consumption of 5 or more servings of fruits and vegetables per day among adults in Wayne County and New York State 146

Table 68b. Consumption of 5 or more servings of fruits and vegetables per day among adults in Wayne County and New York State - Age-adjusted 147

Injury

Table 69a. Had a fall within the past 3 months among adults in Wayne County and New York State 148

Table 69b. Had a fall within the past 3 months among adults in Wayne County and New York State - Age-adjusted 149

Table 70a. Had a fall that caused injury within the past 3 months among adults in Wayne County and New York State 150

Table 70b. Had a fall that caused injury within the past 3 months among adults in Wayne County and New York State - Age-adjusted 151

Introduction

The New York State Expanded Behavioral Risk Factor Surveillance System (Expanded BRFSS) is a random digit dialed telephone survey of adults (18 years of age and older) that is representative of the non-institutionalized civilian population with landline telephones living in New York State. The goal of the Expanded BRFSS is to collect county-specific data on preventive health practices, risk behaviors, injuries and preventable chronic and infectious diseases. Topics assessed by the Expanded BRFSS include tobacco use, physical inactivity, diet, use of cancer screening services and other factors linked to the leading causes of morbidity and mortality. For each topic, health indicators are based on answers to one or more questions in the survey. The survey results are weighted to more accurately represent the adult population living in each county. Weighting procedures adjust for the differential selection probabilities by race, age and gender. More detailed information on weighting procedures is available in the technical brief.

Survey data collection occurred in 58 geographic areas of New York State, including each of the 57 counties outside of New York City and New York City (5 counties) as a single area. A standard questionnaire was used in all areas. Data collection for the 2009 Expanded BRFSS took place during the period of July 2008 through June 2009. The goal was to complete about 650 interviews in each area. The final number of interviews varies because of the sampling design of the survey and differential response rates. More detailed information on the sampling design is available in the technical brief.

The New York State Department of Health contracted with RTI International, which is responsible for the data collection, sampling, weighting and creation of the statistical reports. RTI International subcontracted with Clearwater Research Inc. for data collection and additional expertise in sample design and data quality control.

This report presents both crude estimates and estimates that have been age-adjusted to the Year 2000 US Standard Population when they represent the entire adult population of an area. Estimates for specific age groups are not age-adjusted. Age adjustment is a statistical process applied to rates that allows communities with different age structures to be compared. Counties can compare their data with results from New York State as a whole and with other counties. Confidence intervals at the 95% level associated with each health indicator give an estimated range of values which is likely to include the actual prevalence of the indicator. Confidence intervals are similar to margins of error. When the confidence intervals of two estimates of the same indicator from different areas do not overlap, they may be said to be statistically significantly different, i.e., these differences are unlikely related to chance and are considered true differences. If there is any value that is included in both intervals, the two estimates are not statistically significantly different.

The survey results for each of the 69 health indicators included in the survey are typically provided by gender, age, race and ethnicity, education and income in the Expanded BRFSS county reports. The estimates for some of these groupings may be suppressed in geographic areas with low proportions of adult respondents in these groupings.

**Expanded BRFSS Report: July 2008 - June 2009
Wayne County**

Table 1. Wayne County and New York State adult US Census Bureau population estimates,¹ 2007

	Wayne County		New York State	
	Adult Population ¹	Percent	Adult Population ¹	Percent
Total	69,494	100	14,884,315	100
Gender				
Male	33,954	48.9	7,103,791	47.7
Female	35,540	51.1	7,780,524	52.3
Age (years)				
18-34	17,848	25.7	4,497,479	30.2
35-44	13,203	19.0	2,844,115	19.1
45-54	15,075	21.7	2,854,316	19.2
55-64	11,247	16.2	2,142,000	14.4
≥ 65	12,121	17.4	2,546,405	17.1
Race/Ethnicity				
White Non-Hispanic	65,031	93.6	9,343,884	62.8
Black Non-Hispanic	2,100	3.0	2,156,941	14.5
Hispanic	1,778	2.6	2,258,717	15.2
Other Non-Hispanic	585	0.8	1,124,773	7.6

¹ Source: National Center for Health Statistics. Postcensal estimates of the resident population of the United States for July 1, 2000-July 1, 2007, by year, county, age, bridged race, Hispanic origin, and sex (Vintage 2007). Prepared under a collaborative arrangement with the U.S. Census Bureau; released August 7, 2008. Available from: http://www.cdc.gov/nchs/nvss/bridged_race.htm as of November 25, 2009.

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 2a. Fair or poor general health among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	655	9,200	13.2	(10.4 - 16.1)	38,124	2,531,400	17.1	(15.1 - 19.0)
Gender								
Male	236	4,000	11.7	(7.5 - 15.8)	14,599	1,069,700	15.1	(12.4 - 17.8)
Female	419	5,200	14.7	(10.9 - 18.6)	23,525	1,461,700	18.8	(16.1 - 21.6)
Age (years)								
18-34	*	*	*	*	3,940	535,800	12.0	(7.2 - 16.8)
35-44	*	*	*	*	4,985	289,200	10.0	(6.8 - 13.3)
45-54	140	2,000	13.4	(6.8 - 19.9)	7,819	507,800	17.9	(14.0 - 21.8)
55-64	161	2,000	17.6	(11.3 - 23.8)	8,757	424,700	19.9	(16.0 - 23.9)
≥ 65	203	3,100	25.9	(19.3 - 32.6)	12,623	773,900	30.5	(26.8 - 34.2)
Race/Ethnicity								
White Non-Hispanic	631	8,800	13.6	(10.6 - 16.5)	35,811	1,226,600	13.2	(11.7 - 14.6)
Black Non-Hispanic	*	*	*	*	739	457,300	21.3	(14.8 - 27.9)
Hispanic	*	*	*	*	861	684,100	30.4	(22.2 - 38.6)
Other Non-Hispanic	*	*	*	*	713	163,400	14.5	(5.1 - 24.0)
Education								
≤ High School	309	5,800	18.2	(13.5 - 23.0)	15,682	1,466,500	26.3	(22.6 - 30.0)
Some College	166	2,100	11.3	(6.1 - 16.6)	9,904	508,500	14.3	(10.9 - 17.8)
≥ College Degree	178	1,300	7.0	(2.5 - 11.6)	12,457	493,500	8.8	(6.5 - 11.1)
Income								
≤ \$24,999	160	3,200	24.9	(16.5 - 33.3)	9,633	1,269,200	36.3	(30.1 - 42.4)
\$25,000-\$49,999	179	2,500	14.6	(9.0 - 20.2)	9,649	447,600	15.4	(11.9 - 18.9)
\$50,000-\$74,999	107	1,600	11.5	(4.3 - 18.8)	5,769	237,800	12.1	(8.0 - 16.2)
≥ \$75,000	*	*	*	*	8,562	218,300	4.9	(3.6 - 6.1)

¹ Defined as reporting fair or poor general health

² Sample size from survey

³ Estimated number of adults who have fair or poor health (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 2b. Fair or poor general health among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	655	12.4	(9.5 - 15.3)	38,124	16.7	(14.8 - 18.6)
Gender						
Male	236	10.8	(6.7 - 14.9)	14,599	15.3	(12.6 - 18.0)
Female	419	14.0	(10.0 - 18.0)	23,525	18.0	(15.2 - 20.7)
Race/Ethnicity						
White Non-Hispanic	631	12.8	(9.8 - 15.9)	35,811	12.9	(11.4 - 14.4)
Black Non-Hispanic	*	*	*	739	20.8	(14.2 - 27.3)
Hispanic	*	*	*	861	29.9	(21.8 - 37.9)
Other Non-Hispanic	*	*	*	713	13.8	(4.8 - 22.8)
Education						
≤ High School	309	17.2	(12.3 - 22.1)	15,682	25.8	(22.1 - 29.5)
Some College	166	10.3	(5.2 - 15.4)	9,904	14.1	(10.7 - 17.6)
≥ College Degree	178	7.0	(2.2 - 11.8)	12,457	8.5	(6.3 - 10.8)
Income						
≤ \$24,999	160	23.1	(14.4 - 31.9)	9,633	35.9	(29.7 - 42.0)
\$25,000-\$49,999	179	13.6	(7.9 - 19.3)	9,649	15.0	(11.6 - 18.5)
\$50,000-\$74,999	107	12.0	(4.0 - 20.0)	5,769	11.8	(7.7 - 16.0)
≥ \$75,000	*	*	*	8,562	4.7	(3.5 - 6.0)

¹ Defined as reporting fair or poor general health

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 3a. Poor physical health 14 or more days within the past month among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	649	8,500	12.3	(8.7 - 15.9)	37,327	1,472,900	10.1	(8.7 - 11.6)
Gender								
Male	234	3,600	10.6	(4.3 - 17.0)	14,325	604,200	8.7	(7.0 - 10.5)
Female	415	4,900	13.9	(10.2 - 17.6)	23,002	868,600	11.4	(9.2 - 13.6)
Age (years)								
18-34	*	*	*	*	3,891	268,100	6.1	(2.7 - 9.5)
35-44	*	*	*	*	4,948	232,100	8.1	(5.2 - 11.1)
45-54	141	1,700	11.4	(5.6 - 17.2)	7,764	350,900	12.5	(9.4 - 15.6)
55-64	161	1,700	14.9	(9.1 - 20.7)	8,621	271,200	13.0	(10.3 - 15.8)
≥ 65	197	2,200	18.5	(12.6 - 24.3)	12,103	350,600	14.5	(12.1 - 16.9)
Race/Ethnicity								
White Non-Hispanic	625	6,700	10.5	(8.0 - 13.0)	35,073	929,700	10.2	(8.8 - 11.6)
Black Non-Hispanic	*	*	*	*	724	208,200	9.9	(4.8 - 14.9)
Hispanic	*	*	*	*	835	250,300	11.5	(6.5 - 16.4)
Other Non-Hispanic	*	*	*	*	695	84,600	7.6	(1.3 - 13.9)
Education								
≤ High School	308	5,000	15.7	(9.2 - 22.2)	15,178	692,800	12.9	(10.6 - 15.3)
Some College	163	2,500	14.1	(7.6 - 20.6)	9,751	362,500	10.3	(7.4 - 13.2)
≥ College Degree	177	1,000	5.2	(1.9 - 8.4)	12,327	359,300	6.4	(4.8 - 8.1)
Income								
≤ \$24,999	*	*	*	*	9,294	625,600	18.4	(13.6 - 23.3)
\$25,000-\$49,999	177	1,800	10.9	(5.9 - 16.0)	9,495	279,100	9.8	(7.3 - 12.3)
\$50,000-\$74,999	107	1,200	8.7	(3.0 - 14.4)	5,733	186,400	9.5	(5.8 - 13.2)
≥ \$75,000	*	*	*	*	8,510	219,100	4.9	(3.6 - 6.2)

¹ Defined as 14 or more days within the past month when physical health was not good

² Sample size from survey

³ Estimated number of adults with poor physical health 14 or more days within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 3b. Poor physical health 14 or more days within the past month among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	649	11.7	(7.9 - 15.5)	37,327	9.9	(8.4 - 11.4)
Gender						
Male	234	10.2	(3.4 - 17.1)	14,325	8.7	(6.9 - 10.5)
Female	415	13.0	(9.3 - 16.8)	23,002	11.0	(8.7 - 13.2)
Race/Ethnicity						
White Non-Hispanic	625	9.7	(7.2 - 12.2)	35,073	9.9	(8.5 - 11.3)
Black Non-Hispanic	*	*	*	724	9.7	(4.6 - 14.9)
Hispanic	*	*	*	835	11.3	(6.3 - 16.2)
Other Non-Hispanic	*	*	*	695	7.6	(1.2 - 14.0)
Education						
≤ High School	308	15.7	(8.6 - 22.8)	15,178	12.7	(10.2 - 15.1)
Some College	163	12.7	(6.3 - 19.1)	9,751	10.1	(7.2 - 12.9)
≥ College Degree	177	4.4	(1.6 - 7.2)	12,327	6.3	(4.7 - 8.0)
Income						
≤ \$24,999	*	*	*	9,294	18.4	(13.5 - 23.4)
\$25,000-\$49,999	177	10.7	(5.4 - 16.0)	9,495	9.4	(7.0 - 11.9)
\$50,000-\$74,999	107	7.2	(2.4 - 12.1)	5,733	9.0	(5.3 - 12.6)
≥ \$75,000	*	*	*	8,510	4.9	(3.5 - 6.2)

¹ Defined as 14 or more days within the past month when physical health was not good

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 4a. Poor mental health 14 or more days within the past month among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	649	8,800	12.9	(8.6 - 17.1)	37,528	1,473,200	10.1	(8.6 - 11.5)
Gender								
Male	231	3,700	11.4	(4.5 - 18.3)	14,407	555,200	7.9	(6.3 - 9.6)
Female	418	5,000	14.3	(9.1 - 19.4)	23,121	918,000	12.0	(9.7 - 14.3)
Age (years)								
18-34	*	*	*	*	3,889	496,700	11.2	(7.4 - 15.1)
35-44	*	*	*	*	4,947	254,100	8.9	(6.7 - 11.1)
45-54	138	2,400	16.7	(9.7 - 23.7)	7,749	347,200	12.3	(9.0 - 15.6)
55-64	162	800	7.5	(3.6 - 11.5)	8,623	211,300	10.1	(7.9 - 12.3)
≥ 65	199	700	6.2	(2.4 - 9.9)	12,320	163,900	6.7	(4.7 - 8.7)
Race/Ethnicity								
White Non-Hispanic	626	7,000	10.9	(7.7 - 14.2)	35,266	899,200	9.8	(8.4 - 11.2)
Black Non-Hispanic	*	*	*	*	722	227,800	10.9	(5.4 - 16.4)
Hispanic	*	*	*	*	845	263,700	11.8	(6.8 - 16.8)
Other Non-Hispanic	*	*	*	*	695	82,500	7.4	(1.4 - 13.5)
Education								
≤ High School	308	5,200	16.6	(9.0 - 24.2)	15,335	665,000	12.2	(10.0 - 14.5)
Some College	162	2,300	12.9	(5.9 - 20.0)	9,771	404,500	11.5	(8.3 - 14.8)
≥ College Degree	178	1,300	7.0	(2.1 - 11.9)	12,342	346,600	6.2	(4.5 - 7.9)
Income								
≤ \$24,999	*	*	*	*	9,398	611,500	17.9	(13.0 - 22.8)
\$25,000-\$49,999	178	2,200	13.1	(6.9 - 19.3)	9,547	260,300	9.0	(6.7 - 11.4)
\$50,000-\$74,999	108	1,500	11.0	(4.0 - 17.9)	5,726	187,300	9.6	(6.0 - 13.1)
≥ \$75,000	*	*	*	*	8,520	276,500	6.2	(4.6 - 7.7)

¹ Defined as 14 or more days within the past month when mental health was not good

² Sample size from survey

³ Estimated number of adults with poor mental health 14 or more days within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 4b. Poor mental health 14 or more days within the past month among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	649	13.2	(8.5 - 17.9)	37,528	10.2	(8.7 - 11.7)
Gender						
Male	231	11.6	(4.1 - 19.0)	14,407	7.9	(6.3 - 9.6)
Female	418	14.8	(8.9 - 20.6)	23,121	12.2	(9.8 - 14.6)
Race/Ethnicity						
White Non-Hispanic	626	11.0	(7.5 - 14.6)	35,266	10.0	(8.5 - 11.4)
Black Non-Hispanic	*	*	*	722	11.1	(5.5 - 16.7)
Hispanic	*	*	*	845	11.5	(6.5 - 16.4)
Other Non-Hispanic	*	*	*	695	7.6	(1.5 - 13.7)
Education						
≤ High School	308	17.3	(8.8 - 25.8)	15,335	12.3	(10.0 - 14.6)
Some College	162	12.9	(5.1 - 20.7)	9,771	11.8	(8.5 - 15.1)
≥ College Degree	178	7.3	(1.8 - 12.8)	12,342	6.3	(4.6 - 8.0)
Income						
≤ \$24,999	*	*	*	9,398	18.2	(13.2 - 23.1)
\$25,000-\$49,999	178	13.2	(6.5 - 19.8)	9,547	9.4	(6.9 - 11.9)
\$50,000-\$74,999	108	10.3	(3.2 - 17.3)	5,726	9.5	(5.9 - 13.1)
≥ \$75,000	*	*	*	8,520	6.2	(4.6 - 7.7)

¹ Defined as 14 or more days within the past month when mental health was not good

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 5a. Poor physical or mental health 14 or more days within the past month among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	645	13,900	20.5	(16.0 - 25.0)	37,005	2,385,900	16.5	(14.8 - 18.2)
Gender								
Male	230	5,600	16.9	(9.7 - 24.2)	14,231	937,700	13.6	(11.4 - 15.8)
Female	415	8,400	23.8	(18.2 - 29.3)	22,774	1,448,200	19.1	(16.5 - 21.7)
Age (years)								
18-34	*	*	*	*	3,859	602,600	13.8	(9.7 - 17.9)
35-44	*	*	*	*	4,924	388,300	13.7	(10.4 - 17.0)
45-54	138	3,100	21.4	(13.8 - 29.0)	7,722	551,000	19.6	(15.8 - 23.4)
55-64	162	2,200	19.7	(13.2 - 26.1)	8,545	395,500	19.2	(16.1 - 22.3)
≥ 65	195	2,500	21.7	(15.4 - 28.0)	11,955	448,500	18.8	(15.9 - 21.7)
Race/Ethnicity								
White Non-Hispanic	622	11,500	18.0	(14.3 - 21.6)	34,778	1,485,700	16.3	(14.8 - 17.8)
Black Non-Hispanic	*	*	*	*	715	343,700	16.5	(10.3 - 22.7)
Hispanic	*	*	*	*	832	412,900	19.0	(12.6 - 25.3)
Other Non-Hispanic	*	*	*	*	680	143,500	13.0	(5.0 - 21.0)
Education								
≤ High School	306	8,000	25.7	(18.1 - 33.3)	15,018	1,077,200	20.3	(17.3 - 23.2)
Some College	161	4,000	22.4	(13.9 - 31.0)	9,672	615,500	17.6	(14.1 - 21.1)
≥ College Degree	177	2,000	10.5	(5.1 - 16.0)	12,244	634,200	11.4	(9.2 - 13.6)
Income								
≤ \$24,999	*	*	*	*	9,199	924,000	27.3	(21.9 - 32.8)
\$25,000-\$49,999	177	3,500	20.5	(13.4 - 27.5)	9,425	456,000	16.1	(13.0 - 19.2)
\$50,000-\$74,999	107	1,900	13.9	(6.4 - 21.5)	5,696	312,000	16.0	(11.4 - 20.5)
≥ \$75,000	130	1,500	8.5	(3.3 - 13.7)	8,475	447,600	10.0	(8.1 - 12.0)

¹ Defined as 14 or more days within the past month when physical or mental health prevented usual activities, such as self-care, work or recreation

² Sample size from survey

³ Estimated number of adults with poor physical or mental health 14 or more days within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 5b. Poor physical or mental health 14 or more days within the past month among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	645	20.3	(15.3 - 25.2)	37,005	16.4	(14.7 - 18.2)
Gender						
Male	230	16.6	(8.8 - 24.4)	14,231	13.6	(11.4 - 15.8)
Female	415	23.7	(17.5 - 29.9)	22,774	19.0	(16.3 - 21.6)
Race/Ethnicity						
White Non-Hispanic	622	17.5	(13.6 - 21.5)	34,778	16.3	(14.7 - 17.8)
Black Non-Hispanic	*	*	*	715	16.7	(10.3 - 23.0)
Hispanic	*	*	*	832	18.5	(12.2 - 24.8)
Other Non-Hispanic	*	*	*	680	13.1	(5.0 - 21.3)
Education						
≤ High School	306	26.0	(17.5 - 34.4)	15,018	20.1	(17.1 - 23.1)
Some College	161	21.5	(12.4 - 30.6)	9,672	17.6	(14.1 - 21.1)
≥ College Degree	177	10.4	(4.6 - 16.2)	12,244	11.4	(9.2 - 13.7)
Income						
≤ \$24,999	*	*	*	9,199	27.6	(22.1 - 33.1)
\$25,000-\$49,999	177	20.2	(12.7 - 27.6)	9,425	16.1	(12.9 - 19.3)
\$50,000-\$74,999	107	12.7	(5.2 - 20.1)	5,696	15.6	(11.0 - 20.1)
≥ \$75,000	130	8.6	(3.0 - 14.2)	8,475	10.0	(8.0 - 12.0)

¹ Defined as 14 or more days within the past month when physical or mental health prevented usual activities, such as self-care, work or recreation

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 6a. Health insurance among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	656	62,700	90.9	(87.7 - 94.0)	38,119	12,854,200	86.6	(84.2 - 89.0)
Gender								
Male	236	30,400	89.5	(84.2 - 94.8)	14,597	5,881,100	83.0	(78.9 - 87.1)
Female	420	32,300	92.2	(88.8 - 95.6)	23,522	6,973,100	89.9	(87.2 - 92.6)
Age (years)								
18-34	*	*	*	*	3,910	3,302,100	74.3	(67.8 - 80.7)
35-44	99	17,000	91.4	(84.5 - 98.2)	4,986	2,560,300	88.9	(84.7 - 93.1)
45-54	141	13,600	90.0	(84.3 - 95.6)	7,831	2,530,100	88.9	(84.6 - 93.2)
55-64	162	10,500	93.4	(89.7 - 97.2)	8,764	1,986,700	92.8	(90.2 - 95.5)
≥ 65	204	11,900	98.4	(96.1 - 100)	12,628	2,475,000	97.8	(96.7 - 98.8)
Race/Ethnicity								
White Non-Hispanic	632	58,500	90.6	(87.3 - 93.9)	35,811	8,591,200	92.2	(90.6 - 93.8)
Black Non-Hispanic	*	*	*	*	736	1,781,600	82.7	(75.3 - 90.1)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	311	27,200	85.8	(80.0 - 91.5)	15,673	4,448,900	79.7	(74.9 - 84.5)
Some College	165	16,500	93.3	(87.8 - 98.7)	9,906	3,004,600	85.0	(80.1 - 89.8)
≥ College Degree	178	18,600	96.9	(94.6 - 99.3)	12,459	5,361,700	95.3	(93.3 - 97.2)
Income								
≤ \$24,999	162	11,000	84.7	(76.2 - 93.3)	9,640	2,437,200	69.7	(62.5 - 76.8)
\$25,000-\$49,999	179	15,100	87.4	(80.7 - 94.1)	9,662	2,384,000	82.2	(76.1 - 88.3)
\$50,000-\$74,999	108	12,700	93.0	(86.9 - 99.0)	5,772	1,855,700	94.0	(91.2 - 96.7)
≥ \$75,000	131	17,200	94.9	(89.0 - 100)	8,567	4,388,400	97.5	(96.6 - 98.5)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Sample size from survey

³ Estimated number of adults who have health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 6b. Health insurance among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	656	90.3	(86.8 - 93.9)	38,119	86.7	(84.4 - 89.1)
Gender						
Male	236	89.3	(83.5 - 95.1)	14,597	83.6	(79.7 - 87.5)
Female	420	91.3	(87.3 - 95.4)	23,522	89.6	(86.8 - 92.3)
Race/Ethnicity						
White Non-Hispanic	632	90.0	(86.3 - 93.7)	35,811	92.1	(90.5 - 93.7)
Black Non-Hispanic	*	*	*	736	82.5	(75.0 - 90.0)
Hispanic	*	*	*	860	68.7	(59.0 - 78.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	311	85.1	(78.6 - 91.5)	15,673	80.1	(75.4 - 84.7)
Some College	165	92.2	(85.8 - 98.7)	9,906	84.6	(79.7 - 89.4)
≥ College Degree	178	96.9	(94.4 - 99.4)	12,459	95.4	(93.6 - 97.2)
Income						
≤ \$24,999	162	84.5	(74.8 - 94.1)	9,640	70.2	(63.2 - 77.1)
\$25,000-\$49,999	179	86.9	(79.5 - 94.2)	9,662	81.9	(76.0 - 87.9)
\$50,000-\$74,999	108	92.1	(84.9 - 99.2)	5,772	94.0	(91.3 - 96.7)
≥ \$75,000	131	94.4	(87.7 - 100)	8,567	97.5	(96.6 - 98.5)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 7a. Health insurance among adults aged 18-64¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	445	50,300	89.2	(85.4 - 92.9)	25,165	10,251,300	84.2	(81.3 - 87.1)
Gender								
Male	160	25,100	88.2	(82.0 - 94.4)	10,078	4,839,900	80.3	(75.5 - 85.0)
Female	285	25,200	90.2	(86.0 - 94.4)	15,087	5,411,400	88.0	(84.7 - 91.3)
Age (years)								
18-34	*	*	*	*	3,910	3,302,100	74.3	(67.8 - 80.7)
35-44	99	17,000	91.4	(84.5 - 98.2)	4,981	2,556,600	88.9	(84.7 - 93.1)
45-54	141	13,600	90.0	(84.3 - 95.6)	7,791	2,472,900	88.7	(84.3 - 93.0)
55-64	155	10,000	93.2	(89.2 - 97.1)	8,483	1,919,700	92.8	(90.1 - 95.5)
≥ 65	-	-	-	-	-	-	-	-
Race/Ethnicity								
White Non-Hispanic	424	46,200	88.7	(84.7 - 92.7)	23,354	6,582,200	90.3	(88.3 - 92.4)
Black Non-Hispanic	*	*	*	*	567	1,464,100	80.2	(71.6 - 88.7)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	192	20,100	82.1	(74.8 - 89.4)	9,037	3,191,500	74.6	(68.6 - 80.6)
Some College	123	14,200	92.7	(86.6 - 98.9)	7,031	2,540,400	82.9	(77.4 - 88.4)
≥ College Degree	129	15,600	96.4	(93.6 - 99.2)	9,068	4,502,400	94.5	(92.2 - 96.8)
Income								
≤ \$24,999	*	*	*	*	4,918	1,707,200	62.6	(54.1 - 71.2)
\$25,000-\$49,999	124	11,800	85.1	(76.9 - 93.3)	6,180	1,783,000	77.7	(70.3 - 85.1)
\$50,000-\$74,999	96	11,800	92.5	(86.0 - 99.0)	4,554	1,553,400	92.9	(89.7 - 96.1)
≥ \$75,000	115	16,100	94.5	(88.2 - 100)	7,324	3,987,600	97.3	(96.3 - 98.3)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Sample size from survey

³ Estimated number of adults aged 18-64 who have health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 7b. Health insurance among adults aged 18-64¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	445	88.7	(84.4 - 92.9)	25,165	84.3	(81.5 - 87.2)
Gender						
Male	160	87.8	(80.9 - 94.8)	10,078	80.5	(75.9 - 85.1)
Female	285	89.4	(84.6 - 94.3)	15,087	87.9	(84.6 - 91.2)
Race/Ethnicity						
White Non-Hispanic	424	88.1	(83.6 - 92.6)	23,354	90.3	(88.3 - 92.3)
Black Non-Hispanic	*	*	*	567	80.0	(71.3 - 88.6)
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	192	81.3	(73.1 - 89.4)	9,037	75.1	(69.3 - 80.9)
Some College	123	91.7	(84.4 - 99.0)	7,031	82.6	(77.1 - 88.1)
≥ College Degree	129	96.4	(93.4 - 99.3)	9,068	94.6	(92.5 - 96.8)
Income						
≤ \$24,999	*	*	*	4,918	63.2	(54.9 - 71.6)
\$25,000-\$49,999	124	84.5	(75.6 - 93.5)	6,180	77.5	(70.2 - 84.7)
\$50,000-\$74,999	96	91.5	(83.9 - 99.1)	4,554	93.0	(89.8 - 96.2)
≥ \$75,000	115	94.0	(86.9 - 100)	7,324	97.3	(96.3 - 98.3)

¹ Defined as having any kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wayne County**

Table 8a. No health insurance among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	656	6,300	9.1	(6.0 - 12.3)	38,119	1,989,300	13.4	(11.0 - 15.8)
Gender								
Male	236	3,600	10.5	(5.2 - 15.8)	14,597	1,207,400	17.0	(12.9 - 21.1)
Female	420	2,700	7.8	(4.4 - 11.2)	23,522	781,900	10.1	(7.4 - 12.8)
Age (years)								
18-34	*	*	*	*	3,910	1,144,100	25.7	(19.3 - 32.2)
35-44	*	*	*	*	4,986	319,200	11.1	(6.9 - 15.3)
45-54	141	1,500	10.0	(4.4 - 15.7)	7,831	316,500	11.1	(6.8 - 15.4)
55-64	162	700	6.6	(2.8 - 10.3)	8,764	153,300	7.2	(4.5 - 9.8)
≥ 65	*	*	*	*	12,628	56,200	2.2	(1.2 - 3.3)
Race/Ethnicity								
White Non-Hispanic	632	6,100	9.4	(6.1 - 12.7)	35,811	727,100	7.8	(6.2 - 9.4)
Black Non-Hispanic	*	*	*	*	736	372,800	17.3	(9.9 - 24.7)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	311	4,500	14.2	(8.5 - 20.0)	15,673	1,134,400	20.3	(15.5 - 25.1)
Some College	*	*	*	*	9,906	531,400	15.0	(10.2 - 19.9)
≥ College Degree	*	*	*	*	12,459	266,300	4.7	(2.8 - 6.7)
Income								
≤ \$24,999	162	2,000	15.3	(6.7 - 23.8)	9,640	1,061,200	30.3	(23.2 - 37.5)
\$25,000-\$49,999	179	2,200	12.6	(5.9 - 19.3)	9,662	516,700	17.8	(11.7 - 23.9)
\$50,000-\$74,999	*	*	*	*	5,772	119,100	6.0	(3.3 - 8.8)
≥ \$75,000	*	*	*	*	8,567	111,100	2.5	(1.5 - 3.4)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Sample size from survey

³ Estimated number of adults who have no health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 8b. No health insurance among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	656	9.7	(6.1 - 13.2)	38,119	13.3	(10.9 - 15.6)
Gender						
Male	236	10.7	(4.9 - 16.5)	14,597	16.4	(12.5 - 20.3)
Female	420	8.7	(4.6 - 12.7)	23,522	10.4	(7.7 - 13.2)
Race/Ethnicity						
White Non-Hispanic	632	10.0	(6.3 - 13.7)	35,811	7.9	(6.3 - 9.5)
Black Non-Hispanic	*	*	*	736	17.5	(10.0 - 25.0)
Hispanic	*	*	*	860	31.3	(21.6 - 41.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	311	14.9	(8.5 - 21.4)	15,673	19.9	(15.3 - 24.6)
Some College	*	*	*	9,906	15.4	(10.6 - 20.3)
≥ College Degree	*	*	*	12,459	4.6	(2.8 - 6.4)
Income						
≤ \$24,999	162	15.5	(5.9 - 25.2)	9,640	29.8	(22.9 - 36.8)
\$25,000-\$49,999	179	13.1	(5.8 - 20.5)	9,662	18.1	(12.1 - 24.0)
\$50,000-\$74,999	*	*	*	5,772	6.0	(3.3 - 8.7)
≥ \$75,000	*	*	*	8,567	2.5	(1.5 - 3.4)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 9a. No health insurance among adults aged 18-64¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	445	6,100	10.8	(7.1 - 14.6)	25,165	1,928,400	15.8	(12.9 - 18.7)
Gender								
Male	160	3,400	11.8	(5.6 - 18.0)	10,078	1,190,200	19.7	(15.0 - 24.5)
Female	285	2,700	9.8	(5.6 - 14.0)	15,087	738,200	12.0	(8.7 - 15.3)
Age (years)								
18-34	*	*	*	*	3,910	1,144,100	25.7	(19.3 - 32.2)
35-44	*	*	*	*	4,981	319,200	11.1	(6.9 - 15.3)
45-54	141	1,500	10.0	(4.4 - 15.7)	7,791	316,100	11.3	(7.0 - 15.7)
55-64	155	700	6.8	(2.9 - 10.8)	8,483	148,900	7.2	(4.5 - 9.9)
≥ 65	-	-	-	-	-	-	-	-
Race/Ethnicity								
White Non-Hispanic	424	5,900	11.3	(7.3 - 15.3)	23,354	703,100	9.7	(7.6 - 11.7)
Black Non-Hispanic	*	*	*	*	567	361,800	19.8	(11.3 - 28.4)
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	192	4,400	17.9	(10.6 - 25.2)	9,037	1,087,700	25.4	(19.4 - 31.4)
Some College	*	*	*	*	7,031	522,800	17.1	(11.6 - 22.6)
≥ College Degree	*	*	*	*	9,068	261,200	5.5	(3.2 - 7.8)
Income								
≤ \$24,999	*	*	*	*	4,918	1,018,000	37.4	(28.8 - 45.9)
\$25,000-\$49,999	124	2,100	14.9	(6.7 - 23.1)	6,180	511,100	22.3	(14.9 - 29.7)
\$50,000-\$74,999	*	*	*	*	4,554	118,800	7.1	(3.9 - 10.3)
≥ \$75,000	*	*	*	*	7,324	110,700	2.7	(1.7 - 3.7)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Sample size from survey

³ Estimated number of adults aged 18-64 who have no health insurance (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 9b. No health insurance among adults aged 18-64¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	445	11.3	(7.1 - 15.6)	25,165	15.7	(12.8 - 18.5)
Gender						
Male	160	12.2	(5.2 - 19.1)	10,078	19.5	(14.9 - 24.1)
Female	285	10.6	(5.7 - 15.4)	15,087	12.1	(8.8 - 15.4)
Race/Ethnicity						
White Non-Hispanic	424	11.9	(7.4 - 16.4)	23,354	9.7	(7.7 - 11.7)
Black Non-Hispanic	*	*	*	567	20.0	(11.4 - 28.7)
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	192	18.7	(10.6 - 26.9)	9,037	24.9	(19.1 - 30.7)
Some College	*	*	*	7,031	17.4	(11.9 - 22.9)
≥ College Degree	*	*	*	9,068	5.4	(3.2 - 7.5)
Income						
≤ \$24,999	*	*	*	4,918	36.8	(28.4 - 45.1)
\$25,000-\$49,999	124	15.5	(6.5 - 24.4)	6,180	22.5	(15.3 - 29.8)
\$50,000-\$74,999	*	*	*	4,554	7.0	(3.8 - 10.2)
≥ \$75,000	*	*	*	7,324	2.7	(1.7 - 3.7)

¹ Defined as having no kind of health care coverage, including health insurance, prepaid plans such as health maintenance organizations, or government plans such as Medicare among adults aged 18-64

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 10a. Adults with regular health care provider¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	655	65,700	95.1	(92.6 - 97.6)	38,135	12,324,300	83.0	(80.4 - 85.6)
Gender								
Male	234	31,300	93.0	(88.3 - 97.7)	14,593	5,494,300	77.6	(73.2 - 82.0)
Female	421	34,400	97.1	(95.3 - 98.8)	23,542	6,830,000	87.8	(85.1 - 90.5)
Age (years)								
18-34	50	11,300	94.6	(88.4 - 100)	3,929	2,931,600	65.8	(59.2 - 72.5)
35-44	100	17,900	95.2	(90.3 - 100)	4,988	2,471,000	86.1	(81.9 - 90.3)
45-54	141	13,700	90.8	(83.0 - 98.6)	7,823	2,529,300	88.6	(84.6 - 92.7)
55-64	161	11,000	98.6	(96.5 - 100)	8,770	2,002,000	93.6	(91.7 - 95.4)
≥ 65	203	11,800	97.5	(95.2 - 99.8)	12,625	2,390,400	94.1	(92.2 - 96.0)
Race/Ethnicity								
White Non-Hispanic	631	62,000	95.9	(93.9 - 98.0)	35,821	8,227,700	88.2	(86.4 - 89.9)
Black Non-Hispanic	*	*	*	*	740	1,753,200	81.3	(72.6 - 89.9)
Hispanic	*	*	*	*	860	1,453,100	64.7	(54.7 - 74.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	311	29,200	91.8	(86.9 - 96.6)	15,692	4,249,200	75.9	(70.9 - 80.8)
Some College	164	17,700	99.3	(98.3 - 100)	9,904	2,935,000	82.8	(77.2 - 88.4)
≥ College Degree	178	18,500	96.6	(93.1 - 100)	12,456	5,046,100	89.8	(87.1 - 92.6)
Income								
≤ \$24,999	161	12,100	93.9	(89.5 - 98.3)	9,644	2,554,200	72.7	(65.9 - 79.6)
\$25,000-\$49,999	178	16,000	94.4	(89.2 - 99.5)	9,655	2,243,100	77.3	(70.0 - 84.6)
\$50,000-\$74,999	108	13,100	96.3	(92.5 - 100)	5,771	1,767,500	89.5	(85.2 - 93.9)
≥ \$75,000	130	17,700	97.7	(94.3 - 100)	8,560	4,122,800	91.9	(90.2 - 93.7)

¹ Defined as having one or more persons respondent thinks of as a personal doctor or health care provider

² Sample size from survey

³ Estimated number of adults who have at least one person they think of as their personal doctor or health care provider (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 10b. Adults with regular health care provider¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	655	95.1	(92.6 - 97.6)	38,135	83.0	(80.4 - 85.5)
Gender						
Male	234	92.9	(88.2 - 97.6)	14,593	78.1	(73.9 - 82.3)
Female	421	97.2	(95.3 - 99.0)	23,542	87.4	(84.6 - 90.1)
Race/Ethnicity						
White Non-Hispanic	631	95.7	(93.4 - 98.0)	35,821	88.0	(86.2 - 89.7)
Black Non-Hispanic	*	*	*	740	81.4	(73.0 - 89.8)
Hispanic	*	*	*	860	65.2	(55.5 - 75.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	311	91.7	(86.9 - 96.5)	15,692	75.8	(71.0 - 80.6)
Some College	164	99.4	(98.6 - 100)	9,904	82.9	(77.5 - 88.3)
≥ College Degree	178	96.5	(92.5 - 100)	12,456	89.8	(87.1 - 92.4)
Income						
≤ \$24,999	161	94.1	(89.7 - 98.5)	9,644	72.9	(66.1 - 79.6)
\$25,000-\$49,999	178	93.8	(87.9 - 99.7)	9,655	77.4	(70.5 - 84.3)
\$50,000-\$74,999	108	96.3	(92.2 - 100)	5,771	89.6	(85.3 - 94.0)
≥ \$75,000	130	97.6	(93.7 - 100)	8,560	91.6	(89.8 - 93.4)

¹ Defined as having one or more persons respondent thinks of as a personal doctor or health care provider

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 11a. Cost prevented visit to doctor within the past year among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	657	7,800	11.2	(7.1 - 15.3)	38,143	2,034,800	13.7	(11.6 - 15.8)
Gender								
Male	236	3,500	10.3	(3.5 - 17.1)	14,607	903,300	12.7	(9.5 - 16.0)
Female	421	4,300	12.1	(7.4 - 16.9)	23,536	1,131,500	14.6	(11.8 - 17.4)
Age (years)								
18-34	*	*	*	*	3,939	947,300	21.2	(15.6 - 26.9)
35-44	*	*	*	*	4,987	347,900	12.1	(8.0 - 16.2)
45-54	141	1,800	11.7	(6.4 - 17.0)	7,825	382,200	13.5	(9.9 - 17.1)
55-64	162	900	7.9	(3.9 - 11.9)	8,766	236,800	11.1	(8.0 - 14.2)
≥ 65	*	*	*	*	12,626	120,600	4.8	(2.5 - 7.0)
Race/Ethnicity								
White Non-Hispanic	633	5,600	8.6	(5.7 - 11.5)	35,833	984,600	10.6	(8.9 - 12.2)
Black Non-Hispanic	*	*	*	*	738	280,000	13.1	(7.2 - 19.0)
Hispanic	*	*	*	*	859	597,900	26.5	(17.3 - 35.7)
Other Non-Hispanic	*	*	*	*	713	172,300	15.3	(5.7 - 24.9)
Education								
≤ High School	311	4,300	13.6	(6.4 - 20.8)	15,696	996,600	17.8	(13.8 - 21.9)
Some College	166	2,200	12.3	(4.8 - 19.9)	9,911	512,400	14.5	(10.4 - 18.6)
≥ College Degree	178	1,200	6.4	(1.7 - 11.0)	12,453	466,900	8.3	(6.0 - 10.6)
Income								
≤ \$24,999	*	*	*	*	9,639	854,500	24.3	(18.4 - 30.3)
\$25,000-\$49,999	179	2,200	12.7	(5.2 - 20.1)	9,656	532,900	18.4	(12.5 - 24.3)
\$50,000-\$74,999	*	*	*	*	5,776	202,700	10.3	(6.0 - 14.5)
≥ \$75,000	*	*	*	*	8,561	192,000	4.3	(2.9 - 5.6)

¹ Defined as needing to see a doctor within the past year but not being able to because of cost

² Sample size from survey

³ Estimated number of adults who did not see a doctor because of cost within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 11b. Cost prevented visit to doctor within the past year among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	657	11.6	(7.0 - 16.2)	38,143	13.8	(11.7 - 15.9)
Gender						
Male	236	10.6	(3.1 - 18.2)	14,607	12.5	(9.4 - 15.6)
Female	421	12.5	(7.0 - 17.9)	23,536	15.0	(12.1 - 17.9)
Race/Ethnicity						
White Non-Hispanic	633	8.7	(5.4 - 11.9)	35,833	10.8	(9.2 - 12.5)
Black Non-Hispanic	*	*	*	738	13.2	(7.2 - 19.3)
Hispanic	*	*	*	859	26.1	(17.2 - 35.1)
Other Non-Hispanic	*	*	*	713	14.8	(5.5 - 24.2)
Education						
≤ High School	311	14.1	(6.0 - 22.3)	15,696	17.8	(13.8 - 21.7)
Some College	166	12.8	(4.4 - 21.1)	9,911	15.3	(11.0 - 19.5)
≥ College Degree	178	6.5	(1.2 - 11.8)	12,453	8.2	(6.0 - 10.5)
Income						
≤ \$24,999	*	*	*	9,639	24.5	(18.6 - 30.4)
\$25,000-\$49,999	179	13.8	(5.3 - 22.4)	9,656	18.7	(12.9 - 24.6)
\$50,000-\$74,999	*	*	*	5,776	10.4	(6.1 - 14.7)
≥ \$75,000	*	*	*	8,561	4.4	(3.1 - 5.8)

¹ Defined as needing to see a doctor within the past year but not being able to because of cost

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 12a. Visited a doctor for routine checkup within the past year among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	656	46,300	66.9	(61.8 - 71.9)	38,026	10,809,300	72.9	(70.6 - 75.3)
Gender								
Male	235	21,300	62.9	(54.5 - 71.4)	14,592	4,738,400	66.9	(63.0 - 70.9)
Female	421	25,000	70.6	(65.1 - 76.1)	23,434	6,070,900	78.5	(75.9 - 81.0)
Age (years)								
18-34	*	*	*	*	3,921	2,747,400	61.7	(55.6 - 67.9)
35-44	*	*	*	*	4,973	1,883,100	65.5	(60.2 - 70.7)
45-54	140	9,000	59.9	(50.6 - 69.2)	7,808	2,174,100	76.6	(73.2 - 80.0)
55-64	162	8,700	77.7	(70.7 - 84.7)	8,751	1,728,600	81.4	(78.7 - 84.2)
≥ 65	203	10,200	85.2	(79.9 - 90.6)	12,573	2,276,000	90.0	(87.5 - 92.5)
Race/Ethnicity								
White Non-Hispanic	632	44,000	67.9	(63.1 - 72.7)	35,721	6,746,200	72.4	(70.6 - 74.2)
Black Non-Hispanic	*	*	*	*	736	1,688,900	78.8	(70.4 - 87.3)
Hispanic	*	*	*	*	857	1,550,900	69.3	(60.0 - 78.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	310	20,300	64.1	(56.2 - 72.0)	15,640	4,150,600	74.7	(70.7 - 78.7)
Some College	166	14,100	77.8	(69.9 - 85.7)	9,880	2,524,000	71.3	(66.1 - 76.5)
≥ College Degree	178	11,600	60.5	(51.1 - 69.8)	12,428	4,042,900	71.9	(68.5 - 75.3)
Income								
≤ \$24,999	*	*	*	*	9,616	2,469,200	70.4	(64.2 - 76.6)
\$25,000-\$49,999	177	11,200	65.3	(56.3 - 74.4)	9,631	2,068,400	71.3	(65.3 - 77.3)
\$50,000-\$74,999	*	*	*	*	5,765	1,466,000	74.8	(69.6 - 80.0)
≥ \$75,000	*	*	*	*	8,547	3,176,500	70.7	(67.3 - 74.2)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past year

² Sample size from survey

³ Estimated number of adults who have visited a doctor for a routine checkup within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 12b. Visited a doctor for routine checkup within the past year among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	656	65.9	(60.4 - 71.4)	38,026	72.7	(70.3 - 75.0)
Gender						
Male	235	62.2	(53.1 - 71.4)	14,592	67.1	(63.3 - 70.9)
Female	421	69.5	(63.3 - 75.7)	23,434	77.7	(75.1 - 80.3)
Race/Ethnicity						
White Non-Hispanic	632	67.1	(61.8 - 72.4)	35,721	71.9	(70.1 - 73.7)
Black Non-Hispanic	*	*	*	736	79.3	(71.3 - 87.4)
Hispanic	*	*	*	857	69.2	(60.1 - 78.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	310	63.2	(54.7 - 71.8)	15,640	74.4	(70.4 - 78.4)
Some College	166	77.4	(68.8 - 86.1)	9,880	70.9	(65.9 - 76.0)
≥ College Degree	*	*	*	12,428	71.7	(68.3 - 75.0)
Income						
≤ \$24,999	*	*	*	9,616	70.2	(64.1 - 76.2)
\$25,000-\$49,999	177	63.5	(53.6 - 73.3)	9,631	71.1	(65.3 - 76.9)
\$50,000-\$74,999	*	*	*	5,765	74.5	(69.2 - 79.8)
≥ \$75,000	*	*	*	8,547	70.3	(66.9 - 73.8)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past year

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 13a. Visited a doctor for routine checkup within the past 2 years among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	656	56,400	81.5	(77.4 - 85.5)	38,026	12,726,600	85.9	(83.9 - 87.9)
Gender								
Male	235	26,500	78.4	(71.5 - 85.2)	14,592	5,791,900	81.8	(78.2 - 85.4)
Female	421	29,900	84.4	(79.9 - 88.9)	23,434	6,934,700	89.6	(87.9 - 91.3)
Age (years)								
18-34	*	*	*	*	3,921	3,535,400	79.4	(73.9 - 85.0)
35-44	100	14,600	77.5	(67.9 - 87.1)	4,973	2,355,600	81.9	(78.0 - 85.7)
45-54	140	12,300	81.7	(74.6 - 88.9)	7,808	2,507,900	88.4	(85.6 - 91.2)
55-64	162	9,900	88.0	(82.7 - 93.3)	8,751	1,924,800	90.7	(88.6 - 92.7)
≥ 65	203	10,900	90.9	(86.2 - 95.5)	12,573	2,402,900	95.0	(93.5 - 96.5)
Race/Ethnicity								
White Non-Hispanic	632	52,400	80.8	(76.6 - 85.1)	35,721	8,070,000	86.6	(85.4 - 87.9)
Black Non-Hispanic	*	*	*	*	736	1,914,500	89.4	(82.3 - 96.5)
Hispanic	*	*	*	*	857	1,832,800	81.9	(73.7 - 90.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	310	25,400	80.0	(73.8 - 86.1)	15,640	4,713,700	84.8	(81.2 - 88.4)
Some College	166	16,200	89.6	(84.2 - 95.0)	9,880	2,990,500	84.5	(79.6 - 89.4)
≥ College Degree	178	14,500	75.8	(66.9 - 84.8)	12,428	4,928,300	87.6	(85.3 - 89.9)
Income								
≤ \$24,999	162	11,000	85.3	(78.8 - 91.8)	9,616	2,911,700	83.0	(77.4 - 88.6)
\$25,000-\$49,999	177	13,800	80.8	(72.9 - 88.8)	9,631	2,375,500	81.9	(76.1 - 87.7)
\$50,000-\$74,999	108	11,100	81.5	(72.5 - 90.6)	5,765	1,733,500	88.4	(84.9 - 91.9)
≥ \$75,000	131	14,100	77.8	(68.3 - 87.4)	8,547	3,907,000	87.0	(84.8 - 89.2)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past 2 years

² Sample size from survey

³ Estimated number of adults who have visited a doctor for a routine checkup within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 13b. Visited a doctor for routine checkup within the past 2 years among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	656	80.6	(76.0 - 85.1)	38,026	85.7	(83.7 - 87.6)
Gender						
Male	235	77.2	(69.6 - 84.8)	14,592	81.9	(78.4 - 85.3)
Female	421	83.8	(78.7 - 88.9)	23,434	89.1	(87.4 - 90.9)
Race/Ethnicity						
White Non-Hispanic	632	79.8	(75.1 - 84.6)	35,721	86.2	(85.0 - 87.5)
Black Non-Hispanic	*	*	*	736	89.6	(83.0 - 96.3)
Hispanic	*	*	*	857	81.8	(73.8 - 89.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	310	79.1	(72.3 - 85.9)	15,640	84.6	(81.1 - 88.1)
Some College	166	89.4	(83.5 - 95.4)	9,880	84.4	(79.8 - 89.0)
≥ College Degree	*	*	*	12,428	87.3	(85.0 - 89.6)
Income						
≤ \$24,999	162	85.2	(78.2 - 92.2)	9,616	83.2	(77.8 - 88.5)
\$25,000-\$49,999	177	80.1	(71.5 - 88.8)	9,631	81.9	(76.3 - 87.4)
\$50,000-\$74,999	*	*	*	5,765	88.1	(84.4 - 91.8)
≥ \$75,000	*	*	*	8,547	86.4	(84.2 - 88.7)

¹ Defined as having visited a doctor for a routine checkup (general physical exam, not an exam for a specific injury, illness or condition) within the past 2 years

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 14a. Dentist visit within the past year among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	656	51,500	74.6	(70.1 - 79.1)	38,026	10,540,400	71.1	(68.6 - 73.6)
Gender								
Male	235	25,300	74.6	(67.5 - 81.8)	14,573	4,845,800	68.4	(64.5 - 72.2)
Female	421	26,200	74.6	(69.2 - 79.9)	23,453	5,694,600	73.6	(70.4 - 76.8)
Age (years)								
18-34	*	*	*	*	3,924	2,789,700	62.6	(56.3 - 69.0)
35-44	*	*	*	*	4,979	2,181,000	75.8	(70.9 - 80.7)
45-54	141	11,600	77.0	(69.3 - 84.8)	7,814	2,149,300	75.8	(71.9 - 79.7)
55-64	162	8,900	79.0	(72.3 - 85.7)	8,742	1,690,500	79.2	(75.9 - 82.6)
≥ 65	203	9,000	74.4	(68.0 - 80.7)	12,567	1,729,800	68.5	(64.7 - 72.3)
Race/Ethnicity								
White Non-Hispanic	632	48,500	75.0	(70.7 - 79.4)	35,721	6,959,900	74.7	(72.7 - 76.6)
Black Non-Hispanic	*	*	*	*	735	1,405,800	66.2	(57.9 - 74.5)
Hispanic	*	*	*	*	862	1,425,600	63.1	(53.7 - 72.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	311	20,900	65.6	(58.4 - 72.8)	15,613	3,620,700	65.1	(60.7 - 69.5)
Some College	165	14,000	78.6	(70.2 - 87.1)	9,880	2,504,700	70.8	(66.1 - 75.5)
≥ College Degree	178	16,700	87.0	(80.4 - 93.6)	12,453	4,402,400	78.2	(74.6 - 81.8)
Income								
≤ \$24,999	*	*	*	*	9,594	1,972,200	56.2	(49.8 - 62.6)
\$25,000-\$49,999	179	12,300	71.3	(62.2 - 80.3)	9,649	1,896,000	65.8	(59.5 - 72.1)
\$50,000-\$74,999	108	11,400	83.6	(76.0 - 91.3)	5,763	1,482,400	75.2	(69.4 - 81.0)
≥ \$75,000	131	15,700	86.6	(78.3 - 94.8)	8,564	3,707,900	82.4	(79.5 - 85.3)

¹ Defined as having visited a dentist or a dental clinic for any reason within the past year

² Sample size from survey

³ Estimated number of adults who have visited a dentist or a dental clinic for any reason within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 14b. Dentist visit within the past year among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	656	74.2	(69.2 - 79.1)	38,026	71.1	(68.6 - 73.5)
Gender						
Male	235	73.8	(65.9 - 81.6)	14,573	68.3	(64.6 - 72.1)
Female	421	74.6	(68.6 - 80.6)	23,453	73.6	(70.4 - 76.8)
Race/Ethnicity						
White Non-Hispanic	632	74.7	(70.0 - 79.5)	35,721	74.4	(72.4 - 76.3)
Black Non-Hispanic	*	*	*	735	66.6	(58.3 - 74.9)
Hispanic	*	*	*	862	63.6	(54.4 - 72.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	311	65.2	(57.3 - 73.1)	15,613	65.3	(60.9 - 69.6)
Some College	165	77.8	(68.3 - 87.2)	9,880	70.4	(65.7 - 75.1)
≥ College Degree	178	86.7	(79.4 - 94.0)	12,453	78.1	(74.6 - 81.7)
Income						
≤ \$24,999	*	*	*	9,594	56.4	(50.1 - 62.8)
\$25,000-\$49,999	*	*	*	9,649	65.8	(59.6 - 72.0)
\$50,000-\$74,999	108	83.9	(75.8 - 92.0)	5,763	75.0	(69.1 - 80.8)
≥ \$75,000	131	86.0	(77.0 - 95.1)	8,564	82.1	(79.1 - 85.0)

¹ Defined as having visited a dentist or a dental clinic for any reason within the past year

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 15a. Had teeth cleaned within the past year among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	601	48,500	73.5	(68.6 - 78.3)	34,070	9,985,900	71.7	(69.1 - 74.3)
Gender								
Male	220	23,700	72.5	(64.7 - 80.4)	13,173	4,616,400	69.0	(65.0 - 73.1)
Female	381	24,700	74.4	(68.6 - 80.1)	20,897	5,369,500	74.2	(70.8 - 77.5)
Age (years)								
18-34	*	*	*	*	3,860	2,746,400	63.0	(56.5 - 69.5)
35-44	*	*	*	*	4,875	2,112,100	75.5	(70.7 - 80.3)
45-54	137	10,400	72.0	(62.5 - 81.6)	7,438	2,022,600	73.3	(69.1 - 77.4)
55-64	150	8,500	79.6	(72.7 - 86.5)	7,906	1,560,900	77.7	(73.8 - 81.6)
≥ 65	164	8,100	81.2	(74.9 - 87.4)	9,991	1,543,900	77.1	(73.7 - 80.5)
Race/Ethnicity								
White Non-Hispanic	578	46,300	75.1	(70.5 - 79.7)	31,982	6,601,500	75.4	(73.4 - 77.4)
Black Non-Hispanic	*	*	*	*	680	1,227,200	61.6	(52.1 - 71.0)
Hispanic	*	*	*	*	789	1,421,300	66.6	(57.1 - 76.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	268	18,500	63.6	(55.6 - 71.6)	12,740	3,388,600	67.7	(63.1 - 72.3)
Some College	156	13,400	76.6	(67.6 - 85.5)	9,145	2,286,800	68.5	(62.9 - 74.1)
≥ College Degree	175	16,500	86.8	(80.1 - 93.6)	12,128	4,298,500	78.3	(74.8 - 81.7)
Income								
≤ \$24,999	*	*	*	*	7,525	1,861,800	60.3	(53.5 - 67.1)
\$25,000-\$49,999	168	10,700	65.3	(55.7 - 75.0)	8,709	1,681,700	62.6	(55.4 - 69.8)
\$50,000-\$74,999	104	11,000	83.2	(75.4 - 90.9)	5,534	1,447,800	76.3	(70.7 - 81.8)
≥ \$75,000	131	15,700	86.3	(78.0 - 94.6)	8,448	3,617,500	80.8	(77.8 - 83.8)

¹ Defined as having had teeth cleaned by a dentist or dental hygienist within the past year

² Sample size from survey

³ Estimated number of adults who have had their teeth cleaned by a dentist or dental hygienist within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 15b. Had teeth cleaned within the past year among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	601	72.9	(67.7 - 78.2)	34,070	71.7	(69.1 - 74.2)
Gender						
Male	220	72.0	(63.6 - 80.4)	13,173	69.2	(65.3 - 73.1)
Female	381	73.9	(67.4 - 80.3)	20,897	73.9	(70.5 - 77.3)
Race/Ethnicity						
White Non-Hispanic	578	74.5	(69.4 - 79.5)	31,982	75.1	(73.0 - 77.1)
Black Non-Hispanic	*	*	*	680	62.0	(52.7 - 71.4)
Hispanic	*	*	*	789	66.9	(57.5 - 76.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	268	63.5	(54.9 - 72.1)	12,740	67.7	(63.2 - 72.3)
Some College	156	75.5	(65.6 - 85.5)	9,145	68.1	(62.6 - 73.6)
≥ College Degree	175	86.0	(78.3 - 93.6)	12,128	78.3	(74.9 - 81.7)
Income						
≤ \$24,999	*	*	*	7,525	60.2	(53.4 - 67.0)
\$25,000-\$49,999	*	*	*	8,709	62.7	(55.7 - 69.7)
\$50,000-\$74,999	104	83.8	(75.6 - 91.9)	5,534	76.4	(70.9 - 81.9)
≥ \$75,000	131	85.6	(76.5 - 94.7)	8,448	80.4	(77.4 - 83.4)

¹ Defined as having had teeth cleaned by a dentist or dental hygienist within the past year

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 16a. Permanent teeth extracted due to decay or gum disease among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	647	36,100	52.9	(47.7 - 58.1)	37,652	7,426,500	50.4	(47.8 - 52.9)
Gender								
Male	231	18,600	55.8	(47.2 - 64.4)	14,457	3,367,000	47.8	(43.9 - 51.6)
Female	416	17,500	50.2	(44.2 - 56.1)	23,195	4,059,500	52.7	(49.5 - 56.0)
Age (years)								
18-34	*	*	*	*	3,921	1,121,600	25.4	(19.7 - 31.0)
35-44	*	*	*	*	4,964	1,250,000	43.6	(37.9 - 49.2)
45-54	140	8,500	56.4	(47.0 - 65.8)	7,780	1,595,300	56.1	(51.4 - 60.7)
55-64	160	7,800	70.6	(62.6 - 78.5)	8,648	1,382,800	65.5	(61.6 - 69.4)
≥ 65	198	9,700	82.2	(76.7 - 87.7)	12,339	2,076,900	83.0	(80.6 - 85.4)
Race/Ethnicity								
White Non-Hispanic	623	32,400	50.7	(45.6 - 55.8)	35,364	4,425,700	47.9	(46.0 - 49.9)
Black Non-Hispanic	*	*	*	*	727	1,244,700	58.2	(48.9 - 67.4)
Hispanic	*	*	*	*	853	1,220,500	54.2	(44.9 - 63.5)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	303	20,400	65.5	(58.0 - 73.0)	15,427	3,436,000	61.9	(57.2 - 66.6)
Some College	*	*	*	*	9,798	1,726,200	49.3	(44.3 - 54.2)
≥ College Degree	177	6,800	35.6	(27.1 - 44.1)	12,351	2,242,600	40.1	(36.5 - 43.6)
Income								
≤ \$24,999	157	9,500	76.2	(66.7 - 85.7)	9,483	2,171,900	62.3	(55.8 - 68.8)
\$25,000-\$49,999	178	10,100	58.9	(49.6 - 68.2)	9,552	1,577,400	54.7	(48.3 - 61.0)
\$50,000-\$74,999	*	*	*	*	5,744	1,000,700	51.5	(45.6 - 57.4)
≥ \$75,000	130	5,700	31.7	(22.4 - 41.0)	8,516	1,738,200	38.8	(35.3 - 42.2)

¹ Defined as having had at least one permanent tooth extracted because of tooth decay or gum disease

² Sample size from survey

³ Estimated number of adults who have had at least one permanent tooth extracted because of tooth decay or gum disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 16b. Permanent teeth extracted due to decay or gum disease among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	647	50.1	(44.5 - 55.8)	37,652	49.4	(46.9 - 51.9)
Gender						
Male	231	53.4	(44.0 - 62.7)	14,457	47.9	(44.1 - 51.7)
Female	416	47.0	(40.4 - 53.6)	23,195	50.7	(47.4 - 54.0)
Race/Ethnicity						
White Non-Hispanic	623	47.7	(42.2 - 53.2)	35,364	46.5	(44.5 - 48.5)
Black Non-Hispanic	*	*	*	727	57.7	(48.6 - 66.9)
Hispanic	*	*	*	853	53.8	(44.6 - 62.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	303	63.2	(54.9 - 71.5)	15,427	61.3	(56.7 - 66.0)
Some College	*	*	*	9,798	47.9	(43.0 - 52.9)
≥ College Degree	177	33.8	(24.8 - 42.7)	12,351	39.0	(35.5 - 42.5)
Income						
≤ \$24,999	*	*	*	9,483	61.6	(55.2 - 68.1)
\$25,000-\$49,999	*	*	*	9,552	54.2	(48.0 - 60.3)
\$50,000-\$74,999	*	*	*	5,744	49.9	(44.0 - 55.9)
≥ \$75,000	130	29.2	(19.6 - 38.9)	8,516	37.7	(34.2 - 41.1)

¹ Defined as having had at least one permanent tooth extracted because of tooth decay or gum disease

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 17a. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	198	2,100	17.7	(12.1 - 23.3)	12,339	491,200	19.6	(16.0 - 23.3)
Gender								
Male	72	700	13.3	(5.0 - 21.6)	4,352	167,200	16.4	(11.8 - 20.9)
Female	126	1,400	21.2	(13.6 - 28.8)	7,987	324,000	21.8	(16.6 - 27.0)
Age (years)								
≥ 65	198	2,100	17.7	(12.1 - 23.3)	12,339	491,200	19.6	(16.0 - 23.3)
Race/Ethnicity								
White Non-Hispanic	195	2,100	17.7	(12.0 - 23.4)	11,885	322,600	16.8	(14.3 - 19.2)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	111	1,600	23.9	(15.4 - 32.3)	6,358	314,100	25.3	(19.9 - 30.7)
Some College	*	*	*	*	2,736	90,600	20.2	(12.1 - 28.2)
≥ College Degree	*	*	*	*	3,214	80,500	10.1	(3.6 - 16.5)
Income								
≤ \$24,999	*	*	*	*	4,558	199,600	26.9	(19.3 - 34.4)
\$25,000-\$49,999	*	*	*	*	3,365	123,900	21.4	(13.1 - 29.6)
\$50,000-\$74,999	*	*	*	*	1,179	21,400	7.3	(3.8 - 10.8)
≥ \$75,000	*	*	*	*	1,187	9,600	2.5	(1.0 - 4.0)

¹ Defined as having all permanent teeth extracted because of tooth decay or gum disease among adults aged 65 and older

² Sample size from survey

³ Estimated number of adults aged 65 and older who have had all permanent teeth extracted because of tooth decay or gum disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 17b. Had all permanent teeth extracted due to decay or gum disease among adults aged 65 and older¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	198	17.3	(11.7 - 23.0)	12,339	19.4	(15.7 - 23.2)
Gender						
Male	72	13.3	(5.0 - 21.7)	4,352	16.5	(11.5 - 21.4)
Female	126	21.1	(13.5 - 28.7)	7,987	22.1	(16.6 - 27.7)
Race/Ethnicity						
White Non-Hispanic	195	17.3	(11.6 - 23.0)	11,885	16.6	(13.9 - 19.3)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	111	23.9	(15.2 - 32.5)	6,358	24.9	(19.3 - 30.5)
Some College	*	*	*	2,736	21.0	(12.1 - 29.9)
≥ College Degree	*	*	*	3,214	10.0	(3.6 - 16.3)
Income						
≤ \$24,999	*	*	*	4,558	27.3	(19.1 - 35.5)
\$25,000-\$49,999	*	*	*	3,365	21.0	(12.9 - 29.1)
\$50,000-\$74,999	*	*	*	1,179	7.6	(3.6 - 11.5)
≥ \$75,000	*	*	*	1,187	2.2	(0.9 - 3.5)

¹ Defined as having all permanent teeth extracted because of tooth decay or gum disease among adults aged 65 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wayne County**

Table 18a. Flu shot within the past 12 months¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	655	33,500	48.3	(43.2 - 53.5)	38,079	6,255,900	42.3	(39.9 - 44.8)
Gender								
Male	235	14,500	42.8	(34.7 - 50.9)	14,579	2,798,900	39.6	(35.8 - 43.5)
Female	420	19,000	53.6	(47.6 - 59.6)	23,500	3,456,900	44.8	(41.6 - 47.9)
Age (years)								
18-49	214	12,900	34.5	(26.7 - 42.4)	12,505	2,685,300	31.1	(27.3 - 34.9)
50-64	239	10,500	52.5	(45.0 - 60.1)	12,972	1,662,900	46.0	(42.6 - 49.5)
≥ 65	202	10,100	84.4	(78.9 - 89.9)	12,602	1,907,700	75.2	(72.0 - 78.4)
Race/Ethnicity								
White Non-Hispanic	631	32,300	49.8	(44.8 - 54.9)	35,770	4,029,000	43.3	(41.4 - 45.3)
Black Non-Hispanic	*	*	*	*	738	907,000	42.1	(33.1 - 51.1)
Hispanic	*	*	*	*	860	882,900	39.6	(30.4 - 48.9)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	310	14,500	45.6	(38.0 - 53.1)	15,663	2,457,800	44.3	(39.8 - 48.8)
Some College	*	*	*	*	9,898	1,377,700	38.9	(34.0 - 43.9)
≥ College Degree	178	10,400	54.3	(44.9 - 63.7)	12,436	2,404,400	43.0	(39.5 - 46.4)
Income								
≤ \$24,999	*	*	*	*	9,635	1,656,800	47.6	(41.2 - 54.0)
\$25,000-\$49,999	179	7,600	43.9	(35.2 - 52.6)	9,647	1,200,600	41.4	(35.6 - 47.3)
\$50,000-\$74,999	*	*	*	*	5,763	792,300	40.2	(34.3 - 46.2)
≥ \$75,000	*	*	*	*	8,548	1,776,100	39.6	(36.3 - 42.8)

¹ Defined as having had a flu shot within the past 12 months

² Sample size from survey

³ Estimated number of adults who have had a flu shot within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 18b. Flu shot within the past 12 months¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	655	46.8	(41.2 - 52.3)	38,079	41.7	(39.2 - 44.1)
Gender						
Male	235	42.0	(33.3 - 50.7)	14,579	40.0	(36.2 - 43.8)
Female	420	51.3	(44.7 - 58.0)	23,500	43.1	(40.0 - 46.3)
Race/Ethnicity						
White Non-Hispanic	631	48.0	(42.5 - 53.5)	35,770	42.4	(40.4 - 44.3)
Black Non-Hispanic	*	*	*	738	42.3	(33.3 - 51.3)
Hispanic	*	*	*	860	39.0	(29.9 - 48.1)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	310	44.3	(36.1 - 52.4)	15,663	43.7	(39.1 - 48.2)
Some College	*	*	*	9,898	37.8	(33.0 - 42.7)
≥ College Degree	*	*	*	12,436	42.5	(39.1 - 46.0)
Income						
≤ \$24,999	*	*	*	9,635	47.3	(41.0 - 53.7)
\$25,000-\$49,999	179	41.1	(32.0 - 50.2)	9,647	40.3	(34.5 - 46.0)
\$50,000-\$74,999	*	*	*	5,763	39.4	(33.4 - 45.4)
≥ \$75,000	*	*	*	8,548	39.1	(35.8 - 42.4)

¹ Defined as having had a flu shot within the past 12 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 19a. Flu vaccine that was sprayed in nose within the past 12 months¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	*	*	*	*	38,106	248,300	1.7	(0.6 - 2.8)
Gender								
Male	*	*	*	*	14,586	148,000	2.1	(0.0 - 4.2)
Female	*	*	*	*	23,520	100,300	1.3	(0.5 - 2.1)
Age (years)								
18-49	*	*	*	*	12,535	184,900	2.1	(0.3 - 3.9)
50-64	*	*	*	*	12,968	36,000	1.0	(0.2 - 1.8)
≥ 65	*	*	*	*	12,603	27,400	1.1	(0.4 - 1.8)
Race/Ethnicity								
White Non-Hispanic	*	*	*	*	35,796	66,000	0.7	(0.4 - 1.0)
Black Non-Hispanic	*	*	*	*	740	38,500	1.8	(0.0 - 4.1)
Hispanic	*	*	*	*	860	133,700	5.9	(0.0 - 12.3)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	15,679	172,500	3.1	(0.3 - 5.9)
Some College	*	*	*	*	9,897	43,500	1.2	(0.4 - 2.0)
≥ College Degree	*	*	*	*	12,449	31,600	0.6	(0.3 - 0.8)
Income								
≤ \$24,999	*	*	*	*	9,636	138,400	3.9	(0.0 - 8.1)
\$25,000-\$49,999	*	*	*	*	9,648	41,500	1.4	(0.0 - 3.1)
\$50,000-\$74,999	*	*	*	*	5,766	13,300	0.7	(0.2 - 1.2)
≥ \$75,000	*	*	*	*	8,561	35,300	0.8	(0.3 - 1.3)

¹ Defined as having had flu vaccine sprayed in nose within the past 12 months

² Sample size from survey

³ Estimated number of adults who have had flu vaccine sprayed in nose within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 19b. Flu vaccine that was sprayed in nose within the past 12 months¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	*	*	*	38,106	1.7	(0.6 - 2.7)
Gender						
Male	*	*	*	14,586	2.0	(0.1 - 3.9)
Female	*	*	*	23,520	1.4	(0.5 - 2.3)
Race/Ethnicity						
White Non-Hispanic	*	*	*	35,796	0.7	(0.4 - 1.0)
Black Non-Hispanic	*	*	*	740	1.9	(0.0 - 4.4)
Hispanic	*	*	*	860	5.9	(0.0 - 11.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	15,679	3.1	(0.4 - 5.8)
Some College	*	*	*	9,897	1.2	(0.4 - 2.0)
≥ College Degree	*	*	*	12,449	0.6	(0.3 - 0.8)
Income						
≤ \$24,999	*	*	*	9,636	3.8	(0.0 - 7.7)
\$25,000-\$49,999	*	*	*	9,648	1.5	(0.0 - 3.4)
\$50,000-\$74,999	*	*	*	5,766	0.7	(0.1 - 1.2)
≥ \$75,000	*	*	*	8,561	0.8	(0.3 - 1.3)

¹ Defined as having had flu vaccine sprayed in nose within the past 12 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 20a. Flu shot or flu vaccine sprayed in nose within the past 12 months¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	655	33,500	48.3	(43.2 - 53.5)	38,059	6,302,300	42.7	(40.2 - 45.1)
Gender								
Male	235	14,500	42.8	(34.7 - 50.9)	14,568	2,822,700	40.0	(36.1 - 43.9)
Female	420	19,000	53.6	(47.6 - 59.6)	23,491	3,479,500	45.1	(42.0 - 48.2)
Age (years)								
18-49	214	12,900	34.5	(26.7 - 42.4)	12,501	2,726,400	31.6	(27.8 - 35.4)
50-64	239	10,500	52.5	(45.0 - 60.1)	12,964	1,664,200	46.1	(42.7 - 49.5)
≥ 65	202	10,100	84.4	(78.9 - 89.9)	12,594	1,911,700	75.4	(72.2 - 78.6)
Race/Ethnicity								
White Non-Hispanic	631	32,300	49.8	(44.8 - 54.9)	35,750	4,048,700	43.6	(41.7 - 45.5)
Black Non-Hispanic	*	*	*	*	738	910,000	42.2	(33.2 - 51.2)
Hispanic	*	*	*	*	860	903,300	40.6	(31.3 - 49.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	310	14,500	45.6	(38.0 - 53.1)	15,651	2,480,100	44.7	(40.2 - 49.3)
Some College	*	*	*	*	9,895	1,387,300	39.2	(34.3 - 44.1)
≥ College Degree	178	10,400	54.3	(44.9 - 63.7)	12,431	2,418,300	43.2	(39.8 - 46.7)
Income								
≤ \$24,999	*	*	*	*	9,626	1,675,100	48.2	(41.8 - 54.6)
\$25,000-\$49,999	179	7,600	43.9	(35.2 - 52.6)	9,643	1,208,300	41.7	(35.8 - 47.6)
\$50,000-\$74,999	*	*	*	*	5,761	795,600	40.4	(34.5 - 46.3)
≥ \$75,000	*	*	*	*	8,548	1,791,200	39.9	(36.6 - 43.2)

¹ Defined as having had a flu shot or flu vaccine sprayed in nose within the past 12 months

² Sample size from survey

³ Estimated number of adults who have had a flu shot or flu vaccine sprayed in nose within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 20b. Flu shot or flu vaccine sprayed in nose within the past 12 months¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	655	46.8	(41.2 - 52.3)	38,059	42.0	(39.6 - 44.5)
Gender						
Male	235	42.0	(33.3 - 50.7)	14,568	40.4	(36.6 - 44.2)
Female	420	51.3	(44.7 - 58.0)	23,491	43.5	(40.3 - 46.7)
Race/Ethnicity						
White Non-Hispanic	631	48.0	(42.5 - 53.5)	35,750	42.6	(40.7 - 44.6)
Black Non-Hispanic	*	*	*	738	42.4	(33.4 - 51.4)
Hispanic	*	*	*	860	40.0	(30.9 - 49.1)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	310	44.3	(36.1 - 52.4)	15,651	44.1	(39.6 - 48.6)
Some College	*	*	*	9,895	38.1	(33.2 - 43.0)
≥ College Degree	*	*	*	12,431	42.8	(39.4 - 46.3)
Income						
≤ \$24,999	*	*	*	9,626	48.0	(41.6 - 54.3)
\$25,000-\$49,999	179	41.1	(32.0 - 50.2)	9,643	40.5	(34.8 - 46.3)
\$50,000-\$74,999	*	*	*	5,761	39.6	(33.6 - 45.6)
≥ \$75,000	*	*	*	8,548	39.4	(36.1 - 42.7)

¹ Defined as having had a flu shot or flu vaccine sprayed in nose within the past 12 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wayne County**

Table 21a. Flu shot among adults aged 65 and older¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	202	10,100	84.4	(78.9 - 89.9)	12,602	1,907,700	75.2	(72.0 - 78.4)
Gender								
Male	73	4,400	82.5	(73.1 - 91.9)	4,417	772,900	74.6	(69.3 - 79.8)
Female	129	5,800	85.9	(79.5 - 92.3)	8,185	1,134,800	75.6	(71.6 - 79.6)
Age (years)								
≥ 65	202	10,100	84.4	(78.9 - 89.9)	12,602	1,907,700	75.2	(72.0 - 78.4)
Race/Ethnicity								
White Non-Hispanic	199	10,000	84.2	(78.7 - 89.7)	12,137	1,487,100	76.0	(73.5 - 78.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	115	5,900	83.6	(76.0 - 91.2)	6,518	930,800	74.0	(68.9 - 79.2)
Some College	*	*	*	*	2,790	317,800	69.7	(62.7 - 76.7)
≥ College Degree	*	*	*	*	3,261	654,000	80.5	(76.4 - 84.6)
Income								
≤ \$24,999	86	4,000	85.2	(77.5 - 92.9)	4,665	545,300	72.4	(65.6 - 79.3)
\$25,000-\$49,999	*	*	*	*	3,422	430,600	72.8	(67.0 - 78.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	1,198	301,000	77.5	(70.9 - 84.2)

¹ Defined as having had a flu shot during the past 12 months among adults aged 65 and older

² Sample size from survey

³ Estimated number of adults aged 65 and older who have had a flu shot within the past 12 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 21b. Flu shot among adults aged 65 and older¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	202	84.2	(78.6 - 89.9)	12,602	75.0	(71.5 - 78.5)
Gender						
Male	73	82.5	(73.1 - 91.9)	4,417	74.4	(68.7 - 80.1)
Female	129	85.9	(79.5 - 92.2)	8,185	75.5	(71.3 - 79.8)
Race/Ethnicity						
White Non-Hispanic	199	84.1	(78.4 - 89.7)	12,137	75.9	(73.0 - 78.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	115	83.2	(75.2 - 91.1)	6,518	73.3	(67.5 - 79.1)
Some College	*	*	*	2,790	70.0	(62.7 - 77.2)
≥ College Degree	*	*	*	3,261	80.7	(76.2 - 85.1)
Income						
≤ \$24,999	86	85.5	(77.9 - 93.2)	4,665	72.6	(65.1 - 80.1)
\$25,000-\$49,999	*	*	*	3,422	71.4	(65.0 - 77.9)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	1,198	78.0	(70.8 - 85.1)

¹ Defined as having had a flu shot during the past 12 months among adults aged 65 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wayne County**

Table 22a. Pneumonia shot or pneumococcal vaccine¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	607	17,700	29.2	(25.0 - 33.4)	35,303	3,456,800	26.1	(23.9 - 28.3)
Gender								
Male	211	7,200	25.0	(18.4 - 31.5)	13,052	1,542,200	25.2	(21.5 - 28.8)
Female	396	10,500	33.0	(27.8 - 38.3)	22,251	1,914,600	26.9	(24.2 - 29.6)
Age (years)								
18-49	182	3,800	12.5	(7.1 - 17.9)	10,943	1,097,000	14.6	(11.3 - 17.9)
50-64	228	4,900	26.6	(20.5 - 32.7)	12,179	758,000	23.0	(19.8 - 26.2)
≥ 65	197	9,000	76.5	(70.1 - 82.9)	12,181	1,601,700	65.7	(61.7 - 69.7)
Race/Ethnicity								
White Non-Hispanic	588	17,300	30.2	(25.9 - 34.4)	33,225	2,358,700	28.7	(26.9 - 30.5)
Black Non-Hispanic	*	*	*	*	685	584,400	28.4	(19.5 - 37.3)
Hispanic	*	*	*	*	769	408,000	19.5	(12.6 - 26.4)
Other Non-Hispanic	*	*	*	*	624	105,600	12.2	(2.3 - 22.2)
Education								
≤ High School	292	9,400	33.0	(26.5 - 39.4)	14,700	1,535,300	30.4	(26.4 - 34.3)
Some College	148	3,400	23.3	(16.2 - 30.4)	9,151	836,200	27.2	(21.9 - 32.5)
≥ College Degree	165	4,800	28.5	(20.2 - 36.8)	11,375	1,070,700	21.4	(18.6 - 24.1)
Income								
≤ \$24,999	*	*	*	*	9,082	1,050,900	33.1	(27.6 - 38.6)
\$25,000-\$49,999	170	5,200	33.4	(25.1 - 41.7)	9,033	797,500	30.3	(24.1 - 36.6)
\$50,000-\$74,999	99	2,300	19.5	(11.1 - 28.0)	5,320	378,800	21.8	(17.4 - 26.1)
≥ \$75,000	116	2,300	15.2	(7.6 - 22.8)	7,773	648,700	16.3	(14.1 - 18.5)

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine

² Sample size from survey

³ Estimated number of adults who have ever had a pneumonia shot or pneumococcal vaccine (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 22b. Pneumonia shot or pneumococcal vaccine¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	607	27.7	(23.3 - 32.2)	35,303	25.4	(23.2 - 27.6)
Gender						
Male	211	24.6	(17.6 - 31.6)	13,052	25.8	(22.3 - 29.4)
Female	396	30.6	(25.1 - 36.0)	22,251	25.1	(22.4 - 27.8)
Race/Ethnicity						
White Non-Hispanic	588	28.8	(24.3 - 33.3)	33,225	27.7	(25.9 - 29.5)
Black Non-Hispanic	*	*	*	685	27.9	(19.1 - 36.7)
Hispanic	*	*	*	769	19.6	(12.8 - 26.5)
Other Non-Hispanic	*	*	*	624	11.3	(2.0 - 20.7)
Education						
≤ High School	292	31.5	(24.7 - 38.3)	14,700	29.5	(25.6 - 33.5)
Some College	148	21.3	(14.2 - 28.3)	9,151	26.2	(21.0 - 31.3)
≥ College Degree	165	27.5	(18.6 - 36.4)	11,375	21.0	(18.3 - 23.8)
Income						
≤ \$24,999	*	*	*	9,082	32.3	(26.9 - 37.8)
\$25,000-\$49,999	170	32.9	(24.1 - 41.8)	9,033	29.0	(23.0 - 35.0)
\$50,000-\$74,999	99	18.0	(9.4 - 26.7)	5,320	21.2	(16.8 - 25.7)
≥ \$75,000	116	14.8	(6.6 - 23.0)	7,773	16.2	(13.9 - 18.5)

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 23a. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	197	9,000	76.5	(70.1 - 82.9)	12,181	1,601,700	65.7	(61.7 - 69.7)
Gender								
Male	*	*	*	*	4,180	606,400	62.1	(55.5 - 68.7)
Female	127	5,400	81.2	(74.0 - 88.4)	8,001	995,300	68.0	(63.1 - 73.0)
Age (years)								
≥ 65	197	9,000	76.5	(70.1 - 82.9)	12,181	1,601,700	65.7	(61.7 - 69.7)
Race/Ethnicity								
White Non-Hispanic	194	9,000	76.5	(70.1 - 83.0)	11,735	1,281,300	68.2	(64.9 - 71.5)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	112	5,600	80.5	(72.4 - 88.6)	6,345	806,200	65.6	(59.5 - 71.7)
Some College	*	*	*	*	2,694	289,500	66.8	(60.1 - 73.4)
≥ College Degree	*	*	*	*	3,111	501,100	65.6	(58.5 - 72.7)
Income								
≤ \$24,999	82	3,700	81.3	(72.5 - 90.1)	4,533	497,100	67.7	(59.9 - 75.5)
\$25,000-\$49,999	*	*	*	*	3,324	344,300	61.5	(53.3 - 69.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	1,138	236,300	64.5	(55.2 - 73.8)

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine among adults aged 65 and older

² Sample size from survey

³ Estimated number of adults aged 65 and older who have ever had a pneumonia shot or pneumococcal vaccine (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 23b. Pneumonia shot or pneumococcal vaccine among adults aged 65 and older¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	197	76.0	(69.4 - 82.6)	12,181	64.7	(60.4 - 69.1)
Gender						
Male	*	*	*	4,180	61.5	(54.3 - 68.6)
Female	127	81.3	(74.1 - 88.5)	8,001	67.6	(62.4 - 72.9)
Race/Ethnicity						
White Non-Hispanic	194	76.0	(69.3 - 82.6)	11,735	67.1	(63.2 - 70.9)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	112	79.7	(71.3 - 88.2)	6,345	64.2	(57.4 - 71.0)
Some College	*	*	*	2,694	65.8	(58.4 - 73.3)
≥ College Degree	*	*	*	3,111	65.3	(57.7 - 72.9)
Income						
≤ \$24,999	82	81.6	(72.8 - 90.5)	4,533	66.0	(57.0 - 74.9)
\$25,000-\$49,999	*	*	*	3,324	60.3	(51.9 - 68.6)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	*	*

¹ Defined as having ever had a pneumonia shot or pneumococcal vaccine among adults aged 65 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 24a. Ever had a mammogram among women aged 40 and older¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	355	22,700	93.8	(90.7 - 96.9)	19,491	4,328,800	91.8	(89.7 - 94.0)
Gender								
Male	-	-	-	-	-	-	-	-
Female	355	22,700	93.8	(90.7 - 96.9)	19,491	4,328,800	91.8	(89.7 - 94.0)
Age (years)								
40-49	88	7,900	93.6	(87.4 - 99.7)	3,851	1,287,500	85.0	(79.7 - 90.3)
50-64	137	8,500	94.4	(89.7 - 99.1)	7,452	1,641,700	96.0	(94.1 - 97.9)
≥ 65	130	6,300	93.4	(88.2 - 98.6)	8,188	1,399,600	94.1	(90.5 - 97.6)
Race/Ethnicity								
White Non-Hispanic	344	22,200	93.9	(90.8 - 97.1)	18,548	2,976,100	93.9	(92.7 - 95.1)
Black Non-Hispanic	*	*	*	*	342	746,600	93.8	(87.3 - 100)
Hispanic	*	*	*	*	333	447,600	91.8	(84.6 - 99.0)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	167	10,400	94.6	(90.7 - 98.5)	8,266	1,698,700	91.0	(87.4 - 94.5)
Some College	94	6,400	93.4	(86.5 - 100)	5,322	1,055,500	93.6	(90.2 - 97.0)
≥ College Degree	93	5,900	92.8	(86.5 - 99.1)	5,872	1,560,700	91.6	(87.5 - 95.6)
Income								
≤ \$24,999	101	5,200	96.2	(92.7 - 99.7)	5,691	1,023,400	93.5	(89.7 - 97.4)
\$25,000-\$49,999	94	5,800	88.6	(80.7 - 96.4)	4,857	860,300	89.5	(83.6 - 95.5)
\$50,000-\$74,999	57	4,300	93.8	(85.0 - 100)	2,648	547,100	94.6	(92.2 - 96.9)
≥ \$75,000	51	4,200	97.4	(92.2 - 100)	3,582	1,185,100	92.2	(88.2 - 96.1)

¹ Defined as having ever had a mammogram among women aged 40 and older

² Sample size from survey

³ Estimated number of women aged 40 and older who have ever had a mammogram (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 24b. Ever had a mammogram among women aged 40 and older¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	355	93.5	(90.1 - 97.0)	19,491	91.1	(88.7 - 93.6)
Gender						
Male	-	-	-	-	-	-
Female	355	93.5	(90.1 - 97.0)	19,491	91.1	(88.7 - 93.6)
Race/Ethnicity						
White Non-Hispanic	344	93.6	(90.1 - 97.1)	18,548	93.3	(92.0 - 94.7)
Black Non-Hispanic	*	*	*	342	93.1	(85.7 - 100)
Hispanic	*	*	*	333	91.0	(82.8 - 99.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	167	94.3	(89.9 - 98.7)	8,266	90.2	(86.1 - 94.2)
Some College	94	92.6	(84.5 - 100)	5,322	92.9	(88.9 - 96.9)
≥ College Degree	93	93.0	(86.8 - 99.3)	5,872	90.9	(86.6 - 95.3)
Income						
≤ \$24,999	101	96.2	(92.6 - 99.7)	5,691	93.0	(88.5 - 97.4)
\$25,000-\$49,999	94	88.1	(79.7 - 96.6)	4,857	88.8	(82.5 - 95.1)
\$50,000-\$74,999	*	*	*	2,648	94.3	(91.7 - 96.8)
≥ \$75,000	51	97.5	(92.8 - 100)	3,582	91.2	(86.8 - 95.6)

¹ Defined as having ever had a mammogram among women aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 25a. Had mammogram within the past 2 years among women aged 40 and older¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	355	17,600	72.8	(67.2 - 78.3)	19,376	3,760,000	80.3	(77.8 - 82.9)
Gender								
Male	-	-	-	-	-	-	-	-
Female	355	17,600	72.8	(67.2 - 78.3)	19,376	3,760,000	80.3	(77.8 - 82.9)
Age (years)								
40-49	*	*	*	*	3,842	1,122,700	74.4	(68.5 - 80.2)
50-64	137	7,000	78.1	(70.2 - 85.9)	7,428	1,487,900	87.2	(84.5 - 89.8)
≥ 65	130	5,200	77.3	(69.3 - 85.2)	8,106	1,149,400	78.5	(74.2 - 82.9)
Race/Ethnicity								
White Non-Hispanic	344	17,200	72.9	(67.3 - 78.5)	18,438	2,531,500	80.2	(78.3 - 82.2)
Black Non-Hispanic	*	*	*	*	339	667,100	85.6	(77.5 - 93.8)
Hispanic	*	*	*	*	332	419,700	86.1	(77.5 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	167	7,600	68.9	(60.8 - 77.0)	8,204	1,486,900	80.5	(76.4 - 84.6)
Some College	*	*	*	*	5,297	929,400	83.0	(78.8 - 87.1)
≥ College Degree	93	5,000	78.1	(68.6 - 87.6)	5,845	1,334,800	78.6	(73.9 - 83.2)
Income								
≤ \$24,999	*	*	*	*	5,650	882,300	80.8	(75.9 - 85.7)
\$25,000-\$49,999	*	*	*	*	4,846	732,300	76.3	(69.7 - 82.9)
\$50,000-\$74,999	*	*	*	*	2,641	494,200	85.6	(81.9 - 89.4)
≥ \$75,000	*	*	*	*	3,572	1,025,900	80.0	(75.2 - 84.8)

¹ Defined as having had a mammogram within the past 2 years among women aged 40 and older

² Sample size from survey

³ Estimated number of women aged 40 and older who have had a mammogram within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 25b. Had mammogram within the past 2 years among women aged 40 and older¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	355	71.9	(65.9 - 77.8)	19,376	79.7	(76.9 - 82.5)
Gender						
Male	-	-	-	-	-	-
Female	355	71.9	(65.9 - 77.8)	19,376	79.7	(76.9 - 82.5)
Race/Ethnicity						
White Non-Hispanic	344	71.9	(65.9 - 78.0)	18,438	79.7	(77.6 - 81.8)
Black Non-Hispanic	*	*	*	339	85.3	(76.5 - 94.2)
Hispanic	*	*	*	332	85.2	(75.6 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	167	67.8	(59.3 - 76.4)	8,204	80.1	(75.6 - 84.7)
Some College	*	*	*	5,297	82.3	(77.7 - 86.9)
≥ College Degree	*	*	*	5,845	77.8	(72.9 - 82.8)
Income						
≤ \$24,999	*	*	*	5,650	80.8	(75.4 - 86.2)
\$25,000-\$49,999	*	*	*	4,846	75.7	(68.7 - 82.6)
\$50,000-\$74,999	*	*	*	2,641	85.5	(81.6 - 89.4)
≥ \$75,000	*	*	*	3,572	79.0	(73.8 - 84.2)

¹ Defined as having had a mammogram within the past 2 years among women aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 26a. Had mammogram within the past 2 years among women aged 50 and older¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	267	12,300	77.7	(72.1 - 83.4)	15,534	2,637,300	83.2	(80.7 - 85.7)
Gender								
Male	-	-	-	-	-	-	-	-
Female	267	12,300	77.7	(72.1 - 83.4)	15,534	2,637,300	83.2	(80.7 - 85.7)
Age (years)								
50-64	137	7,000	78.1	(70.2 - 85.9)	7,428	1,487,900	87.2	(84.5 - 89.8)
≥ 65	130	5,200	77.3	(69.3 - 85.2)	8,106	1,149,400	78.5	(74.2 - 82.9)
Race/Ethnicity								
White Non-Hispanic	262	12,100	78.1	(72.4 - 83.7)	14,887	1,907,000	82.3	(80.3 - 84.4)
Black Non-Hispanic	*	*	*	*	249	385,600	89.9	(83.2 - 96.7)
Hispanic	*	*	*	*	220	286,700	93.9	(89.1 - 98.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	136	6,000	75.9	(68.0 - 83.7)	7,026	1,094,500	83.3	(80.0 - 86.6)
Some College	*	*	*	*	4,124	646,700	87.3	(84.2 - 90.3)
≥ College Degree	*	*	*	*	4,356	890,500	80.6	(75.1 - 86.1)
Income								
≤ \$24,999	*	*	*	*	4,991	649,100	81.0	(76.8 - 85.2)
\$25,000-\$49,999	*	*	*	*	3,913	533,300	78.6	(70.9 - 86.2)
\$50,000-\$74,999	*	*	*	*	1,943	359,300	89.6	(86.3 - 93.0)
≥ \$75,000	*	*	*	*	2,303	601,300	84.6	(79.3 - 89.9)

¹ Defined as having had a mammogram within the past 2 years among women aged 50 and older

² Sample size from survey

³ Estimated number of women aged 50 and older who have had a mammogram within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 26b. Had mammogram within the past 2 years among women aged 50 and older¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	267	77.5	(71.8 - 83.2)	15,534	83.1	(80.3 - 85.8)
Gender						
Male	-	-	-	-	-	-
Female	267	77.5	(71.8 - 83.2)	15,534	83.1	(80.3 - 85.8)
Race/Ethnicity						
White Non-Hispanic	262	77.8	(72.1 - 83.6)	14,887	82.1	(79.9 - 84.3)
Black Non-Hispanic	*	*	*	249	90.4	(83.6 - 97.2)
Hispanic	*	*	*	220	94.1	(89.3 - 99.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	136	75.7	(67.8 - 83.6)	7,026	83.6	(80.1 - 87.0)
Some College	*	*	*	4,124	87.3	(84.1 - 90.5)
≥ College Degree	*	*	*	4,356	79.9	(73.9 - 86.0)
Income						
≤ \$24,999	*	*	*	4,991	81.3	(76.9 - 85.6)
\$25,000-\$49,999	*	*	*	3,913	78.1	(69.9 - 86.4)
\$50,000-\$74,999	*	*	*	1,943	90.2	(87.1 - 93.4)
≥ \$75,000	*	*	*	2,303	83.9	(77.9 - 89.9)

¹ Defined as having had a mammogram within the past 2 years among women aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wayne County**

Table 27a. Ever had a Pap test among women¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	421	34,400	97.0	(95.0 - 99.0)	23,381	7,045,300	92.1	(90.0 - 94.3)
Gender								
Male	-	-	-	-	-	-	-	-
Female	421	34,400	97.0	(95.0 - 99.0)	23,381	7,045,300	92.1	(90.0 - 94.3)
Age (years)								
18-34	*	*	*	*	2,377	1,896,600	85.2	(79.3 - 91.0)
35-44	63	6,500	97.1	(92.6 - 100)	3,023	1,301,500	92.6	(87.7 - 97.5)
45-54	90	7,600	99.5	(98.5 - 100)	4,657	1,420,300	98.3	(97.0 - 99.6)
55-64	99	5,600	99.5	(98.4 - 100)	5,187	1,071,200	97.2	(95.0 - 99.3)
≥ 65	129	6,200	92.3	(87.4 - 97.3)	8,137	1,355,700	92.3	(88.5 - 96.1)
Race/Ethnicity								
White Non-Hispanic	407	33,000	96.9	(94.8 - 99.0)	21,976	4,439,700	94.2	(92.1 - 96.3)
Black Non-Hispanic	*	*	*	*	481	1,231,600	91.2	(84.0 - 98.4)
Hispanic	*	*	*	*	533	984,500	90.1	(85.3 - 94.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	188	14,200	95.6	(92.6 - 98.6)	9,526	2,540,500	88.3	(83.8 - 92.7)
Some College	120	10,600	96.7	(91.7 - 100)	6,462	1,782,200	93.5	(90.4 - 96.6)
≥ College Degree	112	9,600	99.6	(98.8 - 100)	7,345	2,643,200	95.0	(92.2 - 97.9)
Income								
≤ \$24,999	112	6,800	95.5	(91.9 - 99.1)	6,526	1,717,200	93.7	(90.9 - 96.6)
\$25,000-\$49,999	109	9,000	98.9	(96.6 - 100)	5,787	1,378,400	91.0	(85.6 - 96.4)
\$50,000-\$74,999	67	6,000	100	(100 - 100)	3,310	933,900	96.0	(91.8 - 100)
≥ \$75,000	73	7,700	96.0	(89.3 - 100)	4,570	1,943,300	95.4	(92.6 - 98.2)

¹ Defined as having ever had a Pap test among women

² Sample size from survey

³ Estimated number of women who have ever had a Pap test (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 27b. Ever had a Pap test among women¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	421	96.9	(94.6 - 99.2)	23,381	91.8	(89.6 - 94.0)
Gender						
Male	-	-	-	-	-	-
Female	421	96.9	(94.6 - 99.2)	23,381	91.8	(89.6 - 94.0)
Race/Ethnicity						
White Non-Hispanic	407	96.8	(94.3 - 99.2)	21,976	94.0	(91.8 - 96.2)
Black Non-Hispanic	*	*	*	481	90.8	(83.5 - 98.1)
Hispanic	*	*	*	533	89.7	(84.6 - 94.7)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	188	95.6	(92.5 - 98.8)	9,526	87.8	(83.2 - 92.4)
Some College	120	96.1	(90.0 - 100)	6,462	93.1	(89.7 - 96.4)
≥ College Degree	112	99.7	(99.0 - 100)	7,345	94.9	(92.0 - 97.7)
Income						
≤ \$24,999	112	95.6	(92.0 - 99.2)	6,526	93.3	(90.1 - 96.5)
\$25,000-\$49,999	109	99.0	(97.0 - 100)	5,787	91.1	(85.7 - 96.5)
\$50,000-\$74,999	67	100	(100 - 100)	3,310	96.0	(91.9 - 100)
≥ \$75,000	73	95.4	(87.5 - 100)	4,570	95.0	(91.9 - 98.1)

¹ Defined as having ever had a Pap test among women

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 28a. Had a Pap test within the past 3 years among women¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	419	28,100	79.5	(75.1 - 83.9)	23,109	6,205,600	82.2	(79.8 - 84.6)
Gender								
Male	-	-	-	-	-	-	-	-
Female	419	28,100	79.5	(75.1 - 83.9)	23,109	6,205,600	82.2	(79.8 - 84.6)
Age (years)								
18-34	*	*	*	*	2,371	1,818,400	81.8	(75.5 - 88.1)
35-44	*	*	*	*	3,012	1,220,300	86.9	(81.8 - 92.0)
45-54	89	5,900	78.2	(68.5 - 87.9)	4,640	1,311,500	92.0	(89.8 - 94.3)
55-64	98	4,400	79.6	(71.0 - 88.3)	5,152	924,300	84.5	(81.0 - 88.0)
≥ 65	129	4,100	60.7	(51.4 - 70.0)	7,934	931,100	66.4	(61.8 - 71.0)
Race/Ethnicity								
White Non-Hispanic	406	26,900	79.2	(74.7 - 83.7)	21,720	3,807,100	81.5	(79.2 - 83.8)
Black Non-Hispanic	*	*	*	*	473	1,142,800	86.7	(79.1 - 94.3)
Hispanic	*	*	*	*	527	894,900	83.6	(76.3 - 91.0)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	187	11,000	74.3	(67.1 - 81.5)	9,384	2,132,600	76.0	(71.1 - 81.0)
Some College	119	8,700	79.7	(71.1 - 88.3)	6,401	1,594,300	84.0	(80.4 - 87.5)
≥ College Degree	112	8,400	87.3	(80.8 - 93.8)	7,281	2,406,000	86.9	(83.7 - 90.1)
Income								
≤ \$24,999	*	*	*	*	6,441	1,453,900	82.0	(77.9 - 86.0)
\$25,000-\$49,999	107	7,400	83.1	(75.1 - 91.0)	5,740	1,182,000	78.3	(72.6 - 83.9)
\$50,000-\$74,999	*	*	*	*	3,296	825,000	85.0	(78.0 - 91.9)
≥ \$75,000	73	7,300	91.2	(83.4 - 99.0)	4,555	1,810,200	89.0	(85.7 - 92.2)

¹ Defined as having had a Pap test within the past 3 years among women

² Sample size from survey

³ Estimated number of women who have had a Pap test within the past 3 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 28b. Had a Pap test within the past 3 years among women¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	419	80.9	(76.3 - 85.4)	23,109	82.7	(80.2 - 85.1)
Gender						
Male	-	-	-	-	-	-
Female	419	80.9	(76.3 - 85.4)	23,109	82.7	(80.2 - 85.1)
Race/Ethnicity						
White Non-Hispanic	406	80.4	(75.7 - 85.1)	21,720	82.3	(80.0 - 84.6)
Black Non-Hispanic	*	*	*	473	87.0	(79.3 - 94.6)
Hispanic	*	*	*	527	83.4	(75.9 - 90.9)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	187	75.1	(67.4 - 82.7)	9,384	76.4	(71.3 - 81.5)
Some College	119	81.1	(72.1 - 90.1)	6,401	84.2	(80.5 - 88.0)
≥ College Degree	112	89.0	(82.9 - 95.1)	7,281	87.5	(84.3 - 90.6)
Income						
≤ \$24,999	*	*	*	6,441	82.7	(78.5 - 86.9)
\$25,000-\$49,999	107	84.5	(76.6 - 92.3)	5,740	79.2	(73.5 - 84.8)
\$50,000-\$74,999	*	*	*	3,296	85.7	(78.7 - 92.8)
≥ \$75,000	73	91.7	(83.2 - 100)	4,555	88.9	(85.5 - 92.3)

¹ Defined as having had a Pap test within the past 3 years among women

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 29a. Ever had a digital rectal exam among men aged 40 and older¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	205	18,500	72.9	(63.0 - 82.8)	12,033	3,176,500	73.8	(69.7 - 77.8)
Gender								
Male	205	18,500	72.9	(63.0 - 82.8)	12,033	3,176,500	73.8	(69.7 - 77.8)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	3,098	922,800	66.8	(59.9 - 73.8)
55-64	62	4,900	88.8	(80.7 - 97.0)	3,504	865,600	87.1	(82.5 - 91.7)
≥ 65	72	4,800	91.4	(84.0 - 98.8)	4,348	953,900	92.9	(91.0 - 94.9)
Race/Ethnicity								
White Non-Hispanic	197	17,500	76.7	(68.3 - 85.1)	11,411	2,452,100	80.3	(77.7 - 82.9)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	5,013	931,600	64.1	(57.3 - 71.0)
Some College	*	*	*	*	2,695	674,200	75.5	(67.8 - 83.2)
≥ College Degree	*	*	*	*	4,302	1,568,000	80.5	(74.0 - 87.0)
Income								
≤ \$24,999	*	*	*	*	2,573	573,300	65.9	(56.0 - 75.9)
\$25,000-\$49,999	*	*	*	*	3,168	530,100	74.6	(67.2 - 82.1)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,309	1,351,100	78.0	(72.6 - 83.5)

¹ Defined as having ever had a digital rectal exam among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have ever had a digital rectal exam (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 29b. Ever had a digital rectal exam among men aged 40 and older¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	*	*	*	12,033	73.7	(69.6 - 77.8)
Gender						
Male	*	*	*	12,033	73.7	(69.6 - 77.8)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	197	74.9	(65.3 - 84.5)	11,411	79.9	(77.2 - 82.5)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	5,013	64.7	(57.8 - 71.7)
Some College	*	*	*	2,695	75.0	(67.1 - 82.9)
≥ College Degree	*	*	*	4,302	80.3	(73.8 - 86.8)
Income						
≤ \$24,999	*	*	*	2,573	66.9	(56.9 - 76.9)
\$25,000-\$49,999	*	*	*	3,168	74.8	(67.3 - 82.2)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,309	77.5	(72.0 - 83.0)

¹ Defined as having ever had a digital rectal exam among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 30a. Had digital rectal exam within the past 2 years among men aged 40 and older¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	203	13,800	54.7	(45.2 - 64.2)	11,923	2,380,300	55.8	(51.8 - 59.8)
Gender								
Male	203	13,800	54.7	(45.2 - 64.2)	11,923	2,380,300	55.8	(51.8 - 59.8)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	1,078	246,800	27.5	(18.5 - 36.5)
45-54	*	*	*	*	3,074	650,000	47.6	(40.8 - 54.4)
55-64	*	*	*	*	3,467	673,500	68.1	(62.4 - 73.8)
≥ 65	*	*	*	*	4,304	809,900	79.7	(75.0 - 84.3)
Race/Ethnicity								
White Non-Hispanic	195	13,000	57.5	(48.5 - 66.4)	11,305	1,875,600	62.2	(59.2 - 65.2)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	4,963	718,700	49.6	(43.2 - 56.1)
Some College	*	*	*	*	2,665	511,500	58.5	(50.8 - 66.3)
≥ College Degree	*	*	*	*	4,277	1,148,400	59.4	(52.8 - 66.0)
Income								
≤ \$24,999	*	*	*	*	2,553	388,000	45.0	(35.0 - 55.0)
\$25,000-\$49,999	*	*	*	*	3,139	422,400	60.5	(53.0 - 68.0)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,295	978,300	56.9	(51.1 - 62.7)

¹ Defined as having had a digital rectal exam within the past 2 years among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have had a digital rectal exam within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 30b. Had digital rectal exam within the past 2 years among men aged 40 and older¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	*	*	*	11,923	55.9	(51.7 - 60.0)
Gender						
Male	*	*	*	11,923	55.9	(51.7 - 60.0)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	195	55.1	(45.3 - 64.9)	11,305	62.0	(58.9 - 65.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	4,963	50.5	(43.7 - 57.3)
Some College	*	*	*	2,665	58.3	(50.4 - 66.2)
≥ College Degree	*	*	*	4,277	59.2	(52.4 - 65.9)
Income						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,139	60.8	(53.1 - 68.4)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,295	56.5	(50.5 - 62.4)

¹ Defined as having had a digital rectal exam within the past 2 years among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 31a. Ever had a prostate specific antigen test among men aged 40 and older¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	194	13,100	55.5	(45.8 - 65.3)	11,557	2,822,600	67.6	(63.9 - 71.2)
Gender								
Male	194	13,100	55.5	(45.8 - 65.3)	11,557	2,822,600	67.6	(63.9 - 71.2)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,938	760,600	57.4	(50.6 - 64.3)
55-64	*	*	*	*	3,373	811,400	83.4	(78.8 - 88.1)
≥ 65	71	4,500	87.0	(78.4 - 95.7)	4,215	905,400	90.4	(87.1 - 93.6)
Race/Ethnicity								
White Non-Hispanic	187	12,400	58.9	(49.9 - 67.9)	10,950	1,965,400	67.0	(63.9 - 70.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	4,775	829,200	59.0	(52.4 - 65.6)
Some College	*	*	*	*	2,584	535,000	61.9	(53.7 - 70.2)
≥ College Degree	*	*	*	*	4,176	1,455,700	76.8	(72.5 - 81.1)
Income								
≤ \$24,999	*	*	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	*	3,048	444,100	64.1	(56.0 - 72.3)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,188	1,211,100	72.3	(67.6 - 77.0)

¹ Defined as having ever had a prostate specific antigen test among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have ever had a prostate specific antigen test (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 31b. Ever had a prostate specific antigen test among men aged 40 and older¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	*	*	*	11,557	67.4	(63.7 - 71.1)
Gender						
Male	*	*	*	11,557	67.4	(63.7 - 71.1)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	187	57.4	(47.6 - 67.2)	10,950	66.6	(63.4 - 69.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	4,775	59.7	(52.9 - 66.4)
Some College	*	*	*	2,584	61.4	(53.0 - 69.9)
≥ College Degree	*	*	*	4,176	76.3	(71.9 - 80.7)
Income						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,048	64.2	(56.1 - 72.4)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,188	71.5	(66.6 - 76.3)

¹ Defined as having ever had a prostate specific antigen test among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 32a. Had prostate specific antigen test within the past 2 years among men aged 40 and older¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	191	11,800	50.5	(41.0 - 60.1)	11,437	2,493,800	59.9	(56.0 - 63.9)
Gender								
Male	191	11,800	50.5	(41.0 - 60.1)	11,437	2,493,800	59.9	(56.0 - 63.9)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,917	659,400	49.9	(42.8 - 57.0)
55-64	*	*	*	*	3,326	719,700	74.5	(69.4 - 79.7)
≥ 65	71	4,300	82.5	(72.8 - 92.1)	4,166	830,000	83.3	(78.4 - 88.3)
Race/Ethnicity								
White Non-Hispanic	184	11,200	54.0	(45.0 - 63.0)	10,836	1,762,200	60.4	(57.3 - 63.5)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	4,727	734,200	52.4	(45.7 - 59.1)
Some College	*	*	*	*	2,558	483,800	56.3	(48.2 - 64.4)
≥ College Degree	*	*	*	*	4,133	1,273,200	67.4	(61.6 - 73.3)
Income								
≤ \$24,999	*	*	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	*	3,021	410,800	59.6	(51.5 - 67.6)
\$50,000-\$74,999	*	*	*	*	*	*	*	*
≥ \$75,000	*	*	*	*	3,170	1,080,100	64.6	(59.1 - 70.0)

¹ Defined as having had a prostate specific antigen test within the past 2 years among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have had a prostate specific antigen test within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 32b. Had prostate specific antigen test within the past 2 years among men aged 40 and older¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	*	*	*	11,437	59.8	(55.8 - 63.9)
Gender						
Male	*	*	*	11,437	59.8	(55.8 - 63.9)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	184	53.0	(43.3 - 62.8)	10,836	60.1	(56.9 - 63.4)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	4,727	52.9	(45.9 - 59.9)
Some College	*	*	*	2,558	56.0	(47.7 - 64.2)
≥ College Degree	*	*	*	4,133	67.1	(61.2 - 73.1)
Income						
≤ \$24,999	*	*	*	*	*	*
\$25,000-\$49,999	*	*	*	3,021	59.8	(51.7 - 67.9)
\$50,000-\$74,999	*	*	*	*	*	*
≥ \$75,000	*	*	*	3,170	63.9	(58.3 - 69.6)

¹ Defined as having had a prostate specific antigen test within the past 2 years among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 33a. Prostate cancer among men aged 40 and older¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	206	800	3.2	(1.2 - 5.1)	12,133	178,200	4.1	(3.0 - 5.2)
Gender								
Male	206	800	3.2	(1.2 - 5.1)	12,133	178,200	4.1	(3.0 - 5.2)
Female	-	-	-	-	-	-	-	-
Age (years)								
40-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	3,120	6,100	0.4	(0.0 - 0.9)
55-64	*	*	*	*	3,532	31,600	3.1	(1.5 - 4.8)
≥ 65	*	*	*	*	4,388	140,400	13.6	(9.7 - 17.4)
Race/Ethnicity								
White Non-Hispanic	198	800	3.5	(1.4 - 5.7)	11,502	143,700	4.7	(3.5 - 5.8)
Black Non-Hispanic	*	*	*	*	184	23,500	5.1	(0.0 - 11.0)
Hispanic	*	*	*	*	197	7,700	1.6	(0.0 - 3.7)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	*	*	*	*	5,065	46,300	3.1	(2.2 - 4.1)
Some College	*	*	*	*	2,709	39,200	4.4	(1.3 - 7.5)
≥ College Degree	*	*	*	*	4,336	92,500	4.7	(3.0 - 6.5)
Income								
≤ \$24,999	*	*	*	*	2,597	35,800	4.1	(2.2 - 6.0)
\$25,000-\$49,999	*	*	*	*	3,196	29,600	4.1	(2.9 - 5.4)
\$50,000-\$74,999	*	*	*	*	2,007	20,300	2.8	(1.2 - 4.4)
≥ \$75,000	*	*	*	*	3,322	56,500	3.2	(1.7 - 4.8)

¹ Defined as having ever been told by a health professional that respondent had prostate cancer among men aged 40 and older

² Sample size from survey

³ Estimated number of men aged 40 and older who have ever been told they had prostate cancer (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 33b. Prostate cancer among men aged 40 and older¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	206	3.3	(1.2 - 5.3)	12,133	4.5	(3.2 - 5.7)
Gender						
Male	206	3.3	(1.2 - 5.3)	12,133	4.5	(3.2 - 5.7)
Female	-	-	-	-	-	-
Race/Ethnicity						
White Non-Hispanic	198	3.6	(1.4 - 5.9)	11,502	5.1	(3.8 - 6.4)
Black Non-Hispanic	*	*	*	184	5.7	(0.0 - 12.8)
Hispanic	*	*	*	197	1.8	(0.0 - 4.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	5,065	3.3	(2.3 - 4.3)
Some College	*	*	*	2,709	5.0	(1.1 - 9.0)
≥ College Degree	*	*	*	4,336	5.2	(3.1 - 7.2)
Income						
≤ \$24,999	*	*	*	2,597	4.5	(2.2 - 6.7)
\$25,000-\$49,999	*	*	*	3,196	4.3	(3.0 - 5.6)
\$50,000-\$74,999	*	*	*	2,007	3.0	(1.1 - 5.0)
≥ \$75,000	*	*	*	3,322	3.5	(1.6 - 5.4)

¹ Defined as having ever been told by a health professional that respondent had prostate cancer among men aged 40 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wayne County**

Table 34a. Home blood stool test ever used among adults aged 50 and older¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	436	10,700	34.2	(29.2 - 39.3)	25,322	2,288,500	38.0	(35.5 - 40.4)
Gender								
Male	167	5,800	37.7	(29.5 - 45.8)	9,571	1,081,900	38.9	(35.1 - 42.6)
Female	269	4,900	30.9	(24.7 - 37.0)	15,751	1,206,600	37.2	(34.0 - 40.4)
Age (years)								
50-64	234	5,400	27.7	(21.2 - 34.2)	12,834	1,192,000	34.0	(30.8 - 37.2)
≥ 65	202	5,400	44.8	(37.4 - 52.3)	12,472	1,093,400	43.6	(39.8 - 47.4)
Race/Ethnicity								
White Non-Hispanic	426	10,700	35.6	(30.5 - 40.7)	24,213	1,754,500	39.1	(37.1 - 41.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	226	5,200	32.0	(25.1 - 38.9)	11,146	823,500	34.9	(30.5 - 39.2)
Some College	*	*	*	*	6,261	480,300	37.0	(32.6 - 41.4)
≥ College Degree	*	*	*	*	7,851	973,700	41.6	(37.8 - 45.3)
Income								
≤ \$24,999	131	2,600	35.8	(26.6 - 45.0)	7,222	505,000	35.4	(29.7 - 41.2)
\$25,000-\$49,999	120	2,800	32.5	(23.2 - 41.8)	6,609	467,600	40.3	(35.2 - 45.4)
\$50,000-\$74,999	*	*	*	*	3,500	313,000	38.2	(32.6 - 43.9)
≥ \$75,000	*	*	*	*	4,647	657,800	37.3	(33.0 - 41.6)

¹ Defined as having ever used a home blood stool test among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have ever used a home blood stool test (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 34b. Home blood stool test ever used among adults aged 50 and older¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	436	35.0	(29.9 - 40.2)	25,322	38.1	(35.5 - 40.8)
Gender						
Male	167	38.4	(30.3 - 46.6)	9,571	39.4	(35.3 - 43.5)
Female	269	31.6	(25.4 - 37.8)	15,751	36.9	(33.5 - 40.3)
Race/Ethnicity						
White Non-Hispanic	426	36.3	(31.2 - 41.5)	24,213	38.9	(36.7 - 41.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	226	32.9	(25.8 - 39.9)	11,146	35.1	(30.3 - 39.9)
Some College	*	*	*	6,261	36.6	(32.0 - 41.3)
≥ College Degree	*	*	*	7,851	41.9	(37.9 - 46.0)
Income						
≤ \$24,999	131	36.5	(27.0 - 45.9)	7,222	35.9	(29.5 - 42.4)
\$25,000-\$49,999	120	32.8	(23.4 - 42.2)	6,609	40.2	(34.8 - 45.6)
\$50,000-\$74,999	*	*	*	3,500	38.1	(31.9 - 44.3)
≥ \$75,000	*	*	*	4,647	37.5	(32.9 - 42.2)

¹ Defined as having ever used a home blood stool test among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 35a. Home blood stool test used within the past year among adults aged 50 and older¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	434	4,000	12.9	(9.3 - 16.4)	25,074	863,300	14.4	(12.6 - 16.2)
Gender								
Male	165	2,200	14.1	(8.3 - 20.0)	9,508	444,400	16.0	(13.2 - 18.9)
Female	269	1,800	11.6	(7.6 - 15.7)	15,566	418,900	13.1	(10.8 - 15.4)
Age (years)								
50-64	233	2,300	12.1	(7.4 - 16.8)	12,759	454,400	13.0	(10.6 - 15.5)
≥ 65	201	1,700	14.2	(9.0 - 19.4)	12,299	406,400	16.4	(13.7 - 19.0)
Race/Ethnicity								
White Non-Hispanic	424	4,000	13.4	(9.8 - 17.0)	23,971	634,300	14.3	(12.8 - 15.7)
Black Non-Hispanic	*	*	*	*	396	136,000	18.6	(9.4 - 27.8)
Hispanic	*	*	*	*	349	53,400	9.9	(2.5 - 17.2)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	224	1,800	11.3	(6.9 - 15.8)	11,026	288,600	12.3	(9.4 - 15.3)
Some College	102	900	13.1	(5.8 - 20.4)	6,208	212,500	16.4	(12.5 - 20.4)
≥ College Degree	107	1,300	15.8	(7.6 - 23.9)	7,778	352,400	15.2	(12.5 - 17.9)
Income								
≤ \$24,999	129	1,200	16.9	(9.7 - 24.2)	7,157	196,300	13.9	(10.1 - 17.6)
\$25,000-\$49,999	120	1,100	12.4	(5.9 - 18.9)	6,562	221,500	19.2	(14.1 - 24.3)
\$50,000-\$74,999	*	*	*	*	3,469	111,900	13.7	(10.1 - 17.4)
≥ \$75,000	*	*	*	*	4,623	205,100	11.7	(8.8 - 14.6)

¹ Defined as having used a home blood stool test within the past year among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have used a home blood stool test within the past year (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 35b. Home blood stool test used within the past year among adults aged 50 and older¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	434	12.8	(9.3 - 16.3)	25,074	14.6	(12.6 - 16.7)
Gender						
Male	165	13.7	(8.1 - 19.4)	9,508	16.4	(13.2 - 19.6)
Female	269	11.8	(7.7 - 16.0)	15,566	12.9	(10.4 - 15.4)
Race/Ethnicity						
White Non-Hispanic	424	13.3	(9.7 - 16.9)	23,971	14.2	(12.7 - 15.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	349	10.2	(1.7 - 18.7)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	224	11.4	(6.9 - 15.8)	11,026	12.7	(9.2 - 16.1)
Some College	102	13.1	(5.8 - 20.4)	6,208	16.6	(12.3 - 20.9)
≥ College Degree	107	15.4	(7.5 - 23.4)	7,778	15.3	(12.4 - 18.3)
Income						
≤ \$24,999	129	16.2	(9.2 - 23.2)	7,157	14.3	(9.8 - 18.7)
\$25,000-\$49,999	120	12.3	(5.8 - 18.7)	6,562	19.3	(13.8 - 24.8)
\$50,000-\$74,999	*	*	*	3,469	13.8	(9.8 - 17.7)
≥ \$75,000	*	*	*	4,623	11.8	(8.6 - 15.0)

¹ Defined as having used a home blood stool test within the past year among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wayne County**

Table 36a. Home blood stool test used within the past 2 years among adults aged 50 and older¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	434	5,800	18.6	(14.5 - 22.7)	25,074	1,218,000	20.4	(18.3 - 22.5)
Gender								
Male	165	3,200	20.6	(13.9 - 27.3)	9,508	593,100	21.4	(18.3 - 24.5)
Female	269	2,600	16.7	(11.8 - 21.5)	15,566	624,900	19.5	(16.6 - 22.4)
Age (years)								
50-64	233	3,300	16.8	(11.4 - 22.2)	12,759	629,900	18.1	(15.4 - 20.7)
≥ 65	201	2,500	21.4	(15.2 - 27.7)	12,299	585,500	23.6	(20.2 - 27.0)
Race/Ethnicity								
White Non-Hispanic	424	5,800	19.3	(15.1 - 23.6)	23,971	877,500	19.7	(18.2 - 21.3)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	224	2,500	15.5	(10.3 - 20.6)	11,026	454,600	19.4	(15.6 - 23.3)
Some College	102	1,400	20.1	(11.4 - 28.8)	6,208	262,200	20.3	(16.3 - 24.2)
≥ College Degree	107	1,900	23.5	(14.3 - 32.7)	7,778	491,400	21.2	(18.1 - 24.2)
Income								
≤ \$24,999	129	1,600	22.7	(14.5 - 30.9)	7,157	263,900	18.6	(14.6 - 22.7)
\$25,000-\$49,999	120	1,400	16.8	(9.4 - 24.1)	6,562	285,900	24.8	(19.6 - 30.0)
\$50,000-\$74,999	64	1,000	18.0	(8.2 - 27.7)	3,469	167,500	20.6	(15.6 - 25.5)
≥ \$75,000	*	*	*	*	4,623	294,900	16.8	(13.7 - 20.0)

¹ Defined as having used a home blood stool test within the past 2 years among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have used a home blood stool test within the past 2 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

**Table 36b. Home blood stool test used within the past 2 years among adults aged 50 and older¹
in Wayne County and New York State - Age-Adjusted²**

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	434	18.6	(14.5 - 22.7)	25,074	20.5	(18.2 - 22.8)
Gender						
Male	165	20.2	(13.6 - 26.7)	9,508	21.7	(18.3 - 25.1)
Female	269	17.1	(12.1 - 22.0)	15,566	19.4	(16.3 - 22.5)
Race/Ethnicity						
White Non-Hispanic	424	19.3	(15.1 - 23.5)	23,971	19.7	(18.0 - 21.3)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	224	15.6	(10.4 - 20.8)	11,026	19.5	(15.4 - 23.7)
Some College	102	19.9	(11.3 - 28.6)	6,208	20.4	(16.1 - 24.7)
≥ College Degree	107	23.8	(14.6 - 33.0)	7,778	21.4	(18.1 - 24.7)
Income						
≤ \$24,999	129	22.4	(14.1 - 30.6)	7,157	18.9	(14.2 - 23.6)
\$25,000-\$49,999	120	16.6	(9.3 - 23.9)	6,562	24.8	(19.3 - 30.3)
\$50,000-\$74,999	64	17.8	(8.1 - 27.6)	3,469	20.6	(15.3 - 25.9)
≥ \$75,000	*	*	*	4,623	16.9	(13.4 - 20.4)

¹ Defined as having used a home blood stool test within the past 2 years among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 37a. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	437	20,600	65.0	(59.5 - 70.4)	25,357	4,019,100	66.6	(64.1 - 69.1)
Gender								
Male	166	10,300	66.4	(57.5 - 75.2)	9,582	1,894,100	68.0	(64.2 - 71.7)
Female	271	10,200	63.6	(57.0 - 70.1)	15,775	2,125,000	65.5	(62.1 - 68.8)
Age (years)								
50-54	*	*	*	*	4,179	718,900	50.6	(44.6 - 56.5)
55-64	161	7,800	70.4	(62.8 - 77.9)	8,695	1,493,500	71.0	(67.3 - 74.8)
≥ 65	198	9,100	77.8	(71.6 - 84.0)	12,466	1,802,000	72.1	(68.6 - 75.7)
Race/Ethnicity								
White Non-Hispanic	427	20,100	66.1	(61.0 - 71.2)	24,248	3,093,900	68.8	(66.8 - 70.8)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	222	9,700	60.7	(52.6 - 68.8)	11,121	1,445,100	61.2	(56.7 - 65.8)
Some College	*	*	*	*	6,287	835,300	64.3	(59.7 - 69.0)
≥ College Degree	110	6,200	75.0	(65.9 - 84.0)	7,884	1,732,400	73.7	(70.0 - 77.4)
Income								
≤ \$24,999	129	4,300	59.7	(50.3 - 69.1)	7,220	892,600	62.7	(56.8 - 68.6)
\$25,000-\$49,999	120	5,900	69.4	(60.4 - 78.5)	6,619	697,300	60.1	(54.6 - 65.7)
\$50,000-\$74,999	*	*	*	*	3,514	613,700	74.2	(68.9 - 79.6)
≥ \$75,000	*	*	*	*	4,665	1,294,400	73.3	(69.2 - 77.5)

¹ Defined as having ever had a sigmoidoscopy or colonoscopy among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have ever had a sigmoidoscopy or colonoscopy (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 37b. Ever had sigmoidoscopy or colonoscopy among adults aged 50 and older¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	437	65.6	(60.2 - 71.0)	25,357	66.6	(63.9 - 69.3)
Gender						
Male	166	67.6	(59.0 - 76.1)	9,582	68.5	(64.5 - 72.5)
Female	271	63.7	(57.1 - 70.3)	15,775	64.8	(61.2 - 68.4)
Race/Ethnicity						
White Non-Hispanic	427	66.8	(61.7 - 71.8)	24,248	68.7	(66.5 - 70.8)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	222	61.6	(53.6 - 69.5)	11,121	61.3	(56.4 - 66.2)
Some College	*	*	*	6,287	63.8	(58.8 - 68.8)
≥ College Degree	110	75.2	(66.2 - 84.2)	7,884	74.0	(70.2 - 77.9)
Income						
≤ \$24,999	129	59.8	(50.3 - 69.3)	7,220	63.0	(56.7 - 69.3)
\$25,000-\$49,999	120	70.5	(61.5 - 79.5)	6,619	59.7	(53.8 - 65.6)
\$50,000-\$74,999	*	*	*	3,514	74.4	(68.3 - 80.5)
≥ \$75,000	*	*	*	4,665	73.6	(69.3 - 78.0)

¹ Defined as having ever had a sigmoidoscopy or colonoscopy among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wayne County**

Table 38a. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	433	19,500	62.1	(56.5 - 67.6)	25,169	3,855,800	64.6	(62.1 - 67.1)
Gender								
Male	163	10,000	65.2	(56.3 - 74.2)	9,521	1,835,400	66.4	(62.6 - 70.1)
Female	270	9,500	59.1	(52.4 - 65.7)	15,648	2,020,400	63.1	(59.8 - 66.4)
Age (years)								
50-54	*	*	*	*	4,169	700,500	49.3	(43.3 - 55.3)
55-64	161	7,700	69.0	(61.4 - 76.7)	8,652	1,455,100	69.4	(65.7 - 73.2)
≥ 65	194	8,500	74.2	(67.7 - 80.8)	12,331	1,695,700	69.5	(65.9 - 73.1)
Race/Ethnicity								
White Non-Hispanic	423	19,000	63.1	(57.9 - 68.3)	24,069	2,981,500	66.6	(64.6 - 68.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	219	9,100	57.5	(49.4 - 65.7)	11,017	1,360,400	59.0	(54.5 - 63.5)
Some College	*	*	*	*	6,251	807,100	62.4	(57.8 - 67.1)
≥ College Degree	109	6,100	74.7	(65.6 - 83.9)	7,841	1,682,500	71.7	(68.0 - 75.4)
Income								
≤ \$24,999	128	4,000	56.3	(46.8 - 65.8)	7,163	858,900	61.0	(55.1 - 67.0)
\$25,000-\$49,999	119	5,600	67.0	(57.7 - 76.3)	6,584	664,900	57.7	(52.2 - 63.2)
\$50,000-\$74,999	*	*	*	*	3,501	602,600	73.1	(67.7 - 78.5)
≥ \$75,000	*	*	*	*	4,655	1,260,400	71.5	(67.3 - 75.7)

¹ Defined as having had a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have had a sigmoidoscopy or colonoscopy within the past 10 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 38b. Sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	433	62.6	(57.1 - 68.1)	25,169	64.6	(61.9 - 67.3)
Gender						
Male	163	66.2	(57.5 - 74.9)	9,521	66.9	(62.9 - 70.9)
Female	270	59.1	(52.4 - 65.8)	15,648	62.4	(58.9 - 66.0)
Race/Ethnicity						
White Non-Hispanic	423	63.6	(58.4 - 68.8)	24,069	66.5	(64.3 - 68.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	219	58.2	(50.2 - 66.3)	11,017	59.0	(54.1 - 63.9)
Some College	*	*	*	6,251	61.9	(56.9 - 66.9)
≥ College Degree	109	74.9	(65.8 - 84.0)	7,841	72.1	(68.2 - 75.9)
Income						
≤ \$24,999	128	56.4	(46.7 - 66.0)	7,163	61.3	(54.9 - 67.6)
\$25,000-\$49,999	119	67.9	(58.7 - 77.1)	6,584	57.2	(51.4 - 63.0)
\$50,000-\$74,999	*	*	*	3,501	73.3	(67.2 - 79.4)
≥ \$75,000	*	*	*	4,655	71.8	(67.4 - 76.2)

¹ Defined as having had a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 39a. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	426	20,100	65.3	(59.8 - 70.9)	24,792	4,049,200	69.1	(66.7 - 71.6)
Gender								
Male	162	10,400	68.4	(59.4 - 77.4)	9,418	1,945,200	70.7	(67.0 - 74.3)
Female	264	9,700	62.3	(55.6 - 68.9)	15,374	2,104,100	67.8	(64.6 - 71.1)
Age (years)								
50-54	*	*	*	*	4,132	751,200	54.1	(48.0 - 60.1)
55-64	154	8,000	74.6	(67.3 - 81.9)	8,366	1,487,600	73.7	(70.0 - 77.4)
≥ 65	195	8,700	75.8	(69.4 - 82.2)	12,294	1,810,500	74.0	(70.6 - 77.4)
Race/Ethnicity								
White Non-Hispanic	416	19,700	66.5	(61.3 - 71.7)	23,726	3,091,400	70.4	(68.4 - 72.4)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	218	9,700	61.6	(53.4 - 69.8)	10,872	1,461,600	64.2	(59.8 - 68.7)
Some College	*	*	*	*	6,170	873,300	68.4	(63.9 - 72.8)
≥ College Degree	107	6,200	76.3	(67.3 - 85.4)	7,708	1,709,000	74.6	(70.9 - 78.3)
Income								
≤ \$24,999	127	4,500	63.2	(53.9 - 72.4)	7,092	923,300	66.0	(60.1 - 71.9)
\$25,000-\$49,999	119	5,700	68.6	(59.5 - 77.8)	6,526	730,000	64.2	(58.9 - 69.6)
\$50,000-\$74,999	*	*	*	*	3,467	621,500	76.6	(71.4 - 81.9)
≥ \$75,000	*	*	*	*	4,615	1,292,500	73.8	(69.8 - 77.9)

¹ Defined as having used a home blood stool test within the past year or a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 years and older

² Sample size from survey

³ Estimated number of adults aged 50 and older who have used a home blood stool test in the past year or sigmoidoscopy or colonoscopy within the past 10 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wayne County**

Table 39b. Home blood stool test within the past year or sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 and older¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	426	65.8	(60.3 - 71.2)	24,792	69.3	(66.7 - 71.9)
Gender						
Male	162	69.3	(60.6 - 78.0)	9,418	71.5	(67.6 - 75.4)
Female	264	62.2	(55.5 - 68.9)	15,374	67.2	(63.7 - 70.7)
Race/Ethnicity						
White Non-Hispanic	416	66.9	(61.8 - 72.1)	23,726	70.3	(68.1 - 72.4)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	218	62.2	(54.2 - 70.2)	10,872	64.5	(59.8 - 69.3)
Some College	*	*	*	6,170	68.2	(63.4 - 72.9)
≥ College Degree	107	76.4	(67.4 - 85.4)	7,708	75.0	(71.2 - 78.8)
Income						
≤ \$24,999	127	62.9	(53.5 - 72.3)	7,092	66.4	(60.2 - 72.6)
\$25,000-\$49,999	119	69.4	(60.3 - 78.6)	6,526	63.8	(58.2 - 69.5)
\$50,000-\$74,999	*	*	*	3,467	76.7	(70.6 - 82.7)
≥ \$75,000	*	*	*	4,615	74.4	(70.2 - 78.6)

¹ Defined as having used a home blood stool test within the past year or a sigmoidoscopy or colonoscopy within the past 10 years among adults aged 50 years and older

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 40a. High blood pressure among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	655	19,000	27.5	(23.4 - 31.5)	38,125	3,974,800	26.8	(24.9 - 28.7)
Gender								
Male	236	9,800	28.9	(22.1 - 35.7)	14,611	1,960,500	27.7	(24.7 - 30.7)
Female	419	9,200	26.1	(21.6 - 30.6)	23,514	2,014,300	26.0	(23.6 - 28.3)
Age (years)								
18-34	*	*	*	*	3,935	312,200	7.0	(4.2 - 9.8)
35-44	100	2,400	12.8	(3.9 - 21.8)	4,984	426,200	14.8	(11.6 - 18.1)
45-54	141	4,100	26.9	(18.9 - 34.9)	7,825	862,400	30.3	(25.9 - 34.8)
55-64	162	5,600	49.6	(41.2 - 58.0)	8,765	890,800	41.6	(37.6 - 45.7)
≥ 65	201	6,600	55.8	(48.3 - 63.3)	12,616	1,483,200	58.5	(54.7 - 62.3)
Race/Ethnicity								
White Non-Hispanic	631	18,100	27.9	(23.9 - 32.0)	35,814	2,650,900	28.4	(26.9 - 30.0)
Black Non-Hispanic	*	*	*	*	736	558,500	26.2	(19.6 - 32.8)
Hispanic	*	*	*	*	863	538,100	23.8	(17.0 - 30.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	310	9,600	30.4	(24.3 - 36.5)	15,688	1,851,200	33.1	(29.4 - 36.9)
Some College	165	4,800	26.8	(18.5 - 35.1)	9,906	851,100	24.1	(20.9 - 27.3)
≥ College Degree	178	4,600	23.8	(16.6 - 31.0)	12,452	1,253,400	22.3	(19.7 - 24.9)
Income								
≤ \$24,999	*	*	*	*	9,640	1,153,000	32.8	(27.6 - 38.1)
\$25,000-\$49,999	179	4,500	25.8	(19.2 - 32.5)	9,657	807,800	27.9	(23.7 - 32.1)
\$50,000-\$74,999	*	*	*	*	5,773	507,600	25.8	(21.4 - 30.1)
≥ \$75,000	131	2,800	15.3	(8.6 - 22.1)	8,560	942,500	21.0	(18.4 - 23.6)

¹ Defined as having ever been told by a health professional that respondent has high blood pressure

² Sample size from survey

³ Estimated number of adults who have ever been told by a health professional that they have high blood pressure (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 40b. High blood pressure among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	655	24.6	(20.5 - 28.7)	38,125	25.7	(23.8 - 27.5)
Gender						
Male	236	26.6	(19.6 - 33.7)	14,611	27.7	(24.7 - 30.7)
Female	419	22.6	(18.3 - 26.9)	23,514	23.8	(21.5 - 26.0)
Race/Ethnicity						
White Non-Hispanic	631	25.0	(20.9 - 29.1)	35,814	27.2	(25.6 - 28.7)
Black Non-Hispanic	*	*	*	736	24.9	(18.4 - 31.3)
Hispanic	*	*	*	863	23.3	(16.6 - 30.0)
Other Non-Hispanic	*	*	*	712	19.3	(9.5 - 29.1)
Education						
≤ High School	310	27.8	(21.6 - 34.0)	15,688	32.1	(28.4 - 35.8)
Some College	165	23.8	(15.3 - 32.4)	9,906	22.8	(19.7 - 25.9)
≥ College Degree	178	20.5	(13.4 - 27.6)	12,452	21.2	(18.6 - 23.8)
Income						
≤ \$24,999	*	*	*	9,640	31.8	(26.7 - 37.0)
\$25,000-\$49,999	179	22.4	(16.2 - 28.6)	9,657	26.9	(22.8 - 30.9)
\$50,000-\$74,999	*	*	*	5,773	24.8	(20.4 - 29.2)
≥ \$75,000	131	12.7	(6.1 - 19.3)	8,560	19.9	(17.3 - 22.5)

¹ Defined as having ever been told by a health professional that respondent has high blood pressure

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 41a. Blood pressure medication among adults with high blood pressure¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	251	15,800	82.9	(76.5 - 89.3)	15,288	3,145,800	79.2	(76.0 - 82.4)
Gender								
Male	*	*	*	*	6,080	1,432,100	73.1	(67.6 - 78.5)
Female	154	7,900	86.1	(79.4 - 92.8)	9,208	1,713,700	85.2	(82.1 - 88.2)
Age (years)								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	*	*	*	*	2,315	645,500	74.9	(66.8 - 82.9)
55-64	*	*	*	*	4,045	782,300	87.8	(84.5 - 91.1)
≥ 65	116	6,100	91.8	(87.1 - 96.6)	7,675	1,383,400	93.4	(91.1 - 95.7)
Race/Ethnicity								
White Non-Hispanic	243	15,100	83.4	(76.8 - 89.9)	14,442	2,117,300	79.9	(77.5 - 82.4)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	135	8,100	83.8	(76.4 - 91.2)	7,354	1,482,800	80.2	(74.7 - 85.6)
Some College	60	4,600	94.0	(88.9 - 99.1)	3,868	632,000	74.3	(67.8 - 80.7)
≥ College Degree	*	*	*	*	4,035	1,013,200	80.9	(76.8 - 84.9)
Income								
≤ \$24,999	85	4,600	86.4	(77.4 - 95.3)	4,898	948,200	82.3	(75.7 - 88.9)
\$25,000-\$49,999	66	3,800	85.4	(76.4 - 94.4)	4,088	668,300	82.8	(77.6 - 88.0)
\$50,000-\$74,999	*	*	*	*	1,971	391,500	77.2	(67.7 - 86.6)
≥ \$75,000	*	*	*	*	2,397	683,700	72.5	(66.1 - 78.9)

¹ Defined as currently taking medicine for high blood pressure among adults with high blood pressure

² Sample size from survey

³ Estimated number of adults with high blood pressure who are taking medicine for high blood pressure (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 41b. Blood pressure medication among adults with high blood pressure¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	251	83.0	(75.8 - 90.2)	15,288	78.5	(75.2 - 81.7)
Gender						
Male	*	*	*	6,080	73.3	(67.9 - 78.7)
Female	154	85.4	(77.9 - 93.0)	9,208	84.0	(80.7 - 87.2)
Race/Ethnicity						
White Non-Hispanic	243	83.2	(75.7 - 90.6)	14,442	79.0	(76.5 - 81.6)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	135	84.4	(76.7 - 92.1)	7,354	79.7	(74.3 - 85.2)
Some College	60	94.5	(89.7 - 99.3)	3,868	72.5	(65.7 - 79.3)
≥ College Degree	*	*	*	4,035	80.4	(76.1 - 84.6)
Income						
≤ \$24,999	*	*	*	4,898	81.6	(75.0 - 88.3)
\$25,000-\$49,999	66	85.2	(76.3 - 94.1)	4,088	82.1	(76.7 - 87.5)
\$50,000-\$74,999	*	*	*	1,971	76.4	(66.7 - 86.1)
≥ \$75,000	*	*	*	2,397	71.6	(64.8 - 78.3)

¹ Defined as currently taking medicine for high blood pressure among adults with high blood pressure

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 42a. Ever had blood cholesterol checked among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	650	57,900	84.7	(80.1 - 89.3)	37,450	11,496,600	80.3	(77.7 - 83.0)
Gender								
Male	233	29,500	87.8	(80.6 - 94.9)	14,389	5,288,400	77.7	(73.3 - 82.0)
Female	417	28,400	81.7	(75.9 - 87.6)	23,061	6,208,200	82.7	(79.6 - 85.8)
Age (years)								
18-34	*	*	*	*	3,684	2,296,400	55.3	(48.7 - 61.9)
35-44	*	*	*	*	4,883	2,264,200	81.6	(76.6 - 86.5)
45-54	139	13,700	92.5	(87.6 - 97.4)	7,731	2,539,400	91.0	(87.0 - 94.9)
55-64	162	11,000	98.1	(96.0 - 100)	8,718	2,014,900	95.4	(93.5 - 97.2)
≥ 65	201	11,800	98.2	(96.6 - 99.9)	12,434	2,381,700	95.9	(94.3 - 97.6)
Race/Ethnicity								
White Non-Hispanic	626	55,600	87.0	(83.2 - 90.8)	35,198	7,832,600	86.3	(84.8 - 87.9)
Black Non-Hispanic	*	*	*	*	720	1,588,900	77.8	(68.4 - 87.2)
Hispanic	*	*	*	*	842	1,224,600	56.8	(46.8 - 66.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	307	24,900	79.0	(71.0 - 87.1)	15,342	3,795,700	70.6	(65.6 - 75.6)
Some College	164	15,400	88.0	(80.7 - 95.4)	9,732	2,599,800	77.4	(71.7 - 83.2)
≥ College Degree	177	17,600	92.4	(87.3 - 97.5)	12,296	5,008,300	91.3	(88.9 - 93.6)
Income								
≤ \$24,999	*	*	*	*	9,437	2,277,500	68.4	(61.5 - 75.3)
\$25,000-\$49,999	176	14,900	89.1	(82.4 - 95.7)	9,492	2,053,000	71.9	(64.5 - 79.3)
\$50,000-\$74,999	108	12,200	89.3	(82.6 - 96.0)	5,721	1,736,100	90.0	(87.7 - 92.3)
≥ \$75,000	130	15,700	86.8	(78.8 - 94.8)	8,460	4,019,300	92.2	(90.8 - 93.6)

¹ Defined as ever having blood cholesterol checked

² Sample size from survey

³ Estimated number of adults who have ever had blood cholesterol checked (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 42b. Ever had blood cholesterol checked among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	650	82.4	(77.1 - 87.7)	37,450	79.9	(77.3 - 82.5)
Gender						
Male	233	86.9	(79.0 - 94.8)	14,389	78.0	(73.9 - 82.1)
Female	417	78.1	(71.1 - 85.0)	23,061	81.6	(78.4 - 84.8)
Race/Ethnicity						
White Non-Hispanic	626	84.9	(80.4 - 89.4)	35,198	85.6	(84.0 - 87.2)
Black Non-Hispanic	*	*	*	720	77.5	(68.2 - 86.7)
Hispanic	*	*	*	842	57.2	(47.4 - 67.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	307	76.2	(67.1 - 85.2)	15,342	70.2	(65.2 - 75.1)
Some College	164	85.5	(76.6 - 94.4)	9,732	76.7	(71.1 - 82.3)
≥ College Degree	177	91.1	(85.1 - 97.2)	12,296	90.9	(88.6 - 93.2)
Income						
≤ \$24,999	*	*	*	9,437	68.1	(61.3 - 74.9)
\$25,000-\$49,999	176	86.9	(79.0 - 94.8)	9,492	71.6	(64.4 - 78.7)
\$50,000-\$74,999	108	87.8	(79.9 - 95.8)	5,721	89.6	(87.2 - 92.0)
≥ \$75,000	130	84.9	(75.8 - 94.0)	8,460	91.7	(90.1 - 93.2)

¹ Defined as ever having blood cholesterol checked

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 43a. Blood cholesterol checked within the past 5 years among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	647	54,900	80.7	(75.9 - 85.6)	37,201	11,035,200	77.8	(75.1 - 80.4)
Gender								
Male	232	27,800	82.6	(74.8 - 90.4)	14,307	5,020,700	74.5	(70.2 - 78.9)
Female	415	27,200	78.9	(73.0 - 84.8)	22,894	6,014,600	80.7	(77.6 - 83.8)
Age (years)								
18-34	*	*	*	*	3,647	2,179,800	53.3	(46.7 - 59.9)
35-44	*	*	*	*	4,862	2,161,200	78.0	(72.8 - 83.2)
45-54	138	13,100	89.2	(83.7 - 94.8)	7,697	2,408,600	86.6	(82.4 - 90.8)
55-64	162	10,900	97.1	(94.6 - 99.6)	8,677	1,957,900	93.3	(91.4 - 95.3)
≥ 65	199	11,400	96.5	(94.0 - 98.9)	12,318	2,327,800	94.8	(93.1 - 96.6)
Race/Ethnicity								
White Non-Hispanic	623	52,700	82.9	(78.6 - 87.1)	34,960	7,486,600	83.4	(81.8 - 85.0)
Black Non-Hispanic	*	*	*	*	719	1,561,700	76.5	(67.1 - 85.9)
Hispanic	*	*	*	*	838	1,184,000	55.6	(45.6 - 65.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	304	23,500	75.5	(67.2 - 83.7)	15,215	3,612,500	68.3	(63.2 - 73.3)
Some College	164	14,500	83.0	(74.0 - 91.9)	9,673	2,524,200	75.4	(69.8 - 81.1)
≥ College Degree	177	16,900	88.5	(82.9 - 94.2)	12,236	4,806,500	88.1	(85.5 - 90.6)
Income								
≤ \$24,999	*	*	*	*	9,350	2,214,800	66.9	(60.1 - 73.8)
\$25,000-\$49,999	176	13,600	80.8	(71.9 - 89.7)	9,454	1,936,600	68.8	(61.5 - 76.1)
\$50,000-\$74,999	108	11,500	84.6	(76.4 - 92.8)	5,701	1,640,400	85.9	(82.4 - 89.5)
≥ \$75,000	130	15,400	84.9	(76.9 - 93.0)	8,435	3,872,400	89.2	(87.5 - 90.9)

¹ Defined as having had blood cholesterol checked within the past 5 years

² Sample size from survey

³ Estimated number of adults who have had blood cholesterol checked within the past 5 years (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 43b. Blood cholesterol checked within the past 5 years among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	647	78.1	(72.6 - 83.6)	37,201	77.3	(74.7 - 79.9)
Gender						
Male	232	80.9	(72.2 - 89.6)	14,307	74.8	(70.7 - 79.0)
Female	415	75.3	(68.4 - 82.2)	22,894	79.5	(76.3 - 82.7)
Race/Ethnicity						
White Non-Hispanic	623	80.4	(75.5 - 85.4)	34,960	82.6	(81.0 - 84.3)
Black Non-Hispanic	*	*	*	719	76.2	(66.9 - 85.5)
Hispanic	*	*	*	838	55.9	(46.1 - 65.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	304	72.2	(63.0 - 81.4)	15,215	67.8	(62.9 - 72.8)
Some College	*	*	*	9,673	74.7	(69.1 - 80.2)
≥ College Degree	177	87.4	(80.9 - 93.9)	12,236	87.5	(85.0 - 90.1)
Income						
≤ \$24,999	*	*	*	9,350	66.6	(59.8 - 73.4)
\$25,000-\$49,999	*	*	*	9,454	68.5	(61.4 - 75.6)
\$50,000-\$74,999	108	82.7	(73.1 - 92.2)	5,701	85.4	(81.7 - 89.0)
≥ \$75,000	130	83.0	(73.8 - 92.2)	8,435	88.5	(86.7 - 90.3)

¹ Defined as having had blood cholesterol checked within the past 5 years

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 44a. Cardiovascular disease among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	651	6,100	8.8	(6.3 - 11.3)	37,810	1,135,000	7.7	(6.5 - 8.8)
Gender								
Male	233	3,300	9.8	(5.4 - 14.2)	14,477	664,600	9.4	(7.5 - 11.4)
Female	418	2,700	7.8	(5.3 - 10.3)	23,333	470,400	6.1	(4.8 - 7.4)
Age (years)								
18-34	*	*	*	*	3,930	100,800	2.3	(0.0 - 4.8)
35-44	*	*	*	*	4,971	103,400	3.6	(1.5 - 5.7)
45-54	*	*	*	*	7,768	131,300	4.6	(2.9 - 6.4)
55-64	160	1,300	11.5	(6.3 - 16.6)	8,699	224,200	10.6	(8.3 - 12.9)
≥ 65	200	2,700	22.5	(16.2 - 28.9)	12,442	575,300	23.0	(19.9 - 26.1)
Race/Ethnicity								
White Non-Hispanic	627	5,400	8.4	(6.0 - 10.8)	35,522	772,200	8.3	(7.4 - 9.2)
Black Non-Hispanic	*	*	*	*	728	140,700	6.6	(3.3 - 9.9)
Hispanic	*	*	*	*	854	163,300	7.3	(2.0 - 12.5)
Other Non-Hispanic	*	*	*	*	706	58,800	5.2	(0.9 - 9.5)
Education								
≤ High School	307	3,300	10.6	(7.1 - 14.1)	15,486	590,800	10.7	(8.2 - 13.2)
Some College	165	2,000	10.8	(4.0 - 17.6)	9,849	227,100	6.4	(4.7 - 8.1)
≥ College Degree	177	800	4.0	(1.5 - 6.5)	12,393	314,600	5.6	(4.3 - 6.9)
Income								
≤ \$24,999	157	2,600	20.2	(11.8 - 28.6)	9,505	448,100	12.9	(9.1 - 16.7)
\$25,000-\$49,999	178	1,200	6.8	(3.3 - 10.2)	9,571	179,300	6.2	(4.8 - 7.6)
\$50,000-\$74,999	*	*	*	*	5,751	166,900	8.5	(5.5 - 11.4)
≥ \$75,000	*	*	*	*	8,538	183,500	4.1	(2.9 - 5.3)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina or a stroke

² Sample size from survey

³ Estimated number of adults with cardiovascular disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 44b. Cardiovascular disease among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	651	8.1	(5.6 - 10.7)	37,810	7.6	(6.4 - 8.7)
Gender						
Male	233	9.7	(5.0 - 14.4)	14,477	9.7	(7.8 - 11.7)
Female	418	6.7	(4.4 - 8.9)	23,333	5.6	(4.3 - 6.9)
Race/Ethnicity						
White Non-Hispanic	627	7.7	(5.3 - 10.1)	35,522	8.2	(7.2 - 9.1)
Black Non-Hispanic	*	*	*	728	6.8	(3.2 - 10.4)
Hispanic	*	*	*	854	7.0	(2.0 - 12.0)
Other Non-Hispanic	*	*	*	706	5.5	(0.9 - 10.0)
Education						
≤ High School	307	9.8	(6.4 - 13.1)	15,486	10.5	(8.0 - 13.0)
Some College	165	10.3	(3.0 - 17.6)	9,849	6.4	(4.5 - 8.2)
≥ College Degree	177	3.6	(1.3 - 5.9)	12,393	5.5	(4.2 - 6.9)
Income						
≤ \$24,999	157	19.1	(10.3 - 27.9)	9,505	12.8	(9.0 - 16.6)
\$25,000-\$49,999	178	6.2	(3.0 - 9.4)	9,571	6.0	(4.7 - 7.4)
\$50,000-\$74,999	*	*	*	5,751	8.5	(5.4 - 11.6)
≥ \$75,000	*	*	*	8,538	4.0	(2.7 - 5.2)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina or a stroke

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wayne County**

Table 45a. Coronary heart disease among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	653	4,800	6.9	(4.9 - 9.0)	37,838	929,400	6.3	(5.3 - 7.3)
Gender								
Male	234	2,600	7.7	(4.2 - 11.1)	14,479	598,600	8.5	(6.6 - 10.4)
Female	419	2,200	6.2	(4.0 - 8.4)	23,359	330,800	4.3	(3.4 - 5.2)
Age (years)								
18-34	*	*	*	*	3,932	63,400	1.4	(0.0 - 3.6)
35-44	*	*	*	*	4,973	95,300	3.3	(1.2 - 5.4)
45-54	*	*	*	*	7,778	108,300	3.8	(2.1 - 5.6)
55-64	160	1,100	9.8	(5.0 - 14.6)	8,703	188,600	8.9	(6.7 - 11.1)
≥ 65	202	2,400	19.9	(13.9 - 25.9)	12,452	473,800	18.9	(16.1 - 21.7)
Race/Ethnicity								
White Non-Hispanic	629	4,200	6.5	(4.7 - 8.3)	35,552	636,700	6.9	(6.0 - 7.7)
Black Non-Hispanic	*	*	*	*	728	113,100	5.3	(2.3 - 8.4)
Hispanic	*	*	*	*	853	128,000	5.7	(1.1 - 10.3)
Other Non-Hispanic	*	*	*	*	705	51,500	4.6	(0.5 - 8.7)
Education								
≤ High School	307	2,800	8.8	(5.6 - 12.0)	15,485	474,500	8.6	(6.3 - 10.8)
Some College	166	1,200	6.9	(2.1 - 11.7)	9,860	193,000	5.5	(3.9 - 7.1)
≥ College Degree	178	800	4.0	(1.5 - 6.4)	12,411	259,700	4.6	(3.4 - 5.8)
Income								
≤ \$24,999	158	2,000	16.1	(8.3 - 23.9)	9,507	372,300	10.7	(7.3 - 14.1)
\$25,000-\$49,999	179	1,200	6.7	(3.3 - 10.1)	9,585	136,300	4.7	(3.8 - 5.7)
\$50,000-\$74,999	*	*	*	*	5,755	154,600	7.8	(4.9 - 10.7)
≥ \$75,000	*	*	*	*	8,541	151,800	3.4	(2.3 - 4.5)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina

² Sample size from survey

³ Estimated number of adults with coronary heart disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 45b. Coronary heart disease among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	653	6.3	(4.3 - 8.2)	37,838	6.3	(5.2 - 7.3)
Gender						
Male	234	7.2	(3.7 - 10.6)	14,479	8.8	(6.9 - 10.8)
Female	419	5.4	(3.4 - 7.3)	23,359	3.9	(3.0 - 4.8)
Race/Ethnicity						
White Non-Hispanic	629	5.8	(4.1 - 7.5)	35,552	6.8	(5.9 - 7.7)
Black Non-Hispanic	*	*	*	728	5.6	(2.2 - 9.0)
Hispanic	*	*	*	853	5.4	(1.1 - 9.6)
Other Non-Hispanic	*	*	*	705	4.8	(0.4 - 9.2)
Education						
≤ High School	307	8.0	(5.0 - 10.9)	15,485	8.5	(6.2 - 10.7)
Some College	166	6.2	(1.3 - 11.1)	9,860	5.5	(3.7 - 7.2)
≥ College Degree	178	3.6	(1.3 - 5.9)	12,411	4.7	(3.4 - 5.9)
Income						
≤ \$24,999	158	15.4	(7.2 - 23.6)	9,507	10.7	(7.2 - 14.1)
\$25,000-\$49,999	179	6.2	(3.0 - 9.3)	9,585	4.7	(3.6 - 5.7)
\$50,000-\$74,999	*	*	*	5,755	8.0	(4.8 - 11.1)
≥ \$75,000	*	*	*	8,541	3.3	(2.1 - 4.5)

¹ Defined as having ever been told by a health professional that respondent had a heart attack or angina

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 46a. Ever diagnosed with diabetes among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	658	6,700	9.7	(7.3 - 12.1)	38,174	1,382,700	9.3	(8.0 - 10.5)
Gender								
Male	236	3,600	10.6	(6.6 - 14.6)	14,626	530,400	7.5	(6.2 - 8.8)
Female	422	3,100	8.8	(6.1 - 11.5)	23,548	852,300	11.0	(8.9 - 13.0)
Age (years)								
18-34	*	*	*	*	3,941	150,200	3.4	(1.0 - 5.8)
35-44	*	*	*	*	4,987	165,100	5.7	(3.6 - 7.9)
45-54	141	1,800	12.1	(6.0 - 18.2)	7,830	279,000	9.8	(6.7 - 12.8)
55-64	162	2,100	18.3	(11.7 - 25.0)	8,771	332,900	15.5	(12.3 - 18.8)
≥ 65	204	1,900	15.7	(10.5 - 21.0)	12,645	455,600	17.9	(15.0 - 20.8)
Race/Ethnicity								
White Non-Hispanic	634	6,600	10.2	(7.7 - 12.7)	35,862	809,300	8.7	(7.8 - 9.5)
Black Non-Hispanic	*	*	*	*	739	261,000	12.1	(6.6 - 17.6)
Hispanic	*	*	*	*	862	227,300	10.1	(5.4 - 14.8)
Other Non-Hispanic	*	*	*	*	711	85,100	7.6	(2.3 - 12.8)
Education								
≤ High School	312	4,100	12.7	(8.7 - 16.7)	15,709	730,000	13.0	(10.5 - 15.6)
Some College	166	1,300	6.9	(3.4 - 10.4)	9,918	303,500	8.6	(5.9 - 11.2)
≥ College Degree	178	1,400	7.4	(2.9 - 12.0)	12,464	337,500	6.0	(4.8 - 7.2)
Income								
≤ \$24,999	162	2,100	16.5	(9.8 - 23.3)	9,652	543,400	15.5	(11.4 - 19.5)
\$25,000-\$49,999	179	1,400	8.3	(4.5 - 12.0)	9,667	291,000	10.0	(7.5 - 12.6)
\$50,000-\$74,999	108	1,400	10.2	(3.8 - 16.7)	5,777	151,000	7.6	(5.0 - 10.3)
≥ \$75,000	*	*	*	*	8,565	220,800	4.9	(3.8 - 6.1)

¹ Defined as having ever been told by a doctor that respondent had diabetes, excluding pre-diabetes and women with diabetes only when pregnant

² Sample size from survey

³ Estimated number of adults who have diabetes (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 46b. Ever diagnosed with diabetes among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	658	8.6	(6.3 - 10.9)	38,174	9.0	(7.8 - 10.3)
Gender						
Male	236	9.5	(5.6 - 13.3)	14,626	7.5	(6.1 - 8.9)
Female	422	7.8	(5.2 - 10.4)	23,548	10.4	(8.4 - 12.4)
Race/Ethnicity						
White Non-Hispanic	634	9.1	(6.7 - 11.5)	35,862	8.4	(7.5 - 9.2)
Black Non-Hispanic	*	*	*	739	11.4	(6.0 - 16.7)
Hispanic	*	*	*	862	10.1	(5.4 - 14.9)
Other Non-Hispanic	*	*	*	711	7.7	(2.3 - 13.0)
Education						
≤ High School	312	11.7	(7.9 - 15.6)	15,709	12.6	(10.0 - 15.2)
Some College	166	5.7	(2.8 - 8.7)	9,918	8.3	(5.6 - 10.9)
≥ College Degree	178	6.5	(2.0 - 10.9)	12,464	5.9	(4.7 - 7.2)
Income						
≤ \$24,999	162	15.3	(8.4 - 22.3)	9,652	15.2	(11.1 - 19.2)
\$25,000-\$49,999	179	7.3	(3.9 - 10.6)	9,667	9.7	(7.1 - 12.2)
\$50,000-\$74,999	108	9.4	(2.9 - 16.0)	5,777	7.6	(4.9 - 10.2)
≥ \$75,000	*	*	*	8,565	4.7	(3.6 - 5.9)

¹ Defined as having ever been told by a doctor that respondent had diabetes, excluding pre-diabetes and women with diabetes only when pregnant

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 47a. Ever diagnosed with asthma (lifetime) among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	658	8,600	12.4	(8.9 - 15.9)	38,096	2,225,100	15.0	(13.2 - 16.8)
Gender								
Male	236	4,400	13.0	(6.9 - 19.1)	14,601	875,800	12.4	(10.1 - 14.7)
Female	422	4,200	11.8	(8.3 - 15.2)	23,495	1,349,300	17.4	(14.6 - 20.1)
Age (years)								
18-34	*	*	*	*	3,931	825,100	18.5	(13.8 - 23.3)
35-44	100	3,300	17.4	(8.0 - 26.8)	4,972	431,700	15.1	(11.2 - 19.1)
45-54	141	1,100	7.1	(2.5 - 11.6)	7,819	365,900	12.8	(9.7 - 16.0)
55-64	162	1,300	11.6	(6.4 - 16.9)	8,756	316,700	14.8	(11.9 - 17.7)
≥ 65	204	1,300	10.3	(5.7 - 15.0)	12,618	285,700	11.2	(8.6 - 13.8)
Race/Ethnicity								
White Non-Hispanic	634	8,100	12.5	(8.9 - 16.1)	35,783	1,365,700	14.6	(13.0 - 16.3)
Black Non-Hispanic	*	*	*	*	737	367,000	17.2	(10.2 - 24.2)
Hispanic	*	*	*	*	863	317,100	14.1	(9.0 - 19.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	312	3,200	10.1	(6.3 - 14.0)	15,675	831,600	14.9	(12.0 - 17.8)
Some College	166	2,000	10.8	(4.0 - 17.5)	9,893	587,500	16.6	(13.1 - 20.1)
≥ College Degree	178	3,400	17.9	(9.4 - 26.3)	12,444	739,000	13.2	(10.4 - 15.9)
Income								
≤ \$24,999	162	1,400	11.0	(5.6 - 16.4)	9,643	708,900	20.2	(14.9 - 25.5)
\$25,000-\$49,999	179	1,700	9.6	(4.0 - 15.2)	9,641	389,800	13.4	(10.0 - 16.9)
\$50,000-\$74,999	108	2,100	15.4	(5.6 - 25.2)	5,762	257,800	13.2	(10.0 - 16.5)
≥ \$75,000	131	2,900	16.1	(7.8 - 24.5)	8,562	577,900	12.8	(10.4 - 15.3)

¹ Defined as adults having ever been told by a health professional that they had asthma

² Sample size from survey

³ Estimated number of adults who have lifetime asthma (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 47b. Ever diagnosed with asthma (lifetime) among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	658	12.8	(8.9 - 16.7)	38,096	15.2	(13.3 - 17.0)
Gender						
Male	236	14.2	(7.3 - 21.1)	14,601	12.6	(10.3 - 14.9)
Female	422	11.5	(7.7 - 15.3)	23,495	17.5	(14.8 - 20.3)
Race/Ethnicity						
White Non-Hispanic	634	12.9	(8.9 - 17.0)	35,783	14.9	(13.2 - 16.6)
Black Non-Hispanic	*	*	*	737	17.1	(10.1 - 24.1)
Hispanic	*	*	*	863	14.5	(9.3 - 19.7)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	312	9.9	(5.9 - 13.9)	15,675	15.1	(12.1 - 18.0)
Some College	166	11.3	(3.6 - 19.0)	9,893	16.9	(13.4 - 20.4)
≥ College Degree	178	19.3	(9.8 - 28.8)	12,444	13.3	(10.6 - 16.0)
Income						
≤ \$24,999	162	10.3	(4.7 - 15.8)	9,643	20.3	(15.0 - 25.6)
\$25,000-\$49,999	179	10.2	(3.9 - 16.5)	9,641	13.7	(10.2 - 17.2)
\$50,000-\$74,999	*	*	*	5,762	13.5	(10.0 - 17.0)
≥ \$75,000	131	17.1	(7.7 - 26.4)	8,562	13.0	(10.6 - 15.4)

¹ Defined as adults having ever been told by a health professional that they had asthma

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 48a. Current asthma among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	656	5,400	7.8	(5.3 - 10.3)	37,933	1,415,000	9.6	(8.1 - 11.0)
Gender								
Male	235	2,200	6.5	(2.6 - 10.5)	14,546	440,700	6.3	(4.8 - 7.7)
Female	421	3,200	9.0	(5.9 - 12.0)	23,387	974,300	12.6	(10.2 - 15.0)
Age (years)								
18-34	*	*	*	*	3,909	467,200	10.5	(6.8 - 14.3)
35-44	99	1,600	8.7	(2.8 - 14.6)	4,947	246,800	8.7	(5.8 - 11.6)
45-54	141	700	4.8	(1.8 - 7.8)	7,787	240,000	8.4	(6.1 - 10.8)
55-64	161	900	7.8	(3.7 - 11.8)	8,726	231,100	10.8	(8.1 - 13.5)
≥ 65	204	1,100	9.1	(4.6 - 13.5)	12,564	229,900	9.1	(6.5 - 11.6)
Race/Ethnicity								
White Non-Hispanic	632	5,100	7.8	(5.3 - 10.4)	35,629	888,400	9.6	(8.2 - 11.0)
Black Non-Hispanic	*	*	*	*	736	246,900	11.6	(5.7 - 17.5)
Hispanic	*	*	*	*	859	186,800	8.3	(4.4 - 12.2)
Other Non-Hispanic	*	*	*	*	709	93,000	8.3	(1.4 - 15.2)
Education								
≤ High School	310	2,100	6.5	(3.9 - 9.1)	15,594	595,100	10.7	(8.2 - 13.2)
Some College	166	1,400	7.5	(2.5 - 12.5)	9,861	322,000	9.1	(7.0 - 11.3)
≥ College Degree	178	2,000	10.3	(4.1 - 16.5)	12,394	430,900	7.7	(5.8 - 9.7)
Income								
≤ \$24,999	161	1,200	9.7	(4.6 - 14.7)	9,600	489,200	14.0	(9.7 - 18.3)
\$25,000-\$49,999	179	1,000	5.9	(1.7 - 10.1)	9,590	268,400	9.3	(6.2 - 12.4)
\$50,000-\$74,999	108	1,400	10.2	(2.8 - 17.6)	5,741	143,400	7.4	(5.1 - 9.6)
≥ \$75,000	131	1,400	7.9	(2.7 - 13.1)	8,541	339,200	7.6	(5.8 - 9.4)

¹ Defined as adults having ever been told by a health professional that they had asthma and still have asthma

² Sample size from survey

³ Estimated number of adults who currently have asthma (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 48b. Current asthma among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	656	7.9	(5.1 - 10.7)	37,933	9.7	(8.2 - 11.2)
Gender						
Male	235	7.1	(2.6 - 11.6)	14,546	6.4	(4.9 - 7.9)
Female	421	8.7	(5.4 - 12.0)	23,387	12.7	(10.2 - 15.1)
Race/Ethnicity						
White Non-Hispanic	632	7.9	(5.1 - 10.8)	35,629	9.7	(8.3 - 11.2)
Black Non-Hispanic	*	*	*	736	11.5	(5.6 - 17.3)
Hispanic	*	*	*	859	8.6	(4.5 - 12.6)
Other Non-Hispanic	*	*	*	709	8.3	(1.3 - 15.3)
Education						
≤ High School	310	6.1	(3.4 - 8.8)	15,594	10.8	(8.3 - 13.3)
Some College	166	7.7	(2.0 - 13.4)	9,861	9.4	(7.2 - 11.6)
≥ College Degree	178	11.1	(4.1 - 18.1)	12,394	7.8	(5.8 - 9.7)
Income						
≤ \$24,999	161	9.2	(3.8 - 14.6)	9,600	14.1	(9.7 - 18.4)
\$25,000-\$49,999	179	6.1	(1.4 - 10.7)	9,590	9.5	(6.3 - 12.7)
\$50,000-\$74,999	108	10.6	(2.3 - 18.9)	5,741	7.5	(5.0 - 10.1)
≥ \$75,000	131	8.2	(2.4 - 14.1)	8,541	7.7	(5.9 - 9.5)

¹ Defined as adults having ever been told by a health professional that they had asthma and still have asthma

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wayne County**

Table 49a. Chronic joint symptoms among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	655	35,400	51.0	(45.9 - 56.2)	38,099	6,377,100	42.9	(40.5 - 45.3)
Gender								
Male	234	16,500	48.8	(40.3 - 57.3)	14,595	3,033,600	42.8	(39.0 - 46.6)
Female	421	18,900	53.2	(47.2 - 59.2)	23,504	3,343,500	43.0	(40.0 - 46.0)
Age (years)								
18-34	*	*	*	*	3,933	1,261,300	28.3	(22.6 - 34.0)
35-44	*	*	*	*	4,984	1,110,900	38.6	(33.1 - 44.0)
45-54	141	8,300	54.9	(45.2 - 64.5)	7,817	1,456,500	51.2	(46.4 - 55.9)
55-64	160	6,600	59.4	(51.1 - 67.6)	8,751	1,167,400	54.6	(50.4 - 58.8)
≥ 65	203	7,500	61.8	(54.5 - 69.0)	12,614	1,381,000	54.3	(50.5 - 58.1)
Race/Ethnicity								
White Non-Hispanic	631	32,800	50.5	(45.5 - 55.6)	35,785	4,267,600	45.8	(43.8 - 47.7)
Black Non-Hispanic	*	*	*	*	740	890,500	41.3	(32.5 - 50.0)
Hispanic	*	*	*	*	860	827,500	36.7	(27.7 - 45.6)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	312	17,200	53.8	(46.2 - 61.5)	15,676	2,450,000	43.8	(39.4 - 48.1)
Some College	*	*	*	*	9,900	1,598,300	45.1	(40.1 - 50.1)
≥ College Degree	177	8,900	46.8	(37.5 - 56.1)	12,440	2,313,100	41.1	(37.8 - 44.5)
Income								
≤ \$24,999	*	*	*	*	9,641	1,483,000	42.2	(36.3 - 48.1)
\$25,000-\$49,999	179	9,300	53.7	(44.6 - 62.8)	9,647	1,332,100	45.9	(39.8 - 52.1)
\$50,000-\$74,999	*	*	*	*	5,768	894,200	45.3	(39.5 - 51.1)
≥ \$75,000	*	*	*	*	8,549	1,874,200	41.7	(38.4 - 45.1)

¹ Defined as adults having symptoms of pain, aching or stiffness in or around a joint during the past 30 days and began more than 3 months ago

² Sample size from survey

³ Estimated number of adults who have chronic joint symptoms (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 49b. Chronic joint symptoms among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	655	49.3	(43.7 - 55.0)	38,099	42.1	(39.7 - 44.5)
Gender						
Male	234	47.4	(38.3 - 56.6)	14,595	42.6	(38.8 - 46.3)
Female	421	51.2	(44.5 - 57.8)	23,504	41.6	(38.6 - 44.7)
Race/Ethnicity						
White Non-Hispanic	631	48.5	(43.0 - 54.0)	35,785	44.7	(42.7 - 46.6)
Black Non-Hispanic	*	*	*	740	40.2	(31.5 - 48.9)
Hispanic	*	*	*	860	36.6	(27.8 - 45.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	312	51.8	(43.4 - 60.1)	15,676	43.1	(38.8 - 47.5)
Some College	*	*	*	9,900	43.9	(39.0 - 48.8)
≥ College Degree	*	*	*	12,440	40.3	(37.0 - 43.7)
Income						
≤ \$24,999	*	*	*	9,641	41.6	(35.7 - 47.4)
\$25,000-\$49,999	179	51.2	(41.3 - 61.0)	9,647	45.0	(39.0 - 51.0)
\$50,000-\$74,999	*	*	*	5,768	44.3	(38.4 - 50.2)
≥ \$75,000	*	*	*	8,549	40.8	(37.4 - 44.2)

¹ Defined as adults having symptoms of pain, aching or stiffness in or around a joint during the past 30 days and began more than 3 months ago

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 50a. Arthritis among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	656	25,500	36.7	(32.0 - 41.4)	38,013	4,102,800	27.7	(25.9 - 29.6)
Gender								
Male	236	10,600	31.2	(24.0 - 38.4)	14,555	1,642,700	23.3	(20.7 - 25.8)
Female	420	14,900	42.0	(36.2 - 47.8)	23,458	2,460,100	31.8	(29.2 - 34.4)
Age (years)								
18-34	*	*	*	*	3,927	312,100	7.0	(4.9 - 9.1)
35-44	*	*	*	*	4,972	412,500	14.5	(11.3 - 17.6)
45-54	141	5,400	35.9	(27.0 - 44.8)	7,789	869,100	30.7	(26.6 - 34.8)
55-64	161	6,200	55.2	(46.8 - 63.6)	8,729	999,300	47.0	(42.9 - 51.1)
≥ 65	203	7,200	59.7	(52.3 - 67.1)	12,596	1,509,800	59.6	(55.8 - 63.3)
Race/Ethnicity								
White Non-Hispanic	632	24,500	37.7	(33.0 - 42.3)	35,710	2,967,700	31.9	(30.3 - 33.6)
Black Non-Hispanic	*	*	*	*	737	588,100	27.3	(20.2 - 34.4)
Hispanic	*	*	*	*	856	380,700	17.1	(11.8 - 22.5)
Other Non-Hispanic	*	*	*	*	710	166,300	14.8	(7.8 - 21.8)
Education								
≤ High School	312	13,800	43.1	(35.7 - 50.5)	15,652	1,742,700	31.3	(27.8 - 34.8)
Some College	165	5,800	32.2	(23.9 - 40.4)	9,874	902,000	25.5	(22.3 - 28.6)
≥ College Degree	178	5,900	30.9	(22.3 - 39.6)	12,407	1,437,800	25.6	(22.9 - 28.4)
Income								
≤ \$24,999	*	*	*	*	9,618	1,112,500	32.0	(27.1 - 36.9)
\$25,000-\$49,999	179	6,300	36.6	(28.5 - 44.8)	9,624	821,900	28.4	(24.2 - 32.6)
\$50,000-\$74,999	*	*	*	*	5,753	542,500	27.5	(23.2 - 31.9)
≥ \$75,000	131	4,400	24.4	(15.7 - 33.0)	8,536	1,012,800	22.6	(19.8 - 25.3)

¹ Defined as adults having ever been told by a health professional that they have some form of arthritis, rheumatoid arthritis, gout, lupus or fibromyalgia (excluding back or neck)

² Sample size from survey

³ Estimated number of adults who have ever been told by a health professional they have some form of arthritis (excluding back or neck) (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 50b. Arthritis among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	656	34.3	(29.3 - 39.3)	38,013	26.6	(24.8 - 28.5)
Gender						
Male	236	29.4	(21.8 - 37.1)	14,555	23.5	(20.9 - 26.1)
Female	420	39.0	(32.6 - 45.3)	23,458	29.5	(27.0 - 32.1)
Race/Ethnicity						
White Non-Hispanic	632	34.9	(30.0 - 39.9)	35,710	30.4	(28.7 - 32.0)
Black Non-Hispanic	*	*	*	737	26.5	(19.3 - 33.6)
Hispanic	*	*	*	856	17.2	(11.8 - 22.6)
Other Non-Hispanic	*	*	*	710	14.9	(7.7 - 22.0)
Education						
≤ High School	312	41.3	(33.3 - 49.3)	15,652	30.4	(26.9 - 33.8)
Some College	165	28.0	(19.9 - 36.0)	9,874	24.1	(21.0 - 27.1)
≥ College Degree	178	29.6	(20.3 - 38.9)	12,407	24.7	(21.9 - 27.5)
Income						
≤ \$24,999	*	*	*	9,618	31.2	(26.3 - 36.1)
\$25,000-\$49,999	179	33.4	(25.0 - 41.7)	9,624	27.3	(23.2 - 31.3)
\$50,000-\$74,999	*	*	*	5,753	26.3	(21.9 - 30.7)
≥ \$75,000	131	22.6	(13.5 - 31.6)	8,536	21.6	(18.8 - 24.3)

¹ Defined as adults having ever been told by a health professional that they have some form of arthritis, rheumatoid arthritis, gout, lupus or fibromyalgia (excluding back or neck)

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 51a. Activity limitations because of physical, mental or emotional problems among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	651	16,000	23.2	(18.8 - 27.6)	37,998	3,053,000	20.6	(18.7 - 22.5)
Gender								
Male	233	7,200	21.3	(14.2 - 28.3)	14,566	1,318,400	18.6	(15.7 - 21.5)
Female	418	8,800	25.1	(19.7 - 30.4)	23,432	1,734,600	22.4	(19.8 - 25.0)
Age (years)								
18-34	*	*	*	*	3,929	580,000	13.0	(8.2 - 17.9)
35-44	*	*	*	*	4,973	418,600	14.5	(11.0 - 18.1)
45-54	139	3,000	20.1	(12.8 - 27.4)	7,812	620,200	21.7	(18.3 - 25.2)
55-64	161	3,600	32.4	(24.7 - 40.1)	8,739	651,600	30.5	(26.5 - 34.6)
≥ 65	200	4,200	35.4	(28.1 - 42.7)	12,545	782,700	31.1	(27.7 - 34.6)
Race/Ethnicity								
White Non-Hispanic	627	14,200	22.1	(18.3 - 25.8)	35,688	2,037,200	21.9	(20.3 - 23.5)
Black Non-Hispanic	*	*	*	*	735	429,700	20.1	(13.5 - 26.7)
Hispanic	*	*	*	*	863	448,200	19.8	(11.9 - 27.8)
Other Non-Hispanic	*	*	*	*	712	137,900	12.3	(5.2 - 19.3)
Education								
≤ High School	307	8,500	27.0	(19.6 - 34.5)	15,626	1,359,600	24.4	(20.6 - 28.1)
Some College	165	5,100	28.2	(19.7 - 36.8)	9,875	700,800	19.8	(16.5 - 23.2)
≥ College Degree	177	2,400	12.6	(7.5 - 17.7)	12,415	924,800	16.5	(14.2 - 18.7)
Income								
≤ \$24,999	*	*	*	*	9,603	1,055,800	30.1	(24.7 - 35.5)
\$25,000-\$49,999	177	4,000	23.3	(16.4 - 30.1)	9,628	630,900	21.8	(16.6 - 27.0)
\$50,000-\$74,999	107	3,300	24.3	(14.4 - 34.2)	5,760	408,100	20.7	(16.1 - 25.2)
≥ \$75,000	131	1,400	7.7	(3.3 - 12.1)	8,544	583,300	13.0	(11.1 - 14.9)

¹ Defined as activity limitations because of physical, mental or emotional problems

² Sample size from survey

³ Estimated number of adults who are limited in any activities because of physical, mental or emotional problems (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 51b. Activity limitations because of physical, mental or emotional problems among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	651	22.2	(17.5 - 27.0)	37,998	20.1	(18.2 - 22.0)
Gender						
Male	233	20.2	(12.7 - 27.7)	14,566	18.6	(15.8 - 21.4)
Female	418	24.2	(18.3 - 30.1)	23,432	21.5	(18.9 - 24.1)
Race/Ethnicity						
White Non-Hispanic	627	20.7	(16.8 - 24.7)	35,688	21.3	(19.7 - 22.9)
Black Non-Hispanic	*	*	*	735	19.9	(13.2 - 26.5)
Hispanic	*	*	*	863	19.1	(11.5 - 26.8)
Other Non-Hispanic	*	*	*	712	12.4	(5.3 - 19.6)
Education						
≤ High School	307	27.0	(18.8 - 35.3)	15,626	23.8	(20.1 - 27.4)
Some College	165	25.7	(16.7 - 34.8)	9,875	19.3	(16.0 - 22.6)
≥ College Degree	177	11.7	(6.8 - 16.5)	12,415	16.1	(13.9 - 18.3)
Income						
≤ \$24,999	*	*	*	9,603	29.7	(24.3 - 35.2)
\$25,000-\$49,999	177	21.8	(14.8 - 28.7)	9,628	21.2	(16.1 - 26.3)
\$50,000-\$74,999	*	*	*	5,760	20.1	(15.6 - 24.6)
≥ \$75,000	131	6.9	(2.9 - 10.9)	8,544	12.7	(10.8 - 14.6)

¹ Defined as activity limitations because of physical, mental or emotional problems

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 52a. Have health problem that requires the use of special equipment among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	658	5,300	7.6	(5.6 - 9.6)	38,171	1,169,400	7.9	(6.9 - 8.9)
Gender								
Male	236	1,800	5.3	(2.7 - 7.9)	14,617	460,700	6.5	(5.3 - 7.7)
Female	422	3,500	9.8	(6.9 - 12.7)	23,554	708,700	9.1	(7.6 - 10.7)
Age (years)								
18-34	*	*	*	*	3,940	68,800	1.5	(0.7 - 2.4)
35-44	*	*	*	*	4,989	122,500	4.3	(2.4 - 6.1)
45-54	*	*	*	*	7,836	182,500	6.4	(4.7 - 8.1)
55-64	162	1,100	9.6	(4.7 - 14.6)	8,773	219,600	10.3	(7.8 - 12.8)
≥ 65	204	2,700	22.6	(16.4 - 28.8)	12,633	576,100	22.7	(19.0 - 26.3)
Race/Ethnicity								
White Non-Hispanic	634	5,100	7.8	(5.8 - 9.9)	35,853	802,000	8.6	(7.6 - 9.6)
Black Non-Hispanic	*	*	*	*	739	235,800	10.9	(6.4 - 15.5)
Hispanic	*	*	*	*	864	104,100	4.6	(2.1 - 7.1)
Other Non-Hispanic	*	*	*	*	715	27,600	2.5	(0.8 - 4.1)
Education								
≤ High School	312	3,100	9.8	(6.5 - 13.1)	15,711	604,200	10.8	(8.7 - 12.8)
Some College	166	1,700	9.2	(4.8 - 13.6)	9,914	260,000	7.3	(5.5 - 9.1)
≥ College Degree	178	500	2.5	(0.8 - 4.2)	12,463	298,900	5.3	(4.1 - 6.5)
Income								
≤ \$24,999	162	2,200	16.8	(10.4 - 23.2)	9,656	430,500	12.2	(9.6 - 14.8)
\$25,000-\$49,999	179	1,500	8.6	(4.1 - 13.1)	9,663	238,900	8.2	(5.9 - 10.5)
\$50,000-\$74,999	*	*	*	*	5,775	103,000	5.2	(3.6 - 6.8)
≥ \$75,000	*	*	*	*	8,566	136,000	3.0	(2.1 - 4.0)

¹ Defined as having any health problem that requires the use of special equipment such as a cane, a wheelchair, a special bed, or a special telephone

² Sample size from survey

³ Estimated number of adults who have any health problem that requires the use of special equipment (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 52b. Have health problem that requires the use of special equipment among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	658	6.9	(5.0 - 8.8)	38,171	7.6	(6.6 - 8.6)
Gender						
Male	236	5.1	(2.5 - 7.6)	14,617	6.8	(5.5 - 8.0)
Female	422	8.7	(6.0 - 11.5)	23,554	8.4	(6.9 - 9.9)
Race/Ethnicity						
White Non-Hispanic	634	7.2	(5.2 - 9.1)	35,853	8.3	(7.3 - 9.3)
Black Non-Hispanic	*	*	*	739	10.5	(6.1 - 15.0)
Hispanic	*	*	*	864	4.5	(2.1 - 6.9)
Other Non-Hispanic	*	*	*	715	2.2	(0.7 - 3.8)
Education						
≤ High School	312	9.3	(6.0 - 12.7)	15,711	10.3	(8.3 - 12.3)
Some College	166	7.9	(3.9 - 11.8)	9,914	7.1	(5.2 - 8.9)
≥ College Degree	178	2.3	(0.7 - 3.9)	12,463	5.3	(4.1 - 6.6)
Income						
≤ \$24,999	162	15.3	(9.1 - 21.6)	9,656	11.9	(9.4 - 14.5)
\$25,000-\$49,999	179	8.2	(3.7 - 12.7)	9,663	7.7	(5.5 - 10.0)
\$50,000-\$74,999	*	*	*	5,775	5.1	(3.4 - 6.8)
≥ \$75,000	*	*	*	8,566	2.9	(2.0 - 3.9)

¹ Defined as having any health problem that requires the use of special equipment such as a cane, a wheelchair, a special bed, or a special telephone

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wayne County**

Table 53a. Disability among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	652	17,200	24.9	(20.5 - 29.3)	38,016	3,394,200	22.9	(20.9 - 24.9)
Gender								
Male	233	7,500	22.3	(15.3 - 29.4)	14,568	1,469,700	20.7	(17.8 - 23.7)
Female	419	9,600	27.4	(21.9 - 32.8)	23,448	1,924,600	24.8	(22.1 - 27.5)
Age (years)								
18-34	*	*	*	*	3,931	590,300	13.3	(8.4 - 18.1)
35-44	*	*	*	*	4,974	447,900	15.6	(11.9 - 19.3)
45-54	139	3,100	20.9	(13.5 - 28.3)	7,811	643,900	22.6	(19.1 - 26.0)
55-64	161	3,800	33.8	(26.0 - 41.6)	8,740	702,100	33.0	(28.9 - 37.0)
≥ 65	201	5,100	42.7	(35.3 - 50.2)	12,560	1,010,000	39.9	(36.1 - 43.7)
Race/Ethnicity								
White Non-Hispanic	628	15,300	23.7	(19.8 - 27.5)	35,703	2,253,400	24.2	(22.6 - 25.9)
Black Non-Hispanic	*	*	*	*	736	511,700	23.8	(16.8 - 30.8)
Hispanic	*	*	*	*	864	484,300	21.4	(13.4 - 29.5)
Other Non-Hispanic	*	*	*	*	713	144,800	12.9	(5.8 - 19.9)
Education								
≤ High School	308	9,300	29.5	(22.0 - 36.9)	15,638	1,544,200	27.6	(23.7 - 31.4)
Some College	165	5,200	29.0	(20.4 - 37.6)	9,876	755,200	21.4	(17.9 - 24.8)
≥ College Degree	177	2,600	13.9	(8.6 - 19.1)	12,419	1,021,300	18.2	(15.9 - 20.5)
Income								
≤ \$24,999	*	*	*	*	9,613	1,179,100	33.6	(28.0 - 39.2)
\$25,000-\$49,999	177	4,300	24.8	(17.8 - 31.8)	9,629	699,000	24.2	(18.9 - 29.4)
\$50,000-\$74,999	107	3,400	25.0	(15.0 - 34.9)	5,757	430,100	21.8	(17.2 - 26.4)
≥ \$75,000	131	1,400	7.9	(3.5 - 12.3)	8,546	632,700	14.1	(12.1 - 16.1)

¹ Defined as having activity limitations due to physical, mental or emotional problems or have health problems that require the use of special equipment

² Sample size from survey

³ Estimated number of adults with a disability (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 53b. Disability among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	652	23.7	(19.0 - 28.5)	38,016	22.3	(20.4 - 24.3)
Gender						
Male	233	21.2	(13.7 - 28.8)	14,568	20.9	(18.0 - 23.8)
Female	419	26.2	(20.2 - 32.1)	23,448	23.6	(21.0 - 26.3)
Race/Ethnicity						
White Non-Hispanic	628	22.2	(18.2 - 26.2)	35,703	23.6	(21.9 - 25.3)
Black Non-Hispanic	*	*	*	736	23.5	(16.5 - 30.5)
Hispanic	*	*	*	864	20.6	(12.9 - 28.3)
Other Non-Hispanic	*	*	*	713	12.9	(5.8 - 20.1)
Education						
≤ High School	308	29.3	(21.1 - 37.6)	15,638	26.8	(23.0 - 30.6)
Some College	165	26.3	(17.3 - 35.4)	9,876	20.9	(17.4 - 24.3)
≥ College Degree	177	12.8	(7.7 - 17.8)	12,419	17.9	(15.5 - 20.3)
Income						
≤ \$24,999	*	*	*	9,613	33.2	(27.6 - 38.8)
\$25,000-\$49,999	177	23.3	(16.1 - 30.5)	9,629	23.5	(18.3 - 28.6)
\$50,000-\$74,999	*	*	*	5,757	21.2	(16.6 - 25.8)
≥ \$75,000	131	7.1	(3.0 - 11.1)	8,546	13.8	(11.7 - 15.8)

¹ Defined as having activity limitations due to physical, mental or emotional problems or have health problems that require the use of special equipment

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 54a. Current smoking among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	656	13,400	19.4	(14.9 - 23.8)	38,029	2,483,500	16.8	(15.1 - 18.6)
Gender								
Male	235	7,200	21.2	(13.7 - 28.6)	14,559	1,356,400	19.3	(16.3 - 22.2)
Female	421	6,300	17.7	(12.7 - 22.6)	23,470	1,127,100	14.6	(12.7 - 16.6)
Age (years)								
18-34	*	*	*	*	3,929	993,500	22.3	(17.5 - 27.1)
35-44	*	*	*	*	4,977	421,100	14.8	(11.6 - 17.9)
45-54	141	3,000	20.0	(12.9 - 27.0)	7,807	508,800	18.3	(15.4 - 21.3)
55-64	161	1,800	16.4	(10.5 - 22.2)	8,744	314,500	14.7	(12.1 - 17.3)
≥ 65	203	900	7.6	(3.8 - 11.4)	12,572	245,600	9.7	(7.2 - 12.3)
Race/Ethnicity								
White Non-Hispanic	632	11,400	17.6	(13.9 - 21.3)	35,729	1,615,500	17.3	(15.9 - 18.8)
Black Non-Hispanic	*	*	*	*	735	284,100	13.6	(8.9 - 18.2)
Hispanic	*	*	*	*	858	350,900	15.8	(9.6 - 21.9)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	311	9,100	28.6	(20.8 - 36.5)	15,654	1,237,300	22.4	(19.1 - 25.6)
Some College	165	2,900	15.9	(9.0 - 22.8)	9,875	670,600	19.1	(15.5 - 22.8)
≥ College Degree	178	1,100	6.0	(2.6 - 9.3)	12,421	563,300	10.0	(7.5 - 12.5)
Income								
≤ \$24,999	*	*	*	*	9,626	835,100	24.1	(19.1 - 29.1)
\$25,000-\$49,999	178	4,400	25.7	(17.5 - 34.0)	9,629	558,100	19.4	(15.6 - 23.2)
\$50,000-\$74,999	108	1,100	8.4	(3.5 - 13.2)	5,756	338,100	17.3	(13.0 - 21.7)
≥ \$75,000	131	1,600	8.8	(3.2 - 14.4)	8,549	468,700	10.4	(8.6 - 12.3)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day or some days

² Sample size from survey

³ Estimated number of adult current smokers (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 54b. Current smoking among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	656	19.9	(15.0 - 24.8)	38,029	17.0	(15.3 - 18.8)
Gender						
Male	235	21.6	(13.4 - 29.7)	14,559	19.2	(16.3 - 22.1)
Female	421	18.3	(12.7 - 23.8)	23,470	15.0	(13.0 - 17.0)
Race/Ethnicity						
White Non-Hispanic	632	17.9	(13.8 - 21.9)	35,729	17.7	(16.2 - 19.2)
Black Non-Hispanic	*	*	*	735	13.5	(8.8 - 18.2)
Hispanic	*	*	*	858	15.9	(9.8 - 22.0)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	311	29.9	(21.2 - 38.5)	15,654	22.8	(19.5 - 26.0)
Some College	165	15.8	(8.3 - 23.2)	9,875	19.4	(15.8 - 22.9)
≥ College Degree	178	6.1	(2.5 - 9.8)	12,421	10.0	(7.6 - 12.5)
Income						
≤ \$24,999	*	*	*	9,626	24.1	(19.2 - 29.1)
\$25,000-\$49,999	178	27.0	(17.9 - 36.2)	9,629	19.9	(16.0 - 23.7)
\$50,000-\$74,999	108	7.4	(2.9 - 11.8)	5,756	17.8	(13.3 - 22.3)
≥ \$75,000	131	8.7	(2.7 - 14.7)	8,549	10.6	(8.6 - 12.5)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day or some days

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wayne County**

Table 55a. Everyday smoking among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	656	10,600	15.3	(11.1 - 19.5)	38,029	1,626,800	11.0	(9.8 - 12.2)
Gender								
Male	235	6,100	17.9	(10.7 - 25.2)	14,559	848,400	12.0	(10.0 - 14.1)
Female	421	4,500	12.8	(8.4 - 17.2)	23,470	778,400	10.1	(8.7 - 11.5)
Age (years)								
18-34	*	*	*	*	3,929	559,800	12.6	(9.6 - 15.6)
35-44	*	*	*	*	4,977	315,200	11.0	(8.4 - 13.7)
45-54	141	2,500	16.4	(9.9 - 22.9)	7,807	377,000	13.6	(11.1 - 16.0)
55-64	161	1,500	13.2	(7.8 - 18.5)	8,744	203,000	9.5	(7.5 - 11.5)
≥ 65	203	900	7.1	(3.4 - 10.9)	12,572	171,700	6.8	(4.8 - 8.8)
Race/Ethnicity								
White Non-Hispanic	632	8,700	13.4	(10.1 - 16.7)	35,729	1,158,100	12.4	(11.2 - 13.7)
Black Non-Hispanic	*	*	*	*	735	197,700	9.4	(5.6 - 13.3)
Hispanic	*	*	*	*	858	176,200	7.9	(4.2 - 11.6)
Other Non-Hispanic	*	*	*	*	707	94,900	8.5	(3.3 - 13.7)
Education								
≤ High School	311	7,600	23.8	(16.1 - 31.5)	15,654	881,900	15.9	(13.4 - 18.5)
Some College	165	2,000	11.2	(5.3 - 17.1)	9,875	410,200	11.7	(9.6 - 13.8)
≥ College Degree	178	700	3.8	(1.4 - 6.1)	12,421	323,500	5.8	(4.3 - 7.2)
Income								
≤ \$24,999	*	*	*	*	9,626	452,800	13.1	(10.3 - 15.8)
\$25,000-\$49,999	178	3,400	20.0	(12.4 - 27.5)	9,629	425,900	14.8	(11.9 - 17.7)
\$50,000-\$74,999	*	*	*	*	5,756	222,700	11.4	(8.5 - 14.4)
≥ \$75,000	131	1,100	6.1	(1.6 - 10.6)	8,549	320,200	7.1	(5.5 - 8.8)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day

² Sample size from survey

³ Estimated number of adults who have smoked at least 100 cigarettes in lifetime and currently smoking everyday (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 55b. Everyday smoking among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	656	15.5	(10.9 - 20.2)	38,029	11.3	(10.0 - 12.5)
Gender						
Male	235	18.3	(10.3 - 26.3)	14,559	12.1	(10.1 - 14.1)
Female	421	12.9	(8.0 - 17.9)	23,470	10.5	(9.0 - 12.0)
Race/Ethnicity						
White Non-Hispanic	632	13.3	(9.7 - 17.0)	35,729	12.8	(11.5 - 14.1)
Black Non-Hispanic	*	*	*	735	9.4	(5.5 - 13.4)
Hispanic	*	*	*	858	8.0	(4.3 - 11.7)
Other Non-Hispanic	*	*	*	707	8.5	(3.2 - 13.8)
Education						
≤ High School	311	24.7	(16.1 - 33.3)	15,654	16.4	(13.8 - 19.0)
Some College	165	10.7	(4.5 - 16.9)	9,875	12.1	(9.9 - 14.3)
≥ College Degree	178	3.8	(1.4 - 6.3)	12,421	5.8	(4.3 - 7.3)
Income						
≤ \$24,999	*	*	*	9,626	13.4	(10.6 - 16.2)
\$25,000-\$49,999	178	20.6	(12.3 - 28.9)	9,629	15.3	(12.3 - 18.3)
\$50,000-\$74,999	*	*	*	5,756	11.7	(8.7 - 14.8)
≥ \$75,000	131	5.8	(1.2 - 10.5)	8,549	7.2	(5.5 - 8.9)

¹ Defined as having smoked at least 100 cigarettes in lifetime and currently smoking every day

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 56a. Adults living in homes in which smoking is prohibited¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	654	55,900	80.9	(77.2 - 84.6)	38,045	11,997,100	80.8	(79.0 - 82.6)
Gender								
Male	234	26,800	79.3	(73.2 - 85.4)	14,577	5,634,400	79.4	(76.6 - 82.2)
Female	420	29,200	82.5	(78.2 - 86.7)	23,468	6,362,800	82.1	(79.9 - 84.3)
Age (years)								
18-34	51	11,100	90.7	(82.3 - 99.1)	3,934	3,655,200	82.1	(77.8 - 86.4)
35-44	100	15,500	82.5	(73.6 - 91.4)	4,985	2,479,700	86.1	(82.5 - 89.7)
45-54	141	11,500	76.4	(69.0 - 83.8)	7,808	2,194,400	77.2	(73.5 - 80.8)
55-64	161	8,300	74.3	(67.0 - 81.5)	8,753	1,732,500	81.0	(78.3 - 83.7)
≥ 65	201	9,500	80.2	(74.2 - 86.2)	12,565	1,935,400	76.5	(73.4 - 79.5)
Race/Ethnicity								
White Non-Hispanic	630	52,100	80.6	(76.8 - 84.3)	35,737	7,426,900	79.8	(78.3 - 81.2)
Black Non-Hispanic	*	*	*	*	738	1,692,100	78.6	(71.9 - 85.3)
Hispanic	*	*	*	*	863	2,003,900	88.7	(84.1 - 93.3)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	308	23,500	74.4	(68.3 - 80.6)	15,647	4,307,200	77.1	(74.0 - 80.2)
Some College	166	15,400	85.2	(79.1 - 91.3)	9,883	2,795,400	78.9	(75.0 - 82.8)
≥ College Degree	178	16,700	87.1	(80.8 - 93.5)	12,437	4,817,300	85.7	(83.3 - 88.2)
Income								
≤ \$24,999	*	*	*	*	9,608	2,627,700	74.9	(70.0 - 79.7)
\$25,000-\$49,999	176	13,800	81.2	(74.7 - 87.7)	9,640	2,314,500	79.8	(76.0 - 83.5)
\$50,000-\$74,999	108	11,100	81.7	(72.7 - 90.6)	5,766	1,551,300	78.7	(74.0 - 83.4)
≥ \$75,000	131	16,800	92.2	(88.1 - 96.4)	8,556	3,883,200	86.5	(84.3 - 88.7)

¹ Defined as having a rule that prohibits smoking anywhere inside the home

² Sample size from survey

³ Estimated number of adults who live in a home with a rule that prohibits smoking anywhere inside the home (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 56b. Adults living in homes in which smoking is prohibited¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	654	81.8	(77.9 - 85.7)	38,045	80.9	(79.2 - 82.7)
Gender						
Male	234	79.9	(73.4 - 86.4)	14,577	79.4	(76.7 - 82.2)
Female	420	83.5	(79.0 - 88.1)	23,468	82.3	(80.0 - 84.5)
Race/Ethnicity						
White Non-Hispanic	630	81.4	(77.4 - 85.4)	35,737	79.9	(78.4 - 81.4)
Black Non-Hispanic	*	*	*	738	78.6	(71.8 - 85.4)
Hispanic	*	*	*	863	88.9	(84.4 - 93.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	308	75.4	(68.9 - 82.0)	15,647	76.9	(73.8 - 80.0)
Some College	166	86.2	(79.7 - 92.7)	9,883	79.1	(75.2 - 83.0)
≥ College Degree	178	87.4	(80.6 - 94.2)	12,437	86.0	(83.5 - 88.5)
Income						
≤ \$24,999	*	*	*	9,608	74.9	(70.1 - 79.7)
\$25,000-\$49,999	176	82.5	(75.8 - 89.2)	9,640	79.8	(76.1 - 83.6)
\$50,000-\$74,999	108	82.6	(72.9 - 92.2)	5,766	78.6	(73.8 - 83.3)
≥ \$75,000	131	93.7	(90.2 - 97.1)	8,556	86.6	(84.3 - 88.9)

¹ Defined as having a rule that prohibits smoking anywhere inside the home

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 57a. Binge drinking within the past month among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	645	11,800	17.3	(12.6 - 22.0)	37,745	2,602,000	17.7	(15.7 - 19.8)
Gender								
Male	230	7,400	22.1	(14.0 - 30.2)	14,420	1,712,400	24.5	(20.9 - 28.1)
Female	415	4,500	12.7	(7.8 - 17.7)	23,325	889,600	11.6	(9.5 - 13.7)
Age (years)								
18-34	*	*	*	*	3,895	1,254,300	28.5	(22.8 - 34.2)
35-44	*	*	*	*	4,934	584,800	20.6	(16.5 - 24.7)
45-54	136	1,700	11.5	(5.3 - 17.8)	7,738	467,600	16.6	(13.3 - 19.9)
55-64	156	1,200	10.6	(5.7 - 15.6)	8,663	207,200	9.8	(7.9 - 11.8)
≥ 65	*	*	*	*	12,515	88,100	3.5	(2.5 - 4.5)
Race/Ethnicity								
White Non-Hispanic	622	9,800	15.3	(11.2 - 19.3)	35,453	1,761,900	19.2	(17.5 - 20.8)
Black Non-Hispanic	*	*	*	*	728	247,000	11.6	(6.3 - 17.0)
Hispanic	*	*	*	*	854	460,600	20.5	(11.3 - 29.8)
Other Non-Hispanic	*	*	*	*	710	132,500	12.1	(3.4 - 20.7)
Education								
≤ High School	303	6,700	21.7	(13.5 - 29.9)	15,491	945,800	17.4	(13.2 - 21.6)
Some College	164	3,000	16.9	(8.2 - 25.7)	9,819	636,600	18.1	(14.6 - 21.5)
≥ College Degree	176	2,000	10.8	(5.6 - 16.0)	12,358	1,018,700	18.2	(15.3 - 21.1)
Income								
≤ \$24,999	*	*	*	*	9,557	469,100	13.5	(8.5 - 18.5)
\$25,000-\$49,999	177	3,400	19.7	(11.3 - 28.2)	9,575	534,300	18.5	(13.0 - 24.0)
\$50,000-\$74,999	106	1,500	10.8	(4.6 - 17.0)	5,726	378,200	19.2	(14.1 - 24.3)
≥ \$75,000	129	3,500	19.3	(10.6 - 27.9)	8,501	1,011,400	22.6	(19.7 - 25.6)

¹ Defined as men having 5 or more drinks or women having 4 or more drinks on 1 or more occasion within the past month

² Sample size from survey

³ Estimated number of adult binge drinkers within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 57b. Binge drinking within the past month among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	645	18.6	(13.3 - 23.9)	37,745	18.1	(16.1 - 20.2)
Gender						
Male	230	23.5	(14.6 - 32.4)	14,420	24.5	(21.0 - 27.9)
Female	415	13.9	(8.1 - 19.6)	23,325	12.4	(10.2 - 14.5)
Race/Ethnicity						
White Non-Hispanic	622	16.2	(11.7 - 20.8)	35,453	19.8	(18.1 - 21.6)
Black Non-Hispanic	*	*	*	728	11.8	(6.4 - 17.1)
Hispanic	*	*	*	854	20.2	(11.3 - 29.1)
Other Non-Hispanic	*	*	*	710	12.1	(3.4 - 20.8)
Education						
≤ High School	303	23.6	(14.4 - 32.8)	15,491	17.5	(13.5 - 21.5)
Some College	164	18.1	(8.2 - 28.0)	9,819	18.6	(15.1 - 22.0)
≥ College Degree	176	11.4	(5.6 - 17.1)	12,358	18.8	(15.9 - 21.7)
Income						
≤ \$24,999	*	*	*	9,557	13.4	(8.6 - 18.2)
\$25,000-\$49,999	177	21.8	(12.2 - 31.5)	9,575	18.9	(13.5 - 24.2)
\$50,000-\$74,999	106	10.9	(4.4 - 17.5)	5,726	19.9	(14.8 - 25.1)
≥ \$75,000	129	19.8	(10.5 - 29.2)	8,501	23.2	(20.2 - 26.1)

¹ Defined as men having 5 or more drinks or women having 4 or more drinks on 1 or more occasion within the past month

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 58a. Heavy drinking within the past month among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	644	4,400	6.4	(2.7 - 10.1)	37,553	718,700	4.9	(4.2 - 5.6)
Gender								
Male	229	2,400	7.2	(0.7 - 13.7)	14,330	333,300	4.8	(3.9 - 5.7)
Female	415	2,000	5.6	(1.9 - 9.4)	23,223	385,400	5.0	(4.0 - 6.1)
Age (years)								
18-34	*	*	*	*	3,866	224,900	5.1	(3.5 - 6.8)
35-44	*	*	*	*	4,924	137,200	4.8	(3.0 - 6.7)
45-54	136	900	6.5	(2.5 - 10.6)	7,715	144,600	5.2	(4.1 - 6.2)
55-64	*	*	*	*	8,624	117,400	5.6	(3.9 - 7.3)
≥ 65	*	*	*	*	12,424	94,600	3.8	(2.7 - 4.9)
Race/Ethnicity								
White Non-Hispanic	621	2,700	4.2	(2.1 - 6.3)	35,279	572,400	6.3	(5.4 - 7.1)
Black Non-Hispanic	*	*	*	*	723	58,100	2.8	(0.6 - 4.9)
Hispanic	*	*	*	*	848	60,900	2.7	(0.5 - 4.9)
Other Non-Hispanic	*	*	*	*	703	27,200	2.5	(0.8 - 4.2)
Education								
≤ High School	301	3,300	10.8	(3.2 - 18.4)	15,411	228,100	4.2	(3.2 - 5.2)
Some College	*	*	*	*	9,759	163,700	4.7	(3.6 - 5.8)
≥ College Degree	*	*	*	*	12,306	326,800	5.9	(4.5 - 7.3)
Income								
≤ \$24,999	*	*	*	*	9,512	104,200	3.0	(1.8 - 4.1)
\$25,000-\$49,999	178	1,000	5.6	(1.5 - 9.6)	9,537	158,700	5.5	(3.9 - 7.2)
\$50,000-\$74,999	*	*	*	*	5,708	95,400	4.9	(3.1 - 6.7)
≥ \$75,000	*	*	*	*	8,481	313,900	7.0	(5.4 - 8.6)

¹ Defined as adult men averaging more than 2 alcoholic drinks per day and adult women averaging more than 1 alcoholic drink per day within the past month

² Sample size from survey

³ Estimated number of adult heavy drinkers within the past month (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 58b. Heavy drinking within the past month among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	644	6.6	(2.5 - 10.8)	37,553	5.0	(4.3 - 5.8)
Gender						
Male	229	7.6	(0.4 - 14.8)	14,330	4.9	(3.9 - 5.9)
Female	415	5.7	(1.4 - 10.0)	23,223	5.1	(4.0 - 6.2)
Race/Ethnicity						
White Non-Hispanic	621	4.2	(1.9 - 6.5)	35,279	6.4	(5.5 - 7.2)
Black Non-Hispanic	*	*	*	723	2.9	(0.6 - 5.1)
Hispanic	*	*	*	848	2.8	(0.5 - 5.1)
Other Non-Hispanic	*	*	*	703	2.5	(0.8 - 4.2)
Education						
≤ High School	301	11.7	(3.1 - 20.3)	15,411	4.3	(3.2 - 5.3)
Some College	*	*	*	9,759	4.8	(3.6 - 5.9)
≥ College Degree	*	*	*	12,306	6.0	(4.5 - 7.4)
Income						
≤ \$24,999	*	*	*	9,512	3.0	(1.9 - 4.1)
\$25,000-\$49,999	178	5.5	(1.0 - 9.9)	9,537	5.8	(4.0 - 7.5)
\$50,000-\$74,999	*	*	*	5,708	5.0	(3.1 - 7.0)
≥ \$75,000	*	*	*	8,481	7.0	(5.4 - 8.7)

¹ Defined as adult men averaging more than 2 alcoholic drinks per day and adult women averaging more than 1 alcoholic drink per day within the past month

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 59a. Never or rarely been asked about sexual history during routine check-up among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	623	46,000	70.5	(65.4 - 75.6)	35,193	8,480,600	61.9	(59.2 - 64.5)
Gender								
Male	218	23,600	76.0	(67.8 - 84.2)	13,299	4,357,400	67.7	(63.6 - 71.7)
Female	405	22,400	65.5	(59.3 - 71.7)	21,894	4,123,300	56.8	(53.3 - 60.2)
Age (years)								
18-34	*	*	*	*	3,507	1,804,600	44.3	(37.8 - 50.8)
35-44	*	*	*	*	4,531	1,635,300	62.2	(56.6 - 67.8)
45-54	133	9,900	71.2	(61.6 - 80.8)	7,215	1,715,600	65.4	(60.5 - 70.3)
55-64	156	8,200	76.3	(69.0 - 83.5)	8,168	1,416,200	70.7	(66.6 - 74.8)
≥ 65	193	9,900	88.0	(82.7 - 93.3)	11,772	1,908,900	80.3	(76.9 - 83.8)
Race/Ethnicity								
White Non-Hispanic	602	43,400	71.2	(66.2 - 76.1)	33,090	5,940,300	68.5	(66.3 - 70.7)
Black Non-Hispanic	*	*	*	*	683	831,800	40.8	(31.8 - 49.9)
Hispanic	*	*	*	*	784	1,071,700	53.4	(43.6 - 63.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	291	20,700	71.3	(63.5 - 79.1)	14,264	3,001,200	60.1	(55.4 - 64.9)
Some College	*	*	*	*	9,156	1,972,800	60.3	(54.9 - 65.7)
≥ College Degree	169	12,600	68.5	(59.7 - 77.3)	11,706	3,476,000	65.0	(61.2 - 68.7)
Income								
≤ \$24,999	*	*	*	*	8,772	1,529,400	49.3	(42.6 - 56.0)
\$25,000-\$49,999	172	12,100	73.7	(65.8 - 81.6)	8,932	1,705,000	64.4	(58.0 - 70.7)
\$50,000-\$74,999	*	*	*	*	5,433	1,217,000	64.5	(58.5 - 70.5)
≥ \$75,000	*	*	*	*	8,108	2,934,300	68.2	(64.7 - 71.7)

¹ Defined as respondents who never or rarely (hardly ever) were asked about their sexual history, such as about sexual partners and sexual practices, during a routine check-up at a doctor's office or clinic

² Sample size from survey

³ Estimated number of adults who were never or rarely asked about their sexual history during a routine check-up at a doctor's office or clinic (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 59b. Never or rarely been asked about sexual history during routine check-up among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	623	69.2	(63.5 - 74.8)	35,193	61.4	(58.7 - 64.0)
Gender						
Male	218	75.6	(66.7 - 84.5)	13,299	67.9	(64.0 - 71.9)
Female	405	63.3	(56.4 - 70.3)	21,894	55.5	(52.0 - 59.0)
Race/Ethnicity						
White Non-Hispanic	602	69.7	(64.2 - 75.3)	33,090	67.8	(65.6 - 70.1)
Black Non-Hispanic	*	*	*	683	41.1	(32.0 - 50.2)
Hispanic	*	*	*	784	53.0	(43.3 - 62.6)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	291	69.4	(60.8 - 78.0)	14,264	59.5	(54.7 - 64.3)
Some College	*	*	*	9,156	59.7	(54.4 - 65.0)
≥ College Degree	169	68.4	(58.8 - 78.0)	11,706	64.5	(60.8 - 68.2)
Income						
≤ \$24,999	*	*	*	8,772	48.7	(42.0 - 55.4)
\$25,000-\$49,999	172	72.3	(63.5 - 81.1)	8,932	63.7	(57.5 - 70.0)
\$50,000-\$74,999	*	*	*	5,433	64.2	(58.2 - 70.2)
≥ \$75,000	*	*	*	8,108	67.5	(64.0 - 71.0)

¹ Defined as respondents who never or rarely (hardly ever) were asked about their sexual history, such as about sexual partners and sexual practices, during a routine check-up at a doctor's office or clinic

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wayne County**

Table 60a. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	400	35,000	78.9	(73.2 - 84.7)	22,788	7,011,100	77.9	(75.2 - 80.5)
Gender								
Male	143	16,500	79.3	(70.0 - 88.6)	9,034	3,618,500	81.0	(77.2 - 84.8)
Female	257	18,500	78.6	(71.8 - 85.5)	13,754	3,392,600	74.8	(71.2 - 78.4)
Age (years)								
18-34	*	*	*	*	3,213	2,362,500	70.8	(65.1 - 76.6)
35-44	*	*	*	*	3,513	1,313,400	71.7	(65.6 - 77.7)
45-54	87	6,900	80.9	(72.4 - 89.3)	4,975	1,290,600	80.3	(75.9 - 84.8)
55-64	104	6,600	92.4	(87.3 - 97.5)	4,955	932,200	87.2	(84.1 - 90.3)
≥ 65	95	5,300	94.0	(88.4 - 99.6)	6,132	1,112,400	95.9	(94.4 - 97.5)
Race/Ethnicity								
White Non-Hispanic	383	33,500	79.9	(74.4 - 85.5)	21,540	4,875,200	78.3	(75.6 - 80.9)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	491	1,002,400	84.8	(77.8 - 91.8)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	182	15,000	78.6	(69.5 - 87.8)	8,879	2,581,700	82.1	(77.9 - 86.3)
Some College	105	10,000	83.7	(75.6 - 91.8)	5,953	1,599,500	77.5	(72.6 - 82.3)
≥ College Degree	*	*	*	*	7,928	2,824,800	75.6	(71.8 - 79.5)
Income								
≤ \$24,999	*	*	*	*	5,491	1,602,900	78.7	(71.9 - 85.6)
\$25,000-\$49,999	121	9,400	79.5	(69.9 - 89.1)	5,912	1,335,900	78.8	(73.8 - 83.9)
\$50,000-\$74,999	75	7,800	83.3	(73.6 - 93.0)	3,717	987,500	76.6	(70.4 - 82.8)
≥ \$75,000	*	*	*	*	5,771	2,321,600	77.6	(74.0 - 81.2)

¹ Defined as respondent estimating there were hardly any (0-1 out of 10) or a few (2-3 out of 10) persons their age in their community who have had a sexually transmitted disease such as gonorrhea, chlamydia, herpes or syphilis

² Sample size from survey

³ Estimated number of adults who believed hardly any or a few people their age in their community have had a sexually transmitted disease (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 60b. Believed hardly any or a few people their age in their community have had a sexually transmitted disease among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	400	77.5	(71.2 - 83.9)	22,788	77.3	(74.7 - 79.9)
Gender						
Male	*	*	*	9,034	81.1	(77.4 - 84.8)
Female	257	77.9	(70.2 - 85.6)	13,754	73.6	(70.0 - 77.3)
Race/Ethnicity						
White Non-Hispanic	383	78.6	(72.4 - 84.8)	21,540	77.5	(74.9 - 80.1)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	491	84.6	(77.6 - 91.5)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	*	*	*	8,879	81.6	(77.4 - 85.7)
Some College	105	84.5	(76.1 - 92.8)	5,953	76.8	(72.0 - 81.6)
≥ College Degree	*	*	*	7,928	75.1	(71.3 - 79.0)
Income						
≤ \$24,999	*	*	*	5,491	77.9	(71.0 - 84.7)
\$25,000-\$49,999	*	*	*	5,912	77.9	(72.8 - 83.0)
\$50,000-\$74,999	*	*	*	3,717	76.1	(70.0 - 82.3)
≥ \$75,000	*	*	*	5,771	77.0	(73.4 - 80.6)

¹ Defined as respondent estimating there were hardly any (0-1 out of 10) or a few (2-3 out of 10) persons their age in their community who have had a sexually transmitted disease such as gonorrhea, chlamydia, herpes or syphilis

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 61a. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	632	58,500	87.1	(83.3 - 90.8)	36,287	12,738,300	89.7	(88.2 - 91.3)
Gender								
Male	224	28,200	86.7	(80.0 - 93.5)	13,893	6,104,600	89.4	(87.2 - 91.5)
Female	408	30,300	87.4	(83.8 - 90.9)	22,394	6,633,700	90.1	(87.9 - 92.2)
Age (years)								
18-34	50	11,100	93.7	(87.6 - 99.8)	3,850	4,007,400	91.3	(87.5 - 95.1)
35-44	*	*	*	*	4,875	2,578,600	93.2	(90.4 - 96.0)
45-54	140	13,300	88.5	(83.0 - 94.1)	7,607	2,416,800	90.5	(87.9 - 93.2)
55-64	157	9,800	89.7	(85.0 - 94.5)	8,411	1,876,500	92.2	(90.0 - 94.5)
≥ 65	187	8,300	74.9	(68.0 - 81.8)	11,544	1,858,900	79.7	(76.6 - 82.8)
Race/Ethnicity								
White Non-Hispanic	610	55,500	88.2	(85.3 - 91.1)	34,102	8,050,700	89.4	(87.7 - 91.0)
Black Non-Hispanic	*	*	*	*	702	1,822,200	91.9	(88.2 - 95.6)
Hispanic	*	*	*	*	820	1,967,200	89.9	(85.2 - 94.7)
Other Non-Hispanic	*	*	*	*	663	898,200	88.6	(79.5 - 97.6)
Education								
≤ High School	294	24,400	80.6	(73.5 - 87.7)	14,507	4,385,300	84.6	(81.9 - 87.2)
Some College	163	16,700	93.2	(89.8 - 96.6)	9,571	3,094,300	90.0	(86.7 - 93.4)
≥ College Degree	173	17,100	91.4	(86.4 - 96.4)	12,142	5,228,800	95.4	(94.0 - 96.7)
Income								
≤ \$24,999	*	*	*	*	9,021	2,833,500	86.5	(82.6 - 90.3)
\$25,000-\$49,999	175	15,800	93.8	(90.1 - 97.5)	9,261	2,529,900	89.5	(86.5 - 92.5)
\$50,000-\$74,999	108	12,100	89.0	(81.7 - 96.3)	5,644	1,740,000	91.2	(87.4 - 95.1)
≥ \$75,000	127	16,200	92.5	(88.1 - 96.9)	8,397	4,184,300	94.4	(92.5 - 96.3)

¹ Defined as believing it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community

² Sample size from survey

³ Estimated number of adults who believed it is at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 61b. Believed it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	632	87.2	(83.2 - 91.3)	36,287	89.8	(88.3 - 91.4)
Gender						
Male	224	86.4	(79.0 - 93.8)	13,893	89.1	(86.9 - 91.3)
Female	408	88.0	(84.4 - 91.7)	22,394	90.5	(88.3 - 92.7)
Race/Ethnicity						
White Non-Hispanic	610	88.5	(85.4 - 91.6)	34,102	89.5	(87.8 - 91.2)
Black Non-Hispanic	*	*	*	702	91.7	(87.8 - 95.6)
Hispanic	*	*	*	820	89.6	(84.7 - 94.5)
Other Non-Hispanic	*	*	*	663	89.0	(80.5 - 97.6)
Education						
≤ High School	294	80.3	(72.5 - 88.1)	14,507	84.6	(81.8 - 87.4)
Some College	163	94.1	(91.1 - 97.2)	9,571	90.2	(86.9 - 93.5)
≥ College Degree	173	91.3	(85.8 - 96.7)	12,142	95.4	(94.0 - 96.7)
Income						
≤ \$24,999	*	*	*	9,021	86.5	(82.6 - 90.4)
\$25,000-\$49,999	175	94.6	(91.3 - 97.9)	9,261	89.6	(86.5 - 92.6)
\$50,000-\$74,999	108	88.0	(79.7 - 96.4)	5,644	91.1	(86.9 - 95.3)
≥ \$75,000	127	93.0	(88.7 - 97.3)	8,397	94.5	(92.6 - 96.4)

¹ Defined as believing it was at least somewhat acceptable to see and hear discussions about sexually transmitted disease risks in public forums in their community

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 62a. Overweight among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	630	26,900	40.0	(34.8 - 45.2)	36,710	5,213,000	36.4	(34.0 - 38.8)
Gender								
Male	233	15,600	46.3	(37.8 - 54.8)	14,502	3,073,200	43.7	(39.9 - 47.6)
Female	397	11,300	33.7	(27.9 - 39.5)	22,208	2,139,800	29.3	(26.4 - 32.2)
Age (years)								
18-34	*	*	*	*	3,789	1,295,200	30.1	(24.7 - 35.6)
35-44	*	*	*	*	4,785	1,052,000	37.5	(31.7 - 43.3)
45-54	*	*	*	*	7,495	1,053,300	38.8	(34.1 - 43.4)
55-64	153	3,900	35.8	(27.4 - 44.2)	8,356	798,100	38.8	(34.8 - 42.7)
≥ 65	198	5,600	47.0	(39.4 - 54.6)	12,285	1,014,400	41.5	(37.6 - 45.4)
Race/Ethnicity								
White Non-Hispanic	606	25,700	40.8	(35.7 - 45.9)	34,501	3,282,200	36.4	(34.4 - 38.3)
Black Non-Hispanic	*	*	*	*	708	740,000	36.7	(28.3 - 45.1)
Hispanic	*	*	*	*	813	781,500	35.6	(27.1 - 44.1)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	300	12,600	40.9	(33.3 - 48.6)	15,079	1,889,700	35.4	(31.2 - 39.5)
Some College	*	*	*	*	9,515	1,314,500	38.7	(33.8 - 43.7)
≥ College Degree	174	7,600	40.2	(31.1 - 49.3)	12,062	1,996,900	36.3	(32.8 - 39.7)
Income								
≤ \$24,999	157	3,700	29.4	(20.1 - 38.7)	9,346	1,170,900	34.0	(28.2 - 39.8)
\$25,000-\$49,999	172	6,700	39.9	(30.8 - 49.0)	9,368	1,136,000	40.4	(34.6 - 46.2)
\$50,000-\$74,999	*	*	*	*	5,615	738,700	38.3	(32.3 - 44.3)
≥ \$75,000	*	*	*	*	8,337	1,659,100	37.5	(34.1 - 40.9)

¹ Defined as having a body mass index (BMI) more than 24.9 but less than 30.0. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Sample size from survey

³ Estimated number of adults who are overweight (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 62b. Overweight among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	630	40.1	(34.5 - 45.8)	36,710	36.2	(33.8 - 38.6)
Gender						
Male	233	46.5	(37.3 - 55.6)	14,502	43.9	(40.1 - 47.6)
Female	397	33.8	(27.4 - 40.3)	22,208	28.9	(25.9 - 31.8)
Race/Ethnicity						
White Non-Hispanic	606	41.1	(35.5 - 46.8)	34,501	36.0	(34.0 - 38.0)
Black Non-Hispanic	*	*	*	708	37.0	(28.5 - 45.5)
Hispanic	*	*	*	813	36.0	(27.5 - 44.4)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	300	40.9	(32.6 - 49.1)	15,079	35.2	(31.1 - 39.4)
Some College	*	*	*	9,515	38.3	(33.3 - 43.2)
≥ College Degree	174	40.0	(30.2 - 49.9)	12,062	36.2	(32.7 - 39.8)
Income						
≤ \$24,999	*	*	*	9,346	33.8	(28.0 - 39.5)
\$25,000-\$49,999	172	40.6	(30.8 - 50.4)	9,368	40.6	(34.8 - 46.4)
\$50,000-\$74,999	*	*	*	5,615	38.6	(32.5 - 44.7)
≥ \$75,000	*	*	*	8,337	37.2	(33.7 - 40.6)

¹ Defined as having a body mass index (BMI) more than 24.9 but less than 30.0. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 63a. Obesity among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	630	21,400	31.7	(26.5 - 36.9)	36,710	3,327,400	23.2	(21.2 - 25.3)
Gender								
Male	233	11,800	35.0	(26.5 - 43.6)	14,502	1,505,800	21.4	(19.0 - 23.9)
Female	397	9,600	28.4	(22.6 - 34.2)	22,208	1,821,600	25.0	(21.8 - 28.1)
Age (years)								
18-34	*	*	*	*	3,789	730,600	17.0	(12.4 - 21.6)
35-44	*	*	*	*	4,785	662,200	23.6	(19.4 - 27.9)
45-54	132	3,800	26.6	(18.1 - 35.1)	7,495	816,800	30.1	(25.3 - 34.8)
55-64	153	4,300	40.2	(31.8 - 48.6)	8,356	563,900	27.4	(23.7 - 31.1)
≥ 65	198	2,500	21.5	(15.4 - 27.5)	12,285	554,000	22.7	(19.2 - 26.1)
Race/Ethnicity								
White Non-Hispanic	606	19,000	30.1	(25.3 - 35.0)	34,501	2,062,700	22.9	(21.2 - 24.6)
Black Non-Hispanic	*	*	*	*	708	703,100	34.9	(26.1 - 43.7)
Hispanic	*	*	*	*	813	456,400	20.8	(14.3 - 27.2)
Other Non-Hispanic	*	*	*	*	688	105,200	9.6	(4.0 - 15.3)
Education								
≤ High School	300	10,600	34.2	(26.0 - 42.5)	15,079	1,561,600	29.2	(25.3 - 33.2)
Some College	155	5,400	31.5	(22.1 - 40.9)	9,515	792,800	23.4	(19.4 - 27.3)
≥ College Degree	174	5,400	28.3	(19.2 - 37.4)	12,062	915,000	16.6	(14.4 - 18.8)
Income								
≤ \$24,999	*	*	*	*	9,346	974,800	28.3	(22.6 - 34.0)
\$25,000-\$49,999	172	4,200	25.0	(17.3 - 32.7)	9,368	584,800	20.8	(17.1 - 24.5)
\$50,000-\$74,999	*	*	*	*	5,615	475,300	24.6	(20.0 - 29.3)
≥ \$75,000	*	*	*	*	8,337	867,700	19.6	(17.0 - 22.3)

¹ Defined as having a body mass index (BMI) of 30.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Sample size from survey

³ Estimated number of adults who are obese (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 63b. Obesity among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	630	31.6	(25.9 - 37.3)	36,710	23.1	(21.0 - 25.1)
Gender						
Male	233	35.6	(26.3 - 44.9)	14,502	21.5	(19.0 - 24.0)
Female	397	27.6	(21.2 - 34.0)	22,208	24.6	(21.5 - 27.8)
Race/Ethnicity						
White Non-Hispanic	606	29.8	(24.5 - 35.1)	34,501	22.8	(21.0 - 24.5)
Black Non-Hispanic	*	*	*	708	34.8	(25.9 - 43.6)
Hispanic	*	*	*	813	20.5	(14.1 - 26.9)
Other Non-Hispanic	*	*	*	688	9.5	(3.8 - 15.2)
Education						
≤ High School	300	34.7	(25.6 - 43.8)	15,079	29.4	(25.4 - 33.3)
Some College	*	*	*	9,515	23.1	(19.2 - 27.1)
≥ College Degree	*	*	*	12,062	16.4	(14.2 - 18.5)
Income						
≤ \$24,999	*	*	*	9,346	28.3	(22.6 - 34.1)
\$25,000-\$49,999	172	24.0	(15.8 - 32.1)	9,368	21.0	(17.2 - 24.8)
\$50,000-\$74,999	*	*	*	5,615	24.0	(19.4 - 28.7)
≥ \$75,000	*	*	*	8,337	19.3	(16.7 - 21.9)

¹ Defined as having a body mass index (BMI) of 30.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 64a. Overweight or obesity among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	630	48,300	71.7	(67.3 - 76.1)	36,710	8,540,500	59.6	(57.0 - 62.2)
Gender								
Male	233	27,400	81.3	(75.4 - 87.2)	14,502	4,579,100	65.2	(61.0 - 69.4)
Female	397	20,900	62.1	(56.1 - 68.0)	22,208	3,961,400	54.3	(51.0 - 57.6)
Age (years)								
18-34	*	*	*	*	3,789	2,025,800	47.1	(40.8 - 53.5)
35-44	97	14,100	76.8	(67.4 - 86.3)	4,785	1,714,200	61.1	(55.6 - 66.6)
45-54	132	9,700	68.3	(59.4 - 77.2)	7,495	1,870,100	68.8	(64.6 - 73.1)
55-64	153	8,200	76.0	(68.8 - 83.2)	8,356	1,362,000	66.2	(61.9 - 70.4)
≥ 65	198	8,100	68.5	(61.5 - 75.5)	12,285	1,568,400	64.1	(60.6 - 67.7)
Race/Ethnicity								
White Non-Hispanic	606	44,600	71.0	(66.5 - 75.4)	34,501	5,344,900	59.3	(57.2 - 61.3)
Black Non-Hispanic	*	*	*	*	708	1,443,100	71.6	(62.5 - 80.7)
Hispanic	*	*	*	*	813	1,237,900	56.4	(46.5 - 66.2)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	300	23,200	75.2	(69.4 - 80.9)	15,079	3,451,400	64.6	(59.8 - 69.4)
Some College	155	12,200	70.3	(60.9 - 79.8)	9,515	2,107,300	62.1	(57.0 - 67.2)
≥ College Degree	174	12,900	68.5	(60.1 - 76.9)	12,062	2,911,900	52.9	(49.2 - 56.6)
Income								
≤ \$24,999	157	9,700	78.2	(70.3 - 86.1)	9,346	2,145,700	62.3	(55.7 - 68.9)
\$25,000-\$49,999	172	10,900	64.9	(56.0 - 73.8)	9,368	1,720,700	61.2	(54.5 - 67.9)
\$50,000-\$74,999	105	10,300	77.2	(68.5 - 85.8)	5,615	1,214,000	62.9	(57.1 - 68.8)
≥ \$75,000	127	13,100	73.5	(64.5 - 82.6)	8,337	2,526,700	57.1	(53.6 - 60.6)

¹ Defined as having a body mass index (BMI) of 25.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Sample size from survey

³ Estimated number of adults who are overweight or obese (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 64b. Overweight or obesity among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	630	71.7	(67.0 - 76.5)	36,710	59.3	(56.8 - 61.9)
Gender						
Male	233	82.1	(75.9 - 88.3)	14,502	65.4	(61.3 - 69.4)
Female	397	61.4	(54.8 - 68.0)	22,208	53.5	(50.1 - 56.8)
Race/Ethnicity						
White Non-Hispanic	606	70.9	(66.0 - 75.8)	34,501	58.8	(56.7 - 60.8)
Black Non-Hispanic	*	*	*	708	71.8	(63.0 - 80.6)
Hispanic	*	*	*	813	56.5	(46.8 - 66.1)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	300	75.6	(69.5 - 81.6)	15,079	64.6	(59.9 - 69.3)
Some College	*	*	*	9,515	61.4	(56.4 - 66.5)
≥ College Degree	174	68.6	(59.4 - 77.7)	12,062	52.6	(48.9 - 56.3)
Income						
≤ \$24,999	157	79.3	(71.1 - 87.5)	9,346	62.1	(55.6 - 68.6)
\$25,000-\$49,999	172	64.5	(54.8 - 74.2)	9,368	61.6	(55.1 - 68.1)
\$50,000-\$74,999	105	78.1	(69.1 - 87.2)	5,615	62.6	(56.7 - 68.6)
≥ \$75,000	127	72.9	(63.1 - 82.7)	8,337	56.5	(52.9 - 60.0)

¹ Defined as having a body mass index (BMI) of 25.0 or greater. BMI is calculated as weight in kilograms divided by the square of height in meters.

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 65a. Received advice about weight by a health professional among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	633	17,600	26.2	(21.4 - 31.0)	36,729	3,929,700	27.4	(25.1 - 29.7)
Gender								
Male	233	8,900	26.5	(18.7 - 34.4)	14,499	1,654,800	23.6	(20.6 - 26.6)
Female	400	8,800	25.9	(20.3 - 31.5)	22,230	2,274,900	31.0	(27.7 - 34.3)
Age (years)								
18-34	*	*	*	*	3,796	895,500	20.7	(15.3 - 26.2)
35-44	*	*	*	*	4,779	722,700	25.8	(20.9 - 30.6)
45-54	132	3,600	24.9	(16.9 - 33.0)	7,490	910,900	33.5	(29.0 - 38.1)
55-64	155	3,800	35.5	(27.4 - 43.7)	8,351	741,100	36.1	(32.0 - 40.3)
≥ 65	200	2,900	24.2	(17.8 - 30.7)	12,313	659,500	26.9	(23.0 - 30.7)
Race/Ethnicity								
White Non-Hispanic	609	15,200	24.2	(20.0 - 28.4)	34,502	2,386,200	26.5	(24.6 - 28.4)
Black Non-Hispanic	*	*	*	*	709	615,700	30.4	(22.0 - 38.8)
Hispanic	*	*	*	*	830	555,400	25.1	(17.8 - 32.4)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	301	8,800	28.4	(20.6 - 36.3)	15,116	1,469,700	27.5	(23.6 - 31.4)
Some College	156	5,100	30.0	(20.5 - 39.5)	9,515	946,800	27.8	(23.2 - 32.4)
≥ College Degree	174	3,700	19.7	(12.7 - 26.7)	12,041	1,445,500	26.2	(23.0 - 29.4)
Income								
≤ \$24,999	*	*	*	*	9,360	1,079,400	31.4	(25.3 - 37.5)
\$25,000-\$49,999	173	3,500	20.9	(14.3 - 27.4)	9,365	696,900	24.6	(19.8 - 29.4)
\$50,000-\$74,999	*	*	*	*	5,603	507,100	26.3	(21.6 - 31.0)
≥ \$75,000	127	4,900	27.7	(18.7 - 36.7)	8,334	1,207,100	27.3	(24.2 - 30.5)

¹ Defined as being advised by a doctor, nurse or other health professional about their weight (lose, gain or maintain)

² Sample size from survey

³ Estimated number of adults provided advice by a health professional about their weight (lose, gain or maintain) (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wayne County**

Table 65b. Received advice about weight by a health professional among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	633	25.5	(20.2 - 30.8)	36,729	27.1	(24.8 - 29.4)
Gender						
Male	233	25.9	(17.4 - 34.4)	14,499	23.4	(20.5 - 26.3)
Female	400	25.1	(18.9 - 31.3)	22,230	30.6	(27.3 - 34.0)
Race/Ethnicity						
White Non-Hispanic	609	23.1	(18.6 - 27.6)	34,502	26.1	(24.2 - 28.0)
Black Non-Hispanic	*	*	*	709	30.4	(21.9 - 38.9)
Hispanic	*	*	*	830	24.9	(17.7 - 32.2)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	301	28.2	(19.6 - 36.9)	15,116	27.3	(23.4 - 31.2)
Some College	*	*	*	9,515	27.4	(22.9 - 32.0)
≥ College Degree	174	18.4	(11.0 - 25.9)	12,041	25.9	(22.7 - 29.0)
Income						
≤ \$24,999	*	*	*	9,360	31.0	(24.9 - 37.0)
\$25,000-\$49,999	173	19.3	(12.7 - 25.8)	9,365	24.9	(20.0 - 29.8)
\$50,000-\$74,999	*	*	*	5,603	25.9	(21.2 - 30.6)
≥ \$75,000	127	26.1	(16.7 - 35.5)	8,334	26.9	(23.7 - 30.0)

¹ Defined as being advised by a doctor, nurse or other health professional about their weight (lose, gain or maintain)

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 66a. Received advice to lose weight by a health professional among adults provided weight advice¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	170	15,900	90.4	(84.1 - 96.7)	9,724	3,412,900	86.8	(83.9 - 89.8)
Gender								
Male	63	8,400	94.1	(84.4 - 100)	3,807	1,468,700	88.8	(85.7 - 91.8)
Female	107	7,600	86.6	(78.5 - 94.6)	5,917	1,944,200	85.5	(80.8 - 90.1)
Age (years)								
18-34	*	*	*	*	681	779,700	87.1	(80.6 - 93.6)
35-44	*	*	*	*	1,215	622,100	86.1	(76.9 - 95.2)
45-54	*	*	*	*	2,238	813,300	89.3	(84.4 - 94.2)
55-64	56	3,500	90.6	(84.1 - 97.2)	2,697	645,500	87.1	(81.2 - 93.0)
≥ 65	*	*	*	*	2,893	552,300	83.7	(76.9 - 90.6)
Race/Ethnicity								
White Non-Hispanic	163	13,500	88.9	(81.8 - 96.0)	9,082	2,076,700	87.0	(84.5 - 89.6)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	251	483,600	87.1	(78.5 - 95.7)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	84	8,200	93.6	(89.0 - 98.3)	3,962	1,239,700	84.3	(78.6 - 90.1)
Some College	*	*	*	*	2,633	814,800	86.1	(80.1 - 92.0)
≥ College Degree	*	*	*	*	3,115	1,290,900	89.3	(85.3 - 93.3)
Income								
≤ \$24,999	*	*	*	*	2,539	922,100	85.4	(78.7 - 92.1)
\$25,000-\$49,999	*	*	*	*	2,482	594,100	85.2	(78.4 - 92.1)
\$50,000-\$74,999	*	*	*	*	1,521	446,100	88.0	(83.2 - 92.7)
≥ \$75,000	*	*	*	*	2,270	1,073,200	88.9	(83.7 - 94.1)

¹ Defined as provided advice by a health professional about their weight and advised to lose weight

² Sample size from survey

³ Estimated number of adults provided advice by a health professional about their weight and advised to lose weight (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 66b. Received advice to lose weight by a health professional among adults provided weight advice¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	170	90.1	(82.8 - 97.4)	9,724	86.9	(83.8 - 89.9)
Gender						
Male	*	*	*	3,807	88.7	(85.7 - 91.8)
Female	107	86.8	(77.7 - 95.9)	5,917	85.5	(80.7 - 90.3)
Race/Ethnicity						
White Non-Hispanic	163	88.3	(79.9 - 96.7)	9,082	87.0	(84.4 - 89.6)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	251	87.6	(79.3 - 95.8)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	84	94.4	(90.0 - 98.8)	3,962	84.6	(78.7 - 90.5)
Some College	*	*	*	2,633	85.6	(79.5 - 91.7)
≥ College Degree	*	*	*	3,115	89.4	(85.4 - 93.3)
Income						
≤ \$24,999	*	*	*	2,539	85.7	(79.0 - 92.3)
\$25,000-\$49,999	*	*	*	2,482	85.3	(78.4 - 92.3)
\$50,000-\$74,999	*	*	*	1,521	88.0	(83.1 - 92.8)
≥ \$75,000	*	*	*	2,270	88.7	(83.1 - 94.2)

¹ Defined as provided advice by a health professional about their weight and advised to lose weight

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 67a. No leisure-time physical activity among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	655	17,100	24.6	(20.0 - 29.2)	38,128	3,541,700	23.9	(21.6 - 26.1)
Gender								
Male	235	8,700	25.8	(18.0 - 33.6)	14,602	1,482,500	20.9	(17.8 - 24.0)
Female	420	8,300	23.5	(18.5 - 28.4)	23,526	2,059,200	26.6	(23.4 - 29.7)
Age (years)								
18-34	*	*	*	*	3,938	982,900	22.0	(16.6 - 27.5)
35-44	*	*	*	*	4,988	556,000	19.3	(15.0 - 23.6)
45-54	141	3,400	22.4	(14.3 - 30.6)	7,827	759,000	26.8	(21.7 - 31.9)
55-64	160	3,400	30.7	(22.9 - 38.4)	8,765	464,900	21.7	(18.2 - 25.2)
≥ 65	203	3,600	30.0	(23.2 - 36.8)	12,610	778,800	30.7	(27.3 - 34.2)
Race/Ethnicity								
White Non-Hispanic	631	15,400	23.7	(19.6 - 27.9)	35,816	1,892,800	20.3	(18.6 - 22.0)
Black Non-Hispanic	*	*	*	*	739	690,000	32.0	(23.4 - 40.6)
Hispanic	*	*	*	*	860	600,800	26.7	(19.1 - 34.4)
Other Non-Hispanic	*	*	*	*	*	*	*	*
Education								
≤ High School	311	11,400	35.7	(28.0 - 43.4)	15,683	1,810,200	32.5	(28.4 - 36.5)
Some College	164	3,600	19.9	(11.8 - 28.0)	9,906	814,200	23.0	(18.6 - 27.3)
≥ College Degree	178	2,100	10.8	(5.5 - 16.0)	12,458	841,000	14.9	(11.9 - 18.0)
Income								
≤ \$24,999	*	*	*	*	9,640	1,332,200	38.0	(31.7 - 44.3)
\$25,000-\$49,999	179	3,900	22.4	(14.8 - 30.0)	9,658	648,000	22.3	(18.3 - 26.3)
\$50,000-\$74,999	108	2,300	16.7	(8.8 - 24.7)	5,773	430,700	21.8	(16.6 - 27.0)
≥ \$75,000	131	3,200	17.4	(8.7 - 26.1)	8,565	560,800	12.5	(10.3 - 14.6)

¹ Defined as having not participated in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise

² Sample size from survey

³ Estimated number of adults who have not participated in any physical activities or exercises (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 67b. No leisure-time physical activity among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	655	23.9	(18.9 - 28.9)	38,128	23.7	(21.5 - 25.9)
Gender						
Male	235	25.6	(17.2 - 34.1)	14,602	20.9	(17.8 - 23.9)
Female	420	22.3	(16.9 - 27.6)	23,526	26.3	(23.1 - 29.5)
Race/Ethnicity						
White Non-Hispanic	631	22.9	(18.4 - 27.3)	35,816	20.0	(18.3 - 21.7)
Black Non-Hispanic	*	*	*	739	32.0	(23.3 - 40.6)
Hispanic	*	*	*	860	26.9	(19.2 - 34.5)
Other Non-Hispanic	*	*	*	*	*	*
Education						
≤ High School	311	34.9	(26.6 - 43.3)	15,683	32.3	(28.3 - 36.4)
Some College	164	19.7	(10.8 - 28.6)	9,906	22.8	(18.5 - 27.2)
≥ College Degree	178	10.4	(4.9 - 15.9)	12,458	14.9	(11.8 - 17.9)
Income						
≤ \$24,999	*	*	*	9,640	37.9	(31.6 - 44.2)
\$25,000-\$49,999	179	21.6	(13.6 - 29.6)	9,658	22.0	(18.0 - 26.0)
\$50,000-\$74,999	108	15.8	(7.5 - 24.0)	5,773	21.6	(16.3 - 26.8)
≥ \$75,000	131	17.4	(7.9 - 26.9)	8,565	12.6	(10.3 - 14.9)

¹ Defined as having not participated in any physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 68a. Consumption of 5 or more servings of fruits and vegetables per day among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	658	17,900	25.7	(21.4 - 30.0)	38,203	4,023,400	27.0	(24.9 - 29.2)
Gender								
Male	236	7,000	20.6	(13.6 - 27.6)	14,634	1,603,900	22.6	(19.4 - 25.8)
Female	422	10,900	30.6	(25.3 - 35.8)	23,569	2,419,500	31.1	(28.2 - 34.0)
Age (years)								
18-34	*	*	*	*	3,940	1,261,700	28.3	(22.7 - 33.9)
35-44	*	*	*	*	4,989	688,000	23.9	(19.5 - 28.3)
45-54	141	3,500	23.0	(15.4 - 30.6)	7,837	746,800	26.2	(22.3 - 30.1)
55-64	162	2,700	24.1	(16.9 - 31.3)	8,777	579,100	27.0	(23.6 - 30.5)
≥ 65	204	3,900	31.9	(25.0 - 38.7)	12,660	747,800	29.4	(26.0 - 32.8)
Race/Ethnicity								
White Non-Hispanic	634	17,500	26.9	(22.4 - 31.3)	35,884	2,538,000	27.2	(25.4 - 28.9)
Black Non-Hispanic	*	*	*	*	740	572,400	26.5	(19.2 - 33.9)
Hispanic	*	*	*	*	864	672,200	29.8	(21.0 - 38.5)
Other Non-Hispanic	*	*	*	*	715	240,800	21.4	(12.0 - 30.8)
Education								
≤ High School	312	7,600	23.8	(17.8 - 29.8)	15,726	1,309,800	23.4	(19.3 - 27.4)
Some College	166	3,600	20.0	(11.3 - 28.6)	9,924	964,300	27.2	(23.0 - 31.3)
≥ College Degree	178	6,700	34.8	(26.0 - 43.5)	12,471	1,747,500	31.0	(27.9 - 34.1)
Income								
≤ \$24,999	162	2,800	22.0	(14.2 - 29.8)	9,667	831,500	23.7	(18.6 - 28.7)
\$25,000-\$49,999	179	4,200	24.3	(17.1 - 31.4)	9,669	917,200	31.6	(25.6 - 37.6)
\$50,000-\$74,999	*	*	*	*	5,777	501,000	25.3	(21.4 - 29.3)
≥ \$75,000	131	5,100	27.9	(18.3 - 37.5)	8,569	1,260,100	28.0	(25.0 - 31.0)

¹ Defined as having consumed 5 or more servings of fruits and vegetables in an average day

² Sample size from survey

³ Estimated number of adults who consume 5 or more servings of fruits and vegetables per day (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 68b. Consumption of 5 or more servings of fruits and vegetables per day among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	658	25.8	(21.0 - 30.5)	38,203	27.1	(24.9 - 29.2)
Gender						
Male	236	20.9	(13.2 - 28.5)	14,634	22.7	(19.6 - 25.9)
Female	422	30.5	(24.7 - 36.2)	23,569	31.0	(28.1 - 34.0)
Race/Ethnicity						
White Non-Hispanic	634	27.0	(22.1 - 31.9)	35,884	27.1	(25.4 - 28.9)
Black Non-Hispanic	*	*	*	740	26.8	(19.4 - 34.2)
Hispanic	*	*	*	864	29.5	(20.9 - 38.1)
Other Non-Hispanic	*	*	*	715	21.8	(12.2 - 31.5)
Education						
≤ High School	312	24.2	(17.5 - 30.8)	15,726	23.3	(19.3 - 27.3)
Some College	166	20.2	(10.7 - 29.7)	9,924	27.1	(22.9 - 31.2)
≥ College Degree	178	34.1	(24.7 - 43.6)	12,471	31.2	(28.0 - 34.3)
Income						
≤ \$24,999	162	20.3	(12.3 - 28.2)	9,667	23.9	(18.8 - 28.9)
\$25,000-\$49,999	179	25.2	(17.3 - 33.1)	9,669	32.0	(26.0 - 38.0)
\$50,000-\$74,999	*	*	*	5,777	25.0	(21.0 - 28.9)
≥ \$75,000	*	*	*	8,569	28.0	(24.9 - 31.1)

¹ Defined as having consumed 5 or more servings of fruits and vegetables in an average day

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 69a. Had a fall within the past 3 months among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	498	5,900	15.5	(11.8 - 19.2)	28,783	1,244,800	17.0	(14.9 - 19.0)
Gender								
Male	184	2,300	12.8	(7.0 - 18.6)	10,966	489,400	14.4	(11.9 - 16.8)
Female	314	3,500	18.0	(13.4 - 22.7)	17,817	755,300	19.2	(16.1 - 22.3)
Age (years)								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	140	2,600	17.5	(10.3 - 24.6)	7,741	445,500	16.0	(12.0 - 20.0)
55-64	154	1,500	14.5	(8.4 - 20.6)	8,454	336,000	16.5	(13.4 - 19.6)
≥ 65	204	1,700	14.0	(9.0 - 19.1)	12,588	463,300	18.4	(15.2 - 21.6)
Race/Ethnicity								
White Non-Hispanic	483	5,600	15.4	(11.7 - 19.2)	27,456	869,000	16.6	(15.1 - 18.1)
Black Non-Hispanic	*	*	*	*	*	*	*	*
Hispanic	*	*	*	*	424	76,100	11.0	(4.5 - 17.4)
Other Non-Hispanic	*	*	*	*	445	44,300	10.0	(3.1 - 16.8)
Education								
≤ High School	252	3,300	17.4	(11.5 - 23.2)	12,417	493,000	17.3	(13.3 - 21.3)
Some College	120	1,100	12.8	(6.3 - 19.3)	7,232	299,400	18.7	(14.6 - 22.7)
≥ College Degree	125	1,400	14.5	(8.4 - 20.5)	9,086	441,500	15.4	(12.9 - 17.9)
Income								
≤ \$24,999	140	1,300	16.0	(9.3 - 22.7)	7,824	423,900	25.0	(18.5 - 31.6)
\$25,000-\$49,999	140	2,100	20.0	(11.8 - 28.3)	7,445	223,600	16.0	(12.8 - 19.2)
\$50,000-\$74,999	76	1,100	15.5	(5.6 - 25.3)	4,141	151,200	14.6	(10.8 - 18.4)
≥ \$75,000	82	700	9.4	(3.6 - 15.1)	5,874	287,100	12.5	(10.0 - 15.1)

¹ Defined as respondent unintentionally coming to rest on the ground or another lower level one or more times within the past 3 months

² Sample size from survey

³ Estimated number of adults who have had a fall within the past 3 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 69b. Had a fall within the past 3 months among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	498	15.4	(11.8 - 19.1)	28,783	17.3	(15.0 - 19.5)
Gender						
Male	184	12.6	(7.0 - 18.3)	10,966	15.0	(12.1 - 17.8)
Female	314	18.0	(13.3 - 22.7)	17,817	19.4	(16.0 - 22.8)
Race/Ethnicity						
White Non-Hispanic	483	15.2	(11.5 - 18.9)	27,456	16.9	(15.2 - 18.7)
Black Non-Hispanic	*	*	*	*	*	*
Hispanic	*	*	*	424	10.7	(4.2 - 17.1)
Other Non-Hispanic	*	*	*	445	10.0	(2.9 - 17.0)
Education						
≤ High School	252	16.8	(11.1 - 22.5)	12,417	17.4	(13.0 - 21.7)
Some College	120	12.5	(6.1 - 18.8)	7,232	19.0	(14.7 - 23.2)
≥ College Degree	125	15.3	(8.9 - 21.7)	9,086	16.0	(13.1 - 18.8)
Income						
≤ \$24,999	140	15.8	(9.0 - 22.6)	7,824	25.3	(18.3 - 32.4)
\$25,000-\$49,999	140	19.6	(11.5 - 27.6)	7,445	15.9	(12.6 - 19.2)
\$50,000-\$74,999	76	15.1	(5.3 - 24.9)	4,141	14.7	(10.6 - 18.8)
≥ \$75,000	82	9.6	(3.7 - 15.5)	5,874	12.9	(9.9 - 15.8)

¹ Defined as respondent unintentionally coming to rest on the ground or another lower level one or more times within the past 3 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

Expanded BRFSS Report: July 2008 - June 2009
Wayne County

Table 70a. Had a fall that caused injury within the past 3 months among adults¹ in Wayne County and New York State

	Wayne County				New York State			
	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴	n ²	Est. # of Adults ³	Percent	(C.I.) ⁴
Total	498	2,400	6.4	(3.9 - 8.8)	28,760	335,700	4.6	(3.7 - 5.5)
Gender								
Male	*	*	*	*	10,957	120,000	3.5	(2.5 - 4.6)
Female	314	1,700	8.5	(5.1 - 12.0)	17,803	215,600	5.5	(4.1 - 6.9)
Age (years)								
18-34	*	*	*	*	*	*	*	*
35-44	*	*	*	*	*	*	*	*
45-54	140	1,300	8.7	(3.6 - 13.8)	7,737	115,100	4.1	(2.9 - 5.4)
55-64	*	*	*	*	8,449	90,300	4.4	(2.7 - 6.1)
≥ 65	*	*	*	*	12,574	130,300	5.2	(3.4 - 6.9)
Race/Ethnicity								
White Non-Hispanic	483	2,200	6.0	(3.5 - 8.6)	27,433	270,800	5.2	(4.2 - 6.1)
Black Non-Hispanic	*	*	*	*	458	47,900	4.9	(0.9 - 9.0)
Hispanic	*	*	*	*	424	12,400	1.8	(0.2 - 3.3)
Other Non-Hispanic	*	*	*	*	445	4,600	1.0	(0.4 - 1.6)
Education								
≤ High School	252	1,300	6.9	(3.0 - 10.8)	12,404	148,900	5.2	(3.6 - 6.8)
Some College	*	*	*	*	7,228	85,400	5.3	(3.1 - 7.5)
≥ College Degree	*	*	*	*	9,081	101,300	3.5	(2.4 - 4.6)
Income								
≤ \$24,999	140	600	7.6	(2.5 - 12.8)	7,812	90,200	5.3	(3.6 - 7.1)
\$25,000-\$49,999	140	700	6.6	(2.3 - 10.9)	7,442	93,200	6.7	(3.9 - 9.4)
\$50,000-\$74,999	*	*	*	*	4,139	37,000	3.6	(1.8 - 5.3)
≥ \$75,000	*	*	*	*	5,872	54,700	2.4	(1.5 - 3.3)

¹ Defined as as respondent unintentionally coming to rest on the ground or another lower level and caused respondent to limit regular activities for at least a day or to go see a doctor one or more times within the past 3 months

² Sample size from survey

³ Estimated number of adults who have had a fall that caused an injury within the past 3 months (rounded to the nearest hundred)

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)

**Expanded BRFSS Report: July 2008 - June 2009
Wayne County**

Table 70b. Had a fall that caused injury within the past 3 months among adults¹ in Wayne County and New York State - Age-Adjusted²

	Wayne County			New York State		
	n ³	Percent	(C.I.) ⁴	n ³	Percent	(C.I.) ⁴
Total	498	6.2	(3.8 - 8.6)	28,760	4.6	(3.6 - 5.5)
Gender						
Male	*	*	*	10,957	3.6	(2.4 - 4.7)
Female	314	8.5	(5.0 - 11.9)	17,803	5.5	(4.1 - 6.9)
Race/Ethnicity						
White Non-Hispanic	483	5.9	(3.4 - 8.3)	27,433	5.2	(4.2 - 6.2)
Black Non-Hispanic	*	*	*	458	4.7	(0.7 - 8.7)
Hispanic	*	*	*	424	1.8	(0.1 - 3.4)
Other Non-Hispanic	*	*	*	445	1.0	(0.4 - 1.6)
Education						
≤ High School	252	6.7	(2.9 - 10.4)	12,404	5.2	(3.5 - 6.8)
Some College	*	*	*	7,228	5.3	(3.1 - 7.5)
≥ College Degree	*	*	*	9,081	3.6	(2.4 - 4.7)
Income						
≤ \$24,999	140	7.3	(2.4 - 12.2)	7,812	5.1	(3.3 - 7.0)
\$25,000-\$49,999	140	6.3	(2.2 - 10.4)	7,442	6.6	(3.9 - 9.2)
\$50,000-\$74,999	*	*	*	4,139	3.7	(1.7 - 5.8)
≥ \$75,000	*	*	*	5,872	2.4	(1.4 - 3.3)

¹ Defined as as respondent unintentionally coming to rest on the ground or another lower level and caused respondent to limit regular activities for at least a day or to go see a doctor one or more times within the past 3 months

² Prevalence estimates are age-adjusted to the 2000 US Standard population

³ Sample size from survey

⁴ 95% confidence interval

* Data do not meet reporting criteria (confidence interval with a half-width greater than 10, denominator less than 50 and/or a numerator less than 10)