New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
All Invasive Malignant Tumors

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	28739	461.7	5.6	30569	359.1	4.1	16784	276.3	4.4	15720	181.5	2.9
1977	28839	464.0	5.6	30466	357.2	4.1	16936	278.3	4.4	15851	182.3	2.9
1978	28978	467.5	5.6	30690	359.5	4.1	16721	275.7	4.4	15696	179.3	2.9
1979	29500	475.6	5.7	31271	365.3	4.2	16522	272.3	4.4	15809	180.1	2.9
1980	30966	497.3	5.8	31714	369.7	4.2	17047	280.7	4.4	16119	182.0	2.9
1981	30639	494.0	5.8	32281	375.6	4.2	16659	274.4	4.4	16006	180.6	2.9
1982	31003	498.8	5.8	32739	379.6	4.2	16614	274.3	4.4	15994	179.9	2.8
1983	31691	507.1	5.8	33344	386.5	4.3	16571	273.3	4.3	16088	179.6	2.8
1984	31736	506.6	5.8	33898	392.2	4.3	16618	272.7	4.3	16354	182.4	2.9
1985	32861	522.1	5.9	34437	399.0	4.3	16682	272.8	4.3	16262	181.3	2.8
1986	32699	520.6	5.8	34595	400.8	4.3	16603	271.2	4.3	16398	182.3	2.9
1987	32663	518.2	5.8	34974	403.8	4.3	16550	270.6	4.3	16299	180.3	2.8
1988	32189	511.4	5.8	34921	403.7	4.3	16728	273.3	4.3	16424	181.9	2.8
1989	32714	517.8	5.8	34553	401.0	4.3	16163	263.8	4.2	16158	179.3	2.8
1990	34455	541.3	5.9	36014	418.1	4.4	16046	261.7	4.2	16159	178.3	2.8
1991	35676	558.9	5.9	36222	421.1	4.4	16200	263.3	4.2	16211	179.0	2.8
1992	37622	585.0	6.0	36265	420.5	4.4	16061	260.5	4.1	16364	179.8	2.8
1993	36821	567.4	5.9	36198	418.5	4.4	15924	256.6	4.1	16168	176.7	2.8
1994	37235	572.2	5.9	37706	435.0	4.5	16718	268.1	4.2	16635	181.5	2.8
1995	36319	554.8	5.8	37915	438.8	4.5	16245	259.7	4.1	16652	181.4	2.8
1996	37627	573.8	5.9	38563	446.9	4.5	15956	253.1	4.0	16508	180.0	2.8
1997	38218	578.8	5.9	39133	450.7	4.5	15645	246.2	3.9	15973	173.2	2.7
1998	38925	582.0	5.8	40087	461.0	4.6	15346	236.9	3.8	15764	169.8	2.7
1999	39311	583.6	5.8	40254	458.3	4.5	15562	238.6	3.8	15928	170.1	2.7
2000	40800	600.8	5.9	40772	460.2	4.5	15390	234.0	3.7	15971	168.1	2.6
2001	40343	588.4	5.8	40427	455.7	4.5	15241	229.6	3.7	15650	164.2	2.6
2002	41378	596.4	5.8	40108	449.1	4.5	15067	224.2	3.6	15547	162.4	2.6
2003	40634	580.5	5.7	40371	450.3	4.5	14562	214.1	3.5	15447	160.1	2.6
2004	40740	576.6	5.6	40633	453.1	4.5	14581	211.9	3.5	15139	156.6	2.5
2005	41362	581.1	5.6	40958	456.2	4.5	14385	207.8	3.4	14804	152.6	2.5
2006	43143	602.1	5.7	41511	464.1	4.6	14350	205.6	3.4	14839	153.3	2.5
2007	43694	600.9	5.7	41740	465.8	4.6	14430	203.3	3.3	14547	149.6	2.5
2008	43738	593.6	5.6	42764	474.9	4.6	14575	202.8	3.3	14534	147.9	2.5
2009	43619	583.0	5.5	42837	471.7	4.6	14353	197.3	3.3	14201	143.9	2.4
2010	42155	555.7	5.4	41807	459.1	4.5	14212	192.2	3.2	14549	145.7	2.4
2011	43180	560.2	5.4	42591	463.7	4.5	14097	187.4	3.1	14278	141.1	2.4
2012	41463	530.5	5.2	41975	455.6	4.5	14416	189.7	3.1	14282	141.0	2.4
2013	41474	523.0	5.1	43081	464.8	4.5	14209	182.7	3.1	14000	136.4	2.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Oral cavity and pharynx

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	1107	16.8	1.0	524	6.0	0.5	507	7.7	0.7	201	2.2	0.3
1977	1069	16.2	1.0	547	6.2	0.5	478	7.4	0.7	220	2.4	0.3
1978	1060	16.1	1.0	552	6.3	0.5	473	7.3	0.7	220	2.5	0.3
1979	1079	16.2	1.0	573	6.5	0.5	420	6.3	0.6	226	2.6	0.3
1980	1047	16.0	1.0	565	6.5	0.5	424	6.5	0.6	221	2.4	0.3
1981	1162	17.5	1.0	526	6.0	0.5	427	6.5	0.6	208	2.3	0.3
1982	1172	18.1	1.1	577	6.7	0.6	437	6.8	0.7	190	2.1	0.3
1983	1108	16.8	1.0	555	6.3	0.5	415	6.3	0.6	188	2.1	0.3
1984	1125	17.3	1.0	546	6.2	0.5	402	6.3	0.6	194	2.2	0.3
1985	1116	17.2	1.0	591	6.7	0.6	379	5.9	0.6	196	2.2	0.3
1986	1066	16.6	1.0	542	6.2	0.5	390	6.0	0.6	201	2.2	0.3
1987	1022	15.8	1.0	572	6.6	0.6	353	5.6	0.6	193	2.1	0.3
1988	990	15.2	1.0	564	6.5	0.5	382	6.2	0.6	209	2.3	0.3
1989	997	15.3	1.0	525	6.1	0.5	304	4.8	0.6	186	2.1	0.3
1990	993	15.3	1.0	550	6.4	0.6	312	4.9	0.6	182	2.0	0.3
1991	1001	15.5	1.0	568	6.6	0.6	300	4.8	0.6	174	1.9	0.3
1992	1041	16.0	1.0	559	6.5	0.6	272	4.2	0.5	172	1.8	0.3
1993	1023	15.9	1.0	528	6.1	0.5	285	4.4	0.5	151	1.7	0.3
1994	983	15.0	0.9	523	6.1	0.5	307	4.7	0.5	131	1.5	0.3
1995	864	13.1	0.9	515	5.9	0.5	294	4.6	0.5	151	1.7	0.3
1996	1007	15.0	0.9	529	6.2	0.5	257	3.9	0.5	156	1.8	0.3
1997	992	14.8	0.9	549	6.3	0.5	274	4.1	0.5	132	1.4	0.2
1998	960	14.2	0.9	565	6.5	0.5	265	4.0	0.5	153	1.6	0.3
1999	1053	15.3	0.9	492	5.6	0.5	251	3.7	0.5	124	1.3	0.2
2000	952	13.8	0.9	506	5.7	0.5	216	3.2	0.4	126	1.3	0.2
2001	983	13.9	0.9	552	6.1	0.5	245	3.6	0.4	138	1.4	0.2
2002	1025	14.4	0.9	551	6.1	0.5	228	3.3	0.4	131	1.3	0.2
2003	1034	14.4	0.9	535	5.9	0.5	214	3.0	0.4	127	1.3	0.2
2004	1084	14.9	0.9	545	6.0	0.5	260	3.6	0.4	145	1.5	0.2
2005	1028	13.9	0.9	535	6.0	0.5	225	3.1	0.4	119	1.2	0.2
2006	1164	15.7	0.9	535	6.0	0.5	227	3.1	0.4	115	1.2	0.2
2007	1221	16.2	0.9	573	6.4	0.5	229	3.1	0.4	113	1.2	0.2
2008	1177	15.5	0.9	589	6.5	0.5	250	3.3	0.4	95	1.0	0.2
2009	1278	16.5	0.9	570	6.2	0.5	247	3.3	0.4	117	1.1	0.2
2010	1250	15.8	0.9	620	6.7	0.5	252	3.3	0.4	129	1.3	0.2
2011	1300	16.2	0.9	567	6.0	0.5	232	3.0	0.4	119	1.2	0.2
2012	1284	15.9	0.9	609	6.4	0.5	271	3.4	0.4	120	1.1	0.2
2013	1354	16.7	0.9	583	6.0	0.5	257	3.1	0.4	114	1.1	0.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Esophagus

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	456	7.0	0.7	175	2.0	0.3	423	6.6	0.7	161	1.8	0.3
1977	422	6.6	0.7	200	2.3	0.3	383	6.1	0.6	185	2.1	0.3
1978	420	6.6	0.7	197	2.2	0.3	434	6.9	0.7	193	2.1	0.3
1979	441	6.8	0.7	194	2.1	0.3	379	5.9	0.6	181	2.0	0.3
1980	451	7.1	0.7	217	2.3	0.3	382	6.0	0.6	181	2.0	0.3
1981	418	6.5	0.7	208	2.3	0.3	429	6.8	0.7	173	1.9	0.3
1982	405	6.3	0.6	224	2.4	0.3	379	6.0	0.6	179	1.9	0.3
1983	447	6.9	0.7	190	2.1	0.3	416	6.7	0.7	159	1.7	0.3
1984	436	6.8	0.7	201	2.2	0.3	394	6.1	0.6	190	2.0	0.3
1985	440	6.9	0.7	172	1.9	0.3	405	6.3	0.6	146	1.6	0.3
1986	449	7.0	0.7	218	2.4	0.3	413	6.6	0.7	151	1.6	0.3
1987	407	6.3	0.6	210	2.3	0.3	396	6.2	0.6	177	1.9	0.3
1988	415	6.4	0.6	189	2.0	0.3	455	7.1	0.7	168	1.8	0.3
1989	457	7.0	0.7	199	2.1	0.3	422	6.6	0.6	172	1.8	0.3
1990	461	7.1	0.7	235	2.5	0.3	411	6.3	0.6	176	1.9	0.3
1991	453	6.9	0.6	217	2.4	0.3	424	6.5	0.6	179	1.9	0.3
1992	493	7.5	0.7	245	2.7	0.3	461	7.0	0.7	175	1.9	0.3
1993	496	7.5	0.7	202	2.2	0.3	455	7.0	0.7	163	1.7	0.3
1994	533	8.1	0.7	212	2.2	0.3	460	7.0	0.6	187	2.0	0.3
1995	479	7.3	0.7	228	2.5	0.3	505	7.8	0.7	178	1.9	0.3
1996	518	7.9	0.7	232	2.5	0.3	466	7.1	0.7	203	2.1	0.3
1997	481	7.3	0.7	220	2.4	0.3	488	7.4	0.7	163	1.7	0.3
1998	570	8.5	0.7	203	2.2	0.3	502	7.5	0.7	177	1.9	0.3
1999	546	8.0	0.7	198	2.1	0.3	506	7.5	0.7	175	1.8	0.3
2000	558	8.2	0.7	223	2.3	0.3	495	7.3	0.7	203	2.1	0.3
2001	544	7.9	0.7	200	2.1	0.3	538	7.8	0.7	163	1.7	0.3
2002	596	8.5	0.7	201	2.1	0.3	507	7.3	0.6	172	1.7	0.3
2003	613	8.6	0.7	217	2.2	0.3	535	7.7	0.7	201	2.0	0.3
2004	625	8.8	0.7	218	2.2	0.3	567	8.0	0.7	179	1.8	0.3
2005	668	9.3	0.7	230	2.4	0.3	623	8.8	0.7	183	1.8	0.3
2006	723	10.0	0.7	210	2.2	0.3	546	7.6	0.6	185	1.9	0.3
2007	655	8.9	0.7	219	2.3	0.3	578	8.0	0.7	197	2.0	0.3
2008	717	9.7	0.7	230	2.4	0.3	586	7.9	0.6	163	1.7	0.3
2009	721	9.5	0.7	212	2.1	0.3	590	7.9	0.6	165	1.6	0.3
2010	642	8.4	0.7	217	2.1	0.3	585	7.7	0.6	161	1.6	0.2
2011	625	8.0	0.6	209	2.1	0.3	585	7.6	0.6	190	1.8	0.3
2012	630	7.9	0.6	170	1.7	0.3	550	7.1	0.6	161	1.5	0.2
2013	675	8.3	0.6	205	2.0	0.3	570	7.1	0.6	167	1.6	0.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Stomach

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	1038	17.2	1.1	775	9.0	0.6	793	13.5	1.0	638	7.4	0.6
1977	1043	17.7	1.1	758	8.8	0.6	797	13.8	1.0	600	6.9	0.6
1978	985	16.7	1.1	782	8.9	0.6	721	12.5	1.0	579	6.6	0.5
1979	1020	16.9	1.1	780	8.7	0.6	746	12.7	1.0	579	6.4	0.5
1980	1043	17.4	1.1	725	8.2	0.6	751	12.8	1.0	571	6.4	0.5
1981	1050	17.5	1.1	756	8.5	0.6	751	12.8	1.0	533	5.9	0.5
1982	994	16.4	1.1	733	8.0	0.6	704	12.0	0.9	493	5.4	0.5
1983	1012	16.8	1.1	773	8.4	0.6	690	11.8	0.9	508	5.5	0.5
1984	985	15.9	1.0	718	7.8	0.6	677	11.0	0.9	540	5.8	0.5
1985	1017	16.7	1.1	673	7.3	0.6	655	10.9	0.9	493	5.4	0.5
1986	965	15.8	1.0	664	7.2	0.6	645	10.7	0.9	461	4.9	0.5
1987	973	15.8	1.0	687	7.3	0.6	652	10.8	0.9	451	4.8	0.4
1988	892	14.5	1.0	689	7.4	0.6	578	9.5	0.8	486	5.2	0.5
1989	933	15.1	1.0	578	6.1	0.5	645	10.7	0.9	464	4.9	0.5
1990	1030	16.7	1.0	674	7.3	0.6	613	10.1	0.8	388	4.1	0.4
1991	964	15.7	1.0	625	6.6	0.5	615	10.2	0.8	390	4.1	0.4
1992	941	15.2	1.0	625	6.6	0.5	614	10.2	0.8	411	4.3	0.4
1993	972	15.7	1.0	663	7.0	0.5	587	9.8	0.8	411	4.3	0.4
1994	935	14.9	1.0	677	7.2	0.5	610	10.0	0.8	434	4.5	0.4
1995	898	14.2	0.9	628	6.6	0.5	559	9.2	0.8	444	4.6	0.4
1996	901	14.3	0.9	646	6.7	0.5	515	8.3	0.7	424	4.4	0.4
1997	898	14.0	0.9	610	6.4	0.5	490	7.7	0.7	398	4.2	0.4
1998	961	14.7	0.9	613	6.5	0.5	495	7.7	0.7	389	4.0	0.4
1999	915	14.0	0.9	571	5.9	0.5	487	7.5	0.7	351	3.6	0.4
2000	857	12.9	0.9	611	6.2	0.5	439	6.8	0.6	372	3.7	0.4
2001	885	13.1	0.9	565	5.9	0.5	434	6.5	0.6	355	3.6	0.4
2002	822	12.0	0.8	559	5.8	0.5	413	6.2	0.6	313	3.1	0.3
2003	877	12.7	0.8	595	6.2	0.5	401	5.9	0.6	323	3.3	0.4
2004	829	12.0	0.8	592	6.2	0.5	428	6.3	0.6	349	3.5	0.4
2005	801	11.4	0.8	543	5.6	0.5	404	5.9	0.6	311	3.1	0.4
2006	785	11.1	0.8	526	5.5	0.5	358	5.1	0.5	273	2.8	0.3
2007	841	11.7	0.8	543	5.7	0.5	347	4.8	0.5	259	2.7	0.3
2008	806	11.1	0.8	469	4.9	0.5	396	5.5	0.5	246	2.4	0.3
2009	751	10.3	0.7	481	4.9	0.5	353	4.9	0.5	240	2.4	0.3
2010	803	10.7	0.8	496	5.1	0.5	313	4.3	0.5	268	2.7	0.3
2011	796	10.6	0.7	513	5.3	0.5	311	4.2	0.5	244	2.4	0.3
2012	809	10.5	0.7	505	5.2	0.5	337	4.4	0.5	241	2.4	0.3
2013	855	10.9	0.7	539	5.5	0.5	353	4.5	0.5	228	2.2	0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Colorectal

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	4915	80.5	2.4	5157	58.9	1.6	2572	44.0	1.8	2680	30.8	1.2
1977	4967	81.6	2.4	5169	58.5	1.6	2475	42.6	1.8	2644	30.1	1.2
1978	4937	81.3	2.4	5159	58.0	1.6	2568	43.8	1.8	2597	29.4	1.1
1979	4941	81.0	2.4	5295	59.1	1.6	2452	42.0	1.7	2679	29.8	1.1
1980	5312	86.4	2.4	5304	58.7	1.6	2530	43.4	1.8	2590	28.7	1.1
1981	5304	87.6	2.5	5387	59.3	1.6	2405	41.0	1.7	2590	28.4	1.1
1982	5234	85.6	2.4	5436	59.5	1.6	2438	41.6	1.7	2498	27.3	1.1
1983	5305	86.4	2.4	5401	58.8	1.6	2373	40.4	1.7	2519	27.2	1.1
1984	5173	84.0	2.4	5508	60.0	1.6	2509	42.3	1.7	2535	27.4	1.1
1985	5490	88.6	2.4	5439	59.0	1.6	2357	39.4	1.7	2475	26.5	1.1
1986	5310	86.1	2.4	5417	58.6	1.6	2360	39.7	1.7	2419	25.9	1.0
1987	5089	82.3	2.3	5171	55.7	1.5	2344	39.3	1.7	2395	25.4	1.0
1988	4829	77.8	2.3	5080	54.5	1.5	2342	39.1	1.6	2303	24.6	1.0
1989	4792	77.2	2.3	4850	52.4	1.5	2059	34.4	1.5	2260	23.9	1.0
1990	4679	74.5	2.2	4968	53.1	1.5	2114	35.1	1.5	2124	22.4	1.0
1991	4782	76.8	2.2	4804	51.6	1.5	2030	33.8	1.5	2141	22.5	1.0
1992	4623	73.7	2.2	4728	50.6	1.5	1913	31.5	1.5	2051	21.4	0.9
1993	4565	71.9	2.1	4857	51.8	1.5	1954	32.0	1.5	2047	21.2	0.9
1994	4772	75.3	2.2	5080	54.2	1.5	2014	33.1	1.5	2179	22.5	1.0
1995	4638	73.0	2.1	4894	52.0	1.5	1977	32.3	1.5	2165	22.3	1.0
1996	4641	72.5	2.1	4675	49.8	1.4	1862	29.9	1.4	1932	19.8	0.9
1997	4753	73.3	2.1	5025	53.4	1.5	1858	29.6	1.4	1900	19.4	0.9
1998	4985	76.1	2.1	5185	55.2	1.5	1754	27.3	1.3	1936	19.7	0.9
1999	4970	75.1	2.1	5111	53.9	1.5	1734	26.9	1.3	1836	18.5	0.9
2000	5018	74.9	2.1	5093	53.2	1.5	1812	27.8	1.3	1821	18.0	0.8
2001	4780	70.8	2.0	4874	50.9	1.5	1651	25.1	1.2	1802	17.9	0.8
2002	4707	68.5	2.0	4797	49.8	1.4	1688	25.3	1.2	1768	17.5	0.8
2003	4475	64.6	1.9	4659	48.2	1.4	1578	23.3	1.2	1685	16.6	0.8
2004	4291	61.4	1.8	4583	47.4	1.4	1478	21.5	1.1	1595	15.5	0.8
2005	4209	59.8	1.8	4243	43.8	1.3	1368	20.0	1.1	1475	14.2	0.7
2006	4024	56.9	1.8	4093	42.4	1.3	1378	20.0	1.1	1461	14.3	0.8
2007	4049	56.2	1.7	4138	43.0	1.3	1332	18.9	1.0	1408	13.6	0.7
2008	3924	53.9	1.7	3989	41.2	1.3	1343	18.9	1.0	1410	13.6	0.7
2009	3714	50.3	1.6	3812	39.1	1.3	1279	17.8	1.0	1367	13.2	0.7
2010	3685	49.2	1.6	3791	38.8	1.3	1324	18.0	1.0	1328	12.6	0.7
2011	3541	46.9	1.6	3612	37.0	1.2	1283	17.0	0.9	1301	12.1	0.7
2012	3517	45.7	1.5	3317	33.9	1.2	1180	15.5	0.9	1293	12.2	0.7
2013	3558	45.5	1.5	3392	34.5	1.2	1258	16.3	0.9	1190	11.0	0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Colon excluding rectum

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	3273	54.4	2.0	3772	43.0	1.4	1938	33.3	1.6	2199	25.3	1.1
1977	3336	56.0	2.0	3783	42.9	1.4	1973	34.0	1.6	2204	25.1	1.1
1978	3355	56.0	2.0	3798	42.8	1.4	2018	34.6	1.6	2176	24.6	1.0
1979	3379	56.4	2.0	3932	43.9	1.4	1971	33.7	1.6	2282	25.4	1.1
1980	3642	60.3	2.0	3887	43.0	1.4	2069	35.6	1.6	2187	24.2	1.0
1981	3632	60.6	2.1	3988	43.9	1.4	2010	34.4	1.6	2233	24.6	1.0
1982	3500	58.2	2.0	4028	44.0	1.4	2030	34.7	1.6	2122	23.2	1.0
1983	3573	59.1	2.0	4024	43.8	1.4	1959	33.4	1.5	2149	23.2	1.0
1984	3546	58.4	2.0	4148	45.0	1.4	2087	35.2	1.6	2182	23.5	1.0
1985	3802	62.2	2.1	4122	44.5	1.4	1981	33.0	1.5	2147	23.0	1.0
1986	3681	60.4	2.0	4133	44.6	1.4	1983	33.4	1.5	2103	22.5	1.0
1987	3502	57.3	2.0	3853	41.2	1.3	2010	33.7	1.5	2044	21.7	1.0
1988	3295	53.6	1.9	3846	41.1	1.3	2008	33.5	1.5	2006	21.4	0.9
1989	3297	54.0	1.9	3582	38.5	1.3	1742	29.2	1.4	1988	21.1	0.9
1990	3225	52.1	1.9	3717	39.6	1.3	1842	30.7	1.4	1862	19.6	0.9
1991	3356	54.4	1.9	3633	38.7	1.3	1708	28.5	1.4	1841	19.3	0.9
1992	3181	51.3	1.8	3556	37.9	1.3	1597	26.4	1.3	1792	18.7	0.9
1993	3176	50.4	1.8	3697	39.2	1.3	1669	27.5	1.4	1784	18.5	0.9
1994	3389	54.0	1.9	3822	40.5	1.3	1724	28.4	1.4	1919	19.8	0.9
1995	3273	51.9	1.8	3692	38.8	1.3	1659	27.3	1.3	1878	19.3	0.9
1996	3193	50.4	1.8	3531	37.2	1.2	1587	25.6	1.3	1665	17.0	0.8
1997	3369	52.4	1.8	3877	40.8	1.3	1580	25.1	1.3	1687	17.2	0.8
1998	3509	54.0	1.8	3900	41.2	1.3	1491	23.3	1.2	1688	17.1	0.8
1999	3459	52.7	1.8	3848	40.1	1.3	1448	22.5	1.2	1574	15.8	0.8
2000	3477	52.3	1.8	3894	40.2	1.3	1529	23.6	1.2	1577	15.5	0.8
2001	3362	50.1	1.7	3681	37.9	1.2	1373	21.0	1.1	1558	15.4	0.8
2002	3282	48.2	1.7	3634	37.4	1.2	1401	21.1	1.1	1493	14.7	0.8
2003	3117	45.4	1.6	3518	36.0	1.2	1324	19.7	1.1	1467	14.4	0.7
2004	2962	42.7	1.5	3431	35.0	1.2	1215	17.8	1.0	1371	13.2	0.7
2005	2955	42.2	1.5	3241	33.1	1.2	1119	16.4	1.0	1257	12.1	0.7
2006	2820	40.2	1.5	3096	31.6	1.1	1144	16.7	1.0	1257	12.2	0.7
2007	2828	39.5	1.5	3114	31.9	1.1	1110	15.8	0.9	1217	11.7	0.7
2008	2713	37.4	1.4	3012	30.8	1.1	1088	15.3	0.9	1174	11.3	0.7
2009	2503	34.1	1.4	2886	29.1	1.1	1044	14.6	0.9	1156	11.1	0.7
2010	2517	33.9	1.3	2837	28.5	1.1	1065	14.5	0.9	1137	10.7	0.6
2011	2422	32.4	1.3	2689	27.1	1.1	1030	13.6	0.8	1102	10.2	0.6
2012	2392	31.3	1.3	2531	25.4	1.0	942	12.5	0.8	1102	10.3	0.6
2013	2399	30.9	1.3	2499	25.0	1.0	1004	13.1	0.8	1005	9.2	0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Rectum & rectosigmoid

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	1642	26.2	1.3	1385	15.8	0.8	634	10.7	0.9	481	5.5	0.5
1977	1631	25.6	1.3	1386	15.6	0.8	502	8.6	0.8	440	5.0	0.5
1978	1582	25.3	1.3	1361	15.2	0.8	550	9.2	0.8	421	4.8	0.5
1979	1562	24.6	1.3	1363	15.1	0.8	481	8.3	0.8	397	4.4	0.4
1980	1670	26.1	1.3	1417	15.7	0.8	461	7.8	0.7	403	4.5	0.4
1981	1672	27.0	1.3	1399	15.4	0.8	395	6.6	0.7	357	3.9	0.4
1982	1734	27.4	1.3	1408	15.5	0.8	408	6.9	0.7	376	4.1	0.4
1983	1732	27.2	1.3	1377	15.0	0.8	414	7.0	0.7	370	4.0	0.4
1984	1627	25.5	1.3	1360	15.0	0.8	422	7.1	0.7	353	3.8	0.4
1985	1688	26.5	1.3	1317	14.4	0.8	376	6.3	0.7	328	3.5	0.4
1986	1629	25.8	1.3	1284	14.0	0.8	377	6.3	0.7	316	3.4	0.4
1987	1587	25.0	1.3	1318	14.5	0.8	334	5.6	0.6	351	3.7	0.4
1988	1534	24.2	1.2	1234	13.5	0.8	334	5.6	0.6	297	3.2	0.4
1989	1495	23.2	1.2	1268	13.9	0.8	317	5.2	0.6	272	2.9	0.3
1990	1454	22.4	1.2	1251	13.5	0.8	272	4.4	0.5	262	2.8	0.3
1991	1426	22.4	1.2	1171	12.9	0.8	322	5.3	0.6	300	3.2	0.4
1992	1442	22.4	1.2	1172	12.7	0.7	316	5.2	0.6	259	2.7	0.3
1993	1389	21.4	1.1	1160	12.6	0.7	285	4.5	0.5	263	2.7	0.3
1994	1383	21.3	1.1	1258	13.8	0.8	290	4.7	0.6	260	2.7	0.3
1995	1365	21.1	1.1	1202	13.2	0.8	318	5.1	0.6	287	3.0	0.4
1996	1448	22.2	1.2	1144	12.6	0.7	275	4.3	0.5	267	2.8	0.3
1997	1384	20.9	1.1	1148	12.5	0.7	278	4.4	0.5	213	2.2	0.3
1998	1476	22.0	1.1	1285	14.0	0.8	263	4.0	0.5	248	2.6	0.3
1999	1511	22.5	1.1	1263	13.8	0.8	286	4.4	0.5	262	2.7	0.3
2000	1541	22.6	1.1	1199	12.9	0.7	283	4.2	0.5	244	2.5	0.3
2001	1418	20.7	1.1	1193	13.0	0.7	278	4.2	0.5	244	2.5	0.3
2002	1425	20.4	1.1	1163	12.4	0.7	287	4.2	0.5	275	2.8	0.3
2003	1358	19.3	1.0	1141	12.2	0.7	254	3.7	0.5	218	2.2	0.3
2004	1329	18.7	1.0	1152	12.4	0.7	263	3.8	0.5	224	2.2	0.3
2005	1254	17.6	1.0	1002	10.7	0.7	249	3.6	0.4	218	2.1	0.3
2006	1204	16.7	0.9	997	10.8	0.7	234	3.4	0.4	204	2.1	0.3
2007	1221	16.6	0.9	1024	11.1	0.7	222	3.1	0.4	191	1.9	0.3
2008	1211	16.4	0.9	977	10.4	0.7	255	3.5	0.4	236	2.3	0.3
2009	1211	16.2	0.9	926	10.0	0.7	235	3.2	0.4	211	2.1	0.3
2010	1168	15.3	0.9	954	10.3	0.7	259	3.5	0.4	191	1.9	0.3
2011	1119	14.6	0.9	923	9.9	0.7	253	3.3	0.4	199	1.9	0.3
2012	1125	14.4	0.9	786	8.5	0.6	238	3.1	0.4	191	1.9	0.3
2013	1159	14.6	0.9	893	9.5	0.6	254	3.2	0.4	185	1.8	0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Liver / intrahepatic bile duct

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	260	4.2	0.5	198	2.2	0.3	229	3.6	0.5	179	2.1	0.3
1977	317	4.9	0.6	177	2.0	0.3	275	4.2	0.5	218	2.5	0.3
1978	261	4.1	0.5	194	2.2	0.3	274	4.3	0.5	193	2.2	0.3
1979	248	3.9	0.5	195	2.3	0.3	241	3.9	0.5	209	2.4	0.3
1980	261	4.3	0.5	190	2.1	0.3	222	3.6	0.5	210	2.3	0.3
1981	258	4.2	0.5	202	2.2	0.3	273	4.5	0.6	224	2.5	0.3
1982	276	4.5	0.5	153	1.7	0.3	280	4.6	0.6	181	2.0	0.3
1983	300	4.7	0.6	172	2.0	0.3	285	4.5	0.5	197	2.2	0.3
1984	269	4.2	0.5	192	2.1	0.3	266	4.3	0.5	211	2.3	0.3
1985	287	4.7	0.6	180	2.0	0.3	302	5.0	0.6	200	2.2	0.3
1986	312	5.0	0.6	185	2.1	0.3	293	4.7	0.6	226	2.5	0.3
1987	294	4.6	0.5	180	2.0	0.3	312	5.2	0.6	227	2.5	0.3
1988	351	5.6	0.6	206	2.3	0.3	329	5.3	0.6	226	2.5	0.3
1989	314	4.9	0.6	188	2.0	0.3	286	4.5	0.5	198	2.1	0.3
1990	369	5.9	0.6	220	2.4	0.3	346	5.7	0.6	238	2.5	0.3
1991	369	5.6	0.6	231	2.5	0.3	347	5.4	0.6	238	2.5	0.3
1992	385	6.1	0.6	269	3.0	0.4	370	5.9	0.6	250	2.7	0.3
1993	402	6.3	0.6	252	2.7	0.3	379	5.9	0.6	287	3.0	0.4
1994	417	6.5	0.6	234	2.5	0.3	362	5.8	0.6	242	2.6	0.3
1995	469	7.2	0.7	265	2.9	0.3	406	6.3	0.6	261	2.7	0.3
1996	467	7.2	0.7	256	2.8	0.3	391	6.0	0.6	254	2.6	0.3
1997	521	7.9	0.7	284	3.0	0.4	388	5.9	0.6	245	2.5	0.3
1998	589	8.8	0.7	309	3.3	0.4	402	6.0	0.6	258	2.7	0.3
1999	566	8.3	0.7	295	3.1	0.4	420	6.3	0.6	269	2.8	0.3
2000	567	8.2	0.7	290	3.1	0.4	422	6.2	0.6	262	2.7	0.3
2001	628	9.1	0.7	280	3.0	0.4	455	6.7	0.6	255	2.6	0.3
2002	629	8.9	0.7	270	2.9	0.4	498	7.2	0.6	274	2.8	0.3
2003	636	8.9	0.7	338	3.7	0.4	447	6.4	0.6	266	2.7	0.3
2004	723	10.0	0.7	289	3.1	0.4	481	6.7	0.6	298	3.0	0.3
2005	758	10.3	0.7	315	3.4	0.4	517	7.1	0.6	281	2.8	0.3
2006	784	10.6	0.7	333	3.6	0.4	520	7.2	0.6	322	3.3	0.4
2007	849	11.2	0.8	329	3.5	0.4	550	7.5	0.6	275	2.8	0.3
2008	873	11.4	0.8	351	3.7	0.4	579	7.7	0.6	305	3.1	0.4
2009	850	10.9	0.7	365	3.8	0.4	594	7.9	0.6	315	3.2	0.4
2010	906	11.5	0.8	306	3.2	0.4	598	7.7	0.6	318	3.2	0.4
2011	876	10.7	0.7	371	3.9	0.4	590	7.4	0.6	345	3.4	0.4
2012	973	11.7	0.7	373	3.9	0.4	640	8.0	0.6	357	3.5	0.4
2013	970	11.6	0.7	422	4.2	0.4	689	8.4	0.6	335	3.2	0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Pancreas

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	977	15.7	1.0	896	10.2	0.7	887	14.4	1.0	873	9.9	0.7
1977	977	16.3	1.1	949	10.7	0.7	928	15.3	1.0	922	10.4	0.7
1978	967	15.4	1.0	932	10.4	0.7	918	14.7	1.0	883	9.9	0.7
1979	1009	16.3	1.0	961	10.6	0.7	966	15.8	1.0	926	10.2	0.7
1980	966	15.6	1.0	1003	11.0	0.7	923	15.0	1.0	909	10.0	0.7
1981	948	15.4	1.0	938	10.4	0.7	883	14.4	1.0	920	10.1	0.7
1982	899	14.9	1.0	1016	11.0	0.7	837	13.7	1.0	864	9.4	0.6
1983	883	14.1	1.0	939	10.3	0.7	868	13.7	0.9	915	9.9	0.7
1984	901	14.8	1.0	993	10.7	0.7	826	13.5	1.0	932	10.0	0.7
1985	902	14.3	1.0	1041	11.2	0.7	811	13.2	0.9	907	9.8	0.6
1986	891	14.4	1.0	1003	10.8	0.7	822	13.1	0.9	923	9.8	0.6
1987	871	14.1	1.0	1023	10.8	0.7	817	13.2	0.9	883	9.4	0.6
1988	891	14.4	1.0	1033	11.0	0.7	825	13.4	0.9	913	9.6	0.6
1989	875	14.0	1.0	965	10.4	0.7	813	13.1	0.9	867	9.3	0.6
1990	890	14.3	1.0	998	10.7	0.7	794	12.8	0.9	901	9.6	0.6
1991	911	14.5	1.0	1039	11.1	0.7	789	12.7	0.9	895	9.5	0.6
1992	909	14.6	1.0	1012	10.8	0.7	766	12.3	0.9	914	9.6	0.6
1993	904	14.2	0.9	1017	10.8	0.7	860	13.6	0.9	938	9.9	0.6
1994	978	15.3	1.0	1070	11.2	0.7	868	13.7	0.9	988	10.3	0.6
1995	869	13.5	0.9	1080	11.4	0.7	821	12.7	0.9	969	10.0	0.6
1996	945	14.8	1.0	985	10.5	0.7	851	13.5	0.9	964	10.1	0.6
1997	941	14.5	0.9	1045	11.0	0.7	871	13.7	0.9	940	9.8	0.6
1998	1001	15.3	1.0	1089	11.5	0.7	861	13.3	0.9	936	9.8	0.6
1999	981	14.8	0.9	1072	11.2	0.7	927	14.0	0.9	940	9.8	0.6
2000	982	14.7	0.9	1129	11.8	0.7	831	12.5	0.9	991	10.1	0.6
2001	935	13.8	0.9	1137	11.7	0.7	849	12.7	0.9	1010	10.2	0.6
2002	990	14.4	0.9	1123	11.5	0.7	850	12.4	0.8	979	9.9	0.6
2003	1046	15.1	0.9	1154	11.8	0.7	855	12.4	0.8	974	9.8	0.6
2004	1024	14.6	0.9	1150	11.8	0.7	906	13.0	0.9	957	9.7	0.6
2005	1071	15.1	0.9	1108	11.3	0.7	900	12.8	0.8	999	10.0	0.6
2006	1057	14.9	0.9	1177	12.0	0.7	914	12.9	0.8	986	9.8	0.6
2007	1133	15.7	0.9	1190	12.0	0.7	922	12.8	0.8	986	9.9	0.6
2008	1191	16.3	0.9	1257	12.6	0.7	938	12.9	0.8	1049	10.2	0.6
2009	1252	16.8	0.9	1194	12.0	0.7	1015	13.7	0.9	996	9.8	0.6
2010	1191	15.9	0.9	1270	12.7	0.7	987	13.0	0.8	1052	10.1	0.6
2011	1256	16.4	0.9	1229	12.1	0.7	986	13.0	0.8	1029	9.9	0.6
2012	1269	16.5	0.9	1231	12.1	0.7	1066	13.8	0.8	1006	9.7	0.6
2013	1258	15.9	0.9	1263	12.2	0.7	1028	13.1	0.8	1030	9.8	0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Larynx

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	718	10.5	0.8	137	1.5	0.3	253	3.9	0.5	59	0.7	0.2
1977	701	10.4	0.8	142	1.6	0.3	266	4.0	0.5	43	0.5	0.1
1978	706	10.6	0.8	154	1.7	0.3	247	3.8	0.5	65	0.7	0.2
1979	735	10.7	0.8	186	2.1	0.3	276	4.2	0.5	64	0.7	0.2
1980	773	11.6	0.8	170	2.0	0.3	252	3.9	0.5	54	0.6	0.2
1981	784	11.7	0.9	185	2.1	0.3	248	3.8	0.5	50	0.5	0.1
1982	715	10.8	0.8	179	2.1	0.3	249	3.8	0.5	63	0.7	0.2
1983	695	10.3	0.8	187	2.2	0.3	222	3.4	0.5	70	0.8	0.2
1984	730	10.8	0.8	200	2.3	0.3	211	3.2	0.5	57	0.6	0.2
1985	732	10.8	0.8	203	2.4	0.3	232	3.5	0.5	64	0.7	0.2
1986	683	10.3	0.8	200	2.4	0.3	197	3.1	0.4	51	0.5	0.2
1987	719	10.9	0.8	185	2.2	0.3	216	3.4	0.5	59	0.6	0.2
1988	660	9.8	0.8	196	2.3	0.3	211	3.3	0.5	56	0.6	0.2
1989	687	10.5	0.8	206	2.5	0.3	186	2.8	0.4	81	0.9	0.2
1990	713	10.7	0.8	211	2.5	0.3	196	3.1	0.4	41	0.4	0.1
1991	674	10.2	0.8	165	1.9	0.3	198	3.0	0.4	58	0.6	0.2
1992	649	9.8	0.8	206	2.5	0.3	239	3.8	0.5	75	0.8	0.2
1993	604	9.2	0.7	208	2.4	0.3	212	3.3	0.5	55	0.6	0.2
1994	597	9.1	0.7	181	2.1	0.3	203	3.2	0.4	61	0.7	0.2
1995	580	8.6	0.7	178	2.1	0.3	195	3.0	0.4	60	0.7	0.2
1996	598	9.0	0.7	168	2.0	0.3	205	3.1	0.4	59	0.7	0.2
1997	586	8.8	0.7	152	1.8	0.3	161	2.5	0.4	55	0.6	0.2
1998	520	7.7	0.7	187	2.2	0.3	178	2.7	0.4	56	0.6	0.2
1999	585	8.5	0.7	147	1.7	0.3	176	2.6	0.4	56	0.6	0.2
2000	568	8.3	0.7	164	1.9	0.3	181	2.7	0.4	44	0.5	0.1
2001	535	7.7	0.7	161	1.8	0.3	163	2.4	0.4	36	0.4	0.1
2002	530	7.6	0.6	138	1.6	0.3	176	2.6	0.4	47	0.5	0.1
2003	527	7.4	0.6	150	1.7	0.3	156	2.2	0.4	51	0.5	0.1
2004	550	7.5	0.6	128	1.4	0.3	159	2.3	0.4	38	0.4	0.1
2005	557	7.6	0.6	159	1.8	0.3	155	2.2	0.3	54	0.6	0.2
2006	521	7.2	0.6	155	1.8	0.3	161	2.3	0.4	46	0.5	0.1
2007	505	6.8	0.6	136	1.5	0.2	154	2.1	0.3	35	0.4	0.1
2008	566	7.5	0.6	156	1.7	0.3	140	1.9	0.3	33	0.3	0.1
2009	506	6.6	0.6	149	1.6	0.3	145	1.9	0.3	51	0.5	0.1
2010	502	6.5	0.6	130	1.4	0.2	153	2.0	0.3	51	0.5	0.1
2011	476	5.9	0.5	112	1.2	0.2	144	1.9	0.3	48	0.5	0.1
2012	518	6.5	0.6	132	1.4	0.2	135	1.7	0.3	39	0.4	0.1
2013	514	6.2	0.6	124	1.3	0.2	170	2.1	0.3	39	0.4	0.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Lung and bronchus

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	6059	92.7	2.4	2433	27.5	1.1	4988	77.5	2.3	1865	21.1	1.0
1977	5962	91.6	2.4	2527	28.7	1.1	5157	80.1	2.3	1972	22.3	1.0
1978	6147	94.7	2.5	2491	28.2	1.1	5103	79.7	2.3	2022	22.8	1.0
1979	6060	93.8	2.5	2778	31.6	1.2	4958	77.5	2.2	2112	23.7	1.0
1980	6285	96.6	2.5	2855	32.1	1.2	5268	82.0	2.3	2289	25.6	1.1
1981	6072	94.3	2.5	3000	33.9	1.2	5159	80.7	2.3	2264	25.4	1.1
1982	6228	96.6	2.5	3240	36.5	1.3	5096	80.1	2.3	2526	28.3	1.1
1983	6323	97.7	2.5	3442	38.5	1.3	5066	79.7	2.3	2697	29.9	1.2
1984	6275	96.8	2.5	3437	39.0	1.3	5095	79.6	2.3	2656	29.6	1.2
1985	6359	97.9	2.5	3674	41.4	1.4	5199	81.4	2.3	2901	32.2	1.2
1986	6202	96.0	2.5	3726	42.0	1.4	5081	79.4	2.3	2970	33.2	1.2
1987	6269	96.6	2.5	3892	43.9	1.4	5135	80.5	2.3	2963	32.9	1.2
1988	6202	96.4	2.5	4059	45.6	1.4	5146	80.6	2.3	3100	34.6	1.2
1989	6162	95.1	2.4	4215	47.7	1.5	5046	79.0	2.2	3193	35.7	1.3
1990	6211	96.1	2.5	4211	47.4	1.5	5022	78.4	2.2	3345	37.1	1.3
1991	6108	94.6	2.4	4462	50.4	1.5	5088	79.7	2.2	3506	39.0	1.3
1992	6174	95.0	2.4	4557	51.4	1.5	4988	77.9	2.2	3579	39.6	1.3
1993	5939	91.3	2.4	4657	52.5	1.5	4832	75.3	2.2	3587	39.8	1.3
1994	6015	92.1	2.4	4843	54.8	1.6	5075	78.8	2.2	3587	39.6	1.3
1995	5975	90.9	2.3	4757	53.3	1.5	4904	75.7	2.2	3694	40.8	1.3
1996	6047	92.2	2.4	4840	54.6	1.6	4782	73.6	2.1	3654	40.4	1.3
1997	5946	90.1	2.3	4980	56.0	1.6	4587	70.3	2.1	3611	39.7	1.3
1998	5905	88.1	2.3	5115	57.1	1.6	4582	69.0	2.0	3666	40.1	1.3
1999	5832	86.6	2.2	5150	57.1	1.6	4450	66.8	2.0	3724	40.6	1.3
2000	5938	87.4	2.2	5320	58.4	1.6	4513	67.1	2.0	3780	40.4	1.3
2001	5669	82.9	2.2	5346	58.5	1.6	4447	65.9	1.9	3757	40.2	1.3
2002	5894	85.2	2.2	5327	57.9	1.6	4411	64.3	1.9	3869	41.2	1.3
2003	5884	84.3	2.2	5299	57.4	1.6	4301	62.1	1.9	3766	39.8	1.3
2004	5718	81.2	2.1	5450	58.6	1.6	4160	59.6	1.8	3807	40.1	1.3
2005	5753	81.4	2.1	5469	58.9	1.6	4174	59.7	1.8	3706	39.0	1.3
2006	5676	79.9	2.1	5773	62.6	1.6	4183	59.3	1.8	3667	38.7	1.3
2007	5819	80.9	2.1	5677	60.9	1.6	4102	57.3	1.8	3828	40.1	1.3
2008	5833	80.2	2.1	5846	62.3	1.6	4082	56.4	1.7	3784	39.2	1.3
2009	5863	79.4	2.1	5697	59.9	1.6	3924	53.6	1.7	3641	37.6	1.2
2010	5598	74.6	2.0	5686	59.9	1.6	3873	52.1	1.7	3722	38.2	1.3
2011	5696	74.7	2.0	5788	59.8	1.6	3829	50.7	1.6	3643	36.6	1.2
2012	5543	71.7	1.9	5728	58.8	1.6	3993	52.1	1.6	3628	36.6	1.2
2013	5509	69.9	1.9	5802	58.8	1.5	3783	48.2	1.6	3624	35.8	1.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Melanoma of the skin

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	469	7.2	0.7	447	5.6	0.5	211	3.3	0.5	168	2.1	0.3
1977	543	8.2	0.7	528	6.6	0.6	231	3.6	0.5	157	1.9	0.3
1978	500	7.7	0.7	503	6.5	0.6	208	3.3	0.5	124	1.5	0.3
1979	554	8.5	0.7	505	6.5	0.6	208	3.3	0.5	168	2.1	0.3
1980	659	10.2	0.8	573	7.3	0.6	222	3.4	0.5	165	2.0	0.3
1981	603	9.3	0.8	568	7.2	0.6	219	3.4	0.5	160	2.0	0.3
1982	611	9.5	0.8	601	7.6	0.6	230	3.6	0.5	152	1.8	0.3
1983	719	11.2	0.8	625	8.1	0.7	213	3.4	0.5	164	2.1	0.3
1984	683	10.4	0.8	601	7.6	0.6	238	3.7	0.5	174	2.1	0.3
1985	728	11.2	0.8	604	7.7	0.6	239	3.8	0.5	178	2.1	0.3
1986	736	11.4	0.8	670	8.6	0.7	230	3.6	0.5	182	2.2	0.3
1987	725	11.2	0.8	612	7.7	0.6	262	4.1	0.5	176	2.2	0.3
1988	717	11.2	0.8	554	6.9	0.6	260	4.1	0.5	156	1.8	0.3
1989	728	11.3	0.8	603	7.5	0.6	266	4.3	0.5	151	1.8	0.3
1990	731	11.1	0.8	623	7.8	0.6	246	3.8	0.5	166	2.0	0.3
1991	743	11.3	0.8	559	6.8	0.6	245	3.9	0.5	165	1.9	0.3
1992	715	10.9	0.8	556	6.7	0.6	230	3.5	0.5	159	1.8	0.3
1993	735	11.1	0.8	553	6.6	0.6	232	3.7	0.5	142	1.6	0.3
1994	836	12.6	0.9	604	7.3	0.6	257	4.0	0.5	148	1.7	0.3
1995	909	13.6	0.9	619	7.6	0.6	248	3.8	0.5	178	2.0	0.3
1996	923	13.9	0.9	725	8.8	0.7	254	3.8	0.5	177	2.1	0.3
1997	1011	15.2	0.9	749	9.1	0.7	264	4.0	0.5	148	1.7	0.3
1998	1035	15.3	0.9	816	10.0	0.7	259	3.9	0.5	159	1.8	0.3
1999	1079	15.8	0.9	819	9.9	0.7	231	3.4	0.4	159	1.8	0.3
2000	1207	17.6	1.0	945	11.4	0.7	268	3.9	0.5	160	1.8	0.3
2001	1294	18.6	1.0	1020	12.3	0.8	289	4.3	0.5	179	2.0	0.3
2002	1310	18.8	1.0	1000	11.9	0.8	218	3.1	0.4	194	2.2	0.3
2003	1305	18.5	1.0	990	11.7	0.7	241	3.5	0.4	178	1.9	0.3
2004	1597	22.5	1.1	1243	14.9	0.8	250	3.5	0.4	171	1.9	0.3
2005	1769	24.9	1.2	1334	16.0	0.9	258	3.7	0.5	173	1.9	0.3
2006	1969	27.5	1.2	1476	17.7	0.9	288	4.1	0.5	162	1.8	0.3
2007	1798	24.9	1.2	1386	16.6	0.9	298	4.2	0.5	135	1.4	0.2
2008	1949	26.9	1.2	1463	17.4	0.9	285	3.9	0.5	160	1.7	0.3
2009	2112	28.6	1.2	1548	18.3	0.9	269	3.7	0.4	178	1.9	0.3
2010	1933	26.0	1.2	1497	17.5	0.9	271	3.6	0.4	177	1.9	0.3
2011	2194	28.9	1.2	1633	18.7	0.9	284	3.8	0.4	166	1.7	0.3
2012	2052	26.6	1.2	1593	18.3	0.9	319	4.2	0.5	199	2.0	0.3
2013	2248	28.7	1.2	1681	19.3	1.0	316	4.1	0.5	164	1.6	0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Female breast

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	8527	103.5	2.3	3310	38.8	1.4
1977	8391	102.0	2.2	3351	39.5	1.4
1978	8543	103.9	2.3	3359	39.4	1.4
1979	8410	102.4	2.3	3196	37.8	1.3
1980	8528	103.8	2.3	3300	38.5	1.3
1981	8739	106.6	2.3	3267	38.2	1.3
1982	8744	106.2	2.3	3212	37.4	1.3
1983	9115	110.7	2.4	3178	37.0	1.3
1984	9489	114.7	2.4	3312	38.5	1.4
1985	9811	119.3	2.4	3315	38.6	1.4
1986	9874	119.7	2.4	3297	38.1	1.3
1987	10180	123.0	2.5	3265	37.5	1.3
1988	10313	124.7	2.5	3327	38.4	1.3
1989	10128	123.1	2.5	3255	37.9	1.3
1990	10746	130.9	2.5	3156	36.4	1.3
1991	10697	129.8	2.5	3040	35.1	1.3
1992	10706	130.0	2.5	3020	34.7	1.3
1993	10520	127.3	2.5	3013	34.2	1.2
1994	10946	132.2	2.5	3007	34.2	1.2
1995	11229	136.0	2.6	2963	33.8	1.2
1996	11866	143.4	2.6	3013	34.0	1.2
1997	11843	142.1	2.6	2750	31.0	1.2
1998	12170	145.1	2.6	2706	30.1	1.2
1999	12078	142.7	2.6	2558	28.1	1.1
2000	12034	140.7	2.5	2571	27.9	1.1
2001	11752	136.9	2.5	2536	27.5	1.1
2002	11681	135.3	2.5	2446	26.1	1.1
2003	11542	133.0	2.5	2420	25.8	1.0
2004	11382	130.8	2.4	2298	24.3	1.0
2005	11425	130.7	2.4	2226	23.5	1.0
2006	11590	133.5	2.5	2155	22.8	1.0
2007	11743	134.4	2.5	2167	22.7	1.0
2008	12044	137.2	2.5	2094	21.8	1.0
2009	12318	138.6	2.5	2090	21.6	1.0
2010	11561	130.1	2.4	2134	21.7	1.0
2011	12073	134.8	2.5	2050	20.6	0.9
2012	11845	131.9	2.5	2001	20.1	0.9
2013	12219	135.4	2.5	1996	20.0	0.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Cervix uteri

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	981	12.5	0.8	363	4.3	0.5
1977	858	10.8	0.7	319	3.8	0.4
1978	861	11.1	0.8	349	4.2	0.5
1979	863	11.3	0.8	318	3.9	0.4
1980	868	11.2	0.8	314	3.9	0.4
1981	766	9.7	0.7	330	4.0	0.4
1982	773	10.2	0.7	300	3.6	0.4
1983	800	10.4	0.7	257	3.1	0.4
1984	820	10.8	0.8	241	3.0	0.4
1985	780	10.2	0.7	242	3.0	0.4
1986	703	9.2	0.7	248	3.0	0.4
1987	821	10.6	0.7	251	3.1	0.4
1988	810	10.6	0.7	247	3.1	0.4
1989	770	10.1	0.7	215	2.6	0.4
1990	832	10.9	0.8	257	3.1	0.4
1991	827	10.8	0.8	210	2.6	0.4
1992	782	10.0	0.7	225	2.8	0.4
1993	790	10.2	0.7	209	2.6	0.4
1994	810	10.4	0.7	252	3.1	0.4
1995	898	11.5	0.8	204	2.5	0.4
1996	879	11.3	0.8	243	3.0	0.4
1997	796	10.1	0.7	240	2.9	0.4
1998	729	9.3	0.7	197	2.4	0.3
1999	720	9.1	0.7	238	2.9	0.4
2000	663	8.4	0.6	202	2.4	0.3
2001	651	8.2	0.6	193	2.2	0.3
2002	613	7.7	0.6	203	2.4	0.3
2003	645	8.2	0.6	191	2.2	0.3
2004	570	7.2	0.6	191	2.2	0.3
2005	643	8.2	0.6	197	2.2	0.3
2006	596	7.6	0.6	189	2.2	0.3
2007	604	7.8	0.6	184	2.2	0.3
2008	602	7.7	0.6	160	1.8	0.3
2009	622	7.9	0.6	150	1.7	0.3
2010	563	7.3	0.6	177	2.0	0.3
2011	501	6.4	0.6	178	2.1	0.3
2012	550	6.9	0.6	201	2.4	0.3
2013	554	7.2	0.6	174	1.9	0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Corpus uterus and uterus, not otherwise specified

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	2362	27.2	1.1	533	6.0	0.5
1977	2131	24.4	1.1	484	5.4	0.5
1978	1910	21.7	1.0	440	4.9	0.5
1979	2103	24.2	1.1	459	5.0	0.5
1980	2138	24.9	1.1	488	5.3	0.5
1981	2198	25.3	1.1	454	5.0	0.5
1982	2134	24.5	1.1	464	5.1	0.5
1983	2089	24.1	1.1	445	4.7	0.4
1984	2078	24.0	1.1	454	4.9	0.5
1985	2029	23.3	1.0	411	4.5	0.4
1986	2016	23.5	1.1	412	4.4	0.4
1987	2104	24.5	1.1	424	4.6	0.4
1988	1983	23.1	1.0	400	4.3	0.4
1989	1966	22.9	1.0	406	4.4	0.4
1990	1996	23.6	1.1	369	3.9	0.4
1991	2097	25.3	1.1	415	4.5	0.4
1992	2158	25.7	1.1	418	4.5	0.4
1993	2064	24.5	1.1	439	4.7	0.4
1994	2241	26.4	1.1	435	4.7	0.4
1995	2246	26.8	1.1	453	4.8	0.5
1996	2339	27.7	1.1	417	4.5	0.4
1997	2334	27.7	1.1	421	4.5	0.4
1998	2526	30.0	1.2	417	4.5	0.4
1999	2459	28.7	1.1	401	4.2	0.4
2000	2398	27.8	1.1	409	4.3	0.4
2001	2609	30.3	1.2	394	4.1	0.4
2002	2642	30.2	1.2	422	4.5	0.4
2003	2627	29.9	1.2	444	4.6	0.4
2004	2682	30.4	1.2	418	4.4	0.4
2005	2720	30.6	1.2	435	4.6	0.4
2006	2683	29.9	1.1	461	4.8	0.4
2007	2715	30.3	1.2	450	4.6	0.4
2008	2994	33.1	1.2	455	4.7	0.4
2009	2803	30.7	1.2	429	4.4	0.4
2010	2816	30.7	1.2	480	4.8	0.4
2011	2960	31.7	1.2	500	5.1	0.5
2012	2980	31.4	1.2	495	4.9	0.4
2013	3181	33.4	1.2	534	5.2	0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Ovary

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	1435	16.9	0.9	1033	11.9	0.7
1977	1353	16.1	0.9	949	10.9	0.7
1978	1366	16.2	0.9	931	10.6	0.7
1979	1403	16.8	0.9	911	10.4	0.7
1980	1395	16.5	0.9	937	10.7	0.7
1981	1492	17.9	0.9	960	10.8	0.7
1982	1434	17.2	0.9	933	10.6	0.7
1983	1490	17.8	0.9	932	10.5	0.7
1984	1422	17.0	0.9	929	10.5	0.7
1985	1476	17.9	0.9	884	10.1	0.7
1986	1421	17.2	0.9	883	10.1	0.7
1987	1447	17.4	0.9	915	10.4	0.7
1988	1427	17.2	0.9	960	10.8	0.7
1989	1425	17.1	0.9	892	10.1	0.7
1990	1484	17.9	0.9	915	10.3	0.7
1991	1528	18.4	0.9	967	11.0	0.7
1992	1475	17.9	0.9	940	10.8	0.7
1993	1455	17.4	0.9	889	9.9	0.7
1994	1503	17.9	0.9	892	10.1	0.7
1995	1557	18.7	0.9	917	10.3	0.7
1996	1523	18.2	0.9	880	9.9	0.7
1997	1481	17.7	0.9	939	10.5	0.7
1998	1362	16.2	0.9	821	9.2	0.6
1999	1399	16.4	0.9	839	9.3	0.6
2000	1413	16.3	0.9	898	9.8	0.6
2001	1448	16.7	0.9	875	9.5	0.6
2002	1407	16.1	0.9	867	9.4	0.6
2003	1366	15.6	0.8	896	9.6	0.6
2004	1352	15.3	0.8	874	9.3	0.6
2005	1263	14.3	0.8	897	9.6	0.6
2006	1307	14.7	0.8	873	9.3	0.6
2007	1299	14.6	0.8	804	8.5	0.6
2008	1286	14.3	0.8	847	8.8	0.6
2009	1229	13.7	0.8	790	8.1	0.6
2010	1243	13.7	0.8	780	8.1	0.6
2011	1256	13.7	0.8	801	8.0	0.6
2012	1155	12.6	0.7	777	7.8	0.6
2013	1239	13.5	0.8	815	8.1	0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Prostate

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	4110	74.0	2.4	1501	29.3	1.6
1977	4184	74.1	2.4	1500	28.8	1.5
1978	4129	73.2	2.3	1566	30.4	1.6
1979	4344	77.4	2.4	1527	29.7	1.6
1980	4787	83.3	2.5	1548	29.9	1.6
1981	4767	83.1	2.5	1508	29.2	1.5
1982	4864	84.3	2.5	1530	29.2	1.5
1983	5081	86.7	2.5	1680	31.7	1.6
1984	5057	86.0	2.5	1746	33.0	1.6
1985	5323	89.3	2.5	1623	30.1	1.5
1986	5415	91.1	2.5	1695	31.0	1.5
1987	5756	95.7	2.6	1687	30.9	1.5
1988	5761	96.4	2.6	1767	32.3	1.6
1989	5977	98.9	2.6	1766	32.4	1.6
1990	7125	114.9	2.8	1802	32.9	1.6
1991	8444	133.5	2.9	1888	34.0	1.6
1992	10576	163.9	3.2	1898	34.1	1.6
1993	10105	154.8	3.1	1902	34.0	1.6
1994	9513	144.7	3.0	1965	34.9	1.6
1995	9284	139.9	2.9	1886	33.3	1.5
1996	10015	150.8	3.0	1852	32.1	1.5
1997	10552	157.9	3.0	1798	30.9	1.5
1998	10435	154.4	3.0	1680	28.3	1.4
1999	10918	160.7	3.0	1597	26.5	1.3
2000	12019	175.7	3.2	1632	26.9	1.3
2001	11856	171.4	3.1	1611	26.1	1.3
2002	12170	173.8	3.1	1531	24.6	1.2
2003	11254	159.4	3.0	1535	24.1	1.2
2004	10950	153.1	2.9	1462	22.8	1.2
2005	11110	153.6	2.9	1364	20.9	1.1
2006	12514	172.0	3.0	1324	20.1	1.1
2007	12736	171.4	3.0	1384	20.5	1.1
2008	12246	161.6	2.9	1317	19.2	1.0
2009	12034	155.3	2.8	1265	18.2	1.0
2010	11253	142.5	2.7	1371	19.3	1.0
2011	11672	145.1	2.7	1304	18.0	1.0
2012	9916	120.4	2.4	1221	16.7	0.9
2013	9466	112.9	2.3	1165	15.6	0.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Testis

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	248	3.3	0.4	48	0.6	0.2
1977	241	3.3	0.4	44	0.6	0.2
1978	263	3.7	0.5	49	0.7	0.2
1979	301	4.3	0.5	47	0.7	0.2
1980	325	4.5	0.5	44	0.6	0.2
1981	317	4.3	0.5	32	0.4	0.1
1982	321	4.4	0.5	32	0.5	0.2
1983	335	4.5	0.5	30	0.4	0.2
1984	379	5.2	0.5	27	0.4	0.2
1985	351	4.7	0.5	34	0.5	0.2
1986	354	4.7	0.5	34	0.5	0.2
1987	380	5.0	0.5	27	0.4	0.1
1988	349	4.6	0.5	23	0.3	0.1
1989	364	4.8	0.5	31	0.4	0.1
1990	406	5.4	0.5	24	0.3	0.1
1991	389	5.2	0.5	22	0.3	0.1
1992	397	5.4	0.5	28	0.4	0.1
1993	419	5.7	0.5	32	0.5	0.2
1994	409	5.6	0.5	21	0.3	0.1
1995	416	5.7	0.6	22	0.3	0.1
1996	410	5.6	0.5	26	0.4	0.1
1997	426	5.9	0.6	12	0.2	0.1
1998	428	6.0	0.6	25	0.3	0.1
1999	470	6.6	0.6	17	0.2	0.1
2000	422	5.9	0.6	15	0.2	0.1
2001	434	6.1	0.6	19	0.3	0.1
2002	426	6.1	0.6	25	0.4	0.1
2003	477	6.9	0.6	19	0.3	0.1
2004	462	6.7	0.6	19	0.3	0.1
2005	488	7.1	0.6	14	0.2	0.1
2006	461	6.8	0.6	23	0.3	0.1
2007	498	7.3	0.6	7	0.1	0.1
2008	517	7.6	0.7	15	0.2	0.1
2009	502	7.4	0.7	15	0.2	0.1
2010	503	7.4	0.7	28	0.4	0.1
2011	501	7.3	0.7	16	0.2	0.1
2012	528	7.8	0.7	18	0.3	0.1
2013	470	6.9	0.6	13	0.2	0.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Urinary bladder (incl. in situ)

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	2241	37.0	1.6	831	9.4	0.6	673	12.4	1.0	277	3.3	0.4
1977	2329	38.0	1.6	896	10.3	0.7	677	12.3	1.0	290	3.4	0.4
1978	2443	40.4	1.7	927	10.6	0.7	653	11.9	1.0	295	3.4	0.4
1979	2492	41.4	1.7	910	10.2	0.7	631	11.4	0.9	294	3.3	0.4
1980	2650	43.4	1.7	930	10.3	0.7	641	11.4	0.9	272	3.0	0.4
1981	2582	42.3	1.7	918	10.3	0.7	638	11.5	0.9	278	3.1	0.4
1982	2577	41.8	1.7	985	10.9	0.7	623	11.1	0.9	258	2.8	0.3
1983	2645	43.5	1.7	975	10.6	0.7	598	10.9	0.9	246	2.6	0.3
1984	2543	40.9	1.7	971	10.6	0.7	575	10.4	0.9	277	3.0	0.4
1985	2546	41.2	1.7	971	10.6	0.7	585	10.5	0.9	269	2.9	0.3
1986	2675	43.0	1.7	1006	11.1	0.7	567	10.4	0.9	267	2.8	0.3
1987	2552	40.7	1.6	1006	10.9	0.7	528	9.4	0.8	245	2.6	0.3
1988	2528	40.4	1.6	994	10.8	0.7	543	9.4	0.8	286	2.9	0.3
1989	2541	40.9	1.6	966	10.5	0.7	563	10.0	0.9	255	2.6	0.3
1990	2584	41.4	1.6	1059	11.5	0.7	520	9.2	0.8	279	2.9	0.3
1991	2626	41.8	1.6	1013	11.0	0.7	525	9.1	0.8	240	2.5	0.3
1992	2598	41.4	1.6	1038	11.4	0.7	490	8.6	0.8	309	3.1	0.4
1993	2610	41.0	1.6	1090	11.8	0.7	561	9.7	0.8	269	2.7	0.3
1994	2607	41.0	1.6	1065	11.6	0.7	558	9.5	0.8	272	2.7	0.3
1995	2699	42.3	1.6	1114	12.1	0.7	548	9.2	0.8	287	2.9	0.3
1996	2835	44.3	1.7	1059	11.5	0.7	547	9.2	0.8	293	2.9	0.3
1997	2998	46.3	1.7	1173	12.6	0.7	572	9.4	0.8	285	2.9	0.3
1998	3112	47.4	1.7	1198	13.0	0.7	569	9.3	0.8	263	2.6	0.3
1999	3046	46.0	1.6	1195	12.8	0.7	573	9.3	0.8	257	2.5	0.3
2000	3092	46.4	1.6	1235	13.0	0.7	569	8.9	0.7	260	2.5	0.3
2001	3105	46.2	1.6	1161	12.3	0.7	597	9.4	0.8	250	2.4	0.3
2002	3243	47.7	1.7	1137	12.0	0.7	543	8.4	0.7	274	2.7	0.3
2003	3309	48.0	1.6	1199	12.6	0.7	512	7.9	0.7	243	2.3	0.3
2004	3397	49.0	1.7	1243	13.0	0.7	576	8.6	0.7	254	2.4	0.3
2005	3404	48.7	1.6	1265	13.1	0.7	526	7.8	0.7	264	2.5	0.3
2006	3371	48.0	1.6	1174	12.3	0.7	550	8.1	0.7	266	2.5	0.3
2007	3380	47.6	1.6	1212	12.7	0.7	619	8.9	0.7	263	2.5	0.3
2008	3552	49.2	1.6	1156	11.9	0.7	605	8.7	0.7	271	2.5	0.3
2009	3502	47.9	1.6	1206	12.2	0.7	628	8.9	0.7	228	2.0	0.3
2010	3351	45.3	1.6	1072	10.9	0.7	637	8.7	0.7	270	2.4	0.3
2011	3477	46.2	1.6	1199	12.2	0.7	587	8.0	0.7	251	2.3	0.3
2012	3540	46.6	1.6	1254	12.4	0.7	602	8.1	0.7	273	2.4	0.3
2013	3524	45.7	1.5	1215	12.0	0.7	621	8.2	0.7	243	2.1	0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Kidney and renal pelvis

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	687	10.5	0.8	418	4.8	0.5	381	6.0	0.6	217	2.4	0.3
1977	727	11.1	0.8	432	5.0	0.5	393	6.2	0.6	240	2.7	0.4
1978	701	10.9	0.8	445	5.1	0.5	361	5.6	0.6	214	2.4	0.3
1979	703	10.9	0.8	441	4.9	0.5	398	6.3	0.6	238	2.7	0.4
1980	696	10.6	0.8	442	5.1	0.5	357	5.7	0.6	241	2.6	0.3
1981	713	11.3	0.9	500	5.8	0.5	358	5.6	0.6	235	2.6	0.3
1982	781	12.0	0.9	516	5.9	0.5	358	5.7	0.6	241	2.6	0.3
1983	832	12.8	0.9	520	6.0	0.5	350	5.5	0.6	238	2.7	0.3
1984	858	13.3	0.9	511	5.8	0.5	371	6.0	0.6	228	2.5	0.3
1985	885	13.7	0.9	570	6.5	0.5	350	5.6	0.6	264	3.0	0.4
1986	889	13.7	0.9	556	6.4	0.5	395	6.2	0.6	259	2.9	0.4
1987	933	14.3	0.9	639	7.3	0.6	404	6.5	0.7	248	2.7	0.3
1988	877	13.6	0.9	572	6.6	0.6	359	5.7	0.6	263	2.9	0.4
1989	908	14.2	0.9	557	6.3	0.5	376	6.1	0.6	250	2.8	0.3
1990	988	15.3	1.0	639	7.4	0.6	355	5.6	0.6	247	2.7	0.3
1991	980	15.0	1.0	670	7.7	0.6	370	5.8	0.6	256	2.8	0.3
1992	999	15.3	1.0	669	7.8	0.6	343	5.6	0.6	234	2.5	0.3
1993	1056	16.0	1.0	661	7.6	0.6	366	5.6	0.6	213	2.3	0.3
1994	1056	16.1	1.0	699	8.1	0.6	410	6.5	0.6	252	2.7	0.3
1995	1036	15.6	1.0	706	8.1	0.6	363	5.7	0.6	265	2.9	0.4
1996	1158	17.5	1.0	750	8.6	0.6	373	5.8	0.6	276	3.0	0.4
1997	1217	18.4	1.0	758	8.7	0.6	392	6.1	0.6	241	2.6	0.3
1998	1271	18.7	1.0	808	9.3	0.6	385	5.8	0.6	249	2.6	0.3
1999	1337	19.6	1.1	828	9.4	0.7	343	5.2	0.6	205	2.1	0.3
2000	1300	18.9	1.0	893	10.0	0.7	370	5.6	0.6	235	2.5	0.3
2001	1415	20.3	1.1	858	9.6	0.7	374	5.5	0.6	264	2.8	0.3
2002	1433	20.4	1.1	865	9.6	0.7	363	5.3	0.6	242	2.5	0.3
2003	1525	21.3	1.1	890	9.9	0.7	361	5.2	0.5	243	2.4	0.3
2004	1618	22.6	1.1	982	11.1	0.7	383	5.5	0.6	252	2.6	0.3
2005	1594	22.2	1.1	1001	11.2	0.7	379	5.4	0.5	223	2.3	0.3
2006	1780	24.5	1.1	1022	11.4	0.7	381	5.5	0.6	251	2.6	0.3
2007	1833	25.0	1.2	1040	11.7	0.7	380	5.3	0.5	232	2.4	0.3
2008	1799	24.1	1.1	1051	11.6	0.7	391	5.4	0.5	215	2.1	0.3
2009	1690	22.5	1.1	1049	11.6	0.7	406	5.5	0.5	215	2.2	0.3
2010	1763	23.0	1.1	1046	11.5	0.7	378	5.0	0.5	215	2.1	0.3
2011	1855	23.9	1.1	1023	11.2	0.7	384	5.1	0.5	229	2.2	0.3
2012	1884	24.0	1.1	1069	11.6	0.7	417	5.4	0.5	214	2.1	0.3
2013	1877	23.5	1.1	1058	11.4	0.7	366	4.7	0.5	213	2.0	0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Brain and other nervous system

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	510	7.1	0.6	396	4.7	0.5	376	5.4	0.6	312	3.7	0.4
1977	514	7.3	0.7	404	4.8	0.5	376	5.4	0.6	316	3.7	0.4
1978	459	6.6	0.6	430	5.2	0.5	354	5.1	0.5	306	3.5	0.4
1979	531	7.7	0.7	421	5.0	0.5	363	5.3	0.6	283	3.3	0.4
1980	522	7.4	0.7	430	5.2	0.5	418	6.1	0.6	305	3.5	0.4
1981	535	8.0	0.7	423	5.0	0.5	368	5.5	0.6	303	3.5	0.4
1982	520	7.7	0.7	478	5.8	0.5	348	5.2	0.6	342	4.0	0.4
1983	481	7.1	0.7	466	5.7	0.5	352	5.3	0.6	305	3.4	0.4
1984	541	8.0	0.7	502	6.0	0.5	350	5.3	0.6	312	3.6	0.4
1985	547	8.2	0.7	469	5.7	0.5	325	5.0	0.6	327	3.8	0.4
1986	560	8.4	0.7	459	5.6	0.5	386	5.9	0.6	316	3.6	0.4
1987	570	8.5	0.7	528	6.3	0.6	351	5.3	0.6	337	3.9	0.4
1988	589	8.8	0.7	497	6.1	0.5	363	5.3	0.6	294	3.3	0.4
1989	608	9.1	0.7	474	5.9	0.5	352	5.3	0.6	292	3.4	0.4
1990	613	9.3	0.8	522	6.4	0.6	356	5.5	0.6	257	2.9	0.4
1991	575	8.7	0.7	503	6.2	0.6	367	5.7	0.6	288	3.5	0.4
1992	579	8.7	0.7	508	6.3	0.6	332	5.0	0.5	316	3.7	0.4
1993	587	8.7	0.7	508	6.2	0.6	343	5.2	0.6	284	3.3	0.4
1994	650	9.7	0.8	540	6.5	0.6	387	5.8	0.6	326	3.8	0.4
1995	648	9.6	0.7	620	7.6	0.6	359	5.4	0.6	307	3.6	0.4
1996	668	10.0	0.8	580	7.2	0.6	378	5.7	0.6	323	3.8	0.4
1997	620	9.1	0.7	545	6.7	0.6	362	5.4	0.6	337	3.9	0.4
1998	694	10.1	0.8	571	7.1	0.6	375	5.5	0.6	316	3.7	0.4
1999	598	8.7	0.7	533	6.5	0.6	380	5.5	0.6	316	3.6	0.4
2000	637	9.2	0.7	609	7.3	0.6	336	4.8	0.5	317	3.6	0.4
2001	614	8.8	0.7	537	6.4	0.6	337	4.8	0.5	297	3.4	0.4
2002	649	9.3	0.7	524	6.3	0.5	373	5.3	0.5	298	3.4	0.4
2003	603	8.6	0.7	571	7.0	0.6	351	4.9	0.5	268	3.1	0.4
2004	636	9.0	0.7	494	6.0	0.5	350	4.9	0.5	307	3.5	0.4
2005	682	9.7	0.7	505	6.1	0.5	385	5.4	0.5	313	3.6	0.4
2006	672	9.5	0.7	518	6.4	0.6	391	5.4	0.5	270	3.0	0.4
2007	601	8.5	0.7	510	6.3	0.6	403	5.5	0.5	287	3.2	0.4
2008	625	8.8	0.7	529	6.4	0.6	369	5.0	0.5	269	3.0	0.4
2009	627	8.7	0.7	548	6.6	0.6	404	5.4	0.5	329	3.7	0.4
2010	615	8.6	0.7	501	6.1	0.6	376	5.0	0.5	324	3.6	0.4
2011	614	8.4	0.7	488	6.0	0.5	361	4.8	0.5	303	3.2	0.4
2012	625	8.3	0.7	528	6.3	0.6	403	5.2	0.5	350	3.8	0.4
2013	639	8.5	0.7	477	5.7	0.5	380	4.8	0.5	321	3.3	0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Thyroid

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	155	2.4	0.4	335	4.3	0.5	41	0.7	0.2	61	0.7	0.2
1977	168	2.5	0.4	399	5.1	0.5	46	0.7	0.2	68	0.8	0.2
1978	157	2.4	0.4	400	5.1	0.5	31	0.5	0.2	64	0.7	0.2
1979	171	2.7	0.4	338	4.3	0.5	28	0.5	0.2	48	0.5	0.2
1980	162	2.5	0.4	376	4.7	0.5	48	0.8	0.2	61	0.7	0.2
1981	157	2.4	0.4	364	4.6	0.5	36	0.6	0.2	56	0.6	0.2
1982	157	2.4	0.4	350	4.6	0.5	41	0.6	0.2	56	0.6	0.2
1983	158	2.3	0.4	436	5.6	0.5	21	0.3	0.1	57	0.6	0.2
1984	186	2.9	0.4	462	5.9	0.6	37	0.6	0.2	64	0.7	0.2
1985	187	2.8	0.4	465	6.0	0.6	43	0.7	0.2	46	0.5	0.2
1986	176	2.6	0.4	454	5.8	0.6	28	0.5	0.2	53	0.6	0.2
1987	194	2.9	0.4	415	5.4	0.5	25	0.4	0.2	55	0.6	0.2
1988	173	2.6	0.4	432	5.6	0.5	29	0.5	0.2	49	0.5	0.1
1989	166	2.5	0.4	434	5.7	0.6	24	0.4	0.2	58	0.6	0.2
1990	184	2.8	0.4	454	5.9	0.6	29	0.4	0.2	48	0.5	0.1
1991	209	3.1	0.4	507	6.6	0.6	23	0.4	0.2	49	0.5	0.2
1992	198	3.0	0.4	533	6.8	0.6	26	0.4	0.2	55	0.6	0.2
1993	209	3.1	0.4	559	7.3	0.6	30	0.5	0.2	53	0.6	0.2
1994	220	3.3	0.4	611	8.0	0.6	35	0.6	0.2	46	0.5	0.1
1995	261	3.8	0.5	668	8.7	0.7	31	0.5	0.2	47	0.5	0.1
1996	257	3.7	0.5	729	9.7	0.7	38	0.6	0.2	74	0.8	0.2
1997	296	4.3	0.5	793	10.4	0.7	32	0.5	0.2	53	0.5	0.1
1998	318	4.6	0.5	800	10.4	0.7	25	0.4	0.1	53	0.5	0.1
1999	318	4.5	0.5	991	13.0	0.8	30	0.4	0.2	46	0.5	0.1
2000	365	5.2	0.5	1066	14.0	0.8	25	0.4	0.1	55	0.6	0.2
2001	356	5.0	0.5	1165	15.2	0.9	33	0.5	0.2	47	0.5	0.1
2002	426	5.9	0.6	1140	14.9	0.9	29	0.4	0.2	51	0.5	0.1
2003	418	5.8	0.6	1173	15.2	0.9	23	0.3	0.1	54	0.5	0.1
2004	433	6.0	0.6	1336	17.4	0.9	30	0.4	0.2	38	0.4	0.1
2005	539	7.5	0.6	1573	20.6	1.0	40	0.6	0.2	43	0.4	0.1
2006	561	7.7	0.6	1652	21.6	1.1	38	0.5	0.2	58	0.6	0.1
2007	621	8.5	0.7	1814	23.6	1.1	37	0.5	0.2	56	0.6	0.2
2008	652	8.9	0.7	2110	27.7	1.2	40	0.6	0.2	44	0.5	0.1
2009	764	10.4	0.7	2195	28.6	1.2	43	0.6	0.2	53	0.5	0.1
2010	744	10.0	0.7	2203	28.8	1.2	39	0.5	0.2	51	0.5	0.1
2011	768	10.3	0.7	2285	29.6	1.3	49	0.7	0.2	52	0.5	0.1
2012	831	11.0	0.8	2317	30.4	1.3	41	0.5	0.2	51	0.5	0.1
2013	897	11.9	0.8	2362	31.0	1.3	46	0.6	0.2	56	0.5	0.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Hodgkin lymphoma

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	306	4.1	0.5	256	3.0	0.4	126	1.8	0.3	91	1.1	0.2
1977	301	4.1	0.5	235	2.8	0.4	128	1.9	0.3	97	1.2	0.2
1978	297	4.1	0.5	271	3.2	0.4	109	1.6	0.3	88	1.1	0.2
1979	313	4.5	0.5	262	3.1	0.4	97	1.5	0.3	73	0.9	0.2
1980	327	4.6	0.5	271	3.2	0.4	112	1.7	0.3	91	1.1	0.2
1981	300	4.1	0.5	264	3.2	0.4	98	1.5	0.3	78	0.9	0.2
1982	298	4.2	0.5	280	3.3	0.4	97	1.4	0.3	85	1.0	0.2
1983	310	4.4	0.5	232	2.8	0.4	88	1.3	0.3	64	0.7	0.2
1984	301	4.3	0.5	265	3.2	0.4	93	1.4	0.3	64	0.7	0.2
1985	329	4.6	0.5	275	3.4	0.4	84	1.2	0.3	71	0.8	0.2
1986	305	4.2	0.5	258	3.2	0.4	99	1.5	0.3	75	0.9	0.2
1987	267	3.8	0.5	253	3.2	0.4	86	1.2	0.3	54	0.7	0.2
1988	313	4.4	0.5	250	3.1	0.4	76	1.1	0.3	43	0.5	0.2
1989	310	4.3	0.5	245	3.1	0.4	95	1.4	0.3	54	0.7	0.2
1990	286	4.1	0.5	237	3.0	0.4	84	1.3	0.3	53	0.6	0.2
1991	289	4.1	0.5	256	3.3	0.4	61	0.9	0.2	49	0.6	0.2
1992	303	4.3	0.5	257	3.3	0.4	64	0.9	0.2	64	0.8	0.2
1993	250	3.6	0.5	256	3.3	0.4	55	0.8	0.2	47	0.6	0.2
1994	296	4.2	0.5	269	3.5	0.4	55	0.8	0.2	58	0.7	0.2
1995	277	4.0	0.5	236	3.2	0.4	50	0.7	0.2	50	0.6	0.2
1996	249	3.6	0.4	243	3.3	0.4	45	0.7	0.2	44	0.6	0.2
1997	269	3.9	0.5	213	2.9	0.4	53	0.8	0.2	43	0.5	0.2
1998	272	3.9	0.5	265	3.6	0.4	48	0.7	0.2	45	0.6	0.2
1999	269	3.8	0.5	231	3.1	0.4	39	0.6	0.2	44	0.5	0.2
2000	291	4.2	0.5	245	3.3	0.4	44	0.7	0.2	42	0.5	0.1
2001	278	4.0	0.5	259	3.5	0.4	45	0.6	0.2	40	0.5	0.1
2002	324	4.6	0.5	254	3.4	0.4	48	0.7	0.2	45	0.5	0.2
2003	282	4.0	0.5	236	3.2	0.4	32	0.5	0.2	38	0.5	0.1
2004	278	4.0	0.5	262	3.6	0.4	44	0.6	0.2	31	0.4	0.1
2005	303	4.4	0.5	249	3.4	0.4	34	0.5	0.2	36	0.4	0.1
2006	294	4.2	0.5	234	3.1	0.4	41	0.6	0.2	30	0.4	0.1
2007	263	3.7	0.5	265	3.6	0.4	42	0.6	0.2	42	0.5	0.1
2008	272	3.8	0.5	243	3.3	0.4	50	0.7	0.2	34	0.4	0.1
2009	290	4.1	0.5	233	3.2	0.4	35	0.5	0.2	28	0.3	0.1
2010	278	3.9	0.5	224	3.0	0.4	38	0.5	0.2	28	0.3	0.1
2011	287	4.1	0.5	228	3.1	0.4	43	0.6	0.2	26	0.3	0.1
2012	294	4.2	0.5	210	2.9	0.4	31	0.4	0.1	36	0.4	0.1
2013	239	3.3	0.4	249	3.4	0.4	27	0.4	0.1	28	0.3	0.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Non-Hodgkin lymphomas

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	900	13.9	1.0	854	9.9	0.7	519	8.2	0.7	451	5.1	0.5
1977	909	13.9	0.9	874	10.0	0.7	475	7.4	0.7	497	5.6	0.5
1978	990	15.1	1.0	911	10.5	0.7	510	8.1	0.7	479	5.4	0.5
1979	913	14.4	1.0	948	10.9	0.7	521	8.3	0.7	473	5.3	0.5
1980	1016	15.6	1.0	1001	11.5	0.7	503	7.8	0.7	516	5.8	0.5
1981	987	15.4	1.0	1042	11.8	0.7	505	8.0	0.7	509	5.6	0.5
1982	1053	16.3	1.0	1072	12.2	0.8	528	8.6	0.8	529	5.9	0.5
1983	1116	17.4	1.1	1110	12.5	0.8	573	9.1	0.8	546	6.1	0.5
1984	1183	18.8	1.1	1101	12.4	0.7	568	9.2	0.8	499	5.4	0.5
1985	1298	20.3	1.1	1151	12.9	0.8	625	10.0	0.8	556	6.1	0.5
1986	1373	21.3	1.2	1202	13.6	0.8	644	10.4	0.8	532	5.8	0.5
1987	1365	21.3	1.2	1187	13.4	0.8	598	9.5	0.8	594	6.5	0.5
1988	1349	20.9	1.1	1240	14.3	0.8	616	9.9	0.8	623	6.8	0.5
1989	1471	22.7	1.2	1218	14.1	0.8	677	10.8	0.8	566	6.2	0.5
1990	1592	24.5	1.2	1309	14.8	0.8	648	10.4	0.8	635	6.9	0.5
1991	1629	25.4	1.3	1372	15.7	0.8	674	10.8	0.8	606	6.5	0.5
1992	1546	23.6	1.2	1332	15.2	0.8	676	10.8	0.8	659	7.1	0.6
1993	1624	24.7	1.2	1342	15.4	0.8	631	10.0	0.8	612	6.6	0.5
1994	1802	27.3	1.3	1448	16.5	0.9	728	11.4	0.8	691	7.4	0.6
1995	1714	26.0	1.2	1484	17.0	0.9	731	11.6	0.9	678	7.3	0.6
1996	1792	27.2	1.3	1553	17.8	0.9	741	11.7	0.9	685	7.4	0.6
1997	1736	26.1	1.2	1609	18.2	0.9	747	11.5	0.8	731	7.8	0.6
1998	1701	25.3	1.2	1614	18.3	0.9	662	10.0	0.8	720	7.6	0.6
1999	1683	24.9	1.2	1666	18.8	0.9	637	9.8	0.8	675	7.1	0.5
2000	1719	25.1	1.2	1645	18.3	0.9	666	10.1	0.8	688	7.2	0.5
2001	1734	25.3	1.2	1613	17.8	0.9	688	10.2	0.8	573	5.8	0.5
2002	1816	26.1	1.2	1579	17.5	0.9	654	9.7	0.7	569	5.9	0.5
2003	1840	26.2	1.2	1752	19.2	0.9	572	8.5	0.7	562	5.6	0.5
2004	1910	27.1	1.2	1708	18.9	0.9	561	8.2	0.7	583	5.8	0.5
2005	1954	27.5	1.2	1775	19.5	0.9	582	8.4	0.7	529	5.3	0.5
2006	2026	28.4	1.2	1813	19.8	0.9	617	8.8	0.7	590	5.8	0.5
2007	2011	27.9	1.2	1801	19.7	0.9	596	8.4	0.7	492	4.9	0.4
2008	2071	28.4	1.2	1823	19.9	0.9	601	8.4	0.7	529	5.3	0.5
2009	2089	28.4	1.2	1864	20.1	0.9	599	8.2	0.7	481	4.7	0.4
2010	2069	27.8	1.2	1846	19.6	0.9	564	7.7	0.6	518	5.0	0.4
2011	2070	27.4	1.2	1780	19.0	0.9	565	7.6	0.6	515	5.0	0.4
2012	2114	27.7	1.2	1787	18.9	0.9	575	7.7	0.6	467	4.4	0.4
2013	2140	27.7	1.2	1797	18.7	0.9	530	7.0	0.6	451	4.2	0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Myeloma

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	323	5.2	0.6	316	3.5	0.4	211	3.6	0.5	198	2.3	0.3
1977	316	5.1	0.6	313	3.4	0.4	203	3.2	0.5	207	2.3	0.3
1978	302	4.9	0.6	341	3.8	0.4	194	3.1	0.5	206	2.3	0.3
1979	326	5.3	0.6	358	4.0	0.4	202	3.3	0.5	229	2.5	0.3
1980	326	5.4	0.6	325	3.6	0.4	228	3.8	0.5	250	2.7	0.3
1981	358	5.8	0.6	394	4.4	0.4	218	3.7	0.5	239	2.6	0.3
1982	357	5.7	0.6	353	3.8	0.4	212	3.5	0.5	233	2.5	0.3
1983	378	6.3	0.7	333	3.6	0.4	223	3.7	0.5	235	2.5	0.3
1984	367	6.0	0.6	346	3.8	0.4	239	3.8	0.5	248	2.7	0.3
1985	378	6.1	0.6	357	3.9	0.4	257	4.2	0.5	231	2.4	0.3
1986	373	6.0	0.6	325	3.5	0.4	237	3.9	0.5	250	2.6	0.3
1987	386	6.3	0.6	366	3.9	0.4	257	4.2	0.5	237	2.6	0.3
1988	379	6.2	0.6	342	3.7	0.4	273	4.6	0.6	229	2.4	0.3
1989	349	5.7	0.6	418	4.6	0.5	236	3.8	0.5	234	2.5	0.3
1990	413	6.5	0.6	394	4.3	0.4	232	3.8	0.5	229	2.4	0.3
1991	381	6.1	0.6	368	4.0	0.4	231	3.7	0.5	248	2.6	0.3
1992	428	6.8	0.7	401	4.3	0.4	259	4.3	0.5	243	2.6	0.3
1993	405	6.4	0.6	413	4.5	0.4	279	4.5	0.5	283	2.9	0.3
1994	403	6.3	0.6	392	4.3	0.4	256	4.1	0.5	309	3.3	0.4
1995	432	6.8	0.7	404	4.4	0.4	282	4.5	0.5	271	2.9	0.3
1996	437	6.9	0.7	426	4.6	0.4	261	4.2	0.5	275	2.9	0.3
1997	408	6.3	0.6	426	4.7	0.5	249	4.0	0.5	259	2.8	0.3
1998	483	7.3	0.7	472	5.1	0.5	261	4.0	0.5	265	2.8	0.3
1999	480	7.3	0.7	480	5.2	0.5	276	4.2	0.5	267	2.7	0.3
2000	492	7.3	0.6	458	4.9	0.5	290	4.4	0.5	288	3.0	0.3
2001	526	7.8	0.7	411	4.3	0.4	261	3.9	0.5	224	2.3	0.3
2002	559	8.2	0.7	454	4.8	0.4	289	4.4	0.5	248	2.5	0.3
2003	542	7.9	0.7	444	4.7	0.4	274	4.1	0.5	232	2.3	0.3
2004	552	7.8	0.7	471	5.0	0.5	256	3.8	0.5	229	2.3	0.3
2005	542	7.6	0.6	471	4.9	0.5	264	3.8	0.5	249	2.5	0.3
2006	573	8.0	0.7	479	5.0	0.5	281	4.0	0.5	205	2.1	0.3
2007	589	8.2	0.7	499	5.3	0.5	261	3.7	0.4	236	2.3	0.3
2008	615	8.4	0.7	493	5.1	0.5	268	3.8	0.5	228	2.3	0.3
2009	641	8.7	0.7	546	5.5	0.5	269	3.7	0.4	228	2.2	0.3
2010	651	8.5	0.7	540	5.6	0.5	271	3.7	0.4	263	2.5	0.3
2011	707	9.3	0.7	523	5.3	0.5	250	3.3	0.4	231	2.2	0.3
2012	675	8.6	0.7	526	5.4	0.5	256	3.4	0.4	261	2.5	0.3
2013	686	8.6	0.7	562	5.7	0.5	266	3.4	0.4	225	2.1	0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Whites
Leukemias

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	1056	17.1	1.1	816	9.6	0.7	678	11.4	0.9	549	6.5	0.6
1977	966	15.5	1.0	789	9.4	0.7	674	11.0	0.9	569	6.8	0.6
1978	1030	16.8	1.1	811	9.7	0.7	636	10.4	0.9	567	6.7	0.6
1979	1047	16.6	1.0	863	10.1	0.7	659	10.9	0.9	587	7.0	0.6
1980	1070	17.3	1.1	934	11.0	0.7	698	11.4	0.9	558	6.4	0.5
1981	1019	16.5	1.1	804	9.4	0.7	689	11.4	0.9	559	6.6	0.6
1982	1062	17.3	1.1	832	9.7	0.7	728	12.2	0.9	567	6.6	0.6
1983	1003	16.1	1.0	827	9.7	0.7	638	10.7	0.9	581	6.6	0.5
1984	1053	17.1	1.1	913	10.5	0.7	605	9.9	0.8	606	6.9	0.6
1985	1114	17.9	1.1	868	10.0	0.7	673	11.1	0.9	533	6.0	0.5
1986	1043	16.7	1.1	892	10.4	0.7	653	10.6	0.8	541	6.1	0.5
1987	932	14.9	1.0	846	9.7	0.7	606	9.9	0.8	538	6.1	0.5
1988	1005	16.2	1.0	823	9.5	0.7	653	10.7	0.9	500	5.6	0.5
1989	1046	16.9	1.1	850	9.6	0.7	625	10.2	0.8	538	6.0	0.5
1990	1110	17.5	1.1	942	10.9	0.7	586	9.6	0.8	565	6.3	0.5
1991	1060	16.9	1.0	876	10.1	0.7	603	9.9	0.8	538	5.9	0.5
1992	1120	17.9	1.1	865	9.9	0.7	642	10.5	0.8	524	5.8	0.5
1993	1033	16.1	1.0	900	10.5	0.7	595	9.6	0.8	488	5.4	0.5
1994	1167	18.2	1.1	981	11.1	0.7	673	10.7	0.8	553	6.0	0.5
1995	1113	17.2	1.0	925	10.6	0.7	671	10.9	0.8	542	5.9	0.5
1996	1217	18.6	1.1	902	10.2	0.7	661	10.3	0.8	574	6.3	0.5
1997	1176	18.0	1.0	999	11.2	0.7	669	10.5	0.8	535	5.7	0.5
1998	1239	18.6	1.0	924	10.4	0.7	659	10.1	0.8	531	5.7	0.5
1999	1290	19.5	1.1	1083	12.1	0.7	685	10.5	0.8	575	6.1	0.5
2000	1341	20.1	1.1	1077	11.8	0.7	655	9.9	0.8	567	6.0	0.5
2001	1301	19.3	1.1	1097	12.1	0.7	648	9.8	0.8	566	5.9	0.5
2002	1283	18.8	1.0	967	10.7	0.7	661	9.9	0.8	547	5.7	0.5
2003	1368	19.9	1.1	1099	12.1	0.7	648	9.7	0.7	507	5.2	0.5
2004	1376	19.9	1.1	1115	12.3	0.7	691	10.2	0.8	569	5.9	0.5
2005	1409	20.1	1.1	1127	12.3	0.7	693	10.1	0.8	525	5.3	0.5
2006	1428	20.5	1.1	1074	11.8	0.7	671	9.7	0.7	559	5.7	0.5
2007	1484	20.9	1.1	1101	12.3	0.8	665	9.6	0.7	513	5.3	0.5
2008	1423	19.8	1.0	1076	12.0	0.7	683	9.7	0.7	536	5.4	0.5
2009	1564	21.5	1.1	1146	12.6	0.8	689	9.7	0.7	532	5.4	0.5
2010	1629	22.2	1.1	1201	13.0	0.8	651	9.0	0.7	503	5.1	0.5
2011	1679	22.6	1.1	1276	13.7	0.8	701	9.5	0.7	522	5.1	0.5
2012	1620	21.5	1.1	1183	12.9	0.8	695	9.5	0.7	501	4.8	0.4
2013	1668	21.9	1.1	1196	12.9	0.8	691	9.2	0.7	514	5.0	0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
All Invasive Malignant Tumors

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	11208	447.8	8.7	12457	359.0	6.6	6744	275.0	6.9	6507	183.0	4.6
1977	11131	454.0	8.8	12154	356.3	6.6	6612	273.5	6.9	6457	182.8	4.6
1978	11072	459.6	8.9	12084	358.7	6.7	6360	267.3	6.8	6130	175.8	4.6
1979	11224	472.9	9.1	12229	368.1	6.8	6199	264.6	6.8	6124	177.1	4.6
1980	11590	496.3	9.3	12328	375.5	7.0	6332	275.8	7.0	6272	181.0	4.6
1981	11119	482.1	9.3	12366	377.5	7.0	5935	261.1	6.9	6150	180.2	4.7
1982	11180	492.5	9.4	12475	385.1	7.1	5831	259.5	6.9	5927	175.1	4.6
1983	11323	499.6	9.5	12427	385.3	7.1	5680	256.1	6.9	5887	173.3	4.6
1984	11327	503.5	9.5	12444	389.8	7.2	5792	261.6	6.9	5870	174.5	4.6
1985	11637	519.5	9.7	12293	386.7	7.2	5653	255.7	6.8	5655	169.8	4.6
1986	11520	520.6	9.7	12317	391.4	7.2	5321	242.7	6.7	5680	170.6	4.6
1987	11250	511.7	9.7	12135	388.8	7.2	5389	249.3	6.8	5533	167.4	4.6
1988	10734	497.4	9.6	11990	389.2	7.3	5374	253.5	6.9	5558	170.9	4.6
1989	10752	504.9	9.7	11755	392.0	7.4	5071	242.9	6.8	5386	168.2	4.6
1990	11235	533.3	10.0	12228	415.4	7.7	4991	242.9	6.8	5345	169.7	4.7
1991	11716	563.2	10.4	12448	428.4	7.8	4890	240.8	6.9	5223	168.3	4.7
1992	12024	579.9	10.5	12233	425.8	7.8	4859	240.6	6.9	5055	163.3	4.6
1993	11581	560.5	10.3	11785	412.9	7.7	4819	240.0	6.9	4984	162.4	4.6
1994	12179	591.5	10.6	12939	454.1	8.0	5061	253.8	7.1	5242	172.8	4.8
1995	11340	554.0	10.3	12427	439.9	7.9	4829	243.4	6.9	5318	176.0	4.9
1996	11270	554.8	10.3	12462	446.2	8.0	4605	232.6	6.8	5259	176.0	4.9
1997	11268	555.2	10.3	12459	444.5	8.0	4478	226.6	6.7	4938	166.5	4.7
1998	11531	562.6	10.3	12564	450.3	8.0	4494	224.8	6.6	4865	163.4	4.7
1999	11313	547.2	10.1	12477	443.8	7.9	4409	218.7	6.5	4741	158.5	4.6
2000	11889	571.4	10.3	12417	440.1	7.9	4319	212.3	6.4	4766	156.7	4.5
2001	11567	553.3	10.1	12174	431.0	7.8	4286	209.6	6.3	4581	150.4	4.4
2002	11520	546.4	10.0	11930	421.3	7.7	4150	201.5	6.2	4452	146.2	4.4
2003	11303	537.3	9.9	11822	419.0	7.7	4037	195.5	6.1	4496	147.9	4.4
2004	11259	532.6	9.9	11932	425.4	7.8	3993	193.2	6.0	4292	142.1	4.3
2005	11460	542.1	10.0	11947	430.8	7.9	3962	192.0	6.0	4182	139.7	4.3
2006	11973	569.3	10.3	12057	439.4	8.0	3891	189.8	6.0	4172	140.7	4.4
2007	11990	567.7	10.2	12145	446.1	8.1	3920	189.8	6.0	4130	139.5	4.4
2008	11892	558.7	10.1	12290	450.9	8.2	3947	189.9	6.0	4035	136.1	4.3
2009	11828	551.5	10.0	12542	459.4	8.3	3823	183.7	5.9	4019	135.1	4.3
2010	11371	525.2	9.8	11815	432.9	8.0	3951	187.1	5.9	4081	135.9	4.3
2011	11459	521.6	9.7	12016	436.9	8.0	3795	177.4	5.7	3931	129.6	4.2
2012	10958	492.9	9.4	11657	424.4	7.9	3698	171.1	5.6	3899	129.7	4.2
2013	10945	486.8	9.3	12057	434.3	8.0	3671	168.0	5.5	3831	125.5	4.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Oral cavity and pharynx

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	435	17.3	1.7	206	5.6	0.8	222	8.7	1.2	90	2.5	0.5
1977	416	16.8	1.7	228	6.2	0.8	208	8.5	1.2	94	2.6	0.5
1978	397	16.1	1.6	207	6.0	0.9	146	5.8	1.0	77	2.1	0.5
1979	409	16.8	1.7	192	5.7	0.8	163	6.6	1.1	70	2.0	0.5
1980	384	16.2	1.7	213	6.6	0.9	166	7.1	1.1	78	2.3	0.5
1981	439	18.5	1.8	191	5.8	0.9	156	6.7	1.1	84	2.4	0.5
1982	439	19.0	1.8	205	6.5	0.9	161	6.8	1.1	49	1.6	0.5
1983	425	18.5	1.8	197	5.7	0.8	154	6.4	1.1	51	1.4	0.4
1984	414	18.3	1.8	201	6.2	0.9	153	6.8	1.1	54	1.7	0.5
1985	418	18.6	1.8	231	7.3	1.0	139	6.1	1.0	59	1.7	0.4
1986	383	17.4	1.8	189	5.8	0.9	124	5.5	1.0	58	1.7	0.5
1987	371	16.9	1.8	188	6.2	0.9	112	5.1	1.0	52	1.5	0.4
1988	340	15.6	1.7	182	5.7	0.9	121	5.8	1.0	71	2.3	0.6
1989	335	16.0	1.7	186	6.3	0.9	103	4.9	1.0	55	1.7	0.5
1990	370	17.9	1.9	188	6.3	0.9	92	4.5	0.9	57	1.7	0.5
1991	356	17.5	1.8	201	6.9	1.0	91	4.5	0.9	40	1.2	0.4
1992	402	19.7	2.0	191	6.7	1.0	90	4.5	0.9	52	1.6	0.4
1993	362	18.0	1.9	188	6.7	1.0	110	5.5	1.0	30	0.9	0.3
1994	333	16.3	1.8	176	6.2	0.9	96	4.8	1.0	37	1.3	0.4
1995	294	14.5	1.7	151	5.3	0.9	80	4.0	0.9	43	1.4	0.4
1996	339	16.7	1.8	168	6.0	0.9	73	3.5	0.8	57	2.0	0.5
1997	289	14.2	1.6	167	6.0	0.9	97	4.8	1.0	37	1.3	0.4
1998	296	14.3	1.6	177	6.3	0.9	75	3.7	0.8	40	1.3	0.4
1999	361	17.2	1.8	158	5.6	0.9	84	4.1	0.9	30	1.0	0.4
2000	276	13.0	1.5	143	5.1	0.9	69	3.3	0.8	37	1.3	0.4
2001	296	13.9	1.6	163	5.6	0.9	73	3.5	0.8	36	1.1	0.4
2002	300	14.1	1.6	168	6.0	0.9	70	3.3	0.8	38	1.2	0.4
2003	301	14.0	1.6	157	5.5	0.9	65	3.1	0.8	36	1.2	0.4
2004	291	13.6	1.6	170	6.0	0.9	81	3.8	0.8	52	1.7	0.5
2005	311	14.3	1.6	135	4.9	0.8	64	3.0	0.7	39	1.3	0.4
2006	316	14.6	1.6	170	5.9	0.9	66	3.2	0.8	27	0.9	0.3
2007	322	15.0	1.7	172	6.2	1.0	70	3.3	0.8	38	1.2	0.4
2008	315	14.7	1.6	179	6.6	1.0	68	3.2	0.8	27	0.9	0.4
2009	370	17.2	1.8	139	5.0	0.9	83	3.8	0.8	34	1.2	0.4
2010	306	13.8	1.6	175	6.4	1.0	82	3.8	0.8	37	1.3	0.4
2011	329	14.8	1.6	143	5.0	0.8	64	2.9	0.7	29	0.9	0.3
2012	317	13.8	1.5	165	5.8	0.9	74	3.3	0.8	38	1.3	0.4
2013	332	14.5	1.6	150	5.3	0.9	66	2.8	0.7	28	1.0	0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Esophagus

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	179	6.8	1.1	90	2.5	0.5	171	6.6	1.0	83	2.4	0.5
1977	167	6.5	1.0	93	2.6	0.5	151	6.1	1.0	94	2.6	0.5
1978	170	6.8	1.1	95	2.6	0.5	163	6.7	1.1	93	2.6	0.5
1979	185	7.6	1.1	76	2.1	0.5	154	6.2	1.0	81	2.3	0.5
1980	191	8.0	1.2	102	2.8	0.5	154	6.6	1.1	70	1.8	0.4
1981	160	6.8	1.1	91	2.7	0.6	167	7.0	1.1	80	2.2	0.5
1982	155	6.8	1.1	99	2.7	0.5	141	6.2	1.1	63	1.8	0.4
1983	167	7.2	1.1	75	2.2	0.5	140	6.2	1.1	67	1.8	0.5
1984	160	6.9	1.1	75	2.1	0.5	138	5.8	1.0	75	2.1	0.5
1985	160	7.0	1.1	79	2.4	0.6	140	6.1	1.0	52	1.6	0.4
1986	150	6.9	1.1	92	2.7	0.6	149	6.9	1.1	45	1.3	0.4
1987	130	5.9	1.0	86	2.7	0.6	112	5.0	1.0	70	2.1	0.5
1988	116	5.2	1.0	71	2.1	0.5	138	6.4	1.1	61	1.9	0.5
1989	158	7.4	1.2	85	2.5	0.6	121	5.6	1.0	68	2.0	0.5
1990	141	6.7	1.1	90	2.8	0.6	124	6.0	1.1	68	2.0	0.5
1991	144	7.0	1.2	77	2.4	0.5	113	5.4	1.0	59	1.8	0.5
1992	143	6.8	1.1	82	2.7	0.6	127	6.1	1.1	49	1.5	0.4
1993	139	6.8	1.1	59	1.9	0.5	120	5.8	1.1	43	1.4	0.4
1994	169	8.3	1.3	69	2.2	0.5	135	6.6	1.1	52	1.6	0.4
1995	130	6.4	1.1	91	3.1	0.7	127	6.3	1.1	54	1.7	0.5
1996	146	7.2	1.2	74	2.3	0.5	129	6.4	1.1	66	2.1	0.5
1997	142	7.1	1.2	78	2.6	0.6	120	6.0	1.1	54	1.7	0.5
1998	159	7.9	1.2	69	2.3	0.5	136	6.8	1.1	58	1.9	0.5
1999	161	7.8	1.2	50	1.6	0.4	139	6.8	1.1	43	1.4	0.4
2000	137	6.6	1.1	66	2.1	0.5	129	6.3	1.1	65	2.0	0.5
2001	144	6.9	1.1	66	2.1	0.5	126	6.1	1.1	36	1.2	0.4
2002	155	7.3	1.2	58	1.9	0.5	123	5.9	1.0	49	1.5	0.4
2003	150	7.1	1.1	74	2.4	0.6	114	5.4	1.0	55	1.7	0.5
2004	146	6.9	1.1	70	2.3	0.5	115	5.5	1.0	45	1.4	0.4
2005	143	6.8	1.1	74	2.5	0.6	152	7.2	1.2	50	1.5	0.4
2006	174	8.4	1.3	57	1.9	0.5	110	5.3	1.0	68	2.2	0.5
2007	143	6.7	1.1	68	2.2	0.5	144	6.9	1.1	46	1.6	0.5
2008	169	7.8	1.2	75	2.5	0.6	126	5.8	1.0	48	1.6	0.5
2009	145	6.7	1.1	52	1.7	0.5	135	6.4	1.1	53	1.7	0.5
2010	129	6.0	1.0	53	1.7	0.5	115	5.3	1.0	46	1.5	0.4
2011	141	6.5	1.1	43	1.5	0.5	108	5.0	1.0	44	1.4	0.4
2012	118	5.3	1.0	43	1.4	0.4	92	4.3	0.9	35	1.1	0.4
2013	148	6.6	1.1	58	1.9	0.5	113	5.0	0.9	35	1.1	0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Stomach

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	484	20.0	1.9	368	10.2	1.1	375	15.8	1.7	294	8.2	1.0
1977	480	20.6	1.9	366	10.6	1.1	354	15.6	1.7	294	8.3	1.0
1978	455	19.8	1.9	370	10.4	1.1	331	14.7	1.6	271	7.6	0.9
1979	448	19.1	1.8	389	11.0	1.1	326	14.3	1.6	271	7.6	0.9
1980	489	21.5	2.0	362	10.4	1.1	332	14.7	1.6	259	7.4	0.9
1981	440	19.7	1.9	352	10.0	1.1	338	15.3	1.7	245	6.9	0.9
1982	403	17.9	1.8	337	9.5	1.0	271	12.4	1.5	224	6.4	0.9
1983	423	19.3	1.9	390	11.2	1.1	278	13.1	1.6	233	6.6	0.9
1984	399	17.7	1.8	342	9.9	1.1	251	11.4	1.5	241	6.8	0.9
1985	400	18.4	1.9	308	9.0	1.0	277	12.6	1.5	221	6.5	0.9
1986	360	16.6	1.8	307	8.9	1.0	220	10.4	1.4	186	5.3	0.8
1987	392	18.0	1.8	299	8.5	1.0	248	11.5	1.5	179	5.0	0.8
1988	359	16.9	1.8	286	8.3	1.0	214	10.0	1.4	204	5.9	0.8
1989	338	16.1	1.7	279	8.4	1.0	247	11.8	1.5	186	5.6	0.8
1990	398	19.4	1.9	310	9.6	1.1	223	11.0	1.5	159	4.8	0.8
1991	403	20.1	2.0	292	8.9	1.0	230	11.5	1.5	179	5.4	0.8
1992	356	17.8	1.9	285	9.1	1.1	226	11.5	1.5	183	5.6	0.8
1993	386	19.5	2.0	287	9.1	1.1	225	11.5	1.5	175	5.5	0.8
1994	376	18.7	1.9	324	10.5	1.2	247	12.6	1.6	183	5.8	0.9
1995	348	17.5	1.8	276	8.7	1.1	229	11.8	1.5	193	6.1	0.9
1996	349	17.7	1.9	300	9.6	1.1	195	10.0	1.4	193	6.1	0.9
1997	334	16.9	1.8	250	8.1	1.0	198	10.1	1.4	169	5.6	0.9
1998	364	18.2	1.9	280	9.3	1.1	208	10.5	1.4	171	5.4	0.8
1999	355	17.7	1.8	247	8.0	1.0	187	9.4	1.4	157	5.0	0.8
2000	297	14.5	1.7	267	8.6	1.1	156	7.7	1.2	157	5.0	0.8
2001	333	16.2	1.7	256	8.5	1.1	163	8.0	1.2	153	5.0	0.8
2002	298	14.3	1.6	228	7.5	1.0	158	7.6	1.2	127	4.0	0.7
2003	312	14.9	1.7	258	8.5	1.1	158	7.7	1.2	141	4.6	0.8
2004	307	14.8	1.7	254	8.6	1.1	151	7.3	1.2	156	5.3	0.8
2005	307	14.9	1.7	239	8.0	1.0	164	8.0	1.2	132	4.2	0.7
2006	293	14.2	1.6	219	7.5	1.0	146	7.1	1.2	114	3.9	0.7
2007	301	14.6	1.7	249	8.5	1.1	129	6.3	1.1	110	3.6	0.7
2008	301	14.3	1.6	200	6.8	1.0	156	7.5	1.2	92	3.1	0.6
2009	268	12.8	1.5	217	7.5	1.0	137	6.7	1.1	107	3.5	0.7
2010	277	13.0	1.5	212	7.3	1.0	109	5.2	1.0	132	4.4	0.8
2011	242	11.4	1.5	188	6.5	1.0	117	5.5	1.0	118	3.9	0.7
2012	283	13.1	1.6	194	6.7	1.0	125	5.8	1.0	98	3.1	0.6
2013	280	12.6	1.5	211	7.0	1.0	139	6.3	1.1	88	2.8	0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Colorectal

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	2018	81.0	3.7	2156	59.1	2.6	1139	47.8	2.9	1107	30.5	1.8
1977	2030	84.0	3.8	2071	57.0	2.5	1037	44.4	2.9	1067	29.5	1.8
1978	1944	81.5	3.8	2016	55.5	2.5	1018	44.1	2.8	949	26.5	1.7
1979	1981	84.2	3.8	2067	57.5	2.5	942	41.1	2.7	1037	28.9	1.8
1980	2092	90.0	4.0	2103	59.1	2.6	1019	46.2	2.9	986	27.4	1.7
1981	1993	87.8	4.0	2101	59.1	2.6	862	39.1	2.7	969	27.0	1.7
1982	1976	87.9	4.0	2131	60.3	2.6	873	39.6	2.7	882	25.1	1.7
1983	1947	86.1	3.9	2048	58.0	2.6	847	38.9	2.7	920	26.0	1.7
1984	1856	83.2	3.9	2034	58.6	2.6	900	41.2	2.8	884	25.0	1.7
1985	1946	87.1	4.0	1938	55.4	2.5	816	36.9	2.6	842	23.9	1.7
1986	1967	89.5	4.1	1980	57.5	2.6	794	37.0	2.6	817	23.6	1.7
1987	1749	79.9	3.8	1783	51.6	2.5	766	35.9	2.6	819	23.3	1.6
1988	1633	76.1	3.8	1796	53.0	2.5	772	37.1	2.7	738	21.7	1.6
1989	1563	74.0	3.7	1646	49.8	2.5	653	31.9	2.5	758	22.2	1.6
1990	1504	71.2	3.7	1723	52.2	2.5	677	33.2	2.5	743	22.1	1.6
1991	1605	78.6	3.9	1602	49.3	2.5	647	32.6	2.5	662	20.0	1.6
1992	1452	71.4	3.7	1626	51.1	2.5	593	29.7	2.4	646	19.3	1.5
1993	1473	72.1	3.7	1591	50.3	2.5	598	30.0	2.4	615	18.8	1.5
1994	1570	77.6	3.9	1802	57.3	2.7	665	33.8	2.6	688	21.2	1.6
1995	1459	73.0	3.8	1668	53.6	2.6	645	33.0	2.6	708	22.1	1.7
1996	1458	73.0	3.8	1562	51.1	2.6	546	27.6	2.3	599	18.6	1.5
1997	1451	72.6	3.8	1696	55.2	2.7	610	31.0	2.5	632	19.9	1.6
1998	1541	76.7	3.8	1674	55.3	2.7	537	27.1	2.3	635	19.9	1.6
1999	1474	72.4	3.7	1591	52.5	2.6	518	25.8	2.2	574	18.2	1.5
2000	1541	75.0	3.8	1564	50.7	2.6	556	27.5	2.3	580	18.1	1.5
2001	1467	71.4	3.7	1511	49.6	2.6	513	25.3	2.2	529	16.4	1.4
2002	1414	67.8	3.5	1485	48.4	2.5	514	25.1	2.2	530	16.7	1.5
2003	1405	67.5	3.5	1415	46.4	2.5	495	24.1	2.1	570	17.8	1.5
2004	1323	63.4	3.4	1367	44.8	2.4	448	21.8	2.0	479	15.0	1.4
2005	1267	60.9	3.4	1321	43.7	2.4	406	19.9	1.9	463	14.4	1.3
2006	1256	60.9	3.4	1267	42.8	2.4	457	22.5	2.1	452	14.7	1.4
2007	1227	58.5	3.3	1241	42.3	2.4	388	18.9	1.9	430	13.7	1.3
2008	1188	56.4	3.2	1174	40.2	2.4	432	21.0	2.0	431	13.9	1.3
2009	1132	53.5	3.1	1205	41.6	2.4	383	18.5	1.9	442	14.0	1.3
2010	1102	51.6	3.1	1090	37.2	2.3	446	21.3	2.0	406	13.0	1.3
2011	1038	48.3	3.0	1022	35.2	2.2	410	19.1	1.9	399	12.6	1.3
2012	1018	46.4	2.9	954	32.5	2.1	341	15.9	1.7	397	12.7	1.3
2013	1082	49.0	3.0	1041	35.3	2.2	351	16.0	1.7	383	11.9	1.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Colon excluding rectum

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	1367	55.3	3.1	1569	42.9	2.2	881	37.4	2.6	901	24.8	1.7
1977	1396	58.4	3.2	1511	41.6	2.2	841	36.0	2.6	894	24.7	1.7
1978	1364	57.5	3.2	1473	40.7	2.1	832	36.2	2.6	799	22.2	1.6
1979	1383	59.4	3.3	1538	42.8	2.2	760	33.1	2.4	896	24.9	1.7
1980	1440	63.2	3.4	1528	42.7	2.2	857	39.1	2.7	857	23.9	1.6
1981	1358	60.3	3.3	1535	42.8	2.2	731	33.1	2.5	849	23.7	1.6
1982	1334	60.3	3.3	1577	44.4	2.2	761	34.6	2.5	766	21.8	1.6
1983	1341	59.7	3.3	1530	43.1	2.2	708	32.3	2.5	782	22.0	1.6
1984	1314	59.5	3.3	1499	42.8	2.2	766	35.1	2.6	768	21.8	1.6
1985	1384	62.6	3.4	1514	43.3	2.2	712	32.2	2.4	751	21.4	1.6
1986	1383	63.3	3.4	1534	44.4	2.3	678	31.4	2.4	719	20.8	1.6
1987	1228	56.4	3.2	1355	38.8	2.1	688	32.3	2.5	705	20.1	1.5
1988	1141	53.5	3.2	1360	39.7	2.2	678	32.6	2.5	641	18.8	1.5
1989	1100	52.5	3.2	1238	37.2	2.1	580	28.5	2.4	682	20.0	1.5
1990	1085	51.7	3.1	1321	39.9	2.2	590	28.9	2.4	667	19.8	1.5
1991	1157	56.8	3.3	1239	37.9	2.2	557	28.1	2.4	576	17.3	1.4
1992	1048	51.9	3.2	1211	37.8	2.2	509	25.3	2.2	568	16.9	1.4
1993	1061	52.0	3.2	1223	38.4	2.2	514	25.9	2.3	549	16.8	1.4
1994	1150	57.1	3.3	1359	42.6	2.3	592	30.1	2.4	614	19.0	1.5
1995	1056	53.3	3.2	1270	40.2	2.3	563	28.9	2.4	619	19.2	1.5
1996	1030	51.7	3.2	1196	38.4	2.2	486	24.7	2.2	521	16.0	1.4
1997	1066	53.7	3.2	1299	41.8	2.3	529	26.9	2.3	563	17.7	1.5
1998	1088	54.5	3.3	1259	41.1	2.3	478	24.2	2.2	570	17.8	1.5
1999	1062	52.5	3.2	1189	38.9	2.3	451	22.5	2.1	498	15.8	1.4
2000	1078	52.8	3.2	1202	38.7	2.2	490	24.3	2.2	510	15.9	1.4
2001	1062	51.9	3.1	1140	36.6	2.2	428	21.1	2.0	463	14.2	1.3
2002	996	48.1	3.0	1140	36.8	2.2	447	21.9	2.0	456	14.4	1.3
2003	991	47.9	3.0	1065	34.5	2.1	430	21.0	2.0	501	15.5	1.4
2004	946	45.6	2.9	1008	32.5	2.1	382	18.6	1.9	431	13.4	1.3
2005	898	43.2	2.8	1023	33.5	2.1	348	17.1	1.8	405	12.5	1.3
2006	857	41.9	2.8	943	31.5	2.1	394	19.4	1.9	405	13.1	1.3
2007	848	40.6	2.7	906	30.5	2.0	338	16.5	1.8	371	11.7	1.2
2008	831	39.7	2.7	866	29.2	2.0	360	17.6	1.8	362	11.7	1.2
2009	769	36.8	2.6	921	31.2	2.1	326	15.8	1.7	376	11.8	1.2
2010	757	35.5	2.6	801	26.8	1.9	374	17.9	1.8	362	11.5	1.2
2011	741	34.6	2.5	740	25.2	1.9	344	16.0	1.7	336	10.6	1.2
2012	703	32.3	2.4	725	24.3	1.8	282	13.2	1.6	336	10.6	1.2
2013	743	34.0	2.5	768	25.8	1.9	292	13.5	1.6	329	10.1	1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Rectum & rectosigmoid

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	651	25.7	2.1	587	16.3	1.4	258	10.4	1.3	206	5.7	0.8
1977	634	25.6	2.1	560	15.4	1.3	196	8.5	1.3	173	4.8	0.7
1978	580	24.0	2.0	543	14.7	1.3	186	7.9	1.2	150	4.3	0.7
1979	598	24.8	2.1	529	14.8	1.3	182	8.0	1.2	141	4.0	0.7
1980	652	26.9	2.1	575	16.4	1.4	162	7.1	1.1	129	3.5	0.6
1981	635	27.6	2.2	566	16.3	1.4	131	6.0	1.1	120	3.3	0.6
1982	642	27.6	2.2	554	15.9	1.4	112	5.0	1.0	116	3.3	0.6
1983	606	26.4	2.2	518	14.9	1.3	139	6.6	1.1	138	4.0	0.7
1984	542	23.7	2.0	535	15.7	1.4	134	6.1	1.1	116	3.3	0.6
1985	562	24.5	2.1	424	12.2	1.2	104	4.7	0.9	91	2.6	0.5
1986	584	26.2	2.2	446	13.1	1.3	116	5.6	1.0	98	2.8	0.6
1987	521	23.6	2.1	428	12.9	1.3	78	3.7	0.8	114	3.2	0.6
1988	492	22.7	2.0	436	13.3	1.3	94	4.5	0.9	97	2.9	0.6
1989	463	21.5	2.0	408	12.5	1.3	73	3.5	0.8	76	2.2	0.5
1990	419	19.5	1.9	402	12.3	1.2	87	4.2	0.9	76	2.3	0.5
1991	448	21.8	2.1	363	11.4	1.2	90	4.5	0.9	86	2.7	0.6
1992	404	19.6	1.9	415	13.3	1.3	84	4.3	0.9	78	2.4	0.5
1993	412	20.1	2.0	368	11.9	1.2	84	4.1	0.9	66	2.0	0.5
1994	420	20.5	2.0	443	14.7	1.4	73	3.8	0.9	74	2.2	0.5
1995	403	19.8	1.9	398	13.4	1.3	82	4.1	0.9	89	2.9	0.6
1996	428	21.3	2.0	366	12.7	1.3	60	2.9	0.7	78	2.6	0.6
1997	385	18.9	1.9	397	13.3	1.3	81	4.1	0.9	69	2.2	0.5
1998	453	22.2	2.1	415	14.2	1.4	59	2.8	0.7	65	2.1	0.5
1999	412	19.9	1.9	402	13.7	1.4	67	3.3	0.8	76	2.4	0.5
2000	463	22.3	2.0	362	12.0	1.3	66	3.2	0.8	70	2.2	0.5
2001	405	19.5	1.9	371	13.0	1.3	85	4.1	0.9	66	2.2	0.5
2002	418	19.6	1.9	345	11.6	1.2	67	3.2	0.8	74	2.3	0.5
2003	414	19.6	1.9	350	11.9	1.3	65	3.1	0.8	69	2.3	0.6
2004	377	17.9	1.8	359	12.3	1.3	66	3.2	0.8	48	1.5	0.4
2005	369	17.6	1.8	298	10.2	1.2	58	2.8	0.7	58	1.9	0.5
2006	399	19.0	1.9	324	11.3	1.3	63	3.1	0.8	47	1.6	0.5
2007	379	17.9	1.8	335	11.8	1.3	50	2.4	0.7	59	1.9	0.5
2008	357	16.7	1.8	308	11.0	1.3	72	3.4	0.8	69	2.2	0.5
2009	363	16.7	1.7	284	10.4	1.2	57	2.7	0.7	66	2.2	0.6
2010	345	16.1	1.7	289	10.4	1.2	72	3.4	0.8	44	1.5	0.5
2011	297	13.6	1.6	282	10.0	1.2	66	3.1	0.8	63	2.0	0.5
2012	315	14.1	1.6	229	8.3	1.1	59	2.7	0.7	61	2.1	0.6
2013	339	15.0	1.6	273	9.5	1.2	59	2.6	0.7	54	1.8	0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Liver / intrahepatic bile duct

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	122	5.0	0.9	101	2.8	0.6	122	4.8	0.9	110	3.1	0.6
1977	131	5.3	0.9	85	2.3	0.5	136	5.4	0.9	120	3.3	0.6
1978	109	4.4	0.9	81	2.3	0.5	136	5.5	1.0	95	2.6	0.5
1979	116	4.9	0.9	75	2.2	0.5	133	5.7	1.0	118	3.4	0.6
1980	112	4.9	0.9	91	2.8	0.6	108	4.7	0.9	107	3.0	0.6
1981	110	4.9	0.9	96	2.7	0.6	142	6.2	1.1	119	3.4	0.6
1982	123	5.5	1.0	52	1.5	0.4	140	6.1	1.0	101	2.9	0.6
1983	114	5.0	0.9	82	2.5	0.6	132	5.8	1.0	119	3.6	0.7
1984	117	5.1	0.9	79	2.3	0.5	130	5.8	1.0	114	3.2	0.6
1985	126	5.8	1.0	85	2.5	0.5	153	7.0	1.1	110	3.1	0.6
1986	142	6.4	1.1	89	2.6	0.6	137	6.2	1.1	129	3.8	0.7
1987	118	5.5	1.0	75	2.2	0.5	144	6.8	1.1	114	3.4	0.6
1988	139	6.5	1.1	88	2.7	0.6	150	7.0	1.1	124	3.7	0.7
1989	118	5.5	1.0	73	2.2	0.5	130	6.0	1.0	97	2.9	0.6
1990	147	7.2	1.2	94	3.0	0.6	152	7.5	1.2	120	3.7	0.7
1991	160	7.6	1.2	97	3.1	0.6	159	7.6	1.2	116	3.5	0.6
1992	158	7.6	1.2	106	3.5	0.7	175	8.5	1.3	121	3.8	0.7
1993	144	7.0	1.2	98	3.2	0.7	139	6.7	1.1	126	4.0	0.7
1994	172	8.5	1.3	101	3.2	0.6	152	7.6	1.2	89	2.9	0.6
1995	184	9.0	1.3	108	3.6	0.7	155	7.7	1.2	114	3.6	0.7
1996	223	11.1	1.5	101	3.4	0.7	167	8.3	1.3	98	3.1	0.6
1997	221	10.9	1.4	121	3.9	0.7	145	7.2	1.2	102	3.2	0.6
1998	255	12.4	1.5	146	5.0	0.8	183	9.0	1.3	108	3.5	0.7
1999	247	11.9	1.5	126	4.1	0.7	170	8.3	1.2	106	3.5	0.7
2000	259	12.4	1.5	112	3.8	0.7	176	8.4	1.2	102	3.4	0.7
2001	270	12.8	1.5	132	4.6	0.8	186	8.9	1.3	110	3.6	0.7
2002	262	12.3	1.5	114	3.9	0.7	193	9.1	1.3	110	3.6	0.7
2003	296	13.9	1.6	147	5.0	0.8	186	8.9	1.3	116	3.7	0.7
2004	306	14.2	1.6	127	4.5	0.8	193	9.0	1.3	120	3.9	0.7
2005	334	15.5	1.7	130	4.5	0.8	204	9.5	1.3	113	3.7	0.7
2006	330	15.3	1.7	167	5.9	0.9	213	10.1	1.4	124	4.2	0.8
2007	362	16.5	1.7	129	4.6	0.8	228	10.6	1.4	113	3.9	0.7
2008	354	16.1	1.7	133	4.8	0.8	206	9.4	1.3	97	3.3	0.7
2009	341	15.4	1.7	147	5.3	0.9	237	11.0	1.4	118	4.2	0.8
2010	383	17.1	1.7	139	4.9	0.8	250	11.4	1.4	112	3.6	0.7
2011	345	15.1	1.6	151	5.2	0.8	214	9.5	1.3	112	3.6	0.7
2012	377	16.2	1.7	158	5.6	0.9	217	9.7	1.3	130	4.3	0.8
2013	334	14.2	1.5	168	5.7	0.9	223	9.7	1.3	116	3.8	0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Pancreas

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	410	16.5	1.7	410	11.1	1.1	366	14.8	1.6	371	10.2	1.1
1977	389	16.2	1.7	427	11.8	1.2	360	15.0	1.6	405	11.0	1.1
1978	392	15.7	1.6	391	10.7	1.1	353	14.3	1.5	362	9.9	1.0
1979	365	15.1	1.6	386	10.6	1.1	351	14.7	1.6	368	10.1	1.1
1980	390	16.7	1.7	410	11.3	1.1	369	15.6	1.6	363	10.0	1.0
1981	376	16.4	1.7	374	10.6	1.1	329	14.2	1.6	372	10.2	1.1
1982	348	15.9	1.7	440	12.3	1.2	304	13.6	1.6	360	10.0	1.1
1983	347	14.9	1.6	388	10.9	1.1	319	13.7	1.5	335	9.5	1.0
1984	343	15.5	1.7	423	11.8	1.1	302	13.4	1.6	367	10.3	1.1
1985	322	14.2	1.6	390	11.1	1.1	283	12.8	1.5	309	8.7	1.0
1986	317	14.6	1.7	412	11.9	1.2	262	11.9	1.5	350	9.7	1.0
1987	300	13.8	1.6	388	11.0	1.1	263	11.9	1.5	305	8.7	1.0
1988	284	13.3	1.6	395	11.3	1.1	257	12.1	1.5	301	8.6	1.0
1989	301	14.2	1.6	373	11.2	1.2	266	12.6	1.5	325	9.5	1.1
1990	298	14.2	1.6	358	10.8	1.1	250	12.0	1.5	301	8.9	1.0
1991	291	14.2	1.7	367	11.1	1.2	248	12.1	1.5	288	8.7	1.0
1992	319	15.8	1.8	361	11.0	1.2	230	11.3	1.5	297	8.9	1.0
1993	286	14.1	1.6	380	11.8	1.2	279	13.8	1.6	302	9.2	1.1
1994	309	15.5	1.7	402	12.6	1.3	244	12.2	1.5	356	11.0	1.2
1995	261	13.1	1.6	360	11.4	1.2	243	12.2	1.5	301	9.2	1.1
1996	297	15.0	1.7	359	11.7	1.2	275	13.9	1.7	348	11.2	1.2
1997	296	14.8	1.7	357	11.6	1.2	230	11.8	1.5	294	9.3	1.1
1998	323	16.2	1.8	355	11.6	1.2	247	12.4	1.6	301	9.7	1.1
1999	328	16.2	1.8	339	11.0	1.2	295	14.6	1.7	278	8.9	1.1
2000	328	16.0	1.7	392	12.7	1.3	237	11.6	1.5	308	9.6	1.1
2001	288	14.0	1.6	365	11.6	1.2	264	13.0	1.6	318	9.9	1.1
2002	289	13.9	1.6	333	10.4	1.1	217	10.5	1.4	293	9.2	1.1
2003	321	15.6	1.7	378	12.0	1.2	264	12.7	1.5	275	8.5	1.0
2004	308	14.8	1.7	387	12.5	1.3	258	12.5	1.5	296	9.2	1.1
2005	323	15.4	1.7	339	11.4	1.2	269	12.9	1.6	288	9.4	1.1
2006	325	15.7	1.7	372	12.2	1.3	253	12.3	1.5	303	9.8	1.1
2007	348	16.6	1.8	368	11.9	1.2	262	12.6	1.5	302	9.8	1.1
2008	366	17.5	1.8	405	13.3	1.3	271	13.0	1.6	328	10.4	1.2
2009	382	18.1	1.8	387	12.9	1.3	317	15.1	1.7	299	9.8	1.1
2010	366	17.1	1.8	377	12.6	1.3	285	13.3	1.6	313	9.9	1.1
2011	363	16.8	1.7	372	12.3	1.3	298	13.9	1.6	306	9.9	1.1
2012	358	16.4	1.7	375	12.6	1.3	295	13.4	1.5	298	9.7	1.1
2013	355	15.9	1.7	365	11.9	1.3	271	12.3	1.5	275	8.7	1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Larynx

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	286	10.5	1.3	65	1.8	0.5	108	4.1	0.8	29	0.8	0.3
1977	286	11.0	1.3	49	1.4	0.4	125	4.8	0.9	15	0.4	0.2
1978	250	10.2	1.3	64	1.8	0.5	102	4.1	0.8	25	0.7	0.3
1979	294	11.5	1.4	64	2.0	0.5	106	4.3	0.9	25	0.7	0.3
1980	303	12.6	1.5	65	2.0	0.5	104	4.4	0.9	18	0.5	0.2
1981	272	11.4	1.4	70	2.1	0.5	93	3.8	0.8	21	0.6	0.3
1982	263	11.1	1.4	70	2.2	0.5	97	4.0	0.8	24	0.7	0.3
1983	241	10.1	1.3	74	2.3	0.6	83	3.5	0.8	26	0.7	0.3
1984	283	12.0	1.4	77	2.5	0.6	85	3.6	0.8	21	0.7	0.3
1985	267	11.5	1.4	79	2.6	0.6	95	4.2	0.9	23	0.7	0.3
1986	242	10.6	1.4	71	2.4	0.6	65	2.9	0.7	16	0.4	0.2
1987	231	10.5	1.4	63	2.3	0.6	77	3.5	0.8	22	0.7	0.3
1988	245	10.9	1.4	65	2.2	0.6	75	3.5	0.8	20	0.7	0.3
1989	216	10.2	1.4	71	2.6	0.6	68	3.1	0.7	28	0.9	0.4
1990	256	12.1	1.5	57	2.0	0.5	75	3.5	0.8	9	0.3	0.2
1991	234	11.3	1.5	54	2.0	0.6	75	3.6	0.8	22	0.7	0.3
1992	219	10.5	1.4	71	2.6	0.6	82	4.1	0.9	27	1.0	0.4
1993	212	10.3	1.4	58	2.1	0.6	81	3.9	0.9	18	0.6	0.3
1994	205	10.2	1.4	55	2.0	0.5	76	3.7	0.8	19	0.7	0.3
1995	170	8.2	1.2	47	1.7	0.5	68	3.4	0.8	22	0.7	0.3
1996	225	11.1	1.5	59	2.2	0.6	83	4.1	0.9	16	0.5	0.3
1997	205	10.1	1.4	41	1.5	0.5	57	2.9	0.7	18	0.7	0.3
1998	184	8.9	1.3	57	2.1	0.5	66	3.2	0.8	17	0.6	0.3
1999	172	8.3	1.2	40	1.5	0.5	60	2.9	0.7	17	0.6	0.3
2000	174	8.4	1.2	57	2.1	0.5	68	3.3	0.8	17	0.6	0.3
2001	169	8.1	1.2	57	2.1	0.5	48	2.3	0.7	13	0.4	0.2
2002	170	8.1	1.2	36	1.3	0.4	61	2.9	0.7	16	0.5	0.3
2003	178	8.4	1.2	50	1.7	0.5	60	2.9	0.7	18	0.6	0.3
2004	191	8.9	1.3	36	1.3	0.4	64	3.0	0.7	5	0.2	0.2
2005	161	7.4	1.2	46	1.7	0.5	52	2.5	0.7	13	0.4	0.2
2006	157	7.4	1.2	33	1.3	0.4	58	2.8	0.7	10	0.3	0.2
2007	151	7.1	1.1	48	1.7	0.5	48	2.2	0.6	15	0.5	0.3
2008	177	8.3	1.2	52	1.9	0.5	53	2.5	0.7	8	0.3	0.2
2009	150	6.8	1.1	48	1.8	0.5	52	2.4	0.7	16	0.5	0.3
2010	159	7.3	1.2	35	1.3	0.4	47	2.2	0.6	15	0.5	0.2
2011	124	5.4	1.0	29	1.0	0.4	48	2.2	0.6	12	0.4	0.2
2012	163	7.1	1.1	33	1.2	0.4	44	1.9	0.6	13	0.5	0.3
2013	145	6.5	1.1	29	1.0	0.4	46	2.2	0.6	10	0.4	0.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Lung and bronchus

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	2207	84.1	3.7	959	26.3	1.7	1825	70.0	3.4	739	20.3	1.5
1977	2152	84.5	3.7	1000	27.5	1.8	1853	72.5	3.4	743	20.4	1.5
1978	2228	89.0	3.8	947	27.1	1.8	1821	73.5	3.5	752	21.4	1.6
1979	2161	87.8	3.8	1058	30.2	1.9	1767	72.2	3.5	787	22.4	1.6
1980	2121	87.7	3.9	1062	30.8	1.9	1784	74.1	3.6	881	25.4	1.7
1981	1974	83.4	3.8	1085	31.3	1.9	1676	71.1	3.5	808	23.4	1.7
1982	2031	87.0	3.9	1196	35.1	2.1	1671	71.4	3.5	877	25.6	1.8
1983	2046	88.1	3.9	1186	35.0	2.1	1535	67.1	3.5	975	28.1	1.8
1984	2065	90.1	4.0	1157	35.2	2.1	1635	72.1	3.6	867	25.5	1.8
1985	2014	87.9	3.9	1204	36.1	2.1	1550	68.4	3.5	958	28.5	1.9
1986	1934	85.5	3.9	1232	37.4	2.2	1469	65.0	3.4	957	28.9	1.9
1987	1950	86.9	3.9	1260	38.7	2.2	1549	69.7	3.5	946	28.7	1.9
1988	1848	84.9	3.9	1278	39.9	2.3	1527	69.9	3.6	927	28.7	1.9
1989	1802	83.5	3.9	1314	42.2	2.4	1433	67.2	3.5	955	29.9	2.0
1990	1793	84.6	4.0	1320	43.0	2.4	1403	66.4	3.5	976	31.1	2.0
1991	1765	84.8	4.0	1385	46.0	2.5	1368	66.1	3.5	1046	33.8	2.1
1992	1882	90.6	4.1	1387	46.7	2.5	1372	66.4	3.6	969	31.7	2.1
1993	1761	85.8	4.0	1386	47.2	2.6	1344	66.0	3.6	1037	34.2	2.1
1994	1694	82.7	4.0	1455	50.1	2.6	1419	70.0	3.7	1003	33.8	2.1
1995	1678	82.1	4.0	1358	46.0	2.5	1306	64.3	3.5	1046	34.8	2.2
1996	1642	81.6	4.0	1385	48.0	2.6	1281	63.6	3.5	1018	34.0	2.1
1997	1641	81.2	3.9	1386	48.1	2.6	1229	61.2	3.4	932	31.6	2.1
1998	1593	78.0	3.8	1406	48.2	2.6	1182	58.3	3.3	1014	34.3	2.1
1999	1481	72.1	3.7	1401	47.9	2.5	1155	56.7	3.3	1000	34.0	2.1
2000	1575	76.2	3.8	1428	48.9	2.6	1151	56.0	3.2	962	31.5	2.0
2001	1516	72.9	3.7	1428	48.4	2.5	1146	55.5	3.2	981	32.4	2.1
2002	1544	73.6	3.7	1396	47.4	2.5	1117	53.5	3.1	964	31.7	2.0
2003	1508	72.3	3.7	1308	44.2	2.4	1095	52.5	3.1	959	31.8	2.0
2004	1480	70.7	3.6	1297	44.3	2.4	1071	51.5	3.1	867	28.8	2.0
2005	1486	71.2	3.6	1329	46.0	2.5	1056	50.9	3.1	839	28.6	2.0
2006	1439	69.5	3.6	1356	47.2	2.6	1007	48.8	3.0	816	27.4	1.9
2007	1513	72.8	3.7	1391	48.8	2.6	1004	48.7	3.0	903	30.7	2.0
2008	1412	67.6	3.6	1394	49.0	2.6	1013	48.7	3.0	867	29.4	2.0
2009	1426	67.9	3.6	1408	48.9	2.6	887	42.6	2.8	860	28.9	2.0
2010	1358	64.2	3.5	1299	45.2	2.5	945	45.0	2.9	878	29.6	2.0
2011	1373	63.8	3.4	1349	46.4	2.5	908	42.5	2.8	824	27.0	1.9
2012	1293	59.6	3.3	1357	47.2	2.6	922	42.6	2.8	781	26.5	1.9
2013	1315	59.7	3.3	1339	45.6	2.5	920	42.3	2.8	832	27.7	1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Melanoma of the skin

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	198	8.0	1.2	187	5.7	0.9	81	3.2	0.7	65	2.0	0.5
1977	172	6.9	1.1	215	6.5	0.9	72	2.9	0.7	70	2.0	0.5
1978	166	7.0	1.1	185	6.2	0.9	69	2.9	0.7	44	1.3	0.4
1979	194	8.0	1.2	197	6.8	1.0	75	3.2	0.7	65	2.0	0.5
1980	204	8.6	1.2	209	7.2	1.0	70	3.0	0.7	66	2.1	0.5
1981	185	7.9	1.2	186	6.3	1.0	87	3.9	0.8	69	2.3	0.6
1982	206	9.2	1.3	173	5.8	0.9	73	3.2	0.7	56	1.7	0.5
1983	217	9.6	1.3	217	7.3	1.0	77	3.5	0.8	56	1.9	0.5
1984	239	10.4	1.4	232	7.9	1.1	94	4.1	0.8	64	2.0	0.5
1985	240	10.6	1.4	183	6.5	1.0	88	4.0	0.9	77	2.6	0.6
1986	214	9.6	1.3	220	7.5	1.0	72	3.2	0.8	67	2.2	0.6
1987	228	10.6	1.4	198	6.8	1.0	83	3.8	0.8	58	2.0	0.5
1988	212	9.9	1.4	182	6.2	0.9	73	3.4	0.8	52	1.7	0.5
1989	203	9.6	1.3	196	6.7	1.0	67	3.3	0.8	38	1.4	0.5
1990	214	10.1	1.4	181	6.5	1.0	65	3.2	0.8	50	1.8	0.5
1991	228	10.7	1.4	186	6.4	1.0	58	2.8	0.7	54	1.8	0.5
1992	179	8.6	1.3	172	6.0	0.9	58	2.7	0.7	36	1.2	0.4
1993	206	10.0	1.4	155	5.4	0.9	49	2.5	0.7	35	1.2	0.4
1994	247	11.9	1.5	184	6.6	1.0	72	3.5	0.8	39	1.4	0.4
1995	248	11.9	1.5	174	6.3	1.0	65	3.2	0.8	50	1.8	0.5
1996	242	11.7	1.5	219	8.1	1.1	57	2.7	0.7	55	1.9	0.5
1997	269	13.3	1.6	222	8.1	1.1	60	2.9	0.7	49	1.8	0.5
1998	263	12.6	1.5	237	8.6	1.1	72	3.6	0.8	54	1.8	0.5
1999	268	12.7	1.5	231	8.4	1.1	64	3.1	0.8	42	1.4	0.4
2000	314	15.0	1.7	281	10.4	1.2	72	3.5	0.8	52	1.8	0.5
2001	323	15.3	1.7	276	10.1	1.2	60	2.9	0.7	52	1.8	0.5
2002	328	15.4	1.7	279	10.2	1.2	50	2.4	0.7	49	1.7	0.5
2003	278	13.0	1.5	252	9.3	1.2	50	2.4	0.7	57	2.0	0.5
2004	394	18.4	1.8	353	13.2	1.4	53	2.5	0.7	46	1.6	0.5
2005	427	20.0	1.9	382	14.4	1.5	49	2.4	0.7	47	1.6	0.5
2006	543	25.3	2.1	443	16.7	1.6	67	3.3	0.8	43	1.5	0.5
2007	421	19.6	1.9	380	14.4	1.5	63	3.0	0.8	43	1.4	0.4
2008	442	20.7	1.9	371	14.1	1.5	57	2.7	0.7	43	1.5	0.5
2009	476	22.1	2.0	412	15.2	1.5	63	3.0	0.7	34	1.2	0.4
2010	408	19.0	1.9	359	13.4	1.4	59	2.7	0.7	42	1.4	0.4
2011	486	22.1	2.0	416	15.5	1.5	43	2.0	0.6	39	1.2	0.4
2012	435	19.5	1.9	364	13.2	1.4	69	3.1	0.7	55	1.8	0.5
2013	472	21.0	1.9	367	13.3	1.4	64	2.8	0.7	37	1.2	0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Female breast

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	3507	106.0	3.7	1353	38.7	2.1
1977	3314	102.2	3.7	1339	39.3	2.2
1978	3359	104.8	3.7	1332	39.6	2.2
1979	3270	104.2	3.8	1199	36.6	2.2
1980	3224	104.4	3.8	1314	39.8	2.3
1981	3363	110.0	3.9	1273	39.3	2.3
1982	3275	108.2	3.9	1190	37.0	2.2
1983	3267	108.2	3.9	1119	35.2	2.2
1984	3478	116.0	4.1	1186	37.7	2.3
1985	3435	115.0	4.1	1153	36.5	2.2
1986	3325	112.7	4.0	1102	35.0	2.2
1987	3427	116.6	4.1	1069	34.3	2.1
1988	3426	118.5	4.1	1162	37.3	2.2
1989	3377	119.6	4.2	1097	36.4	2.2
1990	3599	130.4	4.4	1031	34.6	2.2
1991	3627	132.9	4.5	986	33.5	2.2
1992	3600	133.2	4.5	977	33.3	2.2
1993	3403	126.5	4.4	987	33.6	2.2
1994	3750	138.8	4.6	987	34.0	2.2
1995	3660	136.7	4.5	1030	35.7	2.2
1996	3831	144.0	4.7	1051	36.9	2.3
1997	3658	137.2	4.5	904	32.0	2.1
1998	3723	139.5	4.6	842	29.6	2.0
1999	3715	138.4	4.5	818	28.1	2.0
2000	3604	133.7	4.4	812	27.6	1.9
2001	3451	127.4	4.3	828	28.3	2.0
2002	3385	124.7	4.3	762	25.7	1.9
2003	3292	122.4	4.3	742	25.3	1.9
2004	3273	122.1	4.3	763	25.9	1.9
2005	3191	119.9	4.2	720	24.7	1.8
2006	3252	123.9	4.3	711	24.7	1.9
2007	3402	130.6	4.5	663	22.9	1.8
2008	3407	129.9	4.5	659	22.7	1.8
2009	3473	131.9	4.5	650	22.7	1.8
2010	3304	125.5	4.4	641	22.0	1.8
2011	3383	127.4	4.4	597	20.3	1.7
2012	3226	121.4	4.3	606	20.8	1.7
2013	3369	126.0	4.4	599	20.1	1.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Cervix uteri

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	384	12.7	1.4	129	3.9	0.7
1977	340	11.3	1.3	128	3.9	0.7
1978	342	12.2	1.4	126	4.1	0.8
1979	356	12.9	1.4	103	3.4	0.7
1980	318	12.0	1.4	105	3.3	0.7
1981	310	11.3	1.3	113	4.0	0.8
1982	310	11.8	1.4	109	3.7	0.7
1983	323	12.2	1.4	90	3.0	0.7
1984	304	11.5	1.4	77	2.7	0.6
1985	301	11.6	1.4	81	2.9	0.7
1986	267	10.2	1.3	66	2.3	0.6
1987	321	12.3	1.4	70	2.3	0.6
1988	295	11.4	1.4	77	2.9	0.7
1989	287	11.2	1.3	64	2.4	0.6
1990	321	12.8	1.4	89	3.3	0.7
1991	331	13.6	1.5	69	2.6	0.6
1992	324	12.7	1.4	62	2.3	0.6
1993	303	12.4	1.4	65	2.5	0.6
1994	350	14.2	1.5	95	3.7	0.8
1995	346	13.9	1.5	87	3.4	0.7
1996	339	13.8	1.5	96	3.6	0.7
1997	311	12.3	1.4	85	3.4	0.7
1998	282	11.3	1.3	75	2.9	0.7
1999	279	11.0	1.3	94	3.7	0.8
2000	258	10.2	1.3	81	3.0	0.7
2001	218	8.8	1.2	64	2.4	0.6
2002	240	9.8	1.3	71	2.7	0.6
2003	232	9.1	1.2	78	3.0	0.7
2004	223	8.8	1.2	69	2.7	0.6
2005	248	10.1	1.3	77	2.8	0.6
2006	215	8.6	1.2	76	2.9	0.7
2007	212	8.8	1.2	66	2.6	0.6
2008	230	9.3	1.2	61	2.3	0.6
2009	220	9.0	1.2	59	2.2	0.6
2010	177	7.5	1.1	64	2.4	0.6
2011	188	7.6	1.1	68	2.6	0.6
2012	204	8.3	1.2	62	2.3	0.6
2013	192	7.7	1.1	61	2.1	0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Corpus uterus and uterus, not otherwise specified

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	847	24.1	1.7	248	6.6	0.8
1977	748	21.7	1.6	198	5.4	0.8
1978	681	19.5	1.5	184	5.0	0.8
1979	753	22.4	1.7	169	4.6	0.7
1980	786	23.7	1.7	196	5.4	0.8
1981	733	22.2	1.7	171	4.8	0.7
1982	733	22.7	1.7	164	4.7	0.7
1983	751	23.0	1.7	165	4.6	0.7
1984	684	21.2	1.7	163	4.7	0.7
1985	699	21.9	1.7	130	3.8	0.7
1986	664	21.5	1.7	138	3.9	0.7
1987	665	21.6	1.7	153	4.5	0.7
1988	667	22.1	1.7	125	3.7	0.7
1989	635	21.3	1.7	130	3.9	0.7
1990	607	21.3	1.8	130	3.9	0.7
1991	697	24.9	1.9	144	4.7	0.8
1992	681	24.5	1.9	137	4.4	0.8
1993	654	23.4	1.8	146	4.7	0.8
1994	723	25.9	1.9	146	4.8	0.8
1995	726	26.7	2.0	155	5.1	0.8
1996	717	26.2	2.0	141	4.7	0.8
1997	764	28.2	2.0	148	5.0	0.8
1998	805	30.0	2.1	145	5.0	0.8
1999	813	29.7	2.1	153	5.2	0.8
2000	745	27.3	2.0	132	4.2	0.7
2001	803	29.5	2.1	125	4.2	0.7
2002	806	29.2	2.0	133	4.5	0.8
2003	804	29.1	2.0	140	4.7	0.8
2004	846	31.0	2.1	125	4.2	0.8
2005	822	30.1	2.1	144	5.0	0.8
2006	823	30.1	2.1	144	4.9	0.8
2007	774	28.4	2.0	152	5.2	0.8
2008	927	34.2	2.2	149	5.1	0.8
2009	847	31.4	2.2	125	4.3	0.8
2010	819	30.1	2.1	177	6.0	0.9
2011	884	32.1	2.2	165	5.6	0.9
2012	813	29.1	2.0	152	5.0	0.8
2013	903	32.3	2.2	159	5.4	0.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Ovary

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	565	16.1	1.4	381	10.7	1.1
1977	507	15.6	1.4	395	11.2	1.2
1978	529	16.4	1.5	335	9.7	1.1
1979	528	16.9	1.5	340	9.8	1.1
1980	523	16.5	1.5	354	10.4	1.1
1981	548	17.5	1.5	336	10.0	1.1
1982	554	18.2	1.6	314	9.5	1.1
1983	540	17.5	1.6	319	9.6	1.1
1984	521	17.0	1.5	328	10.0	1.1
1985	555	18.6	1.6	296	9.4	1.1
1986	476	16.0	1.5	316	9.9	1.1
1987	516	17.9	1.6	304	9.7	1.1
1988	481	16.6	1.6	342	10.8	1.2
1989	473	16.4	1.5	269	8.6	1.1
1990	490	17.7	1.6	283	9.3	1.1
1991	525	18.8	1.7	312	10.5	1.2
1992	480	17.7	1.6	311	10.7	1.2
1993	460	17.1	1.6	265	8.8	1.1
1994	487	17.7	1.6	274	9.4	1.1
1995	541	20.3	1.8	277	9.7	1.2
1996	480	17.9	1.6	280	10.0	1.2
1997	484	18.1	1.6	289	10.2	1.2
1998	437	16.4	1.6	248	8.7	1.1
1999	436	16.2	1.5	257	8.9	1.1
2000	414	15.1	1.5	265	9.1	1.1
2001	461	17.1	1.6	239	8.3	1.1
2002	456	16.3	1.5	263	9.1	1.1
2003	395	14.5	1.5	269	9.3	1.1
2004	426	15.7	1.5	265	9.0	1.1
2005	378	14.1	1.4	244	8.7	1.1
2006	373	13.7	1.4	246	8.6	1.1
2007	399	15.0	1.5	230	8.0	1.1
2008	384	14.1	1.4	232	8.0	1.1
2009	392	14.8	1.5	216	7.2	1.0
2010	361	13.4	1.4	223	7.6	1.0
2011	382	14.0	1.4	236	7.8	1.0
2012	333	12.0	1.3	233	8.1	1.1
2013	333	12.3	1.4	230	7.8	1.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Prostate

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	1576	67.8	3.5	563	26.3	2.3
1977	1574	67.5	3.5	572	26.2	2.3
1978	1555	67.6	3.5	591	27.4	2.3
1979	1700	75.0	3.7	533	25.8	2.3
1980	1815	80.6	3.8	552	26.8	2.3
1981	1726	76.7	3.7	556	27.1	2.3
1982	1692	76.6	3.8	519	25.4	2.3
1983	1727	77.2	3.7	587	28.7	2.4
1984	1743	78.9	3.8	612	30.1	2.4
1985	1821	81.8	3.9	520	24.9	2.2
1986	1840	84.7	4.0	507	24.7	2.2
1987	1835	84.0	3.9	524	25.8	2.3
1988	1723	81.6	3.9	583	29.1	2.4
1989	1731	82.1	3.9	548	27.9	2.4
1990	2123	101.7	4.4	568	29.3	2.5
1991	2409	114.8	4.6	534	27.8	2.4
1992	2944	140.7	5.1	596	31.0	2.5
1993	2788	133.8	5.0	573	30.0	2.5
1994	2914	140.3	5.1	606	32.2	2.6
1995	2674	129.5	4.9	596	31.6	2.6
1996	2721	133.3	5.0	559	29.8	2.5
1997	2902	142.2	5.2	494	26.5	2.3
1998	2914	141.7	5.2	531	28.0	2.4
1999	2962	143.5	5.2	431	22.5	2.1
2000	3413	164.0	5.5	484	24.9	2.2
2001	3182	152.0	5.3	462	23.7	2.2
2002	3137	148.9	5.2	463	23.7	2.2
2003	2812	133.6	5.0	452	22.9	2.1
2004	2769	130.5	4.9	424	21.5	2.0
2005	2832	133.3	4.9	403	20.4	2.0
2006	3183	151.1	5.3	382	19.3	1.9
2007	3290	154.5	5.3	392	19.8	2.0
2008	3115	144.6	5.1	377	19.0	1.9
2009	3117	143.0	5.1	393	19.7	2.0
2010	2853	129.1	4.8	426	21.0	2.0
2011	3038	135.3	4.9	391	18.8	1.9
2012	2556	111.8	4.4	339	16.4	1.8
2013	2440	105.4	4.2	327	15.6	1.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Testis

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	76	2.8	0.7	16	0.6	0.3
1977	72	2.8	0.7	16	0.6	0.3
1978	69	2.8	0.7	15	0.6	0.3
1979	80	3.4	0.8	16	0.6	0.3
1980	97	4.1	0.8	16	0.7	0.3
1981	99	3.9	0.8	4	0.1	0.1
1982	108	4.3	0.8	7	0.3	0.2
1983	101	3.9	0.8	7	0.3	0.2
1984	93	3.8	0.8	10	0.4	0.3
1985	96	3.8	0.8	14	0.5	0.3
1986	108	4.3	0.8	9	0.4	0.3
1987	106	4.2	0.8	7	0.3	0.2
1988	89	3.6	0.8	7	0.2	0.2
1989	103	4.2	0.8	10	0.4	0.3
1990	117	4.8	0.9	8	0.3	0.2
1991	121	4.9	0.9	9	0.4	0.3
1992	113	4.8	0.9	9	0.4	0.3
1993	117	4.8	0.9	9	0.4	0.3
1994	115	4.7	0.9	9	0.4	0.3
1995	106	4.4	0.8	7	0.3	0.2
1996	108	4.4	0.8	5	0.2	0.2
1997	101	4.1	0.8	6	0.2	0.2
1998	99	4.1	0.8	7	0.3	0.2
1999	146	5.9	1.0	4	0.2	0.2
2000	128	5.2	0.9	2	0.1	0.1
2001	118	4.7	0.9	6	0.3	0.2
2002	127	5.2	0.9	3	0.1	0.1
2003	138	5.7	1.0	8	0.4	0.3
2004	144	5.9	1.0	8	0.3	0.2
2005	148	6.0	1.0	7	0.3	0.2
2006	149	6.1	1.0	11	0.5	0.3
2007	160	6.6	1.0	1	0.0	0.1
2008	148	6.3	1.0	5	0.2	0.2
2009	169	7.1	1.1	8	0.3	0.2
2010	159	6.7	1.1	14	0.6	0.3
2011	164	6.7	1.1	7	0.3	0.2
2012	169	7.0	1.1	7	0.3	0.2
2013	157	6.4	1.0	6	0.2	0.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Urinary bladder (incl. in situ)

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	843	33.8	2.4	333	8.9	1.0	249	11.2	1.5	116	3.4	0.6
1977	892	36.2	2.5	373	10.3	1.1	267	11.9	1.5	117	3.3	0.6
1978	947	39.8	2.6	369	10.0	1.1	234	10.5	1.4	119	3.4	0.6
1979	923	39.6	2.6	401	11.3	1.1	242	11.1	1.5	104	2.9	0.6
1980	1035	44.5	2.8	353	9.8	1.0	249	11.3	1.5	109	3.0	0.6
1981	987	42.9	2.8	368	10.4	1.1	202	9.3	1.3	99	2.7	0.5
1982	886	38.8	2.6	365	10.5	1.1	226	10.5	1.4	91	2.5	0.5
1983	976	43.8	2.8	411	11.6	1.1	189	9.0	1.3	95	2.6	0.5
1984	910	39.8	2.7	350	10.0	1.1	186	8.7	1.3	96	2.7	0.5
1985	800	36.2	2.6	331	9.5	1.0	201	9.8	1.4	91	2.5	0.5
1986	855	39.2	2.7	355	10.7	1.2	164	7.9	1.2	96	2.6	0.5
1987	879	40.0	2.7	354	10.4	1.1	172	8.4	1.3	71	2.1	0.5
1988	829	38.0	2.6	359	10.5	1.1	169	8.1	1.2	102	2.8	0.5
1989	788	37.5	2.7	323	9.9	1.1	175	8.8	1.3	102	2.9	0.6
1990	792	38.1	2.7	350	10.6	1.1	159	8.1	1.3	90	2.7	0.6
1991	862	41.9	2.8	345	10.8	1.2	148	7.5	1.2	78	2.3	0.5
1992	830	40.7	2.8	351	11.3	1.2	152	7.9	1.3	81	2.4	0.5
1993	780	38.4	2.7	314	10.2	1.2	187	9.6	1.4	86	2.6	0.6
1994	834	41.3	2.8	373	12.1	1.3	175	9.1	1.4	83	2.5	0.5
1995	844	42.2	2.9	333	10.8	1.2	185	9.6	1.4	103	3.0	0.6
1996	737	37.2	2.7	296	9.6	1.1	146	7.7	1.3	93	2.8	0.6
1997	845	42.2	2.9	338	11.1	1.2	154	8.1	1.3	79	2.5	0.6
1998	851	42.3	2.9	370	12.1	1.3	160	8.2	1.3	93	2.8	0.6
1999	783	38.7	2.7	331	10.9	1.2	166	8.6	1.3	74	2.2	0.5
2000	830	40.8	2.8	359	11.8	1.2	154	7.8	1.2	74	2.2	0.5
2001	801	39.1	2.7	323	10.7	1.2	165	8.3	1.3	69	2.1	0.5
2002	778	38.0	2.7	298	9.8	1.1	136	6.8	1.2	72	2.1	0.5
2003	825	40.2	2.8	318	10.4	1.2	131	6.6	1.1	61	1.8	0.5
2004	829	40.0	2.7	340	11.0	1.2	157	7.7	1.2	68	2.0	0.5
2005	809	39.0	2.7	344	11.4	1.2	143	7.2	1.2	82	2.5	0.5
2006	832	40.5	2.8	285	9.6	1.1	156	7.9	1.2	84	2.5	0.6
2007	857	41.9	2.8	316	10.7	1.2	161	8.0	1.2	81	2.4	0.5
2008	883	42.4	2.8	273	9.3	1.1	159	7.9	1.2	68	2.2	0.5
2009	813	39.1	2.7	315	10.5	1.2	143	7.1	1.2	72	2.1	0.5
2010	820	39.1	2.7	255	8.6	1.1	170	8.1	1.2	71	2.1	0.5
2011	794	37.4	2.6	285	9.6	1.1	150	7.2	1.2	71	2.2	0.5
2012	808	37.8	2.6	290	9.6	1.1	159	7.5	1.2	75	2.2	0.5
2013	773	35.4	2.5	274	9.1	1.1	156	7.3	1.2	65	1.8	0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Kidney and renal pelvis

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	285	10.9	1.3	177	5.1	0.8	158	6.4	1.0	83	2.2	0.5
1977	295	11.7	1.4	162	4.7	0.8	150	6.1	1.0	83	2.2	0.5
1978	272	11.0	1.4	175	4.8	0.7	144	5.7	1.0	77	2.3	0.5
1979	268	11.1	1.4	164	4.6	0.7	143	6.1	1.0	84	2.4	0.5
1980	259	10.4	1.3	155	4.7	0.8	117	5.0	0.9	75	2.1	0.5
1981	240	10.2	1.3	177	5.2	0.8	120	5.0	0.9	88	2.5	0.5
1982	273	12.0	1.5	188	5.7	0.8	124	5.5	1.0	85	2.4	0.5
1983	281	12.1	1.5	172	5.3	0.8	116	5.0	0.9	74	2.1	0.5
1984	298	13.0	1.5	180	5.5	0.8	122	5.4	1.0	77	2.3	0.5
1985	313	13.6	1.5	203	6.3	0.9	104	4.6	0.9	78	2.4	0.6
1986	297	13.0	1.5	195	5.8	0.8	126	5.6	1.0	79	2.2	0.5
1987	326	14.4	1.6	221	6.9	0.9	117	5.4	1.0	72	2.2	0.5
1988	283	13.0	1.5	171	5.7	0.9	106	4.9	0.9	70	2.2	0.5
1989	297	13.8	1.6	189	6.1	0.9	114	5.4	1.0	79	2.4	0.5
1990	299	14.0	1.6	230	7.8	1.0	102	4.9	1.0	83	2.6	0.6
1991	303	14.4	1.6	228	7.6	1.0	91	4.3	0.9	71	2.3	0.5
1992	312	15.0	1.7	215	7.3	1.0	86	4.3	0.9	66	2.1	0.5
1993	353	16.9	1.8	189	6.5	1.0	116	5.6	1.0	53	1.7	0.5
1994	362	17.8	1.8	227	7.7	1.0	120	5.9	1.1	75	2.4	0.5
1995	321	15.6	1.7	214	7.6	1.0	113	5.7	1.1	70	2.3	0.6
1996	366	18.1	1.9	257	9.0	1.1	98	4.9	1.0	78	2.5	0.6
1997	353	17.5	1.8	251	9.1	1.1	109	5.5	1.0	75	2.5	0.6
1998	389	18.9	1.9	253	9.1	1.1	95	4.8	1.0	79	2.7	0.6
1999	413	19.8	1.9	246	8.6	1.1	106	5.2	1.0	58	1.9	0.5
2000	401	19.2	1.9	282	9.9	1.2	101	4.9	1.0	57	1.9	0.5
2001	416	19.7	1.9	254	8.9	1.1	96	4.6	0.9	68	2.3	0.6
2002	420	19.8	1.9	255	8.9	1.1	106	5.1	1.0	64	2.1	0.5
2003	478	22.4	2.0	260	9.1	1.1	91	4.4	0.9	58	1.8	0.5
2004	467	22.0	2.0	280	10.2	1.2	100	4.8	0.9	63	2.1	0.5
2005	460	21.7	2.0	282	10.1	1.2	77	3.7	0.8	56	1.8	0.5
2006	507	23.9	2.1	288	10.7	1.3	93	4.6	0.9	67	2.2	0.5
2007	478	22.8	2.1	281	10.3	1.2	101	4.8	0.9	62	2.0	0.5
2008	520	24.2	2.1	286	10.3	1.2	87	4.2	0.9	58	1.9	0.5
2009	433	20.2	1.9	285	10.4	1.2	125	6.0	1.1	56	1.9	0.5
2010	468	21.2	1.9	269	10.0	1.2	98	4.5	0.9	58	1.9	0.5
2011	461	20.8	1.9	279	10.2	1.2	92	4.1	0.9	46	1.4	0.4
2012	485	21.8	2.0	261	9.7	1.2	91	4.3	0.9	45	1.5	0.4
2013	486	21.4	1.9	275	9.9	1.2	75	3.5	0.8	60	1.8	0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Brain and other nervous system

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	188	7.0	1.0	162	4.9	0.8	152	5.9	1.0	121	3.7	0.7
1977	191	7.5	1.1	157	5.0	0.8	137	5.2	0.9	127	3.9	0.7
1978	175	7.0	1.1	175	5.6	0.9	135	5.4	0.9	127	3.6	0.7
1979	177	7.3	1.1	158	5.1	0.8	134	5.5	1.0	108	3.1	0.6
1980	166	6.9	1.1	167	5.3	0.8	144	5.7	1.0	124	3.7	0.7
1981	194	8.5	1.2	153	4.9	0.8	130	5.5	1.0	120	3.7	0.7
1982	186	8.0	1.2	157	5.1	0.8	125	5.3	1.0	130	4.0	0.7
1983	166	7.2	1.1	152	5.2	0.9	127	5.6	1.0	105	3.1	0.6
1984	162	7.0	1.1	182	6.2	1.0	115	4.9	0.9	117	3.7	0.7
1985	194	8.7	1.3	163	5.5	0.9	119	5.3	1.0	120	3.7	0.7
1986	186	8.5	1.3	145	5.2	0.9	135	6.3	1.1	109	3.3	0.6
1987	181	8.2	1.2	178	6.0	0.9	117	5.3	1.0	113	3.8	0.7
1988	190	8.7	1.3	177	6.2	1.0	90	4.0	0.8	79	2.5	0.6
1989	201	9.5	1.3	172	6.2	1.0	98	4.6	0.9	89	2.8	0.6
1990	191	9.1	1.3	181	6.7	1.0	106	5.1	1.0	88	3.0	0.7
1991	181	8.7	1.3	179	6.6	1.0	105	5.0	1.0	105	3.8	0.7
1992	192	9.0	1.3	159	6.0	1.0	91	4.5	0.9	82	2.8	0.6
1993	178	8.5	1.3	184	6.8	1.0	95	4.6	0.9	77	2.7	0.6
1994	195	9.4	1.3	192	7.2	1.1	94	4.6	0.9	90	3.2	0.7
1995	225	10.7	1.4	191	7.1	1.0	106	5.2	1.0	93	3.3	0.7
1996	211	10.1	1.4	202	7.7	1.1	94	4.6	0.9	93	3.3	0.7
1997	177	8.6	1.3	196	7.7	1.1	90	4.5	0.9	119	4.3	0.8
1998	228	10.7	1.4	200	7.9	1.1	104	5.1	1.0	111	4.0	0.8
1999	182	8.5	1.2	169	6.4	1.0	111	5.3	1.0	102	3.6	0.7
2000	177	8.3	1.2	192	7.2	1.0	93	4.4	0.9	92	3.4	0.7
2001	171	8.0	1.2	166	6.3	1.0	95	4.5	0.9	77	2.7	0.6
2002	181	8.4	1.2	170	6.5	1.0	87	4.1	0.9	78	2.9	0.7
2003	168	7.7	1.2	160	6.0	1.0	95	4.4	0.9	67	2.4	0.6
2004	185	8.8	1.3	144	5.7	0.9	76	3.6	0.8	71	2.6	0.6
2005	200	9.4	1.3	152	5.9	1.0	107	5.1	1.0	84	3.0	0.7
2006	197	9.3	1.3	180	7.1	1.1	107	5.1	1.0	81	2.9	0.7
2007	165	7.8	1.2	146	5.8	1.0	109	5.1	1.0	96	3.5	0.7
2008	169	8.0	1.2	172	6.7	1.0	86	4.0	0.9	71	2.6	0.6
2009	182	8.3	1.2	165	6.2	1.0	106	5.0	1.0	89	3.4	0.7
2010	170	8.0	1.2	144	5.9	1.0	91	4.1	0.9	94	3.4	0.7
2011	164	7.5	1.2	139	5.4	0.9	102	4.8	0.9	90	3.3	0.7
2012	162	7.3	1.1	157	6.0	1.0	81	3.6	0.8	102	3.7	0.7
2013	166	7.5	1.2	146	5.6	0.9	85	3.9	0.8	85	2.8	0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Thyroid

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	56	2.3	0.6	137	4.4	0.8	21	0.8	0.4	29	0.8	0.3
1977	63	2.4	0.6	162	5.3	0.9	25	0.9	0.4	22	0.6	0.3
1978	63	2.6	0.7	168	5.7	0.9	11	0.5	0.3	26	0.8	0.3
1979	79	3.4	0.8	125	4.4	0.8	15	0.6	0.3	18	0.5	0.2
1980	57	2.6	0.7	147	5.1	0.9	15	0.7	0.4	25	0.7	0.3
1981	56	2.4	0.6	135	4.7	0.8	12	0.5	0.3	26	0.8	0.3
1982	62	2.9	0.7	136	4.8	0.9	15	0.6	0.3	31	0.8	0.3
1983	59	2.6	0.7	173	6.6	1.0	7	0.3	0.2	20	0.6	0.3
1984	74	3.4	0.8	186	6.8	1.0	13	0.6	0.3	24	0.7	0.3
1985	78	3.4	0.8	185	6.5	1.0	25	1.2	0.5	19	0.6	0.3
1986	57	2.5	0.7	170	6.2	1.0	4	0.2	0.2	25	0.8	0.3
1987	63	3.0	0.8	145	5.4	0.9	10	0.5	0.3	20	0.5	0.2
1988	71	3.2	0.8	141	5.2	0.9	11	0.5	0.3	19	0.6	0.3
1989	51	2.3	0.6	137	5.6	1.0	7	0.3	0.3	21	0.6	0.3
1990	55	2.5	0.7	169	6.5	1.0	8	0.4	0.3	23	0.7	0.3
1991	65	3.0	0.7	177	7.2	1.1	8	0.4	0.3	22	0.7	0.3
1992	64	3.1	0.8	192	7.5	1.1	10	0.5	0.3	16	0.5	0.3
1993	76	3.6	0.8	203	8.1	1.1	9	0.4	0.3	17	0.5	0.2
1994	81	3.8	0.8	208	8.3	1.2	12	0.6	0.3	19	0.6	0.3
1995	90	4.2	0.9	219	8.6	1.2	7	0.3	0.2	16	0.5	0.3
1996	73	3.5	0.8	231	9.4	1.2	11	0.5	0.3	20	0.6	0.3
1997	93	4.3	0.9	254	10.1	1.3	9	0.4	0.3	21	0.7	0.3
1998	106	4.9	0.9	265	10.5	1.3	8	0.4	0.3	15	0.5	0.2
1999	102	4.6	0.9	309	12.3	1.4	10	0.5	0.3	12	0.4	0.2
2000	100	4.5	0.9	300	12.0	1.4	8	0.4	0.3	24	0.8	0.3
2001	124	5.7	1.0	347	13.8	1.5	10	0.5	0.3	16	0.5	0.2
2002	144	6.5	1.1	341	13.8	1.5	11	0.5	0.3	22	0.7	0.3
2003	109	4.9	0.9	351	14.0	1.5	9	0.4	0.3	14	0.4	0.2
2004	133	6.1	1.0	396	15.9	1.6	10	0.5	0.3	11	0.3	0.2
2005	183	8.4	1.2	535	21.8	1.9	16	0.8	0.4	21	0.7	0.3
2006	172	7.6	1.1	520	21.3	1.9	13	0.7	0.4	28	1.0	0.4
2007	181	8.2	1.2	601	24.8	2.0	11	0.5	0.3	28	1.0	0.4
2008	223	9.9	1.3	705	28.9	2.2	14	0.7	0.4	12	0.4	0.2
2009	274	12.1	1.4	784	32.0	2.3	17	0.8	0.4	14	0.5	0.3
2010	243	10.9	1.4	757	31.1	2.3	14	0.6	0.3	17	0.5	0.3
2011	209	9.3	1.3	781	31.8	2.3	10	0.5	0.3	17	0.5	0.3
2012	274	11.9	1.4	813	33.2	2.3	13	0.6	0.3	13	0.5	0.3
2013	312	13.6	1.5	851	35.0	2.4	18	0.8	0.4	20	0.6	0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Hodgkin lymphoma

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	116	4.3	0.8	100	3.1	0.6	51	2.0	0.6	27	0.8	0.3
1977	119	4.5	0.8	89	3.0	0.6	49	1.9	0.5	39	1.3	0.4
1978	112	4.2	0.8	100	3.2	0.7	43	1.7	0.5	35	1.1	0.4
1979	100	4.3	0.9	98	3.4	0.7	37	1.5	0.5	25	0.9	0.4
1980	103	4.4	0.9	95	3.1	0.6	35	1.6	0.5	37	1.1	0.4
1981	101	4.2	0.8	84	2.8	0.6	33	1.5	0.5	31	1.0	0.4
1982	120	5.1	0.9	92	3.2	0.7	26	1.0	0.4	25	0.8	0.3
1983	113	4.9	0.9	79	2.7	0.6	36	1.5	0.5	26	0.8	0.3
1984	103	4.3	0.9	84	3.0	0.7	34	1.5	0.5	29	0.9	0.3
1985	128	5.4	1.0	83	2.8	0.6	30	1.2	0.4	28	1.0	0.4
1986	128	5.5	1.0	81	2.9	0.7	38	1.6	0.5	18	0.6	0.3
1987	87	3.8	0.8	86	3.3	0.7	24	1.0	0.4	16	0.5	0.3
1988	113	4.9	0.9	87	3.2	0.7	22	1.0	0.4	17	0.6	0.3
1989	104	4.5	0.9	76	3.0	0.7	32	1.5	0.5	24	0.9	0.4
1990	97	4.4	0.9	65	2.4	0.6	38	1.7	0.6	17	0.6	0.3
1991	100	4.6	0.9	92	3.8	0.8	25	1.2	0.5	18	0.6	0.3
1992	108	4.7	0.9	73	2.7	0.6	22	0.9	0.4	17	0.7	0.3
1993	81	3.8	0.9	75	2.8	0.7	19	0.8	0.4	15	0.5	0.3
1994	101	4.7	0.9	72	2.9	0.7	18	0.9	0.4	23	0.8	0.3
1995	95	4.3	0.9	86	3.6	0.8	17	0.8	0.4	14	0.6	0.3
1996	74	3.4	0.8	69	2.8	0.7	16	0.7	0.4	15	0.5	0.3
1997	94	4.3	0.9	69	2.9	0.7	18	0.8	0.4	19	0.7	0.3
1998	95	4.2	0.9	87	3.5	0.8	17	0.8	0.4	18	0.7	0.3
1999	77	3.3	0.7	84	3.3	0.7	10	0.5	0.3	11	0.4	0.2
2000	110	4.9	0.9	97	4.0	0.8	19	0.9	0.4	17	0.6	0.3
2001	91	3.9	0.8	89	3.5	0.7	13	0.6	0.3	11	0.4	0.2
2002	105	4.6	0.9	90	3.6	0.8	21	1.0	0.4	8	0.3	0.2
2003	94	4.3	0.9	91	3.9	0.8	11	0.5	0.3	13	0.5	0.3
2004	89	3.9	0.8	82	3.3	0.7	13	0.6	0.3	10	0.4	0.3
2005	103	4.7	0.9	76	3.0	0.7	14	0.7	0.4	9	0.3	0.2
2006	107	4.9	0.9	82	3.3	0.7	12	0.6	0.3	3	0.1	0.2
2007	83	3.7	0.8	104	4.1	0.8	17	0.8	0.4	15	0.5	0.3
2008	81	3.6	0.8	90	3.6	0.8	22	1.0	0.4	11	0.4	0.2
2009	103	4.7	0.9	75	3.1	0.7	15	0.7	0.4	8	0.3	0.2
2010	87	3.8	0.8	85	3.4	0.7	15	0.6	0.3	7	0.3	0.2
2011	99	4.3	0.9	78	3.2	0.7	17	0.8	0.4	15	0.5	0.3
2012	80	3.6	0.8	72	3.1	0.7	11	0.5	0.3	9	0.3	0.2
2013	73	3.2	0.8	96	3.7	0.8	13	0.6	0.3	9	0.3	0.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Non-Hodgkin lymphomas

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	370	14.7	1.6	362	10.2	1.1	217	8.9	1.2	181	5.0	0.8
1977	351	13.8	1.5	344	10.0	1.1	169	6.8	1.1	185	5.2	0.8
1978	351	14.2	1.5	347	10.3	1.1	198	8.1	1.2	173	5.0	0.8
1979	348	14.9	1.6	399	11.6	1.2	181	7.9	1.2	183	5.3	0.8
1980	396	16.7	1.7	419	12.4	1.2	189	7.8	1.1	184	5.3	0.8
1981	382	16.3	1.7	422	12.5	1.2	172	7.5	1.2	203	5.8	0.8
1982	403	17.4	1.8	473	14.2	1.3	190	8.5	1.2	214	6.3	0.9
1983	443	20.2	1.9	426	12.8	1.3	214	9.7	1.3	220	6.5	0.9
1984	447	19.9	1.9	431	13.0	1.3	194	8.8	1.3	182	5.2	0.8
1985	523	23.9	2.1	449	13.5	1.3	240	10.9	1.4	205	6.0	0.8
1986	540	24.1	2.1	486	15.0	1.4	216	10.0	1.4	177	5.2	0.8
1987	561	26.1	2.2	476	14.9	1.4	214	9.8	1.3	197	5.8	0.8
1988	516	23.8	2.1	440	14.0	1.4	218	10.3	1.4	229	6.7	0.9
1989	579	27.5	2.3	410	13.5	1.4	238	11.4	1.5	182	5.7	0.9
1990	638	30.1	2.4	470	15.5	1.4	214	10.2	1.4	214	6.7	0.9
1991	702	33.6	2.5	527	17.6	1.6	222	10.8	1.4	192	6.0	0.9
1992	628	29.8	2.4	492	16.8	1.5	217	10.6	1.4	207	6.6	0.9
1993	642	30.8	2.4	461	15.9	1.5	213	10.5	1.4	186	6.0	0.9
1994	758	36.1	2.6	542	19.0	1.7	236	11.6	1.5	222	7.2	1.0
1995	691	33.3	2.5	538	18.8	1.6	223	11.1	1.5	206	6.7	0.9
1996	664	32.2	2.5	534	19.1	1.7	233	11.8	1.5	203	6.7	0.9
1997	605	29.4	2.4	543	18.8	1.6	229	11.4	1.5	216	7.1	1.0
1998	579	27.7	2.3	496	17.7	1.6	204	10.0	1.4	226	7.5	1.0
1999	553	26.5	2.2	534	18.9	1.6	194	9.6	1.4	206	6.8	0.9
2000	550	25.9	2.2	529	18.6	1.6	197	9.6	1.3	200	6.6	0.9
2001	574	27.2	2.2	484	17.0	1.5	214	10.4	1.4	167	5.2	0.8
2002	577	27.1	2.2	466	16.3	1.5	176	8.5	1.3	162	5.3	0.8
2003	591	27.8	2.3	556	19.6	1.7	166	7.9	1.2	160	5.1	0.8
2004	579	27.3	2.2	537	18.8	1.6	178	8.6	1.3	176	5.7	0.9
2005	580	27.3	2.2	582	20.7	1.7	179	8.7	1.3	171	5.6	0.9
2006	644	30.3	2.4	515	18.3	1.6	187	9.1	1.3	171	5.6	0.9
2007	572	27.1	2.2	530	19.2	1.7	180	8.7	1.3	158	5.4	0.9
2008	627	29.2	2.3	544	19.6	1.7	178	8.5	1.3	135	4.4	0.8
2009	606	28.6	2.3	597	21.3	1.8	172	8.1	1.2	142	4.6	0.8
2010	613	28.3	2.3	551	19.5	1.7	150	7.3	1.2	140	4.6	0.8
2011	575	26.2	2.2	558	19.8	1.7	154	7.2	1.1	148	4.8	0.8
2012	626	28.5	2.3	528	19.0	1.7	162	7.6	1.2	131	4.1	0.7
2013	637	28.5	2.2	531	18.6	1.6	147	6.8	1.1	129	4.1	0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Myeloma

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	123	4.9	0.9	120	3.3	0.6	75	3.2	0.8	76	2.1	0.5
1977	124	5.1	0.9	134	3.6	0.6	81	3.2	0.7	81	2.1	0.5
1978	123	5.1	0.9	130	3.5	0.6	66	2.7	0.7	83	2.2	0.5
1979	139	5.9	1.0	127	3.6	0.6	71	2.9	0.7	102	2.8	0.6
1980	114	5.1	1.0	127	3.6	0.6	78	3.3	0.8	89	2.5	0.5
1981	147	6.3	1.1	155	4.3	0.7	86	3.9	0.8	83	2.3	0.5
1982	121	5.4	1.0	145	4.2	0.7	70	3.0	0.7	89	2.6	0.6
1983	141	6.4	1.1	136	3.9	0.7	68	3.0	0.7	92	2.5	0.5
1984	111	5.2	1.0	121	3.4	0.6	75	3.4	0.8	82	2.3	0.5
1985	136	6.1	1.1	117	3.4	0.6	82	3.8	0.8	63	1.8	0.4
1986	129	5.8	1.0	125	3.6	0.6	68	3.0	0.7	76	2.2	0.5
1987	137	6.4	1.1	111	3.2	0.6	88	4.1	0.9	66	1.9	0.5
1988	128	6.2	1.1	118	3.5	0.6	74	3.5	0.8	67	1.9	0.5
1989	108	5.1	1.0	142	4.6	0.8	72	3.3	0.8	78	2.3	0.5
1990	131	6.1	1.1	132	4.2	0.7	69	3.3	0.8	70	2.1	0.5
1991	111	5.3	1.0	126	4.1	0.7	59	2.8	0.7	68	2.1	0.5
1992	145	7.1	1.2	126	4.0	0.7	74	3.7	0.8	75	2.4	0.6
1993	119	6.0	1.1	140	4.6	0.8	90	4.4	0.9	87	2.7	0.6
1994	113	5.5	1.0	125	4.3	0.8	73	3.5	0.8	81	2.7	0.6
1995	119	5.9	1.1	128	4.4	0.8	71	3.6	0.8	78	2.5	0.6
1996	138	6.9	1.2	146	5.0	0.8	63	3.2	0.8	88	2.9	0.6
1997	120	6.0	1.1	119	4.0	0.7	66	3.3	0.8	67	2.3	0.5
1998	146	7.2	1.2	147	5.0	0.8	69	3.4	0.8	70	2.3	0.6
1999	119	5.8	1.1	160	5.5	0.9	63	3.1	0.8	75	2.5	0.6
2000	131	6.3	1.1	143	4.9	0.8	72	3.5	0.8	96	3.1	0.6
2001	152	7.4	1.2	144	4.9	0.8	70	3.4	0.8	66	2.1	0.5
2002	157	7.6	1.2	145	5.0	0.8	71	3.5	0.8	72	2.4	0.6
2003	169	8.2	1.2	151	5.3	0.9	59	2.9	0.7	74	2.3	0.5
2004	155	7.4	1.2	156	5.4	0.9	56	2.8	0.7	65	2.1	0.5
2005	172	8.2	1.2	164	5.6	0.9	66	3.2	0.8	70	2.3	0.5
2006	161	7.8	1.2	169	5.9	0.9	82	4.0	0.9	44	1.5	0.5
2007	165	7.9	1.2	143	4.9	0.8	68	3.3	0.8	55	1.8	0.5
2008	195	9.3	1.3	136	4.8	0.8	68	3.3	0.8	70	2.4	0.6
2009	202	9.6	1.3	163	5.6	0.9	66	3.2	0.8	51	1.6	0.5
2010	199	9.2	1.3	171	6.0	0.9	82	4.0	0.9	66	2.2	0.5
2011	226	10.4	1.4	160	5.7	0.9	53	2.5	0.7	65	2.2	0.5
2012	212	9.7	1.3	150	5.2	0.9	58	2.8	0.7	65	2.1	0.5
2013	188	8.3	1.2	185	6.2	0.9	85	3.8	0.8	62	2.0	0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Whites
Leukemias

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	383	15.8	1.7	306	9.0	1.1	263	11.2	1.4	237	7.0	0.9
1977	349	14.5	1.6	292	8.8	1.0	254	10.8	1.4	204	6.1	0.9
1978	394	16.8	1.7	293	8.9	1.1	250	10.7	1.4	213	6.4	0.9
1979	328	13.8	1.5	318	9.9	1.1	238	10.2	1.3	229	7.2	1.0
1980	379	16.6	1.7	361	11.4	1.2	252	11.2	1.4	184	5.6	0.8
1981	362	16.0	1.7	293	9.1	1.1	245	10.8	1.4	205	6.4	0.9
1982	383	17.1	1.8	308	9.5	1.1	264	12.0	1.5	210	6.5	0.9
1983	327	14.7	1.6	320	10.7	1.2	224	10.4	1.4	195	6.1	0.9
1984	369	17.2	1.8	319	10.0	1.1	241	11.2	1.5	217	6.5	0.9
1985	354	16.2	1.7	295	9.2	1.1	204	9.4	1.3	203	6.1	0.9
1986	351	15.8	1.7	342	11.4	1.3	202	9.2	1.3	203	6.3	0.9
1987	263	12.4	1.5	294	9.3	1.1	172	8.0	1.2	186	5.9	0.9
1988	311	14.9	1.7	282	9.2	1.1	192	9.2	1.3	161	5.1	0.8
1989	362	17.4	1.8	316	10.3	1.2	192	9.2	1.3	174	5.4	0.8
1990	362	17.3	1.8	327	11.2	1.3	181	9.0	1.3	180	5.9	0.9
1991	345	16.9	1.8	277	9.5	1.2	208	10.4	1.4	147	4.8	0.8
1992	353	17.3	1.8	286	9.8	1.2	178	9.0	1.3	146	4.8	0.8
1993	291	14.2	1.6	301	10.7	1.3	190	9.3	1.3	146	4.9	0.8
1994	353	17.5	1.8	339	11.5	1.3	197	9.7	1.4	190	6.1	0.9
1995	318	15.7	1.7	295	10.3	1.2	199	10.0	1.4	179	6.0	0.9
1996	347	17.0	1.8	276	9.8	1.2	165	8.1	1.2	175	5.9	0.9
1997	352	17.4	1.8	312	11.1	1.3	199	10.0	1.4	147	4.9	0.8
1998	360	17.5	1.8	290	10.2	1.2	205	10.0	1.4	157	5.3	0.9
1999	375	18.3	1.9	349	12.2	1.3	203	9.9	1.4	167	5.6	0.9
2000	341	16.5	1.8	327	11.4	1.3	149	7.2	1.2	180	6.0	0.9
2001	346	16.7	1.8	337	11.7	1.3	176	8.5	1.3	168	5.4	0.8
2002	355	16.9	1.8	274	9.5	1.2	178	8.6	1.3	164	5.4	0.9
2003	366	17.5	1.8	335	11.8	1.3	162	7.8	1.2	147	4.9	0.8
2004	358	17.2	1.8	311	10.8	1.2	185	9.0	1.3	169	5.6	0.9
2005	392	18.7	1.9	305	11.0	1.3	193	9.3	1.3	152	5.1	0.8
2006	366	17.7	1.8	309	11.2	1.3	177	8.6	1.3	167	5.6	0.9
2007	392	18.7	1.9	316	11.8	1.4	178	8.7	1.3	147	5.3	0.9
2008	385	18.7	1.9	307	11.4	1.3	171	8.3	1.3	176	6.2	0.9
2009	411	19.5	1.9	353	12.6	1.4	143	6.9	1.1	173	5.9	0.9
2010	443	20.9	2.0	346	12.4	1.4	178	8.5	1.3	149	5.1	0.8
2011	441	20.3	1.9	361	12.8	1.4	197	9.3	1.3	143	4.5	0.8
2012	421	19.3	1.9	308	11.1	1.3	176	8.3	1.2	135	4.4	0.8
2013	423	19.5	1.9	320	11.4	1.3	196	9.1	1.3	146	5.0	0.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
All Invasive Malignant Tumors

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	17530	471.7	7.4	18112	359.2	5.3	10040	277.4	5.8	9213	180.9	3.7
1977	17708	471.3	7.4	18310	358.1	5.3	10324	281.8	5.8	9394	182.0	3.7
1978	17906	473.7	7.3	18603	360.3	5.3	10361	281.7	5.7	9566	182.0	3.7
1979	18274	477.5	7.3	19039	364.2	5.3	10323	278.0	5.7	9685	182.4	3.7
1980	19373	498.0	7.4	19382	366.7	5.3	10715	284.0	5.7	9847	182.9	3.7
1981	19517	502.0	7.4	19910	374.3	5.3	10724	283.5	5.7	9856	181.1	3.6
1982	19823	503.5	7.4	20260	376.7	5.3	10783	283.7	5.6	10067	183.1	3.6
1983	20365	513.0	7.4	20917	387.6	5.4	10891	284.3	5.6	10201	183.5	3.6
1984	20406	509.3	7.3	21453	394.3	5.4	10826	280.1	5.6	10484	187.4	3.6
1985	21223	524.8	7.4	22139	406.0	5.5	11029	283.5	5.6	10607	188.5	3.6
1986	21177	522.0	7.4	22274	406.4	5.5	11282	287.9	5.6	10718	189.2	3.6
1987	21409	523.4	7.3	22831	412.4	5.5	11161	283.2	5.5	10766	187.9	3.6
1988	21451	520.4	7.3	22924	411.5	5.4	11354	284.4	5.5	10866	188.1	3.6
1989	21962	526.9	7.3	22793	406.0	5.4	11092	275.4	5.4	10772	185.5	3.6
1990	23214	546.2	7.3	23780	419.9	5.4	11055	271.6	5.3	10814	183.0	3.5
1991	23955	557.8	7.3	23771	417.6	5.4	11310	275.2	5.3	10988	184.9	3.5
1992	25590	588.1	7.4	24029	418.2	5.4	11202	270.9	5.2	11309	188.2	3.5
1993	25237	571.1	7.2	24408	421.6	5.4	11105	265.1	5.1	11184	184.0	3.5
1994	25052	564.4	7.2	24763	425.9	5.4	11657	275.7	5.2	11393	186.0	3.5
1995	24977	555.3	7.0	25487	438.5	5.5	11416	268.1	5.0	11334	184.2	3.4
1996	26356	582.4	7.1	26101	447.5	5.5	11351	263.1	4.9	11249	182.1	3.4
1997	26949	589.4	7.1	26674	453.8	5.5	11167	255.5	4.8	11035	176.5	3.3
1998	27393	590.6	7.1	27521	466.3	5.6	10852	242.6	4.6	10899	173.0	3.3
1999	27995	599.3	7.1	27771	465.2	5.5	11153	247.9	4.7	11187	175.8	3.3
2000	28883	613.5	7.1	28338	469.7	5.5	11071	244.3	4.6	11205	173.6	3.3
2001	28744	603.6	7.0	28232	467.2	5.5	10955	238.8	4.5	11069	170.8	3.2
2002	29847	618.4	7.1	28165	462.3	5.5	10917	234.5	4.4	11095	170.1	3.2
2003	29317	599.0	6.9	28531	464.5	5.5	10525	222.7	4.3	10951	165.9	3.2
2004	29466	595.3	6.9	28683	465.4	5.5	10588	220.1	4.2	10847	163.3	3.1
2005	29897	597.2	6.8	29010	468.3	5.5	10423	214.7	4.2	10622	158.6	3.1
2006	31169	615.2	6.9	29453	475.0	5.5	10459	212.3	4.1	10667	159.1	3.1
2007	31703	614.4	6.8	29594	474.9	5.5	10510	209.0	4.0	10417	154.0	3.0
2008	31842	608.3	6.8	30473	485.4	5.6	10628	208.0	4.0	10499	153.2	3.0
2009	31791	595.7	6.7	30294	477.3	5.5	10530	203.0	3.9	10182	147.8	3.0
2010	30782	567.8	6.5	29986	470.4	5.5	10261	194.4	3.8	10468	149.9	3.0
2011	31720	575.6	6.5	30574	475.9	5.5	10302	191.3	3.8	10347	146.3	2.9
2012	30495	545.7	6.3	30316	469.5	5.5	10718	197.2	3.8	10383	145.8	2.9
2013	30505	537.1	6.2	31006	478.1	5.6	10538	188.5	3.7	10169	140.9	2.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Oral cavity and pharynx

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	672	16.7	1.3	318	6.2	0.7	285	7.1	0.9	111	2.1	0.4
1977	653	15.9	1.3	319	6.1	0.7	270	6.8	0.9	126	2.4	0.4
1978	663	16.3	1.3	345	6.5	0.7	327	8.3	1.0	143	2.7	0.4
1979	670	16.0	1.3	381	7.1	0.7	257	6.2	0.8	156	2.9	0.5
1980	663	15.9	1.3	352	6.5	0.7	258	6.3	0.8	143	2.6	0.4
1981	723	17.1	1.3	335	6.1	0.7	271	6.5	0.8	124	2.2	0.4
1982	733	17.6	1.3	372	6.9	0.7	276	6.9	0.9	141	2.5	0.4
1983	683	16.0	1.3	358	6.6	0.7	261	6.2	0.8	137	2.4	0.4
1984	711	16.9	1.3	345	6.3	0.7	249	6.1	0.8	140	2.5	0.4
1985	698	16.5	1.3	360	6.4	0.7	240	5.8	0.8	137	2.5	0.4
1986	683	16.2	1.3	353	6.3	0.7	266	6.3	0.8	143	2.5	0.4
1987	651	15.3	1.2	384	6.8	0.7	241	5.9	0.8	141	2.4	0.4
1988	650	15.1	1.2	382	6.9	0.7	261	6.5	0.8	138	2.3	0.4
1989	662	15.0	1.2	339	6.0	0.7	201	4.8	0.7	131	2.3	0.4
1990	622	14.1	1.1	362	6.5	0.7	220	5.0	0.7	125	2.1	0.4
1991	645	14.6	1.2	367	6.5	0.7	209	4.9	0.7	134	2.3	0.4
1992	637	14.3	1.1	368	6.4	0.7	182	4.1	0.6	120	2.0	0.4
1993	661	14.9	1.2	340	5.8	0.6	175	3.9	0.6	121	2.0	0.4
1994	649	14.4	1.1	347	6.0	0.6	211	4.7	0.7	94	1.6	0.3
1995	570	12.5	1.0	364	6.2	0.6	214	4.9	0.7	108	1.8	0.3
1996	668	14.3	1.1	361	6.3	0.7	184	4.2	0.6	99	1.6	0.3
1997	703	15.1	1.1	382	6.5	0.7	177	3.9	0.6	95	1.5	0.3
1998	664	14.1	1.1	388	6.6	0.7	190	4.1	0.6	113	1.8	0.3
1999	692	14.5	1.1	334	5.6	0.6	167	3.5	0.5	94	1.5	0.3
2000	676	14.1	1.1	363	6.0	0.6	147	3.2	0.5	89	1.4	0.3
2001	687	13.9	1.0	389	6.4	0.6	172	3.6	0.5	102	1.5	0.3
2002	725	14.5	1.1	382	6.2	0.6	158	3.3	0.5	93	1.4	0.3
2003	733	14.6	1.1	378	6.1	0.6	149	3.0	0.5	91	1.4	0.3
2004	793	15.6	1.1	374	6.0	0.6	179	3.5	0.5	93	1.4	0.3
2005	717	13.8	1.0	400	6.5	0.7	161	3.1	0.5	80	1.2	0.3
2006	848	16.1	1.1	365	5.9	0.6	161	3.1	0.5	88	1.3	0.3
2007	898	16.7	1.1	401	6.5	0.6	159	3.0	0.5	75	1.1	0.3
2008	862	15.8	1.1	410	6.5	0.6	182	3.3	0.5	68	1.0	0.2
2009	908	16.2	1.1	431	6.7	0.6	164	3.1	0.5	83	1.1	0.2
2010	944	16.7	1.1	445	6.9	0.7	170	3.1	0.5	92	1.3	0.3
2011	971	16.7	1.1	424	6.4	0.6	168	3.0	0.5	90	1.2	0.3
2012	967	16.7	1.1	444	6.6	0.6	197	3.5	0.5	82	1.1	0.2
2013	1022	17.6	1.1	433	6.4	0.6	191	3.2	0.5	86	1.1	0.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Esophagus

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	277	7.2	0.9	85	1.7	0.4	252	6.6	0.9	78	1.5	0.3
1977	255	6.7	0.9	107	2.0	0.4	232	6.0	0.8	91	1.7	0.4
1978	250	6.6	0.9	102	1.9	0.4	271	7.1	0.9	100	1.9	0.4
1979	256	6.3	0.8	118	2.2	0.4	225	5.8	0.8	100	1.8	0.4
1980	260	6.6	0.8	115	2.1	0.4	228	5.7	0.8	111	2.0	0.4
1981	258	6.4	0.8	117	2.1	0.4	262	6.7	0.9	93	1.7	0.3
1982	250	6.0	0.8	125	2.2	0.4	238	5.9	0.8	116	2.0	0.4
1983	280	6.8	0.8	115	2.0	0.4	276	7.1	0.9	92	1.6	0.3
1984	276	6.7	0.8	126	2.2	0.4	256	6.4	0.8	115	2.0	0.4
1985	280	6.8	0.8	93	1.6	0.3	265	6.4	0.8	94	1.6	0.3
1986	299	7.1	0.8	126	2.2	0.4	264	6.2	0.8	106	1.8	0.4
1987	277	6.6	0.8	123	2.1	0.4	284	6.9	0.8	107	1.8	0.4
1988	299	7.0	0.8	118	2.0	0.4	317	7.5	0.9	107	1.8	0.3
1989	299	6.9	0.8	114	1.9	0.4	301	7.1	0.8	104	1.7	0.3
1990	320	7.2	0.8	144	2.4	0.4	287	6.4	0.8	108	1.8	0.3
1991	309	6.8	0.8	140	2.3	0.4	311	7.1	0.8	120	2.0	0.4
1992	350	7.7	0.8	163	2.6	0.4	334	7.5	0.8	126	2.1	0.4
1993	357	7.9	0.8	143	2.3	0.4	335	7.5	0.8	120	1.9	0.3
1994	364	8.0	0.8	143	2.3	0.4	325	7.2	0.8	135	2.2	0.4
1995	349	7.8	0.8	137	2.2	0.4	378	8.5	0.9	124	1.9	0.3
1996	372	8.1	0.8	158	2.5	0.4	337	7.5	0.8	137	2.2	0.4
1997	339	7.4	0.8	142	2.3	0.4	368	8.1	0.8	109	1.7	0.3
1998	411	8.8	0.9	134	2.2	0.4	366	7.8	0.8	119	1.9	0.3
1999	385	8.2	0.8	148	2.3	0.4	367	7.9	0.8	132	2.0	0.4
2000	421	8.9	0.9	157	2.4	0.4	366	7.8	0.8	138	2.1	0.4
2001	400	8.2	0.8	134	2.1	0.4	412	8.6	0.8	127	1.9	0.3
2002	441	9.0	0.8	143	2.2	0.4	384	7.9	0.8	123	1.8	0.3
2003	463	9.3	0.9	143	2.1	0.3	421	8.6	0.8	146	2.1	0.4
2004	479	9.6	0.9	147	2.2	0.4	452	9.0	0.8	134	1.9	0.3
2005	525	10.4	0.9	156	2.4	0.4	471	9.5	0.9	133	1.9	0.3
2006	549	10.7	0.9	153	2.3	0.4	436	8.6	0.8	117	1.8	0.3
2007	512	9.8	0.9	151	2.3	0.4	434	8.4	0.8	151	2.2	0.4
2008	548	10.4	0.9	154	2.3	0.4	460	8.7	0.8	115	1.7	0.3
2009	576	10.6	0.9	160	2.3	0.4	455	8.4	0.8	112	1.6	0.3
2010	513	9.3	0.8	164	2.3	0.4	470	8.7	0.8	115	1.6	0.3
2011	484	8.5	0.8	166	2.4	0.4	477	8.6	0.8	146	2.0	0.3
2012	512	9.0	0.8	127	1.7	0.3	458	8.2	0.8	126	1.7	0.3
2013	527	9.0	0.8	147	2.0	0.3	457	7.9	0.7	132	1.8	0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Stomach

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	554	15.3	1.3	407	8.1	0.8	418	11.9	1.2	344	6.8	0.7
1977	563	15.7	1.4	392	7.6	0.8	443	12.6	1.3	306	6.0	0.7
1978	530	14.6	1.3	412	7.9	0.8	390	10.9	1.1	308	5.9	0.7
1979	572	15.5	1.3	391	7.3	0.7	420	11.5	1.2	308	5.7	0.6
1980	554	14.7	1.3	363	6.8	0.7	419	11.5	1.2	312	5.8	0.6
1981	609	16.0	1.3	404	7.4	0.7	413	11.1	1.1	288	5.3	0.6
1982	591	15.5	1.3	396	7.1	0.7	433	11.6	1.2	269	4.8	0.6
1983	589	15.3	1.3	383	6.8	0.7	412	10.9	1.1	275	4.9	0.6
1984	586	14.8	1.3	376	6.6	0.7	426	10.7	1.1	299	5.2	0.6
1985	617	15.7	1.3	365	6.4	0.7	378	10.0	1.1	272	4.7	0.6
1986	605	15.4	1.3	357	6.1	0.6	425	10.8	1.1	275	4.7	0.6
1987	581	14.6	1.2	387	6.6	0.7	404	10.5	1.1	272	4.6	0.6
1988	533	13.2	1.2	403	6.9	0.7	364	9.2	1.0	282	4.7	0.6
1989	595	14.7	1.2	299	4.9	0.6	398	10.2	1.1	278	4.6	0.5
1990	632	15.1	1.2	364	6.0	0.6	390	9.7	1.0	229	3.7	0.5
1991	561	13.4	1.2	333	5.5	0.6	385	9.6	1.0	211	3.4	0.5
1992	584	13.9	1.2	340	5.4	0.6	388	9.6	1.0	228	3.6	0.5
1993	586	13.8	1.1	376	5.9	0.6	362	8.9	0.9	236	3.7	0.5
1994	559	13.0	1.1	353	5.6	0.6	363	8.7	0.9	251	3.9	0.5
1995	550	12.6	1.1	352	5.6	0.6	330	7.9	0.9	251	3.9	0.5
1996	552	12.7	1.1	346	5.3	0.6	320	7.5	0.8	231	3.5	0.5
1997	564	12.6	1.1	360	5.6	0.6	292	6.7	0.8	229	3.5	0.5
1998	597	13.0	1.1	333	5.1	0.6	287	6.4	0.8	218	3.3	0.4
1999	560	12.3	1.0	324	5.0	0.6	300	6.6	0.8	194	2.9	0.4
2000	559	12.2	1.0	342	5.0	0.5	283	6.4	0.7	215	3.2	0.4
2001	552	11.8	1.0	309	4.8	0.5	271	5.9	0.7	202	3.0	0.4
2002	524	11.1	1.0	331	5.0	0.5	255	5.5	0.7	186	2.7	0.4
2003	564	11.7	1.0	337	5.1	0.6	243	5.2	0.7	182	2.7	0.4
2004	521	10.8	0.9	337	5.1	0.6	277	5.9	0.7	193	2.8	0.4
2005	494	9.9	0.9	304	4.6	0.5	240	5.0	0.6	179	2.6	0.4
2006	492	9.8	0.9	307	4.7	0.5	212	4.2	0.6	159	2.3	0.4
2007	540	10.5	0.9	294	4.5	0.5	218	4.3	0.6	149	2.2	0.4
2008	505	9.8	0.9	269	4.0	0.5	240	4.7	0.6	154	2.1	0.3
2009	483	9.3	0.8	264	3.8	0.5	216	4.2	0.6	133	1.9	0.3
2010	526	9.8	0.9	284	4.2	0.5	204	3.9	0.5	136	1.9	0.3
2011	554	10.2	0.9	325	4.8	0.5	194	3.6	0.5	126	1.7	0.3
2012	526	9.4	0.8	311	4.6	0.5	212	3.9	0.5	143	2.0	0.3
2013	575	10.2	0.9	328	4.8	0.5	214	3.8	0.5	140	1.9	0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Colorectal

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	2896	80.0	3.1	3001	58.7	2.1	1433	41.2	2.3	1573	30.9	1.5
1977	2937	80.0	3.1	3098	59.4	2.1	1438	41.3	2.3	1577	30.5	1.5
1978	2993	81.0	3.1	3141	59.6	2.1	1550	43.6	2.3	1648	31.4	1.5
1979	2960	79.0	3.0	3227	60.1	2.1	1510	42.7	2.3	1642	30.6	1.5
1980	3220	84.0	3.1	3201	58.6	2.1	1511	41.5	2.2	1604	29.5	1.5
1981	3311	87.4	3.1	3283	59.4	2.1	1543	42.1	2.2	1621	29.3	1.4
1982	3258	84.0	3.0	3304	59.0	2.0	1565	42.8	2.2	1616	28.8	1.4
1983	3358	86.4	3.1	3353	59.4	2.0	1526	41.3	2.2	1599	28.0	1.4
1984	3317	84.3	3.0	3474	61.0	2.1	1609	42.9	2.2	1651	28.9	1.4
1985	3543	89.3	3.1	3501	61.1	2.0	1541	40.9	2.2	1633	28.2	1.4
1986	3343	84.2	3.0	3436	59.4	2.0	1566	41.4	2.2	1602	27.4	1.4
1987	3340	83.7	3.0	3386	58.1	2.0	1578	41.3	2.1	1576	26.6	1.3
1988	3196	78.6	2.9	3284	55.5	1.9	1570	40.1	2.1	1565	26.3	1.3
1989	3229	79.1	2.8	3204	53.9	1.9	1406	35.7	2.0	1502	24.9	1.3
1990	3175	76.3	2.8	3243	53.6	1.9	1437	36.2	2.0	1381	22.6	1.2
1991	3177	75.6	2.7	3201	52.7	1.9	1383	34.3	1.9	1479	23.9	1.2
1992	3171	74.8	2.7	3102	50.5	1.8	1320	32.4	1.8	1405	22.5	1.2
1993	3092	71.7	2.6	3265	52.6	1.8	1356	33.0	1.8	1432	22.5	1.2
1994	3202	74.4	2.6	3278	52.7	1.8	1349	32.8	1.8	1491	23.2	1.2
1995	3179	72.9	2.6	3226	51.3	1.8	1332	32.1	1.8	1457	22.5	1.2
1996	3183	72.5	2.6	3113	49.3	1.8	1316	31.1	1.7	1333	20.4	1.1
1997	3301	73.7	2.6	3329	52.5	1.8	1248	28.9	1.6	1268	19.3	1.1
1998	3443	75.7	2.6	3511	55.1	1.8	1217	27.5	1.6	1301	19.6	1.1
1999	3496	76.4	2.6	3518	54.6	1.8	1216	27.4	1.6	1262	18.7	1.0
2000	3476	74.9	2.5	3528	54.3	1.8	1256	28.0	1.6	1241	18.0	1.0
2001	3312	70.5	2.4	3361	51.5	1.8	1138	25.1	1.5	1273	18.6	1.0
2002	3293	68.9	2.4	3312	50.4	1.7	1174	25.4	1.5	1238	17.9	1.0
2003	3069	63.4	2.3	3243	49.0	1.7	1083	23.1	1.4	1115	16.0	1.0
2004	2968	60.5	2.2	3214	48.5	1.7	1030	21.4	1.3	1116	15.7	0.9
2005	2942	59.4	2.2	2922	43.8	1.6	962	20.1	1.3	1012	14.2	0.9
2006	2768	55.2	2.1	2826	42.2	1.6	921	19.0	1.2	1009	14.1	0.9
2007	2822	55.2	2.1	2897	43.3	1.6	944	18.9	1.2	978	13.6	0.9
2008	2736	52.9	2.0	2815	41.6	1.6	911	18.0	1.2	979	13.4	0.9
2009	2582	49.2	1.9	2607	38.1	1.5	896	17.5	1.2	925	12.8	0.9
2010	2583	48.4	1.9	2701	39.4	1.5	878	16.7	1.1	922	12.4	0.8
2011	2503	46.5	1.9	2590	37.9	1.5	873	16.1	1.1	902	12.0	0.8
2012	2497	45.4	1.8	2363	34.6	1.5	839	15.5	1.1	896	11.9	0.8
2013	2476	44.2	1.8	2350	34.4	1.5	907	16.4	1.1	807	10.6	0.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Colon excluding rectum

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	1906	53.6	2.6	2203	43.1	1.8	1057	30.4	1.9	1298	25.5	1.4
1977	1940	54.3	2.6	2272	43.7	1.8	1132	32.6	2.0	1310	25.4	1.4
1978	1991	54.9	2.6	2324	44.1	1.8	1186	33.5	2.0	1377	26.2	1.4
1979	1996	54.4	2.5	2393	44.7	1.8	1211	34.2	2.0	1386	25.8	1.4
1980	2202	58.2	2.6	2359	43.3	1.8	1212	33.4	2.0	1330	24.4	1.3
1981	2274	60.7	2.6	2451	44.5	1.8	1279	35.1	2.0	1384	25.1	1.3
1982	2166	56.8	2.5	2450	43.7	1.7	1269	34.7	2.0	1356	24.2	1.3
1983	2232	58.7	2.6	2494	44.2	1.8	1251	34.0	2.0	1367	24.0	1.3
1984	2232	57.6	2.5	2649	46.4	1.8	1321	35.2	2.0	1414	24.7	1.3
1985	2417	61.6	2.6	2608	45.4	1.8	1269	33.6	1.9	1396	24.1	1.3
1986	2298	58.7	2.5	2599	44.9	1.7	1305	34.6	2.0	1384	23.7	1.3
1987	2274	57.8	2.5	2496	42.6	1.7	1322	34.6	2.0	1339	22.7	1.2
1988	2154	53.6	2.4	2486	41.9	1.7	1330	34.0	1.9	1365	22.9	1.2
1989	2197	55.0	2.4	2344	39.3	1.6	1162	29.5	1.8	1306	21.7	1.2
1990	2140	52.3	2.3	2394	39.5	1.6	1252	31.7	1.8	1195	19.5	1.1
1991	2199	53.0	2.3	2393	39.2	1.6	1151	28.6	1.7	1265	20.4	1.1
1992	2133	50.9	2.2	2345	38.0	1.6	1088	26.9	1.7	1224	19.6	1.1
1993	2115	49.7	2.2	2473	39.6	1.6	1155	28.3	1.7	1235	19.4	1.1
1994	2239	52.6	2.2	2463	39.4	1.6	1132	27.7	1.7	1305	20.3	1.1
1995	2217	51.2	2.2	2422	38.2	1.5	1096	26.6	1.6	1259	19.4	1.1
1996	2163	50.0	2.2	2335	36.6	1.5	1101	26.1	1.6	1144	17.5	1.0
1997	2302	51.7	2.1	2578	40.3	1.6	1051	24.4	1.5	1124	17.0	1.0
1998	2420	53.8	2.2	2641	41.2	1.6	1013	22.9	1.4	1118	16.7	1.0
1999	2397	52.7	2.1	2658	40.8	1.6	997	22.5	1.4	1076	15.9	1.0
2000	2398	52.2	2.1	2691	41.0	1.6	1039	23.3	1.4	1067	15.4	0.9
2001	2300	49.3	2.0	2539	38.5	1.5	945	20.9	1.3	1095	15.9	1.0
2002	2286	48.2	2.0	2494	37.7	1.5	954	20.7	1.3	1037	14.9	0.9
2003	2126	44.3	1.9	2452	36.7	1.5	894	19.1	1.3	966	13.8	0.9
2004	2016	41.5	1.8	2421	36.1	1.5	833	17.4	1.2	940	13.1	0.9
2005	2057	41.8	1.8	2218	32.9	1.4	771	16.2	1.2	852	11.9	0.8
2006	1963	39.5	1.8	2153	31.6	1.4	750	15.5	1.1	852	11.8	0.8
2007	1980	39.1	1.7	2208	32.5	1.4	772	15.6	1.1	846	11.7	0.8
2008	1882	36.6	1.7	2146	31.5	1.4	728	14.4	1.1	812	11.1	0.8
2009	1734	33.2	1.6	1965	28.3	1.3	718	14.1	1.0	780	10.8	0.8
2010	1760	33.4	1.6	2036	29.2	1.3	691	13.2	1.0	775	10.3	0.8
2011	1681	31.5	1.5	1949	28.0	1.3	686	12.7	1.0	766	10.1	0.7
2012	1687	30.9	1.5	1806	26.0	1.2	660	12.2	0.9	766	10.2	0.8
2013	1656	29.7	1.5	1730	24.7	1.2	712	13.0	1.0	676	8.8	0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Rectum & rectosigmoid

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	990	26.5	1.7	798	15.6	1.1	376	10.8	1.2	275	5.4	0.6
1977	997	25.6	1.7	826	15.8	1.1	306	8.7	1.0	267	5.1	0.6
1978	1002	26.1	1.7	817	15.5	1.1	364	10.2	1.1	271	5.2	0.6
1979	964	24.5	1.6	834	15.4	1.1	299	8.6	1.0	256	4.8	0.6
1980	1018	25.8	1.7	842	15.3	1.0	299	8.2	1.0	274	5.1	0.6
1981	1037	26.7	1.7	832	14.9	1.0	264	7.0	0.9	237	4.2	0.5
1982	1092	27.2	1.7	854	15.3	1.0	296	8.1	1.0	260	4.6	0.6
1983	1126	27.7	1.7	859	15.1	1.0	275	7.3	0.9	232	4.1	0.5
1984	1085	26.7	1.7	825	14.6	1.0	288	7.7	0.9	237	4.2	0.5
1985	1126	27.7	1.7	893	15.7	1.0	272	7.3	0.9	237	4.1	0.5
1986	1045	25.5	1.6	837	14.5	1.0	261	6.7	0.9	218	3.8	0.5
1987	1066	25.9	1.6	890	15.5	1.0	256	6.7	0.9	237	4.0	0.5
1988	1042	25.0	1.6	798	13.7	1.0	240	6.1	0.8	200	3.4	0.5
1989	1032	24.1	1.5	860	14.6	1.0	244	6.2	0.8	196	3.3	0.5
1990	1035	23.9	1.5	849	14.1	1.0	185	4.5	0.7	186	3.1	0.4
1991	978	22.6	1.5	808	13.6	1.0	232	5.7	0.8	214	3.5	0.5
1992	1038	23.8	1.5	757	12.4	0.9	232	5.5	0.7	181	2.9	0.4
1993	977	22.1	1.4	792	13.0	0.9	201	4.7	0.7	197	3.1	0.4
1994	963	21.8	1.4	815	13.3	0.9	217	5.1	0.7	186	2.9	0.4
1995	962	21.7	1.4	804	13.2	0.9	236	5.5	0.7	198	3.1	0.4
1996	1020	22.5	1.4	778	12.6	0.9	215	5.0	0.7	189	2.9	0.4
1997	999	21.9	1.4	751	12.1	0.9	197	4.6	0.7	144	2.2	0.4
1998	1023	21.9	1.4	870	13.9	0.9	204	4.5	0.6	183	2.8	0.4
1999	1099	23.7	1.4	860	13.8	0.9	219	4.9	0.7	186	2.8	0.4
2000	1078	22.7	1.4	837	13.4	0.9	217	4.7	0.6	174	2.6	0.4
2001	1012	21.2	1.3	822	13.0	0.9	193	4.2	0.6	178	2.6	0.4
2002	1007	20.7	1.3	818	12.8	0.9	220	4.7	0.6	201	3.0	0.4
2003	943	19.1	1.2	791	12.3	0.9	189	4.0	0.6	149	2.2	0.4
2004	952	19.0	1.2	793	12.4	0.9	197	4.0	0.6	176	2.5	0.4
2005	885	17.6	1.2	704	10.9	0.8	191	3.9	0.6	160	2.3	0.4
2006	805	15.7	1.1	673	10.6	0.8	171	3.5	0.5	157	2.3	0.4
2007	842	16.1	1.1	689	10.8	0.8	172	3.4	0.5	132	1.9	0.3
2008	854	16.3	1.1	669	10.1	0.8	183	3.6	0.5	167	2.3	0.4
2009	848	16.0	1.1	642	9.8	0.8	178	3.4	0.5	145	2.0	0.3
2010	823	15.0	1.0	665	10.2	0.8	187	3.5	0.5	147	2.1	0.4
2011	822	14.9	1.0	641	9.9	0.8	187	3.4	0.5	136	1.9	0.3
2012	810	14.5	1.0	557	8.6	0.7	179	3.3	0.5	130	1.8	0.3
2013	820	14.5	1.0	620	9.6	0.8	195	3.5	0.5	131	1.8	0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Liver / intrahepatic bile duct

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	138	3.6	0.6	97	1.9	0.4	107	2.9	0.6	69	1.3	0.3
1977	186	4.6	0.7	92	1.7	0.4	139	3.4	0.6	98	1.9	0.4
1978	152	3.9	0.6	113	2.1	0.4	138	3.5	0.6	98	1.8	0.4
1979	132	3.3	0.6	120	2.3	0.4	108	2.7	0.5	91	1.7	0.4
1980	149	4.0	0.7	99	1.8	0.4	114	3.0	0.6	103	1.9	0.4
1981	148	3.9	0.7	106	1.9	0.4	131	3.5	0.6	105	1.9	0.4
1982	153	3.9	0.6	101	1.8	0.4	140	3.7	0.6	80	1.4	0.3
1983	186	4.5	0.7	90	1.7	0.4	153	3.9	0.6	78	1.4	0.3
1984	152	3.8	0.6	113	2.0	0.4	136	3.4	0.6	97	1.7	0.3
1985	161	4.1	0.7	95	1.7	0.3	149	3.8	0.6	90	1.6	0.3
1986	170	4.1	0.6	96	1.7	0.3	156	3.8	0.6	97	1.7	0.3
1987	176	4.2	0.7	105	1.9	0.4	168	4.2	0.7	113	2.0	0.4
1988	212	5.1	0.7	118	2.0	0.4	179	4.3	0.7	102	1.7	0.3
1989	196	4.5	0.7	115	1.9	0.4	156	3.7	0.6	101	1.7	0.3
1990	222	5.2	0.7	126	2.1	0.4	194	4.8	0.7	118	1.9	0.4
1991	209	4.7	0.7	134	2.2	0.4	188	4.3	0.6	122	2.0	0.4
1992	227	5.5	0.7	163	2.7	0.4	195	4.7	0.7	129	2.1	0.4
1993	258	6.0	0.8	154	2.5	0.4	240	5.6	0.7	161	2.5	0.4
1994	245	5.7	0.7	133	2.1	0.4	210	5.0	0.7	153	2.5	0.4
1995	285	6.4	0.8	157	2.5	0.4	251	5.8	0.7	147	2.3	0.4
1996	244	5.5	0.7	155	2.5	0.4	224	5.1	0.7	156	2.4	0.4
1997	300	6.5	0.7	163	2.6	0.4	243	5.4	0.7	143	2.2	0.4
1998	334	7.2	0.8	163	2.6	0.4	219	4.8	0.6	150	2.3	0.4
1999	319	6.9	0.8	169	2.7	0.4	250	5.4	0.7	163	2.5	0.4
2000	308	6.5	0.7	178	2.8	0.4	246	5.3	0.7	160	2.4	0.4
2001	358	7.5	0.8	148	2.3	0.4	269	5.7	0.7	145	2.2	0.4
2002	367	7.6	0.8	156	2.4	0.4	305	6.4	0.7	164	2.4	0.4
2003	340	6.8	0.7	191	3.0	0.4	261	5.4	0.7	150	2.2	0.4
2004	416	8.2	0.8	162	2.5	0.4	288	5.8	0.7	178	2.6	0.4
2005	424	8.2	0.8	185	2.9	0.4	313	6.2	0.7	168	2.5	0.4
2006	454	8.7	0.8	166	2.6	0.4	307	6.1	0.7	198	2.9	0.4
2007	487	9.2	0.8	200	3.1	0.4	322	6.2	0.7	162	2.3	0.4
2008	519	9.5	0.8	218	3.3	0.4	373	7.0	0.7	208	3.0	0.4
2009	509	9.2	0.8	218	3.2	0.4	357	6.6	0.7	197	2.8	0.4
2010	523	9.3	0.8	167	2.5	0.4	348	6.3	0.7	206	3.0	0.4
2011	531	9.0	0.8	220	3.3	0.4	376	6.6	0.7	233	3.3	0.4
2012	596	10.0	0.8	215	3.1	0.4	423	7.4	0.7	227	3.2	0.4
2013	636	10.6	0.8	254	3.7	0.5	466	8.0	0.7	219	3.0	0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Pancreas

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	567	15.2	1.3	486	9.5	0.9	521	14.1	1.3	502	9.7	0.9
1977	588	16.4	1.4	522	10.0	0.9	568	15.5	1.4	517	10.0	0.9
1978	575	15.2	1.3	541	10.2	0.9	565	15.0	1.3	521	9.8	0.8
1979	644	17.1	1.4	575	10.7	0.9	615	16.5	1.4	558	10.3	0.9
1980	576	14.8	1.3	593	10.9	0.9	554	14.6	1.3	546	10.0	0.8
1981	572	14.8	1.3	564	10.3	0.9	554	14.4	1.3	548	10.0	0.8
1982	551	14.2	1.3	576	10.2	0.8	533	13.8	1.2	504	8.9	0.8
1983	536	13.6	1.2	551	9.8	0.8	549	13.7	1.2	580	10.2	0.8
1984	558	14.4	1.3	570	9.9	0.8	524	13.6	1.2	565	9.8	0.8
1985	580	14.3	1.2	651	11.3	0.9	528	13.4	1.2	598	10.4	0.8
1986	574	14.2	1.2	591	10.2	0.8	560	13.7	1.2	573	9.8	0.8
1987	571	14.3	1.2	635	10.7	0.8	554	14.0	1.2	578	9.8	0.8
1988	607	15.0	1.2	638	10.7	0.8	568	14.2	1.2	612	10.2	0.8
1989	574	13.8	1.2	592	9.9	0.8	547	13.4	1.2	542	9.1	0.8
1990	592	14.3	1.2	640	10.6	0.8	544	13.1	1.1	600	9.9	0.8
1991	620	14.6	1.2	672	11.0	0.8	541	13.0	1.1	607	9.9	0.8
1992	590	14.0	1.2	651	10.6	0.8	536	12.9	1.1	617	9.9	0.8
1993	618	14.3	1.2	637	10.3	0.8	581	13.5	1.1	636	10.2	0.8
1994	669	15.3	1.2	668	10.5	0.8	624	14.5	1.2	632	9.9	0.8
1995	608	13.6	1.1	720	11.4	0.8	578	12.9	1.1	668	10.4	0.8
1996	648	14.7	1.1	626	9.9	0.8	576	13.2	1.1	616	9.6	0.8
1997	645	14.3	1.1	688	10.8	0.8	641	14.5	1.1	646	9.9	0.8
1998	678	14.9	1.1	734	11.5	0.8	614	13.7	1.1	635	9.8	0.8
1999	652	14.1	1.1	733	11.3	0.8	632	13.7	1.1	662	10.1	0.8
2000	654	14.0	1.1	737	11.3	0.8	594	13.0	1.1	683	10.3	0.8
2001	647	13.7	1.1	772	11.7	0.8	585	12.6	1.0	692	10.4	0.8
2002	700	14.6	1.1	790	12.0	0.8	633	13.2	1.0	686	10.2	0.8
2003	725	14.9	1.1	776	11.7	0.8	591	12.3	1.0	699	10.3	0.8
2004	716	14.5	1.1	762	11.4	0.8	648	13.2	1.0	661	9.8	0.8
2005	748	14.9	1.1	769	11.3	0.8	631	12.7	1.0	711	10.3	0.8
2006	732	14.5	1.1	805	11.9	0.8	661	13.1	1.0	683	9.8	0.8
2007	785	15.3	1.1	822	12.0	0.8	660	12.9	1.0	684	9.9	0.8
2008	825	15.9	1.1	852	12.4	0.8	667	12.9	1.0	721	10.2	0.8
2009	870	16.3	1.1	807	11.6	0.8	698	13.2	1.0	697	9.9	0.8
2010	825	15.4	1.1	893	12.7	0.9	702	12.9	1.0	739	10.2	0.8
2011	893	16.2	1.1	857	12.1	0.8	688	12.6	1.0	723	9.9	0.7
2012	911	16.5	1.1	856	12.0	0.8	771	14.0	1.0	708	9.7	0.7
2013	902	15.9	1.1	897	12.3	0.8	757	13.4	1.0	755	10.2	0.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Larynx

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	432	10.4	1.0	72	1.3	0.3	145	3.8	0.7	30	0.6	0.2
1977	415	10.0	1.0	93	1.7	0.4	141	3.6	0.6	28	0.5	0.2
1978	456	10.9	1.0	90	1.7	0.3	145	3.7	0.6	40	0.7	0.2
1979	441	10.2	1.0	122	2.2	0.4	170	4.2	0.7	39	0.7	0.2
1980	470	11.1	1.1	105	2.0	0.4	148	3.7	0.6	36	0.7	0.2
1981	512	11.9	1.1	115	2.1	0.4	155	3.9	0.6	29	0.5	0.2
1982	452	10.5	1.0	109	2.0	0.4	152	3.6	0.6	39	0.7	0.2
1983	454	10.5	1.0	113	2.1	0.4	139	3.4	0.6	44	0.8	0.2
1984	447	10.2	1.0	123	2.3	0.4	126	3.1	0.6	36	0.6	0.2
1985	465	10.5	1.0	124	2.3	0.4	137	3.2	0.6	41	0.7	0.2
1986	441	10.2	1.0	129	2.3	0.4	132	3.2	0.6	35	0.6	0.2
1987	488	11.2	1.0	122	2.1	0.4	139	3.4	0.6	37	0.6	0.2
1988	414	9.3	0.9	131	2.3	0.4	136	3.3	0.6	36	0.6	0.2
1989	471	10.7	1.0	135	2.4	0.4	118	2.7	0.5	53	0.9	0.3
1990	456	9.9	0.9	154	2.7	0.4	121	2.9	0.5	32	0.5	0.2
1991	440	9.7	0.9	111	1.9	0.4	123	2.7	0.5	36	0.6	0.2
1992	430	9.5	0.9	135	2.4	0.4	157	3.6	0.6	48	0.8	0.2
1993	392	8.7	0.9	150	2.6	0.4	131	3.0	0.5	37	0.6	0.2
1994	391	8.6	0.9	126	2.2	0.4	127	2.9	0.5	42	0.7	0.2
1995	410	8.9	0.9	131	2.3	0.4	127	2.9	0.5	38	0.6	0.2
1996	373	8.0	0.8	109	1.9	0.4	122	2.7	0.5	43	0.7	0.2
1997	381	8.2	0.8	111	2.0	0.4	104	2.3	0.5	37	0.6	0.2
1998	336	7.1	0.8	130	2.2	0.4	112	2.4	0.5	39	0.6	0.2
1999	413	8.6	0.8	107	1.8	0.3	116	2.5	0.5	39	0.6	0.2
2000	394	8.2	0.8	107	1.8	0.3	113	2.4	0.5	27	0.4	0.2
2001	366	7.5	0.8	104	1.7	0.3	115	2.4	0.5	23	0.4	0.1
2002	360	7.4	0.8	102	1.7	0.3	115	2.4	0.4	31	0.5	0.2
2003	349	6.9	0.7	100	1.6	0.3	96	2.0	0.4	33	0.5	0.2
2004	359	7.0	0.7	92	1.5	0.3	95	2.0	0.4	33	0.5	0.2
2005	396	7.6	0.8	113	1.8	0.3	103	2.1	0.4	41	0.7	0.2
2006	364	7.1	0.7	122	2.0	0.4	103	2.0	0.4	36	0.6	0.2
2007	354	6.7	0.7	88	1.3	0.3	106	2.0	0.4	20	0.3	0.1
2008	389	7.2	0.7	104	1.6	0.3	87	1.6	0.3	25	0.4	0.1
2009	356	6.6	0.7	101	1.6	0.3	93	1.8	0.4	35	0.5	0.2
2010	343	6.2	0.7	95	1.5	0.3	106	2.0	0.4	36	0.5	0.2
2011	352	6.0	0.6	83	1.3	0.3	96	1.7	0.4	36	0.5	0.2
2012	354	6.2	0.7	99	1.4	0.3	91	1.6	0.3	26	0.4	0.1
2013	368	6.1	0.6	95	1.4	0.3	124	2.1	0.4	29	0.4	0.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Lung and bronchus

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	3852	98.7	3.3	1474	28.4	1.5	3163	82.6	3.0	1126	21.7	1.3
1977	3810	96.5	3.2	1527	29.2	1.5	3304	85.3	3.1	1229	23.4	1.3
1978	3919	98.6	3.2	1544	29.0	1.5	3282	83.7	3.0	1270	23.7	1.3
1979	3898	97.6	3.2	1720	32.2	1.6	3191	81.0	2.9	1325	24.5	1.3
1980	4164	102.3	3.3	1792	32.9	1.6	3484	86.9	3.0	1408	25.7	1.4
1981	4096	101.0	3.2	1915	35.3	1.6	3483	87.0	3.0	1456	26.6	1.4
1982	4197	102.4	3.2	2043	37.2	1.6	3425	85.4	3.0	1649	29.9	1.5
1983	4277	103.3	3.2	2256	40.6	1.7	3531	87.2	3.0	1722	30.8	1.5
1984	4210	100.7	3.2	2280	41.2	1.7	3460	83.9	2.9	1789	31.9	1.5
1985	4345	103.7	3.2	2469	44.4	1.8	3649	88.7	3.0	1943	34.4	1.6
1986	4266	102.1	3.2	2494	44.6	1.8	3612	87.6	3.0	2013	35.7	1.6
1987	4318	102.0	3.2	2632	46.7	1.8	3586	86.5	3.0	2017	35.2	1.6
1988	4354	102.6	3.2	2779	48.7	1.8	3619	86.4	2.9	2173	37.8	1.6
1989	4360	101.2	3.1	2900	50.6	1.9	3613	85.0	2.9	2238	38.8	1.6
1990	4417	102.0	3.1	2891	49.7	1.8	3619	84.5	2.9	2369	40.2	1.6
1991	4340	99.6	3.1	3077	52.7	1.9	3720	86.5	2.9	2460	41.7	1.7
1992	4291	97.3	3.0	3170	53.8	1.9	3616	83.7	2.8	2610	43.5	1.7
1993	4178	93.9	2.9	3271	55.2	1.9	3488	79.6	2.7	2550	42.5	1.7
1994	4321	96.7	2.9	3387	57.0	1.9	3656	83.1	2.8	2584	42.5	1.7
1995	4297	94.9	2.9	3399	56.8	1.9	3598	81.1	2.7	2648	43.6	1.7
1996	4404	97.0	2.9	3455	57.7	1.9	3501	78.3	2.6	2636	43.4	1.7
1997	4305	94.1	2.8	3594	59.7	2.0	3358	74.5	2.6	2679	43.5	1.7
1998	4312	92.6	2.8	3708	61.3	2.0	3400	73.8	2.5	2652	42.8	1.6
1999	4351	93.0	2.8	3748	61.3	2.0	3295	71.3	2.5	2724	43.7	1.7
2000	4362	92.3	2.8	3891	62.9	2.0	3362	71.9	2.4	2818	44.6	1.7
2001	4151	87.4	2.7	3918	63.2	2.0	3301	70.6	2.4	2776	43.8	1.6
2002	4350	90.4	2.7	3931	62.8	2.0	3294	69.2	2.4	2905	45.6	1.7
2003	4376	89.6	2.7	3990	63.3	2.0	3206	66.4	2.3	2807	43.5	1.6
2004	4237	85.7	2.6	4151	65.2	2.0	3089	63.1	2.2	2940	45.2	1.7
2005	4266	85.6	2.6	4140	64.6	2.0	3118	63.4	2.2	2867	43.7	1.6
2006	4237	84.2	2.6	4417	69.3	2.1	3176	63.7	2.2	2851	43.7	1.6
2007	4306	84.2	2.5	4286	66.2	2.0	3098	60.9	2.2	2925	44.2	1.6
2008	4421	85.3	2.5	4452	68.1	2.0	3069	59.5	2.1	2917	43.4	1.6
2009	4437	84.0	2.5	4289	64.6	2.0	3037	57.9	2.1	2781	41.3	1.6
2010	4240	78.7	2.4	4387	66.1	2.0	2928	54.9	2.0	2844	41.9	1.6
2011	4323	79.0	2.4	4438	65.6	2.0	2921	54.0	2.0	2819	40.8	1.5
2012	4249	76.5	2.3	4371	63.7	1.9	3071	55.9	2.0	2847	40.8	1.5
2013	4193	73.9	2.3	4462	64.3	1.9	2863	50.5	1.9	2792	39.2	1.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Melanoma of the skin

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	271	6.6	0.8	260	5.4	0.7	130	3.2	0.6	103	2.1	0.4
1977	371	9.0	1.0	313	6.5	0.7	159	4.0	0.7	87	1.8	0.4
1978	334	8.2	0.9	318	6.7	0.8	139	3.5	0.6	80	1.6	0.4
1979	360	8.8	0.9	308	6.3	0.7	133	3.3	0.6	103	2.1	0.4
1980	455	11.1	1.1	364	7.4	0.8	152	3.6	0.6	99	2.0	0.4
1981	418	10.1	1.0	382	7.7	0.8	132	3.2	0.6	91	1.8	0.4
1982	405	9.6	1.0	428	8.5	0.8	157	3.8	0.6	96	1.8	0.4
1983	502	12.0	1.1	408	8.3	0.8	136	3.2	0.6	108	2.2	0.4
1984	444	10.3	1.0	369	7.4	0.8	144	3.5	0.6	110	2.1	0.4
1985	488	11.4	1.1	421	8.4	0.8	151	3.7	0.6	101	1.9	0.4
1986	522	12.3	1.1	449	9.0	0.9	158	3.8	0.6	115	2.2	0.4
1987	496	11.5	1.0	414	8.1	0.8	179	4.2	0.6	118	2.3	0.4
1988	504	11.8	1.1	372	7.1	0.7	187	4.4	0.7	104	1.9	0.4
1989	525	12.1	1.1	407	7.8	0.8	199	4.8	0.7	113	2.1	0.4
1990	517	11.5	1.0	442	8.3	0.8	181	4.2	0.6	116	2.2	0.4
1991	515	11.6	1.0	373	6.9	0.7	187	4.4	0.6	111	2.0	0.4
1992	536	12.0	1.0	384	7.0	0.7	172	3.9	0.6	123	2.2	0.4
1993	529	11.6	1.0	398	7.2	0.7	183	4.2	0.6	107	1.9	0.4
1994	589	12.9	1.1	420	7.7	0.7	185	4.2	0.6	109	1.9	0.4
1995	661	14.4	1.1	445	8.1	0.8	183	4.2	0.6	128	2.2	0.4
1996	681	14.8	1.1	506	9.2	0.8	197	4.3	0.6	122	2.1	0.4
1997	742	15.9	1.2	527	9.6	0.8	204	4.6	0.6	99	1.6	0.3
1998	772	16.4	1.2	579	10.6	0.9	187	4.0	0.6	105	1.8	0.3
1999	810	17.1	1.2	588	10.5	0.9	167	3.6	0.6	117	2.0	0.4
2000	888	18.6	1.2	664	11.9	0.9	196	4.1	0.6	108	1.8	0.3
2001	966	20.1	1.3	743	13.2	1.0	229	4.8	0.6	127	2.1	0.4
2002	980	20.1	1.3	716	12.6	0.9	168	3.5	0.5	145	2.4	0.4
2003	1026	20.9	1.3	733	12.7	0.9	191	3.9	0.6	121	1.9	0.3
2004	1201	24.1	1.4	887	15.5	1.0	197	4.0	0.6	125	2.0	0.4
2005	1342	26.9	1.5	952	16.7	1.1	209	4.2	0.6	126	2.0	0.4
2006	1425	28.2	1.5	1033	18.0	1.1	221	4.4	0.6	119	1.8	0.3
2007	1377	27.0	1.4	1006	17.6	1.1	235	4.6	0.6	92	1.4	0.3
2008	1507	29.3	1.5	1092	19.0	1.2	228	4.4	0.6	117	1.8	0.3
2009	1636	31.2	1.5	1136	19.6	1.2	206	4.0	0.6	144	2.2	0.4
2010	1524	28.9	1.5	1137	19.3	1.2	212	4.0	0.5	135	2.1	0.4
2011	1708	31.6	1.5	1217	20.0	1.2	241	4.5	0.6	127	1.9	0.3
2012	1617	29.5	1.5	1229	20.7	1.2	250	4.7	0.6	144	2.0	0.3
2013	1769	31.7	1.5	1309	22.2	1.3	252	4.7	0.6	127	1.8	0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Female breast

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	5020	102.0	2.9	1957	38.9	1.7
1977	5075	102.2	2.9	2012	39.7	1.8
1978	5183	103.4	2.9	2027	39.2	1.7
1979	5138	101.5	2.8	1997	38.6	1.7
1980	5302	103.7	2.9	1986	37.7	1.7
1981	5374	104.8	2.9	1994	37.5	1.7
1982	5468	105.3	2.9	2022	37.7	1.7
1983	5848	112.4	3.0	2059	38.2	1.7
1984	6010	114.4	3.0	2126	39.1	1.7
1985	6376	121.7	3.1	2162	39.8	1.7
1986	6548	123.8	3.1	2195	39.9	1.7
1987	6753	126.7	3.1	2196	39.4	1.7
1988	6884	128.0	3.1	2165	38.9	1.7
1989	6748	125.0	3.0	2158	38.7	1.7
1990	7144	131.2	3.1	2125	37.3	1.6
1991	7070	128.5	3.1	2054	36.0	1.6
1992	7103	128.5	3.0	2043	35.3	1.6
1993	7116	127.7	3.0	2026	34.5	1.5
1994	7195	128.9	3.0	2020	34.3	1.5
1995	7569	135.7	3.1	1933	32.8	1.5
1996	8035	143.2	3.2	1962	32.7	1.5
1997	8185	144.5	3.2	1846	30.6	1.4
1998	8446	147.6	3.2	1864	30.4	1.4
1999	8362	144.8	3.1	1740	28.0	1.3
2000	8428	144.0	3.1	1759	28.0	1.3
2001	8296	141.0	3.1	1708	27.1	1.3
2002	8295	140.0	3.0	1684	26.3	1.3
2003	8245	137.8	3.0	1678	26.0	1.3
2004	8103	134.6	3.0	1535	23.6	1.2
2005	8233	135.5	3.0	1506	23.0	1.2
2006	8337	137.8	3.0	1444	22.0	1.2
2007	8341	136.2	3.0	1504	22.6	1.2
2008	8637	140.3	3.0	1435	21.4	1.1
2009	8845	141.4	3.0	1440	21.1	1.1
2010	8256	131.7	2.9	1493	21.5	1.1
2011	8690	138.0	3.0	1453	20.8	1.1
2012	8619	136.4	3.0	1395	19.8	1.1
2013	8847	139.2	3.0	1397	19.9	1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Cervix uteri

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	597	12.4	1.0	234	4.6	0.6
1977	518	10.6	0.9	191	3.7	0.5
1978	519	10.6	0.9	223	4.4	0.6
1979	507	10.4	0.9	215	4.3	0.6
1980	550	10.9	0.9	209	4.2	0.6
1981	456	9.0	0.8	217	4.1	0.6
1982	463	9.4	0.9	191	3.6	0.5
1983	477	9.6	0.9	167	3.2	0.5
1984	516	10.4	0.9	164	3.2	0.5
1985	478	9.5	0.9	161	3.1	0.5
1986	435	8.7	0.8	182	3.5	0.5
1987	499	9.7	0.9	181	3.5	0.5
1988	514	10.2	0.9	170	3.2	0.5
1989	483	9.5	0.9	151	2.8	0.5
1990	511	10.0	0.9	168	3.0	0.5
1991	496	9.5	0.8	141	2.6	0.4
1992	458	8.8	0.8	163	3.0	0.5
1993	486	9.2	0.8	144	2.6	0.4
1994	459	8.7	0.8	157	2.9	0.5
1995	552	10.5	0.9	117	2.1	0.4
1996	540	10.2	0.9	147	2.7	0.4
1997	485	9.2	0.8	155	2.7	0.4
1998	447	8.5	0.8	122	2.2	0.4
1999	441	8.3	0.8	144	2.5	0.4
2000	404	7.6	0.7	121	2.1	0.4
2001	432	8.0	0.8	129	2.2	0.4
2002	372	6.9	0.7	132	2.3	0.4
2003	412	7.7	0.8	113	1.9	0.4
2004	346	6.5	0.7	122	2.1	0.4
2005	395	7.5	0.8	120	2.0	0.4
2006	381	7.2	0.7	113	1.9	0.4
2007	392	7.3	0.7	118	2.0	0.4
2008	372	7.0	0.7	99	1.6	0.3
2009	402	7.6	0.8	91	1.5	0.3
2010	386	7.5	0.8	113	1.8	0.3
2011	313	5.8	0.7	110	1.8	0.4
2012	346	6.4	0.7	139	2.4	0.4
2013	362	6.9	0.8	113	1.9	0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Corpus uterus and uterus, not otherwise specified

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	1515	29.3	1.5	285	5.5	0.6
1977	1383	26.2	1.4	286	5.4	0.6
1978	1229	23.1	1.3	256	4.8	0.6
1979	1350	25.3	1.4	290	5.3	0.6
1980	1352	25.6	1.4	292	5.2	0.6
1981	1465	27.2	1.4	283	5.1	0.6
1982	1401	25.7	1.4	300	5.4	0.6
1983	1338	24.8	1.4	280	4.8	0.6
1984	1394	25.6	1.4	291	5.1	0.6
1985	1329	24.1	1.3	281	4.9	0.6
1986	1352	24.7	1.3	274	4.6	0.6
1987	1438	26.1	1.4	271	4.6	0.6
1988	1316	23.8	1.3	275	4.6	0.6
1989	1331	23.7	1.3	276	4.6	0.6
1990	1389	24.8	1.3	239	3.9	0.5
1991	1400	25.5	1.4	271	4.4	0.5
1992	1477	26.3	1.4	281	4.6	0.5
1993	1409	25.0	1.3	293	4.7	0.5
1994	1518	26.6	1.4	289	4.7	0.5
1995	1520	26.9	1.4	298	4.7	0.5
1996	1622	28.4	1.4	276	4.4	0.5
1997	1570	27.5	1.4	273	4.4	0.5
1998	1721	30.0	1.4	272	4.3	0.5
1999	1646	28.3	1.4	248	3.8	0.5
2000	1653	28.0	1.4	277	4.3	0.5
2001	1804	30.6	1.4	269	4.1	0.5
2002	1832	30.6	1.4	289	4.5	0.5
2003	1822	30.2	1.4	304	4.6	0.5
2004	1836	30.0	1.4	293	4.4	0.5
2005	1898	30.9	1.4	291	4.4	0.5
2006	1860	29.8	1.4	317	4.7	0.5
2007	1941	31.1	1.4	298	4.3	0.5
2008	2067	32.7	1.4	306	4.5	0.5
2009	1956	30.5	1.4	304	4.5	0.5
2010	1996	30.9	1.4	303	4.3	0.5
2011	2076	31.6	1.4	335	4.9	0.5
2012	2167	32.4	1.4	343	4.8	0.5
2013	2278	33.8	1.4	375	5.1	0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Ovary

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	870	17.3	1.2	652	12.7	1.0
1977	846	16.5	1.1	554	10.7	0.9
1978	837	16.2	1.1	596	11.2	0.9
1979	875	16.9	1.1	571	10.8	0.9
1980	872	16.7	1.1	583	10.9	0.9
1981	944	18.1	1.2	624	11.4	0.9
1982	879	16.7	1.1	619	11.3	0.9
1983	950	17.9	1.2	613	11.0	0.9
1984	901	16.9	1.1	601	10.9	0.9
1985	921	17.5	1.2	588	10.6	0.9
1986	945	17.9	1.2	567	10.2	0.9
1987	931	17.3	1.1	611	10.9	0.9
1988	946	17.6	1.1	618	10.9	0.9
1989	952	17.5	1.1	623	10.9	0.9
1990	994	18.1	1.1	632	10.8	0.9
1991	1002	18.2	1.2	655	11.2	0.9
1992	995	18.0	1.1	629	10.9	0.9
1993	994	17.6	1.1	624	10.5	0.8
1994	1016	18.0	1.1	618	10.4	0.8
1995	1016	17.9	1.1	640	10.7	0.8
1996	1043	18.3	1.1	600	9.9	0.8
1997	997	17.5	1.1	650	10.7	0.8
1998	925	16.2	1.1	573	9.4	0.8
1999	963	16.5	1.1	582	9.4	0.8
2000	998	16.8	1.1	633	10.1	0.8
2001	986	16.6	1.0	636	10.0	0.8
2002	951	15.9	1.0	604	9.5	0.8
2003	971	16.1	1.0	627	9.7	0.8
2004	926	15.2	1.0	609	9.4	0.8
2005	885	14.5	1.0	653	10.0	0.8
2006	934	15.2	1.0	627	9.7	0.8
2007	900	14.4	1.0	574	8.8	0.7
2008	902	14.4	1.0	615	9.1	0.7
2009	837	13.2	0.9	574	8.5	0.7
2010	882	13.9	0.9	557	8.3	0.7
2011	874	13.5	0.9	565	8.2	0.7
2012	822	12.8	0.9	544	7.8	0.7
2013	906	14.1	1.0	585	8.3	0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Prostate

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	2534	78.5	3.2	938	31.5	2.1
1977	2610	78.7	3.2	928	30.8	2.1
1978	2574	77.3	3.1	975	32.6	2.1
1979	2644	79.1	3.2	994	32.5	2.1
1980	2972	85.2	3.2	996	32.1	2.1
1981	3041	87.4	3.3	952	30.6	2.0
1982	3172	89.3	3.3	1011	31.7	2.1
1983	3353	92.8	3.3	1093	33.6	2.1
1984	3311	90.3	3.2	1134	34.8	2.1
1985	3502	94.0	3.3	1103	33.6	2.1
1986	3575	94.9	3.3	1188	34.9	2.1
1987	3920	102.6	3.4	1163	34.1	2.1
1988	4038	104.8	3.4	1184	34.2	2.0
1989	4246	108.2	3.4	1218	35.1	2.1
1990	5001	121.6	3.5	1234	34.8	2.0
1991	6035	143.1	3.7	1354	37.4	2.1
1992	7630	175.1	4.1	1302	35.9	2.0
1993	7314	164.9	3.9	1329	36.1	2.0
1994	6597	146.9	3.6	1359	36.3	2.0
1995	6609	144.5	3.6	1290	34.3	1.9
1996	7294	158.6	3.7	1293	33.4	1.9
1997	7650	164.9	3.7	1304	33.0	1.8
1998	7521	160.1	3.7	1149	28.4	1.7
1999	7956	168.0	3.7	1166	28.3	1.7
2000	8596	180.6	3.8	1148	28.0	1.6
2001	8662	179.6	3.8	1149	27.4	1.6
2002	9027	184.3	3.8	1068	25.0	1.5
2003	8437	170.3	3.7	1083	24.8	1.5
2004	8180	162.6	3.5	1038	23.4	1.4
2005	8276	162.0	3.5	961	21.2	1.3
2006	9331	180.5	3.7	942	20.4	1.3
2007	9446	178.1	3.6	992	20.8	1.3
2008	9128	168.3	3.5	940	19.3	1.2
2009	8917	160.2	3.4	872	17.7	1.2
2010	8399	147.7	3.2	945	18.6	1.2
2011	8634	148.9	3.2	913	17.7	1.2
2012	7358	123.7	2.9	882	16.9	1.1
2013	7022	115.7	2.8	838	15.5	1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Testis

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	172	3.6	0.6	32	0.6	0.2
1977	169	3.6	0.6	28	0.5	0.2
1978	194	4.2	0.6	34	0.7	0.3
1979	221	4.8	0.6	31	0.7	0.3
1980	228	4.7	0.6	28	0.6	0.2
1981	218	4.5	0.6	28	0.5	0.2
1982	213	4.4	0.6	25	0.5	0.2
1983	234	4.8	0.6	23	0.5	0.2
1984	286	5.9	0.7	17	0.4	0.2
1985	255	5.2	0.7	20	0.4	0.2
1986	246	4.9	0.6	25	0.5	0.2
1987	274	5.5	0.7	20	0.4	0.2
1988	260	5.1	0.6	16	0.4	0.2
1989	261	5.1	0.6	21	0.4	0.2
1990	289	5.7	0.7	16	0.3	0.2
1991	268	5.3	0.6	13	0.2	0.1
1992	284	5.6	0.7	19	0.4	0.2
1993	302	6.1	0.7	23	0.5	0.2
1994	294	6.0	0.7	12	0.2	0.1
1995	310	6.4	0.7	15	0.3	0.2
1996	302	6.3	0.7	21	0.4	0.2
1997	325	6.8	0.7	6	0.1	0.1
1998	329	7.0	0.8	18	0.4	0.2
1999	324	6.9	0.8	13	0.3	0.1
2000	294	6.3	0.7	13	0.3	0.2
2001	315	6.8	0.8	13	0.3	0.2
2002	299	6.5	0.7	22	0.5	0.2
2003	339	7.5	0.8	11	0.2	0.1
2004	318	7.1	0.8	11	0.2	0.1
2005	340	7.6	0.8	7	0.1	0.1
2006	312	6.9	0.8	12	0.2	0.1
2007	338	7.7	0.8	6	0.1	0.1
2008	369	8.5	0.9	10	0.2	0.1
2009	333	7.6	0.8	7	0.1	0.1
2010	344	7.8	0.8	14	0.3	0.2
2011	337	7.6	0.8	9	0.2	0.1
2012	359	8.2	0.9	11	0.2	0.1
2013	313	7.3	0.8	7	0.1	0.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Urinary bladder (incl. in situ)

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	1398	39.1	2.2	498	9.7	0.9	424	13.3	1.3	161	3.2	0.5
1977	1437	39.1	2.1	523	10.2	0.9	410	12.7	1.3	173	3.4	0.5
1978	1496	40.8	2.2	558	10.8	0.9	419	12.9	1.3	176	3.4	0.5
1979	1569	42.4	2.2	509	9.5	0.8	389	11.7	1.2	190	3.6	0.5
1980	1614	42.6	2.2	577	10.6	0.9	392	11.6	1.2	163	3.0	0.5
1981	1595	41.8	2.2	550	10.1	0.9	436	13.0	1.3	179	3.3	0.5
1982	1691	43.7	2.2	620	11.2	0.9	397	11.5	1.2	167	3.0	0.5
1983	1667	43.1	2.2	564	10.0	0.8	409	12.1	1.2	151	2.6	0.4
1984	1633	41.5	2.1	621	11.0	0.9	389	11.6	1.2	181	3.1	0.5
1985	1746	44.1	2.2	640	11.3	0.9	384	10.9	1.2	178	3.1	0.5
1986	1820	45.0	2.2	651	11.4	0.9	403	11.9	1.2	171	2.9	0.4
1987	1673	41.0	2.1	652	11.2	0.9	356	10.0	1.1	174	2.9	0.4
1988	1699	41.7	2.1	635	10.9	0.9	374	10.1	1.1	184	3.0	0.4
1989	1753	42.6	2.1	643	10.8	0.8	388	10.7	1.1	153	2.5	0.4
1990	1792	43.0	2.1	709	12.0	0.9	361	9.9	1.1	189	3.0	0.4
1991	1764	41.7	2.0	668	11.1	0.9	377	10.0	1.1	162	2.6	0.4
1992	1768	41.7	2.0	687	11.4	0.9	338	8.9	1.0	228	3.6	0.5
1993	1830	42.2	2.0	776	12.7	0.9	374	9.8	1.0	183	2.8	0.4
1994	1773	40.8	1.9	692	11.3	0.9	383	9.7	1.0	189	2.8	0.4
1995	1855	42.3	2.0	781	12.7	0.9	363	8.9	0.9	184	2.8	0.4
1996	2098	47.5	2.1	763	12.4	0.9	401	10.0	1.0	200	3.0	0.4
1997	2153	48.2	2.1	835	13.4	0.9	418	10.0	1.0	206	3.1	0.4
1998	2261	49.8	2.1	828	13.4	0.9	409	9.9	1.0	170	2.5	0.4
1999	2262	49.3	2.1	864	13.7	0.9	407	9.6	1.0	183	2.6	0.4
2000	2261	48.8	2.0	875	13.6	0.9	415	9.4	0.9	186	2.7	0.4
2001	2303	49.3	2.0	837	13.0	0.9	432	9.9	0.9	181	2.5	0.4
2002	2465	51.9	2.1	839	13.0	0.9	407	9.1	0.9	202	2.9	0.4
2003	2480	51.3	2.0	881	13.6	0.9	381	8.4	0.9	182	2.5	0.4
2004	2564	52.8	2.1	903	13.9	0.9	419	9.0	0.9	186	2.6	0.4
2005	2594	52.8	2.0	921	13.9	0.9	383	8.1	0.8	182	2.5	0.4
2006	2539	51.1	2.0	889	13.5	0.9	394	8.3	0.8	182	2.5	0.4
2007	2523	49.8	2.0	896	13.6	0.9	458	9.4	0.9	182	2.5	0.4
2008	2669	52.0	2.0	883	13.0	0.9	446	9.0	0.8	203	2.7	0.4
2009	2689	51.4	2.0	891	13.0	0.9	485	9.7	0.9	156	2.0	0.3
2010	2531	47.7	1.9	817	11.8	0.8	467	9.0	0.8	199	2.6	0.4
2011	2683	49.8	1.9	914	13.4	0.9	437	8.3	0.8	180	2.3	0.3
2012	2731	50.0	1.9	964	13.6	0.9	443	8.3	0.8	198	2.5	0.4
2013	2749	49.8	1.9	941	13.2	0.9	465	8.6	0.8	178	2.3	0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Kidney and renal pelvis

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	402	10.2	1.0	241	4.7	0.6	223	5.7	0.8	134	2.6	0.4
1977	432	10.8	1.1	270	5.2	0.6	243	6.3	0.8	157	3.0	0.5
1978	429	10.8	1.1	270	5.2	0.6	217	5.5	0.8	137	2.6	0.4
1979	435	10.7	1.1	277	5.1	0.6	255	6.4	0.8	154	2.9	0.5
1980	436	10.7	1.1	287	5.4	0.6	240	6.1	0.8	166	3.0	0.5
1981	473	11.9	1.1	323	6.1	0.7	238	6.0	0.8	147	2.7	0.4
1982	508	12.0	1.1	328	6.0	0.7	234	5.8	0.8	156	2.8	0.4
1983	551	13.2	1.1	348	6.4	0.7	234	5.7	0.8	164	3.0	0.5
1984	560	13.5	1.2	331	6.1	0.7	249	6.4	0.8	151	2.7	0.4
1985	572	13.7	1.2	367	6.6	0.7	246	6.2	0.8	186	3.3	0.5
1986	592	14.1	1.2	361	6.7	0.7	269	6.5	0.8	180	3.2	0.5
1987	607	14.3	1.2	418	7.5	0.7	287	7.1	0.9	176	3.0	0.5
1988	594	13.9	1.2	401	7.1	0.7	253	6.1	0.8	193	3.3	0.5
1989	611	14.4	1.2	368	6.5	0.7	262	6.4	0.8	171	2.9	0.4
1990	689	16.0	1.2	409	7.1	0.7	253	5.9	0.8	164	2.7	0.4
1991	677	15.4	1.2	442	7.8	0.7	279	6.6	0.8	185	3.0	0.4
1992	687	15.4	1.2	454	8.0	0.7	257	6.1	0.8	168	2.8	0.4
1993	703	15.6	1.2	472	8.1	0.7	250	5.7	0.7	160	2.6	0.4
1994	694	15.4	1.2	472	8.2	0.7	290	6.8	0.8	177	2.9	0.4
1995	715	15.6	1.2	492	8.4	0.8	250	5.8	0.7	195	3.2	0.5
1996	792	17.2	1.2	493	8.4	0.7	275	6.2	0.7	198	3.2	0.5
1997	864	18.8	1.3	507	8.5	0.8	283	6.4	0.8	166	2.7	0.4
1998	882	18.7	1.2	555	9.4	0.8	290	6.3	0.7	170	2.6	0.4
1999	924	19.5	1.3	582	9.8	0.8	237	5.2	0.7	147	2.2	0.4
2000	898	18.7	1.2	610	10.1	0.8	269	5.9	0.7	178	2.7	0.4
2001	999	20.6	1.3	604	10.0	0.8	278	6.0	0.7	196	3.1	0.4
2002	1013	20.8	1.3	610	9.9	0.8	257	5.5	0.7	178	2.7	0.4
2003	1047	20.8	1.3	630	10.2	0.8	270	5.5	0.7	185	2.7	0.4
2004	1151	22.8	1.3	702	11.5	0.9	283	5.8	0.7	189	2.8	0.4
2005	1134	22.5	1.3	719	11.7	0.9	302	6.1	0.7	167	2.5	0.4
2006	1273	24.7	1.4	734	11.7	0.9	288	5.8	0.7	184	2.7	0.4
2007	1355	26.0	1.4	759	12.3	0.9	279	5.5	0.7	170	2.5	0.4
2008	1279	24.1	1.3	765	12.2	0.9	304	5.9	0.7	157	2.2	0.4
2009	1257	23.4	1.3	764	12.2	0.9	281	5.3	0.6	159	2.3	0.4
2010	1295	23.8	1.3	777	12.1	0.9	280	5.1	0.6	157	2.2	0.4
2011	1394	25.2	1.4	744	11.7	0.9	292	5.5	0.6	183	2.5	0.4
2012	1399	24.9	1.3	808	12.4	0.9	326	5.9	0.7	169	2.3	0.4
2013	1391	24.4	1.3	782	12.1	0.9	291	5.2	0.6	153	2.1	0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Brain and other nervous system

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	322	7.2	0.8	234	4.6	0.6	224	5.1	0.7	191	3.7	0.5
1977	323	7.2	0.8	247	4.7	0.6	239	5.4	0.7	189	3.6	0.5
1978	284	6.4	0.8	255	5.0	0.6	219	4.9	0.7	179	3.3	0.5
1979	354	8.0	0.9	263	5.0	0.6	229	5.1	0.7	175	3.3	0.5
1980	356	7.7	0.8	263	5.1	0.6	274	6.2	0.8	181	3.4	0.5
1981	341	7.8	0.9	270	5.1	0.6	238	5.5	0.7	183	3.4	0.5
1982	334	7.6	0.8	321	6.2	0.7	223	5.1	0.7	212	4.0	0.5
1983	315	7.1	0.8	314	6.0	0.7	225	5.1	0.7	200	3.6	0.5
1984	379	8.5	0.9	320	6.0	0.7	235	5.5	0.7	195	3.5	0.5
1985	353	7.9	0.9	306	5.8	0.7	206	4.8	0.7	207	3.9	0.5
1986	374	8.4	0.9	314	5.9	0.7	251	5.7	0.7	207	3.7	0.5
1987	389	8.7	0.9	349	6.5	0.7	234	5.3	0.7	224	4.0	0.5
1988	399	8.8	0.9	320	6.0	0.7	273	6.0	0.7	215	3.8	0.5
1989	407	8.8	0.9	302	5.7	0.7	254	5.7	0.7	203	3.6	0.5
1990	422	9.4	0.9	341	6.3	0.7	250	5.6	0.7	169	2.9	0.4
1991	394	8.8	0.9	324	6.0	0.7	262	6.0	0.7	183	3.4	0.5
1992	387	8.5	0.9	349	6.4	0.7	241	5.3	0.7	234	4.2	0.5
1993	409	8.8	0.9	324	5.9	0.7	248	5.4	0.7	207	3.6	0.5
1994	455	9.9	0.9	348	6.2	0.7	293	6.4	0.8	236	4.1	0.5
1995	423	9.0	0.9	429	7.8	0.7	253	5.5	0.7	214	3.7	0.5
1996	457	9.9	0.9	378	6.9	0.7	284	6.2	0.7	230	4.0	0.5
1997	443	9.4	0.9	349	6.3	0.7	272	5.8	0.7	218	3.8	0.5
1998	466	9.7	0.9	371	6.8	0.7	271	5.7	0.7	205	3.5	0.5
1999	416	8.8	0.8	364	6.5	0.7	269	5.6	0.7	214	3.7	0.5
2000	456	9.6	0.9	415	7.3	0.7	243	5.0	0.6	225	3.7	0.5
2001	442	9.2	0.9	370	6.5	0.7	242	5.0	0.6	220	3.7	0.5
2002	468	9.8	0.9	354	6.3	0.7	286	5.8	0.7	220	3.7	0.5
2003	435	8.9	0.8	411	7.5	0.7	256	5.2	0.6	201	3.4	0.5
2004	451	9.2	0.9	350	6.2	0.7	274	5.5	0.7	236	3.9	0.5
2005	482	9.9	0.9	353	6.3	0.7	278	5.6	0.7	229	3.8	0.5
2006	475	9.5	0.9	338	6.0	0.7	284	5.6	0.7	189	3.1	0.4
2007	436	8.8	0.8	364	6.5	0.7	294	5.7	0.7	191	3.1	0.4
2008	456	9.2	0.9	357	6.3	0.7	283	5.5	0.7	198	3.1	0.4
2009	445	8.8	0.8	383	6.8	0.7	298	5.6	0.7	240	3.8	0.5
2010	445	8.8	0.8	357	6.2	0.7	285	5.5	0.6	230	3.6	0.5
2011	450	8.7	0.8	349	6.2	0.7	259	4.8	0.6	213	3.2	0.4
2012	462	8.8	0.8	371	6.5	0.7	322	6.0	0.7	248	3.9	0.5
2013	473	8.9	0.8	331	5.7	0.6	295	5.1	0.6	236	3.6	0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Thyroid

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	99	2.4	0.5	198	4.2	0.6	20	0.6	0.3	32	0.6	0.2
1977	105	2.5	0.5	237	4.9	0.6	21	0.6	0.3	46	0.9	0.3
1978	94	2.3	0.5	232	4.7	0.6	20	0.6	0.3	38	0.7	0.2
1979	92	2.2	0.5	213	4.3	0.6	13	0.3	0.2	30	0.5	0.2
1980	105	2.4	0.5	229	4.5	0.6	33	0.9	0.3	36	0.7	0.2
1981	101	2.4	0.5	229	4.6	0.6	24	0.6	0.2	30	0.6	0.2
1982	95	2.2	0.4	214	4.4	0.6	26	0.7	0.3	25	0.5	0.2
1983	99	2.2	0.5	263	5.2	0.6	14	0.4	0.2	37	0.7	0.2
1984	112	2.6	0.5	276	5.5	0.7	24	0.7	0.3	40	0.7	0.2
1985	109	2.4	0.5	280	5.6	0.7	18	0.5	0.2	27	0.5	0.2
1986	119	2.7	0.5	284	5.6	0.7	24	0.6	0.3	28	0.5	0.2
1987	131	2.9	0.5	270	5.3	0.7	15	0.3	0.2	35	0.6	0.2
1988	102	2.3	0.5	290	5.8	0.7	18	0.4	0.2	30	0.5	0.2
1989	115	2.6	0.5	297	5.8	0.7	17	0.4	0.2	37	0.6	0.2
1990	129	2.9	0.5	285	5.5	0.7	21	0.5	0.2	25	0.4	0.2
1991	144	3.1	0.5	330	6.4	0.7	15	0.4	0.2	27	0.5	0.2
1992	134	2.9	0.5	341	6.5	0.7	16	0.4	0.2	39	0.6	0.2
1993	133	2.8	0.5	356	7.0	0.7	21	0.5	0.2	36	0.6	0.2
1994	139	3.0	0.5	403	7.9	0.8	23	0.6	0.2	27	0.4	0.2
1995	171	3.7	0.6	449	8.8	0.8	24	0.5	0.2	31	0.5	0.2
1996	184	3.9	0.6	498	9.8	0.9	27	0.6	0.2	54	0.9	0.2
1997	203	4.3	0.6	539	10.5	0.9	23	0.5	0.2	32	0.5	0.2
1998	212	4.4	0.6	535	10.4	0.9	17	0.4	0.2	38	0.6	0.2
1999	216	4.4	0.6	682	13.4	1.0	20	0.4	0.2	34	0.5	0.2
2000	265	5.5	0.7	765	15.0	1.1	17	0.4	0.2	31	0.5	0.2
2001	231	4.7	0.6	816	16.0	1.1	23	0.5	0.2	31	0.5	0.2
2002	282	5.7	0.7	799	15.5	1.1	18	0.4	0.2	29	0.5	0.2
2003	309	6.2	0.7	822	15.9	1.1	14	0.3	0.1	40	0.6	0.2
2004	300	5.9	0.7	940	18.1	1.2	20	0.4	0.2	27	0.4	0.1
2005	356	7.1	0.7	1038	20.2	1.3	24	0.5	0.2	22	0.3	0.1
2006	389	7.8	0.8	1132	21.9	1.3	25	0.5	0.2	30	0.4	0.1
2007	440	8.8	0.8	1213	23.2	1.3	26	0.5	0.2	28	0.4	0.2
2008	429	8.4	0.8	1405	27.2	1.5	26	0.5	0.2	32	0.5	0.2
2009	490	9.7	0.9	1411	27.2	1.5	26	0.5	0.2	39	0.5	0.2
2010	501	9.6	0.9	1446	28.0	1.5	25	0.5	0.2	34	0.4	0.2
2011	559	10.8	0.9	1504	29.1	1.5	39	0.8	0.2	35	0.5	0.2
2012	556	10.5	0.9	1504	29.4	1.6	28	0.5	0.2	38	0.5	0.2
2013	584	11.1	0.9	1511	29.5	1.6	28	0.5	0.2	36	0.5	0.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Hodgkin lymphoma

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	190	3.9	0.6	156	2.9	0.5	75	1.8	0.4	64	1.2	0.3
1977	182	3.9	0.6	146	2.8	0.5	79	1.9	0.4	58	1.1	0.3
1978	185	3.9	0.6	171	3.1	0.5	66	1.6	0.4	53	1.0	0.3
1979	213	4.6	0.6	164	3.0	0.5	60	1.4	0.4	48	0.9	0.3
1980	223	4.7	0.6	176	3.2	0.5	77	1.8	0.4	54	1.0	0.3
1981	199	4.1	0.6	180	3.4	0.5	65	1.5	0.4	47	0.9	0.3
1982	178	3.8	0.6	188	3.4	0.5	71	1.6	0.4	60	1.2	0.3
1983	197	4.2	0.6	153	2.9	0.5	52	1.1	0.3	38	0.7	0.2
1984	198	4.3	0.6	181	3.3	0.5	59	1.4	0.4	35	0.6	0.2
1985	201	4.2	0.6	192	3.6	0.5	54	1.2	0.3	43	0.8	0.2
1986	177	3.6	0.5	177	3.3	0.5	61	1.4	0.4	57	1.0	0.3
1987	179	3.7	0.6	167	3.1	0.5	62	1.3	0.3	38	0.7	0.2
1988	200	4.2	0.6	163	3.1	0.5	54	1.2	0.3	26	0.5	0.2
1989	206	4.3	0.6	169	3.1	0.5	63	1.4	0.4	30	0.6	0.2
1990	187	3.9	0.6	172	3.3	0.5	46	1.0	0.3	36	0.7	0.2
1991	189	3.9	0.6	164	3.1	0.5	36	0.8	0.3	31	0.6	0.2
1992	195	4.1	0.6	184	3.5	0.5	42	0.9	0.3	47	0.8	0.2
1993	169	3.5	0.5	181	3.5	0.5	36	0.8	0.3	32	0.6	0.2
1994	195	4.1	0.6	197	3.8	0.5	37	0.8	0.3	35	0.6	0.2
1995	182	3.8	0.6	150	3.0	0.5	33	0.7	0.2	36	0.6	0.2
1996	175	3.7	0.6	174	3.5	0.5	29	0.7	0.2	29	0.6	0.2
1997	175	3.7	0.6	144	3.0	0.5	35	0.7	0.2	24	0.4	0.2
1998	177	3.8	0.6	178	3.6	0.5	31	0.6	0.2	27	0.5	0.2
1999	192	4.1	0.6	147	3.0	0.5	29	0.6	0.2	33	0.6	0.2
2000	181	3.9	0.6	148	3.0	0.5	25	0.5	0.2	25	0.4	0.2
2001	187	4.0	0.6	170	3.5	0.5	32	0.7	0.2	29	0.5	0.2
2002	219	4.7	0.6	164	3.4	0.5	27	0.6	0.2	37	0.7	0.2
2003	188	3.9	0.6	145	2.9	0.5	21	0.4	0.2	25	0.5	0.2
2004	189	4.0	0.6	180	3.7	0.6	31	0.6	0.2	21	0.3	0.2
2005	200	4.2	0.6	173	3.6	0.5	20	0.4	0.2	27	0.5	0.2
2006	187	3.9	0.6	152	3.1	0.5	29	0.6	0.2	27	0.5	0.2
2007	180	3.7	0.6	161	3.3	0.5	25	0.5	0.2	27	0.4	0.2
2008	191	4.0	0.6	153	3.1	0.5	28	0.6	0.2	23	0.4	0.1
2009	187	3.9	0.6	158	3.3	0.5	20	0.4	0.2	20	0.3	0.2
2010	191	3.9	0.6	139	2.8	0.5	23	0.5	0.2	21	0.4	0.2
2011	188	3.9	0.6	150	3.1	0.5	26	0.5	0.2	11	0.2	0.1
2012	214	4.6	0.6	138	2.9	0.5	20	0.4	0.2	27	0.4	0.2
2013	166	3.4	0.5	153	3.1	0.5	14	0.3	0.2	19	0.3	0.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Non-Hodgkin lymphomas

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	530	13.3	1.2	492	9.7	0.9	302	7.8	0.9	270	5.2	0.6
1977	558	14.0	1.2	530	10.1	0.9	306	7.7	0.9	312	5.9	0.7
1978	639	15.6	1.3	564	10.7	0.9	312	8.2	1.0	306	5.8	0.7
1979	564	14.0	1.2	549	10.4	0.9	340	8.6	1.0	290	5.4	0.6
1980	620	15.0	1.2	582	10.9	0.9	314	7.8	0.9	332	6.2	0.7
1981	605	15.0	1.2	620	11.4	0.9	333	8.4	1.0	306	5.6	0.6
1982	650	15.8	1.3	599	11.0	0.9	338	8.7	1.0	315	5.6	0.6
1983	673	16.0	1.3	684	12.3	0.9	359	8.8	1.0	326	5.9	0.7
1984	736	18.2	1.4	670	12.0	0.9	374	9.6	1.0	317	5.6	0.6
1985	775	18.3	1.3	701	12.5	0.9	385	9.5	1.0	351	6.2	0.7
1986	833	19.9	1.4	716	12.8	1.0	428	10.7	1.1	355	6.1	0.6
1987	804	18.9	1.4	711	12.5	0.9	384	9.4	1.0	397	6.9	0.7
1988	832	19.5	1.4	800	14.4	1.0	398	9.8	1.0	394	6.8	0.7
1989	892	20.6	1.4	807	14.3	1.0	439	10.6	1.0	384	6.5	0.7
1990	954	21.9	1.4	839	14.5	1.0	434	10.5	1.0	421	7.0	0.7
1991	927	21.6	1.4	845	14.7	1.0	452	10.9	1.0	414	6.8	0.7
1992	918	20.8	1.4	840	14.4	1.0	459	11.0	1.0	452	7.4	0.7
1993	982	21.9	1.4	881	15.1	1.0	418	9.7	1.0	426	6.9	0.7
1994	1044	23.3	1.4	906	15.3	1.0	492	11.5	1.0	469	7.6	0.7
1995	1022	22.8	1.4	946	16.1	1.0	508	11.9	1.1	472	7.5	0.7
1996	1128	24.9	1.5	1019	17.2	1.1	508	11.7	1.0	482	7.7	0.7
1997	1131	24.6	1.5	1066	17.9	1.1	518	11.6	1.0	515	8.2	0.7
1998	1122	24.2	1.4	1118	18.7	1.1	458	10.1	0.9	494	7.7	0.7
1999	1130	24.2	1.4	1132	18.7	1.1	443	9.9	0.9	469	7.3	0.7
2000	1169	24.8	1.4	1116	18.2	1.1	469	10.3	0.9	488	7.5	0.7
2001	1159	24.4	1.4	1127	18.3	1.1	474	10.1	0.9	406	6.1	0.6
2002	1239	25.8	1.4	1113	18.1	1.1	478	10.2	0.9	407	6.1	0.6
2003	1249	25.6	1.4	1195	19.1	1.1	406	8.7	0.9	402	5.9	0.6
2004	1330	27.0	1.5	1171	18.9	1.1	383	8.0	0.8	407	5.9	0.6
2005	1374	27.6	1.5	1193	18.9	1.1	403	8.4	0.8	358	5.1	0.5
2006	1382	27.5	1.5	1298	20.4	1.1	430	8.8	0.8	419	5.9	0.6
2007	1439	28.3	1.5	1271	20.0	1.1	416	8.3	0.8	334	4.7	0.5
2008	1444	28.0	1.5	1279	20.1	1.1	423	8.4	0.8	394	5.7	0.6
2009	1483	28.3	1.5	1267	19.5	1.1	427	8.3	0.8	339	4.8	0.5
2010	1456	27.5	1.4	1295	19.5	1.1	414	7.9	0.8	378	5.2	0.5
2011	1495	27.9	1.4	1222	18.8	1.1	411	7.7	0.8	367	5.1	0.5
2012	1488	27.4	1.4	1259	18.8	1.1	413	7.8	0.8	336	4.5	0.5
2013	1500	27.2	1.4	1265	18.9	1.1	383	7.0	0.7	322	4.3	0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Myeloma

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	200	5.4	0.8	196	3.8	0.5	136	3.8	0.7	122	2.4	0.4
1977	192	5.2	0.8	179	3.4	0.5	122	3.1	0.6	126	2.4	0.4
1978	179	4.8	0.7	211	4.0	0.5	128	3.4	0.6	123	2.3	0.4
1979	187	5.0	0.8	231	4.3	0.6	131	3.7	0.7	127	2.4	0.4
1980	212	5.6	0.8	198	3.6	0.5	150	4.1	0.7	161	2.9	0.5
1981	211	5.5	0.8	239	4.4	0.6	132	3.5	0.6	156	2.8	0.4
1982	236	6.0	0.8	208	3.6	0.5	142	3.8	0.7	144	2.5	0.4
1983	237	6.3	0.9	197	3.5	0.5	155	4.2	0.7	143	2.5	0.4
1984	256	6.4	0.8	225	4.0	0.5	164	4.1	0.7	166	2.9	0.4
1985	242	6.1	0.8	240	4.2	0.5	175	4.5	0.7	168	2.8	0.4
1986	244	6.1	0.8	200	3.5	0.5	169	4.4	0.7	174	2.9	0.4
1987	249	6.2	0.8	255	4.3	0.5	169	4.3	0.7	171	2.9	0.4
1988	251	6.2	0.8	224	3.8	0.5	199	5.2	0.7	162	2.7	0.4
1989	241	6.1	0.8	276	4.7	0.6	164	4.1	0.6	156	2.6	0.4
1990	282	6.7	0.8	262	4.4	0.5	163	4.1	0.7	159	2.6	0.4
1991	270	6.5	0.8	242	4.0	0.5	172	4.2	0.6	180	2.9	0.4
1992	283	6.7	0.8	275	4.5	0.5	185	4.6	0.7	168	2.6	0.4
1993	286	6.5	0.8	273	4.5	0.5	189	4.5	0.7	196	3.1	0.4
1994	290	6.7	0.8	267	4.3	0.5	183	4.4	0.7	228	3.6	0.5
1995	313	7.3	0.8	276	4.5	0.5	211	4.9	0.7	193	3.0	0.4
1996	299	6.8	0.8	280	4.5	0.5	198	4.8	0.7	187	2.9	0.4
1997	288	6.4	0.7	307	5.0	0.6	183	4.3	0.6	192	3.0	0.4
1998	337	7.3	0.8	325	5.2	0.6	192	4.3	0.6	195	3.0	0.4
1999	361	7.9	0.8	320	5.0	0.6	213	4.7	0.6	192	2.9	0.4
2000	361	7.7	0.8	315	4.9	0.6	218	4.8	0.7	192	2.9	0.4
2001	374	8.0	0.8	267	4.1	0.5	191	4.1	0.6	158	2.3	0.4
2002	402	8.4	0.8	309	4.7	0.5	218	4.8	0.6	176	2.6	0.4
2003	373	7.7	0.8	292	4.5	0.5	215	4.6	0.6	158	2.3	0.4
2004	397	8.0	0.8	315	4.8	0.5	200	4.2	0.6	164	2.4	0.4
2005	370	7.4	0.8	307	4.6	0.5	198	4.1	0.6	179	2.6	0.4
2006	412	8.1	0.8	310	4.6	0.5	199	4.0	0.6	161	2.4	0.4
2007	424	8.3	0.8	356	5.4	0.6	193	3.8	0.5	181	2.6	0.4
2008	420	8.0	0.8	357	5.3	0.6	200	4.0	0.6	158	2.2	0.4
2009	439	8.3	0.8	383	5.5	0.6	203	3.9	0.5	177	2.5	0.4
2010	452	8.3	0.8	368	5.4	0.6	189	3.6	0.5	197	2.7	0.4
2011	481	8.9	0.8	363	5.2	0.6	197	3.6	0.5	166	2.2	0.3
2012	463	8.2	0.8	375	5.4	0.6	198	3.6	0.5	196	2.6	0.4
2013	498	8.7	0.8	375	5.4	0.6	181	3.3	0.5	163	2.2	0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Whites
Leukemias

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	673	18.2	1.5	510	10.0	0.9	415	11.7	1.2	312	6.1	0.7
1977	617	16.3	1.4	497	9.7	0.9	420	11.2	1.2	365	7.2	0.7
1978	636	17.0	1.4	518	10.2	0.9	386	10.3	1.1	354	6.8	0.7
1979	719	18.4	1.4	545	10.3	0.9	421	11.3	1.1	358	6.9	0.7
1980	691	18.0	1.4	573	10.9	0.9	446	11.7	1.1	374	7.0	0.7
1981	657	17.0	1.4	511	9.6	0.8	444	11.9	1.2	354	6.7	0.7
1982	679	17.6	1.4	524	9.9	0.9	464	12.5	1.2	357	6.7	0.7
1983	676	16.9	1.3	507	9.3	0.8	414	11.1	1.1	386	7.0	0.7
1984	684	17.0	1.3	594	10.8	0.9	364	9.2	1.0	389	7.1	0.7
1985	760	18.8	1.4	573	10.4	0.9	469	12.1	1.2	330	5.9	0.6
1986	692	17.3	1.4	550	10.0	0.8	451	11.5	1.1	338	6.0	0.6
1987	669	16.3	1.3	552	10.0	0.8	434	11.0	1.1	352	6.2	0.7
1988	693	16.8	1.3	541	9.7	0.8	461	11.7	1.1	339	5.9	0.6
1989	684	16.6	1.3	534	9.3	0.8	433	10.8	1.1	364	6.3	0.7
1990	748	17.7	1.3	615	10.8	0.9	405	9.9	1.0	385	6.6	0.7
1991	714	17.0	1.3	598	10.4	0.9	395	9.7	1.0	391	6.6	0.7
1992	766	18.1	1.3	579	10.0	0.8	464	11.3	1.1	378	6.3	0.6
1993	742	17.0	1.3	599	10.5	0.9	405	9.9	1.0	342	5.7	0.6
1994	814	18.6	1.3	641	10.8	0.9	476	11.3	1.0	363	5.9	0.6
1995	795	18.0	1.3	630	10.7	0.8	472	11.3	1.0	363	5.9	0.6
1996	870	19.3	1.3	626	10.5	0.8	496	11.4	1.0	399	6.5	0.7
1997	824	18.3	1.3	687	11.3	0.9	470	10.7	1.0	388	6.1	0.6
1998	879	19.1	1.3	634	10.5	0.8	454	10.1	0.9	374	5.9	0.6
1999	915	20.0	1.3	732	12.0	0.9	482	10.8	1.0	408	6.3	0.6
2000	999	21.7	1.4	749	12.0	0.9	506	11.2	1.0	387	6.0	0.6
2001	954	20.4	1.3	760	12.3	0.9	472	10.4	1.0	398	6.1	0.6
2002	928	19.6	1.3	693	11.2	0.9	483	10.4	0.9	383	5.8	0.6
2003	1002	21.0	1.3	764	12.3	0.9	486	10.5	0.9	360	5.4	0.6
2004	1016	21.0	1.3	804	12.9	0.9	506	10.6	0.9	400	6.0	0.6
2005	1017	20.7	1.3	822	13.0	0.9	500	10.5	0.9	373	5.5	0.6
2006	1062	21.6	1.3	765	12.1	0.9	494	10.3	0.9	392	5.7	0.6
2007	1092	21.7	1.3	785	12.6	0.9	487	9.9	0.9	366	5.4	0.6
2008	1037	20.3	1.3	769	12.3	0.9	512	10.2	0.9	360	5.2	0.6
2009	1153	22.3	1.3	793	12.5	0.9	546	10.9	0.9	359	5.3	0.6
2010	1186	22.7	1.3	854	13.3	0.9	473	9.3	0.8	354	5.1	0.6
2011	1237	23.5	1.3	915	14.1	1.0	504	9.5	0.8	379	5.4	0.6
2012	1199	22.4	1.3	875	13.6	1.0	519	9.9	0.9	366	5.1	0.5
2013	1241	22.9	1.3	874	13.6	1.0	495	9.2	0.8	368	5.1	0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
All Invasive Malignant Tumors

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	2777	491.0	21.3	2841	323.7	12.8	1833	330.6	17.8	1602	194.4	10.3
1977	2950	520.3	21.8	3014	331.3	12.7	1911	343.2	17.8	1587	179.9	9.5
1978	3130	538.1	21.8	3158	345.1	12.9	1979	338.2	17.1	1790	202.7	10.1
1979	3259	549.3	21.6	3258	344.2	12.6	2110	366.3	17.9	1794	198.3	9.8
1980	3561	583.1	21.8	3451	348.1	12.3	2142	362.0	17.5	1871	196.1	9.4
1981	3504	555.4	20.9	3354	330.2	11.7	2219	364.2	17.3	1948	197.6	9.2
1982	3557	549.0	20.2	3516	334.3	11.5	2241	357.8	16.7	1915	186.7	8.7
1983	3590	534.3	19.6	3615	329.5	11.2	2358	358.8	16.2	2020	190.0	8.6
1984	3857	553.8	19.6	3914	349.2	11.4	2282	344.9	15.9	2048	189.4	8.5
1985	3989	560.5	19.4	4018	349.3	11.2	2316	343.7	15.7	2027	182.8	8.2
1986	4180	579.7	19.6	4029	339.9	10.8	2348	342.9	15.5	2131	185.6	8.1
1987	4107	550.4	18.7	4126	338.7	10.6	2334	338.3	15.4	2096	177.6	7.8
1988	4100	548.8	18.7	4287	345.1	10.6	2331	332.0	15.0	2150	178.6	7.8
1989	4397	571.9	18.8	4290	336.9	10.3	2363	328.3	14.7	2114	171.1	7.5
1990	4482	577.7	18.8	4534	350.0	10.4	2340	321.8	14.6	2252	181.1	7.7
1991	4803	604.6	19.0	4808	364.4	10.5	2360	320.7	14.4	2195	171.9	7.4
1992	5087	625.0	19.1	4885	361.7	10.3	2221	300.4	13.9	2319	177.8	7.4
1993	5221	623.2	18.6	4770	346.4	10.0	2291	302.0	13.8	2228	167.6	7.1
1994	5423	631.9	18.4	5170	368.7	10.2	2521	324.3	14.0	2453	181.5	7.3
1995	5576	628.0	18.0	5044	352.6	9.9	2447	305.8	13.3	2526	183.3	7.2
1996	5786	634.9	17.8	5484	375.0	10.1	2434	297.5	12.9	2422	172.4	7.0
1997	5835	638.6	17.7	5536	373.5	10.0	2410	291.7	12.7	2425	169.4	6.8
1998	5838	613.9	17.0	5832	384.1	10.0	2342	272.4	12.0	2498	171.4	6.8
1999	6089	626.8	16.9	5723	368.3	9.6	2335	268.6	11.8	2549	170.1	6.7
2000	6338	634.3	16.7	5777	362.0	9.4	2379	265.5	11.5	2500	161.8	6.4
2001	6401	622.5	16.3	5788	355.5	9.2	2312	255.1	11.2	2611	165.4	6.4
2002	6395	608.4	15.9	5999	360.7	9.2	2368	250.9	10.9	2485	154.3	6.1
2003	6547	607.5	15.7	6161	363.5	9.1	2265	231.8	10.3	2462	149.5	6.0
2004	6666	599.7	15.3	6373	370.5	9.2	2319	234.6	10.3	2465	148.1	5.9
2005	6778	598.3	15.1	6469	371.2	9.1	2237	220.0	9.7	2544	149.8	5.9
2006	6994	606.9	15.1	6571	371.4	9.1	2152	210.2	9.5	2454	142.6	5.7
2007	7379	620.9	15.0	6592	368.1	9.0	2259	213.9	9.4	2558	146.3	5.7
2008	7299	597.1	14.5	6931	380.0	9.0	2185	203.0	9.1	2513	140.3	5.5
2009	7498	593.2	14.2	7031	378.8	9.0	2315	210.4	9.1	2448	134.1	5.4
2010	7490	578.1	13.8	7096	374.4	8.8	2345	209.6	9.0	2503	134.0	5.3
2011	8019	597.3	13.8	7623	391.8	8.9	2380	201.5	8.6	2731	143.5	5.4
2012	7415	539.2	12.9	7528	380.5	8.7	2376	195.6	8.3	2692	138.0	5.3
2013	7507	528.8	12.6	7743	382.8	8.6	2349	189.4	8.1	2581	128.2	5.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Oral cavity and pharynx

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	170	24.9	4.2	78	7.8	1.8	75	12.6	3.3	36	3.7	1.3
1977	181	26.1	4.1	57	5.7	1.5	100	13.9	2.9	27	3.0	1.3
1978	185	24.7	3.8	84	7.9	1.7	100	15.1	3.4	33	3.1	1.1
1979	203	28.3	4.2	68	6.7	1.7	109	15.4	3.2	39	3.7	1.2
1980	196	26.3	3.9	81	7.3	1.6	114	16.0	3.2	26	2.3	0.9
1981	220	27.2	3.8	84	7.7	1.7	85	11.3	2.6	35	3.2	1.1
1982	200	25.4	3.8	92	8.0	1.7	116	15.3	3.0	36	3.1	1.0
1983	208	26.3	3.9	77	6.8	1.6	110	14.7	3.0	35	3.2	1.1
1984	226	26.5	3.6	108	8.9	1.7	92	11.8	2.6	29	2.4	0.9
1985	222	26.6	3.7	103	8.6	1.7	82	10.2	2.4	34	2.9	1.0
1986	229	26.2	3.6	92	7.3	1.5	97	12.5	2.7	38	3.3	1.1
1987	240	26.8	3.6	110	8.6	1.6	77	9.3	2.2	28	2.2	0.8
1988	203	22.6	3.3	103	7.7	1.5	96	10.7	2.3	37	2.9	0.9
1989	226	24.1	3.3	106	7.7	1.5	94	10.3	2.2	40	3.0	0.9
1990	215	24.0	3.5	91	6.8	1.4	70	8.5	2.2	31	2.3	0.8
1991	282	31.4	3.9	113	8.1	1.5	101	11.0	2.2	41	3.2	1.0
1992	242	25.9	3.5	126	8.9	1.6	84	9.0	2.0	34	2.4	0.8
1993	230	23.5	3.2	120	8.4	1.5	73	7.9	1.9	30	2.1	0.8
1994	229	23.2	3.2	102	7.0	1.4	89	9.6	2.2	30	2.1	0.7
1995	224	21.6	3.0	79	5.4	1.2	101	10.2	2.1	38	2.7	0.9
1996	220	21.9	3.1	113	7.4	1.4	76	7.8	1.9	32	2.2	0.8
1997	220	20.3	2.8	106	7.0	1.4	66	7.1	1.9	23	1.6	0.7
1998	193	17.4	2.6	104	6.7	1.3	70	6.9	1.7	27	1.8	0.7
1999	180	16.1	2.5	88	5.6	1.2	57	5.7	1.6	34	2.2	0.7
2000	186	17.1	2.6	93	5.7	1.2	71	6.8	1.7	30	1.9	0.7
2001	182	15.5	2.4	69	4.1	1.0	60	5.5	1.5	26	1.6	0.6
2002	189	16.2	2.5	101	6.1	1.2	54	4.6	1.3	22	1.4	0.6
2003	180	14.4	2.2	88	5.2	1.1	60	5.3	1.4	19	1.2	0.5
2004	193	15.5	2.3	103	5.8	1.1	78	6.6	1.6	24	1.4	0.6
2005	186	15.3	2.3	82	4.6	1.0	60	5.3	1.4	23	1.3	0.5
2006	170	13.3	2.1	88	4.8	1.0	51	4.1	1.2	13	0.7	0.4
2007	203	15.6	2.3	103	5.8	1.1	61	5.1	1.4	21	1.2	0.5
2008	179	13.2	2.0	96	5.2	1.0	54	4.4	1.2	22	1.2	0.5
2009	179	13.0	2.0	99	5.2	1.0	49	3.7	1.1	22	1.2	0.5
2010	193	13.7	2.0	102	5.2	1.0	38	2.9	1.0	29	1.5	0.6
2011	201	13.6	2.0	81	4.1	0.9	68	4.8	1.2	31	1.6	0.6
2012	170	11.6	1.8	111	5.5	1.0	50	3.6	1.1	24	1.2	0.5
2013	184	12.1	1.8	89	4.4	0.9	61	4.4	1.2	27	1.4	0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Esophagus

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	143	21.6	3.9	60	6.3	1.7	155	24.7	4.4	49	5.3	1.6
1977	156	24.1	4.3	65	6.6	1.7	145	22.6	4.2	51	5.0	1.4
1978	178	27.1	4.5	69	7.1	1.8	124	18.0	3.4	55	5.5	1.5
1979	168	24.9	4.2	69	6.7	1.6	161	24.9	4.4	54	5.2	1.4
1980	194	28.3	4.4	64	6.1	1.5	148	22.2	4.0	65	6.2	1.6
1981	186	24.7	3.8	76	6.9	1.6	157	20.6	3.4	55	4.9	1.3
1982	176	24.5	3.9	63	5.5	1.4	155	22.3	3.8	58	5.2	1.4
1983	175	22.4	3.6	72	6.3	1.5	151	19.8	3.4	60	5.4	1.4
1984	179	22.9	3.7	84	7.1	1.6	145	19.6	3.5	62	5.4	1.4
1985	168	21.5	3.5	72	6.1	1.4	163	20.5	3.4	54	4.6	1.3
1986	190	24.7	3.8	64	5.3	1.3	152	19.9	3.5	66	5.5	1.4
1987	146	18.5	3.2	79	6.3	1.4	131	16.4	3.0	48	4.0	1.2
1988	180	22.6	3.6	73	5.8	1.3	149	19.4	3.4	53	4.2	1.2
1989	180	21.9	3.5	77	6.0	1.4	134	16.0	2.9	50	4.0	1.1
1990	158	19.0	3.2	72	5.7	1.3	123	15.3	2.9	68	5.2	1.3
1991	144	17.3	3.1	67	4.9	1.2	120	14.1	2.7	57	4.2	1.1
1992	152	17.8	3.1	69	5.2	1.3	100	11.6	2.5	48	3.6	1.0
1993	160	17.2	2.8	76	5.6	1.3	134	15.6	2.9	48	3.5	1.0
1994	152	17.4	3.0	80	5.8	1.3	122	14.1	2.7	55	4.0	1.1
1995	123	13.2	2.5	62	4.5	1.1	119	13.2	2.6	53	3.9	1.1
1996	122	13.1	2.5	73	5.3	1.2	109	11.9	2.4	57	4.1	1.1
1997	132	13.9	2.5	57	4.0	1.1	99	10.9	2.3	51	3.6	1.0
1998	126	12.8	2.4	56	3.8	1.0	95	9.8	2.1	52	3.6	1.0
1999	135	13.6	2.4	59	3.7	1.0	91	9.1	2.0	40	2.6	0.8
2000	102	10.6	2.2	85	5.6	1.2	103	10.6	2.2	63	4.1	1.0
2001	118	11.4	2.2	55	3.4	0.9	92	9.2	2.0	35	2.2	0.7
2002	98	9.4	2.0	59	3.6	0.9	86	8.3	1.9	46	2.8	0.8
2003	124	11.8	2.2	61	3.7	0.9	80	7.2	1.7	48	2.9	0.8
2004	108	9.4	1.9	65	4.0	1.0	86	8.0	1.8	53	3.2	0.9
2005	92	7.7	1.6	63	3.6	0.9	67	5.9	1.5	41	2.5	0.8
2006	91	8.9	2.0	39	2.3	0.7	71	7.0	1.7	43	2.5	0.7
2007	100	8.3	1.7	39	2.2	0.7	67	5.6	1.4	31	1.8	0.6
2008	101	8.8	1.8	46	2.5	0.7	74	6.4	1.5	46	2.6	0.8
2009	102	8.4	1.7	45	2.5	0.7	81	6.8	1.6	21	1.2	0.5
2010	81	6.5	1.5	46	2.5	0.7	74	6.3	1.5	36	1.9	0.6
2011	104	8.1	1.7	49	2.6	0.7	64	4.9	1.3	28	1.5	0.6
2012	82	6.2	1.4	44	2.1	0.6	59	4.2	1.1	29	1.4	0.5
2013	88	6.9	1.5	40	2.0	0.6	55	4.1	1.1	33	1.7	0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Stomach

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	139	26.5	5.2	105	13.8	2.9	117	23.0	4.9	72	8.9	2.2
1977	149	27.0	5.0	121	15.0	2.9	124	23.8	4.9	88	11.4	2.6
1978	138	23.8	4.5	106	13.5	2.8	106	19.3	4.2	103	13.0	2.7
1979	143	26.0	4.9	110	13.7	2.7	99	17.5	4.0	81	10.0	2.3
1980	173	30.0	5.1	143	16.8	2.9	114	21.6	4.5	94	11.3	2.4
1981	173	28.9	5.0	109	12.5	2.5	149	25.2	4.6	91	10.7	2.3
1982	163	28.5	4.9	118	12.4	2.3	130	25.6	5.0	78	8.5	2.0
1983	144	22.4	4.0	138	13.7	2.4	119	18.8	3.8	95	9.5	2.0
1984	155	24.7	4.4	110	10.4	2.0	102	18.1	3.9	77	7.1	1.7
1985	153	22.3	4.0	134	13.4	2.3	97	14.5	3.3	87	8.2	1.8
1986	183	27.9	4.6	124	11.3	2.1	124	19.1	3.8	85	8.2	1.8
1987	185	25.6	4.1	134	12.9	2.3	107	15.6	3.3	93	9.0	1.9
1988	175	25.4	4.1	130	11.8	2.1	121	18.2	3.6	94	8.8	1.8
1989	163	21.8	3.7	148	13.6	2.2	121	17.0	3.3	96	8.5	1.8
1990	181	25.3	4.1	142	12.0	2.0	113	15.8	3.3	98	8.9	1.8
1991	156	20.4	3.5	139	11.5	1.9	122	16.2	3.2	83	7.0	1.5
1992	163	20.3	3.5	161	13.1	2.1	102	13.2	2.8	127	10.6	1.9
1993	176	22.5	3.7	157	12.5	2.0	98	12.9	2.8	95	7.6	1.6
1994	196	24.7	3.8	168	13.2	2.0	141	18.5	3.4	104	8.3	1.6
1995	165	20.5	3.4	161	12.3	1.9	102	12.6	2.7	111	8.6	1.6
1996	192	23.4	3.6	147	10.8	1.8	122	15.8	3.1	94	7.1	1.5
1997	184	22.9	3.6	148	10.8	1.8	113	13.3	2.7	82	5.8	1.3
1998	184	20.7	3.2	159	11.2	1.8	113	13.4	2.7	92	6.6	1.4
1999	171	19.7	3.2	146	10.0	1.6	107	13.0	2.7	94	6.5	1.3
2000	171	19.3	3.1	165	11.0	1.7	108	12.5	2.5	87	5.8	1.2
2001	159	17.5	2.9	167	11.1	1.7	93	10.1	2.2	74	5.1	1.2
2002	162	16.9	2.8	161	10.1	1.6	101	11.6	2.5	88	5.6	1.2
2003	170	16.8	2.7	157	9.8	1.5	93	8.8	1.9	81	5.1	1.1
2004	153	16.2	2.7	160	9.7	1.5	84	8.9	2.0	90	5.5	1.1
2005	210	21.3	3.1	197	11.9	1.7	122	13.2	2.5	90	5.5	1.1
2006	203	19.6	2.9	165	9.7	1.5	107	11.1	2.2	81	4.9	1.1
2007	171	16.4	2.6	163	9.5	1.5	86	8.1	1.8	92	5.5	1.1
2008	164	15.7	2.5	143	8.1	1.3	93	8.9	1.9	87	5.0	1.1
2009	201	17.6	2.6	156	8.9	1.4	85	8.5	1.9	74	4.2	1.0
2010	180	15.5	2.4	164	9.1	1.4	92	8.8	1.9	67	3.7	0.9
2011	188	15.5	2.3	183	9.7	1.4	98	8.7	1.8	73	3.9	0.9
2012	210	16.6	2.4	173	9.0	1.4	85	6.9	1.6	80	4.3	1.0
2013	197	15.5	2.3	175	8.8	1.3	91	7.4	1.6	81	4.1	0.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Colorectal

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	272	51.3	7.0	370	46.1	5.1	168	35.1	6.2	216	28.9	4.2
1977	321	64.7	8.1	460	57.1	5.7	174	36.1	6.0	201	25.4	3.8
1978	334	65.6	8.2	467	57.4	5.6	202	41.7	6.6	260	33.5	4.4
1979	369	68.8	8.0	472	55.5	5.3	206	40.9	6.4	270	32.3	4.1
1980	391	70.0	7.9	497	55.2	5.1	197	36.4	5.8	266	30.7	3.9
1981	376	67.2	7.7	492	52.8	4.9	201	38.8	6.0	295	33.3	4.0
1982	397	64.9	7.1	475	48.9	4.6	208	35.4	5.5	260	27.6	3.5
1983	383	64.1	7.3	487	47.4	4.4	203	31.6	4.9	240	24.3	3.2
1984	430	66.1	7.0	499	49.1	4.5	211	35.3	5.4	272	26.7	3.3
1985	447	69.9	7.3	542	50.3	4.4	201	33.6	5.2	291	28.4	3.4
1986	472	69.3	6.9	566	51.2	4.4	215	34.0	5.1	259	24.1	3.0
1987	434	63.0	6.6	527	46.1	4.1	222	34.5	5.1	257	23.0	2.9
1988	420	59.9	6.4	558	48.1	4.1	221	34.2	5.1	255	22.8	2.9
1989	485	69.0	6.8	534	44.8	3.9	238	36.3	5.1	231	19.5	2.6
1990	434	60.9	6.3	574	47.6	4.0	212	31.8	4.8	253	21.8	2.8
1991	467	62.3	6.3	613	50.4	4.1	216	32.9	4.9	279	23.4	2.8
1992	447	59.7	6.1	622	49.3	3.9	204	29.6	4.5	273	21.9	2.7
1993	489	62.3	6.1	592	46.4	3.8	211	28.4	4.2	273	21.8	2.6
1994	522	66.8	6.3	642	49.1	3.9	254	32.8	4.4	281	22.0	2.6
1995	525	63.8	6.0	639	47.2	3.7	233	30.0	4.2	303	23.2	2.6
1996	558	66.5	6.0	687	49.9	3.8	263	34.0	4.4	284	21.3	2.5
1997	593	71.1	6.3	698	49.7	3.7	271	34.3	4.5	299	21.7	2.5
1998	548	62.5	5.7	737	51.4	3.7	226	26.9	3.8	287	20.4	2.4
1999	598	65.3	5.6	816	54.7	3.8	255	30.8	4.1	342	23.4	2.5
2000	569	61.1	5.4	710	45.9	3.4	249	27.9	3.8	299	19.9	2.3
2001	603	63.0	5.4	759	48.1	3.4	237	27.2	3.7	306	19.7	2.2
2002	600	59.3	5.1	783	48.7	3.4	248	27.9	3.8	293	18.6	2.1
2003	668	67.0	5.5	785	47.9	3.4	258	28.1	3.7	289	17.9	2.1
2004	643	61.5	5.1	824	49.1	3.4	258	26.7	3.5	278	17.1	2.0
2005	649	60.3	4.9	765	44.9	3.2	220	22.4	3.1	279	16.7	2.0
2006	633	57.6	4.8	838	48.6	3.3	233	23.5	3.2	294	17.3	2.0
2007	684	58.7	4.7	760	43.3	3.1	251	23.3	3.1	296	17.2	2.0
2008	687	58.9	4.7	794	44.3	3.1	231	21.5	2.9	258	14.7	1.8
2009	700	56.8	4.4	753	40.9	3.0	246	22.7	3.0	261	14.5	1.8
2010	667	55.6	4.5	736	39.4	2.9	247	21.7	2.9	234	12.4	1.6
2011	692	53.6	4.2	743	38.4	2.8	214	18.5	2.6	280	14.8	1.8
2012	653	49.7	4.0	686	34.9	2.6	260	21.5	2.8	277	14.3	1.7
2013	650	47.9	3.9	728	36.0	2.6	223	17.1	2.4	258	12.9	1.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Colon excluding rectum

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	191	37.5	6.1	280	34.5	4.4	135	28.6	5.7	177	23.4	3.8
1977	211	44.2	6.9	356	45.1	5.1	124	26.8	5.3	166	21.0	3.4
1978	245	50.3	7.3	357	44.1	4.9	163	34.4	6.1	219	27.8	3.9
1979	284	53.1	7.1	369	43.7	4.7	171	34.2	5.8	233	27.6	3.8
1980	280	51.1	6.8	374	41.8	4.5	158	29.9	5.3	220	25.6	3.6
1981	272	50.4	6.8	360	39.6	4.3	170	33.1	5.6	257	28.9	3.7
1982	293	49.3	6.3	354	36.6	4.0	168	29.8	5.1	222	23.4	3.2
1983	284	48.2	6.3	371	35.9	3.8	165	26.3	4.5	206	20.9	3.0
1984	306	47.3	5.9	382	38.2	4.0	183	30.6	5.0	237	23.2	3.1
1985	322	52.1	6.4	423	39.7	3.9	165	27.8	4.7	256	24.8	3.1
1986	338	50.7	6.0	430	38.7	3.8	186	29.3	4.7	225	20.9	2.8
1987	328	47.5	5.7	406	35.7	3.6	194	29.9	4.7	229	20.5	2.7
1988	329	47.1	5.6	436	38.0	3.7	192	29.1	4.7	224	19.9	2.7
1989	367	53.1	6.0	422	35.7	3.5	209	32.0	4.8	208	17.5	2.4
1990	324	46.2	5.5	443	36.7	3.5	183	27.1	4.4	222	19.1	2.6
1991	362	48.9	5.6	465	38.5	3.6	195	29.5	4.6	251	21.0	2.7
1992	331	45.2	5.4	494	39.2	3.5	176	26.0	4.3	238	19.0	2.5
1993	355	46.9	5.4	457	36.1	3.4	185	24.6	3.9	245	19.5	2.5
1994	394	51.3	5.6	517	39.8	3.5	222	29.1	4.2	247	19.4	2.4
1995	394	49.2	5.3	498	37.0	3.3	212	27.7	4.1	270	20.7	2.5
1996	408	49.4	5.3	527	38.4	3.3	222	29.2	4.1	247	18.5	2.3
1997	437	53.1	5.5	555	39.4	3.3	231	29.9	4.2	261	18.9	2.3
1998	417	48.9	5.1	575	40.2	3.3	197	23.4	3.6	256	18.1	2.2
1999	433	48.5	4.9	637	42.7	3.4	217	26.5	3.8	312	21.3	2.4
2000	420	46.5	4.8	554	36.1	3.0	218	25.0	3.6	266	17.7	2.1
2001	442	47.5	4.8	581	37.0	3.0	209	24.2	3.5	278	18.0	2.1
2002	442	44.4	4.4	635	39.7	3.1	216	24.5	3.5	267	17.0	2.1
2003	488	50.5	4.8	606	37.1	3.0	209	23.2	3.4	257	15.9	2.0
2004	476	46.4	4.5	633	38.0	3.0	218	22.5	3.2	248	15.4	1.9
2005	480	45.7	4.3	594	35.1	2.8	183	18.8	2.9	249	15.0	1.9
2006	462	42.8	4.2	626	36.7	2.9	204	20.8	3.0	253	14.9	1.9
2007	498	43.9	4.1	577	33.0	2.7	208	19.4	2.8	257	14.9	1.8
2008	504	44.3	4.1	618	34.7	2.8	194	18.2	2.7	227	12.9	1.7
2009	497	41.3	3.9	568	31.0	2.6	211	19.9	2.8	230	12.8	1.7
2010	473	40.2	3.9	559	30.2	2.5	209	18.7	2.7	199	10.6	1.5
2011	501	39.7	3.7	568	29.5	2.5	177	15.7	2.4	241	12.7	1.6
2012	453	35.7	3.5	519	26.6	2.3	206	17.1	2.5	244	12.7	1.6
2013	457	34.4	3.3	524	26.0	2.3	186	14.3	2.2	232	11.6	1.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Rectum & rectosigmoid

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	81	13.8	3.4	90	11.6	2.6	33	6.6	2.6	39	5.5	1.9
1977	110	20.5	4.3	104	12.1	2.5	50	9.3	2.8	35	4.4	1.6
1978	89	15.3	3.7	110	13.3	2.7	39	7.3	2.5	41	5.7	1.9
1979	85	15.7	3.8	103	11.8	2.4	35	6.7	2.5	37	4.7	1.6
1980	111	18.8	3.9	123	13.4	2.5	39	6.6	2.3	46	5.1	1.6
1981	104	16.8	3.6	132	13.1	2.4	31	5.8	2.3	38	4.4	1.5
1982	104	15.6	3.3	121	12.3	2.3	40	5.6	1.9	38	4.2	1.4
1983	99	15.9	3.5	116	11.5	2.2	38	5.3	1.9	34	3.4	1.2
1984	124	18.7	3.7	117	10.9	2.0	28	4.8	2.0	35	3.5	1.2
1985	125	17.8	3.5	119	10.6	2.0	36	5.8	2.1	35	3.6	1.2
1986	134	18.6	3.5	136	12.4	2.2	29	4.7	1.9	34	3.2	1.1
1987	106	15.6	3.3	121	10.5	1.9	28	4.6	1.9	28	2.5	1.0
1988	91	12.8	2.9	122	10.2	1.9	29	5.1	2.0	31	2.9	1.0
1989	118	16.0	3.2	112	9.1	1.7	29	4.3	1.8	23	2.0	0.8
1990	110	14.7	3.0	131	10.9	1.9	29	4.7	2.0	31	2.7	1.0
1991	105	13.4	2.8	148	11.9	2.0	21	3.3	1.6	28	2.3	0.9
1992	116	14.5	2.9	128	10.1	1.8	28	3.6	1.5	35	2.9	1.0
1993	134	15.4	2.9	135	10.3	1.8	26	3.8	1.6	28	2.4	0.9
1994	128	15.5	2.9	125	9.2	1.6	32	3.7	1.4	34	2.6	0.9
1995	131	14.6	2.7	141	10.1	1.7	21	2.4	1.1	33	2.5	0.9
1996	150	17.1	2.9	160	11.6	1.8	41	4.8	1.6	37	2.8	0.9
1997	156	18.0	3.1	143	10.3	1.7	40	4.5	1.5	38	2.8	0.9
1998	131	13.6	2.5	162	11.2	1.7	29	3.4	1.3	31	2.3	0.8
1999	165	16.7	2.7	179	12.0	1.8	38	4.3	1.5	30	2.1	0.8
2000	149	14.6	2.5	156	9.9	1.6	31	2.9	1.1	33	2.2	0.8
2001	161	15.5	2.6	178	11.1	1.6	28	3.0	1.2	28	1.7	0.6
2002	158	14.9	2.5	148	9.0	1.5	32	3.4	1.3	26	1.7	0.6
2003	180	16.5	2.6	179	10.8	1.6	49	4.8	1.5	32	2.0	0.7
2004	167	15.2	2.4	191	11.1	1.6	40	4.1	1.4	30	1.8	0.6
2005	169	14.6	2.3	171	9.8	1.5	37	3.5	1.2	30	1.8	0.6
2006	171	14.8	2.4	212	11.9	1.6	29	2.7	1.1	41	2.4	0.7
2007	186	14.8	2.2	183	10.3	1.5	43	3.9	1.2	39	2.3	0.7
2008	183	14.6	2.2	176	9.6	1.4	37	3.2	1.1	31	1.8	0.6
2009	203	15.4	2.2	185	9.9	1.4	35	2.8	1.0	31	1.7	0.6
2010	194	15.4	2.3	177	9.2	1.4	38	3.0	1.0	35	1.8	0.6
2011	191	13.9	2.1	175	8.9	1.3	37	2.8	0.9	39	2.0	0.6
2012	200	14.1	2.1	167	8.3	1.3	54	4.4	1.2	33	1.7	0.6
2013	193	13.4	2.0	204	10.0	1.4	37	2.8	1.0	26	1.2	0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Liver / intrahepatic bile duct

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	55	9.5	2.8	24	2.9	1.3	58	10.5	3.1	24	3.2	1.4
1977	51	8.0	2.5	23	2.8	1.2	66	10.9	3.0	26	2.8	1.1
1978	60	9.7	2.7	41	4.6	1.5	45	7.0	2.3	44	5.0	1.6
1979	44	7.0	2.3	31	3.3	1.2	51	8.9	2.7	42	4.8	1.6
1980	59	8.7	2.5	29	3.2	1.2	55	8.5	2.6	30	3.3	1.2
1981	57	8.4	2.4	33	3.3	1.2	55	8.1	2.4	38	4.3	1.4
1982	73	10.5	2.6	27	2.8	1.1	63	9.0	2.4	32	3.5	1.3
1983	65	9.5	2.6	31	2.9	1.1	77	11.3	2.8	44	4.4	1.3
1984	69	10.4	2.8	33	2.9	1.0	68	10.9	3.0	45	3.9	1.2
1985	62	8.5	2.3	37	3.1	1.0	58	8.2	2.4	44	4.1	1.2
1986	65	8.9	2.4	33	2.8	1.0	66	8.9	2.3	39	3.4	1.1
1987	57	7.1	2.1	44	3.7	1.1	57	8.5	2.5	55	4.9	1.3
1988	64	8.0	2.2	35	2.9	1.0	67	8.8	2.3	37	3.1	1.0
1989	68	9.1	2.5	42	3.6	1.1	61	7.7	2.1	50	4.2	1.2
1990	81	9.6	2.3	50	4.1	1.2	65	8.2	2.2	53	4.3	1.2
1991	90	10.5	2.4	46	3.5	1.0	76	9.4	2.3	53	4.1	1.1
1992	83	10.2	2.4	49	3.9	1.1	72	8.8	2.2	52	4.0	1.1
1993	83	9.6	2.2	47	3.7	1.1	76	8.7	2.1	56	4.5	1.2
1994	84	9.2	2.1	56	4.2	1.1	76	9.3	2.2	44	3.4	1.0
1995	100	10.8	2.3	54	3.9	1.1	79	8.9	2.1	51	3.7	1.0
1996	120	13.3	2.6	50	3.7	1.0	90	10.4	2.3	56	4.1	1.1
1997	110	11.5	2.3	64	4.5	1.1	78	8.3	2.0	51	3.6	1.0
1998	131	12.4	2.2	59	4.1	1.1	105	10.6	2.2	54	3.8	1.0
1999	109	10.3	2.0	72	4.7	1.1	65	6.2	1.6	47	3.2	0.9
2000	126	11.0	2.0	63	4.0	1.0	103	10.1	2.1	54	3.6	1.0
2001	141	12.3	2.2	49	3.1	0.9	86	7.8	1.8	57	3.6	0.9
2002	146	12.6	2.2	58	3.7	0.9	86	8.0	1.8	50	3.2	0.9
2003	162	13.0	2.1	68	4.1	1.0	107	9.4	1.9	47	2.9	0.8
2004	192	15.8	2.4	82	4.9	1.1	119	10.0	1.9	76	4.7	1.1
2005	215	17.1	2.4	70	4.0	1.0	121	10.3	1.9	55	3.3	0.9
2006	190	15.2	2.3	65	3.7	0.9	127	10.3	1.9	47	2.8	0.8
2007	239	18.4	2.5	87	4.9	1.0	131	10.7	1.9	66	3.8	0.9
2008	245	17.8	2.3	82	4.5	1.0	144	11.1	1.9	54	2.9	0.8
2009	269	19.7	2.5	103	5.5	1.1	163	12.3	2.0	51	2.8	0.8
2010	271	18.8	2.4	113	5.8	1.1	172	12.7	2.0	78	3.9	0.9
2011	269	17.9	2.2	109	5.7	1.1	159	10.9	1.8	82	4.3	0.9
2012	301	20.3	2.4	114	5.7	1.1	179	12.2	1.9	76	4.0	0.9
2013	273	17.4	2.1	125	6.0	1.1	173	12.1	1.9	73	3.5	0.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Pancreas

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	85	14.5	3.4	118	15.4	3.0	89	15.0	3.5	100	12.3	2.6
1977	105	17.8	4.0	96	11.7	2.5	99	16.9	3.9	83	10.1	2.4
1978	101	16.3	3.7	117	14.3	2.8	82	13.5	3.3	102	11.8	2.4
1979	113	19.1	4.0	116	14.0	2.7	87	13.4	3.1	116	14.3	2.8
1980	112	18.7	4.0	96	10.6	2.2	96	16.6	3.9	81	9.1	2.1
1981	128	21.1	4.1	112	12.1	2.4	108	17.4	3.8	93	10.2	2.2
1982	102	16.9	3.7	134	13.8	2.4	100	17.4	3.8	126	13.1	2.4
1983	130	19.0	3.6	140	13.7	2.4	126	19.6	3.8	117	11.5	2.2
1984	113	17.8	3.7	147	14.3	2.4	117	18.6	3.8	116	11.2	2.1
1985	126	18.0	3.5	117	11.3	2.1	103	15.9	3.5	103	10.1	2.0
1986	115	15.6	3.1	144	13.2	2.2	104	14.3	3.0	120	11.5	2.1
1987	125	16.6	3.3	156	13.9	2.3	106	15.3	3.3	136	11.9	2.1
1988	130	18.7	3.5	144	12.7	2.1	110	15.0	3.1	106	9.6	1.9
1989	104	14.4	3.0	151	12.7	2.1	99	13.6	2.9	135	11.5	2.0
1990	122	17.8	3.5	176	15.0	2.3	90	13.0	3.0	153	12.8	2.1
1991	123	16.3	3.2	132	11.0	1.9	111	14.2	2.9	110	9.1	1.7
1992	104	14.2	3.1	157	12.5	2.0	81	12.3	3.0	124	10.2	1.8
1993	107	13.8	2.9	138	10.9	1.9	84	11.8	2.8	120	9.3	1.7
1994	124	15.6	3.0	172	13.3	2.0	104	13.6	2.8	153	11.9	1.9
1995	117	14.2	2.8	146	11.1	1.8	108	13.7	2.8	132	9.9	1.7
1996	122	14.3	2.7	183	13.3	1.9	95	11.7	2.6	147	10.7	1.8
1997	128	14.1	2.6	175	12.7	1.9	103	11.8	2.4	137	10.1	1.7
1998	112	12.4	2.5	180	12.8	1.9	102	11.6	2.4	155	11.2	1.8
1999	136	15.1	2.7	200	13.7	1.9	108	11.7	2.4	178	12.2	1.8
2000	112	12.2	2.4	208	13.9	1.9	109	12.2	2.4	147	9.8	1.6
2001	140	14.3	2.5	220	14.3	1.9	105	11.1	2.2	179	11.7	1.7
2002	144	14.0	2.4	200	12.9	1.8	134	13.1	2.4	154	9.9	1.6
2003	158	16.0	2.7	162	10.1	1.6	114	11.9	2.3	144	9.1	1.5
2004	144	13.6	2.4	197	12.1	1.7	133	13.3	2.4	161	9.8	1.5
2005	148	13.9	2.4	198	11.7	1.6	120	11.2	2.1	149	9.0	1.5
2006	170	15.0	2.4	234	13.7	1.8	115	10.7	2.1	162	9.4	1.5
2007	171	15.4	2.5	216	12.6	1.7	136	12.5	2.2	164	9.7	1.5
2008	150	12.8	2.2	227	12.9	1.7	123	11.2	2.1	185	10.6	1.5
2009	188	16.0	2.4	199	11.0	1.6	141	12.4	2.2	180	10.0	1.5
2010	193	15.8	2.4	227	12.4	1.6	131	11.0	2.0	166	9.4	1.4
2011	216	16.7	2.4	280	14.9	1.8	160	13.5	2.2	201	10.7	1.5
2012	194	14.9	2.2	285	14.5	1.7	141	10.5	1.8	231	11.9	1.6
2013	227	16.8	2.3	270	13.5	1.6	161	12.7	2.1	182	9.1	1.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Larynx

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	95	13.3	2.8	20	2.1	1.0	52	8.8	2.7	14	1.5	0.9
1977	92	14.3	3.4	20	2.0	0.9	41	6.8	2.6	13	1.2	0.7
1978	101	14.4	3.1	24	2.2	0.9	71	9.9	2.5	12	1.1	0.6
1979	115	16.1	3.1	33	3.1	1.1	57	8.0	2.2	11	1.0	0.6
1980	108	14.2	2.9	30	2.7	1.0	45	6.4	2.0	21	1.8	0.8
1981	107	13.5	2.7	25	2.3	0.9	51	7.0	2.1	23	1.9	0.8
1982	107	13.3	2.6	14	1.2	0.7	47	6.5	2.0	11	1.0	0.6
1983	116	14.4	2.8	22	2.0	0.9	53	6.7	1.9	11	1.0	0.6
1984	108	13.2	2.6	39	3.3	1.0	59	7.3	2.0	14	1.2	0.7
1985	147	17.9	3.0	41	3.2	1.0	55	7.1	2.0	12	1.1	0.6
1986	140	17.7	3.2	39	3.1	1.0	63	8.1	2.1	18	1.5	0.7
1987	121	14.3	2.7	30	2.4	0.9	57	7.2	2.0	21	1.7	0.8
1988	114	12.7	2.5	37	2.8	0.9	75	9.1	2.2	13	0.9	0.5
1989	115	13.2	2.6	62	4.6	1.2	53	6.5	1.9	18	1.3	0.6
1990	137	15.4	2.8	42	3.2	1.0	49	6.0	1.9	13	1.0	0.5
1991	117	13.3	2.6	34	2.4	0.8	57	6.5	1.8	15	1.1	0.6
1992	150	15.9	2.7	38	2.6	0.8	63	7.2	1.9	19	1.3	0.6
1993	121	13.7	2.6	33	2.4	0.8	62	7.1	1.9	21	1.5	0.7
1994	114	12.1	2.4	33	2.3	0.8	56	6.2	1.7	22	1.6	0.7
1995	113	11.7	2.3	42	2.9	0.9	70	8.2	2.1	20	1.4	0.6
1996	127	13.0	2.4	27	1.8	0.7	69	7.8	2.0	11	0.7	0.4
1997	107	10.6	2.1	35	2.3	0.8	54	5.4	1.5	14	0.9	0.5
1998	110	11.1	2.2	49	3.2	0.9	57	5.7	1.5	10	0.7	0.4
1999	103	10.0	2.1	33	2.1	0.7	50	5.0	1.5	15	1.0	0.5
2000	90	8.6	1.9	38	2.4	0.8	49	5.0	1.5	18	1.2	0.6
2001	95	8.2	1.7	28	1.7	0.6	36	3.6	1.3	9	0.6	0.4
2002	89	7.5	1.6	36	2.2	0.7	52	4.5	1.3	18	1.1	0.5
2003	109	9.7	1.9	38	2.2	0.7	45	4.1	1.3	12	0.7	0.4
2004	111	9.0	1.7	20	1.1	0.5	50	4.5	1.3	11	0.7	0.4
2005	94	8.5	1.8	27	1.5	0.6	44	3.8	1.2	7	0.4	0.3
2006	96	7.6	1.6	29	1.5	0.6	42	3.8	1.2	8	0.5	0.3
2007	107	8.8	1.7	33	1.8	0.6	42	3.6	1.1	12	0.7	0.4
2008	106	8.9	1.8	47	2.6	0.8	41	3.5	1.1	20	1.1	0.5
2009	92	7.5	1.6	29	1.5	0.6	42	3.6	1.2	15	0.8	0.4
2010	95	7.4	1.6	33	1.7	0.6	39	3.4	1.1	8	0.4	0.3
2011	98	7.0	1.4	32	1.5	0.5	30	2.4	0.9	13	0.7	0.4
2012	73	5.4	1.3	29	1.4	0.5	43	3.5	1.1	9	0.4	0.3
2013	86	6.3	1.4	27	1.2	0.5	36	2.8	1.0	11	0.5	0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Lung and bronchus

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	669	105.0	8.8	223	24.9	3.5	517	82.0	8.0	163	18.0	2.9
1977	707	113.7	9.5	251	25.6	3.3	573	92.0	8.5	186	19.0	2.9
1978	730	110.8	8.9	258	28.2	3.6	613	94.0	8.2	214	22.9	3.3
1979	768	119.6	9.5	298	30.9	3.7	639	101.4	8.9	216	23.0	3.3
1980	825	122.5	9.3	298	28.8	3.4	667	100.8	8.5	237	23.7	3.2
1981	824	118.5	8.9	337	31.3	3.4	685	100.4	8.3	272	25.7	3.2
1982	827	116.3	8.6	350	32.0	3.5	667	93.3	7.7	278	26.3	3.2
1983	831	114.0	8.4	362	32.6	3.5	753	106.6	8.3	295	26.5	3.1
1984	844	112.4	8.2	401	35.4	3.6	689	94.5	7.7	289	25.2	3.0
1985	874	116.5	8.4	389	33.1	3.4	714	97.2	7.8	306	26.5	3.1
1986	859	114.1	8.4	448	37.1	3.5	667	89.4	7.4	354	29.5	3.1
1987	897	113.9	8.0	430	34.8	3.4	732	97.1	7.7	365	30.0	3.1
1988	869	112.9	8.2	458	36.4	3.4	698	91.9	7.4	354	28.6	3.1
1989	925	116.5	8.1	478	36.9	3.4	761	99.7	7.7	365	28.5	3.0
1990	918	114.1	8.0	478	36.7	3.4	735	90.8	7.1	376	29.5	3.0
1991	889	110.1	7.9	504	38.4	3.4	707	89.2	7.1	374	28.6	3.0
1992	932	113.4	8.0	504	37.4	3.3	665	84.7	7.1	398	30.1	3.0
1993	850	103.6	7.6	521	38.1	3.3	696	86.8	7.1	389	28.4	2.9
1994	886	103.0	7.3	577	41.0	3.4	769	91.3	7.0	410	30.0	2.9
1995	865	98.0	7.0	541	38.7	3.3	704	82.2	6.5	459	32.7	3.0
1996	908	100.3	7.0	629	44.0	3.5	724	81.1	6.4	463	33.0	3.0
1997	873	96.1	6.8	619	42.5	3.4	696	79.6	6.3	458	31.8	2.9
1998	846	88.6	6.3	655	44.0	3.4	691	75.5	6.0	467	32.0	2.9
1999	813	85.1	6.2	604	39.8	3.2	620	66.7	5.6	462	31.1	2.9
2000	859	86.6	6.1	615	39.3	3.1	654	67.8	5.5	455	29.2	2.7
2001	842	85.7	6.1	629	39.6	3.1	625	65.2	5.4	470	29.8	2.7
2002	826	82.3	6.0	661	40.5	3.1	645	64.8	5.3	478	29.4	2.7
2003	767	73.1	5.5	685	41.3	3.1	613	60.0	5.0	485	29.5	2.6
2004	853	77.5	5.5	702	41.7	3.1	599	57.4	4.9	469	28.0	2.6
2005	856	79.4	5.6	687	40.1	3.0	602	55.7	4.7	510	30.3	2.6
2006	819	73.1	5.3	700	40.6	3.0	546	50.7	4.5	490	28.7	2.6
2007	831	73.1	5.3	728	41.1	3.0	591	52.7	4.5	490	28.1	2.5
2008	814	71.2	5.2	748	41.9	3.0	526	46.8	4.2	479	27.1	2.5
2009	859	72.2	5.1	769	42.0	3.0	568	49.0	4.3	459	25.3	2.3
2010	837	68.0	4.9	808	43.1	3.0	610	52.6	4.4	510	27.4	2.4
2011	864	69.5	4.9	836	43.2	3.0	599	48.6	4.1	530	27.7	2.4
2012	867	67.7	4.8	812	41.1	2.9	599	48.6	4.1	510	26.1	2.3
2013	797	60.2	4.4	768	37.8	2.7	576	45.0	3.9	505	25.1	2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Melanoma of the skin

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	2	0.3	0.5	8	1.0	0.8	4	1.1	1.3	5	0.7	0.7
1977	8	1.9	1.6	6	0.9	0.8	4	1.3	1.5	3	0.3	0.3
1978	5	1.2	1.2	8	0.9	0.6	2	0.2	0.3	4	0.5	0.5
1979	7	0.9	0.7	11	1.1	0.7	2	0.4	0.6	2	0.3	0.5
1980	5	0.6	0.5	9	1.1	0.8	8	0.9	0.6	6	0.7	0.6
1981	7	0.8	0.6	18	1.8	0.9	7	1.0	0.8	8	0.9	0.7
1982	8	0.9	0.6	11	1.0	0.6	6	1.1	1.0	4	0.3	0.3
1983	4	0.5	0.6	8	0.7	0.5	14	1.7	0.9	6	0.6	0.5
1984	7	1.0	0.9	19	1.8	0.8	10	1.1	0.7	3	0.2	0.2
1985	12	1.4	0.9	11	1.2	0.7	10	1.7	1.2	4	0.3	0.3
1986	11	1.7	1.2	9	0.8	0.5	15	1.8	1.0	6	0.6	0.5
1987	8	0.9	0.7	9	0.7	0.5	4	0.5	0.5	5	0.4	0.4
1988	7	1.1	0.9	13	1.0	0.6	4	0.7	0.8	5	0.5	0.4
1989	6	0.8	0.7	11	1.1	0.7	6	0.6	0.6	11	0.9	0.6
1990	10	1.2	0.9	11	0.8	0.5	3	0.5	0.7	3	0.3	0.3
1991	13	2.0	1.3	12	0.7	0.4	6	0.8	0.8	5	0.4	0.4
1992	6	0.5	0.5	8	0.6	0.4	3	0.3	0.3	4	0.3	0.3
1993	12	1.5	0.9	13	1.0	0.6	2	0.2	0.2	6	0.4	0.3
1994	5	0.6	0.7	13	0.9	0.5	1	0.2	0.4	5	0.4	0.4
1995	6	0.5	0.4	18	1.3	0.6	1	0.1	0.1	5	0.4	0.3
1996	8	1.0	0.8	15	1.1	0.6	4	0.5	0.5	9	0.7	0.4
1997	15	1.6	0.9	22	1.5	0.7	4	0.3	0.3	5	0.4	0.4
1998	13	1.5	0.9	15	1.0	0.5	3	0.2	0.3	8	0.6	0.4
1999	5	0.4	0.4	7	0.5	0.3	5	0.6	0.6	3	0.2	0.2
2000	6	0.6	0.5	6	0.4	0.3	7	1.1	0.9	6	0.4	0.3
2001	10	0.8	0.5	19	1.2	0.5	3	0.3	0.3	3	0.2	0.2
2002	9	0.8	0.5	16	1.0	0.5	3	0.4	0.5	4	0.3	0.3
2003	5	0.6	0.6	15	0.9	0.4	3	0.3	0.4	7	0.4	0.3
2004	11	1.1	0.7	13	0.7	0.4	2	0.1	0.2	2	0.1	0.2
2005	12	0.9	0.5	16	0.9	0.5	4	0.3	0.3	2	0.1	0.2
2006	19	1.9	0.9	19	1.1	0.5	6	0.5	0.4	5	0.3	0.3
2007	12	1.1	0.7	30	1.7	0.6	6	0.7	0.6	10	0.6	0.4
2008	15	1.2	0.7	25	1.4	0.6	3	0.2	0.2	7	0.4	0.3
2009	11	0.7	0.4	20	1.0	0.5	5	0.4	0.3	3	0.2	0.2
2010	12	0.9	0.5	19	0.9	0.4	3	0.3	0.4	8	0.4	0.3
2011	18	1.4	0.7	21	1.1	0.5	6	0.5	0.4	9	0.5	0.3
2012	14	1.1	0.6	16	0.8	0.4	3	0.3	0.3	9	0.4	0.3
2013	12	1.0	0.6	27	1.3	0.5	8	0.5	0.3	7	0.3	0.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Female breast

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	689	75.4	6.0	315	36.5	4.4
1977	713	75.7	5.9	295	32.2	3.9
1978	792	82.3	6.0	313	33.2	3.9
1979	802	81.2	5.9	294	30.3	3.6
1980	868	83.8	5.8	348	35.0	3.9
1981	855	81.9	5.7	357	35.6	3.9
1982	905	84.2	5.7	367	34.3	3.6
1983	926	83.2	5.5	376	34.3	3.6
1984	1045	90.9	5.7	413	38.2	3.8
1985	1058	90.2	5.6	342	29.6	3.2
1986	1057	86.6	5.4	392	32.6	3.3
1987	1135	90.6	5.4	396	32.2	3.3
1988	1225	95.4	5.5	427	33.5	3.2
1989	1154	88.7	5.2	425	33.6	3.3
1990	1226	92.4	5.3	430	33.3	3.2
1991	1371	101.5	5.5	434	32.8	3.1
1992	1327	96.8	5.3	444	32.8	3.1
1993	1352	96.7	5.2	423	31.4	3.0
1994	1426	99.4	5.2	496	35.0	3.1
1995	1454	99.4	5.2	511	35.9	3.1
1996	1595	106.7	5.3	489	33.5	3.0
1997	1639	107.7	5.3	490	33.3	3.0
1998	1795	115.5	5.4	491	32.4	2.9
1999	1649	103.6	5.0	473	30.5	2.8
2000	1612	98.5	4.8	463	29.2	2.7
2001	1610	96.6	4.8	516	31.7	2.8
2002	1715	100.6	4.8	467	28.2	2.6
2003	1775	102.7	4.8	468	27.7	2.5
2004	1825	103.9	4.8	476	27.8	2.5
2005	1845	104.0	4.8	518	29.5	2.6
2006	1951	107.9	4.8	473	26.8	2.4
2007	1997	109.5	4.8	469	25.9	2.4
2008	2068	112.1	4.9	467	25.5	2.3
2009	2097	111.9	4.8	483	26.1	2.4
2010	2071	108.3	4.7	446	23.3	2.2
2011	2354	120.1	4.9	503	26.2	2.3
2012	2240	112.5	4.7	503	25.6	2.3
2013	2363	116.9	4.8	454	22.5	2.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Cervix uteri

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	292	29.8	3.6	130	15.0	2.8
1977	303	30.6	3.6	129	14.3	2.6
1978	289	28.3	3.4	110	11.7	2.3
1979	310	29.6	3.4	105	10.8	2.2
1980	278	26.6	3.3	115	11.6	2.2
1981	236	20.7	2.7	108	10.1	2.0
1982	264	23.4	2.9	97	8.5	1.8
1983	251	21.4	2.7	98	8.5	1.7
1984	260	21.7	2.7	77	7.2	1.7
1985	232	19.0	2.5	94	7.8	1.6
1986	269	21.9	2.7	91	7.5	1.6
1987	248	19.0	2.4	74	5.9	1.4
1988	233	17.7	2.3	108	8.5	1.6
1989	230	16.6	2.2	69	5.4	1.3
1990	299	20.8	2.4	83	6.4	1.4
1991	269	18.7	2.3	82	5.9	1.3
1992	259	17.4	2.2	78	5.6	1.3
1993	253	16.9	2.1	101	6.9	1.4
1994	262	17.7	2.2	79	5.4	1.2
1995	270	17.6	2.1	93	6.4	1.3
1996	291	18.5	2.1	78	5.1	1.1
1997	247	15.5	2.0	91	6.0	1.3
1998	257	15.4	1.9	79	5.1	1.1
1999	249	15.1	1.9	88	5.4	1.1
2000	288	17.3	2.0	91	5.4	1.1
2001	245	14.4	1.8	70	4.2	1.0
2002	241	13.9	1.8	74	4.4	1.0
2003	231	13.2	1.7	85	4.9	1.0
2004	214	11.9	1.6	64	3.7	0.9
2005	232	13.0	1.7	71	4.1	0.9
2006	219	12.1	1.6	72	4.1	1.0
2007	235	13.0	1.7	93	5.2	1.1
2008	216	11.7	1.6	77	4.1	0.9
2009	198	10.4	1.5	90	4.7	1.0
2010	210	11.2	1.5	60	3.1	0.8
2011	207	10.8	1.5	81	4.2	0.9
2012	197	10.3	1.5	67	3.4	0.8
2013	211	10.7	1.5	83	4.1	0.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Corpus uterus and uterus, not otherwise specified

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	146	16.4	2.8	68	8.9	2.3
1977	143	16.1	2.8	66	7.4	1.9
1978	140	15.4	2.7	69	8.4	2.1
1979	143	14.3	2.5	66	7.3	1.9
1980	183	18.7	2.9	82	8.4	1.9
1981	141	13.7	2.4	72	6.9	1.7
1982	162	14.7	2.4	58	5.4	1.4
1983	164	14.7	2.3	74	7.1	1.7
1984	187	17.1	2.5	75	6.9	1.6
1985	213	18.3	2.5	87	8.2	1.8
1986	149	12.2	2.0	65	5.7	1.4
1987	210	17.2	2.4	55	4.9	1.3
1988	216	17.0	2.3	81	6.7	1.5
1989	182	13.9	2.1	76	5.9	1.4
1990	205	15.8	2.2	76	6.2	1.4
1991	250	18.8	2.4	77	6.0	1.4
1992	243	17.9	2.3	87	6.8	1.5
1993	265	19.0	2.3	85	6.6	1.4
1994	279	19.4	2.3	92	6.7	1.4
1995	255	17.6	2.2	77	5.6	1.3
1996	305	20.7	2.3	110	7.8	1.5
1997	322	21.7	2.4	109	7.5	1.4
1998	328	21.4	2.3	124	8.5	1.5
1999	353	22.7	2.4	114	7.6	1.4
2000	363	22.9	2.4	110	7.2	1.3
2001	384	23.4	2.4	123	7.8	1.4
2002	364	21.7	2.2	110	6.8	1.3
2003	394	23.0	2.3	140	8.4	1.4
2004	452	26.0	2.4	126	7.5	1.3
2005	452	25.8	2.4	144	8.4	1.4
2006	475	26.5	2.4	139	8.1	1.4
2007	463	25.5	2.3	136	7.9	1.3
2008	570	30.6	2.5	161	9.1	1.4
2009	562	29.6	2.5	150	8.2	1.3
2010	576	29.7	2.5	160	8.6	1.3
2011	597	30.0	2.4	181	9.5	1.4
2012	646	31.7	2.5	178	9.1	1.4
2013	634	30.2	2.4	170	8.3	1.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Ovary

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	104	12.0	2.5	64	7.7	2.0
1977	120	12.1	2.3	73	7.9	1.9
1978	102	10.6	2.2	72	7.6	1.9
1979	129	13.3	2.5	69	6.9	1.7
1980	127	12.6	2.3	96	10.1	2.1
1981	118	10.8	2.0	62	5.7	1.5
1982	115	10.5	2.0	67	6.3	1.6
1983	151	13.1	2.2	86	8.0	1.8
1984	147	12.7	2.1	90	8.1	1.7
1985	146	11.7	2.0	90	7.7	1.6
1986	152	12.5	2.1	84	7.0	1.5
1987	120	9.9	1.8	82	6.8	1.5
1988	150	11.8	1.9	75	6.1	1.4
1989	131	9.8	1.7	85	6.6	1.4
1990	164	12.3	1.9	107	8.4	1.6
1991	138	10.4	1.8	92	7.0	1.5
1992	183	13.3	2.0	123	9.2	1.7
1993	158	10.9	1.7	114	8.6	1.6
1994	192	13.5	1.9	113	8.4	1.6
1995	179	12.1	1.8	122	8.8	1.6
1996	150	10.0	1.6	77	5.3	1.2
1997	199	13.0	1.8	101	7.0	1.4
1998	144	9.4	1.6	102	7.0	1.4
1999	174	11.2	1.7	100	6.6	1.3
2000	163	10.3	1.6	107	7.0	1.3
2001	170	10.3	1.6	108	6.8	1.3
2002	184	10.8	1.6	108	6.7	1.3
2003	158	9.2	1.4	110	6.6	1.2
2004	183	10.5	1.5	106	6.4	1.2
2005	172	9.8	1.5	114	6.7	1.2
2006	155	8.6	1.4	111	6.5	1.2
2007	163	9.0	1.4	107	6.1	1.2
2008	181	9.9	1.5	103	5.6	1.1
2009	191	10.1	1.5	105	5.6	1.1
2010	175	9.2	1.4	114	6.1	1.1
2011	182	9.3	1.4	116	6.2	1.1
2012	209	10.5	1.4	131	6.6	1.1
2013	183	9.0	1.3	105	5.1	1.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Prostate

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	519	119.9	12.0	209	55.0	8.7
1977	550	120.4	11.5	211	53.8	8.2
1978	606	132.3	12.1	226	54.7	8.1
1979	625	130.4	11.5	262	62.3	8.5
1980	758	153.0	12.2	274	65.1	8.6
1981	736	147.0	11.9	314	72.0	8.9
1982	699	134.7	11.1	276	60.7	7.9
1983	711	132.5	10.8	288	60.9	7.7
1984	773	137.7	10.8	286	58.5	7.4
1985	802	135.3	10.4	326	65.8	7.9
1986	844	142.3	10.6	346	68.3	7.9
1987	845	136.1	10.1	348	67.3	7.8
1988	913	149.0	10.7	310	60.1	7.2
1989	953	149.3	10.4	315	56.3	6.8
1990	1088	166.2	10.8	384	69.3	7.6
1991	1254	183.4	11.2	359	64.9	7.3
1992	1551	215.4	11.7	392	68.3	7.4
1993	1722	228.8	11.8	376	66.6	7.3
1994	1843	234.2	11.6	417	71.4	7.4
1995	2033	246.3	11.6	435	69.9	7.1
1996	2065	240.8	11.1	409	63.5	6.6
1997	2187	250.6	11.3	405	61.6	6.5
1998	2269	250.6	11.0	402	60.8	6.3
1999	2575	273.6	11.2	413	60.3	6.2
2000	2850	294.2	11.5	425	60.3	6.1
2001	2717	271.9	10.8	425	59.5	6.0
2002	2769	270.1	10.7	437	57.4	5.7
2003	2768	262.0	10.3	378	48.7	5.2
2004	2786	253.9	10.0	374	48.1	5.2
2005	2815	248.9	9.7	370	44.6	4.8
2006	2989	260.1	9.8	346	42.4	4.7
2007	3240	273.5	9.9	381	45.5	4.8
2008	3149	254.6	9.3	365	42.5	4.5
2009	3149	246.0	9.0	393	44.7	4.6
2010	3116	234.8	8.7	381	41.4	4.3
2011	3392	245.2	8.7	369	38.6	4.1
2012	3040	212.7	7.9	368	37.8	4.0
2013	3038	206.2	7.7	366	36.5	3.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Testis

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	8	0.6	0.4	4	1.0	1.3
1977	9	1.0	0.8	3	0.3	0.3
1978	4	0.5	0.6	4	0.5	0.6
1979	6	0.5	0.4	0	0.0	0.0
1980	15	1.4	0.8	3	0.3	0.3
1981	10	1.0	0.6	2	0.2	0.3
1982	7	0.7	0.5	0	0.0	0.0
1983	7	0.7	0.6	7	1.0	0.9
1984	8	0.6	0.4	1	0.2	0.3
1985	17	1.7	1.0	2	0.1	0.2
1986	17	1.3	0.7	2	0.2	0.2
1987	13	0.8	0.4	2	0.1	0.1
1988	13	0.9	0.5	2	0.4	0.7
1989	15	1.3	0.7	1	0.1	0.1
1990	12	0.9	0.5	3	0.2	0.2
1991	10	0.8	0.6	1	0.1	0.2
1992	22	1.5	0.8	2	0.1	0.2
1993	13	0.9	0.5	3	0.2	0.2
1994	17	1.1	0.6	1	0.1	0.2
1995	7	0.4	0.3	0	0.0	0.0
1996	21	1.3	0.6	0	0.0	0.0
1997	13	0.8	0.4	1	0.1	0.2
1998	21	1.3	0.6	1	0.1	0.1
1999	21	1.3	0.6	1	0.1	0.1
2000	29	2.1	0.8	3	0.3	0.3
2001	23	1.5	0.6	1	0.1	0.3
2002	19	1.2	0.5	2	0.1	0.2
2003	27	1.7	0.6	1	0.1	0.1
2004	16	1.0	0.5	1	0.1	0.1
2005	22	1.4	0.6	2	0.1	0.2
2006	30	1.9	0.7	1	0.1	0.1
2007	30	1.9	0.7	1	0.1	0.1
2008	17	1.1	0.5	1	0.0	0.1
2009	17	1.0	0.5	1	0.1	0.2
2010	23	1.4	0.6	2	0.1	0.2
2011	27	1.6	0.6	3	0.2	0.2
2012	23	1.4	0.6	1	0.1	0.1
2013	27	1.5	0.6	1	0.1	0.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Urinary bladder (incl. in situ)

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	107	24.1	5.4	54	7.7	2.2	39	7.6	2.8	30	4.7	1.8
1977	78	16.2	4.1	54	7.0	2.0	35	9.3	3.6	25	3.5	1.4
1978	94	19.4	4.4	49	5.7	1.7	35	7.5	2.8	27	3.3	1.3
1979	97	19.0	4.3	70	8.8	2.2	40	9.8	3.4	36	4.9	1.7
1980	120	21.4	4.3	61	7.2	1.9	37	8.0	2.9	29	3.7	1.4
1981	90	15.8	3.8	68	7.6	1.9	46	8.8	2.9	32	3.6	1.3
1982	89	16.0	3.7	72	7.5	1.8	37	6.9	2.5	39	4.3	1.4
1983	87	13.9	3.3	75	7.5	1.8	31	5.7	2.2	36	3.9	1.3
1984	109	17.7	3.7	58	5.6	1.5	45	7.8	2.6	28	3.1	1.2
1985	107	17.3	3.7	83	8.4	1.9	43	8.3	2.7	32	3.2	1.2
1986	110	19.0	3.9	59	5.5	1.4	33	6.4	2.4	31	3.0	1.1
1987	104	16.2	3.4	58	5.7	1.5	40	7.2	2.4	32	3.1	1.1
1988	113	16.7	3.4	74	7.1	1.7	40	7.5	2.6	28	2.6	1.0
1989	107	15.2	3.2	71	6.4	1.5	42	7.7	2.5	29	2.9	1.1
1990	116	18.1	3.6	62	5.5	1.4	41	6.3	2.1	33	3.1	1.1
1991	129	18.7	3.6	74	5.9	1.4	40	6.8	2.3	24	2.1	0.9
1992	111	15.6	3.2	80	6.3	1.4	30	4.9	1.9	27	2.3	0.9
1993	100	13.9	3.0	96	7.6	1.5	44	6.6	2.1	35	2.8	0.9
1994	111	15.5	3.1	85	6.5	1.4	36	5.5	2.0	36	3.0	1.0
1995	117	15.7	3.1	60	4.6	1.2	33	5.3	2.0	34	2.7	0.9
1996	118	15.2	3.0	91	6.8	1.4	41	6.1	2.0	33	2.6	0.9
1997	125	16.7	3.2	88	6.6	1.4	35	5.0	1.8	37	2.8	0.9
1998	157	19.4	3.3	101	7.4	1.4	39	5.6	1.9	44	3.3	1.0
1999	148	17.1	3.0	83	5.7	1.2	35	4.4	1.6	34	2.4	0.8
2000	145	16.4	2.9	95	6.4	1.3	43	5.4	1.8	38	2.6	0.8
2001	154	17.0	2.9	98	6.4	1.3	37	4.7	1.7	46	3.1	0.9
2002	141	15.7	2.8	98	6.3	1.3	28	3.3	1.3	38	2.5	0.8
2003	175	17.8	2.8	79	4.9	1.1	39	4.5	1.5	32	2.0	0.7
2004	178	18.3	2.9	99	6.2	1.2	40	4.2	1.4	38	2.4	0.8
2005	187	18.1	2.8	110	6.7	1.3	45	5.5	1.7	35	2.3	0.8
2006	199	20.0	2.9	101	5.9	1.2	39	4.3	1.5	22	1.4	0.6
2007	162	16.0	2.6	100	5.9	1.2	37	4.2	1.4	41	2.5	0.8
2008	197	18.0	2.7	100	5.7	1.1	37	4.2	1.4	30	1.7	0.6
2009	206	18.9	2.7	112	6.4	1.2	42	4.7	1.5	31	1.8	0.6
2010	203	17.9	2.6	96	5.2	1.1	43	4.6	1.4	45	2.5	0.8
2011	250	21.3	2.8	104	5.4	1.1	51	4.9	1.4	37	2.0	0.7
2012	204	17.9	2.6	119	6.1	1.1	53	4.6	1.3	35	1.9	0.6
2013	242	19.9	2.6	133	6.8	1.2	52	4.7	1.3	36	1.8	0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Kidney and renal pelvis

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	58	8.6	2.5	38	5.3	1.8	28	4.4	1.9	19	2.4	1.2
1977	47	7.7	2.6	27	2.5	1.0	20	3.0	1.4	9	0.9	0.7
1978	72	10.4	2.7	39	4.4	1.5	36	4.9	1.9	18	2.1	1.1
1979	65	9.2	2.4	24	2.3	1.0	37	5.7	2.0	10	1.0	0.7
1980	46	6.6	2.1	43	4.1	1.3	28	3.9	1.7	8	0.8	0.5
1981	65	8.4	2.2	28	2.7	1.0	25	3.0	1.2	12	0.9	0.5
1982	65	8.6	2.3	40	3.8	1.2	25	3.4	1.5	21	2.2	1.0
1983	62	8.2	2.3	52	4.6	1.3	30	4.2	1.7	19	1.7	0.8
1984	90	11.5	2.7	51	4.5	1.3	30	4.3	1.7	22	2.0	0.8
1985	70	8.6	2.2	56	4.7	1.3	33	3.9	1.5	25	2.3	0.9
1986	79	9.2	2.2	60	4.9	1.3	43	5.8	1.9	30	2.6	1.0
1987	90	12.1	2.7	57	4.6	1.2	25	3.3	1.4	25	2.2	0.9
1988	75	8.8	2.2	65	5.2	1.3	34	4.6	1.7	22	1.8	0.8
1989	76	9.5	2.4	57	4.2	1.1	20	2.9	1.4	26	2.2	0.9
1990	97	11.0	2.4	72	5.5	1.3	29	3.9	1.6	25	1.9	0.8
1991	119	13.0	2.6	67	5.0	1.2	35	4.5	1.7	25	1.9	0.8
1992	111	12.4	2.6	61	4.6	1.2	32	3.9	1.5	21	1.6	0.7
1993	121	12.7	2.5	64	4.5	1.1	39	4.5	1.6	28	2.2	0.8
1994	97	9.6	2.1	93	6.3	1.3	30	3.4	1.3	21	1.5	0.6
1995	133	14.1	2.6	103	7.3	1.4	46	5.5	1.8	30	2.2	0.8
1996	141	14.0	2.5	76	5.1	1.2	36	3.7	1.3	27	1.9	0.7
1997	131	13.4	2.5	101	6.8	1.3	42	4.8	1.6	29	2.0	0.7
1998	155	14.7	2.5	98	6.4	1.3	34	3.3	1.2	23	1.6	0.7
1999	134	12.8	2.3	130	8.4	1.5	26	2.6	1.1	28	1.9	0.7
2000	129	11.9	2.2	113	7.2	1.3	34	3.4	1.2	28	1.8	0.7
2001	159	14.1	2.3	118	7.2	1.3	37	3.8	1.3	32	2.0	0.7
2002	165	14.6	2.4	123	7.3	1.3	37	3.6	1.2	17	1.1	0.5
2003	185	15.6	2.4	136	8.0	1.4	44	4.4	1.4	26	1.6	0.6
2004	198	15.6	2.3	144	8.3	1.4	43	4.0	1.3	30	1.8	0.7
2005	188	14.9	2.2	168	9.6	1.5	44	3.7	1.2	23	1.3	0.6
2006	241	19.4	2.6	151	8.5	1.4	42	3.8	1.2	36	2.1	0.7
2007	265	20.9	2.7	156	8.7	1.4	57	5.2	1.4	28	1.6	0.6
2008	227	17.5	2.4	193	10.7	1.5	45	3.6	1.1	30	1.7	0.6
2009	271	19.8	2.5	193	10.3	1.5	46	3.8	1.2	25	1.3	0.5
2010	317	23.4	2.7	184	9.8	1.4	42	3.8	1.2	34	1.9	0.6
2011	293	21.6	2.6	182	9.5	1.4	45	3.8	1.2	29	1.5	0.6
2012	272	19.1	2.4	202	10.3	1.4	43	3.6	1.2	33	1.6	0.6
2013	329	21.9	2.5	206	10.1	1.4	39	2.8	0.9	22	1.1	0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Brain and other nervous system

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	35	4.1	1.4	31	2.8	1.0	29	3.5	1.4	23	2.3	1.0
1977	39	4.5	1.5	45	3.6	1.1	23	2.9	1.2	19	1.8	0.9
1978	24	3.1	1.3	27	2.4	1.0	20	2.2	1.0	20	1.9	0.9
1979	37	4.1	1.4	38	3.1	1.0	25	3.2	1.4	26	2.3	0.9
1980	47	5.6	1.8	29	2.7	1.0	30	3.8	1.5	17	1.6	0.8
1981	29	2.7	1.0	25	2.1	0.9	16	2.0	1.0	19	2.0	1.0
1982	45	4.3	1.3	33	2.5	0.9	41	5.1	1.7	24	2.1	0.9
1983	41	4.1	1.4	34	2.7	1.0	23	2.6	1.1	38	3.1	1.0
1984	54	5.1	1.5	45	3.3	1.0	43	4.5	1.5	27	2.1	0.8
1985	43	5.0	1.6	56	4.1	1.1	32	3.6	1.3	24	1.7	0.7
1986	55	5.4	1.6	39	3.0	1.0	33	3.4	1.2	30	2.4	0.9
1987	48	4.6	1.5	53	3.8	1.1	28	3.5	1.5	27	2.2	0.9
1988	51	4.7	1.5	37	2.5	0.9	23	2.4	1.1	25	1.7	0.7
1989	41	3.6	1.2	51	3.5	1.0	19	1.8	0.9	21	1.5	0.7
1990	46	4.3	1.4	56	3.8	1.0	24	2.8	1.3	22	1.6	0.7
1991	63	6.0	1.7	65	4.3	1.1	22	2.2	1.0	31	2.3	0.8
1992	66	6.1	1.6	64	4.3	1.1	25	2.9	1.3	30	2.2	0.8
1993	53	4.5	1.3	39	2.6	0.8	29	2.8	1.1	33	2.1	0.7
1994	57	4.6	1.3	50	3.2	0.9	28	2.6	1.1	31	2.0	0.7
1995	49	3.9	1.2	65	4.2	1.0	23	1.9	0.8	27	1.8	0.7
1996	64	5.5	1.4	74	4.6	1.1	24	2.3	1.0	27	1.7	0.7
1997	58	4.7	1.3	52	3.1	0.9	24	1.9	0.8	25	1.6	0.6
1998	59	4.1	1.1	59	3.6	0.9	25	2.2	0.9	20	1.3	0.6
1999	67	5.4	1.4	58	3.5	0.9	26	2.2	0.9	16	1.0	0.5
2000	58	4.6	1.3	64	3.7	0.9	26	2.2	0.9	28	1.8	0.7
2001	61	4.4	1.1	85	5.0	1.1	28	2.5	1.0	35	2.1	0.7
2002	63	4.8	1.3	63	3.6	0.9	26	1.9	0.7	36	2.1	0.7
2003	73	5.3	1.3	74	4.3	1.0	33	2.6	0.9	22	1.3	0.5
2004	67	5.0	1.2	62	3.5	0.9	31	2.4	0.9	31	1.8	0.6
2005	67	5.4	1.4	61	3.4	0.9	32	2.5	0.9	17	0.9	0.5
2006	56	4.0	1.1	58	3.2	0.8	33	2.6	1.0	26	1.4	0.6
2007	65	4.4	1.1	63	3.4	0.9	21	1.8	0.8	28	1.5	0.6
2008	57	4.0	1.1	65	3.5	0.9	31	2.4	0.9	32	1.7	0.6
2009	66	4.7	1.2	62	3.4	0.8	30	2.4	0.9	26	1.4	0.6
2010	74	5.0	1.2	63	3.3	0.8	28	2.1	0.8	26	1.4	0.5
2011	74	5.0	1.2	70	3.6	0.8	37	2.6	0.9	35	1.8	0.6
2012	78	5.2	1.2	56	2.9	0.8	33	2.3	0.8	32	1.6	0.6
2013	75	4.5	1.1	84	4.3	0.9	38	2.6	0.9	35	1.8	0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Thyroid

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	10	1.4	1.0	33	3.1	1.1	3	0.4	0.5	7	0.8	0.6
1977	3	0.4	0.4	42	3.8	1.2	3	0.4	0.4	3	0.5	0.5
1978	11	1.6	1.1	26	2.5	1.0	2	0.4	0.6	9	1.1	0.7
1979	11	1.6	1.2	36	3.7	1.3	2	0.2	0.3	7	0.8	0.7
1980	5	0.4	0.4	39	3.2	1.1	1	0.2	0.4	7	0.7	0.5
1981	6	0.6	0.5	28	2.7	1.0	1	0.1	0.2	6	0.7	0.6
1982	7	0.8	0.6	36	3.2	1.1	4	0.8	0.9	7	0.6	0.5
1983	5	0.5	0.4	40	3.2	1.0	1	0.2	0.5	5	0.5	0.5
1984	8	0.8	0.5	46	3.6	1.1	2	0.5	0.8	7	0.6	0.5
1985	20	2.1	1.0	48	3.8	1.1	4	0.5	0.5	4	0.4	0.4
1986	13	1.8	1.1	32	2.5	0.9	4	0.7	0.8	7	0.5	0.4
1987	13	1.5	0.9	51	3.8	1.1	3	0.6	0.7	9	0.8	0.5
1988	11	1.3	0.8	38	2.8	0.9	3	0.4	0.4	5	0.5	0.4
1989	5	0.5	0.5	43	3.1	1.0	1	0.2	0.3	10	1.0	0.6
1990	9	0.8	0.6	48	3.3	1.0	1	0.1	0.3	4	0.4	0.4
1991	8	1.1	0.8	58	4.0	1.0	2	0.4	0.6	8	0.6	0.5
1992	22	1.8	0.8	68	4.3	1.0	3	0.3	0.3	8	0.6	0.4
1993	17	1.5	0.7	62	4.1	1.0	3	0.3	0.4	5	0.4	0.4
1994	13	1.2	0.7	76	4.9	1.1	2	0.2	0.3	6	0.4	0.3
1995	15	1.4	0.7	82	5.1	1.1	1	0.1	0.2	7	0.6	0.4
1996	23	2.2	1.0	93	5.5	1.1	9	1.0	0.7	6	0.4	0.4
1997	27	2.4	1.0	97	5.8	1.2	8	0.8	0.6	6	0.4	0.3
1998	26	2.3	0.9	107	6.3	1.2	4	0.4	0.4	6	0.4	0.3
1999	18	1.4	0.7	107	6.3	1.2	2	0.2	0.2	13	0.9	0.5
2000	24	1.7	0.7	139	8.0	1.3	2	0.2	0.2	12	0.8	0.4
2001	35	2.7	0.9	132	7.4	1.3	3	0.2	0.3	14	0.9	0.5
2002	30	2.3	0.9	142	7.9	1.3	2	0.1	0.2	7	0.5	0.3
2003	31	2.5	0.9	135	7.4	1.2	1	0.1	0.1	7	0.5	0.3
2004	33	3.0	1.1	152	8.3	1.3	7	0.8	0.6	9	0.6	0.4
2005	44	3.2	1.0	212	11.5	1.6	9	0.8	0.5	12	0.7	0.4
2006	35	2.8	1.0	188	10.2	1.5	3	0.2	0.2	8	0.5	0.3
2007	50	3.7	1.1	188	10.1	1.5	3	0.3	0.4	9	0.5	0.4
2008	60	4.4	1.2	227	12.0	1.6	5	0.5	0.5	10	0.6	0.4
2009	59	4.0	1.1	240	12.9	1.6	5	0.5	0.4	10	0.5	0.3
2010	49	3.5	1.0	281	14.4	1.7	11	0.9	0.6	10	0.5	0.3
2011	77	5.3	1.2	307	15.2	1.7	1	0.1	0.3	11	0.6	0.3
2012	56	3.8	1.0	294	14.9	1.7	6	0.6	0.5	12	0.6	0.3
2013	53	3.3	0.9	369	18.2	1.9	5	0.4	0.3	14	0.7	0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Hodgkin lymphoma

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	21	3.3	2.0	25	2.4	1.0	10	1.1	0.7	8	0.8	0.5
1977	20	2.1	1.0	18	1.5	0.7	7	1.3	1.3	12	1.1	0.7
1978	24	2.4	1.0	18	1.5	0.8	8	0.9	0.6	6	0.4	0.4
1979	30	3.0	1.2	18	1.4	0.7	14	1.5	0.8	7	0.7	0.6
1980	32	2.7	1.0	28	1.9	0.7	6	0.7	0.6	7	0.5	0.4
1981	33	3.1	1.2	21	1.5	0.7	13	1.6	1.0	9	0.7	0.5
1982	32	3.0	1.1	26	1.9	0.8	9	0.9	0.6	9	0.7	0.4
1983	39	3.7	1.4	28	1.9	0.7	8	0.7	0.5	6	0.5	0.4
1984	39	3.6	1.2	19	1.4	0.6	19	1.7	0.8	2	0.2	0.3
1985	31	2.6	1.0	19	1.1	0.5	18	1.6	0.8	9	0.6	0.4
1986	47	3.9	1.2	20	1.3	0.6	13	1.2	0.7	6	0.5	0.4
1987	36	3.3	1.2	27	1.6	0.6	14	1.4	0.8	3	0.2	0.2
1988	43	2.8	0.9	26	1.7	0.7	9	0.8	0.5	8	0.5	0.3
1989	43	3.3	1.1	29	1.8	0.7	16	1.2	0.6	3	0.3	0.3
1990	38	2.9	1.0	29	1.8	0.7	12	1.0	0.6	8	0.5	0.3
1991	54	4.2	1.3	39	2.2	0.7	8	0.6	0.4	10	0.6	0.4
1992	36	2.5	0.8	30	1.8	0.7	12	0.8	0.5	10	0.6	0.4
1993	38	2.5	0.8	34	1.9	0.6	10	0.7	0.4	7	0.4	0.3
1994	36	2.8	1.0	24	1.4	0.6	13	1.1	0.6	7	0.4	0.3
1995	49	3.3	1.0	33	1.8	0.6	13	0.9	0.5	4	0.2	0.2
1996	34	2.4	0.9	35	2.0	0.7	7	0.6	0.5	6	0.4	0.3
1997	51	3.3	0.9	29	1.6	0.6	6	0.4	0.3	10	0.7	0.4
1998	44	2.9	0.9	38	2.1	0.7	5	0.3	0.3	7	0.4	0.3
1999	43	3.0	1.0	37	2.0	0.7	9	0.7	0.5	5	0.3	0.3
2000	56	3.6	1.0	35	1.8	0.6	3	0.2	0.2	2	0.1	0.2
2001	54	3.3	0.9	29	1.5	0.6	6	0.4	0.4	6	0.3	0.3
2002	49	3.3	1.0	40	2.2	0.7	8	0.5	0.4	2	0.1	0.1
2003	35	2.2	0.8	49	2.6	0.7	8	0.8	0.6	6	0.3	0.3
2004	48	3.0	0.9	46	2.4	0.7	6	0.4	0.4	3	0.2	0.2
2005	60	3.8	1.0	39	2.1	0.7	9	0.6	0.4	2	0.1	0.1
2006	48	3.2	0.9	41	2.2	0.7	5	0.3	0.3	4	0.2	0.2
2007	50	3.4	1.0	37	1.9	0.6	4	0.3	0.3	9	0.5	0.3
2008	71	4.5	1.1	51	2.7	0.7	5	0.3	0.3	3	0.2	0.2
2009	45	3.0	0.9	57	3.0	0.8	6	0.4	0.4	4	0.2	0.2
2010	68	4.4	1.1	50	2.6	0.7	5	0.3	0.2	4	0.2	0.2
2011	68	3.8	0.9	52	2.6	0.7	8	0.6	0.4	6	0.3	0.2
2012	48	2.9	0.9	55	2.8	0.8	4	0.3	0.3	1	0.1	0.1
2013	59	3.7	1.0	54	2.8	0.8	5	0.4	0.3	5	0.2	0.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Non-Hodgkin lymphomas

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	49	7.7	2.5	49	5.1	1.5	34	6.0	2.3	23	2.6	1.2
1977	57	8.5	2.6	61	6.1	1.6	28	4.2	1.9	33	3.5	1.3
1978	84	10.8	2.5	72	7.4	1.8	41	5.6	1.9	24	2.3	1.0
1979	64	8.5	2.3	51	5.4	1.6	42	6.0	2.1	37	4.5	1.6
1980	69	9.6	2.7	64	6.2	1.6	42	5.7	1.9	31	3.3	1.2
1981	66	7.7	2.2	65	7.0	1.8	34	4.8	1.8	44	4.1	1.3
1982	81	9.8	2.3	79	7.2	1.7	30	4.3	1.8	31	3.2	1.2
1983	85	9.8	2.3	87	7.7	1.7	49	5.8	1.8	42	4.0	1.3
1984	96	11.0	2.5	95	8.1	1.7	54	6.3	1.9	41	3.7	1.2
1985	125	13.5	2.7	107	8.8	1.7	55	6.0	1.8	53	4.7	1.3
1986	120	13.8	2.9	120	9.8	1.8	52	6.0	1.9	50	4.1	1.2
1987	150	16.4	3.0	102	7.7	1.5	50	5.6	1.8	45	3.5	1.1
1988	136	14.4	2.7	94	7.2	1.5	76	9.9	2.6	41	3.3	1.0
1989	155	14.4	2.5	117	8.7	1.6	67	7.2	1.9	48	3.7	1.1
1990	184	16.2	2.5	126	9.2	1.7	55	5.9	1.8	51	3.9	1.1
1991	180	16.3	2.7	134	9.9	1.7	62	6.1	1.7	47	3.8	1.1
1992	208	18.9	2.8	150	10.7	1.8	60	6.3	1.8	54	4.0	1.1
1993	225	18.9	2.6	136	8.9	1.5	55	5.1	1.4	62	4.5	1.1
1994	231	20.1	2.8	171	11.8	1.8	57	5.7	1.6	58	4.3	1.1
1995	237	21.0	2.9	145	9.8	1.6	65	5.8	1.5	53	3.9	1.1
1996	226	18.6	2.6	185	12.2	1.8	58	6.0	1.8	67	4.8	1.2
1997	249	22.0	3.0	183	12.0	1.8	60	6.3	1.7	56	3.9	1.0
1998	240	20.8	2.8	193	12.2	1.7	73	7.3	1.8	69	4.7	1.1
1999	253	21.3	2.8	208	12.8	1.8	79	7.5	1.8	67	4.3	1.0
2000	209	16.8	2.4	192	11.3	1.6	84	7.8	1.8	57	3.6	0.9
2001	233	18.8	2.6	200	12.0	1.7	78	7.0	1.7	75	4.7	1.1
2002	232	17.8	2.4	194	11.2	1.6	87	7.6	1.7	66	4.0	1.0
2003	212	16.9	2.4	249	14.2	1.8	71	6.1	1.5	65	3.8	0.9
2004	213	17.5	2.5	204	11.6	1.6	65	5.5	1.4	58	3.4	0.9
2005	240	18.4	2.4	238	13.4	1.7	63	5.7	1.5	60	3.5	0.9
2006	225	17.3	2.4	231	12.9	1.7	54	4.4	1.3	61	3.4	0.9
2007	269	20.1	2.5	210	11.6	1.6	64	5.5	1.4	67	3.8	0.9
2008	282	20.7	2.5	213	11.5	1.6	60	5.0	1.3	54	2.9	0.8
2009	253	19.0	2.5	259	14.2	1.7	63	5.0	1.3	57	3.1	0.8
2010	260	18.4	2.4	239	12.6	1.6	61	4.8	1.3	66	3.6	0.9
2011	304	22.2	2.6	258	13.1	1.6	76	6.0	1.4	58	3.1	0.8
2012	248	17.5	2.3	260	13.3	1.6	63	5.0	1.3	60	3.0	0.8
2013	259	17.5	2.2	250	12.4	1.6	64	4.9	1.3	59	2.9	0.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Myeloma

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	58	10.9	3.1	65	8.1	2.1	30	4.9	1.9	42	5.6	1.8
1977	54	10.4	3.3	68	8.4	2.1	40	8.9	3.2	41	5.1	1.7
1978	64	13.1	3.7	56	6.2	1.7	34	7.3	2.9	27	2.9	1.1
1979	63	10.3	2.9	78	8.6	2.0	31	5.4	2.1	41	4.7	1.5
1980	61	10.7	3.0	92	9.8	2.1	43	7.5	2.5	56	6.1	1.7
1981	64	10.7	2.9	88	9.4	2.1	34	6.5	2.4	48	5.1	1.5
1982	78	12.2	3.0	87	9.2	2.0	58	9.1	2.6	55	5.7	1.6
1983	64	9.4	2.5	81	8.1	1.8	41	7.1	2.4	61	6.2	1.6
1984	81	12.6	3.1	100	9.8	2.0	43	6.8	2.3	73	7.3	1.7
1985	75	11.6	3.0	97	8.7	1.8	54	8.2	2.4	55	5.1	1.4
1986	86	12.9	3.0	101	9.3	1.9	56	8.7	2.5	58	5.7	1.5
1987	79	11.5	2.8	85	7.4	1.6	53	8.2	2.5	46	4.1	1.2
1988	68	9.1	2.4	86	7.3	1.6	39	5.5	1.9	44	3.9	1.2
1989	86	12.3	2.9	105	9.3	1.8	39	6.5	2.3	51	4.5	1.3
1990	85	12.1	2.9	108	8.5	1.6	42	6.5	2.2	56	4.8	1.3
1991	73	10.0	2.6	131	10.5	1.8	45	6.7	2.2	65	5.3	1.3
1992	106	13.8	2.9	129	9.9	1.7	53	7.8	2.4	79	6.1	1.4
1993	81	9.8	2.3	106	8.0	1.5	47	6.5	2.0	67	5.1	1.2
1994	87	11.1	2.5	118	9.1	1.7	43	5.4	1.8	76	5.9	1.4
1995	96	12.2	2.7	119	8.7	1.6	59	7.5	2.1	79	5.9	1.3
1996	96	10.7	2.3	125	8.9	1.6	55	6.9	2.0	54	3.9	1.1
1997	110	12.1	2.4	138	9.8	1.6	70	8.7	2.2	64	4.6	1.1
1998	108	12.0	2.4	138	9.6	1.6	41	4.9	1.7	71	5.0	1.2
1999	103	10.6	2.2	151	10.3	1.7	48	5.3	1.6	73	5.0	1.2
2000	118	11.9	2.3	158	10.2	1.6	42	4.8	1.6	74	5.0	1.2
2001	128	12.8	2.4	170	10.9	1.7	54	5.9	1.7	82	5.3	1.2
2002	118	12.1	2.4	172	10.7	1.6	59	5.6	1.6	88	5.7	1.2
2003	135	13.4	2.4	179	11.0	1.6	55	5.1	1.5	69	4.3	1.0
2004	128	12.0	2.2	174	10.5	1.6	63	6.7	1.8	73	4.6	1.1
2005	156	14.1	2.3	202	11.9	1.7	51	4.7	1.4	74	4.4	1.0
2006	169	15.1	2.4	185	10.7	1.6	60	5.6	1.5	65	3.9	0.9
2007	140	12.8	2.2	166	9.5	1.5	70	6.4	1.6	83	4.8	1.1
2008	188	15.7	2.4	222	12.1	1.6	53	4.9	1.4	65	3.7	0.9
2009	216	17.4	2.4	204	11.3	1.6	71	5.9	1.4	58	3.2	0.8
2010	207	17.6	2.5	202	10.8	1.5	69	6.3	1.6	82	4.5	1.0
2011	203	15.7	2.3	241	12.7	1.6	62	5.2	1.4	79	4.1	0.9
2012	215	16.5	2.3	244	12.3	1.6	72	5.4	1.3	72	3.7	0.9
2013	224	17.0	2.3	223	11.1	1.5	75	6.1	1.5	69	3.5	0.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State
Blacks
Leukemias

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	70	9.9	2.8	62	6.3	1.7	53	7.8	2.7	39	4.4	1.5
1977	67	11.3	3.5	59	6.1	1.7	54	8.4	2.7	47	4.6	1.4
1978	76	10.9	3.0	80	8.3	2.0	62	9.0	2.6	53	6.0	1.8
1979	72	11.3	3.2	71	7.3	1.8	56	9.0	2.8	42	4.0	1.3
1980	74	10.5	2.9	85	8.2	1.9	49	7.6	2.7	63	6.1	1.6
1981	83	12.5	3.2	73	7.1	1.7	46	7.4	2.7	62	6.0	1.6
1982	78	11.0	2.9	78	6.9	1.6	44	6.5	2.3	46	4.2	1.3
1983	90	12.2	2.9	83	7.3	1.6	50	7.1	2.3	55	4.9	1.4
1984	86	9.8	2.4	79	7.0	1.6	54	7.1	2.2	51	4.5	1.3
1985	111	13.4	2.8	82	6.8	1.5	61	8.2	2.4	39	3.2	1.1
1986	96	12.0	2.8	75	6.0	1.4	61	6.9	2.0	49	4.0	1.2
1987	94	10.8	2.5	96	7.8	1.6	51	6.2	2.1	54	4.4	1.2
1988	77	10.7	2.7	79	6.2	1.4	55	7.5	2.3	61	5.2	1.4
1989	105	12.6	2.8	98	7.6	1.6	61	8.4	2.4	39	3.1	1.0
1990	108	11.8	2.6	111	8.6	1.7	59	7.6	2.2	69	5.4	1.3
1991	117	13.1	2.8	121	8.9	1.6	55	6.6	2.1	51	4.0	1.1
1992	104	11.3	2.6	125	9.1	1.6	48	5.4	1.7	59	4.7	1.2
1993	104	11.8	2.6	102	7.4	1.5	55	6.2	1.9	56	4.1	1.1
1994	109	12.0	2.5	102	7.2	1.4	74	8.4	2.2	82	6.1	1.4
1995	122	11.8	2.3	109	7.5	1.4	66	7.4	2.0	69	4.7	1.1
1996	138	13.3	2.5	111	7.5	1.4	66	7.0	1.9	73	5.2	1.2
1997	100	9.6	2.1	124	8.4	1.5	72	8.2	2.1	62	4.3	1.1
1998	105	10.7	2.3	121	7.9	1.4	59	6.1	1.7	64	4.3	1.1
1999	106	10.7	2.3	99	6.1	1.2	65	7.0	1.9	64	4.2	1.0
2000	130	11.9	2.2	128	8.1	1.4	53	5.6	1.7	72	4.6	1.1
2001	108	9.8	2.0	127	7.9	1.4	65	6.4	1.7	72	4.6	1.1
2002	145	12.9	2.3	123	7.5	1.3	61	5.9	1.6	63	3.9	1.0
2003	129	10.5	1.9	133	7.8	1.3	57	4.9	1.4	71	4.4	1.0
2004	160	13.7	2.3	121	7.1	1.3	70	6.9	1.7	63	3.9	1.0
2005	119	10.3	2.0	117	6.7	1.2	62	6.0	1.6	68	4.0	1.0
2006	165	13.7	2.2	118	6.7	1.2	66	5.8	1.5	53	3.1	0.8
2007	131	10.3	1.9	139	7.9	1.3	68	6.3	1.6	60	3.5	0.9
2008	131	11.0	2.0	136	7.5	1.3	74	6.6	1.6	65	3.6	0.9
2009	151	11.8	2.0	124	6.8	1.2	65	5.1	1.3	52	2.9	0.8
2010	189	14.3	2.2	152	8.0	1.3	74	6.5	1.6	63	3.4	0.8
2011	179	13.1	2.0	155	8.2	1.3	85	6.5	1.5	74	3.9	0.9
2012	173	12.8	2.0	160	8.2	1.3	77	6.2	1.5	60	3.1	0.8
2013	171	12.0	1.9	181	9.1	1.3	85	6.7	1.5	56	2.8	0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
All Invasive Malignant Tumors

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	2066	482.0	24.2	2165	313.3	14.2	1398	330.6	20.3	1265	194.3	11.6
1977	2165	508.3	24.9	2296	319.5	14.0	1394	326.4	19.8	1237	176.6	10.6
1978	2293	519.6	24.5	2414	337.2	14.4	1462	327.9	19.3	1423	205.2	11.4
1979	2432	540.4	24.4	2512	337.7	14.0	1579	360.1	20.4	1380	193.3	10.9
1980	2644	574.0	24.8	2650	341.0	13.7	1597	356.6	20.0	1451	194.2	10.6
1981	2596	539.6	23.5	2589	324.9	13.2	1629	347.9	19.2	1504	193.4	10.3
1982	2672	546.3	23.2	2758	335.9	13.1	1662	349.8	18.9	1485	184.8	9.9
1983	2665	523.0	22.3	2823	327.4	12.5	1755	349.1	18.2	1593	190.8	9.8
1984	2875	546.2	22.4	3080	349.8	12.8	1689	337.2	18.0	1580	185.4	9.5
1985	3014	560.0	22.3	3098	342.3	12.5	1734	342.4	18.1	1557	178.3	9.2
1986	3140	573.8	22.4	3104	332.6	12.1	1693	325.9	17.3	1633	180.3	9.0
1987	3137	550.9	21.4	3201	334.1	11.9	1705	323.1	17.1	1617	174.2	8.7
1988	3061	541.4	21.4	3283	336.5	11.8	1671	309.6	16.5	1638	172.8	8.6
1989	3258	564.1	21.5	3280	328.9	11.5	1672	307.1	16.3	1577	162.4	8.2
1990	3309	565.1	21.3	3457	342.1	11.7	1679	309.0	16.4	1704	174.8	8.5
1991	3506	590.6	21.7	3717	363.6	12.0	1677	302.4	16.1	1636	164.8	8.2
1992	3679	604.3	21.5	3776	361.6	11.8	1570	286.3	15.7	1723	169.8	8.2
1993	3745	600.8	21.1	3601	339.0	11.3	1584	281.0	15.3	1661	161.4	7.9
1994	3983	623.3	21.1	3941	366.2	11.6	1749	301.7	15.5	1886	180.8	8.3
1995	3993	606.2	20.3	3782	344.6	11.1	1739	292.2	15.0	1915	180.2	8.2
1996	4066	608.2	20.2	4117	368.1	11.4	1755	290.7	14.8	1812	168.0	7.8
1997	4143	617.8	20.2	4161	367.8	11.3	1683	277.5	14.3	1829	166.8	7.7
1998	4069	585.7	19.2	4351	376.4	11.3	1595	251.1	13.3	1852	165.8	7.6
1999	4310	609.4	19.4	4241	357.9	10.9	1640	257.5	13.4	1912	166.2	7.5
2000	4521	618.8	19.1	4356	358.6	10.7	1619	246.0	12.8	1814	153.5	7.1
2001	4459	596.5	18.6	4297	347.7	10.5	1628	244.0	12.7	1943	161.3	7.2
2002	4441	581.3	18.1	4416	351.7	10.4	1616	235.2	12.3	1811	147.8	6.8
2003	4572	587.6	18.0	4559	356.5	10.4	1525	215.0	11.5	1789	143.0	6.7
2004	4692	587.7	17.7	4630	358.1	10.4	1574	218.4	11.5	1792	142.1	6.6
2005	4766	584.5	17.5	4737	362.5	10.4	1490	204.3	11.0	1860	145.1	6.6
2006	4833	588.6	17.5	4823	365.3	10.4	1499	205.9	11.1	1798	139.2	6.5
2007	5146	608.6	17.5	4778	359.7	10.3	1540	203.8	10.8	1823	139.3	6.4
2008	5030	582.8	16.9	5020	371.9	10.4	1492	197.1	10.6	1815	136.2	6.3
2009	5189	584.9	16.7	5066	369.6	10.3	1556	200.0	10.5	1736	128.0	6.1
2010	5131	562.4	16.2	5171	371.6	10.2	1563	197.3	10.3	1800	130.1	6.1
2011	5536	587.3	16.2	5544	388.7	10.4	1603	193.3	10.0	1977	140.3	6.3
2012	5125	533.8	15.3	5475	378.5	10.2	1585	185.7	9.6	1943	135.1	6.1
2013	5212	526.0	15.0	5615	381.7	10.1	1584	182.0	9.4	1819	122.9	5.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Oral cavity and pharynx

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	132	26.2	5.1	60	7.8	2.1	62	12.4	3.3	30	4.0	1.5
1977	142	27.3	4.9	46	5.9	1.8	83	15.5	3.5	20	2.9	1.4
1978	147	26.0	4.6	58	7.1	1.9	80	15.9	4.0	27	3.2	1.2
1979	156	29.0	4.9	57	7.2	2.0	78	14.6	3.6	31	3.8	1.4
1980	154	27.8	4.7	59	7.0	1.8	85	15.8	3.8	18	2.1	1.0
1981	167	27.4	4.4	57	6.8	1.8	63	11.1	3.0	25	2.9	1.2
1982	156	26.6	4.5	71	7.9	1.9	89	15.9	3.6	28	3.1	1.2
1983	153	25.8	4.4	58	6.6	1.7	86	15.4	3.6	24	2.8	1.2
1984	170	26.2	4.1	88	9.4	2.0	68	11.6	3.0	23	2.4	1.0
1985	165	25.8	4.2	82	8.7	1.9	57	9.7	2.7	25	2.8	1.1
1986	180	27.1	4.2	71	7.3	1.7	72	12.3	3.1	28	3.1	1.2
1987	190	28.1	4.2	80	7.9	1.8	57	8.9	2.5	19	1.9	0.8
1988	150	22.2	3.8	81	7.8	1.7	71	10.4	2.5	25	2.5	1.0
1989	148	21.1	3.6	75	6.9	1.6	65	9.7	2.6	26	2.5	1.0
1990	162	24.2	4.0	76	7.3	1.7	52	9.0	2.8	22	2.1	0.9
1991	211	31.9	4.6	93	8.6	1.8	77	11.1	2.6	30	3.1	1.1
1992	184	26.4	4.0	97	9.0	1.8	53	7.8	2.2	27	2.5	1.0
1993	173	24.1	3.8	94	8.5	1.7	60	8.6	2.3	23	2.2	0.9
1994	175	24.4	3.8	81	7.3	1.6	66	9.9	2.6	25	2.3	0.9
1995	162	21.4	3.4	58	5.1	1.3	71	10.0	2.5	28	2.6	1.0
1996	153	20.2	3.3	86	7.4	1.6	59	8.3	2.3	22	2.0	0.9
1997	150	19.1	3.2	71	6.2	1.5	49	6.4	1.9	16	1.4	0.7
1998	138	17.1	3.0	75	6.4	1.5	45	6.2	1.9	25	2.2	0.9
1999	118	14.4	2.7	68	5.7	1.4	35	4.7	1.6	24	2.1	0.8
2000	141	17.7	3.1	67	5.5	1.3	45	6.0	1.9	18	1.5	0.7
2001	132	15.9	2.8	56	4.4	1.1	51	6.3	1.8	19	1.5	0.7
2002	126	14.9	2.7	72	5.7	1.3	34	3.9	1.3	16	1.3	0.6
2003	130	14.5	2.6	58	4.6	1.2	37	4.6	1.6	13	1.0	0.6
2004	139	15.8	2.8	82	6.2	1.3	53	6.3	1.8	18	1.4	0.7
2005	141	16.2	2.8	59	4.5	1.1	44	5.4	1.7	19	1.4	0.6
2006	119	13.1	2.5	69	5.1	1.2	33	3.7	1.3	9	0.7	0.4
2007	141	15.4	2.7	81	6.1	1.3	46	5.3	1.6	13	1.0	0.5
2008	126	13.2	2.4	62	4.5	1.1	42	4.9	1.6	15	1.1	0.6
2009	124	12.8	2.3	66	4.8	1.2	37	4.0	1.3	17	1.2	0.6
2010	131	13.6	2.4	73	5.1	1.2	26	2.9	1.2	24	1.7	0.7
2011	141	13.8	2.4	64	4.4	1.1	50	5.1	1.5	22	1.5	0.6
2012	130	12.6	2.3	76	5.1	1.2	32	3.4	1.2	20	1.4	0.6
2013	139	13.3	2.3	66	4.5	1.1	45	4.8	1.5	21	1.4	0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Esophagus

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	116	22.6	4.5	45	5.9	1.8	118	24.4	4.9	38	5.2	1.8
1977	119	24.7	5.1	51	6.6	1.9	107	22.1	4.9	41	5.2	1.6
1978	128	26.2	5.1	53	6.6	1.8	83	15.9	3.7	42	5.3	1.7
1979	127	23.7	4.5	61	7.5	1.9	125	24.7	4.9	48	5.9	1.7
1980	150	28.9	5.1	51	6.1	1.7	108	21.3	4.5	58	6.9	1.9
1981	144	25.1	4.4	63	7.2	1.8	123	21.0	3.9	44	5.1	1.5
1982	143	25.8	4.6	55	6.0	1.6	117	22.0	4.4	50	5.6	1.6
1983	134	22.1	4.0	61	6.9	1.8	113	18.8	3.6	49	5.6	1.6
1984	139	23.6	4.3	70	7.5	1.8	112	19.3	3.9	48	5.3	1.5
1985	126	20.7	3.9	62	6.7	1.7	126	21.1	4.0	46	4.9	1.5
1986	141	23.8	4.2	47	4.9	1.4	114	18.7	3.7	51	5.5	1.5
1987	115	19.3	3.8	66	6.7	1.7	99	16.4	3.5	39	4.1	1.3
1988	143	24.0	4.3	58	5.8	1.5	112	19.0	3.8	36	3.7	1.2
1989	132	21.0	3.8	60	5.9	1.5	99	16.0	3.4	43	4.5	1.4
1990	119	19.4	3.8	54	5.6	1.5	86	14.0	3.2	49	4.8	1.4
1991	102	16.5	3.5	52	4.9	1.4	85	13.7	3.2	43	4.1	1.3
1992	110	17.1	3.5	59	5.7	1.5	73	11.0	2.7	37	3.6	1.2
1993	122	17.9	3.4	61	5.8	1.5	95	15.2	3.4	43	4.1	1.2
1994	115	18.2	3.7	60	5.7	1.5	88	13.7	3.1	45	4.3	1.3
1995	95	13.9	3.0	47	4.5	1.3	84	13.0	3.0	40	3.9	1.2
1996	78	11.5	2.7	60	5.7	1.5	80	12.1	2.8	43	4.0	1.2
1997	100	14.0	2.9	44	4.1	1.2	75	11.4	2.7	37	3.5	1.1
1998	88	12.2	2.7	40	3.5	1.1	59	8.4	2.3	36	3.3	1.1
1999	88	12.2	2.7	42	3.5	1.1	70	9.4	2.3	30	2.5	0.9
2000	67	9.6	2.4	67	5.8	1.4	65	9.0	2.3	47	4.0	1.1
2001	83	10.6	2.4	38	3.1	1.0	67	9.2	2.3	24	2.0	0.8
2002	69	9.1	2.3	48	3.9	1.1	47	6.7	2.0	34	2.8	0.9
2003	81	10.9	2.5	48	3.8	1.1	54	6.4	1.8	37	2.9	0.9
2004	77	9.2	2.2	46	3.7	1.1	54	6.6	1.8	42	3.4	1.0
2005	56	6.5	1.8	51	3.9	1.1	43	5.1	1.6	29	2.3	0.8
2006	64	8.2	2.1	28	2.3	0.8	52	6.9	2.0	37	2.8	0.9
2007	63	7.4	1.9	28	2.1	0.8	44	4.9	1.5	21	1.6	0.7
2008	75	9.2	2.2	36	2.6	0.9	52	6.5	1.9	32	2.4	0.8
2009	68	8.0	2.0	36	2.7	0.9	58	6.6	1.8	16	1.2	0.6
2010	49	5.4	1.6	36	2.6	0.9	42	5.4	1.7	25	1.8	0.7
2011	72	7.9	1.9	34	2.4	0.8	45	4.8	1.5	21	1.5	0.7
2012	51	5.8	1.7	33	2.2	0.7	37	3.8	1.3	19	1.2	0.6
2013	54	6.1	1.7	26	1.7	0.7	29	3.1	1.2	20	1.4	0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Stomach

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	101	24.6	5.6	78	13.1	3.2	88	22.6	5.5	60	9.5	2.5
1977	115	27.7	5.8	98	15.5	3.3	93	23.3	5.5	68	11.2	2.9
1978	91	18.8	4.2	85	13.7	3.2	81	18.6	4.6	78	12.6	3.0
1979	112	26.6	5.7	74	12.0	2.9	76	17.1	4.4	62	9.7	2.6
1980	132	29.5	5.7	106	15.6	3.2	92	22.3	5.2	69	10.5	2.7
1981	133	28.5	5.6	88	12.6	2.8	111	24.6	5.2	67	9.6	2.4
1982	130	29.9	5.8	101	13.5	2.8	106	27.8	6.0	61	8.4	2.2
1983	111	22.7	4.6	110	14.0	2.7	96	19.8	4.4	77	9.8	2.3
1984	114	24.6	5.1	88	10.5	2.3	73	16.4	4.2	61	7.3	1.9
1985	115	22.5	4.6	109	13.6	2.6	72	15.0	3.9	66	7.8	2.0
1986	152	30.8	5.5	105	11.9	2.4	96	19.5	4.4	69	8.5	2.1
1987	139	24.9	4.5	104	12.6	2.5	80	15.8	3.9	74	9.1	2.1
1988	131	24.1	4.5	103	11.8	2.3	98	18.4	4.0	71	8.2	2.0
1989	120	21.4	4.3	117	13.7	2.6	82	15.3	3.6	77	8.7	2.0
1990	129	23.3	4.4	119	12.8	2.4	82	15.4	3.7	81	9.3	2.1
1991	118	20.3	4.0	108	11.5	2.2	86	15.1	3.5	65	7.1	1.8
1992	127	21.0	4.0	125	13.2	2.4	78	13.9	3.4	90	9.9	2.1
1993	122	20.9	4.0	126	12.7	2.2	69	12.1	3.1	80	8.1	1.8
1994	142	24.0	4.3	142	14.5	2.4	102	17.6	3.7	87	9.0	1.9
1995	123	20.0	3.8	118	11.8	2.1	71	11.3	2.8	89	8.9	1.9
1996	142	22.9	4.0	118	11.2	2.1	96	16.5	3.6	84	8.2	1.8
1997	139	24.1	4.3	119	11.4	2.1	86	13.9	3.2	71	6.6	1.6
1998	128	20.0	3.7	126	11.6	2.0	76	12.3	3.0	73	6.8	1.6
1999	121	18.8	3.6	116	10.3	1.9	72	11.8	3.0	73	6.5	1.5
2000	130	20.5	3.8	130	11.3	2.0	85	13.2	3.0	63	5.4	1.3
2001	117	17.4	3.4	128	11.2	1.9	69	10.6	2.7	62	5.5	1.4
2002	117	17.0	3.3	129	10.6	1.8	70	11.0	2.8	68	5.6	1.3
2003	125	17.4	3.3	131	10.8	1.9	65	8.6	2.2	71	5.9	1.4
2004	120	17.3	3.3	122	9.8	1.7	64	9.3	2.4	71	5.7	1.3
2005	165	23.4	3.8	141	11.3	1.9	83	12.5	2.9	65	5.2	1.3
2006	143	19.2	3.3	126	9.8	1.7	79	11.6	2.7	64	5.1	1.3
2007	119	15.8	3.0	118	9.2	1.7	57	7.1	2.0	70	5.5	1.3
2008	113	15.6	3.0	115	8.8	1.6	60	8.0	2.2	63	4.9	1.2
2009	149	18.4	3.1	116	8.9	1.6	62	8.6	2.2	54	4.1	1.1
2010	130	15.9	2.9	129	9.5	1.7	60	8.1	2.2	53	3.9	1.1
2011	126	15.0	2.7	136	9.6	1.6	73	9.3	2.2	55	3.9	1.0
2012	152	17.1	2.9	126	8.8	1.6	56	6.3	1.8	59	4.3	1.1
2013	147	16.6	2.8	126	8.7	1.5	60	7.0	1.9	60	4.1	1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Colorectal

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	209	52.6	8.1	273	42.9	5.5	133	37.1	7.4	169	28.3	4.6
1977	240	64.7	9.4	353	56.3	6.4	123	34.7	6.9	156	24.8	4.2
1978	247	65.2	9.4	361	57.6	6.4	148	39.8	7.3	216	35.8	5.2
1979	266	67.5	9.3	374	55.8	6.0	161	42.8	7.5	194	29.2	4.4
1980	294	67.3	8.7	375	53.0	5.7	146	36.7	6.8	210	30.9	4.4
1981	279	64.5	8.6	381	51.7	5.5	140	35.8	6.7	230	32.9	4.5
1982	311	67.2	8.3	368	48.3	5.2	160	34.3	5.9	195	26.4	3.9
1983	297	64.1	8.2	374	45.6	4.8	157	31.6	5.5	187	23.9	3.6
1984	316	62.6	7.7	383	47.7	5.0	168	36.9	6.3	212	26.6	3.7
1985	346	71.0	8.4	425	49.8	4.9	152	33.4	6.0	224	27.5	3.7
1986	357	69.7	8.0	443	50.4	4.8	159	32.4	5.6	196	22.9	3.3
1987	308	58.5	7.2	406	45.0	4.5	151	29.8	5.3	203	23.3	3.3
1988	311	59.1	7.3	423	46.4	4.6	157	31.4	5.5	194	21.8	3.2
1989	361	67.9	7.7	397	42.5	4.3	163	30.9	5.3	179	19.1	2.9
1990	323	61.1	7.3	444	47.0	4.5	148	30.3	5.5	185	20.4	3.0
1991	348	64.0	7.4	489	51.8	4.7	152	31.5	5.6	217	23.4	3.2
1992	311	56.2	6.9	480	49.0	4.5	151	29.8	5.2	210	21.4	3.0
1993	367	64.1	7.2	452	45.7	4.3	141	25.7	4.7	204	21.0	2.9
1994	363	62.9	7.1	500	49.5	4.4	176	30.9	4.9	205	20.9	2.9
1995	379	62.4	6.8	471	45.1	4.1	170	28.8	4.7	239	23.8	3.0
1996	390	63.7	6.9	528	50.0	4.3	203	36.0	5.3	218	21.1	2.8
1997	436	71.5	7.3	536	50.0	4.3	193	34.1	5.2	229	21.6	2.8
1998	388	60.7	6.5	560	51.0	4.3	161	26.4	4.4	223	20.7	2.7
1999	407	61.0	6.3	618	53.9	4.3	182	29.3	4.5	262	23.2	2.8
2000	396	58.5	6.1	529	44.5	3.8	165	25.1	4.1	223	19.4	2.6
2001	411	59.4	6.1	580	48.2	3.9	175	27.7	4.4	242	20.4	2.6
2002	412	55.6	5.7	558	45.7	3.8	179	27.5	4.3	209	17.4	2.4
2003	464	63.7	6.2	592	47.4	3.8	191	28.6	4.3	218	17.8	2.4
2004	425	56.6	5.7	619	48.6	3.9	179	25.6	4.0	202	16.3	2.3
2005	451	57.8	5.6	568	44.3	3.7	153	21.7	3.6	205	16.3	2.2
2006	441	57.3	5.7	635	48.8	3.8	159	22.7	3.7	222	17.2	2.3
2007	492	59.9	5.6	570	43.4	3.6	162	21.2	3.5	208	16.1	2.2
2008	467	56.6	5.4	578	43.4	3.6	170	22.1	3.5	197	15.1	2.1
2009	488	56.5	5.3	551	40.4	3.4	170	22.3	3.5	192	14.5	2.1
2010	461	54.9	5.3	556	40.5	3.4	183	23.0	3.5	172	12.3	1.9
2011	481	53.2	5.0	550	38.4	3.2	143	17.5	3.0	214	15.3	2.1
2012	467	51.3	4.9	514	35.6	3.1	201	22.9	3.3	212	14.7	2.0
2013	461	48.9	4.7	534	36.0	3.1	152	16.8	2.8	192	13.1	1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Colon excluding rectum

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	151	39.4	7.1	211	32.6	4.7	105	29.9	6.6	139	23.1	4.2
1977	156	43.9	8.0	273	44.4	5.7	85	25.2	6.0	132	21.2	3.9
1978	181	51.1	8.6	280	45.1	5.7	120	33.0	6.8	184	30.1	4.7
1979	209	53.0	8.2	292	43.9	5.4	130	34.9	6.8	166	24.6	4.0
1980	213	49.8	7.6	284	40.7	5.0	117	29.9	6.2	178	26.5	4.1
1981	201	48.3	7.6	277	38.7	4.8	119	30.4	6.2	204	29.4	4.3
1982	224	49.9	7.3	277	36.6	4.5	130	28.4	5.5	165	22.3	3.6
1983	224	49.0	7.2	278	33.6	4.1	126	25.8	5.0	158	20.1	3.3
1984	223	44.2	6.4	289	36.6	4.4	147	32.0	5.8	184	23.0	3.5
1985	245	52.2	7.3	327	38.7	4.4	123	27.1	5.4	193	23.5	3.4
1986	261	52.2	7.0	334	38.0	4.2	141	28.9	5.3	171	19.9	3.1
1987	231	43.3	6.2	315	35.0	4.0	128	25.2	4.9	180	20.6	3.1
1988	244	46.9	6.5	325	36.1	4.0	137	26.9	5.1	174	19.5	3.0
1989	278	52.9	6.8	317	34.3	3.9	143	27.4	5.0	162	17.3	2.7
1990	242	46.6	6.4	346	36.7	4.0	128	26.2	5.0	165	18.2	2.8
1991	269	49.9	6.6	369	39.4	4.1	138	28.4	5.3	197	21.2	3.0
1992	228	42.4	6.1	379	38.6	4.0	132	26.9	5.1	186	19.0	2.8
1993	266	47.9	6.3	349	35.6	3.8	124	22.4	4.4	182	18.7	2.8
1994	275	48.5	6.3	411	40.9	4.0	155	27.5	4.7	179	18.3	2.7
1995	287	48.5	6.1	367	35.4	3.7	158	26.8	4.5	217	21.6	2.9
1996	303	50.3	6.2	398	37.7	3.7	178	32.2	5.1	192	18.6	2.7
1997	325	53.8	6.4	426	39.6	3.8	164	29.5	4.9	200	18.9	2.6
1998	291	46.7	5.8	438	40.0	3.8	142	23.4	4.2	199	18.4	2.6
1999	293	45.1	5.5	487	42.6	3.8	156	25.6	4.3	238	21.1	2.7
2000	286	43.5	5.4	413	35.0	3.4	146	22.5	3.9	203	17.6	2.4
2001	304	45.1	5.4	432	36.2	3.4	156	24.7	4.2	221	18.7	2.5
2002	295	39.8	4.8	448	36.9	3.4	159	24.5	4.1	191	15.9	2.3
2003	334	47.3	5.4	459	36.8	3.4	160	24.5	4.0	193	15.8	2.2
2004	310	41.9	4.9	470	37.1	3.4	150	21.5	3.6	178	14.4	2.1
2005	328	42.8	4.9	447	34.9	3.3	129	18.4	3.4	183	14.6	2.1
2006	323	42.7	4.9	472	36.6	3.3	139	20.1	3.5	193	15.0	2.1
2007	351	44.0	4.9	432	32.9	3.1	138	18.1	3.2	186	14.4	2.1
2008	347	43.2	4.8	443	33.5	3.2	142	18.6	3.2	173	13.2	2.0
2009	338	40.4	4.5	418	30.8	3.0	149	19.9	3.4	168	12.8	2.0
2010	324	39.4	4.5	426	31.3	3.0	157	20.1	3.3	148	10.6	1.7
2011	352	39.7	4.4	413	28.9	2.8	116	14.5	2.8	184	13.1	1.9
2012	322	36.6	4.2	395	27.6	2.8	156	18.0	3.0	183	12.7	1.9
2013	327	35.3	4.0	386	26.1	2.6	123	13.8	2.6	174	11.9	1.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Rectum & rectosigmoid

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	58	13.2	3.9	62	10.3	2.8	28	7.2	3.2	30	5.2	2.0
1977	84	20.8	5.0	80	11.9	2.8	38	9.4	3.3	24	3.6	1.6
1978	66	14.1	3.9	81	12.5	2.9	28	6.8	2.8	32	5.7	2.2
1979	57	14.5	4.3	82	11.9	2.8	31	7.9	3.1	28	4.6	1.8
1980	81	17.5	4.3	91	12.3	2.6	29	6.8	2.7	32	4.4	1.6
1981	78	16.2	4.1	104	13.0	2.6	21	5.4	2.6	26	3.6	1.4
1982	87	17.3	3.9	91	11.7	2.5	30	5.9	2.3	30	4.1	1.5
1983	73	15.1	3.9	96	12.0	2.5	31	5.8	2.3	29	3.8	1.4
1984	93	18.3	4.2	94	11.1	2.3	21	5.0	2.3	28	3.6	1.4
1985	101	18.9	4.1	98	11.1	2.3	29	6.4	2.6	31	4.0	1.5
1986	96	17.5	3.8	109	12.4	2.4	18	3.5	1.8	25	3.0	1.2
1987	77	15.2	3.8	91	9.9	2.1	23	4.7	2.1	23	2.7	1.2
1988	67	12.1	3.2	98	10.3	2.1	20	4.4	2.1	20	2.4	1.1
1989	83	15.0	3.6	80	8.2	1.9	20	3.5	1.7	17	1.8	0.9
1990	81	14.5	3.4	98	10.3	2.1	20	4.1	2.1	20	2.2	1.0
1991	79	14.1	3.4	120	12.3	2.3	14	3.1	1.8	20	2.1	1.0
1992	83	13.8	3.3	101	10.4	2.1	19	2.9	1.4	24	2.5	1.0
1993	101	16.2	3.5	103	10.1	2.0	17	3.2	1.7	22	2.3	1.0
1994	88	14.4	3.3	89	8.6	1.8	21	3.4	1.5	26	2.6	1.0
1995	92	13.8	3.1	104	9.7	1.9	12	1.9	1.2	22	2.2	0.9
1996	87	13.4	3.0	130	12.3	2.1	25	3.8	1.6	26	2.5	1.0
1997	111	17.6	3.5	110	10.4	2.0	29	4.6	1.8	29	2.7	1.0
1998	97	14.0	3.0	122	11.1	2.0	19	3.1	1.5	24	2.3	0.9
1999	114	16.0	3.1	131	11.4	2.0	26	3.7	1.5	24	2.1	0.9
2000	110	15.1	3.0	116	9.6	1.8	19	2.6	1.2	20	1.8	0.8
2001	107	14.3	2.9	148	12.1	2.0	19	3.0	1.4	21	1.7	0.7
2002	117	15.9	3.1	110	8.8	1.7	20	3.0	1.4	18	1.5	0.7
2003	130	16.4	3.0	133	10.6	1.8	31	4.1	1.5	25	2.0	0.8
2004	115	14.7	2.8	149	11.5	1.9	29	4.1	1.6	24	1.9	0.8
2005	123	15.0	2.8	121	9.3	1.7	24	3.3	1.4	22	1.7	0.7
2006	118	14.6	2.8	163	12.2	1.9	20	2.6	1.2	29	2.2	0.8
2007	141	15.9	2.7	138	10.5	1.8	24	3.1	1.3	22	1.7	0.7
2008	120	13.4	2.5	135	9.8	1.7	28	3.5	1.4	24	1.9	0.7
2009	150	16.0	2.7	133	9.7	1.7	21	2.4	1.1	24	1.7	0.7
2010	137	15.5	2.7	130	9.2	1.6	26	2.9	1.2	24	1.7	0.7
2011	129	13.5	2.4	137	9.5	1.6	27	2.9	1.2	30	2.1	0.8
2012	145	14.7	2.5	119	8.1	1.5	45	4.9	1.5	29	2.0	0.7
2013	134	13.6	2.4	148	9.9	1.6	29	3.0	1.2	18	1.2	0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Liver / intrahepatic bile duct

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	41	9.9	3.4	22	3.4	1.6	43	11.1	3.7	21	3.5	1.6
1977	42	9.0	3.2	19	2.8	1.4	54	12.4	3.8	26	3.5	1.4
1978	46	10.0	3.1	35	5.0	1.8	40	8.0	2.7	38	5.5	1.9
1979	36	7.8	2.9	26	3.4	1.4	44	10.6	3.5	39	5.5	1.8
1980	45	8.9	2.9	22	3.0	1.3	43	8.8	3.0	26	3.5	1.4
1981	43	8.3	2.7	23	3.0	1.3	41	7.8	2.6	27	3.8	1.5
1982	53	9.6	2.7	24	3.1	1.3	48	8.6	2.6	27	3.7	1.5
1983	48	9.4	3.0	29	3.5	1.3	58	11.1	3.2	39	4.9	1.6
1984	58	12.0	3.5	29	3.2	1.2	54	11.7	3.5	40	4.3	1.4
1985	57	10.4	3.0	27	2.8	1.1	55	10.3	3.0	38	4.4	1.4
1986	53	9.9	2.9	26	3.0	1.2	57	10.5	2.9	33	3.8	1.3
1987	48	8.1	2.6	36	3.8	1.3	45	9.4	3.0	47	5.3	1.6
1988	49	8.2	2.6	27	2.9	1.1	46	7.8	2.5	33	3.6	1.3
1989	45	8.3	2.8	37	4.0	1.3	49	8.0	2.4	41	4.4	1.4
1990	60	9.4	2.6	40	4.2	1.3	46	8.1	2.6	47	4.9	1.4
1991	66	10.9	2.9	37	3.6	1.2	57	10.1	2.9	45	4.6	1.4
1992	68	11.6	3.0	41	4.1	1.3	54	9.3	2.7	41	4.1	1.3
1993	71	11.2	2.8	32	3.2	1.1	68	10.7	2.7	42	4.4	1.3
1994	65	9.4	2.4	50	4.8	1.4	57	9.3	2.6	35	3.6	1.2
1995	70	10.1	2.6	43	4.0	1.2	56	8.2	2.4	44	4.1	1.2
1996	95	14.2	3.1	35	3.3	1.1	65	10.2	2.7	42	4.0	1.2
1997	77	10.6	2.6	46	4.3	1.2	58	8.3	2.3	38	3.5	1.1
1998	94	12.6	2.7	42	3.8	1.2	75	10.3	2.5	37	3.4	1.1
1999	88	11.5	2.5	63	5.4	1.3	51	6.7	1.9	36	3.2	1.0
2000	101	12.0	2.4	53	4.5	1.2	77	10.4	2.5	41	3.5	1.1
2001	111	13.6	2.7	35	2.9	1.0	65	8.3	2.1	41	3.4	1.1
2002	96	11.8	2.5	46	3.8	1.1	64	8.3	2.2	39	3.3	1.0
2003	120	13.7	2.6	48	3.8	1.1	72	9.2	2.3	40	3.2	1.0
2004	143	16.4	2.8	62	4.9	1.2	89	10.4	2.3	58	4.7	1.2
2005	153	16.7	2.8	49	3.7	1.0	81	10.0	2.3	42	3.3	1.0
2006	138	15.8	2.8	51	3.9	1.1	105	11.9	2.4	31	2.4	0.9
2007	167	18.5	3.0	60	4.5	1.2	96	11.0	2.3	48	3.6	1.0
2008	170	17.7	2.8	59	4.4	1.1	98	10.8	2.2	41	3.0	0.9
2009	178	19.1	3.0	82	5.9	1.3	99	10.8	2.2	42	3.1	0.9
2010	187	18.8	2.8	94	6.6	1.3	113	12.2	2.4	58	4.0	1.0
2011	188	18.5	2.8	92	6.5	1.3	99	9.9	2.0	65	4.7	1.2
2012	223	22.1	3.0	92	6.3	1.3	114	11.5	2.2	52	3.7	1.0
2013	205	18.9	2.7	90	5.9	1.2	112	11.3	2.2	59	3.8	1.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Pancreas

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	64	14.7	4.0	85	14.3	3.3	70	15.9	4.2	81	12.6	3.0
1977	72	17.1	4.7	73	11.3	2.8	65	15.5	4.5	60	9.3	2.6
1978	61	12.8	3.7	87	13.6	3.1	55	11.8	3.5	79	11.6	2.7
1979	72	14.8	3.7	87	13.3	3.0	56	11.1	3.1	90	14.2	3.1
1980	85	18.3	4.5	73	10.5	2.5	69	15.9	4.4	61	8.8	2.3
1981	100	21.1	4.6	81	11.1	2.6	82	17.1	4.3	65	8.9	2.3
1982	69	15.8	4.2	103	13.5	2.7	67	15.6	4.2	93	12.4	2.6
1983	89	16.7	3.9	107	13.3	2.6	90	18.3	4.2	88	10.9	2.4
1984	83	16.1	3.9	104	13.0	2.6	81	17.0	4.1	84	10.5	2.3
1985	93	17.1	3.8	83	10.2	2.3	75	14.4	3.6	74	9.3	2.2
1986	83	15.2	3.6	107	12.6	2.5	70	13.0	3.3	88	10.7	2.3
1987	104	17.4	3.7	110	12.7	2.5	80	14.9	3.7	95	10.6	2.2
1988	96	17.7	3.9	113	12.2	2.3	72	12.2	3.1	77	8.7	2.0
1989	79	14.0	3.4	116	12.5	2.3	75	13.1	3.2	105	11.2	2.2
1990	87	16.2	3.8	119	12.9	2.4	70	12.8	3.3	101	10.8	2.2
1991	81	14.4	3.5	98	10.5	2.1	82	13.6	3.2	83	8.6	1.9
1992	71	12.7	3.3	120	12.2	2.2	53	10.6	3.1	90	9.4	2.0
1993	85	14.6	3.4	109	11.0	2.1	58	11.0	3.2	90	9.0	1.9
1994	88	14.6	3.3	129	12.8	2.2	71	11.9	3.0	120	11.9	2.2
1995	83	14.0	3.2	105	10.4	2.0	75	12.8	3.1	98	9.5	1.9
1996	79	12.4	2.9	139	13.2	2.2	62	9.7	2.6	105	10.1	2.0
1997	96	14.0	2.9	128	12.2	2.1	72	11.0	2.7	105	10.1	1.9
1998	82	12.1	2.8	128	11.9	2.1	69	10.3	2.6	102	9.6	1.9
1999	96	15.2	3.3	153	13.6	2.2	71	11.0	2.7	125	11.2	2.0
2000	80	12.2	2.8	156	13.5	2.1	78	12.0	2.8	106	9.1	1.7
2001	91	12.8	2.8	153	12.9	2.1	67	9.7	2.4	133	11.3	1.9
2002	94	12.5	2.7	151	12.8	2.0	92	12.4	2.7	115	9.6	1.8
2003	108	15.0	3.0	119	9.7	1.8	74	10.3	2.5	97	8.1	1.6
2004	102	13.2	2.7	139	11.2	1.9	91	12.4	2.7	119	9.4	1.7
2005	89	11.9	2.6	131	10.2	1.8	76	9.7	2.3	109	8.7	1.6
2006	117	14.0	2.7	174	13.6	2.0	83	10.8	2.4	111	8.7	1.6
2007	131	16.4	3.0	152	11.8	1.9	102	13.1	2.7	122	9.6	1.7
2008	92	11.1	2.4	160	12.1	1.9	88	11.1	2.5	130	9.9	1.7
2009	119	14.8	2.8	144	10.6	1.7	91	11.6	2.5	118	8.7	1.6
2010	136	15.8	2.8	171	12.5	1.9	88	10.6	2.3	120	9.1	1.7
2011	147	16.3	2.8	202	14.5	2.0	117	13.9	2.7	145	10.4	1.7
2012	129	14.2	2.6	208	14.5	2.0	86	9.4	2.1	171	12.0	1.8
2013	164	17.4	2.8	206	14.0	1.9	115	12.7	2.5	131	8.9	1.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Larynx

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	63	11.7	3.0	16	2.1	1.1	39	8.3	3.0	11	1.6	1.0
1977	72	14.8	3.9	18	2.3	1.1	32	7.4	3.3	11	1.3	0.8
1978	81	15.3	3.7	19	2.2	1.0	55	9.7	2.7	9	1.0	0.7
1979	83	15.5	3.6	24	2.9	1.2	47	8.7	2.6	11	1.2	0.7
1980	74	12.8	3.1	26	3.0	1.2	34	6.6	2.3	18	2.0	0.9
1981	75	12.8	3.1	18	2.1	1.0	35	6.1	2.1	20	2.1	0.9
1982	73	12.0	2.9	12	1.4	0.8	31	5.6	2.1	8	0.9	0.6
1983	94	15.6	3.4	20	2.3	1.0	38	6.3	2.2	8	0.9	0.7
1984	81	13.0	2.9	30	3.2	1.2	49	7.6	2.2	13	1.4	0.8
1985	117	19.1	3.6	29	2.8	1.0	42	6.9	2.2	10	1.1	0.7
1986	108	18.2	3.7	32	3.3	1.2	52	8.8	2.6	13	1.4	0.8
1987	97	15.3	3.2	25	2.6	1.0	50	8.3	2.5	16	1.7	0.8
1988	98	14.4	3.0	30	2.8	1.0	58	9.4	2.6	13	1.2	0.7
1989	89	13.6	3.0	49	4.7	1.3	43	7.1	2.4	17	1.6	0.8
1990	110	16.6	3.4	37	3.6	1.2	35	5.7	2.1	11	1.1	0.6
1991	93	14.2	3.1	30	2.7	1.0	40	6.0	2.0	15	1.4	0.7
1992	122	17.2	3.2	27	2.4	0.9	51	7.9	2.3	14	1.3	0.7
1993	88	13.6	3.0	28	2.6	1.0	48	7.7	2.4	19	1.8	0.8
1994	78	11.3	2.6	30	2.7	1.0	42	6.0	1.9	18	1.7	0.8
1995	83	11.5	2.6	35	3.2	1.1	55	8.5	2.4	17	1.5	0.7
1996	92	12.6	2.7	21	1.8	0.8	54	8.2	2.3	9	0.8	0.5
1997	74	9.9	2.3	26	2.3	0.9	45	6.2	1.9	11	1.0	0.6
1998	75	10.5	2.5	35	2.9	1.0	43	5.8	1.8	9	0.8	0.5
1999	74	10.1	2.4	25	2.1	0.8	31	4.2	1.6	14	1.2	0.6
2000	67	9.0	2.3	28	2.3	0.9	42	5.7	1.8	12	1.1	0.6
2001	64	7.7	1.9	22	1.8	0.8	27	3.7	1.4	8	0.6	0.5
2002	61	7.2	1.9	27	2.2	0.8	34	4.2	1.5	12	1.0	0.6
2003	77	9.5	2.3	28	2.2	0.8	31	4.0	1.5	6	0.5	0.4
2004	75	8.4	2.0	16	1.2	0.6	38	4.9	1.6	9	0.7	0.5
2005	68	8.4	2.1	22	1.7	0.7	31	3.7	1.4	6	0.5	0.4
2006	57	6.2	1.6	20	1.4	0.6	32	4.0	1.5	7	0.5	0.4
2007	76	8.5	2.0	18	1.3	0.6	35	4.2	1.4	8	0.6	0.4
2008	69	8.2	2.0	37	2.7	0.9	26	3.3	1.3	15	1.1	0.5
2009	68	7.8	1.9	19	1.3	0.6	29	3.4	1.3	10	0.7	0.4
2010	63	6.7	1.7	23	1.6	0.7	25	3.1	1.3	6	0.4	0.3
2011	62	6.3	1.7	26	1.7	0.6	21	2.5	1.1	11	0.8	0.5
2012	52	5.4	1.5	19	1.3	0.6	24	2.8	1.2	7	0.4	0.3
2013	65	6.7	1.7	17	1.1	0.5	22	2.3	1.0	10	0.6	0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Lung and bronchus

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	495	102.9	10.2	172	24.9	4.0	395	82.5	9.2	129	18.3	3.4
1977	488	102.6	10.3	186	24.0	3.6	408	85.5	9.4	146	18.9	3.2
1978	527	105.2	9.9	191	26.7	4.0	444	88.5	9.0	167	22.7	3.7
1979	568	115.1	10.5	220	29.8	4.2	473	98.1	10.0	165	22.5	3.6
1980	572	110.7	10.0	218	27.1	3.7	474	91.9	9.1	168	21.6	3.4
1981	574	107.1	9.6	255	29.8	3.8	495	93.3	9.0	204	24.6	3.5
1982	584	107.5	9.4	268	31.3	3.9	470	86.3	8.5	210	25.5	3.6
1983	590	106.5	9.4	266	30.7	3.8	528	96.9	9.0	227	25.7	3.4
1984	616	108.7	9.3	304	33.8	3.9	492	89.5	8.7	213	23.6	3.3
1985	624	111.1	9.5	287	31.1	3.7	516	94.0	8.9	230	25.3	3.4
1986	595	103.0	9.0	339	35.7	3.9	468	82.2	8.2	260	27.6	3.4
1987	664	112.1	9.2	327	33.7	3.7	498	87.8	8.4	274	28.7	3.5
1988	594	101.4	8.8	337	34.0	3.7	494	86.3	8.3	266	27.4	3.4
1989	638	107.2	9.0	355	34.9	3.7	503	88.6	8.5	245	24.4	3.1
1990	644	105.4	8.8	368	35.8	3.7	511	83.0	7.7	269	26.7	3.3
1991	626	102.4	8.7	357	35.0	3.7	486	81.1	7.8	249	24.5	3.1
1992	636	102.5	8.7	379	36.4	3.7	438	74.1	7.5	299	29.3	3.4
1993	584	95.1	8.4	383	36.2	3.7	444	74.2	7.5	279	26.1	3.1
1994	633	98.1	8.2	393	36.4	3.6	509	81.8	7.7	301	28.7	3.3
1995	586	90.0	7.8	383	35.4	3.6	492	77.0	7.3	319	29.5	3.3
1996	632	94.1	7.8	451	41.1	3.8	509	77.6	7.2	319	29.5	3.3
1997	579	87.1	7.5	448	40.2	3.8	460	71.7	7.0	323	29.4	3.2
1998	573	81.6	7.0	460	40.4	3.7	452	67.1	6.6	331	29.7	3.2
1999	548	77.8	6.9	409	35.2	3.4	416	60.9	6.2	330	28.9	3.1
2000	553	76.9	6.7	440	36.7	3.5	423	60.0	6.0	312	26.2	2.9
2001	540	76.0	6.8	438	36.3	3.4	414	59.2	6.0	328	27.2	3.0
2002	553	75.5	6.7	469	37.7	3.4	412	57.1	5.8	340	27.5	2.9
2003	497	65.9	6.1	481	38.4	3.4	387	52.3	5.5	338	27.0	2.9
2004	552	70.0	6.1	495	39.1	3.5	375	48.7	5.2	321	25.3	2.8
2005	560	71.9	6.3	483	37.5	3.4	383	49.2	5.2	345	27.3	2.9
2006	531	65.4	5.8	493	37.7	3.4	360	46.7	5.1	352	27.4	2.9
2007	528	64.9	5.8	507	38.5	3.4	382	47.9	5.1	330	25.4	2.8
2008	541	66.6	5.9	515	38.8	3.4	344	43.0	4.8	335	25.5	2.8
2009	554	65.9	5.8	520	38.3	3.3	365	44.4	4.8	306	22.7	2.6
2010	540	61.8	5.5	542	39.1	3.3	387	46.5	4.9	355	25.6	2.7
2011	567	65.2	5.6	589	41.2	3.4	371	42.5	4.5	357	25.2	2.6
2012	533	59.7	5.3	574	39.5	3.3	384	44.3	4.7	346	24.1	2.6
2013	507	54.9	5.0	512	34.9	3.1	365	40.6	4.4	340	23.0	2.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Melanoma of the skin

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	1	0.2	0.3	7	1.1	1.0	3	1.3	1.7	2	0.3	0.4
1977	3	0.7	0.9	5	0.8	0.7	2	0.7	1.0	2	0.2	0.3
1978	3	0.6	0.8	6	0.9	0.7	2	0.3	0.4	3	0.5	0.6
1979	6	1.0	0.8	11	1.4	0.9	1	0.3	0.7	1	0.1	0.3
1980	3	0.5	0.5	7	1.0	0.8	8	1.2	0.8	6	0.9	0.8
1981	5	0.8	0.7	13	1.8	1.0	6	0.9	0.8	8	1.1	0.9
1982	2	0.3	0.5	8	1.0	0.7	4	1.0	1.2	3	0.3	0.3
1983	2	0.3	0.4	8	0.9	0.7	13	2.1	1.2	5	0.6	0.6
1984	5	0.8	0.8	16	1.9	1.0	9	1.4	1.0	3	0.3	0.3
1985	9	1.4	1.1	8	1.1	0.8	9	1.9	1.3	3	0.3	0.4
1986	8	1.8	1.5	6	0.6	0.5	10	1.7	1.2	4	0.5	0.5
1987	5	0.7	0.6	7	0.7	0.5	3	0.5	0.6	5	0.5	0.5
1988	5	1.1	1.1	10	1.0	0.6	2	0.6	0.9	5	0.6	0.5
1989	4	0.8	0.9	9	1.1	0.7	4	0.5	0.5	8	0.9	0.7
1990	9	1.4	1.1	8	0.8	0.5	3	0.6	0.9	3	0.3	0.4
1991	12	2.2	1.5	8	0.6	0.4	6	1.0	1.0	3	0.3	0.4
1992	4	0.5	0.5	7	0.6	0.5	2	0.2	0.3	4	0.4	0.4
1993	11	1.6	1.0	6	0.7	0.5	2	0.2	0.3	3	0.3	0.3
1994	3	0.7	0.8	10	0.9	0.6	1	0.2	0.5	2	0.2	0.3
1995	3	0.4	0.4	15	1.4	0.7	0	0.0	0.0	2	0.2	0.3
1996	5	0.8	0.7	13	1.3	0.7	3	0.5	0.7	6	0.6	0.5
1997	14	2.0	1.1	17	1.5	0.7	3	0.3	0.4	4	0.4	0.4
1998	10	1.6	1.1	10	0.9	0.5	2	0.2	0.3	6	0.5	0.4
1999	3	0.4	0.5	5	0.4	0.4	4	0.7	0.8	3	0.3	0.3
2000	6	0.8	0.7	6	0.5	0.4	5	1.1	1.0	5	0.4	0.4
2001	8	0.9	0.7	15	1.2	0.6	1	0.1	0.2	1	0.1	0.2
2002	5	0.5	0.4	11	0.9	0.5	2	0.2	0.3	3	0.3	0.3
2003	3	0.5	0.6	9	0.7	0.5	1	0.1	0.2	4	0.3	0.3
2004	6	1.0	0.8	9	0.7	0.4	1	0.1	0.2	2	0.2	0.2
2005	7	0.8	0.6	13	1.0	0.6	3	0.3	0.3	1	0.1	0.2
2006	13	1.9	1.1	15	1.1	0.6	4	0.4	0.4	3	0.2	0.3
2007	11	1.4	0.9	21	1.6	0.7	6	0.9	0.8	9	0.7	0.5
2008	11	1.4	0.9	17	1.3	0.6	3	0.3	0.3	5	0.4	0.3
2009	9	0.9	0.6	14	1.0	0.5	3	0.3	0.3	1	0.1	0.1
2010	9	1.0	0.7	12	0.8	0.5	2	0.3	0.5	4	0.3	0.3
2011	14	1.6	0.9	16	1.2	0.6	5	0.6	0.6	6	0.5	0.4
2012	10	1.1	0.7	13	0.9	0.5	3	0.4	0.5	5	0.3	0.3
2013	7	0.8	0.6	17	1.2	0.6	5	0.4	0.4	4	0.3	0.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Female breast

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	538	74.3	6.7	248	36.3	4.9
1977	544	72.5	6.4	230	31.5	4.3
1978	608	80.2	6.7	258	35.1	4.5
1979	629	80.6	6.6	229	30.1	4.1
1980	664	81.9	6.5	278	35.7	4.4
1981	650	79.5	6.4	273	34.5	4.3
1982	703	84.0	6.5	277	33.1	4.1
1983	705	80.7	6.2	299	35.1	4.1
1984	823	91.4	6.4	316	36.8	4.2
1985	811	88.3	6.3	251	27.7	3.5
1986	802	84.0	6.0	303	32.1	3.7
1987	886	89.7	6.0	312	32.1	3.7
1988	947	94.6	6.2	321	32.2	3.6
1989	906	89.0	5.9	319	32.1	3.6
1990	925	89.7	5.9	340	33.9	3.7
1991	1057	101.2	6.2	321	31.3	3.5
1992	999	94.6	6.0	330	31.7	3.5
1993	1014	94.1	5.9	314	30.4	3.4
1994	1079	98.4	5.9	380	34.7	3.5
1995	1109	99.0	5.9	403	36.7	3.6
1996	1201	105.6	6.0	381	34.1	3.5
1997	1210	104.3	5.9	375	33.1	3.4
1998	1352	115.0	6.2	369	32.0	3.3
1999	1204	99.8	5.7	375	31.8	3.2
2000	1215	98.2	5.6	336	27.7	3.0
2001	1196	94.9	5.4	395	32.1	3.2
2002	1254	98.0	5.5	344	27.6	2.9
2003	1312	101.2	5.5	349	27.4	2.9
2004	1316	100.1	5.4	348	27.0	2.9
2005	1335	100.6	5.4	392	29.7	3.0
2006	1404	104.7	5.5	352	26.7	2.8
2007	1444	107.0	5.6	334	24.7	2.7
2008	1491	109.9	5.6	334	24.8	2.7
2009	1509	109.6	5.6	332	24.3	2.6
2010	1481	105.8	5.5	320	22.8	2.5
2011	1678	117.3	5.7	355	25.3	2.7
2012	1625	111.7	5.5	382	26.5	2.7
2013	1721	117.1	5.6	314	21.1	2.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Cervix uteri

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	223	28.3	3.9	101	14.6	3.0
1977	245	31.5	4.1	102	14.1	2.9
1978	233	29.3	3.9	89	12.1	2.7
1979	249	30.6	4.0	81	10.9	2.5
1980	220	26.8	3.7	90	11.5	2.5
1981	199	22.2	3.2	81	9.3	2.1
1982	220	25.3	3.5	81	9.0	2.0
1983	202	22.2	3.2	76	8.5	2.0
1984	208	22.3	3.1	64	7.3	1.8
1985	191	19.9	2.9	71	7.6	1.8
1986	216	22.7	3.1	71	7.3	1.8
1987	199	19.4	2.8	58	5.8	1.5
1988	187	18.2	2.7	90	9.0	1.9
1989	180	16.5	2.5	50	5.0	1.4
1990	234	21.1	2.8	65	6.4	1.6
1991	216	19.5	2.6	63	5.8	1.5
1992	217	19.1	2.6	64	6.0	1.5
1993	201	17.7	2.5	76	6.8	1.6
1994	215	19.1	2.6	64	5.8	1.4
1995	217	18.6	2.5	77	6.9	1.6
1996	218	18.2	2.4	59	5.1	1.3
1997	213	17.6	2.4	67	5.9	1.4
1998	206	16.4	2.3	62	5.2	1.3
1999	197	15.7	2.2	75	6.1	1.4
2000	246	19.6	2.5	72	5.7	1.3
2001	192	15.0	2.1	60	4.8	1.2
2002	195	15.0	2.1	58	4.6	1.2
2003	175	13.3	2.0	66	5.1	1.2
2004	164	12.3	1.9	50	3.8	1.1
2005	182	13.7	2.0	54	4.1	1.1
2006	172	12.9	1.9	57	4.4	1.1
2007	187	14.0	2.0	72	5.4	1.2
2008	164	12.1	1.9	62	4.5	1.1
2009	156	11.2	1.8	64	4.5	1.1
2010	166	12.2	1.9	47	3.3	1.0
2011	162	11.6	1.8	67	4.8	1.2
2012	154	11.2	1.8	48	3.3	1.0
2013	150	10.6	1.7	60	4.1	1.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Corpus uterus and uterus, not otherwise specified

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	115	16.0	3.1	55	8.6	2.5
1977	107	15.2	3.0	51	7.2	2.1
1978	114	15.7	3.0	60	9.5	2.6
1979	106	13.8	2.8	52	7.3	2.1
1980	133	17.3	3.1	61	8.1	2.2
1981	113	14.1	2.7	51	6.2	1.8
1982	130	15.0	2.7	47	5.7	1.7
1983	133	15.3	2.7	61	7.3	1.9
1984	154	17.9	2.9	59	7.0	1.8
1985	164	17.8	2.8	68	7.8	1.9
1986	111	11.3	2.2	53	5.9	1.6
1987	161	16.8	2.7	44	5.0	1.5
1988	170	16.7	2.6	63	6.5	1.6
1989	147	14.2	2.4	52	4.9	1.4
1990	154	15.3	2.5	66	6.7	1.7
1991	208	20.1	2.8	60	6.0	1.6
1992	179	16.8	2.5	64	6.3	1.6
1993	205	18.9	2.6	69	6.8	1.6
1994	213	19.1	2.6	79	7.5	1.7
1995	203	18.2	2.5	59	5.5	1.4
1996	226	19.9	2.6	86	8.0	1.7
1997	252	22.1	2.7	97	8.7	1.7
1998	249	21.3	2.7	96	8.5	1.7
1999	269	22.7	2.7	90	7.8	1.6
2000	282	23.3	2.7	80	6.8	1.5
2001	304	24.2	2.7	98	8.1	1.6
2002	289	22.8	2.6	81	6.5	1.4
2003	310	23.9	2.7	104	8.2	1.6
2004	342	26.2	2.8	106	8.5	1.6
2005	350	26.5	2.8	111	8.5	1.6
2006	373	27.9	2.9	107	8.3	1.6
2007	337	25.0	2.7	95	7.3	1.5
2008	456	32.8	3.0	115	8.7	1.6
2009	427	30.2	2.9	120	8.8	1.6
2010	445	31.0	2.9	114	8.2	1.5
2011	463	31.7	2.9	145	10.2	1.7
2012	489	32.6	2.9	131	8.9	1.6
2013	502	32.8	2.9	125	8.2	1.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Ovary

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	76	10.8	2.6	47	6.7	2.0
1977	91	11.8	2.6	56	7.7	2.1
1978	78	10.5	2.5	59	7.8	2.1
1979	95	12.4	2.7	47	6.0	1.8
1980	101	12.8	2.6	76	10.3	2.5
1981	90	10.3	2.2	48	5.6	1.6
1982	79	9.4	2.1	51	6.3	1.8
1983	122	13.3	2.5	72	8.5	2.0
1984	119	13.2	2.5	66	7.4	1.9
1985	111	11.1	2.1	77	8.4	1.9
1986	115	12.1	2.3	63	6.6	1.7
1987	92	9.6	2.0	55	5.7	1.5
1988	112	11.5	2.2	55	5.7	1.6
1989	93	8.8	1.8	62	6.2	1.6
1990	127	12.3	2.2	79	8.0	1.8
1991	115	11.0	2.0	72	7.0	1.6
1992	152	14.3	2.3	89	8.5	1.8
1993	116	10.5	1.9	95	9.2	1.9
1994	160	14.7	2.3	90	8.7	1.8
1995	137	12.1	2.1	99	9.2	1.8
1996	110	9.5	1.8	58	5.2	1.4
1997	147	12.6	2.1	75	6.8	1.5
1998	107	9.2	1.8	79	7.0	1.6
1999	127	10.8	1.9	74	6.4	1.5
2000	121	10.0	1.8	76	6.5	1.5
2001	129	10.4	1.8	84	6.9	1.5
2002	140	11.0	1.8	79	6.5	1.4
2003	130	10.0	1.7	75	6.0	1.4
2004	137	10.5	1.8	78	6.2	1.4
2005	133	10.2	1.7	88	6.8	1.4
2006	115	8.5	1.6	80	6.2	1.4
2007	128	9.5	1.7	81	6.1	1.3
2008	133	9.7	1.7	82	5.9	1.3
2009	140	10.1	1.7	80	5.8	1.3
2010	134	9.7	1.7	88	6.3	1.3
2011	133	9.3	1.6	86	6.2	1.3
2012	151	10.4	1.7	97	6.7	1.4
2013	143	9.9	1.6	71	4.7	1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Prostate

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	396	118.2	13.6	144	49.8	9.5
1977	422	122.4	13.4	147	45.9	8.3
1978	463	133.5	14.0	167	54.9	9.5
1979	481	131.7	13.2	189	60.6	9.7
1980	585	157.2	14.3	218	67.4	10.0
1981	567	148.4	13.7	232	70.7	10.2
1982	544	139.4	13.0	202	59.6	9.0
1983	526	129.1	12.3	215	59.5	8.7
1984	602	142.2	12.6	219	60.5	8.7
1985	621	137.5	12.0	254	68.3	9.2
1986	638	140.9	12.0	254	66.4	8.9
1987	659	138.3	11.7	265	64.9	8.6
1988	703	152.4	12.4	210	53.0	7.7
1989	743	154.3	12.2	233	55.4	7.8
1990	832	167.9	12.5	288	69.7	8.8
1991	897	174.9	12.6	255	59.5	8.0
1992	1120	208.2	13.2	305	70.7	8.6
1993	1208	214.9	13.1	274	64.7	8.3
1994	1338	229.1	13.2	296	67.6	8.3
1995	1474	239.3	13.1	322	69.7	8.2
1996	1444	231.5	12.8	293	61.2	7.5
1997	1571	245.2	12.9	297	61.0	7.4
1998	1603	242.8	12.6	288	57.5	7.1
1999	1872	272.9	13.1	319	62.3	7.3
2000	2080	290.9	13.2	296	55.7	6.7
2001	1935	264.8	12.4	315	58.4	6.8
2002	1949	260.2	12.2	323	56.2	6.5
2003	1953	255.2	11.9	266	46.1	5.8
2004	2029	257.4	11.8	277	48.4	6.0
2005	2012	247.8	11.3	263	44.1	5.6
2006	2110	257.3	11.5	263	44.8	5.7
2007	2288	270.4	11.6	282	46.4	5.6
2008	2229	253.5	11.0	262	43.2	5.4
2009	2213	245.4	10.7	282	44.7	5.4
2010	2152	229.3	10.1	268	40.3	5.0
2011	2388	244.8	10.3	281	41.5	5.0
2012	2157	216.1	9.5	271	38.9	4.8
2013	2123	206.3	9.2	269	37.5	4.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Testis

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	5	0.5	0.4	3	1.0	1.6
1977	4	0.8	0.9	2	0.2	0.3
1978	2	0.5	0.7	3	0.6	0.8
1979	6	0.7	0.6	0	0.0	0.0
1980	8	1.2	0.8	1	0.1	0.3
1981	6	0.7	0.6	2	0.3	0.5
1982	4	0.5	0.5	0	0.0	0.0
1983	5	0.8	0.8	6	1.2	1.2
1984	3	0.2	0.3	1	0.2	0.4
1985	11	1.7	1.2	1	0.1	0.1
1986	11	1.1	0.7	2	0.2	0.3
1987	8	0.7	0.5	1	0.1	0.1
1988	8	0.8	0.6	2	0.5	0.9
1989	13	1.5	0.9	1	0.1	0.2
1990	9	0.9	0.6	3	0.3	0.3
1991	7	0.9	0.8	0	0.0	0.0
1992	14	1.1	0.6	1	0.1	0.2
1993	10	1.0	0.7	2	0.2	0.3
1994	13	1.2	0.7	1	0.1	0.2
1995	5	0.4	0.3	0	0.0	0.0
1996	17	1.5	0.7	0	0.0	0.0
1997	7	0.6	0.5	1	0.1	0.2
1998	15	1.4	0.7	1	0.1	0.2
1999	13	1.2	0.7	0	0.0	0.0
2000	24	2.3	1.0	3	0.4	0.4
2001	15	1.4	0.7	1	0.2	0.4
2002	13	1.2	0.6	2	0.2	0.3
2003	18	1.7	0.8	1	0.1	0.1
2004	11	1.0	0.6	1	0.1	0.2
2005	20	1.9	0.9	2	0.2	0.2
2006	24	2.3	0.9	1	0.1	0.2
2007	21	2.0	0.9	1	0.1	0.2
2008	11	1.0	0.6	0	0.0	0.0
2009	14	1.3	0.7	1	0.1	0.2
2010	19	1.7	0.8	1	0.1	0.2
2011	20	1.8	0.8	3	0.3	0.3
2012	17	1.6	0.8	1	0.1	0.1
2013	17	1.4	0.7	0	0.0	0.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Urinary bladder (incl. in situ)

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	76	20.9	5.5	41	7.5	2.5	32	8.2	3.3	22	4.6	2.1
1977	65	18.0	5.0	37	5.9	2.0	27	8.7	3.9	19	3.2	1.5
1978	64	16.7	4.6	37	5.5	1.9	27	8.3	3.5	17	2.8	1.4
1979	67	17.7	4.8	45	7.2	2.2	26	7.5	3.2	27	4.7	1.9
1980	91	21.5	4.9	50	7.3	2.2	25	7.3	3.2	21	3.6	1.7
1981	67	15.7	4.4	49	7.0	2.1	33	7.9	3.0	29	4.2	1.6
1982	68	16.1	4.2	60	7.8	2.1	25	6.0	2.6	32	4.4	1.6
1983	69	14.7	3.9	58	7.2	1.9	24	5.9	2.6	25	3.5	1.4
1984	71	14.4	3.7	41	5.0	1.6	31	6.1	2.4	22	3.0	1.3
1985	80	17.0	4.3	65	8.4	2.1	33	8.3	3.0	26	3.3	1.3
1986	85	19.9	4.7	44	5.2	1.6	25	6.6	2.8	23	2.8	1.2
1987	84	17.3	4.1	44	5.4	1.7	34	8.0	2.9	25	3.0	1.2
1988	80	15.9	3.8	54	6.5	1.8	31	7.5	2.9	21	2.3	1.0
1989	73	13.9	3.5	51	6.0	1.7	27	6.1	2.5	22	2.7	1.2
1990	86	17.8	4.1	51	5.7	1.6	25	5.3	2.2	25	2.8	1.1
1991	82	15.8	3.7	57	5.9	1.6	28	6.5	2.6	12	1.3	0.7
1992	81	15.5	3.7	71	7.2	1.7	21	4.9	2.3	24	2.6	1.1
1993	76	15.1	3.7	71	7.2	1.7	34	6.5	2.3	24	2.4	1.0
1994	79	14.4	3.4	72	7.1	1.7	27	5.4	2.2	30	3.1	1.1
1995	84	15.4	3.6	46	4.5	1.3	25	4.8	2.1	28	2.9	1.1
1996	88	15.3	3.5	61	5.9	1.5	28	5.5	2.2	23	2.3	1.0
1997	77	13.3	3.2	71	6.8	1.6	26	5.1	2.1	30	3.0	1.1
1998	99	16.0	3.4	72	6.7	1.6	29	5.6	2.2	34	3.3	1.1
1999	107	16.5	3.3	60	5.3	1.4	23	3.9	1.7	26	2.3	0.9
2000	89	14.3	3.1	73	6.4	1.5	36	5.8	2.0	30	2.7	1.0
2001	102	15.2	3.2	72	6.1	1.4	27	4.5	1.8	37	3.2	1.0
2002	98	15.2	3.2	75	6.3	1.4	18	2.7	1.3	29	2.5	0.9
2003	120	16.9	3.2	51	4.2	1.1	26	3.9	1.6	21	1.7	0.7
2004	118	17.0	3.3	71	5.8	1.4	29	4.2	1.6	28	2.2	0.8
2005	127	16.6	3.0	87	7.0	1.5	32	5.0	1.8	25	2.1	0.8
2006	144	20.2	3.5	74	5.9	1.3	31	4.8	1.8	18	1.5	0.7
2007	105	14.8	3.0	74	5.8	1.3	22	3.6	1.6	25	2.0	0.8
2008	129	16.3	3.0	67	5.1	1.2	25	3.9	1.6	24	1.8	0.7
2009	139	17.9	3.1	79	6.0	1.3	22	3.5	1.5	24	1.8	0.7
2010	135	16.5	2.9	72	5.3	1.2	34	4.9	1.7	31	2.4	0.9
2011	150	18.2	3.1	73	5.2	1.2	32	4.3	1.6	29	2.1	0.8
2012	127	15.6	2.9	82	5.6	1.2	38	4.7	1.6	27	1.9	0.7
2013	166	19.4	3.1	94	6.6	1.4	33	4.2	1.5	25	1.7	0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Kidney and renal pelvis

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	38	7.3	2.7	27	4.8	1.9	22	4.7	2.4	15	2.4	1.3
1977	33	6.5	2.6	14	1.6	0.9	16	2.9	1.5	6	0.9	0.8
1978	55	10.3	3.0	26	3.5	1.4	28	5.4	2.3	11	1.4	0.9
1979	44	8.6	2.8	15	1.6	0.8	24	4.4	1.8	8	1.1	0.8
1980	36	6.6	2.3	32	3.8	1.4	21	3.7	1.7	5	0.6	0.5
1981	43	7.3	2.3	20	2.4	1.1	17	2.8	1.3	8	0.8	0.5
1982	47	8.2	2.5	27	3.1	1.2	16	3.0	1.7	11	1.3	0.8
1983	43	7.7	2.5	45	4.9	1.5	23	4.1	1.8	16	1.9	1.0
1984	62	10.4	2.8	35	3.9	1.3	23	4.2	1.9	17	2.0	1.0
1985	45	6.9	2.2	35	3.8	1.3	22	3.6	1.7	20	2.3	1.0
1986	59	8.9	2.4	41	4.3	1.4	32	5.7	2.1	21	2.4	1.1
1987	67	11.4	3.0	44	4.5	1.4	20	3.4	1.6	13	1.4	0.8
1988	57	8.5	2.3	43	4.5	1.4	25	4.1	1.7	14	1.5	0.8
1989	57	9.5	2.7	43	4.0	1.2	14	2.6	1.5	20	2.1	0.9
1990	60	9.3	2.6	47	4.7	1.4	20	3.8	1.9	18	1.8	0.8
1991	81	12.0	2.9	46	4.5	1.3	23	3.7	1.6	17	1.6	0.8
1992	71	10.9	2.8	41	4.1	1.3	20	3.3	1.6	14	1.5	0.8
1993	79	11.3	2.7	41	3.8	1.2	29	4.4	1.8	16	1.7	0.8
1994	61	8.6	2.3	56	4.9	1.3	17	2.5	1.3	14	1.3	0.7
1995	95	13.1	2.8	64	5.9	1.5	32	4.9	1.8	21	2.0	0.9
1996	93	12.4	2.7	53	4.7	1.3	26	3.3	1.3	15	1.4	0.7
1997	85	11.8	2.7	74	6.5	1.5	28	4.5	1.8	16	1.4	0.7
1998	92	11.8	2.6	59	5.2	1.3	27	3.5	1.5	16	1.5	0.7
1999	96	12.7	2.7	100	8.5	1.7	16	2.2	1.2	18	1.6	0.7
2000	89	11.3	2.5	79	6.7	1.5	22	3.1	1.4	18	1.5	0.7
2001	107	13.4	2.7	86	6.8	1.4	24	3.6	1.5	22	1.8	0.7
2002	105	12.9	2.6	81	6.4	1.4	30	4.0	1.5	14	1.2	0.6
2003	133	15.7	2.8	96	7.4	1.5	24	3.1	1.3	20	1.6	0.7
2004	130	14.7	2.6	96	7.4	1.5	32	4.1	1.5	20	1.6	0.7
2005	130	14.4	2.6	120	9.2	1.7	23	2.8	1.2	16	1.3	0.6
2006	150	17.0	2.8	109	8.3	1.6	30	3.8	1.4	27	2.2	0.8
2007	183	20.5	3.1	92	7.0	1.4	34	4.4	1.6	19	1.4	0.7
2008	142	16.0	2.8	130	9.7	1.7	26	3.2	1.3	21	1.6	0.7
2009	168	17.7	2.8	120	8.4	1.5	26	3.3	1.3	12	0.9	0.5
2010	210	22.3	3.2	119	8.6	1.6	31	4.1	1.5	20	1.5	0.7
2011	185	19.2	2.9	128	9.2	1.6	31	3.8	1.4	19	1.3	0.6
2012	175	17.3	2.7	142	10.0	1.7	24	2.7	1.1	23	1.5	0.6
2013	210	20.3	2.8	149	10.0	1.6	28	2.8	1.1	16	1.1	0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Brain and other nervous system

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	25	3.8	1.6	21	2.4	1.1	21	3.7	1.7	15	1.9	1.0
1977	29	4.3	1.7	29	3.0	1.1	19	3.2	1.5	13	1.6	0.9
1978	18	3.0	1.5	14	1.7	0.9	15	2.2	1.2	15	1.8	0.9
1979	21	2.9	1.4	30	3.2	1.2	19	3.4	1.7	20	2.2	1.0
1980	31	5.1	2.0	26	3.1	1.3	22	3.9	1.8	16	1.9	0.9
1981	22	2.7	1.2	18	2.1	1.0	11	1.8	1.2	18	2.4	1.2
1982	35	4.6	1.6	30	3.0	1.1	36	5.8	2.0	20	2.3	1.0
1983	34	4.7	1.7	26	2.8	1.1	18	2.7	1.3	32	3.4	1.2
1984	40	5.2	1.8	37	3.6	1.2	32	4.7	1.8	20	2.1	1.0
1985	34	5.1	1.9	44	4.3	1.3	27	4.2	1.7	18	1.7	0.8
1986	47	5.8	1.8	28	2.8	1.1	30	4.1	1.5	24	2.4	1.0
1987	41	5.4	1.9	42	3.7	1.2	25	4.3	1.9	21	2.0	0.9
1988	37	4.7	1.8	27	2.4	0.9	16	2.3	1.3	21	1.9	0.8
1989	36	4.5	1.6	38	3.4	1.1	15	2.1	1.1	17	1.6	0.8
1990	36	4.3	1.5	45	4.1	1.2	21	3.3	1.6	16	1.5	0.8
1991	44	6.1	2.0	47	4.1	1.2	19	2.5	1.2	23	2.2	0.9
1992	47	6.1	1.9	45	3.9	1.2	20	3.2	1.6	19	1.9	0.9
1993	42	5.1	1.7	26	2.3	0.9	18	2.4	1.2	20	1.7	0.7
1994	48	5.4	1.6	39	3.2	1.0	24	3.2	1.4	27	2.3	0.9
1995	34	3.5	1.3	49	4.2	1.2	14	1.5	0.8	20	1.7	0.7
1996	41	5.0	1.6	58	4.8	1.3	12	1.6	1.0	21	1.7	0.7
1997	36	4.0	1.4	43	3.5	1.0	17	1.9	1.0	21	1.8	0.8
1998	45	4.5	1.4	52	4.2	1.2	15	1.8	1.0	17	1.4	0.7
1999	43	5.2	1.7	47	3.8	1.1	18	2.1	1.0	11	0.9	0.5
2000	47	5.4	1.6	48	3.8	1.1	18	2.1	1.0	23	1.9	0.8
2001	39	3.8	1.2	55	4.3	1.2	19	2.1	1.1	22	1.7	0.7
2002	45	4.9	1.5	39	3.1	1.0	16	1.7	0.9	26	2.0	0.8
2003	55	5.6	1.5	54	4.2	1.1	21	2.4	1.1	13	1.0	0.5
2004	49	5.3	1.5	39	3.0	0.9	27	2.9	1.1	22	1.7	0.7
2005	44	5.0	1.6	51	3.8	1.1	24	2.8	1.2	12	0.9	0.5
2006	41	4.3	1.4	42	3.2	1.0	22	2.6	1.2	23	1.7	0.7
2007	52	5.1	1.4	49	3.7	1.0	14	1.5	0.8	19	1.4	0.6
2008	41	4.2	1.4	42	3.1	1.0	22	2.6	1.1	22	1.6	0.7
2009	44	4.5	1.4	45	3.4	1.0	22	2.4	1.1	17	1.2	0.6
2010	44	4.4	1.3	42	3.1	1.0	16	1.7	0.9	17	1.3	0.6
2011	51	5.3	1.5	49	3.5	1.0	22	2.3	1.0	27	2.0	0.8
2012	53	5.3	1.5	34	2.5	0.8	20	2.1	1.0	20	1.4	0.6
2013	55	4.9	1.4	61	4.5	1.2	23	2.3	1.0	24	1.7	0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Thyroid

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	5	1.0	0.9	23	2.8	1.2	2	0.4	0.5	6	0.8	0.6
1977	1	0.2	0.3	32	3.5	1.3	2	0.3	0.5	2	0.3	0.5
1978	8	1.7	1.4	18	2.0	0.9	0	0.0	0.0	7	0.9	0.7
1979	9	2.0	1.6	27	3.7	1.5	2	0.3	0.5	4	0.4	0.4
1980	3	0.4	0.5	30	3.1	1.2	1	0.2	0.5	6	0.8	0.7
1981	4	0.6	0.6	23	3.0	1.3	0	0.0	0.0	5	0.8	0.7
1982	6	1.0	0.8	26	3.1	1.2	3	0.9	1.2	6	0.7	0.5
1983	3	0.4	0.4	26	2.7	1.1	1	0.3	0.6	5	0.6	0.6
1984	6	0.8	0.6	38	4.0	1.3	2	0.6	1.0	7	0.8	0.6
1985	17	2.5	1.2	32	3.2	1.1	3	0.5	0.6	4	0.5	0.5
1986	10	1.9	1.4	26	2.6	1.1	3	0.7	1.0	5	0.5	0.4
1987	10	1.5	1.1	42	4.0	1.3	2	0.6	0.9	6	0.7	0.6
1988	10	1.6	1.0	31	3.0	1.1	1	0.1	0.2	4	0.5	0.5
1989	4	0.7	0.7	33	3.1	1.1	1	0.2	0.4	7	0.9	0.7
1990	5	0.7	0.6	29	2.6	1.0	1	0.2	0.4	2	0.2	0.3
1991	4	0.7	0.8	36	3.1	1.1	1	0.3	0.6	6	0.7	0.5
1992	16	1.6	0.8	53	4.4	1.2	2	0.2	0.3	7	0.7	0.5
1993	10	1.3	0.9	43	3.8	1.1	1	0.1	0.3	4	0.4	0.4
1994	11	1.5	0.9	49	4.2	1.2	2	0.3	0.4	5	0.4	0.4
1995	11	1.4	0.9	54	4.4	1.2	0	0.0	0.0	5	0.5	0.5
1996	17	2.3	1.2	71	5.6	1.3	7	1.1	0.9	5	0.5	0.4
1997	19	2.4	1.2	71	5.6	1.3	4	0.6	0.7	5	0.4	0.4
1998	20	2.4	1.1	85	6.6	1.4	3	0.3	0.4	5	0.5	0.4
1999	15	1.6	0.8	83	6.6	1.4	2	0.2	0.3	9	0.8	0.5
2000	14	1.4	0.7	98	7.5	1.5	2	0.2	0.3	10	0.9	0.5
2001	24	2.5	1.0	93	7.0	1.4	3	0.3	0.4	12	1.0	0.6
2002	21	2.3	1.0	104	7.7	1.5	1	0.1	0.2	5	0.4	0.4
2003	16	1.9	1.0	96	7.0	1.4	0	0.0	0.0	4	0.3	0.3
2004	25	3.0	1.2	99	7.2	1.4	5	0.8	0.8	5	0.4	0.4
2005	32	3.4	1.2	155	11.4	1.8	5	0.6	0.5	11	0.8	0.5
2006	19	2.2	1.0	137	10.1	1.7	2	0.2	0.3	7	0.5	0.4
2007	32	3.3	1.2	125	9.1	1.6	1	0.2	0.3	8	0.6	0.4
2008	42	4.4	1.4	168	12.0	1.8	3	0.5	0.6	8	0.6	0.4
2009	37	3.5	1.2	169	12.5	1.9	4	0.5	0.6	6	0.4	0.4
2010	32	3.5	1.3	205	14.5	2.0	7	0.9	0.7	7	0.5	0.4
2011	52	5.3	1.5	220	15.0	2.0	1	0.2	0.4	7	0.5	0.4
2012	33	3.4	1.2	209	14.6	2.0	3	0.4	0.5	6	0.4	0.3
2013	38	3.4	1.1	267	18.3	2.2	5	0.5	0.5	7	0.5	0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Hodgkin lymphoma

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	16	3.8	2.6	17	2.0	1.1	9	1.2	0.8	6	0.7	0.6
1977	16	2.3	1.2	14	1.5	0.8	4	1.2	1.6	11	1.4	0.8
1978	14	1.9	1.1	16	1.8	1.0	6	0.8	0.7	6	0.6	0.5
1979	25	3.6	1.5	15	1.6	0.8	11	1.6	1.0	6	0.6	0.5
1980	22	2.5	1.1	24	2.2	0.9	4	0.6	0.6	5	0.4	0.4
1981	25	3.3	1.5	16	1.5	0.8	9	1.6	1.2	8	0.8	0.5
1982	25	2.9	1.2	15	1.4	0.7	7	0.9	0.7	8	0.7	0.5
1983	32	4.3	1.8	23	2.1	0.9	6	0.8	0.6	6	0.6	0.5
1984	28	3.2	1.3	15	1.5	0.8	16	1.9	0.9	2	0.3	0.4
1985	22	2.4	1.1	11	0.8	0.5	16	1.9	1.0	5	0.4	0.4
1986	31	3.7	1.3	14	1.3	0.7	6	0.8	0.7	5	0.5	0.5
1987	27	3.4	1.5	12	1.0	0.6	11	1.4	0.9	2	0.1	0.2
1988	32	3.0	1.1	23	2.0	0.8	4	0.4	0.4	6	0.5	0.4
1989	33	3.7	1.4	19	1.5	0.7	15	1.6	0.8	1	0.1	0.2
1990	25	2.9	1.2	18	1.4	0.6	10	1.2	0.8	5	0.4	0.4
1991	41	4.6	1.6	27	2.0	0.8	5	0.5	0.4	6	0.5	0.4
1992	25	2.3	0.9	28	2.3	0.9	9	0.8	0.6	9	0.7	0.5
1993	22	2.1	0.9	26	1.9	0.7	8	0.7	0.5	6	0.5	0.4
1994	30	3.2	1.3	17	1.3	0.6	9	1.1	0.7	6	0.4	0.4
1995	31	3.1	1.1	24	1.7	0.7	5	0.5	0.5	4	0.3	0.3
1996	23	2.3	1.0	22	1.7	0.7	6	0.8	0.6	6	0.5	0.4
1997	37	3.4	1.1	25	1.9	0.7	4	0.4	0.4	8	0.7	0.5
1998	28	2.5	1.0	28	2.1	0.8	5	0.4	0.4	5	0.4	0.3
1999	27	2.8	1.1	23	1.7	0.7	7	0.8	0.6	3	0.2	0.3
2000	35	3.2	1.1	23	1.7	0.7	3	0.3	0.3	2	0.2	0.2
2001	35	3.0	1.0	17	1.2	0.6	4	0.4	0.4	4	0.3	0.3
2002	33	3.3	1.2	31	2.3	0.8	5	0.5	0.4	2	0.1	0.2
2003	25	2.4	0.9	36	2.6	0.9	7	1.0	0.7	6	0.4	0.3
2004	35	3.2	1.1	30	2.2	0.8	5	0.5	0.5	3	0.2	0.2
2005	45	4.1	1.2	29	2.2	0.8	4	0.4	0.3	1	0.1	0.1
2006	29	2.7	1.0	27	1.9	0.7	1	0.1	0.2	4	0.3	0.3
2007	29	2.9	1.1	26	1.9	0.7	2	0.2	0.2	6	0.5	0.4
2008	53	4.9	1.3	40	3.0	0.9	3	0.3	0.3	2	0.2	0.2
2009	26	2.6	1.1	44	3.2	1.0	4	0.4	0.4	3	0.2	0.2
2010	48	4.6	1.4	33	2.4	0.8	4	0.3	0.3	4	0.3	0.3
2011	47	4.0	1.2	38	2.7	0.9	4	0.5	0.5	3	0.2	0.2
2012	31	2.7	1.0	34	2.5	0.8	3	0.3	0.4	1	0.1	0.1
2013	40	3.7	1.2	41	3.0	0.9	4	0.4	0.4	4	0.3	0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Non-Hodgkin lymphomas

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	32	7.1	2.9	37	4.8	1.6	28	6.2	2.6	17	2.2	1.1
1977	29	5.7	2.5	43	5.6	1.8	14	2.2	1.3	23	3.2	1.5
1978	60	10.4	2.9	49	6.2	1.9	33	6.1	2.3	19	2.2	1.0
1979	48	9.2	2.9	37	5.2	1.8	24	5.2	2.4	26	4.0	1.7
1980	53	9.1	2.7	54	6.7	1.9	34	5.6	2.0	21	2.9	1.4
1981	46	7.1	2.4	53	7.2	2.1	26	4.8	2.1	34	4.2	1.5
1982	57	9.1	2.6	69	8.2	2.0	24	4.8	2.2	29	3.8	1.5
1983	62	9.9	2.8	69	7.8	1.9	39	6.3	2.2	34	4.2	1.5
1984	72	11.4	3.0	78	8.4	1.9	34	5.3	2.1	26	2.9	1.2
1985	98	13.8	3.1	89	9.5	2.1	41	5.9	2.1	42	4.8	1.5
1986	96	15.2	3.5	94	9.8	2.1	40	6.2	2.3	42	4.5	1.4
1987	121	17.8	3.6	80	7.8	1.8	38	5.5	2.0	39	3.8	1.2
1988	101	14.0	3.0	72	7.1	1.7	55	8.9	2.7	30	3.1	1.1
1989	118	15.1	3.0	85	8.3	1.8	51	7.5	2.3	36	3.6	1.2
1990	139	16.4	2.9	93	8.5	1.8	39	5.6	2.0	36	3.6	1.2
1991	135	16.8	3.2	105	10.0	2.0	43	5.8	2.0	32	3.3	1.2
1992	155	19.2	3.2	127	11.7	2.1	41	5.6	1.9	44	4.2	1.3
1993	168	18.9	3.0	99	8.5	1.7	39	5.0	1.7	39	3.6	1.2
1994	184	22.1	3.4	134	12.1	2.1	38	4.8	1.6	49	4.6	1.3
1995	167	20.1	3.3	115	10.3	1.9	49	6.1	1.8	36	3.4	1.1
1996	157	18.1	3.0	139	11.9	2.0	39	5.4	1.9	54	5.0	1.3
1997	188	22.7	3.5	142	12.3	2.0	49	6.8	2.0	46	4.2	1.2
1998	166	20.1	3.3	149	12.5	2.0	53	7.2	2.1	55	4.9	1.3
1999	177	20.7	3.2	155	12.5	2.0	64	8.5	2.3	50	4.2	1.2
2000	154	17.2	2.9	145	11.4	1.9	53	6.8	1.9	46	3.8	1.1
2001	159	18.2	3.0	149	11.9	1.9	57	6.7	1.8	54	4.4	1.2
2002	158	16.9	2.7	140	10.7	1.8	61	7.5	2.0	44	3.5	1.0
2003	156	17.3	2.8	183	13.9	2.0	57	6.8	1.9	52	4.0	1.1
2004	154	17.4	2.9	156	11.9	1.9	46	5.4	1.6	41	3.2	1.0
2005	172	18.3	2.9	188	14.1	2.0	45	5.5	1.7	46	3.6	1.0
2006	150	16.3	2.7	167	12.7	1.9	35	4.2	1.5	46	3.5	1.0
2007	194	20.2	2.9	158	11.9	1.9	45	5.3	1.6	55	4.1	1.1
2008	188	19.6	2.9	149	11.0	1.8	39	4.5	1.5	43	3.2	1.0
2009	185	19.9	3.0	194	14.5	2.1	40	4.5	1.5	46	3.4	1.0
2010	185	19.0	2.9	174	12.5	1.9	39	4.4	1.5	46	3.3	1.0
2011	208	21.8	3.1	197	13.7	1.9	49	5.4	1.6	43	3.1	0.9
2012	177	18.0	2.8	194	13.7	2.0	47	5.2	1.6	43	3.0	0.9
2013	189	18.3	2.7	188	12.8	1.9	50	5.5	1.6	45	3.0	0.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Myeloma

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	45	11.7	3.8	47	7.6	2.4	19	4.2	2.0	31	5.5	2.1
1977	40	10.3	3.8	51	7.8	2.3	28	7.9	3.4	34	5.2	1.9
1978	52	14.0	4.4	40	5.8	1.9	25	6.8	3.1	20	2.7	1.2
1979	48	9.2	2.8	58	8.3	2.2	23	5.1	2.3	33	4.9	1.8
1980	48	11.4	3.7	72	9.7	2.4	34	8.1	3.1	43	5.9	1.9
1981	49	10.2	3.1	63	8.5	2.2	21	5.1	2.5	35	4.8	1.7
1982	52	10.7	3.2	66	8.9	2.3	42	8.9	3.0	43	5.8	1.8
1983	48	9.0	2.7	71	9.1	2.2	28	6.0	2.4	48	6.2	1.9
1984	54	11.8	3.6	78	9.7	2.2	30	5.9	2.3	59	7.5	2.0
1985	54	11.6	3.5	80	9.0	2.0	38	8.0	2.8	42	4.8	1.5
1986	61	11.8	3.3	74	8.4	2.0	34	6.9	2.5	45	5.4	1.6
1987	60	10.7	2.9	65	7.1	1.8	40	7.6	2.6	37	4.1	1.4
1988	48	8.4	2.6	61	6.6	1.7	29	5.6	2.3	34	3.8	1.3
1989	62	12.0	3.4	72	8.1	1.9	28	6.2	2.5	33	3.7	1.3
1990	56	10.6	3.2	83	8.2	1.8	26	5.7	2.4	39	4.2	1.3
1991	47	8.7	2.8	98	10.2	2.1	33	6.6	2.5	51	5.4	1.5
1992	79	13.8	3.4	98	9.6	1.9	30	5.7	2.3	52	5.0	1.4
1993	55	8.4	2.3	75	7.4	1.7	32	6.0	2.3	43	4.3	1.3
1994	60	10.4	2.8	85	8.3	1.8	27	4.4	1.8	51	5.1	1.4
1995	68	11.5	3.0	95	9.0	1.8	45	7.6	2.4	63	6.0	1.5
1996	67	10.1	2.6	93	8.6	1.8	38	6.4	2.2	43	4.0	1.2
1997	74	10.9	2.7	102	9.4	1.8	39	6.5	2.2	48	4.4	1.3
1998	77	11.5	2.8	102	9.2	1.8	26	4.1	1.7	50	4.5	1.3
1999	68	9.5	2.4	119	10.5	1.9	31	4.4	1.6	53	4.7	1.3
2000	77	10.6	2.6	113	9.5	1.8	17	2.8	1.4	54	4.8	1.3
2001	87	12.1	2.7	124	10.4	1.8	36	5.4	1.9	51	4.3	1.2
2002	88	12.2	2.7	130	10.6	1.8	43	5.7	1.8	64	5.4	1.3
2003	94	13.1	2.8	134	10.9	1.9	38	4.8	1.7	51	4.2	1.2
2004	86	10.6	2.4	123	9.7	1.7	39	5.3	1.8	55	4.6	1.2
2005	106	13.2	2.6	140	10.8	1.8	35	4.6	1.6	52	4.1	1.1
2006	117	14.6	2.8	134	10.4	1.8	34	4.4	1.6	46	3.6	1.1
2007	100	13.0	2.7	119	9.1	1.7	48	6.0	1.8	61	4.7	1.2
2008	133	16.2	2.9	155	11.4	1.8	31	4.0	1.5	45	3.4	1.0
2009	151	17.2	2.9	148	11.1	1.8	52	6.2	1.8	41	3.1	1.0
2010	148	17.4	2.9	137	9.9	1.7	45	5.7	1.7	56	4.2	1.1
2011	160	17.6	2.9	171	12.3	1.9	47	5.5	1.6	51	3.5	1.0
2012	143	15.9	2.7	186	12.8	1.9	47	5.0	1.5	48	3.3	1.0
2013	141	15.2	2.6	159	10.7	1.7	52	5.7	1.6	44	3.0	0.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York City
Blacks
Leukemias

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	51	9.6	3.0	50	6.7	2.0	41	8.1	3.2	33	4.8	1.8
1977	46	9.5	3.5	45	6.0	1.8	40	8.5	3.0	41	5.0	1.6
1978	60	11.9	3.6	57	7.5	2.1	47	8.6	2.7	36	5.1	1.8
1979	57	12.6	4.0	50	6.6	1.9	44	9.6	3.4	32	4.0	1.5
1980	53	10.1	3.3	62	7.9	2.1	36	7.9	3.2	45	5.8	1.8
1981	61	12.1	3.6	55	6.9	1.9	33	6.4	2.7	48	6.0	1.8
1982	63	12.1	3.5	61	7.1	1.9	36	7.5	3.0	36	4.3	1.5
1983	65	11.6	3.2	64	7.2	1.8	38	6.5	2.3	41	4.8	1.6
1984	70	10.7	2.9	62	7.2	1.9	41	7.2	2.6	42	4.7	1.5
1985	82	13.4	3.3	53	5.7	1.6	49	8.9	2.9	29	3.1	1.2
1986	68	11.7	3.2	61	6.3	1.6	40	6.6	2.3	35	3.5	1.2
1987	67	10.3	2.9	73	7.6	1.8	36	5.5	2.2	38	3.9	1.3
1988	58	10.6	3.1	53	5.4	1.5	35	6.2	2.4	41	4.6	1.5
1989	79	12.5	3.2	76	7.5	1.7	47	8.4	2.7	35	3.6	1.2
1990	75	10.9	2.9	80	8.0	1.8	45	7.5	2.5	50	5.0	1.4
1991	89	13.5	3.2	91	8.6	1.8	32	5.3	2.1	39	3.9	1.2
1992	73	11.1	3.0	98	9.3	1.9	36	5.4	2.0	50	5.1	1.5
1993	75	11.3	2.9	77	7.3	1.7	39	5.9	2.1	45	4.4	1.3
1994	79	11.7	2.9	76	7.1	1.6	51	7.4	2.3	62	6.0	1.5
1995	80	10.2	2.4	84	7.6	1.7	43	6.4	2.2	50	4.6	1.3
1996	94	12.4	2.8	82	7.4	1.6	45	6.5	2.1	52	4.9	1.3
1997	64	8.8	2.4	96	8.5	1.7	49	7.8	2.4	46	4.1	1.2
1998	67	9.3	2.5	86	7.4	1.6	34	4.6	1.7	42	3.7	1.1
1999	75	10.3	2.6	69	5.6	1.3	47	7.0	2.2	45	3.9	1.1
2000	91	11.6	2.6	101	8.4	1.7	35	5.4	2.0	58	4.9	1.3
2001	70	9.0	2.3	93	7.7	1.6	41	5.6	1.8	49	4.1	1.2
2002	99	12.2	2.6	85	6.9	1.5	47	6.4	2.0	46	3.7	1.1
2003	90	10.1	2.2	93	7.3	1.5	40	4.8	1.6	50	4.1	1.1
2004	99	11.8	2.5	83	6.5	1.4	42	5.5	1.8	39	3.2	1.0
2005	86	10.3	2.3	84	6.5	1.4	41	5.6	1.8	48	3.8	1.1
2006	107	12.9	2.6	77	5.9	1.3	47	5.9	1.8	33	2.5	0.9
2007	85	9.4	2.1	98	7.5	1.5	45	6.0	1.9	40	3.2	1.0
2008	80	10.3	2.4	97	7.3	1.5	51	6.5	1.9	44	3.3	1.0
2009	107	12.2	2.4	96	7.1	1.4	43	4.9	1.5	40	3.0	0.9
2010	128	14.2	2.6	115	8.4	1.6	51	6.3	1.8	51	3.8	1.0
2011	116	11.9	2.3	112	8.3	1.6	57	5.9	1.6	55	4.0	1.1
2012	111	11.5	2.3	101	7.2	1.4	46	5.4	1.6	44	3.2	1.0
2013	108	10.8	2.1	127	8.9	1.6	57	6.4	1.7	35	2.4	0.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
All Invasive Malignant Tumors

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	711	520.3	44.7	676	363.1	29.5	435	330.5	36.8	337	195.2	22.6
1977	785	559.0	45.1	718	375.1	29.5	517	397.0	39.8	350	192.6	21.7
1978	837	597.1	47.2	744	376.1	28.7	517	371.4	37.1	367	194.0	21.2
1979	827	577.7	45.7	746	368.9	28.2	531	386.8	37.9	414	217.0	22.4
1980	917	611.7	45.3	801	375.4	27.4	545	379.6	36.5	420	204.4	20.6
1981	908	606.2	45.2	765	349.1	26.0	590	416.5	38.6	444	213.0	20.8
1982	885	557.5	41.1	758	327.9	24.4	579	383.6	35.3	430	193.6	19.1
1983	925	570.6	41.6	792	337.0	24.4	603	390.0	35.0	427	187.3	18.4
1984	981	577.9	40.6	833	347.9	24.6	593	369.7	33.7	468	205.1	19.3
1985	975	562.3	39.5	920	374.4	25.1	582	347.4	31.4	470	199.3	18.7
1986	1039	598.0	41.1	925	367.1	24.5	655	397.4	34.6	498	206.6	18.7
1987	970	550.5	39.0	925	355.9	23.7	629	387.5	34.4	479	191.3	17.7
1988	1039	573.5	39.1	1004	376.3	23.9	660	404.5	34.8	512	199.3	17.8
1989	1139	596.1	39.1	1010	365.0	23.2	691	395.3	33.0	537	203.1	17.6
1990	1173	618.0	40.0	1075	377.3	23.2	661	362.1	31.2	548	203.6	17.5
1991	1296	647.3	39.9	1091	367.3	22.4	683	377.2	31.9	559	197.1	16.8
1992	1408	688.4	40.8	1108	361.7	21.8	651	341.9	29.7	596	205.5	16.9
1993	1476	690.4	39.5	1169	371.2	21.8	707	364.3	30.2	567	189.1	15.9
1994	1440	659.9	38.0	1229	377.0	21.5	772	391.4	31.0	567	184.4	15.5
1995	1582	692.0	38.0	1262	379.9	21.4	708	346.1	28.5	611	194.0	15.7
1996	1720	710.3	37.0	1367	397.5	21.5	679	317.2	26.6	610	187.7	15.2
1997	1692	697.8	36.7	1375	392.4	21.1	727	332.5	26.8	596	178.4	14.6
1998	1768	693.0	35.5	1481	408.5	21.2	747	333.8	26.4	646	190.5	14.9
1999	1777	673.5	34.2	1482	402.9	20.9	695	300.3	24.9	637	184.0	14.5
2000	1812	676.8	34.0	1418	373.5	19.8	760	321.0	25.2	686	188.9	14.4
2001	1935	691.5	33.7	1488	379.8	19.6	684	287.8	23.9	668	178.8	13.8
2002	1951	683.1	33.2	1579	386.8	19.4	752	294.6	23.5	674	175.4	13.5
2003	1971	660.6	31.8	1597	384.2	19.2	740	278.2	22.1	673	170.7	13.1
2004	1970	630.1	30.3	1739	407.1	19.5	745	280.0	22.1	673	167.3	12.9
2005	2011	636.5	30.2	1731	398.1	19.1	747	261.4	20.5	684	164.5	12.5
2006	2159	653.4	29.8	1748	389.6	18.6	653	220.5	18.4	656	153.5	11.9
2007	2231	652.7	29.1	1813	392.1	18.3	719	240.4	19.0	735	167.3	12.3
2008	2268	631.9	27.9	1911	403.0	18.3	693	217.3	17.5	698	152.8	11.5
2009	2309	611.5	26.7	1965	405.9	18.2	759	235.3	18.2	712	151.9	11.3
2010	2358	617.6	26.8	1925	382.8	17.4	782	240.3	18.2	703	145.0	10.9
2011	2483	622.7	26.3	2078	401.6	17.6	777	221.1	16.8	754	152.7	11.1
2012	2290	550.7	24.1	2052	384.5	16.9	791	220.2	16.5	749	146.7	10.7
2013	2292	534.8	23.3	2126	387.0	16.7	765	207.2	15.7	762	142.7	10.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Oral cavity and pharynx

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	38	20.9	6.8	18	7.7	3.7	13	13.2	8.7	6	2.6	2.1
1977	39	22.1	7.3	11	4.8	2.9	17	9.0	4.3	7	3.5	2.7
1978	38	20.6	6.8	26	10.8	4.2	20	12.4	5.7	6	2.5	2.0
1979	47	26.0	7.8	11	4.7	2.9	31	17.9	6.8	8	3.2	2.2
1980	42	21.4	6.7	22	8.5	3.6	29	16.9	6.4	8	3.3	2.3
1981	53	26.6	7.8	27	11.2	4.4	22	11.7	5.1	10	4.3	2.7
1982	44	21.5	6.5	21	8.4	3.7	27	13.4	5.3	8	2.9	2.0
1983	55	28.2	8.2	19	7.5	3.5	24	12.3	5.5	11	4.4	2.6
1984	56	27.7	7.6	20	7.3	3.2	24	12.5	5.3	6	2.4	1.9
1985	57	29.0	8.1	21	8.2	3.6	25	12.0	5.0	9	3.3	2.3
1986	49	23.5	7.1	21	7.5	3.3	25	12.8	5.5	10	4.0	2.5
1987	50	23.0	6.7	30	11.3	4.2	20	10.4	5.0	9	3.4	2.3
1988	53	24.0	7.0	22	7.4	3.1	25	12.0	5.3	12	4.2	2.4
1989	78	32.8	7.7	31	10.3	3.7	29	12.0	4.5	14	4.8	2.6
1990	53	23.5	6.9	15	5.1	2.6	18	6.9	3.3	9	3.0	2.0
1991	71	30.2	7.3	20	6.4	2.8	24	10.5	4.5	11	3.8	2.3
1992	58	24.5	7.2	29	8.7	3.2	31	12.2	4.5	7	2.2	1.6
1993	57	21.8	6.0	26	8.1	3.1	13	5.8	3.4	7	2.0	1.5
1994	54	19.6	5.5	21	5.9	2.6	23	8.9	3.8	5	1.4	1.2
1995	62	22.3	5.9	21	6.6	2.9	30	10.5	3.9	10	3.0	1.9
1996	67	27.2	7.1	27	7.7	3.0	17	6.7	3.4	10	3.0	1.9
1997	70	23.6	5.8	35	9.3	3.1	17	9.4	5.0	7	2.3	1.7
1998	55	18.2	5.1	29	7.7	2.8	25	8.6	3.6	2	0.6	0.8
1999	62	20.8	5.6	20	5.5	2.5	22	8.3	3.8	10	2.5	1.6
2000	45	15.4	4.8	26	6.5	2.5	26	9.0	3.7	12	3.4	1.9
2001	50	14.3	4.1	13	3.5	1.9	9	3.7	2.8	7	2.1	1.5
2002	63	19.9	5.3	29	7.5	2.8	20	6.7	3.2	6	1.7	1.4
2003	50	14.0	4.2	30	7.2	2.6	23	6.9	2.9	6	1.7	1.3
2004	54	14.7	4.1	21	4.8	2.1	25	7.1	2.9	6	1.4	1.2
2005	45	13.0	4.0	23	5.1	2.1	16	5.2	2.7	4	1.0	1.0
2006	51	13.7	4.0	19	4.0	1.8	18	5.2	2.5	4	0.9	0.9
2007	62	16.3	4.3	22	4.7	2.0	15	4.8	2.7	8	1.8	1.2
2008	53	12.9	3.7	34	7.1	2.4	12	3.0	1.8	7	1.5	1.1
2009	55	13.3	3.8	33	6.4	2.2	12	3.3	2.0	5	1.0	0.9
2010	62	14.2	3.7	29	5.3	1.9	12	2.7	1.6	5	1.0	0.9
2011	60	13.2	3.5	17	3.4	1.7	18	4.1	2.0	9	1.8	1.2
2012	40	9.3	3.1	35	6.2	2.1	18	4.3	2.1	4	0.7	0.7
2013	45	9.4	2.9	23	4.0	1.7	16	3.6	1.8	6	1.1	0.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Esophagus

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	27	18.6	7.4	15	7.9	4.3	37	25.5	9.6	11	5.6	3.6
1977	37	22.3	7.6	14	6.7	3.8	38	24.7	8.4	10	4.3	2.7
1978	50	30.2	9.3	16	9.1	4.8	41	24.3	8.3	13	6.4	3.7
1979	41	29.0	9.9	8	4.0	2.9	36	25.3	9.4	6	2.6	2.1
1980	44	26.2	8.3	13	6.0	3.4	40	25.4	8.7	7	3.7	2.8
1981	42	23.8	7.9	13	5.7	3.2	34	19.6	7.1	11	4.4	2.6
1982	33	20.5	7.8	8	3.4	2.3	38	23.2	8.1	8	3.8	2.7
1983	41	23.3	8.0	11	4.2	2.5	38	23.1	8.4	11	4.6	2.8
1984	40	20.7	6.8	14	5.8	3.1	33	20.8	8.0	14	5.7	3.1
1985	42	24.3	8.1	10	4.0	2.5	37	18.4	6.4	8	3.2	2.3
1986	49	27.5	8.6	17	6.4	3.1	38	23.8	8.7	15	5.8	3.0
1987	31	15.8	6.0	13	5.0	2.8	32	16.4	6.2	9	3.7	2.5
1988	37	18.4	6.3	15	5.5	2.8	37	20.7	7.3	17	6.4	3.1
1989	48	25.0	7.8	17	6.1	3.0	35	16.0	5.7	7	2.5	1.9
1990	39	17.6	6.1	18	6.0	2.8	37	19.3	6.9	19	6.7	3.1
1991	42	19.8	6.5	15	4.8	2.5	35	15.2	5.3	14	4.6	2.4
1992	42	19.8	6.6	10	3.5	2.2	27	13.3	5.6	11	3.8	2.3
1993	38	15.1	5.2	15	4.5	2.3	39	16.8	5.8	5	1.6	1.4
1994	37	14.8	5.0	20	6.3	2.8	34	15.2	5.4	10	3.2	2.0
1995	28	10.9	4.3	15	4.7	2.4	35	13.8	4.8	13	4.1	2.3
1996	44	17.4	5.6	13	4.1	2.3	29	11.4	4.4	14	4.2	2.3
1997	32	13.9	5.3	13	3.7	2.0	24	9.8	4.3	14	3.9	2.1
1998	38	14.7	5.0	16	4.9	2.4	36	13.7	4.8	16	5.0	2.5
1999	47	17.5	5.3	17	4.5	2.2	21	8.6	4.0	10	2.9	1.8
2000	35	13.2	4.7	18	5.1	2.4	38	15.4	5.2	16	4.3	2.1
2001	35	13.7	5.0	17	4.3	2.0	25	9.1	3.9	11	2.9	1.7
2002	29	10.3	4.1	11	2.8	1.7	39	12.7	4.3	12	2.9	1.7
2003	43	14.1	4.5	13	3.6	2.0	26	9.6	4.2	11	3.0	1.8
2004	31	10.0	3.6	19	5.1	2.3	32	11.9	4.5	11	2.8	1.7
2005	36	10.6	3.7	12	2.7	1.5	24	8.0	3.4	12	3.1	1.8
2006	27	11.0	4.6	11	2.4	1.4	19	7.4	3.6	6	1.3	1.1
2007	37	10.5	3.6	11	2.5	1.5	23	7.7	3.4	10	2.4	1.5
2008	26	7.7	3.3	10	2.1	1.3	22	5.8	2.6	14	3.1	1.6
2009	34	9.5	3.5	9	2.0	1.3	23	7.4	3.3	5	1.0	0.9
2010	32	9.2	3.4	10	2.1	1.3	32	8.4	3.2	11	2.4	1.4
2011	32	8.6	3.3	15	2.9	1.5	19	5.3	2.6	7	1.5	1.1
2012	31	6.7	2.5	11	2.0	1.2	22	5.3	2.4	10	2.0	1.3
2013	34	8.9	3.2	14	2.6	1.4	26	6.4	2.6	13	2.5	1.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Stomach

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	38	32.3	12.1	27	16.3	6.7	29	24.3	10.3	12	6.8	4.2
1977	34	25.1	9.4	23	12.9	5.7	31	25.5	10.2	20	12.1	5.7
1978	47	39.8	13.1	21	13.0	5.9	25	21.5	9.9	25	14.6	6.1
1979	31	24.0	10.2	36	20.3	7.0	23	18.9	9.1	19	10.8	5.1
1980	41	31.8	11.1	37	21.4	7.2	22	19.5	9.0	25	13.9	5.7
1981	40	29.8	10.7	21	12.0	5.3	38	27.0	9.6	24	14.8	6.1
1982	33	24.1	9.0	17	8.3	4.1	24	18.9	8.4	17	9.1	4.4
1983	33	21.4	8.0	28	12.5	4.9	23	15.7	7.2	18	8.4	4.1
1984	41	25.1	8.6	22	10.1	4.3	29	23.7	9.5	16	6.6	3.3
1985	38	21.6	7.9	25	12.3	5.0	25	13.1	5.5	21	9.5	4.2
1986	31	18.7	7.6	19	8.9	4.1	28	18.0	7.6	16	7.3	3.7
1987	46	27.9	9.1	30	13.9	5.1	27	14.9	6.0	19	8.6	4.0
1988	44	29.6	9.8	27	12.1	4.7	23	17.4	8.0	23	10.6	4.4
1989	43	22.7	7.7	31	13.1	4.7	39	22.2	7.7	19	8.0	3.7
1990	52	31.3	9.8	23	9.0	3.8	31	16.7	7.0	17	7.3	3.5
1991	38	20.8	7.6	31	11.3	4.1	36	19.5	7.3	18	6.5	3.1
1992	36	18.4	6.9	36	12.6	4.2	24	11.0	4.7	37	12.9	4.3
1993	54	27.3	8.2	31	12.0	4.3	29	15.0	6.3	15	5.9	3.0
1994	54	27.0	8.3	26	8.5	3.3	39	21.0	7.7	17	5.7	2.8
1995	42	22.0	7.6	43	13.9	4.2	31	16.6	6.6	22	7.3	3.1
1996	50	24.9	7.7	29	9.1	3.4	26	13.6	6.0	10	3.4	2.2
1997	45	19.4	6.4	29	8.7	3.2	27	11.7	4.7	11	3.2	1.9
1998	56	22.5	6.6	33	9.9	3.4	37	16.4	5.9	19	6.0	2.8
1999	50	22.5	7.2	30	9.1	3.3	35	16.3	6.2	21	6.4	2.8
2000	41	16.0	5.2	35	10.0	3.4	23	10.6	4.7	24	7.0	2.8
2001	42	18.1	6.2	39	11.0	3.5	24	8.4	3.8	12	3.5	2.0
2002	45	16.7	5.5	32	8.6	3.0	31	13.6	5.5	20	5.7	2.5
2003	45	14.9	4.7	26	6.6	2.6	28	9.3	3.8	10	2.7	1.7
2004	33	13.2	4.9	38	9.5	3.0	20	7.9	3.8	19	4.7	2.2
2005	45	15.3	4.9	56	13.6	3.6	39	15.0	5.2	25	6.7	2.7
2006	60	20.9	5.7	39	9.4	3.0	28	9.4	3.7	17	4.4	2.1
2007	52	18.0	5.3	45	10.3	3.0	29	10.8	4.3	22	5.4	2.3
2008	51	16.1	4.7	28	6.0	2.3	33	11.2	4.2	24	5.3	2.2
2009	52	15.7	4.6	40	9.0	2.8	23	8.2	3.6	20	4.5	2.0
2010	50	14.5	4.3	35	7.8	2.6	32	10.5	3.9	14	3.1	1.6
2011	62	16.8	4.5	47	9.9	2.9	25	7.0	2.9	18	3.9	1.8
2012	58	15.2	4.2	47	9.6	2.8	29	8.4	3.3	21	4.6	2.0
2013	50	12.9	3.9	49	9.2	2.6	31	8.2	3.1	21	4.1	1.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Colorectal

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	63	47.4	13.8	97	58.1	12.5	35	28.6	11.3	47	30.7	9.6
1977	81	64.9	16.2	107	60.7	12.4	51	41.0	12.4	45	27.8	8.7
1978	87	66.5	16.4	106	57.2	11.5	54	47.9	15.0	44	25.4	7.9
1979	103	72.4	16.0	98	54.5	11.4	45	34.9	11.6	76	44.1	10.6
1980	97	78.6	18.0	122	63.3	11.8	51	35.8	10.9	56	30.2	8.4
1981	97	75.7	16.9	111	56.8	11.1	61	48.3	13.4	65	34.3	8.8
1982	86	57.9	13.9	107	51.3	10.1	48	39.2	12.8	65	32.2	8.1
1983	86	63.9	15.4	113	54.5	10.4	46	31.8	10.6	53	26.3	7.4
1984	114	77.5	16.0	116	53.9	10.2	43	30.3	10.3	60	27.5	7.2
1985	101	66.1	14.6	117	52.5	9.9	49	34.2	10.6	67	31.6	7.8
1986	115	67.8	13.8	123	54.2	9.9	56	39.1	11.7	63	28.7	7.3
1987	126	77.4	15.4	121	50.6	9.3	71	49.7	13.0	54	21.9	6.0
1988	109	62.1	13.1	135	54.3	9.4	64	43.2	12.1	61	26.5	6.8
1989	124	72.4	14.5	137	53.2	9.1	75	53.8	13.6	52	20.9	5.8
1990	111	60.0	12.5	130	49.5	8.8	64	36.6	10.1	68	26.8	6.5
1991	119	56.5	11.3	124	45.5	8.2	64	36.8	10.3	62	23.3	5.9
1992	136	69.7	13.2	142	50.5	8.5	53	28.8	8.7	63	23.6	6.0
1993	122	56.5	11.2	140	49.0	8.3	70	36.2	9.5	69	24.8	6.0
1994	159	78.2	13.7	142	47.5	8.0	78	38.2	9.6	76	25.7	5.9
1995	146	67.5	12.3	168	54.0	8.3	63	34.1	9.6	64	21.4	5.3
1996	168	74.1	12.3	159	49.8	7.9	60	27.8	7.7	66	22.2	5.4
1997	157	70.0	12.1	162	48.5	7.6	78	34.6	8.7	70	22.2	5.3
1998	160	67.3	11.6	177	52.0	7.8	65	27.9	7.5	64	19.2	4.8
1999	191	77.1	12.1	198	57.3	8.1	73	35.5	9.1	80	24.3	5.4
2000	173	68.0	11.2	180	50.6	7.5	84	36.0	8.6	76	21.1	4.8
2001	192	72.6	11.2	178	47.0	7.0	62	25.7	7.0	64	17.6	4.4
2002	188	69.3	10.9	224	57.9	7.7	69	29.1	7.7	84	22.6	4.9
2003	203	76.1	11.5	193	49.7	7.1	67	26.4	7.0	71	18.4	4.3
2004	218	74.8	11.0	205	51.2	7.1	79	29.8	7.2	76	19.9	4.5
2005	198	67.1	10.2	197	46.7	6.6	67	23.9	6.2	74	18.1	4.2
2006	192	57.9	9.0	203	48.3	6.7	74	25.7	6.4	72	17.6	4.1
2007	192	55.3	8.5	190	43.0	6.2	89	28.3	6.4	88	20.5	4.3
2008	220	64.7	9.3	216	46.7	6.3	61	19.8	5.4	61	13.6	3.5
2009	212	57.3	8.2	202	41.7	5.8	76	23.1	5.6	69	14.4	3.4
2010	206	57.2	8.4	180	36.5	5.4	64	18.3	4.9	62	12.7	3.2
2011	211	54.8	7.9	193	38.5	5.5	71	20.8	5.2	66	13.3	3.3
2012	186	45.5	7.1	172	32.9	5.0	59	18.5	5.1	65	13.1	3.2
2013	189	45.5	7.0	193	35.5	5.1	71	17.9	4.5	66	12.0	2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Colon excluding rectum

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	40	31.1	11.8	69	41.3	10.6	30	24.1	10.6	38	24.1	8.5
1977	55	45.3	13.8	83	47.7	11.0	39	32.0	11.1	34	20.5	7.3
1978	64	47.7	13.5	77	41.1	9.6	43	39.0	13.8	35	19.7	6.9
1979	75	53.1	13.6	77	42.7	10.1	41	31.9	11.0	67	38.9	9.9
1980	67	55.5	15.3	90	45.8	10.0	41	30.0	10.2	42	22.5	7.2
1981	71	57.2	15.0	83	43.2	9.7	51	41.4	12.5	53	27.0	7.6
1982	69	47.5	12.6	77	36.7	8.5	38	34.4	12.4	57	27.7	7.5
1983	60	45.6	13.2	93	44.8	9.5	39	28.1	10.2	48	24.1	7.1
1984	83	57.5	13.7	93	43.6	9.2	36	26.2	9.7	53	24.2	6.7
1985	77	51.8	13.1	96	43.5	9.0	42	30.1	10.1	63	29.6	7.6
1986	77	45.7	11.4	96	41.5	8.6	45	30.6	10.1	54	24.9	6.9
1987	97	60.7	13.7	91	38.3	8.1	66	45.3	12.3	49	19.9	5.7
1988	85	47.4	11.3	111	44.6	8.5	55	36.1	10.9	50	21.7	6.2
1989	89	53.7	12.7	105	40.7	8.0	66	46.7	12.7	46	18.2	5.4
1990	82	44.6	11.0	97	36.6	7.5	55	30.1	8.8	57	22.5	6.0
1991	93	45.5	10.3	96	35.1	7.2	57	32.7	9.7	54	20.3	5.6
1992	103	53.2	11.5	115	41.4	7.7	44	23.1	7.6	52	19.2	5.4
1993	89	43.7	10.3	108	38.0	7.3	61	30.7	8.6	63	22.3	5.6
1994	119	59.5	12.0	106	36.3	7.1	67	33.7	9.1	68	23.0	5.6
1995	107	50.9	10.9	131	42.7	7.4	54	30.6	9.3	53	18.0	4.9
1996	105	46.7	9.8	129	40.4	7.1	44	20.2	6.5	55	18.2	4.9
1997	112	50.9	10.5	129	38.5	6.8	67	30.7	8.3	61	19.3	4.9
1998	126	54.9	10.6	137	40.8	6.9	55	23.4	6.9	57	17.1	4.5
1999	140	58.4	10.8	150	43.2	7.0	61	29.5	8.3	74	22.3	5.2
2000	134	54.6	10.3	140	39.5	6.7	72	32.5	8.4	63	17.6	4.4
2001	138	54.2	9.8	148	39.1	6.4	53	22.5	6.6	57	15.9	4.2
2002	147	57.8	10.3	186	48.3	7.0	57	24.6	7.1	76	20.5	4.7
2003	153	59.3	10.3	147	38.1	6.3	49	19.6	6.0	64	16.4	4.1
2004	166	58.5	9.9	163	41.3	6.4	68	25.3	6.6	70	18.3	4.4
2005	152	53.7	9.3	147	35.6	5.8	54	19.8	5.8	66	16.1	3.9
2006	139	42.7	7.8	154	37.1	5.9	65	22.4	5.9	60	14.7	3.8
2007	147	43.0	7.5	145	33.1	5.5	70	22.7	5.8	71	16.4	3.9
2008	157	47.5	8.0	175	38.0	5.7	52	17.4	5.1	54	12.1	3.3
2009	159	43.1	7.2	150	31.2	5.1	62	19.5	5.2	62	12.9	3.3
2010	149	41.9	7.2	133	27.3	4.7	52	15.1	4.4	51	10.5	2.9
2011	149	39.8	6.9	155	30.8	4.9	61	18.6	5.0	57	11.5	3.0
2012	131	33.1	6.1	124	24.1	4.3	50	15.1	4.5	61	12.4	3.1
2013	130	32.1	5.9	137	25.1	4.3	63	15.5	4.1	58	10.7	2.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Rectum & rectosigmoid

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	23	16.3	7.2	28	16.8	6.6	5	4.5	4.1	9	6.6	4.6
1977	26	19.6	8.4	24	13.0	5.7	12	9.0	5.7	11	7.3	4.7
1978	23	18.8	9.3	29	16.1	6.3	11	8.9	5.8	9	5.8	3.8
1979	28	19.3	8.4	21	11.8	5.3	4	3.1	3.5	9	5.2	3.6
1980	30	23.1	9.4	32	17.5	6.4	10	5.8	3.8	14	7.8	4.3
1981	26	18.5	7.9	28	13.7	5.3	10	6.9	4.7	12	7.3	4.4
1982	17	10.3	5.8	30	14.6	5.5	10	4.8	3.0	8	4.4	3.2
1983	26	18.4	7.9	20	9.7	4.4	7	3.7	2.9	5	2.2	2.0
1984	31	20.0	8.1	23	10.4	4.3	7	4.1	3.4	7	3.3	2.5
1985	24	14.3	6.3	21	9.0	4.0	7	4.1	3.2	4	2.0	2.0
1986	38	22.2	7.8	27	12.6	4.9	11	8.5	5.9	9	3.8	2.6
1987	29	16.7	7.0	30	12.4	4.6	5	4.3	4.2	5	2.0	1.8
1988	24	14.8	6.6	24	9.6	4.0	9	7.1	5.1	11	4.8	2.9
1989	35	18.8	7.1	32	12.4	4.4	9	7.1	5.0	6	2.7	2.2
1990	29	15.4	6.1	33	12.9	4.5	9	6.5	4.9	11	4.3	2.6
1991	26	11.1	4.6	28	10.4	3.9	7	4.1	3.4	8	3.0	2.1
1992	33	16.5	6.5	27	9.1	3.5	9	5.7	4.3	11	4.4	2.6
1993	33	12.8	4.6	32	11.1	3.9	9	5.4	4.1	6	2.5	2.0
1994	40	18.8	6.5	36	11.1	3.7	11	4.5	2.9	8	2.6	1.9
1995	39	16.6	5.7	37	11.3	3.7	9	3.5	2.4	11	3.4	2.0
1996	63	27.4	7.5	30	9.4	3.4	16	7.5	4.1	11	4.0	2.4
1997	45	19.1	6.1	33	9.9	3.5	11	4.0	2.5	9	3.0	2.0
1998	34	12.5	4.6	40	11.3	3.5	10	4.5	3.0	7	2.0	1.5
1999	51	18.7	5.6	48	14.1	4.1	12	6.0	3.7	6	2.0	1.6
2000	39	13.4	4.6	40	11.0	3.5	12	3.6	2.0	13	3.5	1.9
2001	54	18.3	5.4	30	7.9	2.9	9	3.2	2.2	7	1.7	1.2
2002	41	11.5	3.8	38	9.6	3.1	12	4.4	2.8	8	2.1	1.5
2003	50	16.8	5.2	46	11.6	3.4	18	6.7	3.5	7	2.0	1.5
2004	52	16.3	4.7	42	9.9	3.0	11	4.5	2.9	6	1.5	1.3
2005	46	13.3	4.1	50	11.1	3.1	13	4.1	2.3	8	2.1	1.4
2006	53	15.1	4.4	49	11.2	3.2	9	3.3	2.5	12	2.9	1.6
2007	45	12.3	3.9	45	9.9	2.9	19	5.7	2.8	17	4.1	2.0
2008	63	17.2	4.6	41	8.7	2.7	9	2.5	1.7	7	1.5	1.1
2009	53	14.1	4.0	52	10.6	2.9	14	3.6	2.0	7	1.5	1.1
2010	57	15.3	4.3	47	9.2	2.7	12	3.1	2.0	11	2.2	1.3
2011	62	15.0	4.0	38	7.7	2.5	10	2.3	1.4	9	1.8	1.2
2012	55	12.4	3.5	48	8.8	2.5	9	3.4	2.3	4	0.7	0.7
2013	59	13.5	3.7	56	10.4	2.8	8	2.4	1.8	8	1.4	1.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Liver / intrahepatic bile duct

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	14	8.2	4.7	2	0.9	1.2	15	8.3	4.6	3	2.2	2.7
1977	9	4.9	3.3	4	2.8	2.9	12	6.2	3.6	0	0.0	0.0
1978	14	9.0	5.6	6	3.2	2.7	5	3.9	3.9	6	2.9	2.4
1979	8	4.5	3.3	5	2.8	2.6	7	3.4	2.6	3	2.3	2.7
1980	14	8.0	4.7	7	4.1	3.2	12	7.4	4.7	4	2.6	2.7
1981	14	9.0	5.2	10	4.4	2.8	14	9.2	5.4	11	5.9	3.7
1982	20	13.3	6.7	3	1.5	1.8	15	10.2	5.8	5	2.7	2.4
1983	17	10.1	5.4	2	0.9	1.3	19	11.8	6.0	5	2.3	2.0
1984	11	5.4	3.3	4	1.6	1.6	14	8.7	5.1	5	2.2	2.0
1985	5	2.2	2.0	10	4.0	2.5	3	1.9	2.3	6	2.8	2.3
1986	11	5.5	3.6	7	2.4	1.8	9	3.8	2.5	6	1.9	1.6
1987	9	3.7	2.5	8	3.2	2.3	12	5.7	3.6	8	3.6	2.5
1988	15	7.5	4.4	8	3.0	2.1	21	12.0	5.8	4	1.2	1.2
1989	23	11.4	5.2	5	2.3	2.0	12	7.1	4.6	9	3.6	2.4
1990	21	9.9	4.6	10	3.7	2.4	19	8.3	4.0	6	2.0	1.6
1991	24	9.5	4.0	9	2.9	2.0	19	6.9	3.2	8	2.4	1.7
1992	15	6.1	3.3	8	2.8	2.0	18	7.1	3.4	11	3.5	2.2
1993	12	4.9	2.9	15	5.4	2.8	8	2.7	2.0	14	5.0	2.7
1994	19	8.7	4.5	6	2.0	1.6	19	9.5	4.7	9	2.8	1.9
1995	30	13.0	5.2	11	3.6	2.2	23	10.8	4.8	7	2.1	1.6
1996	25	11.0	5.0	15	4.9	2.5	25	10.8	4.8	14	4.6	2.5
1997	33	14.2	5.3	18	5.3	2.5	20	8.5	4.1	13	3.9	2.2
1998	37	11.8	3.9	17	5.2	2.5	30	11.4	4.5	17	5.3	2.6
1999	21	6.6	2.9	9	2.6	1.7	14	5.2	2.9	11	3.4	2.0
2000	25	8.6	3.6	10	2.6	1.7	26	9.0	3.9	13	3.8	2.1
2001	30	8.8	3.3	14	3.5	1.9	21	6.7	3.2	16	4.2	2.1
2002	50	14.6	4.3	12	3.3	1.9	22	7.2	3.3	11	2.8	1.7
2003	42	11.1	3.5	20	5.4	2.4	35	9.4	3.2	7	1.8	1.4
2004	49	14.5	4.3	20	4.6	2.0	30	9.1	3.5	18	4.4	2.1
2005	62	18.4	4.9	21	5.2	2.2	40	10.6	3.5	13	3.0	1.6
2006	52	13.9	3.9	14	3.3	1.8	22	6.4	2.8	16	4.0	2.0
2007	72	18.2	4.5	27	5.9	2.3	35	10.0	3.5	18	4.2	2.0
2008	75	18.2	4.4	23	4.4	1.8	46	12.0	3.8	13	2.8	1.5
2009	91	20.9	4.5	21	4.1	1.8	64	15.9	4.2	9	1.9	1.3
2010	84	18.6	4.1	19	3.7	1.7	59	13.8	3.7	20	3.7	1.7
2011	81	16.3	3.7	17	3.3	1.6	60	13.0	3.4	17	3.2	1.5
2012	78	15.9	3.7	22	4.1	1.8	65	14.0	3.6	24	5.0	2.0
2013	68	14.1	3.5	35	6.3	2.1	61	14.1	3.8	14	2.6	1.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Pancreas

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	21	13.9	6.5	33	19.7	7.2	19	12.5	5.8	19	11.3	5.4
1977	33	19.9	7.2	23	13.6	5.9	34	21.4	7.8	23	13.4	5.8
1978	40	27.5	9.9	30	17.2	6.6	27	19.1	8.3	23	12.6	5.5
1979	41	33.1	12.0	29	16.9	6.6	31	20.9	8.7	26	15.1	6.2
1980	27	20.1	8.6	23	11.1	4.7	27	18.6	8.2	20	10.5	4.9
1981	28	21.2	8.8	31	15.9	5.9	26	18.1	8.3	28	14.7	5.7
1982	33	20.1	7.6	31	14.8	5.4	33	23.2	8.9	33	15.6	5.5
1983	41	26.4	9.0	33	15.4	5.5	36	23.7	8.7	29	13.7	5.2
1984	30	23.4	9.5	43	19.2	5.9	36	23.9	8.8	32	13.7	4.9
1985	33	21.1	8.1	34	15.7	5.4	28	20.8	8.7	29	13.0	4.9
1986	32	16.9	6.4	37	15.3	5.1	34	18.2	6.8	32	14.3	5.1
1987	21	14.2	7.1	46	18.5	5.5	26	16.4	7.3	41	16.9	5.3
1988	34	22.1	8.2	31	14.2	5.1	38	24.4	8.5	29	12.9	4.8
1989	25	15.4	6.8	35	13.7	4.7	24	15.2	6.8	30	12.6	4.6
1990	35	22.9	8.5	57	22.5	6.0	20	13.7	6.9	52	20.1	5.6
1991	42	21.9	7.6	34	12.7	4.4	29	16.1	6.6	27	10.9	4.2
1992	33	18.6	7.3	37	13.8	4.5	28	17.7	7.6	34	13.2	4.5
1993	22	11.4	5.3	29	10.7	4.0	26	13.9	6.0	30	10.6	3.9
1994	36	18.7	6.7	43	15.2	4.6	33	18.5	7.0	33	12.2	4.2
1995	34	14.8	5.3	41	13.5	4.2	33	16.6	6.3	34	11.2	3.8
1996	43	19.7	6.5	44	13.4	4.0	33	17.7	6.7	42	12.7	3.9
1997	32	14.7	5.6	47	14.5	4.2	31	14.3	5.4	32	10.2	3.6
1998	30	13.4	5.4	52	15.6	4.3	33	15.6	5.9	53	16.6	4.6
1999	40	14.7	4.8	47	13.9	4.0	37	13.6	4.8	53	15.5	4.2
2000	32	12.0	4.5	52	15.3	4.2	31	12.7	4.8	41	11.9	3.7
2001	49	18.3	5.6	67	18.9	4.6	38	15.2	5.2	46	13.3	3.9
2002	50	18.3	5.4	49	13.0	3.7	42	15.3	5.0	39	11.0	3.5
2003	50	19.0	5.8	43	11.3	3.4	40	16.4	5.6	47	12.4	3.6
2004	42	14.7	5.0	57	14.8	3.9	42	15.9	5.4	42	11.0	3.4
2005	59	19.2	5.2	67	16.6	4.0	44	15.0	4.9	40	10.0	3.1
2006	53	17.5	5.0	60	13.9	3.6	32	10.5	3.9	51	11.7	3.3
2007	40	12.9	4.3	64	14.8	3.7	34	10.7	3.8	42	9.7	3.0
2008	58	17.1	4.7	67	15.1	3.7	35	11.7	4.2	55	12.6	3.4
2009	69	18.7	4.7	55	12.4	3.3	50	14.2	4.2	62	13.7	3.5
2010	57	15.9	4.5	56	11.9	3.2	43	11.9	3.8	46	9.9	2.9
2011	69	17.5	4.5	78	16.0	3.6	43	12.4	4.0	56	11.8	3.1
2012	65	16.5	4.3	77	14.6	3.3	55	13.1	3.7	60	11.8	3.0
2013	63	15.5	4.1	64	12.1	3.0	46	12.7	3.9	51	9.6	2.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Larynx

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	32	18.6	6.6	4	1.9	1.8	13	10.1	6.2	3	1.4	1.6
1977	20	12.8	6.7	2	1.1	1.5	9	4.7	3.2	2	0.9	1.2
1978	20	11.4	5.3	5	2.2	1.9	16	10.4	6.2	3	1.3	1.4
1979	32	17.9	6.6	9	3.9	2.6	10	6.1	4.0	0	0.0	0.0
1980	34	18.8	6.6	4	1.6	1.6	11	6.0	3.7	3	1.2	1.4
1981	32	16.0	5.7	7	3.1	2.4	16	10.0	5.7	3	1.2	1.3
1982	34	17.3	6.2	2	0.8	1.1	16	9.2	5.2	3	1.4	1.6
1983	22	10.5	4.4	2	1.1	1.5	15	7.7	4.0	3	1.2	1.3
1984	27	14.1	5.8	9	3.5	2.3	10	6.3	4.5	1	0.6	1.1
1985	30	14.0	5.2	12	4.7	2.7	13	8.0	4.7	2	0.8	1.1
1986	32	16.2	6.0	7	2.5	1.9	11	5.8	3.6	5	1.9	1.7
1987	24	11.1	4.6	5	1.8	1.5	7	3.8	3.0	5	1.9	1.7
1988	16	7.6	4.0	7	2.7	2.0	17	8.2	4.1	0	0.0	0.0
1989	26	11.9	5.1	13	4.2	2.3	10	4.8	3.0	1	0.4	0.7
1990	27	11.6	4.7	5	1.7	1.5	14	6.8	4.1	2	0.7	1.0
1991	24	10.6	4.5	4	1.4	1.4	17	8.0	4.2	0	0.0	0.0
1992	28	12.1	4.8	11	3.5	2.1	12	5.3	3.3	5	1.5	1.3
1993	33	13.8	5.0	5	1.5	1.3	14	5.3	2.8	2	0.7	1.0
1994	36	14.5	5.2	3	0.9	1.0	14	6.8	4.0	4	1.3	1.3
1995	30	12.4	4.7	7	2.1	1.5	15	7.4	4.0	3	0.9	1.0
1996	35	14.3	5.2	6	1.7	1.4	15	6.8	4.0	2	0.6	0.8
1997	33	12.7	4.7	9	2.4	1.6	9	3.2	2.2	3	0.8	0.9
1998	35	12.7	4.6	14	3.8	2.0	14	5.3	3.0	1	0.3	0.6
1999	29	9.7	3.8	8	2.0	1.4	19	7.2	3.5	1	0.2	0.5
2000	23	7.4	3.2	10	2.4	1.5	7	3.3	2.7	6	1.5	1.2
2001	31	9.4	3.4	6	1.5	1.2	9	3.5	2.6	1	0.2	0.4
2002	28	8.3	3.2	9	2.0	1.4	18	5.4	2.5	6	1.6	1.3
2003	32	10.3	3.8	10	2.5	1.6	14	4.5	2.5	6	1.4	1.2
2004	36	10.5	3.6	4	0.9	0.9	12	3.5	2.0	2	0.5	0.7
2005	26	8.8	3.7	5	1.1	1.0	13	3.9	2.3	1	0.2	0.4
2006	39	11.2	3.7	9	1.9	1.3	10	3.4	2.2	1	0.3	0.5
2007	31	9.5	3.6	15	3.2	1.6	7	2.1	1.7	4	0.9	0.9
2008	37	10.6	3.7	10	2.2	1.4	15	4.0	2.1	5	1.1	1.0
2009	24	6.7	2.9	10	2.1	1.3	13	3.9	2.3	5	1.0	0.9
2010	32	9.0	3.3	10	2.0	1.3	14	4.3	2.4	2	0.4	0.6
2011	36	8.5	2.9	6	1.1	0.9	9	2.0	1.4	2	0.4	0.6
2012	21	5.5	2.5	10	1.8	1.1	19	5.4	2.6	2	0.4	0.5
2013	21	5.3	2.4	10	1.6	1.0	14	4.1	2.3	1	0.2	0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Lung and bronchus

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	174	111.8	17.8	51	25.1	7.1	122	80.3	15.8	34	16.6	5.7
1977	219	149.3	22.2	65	32.0	8.2	165	112.9	19.1	40	19.8	6.4
1978	203	128.7	19.4	67	33.7	8.5	169	111.8	18.9	47	23.6	7.1
1979	200	134.3	21.2	78	35.5	8.2	166	112.4	19.2	51	25.1	7.3
1980	253	159.8	21.9	80	35.4	8.0	193	129.2	20.7	69	31.6	7.7
1981	250	155.0	21.2	82	36.3	8.1	190	123.5	19.8	68	29.9	7.3
1982	243	144.5	20.2	82	34.5	7.7	197	115.5	17.7	68	29.5	7.2
1983	241	138.2	18.8	96	39.7	8.2	225	137.7	19.7	68	29.2	7.2
1984	228	124.2	17.4	97	41.2	8.5	197	110.4	16.8	76	31.1	7.2
1985	250	133.2	17.8	102	40.3	8.0	198	107.2	16.1	76	31.1	7.2
1986	264	149.7	20.2	109	42.6	8.2	199	112.5	17.2	94	36.7	7.6
1987	233	119.4	16.4	103	39.1	7.7	234	126.7	17.9	91	34.6	7.3
1988	275	149.3	19.4	121	45.0	8.2	204	110.0	16.4	88	32.8	7.0
1989	287	145.5	18.4	123	44.1	7.9	258	134.4	17.9	120	43.1	7.9
1990	274	141.4	18.4	110	40.0	7.6	224	115.3	16.6	107	39.2	7.6
1991	263	133.5	17.9	147	50.8	8.4	221	114.1	16.5	125	43.3	7.8
1992	296	146.1	18.6	125	41.0	7.3	227	116.6	16.9	99	33.4	6.7
1993	266	128.8	17.2	138	44.9	7.6	252	124.5	17.1	110	36.3	6.9
1994	253	117.8	16.0	184	56.3	8.3	260	118.9	15.9	109	34.4	6.5
1995	279	121.2	15.4	158	50.0	7.9	212	97.8	14.3	140	43.3	7.3
1996	276	118.2	15.2	178	53.6	8.0	215	91.1	13.3	144	44.7	7.4
1997	294	121.2	15.0	171	50.2	7.6	236	102.0	14.1	135	39.5	6.7
1998	273	108.1	14.0	195	55.8	7.9	239	99.5	13.6	136	39.4	6.7
1999	265	105.8	13.8	195	55.3	7.9	204	82.8	12.5	132	38.6	6.7
2000	305	112.9	13.7	175	47.9	7.2	231	89.7	12.5	143	38.6	6.4
2001	301	112.0	13.6	191	49.9	7.2	211	82.1	12.0	142	38.4	6.4
2002	273	101.3	13.0	192	49.5	7.1	233	86.1	12.0	138	35.8	6.1
2003	270	92.4	11.9	204	50.5	7.0	226	80.8	11.4	147	37.3	6.1
2004	299	97.0	11.8	207	49.3	6.8	224	81.9	11.6	148	36.3	5.9
2005	296	99.5	12.4	203	48.1	6.7	219	73.0	10.6	165	39.5	6.1
2006	288	93.5	11.7	207	49.4	6.8	186	61.1	9.5	138	32.9	5.6
2007	303	94.1	11.4	221	49.1	6.6	209	65.0	9.5	160	36.1	5.7
2008	273	83.0	10.6	233	51.1	6.7	182	56.4	8.8	144	32.0	5.3
2009	305	88.1	10.7	249	52.7	6.7	203	60.7	9.0	153	32.9	5.3
2010	297	83.3	10.2	266	54.7	6.7	223	68.3	9.6	155	32.6	5.2
2011	297	79.6	9.7	247	48.7	6.2	228	63.8	8.9	173	35.0	5.3
2012	334	86.8	9.9	238	45.4	5.9	215	59.1	8.5	164	31.8	5.0
2013	290	72.5	8.9	256	45.3	5.6	211	55.4	8.0	165	31.0	4.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Melanoma of the skin

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	1	0.9	1.7	1	0.4	0.8	1	0.5	1.0	3	2.4	2.9
1977	5	5.9	5.9	1	1.1	2.2	2	3.3	5.1	1	0.4	0.8
1978	2	2.8	4.6	2	0.8	1.2	0	0.0	0.0	1	0.7	1.4
1979	1	0.5	0.9	0	0.0	0.0	1	0.5	1.0	1	1.0	1.9
1980	2	1.0	1.4	2	1.5	2.1	0	0.0	0.0	0	0.0	0.0
1981	2	0.7	1.1	5	1.9	1.8	1	1.0	2.0	0	0.0	0.0
1982	6	2.6	2.1	3	0.9	1.1	2	1.2	1.7	1	0.4	0.8
1983	2	1.4	2.1	0	0.0	0.0	1	0.4	0.8	1	0.6	1.1
1984	2	1.7	2.7	3	1.4	1.7	1	0.2	0.4	0	0.0	0.0
1985	3	1.4	1.6	3	1.6	1.9	1	1.2	2.4	1	0.5	0.9
1986	3	1.1	1.3	3	1.2	1.4	5	2.1	1.9	2	0.7	0.9
1987	3	1.7	2.1	2	0.8	1.2	1	0.5	0.9	0	0.0	0.0
1988	2	1.1	1.6	3	1.0	1.1	2	1.0	1.5	0	0.0	0.0
1989	2	0.6	0.8	2	1.0	1.4	2	1.1	1.7	3	1.1	1.3
1990	1	0.4	0.8	3	1.0	1.2	0	0.0	0.0	0	0.0	0.0
1991	1	1.4	2.7	4	1.1	1.1	0	0.0	0.0	2	0.8	1.2
1992	2	0.7	0.9	1	0.3	0.6	1	0.3	0.6	0	0.0	0.0
1993	1	1.2	2.4	7	2.3	1.7	0	0.0	0.0	3	0.8	0.9
1994	2	0.5	0.8	3	1.0	1.2	0	0.0	0.0	3	1.0	1.1
1995	3	0.8	1.0	3	0.9	1.0	1	0.2	0.3	3	0.7	0.8
1996	3	1.6	2.3	2	0.5	0.8	1	0.3	0.6	3	0.8	0.9
1997	1	0.3	0.6	5	1.5	1.4	1	0.2	0.4	1	0.4	0.7
1998	3	0.9	1.1	5	1.5	1.3	1	0.2	0.3	2	0.6	0.8
1999	2	0.5	0.7	2	0.5	0.7	1	0.3	0.7	0	0.0	0.0
2000	0	0.0	0.0	0	0.0	0.0	2	1.1	1.6	1	0.3	0.7
2001	2	0.5	0.6	4	0.9	0.9	2	0.6	0.9	2	0.6	0.9
2002	4	1.5	1.5	4	1.1	1.1	1	0.9	1.8	1	0.3	0.6
2003	2	0.9	1.3	6	1.3	1.1	2	0.9	1.3	3	0.6	0.7
2004	5	1.3	1.2	4	0.9	0.8	1	0.2	0.4	0	0.0	0.0
2005	5	1.1	0.9	3	0.7	0.8	1	0.2	0.4	1	0.2	0.4
2006	6	1.7	1.5	4	1.1	1.0	2	0.7	1.0	2	0.5	0.7
2007	1	0.2	0.4	9	2.0	1.3	0	0.0	0.0	1	0.2	0.4
2008	4	0.9	0.9	8	1.8	1.3	0	0.0	0.0	2	0.5	0.7
2009	2	0.4	0.5	6	1.1	0.9	2	0.6	0.9	2	0.4	0.6
2010	3	0.7	0.8	7	1.2	0.9	1	0.2	0.4	4	0.8	0.8
2011	4	1.1	1.1	5	1.0	0.9	1	0.2	0.5	3	0.7	0.8
2012	4	1.1	1.2	3	0.5	0.5	0	0.0	0.0	4	0.8	0.8
2013	5	1.5	1.5	10	1.8	1.1	3	0.6	0.7	3	0.5	0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Female breast

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	151	79.1	13.5	67	37.3	9.6
1977	169	87.6	14.0	65	34.8	9.1
1978	184	90.2	13.7	55	26.1	7.2
1979	173	83.4	13.2	65	30.7	7.9
1980	204	91.0	13.0	70	32.5	8.0
1981	205	90.0	12.8	84	39.6	8.9
1982	202	84.9	12.1	90	38.1	8.2
1983	221	91.8	12.5	77	31.5	7.2
1984	221	89.1	12.2	97	43.2	8.9
1985	247	97.2	12.5	91	36.4	7.7
1986	255	96.2	12.1	89	34.6	7.4
1987	249	93.9	12.0	84	32.6	7.2
1988	278	98.8	11.8	106	38.3	7.4
1989	248	87.0	11.1	106	39.1	7.6
1990	301	101.6	11.7	90	30.8	6.5
1991	314	102.6	11.6	113	37.9	7.1
1992	327	103.9	11.5	114	36.6	6.8
1993	338	105.1	11.4	109	34.6	6.6
1994	347	102.9	11.0	116	36.2	6.7
1995	345	100.4	10.8	108	33.0	6.3
1996	394	110.1	11.1	108	31.5	6.0
1997	429	118.6	11.4	115	34.0	6.3
1998	443	116.5	11.0	122	33.7	6.1
1999	445	115.3	10.9	98	25.9	5.2
2000	396	99.6	10.0	127	34.2	6.1
2001	414	101.6	9.9	121	30.3	5.5
2002	460	107.4	9.9	123	29.8	5.3
2003	461	106.7	9.9	119	28.1	5.1
2004	508	114.5	10.1	128	30.5	5.4
2005	510	113.9	10.0	126	28.7	5.1
2006	547	117.3	10.0	121	26.7	4.8
2007	553	116.3	9.8	135	29.2	5.0
2008	577	118.0	9.7	133	27.3	4.7
2009	588	119.0	9.8	151	31.5	5.1
2010	590	115.2	9.5	126	24.7	4.4
2011	676	127.8	9.8	148	28.7	4.7
2012	615	114.1	9.1	121	23.2	4.2
2013	641	116.9	9.2	140	25.8	4.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Cervix uteri

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	69	35.4	9.0	29	16.5	6.6
1977	58	27.2	7.5	27	14.6	6.0
1978	56	25.3	7.0	21	10.3	4.7
1979	61	25.8	6.8	24	10.7	4.4
1980	58	26.1	7.1	25	12.2	5.1
1981	37	15.3	5.1	27	13.0	5.1
1982	44	16.8	5.1	16	7.0	3.6
1983	49	18.8	5.4	22	8.6	3.6
1984	52	19.3	5.5	13	6.9	3.9
1985	41	15.8	5.0	23	8.9	3.8
1986	53	18.8	5.3	20	8.0	3.6
1987	49	17.5	5.1	16	6.0	3.1
1988	46	15.7	4.7	18	6.6	3.2
1989	50	17.2	4.9	19	6.9	3.2
1990	63	19.0	4.8	18	6.2	3.0
1991	53	16.2	4.5	19	6.2	2.8
1992	42	11.4	3.5	14	4.2	2.2
1993	52	14.3	4.0	25	7.4	3.0
1994	47	12.6	3.7	15	4.2	2.1
1995	53	14.2	4.0	16	4.7	2.4
1996	73	19.3	4.5	19	5.2	2.4
1997	34	8.9	3.0	24	6.4	2.6
1998	51	12.0	3.4	17	4.6	2.2
1999	52	13.4	3.7	13	3.3	1.8
2000	42	10.2	3.1	19	4.6	2.1
2001	52	12.7	3.5	10	2.5	1.6
2002	45	10.4	3.1	16	4.0	2.0
2003	56	12.8	3.4	19	4.4	2.0
2004	49	10.6	3.0	14	3.0	1.6
2005	50	10.9	3.1	17	3.6	1.7
2006	47	9.8	2.8	15	3.3	1.7
2007	48	10.1	2.9	21	4.6	2.0
2008	52	10.3	2.8	15	3.1	1.6
2009	42	8.5	2.6	26	5.0	2.0
2010	44	8.6	2.6	13	2.5	1.4
2011	45	8.9	2.6	14	2.6	1.4
2012	43	7.9	2.4	19	3.5	1.6
2013	61	11.0	2.8	23	4.2	1.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Corpus uterus and uterus, not otherwise specified

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	31	17.9	6.7	13	9.6	5.6
1977	36	19.4	6.8	15	8.2	4.5
1978	26	14.3	5.9	9	4.6	3.2
1979	37	16.3	5.4	14	7.6	4.2
1980	50	23.8	6.9	21	9.8	4.3
1981	28	12.2	4.7	21	9.9	4.5
1982	32	13.8	5.0	11	4.2	2.5
1983	31	12.9	4.7	13	6.0	3.4
1984	33	14.2	5.1	16	6.7	3.4
1985	49	19.8	5.7	19	9.5	4.4
1986	38	15.4	5.0	12	4.9	2.8
1987	49	19.1	5.5	11	4.7	2.9
1988	46	17.9	5.3	18	7.2	3.4
1989	35	12.6	4.3	24	9.4	3.8
1990	51	17.8	5.0	10	4.0	2.6
1991	42	14.5	4.5	17	5.8	2.8
1992	64	21.9	5.5	23	8.7	3.6
1993	60	19.4	5.0	16	6.0	3.0
1994	66	20.3	5.0	13	4.2	2.3
1995	52	15.8	4.4	18	5.9	2.8
1996	79	23.2	5.2	24	7.4	3.0
1997	70	20.8	4.9	12	3.7	2.1
1998	79	21.5	4.8	28	8.5	3.2
1999	84	22.7	4.9	24	7.0	2.8
2000	81	21.8	4.8	30	8.5	3.1
2001	80	20.8	4.6	25	7.0	2.8
2002	75	18.2	4.2	29	7.8	2.9
2003	83	19.9	4.3	36	9.1	3.0
2004	110	25.6	4.9	20	4.4	2.0
2005	102	23.8	4.7	33	8.3	2.9
2006	102	22.1	4.3	32	7.3	2.6
2007	126	26.8	4.7	41	9.7	3.0
2008	114	24.1	4.5	46	10.2	3.0
2009	135	28.2	4.8	30	6.7	2.4
2010	131	26.1	4.6	46	9.6	2.8
2011	134	24.8	4.3	36	7.3	2.4
2012	156	28.9	4.6	47	9.5	2.8
2013	132	23.0	4.0	45	8.4	2.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Ovary

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	28	16.5	6.6	17	11.7	6.0
1977	29	13.2	5.2	17	8.4	4.2
1978	24	10.7	4.4	13	6.9	4.0
1979	34	16.4	5.9	22	10.1	4.5
1980	26	12.0	4.8	20	9.5	4.3
1981	28	12.3	4.8	14	6.2	3.4
1982	36	14.3	4.8	16	6.7	3.4
1983	29	12.2	4.6	14	6.1	3.3
1984	28	10.8	4.1	24	10.5	4.4
1985	35	14.3	4.9	13	5.4	3.0
1986	37	14.3	4.8	21	8.5	3.7
1987	28	11.1	4.3	27	10.9	4.3
1988	38	13.3	4.3	20	7.5	3.4
1989	38	13.2	4.3	23	8.3	3.5
1990	37	12.2	4.1	28	10.2	3.9
1991	23	8.8	3.7	20	7.2	3.2
1992	31	9.7	3.5	34	11.9	4.1
1993	42	12.3	3.8	19	6.5	3.0
1994	32	9.7	3.4	23	7.4	3.1
1995	42	11.8	3.7	23	7.9	3.3
1996	40	11.8	3.7	19	5.7	2.6
1997	52	14.5	4.0	26	7.6	3.0
1998	37	9.9	3.3	23	7.0	2.9
1999	47	12.7	3.7	26	7.2	2.8
2000	42	11.1	3.4	31	8.5	3.0
2001	41	10.1	3.1	24	6.6	2.7
2002	44	10.4	3.2	29	7.5	2.8
2003	28	6.4	2.4	35	8.5	2.9
2004	46	10.9	3.2	28	7.1	2.7
2005	39	8.5	2.7	26	6.0	2.3
2006	40	9.1	2.9	31	7.3	2.6
2007	35	7.4	2.5	26	6.1	2.4
2008	48	10.4	3.0	21	4.9	2.1
2009	51	10.2	2.8	25	5.3	2.1
2010	41	8.0	2.5	26	5.3	2.1
2011	49	9.2	2.6	30	6.3	2.3
2012	58	10.6	2.8	34	6.3	2.2
2013	40	6.9	2.2	34	6.3	2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Prostate

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	123	126.0	25.8	65	71.9	20.2
1977	128	114.2	22.6	64	79.0	21.7
1978	143	128.1	24.3	59	54.2	15.6
1979	144	126.1	23.7	73	67.8	17.4
1980	173	139.5	23.4	56	57.7	16.8
1981	169	142.7	24.3	82	76.0	18.5
1982	155	120.1	20.7	74	64.2	16.2
1983	185	143.2	22.9	73	65.3	16.4
1984	171	123.2	20.4	67	51.9	13.8
1985	181	127.9	20.7	72	58.0	14.6
1986	206	146.6	22.1	92	74.5	16.9
1987	186	129.3	20.3	83	75.3	17.8
1988	210	138.3	20.7	100	83.0	17.7
1989	210	133.0	19.9	82	59.1	13.8
1990	256	160.7	21.8	96	67.9	15.0
1991	357	209.5	24.1	104	82.1	17.1
1992	431	236.9	24.9	87	60.9	14.0
1993	514	270.5	25.8	102	72.5	15.4
1994	505	249.0	23.8	121	83.3	16.2
1995	558	266.3	24.3	113	70.2	14.2
1996	621	266.3	22.7	116	70.6	13.9
1997	616	265.5	22.9	108	63.4	13.1
1998	666	272.0	22.2	114	70.9	13.9
1999	703	274.3	21.7	94	54.6	11.9
2000	766	302.8	23.2	129	74.3	13.7
2001	779	290.3	22.1	110	63.3	12.7
2002	817	296.8	22.2	114	61.5	12.2
2003	812	278.9	20.6	112	56.5	11.2
2004	756	242.9	18.5	97	47.0	10.2
2005	802	250.9	18.6	107	45.2	9.1
2006	877	266.0	18.8	83	35.1	8.0
2007	952	281.6	19.0	99	43.3	9.0
2008	920	256.5	17.6	103	40.2	8.2
2009	936	246.4	16.8	111	44.8	8.8
2010	964	248.3	16.8	113	44.3	8.6
2011	1004	245.8	16.1	88	30.8	6.8
2012	883	203.5	14.2	97	35.2	7.3
2013	912	204.7	14.0	97	33.9	7.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Testis

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	3	0.9	1.0	1	0.9	1.7
1977	5	1.5	1.3	1	0.4	0.8
1978	2	0.5	0.7	1	0.3	0.6
1980	7	2.1	1.6	2	0.6	0.9
1981	4	1.6	1.6	0	0.0	0.0
1982	3	1.2	1.4	0	0.0	0.0
1983	2	0.4	0.6	1	0.2	0.4
1984	5	1.5	1.4	0	0.0	0.0
1985	6	1.9	1.6	1	0.4	0.8
1986	6	1.8	1.5	0	0.0	0.0
1987	5	1.2	1.1	1	0.2	0.3
1988	5	1.2	1.1	0	0.0	0.0
1989	2	0.5	0.8	0	0.0	0.0
1990	3	0.7	0.8	0	0.0	0.0
1991	3	0.6	0.8	1	0.4	0.7
1992	8	2.6	2.7	1	0.3	0.5
1993	3	0.6	0.7	1	0.2	0.4
1994	4	0.8	0.8	0	0.0	0.0
1995	2	0.5	0.7	0	0.0	0.0
1996	4	0.9	0.9	0	0.0	0.0
1997	6	1.2	0.9	0	0.0	0.0
1998	6	1.2	1.0	0	0.0	0.0
1999	8	1.5	1.0	1	0.2	0.5
2000	5	1.5	1.5	0	0.0	0.0
2001	8	1.7	1.2	0	0.0	0.0
2002	6	1.2	1.0	0	0.0	0.0
2003	9	1.8	1.2	0	0.0	0.0
2004	5	1.0	0.9	0	0.0	0.0
2005	2	0.4	0.5	0	0.0	0.0
2006	6	1.2	1.0	0	0.0	0.0
2007	9	1.7	1.1	0	0.0	0.0
2008	6	1.2	1.0	1	0.1	0.3
2009	3	0.5	0.6	0	0.0	0.0
2010	4	0.8	0.8	1	0.2	0.4
2011	7	1.3	1.0	0	0.0	0.0
2012	6	1.2	1.0	0	0.0	0.0
2013	10	1.8	1.1	1	0.2	0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Urinary bladder (incl. in situ)

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	31	34.4	14.6	13	8.8	5.1	7	5.8	5.7	8	5.5	4.0
1977	13	10.4	6.3	17	11.1	5.7	8	11.1	8.6	6	4.2	3.7
1978	30	28.1	11.6	12	6.4	3.8	8	4.6	3.2	10	5.6	3.6
1979	30	23.0	9.0	25	14.6	6.1	14	17.3	10.1	9	5.9	4.0
1980	29	21.1	8.8	11	6.8	4.2	12	10.2	6.4	8	4.1	3.0
1981	23	16.2	7.4	19	10.2	4.8	13	11.7	7.6	3	1.3	1.5
1982	21	15.8	7.6	12	6.4	3.8	12	9.6	6.1	7	3.8	3.0
1983	18	11.1	5.7	17	8.5	4.2	7	5.3	4.1	11	5.4	3.3
1984	38	28.1	10.1	17	7.9	3.9	14	13.3	7.7	6	3.4	2.8
1985	27	18.2	7.5	18	8.1	3.9	10	8.4	5.7	6	2.8	2.3
1986	25	16.3	6.9	15	6.9	3.6	8	5.7	4.3	8	4.0	2.8
1987	20	12.6	6.1	14	6.6	3.6	6	4.9	4.1	7	3.3	2.5
1988	33	19.4	7.0	20	9.5	4.3	9	7.5	5.3	7	3.5	2.7
1989	34	19.2	7.2	20	7.6	3.4	15	13.0	7.1	7	3.4	2.5
1990	30	18.9	7.6	11	4.9	3.0	16	9.7	5.1	8	4.0	2.8
1991	47	27.8	8.9	17	6.0	2.9	12	7.5	4.9	12	5.1	2.9
1992	30	16.2	6.3	9	3.3	2.2	9	4.8	3.4	3	1.1	1.3
1993	24	10.2	4.5	25	9.2	3.7	10	7.0	4.5	11	4.0	2.4
1994	32	19.0	7.3	13	4.5	2.5	9	6.0	4.2	6	2.5	2.0
1995	33	16.3	6.0	14	4.6	2.5	8	6.8	5.1	6	2.3	1.8
1996	30	14.8	6.0	30	9.7	3.5	13	7.7	4.5	10	3.3	2.1
1997	48	26.6	8.5	17	5.6	2.7	9	4.5	3.3	7	2.3	1.7
1998	58	29.7	8.6	29	9.6	3.5	10	5.7	3.8	10	3.3	2.1
1999	41	19.2	6.5	23	6.7	2.8	12	5.9	3.7	8	2.6	1.8
2000	56	21.9	6.3	22	6.5	2.8	7	4.6	3.7	8	2.4	1.7
2001	51	21.7	6.7	26	7.5	2.9	10	5.3	3.8	9	2.8	1.8
2002	43	17.0	5.5	23	6.4	2.6	10	5.2	3.6	9	2.6	1.7
2003	55	20.2	5.9	28	7.2	2.7	13	6.3	3.7	11	3.0	1.8
2004	60	21.5	5.8	28	7.6	2.8	11	4.2	2.7	10	2.8	1.8
2005	60	22.5	6.3	23	5.7	2.4	13	7.1	4.0	10	2.8	1.7
2006	55	19.5	5.5	27	6.1	2.4	8	2.8	2.2	4	1.1	1.1
2007	57	18.8	5.1	26	6.2	2.4	15	5.7	3.0	16	3.9	1.9
2008	68	22.3	5.8	33	7.5	2.6	12	4.9	2.8	6	1.4	1.1
2009	67	21.1	5.4	33	7.5	2.6	20	7.6	3.6	7	1.7	1.3
2010	68	21.7	5.6	24	4.8	1.9	9	3.6	2.4	14	2.9	1.6
2011	100	28.9	6.1	31	6.0	2.2	19	6.6	3.2	8	1.9	1.3
2012	77	23.5	5.6	37	7.7	2.5	15	4.5	2.5	8	1.6	1.1
2013	76	21.0	5.0	39	7.5	2.4	19	5.8	2.8	11	2.2	1.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Kidney and renal pelvis

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	20	12.7	6.2	11	7.3	4.5	6	3.4	2.8	4	2.2	2.4
1977	14	11.7	7.4	13	5.8	3.3	4	3.3	3.7	3	1.1	1.4
1978	17	10.7	5.9	13	8.0	4.8	8	3.5	2.5	7	4.7	3.7
1979	21	11.1	4.9	9	4.6	3.2	13	10.0	6.2	2	0.8	1.1
1980	10	6.5	4.9	11	4.9	3.1	7	4.8	4.5	3	1.3	1.5
1981	22	11.6	5.8	8	3.7	2.6	8	3.8	2.7	4	1.5	1.5
1982	18	10.0	5.3	13	6.1	3.5	9	4.5	3.2	10	5.8	3.7
1983	19	9.9	5.1	7	3.2	2.5	7	4.3	3.9	3	1.3	1.5
1984	28	14.8	6.6	16	6.7	3.4	7	4.7	3.8	5	2.0	1.8
1985	25	13.9	6.2	21	8.1	3.6	11	4.8	2.9	5	2.6	2.3
1986	20	10.3	4.7	19	7.3	3.4	11	6.1	4.2	9	3.5	2.3
1987	23	14.4	6.4	13	4.9	2.8	5	3.1	3.0	12	5.1	3.0
1988	18	9.9	5.1	22	7.7	3.3	9	6.4	4.6	8	2.9	2.1
1989	19	9.5	5.0	14	5.0	2.7	6	4.0	3.5	6	2.4	2.0
1990	37	15.8	5.5	25	8.3	3.3	9	4.1	3.0	7	2.5	1.9
1991	38	15.8	5.5	21	6.6	2.9	12	7.2	4.6	8	3.0	2.1
1992	40	16.9	6.0	20	6.3	2.8	12	5.4	3.4	7	2.0	1.6
1993	42	16.7	5.5	23	6.8	2.8	10	4.9	3.3	12	3.9	2.2
1994	36	12.5	4.4	37	11.1	3.7	13	5.9	3.6	7	2.1	1.6
1995	38	17.4	6.4	39	11.9	3.8	14	7.6	4.7	9	3.0	2.0
1996	48	18.4	5.9	23	6.4	2.7	10	5.1	3.7	12	3.4	1.9
1997	46	18.0	5.7	27	7.7	3.0	14	5.6	3.3	13	4.2	2.3
1998	63	22.8	6.1	39	10.3	3.3	7	2.7	2.1	7	2.0	1.5
1999	38	13.1	4.4	30	8.2	3.0	10	3.7	2.6	10	3.2	2.0
2000	40	13.3	4.5	34	8.4	2.9	12	4.2	2.6	10	2.6	1.6
2001	52	15.6	4.6	31	8.1	2.9	13	4.0	2.3	10	2.9	1.8
2002	60	19.1	5.2	42	10.0	3.1	7	2.2	1.7	3	0.6	0.7
2003	52	15.5	4.5	40	10.3	3.3	20	7.9	4.0	6	1.6	1.3
2004	68	17.7	4.4	48	10.9	3.1	11	3.8	2.5	10	2.5	1.6
2005	58	16.4	4.5	48	10.6	3.0	21	6.2	2.9	7	1.7	1.2
2006	91	25.3	5.8	42	9.4	2.9	12	3.7	2.4	9	1.9	1.3
2007	82	22.0	5.2	64	13.8	3.4	23	7.3	3.2	9	2.1	1.4
2008	85	20.8	4.7	63	13.6	3.4	19	4.7	2.2	9	2.0	1.3
2009	103	24.7	5.1	73	15.4	3.6	20	5.0	2.3	13	2.8	1.5
2010	107	25.7	5.2	65	13.3	3.3	11	3.0	1.9	14	3.2	1.7
2011	108	27.6	5.6	54	10.6	2.9	14	3.8	2.2	10	2.1	1.3
2012	97	23.2	4.9	60	10.9	2.8	19	6.2	3.0	10	1.8	1.1
2013	119	25.6	4.8	57	10.5	2.8	11	2.6	1.6	6	1.1	0.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Brain and other nervous system

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	10	4.9	3.2	10	4.0	2.8	8	2.9	2.2	8	3.5	2.7
1977	10	5.2	3.4	16	5.5	2.8	4	1.9	1.9	6	2.3	1.9
1978	6	3.4	2.8	13	4.7	2.8	5	2.3	2.1	5	2.2	2.0
1979	16	7.5	3.8	8	2.6	1.9	6	2.5	2.1	6	2.3	1.9
1980	16	6.8	3.7	3	1.3	1.5	8	3.4	2.4	1	0.4	0.8
1981	7	2.6	2.0	7	2.1	1.6	5	2.4	2.1	1	0.2	0.4
1982	10	3.7	2.4	3	0.9	1.1	5	3.3	3.3	4	1.3	1.3
1983	7	2.2	1.8	8	2.4	1.7	5	2.4	2.2	6	2.1	1.9
1984	14	4.5	2.5	8	2.3	1.6	11	3.8	2.4	7	1.9	1.5
1985	9	4.4	3.4	12	3.6	2.1	5	1.8	1.6	6	1.8	1.5
1986	8	4.2	3.6	11	4.0	2.6	3	1.3	1.4	6	2.5	2.1
1987	7	2.3	1.8	11	4.1	2.6	3	1.2	1.3	6	2.7	2.2
1988	14	4.5	2.5	10	3.0	2.0	7	2.6	2.1	4	0.9	0.9
1989	5	1.2	1.1	13	3.7	2.1	4	1.0	1.1	4	1.3	1.3
1990	10	4.7	3.6	11	2.7	1.7	3	1.2	1.5	6	1.7	1.4
1991	19	5.7	2.9	18	5.1	2.4	3	1.4	1.8	8	2.4	1.8
1992	19	6.0	2.9	19	5.6	2.6	5	2.1	2.1	11	3.1	1.9
1993	11	2.7	1.7	13	3.5	2.0	11	3.7	2.3	13	3.7	2.1
1994	9	2.4	1.9	11	3.3	2.0	4	1.0	1.0	4	1.1	1.1
1995	15	4.9	2.7	16	4.5	2.3	9	3.2	2.3	7	2.1	1.6
1996	23	6.9	3.0	16	4.0	2.0	12	4.2	2.7	6	1.8	1.5
1997	22	6.4	3.1	9	2.2	1.5	7	1.9	1.5	4	1.1	1.1
1998	14	3.3	1.8	7	1.6	1.2	10	3.0	1.9	3	0.8	0.9
1999	24	5.8	2.4	11	2.7	1.6	8	2.4	1.8	5	1.1	1.0
2000	11	2.3	1.4	16	3.6	1.8	8	2.3	1.6	5	1.2	1.1
2001	22	5.8	2.7	30	7.4	2.7	9	3.6	2.6	13	3.2	1.8
2002	18	4.6	2.3	24	5.0	2.0	10	2.0	1.3	10	2.4	1.5
2003	18	4.5	2.2	20	4.3	2.0	12	3.1	1.8	9	2.0	1.4
2004	17	3.9	2.0	22	4.4	1.9	4	1.3	1.4	9	1.9	1.3
2005	23	6.4	2.8	10	2.0	1.3	8	1.9	1.3	5	1.1	1.0
2006	15	3.1	1.8	16	3.2	1.6	11	2.7	1.7	3	0.7	0.8
2007	13	2.9	1.7	14	2.9	1.6	7	2.6	2.0	9	2.0	1.3
2008	16	3.3	1.7	23	4.7	2.0	9	1.7	1.1	10	2.1	1.3
2009	22	5.3	2.5	17	3.5	1.7	8	2.6	1.9	9	2.0	1.3
2010	30	6.6	2.5	21	4.1	1.8	12	2.7	1.7	9	1.7	1.1
2011	23	4.2	1.7	21	3.9	1.7	15	3.2	1.7	8	1.5	1.1
2012	25	4.8	1.9	22	3.8	1.6	13	2.5	1.4	12	2.4	1.4
2013	20	3.6	1.6	23	4.0	1.7	15	3.1	1.6	11	1.9	1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Thyroid

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	5	2.9	3.0	10	3.9	2.5	1	0.5	1.0	1	0.8	1.6
1977	2	0.9	1.3	10	4.7	3.2	1	0.5	1.0	1	1.0	2.0
1978	3	1.3	1.6	8	4.1	3.0	2	1.7	2.6	2	1.5	2.2
1979	2	0.5	0.7	9	4.0	2.9	0	0.0	0.0	3	2.3	2.7
1980	2	0.6	0.8	9	3.4	2.5	0	0.0	0.0	1	0.4	0.8
1981	2	0.7	1.0	5	1.4	1.3	1	0.5	0.9	1	0.4	0.8
1982	1	0.4	0.9	10	3.5	2.2	1	0.4	0.8	1	0.4	0.8
1983	2	0.8	1.1	14	4.7	2.6	0	0.0	0.0	0	0.0	0.0
1984	2	0.8	1.2	8	2.3	1.7	0	0.0	0.0	0	0.0	0.0
1985	3	1.2	1.4	16	5.7	2.9	1	0.5	0.9	0	0.0	0.0
1986	3	1.3	1.5	6	2.1	1.7	1	0.5	0.9	2	0.8	1.1
1987	3	1.4	1.7	9	2.9	1.9	1	0.4	0.8	3	1.1	1.2
1988	1	0.4	0.8	7	2.2	1.7	2	1.1	1.5	1	0.3	0.6
1989	1	0.2	0.4	10	3.1	2.0	0	0.0	0.0	3	1.3	1.5
1990	4	1.2	1.2	19	5.8	2.7	0	0.0	0.0	2	1.0	1.4
1991	4	2.3	2.4	22	6.6	2.8	1	0.8	1.6	2	0.7	0.9
1992	6	2.4	2.4	15	3.7	1.9	1	0.3	0.7	1	0.3	0.6
1993	7	1.8	1.4	19	5.1	2.4	2	0.9	1.2	1	0.4	0.8
1994	2	0.6	0.9	27	7.3	2.8	0	0.0	0.0	1	0.3	0.5
1995	4	1.5	1.5	28	7.5	2.8	1	0.3	0.5	2	0.7	1.0
1996	6	2.0	1.6	22	5.5	2.3	2	0.7	1.1	1	0.3	0.5
1997	8	2.3	1.7	26	6.5	2.6	4	1.3	1.3	1	0.3	0.5
1998	6	1.9	1.5	22	5.3	2.2	1	0.6	1.1	1	0.2	0.5
1999	3	0.9	1.1	24	5.6	2.3	0	0.0	0.0	4	1.1	1.1
2000	10	2.5	1.6	41	9.6	3.0	0	0.0	0.0	2	0.6	0.8
2001	11	3.1	2.1	39	8.6	2.7	0	0.0	0.0	2	0.6	0.8
2002	9	2.5	1.7	38	8.2	2.6	1	0.3	0.5	2	0.5	0.8
2003	15	4.1	2.2	39	8.3	2.6	1	0.2	0.5	3	0.8	0.9
2004	8	3.1	2.4	53	11.3	3.1	2	0.5	0.7	4	1.1	1.1
2005	12	2.7	1.5	57	11.6	3.0	4	1.3	1.4	1	0.3	0.5
2006	16	4.6	2.4	51	10.2	2.8	1	0.3	0.6	1	0.2	0.4
2007	18	4.6	2.2	63	12.9	3.2	2	0.7	1.1	1	0.3	0.6
2008	18	4.5	2.2	59	11.8	3.0	2	0.4	0.5	2	0.4	0.6
2009	22	5.1	2.2	71	13.9	3.3	1	0.3	0.5	4	0.9	0.9
2010	17	3.5	1.7	76	14.0	3.2	4	1.0	1.1	3	0.6	0.7
2011	25	5.5	2.3	87	15.8	3.4	0	0.0	0.0	4	0.8	0.8
2012	23	4.5	1.9	85	15.6	3.4	3	0.8	1.0	6	1.1	0.9
2013	15	3.1	1.6	102	18.0	3.6	0	0.0	0.0	7	1.4	1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Hodgkin lymphoma

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	5	1.8	1.7	8	4.1	3.0	1	0.6	1.2	2	0.8	1.2
1977	4	1.4	1.5	4	1.5	1.5	3	1.6	1.8	1	0.2	0.5
1978	10	3.9	2.5	2	0.6	1.0	2	0.9	1.3	0	0.0	0.0
1979	5	1.1	0.9	3	0.7	0.8	3	0.8	1.0	1	0.9	1.8
1980	10	2.9	1.9	4	0.9	0.9	2	1.0	1.3	2	0.6	0.8
1981	8	2.0	1.5	5	1.6	1.6	4	1.4	1.5	1	0.6	1.2
1982	7	3.2	2.4	11	3.5	2.3	2	1.0	1.4	1	0.4	0.8
1983	7	2.0	1.6	5	1.4	1.3	2	0.4	0.6	0	0.0	0.0
1984	11	5.1	3.3	4	1.0	1.0	3	1.4	2.0	0	0.0	0.0
1985	9	3.2	2.2	8	2.2	1.6	2	1.0	1.5	4	1.0	1.0
1986	16	4.5	2.3	6	1.5	1.2	7	2.2	1.8	1	0.3	0.7
1987	9	2.9	2.0	15	3.7	1.9	3	1.2	1.4	1	0.2	0.4
1988	11	2.3	1.4	3	0.6	0.7	5	2.0	1.9	2	0.4	0.6
1989	10	2.0	1.3	10	2.7	1.8	1	0.4	0.7	2	0.9	1.3
1990	13	3.0	1.7	11	3.3	2.1	2	0.5	0.7	3	0.8	0.9
1991	13	3.0	1.7	12	2.8	1.7	3	0.7	0.8	4	0.9	0.9
1992	11	3.0	1.8	2	0.4	0.5	3	0.8	0.9	1	0.2	0.3
1993	16	3.4	1.8	8	1.8	1.3	2	0.5	0.8	1	0.3	0.6
1994	6	1.5	1.3	7	1.7	1.3	4	1.2	1.2	1	0.3	0.5
1995	18	3.8	1.8	9	2.0	1.3	8	1.8	1.3	0	0.0	0.0
1996	11	2.7	1.7	13	3.0	1.6	1	0.4	0.7	0	0.0	0.0
1997	14	2.9	1.6	4	0.8	0.8	2	0.4	0.5	2	0.5	0.7
1998	16	3.9	2.1	10	2.0	1.3	0	0.0	0.0	2	0.4	0.5
1999	16	3.5	1.8	14	3.0	1.6	2	0.6	0.8	2	0.6	0.8
2000	21	4.4	2.0	12	2.4	1.4	0	0.0	0.0	0	0.0	0.0
2001	19	4.0	1.8	12	2.6	1.5	2	0.6	0.9	2	0.4	0.6
2002	16	3.1	1.5	9	2.0	1.3	3	0.7	0.8	0	0.0	0.0
2003	10	1.8	1.1	13	2.5	1.3	1	0.2	0.4	0	0.0	0.0
2004	13	2.5	1.4	16	3.0	1.5	1	0.2	0.4	0	0.0	0.0
2005	15	3.4	1.8	10	1.9	1.2	5	1.1	0.9	1	0.2	0.3
2006	19	4.3	2.1	14	3.0	1.6	4	0.9	0.9	0	0.0	0.0
2007	21	4.4	2.0	11	1.8	1.1	2	0.7	1.0	3	0.6	0.7
2008	18	3.9	1.8	11	1.9	1.2	2	0.4	0.6	1	0.2	0.4
2009	19	3.6	1.7	13	2.3	1.3	2	0.4	0.6	1	0.1	0.3
2010	20	3.9	1.8	17	3.0	1.4	1	0.2	0.4	0	0.0	0.0
2011	21	3.5	1.5	14	2.3	1.2	4	0.7	0.7	3	0.5	0.6
2012	17	3.6	1.8	21	3.6	1.6	1	0.2	0.3	0	0.0	0.0
2013	19	3.5	1.6	13	2.3	1.3	1	0.2	0.4	1	0.2	0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Non-Hodgkin lymphomas

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	17	9.6	5.0	12	6.0	3.7	6	5.4	5.1	6	4.0	3.5
1977	28	17.7	7.4	18	8.0	3.9	14	10.8	7.0	10	4.6	2.9
1978	24	12.0	5.0	23	11.6	5.2	8	4.0	3.0	5	2.6	2.5
1979	16	6.3	3.2	14	6.3	3.4	18	8.3	4.1	11	6.4	4.1
1980	16	11.0	6.8	10	4.5	3.0	8	5.9	5.0	10	4.8	3.1
1981	20	9.6	4.8	12	6.3	3.8	8	4.7	3.8	10	3.9	2.4
1982	24	12.3	5.4	10	3.7	2.4	6	3.0	2.5	2	0.6	0.9
1983	23	9.3	4.0	18	7.4	3.6	10	4.0	2.6	8	3.4	2.4
1984	24	9.7	4.2	17	7.3	3.6	20	9.1	4.2	15	6.5	3.5
1985	27	12.9	5.6	18	6.4	3.0	14	6.6	3.7	11	4.5	2.7
1986	24	9.3	4.2	26	9.7	3.9	12	5.3	3.3	8	2.9	2.2
1987	29	12.2	5.2	22	7.3	3.1	12	5.9	3.7	6	2.6	2.1
1988	35	15.8	6.2	22	7.5	3.3	21	13.1	6.6	11	3.8	2.4
1989	37	12.3	4.3	32	10.0	3.5	16	6.2	3.4	12	4.2	2.4
1990	45	15.8	5.0	33	11.5	4.1	16	6.7	3.6	15	4.9	2.5
1991	45	14.8	5.3	29	9.4	3.6	19	6.8	3.3	15	5.3	2.8
1992	53	18.6	5.6	23	7.3	3.1	19	8.4	4.1	10	3.5	2.3
1993	57	19.2	5.4	37	10.4	3.5	16	5.2	2.8	23	7.8	3.3
1994	47	14.7	4.5	37	10.9	3.6	19	8.3	4.3	9	3.2	2.1
1995	70	23.6	6.5	30	8.1	3.0	16	4.9	2.5	17	5.8	2.8
1996	69	19.8	5.0	46	13.3	4.0	19	7.9	4.1	13	4.0	2.2
1997	61	20.3	5.8	41	10.8	3.4	11	5.3	3.6	10	3.0	1.9
1998	74	22.8	5.8	44	11.3	3.4	20	7.6	3.8	14	4.0	2.2
1999	75	22.4	5.5	53	13.6	3.8	15	4.7	2.5	17	4.9	2.4
2000	55	16.1	4.7	47	11.1	3.2	31	10.6	4.2	11	2.9	1.8
2001	74	20.1	4.9	51	12.4	3.5	21	8.5	4.0	21	5.5	2.4
2002	74	20.2	5.0	54	12.8	3.5	26	7.7	3.2	22	5.5	2.4
2003	56	16.3	4.6	66	15.4	3.8	14	4.1	2.2	13	3.3	1.8
2004	59	18.4	5.3	48	11.0	3.2	19	5.8	2.9	17	4.5	2.1
2005	68	18.8	4.9	50	11.6	3.3	18	6.6	3.3	14	3.5	1.9
2006	75	19.7	4.9	64	13.9	3.5	19	4.8	2.4	15	3.5	1.8
2007	75	20.7	5.1	52	10.7	2.9	19	6.0	3.0	12	2.7	1.6
2008	94	23.9	5.2	64	13.0	3.2	21	6.3	2.9	11	2.2	1.3
2009	68	17.2	4.4	65	13.0	3.2	23	6.1	2.7	11	2.2	1.3
2010	75	17.1	4.1	65	13.0	3.2	22	5.9	2.7	20	4.4	2.0
2011	96	23.6	5.2	60	11.5	3.0	27	7.6	3.1	15	3.2	1.6
2012	71	16.3	4.0	66	12.1	3.0	16	4.8	2.5	17	3.2	1.5
2013	70	15.9	4.0	62	11.4	2.9	14	3.3	1.8	14	2.8	1.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Myeloma

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	13	8.6	5.1	18	9.7	4.6	11	7.1	4.5	11	5.8	3.6
1977	14	10.6	6.7	17	10.5	5.3	12	11.8	7.8	7	4.7	3.7
1978	12	10.1	6.8	16	7.9	4.0	9	8.8	6.7	7	3.6	2.8
1979	15	14.2	8.4	20	9.8	4.4	8	6.2	5.2	8	4.1	3.0
1980	13	8.3	4.8	20	10.1	4.7	9	5.6	3.8	13	6.8	3.9
1981	15	12.2	7.1	25	13.0	5.3	13	10.9	6.5	13	6.3	3.6
1982	26	17.1	7.3	21	10.1	4.6	16	9.9	5.2	12	5.2	3.1
1983	16	10.5	5.8	10	4.4	2.8	13	10.5	6.4	13	6.3	3.5
1984	27	14.9	6.3	22	10.4	4.5	13	9.9	6.0	14	6.4	3.5
1985	21	11.7	5.5	17	7.6	3.7	16	8.6	4.4	13	6.1	3.4
1986	25	16.4	7.2	27	12.5	4.9	22	14.7	6.8	13	6.6	3.7
1987	19	14.1	7.2	20	8.0	3.6	13	9.9	6.1	9	4.1	2.7
1988	20	11.5	5.5	25	9.8	4.0	10	5.4	3.5	10	4.1	2.6
1989	24	13.5	6.0	33	13.5	4.7	11	7.4	4.9	18	7.5	3.5
1990	29	16.8	6.7	25	9.5	3.8	16	9.0	5.0	17	6.8	3.3
1991	26	14.0	6.1	33	11.8	4.1	12	6.8	4.1	14	5.0	2.7
1992	27	13.8	5.8	31	10.9	3.9	23	14.2	6.6	27	9.9	3.8
1993	26	14.5	6.3	31	10.1	3.6	15	8.2	4.4	24	7.9	3.2
1994	27	13.1	5.5	33	11.8	4.1	16	8.4	4.5	25	8.6	3.5
1995	28	14.2	6.0	24	7.5	3.1	14	7.2	4.4	16	5.4	2.7
1996	29	12.3	5.1	32	9.9	3.5	17	8.2	4.6	11	3.5	2.1
1997	36	15.6	5.5	36	11.1	3.7	31	15.1	5.9	16	5.1	2.5
1998	31	13.2	5.1	36	10.6	3.5	15	7.4	4.3	21	6.6	2.9
1999	35	13.6	4.9	32	9.5	3.3	17	8.0	4.0	20	6.2	2.8
2000	41	15.5	5.3	45	12.6	3.8	25	10.3	4.4	20	5.9	2.6
2001	41	14.4	4.8	46	13.0	3.8	18	7.4	3.9	31	8.9	3.2
2002	30	12.0	4.7	42	11.2	3.4	16	5.4	2.9	24	6.5	2.6
2003	41	13.8	4.5	45	11.1	3.3	17	6.1	3.1	18	4.6	2.2
2004	42	16.1	5.4	51	13.1	3.7	24	10.6	4.6	18	4.8	2.3
2005	50	16.6	5.0	62	15.3	3.9	16	4.9	2.6	22	5.6	2.4
2006	52	16.4	4.8	51	11.5	3.2	26	8.6	3.5	19	4.7	2.1
2007	40	12.3	4.1	47	10.3	3.0	22	7.7	3.4	22	5.4	2.3
2008	54	14.2	4.0	67	14.1	3.4	22	7.2	3.2	20	4.6	2.1
2009	65	17.7	4.6	56	11.9	3.2	19	5.1	2.4	17	3.7	1.8
2010	58	18.2	5.0	65	13.2	3.3	24	7.7	3.3	26	5.5	2.2
2011	43	11.1	3.5	70	14.2	3.4	15	4.7	2.5	28	5.7	2.1
2012	72	17.9	4.5	58	10.9	2.8	25	6.3	2.6	24	4.9	2.0
2013	83	21.0	4.8	64	12.6	3.1	23	7.2	3.1	25	4.9	1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2013
New York State Excl New York City
Blacks
Leukemias

	Incidence						Mortality					
	Males			Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI (+/-)	Cases	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)	Deaths	Rate per 100,000	95% CI (+/-)
1976	19	10.9	6.5	12	4.6	2.8	12	6.8	4.7	6	3.1	2.8
1977	21	17.2	9.4	14	6.8	3.7	14	8.1	5.8	6	3.0	2.7
1978	16	7.8	4.2	23	11.4	5.0	15	10.1	6.3	17	9.2	4.9
1979	15	6.9	3.6	21	10.2	4.7	12	7.1	4.3	10	4.1	2.6
1980	21	11.6	6.0	23	9.2	4.0	13	6.7	4.3	18	7.2	3.5
1981	22	13.9	7.1	18	8.1	3.9	13	10.7	7.2	14	6.1	3.4
1982	15	7.4	4.2	17	6.3	3.2	8	3.3	2.7	10	3.6	2.4
1983	25	13.7	6.7	19	7.6	3.6	12	9.1	6.0	14	5.3	2.9
1984	16	7.2	4.2	17	6.2	3.3	13	6.9	4.6	9	4.2	2.8
1985	29	13.6	5.7	29	10.4	4.1	12	6.0	4.1	10	3.7	2.4
1986	28	13.0	5.7	14	5.2	2.9	21	7.6	3.6	14	5.9	3.3
1987	27	12.2	5.3	23	8.3	3.6	15	8.5	5.2	16	6.4	3.2
1988	19	11.0	5.5	26	8.9	3.5	20	11.3	5.9	20	7.2	3.3
1989	26	12.8	5.8	22	8.0	3.5	14	8.3	4.9	4	1.4	1.4
1990	33	14.4	5.8	31	10.9	4.0	14	8.1	5.0	19	6.9	3.2
1991	28	12.2	5.4	30	9.7	3.6	23	10.7	5.3	12	4.3	2.5
1992	31	11.8	5.0	27	8.3	3.2	12	5.5	3.7	9	3.3	2.2
1993	29	13.3	5.6	25	7.6	3.1	16	7.0	4.1	11	3.2	2.0
1994	30	13.1	5.3	26	7.5	3.0	23	11.5	5.3	20	6.6	3.0
1995	42	16.4	5.8	25	7.0	2.8	23	10.1	4.7	19	5.2	2.4
1996	44	15.8	5.3	29	7.7	2.9	21	8.4	4.0	21	6.1	2.7
1997	36	11.6	4.1	28	8.2	3.1	23	9.5	4.3	16	4.8	2.4
1998	38	14.5	5.5	35	9.7	3.3	25	10.6	4.6	22	6.4	2.8
1999	30	11.5	4.8	30	7.4	2.7	18	6.8	3.6	19	5.3	2.4
2000	39	12.7	4.4	27	7.0	2.7	18	5.9	3.1	14	3.6	1.9
2001	38	11.9	4.2	34	8.5	2.9	24	8.5	3.8	23	5.7	2.4
2002	46	15.0	4.7	38	9.7	3.2	14	4.5	2.6	17	5.0	2.4
2003	39	11.8	4.1	40	9.0	2.9	17	5.6	2.8	21	5.5	2.4
2004	61	18.8	5.4	38	8.8	2.9	28	11.0	4.5	24	6.1	2.5
2005	33	10.5	4.0	33	7.5	2.6	21	7.0	3.3	20	4.8	2.2
2006	58	15.4	4.4	41	9.1	2.8	19	5.9	2.9	20	4.6	2.1
2007	46	13.0	4.2	40	8.5	2.7	23	7.2	3.2	20	4.4	2.0
2008	51	11.9	3.5	39	8.0	2.5	23	6.8	3.1	21	4.5	1.9
2009	44	10.9	3.5	28	6.0	2.2	22	5.5	2.5	12	2.5	1.4
2010	61	14.4	3.9	37	7.2	2.4	23	6.9	3.1	12	2.3	1.3
2011	63	16.4	4.4	43	8.2	2.5	28	8.3	3.3	19	3.8	1.7
2012	62	16.2	4.3	59	11.0	2.8	31	8.1	3.1	16	3.0	1.5
2013	63	14.7	3.8	54	9.9	2.7	28	7.4	2.9	21	3.9	1.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2015.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2009-2013 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.