New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
All Invasive Malignant Tumors

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	59295	393.5	390.3-396.8	28733	461.5	455.9-467.2	30562	359.0	354.9-363.1	32504	217.1	214.7-219.5	16784	276.3	271.9-280.7	15720	181.5	178.7-184.5
1977	59281	393.4	390.2-396.7	28831	463.7	458.1-469.4	30450	357.0	352.9-361.2	32787	218.5	216.1-220.9	16936	278.3	273.9-282.8	15851	182.3	179.4-185.2
1978	59655	396.0	392.8-399.3	28981	467.5	461.9-473.2	30674	359.3	355.2-363.4	32417	215.2	212.8-217.6	16721	275.7	271.3-280.1	15696	179.3	176.5-182.2
1979	60778	402.6	399.4-405.9	29504	475.7	470.0-481.4	31274	365.4	361.2-369.5	32331	214.4	212.0-216.8	16522	272.3	268.0-276.7	15809	180.1	177.3-183.0
1980	62680	413.8	410.5-417.1	30965	497.1	491.4-502.9	31715	369.7	365.5-373.9	33166	218.7	216.3-221.1	17047	280.7	276.3-285.1	16119	182.0	179.2-184.9
1981	62911	415.5	412.2-418.8	30637	494.0	488.2-499.8	32274	375.5	371.3-379.7	32665	215.2	212.8-217.5	16659	274.4	270.1-278.8	16006	180.6	177.8-183.5
1982	63752	419.7	416.4-423.1	31004	498.9	493.1-504.7	32748	379.8	375.5-384.0	32608	214.4	212.0-216.8	16614	274.3	270.0-278.7	15994	179.9	177.0-182.7
1983	65045	427.2	423.8-430.5	31688	507.0	501.2-512.9	33357	386.7	382.4-391.0	32659	213.7	211.4-216.1	16571	273.3	269.0-277.7	16088	179.6	176.8-182.5
1984	65641	430.6	427.3-434.0	31746	506.7	500.9-512.5	33895	392.2	387.9-396.5	32972	215.3	212.9-217.6	16618	272.7	268.4-277.1	16354	182.4	179.5-185.3
1985	67298	441.3	438.0-444.7	32858	521.8	516.0-527.7	34440	399.0	394.7-403.4	32944	214.8	212.4-217.1	16682	272.8	268.5-277.2	16262	181.3	178.5-184.2
1986	67283	441.3	437.9-444.7	32693	520.4	514.6-526.3	34590	400.8	396.5-405.2	33001	214.8	212.4-217.1	16603	271.2	266.9-275.5	16398	182.3	179.4-185.1
1987	67631	442.4	439.0-445.8	32658	518.1	512.3-524.0	34973	403.9	399.5-408.2	32849	213.6	211.2-215.9	16550	270.6	266.3-274.9	16299	180.3	177.5-183.2
1988	67107	439.1	435.7-442.4	32192	511.4	505.7-517.2	34915	403.7	399.4-408.1	33152	215.2	212.9-217.6	16728	273.3	269.0-277.6	16424	181.9	179.1-184.8
1989	67228	440.6	437.3-444.0	32702	517.6	511.8-523.4	34526	400.7	396.3-405.0	32321	210.2	207.9-212.5	16163	263.8	259.6-268.0	16158	179.3	176.5-182.2
1990	70477	460.7	457.3-464.2	34448	541.1	535.2-547.0	36029	418.3	413.9-422.8	32205	208.7	206.4-211.0	16046	261.7	257.5-265.9	16159	178.3	175.5-181.1
1991	71912	469.7	466.2-473.2	35684	559.0	553.1-565.0	36228	421.2	416.8-425.7	32411	209.8	207.5-212.2	16200	263.3	259.1-267.5	16211	179.0	176.2-181.9
1992	73954	480.6	477.1-484.1	37646	585.3	579.2-591.3	36308	421.1	416.6-425.5	32425	208.8	206.5-211.1	16061	260.5	256.3-264.6	16364	179.8	177.0-182.6
1993	73070	473.9	470.4-477.3	36848	567.7	561.8-573.7	36222	418.8	414.4-423.3	32092	205.8	203.5-208.1	15924	256.6	252.5-260.7	16168	176.7	174.0-179.5
1994	75007	486.3	482.8-489.8	37301	573.1	567.2-579.1	37706	435.1	430.6-439.6	33353	213.5	211.2-215.8	16718	268.1	264.0-272.3	16635	181.5	178.7-184.4
1995	74311	481.7	478.3-485.2	36368	555.5	549.7-561.3	37943	439.1	434.7-443.7	32897	210.0	207.8-212.3	16245	259.7	255.7-263.8	16652	181.4	178.6-184.2
1996	76249	494.4	490.9-498.0	37653	574.2	568.3-580.1	38596	447.3	442.8-451.9	32464	207.0	204.8-209.3	15956	253.2	249.2-257.2	16508	180.0	177.2-182.8
1997	77444	499.9	496.3-503.4	38270	579.5	573.7-585.4	39174	451.3	446.8-455.9	31618	200.5	198.3-202.8	15645	246.2	242.3-250.2	15973	173.2	170.5-175.9
1998	79117	507.5	503.9-511.0	38971	582.6	576.8-588.5	40146	461.8	457.2-466.4	31110	195.8	193.6-198.0	15346	236.9	233.2-240.8	15764	169.8	167.1-172.5
1999	79688	506.7	503.2-510.2	39402	584.8	579.0-590.7	40286	458.7	454.1-463.2	31490	196.0	193.8-198.2	15562	238.6	234.8-242.4	15928	170.1	167.4-172.8
2000	81773	515.6	512.0-519.1	40912	602.4	596.5-608.3	40861	461.3	456.8-465.9	31361	192.9	190.8-195.1	15390	234.0	230.3-237.8	15971	168.1	165.4-170.8
2001	80931	507.8	504.3-511.3	40416	589.4	583.6-595.2	40515	456.7	452.2-461.3	30891	189.0	186.9-191.1	15242	229.6	225.9-233.3	15649	164.2	161.6-166.9
2002	81624	508.2	504.7-511.7	41453	597.4	591.7-603.2	40171	449.9	445.4-454.4	30614	185.7	183.6-187.8	15067	224.2	220.6-227.8	15547	162.4	159.8-165.0
2003	81146	502.1	498.7-505.6	40710	581.6	575.9-587.3	40436	451.1	446.6-455.6	30009	180.6	178.6-182.7	14562	214.1	210.6-217.6	15447	160.1	157.6-162.7
2004	81577	502.6	499.1-506.1	40841	578.0	572.4-583.6	40736	454.4	449.9-458.9	29720	177.9	175.9-180.0	14581	211.9	208.5-215.4	15139	156.6	154.1-159.2
2005	82543	507.2	503.7-510.7	41479	582.7	577.0-588.3	41064	457.5	453.0-462.0	29189	174.0	172.0-176.1	14385	207.8	204.4-211.2	14804	152.6	150.1-155.2
2006	84897	521.2	517.7-524.8	43271	603.9	598.1-609.6	41626	465.5	460.9-470.1	29189	173.8	171.8-175.8	14350	205.6	202.2-209.0	14839	153.3	150.8-155.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
All Invasive Malignant Tumors

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	85725	521.8	518.3-525.4	43855	603.0	597.4-608.8	41870	467.4	462.8-472.0	28977	170.7	168.8-172.7	14431	203.3	200.0-206.7	14546	149.6	147.1-152.1
2008	86844	524.0	520.4-527.5	43915	596.1	590.4-601.7	42929	476.8	472.1-481.4	29109	169.7	167.7-171.7	14576	202.8	199.5-206.1	14533	147.9	145.4-150.4
2009	86850	518.7	515.2-522.2	43838	585.8	580.3-591.4	43012	473.8	469.2-478.4	28555	165.1	163.1-167.0	14353	197.3	194.1-200.6	14202	144.0	141.5-146.4
2010	84574	500.8	497.4-504.3	42501	559.9	554.5-565.4	42073	461.9	457.4-466.5	28761	164.0	162.1-166.0	14212	192.0	188.8-195.3	14549	145.6	143.1-148.0
2011	86549	506.5	503.1-510.0	43615	565.6	560.2-571.1	42934	467.1	462.6-471.7	28375	159.7	157.8-161.6	14097	187.3	184.1-190.4	14278	140.9	138.6-143.3
2012	84482	488.6	485.2-492.0	42100	537.0	531.8-542.3	42382	458.6	454.1-463.1	28698	160.0	158.1-161.9	14417	189.0	185.9-192.2	14281	140.3	138.0-142.7
2013	86627	495.9	492.5-499.3	42748	537.4	532.2-542.6	43879	471.7	467.2-476.3	28209	154.7	152.8-156.5	14208	182.0	179.0-185.1	14001	135.8	133.5-138.1
2014	85939	485.4	482.0-488.7	42257	521.5	516.5-526.7	43682	464.9	460.4-469.4	27598	149.2	147.4-151.0	14009	176.5	173.6-179.5	13589	130.3	128.0-132.6
2015	86471	484.1	480.8-487.4	42538	517.5	512.5-522.6	43933	466.1	461.6-470.7	27530	147.1	145.3-148.9	13828	172.0	169.1-174.9	13702	129.9	127.7-132.2
2016	85798	475.6	472.3-478.9	42767	512.8	507.9-517.9	43031	454.2	449.7-458.7	27767	147.1	145.3-148.8	13986	171.3	168.5-174.3	13781	130.6	128.3-132.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Oral cavity and pharynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1627	10.5	10.0-11.1	1105	16.8	15.8-17.9	522	5.9	5.4-6.5	708	4.6	4.2-4.9	507	7.7	7.0-8.5	201	2.2	1.9-2.6
1977	1616	10.4	9.9-11.0	1068	16.2	15.2-17.3	548	6.2	5.7-6.8	698	4.5	4.2-4.9	478	7.4	6.8-8.2	220	2.4	2.1-2.8
1978	1610	10.4	9.9-11.0	1060	16.1	15.1-17.1	550	6.3	5.8-6.9	693	4.5	4.2-4.8	473	7.3	6.6-8.0	220	2.5	2.2-2.8
1979	1650	10.6	10.1-11.1	1078	16.2	15.2-17.2	572	6.5	6.0-7.1	646	4.2	3.8-4.5	420	6.3	5.7-7.0	226	2.6	2.2-2.9
1980	1611	10.5	9.9-11.0	1046	15.9	15.0-17.0	565	6.5	5.9-7.0	645	4.1	3.8-4.5	424	6.5	5.9-7.2	221	2.4	2.1-2.8
1981	1686	10.9	10.4-11.5	1161	17.5	16.5-18.6	525	6.0	5.5-6.5	635	4.1	3.7-4.4	427	6.5	5.9-7.2	208	2.3	2.0-2.6
1982	1749	11.5	10.9-12.0	1172	18.1	17.0-19.2	577	6.7	6.1-7.3	627	4.1	3.7-4.4	437	6.8	6.2-7.5	190	2.1	1.8-2.5
1983	1663	10.8	10.2-11.3	1109	16.8	15.8-17.8	554	6.3	5.8-6.9	603	3.8	3.5-4.2	415	6.3	5.7-6.9	188	2.1	1.8-2.4
1984	1672	10.9	10.4-11.5	1126	17.3	16.3-18.4	546	6.2	5.7-6.8	596	3.9	3.5-4.2	402	6.3	5.6-6.9	194	2.2	1.9-2.5
1985	1705	11.2	10.6-11.7	1115	17.2	16.1-18.2	590	6.7	6.2-7.3	575	3.7	3.4-4.0	379	5.9	5.3-6.5	196	2.2	1.9-2.5
1986	1607	10.6	10.1-11.1	1066	16.6	15.6-17.7	541	6.1	5.6-6.7	591	3.8	3.5-4.1	390	6.0	5.4-6.6	201	2.2	1.9-2.6
1987	1594	10.6	10.0-11.1	1022	15.8	14.8-16.8	572	6.6	6.0-7.1	546	3.6	3.3-3.9	353	5.6	5.0-6.2	193	2.1	1.8-2.4
1988	1551	10.2	9.7-10.8	988	15.1	14.2-16.1	563	6.5	5.9-7.0	591	3.9	3.6-4.2	382	6.2	5.6-6.9	209	2.3	2.0-2.7
1989	1520	10.0	9.5-10.6	998	15.3	14.4-16.3	522	6.1	5.6-6.6	490	3.2	2.9-3.5	304	4.8	4.3-5.4	186	2.1	1.8-2.4
1990	1544	10.3	9.8-10.8	993	15.3	14.4-16.4	551	6.5	5.9-7.0	494	3.2	2.9-3.5	312	4.9	4.3-5.5	182	2.0	1.7-2.3
1991	1568	10.5	10.0-11.1	1001	15.5	14.5-16.5	567	6.6	6.1-7.2	474	3.1	2.8-3.4	300	4.8	4.2-5.4	174	1.9	1.6-2.2
1992	1602	10.7	10.1-11.2	1042	16.0	15.0-17.0	560	6.5	6.0-7.1	444	2.9	2.6-3.1	272	4.2	3.7-4.7	172	1.8	1.6-2.1
1993	1555	10.4	9.8-10.9	1025	15.9	14.9-16.9	530	6.1	5.6-6.7	436	2.8	2.6-3.1	285	4.4	3.9-4.9	151	1.7	1.4-2.0
1994	1505	10.0	9.5-10.5	986	15.0	14.1-16.0	519	6.1	5.5-6.6	438	2.9	2.6-3.1	307	4.7	4.2-5.3	131	1.5	1.2-1.7
1995	1379	9.1	8.6-9.6	864	13.1	12.2-14.0	515	5.9	5.4-6.5	445	2.9	2.6-3.2	294	4.6	4.1-5.2	151	1.7	1.4-2.0
1996	1536	10.1	9.6-10.7	1007	15.0	14.1-16.0	529	6.2	5.6-6.7	413	2.7	2.5-3.0	257	3.9	3.5-4.5	156	1.8	1.5-2.1
1997	1542	10.1	9.6-10.6	992	14.8	13.9-15.7	550	6.3	5.8-6.9	406	2.6	2.4-2.9	274	4.1	3.7-4.7	132	1.4	1.2-1.7
1998	1529	10.0	9.5-10.5	960	14.2	13.3-15.1	569	6.6	6.0-7.1	418	2.6	2.4-2.9	265	4.0	3.5-4.5	153	1.6	1.4-1.9
1999	1548	10.0	9.5-10.5	1058	15.4	14.5-16.3	490	5.5	5.1-6.1	375	2.4	2.1-2.6	251	3.7	3.3-4.2	124	1.3	1.1-1.6
2000	1460	9.3	8.8-9.8	956	13.8	13.0-14.8	504	5.7	5.2-6.2	342	2.1	1.9-2.4	216	3.2	2.8-3.7	126	1.3	1.1-1.6
2001	1537	9.7	9.3-10.2	986	14.0	13.1-14.9	551	6.1	5.6-6.7	383	2.4	2.1-2.6	245	3.6	3.1-4.0	138	1.4	1.1-1.6
2002	1580	9.9	9.4-10.4	1027	14.4	13.5-15.3	553	6.1	5.6-6.7	359	2.2	2.0-2.4	228	3.3	2.9-3.7	131	1.3	1.1-1.6
2003	1572	9.8	9.3-10.3	1035	14.4	13.5-15.3	537	6.0	5.5-6.5	341	2.1	1.9-2.3	214	3.0	2.6-3.5	127	1.3	1.1-1.6
2004	1632	10.1	9.6-10.6	1086	15.0	14.1-15.9	546	6.0	5.5-6.6	405	2.4	2.2-2.7	260	3.6	3.2-4.1	145	1.5	1.3-1.8
2005	1566	9.6	9.2-10.1	1027	13.9	13.1-14.8	539	6.1	5.6-6.6	344	2.1	1.8-2.3	225	3.1	2.7-3.5	119	1.2	1.0-1.4
2006	1700	10.4	9.9-10.9	1167	15.7	14.8-16.6	533	5.9	5.4-6.5	342	2.1	1.8-2.3	227	3.1	2.7-3.6	115	1.2	1.0-1.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Oral cavity and pharynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	1802	10.9	10.4-11.4	1229	16.3	15.4-17.3	573	6.4	5.9-6.9	342	2.0	1.8-2.2	229	3.1	2.7-3.5	113	1.2	1.0-1.4
2008	1771	10.7	10.2-11.2	1181	15.6	14.7-16.5	590	6.5	6.0-7.1	345	2.0	1.8-2.2	250	3.3	2.9-3.7	95	1.0	0.8-1.2
2009	1852	10.9	10.4-11.5	1282	16.5	15.6-17.5	570	6.2	5.6-6.7	364	2.1	1.9-2.3	247	3.3	2.9-3.7	117	1.1	0.9-1.4
2010	1875	10.9	10.4-11.4	1251	15.9	15.0-16.8	624	6.8	6.2-7.4	381	2.2	2.0-2.4	252	3.3	2.9-3.7	129	1.3	1.1-1.5
2011	1877	10.8	10.3-11.3	1307	16.3	15.4-17.2	570	6.1	5.6-6.6	351	2.0	1.8-2.2	232	3.0	2.6-3.4	119	1.2	1.0-1.4
2012	1914	10.9	10.4-11.4	1304	16.1	15.2-17.0	610	6.4	5.9-6.9	391	2.2	2.0-2.4	271	3.4	3.0-3.8	120	1.1	0.9-1.4
2013	1970	11.2	10.7-11.7	1383	17.0	16.1-17.9	587	6.1	5.6-6.6	371	2.0	1.8-2.2	257	3.1	2.7-3.5	114	1.1	0.9-1.3
2014	2060	11.4	10.9-12.0	1385	16.6	15.7-17.5	675	7.0	6.4-7.6	330	1.8	1.6-2.0	231	2.8	2.5-3.2	99	0.9	0.7-1.1
2015	2111	11.7	11.2-12.2	1501	17.8	16.9-18.8	610	6.4	5.9-6.9	397	2.1	1.9-2.4	273	3.3	2.9-3.7	124	1.1	0.9-1.4
2016	2205	12.0	11.5-12.5	1528	17.9	17.0-18.9	677	7.0	6.4-7.5	421	2.2	2.0-2.4	297	3.5	3.1-3.9	124	1.1	0.9-1.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Esophagus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	630	4.1	3.8-4.4	455	7.0	6.3-7.7	175	2.0	1.7-2.3	584	3.8	3.5-4.1	423	6.6	5.9-7.3	161	1.8	1.6-2.2
1977	622	4.1	3.8-4.4	422	6.6	6.0-7.3	200	2.3	2.0-2.6	568	3.7	3.4-4.0	383	6.1	5.4-6.7	185	2.1	1.8-2.4
1978	618	4.0	3.7-4.4	421	6.6	6.0-7.3	197	2.2	1.9-2.5	627	4.1	3.8-4.4	434	6.9	6.3-7.6	193	2.1	1.9-2.5
1979	635	4.1	3.8-4.4	441	6.8	6.2-7.5	194	2.1	1.8-2.5	560	3.6	3.3-4.0	379	5.9	5.3-6.5	181	2.0	1.7-2.4
1980	668	4.3	4.0-4.6	451	7.1	6.5-7.8	217	2.3	2.0-2.7	563	3.6	3.3-3.9	382	6.0	5.4-6.7	181	2.0	1.7-2.3
1981	626	4.0	3.7-4.4	418	6.5	5.9-7.2	208	2.3	2.0-2.6	602	3.9	3.6-4.2	429	6.8	6.1-7.5	173	1.9	1.6-2.2
1982	629	4.0	3.7-4.3	405	6.3	5.7-7.0	224	2.4	2.1-2.7	558	3.6	3.3-3.9	379	6.0	5.4-6.7	179	1.9	1.7-2.2
1983	637	4.1	3.8-4.4	447	6.9	6.3-7.6	190	2.1	1.8-2.4	575	3.7	3.4-4.1	416	6.7	6.1-7.4	159	1.7	1.5-2.0
1984	637	4.0	3.7-4.4	436	6.8	6.2-7.5	201	2.2	1.9-2.5	584	3.7	3.4-4.0	394	6.1	5.5-6.8	190	2.0	1.7-2.3
1985	612	4.0	3.7-4.3	440	6.9	6.2-7.6	172	1.9	1.6-2.2	551	3.5	3.2-3.8	405	6.3	5.6-6.9	146	1.6	1.3-1.8
1986	667	4.3	4.0-4.7	449	7.0	6.4-7.7	218	2.4	2.1-2.8	564	3.6	3.3-4.0	413	6.6	5.9-7.2	151	1.6	1.4-1.9
1987	617	4.0	3.7-4.3	407	6.3	5.7-6.9	210	2.3	2.0-2.6	573	3.7	3.4-4.0	396	6.2	5.6-6.9	177	1.9	1.6-2.2
1988	604	3.9	3.6-4.2	414	6.4	5.8-7.0	190	2.0	1.8-2.4	623	4.0	3.7-4.3	455	7.1	6.5-7.8	168	1.8	1.5-2.1
1989	655	4.2	3.9-4.5	456	7.0	6.4-7.7	199	2.1	1.9-2.5	594	3.8	3.5-4.2	422	6.6	5.9-7.2	172	1.8	1.6-2.1
1990	695	4.5	4.2-4.8	461	7.1	6.4-7.7	234	2.5	2.2-2.9	587	3.7	3.4-4.1	411	6.3	5.7-7.0	176	1.9	1.6-2.2
1991	671	4.3	4.0-4.7	454	6.9	6.3-7.6	217	2.4	2.0-2.7	603	3.8	3.5-4.2	424	6.5	5.9-7.2	179	1.9	1.6-2.2
1992	738	4.7	4.4-5.1	493	7.5	6.8-8.2	245	2.7	2.3-3.0	636	4.1	3.8-4.4	461	7.0	6.4-7.7	175	1.9	1.6-2.2
1993	698	4.5	4.2-4.8	496	7.5	6.9-8.2	202	2.2	1.9-2.5	618	4.0	3.6-4.3	455	7.0	6.3-7.7	163	1.7	1.5-2.0
1994	745	4.8	4.5-5.2	534	8.1	7.5-8.9	211	2.2	1.9-2.6	647	4.1	3.8-4.5	460	7.0	6.4-7.7	187	2.0	1.7-2.3
1995	708	4.6	4.3-4.9	480	7.4	6.7-8.1	228	2.5	2.1-2.8	683	4.4	4.1-4.8	505	7.8	7.1-8.5	178	1.9	1.6-2.2
1996	751	4.8	4.5-5.2	519	7.9	7.2-8.6	232	2.5	2.2-2.8	669	4.3	4.0-4.7	466	7.1	6.5-7.8	203	2.1	1.9-2.5
1997	701	4.5	4.2-4.8	481	7.3	6.7-8.0	220	2.4	2.1-2.7	651	4.2	3.8-4.5	488	7.4	6.8-8.1	163	1.7	1.4-2.0
1998	773	4.9	4.6-5.3	570	8.5	7.8-9.3	203	2.2	1.9-2.5	679	4.3	4.0-4.7	502	7.5	6.9-8.2	177	1.9	1.6-2.2
1999	744	4.7	4.4-5.0	547	8.1	7.4-8.8	197	2.0	1.8-2.4	681	4.3	4.0-4.6	506	7.5	6.9-8.2	175	1.8	1.6-2.1
2000	783	4.9	4.5-5.2	561	8.2	7.6-8.9	222	2.3	2.0-2.6	698	4.3	4.0-4.7	495	7.3	6.7-8.0	203	2.1	1.8-2.4
2001	745	4.6	4.3-5.0	546	7.9	7.2-8.6	199	2.1	1.8-2.4	701	4.3	4.0-4.7	538	7.8	7.2-8.5	163	1.7	1.4-1.9
2002	798	4.9	4.6-5.3	596	8.5	7.8-9.2	202	2.1	1.8-2.4	679	4.1	3.8-4.5	507	7.3	6.7-8.0	172	1.7	1.5-2.0
2003	831	5.1	4.7-5.4	612	8.6	8.0-9.3	219	2.2	1.9-2.5	736	4.5	4.1-4.8	535	7.7	7.0-8.3	201	2.0	1.7-2.3
2004	842	5.1	4.8-5.5	624	8.8	8.1-9.5	218	2.2	1.9-2.5	746	4.5	4.2-4.9	567	8.0	7.3-8.7	179	1.8	1.5-2.1
2005	899	5.5	5.1-5.8	669	9.4	8.7-10.1	230	2.4	2.1-2.7	806	4.8	4.5-5.2	623	8.8	8.1-9.5	183	1.8	1.6-2.1
2006	933	5.6	5.3-6.0	724	10.0	9.3-10.8	209	2.2	1.9-2.5	731	4.4	4.1-4.8	546	7.6	7.0-8.3	185	1.9	1.7-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Esophagus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	874	5.2	4.9-5.6	655	8.9	8.2-9.6	219	2.3	2.0-2.6	775	4.6	4.3-4.9	578	8.0	7.3-8.7	197	2.0	1.7-2.3
2008	947	5.6	5.3-6.0	717	9.7	9.0-10.4	230	2.4	2.1-2.7	749	4.4	4.1-4.7	586	7.9	7.2-8.5	163	1.7	1.4-1.9
2009	932	5.4	5.1-5.8	720	9.5	8.8-10.2	212	2.1	1.8-2.4	755	4.4	4.1-4.7	590	7.9	7.2-8.5	165	1.6	1.4-1.9
2010	862	4.9	4.6-5.3	644	8.4	7.8-9.1	218	2.1	1.9-2.5	746	4.3	4.0-4.6	585	7.7	7.1-8.4	161	1.6	1.3-1.8
2011	835	4.8	4.4-5.1	627	8.0	7.4-8.7	208	2.1	1.8-2.4	775	4.4	4.1-4.7	585	7.6	7.0-8.2	190	1.8	1.6-2.1
2012	804	4.5	4.2-4.8	631	7.9	7.3-8.6	173	1.7	1.4-2.0	711	4.0	3.7-4.3	550	7.1	6.5-7.7	161	1.5	1.3-1.8
2013	891	4.9	4.6-5.3	684	8.4	7.8-9.1	207	2.0	1.7-2.3	737	4.0	3.7-4.3	570	7.0	6.5-7.7	167	1.6	1.4-1.9
2014	894	4.8	4.5-5.1	678	8.1	7.5-8.8	216	2.1	1.8-2.4	752	4.1	3.8-4.5	603	7.5	6.9-8.2	149	1.4	1.2-1.7
2015	931	5.0	4.7-5.4	700	8.3	7.7-9.0	231	2.3	2.0-2.6	707	3.8	3.5-4.0	530	6.4	5.8-6.9	177	1.6	1.4-1.9
2016	925	4.9	4.6-5.2	711	8.4	7.7-9.0	214	2.0	1.8-2.4	757	4.0	3.7-4.3	594	7.0	6.4-7.6	163	1.5	1.3-1.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Stomach

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1813	12.3	11.8-12.9	1038	17.2	16.2-18.4	775	9.0	8.3-9.6	1431	9.8	9.3-10.3	793	13.5	12.5-14.5	638	7.4	6.8-8.0
1977	1799	12.3	11.8-12.9	1042	17.6	16.5-18.8	757	8.8	8.2-9.4	1397	9.6	9.1-10.2	797	13.8	12.8-14.9	600	6.9	6.4-7.5
1978	1765	12.0	11.4-12.5	984	16.7	15.6-17.8	781	8.9	8.3-9.5	1300	8.9	8.5-9.5	721	12.5	11.5-13.5	579	6.6	6.1-7.2
1979	1800	12.0	11.4-12.6	1020	16.9	15.8-18.0	780	8.7	8.1-9.4	1325	8.9	8.4-9.4	746	12.7	11.7-13.6	579	6.4	5.9-7.0
1980	1767	11.8	11.3-12.4	1042	17.4	16.3-18.5	725	8.2	7.6-8.8	1322	8.9	8.4-9.4	751	12.8	11.8-13.8	571	6.4	5.9-7.0
1981	1805	12.0	11.5-12.6	1049	17.5	16.4-18.6	756	8.4	7.8-9.1	1284	8.6	8.1-9.1	751	12.8	11.8-13.7	533	5.9	5.4-6.4
1982	1727	11.3	10.8-11.9	994	16.4	15.4-17.5	733	8.0	7.4-8.6	1197	8.0	7.5-8.4	704	12.0	11.1-12.9	493	5.4	4.9-5.9
1983	1785	11.8	11.3-12.4	1012	16.8	15.7-17.9	773	8.4	7.8-9.1	1198	8.0	7.5-8.4	690	11.8	10.9-12.7	508	5.5	5.1-6.0
1984	1703	11.1	10.6-11.7	985	15.9	14.9-17.0	718	7.8	7.2-8.4	1217	8.0	7.5-8.4	677	11.0	10.2-11.9	540	5.8	5.3-6.3
1985	1690	11.1	10.5-11.6	1017	16.7	15.6-17.8	673	7.3	6.8-7.9	1148	7.6	7.1-8.0	655	10.9	10.1-11.8	493	5.4	4.9-5.9
1986	1627	10.6	10.1-11.2	964	15.8	14.8-16.9	663	7.2	6.6-7.7	1106	7.2	6.8-7.7	645	10.7	9.9-11.6	461	4.9	4.5-5.4
1987	1659	10.7	10.2-11.3	973	15.8	14.8-16.9	686	7.3	6.7-7.9	1103	7.2	6.7-7.6	652	10.8	10.0-11.7	451	4.8	4.4-5.3
1988	1579	10.3	9.8-10.8	891	14.5	13.5-15.5	688	7.4	6.8-8.0	1064	6.9	6.5-7.4	578	9.5	8.7-10.3	486	5.2	4.7-5.7
1989	1510	9.8	9.3-10.3	933	15.1	14.1-16.2	577	6.1	5.6-6.7	1109	7.2	6.8-7.6	645	10.7	9.9-11.6	464	4.9	4.5-5.4
1990	1703	11.1	10.6-11.7	1029	16.6	15.6-17.7	674	7.3	6.7-7.8	1001	6.6	6.2-7.0	613	10.1	9.3-11.0	388	4.1	3.7-4.5
1991	1589	10.3	9.8-10.8	964	15.7	14.7-16.7	625	6.6	6.1-7.2	1005	6.5	6.1-6.9	615	10.2	9.4-11.1	390	4.1	3.7-4.5
1992	1566	10.1	9.6-10.6	941	15.2	14.3-16.3	625	6.6	6.1-7.2	1025	6.6	6.2-7.0	614	10.2	9.4-11.1	411	4.3	3.9-4.7
1993	1632	10.5	10.0-11.0	970	15.6	14.6-16.7	662	6.9	6.4-7.5	998	6.4	6.0-6.8	587	9.8	9.0-10.6	411	4.3	3.9-4.7
1994	1612	10.4	9.9-10.9	934	14.8	13.9-15.8	678	7.2	6.6-7.8	1044	6.7	6.3-7.1	610	10.0	9.2-10.8	434	4.5	4.1-5.0
1995	1526	9.7	9.2-10.2	896	14.1	13.2-15.1	630	6.7	6.1-7.2	1003	6.4	6.0-6.8	559	9.2	8.4-10.0	444	4.6	4.2-5.1
1996	1546	9.9	9.4-10.4	900	14.2	13.3-15.2	646	6.7	6.2-7.3	939	6.0	5.6-6.4	515	8.3	7.6-9.1	424	4.4	4.0-4.8
1997	1508	9.5	9.1-10.0	896	14.0	13.1-14.9	612	6.4	5.9-7.0	888	5.6	5.3-6.0	490	7.7	7.1-8.5	398	4.2	3.7-4.6
1998	1576	9.9	9.4-10.4	958	14.6	13.7-15.6	618	6.5	6.0-7.1	884	5.5	5.2-5.9	495	7.7	7.1-8.5	389	4.0	3.6-4.4
1999	1486	9.3	8.8-9.8	914	13.9	13.0-14.9	572	5.9	5.5-6.5	838	5.2	4.8-5.5	487	7.5	6.9-8.2	351	3.6	3.2-4.0
2000	1470	9.0	8.6-9.5	860	13.0	12.1-13.9	610	6.2	5.7-6.7	811	4.9	4.6-5.3	439	6.8	6.2-7.4	372	3.7	3.4-4.2
2001	1449	8.9	8.5-9.4	882	13.1	12.2-14.0	567	6.0	5.5-6.5	789	4.8	4.5-5.2	434	6.5	5.9-7.2	355	3.6	3.2-4.0
2002	1383	8.4	8.0-8.9	823	12.0	11.2-12.9	560	5.8	5.4-6.4	726	4.4	4.1-4.7	413	6.2	5.6-6.8	313	3.1	2.7-3.5
2003	1473	8.9	8.5-9.4	875	12.6	11.8-13.5	598	6.2	5.7-6.7	724	4.3	4.0-4.7	401	5.9	5.3-6.5	323	3.3	2.9-3.7
2004	1421	8.6	8.1-9.1	829	12.0	11.2-12.8	592	6.2	5.7-6.7	777	4.6	4.3-5.0	428	6.3	5.7-6.9	349	3.5	3.2-3.9
2005	1348	8.1	7.7-8.6	801	11.4	10.6-12.2	547	5.7	5.2-6.2	715	4.2	3.9-4.6	404	5.9	5.3-6.5	311	3.1	2.7-3.5
2006	1312	7.9	7.5-8.4	785	11.1	10.3-11.9	527	5.6	5.1-6.1	631	3.8	3.5-4.1	358	5.1	4.6-5.7	273	2.8	2.5-3.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Stomach

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	1386	8.3	7.9-8.8	846	11.7	11.0-12.6	540	5.7	5.2-6.2	606	3.6	3.3-3.9	347	4.8	4.3-5.4	259	2.7	2.3-3.0
2008	1278	7.5	7.1-8.0	808	11.2	10.4-12.0	470	4.9	4.4-5.4	642	3.7	3.4-4.0	396	5.5	5.0-6.1	246	2.4	2.1-2.7
2009	1236	7.3	6.9-7.7	753	10.3	9.6-11.1	483	4.9	4.5-5.4	593	3.5	3.2-3.8	353	4.9	4.4-5.5	240	2.4	2.1-2.7
2010	1309	7.7	7.3-8.1	809	10.8	10.0-11.6	500	5.2	4.7-5.7	581	3.3	3.0-3.6	313	4.3	3.8-4.8	268	2.7	2.3-3.0
2011	1313	7.6	7.2-8.0	797	10.6	9.9-11.4	516	5.3	4.8-5.8	555	3.1	2.9-3.4	311	4.2	3.7-4.7	244	2.4	2.1-2.7
2012	1329	7.6	7.2-8.0	816	10.6	9.8-11.3	513	5.3	4.8-5.8	578	3.2	3.0-3.5	337	4.4	4.0-4.9	241	2.3	2.0-2.7
2013	1424	8.0	7.6-8.5	870	11.1	10.3-11.9	554	5.6	5.1-6.1	581	3.2	2.9-3.4	353	4.5	4.0-5.0	228	2.2	1.9-2.5
2014	1315	7.4	7.0-7.8	852	10.7	10.0-11.5	463	4.8	4.3-5.2	588	3.2	3.0-3.5	354	4.5	4.0-5.0	234	2.3	2.0-2.6
2015	1306	7.2	6.8-7.6	787	9.7	9.0-10.4	519	5.2	4.7-5.7	516	2.8	2.6-3.1	301	3.8	3.3-4.2	215	2.1	1.8-2.4
2016	1349	7.4	7.0-7.9	818	10.0	9.3-10.7	531	5.5	5.0-6.0	536	2.9	2.6-3.1	339	4.1	3.7-4.6	197	1.9	1.7-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Colorectal

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	10070	67.3	66.0-68.7	4916	80.5	78.2-83.0	5154	58.8	57.2-60.5	5252	35.8	34.8-36.8	2572	44.0	42.2-45.8	2680	30.8	29.6-32.0
1977	10133	67.4	66.1-68.8	4965	81.5	79.2-84.0	5168	58.5	56.9-60.1	5119	34.9	33.9-35.8	2475	42.6	40.8-44.4	2644	30.1	28.9-31.3
1978	10094	67.1	65.7-68.4	4936	81.3	78.9-83.7	5158	58.0	56.4-59.6	5165	34.9	34.0-35.9	2568	43.8	42.1-45.7	2597	29.4	28.3-30.6
1979	10238	67.7	66.4-69.0	4942	81.1	78.7-83.5	5296	59.1	57.5-60.7	5131	34.4	33.4-35.4	2452	42.0	40.2-43.7	2679	29.8	28.7-31.0
1980	10616	69.7	68.3-71.0	5313	86.4	84.0-88.9	5303	58.7	57.1-60.3	5120	34.2	33.3-35.2	2530	43.4	41.6-45.2	2590	28.7	27.6-29.8
1981	10686	70.3	68.9-71.6	5301	87.6	85.1-90.0	5385	59.3	57.7-60.9	4995	33.3	32.3-34.2	2405	41.0	39.3-42.7	2590	28.4	27.3-29.6
1982	10670	69.7	68.4-71.1	5231	85.5	83.1-88.0	5439	59.5	57.9-61.2	4936	32.7	31.8-33.6	2438	41.6	39.9-43.4	2498	27.3	26.2-28.4
1983	10709	69.6	68.3-71.0	5308	86.4	84.0-88.9	5401	58.8	57.3-60.5	4892	32.3	31.3-33.2	2373	40.4	38.7-42.1	2519	27.2	26.2-28.3
1984	10682	69.5	68.1-70.8	5172	84.0	81.6-86.4	5510	60.0	58.4-61.6	5044	33.0	32.1-34.0	2509	42.3	40.6-44.1	2535	27.4	26.3-28.5
1985	10925	70.6	69.2-71.9	5489	88.6	86.2-91.0	5436	58.9	57.3-60.5	4832	31.6	30.7-32.5	2357	39.4	37.7-41.1	2475	26.5	25.5-27.6
1986	10725	69.3	68.0-70.6	5310	86.1	83.7-88.5	5415	58.6	57.0-60.2	4779	31.2	30.3-32.1	2360	39.7	38.1-41.4	2419	25.9	24.9-27.0
1987	10257	66.1	64.8-67.4	5087	82.3	80.0-84.7	5170	55.7	54.2-57.3	4739	30.7	29.9-31.6	2344	39.3	37.7-41.0	2395	25.4	24.4-26.5
1988	9907	63.8	62.5-65.1	4830	77.8	75.6-80.1	5077	54.5	53.0-56.1	4645	30.2	29.3-31.0	2342	39.1	37.5-40.8	2303	24.6	23.6-25.6
1989	9640	62.2	61.0-63.5	4794	77.2	75.0-79.5	4846	52.3	50.8-53.8	4319	27.9	27.1-28.8	2059	34.4	32.9-36.0	2260	23.9	22.9-25.0
1990	9645	61.8	60.6-63.1	4678	74.4	72.3-76.7	4967	53.1	51.6-54.6	4238	27.3	26.5-28.1	2114	35.1	33.6-36.7	2124	22.4	21.4-23.4
1991	9583	61.5	60.3-62.8	4780	76.8	74.6-79.1	4803	51.6	50.1-53.1	4171	26.8	26.0-27.7	2030	33.8	32.3-35.3	2141	22.5	21.6-23.5
1992	9352	59.9	58.7-61.1	4621	73.7	71.5-75.9	4731	50.7	49.2-52.2	3964	25.3	24.6-26.1	1913	31.5	30.1-33.0	2051	21.4	20.5-22.4
1993	9416	60.0	58.8-61.2	4566	71.9	69.8-74.0	4850	51.8	50.3-53.3	4001	25.4	24.6-26.2	1954	32.0	30.5-33.5	2047	21.2	20.3-22.2
1994	9860	62.7	61.5-64.0	4775	75.3	73.1-77.5	5085	54.3	52.8-55.8	4193	26.6	25.8-27.4	2014	33.1	31.7-34.7	2179	22.5	21.6-23.5
1995	9538	60.6	59.4-61.8	4648	73.1	71.0-75.3	4890	52.0	50.5-53.5	4142	26.1	25.3-26.9	1977	32.3	30.9-33.8	2165	22.3	21.4-23.3
1996	9319	59.2	58.0-60.4	4637	72.5	70.4-74.6	4682	49.9	48.5-51.4	3794	23.9	23.1-24.6	1862	29.9	28.6-31.4	1932	19.8	18.9-20.8
1997	9781	61.8	60.6-63.0	4765	73.5	71.4-75.7	5016	53.3	51.8-54.8	3758	23.5	22.8-24.3	1858	29.6	28.2-31.0	1900	19.4	18.6-20.3
1998	10172	63.9	62.6-65.1	4985	76.0	73.9-78.2	5187	55.2	53.7-56.8	3690	22.9	22.2-23.7	1754	27.3	26.1-28.7	1936	19.7	18.8-20.6
1999	10080	62.7	61.5-63.9	4966	75.0	72.9-77.1	5114	53.9	52.4-55.5	3570	21.9	21.2-22.6	1734	26.9	25.6-28.2	1836	18.5	17.7-19.4
2000	10121	62.4	61.2-63.7	5028	75.0	73.0-77.2	5093	53.2	51.7-54.7	3633	22.0	21.3-22.8	1812	27.8	26.5-29.1	1821	18.0	17.2-18.9
2001	9656	59.2	58.0-60.4	4769	70.6	68.6-72.7	4887	51.0	49.5-52.5	3453	20.8	20.1-21.5	1651	25.1	23.9-26.4	1802	17.9	17.0-18.7
2002	9517	58.0	56.9-59.2	4717	68.7	66.7-70.7	4800	49.8	48.4-51.2	3456	20.7	20.0-21.4	1688	25.3	24.1-26.5	1768	17.5	16.7-18.4
2003	9142	55.3	54.1-56.4	4477	64.6	62.7-66.6	4665	48.3	46.9-49.7	3263	19.4	18.7-20.0	1578	23.3	22.2-24.5	1685	16.6	15.8-17.4
2004	8887	53.5	52.4-54.6	4300	61.5	59.7-63.4	4587	47.4	46.0-48.9	3073	18.1	17.5-18.8	1478	21.5	20.4-22.6	1595	15.5	14.7-16.3
2005	8469	50.7	49.6-51.8	4220	59.9	58.1-61.8	4249	43.8	42.5-45.2	2843	16.6	16.0-17.2	1368	20.0	18.9-21.1	1475	14.2	13.5-15.0
2006	8121	48.7	47.6-49.8	4026	56.9	55.2-58.7	4095	42.4	41.1-43.8	2839	16.6	16.0-17.3	1378	20.0	19.0-21.1	1461	14.3	13.5-15.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Colorectal

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	8204	48.8	47.8-49.9	4056	56.3	54.5-58.0	4148	43.1	41.8-44.5	2740	15.9	15.3-16.5	1332	18.9	17.9-20.0	1408	13.6	12.9-14.4
2008	7929	46.8	45.8-47.8	3930	53.9	52.2-55.7	3999	41.3	40.0-42.7	2753	15.8	15.3-16.5	1343	18.9	17.9-19.9	1410	13.6	12.8-14.3
2009	7550	44.2	43.2-45.2	3723	50.4	48.8-52.1	3827	39.2	38.0-40.5	2646	15.1	14.5-15.7	1279	17.8	16.8-18.8	1367	13.2	12.5-13.9
2010	7505	43.6	42.6-44.6	3702	49.4	47.8-51.1	3803	38.9	37.6-40.2	2652	14.9	14.3-15.5	1324	18.0	17.0-19.0	1328	12.6	11.9-13.3
2011	7181	41.5	40.5-42.5	3554	47.0	45.5-48.6	3627	37.1	35.9-38.4	2584	14.3	13.7-14.8	1283	16.9	16.0-17.9	1301	12.1	11.4-12.8
2012	6904	39.4	38.5-40.4	3556	46.1	44.5-47.6	3348	34.1	32.9-35.4	2473	13.6	13.0-14.1	1180	15.5	14.6-16.4	1293	12.1	11.4-12.8
2013	7082	40.0	39.1-41.0	3635	46.4	44.9-48.0	3447	34.9	33.7-36.1	2448	13.3	12.8-13.8	1258	16.3	15.4-17.2	1190	10.9	10.3-11.6
2014	6851	38.3	37.4-39.2	3526	44.2	42.8-45.8	3325	33.4	32.2-34.6	2378	12.7	12.2-13.2	1201	15.2	14.3-16.1	1177	10.8	10.2-11.5
2015	6851	38.2	37.3-39.1	3442	42.8	41.4-44.3	3409	34.4	33.2-35.6	2378	12.6	12.1-13.1	1219	15.3	14.4-16.2	1159	10.5	9.9-11.2
2016	6821	37.8	36.9-38.8	3532	43.4	42.0-44.9	3289	33.3	32.1-34.5	2275	12.0	11.5-12.5	1129	13.9	13.1-14.7	1146	10.5	9.8-11.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Colon excluding rectum

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	7043	47.4	46.2-48.5	3274	54.4	52.4-56.4	3769	43.0	41.6-44.4	4137	28.3	27.4-29.2	1938	33.3	31.8-34.9	2199	25.3	24.2-26.4
1977	7118	47.7	46.6-48.9	3335	55.9	53.9-58.0	3783	42.9	41.5-44.3	4177	28.5	27.6-29.4	1973	34.0	32.4-35.6	2204	25.1	24.1-26.2
1978	7149	47.8	46.6-48.9	3355	56.0	54.0-58.0	3794	42.7	41.4-44.1	4194	28.4	27.5-29.3	2018	34.6	33.0-36.2	2176	24.6	23.5-25.6
1979	7312	48.7	47.6-49.8	3379	56.4	54.5-58.5	3933	43.9	42.6-45.3	4253	28.5	27.6-29.4	1971	33.7	32.1-35.3	2282	25.4	24.4-26.5
1980	7528	49.7	48.6-50.9	3642	60.3	58.2-62.4	3886	43.0	41.7-44.4	4256	28.5	27.6-29.3	2069	35.6	34.0-37.2	2187	24.2	23.2-25.2
1981	7616	50.2	49.1-51.4	3629	60.6	58.5-62.7	3987	43.9	42.6-45.3	4243	28.3	27.5-29.2	2010	34.4	32.8-36.0	2233	24.6	23.5-25.6
1982	7527	49.4	48.3-50.6	3498	58.2	56.2-60.3	4029	44.0	42.6-45.4	4152	27.5	26.7-28.4	2030	34.7	33.2-36.3	2122	23.2	22.2-24.2
1983	7601	49.7	48.5-50.8	3575	59.2	57.2-61.2	4026	43.8	42.5-45.2	4108	27.1	26.2-27.9	1959	33.4	31.8-34.9	2149	23.2	22.2-24.2
1984	7694	50.2	49.0-51.3	3544	58.4	56.4-60.4	4150	45.0	43.6-46.4	4269	27.9	27.1-28.8	2087	35.2	33.7-36.8	2182	23.5	22.6-24.6
1985	7922	51.3	50.2-52.5	3801	62.1	60.1-64.2	4121	44.5	43.2-45.9	4128	27.0	26.1-27.8	1981	33.0	31.5-34.6	2147	23.0	22.1-24.0
1986	7815	50.6	49.5-51.8	3681	60.3	58.3-62.4	4134	44.6	43.3-46.1	4086	26.6	25.8-27.5	1983	33.4	31.9-35.0	2103	22.5	21.6-23.5
1987	7355	47.4	46.3-48.5	3501	57.3	55.4-59.3	3854	41.2	39.9-42.6	4054	26.3	25.5-27.1	2010	33.7	32.2-35.3	2044	21.7	20.8-22.7
1988	7138	46.0	44.9-47.1	3296	53.6	51.7-55.6	3842	41.0	39.7-42.3	4014	26.0	25.2-26.9	2008	33.5	32.0-35.1	2006	21.4	20.4-22.4
1989	6878	44.5	43.4-45.5	3299	54.0	52.2-56.0	3579	38.4	37.2-39.7	3730	24.1	23.3-24.9	1742	29.2	27.8-30.7	1988	21.1	20.1-22.0
1990	6940	44.5	43.5-45.6	3223	52.0	50.2-53.9	3717	39.6	38.3-40.9	3704	23.9	23.1-24.6	1842	30.7	29.3-32.2	1862	19.6	18.7-20.5
1991	6988	44.8	43.7-45.9	3355	54.4	52.5-56.3	3633	38.7	37.5-40.0	3549	22.9	22.1-23.6	1708	28.5	27.1-29.9	1841	19.3	18.4-20.2
1992	6737	43.1	42.1-44.2	3178	51.2	49.4-53.1	3559	37.9	36.7-39.2	3389	21.6	20.9-22.4	1597	26.4	25.0-27.7	1792	18.7	17.8-19.6
1993	6868	43.7	42.7-44.8	3174	50.4	48.6-52.2	3694	39.2	37.9-40.5	3453	22.0	21.2-22.7	1669	27.5	26.1-28.9	1784	18.5	17.7-19.4
1994	7214	45.8	44.8-46.9	3388	53.9	52.1-55.8	3826	40.5	39.2-41.9	3643	23.0	22.3-23.8	1724	28.4	27.1-29.8	1919	19.8	18.9-20.7
1995	6973	44.1	43.1-45.2	3280	52.0	50.2-53.8	3693	38.8	37.6-40.1	3537	22.3	21.5-23.0	1659	27.3	25.9-28.6	1878	19.3	18.4-20.2
1996	6725	42.5	41.5-43.5	3193	50.4	48.6-52.2	3532	37.2	36.0-38.5	3252	20.4	19.7-21.1	1587	25.6	24.4-26.9	1665	17.0	16.2-17.9
1997	7250	45.7	44.6-46.7	3379	52.5	50.8-54.4	3871	40.8	39.5-42.1	3267	20.4	19.8-21.2	1580	25.1	23.9-26.4	1687	17.2	16.4-18.1
1998	7411	46.4	45.3-47.5	3508	54.0	52.2-55.9	3903	41.2	39.9-42.5	3179	19.7	19.0-20.4	1491	23.3	22.2-24.6	1688	17.1	16.3-17.9
1999	7306	45.2	44.2-46.3	3456	52.6	50.8-54.4	3850	40.1	38.9-41.4	3022	18.5	17.9-19.2	1448	22.5	21.4-23.7	1574	15.8	15.0-16.6
2000	7382	45.3	44.3-46.4	3484	52.4	50.7-54.2	3898	40.3	39.0-41.6	3106	18.8	18.1-19.5	1529	23.6	22.4-24.8	1577	15.5	14.8-16.3
2001	7044	43.0	42.0-44.0	3356	50.0	48.3-51.7	3688	37.9	36.7-39.2	2931	17.6	17.0-18.3	1373	21.0	19.9-22.1	1558	15.4	14.6-16.2
2002	6928	42.0	41.1-43.0	3291	48.3	46.7-50.0	3637	37.4	36.2-38.7	2894	17.3	16.7-18.0	1401	21.1	20.0-22.2	1493	14.7	14.0-15.5
2003	6643	40.0	39.0-40.9	3119	45.4	43.8-47.0	3524	36.1	34.9-37.3	2791	16.5	15.9-17.2	1324	19.7	18.6-20.8	1467	14.4	13.6-15.1
2004	6405	38.4	37.4-39.3	2969	42.8	41.3-44.4	3436	35.1	33.9-36.3	2586	15.2	14.6-15.8	1215	17.8	16.8-18.8	1371	13.2	12.5-14.0
2005	6214	37.0	36.1-38.0	2967	42.4	40.9-43.9	3247	33.2	32.0-34.3	2376	13.9	13.3-14.4	1119	16.4	15.5-17.4	1257	12.1	11.4-12.8
2006	5919	35.3	34.4-36.2	2823	40.2	38.8-41.8	3096	31.6	30.5-32.7	2401	14.0	13.5-14.6	1144	16.7	15.7-17.7	1257	12.2	11.5-12.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Colon excluding rectum

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	5956	35.3	34.4-36.2	2832	39.6	38.1-41.1	3124	32.0	30.9-33.2	2327	13.4	12.9-14.0	1110	15.8	14.9-16.8	1217	11.7	11.1-12.4
2008	5737	33.7	32.8-34.6	2716	37.5	36.1-38.9	3021	30.9	29.8-32.1	2262	13.0	12.5-13.6	1088	15.3	14.4-16.3	1174	11.3	10.6-12.0
2009	5406	31.4	30.6-32.3	2512	34.3	32.9-35.6	2894	29.2	28.1-30.3	2200	12.6	12.0-13.1	1044	14.6	13.7-15.5	1156	11.1	10.5-11.8
2010	5376	31.1	30.2-31.9	2530	34.1	32.7-35.5	2846	28.6	27.5-29.7	2202	12.3	11.8-12.8	1065	14.5	13.6-15.4	1137	10.6	10.0-11.3
2011	5132	29.5	28.7-30.3	2431	32.5	31.2-33.8	2701	27.2	26.1-28.3	2132	11.7	11.2-12.3	1030	13.6	12.8-14.5	1102	10.2	9.6-10.8
2012	4981	28.3	27.5-29.1	2420	31.6	30.3-32.9	2561	25.7	24.6-26.7	2044	11.2	10.7-11.7	942	12.4	11.6-13.2	1102	10.2	9.6-10.9
2013	5004	28.1	27.3-28.9	2456	31.6	30.3-32.9	2548	25.4	24.3-26.4	2009	10.9	10.4-11.4	1004	13.1	12.3-13.9	1005	9.2	8.6-9.8
2014	4859	26.9	26.2-27.7	2376	30.1	28.8-31.3	2483	24.5	23.5-25.5	1924	10.2	9.8-10.7	948	12.1	11.3-12.9	976	8.9	8.3-9.5
2015	4841	26.9	26.1-27.7	2300	28.9	27.7-30.2	2541	25.3	24.3-26.3	1936	10.2	9.7-10.7	964	12.1	11.3-12.9	972	8.8	8.2-9.4
2016	4894	27.0	26.2-27.8	2416	29.9	28.6-31.1	2478	24.7	23.7-25.7	1803	9.4	9.0-9.9	867	10.7	10.0-11.4	936	8.5	7.9-9.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Rectum & rectosigmoid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	3027	20.0	19.2-20.7	1642	26.2	24.8-27.5	1385	15.8	15.0-16.7	1115	7.5	7.1-8.0	634	10.7	9.8-11.6	481	5.5	5.0-6.0
1977	3015	19.7	19.0-20.4	1630	25.6	24.3-27.0	1385	15.6	14.8-16.5	942	6.4	6.0-6.8	502	8.6	7.8-9.4	440	5.0	4.5-5.5
1978	2945	19.3	18.6-20.0	1581	25.3	24.0-26.6	1364	15.3	14.5-16.1	971	6.6	6.2-7.0	550	9.2	8.4-10.1	421	4.8	4.4-5.3
1979	2926	19.0	18.3-19.7	1563	24.6	23.4-25.9	1363	15.1	14.3-16.0	878	5.9	5.5-6.3	481	8.3	7.5-9.1	397	4.4	4.0-4.9
1980	3088	20.0	19.3-20.7	1671	26.2	24.9-27.5	1417	15.7	14.9-16.5	864	5.8	5.4-6.2	461	7.8	7.0-8.5	403	4.5	4.0-4.9
1981	3070	20.0	19.3-20.8	1672	27.0	25.6-28.3	1398	15.4	14.6-16.2	752	5.0	4.6-5.3	395	6.6	6.0-7.3	357	3.9	3.5-4.3
1982	3143	20.3	19.6-21.0	1733	27.3	26.0-28.7	1410	15.5	14.7-16.4	784	5.2	4.8-5.6	408	6.9	6.2-7.6	376	4.1	3.7-4.5
1983	3108	20.0	19.3-20.7	1733	27.3	25.9-28.6	1375	15.0	14.2-15.8	784	5.2	4.8-5.6	414	7.0	6.3-7.8	370	4.0	3.6-4.5
1984	2988	19.3	18.6-20.0	1628	25.6	24.3-26.9	1360	15.0	14.2-15.8	775	5.1	4.7-5.5	422	7.1	6.4-7.9	353	3.8	3.4-4.2
1985	3003	19.3	18.6-20.0	1688	26.5	25.2-27.8	1315	14.4	13.6-15.2	704	4.6	4.3-5.0	376	6.3	5.7-7.0	328	3.5	3.1-3.9
1986	2910	18.7	18.0-19.4	1629	25.7	24.5-27.1	1281	13.9	13.2-14.7	693	4.5	4.2-4.9	377	6.3	5.7-7.0	316	3.4	3.0-3.8
1987	2902	18.7	18.1-19.4	1586	25.0	23.7-26.3	1316	14.5	13.7-15.3	685	4.5	4.1-4.8	334	5.6	5.0-6.3	351	3.7	3.3-4.1
1988	2769	17.8	17.2-18.5	1534	24.2	22.9-25.5	1235	13.5	12.8-14.3	631	4.1	3.8-4.4	334	5.6	5.0-6.2	297	3.2	2.8-3.6
1989	2762	17.8	17.1-18.5	1495	23.2	22.0-24.4	1267	13.9	13.1-14.7	589	3.8	3.5-4.2	317	5.2	4.6-5.8	272	2.9	2.5-3.3
1990	2705	17.3	16.7-18.0	1455	22.4	21.3-23.6	1250	13.5	12.7-14.3	534	3.4	3.2-3.8	272	4.4	3.9-5.0	262	2.8	2.5-3.2
1991	2595	16.7	16.1-17.4	1425	22.4	21.2-23.6	1170	12.8	12.1-13.6	622	4.0	3.7-4.3	322	5.3	4.7-5.9	300	3.2	2.9-3.6
1992	2615	16.8	16.1-17.4	1443	22.4	21.3-23.6	1172	12.7	12.0-13.5	575	3.7	3.4-4.0	316	5.2	4.6-5.8	259	2.7	2.4-3.1
1993	2548	16.3	15.7-16.9	1392	21.5	20.3-22.7	1156	12.6	11.9-13.3	548	3.5	3.2-3.8	285	4.5	4.0-5.1	263	2.7	2.4-3.1
1994	2646	16.9	16.3-17.6	1387	21.4	20.2-22.5	1259	13.8	13.0-14.6	550	3.5	3.2-3.8	290	4.7	4.2-5.3	260	2.7	2.4-3.1
1995	2565	16.5	15.8-17.1	1368	21.1	20.0-22.3	1197	13.2	12.4-13.9	605	3.8	3.5-4.2	318	5.1	4.5-5.7	287	3.0	2.7-3.4
1996	2594	16.7	16.0-17.3	1444	22.1	21.0-23.3	1150	12.7	11.9-13.4	542	3.4	3.1-3.7	275	4.3	3.8-4.9	267	2.8	2.5-3.2
1997	2531	16.1	15.5-16.8	1386	21.0	19.9-22.1	1145	12.5	11.8-13.2	491	3.1	2.8-3.4	278	4.4	3.9-5.0	213	2.2	1.9-2.5
1998	2761	17.5	16.8-18.2	1477	22.0	20.9-23.2	1284	14.0	13.2-14.8	511	3.2	2.9-3.5	263	4.0	3.5-4.5	248	2.6	2.3-2.9
1999	2774	17.5	16.8-18.1	1510	22.4	21.3-23.6	1264	13.8	13.0-14.6	548	3.4	3.1-3.7	286	4.4	3.9-4.9	262	2.7	2.4-3.0
2000	2739	17.1	16.5-17.8	1544	22.6	21.5-23.8	1195	12.9	12.2-13.7	527	3.2	3.0-3.5	283	4.2	3.8-4.8	244	2.5	2.2-2.8
2001	2612	16.2	15.6-16.9	1413	20.6	19.6-21.7	1199	13.1	12.3-13.8	522	3.2	2.9-3.5	278	4.2	3.7-4.7	244	2.5	2.2-2.8
2002	2589	16.0	15.4-16.6	1426	20.4	19.3-21.5	1163	12.4	11.7-13.1	562	3.4	3.1-3.7	287	4.2	3.8-4.8	275	2.8	2.5-3.2
2003	2499	15.3	14.7-15.9	1358	19.3	18.2-20.3	1141	12.2	11.5-13.0	472	2.8	2.6-3.1	254	3.7	3.2-4.2	218	2.2	1.9-2.6
2004	2482	15.1	14.5-15.7	1331	18.7	17.7-19.7	1151	12.4	11.7-13.1	487	2.9	2.7-3.2	263	3.8	3.3-4.3	224	2.2	1.9-2.5
2005	2255	13.6	13.1-14.2	1253	17.6	16.6-18.6	1002	10.7	10.0-11.4	467	2.8	2.5-3.0	249	3.6	3.1-4.1	218	2.1	1.9-2.5
2006	2202	13.4	12.8-14.0	1203	16.7	15.7-17.6	999	10.8	10.1-11.5	438	2.6	2.4-2.9	234	3.4	2.9-3.8	204	2.1	1.8-2.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Rectum & rectosigmoid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	2248	13.5	13.0-14.1	1224	16.7	15.7-17.6	1024	11.1	10.4-11.8	413	2.4	2.2-2.7	222	3.1	2.7-3.5	191	1.9	1.6-2.2
2008	2192	13.1	12.5-13.7	1214	16.5	15.5-17.4	978	10.4	9.7-11.1	491	2.8	2.6-3.1	255	3.5	3.1-4.0	236	2.3	2.0-2.6
2009	2144	12.8	12.2-13.3	1211	16.2	15.3-17.1	933	10.0	9.4-10.7	446	2.6	2.3-2.8	235	3.2	2.8-3.6	211	2.1	1.8-2.4
2010	2129	12.5	12.0-13.1	1172	15.3	14.5-16.3	957	10.3	9.6-11.0	450	2.6	2.4-2.9	259	3.5	3.1-3.9	191	1.9	1.6-2.2
2011	2049	12.0	11.5-12.5	1123	14.6	13.7-15.5	926	9.9	9.3-10.6	452	2.5	2.3-2.8	253	3.3	2.9-3.8	199	1.9	1.6-2.2
2012	1923	11.2	10.7-11.7	1136	14.5	13.6-15.4	787	8.5	7.9-9.1	429	2.4	2.2-2.6	238	3.1	2.7-3.5	191	1.9	1.6-2.2
2013	2078	11.9	11.4-12.5	1179	14.9	14.0-15.8	899	9.5	8.9-10.2	439	2.4	2.2-2.7	254	3.2	2.8-3.6	185	1.8	1.5-2.1
2014	1992	11.3	10.8-11.9	1150	14.2	13.4-15.0	842	8.9	8.3-9.6	454	2.5	2.2-2.7	253	3.1	2.7-3.5	201	1.9	1.7-2.2
2015	2010	11.3	10.8-11.8	1142	13.9	13.1-14.7	868	9.1	8.5-9.7	442	2.4	2.2-2.6	255	3.2	2.8-3.6	187	1.8	1.5-2.0
2016	1927	10.9	10.4-11.4	1116	13.6	12.8-14.4	811	8.6	8.0-9.2	472	2.5	2.3-2.8	262	3.2	2.8-3.6	210	2.0	1.7-2.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Liver / intrahepatic bile duct

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	458	3.0	2.8-3.3	260	4.2	3.6-4.7	198	2.2	1.9-2.6	408	2.7	2.4-3.0	229	3.6	3.1-4.1	179	2.1	1.8-2.4
1977	494	3.2	2.9-3.5	317	4.9	4.3-5.4	177	2.0	1.7-2.3	493	3.2	2.9-3.5	275	4.2	3.7-4.8	218	2.5	2.1-2.8
1978	455	3.0	2.7-3.2	261	4.1	3.6-4.6	194	2.2	1.9-2.5	467	3.0	2.8-3.3	274	4.3	3.8-4.9	193	2.2	1.9-2.5
1979	443	2.9	2.7-3.2	248	3.9	3.4-4.4	195	2.3	2.0-2.6	450	3.0	2.7-3.3	241	3.9	3.4-4.5	209	2.4	2.1-2.7
1980	451	3.0	2.7-3.3	261	4.3	3.8-4.9	190	2.1	1.8-2.5	432	2.8	2.6-3.1	222	3.6	3.2-4.2	210	2.3	2.0-2.7
1981	460	3.0	2.8-3.3	258	4.2	3.7-4.8	202	2.2	1.9-2.6	497	3.3	3.0-3.6	273	4.5	4.0-5.1	224	2.5	2.2-2.8
1982	429	2.8	2.5-3.1	276	4.5	3.9-5.1	153	1.7	1.4-2.0	461	3.0	2.7-3.3	280	4.6	4.0-5.2	181	2.0	1.7-2.3
1983	472	3.1	2.8-3.4	300	4.7	4.2-5.3	172	2.0	1.7-2.3	482	3.1	2.9-3.4	285	4.5	4.0-5.1	197	2.2	1.9-2.5
1984	461	3.0	2.7-3.3	269	4.2	3.7-4.8	192	2.1	1.8-2.5	477	3.1	2.8-3.4	266	4.3	3.7-4.8	211	2.3	2.0-2.6
1985	466	3.1	2.8-3.4	287	4.7	4.2-5.3	179	2.0	1.7-2.3	502	3.3	3.0-3.6	302	5.0	4.4-5.6	200	2.2	1.9-2.5
1986	497	3.3	3.0-3.6	312	5.0	4.5-5.6	185	2.1	1.8-2.4	519	3.4	3.1-3.7	293	4.7	4.2-5.3	226	2.5	2.2-2.9
1987	474	3.1	2.8-3.4	294	4.6	4.1-5.2	180	2.0	1.7-2.3	539	3.5	3.2-3.9	312	5.2	4.6-5.8	227	2.5	2.2-2.8
1988	557	3.6	3.3-4.0	351	5.6	5.0-6.2	206	2.3	2.0-2.6	555	3.6	3.3-3.9	329	5.3	4.7-5.9	226	2.5	2.1-2.8
1989	502	3.2	3.0-3.5	314	4.9	4.3-5.4	188	2.0	1.8-2.4	484	3.1	2.8-3.4	286	4.5	3.9-5.0	198	2.1	1.8-2.4
1990	589	3.8	3.5-4.2	369	5.9	5.3-6.6	220	2.4	2.1-2.7	584	3.8	3.5-4.1	346	5.7	5.1-6.4	238	2.5	2.2-2.9
1991	599	3.9	3.6-4.2	369	5.6	5.0-6.2	230	2.5	2.2-2.9	585	3.7	3.4-4.0	347	5.4	4.8-6.0	238	2.5	2.2-2.9
1992	654	4.2	3.9-4.6	385	6.1	5.5-6.8	269	3.0	2.6-3.4	620	4.0	3.7-4.3	370	5.9	5.3-6.6	250	2.7	2.4-3.1
1993	654	4.2	3.9-4.5	402	6.3	5.6-6.9	252	2.7	2.4-3.1	666	4.2	3.9-4.6	379	5.9	5.3-6.6	287	3.0	2.7-3.4
1994	651	4.2	3.9-4.5	417	6.5	5.9-7.2	234	2.5	2.2-2.9	604	3.9	3.6-4.2	362	5.8	5.2-6.4	242	2.6	2.3-3.0
1995	734	4.7	4.4-5.1	469	7.2	6.5-7.9	265	2.9	2.5-3.2	667	4.2	3.9-4.6	406	6.3	5.7-7.0	261	2.7	2.4-3.1
1996	723	4.7	4.3-5.0	467	7.2	6.5-7.9	256	2.8	2.4-3.1	645	4.1	3.8-4.4	391	6.0	5.4-6.7	254	2.6	2.3-3.0
1997	805	5.2	4.9-5.6	520	7.9	7.2-8.6	285	3.1	2.7-3.4	633	4.0	3.7-4.4	388	5.9	5.4-6.6	245	2.5	2.2-2.9
1998	898	5.7	5.4-6.1	589	8.8	8.1-9.5	309	3.3	3.0-3.7	660	4.2	3.9-4.5	402	6.0	5.5-6.7	258	2.7	2.4-3.1
1999	862	5.5	5.1-5.8	567	8.4	7.7-9.1	295	3.1	2.8-3.5	689	4.3	4.0-4.7	420	6.3	5.7-6.9	269	2.8	2.5-3.2
2000	857	5.4	5.0-5.8	567	8.2	7.6-9.0	290	3.1	2.8-3.5	684	4.2	3.9-4.6	422	6.2	5.6-6.8	262	2.7	2.4-3.1
2001	910	5.7	5.3-6.1	629	9.1	8.4-9.8	281	3.0	2.7-3.4	710	4.4	4.1-4.7	455	6.7	6.1-7.3	255	2.6	2.3-3.0
2002	900	5.6	5.2-6.0	629	8.9	8.2-9.7	271	2.9	2.6-3.3	772	4.7	4.4-5.1	498	7.2	6.6-7.9	274	2.8	2.5-3.1
2003	974	6.0	5.7-6.4	636	8.9	8.2-9.6	338	3.7	3.3-4.1	713	4.3	4.0-4.7	447	6.4	5.8-7.0	266	2.7	2.4-3.1
2004	1013	6.2	5.8-6.6	725	10.0	9.3-10.8	288	3.1	2.7-3.5	779	4.7	4.4-5.0	481	6.7	6.1-7.3	298	3.0	2.7-3.4
2005	1076	6.6	6.2-7.0	760	10.4	9.6-11.1	316	3.4	3.0-3.8	798	4.8	4.4-5.1	517	7.1	6.5-7.8	281	2.8	2.5-3.2
2006	1123	6.8	6.4-7.2	789	10.6	9.9-11.4	334	3.6	3.2-4.0	842	5.0	4.7-5.4	520	7.2	6.6-7.9	322	3.3	3.0-3.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Liver / intrahepatic bile duct

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	1180	7.1	6.7-7.5	852	11.3	10.5-12.0	328	3.5	3.1-3.9	825	4.9	4.6-5.2	550	7.5	6.9-8.1	275	2.8	2.5-3.2
2008	1224	7.2	6.8-7.6	871	11.3	10.6-12.1	353	3.7	3.4-4.2	884	5.2	4.8-5.5	579	7.7	7.1-8.3	305	3.1	2.7-3.4
2009	1223	7.1	6.7-7.5	856	11.0	10.3-11.8	367	3.8	3.4-4.2	909	5.3	4.9-5.6	594	7.9	7.2-8.5	315	3.2	2.8-3.6
2010	1214	7.0	6.6-7.4	907	11.5	10.7-12.3	307	3.2	2.8-3.6	916	5.2	4.9-5.6	598	7.7	7.1-8.4	318	3.2	2.8-3.5
2011	1261	7.1	6.7-7.5	889	10.8	10.1-11.6	372	3.9	3.5-4.3	935	5.2	4.9-5.6	590	7.4	6.8-8.0	345	3.4	3.0-3.7
2012	1377	7.6	7.2-8.0	998	12.0	11.2-12.8	379	3.9	3.5-4.3	997	5.5	5.2-5.9	640	8.0	7.3-8.6	357	3.5	3.1-3.9
2013	1445	7.9	7.4-8.3	1008	12.0	11.2-12.8	437	4.4	4.0-4.8	1024	5.5	5.2-5.9	689	8.4	7.8-9.1	335	3.2	2.9-3.6
2014	1480	8.0	7.6-8.4	1053	12.3	11.5-13.1	427	4.3	3.8-4.7	991	5.3	5.0-5.6	668	7.8	7.2-8.5	323	3.2	2.8-3.5
2015	1476	7.8	7.4-8.2	1048	12.1	11.4-12.9	428	4.2	3.8-4.6	1037	5.5	5.1-5.8	679	8.1	7.4-8.7	358	3.3	3.0-3.7
2016	1500	7.9	7.5-8.3	1081	12.3	11.5-13.0	419	4.1	3.7-4.6	1056	5.5	5.2-5.9	691	8.0	7.4-8.7	365	3.4	3.1-3.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Pancreas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1872	12.4	11.8-13.0	976	15.7	14.7-16.8	896	10.2	9.5-10.9	1760	11.7	11.2-12.3	887	14.4	13.4-15.5	873	9.9	9.2-10.6
1977	1924	12.9	12.3-13.5	976	16.3	15.2-17.4	948	10.7	10.0-11.4	1850	12.4	11.8-13.0	928	15.3	14.3-16.4	922	10.4	9.8-11.1
1978	1898	12.5	11.9-13.1	967	15.4	14.4-16.4	931	10.4	9.8-11.1	1801	11.8	11.3-12.4	918	14.7	13.8-15.8	883	9.9	9.2-10.5
1979	1970	13.0	12.4-13.6	1008	16.3	15.3-17.4	962	10.6	10.0-11.4	1892	12.5	11.9-13.0	966	15.8	14.8-16.9	926	10.2	9.5-10.9
1980	1969	12.9	12.3-13.5	966	15.6	14.6-16.7	1003	11.0	10.4-11.8	1832	12.0	11.5-12.6	923	15.0	14.0-16.1	909	10.0	9.3-10.7
1981	1886	12.4	11.9-13.0	948	15.4	14.4-16.4	938	10.4	9.8-11.1	1803	11.9	11.3-12.4	883	14.4	13.4-15.4	920	10.1	9.5-10.8
1982	1915	12.5	12.0-13.1	900	14.9	13.9-15.9	1015	11.0	10.4-11.7	1701	11.1	10.6-11.6	837	13.7	12.7-14.7	864	9.4	8.7-10.0
1983	1820	11.8	11.3-12.4	881	14.1	13.1-15.1	939	10.2	9.6-10.9	1783	11.5	11.0-12.1	868	13.7	12.8-14.7	915	9.9	9.3-10.6
1984	1892	12.3	11.7-12.8	900	14.8	13.8-15.8	992	10.6	10.0-11.3	1758	11.4	10.9-11.9	826	13.5	12.5-14.5	932	10.0	9.4-10.7
1985	1943	12.5	12.0-13.1	902	14.3	13.3-15.3	1041	11.2	10.5-11.9	1718	11.1	10.6-11.6	811	13.2	12.2-14.1	907	9.8	9.1-10.4
1986	1895	12.4	11.8-12.9	891	14.4	13.4-15.4	1004	10.8	10.2-11.6	1745	11.2	10.7-11.8	822	13.1	12.2-14.1	923	9.8	9.2-10.5
1987	1893	12.2	11.6-12.7	870	14.1	13.2-15.1	1023	10.8	10.2-11.5	1700	11.0	10.4-11.5	817	13.2	12.3-14.1	883	9.4	8.7-10.0
1988	1923	12.4	11.9-13.0	891	14.4	13.4-15.4	1032	11.0	10.3-11.7	1738	11.2	10.6-11.7	825	13.4	12.5-14.4	913	9.6	9.0-10.2
1989	1838	11.8	11.3-12.4	875	14.0	13.0-15.0	963	10.4	9.7-11.0	1680	10.8	10.3-11.4	813	13.1	12.2-14.1	867	9.3	8.6-9.9
1990	1888	12.2	11.6-12.7	890	14.3	13.3-15.3	998	10.7	10.0-11.4	1695	10.9	10.4-11.4	794	12.8	11.9-13.7	901	9.6	8.9-10.2
1991	1948	12.5	12.0-13.1	909	14.4	13.5-15.4	1039	11.1	10.4-11.8	1684	10.8	10.3-11.4	789	12.7	11.8-13.6	895	9.5	8.9-10.1
1992	1921	12.4	11.8-12.9	909	14.6	13.7-15.6	1012	10.8	10.1-11.5	1680	10.8	10.2-11.3	766	12.3	11.4-13.2	914	9.6	8.9-10.2
1993	1919	12.3	11.7-12.8	901	14.2	13.2-15.1	1018	10.8	10.1-11.5	1798	11.4	10.9-12.0	860	13.6	12.7-14.5	938	9.9	9.2-10.5
1994	2048	13.1	12.5-13.6	977	15.3	14.3-16.3	1071	11.2	10.5-11.9	1856	11.8	11.3-12.3	868	13.7	12.8-14.7	988	10.3	9.6-10.9
1995	1948	12.3	11.8-12.9	868	13.5	12.6-14.5	1080	11.4	10.7-12.1	1790	11.3	10.8-11.8	821	12.7	11.9-13.7	969	10.0	9.4-10.7
1996	1930	12.3	11.7-12.8	942	14.8	13.8-15.7	988	10.5	9.9-11.2	1815	11.5	11.0-12.1	851	13.5	12.6-14.4	964	10.1	9.5-10.8
1997	1988	12.6	12.0-13.1	943	14.5	13.6-15.5	1045	11.0	10.4-11.7	1811	11.4	10.9-11.9	871	13.7	12.8-14.6	940	9.8	9.1-10.4
1998	2089	13.1	12.6-13.7	1000	15.3	14.3-16.2	1089	11.5	10.8-12.2	1797	11.2	10.7-11.8	861	13.3	12.4-14.2	936	9.8	9.1-10.4
1999	2054	12.8	12.2-13.3	982	14.8	13.9-15.8	1072	11.2	10.5-11.9	1867	11.6	11.1-12.1	927	14.0	13.1-14.9	940	9.8	9.1-10.4
2000	2112	13.0	12.5-13.6	982	14.7	13.8-15.6	1130	11.8	11.1-12.5	1822	11.1	10.6-11.7	831	12.5	11.7-13.4	991	10.1	9.4-10.7
2001	2073	12.7	12.1-13.2	938	13.8	12.9-14.7	1135	11.7	11.0-12.4	1859	11.3	10.8-11.8	849	12.7	11.9-13.6	1010	10.2	9.6-10.9
2002	2114	12.8	12.3-13.4	991	14.4	13.5-15.3	1123	11.5	10.8-12.2	1829	11.1	10.5-11.6	850	12.4	11.6-13.3	979	9.9	9.3-10.5
2003	2199	13.2	12.7-13.8	1045	15.1	14.2-16.0	1154	11.8	11.1-12.5	1829	10.9	10.5-11.5	855	12.4	11.6-13.3	974	9.8	9.1-10.4
2004	2175	13.0	12.5-13.6	1024	14.6	13.7-15.6	1151	11.8	11.1-12.5	1863	11.1	10.6-11.6	906	13.0	12.2-13.9	957	9.7	9.0-10.3
2005	2182	13.0	12.5-13.6	1072	15.1	14.2-16.0	1110	11.3	10.7-12.0	1899	11.2	10.7-11.8	900	12.8	12.0-13.7	999	10.0	9.4-10.7
2006	2235	13.3	12.7-13.9	1059	14.9	14.0-15.8	1176	12.0	11.3-12.7	1900	11.2	10.7-11.7	914	12.9	12.1-13.8	986	9.8	9.2-10.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Pancreas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	2323	13.6	13.1-14.2	1133	15.7	14.8-16.6	1190	12.0	11.3-12.7	1908	11.1	10.6-11.6	922	12.8	12.0-13.7	986	9.9	9.3-10.5
2008	2451	14.3	13.8-14.9	1195	16.4	15.5-17.4	1256	12.6	11.9-13.4	1987	11.5	11.0-12.0	938	12.9	12.1-13.8	1049	10.2	9.6-10.9
2009	2448	14.1	13.6-14.7	1251	16.8	15.9-17.8	1197	12.0	11.3-12.7	2011	11.5	11.0-12.0	1015	13.7	12.9-14.6	996	9.8	9.2-10.5
2010	2467	14.2	13.6-14.7	1192	15.9	15.0-16.8	1275	12.7	12.0-13.5	2039	11.5	11.0-12.0	987	13.0	12.2-13.9	1052	10.1	9.5-10.8
2011	2491	14.1	13.5-14.7	1257	16.4	15.5-17.4	1234	12.2	11.5-12.9	2015	11.3	10.8-11.8	986	13.0	12.1-13.8	1029	9.9	9.3-10.5
2012	2511	14.1	13.5-14.6	1272	16.4	15.5-17.4	1239	12.2	11.5-12.9	2072	11.5	11.0-12.0	1066	13.8	12.9-14.6	1006	9.7	9.1-10.3
2013	2551	14.0	13.4-14.5	1276	16.0	15.1-16.9	1275	12.3	11.6-13.0	2058	11.2	10.7-11.7	1028	13.0	12.2-13.9	1030	9.7	9.1-10.3
2014	2665	14.5	13.9-15.1	1365	16.8	15.9-17.7	1300	12.5	11.8-13.3	2094	11.2	10.7-11.7	1066	13.3	12.5-14.1	1028	9.5	8.9-10.1
2015	2633	14.1	13.6-14.7	1326	16.1	15.2-17.0	1307	12.4	11.8-13.2	1998	10.5	10.1-11.0	1004	12.3	11.5-13.1	994	9.1	8.5-9.7
2016	2690	14.4	13.8-15.0	1364	16.4	15.5-17.3	1326	12.8	12.1-13.5	2155	11.4	10.9-11.9	1066	12.8	12.1-13.6	1089	10.2	9.6-10.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Larynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	855	5.3	4.9-5.7	718	10.5	9.7-11.3	137	1.5	1.3-1.8	312	2.0	1.8-2.2	253	3.9	3.4-4.4	59	0.7	0.5-0.9
1977	843	5.3	5.0-5.7	701	10.4	9.6-11.2	142	1.6	1.3-1.9	309	1.9	1.7-2.2	266	4.0	3.6-4.6	43	0.5	0.3-0.6
1978	861	5.4	5.1-5.8	706	10.6	9.8-11.4	155	1.7	1.5-2.0	312	2.0	1.8-2.2	247	3.8	3.3-4.4	65	0.7	0.6-0.9
1979	920	5.7	5.4-6.1	734	10.7	9.9-11.5	186	2.1	1.8-2.5	340	2.1	1.9-2.4	276	4.2	3.7-4.8	64	0.7	0.5-0.9
1980	944	6.0	5.6-6.4	774	11.6	10.8-12.5	170	2.0	1.7-2.3	306	1.9	1.7-2.2	252	3.9	3.4-4.5	54	0.6	0.5-0.8
1981	969	6.2	5.8-6.6	784	11.8	10.9-12.6	185	2.1	1.8-2.4	298	1.9	1.6-2.1	248	3.8	3.3-4.3	50	0.5	0.4-0.7
1982	893	5.7	5.3-6.1	714	10.7	9.9-11.6	179	2.1	1.8-2.4	312	1.9	1.7-2.2	249	3.8	3.3-4.3	63	0.7	0.5-0.9
1983	881	5.7	5.3-6.1	694	10.3	9.5-11.1	187	2.2	1.9-2.6	292	1.8	1.6-2.1	222	3.4	3.0-3.9	70	0.8	0.6-1.0
1984	929	5.9	5.5-6.3	729	10.8	10.0-11.7	200	2.3	2.0-2.7	268	1.7	1.5-1.9	211	3.2	2.8-3.7	57	0.6	0.5-0.8
1985	934	6.0	5.6-6.4	732	10.8	10.0-11.7	202	2.4	2.0-2.7	296	1.8	1.6-2.1	232	3.5	3.0-4.0	64	0.7	0.5-0.9
1986	882	5.7	5.3-6.1	681	10.3	9.5-11.1	201	2.4	2.0-2.7	248	1.6	1.4-1.8	197	3.1	2.7-3.6	51	0.5	0.4-0.7
1987	904	5.9	5.5-6.3	719	10.9	10.1-11.8	185	2.2	1.9-2.5	275	1.8	1.6-2.0	216	3.4	2.9-3.9	59	0.6	0.5-0.8
1988	856	5.5	5.2-5.9	661	9.8	9.1-10.6	195	2.3	2.0-2.6	267	1.7	1.5-2.0	211	3.3	2.9-3.8	56	0.6	0.5-0.8
1989	895	5.9	5.5-6.3	687	10.5	9.7-11.3	208	2.5	2.2-2.9	267	1.7	1.5-1.9	186	2.8	2.4-3.3	81	0.9	0.7-1.1
1990	922	6.0	5.6-6.4	712	10.7	9.9-11.5	210	2.5	2.1-2.8	237	1.6	1.4-1.8	196	3.1	2.7-3.6	41	0.4	0.3-0.6
1991	839	5.5	5.2-5.9	674	10.2	9.5-11.1	165	1.9	1.6-2.3	256	1.6	1.4-1.9	198	3.0	2.6-3.5	58	0.6	0.5-0.8
1992	856	5.6	5.3-6.0	651	9.8	9.1-10.6	205	2.5	2.1-2.8	314	2.0	1.8-2.3	239	3.8	3.3-4.3	75	0.8	0.7-1.1
1993	810	5.3	4.9-5.7	603	9.2	8.4-9.9	207	2.4	2.1-2.8	267	1.7	1.5-1.9	212	3.3	2.8-3.8	55	0.6	0.5-0.8
1994	780	5.1	4.8-5.5	601	9.2	8.4-9.9	179	2.1	1.8-2.4	264	1.7	1.5-1.9	203	3.2	2.7-3.6	61	0.7	0.5-0.9
1995	758	5.0	4.6-5.3	581	8.7	8.0-9.4	177	2.1	1.8-2.4	255	1.7	1.5-1.9	195	3.0	2.6-3.4	60	0.7	0.5-0.8
1996	765	5.0	4.7-5.4	595	8.9	8.2-9.7	170	2.0	1.7-2.4	264	1.7	1.5-1.9	205	3.1	2.7-3.6	59	0.7	0.5-0.8
1997	738	4.9	4.5-5.2	586	8.8	8.1-9.5	152	1.8	1.5-2.1	216	1.4	1.2-1.6	161	2.5	2.1-2.9	55	0.6	0.5-0.8
1998	708	4.6	4.2-4.9	521	7.7	7.0-8.4	187	2.2	1.9-2.5	234	1.5	1.3-1.7	178	2.7	2.3-3.1	56	0.6	0.5-0.8
1999	735	4.7	4.4-5.1	589	8.5	7.9-9.3	146	1.7	1.4-2.0	232	1.5	1.3-1.7	176	2.6	2.2-3.0	56	0.6	0.5-0.8
2000	734	4.7	4.3-5.0	571	8.3	7.6-9.0	163	1.9	1.6-2.2	225	1.4	1.2-1.6	181	2.7	2.3-3.1	44	0.5	0.4-0.7
2001	697	4.4	4.1-4.7	536	7.7	7.0-8.4	161	1.8	1.6-2.2	199	1.2	1.1-1.4	163	2.4	2.0-2.8	36	0.4	0.3-0.5
2002	671	4.2	3.9-4.5	533	7.6	7.0-8.3	138	1.6	1.3-1.9	223	1.4	1.2-1.5	176	2.6	2.2-3.0	47	0.5	0.4-0.7
2003	677	4.2	3.9-4.5	526	7.4	6.7-8.0	151	1.7	1.4-2.0	207	1.3	1.1-1.4	156	2.2	1.9-2.6	51	0.5	0.4-0.7
2004	682	4.2	3.9-4.5	554	7.6	7.0-8.2	128	1.4	1.2-1.7	197	1.2	1.0-1.4	159	2.3	1.9-2.6	38	0.4	0.3-0.5
2005	719	4.4	4.1-4.7	560	7.6	7.0-8.3	159	1.8	1.5-2.1	209	1.3	1.1-1.5	155	2.2	1.9-2.6	54	0.6	0.4-0.8
2006	676	4.1	3.8-4.5	522	7.2	6.6-7.8	154	1.8	1.5-2.1	207	1.2	1.1-1.4	161	2.3	1.9-2.6	46	0.5	0.4-0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Larynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	640	3.8	3.5-4.1	503	6.8	6.2-7.4	137	1.5	1.2-1.7	189	1.1	1.0-1.3	154	2.1	1.8-2.4	35	0.4	0.2-0.5
2008	722	4.3	4.0-4.6	566	7.5	6.9-8.2	156	1.7	1.5-2.0	173	1.0	0.9-1.2	140	1.9	1.6-2.2	33	0.3	0.2-0.5
2009	656	3.9	3.6-4.2	506	6.6	6.1-7.2	150	1.6	1.4-1.9	196	1.1	1.0-1.3	145	1.9	1.6-2.3	51	0.5	0.4-0.7
2010	632	3.7	3.4-4.0	502	6.5	5.9-7.1	130	1.4	1.2-1.7	204	1.1	1.0-1.3	153	2.0	1.7-2.4	51	0.5	0.4-0.7
2011	593	3.3	3.1-3.6	481	5.9	5.4-6.5	112	1.2	1.0-1.5	192	1.1	0.9-1.3	144	1.9	1.6-2.2	48	0.5	0.3-0.6
2012	653	3.7	3.4-4.0	521	6.5	5.9-7.1	132	1.4	1.1-1.6	174	1.0	0.8-1.1	135	1.7	1.4-2.0	39	0.4	0.3-0.6
2013	648	3.6	3.3-3.8	523	6.3	5.8-6.9	125	1.3	1.1-1.6	209	1.1	1.0-1.3	170	2.1	1.8-2.4	39	0.4	0.3-0.6
2014	614	3.4	3.1-3.6	478	5.7	5.2-6.3	136	1.4	1.2-1.7	179	1.0	0.8-1.1	143	1.7	1.5-2.0	36	0.4	0.2-0.5
2015	625	3.4	3.1-3.7	494	5.8	5.3-6.4	131	1.3	1.1-1.6	165	0.9	0.7-1.0	126	1.5	1.2-1.8	39	0.4	0.3-0.5
2016	559	3.0	2.8-3.3	455	5.4	4.9-5.9	104	1.1	0.9-1.3	181	1.0	0.8-1.1	151	1.8	1.5-2.1	30	0.3	0.2-0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Lung and bronchus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	8487	54.1	52.9-55.3	6057	92.6	90.2-95.1	2430	27.5	26.4-28.7	6853	43.9	42.8-44.9	4988	77.5	75.2-79.8	1865	21.1	20.1-22.1
1977	8485	54.1	52.9-55.3	5960	91.6	89.2-94.0	2525	28.6	27.5-29.8	7129	45.6	44.6-46.7	5157	80.1	77.9-82.4	1972	22.3	21.3-23.3
1978	8636	55.2	54.0-56.4	6146	94.7	92.2-97.2	2490	28.2	27.1-29.4	7125	45.7	44.6-46.8	5103	79.7	77.4-82.0	2022	22.8	21.8-23.8
1979	8837	56.5	55.3-57.7	6058	93.7	91.3-96.2	2779	31.6	30.4-32.8	7070	45.3	44.2-46.3	4958	77.5	75.3-79.8	2112	23.7	22.7-24.8
1980	9136	58.2	57.0-59.4	6284	96.6	94.1-99.1	2852	32.1	30.9-33.3	7557	48.2	47.1-49.4	5268	82.0	79.7-84.4	2289	25.6	24.5-26.7
1981	9071	58.0	56.8-59.2	6073	94.3	91.8-96.8	2998	33.9	32.7-35.2	7423	47.5	46.4-48.6	5159	80.7	78.4-83.0	2264	25.4	24.4-26.5
1982	9468	60.5	59.2-61.7	6229	96.7	94.2-99.2	3239	36.5	35.2-37.8	7622	48.8	47.7-49.9	5096	80.1	77.9-82.4	2526	28.3	27.2-29.5
1983	9761	62.2	60.9-63.4	6321	97.6	95.1-100.1	3440	38.5	37.2-39.9	7763	49.5	48.4-50.6	5066	79.7	77.5-82.0	2697	29.9	28.7-31.0
1984	9708	62.1	60.9-63.4	6274	96.8	94.3-99.3	3434	38.9	37.6-40.3	7751	49.5	48.4-50.6	5095	79.6	77.3-81.9	2656	29.6	28.4-30.8
1985	10028	63.9	62.7-65.2	6355	97.8	95.3-100.3	3673	41.4	40.1-42.8	8100	51.7	50.5-52.8	5199	81.4	79.1-83.7	2901	32.2	31.0-33.5
1986	9924	63.6	62.4-64.9	6200	96.0	93.5-98.5	3724	42.0	40.7-43.4	8051	51.3	50.2-52.4	5081	79.4	77.2-81.7	2970	33.2	32.0-34.5
1987	10159	65.0	63.7-66.3	6268	96.6	94.2-99.1	3891	43.9	42.5-45.3	8098	51.8	50.6-52.9	5135	80.5	78.2-82.8	2963	32.9	31.7-34.1
1988	10264	65.7	64.4-67.0	6203	96.4	93.9-98.9	4061	45.7	44.2-47.1	8246	52.7	51.6-53.9	5146	80.6	78.3-82.9	3100	34.6	33.4-35.9
1989	10370	66.7	65.4-68.0	6161	95.0	92.6-97.5	4209	47.7	46.2-49.2	8239	52.9	51.7-54.0	5046	79.0	76.7-81.2	3193	35.7	34.4-37.0
1990	10422	66.9	65.6-68.2	6207	96.1	93.6-98.6	4215	47.4	46.0-48.9	8367	53.5	52.4-54.7	5022	78.4	76.2-80.7	3345	37.1	35.8-38.4
1991	10563	68.1	66.8-69.4	6106	94.6	92.1-97.0	4457	50.4	48.9-51.9	8594	55.1	54.0-56.3	5088	79.7	77.4-81.9	3506	39.0	37.7-40.3
1992	10720	68.8	67.5-70.1	6167	94.9	92.5-97.4	4553	51.3	49.8-52.9	8567	54.8	53.6-56.0	4988	77.9	75.7-80.2	3579	39.6	38.3-40.9
1993	10593	68.0	66.7-69.4	5933	91.2	88.8-93.6	4660	52.6	51.1-54.1	8419	53.8	52.6-54.9	4832	75.3	73.1-77.4	3587	39.8	38.5-41.1
1994	10854	69.6	68.3-70.9	6010	92.0	89.7-94.4	4844	54.8	53.2-56.4	8662	55.2	54.1-56.4	5075	78.8	76.6-81.0	3587	39.6	38.3-41.0
1995	10739	68.7	67.4-70.0	5979	90.9	88.6-93.3	4760	53.3	51.8-54.9	8598	54.7	53.6-55.9	4904	75.7	73.5-77.9	3694	40.8	39.4-42.1
1996	10890	69.8	68.5-71.1	6047	92.2	89.9-94.6	4843	54.6	53.1-56.2	8436	53.8	52.7-55.0	4782	73.6	71.5-75.8	3654	40.4	39.1-41.7
1997	10926	69.8	68.5-71.1	5940	90.0	87.7-92.3	4986	56.1	54.5-57.7	8198	52.0	50.9-53.2	4587	70.3	68.3-72.4	3611	39.7	38.4-41.0
1998	11022	69.9	68.6-71.2	5903	88.1	85.9-90.4	5119	57.2	55.6-58.8	8248	52.0	50.9-53.1	4582	69.0	67.0-71.1	3666	40.1	38.8-41.4
1999	10985	69.1	67.8-70.4	5838	86.7	84.4-88.9	5147	57.1	55.5-58.7	8174	51.1	50.0-52.2	4450	66.8	64.8-68.8	3724	40.6	39.3-41.9
2000	11261	70.2	68.9-71.5	5935	87.3	85.1-89.6	5326	58.5	56.9-60.1	8293	51.3	50.2-52.4	4513	67.1	65.1-69.1	3780	40.4	39.1-41.7
2001	11018	68.3	67.1-69.6	5674	83.0	80.9-85.2	5344	58.5	56.9-60.1	8204	50.5	49.4-51.6	4447	65.9	63.9-67.9	3757	40.2	38.9-41.5
2002	11222	69.0	67.7-70.3	5893	85.2	83.0-87.4	5329	58.0	56.4-59.6	8280	50.6	49.5-51.7	4411	64.3	62.4-66.3	3869	41.2	39.9-42.5
2003	11181	68.4	67.1-69.7	5880	84.3	82.1-86.4	5301	57.4	55.8-59.0	8067	49.0	47.9-50.0	4301	62.1	60.3-64.0	3766	39.8	38.6-41.1
2004	11172	67.9	66.6-69.2	5715	81.2	79.1-83.3	5457	58.7	57.2-60.3	7967	48.1	47.0-49.1	4160	59.6	57.8-61.5	3807	40.1	38.8-41.4
2005	11232	68.1	66.9-69.4	5750	81.3	79.2-83.5	5482	59.0	57.4-60.6	7880	47.5	46.4-48.5	4174	59.7	57.9-61.5	3706	39.0	37.7-40.3
2006	11453	69.5	68.3-70.8	5683	80.0	77.9-82.1	5770	62.5	60.9-64.2	7850	47.2	46.2-48.3	4183	59.3	57.5-61.1	3667	38.7	37.5-40.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Lung and bronchus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	11500	69.1	67.8-70.4	5816	80.9	78.8-83.0	5684	61.0	59.4-62.6	7930	47.2	46.1-48.2	4102	57.3	55.6-59.1	3828	40.1	38.8-41.4
2008	11697	69.6	68.3-70.9	5837	80.3	78.2-82.4	5860	62.4	60.8-64.1	7866	46.2	45.2-47.2	4082	56.4	54.7-58.2	3784	39.2	37.9-40.5
2009	11573	68.0	66.8-69.3	5865	79.5	77.4-81.6	5708	60.0	58.4-61.6	7565	44.2	43.2-45.2	3924	53.6	51.9-55.3	3641	37.6	36.3-38.8
2010	11301	65.9	64.6-67.1	5607	74.7	72.7-76.7	5694	59.9	58.3-61.5	7595	43.9	42.9-44.9	3873	52.1	50.4-53.8	3722	38.1	36.9-39.4
2011	11534	66.1	64.9-67.4	5717	74.9	73.0-76.9	5817	60.1	58.5-61.7	7472	42.5	41.5-43.5	3829	50.7	49.1-52.3	3643	36.6	35.4-37.8
2012	11378	64.3	63.1-65.5	5600	72.2	70.3-74.1	5778	59.1	57.5-60.7	7621	42.9	41.9-43.9	3993	52.0	50.3-53.6	3628	36.4	35.2-37.7
2013	11516	64.2	63.0-65.4	5625	71.1	69.2-73.0	5891	59.4	57.9-61.0	7407	40.8	39.9-41.8	3783	48.0	46.5-49.6	3624	35.7	34.5-36.9
2014	11179	61.1	59.9-62.2	5455	67.5	65.7-69.4	5724	56.6	55.1-58.1	6961	37.9	37.0-38.8	3580	44.8	43.3-46.3	3381	32.8	31.7-34.0
2015	11572	62.6	61.4-63.8	5606	68.6	66.8-70.5	5966	58.6	57.1-60.1	6970	37.3	36.4-38.2	3559	43.8	42.4-45.3	3411	32.6	31.4-33.7
2016	11312	60.2	59.1-61.3	5450	65.4	63.6-67.2	5862	56.9	55.4-58.4	6568	34.8	33.9-35.6	3318	40.4	39.0-41.9	3250	30.6	29.6-31.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Melanoma of the skin

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	916	6.2	5.8-6.7	469	7.2	6.5-7.9	447	5.6	5.1-6.1	379	2.6	2.3-2.9	211	3.3	2.8-3.7	168	2.1	1.8-2.4
1977	1072	7.3	6.8-7.7	545	8.2	7.5-9.0	527	6.5	6.0-7.1	388	2.6	2.4-2.9	231	3.6	3.2-4.1	157	1.9	1.6-2.3
1978	1006	7.0	6.6-7.4	502	7.8	7.1-8.5	504	6.5	6.0-7.1	332	2.3	2.0-2.5	208	3.3	2.9-3.8	124	1.5	1.2-1.8
1979	1064	7.4	6.9-7.8	555	8.5	7.8-9.3	509	6.6	6.0-7.2	376	2.6	2.3-2.9	208	3.3	2.8-3.8	168	2.1	1.8-2.4
1980	1237	8.6	8.1-9.1	660	10.2	9.5-11.1	577	7.4	6.8-8.0	387	2.6	2.4-2.9	222	3.4	3.0-3.9	165	2.0	1.7-2.4
1981	1175	8.1	7.6-8.6	605	9.3	8.6-10.1	570	7.2	6.6-7.9	379	2.6	2.3-2.9	219	3.4	3.0-3.9	160	2.0	1.7-2.3
1982	1218	8.4	7.9-8.9	614	9.5	8.8-10.4	604	7.6	7.0-8.3	382	2.6	2.3-2.9	230	3.6	3.1-4.1	152	1.8	1.5-2.1
1983	1345	9.3	8.7-9.8	720	11.2	10.4-12.1	625	8.0	7.4-8.7	377	2.6	2.3-2.8	213	3.4	2.9-3.9	164	2.1	1.8-2.4
1984	1285	8.8	8.3-9.3	684	10.4	9.6-11.3	601	7.6	7.0-8.3	412	2.8	2.5-3.1	238	3.7	3.2-4.2	174	2.1	1.8-2.4
1985	1336	9.2	8.7-9.7	730	11.2	10.4-12.1	606	7.7	7.1-8.4	417	2.8	2.6-3.1	239	3.8	3.3-4.3	178	2.1	1.8-2.5
1986	1406	9.6	9.1-10.1	736	11.4	10.5-12.2	670	8.6	7.9-9.3	412	2.8	2.5-3.1	230	3.6	3.1-4.1	182	2.2	1.9-2.6
1987	1338	9.1	8.6-9.7	727	11.3	10.4-12.1	611	7.7	7.0-8.3	438	3.0	2.7-3.3	262	4.1	3.6-4.6	176	2.2	1.9-2.6
1988	1271	8.6	8.1-9.1	716	11.1	10.3-12.0	555	6.9	6.3-7.5	416	2.8	2.5-3.1	260	4.1	3.6-4.6	156	1.8	1.6-2.2
1989	1323	9.0	8.5-9.5	725	11.2	10.4-12.1	598	7.4	6.8-8.1	417	2.8	2.6-3.1	266	4.3	3.8-4.8	151	1.8	1.5-2.1
1990	1357	9.1	8.7-9.7	732	11.1	10.3-12.0	625	7.8	7.2-8.4	412	2.8	2.5-3.1	246	3.8	3.4-4.4	166	2.0	1.7-2.4
1991	1304	8.7	8.2-9.2	744	11.3	10.5-12.2	560	6.8	6.3-7.4	410	2.7	2.5-3.0	245	3.9	3.4-4.4	165	1.9	1.6-2.3
1992	1275	8.5	8.0-9.0	720	10.9	10.1-11.8	555	6.7	6.1-7.3	389	2.6	2.3-2.8	230	3.5	3.1-4.0	159	1.8	1.5-2.1
1993	1291	8.5	8.0-9.0	736	11.1	10.3-11.9	555	6.7	6.1-7.3	374	2.5	2.2-2.7	232	3.7	3.2-4.2	142	1.6	1.4-2.0
1994	1446	9.6	9.1-10.1	837	12.6	11.8-13.5	609	7.4	6.8-8.0	405	2.7	2.4-2.9	257	4.0	3.5-4.5	148	1.7	1.4-2.0
1995	1536	10.2	9.7-10.7	915	13.7	12.8-14.6	621	7.6	7.0-8.2	426	2.8	2.5-3.1	248	3.8	3.4-4.4	178	2.0	1.7-2.4
1996	1651	10.9	10.4-11.5	928	13.9	13.0-14.9	723	8.8	8.1-9.5	431	2.8	2.6-3.1	254	3.8	3.3-4.3	177	2.1	1.8-2.4
1997	1770	11.6	11.1-12.2	1021	15.3	14.4-16.3	749	9.1	8.4-9.8	412	2.7	2.4-2.9	264	4.0	3.6-4.6	148	1.7	1.4-2.0
1998	1864	12.2	11.7-12.8	1043	15.4	14.5-16.4	821	10.0	9.4-10.8	418	2.7	2.4-2.9	259	3.9	3.4-4.4	159	1.8	1.5-2.1
1999	1900	12.4	11.8-12.9	1083	15.9	14.9-16.9	817	9.8	9.2-10.5	390	2.5	2.3-2.8	231	3.4	3.0-3.9	159	1.8	1.5-2.1
2000	2160	13.9	13.4-14.5	1210	17.6	16.6-18.7	950	11.5	10.7-12.2	428	2.7	2.4-3.0	268	3.9	3.5-4.4	160	1.8	1.5-2.1
2001	2335	15.0	14.4-15.6	1301	18.7	17.7-19.8	1034	12.4	11.7-13.2	468	2.9	2.7-3.2	289	4.3	3.8-4.8	179	2.0	1.7-2.3
2002	2323	14.8	14.2-15.4	1317	18.8	17.8-19.9	1006	12.0	11.3-12.8	412	2.6	2.3-2.8	218	3.1	2.7-3.6	194	2.2	1.9-2.5
2003	2309	14.6	14.0-15.2	1314	18.6	17.6-19.7	995	11.8	11.0-12.5	419	2.6	2.3-2.8	241	3.5	3.0-3.9	178	1.9	1.7-2.3
2004	2855	18.1	17.4-18.8	1607	22.6	21.5-23.7	1248	15.0	14.2-15.9	421	2.6	2.3-2.8	250	3.5	3.1-4.0	171	1.9	1.6-2.2
2005	3124	19.8	19.1-20.5	1780	25.0	23.9-26.2	1344	16.2	15.3-17.1	431	2.6	2.4-2.9	258	3.7	3.2-4.2	173	1.9	1.6-2.2
2006	3470	21.9	21.2-22.7	1974	27.5	26.3-28.8	1496	18.0	17.0-18.9	450	2.7	2.5-3.0	288	4.1	3.6-4.6	162	1.8	1.5-2.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Melanoma of the skin

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	3202	20.1	19.4-20.8	1805	25.0	23.8-26.2	1397	16.8	15.9-17.7	433	2.6	2.4-2.9	298	4.2	3.7-4.7	135	1.4	1.2-1.7
2008	3447	21.4	20.7-22.2	1970	27.1	25.9-28.4	1477	17.6	16.7-18.5	445	2.7	2.4-2.9	285	3.9	3.5-4.4	160	1.7	1.4-2.0
2009	3699	22.7	22.0-23.5	2133	28.9	27.7-30.2	1566	18.5	17.6-19.5	447	2.7	2.4-2.9	269	3.7	3.3-4.2	178	1.9	1.6-2.2
2010	3471	21.3	20.5-22.0	1961	26.4	25.2-27.6	1510	17.7	16.7-18.6	448	2.6	2.4-2.9	271	3.6	3.2-4.1	177	1.9	1.6-2.2
2011	3896	23.3	22.6-24.1	2231	29.3	28.1-30.6	1665	19.1	18.2-20.1	450	2.6	2.3-2.8	284	3.8	3.3-4.2	166	1.7	1.4-2.0
2012	3721	22.1	21.4-22.9	2092	27.1	25.9-28.3	1629	18.7	17.8-19.7	518	2.9	2.7-3.2	319	4.2	3.8-4.7	199	2.0	1.7-2.3
2013	4081	24.0	23.2-24.8	2349	29.9	28.7-31.2	1732	19.8	18.9-20.8	480	2.7	2.5-3.0	316	4.1	3.7-4.6	164	1.6	1.4-1.9
2014	4025	23.5	22.7-24.2	2315	29.3	28.1-30.6	1710	19.3	18.4-20.3	434	2.4	2.2-2.6	284	3.6	3.2-4.0	150	1.4	1.2-1.7
2015	3930	22.5	21.7-23.2	2298	28.7	27.5-29.9	1632	18.1	17.2-19.0	359	2.0	1.8-2.2	233	2.9	2.5-3.3	126	1.3	1.1-1.6
2016	3909	22.2	21.5-23.0	2297	28.2	27.1-29.4	1612	18.0	17.1-19.0	407	2.2	2.0-2.4	256	3.1	2.8-3.6	151	1.5	1.2-1.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Female breast

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	8525	103.5	101.2-105.8	3310	38.8	37.5-40.2
1977	8388	102.0	99.8-104.3	3351	39.5	38.2-40.9
1978	8537	103.8	101.5-106.1	3359	39.4	38.0-40.8
1979	8407	102.4	100.1-104.6	3196	37.8	36.4-39.1
1980	8527	103.8	101.5-106.1	3300	38.5	37.1-39.9
1981	8734	106.5	104.2-108.8	3267	38.2	36.8-39.5
1982	8748	106.2	103.9-108.5	3212	37.4	36.1-38.8
1983	9119	110.8	108.4-113.1	3178	37.0	35.7-38.4
1984	9489	114.8	112.4-117.2	3312	38.5	37.2-39.9
1985	9814	119.3	116.9-121.8	3315	38.6	37.2-40.0
1986	9874	119.8	117.3-122.2	3297	38.1	36.8-39.5
1987	10181	123.1	120.6-125.5	3265	37.5	36.2-38.9
1988	10311	124.7	122.2-127.2	3327	38.4	37.1-39.8
1989	10121	123.0	120.6-125.5	3255	37.9	36.6-39.2
1990	10753	131.0	128.4-133.5	3156	36.4	35.1-37.7
1991	10699	129.9	127.3-132.4	3040	35.1	33.9-36.4
1992	10729	130.3	127.8-132.8	3020	34.7	33.4-35.9
1993	10535	127.5	125.0-130.0	3013	34.2	33.0-35.5
1994	10953	132.3	129.8-134.8	3007	34.2	33.0-35.5
1995	11232	136.0	133.5-138.6	2963	33.8	32.5-35.0
1996	11875	143.5	140.9-146.2	3013	34.0	32.8-35.3
1997	11855	142.3	139.7-144.9	2750	31.0	29.8-32.2
1998	12191	145.4	142.8-148.0	2706	30.1	29.0-31.3
1999	12099	142.9	140.3-145.5	2558	28.1	27.0-29.2
2000	12064	141.1	138.5-143.6	2571	27.9	26.8-29.0
2001	11774	137.1	134.6-139.7	2536	27.5	26.4-28.6
2002	11699	135.5	133.0-138.0	2446	26.1	25.1-27.2
2003	11559	133.3	130.8-135.7	2420	25.8	24.8-26.9
2004	11399	131.1	128.6-133.5	2298	24.3	23.3-25.3
2005	11443	130.9	128.5-133.4	2226	23.5	22.5-24.5
2006	11624	133.9	131.5-136.4	2155	22.8	21.8-23.8
2007	11760	134.6	132.1-137.1	2167	22.7	21.7-23.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Female breast

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	12088	137.7	135.2-140.2	2094	21.8	20.8-22.8
2009	12348	139.0	136.5-141.5	2090	21.6	20.6-22.6
2010	11629	130.8	128.4-133.3	2134	21.7	20.7-22.7
2011	12150	135.5	133.1-138.0	2050	20.6	19.7-21.5
2012	11925	132.4	130.0-134.9	2001	20.0	19.1-20.9
2013	12384	136.9	134.4-139.4	1996	19.9	19.0-20.8
2014	12615	138.3	135.8-140.8	1813	17.8	16.9-18.7
2015	12661	138.6	136.1-141.2	1882	18.5	17.6-19.3
2016	12260	133.4	131.0-135.9	1954	19.0	18.2-20.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Cervix uteri

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	980	12.5	11.7-13.3	363	4.3	3.9-4.8
1977	856	10.8	10.1-11.6	319	3.8	3.4-4.2
1978	859	11.1	10.3-11.9	349	4.2	3.8-4.7
1979	864	11.3	10.5-12.1	318	3.9	3.5-4.4
1980	871	11.2	10.5-12.0	314	3.9	3.4-4.3
1981	766	9.7	9.0-10.5	330	4.0	3.6-4.5
1982	773	10.2	9.4-11.0	300	3.6	3.2-4.0
1983	801	10.5	9.7-11.2	257	3.1	2.7-3.5
1984	821	10.8	10.0-11.6	241	3.0	2.6-3.4
1985	781	10.2	9.5-11.0	242	3.0	2.6-3.5
1986	704	9.2	8.5-9.9	248	3.0	2.7-3.4
1987	821	10.6	9.9-11.4	251	3.1	2.7-3.5
1988	808	10.6	9.8-11.4	247	3.1	2.7-3.5
1989	772	10.1	9.4-10.8	215	2.6	2.3-3.0
1990	834	11.0	10.2-11.7	257	3.1	2.7-3.5
1991	829	10.8	10.1-11.6	210	2.6	2.3-3.0
1992	783	10.1	9.4-10.8	225	2.8	2.4-3.2
1993	790	10.2	9.5-11.0	209	2.6	2.2-3.0
1994	808	10.4	9.7-11.2	252	3.1	2.7-3.5
1995	899	11.6	10.8-12.3	204	2.5	2.2-2.9
1996	880	11.3	10.6-12.1	243	3.0	2.6-3.4
1997	800	10.2	9.5-10.9	240	2.9	2.6-3.3
1998	735	9.4	8.7-10.1	197	2.4	2.1-2.8
1999	720	9.1	8.5-9.8	238	2.9	2.5-3.3
2000	666	8.5	7.8-9.1	202	2.4	2.1-2.8
2001	652	8.2	7.6-8.9	193	2.2	1.9-2.6
2002	615	7.8	7.2-8.4	203	2.4	2.1-2.8
2003	645	8.2	7.5-8.8	191	2.2	1.9-2.6
2004	570	7.2	6.7-7.9	191	2.2	1.9-2.6
2005	646	8.3	7.6-8.9	197	2.2	1.9-2.6
2006	600	7.6	7.0-8.3	189	2.2	1.9-2.6
2007	606	7.8	7.2-8.5	184	2.2	1.9-2.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Cervix uteri

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	605	7.8	7.2-8.4	160	1.8	1.5-2.1
2009	624	8.0	7.3-8.6	150	1.7	1.5-2.0
2010	566	7.4	6.8-8.0	177	2.0	1.7-2.3
2011	501	6.4	5.8-7.0	178	2.1	1.8-2.4
2012	555	7.0	6.4-7.6	201	2.3	2.0-2.7
2013	570	7.3	6.7-8.0	174	1.9	1.7-2.3
2014	538	6.7	6.2-7.4	191	2.1	1.8-2.5
2015	560	7.2	6.6-7.9	162	1.8	1.5-2.1
2016	523	6.7	6.1-7.4	181	2.0	1.7-2.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Corpus uterus and uterus, not otherwise specified

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	2359	27.2	26.0-28.3	533	6.0	5.5-6.5
1977	2133	24.4	23.4-25.5	484	5.4	4.9-5.9
1978	1911	21.7	20.7-22.7	440	4.9	4.4-5.4
1979	2103	24.2	23.1-25.2	459	5.0	4.6-5.5
1980	2136	24.9	23.8-26.0	488	5.3	4.8-5.8
1981	2199	25.4	24.3-26.5	454	5.0	4.5-5.5
1982	2135	24.5	23.5-25.6	464	5.1	4.6-5.6
1983	2090	24.1	23.1-25.2	445	4.7	4.3-5.2
1984	2077	23.9	22.9-25.0	454	4.9	4.5-5.4
1985	2029	23.3	22.3-24.4	411	4.5	4.0-4.9
1986	2016	23.5	22.5-24.6	412	4.4	3.9-4.8
1987	2102	24.5	23.4-25.6	424	4.6	4.1-5.0
1988	1980	23.1	22.1-24.2	400	4.3	3.9-4.7
1989	1966	22.9	21.9-23.9	406	4.4	3.9-4.8
1990	1995	23.6	22.5-24.7	369	3.9	3.5-4.4
1991	2098	25.3	24.2-26.5	415	4.5	4.1-5.0
1992	2161	25.7	24.6-26.9	418	4.5	4.1-5.0
1993	2066	24.5	23.5-25.6	439	4.7	4.2-5.1
1994	2244	26.4	25.3-27.5	435	4.7	4.3-5.2
1995	2250	26.8	25.7-28.0	453	4.8	4.4-5.3
1996	2341	27.7	26.6-28.9	417	4.5	4.0-4.9
1997	2339	27.8	26.7-28.9	421	4.5	4.1-5.0
1998	2531	30.0	28.9-31.2	417	4.5	4.1-5.0
1999	2463	28.8	27.7-30.0	401	4.2	3.8-4.7
2000	2400	27.8	26.7-29.0	409	4.3	3.8-4.7
2001	2613	30.3	29.2-31.5	394	4.1	3.7-4.5
2002	2643	30.2	29.1-31.4	422	4.5	4.1-4.9
2003	2626	29.9	28.7-31.1	444	4.6	4.2-5.1
2004	2686	30.4	29.3-31.6	418	4.4	3.9-4.8
2005	2723	30.6	29.5-31.8	435	4.6	4.2-5.1
2006	2686	30.0	28.8-31.2	461	4.8	4.3-5.2
2007	2716	30.3	29.2-31.5	450	4.6	4.2-5.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Corpus uterus and uterus, not otherwise specified

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	2994	33.1	31.9-34.4	455	4.7	4.2-5.1
2009	2809	30.8	29.6-31.9	429	4.4	4.0-4.8
2010	2827	30.8	29.6-31.9	480	4.8	4.4-5.3
2011	2973	31.8	30.7-33.0	500	5.1	4.6-5.6
2012	2994	31.5	30.3-32.6	495	4.9	4.4-5.3
2013	3223	33.7	32.5-34.9	534	5.2	4.8-5.7
2014	3133	32.3	31.2-33.5	511	4.9	4.5-5.4
2015	3115	32.0	30.8-33.2	508	4.9	4.4-5.3
2016	3191	32.8	31.7-34.0	562	5.4	4.9-5.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Ovary

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1435	16.9	16.0-17.8	1033	11.9	11.1-12.6
1977	1355	16.1	15.2-17.0	949	10.9	10.2-11.6
1978	1366	16.2	15.4-17.1	931	10.6	9.9-11.3
1979	1403	16.8	15.9-17.8	911	10.4	9.7-11.1
1980	1394	16.5	15.6-17.4	937	10.7	10.0-11.4
1981	1492	17.9	16.9-18.8	960	10.8	10.2-11.6
1982	1434	17.2	16.3-18.2	933	10.6	9.9-11.3
1983	1491	17.8	16.9-18.7	932	10.5	9.8-11.2
1984	1425	17.0	16.1-17.9	929	10.5	9.8-11.2
1985	1480	18.0	17.0-18.9	884	10.1	9.5-10.8
1986	1422	17.2	16.3-18.2	883	10.1	9.4-10.8
1987	1447	17.4	16.5-18.4	915	10.4	9.8-11.2
1988	1428	17.2	16.3-18.1	960	10.8	10.1-11.6
1989	1425	17.1	16.2-18.0	892	10.1	9.4-10.8
1990	1487	17.9	17.0-18.9	915	10.3	9.6-11.0
1991	1528	18.4	17.5-19.4	967	11.0	10.3-11.7
1992	1482	17.9	17.0-18.9	940	10.8	10.1-11.6
1993	1456	17.4	16.5-18.4	889	9.9	9.3-10.6
1994	1504	17.9	17.0-18.9	892	10.1	9.4-10.8
1995	1562	18.7	17.8-19.7	917	10.3	9.7-11.0
1996	1525	18.2	17.3-19.1	880	9.9	9.3-10.6
1997	1484	17.7	16.8-18.7	939	10.5	9.8-11.2
1998	1362	16.2	15.4-17.1	821	9.2	8.5-9.8
1999	1400	16.4	15.6-17.3	839	9.3	8.6-9.9
2000	1417	16.4	15.5-17.2	898	9.8	9.1-10.4
2001	1449	16.7	15.8-17.6	875	9.5	8.8-10.1
2002	1407	16.1	15.2-17.0	867	9.4	8.7-10.0
2003	1368	15.6	14.8-16.5	896	9.6	8.9-10.2
2004	1355	15.4	14.6-16.2	874	9.3	8.7-9.9
2005	1266	14.4	13.6-15.2	897	9.6	8.9-10.2
2006	1305	14.7	13.9-15.6	873	9.3	8.7-10.0
2007	1302	14.6	13.8-15.5	804	8.5	7.9-9.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Ovary

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	1287	14.3	13.6-15.2	847	8.8	8.2-9.4
2009	1233	13.7	13.0-14.6	790	8.1	7.5-8.7
2010	1261	13.9	13.1-14.7	780	8.1	7.5-8.7
2011	1273	13.8	13.1-14.7	802	8.0	7.5-8.6
2012	1166	12.6	11.9-13.4	777	7.8	7.2-8.4
2013	1281	14.0	13.2-14.8	816	8.1	7.5-8.7
2014	1138	12.3	11.5-13.0	717	7.0	6.5-7.5
2015	1184	12.9	12.1-13.6	719	7.1	6.5-7.6
2016	1142	12.2	11.5-13.0	726	7.0	6.5-7.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Prostate

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	4110	74.0	71.6-76.4	1501	29.3	27.8-30.9
1977	4183	74.0	71.7-76.4	1500	28.8	27.3-30.4
1978	4127	73.2	70.9-75.6	1566	30.4	28.8-32.0
1979	4344	77.4	75.0-79.8	1527	29.7	28.2-31.3
1980	4788	83.3	80.9-85.8	1548	29.9	28.4-31.5
1981	4764	83.0	80.6-85.5	1508	29.2	27.7-30.8
1982	4861	84.2	81.8-86.7	1530	29.2	27.7-30.7
1983	5079	86.6	84.2-89.2	1680	31.7	30.1-33.3
1984	5059	86.0	83.6-88.5	1746	33.0	31.4-34.6
1985	5319	89.2	86.7-91.7	1623	30.1	28.6-31.7
1986	5416	91.1	88.6-93.7	1695	31.0	29.5-32.5
1987	5755	95.6	93.1-98.2	1687	30.9	29.4-32.5
1988	5763	96.5	93.9-99.1	1767	32.3	30.8-33.9
1989	5979	98.9	96.3-101.5	1766	32.4	30.8-34.0
1990	7129	114.9	112.2-117.7	1802	32.9	31.3-34.5
1991	8447	133.6	130.7-136.5	1888	34.0	32.4-35.6
1992	10604	164.2	161.1-167.5	1898	34.1	32.5-35.7
1993	10126	155.1	152.0-158.2	1902	34.0	32.4-35.6
1994	9550	145.2	142.3-148.2	1965	34.9	33.3-36.5
1995	9303	140.2	137.3-143.1	1886	33.3	31.8-34.9
1996	10026	150.9	148.0-154.0	1852	32.1	30.7-33.7
1997	10573	158.2	155.2-161.3	1798	30.9	29.5-32.4
1998	10457	154.8	151.8-157.8	1680	28.3	26.9-29.7
1999	10956	161.2	158.2-164.3	1597	26.5	25.2-27.8
2000	12067	176.4	173.2-179.5	1632	26.9	25.6-28.3
2001	11894	171.9	168.8-175.0	1612	26.2	24.9-27.5
2002	12189	174.0	170.9-177.2	1531	24.6	23.4-25.9
2003	11287	159.8	156.9-162.8	1535	24.1	22.9-25.4
2004	10992	153.6	150.8-156.5	1462	22.8	21.6-24.0
2005	11157	154.3	151.4-157.2	1364	20.9	19.8-22.1
2006	12551	172.5	169.5-175.6	1324	20.1	19.0-21.2
2007	12782	172.0	169.0-175.1	1385	20.5	19.5-21.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Prostate

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	12317	162.6	159.7-165.5	1318	19.2	18.2-20.3
2009	12109	156.3	153.5-159.1	1265	18.2	17.2-19.3
2010	11375	144.0	141.3-146.7	1371	19.3	18.3-20.3
2011	11800	146.7	144.0-149.4	1305	18.0	17.0-19.0
2012	10090	122.1	119.7-124.6	1222	16.7	15.8-17.7
2013	9799	116.5	114.2-118.9	1165	15.5	14.6-16.4
2014	9509	110.2	107.9-112.5	1266	16.5	15.6-17.5
2015	9847	112.1	109.9-114.4	1227	15.8	14.9-16.7
2016	10049	112.8	110.5-115.0	1311	16.5	15.6-17.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Testis

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	248	3.3	2.9-3.8	48	0.6	0.4-0.8
1977	241	3.3	2.9-3.8	44	0.6	0.4-0.8
1978	263	3.7	3.3-4.2	49	0.7	0.5-0.9
1979	303	4.3	3.8-4.8	47	0.7	0.5-0.9
1980	325	4.5	4.0-5.1	44	0.6	0.4-0.8
1981	317	4.3	3.8-4.8	32	0.4	0.3-0.6
1982	321	4.4	3.9-4.9	32	0.5	0.3-0.7
1983	336	4.5	4.0-5.0	30	0.4	0.3-0.6
1984	381	5.2	4.7-5.8	27	0.4	0.3-0.6
1985	352	4.7	4.3-5.3	34	0.5	0.3-0.7
1986	355	4.7	4.2-5.2	34	0.5	0.3-0.7
1987	381	5.1	4.6-5.6	27	0.4	0.2-0.5
1988	349	4.6	4.1-5.1	23	0.3	0.2-0.5
1989	363	4.8	4.3-5.4	31	0.4	0.3-0.6
1990	407	5.4	4.9-6.0	24	0.3	0.2-0.5
1991	389	5.2	4.7-5.7	22	0.3	0.2-0.4
1992	399	5.4	4.9-5.9	28	0.4	0.3-0.6
1993	419	5.6	5.1-6.2	32	0.5	0.3-0.7
1994	409	5.6	5.0-6.1	21	0.3	0.2-0.5
1995	416	5.7	5.2-6.3	22	0.3	0.2-0.5
1996	410	5.6	5.1-6.2	26	0.4	0.2-0.5
1997	427	5.9	5.4-6.5	12	0.2	0.1-0.3
1998	428	6.0	5.4-6.6	25	0.3	0.2-0.5
1999	470	6.6	6.0-7.2	17	0.2	0.1-0.4
2000	423	6.0	5.4-6.6	15	0.2	0.1-0.3
2001	435	6.1	5.6-6.7	19	0.3	0.2-0.4
2002	427	6.1	5.5-6.7	25	0.4	0.2-0.5
2003	478	6.9	6.3-7.5	19	0.3	0.2-0.4
2004	462	6.7	6.1-7.3	19	0.3	0.2-0.4
2005	489	7.1	6.5-7.8	14	0.2	0.1-0.3
2006	464	6.8	6.2-7.4	23	0.3	0.2-0.5
2007	500	7.3	6.7-8.0	7	0.1	0.0-0.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Testis

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	518	7.6	7.0-8.3	15	0.2	0.1-0.4
2009	504	7.4	6.8-8.1	15	0.2	0.1-0.4
2010	509	7.5	6.9-8.2	28	0.4	0.3-0.6
2011	507	7.4	6.8-8.1	16	0.2	0.1-0.3
2012	535	7.9	7.2-8.6	18	0.3	0.2-0.4
2013	474	7.0	6.4-7.7	13	0.2	0.1-0.3
2014	496	7.3	6.7-8.0	17	0.3	0.1-0.4
2015	492	7.2	6.5-7.8	18	0.2	0.1-0.4
2016	498	7.3	6.7-8.0	16	0.2	0.1-0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Urinary bladder (incl. in situ)

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	3071	20.4	19.7-21.1	2239	36.9	35.3-38.6	832	9.4	8.8-10.1	950	6.7	6.3-7.2	673	12.4	11.4-13.4	277	3.3	2.9-3.7
1977	3224	21.4	20.6-22.1	2329	38.0	36.4-39.6	895	10.3	9.6-11.0	967	6.8	6.3-7.2	677	12.3	11.4-13.3	290	3.4	3.0-3.8
1978	3368	22.4	21.6-23.2	2443	40.4	38.7-42.1	925	10.6	9.9-11.3	948	6.6	6.1-7.0	653	11.9	10.9-12.9	295	3.4	3.0-3.8
1979	3401	22.5	21.7-23.3	2493	41.4	39.7-43.1	908	10.2	9.5-10.9	925	6.4	6.0-6.8	631	11.4	10.5-12.4	294	3.3	3.0-3.7
1980	3582	23.5	22.7-24.3	2652	43.4	41.7-45.2	930	10.3	9.7-11.0	913	6.1	5.7-6.6	641	11.4	10.5-12.4	272	3.0	2.7-3.4
1981	3500	22.9	22.1-23.7	2582	42.3	40.6-44.0	918	10.3	9.6-11.0	916	6.2	5.8-6.6	638	11.5	10.6-12.5	278	3.1	2.7-3.5
1982	3563	23.2	22.4-24.0	2578	41.8	40.2-43.6	985	10.9	10.2-11.6	881	5.9	5.5-6.3	623	11.1	10.2-12.1	258	2.8	2.5-3.2
1983	3622	23.5	22.7-24.3	2644	43.5	41.7-45.2	978	10.7	10.0-11.4	844	5.6	5.2-6.0	598	10.9	10.0-11.9	246	2.6	2.3-3.0
1984	3513	22.6	21.9-23.4	2543	40.9	39.3-42.6	970	10.6	10.0-11.3	852	5.7	5.3-6.1	575	10.4	9.6-11.4	277	3.0	2.6-3.3
1985	3515	22.7	22.0-23.5	2545	41.2	39.6-42.9	970	10.6	10.0-11.3	854	5.7	5.3-6.0	585	10.5	9.7-11.4	269	2.9	2.5-3.2
1986	3679	23.7	22.9-24.5	2674	43.0	41.3-44.7	1005	11.1	10.4-11.8	834	5.5	5.1-5.9	567	10.4	9.5-11.3	267	2.8	2.5-3.2
1987	3556	22.7	22.0-23.5	2550	40.7	39.1-42.4	1006	10.9	10.2-11.6	773	5.1	4.7-5.5	528	9.4	8.6-10.3	245	2.6	2.3-2.9
1988	3524	22.6	21.9-23.4	2529	40.4	38.8-42.1	995	10.8	10.1-11.5	829	5.3	5.0-5.7	543	9.4	8.6-10.2	286	2.9	2.6-3.3
1989	3509	22.6	21.8-23.3	2544	40.9	39.3-42.6	965	10.5	9.8-11.2	818	5.3	4.9-5.7	563	10.0	9.2-10.9	255	2.6	2.3-3.0
1990	3640	23.4	22.7-24.2	2584	41.4	39.8-43.1	1056	11.5	10.8-12.2	799	5.2	4.8-5.6	520	9.2	8.4-10.1	279	2.9	2.5-3.2
1991	3638	23.3	22.6-24.1	2627	41.8	40.2-43.5	1011	11.0	10.3-11.7	765	4.9	4.6-5.3	525	9.1	8.3-9.9	240	2.5	2.2-2.8
1992	3639	23.3	22.5-24.1	2602	41.5	39.9-43.1	1037	11.4	10.7-12.1	799	5.1	4.7-5.4	490	8.6	7.8-9.4	309	3.1	2.8-3.5
1993	3702	23.6	22.8-24.4	2615	41.1	39.5-42.7	1087	11.8	11.1-12.5	830	5.3	4.9-5.6	561	9.7	8.9-10.6	269	2.7	2.4-3.1
1994	3678	23.5	22.7-24.3	2617	41.2	39.6-42.8	1061	11.6	10.9-12.3	830	5.2	4.9-5.6	558	9.5	8.7-10.4	272	2.7	2.4-3.1
1995	3813	24.3	23.5-25.1	2700	42.3	40.7-44.0	1113	12.1	11.4-12.8	835	5.2	4.9-5.6	548	9.2	8.4-10.0	287	2.9	2.5-3.2
1996	3898	24.8	24.0-25.6	2846	44.4	42.8-46.1	1052	11.4	10.7-12.1	840	5.2	4.9-5.6	547	9.2	8.5-10.1	293	2.9	2.6-3.3
1997	4172	26.5	25.7-27.3	2997	46.2	44.6-48.0	1175	12.6	11.9-13.4	857	5.3	5.0-5.7	572	9.4	8.7-10.2	285	2.9	2.6-3.2
1998	4314	27.2	26.4-28.0	3112	47.4	45.8-49.2	1202	13.0	12.3-13.8	832	5.1	4.8-5.5	569	9.3	8.5-10.1	263	2.6	2.3-2.9
1999	4245	26.5	25.7-27.3	3053	46.2	44.5-47.8	1192	12.8	12.1-13.6	830	5.0	4.7-5.4	573	9.3	8.5-10.1	257	2.5	2.2-2.8
2000	4333	26.8	26.0-27.6	3106	46.6	44.9-48.2	1227	13.0	12.2-13.7	829	5.0	4.7-5.3	569	8.9	8.2-9.7	260	2.5	2.2-2.9
2001	4272	26.2	25.5-27.0	3113	46.3	44.7-48.0	1159	12.3	11.6-13.0	847	5.1	4.7-5.4	597	9.4	8.6-10.1	250	2.4	2.1-2.7
2002	4385	26.8	26.0-27.6	3247	47.7	46.1-49.4	1138	12.0	11.3-12.7	817	4.8	4.5-5.2	543	8.4	7.7-9.2	274	2.7	2.4-3.0
2003	4518	27.5	26.7-28.3	3321	48.2	46.6-49.9	1197	12.6	11.9-13.3	755	4.4	4.1-4.7	512	7.9	7.2-8.6	243	2.3	2.0-2.6
2004	4651	28.1	27.3-28.9	3408	49.1	47.5-50.8	1243	13.0	12.3-13.7	830	4.8	4.5-5.2	576	8.6	7.9-9.3	254	2.4	2.1-2.7
2005	4677	28.1	27.3-29.0	3410	48.8	47.2-50.5	1267	13.2	12.4-13.9	790	4.6	4.3-4.9	526	7.8	7.2-8.5	264	2.5	2.2-2.8
2006	4565	27.3	26.5-28.1	3386	48.2	46.6-49.9	1179	12.4	11.7-13.1	816	4.7	4.4-5.1	550	8.1	7.5-8.9	266	2.5	2.2-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Urinary bladder (incl. in situ)

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	4618	27.5	26.7-28.3	3396	47.7	46.1-49.4	1222	12.8	12.1-13.6	882	5.0	4.7-5.4	619	8.9	8.3-9.7	263	2.5	2.2-2.8
2008	4721	27.7	26.9-28.5	3565	49.4	47.8-51.0	1156	11.9	11.2-12.6	876	5.0	4.6-5.3	605	8.7	8.0-9.4	271	2.5	2.2-2.9
2009	4738	27.5	26.8-28.3	3524	48.2	46.6-49.9	1214	12.3	11.6-13.0	856	4.8	4.4-5.1	628	8.9	8.2-9.6	228	2.0	1.8-2.3
2010	4460	25.7	24.9-26.5	3382	45.7	44.2-47.3	1078	10.9	10.3-11.6	907	5.0	4.6-5.3	637	8.7	8.1-9.4	270	2.4	2.1-2.7
2011	4732	27.1	26.3-27.9	3521	46.8	45.2-48.4	1211	12.3	11.6-13.1	838	4.5	4.2-4.9	587	8.0	7.4-8.7	251	2.3	2.0-2.6
2012	4852	27.2	26.4-28.0	3587	47.0	45.5-48.6	1265	12.4	11.7-13.2	875	4.7	4.4-5.0	602	8.0	7.4-8.7	273	2.4	2.1-2.7
2013	4856	27.0	26.2-27.8	3615	46.8	45.2-48.3	1241	12.2	11.5-12.9	864	4.6	4.3-4.9	621	8.2	7.6-8.9	243	2.1	1.8-2.4
2014	4793	26.1	25.3-26.8	3576	45.2	43.7-46.7	1217	11.6	11.0-12.3	923	4.8	4.5-5.1	631	8.2	7.5-8.8	292	2.6	2.3-2.9
2015	4768	25.7	25.0-26.5	3593	44.6	43.1-46.1	1175	11.5	10.8-12.2	863	4.5	4.2-4.8	592	7.5	6.9-8.1	271	2.4	2.1-2.7
2016	4758	25.3	24.6-26.0	3599	44.0	42.5-45.5	1159	10.9	10.3-11.6	896	4.6	4.3-4.9	634	8.0	7.4-8.7	262	2.3	2.0-2.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Kidney and renal pelvis

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1106	7.2	6.8-7.6	688	10.5	9.7-11.3	418	4.8	4.3-5.3	598	3.9	3.6-4.2	381	6.0	5.4-6.6	217	2.4	2.1-2.8
1977	1159	7.6	7.1-8.0	727	11.1	10.3-12.0	432	5.0	4.5-5.5	633	4.2	3.8-4.5	393	6.2	5.5-6.8	240	2.7	2.4-3.1
1978	1147	7.5	7.1-8.0	701	10.9	10.1-11.8	446	5.1	4.6-5.6	575	3.7	3.4-4.1	361	5.6	5.0-6.3	214	2.4	2.1-2.8
1979	1144	7.5	7.1-8.0	703	10.9	10.1-11.8	441	4.9	4.4-5.4	636	4.2	3.9-4.6	398	6.3	5.6-6.9	238	2.7	2.4-3.1
1980	1138	7.4	7.0-7.9	695	10.6	9.8-11.4	443	5.1	4.7-5.7	598	3.9	3.5-4.2	357	5.7	5.1-6.3	241	2.6	2.3-3.0
1981	1213	8.1	7.6-8.5	713	11.3	10.5-12.2	500	5.8	5.3-6.3	593	3.9	3.6-4.2	358	5.6	5.0-6.3	235	2.6	2.3-3.0
1982	1296	8.5	8.0-9.0	781	12.0	11.2-12.9	515	5.9	5.4-6.4	599	3.8	3.5-4.2	358	5.7	5.1-6.3	241	2.6	2.3-3.0
1983	1354	8.8	8.4-9.3	833	12.8	11.9-13.7	521	6.0	5.5-6.5	588	3.8	3.5-4.1	350	5.5	4.9-6.1	238	2.7	2.3-3.0
1984	1372	9.0	8.5-9.5	861	13.3	12.4-14.3	511	5.8	5.3-6.4	599	3.9	3.6-4.3	371	6.0	5.4-6.7	228	2.5	2.2-2.9
1985	1456	9.5	9.0-10.0	885	13.7	12.8-14.6	571	6.5	5.9-7.0	614	4.0	3.7-4.3	350	5.6	5.0-6.3	264	3.0	2.6-3.3
1986	1446	9.4	9.0-10.0	889	13.7	12.7-14.6	557	6.4	5.9-7.0	654	4.2	3.9-4.6	395	6.2	5.6-6.9	259	2.9	2.5-3.2
1987	1571	10.2	9.7-10.8	932	14.3	13.4-15.3	639	7.3	6.7-7.9	652	4.2	3.9-4.6	404	6.5	5.9-7.2	248	2.7	2.4-3.1
1988	1449	9.5	9.0-10.0	877	13.6	12.7-14.6	572	6.6	6.0-7.1	622	4.0	3.7-4.4	359	5.7	5.1-6.3	263	2.9	2.5-3.3
1989	1464	9.7	9.2-10.2	910	14.2	13.3-15.2	554	6.3	5.8-6.9	626	4.1	3.8-4.4	376	6.1	5.4-6.7	250	2.8	2.4-3.1
1990	1626	10.6	10.1-11.2	986	15.2	14.3-16.3	640	7.4	6.8-8.0	602	3.9	3.6-4.2	355	5.6	5.0-6.2	247	2.7	2.3-3.0
1991	1653	10.9	10.3-11.4	981	15.1	14.1-16.1	672	7.8	7.2-8.4	626	4.0	3.7-4.4	370	5.8	5.2-6.5	256	2.8	2.4-3.1
1992	1669	11.0	10.4-11.5	1000	15.3	14.3-16.3	669	7.8	7.2-8.4	577	3.7	3.4-4.1	343	5.6	5.0-6.2	234	2.5	2.2-2.9
1993	1722	11.2	10.7-11.8	1055	16.0	15.0-17.0	667	7.7	7.1-8.3	579	3.7	3.4-4.1	366	5.6	5.1-6.3	213	2.3	2.0-2.7
1994	1757	11.5	11.0-12.1	1059	16.1	15.2-17.2	698	8.0	7.4-8.7	662	4.2	3.9-4.6	410	6.5	5.9-7.2	252	2.7	2.4-3.1
1995	1743	11.4	10.8-11.9	1037	15.6	14.7-16.6	706	8.1	7.5-8.8	628	4.0	3.7-4.4	363	5.7	5.2-6.4	265	2.9	2.5-3.3
1996	1910	12.4	11.9-13.0	1162	17.6	16.6-18.6	748	8.6	8.0-9.2	649	4.2	3.8-4.5	373	5.8	5.2-6.4	276	3.0	2.6-3.3
1997	1976	12.8	12.3-13.4	1218	18.4	17.4-19.5	758	8.7	8.1-9.4	633	4.0	3.7-4.4	392	6.1	5.5-6.7	241	2.6	2.3-3.0
1998	2082	13.5	12.9-14.1	1271	18.7	17.7-19.8	811	9.3	8.7-10.0	634	4.0	3.7-4.3	385	5.8	5.3-6.5	249	2.6	2.3-3.0
1999	2171	13.9	13.3-14.5	1342	19.7	18.6-20.7	829	9.4	8.8-10.1	548	3.4	3.1-3.7	343	5.2	4.6-5.8	205	2.1	1.8-2.4
2000	2196	13.9	13.3-14.5	1300	18.9	17.9-19.9	896	10.1	9.4-10.8	605	3.7	3.5-4.1	370	5.6	5.0-6.2	235	2.5	2.2-2.8
2001	2279	14.4	13.8-15.0	1413	20.3	19.2-21.3	866	9.7	9.1-10.4	638	3.9	3.6-4.2	374	5.5	5.0-6.1	264	2.8	2.5-3.2
2002	2299	14.4	13.8-15.0	1435	20.4	19.4-21.5	864	9.6	9.0-10.3	605	3.7	3.4-4.0	363	5.3	4.8-5.9	242	2.5	2.2-2.8
2003	2419	15.1	14.5-15.7	1525	21.3	20.3-22.4	894	9.9	9.3-10.6	604	3.6	3.4-4.0	361	5.2	4.7-5.7	243	2.4	2.1-2.7
2004	2609	16.2	15.6-16.9	1624	22.6	21.5-23.8	985	11.1	10.4-11.8	635	3.8	3.5-4.1	383	5.5	5.0-6.1	252	2.6	2.3-2.9
2005	2603	16.1	15.5-16.8	1599	22.3	21.2-23.4	1004	11.2	10.6-12.0	602	3.6	3.3-3.9	379	5.4	4.9-6.0	223	2.3	2.0-2.6
2006	2809	17.3	16.6-17.9	1785	24.5	23.4-25.7	1024	11.4	10.7-12.1	632	3.7	3.5-4.0	381	5.5	4.9-6.0	251	2.6	2.2-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Kidney and renal pelvis

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	2880	17.6	17.0-18.3	1833	25.0	23.8-26.2	1047	11.8	11.1-12.5	612	3.6	3.3-3.9	380	5.3	4.8-5.8	232	2.4	2.1-2.7
2008	2857	17.2	16.6-17.9	1802	24.1	23.0-25.3	1055	11.6	10.9-12.3	606	3.5	3.2-3.8	391	5.4	4.9-5.9	215	2.1	1.8-2.4
2009	2744	16.6	15.9-17.2	1692	22.5	21.4-23.6	1052	11.7	11.0-12.4	621	3.6	3.3-3.9	406	5.5	5.0-6.1	215	2.2	1.9-2.5
2010	2826	16.8	16.2-17.5	1775	23.2	22.1-24.3	1051	11.5	10.8-12.3	593	3.4	3.1-3.6	378	5.0	4.5-5.5	215	2.1	1.8-2.4
2011	2902	17.2	16.5-17.8	1874	24.1	23.0-25.3	1028	11.3	10.6-12.0	613	3.4	3.2-3.7	384	5.1	4.6-5.6	229	2.2	1.9-2.5
2012	3007	17.5	16.9-18.2	1918	24.4	23.3-25.5	1089	11.8	11.0-12.5	631	3.5	3.2-3.8	417	5.4	4.9-6.0	214	2.1	1.8-2.4
2013	3006	17.3	16.7-17.9	1937	24.2	23.1-25.3	1069	11.5	10.8-12.2	579	3.2	2.9-3.5	366	4.7	4.2-5.2	213	2.0	1.7-2.3
2014	3053	17.4	16.7-18.0	1975	24.3	23.2-25.4	1078	11.4	10.7-12.2	580	3.1	2.8-3.3	368	4.5	4.1-5.0	212	2.0	1.7-2.3
2015	3104	17.4	16.8-18.0	1997	24.3	23.2-25.4	1107	11.4	10.7-12.2	629	3.3	3.1-3.6	401	4.9	4.5-5.5	228	2.1	1.8-2.4
2016	3021	16.9	16.3-17.6	1985	23.9	22.8-25.0	1036	10.9	10.3-11.7	531	2.8	2.6-3.1	350	4.2	3.8-4.7	181	1.7	1.5-2.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Brain and other central nervous system

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	906	5.8	5.4-6.2	510	7.1	6.5-7.8	396	4.7	4.3-5.2	688	4.4	4.1-4.8	376	5.4	4.9-6.0	312	3.7	3.3-4.1
1977	917	6.0	5.6-6.4	514	7.3	6.7-8.0	403	4.8	4.3-5.3	692	4.4	4.1-4.8	376	5.4	4.8-5.9	316	3.7	3.3-4.1
1978	890	5.8	5.4-6.2	460	6.6	6.0-7.3	430	5.2	4.7-5.7	660	4.2	3.9-4.5	354	5.1	4.6-5.7	306	3.5	3.1-3.9
1979	953	6.2	5.8-6.6	532	7.8	7.1-8.5	421	5.0	4.5-5.6	646	4.1	3.8-4.5	363	5.3	4.8-5.9	283	3.3	2.9-3.7
1980	952	6.2	5.8-6.6	522	7.4	6.8-8.1	430	5.2	4.7-5.7	723	4.6	4.3-5.0	418	6.1	5.5-6.7	305	3.5	3.1-4.0
1981	959	6.4	6.0-6.8	536	8.0	7.3-8.8	423	5.0	4.6-5.6	671	4.4	4.0-4.7	368	5.5	4.9-6.1	303	3.5	3.1-3.9
1982	995	6.6	6.2-7.0	519	7.7	7.0-8.4	476	5.8	5.3-6.3	690	4.5	4.2-4.9	348	5.2	4.6-5.8	342	4.0	3.6-4.5
1983	947	6.4	6.0-6.8	481	7.1	6.5-7.8	466	5.7	5.2-6.3	657	4.3	3.9-4.6	352	5.3	4.7-5.9	305	3.4	3.1-3.9
1984	1042	6.9	6.4-7.3	541	8.0	7.3-8.7	501	6.0	5.5-6.6	662	4.3	4.0-4.6	350	5.3	4.7-5.9	312	3.6	3.2-4.0
1985	1016	6.8	6.4-7.3	548	8.2	7.5-8.9	468	5.7	5.2-6.2	652	4.3	4.0-4.7	325	5.0	4.4-5.6	327	3.8	3.4-4.3
1986	1021	6.8	6.4-7.3	562	8.5	7.8-9.2	459	5.6	5.1-6.1	702	4.6	4.3-5.0	386	5.9	5.3-6.6	316	3.6	3.2-4.0
1987	1098	7.4	6.9-7.8	570	8.5	7.8-9.2	528	6.3	5.8-6.9	688	4.5	4.2-4.9	351	5.3	4.8-5.9	337	3.9	3.5-4.3
1988	1084	7.3	6.8-7.7	588	8.8	8.1-9.5	496	6.1	5.5-6.6	657	4.2	3.9-4.6	363	5.3	4.8-5.9	294	3.3	3.0-3.7
1989	1083	7.3	6.9-7.8	610	9.1	8.4-9.9	473	5.9	5.3-6.4	644	4.2	3.9-4.6	352	5.3	4.8-5.9	292	3.4	3.0-3.8
1990	1137	7.7	7.3-8.2	614	9.3	8.6-10.1	523	6.5	5.9-7.1	613	4.1	3.7-4.4	356	5.5	4.9-6.1	257	2.9	2.6-3.3
1991	1079	7.3	6.9-7.8	576	8.7	8.0-9.5	503	6.2	5.7-6.8	655	4.4	4.1-4.8	367	5.7	5.1-6.3	288	3.5	3.1-3.9
1992	1088	7.3	6.9-7.8	577	8.6	7.9-9.4	511	6.3	5.8-6.9	648	4.3	3.9-4.6	332	5.0	4.5-5.6	316	3.7	3.3-4.2
1993	1099	7.4	7.0-7.8	587	8.7	8.0-9.5	512	6.3	5.7-6.9	627	4.1	3.8-4.5	343	5.2	4.6-5.8	284	3.3	2.9-3.8
1994	1190	8.0	7.5-8.5	648	9.7	9.0-10.5	542	6.5	6.0-7.1	713	4.7	4.4-5.1	387	5.8	5.3-6.5	326	3.8	3.4-4.2
1995	1267	8.5	8.0-9.0	647	9.6	8.8-10.3	620	7.6	7.0-8.2	666	4.4	4.1-4.8	359	5.4	4.9-6.0	307	3.6	3.2-4.0
1996	1249	8.4	8.0-8.9	670	10.0	9.2-10.8	579	7.1	6.6-7.7	701	4.7	4.3-5.0	378	5.7	5.1-6.3	323	3.8	3.4-4.2
1997	1165	7.8	7.4-8.3	623	9.2	8.5-9.9	542	6.7	6.1-7.3	699	4.6	4.3-5.0	362	5.4	4.8-6.0	337	3.9	3.5-4.4
1998	1268	8.5	8.0-9.0	698	10.1	9.4-10.9	570	7.1	6.5-7.7	691	4.5	4.2-4.9	375	5.5	4.9-6.1	316	3.7	3.3-4.1
1999	1131	7.5	7.1-8.0	594	8.7	8.0-9.4	537	6.5	6.0-7.1	696	4.5	4.2-4.9	380	5.5	5.0-6.1	316	3.6	3.3-4.1
2000	1251	8.3	7.8-8.7	640	9.3	8.6-10.0	611	7.4	6.8-8.0	653	4.2	3.9-4.5	336	4.8	4.3-5.4	317	3.6	3.2-4.0
2001	1156	7.6	7.2-8.0	615	8.8	8.1-9.6	541	6.5	6.0-7.1	634	4.0	3.7-4.4	337	4.8	4.3-5.4	297	3.4	3.0-3.8
2002	1176	7.7	7.3-8.1	647	9.3	8.6-10.1	529	6.4	5.8-7.0	671	4.2	3.9-4.6	373	5.3	4.8-5.9	298	3.4	3.0-3.8
2003	1178	7.7	7.3-8.2	607	8.6	7.9-9.3	571	7.0	6.4-7.6	619	3.9	3.6-4.3	351	4.9	4.4-5.5	268	3.1	2.7-3.5
2004	1135	7.4	7.0-7.9	635	9.0	8.3-9.8	500	6.1	5.6-6.7	657	4.1	3.8-4.5	350	4.9	4.4-5.5	307	3.5	3.1-3.9
2005	1192	7.8	7.4-8.3	681	9.7	9.0-10.5	511	6.2	5.7-6.8	698	4.4	4.1-4.7	385	5.4	4.9-6.0	313	3.6	3.2-4.0
2006	1196	7.9	7.4-8.3	674	9.5	8.8-10.3	522	6.4	5.9-7.0	661	4.1	3.8-4.4	391	5.4	4.9-6.0	270	3.0	2.6-3.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Brain and other central nervous system

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	1119	7.3	6.9-7.8	601	8.5	7.8-9.2	518	6.4	5.8-7.0	690	4.3	3.9-4.6	403	5.5	5.0-6.1	287	3.2	2.8-3.6
2008	1161	7.6	7.1-8.0	630	8.9	8.2-9.6	531	6.5	5.9-7.1	638	3.9	3.6-4.2	369	5.0	4.5-5.6	269	3.0	2.6-3.4
2009	1180	7.6	7.1-8.0	630	8.7	8.0-9.4	550	6.6	6.0-7.2	733	4.5	4.1-4.8	404	5.4	4.9-6.0	329	3.7	3.3-4.1
2010	1122	7.2	6.8-7.7	617	8.6	7.9-9.3	505	6.1	5.6-6.7	700	4.2	3.9-4.6	376	5.0	4.5-5.6	324	3.6	3.2-4.0
2011	1111	7.1	6.7-7.6	615	8.4	7.7-9.1	496	6.1	5.5-6.7	664	4.0	3.7-4.3	361	4.8	4.3-5.3	303	3.2	2.9-3.6
2012	1168	7.3	6.9-7.7	631	8.4	7.7-9.1	537	6.4	5.8-7.0	753	4.4	4.1-4.8	403	5.2	4.7-5.8	350	3.8	3.4-4.2
2013	1141	7.2	6.8-7.6	655	8.7	8.0-9.4	486	5.8	5.3-6.3	701	4.0	3.7-4.3	380	4.8	4.3-5.3	321	3.3	3.0-3.7
2014	1259	7.8	7.4-8.3	695	9.1	8.4-9.8	564	6.7	6.2-7.3	762	4.4	4.1-4.7	419	5.2	4.7-5.7	343	3.7	3.3-4.1
2015	1190	7.4	6.9-7.8	623	8.1	7.5-8.8	567	6.7	6.1-7.3	730	4.2	3.9-4.5	378	4.7	4.2-5.2	352	3.7	3.3-4.1
2016	1166	7.1	6.7-7.6	671	8.7	8.0-9.4	495	5.8	5.2-6.3	764	4.3	4.0-4.6	442	5.4	4.9-6.0	322	3.4	3.0-3.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Thyroid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	490	3.4	3.1-3.7	154	2.4	2.0-2.8	336	4.3	3.8-4.8	102	0.7	0.6-0.9	41	0.7	0.5-0.9	61	0.7	0.5-0.9
1977	565	3.9	3.5-4.2	168	2.5	2.1-2.9	397	5.1	4.6-5.6	114	0.8	0.6-0.9	46	0.7	0.5-1.0	68	0.8	0.6-1.0
1978	558	3.8	3.5-4.2	157	2.4	2.0-2.8	401	5.1	4.6-5.6	95	0.7	0.5-0.8	31	0.5	0.4-0.8	64	0.7	0.6-0.9
1979	511	3.6	3.2-3.9	171	2.7	2.3-3.1	340	4.4	3.9-4.9	76	0.5	0.4-0.6	28	0.5	0.3-0.7	48	0.5	0.4-0.7
1980	538	3.7	3.4-4.0	161	2.5	2.1-2.9	377	4.8	4.3-5.3	109	0.7	0.6-0.9	48	0.8	0.6-1.1	61	0.7	0.5-0.9
1981	520	3.6	3.3-3.9	157	2.4	2.0-2.8	363	4.6	4.2-5.2	92	0.6	0.5-0.8	36	0.6	0.4-0.8	56	0.6	0.5-0.8
1982	508	3.5	3.2-3.9	157	2.4	2.0-2.8	351	4.6	4.1-5.1	97	0.6	0.5-0.8	41	0.6	0.5-0.9	56	0.6	0.5-0.8
1983	596	4.1	3.8-4.5	159	2.4	2.0-2.8	437	5.7	5.1-6.2	78	0.5	0.4-0.7	21	0.3	0.2-0.5	57	0.6	0.5-0.8
1984	648	4.5	4.1-4.8	186	2.9	2.5-3.4	462	5.9	5.4-6.5	101	0.7	0.5-0.8	37	0.6	0.4-0.9	64	0.7	0.5-0.9
1985	654	4.5	4.1-4.9	188	2.8	2.4-3.2	466	6.0	5.4-6.6	89	0.6	0.5-0.7	43	0.7	0.5-1.0	46	0.5	0.4-0.7
1986	629	4.3	3.9-4.6	176	2.6	2.2-3.0	453	5.8	5.3-6.4	81	0.5	0.4-0.7	28	0.5	0.3-0.7	53	0.6	0.4-0.8
1987	609	4.2	3.9-4.5	195	2.9	2.5-3.4	414	5.4	4.8-5.9	80	0.5	0.4-0.6	25	0.4	0.2-0.6	55	0.6	0.4-0.8
1988	606	4.2	3.8-4.5	173	2.6	2.2-3.1	433	5.6	5.1-6.2	78	0.5	0.4-0.6	29	0.5	0.3-0.7	49	0.5	0.4-0.7
1989	598	4.1	3.8-4.5	165	2.5	2.1-2.9	433	5.7	5.2-6.3	82	0.5	0.4-0.7	24	0.4	0.2-0.6	58	0.6	0.5-0.8
1990	640	4.4	4.1-4.8	185	2.8	2.4-3.3	455	5.9	5.3-6.4	77	0.5	0.4-0.6	29	0.4	0.3-0.6	48	0.5	0.4-0.7
1991	718	4.9	4.6-5.3	209	3.1	2.7-3.5	509	6.7	6.1-7.3	72	0.5	0.4-0.6	23	0.4	0.2-0.6	49	0.5	0.4-0.7
1992	739	5.0	4.7-5.4	199	3.0	2.6-3.5	540	6.9	6.3-7.5	81	0.5	0.4-0.7	26	0.4	0.3-0.6	55	0.6	0.4-0.7
1993	769	5.3	4.9-5.6	210	3.1	2.7-3.5	559	7.4	6.7-8.0	83	0.5	0.4-0.7	30	0.5	0.3-0.7	53	0.6	0.4-0.7
1994	832	5.7	5.3-6.1	220	3.3	2.8-3.7	612	8.0	7.4-8.7	81	0.5	0.4-0.7	35	0.6	0.4-0.8	46	0.5	0.3-0.6
1995	932	6.3	5.9-6.7	261	3.8	3.4-4.3	671	8.7	8.1-9.4	78	0.5	0.4-0.6	31	0.5	0.3-0.7	47	0.5	0.4-0.7
1996	987	6.8	6.3-7.2	253	3.7	3.2-4.2	734	9.7	9.0-10.5	112	0.7	0.6-0.9	38	0.6	0.4-0.8	74	0.8	0.6-1.0
1997	1092	7.4	7.0-7.9	294	4.3	3.8-4.8	798	10.4	9.7-11.2	85	0.5	0.4-0.7	32	0.5	0.3-0.7	53	0.5	0.4-0.7
1998	1121	7.6	7.2-8.1	316	4.6	4.1-5.1	805	10.5	9.8-11.2	78	0.5	0.4-0.6	25	0.4	0.2-0.6	53	0.5	0.4-0.7
1999	1314	8.9	8.4-9.4	314	4.5	4.0-5.0	1000	13.1	12.3-14.0	76	0.5	0.4-0.6	30	0.4	0.3-0.6	46	0.5	0.3-0.6
2000	1438	9.7	9.2-10.2	369	5.2	4.7-5.8	1069	14.0	13.2-14.9	80	0.5	0.4-0.6	25	0.4	0.2-0.6	55	0.6	0.4-0.7
2001	1521	10.2	9.7-10.7	349	4.9	4.4-5.4	1172	15.3	14.5-16.2	80	0.5	0.4-0.6	33	0.5	0.3-0.7	47	0.5	0.4-0.7
2002	1570	10.5	10.0-11.0	424	5.9	5.4-6.5	1146	14.9	14.1-15.8	80	0.5	0.4-0.6	29	0.4	0.3-0.6	51	0.5	0.4-0.7
2003	1593	10.6	10.1-11.2	416	5.8	5.2-6.4	1177	15.3	14.4-16.2	77	0.5	0.4-0.6	23	0.3	0.2-0.5	54	0.5	0.4-0.7
2004	1779	11.9	11.3-12.4	435	6.0	5.5-6.6	1344	17.5	16.6-18.5	68	0.4	0.3-0.5	30	0.4	0.3-0.6	38	0.4	0.2-0.5
2005	2114	14.1	13.5-14.8	536	7.4	6.8-8.1	1578	20.7	19.7-21.8	83	0.5	0.4-0.6	40	0.6	0.4-0.8	43	0.4	0.3-0.6
2006	2223	14.8	14.2-15.5	560	7.7	7.1-8.4	1663	21.7	20.7-22.8	96	0.6	0.5-0.7	38	0.5	0.4-0.8	58	0.6	0.4-0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Thyroid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	2439	16.2	15.6-16.9	616	8.5	7.8-9.2	1823	23.7	22.6-24.8	93	0.6	0.4-0.7	37	0.5	0.4-0.7	56	0.6	0.4-0.8
2008	2774	18.5	17.8-19.2	657	8.9	8.2-9.6	2117	27.8	26.6-29.0	84	0.5	0.4-0.6	40	0.6	0.4-0.8	44	0.5	0.3-0.6
2009	2968	19.7	19.0-20.4	767	10.4	9.7-11.2	2201	28.7	27.5-29.9	96	0.5	0.4-0.7	43	0.6	0.4-0.8	53	0.5	0.4-0.7
2010	2968	19.6	18.9-20.4	749	10.0	9.3-10.8	2219	29.0	27.8-30.3	90	0.5	0.4-0.6	39	0.5	0.4-0.7	51	0.5	0.3-0.6
2011	3080	20.3	19.5-21.0	776	10.4	9.7-11.2	2304	29.8	28.6-31.1	101	0.6	0.5-0.7	49	0.7	0.5-0.9	52	0.5	0.4-0.6
2012	3170	20.9	20.1-21.6	835	11.0	10.3-11.8	2335	30.6	29.3-31.9	92	0.5	0.4-0.6	41	0.5	0.4-0.7	51	0.5	0.4-0.7
2013	3308	21.8	21.0-22.5	914	12.1	11.3-12.9	2394	31.4	30.1-32.7	102	0.6	0.4-0.7	46	0.6	0.4-0.8	56	0.5	0.4-0.7
2014	3238	21.2	20.5-22.0	871	11.4	10.6-12.2	2367	31.0	29.7-32.3	86	0.4	0.4-0.5	38	0.5	0.3-0.6	48	0.4	0.3-0.6
2015	3217	21.3	20.5-22.1	860	11.2	10.5-12.0	2357	31.3	30.0-32.6	97	0.5	0.4-0.6	45	0.5	0.4-0.7	52	0.5	0.4-0.6
2016	2989	19.7	19.0-20.5	797	10.3	9.6-11.1	2192	29.0	27.8-30.3	102	0.6	0.4-0.7	39	0.5	0.4-0.7	63	0.6	0.5-0.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Hodgkin lymphoma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	564	3.5	3.2-3.8	307	4.1	3.6-4.6	257	3.0	2.7-3.4	217	1.4	1.2-1.6	126	1.8	1.5-2.2	91	1.1	0.9-1.3
1977	534	3.4	3.1-3.7	301	4.1	3.6-4.6	233	2.8	2.5-3.2	225	1.5	1.3-1.7	128	1.9	1.6-2.3	97	1.2	1.0-1.5
1978	568	3.6	3.3-3.9	297	4.1	3.6-4.6	271	3.2	2.8-3.6	197	1.3	1.1-1.5	109	1.6	1.3-1.9	88	1.1	0.8-1.3
1979	576	3.8	3.4-4.1	314	4.5	4.0-5.0	262	3.1	2.8-3.6	170	1.1	1.0-1.3	97	1.5	1.2-1.8	73	0.9	0.7-1.1
1980	597	3.9	3.5-4.2	327	4.6	4.1-5.2	270	3.2	2.8-3.6	203	1.4	1.2-1.6	112	1.7	1.4-2.1	91	1.1	0.8-1.3
1981	564	3.6	3.3-4.0	300	4.1	3.7-4.6	264	3.2	2.8-3.6	176	1.2	1.0-1.4	98	1.5	1.2-1.8	78	0.9	0.7-1.2
1982	580	3.8	3.4-4.1	300	4.3	3.8-4.8	280	3.3	2.9-3.8	182	1.2	1.0-1.4	97	1.4	1.1-1.7	85	1.0	0.8-1.3
1983	543	3.6	3.3-3.9	310	4.4	3.9-4.9	233	2.8	2.5-3.2	152	1.0	0.8-1.2	88	1.3	1.0-1.6	64	0.7	0.5-0.9
1984	567	3.7	3.4-4.0	302	4.3	3.8-4.8	265	3.2	2.8-3.6	157	1.1	0.9-1.2	93	1.4	1.1-1.7	64	0.7	0.6-1.0
1985	605	4.0	3.6-4.3	330	4.6	4.1-5.2	275	3.4	3.0-3.8	155	1.0	0.8-1.2	84	1.2	0.9-1.5	71	0.8	0.6-1.1
1986	562	3.6	3.3-3.9	306	4.2	3.7-4.7	256	3.1	2.7-3.5	174	1.1	1.0-1.3	99	1.5	1.2-1.8	75	0.9	0.7-1.1
1987	522	3.4	3.1-3.8	268	3.8	3.3-4.3	254	3.2	2.8-3.6	140	0.9	0.8-1.1	86	1.2	1.0-1.5	54	0.7	0.5-0.9
1988	561	3.7	3.4-4.0	313	4.4	3.9-5.0	248	3.1	2.7-3.5	119	0.8	0.7-1.0	76	1.1	0.9-1.4	43	0.5	0.4-0.7
1989	555	3.7	3.4-4.0	310	4.3	3.9-4.9	245	3.1	2.7-3.5	149	1.0	0.9-1.2	95	1.4	1.2-1.8	54	0.7	0.5-0.9
1990	525	3.6	3.3-3.9	286	4.1	3.6-4.6	239	3.0	2.7-3.5	137	0.9	0.8-1.1	84	1.3	1.0-1.6	53	0.6	0.5-0.8
1991	545	3.7	3.4-4.0	290	4.1	3.7-4.7	255	3.3	2.9-3.7	110	0.7	0.6-0.9	61	0.9	0.7-1.2	49	0.6	0.4-0.8
1992	559	3.8	3.5-4.1	304	4.3	3.9-4.9	255	3.2	2.9-3.7	128	0.8	0.7-1.0	64	0.9	0.7-1.2	64	0.8	0.6-1.0
1993	505	3.5	3.2-3.8	250	3.6	3.2-4.1	255	3.3	2.9-3.8	102	0.7	0.5-0.8	55	0.8	0.6-1.1	47	0.6	0.4-0.8
1994	566	3.8	3.5-4.2	297	4.3	3.8-4.8	269	3.5	3.1-3.9	113	0.7	0.6-0.9	55	0.8	0.6-1.1	58	0.7	0.5-0.9
1995	513	3.5	3.2-3.9	276	3.9	3.5-4.4	237	3.2	2.8-3.6	100	0.7	0.5-0.8	50	0.7	0.5-1.0	50	0.6	0.4-0.8
1996	493	3.4	3.1-3.7	251	3.6	3.2-4.1	242	3.3	2.8-3.7	89	0.6	0.5-0.7	45	0.7	0.5-0.9	44	0.6	0.4-0.7
1997	482	3.4	3.1-3.7	267	3.8	3.4-4.3	215	2.9	2.6-3.4	96	0.6	0.5-0.8	53	0.8	0.6-1.0	43	0.5	0.4-0.7
1998	538	3.7	3.4-4.1	273	3.9	3.5-4.4	265	3.6	3.1-4.0	93	0.6	0.5-0.8	48	0.7	0.5-0.9	45	0.6	0.4-0.8
1999	501	3.5	3.2-3.8	269	3.8	3.4-4.3	232	3.1	2.7-3.5	83	0.5	0.4-0.7	39	0.6	0.4-0.8	44	0.5	0.4-0.7
2000	535	3.7	3.4-4.0	290	4.2	3.7-4.7	245	3.3	2.9-3.7	86	0.6	0.4-0.7	44	0.7	0.5-0.9	42	0.5	0.3-0.7
2001	536	3.7	3.4-4.0	279	4.0	3.5-4.5	257	3.5	3.0-3.9	85	0.6	0.4-0.7	45	0.6	0.5-0.9	40	0.5	0.3-0.6
2002	583	4.0	3.7-4.4	325	4.7	4.2-5.2	258	3.5	3.0-3.9	93	0.6	0.5-0.7	48	0.7	0.5-0.9	45	0.5	0.4-0.7
2003	517	3.6	3.3-3.9	283	4.0	3.6-4.5	234	3.1	2.7-3.6	70	0.5	0.4-0.6	32	0.5	0.3-0.6	38	0.5	0.3-0.6
2004	542	3.8	3.4-4.1	278	4.0	3.5-4.5	264	3.6	3.2-4.0	75	0.5	0.4-0.6	44	0.6	0.4-0.8	31	0.4	0.2-0.5
2005	553	3.9	3.5-4.2	304	4.4	3.9-4.9	249	3.4	3.0-3.9	70	0.4	0.3-0.6	34	0.5	0.3-0.7	36	0.4	0.3-0.6
2006	527	3.6	3.3-4.0	294	4.2	3.7-4.7	233	3.1	2.7-3.6	71	0.5	0.4-0.6	41	0.6	0.4-0.8	30	0.4	0.2-0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Hodgkin lymphoma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	534	3.7	3.4-4.0	266	3.8	3.3-4.3	268	3.6	3.2-4.1	84	0.5	0.4-0.7	42	0.6	0.4-0.8	42	0.5	0.3-0.6
2008	519	3.6	3.3-3.9	273	3.8	3.4-4.3	246	3.3	2.9-3.8	84	0.5	0.4-0.6	50	0.7	0.5-0.9	34	0.4	0.2-0.5
2009	524	3.6	3.3-4.0	291	4.1	3.6-4.6	233	3.2	2.8-3.7	63	0.4	0.3-0.5	35	0.5	0.3-0.7	28	0.3	0.2-0.5
2010	509	3.4	3.2-3.8	283	3.9	3.5-4.4	226	3.0	2.6-3.4	66	0.4	0.3-0.5	38	0.5	0.4-0.7	28	0.3	0.2-0.5
2011	519	3.6	3.3-3.9	288	4.1	3.6-4.6	231	3.2	2.8-3.6	69	0.4	0.3-0.5	43	0.6	0.4-0.8	26	0.3	0.2-0.4
2012	517	3.6	3.3-3.9	298	4.2	3.7-4.7	219	3.1	2.7-3.5	67	0.4	0.3-0.5	31	0.4	0.3-0.6	36	0.4	0.3-0.5
2013	493	3.4	3.1-3.7	241	3.4	2.9-3.8	252	3.4	3.0-3.9	55	0.3	0.2-0.4	27	0.4	0.2-0.5	28	0.3	0.2-0.4
2014	548	3.7	3.4-4.0	293	4.1	3.6-4.6	255	3.3	2.9-3.8	59	0.3	0.3-0.5	25	0.3	0.2-0.5	34	0.4	0.2-0.5
2015	472	3.3	3.0-3.6	271	3.8	3.4-4.3	201	2.8	2.4-3.2	53	0.3	0.2-0.4	26	0.4	0.2-0.5	27	0.3	0.2-0.4
2016	514	3.5	3.2-3.8	257	3.5	3.1-4.0	257	3.5	3.0-3.9	64	0.4	0.3-0.5	37	0.5	0.3-0.6	27	0.3	0.2-0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Non-Hodgkin lymphomas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1755	11.6	11.0-12.1	901	13.9	13.0-14.9	854	9.9	9.2-10.6	970	6.4	6.0-6.8	519	8.2	7.5-9.0	451	5.1	4.7-5.7
1977	1782	11.7	11.1-12.2	908	13.9	12.9-14.9	874	10.0	9.4-10.7	972	6.4	6.0-6.8	475	7.4	6.7-8.1	497	5.6	5.1-6.1
1978	1900	12.5	11.9-13.1	991	15.1	14.1-16.1	909	10.4	9.8-11.2	989	6.5	6.1-7.0	510	8.1	7.4-8.9	479	5.4	5.0-6.0
1979	1860	12.4	11.8-13.0	913	14.4	13.4-15.4	947	10.8	10.2-11.6	994	6.6	6.2-7.0	521	8.3	7.6-9.1	473	5.3	4.9-5.9
1980	2018	13.3	12.7-13.9	1016	15.6	14.6-16.6	1002	11.5	10.8-12.2	1019	6.7	6.3-7.1	503	7.8	7.1-8.5	516	5.8	5.3-6.4
1981	2028	13.4	12.8-14.0	986	15.4	14.4-16.4	1042	11.8	11.1-12.6	1014	6.6	6.2-7.1	505	8.0	7.3-8.8	509	5.6	5.1-6.1
1982	2125	14.0	13.4-14.6	1053	16.3	15.3-17.3	1072	12.2	11.5-13.0	1057	7.0	6.6-7.4	528	8.6	7.9-9.4	529	5.9	5.4-6.4
1983	2225	14.6	14.0-15.3	1117	17.4	16.4-18.5	1108	12.5	11.7-13.2	1119	7.4	7.0-7.8	573	9.1	8.3-9.9	546	6.1	5.6-6.7
1984	2284	15.1	14.4-15.7	1183	18.8	17.7-19.9	1101	12.3	11.6-13.1	1067	7.0	6.6-7.4	568	9.2	8.5-10.1	499	5.4	5.0-5.9
1985	2450	16.2	15.5-16.8	1300	20.3	19.2-21.5	1150	12.9	12.2-13.7	1181	7.7	7.3-8.2	625	10.0	9.2-10.8	556	6.1	5.6-6.6
1986	2574	17.0	16.3-17.7	1373	21.3	20.1-22.4	1201	13.6	12.8-14.4	1176	7.8	7.3-8.2	644	10.4	9.6-11.2	532	5.8	5.3-6.3
1987	2552	16.9	16.2-17.6	1365	21.3	20.1-22.5	1187	13.4	12.6-14.2	1192	7.8	7.4-8.3	598	9.5	8.7-10.3	594	6.5	6.0-7.0
1988	2590	17.2	16.5-17.9	1349	20.9	19.8-22.1	1241	14.3	13.5-15.1	1239	8.1	7.7-8.6	616	9.9	9.1-10.7	623	6.8	6.2-7.3
1989	2687	18.0	17.3-18.7	1470	22.7	21.5-23.9	1217	14.1	13.3-14.9	1243	8.2	7.7-8.6	677	10.8	10.0-11.7	566	6.2	5.7-6.7
1990	2893	19.1	18.4-19.9	1587	24.4	23.2-25.7	1306	14.8	14.0-15.6	1283	8.4	7.9-8.9	648	10.4	9.6-11.2	635	6.9	6.4-7.5
1991	2998	19.9	19.2-20.6	1628	25.4	24.2-26.7	1370	15.7	14.8-16.5	1280	8.4	7.9-8.8	674	10.8	10.0-11.7	606	6.5	6.0-7.1
1992	2881	19.1	18.4-19.8	1548	23.7	22.5-24.9	1333	15.2	14.4-16.1	1335	8.7	8.2-9.1	676	10.8	10.0-11.7	659	7.1	6.6-7.7
1993	2963	19.6	18.9-20.3	1622	24.6	23.4-25.9	1341	15.3	14.5-16.2	1243	8.0	7.6-8.5	631	10.0	9.2-10.8	612	6.6	6.0-7.1
1994	3245	21.4	20.6-22.1	1801	27.3	26.0-28.6	1444	16.4	15.6-17.3	1419	9.1	8.7-9.6	728	11.4	10.6-12.3	691	7.4	6.9-8.0
1995	3200	21.1	20.3-21.8	1715	26.1	24.8-27.3	1485	17.0	16.1-17.9	1409	9.1	8.6-9.5	731	11.6	10.8-12.5	678	7.3	6.7-7.8
1996	3349	21.9	21.2-22.7	1792	27.2	25.9-28.5	1557	17.8	16.9-18.8	1426	9.1	8.7-9.6	741	11.7	10.9-12.6	685	7.4	6.8-7.9
1997	3353	21.8	21.1-22.6	1735	26.1	24.9-27.3	1618	18.3	17.4-19.2	1478	9.4	8.9-9.9	747	11.5	10.7-12.4	731	7.8	7.3-8.4
1998	3315	21.4	20.7-22.2	1703	25.3	24.1-26.5	1612	18.3	17.4-19.3	1382	8.7	8.3-9.2	662	10.0	9.3-10.8	720	7.6	7.1-8.2
1999	3358	21.5	20.8-22.2	1694	25.1	23.9-26.3	1664	18.7	17.8-19.7	1312	8.2	7.7-8.6	637	9.8	9.0-10.6	675	7.1	6.6-7.7
2000	3375	21.4	20.7-22.1	1720	25.2	24.0-26.4	1655	18.4	17.5-19.3	1354	8.4	7.9-8.8	666	10.1	9.3-10.9	688	7.2	6.7-7.8
2001	3350	21.1	20.4-21.8	1731	25.3	24.1-26.5	1619	17.9	17.0-18.8	1261	7.7	7.3-8.2	688	10.2	9.5-11.0	573	5.8	5.3-6.3
2002	3403	21.3	20.6-22.1	1825	26.2	25.0-27.5	1578	17.4	16.6-18.3	1223	7.4	7.0-7.9	654	9.7	9.0-10.5	569	5.9	5.4-6.4
2003	3598	22.4	21.7-23.1	1844	26.3	25.1-27.5	1754	19.2	18.3-20.2	1134	6.8	6.4-7.2	572	8.5	7.8-9.2	562	5.6	5.1-6.1
2004	3635	22.5	21.8-23.3	1917	27.2	26.0-28.4	1718	19.0	18.1-19.9	1144	6.8	6.4-7.2	561	8.2	7.5-8.9	583	5.8	5.3-6.3
2005	3735	23.1	22.3-23.8	1958	27.6	26.4-28.8	1777	19.5	18.6-20.4	1111	6.6	6.2-7.0	582	8.4	7.8-9.2	529	5.3	4.8-5.8
2006	3856	23.8	23.0-24.5	2037	28.5	27.3-29.8	1819	19.8	18.9-20.8	1207	7.2	6.8-7.6	617	8.8	8.1-9.6	590	5.8	5.3-6.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Non-Hodgkin lymphomas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	3839	23.5	22.7-24.2	2035	28.2	27.0-29.5	1804	19.8	18.8-20.7	1088	6.4	6.0-6.8	596	8.4	7.7-9.1	492	4.9	4.5-5.4
2008	3898	23.6	22.9-24.4	2069	28.3	27.1-29.6	1829	20.0	19.1-20.9	1130	6.6	6.2-7.0	601	8.4	7.7-9.1	529	5.3	4.8-5.8
2009	3982	23.9	23.2-24.7	2109	28.6	27.4-29.9	1873	20.2	19.3-21.2	1080	6.2	5.8-6.6	599	8.2	7.6-8.9	481	4.7	4.3-5.2
2010	3946	23.5	22.7-24.2	2080	28.0	26.8-29.2	1866	19.8	18.8-20.7	1082	6.1	5.8-6.5	564	7.7	7.1-8.4	518	5.0	4.6-5.5
2011	3915	23.2	22.4-23.9	2108	27.9	26.7-29.2	1807	19.3	18.4-20.3	1080	6.1	5.7-6.5	565	7.6	6.9-8.2	515	5.0	4.6-5.5
2012	3961	23.1	22.3-23.8	2154	28.2	27.0-29.4	1807	19.0	18.1-20.0	1042	5.8	5.5-6.2	575	7.7	7.1-8.4	467	4.4	4.0-4.8
2013	4062	23.4	22.7-24.2	2206	28.5	27.3-29.8	1856	19.3	18.4-20.2	981	5.4	5.1-5.8	530	6.9	6.4-7.6	451	4.2	3.8-4.7
2014	4080	23.2	22.5-24.0	2229	28.5	27.3-29.7	1851	19.0	18.1-19.9	1051	5.7	5.3-6.0	579	7.5	6.9-8.1	472	4.3	3.9-4.7
2015	4050	22.9	22.2-23.6	2222	27.8	26.6-29.0	1828	18.9	18.0-19.8	990	5.4	5.0-5.7	562	7.2	6.6-7.8	428	4.0	3.6-4.4
2016	4055	22.7	22.0-23.4	2145	26.5	25.4-27.7	1910	19.6	18.7-20.6	1064	5.6	5.2-5.9	571	7.1	6.5-7.7	493	4.5	4.1-4.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Myeloma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	638	4.2	3.8-4.5	322	5.2	4.6-5.8	316	3.5	3.1-3.9	409	2.8	2.5-3.0	211	3.6	3.1-4.1	198	2.3	2.0-2.6
1977	629	4.1	3.8-4.5	316	5.1	4.6-5.8	313	3.4	3.1-3.9	410	2.6	2.4-2.9	203	3.2	2.7-3.7	207	2.3	2.0-2.6
1978	643	4.2	3.9-4.6	302	4.9	4.3-5.5	341	3.8	3.4-4.3	400	2.6	2.4-2.9	194	3.1	2.7-3.6	206	2.3	2.0-2.6
1979	684	4.5	4.2-4.9	326	5.3	4.7-6.0	358	4.0	3.6-4.4	431	2.9	2.6-3.1	202	3.3	2.9-3.9	229	2.5	2.2-2.9
1980	651	4.3	3.9-4.6	326	5.4	4.8-6.0	325	3.6	3.2-4.0	478	3.1	2.8-3.4	228	3.8	3.3-4.3	250	2.7	2.4-3.1
1981	751	4.9	4.6-5.3	358	5.8	5.2-6.5	393	4.4	3.9-4.8	457	3.0	2.7-3.3	218	3.7	3.2-4.2	239	2.6	2.3-3.0
1982	711	4.6	4.2-4.9	358	5.8	5.2-6.4	353	3.8	3.4-4.3	445	2.9	2.7-3.2	212	3.5	3.0-4.0	233	2.5	2.2-2.9
1983	710	4.6	4.3-5.0	377	6.3	5.6-7.0	333	3.6	3.2-4.0	458	3.0	2.7-3.3	223	3.7	3.2-4.3	235	2.5	2.2-2.9
1984	713	4.7	4.3-5.0	367	6.0	5.4-6.6	346	3.8	3.4-4.2	487	3.1	2.9-3.4	239	3.8	3.4-4.4	248	2.7	2.3-3.0
1985	735	4.8	4.4-5.1	378	6.1	5.5-6.8	357	3.9	3.5-4.3	488	3.1	2.9-3.4	257	4.2	3.7-4.8	231	2.4	2.1-2.8
1986	698	4.5	4.2-4.9	373	6.0	5.4-6.6	325	3.5	3.1-3.9	487	3.1	2.8-3.4	237	3.9	3.4-4.5	250	2.6	2.3-3.0
1987	752	4.9	4.5-5.3	386	6.3	5.7-7.0	366	3.9	3.5-4.3	494	3.2	2.9-3.5	257	4.2	3.7-4.8	237	2.6	2.2-2.9
1988	721	4.7	4.4-5.1	379	6.2	5.6-6.9	342	3.7	3.3-4.1	502	3.2	2.9-3.5	273	4.6	4.0-5.1	229	2.4	2.1-2.7
1989	767	5.0	4.7-5.4	349	5.7	5.1-6.4	418	4.6	4.2-5.1	470	3.0	2.7-3.3	236	3.8	3.3-4.3	234	2.5	2.2-2.8
1990	807	5.2	4.8-5.5	413	6.5	5.9-7.2	394	4.3	3.9-4.8	461	2.9	2.7-3.2	232	3.8	3.4-4.4	229	2.4	2.1-2.7
1991	749	4.8	4.5-5.2	379	6.0	5.4-6.7	370	4.0	3.6-4.5	479	3.1	2.8-3.4	231	3.7	3.3-4.3	248	2.6	2.3-3.0
1992	829	5.3	4.9-5.7	427	6.8	6.2-7.5	402	4.3	3.9-4.8	502	3.2	2.9-3.5	259	4.3	3.8-4.8	243	2.6	2.2-2.9
1993	818	5.3	4.9-5.6	403	6.4	5.7-7.0	415	4.5	4.1-5.0	562	3.6	3.3-3.9	279	4.5	4.0-5.1	283	2.9	2.6-3.3
1994	794	5.1	4.8-5.5	403	6.3	5.7-7.0	391	4.3	3.9-4.7	565	3.6	3.3-3.9	256	4.1	3.6-4.6	309	3.3	3.0-3.7
1995	837	5.4	5.0-5.8	430	6.8	6.1-7.5	407	4.5	4.0-4.9	553	3.5	3.2-3.8	282	4.5	4.0-5.0	271	2.9	2.5-3.2
1996	865	5.5	5.1-5.9	439	6.9	6.3-7.6	426	4.6	4.2-5.1	536	3.4	3.1-3.7	261	4.2	3.7-4.8	275	2.9	2.6-3.3
1997	834	5.3	5.0-5.7	408	6.3	5.7-6.9	426	4.7	4.2-5.2	508	3.2	2.9-3.5	249	4.0	3.5-4.5	259	2.8	2.4-3.1
1998	955	6.0	5.7-6.4	483	7.3	6.7-8.0	472	5.1	4.7-5.6	526	3.3	3.0-3.6	261	4.0	3.6-4.6	265	2.8	2.4-3.1
1999	960	6.0	5.6-6.4	481	7.3	6.6-8.0	479	5.2	4.7-5.7	543	3.3	3.1-3.6	276	4.2	3.8-4.8	267	2.7	2.4-3.1
2000	955	5.9	5.6-6.3	494	7.3	6.7-8.0	461	5.0	4.5-5.4	578	3.5	3.2-3.8	290	4.4	3.9-5.0	288	3.0	2.6-3.3
2001	940	5.8	5.4-6.2	528	7.8	7.2-8.5	412	4.3	3.9-4.8	485	2.9	2.7-3.2	261	3.9	3.5-4.4	224	2.3	2.0-2.6
2002	1021	6.2	5.9-6.6	563	8.2	7.6-8.9	458	4.8	4.4-5.3	537	3.2	3.0-3.5	289	4.4	3.9-4.9	248	2.5	2.2-2.9
2003	990	6.0	5.6-6.4	544	7.9	7.2-8.6	446	4.7	4.3-5.2	506	3.0	2.8-3.3	274	4.1	3.6-4.6	232	2.3	2.0-2.6
2004	1025	6.2	5.8-6.6	552	7.8	7.2-8.5	473	5.0	4.5-5.5	485	2.9	2.6-3.1	256	3.8	3.3-4.2	229	2.3	2.0-2.6
2005	1017	6.1	5.7-6.5	545	7.7	7.0-8.4	472	4.9	4.5-5.4	513	3.0	2.8-3.3	264	3.8	3.4-4.3	249	2.5	2.2-2.8
2006	1054	6.3	6.0-6.7	574	8.0	7.4-8.7	480	5.0	4.6-5.5	486	2.9	2.6-3.2	281	4.0	3.6-4.5	205	2.1	1.8-2.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Myeloma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	1098	6.6	6.2-7.0	597	8.3	7.7-9.0	501	5.3	4.8-5.8	497	2.9	2.7-3.2	261	3.7	3.2-4.1	236	2.3	2.0-2.7
2008	1126	6.6	6.2-7.0	624	8.5	7.9-9.2	502	5.2	4.8-5.7	496	2.9	2.6-3.1	268	3.8	3.3-4.3	228	2.3	2.0-2.6
2009	1204	7.0	6.6-7.4	649	8.8	8.1-9.5	555	5.6	5.2-6.2	497	2.8	2.6-3.1	269	3.7	3.3-4.2	228	2.2	1.9-2.5
2010	1215	7.0	6.6-7.4	662	8.7	8.0-9.4	553	5.7	5.3-6.3	534	3.0	2.8-3.3	271	3.7	3.3-4.2	263	2.5	2.2-2.9
2011	1269	7.3	6.9-7.7	728	9.5	8.8-10.2	541	5.5	5.0-6.0	481	2.7	2.4-2.9	250	3.3	2.9-3.7	231	2.2	1.9-2.5
2012	1235	6.9	6.5-7.3	697	8.8	8.2-9.5	538	5.5	5.0-6.0	517	2.8	2.6-3.1	256	3.4	3.0-3.8	261	2.5	2.2-2.8
2013	1314	7.3	6.9-7.7	731	9.2	8.5-9.9	583	5.9	5.4-6.4	491	2.7	2.4-2.9	266	3.4	3.0-3.9	225	2.1	1.9-2.4
2014	1413	7.8	7.4-8.2	777	9.6	9.0-10.4	636	6.3	5.8-6.9	538	2.9	2.6-3.1	291	3.6	3.2-4.1	247	2.3	2.0-2.6
2015	1279	7.0	6.6-7.4	704	8.7	8.0-9.4	575	5.7	5.2-6.2	519	2.8	2.5-3.0	298	3.7	3.3-4.2	221	2.1	1.8-2.4
2016	1317	7.2	6.8-7.6	770	9.3	8.6-10.0	547	5.5	5.0-6.0	536	2.8	2.6-3.0	275	3.3	2.9-3.8	261	2.4	2.1-2.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Leukemias

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1871	12.6	12.0-13.2	1055	17.1	16.0-18.2	816	9.6	9.0-10.3	1227	8.4	7.9-8.9	678	11.4	10.5-12.4	549	6.5	5.9-7.0
1977	1755	11.9	11.4-12.5	966	15.5	14.5-16.6	789	9.4	8.7-10.1	1243	8.5	8.0-9.0	674	11.0	10.2-12.0	569	6.8	6.2-7.3
1978	1840	12.6	12.0-13.2	1030	16.8	15.8-17.9	810	9.7	9.0-10.4	1203	8.1	7.7-8.6	636	10.4	9.6-11.3	567	6.7	6.1-7.3
1979	1911	12.8	12.2-13.4	1048	16.6	15.6-17.7	863	10.1	9.5-10.8	1246	8.5	8.0-9.0	659	10.9	10.0-11.8	587	7.0	6.4-7.6
1980	2005	13.5	12.9-14.1	1069	17.3	16.2-18.4	936	11.1	10.4-11.8	1256	8.4	8.0-8.9	698	11.4	10.6-12.3	558	6.4	5.9-7.0
1981	1823	12.3	11.7-12.9	1019	16.5	15.4-17.5	804	9.4	8.8-10.1	1248	8.5	8.0-9.0	689	11.4	10.5-12.3	559	6.6	6.0-7.2
1982	1894	12.8	12.2-13.4	1062	17.3	16.3-18.4	832	9.7	9.1-10.4	1295	8.7	8.3-9.2	728	12.2	11.2-13.1	567	6.6	6.1-7.2
1983	1831	12.3	11.7-12.9	1004	16.1	15.1-17.2	827	9.7	9.1-10.4	1219	8.2	7.8-8.7	638	10.7	9.9-11.6	581	6.6	6.1-7.2
1984	1965	13.1	12.6-13.7	1054	17.1	16.1-18.2	911	10.4	9.8-11.2	1211	8.1	7.6-8.5	605	9.9	9.1-10.8	606	6.9	6.3-7.4
1985	1986	13.2	12.6-13.8	1113	17.9	16.8-19.0	873	10.0	9.4-10.7	1206	8.0	7.5-8.5	673	11.1	10.2-12.0	533	6.0	5.4-6.5
1986	1938	13.0	12.4-13.6	1043	16.7	15.7-17.8	895	10.4	9.7-11.1	1194	8.0	7.5-8.5	653	10.6	9.8-11.5	541	6.1	5.6-6.7
1987	1781	11.8	11.3-12.4	934	14.9	13.9-15.9	847	9.8	9.1-10.5	1144	7.6	7.1-8.0	606	9.9	9.1-10.8	538	6.1	5.5-6.6
1988	1835	12.2	11.7-12.8	1009	16.2	15.2-17.3	826	9.6	8.9-10.3	1153	7.6	7.2-8.1	653	10.7	9.9-11.6	500	5.6	5.1-6.1
1989	1898	12.6	12.0-13.1	1045	16.9	15.9-18.0	853	9.7	9.0-10.4	1163	7.7	7.2-8.1	625	10.2	9.4-11.1	538	6.0	5.5-6.6
1990	2051	13.7	13.1-14.3	1109	17.5	16.4-18.6	942	10.9	10.2-11.6	1151	7.6	7.1-8.0	586	9.6	8.8-10.4	565	6.3	5.8-6.9
1991	1941	12.9	12.4-13.5	1067	17.0	15.9-18.0	874	10.1	9.4-10.8	1141	7.5	7.1-7.9	603	9.9	9.1-10.8	538	5.9	5.4-6.5
1992	1985	13.2	12.6-13.8	1116	17.8	16.8-18.9	869	10.0	9.3-10.7	1166	7.6	7.2-8.1	642	10.5	9.7-11.3	524	5.8	5.3-6.3
1993	1938	12.8	12.2-13.4	1039	16.2	15.2-17.2	899	10.5	9.8-11.2	1083	7.1	6.6-7.5	595	9.6	8.8-10.4	488	5.4	5.0-6.0
1994	2150	14.1	13.5-14.7	1170	18.3	17.2-19.4	980	11.1	10.4-11.8	1226	7.9	7.5-8.4	673	10.7	9.9-11.6	553	6.0	5.5-6.5
1995	2048	13.4	12.8-13.9	1114	17.3	16.2-18.3	934	10.7	10.0-11.4	1213	7.8	7.4-8.3	671	10.9	10.0-11.7	542	5.9	5.4-6.4
1996	2127	13.9	13.3-14.5	1223	18.7	17.7-19.8	904	10.3	9.6-11.0	1235	7.9	7.5-8.4	661	10.3	9.6-11.2	574	6.3	5.8-6.9
1997	2183	14.2	13.6-14.8	1180	18.1	17.1-19.2	1003	11.3	10.6-12.0	1204	7.7	7.2-8.1	669	10.5	9.7-11.3	535	5.7	5.2-6.2
1998	2171	14.0	13.4-14.6	1246	18.7	17.7-19.8	925	10.4	9.7-11.1	1190	7.5	7.1-8.0	659	10.1	9.3-10.9	531	5.7	5.2-6.2
1999	2389	15.3	14.7-15.9	1306	19.7	18.7-20.8	1083	12.1	11.4-12.8	1260	7.9	7.4-8.3	685	10.5	9.7-11.3	575	6.1	5.6-6.6
2000	2443	15.5	14.8-16.1	1348	20.2	19.1-21.3	1095	12.0	11.3-12.8	1222	7.6	7.2-8.0	655	9.9	9.2-10.7	567	6.0	5.5-6.5
2001	2425	15.3	14.7-15.9	1319	19.5	18.5-20.6	1106	12.2	11.5-13.0	1214	7.5	7.0-7.9	648	9.8	9.1-10.6	566	5.9	5.4-6.4
2002	2271	14.3	13.7-14.9	1295	18.9	17.9-20.0	976	10.8	10.1-11.5	1208	7.4	7.0-7.8	661	9.9	9.1-10.7	547	5.7	5.2-6.2
2003	2497	15.6	15.0-16.2	1380	20.1	19.0-21.2	1117	12.3	11.6-13.1	1155	7.0	6.6-7.4	648	9.7	8.9-10.4	507	5.2	4.8-5.7
2004	2525	15.7	15.1-16.3	1389	20.0	19.0-21.1	1136	12.5	11.8-13.3	1260	7.6	7.2-8.0	691	10.2	9.4-11.0	569	5.9	5.4-6.4
2005	2578	16.0	15.4-16.6	1433	20.4	19.4-21.5	1145	12.5	11.7-13.2	1218	7.3	6.9-7.7	693	10.1	9.4-10.9	525	5.3	4.9-5.8
2006	2545	15.8	15.2-16.5	1448	20.7	19.7-21.9	1097	12.1	11.4-12.8	1230	7.4	7.0-7.8	671	9.7	9.0-10.5	559	5.7	5.2-6.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Whites
Leukemias

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	2650	16.4	15.8-17.1	1522	21.4	20.3-22.5	1128	12.6	11.9-13.4	1178	7.1	6.7-7.5	665	9.6	8.9-10.3	513	5.3	4.9-5.8
2008	2581	15.9	15.3-16.5	1469	20.5	19.4-21.6	1112	12.4	11.7-13.2	1219	7.2	6.8-7.6	683	9.7	9.0-10.4	536	5.4	5.0-6.0
2009	2774	16.8	16.2-17.5	1594	21.9	20.8-23.0	1180	12.9	12.2-13.7	1221	7.2	6.8-7.7	689	9.7	9.0-10.5	532	5.4	5.0-5.9
2010	2922	17.6	17.0-18.3	1685	23.0	21.9-24.1	1237	13.4	12.6-14.2	1154	6.7	6.3-7.1	651	9.0	8.3-9.8	503	5.0	4.6-5.5
2011	3049	18.1	17.5-18.8	1732	23.3	22.2-24.4	1317	14.1	13.3-14.9	1223	7.0	6.6-7.4	701	9.5	8.8-10.2	522	5.1	4.7-5.6
2012	2910	17.1	16.5-17.8	1695	22.4	21.3-23.5	1215	13.1	12.4-13.9	1196	6.8	6.4-7.2	695	9.4	8.7-10.2	501	4.8	4.4-5.3
2013	3106	18.1	17.5-18.8	1823	23.9	22.7-25.0	1283	13.7	12.9-14.5	1205	6.7	6.4-7.1	691	9.2	8.5-9.9	514	5.0	4.6-5.5
2014	3108	17.9	17.3-18.6	1795	23.1	22.0-24.2	1313	13.8	13.1-14.7	1149	6.3	5.9-6.7	631	8.2	7.6-8.9	518	4.9	4.5-5.4
2015	3107	17.8	17.1-18.4	1795	22.9	21.8-24.0	1312	13.7	12.9-14.5	1187	6.4	6.1-6.8	680	8.7	8.1-9.4	507	4.7	4.3-5.2
2016	2973	16.7	16.1-17.3	1757	21.9	20.8-23.0	1216	12.4	11.7-13.2	1177	6.4	6.0-6.8	661	8.4	7.8-9.1	516	4.9	4.5-5.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
All Invasive Malignant Tumors

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	23673	388.5	383.4-393.7	11209	447.8	439.1-456.6	12464	359.2	352.6-365.9	13251	217.6	213.8-221.4	6744	275.0	268.1-282.0	6507	183.0	178.4-187.7
1977	23287	389.3	384.2-394.5	11134	454.1	445.3-463.0	12153	356.3	349.7-363.0	13069	216.7	212.9-220.6	6612	273.5	266.7-280.5	6457	182.8	178.2-187.5
1978	23145	392.4	387.2-397.7	11072	459.5	450.6-468.5	12073	358.4	351.7-365.1	12490	210.0	206.2-213.8	6360	267.3	260.5-274.3	6130	175.8	171.3-180.4
1979	23453	402.7	397.4-408.1	11225	473.0	464.0-482.2	12228	368.0	361.2-374.9	12323	209.5	205.7-213.3	6199	264.6	257.8-271.5	6124	177.1	172.5-181.8
1980	23912	415.9	410.5-421.4	11586	496.0	486.7-505.5	12326	375.4	368.5-382.5	12604	215.8	211.9-219.7	6332	275.9	268.9-283.0	6272	181.0	176.4-185.7
1981	23476	412.2	406.8-417.7	11111	481.8	472.6-491.2	12365	377.5	370.6-384.6	12085	209.7	205.9-213.6	5935	261.1	254.3-268.1	6150	180.2	175.5-184.9
1982	23652	420.5	415.0-426.1	11177	492.3	483.0-501.8	12475	385.1	378.1-392.3	11758	205.5	201.7-209.4	5831	259.5	252.7-266.5	5927	175.1	170.5-179.8
1983	23749	424.0	418.5-429.6	11321	499.4	490.0-509.0	12428	385.4	378.3-392.6	11567	203.5	199.7-207.3	5680	256.1	249.3-263.1	5887	173.3	168.7-178.0
1984	23766	427.9	422.3-433.6	11328	503.5	494.0-513.2	12438	389.8	382.6-397.0	11662	206.3	202.5-210.2	5792	261.6	254.7-268.6	5870	174.5	169.9-179.2
1985	23931	433.9	428.3-439.6	11636	519.2	509.6-529.0	12295	386.8	379.6-394.0	11308	201.2	197.5-205.1	5653	255.7	248.9-262.7	5655	169.8	165.3-174.5
1986	23837	436.8	431.2-442.6	11520	520.5	510.8-530.3	12317	391.3	384.1-398.6	11001	196.7	193.0-200.5	5321	242.7	236.1-249.5	5680	170.6	166.1-175.3
1987	23389	432.4	426.7-438.1	11249	511.6	502.0-521.4	12140	389.1	381.9-396.4	10922	197.5	193.8-201.3	5389	249.3	242.5-256.2	5533	167.4	162.9-172.0
1988	22732	426.9	421.3-432.7	10734	497.5	487.9-507.2	11998	389.5	382.2-396.8	10932	200.8	197.0-204.7	5374	253.5	246.6-260.5	5558	170.9	166.3-175.6
1989	22475	431.9	426.2-437.7	10734	504.0	494.4-513.8	11741	391.5	384.2-399.0	10457	195.3	191.5-199.2	5071	242.9	236.2-249.8	5386	168.2	163.5-172.9
1990	23461	456.4	450.5-462.4	11231	533.1	523.1-543.2	12230	415.6	408.0-423.3	10336	196.0	192.1-199.8	4991	242.9	236.1-249.9	5345	169.7	165.1-174.5
1991	24186	477.2	471.1-483.4	11722	563.6	553.3-574.0	12464	429.1	421.3-437.0	10113	194.8	190.9-198.6	4890	240.8	234.0-247.7	5223	168.3	163.6-173.1
1992	24308	482.6	476.4-488.8	12037	580.5	570.0-591.1	12271	427.4	419.6-435.3	9914	191.5	187.7-195.4	4859	240.6	233.8-247.6	5055	163.3	158.7-168.0
1993	23395	468.0	462.0-474.1	11599	561.4	551.1-571.8	11796	413.3	405.7-421.1	9803	190.8	187.0-194.7	4819	240.0	233.2-247.0	4984	162.4	157.8-167.1
1994	25147	506.6	500.3-512.9	12202	592.5	582.0-603.2	12945	454.2	446.2-462.4	10303	202.7	198.7-206.7	5061	253.8	246.8-261.0	5242	172.8	168.0-177.7
1995	23789	482.3	476.1-488.5	11362	555.1	544.8-565.5	12427	439.9	432.0-447.9	10147	200.8	196.9-204.8	4829	243.4	236.5-250.4	5318	176.0	171.1-180.9
1996	23749	485.5	479.3-491.8	11278	555.2	544.9-565.6	12471	446.5	438.5-454.6	9864	196.6	192.7-200.5	4605	232.6	225.8-239.4	5259	176.0	171.2-181.0
1997	23754	485.8	479.6-492.1	11296	556.6	546.3-567.0	12458	444.6	436.7-452.6	9416	188.5	184.7-192.3	4478	226.6	219.9-233.3	4938	166.5	161.8-171.3
1998	24137	492.8	486.6-499.1	11550	563.5	553.2-573.9	12587	451.1	443.1-459.2	9359	186.8	183.0-190.6	4494	224.8	218.2-231.5	4865	163.4	158.7-168.1
1999	23828	482.1	476.0-488.3	11348	548.8	538.7-559.1	12480	444.0	436.1-452.0	9150	180.9	177.2-184.7	4409	218.7	212.2-225.2	4741	158.5	153.9-163.1
2000	24378	490.6	484.5-496.9	11929	573.3	563.0-583.7	12449	441.3	433.5-449.3	9085	177.8	174.2-181.5	4319	212.3	206.0-218.8	4766	156.7	152.2-161.3
2001	23813	478.2	472.1-484.4	11606	555.0	545.0-565.3	12207	432.3	424.6-440.2	8867	173.0	169.4-176.7	4287	209.7	203.4-216.1	4580	150.4	146.0-154.9
2002	23510	471.1	465.1-477.2	11543	547.4	537.5-557.5	11967	422.7	415.0-430.5	8602	167.4	163.9-171.0	4150	201.5	195.4-207.8	4452	146.2	141.9-150.7
2003	23213	465.8	459.8-471.9	11334	538.8	528.9-548.9	11879	421.3	413.6-429.1	8533	166.3	162.8-169.9	4037	195.5	189.5-201.6	4496	147.9	143.5-152.4
2004	23289	468.0	461.9-474.1	11314	535.2	525.3-545.2	11975	427.1	419.3-435.0	8285	161.6	158.1-165.2	3993	193.2	187.3-199.4	4292	142.1	137.8-146.5
2005	23489	474.6	468.5-480.8	11498	543.8	533.9-553.9	11991	432.4	424.6-440.4	8144	159.9	156.4-163.5	3962	192.0	186.0-198.1	4182	139.7	135.4-144.1
2006	24136	492.3	486.1-498.6	12028	571.9	561.7-582.3	12108	441.5	433.5-449.7	8063	159.7	156.2-163.3	3891	189.8	183.9-195.9	4172	140.7	136.3-145.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
All Invasive Malignant Tumors

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	24266	495.4	489.1-501.7	12065	571.2	560.9-581.5	12201	448.3	440.2-456.6	8050	159.0	155.5-162.6	3920	189.8	183.9-195.9	4130	139.5	135.1-143.9
2008	24330	494.8	488.5-501.1	11964	562.1	552.0-572.4	12366	453.7	445.5-462.0	7982	157.4	153.9-160.9	3947	189.9	184.0-196.0	4035	136.1	131.8-140.5
2009	24548	497.5	491.2-503.8	11933	556.3	546.2-566.4	12615	462.3	454.0-470.6	7842	154.1	150.7-157.6	3823	183.7	177.9-189.7	4019	135.1	130.8-139.4
2010	23437	472.4	466.3-478.6	11515	531.4	521.6-541.3	11922	436.5	428.5-444.6	8032	155.9	152.5-159.4	3951	186.9	181.0-192.9	4081	135.7	131.5-140.1
2011	23760	472.2	466.1-478.3	11603	526.0	516.4-535.8	12157	440.4	432.4-448.6	7726	148.1	144.8-151.5	3795	176.6	170.9-182.3	3931	128.9	124.8-133.2
2012	23023	452.5	446.6-458.5	11188	497.9	488.6-507.3	11835	427.0	419.1-435.0	7597	144.0	140.7-147.4	3698	168.7	163.2-174.2	3899	128.1	124.0-132.3
2013	23732	460.2	454.3-466.2	11341	497.9	488.7-507.3	12391	441.3	433.3-449.4	7502	140.0	136.8-143.2	3671	165.0	159.6-170.5	3831	123.5	119.5-127.7
2014	23261	445.0	439.2-450.9	11006	473.9	465.0-483.0	12255	432.0	424.1-439.9	7422	136.2	133.1-139.4	3653	161.0	155.7-166.3	3769	119.3	115.4-123.4
2015	23115	437.5	431.8-443.3	11072	469.1	460.3-478.1	12043	423.5	415.7-431.4	7364	133.7	130.6-136.9	3620	156.6	151.5-161.9	3744	118.5	114.6-122.5
2016	22595	423.9	418.2-429.5	10808	452.1	443.5-460.8	11787	412.7	405.1-420.5	8551	153.6	150.3-157.0	4291	182.8	177.3-188.4	4260	134.4	130.3-138.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Oral cavity and pharynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	640	10.4	9.6-11.3	435	17.2	15.6-19.0	205	5.6	4.8-6.4	312	5.1	4.5-5.7	222	8.7	7.5-10.0	90	2.5	2.0-3.1
1977	645	10.6	9.8-11.5	417	16.9	15.3-18.7	228	6.2	5.4-7.1	302	5.0	4.5-5.6	208	8.5	7.4-9.8	94	2.6	2.1-3.2
1978	604	10.3	9.5-11.2	397	16.1	14.5-17.9	207	6.0	5.2-7.0	223	3.7	3.2-4.2	146	5.8	4.9-6.9	77	2.1	1.7-2.7
1979	600	10.3	9.5-11.2	408	16.8	15.1-18.6	192	5.7	4.9-6.6	233	4.0	3.4-4.5	163	6.6	5.6-7.8	70	2.0	1.6-2.6
1980	597	10.6	9.7-11.5	384	16.2	14.6-18.0	213	6.6	5.7-7.6	244	4.3	3.7-4.8	166	7.1	6.0-8.3	78	2.3	1.8-2.9
1981	628	11.1	10.2-12.0	438	18.4	16.7-20.3	190	5.7	4.9-6.7	240	4.2	3.7-4.8	156	6.7	5.6-7.8	84	2.4	1.9-3.0
1982	645	11.6	10.7-12.6	440	19.0	17.2-20.9	205	6.5	5.6-7.5	210	3.7	3.2-4.3	161	6.8	5.8-8.0	49	1.6	1.1-2.1
1983	621	11.2	10.3-12.1	425	18.5	16.7-20.4	196	5.7	4.9-6.6	205	3.5	3.1-4.1	154	6.4	5.4-7.6	51	1.4	1.0-1.9
1984	615	11.3	10.4-12.3	414	18.3	16.6-20.3	201	6.2	5.3-7.2	207	3.8	3.3-4.3	153	6.8	5.7-8.0	54	1.7	1.2-2.2
1985	647	12.0	11.1-13.0	417	18.6	16.8-20.5	230	7.3	6.3-8.4	198	3.5	3.0-4.1	139	6.1	5.1-7.3	59	1.7	1.3-2.2
1986	573	10.7	9.8-11.7	383	17.4	15.7-19.3	190	5.8	5.0-6.8	182	3.3	2.8-3.8	124	5.5	4.6-6.6	58	1.7	1.3-2.3
1987	560	10.8	9.9-11.7	372	16.9	15.2-18.8	188	6.2	5.3-7.2	164	3.1	2.6-3.6	112	5.1	4.2-6.2	52	1.5	1.1-2.1
1988	520	10.1	9.2-11.0	339	15.6	14.0-17.4	181	5.7	4.9-6.6	192	3.8	3.2-4.3	121	5.8	4.8-6.9	71	2.3	1.8-2.9
1989	522	10.4	9.5-11.4	337	16.1	14.4-17.9	185	6.3	5.4-7.3	158	3.1	2.6-3.6	103	4.9	4.0-6.0	55	1.7	1.3-2.2
1990	560	11.3	10.4-12.3	370	17.9	16.1-19.9	190	6.4	5.5-7.4	149	2.9	2.5-3.4	92	4.5	3.6-5.6	57	1.7	1.3-2.3
1991	557	11.5	10.5-12.5	356	17.5	15.7-19.4	201	6.9	6.0-8.0	131	2.5	2.1-3.0	91	4.5	3.6-5.6	40	1.2	0.9-1.7
1992	594	12.4	11.4-13.5	403	19.7	17.8-21.8	191	6.7	5.8-7.8	142	2.8	2.4-3.3	90	4.5	3.6-5.5	52	1.6	1.2-2.1
1993	552	11.6	10.6-12.6	364	18.1	16.3-20.1	188	6.7	5.8-7.8	140	2.9	2.5-3.5	110	5.5	4.5-6.6	30	0.9	0.6-1.4
1994	508	10.6	9.7-11.6	333	16.3	14.6-18.2	175	6.2	5.3-7.2	133	2.8	2.3-3.3	96	4.8	3.9-5.9	37	1.3	0.9-1.8
1995	446	9.4	8.5-10.3	295	14.5	12.9-16.3	151	5.3	4.5-6.3	123	2.5	2.1-3.0	80	4.0	3.2-5.0	43	1.4	1.0-1.9
1996	507	10.7	9.8-11.7	341	16.8	15.0-18.7	166	5.9	5.0-6.9	130	2.7	2.3-3.3	73	3.5	2.8-4.4	57	2.0	1.5-2.7
1997	456	9.6	8.7-10.5	289	14.2	12.6-15.9	167	6.0	5.1-7.0	134	2.8	2.3-3.3	97	4.8	3.9-5.8	37	1.3	0.9-1.8
1998	477	9.9	9.1-10.9	296	14.3	12.7-16.1	181	6.5	5.5-7.5	115	2.3	1.9-2.8	75	3.7	2.9-4.6	40	1.3	0.9-1.7
1999	520	10.7	9.8-11.7	363	17.3	15.6-19.2	157	5.6	4.7-6.5	114	2.3	1.9-2.8	84	4.1	3.2-5.0	30	1.0	0.7-1.4
2000	420	8.6	7.8-9.5	279	13.1	11.6-14.8	141	5.0	4.2-6.0	106	2.2	1.8-2.6	69	3.3	2.6-4.2	37	1.3	0.9-1.7
2001	460	9.3	8.5-10.2	298	14.0	12.5-15.7	162	5.6	4.7-6.5	109	2.2	1.8-2.6	73	3.5	2.7-4.4	36	1.1	0.7-1.5
2002	471	9.6	8.7-10.5	303	14.2	12.6-15.9	168	6.0	5.1-7.0	108	2.2	1.8-2.6	70	3.3	2.6-4.2	38	1.2	0.9-1.7
2003	459	9.3	8.4-10.2	301	14.0	12.5-15.7	158	5.6	4.7-6.5	101	2.0	1.6-2.5	65	3.1	2.4-3.9	36	1.2	0.8-1.7
2004	462	9.3	8.5-10.2	291	13.6	12.0-15.2	171	6.0	5.1-7.0	133	2.6	2.2-3.1	81	3.8	3.0-4.7	52	1.7	1.3-2.2
2005	445	9.0	8.2-9.9	309	14.2	12.6-15.9	136	5.0	4.1-5.9	103	2.0	1.7-2.5	64	3.0	2.3-3.9	39	1.3	0.9-1.7
2006	486	9.9	9.0-10.8	317	14.7	13.1-16.4	169	5.9	5.0-6.9	93	1.9	1.5-2.3	66	3.2	2.5-4.1	27	0.9	0.6-1.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Oral cavity and pharynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	499	10.2	9.3-11.1	325	15.2	13.6-16.9	174	6.3	5.4-7.3	108	2.2	1.8-2.6	70	3.3	2.6-4.2	38	1.2	0.9-1.7
2008	495	10.2	9.3-11.2	315	14.7	13.1-16.4	180	6.7	5.7-7.7	95	1.9	1.6-2.4	68	3.2	2.5-4.0	27	0.9	0.6-1.4
2009	512	10.5	9.6-11.5	372	17.3	15.5-19.1	140	5.1	4.2-6.0	117	2.3	1.9-2.8	83	3.8	3.0-4.7	34	1.2	0.8-1.7
2010	484	9.7	8.9-10.6	308	13.9	12.4-15.6	176	6.4	5.4-7.4	119	2.4	1.9-2.8	82	3.8	3.0-4.7	37	1.3	0.9-1.8
2011	476	9.5	8.6-10.4	330	14.9	13.3-16.6	146	5.1	4.3-6.0	93	1.8	1.4-2.2	64	2.9	2.2-3.7	29	0.9	0.6-1.4
2012	488	9.5	8.7-10.4	322	14.0	12.5-15.6	166	5.8	4.9-6.8	112	2.1	1.8-2.6	74	3.3	2.6-4.1	38	1.3	0.9-1.8
2013	492	9.6	8.7-10.5	340	14.7	13.2-16.4	152	5.3	4.5-6.2	94	1.8	1.4-2.2	66	2.8	2.1-3.6	28	0.9	0.6-1.4
2014	542	10.3	9.4-11.2	335	14.1	12.6-15.7	207	7.1	6.2-8.2	104	2.0	1.6-2.4	72	3.2	2.5-4.0	32	1.0	0.7-1.5
2015	512	9.7	8.9-10.6	349	14.6	13.1-16.2	163	5.7	4.8-6.6	116	2.1	1.8-2.6	84	3.6	2.9-4.5	32	1.0	0.6-1.4
2016	530	9.9	9.1-10.8	363	15.1	13.6-16.8	167	5.7	4.8-6.7	136	2.5	2.1-3.0	94	3.9	3.2-4.8	42	1.4	1.0-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Esophagus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	269	4.3	3.8-4.8	179	6.8	5.8-8.0	90	2.5	2.0-3.0	254	4.1	3.6-4.7	171	6.6	5.6-7.7	83	2.4	1.9-3.0
1977	260	4.2	3.7-4.8	167	6.5	5.5-7.6	93	2.6	2.1-3.2	245	4.0	3.5-4.5	151	6.1	5.1-7.2	94	2.6	2.1-3.2
1978	265	4.4	3.9-5.0	170	6.8	5.8-7.9	95	2.6	2.1-3.2	256	4.2	3.7-4.8	163	6.7	5.7-7.8	93	2.6	2.1-3.2
1979	261	4.3	3.8-4.9	185	7.6	6.5-8.8	76	2.1	1.6-2.6	235	3.9	3.4-4.5	154	6.2	5.3-7.3	81	2.3	1.8-2.9
1980	293	4.9	4.3-5.5	191	8.1	6.9-9.3	102	2.8	2.3-3.4	224	3.8	3.3-4.3	154	6.6	5.5-7.7	70	1.8	1.4-2.3
1981	251	4.3	3.8-4.9	160	6.8	5.7-8.0	91	2.7	2.1-3.3	247	4.2	3.6-4.7	167	7.0	6.0-8.2	80	2.2	1.7-2.8
1982	254	4.4	3.9-5.0	155	6.8	5.7-8.0	99	2.7	2.2-3.3	204	3.5	3.0-4.0	141	6.2	5.2-7.3	63	1.8	1.3-2.3
1983	243	4.3	3.7-4.8	167	7.2	6.1-8.4	76	2.2	1.7-2.8	207	3.6	3.1-4.1	140	6.2	5.2-7.3	67	1.8	1.4-2.4
1984	236	4.1	3.5-4.6	161	7.0	5.9-8.2	75	2.1	1.6-2.6	213	3.7	3.2-4.3	138	5.8	4.9-6.9	75	2.1	1.7-2.7
1985	239	4.3	3.8-4.9	160	7.0	6.0-8.3	79	2.4	1.9-3.1	192	3.4	3.0-4.0	140	6.1	5.1-7.2	52	1.6	1.1-2.1
1986	243	4.4	3.9-5.0	150	6.9	5.8-8.1	93	2.8	2.2-3.4	194	3.5	3.0-4.0	149	6.9	5.8-8.2	45	1.3	0.9-1.8
1987	217	4.0	3.5-4.6	131	5.9	4.9-7.1	86	2.7	2.1-3.3	182	3.3	2.8-3.8	112	5.0	4.1-6.1	70	2.1	1.6-2.6
1988	187	3.4	2.9-3.9	115	5.2	4.3-6.2	72	2.1	1.7-2.7	199	3.7	3.2-4.3	138	6.4	5.4-7.6	61	1.9	1.4-2.4
1989	242	4.5	4.0-5.2	157	7.4	6.2-8.6	85	2.5	2.0-3.2	189	3.5	3.0-4.1	121	5.6	4.7-6.8	68	2.0	1.5-2.5
1990	230	4.4	3.9-5.1	141	6.7	5.6-7.9	89	2.8	2.2-3.5	192	3.6	3.1-4.2	124	6.0	5.0-7.2	68	2.0	1.5-2.5
1991	221	4.3	3.7-4.9	144	7.0	5.9-8.2	77	2.4	1.9-3.0	172	3.3	2.8-3.8	113	5.4	4.4-6.5	59	1.8	1.3-2.3
1992	225	4.5	3.9-5.1	143	6.8	5.7-8.1	82	2.7	2.1-3.4	176	3.4	2.9-4.0	127	6.1	5.1-7.3	49	1.5	1.1-2.0
1993	198	3.9	3.4-4.5	139	6.8	5.7-8.0	59	1.9	1.4-2.5	163	3.2	2.7-3.7	120	5.8	4.8-7.0	43	1.4	1.0-1.9
1994	238	4.8	4.2-5.5	169	8.3	7.1-9.7	69	2.2	1.7-2.8	187	3.7	3.2-4.3	135	6.6	5.5-7.9	52	1.6	1.2-2.1
1995	221	4.5	4.0-5.2	130	6.4	5.4-7.6	91	3.1	2.5-3.8	181	3.7	3.1-4.2	127	6.3	5.2-7.4	54	1.7	1.3-2.3
1996	219	4.4	3.8-5.0	146	7.2	6.1-8.5	73	2.3	1.8-2.9	195	3.9	3.4-4.5	129	6.4	5.3-7.6	66	2.1	1.6-2.7
1997	220	4.4	3.9-5.1	142	7.1	6.0-8.3	78	2.6	2.0-3.2	174	3.5	3.0-4.1	120	6.0	5.0-7.2	54	1.7	1.3-2.3
1998	228	4.6	4.0-5.2	159	7.9	6.7-9.2	69	2.3	1.8-2.9	194	3.9	3.4-4.5	136	6.8	5.7-8.0	58	1.9	1.4-2.5
1999	211	4.2	3.7-4.8	161	7.8	6.6-9.1	50	1.6	1.1-2.1	182	3.6	3.1-4.2	139	6.8	5.7-8.0	43	1.4	1.0-1.9
2000	203	4.0	3.5-4.6	137	6.6	5.5-7.8	66	2.1	1.6-2.6	194	3.8	3.3-4.4	129	6.3	5.2-7.4	65	2.0	1.6-2.6
2001	210	4.1	3.6-4.8	145	7.0	5.9-8.2	65	2.1	1.6-2.7	162	3.2	2.7-3.7	126	6.1	5.0-7.2	36	1.2	0.8-1.7
2002	214	4.3	3.7-4.9	155	7.3	6.2-8.6	59	1.9	1.5-2.5	172	3.3	2.9-3.9	123	5.9	4.9-7.1	49	1.5	1.1-2.0
2003	224	4.4	3.9-5.0	150	7.1	6.0-8.3	74	2.4	1.9-3.0	169	3.3	2.8-3.9	114	5.4	4.4-6.4	55	1.7	1.3-2.3
2004	215	4.2	3.7-4.9	145	6.8	5.7-8.0	70	2.3	1.8-2.9	160	3.1	2.7-3.7	115	5.5	4.5-6.6	45	1.4	1.0-1.9
2005	217	4.4	3.8-5.0	143	6.8	5.7-8.0	74	2.5	1.9-3.1	202	4.0	3.5-4.6	152	7.2	6.1-8.5	50	1.5	1.1-2.1
2006	231	4.7	4.1-5.3	174	8.4	7.2-9.7	57	1.9	1.5-2.5	178	3.6	3.1-4.2	110	5.3	4.4-6.4	68	2.2	1.7-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Esophagus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	211	4.2	3.6-4.8	143	6.7	5.6-7.9	68	2.2	1.7-2.8	190	3.8	3.3-4.4	144	6.9	5.8-8.1	46	1.6	1.1-2.1
2008	244	4.9	4.3-5.5	169	7.8	6.7-9.1	75	2.5	2.0-3.2	174	3.4	2.9-4.0	126	5.8	4.9-7.0	48	1.6	1.2-2.1
2009	197	3.9	3.4-4.5	145	6.7	5.6-7.9	52	1.7	1.2-2.2	188	3.8	3.2-4.3	135	6.4	5.4-7.6	53	1.7	1.3-2.2
2010	184	3.6	3.1-4.1	130	6.0	5.0-7.1	54	1.8	1.3-2.3	161	3.1	2.6-3.6	115	5.3	4.4-6.4	46	1.5	1.1-2.0
2011	184	3.6	3.1-4.2	141	6.5	5.4-7.7	43	1.5	1.1-2.0	152	3.0	2.5-3.5	108	5.0	4.1-6.0	44	1.4	1.0-1.9
2012	165	3.2	2.7-3.7	121	5.3	4.4-6.4	44	1.5	1.0-2.0	127	2.4	2.0-2.9	92	4.2	3.4-5.2	35	1.1	0.7-1.5
2013	209	4.0	3.5-4.6	150	6.6	5.6-7.8	59	1.9	1.4-2.5	148	2.8	2.3-3.3	113	4.9	4.0-5.9	35	1.1	0.8-1.6
2014	209	3.9	3.4-4.5	143	6.2	5.2-7.4	66	2.1	1.6-2.7	176	3.3	2.8-3.8	128	5.7	4.7-6.8	48	1.5	1.1-2.0
2015	186	3.4	3.0-4.0	129	5.4	4.5-6.5	57	1.8	1.4-2.4	148	2.7	2.3-3.2	94	4.1	3.3-5.0	54	1.6	1.2-2.1
2016	190	3.5	3.0-4.0	143	5.9	4.9-6.9	47	1.5	1.1-2.0	176	3.2	2.7-3.7	132	5.5	4.6-6.5	44	1.3	1.0-1.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Stomach

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	852	14.1	13.2-15.1	484	20.0	18.2-22.0	368	10.2	9.2-11.3	669	11.2	10.3-12.1	375	15.8	14.2-17.6	294	8.2	7.3-9.2
1977	846	14.5	13.5-15.5	480	20.6	18.7-22.6	366	10.6	9.5-11.7	648	11.1	10.2-12.0	354	15.6	13.9-17.4	294	8.3	7.4-9.3
1978	822	14.0	13.0-15.0	454	19.7	17.9-21.7	368	10.3	9.2-11.4	602	10.4	9.5-11.2	331	14.7	13.1-16.4	271	7.6	6.7-8.6
1979	837	14.2	13.2-15.2	448	19.1	17.3-21.0	389	11.0	9.9-12.2	597	10.2	9.4-11.1	326	14.3	12.7-16.0	271	7.6	6.7-8.6
1980	850	14.7	13.7-15.7	488	21.4	19.5-23.5	362	10.4	9.3-11.6	591	10.2	9.4-11.1	332	14.7	13.1-16.5	259	7.4	6.5-8.4
1981	792	13.8	12.8-14.8	439	19.7	17.8-21.7	353	10.1	9.0-11.2	583	10.1	9.3-11.0	338	15.3	13.7-17.1	245	6.9	6.1-7.9
1982	740	12.8	11.9-13.8	403	17.9	16.2-19.8	337	9.5	8.5-10.6	495	8.7	7.9-9.5	271	12.4	11.0-14.1	224	6.4	5.6-7.3
1983	814	14.5	13.5-15.5	423	19.3	17.5-21.3	391	11.3	10.2-12.5	511	9.1	8.3-9.9	278	13.1	11.6-14.8	233	6.6	5.8-7.5
1984	741	13.0	12.1-14.0	399	17.7	16.0-19.6	342	9.9	8.8-11.0	492	8.6	7.9-9.5	251	11.4	10.0-13.0	241	6.8	6.0-7.8
1985	708	12.6	11.7-13.6	400	18.4	16.6-20.3	308	9.0	8.0-10.1	498	8.9	8.1-9.8	277	12.6	11.1-14.2	221	6.5	5.7-7.5
1986	666	11.9	11.0-12.9	360	16.6	14.9-18.5	306	8.9	7.9-10.0	406	7.2	6.5-8.0	220	10.4	9.0-11.9	186	5.3	4.6-6.2
1987	691	12.3	11.4-13.3	392	18.0	16.3-20.0	299	8.5	7.5-9.5	427	7.7	6.9-8.4	248	11.5	10.1-13.1	179	5.0	4.3-5.9
1988	645	11.8	10.9-12.7	358	16.8	15.1-18.7	287	8.3	7.4-9.4	418	7.6	6.9-8.3	214	10.0	8.7-11.4	204	5.9	5.1-6.8
1989	616	11.6	10.6-12.5	338	16.1	14.4-17.9	278	8.3	7.4-9.4	433	8.0	7.2-8.8	247	11.8	10.3-13.4	186	5.6	4.8-6.5
1990	707	13.4	12.5-14.5	397	19.4	17.5-21.4	310	9.6	8.5-10.7	382	7.3	6.5-8.0	223	11.0	9.6-12.6	159	4.8	4.1-5.7
1991	696	13.4	12.4-14.4	404	20.2	18.2-22.3	292	8.9	7.9-10.0	409	7.8	7.0-8.6	230	11.5	10.1-13.1	179	5.4	4.6-6.3
1992	642	12.5	11.5-13.5	357	17.9	16.0-19.8	285	9.1	8.0-10.2	409	7.9	7.1-8.7	226	11.5	10.0-13.1	183	5.6	4.8-6.6
1993	672	13.1	12.1-14.1	385	19.4	17.5-21.5	287	9.1	8.0-10.2	400	7.8	7.0-8.6	225	11.5	10.1-13.2	175	5.5	4.7-6.4
1994	699	13.8	12.7-14.8	374	18.6	16.7-20.6	325	10.5	9.4-11.8	430	8.4	7.6-9.2	247	12.6	11.0-14.2	183	5.8	5.0-6.8
1995	624	12.2	11.3-13.2	348	17.5	15.7-19.4	276	8.7	7.7-9.9	422	8.3	7.5-9.1	229	11.8	10.3-13.4	193	6.1	5.3-7.1
1996	649	12.8	11.9-13.9	348	17.6	15.8-19.6	301	9.6	8.6-10.8	388	7.6	6.9-8.5	195	10.0	8.7-11.5	193	6.1	5.3-7.1
1997	583	11.6	10.6-12.6	333	16.9	15.1-18.8	250	8.1	7.1-9.2	367	7.3	6.6-8.1	198	10.1	8.7-11.6	169	5.6	4.8-6.5
1998	646	12.9	11.9-13.9	363	18.2	16.4-20.2	283	9.4	8.3-10.6	379	7.5	6.7-8.2	208	10.5	9.1-12.1	171	5.4	4.6-6.4
1999	602	11.9	10.9-12.8	354	17.6	15.8-19.5	248	8.0	7.0-9.1	344	6.7	6.0-7.4	187	9.4	8.1-10.9	157	5.0	4.3-5.9
2000	565	11.0	10.2-12.0	297	14.5	12.9-16.3	268	8.7	7.6-9.8	313	6.0	5.4-6.8	156	7.7	6.5-9.0	157	5.0	4.2-5.9
2001	589	11.6	10.7-12.6	332	16.1	14.4-17.9	257	8.5	7.5-9.6	316	6.2	5.5-6.9	163	8.0	6.8-9.3	153	5.0	4.2-5.9
2002	527	10.3	9.5-11.3	298	14.3	12.7-16.0	229	7.6	6.6-8.6	285	5.5	4.9-6.2	158	7.6	6.5-8.9	127	4.0	3.3-4.8
2003	570	11.1	10.2-12.1	310	14.8	13.2-16.6	260	8.6	7.6-9.8	299	5.8	5.2-6.5	158	7.7	6.5-9.0	141	4.6	3.9-5.5
2004	561	11.1	10.2-12.1	307	14.8	13.2-16.5	254	8.6	7.6-9.8	307	6.0	5.3-6.7	151	7.3	6.2-8.6	156	5.3	4.5-6.2
2005	548	10.9	10.0-11.8	306	14.8	13.2-16.6	242	8.1	7.1-9.2	296	5.7	5.1-6.4	164	8.0	6.8-9.3	132	4.2	3.5-5.0
2006	513	10.3	9.4-11.2	293	14.2	12.6-15.9	220	7.6	6.6-8.7	260	5.2	4.5-5.8	146	7.1	6.0-8.3	114	3.9	3.2-4.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Stomach

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	549	11.0	10.1-12.0	302	14.6	13.0-16.4	247	8.4	7.4-9.6	239	4.7	4.1-5.3	129	6.3	5.2-7.4	110	3.6	2.9-4.4
2008	503	10.0	9.1-10.9	303	14.4	12.8-16.2	200	6.8	5.9-7.9	248	4.9	4.3-5.6	156	7.5	6.4-8.8	92	3.1	2.5-3.8
2009	488	9.8	8.9-10.7	271	12.9	11.4-14.6	217	7.5	6.5-8.6	244	4.8	4.2-5.5	137	6.7	5.6-7.9	107	3.5	2.8-4.3
2010	490	9.8	8.9-10.7	279	13.1	11.6-14.8	211	7.3	6.3-8.4	241	4.6	4.0-5.2	109	5.2	4.2-6.3	132	4.4	3.6-5.2
2011	431	8.5	7.7-9.4	243	11.4	10.0-12.9	188	6.4	5.5-7.5	235	4.5	4.0-5.2	117	5.5	4.5-6.6	118	3.9	3.2-4.7
2012	484	9.4	8.6-10.3	284	13.0	11.6-14.7	200	6.8	5.9-7.9	223	4.2	3.7-4.8	125	5.7	4.8-6.8	98	3.1	2.5-3.8
2013	502	9.5	8.7-10.4	282	12.6	11.1-14.1	220	7.2	6.2-8.2	227	4.2	3.6-4.8	139	6.2	5.2-7.4	88	2.7	2.2-3.4
2014	473	9.0	8.2-9.9	288	12.7	11.2-14.2	185	6.2	5.3-7.2	250	4.6	4.1-5.3	139	6.1	5.1-7.2	111	3.5	2.8-4.3
2015	464	8.7	7.9-9.5	249	10.7	9.4-12.2	215	7.1	6.1-8.1	213	3.9	3.4-4.5	114	5.0	4.1-6.0	99	3.0	2.5-3.8
2016	475	8.9	8.1-9.7	278	11.8	10.4-13.3	197	6.8	5.8-7.8	250	4.6	4.0-5.2	162	6.9	5.8-8.0	88	2.8	2.3-3.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Colorectal

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	4172	67.6	65.5-69.7	2018	81.0	77.3-84.8	2154	59.1	56.5-61.7	2246	37.1	35.5-38.7	1139	47.8	44.9-50.8	1107	30.5	28.7-32.4
1977	4103	67.4	65.3-69.6	2032	84.1	80.3-88.0	2071	56.9	54.4-59.5	2104	35.1	33.6-36.7	1037	44.4	41.6-47.4	1067	29.5	27.7-31.4
1978	3961	65.4	63.3-67.5	1944	81.5	77.8-85.4	2017	55.5	53.0-58.0	1967	33.2	31.7-34.7	1018	44.1	41.3-47.0	949	26.5	24.8-28.3
1979	4049	67.7	65.6-69.9	1982	84.2	80.4-88.2	2067	57.5	55.0-60.1	1979	33.5	32.0-35.0	942	41.1	38.4-43.9	1037	28.9	27.1-30.7
1980	4197	70.9	68.7-73.1	2093	90.0	86.1-94.1	2104	59.1	56.5-61.8	2005	34.4	32.9-36.0	1019	46.2	43.3-49.2	986	27.4	25.7-29.3
1981	4090	70.0	67.8-72.2	1989	87.7	83.8-91.8	2101	59.1	56.5-61.8	1831	31.5	30.0-33.0	862	39.1	36.4-41.9	969	27.0	25.3-28.8
1982	4106	70.7	68.5-72.9	1975	87.9	83.9-92.0	2131	60.3	57.7-63.1	1755	30.4	29.0-31.9	873	39.6	36.9-42.4	882	25.1	23.4-26.9
1983	3996	68.7	66.5-70.9	1948	86.1	82.2-90.1	2048	58.0	55.5-60.6	1767	30.9	29.4-32.4	847	38.9	36.2-41.7	920	26.0	24.3-27.8
1984	3888	67.9	65.7-70.1	1854	83.1	79.3-87.1	2034	58.6	56.0-61.2	1784	31.0	29.5-32.5	900	41.2	38.5-44.1	884	25.0	23.4-26.8
1985	3887	67.5	65.3-69.7	1949	87.2	83.3-91.3	1938	55.4	52.9-58.0	1658	29.0	27.6-30.4	816	36.9	34.4-39.6	842	23.9	22.3-25.7
1986	3949	69.6	67.4-71.9	1970	89.6	85.5-93.7	1979	57.4	54.9-60.1	1611	28.5	27.1-30.0	794	37.0	34.4-39.7	817	23.6	22.0-25.4
1987	3531	62.6	60.5-64.7	1750	80.0	76.2-83.9	1781	51.6	49.2-54.1	1585	28.0	26.6-29.4	766	35.9	33.4-38.6	819	23.3	21.7-25.0
1988	3429	62.0	59.9-64.1	1633	76.2	72.4-80.0	1796	52.9	50.5-55.5	1510	27.4	26.0-28.8	772	37.1	34.5-39.9	738	21.7	20.1-23.4
1989	3206	59.2	57.1-61.3	1563	74.0	70.3-77.8	1643	49.7	47.2-52.2	1411	25.8	24.4-27.2	653	31.9	29.5-34.5	758	22.2	20.7-23.9
1990	3227	59.9	57.8-62.0	1504	71.2	67.6-74.9	1723	52.2	49.7-54.8	1420	26.3	25.0-27.8	677	33.2	30.7-35.8	743	22.1	20.5-23.8
1991	3208	60.5	58.4-62.6	1607	78.7	74.9-82.8	1601	49.2	46.8-51.8	1309	24.6	23.3-26.0	647	32.6	30.1-35.2	662	20.0	18.5-21.6
1992	3080	58.9	56.8-61.1	1448	71.2	67.6-75.1	1632	51.3	48.8-53.9	1239	23.3	22.0-24.6	593	29.7	27.3-32.2	646	19.3	17.8-20.9
1993	3064	59.0	56.9-61.1	1477	72.3	68.7-76.2	1587	50.2	47.7-52.8	1213	23.1	21.8-24.4	598	30.0	27.6-32.6	615	18.8	17.3-20.4
1994	3376	65.5	63.3-67.8	1570	77.6	73.7-81.5	1806	57.4	54.7-60.2	1353	25.9	24.5-27.3	665	33.8	31.3-36.5	688	21.2	19.6-22.9
1995	3127	61.4	59.2-63.6	1463	73.2	69.5-77.1	1664	53.5	50.9-56.2	1353	26.1	24.8-27.6	645	33.0	30.5-35.6	708	22.1	20.4-23.8
1996	3022	60.0	57.8-62.2	1457	72.9	69.2-76.8	1565	51.2	48.6-53.9	1145	22.2	20.9-23.5	546	27.6	25.3-30.0	599	18.6	17.1-20.2
1997	3148	62.5	60.3-64.7	1459	73.0	69.3-76.9	1689	55.0	52.3-57.7	1242	24.3	23.0-25.7	610	31.0	28.6-33.6	632	19.9	18.3-21.5
1998	3212	63.8	61.6-66.1	1537	76.5	72.7-80.4	1675	55.4	52.7-58.1	1172	22.9	21.6-24.2	537	27.1	24.8-29.4	635	19.9	18.4-21.6
1999	3063	60.5	58.4-62.7	1475	72.4	68.7-76.2	1588	52.4	49.8-55.1	1092	21.2	19.9-22.5	518	25.8	23.6-28.1	574	18.2	16.7-19.8
2000	3110	61.0	58.9-63.2	1548	75.4	71.7-79.2	1562	50.6	48.1-53.2	1136	21.9	20.6-23.2	556	27.5	25.2-29.9	580	18.1	16.6-19.7
2001	2980	58.3	56.2-60.4	1469	71.5	67.9-75.3	1511	49.6	47.1-52.2	1042	20.0	18.8-21.2	513	25.3	23.1-27.5	529	16.4	15.0-17.9
2002	2904	56.8	54.7-58.9	1417	67.9	64.4-71.5	1487	48.4	46.0-51.0	1044	20.1	18.8-21.3	514	25.1	23.0-27.4	530	16.7	15.3-18.2
2003	2828	55.4	53.3-57.5	1410	67.7	64.2-71.4	1418	46.6	44.1-49.1	1065	20.4	19.2-21.7	495	24.1	22.0-26.3	570	17.8	16.3-19.3
2004	2695	52.7	50.7-54.8	1327	63.6	60.2-67.1	1368	44.8	42.5-47.3	927	17.8	16.7-19.0	448	21.8	19.8-23.9	479	15.0	13.6-16.4
2005	2600	51.2	49.2-53.2	1273	61.1	57.8-64.6	1327	43.9	41.5-46.4	869	16.7	15.6-17.8	406	19.9	18.0-22.0	463	14.4	13.1-15.8
2006	2527	50.4	48.5-52.5	1256	60.9	57.5-64.3	1271	42.9	40.5-45.4	909	17.8	16.6-19.0	457	22.5	20.5-24.7	452	14.7	13.4-16.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Colorectal

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	2478	49.4	47.4-51.4	1235	58.9	55.6-62.3	1243	42.3	40.0-44.8	818	15.9	14.8-17.0	388	18.9	17.1-20.9	430	13.7	12.4-15.1
2008	2366	47.3	45.3-49.2	1191	56.6	53.4-59.9	1175	40.2	37.9-42.6	863	16.8	15.7-18.0	432	21.0	19.1-23.1	431	13.9	12.6-15.3
2009	2346	46.8	44.9-48.8	1134	53.5	50.4-56.8	1212	41.8	39.5-44.3	825	15.9	14.8-17.1	383	18.5	16.7-20.5	442	14.0	12.7-15.4
2010	2212	43.8	42.0-45.7	1116	52.2	49.1-55.4	1096	37.4	35.1-39.7	852	16.4	15.3-17.5	446	21.2	19.3-23.3	406	13.0	11.7-14.4
2011	2071	40.7	38.9-42.5	1044	48.3	45.4-51.4	1027	35.2	33.0-37.5	809	15.3	14.3-16.4	410	19.0	17.2-21.0	399	12.5	11.3-13.8
2012	2006	38.8	37.1-40.6	1035	46.7	43.9-49.7	971	32.7	30.6-34.9	738	13.8	12.8-14.9	341	15.7	14.0-17.4	397	12.5	11.3-13.9
2013	2161	41.4	39.7-43.2	1097	49.0	46.1-52.0	1064	35.5	33.3-37.8	734	13.5	12.5-14.5	351	15.7	14.1-17.5	383	11.7	10.5-13.0
2014	1995	37.9	36.2-39.6	1037	45.1	42.3-48.0	958	32.2	30.1-34.4	711	12.9	12.0-13.9	347	15.3	13.7-17.0	364	11.1	10.0-12.4
2015	1966	37.0	35.3-38.7	975	42.1	39.5-44.9	991	33.1	31.0-35.3	705	12.6	11.6-13.6	360	15.7	14.1-17.4	345	10.3	9.2-11.5
2016	1903	35.7	34.1-37.4	1023	43.7	41.0-46.5	880	29.7	27.7-31.8	785	13.8	12.8-14.8	405	17.2	15.5-18.9	380	11.3	10.2-12.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Colon excluding rectum

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	2933	47.5	45.7-49.3	1367	55.2	52.2-58.4	1566	42.8	40.6-45.0	1782	29.5	28.1-31.0	881	37.4	34.8-40.1	901	24.8	23.2-26.6
1977	2910	48.0	46.2-49.8	1398	58.5	55.4-61.8	1512	41.6	39.5-43.8	1735	28.9	27.5-30.3	841	36.0	33.5-38.6	894	24.7	23.1-26.5
1978	2838	47.0	45.3-48.9	1365	57.6	54.4-60.8	1473	40.7	38.6-42.9	1631	27.5	26.1-28.9	832	36.2	33.6-38.8	799	22.2	20.6-23.8
1979	2921	49.0	47.2-50.8	1383	59.4	56.2-62.7	1538	42.7	40.6-45.0	1656	28.0	26.6-29.4	760	33.1	30.7-35.6	896	24.9	23.3-26.6
1980	2969	50.3	48.5-52.2	1439	63.1	59.7-66.5	1530	42.8	40.6-45.0	1714	29.5	28.1-31.0	857	39.1	36.4-41.9	857	23.9	22.3-25.6
1981	2891	49.3	47.5-51.2	1355	60.2	56.9-63.6	1536	42.9	40.7-45.1	1580	27.2	25.8-28.6	731	33.1	30.7-35.7	849	23.7	22.1-25.4
1982	2909	50.1	48.3-52.0	1333	60.2	56.9-63.7	1576	44.4	42.2-46.7	1527	26.4	25.1-27.8	761	34.6	32.1-37.2	766	21.8	20.3-23.5
1983	2872	49.4	47.5-51.2	1342	59.7	56.4-63.1	1530	43.1	40.9-45.4	1490	25.9	24.6-27.3	708	32.3	29.9-34.8	782	22.0	20.5-23.7
1984	2811	48.9	47.1-50.8	1312	59.5	56.2-62.9	1499	42.8	40.6-45.1	1534	26.6	25.2-27.9	766	35.1	32.6-37.7	768	21.8	20.2-23.4
1985	2900	50.4	48.5-52.3	1386	62.7	59.3-66.2	1514	43.3	41.1-45.6	1463	25.6	24.2-26.9	712	32.2	29.8-34.7	751	21.4	19.9-23.0
1986	2921	51.5	49.6-53.4	1386	63.4	60.0-66.9	1535	44.4	42.2-46.8	1397	24.7	23.4-26.1	678	31.4	29.1-34.0	719	20.8	19.3-22.5
1987	2583	45.4	43.7-47.3	1229	56.4	53.3-59.8	1354	38.7	36.6-40.9	1393	24.6	23.3-26.0	688	32.3	29.8-34.8	705	20.1	18.6-21.6
1988	2500	44.9	43.2-46.7	1141	53.5	50.3-56.7	1359	39.6	37.5-41.9	1319	23.9	22.6-25.3	678	32.6	30.1-35.2	641	18.8	17.4-20.4
1989	2335	43.0	41.2-44.8	1100	52.5	49.4-55.7	1235	37.1	35.0-39.3	1262	23.0	21.8-24.4	580	28.5	26.1-30.9	682	20.0	18.5-21.6
1990	2408	44.6	42.8-46.4	1085	51.7	48.6-54.9	1323	40.0	37.8-42.3	1257	23.3	22.0-24.6	590	28.9	26.6-31.4	667	19.8	18.2-21.4
1991	2395	44.9	43.1-46.8	1158	56.9	53.6-60.3	1237	37.8	35.7-40.0	1133	21.2	20.0-22.5	557	28.1	25.8-30.6	576	17.3	15.9-18.8
1992	2261	43.0	41.2-44.9	1045	51.7	48.6-55.0	1216	38.0	35.8-40.2	1077	20.2	19.0-21.4	509	25.3	23.1-27.7	568	16.9	15.5-18.4
1993	2282	43.7	41.9-45.6	1061	52.0	48.9-55.3	1221	38.3	36.2-40.6	1063	20.2	19.0-21.5	514	25.9	23.7-28.3	549	16.8	15.3-18.3
1994	2512	48.5	46.6-50.4	1149	57.0	53.7-60.5	1363	42.7	40.5-45.1	1206	23.0	21.8-24.4	592	30.1	27.7-32.6	614	19.0	17.5-20.6
1995	2328	45.4	43.6-47.3	1060	53.4	50.2-56.8	1268	40.1	37.9-42.5	1182	22.8	21.5-24.1	563	28.9	26.5-31.4	619	19.2	17.7-20.8
1996	2227	43.9	42.0-45.7	1032	51.8	48.7-55.1	1195	38.4	36.2-40.7	1007	19.5	18.3-20.7	486	24.7	22.5-27.0	521	16.0	14.6-17.5
1997	2367	46.7	44.8-48.6	1072	54.0	50.8-57.3	1295	41.7	39.4-44.1	1092	21.3	20.1-22.6	529	26.9	24.6-29.3	563	17.7	16.2-19.2
1998	2343	46.3	44.4-48.2	1084	54.3	51.1-57.7	1259	41.1	38.8-43.5	1048	20.4	19.2-21.7	478	24.2	22.1-26.5	570	17.8	16.4-19.4
1999	2249	44.2	42.3-46.0	1062	52.5	49.3-55.7	1187	38.8	36.6-41.1	949	18.4	17.2-19.6	451	22.5	20.5-24.7	498	15.8	14.4-17.3
2000	2285	44.6	42.8-46.4	1083	53.0	49.9-56.3	1202	38.7	36.5-41.0	1000	19.2	18.1-20.5	490	24.3	22.2-26.5	510	15.9	14.5-17.4
2001	2203	42.8	41.0-44.6	1063	51.9	48.9-55.2	1140	36.6	34.5-38.9	891	17.0	15.9-18.2	428	21.1	19.1-23.2	463	14.2	13.0-15.6
2002	2139	41.5	39.8-43.4	999	48.2	45.3-51.3	1140	36.8	34.6-39.0	903	17.3	16.2-18.5	447	21.9	19.9-24.0	456	14.4	13.0-15.8
2003	2063	40.1	38.4-41.9	996	48.2	45.2-51.2	1067	34.6	32.5-36.8	931	17.8	16.6-19.0	430	21.0	19.1-23.1	501	15.5	14.1-16.9
2004	1960	38.2	36.5-39.9	950	45.7	42.9-48.8	1010	32.6	30.6-34.7	813	15.6	14.5-16.7	382	18.6	16.7-20.5	431	13.4	12.2-14.8
2005	1931	37.7	36.1-39.5	903	43.4	40.6-46.4	1028	33.6	31.6-35.8	753	14.4	13.4-15.5	348	17.1	15.4-19.0	405	12.5	11.3-13.9
2006	1803	35.8	34.2-37.5	858	41.9	39.1-44.8	945	31.6	29.5-33.7	799	15.6	14.5-16.7	394	19.4	17.6-21.5	405	13.1	11.8-14.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Colon excluding rectum

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	1764	34.9	33.3-36.6	856	41.0	38.2-43.8	908	30.6	28.6-32.7	709	13.7	12.7-14.8	338	16.5	14.8-18.3	371	11.7	10.5-13.0
2008	1700	33.7	32.1-35.4	834	39.8	37.2-42.7	866	29.2	27.2-31.3	722	14.1	13.0-15.1	360	17.6	15.8-19.5	362	11.7	10.5-13.0
2009	1695	33.6	32.0-35.2	770	36.8	34.2-39.5	925	31.3	29.3-33.5	702	13.5	12.5-14.6	326	15.8	14.1-17.6	376	11.8	10.6-13.1
2010	1573	30.9	29.4-32.5	767	35.9	33.4-38.6	806	26.9	25.0-28.9	736	14.1	13.0-15.1	374	17.9	16.1-19.8	362	11.4	10.3-12.7
2011	1488	29.1	27.6-30.7	746	34.7	32.2-37.3	742	25.1	23.3-27.1	680	12.9	11.9-13.9	344	15.9	14.3-17.7	336	10.5	9.4-11.8
2012	1451	27.8	26.4-29.3	713	32.4	30.0-34.9	738	24.4	22.6-26.3	618	11.5	10.6-12.5	282	13.0	11.5-14.7	336	10.4	9.3-11.6
2013	1546	29.5	28.0-31.0	756	34.0	31.6-36.5	790	26.1	24.2-28.0	621	11.3	10.4-12.3	292	13.2	11.7-14.8	329	9.9	8.9-11.1
2014	1426	26.8	25.4-28.3	725	31.7	29.4-34.2	701	23.0	21.2-24.8	590	10.6	9.8-11.5	281	12.4	11.0-14.0	309	9.3	8.3-10.5
2015	1362	25.5	24.1-26.9	652	28.4	26.2-30.7	710	23.4	21.7-25.3	575	10.2	9.4-11.1	291	12.6	11.2-14.2	284	8.5	7.5-9.6
2016	1359	25.3	23.9-26.7	720	30.9	28.6-33.3	639	21.2	19.5-23.0	643	11.3	10.4-12.2	328	13.9	12.4-15.5	315	9.4	8.3-10.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Rectum & rectosigmoid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1239	20.1	19.0-21.3	651	25.8	23.7-27.9	588	16.3	15.0-17.8	464	7.5	6.8-8.3	258	10.4	9.1-11.9	206	5.7	4.9-6.5
1977	1193	19.4	18.3-20.6	634	25.6	23.5-27.7	559	15.4	14.1-16.7	369	6.2	5.6-6.9	196	8.5	7.2-9.8	173	4.8	4.1-5.6
1978	1123	18.3	17.3-19.5	579	24.0	22.0-26.1	544	14.8	13.5-16.1	336	5.7	5.1-6.4	186	7.9	6.8-9.2	150	4.3	3.6-5.1
1979	1128	18.8	17.7-19.9	599	24.8	22.8-27.0	529	14.8	13.5-16.2	323	5.5	4.9-6.2	182	8.0	6.8-9.3	141	4.0	3.3-4.7
1980	1228	20.6	19.5-21.8	654	27.0	24.9-29.2	574	16.3	15.0-17.8	291	4.9	4.4-5.5	162	7.1	6.0-8.3	129	3.5	3.0-4.2
1981	1199	20.7	19.5-21.9	634	27.5	25.4-29.8	565	16.2	14.9-17.7	251	4.3	3.8-4.9	131	6.0	5.0-7.1	120	3.3	2.8-4.0
1982	1197	20.6	19.4-21.8	642	27.6	25.5-29.9	555	15.9	14.6-17.4	228	4.0	3.5-4.5	112	5.0	4.1-6.1	116	3.3	2.7-4.0
1983	1124	19.3	18.2-20.5	606	26.4	24.3-28.6	518	14.9	13.6-16.3	277	4.9	4.4-5.6	139	6.6	5.5-7.8	138	4.0	3.3-4.7
1984	1077	18.9	17.8-20.1	542	23.7	21.7-25.8	535	15.7	14.4-17.2	250	4.4	3.9-5.0	134	6.1	5.1-7.3	116	3.3	2.7-4.0
1985	987	17.1	16.0-18.2	563	24.5	22.5-26.7	424	12.1	11.0-13.4	195	3.4	3.0-4.0	104	4.7	3.8-5.8	91	2.6	2.0-3.2
1986	1028	18.1	17.0-19.3	584	26.2	24.0-28.4	444	13.0	11.8-14.3	214	3.8	3.3-4.3	116	5.6	4.6-6.7	98	2.8	2.3-3.4
1987	948	17.2	16.1-18.3	521	23.6	21.5-25.7	427	12.9	11.6-14.2	192	3.4	2.9-3.9	78	3.7	2.9-4.6	114	3.2	2.6-3.9
1988	929	17.0	15.9-18.2	492	22.7	20.7-24.8	437	13.3	12.0-14.7	191	3.5	3.0-4.0	94	4.5	3.6-5.6	97	2.9	2.3-3.5
1989	871	16.2	15.1-17.4	463	21.5	19.6-23.6	408	12.5	11.3-13.9	149	2.7	2.3-3.2	73	3.5	2.7-4.4	76	2.2	1.7-2.8
1990	819	15.3	14.2-16.4	419	19.5	17.6-21.4	400	12.2	11.0-13.5	163	3.0	2.6-3.6	87	4.2	3.4-5.2	76	2.3	1.8-3.0
1991	813	15.5	14.5-16.7	449	21.9	19.9-24.0	364	11.5	10.3-12.7	176	3.4	2.9-3.9	90	4.5	3.6-5.5	86	2.7	2.1-3.4
1992	819	15.9	14.8-17.1	403	19.5	17.7-21.6	416	13.3	12.0-14.7	162	3.1	2.7-3.7	84	4.3	3.4-5.4	78	2.4	1.9-3.0
1993	782	15.3	14.2-16.4	416	20.3	18.4-22.4	366	11.8	10.6-13.1	150	2.9	2.4-3.4	84	4.1	3.3-5.1	66	2.0	1.6-2.6
1994	864	17.0	15.9-18.2	421	20.5	18.6-22.6	443	14.7	13.3-16.1	147	2.8	2.4-3.4	73	3.8	2.9-4.7	74	2.2	1.8-2.8
1995	799	15.9	14.8-17.1	403	19.8	17.9-21.8	396	13.3	12.0-14.8	171	3.4	2.9-3.9	82	4.1	3.3-5.1	89	2.9	2.3-3.6
1996	795	16.1	15.0-17.3	425	21.1	19.2-23.3	370	12.8	11.5-14.2	138	2.7	2.3-3.2	60	2.9	2.2-3.8	78	2.6	2.0-3.3
1997	781	15.8	14.7-16.9	387	19.0	17.2-21.0	394	13.2	11.9-14.7	150	3.0	2.5-3.5	81	4.1	3.3-5.1	69	2.2	1.7-2.8
1998	869	17.5	16.4-18.7	453	22.2	20.2-24.3	416	14.3	12.9-15.8	124	2.5	2.0-2.9	59	2.8	2.2-3.7	65	2.1	1.6-2.7
1999	814	16.3	15.2-17.5	413	19.9	18.0-22.0	401	13.6	12.3-15.1	143	2.8	2.3-3.3	67	3.3	2.5-4.2	76	2.4	1.9-3.0
2000	825	16.4	15.3-17.6	465	22.4	20.4-24.5	360	11.9	10.7-13.3	136	2.7	2.2-3.1	66	3.2	2.5-4.1	70	2.2	1.7-2.8
2001	777	15.5	14.4-16.6	406	19.6	17.7-21.6	371	13.0	11.7-14.4	151	3.0	2.5-3.5	85	4.1	3.3-5.1	66	2.2	1.7-2.8
2002	765	15.3	14.2-16.4	418	19.6	17.8-21.6	347	11.7	10.4-13.0	141	2.7	2.3-3.2	67	3.2	2.5-4.1	74	2.3	1.8-3.0
2003	765	15.2	14.2-16.4	414	19.6	17.7-21.6	351	12.0	10.7-13.3	134	2.6	2.2-3.1	65	3.1	2.4-4.0	69	2.3	1.8-2.9
2004	735	14.6	13.5-15.7	377	17.9	16.1-19.8	358	12.2	11.0-13.6	114	2.2	1.8-2.7	66	3.2	2.5-4.1	48	1.5	1.1-2.1
2005	669	13.5	12.4-14.5	370	17.7	15.9-19.6	299	10.3	9.1-11.6	116	2.2	1.9-2.7	58	2.8	2.1-3.6	58	1.9	1.4-2.4
2006	724	14.6	13.6-15.7	398	19.0	17.1-20.9	326	11.4	10.1-12.7	110	2.2	1.8-2.6	63	3.1	2.4-4.0	47	1.6	1.2-2.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Rectum & rectosigmoid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	714	14.5	13.4-15.6	379	17.9	16.1-19.8	335	11.8	10.5-13.2	109	2.2	1.8-2.6	50	2.4	1.8-3.2	59	1.9	1.4-2.5
2008	666	13.5	12.5-14.6	357	16.7	15.0-18.6	309	11.0	9.8-12.3	141	2.7	2.3-3.2	72	3.4	2.7-4.3	69	2.2	1.7-2.8
2009	651	13.3	12.3-14.3	364	16.8	15.1-18.6	287	10.5	9.3-11.8	123	2.4	2.0-2.9	57	2.7	2.0-3.5	66	2.2	1.7-2.9
2010	639	12.9	11.9-14.0	349	16.3	14.6-18.1	290	10.4	9.2-11.8	116	2.3	1.9-2.8	72	3.4	2.7-4.3	44	1.5	1.1-2.1
2011	583	11.6	10.6-12.6	298	13.6	12.1-15.3	285	10.1	8.9-11.3	129	2.5	2.0-2.9	66	3.1	2.4-3.9	63	2.0	1.5-2.6
2012	555	11.0	10.1-11.9	322	14.3	12.8-16.0	233	8.3	7.3-9.5	120	2.3	1.9-2.8	59	2.6	2.0-3.4	61	2.1	1.6-2.7
2013	615	11.9	11.0-12.9	341	15.0	13.4-16.7	274	9.4	8.3-10.6	113	2.2	1.8-2.6	59	2.5	1.9-3.3	54	1.8	1.3-2.4
2014	569	11.1	10.2-12.0	312	13.4	11.9-15.0	257	9.2	8.1-10.4	121	2.3	1.9-2.7	66	2.9	2.2-3.7	55	1.8	1.3-2.3
2015	604	11.5	10.6-12.5	323	13.7	12.2-15.3	281	9.7	8.5-10.9	130	2.4	2.0-2.9	69	3.0	2.4-3.9	61	1.8	1.4-2.4
2016	544	10.4	9.6-11.4	303	12.8	11.4-14.4	241	8.6	7.5-9.7	142	2.5	2.1-3.0	77	3.2	2.6-4.1	65	1.9	1.5-2.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Liver / intrahepatic bile duct

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	223	3.7	3.2-4.2	122	5.0	4.1-6.0	101	2.8	2.2-3.4	232	3.8	3.3-4.3	122	4.8	3.9-5.7	110	3.1	2.5-3.7
1977	216	3.6	3.1-4.1	131	5.3	4.4-6.3	85	2.3	1.9-2.9	256	4.2	3.7-4.7	136	5.4	4.5-6.5	120	3.3	2.7-4.0
1978	190	3.2	2.7-3.7	109	4.4	3.6-5.4	81	2.3	1.8-2.8	231	3.8	3.3-4.3	136	5.5	4.6-6.6	95	2.6	2.1-3.2
1979	191	3.3	2.8-3.8	116	4.9	4.0-5.9	75	2.2	1.7-2.8	251	4.3	3.8-4.9	133	5.7	4.7-6.8	118	3.4	2.8-4.1
1980	203	3.6	3.1-4.1	112	4.9	4.0-5.9	91	2.8	2.2-3.4	215	3.7	3.2-4.2	108	4.7	3.8-5.7	107	3.0	2.4-3.6
1981	206	3.6	3.1-4.2	110	4.9	4.0-5.9	96	2.7	2.2-3.3	261	4.5	4.0-5.1	142	6.2	5.2-7.4	119	3.4	2.8-4.1
1982	175	3.1	2.6-3.6	123	5.5	4.5-6.6	52	1.5	1.1-2.0	241	4.1	3.6-4.7	140	6.1	5.1-7.2	101	2.9	2.3-3.6
1983	196	3.5	3.0-4.0	114	5.0	4.1-6.0	82	2.5	2.0-3.1	251	4.5	3.9-5.1	132	5.8	4.8-6.9	119	3.6	2.9-4.3
1984	196	3.4	3.0-4.0	117	5.1	4.2-6.1	79	2.3	1.8-3.0	244	4.3	3.7-4.9	130	5.8	4.8-6.9	114	3.2	2.6-3.9
1985	211	3.9	3.4-4.5	126	5.8	4.8-7.0	85	2.5	2.0-3.1	263	4.7	4.1-5.3	153	7.0	6.0-8.3	110	3.1	2.6-3.8
1986	232	4.1	3.6-4.7	143	6.5	5.4-7.6	89	2.6	2.1-3.3	266	4.8	4.2-5.4	137	6.2	5.2-7.4	129	3.8	3.2-4.6
1987	193	3.5	3.1-4.1	118	5.5	4.5-6.6	75	2.2	1.7-2.8	258	4.7	4.1-5.3	144	6.8	5.7-8.0	114	3.4	2.8-4.1
1988	227	4.3	3.8-4.9	139	6.5	5.5-7.7	88	2.7	2.2-3.4	274	5.1	4.5-5.7	150	7.0	5.9-8.2	124	3.7	3.1-4.4
1989	191	3.6	3.1-4.1	118	5.5	4.5-6.6	73	2.2	1.7-2.8	227	4.2	3.6-4.7	130	6.0	5.0-7.1	97	2.9	2.3-3.6
1990	241	4.7	4.1-5.4	147	7.2	6.1-8.5	94	3.0	2.4-3.7	272	5.1	4.5-5.8	152	7.5	6.3-8.8	120	3.7	3.0-4.4
1991	257	5.0	4.4-5.7	160	7.6	6.4-8.9	97	3.1	2.5-3.9	275	5.2	4.6-5.8	159	7.6	6.4-8.9	116	3.5	2.9-4.2
1992	264	5.3	4.6-5.9	158	7.6	6.5-8.9	106	3.5	2.9-4.3	296	5.8	5.1-6.5	175	8.5	7.3-9.9	121	3.8	3.1-4.6
1993	242	4.8	4.2-5.5	144	7.0	5.9-8.2	98	3.2	2.6-3.9	265	5.1	4.5-5.8	139	6.7	5.6-7.9	126	4.0	3.3-4.7
1994	273	5.5	4.9-6.2	172	8.5	7.3-9.9	101	3.2	2.6-4.0	241	4.8	4.2-5.5	152	7.6	6.4-8.9	89	2.9	2.3-3.6
1995	292	6.0	5.3-6.7	184	9.0	7.8-10.5	108	3.6	2.9-4.4	269	5.4	4.7-6.1	155	7.7	6.5-9.0	114	3.6	2.9-4.3
1996	324	6.7	6.0-7.5	223	11.1	9.7-12.6	101	3.4	2.8-4.2	265	5.3	4.7-6.0	167	8.3	7.1-9.6	98	3.1	2.5-3.9
1997	342	7.0	6.3-7.8	220	10.9	9.5-12.4	122	4.0	3.3-4.8	247	4.9	4.3-5.6	145	7.2	6.1-8.5	102	3.2	2.6-3.9
1998	400	8.2	7.4-9.1	254	12.4	10.9-14.0	146	5.0	4.2-5.9	291	5.9	5.2-6.6	183	9.0	7.7-10.4	108	3.5	2.9-4.3
1999	374	7.7	6.9-8.5	248	11.9	10.5-13.5	126	4.1	3.4-4.9	276	5.5	4.9-6.2	170	8.3	7.1-9.6	106	3.5	2.8-4.2
2000	371	7.5	6.8-8.4	259	12.4	10.9-14.0	112	3.8	3.1-4.6	278	5.6	5.0-6.3	176	8.4	7.2-9.8	102	3.4	2.7-4.1
2001	402	8.2	7.4-9.0	270	12.8	11.3-14.4	132	4.5	3.8-5.4	296	5.9	5.3-6.6	186	8.9	7.6-10.2	110	3.6	2.9-4.3
2002	376	7.7	6.9-8.5	261	12.2	10.8-13.8	115	4.0	3.3-4.8	303	6.1	5.4-6.8	193	9.1	7.9-10.5	110	3.6	2.9-4.3
2003	443	9.0	8.2-9.9	296	13.9	12.4-15.6	147	5.0	4.3-6.0	302	5.9	5.3-6.7	186	8.9	7.6-10.2	116	3.7	3.1-4.5
2004	433	8.8	8.0-9.6	307	14.3	12.7-16.0	126	4.4	3.7-5.3	313	6.2	5.5-6.9	193	9.0	7.8-10.4	120	3.9	3.2-4.7
2005	465	9.5	8.6-10.4	335	15.6	14.0-17.4	130	4.5	3.7-5.3	317	6.3	5.6-7.1	204	9.5	8.2-10.9	113	3.7	3.0-4.5
2006	499	10.2	9.3-11.1	333	15.4	13.8-17.2	166	5.8	5.0-6.8	337	6.8	6.1-7.6	213	10.1	8.8-11.5	124	4.2	3.5-5.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Liver / intrahepatic bile duct

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	494	10.0	9.1-10.9	365	16.6	14.9-18.4	129	4.6	3.8-5.5	341	6.9	6.2-7.7	228	10.6	9.3-12.1	113	3.9	3.2-4.8
2008	486	9.8	9.0-10.8	353	16.0	14.4-17.8	133	4.8	4.0-5.7	303	6.0	5.3-6.7	206	9.4	8.1-10.7	97	3.3	2.7-4.0
2009	495	9.9	9.1-10.9	347	15.7	14.0-17.4	148	5.3	4.4-6.2	355	7.1	6.4-7.9	237	11.0	9.6-12.5	118	4.2	3.4-5.0
2010	523	10.4	9.5-11.3	383	17.1	15.4-19.0	140	4.9	4.1-5.8	362	7.1	6.4-7.9	250	11.4	10.0-12.9	112	3.6	3.0-4.4
2011	505	9.8	9.0-10.8	354	15.4	13.8-17.1	151	5.2	4.4-6.1	326	6.2	5.6-7.0	214	9.4	8.2-10.8	112	3.6	2.9-4.3
2012	549	10.6	9.7-11.5	389	16.6	15.0-18.4	160	5.6	4.7-6.6	347	6.6	5.9-7.3	217	9.6	8.3-10.9	130	4.2	3.5-5.1
2013	532	10.0	9.1-10.9	354	14.8	13.3-16.5	178	6.0	5.1-6.9	339	6.3	5.7-7.1	223	9.5	8.3-10.9	116	3.8	3.1-4.6
2014	539	9.9	9.1-10.8	367	15.0	13.5-16.6	172	5.7	4.8-6.6	356	6.4	5.8-7.2	230	9.4	8.2-10.7	126	4.0	3.3-4.8
2015	550	10.0	9.2-10.9	375	15.4	13.8-17.0	175	5.6	4.8-6.6	381	6.9	6.2-7.6	250	10.4	9.1-11.7	131	4.0	3.3-4.8
2016	522	9.4	8.6-10.2	369	14.7	13.2-16.3	153	5.0	4.2-5.9	417	7.6	6.8-8.3	275	11.4	10.1-12.9	142	4.4	3.7-5.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Pancreas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	821	13.3	12.3-14.2	410	16.5	14.8-18.2	411	11.1	10.1-12.3	737	12.0	11.2-13.0	366	14.8	13.3-16.5	371	10.2	9.2-11.3
1977	814	13.5	12.6-14.5	388	16.2	14.5-17.9	426	11.8	10.6-13.0	765	12.6	11.7-13.6	360	15.0	13.4-16.7	405	11.0	9.9-12.2
1978	783	12.7	11.8-13.7	392	15.7	14.1-17.4	391	10.7	9.6-11.8	715	11.6	10.8-12.6	353	14.3	12.8-15.9	362	9.9	8.9-11.1
1979	751	12.5	11.6-13.4	365	15.1	13.6-16.8	386	10.6	9.5-11.7	719	12.0	11.1-12.9	351	14.7	13.2-16.4	368	10.1	9.1-11.3
1980	800	13.4	12.5-14.4	390	16.7	15.0-18.5	410	11.3	10.2-12.5	732	12.2	11.3-13.2	369	15.6	14.0-17.3	363	10.0	8.9-11.1
1981	750	12.9	12.0-13.9	376	16.4	14.7-18.2	374	10.6	9.5-11.8	701	11.9	11.0-12.8	329	14.2	12.7-15.9	372	10.2	9.2-11.3
1982	787	13.7	12.7-14.7	348	15.8	14.2-17.7	439	12.3	11.1-13.5	664	11.4	10.5-12.3	304	13.6	12.0-15.2	360	10.0	9.0-11.1
1983	734	12.6	11.7-13.5	346	14.9	13.3-16.6	388	10.9	9.8-12.1	654	11.2	10.4-12.1	319	13.7	12.2-15.4	335	9.5	8.5-10.6
1984	764	13.2	12.3-14.2	342	15.4	13.8-17.2	422	11.8	10.7-13.0	669	11.6	10.7-12.5	302	13.4	11.9-15.0	367	10.3	9.2-11.4
1985	712	12.4	11.5-13.4	322	14.2	12.7-15.9	390	11.1	10.0-12.2	592	10.3	9.5-11.2	283	12.8	11.3-14.4	309	8.7	7.8-9.8
1986	729	13.0	12.1-14.0	317	14.6	13.0-16.4	412	11.9	10.8-13.2	612	10.6	9.8-11.5	262	11.9	10.5-13.5	350	9.7	8.7-10.8
1987	687	12.2	11.3-13.2	299	13.8	12.2-15.5	388	11.0	9.9-12.2	568	10.0	9.2-10.9	263	11.9	10.4-13.4	305	8.7	7.7-9.8
1988	678	12.2	11.3-13.2	284	13.3	11.7-14.9	394	11.3	10.2-12.5	558	10.0	9.2-10.9	257	12.1	10.6-13.7	301	8.6	7.6-9.6
1989	674	12.4	11.5-13.4	302	14.3	12.7-16.0	372	11.2	10.0-12.4	591	10.8	10.0-11.7	266	12.6	11.1-14.3	325	9.5	8.5-10.7
1990	656	12.2	11.3-13.2	298	14.2	12.6-15.9	358	10.8	9.7-12.0	551	10.2	9.3-11.1	250	12.0	10.6-13.7	301	8.9	7.9-10.0
1991	658	12.4	11.5-13.4	291	14.2	12.6-16.0	367	11.1	9.9-12.3	536	10.1	9.3-11.0	248	12.1	10.6-13.7	288	8.7	7.7-9.8
1992	680	13.0	12.0-14.0	318	15.7	14.0-17.6	362	11.0	9.9-12.3	527	10.0	9.1-10.9	230	11.3	9.9-12.9	297	8.9	7.9-10.0
1993	665	12.9	11.9-13.9	285	14.0	12.4-15.8	380	11.8	10.7-13.1	581	11.1	10.3-12.1	279	13.8	12.2-15.5	302	9.2	8.2-10.4
1994	710	13.9	12.9-15.0	309	15.5	13.8-17.3	401	12.6	11.3-13.9	600	11.6	10.7-12.6	244	12.2	10.7-13.9	356	11.0	9.9-12.3
1995	620	12.0	11.1-13.0	261	13.1	11.6-14.9	359	11.4	10.2-12.7	544	10.5	9.6-11.4	243	12.2	10.7-13.9	301	9.2	8.2-10.3
1996	656	13.0	12.0-14.0	296	14.9	13.3-16.7	360	11.7	10.5-13.0	623	12.2	11.3-13.3	275	13.9	12.3-15.7	348	11.2	10.0-12.5
1997	654	13.0	12.0-14.0	297	14.9	13.2-16.7	357	11.6	10.4-12.9	524	10.4	9.5-11.3	230	11.8	10.3-13.4	294	9.3	8.3-10.5
1998	677	13.5	12.5-14.5	323	16.2	14.4-18.0	354	11.5	10.4-12.8	548	10.8	9.9-11.7	247	12.4	10.9-14.1	301	9.7	8.7-10.9
1999	667	13.2	12.2-14.2	327	16.1	14.4-18.0	340	11.1	9.9-12.3	573	11.3	10.4-12.3	295	14.6	12.9-16.3	278	8.9	7.9-10.1
2000	721	14.1	13.1-15.2	327	16.0	14.3-17.8	394	12.7	11.5-14.1	545	10.5	9.7-11.5	237	11.6	10.1-13.1	308	9.6	8.6-10.8
2001	653	12.7	11.7-13.7	288	14.0	12.4-15.7	365	11.6	10.4-12.9	582	11.2	10.3-12.1	264	13.0	11.5-14.7	318	9.9	8.9-11.1
2002	624	12.1	11.2-13.1	290	13.9	12.4-15.6	334	10.5	9.4-11.7	510	9.8	9.0-10.7	217	10.5	9.2-12.0	293	9.2	8.1-10.3
2003	698	13.6	12.6-14.6	320	15.5	13.9-17.3	378	12.0	10.8-13.3	539	10.4	9.6-11.3	264	12.7	11.2-14.3	275	8.5	7.5-9.6
2004	695	13.5	12.5-14.6	308	14.8	13.2-16.6	387	12.5	11.3-13.9	554	10.6	9.8-11.6	258	12.5	11.0-14.1	296	9.2	8.2-10.4
2005	664	13.1	12.1-14.1	324	15.4	13.8-17.2	340	11.4	10.2-12.7	557	10.8	10.0-11.8	269	12.9	11.4-14.6	288	9.4	8.3-10.6
2006	697	13.7	12.7-14.8	325	15.7	14.0-17.5	372	12.2	11.0-13.6	556	10.9	10.0-11.8	253	12.3	10.8-13.9	303	9.8	8.7-11.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Pancreas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	716	13.9	12.9-15.0	348	16.6	14.9-18.4	368	11.9	10.7-13.3	564	10.9	10.0-11.9	262	12.6	11.1-14.2	302	9.8	8.7-11.0
2008	773	15.2	14.1-16.3	368	17.6	15.8-19.5	405	13.3	12.0-14.7	599	11.7	10.7-12.6	271	13.0	11.5-14.7	328	10.4	9.3-11.7
2009	770	15.1	14.0-16.2	381	18.0	16.2-19.9	389	13.0	11.7-14.4	616	12.0	11.1-13.0	317	15.1	13.5-16.9	299	9.8	8.7-11.0
2010	747	14.7	13.6-15.8	368	17.2	15.5-19.1	379	12.7	11.4-14.1	598	11.5	10.6-12.5	285	13.3	11.8-15.0	313	9.9	8.8-11.1
2011	737	14.2	13.2-15.3	363	16.7	15.0-18.5	374	12.3	11.1-13.7	604	11.5	10.6-12.5	298	13.8	12.3-15.5	306	9.8	8.7-11.1
2012	739	14.2	13.2-15.3	358	16.2	14.6-18.0	381	12.6	11.3-14.0	593	11.1	10.2-12.1	295	13.2	11.7-14.8	298	9.6	8.5-10.8
2013	726	13.6	12.6-14.6	358	15.7	14.1-17.5	368	11.8	10.6-13.1	546	10.1	9.2-11.0	271	12.1	10.7-13.7	275	8.5	7.5-9.7
2014	742	13.7	12.7-14.8	380	16.7	15.0-18.5	362	11.4	10.2-12.7	595	10.8	10.0-11.8	300	13.3	11.8-14.9	295	8.9	7.9-10.0
2015	751	13.7	12.7-14.7	383	16.3	14.7-18.1	368	11.6	10.4-12.9	556	9.9	9.1-10.8	270	11.5	10.1-13.0	286	8.7	7.7-9.8
2016	707	12.7	11.8-13.7	338	14.2	12.7-15.8	369	11.5	10.3-12.8	709	12.7	11.8-13.7	345	14.5	13.0-16.2	364	11.4	10.2-12.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Larynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	352	5.4	4.9-6.1	287	10.6	9.3-11.9	65	1.8	1.4-2.4	137	2.1	1.8-2.5	108	4.1	3.3-5.0	29	0.8	0.5-1.2
1977	336	5.5	4.9-6.1	287	11.1	9.8-12.5	49	1.4	1.0-1.9	140	2.2	1.8-2.6	125	4.8	4.0-5.8	15	0.4	0.2-0.7
1978	315	5.3	4.7-5.9	250	10.2	8.9-11.6	65	1.8	1.4-2.4	127	2.1	1.7-2.5	102	4.1	3.3-5.0	25	0.7	0.4-1.1
1979	357	5.9	5.3-6.6	293	11.5	10.2-12.9	64	2.0	1.5-2.6	131	2.1	1.8-2.5	106	4.3	3.5-5.3	25	0.7	0.4-1.1
1980	367	6.4	5.7-7.1	302	12.5	11.1-14.1	65	2.0	1.5-2.6	122	2.1	1.7-2.5	104	4.4	3.5-5.3	18	0.5	0.3-0.8
1981	342	6.0	5.3-6.7	272	11.4	10.1-12.9	70	2.1	1.6-2.7	114	1.9	1.6-2.3	93	3.8	3.1-4.7	21	0.6	0.4-0.9
1982	333	5.8	5.2-6.5	263	11.1	9.8-12.6	70	2.2	1.7-2.8	121	2.0	1.7-2.4	97	4.0	3.3-5.0	24	0.7	0.4-1.1
1983	314	5.6	5.0-6.3	240	10.0	8.8-11.4	74	2.3	1.8-3.0	109	1.9	1.5-2.3	83	3.5	2.8-4.4	26	0.7	0.5-1.1
1984	360	6.4	5.7-7.1	283	12.0	10.6-13.5	77	2.5	1.9-3.1	106	1.9	1.5-2.3	85	3.6	2.8-4.5	21	0.7	0.4-1.1
1985	346	6.3	5.7-7.1	267	11.5	10.1-13.0	79	2.6	2.0-3.3	118	2.1	1.7-2.5	95	4.2	3.3-5.1	23	0.7	0.4-1.1
1986	311	5.7	5.1-6.4	240	10.5	9.2-12.0	71	2.4	1.9-3.1	81	1.4	1.1-1.7	65	2.9	2.2-3.7	16	0.4	0.3-0.7
1987	295	5.7	5.1-6.4	232	10.5	9.2-12.0	63	2.3	1.7-3.0	99	1.8	1.5-2.2	77	3.5	2.7-4.4	22	0.7	0.4-1.0
1988	311	5.9	5.3-6.7	246	11.0	9.6-12.5	65	2.2	1.7-2.9	95	1.8	1.5-2.2	75	3.5	2.7-4.4	20	0.7	0.4-1.1
1989	288	5.8	5.2-6.6	216	10.2	8.8-11.7	72	2.7	2.1-3.4	96	1.8	1.5-2.2	68	3.1	2.4-3.9	28	0.9	0.6-1.4
1990	313	6.2	5.6-7.0	256	12.1	10.7-13.7	57	2.0	1.5-2.6	84	1.6	1.3-2.0	75	3.5	2.8-4.5	9	0.3	0.1-0.5
1991	288	5.9	5.3-6.7	234	11.3	9.9-12.9	54	2.0	1.5-2.6	97	1.9	1.6-2.4	75	3.6	2.8-4.5	22	0.7	0.5-1.1
1992	291	6.0	5.3-6.7	221	10.6	9.2-12.1	70	2.6	2.0-3.3	109	2.2	1.8-2.7	82	4.1	3.3-5.1	27	1.0	0.6-1.4
1993	268	5.5	4.9-6.3	210	10.1	8.8-11.6	58	2.1	1.6-2.8	99	1.9	1.6-2.4	81	3.9	3.1-4.8	18	0.6	0.3-0.9
1994	262	5.4	4.8-6.1	208	10.3	8.9-11.8	54	2.0	1.5-2.6	95	1.9	1.5-2.3	76	3.7	2.9-4.6	19	0.7	0.4-1.1
1995	217	4.5	3.9-5.1	171	8.2	7.0-9.5	46	1.6	1.2-2.2	90	1.9	1.5-2.3	68	3.4	2.6-4.3	22	0.7	0.5-1.1
1996	282	6.0	5.3-6.7	223	11.0	9.6-12.6	59	2.2	1.6-2.8	99	2.0	1.6-2.5	83	4.1	3.3-5.1	16	0.5	0.3-0.9
1997	246	5.2	4.6-5.9	205	10.1	8.8-11.6	41	1.5	1.1-2.1	75	1.6	1.2-2.0	57	2.9	2.2-3.7	18	0.7	0.4-1.1
1998	242	5.0	4.4-5.7	185	9.0	7.7-10.4	57	2.1	1.6-2.7	83	1.7	1.4-2.1	66	3.2	2.5-4.1	17	0.6	0.3-1.0
1999	214	4.4	3.8-5.1	175	8.4	7.2-9.8	39	1.4	1.0-2.0	77	1.6	1.2-2.0	60	2.9	2.2-3.7	17	0.6	0.3-1.0
2000	234	4.8	4.2-5.5	178	8.6	7.3-9.9	56	2.0	1.5-2.7	85	1.7	1.4-2.1	68	3.3	2.6-4.2	17	0.6	0.4-1.0
2001	225	4.6	4.0-5.2	168	8.0	6.9-9.4	57	2.1	1.6-2.7	61	1.2	0.9-1.5	48	2.3	1.7-3.1	13	0.4	0.2-0.7
2002	209	4.2	3.7-4.8	173	8.2	7.0-9.5	36	1.3	0.9-1.8	77	1.5	1.2-1.9	61	2.9	2.2-3.8	16	0.5	0.3-0.8
2003	227	4.6	4.0-5.2	176	8.3	7.1-9.7	51	1.8	1.3-2.4	78	1.5	1.2-1.9	60	2.9	2.2-3.7	18	0.6	0.3-0.9
2004	229	4.6	4.0-5.3	192	8.9	7.7-10.3	37	1.4	0.9-1.9	69	1.4	1.1-1.8	64	3.0	2.3-3.8	5	0.2	0.1-0.4
2005	209	4.2	3.7-4.9	163	7.5	6.4-8.8	46	1.7	1.2-2.2	65	1.3	1.0-1.6	52	2.5	1.8-3.2	13	0.4	0.2-0.7
2006	190	3.9	3.4-4.5	157	7.4	6.3-8.6	33	1.3	0.9-1.8	68	1.4	1.1-1.7	58	2.8	2.1-3.6	10	0.3	0.1-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Larynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	199	4.1	3.5-4.7	151	7.1	6.0-8.3	48	1.7	1.3-2.3	63	1.3	1.0-1.6	48	2.2	1.7-3.0	15	0.5	0.3-0.9
2008	229	4.7	4.1-5.4	177	8.3	7.1-9.7	52	1.9	1.4-2.6	61	1.3	1.0-1.6	53	2.5	1.9-3.3	8	0.3	0.1-0.6
2009	199	4.0	3.5-4.6	151	6.8	5.8-8.0	48	1.8	1.3-2.4	68	1.3	1.0-1.7	52	2.4	1.8-3.2	16	0.5	0.3-0.9
2010	195	3.9	3.4-4.5	160	7.3	6.2-8.6	35	1.3	0.9-1.8	62	1.2	0.9-1.5	47	2.2	1.6-2.9	15	0.5	0.3-0.8
2011	154	3.0	2.5-3.5	125	5.5	4.5-6.5	29	1.0	0.7-1.5	60	1.2	0.9-1.5	48	2.2	1.6-3.0	12	0.4	0.2-0.7
2012	198	3.8	3.3-4.4	165	7.2	6.1-8.4	33	1.2	0.8-1.7	57	1.1	0.8-1.4	44	1.9	1.4-2.6	13	0.5	0.3-0.8
2013	177	3.4	2.9-3.9	149	6.5	5.5-7.7	28	1.0	0.6-1.4	56	1.1	0.8-1.4	46	2.1	1.5-2.8	10	0.4	0.2-0.7
2014	173	3.3	2.8-3.8	133	5.7	4.8-6.8	40	1.3	0.9-1.8	42	0.8	0.6-1.1	35	1.5	1.0-2.1	7	0.2	0.1-0.5
2015	163	3.0	2.6-3.5	127	5.3	4.4-6.3	36	1.2	0.8-1.7	51	1.0	0.7-1.3	39	1.6	1.2-2.3	12	0.4	0.2-0.8
2016	148	2.7	2.3-3.2	122	5.0	4.2-6.0	26	0.9	0.6-1.3	64	1.1	0.9-1.5	59	2.4	1.8-3.2	5	0.1	0.0-0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Lung and bronchus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	3166	49.6	47.9-51.4	2207	84.1	80.4-87.8	959	26.3	24.6-28.1	2564	40.1	38.6-41.8	1825	70.0	66.7-73.5	739	20.3	18.8-21.9
1977	3153	50.2	48.4-52.1	2153	84.5	80.8-88.3	1000	27.5	25.8-29.4	2596	41.2	39.6-42.8	1853	72.5	69.1-76.0	743	20.4	18.9-22.0
1978	3175	51.9	50.0-53.8	2228	89.0	85.2-92.9	947	27.1	25.4-29.0	2573	42.1	40.4-43.8	1821	73.5	70.0-77.0	752	21.4	19.8-23.0
1979	3218	53.0	51.1-54.9	2160	87.7	84.0-91.7	1058	30.2	28.3-32.2	2554	42.0	40.4-43.7	1767	72.2	68.8-75.8	787	22.4	20.8-24.1
1980	3184	53.3	51.5-55.3	2122	87.7	83.9-91.6	1062	30.8	28.9-32.8	2665	44.5	42.8-46.2	1784	74.1	70.6-77.7	881	25.4	23.7-27.2
1981	3059	51.9	50.0-53.8	1975	83.5	79.8-87.4	1084	31.3	29.4-33.3	2484	42.2	40.5-44.0	1676	71.1	67.6-74.7	808	23.4	21.8-25.2
1982	3229	55.5	53.6-57.5	2033	87.1	83.2-91.1	1196	35.1	33.1-37.3	2548	43.4	41.7-45.2	1671	71.4	67.9-75.0	877	25.6	23.8-27.4
1983	3232	55.8	53.9-57.8	2045	88.0	84.1-92.0	1187	35.0	33.0-37.2	2510	43.2	41.4-44.9	1535	67.1	63.7-70.7	975	28.1	26.3-30.0
1984	3220	56.8	54.8-58.8	2064	90.0	86.0-94.1	1156	35.2	33.1-37.4	2502	43.8	42.1-45.6	1635	72.1	68.6-75.8	867	25.5	23.8-27.4
1985	3217	56.5	54.5-58.5	2012	87.7	83.8-91.7	1205	36.2	34.1-38.4	2508	43.9	42.2-45.7	1550	68.4	65.0-72.0	958	28.5	26.6-30.4
1986	3163	56.5	54.5-58.6	1932	85.5	81.6-89.4	1231	37.4	35.2-39.6	2426	42.8	41.1-44.6	1469	65.0	61.6-68.5	957	28.9	27.0-30.9
1987	3209	57.7	55.7-59.8	1947	86.8	82.9-90.8	1262	38.7	36.5-41.0	2495	44.7	42.9-46.5	1549	69.7	66.2-73.4	946	28.7	26.8-30.6
1988	3128	57.6	55.5-59.7	1847	84.8	80.9-88.9	1281	40.0	37.8-42.4	2454	44.7	42.9-46.6	1527	69.9	66.4-73.6	927	28.7	26.8-30.7
1989	3113	58.6	56.5-60.7	1800	83.4	79.5-87.4	1313	42.2	39.9-44.7	2388	44.4	42.6-46.3	1433	67.2	63.7-70.8	955	29.9	28.0-32.0
1990	3113	59.4	57.3-61.6	1793	84.7	80.7-88.7	1320	43.0	40.6-45.5	2379	44.8	43.0-46.7	1403	66.4	62.9-70.0	976	31.1	29.1-33.2
1991	3149	61.5	59.3-63.7	1764	84.7	80.8-88.8	1385	46.0	43.5-48.6	2414	46.5	44.6-48.4	1368	66.1	62.6-69.7	1046	33.8	31.8-36.0
1992	3264	64.0	61.8-66.2	1879	90.4	86.4-94.7	1385	46.7	44.2-49.3	2341	45.4	43.5-47.3	1372	66.4	62.9-70.0	969	31.7	29.7-33.8
1993	3145	62.4	60.2-64.6	1758	85.6	81.6-89.8	1387	47.2	44.7-49.9	2381	46.6	44.7-48.5	1344	66.0	62.5-69.7	1037	34.2	32.1-36.4
1994	3148	63.0	60.8-65.3	1692	82.6	78.7-86.7	1456	50.1	47.5-52.8	2422	48.2	46.3-50.1	1419	70.0	66.4-73.8	1003	33.8	31.7-36.1
1995	3039	60.7	58.6-63.0	1678	82.1	78.2-86.2	1361	46.1	43.6-48.6	2352	46.7	44.8-48.6	1306	64.3	60.8-67.9	1046	34.8	32.7-37.1
1996	3028	61.3	59.2-63.6	1642	81.6	77.7-85.6	1386	48.1	45.5-50.7	2299	46.0	44.2-48.0	1281	63.6	60.1-67.2	1018	34.0	31.9-36.2
1997	3021	61.4	59.2-63.6	1638	81.0	77.1-85.1	1383	48.0	45.4-50.6	2161	43.5	41.7-45.4	1229	61.2	57.8-64.7	932	31.6	29.6-33.8
1998	2998	60.6	58.5-62.8	1595	78.1	74.3-82.1	1403	48.1	45.6-50.7	2196	44.1	42.2-46.0	1182	58.3	55.0-61.8	1014	34.3	32.2-36.5
1999	2881	57.7	55.6-59.9	1481	72.1	68.5-75.9	1400	47.9	45.4-50.5	2155	43.0	41.2-44.9	1155	56.7	53.4-60.0	1000	34.0	31.9-36.2
2000	3003	59.9	57.8-62.1	1576	76.3	72.5-80.1	1427	48.9	46.3-51.5	2113	41.6	39.8-43.4	1151	56.0	52.8-59.4	962	31.5	29.5-33.6
2001	2947	58.6	56.5-60.7	1517	72.9	69.3-76.7	1430	48.5	46.0-51.1	2127	41.8	40.1-43.7	1146	55.5	52.4-58.8	981	32.4	30.3-34.5
2002	2939	58.3	56.2-60.4	1540	73.4	69.8-77.2	1399	47.5	45.0-50.1	2081	40.8	39.1-42.6	1117	53.5	50.4-56.7	964	31.7	29.7-33.8
2003	2816	55.9	53.8-58.0	1504	72.1	68.5-75.8	1312	44.3	41.9-46.8	2054	40.4	38.7-42.2	1095	52.5	49.5-55.7	959	31.8	29.8-33.9
2004	2783	55.2	53.2-57.3	1484	70.9	67.3-74.6	1299	44.4	42.0-46.9	1938	38.1	36.4-39.8	1071	51.5	48.4-54.6	867	28.8	26.9-30.9
2005	2818	56.2	54.1-58.3	1485	71.2	67.6-74.9	1333	46.1	43.7-48.7	1895	37.6	35.9-39.3	1056	50.9	47.9-54.1	839	28.6	26.6-30.6
2006	2798	56.4	54.3-58.6	1444	69.7	66.2-73.5	1354	47.2	44.7-49.8	1823	36.3	34.7-38.0	1007	48.8	45.8-51.9	816	27.4	25.5-29.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Lung and bronchus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	2906	58.7	56.6-60.9	1509	72.6	69.0-76.4	1397	49.0	46.4-51.7	1907	37.9	36.2-39.7	1004	48.7	45.7-51.8	903	30.7	28.6-32.8
2008	2814	56.7	54.6-58.8	1413	67.6	64.1-71.3	1401	49.2	46.6-51.9	1880	37.4	35.7-39.1	1013	48.7	45.7-51.8	867	29.4	27.4-31.4
2009	2839	56.7	54.6-58.8	1426	67.9	64.4-71.6	1413	49.1	46.5-51.8	1747	34.5	32.9-36.1	887	42.6	39.8-45.5	860	28.9	26.9-30.9
2010	2666	53.3	51.2-55.4	1364	64.4	61.0-67.9	1302	45.2	42.8-47.8	1823	35.9	34.2-37.6	945	44.9	42.1-47.9	878	29.6	27.6-31.7
2011	2742	53.8	51.7-55.8	1378	63.8	60.4-67.3	1364	46.7	44.2-49.3	1732	33.3	31.8-35.0	908	42.3	39.5-45.1	824	26.8	25.0-28.8
2012	2678	51.9	49.9-53.9	1306	59.3	56.1-62.7	1372	47.1	44.6-49.8	1703	32.7	31.1-34.3	922	42.0	39.3-44.8	781	26.1	24.3-28.1
2013	2726	51.9	50.0-53.9	1345	60.0	56.8-63.3	1381	46.4	43.9-49.0	1752	33.0	31.5-34.6	920	41.5	38.8-44.3	832	27.3	25.4-29.3
2014	2628	49.0	47.1-50.9	1285	55.7	52.7-58.9	1343	44.3	41.9-46.8	1527	28.1	26.7-29.5	797	34.9	32.5-37.5	730	23.1	21.4-24.9
2015	2558	47.1	45.3-49.0	1321	56.8	53.7-60.0	1237	40.3	38.0-42.6	1550	28.0	26.6-29.5	804	34.7	32.3-37.2	746	23.3	21.6-25.1
2016	2543	46.4	44.6-48.2	1215	51.5	48.6-54.6	1328	43.0	40.7-45.4	1600	28.8	27.4-30.2	828	35.6	33.2-38.1	772	24.1	22.4-25.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Melanoma of the skin

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	386	6.6	6.0-7.4	199	8.0	6.9-9.3	187	5.7	4.9-6.6	146	2.5	2.1-3.0	81	3.2	2.5-4.1	65	2.0	1.5-2.6
1977	387	6.6	6.0-7.4	172	6.9	5.8-8.0	215	6.5	5.6-7.4	142	2.4	2.0-2.8	72	2.9	2.3-3.7	70	2.0	1.5-2.5
1978	353	6.5	5.8-7.3	168	7.1	6.1-8.3	185	6.1	5.2-7.2	113	1.9	1.6-2.3	69	2.9	2.3-3.8	44	1.3	0.9-1.7
1979	391	7.3	6.5-8.0	194	8.0	6.9-9.2	197	6.8	5.8-7.8	140	2.6	2.1-3.0	75	3.2	2.5-4.0	65	2.0	1.6-2.6
1980	413	7.7	7.0-8.5	204	8.6	7.4-9.9	209	7.2	6.2-8.3	136	2.5	2.0-2.9	70	3.0	2.3-3.8	66	2.1	1.6-2.7
1981	372	7.0	6.3-7.8	185	7.9	6.8-9.2	187	6.4	5.4-7.4	156	3.0	2.5-3.5	87	3.9	3.1-4.8	69	2.3	1.7-2.9
1982	381	7.2	6.5-8.0	206	9.2	7.9-10.6	175	5.8	5.0-6.8	129	2.3	1.9-2.8	73	3.2	2.5-4.0	56	1.7	1.3-2.2
1983	432	8.2	7.4-9.0	217	9.6	8.4-11.1	215	7.3	6.3-8.4	133	2.5	2.1-3.0	77	3.5	2.8-4.4	56	1.9	1.4-2.4
1984	471	8.9	8.0-9.7	239	10.4	9.1-11.8	232	7.9	6.9-9.1	158	2.9	2.4-3.4	94	4.1	3.3-5.0	64	2.0	1.5-2.6
1985	423	8.1	7.3-9.0	241	10.6	9.3-12.1	182	6.4	5.5-7.5	165	3.1	2.7-3.7	88	4.0	3.2-4.9	77	2.6	2.0-3.3
1986	434	8.3	7.5-9.2	213	9.6	8.3-11.0	221	7.6	6.6-8.7	139	2.6	2.1-3.0	72	3.2	2.5-4.0	67	2.2	1.7-2.8
1987	425	8.3	7.5-9.2	227	10.5	9.2-12.0	198	6.8	5.9-7.9	141	2.7	2.2-3.2	83	3.8	3.0-4.7	58	2.0	1.5-2.6
1988	393	7.7	7.0-8.6	212	9.9	8.6-11.4	181	6.1	5.2-7.1	125	2.4	2.0-2.9	73	3.4	2.7-4.3	52	1.7	1.3-2.3
1989	396	7.9	7.1-8.7	202	9.5	8.2-11.0	194	6.7	5.7-7.7	105	2.1	1.7-2.6	67	3.3	2.5-4.2	38	1.4	1.0-2.0
1990	395	8.0	7.2-8.8	214	10.1	8.8-11.6	181	6.5	5.6-7.6	115	2.3	1.9-2.8	65	3.2	2.4-4.0	50	1.8	1.3-2.4
1991	414	8.3	7.5-9.1	227	10.7	9.3-12.2	187	6.5	5.6-7.5	112	2.2	1.8-2.6	58	2.8	2.1-3.7	54	1.8	1.3-2.3
1992	351	7.1	6.4-7.9	180	8.6	7.4-10.0	171	6.0	5.1-7.0	94	1.8	1.5-2.3	58	2.7	2.0-3.5	36	1.2	0.8-1.6
1993	364	7.3	6.6-8.1	206	10.0	8.7-11.5	158	5.5	4.7-6.5	84	1.7	1.3-2.1	49	2.5	1.8-3.3	35	1.2	0.8-1.7
1994	433	8.8	8.0-9.7	245	11.8	10.4-13.4	188	6.7	5.8-7.8	111	2.2	1.8-2.7	72	3.5	2.7-4.4	39	1.4	1.0-1.9
1995	426	8.8	8.0-9.7	250	12.0	10.5-13.6	176	6.4	5.5-7.5	115	2.4	2.0-2.9	65	3.2	2.5-4.1	50	1.8	1.3-2.4
1996	460	9.5	8.6-10.4	242	11.7	10.3-13.3	218	8.0	7.0-9.2	112	2.3	1.9-2.7	57	2.7	2.1-3.5	55	1.9	1.4-2.5
1997	492	10.1	9.3-11.1	269	13.3	11.8-15.0	223	8.1	7.1-9.3	109	2.3	1.9-2.7	60	2.9	2.2-3.8	49	1.8	1.3-2.4
1998	505	10.4	9.5-11.3	266	12.8	11.3-14.4	239	8.7	7.6-9.9	126	2.5	2.1-3.0	72	3.6	2.8-4.5	54	1.8	1.4-2.4
1999	499	10.1	9.2-11.0	272	12.9	11.4-14.5	227	8.3	7.2-9.4	106	2.2	1.8-2.6	64	3.1	2.4-3.9	42	1.4	1.0-2.0
2000	596	12.1	11.1-13.1	312	14.9	13.2-16.6	284	10.5	9.3-11.8	124	2.5	2.1-3.0	72	3.5	2.8-4.5	52	1.8	1.4-2.4
2001	607	12.4	11.4-13.4	327	15.5	13.9-17.3	280	10.2	9.1-11.6	112	2.2	1.8-2.7	60	2.9	2.2-3.8	52	1.8	1.3-2.4
2002	616	12.4	11.4-13.4	333	15.7	14.0-17.4	283	10.4	9.2-11.7	99	2.0	1.6-2.4	50	2.4	1.8-3.2	49	1.7	1.3-2.3
2003	538	10.9	10.0-11.9	281	13.2	11.7-14.8	257	9.5	8.3-10.8	107	2.1	1.7-2.6	50	2.4	1.8-3.2	57	2.0	1.5-2.6
2004	751	15.3	14.2-16.4	397	18.6	16.8-20.5	354	13.3	11.9-14.7	99	1.9	1.6-2.4	53	2.5	1.9-3.3	46	1.6	1.1-2.1
2005	819	16.7	15.6-17.9	432	20.2	18.3-22.2	387	14.6	13.2-16.2	96	1.9	1.5-2.3	49	2.4	1.8-3.2	47	1.6	1.2-2.2
2006	995	20.5	19.2-21.8	546	25.4	23.3-27.6	449	17.0	15.4-18.6	110	2.2	1.8-2.7	67	3.3	2.5-4.2	43	1.5	1.1-2.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Melanoma of the skin

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	804	16.5	15.4-17.7	421	19.6	17.8-21.6	383	14.5	13.1-16.1	106	2.0	1.7-2.5	63	3.0	2.3-3.9	43	1.4	1.0-2.0
2008	826	16.9	15.8-18.1	450	21.1	19.2-23.1	376	14.3	12.8-15.8	100	2.0	1.6-2.4	57	2.7	2.1-3.5	43	1.5	1.1-2.1
2009	900	18.1	17.0-19.4	485	22.5	20.5-24.6	415	15.3	13.8-16.9	97	2.0	1.6-2.4	63	3.0	2.3-3.8	34	1.2	0.8-1.7
2010	783	15.9	14.8-17.1	421	19.6	17.7-21.5	362	13.5	12.1-15.0	101	2.0	1.6-2.4	59	2.7	2.0-3.5	42	1.4	1.0-1.9
2011	925	18.4	17.2-19.7	498	22.5	20.6-24.6	427	15.8	14.3-17.5	82	1.6	1.2-1.9	43	2.0	1.4-2.7	39	1.2	0.9-1.7
2012	816	16.0	14.9-17.1	443	19.7	17.9-21.6	373	13.5	12.1-15.0	124	2.3	1.9-2.8	69	3.1	2.4-3.9	55	1.8	1.3-2.3
2013	888	17.1	16.0-18.3	505	22.2	20.2-24.2	383	13.7	12.3-15.2	101	1.9	1.5-2.3	64	2.8	2.1-3.6	37	1.2	0.9-1.7
2014	853	16.3	15.2-17.5	459	20.2	18.4-22.2	394	13.8	12.4-15.3	98	1.8	1.5-2.2	67	2.9	2.3-3.7	31	0.9	0.6-1.3
2015	822	15.6	14.5-16.7	461	20.0	18.2-22.0	361	12.4	11.2-13.8	66	1.3	1.0-1.6	40	1.8	1.3-2.4	26	1.0	0.6-1.4
2016	823	15.5	14.5-16.7	460	19.6	17.8-21.5	363	12.7	11.4-14.1	101	1.8	1.5-2.2	61	2.6	2.0-3.3	40	1.3	0.9-1.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Female breast

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	3506	106.0	102.3-109.8	1353	38.7	36.5-40.9
1977	3318	102.4	98.7-106.1	1339	39.3	37.2-41.6
1978	3352	104.6	100.9-108.4	1332	39.6	37.4-41.9
1979	3266	104.1	100.3-107.9	1199	36.6	34.5-38.9
1980	3218	104.1	100.3-108.0	1314	39.8	37.6-42.1
1981	3360	109.9	106.0-113.9	1273	39.3	37.1-41.7
1982	3274	108.2	104.3-112.2	1190	37.0	34.8-39.2
1983	3266	108.2	104.3-112.2	1119	35.2	33.0-37.4
1984	3474	115.9	111.9-120.1	1186	37.7	35.5-40.0
1985	3436	115.1	111.1-119.2	1153	36.5	34.3-38.8
1986	3326	112.8	108.8-116.9	1102	35.0	32.9-37.3
1987	3426	116.6	112.5-120.8	1069	34.3	32.2-36.5
1988	3429	118.6	114.5-122.8	1162	37.3	35.1-39.7
1989	3373	119.5	115.3-123.8	1097	36.4	34.2-38.7
1990	3599	130.4	126.0-135.0	1031	34.6	32.4-36.9
1991	3632	133.1	128.7-137.7	986	33.5	31.4-35.8
1992	3616	133.9	129.4-138.5	977	33.3	31.2-35.5
1993	3410	126.8	122.4-131.2	987	33.6	31.5-35.9
1994	3752	138.8	134.3-143.5	987	34.0	31.8-36.2
1995	3657	136.6	132.1-141.2	1030	35.7	33.5-38.0
1996	3833	144.2	139.5-148.9	1051	36.9	34.6-39.3
1997	3665	137.5	133.0-142.1	904	32.0	29.9-34.2
1998	3728	139.7	135.2-144.3	842	29.6	27.6-31.7
1999	3725	138.7	134.3-143.4	818	28.1	26.2-30.2
2000	3612	134.0	129.6-138.5	812	27.6	25.7-29.6
2001	3466	128.0	123.7-132.4	828	28.3	26.3-30.3
2002	3394	125.1	120.8-129.4	762	25.7	23.9-27.7
2003	3308	123.0	118.8-127.4	742	25.3	23.5-27.3
2004	3285	122.7	118.4-127.0	763	25.9	24.1-27.9
2005	3199	120.2	116.0-124.5	720	24.7	22.9-26.6
2006	3274	124.9	120.5-129.3	711	24.7	22.9-26.7
2007	3414	131.1	126.7-135.7	663	22.9	21.1-24.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Female breast

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	3428	130.8	126.4-135.4	659	22.7	21.0-24.6
2009	3483	132.3	127.9-136.9	650	22.7	20.9-24.6
2010	3333	126.6	122.2-131.1	641	22.0	20.3-23.8
2011	3408	127.9	123.5-132.4	597	20.2	18.6-22.0
2012	3266	122.0	117.7-126.3	606	20.6	18.9-22.4
2013	3437	127.4	123.0-131.8	599	19.8	18.2-21.6
2014	3485	127.7	123.3-132.1	567	18.6	17.0-20.2
2015	3461	126.8	122.5-131.3	565	18.9	17.3-20.6
2016	3416	124.0	119.8-128.4	643	20.6	19.0-22.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Cervix uteri

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	383	12.7	11.4-14.1	129	3.9	3.2-4.7
1977	339	11.2	10.0-12.6	128	3.9	3.2-4.7
1978	341	12.2	10.9-13.6	126	4.1	3.4-4.9
1979	357	13.0	11.6-14.5	103	3.4	2.7-4.2
1980	319	12.0	10.7-13.5	105	3.3	2.7-4.0
1981	310	11.3	10.0-12.7	113	4.0	3.2-4.8
1982	310	11.8	10.5-13.3	109	3.7	3.0-4.5
1983	324	12.3	10.9-13.7	90	3.0	2.4-3.8
1984	304	11.5	10.2-13.0	77	2.7	2.1-3.4
1985	302	11.6	10.3-13.1	81	2.9	2.3-3.6
1986	268	10.3	9.0-11.6	66	2.3	1.7-2.9
1987	322	12.3	11.0-13.8	70	2.3	1.8-2.9
1988	295	11.4	10.1-12.9	77	2.9	2.2-3.6
1989	289	11.3	10.0-12.7	64	2.4	1.8-3.1
1990	321	12.8	11.4-14.4	89	3.3	2.7-4.1
1991	331	13.5	12.1-15.1	69	2.6	2.0-3.3
1992	325	12.8	11.4-14.3	62	2.3	1.8-3.0
1993	303	12.4	11.0-13.9	65	2.5	1.9-3.2
1994	346	14.1	12.6-15.7	95	3.7	3.0-4.5
1995	347	14.0	12.5-15.6	87	3.4	2.7-4.2
1996	339	13.8	12.4-15.4	96	3.6	2.9-4.4
1997	313	12.4	11.1-13.9	85	3.4	2.7-4.2
1998	287	11.5	10.2-12.9	75	2.9	2.3-3.6
1999	279	11.0	9.7-12.4	94	3.7	3.0-4.5
2000	260	10.3	9.1-11.7	81	3.0	2.4-3.7
2001	218	8.8	7.6-10.0	64	2.4	1.9-3.1
2002	242	9.8	8.6-11.2	71	2.7	2.1-3.4
2003	232	9.1	8.0-10.4	78	3.0	2.4-3.7
2004	223	8.8	7.6-10.0	69	2.7	2.1-3.4
2005	249	10.1	8.9-11.5	77	2.8	2.2-3.5
2006	217	8.7	7.5-9.9	76	2.9	2.3-3.6
2007	213	8.9	7.7-10.2	66	2.6	2.0-3.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Cervix uteri

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	231	9.4	8.2-10.7	61	2.3	1.7-3.0
2009	221	9.0	7.9-10.3	59	2.2	1.7-2.9
2010	177	7.4	6.4-8.7	64	2.4	1.8-3.1
2011	187	7.6	6.5-8.8	68	2.6	2.0-3.3
2012	206	8.4	7.2-9.6	62	2.3	1.8-3.0
2013	204	8.1	7.0-9.4	61	2.1	1.6-2.7
2014	189	7.5	6.5-8.7	75	2.6	2.0-3.3
2015	198	7.8	6.8-9.1	61	2.2	1.7-2.9
2016	191	7.5	6.5-8.7	67	2.5	1.9-3.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Corpus uterus and uterus, not otherwise specified

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	849	24.1	22.5-25.9	248	6.6	5.8-7.5
1977	749	21.7	20.1-23.4	198	5.4	4.7-6.3
1978	681	19.5	18.0-21.1	184	5.0	4.3-5.9
1979	754	22.4	20.8-24.1	169	4.6	3.9-5.4
1980	786	23.8	22.1-25.6	196	5.4	4.6-6.2
1981	734	22.3	20.6-24.0	171	4.8	4.1-5.6
1982	732	22.7	21.0-24.5	164	4.7	4.0-5.5
1983	752	23.0	21.3-24.8	165	4.6	3.9-5.4
1984	684	21.2	19.6-22.9	163	4.7	4.0-5.5
1985	698	21.9	20.2-23.6	130	3.8	3.1-4.5
1986	664	21.5	19.8-23.3	138	3.9	3.3-4.6
1987	665	21.6	19.9-23.4	153	4.5	3.8-5.4
1988	665	22.0	20.3-23.8	125	3.7	3.0-4.4
1989	635	21.3	19.6-23.1	130	3.9	3.3-4.7
1990	607	21.3	19.6-23.2	130	3.9	3.2-4.6
1991	696	24.8	23.0-26.8	144	4.7	3.9-5.5
1992	684	24.7	22.8-26.7	137	4.4	3.7-5.2
1993	655	23.5	21.7-25.4	146	4.7	3.9-5.5
1994	726	26.1	24.2-28.1	146	4.8	4.0-5.7
1995	727	26.7	24.7-28.7	155	5.1	4.3-6.0
1996	718	26.2	24.3-28.3	141	4.7	3.9-5.5
1997	767	28.3	26.3-30.4	148	5.0	4.2-5.8
1998	807	30.1	28.0-32.3	145	5.0	4.2-5.9
1999	814	29.7	27.7-31.9	153	5.2	4.4-6.1
2000	747	27.4	25.4-29.4	132	4.2	3.5-5.0
2001	805	29.5	27.5-31.7	125	4.2	3.5-5.0
2002	807	29.3	27.3-31.4	133	4.5	3.7-5.3
2003	805	29.2	27.2-31.3	140	4.7	4.0-5.6
2004	848	31.1	29.0-33.3	125	4.2	3.5-5.1
2005	824	30.2	28.1-32.4	144	5.0	4.2-5.9
2006	824	30.1	28.1-32.3	144	4.9	4.1-5.8
2007	773	28.3	26.3-30.4	152	5.2	4.4-6.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Corpus uterus and uterus, not otherwise specified

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	927	34.2	32.0-36.5	149	5.1	4.3-6.0
2009	849	31.4	29.3-33.7	125	4.3	3.5-5.1
2010	824	30.3	28.2-32.5	177	6.0	5.1-7.0
2011	890	32.2	30.1-34.5	165	5.6	4.7-6.5
2012	822	29.2	27.2-31.3	152	5.0	4.2-5.9
2013	917	32.5	30.4-34.7	159	5.3	4.5-6.2
2014	857	29.7	27.7-31.9	151	5.1	4.3-6.0
2015	892	31.3	29.2-33.5	188	6.0	5.2-7.0
2016	896	31.0	29.0-33.2	221	7.1	6.2-8.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Ovary

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	565	16.2	14.8-17.6	381	10.7	9.6-11.9
1977	507	15.6	14.2-17.1	395	11.2	10.1-12.5
1978	529	16.4	14.9-17.9	335	9.7	8.7-10.9
1979	529	16.9	15.4-18.5	340	9.8	8.7-11.0
1980	523	16.5	15.0-18.0	354	10.4	9.3-11.6
1981	548	17.5	16.0-19.2	336	10.0	8.9-11.2
1982	554	18.2	16.6-19.9	314	9.5	8.4-10.6
1983	541	17.6	16.1-19.2	319	9.6	8.5-10.8
1984	521	17.0	15.5-18.6	328	10.0	8.9-11.1
1985	555	18.6	17.0-20.3	296	9.4	8.3-10.6
1986	475	15.9	14.5-17.5	316	9.9	8.8-11.1
1987	516	17.9	16.3-19.6	304	9.7	8.6-10.9
1988	481	16.6	15.0-18.2	342	10.8	9.6-12.1
1989	473	16.4	14.9-18.0	269	8.6	7.6-9.8
1990	489	17.7	16.1-19.4	283	9.3	8.2-10.5
1991	526	18.9	17.2-20.6	312	10.5	9.4-11.8
1992	481	17.8	16.2-19.5	311	10.7	9.5-12.0
1993	460	17.1	15.6-18.8	265	8.8	7.8-10.0
1994	485	17.7	16.1-19.4	274	9.4	8.3-10.6
1995	542	20.3	18.6-22.2	277	9.7	8.5-10.9
1996	481	17.9	16.3-19.6	280	10.0	8.8-11.2
1997	484	18.1	16.5-19.8	289	10.2	9.0-11.5
1998	439	16.5	14.9-18.1	248	8.7	7.6-9.9
1999	435	16.2	14.7-17.8	257	8.9	7.9-10.1
2000	417	15.2	13.8-16.8	265	9.1	8.0-10.3
2001	462	17.1	15.6-18.8	239	8.3	7.3-9.4
2002	456	16.3	14.9-17.9	263	9.1	8.0-10.2
2003	396	14.5	13.1-16.1	269	9.3	8.2-10.5
2004	426	15.7	14.2-17.3	265	9.0	8.0-10.2
2005	378	14.1	12.7-15.6	244	8.7	7.6-9.8
2006	372	13.7	12.3-15.2	246	8.6	7.5-9.7
2007	400	15.0	13.5-16.6	230	8.0	6.9-9.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Ovary

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	385	14.2	12.7-15.7	232	8.0	7.0-9.1
2009	393	14.9	13.4-16.5	216	7.2	6.2-8.3
2010	368	13.7	12.3-15.2	223	7.6	6.6-8.7
2011	392	14.3	12.9-15.9	236	7.8	6.8-8.9
2012	338	12.1	10.8-13.6	233	8.0	7.0-9.1
2013	354	12.9	11.6-14.4	230	7.7	6.7-8.8
2014	362	12.9	11.6-14.4	201	6.5	5.6-7.5
2015	369	13.4	12.0-14.9	195	6.5	5.6-7.6
2016	333	11.9	10.6-13.3	245	7.9	7.0-9.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Prostate

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1576	67.8	64.3-71.5	563	26.3	24.1-28.7
1977	1573	67.4	64.0-71.0	572	26.2	24.0-28.6
1978	1553	67.5	64.0-71.1	591	27.4	25.2-29.8
1979	1701	75.1	71.4-78.9	533	25.8	23.5-28.1
1980	1814	80.5	76.7-84.4	552	26.8	24.5-29.2
1981	1724	76.6	72.9-80.4	556	27.1	24.9-29.6
1982	1691	76.4	72.7-80.3	519	25.4	23.2-27.7
1983	1724	77.1	73.4-80.9	587	28.7	26.4-31.2
1984	1744	79.0	75.2-82.9	612	30.1	27.7-32.7
1985	1819	81.7	77.9-85.6	520	24.9	22.8-27.2
1986	1841	84.8	80.9-88.9	507	24.7	22.5-27.0
1987	1835	84.1	80.2-88.1	524	25.8	23.6-28.2
1988	1727	81.8	77.9-85.8	583	29.1	26.8-31.7
1989	1730	82.1	78.2-86.1	548	27.9	25.6-30.4
1990	2123	101.6	97.3-106.1	568	29.3	26.9-31.9
1991	2407	114.7	110.1-119.4	534	27.8	25.5-30.3
1992	2957	141.2	136.1-146.5	596	31.0	28.5-33.6
1993	2797	134.1	129.2-139.3	573	30.0	27.6-32.6
1994	2931	141.0	135.9-146.3	606	32.2	29.7-34.9
1995	2681	129.9	124.9-134.9	596	31.6	29.1-34.2
1996	2722	133.3	128.3-138.5	559	29.8	27.4-32.4
1997	2909	142.5	137.4-147.8	494	26.5	24.2-28.9
1998	2923	142.1	137.0-147.4	531	28.0	25.7-30.5
1999	2975	144.1	139.0-149.4	431	22.5	20.4-24.8
2000	3427	164.7	159.2-170.3	484	24.9	22.8-27.3
2001	3202	152.9	147.6-158.3	463	23.7	21.6-26.0
2002	3145	149.3	144.1-154.6	463	23.7	21.6-25.9
2003	2830	134.5	129.6-139.6	452	22.9	20.8-25.1
2004	2789	131.5	126.6-136.4	424	21.5	19.5-23.6
2005	2845	133.9	129.0-138.9	403	20.4	18.4-22.5
2006	3202	152.0	146.7-157.4	382	19.3	17.4-21.4
2007	3315	155.7	150.4-161.2	392	19.8	17.8-21.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Prostate

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	3142	145.9	140.8-151.1	377	19.0	17.1-21.0
2009	3150	144.5	139.4-149.7	393	19.7	17.8-21.7
2010	2898	131.1	126.3-136.1	426	21.0	19.0-23.1
2011	3076	136.5	131.6-141.4	391	18.7	16.9-20.7
2012	2626	113.6	109.3-118.1	339	16.0	14.4-17.8
2013	2542	108.5	104.3-112.9	327	15.2	13.6-17.0
2014	2422	100.6	96.5-104.7	364	16.6	15.0-18.4
2015	2394	97.8	93.8-101.8	358	15.7	14.1-17.5
2016	2411	97.3	93.4-101.4	427	18.6	16.9-20.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Testis

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	76	2.8	2.2-3.6	16	0.6	0.3-1.0
1977	72	2.8	2.1-3.5	16	0.6	0.3-1.0
1978	70	2.8	2.2-3.6	15	0.6	0.3-1.0
1979	80	3.4	2.6-4.2	16	0.6	0.3-1.0
1980	97	4.1	3.3-5.0	16	0.7	0.4-1.1
1981	99	3.9	3.1-4.8	4	0.1	0.0-0.4
1982	108	4.3	3.5-5.3	7	0.3	0.1-0.6
1983	101	3.9	3.2-4.8	7	0.3	0.1-0.6
1984	94	3.9	3.1-4.8	10	0.4	0.2-0.8
1985	96	3.8	3.1-4.7	14	0.5	0.3-0.9
1986	108	4.3	3.5-5.2	9	0.4	0.2-0.8
1987	106	4.2	3.4-5.2	7	0.3	0.1-0.6
1988	89	3.6	2.8-4.4	7	0.2	0.1-0.5
1989	103	4.2	3.4-5.2	10	0.4	0.2-0.8
1990	117	4.8	4.0-5.8	8	0.3	0.1-0.7
1991	121	4.9	4.1-5.9	9	0.4	0.2-0.8
1992	114	4.9	4.0-5.9	9	0.4	0.2-0.8
1993	117	4.8	3.9-5.8	9	0.4	0.2-0.8
1994	115	4.7	3.9-5.7	9	0.4	0.2-0.8
1995	107	4.4	3.6-5.4	7	0.3	0.1-0.7
1996	108	4.4	3.6-5.3	5	0.2	0.1-0.5
1997	102	4.2	3.4-5.1	6	0.2	0.1-0.5
1998	99	4.1	3.3-5.0	7	0.3	0.1-0.6
1999	146	5.9	4.9-6.9	4	0.2	0.1-0.5
2000	128	5.2	4.3-6.2	2	0.1	0.0-0.3
2001	118	4.7	3.9-5.7	6	0.3	0.1-0.6
2002	127	5.2	4.3-6.2	3	0.1	0.0-0.4
2003	139	5.8	4.8-6.8	8	0.4	0.2-0.7
2004	145	6.0	5.0-7.0	8	0.3	0.1-0.7
2005	150	6.1	5.2-7.2	7	0.3	0.1-0.6
2006	150	6.1	5.2-7.2	11	0.5	0.2-0.9
2007	161	6.7	5.7-7.8	1	0.0	0.0-0.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Testis

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	148	6.3	5.3-7.4	5	0.2	0.1-0.5
2009	170	7.1	6.1-8.3	8	0.3	0.1-0.7
2010	161	6.8	5.7-7.9	14	0.6	0.3-1.0
2011	166	6.8	5.8-8.0	7	0.3	0.1-0.6
2012	171	7.1	6.0-8.2	7	0.3	0.1-0.6
2013	157	6.4	5.4-7.5	6	0.2	0.1-0.5
2014	152	6.2	5.3-7.3	2	0.1	0.0-0.3
2015	160	6.4	5.4-7.5	5	0.2	0.1-0.5
2016	150	6.1	5.1-7.2	7	0.3	0.1-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Urinary bladder (incl. in situ)

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1176	18.7	17.6-19.9	842	33.8	31.4-36.3	334	9.0	8.0-10.0	365	6.3	5.7-7.0	249	11.2	9.7-12.7	116	3.4	2.8-4.1
1977	1263	20.4	19.3-21.6	891	36.2	33.7-38.7	372	10.3	9.2-11.4	384	6.5	5.9-7.2	267	11.9	10.4-13.4	117	3.3	2.7-4.0
1978	1313	21.7	20.5-23.0	946	39.7	37.1-42.5	367	10.0	9.0-11.1	353	6.0	5.4-6.7	234	10.5	9.1-12.0	119	3.4	2.8-4.1
1979	1325	22.2	21.0-23.5	924	39.6	37.0-42.3	401	11.3	10.2-12.5	346	5.9	5.3-6.6	242	11.1	9.7-12.6	104	2.9	2.3-3.5
1980	1388	23.3	22.1-24.6	1035	44.5	41.8-47.4	353	9.8	8.8-10.9	358	6.1	5.5-6.8	249	11.3	9.9-12.9	109	3.0	2.5-3.7
1981	1355	23.1	21.8-24.4	987	42.9	40.2-45.8	368	10.4	9.4-11.6	301	5.2	4.6-5.8	202	9.3	8.0-10.7	99	2.7	2.2-3.3
1982	1252	21.5	20.3-22.7	887	38.8	36.2-41.5	365	10.5	9.4-11.7	317	5.4	4.8-6.1	226	10.5	9.1-12.0	91	2.5	2.0-3.1
1983	1387	23.9	22.6-25.2	976	43.8	41.0-46.7	411	11.6	10.5-12.8	284	4.9	4.3-5.5	189	9.0	7.7-10.4	95	2.6	2.1-3.2
1984	1259	21.6	20.4-22.8	910	39.8	37.2-42.5	349	9.9	8.9-11.1	282	4.9	4.3-5.5	186	8.7	7.4-10.1	96	2.7	2.2-3.3
1985	1131	19.9	18.7-21.1	800	36.2	33.7-38.9	331	9.5	8.5-10.6	292	5.1	4.5-5.8	201	9.8	8.5-11.3	91	2.5	2.0-3.1
1986	1208	21.6	20.3-22.8	854	39.2	36.6-42.0	354	10.6	9.5-11.8	260	4.5	4.0-5.1	164	7.9	6.7-9.3	96	2.6	2.1-3.2
1987	1231	21.8	20.5-23.0	877	39.9	37.3-42.7	354	10.4	9.3-11.6	243	4.3	3.8-4.9	172	8.4	7.2-9.8	71	2.1	1.6-2.6
1988	1188	21.3	20.0-22.5	830	38.1	35.5-40.8	358	10.4	9.4-11.6	271	4.7	4.2-5.4	169	8.1	6.9-9.5	102	2.8	2.3-3.4
1989	1112	20.6	19.4-21.9	789	37.6	34.9-40.3	323	9.9	8.8-11.1	277	5.0	4.4-5.7	175	8.8	7.5-10.3	102	2.9	2.4-3.6
1990	1142	21.3	20.1-22.6	792	38.1	35.5-40.9	350	10.6	9.5-11.8	249	4.6	4.0-5.2	159	8.1	6.8-9.4	90	2.7	2.1-3.3
1991	1206	22.9	21.6-24.3	862	41.9	39.2-44.9	344	10.8	9.6-12.0	226	4.2	3.7-4.8	148	7.5	6.4-8.9	78	2.3	1.8-2.8
1992	1183	22.6	21.4-24.0	832	40.8	38.1-43.8	351	11.3	10.1-12.6	233	4.3	3.8-4.9	152	7.9	6.7-9.2	81	2.4	1.9-3.0
1993	1095	21.2	20.0-22.5	782	38.5	35.8-41.3	313	10.1	9.0-11.4	273	5.2	4.6-5.8	187	9.6	8.3-11.1	86	2.6	2.1-3.3
1994	1209	23.5	22.2-24.9	838	41.4	38.7-44.4	371	12.0	10.8-13.4	258	4.8	4.3-5.5	175	9.1	7.8-10.6	83	2.5	2.0-3.1
1995	1177	23.1	21.8-24.5	844	42.2	39.4-45.1	333	10.8	9.7-12.1	288	5.5	4.9-6.2	185	9.6	8.3-11.1	103	3.0	2.5-3.7
1996	1033	20.5	19.2-21.7	739	37.3	34.6-40.1	294	9.5	8.4-10.7	239	4.6	4.0-5.2	146	7.7	6.5-9.1	93	2.8	2.2-3.4
1997	1184	23.7	22.3-25.1	849	42.4	39.6-45.3	335	11.0	9.8-12.3	233	4.5	3.9-5.1	154	8.1	6.9-9.5	79	2.5	2.0-3.1
1998	1221	24.3	23.0-25.7	849	42.1	39.3-45.1	372	12.2	10.9-13.5	253	4.9	4.3-5.5	160	8.2	7.0-9.6	93	2.8	2.2-3.4
1999	1116	22.0	20.8-23.4	784	38.8	36.1-41.6	332	11.0	9.8-12.2	240	4.5	4.0-5.2	166	8.6	7.3-10.0	74	2.2	1.7-2.8
2000	1193	23.4	22.1-24.8	836	41.1	38.4-44.0	357	11.8	10.6-13.1	228	4.3	3.8-4.9	154	7.8	6.6-9.1	74	2.2	1.7-2.8
2001	1126	22.1	20.8-23.4	805	39.3	36.6-42.1	321	10.6	9.5-11.9	234	4.4	3.9-5.1	165	8.3	7.0-9.6	69	2.1	1.6-2.7
2002	1076	21.0	19.8-22.3	777	37.9	35.3-40.7	299	9.8	8.7-11.0	208	3.9	3.4-4.4	136	6.8	5.7-8.1	72	2.1	1.6-2.7
2003	1150	22.5	21.2-23.9	830	40.4	37.7-43.3	320	10.4	9.3-11.7	192	3.6	3.1-4.1	131	6.6	5.5-7.8	61	1.8	1.4-2.4
2004	1178	23.0	21.7-24.4	837	40.3	37.6-43.2	341	11.0	9.9-12.3	225	4.2	3.7-4.8	157	7.7	6.6-9.0	68	2.0	1.6-2.6
2005	1155	22.8	21.5-24.2	811	39.1	36.5-41.9	344	11.4	10.2-12.8	225	4.3	3.7-4.9	143	7.2	6.0-8.4	82	2.5	1.9-3.1
2006	1120	22.2	20.9-23.5	835	40.6	37.9-43.5	285	9.6	8.5-10.8	240	4.6	4.0-5.2	156	7.9	6.7-9.2	84	2.5	2.0-3.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Urinary bladder (incl. in situ)

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	1178	23.5	22.1-24.9	857	41.9	39.1-44.8	321	10.8	9.6-12.1	242	4.6	4.0-5.2	161	8.0	6.8-9.3	81	2.4	1.9-3.1
2008	1163	23.1	21.8-24.5	886	42.5	39.8-45.5	277	9.4	8.3-10.7	227	4.4	3.8-5.0	159	7.9	6.7-9.2	68	2.2	1.7-2.8
2009	1139	22.5	21.2-23.9	822	39.6	36.9-42.4	317	10.6	9.4-11.9	215	4.0	3.5-4.6	143	7.1	5.9-8.3	72	2.1	1.6-2.7
2010	1089	21.4	20.1-22.7	831	39.6	36.9-42.4	258	8.7	7.7-9.9	241	4.5	3.9-5.1	170	8.1	6.9-9.4	71	2.1	1.6-2.7
2011	1091	21.2	19.9-22.5	803	37.6	35.0-40.3	288	9.6	8.5-10.8	221	4.1	3.6-4.7	150	7.1	6.0-8.4	71	2.2	1.7-2.8
2012	1113	21.3	20.0-22.6	819	37.7	35.2-40.4	294	9.6	8.5-10.8	234	4.2	3.7-4.8	159	7.3	6.2-8.6	75	2.2	1.7-2.8
2013	1083	20.5	19.3-21.8	798	35.9	33.5-38.5	285	9.3	8.2-10.5	221	3.9	3.4-4.4	156	7.1	6.1-8.4	65	1.8	1.4-2.3
2014	1103	20.4	19.2-21.6	810	35.9	33.5-38.5	293	9.2	8.2-10.4	243	4.2	3.7-4.8	170	7.7	6.6-8.9	73	2.0	1.6-2.6
2015	1203	22.1	20.8-23.4	932	40.6	38.0-43.3	271	8.8	7.7-9.9	230	4.0	3.5-4.6	154	6.8	5.7-8.0	76	2.2	1.7-2.7
2016	1076	19.5	18.3-20.7	824	35.4	33.0-38.0	252	8.0	7.0-9.1	260	4.5	4.0-5.1	178	7.8	6.7-9.1	82	2.4	1.9-3.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Kidney and renal pelvis

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	463	7.5	6.8-8.2	286	10.9	9.6-12.3	177	5.1	4.3-5.9	241	3.9	3.4-4.4	158	6.4	5.4-7.5	83	2.2	1.8-2.8
1977	457	7.6	6.9-8.3	295	11.7	10.3-13.1	162	4.7	4.0-5.6	233	3.8	3.3-4.4	150	6.1	5.1-7.2	83	2.2	1.8-2.8
1978	448	7.4	6.7-8.2	272	11.0	9.7-12.5	176	4.9	4.2-5.7	221	3.7	3.2-4.2	144	5.7	4.8-6.8	77	2.3	1.8-2.9
1979	432	7.3	6.6-8.1	268	11.1	9.8-12.6	164	4.6	3.9-5.3	227	3.9	3.4-4.5	143	6.1	5.1-7.3	84	2.4	1.9-3.0
1980	413	7.0	6.3-7.7	258	10.4	9.1-11.7	155	4.7	4.0-5.6	192	3.2	2.8-3.7	117	5.0	4.1-6.0	75	2.1	1.6-2.6
1981	417	7.2	6.5-8.0	240	10.2	9.0-11.7	177	5.2	4.4-6.1	208	3.5	3.1-4.1	120	5.0	4.1-6.0	88	2.5	2.0-3.1
1982	461	8.3	7.5-9.1	273	12.0	10.6-13.5	188	5.7	4.9-6.6	209	3.6	3.1-4.2	124	5.5	4.6-6.6	85	2.4	1.9-3.0
1983	453	8.1	7.3-8.9	281	12.1	10.7-13.6	172	5.3	4.5-6.3	190	3.3	2.8-3.8	116	5.0	4.1-6.1	74	2.1	1.7-2.7
1984	480	8.6	7.8-9.5	300	13.1	11.6-14.7	180	5.5	4.7-6.4	199	3.5	3.1-4.1	122	5.4	4.5-6.5	77	2.3	1.8-2.9
1985	517	9.3	8.5-10.2	314	13.7	12.2-15.3	203	6.3	5.4-7.3	182	3.2	2.8-3.7	104	4.6	3.7-5.6	78	2.4	1.9-3.0
1986	493	8.8	8.0-9.6	297	13.0	11.5-14.6	196	5.8	5.0-6.7	205	3.6	3.1-4.1	126	5.6	4.6-6.7	79	2.2	1.8-2.8
1987	547	10.0	9.1-10.9	326	14.4	12.8-16.1	221	6.9	6.0-7.9	189	3.4	2.9-4.0	117	5.4	4.4-6.5	72	2.2	1.7-2.7
1988	454	8.6	7.8-9.5	283	13.0	11.5-14.6	171	5.7	4.8-6.6	176	3.3	2.8-3.8	106	4.9	4.0-5.9	70	2.2	1.7-2.9
1989	484	9.3	8.5-10.2	297	13.8	12.3-15.5	187	6.1	5.2-7.0	193	3.6	3.1-4.1	114	5.4	4.5-6.5	79	2.4	1.9-3.0
1990	529	10.2	9.4-11.2	299	14.0	12.4-15.7	230	7.8	6.8-8.9	185	3.5	3.0-4.1	102	4.9	4.0-6.0	83	2.6	2.0-3.2
1991	534	10.5	9.6-11.4	304	14.4	12.8-16.1	230	7.7	6.7-8.8	162	3.1	2.6-3.6	91	4.3	3.4-5.3	71	2.3	1.8-2.9
1992	530	10.6	9.7-11.5	314	15.1	13.5-16.9	216	7.3	6.4-8.4	152	2.9	2.5-3.4	86	4.3	3.4-5.3	66	2.1	1.6-2.7
1993	546	10.9	10.0-11.9	354	17.0	15.2-18.8	192	6.6	5.7-7.6	169	3.3	2.8-3.9	116	5.6	4.6-6.7	53	1.7	1.3-2.3
1994	592	12.1	11.1-13.1	364	17.9	16.1-19.8	228	7.8	6.8-8.9	195	3.8	3.3-4.4	120	5.9	4.9-7.0	75	2.4	1.8-3.0
1995	535	10.9	10.0-11.9	321	15.6	13.9-17.4	214	7.6	6.6-8.7	183	3.7	3.1-4.2	113	5.7	4.7-6.8	70	2.3	1.8-2.9
1996	625	12.8	11.8-13.8	368	18.2	16.3-20.1	257	9.0	7.9-10.2	176	3.5	3.0-4.1	98	4.9	4.0-6.0	78	2.5	1.9-3.1
1997	604	12.5	11.5-13.5	353	17.5	15.7-19.4	251	9.1	8.0-10.3	184	3.7	3.2-4.3	109	5.5	4.5-6.6	75	2.5	2.0-3.2
1998	644	13.3	12.3-14.4	390	19.0	17.1-20.9	254	9.2	8.1-10.4	174	3.5	3.0-4.0	95	4.8	3.9-5.8	79	2.7	2.1-3.4
1999	660	13.5	12.5-14.6	415	19.9	18.1-22.0	245	8.6	7.5-9.8	164	3.3	2.8-3.8	106	5.2	4.2-6.3	58	1.9	1.4-2.4
2000	683	13.9	12.9-15.0	401	19.2	17.3-21.2	282	10.0	8.8-11.2	158	3.1	2.7-3.7	101	4.9	4.0-6.0	57	1.9	1.4-2.4
2001	671	13.7	12.7-14.8	416	19.7	17.9-21.7	255	9.0	7.9-10.2	164	3.3	2.8-3.8	96	4.6	3.7-5.6	68	2.3	1.8-2.9
2002	677	13.8	12.8-14.9	421	19.9	18.0-21.8	256	8.9	7.8-10.1	170	3.3	2.8-3.9	106	5.1	4.2-6.2	64	2.1	1.6-2.7
2003	741	15.1	14.0-16.2	477	22.4	20.4-24.5	264	9.3	8.2-10.5	149	2.9	2.5-3.5	91	4.4	3.5-5.4	58	1.8	1.4-2.4
2004	753	15.4	14.3-16.5	472	22.2	20.2-24.3	281	10.2	9.0-11.5	163	3.2	2.7-3.8	100	4.8	3.9-5.9	63	2.1	1.6-2.7
2005	746	15.3	14.2-16.4	461	21.8	19.8-23.9	285	10.2	9.1-11.5	133	2.6	2.2-3.1	77	3.7	2.9-4.6	56	1.8	1.3-2.4
2006	796	16.5	15.3-17.7	507	23.9	21.9-26.1	289	10.7	9.5-12.0	160	3.2	2.7-3.7	93	4.6	3.7-5.6	67	2.2	1.7-2.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Kidney and renal pelvis

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	762	15.8	14.7-17.0	479	22.8	20.8-25.0	283	10.3	9.1-11.6	163	3.2	2.7-3.7	101	4.8	3.9-5.8	62	2.0	1.5-2.6
2008	810	16.5	15.4-17.7	521	24.3	22.2-26.5	289	10.4	9.2-11.7	145	2.8	2.4-3.3	87	4.2	3.3-5.2	58	1.9	1.4-2.4
2009	721	14.8	13.8-16.0	435	20.3	18.4-22.3	286	10.5	9.3-11.8	181	3.6	3.1-4.2	125	6.0	5.0-7.1	56	1.9	1.4-2.5
2010	746	15.2	14.1-16.4	473	21.5	19.5-23.5	273	10.2	9.0-11.5	156	3.0	2.5-3.5	98	4.5	3.6-5.5	58	1.9	1.4-2.5
2011	748	15.2	14.1-16.3	467	21.0	19.1-23.0	281	10.2	9.0-11.5	138	2.6	2.1-3.0	92	4.1	3.3-5.1	46	1.4	1.0-1.9
2012	767	15.4	14.3-16.6	497	22.2	20.2-24.2	270	9.9	8.8-11.2	136	2.6	2.2-3.1	91	4.2	3.4-5.2	45	1.4	1.0-2.0
2013	786	15.3	14.3-16.5	507	22.1	20.2-24.1	279	9.8	8.7-11.1	135	2.5	2.1-3.0	75	3.4	2.7-4.3	60	1.8	1.3-2.3
2014	762	14.7	13.6-15.8	500	21.2	19.4-23.2	262	9.3	8.2-10.5	153	2.8	2.4-3.3	91	4.0	3.2-4.9	62	1.9	1.5-2.5
2015	789	15.0	13.9-16.1	498	20.7	18.9-22.7	291	10.0	8.8-11.3	179	3.3	2.8-3.8	118	5.1	4.2-6.1	61	1.8	1.4-2.4
2016	759	14.4	13.4-15.5	480	20.1	18.3-22.0	279	9.9	8.7-11.1	146	2.6	2.2-3.1	102	4.3	3.5-5.2	44	1.4	1.0-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Brain and other central nervous system

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	351	5.8	5.2-6.5	188	7.0	6.0-8.1	163	4.9	4.1-5.7	273	4.6	4.0-5.2	152	5.9	5.0-6.9	121	3.7	3.0-4.4
1977	347	6.1	5.4-6.8	191	7.5	6.4-8.7	156	5.0	4.2-5.9	264	4.4	3.9-5.0	137	5.2	4.3-6.1	127	3.9	3.2-4.6
1978	350	6.2	5.6-7.0	175	7.0	6.0-8.2	175	5.6	4.8-6.5	262	4.4	3.9-5.0	135	5.4	4.5-6.4	127	3.6	3.0-4.4
1979	335	6.0	5.3-6.7	177	7.3	6.2-8.5	158	5.0	4.2-6.0	242	4.1	3.6-4.7	134	5.5	4.6-6.5	108	3.1	2.6-3.8
1980	333	6.0	5.3-6.7	166	6.9	5.9-8.1	167	5.3	4.5-6.2	268	4.5	4.0-5.1	144	5.7	4.8-6.8	124	3.7	3.0-4.4
1981	347	6.5	5.8-7.3	194	8.5	7.3-9.8	153	4.9	4.1-5.8	250	4.4	3.9-5.1	130	5.5	4.5-6.5	120	3.7	3.0-4.4
1982	342	6.3	5.6-7.0	185	7.9	6.8-9.2	157	5.1	4.3-6.0	255	4.6	4.0-5.2	125	5.3	4.4-6.4	130	4.0	3.3-4.8
1983	318	6.1	5.5-6.9	166	7.2	6.1-8.4	152	5.2	4.4-6.2	232	4.2	3.7-4.8	127	5.6	4.7-6.7	105	3.1	2.6-3.8
1984	344	6.6	5.9-7.3	161	7.0	5.9-8.2	183	6.3	5.4-7.3	232	4.1	3.6-4.7	115	4.9	4.0-5.9	117	3.7	3.0-4.4
1985	356	6.9	6.2-7.7	194	8.7	7.5-10.1	162	5.5	4.6-6.4	239	4.4	3.9-5.0	119	5.3	4.4-6.4	120	3.7	3.1-4.5
1986	333	6.6	5.9-7.3	188	8.6	7.4-9.9	145	5.2	4.3-6.1	244	4.6	4.0-5.2	135	6.3	5.3-7.5	109	3.3	2.7-4.0
1987	360	7.1	6.4-7.9	181	8.2	7.0-9.5	179	6.0	5.2-7.1	230	4.4	3.9-5.1	117	5.3	4.4-6.4	113	3.8	3.1-4.6
1988	367	7.3	6.5-8.1	190	8.7	7.5-10.1	177	6.2	5.3-7.3	169	3.2	2.7-3.7	90	4.0	3.2-5.0	79	2.5	2.0-3.2
1989	374	7.7	6.9-8.5	202	9.6	8.3-11.0	172	6.2	5.2-7.2	187	3.6	3.1-4.2	98	4.6	3.7-5.6	89	2.8	2.2-3.4
1990	373	7.8	7.0-8.7	191	9.1	7.9-10.6	182	6.8	5.8-7.9	194	3.9	3.4-4.5	106	5.1	4.2-6.2	88	3.0	2.4-3.8
1991	360	7.6	6.8-8.4	181	8.7	7.5-10.1	179	6.6	5.7-7.7	210	4.3	3.7-5.0	105	5.0	4.1-6.1	105	3.8	3.1-4.6
1992	350	7.4	6.6-8.2	191	9.0	7.7-10.3	159	6.0	5.1-7.1	173	3.5	3.0-4.1	91	4.5	3.6-5.5	82	2.8	2.2-3.5
1993	362	7.5	6.7-8.3	178	8.5	7.3-9.8	184	6.8	5.9-7.9	172	3.5	3.0-4.0	95	4.6	3.7-5.6	77	2.7	2.1-3.4
1994	386	8.2	7.4-9.1	194	9.4	8.1-10.8	192	7.2	6.2-8.3	184	3.8	3.3-4.4	94	4.6	3.7-5.6	90	3.2	2.6-4.0
1995	415	8.7	7.9-9.6	226	10.7	9.3-12.2	189	7.0	6.0-8.2	199	4.1	3.5-4.7	106	5.2	4.2-6.2	93	3.3	2.6-4.0
1996	413	8.8	8.0-9.7	211	10.1	8.7-11.5	202	7.7	6.6-8.8	187	3.9	3.4-4.6	94	4.6	3.7-5.7	93	3.3	2.7-4.1
1997	374	8.0	7.2-8.9	181	8.7	7.5-10.1	193	7.5	6.5-8.7	209	4.4	3.8-5.0	90	4.5	3.6-5.5	119	4.3	3.6-5.2
1998	430	9.2	8.3-10.1	229	10.8	9.4-12.3	201	8.0	6.9-9.2	215	4.5	3.9-5.1	104	5.1	4.1-6.1	111	4.0	3.3-4.8
1999	351	7.4	6.7-8.3	181	8.5	7.3-9.8	170	6.4	5.5-7.5	213	4.4	3.8-5.0	111	5.3	4.4-6.4	102	3.6	2.9-4.4
2000	371	7.8	7.0-8.6	178	8.4	7.2-9.7	193	7.3	6.2-8.4	185	3.9	3.3-4.5	93	4.4	3.6-5.4	92	3.4	2.8-4.3
2001	340	7.1	6.4-7.9	171	8.0	6.9-9.3	169	6.4	5.5-7.5	172	3.5	3.0-4.1	95	4.5	3.6-5.5	77	2.7	2.1-3.4
2002	353	7.4	6.6-8.2	180	8.4	7.2-9.7	173	6.6	5.7-7.7	165	3.4	2.9-4.0	87	4.1	3.3-5.1	78	2.9	2.3-3.6
2003	330	6.9	6.1-7.7	169	7.8	6.6-9.0	161	6.1	5.1-7.1	162	3.3	2.8-3.9	95	4.4	3.6-5.4	67	2.4	1.8-3.1
2004	332	7.1	6.3-7.9	186	8.8	7.6-10.2	146	5.7	4.8-6.7	147	3.0	2.5-3.6	76	3.6	2.8-4.5	71	2.6	2.0-3.3
2005	353	7.5	6.7-8.3	199	9.3	8.1-10.7	154	5.9	5.0-7.0	191	3.9	3.4-4.5	107	5.1	4.2-6.1	84	3.0	2.4-3.8
2006	379	8.1	7.3-9.0	199	9.4	8.1-10.8	180	7.1	6.1-8.3	188	3.9	3.4-4.5	107	5.1	4.2-6.1	81	2.9	2.3-3.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Brain and other central nervous system

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	317	6.8	6.1-7.7	165	7.8	6.7-9.1	152	6.1	5.1-7.1	205	4.2	3.7-4.8	109	5.1	4.2-6.2	96	3.5	2.8-4.3
2008	344	7.4	6.6-8.2	172	8.1	6.9-9.4	172	6.7	5.8-7.9	157	3.2	2.7-3.8	86	4.0	3.2-5.0	71	2.6	2.0-3.3
2009	348	7.2	6.5-8.0	183	8.4	7.2-9.7	165	6.2	5.3-7.3	195	4.0	3.5-4.7	106	5.0	4.1-6.1	89	3.4	2.7-4.2
2010	315	6.8	6.0-7.6	171	8.0	6.8-9.3	144	5.9	4.9-7.0	185	3.7	3.2-4.3	91	4.1	3.3-5.1	94	3.4	2.7-4.2
2011	308	6.4	5.7-7.2	164	7.5	6.4-8.8	144	5.6	4.7-6.6	192	4.0	3.4-4.6	102	4.8	3.9-5.8	90	3.3	2.6-4.1
2012	328	6.8	6.0-7.5	166	7.5	6.3-8.7	162	6.1	5.2-7.2	183	3.6	3.1-4.1	81	3.6	2.8-4.4	102	3.6	2.9-4.4
2013	321	6.7	5.9-7.4	172	7.7	6.6-9.0	149	5.7	4.8-6.7	170	3.3	2.8-3.8	85	3.9	3.1-4.8	85	2.8	2.2-3.5
2014	366	7.3	6.6-8.2	208	9.0	7.8-10.4	158	6.0	5.1-7.0	210	4.0	3.5-4.6	108	4.6	3.8-5.6	102	3.6	2.9-4.3
2015	355	7.1	6.4-8.0	182	7.9	6.8-9.2	173	6.4	5.5-7.5	193	3.7	3.1-4.2	96	4.1	3.3-5.0	97	3.3	2.7-4.1
2016	349	7.0	6.3-7.8	191	8.3	7.1-9.6	158	6.0	5.0-7.0	216	4.0	3.5-4.6	118	5.1	4.2-6.1	98	3.2	2.6-4.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Thyroid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	193	3.4	3.0-4.0	55	2.3	1.7-3.0	138	4.4	3.7-5.3	50	0.8	0.6-1.1	21	0.8	0.5-1.3	29	0.8	0.6-1.2
1977	223	3.9	3.4-4.5	62	2.4	1.8-3.1	161	5.2	4.4-6.2	47	0.8	0.6-1.0	25	0.9	0.6-1.4	22	0.6	0.4-0.9
1978	231	4.3	3.8-4.9	63	2.6	2.0-3.4	168	5.7	4.9-6.7	37	0.7	0.5-0.9	11	0.5	0.2-0.9	26	0.8	0.5-1.2
1979	203	3.9	3.3-4.5	79	3.4	2.6-4.2	124	4.4	3.6-5.3	33	0.6	0.4-0.8	15	0.6	0.4-1.1	18	0.5	0.3-0.8
1980	204	4.0	3.4-4.6	57	2.6	1.9-3.4	147	5.1	4.3-6.1	40	0.7	0.5-1.0	15	0.7	0.4-1.1	25	0.7	0.4-1.1
1981	191	3.6	3.1-4.2	56	2.4	1.8-3.1	135	4.7	3.9-5.7	38	0.7	0.5-0.9	12	0.5	0.2-0.9	26	0.8	0.5-1.1
1982	198	3.9	3.4-4.5	61	2.8	2.1-3.6	137	4.9	4.0-5.8	46	0.8	0.6-1.0	15	0.6	0.3-1.0	31	0.8	0.6-1.2
1983	232	4.7	4.1-5.4	59	2.6	1.9-3.3	173	6.6	5.6-7.7	27	0.5	0.3-0.7	7	0.3	0.1-0.6	20	0.6	0.4-1.0
1984	260	5.2	4.5-5.9	74	3.4	2.6-4.3	186	6.8	5.8-7.9	37	0.6	0.4-0.9	13	0.6	0.3-1.0	24	0.7	0.4-1.0
1985	264	5.2	4.5-5.8	79	3.4	2.7-4.3	185	6.5	5.6-7.6	44	0.8	0.6-1.1	25	1.2	0.7-1.7	19	0.6	0.3-0.9
1986	227	4.4	3.9-5.1	57	2.5	1.9-3.2	170	6.2	5.3-7.3	29	0.5	0.4-0.8	4	0.2	0.0-0.5	25	0.8	0.5-1.1
1987	208	4.2	3.7-4.9	64	3.0	2.3-3.8	144	5.4	4.5-6.4	30	0.5	0.4-0.8	10	0.5	0.2-0.9	20	0.5	0.3-0.9
1988	214	4.3	3.7-4.9	71	3.2	2.5-4.1	143	5.3	4.4-6.3	30	0.5	0.4-0.8	11	0.5	0.2-0.9	19	0.6	0.3-0.9
1989	188	4.0	3.4-4.6	51	2.3	1.7-3.0	137	5.6	4.6-6.6	28	0.5	0.3-0.8	7	0.3	0.1-0.7	21	0.6	0.4-1.0
1990	226	4.7	4.1-5.4	56	2.6	2.0-3.4	170	6.6	5.6-7.7	31	0.6	0.4-0.8	8	0.4	0.2-0.8	23	0.7	0.4-1.1
1991	243	5.2	4.5-5.9	65	3.0	2.3-3.9	178	7.2	6.1-8.4	30	0.6	0.4-0.8	8	0.4	0.2-0.8	22	0.7	0.4-1.1
1992	259	5.5	4.8-6.2	64	3.1	2.4-4.0	195	7.7	6.6-8.9	26	0.5	0.3-0.8	10	0.5	0.3-1.0	16	0.5	0.3-0.8
1993	280	6.0	5.3-6.7	77	3.7	2.9-4.6	203	8.2	7.1-9.4	26	0.5	0.3-0.7	9	0.4	0.2-0.8	17	0.5	0.3-0.8
1994	290	6.2	5.5-7.0	82	3.9	3.1-4.8	208	8.4	7.2-9.6	31	0.6	0.4-0.9	12	0.6	0.3-1.1	19	0.6	0.4-1.0
1995	312	6.5	5.8-7.3	91	4.2	3.4-5.2	221	8.7	7.5-9.9	23	0.4	0.3-0.7	7	0.3	0.1-0.7	16	0.5	0.3-0.8
1996	306	6.6	5.8-7.4	71	3.4	2.7-4.3	235	9.5	8.3-10.9	31	0.6	0.4-0.9	11	0.5	0.3-1.0	20	0.6	0.4-1.0
1997	347	7.3	6.6-8.1	93	4.3	3.5-5.3	254	10.1	8.9-11.4	30	0.6	0.4-0.9	9	0.4	0.2-0.8	21	0.7	0.4-1.1
1998	375	7.9	7.1-8.7	105	4.9	4.0-5.9	270	10.6	9.4-12.0	23	0.5	0.3-0.7	8	0.4	0.2-0.8	15	0.5	0.3-0.8
1999	413	8.6	7.8-9.5	97	4.3	3.5-5.3	316	12.6	11.2-14.1	22	0.4	0.3-0.7	10	0.5	0.2-0.9	12	0.4	0.2-0.7
2000	402	8.4	7.6-9.3	102	4.6	3.7-5.5	300	12.0	10.6-13.4	32	0.6	0.4-0.9	8	0.4	0.2-0.8	24	0.8	0.5-1.2
2001	471	9.9	9.0-10.8	123	5.7	4.7-6.8	348	13.8	12.4-15.4	26	0.5	0.3-0.7	10	0.5	0.2-0.9	16	0.5	0.3-0.8
2002	487	10.3	9.4-11.2	143	6.4	5.4-7.6	344	13.9	12.5-15.5	33	0.6	0.4-0.9	11	0.5	0.3-0.9	22	0.7	0.4-1.1
2003	462	9.7	8.8-10.6	109	4.9	4.0-5.9	353	14.1	12.6-15.6	23	0.4	0.3-0.7	9	0.4	0.2-0.8	14	0.4	0.2-0.7
2004	532	11.2	10.2-12.2	136	6.2	5.2-7.4	396	15.9	14.3-17.5	21	0.4	0.2-0.6	10	0.5	0.2-0.9	11	0.3	0.2-0.6
2005	719	15.2	14.1-16.4	182	8.3	7.1-9.6	537	21.9	20.1-23.9	37	0.7	0.5-1.0	16	0.8	0.4-1.3	21	0.7	0.4-1.1
2006	697	14.8	13.7-15.9	173	7.7	6.6-8.9	524	21.5	19.6-23.4	41	0.8	0.6-1.1	13	0.7	0.4-1.1	28	1.0	0.6-1.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Thyroid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	782	16.8	15.6-18.0	181	8.2	7.0-9.5	601	24.9	22.9-27.0	39	0.8	0.6-1.1	11	0.5	0.3-1.0	28	1.0	0.6-1.4
2008	936	19.7	18.5-21.0	225	10.0	8.7-11.4	711	29.1	26.9-31.4	26	0.5	0.3-0.8	14	0.7	0.4-1.2	12	0.4	0.2-0.7
2009	1063	22.4	21.0-23.8	276	12.1	10.7-13.7	787	32.1	29.9-34.5	31	0.6	0.4-0.9	17	0.8	0.5-1.3	14	0.5	0.3-0.8
2010	1014	21.4	20.1-22.8	246	11.0	9.6-12.5	768	31.5	29.3-33.9	31	0.6	0.4-0.8	14	0.6	0.3-1.1	17	0.5	0.3-0.9
2011	1002	21.1	19.8-22.4	214	9.5	8.2-10.9	788	32.0	29.8-34.4	27	0.5	0.3-0.8	10	0.5	0.2-0.9	17	0.5	0.3-0.9
2012	1097	22.8	21.5-24.2	276	11.9	10.5-13.4	821	33.4	31.1-35.8	26	0.5	0.3-0.8	13	0.6	0.3-1.0	13	0.5	0.2-0.8
2013	1186	24.8	23.3-26.2	318	13.8	12.3-15.4	868	35.6	33.2-38.1	38	0.7	0.5-1.0	18	0.8	0.5-1.3	20	0.6	0.4-1.0
2014	1130	23.1	21.8-24.6	288	12.4	11.0-13.9	842	33.7	31.4-36.2	34	0.6	0.4-0.9	12	0.5	0.3-0.9	22	0.7	0.4-1.1
2015	1074	22.0	20.7-23.4	283	11.9	10.6-13.4	791	32.0	29.7-34.3	29	0.5	0.3-0.7	12	0.5	0.2-0.8	17	0.5	0.3-0.8
2016	1057	21.6	20.3-23.0	254	10.6	9.3-12.1	803	32.4	30.1-34.8	38	0.7	0.5-1.0	15	0.7	0.4-1.1	23	0.7	0.5-1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Hodgkin lymphoma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	217	3.7	3.2-4.2	116	4.3	3.6-5.2	101	3.1	2.5-3.8	78	1.4	1.1-1.8	51	2.0	1.5-2.7	27	0.8	0.5-1.3
1977	208	3.6	3.1-4.2	119	4.5	3.7-5.4	89	3.0	2.3-3.7	88	1.6	1.3-2.0	49	1.9	1.4-2.5	39	1.3	0.9-1.8
1978	213	3.7	3.2-4.2	112	4.2	3.4-5.1	101	3.3	2.6-4.0	78	1.4	1.1-1.7	43	1.7	1.2-2.3	35	1.1	0.8-1.6
1979	199	3.8	3.3-4.4	101	4.3	3.5-5.3	98	3.4	2.7-4.1	62	1.1	0.9-1.5	37	1.5	1.1-2.1	25	0.9	0.5-1.3
1980	199	3.7	3.2-4.3	103	4.4	3.5-5.3	96	3.2	2.5-3.9	72	1.3	1.0-1.6	35	1.6	1.1-2.2	37	1.1	0.8-1.5
1981	185	3.5	3.0-4.0	101	4.2	3.4-5.1	84	2.8	2.2-3.6	64	1.2	1.0-1.6	33	1.5	1.0-2.1	31	1.0	0.7-1.5
1982	212	4.0	3.5-4.7	120	5.1	4.2-6.1	92	3.2	2.6-4.0	51	0.9	0.6-1.2	26	1.0	0.6-1.5	25	0.8	0.5-1.2
1983	192	3.7	3.2-4.3	113	4.9	4.0-5.9	79	2.7	2.1-3.4	62	1.1	0.8-1.4	36	1.5	1.0-2.1	26	0.8	0.5-1.2
1984	187	3.6	3.1-4.2	103	4.3	3.5-5.3	84	3.0	2.4-3.8	63	1.2	0.9-1.6	34	1.5	1.0-2.1	29	0.9	0.6-1.4
1985	211	4.1	3.5-4.7	128	5.4	4.5-6.5	83	2.9	2.3-3.6	58	1.1	0.8-1.4	30	1.2	0.8-1.7	28	1.0	0.6-1.4
1986	209	4.1	3.5-4.7	129	5.5	4.5-6.5	80	2.9	2.2-3.6	56	1.0	0.8-1.4	38	1.6	1.1-2.2	18	0.6	0.3-1.0
1987	174	3.5	3.0-4.1	88	3.9	3.1-4.8	86	3.3	2.6-4.1	40	0.8	0.5-1.1	24	1.0	0.7-1.6	16	0.5	0.3-0.9
1988	199	4.0	3.4-4.6	113	4.9	4.0-6.0	86	3.1	2.5-3.9	39	0.7	0.5-1.0	22	1.0	0.6-1.6	17	0.6	0.3-0.9
1989	180	3.8	3.2-4.4	104	4.5	3.7-5.5	76	3.0	2.4-3.8	56	1.2	0.9-1.5	32	1.5	1.0-2.1	24	0.9	0.6-1.3
1990	162	3.4	2.9-4.0	97	4.4	3.6-5.4	65	2.4	1.9-3.2	55	1.1	0.8-1.4	38	1.7	1.2-2.4	17	0.6	0.3-0.9
1991	193	4.2	3.6-4.8	101	4.7	3.8-5.7	92	3.8	3.0-4.7	43	0.9	0.6-1.2	25	1.2	0.8-1.8	18	0.6	0.4-1.0
1992	181	3.7	3.2-4.3	108	4.7	3.9-5.7	73	2.7	2.1-3.4	39	0.8	0.6-1.1	22	0.9	0.6-1.4	17	0.7	0.4-1.1
1993	156	3.3	2.8-3.9	81	3.8	3.0-4.8	75	2.8	2.2-3.6	34	0.7	0.5-0.9	19	0.8	0.5-1.3	15	0.5	0.3-0.8
1994	173	3.7	3.1-4.3	101	4.7	3.8-5.7	72	2.9	2.2-3.6	41	0.8	0.6-1.1	18	0.9	0.5-1.4	23	0.8	0.5-1.3
1995	182	4.0	3.4-4.6	94	4.3	3.4-5.2	88	3.7	2.9-4.6	31	0.7	0.5-1.0	17	0.8	0.5-1.3	14	0.6	0.3-1.0
1996	144	3.1	2.6-3.6	76	3.5	2.7-4.4	68	2.7	2.1-3.5	31	0.6	0.4-0.9	16	0.7	0.4-1.2	15	0.5	0.3-0.8
1997	162	3.5	3.0-4.1	93	4.3	3.4-5.3	69	2.9	2.3-3.7	37	0.7	0.5-1.0	18	0.8	0.5-1.3	19	0.7	0.4-1.1
1998	183	3.9	3.3-4.5	97	4.3	3.5-5.3	86	3.5	2.8-4.3	35	0.7	0.5-1.0	17	0.8	0.4-1.2	18	0.7	0.4-1.1
1999	162	3.4	2.9-3.9	77	3.3	2.6-4.1	85	3.4	2.7-4.2	21	0.4	0.3-0.7	10	0.5	0.2-0.9	11	0.4	0.2-0.7
2000	206	4.4	3.8-5.0	108	4.8	3.9-5.8	98	4.1	3.3-5.0	36	0.7	0.5-1.0	19	0.9	0.5-1.4	17	0.6	0.4-1.0
2001	179	3.7	3.1-4.2	91	3.9	3.1-4.8	88	3.4	2.7-4.2	24	0.5	0.3-0.7	13	0.6	0.3-1.0	11	0.4	0.2-0.7
2002	197	4.1	3.5-4.7	104	4.5	3.7-5.5	93	3.7	3.0-4.5	29	0.6	0.4-0.9	21	1.0	0.6-1.5	8	0.3	0.1-0.6
2003	185	4.0	3.5-4.7	94	4.3	3.4-5.2	91	3.9	3.1-4.8	24	0.5	0.3-0.7	11	0.5	0.3-0.9	13	0.5	0.3-0.9
2004	172	3.6	3.1-4.2	89	3.9	3.1-4.8	83	3.4	2.7-4.2	23	0.5	0.3-0.7	13	0.6	0.3-1.0	10	0.4	0.2-0.7
2005	181	3.9	3.3-4.5	104	4.8	3.9-5.8	77	3.1	2.4-3.8	23	0.5	0.3-0.7	14	0.7	0.4-1.1	9	0.3	0.1-0.6
2006	188	4.1	3.5-4.7	106	4.9	4.0-5.9	82	3.3	2.6-4.1	15	0.3	0.2-0.5	12	0.6	0.3-1.0	3	0.1	0.0-0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Hodgkin lymphoma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	191	4.0	3.5-4.6	84	3.7	3.0-4.7	107	4.2	3.4-5.1	32	0.7	0.4-0.9	17	0.8	0.5-1.3	15	0.5	0.3-0.9
2008	173	3.7	3.2-4.3	82	3.7	2.9-4.6	91	3.7	2.9-4.5	33	0.6	0.4-0.9	22	1.0	0.6-1.5	11	0.4	0.2-0.7
2009	178	3.9	3.3-4.5	103	4.7	3.8-5.7	75	3.1	2.4-3.9	23	0.5	0.3-0.7	15	0.7	0.4-1.2	8	0.3	0.1-0.6
2010	176	3.7	3.2-4.3	90	3.9	3.1-4.8	86	3.5	2.8-4.3	22	0.4	0.3-0.7	15	0.6	0.3-1.0	7	0.3	0.1-0.6
2011	181	3.8	3.3-4.4	100	4.3	3.5-5.3	81	3.3	2.6-4.2	32	0.7	0.4-0.9	17	0.8	0.5-1.3	15	0.5	0.3-0.9
2012	162	3.5	3.0-4.1	82	3.7	2.9-4.6	80	3.5	2.7-4.3	20	0.4	0.2-0.6	11	0.5	0.2-0.9	9	0.3	0.1-0.6
2013	172	3.5	3.0-4.1	73	3.2	2.5-4.0	99	3.8	3.1-4.7	22	0.4	0.3-0.7	13	0.6	0.3-1.0	9	0.3	0.1-0.7
2014	178	3.8	3.3-4.4	97	4.3	3.5-5.3	81	3.3	2.6-4.2	25	0.5	0.3-0.8	10	0.4	0.2-0.8	15	0.5	0.3-0.9
2015	179	3.8	3.2-4.4	107	4.6	3.8-5.6	72	3.0	2.3-3.8	24	0.5	0.3-0.7	13	0.6	0.3-1.0	11	0.4	0.2-0.7
2016	175	3.7	3.1-4.3	87	3.8	3.0-4.7	88	3.5	2.8-4.4	24	0.4	0.3-0.6	14	0.6	0.3-1.0	10	0.3	0.2-0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Non-Hodgkin lymphomas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	733	12.0	11.1-12.9	370	14.7	13.2-16.3	363	10.3	9.2-11.4	398	6.5	5.9-7.2	217	8.9	7.7-10.2	181	5.0	4.3-5.8
1977	694	11.6	10.7-12.5	350	13.7	12.3-15.3	344	10.0	8.9-11.2	354	5.8	5.2-6.5	169	6.8	5.7-7.9	185	5.2	4.4-6.0
1978	697	12.0	11.1-12.9	351	14.2	12.7-15.8	346	10.3	9.2-11.5	371	6.3	5.6-6.9	198	8.1	6.9-9.3	173	5.0	4.3-5.8
1979	746	12.9	12.0-13.9	347	14.8	13.3-16.5	399	11.6	10.4-12.8	364	6.4	5.7-7.1	181	7.9	6.7-9.2	183	5.3	4.5-6.2
1980	814	14.3	13.3-15.3	395	16.6	15.0-18.4	419	12.4	11.2-13.7	373	6.4	5.7-7.1	189	7.8	6.7-9.0	184	5.3	4.6-6.2
1981	803	14.1	13.1-15.2	381	16.2	14.6-18.0	422	12.5	11.3-13.8	375	6.5	5.8-7.2	172	7.5	6.4-8.8	203	5.8	5.0-6.7
1982	875	15.7	14.6-16.8	402	17.4	15.7-19.2	473	14.2	12.9-15.6	404	7.2	6.5-8.0	190	8.5	7.3-9.8	214	6.3	5.5-7.2
1983	869	16.1	15.0-17.2	446	20.3	18.4-22.3	423	12.8	11.5-14.1	434	7.9	7.1-8.7	214	9.7	8.4-11.1	220	6.5	5.6-7.4
1984	879	16.0	14.9-17.1	447	19.9	18.1-21.9	432	13.0	11.8-14.3	376	6.8	6.1-7.5	194	8.8	7.5-10.1	182	5.2	4.5-6.1
1985	972	18.0	16.9-19.2	522	23.8	21.8-26.0	450	13.6	12.3-15.0	445	8.0	7.3-8.8	240	10.9	9.5-12.4	205	6.0	5.2-6.9
1986	1026	19.3	18.1-20.5	540	24.1	22.1-26.3	486	15.0	13.7-16.5	393	7.3	6.6-8.1	216	10.0	8.7-11.4	177	5.2	4.5-6.1
1987	1037	20.0	18.8-21.3	560	26.1	23.9-28.4	477	15.0	13.6-16.4	411	7.6	6.9-8.4	214	9.8	8.5-11.3	197	5.8	5.0-6.7
1988	956	18.5	17.3-19.7	515	23.8	21.7-25.9	441	14.1	12.7-15.5	447	8.3	7.6-9.2	218	10.3	8.9-11.8	229	6.7	5.8-7.7
1989	986	19.9	18.7-21.3	578	27.5	25.2-29.8	408	13.5	12.1-14.9	420	8.1	7.3-8.9	238	11.4	10.0-13.0	182	5.7	4.9-6.6
1990	1105	22.1	20.8-23.4	635	30.0	27.6-32.4	470	15.4	14.0-16.9	428	8.2	7.5-9.1	214	10.2	8.8-11.6	214	6.7	5.8-7.7
1991	1229	24.8	23.4-26.3	702	33.6	31.1-36.2	527	17.6	16.1-19.2	414	8.3	7.5-9.1	222	10.8	9.4-12.4	192	6.0	5.2-7.0
1992	1123	22.9	21.5-24.3	628	29.8	27.5-32.3	495	16.9	15.4-18.5	424	8.2	7.5-9.1	217	10.6	9.2-12.1	207	6.6	5.7-7.6
1993	1101	22.7	21.3-24.1	642	30.8	28.4-33.3	459	15.8	14.3-17.3	399	7.8	7.1-8.7	213	10.5	9.1-12.0	186	6.0	5.2-7.0
1994	1298	27.0	25.5-28.5	755	36.0	33.4-38.7	543	19.1	17.5-20.8	458	9.2	8.3-10.1	236	11.6	10.1-13.2	222	7.2	6.3-8.3
1995	1231	25.6	24.1-27.1	693	33.4	30.9-36.0	538	18.8	17.2-20.6	429	8.6	7.8-9.5	223	11.1	9.7-12.6	206	6.7	5.8-7.8
1996	1199	24.9	23.5-26.3	664	32.2	29.8-34.8	535	19.1	17.5-20.8	436	8.8	7.9-9.6	233	11.8	10.3-13.4	203	6.7	5.8-7.8
1997	1148	23.7	22.3-25.1	603	29.3	27.0-31.8	545	18.9	17.3-20.6	445	8.9	8.1-9.8	229	11.4	10.0-13.0	216	7.1	6.2-8.2
1998	1073	22.1	20.8-23.5	579	27.7	25.5-30.0	494	17.6	16.1-19.3	430	8.6	7.8-9.5	204	10.0	8.7-11.5	226	7.5	6.5-8.5
1999	1086	22.2	20.9-23.5	558	26.7	24.5-29.0	528	18.6	17.0-20.3	400	8.0	7.2-8.8	194	9.6	8.3-11.1	206	6.8	5.9-7.8
2000	1085	22.1	20.8-23.4	551	26.0	23.8-28.3	534	18.7	17.1-20.4	397	7.8	7.1-8.7	197	9.6	8.3-11.1	200	6.6	5.7-7.6
2001	1058	21.4	20.1-22.7	572	27.1	24.9-29.5	486	17.0	15.5-18.6	381	7.4	6.7-8.2	214	10.4	9.0-11.9	167	5.2	4.4-6.1
2002	1043	21.2	19.9-22.5	579	27.2	25.0-29.5	464	16.2	14.7-17.8	338	6.6	5.9-7.4	176	8.5	7.3-9.8	162	5.3	4.5-6.2
2003	1152	23.3	21.9-24.7	595	28.0	25.7-30.3	557	19.6	18.0-21.3	326	6.4	5.7-7.1	166	7.9	6.8-9.2	160	5.1	4.3-6.0
2004	1122	22.7	21.4-24.1	580	27.4	25.2-29.7	542	19.0	17.4-20.8	354	6.9	6.2-7.7	178	8.6	7.4-10.0	176	5.7	4.9-6.7
2005	1160	23.6	22.2-25.0	578	27.2	25.0-29.5	582	20.7	19.0-22.5	350	6.9	6.2-7.7	179	8.7	7.4-10.0	171	5.6	4.8-6.6
2006	1166	23.9	22.5-25.3	648	30.5	28.2-33.0	518	18.4	16.8-20.1	358	7.1	6.4-7.9	187	9.1	7.8-10.5	171	5.6	4.8-6.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Non-Hodgkin lymphomas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	1113	22.9	21.6-24.3	580	27.5	25.2-29.8	533	19.3	17.7-21.1	338	6.8	6.1-7.6	180	8.7	7.4-10.1	158	5.4	4.6-6.4
2008	1173	23.8	22.5-25.2	626	29.2	27.0-31.6	547	19.7	18.1-21.5	313	6.1	5.5-6.8	178	8.5	7.3-9.8	135	4.4	3.6-5.2
2009	1223	24.9	23.5-26.4	618	29.1	26.8-31.5	605	21.7	19.9-23.5	314	6.1	5.5-6.9	172	8.1	6.9-9.4	142	4.6	3.9-5.5
2010	1173	23.7	22.3-25.1	615	28.4	26.2-30.8	558	19.7	18.1-21.5	290	5.7	5.1-6.4	150	7.3	6.1-8.5	140	4.6	3.8-5.4
2011	1164	23.3	22.0-24.7	592	26.9	24.7-29.1	572	20.3	18.6-22.0	302	5.8	5.1-6.5	154	7.1	6.0-8.4	148	4.8	4.0-5.7
2012	1175	23.4	22.0-24.8	645	29.1	26.9-31.5	530	18.9	17.3-20.6	293	5.5	4.9-6.2	162	7.5	6.3-8.7	131	4.0	3.3-4.8
2013	1211	23.6	22.3-25.0	655	29.1	26.9-31.5	556	19.2	17.6-20.9	276	5.2	4.5-5.8	147	6.7	5.6-7.9	129	4.0	3.3-4.8
2014	1170	22.6	21.3-23.9	630	27.8	25.6-30.1	540	18.5	16.9-20.2	298	5.4	4.8-6.1	159	7.1	6.0-8.3	139	4.2	3.5-5.0
2015	1137	21.7	20.4-23.0	612	26.1	24.0-28.2	525	18.1	16.5-19.7	255	4.7	4.1-5.4	152	6.7	5.7-7.9	103	3.3	2.7-4.0
2016	1128	21.1	19.9-22.4	600	25.3	23.3-27.5	528	17.8	16.3-19.5	373	6.6	6.0-7.3	203	8.6	7.5-9.9	170	5.1	4.3-6.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Myeloma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	243	3.9	3.4-4.5	123	4.9	4.1-5.9	120	3.3	2.7-4.0	151	2.5	2.1-3.0	75	3.2	2.5-4.1	76	2.1	1.6-2.6
1977	258	4.2	3.7-4.7	124	5.1	4.2-6.1	134	3.6	3.0-4.3	162	2.5	2.2-3.0	81	3.2	2.5-4.0	81	2.1	1.7-2.6
1978	253	4.2	3.7-4.7	123	5.1	4.2-6.1	130	3.5	2.9-4.2	149	2.4	2.0-2.9	66	2.7	2.0-3.4	83	2.2	1.8-2.8
1979	267	4.6	4.0-5.2	139	5.9	5.0-7.1	128	3.6	3.0-4.4	173	2.9	2.5-3.4	71	2.9	2.2-3.7	102	2.8	2.3-3.5
1980	241	4.2	3.7-4.8	114	5.1	4.2-6.1	127	3.6	3.0-4.3	167	2.8	2.4-3.3	78	3.3	2.6-4.2	89	2.5	2.0-3.1
1981	302	5.1	4.5-5.7	147	6.3	5.3-7.5	155	4.3	3.6-5.1	169	2.9	2.5-3.4	86	3.9	3.1-4.8	83	2.3	1.8-2.9
1982	267	4.6	4.0-5.2	122	5.4	4.5-6.5	145	4.2	3.5-4.9	159	2.7	2.3-3.2	70	3.0	2.3-3.9	89	2.6	2.1-3.2
1983	276	4.8	4.3-5.5	140	6.3	5.3-7.5	136	3.9	3.3-4.7	160	2.8	2.3-3.2	68	3.0	2.3-3.8	92	2.5	2.0-3.1
1984	232	4.1	3.6-4.7	111	5.2	4.2-6.3	121	3.4	2.8-4.1	157	2.7	2.3-3.2	75	3.4	2.6-4.2	82	2.3	1.8-2.9
1985	253	4.5	3.9-5.1	136	6.1	5.1-7.3	117	3.4	2.8-4.1	145	2.5	2.1-2.9	82	3.8	3.0-4.7	63	1.8	1.3-2.3
1986	254	4.5	4.0-5.1	129	5.8	4.8-6.9	125	3.6	3.0-4.3	144	2.5	2.1-3.0	68	3.0	2.3-3.9	76	2.2	1.7-2.8
1987	248	4.4	3.9-5.0	137	6.4	5.3-7.6	111	3.2	2.6-3.9	154	2.8	2.4-3.3	88	4.1	3.3-5.1	66	1.9	1.4-2.4
1988	246	4.5	3.9-5.1	128	6.2	5.1-7.4	118	3.5	2.9-4.2	141	2.5	2.1-2.9	74	3.5	2.7-4.4	67	1.9	1.4-2.4
1989	251	4.8	4.2-5.4	108	5.1	4.2-6.2	143	4.6	3.8-5.5	150	2.7	2.3-3.2	72	3.3	2.6-4.2	78	2.3	1.8-2.9
1990	263	5.0	4.4-5.6	131	6.1	5.1-7.3	132	4.2	3.5-5.0	139	2.5	2.1-3.0	69	3.3	2.5-4.2	70	2.1	1.6-2.7
1991	238	4.6	4.0-5.2	111	5.3	4.3-6.4	127	4.1	3.4-5.0	127	2.4	2.0-2.8	59	2.8	2.2-3.7	68	2.1	1.6-2.7
1992	271	5.2	4.6-5.9	145	7.1	6.0-8.4	126	4.0	3.3-4.8	149	2.8	2.4-3.3	74	3.7	2.9-4.6	75	2.4	1.9-3.0
1993	259	5.1	4.5-5.8	119	6.1	5.0-7.3	140	4.6	3.8-5.4	177	3.4	2.9-3.9	90	4.4	3.6-5.5	87	2.7	2.2-3.4
1994	238	4.8	4.2-5.4	113	5.5	4.5-6.6	125	4.3	3.5-5.1	154	3.0	2.6-3.5	73	3.5	2.8-4.5	81	2.7	2.1-3.3
1995	247	5.0	4.4-5.6	118	5.8	4.8-7.0	129	4.4	3.7-5.3	149	2.9	2.5-3.5	71	3.6	2.8-4.5	78	2.5	2.0-3.2
1996	285	5.7	5.1-6.4	139	7.0	5.9-8.3	146	5.0	4.2-5.9	151	3.0	2.5-3.5	63	3.2	2.4-4.1	88	2.9	2.3-3.6
1997	239	4.8	4.2-5.5	120	6.0	5.0-7.2	119	4.0	3.3-4.8	133	2.7	2.2-3.2	66	3.3	2.6-4.2	67	2.3	1.7-2.9
1998	293	5.9	5.2-6.6	146	7.2	6.1-8.5	147	5.0	4.3-6.0	139	2.8	2.3-3.3	69	3.4	2.6-4.3	70	2.3	1.8-2.9
1999	279	5.6	4.9-6.3	119	5.8	4.8-7.0	160	5.5	4.7-6.4	138	2.7	2.3-3.2	63	3.1	2.4-4.0	75	2.5	1.9-3.1
2000	276	5.5	4.9-6.2	130	6.3	5.3-7.5	146	5.0	4.2-5.9	168	3.3	2.8-3.8	72	3.5	2.7-4.4	96	3.1	2.5-3.8
2001	296	5.9	5.2-6.6	152	7.4	6.3-8.7	144	4.9	4.1-5.7	136	2.6	2.2-3.1	70	3.4	2.7-4.3	66	2.1	1.6-2.7
2002	305	6.0	5.4-6.8	160	7.7	6.6-9.0	145	5.0	4.2-5.9	143	2.8	2.3-3.3	71	3.5	2.8-4.5	72	2.4	1.8-3.0
2003	322	6.3	5.7-7.1	170	8.2	7.0-9.5	152	5.3	4.5-6.2	133	2.5	2.1-3.0	59	2.9	2.2-3.7	74	2.3	1.8-2.9
2004	311	6.2	5.5-6.9	155	7.4	6.3-8.6	156	5.4	4.6-6.3	121	2.3	1.9-2.8	56	2.8	2.1-3.6	65	2.1	1.6-2.7
2005	335	6.7	6.0-7.4	171	8.2	7.0-9.5	164	5.6	4.8-6.6	136	2.7	2.2-3.1	66	3.2	2.5-4.1	70	2.3	1.7-2.9
2006	331	6.7	6.0-7.5	161	7.8	6.6-9.1	170	6.0	5.1-7.0	126	2.5	2.1-3.0	82	4.0	3.2-5.0	44	1.5	1.1-2.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Myeloma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	312	6.3	5.6-7.0	169	8.1	6.9-9.4	143	5.0	4.2-5.9	123	2.4	2.0-2.9	68	3.3	2.6-4.2	55	1.8	1.3-2.3
2008	338	6.8	6.1-7.6	197	9.4	8.2-10.9	141	5.0	4.2-5.9	138	2.7	2.3-3.2	68	3.3	2.6-4.2	70	2.4	1.8-3.0
2009	376	7.5	6.8-8.3	205	9.7	8.4-11.2	171	5.9	5.0-6.9	117	2.3	1.9-2.7	66	3.2	2.5-4.0	51	1.6	1.2-2.2
2010	375	7.4	6.7-8.2	203	9.4	8.1-10.8	172	6.0	5.1-7.0	148	2.9	2.4-3.4	82	4.0	3.1-4.9	66	2.2	1.7-2.8
2011	404	7.9	7.2-8.8	235	10.7	9.4-12.2	169	6.0	5.1-7.0	118	2.2	1.9-2.7	53	2.5	1.9-3.3	65	2.2	1.7-2.8
2012	376	7.3	6.5-8.1	220	9.9	8.6-11.3	156	5.3	4.5-6.3	123	2.3	1.9-2.8	58	2.7	2.1-3.5	65	2.1	1.6-2.7
2013	399	7.5	6.8-8.3	202	8.9	7.7-10.2	197	6.5	5.6-7.5	147	2.7	2.3-3.2	85	3.7	3.0-4.6	62	1.9	1.5-2.5
2014	424	8.1	7.3-8.9	248	10.8	9.5-12.3	176	5.9	5.1-6.9	146	2.7	2.2-3.2	71	3.1	2.4-4.0	75	2.3	1.8-2.9
2015	387	7.2	6.5-8.0	204	8.8	7.6-10.1	183	6.0	5.2-7.0	135	2.5	2.1-3.0	77	3.3	2.6-4.2	58	1.8	1.4-2.4
2016	368	6.7	6.1-7.5	233	9.7	8.5-11.1	135	4.5	3.7-5.3	199	3.6	3.1-4.1	120	5.1	4.2-6.1	79	2.4	1.9-3.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Leukemias

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	689	11.9	11.0-12.9	383	15.8	14.2-17.6	306	9.1	8.0-10.2	500	8.7	8.0-9.6	263	11.2	9.8-12.7	237	7.0	6.1-8.0
1977	641	11.2	10.3-12.1	349	14.5	13.0-16.2	292	8.8	7.7-9.9	458	8.0	7.3-8.8	254	10.8	9.5-12.3	204	6.1	5.3-7.1
1978	686	12.1	11.2-13.1	394	16.8	15.1-18.7	292	8.9	7.8-10.0	463	8.1	7.4-8.9	250	10.7	9.4-12.1	213	6.4	5.6-7.4
1979	646	11.5	10.6-12.5	328	13.8	12.3-15.4	318	9.9	8.8-11.2	467	8.3	7.6-9.2	238	10.2	8.9-11.7	229	7.2	6.2-8.2
1980	739	13.4	12.4-14.4	378	16.5	14.9-18.4	361	11.4	10.2-12.7	436	7.8	7.1-8.6	252	11.2	9.8-12.7	184	5.6	4.7-6.5
1981	655	11.9	11.0-12.9	362	16.0	14.3-17.7	293	9.1	8.0-10.2	450	8.2	7.4-9.0	245	10.8	9.5-12.3	205	6.4	5.5-7.4
1982	691	12.6	11.7-13.6	383	17.1	15.4-19.0	308	9.5	8.4-10.7	474	8.7	7.9-9.6	264	12.0	10.5-13.5	210	6.5	5.6-7.5
1983	648	12.2	11.3-13.2	328	14.8	13.2-16.5	320	10.7	9.5-12.1	419	7.8	7.1-8.6	224	10.4	9.0-11.9	195	6.1	5.2-7.1
1984	687	12.7	11.8-13.7	370	17.2	15.5-19.1	317	9.9	8.8-11.1	458	8.4	7.6-9.2	241	11.2	9.8-12.8	217	6.5	5.6-7.5
1985	649	11.9	11.0-12.9	353	16.1	14.4-17.9	296	9.2	8.1-10.3	407	7.4	6.7-8.2	204	9.4	8.1-10.8	203	6.1	5.3-7.1
1986	694	13.1	12.2-14.2	352	15.9	14.2-17.7	342	11.4	10.1-12.7	405	7.5	6.8-8.3	202	9.2	7.9-10.6	203	6.3	5.5-7.3
1987	559	10.6	9.7-11.5	265	12.4	10.9-14.1	294	9.3	8.3-10.5	358	6.6	5.9-7.3	172	8.0	6.8-9.3	186	5.9	5.0-6.8
1988	595	11.5	10.6-12.5	312	15.0	13.3-16.8	283	9.3	8.2-10.5	353	6.7	6.0-7.5	192	9.2	7.9-10.6	161	5.1	4.3-6.0
1989	679	13.1	12.1-14.2	362	17.3	15.6-19.3	317	10.4	9.2-11.6	366	6.9	6.2-7.7	192	9.2	8.0-10.7	174	5.4	4.6-6.3
1990	687	13.7	12.7-14.8	361	17.3	15.5-19.2	326	11.2	9.9-12.5	361	7.0	6.3-7.8	181	9.0	7.7-10.4	180	5.9	5.0-6.9
1991	624	12.7	11.7-13.8	348	17.0	15.3-19.0	276	9.4	8.3-10.7	355	7.0	6.3-7.8	208	10.4	9.0-12.0	147	4.8	4.0-5.7
1992	639	12.9	11.9-13.9	352	17.3	15.5-19.2	287	9.9	8.7-11.1	324	6.4	5.7-7.2	178	9.0	7.7-10.4	146	4.8	4.0-5.7
1993	594	12.1	11.1-13.1	293	14.3	12.7-16.1	301	10.7	9.5-12.0	336	6.6	5.9-7.4	190	9.3	8.0-10.7	146	4.9	4.2-5.9
1994	694	14.0	13.0-15.1	355	17.6	15.8-19.6	339	11.4	10.2-12.8	387	7.6	6.9-8.4	197	9.7	8.4-11.2	190	6.1	5.2-7.0
1995	615	12.6	11.6-13.7	317	15.7	14.0-17.5	298	10.4	9.3-11.8	378	7.6	6.8-8.4	199	10.0	8.7-11.5	179	6.0	5.1-6.9
1996	626	12.9	11.9-13.9	350	17.2	15.4-19.1	276	9.8	8.6-11.0	340	6.8	6.1-7.5	165	8.1	6.9-9.5	175	5.9	5.1-6.9
1997	668	13.7	12.7-14.8	356	17.6	15.8-19.6	312	11.1	9.9-12.4	346	7.0	6.3-7.8	199	10.0	8.7-11.5	147	4.9	4.1-5.8
1998	652	13.3	12.3-14.4	362	17.5	15.8-19.5	290	10.2	9.0-11.4	362	7.3	6.6-8.1	205	10.0	8.7-11.5	157	5.3	4.5-6.3
1999	729	14.8	13.7-15.9	380	18.5	16.7-20.5	349	12.1	10.9-13.5	370	7.3	6.6-8.1	203	9.9	8.6-11.4	167	5.6	4.7-6.5
2000	674	13.7	12.6-14.7	344	16.7	15.0-18.5	330	11.5	10.3-12.9	329	6.5	5.8-7.3	149	7.2	6.1-8.4	180	6.0	5.1-7.0
2001	695	14.0	13.0-15.1	355	17.1	15.4-19.0	340	11.8	10.6-13.2	344	6.7	6.0-7.5	176	8.5	7.3-9.9	168	5.4	4.6-6.3
2002	633	12.8	11.8-13.8	358	17.1	15.3-18.9	275	9.6	8.4-10.8	342	6.7	6.0-7.4	178	8.6	7.4-10.0	164	5.4	4.6-6.3
2003	716	14.5	13.5-15.6	369	17.6	15.9-19.5	347	12.2	10.9-13.6	309	6.1	5.4-6.8	162	7.8	6.6-9.1	147	4.9	4.2-5.9
2004	683	13.7	12.7-14.8	360	17.3	15.5-19.2	323	11.2	10.0-12.6	354	6.9	6.2-7.7	185	9.0	7.8-10.4	169	5.6	4.8-6.5
2005	710	14.5	13.4-15.6	400	19.0	17.2-21.0	310	11.1	9.9-12.5	345	6.8	6.1-7.6	193	9.3	8.0-10.7	152	5.1	4.3-6.0
2006	692	14.3	13.3-15.4	375	18.2	16.4-20.1	317	11.5	10.2-12.9	344	6.8	6.1-7.6	177	8.6	7.4-10.0	167	5.6	4.8-6.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Whites
Leukemias

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	736	15.4	14.3-16.6	411	19.6	17.8-21.7	325	12.2	10.8-13.6	325	6.5	5.8-7.3	178	8.7	7.4-10.1	147	5.3	4.4-6.2
2008	723	15.1	14.0-16.2	403	19.5	17.7-21.6	320	11.8	10.5-13.3	347	7.0	6.3-7.8	171	8.3	7.1-9.7	176	6.2	5.3-7.2
2009	789	16.2	15.1-17.4	425	20.2	18.3-22.3	364	13.1	11.7-14.6	316	6.3	5.6-7.1	143	6.9	5.8-8.1	173	5.9	5.0-6.9
2010	822	16.8	15.6-18.0	459	21.6	19.6-23.7	363	13.0	11.6-14.4	327	6.4	5.7-7.2	178	8.5	7.3-9.9	149	5.0	4.2-6.0
2011	835	16.5	15.4-17.7	457	20.9	19.0-23.0	378	13.3	12.0-14.8	340	6.5	5.8-7.2	197	9.3	8.0-10.7	143	4.5	3.8-5.4
2012	772	15.2	14.2-16.4	449	20.3	18.5-22.3	323	11.5	10.3-12.9	311	5.9	5.3-6.6	176	8.2	7.0-9.5	135	4.3	3.6-5.1
2013	812	15.9	14.8-17.0	473	21.4	19.5-23.5	339	11.8	10.5-13.2	342	6.6	5.9-7.3	196	8.9	7.7-10.3	146	4.9	4.1-5.8
2014	847	16.6	15.5-17.8	435	19.4	17.6-21.3	412	14.7	13.2-16.2	293	5.5	4.8-6.1	148	6.7	5.7-7.9	145	4.6	3.9-5.5
2015	858	16.5	15.4-17.6	486	21.3	19.4-23.3	372	12.7	11.4-14.1	330	6.1	5.4-6.8	182	8.1	6.9-9.4	148	4.6	3.9-5.5
2016	784	14.8	13.7-15.9	456	19.4	17.6-21.3	328	11.1	9.9-12.4	441	8.1	7.4-8.9	237	10.2	9.0-11.7	204	6.7	5.7-7.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
All Invasive Malignant Tumors

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	35622	397.3	393.1-401.6	17524	471.4	464.1-478.9	18098	358.9	353.6-364.2	19253	217.2	214.0-220.3	10040	277.4	271.7-283.2	9213	180.9	177.1-184.6
1977	35994	396.6	392.5-400.8	17697	470.8	463.5-478.2	18297	357.8	352.6-363.1	19718	219.9	216.7-223.0	10324	281.8	276.1-287.6	9394	182.0	178.3-185.7
1978	36509	399.1	394.9-403.3	17909	473.7	466.4-481.1	18600	360.2	355.0-365.5	19927	219.0	215.9-222.1	10361	281.7	276.0-287.5	9566	182.0	178.3-185.7
1979	37321	403.1	398.9-407.3	18276	477.5	470.2-484.8	19045	364.3	359.1-369.6	20008	218.1	215.0-221.2	10323	278.0	272.4-283.7	9685	182.4	178.7-186.1
1980	38760	413.1	408.9-417.3	19375	497.9	490.6-505.4	19385	366.8	361.6-372.1	20562	220.9	217.8-224.0	10715	284.0	278.3-289.7	9847	182.9	179.2-186.6
1981	39429	417.8	413.6-422.1	19523	502.1	494.7-509.6	19906	374.1	368.8-379.5	20580	219.0	216.0-222.1	10724	283.5	277.8-289.2	9856	181.1	177.5-184.8
1982	40097	420.1	415.9-424.3	19827	503.7	496.3-511.1	20270	376.9	371.6-382.2	20850	220.1	217.1-223.2	10783	283.7	278.0-289.4	10067	183.1	179.4-186.7
1983	41295	430.0	425.8-434.2	20367	513.0	505.6-520.5	20928	387.8	382.5-393.2	21092	220.3	217.3-223.3	10891	284.3	278.7-289.9	10201	183.5	179.9-187.2
1984	41874	433.1	428.9-437.3	20417	509.5	502.2-516.9	21457	394.4	389.0-399.8	21310	221.0	218.0-224.1	10826	280.1	274.5-285.7	10484	187.4	183.8-191.1
1985	43359	446.1	441.8-450.4	21220	524.6	517.2-532.1	22139	406.0	400.6-411.5	21636	223.1	220.1-226.2	11029	283.5	278.0-289.1	10607	188.5	184.9-192.2
1986	43439	444.5	440.3-448.8	21171	521.8	514.5-529.2	22268	406.3	400.9-411.8	22000	225.3	222.3-228.4	11282	287.9	282.3-293.5	10718	189.2	185.6-192.9
1987	44231	448.6	444.4-452.9	21405	523.2	515.9-530.6	22826	412.4	406.9-417.9	21927	222.9	219.9-225.9	11161	283.2	277.7-288.7	10766	187.9	184.3-191.5
1988	44366	446.2	442.0-450.4	21455	520.5	513.2-527.8	22911	411.3	405.9-416.8	22220	223.3	220.4-226.3	11354	284.4	279.0-289.9	10866	188.1	184.5-191.8
1989	44746	446.3	442.1-450.5	21967	527.0	519.7-534.3	22779	405.8	400.5-411.2	21864	218.4	215.5-221.3	11092	275.4	270.0-280.8	10772	185.5	182.0-189.1
1990	47006	463.6	459.3-467.8	23213	546.1	538.8-553.4	23793	420.2	414.7-425.6	21869	215.5	212.6-218.4	11055	271.6	266.3-276.9	10814	183.0	179.6-186.6
1991	47717	466.7	462.4-470.9	23956	557.7	550.4-565.1	23761	417.4	412.0-422.9	22298	218.0	215.1-220.9	11310	275.2	270.0-280.6	10988	184.9	181.4-188.4
1992	49639	480.3	476.1-484.6	25606	588.4	581.0-595.9	24033	418.3	412.9-423.7	22511	217.6	214.8-220.5	11202	270.9	265.7-276.1	11309	188.2	184.7-191.7
1993	49666	477.1	472.9-481.3	25245	571.2	564.0-578.5	24421	421.8	416.5-427.3	22289	213.3	210.5-216.2	11105	265.1	260.0-270.3	11184	184.0	180.5-187.5
1994	49854	477.1	472.9-481.3	25095	565.2	558.1-572.4	24759	425.9	420.6-431.3	23050	219.0	216.2-221.9	11657	275.7	270.5-280.9	11393	186.0	182.6-189.5
1995	50516	481.6	477.4-485.9	25004	555.7	548.7-562.8	25512	439.0	433.5-444.5	22750	214.7	211.9-217.5	11416	268.1	263.1-273.2	11334	184.2	180.8-187.7
1996	52499	498.8	494.5-503.1	26374	582.7	575.6-589.9	26125	448.0	442.5-453.5	22600	212.1	209.4-214.9	11351	263.1	258.2-268.1	11249	182.1	178.7-185.5
1997	53688	506.5	502.2-510.8	26974	589.9	582.8-597.0	26714	454.5	449.0-460.1	22202	206.3	203.6-209.0	11167	255.5	250.7-260.4	11035	176.5	173.2-179.9
1998	54976	514.2	509.9-518.5	27420	591.1	584.1-598.2	27556	466.9	461.3-472.5	21751	200.0	197.4-202.7	10852	242.6	238.0-247.3	10899	173.0	169.7-176.3
1999	55852	517.9	513.6-522.2	28052	600.4	593.3-607.5	27800	465.7	460.2-471.3	22340	203.2	200.5-205.9	11153	247.9	243.3-252.6	11187	175.8	172.5-179.1
2000	57355	526.8	522.5-531.1	28960	615.1	607.9-622.2	28395	470.7	465.1-476.2	22276	200.1	197.5-202.8	11071	244.3	239.7-248.9	11205	173.6	170.3-176.9
2001	57075	521.1	516.9-525.5	28786	604.4	597.4-611.5	28289	468.1	462.6-473.7	22024	196.4	193.8-199.0	10955	238.8	234.3-243.4	11069	170.8	167.6-174.1
2002	58096	525.2	520.9-529.5	29901	619.4	612.4-626.5	28195	462.9	457.4-468.4	22012	194.2	191.6-196.8	10917	234.5	230.1-239.0	11095	170.1	166.9-173.3
2003	57910	518.3	514.1-522.6	29365	600.0	593.1-606.9	28545	464.7	459.3-470.2	21476	187.2	184.7-189.7	10525	222.7	218.4-227.0	10951	165.9	162.7-169.1
2004	58273	517.8	513.6-522.1	29520	596.3	589.5-603.2	28753	466.6	461.1-472.1	21435	185.3	182.8-187.8	10588	220.1	215.9-224.4	10847	163.3	160.2-166.4
2005	59052	521.7	517.5-526.0	29979	598.8	592.0-605.7	29073	469.3	463.9-474.9	21045	180.4	178.0-182.9	10423	214.7	210.6-218.9	10622	158.6	155.6-161.8
2006	60759	533.5	529.2-537.8	31242	616.5	609.6-623.5	29517	476.2	470.6-481.8	21126	180.0	177.6-182.5	10459	212.3	208.2-216.5	10667	159.1	156.0-162.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
All Invasive Malignant Tumors

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	61458	533.3	529.0-537.6	31790	616.1	609.3-623.0	29668	476.1	470.6-481.7	20927	175.8	173.4-178.2	10511	209.0	205.0-213.1	10416	154.0	151.0-157.1
2008	62509	536.5	532.2-540.8	31947	610.3	603.5-617.1	30562	486.8	481.2-492.4	21127	174.9	172.5-177.3	10629	208.0	204.0-212.1	10498	153.2	150.2-156.2
2009	62300	527.8	523.6-532.1	31903	597.8	591.2-604.5	30397	479.1	473.5-484.7	20713	169.8	167.4-172.2	10530	203.0	199.1-206.9	10183	147.9	144.9-150.8
2010	61137	512.7	508.6-516.9	30986	571.4	565.0-577.9	30151	472.9	467.4-478.5	20729	167.5	165.2-169.8	10261	194.3	190.5-198.1	10468	149.8	146.9-152.8
2011	62788	521.1	517.0-525.3	32012	581.5	575.0-588.0	30776	479.2	473.7-484.8	20649	164.6	162.3-166.9	10302	191.5	187.8-195.3	10347	146.4	143.5-149.3
2012	61455	504.2	500.1-508.3	30910	553.1	546.8-559.5	30545	472.7	467.2-478.3	21101	166.7	164.4-169.0	10719	197.4	193.6-201.2	10382	145.7	142.8-148.6
2013	62894	511.1	507.0-515.2	31407	553.3	547.0-559.6	31487	485.4	479.8-491.0	20707	160.8	158.6-163.1	10537	188.9	185.3-192.6	10170	141.0	138.2-143.9
2014	62675	502.9	498.8-507.0	31249	541.1	535.0-547.3	31426	480.0	474.4-485.6	20176	154.7	152.5-156.9	10356	182.9	179.4-186.6	9820	135.1	132.4-137.9
2015	63355	504.1	500.1-508.2	31466	537.1	531.0-543.2	31889	485.5	479.9-491.2	20166	152.7	150.5-154.9	10208	178.2	174.7-181.7	9958	135.0	132.2-137.8
2016	63194	498.0	494.0-502.1	31955	537.8	531.7-543.9	31239	473.3	467.7-478.9	19216	144.4	142.3-146.5	9695	166.8	163.4-170.2	9521	129.0	126.3-131.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Oral cavity and pharynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	987	10.6	10.0-11.3	670	16.6	15.3-18.0	317	6.1	5.5-6.9	396	4.3	3.8-4.7	285	7.1	6.3-8.1	111	2.1	1.7-2.5
1977	971	10.3	9.6-11.0	651	15.9	14.6-17.2	320	6.1	5.5-6.9	396	4.2	3.8-4.7	270	6.8	6.0-7.7	126	2.4	2.0-2.8
1978	1006	10.6	9.9-11.3	663	16.3	15.0-17.6	343	6.5	5.8-7.2	470	5.0	4.6-5.5	327	8.3	7.4-9.3	143	2.7	2.3-3.2
1979	1050	10.8	10.2-11.5	670	16.0	14.8-17.3	380	7.0	6.3-7.8	413	4.3	3.9-4.8	257	6.2	5.4-7.0	156	2.9	2.5-3.4
1980	1014	10.5	9.8-11.1	662	15.9	14.6-17.2	352	6.5	5.8-7.2	401	4.1	3.7-4.6	258	6.3	5.5-7.2	143	2.6	2.1-3.0
1981	1058	10.8	10.2-11.5	723	17.1	15.8-18.4	335	6.1	5.5-6.8	395	4.0	3.6-4.4	271	6.5	5.7-7.4	124	2.2	1.8-2.6
1982	1104	11.4	10.7-12.1	732	17.6	16.3-19.0	372	6.9	6.2-7.6	417	4.3	3.9-4.7	276	6.9	6.1-7.8	141	2.5	2.1-3.0
1983	1042	10.6	9.9-11.3	684	16.0	14.8-17.3	358	6.6	5.9-7.4	398	4.0	3.6-4.5	261	6.2	5.5-7.1	137	2.4	2.0-2.9
1984	1056	10.8	10.1-11.4	711	16.9	15.7-18.3	345	6.3	5.6-7.0	389	4.0	3.6-4.4	249	6.1	5.3-6.9	140	2.5	2.1-3.0
1985	1058	10.7	10.1-11.4	698	16.5	15.2-17.8	360	6.4	5.8-7.2	377	3.8	3.4-4.2	240	5.8	5.1-6.6	137	2.5	2.1-2.9
1986	1034	10.5	9.9-11.2	683	16.2	15.0-17.5	351	6.3	5.6-7.0	409	4.1	3.7-4.6	266	6.3	5.5-7.1	143	2.5	2.1-3.0
1987	1034	10.5	9.9-11.2	650	15.3	14.1-16.6	384	6.8	6.1-7.6	382	3.9	3.5-4.3	241	5.9	5.2-6.8	141	2.4	2.1-2.9
1988	1031	10.4	9.7-11.0	649	15.1	13.9-16.3	382	6.9	6.2-7.6	399	4.0	3.6-4.4	261	6.5	5.7-7.4	138	2.3	2.0-2.8
1989	998	9.9	9.3-10.5	661	15.0	13.8-16.2	337	6.0	5.4-6.7	332	3.3	3.0-3.7	201	4.8	4.1-5.5	131	2.3	1.9-2.7
1990	983	9.8	9.2-10.4	622	14.1	12.9-15.2	361	6.5	5.8-7.2	345	3.4	3.0-3.7	220	5.0	4.4-5.8	125	2.1	1.7-2.5
1991	1011	10.1	9.5-10.7	645	14.6	13.5-15.8	366	6.5	5.8-7.2	343	3.4	3.0-3.8	209	4.9	4.2-5.6	134	2.3	1.9-2.7
1992	1006	9.8	9.2-10.5	637	14.3	13.1-15.4	369	6.4	5.8-7.2	302	2.9	2.6-3.2	182	4.1	3.5-4.7	120	2.0	1.6-2.4
1993	1003	9.8	9.2-10.4	661	14.9	13.7-16.1	342	5.9	5.3-6.5	296	2.8	2.5-3.2	175	3.9	3.3-4.6	121	2.0	1.7-2.4
1994	996	9.7	9.1-10.3	652	14.5	13.4-15.7	344	6.0	5.4-6.7	305	2.9	2.6-3.3	211	4.7	4.1-5.4	94	1.6	1.3-1.9
1995	933	9.0	8.4-9.6	569	12.5	11.5-13.6	364	6.2	5.6-6.9	322	3.1	2.8-3.4	214	4.9	4.2-5.6	108	1.8	1.5-2.2
1996	1029	9.9	9.3-10.5	666	14.3	13.2-15.4	363	6.3	5.7-7.0	283	2.7	2.4-3.0	184	4.2	3.6-4.8	99	1.6	1.3-2.0
1997	1086	10.4	9.8-11.0	703	15.1	13.9-16.2	383	6.5	5.9-7.2	272	2.6	2.3-2.9	177	3.9	3.3-4.5	95	1.5	1.2-1.9
1998	1052	10.0	9.4-10.6	664	14.1	13.1-15.3	388	6.6	5.9-7.3	303	2.8	2.5-3.1	190	4.1	3.6-4.8	113	1.8	1.5-2.1
1999	1028	9.6	9.1-10.2	695	14.6	13.5-15.7	333	5.5	4.9-6.2	261	2.4	2.1-2.7	167	3.5	3.0-4.1	94	1.5	1.2-1.8
2000	1040	9.6	9.1-10.2	677	14.1	13.1-15.3	363	6.0	5.4-6.7	236	2.1	1.9-2.4	147	3.2	2.7-3.7	89	1.4	1.1-1.7
2001	1077	9.9	9.3-10.5	688	13.9	12.9-15.0	389	6.4	5.7-7.1	274	2.5	2.2-2.8	172	3.6	3.1-4.2	102	1.5	1.2-1.8
2002	1108	10.1	9.5-10.7	724	14.5	13.5-15.6	384	6.2	5.6-6.9	251	2.2	2.0-2.5	158	3.3	2.8-3.8	93	1.4	1.1-1.7
2003	1113	10.0	9.4-10.6	734	14.6	13.5-15.7	379	6.2	5.5-6.8	240	2.1	1.9-2.4	149	3.0	2.5-3.5	91	1.4	1.1-1.7
2004	1170	10.4	9.8-11.0	795	15.6	14.5-16.7	375	6.0	5.4-6.7	272	2.3	2.1-2.6	179	3.5	3.0-4.1	93	1.4	1.1-1.7
2005	1121	9.9	9.3-10.5	718	13.8	12.8-14.8	403	6.6	6.0-7.3	241	2.1	1.8-2.3	161	3.1	2.7-3.7	80	1.2	0.9-1.5
2006	1214	10.6	10.0-11.2	850	16.1	15.1-17.3	364	5.9	5.3-6.6	249	2.1	1.9-2.4	161	3.1	2.7-3.7	88	1.3	1.0-1.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Oral cavity and pharynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	1303	11.2	10.6-11.8	904	16.8	15.7-18.0	399	6.4	5.8-7.1	234	2.0	1.7-2.2	159	3.0	2.5-3.5	75	1.1	0.9-1.4
2008	1276	10.8	10.2-11.5	866	15.9	14.9-17.0	410	6.5	5.9-7.2	250	2.0	1.8-2.3	182	3.3	2.8-3.8	68	1.0	0.8-1.3
2009	1340	11.1	10.5-11.7	910	16.2	15.1-17.3	430	6.6	6.0-7.3	247	2.0	1.7-2.3	164	3.1	2.6-3.6	83	1.1	0.9-1.4
2010	1391	11.4	10.8-12.1	943	16.7	15.6-17.8	448	6.9	6.3-7.6	262	2.1	1.9-2.4	170	3.1	2.6-3.6	92	1.3	1.0-1.6
2011	1401	11.3	10.7-11.9	977	16.8	15.7-17.9	424	6.4	5.8-7.1	258	2.0	1.8-2.3	168	3.0	2.5-3.5	90	1.3	1.0-1.6
2012	1426	11.5	10.9-12.1	982	17.0	15.9-18.1	444	6.6	6.0-7.3	279	2.2	1.9-2.4	197	3.5	3.0-4.0	82	1.1	0.8-1.3
2013	1478	11.8	11.2-12.5	1043	18.0	16.9-19.1	435	6.4	5.8-7.1	277	2.1	1.9-2.4	191	3.2	2.8-3.8	86	1.1	0.9-1.4
2014	1518	11.9	11.3-12.6	1050	17.6	16.5-18.7	468	6.9	6.3-7.6	226	1.7	1.5-2.0	159	2.7	2.3-3.2	67	0.9	0.7-1.2
2015	1599	12.5	11.9-13.2	1152	19.1	18.0-20.3	447	6.8	6.1-7.5	281	2.1	1.9-2.4	189	3.2	2.8-3.7	92	1.2	1.0-1.5
2016	1675	12.8	12.2-13.5	1165	19.0	17.9-20.2	510	7.4	6.7-8.1	285	2.1	1.8-2.3	203	3.3	2.8-3.8	82	1.0	0.8-1.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Esophagus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	361	4.0	3.6-4.4	276	7.2	6.3-8.1	85	1.7	1.3-2.1	330	3.6	3.2-4.1	252	6.6	5.8-7.5	78	1.5	1.2-1.9
1977	362	4.0	3.6-4.4	255	6.7	5.9-7.7	107	2.0	1.7-2.5	323	3.5	3.1-3.9	232	6.0	5.2-6.9	91	1.7	1.4-2.1
1978	353	3.8	3.4-4.2	251	6.6	5.8-7.5	102	1.9	1.6-2.3	371	4.0	3.6-4.5	271	7.1	6.3-8.1	100	1.9	1.5-2.3
1979	374	3.9	3.5-4.4	256	6.3	5.5-7.2	118	2.2	1.8-2.6	325	3.5	3.1-3.9	225	5.8	5.0-6.6	100	1.8	1.5-2.3
1980	375	3.9	3.5-4.3	260	6.6	5.8-7.5	115	2.1	1.7-2.5	339	3.5	3.1-3.9	228	5.7	4.9-6.5	111	2.0	1.6-2.4
1981	375	3.9	3.5-4.3	258	6.4	5.6-7.3	117	2.1	1.7-2.5	355	3.7	3.3-4.1	262	6.7	5.8-7.6	93	1.7	1.4-2.1
1982	375	3.8	3.4-4.2	250	6.0	5.2-6.8	125	2.2	1.8-2.6	354	3.6	3.3-4.0	238	5.9	5.2-6.8	116	2.0	1.7-2.4
1983	394	4.0	3.6-4.4	280	6.8	6.0-7.7	114	2.0	1.6-2.4	368	3.8	3.4-4.3	276	7.1	6.2-8.0	92	1.6	1.3-2.0
1984	401	4.0	3.6-4.5	275	6.7	5.9-7.6	126	2.2	1.8-2.6	371	3.7	3.4-4.1	256	6.4	5.6-7.3	115	2.0	1.6-2.4
1985	373	3.8	3.4-4.2	280	6.8	6.0-7.6	93	1.6	1.3-2.0	359	3.6	3.2-4.0	265	6.4	5.6-7.3	94	1.6	1.3-2.0
1986	424	4.3	3.9-4.7	299	7.1	6.3-8.0	125	2.2	1.8-2.6	370	3.7	3.3-4.1	264	6.2	5.5-7.1	106	1.8	1.5-2.2
1987	399	4.0	3.6-4.4	276	6.5	5.8-7.4	123	2.1	1.7-2.5	391	3.9	3.5-4.3	284	6.9	6.0-7.7	107	1.8	1.5-2.2
1988	417	4.1	3.7-4.6	299	7.0	6.2-7.9	118	2.0	1.6-2.4	424	4.2	3.8-4.6	317	7.5	6.7-8.4	107	1.8	1.5-2.2
1989	413	4.0	3.6-4.4	299	6.9	6.1-7.7	114	1.9	1.6-2.3	405	4.0	3.6-4.4	301	7.1	6.3-7.9	104	1.7	1.4-2.1
1990	464	4.5	4.1-4.9	320	7.2	6.4-8.1	144	2.4	2.0-2.8	395	3.8	3.4-4.2	287	6.4	5.7-7.2	108	1.8	1.4-2.1
1991	450	4.3	3.9-4.7	310	6.9	6.1-7.7	140	2.3	2.0-2.8	431	4.1	3.8-4.5	311	7.1	6.3-7.9	120	2.0	1.6-2.3
1992	513	4.9	4.5-5.3	350	7.7	6.9-8.6	163	2.6	2.2-3.1	460	4.4	4.0-4.8	334	7.5	6.7-8.4	126	2.1	1.7-2.5
1993	500	4.7	4.3-5.2	357	7.9	7.1-8.8	143	2.3	1.9-2.7	455	4.3	3.9-4.7	335	7.5	6.7-8.3	120	1.9	1.6-2.3
1994	507	4.8	4.4-5.2	365	8.1	7.3-9.0	142	2.3	1.9-2.7	460	4.3	3.9-4.7	325	7.2	6.4-8.0	135	2.2	1.8-2.6
1995	487	4.6	4.2-5.1	350	7.8	7.0-8.7	137	2.2	1.8-2.6	502	4.8	4.4-5.2	378	8.5	7.6-9.4	124	1.9	1.6-2.3
1996	532	5.0	4.6-5.5	373	8.1	7.3-9.0	159	2.5	2.1-3.0	474	4.5	4.1-4.9	337	7.5	6.7-8.4	137	2.2	1.8-2.6
1997	481	4.5	4.1-4.9	339	7.4	6.6-8.3	142	2.3	1.9-2.7	477	4.4	4.0-4.9	368	8.1	7.3-8.9	109	1.7	1.4-2.0
1998	545	5.1	4.6-5.5	411	8.8	7.9-9.7	134	2.2	1.8-2.6	485	4.5	4.1-4.9	366	7.8	7.1-8.7	119	1.9	1.6-2.3
1999	533	4.9	4.5-5.3	386	8.2	7.4-9.1	147	2.3	1.9-2.7	499	4.6	4.2-5.0	367	7.9	7.1-8.7	132	2.0	1.7-2.4
2000	580	5.3	4.9-5.7	424	9.0	8.1-9.9	156	2.4	2.0-2.8	504	4.6	4.2-5.0	366	7.8	7.0-8.7	138	2.1	1.8-2.5
2001	535	4.8	4.4-5.3	401	8.3	7.5-9.1	134	2.1	1.7-2.4	539	4.8	4.4-5.3	412	8.6	7.8-9.5	127	1.9	1.6-2.3
2002	584	5.2	4.8-5.7	441	9.0	8.2-9.9	143	2.2	1.8-2.5	507	4.5	4.1-4.9	384	7.9	7.1-8.7	123	1.8	1.5-2.2
2003	607	5.4	4.9-5.8	462	9.3	8.5-10.2	145	2.1	1.8-2.5	567	5.0	4.6-5.4	421	8.6	7.8-9.5	146	2.1	1.8-2.5
2004	626	5.5	5.1-5.9	479	9.6	8.8-10.6	147	2.2	1.8-2.6	586	5.1	4.7-5.6	452	9.0	8.2-9.9	134	1.9	1.6-2.3
2005	682	5.9	5.5-6.4	526	10.4	9.5-11.4	156	2.4	2.0-2.8	604	5.2	4.8-5.6	471	9.5	8.6-10.4	133	1.9	1.6-2.3
2006	702	6.1	5.6-6.5	550	10.7	9.8-11.7	152	2.3	2.0-2.7	553	4.8	4.4-5.2	436	8.6	7.8-9.5	117	1.8	1.5-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Esophagus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	663	5.6	5.2-6.1	512	9.8	8.9-10.7	151	2.3	1.9-2.7	585	4.9	4.5-5.4	434	8.4	7.7-9.3	151	2.2	1.9-2.6
2008	702	5.9	5.5-6.4	548	10.4	9.6-11.3	154	2.3	1.9-2.7	575	4.8	4.4-5.2	460	8.7	7.9-9.5	115	1.7	1.4-2.0
2009	735	6.1	5.6-6.5	575	10.6	9.8-11.6	160	2.3	2.0-2.7	567	4.6	4.2-5.0	455	8.4	7.6-9.2	112	1.6	1.3-1.9
2010	678	5.5	5.1-5.9	514	9.3	8.5-10.2	164	2.3	2.0-2.7	585	4.8	4.4-5.2	470	8.7	7.9-9.5	115	1.6	1.3-1.9
2011	651	5.2	4.8-5.6	486	8.6	7.8-9.4	165	2.4	2.0-2.8	623	5.0	4.6-5.4	477	8.6	7.8-9.4	146	2.0	1.7-2.4
2012	639	5.0	4.6-5.5	510	9.0	8.2-9.8	129	1.8	1.5-2.1	584	4.6	4.2-5.0	458	8.2	7.5-9.0	126	1.7	1.4-2.0
2013	682	5.3	4.9-5.7	534	9.1	8.4-10.0	148	2.0	1.7-2.4	589	4.5	4.2-4.9	457	7.9	7.2-8.7	132	1.8	1.5-2.2
2014	685	5.2	4.8-5.6	535	8.8	8.1-9.6	150	2.1	1.7-2.4	576	4.5	4.1-4.9	475	8.3	7.5-9.1	101	1.4	1.1-1.7
2015	745	5.7	5.3-6.1	571	9.4	8.7-10.3	174	2.5	2.1-2.9	559	4.2	3.8-4.5	436	7.3	6.6-8.0	123	1.6	1.4-2.0
2016	735	5.5	5.1-5.9	568	9.3	8.6-10.2	167	2.3	1.9-2.7	581	4.3	4.0-4.7	462	7.6	6.9-8.3	119	1.6	1.3-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Stomach

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	961	11.1	10.4-11.8	554	15.3	14.0-16.7	407	8.1	7.3-8.9	762	8.9	8.3-9.6	418	11.9	10.7-13.2	344	6.8	6.1-7.6
1977	953	10.9	10.2-11.6	562	15.7	14.3-17.1	391	7.6	6.9-8.4	749	8.6	8.0-9.3	443	12.6	11.4-13.9	306	6.0	5.3-6.7
1978	943	10.6	9.9-11.3	530	14.6	13.3-16.0	413	7.9	7.1-8.7	698	8.0	7.4-8.6	390	10.9	9.8-12.1	308	5.9	5.3-6.6
1979	963	10.6	9.9-11.3	572	15.5	14.2-16.9	391	7.3	6.6-8.0	728	8.1	7.5-8.7	420	11.5	10.4-12.7	308	5.7	5.1-6.4
1980	917	10.0	9.4-10.7	554	14.7	13.4-16.1	363	6.8	6.1-7.5	731	8.1	7.5-8.7	419	11.5	10.4-12.8	312	5.8	5.2-6.5
1981	1012	10.9	10.2-11.6	609	16.0	14.7-17.4	403	7.4	6.7-8.2	701	7.6	7.0-8.2	413	11.1	10.0-12.2	288	5.3	4.7-5.9
1982	987	10.4	9.8-11.1	591	15.5	14.2-16.9	396	7.1	6.4-7.8	702	7.5	7.0-8.1	433	11.6	10.5-12.8	269	4.8	4.3-5.5
1983	971	10.2	9.6-10.9	589	15.3	14.0-16.7	382	6.8	6.1-7.5	687	7.3	6.8-7.9	412	10.9	9.8-12.1	275	4.9	4.3-5.5
1984	962	10.0	9.4-10.7	586	14.9	13.6-16.2	376	6.6	5.9-7.3	725	7.5	7.0-8.1	426	10.7	9.7-11.8	299	5.2	4.6-5.8
1985	982	10.1	9.5-10.8	617	15.7	14.4-17.0	365	6.4	5.7-7.1	650	6.8	6.3-7.3	378	10.0	9.0-11.1	272	4.7	4.2-5.3
1986	961	9.8	9.2-10.5	604	15.4	14.1-16.7	357	6.1	5.5-6.8	700	7.2	6.7-7.8	425	10.8	9.8-12.0	275	4.7	4.1-5.3
1987	967	9.8	9.2-10.5	581	14.6	13.4-15.9	386	6.6	5.9-7.3	676	6.9	6.4-7.4	404	10.5	9.4-11.6	272	4.6	4.1-5.2
1988	934	9.5	8.9-10.1	533	13.2	12.1-14.5	401	6.8	6.2-7.6	646	6.6	6.1-7.1	364	9.2	8.3-10.3	282	4.7	4.2-5.3
1989	894	8.9	8.3-9.5	595	14.7	13.5-16.0	299	4.9	4.4-5.5	676	6.8	6.3-7.3	398	10.2	9.2-11.3	278	4.6	4.0-5.1
1990	996	9.9	9.2-10.5	632	15.1	13.9-16.4	364	6.0	5.4-6.7	619	6.2	5.7-6.7	390	9.7	8.7-10.7	229	3.7	3.3-4.3
1991	893	8.7	8.1-9.3	560	13.3	12.2-14.5	333	5.5	4.9-6.1	596	5.8	5.4-6.3	385	9.6	8.6-10.6	211	3.4	3.0-3.9
1992	924	8.9	8.4-9.5	584	13.9	12.8-15.1	340	5.4	4.9-6.0	616	6.0	5.5-6.5	388	9.6	8.6-10.6	228	3.6	3.1-4.1
1993	960	9.2	8.6-9.8	585	13.7	12.6-14.9	375	5.9	5.3-6.5	598	5.7	5.3-6.2	362	8.9	8.0-9.9	236	3.7	3.2-4.2
1994	913	8.7	8.2-9.3	560	13.1	12.0-14.2	353	5.6	5.0-6.2	614	5.9	5.4-6.4	363	8.7	7.8-9.6	251	3.9	3.4-4.4
1995	902	8.5	8.0-9.1	548	12.6	11.5-13.7	354	5.6	5.1-6.3	581	5.5	5.1-6.0	330	7.9	7.1-8.8	251	3.9	3.4-4.4
1996	897	8.4	7.9-9.0	552	12.7	11.6-13.8	345	5.3	4.8-5.9	551	5.2	4.7-5.6	320	7.5	6.7-8.4	231	3.5	3.1-4.0
1997	925	8.6	8.0-9.2	563	12.6	11.6-13.7	362	5.6	5.0-6.2	521	4.8	4.4-5.3	292	6.7	5.9-7.5	229	3.5	3.0-4.0
1998	930	8.6	8.0-9.1	595	13.0	11.9-14.1	335	5.2	4.6-5.8	505	4.6	4.2-5.0	287	6.4	5.7-7.3	218	3.3	2.9-3.8
1999	884	8.1	7.6-8.6	560	12.3	11.3-13.3	324	5.0	4.5-5.6	494	4.5	4.1-4.9	300	6.6	5.9-7.4	194	2.9	2.5-3.4
2000	902	8.1	7.6-8.6	561	12.2	11.2-13.3	341	5.0	4.5-5.6	498	4.4	4.1-4.9	283	6.4	5.6-7.1	215	3.2	2.8-3.6
2001	860	7.7	7.2-8.3	550	11.7	10.8-12.8	310	4.8	4.3-5.4	473	4.2	3.8-4.6	271	5.9	5.2-6.7	202	3.0	2.6-3.4
2002	856	7.6	7.1-8.1	525	11.1	10.2-12.1	331	5.0	4.5-5.6	441	3.9	3.5-4.2	255	5.5	4.9-6.3	186	2.7	2.3-3.1
2003	902	7.9	7.4-8.5	564	11.7	10.7-12.7	338	5.1	4.6-5.7	425	3.7	3.3-4.0	243	5.2	4.5-5.9	182	2.7	2.3-3.1
2004	859	7.5	7.0-8.0	521	10.8	9.9-11.8	338	5.1	4.5-5.7	470	4.0	3.7-4.4	277	5.9	5.2-6.6	193	2.8	2.4-3.2
2005	800	6.9	6.4-7.4	495	9.9	9.0-10.8	305	4.6	4.1-5.1	419	3.6	3.2-3.9	240	5.0	4.3-5.6	179	2.6	2.2-3.0
2006	799	6.9	6.4-7.4	492	9.8	8.9-10.7	307	4.7	4.1-5.2	371	3.1	2.8-3.5	212	4.2	3.7-4.9	159	2.3	2.0-2.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Stomach

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	837	7.2	6.7-7.7	544	10.6	9.7-11.5	293	4.5	4.0-5.0	367	3.1	2.8-3.5	218	4.3	3.7-4.9	149	2.2	1.9-2.6
2008	775	6.5	6.0-7.0	505	9.8	9.0-10.7	270	4.0	3.6-4.6	394	3.2	2.9-3.5	240	4.7	4.1-5.3	154	2.1	1.8-2.5
2009	748	6.3	5.8-6.7	482	9.2	8.4-10.1	266	3.8	3.4-4.3	349	2.9	2.6-3.2	216	4.2	3.7-4.8	133	1.9	1.6-2.2
2010	819	6.8	6.3-7.3	530	9.9	9.0-10.8	289	4.3	3.8-4.8	340	2.8	2.5-3.1	204	3.9	3.4-4.5	136	1.9	1.6-2.3
2011	882	7.2	6.7-7.7	554	10.2	9.4-11.1	328	4.8	4.3-5.4	320	2.5	2.3-2.8	194	3.6	3.1-4.2	126	1.7	1.4-2.1
2012	845	6.8	6.4-7.3	532	9.5	8.7-10.4	313	4.7	4.1-5.2	355	2.8	2.5-3.1	212	3.9	3.4-4.5	143	2.0	1.7-2.4
2013	922	7.4	6.9-7.9	588	10.5	9.6-11.4	334	4.9	4.4-5.5	354	2.7	2.4-3.0	214	3.8	3.3-4.4	140	1.9	1.6-2.3
2014	842	6.7	6.2-7.1	564	9.9	9.1-10.8	278	4.1	3.6-4.6	338	2.7	2.4-3.0	215	3.8	3.3-4.4	123	1.8	1.5-2.1
2015	842	6.6	6.1-7.0	538	9.3	8.5-10.1	304	4.4	3.9-4.9	303	2.4	2.1-2.7	187	3.3	2.8-3.8	116	1.7	1.4-2.0
2016	874	6.8	6.4-7.3	540	9.2	8.4-10.0	334	4.9	4.4-5.5	286	2.2	1.9-2.4	177	3.0	2.6-3.5	109	1.5	1.2-1.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Colorectal

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	5898	67.1	65.4-68.9	2898	80.1	77.0-83.2	3000	58.7	56.6-60.8	3006	34.9	33.6-36.2	1433	41.2	38.9-43.5	1573	30.9	29.4-32.5
1977	6030	67.4	65.6-69.1	2933	79.8	76.8-82.9	3097	59.4	57.3-61.6	3015	34.7	33.5-36.0	1438	41.3	39.0-43.6	1577	30.5	29.0-32.1
1978	6132	68.1	66.4-69.9	2992	80.9	77.9-84.1	3140	59.6	57.5-61.7	3198	36.1	34.9-37.4	1550	43.6	41.4-46.0	1648	31.4	29.8-32.9
1979	6188	67.7	66.0-69.4	2960	79.0	76.0-82.0	3228	60.1	58.0-62.2	3152	35.1	33.9-36.4	1510	42.7	40.5-45.1	1642	30.6	29.1-32.1
1980	6419	69.0	67.3-70.7	3220	84.0	81.0-87.1	3199	58.6	56.6-60.7	3115	34.2	33.0-35.4	1511	41.5	39.3-43.8	1604	29.5	28.0-31.0
1981	6595	70.4	68.7-72.1	3312	87.4	84.3-90.6	3283	59.4	57.3-61.5	3164	34.4	33.2-35.6	1543	42.1	39.9-44.4	1621	29.3	27.9-30.8
1982	6564	69.2	67.5-70.9	3256	83.9	80.9-87.0	3308	59.1	57.1-61.2	3181	34.2	33.0-35.4	1565	42.8	40.6-45.1	1616	28.8	27.4-30.2
1983	6713	70.2	68.5-71.9	3360	86.4	83.4-89.5	3353	59.4	57.4-61.5	3125	33.1	31.9-34.3	1526	41.3	39.1-43.5	1599	28.0	26.7-29.5
1984	6794	70.5	68.8-72.2	3318	84.3	81.4-87.4	3476	61.0	59.0-63.1	3260	34.3	33.1-35.5	1609	42.9	40.8-45.2	1651	28.9	27.5-30.3
1985	7037	72.4	70.7-74.1	3539	89.1	86.1-92.3	3498	61.0	59.0-63.1	3174	33.2	32.0-34.4	1541	40.9	38.8-43.1	1633	28.2	26.8-29.6
1986	6775	69.2	67.5-70.9	3340	84.1	81.1-87.1	3435	59.4	57.4-61.4	3168	32.8	31.6-33.9	1566	41.4	39.2-43.6	1602	27.4	26.1-28.8
1987	6724	68.1	66.5-69.8	3337	83.6	80.7-86.7	3387	58.1	56.1-60.1	3154	32.3	31.2-33.5	1578	41.3	39.2-43.4	1576	26.6	25.3-28.0
1988	6478	64.8	63.2-66.4	3197	78.7	75.8-81.6	3281	55.5	53.6-57.5	3135	31.8	30.7-32.9	1570	40.1	38.1-42.3	1565	26.3	25.0-27.7
1989	6434	64.0	62.4-65.6	3231	79.2	76.3-82.1	3203	53.9	52.0-55.8	2908	29.1	28.1-30.2	1406	35.7	33.8-37.7	1502	24.9	23.7-26.2
1990	6418	62.9	61.3-64.4	3174	76.2	73.5-79.0	3244	53.6	51.8-55.5	2818	27.9	26.8-28.9	1437	36.2	34.3-38.2	1381	22.6	21.4-23.8
1991	6374	62.0	60.5-63.6	3173	75.5	72.8-78.3	3201	52.7	50.9-54.6	2862	28.0	27.0-29.1	1383	34.3	32.4-36.2	1479	23.9	22.7-25.2
1992	6272	60.4	58.9-61.9	3173	74.8	72.1-77.6	3099	50.4	48.6-52.2	2725	26.3	25.4-27.4	1320	32.4	30.6-34.3	1405	22.5	21.3-23.7
1993	6351	60.5	59.0-62.0	3089	71.7	69.1-74.3	3262	52.6	50.8-54.4	2788	26.6	25.6-27.6	1356	33.0	31.2-34.9	1432	22.5	21.3-23.7
1994	6484	61.4	59.9-62.9	3205	74.5	71.8-77.1	3279	52.7	50.9-54.6	2840	26.9	26.0-28.0	1349	32.8	31.0-34.7	1491	23.2	22.0-24.4
1995	6410	60.3	58.8-61.8	3184	73.0	70.5-75.7	3226	51.3	49.5-53.1	2789	26.1	25.2-27.1	1332	32.1	30.3-33.9	1457	22.5	21.4-23.7
1996	6297	58.9	57.4-60.4	3180	72.5	69.9-75.1	3117	49.3	47.6-51.1	2649	24.6	23.7-25.6	1316	31.1	29.4-32.8	1333	20.4	19.3-21.6
1997	6633	61.5	60.0-63.0	3306	73.7	71.2-76.3	3327	52.5	50.7-54.3	2516	23.2	22.3-24.1	1248	28.9	27.3-30.6	1268	19.3	18.2-20.4
1998	6959	63.9	62.4-65.4	3447	75.8	73.2-78.4	3512	55.1	53.3-57.0	2518	22.9	22.0-23.8	1217	27.5	25.9-29.1	1301	19.6	18.5-20.7
1999	7015	63.7	62.3-65.3	3491	76.2	73.7-78.8	3524	54.7	52.9-56.6	2478	22.3	21.4-23.2	1216	27.4	25.9-29.0	1262	18.7	17.7-19.8
2000	7010	63.1	61.7-64.6	3480	75.0	72.5-77.5	3530	54.4	52.6-56.2	2497	22.2	21.3-23.0	1256	28.0	26.5-29.6	1241	18.0	17.0-19.0
2001	6674	59.7	58.2-61.1	3299	70.2	67.8-72.7	3375	51.7	49.9-53.5	2411	21.2	20.4-22.1	1138	25.1	23.7-26.7	1273	18.6	17.5-19.6
2002	6613	58.6	57.2-60.0	3300	69.0	66.7-71.4	3313	50.4	48.7-52.2	2412	21.0	20.2-21.9	1174	25.4	24.0-26.9	1238	17.9	16.9-19.0
2003	6313	55.2	53.9-56.6	3067	63.3	61.1-65.6	3246	49.0	47.3-50.8	2198	18.9	18.1-19.7	1083	23.1	21.7-24.5	1115	16.0	15.1-17.0
2004	6192	53.8	52.4-55.1	2973	60.7	58.5-62.9	3219	48.5	46.9-50.3	2146	18.3	17.5-19.0	1030	21.4	20.1-22.8	1116	15.7	14.7-16.6
2005	5869	50.5	49.2-51.8	2947	59.5	57.3-61.7	2922	43.8	42.2-45.5	1974	16.6	15.9-17.4	962	20.1	18.8-21.4	1012	14.2	13.3-15.1
2006	5594	47.9	46.7-49.2	2770	55.3	53.2-57.4	2824	42.2	40.6-43.8	1930	16.2	15.4-16.9	921	19.0	17.8-20.3	1009	14.1	13.2-15.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Colorectal

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	5726	48.6	47.3-49.9	2821	55.2	53.2-57.3	2905	43.4	41.8-45.1	1922	15.9	15.2-16.6	944	18.9	17.7-20.2	978	13.6	12.8-14.5
2008	5563	46.6	45.3-47.8	2739	52.9	50.9-55.0	2824	41.8	40.2-43.4	1890	15.4	14.7-16.2	911	18.0	16.8-19.2	979	13.4	12.6-14.3
2009	5204	43.2	42.0-44.4	2589	49.3	47.4-51.2	2615	38.2	36.7-39.8	1821	14.8	14.1-15.5	896	17.5	16.4-18.7	925	12.8	12.0-13.7
2010	5293	43.5	42.3-44.7	2586	48.4	46.5-50.3	2707	39.5	38.0-41.1	1800	14.3	13.6-15.0	878	16.7	15.6-17.8	922	12.4	11.6-13.3
2011	5110	41.9	40.8-43.1	2510	46.6	44.8-48.5	2600	38.1	36.6-39.6	1775	13.9	13.2-14.5	873	16.2	15.1-17.3	902	12.0	11.2-12.8
2012	4898	39.7	38.6-40.9	2521	45.8	44.0-47.7	2377	34.8	33.3-36.3	1735	13.5	12.8-14.1	839	15.5	14.4-16.6	896	11.9	11.1-12.8
2013	4921	39.5	38.4-40.7	2538	45.4	43.6-47.3	2383	34.8	33.4-36.3	1714	13.3	12.6-13.9	907	16.5	15.4-17.6	807	10.6	9.9-11.4
2014	4856	38.5	37.4-39.6	2489	44.0	42.2-45.8	2367	34.0	32.6-35.5	1667	12.7	12.0-13.3	854	15.2	14.2-16.3	813	10.7	9.9-11.5
2015	4885	38.8	37.7-40.0	2467	43.2	41.5-45.0	2418	35.1	33.6-36.6	1673	12.6	12.0-13.2	859	15.1	14.1-16.2	814	10.7	9.9-11.5
2016	4918	38.8	37.7-40.0	2509	43.4	41.7-45.2	2409	34.8	33.4-36.3	1490	11.2	10.6-11.8	724	12.5	11.6-13.5	766	10.1	9.4-10.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Colon excluding rectum

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	4110	47.2	45.7-48.6	1907	53.6	51.1-56.2	2203	43.1	41.3-45.0	2355	27.4	26.2-28.5	1057	30.4	28.5-32.4	1298	25.5	24.2-27.0
1977	4208	47.5	46.1-49.0	1937	54.2	51.7-56.8	2271	43.6	41.9-45.5	2442	28.2	27.1-29.4	1132	32.6	30.6-34.6	1310	25.4	24.0-26.8
1978	4310	48.2	46.8-49.7	1990	54.9	52.4-57.5	2320	44.0	42.3-45.9	2563	29.0	27.8-30.1	1186	33.5	31.5-35.5	1377	26.2	24.8-27.6
1979	4390	48.5	47.0-50.0	1996	54.4	51.9-57.0	2394	44.7	42.9-46.5	2597	28.9	27.8-30.0	1211	34.2	32.2-36.3	1386	25.8	24.4-27.2
1980	4559	49.3	47.9-50.8	2203	58.3	55.7-60.9	2356	43.3	41.5-45.1	2542	27.9	26.8-29.0	1212	33.4	31.4-35.4	1330	24.4	23.1-25.8
1981	4725	50.7	49.3-52.2	2274	60.7	58.1-63.4	2451	44.5	42.7-46.3	2663	29.0	27.9-30.2	1279	35.1	33.1-37.2	1384	25.1	23.8-26.5
1982	4618	49.1	47.6-50.5	2165	56.7	54.2-59.3	2453	43.8	42.1-45.6	2625	28.2	27.1-29.3	1269	34.7	32.7-36.8	1356	24.2	22.9-25.5
1983	4729	49.8	48.4-51.3	2233	58.7	56.2-61.4	2496	44.3	42.6-46.1	2618	27.7	26.7-28.8	1251	34.0	32.0-36.0	1367	24.0	22.7-25.3
1984	4883	51.0	49.5-52.4	2232	57.6	55.2-60.2	2651	46.4	44.7-48.3	2735	28.8	27.7-29.9	1321	35.2	33.2-37.2	1414	24.7	23.4-26.0
1985	5021	51.8	50.4-53.3	2414	61.5	59.0-64.1	2607	45.3	43.6-47.1	2665	27.8	26.8-28.9	1269	33.6	31.7-35.6	1396	24.1	22.8-25.4
1986	4894	50.2	48.8-51.6	2295	58.6	56.1-61.1	2599	44.9	43.2-46.7	2689	27.8	26.7-28.9	1305	34.6	32.7-36.7	1384	23.7	22.4-25.0
1987	4770	48.5	47.1-49.9	2272	57.8	55.3-60.3	2498	42.7	41.0-44.4	2661	27.3	26.2-28.3	1322	34.6	32.6-36.6	1339	22.7	21.5-23.9
1988	4638	46.5	45.2-47.9	2155	53.6	51.3-56.0	2483	41.8	40.2-43.5	2695	27.3	26.3-28.4	1330	34.0	32.1-35.9	1365	22.9	21.7-24.2
1989	4543	45.4	44.1-46.7	2199	55.1	52.7-57.5	2344	39.3	37.7-40.9	2468	24.7	23.7-25.7	1162	29.5	27.8-31.3	1306	21.7	20.5-22.9
1990	4532	44.5	43.2-45.8	2138	52.3	50.0-54.6	2394	39.5	37.9-41.1	2447	24.2	23.2-25.2	1252	31.7	29.9-33.6	1195	19.5	18.4-20.7
1991	4592	44.7	43.4-46.0	2197	53.0	50.7-55.3	2395	39.2	37.6-40.8	2416	23.7	22.8-24.7	1151	28.6	26.9-30.4	1265	20.4	19.3-21.6
1992	4476	43.2	41.9-44.5	2133	50.9	48.7-53.2	2343	38.0	36.4-39.6	2312	22.4	21.5-23.3	1088	26.9	25.3-28.6	1224	19.6	18.5-20.7
1993	4585	43.7	42.5-45.0	2113	49.6	47.5-51.9	2472	39.6	38.0-41.2	2390	22.8	21.9-23.8	1155	28.3	26.6-30.0	1235	19.4	18.3-20.6
1994	4702	44.5	43.3-45.8	2239	52.6	50.4-54.9	2463	39.4	37.8-41.0	2437	23.1	22.2-24.0	1132	27.7	26.1-29.4	1305	20.3	19.2-21.4
1995	4644	43.5	42.3-44.8	2219	51.3	49.1-53.5	2425	38.2	36.7-39.8	2355	22.1	21.2-23.0	1096	26.6	25.0-28.2	1259	19.4	18.3-20.5
1996	4498	41.9	40.7-43.2	2161	50.0	47.8-52.2	2337	36.7	35.2-38.2	2245	20.9	20.0-21.8	1101	26.1	24.6-27.7	1144	17.5	16.5-18.5
1997	4883	45.2	43.9-46.5	2307	51.8	49.7-54.0	2576	40.3	38.8-41.9	2175	20.1	19.2-20.9	1051	24.4	22.9-25.9	1124	17.0	16.0-18.1
1998	5067	46.4	45.1-47.7	2423	53.8	51.7-56.1	2644	41.3	39.7-42.9	2131	19.4	18.6-20.2	1013	22.9	21.5-24.4	1118	16.7	15.8-17.8
1999	5056	45.8	44.5-47.0	2394	52.6	50.5-54.8	2662	40.8	39.3-42.4	2073	18.6	17.8-19.4	997	22.5	21.2-24.0	1076	15.9	14.9-16.9
2000	5096	45.7	44.4-47.0	2401	52.2	50.1-54.4	2695	41.0	39.5-42.6	2106	18.6	17.9-19.5	1039	23.3	21.9-24.8	1067	15.4	14.5-16.4
2001	4840	43.1	41.9-44.4	2293	49.1	47.1-51.2	2547	38.6	37.1-40.1	2040	17.9	17.2-18.7	945	20.9	19.6-22.3	1095	15.9	15.0-16.9
2002	4789	42.3	41.1-43.5	2292	48.3	46.3-50.3	2497	37.7	36.2-39.2	1991	17.3	16.6-18.1	954	20.7	19.4-22.1	1037	14.9	14.0-15.9
2003	4579	39.9	38.7-41.1	2123	44.2	42.3-46.1	2456	36.7	35.3-38.2	1860	16.0	15.3-16.7	894	19.1	17.9-20.4	966	13.8	13.0-14.8
2004	4445	38.4	37.3-39.6	2019	41.6	39.8-43.5	2426	36.2	34.7-37.7	1773	15.0	14.3-15.8	833	17.4	16.2-18.6	940	13.1	12.3-14.0
2005	4283	36.7	35.6-37.9	2064	41.9	40.1-43.8	2219	33.0	31.6-34.4	1623	13.6	13.0-14.3	771	16.2	15.0-17.4	852	11.9	11.1-12.8
2006	4116	35.1	34.0-36.2	1965	39.6	37.8-41.4	2151	31.6	30.2-33.0	1602	13.4	12.7-14.1	750	15.5	14.4-16.7	852	11.8	11.0-12.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Colon excluding rectum

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	4192	35.5	34.4-36.6	1976	39.0	37.3-40.8	2216	32.6	31.2-34.0	1618	13.3	12.7-14.0	772	15.6	14.5-16.7	846	11.7	10.9-12.6
2008	4037	33.7	32.6-34.8	1882	36.6	34.9-38.3	2155	31.7	30.3-33.1	1540	12.6	11.9-13.2	728	14.4	13.4-15.5	812	11.1	10.3-11.9
2009	3711	30.6	29.6-31.6	1742	33.3	31.7-34.9	1969	28.3	27.1-29.7	1498	12.2	11.6-12.8	718	14.1	13.1-15.2	780	10.8	10.0-11.6
2010	3803	31.2	30.2-32.2	1763	33.4	31.9-35.1	2040	29.3	28.0-30.6	1466	11.6	11.0-12.2	691	13.2	12.2-14.2	775	10.3	9.6-11.1
2011	3644	29.7	28.7-30.7	1685	31.6	30.1-33.2	1959	28.1	26.8-29.4	1452	11.3	10.7-11.9	686	12.8	11.8-13.8	766	10.1	9.4-10.9
2012	3530	28.5	27.5-29.4	1707	31.2	29.7-32.8	1823	26.2	25.0-27.5	1426	11.1	10.5-11.7	660	12.2	11.3-13.2	766	10.2	9.4-11.0
2013	3458	27.5	26.6-28.5	1700	30.6	29.1-32.1	1758	25.1	23.9-26.4	1388	10.7	10.1-11.3	712	13.0	12.0-14.0	676	8.8	8.2-9.6
2014	3433	27.1	26.1-28.0	1651	29.4	28.0-30.9	1782	25.2	24.0-26.5	1334	10.1	9.6-10.7	667	12.0	11.1-12.9	667	8.7	8.0-9.4
2015	3479	27.7	26.7-28.6	1648	29.4	27.9-30.9	1831	26.2	25.0-27.5	1361	10.2	9.7-10.8	673	11.8	10.9-12.8	688	8.9	8.3-9.7
2016	3535	27.8	26.8-28.7	1696	29.5	28.1-31.0	1839	26.2	25.0-27.5	1160	8.7	8.2-9.2	539	9.3	8.5-10.2	621	8.1	7.4-8.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Rectum & rectosigmoid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1788	19.9	19.0-20.9	991	26.5	24.8-28.3	797	15.5	14.5-16.7	651	7.5	6.9-8.1	376	10.8	9.6-12.0	275	5.4	4.8-6.1
1977	1822	19.9	18.9-20.8	996	25.6	24.0-27.3	826	15.8	14.7-16.9	573	6.5	6.0-7.1	306	8.7	7.7-9.8	267	5.1	4.5-5.8
1978	1822	19.9	19.0-20.8	1002	26.1	24.4-27.8	820	15.5	14.5-16.6	635	7.2	6.6-7.8	364	10.2	9.1-11.4	271	5.2	4.6-5.8
1979	1798	19.2	18.3-20.1	964	24.5	22.9-26.2	834	15.4	14.4-16.5	555	6.2	5.7-6.8	299	8.6	7.6-9.6	256	4.8	4.2-5.4
1980	1860	19.6	18.7-20.6	1017	25.8	24.1-27.5	843	15.4	14.3-16.4	573	6.3	5.8-6.8	299	8.2	7.2-9.2	274	5.1	4.5-5.7
1981	1870	19.7	18.8-20.6	1038	26.7	25.0-28.5	832	14.9	13.9-16.0	501	5.3	4.9-5.8	264	7.0	6.1-7.9	237	4.2	3.7-4.8
1982	1946	20.1	19.2-21.0	1091	27.2	25.5-28.9	855	15.3	14.3-16.4	556	6.0	5.5-6.5	296	8.1	7.2-9.2	260	4.6	4.1-5.2
1983	1984	20.4	19.5-21.3	1127	27.7	26.0-29.4	857	15.1	14.1-16.2	507	5.4	4.9-5.9	275	7.3	6.4-8.3	232	4.1	3.6-4.6
1984	1911	19.6	18.7-20.5	1086	26.7	25.1-28.4	825	14.6	13.6-15.6	525	5.5	5.1-6.0	288	7.7	6.8-8.7	237	4.2	3.6-4.7
1985	2016	20.5	19.6-21.4	1125	27.6	26.0-29.4	891	15.7	14.7-16.8	509	5.3	4.9-5.8	272	7.3	6.5-8.3	237	4.1	3.6-4.6
1986	1881	19.0	18.1-19.9	1045	25.5	23.9-27.2	836	14.5	13.5-15.5	479	5.0	4.5-5.4	261	6.7	5.9-7.6	218	3.8	3.3-4.3
1987	1954	19.6	18.8-20.6	1065	25.9	24.3-27.5	889	15.4	14.4-16.5	493	5.1	4.6-5.5	256	6.7	5.9-7.6	237	4.0	3.5-4.5
1988	1840	18.3	17.5-19.2	1042	25.0	23.5-26.7	798	13.7	12.7-14.7	440	4.5	4.1-4.9	240	6.1	5.4-7.0	200	3.4	2.9-3.9
1989	1891	18.6	17.8-19.5	1032	24.1	22.6-25.7	859	14.6	13.6-15.6	440	4.4	4.0-4.9	244	6.2	5.4-7.0	196	3.3	2.8-3.7
1990	1886	18.4	17.5-19.2	1036	23.9	22.5-25.5	850	14.2	13.2-15.2	371	3.7	3.3-4.1	185	4.5	3.8-5.2	186	3.1	2.6-3.5
1991	1782	17.3	16.5-18.1	976	22.5	21.1-24.1	806	13.5	12.6-14.5	446	4.3	3.9-4.7	232	5.7	4.9-6.5	214	3.5	3.0-4.0
1992	1796	17.2	16.4-18.0	1040	23.9	22.4-25.4	756	12.4	11.6-13.4	413	4.0	3.6-4.4	232	5.5	4.8-6.3	181	2.9	2.5-3.4
1993	1766	16.8	16.0-17.6	976	22.0	20.6-23.5	790	13.0	12.1-13.9	398	3.8	3.4-4.2	201	4.7	4.1-5.4	197	3.1	2.7-3.5
1994	1782	16.9	16.1-17.7	966	21.8	20.4-23.3	816	13.4	12.4-14.3	403	3.9	3.5-4.3	217	5.1	4.5-5.9	186	2.9	2.5-3.4
1995	1766	16.7	16.0-17.5	965	21.8	20.4-23.2	801	13.1	12.2-14.1	434	4.1	3.7-4.5	236	5.5	4.8-6.3	198	3.1	2.7-3.6
1996	1799	17.0	16.2-17.8	1019	22.5	21.1-24.0	780	12.7	11.8-13.6	404	3.8	3.4-4.2	215	5.0	4.3-5.7	189	2.9	2.5-3.4
1997	1750	16.3	15.6-17.1	999	21.9	20.6-23.3	751	12.1	11.3-13.0	341	3.2	2.8-3.5	197	4.6	3.9-5.3	144	2.2	1.9-2.6
1998	1892	17.5	16.7-18.3	1024	21.9	20.6-23.4	868	13.9	13.0-14.9	387	3.5	3.2-3.9	204	4.5	3.9-5.2	183	2.8	2.4-3.3
1999	1959	18.0	17.2-18.8	1097	23.6	22.2-25.1	862	13.9	12.9-14.8	405	3.7	3.3-4.0	219	4.9	4.3-5.6	186	2.8	2.4-3.3
2000	1914	17.4	16.7-18.2	1079	22.7	21.4-24.2	835	13.3	12.4-14.3	391	3.5	3.2-3.9	217	4.7	4.1-5.4	174	2.6	2.2-3.0
2001	1834	16.6	15.8-17.3	1006	21.1	19.8-22.4	828	13.1	12.2-14.1	371	3.3	3.0-3.6	193	4.2	3.6-4.9	178	2.6	2.3-3.1
2002	1824	16.3	15.6-17.1	1008	20.7	19.5-22.1	816	12.7	11.8-13.6	421	3.7	3.4-4.1	220	4.7	4.1-5.4	201	3.0	2.6-3.5
2003	1734	15.3	14.6-16.1	944	19.1	17.9-20.4	790	12.3	11.4-13.2	338	2.9	2.6-3.3	189	4.0	3.4-4.6	149	2.2	1.9-2.6
2004	1747	15.3	14.6-16.1	954	19.0	17.8-20.3	793	12.4	11.5-13.3	373	3.2	2.9-3.6	197	4.0	3.5-4.6	176	2.5	2.1-2.9
2005	1586	13.7	13.1-14.4	883	17.5	16.4-18.7	703	10.9	10.1-11.7	351	3.0	2.7-3.3	191	3.9	3.4-4.5	160	2.3	1.9-2.7
2006	1478	12.9	12.2-13.6	805	15.7	14.6-16.8	673	10.6	9.8-11.4	328	2.8	2.5-3.1	171	3.5	3.0-4.0	157	2.3	1.9-2.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Rectum & rectosigmoid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	1534	13.1	12.5-13.8	845	16.2	15.1-17.3	689	10.8	10.0-11.7	304	2.6	2.3-2.9	172	3.4	2.9-3.9	132	1.9	1.6-2.3
2008	1526	12.9	12.2-13.6	857	16.4	15.3-17.5	669	10.1	9.3-10.9	350	2.9	2.6-3.2	183	3.6	3.1-4.1	167	2.3	2.0-2.7
2009	1493	12.6	12.0-13.3	847	16.0	14.9-17.1	646	9.9	9.1-10.7	323	2.6	2.3-2.9	178	3.4	2.9-3.9	145	2.0	1.7-2.4
2010	1490	12.4	11.7-13.0	823	15.0	13.9-16.0	667	10.2	9.4-11.1	334	2.7	2.4-3.0	187	3.5	3.0-4.0	147	2.1	1.8-2.5
2011	1466	12.2	11.6-12.9	825	15.0	14.0-16.1	641	10.0	9.2-10.8	323	2.6	2.3-2.9	187	3.4	2.9-3.9	136	1.9	1.6-2.3
2012	1368	11.3	10.7-11.9	814	14.6	13.6-15.7	554	8.6	7.8-9.3	309	2.4	2.2-2.7	179	3.3	2.8-3.8	130	1.8	1.5-2.1
2013	1463	12.0	11.4-12.7	838	14.8	13.8-15.9	625	9.7	8.9-10.5	326	2.6	2.3-2.9	195	3.5	3.0-4.0	131	1.8	1.5-2.2
2014	1423	11.5	10.8-12.1	838	14.5	13.6-15.6	585	8.8	8.1-9.6	333	2.5	2.3-2.8	187	3.2	2.8-3.7	146	2.0	1.7-2.4
2015	1406	11.2	10.6-11.8	819	13.8	12.9-14.9	587	8.9	8.1-9.7	312	2.4	2.1-2.7	186	3.3	2.8-3.8	126	1.7	1.4-2.1
2016	1383	11.1	10.5-11.7	813	13.9	13.0-15.0	570	8.6	7.9-9.4	330	2.5	2.2-2.8	185	3.2	2.7-3.7	145	2.0	1.7-2.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Liver / intrahepatic bile duct

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	235	2.6	2.3-3.0	138	3.6	3.0-4.3	97	1.9	1.5-2.3	176	2.0	1.7-2.3	107	2.9	2.3-3.5	69	1.3	1.0-1.7
1977	278	2.9	2.6-3.3	186	4.6	3.9-5.4	92	1.7	1.4-2.1	237	2.5	2.2-2.9	139	3.4	2.8-4.1	98	1.9	1.5-2.3
1978	265	2.9	2.5-3.2	152	3.9	3.3-4.6	113	2.1	1.7-2.5	236	2.5	2.2-2.9	138	3.5	2.9-4.2	98	1.8	1.5-2.2
1979	252	2.7	2.4-3.1	132	3.3	2.7-3.9	120	2.3	1.9-2.8	199	2.2	1.9-2.5	108	2.7	2.2-3.3	91	1.7	1.4-2.1
1980	248	2.7	2.3-3.0	149	4.0	3.4-4.7	99	1.8	1.5-2.2	217	2.3	2.0-2.6	114	3.0	2.5-3.6	103	1.9	1.5-2.3
1981	254	2.7	2.4-3.1	148	3.9	3.2-4.6	106	1.9	1.6-2.3	236	2.5	2.2-2.9	131	3.5	2.9-4.2	105	1.9	1.6-2.3
1982	254	2.6	2.3-3.0	153	3.9	3.3-4.6	101	1.8	1.5-2.2	220	2.3	2.0-2.7	140	3.7	3.1-4.4	80	1.4	1.1-1.8
1983	276	2.9	2.5-3.2	186	4.5	3.9-5.3	90	1.7	1.3-2.1	231	2.4	2.1-2.7	153	3.9	3.2-4.6	78	1.4	1.1-1.7
1984	265	2.7	2.4-3.1	152	3.8	3.2-4.5	113	2.0	1.7-2.4	233	2.4	2.1-2.8	136	3.4	2.8-4.1	97	1.7	1.4-2.1
1985	255	2.7	2.3-3.0	161	4.1	3.5-4.8	94	1.7	1.4-2.1	239	2.5	2.2-2.8	149	3.8	3.2-4.5	90	1.6	1.3-2.0
1986	265	2.7	2.4-3.1	169	4.1	3.5-4.8	96	1.7	1.4-2.1	253	2.6	2.3-2.9	156	3.8	3.2-4.5	97	1.7	1.4-2.1
1987	281	2.8	2.5-3.2	176	4.2	3.6-4.9	105	1.9	1.5-2.3	281	2.9	2.5-3.2	168	4.2	3.6-5.0	113	2.0	1.6-2.4
1988	330	3.3	3.0-3.7	212	5.1	4.4-5.9	118	2.0	1.7-2.4	281	2.8	2.5-3.2	179	4.3	3.7-5.1	102	1.7	1.4-2.1
1989	311	3.1	2.7-3.4	196	4.5	3.9-5.2	115	1.9	1.6-2.3	257	2.5	2.2-2.9	156	3.7	3.1-4.4	101	1.7	1.4-2.1
1990	348	3.4	3.1-3.8	222	5.2	4.5-6.0	126	2.1	1.7-2.5	312	3.1	2.7-3.4	194	4.8	4.1-5.6	118	1.9	1.6-2.3
1991	342	3.3	3.0-3.7	209	4.7	4.1-5.4	133	2.2	1.9-2.6	310	3.0	2.6-3.3	188	4.3	3.7-5.0	122	2.0	1.7-2.4
1992	390	3.8	3.4-4.2	227	5.5	4.8-6.3	163	2.7	2.3-3.2	324	3.1	2.8-3.5	195	4.7	4.0-5.4	129	2.1	1.8-2.5
1993	412	3.9	3.5-4.3	258	6.0	5.3-6.8	154	2.5	2.1-2.9	401	3.8	3.4-4.2	240	5.6	4.9-6.4	161	2.5	2.2-3.0
1994	378	3.6	3.2-4.0	245	5.7	5.0-6.4	133	2.1	1.8-2.5	363	3.4	3.1-3.8	210	5.0	4.3-5.7	153	2.5	2.1-2.9
1995	442	4.2	3.8-4.6	285	6.4	5.7-7.2	157	2.5	2.1-3.0	398	3.7	3.4-4.1	251	5.8	5.1-6.6	147	2.3	1.9-2.7
1996	399	3.7	3.4-4.1	244	5.5	4.8-6.2	155	2.5	2.1-2.9	380	3.5	3.2-3.9	224	5.1	4.4-5.8	156	2.4	2.0-2.8
1997	463	4.4	4.0-4.8	300	6.5	5.8-7.3	163	2.6	2.2-3.1	386	3.6	3.3-4.0	243	5.4	4.7-6.1	143	2.2	1.9-2.6
1998	498	4.6	4.2-5.0	335	7.2	6.5-8.1	163	2.6	2.2-3.0	369	3.4	3.1-3.8	219	4.8	4.2-5.4	150	2.3	2.0-2.8
1999	488	4.5	4.1-4.9	319	6.9	6.1-7.7	169	2.7	2.3-3.1	413	3.8	3.4-4.2	250	5.4	4.7-6.1	163	2.5	2.1-2.9
2000	486	4.4	4.0-4.8	308	6.5	5.8-7.3	178	2.8	2.4-3.3	406	3.7	3.3-4.0	246	5.3	4.6-6.0	160	2.4	2.0-2.8
2001	508	4.6	4.2-5.0	359	7.5	6.8-8.4	149	2.3	2.0-2.7	414	3.7	3.4-4.1	269	5.7	5.1-6.5	145	2.2	1.8-2.6
2002	524	4.7	4.3-5.1	368	7.6	6.8-8.4	156	2.4	2.0-2.8	469	4.1	3.8-4.5	305	6.4	5.7-7.2	164	2.4	2.1-2.8
2003	531	4.7	4.3-5.2	340	6.8	6.1-7.6	191	3.0	2.6-3.5	411	3.6	3.3-4.0	261	5.4	4.7-6.0	150	2.2	1.9-2.6
2004	579	5.1	4.7-5.5	417	8.2	7.4-9.0	162	2.5	2.1-2.9	466	4.1	3.7-4.4	288	5.8	5.1-6.5	178	2.6	2.2-3.0
2005	611	5.4	4.9-5.8	425	8.2	7.5-9.1	186	2.9	2.5-3.4	481	4.1	3.8-4.5	313	6.2	5.5-7.0	168	2.5	2.1-2.9
2006	624	5.4	4.9-5.8	456	8.7	7.9-9.6	168	2.6	2.2-3.0	505	4.3	3.9-4.7	307	6.1	5.4-6.8	198	2.9	2.5-3.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Liver / intrahepatic bile duct

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	686	5.8	5.4-6.3	487	9.2	8.4-10.0	199	3.1	2.6-3.5	484	4.1	3.7-4.4	322	6.2	5.6-6.9	162	2.3	2.0-2.7
2008	738	6.2	5.7-6.6	518	9.5	8.7-10.4	220	3.3	2.9-3.8	581	4.8	4.4-5.2	373	7.0	6.3-7.8	208	3.0	2.6-3.4
2009	728	6.0	5.6-6.5	509	9.2	8.4-10.0	219	3.2	2.8-3.7	554	4.5	4.2-4.9	357	6.6	6.0-7.4	197	2.8	2.4-3.2
2010	691	5.6	5.2-6.0	524	9.3	8.5-10.1	167	2.5	2.1-2.9	554	4.4	4.1-4.8	348	6.3	5.7-7.0	206	3.0	2.6-3.4
2011	756	6.0	5.6-6.5	535	9.1	8.3-9.9	221	3.3	2.9-3.8	609	4.8	4.4-5.2	376	6.6	5.9-7.3	233	3.3	2.8-3.7
2012	828	6.4	6.0-6.9	609	10.2	9.4-11.1	219	3.2	2.8-3.7	650	5.1	4.7-5.5	423	7.4	6.7-8.1	227	3.2	2.8-3.7
2013	913	7.0	6.6-7.5	654	10.9	10.0-11.8	259	3.7	3.3-4.2	685	5.2	4.8-5.7	466	8.0	7.3-8.8	219	3.0	2.6-3.4
2014	941	7.2	6.7-7.6	686	11.2	10.4-12.1	255	3.6	3.2-4.1	635	4.8	4.4-5.2	438	7.2	6.6-8.0	197	2.8	2.4-3.2
2015	926	6.9	6.4-7.4	673	10.8	10.0-11.7	253	3.5	3.1-4.0	656	4.9	4.5-5.3	429	7.1	6.5-7.9	227	3.0	2.6-3.5
2016	978	7.3	6.8-7.7	712	11.3	10.5-12.2	266	3.8	3.3-4.3	639	4.7	4.3-5.1	416	6.7	6.0-7.4	223	3.0	2.6-3.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Pancreas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1051	11.8	11.1-12.5	566	15.1	13.8-16.5	485	9.5	8.6-10.4	1023	11.5	10.8-12.3	521	14.1	12.9-15.5	502	9.7	8.9-10.6
1977	1110	12.5	11.8-13.3	588	16.4	15.0-17.8	522	10.0	9.1-10.9	1085	12.2	11.5-13.0	568	15.5	14.2-16.9	517	10.0	9.1-10.9
1978	1115	12.3	11.6-13.1	575	15.2	13.9-16.5	540	10.2	9.4-11.1	1086	11.9	11.2-12.7	565	15.0	13.7-16.4	521	9.8	9.0-10.7
1979	1219	13.3	12.6-14.1	643	17.0	15.7-18.5	576	10.7	9.8-11.6	1173	12.8	12.0-13.5	615	16.5	15.1-17.9	558	10.3	9.4-11.1
1980	1169	12.5	11.8-13.3	576	14.8	13.6-16.2	593	10.9	10.0-11.8	1100	11.9	11.2-12.6	554	14.6	13.4-16.0	546	10.0	9.1-10.9
1981	1136	12.1	11.4-12.9	572	14.8	13.6-16.2	564	10.3	9.5-11.2	1102	11.8	11.1-12.5	554	14.4	13.2-15.7	548	10.0	9.2-10.9
1982	1128	11.8	11.2-12.6	552	14.3	13.1-15.6	576	10.2	9.4-11.1	1037	10.9	10.2-11.6	533	13.8	12.6-15.0	504	8.9	8.2-9.8
1983	1086	11.4	10.7-12.1	535	13.6	12.4-14.9	551	9.8	9.0-10.7	1129	11.8	11.1-12.5	549	13.7	12.6-15.0	580	10.2	9.4-11.1
1984	1128	11.7	11.0-12.4	558	14.4	13.1-15.7	570	9.9	9.1-10.8	1089	11.3	10.6-12.0	524	13.6	12.4-14.8	565	9.8	9.0-10.7
1985	1231	12.6	11.9-13.3	580	14.3	13.1-15.5	651	11.3	10.4-12.2	1126	11.5	10.9-12.3	528	13.4	12.2-14.6	598	10.4	9.6-11.3
1986	1166	12.0	11.3-12.7	574	14.2	13.0-15.5	592	10.2	9.4-11.1	1133	11.6	10.9-12.3	560	13.7	12.5-14.9	573	9.8	9.0-10.7
1987	1206	12.1	11.5-12.9	571	14.3	13.1-15.6	635	10.7	9.9-11.6	1132	11.5	10.8-12.2	554	14.0	12.8-15.2	578	9.8	9.0-10.6
1988	1245	12.5	11.8-13.3	607	15.0	13.8-16.3	638	10.7	9.9-11.6	1180	11.8	11.1-12.5	568	14.2	13.0-15.4	612	10.2	9.4-11.0
1989	1164	11.5	10.8-12.2	573	13.8	12.7-15.1	591	9.9	9.1-10.8	1089	10.8	10.2-11.5	547	13.4	12.3-14.6	542	9.1	8.3-9.9
1990	1232	12.2	11.5-12.9	592	14.3	13.1-15.6	640	10.6	9.8-11.4	1144	11.3	10.6-11.9	544	13.1	12.0-14.3	600	9.9	9.1-10.7
1991	1290	12.6	11.9-13.3	618	14.5	13.4-15.8	672	11.0	10.2-11.9	1148	11.2	10.6-11.9	541	13.0	11.9-14.2	607	9.9	9.1-10.7
1992	1241	12.0	11.4-12.7	591	14.1	12.9-15.3	650	10.6	9.8-11.5	1153	11.1	10.5-11.8	536	12.9	11.8-14.0	617	9.9	9.1-10.7
1993	1254	12.0	11.3-12.7	616	14.2	13.1-15.4	638	10.3	9.5-11.1	1217	11.6	10.9-12.3	581	13.5	12.4-14.6	636	10.2	9.4-11.0
1994	1338	12.6	12.0-13.3	668	15.2	14.1-16.5	670	10.6	9.8-11.4	1256	11.9	11.2-12.5	624	14.5	13.3-15.7	632	9.9	9.1-10.7
1995	1328	12.4	11.8-13.1	607	13.6	12.5-14.7	721	11.4	10.5-12.2	1246	11.7	11.0-12.3	578	12.9	11.8-14.0	668	10.4	9.6-11.3
1996	1274	11.9	11.3-12.6	646	14.6	13.5-15.8	628	10.0	9.2-10.8	1192	11.1	10.5-11.8	576	13.2	12.1-14.4	616	9.6	8.9-10.4
1997	1334	12.4	11.7-13.1	646	14.3	13.2-15.5	688	10.8	10.0-11.6	1287	11.9	11.2-12.5	641	14.5	13.4-15.7	646	9.9	9.2-10.8
1998	1412	13.0	12.3-13.7	677	14.8	13.7-16.0	735	11.5	10.7-12.4	1249	11.4	10.8-12.1	614	13.7	12.6-14.8	635	9.8	9.0-10.6
1999	1386	12.6	11.9-13.3	654	14.1	13.1-15.3	732	11.3	10.5-12.1	1294	11.7	11.1-12.4	632	13.7	12.7-14.9	662	10.1	9.4-11.0
2000	1391	12.5	11.9-13.2	655	14.1	13.0-15.2	736	11.3	10.5-12.2	1277	11.4	10.8-12.1	594	13.0	12.0-14.1	683	10.3	9.5-11.1
2001	1420	12.7	12.0-13.3	650	13.7	12.7-14.8	770	11.7	10.9-12.6	1277	11.3	10.7-12.0	585	12.6	11.6-13.6	692	10.4	9.6-11.2
2002	1490	13.2	12.5-13.9	701	14.6	13.5-15.7	789	12.0	11.2-12.9	1319	11.6	11.0-12.3	633	13.2	12.2-14.3	686	10.2	9.5-11.0
2003	1501	13.1	12.4-13.8	725	14.9	13.8-16.0	776	11.7	10.9-12.5	1290	11.2	10.6-11.8	591	12.3	11.3-13.3	699	10.3	9.5-11.1
2004	1478	12.8	12.1-13.4	716	14.5	13.5-15.6	762	11.4	10.6-12.3	1309	11.3	10.7-11.9	648	13.2	12.2-14.3	661	9.8	9.1-10.6
2005	1518	13.0	12.4-13.7	748	14.9	13.9-16.0	770	11.3	10.5-12.2	1342	11.4	10.8-12.1	631	12.7	11.8-13.8	711	10.3	9.6-11.2
2006	1538	13.1	12.5-13.8	734	14.6	13.5-15.7	804	11.9	11.1-12.8	1344	11.4	10.8-12.0	661	13.1	12.1-14.1	683	9.8	9.1-10.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Pancreas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	1607	13.5	12.9-14.2	785	15.3	14.2-16.4	822	12.0	11.2-12.9	1344	11.2	10.6-11.8	660	12.9	11.9-14.0	684	9.9	9.2-10.7
2008	1678	14.0	13.3-14.7	827	15.9	14.8-17.1	851	12.3	11.5-13.2	1388	11.5	10.9-12.1	667	12.9	12.0-14.0	721	10.2	9.4-11.0
2009	1678	13.7	13.1-14.4	870	16.3	15.3-17.5	808	11.6	10.8-12.5	1395	11.3	10.7-11.9	698	13.2	12.2-14.2	697	9.9	9.1-10.6
2010	1720	13.9	13.3-14.6	824	15.3	14.3-16.4	896	12.8	12.0-13.7	1441	11.4	10.9-12.1	702	12.9	12.0-13.9	739	10.2	9.5-11.0
2011	1754	14.0	13.4-14.7	894	16.2	15.2-17.4	860	12.1	11.3-13.0	1411	11.1	10.5-11.7	688	12.6	11.7-13.6	723	9.9	9.2-10.7
2012	1772	14.0	13.3-14.7	914	16.5	15.5-17.7	858	12.0	11.2-12.9	1479	11.6	11.0-12.3	771	14.0	13.0-15.1	708	9.7	9.0-10.5
2013	1825	14.2	13.5-14.9	918	16.2	15.1-17.3	907	12.5	11.6-13.3	1512	11.7	11.1-12.3	757	13.4	12.5-14.4	755	10.2	9.5-11.0
2014	1923	14.8	14.1-15.5	985	16.9	15.8-18.0	938	13.0	12.2-13.9	1499	11.4	10.8-12.0	766	13.3	12.3-14.3	733	9.8	9.1-10.6
2015	1882	14.3	13.6-14.9	943	15.9	14.9-17.0	939	12.8	12.0-13.7	1442	10.8	10.3-11.4	734	12.6	11.6-13.5	708	9.3	8.6-10.0
2016	1983	15.1	14.4-15.8	1026	17.3	16.3-18.5	957	13.4	12.5-14.3	1446	10.8	10.2-11.4	721	12.1	11.2-13.1	725	9.7	8.9-10.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Larynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	503	5.2	4.8-5.7	431	10.4	9.4-11.5	72	1.3	1.0-1.7	175	1.9	1.6-2.2	145	3.8	3.1-4.5	30	0.6	0.4-0.8
1977	507	5.3	4.8-5.8	414	10.0	9.0-11.1	93	1.7	1.4-2.1	169	1.8	1.5-2.1	141	3.6	3.0-4.3	28	0.5	0.3-0.7
1978	546	5.6	5.1-6.1	456	10.9	9.9-12.0	90	1.7	1.3-2.0	185	1.9	1.7-2.2	145	3.7	3.1-4.4	40	0.7	0.5-1.0
1979	563	5.7	5.2-6.2	441	10.2	9.3-11.3	122	2.2	1.8-2.7	209	2.2	1.9-2.5	170	4.2	3.6-5.0	39	0.7	0.5-1.0
1980	577	5.9	5.4-6.4	472	11.1	10.1-12.2	105	2.0	1.6-2.4	184	1.9	1.6-2.2	148	3.7	3.1-4.4	36	0.7	0.5-0.9
1981	627	6.3	5.8-6.8	512	11.9	10.9-13.1	115	2.1	1.7-2.5	184	1.8	1.6-2.1	155	3.9	3.2-4.6	29	0.5	0.3-0.7
1982	560	5.6	5.2-6.1	451	10.5	9.5-11.5	109	2.0	1.6-2.4	191	1.9	1.6-2.2	152	3.6	3.0-4.3	39	0.7	0.5-0.9
1983	567	5.7	5.3-6.2	454	10.5	9.5-11.6	113	2.1	1.8-2.6	183	1.8	1.6-2.1	139	3.4	2.8-4.0	44	0.8	0.6-1.0
1984	569	5.7	5.2-6.2	446	10.2	9.2-11.2	123	2.3	1.9-2.7	162	1.6	1.4-1.9	126	3.1	2.5-3.7	36	0.6	0.4-0.9
1985	588	5.8	5.4-6.3	465	10.5	9.5-11.5	123	2.2	1.9-2.7	178	1.7	1.5-2.0	137	3.2	2.6-3.8	41	0.7	0.5-1.0
1986	571	5.7	5.2-6.2	441	10.2	9.2-11.2	130	2.3	1.9-2.8	167	1.7	1.4-2.0	132	3.2	2.7-3.8	35	0.6	0.4-0.8
1987	609	6.0	5.5-6.5	487	11.2	10.2-12.3	122	2.1	1.8-2.6	176	1.7	1.5-2.0	139	3.4	2.8-4.0	37	0.6	0.4-0.9
1988	544	5.3	4.9-5.8	414	9.3	8.4-10.3	130	2.3	1.9-2.8	172	1.7	1.5-2.0	136	3.3	2.7-3.9	36	0.6	0.4-0.9
1989	607	6.0	5.5-6.5	471	10.7	9.7-11.8	136	2.4	2.0-2.9	171	1.7	1.4-1.9	118	2.7	2.2-3.3	53	0.9	0.7-1.2
1990	608	5.8	5.4-6.3	455	9.9	9.0-10.9	153	2.7	2.3-3.2	153	1.5	1.3-1.8	121	2.9	2.4-3.5	32	0.5	0.4-0.8
1991	551	5.4	4.9-5.8	440	9.7	8.8-10.7	111	1.9	1.6-2.3	159	1.5	1.3-1.8	123	2.7	2.3-3.3	36	0.6	0.4-0.8
1992	565	5.5	5.0-6.0	430	9.5	8.6-10.4	135	2.4	2.0-2.8	205	1.9	1.7-2.2	157	3.6	3.1-4.3	48	0.8	0.6-1.1
1993	542	5.2	4.8-5.7	393	8.8	7.9-9.7	149	2.6	2.2-3.0	168	1.6	1.4-1.9	131	3.0	2.5-3.6	37	0.6	0.4-0.9
1994	518	5.0	4.5-5.4	393	8.6	7.8-9.5	125	2.2	1.8-2.6	169	1.6	1.3-1.8	127	2.9	2.4-3.5	42	0.7	0.5-0.9
1995	541	5.2	4.8-5.6	410	8.9	8.0-9.8	131	2.3	1.9-2.8	165	1.6	1.3-1.8	127	2.9	2.4-3.4	38	0.6	0.4-0.9
1996	483	4.6	4.2-5.0	372	8.0	7.2-8.9	111	2.0	1.6-2.4	165	1.6	1.3-1.8	122	2.7	2.2-3.2	43	0.7	0.5-1.0
1997	492	4.7	4.3-5.1	381	8.2	7.4-9.1	111	2.0	1.6-2.4	141	1.3	1.1-1.6	104	2.3	1.9-2.8	37	0.6	0.4-0.9
1998	466	4.4	4.0-4.8	336	7.1	6.4-7.9	130	2.2	1.9-2.7	151	1.4	1.2-1.6	112	2.4	2.0-2.9	39	0.6	0.4-0.9
1999	521	4.9	4.5-5.3	414	8.6	7.8-9.5	107	1.8	1.5-2.2	155	1.4	1.2-1.7	116	2.5	2.0-3.0	39	0.6	0.5-0.9
2000	500	4.6	4.2-5.0	393	8.2	7.4-9.0	107	1.8	1.5-2.2	140	1.3	1.1-1.5	113	2.4	2.0-2.9	27	0.4	0.3-0.7
2001	472	4.3	3.9-4.7	368	7.5	6.7-8.3	104	1.7	1.4-2.1	138	1.2	1.0-1.5	115	2.4	2.0-2.9	23	0.4	0.2-0.5
2002	462	4.2	3.8-4.6	360	7.4	6.6-8.2	102	1.7	1.4-2.1	146	1.3	1.1-1.5	115	2.4	2.0-2.9	31	0.5	0.3-0.7
2003	450	4.0	3.7-4.4	350	7.0	6.2-7.7	100	1.6	1.3-2.0	129	1.1	0.9-1.3	96	2.0	1.6-2.4	33	0.5	0.3-0.7
2004	453	4.0	3.6-4.4	362	7.0	6.3-7.8	91	1.5	1.2-1.8	128	1.1	0.9-1.3	95	2.0	1.6-2.4	33	0.5	0.3-0.7
2005	510	4.4	4.1-4.8	397	7.6	6.9-8.4	113	1.8	1.5-2.2	144	1.3	1.1-1.5	103	2.1	1.7-2.5	41	0.7	0.5-0.9
2006	486	4.2	3.9-4.6	365	7.1	6.4-7.9	121	2.0	1.6-2.3	139	1.2	1.0-1.4	103	2.0	1.7-2.5	36	0.6	0.4-0.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Larynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	441	3.7	3.4-4.1	352	6.7	6.0-7.4	89	1.4	1.1-1.7	126	1.0	0.9-1.3	106	2.0	1.6-2.4	20	0.3	0.2-0.5
2008	493	4.2	3.8-4.5	389	7.2	6.5-8.0	104	1.6	1.3-2.0	112	0.9	0.8-1.1	87	1.6	1.3-2.0	25	0.4	0.2-0.6
2009	457	3.8	3.5-4.2	355	6.6	5.9-7.3	102	1.6	1.3-1.9	128	1.0	0.9-1.2	93	1.8	1.4-2.2	35	0.5	0.3-0.7
2010	437	3.6	3.3-4.0	342	6.2	5.5-6.9	95	1.5	1.2-1.8	142	1.1	1.0-1.4	106	2.0	1.6-2.4	36	0.5	0.4-0.7
2011	439	3.5	3.2-3.9	356	6.1	5.5-6.8	83	1.3	1.0-1.6	132	1.1	0.9-1.3	96	1.7	1.4-2.1	36	0.5	0.4-0.7
2012	455	3.6	3.3-4.0	356	6.2	5.6-7.0	99	1.4	1.2-1.8	117	0.9	0.7-1.1	91	1.6	1.3-2.0	26	0.4	0.2-0.6
2013	471	3.6	3.3-4.0	374	6.2	5.6-6.9	97	1.5	1.2-1.8	153	1.1	1.0-1.3	124	2.1	1.7-2.5	29	0.4	0.3-0.6
2014	441	3.4	3.1-3.7	345	5.7	5.1-6.4	96	1.4	1.1-1.7	137	1.1	0.9-1.3	108	1.8	1.5-2.2	29	0.4	0.3-0.6
2015	462	3.5	3.2-3.9	367	6.1	5.4-6.7	95	1.4	1.1-1.7	114	0.8	0.7-1.0	87	1.5	1.2-1.8	27	0.4	0.2-0.5
2016	410	3.1	2.8-3.5	332	5.5	4.9-6.1	78	1.1	0.9-1.4	117	0.9	0.7-1.1	92	1.6	1.3-2.0	25	0.4	0.2-0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Lung and bronchus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	5321	57.3	55.7-58.8	3850	98.6	95.4-102.0	1471	28.3	26.9-29.8	4289	46.5	45.1-47.9	3163	82.6	79.6-85.7	1126	21.7	20.4-23.0
1977	5332	56.7	55.1-58.2	3807	96.4	93.2-99.7	1525	29.2	27.7-30.7	4533	48.6	47.2-50.1	3304	85.3	82.2-88.4	1229	23.4	22.1-24.8
1978	5461	57.5	55.9-59.0	3918	98.6	95.3-101.8	1543	29.0	27.6-30.5	4552	48.0	46.6-49.5	3282	83.7	80.8-86.8	1270	23.7	22.4-25.1
1979	5618	58.7	57.2-60.3	3897	97.6	94.4-100.8	1721	32.3	30.7-33.9	4516	47.3	45.9-48.7	3191	81.0	78.0-84.0	1325	24.5	23.2-25.9
1980	5951	61.3	59.7-62.9	4162	102.2	99.0-105.5	1789	32.9	31.3-34.5	4892	50.6	49.2-52.1	3484	86.9	83.9-90.0	1408	25.7	24.4-27.1
1981	6010	61.7	60.2-63.4	4096	101.0	97.8-104.3	1914	35.3	33.7-37.0	4939	50.8	49.4-52.3	3483	87.0	84.0-90.1	1456	26.6	25.2-28.0
1982	6238	63.4	61.8-65.1	4196	102.4	99.2-105.7	2042	37.1	35.5-38.8	5074	52.0	50.6-53.5	3425	85.4	82.4-88.4	1649	29.9	28.4-31.4
1983	6529	65.9	64.3-67.6	4276	103.2	100.0-106.5	2253	40.5	38.8-42.3	5253	53.2	51.8-54.7	3531	87.2	84.2-90.3	1722	30.8	29.3-32.3
1984	6488	65.3	63.7-66.9	4210	100.7	97.5-103.9	2278	41.1	39.4-42.9	5249	52.9	51.4-54.3	3460	83.9	81.0-86.8	1789	31.9	30.4-33.5
1985	6810	68.2	66.5-69.8	4343	103.6	100.4-106.8	2467	44.4	42.6-46.2	5592	56.1	54.6-57.6	3649	88.7	85.7-91.8	1943	34.4	32.9-36.0
1986	6759	67.6	66.0-69.3	4266	102.1	98.9-105.3	2493	44.6	42.8-46.4	5625	56.2	54.7-57.7	3612	87.6	84.7-90.7	2013	35.7	34.2-37.4
1987	6949	69.0	67.3-70.6	4320	102.0	98.9-105.2	2629	46.7	44.9-48.5	5603	55.7	54.3-57.2	3586	86.5	83.6-89.5	2017	35.2	33.7-36.9
1988	7134	70.1	68.5-71.8	4356	102.7	99.5-105.9	2778	48.7	46.8-50.6	5792	57.0	55.6-58.6	3619	86.4	83.5-89.3	2173	37.8	36.2-39.5
1989	7256	70.9	69.3-72.6	4361	101.2	98.1-104.4	2895	50.5	48.7-52.4	5851	57.3	55.8-58.8	3613	85.0	82.1-87.9	2238	38.8	37.1-40.4
1990	7308	70.7	69.1-72.4	4413	101.9	98.8-105.0	2895	49.8	47.9-51.6	5988	58.0	56.5-59.5	3619	84.5	81.7-87.5	2369	40.2	38.6-41.9
1991	7411	71.5	69.8-73.1	4339	99.6	96.5-102.7	3072	52.6	50.8-54.6	6180	59.5	58.1-61.1	3720	86.5	83.7-89.4	2460	41.7	40.0-43.4
1992	7456	71.3	69.6-72.9	4288	97.2	94.2-100.3	3168	53.7	51.9-55.7	6226	59.5	58.0-61.0	3616	83.7	80.9-86.6	2610	43.5	41.9-45.3
1993	7448	70.8	69.2-72.4	4175	93.8	90.9-96.8	3273	55.3	53.4-57.2	6038	57.3	55.8-58.7	3488	79.6	76.9-82.4	2550	42.5	40.9-44.2
1994	7705	72.8	71.2-74.4	4318	96.6	93.7-99.6	3387	57.0	55.1-59.0	6240	58.7	57.3-60.2	3656	83.1	80.4-85.9	2584	42.5	40.9-44.2
1995	7700	72.4	70.8-74.0	4301	95.0	92.1-97.9	3399	56.8	54.9-58.8	6246	58.6	57.1-60.0	3598	81.1	78.4-83.9	2648	43.6	41.9-45.3
1996	7861	73.7	72.1-75.4	4404	97.0	94.1-99.9	3457	57.7	55.8-59.7	6137	57.4	56.0-58.9	3501	78.3	75.7-81.0	2636	43.4	41.7-45.1
1997	7905	73.6	72.0-75.3	4302	94.1	91.2-97.0	3603	59.9	57.9-61.9	6037	56.0	54.6-57.4	3358	74.5	71.9-77.1	2679	43.5	41.8-45.2
1998	8024	74.1	72.5-75.8	4308	92.5	89.7-95.3	3716	61.5	59.5-63.5	6052	55.6	54.2-57.0	3400	73.8	71.3-76.4	2652	42.8	41.1-44.4
1999	8103	74.2	72.6-75.9	4357	93.1	90.4-96.0	3746	61.3	59.3-63.3	6019	54.9	53.5-56.3	3295	71.3	68.9-73.8	2724	43.7	42.1-45.4
2000	8256	74.9	73.3-76.5	4358	92.2	89.5-95.0	3898	63.0	61.0-65.0	6180	55.7	54.3-57.1	3362	71.9	69.5-74.4	2818	44.6	42.9-46.3
2001	8068	72.7	71.2-74.4	4154	87.5	84.8-90.2	3914	63.2	61.2-65.2	6077	54.4	53.0-55.8	3301	70.6	68.2-73.0	2776	43.8	42.2-45.5
2002	8283	73.9	72.3-75.5	4353	90.5	87.8-93.2	3930	62.8	60.9-64.8	6199	54.9	53.6-56.3	3294	69.2	66.8-71.6	2905	45.6	43.9-47.3
2003	8364	74.0	72.4-75.6	4376	89.6	86.9-92.3	3988	63.3	61.3-65.3	6013	52.8	51.4-54.1	3206	66.4	64.1-68.7	2807	43.5	41.9-45.2
2004	8387	73.5	71.9-75.1	4230	85.6	83.0-88.2	4157	65.3	63.3-67.3	6029	52.5	51.2-53.8	3089	63.1	60.9-65.4	2940	45.2	43.6-46.9
2005	8414	73.3	71.8-74.9	4265	85.6	83.0-88.2	4149	64.8	62.8-66.8	5985	51.8	50.5-53.2	3118	63.4	61.2-65.7	2867	43.7	42.1-45.4
2006	8655	75.1	73.6-76.8	4239	84.2	81.7-86.8	4416	69.3	67.2-71.4	6027	51.9	50.6-53.2	3176	63.7	61.5-66.0	2851	43.7	42.1-45.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Lung and bronchus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	8594	73.5	71.9-75.1	4307	84.3	81.7-86.9	4287	66.2	64.2-68.3	6023	51.1	49.8-52.4	3098	60.9	58.8-63.1	2925	44.2	42.6-45.9
2008	8883	75.0	73.4-76.6	4424	85.4	82.8-88.0	4459	68.2	66.2-70.3	5986	49.9	48.6-51.2	3069	59.5	57.4-61.6	2917	43.4	41.8-45.1
2009	8734	72.7	71.2-74.2	4439	84.0	81.5-86.6	4295	64.7	62.7-66.7	5818	48.2	47.0-49.5	3037	57.9	55.9-60.1	2781	41.3	39.8-42.9
2010	8635	71.1	69.5-72.6	4243	78.7	76.4-81.2	4392	66.2	64.2-68.2	5772	47.2	46.0-48.5	2928	54.9	52.9-56.9	2844	41.8	40.3-43.4
2011	8792	71.3	69.8-72.8	4339	79.3	77.0-81.8	4453	65.8	63.9-67.8	5740	46.3	45.1-47.5	2921	54.0	52.0-56.0	2819	40.8	39.2-42.4
2012	8700	69.4	68.0-70.9	4294	77.4	75.0-79.7	4406	64.1	62.2-66.1	5918	47.1	45.9-48.3	3071	55.9	53.9-58.0	2847	40.8	39.3-42.4
2013	8790	69.3	67.8-70.7	4280	75.4	73.1-77.8	4510	65.0	63.0-66.9	5655	44.1	42.9-45.2	2863	50.6	48.7-52.5	2792	39.3	37.8-40.8
2014	8551	66.0	64.6-67.5	4170	72.3	70.1-74.6	4381	61.8	59.9-63.7	5434	41.9	40.8-43.1	2783	48.7	46.9-50.6	2651	37.0	35.6-38.5
2015	9014	69.0	67.5-70.5	4285	73.3	71.1-75.6	4729	66.4	64.5-68.4	5420	41.1	40.0-42.3	2755	47.5	45.7-49.4	2665	36.5	35.1-38.0
2016	8769	65.9	64.5-67.3	4235	70.9	68.7-73.1	4534	62.8	60.9-64.7	4968	37.2	36.2-38.3	2490	42.4	40.7-44.1	2478	33.4	32.1-34.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Melanoma of the skin

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	530	5.9	5.4-6.5	270	6.5	5.8-7.4	260	5.4	4.8-6.1	233	2.6	2.3-3.0	130	3.2	2.7-3.9	103	2.1	1.7-2.6
1977	685	7.6	7.0-8.2	373	9.0	8.1-10.0	312	6.5	5.8-7.3	246	2.8	2.4-3.2	159	4.0	3.4-4.7	87	1.8	1.5-2.3
1978	653	7.3	6.7-7.8	334	8.2	7.3-9.2	319	6.7	6.0-7.5	219	2.5	2.1-2.8	139	3.5	2.9-4.1	80	1.6	1.3-2.0
1979	673	7.4	6.9-8.0	361	8.8	7.9-9.8	312	6.4	5.7-7.2	236	2.6	2.3-3.0	133	3.3	2.8-3.9	103	2.1	1.7-2.5
1980	824	9.0	8.4-9.7	456	11.1	10.1-12.3	368	7.5	6.7-8.3	251	2.7	2.4-3.1	152	3.6	3.1-4.3	99	2.0	1.6-2.5
1981	803	8.7	8.1-9.3	420	10.1	9.1-11.2	383	7.7	6.9-8.5	223	2.4	2.1-2.7	132	3.2	2.6-3.8	91	1.8	1.4-2.2
1982	837	8.9	8.3-9.6	408	9.7	8.7-10.7	429	8.5	7.7-9.4	253	2.7	2.4-3.1	157	3.8	3.2-4.5	96	1.8	1.5-2.3
1983	913	9.8	9.1-10.4	503	12.1	11.0-13.2	410	8.4	7.6-9.2	244	2.6	2.3-2.9	136	3.2	2.7-3.9	108	2.2	1.8-2.6
1984	814	8.6	8.0-9.2	445	10.4	9.4-11.4	369	7.4	6.6-8.2	254	2.7	2.4-3.0	144	3.5	2.9-4.1	110	2.1	1.7-2.6
1985	913	9.7	9.1-10.3	489	11.5	10.4-12.6	424	8.4	7.6-9.3	252	2.6	2.3-3.0	151	3.7	3.1-4.3	101	1.9	1.5-2.3
1986	971	10.2	9.6-10.9	523	12.3	11.2-13.4	448	9.0	8.1-9.9	273	2.9	2.5-3.3	158	3.8	3.2-4.5	115	2.2	1.8-2.7
1987	913	9.5	8.9-10.2	500	11.6	10.5-12.6	413	8.1	7.3-8.9	297	3.1	2.8-3.5	179	4.2	3.6-4.9	118	2.3	1.9-2.8
1988	877	9.0	8.4-9.7	503	11.7	10.7-12.8	374	7.2	6.5-8.0	291	3.0	2.6-3.3	187	4.4	3.8-5.1	104	1.9	1.5-2.3
1989	927	9.6	8.9-10.2	523	12.0	11.0-13.1	404	7.7	7.0-8.5	312	3.2	2.9-3.6	199	4.8	4.1-5.6	113	2.1	1.7-2.5
1990	962	9.7	9.1-10.3	518	11.6	10.6-12.6	444	8.4	7.6-9.2	297	3.0	2.7-3.4	181	4.2	3.6-4.9	116	2.2	1.8-2.6
1991	890	8.9	8.3-9.5	517	11.6	10.6-12.7	373	6.9	6.2-7.7	298	3.0	2.6-3.3	187	4.4	3.7-5.0	111	2.0	1.6-2.4
1992	924	9.2	8.6-9.8	540	12.1	11.0-13.1	384	7.0	6.3-7.8	295	2.9	2.6-3.3	172	3.9	3.4-4.6	123	2.2	1.8-2.6
1993	927	9.0	8.5-9.7	530	11.6	10.6-12.6	397	7.2	6.5-7.9	290	2.8	2.5-3.2	183	4.2	3.6-4.9	107	1.9	1.5-2.3
1994	1013	9.9	9.3-10.6	592	13.0	11.9-14.1	421	7.7	7.0-8.5	294	2.9	2.5-3.2	185	4.2	3.6-4.8	109	1.9	1.5-2.3
1995	1110	10.9	10.2-11.5	665	14.5	13.4-15.6	445	8.1	7.4-8.9	311	3.0	2.7-3.4	183	4.2	3.6-4.8	128	2.2	1.8-2.6
1996	1191	11.6	11.0-12.3	686	14.9	13.8-16.1	505	9.2	8.4-10.0	319	3.1	2.7-3.4	197	4.3	3.7-4.9	122	2.1	1.8-2.6
1997	1278	12.3	11.7-13.0	752	16.1	15.0-17.3	526	9.5	8.7-10.4	303	2.9	2.6-3.2	204	4.6	3.9-5.2	99	1.6	1.3-2.0
1998	1359	13.1	12.4-13.8	777	16.5	15.4-17.8	582	10.6	9.8-11.6	292	2.7	2.4-3.1	187	4.0	3.5-4.6	105	1.8	1.4-2.1
1999	1401	13.3	12.6-14.1	811	17.1	16.0-18.4	590	10.5	9.7-11.4	284	2.6	2.3-3.0	167	3.6	3.1-4.2	117	2.0	1.6-2.4
2000	1557	14.7	14.0-15.5	894	18.8	17.5-20.0	663	11.9	11.0-12.8	304	2.8	2.5-3.1	196	4.1	3.5-4.7	108	1.8	1.5-2.2
2001	1724	16.2	15.4-17.0	972	20.2	18.9-21.5	752	13.4	12.4-14.4	356	3.2	2.9-3.6	229	4.8	4.2-5.5	127	2.1	1.7-2.5
2002	1702	15.8	15.0-16.6	982	20.2	18.9-21.5	720	12.7	11.8-13.7	313	2.8	2.5-3.2	168	3.5	3.0-4.0	145	2.4	2.0-2.8
2003	1768	16.2	15.5-17.0	1033	21.0	19.7-22.3	735	12.8	11.9-13.8	312	2.8	2.5-3.1	191	3.9	3.4-4.5	121	1.9	1.6-2.3
2004	2103	19.3	18.4-20.1	1210	24.3	22.9-25.7	893	15.7	14.6-16.7	322	2.8	2.5-3.2	197	4.0	3.4-4.6	125	2.0	1.7-2.4
2005	2305	21.1	20.2-22.0	1348	27.0	25.6-28.5	957	16.8	15.7-17.9	335	2.9	2.6-3.3	209	4.2	3.6-4.8	126	2.0	1.7-2.4
2006	2475	22.4	21.5-23.3	1428	28.2	26.7-29.7	1047	18.3	17.1-19.4	340	3.0	2.7-3.3	221	4.4	3.8-5.0	119	1.8	1.5-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Melanoma of the skin

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	2398	21.6	20.7-22.5	1384	27.1	25.7-28.6	1014	17.8	16.7-18.9	327	2.8	2.5-3.1	235	4.6	4.1-5.3	92	1.4	1.1-1.8
2008	2621	23.4	22.5-24.3	1520	29.6	28.1-31.1	1101	19.1	18.0-20.3	345	2.9	2.6-3.3	228	4.4	3.8-5.0	117	1.8	1.5-2.2
2009	2798	24.6	23.7-25.6	1647	31.5	29.9-33.1	1151	19.8	18.7-21.1	350	3.0	2.7-3.3	206	4.0	3.5-4.6	144	2.2	1.9-2.6
2010	2688	23.5	22.6-24.5	1540	29.2	27.7-30.7	1148	19.5	18.3-20.7	347	2.9	2.6-3.2	212	4.0	3.5-4.6	135	2.1	1.7-2.5
2011	2971	25.3	24.4-26.3	1733	32.0	30.5-33.6	1238	20.4	19.3-21.7	368	3.0	2.7-3.4	241	4.5	3.9-5.1	127	1.9	1.6-2.3
2012	2903	24.7	23.8-25.7	1648	30.1	28.6-31.6	1255	21.1	19.9-22.4	394	3.2	2.8-3.5	250	4.7	4.1-5.3	144	2.0	1.7-2.4
2013	3193	27.0	26.1-28.0	1844	33.1	31.6-34.7	1349	22.8	21.6-24.2	379	3.1	2.8-3.4	252	4.7	4.1-5.3	127	1.8	1.5-2.2
2014	3172	26.7	25.7-27.7	1856	33.1	31.6-34.7	1316	22.0	20.7-23.3	336	2.6	2.4-2.9	217	3.8	3.3-4.4	119	1.7	1.4-2.0
2015	3108	25.5	24.6-26.5	1837	32.2	30.7-33.8	1271	20.8	19.6-22.0	293	2.3	2.0-2.6	193	3.4	2.9-3.9	100	1.4	1.2-1.8
2016	3086	25.2	24.2-26.1	1837	31.7	30.2-33.3	1249	20.5	19.3-21.8	306	2.3	2.0-2.6	195	3.4	2.9-3.9	111	1.5	1.2-1.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Female breast

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	5019	102.0	99.1-104.9	1957	38.9	37.2-40.7
1977	5070	102.1	99.2-105.0	2012	39.7	37.9-41.5
1978	5185	103.4	100.5-106.3	2027	39.2	37.5-41.0
1979	5141	101.6	98.7-104.4	1997	38.6	36.9-40.4
1980	5307	103.7	100.9-106.6	1986	37.7	36.0-39.4
1981	5372	104.7	101.8-107.6	1994	37.5	35.8-39.2
1982	5472	105.3	102.5-108.2	2022	37.7	36.1-39.5
1983	5852	112.4	109.5-115.4	2059	38.2	36.6-40.0
1984	6015	114.5	111.5-117.5	2126	39.1	37.4-40.8
1985	6377	121.7	118.7-124.8	2162	39.8	38.1-41.6
1986	6547	123.8	120.7-126.9	2195	39.9	38.2-41.6
1987	6755	126.8	123.7-129.9	2196	39.4	37.7-41.1
1988	6880	128.0	124.9-131.1	2165	38.9	37.3-40.6
1989	6744	124.9	121.9-128.0	2158	38.7	37.0-40.4
1990	7151	131.3	128.2-134.4	2125	37.3	35.7-39.0
1991	7066	128.5	125.4-131.5	2054	36.0	34.4-37.6
1992	7110	128.6	125.6-131.7	2043	35.3	33.8-36.9
1993	7124	127.9	124.9-131.0	2026	34.5	33.0-36.0
1994	7201	129.1	126.0-132.1	2020	34.3	32.8-35.9
1995	7573	135.7	132.7-138.9	1933	32.8	31.4-34.4
1996	8042	143.3	140.1-146.5	1962	32.7	31.2-34.2
1997	8190	144.5	141.4-147.7	1846	30.6	29.2-32.0
1998	8461	147.9	144.8-151.2	1864	30.4	29.1-31.9
1999	8373	145.0	141.8-148.1	1740	28.0	26.7-29.4
2000	8451	144.4	141.3-147.5	1759	28.0	26.7-29.3
2001	8303	141.2	138.1-144.3	1708	27.1	25.8-28.4
2002	8305	140.2	137.2-143.3	1684	26.3	25.0-27.6
2003	8247	137.8	134.8-140.8	1678	26.0	24.8-27.3
2004	8112	134.8	131.8-137.8	1535	23.6	22.4-24.8
2005	8244	135.7	132.7-138.7	1506	23.0	21.8-24.2
2006	8350	138.0	135.0-141.1	1444	22.0	20.9-23.2
2007	8345	136.3	133.3-139.3	1504	22.6	21.5-23.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Female breast

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	8660	140.6	137.6-143.7	1435	21.4	20.3-22.6
2009	8865	141.8	138.8-144.9	1440	21.1	20.0-22.3
2010	8296	132.3	129.4-135.3	1493	21.5	20.4-22.7
2011	8742	138.8	135.8-141.8	1453	20.8	19.7-22.0
2012	8659	137.0	134.0-140.0	1395	19.8	18.7-20.9
2013	8946	140.8	137.8-143.9	1397	19.9	18.8-21.0
2014	9129	142.9	139.9-146.0	1246	17.4	16.4-18.5
2015	9199	143.8	140.7-146.9	1317	18.3	17.3-19.4
2016	8843	137.7	134.7-140.8	1311	18.4	17.4-19.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Cervix uteri

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	597	12.4	11.4-13.5	234	4.6	4.1-5.3
1977	517	10.6	9.7-11.5	191	3.7	3.2-4.3
1978	518	10.6	9.7-11.6	223	4.4	3.8-5.0
1979	507	10.4	9.5-11.4	215	4.3	3.7-4.9
1980	552	10.9	10.0-11.9	209	4.2	3.6-4.8
1981	456	9.0	8.2-9.9	217	4.1	3.6-4.7
1982	463	9.4	8.5-10.3	191	3.6	3.1-4.1
1983	477	9.6	8.7-10.5	167	3.2	2.7-3.7
1984	517	10.4	9.5-11.4	164	3.2	2.7-3.7
1985	478	9.5	8.7-10.4	161	3.1	2.7-3.7
1986	435	8.7	7.9-9.6	182	3.5	3.0-4.0
1987	499	9.8	8.9-10.7	181	3.5	3.0-4.1
1988	512	10.1	9.3-11.1	170	3.2	2.7-3.7
1989	483	9.5	8.6-10.4	151	2.8	2.3-3.3
1990	513	10.1	9.2-11.0	168	3.0	2.6-3.5
1991	498	9.5	8.7-10.4	141	2.6	2.2-3.1
1992	458	8.8	8.0-9.6	163	3.0	2.6-3.5
1993	486	9.2	8.4-10.1	144	2.6	2.2-3.1
1994	462	8.7	7.9-9.6	157	2.9	2.4-3.3
1995	552	10.5	9.6-11.4	117	2.1	1.8-2.6
1996	541	10.2	9.4-11.1	147	2.7	2.3-3.2
1997	487	9.2	8.4-10.1	155	2.7	2.3-3.2
1998	448	8.5	7.7-9.3	122	2.2	1.8-2.6
1999	441	8.3	7.5-9.1	144	2.5	2.1-3.0
2000	406	7.6	6.9-8.4	121	2.1	1.7-2.5
2001	433	8.0	7.2-8.8	129	2.2	1.8-2.6
2002	372	6.9	6.2-7.7	132	2.3	1.9-2.7
2003	412	7.7	7.0-8.5	113	1.9	1.5-2.3
2004	346	6.5	5.8-7.3	122	2.1	1.7-2.5
2005	397	7.5	6.8-8.3	120	2.0	1.7-2.4
2006	383	7.2	6.5-8.0	113	1.9	1.6-2.3
2007	393	7.3	6.6-8.1	118	2.0	1.6-2.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Cervix uteri

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	374	7.1	6.3-7.8	99	1.6	1.3-2.0
2009	403	7.6	6.8-8.4	91	1.5	1.2-1.9
2010	389	7.5	6.8-8.4	113	1.8	1.5-2.2
2011	314	5.8	5.2-6.6	110	1.8	1.5-2.2
2012	349	6.4	5.7-7.2	139	2.4	2.0-2.8
2013	366	7.0	6.2-7.8	113	1.9	1.5-2.3
2014	349	6.7	5.9-7.4	116	2.0	1.6-2.4
2015	362	7.0	6.3-7.8	101	1.6	1.3-1.9
2016	331	6.6	5.8-7.4	114	1.9	1.5-2.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Corpus uterus and uterus, not otherwise specified

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1510	29.3	27.8-30.8	285	5.5	4.9-6.2
1977	1384	26.3	24.9-27.7	286	5.4	4.8-6.1
1978	1230	23.1	21.8-24.4	256	4.8	4.2-5.4
1979	1349	25.3	24.0-26.7	290	5.3	4.7-5.9
1980	1350	25.5	24.2-27.0	292	5.2	4.6-5.9
1981	1465	27.3	25.8-28.7	283	5.1	4.5-5.8
1982	1403	25.7	24.4-27.1	300	5.4	4.8-6.0
1983	1338	24.8	23.4-26.2	280	4.8	4.3-5.5
1984	1393	25.5	24.2-26.9	291	5.1	4.5-5.7
1985	1330	24.1	22.8-25.5	281	4.9	4.3-5.5
1986	1352	24.7	23.3-26.0	274	4.6	4.1-5.2
1987	1436	26.0	24.7-27.4	271	4.6	4.0-5.2
1988	1315	23.7	22.4-25.1	275	4.6	4.1-5.2
1989	1331	23.7	22.4-25.0	276	4.6	4.1-5.2
1990	1387	24.8	23.5-26.1	239	3.9	3.5-4.5
1991	1402	25.5	24.2-26.9	271	4.4	3.9-5.0
1992	1477	26.3	24.9-27.7	281	4.6	4.1-5.2
1993	1410	25.0	23.7-26.4	293	4.7	4.2-5.3
1994	1518	26.6	25.2-28.0	289	4.7	4.1-5.2
1995	1523	26.9	25.6-28.3	298	4.7	4.2-5.3
1996	1623	28.4	27.0-29.8	276	4.4	3.9-4.9
1997	1572	27.6	26.2-29.0	273	4.4	3.8-4.9
1998	1724	30.0	28.6-31.5	272	4.3	3.8-4.9
1999	1649	28.3	27.0-29.8	248	3.8	3.3-4.3
2000	1653	28.0	26.7-29.4	277	4.3	3.8-4.8
2001	1806	30.7	29.3-32.1	269	4.1	3.6-4.6
2002	1833	30.6	29.2-32.1	289	4.5	4.0-5.0
2003	1821	30.2	28.8-31.6	304	4.6	4.1-5.2
2004	1838	30.1	28.7-31.5	293	4.4	3.9-5.0
2005	1899	30.9	29.5-32.3	291	4.4	3.9-5.0
2006	1862	29.9	28.5-31.3	317	4.7	4.2-5.3
2007	1943	31.1	29.7-32.6	298	4.3	3.9-4.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Corpus uterus and uterus, not otherwise specified

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	2067	32.7	31.3-34.2	306	4.5	4.0-5.1
2009	1960	30.6	29.2-32.0	304	4.5	4.0-5.0
2010	2003	31.0	29.7-32.5	303	4.3	3.8-4.8
2011	2083	31.7	30.3-33.1	335	4.9	4.4-5.5
2012	2171	32.4	31.0-33.8	343	4.8	4.3-5.4
2013	2306	34.2	32.8-35.7	375	5.1	4.6-5.7
2014	2276	33.5	32.1-34.9	360	4.8	4.3-5.4
2015	2223	32.3	30.9-33.7	320	4.4	3.9-4.9
2016	2295	33.7	32.3-35.2	341	4.6	4.1-5.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Ovary

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	870	17.3	16.2-18.5	652	12.7	11.7-13.7
1977	848	16.6	15.4-17.7	554	10.7	9.8-11.6
1978	837	16.2	15.1-17.4	596	11.2	10.3-12.1
1979	874	16.9	15.8-18.1	571	10.8	9.9-11.7
1980	871	16.7	15.6-17.8	583	10.9	10.0-11.8
1981	944	18.1	16.9-19.3	624	11.4	10.5-12.4
1982	880	16.8	15.7-17.9	619	11.3	10.4-12.3
1983	950	17.9	16.8-19.1	613	11.0	10.1-11.9
1984	904	17.0	15.9-18.2	601	10.9	10.0-11.8
1985	925	17.6	16.5-18.9	588	10.6	9.8-11.5
1986	947	17.9	16.8-19.1	567	10.2	9.4-11.1
1987	931	17.3	16.2-18.5	611	10.9	10.1-11.8
1988	947	17.6	16.4-18.7	618	10.9	10.0-11.8
1989	952	17.5	16.3-18.6	623	10.9	10.0-11.8
1990	998	18.1	17.0-19.3	632	10.8	10.0-11.7
1991	1002	18.2	17.1-19.4	655	11.2	10.4-12.1
1992	1001	18.1	17.0-19.3	629	10.9	10.0-11.8
1993	995	17.6	16.5-18.8	624	10.5	9.6-11.3
1994	1019	18.0	16.9-19.2	618	10.4	9.6-11.3
1995	1020	18.0	16.9-19.2	640	10.7	9.8-11.5
1996	1044	18.3	17.2-19.5	600	9.9	9.1-10.8
1997	999	17.6	16.5-18.7	650	10.7	9.9-11.6
1998	923	16.1	15.1-17.2	573	9.4	8.6-10.2
1999	965	16.5	15.5-17.6	582	9.4	8.7-10.2
2000	999	16.9	15.8-17.9	633	10.1	9.3-10.9
2001	986	16.6	15.5-17.7	636	10.0	9.2-10.8
2002	951	15.9	14.9-17.0	604	9.5	8.8-10.3
2003	972	16.1	15.1-17.1	627	9.7	9.0-10.5
2004	929	15.2	14.3-16.3	609	9.4	8.6-10.2
2005	888	14.5	13.6-15.5	653	10.0	9.2-10.8
2006	933	15.2	14.2-16.2	627	9.7	8.9-10.5
2007	902	14.5	13.5-15.5	574	8.8	8.0-9.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Ovary

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	902	14.4	13.4-15.4	615	9.1	8.4-9.9
2009	840	13.2	12.3-14.2	574	8.5	7.8-9.2
2010	893	14.1	13.1-15.1	557	8.3	7.6-9.0
2011	881	13.6	12.7-14.6	566	8.2	7.5-8.9
2012	828	12.9	12.0-13.8	544	7.7	7.1-8.5
2013	927	14.4	13.5-15.4	586	8.3	7.6-9.0
2014	776	11.9	11.1-12.9	516	7.2	6.6-7.9
2015	815	12.6	11.7-13.6	524	7.3	6.7-8.0
2016	809	12.4	11.5-13.3	481	6.7	6.1-7.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Prostate

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	2534	78.5	75.3-81.8	938	31.5	29.4-33.7
1977	2610	78.7	75.6-82.0	928	30.8	28.7-32.9
1978	2574	77.3	74.2-80.5	975	32.6	30.5-34.8
1979	2643	79.0	75.8-82.2	994	32.5	30.4-34.7
1980	2973	85.2	82.0-88.5	996	32.1	30.0-34.2
1981	3040	87.4	84.1-90.7	952	30.6	28.6-32.7
1982	3170	89.3	86.0-92.6	1011	31.7	29.6-33.8
1983	3355	92.8	89.5-96.2	1093	33.6	31.6-35.8
1984	3315	90.4	87.2-93.7	1134	34.8	32.7-37.0
1985	3500	94.0	90.7-97.3	1103	33.6	31.5-35.7
1986	3575	94.9	91.7-98.2	1188	34.9	32.8-37.0
1987	3919	102.6	99.2-106.0	1163	34.1	32.1-36.2
1988	4036	104.7	101.4-108.2	1184	34.2	32.2-36.3
1989	4249	108.3	104.9-111.8	1218	35.1	33.1-37.2
1990	5005	121.7	118.2-125.3	1234	34.8	32.8-36.9
1991	6040	143.2	139.5-147.0	1354	37.4	35.3-39.5
1992	7646	175.4	171.4-179.5	1302	35.9	33.9-38.0
1993	7325	165.1	161.2-169.0	1329	36.1	34.2-38.2
1994	6617	147.3	143.7-151.0	1359	36.3	34.4-38.4
1995	6622	144.8	141.3-148.4	1290	34.3	32.4-36.3
1996	7304	158.8	155.1-162.5	1293	33.4	31.5-35.3
1997	7664	165.2	161.5-169.0	1304	33.0	31.2-34.9
1998	7534	160.4	156.7-164.1	1149	28.4	26.8-30.2
1999	7981	168.6	164.8-172.3	1166	28.3	26.7-30.0
2000	8633	181.4	177.5-185.3	1148	28.0	26.4-29.7
2001	8684	180.0	176.2-183.8	1149	27.4	25.8-29.1
2002	9041	184.6	180.8-188.5	1068	25.0	23.5-26.6
2003	8455	170.6	167.0-174.3	1083	24.8	23.3-26.3
2004	8201	163.0	159.5-166.6	1038	23.4	22.0-24.9
2005	8311	162.7	159.2-166.3	961	21.2	19.9-22.6
2006	9349	180.9	177.2-184.6	942	20.4	19.1-21.8
2007	9467	178.5	174.9-182.2	993	20.9	19.6-22.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Prostate

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	9173	169.2	165.7-172.7	941	19.3	18.1-20.6
2009	8958	160.9	157.5-164.3	872	17.7	16.5-18.9
2010	8477	149.0	145.8-152.3	945	18.6	17.4-19.8
2011	8724	150.7	147.5-154.0	914	17.7	16.6-18.9
2012	7463	125.4	122.5-128.3	883	17.0	15.8-18.1
2013	7257	119.6	116.8-122.4	838	15.6	14.6-16.7
2014	7086	114.0	111.3-116.7	902	16.5	15.4-17.6
2015	7453	117.7	115.0-120.5	869	15.8	14.8-17.0
2016	7638	118.8	116.0-121.5	884	15.6	14.6-16.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Testis

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	172	3.6	3.1-4.3	32	0.6	0.4-0.9
1977	169	3.6	3.0-4.2	28	0.5	0.4-0.8
1978	193	4.2	3.6-4.9	34	0.7	0.5-1.0
1979	223	4.8	4.2-5.5	31	0.7	0.5-1.0
1980	228	4.7	4.1-5.4	28	0.6	0.4-0.8
1981	218	4.5	3.9-5.1	28	0.5	0.4-0.8
1982	213	4.4	3.8-5.1	25	0.5	0.3-0.8
1983	235	4.8	4.2-5.5	23	0.5	0.3-0.8
1984	287	5.9	5.2-6.6	17	0.4	0.2-0.6
1985	255	5.2	4.6-5.9	20	0.4	0.3-0.7
1986	247	4.9	4.3-5.6	25	0.5	0.3-0.8
1987	274	5.5	4.8-6.2	20	0.4	0.2-0.6
1988	260	5.1	4.5-5.8	16	0.4	0.2-0.6
1989	260	5.1	4.5-5.8	21	0.4	0.3-0.6
1990	290	5.7	5.1-6.4	16	0.3	0.2-0.5
1991	268	5.3	4.7-6.0	13	0.2	0.1-0.4
1992	285	5.7	5.0-6.4	19	0.4	0.2-0.6
1993	302	6.1	5.4-6.8	23	0.5	0.3-0.8
1994	294	6.0	5.3-6.7	12	0.2	0.1-0.4
1995	309	6.4	5.7-7.1	15	0.3	0.2-0.5
1996	302	6.3	5.6-7.0	21	0.4	0.3-0.7
1997	325	6.8	6.1-7.6	6	0.1	0.1-0.3
1998	329	7.0	6.2-7.8	18	0.4	0.2-0.6
1999	324	6.9	6.2-7.7	13	0.3	0.1-0.5
2000	295	6.3	5.6-7.1	13	0.3	0.1-0.5
2001	316	6.8	6.1-7.6	13	0.3	0.2-0.5
2002	300	6.5	5.8-7.3	22	0.5	0.3-0.7
2003	339	7.5	6.7-8.3	11	0.2	0.1-0.4
2004	317	7.1	6.3-7.9	11	0.2	0.1-0.4
2005	339	7.6	6.8-8.4	7	0.1	0.1-0.3
2006	314	7.0	6.2-7.8	12	0.2	0.1-0.4
2007	339	7.7	6.9-8.6	6	0.1	0.0-0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Testis

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	370	8.5	7.6-9.4	10	0.2	0.1-0.4
2009	334	7.6	6.8-8.5	7	0.1	0.1-0.3
2010	348	7.9	7.1-8.8	14	0.3	0.2-0.5
2011	341	7.7	6.9-8.6	9	0.2	0.1-0.3
2012	364	8.3	7.5-9.3	11	0.2	0.1-0.4
2013	317	7.3	6.5-8.2	7	0.1	0.1-0.3
2014	344	8.0	7.1-8.9	15	0.3	0.2-0.6
2015	332	7.6	6.8-8.5	13	0.3	0.1-0.5
2016	348	8.1	7.3-9.0	9	0.2	0.1-0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Urinary bladder (incl. in situ)

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1895	21.4	20.5-22.4	1397	39.0	36.9-41.3	498	9.7	8.8-10.6	585	7.0	6.5-7.6	424	13.3	12.0-14.7	161	3.2	2.8-3.8
1977	1961	21.9	20.9-22.9	1438	39.2	37.0-41.3	523	10.2	9.3-11.1	583	6.9	6.4-7.5	410	12.7	11.4-14.0	173	3.4	2.9-4.0
1978	2055	22.8	21.8-23.8	1497	40.8	38.7-43.1	558	10.8	9.9-11.7	595	6.9	6.4-7.5	419	12.9	11.6-14.3	176	3.4	2.9-3.9
1979	2076	22.6	21.6-23.6	1569	42.4	40.2-44.6	507	9.4	8.6-10.3	579	6.7	6.2-7.3	389	11.7	10.5-13.0	190	3.6	3.1-4.2
1980	2192	23.5	22.5-24.5	1615	42.6	40.5-44.9	577	10.6	9.8-11.5	555	6.2	5.7-6.7	392	11.6	10.4-12.9	163	3.0	2.6-3.5
1981	2145	22.7	21.7-23.7	1595	41.8	39.6-44.0	550	10.1	9.3-11.0	615	6.8	6.3-7.4	436	13.0	11.7-14.3	179	3.3	2.8-3.8
1982	2311	24.3	23.3-25.3	1691	43.7	41.5-45.9	620	11.2	10.3-12.1	564	6.2	5.7-6.7	397	11.5	10.3-12.7	167	3.0	2.5-3.5
1983	2235	23.2	22.2-24.2	1668	43.1	41.0-45.3	567	10.1	9.3-11.0	560	6.0	5.6-6.6	409	12.1	10.9-13.4	151	2.6	2.2-3.1
1984	2254	23.2	22.2-24.2	1633	41.5	39.4-43.7	621	11.0	10.2-11.9	570	6.2	5.7-6.7	389	11.6	10.4-12.9	181	3.1	2.7-3.6
1985	2384	24.4	23.4-25.4	1745	44.1	41.9-46.3	639	11.3	10.4-12.2	562	6.0	5.5-6.5	384	10.9	9.8-12.1	178	3.1	2.6-3.6
1986	2471	24.9	23.9-25.9	1820	45.0	42.9-47.2	651	11.4	10.5-12.3	574	6.1	5.6-6.6	403	11.9	10.7-13.2	171	2.9	2.5-3.4
1987	2325	23.2	22.3-24.2	1673	41.0	38.9-43.1	652	11.2	10.3-12.1	530	5.5	5.1-6.0	356	10.0	9.0-11.2	174	2.9	2.5-3.4
1988	2336	23.3	22.4-24.3	1699	41.7	39.7-43.9	637	10.9	10.1-11.8	558	5.7	5.2-6.2	374	10.1	9.1-11.3	184	3.0	2.6-3.5
1989	2397	23.6	22.7-24.6	1755	42.6	40.5-44.7	642	10.8	10.0-11.7	541	5.5	5.0-5.9	388	10.7	9.6-11.8	153	2.5	2.1-2.9
1990	2498	24.5	23.5-25.5	1792	43.0	41.0-45.1	706	12.0	11.1-12.9	550	5.5	5.1-6.0	361	9.9	8.8-11.0	189	3.0	2.6-3.5
1991	2432	23.5	22.6-24.5	1765	41.7	39.7-43.7	667	11.1	10.3-12.0	539	5.3	4.8-5.8	377	10.0	9.0-11.1	162	2.6	2.2-3.0
1992	2456	23.6	22.7-24.5	1770	41.7	39.8-43.8	686	11.4	10.6-12.3	566	5.5	5.0-5.9	338	8.9	7.9-9.9	228	3.6	3.1-4.1
1993	2607	24.7	23.8-25.7	1833	42.3	40.3-44.3	774	12.6	11.7-13.6	557	5.3	4.9-5.8	374	9.8	8.8-10.9	183	2.8	2.4-3.2
1994	2468	23.4	22.5-24.4	1778	40.9	39.0-42.9	690	11.3	10.5-12.2	572	5.4	5.0-5.9	383	9.7	8.7-10.8	189	2.8	2.5-3.3
1995	2636	24.8	23.9-25.8	1856	42.3	40.4-44.3	780	12.7	11.8-13.6	547	5.1	4.7-5.6	363	8.9	8.0-9.9	184	2.8	2.4-3.2
1996	2865	26.8	25.8-27.8	2107	47.7	45.6-49.8	758	12.3	11.4-13.2	601	5.6	5.1-6.0	401	10.0	9.0-11.0	200	3.0	2.6-3.5
1997	2988	27.7	26.8-28.8	2148	48.0	46.0-50.1	840	13.4	12.5-14.4	624	5.7	5.3-6.2	418	10.0	9.1-11.0	206	3.1	2.7-3.5
1998	3092	28.5	27.5-29.5	2263	49.9	47.8-52.0	829	13.4	12.5-14.4	579	5.2	4.8-5.7	409	9.9	8.9-10.9	170	2.5	2.1-2.9
1999	3128	28.6	27.6-29.6	2268	49.4	47.4-51.5	860	13.7	12.8-14.6	590	5.3	4.8-5.7	407	9.6	8.7-10.6	183	2.6	2.3-3.1
2000	3139	28.3	27.3-29.3	2270	48.9	46.9-51.0	869	13.5	12.6-14.4	601	5.3	4.9-5.7	415	9.4	8.5-10.4	186	2.7	2.3-3.1
2001	3144	28.1	27.1-29.1	2307	49.4	47.4-51.5	837	13.0	12.2-14.0	613	5.3	4.9-5.8	432	9.9	9.0-10.9	181	2.5	2.2-2.9
2002	3309	29.3	28.4-30.4	2470	52.0	49.9-54.1	839	13.0	12.1-14.0	609	5.3	4.9-5.7	407	9.1	8.2-10.1	202	2.9	2.5-3.4
2003	3364	29.6	28.6-30.6	2487	51.4	49.4-53.5	877	13.5	12.6-14.5	563	4.8	4.4-5.2	381	8.4	7.6-9.3	182	2.5	2.2-3.0
2004	3473	30.3	29.3-31.3	2571	53.0	50.9-55.1	902	13.9	13.0-14.8	605	5.1	4.7-5.6	419	9.0	8.1-9.9	186	2.6	2.2-3.0
2005	3521	30.5	29.5-31.5	2598	52.9	50.8-55.0	923	13.9	13.0-14.9	565	4.7	4.3-5.1	383	8.1	7.3-9.0	182	2.5	2.1-2.9
2006	3444	29.5	28.5-30.5	2550	51.3	49.3-53.3	894	13.6	12.7-14.5	576	4.8	4.4-5.2	394	8.3	7.5-9.1	182	2.5	2.2-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Urinary bladder (incl. in situ)

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	3440	29.2	28.2-30.2	2539	50.1	48.1-52.1	901	13.7	12.8-14.6	640	5.2	4.8-5.7	458	9.4	8.5-10.3	182	2.5	2.1-2.9
2008	3558	29.7	28.7-30.7	2679	52.2	50.2-54.2	879	12.9	12.1-13.8	649	5.2	4.8-5.6	446	9.0	8.2-9.9	203	2.7	2.3-3.1
2009	3599	29.6	28.7-30.6	2702	51.7	49.7-53.7	897	13.0	12.2-14.0	641	5.1	4.7-5.5	485	9.7	8.8-10.6	156	2.0	1.7-2.4
2010	3371	27.4	26.5-28.4	2551	48.1	46.2-50.0	820	11.8	11.0-12.7	666	5.2	4.8-5.6	467	9.0	8.2-9.9	199	2.6	2.2-3.0
2011	3641	29.5	28.6-30.5	2718	50.5	48.6-52.5	923	13.5	12.6-14.4	617	4.7	4.3-5.1	437	8.4	7.6-9.2	180	2.3	2.0-2.7
2012	3739	29.6	28.7-30.6	2768	50.7	48.8-52.7	971	13.7	12.8-14.6	641	4.9	4.5-5.3	443	8.3	7.5-9.1	198	2.5	2.2-2.9
2013	3773	29.7	28.7-30.7	2817	51.1	49.2-53.1	956	13.5	12.6-14.4	643	4.9	4.5-5.3	465	8.6	7.8-9.5	178	2.3	1.9-2.6
2014	3690	28.4	27.5-29.4	2766	48.9	47.0-50.8	924	12.6	11.8-13.5	680	5.0	4.7-5.4	461	8.4	7.6-9.2	219	2.8	2.5-3.3
2015	3565	27.2	26.3-28.2	2661	46.2	44.5-48.1	904	12.6	11.8-13.5	633	4.6	4.3-5.0	438	7.8	7.1-8.6	195	2.5	2.2-2.9
2016	3681	27.7	26.8-28.7	2774	47.4	45.6-49.3	907	12.2	11.4-13.1	636	4.6	4.3-5.0	456	8.1	7.4-8.9	180	2.2	1.9-2.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Kidney and renal pelvis

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	643	7.0	6.4-7.6	402	10.2	9.2-11.3	241	4.7	4.1-5.3	357	3.9	3.5-4.3	223	5.7	4.9-6.5	134	2.6	2.1-3.1
1977	702	7.6	7.1-8.2	432	10.8	9.8-12.0	270	5.2	4.6-5.8	400	4.4	4.0-4.9	243	6.3	5.5-7.1	157	3.0	2.6-3.5
1978	699	7.6	7.0-8.2	429	10.8	9.7-11.9	270	5.2	4.6-5.9	354	3.8	3.4-4.2	217	5.5	4.8-6.4	137	2.6	2.1-3.0
1979	712	7.6	7.0-8.2	435	10.7	9.7-11.8	277	5.1	4.5-5.8	409	4.4	4.0-4.8	255	6.4	5.6-7.2	154	2.9	2.5-3.4
1980	725	7.7	7.1-8.3	437	10.7	9.7-11.9	288	5.4	4.8-6.1	406	4.3	3.8-4.7	240	6.1	5.3-7.0	166	3.0	2.5-3.5
1981	796	8.5	7.9-9.1	473	11.9	10.8-13.1	323	6.1	5.4-6.8	385	4.1	3.7-4.5	238	6.0	5.2-6.9	147	2.7	2.3-3.1
1982	835	8.6	8.0-9.2	508	12.0	10.9-13.2	327	6.0	5.4-6.7	390	4.0	3.6-4.4	234	5.8	5.0-6.6	156	2.8	2.4-3.3
1983	901	9.3	8.7-9.9	552	13.2	12.1-14.4	349	6.4	5.7-7.1	398	4.1	3.7-4.6	234	5.7	5.0-6.5	164	3.0	2.5-3.5
1984	892	9.2	8.6-9.8	561	13.5	12.4-14.7	331	6.0	5.4-6.8	400	4.2	3.8-4.6	249	6.4	5.6-7.3	151	2.7	2.3-3.1
1985	939	9.6	9.0-10.2	571	13.6	12.5-14.8	368	6.6	5.9-7.3	432	4.5	4.0-4.9	246	6.2	5.4-7.0	186	3.3	2.8-3.8
1986	953	9.8	9.2-10.4	592	14.1	12.9-15.3	361	6.7	6.0-7.4	449	4.6	4.1-5.0	269	6.5	5.7-7.4	180	3.2	2.7-3.7
1987	1024	10.4	9.7-11.0	606	14.3	13.1-15.5	418	7.5	6.8-8.3	463	4.7	4.3-5.2	287	7.1	6.3-8.1	176	3.0	2.6-3.5
1988	995	10.0	9.4-10.6	594	13.9	12.8-15.1	401	7.1	6.4-7.8	446	4.4	4.0-4.9	253	6.1	5.3-6.9	193	3.3	2.8-3.8
1989	980	9.8	9.2-10.5	613	14.5	13.3-15.7	367	6.4	5.8-7.2	433	4.3	3.9-4.8	262	6.4	5.6-7.3	171	2.9	2.5-3.4
1990	1097	10.9	10.2-11.5	687	15.9	14.7-17.2	410	7.2	6.5-7.9	417	4.1	3.7-4.5	253	5.9	5.2-6.7	164	2.7	2.3-3.2
1991	1119	11.0	10.4-11.7	677	15.4	14.2-16.6	442	7.8	7.1-8.5	464	4.5	4.1-5.0	279	6.6	5.8-7.5	185	3.0	2.6-3.5
1992	1139	11.1	10.5-11.8	686	15.3	14.2-16.6	453	8.0	7.2-8.7	425	4.1	3.7-4.5	257	6.1	5.4-7.0	168	2.8	2.4-3.2
1993	1176	11.4	10.7-12.0	701	15.6	14.4-16.8	475	8.2	7.4-8.9	410	3.9	3.6-4.3	250	5.7	5.0-6.4	160	2.6	2.2-3.0
1994	1165	11.2	10.6-11.9	695	15.4	14.3-16.6	470	8.1	7.4-8.9	467	4.5	4.1-4.9	290	6.8	6.0-7.7	177	2.9	2.5-3.3
1995	1208	11.6	10.9-12.3	716	15.6	14.4-16.8	492	8.4	7.7-9.2	445	4.2	3.8-4.6	250	5.8	5.1-6.6	195	3.2	2.7-3.7
1996	1285	12.3	11.6-13.0	794	17.3	16.1-18.5	491	8.3	7.6-9.1	473	4.5	4.1-4.9	275	6.2	5.5-7.0	198	3.2	2.8-3.7
1997	1372	13.0	12.3-13.7	865	18.8	17.5-20.1	507	8.5	7.8-9.3	449	4.2	3.8-4.6	283	6.4	5.6-7.2	166	2.7	2.3-3.1
1998	1438	13.6	12.9-14.3	881	18.7	17.4-19.9	557	9.4	8.6-10.3	460	4.2	3.9-4.7	290	6.3	5.6-7.1	170	2.6	2.2-3.1
1999	1511	14.1	13.4-14.8	927	19.5	18.3-20.9	584	9.8	9.0-10.7	384	3.5	3.2-3.9	237	5.2	4.5-5.9	147	2.2	1.9-2.7
2000	1511	13.9	13.2-14.7	898	18.7	17.5-20.0	613	10.1	9.3-11.0	447	4.0	3.7-4.4	269	5.9	5.2-6.6	178	2.7	2.4-3.2
2001	1608	14.7	14.0-15.5	997	20.6	19.3-21.9	611	10.1	9.3-10.9	474	4.2	3.9-4.6	278	6.0	5.3-6.7	196	3.1	2.7-3.6
2002	1622	14.7	14.0-15.4	1014	20.8	19.5-22.1	608	9.9	9.1-10.7	435	3.8	3.5-4.2	257	5.5	4.8-6.2	178	2.7	2.3-3.1
2003	1678	15.1	14.4-15.8	1048	20.9	19.6-22.2	630	10.2	9.4-11.1	455	4.0	3.6-4.4	270	5.5	4.9-6.2	185	2.7	2.3-3.1
2004	1856	16.6	15.8-17.4	1152	22.8	21.5-24.2	704	11.5	10.7-12.4	472	4.1	3.7-4.5	283	5.8	5.1-6.5	189	2.8	2.4-3.3
2005	1857	16.5	15.8-17.3	1138	22.5	21.2-23.9	719	11.7	10.9-12.6	469	4.0	3.7-4.4	302	6.1	5.4-6.8	167	2.5	2.1-2.9
2006	2013	17.6	16.9-18.4	1278	24.8	23.4-26.2	735	11.7	10.9-12.7	472	4.0	3.6-4.4	288	5.8	5.2-6.5	184	2.7	2.3-3.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Kidney and renal pelvis

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	2118	18.5	17.7-19.3	1354	25.9	24.6-27.4	764	12.4	11.5-13.3	449	3.8	3.4-4.2	279	5.5	4.9-6.2	170	2.5	2.1-2.9
2008	2047	17.6	16.8-18.4	1281	24.1	22.8-25.5	766	12.2	11.3-13.1	461	3.8	3.5-4.2	304	5.9	5.2-6.6	157	2.2	1.9-2.6
2009	2023	17.3	16.5-18.1	1257	23.4	22.1-24.7	766	12.2	11.3-13.2	440	3.6	3.3-4.0	281	5.3	4.7-6.0	159	2.3	1.9-2.7
2010	2080	17.5	16.7-18.3	1302	23.9	22.6-25.3	778	12.1	11.3-13.0	437	3.5	3.2-3.9	280	5.1	4.5-5.8	157	2.2	1.9-2.6
2011	2154	18.0	17.3-18.8	1407	25.4	24.1-26.8	747	11.7	10.9-12.6	475	3.8	3.4-4.2	292	5.5	4.8-6.1	183	2.5	2.2-3.0
2012	2240	18.4	17.6-19.2	1421	25.3	24.0-26.7	819	12.5	11.7-13.5	495	3.9	3.5-4.2	326	5.9	5.2-6.6	169	2.3	2.0-2.7
2013	2220	18.2	17.4-19.0	1430	25.1	23.8-26.5	790	12.2	11.3-13.1	444	3.4	3.1-3.8	291	5.2	4.6-5.9	153	2.1	1.8-2.5
2014	2291	18.5	17.7-19.3	1475	25.5	24.2-26.9	816	12.4	11.5-13.3	427	3.2	2.9-3.5	277	4.7	4.2-5.3	150	2.0	1.7-2.4
2015	2315	18.4	17.7-19.2	1499	25.7	24.4-27.1	816	12.1	11.2-13.0	450	3.4	3.0-3.7	283	4.9	4.3-5.5	167	2.2	1.9-2.6
2016	2262	18.1	17.3-18.9	1505	25.7	24.3-27.1	757	11.5	10.6-12.4	385	2.9	2.6-3.2	248	4.1	3.6-4.7	137	1.9	1.6-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Brain and other central nervous system

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	555	5.8	5.3-6.3	322	7.2	6.4-8.0	233	4.6	4.0-5.3	415	4.4	3.9-4.8	224	5.1	4.4-5.8	191	3.7	3.2-4.3
1977	570	5.9	5.4-6.4	323	7.3	6.5-8.1	247	4.7	4.2-5.4	428	4.4	4.0-4.8	239	5.4	4.7-6.2	189	3.6	3.1-4.1
1978	540	5.6	5.2-6.1	285	6.4	5.7-7.3	255	5.0	4.4-5.7	398	4.1	3.7-4.5	219	4.9	4.3-5.7	179	3.3	2.9-3.9
1979	617	6.4	5.9-6.9	354	8.0	7.1-8.9	263	5.0	4.4-5.7	404	4.1	3.7-4.6	229	5.1	4.5-5.9	175	3.3	2.8-3.9
1980	619	6.3	5.8-6.8	356	7.7	6.9-8.6	263	5.1	4.5-5.8	455	4.7	4.2-5.1	274	6.2	5.5-7.0	181	3.4	2.9-4.0
1981	612	6.3	5.8-6.9	342	7.8	7.0-8.7	270	5.1	4.5-5.8	421	4.3	3.9-4.7	238	5.5	4.8-6.3	183	3.4	2.9-4.0
1982	653	6.8	6.2-7.3	334	7.6	6.8-8.5	319	6.1	5.5-6.8	435	4.5	4.0-4.9	223	5.1	4.4-5.8	212	4.0	3.4-4.5
1983	629	6.5	6.0-7.1	315	7.1	6.3-8.0	314	6.0	5.4-6.8	425	4.3	3.9-4.7	225	5.1	4.4-5.8	200	3.6	3.1-4.1
1984	698	7.1	6.6-7.6	380	8.6	7.7-9.5	318	5.9	5.3-6.7	430	4.4	4.0-4.8	235	5.5	4.8-6.2	195	3.5	3.1-4.1
1985	660	6.8	6.2-7.3	354	7.9	7.1-8.8	306	5.8	5.1-6.5	413	4.3	3.9-4.7	206	4.8	4.2-5.6	207	3.9	3.4-4.5
1986	688	7.0	6.5-7.6	374	8.4	7.5-9.3	314	5.9	5.2-6.6	458	4.6	4.2-5.1	251	5.7	5.0-6.5	207	3.7	3.2-4.3
1987	737	7.5	7.0-8.1	389	8.7	7.8-9.6	348	6.5	5.8-7.2	458	4.6	4.2-5.1	234	5.3	4.6-6.1	224	4.0	3.5-4.5
1988	717	7.2	6.7-7.8	398	8.7	7.9-9.7	319	6.0	5.3-6.7	488	4.8	4.3-5.2	273	6.0	5.3-6.8	215	3.8	3.3-4.3
1989	709	7.1	6.6-7.7	408	8.8	8.0-9.8	301	5.7	5.1-6.4	457	4.5	4.1-5.0	254	5.7	5.0-6.4	203	3.6	3.2-4.2
1990	764	7.7	7.1-8.2	423	9.4	8.5-10.4	341	6.3	5.6-7.0	419	4.1	3.8-4.6	250	5.6	4.9-6.4	169	2.9	2.5-3.4
1991	719	7.2	6.7-7.7	395	8.8	7.9-9.7	324	6.0	5.4-6.7	445	4.5	4.0-4.9	262	6.0	5.3-6.8	183	3.4	2.9-3.9
1992	738	7.3	6.8-7.9	386	8.5	7.7-9.4	352	6.5	5.8-7.2	475	4.6	4.2-5.1	241	5.3	4.6-6.0	234	4.2	3.6-4.7
1993	737	7.3	6.8-7.9	409	8.8	8.0-9.7	328	6.0	5.4-6.7	455	4.4	4.0-4.9	248	5.4	4.8-6.1	207	3.6	3.1-4.2
1994	804	7.9	7.3-8.5	454	9.9	9.0-10.8	350	6.2	5.6-6.9	529	5.1	4.7-5.6	293	6.4	5.7-7.2	236	4.1	3.6-4.6
1995	852	8.4	7.8-9.0	421	9.0	8.1-9.9	431	7.8	7.1-8.6	467	4.5	4.1-5.0	253	5.5	4.8-6.2	214	3.7	3.2-4.3
1996	836	8.2	7.7-8.8	459	10.0	9.1-10.9	377	6.9	6.2-7.6	514	5.0	4.6-5.4	284	6.2	5.5-7.0	230	4.0	3.5-4.6
1997	791	7.7	7.2-8.3	442	9.4	8.5-10.3	349	6.3	5.6-7.0	490	4.7	4.3-5.1	272	5.8	5.1-6.5	218	3.8	3.3-4.3
1998	838	8.2	7.6-8.7	469	9.8	8.9-10.7	369	6.7	6.1-7.5	476	4.5	4.1-5.0	271	5.7	5.0-6.4	205	3.5	3.0-4.0
1999	780	7.6	7.0-8.1	413	8.7	7.9-9.6	367	6.6	5.9-7.3	483	4.6	4.2-5.0	269	5.6	5.0-6.4	214	3.7	3.2-4.2
2000	874	8.4	7.9-9.0	458	9.6	8.7-10.5	416	7.3	6.6-8.1	468	4.4	4.0-4.8	243	5.0	4.4-5.7	225	3.7	3.2-4.2
2001	815	7.8	7.3-8.4	444	9.3	8.4-10.2	371	6.6	5.9-7.3	462	4.3	3.9-4.7	242	5.0	4.4-5.7	220	3.7	3.2-4.2
2002	823	7.9	7.3-8.4	467	9.7	8.9-10.7	356	6.3	5.7-7.0	506	4.6	4.2-5.1	286	5.8	5.2-6.6	220	3.7	3.2-4.2
2003	848	8.1	7.6-8.7	438	9.0	8.1-9.9	410	7.5	6.7-8.2	457	4.2	3.8-4.6	256	5.2	4.5-5.8	201	3.4	3.0-4.0
2004	803	7.6	7.1-8.2	449	9.2	8.4-10.1	354	6.3	5.6-7.0	510	4.6	4.2-5.1	274	5.5	4.8-6.2	236	3.9	3.4-4.4
2005	839	8.0	7.5-8.6	482	9.9	9.0-10.8	357	6.4	5.7-7.1	507	4.6	4.2-5.0	278	5.6	4.9-6.3	229	3.8	3.3-4.3
2006	817	7.7	7.2-8.3	475	9.5	8.7-10.4	342	6.1	5.4-6.8	473	4.2	3.8-4.6	284	5.6	4.9-6.3	189	3.1	2.6-3.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Brain and other central nervous system

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	802	7.6	7.1-8.1	436	8.8	8.0-9.7	366	6.6	5.9-7.3	485	4.3	3.9-4.7	294	5.7	5.1-6.4	191	3.1	2.7-3.6
2008	817	7.7	7.2-8.3	458	9.3	8.4-10.2	359	6.3	5.7-7.1	481	4.2	3.8-4.6	283	5.5	4.9-6.2	198	3.1	2.7-3.6
2009	832	7.7	7.2-8.3	447	8.8	8.0-9.7	385	6.8	6.1-7.5	538	4.6	4.3-5.1	298	5.6	5.0-6.3	240	3.8	3.3-4.4
2010	807	7.5	6.9-8.0	446	8.9	8.0-9.7	361	6.3	5.6-7.0	515	4.5	4.1-4.9	285	5.5	4.8-6.1	230	3.6	3.2-4.2
2011	802	7.4	6.9-7.9	451	8.8	8.0-9.6	351	6.2	5.5-6.9	472	4.0	3.6-4.3	259	4.8	4.2-5.4	213	3.2	2.8-3.7
2012	840	7.6	7.1-8.2	465	8.8	8.0-9.7	375	6.6	5.9-7.3	570	4.8	4.4-5.2	322	6.0	5.3-6.7	248	3.9	3.4-4.4
2013	820	7.4	6.9-8.0	483	9.1	8.3-10.0	337	5.8	5.1-6.5	531	4.3	3.9-4.7	295	5.2	4.6-5.8	236	3.6	3.1-4.1
2014	893	8.0	7.5-8.6	487	9.2	8.3-10.1	406	7.1	6.4-7.9	552	4.5	4.1-4.9	311	5.4	4.8-6.1	241	3.7	3.3-4.3
2015	835	7.5	7.0-8.1	441	8.3	7.5-9.2	394	6.8	6.1-7.6	537	4.4	4.0-4.8	282	5.0	4.4-5.6	255	3.8	3.4-4.4
2016	816	7.2	6.7-7.7	479	8.8	8.0-9.7	337	5.7	5.1-6.4	548	4.4	4.0-4.8	324	5.6	5.0-6.3	224	3.4	3.0-3.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Thyroid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	297	3.3	3.0-3.8	99	2.4	2.0-3.0	198	4.2	3.6-4.8	52	0.6	0.5-0.8	20	0.6	0.3-0.9	32	0.6	0.4-0.9
1977	342	3.8	3.4-4.2	106	2.5	2.0-3.1	236	4.9	4.3-5.6	67	0.8	0.6-1.0	21	0.6	0.3-0.9	46	0.9	0.6-1.2
1978	327	3.5	3.2-4.0	94	2.3	1.8-2.8	233	4.7	4.1-5.4	58	0.7	0.5-0.9	20	0.6	0.3-0.9	38	0.7	0.5-1.0
1979	308	3.3	3.0-3.8	92	2.2	1.8-2.8	216	4.3	3.8-5.0	43	0.5	0.3-0.6	13	0.3	0.2-0.6	30	0.5	0.4-0.8
1980	334	3.5	3.2-4.0	104	2.4	2.0-3.0	230	4.5	4.0-5.2	69	0.8	0.6-1.0	33	0.9	0.6-1.2	36	0.7	0.5-0.9
1981	329	3.5	3.1-3.9	101	2.4	2.0-3.0	228	4.5	4.0-5.2	54	0.6	0.4-0.8	24	0.6	0.4-0.9	30	0.6	0.4-0.8
1982	310	3.3	3.0-3.7	96	2.2	1.7-2.7	214	4.4	3.8-5.0	51	0.6	0.4-0.7	26	0.7	0.4-1.0	25	0.5	0.3-0.7
1983	364	3.8	3.4-4.3	100	2.3	1.8-2.8	264	5.2	4.6-5.9	51	0.6	0.4-0.7	14	0.4	0.2-0.6	37	0.7	0.5-0.9
1984	388	4.1	3.7-4.5	112	2.6	2.2-3.2	276	5.5	4.8-6.2	64	0.7	0.5-0.9	24	0.7	0.4-1.0	40	0.7	0.5-0.9
1985	390	4.1	3.7-4.5	109	2.4	2.0-3.0	281	5.6	5.0-6.4	45	0.5	0.4-0.7	18	0.5	0.3-0.8	27	0.5	0.3-0.7
1986	402	4.2	3.8-4.6	119	2.7	2.2-3.2	283	5.6	5.0-6.3	52	0.6	0.4-0.7	24	0.6	0.4-1.0	28	0.5	0.3-0.7
1987	401	4.2	3.8-4.6	131	2.9	2.4-3.4	270	5.3	4.7-6.0	50	0.5	0.4-0.7	15	0.3	0.2-0.5	35	0.6	0.4-0.8
1988	391	4.1	3.7-4.5	102	2.3	1.9-2.8	289	5.8	5.1-6.5	48	0.5	0.4-0.6	18	0.4	0.2-0.7	30	0.5	0.3-0.7
1989	410	4.2	3.8-4.7	114	2.5	2.1-3.1	296	5.8	5.2-6.5	54	0.5	0.4-0.7	17	0.4	0.2-0.7	37	0.6	0.4-0.8
1990	414	4.3	3.9-4.7	129	2.9	2.4-3.5	285	5.5	4.9-6.2	46	0.4	0.3-0.6	21	0.5	0.3-0.7	25	0.4	0.3-0.6
1991	475	4.8	4.4-5.3	144	3.1	2.6-3.7	331	6.4	5.7-7.1	42	0.4	0.3-0.6	15	0.4	0.2-0.6	27	0.5	0.3-0.7
1992	479	4.8	4.4-5.3	135	3.0	2.5-3.5	344	6.5	5.9-7.3	55	0.5	0.4-0.7	16	0.4	0.2-0.6	39	0.6	0.4-0.8
1993	489	4.9	4.5-5.4	133	2.8	2.4-3.4	356	7.0	6.3-7.8	57	0.6	0.4-0.7	21	0.5	0.3-0.8	36	0.6	0.4-0.8
1994	542	5.5	5.0-5.9	138	3.0	2.5-3.5	404	7.9	7.1-8.7	50	0.5	0.4-0.6	23	0.6	0.4-0.8	27	0.4	0.3-0.6
1995	620	6.2	5.7-6.7	170	3.7	3.1-4.3	450	8.8	8.0-9.6	55	0.5	0.4-0.7	24	0.5	0.3-0.8	31	0.5	0.3-0.7
1996	681	6.9	6.4-7.4	182	3.8	3.3-4.4	499	9.8	9.0-10.7	81	0.8	0.6-1.0	27	0.6	0.4-0.9	54	0.9	0.7-1.2
1997	744	7.5	7.0-8.0	201	4.2	3.7-4.9	543	10.6	9.7-11.5	55	0.5	0.4-0.7	23	0.5	0.3-0.8	32	0.5	0.3-0.7
1998	746	7.5	7.0-8.1	211	4.4	3.8-5.1	535	10.4	9.6-11.4	55	0.5	0.4-0.7	17	0.4	0.2-0.6	38	0.6	0.4-0.8
1999	901	9.0	8.4-9.6	217	4.5	3.9-5.1	684	13.5	12.5-14.5	54	0.5	0.4-0.6	20	0.4	0.3-0.7	34	0.5	0.3-0.7
2000	1034	10.3	9.7-11.0	266	5.5	4.9-6.2	768	15.1	14.0-16.2	48	0.4	0.3-0.6	17	0.4	0.2-0.6	31	0.5	0.3-0.7
2001	1047	10.4	9.8-11.1	225	4.6	4.0-5.2	822	16.1	15.0-17.3	54	0.5	0.4-0.6	23	0.5	0.3-0.7	31	0.5	0.3-0.7
2002	1083	10.7	10.1-11.4	281	5.7	5.0-6.4	802	15.6	14.5-16.7	47	0.4	0.3-0.6	18	0.4	0.2-0.6	29	0.5	0.3-0.7
2003	1131	11.1	10.5-11.8	307	6.2	5.5-6.9	824	15.9	14.8-17.0	54	0.5	0.4-0.6	14	0.3	0.2-0.5	40	0.6	0.4-0.8
2004	1247	12.2	11.5-12.9	299	5.9	5.2-6.6	948	18.3	17.2-19.5	47	0.4	0.3-0.5	20	0.4	0.3-0.7	27	0.4	0.2-0.6
2005	1395	13.7	13.0-14.5	354	7.0	6.3-7.8	1041	20.3	19.0-21.6	46	0.4	0.3-0.5	24	0.5	0.3-0.7	22	0.3	0.2-0.5
2006	1526	15.0	14.2-15.8	387	7.8	7.0-8.6	1139	22.1	20.8-23.4	55	0.5	0.3-0.6	25	0.5	0.3-0.7	30	0.4	0.3-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Thyroid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	1657	16.1	15.3-16.9	435	8.7	7.9-9.5	1222	23.3	22.0-24.7	54	0.5	0.3-0.6	26	0.5	0.3-0.7	28	0.4	0.3-0.6
2008	1838	17.9	17.1-18.8	432	8.5	7.7-9.3	1406	27.2	25.7-28.7	58	0.5	0.4-0.6	26	0.5	0.3-0.7	32	0.5	0.3-0.7
2009	1905	18.6	17.8-19.5	491	9.7	8.8-10.6	1414	27.3	25.8-28.8	65	0.5	0.4-0.7	26	0.5	0.3-0.7	39	0.5	0.4-0.7
2010	1954	19.0	18.1-19.8	503	9.6	8.8-10.6	1451	28.1	26.6-29.6	59	0.5	0.3-0.6	25	0.5	0.3-0.7	34	0.4	0.3-0.6
2011	2078	20.2	19.3-21.1	562	10.9	10.0-11.9	1516	29.3	27.8-30.9	74	0.6	0.5-0.8	39	0.8	0.5-1.0	35	0.5	0.3-0.7
2012	2073	20.1	19.2-21.0	559	10.6	9.7-11.5	1514	29.6	28.1-31.2	66	0.5	0.4-0.7	28	0.5	0.3-0.7	38	0.5	0.4-0.7
2013	2122	20.5	19.6-21.5	596	11.3	10.4-12.3	1526	29.8	28.2-31.4	64	0.5	0.4-0.6	28	0.5	0.3-0.7	36	0.5	0.3-0.7
2014	2108	20.5	19.6-21.5	583	11.0	10.1-12.0	1525	30.0	28.5-31.7	52	0.4	0.3-0.5	26	0.4	0.3-0.6	26	0.3	0.2-0.5
2015	2143	21.1	20.2-22.1	577	11.0	10.1-12.0	1566	31.2	29.6-32.8	68	0.5	0.4-0.7	33	0.6	0.4-0.8	35	0.5	0.3-0.7
2016	1932	19.1	18.2-20.0	543	10.4	9.5-11.3	1389	27.7	26.2-29.3	64	0.5	0.4-0.6	24	0.4	0.3-0.7	40	0.5	0.4-0.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Hodgkin lymphoma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	347	3.4	3.1-3.8	191	4.0	3.4-4.6	156	2.9	2.5-3.4	139	1.5	1.2-1.8	75	1.8	1.4-2.2	64	1.2	1.0-1.6
1977	326	3.3	2.9-3.7	182	3.9	3.3-4.5	144	2.8	2.3-3.3	137	1.5	1.2-1.8	79	1.9	1.5-2.4	58	1.1	0.8-1.4
1978	355	3.5	3.1-3.9	185	3.9	3.4-4.6	170	3.1	2.7-3.6	119	1.3	1.0-1.5	66	1.6	1.2-2.0	53	1.0	0.8-1.4
1979	377	3.7	3.4-4.2	213	4.6	3.9-5.2	164	3.0	2.6-3.5	108	1.1	0.9-1.4	60	1.4	1.1-1.8	48	0.9	0.7-1.2
1980	397	3.9	3.5-4.3	223	4.7	4.1-5.4	174	3.2	2.7-3.7	131	1.4	1.1-1.6	77	1.8	1.4-2.3	54	1.0	0.8-1.3
1981	379	3.7	3.4-4.1	199	4.1	3.6-4.8	180	3.4	2.9-3.9	112	1.1	0.9-1.4	65	1.5	1.1-1.9	47	0.9	0.6-1.2
1982	368	3.6	3.2-4.0	180	3.8	3.3-4.5	188	3.4	2.9-3.9	131	1.4	1.1-1.6	71	1.6	1.2-2.1	60	1.2	0.9-1.5
1983	351	3.5	3.1-3.9	197	4.2	3.6-4.8	154	2.9	2.4-3.4	90	0.9	0.7-1.1	52	1.1	0.8-1.5	38	0.7	0.5-0.9
1984	380	3.8	3.4-4.2	199	4.3	3.7-5.0	181	3.3	2.8-3.8	94	1.0	0.8-1.2	59	1.4	1.0-1.8	35	0.6	0.4-0.9
1985	394	3.9	3.5-4.3	202	4.2	3.6-4.8	192	3.6	3.1-4.2	97	1.0	0.8-1.2	54	1.2	0.9-1.5	43	0.8	0.6-1.0
1986	353	3.4	3.1-3.8	177	3.6	3.0-4.1	176	3.3	2.8-3.8	118	1.2	1.0-1.4	61	1.4	1.1-1.8	57	1.0	0.8-1.3
1987	347	3.4	3.0-3.8	179	3.7	3.2-4.3	168	3.1	2.7-3.7	100	1.0	0.8-1.2	62	1.3	1.0-1.7	38	0.7	0.5-1.0
1988	362	3.6	3.2-4.0	200	4.2	3.6-4.8	162	3.1	2.6-3.6	80	0.8	0.6-1.0	54	1.2	0.9-1.5	26	0.5	0.3-0.7
1989	375	3.7	3.3-4.1	206	4.3	3.7-4.9	169	3.1	2.7-3.7	93	1.0	0.8-1.2	63	1.4	1.1-1.9	30	0.6	0.4-0.8
1990	362	3.6	3.2-4.0	189	4.0	3.4-4.6	173	3.3	2.8-3.8	82	0.8	0.7-1.0	46	1.0	0.8-1.4	36	0.7	0.5-0.9
1991	352	3.5	3.1-3.9	189	3.9	3.3-4.5	163	3.1	2.6-3.6	67	0.7	0.5-0.9	36	0.8	0.6-1.1	31	0.6	0.4-0.8
1992	378	3.8	3.4-4.2	196	4.2	3.6-4.8	182	3.5	3.0-4.0	89	0.9	0.7-1.1	42	0.9	0.6-1.2	47	0.8	0.6-1.1
1993	349	3.5	3.2-3.9	169	3.5	3.0-4.1	180	3.5	3.0-4.1	68	0.7	0.5-0.8	36	0.8	0.5-1.1	32	0.6	0.4-0.8
1994	393	4.0	3.6-4.4	196	4.1	3.6-4.7	197	3.8	3.3-4.4	72	0.7	0.6-0.9	37	0.8	0.6-1.1	35	0.6	0.4-0.9
1995	331	3.4	3.0-3.8	182	3.8	3.3-4.4	149	2.9	2.5-3.5	69	0.7	0.5-0.9	33	0.7	0.5-1.0	36	0.6	0.4-0.9
1996	349	3.6	3.2-4.0	175	3.7	3.2-4.3	174	3.5	3.0-4.1	58	0.6	0.4-0.8	29	0.7	0.4-0.9	29	0.6	0.4-0.8
1997	320	3.3	3.0-3.7	174	3.7	3.2-4.3	146	3.0	2.5-3.5	59	0.6	0.4-0.7	35	0.7	0.5-1.0	24	0.4	0.3-0.6
1998	355	3.7	3.3-4.1	176	3.8	3.2-4.4	179	3.6	3.1-4.2	58	0.6	0.4-0.7	31	0.6	0.4-0.9	27	0.5	0.3-0.8
1999	339	3.5	3.1-3.9	192	4.1	3.5-4.7	147	3.0	2.5-3.5	62	0.6	0.5-0.8	29	0.6	0.4-0.9	33	0.6	0.4-0.8
2000	329	3.4	3.0-3.8	182	3.9	3.4-4.5	147	2.9	2.5-3.5	50	0.5	0.4-0.6	25	0.5	0.3-0.8	25	0.4	0.3-0.6
2001	357	3.7	3.4-4.1	188	4.0	3.5-4.6	169	3.5	3.0-4.1	61	0.6	0.5-0.8	32	0.7	0.5-1.0	29	0.5	0.3-0.8
2002	385	4.0	3.6-4.4	220	4.7	4.1-5.4	165	3.4	2.9-4.0	64	0.6	0.5-0.8	27	0.6	0.4-0.8	37	0.7	0.5-0.9
2003	332	3.4	3.0-3.8	189	4.0	3.4-4.6	143	2.9	2.4-3.4	46	0.5	0.3-0.6	21	0.4	0.3-0.7	25	0.5	0.3-0.7
2004	370	3.8	3.4-4.2	189	4.0	3.4-4.6	181	3.7	3.2-4.3	52	0.5	0.4-0.6	31	0.6	0.4-0.9	21	0.3	0.2-0.5
2005	372	3.9	3.5-4.3	200	4.2	3.7-4.9	172	3.6	3.1-4.2	47	0.4	0.3-0.6	20	0.4	0.3-0.6	27	0.5	0.3-0.7
2006	339	3.5	3.1-3.9	188	3.9	3.4-4.5	151	3.0	2.6-3.6	56	0.5	0.4-0.7	29	0.6	0.4-0.8	27	0.5	0.3-0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Hodgkin lymphoma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	343	3.5	3.2-3.9	182	3.8	3.2-4.4	161	3.3	2.8-3.9	52	0.5	0.4-0.6	25	0.5	0.3-0.7	27	0.4	0.3-0.6
2008	346	3.5	3.2-3.9	191	4.0	3.4-4.6	155	3.1	2.6-3.7	51	0.5	0.3-0.6	28	0.6	0.4-0.8	23	0.4	0.2-0.5
2009	346	3.6	3.2-4.0	188	3.9	3.4-4.5	158	3.3	2.8-3.8	40	0.4	0.3-0.5	20	0.4	0.2-0.6	20	0.3	0.2-0.5
2010	333	3.3	3.0-3.7	193	4.0	3.4-4.6	140	2.8	2.3-3.3	44	0.4	0.3-0.5	23	0.5	0.3-0.7	21	0.4	0.2-0.6
2011	338	3.5	3.1-3.9	188	3.9	3.4-4.6	150	3.1	2.6-3.7	37	0.3	0.2-0.4	26	0.5	0.3-0.8	11	0.2	0.1-0.3
2012	355	3.7	3.3-4.2	216	4.6	4.0-5.3	139	2.9	2.4-3.5	47	0.4	0.3-0.5	20	0.4	0.2-0.6	27	0.4	0.3-0.6
2013	321	3.3	2.9-3.6	168	3.4	2.9-4.0	153	3.1	2.6-3.7	33	0.3	0.2-0.4	14	0.3	0.2-0.5	19	0.3	0.2-0.5
2014	370	3.7	3.4-4.2	196	4.1	3.5-4.7	174	3.4	2.9-4.0	34	0.3	0.2-0.4	15	0.3	0.1-0.5	19	0.3	0.2-0.5
2015	293	3.1	2.7-3.5	164	3.5	2.9-4.0	129	2.7	2.3-3.3	29	0.2	0.2-0.3	13	0.2	0.1-0.4	16	0.2	0.1-0.4
2016	339	3.5	3.1-3.9	170	3.5	3.0-4.1	169	3.5	2.9-4.0	40	0.3	0.2-0.5	23	0.4	0.3-0.6	17	0.3	0.2-0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Non-Hodgkin lymphomas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1022	11.2	10.5-12.0	531	13.4	12.2-14.6	491	9.6	8.8-10.5	572	6.3	5.8-6.9	302	7.8	6.9-8.7	270	5.2	4.6-5.9
1977	1088	11.7	11.0-12.5	558	14.0	12.8-15.3	530	10.1	9.3-11.0	618	6.7	6.2-7.3	306	7.7	6.9-8.7	312	5.9	5.3-6.6
1978	1203	12.9	12.2-13.7	640	15.6	14.4-16.9	563	10.7	9.8-11.6	618	6.8	6.2-7.3	312	8.2	7.3-9.2	306	5.8	5.1-6.5
1979	1113	12.0	11.3-12.7	565	14.1	12.9-15.3	548	10.3	9.5-11.3	630	6.8	6.3-7.4	340	8.6	7.7-9.6	290	5.4	4.8-6.1
1980	1204	12.8	12.0-13.5	621	15.0	13.8-16.3	583	10.9	10.0-11.9	646	6.9	6.4-7.5	314	7.8	6.9-8.8	332	6.2	5.5-6.9
1981	1225	12.9	12.2-13.7	605	15.0	13.7-16.3	620	11.4	10.5-12.4	639	6.8	6.2-7.3	333	8.4	7.5-9.4	306	5.6	4.9-6.2
1982	1250	13.0	12.3-13.8	651	15.8	14.6-17.2	599	11.0	10.1-12.0	653	6.9	6.4-7.5	338	8.7	7.8-9.7	315	5.6	5.0-6.3
1983	1356	13.9	13.2-14.7	671	16.0	14.8-17.3	685	12.3	11.4-13.3	685	7.1	6.6-7.7	359	8.8	7.9-9.9	326	5.9	5.3-6.6
1984	1405	14.6	13.8-15.4	736	18.2	16.9-19.7	669	12.0	11.1-13.0	691	7.1	6.6-7.7	374	9.6	8.6-10.7	317	5.6	5.0-6.2
1985	1477	15.1	14.4-15.9	778	18.4	17.0-19.7	699	12.5	11.6-13.5	736	7.6	7.0-8.2	385	9.5	8.5-10.5	351	6.2	5.5-6.9
1986	1548	15.8	15.0-16.7	833	19.9	18.5-21.4	715	12.8	11.9-13.8	783	8.1	7.5-8.7	428	10.7	9.7-11.8	355	6.1	5.5-6.8
1987	1515	15.3	14.6-16.1	805	18.9	17.6-20.3	710	12.5	11.6-13.5	781	8.0	7.4-8.5	384	9.4	8.4-10.4	397	6.9	6.2-7.6
1988	1634	16.6	15.8-17.4	834	19.6	18.2-21.0	800	14.4	13.4-15.4	792	8.0	7.5-8.6	398	9.8	8.8-10.8	394	6.8	6.1-7.5
1989	1701	17.1	16.3-18.0	892	20.5	19.2-22.0	809	14.4	13.4-15.4	823	8.2	7.7-8.8	439	10.6	9.6-11.7	384	6.5	5.9-7.2
1990	1788	17.7	16.9-18.6	952	21.8	20.4-23.3	836	14.5	13.5-15.5	855	8.5	7.9-9.1	434	10.5	9.5-11.6	421	7.0	6.4-7.8
1991	1769	17.5	16.7-18.3	926	21.6	20.2-23.1	843	14.6	13.6-15.7	866	8.5	7.9-9.1	452	10.9	9.9-12.0	414	6.8	6.1-7.5
1992	1758	17.3	16.5-18.1	920	20.8	19.5-22.2	838	14.4	13.4-15.4	911	8.9	8.3-9.5	459	11.0	10.0-12.1	452	7.4	6.8-8.2
1993	1862	18.2	17.4-19.1	980	21.8	20.5-23.3	882	15.1	14.1-16.2	844	8.1	7.6-8.7	418	9.7	8.8-10.7	426	6.9	6.2-7.6
1994	1947	18.8	18.0-19.7	1046	23.3	21.9-24.8	901	15.2	14.2-16.2	961	9.2	8.6-9.8	492	11.5	10.5-12.6	469	7.6	6.9-8.3
1995	1968	19.0	18.2-19.8	1021	22.8	21.4-24.2	947	16.1	15.1-17.2	980	9.3	8.7-9.9	508	11.9	10.9-13.0	472	7.5	6.9-8.3
1996	2150	20.6	19.7-21.5	1128	24.9	23.4-26.4	1022	17.3	16.2-18.4	990	9.3	8.7-9.9	508	11.7	10.7-12.7	482	7.7	7.0-8.4
1997	2205	20.9	20.1-21.8	1132	24.6	23.2-26.1	1073	18.0	16.9-19.1	1033	9.7	9.1-10.3	518	11.6	10.6-12.7	515	8.2	7.5-8.9
1998	2242	21.1	20.2-22.0	1124	24.2	22.8-25.7	1118	18.7	17.6-19.8	952	8.8	8.2-9.3	458	10.1	9.2-11.1	494	7.7	7.0-8.4
1999	2272	21.2	20.3-22.1	1136	24.3	22.9-25.8	1136	18.8	17.7-20.0	912	8.3	7.8-8.9	443	9.9	9.0-10.9	469	7.3	6.6-8.0
2000	2290	21.1	20.3-22.0	1169	24.8	23.4-26.3	1121	18.3	17.2-19.4	957	8.6	8.1-9.2	469	10.3	9.4-11.3	488	7.5	6.8-8.2
2001	2291	21.0	20.1-21.8	1159	24.4	23.0-25.9	1132	18.3	17.3-19.5	880	7.9	7.4-8.4	474	10.1	9.2-11.0	406	6.1	5.5-6.7
2002	2359	21.5	20.6-22.4	1245	25.9	24.4-27.4	1114	18.1	17.0-19.2	885	7.8	7.3-8.4	478	10.2	9.3-11.2	407	6.1	5.6-6.8
2003	2445	22.0	21.1-22.9	1249	25.6	24.2-27.1	1196	19.1	18.0-20.2	808	7.0	6.5-7.5	406	8.7	7.9-9.6	402	5.9	5.3-6.5
2004	2512	22.5	21.6-23.4	1336	27.1	25.6-28.6	1176	19.0	17.9-20.1	790	6.8	6.3-7.3	383	8.0	7.2-8.8	407	5.9	5.3-6.5
2005	2575	22.9	22.0-23.8	1380	27.7	26.3-29.3	1195	18.9	17.9-20.1	761	6.5	6.0-7.0	403	8.4	7.6-9.2	358	5.1	4.6-5.7
2006	2690	23.7	22.8-24.7	1389	27.7	26.2-29.2	1301	20.4	19.3-21.6	849	7.2	6.7-7.7	430	8.8	7.9-9.6	419	5.9	5.3-6.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Non-Hodgkin lymphomas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	2726	23.7	22.8-24.7	1455	28.6	27.1-30.1	1271	20.0	18.9-21.2	750	6.2	5.8-6.7	416	8.3	7.5-9.1	334	4.7	4.2-5.3
2008	2724	23.6	22.7-24.5	1442	28.0	26.6-29.5	1282	20.1	19.0-21.3	817	6.8	6.3-7.3	423	8.4	7.6-9.2	394	5.7	5.1-6.3
2009	2759	23.5	22.6-24.4	1491	28.4	27.0-29.9	1268	19.5	18.4-20.7	766	6.2	5.8-6.7	427	8.3	7.5-9.1	339	4.8	4.3-5.3
2010	2773	23.3	22.4-24.2	1465	27.7	26.3-29.2	1308	19.7	18.6-20.9	792	6.4	5.9-6.8	414	7.9	7.2-8.8	378	5.2	4.7-5.8
2011	2751	23.1	22.3-24.0	1516	28.4	26.9-29.9	1235	19.0	17.9-20.1	778	6.2	5.8-6.7	411	7.7	7.0-8.5	367	5.1	4.6-5.7
2012	2786	23.0	22.1-23.9	1509	27.9	26.4-29.3	1277	19.0	18.0-20.2	749	6.0	5.5-6.4	413	7.8	7.1-8.6	336	4.5	4.0-5.1
2013	2851	23.3	22.4-24.2	1551	28.2	26.7-29.6	1300	19.4	18.3-20.5	705	5.5	5.1-5.9	383	7.0	6.3-7.8	322	4.3	3.8-4.8
2014	2910	23.4	22.5-24.3	1599	28.6	27.2-30.1	1311	19.1	18.0-20.2	753	5.7	5.3-6.2	420	7.6	6.9-8.4	333	4.3	3.8-4.8
2015	2913	23.3	22.4-24.2	1610	28.3	26.9-29.8	1303	19.2	18.1-20.3	735	5.6	5.2-6.1	410	7.3	6.6-8.1	325	4.3	3.8-4.8
2016	2925	23.3	22.5-24.2	1545	27.0	25.6-28.4	1380	20.4	19.3-21.6	691	5.2	4.8-5.6	368	6.4	5.8-7.2	323	4.2	3.8-4.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Myeloma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	395	4.4	4.0-4.8	199	5.3	4.6-6.2	196	3.8	3.2-4.3	258	3.0	2.6-3.3	136	3.8	3.1-4.5	122	2.4	2.0-2.9
1977	371	4.1	3.7-4.5	192	5.2	4.4-6.0	179	3.4	2.9-3.9	248	2.7	2.4-3.1	122	3.1	2.6-3.8	126	2.4	2.0-2.8
1978	390	4.3	3.9-4.8	179	4.8	4.1-5.6	211	4.0	3.5-4.6	251	2.8	2.4-3.1	128	3.4	2.8-4.1	123	2.3	1.9-2.7
1979	417	4.5	4.1-5.0	187	5.0	4.3-5.8	230	4.2	3.7-4.8	258	2.9	2.5-3.2	131	3.7	3.0-4.4	127	2.4	2.0-2.8
1980	410	4.4	3.9-4.8	212	5.6	4.8-6.4	198	3.6	3.1-4.2	311	3.3	3.0-3.7	150	4.1	3.4-4.8	161	2.9	2.5-3.4
1981	449	4.8	4.3-5.3	211	5.5	4.8-6.4	238	4.4	3.8-5.0	288	3.0	2.7-3.4	132	3.5	2.9-4.2	156	2.8	2.4-3.3
1982	444	4.6	4.2-5.0	236	6.0	5.2-6.8	208	3.7	3.2-4.2	286	3.0	2.7-3.4	142	3.8	3.2-4.5	144	2.5	2.1-3.0
1983	434	4.5	4.1-5.0	237	6.3	5.5-7.2	197	3.5	3.0-4.0	298	3.1	2.8-3.5	155	4.2	3.5-4.9	143	2.5	2.1-3.0
1984	481	5.0	4.5-5.5	256	6.4	5.6-7.3	225	3.9	3.4-4.5	330	3.4	3.0-3.8	164	4.1	3.5-4.9	166	2.9	2.5-3.4
1985	482	5.0	4.5-5.4	242	6.1	5.3-6.9	240	4.2	3.7-4.8	343	3.5	3.1-3.9	175	4.5	3.8-5.3	168	2.8	2.4-3.3
1986	444	4.5	4.1-5.0	244	6.1	5.3-6.9	200	3.5	3.0-4.0	343	3.5	3.1-3.9	169	4.4	3.8-5.2	174	2.9	2.5-3.4
1987	504	5.1	4.7-5.6	249	6.2	5.4-7.0	255	4.3	3.8-4.9	340	3.4	3.1-3.8	169	4.3	3.7-5.0	171	2.9	2.5-3.4
1988	475	4.8	4.4-5.3	251	6.2	5.4-7.0	224	3.8	3.3-4.4	361	3.6	3.3-4.0	199	5.2	4.4-6.0	162	2.7	2.3-3.1
1989	516	5.2	4.7-5.7	241	6.1	5.3-6.9	275	4.7	4.1-5.3	320	3.2	2.8-3.6	164	4.1	3.5-4.8	156	2.6	2.2-3.0
1990	544	5.3	4.9-5.8	282	6.7	6.0-7.6	262	4.4	3.9-5.0	322	3.2	2.8-3.5	163	4.1	3.5-4.8	159	2.6	2.2-3.0
1991	511	5.0	4.5-5.4	268	6.5	5.7-7.3	243	4.0	3.5-4.5	352	3.4	3.1-3.8	172	4.2	3.6-4.9	180	2.9	2.5-3.4
1992	558	5.4	4.9-5.8	282	6.7	5.9-7.6	276	4.5	4.0-5.1	353	3.4	3.0-3.8	185	4.6	3.9-5.3	168	2.6	2.3-3.1
1993	559	5.3	4.9-5.8	284	6.5	5.7-7.3	275	4.5	4.0-5.1	385	3.7	3.3-4.1	189	4.5	3.9-5.2	196	3.1	2.6-3.5
1994	556	5.3	4.9-5.8	290	6.7	6.0-7.6	266	4.3	3.8-4.8	411	3.9	3.5-4.3	183	4.4	3.7-5.1	228	3.6	3.2-4.2
1995	590	5.6	5.1-6.1	312	7.3	6.5-8.2	278	4.5	4.0-5.1	404	3.8	3.4-4.2	211	4.9	4.3-5.6	193	3.0	2.6-3.5
1996	580	5.4	5.0-5.9	300	6.9	6.1-7.7	280	4.5	4.0-5.1	385	3.6	3.2-4.0	198	4.8	4.1-5.5	187	2.9	2.5-3.4
1997	595	5.5	5.1-6.0	288	6.4	5.7-7.2	307	5.0	4.4-5.6	375	3.5	3.1-3.8	183	4.3	3.7-5.0	192	3.0	2.6-3.5
1998	662	6.1	5.6-6.6	337	7.3	6.6-8.2	325	5.2	4.6-5.8	387	3.5	3.2-3.9	192	4.3	3.7-5.0	195	3.0	2.6-3.4
1999	681	6.2	5.7-6.7	362	7.9	7.1-8.8	319	5.0	4.5-5.6	405	3.6	3.3-4.0	213	4.7	4.1-5.4	192	2.9	2.5-3.3
2000	679	6.1	5.7-6.6	364	7.8	7.0-8.6	315	4.9	4.4-5.5	410	3.6	3.3-4.0	218	4.8	4.2-5.5	192	2.9	2.5-3.4
2001	644	5.8	5.3-6.2	376	8.0	7.2-8.9	268	4.1	3.6-4.6	349	3.1	2.8-3.4	191	4.1	3.6-4.8	158	2.3	2.0-2.7
2002	716	6.3	5.9-6.8	403	8.5	7.6-9.3	313	4.8	4.3-5.4	394	3.4	3.1-3.8	218	4.8	4.1-5.4	176	2.6	2.2-3.0
2003	668	5.9	5.4-6.3	374	7.7	7.0-8.6	294	4.5	4.0-5.1	373	3.2	2.9-3.6	215	4.6	4.0-5.2	158	2.3	2.0-2.7
2004	714	6.2	5.8-6.7	397	8.0	7.2-8.9	317	4.8	4.3-5.4	364	3.1	2.8-3.5	200	4.2	3.6-4.8	164	2.4	2.0-2.8
2005	682	5.9	5.4-6.3	374	7.5	6.7-8.3	308	4.6	4.1-5.2	377	3.2	2.9-3.6	198	4.1	3.5-4.7	179	2.6	2.2-3.0
2006	723	6.2	5.7-6.6	413	8.1	7.3-8.9	310	4.6	4.1-5.2	360	3.0	2.7-3.4	199	4.0	3.5-4.6	161	2.4	2.0-2.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Myeloma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	786	6.7	6.2-7.2	428	8.4	7.6-9.2	358	5.4	4.8-6.0	374	3.1	2.8-3.4	193	3.8	3.3-4.4	181	2.6	2.2-3.0
2008	788	6.5	6.1-7.0	427	8.2	7.4-9.0	361	5.3	4.8-5.9	358	2.9	2.6-3.3	200	4.0	3.5-4.6	158	2.2	1.9-2.6
2009	828	6.8	6.3-7.3	444	8.4	7.6-9.2	384	5.6	5.0-6.2	380	3.1	2.7-3.4	203	3.9	3.4-4.5	177	2.5	2.1-2.9
2010	840	6.9	6.4-7.3	459	8.4	7.7-9.2	381	5.6	5.0-6.2	386	3.1	2.8-3.4	189	3.6	3.1-4.2	197	2.7	2.3-3.1
2011	865	7.0	6.5-7.5	493	9.1	8.3-9.9	372	5.3	4.8-5.9	363	2.8	2.5-3.1	197	3.6	3.1-4.2	166	2.2	1.9-2.6
2012	859	6.8	6.3-7.2	477	8.4	7.7-9.2	382	5.5	5.0-6.1	394	3.0	2.7-3.4	198	3.6	3.1-4.2	196	2.6	2.3-3.0
2013	915	7.2	6.7-7.7	529	9.3	8.5-10.2	386	5.5	5.0-6.1	344	2.7	2.4-3.0	181	3.3	2.8-3.8	163	2.2	1.9-2.6
2014	989	7.7	7.2-8.2	529	9.2	8.4-10.0	460	6.5	5.9-7.1	392	2.9	2.7-3.3	220	3.8	3.3-4.4	172	2.3	1.9-2.6
2015	892	6.9	6.5-7.4	500	8.6	7.9-9.4	392	5.6	5.0-6.2	384	2.9	2.6-3.2	221	3.9	3.4-4.5	163	2.1	1.8-2.5
2016	949	7.4	6.9-7.9	537	9.1	8.4-10.0	412	6.0	5.4-6.6	337	2.5	2.2-2.8	155	2.6	2.2-3.1	182	2.4	2.0-2.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Leukemias

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1182	13.2	12.4-14.0	672	18.1	16.7-19.6	510	10.0	9.1-10.9	727	8.2	7.6-8.9	415	11.7	10.5-13.0	312	6.1	5.5-6.9
1977	1114	12.4	11.7-13.2	617	16.3	14.9-17.7	497	9.7	8.9-10.6	785	8.8	8.2-9.5	420	11.2	10.1-12.4	365	7.2	6.5-8.0
1978	1154	12.9	12.2-13.7	636	17.0	15.6-18.4	518	10.2	9.3-11.1	740	8.2	7.6-8.8	386	10.3	9.3-11.5	354	6.8	6.1-7.6
1979	1265	13.7	12.9-14.5	720	18.4	17.0-19.9	545	10.3	9.5-11.3	779	8.6	8.0-9.2	421	11.3	10.1-12.5	358	6.9	6.2-7.6
1980	1266	13.7	12.9-14.5	691	17.9	16.6-19.4	575	11.0	10.1-11.9	820	8.9	8.3-9.5	446	11.7	10.5-12.9	374	7.0	6.3-7.7
1981	1168	12.6	11.9-13.4	657	17.0	15.6-18.4	511	9.6	8.8-10.5	798	8.7	8.1-9.4	444	11.9	10.8-13.1	354	6.7	6.0-7.5
1982	1203	12.9	12.2-13.7	679	17.6	16.2-19.0	524	9.9	9.0-10.8	821	8.9	8.3-9.5	464	12.5	11.3-13.7	357	6.7	6.0-7.4
1983	1183	12.5	11.8-13.2	676	16.9	15.6-18.3	507	9.3	8.5-10.2	800	8.6	8.0-9.2	414	11.1	10.0-12.2	386	7.0	6.3-7.7
1984	1278	13.4	12.6-14.1	684	17.0	15.6-18.3	594	10.8	9.9-11.7	753	7.9	7.4-8.5	364	9.2	8.3-10.3	389	7.1	6.4-7.8
1985	1337	13.9	13.2-14.7	760	18.8	17.4-20.2	577	10.5	9.6-11.4	799	8.3	7.8-9.0	469	12.1	10.9-13.3	330	5.9	5.2-6.5
1986	1243	13.0	12.3-13.7	691	17.3	16.0-18.7	552	10.0	9.2-10.9	789	8.3	7.7-8.9	451	11.5	10.4-12.7	338	6.0	5.4-6.7
1987	1222	12.5	11.8-13.2	669	16.3	15.0-17.6	553	10.0	9.2-10.9	786	8.2	7.6-8.8	434	11.0	9.9-12.1	352	6.2	5.6-6.9
1988	1239	12.6	11.9-13.3	696	16.9	15.6-18.2	543	9.7	8.9-10.6	800	8.1	7.6-8.7	461	11.7	10.6-12.8	339	5.9	5.2-6.5
1989	1219	12.3	11.6-13.0	683	16.6	15.4-18.0	536	9.3	8.6-10.2	797	8.1	7.5-8.6	433	10.8	9.8-11.9	364	6.3	5.7-7.0
1990	1364	13.7	13.0-14.5	748	17.7	16.4-19.0	616	10.8	10.0-11.7	790	7.9	7.4-8.5	405	9.9	8.9-11.0	385	6.6	5.9-7.3
1991	1314	13.1	12.4-13.8	717	17.0	15.8-18.4	597	10.4	9.6-11.3	786	7.8	7.2-8.3	395	9.7	8.7-10.8	391	6.6	5.9-7.2
1992	1346	13.3	12.6-14.0	764	18.1	16.8-19.5	582	10.0	9.2-10.9	842	8.2	7.7-8.8	464	11.3	10.2-12.4	378	6.3	5.7-7.0
1993	1344	13.2	12.5-13.9	746	17.1	15.9-18.4	598	10.4	9.6-11.3	747	7.3	6.8-7.8	405	9.9	8.9-10.9	342	5.7	5.1-6.3
1994	1455	14.1	13.4-14.8	815	18.7	17.4-20.0	640	10.8	10.0-11.7	839	8.1	7.5-8.6	476	11.3	10.3-12.4	363	5.9	5.3-6.6
1995	1433	13.7	13.0-14.5	797	18.0	16.8-19.4	636	10.8	10.0-11.7	835	7.9	7.4-8.5	472	11.3	10.3-12.4	363	5.9	5.3-6.5
1996	1501	14.3	13.6-15.1	873	19.4	18.1-20.7	628	10.5	9.7-11.4	895	8.5	7.9-9.0	496	11.4	10.4-12.5	399	6.5	5.9-7.2
1997	1515	14.4	13.7-15.1	824	18.3	17.0-19.6	691	11.4	10.6-12.3	858	8.0	7.4-8.5	470	10.7	9.7-11.7	388	6.1	5.5-6.7
1998	1519	14.3	13.6-15.1	884	19.2	17.9-20.5	635	10.6	9.7-11.4	828	7.7	7.2-8.2	454	10.1	9.2-11.1	374	5.9	5.3-6.5
1999	1658	15.5	14.7-16.2	926	20.3	19.0-21.6	732	12.0	11.1-12.9	890	8.1	7.6-8.7	482	10.8	9.8-11.8	408	6.3	5.7-7.0
2000	1766	16.3	15.6-17.1	1003	21.8	20.4-23.2	763	12.2	11.4-13.2	893	8.1	7.6-8.7	506	11.2	10.2-12.2	387	6.0	5.4-6.6
2001	1729	15.9	15.2-16.7	963	20.6	19.3-22.0	766	12.5	11.6-13.4	870	7.8	7.3-8.4	472	10.4	9.5-11.4	398	6.1	5.5-6.8
2002	1637	15.0	14.2-15.7	937	19.7	18.5-21.0	700	11.4	10.5-12.3	866	7.7	7.2-8.2	483	10.4	9.5-11.4	383	5.8	5.2-6.4
2003	1780	16.1	15.4-16.9	1010	21.2	19.9-22.5	770	12.4	11.5-13.4	846	7.4	6.9-8.0	486	10.5	9.5-11.4	360	5.4	4.8-6.0
2004	1842	16.6	15.8-17.3	1029	21.2	19.9-22.6	813	13.1	12.2-14.0	906	7.9	7.4-8.4	506	10.6	9.7-11.6	400	6.0	5.4-6.6
2005	1868	16.7	16.0-17.5	1033	21.0	19.8-22.4	835	13.2	12.3-14.2	873	7.5	7.0-8.0	500	10.5	9.6-11.5	373	5.5	4.9-6.1
2006	1853	16.5	15.7-17.3	1073	21.8	20.5-23.2	780	12.4	11.5-13.3	886	7.7	7.2-8.2	494	10.3	9.4-11.2	392	5.7	5.1-6.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Whites
Leukemias

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	1914	16.9	16.1-17.7	1111	22.1	20.8-23.5	803	12.9	12.0-13.9	853	7.3	6.8-7.8	487	9.9	9.0-10.8	366	5.4	4.8-6.0
2008	1857	16.2	15.5-17.0	1065	20.8	19.5-22.1	792	12.7	11.8-13.7	872	7.3	6.8-7.8	512	10.2	9.3-11.1	360	5.2	4.6-5.8
2009	1985	17.1	16.3-17.9	1169	22.6	21.3-23.9	816	12.9	12.0-13.8	905	7.7	7.2-8.2	546	10.9	10.0-11.8	359	5.3	4.7-5.9
2010	2100	17.9	17.2-18.7	1226	23.4	22.1-24.8	874	13.6	12.6-14.5	827	6.8	6.3-7.3	473	9.3	8.4-10.1	354	5.0	4.5-5.6
2011	2214	18.8	18.0-19.6	1275	24.3	22.9-25.7	939	14.5	13.5-15.5	883	7.2	6.7-7.7	504	9.5	8.7-10.4	379	5.4	4.9-6.1
2012	2138	17.9	17.2-18.8	1246	23.2	21.9-24.6	892	13.8	12.9-14.8	885	7.1	6.6-7.6	519	9.9	9.1-10.8	366	5.1	4.5-5.6
2013	2294	19.1	18.3-20.0	1350	24.9	23.6-26.3	944	14.5	13.6-15.5	863	6.8	6.4-7.3	495	9.2	8.4-10.1	368	5.1	4.5-5.6
2014	2261	18.5	17.7-19.3	1360	24.6	23.3-26.0	901	13.5	12.6-14.5	856	6.6	6.2-7.1	483	8.9	8.1-9.7	373	5.1	4.5-5.7
2015	2249	18.4	17.6-19.2	1309	23.7	22.3-25.0	940	14.2	13.3-15.2	857	6.6	6.1-7.0	498	9.0	8.2-9.8	359	4.8	4.3-5.3
2016	2189	17.5	16.7-18.2	1301	22.9	21.6-24.2	888	12.9	12.0-13.9	736	5.7	5.2-6.1	424	7.7	7.0-8.5	312	4.2	3.7-4.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
All Invasive Malignant Tumors

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	5617	386.1	375.0-397.4	2779	491.4	470.3-513.0	2838	323.3	310.7-336.4	3435	246.8	237.7-256.1	1833	330.6	313.0-348.8	1602	194.4	184.3-205.0
1977	5956	400.1	389.0-411.4	2945	519.4	497.8-541.5	3011	330.8	318.3-343.7	3498	241.5	232.7-250.4	1911	343.2	325.5-361.4	1587	179.9	170.5-189.6
1978	6285	415.9	404.7-427.3	3131	537.6	516.1-559.7	3154	344.9	332.1-357.9	3769	254.2	245.4-263.2	1979	338.2	321.3-355.7	1790	202.7	192.7-212.9
1979	6515	419.2	408.2-430.4	3256	548.5	527.1-570.3	3259	344.2	331.8-357.0	3904	261.1	252.2-270.2	2110	366.3	348.6-384.7	1794	198.3	188.6-208.3
1980	7010	433.3	422.4-444.3	3562	582.9	561.3-605.0	3448	347.8	335.6-360.2	4013	257.3	248.7-266.0	2142	362.0	344.7-379.9	1871	196.1	186.8-205.7
1981	6853	411.5	401.1-422.1	3500	554.7	534.0-575.9	3353	330.1	318.5-342.0	4167	258.6	250.2-267.2	2219	364.2	347.1-381.9	1948	197.6	188.4-207.0
1982	7078	412.0	401.8-422.3	3558	548.9	528.9-569.4	3520	334.5	323.1-346.2	4156	249.0	241.0-257.3	2241	357.8	341.3-374.8	1915	186.7	178.0-195.6
1983	7211	402.3	392.5-412.3	3593	534.5	515.0-554.5	3618	329.8	318.7-341.1	4378	252.0	244.2-260.1	2358	358.8	342.8-375.3	2020	190.0	181.5-198.8
1984	7764	421.8	411.9-431.8	3858	553.9	534.4-573.8	3906	348.7	337.4-360.2	4330	245.5	237.8-253.4	2282	344.9	329.2-361.1	2048	189.4	181.0-198.1
1985	8005	424.2	414.5-434.2	3988	560.1	540.8-579.8	4017	349.1	338.0-360.4	4343	240.5	233.0-248.2	2316	343.7	328.2-359.6	2027	182.8	174.6-191.2
1986	8205	424.0	414.3-433.7	4179	579.5	560.0-599.5	4026	339.7	329.0-350.7	4479	241.2	233.8-248.7	2348	342.9	327.6-358.8	2131	185.6	177.6-193.9
1987	8237	413.3	403.9-422.7	4110	550.4	531.8-569.4	4127	338.6	328.0-349.4	4430	233.5	226.3-240.8	2334	338.3	323.1-354.0	2096	177.6	169.9-185.6
1988	8388	414.9	405.6-424.3	4098	548.4	529.8-567.4	4290	345.3	334.8-356.1	4481	232.2	225.1-239.4	2331	332.0	317.2-347.3	2150	178.6	170.9-186.6
1989	8682	418.2	409.0-427.5	4393	571.4	552.7-590.4	4289	336.7	326.5-347.2	4477	225.9	219.1-233.0	2363	328.3	313.7-343.2	2114	171.1	163.7-178.8
1990	9016	428.5	419.3-437.8	4480	577.5	558.8-596.6	4536	350.2	339.8-360.8	4592	229.8	222.8-236.8	2340	321.8	307.5-336.7	2252	181.1	173.5-188.9
1991	9613	447.5	438.2-457.0	4796	603.9	585.0-623.2	4817	364.9	354.5-375.6	4555	223.0	216.2-229.8	2360	320.7	306.5-335.4	2195	171.9	164.6-179.4
1992	9980	453.5	444.3-462.9	5095	625.7	606.8-645.0	4885	361.7	351.4-372.2	4540	218.4	211.8-225.1	2221	300.4	286.6-314.6	2319	177.8	170.5-185.3
1993	10002	444.7	435.7-453.9	5226	623.4	604.9-642.3	4776	346.8	336.8-356.9	4519	212.5	206.1-219.0	2291	302.0	288.3-316.0	2228	167.6	160.6-174.8
1994	10613	463.0	454.0-472.2	5433	633.0	614.8-651.7	5180	369.4	359.2-379.8	4974	230.1	223.5-236.8	2521	324.3	310.5-338.5	2453	181.5	174.3-188.9
1995	10635	453.1	444.2-462.0	5581	628.3	610.5-646.5	5054	353.2	343.4-363.3	4973	225.1	218.7-231.6	2447	305.8	292.6-319.3	2526	183.3	176.1-190.6
1996	11298	470.2	461.4-479.3	5803	636.6	618.9-654.6	5495	375.7	365.7-385.9	4856	215.5	209.3-221.9	2434	297.5	284.7-310.6	2422	172.4	165.5-179.5
1997	11383	469.4	460.6-478.3	5836	638.2	620.6-656.2	5547	374.3	364.4-384.4	4835	211.3	205.2-217.5	2410	291.7	279.2-304.6	2425	169.4	162.7-176.4
1998	11689	468.1	459.4-476.9	5839	613.7	596.9-630.9	5850	385.2	375.3-395.3	4840	206.3	200.4-212.4	2342	272.4	260.5-284.6	2498	171.4	164.7-178.3
1999	11836	463.8	455.3-472.4	6099	627.0	610.2-644.0	5737	369.2	359.7-379.0	4884	203.1	197.3-209.0	2335	268.6	256.9-280.7	2549	170.1	163.5-176.9
2000	12156	463.5	455.1-472.0	6355	635.7	619.1-652.5	5801	363.5	354.1-373.1	4879	197.1	191.5-202.9	2379	265.5	254.1-277.2	2500	161.8	155.5-168.3
2001	12232	455.3	447.1-463.6	6417	623.4	607.2-639.9	5815	357.1	347.9-366.5	4923	195.2	189.6-200.8	2311	255.0	243.9-266.4	2612	165.5	159.1-172.0
2002	12454	453.2	445.2-461.4	6423	610.8	594.9-626.9	6031	362.6	353.5-372.0	4853	187.3	181.9-192.7	2368	250.9	240.1-262.0	2485	154.3	148.2-160.5
2003	12748	454.5	446.5-462.7	6573	609.9	594.3-625.8	6175	364.3	355.2-373.6	4727	178.1	172.9-183.3	2265	231.8	221.6-242.2	2462	149.5	143.6-155.6
2004	13105	457.8	449.9-465.9	6703	602.5	587.3-618.1	6402	372.1	363.0-381.4	4784	178.2	173.1-183.5	2319	234.6	224.4-245.0	2465	148.1	142.3-154.1
2005	13304	457.5	449.6-465.5	6811	601.1	586.0-616.4	6493	372.5	363.4-381.7	4781	173.8	168.9-179.0	2237	220.0	210.3-229.9	2544	149.8	144.0-155.7
2006	13626	461.4	453.5-469.4	7035	610.3	595.2-625.5	6591	372.5	363.5-381.7	4606	165.4	160.6-170.4	2152	210.2	200.8-219.8	2454	142.6	136.9-148.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
All Invasive Malignant Tumors

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	14047	466.6	458.8-474.6	7426	624.7	609.8-639.9	6621	369.7	360.8-378.8	4817	169.3	164.5-174.3	2259	213.9	204.6-223.5	2558	146.3	140.6-152.1
2008	14329	465.6	457.8-473.4	7356	601.8	587.3-616.5	6973	382.3	373.3-391.5	4698	161.6	156.9-166.4	2185	203.0	194.1-212.3	2513	140.3	134.8-145.9
2009	14641	465.1	457.5-472.9	7559	597.7	583.5-612.0	7082	381.5	372.6-390.6	4763	160.7	156.0-165.4	2315	210.4	201.3-219.6	2448	134.1	128.8-139.6
2010	14716	455.8	448.3-463.4	7569	583.6	569.8-597.7	7147	376.7	367.9-385.6	4848	159.7	155.1-164.4	2345	209.4	200.5-218.6	2503	133.9	128.6-139.3
2011	15813	474.5	467.0-482.1	8118	602.4	588.7-616.4	7695	394.3	385.4-403.4	5111	163.3	158.7-167.9	2380	200.6	192.2-209.3	2731	143.0	137.6-148.5
2012	15233	446.3	439.1-453.6	7566	546.3	533.4-559.4	7667	385.0	376.3-393.8	5068	156.7	152.4-161.2	2376	193.5	185.4-201.9	2692	137.0	131.8-142.3
2013	15646	446.0	438.9-453.2	7737	540.1	527.6-552.9	7909	387.9	379.3-396.7	4930	148.5	144.3-152.8	2349	187.0	179.1-195.1	2581	127.0	122.1-132.1
2014	15718	436.9	429.9-443.9	7704	519.4	507.3-531.6	8014	385.5	377.0-394.2	5119	149.6	145.4-153.8	2390	182.9	175.3-190.8	2729	131.5	126.5-136.6
2015	15863	429.6	422.8-436.5	7745	508.9	497.1-520.9	8118	381.2	372.8-389.7	4913	139.6	135.6-143.6	2261	167.2	160.0-174.5	2652	124.4	119.6-129.3
2016	16069	426.6	419.9-433.4	7895	507.2	495.6-519.0	8174	376.9	368.6-385.3	5058	139.0	135.1-143.0	2398	169.2	162.2-176.5	2660	121.6	117.0-126.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Oral cavity and pharynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	249	14.8	12.9-16.9	170	24.9	20.8-29.5	79	8.0	6.3-10.1	111	7.2	5.8-8.7	75	12.6	9.5-16.2	36	3.7	2.6-5.2
1977	237	13.9	12.1-15.8	181	26.1	22.1-30.5	56	5.5	4.1-7.2	127	7.6	6.3-9.2	100	13.9	11.2-17.1	27	3.0	1.9-4.5
1978	268	14.7	13.0-16.7	185	24.7	21.0-28.9	83	7.8	6.2-9.8	133	7.7	6.4-9.3	100	15.1	11.9-18.8	33	3.1	2.1-4.3
1979	271	15.5	13.6-17.6	203	28.3	24.3-32.7	68	6.7	5.1-8.6	148	8.4	7.0-9.9	109	15.4	12.4-18.9	39	3.7	2.6-5.1
1980	277	15.0	13.2-16.9	196	26.3	22.5-30.5	81	7.3	5.8-9.1	140	7.8	6.5-9.3	114	16.0	12.9-19.6	26	2.3	1.5-3.4
1981	303	15.8	14.0-17.8	219	27.1	23.4-31.2	84	7.7	6.1-9.6	120	6.4	5.3-7.7	85	11.3	8.9-14.2	35	3.2	2.2-4.5
1982	292	15.0	13.3-16.9	200	25.4	21.8-29.4	92	8.0	6.4-9.9	152	8.0	6.7-9.4	116	15.3	12.4-18.6	36	3.1	2.1-4.3
1983	285	14.7	13.0-16.6	208	26.3	22.6-30.5	77	6.8	5.4-8.6	145	7.7	6.4-9.1	110	14.7	11.8-18.0	35	3.2	2.2-4.5
1984	334	16.1	14.4-18.0	226	26.5	23.0-30.3	108	8.9	7.3-10.8	121	6.1	5.0-7.3	92	11.8	9.3-14.7	29	2.4	1.6-3.5
1985	325	15.9	14.1-17.7	222	26.6	23.0-30.6	103	8.6	7.0-10.5	116	5.8	4.8-7.0	82	10.2	8.0-12.9	34	2.9	2.0-4.0
1986	321	15.0	13.3-16.8	229	26.2	22.7-30.0	92	7.3	5.8-9.0	135	6.9	5.7-8.2	97	12.5	9.9-15.4	38	3.3	2.3-4.6
1987	350	16.1	14.4-17.9	240	26.8	23.4-30.6	110	8.6	7.1-10.4	105	5.0	4.1-6.1	77	9.3	7.2-11.8	28	2.2	1.4-3.2
1988	306	13.7	12.1-15.3	203	22.6	19.4-26.2	103	7.7	6.2-9.3	133	6.0	5.0-7.2	96	10.7	8.6-13.3	37	2.9	2.0-3.9
1989	332	14.3	12.7-15.9	226	24.1	20.9-27.6	106	7.7	6.3-9.3	134	5.9	4.9-7.0	94	10.3	8.2-12.8	40	3.0	2.1-4.1
1990	305	13.5	12.0-15.2	214	23.9	20.6-27.6	91	6.8	5.5-8.4	101	4.6	3.7-5.7	70	8.5	6.4-11.1	31	2.3	1.5-3.3
1991	397	17.3	15.6-19.1	282	31.4	27.6-35.6	115	8.3	6.8-9.9	142	6.4	5.3-7.5	101	11.0	8.8-13.4	41	3.2	2.3-4.4
1992	368	15.5	14.0-17.3	241	25.7	22.3-29.4	127	9.0	7.5-10.7	118	5.0	4.1-6.0	84	9.0	7.0-11.2	34	2.4	1.7-3.4
1993	349	14.5	12.9-16.1	229	23.5	20.4-26.9	120	8.4	6.9-10.0	103	4.4	3.6-5.3	73	7.9	6.1-10.0	30	2.1	1.4-3.1
1994	331	13.5	12.0-15.0	228	23.1	20.0-26.4	103	7.1	5.7-8.6	119	5.0	4.1-5.9	89	9.6	7.6-12.0	30	2.1	1.4-2.9
1995	303	12.1	10.7-13.5	224	21.6	18.7-24.8	79	5.4	4.3-6.8	139	5.7	4.8-6.8	101	10.2	8.2-12.5	38	2.7	1.9-3.7
1996	333	13.1	11.7-14.6	219	21.8	18.8-25.0	114	7.5	6.2-9.0	108	4.4	3.6-5.4	76	7.8	6.1-9.9	32	2.2	1.5-3.2
1997	328	12.6	11.2-14.0	220	20.3	17.6-23.3	108	7.1	5.9-8.6	89	3.7	2.9-4.5	66	7.1	5.3-9.1	23	1.6	1.0-2.4
1998	295	11.0	9.8-12.4	195	17.6	15.1-20.4	100	6.5	5.2-7.9	97	3.8	3.1-4.6	70	6.9	5.3-8.8	27	1.8	1.2-2.6
1999	268	9.9	8.7-11.2	180	16.1	13.7-18.7	88	5.6	4.5-6.9	91	3.5	2.8-4.3	57	5.7	4.2-7.4	34	2.2	1.5-3.0
2000	278	10.2	9.0-11.5	184	16.9	14.4-19.6	94	5.8	4.7-7.1	101	3.9	3.1-4.7	71	6.8	5.3-8.7	30	1.9	1.3-2.8
2001	251	8.7	7.7-9.9	183	15.6	13.3-18.1	68	4.0	3.1-5.1	86	3.2	2.5-3.9	60	5.5	4.1-7.2	26	1.6	1.1-2.4
2002	292	10.3	9.1-11.5	188	16.2	13.8-18.8	104	6.3	5.1-7.6	76	2.7	2.1-3.4	54	4.6	3.4-6.1	22	1.4	0.8-2.1
2003	268	9.0	8.0-10.2	180	14.4	12.3-16.7	88	5.2	4.2-6.4	79	2.8	2.2-3.5	60	5.3	4.0-6.8	19	1.2	0.7-1.8
2004	296	9.7	8.6-10.9	193	15.5	13.3-18.0	103	5.8	4.7-7.0	102	3.5	2.8-4.2	78	6.6	5.1-8.3	24	1.4	0.9-2.1
2005	267	8.8	7.7-9.9	185	15.2	13.0-17.7	82	4.6	3.7-5.7	83	2.8	2.2-3.5	60	5.3	4.0-6.9	23	1.3	0.8-1.9
2006	260	8.3	7.3-9.4	173	13.5	11.5-15.8	87	4.8	3.8-5.9	64	2.1	1.6-2.7	51	4.1	3.0-5.5	13	0.7	0.4-1.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Oral cavity and pharynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	306	9.8	8.7-11.0	202	15.6	13.4-18.0	104	5.8	4.8-7.1	82	2.7	2.1-3.3	61	5.1	3.8-6.7	21	1.2	0.7-1.8
2008	275	8.6	7.6-9.7	179	13.2	11.3-15.4	96	5.2	4.2-6.3	76	2.5	1.9-3.1	54	4.4	3.2-5.8	22	1.2	0.8-1.9
2009	279	8.5	7.5-9.6	179	13.0	11.1-15.1	100	5.3	4.3-6.4	71	2.2	1.7-2.8	49	3.7	2.7-5.0	22	1.2	0.7-1.8
2010	295	8.7	7.7-9.8	193	13.7	11.8-15.9	102	5.2	4.2-6.3	67	2.1	1.6-2.6	38	2.9	2.0-4.0	29	1.5	1.0-2.2
2011	283	8.1	7.2-9.1	201	13.5	11.7-15.6	82	4.2	3.3-5.2	99	2.9	2.4-3.6	68	4.8	3.7-6.1	31	1.6	1.1-2.3
2012	287	8.1	7.1-9.1	174	11.7	10.0-13.7	113	5.5	4.5-6.6	74	2.2	1.7-2.8	50	3.6	2.6-4.8	24	1.2	0.8-1.9
2013	275	7.7	6.8-8.6	184	12.1	10.3-14.0	91	4.5	3.6-5.5	88	2.6	2.1-3.2	61	4.4	3.3-5.7	27	1.3	0.9-2.0
2014	280	7.5	6.6-8.4	177	11.0	9.4-12.9	103	4.9	4.0-6.0	65	1.8	1.4-2.3	40	2.7	1.9-3.8	25	1.2	0.8-1.8
2015	272	7.3	6.4-8.2	176	11.3	9.7-13.2	96	4.5	3.6-5.5	70	1.9	1.5-2.4	47	3.0	2.2-4.0	23	1.1	0.7-1.7
2016	307	8.1	7.2-9.1	199	12.5	10.8-14.5	108	5.0	4.1-6.1	61	1.6	1.2-2.1	39	2.6	1.8-3.6	22	0.9	0.6-1.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Esophagus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	202	12.5	10.7-14.5	143	21.6	17.9-25.8	59	6.2	4.7-8.2	204	13.0	11.2-15.1	155	24.7	20.5-29.4	49	5.3	3.8-7.2
1977	220	13.4	11.6-15.5	155	23.9	19.8-28.5	65	6.6	5.1-8.5	196	11.9	10.2-13.8	145	22.6	18.6-27.2	51	5.0	3.7-6.7
1978	247	15.1	13.1-17.2	178	27.1	22.9-31.9	69	7.1	5.5-9.1	179	10.5	9.0-12.3	124	18.0	14.7-21.7	55	5.5	4.1-7.3
1979	237	13.9	12.1-15.9	168	24.9	20.9-29.4	69	6.7	5.2-8.6	215	12.8	11.0-14.7	161	24.9	20.7-29.6	54	5.2	3.9-6.8
1980	258	14.8	12.9-16.8	194	28.3	24.1-33.0	64	6.1	4.6-7.8	213	12.4	10.7-14.4	148	22.2	18.4-26.6	65	6.2	4.8-8.0
1981	262	14.0	12.3-15.9	186	24.7	21.1-28.8	76	6.9	5.4-8.6	212	11.3	9.7-12.9	157	20.6	17.3-24.3	55	4.9	3.7-6.5
1982	239	12.9	11.2-14.7	176	24.5	20.7-28.7	63	5.5	4.2-7.0	213	11.8	10.2-13.5	155	22.3	18.6-26.4	58	5.2	3.9-6.7
1983	247	12.7	11.1-14.4	175	22.4	18.9-26.2	72	6.3	4.9-8.0	211	11.1	9.6-12.7	151	19.8	16.6-23.5	60	5.4	4.1-7.0
1984	263	13.3	11.7-15.1	179	22.9	19.4-26.9	84	7.1	5.7-8.9	207	10.8	9.3-12.5	145	19.6	16.3-23.4	62	5.4	4.1-6.9
1985	240	12.1	10.6-13.8	168	21.5	18.1-25.3	72	6.1	4.7-7.7	217	10.9	9.4-12.5	163	20.5	17.2-24.1	54	4.6	3.4-6.0
1986	254	12.7	11.1-14.4	190	24.7	21.0-28.7	64	5.3	4.0-6.7	218	11.0	9.5-12.6	152	19.9	16.6-23.7	66	5.5	4.3-7.1
1987	225	11.0	9.6-12.6	146	18.5	15.4-21.9	79	6.3	5.0-7.9	179	8.8	7.5-10.2	131	16.4	13.5-19.6	48	4.0	2.9-5.3
1988	253	12.2	10.7-13.8	180	22.6	19.1-26.4	73	5.8	4.5-7.3	202	10.0	8.6-11.5	149	19.4	16.2-23.1	53	4.2	3.2-5.6
1989	257	12.1	10.6-13.7	180	21.9	18.6-25.6	77	6.0	4.7-7.5	184	8.7	7.4-10.1	134	16.0	13.3-19.2	50	4.0	3.0-5.3
1990	230	10.7	9.3-12.3	158	19.0	15.9-22.4	72	5.7	4.4-7.1	191	9.0	7.7-10.4	123	15.3	12.5-18.4	68	5.2	4.0-6.6
1991	211	9.6	8.3-11.0	144	17.3	14.4-20.6	67	4.9	3.8-6.2	177	8.0	6.8-9.3	120	14.1	11.5-17.1	57	4.2	3.2-5.5
1992	221	10.1	8.7-11.5	152	17.8	14.8-21.1	69	5.2	4.1-6.6	148	6.6	5.6-7.8	100	11.6	9.3-14.3	48	3.6	2.6-4.8
1993	236	10.2	8.9-11.6	159	17.1	14.4-20.2	77	5.6	4.4-7.0	182	8.0	6.9-9.3	134	15.6	12.9-18.8	48	3.5	2.6-4.7
1994	232	10.1	8.8-11.5	152	17.4	14.5-20.6	80	5.8	4.6-7.2	177	7.8	6.7-9.0	122	14.1	11.5-17.0	55	4.0	3.0-5.2
1995	185	7.9	6.8-9.2	122	13.1	10.7-15.8	63	4.6	3.5-5.9	172	7.5	6.4-8.8	119	13.2	10.8-16.0	53	3.9	2.9-5.1
1996	195	8.4	7.3-9.7	122	13.1	10.8-15.8	73	5.3	4.2-6.7	166	7.1	6.1-8.3	109	11.9	9.7-14.5	57	4.1	3.1-5.3
1997	189	7.9	6.8-9.1	132	13.9	11.5-16.7	57	4.0	3.0-5.2	150	6.4	5.4-7.6	99	10.9	8.8-13.5	51	3.6	2.7-4.8
1998	182	7.3	6.3-8.5	126	12.8	10.6-15.4	56	3.8	2.9-4.9	147	6.0	5.1-7.1	95	9.8	7.8-12.2	52	3.6	2.7-4.8
1999	194	7.5	6.5-8.7	135	13.6	11.3-16.2	59	3.7	2.8-4.8	131	5.1	4.3-6.1	91	9.1	7.2-11.3	40	2.6	1.9-3.6
2000	187	7.5	6.5-8.7	102	10.6	8.5-12.9	85	5.6	4.5-7.0	166	6.6	5.6-7.6	103	10.6	8.6-13.0	63	4.1	3.1-5.2
2001	173	6.4	5.5-7.5	118	11.4	9.3-13.7	55	3.4	2.5-4.4	127	4.8	4.0-5.8	92	9.2	7.3-11.4	35	2.2	1.5-3.1
2002	157	5.9	5.0-6.9	98	9.4	7.5-11.6	59	3.6	2.8-4.7	132	4.9	4.1-5.9	86	8.3	6.6-10.4	46	2.8	2.1-3.8
2003	185	6.8	5.8-7.9	124	11.8	9.7-14.2	61	3.7	2.9-4.8	128	4.6	3.9-5.5	80	7.2	5.6-9.1	48	2.9	2.2-3.9
2004	173	6.2	5.3-7.2	108	9.4	7.7-11.5	65	4.0	3.1-5.1	139	5.1	4.2-6.0	86	8.0	6.3-10.0	53	3.2	2.4-4.2
2005	155	5.3	4.4-6.2	92	7.7	6.1-9.5	63	3.6	2.8-4.6	108	3.9	3.2-4.7	67	5.9	4.5-7.6	41	2.5	1.8-3.3
2006	130	4.7	3.9-5.6	91	8.9	7.0-11.0	39	2.3	1.6-3.2	114	4.1	3.3-4.9	71	7.0	5.4-8.9	43	2.5	1.8-3.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Esophagus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	139	4.6	3.8-5.4	100	8.2	6.6-10.0	39	2.2	1.6-3.0	98	3.3	2.7-4.0	67	5.6	4.3-7.2	31	1.8	1.2-2.5
2008	148	4.9	4.1-5.7	102	8.9	7.1-10.9	46	2.5	1.8-3.3	120	4.0	3.3-4.8	74	6.4	4.9-8.1	46	2.6	1.9-3.4
2009	147	4.7	4.0-5.6	102	8.4	6.7-10.3	45	2.5	1.8-3.3	102	3.3	2.7-4.0	81	6.8	5.3-8.6	21	1.2	0.7-1.8
2010	127	4.0	3.3-4.8	81	6.5	5.1-8.1	46	2.5	1.8-3.3	110	3.5	2.9-4.3	74	6.3	4.8-7.9	36	1.9	1.3-2.7
2011	155	4.8	4.0-5.6	104	8.1	6.5-9.9	51	2.6	2.0-3.5	92	2.8	2.3-3.5	64	4.9	3.7-6.3	28	1.5	1.0-2.2
2012	127	3.8	3.1-4.5	83	6.3	4.9-7.8	44	2.1	1.5-2.9	88	2.5	2.0-3.2	59	4.2	3.1-5.5	29	1.4	0.9-2.0
2013	128	3.8	3.2-4.6	89	7.0	5.5-8.6	39	1.9	1.3-2.6	88	2.7	2.1-3.3	55	4.1	3.0-5.4	33	1.7	1.2-2.4
2014	128	3.7	3.1-4.4	91	6.8	5.4-8.4	37	1.8	1.2-2.5	99	2.9	2.4-3.6	64	5.0	3.8-6.4	35	1.7	1.2-2.4
2015	125	3.4	2.8-4.1	80	5.7	4.4-7.1	45	2.0	1.5-2.7	72	2.1	1.6-2.6	47	3.5	2.5-4.7	25	1.1	0.7-1.7
2016	143	3.9	3.3-4.6	91	6.4	5.1-7.9	52	2.4	1.8-3.2	92	2.5	2.0-3.1	64	4.6	3.5-5.9	28	1.3	0.8-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Stomach

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	243	18.6	16.1-21.4	138	26.3	21.4-31.9	105	13.8	11.1-16.9	189	14.3	12.1-16.6	117	23.0	18.4-28.3	72	8.9	6.9-11.4
1977	270	19.8	17.3-22.5	149	27.0	22.3-32.4	121	15.0	12.3-18.1	212	16.2	13.8-18.7	124	23.8	19.2-29.1	88	11.4	8.9-14.2
1978	244	17.7	15.3-20.2	138	23.8	19.5-28.6	106	13.5	10.9-16.5	209	15.4	13.2-17.9	106	19.3	15.3-23.9	103	13.0	10.4-15.9
1979	253	18.4	16.0-21.0	143	26.0	21.3-31.3	110	13.7	11.1-16.7	180	12.9	10.9-15.0	99	17.5	13.8-21.8	81	10.0	7.8-12.5
1980	316	21.9	19.3-24.6	173	30.0	25.1-35.4	143	16.8	14.0-19.9	208	15.1	13.0-17.5	114	21.6	17.4-26.5	94	11.3	9.0-13.9
1981	282	18.7	16.4-21.2	173	28.8	24.1-34.2	109	12.5	10.1-15.2	240	16.2	14.0-18.5	149	25.2	20.8-30.2	91	10.7	8.5-13.2
1982	281	18.2	16.0-20.6	163	28.5	23.8-33.7	118	12.4	10.2-14.9	208	14.5	12.5-16.8	130	25.6	20.9-31.0	78	8.5	6.7-10.7
1983	283	17.0	15.0-19.2	145	22.5	18.6-26.8	138	13.7	11.4-16.3	214	13.0	11.2-15.0	119	18.8	15.3-22.9	95	9.5	7.6-11.7
1984	264	15.5	13.6-17.6	154	24.6	20.5-29.3	110	10.4	8.4-12.6	179	10.9	9.2-12.7	102	18.1	14.4-22.4	77	7.1	5.6-9.0
1985	287	16.9	14.9-19.1	153	22.3	18.5-26.5	134	13.4	11.1-15.9	184	10.6	9.0-12.3	97	14.5	11.5-18.1	87	8.2	6.5-10.1
1986	307	17.2	15.2-19.3	183	27.9	23.5-32.8	124	11.2	9.3-13.5	209	12.2	10.5-14.1	124	19.1	15.5-23.3	85	8.2	6.5-10.2
1987	319	17.9	15.9-20.1	185	25.6	21.7-30.0	134	12.9	10.8-15.3	200	11.5	9.9-13.3	107	15.6	12.4-19.2	93	9.0	7.2-11.0
1988	305	16.8	14.9-18.9	175	25.5	21.5-29.9	130	11.8	9.8-14.1	215	12.2	10.5-14.0	121	18.2	14.8-22.0	94	8.8	7.0-10.8
1989	309	16.8	14.9-18.9	162	21.7	18.1-25.7	147	13.5	11.3-15.9	217	11.7	10.2-13.5	121	17.0	13.8-20.6	96	8.5	6.9-10.5
1990	323	16.8	14.9-18.8	181	25.3	21.3-29.7	142	12.0	10.1-14.2	211	11.5	10.0-13.3	113	15.8	12.7-19.3	98	8.9	7.2-10.9
1991	295	14.8	13.1-16.6	156	20.4	17.0-24.2	139	11.5	9.6-13.6	205	10.4	8.9-12.0	122	16.2	13.1-19.7	83	7.0	5.5-8.7
1992	324	15.9	14.2-17.9	163	20.3	17.0-24.1	161	13.1	11.1-15.3	229	11.7	10.2-13.4	102	13.2	10.5-16.3	127	10.6	8.8-12.6
1993	333	16.3	14.5-18.2	176	22.5	19.0-26.4	157	12.5	10.6-14.6	193	9.5	8.2-11.0	98	12.9	10.2-15.9	95	7.6	6.1-9.3
1994	364	17.5	15.7-19.4	196	24.7	21.0-28.7	168	13.2	11.2-15.3	245	11.9	10.5-13.6	141	18.5	15.3-22.1	104	8.3	6.8-10.1
1995	325	15.3	13.6-17.1	164	20.2	17.0-23.9	161	12.3	10.4-14.3	213	10.2	8.8-11.7	102	12.6	10.1-15.5	111	8.6	7.0-10.3
1996	339	15.4	13.8-17.2	193	23.5	20.1-27.3	146	10.7	9.0-12.6	216	10.2	8.9-11.7	122	15.8	12.8-19.1	94	7.1	5.7-8.7
1997	332	15.1	13.4-16.8	185	23.0	19.5-26.9	147	10.7	9.0-12.6	195	8.6	7.4-9.9	113	13.3	10.8-16.2	82	5.8	4.6-7.2
1998	342	14.8	13.2-16.5	183	20.6	17.5-24.1	159	11.2	9.5-13.1	205	9.1	7.9-10.5	113	13.4	10.8-16.2	92	6.6	5.3-8.1
1999	317	13.5	12.0-15.1	171	19.7	16.7-23.2	146	10.0	8.4-11.8	201	8.7	7.5-10.1	107	13.0	10.4-15.9	94	6.5	5.2-7.9
2000	337	14.1	12.6-15.7	172	19.4	16.4-22.7	165	11.0	9.4-12.9	195	8.2	7.0-9.4	108	12.5	10.1-15.3	87	5.8	4.6-7.1
2001	326	13.5	12.1-15.1	159	17.5	14.7-20.7	167	11.1	9.5-13.0	167	7.0	6.0-8.2	93	10.1	8.0-12.5	74	5.1	4.0-6.3
2002	323	12.6	11.2-14.1	162	16.9	14.2-20.0	161	10.1	8.6-11.8	189	7.6	6.5-8.8	101	11.6	9.3-14.3	88	5.6	4.5-6.9
2003	329	12.6	11.2-14.0	170	16.8	14.2-19.7	159	9.9	8.4-11.6	174	6.7	5.7-7.7	93	8.8	7.0-11.0	81	5.1	4.1-6.4
2004	314	12.0	10.7-13.5	153	16.2	13.5-19.1	161	9.8	8.3-11.4	174	6.7	5.8-7.8	84	8.9	7.0-11.1	90	5.5	4.4-6.8
2005	407	15.3	13.9-16.9	210	21.3	18.3-24.5	197	11.9	10.3-13.7	212	8.2	7.1-9.4	122	13.2	10.8-15.9	90	5.5	4.4-6.8
2006	369	13.5	12.1-15.0	203	19.6	16.9-22.7	166	9.8	8.4-11.4	188	7.2	6.1-8.3	107	11.1	8.9-13.5	81	4.9	3.9-6.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Stomach

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	335	11.9	10.7-13.3	172	16.5	14.0-19.4	163	9.4	8.0-11.0	178	6.5	5.6-7.6	86	8.1	6.4-10.1	92	5.5	4.4-6.7
2008	307	10.9	9.7-12.2	164	15.7	13.3-18.5	143	8.1	6.8-9.5	180	6.4	5.4-7.4	93	8.9	7.1-11.0	87	5.0	4.0-6.1
2009	359	12.4	11.1-13.7	201	17.6	15.1-20.3	158	9.0	7.7-10.6	159	5.7	4.9-6.7	85	8.5	6.7-10.5	74	4.2	3.3-5.3
2010	346	11.6	10.4-12.9	182	15.6	13.3-18.2	164	9.1	7.7-10.6	159	5.6	4.7-6.5	92	8.8	7.0-10.8	67	3.7	2.9-4.7
2011	375	12.0	10.8-13.3	189	15.5	13.3-18.0	186	9.8	8.4-11.3	171	5.7	4.9-6.6	98	8.6	6.9-10.6	73	3.9	3.0-4.9
2012	383	11.9	10.7-13.2	210	16.5	14.2-19.0	173	8.9	7.6-10.3	165	5.4	4.6-6.3	85	6.9	5.4-8.6	80	4.3	3.4-5.3
2013	379	11.4	10.3-12.6	203	15.7	13.5-18.2	176	8.8	7.5-10.2	172	5.3	4.5-6.2	91	7.3	5.8-9.0	81	4.1	3.2-5.1
2014	375	10.8	9.7-12.0	207	15.1	13.0-17.4	168	8.1	6.9-9.4	189	5.7	4.9-6.6	111	8.4	6.8-10.2	78	4.0	3.2-5.0
2015	403	11.4	10.3-12.6	215	15.2	13.1-17.5	188	9.0	7.7-10.4	149	4.3	3.6-5.0	90	6.9	5.5-8.5	59	2.7	2.0-3.5
2016	398	11.1	10.0-12.2	219	15.3	13.2-17.5	179	8.5	7.3-9.8	196	5.4	4.6-6.2	123	8.4	6.9-10.1	73	3.3	2.6-4.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Colorectal

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	641	48.1	44.1-52.4	273	51.5	44.7-58.9	368	45.9	41.0-51.3	384	31.2	27.8-34.8	168	35.1	29.2-41.8	216	28.9	24.8-33.3
1977	778	59.8	55.2-64.6	319	64.3	56.5-72.9	459	57.0	51.5-63.0	375	29.5	26.3-32.9	174	36.1	30.4-42.5	201	25.4	21.8-29.5
1978	802	60.3	55.8-65.1	334	65.6	57.7-74.2	468	57.5	52.0-63.3	462	36.5	32.9-40.3	202	41.7	35.3-48.7	260	33.5	29.2-38.1
1979	841	60.4	56.0-65.0	369	68.8	61.0-77.2	472	55.5	50.3-61.1	476	35.4	32.0-39.0	206	40.9	34.8-47.7	270	32.3	28.3-36.7
1980	887	60.4	56.2-64.9	391	69.9	62.3-78.2	496	55.1	50.1-60.4	463	32.7	29.6-36.1	197	36.4	30.9-42.6	266	30.7	26.9-34.8
1981	866	57.6	53.6-61.9	375	67.0	59.6-75.1	491	52.7	47.9-57.8	496	35.1	31.9-38.6	201	38.8	33.0-45.2	295	33.3	29.4-37.5
1982	873	54.9	51.1-58.9	398	65.0	58.0-72.5	475	48.9	44.4-53.7	468	30.5	27.6-33.5	208	35.4	30.2-41.2	260	27.6	24.2-31.3
1983	870	52.9	49.2-56.8	383	64.0	56.9-71.5	487	47.5	43.2-52.0	443	27.2	24.5-30.0	203	31.6	26.9-36.8	240	24.3	21.2-27.8
1984	929	55.5	51.8-59.4	430	66.1	59.3-73.4	499	49.1	44.7-53.7	483	29.6	26.8-32.5	211	35.3	30.2-41.0	272	26.7	23.5-30.2
1985	987	57.0	53.3-60.9	447	69.9	62.8-77.4	540	50.1	45.9-54.7	492	30.1	27.3-33.0	201	33.6	28.6-39.1	291	28.4	25.1-32.0
1986	1039	57.8	54.1-61.6	472	69.3	62.5-76.5	567	51.3	47.0-55.8	474	27.5	25.0-30.3	215	34.0	29.1-39.4	259	24.1	21.2-27.3
1987	962	52.2	48.8-55.7	435	63.1	56.7-70.0	527	46.1	42.2-50.3	479	26.9	24.4-29.6	222	34.5	29.6-39.9	257	23.0	20.2-26.1
1988	980	52.4	49.0-56.0	421	59.9	53.7-66.5	559	48.2	44.2-52.5	476	26.7	24.3-29.4	221	34.2	29.3-39.5	255	22.8	20.0-25.8
1989	1019	53.2	49.8-56.7	485	69.0	62.4-76.1	534	44.8	41.0-48.9	469	25.2	22.8-27.6	238	36.3	31.3-41.7	231	19.5	17.0-22.2
1990	1009	52.5	49.1-56.0	434	60.9	54.7-67.4	575	47.7	43.8-51.8	465	25.2	22.9-27.7	212	31.8	27.2-36.9	253	21.8	19.1-24.7
1991	1079	55.0	51.6-58.5	467	62.3	56.2-68.8	612	50.4	46.4-54.6	495	26.5	24.1-29.0	216	32.9	28.2-38.1	279	23.4	20.6-26.3
1992	1073	52.9	49.7-56.3	448	59.8	53.8-66.2	625	49.5	45.6-53.6	477	24.4	22.2-26.7	204	29.6	25.3-34.4	273	21.9	19.4-24.7
1993	1084	52.4	49.2-55.7	491	62.4	56.5-68.8	593	46.5	42.7-50.4	484	24.2	22.0-26.5	211	28.4	24.3-32.9	273	21.8	19.3-24.6
1994	1168	55.5	52.3-58.9	528	67.6	61.4-74.2	640	48.9	45.1-52.9	535	26.1	23.8-28.5	254	32.8	28.6-37.5	281	22.0	19.5-24.7
1995	1164	53.1	50.0-56.4	523	63.6	57.8-69.8	641	47.3	43.7-51.1	536	25.8	23.6-28.1	233	30.0	26.0-34.5	303	23.2	20.6-26.0
1996	1248	56.1	52.9-59.4	558	66.4	60.5-72.7	690	50.1	46.4-54.1	547	25.9	23.7-28.2	263	34.0	29.7-38.6	284	21.3	18.9-24.0
1997	1290	57.3	54.1-60.6	590	70.6	64.5-77.1	700	49.9	46.2-53.8	570	26.1	23.9-28.4	271	34.3	30.0-39.0	299	21.7	19.3-24.4
1998	1289	55.6	52.5-58.8	547	62.2	56.7-68.1	742	51.7	48.0-55.6	513	22.7	20.8-24.8	226	26.9	23.2-30.9	287	20.4	18.1-22.9
1999	1420	58.8	55.7-62.0	602	65.6	60.0-71.4	818	54.9	51.1-58.8	597	25.9	23.8-28.1	255	30.8	26.9-35.1	342	23.4	21.0-26.0
2000	1281	51.3	48.5-54.3	571	61.4	56.1-67.0	710	45.9	42.6-49.5	548	22.7	20.8-24.8	249	27.9	24.3-31.9	299	19.9	17.6-22.2
2001	1365	53.5	50.6-56.4	602	62.9	57.6-68.5	763	48.4	45.0-52.0	543	22.0	20.2-24.0	237	27.2	23.6-31.1	306	19.7	17.6-22.1
2002	1387	52.8	50.0-55.7	599	59.2	54.2-64.4	788	49.0	45.6-52.6	541	21.7	19.9-23.7	248	27.9	24.2-31.8	293	18.6	16.5-20.9
2003	1456	54.6	51.8-57.6	668	67.1	61.7-72.7	788	48.0	44.7-51.5	547	21.4	19.6-23.3	258	28.1	24.5-32.0	289	17.9	15.9-20.1
2004	1479	53.9	51.2-56.8	647	61.8	56.8-67.1	832	49.6	46.2-53.1	536	20.6	18.9-22.5	258	26.7	23.3-30.4	278	17.1	15.2-19.3
2005	1415	50.6	48.0-53.4	650	60.3	55.5-65.4	765	44.9	41.8-48.3	499	18.8	17.2-20.6	220	22.4	19.3-25.7	279	16.7	14.8-18.8
2006	1475	52.1	49.4-54.8	634	57.7	53.0-62.7	841	48.8	45.5-52.3	527	19.4	17.8-21.2	233	23.5	20.4-26.9	294	17.3	15.3-19.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Colorectal

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	1451	49.7	47.2-52.4	688	59.0	54.4-63.8	763	43.5	40.4-46.7	547	19.5	17.9-21.2	251	23.3	20.3-26.5	296	17.2	15.3-19.3
2008	1489	50.1	47.6-52.8	693	59.4	54.8-64.3	796	44.4	41.3-47.6	489	17.2	15.7-18.8	231	21.5	18.6-24.6	258	14.7	12.9-16.6
2009	1463	47.6	45.2-50.2	706	57.2	52.8-61.8	757	41.1	38.2-44.2	507	17.6	16.0-19.2	246	22.7	19.7-25.8	261	14.5	12.8-16.4
2010	1404	45.2	42.8-47.7	667	55.4	51.0-60.1	737	39.4	36.6-42.4	481	15.9	14.5-17.4	247	21.7	18.9-24.7	234	12.4	10.8-14.1
2011	1448	44.8	42.5-47.2	703	54.2	50.1-58.6	745	38.4	35.7-41.3	494	16.1	14.7-17.6	214	18.4	15.9-21.2	280	14.7	13.0-16.6
2012	1360	40.9	38.7-43.2	663	50.0	46.1-54.2	697	35.2	32.6-37.9	537	16.9	15.5-18.4	260	21.3	18.6-24.1	277	14.2	12.6-16.0
2013	1406	41.0	38.9-43.3	662	48.3	44.5-52.3	744	36.4	33.8-39.2	481	14.5	13.2-15.9	223	16.9	14.7-19.4	258	12.7	11.2-14.4
2014	1493	42.7	40.5-44.9	703	49.8	46.0-53.8	790	38.2	35.5-41.0	510	15.0	13.7-16.3	246	18.3	16.0-20.9	264	12.8	11.3-14.5
2015	1415	38.9	36.9-41.1	706	48.1	44.4-51.9	709	33.0	30.5-35.5	465	13.4	12.2-14.7	205	14.9	12.8-17.2	260	12.3	10.8-13.9
2016	1452	39.3	37.2-41.4	726	48.8	45.2-52.7	726	33.1	30.7-35.7	501	13.7	12.5-15.0	231	15.7	13.7-18.0	270	12.2	10.8-13.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Colon excluding rectum

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	470	35.5	32.0-39.2	192	37.6	31.7-44.2	278	34.3	30.1-38.9	312	25.2	22.2-28.5	135	28.6	23.2-34.7	177	23.4	19.8-27.4
1977	565	44.5	40.5-48.7	210	44.0	37.4-51.4	355	45.0	40.0-50.3	290	23.1	20.3-26.2	124	26.8	21.8-32.5	166	21.0	17.7-24.7
1978	603	46.2	42.2-50.4	245	50.3	43.3-58.1	358	44.2	39.4-49.4	382	30.1	26.9-33.6	163	34.4	28.5-40.9	219	27.8	24.0-32.0
1979	653	47.1	43.3-51.2	284	53.1	46.3-60.5	369	43.7	39.0-48.6	404	29.9	26.8-33.2	171	34.2	28.6-40.4	233	27.6	24.0-31.6
1980	653	45.0	41.3-48.9	280	51.1	44.5-58.3	373	41.7	37.3-46.4	378	27.1	24.2-30.2	158	29.9	24.8-35.6	220	25.6	22.1-29.4
1981	630	43.2	39.7-47.0	271	50.3	43.7-57.4	359	39.6	35.4-44.1	427	30.2	27.2-33.4	170	33.1	27.7-39.0	257	28.9	25.3-32.8
1982	648	41.2	37.9-44.7	294	49.4	43.2-56.0	354	36.6	32.7-40.7	390	25.6	23.0-28.5	168	29.8	24.9-35.3	222	23.4	20.3-26.9
1983	655	39.8	36.6-43.2	284	48.2	42.1-54.9	371	35.9	32.2-39.9	371	23.0	20.5-25.6	165	26.3	22.0-31.1	206	20.9	18.0-24.1
1984	688	41.7	38.5-45.2	307	47.5	41.8-53.7	381	38.1	34.2-42.2	420	25.6	23.1-28.4	183	30.6	25.8-35.9	237	23.2	20.3-26.5
1985	743	43.8	40.5-47.2	322	52.1	45.9-58.8	421	39.5	35.7-43.6	421	25.6	23.1-28.4	165	27.8	23.2-32.8	256	24.8	21.7-28.1
1986	769	42.9	39.8-46.2	338	50.7	44.8-57.0	431	38.8	35.1-42.8	411	23.9	21.5-26.4	186	29.3	24.8-34.3	225	20.9	18.2-24.0
1987	735	40.0	37.0-43.2	329	47.6	42.0-53.6	406	35.7	32.2-39.4	423	23.7	21.4-26.2	194	29.9	25.4-34.9	229	20.5	17.8-23.4
1988	767	41.3	38.3-44.5	330	47.1	41.7-53.1	437	38.0	34.5-41.9	416	23.1	20.8-25.6	192	29.1	24.6-34.1	224	19.9	17.3-22.8
1989	789	41.6	38.6-44.7	367	53.1	47.2-59.4	422	35.7	32.3-39.3	417	22.3	20.1-24.7	209	32.0	27.3-37.1	208	17.5	15.1-20.1
1990	768	40.1	37.2-43.2	324	46.1	40.8-51.9	444	36.8	33.4-40.5	405	21.9	19.7-24.3	183	27.1	22.9-31.8	222	19.1	16.6-21.9
1991	827	42.5	39.5-45.6	362	48.9	43.5-54.8	465	38.5	35.0-42.3	446	23.9	21.6-26.3	195	29.5	25.1-34.5	251	21.0	18.5-23.9
1992	828	41.1	38.3-44.2	332	45.3	40.1-51.0	496	39.3	35.9-43.0	414	21.2	19.1-23.4	176	26.0	22.0-30.6	238	19.0	16.7-21.7
1993	814	40.1	37.3-43.0	356	47.0	41.8-52.7	458	36.2	32.9-39.7	430	21.3	19.3-23.5	185	24.6	20.8-28.8	245	19.5	17.1-22.1
1994	914	44.0	41.1-47.0	397	51.6	46.2-57.5	517	39.8	36.4-43.5	469	23.0	20.9-25.2	222	29.1	25.1-33.6	247	19.4	17.0-22.0
1995	892	41.4	38.6-44.3	393	49.1	44.0-54.6	499	37.1	33.9-40.5	482	23.3	21.2-25.5	212	27.7	23.8-32.0	270	20.7	18.3-23.4
1996	935	42.2	39.5-45.1	408	49.3	44.2-54.8	527	38.4	35.1-41.8	469	22.3	20.3-24.5	222	29.2	25.2-33.6	247	18.5	16.2-21.0
1997	991	44.0	41.2-46.9	435	52.7	47.5-58.4	556	39.5	36.2-42.9	492	22.6	20.6-24.7	231	29.9	25.8-34.3	261	18.9	16.7-21.4
1998	995	43.3	40.6-46.2	416	48.6	43.7-53.8	579	40.5	37.2-44.0	453	20.0	18.2-22.0	197	23.4	20.0-27.2	256	18.1	15.9-20.5
1999	1073	44.8	42.1-47.6	434	48.6	43.7-53.7	639	42.9	39.6-46.4	529	23.0	21.0-25.0	217	26.5	22.9-30.6	312	21.3	19.0-23.8
2000	976	39.6	37.1-42.2	423	46.8	42.1-51.8	553	36.0	33.0-39.1	484	20.2	18.4-22.1	218	25.0	21.6-28.9	266	17.7	15.6-19.9
2001	1024	40.7	38.2-43.3	440	47.3	42.7-52.2	584	37.2	34.3-40.4	487	19.9	18.1-21.8	209	24.2	20.8-27.9	278	18.0	15.9-20.3
2002	1079	41.5	39.0-44.1	441	44.3	40.0-48.9	638	39.9	36.8-43.1	483	19.5	17.7-21.3	216	24.5	21.1-28.2	267	17.0	15.0-19.1
2003	1096	41.6	39.1-44.2	486	50.3	45.6-55.3	610	37.3	34.4-40.4	466	18.4	16.7-20.2	209	23.2	20.0-26.8	257	15.9	14.0-18.0
2004	1115	41.2	38.8-43.7	479	46.6	42.2-51.2	636	38.2	35.3-41.3	466	18.0	16.4-19.8	218	22.5	19.4-25.9	248	15.4	13.5-17.4
2005	1074	38.9	36.6-41.3	480	45.6	41.4-50.1	594	35.1	32.3-38.1	432	16.4	14.9-18.1	183	18.8	16.1-21.9	249	15.0	13.2-17.0
2006	1093	39.1	36.8-41.5	463	42.9	38.8-47.2	630	37.0	34.1-40.0	457	16.9	15.4-18.6	204	20.8	17.8-24.0	253	14.9	13.1-16.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Colon excluding rectum

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	1078	37.4	35.1-39.7	501	44.1	40.1-48.3	577	33.0	30.3-35.8	465	16.6	15.1-18.2	208	19.4	16.7-22.4	257	14.9	13.1-16.9
2008	1130	38.6	36.3-41.0	510	44.9	40.8-49.2	620	34.8	32.1-37.7	421	14.9	13.4-16.4	194	18.2	15.6-21.1	227	12.9	11.2-14.7
2009	1070	35.4	33.2-37.6	499	41.5	37.7-45.5	571	31.1	28.6-33.8	441	15.4	14.0-17.0	211	19.9	17.2-22.9	230	12.8	11.2-14.6
2010	1031	33.7	31.6-35.9	472	40.0	36.2-44.0	559	30.2	27.7-32.8	408	13.6	12.3-15.0	209	18.7	16.1-21.6	199	10.6	9.1-12.2
2011	1073	33.6	31.5-35.7	506	40.0	36.4-43.8	567	29.3	26.9-31.9	418	13.8	12.5-15.2	177	15.7	13.3-18.3	241	12.7	11.1-14.4
2012	989	30.2	28.4-32.2	462	36.0	32.6-39.6	527	26.8	24.5-29.2	450	14.2	12.9-15.6	206	16.9	14.5-19.5	244	12.6	11.0-14.3
2013	1001	29.6	27.7-31.5	464	34.6	31.4-38.1	537	26.4	24.2-28.7	418	12.7	11.5-14.0	186	14.1	12.1-16.4	232	11.5	10.1-13.1
2014	1097	31.7	29.8-33.7	501	36.2	33.0-39.7	596	29.0	26.7-31.5	420	12.3	11.1-13.5	198	14.7	12.6-17.0	222	10.8	9.4-12.3
2015	1003	27.8	26.1-29.7	482	33.5	30.4-36.7	521	24.2	22.1-26.4	389	11.2	10.1-12.4	173	12.5	10.7-14.6	216	10.2	8.8-11.7
2016	1044	28.3	26.6-30.2	497	33.7	30.7-36.9	547	25.0	22.9-27.3	395	10.8	9.7-11.9	173	11.6	9.9-13.6	222	10.1	8.8-11.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Rectum & rectosigmoid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	171	12.6	10.6-14.9	81	13.8	10.6-17.6	90	11.6	9.1-14.5	72	6.0	4.5-7.7	33	6.6	4.2-9.6	39	5.5	3.8-7.6
1977	213	15.3	13.1-17.7	109	20.3	16.2-25.0	104	12.1	9.7-14.9	85	6.4	5.0-8.0	50	9.3	6.7-12.5	35	4.4	3.0-6.3
1978	199	14.1	12.0-16.4	89	15.3	11.8-19.3	110	13.3	10.7-16.2	80	6.4	5.0-8.1	39	7.3	5.0-10.2	41	5.7	3.9-7.8
1979	188	13.2	11.2-15.4	85	15.7	12.1-19.9	103	11.8	9.5-14.5	72	5.5	4.2-7.0	35	6.7	4.4-9.6	37	4.7	3.2-6.6
1980	234	15.4	13.4-17.7	111	18.8	15.1-23.1	123	13.4	11.0-16.1	85	5.7	4.5-7.1	39	6.6	4.5-9.2	46	5.1	3.6-6.9
1981	236	14.4	12.5-16.5	104	16.8	13.3-20.7	132	13.1	10.9-15.7	69	4.9	3.7-6.3	31	5.8	3.7-8.4	38	4.4	3.1-6.1
1982	225	13.7	11.9-15.7	104	15.6	12.5-19.2	121	12.3	10.1-14.8	78	4.8	3.7-6.1	40	5.6	3.9-7.9	38	4.2	2.9-5.8
1983	215	13.1	11.3-15.1	99	15.8	12.4-19.6	116	11.5	9.5-13.9	72	4.2	3.2-5.4	38	5.3	3.6-7.5	34	3.4	2.3-4.9
1984	241	13.8	12.0-15.7	123	18.7	15.1-22.7	118	11.0	9.1-13.2	63	3.9	3.0-5.1	28	4.8	3.0-7.0	35	3.5	2.4-4.9
1985	244	13.2	11.5-15.1	125	17.8	14.5-21.6	119	10.6	8.7-12.7	71	4.4	3.4-5.6	36	5.8	3.9-8.3	35	3.6	2.5-5.0
1986	270	14.8	13.0-16.8	134	18.6	15.3-22.4	136	12.4	10.4-14.8	63	3.7	2.8-4.7	29	4.7	2.9-6.9	34	3.2	2.2-4.4
1987	227	12.2	10.6-13.9	106	15.6	12.4-19.2	121	10.5	8.6-12.5	56	3.3	2.4-4.3	28	4.6	2.9-6.8	28	2.5	1.7-3.7
1988	213	11.1	9.6-12.8	91	12.8	10.0-16.0	122	10.2	8.4-12.2	60	3.6	2.7-4.7	29	5.1	3.3-7.4	31	2.9	1.9-4.1
1989	230	11.6	10.1-13.3	118	16.0	12.9-19.4	112	9.1	7.5-11.0	52	2.8	2.1-3.7	29	4.3	2.7-6.4	23	2.0	1.3-3.0
1990	241	12.4	10.8-14.1	110	14.7	11.9-18.0	131	10.9	9.0-12.9	60	3.3	2.5-4.3	29	4.7	2.9-7.0	31	2.7	1.8-3.8
1991	252	12.5	10.9-14.2	105	13.4	10.8-16.5	147	11.8	10.0-13.9	49	2.6	1.9-3.5	21	3.3	1.9-5.2	28	2.3	1.5-3.4
1992	245	11.8	10.3-13.4	116	14.5	11.7-17.7	129	10.2	8.5-12.1	63	3.2	2.4-4.1	28	3.6	2.3-5.3	35	2.9	2.0-4.0
1993	270	12.3	10.9-13.9	135	15.4	12.7-18.5	135	10.3	8.6-12.2	54	2.8	2.1-3.7	26	3.8	2.3-5.7	28	2.4	1.6-3.4
1994	254	11.6	10.1-13.1	131	16.0	13.1-19.2	123	9.1	7.5-10.8	66	3.1	2.4-4.0	32	3.7	2.5-5.3	34	2.6	1.8-3.7
1995	272	11.8	10.4-13.3	130	14.5	11.9-17.4	142	10.2	8.6-12.0	54	2.5	1.9-3.3	21	2.4	1.4-3.6	33	2.5	1.7-3.5
1996	313	13.9	12.3-15.5	150	17.1	14.3-20.3	163	11.8	10.0-13.7	78	3.6	2.8-4.5	41	4.8	3.3-6.6	37	2.8	2.0-3.9
1997	299	13.3	11.8-14.9	155	17.9	15.0-21.1	144	10.4	8.8-12.3	78	3.5	2.8-4.4	40	4.5	3.1-6.2	38	2.8	2.0-3.8
1998	294	12.3	10.9-13.8	131	13.6	11.2-16.3	163	11.2	9.6-13.1	60	2.7	2.1-3.5	29	3.4	2.2-5.0	31	2.3	1.5-3.2
1999	347	14.0	12.5-15.6	168	17.0	14.4-19.9	179	12.0	10.3-13.9	68	2.9	2.2-3.7	38	4.3	2.9-5.9	30	2.1	1.4-3.0
2000	305	11.7	10.4-13.1	148	14.6	12.2-17.3	157	9.9	8.4-11.6	64	2.5	1.9-3.3	31	2.9	1.9-4.1	33	2.2	1.5-3.1
2001	341	12.8	11.5-14.3	162	15.6	13.1-18.4	179	11.1	9.6-12.9	56	2.2	1.6-2.8	28	3.0	1.9-4.5	28	1.7	1.1-2.5
2002	308	11.3	10.0-12.6	158	14.9	12.5-17.5	150	9.1	7.7-10.7	58	2.2	1.7-2.9	32	3.4	2.2-4.9	26	1.7	1.1-2.4
2003	360	13.0	11.7-14.5	182	16.7	14.2-19.5	178	10.7	9.2-12.4	81	3.0	2.4-3.8	49	4.8	3.5-6.5	32	2.0	1.3-2.8
2004	364	12.7	11.4-14.2	168	15.2	12.9-17.9	196	11.4	9.8-13.1	70	2.6	2.0-3.3	40	4.1	2.9-5.7	30	1.8	1.2-2.5
2005	341	11.7	10.5-13.1	170	14.7	12.4-17.2	171	9.8	8.4-11.4	67	2.4	1.9-3.1	37	3.5	2.4-4.9	30	1.8	1.2-2.5
2006	382	13.0	11.7-14.4	171	14.9	12.6-17.4	211	11.9	10.3-13.6	70	2.5	2.0-3.2	29	2.7	1.8-4.0	41	2.4	1.7-3.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Rectum & rectosigmoid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	373	12.4	11.1-13.7	187	14.9	12.7-17.3	186	10.5	9.0-12.1	82	2.9	2.3-3.6	43	3.9	2.7-5.3	39	2.3	1.6-3.1
2008	359	11.6	10.4-12.8	183	14.6	12.4-17.0	176	9.6	8.2-11.1	68	2.4	1.8-3.0	37	3.2	2.2-4.5	31	1.8	1.2-2.5
2009	393	12.3	11.1-13.6	207	15.7	13.5-18.1	186	10.0	8.6-11.5	66	2.1	1.6-2.7	35	2.8	1.9-3.9	31	1.7	1.1-2.4
2010	373	11.5	10.3-12.7	195	15.5	13.2-17.9	178	9.2	7.9-10.7	73	2.3	1.8-2.9	38	3.0	2.0-4.1	35	1.8	1.3-2.6
2011	375	11.2	10.1-12.5	197	14.3	12.3-16.5	178	9.1	7.8-10.5	76	2.3	1.8-2.9	37	2.7	1.9-3.8	39	2.0	1.4-2.8
2012	371	10.7	9.6-11.9	201	14.0	12.1-16.2	170	8.4	7.1-9.8	87	2.7	2.1-3.3	54	4.4	3.2-5.8	33	1.7	1.1-2.3
2013	405	11.5	10.4-12.7	198	13.6	11.7-15.8	207	10.1	8.7-11.6	63	1.8	1.4-2.4	37	2.8	1.9-3.9	26	1.2	0.8-1.8
2014	396	11.0	9.9-12.1	202	13.6	11.7-15.7	194	9.2	7.9-10.6	90	2.7	2.2-3.3	48	3.6	2.6-4.9	42	2.1	1.5-2.8
2015	412	11.1	10.0-12.2	224	14.6	12.7-16.8	188	8.8	7.6-10.2	76	2.2	1.7-2.7	32	2.4	1.6-3.4	44	2.1	1.5-2.8
2016	408	10.9	9.9-12.1	229	15.1	13.2-17.3	179	8.1	6.9-9.4	106	2.9	2.4-3.6	58	4.1	3.1-5.3	48	2.2	1.6-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Liver / intrahepatic bile duct

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	79	5.5	4.2-7.0	55	9.5	6.9-12.7	24	2.9	1.8-4.4	82	6.1	4.7-7.7	58	10.5	7.6-13.9	24	3.2	1.9-4.9
1977	74	4.9	3.7-6.2	51	8.0	5.6-10.9	23	2.8	1.7-4.3	92	5.9	4.7-7.4	66	10.9	8.1-14.3	26	2.8	1.8-4.1
1978	101	6.7	5.3-8.2	60	9.7	7.2-12.8	41	4.6	3.2-6.4	89	5.8	4.6-7.3	45	7.0	4.9-9.7	44	5.0	3.5-6.8
1979	75	4.7	3.6-6.1	44	7.0	4.8-9.7	31	3.3	2.2-4.8	93	6.4	5.0-7.9	51	8.9	6.4-12.0	42	4.8	3.4-6.6
1980	88	5.4	4.2-6.7	59	8.7	6.4-11.6	29	3.2	2.1-4.6	85	5.3	4.1-6.6	55	8.5	6.2-11.4	30	3.3	2.2-4.7
1981	90	5.3	4.2-6.5	57	8.4	6.2-11.2	33	3.3	2.2-4.6	93	5.9	4.7-7.2	55	8.1	5.9-10.8	38	4.3	3.0-5.9
1982	100	5.8	4.7-7.1	73	10.5	8.0-13.5	27	2.7	1.8-4.0	95	5.7	4.6-7.0	63	9.0	6.8-11.7	32	3.5	2.3-5.0
1983	96	5.5	4.4-6.7	65	9.5	7.1-12.5	31	2.9	1.9-4.2	121	7.0	5.8-8.5	77	11.3	8.7-14.4	44	4.4	3.1-5.9
1984	101	5.6	4.5-6.9	69	10.4	7.8-13.6	32	2.8	1.9-4.0	113	6.4	5.2-7.7	68	10.9	8.2-14.2	45	3.9	2.8-5.2
1985	99	5.1	4.1-6.2	62	8.5	6.3-11.1	37	3.1	2.1-4.2	102	5.7	4.6-6.9	58	8.2	6.1-10.9	44	4.1	2.9-5.5
1986	98	5.1	4.1-6.3	65	8.9	6.7-11.6	33	2.8	1.9-4.0	105	5.5	4.5-6.7	66	8.9	6.7-11.5	39	3.4	2.4-4.7
1987	101	5.0	4.0-6.1	57	7.1	5.1-9.4	44	3.7	2.7-5.0	112	6.2	5.0-7.5	57	8.5	6.3-11.3	55	4.9	3.7-6.4
1988	100	4.9	3.9-6.0	65	8.1	6.0-10.6	35	2.9	2.0-4.1	104	5.2	4.2-6.4	67	8.8	6.6-11.4	37	3.1	2.2-4.3
1989	110	5.6	4.6-6.8	68	9.1	6.8-11.8	42	3.6	2.6-4.9	111	5.6	4.5-6.8	61	7.7	5.7-10.2	50	4.2	3.1-5.6
1990	131	6.3	5.2-7.5	81	9.6	7.5-12.1	50	4.1	3.1-5.5	118	5.7	4.7-6.9	65	8.2	6.2-10.7	53	4.3	3.2-5.6
1991	136	6.2	5.1-7.3	90	10.5	8.2-13.2	46	3.5	2.5-4.7	129	6.1	5.1-7.3	76	9.4	7.2-12.0	53	4.1	3.1-5.4
1992	132	6.3	5.2-7.5	83	10.2	8.0-12.9	49	3.9	2.8-5.1	124	5.8	4.8-6.9	72	8.8	6.8-11.2	52	4.0	2.9-5.2
1993	129	6.0	5.0-7.2	82	9.4	7.4-11.9	47	3.7	2.7-4.9	132	6.2	5.2-7.4	76	8.7	6.7-11.0	56	4.5	3.4-5.9
1994	140	6.2	5.2-7.3	84	9.2	7.2-11.6	56	4.2	3.1-5.4	120	5.6	4.6-6.8	76	9.3	7.2-11.8	44	3.4	2.5-4.6
1995	154	6.6	5.6-7.7	100	10.8	8.6-13.4	54	3.9	2.9-5.1	130	5.6	4.7-6.7	79	8.9	6.9-11.2	51	3.7	2.7-4.9
1996	170	7.3	6.2-8.6	120	13.3	10.8-16.2	50	3.7	2.7-4.9	146	6.5	5.5-7.7	90	10.4	8.2-12.9	56	4.1	3.1-5.4
1997	174	7.2	6.2-8.4	110	11.5	9.3-14.0	64	4.5	3.5-5.7	129	5.4	4.5-6.5	78	8.3	6.5-10.6	51	3.6	2.7-4.8
1998	190	7.6	6.5-8.7	131	12.4	10.3-14.9	59	4.1	3.1-5.3	159	6.5	5.5-7.6	105	10.6	8.5-12.9	54	3.8	2.9-5.0
1999	181	7.1	6.0-8.2	110	10.3	8.4-12.6	71	4.7	3.6-5.9	112	4.5	3.7-5.4	65	6.2	4.8-8.0	47	3.2	2.3-4.3
2000	189	7.0	6.0-8.1	126	11.0	9.1-13.2	63	4.0	3.1-5.2	157	6.1	5.2-7.1	103	10.1	8.1-12.4	54	3.6	2.7-4.6
2001	190	6.8	5.9-7.9	141	12.3	10.2-14.6	49	3.1	2.3-4.1	143	5.3	4.5-6.3	86	7.8	6.2-9.8	57	3.6	2.7-4.7
2002	205	7.3	6.4-8.4	147	12.7	10.6-15.1	58	3.7	2.8-4.7	136	5.1	4.3-6.1	86	8.0	6.3-10.1	50	3.2	2.4-4.2
2003	230	7.9	6.9-9.0	162	13.0	11.0-15.3	68	4.1	3.2-5.3	154	5.5	4.6-6.4	107	9.4	7.6-11.5	47	2.9	2.1-3.8
2004	277	9.5	8.4-10.7	195	16.0	13.8-18.6	82	4.9	3.9-6.1	195	6.9	6.0-8.0	119	10.0	8.2-12.1	76	4.7	3.7-5.8
2005	285	9.4	8.3-10.5	215	17.1	14.8-19.6	70	4.0	3.1-5.1	176	6.0	5.2-7.0	121	10.3	8.4-12.3	55	3.3	2.4-4.2
2006	255	8.4	7.4-9.5	190	15.2	13.1-17.7	65	3.7	2.9-4.7	174	5.9	5.0-6.8	127	10.3	8.5-12.4	47	2.8	2.1-3.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Liver / intrahepatic bile duct

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	325	10.3	9.2-11.5	238	18.4	16.0-21.0	87	4.9	3.9-6.0	197	6.6	5.7-7.6	131	10.7	8.8-12.8	66	3.8	2.9-4.8
2008	328	10.0	9.0-11.2	245	17.8	15.5-20.3	83	4.5	3.6-5.6	198	6.3	5.4-7.2	144	11.1	9.3-13.2	54	2.9	2.2-3.8
2009	372	11.3	10.2-12.6	268	19.6	17.2-22.3	104	5.5	4.5-6.7	214	6.7	5.8-7.7	163	12.3	10.4-14.5	51	2.8	2.1-3.7
2010	388	11.3	10.2-12.5	273	18.9	16.7-21.4	115	5.9	4.9-7.1	250	7.5	6.6-8.6	172	12.7	10.8-14.9	78	3.9	3.1-4.9
2011	379	10.9	9.8-12.1	269	17.9	15.7-20.3	110	5.7	4.7-6.9	241	7.1	6.2-8.1	159	10.9	9.2-12.8	82	4.3	3.4-5.4
2012	428	12.0	10.8-13.2	308	20.6	18.2-23.1	120	6.0	4.9-7.1	255	7.4	6.5-8.4	179	12.1	10.4-14.2	76	4.0	3.1-5.0
2013	414	11.1	10.0-12.2	279	17.6	15.5-19.9	135	6.4	5.4-7.6	246	6.9	6.0-7.8	173	11.9	10.1-14.0	73	3.5	2.7-4.4
2014	439	11.4	10.3-12.5	309	18.7	16.6-21.0	130	6.0	5.0-7.2	244	6.6	5.8-7.5	156	10.1	8.5-12.0	88	4.2	3.4-5.2
2015	402	10.3	9.3-11.4	284	17.4	15.3-19.6	118	5.3	4.3-6.3	246	6.5	5.7-7.4	160	10.1	8.6-11.9	86	4.0	3.2-4.9
2016	445	11.0	10.0-12.1	331	19.4	17.3-21.7	114	4.9	4.0-5.9	261	6.8	6.0-7.7	191	12.2	10.5-14.2	70	3.1	2.4-4.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Pancreas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	203	15.3	13.0-17.7	85	14.5	11.3-18.3	118	15.4	12.6-18.7	189	13.4	11.4-15.7	89	15.0	11.7-18.9	100	12.3	9.8-15.1
1977	200	14.1	12.0-16.4	105	17.8	14.0-22.2	95	11.6	9.2-14.3	182	12.8	10.8-15.0	99	16.9	13.3-21.2	83	10.1	7.9-12.7
1978	218	15.3	13.2-17.7	101	16.3	12.9-20.4	117	14.3	11.7-17.4	184	12.5	10.6-14.7	82	13.5	10.4-17.2	102	11.8	9.5-14.5
1979	229	16.1	13.9-18.6	113	19.1	15.3-23.5	116	14.1	11.5-17.0	203	14.3	12.2-16.6	87	13.4	10.5-16.9	116	14.3	11.7-17.4
1980	208	13.7	11.8-15.9	112	18.7	15.0-23.1	96	10.6	8.5-13.0	177	11.9	10.1-13.9	96	16.6	13.0-20.8	81	9.1	7.1-11.4
1981	240	15.5	13.5-17.8	128	21.1	17.2-25.5	112	12.1	9.8-14.7	201	12.9	11.1-15.0	108	17.4	13.8-21.5	93	10.2	8.1-12.5
1982	236	14.9	13.0-17.1	102	16.9	13.4-20.9	134	13.8	11.4-16.4	226	14.6	12.6-16.7	100	17.4	13.8-21.6	126	13.1	10.8-15.7
1983	270	15.8	13.8-17.9	131	19.0	15.6-23.0	139	13.6	11.4-16.2	243	14.5	12.7-16.6	126	19.6	15.9-23.7	117	11.5	9.4-13.9
1984	261	15.6	13.7-17.7	114	18.1	14.5-22.1	147	14.3	12.0-16.9	233	13.9	12.1-15.9	117	18.6	15.0-22.7	116	11.2	9.2-13.5
1985	242	13.9	12.1-15.8	125	17.7	14.5-21.5	117	11.3	9.3-13.6	206	12.2	10.5-14.1	103	15.9	12.7-19.7	103	10.1	8.1-12.2
1986	259	14.3	12.5-16.2	115	15.5	12.6-18.9	144	13.2	11.1-15.6	224	12.7	11.0-14.6	104	14.3	11.4-17.6	120	11.5	9.5-13.7
1987	281	15.1	13.3-17.1	125	16.6	13.5-20.1	156	13.9	11.8-16.3	242	13.1	11.4-14.9	106	15.3	12.1-18.9	136	11.9	9.9-14.1
1988	274	14.9	13.2-16.9	131	18.8	15.5-22.7	143	12.6	10.6-14.9	216	11.8	10.2-13.6	110	15.0	12.1-18.4	106	9.6	7.8-11.7
1989	255	13.4	11.7-15.2	104	14.4	11.5-17.7	151	12.7	10.7-15.0	234	12.4	10.8-14.1	99	13.6	10.8-16.8	135	11.5	9.6-13.6
1990	298	15.9	14.1-17.9	122	17.8	14.4-21.5	176	15.0	12.8-17.4	243	12.8	11.2-14.6	90	13.0	10.1-16.3	153	12.8	10.8-15.0
1991	255	13.0	11.4-14.8	123	16.3	13.3-19.9	132	11.0	9.2-13.1	221	11.1	9.6-12.7	111	14.2	11.4-17.4	110	9.1	7.5-11.0
1992	261	13.1	11.5-14.8	104	14.2	11.3-17.5	157	12.5	10.6-14.7	205	10.8	9.3-12.4	81	12.3	9.5-15.6	124	10.2	8.5-12.2
1993	244	12.0	10.5-13.6	106	13.7	10.9-16.8	138	10.9	9.1-12.9	204	10.1	8.7-11.6	84	11.8	9.1-14.9	120	9.3	7.7-11.2
1994	296	14.2	12.6-16.0	124	15.6	12.8-18.9	172	13.3	11.4-15.5	257	12.6	11.0-14.2	104	13.6	10.9-16.6	153	11.9	10.1-14.0
1995	263	12.4	10.9-14.1	117	14.2	11.6-17.2	146	11.1	9.3-13.1	240	11.2	9.8-12.8	108	13.7	11.0-16.8	132	9.9	8.2-11.7
1996	305	13.7	12.2-15.4	122	14.3	11.7-17.2	183	13.3	11.4-15.4	242	11.1	9.7-12.6	95	11.7	9.3-14.5	147	10.7	9.1-12.6
1997	303	13.5	12.0-15.2	128	14.1	11.7-17.0	175	12.7	10.9-14.8	240	11.0	9.6-12.5	103	11.8	9.5-14.4	137	10.1	8.5-12.0
1998	292	12.8	11.3-14.3	111	12.3	10.0-15.1	181	12.8	11.0-14.8	257	11.5	10.1-13.0	102	11.6	9.3-14.3	155	11.2	9.5-13.1
1999	336	14.3	12.8-15.9	136	15.1	12.5-18.0	200	13.7	11.8-15.7	286	12.2	10.8-13.7	108	11.7	9.5-14.3	178	12.2	10.5-14.1
2000	320	13.5	12.0-15.0	110	12.0	9.7-14.5	210	14.0	12.2-16.1	256	10.7	9.4-12.1	109	12.2	9.9-14.8	147	9.8	8.2-11.5
2001	360	14.4	12.9-16.0	140	14.3	11.9-17.0	220	14.3	12.4-16.3	284	11.7	10.3-13.1	105	11.1	9.0-13.6	179	11.7	10.1-13.6
2002	344	13.6	12.1-15.1	144	14.0	11.7-16.7	200	12.9	11.2-14.8	288	11.4	10.1-12.8	134	13.1	10.9-15.7	154	9.9	8.4-11.6
2003	320	12.4	11.0-13.8	158	16.1	13.5-19.0	162	10.1	8.6-11.8	258	10.2	9.0-11.5	114	11.9	9.6-14.4	144	9.1	7.7-10.8
2004	341	12.8	11.5-14.3	144	13.6	11.3-16.2	197	12.1	10.4-13.9	294	11.1	9.8-12.5	133	13.3	11.0-15.9	161	9.8	8.3-11.4
2005	347	12.6	11.3-14.0	148	13.9	11.7-16.5	199	11.7	10.1-13.5	269	10.0	8.8-11.2	120	11.2	9.1-13.5	149	9.0	7.6-10.6
2006	405	14.4	13.0-15.9	171	15.0	12.7-17.6	234	13.7	12.0-15.6	277	9.9	8.8-11.2	115	10.7	8.7-12.9	162	9.4	8.0-11.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Pancreas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	389	13.8	12.5-15.3	173	15.5	13.2-18.2	216	12.6	11.0-14.4	300	10.8	9.6-12.1	136	12.5	10.3-14.9	164	9.7	8.2-11.3
2008	377	13.1	11.8-14.5	149	12.7	10.7-15.1	228	12.9	11.3-14.7	308	10.8	9.6-12.1	123	11.2	9.2-13.5	185	10.6	9.1-12.2
2009	390	13.2	11.8-14.6	189	16.1	13.7-18.7	201	11.1	9.6-12.8	321	11.0	9.8-12.3	141	12.4	10.3-14.7	180	10.0	8.6-11.6
2010	421	13.8	12.5-15.2	193	15.8	13.5-18.3	228	12.4	10.8-14.2	297	10.1	9.0-11.3	131	11.0	9.1-13.1	166	9.3	8.0-10.9
2011	500	15.8	14.4-17.3	219	16.8	14.6-19.4	281	14.9	13.2-16.7	361	11.8	10.6-13.1	160	13.5	11.3-15.8	201	10.7	9.2-12.3
2012	481	14.5	13.2-15.9	196	14.9	12.8-17.2	285	14.4	12.8-16.2	372	11.3	10.2-12.5	141	10.4	8.7-12.4	231	11.9	10.4-13.5
2013	504	14.9	13.6-16.3	229	16.8	14.5-19.2	275	13.6	12.0-15.4	343	10.3	9.3-11.5	161	12.5	10.6-14.7	182	9.0	7.7-10.4
2014	501	14.3	13.0-15.6	225	15.3	13.3-17.5	276	13.4	11.8-15.1	349	10.2	9.1-11.4	151	10.8	9.0-12.7	198	9.7	8.4-11.2
2015	557	15.6	14.3-17.0	253	17.1	15.0-19.5	304	14.4	12.8-16.2	381	11.0	9.9-12.2	173	12.1	10.3-14.1	208	10.1	8.7-11.6
2016	529	14.4	13.2-15.8	247	16.4	14.3-18.7	282	12.9	11.4-14.5	409	11.2	10.1-12.3	202	13.5	11.6-15.6	207	9.5	8.2-10.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Larynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	115	6.7	5.5-8.2	95	13.3	10.6-16.4	20	2.1	1.2-3.3	66	4.4	3.3-5.7	52	8.8	6.2-11.9	14	1.5	0.8-2.7
1977	112	6.9	5.6-8.4	92	14.3	11.1-18.0	20	2.1	1.3-3.2	54	3.3	2.4-4.4	41	6.8	4.4-9.8	13	1.2	0.7-2.1
1978	124	7.0	5.8-8.4	101	14.2	11.3-17.5	23	2.2	1.4-3.3	83	4.7	3.7-5.8	71	9.9	7.5-12.8	12	1.1	0.6-1.9
1979	147	8.2	6.9-9.7	114	16.0	13.0-19.4	33	3.1	2.1-4.4	68	3.8	2.9-4.8	57	8.0	6.0-10.6	11	1.0	0.5-1.7
1980	138	7.3	6.1-8.7	108	14.2	11.5-17.4	30	2.7	1.8-3.9	66	3.6	2.8-4.6	45	6.4	4.6-8.8	21	1.8	1.1-2.8
1981	132	6.9	5.7-8.2	107	13.5	10.9-16.5	25	2.3	1.5-3.5	74	3.9	3.0-4.9	51	7.0	5.1-9.4	23	1.9	1.2-2.9
1982	121	6.1	5.1-7.4	107	13.3	10.8-16.2	14	1.2	0.7-2.1	58	3.1	2.3-4.1	47	6.5	4.6-8.8	11	1.0	0.5-1.8
1983	138	7.0	5.9-8.3	116	14.4	11.7-17.5	22	2.0	1.3-3.1	64	3.3	2.5-4.2	53	6.7	4.9-8.9	11	1.0	0.5-1.8
1984	147	7.3	6.2-8.6	108	13.2	10.7-16.2	39	3.3	2.3-4.5	73	3.7	2.8-4.6	59	7.3	5.4-9.5	14	1.2	0.7-2.1
1985	188	9.0	7.8-10.4	147	17.9	15.0-21.1	41	3.2	2.3-4.4	67	3.4	2.6-4.4	55	7.1	5.3-9.4	12	1.1	0.5-1.9
1986	179	8.8	7.5-10.2	141	17.8	14.8-21.3	38	3.0	2.1-4.2	81	4.0	3.2-5.1	63	8.1	6.1-10.5	18	1.5	0.9-2.4
1987	152	7.2	6.1-8.4	122	14.4	11.8-17.3	30	2.4	1.6-3.5	78	3.8	3.0-4.8	57	7.2	5.3-9.5	21	1.7	1.1-2.6
1988	150	6.7	5.6-7.9	112	12.5	10.1-15.1	38	2.9	2.0-3.9	88	4.1	3.3-5.1	75	9.1	7.0-11.6	13	0.9	0.5-1.6
1989	178	7.9	6.8-9.2	115	13.1	10.7-16.0	63	4.7	3.6-6.0	71	3.3	2.5-4.1	53	6.5	4.7-8.7	18	1.3	0.8-2.1
1990	180	7.9	6.8-9.2	138	15.5	12.8-18.5	42	3.2	2.3-4.3	62	2.9	2.2-3.7	49	6.0	4.3-8.1	13	1.0	0.5-1.7
1991	150	6.6	5.5-7.7	116	13.2	10.7-16.0	34	2.4	1.7-3.4	72	3.2	2.5-4.1	57	6.5	4.8-8.5	15	1.1	0.6-1.8
1992	189	7.9	6.8-9.1	150	15.9	13.3-18.7	39	2.7	1.9-3.7	82	3.6	2.8-4.4	63	7.2	5.5-9.4	19	1.3	0.8-2.1
1993	154	6.7	5.6-7.8	123	13.9	11.4-16.7	31	2.2	1.5-3.2	83	3.6	2.9-4.5	62	7.1	5.3-9.3	21	1.5	0.9-2.3
1994	148	6.2	5.2-7.3	114	12.1	9.9-14.7	34	2.4	1.6-3.3	78	3.4	2.6-4.2	56	6.2	4.5-8.1	22	1.6	1.0-2.4
1995	155	6.4	5.4-7.5	114	11.8	9.6-14.3	41	2.9	2.0-3.9	90	3.9	3.1-4.8	70	8.2	6.2-10.5	20	1.4	0.8-2.1
1996	154	6.2	5.2-7.2	127	13.0	10.7-15.6	27	1.8	1.2-2.6	80	3.3	2.6-4.2	69	7.8	5.9-10.0	11	0.7	0.4-1.3
1997	142	5.6	4.7-6.6	107	10.6	8.6-12.9	35	2.3	1.6-3.2	68	2.7	2.1-3.4	54	5.4	4.0-7.1	14	0.9	0.5-1.6
1998	159	6.2	5.2-7.2	110	11.1	9.0-13.5	49	3.2	2.3-4.2	67	2.7	2.1-3.4	57	5.7	4.3-7.4	10	0.7	0.3-1.2
1999	137	5.2	4.3-6.1	103	10.0	8.1-12.3	34	2.1	1.5-3.0	65	2.5	1.9-3.2	50	5.0	3.6-6.7	15	1.0	0.5-1.6
2000	129	4.8	4.0-5.7	92	8.8	7.0-10.9	37	2.3	1.6-3.2	67	2.7	2.0-3.4	49	5.0	3.7-6.7	18	1.2	0.7-1.9
2001	124	4.4	3.6-5.2	96	8.3	6.7-10.1	28	1.7	1.1-2.5	45	1.7	1.2-2.3	36	3.6	2.5-5.1	9	0.6	0.3-1.0
2002	124	4.3	3.6-5.1	89	7.6	6.0-9.4	35	2.1	1.4-2.9	70	2.5	1.9-3.2	52	4.5	3.3-6.0	18	1.1	0.7-1.8
2003	147	5.1	4.3-6.0	109	9.7	7.9-11.8	38	2.2	1.6-3.1	57	2.0	1.5-2.6	45	4.1	3.0-5.6	12	0.7	0.4-1.2
2004	131	4.3	3.6-5.1	110	8.9	7.3-10.8	21	1.2	0.7-1.8	61	2.2	1.6-2.8	50	4.5	3.3-6.1	11	0.7	0.3-1.2
2005	123	4.3	3.5-5.1	95	8.5	6.8-10.5	28	1.6	1.0-2.3	51	1.7	1.3-2.3	44	3.8	2.7-5.1	7	0.4	0.2-0.8
2006	125	4.0	3.3-4.7	96	7.6	6.1-9.3	29	1.5	1.0-2.2	50	1.8	1.3-2.3	42	3.8	2.7-5.2	8	0.5	0.2-0.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Larynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	140	4.6	3.8-5.4	107	8.8	7.1-10.7	33	1.8	1.2-2.5	54	1.8	1.3-2.3	42	3.6	2.5-4.9	12	0.7	0.3-1.2
2008	153	5.0	4.2-5.9	106	8.9	7.2-10.8	47	2.6	1.9-3.5	61	2.0	1.5-2.6	41	3.5	2.5-4.8	20	1.1	0.6-1.6
2009	121	3.8	3.2-4.6	92	7.5	6.0-9.3	29	1.5	1.0-2.2	57	1.8	1.4-2.4	42	3.6	2.5-4.9	15	0.8	0.4-1.3
2010	128	3.9	3.2-4.7	96	7.4	6.0-9.2	32	1.6	1.1-2.3	47	1.6	1.1-2.1	39	3.4	2.4-4.7	8	0.4	0.2-0.8
2011	129	3.7	3.1-4.4	97	6.9	5.5-8.5	32	1.5	1.0-2.1	43	1.3	0.9-1.8	30	2.4	1.6-3.4	13	0.7	0.4-1.2
2012	103	2.9	2.4-3.6	74	5.4	4.2-6.8	29	1.4	0.9-2.0	52	1.6	1.2-2.1	43	3.5	2.5-4.7	9	0.4	0.2-0.8
2013	116	3.2	2.7-3.9	89	6.4	5.1-8.0	27	1.2	0.8-1.8	47	1.4	1.0-1.8	36	2.7	1.9-3.8	11	0.5	0.3-0.9
2014	134	3.7	3.1-4.4	105	7.1	5.8-8.7	29	1.4	0.9-2.0	54	1.5	1.1-2.0	46	3.4	2.4-4.5	8	0.4	0.2-0.7
2015	153	4.0	3.4-4.7	125	8.4	6.9-10.1	28	1.2	0.8-1.8	35	1.0	0.7-1.4	26	2.0	1.3-3.0	9	0.4	0.2-0.8
2016	115	3.0	2.5-3.7	87	5.8	4.6-7.2	28	1.2	0.8-1.7	49	1.3	1.0-1.7	37	2.6	1.8-3.7	12	0.5	0.3-0.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Lung and bronchus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	891	57.2	53.2-61.3	668	104.9	96.3-114.1	223	24.9	21.5-28.6	680	43.7	40.2-47.4	517	82.0	74.2-90.3	163	18.0	15.2-21.2
1977	956	60.0	56.0-64.3	706	113.4	104.2-123.2	250	25.5	22.3-29.0	759	47.7	44.1-51.5	573	92.0	83.7-100.8	186	19.0	16.3-22.1
1978	988	61.1	57.1-65.3	730	110.7	102.0-119.9	258	28.2	24.7-32.0	827	51.1	47.5-55.0	613	94.0	86.0-102.5	214	22.9	19.8-26.4
1979	1065	65.4	61.3-69.7	767	119.4	110.1-129.2	298	30.9	27.3-34.8	855	53.5	49.7-57.4	639	101.4	92.7-110.6	216	23.0	19.9-26.5
1980	1123	65.3	61.3-69.5	825	122.5	113.4-132.1	298	28.8	25.5-32.4	904	53.6	49.9-57.4	667	100.8	92.5-109.7	237	23.7	20.6-27.0
1981	1160	65.1	61.3-69.2	824	118.4	109.7-127.6	336	31.1	27.8-34.8	957	54.7	51.1-58.4	685	100.4	92.3-109.1	272	25.7	22.7-29.1
1982	1176	64.7	60.9-68.7	826	116.2	107.7-125.1	350	32.0	28.6-35.6	945	52.5	49.0-56.1	667	93.3	85.8-101.3	278	26.3	23.2-29.7
1983	1193	64.2	60.4-68.1	831	114.0	105.8-122.7	362	32.6	29.2-36.3	1048	57.3	53.7-61.0	753	106.6	98.5-115.2	295	26.5	23.5-29.8
1984	1243	65.1	61.4-69.0	844	112.4	104.3-120.8	399	35.2	31.7-39.0	978	51.6	48.3-55.1	689	94.5	86.9-102.5	289	25.2	22.3-28.3
1985	1263	64.9	61.2-68.7	873	116.2	108.0-124.9	390	33.2	29.9-36.7	1020	53.3	50.0-56.9	714	97.2	89.6-105.3	306	26.5	23.5-29.7
1986	1304	65.9	62.2-69.8	857	114.0	105.8-122.7	447	37.0	33.6-40.7	1021	52.0	48.7-55.4	667	89.4	82.1-97.1	354	29.5	26.4-32.8
1987	1328	65.0	61.4-68.7	898	113.9	106.0-122.1	430	34.8	31.5-38.3	1097	55.0	51.6-58.4	732	97.1	89.6-105.1	365	30.0	26.9-33.3
1988	1325	64.6	61.1-68.3	867	112.7	104.7-121.1	458	36.4	33.0-39.9	1052	52.1	48.8-55.4	698	91.9	84.6-99.6	354	28.6	25.6-31.8
1989	1402	66.4	62.8-70.1	924	116.4	108.5-124.8	478	36.9	33.6-40.5	1126	54.5	51.3-57.9	761	99.7	92.1-107.7	365	28.5	25.6-31.6
1990	1396	65.6	62.1-69.2	918	114.0	106.1-122.3	478	36.7	33.5-40.3	1111	52.6	49.4-55.8	735	90.8	83.8-98.2	376	29.5	26.5-32.6
1991	1394	65.0	61.6-68.7	890	110.3	102.6-118.4	504	38.4	35.1-42.0	1081	50.9	47.9-54.2	707	89.2	82.2-96.6	374	28.6	25.7-31.7
1992	1434	65.2	61.8-68.8	930	113.1	105.2-121.3	504	37.4	34.1-40.8	1063	49.9	46.8-53.1	665	84.7	77.8-92.0	398	30.1	27.2-33.3
1993	1372	61.8	58.5-65.3	851	103.6	96.1-111.5	521	38.1	34.9-41.6	1085	49.2	46.3-52.4	696	86.8	79.9-94.1	389	28.4	25.6-31.4
1994	1461	63.7	60.4-67.2	883	102.7	95.5-110.2	578	41.0	37.7-44.5	1179	52.5	49.5-55.7	769	91.3	84.4-98.6	410	30.0	27.1-33.0
1995	1406	61.1	57.9-64.4	863	97.8	90.9-105.0	543	38.9	35.6-42.3	1163	51.0	48.0-54.1	704	82.2	75.8-88.9	459	32.7	29.8-35.9
1996	1538	65.2	61.9-68.6	910	100.5	93.6-107.7	628	44.0	40.6-47.6	1187	51.1	48.1-54.1	724	81.1	74.8-87.7	463	33.0	30.0-36.2
1997	1492	62.6	59.4-65.9	872	96.0	89.3-103.0	620	42.6	39.3-46.1	1154	49.4	46.5-52.3	696	79.6	73.4-86.2	458	31.8	28.9-34.9
1998	1502	61.0	57.9-64.2	846	88.6	82.3-95.1	656	44.1	40.7-47.6	1158	48.3	45.5-51.2	691	75.5	69.7-81.8	467	32.0	29.1-35.0
1999	1416	56.7	53.7-59.8	811	84.6	78.5-91.0	605	39.9	36.7-43.2	1082	44.3	41.6-47.1	620	66.7	61.2-72.5	462	31.1	28.3-34.1
2000	1477	57.2	54.3-60.3	859	86.6	80.6-92.9	618	39.5	36.4-42.7	1109	43.5	41.0-46.2	654	67.8	62.4-73.5	455	29.2	26.5-32.0
2001	1472	56.6	53.7-59.7	842	85.7	79.7-92.0	630	39.6	36.6-42.8	1095	42.7	40.2-45.4	625	65.2	59.9-70.9	470	29.8	27.1-32.6
2002	1489	55.8	53.0-58.8	826	82.3	76.4-88.4	663	40.6	37.6-43.9	1123	42.5	40.0-45.1	645	64.8	59.6-70.3	478	29.4	26.8-32.2
2003	1455	53.3	50.5-56.2	770	73.4	68.0-79.1	685	41.3	38.3-44.6	1098	40.9	38.4-43.4	613	60.0	55.0-65.2	485	29.5	26.9-32.2
2004	1555	55.7	52.9-58.6	853	77.5	72.1-83.1	702	41.7	38.7-45.0	1068	39.1	36.7-41.5	599	57.4	52.7-62.5	469	28.0	25.5-30.7
2005	1544	54.7	51.9-57.5	857	79.5	74.0-85.4	687	40.1	37.2-43.3	1112	39.9	37.5-42.4	602	55.7	51.1-60.6	510	30.3	27.7-33.0
2006	1518	53.1	50.4-55.8	818	73.1	67.9-78.5	700	40.6	37.6-43.7	1036	37.0	34.7-39.3	546	50.7	46.3-55.3	490	28.7	26.2-31.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Lung and bronchus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	1560	53.2	50.5-56.0	833	73.2	68.1-78.7	727	41.1	38.1-44.2	1081	37.4	35.2-39.8	591	52.7	48.3-57.4	490	28.1	25.7-30.8
2008	1560	52.8	50.1-55.5	813	71.0	65.9-76.3	747	41.9	38.9-45.0	1005	34.6	32.5-36.9	526	46.8	42.7-51.2	479	27.1	24.7-29.7
2009	1634	53.7	51.1-56.4	862	72.5	67.5-77.8	772	42.1	39.2-45.2	1027	34.4	32.2-36.6	568	49.0	44.8-53.4	459	25.3	23.0-27.8
2010	1648	52.5	50.0-55.2	840	68.2	63.4-73.2	808	43.1	40.1-46.2	1120	36.5	34.4-38.8	610	52.5	48.2-57.1	510	27.4	25.0-29.9
2011	1703	53.0	50.4-55.6	867	69.4	64.6-74.5	836	43.1	40.2-46.1	1129	35.6	33.5-37.8	599	48.5	44.5-52.7	530	27.6	25.3-30.1
2012	1704	51.4	49.0-54.0	877	67.9	63.2-72.8	827	41.5	38.7-44.5	1109	34.2	32.2-36.3	599	48.1	44.1-52.3	510	25.9	23.7-28.3
2013	1591	46.7	44.4-49.1	815	61.0	56.7-65.6	776	37.9	35.2-40.7	1081	32.4	30.5-34.5	576	44.5	40.7-48.5	505	24.9	22.8-27.2
2014	1665	47.5	45.2-49.9	858	61.8	57.6-66.3	807	38.3	35.7-41.1	1019	29.6	27.7-31.5	543	40.3	36.8-44.0	476	22.7	20.7-24.9
2015	1712	46.9	44.6-49.2	880	60.9	56.8-65.3	832	38.1	35.5-40.8	1017	28.6	26.9-30.5	514	37.2	33.9-40.7	503	23.4	21.4-25.6
2016	1676	45.3	43.1-47.5	836	56.0	52.1-60.1	840	38.1	35.5-40.8	955	25.9	24.3-27.7	523	36.1	32.9-39.4	432	19.4	17.6-21.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Melanoma of the skin

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	10	0.7	0.3-1.4	2	0.3	0.0-1.3	8	1.0	0.4-2.0	9	0.8	0.3-1.6	4	1.1	0.2-3.0	5	0.7	0.2-1.7
1977	14	1.2	0.6-2.1	8	1.9	0.7-4.0	6	0.9	0.3-1.9	7	0.6	0.2-1.3	4	1.3	0.3-3.3	3	0.3	0.1-0.8
1978	13	0.9	0.4-1.6	5	1.2	0.3-2.9	8	0.9	0.3-1.7	6	0.4	0.1-0.9	2	0.2	0.0-1.0	4	0.5	0.1-1.3
1979	18	1.0	0.6-1.6	7	0.9	0.3-1.9	11	1.1	0.5-2.0	4	0.4	0.1-0.9	2	0.4	0.0-1.4	2	0.3	0.0-1.1
1980	14	0.9	0.5-1.6	5	0.6	0.2-1.5	9	1.1	0.5-2.1	14	0.9	0.4-1.5	8	0.9	0.4-1.9	6	0.7	0.3-1.6
1981	25	1.4	0.9-2.2	7	0.8	0.3-1.8	18	1.8	1.0-2.9	15	0.9	0.5-1.6	7	1.0	0.4-2.1	8	0.9	0.3-1.8
1982	19	1.0	0.6-1.5	8	0.9	0.4-1.8	11	1.0	0.5-1.8	10	0.5	0.2-1.0	6	1.1	0.3-2.5	4	0.3	0.1-0.8
1983	12	0.7	0.3-1.2	4	0.5	0.1-1.5	8	0.7	0.3-1.4	20	1.1	0.6-1.7	14	1.7	0.9-2.8	6	0.6	0.2-1.4
1984	26	1.5	0.9-2.2	7	1.0	0.3-2.2	19	1.8	1.0-2.8	13	0.6	0.3-1.0	10	1.1	0.5-2.1	3	0.2	0.0-0.7
1985	23	1.3	0.8-2.0	12	1.4	0.6-2.6	11	1.2	0.6-2.1	14	0.8	0.4-1.4	10	1.7	0.8-3.2	4	0.3	0.1-0.9
1986	20	1.0	0.6-1.6	11	1.7	0.7-3.2	9	0.8	0.3-1.5	21	1.1	0.7-1.7	15	1.8	1.0-3.1	6	0.6	0.2-1.2
1987	17	0.8	0.5-1.3	8	0.9	0.4-1.9	9	0.7	0.3-1.4	9	0.5	0.2-0.9	4	0.5	0.1-1.2	5	0.4	0.1-1.0
1988	20	1.0	0.6-1.5	7	1.1	0.4-2.3	13	1.0	0.5-1.7	9	0.5	0.2-1.0	4	0.7	0.1-1.8	5	0.5	0.1-1.0
1989	17	1.0	0.6-1.6	6	0.8	0.2-1.7	11	1.1	0.5-1.9	17	0.9	0.5-1.4	6	0.6	0.2-1.5	11	0.9	0.5-1.7
1990	21	0.9	0.6-1.5	10	1.2	0.5-2.3	11	0.8	0.4-1.5	6	0.3	0.1-0.7	3	0.5	0.1-1.5	3	0.3	0.1-0.7
1991	26	1.1	0.7-1.7	13	2.0	0.9-3.6	13	0.8	0.4-1.4	11	0.6	0.3-1.0	6	0.8	0.2-1.8	5	0.4	0.1-1.0
1992	14	0.6	0.3-0.9	6	0.5	0.2-1.2	8	0.6	0.2-1.1	7	0.3	0.1-0.6	3	0.3	0.1-0.8	4	0.3	0.1-0.7
1993	25	1.2	0.8-1.8	12	1.5	0.7-2.7	13	1.0	0.5-1.7	8	0.3	0.1-0.6	2	0.2	0.0-0.7	6	0.4	0.1-0.9
1994	18	0.8	0.4-1.2	5	0.6	0.1-1.6	13	0.9	0.5-1.6	6	0.3	0.1-0.7	1	0.2	0.0-0.8	5	0.4	0.1-0.9
1995	25	1.1	0.7-1.6	7	0.6	0.2-1.3	18	1.3	0.8-2.1	6	0.3	0.1-0.6	1	0.1	0.0-0.5	5	0.4	0.1-0.8
1996	23	1.1	0.7-1.6	8	1.0	0.4-2.0	15	1.1	0.6-1.8	13	0.6	0.3-1.1	4	0.5	0.1-1.3	9	0.7	0.3-1.3
1997	37	1.6	1.1-2.2	15	1.6	0.8-2.7	22	1.5	1.0-2.3	9	0.4	0.2-0.8	4	0.3	0.1-0.8	5	0.4	0.1-0.9
1998	28	1.2	0.8-1.7	13	1.5	0.7-2.6	15	1.0	0.6-1.7	11	0.4	0.2-0.8	3	0.2	0.0-0.7	8	0.6	0.2-1.1
1999	12	0.5	0.2-0.8	5	0.4	0.1-1.1	7	0.5	0.2-0.9	8	0.3	0.1-0.6	5	0.6	0.2-1.4	3	0.2	0.0-0.6
2000	12	0.5	0.2-0.8	6	0.6	0.2-1.3	6	0.4	0.1-0.8	13	0.6	0.3-1.0	7	1.1	0.4-2.2	6	0.4	0.2-0.9
2001	29	1.0	0.7-1.5	11	0.9	0.4-1.6	18	1.1	0.7-1.8	6	0.2	0.1-0.5	3	0.3	0.1-0.8	3	0.2	0.0-0.6
2002	25	0.9	0.6-1.4	9	0.8	0.3-1.5	16	1.0	0.6-1.6	7	0.3	0.1-0.6	3	0.4	0.0-1.1	4	0.3	0.1-0.7
2003	20	0.7	0.4-1.1	5	0.6	0.2-1.4	15	0.9	0.5-1.4	10	0.3	0.2-0.6	3	0.3	0.1-0.9	7	0.4	0.2-0.8
2004	24	0.8	0.5-1.3	12	1.2	0.6-2.1	12	0.7	0.3-1.2	4	0.1	0.0-0.4	2	0.1	0.0-0.5	2	0.1	0.0-0.4
2005	28	0.9	0.6-1.4	12	0.9	0.4-1.5	16	0.9	0.5-1.5	6	0.2	0.1-0.4	4	0.3	0.1-0.7	2	0.1	0.0-0.4
2006	38	1.4	1.0-1.9	19	1.9	1.1-3.0	19	1.1	0.7-1.7	11	0.4	0.2-0.7	6	0.5	0.2-1.1	5	0.3	0.1-0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Melanoma of the skin

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	42	1.4	1.0-1.9	12	1.1	0.5-2.0	30	1.7	1.2-2.5	16	0.6	0.3-0.9	6	0.7	0.2-1.5	10	0.6	0.3-1.1
2008	39	1.3	0.9-1.7	15	1.2	0.7-2.1	24	1.3	0.9-2.0	10	0.3	0.2-0.6	3	0.2	0.0-0.6	7	0.4	0.2-0.8
2009	30	0.9	0.6-1.3	11	0.7	0.4-1.3	19	1.0	0.6-1.6	8	0.3	0.1-0.5	5	0.4	0.1-0.9	3	0.2	0.0-0.5
2010	30	0.9	0.6-1.3	11	0.8	0.4-1.5	19	0.9	0.6-1.5	11	0.4	0.2-0.6	3	0.3	0.1-0.9	8	0.4	0.2-0.8
2011	40	1.2	0.9-1.7	19	1.5	0.8-2.3	21	1.1	0.7-1.7	15	0.5	0.3-0.9	6	0.5	0.2-1.1	9	0.5	0.2-1.0
2012	30	0.9	0.6-1.3	14	1.1	0.6-1.8	16	0.8	0.4-1.2	12	0.4	0.2-0.7	3	0.3	0.1-0.8	9	0.4	0.2-0.8
2013	40	1.3	0.9-1.7	13	1.1	0.6-1.9	27	1.3	0.9-1.9	15	0.4	0.2-0.6	8	0.5	0.2-1.0	7	0.3	0.1-0.7
2014	35	1.0	0.7-1.3	16	1.0	0.5-1.6	19	0.9	0.6-1.5	7	0.2	0.1-0.4	2	0.1	0.0-0.4	5	0.3	0.1-0.6
2015	33	0.9	0.6-1.3	17	1.3	0.7-2.0	16	0.8	0.4-1.3	12	0.3	0.2-0.6	7	0.5	0.2-1.1	5	0.2	0.1-0.6
2016	32	0.9	0.6-1.2	15	1.0	0.5-1.6	17	0.7	0.4-1.2	5	0.1	0.0-0.3	2	0.1	0.0-0.5	3	0.1	0.0-0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Female breast

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	688	75.3	69.4-81.4	315	36.5	32.3-41.1
1977	713	75.7	70.0-81.8	295	32.2	28.4-36.3
1978	792	82.3	76.4-88.5	313	33.2	29.4-37.3
1979	803	81.3	75.5-87.5	294	30.3	26.7-34.1
1980	866	83.6	77.9-89.6	348	35.0	31.2-39.1
1981	855	81.9	76.2-87.8	357	35.6	31.8-39.7
1982	904	84.1	78.5-90.0	367	34.3	30.7-38.1
1983	931	83.6	78.2-89.4	376	34.3	30.8-38.0
1984	1042	90.6	85.1-96.5	413	38.2	34.5-42.2
1985	1056	90.1	84.5-95.8	342	29.6	26.5-33.0
1986	1057	86.6	81.3-92.1	392	32.6	29.4-36.1
1987	1137	90.7	85.4-96.2	396	32.2	29.0-35.6
1988	1226	95.5	90.1-101.1	427	33.5	30.3-36.9
1989	1152	88.5	83.4-93.9	425	33.6	30.4-37.0
1990	1229	92.6	87.4-98.0	430	33.3	30.1-36.6
1991	1374	101.7	96.3-107.3	434	32.8	29.7-36.1
1992	1326	96.7	91.5-102.1	444	32.8	29.8-36.0
1993	1352	96.7	91.5-102.0	423	31.4	28.5-34.6
1994	1431	99.7	94.6-105.1	496	35.0	32.0-38.3
1995	1458	99.6	94.5-104.9	511	35.9	32.8-39.1
1996	1604	107.3	102.0-112.7	489	33.5	30.5-36.6
1997	1643	107.9	102.7-113.3	490	33.3	30.4-36.4
1998	1798	115.7	110.4-121.2	491	32.4	29.6-35.4
1999	1650	103.6	98.6-108.7	473	30.5	27.8-33.4
2000	1626	99.4	94.5-104.3	463	29.2	26.6-32.0
2001	1620	97.2	92.5-102.1	516	31.7	29.0-34.6
2002	1724	101.2	96.4-106.1	467	28.2	25.7-30.9
2003	1780	103.0	98.3-107.9	468	27.7	25.2-30.3
2004	1834	104.3	99.6-109.3	476	27.8	25.3-30.4
2005	1851	104.3	99.5-109.2	518	29.5	27.0-32.2
2006	1962	108.5	103.7-113.5	473	26.8	24.4-29.3
2007	2004	109.9	105.1-114.8	469	25.9	23.6-28.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Female breast

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	2086	113.1	108.3-118.1	467	25.5	23.2-27.9
2009	2114	112.8	108.0-117.8	483	26.1	23.8-28.6
2010	2081	108.7	104.0-113.5	446	23.3	21.1-25.6
2011	2380	121.1	116.2-126.2	503	26.1	23.8-28.5
2012	2267	113.2	108.5-118.0	503	25.4	23.2-27.8
2013	2411	118.5	113.7-123.4	454	22.3	20.3-24.5
2014	2457	119.0	114.2-123.8	500	24.1	22.0-26.3
2015	2539	119.6	114.9-124.5	457	21.4	19.4-23.4
2016	2521	117.3	112.6-122.0	473	21.9	20.0-24.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Cervix uteri

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	293	29.9	26.4-33.8	130	15.0	12.4-18.1
1977	304	30.7	27.2-34.5	129	14.3	11.8-17.1
1978	290	28.4	25.1-32.1	110	11.7	9.5-14.3
1979	310	29.6	26.2-33.2	105	10.8	8.7-13.2
1980	278	26.6	23.4-30.0	115	11.6	9.5-14.1
1981	236	20.7	18.0-23.6	108	10.1	8.2-12.3
1982	264	23.4	20.5-26.5	97	8.5	6.9-10.5
1983	251	21.4	18.8-24.3	98	8.5	6.9-10.4
1984	260	21.6	19.0-24.5	77	7.2	5.6-9.1
1985	232	19.1	16.6-21.8	94	7.8	6.3-9.7
1986	269	21.9	19.3-24.8	91	7.5	6.0-9.2
1987	247	18.9	16.6-21.5	74	5.9	4.6-7.4
1988	233	17.7	15.4-20.1	108	8.5	6.9-10.2
1989	230	16.6	14.5-19.0	69	5.4	4.2-6.9
1990	299	20.8	18.5-23.3	83	6.4	5.0-7.9
1991	268	18.7	16.5-21.1	82	5.9	4.6-7.3
1992	256	17.2	15.1-19.5	78	5.6	4.4-7.0
1993	253	16.9	14.9-19.2	101	6.9	5.6-8.4
1994	263	17.7	15.6-20.1	79	5.4	4.3-6.8
1995	270	17.6	15.5-19.9	93	6.4	5.1-7.8
1996	292	18.5	16.4-20.8	78	5.1	4.0-6.4
1997	246	15.5	13.6-17.6	91	6.0	4.8-7.4
1998	257	15.4	13.5-17.4	79	5.1	4.0-6.3
1999	248	15.0	13.2-17.0	88	5.4	4.3-6.7
2000	289	17.4	15.4-19.5	91	5.4	4.4-6.7
2001	245	14.4	12.6-16.3	70	4.2	3.3-5.3
2002	244	14.1	12.3-16.0	74	4.4	3.5-5.6
2003	232	13.2	11.5-15.0	85	4.9	3.9-6.0
2004	216	12.0	10.5-13.8	64	3.7	2.8-4.7
2005	232	13.0	11.3-14.7	71	4.1	3.2-5.1
2006	220	12.2	10.6-13.9	72	4.1	3.2-5.2
2007	237	13.1	11.4-14.8	93	5.2	4.2-6.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Cervix uteri

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	217	11.7	10.2-13.4	77	4.1	3.3-5.2
2009	201	10.6	9.2-12.2	90	4.7	3.7-5.8
2010	212	11.3	9.8-13.0	60	3.1	2.4-4.0
2011	209	10.9	9.4-12.5	81	4.2	3.3-5.2
2012	203	10.5	9.0-12.0	67	3.4	2.6-4.3
2013	214	10.7	9.3-12.3	83	4.1	3.2-5.0
2014	194	9.6	8.3-11.1	77	3.7	2.9-4.7
2015	202	9.8	8.5-11.3	60	2.9	2.2-3.7
2016	201	9.6	8.3-11.1	58	2.8	2.1-3.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Corpus uterus and uterus, not otherwise specified

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	146	16.4	13.7-19.4	68	8.9	6.8-11.4
1977	143	16.0	13.4-19.0	66	7.4	5.6-9.5
1978	139	15.3	12.7-18.2	69	8.4	6.4-10.8
1979	143	14.3	11.9-17.0	66	7.3	5.6-9.4
1980	184	18.8	16.0-21.8	82	8.4	6.6-10.5
1981	141	13.7	11.4-16.3	72	6.9	5.4-8.8
1982	162	14.7	12.5-17.3	58	5.4	4.0-7.0
1983	163	14.7	12.4-17.2	74	7.1	5.5-8.9
1984	187	17.1	14.7-19.8	75	6.9	5.4-8.7
1985	213	18.3	15.8-20.9	87	8.2	6.5-10.1
1986	149	12.2	10.3-14.4	65	5.7	4.4-7.3
1987	209	17.1	14.8-19.7	55	4.9	3.7-6.4
1988	216	17.0	14.7-19.5	81	6.7	5.2-8.3
1989	182	13.9	11.9-16.1	76	5.9	4.6-7.4
1990	205	15.8	13.7-18.1	76	6.2	4.8-7.7
1991	250	18.8	16.5-21.4	77	6.0	4.7-7.5
1992	242	17.8	15.6-20.2	87	6.8	5.4-8.4
1993	265	19.0	16.7-21.4	85	6.6	5.2-8.1
1994	279	19.4	17.1-21.8	92	6.7	5.4-8.2
1995	257	17.8	15.7-20.1	77	5.6	4.4-7.0
1996	304	20.6	18.3-23.1	110	7.8	6.4-9.5
1997	323	21.8	19.5-24.3	109	7.5	6.2-9.1
1998	329	21.4	19.2-23.9	124	8.5	7.1-10.2
1999	353	22.7	20.4-25.3	114	7.6	6.2-9.1
2000	362	22.8	20.5-25.3	110	7.2	5.9-8.6
2001	384	23.3	21.1-25.8	124	7.9	6.6-9.4
2002	365	21.8	19.6-24.2	110	6.8	5.6-8.2
2003	394	23.0	20.7-25.4	140	8.4	7.0-9.9
2004	452	26.0	23.7-28.6	126	7.5	6.3-9.0
2005	454	25.9	23.6-28.4	144	8.4	7.1-10.0
2006	476	26.5	24.2-29.0	139	8.1	6.8-9.6
2007	465	25.6	23.3-28.1	136	7.9	6.6-9.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Corpus uterus and uterus, not otherwise specified

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	573	30.8	28.3-33.5	161	9.1	7.7-10.6
2009	565	29.8	27.4-32.4	150	8.2	6.9-9.6
2010	579	29.8	27.4-32.4	160	8.6	7.3-10.0
2011	600	30.0	27.7-32.6	181	9.5	8.1-11.0
2012	654	31.9	29.4-34.4	178	9.0	7.7-10.5
2013	641	30.3	27.9-32.7	170	8.2	7.0-9.6
2014	704	32.3	29.9-34.8	231	10.9	9.5-12.4
2015	722	32.5	30.1-35.0	206	9.3	8.1-10.7
2016	752	33.5	31.1-36.0	220	9.8	8.6-11.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Ovary

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	104	12.0	9.7-14.8	64	7.7	5.8-10.0
1977	120	12.1	9.9-14.6	73	7.9	6.1-10.0
1978	101	10.5	8.4-12.9	72	7.6	5.9-9.7
1979	129	13.2	10.9-15.9	69	6.9	5.3-8.8
1980	128	12.7	10.5-15.2	96	10.1	8.1-12.5
1981	119	10.9	9.0-13.1	62	5.7	4.3-7.4
1982	115	10.5	8.6-12.6	67	6.3	4.9-8.1
1983	151	13.1	11.0-15.4	86	8.0	6.3-9.9
1984	147	12.7	10.7-15.0	90	8.1	6.4-10.0
1985	147	11.8	9.9-13.9	90	7.7	6.2-9.6
1986	151	12.5	10.5-14.7	84	7.0	5.5-8.7
1987	120	9.9	8.2-11.9	82	6.8	5.4-8.5
1988	150	11.8	10.0-13.9	75	6.1	4.8-7.7
1989	133	9.9	8.3-11.8	85	6.6	5.3-8.2
1990	165	12.4	10.5-14.5	107	8.4	6.9-10.2
1991	138	10.4	8.7-12.3	92	7.0	5.6-8.7
1992	184	13.3	11.4-15.4	123	9.2	7.6-11.0
1993	157	10.8	9.2-12.7	114	8.6	7.0-10.3
1994	193	13.6	11.7-15.7	113	8.4	6.9-10.1
1995	179	12.1	10.4-14.0	122	8.8	7.3-10.6
1996	151	10.1	8.5-11.9	77	5.3	4.2-6.6
1997	199	13.0	11.3-15.0	101	7.0	5.7-8.5
1998	144	9.4	7.9-11.1	102	7.0	5.7-8.5
1999	175	11.3	9.7-13.1	100	6.6	5.4-8.1
2000	163	10.3	8.7-12.0	107	7.0	5.7-8.4
2001	170	10.3	8.8-12.0	108	6.8	5.6-8.2
2002	185	10.9	9.4-12.6	108	6.7	5.5-8.1
2003	158	9.2	7.8-10.7	110	6.6	5.4-7.9
2004	183	10.5	9.1-12.2	106	6.4	5.2-7.7
2005	174	9.9	8.5-11.5	114	6.7	5.5-8.0
2006	157	8.7	7.4-10.2	111	6.5	5.3-7.8
2007	163	9.0	7.7-10.5	107	6.1	5.0-7.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Ovary

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	182	9.9	8.5-11.5	103	5.6	4.6-6.8
2009	191	10.1	8.7-11.7	105	5.6	4.6-6.8
2010	181	9.5	8.2-11.0	114	6.1	5.0-7.3
2011	187	9.5	8.2-11.0	116	6.2	5.1-7.4
2012	218	10.9	9.4-12.4	131	6.6	5.5-7.8
2013	185	9.0	7.8-10.4	105	5.1	4.2-6.2
2014	216	10.3	9.0-11.8	121	5.7	4.7-6.9
2015	182	8.6	7.4-10.0	122	5.7	4.8-6.9
2016	188	8.4	7.2-9.8	122	5.5	4.5-6.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Prostate

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	519	120.0	108.3-132.5	209	55.0	46.6-64.2
1977	550	120.4	109.1-132.4	211	53.8	45.9-62.4
1978	607	132.2	120.4-144.7	226	54.7	46.9-63.3
1979	624	130.0	118.7-141.9	262	62.3	54.1-71.2
1980	758	152.6	140.7-165.2	274	65.1	56.8-74.1
1981	736	146.9	135.3-159.2	314	72.0	63.3-81.3
1982	699	134.6	123.8-146.0	276	60.7	53.1-69.0
1983	711	132.5	121.9-143.6	288	60.9	53.4-68.9
1984	773	137.7	127.1-148.8	286	58.5	51.3-66.3
1985	803	135.4	125.2-146.0	326	65.8	58.2-74.0
1986	845	142.4	132.1-153.3	346	68.3	60.7-76.6
1987	845	136.0	126.1-146.4	348	67.3	59.8-75.4
1988	910	148.7	138.2-159.6	310	60.1	53.1-67.7
1989	951	149.0	138.8-159.7	315	56.3	49.7-63.4
1990	1091	166.4	155.7-177.5	384	69.3	61.9-77.1
1991	1250	182.9	171.9-194.3	359	64.9	57.8-72.5
1992	1555	215.9	204.3-227.8	392	68.3	61.2-76.0
1993	1726	229.2	217.6-241.2	376	66.6	59.5-74.3
1994	1850	234.9	223.4-246.7	417	71.4	64.2-79.2
1995	2043	247.2	235.8-259.0	435	69.9	63.0-77.3
1996	2077	242.0	231.0-253.4	409	63.5	57.1-70.4
1997	2193	251.2	240.1-262.7	405	61.6	55.3-68.4
1998	2274	251.1	240.2-262.3	402	60.8	54.6-67.3
1999	2581	274.0	262.9-285.4	413	60.3	54.3-66.8
2000	2865	295.5	284.2-307.2	425	60.3	54.4-66.6
2001	2738	273.6	262.8-284.6	425	59.5	53.7-65.8
2002	2794	272.3	261.6-283.2	437	57.4	51.8-63.3
2003	2785	263.4	253.1-273.9	378	48.7	43.7-54.1
2004	2808	255.7	245.8-265.8	374	48.1	43.1-53.5
2005	2832	250.4	240.7-260.2	370	44.6	39.9-49.6
2006	3021	262.7	252.9-272.7	346	42.4	37.8-47.2
2007	3273	276.3	266.4-286.4	381	45.5	40.9-50.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Prostate

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	3189	258.0	248.6-267.5	365	42.5	38.0-47.2
2009	3188	248.8	239.8-258.0	393	44.7	40.2-49.5
2010	3155	237.7	229.1-246.6	381	41.4	37.1-45.9
2011	3434	247.5	238.9-256.3	369	38.4	34.4-42.6
2012	3110	216.4	208.5-224.5	368	37.3	33.4-41.4
2013	3149	212.3	204.6-220.2	366	35.8	32.2-39.8
2014	2901	188.9	181.8-196.2	385	34.2	30.7-37.9
2015	2945	184.7	177.8-191.7	336	29.2	26.0-32.6
2016	2988	184.0	177.2-191.0	366	30.2	27.1-33.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Testis

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	8	0.6	0.2-1.4	4	1.0	0.1-2.8
1977	9	1.0	0.4-2.2	3	0.3	0.1-1.1
1978	4	0.5	0.1-1.5	4	0.5	0.1-1.5
1979	6	0.5	0.2-1.3	0	0.0	0.0-0.7
1980	15	1.4	0.8-2.5	3	0.3	0.1-1.0
1981	10	1.0	0.4-1.9	2	0.2	0.0-1.0
1982	7	0.7	0.3-1.5	0	0.0	0.0-0.6
1983	7	0.7	0.2-1.6	7	1.0	0.3-2.3
1984	8	0.6	0.2-1.3	1	0.2	0.0-0.9
1985	17	1.7	0.9-3.0	2	0.1	0.0-0.7
1986	17	1.3	0.7-2.2	2	0.2	0.0-0.7
1987	13	0.8	0.4-1.5	2	0.1	0.0-0.6
1988	13	0.9	0.5-1.6	2	0.4	0.0-1.5
1989	15	1.3	0.7-2.2	1	0.1	0.0-0.6
1990	13	1.1	0.5-1.9	3	0.2	0.0-0.7
1991	10	0.8	0.3-1.7	1	0.1	0.0-0.6
1992	22	1.5	0.8-2.6	2	0.1	0.0-0.6
1993	13	0.9	0.5-1.7	3	0.2	0.0-0.7
1994	18	1.2	0.7-2.0	1	0.1	0.0-0.6
1995	7	0.4	0.2-1.0	0	0.0	0.0-0.4
1996	21	1.3	0.8-2.1	0	0.0	0.0-0.4
1997	13	0.8	0.4-1.4	1	0.1	0.0-0.5
1998	21	1.3	0.8-2.1	1	0.1	0.0-0.5
1999	21	1.3	0.8-2.1	1	0.1	0.0-0.5
2000	29	2.1	1.3-3.0	3	0.3	0.0-0.8
2001	23	1.5	0.9-2.2	1	0.1	0.0-0.6
2002	19	1.2	0.7-1.9	2	0.1	0.0-0.5
2003	27	1.7	1.1-2.5	1	0.1	0.0-0.4
2004	16	1.0	0.5-1.6	1	0.1	0.0-0.4
2005	23	1.5	0.9-2.3	2	0.1	0.0-0.5
2006	30	1.9	1.3-2.7	1	0.1	0.0-0.4
2007	30	1.9	1.3-2.8	1	0.1	0.0-0.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Testis

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	17	1.1	0.6-1.7	1	0.0	0.0-0.4
2009	18	1.1	0.6-1.7	1	0.1	0.0-0.4
2010	24	1.5	0.9-2.2	2	0.1	0.0-0.5
2011	27	1.6	1.1-2.4	3	0.2	0.0-0.6
2012	24	1.5	0.9-2.2	1	0.1	0.0-0.4
2013	27	1.5	1.0-2.2	1	0.1	0.0-0.4
2014	35	2.1	1.5-3.0	1	0.1	0.0-0.3
2015	33	1.9	1.3-2.7	1	0.1	0.0-0.4
2016	35	2.0	1.4-2.8	2	0.2	0.0-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Urinary bladder (incl. in situ)

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	161	13.8	11.5-16.4	107	24.1	19.0-30.0	54	7.7	5.7-10.3	69	5.9	4.5-7.7	39	7.6	5.1-10.9	30	4.7	3.1-6.9
1977	131	10.5	8.6-12.6	78	16.2	12.4-20.7	53	6.9	5.0-9.2	60	5.5	4.0-7.1	35	9.3	6.0-13.4	25	3.5	2.2-5.2
1978	141	10.5	8.7-12.6	94	19.3	15.1-24.1	47	5.5	3.9-7.4	62	4.8	3.6-6.3	35	7.5	4.9-10.7	27	3.3	2.1-4.9
1979	167	12.6	10.6-14.8	97	19.0	14.9-23.6	70	8.8	6.7-11.2	76	6.6	5.1-8.4	40	9.8	6.7-13.6	36	4.9	3.4-6.9
1980	181	12.6	10.7-14.7	120	21.4	17.4-26.0	61	7.2	5.4-9.4	66	5.2	4.0-6.8	37	8.0	5.3-11.3	29	3.7	2.4-5.4
1981	158	10.7	8.9-12.6	90	15.8	12.3-19.9	68	7.6	5.8-9.8	78	5.5	4.2-6.9	46	8.8	6.2-12.2	32	3.6	2.4-5.1
1982	161	10.6	8.9-12.5	89	16.0	12.5-20.0	72	7.5	5.8-9.6	76	5.2	4.1-6.6	37	6.9	4.6-9.7	39	4.3	3.0-5.9
1983	162	9.9	8.3-11.6	87	13.9	10.8-17.5	75	7.5	5.9-9.5	67	4.5	3.5-5.8	31	5.7	3.7-8.3	36	3.9	2.7-5.4
1984	167	10.0	8.4-11.7	109	17.7	14.2-21.7	58	5.6	4.2-7.3	73	4.8	3.7-6.1	45	7.8	5.5-10.7	28	3.1	2.0-4.4
1985	191	11.6	9.9-13.5	108	17.4	13.9-21.4	83	8.4	6.6-10.4	75	4.9	3.8-6.2	43	8.3	5.9-11.3	32	3.2	2.2-4.6
1986	169	10.1	8.6-11.9	110	19.0	15.3-23.3	59	5.5	4.1-7.1	64	4.1	3.1-5.3	33	6.4	4.2-9.1	31	3.0	2.0-4.3
1987	162	9.5	8.0-11.1	104	16.2	12.9-19.9	58	5.7	4.3-7.4	72	4.5	3.5-5.7	40	7.2	5.0-10.0	32	3.1	2.1-4.4
1988	187	10.7	9.2-12.4	113	16.7	13.5-20.3	74	7.1	5.6-9.0	68	4.1	3.2-5.3	40	7.5	5.1-10.4	28	2.6	1.7-3.7
1989	177	9.6	8.2-11.2	106	15.1	12.1-18.6	71	6.4	4.9-8.1	71	4.5	3.5-5.7	42	7.7	5.4-10.6	29	2.9	1.9-4.1
1990	177	9.8	8.4-11.4	115	18.0	14.6-21.9	62	5.5	4.2-7.1	74	4.3	3.3-5.4	41	6.3	4.4-8.7	33	3.1	2.1-4.4
1991	202	10.3	8.9-11.9	129	18.7	15.4-22.6	73	5.9	4.6-7.4	64	3.6	2.8-4.7	40	6.8	4.7-9.4	24	2.1	1.3-3.1
1992	192	9.6	8.3-11.1	112	15.7	12.7-19.2	80	6.3	5.0-7.9	57	3.1	2.3-4.1	30	4.9	3.2-7.1	27	2.3	1.5-3.3
1993	196	9.8	8.4-11.3	100	13.9	11.1-17.1	96	7.6	6.1-9.3	79	4.0	3.2-5.0	44	6.6	4.6-8.9	35	2.8	1.9-3.8
1994	196	9.6	8.3-11.1	111	15.5	12.5-18.9	85	6.5	5.2-8.1	72	3.9	3.0-4.9	36	5.5	3.8-7.8	36	3.0	2.1-4.1
1995	177	8.5	7.2-9.8	118	15.8	12.9-19.2	59	4.5	3.4-5.8	67	3.5	2.7-4.4	33	5.3	3.4-7.6	34	2.7	1.9-3.8
1996	209	9.7	8.4-11.2	118	15.2	12.3-18.4	91	6.8	5.5-8.3	74	3.7	2.9-4.7	41	6.1	4.2-8.3	33	2.6	1.8-3.6
1997	213	9.9	8.6-11.4	124	16.5	13.5-20.0	89	6.6	5.3-8.2	72	3.5	2.8-4.5	35	5.0	3.3-7.0	37	2.8	2.0-3.9
1998	258	11.6	10.2-13.2	157	19.4	16.3-23.0	101	7.4	6.0-8.9	83	4.0	3.2-5.0	39	5.6	3.9-7.7	44	3.3	2.4-4.4
1999	232	9.8	8.6-11.2	149	17.2	14.3-20.4	83	5.7	4.5-7.0	69	3.1	2.4-3.9	35	4.4	3.0-6.2	34	2.4	1.7-3.3
2000	244	10.2	8.9-11.6	147	16.6	13.8-19.7	97	6.5	5.3-8.0	81	3.6	2.8-4.4	43	5.4	3.8-7.4	38	2.6	1.9-3.6
2001	251	10.2	9.0-11.6	152	16.8	14.1-19.9	99	6.5	5.3-7.9	83	3.6	2.8-4.4	37	4.7	3.2-6.6	46	3.1	2.2-4.1
2002	239	9.7	8.5-11.1	141	15.7	13.1-18.7	98	6.3	5.1-7.7	66	2.8	2.2-3.6	28	3.3	2.1-4.8	38	2.5	1.8-3.4
2003	256	9.7	8.5-11.0	177	18.0	15.3-21.0	79	4.9	3.9-6.1	71	2.9	2.2-3.6	39	4.5	3.1-6.2	32	2.0	1.4-2.9
2004	278	10.7	9.5-12.1	178	18.3	15.5-21.4	100	6.3	5.1-7.6	78	3.1	2.4-3.8	40	4.2	2.9-5.8	38	2.4	1.7-3.3
2005	301	11.2	9.9-12.5	190	18.4	15.7-21.4	111	6.8	5.6-8.2	80	3.4	2.7-4.2	45	5.5	3.9-7.4	35	2.3	1.6-3.2
2006	300	11.1	9.8-12.4	199	19.9	17.1-23.1	101	5.9	4.8-7.2	61	2.4	1.8-3.1	39	4.3	3.0-6.0	22	1.4	0.9-2.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Urinary bladder (incl. in situ)

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	261	9.5	8.4-10.7	161	15.9	13.4-18.7	100	5.9	4.8-7.2	78	3.1	2.4-3.8	37	4.2	2.9-5.8	41	2.5	1.8-3.3
2008	299	10.4	9.2-11.7	199	18.2	15.6-21.0	100	5.7	4.6-7.0	67	2.5	1.9-3.2	37	4.2	2.9-5.8	30	1.7	1.1-2.4
2009	317	11.1	9.8-12.4	204	18.9	16.2-21.8	113	6.5	5.3-7.8	73	2.7	2.1-3.5	42	4.7	3.3-6.3	31	1.8	1.2-2.5
2010	302	10.0	8.9-11.3	205	18.1	15.6-20.9	97	5.2	4.2-6.4	88	3.1	2.5-3.9	43	4.6	3.2-6.2	45	2.5	1.8-3.4
2011	358	11.4	10.3-12.7	253	21.5	18.8-24.5	105	5.5	4.5-6.6	88	3.0	2.4-3.7	51	4.9	3.6-6.5	37	2.0	1.4-2.8
2012	331	10.6	9.5-11.8	209	18.1	15.6-20.9	122	6.2	5.2-7.5	88	2.9	2.3-3.5	53	4.6	3.4-6.0	35	1.8	1.3-2.6
2013	383	11.9	10.7-13.1	251	20.2	17.7-23.0	132	6.7	5.6-8.0	88	2.8	2.2-3.4	52	4.6	3.4-6.1	36	1.8	1.2-2.5
2014	358	10.5	9.4-11.7	241	18.2	15.9-20.8	117	5.6	4.7-6.8	91	2.9	2.3-3.6	56	5.1	3.8-6.7	35	1.7	1.2-2.4
2015	378	10.9	9.8-12.1	249	18.6	16.3-21.2	129	6.1	5.1-7.3	84	2.5	1.9-3.0	45	3.6	2.6-4.8	39	1.8	1.3-2.4
2016	351	9.8	8.8-10.9	234	16.9	14.7-19.3	117	5.3	4.4-6.4	90	2.6	2.1-3.2	52	4.3	3.1-5.6	38	1.7	1.2-2.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Kidney and renal pelvis

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	96	6.7	5.3-8.4	58	8.6	6.2-11.5	38	5.3	3.6-7.3	47	3.2	2.2-4.4	28	4.4	2.7-6.7	19	2.4	1.3-3.8
1977	74	4.5	3.4-5.7	47	7.7	5.3-10.8	27	2.5	1.7-3.7	29	1.8	1.1-2.6	20	3.0	1.7-4.8	9	0.9	0.4-1.9
1978	111	6.9	5.6-8.4	72	10.4	7.8-13.4	39	4.4	3.1-6.2	54	3.3	2.4-4.4	36	4.9	3.2-7.2	18	2.1	1.2-3.4
1979	88	5.0	4.0-6.2	65	9.2	6.9-11.9	23	2.2	1.4-3.4	47	2.9	2.1-3.9	37	5.7	3.9-8.1	10	1.0	0.5-1.9
1980	89	5.0	3.9-6.2	47	6.7	4.7-9.2	42	4.0	2.8-5.4	36	2.0	1.4-2.8	28	3.9	2.5-5.9	8	0.8	0.3-1.5
1981	93	5.0	4.0-6.2	65	8.4	6.3-10.9	28	2.7	1.8-3.9	37	1.8	1.2-2.5	25	3.0	1.9-4.5	12	0.9	0.5-1.7
1982	105	5.7	4.6-7.0	65	8.6	6.5-11.2	40	3.8	2.6-5.2	46	2.7	2.0-3.7	25	3.4	2.0-5.1	21	2.2	1.3-3.5
1983	114	5.9	4.9-7.2	62	8.2	6.1-10.8	52	4.6	3.4-6.0	49	2.7	1.9-3.6	30	4.2	2.7-6.2	19	1.7	1.0-2.8
1984	141	7.2	6.0-8.5	90	11.5	9.0-14.5	51	4.5	3.3-5.9	52	2.8	2.1-3.8	30	4.3	2.8-6.3	22	2.0	1.2-3.0
1985	126	6.2	5.1-7.5	70	8.6	6.5-11.1	56	4.7	3.5-6.2	58	3.0	2.3-3.9	33	3.9	2.6-5.7	25	2.3	1.5-3.4
1986	139	6.7	5.6-7.9	79	9.2	7.2-11.6	60	4.9	3.7-6.4	73	3.8	3.0-4.9	43	5.8	4.0-8.0	30	2.6	1.7-3.7
1987	147	7.3	6.1-8.6	90	12.1	9.5-15.1	57	4.6	3.4-6.0	50	2.6	1.9-3.5	25	3.3	2.1-5.0	25	2.2	1.4-3.2
1988	140	6.7	5.6-7.9	75	8.8	6.8-11.2	65	5.2	4.0-6.7	56	2.8	2.1-3.7	34	4.6	3.1-6.6	22	1.8	1.1-2.7
1989	134	6.1	5.1-7.3	77	9.6	7.3-12.3	57	4.2	3.2-5.4	46	2.4	1.8-3.3	20	2.9	1.7-4.6	26	2.2	1.4-3.2
1990	168	7.6	6.4-8.9	96	10.9	8.6-13.5	72	5.5	4.3-7.0	54	2.6	1.9-3.4	29	3.9	2.5-5.8	25	1.9	1.2-2.9
1991	187	8.1	7.0-9.4	119	13.0	10.5-15.7	68	5.0	3.9-6.4	60	2.8	2.1-3.7	35	4.5	3.0-6.4	25	1.9	1.2-2.8
1992	174	7.7	6.5-8.9	113	12.6	10.2-15.4	61	4.6	3.5-5.9	53	2.4	1.8-3.2	32	3.9	2.5-5.6	21	1.6	1.0-2.5
1993	185	7.7	6.6-8.9	121	12.7	10.3-15.4	64	4.5	3.4-5.7	67	3.1	2.4-3.9	39	4.5	3.1-6.3	28	2.2	1.4-3.1
1994	191	7.7	6.6-8.9	96	9.6	7.6-11.8	95	6.5	5.2-7.9	51	2.2	1.6-2.9	30	3.4	2.2-4.9	21	1.5	0.9-2.3
1995	235	9.8	8.6-11.2	134	14.2	11.7-17.1	101	7.1	5.8-8.7	76	3.4	2.7-4.3	46	5.5	3.9-7.5	30	2.2	1.5-3.1
1996	218	8.6	7.5-9.9	142	14.1	11.7-16.8	76	5.1	4.0-6.4	63	2.6	2.0-3.4	36	3.7	2.5-5.3	27	1.9	1.2-2.8
1997	231	9.3	8.1-10.6	131	13.5	11.1-16.1	100	6.7	5.4-8.1	71	3.0	2.3-3.8	42	4.8	3.4-6.7	29	2.0	1.3-2.9
1998	254	9.8	8.6-11.1	154	14.5	12.2-17.2	100	6.6	5.3-8.0	57	2.3	1.7-3.0	34	3.3	2.2-4.7	23	1.6	1.0-2.4
1999	269	10.3	9.1-11.7	137	13.1	10.9-15.6	132	8.5	7.1-10.1	54	2.3	1.7-3.0	26	2.6	1.7-4.0	28	1.9	1.3-2.8
2000	240	9.0	7.9-10.3	125	11.5	9.5-13.8	115	7.3	6.0-8.7	62	2.4	1.8-3.1	34	3.4	2.3-4.8	28	1.8	1.2-2.6
2001	279	10.0	8.8-11.2	158	13.9	11.7-16.4	121	7.3	6.1-8.8	69	2.7	2.1-3.4	37	3.8	2.6-5.3	32	2.0	1.4-2.9
2002	291	10.2	9.1-11.5	167	14.8	12.5-17.3	124	7.4	6.1-8.8	54	2.0	1.5-2.6	37	3.6	2.4-5.0	17	1.1	0.6-1.7
2003	324	11.3	10.0-12.6	189	16.0	13.7-18.5	135	8.0	6.7-9.4	70	2.7	2.1-3.4	44	4.4	3.1-6.0	26	1.6	1.1-2.4
2004	342	11.4	10.2-12.7	198	15.7	13.5-18.2	144	8.3	7.0-9.8	73	2.7	2.1-3.4	43	4.0	2.8-5.5	30	1.8	1.2-2.6
2005	358	11.9	10.7-13.2	189	14.9	12.8-17.3	169	9.7	8.3-11.2	67	2.3	1.8-2.9	44	3.7	2.6-5.1	23	1.3	0.9-2.0
2006	391	12.9	11.6-14.3	241	19.4	16.9-22.2	150	8.5	7.2-10.0	78	2.7	2.1-3.4	42	3.8	2.7-5.2	36	2.1	1.5-2.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Kidney and renal pelvis

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	423	13.6	12.3-15.0	267	21.1	18.5-23.9	156	8.7	7.4-10.2	85	3.0	2.4-3.7	57	5.2	3.9-6.8	28	1.6	1.1-2.3
2008	421	13.5	12.2-14.9	228	17.6	15.3-20.2	193	10.7	9.2-12.3	75	2.4	1.9-3.1	45	3.6	2.6-5.0	30	1.7	1.1-2.4
2009	467	14.4	13.1-15.8	274	20.0	17.6-22.6	193	10.2	8.8-11.8	71	2.3	1.8-2.9	46	3.8	2.7-5.2	25	1.3	0.9-2.0
2010	503	15.4	14.0-16.8	318	23.4	20.8-26.3	185	9.9	8.5-11.4	76	2.6	2.0-3.2	42	3.8	2.7-5.2	34	1.9	1.3-2.6
2011	481	14.5	13.2-15.9	298	21.8	19.3-24.6	183	9.5	8.2-11.0	74	2.3	1.8-2.9	45	3.8	2.7-5.1	29	1.5	1.0-2.2
2012	484	14.1	12.9-15.5	279	19.5	17.2-22.0	205	10.4	9.0-12.0	76	2.3	1.8-2.9	43	3.6	2.5-4.9	33	1.6	1.1-2.2
2013	545	15.2	14.0-16.6	333	21.9	19.5-24.5	212	10.3	9.0-11.9	61	1.8	1.4-2.3	39	2.7	1.9-3.8	22	1.1	0.7-1.6
2014	563	15.3	14.0-16.7	366	23.2	20.8-25.8	197	9.5	8.2-10.9	74	2.2	1.7-2.7	50	3.8	2.8-5.1	24	1.2	0.7-1.7
2015	582	15.5	14.2-16.8	360	22.8	20.4-25.4	222	10.2	8.9-11.7	99	2.8	2.3-3.4	68	4.9	3.8-6.3	31	1.4	1.0-2.0
2016	589	15.5	14.3-16.9	365	23.1	20.7-25.8	224	10.3	9.0-11.8	75	2.1	1.6-2.6	51	3.4	2.5-4.6	24	1.1	0.7-1.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Brain and other central nervous system

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	66	3.3	2.6-4.3	35	4.1	2.8-5.8	31	2.8	1.9-4.0	52	2.8	2.0-3.8	29	3.5	2.3-5.3	23	2.3	1.4-3.5
1977	84	3.9	3.1-4.9	39	4.5	3.1-6.4	45	3.6	2.6-4.8	42	2.2	1.6-3.1	23	2.9	1.8-4.5	19	1.8	1.1-2.9
1978	52	2.7	2.0-3.6	24	3.1	1.9-4.8	28	2.5	1.6-3.7	40	2.1	1.4-2.8	20	2.2	1.3-3.6	20	1.9	1.1-3.0
1979	76	3.5	2.7-4.4	37	4.1	2.8-5.8	39	3.2	2.2-4.4	51	2.6	1.9-3.5	25	3.2	2.0-4.9	26	2.3	1.5-3.3
1980	76	3.9	3.0-5.0	47	5.6	3.9-7.7	29	2.7	1.8-3.9	47	2.5	1.8-3.3	30	3.8	2.4-5.6	17	1.6	0.9-2.5
1981	54	2.4	1.8-3.2	29	2.7	1.7-4.0	25	2.1	1.3-3.2	35	2.0	1.4-2.9	16	2.0	1.1-3.3	19	2.0	1.1-3.2
1982	78	3.3	2.6-4.2	45	4.3	3.1-6.0	33	2.5	1.7-3.6	65	3.3	2.5-4.3	41	5.1	3.6-7.1	24	2.1	1.3-3.1
1983	75	3.3	2.6-4.2	41	4.1	2.9-5.8	34	2.7	1.8-3.8	61	3.0	2.2-3.8	23	2.6	1.6-4.1	38	3.1	2.2-4.4
1984	99	4.1	3.3-5.1	54	5.1	3.7-6.8	45	3.3	2.4-4.5	70	3.1	2.4-4.0	43	4.5	3.1-6.2	27	2.1	1.4-3.1
1985	100	4.4	3.5-5.4	43	5.0	3.5-6.9	57	4.2	3.1-5.5	56	2.5	1.8-3.2	32	3.6	2.4-5.2	24	1.7	1.1-2.6
1986	94	4.1	3.2-5.0	55	5.4	3.9-7.3	39	3.0	2.1-4.2	63	2.8	2.1-3.7	33	3.4	2.3-4.9	30	2.4	1.6-3.4
1987	101	4.1	3.3-5.1	48	4.6	3.2-6.4	53	3.7	2.8-4.9	55	2.7	2.0-3.5	28	3.5	2.2-5.3	27	2.2	1.4-3.2
1988	88	3.4	2.7-4.3	51	4.7	3.3-6.4	37	2.5	1.8-3.6	48	2.0	1.4-2.6	23	2.4	1.4-3.8	25	1.7	1.1-2.5
1989	92	3.5	2.8-4.3	41	3.6	2.5-5.1	51	3.5	2.6-4.6	40	1.6	1.2-2.2	19	1.8	1.0-3.0	21	1.5	0.9-2.3
1990	102	4.0	3.2-4.9	46	4.4	3.1-6.0	56	3.8	2.9-5.0	46	2.0	1.4-2.7	24	2.8	1.7-4.3	22	1.6	1.0-2.4
1991	129	5.0	4.1-6.0	63	6.0	4.4-7.9	66	4.4	3.4-5.6	53	2.2	1.6-3.0	22	2.2	1.3-3.4	31	2.3	1.5-3.2
1992	130	5.0	4.2-6.0	66	6.1	4.6-8.0	64	4.3	3.3-5.5	55	2.4	1.8-3.2	25	2.9	1.7-4.4	30	2.2	1.5-3.2
1993	93	3.4	2.7-4.2	54	4.6	3.3-6.2	39	2.6	1.8-3.5	62	2.3	1.8-3.0	29	2.8	1.8-4.1	33	2.1	1.5-3.0
1994	108	3.8	3.1-4.6	58	4.7	3.5-6.2	50	3.2	2.3-4.2	59	2.2	1.7-2.9	28	2.6	1.7-3.9	31	2.0	1.4-2.8
1995	114	4.1	3.4-5.0	48	3.8	2.7-5.2	66	4.3	3.3-5.5	50	1.8	1.3-2.4	23	1.9	1.2-3.0	27	1.8	1.1-2.6
1996	137	5.0	4.1-5.9	64	5.5	4.2-7.1	73	4.6	3.6-5.8	51	1.9	1.4-2.5	24	2.3	1.4-3.5	27	1.7	1.1-2.5
1997	110	3.7	3.1-4.5	59	4.8	3.5-6.3	51	3.1	2.3-4.0	49	1.8	1.3-2.4	24	1.9	1.2-2.9	25	1.6	1.0-2.4
1998	118	3.9	3.2-4.7	59	4.1	3.1-5.4	59	3.6	2.7-4.6	45	1.7	1.2-2.2	25	2.2	1.4-3.3	20	1.3	0.8-2.0
1999	125	4.3	3.5-5.1	66	5.4	4.0-7.0	59	3.6	2.7-4.6	42	1.5	1.0-2.0	26	2.2	1.4-3.3	16	1.0	0.5-1.6
2000	122	4.1	3.4-4.9	59	4.7	3.5-6.1	63	3.7	2.8-4.7	54	2.0	1.5-2.6	26	2.2	1.4-3.2	28	1.8	1.2-2.6
2001	146	4.9	4.1-5.8	61	4.4	3.3-5.7	85	5.0	4.0-6.2	63	2.2	1.7-2.9	28	2.5	1.6-3.7	35	2.1	1.4-2.9
2002	127	4.1	3.4-4.9	65	5.0	3.8-6.5	62	3.5	2.7-4.5	62	2.1	1.6-2.6	26	1.9	1.2-2.8	36	2.1	1.5-2.9
2003	147	4.8	4.0-5.7	73	5.3	4.1-6.7	74	4.3	3.4-5.4	55	1.8	1.4-2.4	33	2.6	1.7-3.7	22	1.3	0.8-1.9
2004	132	4.2	3.5-5.0	69	5.1	3.9-6.5	63	3.6	2.7-4.6	62	2.0	1.6-2.6	31	2.4	1.6-3.5	31	1.8	1.2-2.5
2005	130	4.2	3.5-5.0	67	5.4	4.1-7.0	63	3.5	2.7-4.5	49	1.6	1.2-2.1	32	2.5	1.7-3.6	17	0.9	0.5-1.5
2006	114	3.6	2.9-4.3	56	4.0	2.9-5.2	58	3.2	2.4-4.2	59	1.9	1.4-2.5	33	2.6	1.8-3.8	26	1.4	0.9-2.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Brain and other central nervous system

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	129	3.9	3.3-4.7	66	4.5	3.4-5.8	63	3.5	2.6-4.4	49	1.6	1.2-2.1	21	1.8	1.1-2.7	28	1.5	1.0-2.2
2008	124	3.8	3.2-4.6	58	4.1	3.1-5.4	66	3.6	2.8-4.6	63	2.0	1.5-2.6	31	2.4	1.6-3.4	32	1.7	1.2-2.5
2009	129	4.0	3.3-4.8	66	4.7	3.6-6.0	63	3.4	2.6-4.4	56	1.8	1.4-2.4	30	2.4	1.6-3.5	26	1.4	0.9-2.1
2010	142	4.2	3.5-5.0	76	5.1	4.0-6.5	66	3.5	2.7-4.5	54	1.7	1.2-2.2	28	2.0	1.3-3.0	26	1.4	0.9-2.1
2011	147	4.3	3.6-5.0	75	5.1	3.9-6.4	72	3.6	2.8-4.6	72	2.2	1.7-2.7	37	2.6	1.8-3.6	35	1.8	1.3-2.6
2012	140	4.1	3.4-4.8	82	5.4	4.2-6.7	58	3.0	2.3-3.9	65	1.9	1.4-2.4	33	2.2	1.5-3.2	32	1.6	1.1-2.3
2013	159	4.4	3.7-5.1	77	4.6	3.6-5.8	82	4.2	3.3-5.2	73	2.1	1.6-2.6	38	2.6	1.8-3.6	35	1.7	1.2-2.4
2014	120	3.3	2.7-4.0	72	4.4	3.4-5.6	48	2.4	1.8-3.2	66	1.9	1.4-2.4	38	2.7	1.9-3.8	28	1.3	0.9-1.9
2015	136	3.8	3.2-4.5	75	4.7	3.6-5.9	61	3.2	2.4-4.1	65	1.8	1.4-2.3	32	2.2	1.5-3.1	33	1.6	1.1-2.3
2016	144	3.9	3.3-4.7	89	5.5	4.4-6.8	55	2.7	2.1-3.6	81	2.2	1.7-2.7	43	2.6	1.8-3.5	38	1.8	1.3-2.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Thyroid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	45	2.4	1.7-3.3	12	1.7	0.8-3.1	33	3.1	2.1-4.3	10	0.6	0.3-1.2	3	0.4	0.1-1.3	7	0.8	0.3-1.6
1977	45	2.3	1.6-3.3	3	0.4	0.1-1.2	42	3.7	2.6-5.2	6	0.5	0.1-1.0	3	0.4	0.1-1.3	3	0.5	0.1-1.3
1978	37	2.1	1.4-2.9	11	1.6	0.7-3.1	26	2.5	1.6-3.7	11	0.8	0.4-1.4	2	0.4	0.0-1.4	9	1.1	0.5-2.0
1979	48	2.9	2.1-4.0	11	1.6	0.7-3.2	37	3.8	2.6-5.4	9	0.6	0.2-1.2	2	0.2	0.0-1.0	7	0.8	0.3-1.8
1980	44	2.1	1.5-2.9	5	0.4	0.1-1.2	39	3.2	2.2-4.5	8	0.5	0.2-1.0	1	0.2	0.0-1.0	7	0.7	0.3-1.5
1981	34	1.9	1.3-2.7	6	0.6	0.2-1.5	28	2.7	1.7-3.9	7	0.5	0.2-1.0	1	0.1	0.0-0.8	6	0.7	0.2-1.5
1982	43	2.2	1.6-3.0	7	0.8	0.3-1.8	36	3.2	2.2-4.5	11	0.7	0.3-1.2	4	0.8	0.2-2.2	7	0.6	0.2-1.3
1983	45	2.0	1.5-2.8	6	0.6	0.2-1.4	39	3.1	2.2-4.3	6	0.4	0.1-0.9	1	0.2	0.0-1.1	5	0.5	0.2-1.2
1984	54	2.5	1.8-3.3	8	0.8	0.3-1.6	46	3.6	2.6-4.9	9	0.5	0.2-1.0	2	0.5	0.0-1.7	7	0.6	0.2-1.3
1985	68	3.1	2.4-4.0	19	2.0	1.2-3.3	49	3.9	2.8-5.1	8	0.4	0.2-0.8	4	0.5	0.1-1.3	4	0.4	0.1-0.9
1986	45	2.2	1.6-3.0	13	1.8	0.9-3.2	32	2.5	1.7-3.6	11	0.6	0.3-1.0	4	0.7	0.1-1.8	7	0.5	0.2-1.1
1987	65	2.9	2.2-3.7	13	1.5	0.8-2.7	52	3.8	2.8-5.1	12	0.7	0.4-1.2	3	0.6	0.1-1.6	9	0.8	0.4-1.5
1988	49	2.2	1.6-2.9	11	1.3	0.6-2.4	38	2.8	2.0-3.9	8	0.5	0.2-0.9	3	0.4	0.1-1.1	5	0.5	0.1-1.1
1989	48	2.1	1.5-2.8	5	0.5	0.2-1.4	43	3.1	2.2-4.3	11	0.7	0.4-1.3	1	0.2	0.0-0.8	10	1.0	0.5-1.8
1990	57	2.3	1.7-3.0	9	0.8	0.4-1.7	48	3.3	2.4-4.4	5	0.3	0.1-0.7	1	0.1	0.0-0.8	4	0.4	0.1-0.9
1991	67	2.8	2.1-3.6	8	1.1	0.4-2.2	59	4.0	3.0-5.2	10	0.6	0.3-1.0	2	0.4	0.1-1.4	8	0.6	0.3-1.3
1992	90	3.2	2.6-4.0	22	1.8	1.1-2.9	68	4.3	3.3-5.5	11	0.5	0.2-0.9	3	0.3	0.1-0.8	8	0.6	0.3-1.2
1993	79	3.0	2.4-3.8	17	1.5	0.8-2.4	62	4.1	3.1-5.3	8	0.4	0.2-0.8	3	0.3	0.1-0.9	5	0.4	0.1-1.0
1994	90	3.4	2.7-4.2	13	1.2	0.6-2.2	77	5.0	3.9-6.3	8	0.3	0.1-0.7	2	0.2	0.0-0.8	6	0.4	0.1-0.9
1995	98	3.6	2.9-4.4	15	1.4	0.8-2.4	83	5.2	4.1-6.4	8	0.4	0.2-0.8	1	0.1	0.0-0.5	7	0.6	0.2-1.1
1996	119	4.1	3.4-4.9	23	2.2	1.3-3.5	96	5.7	4.6-7.0	15	0.7	0.4-1.1	9	1.0	0.4-2.0	6	0.4	0.2-0.9
1997	123	4.2	3.5-5.1	26	2.1	1.3-3.1	97	5.8	4.7-7.1	14	0.5	0.3-0.9	8	0.8	0.3-1.7	6	0.4	0.1-0.8
1998	134	4.6	3.8-5.4	25	2.2	1.4-3.3	109	6.4	5.2-7.7	10	0.4	0.2-0.8	4	0.4	0.1-1.0	6	0.4	0.2-0.9
1999	127	4.4	3.6-5.2	17	1.3	0.8-2.2	110	6.5	5.4-7.9	15	0.6	0.3-1.0	2	0.2	0.0-0.6	13	0.9	0.5-1.5
2000	164	5.4	4.6-6.3	24	1.7	1.1-2.5	140	8.1	6.8-9.6	14	0.6	0.3-1.0	2	0.2	0.0-0.6	12	0.8	0.4-1.4
2001	167	5.3	4.6-6.2	34	2.6	1.8-3.7	133	7.5	6.2-8.8	17	0.6	0.4-1.0	3	0.2	0.0-0.7	14	0.9	0.5-1.5
2002	172	5.5	4.7-6.3	31	2.4	1.6-3.4	141	7.8	6.6-9.2	9	0.4	0.2-0.7	2	0.1	0.0-0.6	7	0.5	0.2-0.9
2003	167	5.2	4.5-6.1	31	2.5	1.6-3.6	136	7.4	6.2-8.8	8	0.3	0.1-0.6	1	0.1	0.0-0.4	7	0.5	0.2-0.9
2004	187	5.9	5.1-6.8	33	3.0	2.0-4.3	154	8.4	7.1-9.8	16	0.6	0.4-1.0	7	0.8	0.3-1.6	9	0.6	0.3-1.1
2005	257	7.9	7.0-8.9	45	3.3	2.4-4.5	212	11.5	10.0-13.1	21	0.7	0.4-1.1	9	0.8	0.3-1.5	12	0.7	0.4-1.2
2006	223	7.0	6.1-7.9	36	2.9	2.0-4.1	187	10.1	8.7-11.7	11	0.4	0.2-0.7	3	0.2	0.0-0.7	8	0.5	0.2-0.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Thyroid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	242	7.4	6.5-8.4	50	3.7	2.7-4.9	192	10.3	8.9-11.9	12	0.5	0.2-0.8	3	0.3	0.1-0.9	9	0.5	0.2-1.0
2008	290	8.7	7.8-9.8	60	4.4	3.3-5.8	230	12.1	10.6-13.8	15	0.5	0.3-0.9	5	0.5	0.1-1.2	10	0.6	0.3-1.1
2009	299	9.0	8.0-10.2	59	4.0	3.0-5.2	240	12.9	11.3-14.6	15	0.5	0.3-0.9	5	0.5	0.1-1.1	10	0.5	0.3-1.0
2010	338	9.8	8.8-11.0	52	3.7	2.8-5.0	286	14.6	13.0-16.5	21	0.7	0.4-1.0	11	0.9	0.4-1.7	10	0.5	0.3-1.0
2011	390	10.9	9.9-12.1	79	5.5	4.3-6.9	311	15.4	13.7-17.2	12	0.4	0.2-0.7	1	0.1	0.0-0.6	11	0.6	0.3-1.0
2012	361	10.3	9.2-11.4	58	4.0	3.0-5.2	303	15.3	13.6-17.1	18	0.6	0.3-0.9	6	0.5	0.2-1.2	12	0.6	0.3-1.0
2013	427	11.7	10.6-12.9	56	3.4	2.6-4.5	371	18.2	16.3-20.1	19	0.6	0.4-1.0	5	0.4	0.1-0.9	14	0.7	0.4-1.2
2014	409	11.3	10.2-12.5	62	4.1	3.1-5.3	347	17.2	15.4-19.1	20	0.6	0.4-0.9	3	0.2	0.0-0.7	17	0.8	0.5-1.3
2015	426	11.8	10.7-13.0	81	5.2	4.1-6.5	345	17.1	15.3-19.1	18	0.5	0.3-0.8	6	0.4	0.1-0.9	12	0.6	0.3-1.0
2016	371	10.0	9.0-11.1	70	4.4	3.4-5.5	301	14.6	12.9-16.3	17	0.5	0.3-0.7	5	0.3	0.1-0.7	12	0.5	0.3-0.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Hodgkin lymphoma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	46	2.7	1.8-3.8	21	3.3	1.6-5.8	25	2.4	1.5-3.6	18	0.9	0.5-1.5	10	1.1	0.5-2.1	8	0.8	0.3-1.5
1977	38	1.7	1.2-2.4	20	2.1	1.2-3.4	18	1.5	0.9-2.4	19	1.1	0.6-1.9	7	1.3	0.3-3.0	12	1.1	0.6-2.0
1978	41	1.9	1.3-2.6	24	2.4	1.5-3.7	17	1.5	0.8-2.5	14	0.6	0.3-1.1	8	0.9	0.3-1.9	6	0.4	0.2-1.0
1979	48	2.1	1.5-2.8	30	3.0	1.9-4.5	18	1.4	0.8-2.3	21	1.0	0.6-1.6	14	1.5	0.8-2.6	7	0.7	0.2-1.5
1980	60	2.3	1.7-3.0	32	2.7	1.8-3.9	28	1.9	1.2-2.8	13	0.6	0.3-1.0	6	0.7	0.3-1.6	7	0.5	0.2-1.0
1981	54	2.2	1.6-2.9	33	3.0	1.9-4.5	21	1.5	0.9-2.4	22	1.1	0.6-1.7	13	1.6	0.7-2.9	9	0.7	0.3-1.4
1982	59	2.4	1.8-3.1	32	3.0	2.0-4.3	27	1.9	1.3-2.9	18	0.8	0.4-1.2	9	0.9	0.4-1.9	9	0.7	0.3-1.3
1983	67	2.6	2.0-3.4	39	3.7	2.5-5.4	28	1.9	1.2-2.8	14	0.6	0.3-1.1	8	0.7	0.3-1.5	6	0.5	0.2-1.1
1984	58	2.3	1.7-3.0	39	3.6	2.5-5.1	19	1.4	0.8-2.2	21	0.9	0.5-1.4	19	1.7	1.0-2.9	2	0.2	0.0-0.7
1985	50	1.8	1.3-2.4	31	2.6	1.7-3.8	19	1.1	0.7-1.8	27	1.0	0.7-1.5	18	1.6	0.9-2.7	9	0.6	0.3-1.1
1986	66	2.5	1.9-3.2	46	3.9	2.8-5.3	20	1.3	0.8-2.1	19	0.8	0.4-1.2	13	1.2	0.6-2.1	6	0.5	0.2-1.0
1987	63	2.2	1.7-2.9	36	3.3	2.1-4.8	27	1.6	1.0-2.4	17	0.6	0.4-1.1	14	1.4	0.7-2.4	3	0.2	0.0-0.5
1988	69	2.2	1.7-2.9	43	2.8	2.0-3.9	26	1.7	1.1-2.5	17	0.6	0.3-1.0	9	0.8	0.3-1.6	8	0.5	0.2-1.0
1989	72	2.4	1.9-3.1	43	3.3	2.3-4.6	29	1.8	1.2-2.6	19	0.7	0.4-1.2	16	1.2	0.7-2.1	3	0.3	0.1-0.8
1990	68	2.3	1.8-3.0	39	3.0	2.1-4.3	29	1.8	1.2-2.6	20	0.7	0.4-1.1	12	1.0	0.5-1.9	8	0.5	0.2-1.0
1991	92	3.0	2.4-3.7	53	4.1	3.0-5.6	39	2.2	1.6-3.1	18	0.6	0.3-1.0	8	0.6	0.2-1.2	10	0.6	0.3-1.2
1992	66	2.2	1.7-2.8	36	2.5	1.7-3.6	30	1.8	1.2-2.7	22	0.7	0.4-1.1	12	0.8	0.4-1.6	10	0.6	0.3-1.1
1993	72	2.2	1.7-2.8	38	2.5	1.7-3.5	34	1.9	1.3-2.7	17	0.5	0.3-0.9	10	0.7	0.3-1.4	7	0.4	0.2-0.9
1994	60	2.0	1.5-2.6	36	2.8	1.9-4.0	24	1.4	0.9-2.1	20	0.7	0.4-1.1	13	1.1	0.6-2.0	7	0.4	0.2-0.8
1995	80	2.4	1.9-3.1	48	3.2	2.4-4.4	32	1.8	1.2-2.5	17	0.5	0.3-0.8	13	0.9	0.5-1.6	4	0.2	0.1-0.6
1996	69	2.2	1.7-2.8	34	2.4	1.6-3.5	35	2.0	1.4-2.8	13	0.5	0.2-0.8	7	0.6	0.2-1.4	6	0.4	0.1-0.8
1997	80	2.4	1.9-3.0	51	3.3	2.4-4.4	29	1.6	1.1-2.3	16	0.6	0.3-1.0	6	0.4	0.1-0.9	10	0.7	0.3-1.2
1998	82	2.4	1.9-3.0	43	2.9	2.0-3.9	39	2.1	1.5-2.9	12	0.3	0.2-0.6	5	0.3	0.1-0.8	7	0.4	0.1-0.8
1999	80	2.5	1.9-3.1	43	3.0	2.1-4.2	37	2.0	1.4-2.8	14	0.5	0.3-0.8	9	0.7	0.3-1.4	5	0.3	0.1-0.7
2000	91	2.6	2.1-3.2	55	3.5	2.6-4.6	36	1.9	1.3-2.6	5	0.2	0.0-0.4	3	0.2	0.0-0.6	2	0.1	0.0-0.4
2001	84	2.4	1.9-3.0	55	3.4	2.5-4.5	29	1.5	1.0-2.2	12	0.4	0.2-0.7	6	0.4	0.2-1.0	6	0.3	0.1-0.7
2002	89	2.7	2.2-3.3	49	3.3	2.4-4.4	40	2.2	1.6-3.0	10	0.3	0.1-0.5	8	0.5	0.2-1.1	2	0.1	0.0-0.4
2003	84	2.4	1.9-3.0	35	2.2	1.6-3.2	49	2.6	1.9-3.4	14	0.5	0.3-0.8	8	0.8	0.3-1.5	6	0.3	0.1-0.7
2004	94	2.7	2.2-3.3	48	3.0	2.2-4.0	46	2.4	1.8-3.2	9	0.3	0.1-0.5	6	0.4	0.2-1.0	3	0.2	0.0-0.5
2005	99	2.9	2.4-3.6	60	3.8	2.9-5.0	39	2.1	1.5-2.9	11	0.3	0.2-0.5	9	0.6	0.3-1.1	2	0.1	0.0-0.4
2006	90	2.7	2.1-3.3	49	3.2	2.4-4.3	41	2.2	1.6-3.0	9	0.3	0.1-0.6	5	0.3	0.1-0.8	4	0.2	0.1-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Hodgkin lymphoma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	86	2.5	2.0-3.1	50	3.4	2.5-4.5	36	1.9	1.3-2.6	13	0.4	0.2-0.7	4	0.3	0.1-0.8	9	0.5	0.2-0.9
2008	121	3.5	2.9-4.2	70	4.5	3.5-5.7	51	2.7	2.0-3.5	8	0.2	0.1-0.5	5	0.3	0.1-0.8	3	0.2	0.0-0.5
2009	104	3.0	2.5-3.7	47	3.2	2.3-4.3	57	3.0	2.3-3.9	10	0.3	0.1-0.6	6	0.4	0.2-1.0	4	0.2	0.1-0.5
2010	119	3.4	2.8-4.0	69	4.4	3.4-5.6	50	2.6	1.9-3.4	9	0.2	0.1-0.5	5	0.3	0.1-0.7	4	0.2	0.1-0.5
2011	120	3.2	2.6-3.8	67	3.8	2.9-4.8	53	2.7	2.0-3.5	14	0.4	0.2-0.7	8	0.6	0.2-1.2	6	0.3	0.1-0.6
2012	103	2.9	2.3-3.5	48	2.9	2.1-3.9	55	2.8	2.1-3.7	5	0.2	0.0-0.4	4	0.3	0.1-0.7	1	0.1	0.0-0.3
2013	121	3.4	2.8-4.1	64	4.0	3.0-5.2	57	2.9	2.2-3.8	10	0.3	0.1-0.5	5	0.4	0.1-0.8	5	0.2	0.1-0.6
2014	119	3.2	2.6-3.8	61	3.6	2.7-4.6	58	2.9	2.2-3.7	7	0.2	0.1-0.4	4	0.3	0.1-0.7	3	0.1	0.0-0.4
2015	116	3.2	2.6-3.8	59	3.5	2.6-4.6	57	3.0	2.3-3.9	11	0.3	0.2-0.6	6	0.4	0.2-1.0	5	0.3	0.1-0.6
2016	95	2.5	2.0-3.1	40	2.5	1.8-3.4	55	2.6	2.0-3.5	3	0.1	0.0-0.2	3	0.2	0.0-0.6	0	0.0	0.0-0.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Non-Hodgkin lymphomas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	98	6.1	4.8-7.6	49	7.7	5.4-10.5	49	5.1	3.7-6.8	57	4.0	2.9-5.3	34	6.0	4.0-8.7	23	2.6	1.6-4.0
1977	118	7.1	5.7-8.6	57	8.5	6.2-11.5	61	6.1	4.6-8.0	61	3.8	2.8-5.1	28	4.2	2.5-6.6	33	3.5	2.3-5.1
1978	156	8.9	7.5-10.6	84	10.8	8.4-13.6	72	7.4	5.7-9.5	65	3.7	2.8-4.7	41	5.6	3.9-7.9	24	2.3	1.4-3.5
1979	115	6.8	5.5-8.2	64	8.5	6.3-11.2	51	5.4	3.9-7.2	79	5.2	4.0-6.7	42	6.0	4.1-8.4	37	4.5	3.1-6.4
1980	133	7.5	6.2-9.0	69	9.6	7.1-12.5	64	6.2	4.7-8.0	73	4.3	3.2-5.5	42	5.7	3.9-8.0	31	3.3	2.1-4.7
1981	131	7.5	6.1-9.0	66	7.7	5.7-10.2	65	7.0	5.3-9.0	78	4.4	3.4-5.5	34	4.8	3.2-7.0	44	4.1	2.9-5.6
1982	161	8.4	7.1-9.9	81	9.8	7.6-12.5	80	7.3	5.7-9.2	61	3.6	2.7-4.7	30	4.3	2.7-6.5	31	3.2	2.1-4.5
1983	172	8.6	7.3-10.1	85	9.8	7.6-12.4	87	7.7	6.1-9.6	91	4.8	3.8-5.9	49	5.8	4.1-7.9	42	4.0	2.8-5.4
1984	190	9.3	7.9-10.8	96	11.0	8.7-13.8	94	8.1	6.4-9.9	95	4.8	3.8-5.9	54	6.3	4.5-8.4	41	3.7	2.6-5.0
1985	231	10.8	9.3-12.3	124	13.5	10.9-16.5	107	8.8	7.2-10.7	108	5.3	4.3-6.5	55	6.0	4.3-8.2	53	4.7	3.5-6.2
1986	241	11.4	9.9-13.0	121	13.8	11.1-17.0	120	9.8	8.0-11.8	102	4.9	3.9-6.0	52	6.0	4.2-8.1	50	4.1	3.0-5.5
1987	252	11.2	9.8-12.8	150	16.4	13.5-19.6	102	7.7	6.3-9.4	95	4.4	3.5-5.4	50	5.6	4.0-7.6	45	3.5	2.6-4.8
1988	231	10.1	8.8-11.6	137	14.5	11.9-17.4	94	7.2	5.8-8.8	117	5.7	4.7-6.9	76	9.9	7.5-12.8	41	3.3	2.3-4.5
1989	272	11.2	9.9-12.7	155	14.4	12.0-17.1	117	8.7	7.2-10.5	115	5.1	4.2-6.2	67	7.2	5.3-9.4	48	3.7	2.7-5.0
1990	309	12.3	10.9-13.9	183	16.1	13.7-18.8	126	9.2	7.6-11.0	106	4.7	3.8-5.7	55	5.9	4.2-7.9	51	3.9	2.9-5.2
1991	312	12.7	11.3-14.3	179	16.3	13.7-19.2	133	9.8	8.2-11.7	109	4.8	3.9-5.9	62	6.1	4.5-8.0	47	3.8	2.7-5.0
1992	357	14.3	12.8-16.0	208	19.0	16.3-22.0	149	10.6	9.0-12.5	114	5.0	4.1-6.0	60	6.3	4.6-8.3	54	4.0	3.0-5.3
1993	364	13.4	12.0-14.9	226	18.9	16.4-21.8	138	9.1	7.6-10.7	117	4.9	4.0-5.9	55	5.1	3.7-6.7	62	4.5	3.4-5.8
1994	401	15.5	14.0-17.2	229	19.9	17.3-22.9	172	11.9	10.1-13.8	115	4.9	4.0-6.0	57	5.7	4.2-7.5	58	4.3	3.2-5.5
1995	383	14.6	13.1-16.2	238	21.0	18.2-24.2	145	9.8	8.2-11.6	118	4.9	4.0-5.9	65	5.8	4.4-7.5	53	3.9	2.9-5.1
1996	412	15.3	13.8-16.9	227	18.7	16.2-21.5	185	12.2	10.5-14.1	125	5.3	4.3-6.3	58	6.0	4.4-8.0	67	4.8	3.7-6.0
1997	432	16.1	14.6-17.8	248	21.9	19.0-25.0	184	12.0	10.3-13.9	116	4.9	4.0-5.9	60	6.3	4.7-8.3	56	3.9	2.9-5.1
1998	435	15.8	14.3-17.4	242	21.0	18.2-24.0	193	12.2	10.5-14.0	142	5.8	4.8-6.8	73	7.3	5.6-9.3	69	4.7	3.7-6.0
1999	461	16.3	14.8-17.9	250	21.0	18.3-23.9	211	13.0	11.2-14.8	146	5.5	4.6-6.5	79	7.5	5.8-9.5	67	4.3	3.3-5.5
2000	401	13.6	12.3-15.0	210	16.9	14.5-19.5	191	11.3	9.7-13.0	141	5.2	4.4-6.2	84	7.8	6.1-9.8	57	3.6	2.7-4.6
2001	433	14.9	13.5-16.4	234	18.9	16.4-21.6	199	11.9	10.3-13.7	153	5.7	4.8-6.7	78	7.0	5.4-8.9	75	4.7	3.7-5.9
2002	431	14.2	12.9-15.6	233	17.8	15.5-20.4	198	11.4	9.9-13.1	153	5.4	4.6-6.4	87	7.6	6.0-9.5	66	4.0	3.1-5.0
2003	461	15.3	13.9-16.8	212	16.9	14.6-19.5	249	14.2	12.5-16.1	136	4.7	3.9-5.5	71	6.1	4.7-7.8	65	3.8	2.9-4.8
2004	419	14.0	12.7-15.4	214	17.6	15.1-20.2	205	11.7	10.1-13.4	123	4.3	3.6-5.2	65	5.5	4.2-7.1	58	3.4	2.6-4.5
2005	481	15.6	14.2-17.1	241	18.5	16.1-21.1	240	13.5	11.9-15.4	123	4.4	3.6-5.2	63	5.7	4.3-7.4	60	3.5	2.7-4.5
2006	460	14.8	13.5-16.3	228	17.5	15.2-20.1	232	13.0	11.4-14.8	115	3.8	3.1-4.5	54	4.4	3.2-5.8	61	3.4	2.6-4.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Non-Hodgkin lymphomas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	483	15.2	13.9-16.7	270	20.1	17.7-22.8	213	11.7	10.2-13.4	131	4.4	3.6-5.2	64	5.5	4.2-7.1	67	3.8	2.9-4.8
2008	502	15.6	14.2-17.0	284	20.8	18.4-23.5	218	11.8	10.3-13.5	114	3.7	3.0-4.5	60	5.0	3.7-6.5	54	2.9	2.2-3.8
2009	517	16.2	14.9-17.7	256	19.2	16.8-21.8	261	14.3	12.6-16.1	120	3.8	3.1-4.6	63	5.0	3.8-6.5	57	3.1	2.3-4.0
2010	501	15.2	13.9-16.7	260	18.5	16.2-21.0	241	12.7	11.1-14.4	127	4.1	3.4-4.9	61	4.8	3.6-6.3	66	3.6	2.8-4.6
2011	574	17.1	15.7-18.6	314	23.0	20.4-25.8	260	13.2	11.6-14.9	134	4.3	3.5-5.1	76	6.0	4.6-7.6	58	3.1	2.3-4.0
2012	521	15.2	13.9-16.6	254	17.7	15.5-20.2	267	13.5	11.9-15.3	123	3.7	3.1-4.5	63	5.0	3.8-6.4	60	3.0	2.3-3.9
2013	525	14.9	13.7-16.3	262	17.7	15.5-20.1	263	12.9	11.4-14.6	123	3.7	3.0-4.4	64	4.8	3.6-6.2	59	2.9	2.2-3.8
2014	562	15.5	14.2-16.8	300	19.6	17.4-22.1	262	12.5	11.0-14.1	136	3.9	3.3-4.7	72	5.5	4.2-6.9	64	3.1	2.4-3.9
2015	564	15.4	14.1-16.8	269	17.5	15.4-19.8	295	13.9	12.3-15.6	127	3.6	3.0-4.3	62	4.7	3.5-6.0	65	3.1	2.4-3.9
2016	592	16.1	14.8-17.5	302	19.8	17.5-22.2	290	13.7	12.1-15.4	130	3.6	3.0-4.3	64	4.5	3.4-5.7	66	3.1	2.4-3.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Myeloma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	122	9.0	7.3-10.9	58	10.9	8.0-14.5	64	7.8	5.9-10.1	72	5.4	4.1-7.0	30	4.9	3.2-7.1	42	5.6	3.9-7.7
1977	122	9.1	7.4-11.0	54	10.4	7.3-14.1	68	8.4	6.4-10.7	81	6.4	4.9-8.1	40	8.9	6.0-12.5	41	5.1	3.5-7.0
1978	120	8.6	7.0-10.5	64	13.1	9.6-17.2	56	6.2	4.6-8.2	61	4.4	3.3-5.7	34	7.3	4.7-10.6	27	2.9	1.9-4.2
1979	141	9.3	7.7-11.0	63	10.3	7.7-13.6	78	8.6	6.7-10.8	72	5.0	3.8-6.3	31	5.4	3.4-7.9	41	4.7	3.3-6.5
1980	153	10.1	8.4-12.0	61	10.7	7.9-14.1	92	9.8	7.8-12.1	99	6.6	5.3-8.2	43	7.5	5.2-10.4	56	6.1	4.5-8.0
1981	152	9.9	8.3-11.7	64	10.7	8.0-13.9	88	9.4	7.5-11.7	82	5.6	4.3-7.0	34	6.5	4.3-9.3	48	5.1	3.7-6.8
1982	166	10.5	8.9-12.3	78	12.2	9.4-15.5	88	9.3	7.3-11.5	113	7.0	5.7-8.5	58	9.1	6.7-12.1	55	5.7	4.2-7.5
1983	146	8.7	7.3-10.3	65	9.5	7.1-12.3	81	8.1	6.4-10.1	102	6.5	5.2-8.0	41	7.1	4.9-9.8	61	6.2	4.7-8.1
1984	181	10.8	9.2-12.6	81	12.6	9.7-16.0	100	9.8	7.9-12.0	116	7.1	5.8-8.6	43	6.8	4.8-9.4	73	7.3	5.6-9.2
1985	172	9.7	8.2-11.3	75	11.6	8.8-14.9	97	8.7	7.0-10.7	109	6.2	5.0-7.5	54	8.2	5.9-10.9	55	5.1	3.8-6.7
1986	186	10.5	9.0-12.2	85	12.8	9.9-16.1	101	9.2	7.5-11.3	114	6.9	5.6-8.3	56	8.7	6.4-11.6	58	5.7	4.3-7.4
1987	164	8.8	7.5-10.3	79	11.5	8.9-14.6	85	7.3	5.8-9.1	99	5.5	4.5-6.8	53	8.2	5.9-10.9	46	4.1	3.0-5.5
1988	154	8.0	6.7-9.4	68	9.1	6.8-11.7	86	7.3	5.8-9.0	83	4.5	3.5-5.6	39	5.5	3.8-7.7	44	3.9	2.8-5.2
1989	191	10.4	8.9-12.0	86	12.3	9.6-15.6	105	9.3	7.6-11.3	90	5.1	4.1-6.3	39	6.5	4.4-9.1	51	4.5	3.3-5.9
1990	193	9.5	8.2-11.0	85	12.1	9.3-15.2	108	8.5	6.9-10.3	98	5.3	4.3-6.5	42	6.5	4.5-8.9	56	4.8	3.6-6.2
1991	204	10.3	8.9-11.9	73	10.0	7.6-12.8	131	10.5	8.8-12.5	110	5.7	4.6-6.9	45	6.7	4.7-9.1	65	5.3	4.1-6.8
1992	237	11.3	9.9-12.9	108	14.0	11.2-17.2	129	9.9	8.2-11.8	132	6.5	5.4-7.7	53	7.8	5.6-10.4	79	6.1	4.8-7.6
1993	187	8.7	7.4-10.1	81	9.8	7.6-12.4	106	8.0	6.5-9.7	114	5.5	4.5-6.7	47	6.5	4.6-8.8	67	5.1	3.9-6.5
1994	205	9.8	8.5-11.3	88	11.2	8.8-14.0	117	9.0	7.4-10.8	119	5.8	4.8-7.0	43	5.4	3.8-7.4	76	5.9	4.6-7.4
1995	215	9.8	8.5-11.3	96	12.2	9.7-15.1	119	8.7	7.2-10.4	138	6.4	5.4-7.6	59	7.5	5.6-9.9	79	5.9	4.6-7.3
1996	221	9.6	8.4-11.0	97	10.8	8.6-13.4	124	8.9	7.4-10.6	109	4.9	4.0-6.0	55	6.9	5.0-9.1	54	3.9	2.9-5.1
1997	249	10.8	9.4-12.2	111	12.3	10.0-15.0	138	9.7	8.2-11.5	134	6.0	5.0-7.1	70	8.7	6.7-11.2	64	4.6	3.5-5.8
1998	246	10.4	9.1-11.8	108	12.0	9.7-14.7	138	9.6	8.0-11.3	112	4.9	4.0-5.9	41	4.9	3.4-6.8	71	5.0	3.9-6.3
1999	254	10.5	9.3-12.0	102	10.3	8.3-12.7	152	10.3	8.8-12.1	121	5.2	4.3-6.2	48	5.3	3.8-7.1	73	5.0	3.9-6.3
2000	279	10.9	9.7-12.3	117	11.6	9.4-14.1	162	10.5	8.9-12.2	116	5.0	4.1-6.0	42	4.8	3.4-6.6	74	5.0	3.9-6.3
2001	300	11.7	10.4-13.1	125	12.4	10.1-14.9	175	11.2	9.6-13.0	136	5.5	4.6-6.6	54	5.9	4.3-7.8	82	5.3	4.2-6.6
2002	289	11.1	9.8-12.5	116	11.9	9.7-14.4	173	10.8	9.2-12.5	147	5.7	4.8-6.8	59	5.6	4.2-7.4	88	5.7	4.6-7.0
2003	314	11.8	10.5-13.2	135	13.4	11.1-16.0	179	11.0	9.4-12.7	124	4.6	3.8-5.5	55	5.1	3.8-6.8	69	4.3	3.3-5.4
2004	304	11.1	9.8-12.4	128	12.0	9.8-14.4	176	10.6	9.1-12.3	136	5.3	4.5-6.3	63	6.7	5.0-8.6	73	4.6	3.6-5.8
2005	364	13.1	11.7-14.5	159	14.3	12.1-16.9	205	12.1	10.5-13.9	125	4.6	3.8-5.5	51	4.7	3.4-6.3	74	4.4	3.5-5.6
2006	357	12.6	11.3-14.0	171	15.3	13.0-17.9	186	10.8	9.3-12.4	125	4.6	3.8-5.4	60	5.6	4.2-7.3	65	3.9	3.0-4.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Myeloma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	311	10.9	9.7-12.2	141	12.9	10.7-15.3	170	9.7	8.3-11.3	153	5.5	4.6-6.4	70	6.4	4.9-8.2	83	4.8	3.8-6.0
2008	409	13.5	12.2-14.9	188	15.7	13.5-18.3	221	12.1	10.5-13.8	118	4.2	3.5-5.0	53	4.9	3.6-6.4	65	3.7	2.9-4.8
2009	434	14.2	12.9-15.7	221	17.7	15.3-20.4	213	11.7	10.2-13.4	129	4.3	3.6-5.2	71	5.9	4.5-7.5	58	3.2	2.5-4.2
2010	425	13.7	12.4-15.1	217	18.5	16.0-21.2	208	11.0	9.6-12.7	151	5.2	4.4-6.1	69	6.3	4.8-8.0	82	4.5	3.6-5.6
2011	459	14.3	13.0-15.7	213	16.4	14.2-18.9	246	13.0	11.4-14.7	141	4.5	3.8-5.3	62	5.2	3.9-6.8	79	4.1	3.2-5.1
2012	488	14.7	13.4-16.1	229	17.5	15.2-20.0	259	13.0	11.4-14.7	144	4.4	3.7-5.2	72	5.3	4.1-6.8	72	3.7	2.9-4.7
2013	489	14.4	13.1-15.8	244	18.2	15.9-20.8	245	12.0	10.6-13.7	144	4.4	3.7-5.2	75	6.0	4.6-7.6	69	3.5	2.7-4.4
2014	495	14.2	12.9-15.5	247	17.5	15.2-19.9	248	12.1	10.6-13.7	148	4.6	3.9-5.4	68	5.5	4.2-7.0	80	4.1	3.2-5.1
2015	464	12.9	11.8-14.2	227	15.7	13.6-17.9	237	11.1	9.7-12.7	181	5.1	4.4-6.0	87	6.4	5.1-8.0	94	4.3	3.5-5.3
2016	505	13.6	12.4-14.8	248	16.7	14.6-19.0	257	11.6	10.2-13.2	165	4.7	4.0-5.5	75	5.5	4.3-6.9	90	4.3	3.5-5.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Leukemias

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	132	7.7	6.3-9.4	70	9.9	7.3-13.1	62	6.3	4.7-8.2	92	5.8	4.5-7.3	53	7.8	5.4-11.0	39	4.4	3.0-6.2
1977	126	8.0	6.5-9.7	67	11.3	8.1-15.2	59	6.1	4.6-8.0	101	6.0	4.8-7.5	54	8.4	6.0-11.5	47	4.6	3.3-6.2
1978	157	9.4	7.8-11.2	76	10.9	8.2-14.2	81	8.4	6.6-10.6	115	7.3	5.9-8.9	62	9.0	6.6-11.9	53	6.0	4.4-8.0
1979	142	8.7	7.2-10.4	72	11.3	8.3-14.9	70	7.2	5.5-9.3	98	5.8	4.6-7.2	56	9.0	6.5-12.1	42	4.0	2.8-5.5
1980	158	9.0	7.5-10.7	74	10.5	7.8-13.7	84	8.1	6.4-10.2	112	6.5	5.3-8.0	49	7.6	5.2-10.6	63	6.1	4.6-7.9
1981	156	9.1	7.6-10.8	83	12.5	9.5-16.1	73	7.1	5.5-9.1	108	6.4	5.1-7.9	46	7.4	5.0-10.5	62	6.0	4.5-7.8
1982	156	8.4	7.0-9.9	78	11.0	8.3-14.2	78	6.9	5.4-8.7	90	5.0	3.9-6.2	44	6.5	4.4-9.2	46	4.2	3.0-5.6
1983	172	9.0	7.6-10.6	89	12.1	9.3-15.3	83	7.3	5.7-9.1	105	5.7	4.6-7.0	50	7.1	5.0-9.7	55	4.9	3.6-6.5
1984	164	8.2	6.9-9.7	86	9.8	7.5-12.5	78	6.9	5.4-8.8	105	5.5	4.4-6.7	54	7.1	5.0-9.6	51	4.5	3.3-6.0
1985	193	9.3	8.0-10.8	111	13.4	10.7-16.5	82	6.8	5.4-8.5	100	5.1	4.1-6.3	61	8.2	6.0-10.9	39	3.2	2.3-4.5
1986	171	8.2	7.0-9.7	96	12.0	9.4-15.1	75	6.0	4.7-7.6	110	5.2	4.2-6.3	61	6.9	5.0-9.1	49	4.0	2.9-5.4
1987	190	9.0	7.7-10.4	94	10.8	8.4-13.6	96	7.8	6.3-9.6	105	5.0	4.0-6.1	51	6.2	4.3-8.5	54	4.4	3.3-5.8
1988	155	7.6	6.4-9.0	77	10.7	8.2-13.6	78	6.1	4.8-7.7	116	6.0	4.9-7.3	55	7.5	5.4-10.0	61	5.2	3.9-6.7
1989	203	9.5	8.1-10.9	105	12.6	9.9-15.7	98	7.6	6.2-9.4	100	4.9	4.0-6.1	61	8.4	6.2-11.0	39	3.1	2.2-4.3
1990	218	9.9	8.5-11.4	107	11.7	9.3-14.6	111	8.6	7.0-10.4	128	6.2	5.1-7.4	59	7.6	5.6-10.1	69	5.4	4.1-6.8
1991	238	10.5	9.1-12.0	115	13.0	10.4-16.0	123	9.1	7.5-10.9	106	4.9	3.9-6.0	55	6.6	4.7-9.0	51	4.0	2.9-5.2
1992	229	9.9	8.6-11.3	104	11.3	8.9-14.2	125	9.1	7.5-10.9	107	5.1	4.1-6.2	48	5.4	3.8-7.4	59	4.7	3.5-6.1
1993	205	8.9	7.7-10.3	103	11.7	9.2-14.5	102	7.4	6.0-9.0	111	4.9	4.0-5.9	55	6.2	4.4-8.3	56	4.1	3.1-5.4
1994	211	9.0	7.7-10.3	109	12.0	9.6-14.8	102	7.2	5.8-8.7	156	7.0	5.9-8.3	74	8.4	6.4-10.9	82	6.1	4.8-7.6
1995	230	9.2	8.0-10.5	121	11.7	9.5-14.3	109	7.5	6.1-9.1	135	5.6	4.7-6.7	66	7.4	5.5-9.6	69	4.7	3.7-6.0
1996	249	9.9	8.6-11.2	139	13.4	11.0-16.1	110	7.5	6.1-9.0	139	5.9	4.9-7.0	66	7.0	5.2-9.1	73	5.2	4.0-6.5
1997	224	9.0	7.8-10.3	100	9.6	7.6-11.9	124	8.4	7.0-10.1	134	5.7	4.7-6.8	72	8.2	6.3-10.6	62	4.3	3.2-5.5
1998	227	8.8	7.7-10.1	103	10.5	8.3-13.0	124	8.1	6.7-9.7	123	4.9	4.1-5.9	59	6.1	4.5-8.1	64	4.3	3.3-5.5
1999	207	7.7	6.6-8.8	106	10.7	8.5-13.2	101	6.2	5.0-7.5	129	5.2	4.3-6.2	65	7.0	5.2-9.1	64	4.2	3.2-5.4
2000	257	9.7	8.5-11.0	130	11.9	9.8-14.3	127	8.1	6.7-9.6	125	4.9	4.1-5.9	53	5.6	4.0-7.5	72	4.6	3.6-5.8
2001	236	8.7	7.6-9.9	108	9.8	7.9-12.0	128	7.9	6.6-9.4	137	5.3	4.4-6.2	65	6.4	4.8-8.2	72	4.6	3.6-5.7
2002	271	9.8	8.6-11.0	145	12.9	10.7-15.4	126	7.7	6.4-9.2	124	4.7	3.9-5.6	61	5.9	4.4-7.8	63	3.9	3.0-5.0
2003	264	9.0	7.9-10.2	130	10.6	8.7-12.7	134	7.8	6.5-9.3	128	4.7	3.9-5.6	57	4.9	3.7-6.5	71	4.4	3.4-5.5
2004	286	9.8	8.7-11.1	164	14.0	11.7-16.4	122	7.1	5.9-8.5	133	5.0	4.2-5.9	70	6.9	5.3-8.9	63	3.9	3.0-5.0
2005	239	8.1	7.1-9.3	122	10.5	8.6-12.7	117	6.7	5.5-8.0	130	4.7	3.9-5.6	62	6.0	4.5-7.8	68	4.0	3.1-5.1
2006	282	9.4	8.3-10.6	163	13.6	11.4-16.0	119	6.8	5.6-8.1	119	4.1	3.4-4.9	66	5.8	4.4-7.5	53	3.1	2.3-4.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State
Blacks
Leukemias

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	274	9.1	8.0-10.3	131	10.4	8.6-12.5	143	8.1	6.8-9.6	128	4.5	3.8-5.4	68	6.3	4.8-8.1	60	3.5	2.7-4.5
2008	271	8.8	7.8-10.0	133	11.1	9.2-13.3	138	7.6	6.4-9.0	139	4.7	3.9-5.5	74	6.6	5.1-8.4	65	3.6	2.8-4.6
2009	275	8.8	7.7-9.9	151	11.8	9.9-14.0	124	6.8	5.6-8.1	117	3.8	3.1-4.6	65	5.1	3.9-6.6	52	2.9	2.1-3.7
2010	358	11.0	9.9-12.2	198	14.9	12.8-17.3	160	8.5	7.2-9.9	137	4.5	3.7-5.3	74	6.5	5.0-8.2	63	3.4	2.6-4.3
2011	341	10.4	9.3-11.6	182	13.3	11.3-15.4	159	8.3	7.1-9.7	159	5.0	4.2-5.8	85	6.5	5.1-8.1	74	3.9	3.0-4.9
2012	344	10.2	9.1-11.4	178	13.2	11.2-15.3	166	8.4	7.2-9.8	137	4.3	3.6-5.1	77	6.1	4.8-7.8	60	3.1	2.4-4.0
2013	387	11.2	10.1-12.4	191	13.3	11.4-15.4	196	9.8	8.5-11.3	141	4.2	3.6-5.0	85	6.6	5.2-8.2	56	2.7	2.1-3.6
2014	405	11.5	10.4-12.7	220	15.4	13.4-17.7	185	9.1	7.8-10.5	150	4.5	3.8-5.3	76	5.8	4.5-7.4	74	3.7	2.9-4.6
2015	337	9.5	8.5-10.6	170	12.1	10.3-14.2	167	8.0	6.8-9.3	153	4.4	3.7-5.2	75	5.4	4.2-6.8	78	3.8	3.0-4.8
2016	393	10.9	9.8-12.0	203	13.9	12.0-16.1	190	9.0	7.8-10.4	178	4.9	4.2-5.7	88	6.1	4.8-7.6	90	4.1	3.3-5.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
All Invasive Malignant Tumors

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	4230	374.6	362.2-387.2	2066	482.1	458.1-506.8	2164	312.9	298.9-327.4	2663	245.5	235.2-256.0	1398	330.6	310.6-351.5	1265	194.3	182.8-206.2
1977	4456	386.0	373.7-398.7	2161	507.3	482.7-532.7	2295	319.3	305.5-333.5	2631	231.9	222.2-241.8	1394	326.4	306.9-346.8	1237	176.6	166.2-187.4
1978	4704	402.5	390.0-415.2	2294	519.0	494.8-543.9	2410	336.8	322.6-351.5	2885	250.9	240.9-261.1	1462	327.9	308.9-347.7	1423	205.2	193.9-216.9
1979	4941	410.1	397.8-422.7	2429	539.4	515.3-564.3	2512	337.6	323.7-351.9	2959	254.3	244.4-264.5	1579	360.1	340.1-381.0	1380	193.3	182.6-204.5
1980	5293	423.3	411.1-435.8	2643	573.3	548.7-598.6	2650	341.0	327.4-355.0	3048	252.8	243.1-262.7	1597	356.6	336.9-377.0	1451	194.2	183.7-205.0
1981	5183	401.1	389.4-413.0	2594	539.1	515.9-563.1	2589	324.8	311.8-338.3	3133	249.1	239.8-258.7	1629	347.9	328.9-367.6	1504	193.4	183.2-204.0
1982	5433	410.2	398.6-422.0	2671	545.9	522.9-569.5	2762	336.2	323.2-349.5	3147	243.8	234.8-253.1	1662	349.8	331.2-369.2	1485	184.8	175.1-194.9
1983	5493	395.4	384.3-406.6	2668	523.3	501.3-546.0	2825	327.7	315.2-340.4	3348	248.1	239.2-257.1	1755	349.1	331.1-367.8	1593	190.8	181.2-200.8
1984	5949	417.2	406.1-428.6	2877	546.4	524.3-569.2	3072	349.1	336.4-362.1	3269	238.7	230.1-247.5	1689	337.2	319.5-355.7	1580	185.4	176.0-195.1
1985	6112	417.8	406.8-429.1	3011	559.1	537.0-581.7	3101	342.5	330.1-355.2	3291	235.5	227.1-244.2	1734	342.4	324.6-360.9	1557	178.3	169.3-187.7
1986	6240	415.1	404.4-426.1	3140	573.7	551.6-596.4	3100	332.2	320.3-344.5	3326	230.5	222.3-238.9	1693	325.9	308.8-343.6	1633	180.3	171.3-189.5
1987	6340	409.3	398.8-420.0	3138	550.7	529.5-572.4	3202	334.1	322.3-346.2	3322	225.0	217.1-233.2	1705	323.1	306.2-340.6	1617	174.2	165.5-183.1
1988	6345	405.3	394.9-415.9	3059	540.9	519.7-562.5	3286	336.8	325.0-348.8	3309	220.1	212.3-228.1	1671	309.6	293.4-326.5	1638	172.8	164.3-181.7
1989	6537	408.5	398.2-419.0	3259	564.2	542.9-586.0	3278	328.6	317.2-340.3	3249	211.7	204.2-219.5	1672	307.1	291.0-323.8	1577	162.4	154.3-170.8
1990	6764	417.4	407.1-428.0	3306	564.7	543.6-586.4	3458	342.2	330.6-354.1	3383	220.0	212.3-227.8	1679	309.0	292.8-325.8	1704	174.8	166.4-183.5
1991	7224	440.3	429.8-451.0	3498	589.3	567.9-611.3	3726	364.4	352.5-376.6	3313	211.0	203.6-218.6	1677	302.4	286.5-318.8	1636	164.8	156.7-173.1
1992	7465	444.7	434.3-455.3	3689	605.8	584.4-627.6	3776	361.5	349.8-373.4	3293	207.0	199.7-214.5	1570	286.3	270.9-302.4	1723	169.8	161.7-178.1
1993	7351	429.8	419.7-440.1	3747	601.0	580.1-622.4	3604	339.2	328.0-350.7	3245	200.0	192.9-207.2	1584	281.0	265.8-296.6	1661	161.4	153.6-169.5
1994	7939	456.9	446.6-467.4	3987	623.9	603.0-645.3	3952	367.2	355.6-379.0	3635	220.7	213.4-228.2	1749	301.7	286.3-317.5	1886	180.8	172.6-189.2
1995	7783	438.1	428.2-448.2	3996	606.9	586.7-627.5	3787	344.9	333.9-356.2	3654	217.6	210.4-225.0	1739	292.2	277.4-307.5	1915	180.2	172.1-188.6
1996	8197	453.4	443.4-463.6	4076	609.5	589.4-630.0	4121	368.5	357.2-380.0	3567	209.3	202.3-216.5	1755	290.7	276.2-305.8	1812	168.0	160.2-176.0
1997	8314	456.2	446.2-466.3	4141	616.9	596.9-637.4	4173	368.9	357.7-380.4	3512	203.6	196.7-210.6	1683	277.5	263.3-292.1	1829	166.8	159.1-174.7
1998	8432	450.9	441.1-460.8	4070	585.8	566.7-605.3	4362	377.3	366.1-388.8	3447	194.5	187.9-201.3	1595	251.1	238.0-264.7	1852	165.8	158.3-173.6
1999	8568	448.2	438.6-458.0	4318	609.9	590.7-629.5	4250	358.7	347.9-369.8	3552	195.6	189.1-202.3	1640	257.5	244.2-271.2	1912	166.2	158.8-173.9
2000	8904	453.3	443.7-462.9	4528	619.4	600.5-638.8	4376	360.2	349.6-371.1	3433	184.4	178.2-190.8	1619	246.0	233.4-259.1	1814	153.5	146.4-160.7
2001	8795	438.8	429.5-448.2	4474	597.7	579.3-616.6	4321	349.6	339.2-360.2	3571	188.3	182.1-194.7	1627	243.8	231.3-256.7	1944	161.4	154.2-168.8
2002	8900	435.8	426.7-445.1	4458	583.4	565.4-601.8	4442	353.7	343.3-364.3	3427	177.0	171.0-183.1	1616	235.2	223.1-247.7	1811	147.8	141.0-154.8
2003	9157	440.5	431.4-449.8	4587	589.5	571.6-607.8	4570	357.3	347.0-367.9	3314	167.1	161.4-173.0	1525	215.0	203.6-226.8	1789	143.0	136.4-149.8
2004	9379	443.9	434.8-453.1	4726	591.4	573.7-609.4	4653	359.8	349.5-370.3	3366	168.1	162.4-174.0	1574	218.4	207.1-230.2	1792	142.1	135.5-148.9
2005	9544	445.0	436.0-454.2	4785	586.7	569.4-604.5	4759	364.2	353.8-374.7	3350	164.3	158.7-170.1	1490	204.3	193.4-215.6	1860	145.1	138.6-151.9
2006	9699	448.4	439.4-457.6	4861	591.5	574.2-609.3	4838	366.4	356.1-377.0	3297	160.7	155.2-166.4	1499	205.9	195.0-217.3	1798	139.2	132.8-145.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
All Invasive Malignant Tumors

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	9976	454.5	445.5-463.7	5175	612.1	594.7-629.9	4801	361.5	351.3-372.0	3363	160.7	155.2-166.3	1540	203.8	193.1-214.8	1823	139.3	132.9-145.9
2008	10125	453.1	444.1-462.2	5073	588.0	571.1-605.2	5052	374.3	364.0-384.9	3307	156.0	150.7-161.5	1492	197.1	186.7-208.0	1815	136.2	130.0-142.7
2009	10336	453.6	444.7-462.6	5233	589.2	572.6-606.2	5103	372.3	362.0-382.8	3292	152.2	147.0-157.6	1556	200.0	189.7-210.8	1736	128.0	122.0-134.3
2010	10396	446.3	437.6-455.2	5182	567.6	551.5-584.1	5214	374.3	364.1-384.7	3363	152.0	146.8-157.3	1563	197.1	186.9-207.6	1800	130.0	124.0-136.2
2011	11213	466.9	458.1-475.8	5606	592.2	576.1-608.7	5607	391.6	381.3-402.1	3580	157.5	152.2-162.8	1603	192.3	182.5-202.4	1977	139.8	133.6-146.2
2012	10819	440.5	432.1-449.1	5231	538.9	523.7-554.4	5588	382.5	372.4-392.8	3528	150.2	145.2-155.3	1585	182.8	173.5-192.5	1943	133.5	127.6-139.7
2013	11107	440.0	431.7-448.5	5377	534.3	519.4-549.4	5730	384.6	374.6-394.9	3403	141.0	136.3-146.0	1584	178.5	169.4-187.9	1819	121.1	115.6-126.9
2014	10979	426.3	418.2-434.5	5208	501.0	486.9-515.4	5771	381.5	371.5-391.6	3590	144.8	140.1-149.8	1616	174.5	165.8-183.7	1974	129.5	123.8-135.5
2015	10928	413.8	405.9-421.8	5178	486.3	472.6-500.3	5750	371.1	361.4-381.0	3407	133.5	129.0-138.2	1513	158.6	150.4-167.1	1894	120.7	115.2-126.3
2016	11154	414.4	406.6-422.4	5363	493.2	479.6-507.1	5791	367.5	357.9-377.3	3740	142.3	137.7-147.0	1726	172.7	164.3-181.4	2014	125.5	120.0-131.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Oral cavity and pharynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	193	15.1	12.9-17.6	132	26.2	21.3-31.8	61	8.1	6.1-10.4	92	7.3	5.8-9.1	62	12.4	9.3-16.1	30	4.0	2.6-5.8
1977	187	14.3	12.2-16.6	142	27.3	22.6-32.7	45	5.6	4.1-7.6	103	8.1	6.5-9.9	83	15.5	12.2-19.4	20	2.9	1.7-4.7
1978	205	14.7	12.7-17.0	147	26.0	21.7-31.0	58	7.1	5.4-9.3	107	8.0	6.5-9.8	80	15.9	12.1-20.4	27	3.2	2.1-4.7
1979	213	15.9	13.8-18.3	156	29.0	24.3-34.3	57	7.2	5.4-9.4	109	8.1	6.5-9.8	78	14.6	11.2-18.7	31	3.8	2.6-5.5
1980	213	15.2	13.2-17.5	154	27.8	23.3-33.0	59	7.0	5.3-9.1	103	7.5	6.0-9.1	85	15.8	12.2-20.0	18	2.1	1.2-3.3
1981	224	15.2	13.2-17.4	167	27.4	23.2-32.1	57	6.8	5.1-8.9	88	6.1	4.8-7.6	63	11.1	8.4-14.5	25	2.9	1.8-4.3
1982	227	15.2	13.2-17.4	156	26.6	22.3-31.5	71	7.9	6.1-10.0	117	8.1	6.7-9.8	89	15.9	12.5-19.9	28	3.1	2.1-4.6
1983	211	14.2	12.3-16.3	153	25.8	21.6-30.6	58	6.6	5.0-8.6	110	7.7	6.2-9.3	86	15.4	12.1-19.4	24	2.8	1.8-4.3
1984	258	16.2	14.3-18.4	170	26.2	22.3-30.6	88	9.4	7.5-11.6	91	6.0	4.8-7.4	68	11.6	8.8-15.0	23	2.4	1.5-3.7
1985	247	15.5	13.6-17.6	165	25.8	21.8-30.4	82	8.7	6.8-10.8	82	5.4	4.3-6.8	57	9.7	7.2-12.7	25	2.8	1.8-4.1
1986	251	15.2	13.3-17.2	180	27.1	23.1-31.6	71	7.3	5.6-9.2	100	6.6	5.3-8.1	72	12.3	9.4-15.8	28	3.1	2.0-4.6
1987	270	15.9	14.0-18.0	190	28.1	24.1-32.6	80	7.9	6.2-9.8	76	4.6	3.6-5.8	57	8.9	6.6-11.8	19	1.9	1.1-2.9
1988	231	13.4	11.7-15.3	150	22.2	18.5-26.3	81	7.8	6.1-9.7	96	5.6	4.5-6.9	71	10.4	8.0-13.2	25	2.5	1.6-3.7
1989	223	12.5	10.9-14.3	148	21.1	17.7-25.0	75	6.9	5.4-8.7	91	5.2	4.2-6.5	65	9.7	7.3-12.6	26	2.5	1.6-3.7
1990	237	13.7	12.0-15.6	161	24.1	20.2-28.4	76	7.3	5.7-9.1	74	4.6	3.6-5.8	52	9.0	6.5-12.2	22	2.1	1.3-3.2
1991	306	17.5	15.6-19.7	211	31.9	27.4-36.9	95	8.8	7.1-10.8	107	6.3	5.1-7.6	77	11.1	8.7-14.0	30	3.1	2.1-4.4
1992	281	15.7	13.9-17.7	184	26.4	22.5-30.7	97	9.0	7.3-11.0	80	4.5	3.6-5.6	53	7.8	5.7-10.4	27	2.5	1.6-3.6
1993	266	14.6	12.8-16.5	172	24.0	20.4-28.0	94	8.5	6.9-10.4	83	4.7	3.7-5.8	60	8.6	6.5-11.3	23	2.2	1.4-3.3
1994	256	14.0	12.3-15.8	174	24.3	20.6-28.4	82	7.4	5.9-9.2	91	5.1	4.1-6.2	66	9.9	7.4-12.8	25	2.3	1.5-3.3
1995	220	11.6	10.1-13.2	162	21.4	18.1-25.1	58	5.1	3.9-6.6	99	5.5	4.4-6.7	71	10.0	7.7-12.8	28	2.6	1.7-3.7
1996	239	12.4	10.8-14.1	152	19.9	16.8-23.5	87	7.5	6.0-9.2	81	4.4	3.5-5.5	59	8.3	6.2-10.8	22	2.0	1.3-3.1
1997	222	11.5	10.0-13.1	149	19.0	16.0-22.4	73	6.4	5.0-8.1	65	3.4	2.6-4.3	49	6.4	4.6-8.5	16	1.4	0.8-2.3
1998	210	10.5	9.1-12.1	139	17.3	14.4-20.5	71	6.1	4.7-7.6	70	3.7	2.9-4.7	45	6.2	4.4-8.4	25	2.2	1.4-3.2
1999	186	9.2	7.9-10.6	118	14.4	11.8-17.3	68	5.7	4.4-7.2	59	3.1	2.3-4.0	35	4.7	3.2-6.6	24	2.1	1.3-3.1
2000	207	10.2	8.8-11.7	139	17.5	14.5-20.8	68	5.6	4.3-7.1	63	3.2	2.5-4.1	45	6.0	4.3-8.2	18	1.5	0.9-2.4
2001	188	8.8	7.6-10.2	133	16.0	13.3-19.0	55	4.3	3.2-5.6	70	3.4	2.6-4.3	51	6.3	4.6-8.3	19	1.5	0.9-2.4
2002	199	9.3	8.1-10.8	125	14.8	12.2-17.8	74	5.8	4.6-7.3	50	2.3	1.7-3.1	34	3.9	2.6-5.5	16	1.3	0.7-2.1
2003	188	8.6	7.4-9.9	130	14.5	12.1-17.3	58	4.6	3.5-5.9	50	2.4	1.8-3.1	37	4.6	3.2-6.4	13	1.0	0.6-1.8
2004	221	9.9	8.6-11.3	139	15.8	13.2-18.8	82	6.2	4.9-7.7	71	3.3	2.6-4.2	53	6.3	4.6-8.4	18	1.4	0.9-2.3
2005	200	8.9	7.7-10.3	140	16.1	13.4-19.1	60	4.5	3.5-5.9	63	2.9	2.2-3.7	44	5.4	3.8-7.3	19	1.4	0.8-2.2
2006	190	8.3	7.2-9.6	121	13.3	11.0-16.0	69	5.1	3.9-6.4	42	1.8	1.3-2.5	33	3.7	2.5-5.2	9	0.7	0.3-1.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Oral cavity and pharynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	222	9.8	8.5-11.2	140	15.3	12.8-18.2	82	6.2	4.9-7.7	59	2.6	2.0-3.4	46	5.3	3.8-7.2	13	1.0	0.5-1.6
2008	188	8.1	6.9-9.3	126	13.3	11.0-15.9	62	4.5	3.4-5.7	57	2.6	1.9-3.4	42	4.9	3.5-6.7	15	1.1	0.6-1.9
2009	191	8.1	6.9-9.3	124	12.8	10.6-15.3	67	4.8	3.7-6.2	54	2.3	1.7-3.0	37	4.0	2.7-5.5	17	1.2	0.7-1.9
2010	204	8.5	7.4-9.8	131	13.5	11.2-16.2	73	5.1	4.0-6.5	50	2.1	1.6-2.8	26	2.9	1.9-4.3	24	1.7	1.1-2.5
2011	206	8.2	7.1-9.5	141	13.7	11.5-16.3	65	4.5	3.5-5.8	72	2.9	2.3-3.7	50	5.1	3.7-6.8	22	1.5	0.9-2.3
2012	211	8.3	7.2-9.5	133	12.8	10.6-15.3	78	5.2	4.1-6.5	52	2.2	1.6-2.8	32	3.3	2.2-4.8	20	1.4	0.9-2.2
2013	208	8.1	7.0-9.3	140	13.3	11.1-15.7	68	4.6	3.5-5.8	66	2.7	2.1-3.5	45	4.7	3.4-6.3	21	1.4	0.9-2.2
2014	195	7.2	6.2-8.3	117	10.3	8.5-12.5	78	5.1	4.0-6.3	46	1.8	1.3-2.4	28	2.8	1.8-4.1	18	1.2	0.7-1.9
2015	193	7.2	6.2-8.3	116	10.6	8.7-12.9	77	4.9	3.9-6.2	50	1.9	1.4-2.5	35	3.1	2.2-4.4	15	1.0	0.5-1.7
2016	212	7.9	6.8-9.0	140	12.8	10.7-15.2	72	4.6	3.6-5.8	44	1.6	1.1-2.1	27	2.5	1.6-3.7	17	1.0	0.6-1.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Esophagus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	161	12.5	10.6-14.8	116	22.6	18.3-27.6	45	5.9	4.2-8.0	156	12.7	10.7-15.1	118	24.4	19.7-29.8	38	5.2	3.6-7.3
1977	169	13.4	11.3-15.7	118	24.4	19.7-29.9	51	6.6	4.9-8.7	148	11.6	9.7-13.8	107	22.1	17.4-27.4	41	5.2	3.7-7.1
1978	181	14.1	12.0-16.5	128	26.2	21.3-31.7	53	6.6	4.9-8.7	125	9.5	7.8-11.4	83	15.9	12.5-20.0	42	5.3	3.8-7.2
1979	188	13.7	11.7-16.0	127	23.7	19.3-28.6	61	7.5	5.6-9.7	173	13.1	11.1-15.3	125	24.7	20.0-30.1	48	5.9	4.3-7.9
1980	201	14.8	12.7-17.1	150	28.9	24.0-34.5	51	6.1	4.5-8.1	166	12.4	10.4-14.5	108	21.3	17.0-26.2	58	6.9	5.2-9.1
1981	207	14.2	12.3-16.4	144	25.1	20.9-29.8	63	7.2	5.5-9.2	167	11.4	9.7-13.3	123	21.0	17.3-25.3	44	5.1	3.6-6.8
1982	198	13.6	11.7-15.7	143	25.8	21.4-30.7	55	6.0	4.5-7.9	167	11.7	10.0-13.7	117	22.0	17.9-26.8	50	5.6	4.1-7.4
1983	195	12.8	11.0-14.9	134	22.1	18.3-26.5	61	6.9	5.2-8.9	162	10.8	9.1-12.6	113	18.8	15.3-22.8	49	5.6	4.1-7.5
1984	209	13.7	11.8-15.7	139	23.6	19.5-28.3	70	7.5	5.8-9.5	160	10.6	9.0-12.5	112	19.3	15.6-23.6	48	5.3	3.9-7.1
1985	188	12.1	10.4-14.0	126	20.7	17.0-24.9	62	6.7	5.1-8.6	172	11.1	9.4-13.0	126	21.1	17.3-25.4	46	4.9	3.5-6.6
1986	188	12.0	10.3-13.9	141	23.7	19.7-28.3	47	4.9	3.6-6.6	165	10.5	8.9-12.3	114	18.7	15.2-22.7	51	5.5	4.0-7.2
1987	181	11.4	9.8-13.3	115	19.3	15.7-23.4	66	6.7	5.1-8.5	138	8.7	7.3-10.3	99	16.4	13.1-20.2	39	4.1	2.9-5.6
1988	201	12.6	10.8-14.5	143	24.0	19.9-28.6	58	5.8	4.4-7.6	148	9.4	7.9-11.1	112	19.0	15.4-23.2	36	3.7	2.5-5.1
1989	192	11.6	10.0-13.4	132	21.0	17.3-25.2	60	5.9	4.5-7.7	142	8.8	7.4-10.4	99	16.0	12.8-19.7	43	4.5	3.2-6.0
1990	173	10.7	9.1-12.4	119	19.3	15.8-23.4	54	5.6	4.2-7.3	135	8.2	6.8-9.7	86	14.0	11.0-17.5	49	4.8	3.5-6.4
1991	154	9.1	7.7-10.7	102	16.5	13.2-20.3	52	4.9	3.6-6.4	128	7.7	6.4-9.1	85	13.7	10.7-17.2	43	4.1	3.0-5.6
1992	169	10.0	8.5-11.7	110	17.1	13.9-20.9	59	5.7	4.4-7.4	110	6.4	5.2-7.7	73	11.0	8.5-14.1	37	3.6	2.5-4.9
1993	183	10.5	9.0-12.1	121	17.7	14.5-21.4	62	5.9	4.5-7.6	138	8.1	6.8-9.6	95	15.2	12.0-18.9	43	4.1	2.9-5.5
1994	175	10.2	8.7-11.8	115	18.2	14.7-22.2	60	5.7	4.3-7.3	133	7.7	6.4-9.2	88	13.7	10.8-17.1	45	4.3	3.1-5.7
1995	142	8.1	6.8-9.6	94	13.8	11.0-17.1	48	4.6	3.4-6.1	124	7.2	6.0-8.7	84	13.0	10.2-16.3	40	3.9	2.8-5.3
1996	138	8.0	6.7-9.5	78	11.5	9.0-14.5	60	5.7	4.3-7.3	123	7.1	5.8-8.4	80	12.1	9.4-15.2	43	4.0	2.9-5.4
1997	144	8.0	6.7-9.4	100	14.0	11.3-17.2	44	4.1	3.0-5.5	112	6.5	5.3-7.8	75	11.4	8.8-14.4	37	3.5	2.5-4.8
1998	128	6.8	5.7-8.1	88	12.2	9.7-15.1	40	3.5	2.5-4.8	95	5.2	4.2-6.3	59	8.4	6.3-11.0	36	3.3	2.3-4.5
1999	130	6.7	5.6-8.0	88	12.2	9.7-15.2	42	3.5	2.5-4.7	100	5.1	4.2-6.3	70	9.4	7.2-11.9	30	2.5	1.7-3.6
2000	134	7.2	6.0-8.5	67	9.6	7.3-12.3	67	5.8	4.5-7.4	112	5.9	4.8-7.1	65	9.0	6.8-11.5	47	4.0	2.9-5.3
2001	121	6.0	4.9-7.1	83	10.6	8.4-13.3	38	3.1	2.2-4.2	91	4.6	3.7-5.7	67	9.2	7.0-11.8	24	2.0	1.3-3.0
2002	117	5.8	4.8-7.0	69	9.1	7.0-11.6	48	3.9	2.9-5.2	81	4.2	3.3-5.2	47	6.7	4.8-8.9	34	2.8	1.9-3.9
2003	129	6.4	5.3-7.6	81	10.9	8.5-13.7	48	3.8	2.8-5.1	91	4.3	3.5-5.3	54	6.4	4.8-8.4	37	2.9	2.1-4.0
2004	123	5.8	4.8-7.0	77	9.2	7.2-11.6	46	3.7	2.7-4.9	96	4.7	3.8-5.7	54	6.6	4.9-8.7	42	3.4	2.4-4.5
2005	107	5.0	4.0-6.0	56	6.5	4.9-8.5	51	3.9	2.9-5.1	72	3.4	2.7-4.3	43	5.1	3.7-7.0	29	2.3	1.5-3.3
2006	92	4.5	3.6-5.5	64	8.2	6.2-10.6	28	2.3	1.5-3.3	89	4.3	3.4-5.3	52	6.9	5.1-9.2	37	2.8	2.0-3.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Esophagus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	91	4.1	3.3-5.1	64	7.5	5.7-9.7	27	2.0	1.3-3.0	65	2.9	2.2-3.7	44	4.9	3.5-6.6	21	1.6	1.0-2.4
2008	111	5.0	4.1-6.1	75	9.2	7.2-11.7	36	2.6	1.8-3.6	84	3.9	3.1-4.8	52	6.5	4.8-8.6	32	2.4	1.6-3.4
2009	104	4.6	3.8-5.6	68	8.0	6.1-10.2	36	2.7	1.9-3.7	74	3.3	2.5-4.1	58	6.6	4.9-8.7	16	1.2	0.7-1.9
2010	85	3.6	2.9-4.5	49	5.4	3.9-7.2	36	2.6	1.8-3.6	67	3.0	2.3-3.8	42	5.4	3.8-7.3	25	1.8	1.1-2.7
2011	107	4.5	3.7-5.5	72	7.9	6.1-10.1	35	2.5	1.7-3.5	66	2.8	2.1-3.6	45	4.7	3.4-6.4	21	1.5	0.9-2.3
2012	85	3.5	2.8-4.4	52	5.9	4.3-7.8	33	2.1	1.5-3.0	56	2.2	1.6-2.9	37	3.7	2.6-5.2	19	1.2	0.7-1.9
2013	80	3.3	2.6-4.1	55	6.2	4.6-8.1	25	1.7	1.1-2.5	49	2.1	1.5-2.7	29	3.1	2.0-4.5	20	1.4	0.8-2.2
2014	92	3.7	2.9-4.5	68	7.0	5.4-9.0	24	1.6	1.0-2.4	61	2.5	1.9-3.3	39	4.3	3.0-6.0	22	1.5	0.9-2.3
2015	87	3.3	2.6-4.0	52	5.2	3.9-6.9	35	2.2	1.5-3.0	49	2.0	1.4-2.6	29	3.1	2.0-4.4	20	1.3	0.8-2.0
2016	101	3.7	3.0-4.5	65	6.2	4.7-8.0	36	2.2	1.5-3.1	58	2.1	1.6-2.7	38	3.6	2.5-5.0	20	1.2	0.7-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Stomach

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	178	17.4	14.7-20.5	100	24.4	19.1-30.5	78	13.1	10.2-16.6	148	14.3	11.9-17.0	88	22.6	17.4-28.6	60	9.5	7.1-12.3
1977	213	20.3	17.4-23.4	115	27.7	22.1-34.0	98	15.6	12.4-19.2	161	15.8	13.2-18.7	93	23.3	18.1-29.4	68	11.2	8.5-14.5
1978	176	15.9	13.4-18.7	91	18.8	14.8-23.5	85	13.7	10.7-17.1	159	14.9	12.5-17.7	81	18.6	14.2-23.7	78	12.6	9.8-15.9
1979	186	17.5	14.8-20.4	112	26.6	21.2-32.7	74	12.0	9.2-15.2	138	12.6	10.4-15.0	76	17.1	13.0-21.9	62	9.7	7.3-12.6
1980	238	20.8	18.0-23.9	132	29.5	24.0-35.6	106	15.6	12.6-19.0	161	14.9	12.5-17.6	92	22.3	17.5-28.0	69	10.5	8.1-13.5
1981	221	18.6	16.0-21.4	133	28.5	23.2-34.6	88	12.6	10.0-15.7	178	15.1	12.8-17.7	111	24.6	19.7-30.3	67	9.6	7.3-12.3
1982	231	19.3	16.7-22.1	130	29.9	24.3-36.1	101	13.5	10.9-16.5	167	15.1	12.7-17.7	106	27.8	22.1-34.3	61	8.4	6.3-10.9
1983	222	17.3	15.0-19.9	112	22.9	18.4-27.9	110	14.0	11.4-17.0	173	13.5	11.5-15.8	96	19.8	15.7-24.6	77	9.8	7.7-12.4
1984	201	15.3	13.2-17.8	113	24.5	19.7-30.0	88	10.5	8.3-13.0	134	10.3	8.6-12.3	73	16.4	12.4-21.0	61	7.3	5.5-9.4
1985	224	17.0	14.7-19.5	115	22.4	18.1-27.3	109	13.6	11.1-16.5	138	10.4	8.6-12.3	72	15.0	11.3-19.3	66	7.8	6.0-10.0
1986	257	18.4	16.1-21.0	152	30.8	25.5-36.7	105	11.9	9.7-14.5	165	12.4	10.5-14.6	96	19.5	15.3-24.3	69	8.5	6.6-10.8
1987	243	17.4	15.2-19.8	139	24.9	20.6-29.8	104	12.6	10.2-15.3	154	11.5	9.7-13.6	80	15.8	12.1-20.2	74	9.1	7.1-11.4
1988	234	16.3	14.2-18.6	131	24.2	19.9-29.1	103	11.8	9.5-14.3	169	11.9	10.1-14.0	98	18.4	14.6-22.8	71	8.2	6.4-10.4
1989	235	16.6	14.4-19.0	119	21.3	17.3-25.9	116	13.5	11.1-16.3	159	11.2	9.4-13.1	82	15.3	11.9-19.3	77	8.7	6.8-10.9
1990	248	16.6	14.5-18.9	129	23.3	19.1-28.1	119	12.8	10.6-15.4	163	11.6	9.8-13.7	82	15.4	11.9-19.5	81	9.3	7.4-11.6
1991	226	14.8	12.8-16.9	118	20.3	16.5-24.6	108	11.5	9.4-13.9	151	10.0	8.4-11.8	86	15.1	11.8-19.0	65	7.1	5.4-9.1
1992	252	16.3	14.3-18.5	127	21.0	17.2-25.4	125	13.2	10.9-15.8	168	11.5	9.8-13.5	78	13.9	10.7-17.7	90	9.9	7.9-12.1
1993	248	15.6	13.7-17.7	122	20.8	17.0-25.2	126	12.7	10.5-15.1	149	9.5	8.0-11.2	69	12.1	9.2-15.5	80	8.1	6.4-10.1
1994	284	18.1	16.0-20.4	142	24.0	19.9-28.5	142	14.5	12.2-17.1	189	12.2	10.5-14.1	102	17.6	14.1-21.7	87	9.0	7.2-11.1
1995	240	14.9	13.0-16.9	123	20.1	16.4-24.2	117	11.6	9.6-13.9	160	10.0	8.5-11.8	71	11.3	8.7-14.4	89	8.9	7.2-11.0
1996	260	15.6	13.7-17.6	143	23.1	19.2-27.4	117	11.2	9.2-13.4	180	11.2	9.6-12.9	96	16.5	13.1-20.4	84	8.2	6.5-10.2
1997	258	15.7	13.8-17.8	139	24.1	19.9-28.7	119	11.4	9.4-13.6	157	9.2	7.8-10.7	86	13.9	10.9-17.3	71	6.6	5.2-8.4
1998	253	14.6	12.8-16.5	127	19.9	16.3-23.9	126	11.6	9.7-13.8	149	8.8	7.4-10.3	76	12.3	9.5-15.5	73	6.8	5.3-8.5
1999	237	13.3	11.7-15.2	121	18.8	15.4-22.7	116	10.3	8.5-12.3	145	8.3	7.0-9.8	72	11.8	9.1-15.1	73	6.5	5.1-8.2
2000	261	14.5	12.8-16.4	131	20.6	17.0-24.7	130	11.3	9.5-13.5	148	8.1	6.8-9.6	85	13.2	10.4-16.6	63	5.4	4.2-6.9
2001	245	13.5	11.9-15.3	117	17.4	14.2-21.0	128	11.2	9.3-13.3	131	7.4	6.2-8.8	69	10.6	8.1-13.5	62	5.5	4.2-7.0
2002	246	12.8	11.2-14.6	117	17.0	13.8-20.6	129	10.6	8.8-12.6	138	7.3	6.1-8.7	70	11.0	8.4-14.0	68	5.6	4.3-7.1
2003	258	13.3	11.7-15.1	125	17.4	14.3-20.9	133	10.9	9.1-13.0	136	7.0	5.9-8.4	65	8.6	6.6-11.1	71	5.9	4.6-7.4
2004	242	12.4	10.9-14.1	120	17.3	14.2-20.8	122	9.8	8.1-11.7	135	7.0	5.9-8.3	64	9.3	7.0-11.9	71	5.7	4.5-7.2
2005	306	15.6	13.9-17.5	165	23.4	19.8-27.5	141	11.3	9.5-13.3	148	7.6	6.4-8.9	83	12.5	9.7-15.6	65	5.2	4.0-6.6
2006	269	13.3	11.7-15.0	143	19.2	16.0-22.7	126	9.8	8.2-11.7	143	7.4	6.2-8.7	79	11.6	9.1-14.6	64	5.1	3.9-6.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Stomach

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	238	11.5	10.1-13.1	120	16.0	13.1-19.3	118	9.2	7.6-11.0	127	6.3	5.2-7.5	57	7.1	5.3-9.3	70	5.5	4.3-7.0
2008	228	11.1	9.7-12.6	113	15.6	12.7-18.8	115	8.8	7.2-10.6	123	5.9	4.9-7.1	60	8.0	6.0-10.5	63	4.9	3.7-6.2
2009	265	12.5	11.0-14.2	149	18.4	15.4-21.8	116	8.9	7.3-10.7	116	5.7	4.7-6.8	62	8.6	6.5-11.1	54	4.1	3.1-5.4
2010	262	12.1	10.6-13.6	132	16.1	13.3-19.2	130	9.5	7.9-11.3	113	5.4	4.4-6.5	60	8.1	6.1-10.5	53	3.9	2.9-5.2
2011	264	11.6	10.3-13.2	126	15.0	12.4-18.0	138	9.7	8.1-11.5	128	5.9	4.9-7.0	73	9.2	7.2-11.7	55	3.9	2.9-5.1
2012	280	11.9	10.5-13.4	152	16.9	14.2-19.9	128	8.8	7.4-10.5	115	5.1	4.2-6.1	56	6.2	4.6-8.2	59	4.2	3.2-5.4
2013	278	11.6	10.2-13.0	151	16.6	13.9-19.6	127	8.6	7.1-10.2	120	5.1	4.2-6.1	60	6.9	5.2-8.9	60	4.1	3.1-5.3
2014	262	10.6	9.3-12.0	136	14.4	12.0-17.2	126	8.3	6.9-9.9	134	5.6	4.7-6.7	73	7.8	6.0-9.8	61	4.3	3.3-5.5
2015	278	10.8	9.6-12.2	144	14.4	12.1-17.1	134	8.6	7.2-10.3	96	3.8	3.1-4.7	52	5.6	4.1-7.4	44	2.7	2.0-3.7
2016	283	10.9	9.6-12.3	156	15.6	13.1-18.3	127	8.2	6.8-9.8	159	6.1	5.2-7.1	100	9.8	7.9-12.0	59	3.7	2.8-4.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Colorectal

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	481	46.5	42.0-51.2	209	52.5	44.6-61.1	272	42.8	37.4-48.6	302	31.5	27.6-35.6	133	37.1	30.2-45.1	169	28.3	23.9-33.3
1977	590	59.1	53.9-64.6	238	64.2	55.2-74.2	352	56.1	49.9-62.9	279	28.4	24.8-32.2	123	34.7	28.1-42.1	156	24.8	20.8-29.3
1978	609	60.3	55.1-65.8	247	65.2	56.1-75.2	362	57.7	51.5-64.4	364	37.3	33.2-41.8	148	39.8	32.8-47.7	216	35.8	30.8-41.3
1979	640	59.8	54.9-65.1	266	67.6	58.6-77.3	374	55.8	49.9-62.1	355	33.9	30.1-37.9	161	42.8	35.6-50.8	194	29.2	25.0-33.9
1980	668	58.1	53.4-63.0	294	67.3	58.8-76.5	374	52.9	47.4-58.9	356	32.8	29.2-36.7	146	36.7	30.2-44.0	210	30.9	26.6-35.6
1981	659	56.0	51.5-60.8	278	64.3	56.0-73.4	381	51.7	46.4-57.5	370	33.7	30.1-37.6	140	35.8	29.4-43.1	230	32.9	28.6-37.7
1982	680	55.3	50.9-59.9	312	67.2	59.2-76.0	368	48.3	43.3-53.7	355	29.4	26.2-32.9	160	34.3	28.6-40.7	195	26.4	22.6-30.5
1983	671	51.6	47.5-55.9	297	64.0	56.1-72.7	374	45.6	40.9-50.6	344	26.8	23.9-30.0	157	31.6	26.3-37.5	187	23.9	20.4-27.7
1984	700	53.4	49.3-57.7	317	62.7	55.3-70.8	383	47.7	42.9-52.9	380	30.0	26.9-33.4	168	36.9	30.9-43.6	212	26.6	23.0-30.6
1985	771	57.0	52.8-61.4	346	71.0	62.9-79.8	425	49.8	45.0-54.9	376	29.4	26.3-32.7	152	33.4	27.7-39.8	224	27.5	23.9-31.5
1986	800	56.9	52.8-61.2	356	69.5	61.7-77.9	444	50.4	45.7-55.5	355	26.2	23.4-29.2	159	32.4	27.1-38.4	196	22.9	19.7-26.4
1987	715	49.8	46.0-53.7	309	58.6	51.6-66.3	406	45.0	40.6-49.7	354	25.6	22.9-28.6	151	29.8	24.7-35.6	203	23.3	20.1-26.9
1988	736	50.9	47.1-54.9	312	59.1	52.1-66.7	424	46.5	42.1-51.3	351	25.1	22.4-28.0	157	31.4	26.1-37.2	194	21.8	18.8-25.2
1989	758	51.3	47.5-55.2	361	67.9	60.4-76.0	397	42.6	38.4-47.1	342	23.2	20.7-25.9	163	30.9	25.9-36.6	179	19.1	16.3-22.2
1990	768	52.0	48.2-55.9	323	61.0	53.9-68.7	445	47.1	42.7-51.8	333	23.6	21.1-26.4	148	30.3	25.1-36.2	185	20.4	17.5-23.6
1991	836	56.1	52.2-60.2	348	64.0	56.8-71.8	488	51.7	47.1-56.6	369	25.9	23.2-28.8	152	31.5	26.2-37.5	217	23.4	20.3-26.7
1992	794	51.3	47.7-55.2	312	56.4	49.7-63.7	482	49.1	44.8-53.8	361	23.9	21.4-26.6	151	29.8	24.8-35.4	210	21.4	18.6-24.6
1993	819	52.2	48.6-56.0	367	64.1	57.1-71.7	452	45.7	41.5-50.1	345	22.6	20.2-25.2	141	25.7	21.2-30.7	204	21.0	18.2-24.1
1994	867	54.3	50.6-58.1	367	63.6	56.7-71.0	500	49.5	45.2-54.0	381	24.6	22.1-27.3	176	30.9	26.2-36.2	205	20.9	18.1-24.0
1995	850	51.3	47.8-54.9	377	62.1	55.5-69.3	473	45.3	41.3-49.6	409	25.8	23.3-28.5	170	28.8	24.3-33.7	239	23.8	20.8-27.0
1996	920	54.8	51.3-58.6	391	63.7	57.0-70.9	529	50.1	45.9-54.6	421	26.2	23.7-28.8	203	36.0	30.9-41.7	218	21.1	18.3-24.1
1997	971	57.4	53.8-61.2	433	70.8	63.7-78.3	538	50.2	46.0-54.7	422	25.7	23.2-28.3	193	34.1	29.0-39.6	229	21.6	18.9-24.6
1998	951	54.8	51.3-58.5	389	60.8	54.5-67.6	562	51.2	47.1-55.7	384	22.7	20.4-25.1	161	26.4	22.2-31.2	223	20.7	18.1-23.6
1999	1029	56.7	53.2-60.3	409	61.2	55.1-67.9	620	54.1	49.9-58.6	444	25.2	22.9-27.7	182	29.3	24.9-34.1	262	23.2	20.5-26.2
2000	927	49.3	46.1-52.7	398	58.9	52.9-65.4	529	44.5	40.8-48.5	388	21.5	19.4-23.8	165	25.1	21.1-29.5	223	19.4	16.9-22.1
2001	994	52.1	48.8-55.5	410	59.2	53.2-65.6	584	48.6	44.7-52.7	417	22.6	20.4-24.9	175	27.7	23.5-32.4	242	20.4	17.9-23.1
2002	974	49.6	46.5-52.9	412	55.6	50.1-61.6	562	46.0	42.2-50.0	388	20.8	18.7-23.0	179	27.5	23.3-32.1	209	17.4	15.1-19.9
2003	1059	53.1	49.9-56.4	463	63.7	57.7-70.1	596	47.7	43.9-51.7	409	21.4	19.4-23.6	191	28.6	24.5-33.2	218	17.8	15.5-20.3
2004	1056	51.8	48.6-55.0	429	56.9	51.4-62.9	627	49.2	45.4-53.3	381	19.6	17.7-21.7	179	25.6	21.8-29.8	202	16.3	14.1-18.7
2005	1018	49.1	46.1-52.3	451	57.7	52.2-63.6	567	44.2	40.6-48.0	358	18.2	16.4-20.2	153	21.7	18.2-25.6	205	16.3	14.1-18.7
2006	1079	51.7	48.6-55.0	440	57.2	51.6-63.1	639	49.2	45.4-53.2	381	19.0	17.1-21.0	159	22.7	19.1-26.6	222	17.2	15.0-19.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Colorectal

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	1066	50.0	47.0-53.1	495	60.2	54.7-66.0	571	43.5	40.0-47.2	370	17.9	16.1-19.9	162	21.2	17.9-24.9	208	16.1	14.0-18.5
2008	1053	48.8	45.8-51.8	473	57.4	52.0-63.1	580	43.5	40.0-47.3	367	17.7	15.9-19.6	170	22.1	18.7-25.9	197	15.1	13.0-17.3
2009	1049	47.2	44.4-50.2	494	57.0	51.9-62.6	555	40.8	37.4-44.3	362	17.3	15.5-19.2	170	22.3	18.9-26.1	192	14.5	12.5-16.7
2010	1016	45.2	42.4-48.2	459	54.4	49.3-59.9	557	40.5	37.2-44.1	355	16.2	14.5-18.0	183	22.9	19.6-26.7	172	12.3	10.5-14.3
2011	1038	44.3	41.6-47.2	488	53.8	48.9-59.0	550	38.2	35.1-41.6	357	16.0	14.4-17.8	143	17.4	14.5-20.6	214	15.2	13.2-17.4
2012	997	41.7	39.1-44.4	474	51.4	46.7-56.5	523	35.8	32.8-39.1	413	17.7	16.0-19.5	201	22.6	19.4-26.1	212	14.5	12.6-16.7
2013	1018	41.2	38.7-43.9	470	49.0	44.5-53.8	548	36.4	33.4-39.7	344	14.4	12.9-16.0	152	16.5	13.8-19.4	192	12.9	11.1-14.9
2014	1066	42.4	39.8-45.0	487	49.3	44.8-54.1	579	38.3	35.2-41.6	362	14.6	13.1-16.3	167	17.9	15.1-20.9	195	12.6	10.9-14.6
2015	982	37.8	35.4-40.3	481	47.5	43.1-52.1	501	31.9	29.1-34.9	309	12.3	10.9-13.7	133	13.8	11.4-16.4	176	11.2	9.6-13.0
2016	1069	40.2	37.8-42.8	520	49.7	45.4-54.4	549	34.3	31.4-37.3	391	14.9	13.4-16.5	172	17.1	14.5-19.9	219	13.4	11.7-15.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Colon excluding rectum

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	361	34.9	31.1-39.1	151	39.3	32.5-47.0	210	32.5	27.9-37.6	244	25.5	22.0-29.2	105	29.9	23.6-37.2	139	23.1	19.2-27.6
1977	427	44.0	39.4-48.8	155	43.7	36.1-52.2	272	44.3	38.7-50.3	217	22.5	19.4-26.0	85	25.2	19.5-31.8	132	21.2	17.4-25.4
1978	462	47.0	42.4-52.0	181	51.1	42.9-60.3	281	45.2	39.7-51.2	304	31.0	27.3-35.1	120	33.0	26.5-40.3	184	30.1	25.6-35.1
1979	501	47.1	42.7-51.8	209	53.0	45.1-61.8	292	43.9	38.7-49.6	296	28.1	24.7-31.7	130	34.9	28.4-42.3	166	24.6	20.8-28.9
1980	496	43.9	39.8-48.3	213	49.8	42.4-57.8	283	40.6	35.7-45.9	295	27.5	24.2-31.2	117	29.9	24.0-36.6	178	26.5	22.5-30.9
1981	477	41.8	37.9-46.1	200	48.1	40.8-56.2	277	38.7	34.1-43.8	323	29.6	26.2-33.2	119	30.4	24.6-37.1	204	29.4	25.3-33.9
1982	502	41.4	37.6-45.4	225	50.0	42.9-57.7	277	36.6	32.2-41.3	295	24.6	21.7-27.8	130	28.4	23.2-34.3	165	22.3	18.9-26.1
1983	502	38.5	35.0-42.3	224	49.0	42.1-56.7	278	33.6	29.6-37.9	284	22.3	19.6-25.2	126	25.8	21.0-31.2	158	20.1	16.9-23.6
1984	512	39.7	36.1-43.5	224	44.4	38.2-51.2	288	36.5	32.3-41.1	331	26.0	23.1-29.1	147	32.0	26.4-38.2	184	23.0	19.7-26.7
1985	572	43.2	39.5-47.1	245	52.2	45.2-59.9	327	38.7	34.5-43.3	316	24.5	21.7-27.5	123	27.1	21.9-32.9	193	23.5	20.2-27.1
1986	596	42.7	39.2-46.5	261	52.2	45.4-59.6	335	38.1	34.0-42.5	312	23.0	20.4-25.8	141	28.9	23.9-34.7	171	19.9	16.9-23.2
1987	547	38.1	34.8-41.6	232	43.4	37.5-50.0	315	35.0	31.1-39.2	308	22.2	19.7-25.0	128	25.2	20.5-30.4	180	20.6	17.6-24.0
1988	571	40.0	36.6-43.5	245	47.0	40.7-53.8	326	36.2	32.3-40.5	311	22.0	19.5-24.7	137	26.9	22.1-32.4	174	19.5	16.6-22.6
1989	595	40.6	37.3-44.2	278	52.9	46.3-60.1	317	34.3	30.6-38.4	305	20.8	18.4-23.3	143	27.4	22.7-32.8	162	17.3	14.7-20.2
1990	589	40.2	36.9-43.7	242	46.5	40.3-53.3	347	36.9	33.0-41.0	293	20.8	18.4-23.5	128	26.2	21.4-31.6	165	18.2	15.5-21.3
1991	638	43.2	39.8-46.9	269	49.9	43.6-56.9	369	39.4	35.4-43.8	335	23.5	20.9-26.2	138	28.4	23.4-34.1	197	21.2	18.3-24.4
1992	610	39.7	36.5-43.1	229	42.6	36.7-49.1	381	38.7	34.9-42.9	318	21.2	18.9-23.8	132	26.9	22.1-32.3	186	19.0	16.3-21.9
1993	615	39.9	36.7-43.2	266	47.9	41.8-54.6	349	35.6	31.9-39.6	306	20.0	17.7-22.4	124	22.4	18.3-27.2	182	18.7	16.0-21.6
1994	690	43.6	40.3-47.0	277	48.8	42.7-55.4	413	41.0	37.1-45.2	334	21.6	19.3-24.2	155	27.5	23.0-32.6	179	18.3	15.6-21.2
1995	654	40.0	36.9-43.3	286	48.4	42.6-54.8	368	35.5	31.9-39.3	375	23.7	21.3-26.3	158	26.8	22.5-31.7	217	21.6	18.8-24.7
1996	701	41.9	38.8-45.2	303	50.2	44.2-56.7	398	37.7	34.1-41.6	370	23.1	20.8-25.7	178	32.2	27.3-37.6	192	18.6	16.0-21.4
1997	750	44.3	41.1-47.6	323	53.4	47.2-60.0	427	39.7	36.0-43.7	364	22.2	19.9-24.6	164	29.5	24.8-34.7	200	18.9	16.3-21.7
1998	731	42.4	39.3-45.7	292	46.8	41.2-52.9	439	40.1	36.4-44.0	341	20.1	17.9-22.4	142	23.4	19.4-27.8	199	18.4	15.9-21.1
1999	782	43.5	40.4-46.7	293	45.0	39.7-50.8	489	42.8	39.1-46.8	394	22.5	20.3-24.8	156	25.6	21.5-30.2	238	21.1	18.5-24.0
2000	701	37.8	35.0-40.8	289	43.9	38.7-49.6	412	34.9	31.6-38.4	349	19.3	17.3-21.5	146	22.5	18.8-26.7	203	17.6	15.3-20.2
2001	737	39.1	36.3-42.1	302	44.7	39.5-50.4	435	36.4	33.1-40.0	377	20.4	18.4-22.6	156	24.7	20.7-29.1	221	18.7	16.3-21.3
2002	746	38.3	35.5-41.2	295	39.7	35.1-44.8	451	37.1	33.7-40.7	350	18.8	16.8-20.9	159	24.5	20.6-28.8	191	15.9	13.7-18.3
2003	795	40.3	37.5-43.2	332	47.1	41.8-52.7	463	37.1	33.8-40.6	353	18.6	16.7-20.7	160	24.5	20.6-28.8	193	15.8	13.6-18.2
2004	786	38.9	36.2-41.8	313	42.2	37.3-47.4	473	37.4	34.0-40.9	328	17.0	15.2-19.0	150	21.5	18.0-25.4	178	14.4	12.4-16.7
2005	774	37.7	35.0-40.5	328	42.7	38.0-47.9	446	34.9	31.7-38.3	312	16.0	14.3-17.9	129	18.4	15.2-22.1	183	14.6	12.5-16.9
2006	798	38.7	36.0-41.5	322	42.5	37.8-47.7	476	37.0	33.7-40.5	332	16.6	14.8-18.5	139	20.1	16.7-23.9	193	15.0	13.0-17.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Colon excluding rectum

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	784	37.1	34.5-39.9	353	44.2	39.5-49.3	431	32.8	29.8-36.1	324	15.7	14.0-17.6	138	18.1	15.0-21.6	186	14.4	12.4-16.7
2008	798	37.5	34.9-40.2	353	43.9	39.2-49.0	445	33.7	30.6-37.0	315	15.2	13.5-17.0	142	18.6	15.5-22.1	173	13.2	11.3-15.3
2009	761	34.8	32.4-37.5	340	40.6	36.2-45.4	421	31.0	28.1-34.1	317	15.3	13.6-17.1	149	19.9	16.7-23.5	168	12.8	10.9-14.9
2010	748	33.8	31.3-36.3	322	38.9	34.5-43.6	426	31.2	28.3-34.4	305	14.0	12.4-15.7	157	20.0	16.9-23.6	148	10.6	8.9-12.5
2011	767	33.1	30.7-35.6	355	39.9	35.7-44.5	412	28.7	26.0-31.7	300	13.6	12.1-15.2	116	14.5	11.8-17.4	184	13.1	11.2-15.1
2012	731	31.0	28.7-33.4	329	36.9	32.8-41.2	402	27.7	25.1-30.6	339	14.6	13.1-16.3	156	17.7	14.9-20.9	183	12.6	10.8-14.6
2013	730	29.8	27.7-32.1	332	35.3	31.4-39.5	398	26.5	24.0-29.3	297	12.5	11.1-14.1	123	13.5	11.1-16.2	174	11.7	10.0-13.7
2014	790	31.7	29.5-34.0	349	36.0	32.2-40.2	441	29.3	26.6-32.3	296	11.9	10.5-13.3	136	14.4	12.0-17.2	160	10.3	8.7-12.0
2015	693	26.9	24.9-29.0	323	32.7	29.1-36.6	370	23.4	21.0-26.0	263	10.4	9.2-11.8	114	11.7	9.6-14.2	149	9.5	8.0-11.2
2016	764	28.9	26.8-31.0	355	34.1	30.5-38.0	409	25.6	23.2-28.3	299	11.3	10.1-12.7	122	12.1	9.9-14.5	177	10.8	9.3-12.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Rectum & rectosigmoid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	120	11.5	9.3-14.1	58	13.2	9.6-17.6	62	10.3	7.6-13.4	58	6.0	4.4-8.0	28	7.2	4.4-11.0	30	5.2	3.3-7.6
1977	163	15.1	12.7-17.8	83	20.6	15.9-26.1	80	11.9	9.2-15.0	62	5.8	4.4-7.6	38	9.4	6.5-13.2	24	3.6	2.2-5.5
1978	147	13.2	11.0-15.8	66	14.1	10.5-18.5	81	12.5	9.7-15.7	60	6.3	4.6-8.3	28	6.8	4.3-10.1	32	5.7	3.7-8.2
1979	139	12.7	10.5-15.3	57	14.5	10.6-19.3	82	11.9	9.3-14.9	59	5.8	4.3-7.6	31	7.9	5.1-11.5	28	4.6	2.9-6.7
1980	172	14.2	12.0-16.6	81	17.5	13.5-22.3	91	12.3	9.8-15.2	61	5.2	3.9-6.8	29	6.8	4.3-10.0	32	4.4	2.9-6.2
1981	182	14.1	12.0-16.5	78	16.2	12.4-20.7	104	13.0	10.5-15.9	47	4.2	3.0-5.6	21	5.4	3.1-8.6	26	3.6	2.3-5.2
1982	178	13.9	11.8-16.2	87	17.3	13.5-21.6	91	11.7	9.3-14.5	60	4.8	3.6-6.3	30	5.9	3.8-8.7	30	4.1	2.7-5.9
1983	169	13.1	11.1-15.4	73	15.0	11.3-19.3	96	12.0	9.7-14.8	60	4.6	3.4-6.0	31	5.8	3.7-8.5	29	3.8	2.5-5.5
1984	188	13.7	11.7-15.9	93	18.3	14.4-22.9	95	11.2	9.0-13.7	49	4.0	2.9-5.4	21	5.0	2.9-7.7	28	3.6	2.3-5.2
1985	199	13.8	11.9-16.0	101	18.9	15.0-23.4	98	11.1	8.9-13.6	60	4.9	3.6-6.3	29	6.4	4.0-9.4	31	4.0	2.7-5.7
1986	204	14.2	12.2-16.4	95	17.3	13.7-21.5	109	12.4	10.1-15.0	43	3.2	2.3-4.3	18	3.5	1.9-5.7	25	3.0	1.9-4.4
1987	168	11.6	9.9-13.6	77	15.2	11.6-19.4	91	9.9	7.9-12.3	46	3.4	2.4-4.6	23	4.7	2.8-7.2	23	2.7	1.7-4.1
1988	165	11.0	9.3-12.9	67	12.1	9.1-15.7	98	10.3	8.3-12.6	40	3.1	2.2-4.3	20	4.4	2.6-6.9	20	2.4	1.4-3.7
1989	163	10.6	9.0-12.5	83	15.0	11.7-19.0	80	8.2	6.5-10.3	37	2.4	1.7-3.4	20	3.5	2.0-5.6	17	1.8	1.0-2.9
1990	179	11.8	10.1-13.7	81	14.5	11.2-18.3	98	10.3	8.3-12.5	40	2.8	2.0-3.8	20	4.1	2.3-6.6	20	2.2	1.3-3.4
1991	198	12.9	11.1-14.9	79	14.1	10.9-17.8	119	12.2	10.1-14.7	34	2.4	1.6-3.4	14	3.1	1.6-5.3	20	2.1	1.3-3.3
1992	184	11.7	10.0-13.5	83	13.8	10.7-17.4	101	10.4	8.5-12.7	43	2.7	1.9-3.6	19	2.9	1.7-4.6	24	2.5	1.6-3.7
1993	204	12.3	10.6-14.2	101	16.2	12.9-20.0	103	10.1	8.2-12.2	39	2.6	1.8-3.6	17	3.2	1.7-5.3	22	2.3	1.4-3.5
1994	177	10.7	9.2-12.5	90	14.8	11.7-18.5	87	8.4	6.7-10.4	47	3.0	2.2-4.0	21	3.4	2.0-5.3	26	2.6	1.7-3.8
1995	196	11.2	9.7-13.0	91	13.7	10.8-17.1	105	9.8	8.0-11.9	34	2.2	1.5-3.0	12	1.9	1.0-3.4	22	2.2	1.4-3.3
1996	219	12.9	11.2-14.8	88	13.6	10.7-16.9	131	12.3	10.3-14.7	51	3.0	2.2-4.0	25	3.8	2.4-5.8	26	2.5	1.6-3.6
1997	221	13.2	11.4-15.0	110	17.4	14.1-21.3	111	10.5	8.7-12.7	58	3.5	2.6-4.5	29	4.6	3.0-6.8	29	2.7	1.8-3.9
1998	220	12.4	10.8-14.2	97	14.0	11.2-17.3	123	11.2	9.3-13.3	43	2.6	1.9-3.5	19	3.1	1.8-4.9	24	2.3	1.5-3.4
1999	247	13.2	11.6-15.0	116	16.2	13.3-19.7	131	11.3	9.5-13.5	50	2.7	2.0-3.6	26	3.7	2.3-5.5	24	2.1	1.4-3.2
2000	226	11.5	10.0-13.1	109	15.0	12.1-18.2	117	9.7	8.0-11.6	39	2.1	1.5-2.9	19	2.6	1.5-4.0	20	1.8	1.1-2.7
2001	257	12.9	11.4-14.6	108	14.5	11.7-17.7	149	12.1	10.3-14.3	40	2.1	1.5-2.9	19	3.0	1.7-4.7	21	1.7	1.1-2.6
2002	228	11.4	9.9-13.0	117	15.9	13.0-19.2	111	8.9	7.3-10.7	38	2.0	1.4-2.7	20	3.0	1.7-4.7	18	1.5	0.9-2.4
2003	264	12.8	11.3-14.5	131	16.6	13.7-19.9	133	10.6	8.8-12.5	56	2.8	2.1-3.6	31	4.1	2.8-5.9	25	2.0	1.3-2.9
2004	270	12.8	11.3-14.5	116	14.8	12.1-17.9	154	11.9	10.1-13.9	53	2.6	1.9-3.4	29	4.1	2.6-5.9	24	1.9	1.2-2.8
2005	244	11.5	10.0-13.0	123	15.0	12.3-18.0	121	9.3	7.7-11.2	46	2.2	1.6-3.0	24	3.3	2.0-4.9	22	1.7	1.1-2.6
2006	281	13.0	11.5-14.7	118	14.6	12.0-17.6	163	12.2	10.4-14.3	49	2.4	1.8-3.2	20	2.6	1.5-4.0	29	2.2	1.5-3.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Rectum & rectosigmoid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	282	12.8	11.4-14.5	142	16.0	13.3-18.9	140	10.6	8.9-12.6	46	2.2	1.6-2.9	24	3.1	2.0-4.7	22	1.7	1.0-2.5
2008	255	11.3	9.9-12.8	120	13.4	11.0-16.2	135	9.8	8.2-11.7	52	2.5	1.9-3.3	28	3.5	2.3-5.2	24	1.9	1.2-2.8
2009	288	12.4	11.0-13.9	154	16.4	13.8-19.3	134	9.8	8.2-11.6	45	2.0	1.4-2.7	21	2.4	1.4-3.7	24	1.7	1.1-2.5
2010	268	11.5	10.1-13.0	137	15.5	12.9-18.5	131	9.3	7.7-11.0	50	2.2	1.6-2.9	26	2.9	1.8-4.3	24	1.7	1.1-2.5
2011	271	11.2	9.9-12.7	133	13.9	11.5-16.6	138	9.5	8.0-11.3	57	2.4	1.8-3.2	27	2.9	1.9-4.3	30	2.1	1.4-3.0
2012	266	10.7	9.4-12.0	145	14.6	12.2-17.3	121	8.1	6.7-9.7	74	3.1	2.4-3.9	45	4.9	3.5-6.6	29	2.0	1.3-2.9
2013	288	11.4	10.1-12.8	138	13.7	11.4-16.3	150	9.9	8.4-11.6	47	1.9	1.4-2.5	29	3.0	1.9-4.3	18	1.2	0.7-1.8
2014	276	10.7	9.4-12.0	138	13.3	11.1-15.8	138	9.0	7.5-10.6	66	2.8	2.1-3.5	31	3.4	2.3-4.9	35	2.3	1.6-3.3
2015	289	10.9	9.7-12.3	158	14.8	12.5-17.4	131	8.5	7.1-10.1	46	1.8	1.3-2.4	19	2.0	1.2-3.2	27	1.7	1.1-2.5
2016	305	11.4	10.1-12.8	165	15.6	13.2-18.3	140	8.6	7.2-10.2	92	3.5	2.8-4.3	50	5.0	3.7-6.6	42	2.6	1.9-3.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Liver / intrahepatic bile duct

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	63	5.9	4.4-7.7	41	9.9	6.7-13.8	22	3.4	2.0-5.3	64	6.3	4.7-8.2	43	11.1	7.7-15.4	21	3.5	2.0-5.4
1977	61	5.1	3.8-6.8	42	9.0	6.1-12.7	19	2.8	1.6-4.5	80	6.8	5.3-8.6	54	12.4	8.9-16.7	26	3.5	2.3-5.3
1978	81	7.0	5.4-8.8	46	10.0	7.1-13.6	35	5.0	3.4-7.1	78	6.6	5.1-8.3	40	8.0	5.5-11.3	38	5.5	3.8-7.7
1979	62	5.1	3.8-6.6	36	7.8	5.1-11.1	26	3.4	2.2-5.1	83	7.3	5.7-9.2	44	10.6	7.4-14.6	39	5.5	3.8-7.6
1980	67	5.2	4.0-6.7	45	8.9	6.3-12.3	22	3.0	1.8-4.6	69	5.5	4.2-7.0	43	8.8	6.1-12.3	26	3.5	2.2-5.1
1981	66	5.0	3.8-6.4	43	8.3	5.7-11.4	23	3.0	1.9-4.5	68	5.4	4.2-7.0	41	7.8	5.4-10.9	27	3.8	2.5-5.6
1982	77	5.6	4.4-7.0	53	9.6	7.0-12.7	24	3.1	1.9-4.6	75	5.7	4.4-7.2	48	8.6	6.2-11.6	27	3.7	2.4-5.4
1983	77	5.7	4.4-7.2	48	9.4	6.6-12.8	29	3.5	2.3-5.0	97	7.3	5.8-9.0	58	11.1	8.2-14.7	39	4.9	3.5-6.8
1984	86	6.2	4.9-7.8	58	12.0	8.7-16.0	28	3.1	2.0-4.6	94	6.8	5.4-8.5	54	11.7	8.4-15.6	40	4.3	3.1-6.0
1985	84	5.6	4.4-7.0	57	10.4	7.7-13.8	27	2.8	1.8-4.1	93	6.6	5.2-8.1	55	10.3	7.5-13.7	38	4.4	3.1-6.1
1986	79	5.5	4.3-6.9	53	9.9	7.2-13.2	26	3.0	1.9-4.4	90	6.3	5.0-7.8	57	10.5	7.7-13.8	33	3.8	2.6-5.4
1987	84	5.4	4.3-6.8	48	8.1	5.7-11.1	36	3.8	2.6-5.3	92	6.6	5.3-8.2	45	9.4	6.6-12.8	47	5.3	3.9-7.1
1988	76	4.9	3.8-6.1	49	8.2	5.8-11.1	27	2.9	1.9-4.2	79	5.2	4.1-6.5	46	7.8	5.5-10.6	33	3.6	2.5-5.1
1989	82	5.5	4.3-6.9	45	8.3	5.8-11.4	37	4.0	2.8-5.5	90	5.8	4.6-7.2	49	8.0	5.7-10.7	41	4.4	3.1-6.0
1990	100	6.3	5.1-7.7	60	9.4	7.0-12.4	40	4.2	3.0-5.8	93	6.1	4.8-7.5	46	8.1	5.8-11.1	47	4.9	3.5-6.5
1991	103	6.3	5.1-7.6	66	10.9	8.1-14.1	37	3.6	2.5-5.0	102	6.6	5.3-8.0	57	10.1	7.4-13.4	45	4.6	3.3-6.2
1992	109	6.8	5.6-8.3	68	11.6	8.8-14.9	41	4.1	3.0-5.6	95	6.0	4.8-7.3	54	9.3	6.9-12.3	41	4.1	2.9-5.5
1993	103	6.2	5.0-7.5	71	11.2	8.5-14.3	32	3.2	2.2-4.5	110	6.8	5.5-8.2	68	10.7	8.1-13.7	42	4.4	3.1-5.9
1994	115	6.7	5.5-8.1	65	9.4	7.1-12.2	50	4.8	3.6-6.4	92	5.7	4.6-7.1	57	9.3	6.9-12.2	35	3.6	2.5-4.9
1995	113	6.3	5.2-7.6	70	10.1	7.7-13.0	43	4.0	2.9-5.4	100	5.7	4.6-7.0	56	8.2	6.0-10.9	44	4.1	3.0-5.6
1996	130	7.4	6.1-8.8	95	14.2	11.3-17.6	35	3.3	2.3-4.6	107	6.3	5.1-7.6	65	10.2	7.7-13.2	42	4.0	2.9-5.4
1997	123	6.7	5.6-8.0	77	10.6	8.2-13.4	46	4.3	3.1-5.7	96	5.4	4.3-6.6	58	8.3	6.2-11.0	38	3.5	2.5-4.8
1998	136	7.3	6.1-8.6	94	12.6	10.1-15.6	42	3.8	2.7-5.1	112	6.1	5.0-7.3	75	10.3	7.9-13.0	37	3.4	2.4-4.6
1999	151	7.9	6.6-9.2	89	11.7	9.3-14.4	62	5.3	4.1-6.8	87	4.6	3.7-5.7	51	6.7	4.9-8.8	36	3.2	2.2-4.4
2000	154	7.6	6.5-9.0	101	12.0	9.7-14.7	53	4.5	3.3-5.8	118	6.1	5.0-7.3	77	10.4	8.1-13.2	41	3.5	2.5-4.7
2001	146	7.1	6.0-8.4	110	13.5	10.9-16.4	36	3.0	2.1-4.2	106	5.3	4.4-6.5	65	8.3	6.3-10.6	41	3.4	2.5-4.7
2002	143	7.0	5.9-8.2	97	12.0	9.6-14.7	46	3.8	2.8-5.1	103	5.2	4.3-6.4	64	8.3	6.3-10.8	39	3.3	2.3-4.5
2003	168	7.7	6.6-9.0	120	13.7	11.2-16.5	48	3.8	2.8-5.0	112	5.4	4.5-6.6	72	9.2	7.1-11.8	40	3.2	2.3-4.3
2004	208	9.7	8.4-11.1	146	16.7	14.0-19.8	62	4.9	3.8-6.3	147	7.1	6.0-8.3	89	10.4	8.3-12.9	58	4.7	3.6-6.1
2005	202	8.9	7.7-10.2	153	16.7	14.0-19.7	49	3.7	2.7-4.9	123	5.9	4.9-7.0	81	10.0	7.9-12.6	42	3.3	2.4-4.5
2006	189	8.5	7.3-9.8	138	15.8	13.1-18.7	51	3.9	2.9-5.1	136	6.1	5.1-7.3	105	11.9	9.6-14.6	31	2.4	1.6-3.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Liver / intrahepatic bile duct

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	226	9.9	8.7-11.3	166	18.4	15.6-21.6	60	4.5	3.4-5.8	144	6.5	5.4-7.7	96	11.0	8.8-13.6	48	3.6	2.7-4.8
2008	230	9.9	8.6-11.2	171	17.8	15.1-20.8	59	4.4	3.4-5.7	139	6.1	5.1-7.2	98	10.8	8.7-13.3	41	3.0	2.1-4.0
2009	260	11.1	9.8-12.6	177	19.0	16.1-22.2	83	6.0	4.7-7.4	141	6.2	5.2-7.3	99	10.8	8.7-13.2	42	3.1	2.2-4.2
2010	285	11.6	10.3-13.1	189	19.0	16.3-22.0	96	6.7	5.4-8.2	171	7.2	6.2-8.4	113	12.2	9.9-14.8	58	4.0	3.0-5.2
2011	281	11.4	10.1-12.8	188	18.4	15.8-21.4	93	6.5	5.3-8.0	164	6.9	5.8-8.0	99	9.9	7.9-12.1	65	4.7	3.6-6.0
2012	326	12.9	11.5-14.4	230	22.5	19.6-25.8	96	6.5	5.2-7.9	166	6.8	5.7-7.9	114	11.3	9.3-13.7	52	3.6	2.7-4.8
2013	307	11.5	10.2-12.9	209	19.0	16.4-21.9	98	6.3	5.1-7.7	171	6.7	5.7-7.8	112	11.1	9.0-13.4	59	3.8	2.9-4.9
2014	294	10.6	9.4-11.9	192	16.4	14.1-19.0	102	6.4	5.2-7.9	172	6.5	5.5-7.6	106	9.7	7.9-11.8	66	4.3	3.3-5.5
2015	289	10.4	9.2-11.7	207	18.2	15.8-21.0	82	5.0	4.0-6.2	160	5.8	4.9-6.8	99	8.8	7.1-10.7	61	3.8	2.9-4.9
2016	294	10.3	9.1-11.6	215	18.5	16.0-21.2	79	4.6	3.7-5.8	187	6.7	5.8-7.8	134	12.1	10.1-14.4	53	3.2	2.4-4.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Pancreas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	149	14.6	12.2-17.4	64	14.7	10.9-19.2	85	14.3	11.2-18.0	151	13.9	11.6-16.5	70	15.9	11.9-20.6	81	12.6	9.8-15.9
1977	145	13.4	11.1-16.1	72	17.1	12.7-22.4	73	11.3	8.7-14.4	125	11.6	9.5-14.1	65	15.5	11.4-20.6	60	9.3	6.9-12.2
1978	148	13.5	11.2-16.1	61	12.8	9.4-17.0	87	13.6	10.7-17.0	134	11.7	9.7-14.1	55	11.8	8.5-15.8	79	11.6	9.0-14.6
1979	159	14.1	11.8-16.7	72	14.8	11.3-19.0	87	13.3	10.5-16.6	146	13.3	11.0-15.8	56	11.1	8.2-14.6	90	14.2	11.2-17.6
1980	158	13.5	11.3-16.0	85	18.3	14.2-23.3	73	10.5	8.1-13.3	130	11.4	9.4-13.7	69	15.9	11.9-20.8	61	8.8	6.6-11.4
1981	181	14.9	12.6-17.4	100	21.1	16.7-26.1	81	11.1	8.6-13.9	147	12.0	10.0-14.3	82	17.1	13.1-21.9	65	8.9	6.8-11.5
1982	172	14.3	12.1-16.7	69	15.8	11.9-20.5	103	13.5	10.9-16.5	160	13.5	11.4-15.9	67	15.6	11.7-20.2	93	12.4	9.9-15.3
1983	196	14.7	12.6-17.0	89	16.7	13.1-20.9	107	13.3	10.8-16.1	178	13.6	11.6-15.9	90	18.3	14.3-22.9	88	10.9	8.7-13.5
1984	188	14.3	12.2-16.6	84	16.4	12.7-20.8	104	13.0	10.5-15.8	165	12.9	10.9-15.1	81	17.0	13.1-21.5	84	10.5	8.3-13.1
1985	175	12.8	10.9-14.9	92	16.7	13.2-20.8	83	10.2	8.1-12.7	149	11.3	9.4-13.3	75	14.4	11.0-18.5	74	9.3	7.2-11.7
1986	190	13.7	11.8-15.9	83	15.1	11.8-19.0	107	12.6	10.3-15.3	158	11.7	9.9-13.7	70	13.0	9.9-16.7	88	10.7	8.5-13.2
1987	214	14.8	12.8-17.0	104	17.4	13.9-21.4	110	12.7	10.4-15.4	175	12.2	10.3-14.2	80	14.9	11.5-19.0	95	10.6	8.5-13.0
1988	209	14.2	12.3-16.4	97	17.8	14.1-22.1	112	12.2	10.0-14.7	149	10.2	8.5-12.0	72	12.2	9.3-15.6	77	8.7	6.8-10.9
1989	195	13.2	11.3-15.2	79	14.0	10.9-17.7	116	12.5	10.3-15.0	180	12.0	10.3-14.0	75	13.1	10.1-16.7	105	11.2	9.1-13.6
1990	206	14.1	12.2-16.2	87	16.2	12.6-20.3	119	12.9	10.6-15.5	171	11.6	9.8-13.5	70	12.8	9.7-16.5	101	10.8	8.8-13.2
1991	179	11.9	10.2-13.8	81	14.4	11.2-18.3	98	10.5	8.5-12.8	165	10.5	8.9-12.3	82	13.6	10.6-17.1	83	8.6	6.8-10.7
1992	191	12.4	10.6-14.3	71	12.7	9.6-16.4	120	12.2	10.1-14.6	143	9.7	8.1-11.4	53	10.6	7.7-14.1	90	9.4	7.5-11.5
1993	193	12.2	10.5-14.1	84	14.4	11.2-18.2	109	11.0	9.0-13.3	148	9.5	8.0-11.2	58	11.0	8.1-14.6	90	9.0	7.2-11.1
1994	217	13.5	11.8-15.5	88	14.6	11.5-18.3	129	12.8	10.7-15.2	191	12.0	10.3-13.9	71	11.9	9.1-15.3	120	11.9	9.8-14.2
1995	187	11.7	10.1-13.6	82	13.9	10.8-17.4	105	10.4	8.5-12.6	173	10.7	9.1-12.4	75	12.8	9.8-16.3	98	9.5	7.7-11.6
1996	218	13.1	11.3-14.9	79	12.4	9.6-15.6	139	13.2	11.1-15.6	167	10.1	8.6-11.8	62	9.7	7.3-12.6	105	10.1	8.3-12.3
1997	224	13.2	11.5-15.0	96	14.0	11.2-17.2	128	12.2	10.1-14.5	177	10.7	9.1-12.4	72	11.0	8.4-13.9	105	10.1	8.2-12.2
1998	210	12.2	10.6-14.0	81	12.0	9.4-15.1	129	12.0	10.0-14.2	171	10.0	8.5-11.7	69	10.3	7.8-13.1	102	9.6	7.8-11.6
1999	249	14.2	12.4-16.1	97	15.3	12.2-18.9	152	13.5	11.4-15.9	196	11.3	9.7-13.0	71	11.0	8.4-14.0	125	11.2	9.3-13.4
2000	237	13.2	11.5-15.0	78	11.8	9.2-14.9	159	13.8	11.7-16.1	184	10.2	8.7-11.8	78	12.0	9.3-15.0	106	9.1	7.5-11.0
2001	244	12.9	11.3-14.7	91	12.8	10.2-15.9	153	12.9	10.9-15.1	200	10.8	9.4-12.5	67	9.7	7.4-12.4	133	11.3	9.5-13.4
2002	245	12.9	11.3-14.7	94	12.5	10.0-15.5	151	12.8	10.8-15.0	207	10.8	9.4-12.4	92	12.4	9.8-15.3	115	9.6	8.0-11.6
2003	227	11.7	10.2-13.4	109	15.2	12.3-18.5	118	9.6	8.0-11.5	171	9.0	7.7-10.5	74	10.3	7.9-13.1	97	8.1	6.6-9.9
2004	241	12.1	10.6-13.7	102	13.2	10.6-16.2	139	11.2	9.4-13.2	210	10.5	9.1-12.0	91	12.4	9.8-15.3	119	9.4	7.8-11.3
2005	221	10.7	9.3-12.3	89	11.9	9.4-14.8	132	10.2	8.6-12.2	185	9.2	7.9-10.7	76	9.7	7.6-12.3	109	8.7	7.2-10.5
2006	291	14.0	12.4-15.7	118	14.1	11.6-17.1	173	13.6	11.6-15.8	194	9.5	8.2-10.9	83	10.8	8.5-13.5	111	8.7	7.1-10.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Pancreas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	283	13.6	12.0-15.3	131	16.4	13.5-19.6	152	11.8	10.0-13.9	224	10.9	9.5-12.5	102	13.1	10.5-16.1	122	9.6	8.0-11.5
2008	253	12.0	10.5-13.6	91	11.0	8.8-13.6	162	12.3	10.4-14.3	218	10.4	9.1-11.9	88	11.1	8.8-13.9	130	9.9	8.3-11.8
2009	266	12.3	10.8-13.9	120	14.9	12.2-17.9	146	10.7	9.0-12.6	209	9.8	8.5-11.3	91	11.6	9.2-14.3	118	8.7	7.2-10.5
2010	307	13.8	12.3-15.5	136	15.8	13.1-18.8	171	12.5	10.6-14.5	208	9.7	8.4-11.2	88	10.5	8.3-13.1	120	9.1	7.6-10.9
2011	353	15.4	13.8-17.1	150	16.5	13.9-19.5	203	14.5	12.6-16.7	262	11.7	10.3-13.2	117	13.9	11.4-16.7	145	10.3	8.7-12.2
2012	339	14.2	12.7-15.9	130	14.1	11.7-16.9	209	14.4	12.5-16.6	257	10.9	9.5-12.3	86	9.3	7.3-11.5	171	11.9	10.1-13.8
2013	376	15.4	13.8-17.0	166	17.2	14.6-20.2	210	14.1	12.2-16.2	246	10.1	8.9-11.5	115	12.5	10.2-15.1	131	8.8	7.3-10.4
2014	337	13.2	11.8-14.8	148	14.1	11.8-16.6	189	12.4	10.7-14.4	246	9.8	8.6-11.1	103	10.2	8.2-12.4	143	9.5	8.0-11.2
2015	391	15.2	13.7-16.9	185	18.1	15.5-21.0	206	13.3	11.5-15.3	269	10.7	9.4-12.1	123	12.3	10.2-14.8	146	9.5	8.0-11.2
2016	367	13.8	12.4-15.3	172	16.2	13.8-18.9	195	12.0	10.4-13.9	295	11.1	9.8-12.5	145	13.7	11.5-16.3	150	9.3	7.8-10.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Larynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	79	6.0	4.7-7.6	63	11.7	8.8-15.1	16	2.1	1.1-3.5	50	4.2	3.0-5.7	39	8.3	5.6-11.8	11	1.6	0.7-2.9
1977	90	7.1	5.6-8.8	72	14.8	11.1-19.1	18	2.3	1.4-3.7	43	3.5	2.4-4.9	32	7.4	4.5-11.2	11	1.3	0.7-2.4
1978	99	7.3	5.9-9.0	81	15.1	11.7-19.1	18	2.2	1.3-3.4	64	4.5	3.5-5.8	55	9.7	7.2-12.9	9	1.0	0.5-2.0
1979	106	7.8	6.3-9.4	82	15.4	12.0-19.3	24	2.9	1.8-4.3	58	4.1	3.1-5.4	47	8.7	6.2-11.7	11	1.2	0.6-2.2
1980	100	6.9	5.6-8.5	74	12.8	9.9-16.4	26	3.0	1.9-4.5	52	3.7	2.8-4.9	34	6.6	4.4-9.4	18	2.0	1.2-3.2
1981	93	6.3	5.1-7.8	75	12.7	9.8-16.2	18	2.1	1.2-3.4	55	3.6	2.7-4.8	35	6.1	4.2-8.5	20	2.1	1.3-3.3
1982	85	5.6	4.4-6.9	73	12.0	9.4-15.3	12	1.4	0.7-2.4	39	2.7	1.9-3.7	31	5.6	3.7-8.1	8	0.9	0.4-1.8
1983	114	7.5	6.1-9.1	94	15.7	12.4-19.5	20	2.3	1.4-3.5	46	3.1	2.2-4.1	38	6.3	4.3-8.9	8	0.9	0.4-1.9
1984	111	7.1	5.8-8.6	81	13.0	10.2-16.2	30	3.2	2.2-4.6	62	3.9	3.0-5.0	49	7.6	5.6-10.2	13	1.4	0.7-2.4
1985	146	9.1	7.6-10.7	117	19.1	15.6-23.0	29	2.8	1.9-4.0	52	3.4	2.5-4.5	42	6.9	4.9-9.4	10	1.1	0.5-2.1
1986	140	8.9	7.4-10.5	109	18.3	14.8-22.4	31	3.2	2.1-4.5	65	4.2	3.3-5.4	52	8.8	6.5-11.8	13	1.4	0.8-2.4
1987	123	7.6	6.2-9.0	98	15.4	12.3-19.0	25	2.6	1.7-3.8	66	4.2	3.2-5.4	50	8.3	6.0-11.2	16	1.7	0.9-2.7
1988	126	7.2	6.0-8.6	95	13.9	11.1-17.2	31	2.9	2.0-4.2	71	4.3	3.3-5.4	58	9.4	7.0-12.3	13	1.2	0.6-2.0
1989	140	8.1	6.8-9.6	90	13.6	10.8-17.0	50	4.8	3.5-6.3	60	3.6	2.7-4.7	43	7.1	5.0-9.8	17	1.6	0.9-2.6
1990	147	8.5	7.1-10.0	110	16.6	13.5-20.3	37	3.6	2.5-4.9	46	2.8	2.0-3.7	35	5.7	3.9-8.2	11	1.1	0.5-1.9
1991	122	6.9	5.7-8.3	92	14.1	11.2-17.4	30	2.7	1.9-3.9	55	3.2	2.4-4.2	40	6.0	4.2-8.3	15	1.4	0.8-2.4
1992	150	8.2	6.9-9.7	122	17.2	14.2-20.7	28	2.5	1.6-3.6	65	3.7	2.9-4.8	51	7.9	5.7-10.5	14	1.3	0.7-2.2
1993	116	6.7	5.5-8.0	89	13.8	10.9-17.1	27	2.5	1.7-3.7	67	3.9	3.0-4.9	48	7.7	5.5-10.4	19	1.8	1.1-2.8
1994	108	6.0	4.9-7.2	77	11.1	8.7-14.1	31	2.8	1.9-4.0	60	3.4	2.5-4.3	42	6.0	4.2-8.2	18	1.7	1.0-2.6
1995	118	6.4	5.3-7.7	84	11.7	9.2-14.6	34	3.1	2.1-4.3	72	4.1	3.2-5.1	55	8.5	6.2-11.2	17	1.5	0.9-2.5
1996	113	6.0	4.9-7.2	92	12.6	10.0-15.6	21	1.8	1.1-2.8	63	3.5	2.7-4.5	54	8.2	6.0-10.8	9	0.8	0.4-1.5
1997	100	5.2	4.2-6.4	74	9.9	7.7-12.5	26	2.3	1.5-3.3	56	3.0	2.2-3.9	45	6.2	4.4-8.4	11	1.0	0.5-1.7
1998	110	5.7	4.7-6.9	75	10.5	8.1-13.3	35	2.9	2.0-4.1	52	2.7	2.0-3.6	43	5.8	4.2-7.9	9	0.8	0.4-1.5
1999	100	5.1	4.1-6.2	74	10.1	7.8-12.8	26	2.2	1.4-3.2	45	2.3	1.7-3.1	31	4.2	2.8-6.1	14	1.2	0.6-2.0
2000	96	4.9	3.9-5.9	69	9.3	7.1-11.9	27	2.2	1.5-3.3	54	2.8	2.1-3.7	42	5.7	4.0-7.8	12	1.1	0.6-1.9
2001	87	4.2	3.3-5.2	65	7.8	6.0-10.0	22	1.8	1.1-2.7	35	1.8	1.2-2.5	27	3.7	2.4-5.4	8	0.6	0.3-1.3
2002	87	4.1	3.3-5.1	61	7.3	5.5-9.4	26	2.1	1.3-3.0	46	2.2	1.6-3.0	34	4.2	2.9-5.9	12	1.0	0.5-1.7
2003	105	4.9	4.0-5.9	77	9.5	7.4-12.0	28	2.2	1.4-3.1	37	1.8	1.3-2.5	31	4.0	2.7-5.7	6	0.5	0.2-1.0
2004	91	4.0	3.2-5.0	74	8.3	6.5-10.5	17	1.2	0.7-2.0	47	2.3	1.6-3.0	38	4.9	3.4-6.8	9	0.7	0.3-1.4
2005	91	4.2	3.4-5.2	68	8.4	6.4-10.7	23	1.7	1.1-2.6	37	1.7	1.2-2.4	31	3.7	2.5-5.3	6	0.5	0.2-1.0
2006	77	3.3	2.6-4.1	57	6.2	4.6-8.1	20	1.4	0.8-2.1	39	1.9	1.3-2.5	32	4.0	2.7-5.7	7	0.5	0.2-1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Larynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	94	4.1	3.3-5.1	76	8.5	6.7-10.7	18	1.3	0.8-2.1	43	1.9	1.4-2.6	35	4.2	2.9-5.9	8	0.6	0.2-1.2
2008	106	4.8	3.9-5.8	69	8.2	6.3-10.4	37	2.7	1.9-3.8	41	1.8	1.3-2.5	26	3.3	2.1-4.8	15	1.1	0.6-1.8
2009	87	3.7	3.0-4.6	68	7.8	6.0-10.0	19	1.3	0.8-2.1	39	1.7	1.2-2.3	29	3.4	2.2-5.0	10	0.7	0.3-1.3
2010	85	3.5	2.8-4.4	63	6.8	5.1-8.7	22	1.5	1.0-2.3	31	1.4	0.9-2.0	25	3.1	1.9-4.6	6	0.4	0.2-0.9
2011	87	3.4	2.7-4.3	61	6.2	4.7-8.1	26	1.6	1.1-2.4	32	1.4	0.9-2.0	21	2.5	1.5-3.8	11	0.8	0.4-1.4
2012	72	2.8	2.2-3.6	53	5.4	4.0-7.1	19	1.2	0.7-2.0	31	1.3	0.9-1.8	24	2.7	1.7-4.1	7	0.4	0.2-0.9
2013	83	3.2	2.5-4.0	66	6.7	5.1-8.6	17	1.1	0.6-1.7	32	1.3	0.9-1.8	22	2.2	1.4-3.4	10	0.6	0.3-1.2
2014	89	3.4	2.8-4.3	67	6.4	4.9-8.1	22	1.5	0.9-2.3	37	1.4	1.0-2.0	31	3.1	2.1-4.5	6	0.4	0.1-0.8
2015	108	3.9	3.2-4.8	86	8.3	6.6-10.3	22	1.3	0.8-2.0	22	0.9	0.5-1.3	14	1.5	0.8-2.5	8	0.5	0.2-1.0
2016	71	2.6	2.0-3.3	53	5.1	3.8-6.8	18	1.1	0.6-1.7	35	1.3	0.9-1.8	27	2.7	1.7-4.0	8	0.4	0.2-0.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Lung and bronchus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	667	55.6	51.1-60.3	494	102.8	92.9-113.4	173	25.0	21.2-29.3	524	43.5	39.6-47.8	395	82.5	73.5-92.2	129	18.3	15.1-22.0
1977	672	54.0	49.7-58.5	487	102.3	92.3-113.1	185	23.8	20.4-27.7	554	44.5	40.6-48.6	408	85.5	76.3-95.3	146	18.9	15.8-22.4
1978	718	57.3	52.9-62.0	527	105.2	95.5-115.5	191	26.7	22.8-31.0	611	48.4	44.4-52.6	444	88.5	79.7-97.9	167	22.7	19.2-26.7
1979	787	62.4	57.8-67.2	567	114.8	104.5-125.8	220	29.7	25.7-34.1	638	51.3	47.1-55.8	473	98.1	88.4-108.6	165	22.5	19.0-26.4
1980	790	59.1	54.8-63.6	572	110.7	100.9-121.2	218	27.1	23.5-31.1	642	48.6	44.7-52.7	474	91.9	83.0-101.4	168	21.6	18.3-25.3
1981	828	59.3	55.2-63.7	574	107.1	97.8-117.1	254	29.6	26.0-33.6	699	50.9	47.0-55.0	495	93.3	84.5-102.7	204	24.6	21.2-28.3
1982	851	60.4	56.2-64.8	583	107.3	98.1-117.1	268	31.3	27.5-35.4	680	48.7	45.0-52.7	470	86.3	78.0-95.1	210	25.5	22.0-29.3
1983	856	59.5	55.4-63.8	590	106.5	97.3-116.2	266	30.7	27.0-34.7	755	52.6	48.8-56.7	528	96.9	88.1-106.3	227	25.7	22.4-29.4
1984	918	61.9	57.8-66.2	616	108.6	99.5-118.2	302	33.6	29.8-37.7	705	48.2	44.5-52.0	492	89.5	81.1-98.6	213	23.6	20.5-27.1
1985	910	60.6	56.6-64.9	623	110.8	101.5-120.7	287	31.1	27.5-35.0	746	50.6	46.8-54.5	516	94.0	85.4-103.3	230	25.3	22.0-28.9
1986	934	60.6	56.6-64.8	595	103.0	94.2-112.4	339	35.7	31.9-39.8	728	47.8	44.2-51.5	468	82.2	74.3-90.8	260	27.6	24.3-31.3
1987	991	62.7	58.7-66.9	664	111.9	102.9-121.5	327	33.6	30.0-37.6	772	50.1	46.5-53.9	498	87.8	79.5-96.5	274	28.7	25.4-32.4
1988	931	58.5	54.6-62.5	594	101.4	92.8-110.6	337	34.0	30.4-37.9	760	48.7	45.2-52.4	494	86.3	78.2-95.0	266	27.4	24.1-31.0
1989	992	61.0	57.1-65.0	638	107.2	98.4-116.5	354	34.9	31.3-38.8	748	47.3	43.8-50.9	503	88.6	80.4-97.4	245	24.4	21.4-27.8
1990	1011	61.1	57.3-65.1	644	105.4	96.7-114.5	367	35.7	32.1-39.7	780	47.6	44.2-51.1	511	83.0	75.4-91.1	269	26.7	23.6-30.2
1991	983	59.6	55.8-63.5	626	102.4	94.0-111.4	357	35.0	31.4-38.9	735	44.9	41.6-48.4	486	81.1	73.5-89.2	249	24.5	21.5-27.8
1992	1014	60.2	56.5-64.1	635	102.4	93.9-111.4	379	36.4	32.7-40.2	737	45.2	41.9-48.6	438	74.1	66.8-82.0	299	29.3	26.0-32.8
1993	968	57.1	53.5-60.9	585	95.2	87.0-103.9	383	36.2	32.6-40.0	723	42.8	39.6-46.1	444	74.2	66.8-82.0	279	26.1	23.1-29.4
1994	1025	58.9	55.3-62.7	631	97.9	89.8-106.3	394	36.4	32.9-40.3	810	47.7	44.4-51.2	509	81.8	74.3-89.9	301	28.7	25.5-32.1
1995	968	55.4	51.9-59.1	585	89.9	82.3-97.9	383	35.4	31.9-39.2	811	46.8	43.5-50.2	492	77.0	69.9-84.5	319	29.5	26.4-33.0
1996	1083	60.6	57.0-64.4	633	94.3	86.6-102.4	450	41.0	37.3-45.0	828	47.1	43.8-50.5	509	77.6	70.5-85.1	319	29.5	26.3-33.0
1997	1028	57.3	53.8-61.0	579	87.0	79.7-94.9	449	40.3	36.6-44.2	783	44.6	41.5-47.9	460	71.7	64.9-79.0	323	29.4	26.3-32.8
1998	1035	55.9	52.5-59.5	573	81.6	74.7-88.9	462	40.6	37.0-44.5	783	43.4	40.3-46.6	452	67.1	60.7-73.9	331	29.7	26.6-33.1
1999	956	50.7	47.5-54.1	546	77.4	70.7-84.5	410	35.3	31.9-38.9	746	40.5	37.6-43.5	416	60.9	54.9-67.4	330	28.9	25.8-32.2
2000	993	51.4	48.2-54.7	551	76.6	70.1-83.7	442	36.9	33.5-40.5	735	38.5	35.7-41.4	423	60.0	54.1-66.3	312	26.2	23.4-29.3
2001	979	50.6	47.4-53.9	541	76.1	69.5-83.2	438	36.3	32.9-39.8	742	38.5	35.8-41.5	414	59.2	53.3-65.5	328	27.2	24.3-30.3
2002	1024	51.2	48.1-54.5	553	75.5	69.0-82.4	471	37.9	34.5-41.5	752	38.1	35.4-40.9	412	57.1	51.4-63.1	340	27.5	24.6-30.6
2003	980	48.4	45.4-51.6	499	66.1	60.2-72.5	481	38.4	35.0-42.0	725	36.1	33.5-38.9	387	52.3	47.0-58.1	338	27.0	24.2-30.1
2004	1047	50.9	47.9-54.2	552	69.9	63.9-76.3	495	39.1	35.7-42.8	696	34.0	31.5-36.7	375	48.7	43.6-54.1	321	25.3	22.6-28.2
2005	1044	50.0	46.9-53.1	560	71.9	65.8-78.5	484	37.6	34.3-41.1	728	35.4	32.9-38.1	383	49.2	44.1-54.6	345	27.3	24.4-30.3
2006	1022	48.1	45.2-51.2	529	65.2	59.5-71.3	493	37.7	34.4-41.2	712	34.4	31.9-37.1	360	46.7	41.8-52.0	352	27.4	24.6-30.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Lung and bronchus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	1036	48.2	45.2-51.2	530	65.1	59.4-71.2	506	38.4	35.1-42.0	712	33.7	31.2-36.3	382	47.9	43.0-53.2	330	25.4	22.7-28.4
2008	1056	49.0	46.0-52.1	541	66.6	60.8-72.7	515	38.9	35.6-42.4	679	32.0	29.6-34.5	344	43.0	38.3-48.0	335	25.5	22.8-28.4
2009	1079	48.8	45.9-51.9	556	66.2	60.5-72.2	523	38.5	35.2-42.0	671	30.8	28.5-33.3	365	44.4	39.7-49.4	306	22.7	20.2-25.5
2010	1084	47.5	44.7-50.5	542	62.0	56.6-67.8	542	39.0	35.8-42.5	742	33.0	30.6-35.5	387	46.4	41.7-51.6	355	25.6	23.0-28.4
2011	1160	49.9	47.0-53.0	569	65.1	59.6-70.9	591	41.3	38.0-44.8	728	31.6	29.3-34.0	371	42.3	37.9-47.0	357	25.1	22.5-27.9
2012	1126	47.0	44.2-49.9	538	59.5	54.3-65.0	588	40.0	36.8-43.4	730	31.1	28.8-33.4	384	43.7	39.2-48.5	346	23.8	21.4-26.5
2013	1034	42.3	39.7-45.0	518	55.1	50.3-60.3	516	34.7	31.7-37.8	705	29.1	26.9-31.4	365	39.8	35.7-44.4	340	22.7	20.3-25.3
2014	1102	43.6	41.0-46.3	550	56.4	51.6-61.5	552	35.7	32.8-38.9	658	26.5	24.5-28.7	338	35.9	32.0-40.1	320	20.8	18.5-23.2
2015	1084	41.1	38.6-43.6	531	51.5	47.0-56.2	553	34.7	31.9-37.8	656	25.3	23.4-27.4	320	32.5	28.9-36.4	336	21.1	18.9-23.6
2016	1072	40.1	37.7-42.6	510	48.3	44.0-52.8	562	34.7	31.9-37.8	649	24.4	22.5-26.4	345	33.2	29.7-37.1	304	18.7	16.6-21.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Melanoma of the skin

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	8	0.8	0.3-1.6	1	0.2	0.0-1.2	7	1.1	0.4-2.4	5	0.6	0.1-1.4	3	1.3	0.2-3.8	2	0.3	0.0-1.0
1977	8	0.7	0.3-1.5	3	0.7	0.1-2.1	5	0.8	0.2-1.8	4	0.4	0.1-1.0	2	0.7	0.1-2.4	2	0.2	0.0-0.9
1978	9	0.8	0.3-1.5	3	0.6	0.1-2.0	6	0.9	0.3-1.9	5	0.4	0.1-1.0	2	0.3	0.0-1.3	3	0.5	0.1-1.4
1979	17	1.2	0.7-2.0	6	1.0	0.4-2.3	11	1.4	0.6-2.5	2	0.2	0.0-0.7	1	0.3	0.0-1.6	1	0.1	0.0-0.7
1980	10	0.8	0.3-1.5	3	0.5	0.1-1.5	7	1.0	0.4-2.0	14	1.1	0.6-1.9	8	1.2	0.5-2.5	6	0.9	0.3-2.1
1981	18	1.4	0.8-2.3	5	0.8	0.3-2.0	13	1.8	0.9-3.1	14	1.1	0.5-1.9	6	0.9	0.3-2.2	8	1.1	0.4-2.3
1982	10	0.7	0.4-1.4	2	0.3	0.0-1.3	8	1.0	0.4-1.9	7	0.5	0.2-1.1	4	1.0	0.2-2.8	3	0.3	0.1-0.8
1983	10	0.7	0.3-1.3	2	0.3	0.0-1.2	8	0.9	0.4-1.8	18	1.2	0.7-2.0	13	2.1	1.1-3.6	5	0.6	0.2-1.5
1984	21	1.5	0.9-2.3	5	0.8	0.2-2.0	16	1.9	1.1-3.1	12	0.7	0.4-1.3	9	1.4	0.6-2.7	3	0.3	0.1-0.8
1985	17	1.2	0.7-2.0	9	1.4	0.5-2.9	8	1.1	0.4-2.1	12	0.9	0.4-1.5	9	1.9	0.8-3.6	3	0.3	0.1-0.9
1986	14	1.0	0.5-1.7	8	1.8	0.6-3.8	6	0.6	0.2-1.4	14	1.0	0.5-1.7	10	1.7	0.8-3.3	4	0.5	0.1-1.3
1987	12	0.7	0.4-1.3	5	0.7	0.2-1.7	7	0.7	0.3-1.5	8	0.5	0.2-1.1	3	0.5	0.1-1.4	5	0.5	0.2-1.3
1988	15	1.0	0.5-1.6	5	1.1	0.3-2.6	10	1.0	0.5-1.8	7	0.5	0.2-1.1	2	0.6	0.0-2.0	5	0.6	0.2-1.3
1989	13	1.0	0.5-1.7	4	0.8	0.2-2.1	9	1.1	0.5-2.1	12	0.8	0.4-1.4	4	0.5	0.1-1.4	8	0.9	0.4-1.8
1990	17	1.0	0.5-1.6	9	1.4	0.5-2.9	8	0.8	0.3-1.5	6	0.4	0.1-0.9	3	0.6	0.1-2.0	3	0.3	0.1-0.9
1991	21	1.2	0.7-1.8	12	2.2	1.0-4.1	9	0.7	0.3-1.4	9	0.6	0.3-1.1	6	1.0	0.3-2.4	3	0.3	0.1-0.9
1992	11	0.6	0.3-1.0	4	0.5	0.1-1.4	7	0.6	0.2-1.3	6	0.3	0.1-0.7	2	0.2	0.0-1.0	4	0.4	0.1-0.9
1993	17	1.1	0.6-1.7	11	1.6	0.7-2.9	6	0.7	0.2-1.4	5	0.3	0.1-0.7	2	0.2	0.0-1.0	3	0.3	0.1-0.8
1994	13	0.7	0.4-1.3	3	0.7	0.1-1.9	10	0.9	0.4-1.6	3	0.3	0.1-0.7	1	0.2	0.0-1.1	2	0.2	0.0-0.8
1995	19	1.1	0.7-1.8	4	0.5	0.1-1.4	15	1.4	0.8-2.4	2	0.2	0.0-0.6	0	0.0	0.0-0.5	2	0.2	0.0-0.8
1996	18	1.2	0.7-1.8	5	0.8	0.2-1.8	13	1.3	0.7-2.2	9	0.6	0.3-1.2	3	0.5	0.1-1.6	6	0.6	0.2-1.3
1997	31	1.7	1.2-2.5	14	2.0	1.1-3.5	17	1.5	0.9-2.5	7	0.4	0.2-0.9	3	0.3	0.1-1.1	4	0.4	0.1-1.0
1998	20	1.1	0.7-1.8	10	1.6	0.7-3.1	10	0.9	0.4-1.6	8	0.4	0.2-0.9	2	0.2	0.0-0.9	6	0.5	0.2-1.2
1999	8	0.4	0.2-0.9	3	0.4	0.1-1.2	5	0.4	0.1-1.0	7	0.4	0.1-0.8	4	0.7	0.1-1.8	3	0.3	0.1-0.7
2000	12	0.6	0.3-1.1	6	0.8	0.3-1.8	6	0.5	0.2-1.1	10	0.6	0.3-1.1	5	1.1	0.3-2.5	5	0.4	0.1-1.0
2001	23	1.1	0.7-1.7	9	1.0	0.4-2.0	14	1.2	0.6-1.9	2	0.1	0.0-0.4	1	0.1	0.0-0.6	1	0.1	0.0-0.5
2002	16	0.8	0.4-1.3	5	0.5	0.2-1.2	11	0.9	0.4-1.6	5	0.3	0.1-0.6	2	0.2	0.0-0.8	3	0.3	0.1-0.7
2003	12	0.6	0.3-1.0	3	0.5	0.1-1.4	9	0.7	0.3-1.3	5	0.2	0.1-0.5	1	0.1	0.0-0.6	4	0.3	0.1-0.8
2004	15	0.7	0.4-1.2	7	1.1	0.4-2.2	8	0.6	0.3-1.2	3	0.1	0.0-0.4	1	0.1	0.0-0.6	2	0.2	0.0-0.6
2005	20	0.9	0.6-1.5	7	0.8	0.3-1.6	13	1.0	0.5-1.7	4	0.2	0.0-0.5	3	0.3	0.1-0.9	1	0.1	0.0-0.5
2006	28	1.3	0.9-2.0	13	1.9	0.9-3.3	15	1.1	0.6-1.9	7	0.3	0.1-0.6	4	0.4	0.1-1.1	3	0.2	0.0-0.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Melanoma of the skin

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	32	1.5	1.0-2.1	11	1.4	0.7-2.6	21	1.6	1.0-2.5	15	0.7	0.4-1.2	6	0.9	0.3-2.0	9	0.7	0.3-1.3
2008	27	1.2	0.8-1.8	11	1.4	0.6-2.5	16	1.2	0.7-2.0	8	0.4	0.1-0.7	3	0.3	0.1-0.9	5	0.4	0.1-0.9
2009	22	0.9	0.6-1.4	9	0.9	0.4-1.7	13	0.9	0.5-1.6	4	0.2	0.0-0.4	3	0.3	0.1-0.9	1	0.1	0.0-0.4
2010	20	0.8	0.5-1.3	8	0.9	0.4-1.8	12	0.8	0.4-1.5	6	0.3	0.1-0.6	2	0.3	0.0-1.1	4	0.3	0.1-0.7
2011	31	1.3	0.9-1.9	15	1.7	0.9-2.8	16	1.2	0.7-1.9	11	0.5	0.3-0.9	5	0.6	0.2-1.5	6	0.5	0.2-1.0
2012	23	1.0	0.6-1.4	10	1.1	0.5-2.0	13	0.9	0.5-1.5	8	0.3	0.1-0.7	3	0.4	0.1-1.1	5	0.3	0.1-0.8
2013	25	1.1	0.7-1.6	8	1.0	0.4-1.9	17	1.1	0.7-1.8	9	0.3	0.1-0.6	5	0.4	0.1-1.0	4	0.2	0.1-0.7
2014	22	0.9	0.5-1.3	10	0.9	0.4-1.7	12	0.8	0.4-1.5	6	0.3	0.1-0.6	2	0.2	0.0-0.6	4	0.3	0.1-0.7
2015	20	0.8	0.5-1.2	10	1.2	0.5-2.1	10	0.6	0.3-1.2	9	0.3	0.1-0.6	5	0.5	0.1-1.2	4	0.2	0.1-0.7
2016	21	0.8	0.5-1.2	9	0.9	0.4-1.7	12	0.7	0.4-1.3	4	0.1	0.0-0.4	1	0.1	0.0-0.5	3	0.2	0.0-0.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Female breast

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	537	74.1	67.6-81.1	248	36.3	31.6-41.5
1977	544	72.5	66.2-79.2	230	31.5	27.3-36.1
1978	607	80.1	73.6-87.1	258	35.1	30.7-39.9
1979	629	80.6	74.1-87.5	229	30.1	26.2-34.5
1980	663	81.8	75.4-88.5	278	35.7	31.4-40.3
1981	650	79.4	73.2-86.1	273	34.5	30.3-39.1
1982	702	83.8	77.5-90.5	277	33.1	29.2-37.4
1983	708	81.1	75.0-87.5	299	35.1	31.0-39.4
1984	820	91.1	84.8-97.7	316	36.8	32.7-41.3
1985	811	88.3	82.1-94.8	251	27.7	24.3-31.4
1986	801	83.9	78.0-90.0	303	32.1	28.5-36.0
1987	887	89.8	83.8-96.0	312	32.1	28.6-36.0
1988	949	94.8	88.7-101.2	321	32.2	28.7-36.0
1989	904	88.8	83.0-94.9	319	32.1	28.6-35.9
1990	927	89.9	84.1-96.0	340	33.9	30.4-37.8
1991	1060	101.5	95.4-107.9	321	31.3	27.9-35.0
1992	997	94.4	88.5-100.5	330	31.7	28.3-35.3
1993	1013	93.9	88.1-100.0	314	30.4	27.1-34.0
1994	1082	98.6	92.8-104.8	380	34.7	31.2-38.4
1995	1112	99.3	93.5-105.4	403	36.7	33.2-40.5
1996	1207	106.1	100.1-112.3	381	34.1	30.7-37.7
1997	1214	104.6	98.8-110.7	375	33.1	29.8-36.7
1998	1357	115.4	109.3-121.7	369	32.0	28.8-35.4
1999	1204	99.8	94.2-105.6	375	31.8	28.7-35.2
2000	1226	99.0	93.5-104.7	336	27.7	24.8-30.9
2001	1205	95.6	90.2-101.2	395	32.1	29.0-35.4
2002	1262	98.6	93.2-104.3	344	27.6	24.7-30.7
2003	1313	101.2	95.8-106.9	349	27.4	24.6-30.5
2004	1325	100.8	95.4-106.4	348	27.0	24.2-30.0
2005	1340	101.0	95.6-106.6	392	29.7	26.8-32.8
2006	1412	105.3	99.8-110.9	352	26.7	24.0-29.7
2007	1452	107.6	102.1-113.4	334	24.7	22.1-27.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Female breast

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	1506	111.0	105.4-116.8	334	24.8	22.2-27.6
2009	1524	110.6	105.1-116.4	332	24.3	21.8-27.1
2010	1490	106.3	100.9-111.9	320	22.7	20.3-25.4
2011	1700	118.4	112.8-124.2	355	25.2	22.6-28.0
2012	1647	112.2	106.8-117.8	382	26.2	23.6-29.0
2013	1756	118.2	112.6-123.9	314	20.9	18.6-23.3
2014	1746	116.6	111.1-122.3	362	23.9	21.5-26.5
2015	1819	118.0	112.5-123.7	333	21.6	19.3-24.1
2016	1750	112.7	107.4-118.2	356	22.6	20.3-25.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Cervix uteri

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	223	28.3	24.6-32.5	101	14.6	11.8-18.0
1977	246	31.6	27.6-36.0	102	14.1	11.4-17.4
1978	233	29.3	25.5-33.5	89	12.1	9.5-15.0
1979	249	30.6	26.8-34.8	81	10.9	8.5-13.7
1980	220	26.8	23.2-30.8	90	11.5	9.1-14.2
1981	199	22.2	19.1-25.6	81	9.3	7.3-11.7
1982	220	25.3	21.9-29.0	81	9.0	7.1-11.2
1983	202	22.2	19.1-25.6	76	8.5	6.6-10.7
1984	208	22.3	19.3-25.6	64	7.3	5.6-9.4
1985	192	20.0	17.2-23.1	71	7.6	5.9-9.6
1986	216	22.7	19.7-26.0	71	7.3	5.7-9.3
1987	198	19.3	16.7-22.3	58	5.8	4.4-7.6
1988	186	18.1	15.5-21.0	90	9.0	7.2-11.0
1989	180	16.5	14.1-19.2	50	5.0	3.7-6.7
1990	235	21.2	18.5-24.1	65	6.4	4.9-8.1
1991	215	19.5	16.9-22.3	63	5.8	4.4-7.5
1992	214	18.8	16.4-21.6	64	6.0	4.6-7.6
1993	201	17.7	15.3-20.3	76	6.8	5.4-8.5
1994	216	19.2	16.7-22.0	64	5.8	4.4-7.4
1995	217	18.6	16.2-21.3	77	6.9	5.4-8.6
1996	219	18.3	15.9-20.9	59	5.1	3.8-6.5
1997	212	17.5	15.2-20.1	67	5.9	4.5-7.5
1998	206	16.4	14.2-18.8	62	5.2	4.0-6.7
1999	197	15.7	13.5-18.0	75	6.1	4.8-7.6
2000	247	19.7	17.3-22.3	72	5.7	4.5-7.2
2001	192	15.0	12.9-17.2	60	4.8	3.7-6.2
2002	198	15.2	13.1-17.5	58	4.6	3.5-5.9
2003	176	13.4	11.5-15.5	66	5.1	3.9-6.4
2004	166	12.4	10.6-14.5	50	3.8	2.8-5.1
2005	182	13.7	11.7-15.8	54	4.1	3.1-5.4
2006	173	12.9	11.1-15.0	57	4.4	3.3-5.7
2007	189	14.2	12.2-16.3	72	5.4	4.2-6.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Cervix uteri

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	164	12.1	10.3-14.2	62	4.5	3.4-5.8
2009	158	11.3	9.6-13.3	64	4.5	3.4-5.7
2010	167	12.2	10.4-14.3	47	3.3	2.4-4.4
2011	164	11.7	9.9-13.6	67	4.8	3.7-6.1
2012	159	11.3	9.6-13.3	48	3.3	2.4-4.4
2013	152	10.6	8.9-12.4	60	4.0	3.1-5.2
2014	144	10.1	8.5-11.9	62	4.2	3.2-5.4
2015	149	10.0	8.4-11.8	43	2.8	2.0-3.8
2016	156	10.5	8.9-12.3	52	3.5	2.6-4.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Corpus uterus and uterus, not otherwise specified

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	115	16.0	13.1-19.4	55	8.7	6.4-11.4
1977	107	15.2	12.3-18.5	51	7.2	5.3-9.5
1978	114	15.7	12.8-19.0	60	9.5	7.1-12.3
1979	106	13.8	11.2-16.8	52	7.3	5.3-9.6
1980	134	17.4	14.5-20.8	61	8.1	6.1-10.5
1981	113	14.1	11.5-17.1	51	6.2	4.5-8.2
1982	131	15.1	12.5-18.0	47	5.7	4.1-7.6
1983	132	15.2	12.7-18.2	61	7.3	5.6-9.5
1984	154	17.9	15.1-21.1	59	7.0	5.2-9.0
1985	164	17.8	15.1-20.8	68	7.8	6.0-10.0
1986	111	11.3	9.3-13.7	53	5.9	4.4-7.8
1987	160	16.7	14.1-19.5	44	5.0	3.6-6.7
1988	170	16.7	14.3-19.5	63	6.5	4.9-8.3
1989	147	14.2	12.0-16.8	52	4.9	3.7-6.5
1990	154	15.3	12.9-17.9	66	6.7	5.2-8.6
1991	208	20.1	17.4-23.0	60	6.0	4.6-7.8
1992	179	16.8	14.4-19.4	64	6.3	4.8-8.0
1993	205	18.9	16.4-21.7	69	6.8	5.2-8.6
1994	213	19.1	16.6-21.9	79	7.5	5.9-9.3
1995	205	18.4	16.0-21.1	59	5.5	4.2-7.1
1996	225	19.8	17.3-22.6	86	8.0	6.4-9.9
1997	253	22.2	19.5-25.1	97	8.7	7.0-10.6
1998	249	21.3	18.7-24.1	96	8.5	6.9-10.4
1999	269	22.7	20.1-25.6	90	7.8	6.2-9.6
2000	281	23.2	20.5-26.0	80	6.8	5.4-8.4
2001	304	24.2	21.6-27.1	99	8.2	6.7-10.0
2002	289	22.8	20.2-25.6	81	6.5	5.2-8.1
2003	311	24.0	21.4-26.8	104	8.2	6.7-9.9
2004	342	26.1	23.4-29.1	106	8.5	6.9-10.2
2005	352	26.7	23.9-29.6	111	8.5	7.0-10.3
2006	374	28.0	25.2-31.0	107	8.3	6.8-10.1
2007	339	25.2	22.5-28.0	95	7.3	5.9-8.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Corpus uterus and uterus, not otherwise specified

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	459	33.1	30.1-36.3	115	8.7	7.1-10.4
2009	428	30.3	27.5-33.3	120	8.8	7.2-10.5
2010	448	31.2	28.3-34.3	114	8.2	6.7-9.8
2011	465	31.7	28.9-34.8	145	10.2	8.6-12.0
2012	495	32.7	29.9-35.8	131	8.8	7.4-10.5
2013	506	32.6	29.8-35.7	125	8.1	6.7-9.7
2014	531	33.2	30.4-36.2	174	11.0	9.5-12.9
2015	553	34.0	31.2-37.0	166	10.1	8.6-11.8
2016	566	34.2	31.4-37.2	178	10.8	9.3-12.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Ovary

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	76	10.8	8.4-13.7	47	6.7	4.8-9.0
1977	91	11.8	9.3-14.6	56	7.7	5.7-10.2
1978	77	10.4	8.1-13.2	59	7.8	5.8-10.2
1979	95	12.4	9.9-15.4	47	6.0	4.4-8.1
1980	102	12.9	10.4-15.9	76	10.3	8.0-13.1
1981	91	10.5	8.4-12.9	48	5.6	4.1-7.5
1982	79	9.4	7.4-11.7	51	6.3	4.6-8.3
1983	122	13.3	10.9-16.0	72	8.5	6.6-10.8
1984	119	13.2	10.9-15.9	66	7.4	5.7-9.5
1985	112	11.1	9.1-13.5	77	8.4	6.6-10.6
1986	114	12.0	9.8-14.5	63	6.6	5.0-8.5
1987	92	9.6	7.7-11.8	55	5.7	4.3-7.5
1988	112	11.5	9.4-13.8	55	5.7	4.3-7.5
1989	95	9.0	7.3-11.1	62	6.2	4.7-7.9
1990	128	12.5	10.4-14.9	79	8.0	6.3-10.0
1991	115	11.0	9.0-13.2	72	7.0	5.4-8.8
1992	153	14.4	12.1-16.9	89	8.5	6.8-10.4
1993	115	10.4	8.5-12.5	95	9.2	7.4-11.3
1994	161	14.7	12.5-17.2	90	8.7	6.9-10.7
1995	137	12.1	10.2-14.4	99	9.2	7.4-11.2
1996	111	9.6	7.9-11.6	58	5.2	3.9-6.7
1997	147	12.6	10.6-14.9	75	6.8	5.3-8.5
1998	107	9.2	7.5-11.2	79	7.0	5.5-8.7
1999	128	10.9	9.0-12.9	74	6.4	5.0-8.1
2000	121	10.0	8.3-12.0	76	6.5	5.1-8.1
2001	129	10.4	8.7-12.4	84	6.9	5.5-8.5
2002	141	11.1	9.3-13.1	79	6.5	5.1-8.1
2003	130	10.0	8.4-11.9	75	6.0	4.7-7.5
2004	137	10.5	8.8-12.4	78	6.2	4.9-7.8
2005	134	10.3	8.6-12.2	88	6.8	5.5-8.4
2006	117	8.7	7.2-10.4	80	6.2	4.9-7.8
2007	128	9.5	7.9-11.4	81	6.1	4.8-7.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Ovary

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	134	9.8	8.2-11.6	82	5.9	4.7-7.3
2009	139	10.0	8.4-11.8	80	5.8	4.6-7.3
2010	138	10.0	8.4-11.9	88	6.3	5.1-7.8
2011	137	9.5	8.0-11.3	86	6.2	4.9-7.6
2012	155	10.6	9.0-12.4	97	6.6	5.4-8.1
2013	144	9.8	8.2-11.6	71	4.6	3.6-5.9
2014	156	10.3	8.7-12.0	84	5.5	4.4-6.8
2015	135	8.7	7.3-10.4	92	5.9	4.7-7.2
2016	138	8.7	7.3-10.3	100	6.1	5.0-7.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Prostate

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	396	118.4	105.2-132.6	144	49.8	40.7-60.0
1977	422	122.4	109.3-136.4	147	45.9	38.0-54.7
1978	464	133.4	119.8-147.9	167	54.9	45.8-64.9
1979	480	131.2	118.4-144.9	189	60.6	51.2-70.9
1980	584	156.6	142.6-171.4	218	67.4	57.8-77.9
1981	567	148.3	134.9-162.5	232	70.7	60.9-81.4
1982	542	138.8	126.1-152.3	202	59.6	50.9-69.1
1983	526	129.1	117.1-141.9	215	59.5	51.1-68.7
1984	602	142.2	129.8-155.3	219	60.5	52.1-69.7
1985	621	137.4	125.7-149.8	254	68.3	59.4-78.0
1986	639	141.1	129.3-153.5	254	66.4	57.8-75.8
1987	658	138.1	126.7-150.1	265	64.9	56.6-73.8
1988	700	151.9	139.7-164.7	210	53.0	45.6-61.2
1989	744	154.4	142.5-167.0	233	55.4	47.9-63.6
1990	834	168.0	155.8-180.9	288	69.7	61.2-78.9
1991	893	174.2	161.9-187.1	255	59.5	51.8-67.9
1992	1124	208.9	195.8-222.5	305	70.7	62.3-79.8
1993	1212	215.5	202.5-229.0	274	64.7	56.7-73.4
1994	1341	229.4	216.4-243.0	296	67.6	59.5-76.3
1995	1481	240.3	227.3-253.7	322	69.7	61.8-78.3
1996	1450	232.2	219.6-245.3	293	61.2	53.9-69.0
1997	1575	245.8	233.0-259.0	297	61.0	53.8-68.8
1998	1605	243.0	230.6-255.9	288	57.5	50.7-64.9
1999	1876	273.3	260.4-286.7	319	62.3	55.2-69.9
2000	2088	291.9	278.9-305.3	296	55.7	49.2-62.6
2001	1951	266.7	254.3-279.4	315	58.4	51.8-65.5
2002	1967	262.4	250.3-274.9	323	56.2	49.9-62.9
2003	1962	256.1	244.3-268.3	266	46.1	40.5-52.3
2004	2048	259.5	247.8-271.6	277	48.4	42.6-54.7
2005	2025	249.3	238.1-260.9	263	44.1	38.7-50.0
2006	2132	259.8	248.3-271.6	263	44.8	39.4-50.8
2007	2309	272.9	261.3-284.7	282	46.4	40.9-52.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Prostate

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	2257	256.8	245.9-268.1	262	43.2	37.9-48.9
2009	2240	248.0	237.4-259.0	282	44.7	39.4-50.3
2010	2178	232.1	222.0-242.5	268	40.2	35.4-45.5
2011	2420	247.2	237.0-257.7	281	41.2	36.4-46.4
2012	2209	219.2	209.8-229.0	271	38.1	33.5-43.0
2013	2203	211.3	202.2-220.6	269	36.6	32.2-41.3
2014	1980	184.0	175.6-192.6	283	34.8	30.7-39.3
2015	2001	179.4	171.4-187.8	259	31.1	27.3-35.2
2016	2052	180.7	172.6-189.0	285	32.9	29.1-37.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Testis

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	5	0.5	0.2-1.5	3	1.0	0.1-3.4
1977	4	0.8	0.2-2.3	2	0.2	0.0-1.2
1978	2	0.5	0.0-1.8	3	0.6	0.1-1.9
1979	6	0.7	0.2-1.8	0	0.0	0.0-0.9
1980	8	1.2	0.5-2.5	1	0.1	0.0-1.0
1981	6	0.7	0.3-1.8	2	0.3	0.0-1.3
1982	4	0.5	0.1-1.5	0	0.0	0.0-0.8
1983	5	0.8	0.2-2.0	6	1.2	0.3-2.9
1984	3	0.2	0.0-1.0	1	0.2	0.0-1.1
1985	11	1.7	0.7-3.3	1	0.1	0.0-0.8
1986	11	1.1	0.6-2.2	2	0.2	0.0-1.0
1987	8	0.7	0.3-1.5	1	0.1	0.0-0.7
1988	8	0.8	0.3-1.7	2	0.5	0.0-1.9
1989	13	1.5	0.8-2.7	1	0.1	0.0-0.8
1990	10	1.2	0.5-2.2	3	0.3	0.1-1.0
1991	7	0.9	0.3-2.0	0	0.0	0.0-0.6
1992	14	1.1	0.6-2.1	1	0.1	0.0-0.7
1993	10	1.0	0.5-2.0	2	0.2	0.0-0.8
1994	14	1.3	0.7-2.4	1	0.1	0.0-0.8
1995	5	0.4	0.1-1.1	0	0.0	0.0-0.5
1996	17	1.5	0.9-2.5	0	0.0	0.0-0.5
1997	7	0.6	0.3-1.4	1	0.1	0.0-0.7
1998	15	1.4	0.8-2.4	1	0.1	0.0-0.6
1999	13	1.2	0.7-2.2	0	0.0	0.0-0.5
2000	24	2.3	1.5-3.5	3	0.4	0.1-1.1
2001	15	1.3	0.8-2.3	1	0.2	0.0-0.9
2002	13	1.2	0.6-2.1	2	0.2	0.0-0.8
2003	18	1.7	1.0-2.7	1	0.1	0.0-0.6
2004	11	1.0	0.5-1.8	1	0.1	0.0-0.6
2005	20	1.9	1.2-3.0	2	0.2	0.0-0.7
2006	24	2.3	1.4-3.4	1	0.1	0.0-0.6
2007	21	2.0	1.2-3.1	1	0.1	0.0-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Testis

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	11	1.0	0.5-1.8	0	0.0	0.0-0.4
2009	14	1.3	0.7-2.2	1	0.1	0.0-0.6
2010	19	1.7	1.0-2.8	1	0.1	0.0-0.5
2011	20	1.8	1.1-2.8	3	0.3	0.1-0.8
2012	18	1.6	1.0-2.6	1	0.1	0.0-0.5
2013	17	1.4	0.8-2.3	0	0.0	0.0-0.4
2014	27	2.5	1.6-3.7	1	0.1	0.0-0.5
2015	25	2.1	1.4-3.2	1	0.1	0.0-0.6
2016	27	2.3	1.5-3.5	2	0.3	0.0-0.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Urinary bladder (incl. in situ)

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	117	12.5	10.1-15.2	76	20.9	15.8-26.9	41	7.5	5.2-10.3	54	6.0	4.4-8.1	32	8.2	5.3-12.0	22	4.6	2.7-7.0
1977	102	10.3	8.2-12.7	65	18.0	13.3-23.5	37	5.9	4.0-8.2	46	5.1	3.6-6.9	27	8.7	5.2-13.3	19	3.2	1.9-5.1
1978	99	9.4	7.5-11.5	64	16.5	12.3-21.6	35	5.3	3.6-7.4	44	4.7	3.3-6.4	27	8.3	5.2-12.4	17	2.8	1.6-4.5
1979	112	11.0	8.9-13.4	67	17.7	13.2-23.1	45	7.2	5.2-9.7	53	5.7	4.2-7.6	26	7.5	4.7-11.2	27	4.7	3.0-6.8
1980	141	12.5	10.4-14.9	91	21.5	16.9-26.8	50	7.3	5.3-9.8	46	4.9	3.5-6.7	25	7.3	4.4-11.1	21	3.6	2.2-5.6
1981	116	10.1	8.2-12.3	67	15.7	11.6-20.5	49	7.0	5.1-9.3	62	5.6	4.2-7.3	33	7.9	5.2-11.5	29	4.2	2.8-6.2
1982	128	10.7	8.8-12.9	68	16.1	12.1-20.7	60	7.8	5.9-10.2	57	5.0	3.7-6.5	25	6.0	3.7-9.1	32	4.4	2.9-6.3
1983	127	10.0	8.2-12.0	69	14.7	11.1-19.1	58	7.2	5.4-9.4	49	4.3	3.1-5.8	24	5.9	3.6-8.9	25	3.5	2.2-5.1
1984	112	8.4	6.8-10.1	71	14.4	11.0-18.5	41	5.0	3.5-6.8	53	4.2	3.1-5.5	31	6.1	4.0-8.9	22	3.0	1.8-4.5
1985	146	11.5	9.6-13.7	81	17.2	13.2-21.9	65	8.4	6.4-10.8	59	4.9	3.7-6.4	33	8.3	5.5-11.8	26	3.3	2.1-4.9
1986	129	10.1	8.3-12.0	85	19.9	15.5-25.0	44	5.2	3.7-7.0	48	3.9	2.9-5.3	25	6.6	4.0-9.8	23	2.8	1.7-4.2
1987	128	9.6	7.9-11.5	84	17.3	13.4-21.8	44	5.4	3.9-7.3	59	4.7	3.5-6.0	34	8.0	5.3-11.3	25	3.0	1.9-4.5
1988	134	9.9	8.2-11.8	80	15.9	12.3-20.1	54	6.5	4.9-8.5	52	3.9	2.9-5.1	31	7.5	4.8-10.9	21	2.3	1.4-3.5
1989	123	8.9	7.3-10.6	72	13.8	10.5-17.7	51	6.0	4.4-7.9	49	3.8	2.8-5.1	27	6.1	3.8-9.0	22	2.7	1.7-4.1
1990	136	9.7	8.0-11.5	85	17.7	13.8-22.1	51	5.7	4.2-7.4	50	3.7	2.7-4.9	25	5.3	3.3-7.9	25	2.8	1.8-4.2
1991	138	9.2	7.7-10.9	82	15.8	12.3-19.9	56	5.9	4.4-7.6	40	2.9	2.1-4.0	28	6.5	4.2-9.5	12	1.3	0.7-2.2
1992	153	10.0	8.4-11.8	82	15.6	12.1-19.6	71	7.2	5.6-9.1	45	3.3	2.4-4.4	21	4.9	2.9-7.6	24	2.6	1.7-3.9
1993	147	9.7	8.2-11.5	76	15.1	11.6-19.1	71	7.2	5.6-9.0	58	3.8	2.8-4.9	34	6.5	4.3-9.2	24	2.4	1.5-3.6
1994	151	9.6	8.1-11.3	79	14.4	11.2-18.1	72	7.1	5.5-9.0	57	3.9	2.9-5.1	27	5.4	3.4-8.0	30	3.1	2.1-4.5
1995	130	8.2	6.8-9.8	85	15.6	12.2-19.6	45	4.4	3.2-5.9	53	3.5	2.6-4.6	25	4.8	2.9-7.3	28	2.9	1.9-4.1
1996	149	9.2	7.7-10.8	88	15.3	12.1-19.1	61	5.9	4.5-7.6	51	3.4	2.5-4.4	28	5.5	3.5-8.0	23	2.3	1.5-3.5
1997	148	9.1	7.6-10.7	76	13.1	10.1-16.6	72	6.9	5.4-8.7	56	3.7	2.8-4.8	26	5.1	3.2-7.6	30	3.0	2.0-4.3
1998	171	10.0	8.5-11.6	99	16.0	12.8-19.6	72	6.7	5.2-8.4	63	4.0	3.1-5.1	29	5.6	3.6-8.1	34	3.3	2.3-4.6
1999	168	9.4	8.0-10.9	108	16.6	13.4-20.2	60	5.3	4.1-6.9	49	2.9	2.1-3.8	23	3.9	2.4-6.0	26	2.3	1.5-3.4
2000	165	9.3	7.9-10.8	90	14.5	11.5-17.9	75	6.6	5.2-8.2	66	3.7	2.9-4.8	36	5.8	3.9-8.1	30	2.7	1.8-3.8
2001	174	9.4	8.0-10.9	102	15.2	12.2-18.6	72	6.1	4.8-7.7	64	3.6	2.8-4.6	27	4.5	2.8-6.6	37	3.2	2.2-4.4
2002	173	9.4	8.0-10.9	98	15.2	12.2-18.7	75	6.3	4.9-7.9	47	2.6	1.9-3.5	18	2.7	1.5-4.3	29	2.5	1.7-3.6
2003	172	8.8	7.5-10.3	121	17.1	14.0-20.6	51	4.2	3.1-5.5	47	2.5	1.8-3.3	26	3.9	2.5-5.8	21	1.7	1.1-2.6
2004	190	9.8	8.4-11.3	118	17.0	13.9-20.5	72	5.9	4.6-7.4	57	3.0	2.2-3.8	29	4.2	2.7-6.2	28	2.2	1.5-3.3
2005	215	10.7	9.3-12.2	127	16.6	13.6-19.9	88	7.1	5.7-8.7	57	3.1	2.4-4.0	32	5.0	3.3-7.1	25	2.1	1.4-3.1
2006	218	10.9	9.5-12.5	144	20.0	16.7-23.7	74	5.9	4.6-7.4	49	2.6	1.9-3.4	31	4.8	3.2-6.9	18	1.5	0.9-2.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Urinary bladder (incl. in situ)

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	177	8.8	7.5-10.2	103	14.6	11.7-17.8	74	5.8	4.6-7.3	47	2.5	1.8-3.3	22	3.6	2.2-5.5	25	2.0	1.3-3.0
2008	196	9.3	8.0-10.7	129	16.3	13.5-19.6	67	5.1	4.0-6.5	49	2.4	1.8-3.2	25	3.9	2.5-5.8	24	1.8	1.1-2.6
2009	217	10.3	8.9-11.8	137	17.8	14.8-21.2	80	6.1	4.8-7.6	46	2.4	1.7-3.1	22	3.5	2.2-5.4	24	1.8	1.1-2.7
2010	210	9.6	8.3-11.0	137	16.8	14.0-20.0	73	5.4	4.2-6.8	65	3.2	2.4-4.0	34	4.9	3.3-6.9	31	2.4	1.6-3.4
2011	226	10.0	8.7-11.4	151	18.3	15.4-21.6	75	5.3	4.2-6.7	61	2.8	2.1-3.6	32	4.3	2.9-6.1	29	2.1	1.4-3.0
2012	211	9.2	7.9-10.5	129	15.6	12.9-18.7	82	5.5	4.4-6.9	65	2.9	2.2-3.7	38	4.6	3.2-6.4	27	1.9	1.2-2.8
2013	260	11.0	9.7-12.5	168	19.1	16.2-22.4	92	6.3	5.1-7.8	58	2.5	1.9-3.2	33	4.1	2.8-5.9	25	1.6	1.1-2.4
2014	246	9.9	8.7-11.2	166	17.6	14.9-20.6	80	5.2	4.1-6.5	67	2.9	2.3-3.7	39	5.0	3.5-6.9	28	1.8	1.2-2.7
2015	254	10.0	8.8-11.4	163	17.1	14.4-20.0	91	5.9	4.7-7.2	58	2.3	1.8-3.0	31	3.5	2.4-5.0	27	1.6	1.1-2.4
2016	235	9.1	7.9-10.4	161	16.7	14.1-19.6	74	4.6	3.6-5.8	71	2.8	2.2-3.6	43	4.9	3.5-6.6	28	1.7	1.1-2.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Kidney and renal pelvis

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	65	5.9	4.4-7.6	38	7.3	4.8-10.5	27	4.8	3.0-7.0	37	3.3	2.2-4.7	22	4.7	2.6-7.6	15	2.4	1.3-4.1
1977	47	3.5	2.5-4.7	33	6.5	4.2-9.6	14	1.6	0.9-2.8	22	1.7	1.0-2.7	16	2.9	1.6-4.9	6	0.9	0.3-2.0
1978	81	6.2	4.8-7.8	55	10.3	7.5-13.7	26	3.5	2.2-5.2	39	3.0	2.0-4.2	28	5.4	3.3-8.2	11	1.4	0.7-2.6
1979	58	4.2	3.2-5.6	44	8.6	6.0-11.8	14	1.5	0.8-2.6	32	2.4	1.6-3.4	24	4.4	2.7-6.6	8	1.1	0.5-2.2
1980	68	4.9	3.7-6.2	36	6.6	4.5-9.3	32	3.8	2.6-5.5	26	1.8	1.2-2.7	21	3.7	2.2-5.7	5	0.6	0.2-1.4
1981	63	4.4	3.3-5.7	43	7.3	5.2-9.9	20	2.4	1.4-3.8	25	1.6	1.0-2.3	17	2.8	1.6-4.5	8	0.8	0.3-1.6
1982	74	5.1	4.0-6.5	47	8.2	5.9-11.0	27	3.1	2.0-4.6	27	1.9	1.2-2.9	16	3.0	1.6-5.1	11	1.3	0.6-2.3
1983	88	5.9	4.7-7.3	43	7.7	5.4-10.6	45	4.9	3.6-6.6	39	2.7	1.9-3.8	23	4.1	2.5-6.4	16	1.9	1.0-3.1
1984	97	6.4	5.1-7.8	62	10.4	7.8-13.6	35	3.9	2.7-5.4	40	2.8	2.0-3.9	23	4.2	2.6-6.5	17	2.0	1.1-3.2
1985	80	5.1	3.9-6.4	45	6.9	4.9-9.4	35	3.8	2.6-5.3	42	2.8	2.0-3.8	22	3.6	2.1-5.7	20	2.3	1.4-3.5
1986	100	6.1	4.9-7.5	59	8.9	6.6-11.6	41	4.3	3.1-5.9	53	3.6	2.7-4.8	32	5.7	3.7-8.2	21	2.4	1.4-3.7
1987	111	7.0	5.7-8.5	67	11.4	8.6-14.7	44	4.5	3.2-6.1	33	2.1	1.4-3.0	20	3.4	2.0-5.4	13	1.4	0.7-2.4
1988	100	6.1	5.0-7.5	57	8.5	6.3-11.2	43	4.5	3.2-6.1	39	2.5	1.8-3.4	25	4.1	2.6-6.1	14	1.5	0.8-2.5
1989	101	5.9	4.8-7.3	58	9.6	7.1-12.7	43	4.0	2.9-5.4	34	2.3	1.5-3.2	14	2.6	1.3-4.4	20	2.1	1.3-3.2
1990	106	6.4	5.2-7.8	59	9.2	6.8-12.1	47	4.7	3.5-6.3	38	2.4	1.7-3.3	20	3.8	2.2-6.1	18	1.8	1.0-2.8
1991	128	7.4	6.1-8.9	81	12.0	9.3-15.2	47	4.6	3.3-6.1	40	2.3	1.6-3.2	23	3.7	2.2-5.6	17	1.6	0.9-2.6
1992	114	6.7	5.5-8.1	73	11.1	8.5-14.3	41	4.1	2.9-5.6	34	2.1	1.5-3.0	20	3.3	1.9-5.3	14	1.5	0.8-2.5
1993	120	6.6	5.5-8.0	79	11.3	8.7-14.3	41	3.8	2.7-5.2	45	2.7	2.0-3.7	29	4.4	2.8-6.5	16	1.7	0.9-2.7
1994	118	6.4	5.2-7.6	60	8.5	6.3-11.0	58	5.1	3.9-6.6	31	1.7	1.2-2.5	17	2.5	1.4-4.1	14	1.3	0.7-2.2
1995	158	8.7	7.3-10.2	96	13.2	10.6-16.3	62	5.7	4.4-7.3	53	3.1	2.3-4.0	32	4.9	3.3-6.9	21	2.0	1.2-3.0
1996	147	7.7	6.5-9.1	94	12.5	9.9-15.4	53	4.7	3.5-6.1	41	2.2	1.6-3.0	26	3.3	2.1-4.9	15	1.4	0.8-2.4
1997	159	8.5	7.2-9.9	86	12.0	9.4-14.9	73	6.4	5.0-8.0	44	2.4	1.8-3.3	28	4.5	2.9-6.7	16	1.4	0.8-2.2
1998	151	7.8	6.6-9.2	91	11.6	9.2-14.3	60	5.3	4.0-6.8	43	2.3	1.6-3.1	27	3.5	2.2-5.2	16	1.5	0.8-2.4
1999	199	10.2	8.8-11.8	99	13.0	10.5-16.0	100	8.5	6.9-10.3	34	1.9	1.3-2.6	16	2.2	1.2-3.7	18	1.6	0.9-2.5
2000	167	8.5	7.3-10.0	87	11.1	8.8-13.8	80	6.8	5.3-8.4	40	2.1	1.5-2.9	22	3.1	1.8-4.8	18	1.5	0.9-2.4
2001	195	9.4	8.1-10.8	107	13.4	10.8-16.3	88	7.0	5.6-8.6	46	2.4	1.7-3.2	24	3.6	2.3-5.5	22	1.8	1.1-2.7
2002	188	8.9	7.7-10.3	107	13.1	10.6-16.0	81	6.4	5.1-8.0	44	2.2	1.6-3.0	30	4.0	2.6-5.8	14	1.2	0.7-2.0
2003	231	10.7	9.4-12.2	136	16.0	13.4-19.1	95	7.3	5.9-8.9	44	2.2	1.6-3.0	24	3.1	2.0-4.7	20	1.6	1.0-2.5
2004	227	10.4	9.1-11.9	131	14.9	12.4-17.8	96	7.4	6.0-9.0	52	2.6	1.9-3.4	32	4.1	2.8-5.8	20	1.6	1.0-2.5
2005	251	11.5	10.1-13.0	130	14.3	11.9-17.2	121	9.3	7.7-11.2	39	1.8	1.3-2.5	23	2.8	1.7-4.3	16	1.3	0.7-2.1
2006	258	11.7	10.3-13.3	150	17.0	14.3-20.1	108	8.2	6.7-9.9	57	2.8	2.1-3.6	30	3.8	2.5-5.5	27	2.2	1.4-3.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Kidney and renal pelvis

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	276	12.2	10.8-13.8	184	20.6	17.6-23.9	92	7.0	5.6-8.6	53	2.6	1.9-3.4	34	4.4	3.0-6.2	19	1.4	0.9-2.3
2008	273	12.2	10.7-13.7	143	16.2	13.5-19.2	130	9.7	8.1-11.5	47	2.1	1.6-2.9	26	3.2	2.0-4.7	21	1.6	1.0-2.4
2009	290	12.2	10.8-13.8	170	17.8	15.1-20.8	120	8.4	7.0-10.1	38	1.7	1.2-2.4	26	3.3	2.1-4.9	12	0.9	0.4-1.5
2010	331	14.0	12.5-15.6	210	22.3	19.2-25.6	121	8.7	7.2-10.5	51	2.4	1.8-3.1	31	4.0	2.7-5.8	20	1.5	0.9-2.3
2011	316	13.3	11.8-14.8	187	19.3	16.6-22.4	129	9.2	7.7-11.0	50	2.2	1.6-2.9	31	3.7	2.5-5.4	19	1.3	0.8-2.1
2012	321	13.1	11.7-14.6	176	17.3	14.8-20.2	145	10.1	8.5-12.0	47	1.9	1.4-2.5	24	2.6	1.6-3.9	23	1.5	1.0-2.3
2013	369	14.4	12.9-16.0	214	20.3	17.6-23.3	155	10.3	8.7-12.1	44	1.8	1.3-2.4	28	2.8	1.8-4.1	16	1.1	0.6-1.8
2014	370	14.1	12.7-15.7	241	22.3	19.4-25.4	129	8.4	7.0-10.1	46	1.8	1.3-2.5	27	2.9	1.9-4.2	19	1.2	0.7-2.0
2015	357	13.3	11.9-14.8	215	19.5	16.9-22.5	142	9.0	7.5-10.6	59	2.3	1.8-3.0	43	4.6	3.3-6.2	16	0.9	0.5-1.5
2016	391	14.4	13.0-15.9	233	21.3	18.6-24.3	158	9.9	8.4-11.6	53	2.1	1.5-2.7	34	3.3	2.3-4.7	19	1.3	0.8-2.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Brain and other central nervous system

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	46	3.0	2.2-4.1	25	3.8	2.4-5.9	21	2.4	1.5-3.8	36	2.6	1.8-3.7	21	3.7	2.2-5.9	15	1.9	1.0-3.3
1977	58	3.5	2.6-4.6	29	4.3	2.8-6.4	29	3.0	2.0-4.4	32	2.3	1.5-3.3	19	3.2	1.8-5.2	13	1.6	0.8-2.9
1978	33	2.3	1.6-3.3	18	3.0	1.7-5.0	15	1.9	1.0-3.1	30	2.0	1.3-2.9	15	2.2	1.2-3.9	15	1.8	1.0-3.1
1979	52	3.1	2.3-4.1	21	2.9	1.7-4.7	31	3.3	2.2-4.8	39	2.7	1.9-3.7	19	3.4	2.0-5.5	20	2.2	1.4-3.5
1980	57	3.9	2.9-5.2	31	5.1	3.3-7.6	26	3.1	2.0-4.6	38	2.6	1.8-3.7	22	3.9	2.3-6.2	16	1.9	1.0-3.0
1981	40	2.4	1.7-3.4	22	2.7	1.6-4.3	18	2.1	1.2-3.4	29	2.3	1.5-3.4	11	1.8	0.9-3.4	18	2.5	1.4-4.0
1982	65	3.7	2.8-4.7	35	4.6	3.1-6.5	30	3.0	2.0-4.3	56	3.7	2.8-4.9	36	5.8	3.9-8.1	20	2.3	1.4-3.6
1983	60	3.6	2.7-4.7	34	4.7	3.2-6.8	26	2.8	1.8-4.1	50	3.2	2.3-4.2	18	2.7	1.6-4.4	32	3.4	2.3-4.9
1984	77	4.3	3.4-5.5	40	5.2	3.5-7.4	37	3.6	2.5-5.1	52	3.2	2.3-4.2	32	4.7	3.0-6.8	20	2.1	1.3-3.3
1985	79	4.6	3.6-5.7	34	5.1	3.4-7.4	45	4.4	3.1-5.9	45	2.6	1.9-3.6	27	4.2	2.7-6.2	18	1.7	1.0-2.7
1986	75	4.1	3.2-5.2	47	5.8	4.2-8.0	28	2.8	1.8-4.1	54	3.1	2.3-4.0	30	4.1	2.7-5.9	24	2.4	1.5-3.6
1987	83	4.3	3.4-5.5	41	5.4	3.7-7.6	42	3.7	2.6-5.1	46	2.8	2.0-3.8	25	4.3	2.6-6.5	21	2.0	1.2-3.1
1988	64	3.3	2.5-4.3	37	4.7	3.1-6.8	27	2.4	1.6-3.5	37	2.0	1.4-2.8	16	2.3	1.2-4.0	21	1.9	1.2-2.9
1989	74	3.8	2.9-4.8	36	4.5	3.0-6.4	38	3.4	2.4-4.7	32	1.7	1.2-2.5	15	2.1	1.1-3.5	17	1.6	0.9-2.5
1990	81	4.3	3.3-5.3	36	4.3	3.0-6.1	45	4.1	3.0-5.6	37	2.1	1.5-3.0	21	3.3	1.9-5.3	16	1.5	0.9-2.5
1991	92	4.8	3.9-6.0	44	6.1	4.2-8.4	48	4.2	3.0-5.6	42	2.3	1.6-3.2	19	2.5	1.4-4.0	23	2.2	1.4-3.3
1992	92	4.8	3.8-5.9	47	6.1	4.4-8.4	45	3.9	2.8-5.3	39	2.4	1.7-3.3	20	3.2	1.8-5.1	19	1.9	1.1-3.0
1993	69	3.4	2.6-4.4	43	5.2	3.6-7.2	26	2.3	1.5-3.3	38	1.9	1.3-2.6	18	2.4	1.4-4.0	20	1.7	1.0-2.6
1994	88	4.1	3.3-5.1	49	5.5	4.0-7.4	39	3.2	2.3-4.4	51	2.6	1.9-3.4	24	3.2	1.9-4.9	27	2.3	1.5-3.3
1995	82	3.9	3.1-4.9	33	3.4	2.3-4.9	49	4.2	3.1-5.5	34	1.6	1.1-2.3	14	1.5	0.8-2.7	20	1.7	1.0-2.6
1996	98	4.8	3.9-5.9	42	5.0	3.6-7.0	56	4.7	3.5-6.1	33	1.6	1.1-2.3	12	1.6	0.7-2.9	21	1.7	1.1-2.6
1997	79	3.7	2.9-4.6	37	4.1	2.8-5.8	42	3.4	2.4-4.6	38	1.9	1.3-2.6	17	1.9	1.1-3.2	21	1.8	1.1-2.7
1998	97	4.4	3.5-5.4	45	4.5	3.2-6.1	52	4.2	3.1-5.5	32	1.6	1.1-2.3	15	1.8	1.0-3.1	17	1.4	0.8-2.3
1999	90	4.3	3.4-5.3	43	5.2	3.6-7.2	47	3.8	2.8-5.0	29	1.4	0.9-2.0	18	2.1	1.2-3.4	11	0.9	0.5-1.6
2000	95	4.4	3.5-5.4	48	5.5	4.0-7.4	47	3.7	2.7-5.0	41	2.0	1.4-2.8	18	2.1	1.2-3.4	23	1.9	1.2-2.9
2001	94	4.2	3.4-5.2	39	3.8	2.7-5.3	55	4.3	3.3-5.6	41	1.9	1.3-2.6	19	2.1	1.2-3.5	22	1.7	1.1-2.6
2002	84	3.8	3.0-4.8	45	4.9	3.5-6.7	39	3.1	2.2-4.2	42	1.9	1.4-2.6	16	1.7	0.9-2.9	26	2.0	1.3-3.0
2003	109	4.9	4.0-6.0	55	5.6	4.2-7.4	54	4.2	3.2-5.5	34	1.6	1.1-2.2	21	2.4	1.4-3.7	13	1.0	0.5-1.7
2004	89	4.0	3.2-4.9	50	5.4	4.0-7.2	39	3.0	2.1-4.1	49	2.2	1.6-2.9	27	2.9	1.9-4.2	22	1.7	1.1-2.6
2005	97	4.3	3.5-5.3	44	5.0	3.6-6.9	53	4.0	3.0-5.3	36	1.6	1.1-2.2	24	2.8	1.7-4.2	12	0.9	0.5-1.6
2006	83	3.7	2.9-4.6	41	4.3	3.0-5.9	42	3.2	2.3-4.3	45	2.0	1.5-2.7	22	2.6	1.6-4.0	23	1.7	1.1-2.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Brain and other central nervous system

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	102	4.3	3.5-5.3	53	5.2	3.9-6.9	49	3.7	2.7-4.9	33	1.5	1.0-2.1	14	1.5	0.8-2.6	19	1.4	0.8-2.2
2008	85	3.7	2.9-4.6	42	4.4	3.1-6.0	43	3.2	2.3-4.3	44	2.0	1.4-2.7	22	2.6	1.6-3.9	22	1.6	1.0-2.5
2009	90	3.9	3.1-4.8	44	4.5	3.2-6.1	46	3.5	2.5-4.6	39	1.7	1.2-2.3	22	2.4	1.5-3.7	17	1.2	0.7-2.0
2010	89	3.7	3.0-4.6	45	4.4	3.2-6.0	44	3.2	2.3-4.4	33	1.4	1.0-2.0	16	1.7	0.9-2.9	17	1.3	0.7-2.0
2011	103	4.3	3.5-5.2	52	5.4	4.0-7.1	51	3.6	2.6-4.7	49	2.1	1.6-2.8	22	2.3	1.4-3.5	27	2.0	1.3-2.9
2012	93	3.9	3.1-4.8	57	5.5	4.1-7.3	36	2.6	1.8-3.7	40	1.6	1.1-2.2	20	2.0	1.2-3.2	20	1.4	0.8-2.1
2013	117	4.7	3.9-5.6	57	5.1	3.8-6.6	60	4.4	3.3-5.6	47	1.9	1.4-2.6	23	2.3	1.4-3.5	24	1.7	1.1-2.5
2014	82	3.3	2.6-4.1	51	4.6	3.4-6.1	31	2.3	1.5-3.2	47	1.8	1.3-2.5	26	2.6	1.7-3.9	21	1.3	0.8-2.1
2015	93	3.8	3.0-4.6	50	4.7	3.5-6.2	43	3.2	2.3-4.3	37	1.4	1.0-2.0	21	2.0	1.2-3.0	16	1.1	0.6-1.8
2016	106	4.2	3.4-5.1	67	6.1	4.7-7.8	39	2.7	1.9-3.8	59	2.2	1.7-2.9	32	2.8	1.9-4.1	27	1.8	1.2-2.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Thyroid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	29	2.1	1.4-3.0	6	1.1	0.3-2.6	23	2.8	1.8-4.2	8	0.6	0.2-1.2	2	0.4	0.0-1.5	6	0.8	0.3-1.7
1977	33	2.1	1.4-3.1	1	0.2	0.0-1.2	32	3.5	2.3-5.1	4	0.3	0.1-0.9	2	0.3	0.0-1.4	2	0.3	0.0-1.1
1978	26	1.8	1.2-2.8	8	1.7	0.6-3.6	18	2.0	1.2-3.2	7	0.6	0.2-1.2	0	0.0	0.0-0.9	7	0.9	0.4-1.9
1979	37	3.1	2.1-4.3	9	2.0	0.7-4.1	28	3.8	2.4-5.6	6	0.4	0.1-0.9	2	0.3	0.0-1.4	4	0.4	0.1-1.1
1980	33	2.0	1.3-2.9	3	0.4	0.1-1.4	30	3.1	2.1-4.6	7	0.6	0.2-1.2	1	0.2	0.0-1.3	6	0.8	0.3-1.7
1981	27	2.1	1.3-3.1	4	0.6	0.2-1.7	23	3.0	1.9-4.6	5	0.5	0.1-1.2	0	0.0	0.0-0.8	5	0.8	0.2-1.8
1982	32	2.3	1.5-3.2	6	1.0	0.3-2.2	26	3.1	2.0-4.6	9	0.7	0.3-1.4	3	0.9	0.1-2.7	6	0.7	0.2-1.5
1983	29	1.8	1.2-2.6	4	0.5	0.1-1.5	25	2.6	1.7-3.9	6	0.5	0.2-1.1	1	0.3	0.0-1.4	5	0.6	0.2-1.5
1984	44	2.7	1.9-3.7	6	0.8	0.3-1.8	38	4.0	2.8-5.6	9	0.7	0.3-1.3	2	0.6	0.0-2.2	7	0.8	0.3-1.6
1985	48	2.8	2.1-3.8	16	2.3	1.3-3.8	32	3.2	2.2-4.5	7	0.5	0.2-1.0	3	0.5	0.1-1.5	4	0.5	0.1-1.2
1986	35	2.3	1.5-3.2	10	1.9	0.8-3.7	25	2.5	1.6-3.8	8	0.5	0.2-1.1	3	0.7	0.1-2.2	5	0.5	0.2-1.2
1987	53	3.1	2.3-4.1	10	1.5	0.7-3.0	43	4.1	2.9-5.6	8	0.7	0.3-1.3	2	0.6	0.1-2.0	6	0.7	0.3-1.5
1988	41	2.4	1.7-3.3	10	1.6	0.7-3.0	31	3.0	2.0-4.3	5	0.4	0.1-0.9	1	0.1	0.0-0.8	4	0.5	0.1-1.3
1989	37	2.2	1.5-3.0	4	0.7	0.2-1.7	33	3.1	2.1-4.5	8	0.7	0.3-1.4	1	0.2	0.0-1.0	7	0.9	0.4-1.9
1990	34	1.8	1.2-2.6	5	0.7	0.2-1.7	29	2.6	1.7-3.8	3	0.2	0.0-0.6	1	0.2	0.0-1.0	2	0.2	0.0-0.8
1991	41	2.2	1.6-3.1	4	0.7	0.2-1.9	37	3.2	2.3-4.5	7	0.5	0.2-1.1	1	0.3	0.0-1.4	6	0.7	0.2-1.4
1992	70	3.4	2.6-4.3	16	1.6	0.9-2.8	54	4.5	3.4-5.9	9	0.5	0.2-1.0	2	0.2	0.0-0.9	7	0.7	0.3-1.4
1993	53	2.8	2.1-3.7	10	1.3	0.6-2.5	43	3.8	2.7-5.1	5	0.4	0.1-0.8	1	0.1	0.0-0.8	4	0.4	0.1-1.1
1994	61	3.1	2.3-4.0	11	1.5	0.7-2.7	50	4.2	3.1-5.6	7	0.4	0.2-0.8	2	0.3	0.0-1.1	5	0.4	0.1-1.0
1995	66	3.2	2.5-4.1	11	1.4	0.7-2.6	55	4.5	3.3-5.8	5	0.3	0.1-0.8	0	0.0	0.0-0.5	5	0.5	0.2-1.2
1996	89	4.1	3.3-5.1	17	2.3	1.2-3.8	72	5.6	4.4-7.1	12	0.7	0.4-1.2	7	1.1	0.4-2.3	5	0.5	0.2-1.1
1997	89	4.1	3.3-5.1	18	2.0	1.2-3.2	71	5.6	4.3-7.1	9	0.5	0.2-0.9	4	0.6	0.2-1.7	5	0.4	0.1-1.0
1998	105	4.8	3.9-5.9	19	2.4	1.4-3.7	86	6.7	5.3-8.3	8	0.4	0.2-0.9	3	0.3	0.1-1.0	5	0.5	0.2-1.1
1999	100	4.7	3.8-5.7	14	1.5	0.8-2.6	86	6.8	5.4-8.4	11	0.6	0.3-1.1	2	0.2	0.0-0.8	9	0.8	0.4-1.5
2000	113	5.1	4.2-6.1	14	1.4	0.7-2.3	99	7.6	6.2-9.3	12	0.6	0.3-1.1	2	0.2	0.0-0.8	10	0.9	0.4-1.6
2001	117	5.1	4.2-6.2	23	2.4	1.5-3.7	94	7.1	5.7-8.6	15	0.7	0.4-1.2	3	0.3	0.1-1.0	12	1.0	0.5-1.7
2002	124	5.4	4.5-6.4	21	2.3	1.4-3.5	103	7.6	6.2-9.3	6	0.3	0.1-0.7	1	0.1	0.0-0.6	5	0.4	0.1-1.0
2003	113	4.9	4.0-5.9	16	1.9	1.0-3.1	97	7.1	5.8-8.7	4	0.2	0.1-0.6	0	0.0	0.0-0.5	4	0.3	0.1-0.9
2004	126	5.4	4.5-6.5	25	3.0	1.9-4.5	101	7.4	6.0-9.0	10	0.5	0.2-0.9	5	0.8	0.2-2.0	5	0.4	0.1-0.9
2005	187	8.0	6.9-9.3	33	3.5	2.4-5.0	154	11.4	9.6-13.3	16	0.7	0.4-1.2	5	0.6	0.2-1.4	11	0.8	0.4-1.5
2006	156	6.8	5.8-8.0	20	2.3	1.4-3.5	136	10.0	8.4-11.9	9	0.4	0.2-0.8	2	0.2	0.0-0.8	7	0.5	0.2-1.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Thyroid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	161	6.8	5.8-7.9	32	3.3	2.2-4.7	129	9.4	7.9-11.2	9	0.5	0.2-0.9	1	0.2	0.0-0.8	8	0.6	0.3-1.2
2008	212	8.9	7.7-10.1	42	4.4	3.1-6.1	170	12.2	10.4-14.2	11	0.6	0.3-1.0	3	0.5	0.1-1.5	8	0.6	0.3-1.2
2009	205	8.7	7.5-10.0	37	3.5	2.4-4.9	168	12.4	10.6-14.5	10	0.5	0.2-0.9	4	0.5	0.1-1.4	6	0.4	0.2-1.0
2010	243	10.0	8.8-11.4	34	3.7	2.5-5.2	209	14.8	12.9-17.0	14	0.6	0.3-1.1	7	0.9	0.3-1.8	7	0.5	0.2-1.1
2011	276	10.9	9.6-12.2	53	5.3	3.9-7.0	223	15.2	13.2-17.3	8	0.3	0.1-0.7	1	0.2	0.0-0.8	7	0.5	0.2-1.0
2012	250	10.0	8.8-11.4	33	3.4	2.3-4.8	217	15.1	13.1-17.2	9	0.4	0.2-0.8	3	0.4	0.1-1.2	6	0.4	0.1-0.9
2013	310	12.0	10.7-13.5	39	3.4	2.4-4.7	271	18.4	16.3-20.8	12	0.5	0.3-0.9	5	0.5	0.2-1.2	7	0.5	0.2-1.0
2014	286	11.3	10.0-12.7	43	4.2	3.0-5.7	243	16.7	14.6-19.0	17	0.7	0.4-1.1	2	0.2	0.0-0.7	15	1.0	0.5-1.6
2015	281	11.2	9.9-12.6	47	4.4	3.2-5.9	234	16.3	14.2-18.6	12	0.5	0.3-0.9	3	0.3	0.1-0.8	9	0.6	0.3-1.2
2016	261	10.1	8.9-11.4	47	4.3	3.1-5.7	214	14.4	12.5-16.6	13	0.5	0.3-0.8	5	0.4	0.1-1.0	8	0.5	0.2-1.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Hodgkin lymphoma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	33	2.5	1.6-3.8	16	3.8	1.6-7.0	17	2.0	1.1-3.3	15	0.9	0.5-1.6	9	1.2	0.5-2.6	6	0.7	0.3-1.6
1977	30	1.8	1.2-2.7	16	2.3	1.3-4.0	14	1.5	0.8-2.6	15	1.2	0.6-2.1	4	1.2	0.2-3.5	11	1.4	0.7-2.5
1978	29	1.8	1.1-2.7	14	1.9	0.9-3.5	15	1.7	0.9-2.9	12	0.7	0.3-1.2	6	0.8	0.3-2.1	6	0.6	0.2-1.4
1979	40	2.4	1.7-3.4	25	3.6	2.2-5.5	15	1.6	0.9-2.7	17	1.0	0.6-1.7	11	1.6	0.8-3.1	6	0.6	0.2-1.4
1980	46	2.4	1.7-3.2	22	2.5	1.6-4.0	24	2.2	1.4-3.3	9	0.5	0.2-1.0	4	0.6	0.2-1.8	5	0.4	0.1-1.1
1981	41	2.2	1.6-3.1	25	3.3	2.0-5.2	16	1.5	0.8-2.5	17	1.1	0.6-1.8	9	1.6	0.6-3.3	8	0.8	0.3-1.5
1982	41	2.1	1.5-2.9	25	2.9	1.8-4.5	16	1.5	0.8-2.4	15	0.8	0.4-1.4	7	0.9	0.3-2.0	8	0.7	0.3-1.5
1983	55	2.9	2.2-3.9	32	4.3	2.7-6.4	23	2.1	1.3-3.1	12	0.7	0.4-1.3	6	0.8	0.3-1.9	6	0.6	0.2-1.4
1984	43	2.2	1.6-3.0	28	3.2	2.0-4.8	15	1.5	0.8-2.5	18	1.0	0.6-1.6	16	1.9	1.0-3.2	2	0.3	0.0-0.9
1985	33	1.5	1.0-2.1	22	2.4	1.4-3.8	11	0.8	0.4-1.6	21	1.1	0.6-1.7	16	1.9	1.0-3.2	5	0.4	0.1-1.1
1986	44	2.3	1.6-3.1	30	3.6	2.4-5.3	14	1.3	0.7-2.2	11	0.6	0.3-1.1	6	0.8	0.3-1.8	5	0.5	0.1-1.2
1987	39	1.9	1.3-2.7	27	3.4	2.1-5.3	12	1.0	0.5-1.7	13	0.6	0.3-1.1	11	1.4	0.7-2.7	2	0.1	0.0-0.6
1988	55	2.5	1.8-3.2	32	3.0	2.0-4.4	23	2.0	1.2-3.0	10	0.5	0.2-0.9	4	0.4	0.1-1.2	6	0.5	0.2-1.1
1989	52	2.4	1.8-3.2	33	3.7	2.5-5.4	19	1.5	0.9-2.4	16	0.7	0.4-1.2	15	1.6	0.9-2.7	1	0.1	0.0-0.6
1990	44	2.0	1.4-2.7	26	3.0	1.9-4.6	18	1.4	0.8-2.2	15	0.7	0.4-1.2	10	1.2	0.5-2.4	5	0.4	0.1-1.0
1991	67	3.0	2.3-3.9	40	4.5	3.1-6.4	27	2.0	1.3-3.0	11	0.5	0.2-1.0	5	0.5	0.2-1.3	6	0.5	0.2-1.1
1992	53	2.4	1.8-3.1	25	2.3	1.5-3.6	28	2.3	1.5-3.3	18	0.8	0.5-1.3	9	0.8	0.4-1.7	9	0.7	0.3-1.4
1993	47	1.9	1.4-2.6	22	2.1	1.3-3.3	25	1.8	1.2-2.7	14	0.6	0.3-1.0	8	0.7	0.3-1.6	6	0.5	0.2-1.1
1994	47	2.1	1.5-2.9	30	3.2	2.1-4.8	17	1.3	0.7-2.1	15	0.7	0.4-1.2	9	1.1	0.5-2.1	6	0.4	0.2-1.0
1995	54	2.3	1.7-3.0	30	3.0	2.0-4.4	24	1.7	1.1-2.6	9	0.4	0.2-0.7	5	0.5	0.2-1.3	4	0.3	0.1-0.7
1996	45	2.0	1.4-2.7	23	2.3	1.4-3.6	22	1.7	1.0-2.6	12	0.6	0.3-1.0	6	0.8	0.3-1.7	6	0.5	0.2-1.0
1997	62	2.6	2.0-3.3	37	3.4	2.4-4.8	25	1.9	1.2-2.8	12	0.6	0.3-1.1	4	0.4	0.1-1.1	8	0.7	0.3-1.4
1998	56	2.3	1.7-3.0	27	2.4	1.6-3.6	29	2.1	1.4-3.1	10	0.4	0.2-0.7	5	0.4	0.1-1.1	5	0.4	0.1-0.8
1999	50	2.1	1.6-2.8	27	2.8	1.8-4.2	23	1.7	1.1-2.6	10	0.4	0.2-0.8	7	0.8	0.3-1.6	3	0.2	0.0-0.7
2000	58	2.3	1.8-3.0	35	3.2	2.2-4.5	23	1.7	1.1-2.5	5	0.2	0.1-0.5	3	0.3	0.1-0.9	2	0.2	0.0-0.6
2001	52	2.0	1.5-2.7	35	3.0	2.1-4.3	17	1.2	0.7-1.9	8	0.3	0.1-0.7	4	0.4	0.1-1.1	4	0.3	0.1-0.8
2002	63	2.7	2.0-3.4	32	3.3	2.2-4.7	31	2.3	1.5-3.2	7	0.3	0.1-0.6	5	0.5	0.2-1.2	2	0.1	0.0-0.5
2003	61	2.5	1.9-3.2	25	2.4	1.5-3.6	36	2.6	1.8-3.6	13	0.6	0.3-1.0	7	1.0	0.4-2.0	6	0.4	0.2-0.9
2004	65	2.6	2.0-3.4	35	3.2	2.2-4.5	30	2.2	1.5-3.1	8	0.3	0.1-0.7	5	0.5	0.2-1.3	3	0.2	0.0-0.6
2005	74	3.1	2.4-3.9	45	4.1	3.0-5.5	29	2.2	1.5-3.1	5	0.2	0.1-0.5	4	0.4	0.1-1.0	1	0.1	0.0-0.4
2006	57	2.3	1.8-3.0	30	2.8	1.9-4.1	27	1.9	1.3-2.8	5	0.2	0.1-0.6	1	0.1	0.0-0.6	4	0.3	0.1-0.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Hodgkin lymphoma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	54	2.3	1.7-3.0	29	2.9	1.9-4.2	25	1.8	1.2-2.7	8	0.4	0.2-0.7	2	0.2	0.0-0.7	6	0.5	0.2-1.0
2008	92	3.8	3.0-4.6	52	4.8	3.6-6.4	40	3.0	2.1-4.1	5	0.2	0.1-0.5	3	0.3	0.1-0.8	2	0.2	0.0-0.6
2009	71	3.0	2.3-3.8	27	2.7	1.7-4.0	44	3.2	2.4-4.4	7	0.3	0.1-0.6	4	0.4	0.1-1.1	3	0.2	0.0-0.6
2010	81	3.3	2.6-4.2	48	4.6	3.3-6.2	33	2.4	1.6-3.4	8	0.3	0.1-0.6	4	0.3	0.1-0.9	4	0.3	0.1-0.8
2011	85	3.3	2.6-4.1	46	3.9	2.8-5.2	39	2.8	2.0-3.8	7	0.3	0.1-0.6	4	0.5	0.1-1.3	3	0.2	0.0-0.6
2012	65	2.6	2.0-3.3	31	2.7	1.8-3.8	34	2.5	1.7-3.5	4	0.2	0.0-0.5	3	0.3	0.1-0.9	1	0.1	0.0-0.4
2013	87	3.5	2.8-4.3	45	4.2	3.0-5.6	42	3.1	2.2-4.2	8	0.3	0.1-0.7	4	0.4	0.1-1.1	4	0.3	0.1-0.7
2014	89	3.4	2.7-4.2	46	4.0	2.9-5.3	43	3.0	2.1-4.0	6	0.2	0.1-0.5	3	0.3	0.1-1.0	3	0.2	0.0-0.5
2015	71	2.9	2.2-3.6	36	3.3	2.2-4.6	35	2.6	1.8-3.6	6	0.2	0.1-0.5	2	0.2	0.0-0.8	4	0.3	0.1-0.7
2016	64	2.5	1.9-3.2	28	2.5	1.6-3.7	36	2.4	1.7-3.4	2	0.1	0.0-0.3	2	0.2	0.0-0.8	0	0.0	0.0-0.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Non-Hodgkin lymphomas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	69	5.6	4.3-7.3	32	7.1	4.5-10.5	37	4.8	3.3-6.8	45	3.8	2.7-5.1	28	6.2	3.9-9.3	17	2.2	1.3-3.7
1977	72	5.6	4.3-7.2	29	5.7	3.5-8.6	43	5.6	4.0-7.7	37	3.0	1.9-4.3	14	2.2	1.1-3.9	23	3.2	1.9-5.1
1978	109	8.0	6.4-9.7	60	10.4	7.7-13.7	49	6.2	4.5-8.4	52	3.7	2.8-5.0	33	6.1	4.1-8.9	19	2.2	1.3-3.5
1979	85	6.8	5.3-8.6	48	9.2	6.5-12.5	37	5.2	3.5-7.3	50	4.5	3.2-6.1	24	5.2	3.0-8.1	26	4.0	2.5-6.0
1980	107	7.7	6.2-9.4	53	9.1	6.6-12.2	54	6.7	4.9-8.9	55	4.1	3.0-5.4	34	5.6	3.8-8.1	21	2.9	1.7-4.6
1981	99	7.4	5.9-9.2	46	7.1	4.9-10.0	53	7.2	5.3-9.6	60	4.4	3.3-5.8	26	4.8	3.0-7.3	34	4.2	2.8-5.9
1982	127	8.8	7.2-10.5	57	9.1	6.7-12.0	70	8.3	6.4-10.6	53	4.2	3.1-5.6	24	4.8	2.8-7.4	29	3.8	2.5-5.6
1983	131	8.6	7.1-10.3	62	9.9	7.4-13.0	69	7.8	6.0-9.9	73	5.1	3.9-6.4	39	6.3	4.3-9.0	34	4.2	2.8-5.9
1984	149	9.5	7.9-11.3	72	11.4	8.6-14.8	77	8.3	6.5-10.4	60	3.9	2.9-5.1	34	5.3	3.5-7.8	26	2.9	1.9-4.3
1985	186	11.3	9.6-13.1	97	13.7	10.8-17.1	89	9.5	7.6-11.8	83	5.3	4.1-6.7	41	5.9	4.0-8.4	42	4.8	3.4-6.5
1986	191	11.9	10.1-13.8	97	15.3	12.0-19.2	94	9.8	7.9-12.1	82	5.1	4.0-6.4	40	6.2	4.1-8.8	42	4.5	3.2-6.1
1987	202	11.8	10.1-13.6	121	17.8	14.4-21.7	81	7.9	6.2-9.9	77	4.5	3.5-5.6	38	5.5	3.7-7.9	39	3.8	2.7-5.3
1988	174	9.9	8.4-11.5	102	14.1	11.3-17.4	72	7.1	5.5-8.9	85	5.3	4.2-6.6	55	8.9	6.5-12.0	30	3.1	2.1-4.5
1989	203	11.3	9.7-13.0	118	15.1	12.2-18.4	85	8.3	6.6-10.3	87	5.1	4.1-6.4	51	7.5	5.3-10.1	36	3.6	2.5-5.0
1990	231	11.9	10.4-13.6	138	16.3	13.5-19.4	93	8.5	6.8-10.5	75	4.4	3.4-5.6	39	5.6	3.7-8.0	36	3.6	2.5-5.0
1991	238	12.9	11.2-14.8	134	16.8	13.7-20.2	104	9.9	8.0-12.1	75	4.4	3.4-5.6	43	5.8	4.0-8.1	32	3.3	2.2-4.7
1992	281	14.9	13.2-16.9	154	19.0	16.0-22.6	127	11.7	9.7-14.0	85	4.8	3.8-6.0	41	5.6	3.8-7.8	44	4.2	3.0-5.7
1993	269	13.1	11.5-14.8	169	19.0	16.1-22.3	100	8.5	6.9-10.4	78	4.2	3.3-5.3	39	5.0	3.5-7.0	39	3.6	2.5-5.0
1994	317	16.4	14.6-18.4	183	22.0	18.8-25.7	134	12.1	10.1-14.4	87	4.8	3.8-6.0	38	4.8	3.3-6.7	49	4.6	3.4-6.1
1995	282	14.5	12.8-16.3	167	20.1	17.0-23.7	115	10.3	8.5-12.4	85	4.6	3.6-5.7	49	6.1	4.4-8.2	36	3.4	2.4-4.7
1996	296	14.7	13.1-16.6	157	18.1	15.3-21.4	139	11.9	10.0-14.1	93	5.2	4.2-6.4	39	5.4	3.6-7.6	54	5.0	3.7-6.5
1997	330	16.6	14.8-18.5	187	22.6	19.2-26.3	143	12.4	10.4-14.6	95	5.3	4.2-6.5	49	6.8	4.9-9.1	46	4.2	3.1-5.6
1998	317	15.6	13.9-17.4	169	20.4	17.2-23.9	148	12.4	10.4-14.6	108	5.9	4.8-7.1	53	7.2	5.3-9.5	55	4.9	3.7-6.4
1999	331	15.8	14.1-17.6	174	20.3	17.2-23.7	157	12.7	10.8-14.9	114	5.7	4.7-6.9	64	8.5	6.4-11.0	50	4.2	3.1-5.5
2000	299	13.8	12.3-15.5	154	17.2	14.5-20.3	145	11.4	9.6-13.4	99	4.9	4.0-6.0	53	6.8	5.0-9.0	46	3.8	2.7-5.0
2001	309	14.5	12.9-16.2	160	18.3	15.5-21.6	149	11.9	10.0-13.9	111	5.4	4.5-6.6	57	6.7	5.0-8.7	54	4.4	3.3-5.8
2002	302	13.5	12.0-15.1	159	17.0	14.4-20.0	143	10.9	9.2-12.9	105	5.0	4.1-6.1	61	7.5	5.6-9.7	44	3.5	2.5-4.6
2003	339	15.3	13.6-17.0	156	17.3	14.6-20.4	183	13.9	11.9-16.1	109	5.0	4.1-6.1	57	6.8	5.1-9.0	52	4.0	3.0-5.2
2004	311	14.2	12.6-15.9	154	17.4	14.7-20.6	157	12.0	10.2-14.0	87	4.1	3.2-5.0	46	5.4	3.9-7.3	41	3.2	2.3-4.3
2005	363	16.0	14.4-17.8	173	18.5	15.7-21.6	190	14.3	12.3-16.5	91	4.3	3.4-5.3	45	5.5	3.9-7.4	46	3.6	2.6-4.8
2006	320	14.3	12.7-15.9	153	16.7	14.0-19.7	167	12.7	10.8-14.8	81	3.7	2.9-4.6	35	4.2	2.8-5.9	46	3.5	2.5-4.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Non-Hodgkin lymphomas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	355	15.5	13.9-17.3	194	20.1	17.3-23.2	161	12.1	10.3-14.1	100	4.5	3.7-5.5	45	5.3	3.8-7.2	55	4.1	3.1-5.4
2008	342	14.8	13.2-16.5	189	19.6	16.9-22.8	153	11.3	9.6-13.3	82	3.7	2.9-4.6	39	4.5	3.2-6.3	43	3.2	2.3-4.3
2009	383	16.9	15.2-18.7	188	20.2	17.3-23.4	195	14.6	12.6-16.8	86	3.8	3.0-4.7	40	4.5	3.2-6.2	46	3.4	2.5-4.5
2010	360	15.3	13.7-17.0	184	18.9	16.2-22.0	176	12.6	10.8-14.7	85	3.8	3.0-4.7	39	4.4	3.1-6.1	46	3.3	2.4-4.5
2011	413	17.2	15.5-18.9	213	22.4	19.3-25.7	200	13.8	11.9-15.9	92	4.0	3.2-4.9	49	5.3	3.9-7.2	43	3.1	2.2-4.2
2012	381	15.7	14.1-17.3	180	18.1	15.4-21.1	201	14.1	12.2-16.2	90	3.8	3.0-4.6	47	5.1	3.7-6.9	43	2.9	2.1-4.0
2013	385	15.2	13.7-16.8	189	18.1	15.5-21.0	196	13.1	11.3-15.1	95	3.9	3.1-4.8	50	5.4	3.9-7.2	45	3.0	2.2-4.0
2014	413	15.9	14.4-17.5	214	20.0	17.3-23.0	199	13.1	11.3-15.0	93	3.6	2.9-4.5	47	4.8	3.5-6.5	46	3.0	2.2-4.0
2015	389	15.0	13.5-16.6	183	17.3	14.8-20.1	206	13.3	11.5-15.3	88	3.6	2.8-4.4	41	4.5	3.2-6.2	47	3.1	2.3-4.1
2016	425	16.3	14.7-18.0	211	19.9	17.2-22.9	214	14.0	12.2-16.1	99	3.9	3.1-4.7	45	4.4	3.2-5.9	54	3.5	2.6-4.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Myeloma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	91	9.0	7.1-11.2	45	11.7	8.2-16.0	46	7.3	5.2-9.9	50	5.2	3.7-6.9	19	4.2	2.4-6.7	31	5.5	3.6-8.0
1977	92	8.7	6.9-10.9	41	10.4	7.0-14.9	51	7.8	5.7-10.4	62	6.1	4.5-8.0	28	7.9	4.9-11.9	34	5.2	3.5-7.3
1978	92	8.7	6.9-10.8	52	14.0	10.0-19.0	40	5.8	4.0-7.9	45	4.1	2.9-5.6	25	6.8	4.1-10.5	20	2.7	1.6-4.2
1979	106	8.8	7.1-10.7	48	9.2	6.6-12.4	58	8.3	6.2-10.8	56	5.0	3.7-6.6	23	5.1	3.1-7.9	33	4.9	3.2-6.9
1980	120	10.2	8.3-12.4	48	11.4	8.1-15.6	72	9.7	7.5-12.4	77	6.7	5.2-8.5	34	8.1	5.3-11.6	43	5.9	4.2-8.0
1981	112	9.1	7.4-11.1	49	10.2	7.3-13.7	63	8.5	6.4-11.0	56	4.9	3.6-6.5	21	5.1	2.9-8.1	35	4.8	3.3-6.8
1982	119	9.7	7.9-11.7	52	10.7	7.7-14.3	67	9.0	6.8-11.5	85	7.0	5.5-8.7	42	8.9	6.1-12.4	43	5.8	4.2-7.9
1983	120	9.2	7.6-11.2	49	9.2	6.6-12.3	71	9.1	7.0-11.5	76	6.2	4.8-7.8	28	6.0	3.8-8.9	48	6.2	4.5-8.3
1984	132	10.4	8.6-12.4	54	11.8	8.5-15.9	78	9.7	7.6-12.1	89	7.0	5.5-8.7	30	5.9	3.8-8.6	59	7.5	5.6-9.8
1985	134	9.7	8.1-11.6	54	11.6	8.3-15.5	80	9.0	7.1-11.3	80	5.9	4.6-7.4	38	8.0	5.4-11.3	42	4.8	3.4-6.6
1986	134	9.5	7.9-11.3	60	11.7	8.6-15.3	74	8.3	6.5-10.5	79	6.0	4.7-7.5	34	6.9	4.6-9.8	45	5.4	3.9-7.3
1987	125	8.5	7.0-10.2	60	10.7	8.0-14.0	65	7.1	5.5-9.1	77	5.4	4.2-6.8	40	7.6	5.2-10.7	37	4.1	2.9-5.7
1988	109	7.2	5.9-8.8	48	8.3	5.9-11.2	61	6.6	5.0-8.5	63	4.4	3.4-5.7	29	5.6	3.5-8.2	34	3.8	2.6-5.3
1989	134	9.5	7.9-11.3	62	12.0	8.9-15.8	72	8.1	6.3-10.2	61	4.5	3.4-5.8	28	6.2	3.9-9.1	33	3.7	2.5-5.2
1990	139	8.8	7.3-10.4	56	10.6	7.7-14.1	83	8.2	6.5-10.2	65	4.6	3.5-5.9	26	5.7	3.6-8.5	39	4.2	3.0-5.7
1991	145	9.7	8.1-11.4	47	8.7	6.2-11.8	98	10.2	8.2-12.5	84	5.7	4.5-7.1	33	6.6	4.3-9.5	51	5.4	4.0-7.1
1992	179	11.1	9.5-12.9	81	14.1	10.9-17.8	98	9.6	7.8-11.8	82	5.1	4.0-6.3	30	5.7	3.6-8.4	52	5.0	3.7-6.6
1993	130	7.9	6.6-9.4	55	8.4	6.2-11.0	75	7.4	5.8-9.3	75	4.8	3.8-6.1	32	6.0	3.9-8.6	43	4.3	3.1-5.8
1994	145	9.0	7.5-10.6	61	10.5	7.9-13.7	84	8.2	6.6-10.2	78	4.9	3.9-6.2	27	4.4	2.8-6.6	51	5.1	3.8-6.7
1995	163	9.8	8.3-11.5	68	11.5	8.7-14.8	95	9.0	7.3-11.0	108	6.6	5.4-8.0	45	7.6	5.4-10.4	63	6.0	4.6-7.7
1996	160	9.2	7.8-10.8	67	10.1	7.7-13.0	93	8.6	7.0-10.6	81	4.8	3.8-6.0	38	6.4	4.4-8.9	43	4.0	2.9-5.4
1997	177	10.1	8.6-11.7	75	11.1	8.6-14.1	102	9.4	7.6-11.4	87	5.1	4.1-6.3	39	6.5	4.5-9.0	48	4.4	3.3-5.9
1998	179	10.1	8.6-11.7	77	11.6	9.0-14.7	102	9.2	7.5-11.2	76	4.4	3.4-5.5	26	4.1	2.6-6.1	50	4.5	3.3-6.0
1999	187	10.3	8.8-11.9	67	9.2	7.0-11.8	120	10.6	8.8-12.7	84	4.6	3.7-5.8	31	4.4	2.9-6.3	53	4.7	3.5-6.1
2000	193	10.1	8.7-11.6	76	10.3	7.9-13.0	117	9.8	8.1-11.8	71	4.1	3.2-5.1	17	2.8	1.6-4.6	54	4.8	3.6-6.2
2001	213	11.0	9.5-12.6	84	11.6	9.1-14.5	129	10.8	9.0-12.8	87	4.7	3.7-5.8	36	5.4	3.7-7.6	51	4.3	3.2-5.6
2002	217	11.0	9.6-12.6	86	11.9	9.4-14.9	131	10.7	8.9-12.7	107	5.6	4.6-6.8	43	5.7	4.0-7.8	64	5.4	4.2-6.9
2003	228	11.6	10.2-13.3	94	13.1	10.4-16.2	134	10.9	9.1-12.9	89	4.5	3.6-5.5	38	4.8	3.3-6.8	51	4.2	3.1-5.5
2004	211	10.2	8.8-11.7	87	10.7	8.5-13.3	124	9.7	8.1-11.6	94	4.8	3.9-5.9	39	5.3	3.7-7.3	55	4.6	3.4-6.0
2005	252	12.1	10.6-13.7	109	13.5	11.0-16.4	143	11.1	9.3-13.1	87	4.3	3.4-5.3	35	4.6	3.1-6.5	52	4.1	3.0-5.4
2006	253	12.2	10.7-13.8	118	14.7	12.1-17.7	135	10.5	8.8-12.4	80	3.9	3.1-4.9	34	4.4	3.0-6.2	46	3.6	2.6-4.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Myeloma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	221	10.6	9.3-12.2	100	13.0	10.4-15.9	121	9.3	7.7-11.1	109	5.2	4.3-6.3	48	6.0	4.3-8.0	61	4.7	3.6-6.0
2008	289	13.2	11.7-14.9	134	16.3	13.5-19.4	155	11.4	9.7-13.4	76	3.7	2.9-4.6	31	4.0	2.7-5.7	45	3.4	2.5-4.6
2009	310	14.0	12.5-15.7	156	17.7	14.9-20.8	154	11.5	9.7-13.5	93	4.3	3.5-5.3	52	6.2	4.6-8.2	41	3.1	2.2-4.2
2010	297	13.1	11.7-14.8	157	18.5	15.6-21.8	140	10.1	8.5-11.9	101	4.7	3.8-5.7	45	5.7	4.1-7.7	56	4.1	3.1-5.4
2011	345	14.7	13.2-16.4	170	18.5	15.7-21.7	175	12.5	10.7-14.6	98	4.3	3.4-5.2	47	5.5	4.0-7.4	51	3.5	2.6-4.6
2012	349	14.6	13.1-16.3	152	16.7	14.1-19.7	197	13.4	11.6-15.5	95	4.0	3.2-4.9	47	4.9	3.6-6.6	48	3.3	2.4-4.4
2013	328	13.1	11.7-14.7	158	16.7	14.0-19.6	170	11.1	9.5-12.9	96	3.9	3.2-4.8	52	5.6	4.1-7.4	44	3.0	2.1-4.0
2014	355	14.2	12.7-15.8	165	16.7	14.2-19.6	190	12.6	10.9-14.6	100	4.2	3.4-5.2	49	5.5	4.0-7.4	51	3.5	2.6-4.6
2015	319	12.4	11.0-13.8	155	15.1	12.8-17.8	164	10.6	9.0-12.4	127	5.0	4.1-5.9	59	6.3	4.8-8.2	68	4.2	3.3-5.4
2016	352	13.2	11.9-14.7	164	15.6	13.2-18.3	188	11.7	10.1-13.6	131	5.2	4.3-6.2	59	6.1	4.6-7.9	72	4.7	3.6-5.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Leukemias

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	101	7.9	6.2-9.8	51	9.6	6.8-13.1	50	6.7	4.8-9.1	74	6.1	4.6-7.9	41	8.1	5.2-11.9	33	4.8	3.2-6.9
1977	91	7.2	5.6-9.0	46	9.5	6.3-13.5	45	6.0	4.3-8.1	81	6.3	4.9-8.0	40	8.5	5.7-12.0	41	5.0	3.5-6.9
1978	118	9.3	7.5-11.3	60	11.9	8.5-16.0	58	7.7	5.7-10.1	83	6.6	5.1-8.3	47	8.6	6.1-11.8	36	5.1	3.5-7.3
1979	106	8.6	6.9-10.6	57	12.6	8.9-17.2	49	6.4	4.7-8.6	76	6.0	4.6-7.6	44	9.6	6.5-13.4	32	4.0	2.6-5.7
1980	115	8.8	7.1-10.7	53	10.1	7.1-13.9	62	7.9	6.0-10.3	81	6.5	5.0-8.2	36	7.9	5.0-11.6	45	5.8	4.1-7.9
1981	116	8.7	7.1-10.6	61	12.1	8.8-16.1	55	6.9	5.1-9.0	81	6.1	4.7-7.7	33	6.4	4.0-9.6	48	6.0	4.3-8.0
1982	124	8.8	7.2-10.7	63	12.1	8.8-16.1	61	7.1	5.3-9.2	72	5.4	4.1-6.9	36	7.5	4.9-11.0	36	4.3	3.0-6.1
1983	129	8.8	7.2-10.5	65	11.6	8.6-15.3	64	7.2	5.5-9.3	79	5.5	4.2-6.9	38	6.5	4.3-9.2	41	4.8	3.4-6.6
1984	131	8.7	7.1-10.4	70	10.7	8.0-13.9	61	7.1	5.4-9.2	83	5.5	4.3-7.0	41	7.2	4.8-10.1	42	4.7	3.3-6.4
1985	135	8.6	7.1-10.3	82	13.4	10.3-17.0	53	5.7	4.2-7.6	78	5.2	4.1-6.6	49	8.9	6.3-12.2	29	3.1	2.0-4.5
1986	129	8.2	6.7-9.8	68	11.7	8.7-15.3	61	6.3	4.8-8.1	75	4.6	3.6-5.9	40	6.6	4.5-9.3	35	3.5	2.4-5.0
1987	140	8.7	7.2-10.4	67	10.4	7.7-13.6	73	7.6	5.9-9.6	74	4.4	3.4-5.6	36	5.5	3.5-8.0	38	3.9	2.7-5.3
1988	110	7.1	5.8-8.7	58	10.6	7.7-14.0	52	5.3	3.9-7.0	76	5.2	4.0-6.6	35	6.2	4.1-9.0	41	4.6	3.3-6.3
1989	155	9.3	7.8-11.0	79	12.5	9.5-16.1	76	7.5	5.9-9.5	82	5.2	4.1-6.5	47	8.4	5.9-11.5	35	3.6	2.4-5.0
1990	154	9.1	7.7-10.8	74	10.9	8.2-14.1	80	8.0	6.3-10.1	95	5.9	4.7-7.3	45	7.5	5.2-10.3	50	5.0	3.6-6.6
1991	180	10.4	8.9-12.2	87	13.2	10.3-16.8	93	8.8	7.1-10.9	71	4.3	3.3-5.5	32	5.3	3.4-7.8	39	3.9	2.7-5.3
1992	172	9.9	8.4-11.6	74	11.2	8.4-14.5	98	9.3	7.5-11.4	86	5.4	4.2-6.7	36	5.4	3.6-7.8	50	5.1	3.8-6.8
1993	151	8.6	7.3-10.2	74	11.2	8.5-14.4	77	7.3	5.7-9.1	84	5.0	3.9-6.2	39	5.9	4.0-8.4	45	4.4	3.2-5.9
1994	155	8.8	7.4-10.3	79	11.7	9.0-14.9	76	7.1	5.5-8.9	113	6.6	5.4-8.0	51	7.4	5.3-10.1	62	6.0	4.6-7.7
1995	163	8.7	7.4-10.2	80	10.2	7.9-13.0	83	7.5	6.0-9.4	93	5.2	4.1-6.4	43	6.4	4.5-8.9	50	4.6	3.4-6.0
1996	176	9.4	8.0-11.0	95	12.6	9.9-15.7	81	7.3	5.8-9.1	97	5.5	4.4-6.8	45	6.5	4.5-9.0	52	4.9	3.6-6.4
1997	160	8.6	7.3-10.1	64	8.8	6.6-11.5	96	8.5	6.9-10.4	95	5.4	4.3-6.6	49	7.8	5.6-10.5	46	4.1	3.0-5.5
1998	154	8.0	6.8-9.5	67	9.3	7.0-12.1	87	7.4	5.9-9.2	76	4.0	3.1-5.0	34	4.6	3.0-6.6	42	3.7	2.6-4.9
1999	146	7.3	6.1-8.6	75	10.3	7.9-13.1	71	5.8	4.5-7.3	92	5.0	4.0-6.1	47	7.0	5.0-9.6	45	3.9	2.8-5.2
2000	191	9.7	8.4-11.2	91	11.6	9.2-14.4	100	8.4	6.8-10.2	93	5.0	4.0-6.2	35	5.4	3.6-7.7	58	4.9	3.7-6.4
2001	164	8.2	7.0-9.6	70	9.0	6.8-11.6	94	7.7	6.2-9.5	90	4.7	3.8-5.8	41	5.6	3.9-7.7	49	4.1	3.0-5.5
2002	186	9.0	7.7-10.4	99	12.2	9.7-15.0	87	7.0	5.6-8.7	93	4.6	3.7-5.7	47	6.4	4.6-8.7	46	3.7	2.7-4.9
2003	185	8.6	7.4-10.0	91	10.2	8.1-12.6	94	7.4	6.0-9.1	90	4.4	3.5-5.4	40	4.8	3.3-6.6	50	4.1	3.0-5.4
2004	186	8.8	7.5-10.2	102	12.1	9.7-14.8	84	6.6	5.3-8.2	81	4.1	3.2-5.1	42	5.5	3.9-7.5	39	3.2	2.3-4.3
2005	171	7.9	6.8-9.3	87	10.4	8.2-13.0	84	6.5	5.2-8.0	89	4.4	3.5-5.5	41	5.6	3.9-7.7	48	3.8	2.8-5.0
2006	184	8.6	7.3-9.9	107	12.9	10.5-15.8	77	5.9	4.7-7.4	80	3.8	3.0-4.7	47	5.9	4.2-7.9	33	2.5	1.7-3.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York City
Blacks
Leukemias

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	185	8.5	7.3-9.9	84	9.4	7.4-11.7	101	7.7	6.3-9.4	85	4.2	3.3-5.2	45	6.0	4.3-8.1	40	3.2	2.3-4.4
2008	178	8.2	7.0-9.5	81	10.4	8.1-13.1	97	7.3	5.9-8.9	95	4.4	3.6-5.4	51	6.5	4.8-8.7	44	3.3	2.4-4.4
2009	202	8.9	7.7-10.3	106	12.1	9.8-14.7	96	7.1	5.8-8.7	83	3.8	3.0-4.7	43	4.9	3.5-6.6	40	3.0	2.1-4.1
2010	256	11.1	9.7-12.6	133	14.7	12.2-17.5	123	9.0	7.5-10.8	102	4.6	3.7-5.6	51	6.3	4.6-8.4	51	3.8	2.8-5.0
2011	233	10.0	8.7-11.4	118	12.1	9.9-14.5	115	8.5	7.0-10.2	112	4.8	3.9-5.7	57	5.9	4.4-7.7	55	3.9	3.0-5.2
2012	221	9.1	8.0-10.5	116	12.0	9.9-14.6	105	7.4	6.0-9.0	90	4.0	3.2-4.9	46	5.3	3.8-7.2	44	3.2	2.3-4.3
2013	262	10.6	9.3-12.0	124	12.3	10.2-14.8	138	9.6	8.0-11.3	92	3.8	3.1-4.7	57	6.3	4.7-8.2	35	2.3	1.6-3.3
2014	295	11.8	10.5-13.2	150	14.8	12.4-17.5	145	9.9	8.3-11.7	101	4.3	3.4-5.2	52	5.7	4.2-7.5	49	3.4	2.5-4.5
2015	239	9.6	8.4-10.9	115	11.8	9.7-14.3	124	8.3	6.8-9.9	113	4.6	3.8-5.6	55	5.6	4.2-7.3	58	4.0	3.0-5.2
2016	263	10.3	9.0-11.6	129	12.5	10.4-15.0	134	8.9	7.4-10.5	128	4.9	4.0-5.8	64	6.2	4.8-8.0	64	4.0	3.1-5.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
All Invasive Malignant Tumors

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	1387	426.5	402.1-452.0	713	521.6	477.8-568.0	674	362.4	333.5-392.9	772	251.2	231.8-271.6	435	330.5	294.7-369.0	337	195.2	173.3-219.0
1977	1500	449.7	424.9-475.3	784	558.1	514.0-604.7	716	374.1	345.2-404.6	867	275.8	255.9-296.7	517	397.0	358.2-438.5	350	192.6	171.5-215.4
1978	1581	463.4	438.7-489.0	837	597.1	550.9-645.8	744	376.1	347.9-405.8	884	266.5	247.6-286.3	517	371.4	335.3-410.0	367	194.0	173.4-216.2
1979	1574	451.6	427.5-476.6	827	577.4	532.6-624.5	747	369.3	341.7-398.5	945	285.6	266.0-306.2	531	386.8	349.8-426.2	414	217.0	195.2-240.4
1980	1717	468.4	444.7-493.0	919	612.6	568.2-659.3	798	374.2	347.3-402.5	965	273.8	255.4-293.0	545	379.6	344.0-417.5	420	204.4	184.3-225.9
1981	1670	447.8	424.9-471.6	906	604.8	560.4-651.3	764	348.7	323.2-375.6	1034	291.9	272.9-311.7	590	416.5	378.8-456.5	444	213.0	192.6-234.7
1982	1645	418.2	396.9-440.3	887	558.6	518.2-601.1	758	327.9	304.0-353.1	1009	267.7	250.3-285.9	579	383.6	349.1-420.2	430	193.6	175.0-213.5
1983	1718	427.1	405.7-449.2	925	570.5	529.7-613.4	793	337.4	313.4-362.7	1030	266.2	249.2-284.0	603	390.0	355.8-426.3	427	187.3	169.3-206.5
1984	1813	438.5	417.2-460.5	980	577.6	537.6-619.4	833	347.8	323.6-373.2	1061	269.9	252.8-287.7	593	369.7	336.8-404.6	468	205.1	186.3-225.3
1985	1893	446.4	425.3-468.2	977	563.2	524.3-604.0	916	373.0	348.3-398.8	1052	258.3	242.0-275.3	582	347.4	316.7-380.0	470	199.3	181.0-218.7
1986	1964	454.8	433.7-476.6	1038	597.8	557.5-640.1	926	367.4	343.3-392.6	1153	279.2	262.3-296.7	655	397.4	363.6-433.2	498	206.6	188.3-226.0
1987	1897	428.0	407.8-448.8	972	550.8	512.6-590.9	925	355.6	332.3-379.9	1108	263.8	247.6-280.7	629	387.5	353.9-423.1	479	191.3	174.0-209.7
1988	2042	448.4	428.1-469.4	1039	573.4	534.9-613.6	1003	376.0	352.5-400.6	1172	274.5	258.2-291.6	660	404.5	370.5-440.4	512	199.3	181.9-217.8
1989	2145	451.0	431.0-471.6	1134	593.7	555.4-633.8	1011	365.4	342.6-389.2	1228	275.4	259.4-292.0	691	395.3	362.9-429.4	537	203.1	185.9-221.5
1990	2251	465.7	445.6-486.5	1174	618.1	578.8-659.2	1077	377.8	355.0-401.7	1209	263.3	247.9-279.4	661	362.1	331.6-394.4	548	203.6	186.5-221.8
1991	2387	472.0	452.2-492.3	1297	648.3	609.0-689.3	1090	366.9	344.8-389.9	1242	263.2	248.0-279.1	683	377.2	346.0-410.3	559	197.1	180.7-214.5
1992	2514	482.7	463.0-502.9	1406	686.8	646.7-728.5	1108	362.1	340.5-384.5	1247	255.5	240.8-270.8	651	341.9	312.8-372.6	596	205.5	189.0-223.1
1993	2650	493.4	473.9-513.5	1479	690.8	651.9-731.3	1171	371.8	350.3-394.2	1274	253.5	239.1-268.5	707	364.3	334.7-395.6	567	189.1	173.5-205.6
1994	2673	483.6	464.6-503.2	1445	662.3	624.9-701.3	1228	376.7	355.5-398.8	1339	260.9	246.5-275.9	772	391.4	361.1-423.4	567	184.4	169.2-200.4
1995	2851	501.1	482.0-520.7	1584	691.5	654.1-730.3	1267	381.7	360.5-403.8	1319	249.7	235.8-264.1	708	346.1	318.2-375.6	611	194.0	178.6-210.2
1996	3101	522.6	503.6-542.1	1727	713.3	676.7-751.2	1374	399.2	377.9-421.3	1289	235.6	222.4-249.3	679	317.2	291.2-344.6	610	187.7	172.8-203.4
1997	3069	510.5	491.9-529.6	1695	698.9	662.6-736.5	1374	392.0	371.2-413.7	1323	235.5	222.5-249.0	727	332.5	306.2-360.2	596	178.4	164.2-193.6
1998	3256	520.5	502.1-539.4	1769	692.2	657.2-728.4	1487	410.3	389.3-432.0	1393	243.9	230.8-257.5	747	333.8	307.9-361.1	646	190.5	175.9-206.0
1999	3266	511.1	493.1-529.6	1779	672.7	638.9-707.6	1487	404.1	383.5-425.6	1332	227.1	214.6-240.1	695	300.3	276.0-326.0	637	184.0	169.8-199.1
2000	3243	495.1	477.6-513.1	1821	680.0	646.3-714.8	1422	374.5	355.0-394.9	1446	236.6	224.1-249.5	760	321.0	296.3-347.1	686	188.9	174.8-203.8
2001	3432	504.6	487.2-522.4	1940	692.7	659.4-727.1	1492	380.9	361.5-401.0	1352	216.9	205.0-229.1	684	287.8	264.4-312.5	668	178.9	165.3-193.1
2002	3549	504.4	487.4-521.9	1963	686.3	653.4-720.2	1586	388.5	369.3-408.4	1426	218.5	206.9-230.6	752	294.6	271.6-318.8	674	175.4	162.2-189.4
2003	3587	495.9	479.2-513.0	1984	665.1	633.6-697.6	1603	385.5	366.6-405.2	1413	211.4	200.1-223.1	740	278.2	256.5-301.0	673	170.7	157.8-184.2
2004	3719	497.0	480.6-513.8	1974	631.1	601.2-662.1	1745	408.4	389.1-428.3	1418	209.2	198.0-220.7	745	280.0	258.3-302.8	673	167.3	154.7-180.6
2005	3759	495.1	478.8-511.8	2026	640.7	610.8-671.6	1733	398.6	379.8-418.1	1431	202.0	191.3-213.2	747	261.4	241.3-282.6	684	164.5	152.2-177.4
2006	3925	498.1	482.2-514.5	2172	658.1	628.6-688.6	1753	390.7	372.3-409.7	1309	179.3	169.4-189.6	653	220.5	202.5-239.6	656	153.5	141.8-165.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
All Invasive Malignant Tumors

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	4070	500.8	485.1-517.0	2251	658.0	629.1-687.7	1819	393.3	375.1-412.0	1454	194.2	184.0-204.7	719	240.4	221.7-260.1	735	167.3	155.3-180.0
2008	4203	499.1	483.7-515.0	2282	635.4	607.7-663.9	1921	405.1	386.9-423.9	1391	177.4	167.8-187.2	693	217.3	200.2-235.5	698	152.8	141.5-164.7
2009	4305	496.5	481.3-512.1	2326	616.6	590.1-644.0	1979	408.8	390.7-427.5	1471	183.6	174.0-193.5	759	235.3	217.5-254.1	712	151.9	140.8-163.6
2010	4320	482.1	467.4-497.2	2387	624.1	597.4-651.6	1933	384.3	367.1-402.1	1485	180.6	171.2-190.3	782	240.3	222.4-259.1	703	145.0	134.3-156.3
2011	4600	496.3	481.6-511.3	2512	628.4	602.4-655.3	2088	403.1	385.7-421.0	1531	179.0	169.9-188.5	777	220.7	204.3-238.0	754	152.6	141.7-164.0
2012	4413	460.9	447.0-475.2	2334	562.8	538.7-587.7	2079	390.8	373.9-408.2	1540	174.9	166.0-184.2	791	220.4	204.1-237.5	749	147.3	136.8-158.4
2013	4539	463.9	450.1-478.0	2360	555.0	531.4-579.3	2179	398.9	382.1-416.3	1527	168.4	159.8-177.4	765	208.1	192.6-224.4	762	143.5	133.4-154.3
2014	4737	465.3	451.7-479.1	2495	564.5	541.2-588.5	2242	397.1	380.6-414.2	1529	162.6	154.3-171.2	774	204.2	189.2-220.1	755	137.0	127.3-147.3
2015	4935	472.6	459.2-486.4	2567	565.3	542.4-589.0	2368	409.8	393.2-426.9	1506	156.9	148.8-165.2	748	188.9	174.7-203.8	758	135.9	126.2-146.1
2016	4910	459.3	446.2-472.7	2527	541.9	519.7-564.7	2383	403.1	386.8-419.9	1318	132.6	125.3-140.2	672	163.1	150.2-176.7	646	112.5	103.8-121.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Oral cavity and pharynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	56	13.7	10.3-18.0	38	20.9	14.6-29.1	18	7.7	4.5-12.5	19	6.6	3.7-10.6	13	13.2	5.9-24.0	6	2.6	1.0-5.9
1977	50	12.4	9.1-16.5	39	22.2	15.4-30.9	11	4.8	2.3-8.7	24	6.1	3.8-9.1	17	9.0	5.2-14.8	7	3.5	1.4-7.4
1978	63	14.9	11.4-19.2	38	20.6	14.4-28.7	25	10.4	6.7-15.5	26	6.8	4.4-10.0	20	12.4	7.4-19.5	6	2.5	0.9-5.6
1979	58	14.1	10.6-18.3	47	26.0	18.8-35.1	11	4.7	2.3-8.5	39	9.5	6.7-13.2	31	17.9	11.7-26.0	8	3.2	1.4-6.5
1980	64	14.2	10.9-18.3	42	21.4	15.2-29.3	22	8.5	5.3-13.0	37	9.1	6.3-12.6	29	16.9	11.1-24.6	8	3.3	1.4-6.5
1981	79	17.8	13.9-22.3	52	26.1	18.9-35.1	27	11.2	7.3-16.4	32	7.5	5.1-10.7	22	11.7	7.2-18.0	10	4.3	2.0-7.9
1982	65	14.3	11.0-18.3	44	21.5	15.5-29.2	21	8.4	5.2-13.0	35	7.5	5.2-10.5	27	13.4	8.6-19.8	8	2.9	1.2-5.8
1983	74	16.4	12.7-20.8	55	28.2	20.6-37.5	19	7.5	4.5-11.9	35	7.7	5.3-10.9	24	12.3	7.4-19.1	11	4.4	2.2-7.9
1984	76	16.0	12.6-20.2	56	27.7	20.6-36.4	20	7.3	4.4-11.3	30	6.7	4.5-9.6	24	12.5	7.7-18.9	6	2.4	0.9-5.2
1985	78	17.1	13.4-21.5	57	29.0	21.5-38.1	21	8.2	5.0-12.5	34	7.1	4.8-10.0	25	12.0	7.5-18.0	9	3.3	1.5-6.4
1986	70	14.4	11.1-18.3	49	23.5	16.9-31.7	21	7.5	4.6-11.6	35	7.8	5.3-11.0	25	12.8	7.9-19.4	10	4.0	1.9-7.3
1987	80	16.5	13.0-20.7	50	23.0	16.8-30.7	30	11.3	7.6-16.2	29	6.2	4.1-9.1	20	10.4	6.0-16.6	9	3.4	1.5-6.4
1988	75	14.5	11.3-18.2	53	24.0	17.5-32.0	22	7.4	4.6-11.2	37	7.3	5.1-10.1	25	12.0	7.3-18.4	12	4.2	2.1-7.2
1989	109	20.2	16.5-24.5	78	32.8	25.6-41.5	31	10.3	6.9-14.7	43	8.0	5.8-10.9	29	12.0	7.9-17.4	14	4.8	2.6-8.1
1990	68	12.9	9.9-16.5	53	23.5	17.1-31.4	15	5.1	2.8-8.4	27	4.8	3.2-7.0	18	6.9	4.1-11.1	9	3.0	1.4-5.7
1991	91	16.6	13.3-20.5	71	30.2	23.3-38.4	20	6.4	3.9-9.9	35	6.7	4.6-9.3	24	10.5	6.5-16.0	11	3.8	1.9-6.7
1992	87	15.1	11.9-18.7	57	23.7	17.2-31.6	30	9.2	6.1-13.1	38	6.6	4.6-9.1	31	12.2	8.1-17.5	7	2.2	0.9-4.5
1993	83	14.1	11.2-17.6	57	21.8	16.2-28.7	26	8.1	5.2-11.8	20	3.5	2.1-5.4	13	5.8	2.9-10.2	7	2.0	0.8-4.2
1994	75	11.9	9.3-15.0	54	19.6	14.4-25.9	21	5.9	3.6-9.0	28	4.6	3.0-6.7	23	8.9	5.5-13.6	5	1.4	0.5-3.3
1995	83	13.7	10.8-17.1	62	22.3	16.8-29.0	21	6.6	4.1-10.1	40	6.4	4.5-8.8	30	10.5	7.0-15.2	10	3.0	1.4-5.5
1996	94	15.7	12.5-19.3	67	27.2	20.6-35.2	27	7.7	5.0-11.2	27	4.6	3.0-6.7	17	6.7	3.7-10.9	10	3.0	1.4-5.4
1997	106	15.7	12.8-19.0	71	23.9	18.4-30.5	35	9.3	6.5-13.0	24	4.8	3.0-7.1	17	9.4	5.1-15.4	7	2.3	0.9-4.6
1998	85	12.4	9.8-15.4	56	18.4	13.6-24.3	29	7.7	5.1-11.0	27	4.1	2.7-6.0	25	8.6	5.4-13.0	2	0.6	0.1-2.0
1999	82	12.1	9.5-15.0	62	20.8	15.5-27.1	20	5.5	3.3-8.5	32	4.8	3.2-6.8	22	8.3	4.9-12.9	10	2.5	1.2-4.6
2000	71	10.2	7.9-13.0	45	15.4	11.0-20.9	26	6.5	4.2-9.5	38	5.8	4.1-8.1	26	9.0	5.7-13.5	12	3.4	1.7-5.8
2001	63	8.5	6.5-10.9	50	14.3	10.5-19.0	13	3.5	1.8-5.9	16	2.7	1.5-4.4	9	3.7	1.5-7.4	7	2.1	0.8-4.1
2002	93	13.2	10.5-16.2	63	19.9	14.9-25.9	30	7.8	5.2-11.2	26	3.8	2.4-5.6	20	6.7	3.9-10.7	6	1.7	0.6-3.6
2003	80	10.3	8.1-12.9	50	14.0	10.1-18.8	30	7.2	4.8-10.3	29	4.1	2.7-5.9	23	6.9	4.3-10.5	6	1.7	0.6-3.5
2004	75	9.2	7.2-11.6	54	14.7	10.9-19.4	21	4.8	2.9-7.3	31	3.9	2.6-5.6	25	7.1	4.5-10.6	6	1.4	0.5-3.1
2005	67	8.4	6.4-10.7	45	13.0	9.3-17.6	22	4.9	3.0-7.4	20	2.8	1.7-4.3	16	5.2	2.9-8.6	4	1.0	0.3-2.5
2006	70	8.2	6.3-10.4	52	13.9	10.2-18.5	18	3.8	2.2-6.0	22	2.8	1.7-4.2	18	5.2	3.0-8.4	4	0.9	0.2-2.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Oral cavity and pharynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	84	9.7	7.7-12.1	62	16.3	12.2-21.2	22	4.7	2.9-7.2	23	2.9	1.8-4.4	15	4.8	2.4-8.1	8	1.8	0.7-3.5
2008	87	9.9	7.9-12.3	53	12.9	9.5-17.2	34	7.1	4.9-10.0	19	2.2	1.3-3.5	12	3.0	1.5-5.4	7	1.5	0.6-3.1
2009	88	9.5	7.5-11.8	55	13.3	9.8-17.6	33	6.4	4.4-9.1	17	1.9	1.1-3.1	12	3.3	1.6-5.9	5	1.0	0.3-2.4
2010	91	9.2	7.3-11.3	62	14.2	10.7-18.4	29	5.3	3.5-7.6	17	1.8	1.0-3.0	12	2.7	1.3-4.8	5	1.0	0.3-2.4
2011	77	7.9	6.2-9.9	60	13.2	9.9-17.2	17	3.4	2.0-5.5	27	2.9	1.9-4.2	18	4.1	2.4-6.6	9	1.8	0.8-3.4
2012	76	7.4	5.8-9.4	41	9.5	6.6-13.1	35	6.2	4.3-8.7	22	2.3	1.4-3.5	18	4.3	2.4-6.9	4	0.8	0.2-1.9
2013	67	6.4	4.9-8.2	44	9.3	6.7-12.6	23	4.1	2.5-6.1	22	2.2	1.3-3.3	16	3.6	2.0-5.9	6	1.1	0.4-2.3
2014	85	8.1	6.4-10.1	60	12.7	9.6-16.5	25	4.4	2.8-6.6	19	1.8	1.1-2.8	12	2.5	1.2-4.3	7	1.2	0.5-2.5
2015	79	7.4	5.8-9.3	60	13.0	9.7-16.9	19	3.2	1.9-5.0	20	2.0	1.2-3.1	12	2.7	1.4-4.8	8	1.4	0.6-2.8
2016	95	8.7	7.0-10.7	59	12.0	9.0-15.8	36	6.2	4.3-8.6	17	1.7	0.9-2.7	12	2.8	1.4-4.9	5	0.8	0.3-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Esophagus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	41	12.4	8.7-17.1	27	18.6	11.9-27.6	14	7.5	3.9-12.9	48	14.1	10.0-19.1	37	25.5	16.8-36.8	11	5.6	2.6-10.4
1977	51	13.6	9.9-18.1	37	22.3	15.4-31.2	14	6.7	3.5-11.5	48	13.0	9.5-17.4	38	24.7	17.0-34.5	10	4.3	2.1-8.1
1978	66	18.5	14.0-23.9	50	30.2	21.7-40.9	16	9.1	5.0-15.1	54	14.2	10.5-18.9	41	24.3	16.7-34.1	13	6.4	3.2-11.1
1979	49	14.4	10.4-19.3	41	29.0	20.0-40.4	8	4.0	1.7-8.0	42	11.9	8.4-16.3	36	25.3	16.8-36.3	6	2.6	0.9-5.7
1980	57	14.7	11.0-19.2	44	26.2	18.6-35.8	13	6.0	3.1-10.4	47	12.8	9.2-17.3	40	25.4	17.4-35.5	7	3.7	1.4-7.6
1981	55	13.4	10.0-17.6	42	23.8	16.6-33.0	13	5.7	3.0-9.8	45	10.8	7.8-14.6	34	19.6	13.2-28.0	11	4.4	2.2-7.9
1982	41	10.3	7.3-14.2	33	20.5	13.4-29.6	8	3.4	1.4-6.6	46	11.9	8.6-16.0	38	23.2	15.9-32.6	8	3.8	1.6-7.5
1983	52	12.0	8.8-15.9	41	23.3	16.0-32.5	11	4.2	2.1-7.5	49	12.2	8.8-16.3	38	23.1	15.5-32.8	11	4.6	2.3-8.3
1984	54	12.3	9.1-16.1	40	20.7	14.5-28.7	14	5.8	3.1-9.8	47	11.6	8.4-15.6	33	20.8	13.6-30.1	14	5.7	3.1-9.6
1985	52	12.3	9.0-16.2	42	24.3	16.8-33.6	10	4.0	1.9-7.4	45	9.9	7.1-13.3	37	18.4	12.6-25.9	8	3.2	1.4-6.4
1986	66	15.0	11.4-19.2	49	27.3	19.5-37.0	17	6.4	3.7-10.4	53	12.7	9.3-16.7	38	23.8	15.9-33.7	15	5.8	3.2-9.6
1987	44	9.6	6.9-13.0	31	15.8	10.4-22.9	13	5.0	2.6-8.6	41	9.1	6.4-12.4	32	16.4	10.7-23.7	9	3.7	1.7-7.1
1988	52	10.8	8.0-14.2	37	18.4	12.6-25.7	15	5.5	3.0-9.0	54	12.0	8.9-15.8	37	20.7	14.0-29.2	17	6.4	3.7-10.2
1989	65	13.7	10.4-17.5	48	25.0	17.8-34.0	17	6.1	3.6-9.8	42	8.3	5.9-11.2	35	16.0	10.8-22.7	7	2.5	1.0-5.1
1990	57	10.8	8.1-14.1	39	17.6	12.1-24.7	18	6.0	3.5-9.5	56	11.6	8.6-15.1	37	19.3	13.0-27.2	19	6.7	4.0-10.5
1991	57	11.0	8.2-14.3	42	19.8	13.8-27.2	15	4.8	2.7-8.0	49	9.1	6.7-12.1	35	15.2	10.3-21.4	14	4.6	2.5-7.7
1992	52	10.1	7.5-13.3	42	19.8	13.7-27.4	10	3.5	1.6-6.3	38	7.6	5.3-10.5	27	13.3	8.3-19.9	11	3.8	1.9-6.7
1993	53	9.1	6.7-11.9	38	15.1	10.4-21.2	15	4.5	2.5-7.5	44	7.9	5.7-10.7	39	16.8	11.5-23.5	5	1.6	0.5-3.7
1994	57	10.0	7.5-13.0	37	14.8	10.2-20.6	20	6.3	3.8-9.8	44	8.1	5.8-10.9	34	15.2	10.3-21.5	10	3.2	1.5-5.8
1995	43	7.4	5.3-10.1	28	10.9	7.1-16.1	15	4.7	2.6-7.7	48	8.3	6.1-11.1	35	13.8	9.4-19.4	13	4.1	2.2-6.9
1996	57	9.7	7.3-12.7	44	17.4	12.3-23.8	13	4.1	2.2-6.9	43	7.3	5.2-9.9	29	11.4	7.4-16.7	14	4.2	2.3-7.1
1997	45	7.7	5.5-10.3	32	13.9	9.1-20.1	13	3.7	2.0-6.3	38	6.3	4.4-8.7	24	9.8	6.0-15.0	14	3.9	2.1-6.6
1998	54	9.1	6.8-11.9	38	14.7	10.2-20.5	16	4.9	2.8-7.8	52	8.7	6.5-11.5	36	13.7	9.3-19.3	16	5.0	2.8-8.1
1999	64	9.9	7.5-12.6	47	17.5	12.6-23.6	17	4.5	2.6-7.2	31	5.2	3.4-7.4	21	8.6	5.0-13.4	10	2.9	1.4-5.3
2000	53	8.5	6.3-11.2	35	13.2	8.9-18.6	18	5.1	3.0-8.0	54	8.7	6.5-11.4	38	15.4	10.6-21.4	16	4.3	2.4-6.9
2001	52	7.9	5.8-10.4	35	13.7	9.2-19.5	17	4.3	2.5-6.8	36	5.4	3.7-7.5	25	9.1	5.6-13.7	11	2.9	1.4-5.1
2002	40	5.9	4.1-8.0	29	10.3	6.6-15.2	11	2.8	1.4-5.0	51	7.0	5.2-9.3	39	12.7	8.7-17.7	12	2.9	1.5-5.1
2003	56	8.2	6.1-10.7	43	14.1	10.0-19.3	13	3.6	1.9-6.2	37	5.6	3.9-7.8	26	9.6	5.9-14.6	11	3.0	1.5-5.4
2004	50	7.3	5.4-9.7	31	10.0	6.7-14.3	19	5.1	3.0-7.9	43	6.3	4.5-8.6	32	11.9	7.8-17.1	11	2.8	1.4-5.0
2005	48	6.1	4.4-8.1	36	10.6	7.2-14.9	12	2.7	1.4-4.7	36	5.3	3.7-7.3	24	8.0	5.0-12.0	12	3.1	1.6-5.3
2006	38	5.3	3.7-7.3	27	11.0	6.9-16.3	11	2.4	1.2-4.3	25	3.5	2.2-5.2	19	7.4	4.2-11.7	6	1.3	0.5-2.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Esophagus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	48	5.9	4.3-7.9	36	9.9	6.8-13.8	12	2.7	1.4-4.8	33	4.5	3.0-6.3	23	7.7	4.7-11.7	10	2.4	1.1-4.3
2008	37	4.4	3.0-6.0	27	7.9	5.0-11.8	10	2.1	1.0-3.8	36	4.2	2.9-5.9	22	5.8	3.5-9.0	14	3.1	1.7-5.2
2009	43	5.0	3.5-6.7	34	9.5	6.3-13.6	9	2.0	0.9-3.7	28	3.4	2.2-5.0	23	7.4	4.5-11.4	5	1.0	0.3-2.4
2010	42	4.9	3.5-6.7	32	9.2	6.1-13.2	10	2.1	1.0-3.8	43	4.9	3.5-6.6	32	8.4	5.5-12.1	11	2.4	1.2-4.3
2011	48	5.4	3.9-7.2	32	8.6	5.6-12.4	16	3.2	1.8-5.1	26	3.0	1.9-4.4	19	5.3	3.0-8.4	7	1.5	0.6-3.0
2012	42	4.2	3.0-5.7	31	6.7	4.5-9.7	11	2.0	1.0-3.7	32	3.5	2.4-5.0	22	5.3	3.2-8.2	10	2.0	1.0-3.7
2013	48	5.2	3.8-7.0	34	9.0	6.1-12.7	14	2.7	1.4-4.5	39	4.2	3.0-5.8	26	6.5	4.1-9.6	13	2.5	1.3-4.4
2014	36	3.9	2.7-5.4	23	6.2	3.8-9.3	13	2.3	1.2-4.0	38	4.1	2.9-5.7	25	6.6	4.1-9.8	13	2.4	1.3-4.2
2015	38	3.8	2.6-5.2	28	6.7	4.3-9.8	10	1.7	0.8-3.2	23	2.3	1.4-3.5	18	4.5	2.5-7.2	5	0.8	0.2-1.9
2016	42	4.6	3.3-6.2	26	6.6	4.2-9.7	16	3.0	1.7-4.9	34	3.8	2.6-5.3	26	6.9	4.4-10.2	8	1.6	0.7-3.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Stomach

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	65	22.9	17.1-29.7	38	32.3	21.4-46.2	27	16.3	10.3-24.2	41	13.9	9.6-19.4	29	24.3	15.1-36.4	12	6.8	3.3-12.1
1977	57	18.1	13.4-23.9	34	25.1	16.6-36.1	23	12.9	7.9-19.8	51	17.5	12.6-23.4	31	25.5	16.3-37.4	20	12.1	7.0-18.9
1978	68	23.8	18.0-30.6	47	39.8	27.9-54.6	21	13.0	7.7-20.0	50	17.2	12.4-23.2	25	21.5	12.8-33.3	25	14.6	9.1-21.8
1979	67	21.5	16.2-27.8	31	24.0	14.9-35.9	36	20.3	13.9-28.3	42	13.9	9.7-19.1	23	18.9	11.0-29.7	19	10.8	6.3-17.0
1980	78	25.5	19.8-32.2	41	31.8	21.7-44.4	37	21.3	14.7-29.5	47	16.0	11.5-21.5	22	19.5	11.6-30.1	25	13.9	8.8-20.7
1981	61	19.1	14.3-24.8	40	29.8	20.1-42.1	21	12.0	7.3-18.4	62	20.0	15.1-26.0	38	27.0	18.2-38.0	24	14.8	9.3-22.0
1982	50	14.5	10.6-19.3	33	24.1	15.9-34.5	17	8.3	4.7-13.3	41	12.8	9.0-17.5	24	18.9	11.4-28.8	17	9.1	5.2-14.5
1983	61	16.0	12.0-20.7	33	21.4	14.2-30.8	28	12.5	8.1-18.3	41	11.2	7.8-15.4	23	15.7	9.4-24.4	18	8.4	4.8-13.4
1984	63	16.2	12.3-21.0	41	25.1	17.3-35.0	22	10.1	6.2-15.3	45	12.8	9.1-17.2	29	23.7	15.2-34.8	16	6.6	3.7-10.8
1985	63	16.5	12.4-21.4	38	21.6	14.4-30.8	25	12.3	7.8-18.1	46	11.3	8.2-15.2	25	13.1	8.2-19.7	21	9.5	5.8-14.6
1986	50	12.8	9.3-17.1	31	18.7	11.9-27.6	19	8.8	5.2-13.7	44	11.5	8.1-15.6	28	18.0	11.2-27.0	16	7.3	4.1-11.8
1987	76	19.7	15.3-24.8	46	27.9	19.6-38.2	30	13.9	9.3-19.8	46	11.5	8.2-15.5	27	14.9	9.5-22.0	19	8.6	5.1-13.5
1988	71	18.6	14.4-23.7	44	29.6	20.6-40.6	27	12.1	7.9-17.5	46	13.0	9.3-17.4	23	17.4	10.3-26.8	23	10.6	6.7-15.9
1989	74	17.3	13.4-21.9	43	22.7	15.6-31.5	31	13.1	8.8-18.6	58	13.8	10.3-17.9	39	22.2	15.2-31.1	19	8.0	4.7-12.4
1990	75	17.3	13.4-21.9	52	31.3	22.3-42.4	23	9.0	5.6-13.5	48	11.1	8.0-14.9	31	16.7	10.5-24.8	17	7.3	4.2-11.6
1991	69	14.8	11.3-18.8	38	20.8	13.9-29.6	31	11.3	7.6-16.1	54	11.4	8.4-15.1	36	19.5	12.9-28.0	18	6.5	3.8-10.2
1992	72	14.7	11.3-18.7	36	18.4	12.2-26.4	36	12.6	8.7-17.4	61	12.3	9.3-15.9	24	11.0	6.8-16.6	37	12.9	9.0-17.8
1993	85	18.4	14.6-23.0	54	27.3	19.8-36.6	31	12.0	8.1-17.0	44	9.6	6.8-12.9	29	15.0	9.4-22.4	15	5.9	3.3-9.7
1994	80	15.4	12.0-19.3	54	27.0	19.3-36.3	26	8.5	5.5-12.5	56	11.1	8.2-14.5	39	21.0	14.1-29.8	17	5.7	3.3-9.1
1995	85	16.7	13.2-20.8	41	20.8	14.2-29.0	44	14.3	10.3-19.2	53	10.7	7.9-14.1	31	16.6	10.6-24.3	22	7.3	4.5-11.0
1996	79	15.0	11.8-18.8	50	24.9	17.8-33.6	29	9.1	6.0-13.0	36	7.2	4.9-10.0	26	13.6	8.3-20.6	10	3.4	1.6-6.2
1997	74	12.7	9.9-16.1	46	19.8	13.9-27.0	28	8.4	5.5-12.1	38	6.6	4.6-9.1	27	11.7	7.5-17.2	11	3.2	1.6-5.7
1998	89	15.2	12.1-18.8	56	22.5	16.4-30.0	33	9.9	6.7-13.9	56	10.3	7.7-13.4	37	16.4	11.1-23.2	19	6.0	3.6-9.4
1999	80	13.9	10.9-17.4	50	22.5	15.9-30.5	30	9.1	6.1-13.0	56	10.0	7.4-13.0	35	16.3	10.7-23.5	21	6.4	3.9-9.7
2000	76	12.7	9.9-16.0	41	16.0	11.1-22.0	35	10.0	6.9-13.9	47	8.3	6.1-11.1	23	10.6	6.5-16.1	24	7.0	4.5-10.4
2001	81	13.5	10.6-16.8	42	18.1	12.5-25.1	39	11.0	7.8-15.0	36	5.7	3.9-7.9	24	8.4	5.1-13.0	12	3.5	1.8-6.1
2002	77	11.9	9.3-14.9	45	16.7	11.7-22.9	32	8.6	5.8-12.1	51	8.6	6.3-11.3	31	13.6	8.7-19.9	20	5.7	3.5-8.8
2003	71	10.0	7.7-12.7	45	14.9	10.5-20.3	26	6.6	4.3-9.7	38	5.4	3.8-7.4	28	9.3	5.9-13.9	10	2.7	1.3-4.9
2004	72	10.9	8.4-13.8	33	13.2	8.8-18.8	39	9.7	6.8-13.2	39	5.8	4.1-8.0	20	7.9	4.6-12.5	19	4.7	2.8-7.4
2005	101	14.3	11.6-17.5	45	15.3	10.8-20.9	56	13.6	10.2-17.7	64	10.1	7.7-12.9	39	15.0	10.3-20.9	25	6.7	4.3-9.9
2006	100	14.1	11.4-17.2	60	20.9	15.6-27.3	40	9.6	6.9-13.1	45	6.5	4.7-8.7	28	9.4	6.0-13.8	17	4.4	2.5-6.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Stomach

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	97	13.2	10.6-16.2	52	18.0	13.1-24.0	45	10.3	7.5-13.8	51	7.4	5.4-9.7	29	10.8	6.9-15.9	22	5.4	3.4-8.1
2008	79	10.2	8.0-12.8	51	16.1	11.7-21.4	28	6.0	4.0-8.7	57	7.6	5.7-9.9	33	11.2	7.5-16.0	24	5.3	3.4-7.9
2009	94	12.0	9.6-14.7	52	15.7	11.4-20.8	42	9.4	6.7-12.7	43	5.9	4.2-7.9	23	8.2	5.0-12.5	20	4.5	2.7-6.9
2010	84	10.4	8.2-12.9	50	14.5	10.4-19.4	34	7.6	5.2-10.6	46	6.0	4.3-8.0	32	10.5	7.0-15.0	14	3.1	1.7-5.1
2011	111	13.1	10.7-15.8	63	17.0	12.8-22.0	48	10.1	7.4-13.5	43	5.2	3.7-7.0	25	7.0	4.4-10.5	18	3.9	2.3-6.1
2012	103	12.1	9.8-14.7	58	15.3	11.4-20.0	45	9.2	6.7-12.3	50	6.3	4.6-8.3	29	8.4	5.4-12.3	21	4.6	2.8-7.0
2013	101	11.0	8.9-13.4	52	13.7	10.0-18.2	49	9.3	6.8-12.3	52	5.9	4.3-7.7	31	8.2	5.4-11.9	21	4.1	2.5-6.3
2014	113	11.5	9.5-13.9	71	16.9	13.0-21.6	42	7.6	5.5-10.3	55	6.0	4.5-7.8	38	10.1	7.0-14.1	17	3.3	1.9-5.2
2015	125	12.9	10.7-15.4	71	17.2	13.2-22.0	54	9.9	7.4-12.9	53	5.6	4.2-7.4	38	10.4	7.2-14.4	15	2.6	1.5-4.4
2016	115	11.5	9.4-13.9	63	14.4	10.9-18.6	52	9.3	6.9-12.3	37	3.5	2.5-4.9	23	4.9	3.0-7.5	14	2.4	1.3-4.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Colorectal

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	160	54.0	45.1-64.0	64	48.3	35.4-63.9	96	57.8	46.0-71.4	82	30.2	23.2-38.3	35	28.6	18.5-41.8	47	30.7	21.9-41.6
1977	188	62.3	52.9-72.8	81	64.9	49.7-82.8	107	60.7	49.0-74.2	96	33.5	26.6-41.5	51	41.0	29.5-55.0	45	27.8	19.7-37.6
1978	193	60.5	51.5-70.4	87	66.5	51.1-84.6	106	57.2	46.3-69.8	98	33.8	26.9-41.7	54	47.9	34.1-64.7	44	25.4	18.2-34.4
1979	201	62.2	53.2-72.2	103	72.4	57.3-89.9	98	54.5	43.7-67.0	121	40.8	33.2-49.4	45	34.9	24.3-48.1	76	44.1	34.2-55.8
1980	219	68.6	59.0-79.0	97	78.6	61.6-98.2	122	63.3	52.0-76.1	107	32.8	26.5-40.1	51	35.8	25.7-48.1	56	30.2	22.5-39.6
1981	207	63.5	54.6-73.3	97	75.7	59.7-94.1	110	56.4	45.9-68.4	126	39.8	32.7-47.8	61	48.3	35.9-63.2	65	34.3	26.1-44.1
1982	193	53.6	45.9-62.3	86	57.9	44.8-73.1	107	51.3	41.6-62.3	113	34.1	27.7-41.4	48	39.2	27.4-53.5	65	32.2	24.5-41.2
1983	199	57.7	49.4-66.8	86	63.9	49.5-80.8	113	54.6	44.6-65.9	99	28.5	22.8-35.0	46	31.8	22.1-43.8	53	26.3	19.5-34.6
1984	229	62.8	54.5-71.9	113	77.3	62.2-94.6	116	53.9	44.3-65.0	103	28.2	22.7-34.5	43	30.3	20.9-41.9	60	27.5	20.8-35.5
1985	216	57.2	49.4-65.9	101	66.1	52.3-82.0	115	51.7	42.4-62.3	116	32.6	26.6-39.4	49	34.2	24.4-46.0	67	31.6	24.3-40.3
1986	239	60.7	52.8-69.3	116	68.5	55.4-83.6	123	54.2	44.7-64.9	119	32.5	26.6-39.2	56	39.1	28.3-52.2	63	28.7	21.9-36.9
1987	247	60.6	52.9-69.0	126	77.4	62.8-94.1	121	50.5	41.7-60.6	125	31.6	26.0-38.0	71	49.7	37.5-64.0	54	21.9	16.3-28.7
1988	244	57.3	50.0-65.4	109	62.1	49.7-76.4	135	54.3	45.3-64.4	125	32.5	26.7-38.9	64	43.2	32.0-56.6	61	26.5	20.1-34.1
1989	261	59.8	52.4-67.9	124	72.4	58.6-88.1	137	53.2	44.4-63.0	127	32.1	26.5-38.4	75	53.8	41.0-68.7	52	20.9	15.5-27.5
1990	241	54.0	47.0-61.6	111	60.0	48.2-73.7	130	49.5	41.2-59.0	132	30.7	25.4-36.6	64	36.6	27.2-47.9	68	26.8	20.7-34.1
1991	243	50.6	44.2-57.7	119	56.5	45.8-68.9	124	45.5	37.7-54.4	126	28.3	23.3-33.9	64	36.8	27.3-48.3	62	23.3	17.8-29.9
1992	279	58.0	51.1-65.5	136	69.7	57.1-83.9	143	50.8	42.6-59.9	116	25.8	21.1-31.2	53	28.8	20.8-38.6	63	23.6	18.0-30.2
1993	265	53.1	46.6-60.2	124	57.0	46.3-69.2	141	49.3	41.4-58.3	139	29.3	24.4-34.8	70	36.2	27.3-46.7	69	24.8	19.2-31.5
1994	301	59.4	52.6-66.8	161	79.3	66.1-94.0	140	46.9	39.3-55.4	154	30.6	25.8-36.1	78	38.2	29.3-48.8	76	25.7	20.1-32.2
1995	314	59.0	52.4-66.1	146	67.5	55.8-80.7	168	54.0	46.0-62.9	127	25.7	21.2-30.7	63	34.1	25.2-44.7	64	21.4	16.4-27.3
1996	328	59.8	53.2-66.9	167	73.8	62.0-87.0	161	50.2	42.6-58.7	126	25.1	20.7-30.0	60	27.8	20.6-36.4	66	22.2	17.1-28.3
1997	319	56.5	50.3-63.3	157	70.0	58.3-83.0	162	48.5	41.2-56.6	148	27.2	22.8-32.1	78	34.6	26.5-44.2	70	22.2	17.3-28.1
1998	338	57.7	51.5-64.4	158	65.9	55.0-78.1	180	53.0	45.4-61.4	129	22.7	18.8-27.0	65	27.9	20.9-36.2	64	19.2	14.7-24.5
1999	391	65.1	58.6-72.1	193	77.6	66.0-90.6	198	57.3	49.5-65.9	153	28.3	23.8-33.2	73	35.5	27.0-45.5	80	24.3	19.2-30.2
2000	353	57.2	51.2-63.7	173	68.0	57.2-80.0	180	50.6	43.3-58.6	160	26.4	22.3-31.0	84	36.0	27.9-45.5	76	21.1	16.6-26.5
2001	371	57.3	51.4-63.6	192	72.6	61.8-84.5	179	47.3	40.5-54.8	126	20.4	16.9-24.4	62	25.7	19.2-33.4	64	17.6	13.5-22.4
2002	413	62.2	56.2-68.7	187	69.0	58.5-80.7	226	58.4	50.9-66.6	153	24.7	20.8-29.0	69	29.1	21.9-37.6	84	22.6	18.0-28.0
2003	397	59.5	53.6-65.9	205	76.6	65.5-88.9	192	49.5	42.6-57.1	138	21.4	17.9-25.4	67	26.4	19.9-34.1	71	18.4	14.3-23.2
2004	423	60.5	54.7-66.8	218	74.8	64.2-86.5	205	51.2	44.3-58.7	155	23.8	20.1-28.0	79	29.8	23.0-37.8	76	19.9	15.6-24.9
2005	397	55.0	49.5-60.8	199	67.3	57.5-78.1	198	47.0	40.6-54.1	141	20.6	17.2-24.4	67	23.9	18.1-30.8	74	18.1	14.2-22.8
2006	396	52.9	47.6-58.5	194	58.6	49.9-68.3	202	48.0	41.5-55.2	146	20.9	17.5-24.7	74	25.7	19.7-32.8	72	17.6	13.7-22.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Colorectal

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	385	48.9	43.9-54.2	193	55.6	47.4-64.6	192	43.4	37.4-50.0	177	23.8	20.3-27.7	89	28.3	22.3-35.4	88	20.5	16.4-25.3
2008	436	53.8	48.7-59.2	220	64.7	55.8-74.6	216	46.7	40.6-53.5	122	16.0	13.2-19.2	61	19.8	14.8-25.8	61	13.6	10.3-17.4
2009	414	48.4	43.7-53.4	212	57.3	49.3-66.1	202	41.7	36.1-48.0	145	18.0	15.1-21.2	76	23.1	17.9-29.3	69	14.4	11.1-18.2
2010	388	45.3	40.7-50.2	208	57.8	49.7-66.9	180	36.5	31.3-42.3	126	15.0	12.4-18.0	64	18.3	13.7-23.7	62	12.7	9.7-16.3
2011	410	46.1	41.6-50.9	215	55.5	47.8-64.0	195	38.8	33.5-44.7	137	16.2	13.5-19.3	71	20.8	15.9-26.6	66	13.3	10.2-16.9
2012	363	38.7	34.7-43.1	189	46.1	39.3-53.7	174	33.3	28.4-38.7	124	14.8	12.3-17.7	59	18.5	13.8-24.1	65	13.1	10.1-16.8
2013	388	40.6	36.6-45.0	192	46.7	39.9-54.3	196	36.4	31.4-41.9	137	14.6	12.2-17.3	71	17.9	13.8-22.9	66	12.1	9.3-15.4
2014	427	43.4	39.3-47.8	216	51.0	44.0-58.7	211	37.8	32.8-43.3	148	15.7	13.2-18.5	79	19.5	15.1-24.5	69	13.1	10.1-16.6
2015	433	41.8	37.8-46.0	225	49.3	42.7-56.6	208	35.8	31.0-41.1	156	16.4	13.9-19.3	72	17.7	13.6-22.6	84	15.2	12.1-18.9
2016	383	36.7	33.0-40.7	206	46.9	40.4-54.2	177	29.7	25.4-34.5	110	10.7	8.7-12.9	59	12.6	9.4-16.5	51	9.0	6.6-11.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Colon excluding rectum

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	109	37.3	29.8-45.8	41	32.0	21.2-45.7	68	41.0	31.1-52.7	68	24.3	18.1-31.6	30	24.1	14.8-36.6	38	24.1	16.4-33.8
1977	138	46.5	38.3-55.7	55	45.3	32.6-60.9	83	47.7	37.4-59.8	73	25.3	19.4-32.3	39	32.0	21.9-44.7	34	20.5	13.8-29.0
1978	141	43.5	36.1-51.9	64	47.7	35.1-62.9	77	41.1	32.1-51.8	78	26.7	20.6-33.9	43	39.0	26.4-54.7	35	19.7	13.4-27.6
1979	152	47.3	39.5-56.1	75	53.1	40.4-68.3	77	42.7	33.2-53.9	108	36.4	29.3-44.6	41	31.9	21.8-44.5	67	38.9	29.6-49.9
1980	157	48.8	40.8-57.6	67	55.5	41.2-72.4	90	45.8	36.4-56.8	83	25.6	20.0-32.2	41	30.0	20.6-41.7	42	22.5	15.9-30.6
1981	153	48.0	40.2-56.7	71	57.2	43.2-73.7	82	42.8	33.7-53.4	104	32.3	26.0-39.5	51	41.4	29.9-55.5	53	27.0	19.9-35.6
1982	146	40.7	34.0-48.3	69	47.5	35.8-61.5	77	36.7	28.6-46.1	95	29.2	23.3-36.1	38	34.4	23.1-48.4	57	27.7	20.8-36.1
1983	153	44.5	37.3-52.6	60	45.6	33.3-60.2	93	44.8	35.8-55.1	87	25.6	20.2-31.9	39	28.1	18.8-39.7	48	24.1	17.5-32.1
1984	176	48.9	41.5-57.1	83	57.5	44.6-72.6	93	43.6	34.9-53.6	89	24.5	19.4-30.4	36	26.2	17.4-37.3	53	24.2	18.0-31.7
1985	171	46.1	39.0-53.9	77	51.8	39.5-66.2	94	42.8	34.3-52.5	105	29.7	24.0-36.2	42	30.1	20.9-41.5	63	29.6	22.6-38.0
1986	173	43.6	36.9-50.9	77	45.7	35.0-58.3	96	41.5	33.4-51.0	99	27.2	21.8-33.4	45	30.6	21.4-42.0	54	24.9	18.5-32.6
1987	188	46.6	39.9-54.1	97	60.7	47.8-75.7	91	38.2	30.5-47.0	115	28.9	23.6-35.0	66	45.3	33.9-59.0	49	19.9	14.6-26.4
1988	196	45.7	39.2-52.9	85	47.4	36.7-59.9	111	44.7	36.5-53.9	105	27.0	21.8-32.9	55	36.1	26.0-48.3	50	21.7	16.0-28.7
1989	194	44.9	38.5-52.1	89	53.7	41.7-67.6	105	40.7	33.1-49.4	112	27.8	22.7-33.7	66	46.7	34.9-60.7	46	18.2	13.2-24.3
1990	179	39.7	33.8-46.2	82	44.6	34.4-56.7	97	36.6	29.5-44.8	112	25.8	21.0-31.2	55	30.1	21.9-40.0	57	22.5	16.9-29.3
1991	189	39.7	33.9-46.0	93	45.5	35.7-56.9	96	35.1	28.3-43.0	111	24.9	20.3-30.2	57	32.7	23.7-43.6	54	20.3	15.2-26.6
1992	218	46.0	39.8-52.7	103	53.2	42.3-65.7	115	41.4	34.0-49.7	96	21.0	16.8-25.8	44	23.1	16.2-31.8	52	19.2	14.3-25.2
1993	199	40.7	35.0-47.1	90	44.1	34.4-55.3	109	38.3	31.3-46.2	124	25.7	21.2-30.8	61	30.7	22.8-40.3	63	22.3	17.1-28.6
1994	224	45.2	39.2-51.8	120	60.0	48.5-73.0	104	35.8	29.1-43.4	135	27.1	22.5-32.3	67	33.7	25.2-43.8	68	23.0	17.8-29.2
1995	238	45.6	39.8-52.0	107	50.9	40.6-62.7	131	42.7	35.6-50.7	107	22.2	18.1-27.0	54	30.6	22.1-41.0	53	18.0	13.4-23.6
1996	234	43.1	37.5-49.2	105	46.7	37.3-57.4	129	40.4	33.6-48.1	99	19.5	15.7-23.9	44	20.2	14.3-27.6	55	18.2	13.7-23.7
1997	241	42.9	37.5-48.9	112	50.9	40.9-62.3	129	38.5	32.1-45.9	128	23.6	19.5-28.2	67	30.7	22.9-39.8	61	19.3	14.7-24.7
1998	264	46.0	40.4-52.1	124	53.4	43.6-64.6	140	41.7	35.0-49.3	112	19.7	16.1-23.8	55	23.4	17.1-31.1	57	17.1	12.9-22.2
1999	291	48.7	43.1-54.8	141	58.6	48.4-70.2	150	43.2	36.4-50.7	135	24.7	20.6-29.4	61	29.5	21.8-38.7	74	22.3	17.5-28.0
2000	274	45.0	39.6-50.8	134	54.6	44.9-65.7	140	39.5	33.1-46.7	135	22.7	18.9-27.0	72	32.5	24.6-41.7	63	17.6	13.5-22.6
2001	287	45.1	39.8-50.8	138	54.2	44.9-64.7	149	39.4	33.2-46.3	110	18.2	14.8-22.0	53	22.5	16.4-29.9	57	15.9	12.0-20.6
2002	333	51.5	45.9-57.5	146	57.5	47.7-68.5	187	48.5	41.7-56.1	133	21.6	18.0-25.7	57	24.6	18.0-32.6	76	20.5	16.1-25.6
2003	301	45.9	40.7-51.5	154	59.6	49.7-70.6	147	38.1	32.1-44.9	113	17.5	14.4-21.2	49	19.6	14.1-26.4	64	16.4	12.6-21.0
2004	329	48.0	42.8-53.6	166	58.5	49.0-69.2	163	41.3	35.1-48.1	138	21.2	17.7-25.1	68	25.3	19.1-32.7	70	18.3	14.2-23.2
2005	300	42.7	37.8-48.0	152	53.7	44.8-63.6	148	35.9	30.2-42.2	120	17.6	14.5-21.1	54	19.8	14.5-26.3	66	16.1	12.4-20.5
2006	295	40.0	35.4-45.0	141	43.2	35.7-51.7	154	37.1	31.4-43.5	125	18.0	14.9-21.5	65	22.4	16.9-29.0	60	14.7	11.2-19.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Colon excluding rectum

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	294	37.8	33.5-42.6	148	43.3	36.1-51.4	146	33.3	28.1-39.3	141	19.0	15.9-22.5	70	22.7	17.3-29.1	71	16.4	12.8-20.7
2008	332	41.5	37.0-46.4	157	47.5	39.8-56.1	175	38.0	32.5-44.1	106	14.1	11.5-17.1	52	17.4	12.6-23.1	54	12.1	9.0-15.8
2009	309	36.3	32.2-40.7	159	43.1	36.2-50.9	150	31.2	26.3-36.7	124	15.6	12.8-18.6	62	19.5	14.7-25.4	62	12.9	9.8-16.5
2010	283	33.6	29.6-37.8	150	42.3	35.3-50.1	133	27.3	22.8-32.4	103	12.5	10.1-15.2	52	15.1	11.0-20.2	51	10.5	7.8-13.8
2011	306	34.8	30.8-39.0	151	40.1	33.6-47.5	155	30.8	26.1-36.1	118	14.1	11.6-17.0	61	18.5	13.9-24.1	57	11.4	8.6-14.9
2012	258	28.1	24.6-31.8	133	33.5	27.6-40.2	125	24.3	20.1-29.0	111	13.2	10.8-15.9	50	15.1	10.9-20.2	61	12.4	9.5-16.0
2013	271	28.7	25.3-32.4	132	33.0	27.3-39.5	139	25.7	21.5-30.4	121	12.8	10.6-15.4	63	15.5	11.7-20.2	58	10.7	8.1-13.9
2014	307	31.7	28.2-35.6	152	36.7	30.7-43.4	155	28.2	23.9-33.0	124	13.2	10.9-15.8	62	15.2	11.4-19.7	62	11.9	9.1-15.3
2015	310	30.2	26.9-33.9	159	35.0	29.5-41.3	151	26.2	22.1-30.8	126	13.3	11.0-15.9	59	14.6	10.9-19.1	67	12.1	9.4-15.5
2016	280	26.8	23.7-30.3	142	32.8	27.3-39.0	138	23.0	19.3-27.2	96	9.2	7.4-11.3	51	10.6	7.7-14.1	45	7.9	5.7-10.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Rectum & rectosigmoid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	51	16.7	12.1-22.3	23	16.3	9.9-25.1	28	16.8	10.8-24.6	14	5.9	3.1-9.9	5	4.5	1.4-10.4	9	6.6	2.9-12.5
1977	50	15.9	11.4-21.4	26	19.6	12.1-29.6	24	13.0	8.0-19.8	23	8.2	4.9-12.7	12	9.0	4.2-16.3	11	7.3	3.4-13.2
1978	52	17.0	12.2-22.8	23	18.8	10.7-29.9	29	16.1	10.4-23.4	20	7.1	4.2-11.0	11	8.9	4.1-16.4	9	5.8	2.6-10.7
1979	49	14.9	10.7-20.1	28	19.3	11.9-29.2	21	11.8	7.1-18.2	13	4.4	2.1-7.7	4	3.1	0.6-8.3	9	5.2	2.2-10.0
1980	62	19.8	14.8-25.8	30	23.1	14.7-34.1	32	17.5	11.7-24.9	24	7.2	4.4-10.9	10	5.8	2.7-11.0	14	7.8	4.1-13.1
1981	54	15.5	11.4-20.5	26	18.5	11.4-27.8	28	13.7	8.9-19.9	22	7.4	4.4-11.5	10	6.9	3.0-13.0	12	7.3	3.6-12.8
1982	47	12.9	9.3-17.5	17	10.3	5.4-17.6	30	14.6	9.7-21.0	18	4.9	2.7-7.9	10	4.8	2.3-9.1	8	4.4	1.8-8.7
1983	46	13.1	9.4-17.7	26	18.4	11.3-27.7	20	9.8	5.9-15.1	12	2.9	1.5-5.1	7	3.7	1.4-7.9	5	2.2	0.7-5.1
1984	53	13.9	10.2-18.3	30	19.8	12.5-29.2	23	10.4	6.5-15.5	14	3.7	1.9-6.2	7	4.1	1.4-8.9	7	3.3	1.3-6.8
1985	45	11.2	8.0-15.1	24	14.3	8.7-21.9	21	9.0	5.4-13.8	11	2.9	1.4-5.3	7	4.1	1.5-8.6	4	2.0	0.5-4.9
1986	66	17.1	13.0-22.0	39	22.8	15.6-32.0	27	12.6	8.2-18.4	20	5.3	3.1-8.3	11	8.5	3.6-15.9	9	3.8	1.7-7.2
1987	59	14.0	10.4-18.2	29	16.7	10.5-25.0	30	12.4	8.2-17.7	10	2.7	1.2-5.0	5	4.3	1.2-10.1	5	2.0	0.6-4.6
1988	48	11.6	8.4-15.6	24	14.8	8.9-22.6	24	9.6	6.1-14.4	20	5.5	3.3-8.5	9	7.1	3.0-13.6	11	4.8	2.3-8.4
1989	67	14.9	11.4-19.0	35	18.8	12.3-27.0	32	12.4	8.4-17.5	15	4.3	2.3-7.0	9	7.1	3.0-13.4	6	2.7	1.0-5.7
1990	62	14.3	10.8-18.5	29	15.4	9.9-22.5	33	12.9	8.8-18.2	20	4.9	2.9-7.6	9	6.5	2.6-12.8	11	4.3	2.1-7.6
1991	54	11.0	8.1-14.4	26	11.1	7.0-16.6	28	10.4	6.8-15.0	15	3.4	1.8-5.6	7	4.1	1.4-8.7	8	3.0	1.3-5.8
1992	61	12.0	9.0-15.5	33	16.5	10.7-24.1	28	9.4	6.2-13.6	20	4.8	2.9-7.5	9	5.7	2.2-11.1	11	4.4	2.1-7.7
1993	66	12.4	9.4-15.8	34	12.9	8.8-18.3	32	11.1	7.5-15.6	15	3.6	2.0-6.0	9	5.4	2.1-10.7	6	2.5	0.9-5.3
1994	77	14.2	11.1-17.8	41	19.3	13.3-26.8	36	11.1	7.8-15.4	19	3.5	2.1-5.5	11	4.5	2.1-8.3	8	2.6	1.1-5.1
1995	76	13.4	10.5-16.9	39	16.6	11.4-23.2	37	11.3	7.9-15.6	20	3.5	2.1-5.4	9	3.5	1.5-6.8	11	3.4	1.7-6.0
1996	94	16.7	13.4-20.5	62	27.1	20.2-35.4	32	9.8	6.7-13.8	27	5.5	3.6-8.1	16	7.5	4.0-12.6	11	4.0	2.0-7.0
1997	78	13.6	10.7-17.1	45	19.1	13.5-26.0	33	9.9	6.8-13.9	20	3.6	2.1-5.6	11	4.0	1.9-7.4	9	3.0	1.3-5.5
1998	74	11.7	9.1-14.8	34	12.5	8.3-17.9	40	11.3	8.0-15.4	17	3.0	1.7-4.8	10	4.5	2.0-8.3	7	2.0	0.8-4.1
1999	100	16.4	13.2-20.0	52	19.0	13.8-25.3	48	14.1	10.4-18.7	18	3.6	2.1-5.6	12	6.0	2.9-10.7	6	2.0	0.7-4.1
2000	79	12.2	9.6-15.3	39	13.4	9.2-18.7	40	11.0	7.8-15.0	25	3.7	2.4-5.5	12	3.6	1.8-6.3	13	3.5	1.9-6.0
2001	84	12.3	9.7-15.2	54	18.3	13.4-24.5	30	7.9	5.3-11.3	16	2.2	1.3-3.6	9	3.2	1.3-6.2	7	1.7	0.7-3.4
2002	80	10.8	8.5-13.5	41	11.5	8.0-15.9	39	9.8	6.9-13.4	20	3.1	1.8-4.7	12	4.4	2.1-8.1	8	2.1	0.9-4.2
2003	96	13.7	11.0-16.8	51	17.0	12.2-22.9	45	11.4	8.3-15.2	25	3.9	2.4-5.7	18	6.7	3.7-11.1	7	2.0	0.8-4.0
2004	94	12.5	10.1-15.4	52	16.3	11.9-21.7	42	9.9	7.1-13.4	17	2.6	1.5-4.2	11	4.5	2.0-8.2	6	1.5	0.6-3.3
2005	97	12.3	9.9-15.0	47	13.6	9.8-18.4	50	11.1	8.2-14.7	21	3.0	1.8-4.7	13	4.1	2.1-7.1	8	2.1	0.9-4.0
2006	101	12.9	10.4-15.7	53	15.4	11.2-20.5	48	10.9	8.0-14.5	21	2.9	1.8-4.5	9	3.3	1.3-6.5	12	2.9	1.5-5.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Rectum & rectosigmoid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	91	11.0	8.8-13.6	45	12.3	8.7-16.8	46	10.0	7.3-13.4	36	4.8	3.3-6.7	19	5.7	3.2-9.1	17	4.1	2.3-6.5
2008	104	12.2	9.9-14.9	63	17.2	12.9-22.4	41	8.7	6.2-11.9	16	1.9	1.1-3.2	9	2.5	1.0-4.8	7	1.5	0.6-3.0
2009	105	12.1	9.8-14.7	53	14.1	10.4-18.7	52	10.6	7.8-13.9	21	2.4	1.5-3.7	14	3.6	1.9-6.2	7	1.5	0.6-3.1
2010	105	11.7	9.5-14.3	58	15.5	11.5-20.4	47	9.2	6.7-12.3	23	2.6	1.6-3.9	12	3.1	1.4-5.7	11	2.2	1.1-4.0
2011	104	11.3	9.2-13.8	64	15.4	11.6-19.9	40	8.0	5.7-10.9	19	2.1	1.3-3.3	10	2.3	1.1-4.2	9	1.8	0.8-3.5
2012	105	10.6	8.6-13.0	56	12.6	9.3-16.6	49	9.0	6.7-12.0	13	1.6	0.9-2.8	9	3.4	1.5-6.2	4	0.7	0.2-1.9
2013	117	11.9	9.8-14.4	60	13.7	10.2-18.0	57	10.7	8.0-13.9	16	1.8	1.0-2.9	8	2.4	1.0-4.7	8	1.4	0.6-2.8
2014	120	11.7	9.6-14.0	64	14.3	10.8-18.5	56	9.6	7.2-12.5	24	2.5	1.6-3.7	17	4.3	2.4-6.9	7	1.2	0.5-2.5
2015	123	11.6	9.6-13.8	66	14.3	10.9-18.4	57	9.6	7.2-12.5	30	3.1	2.1-4.4	13	3.1	1.5-5.4	17	3.1	1.8-4.9
2016	103	9.9	8.0-12.0	64	14.1	10.7-18.2	39	6.7	4.7-9.2	14	1.5	0.8-2.5	8	2.0	0.8-4.0	6	1.0	0.4-2.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Liver / intrahepatic bile duct

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	16	4.1	2.3-6.9	14	8.2	4.2-14.6	2	0.9	0.1-3.4	18	5.0	2.7-8.4	15	8.3	4.4-14.5	3	2.2	0.3-6.3
1977	13	3.9	1.9-7.0	9	4.9	2.2-9.7	4	2.8	0.6-7.0	12	2.8	1.4-5.0	12	6.2	3.1-11.3	0	0.0	0.0-1.9
1978	20	5.6	3.2-9.0	14	9.0	4.3-16.1	6	3.2	1.1-6.9	11	3.3	1.5-6.0	5	3.9	1.0-9.6	6	2.9	1.0-6.4
1979	13	3.7	1.8-6.5	8	4.5	1.8-9.4	5	2.8	0.8-6.6	10	3.0	1.2-5.8	7	3.4	1.3-7.5	3	2.3	0.4-6.3
1980	21	5.9	3.5-9.2	14	8.0	4.0-14.2	7	4.1	1.5-8.3	16	4.7	2.5-7.8	12	7.4	3.5-13.5	4	2.6	0.7-6.4
1981	24	6.2	3.9-9.4	14	9.0	4.5-15.6	10	4.4	2.1-8.2	25	7.4	4.6-11.1	14	9.2	4.6-16.1	11	5.9	2.8-10.6
1982	23	6.3	3.8-9.6	20	13.3	7.5-21.5	3	1.5	0.3-4.4	20	5.7	3.4-8.9	15	10.2	5.2-17.5	5	2.7	0.8-6.1
1983	19	4.7	2.7-7.5	17	10.1	5.5-16.8	2	0.9	0.1-3.2	24	6.1	3.8-9.2	19	11.8	6.6-19.2	5	2.3	0.7-5.2
1984	15	3.3	1.8-5.4	11	5.4	2.6-9.9	4	1.6	0.4-4.1	19	4.8	2.8-7.7	14	8.7	4.4-15.1	5	2.2	0.7-5.1
1985	15	3.3	1.8-5.5	5	2.2	0.7-5.5	10	4.0	1.9-7.3	9	2.4	1.1-4.6	3	1.9	0.3-5.5	6	2.8	1.0-5.9
1986	18	3.6	2.1-5.8	11	5.5	2.5-10.2	7	2.4	0.9-5.0	15	2.8	1.5-4.6	9	3.8	1.7-7.5	6	1.9	0.7-4.3
1987	17	3.5	2.0-5.7	9	3.7	1.6-7.3	8	3.2	1.3-6.3	20	4.5	2.7-7.1	12	5.7	2.7-10.4	8	3.6	1.5-6.9
1988	24	5.0	3.1-7.5	16	7.9	4.1-13.5	8	3.0	1.2-5.9	25	5.4	3.4-8.0	21	12.0	6.9-19.0	4	1.2	0.3-3.2
1989	28	6.1	4.0-8.9	23	11.4	6.8-17.8	5	2.3	0.7-5.1	21	4.9	2.9-7.5	12	7.1	3.3-12.9	9	3.6	1.6-6.7
1990	31	6.4	4.3-9.2	21	9.9	5.9-15.6	10	3.7	1.7-6.9	25	4.6	3.0-6.9	19	8.3	4.8-13.3	6	2.0	0.7-4.3
1991	33	5.9	3.9-8.4	24	9.5	5.9-14.5	9	2.9	1.3-5.7	27	4.5	2.9-6.6	19	6.9	4.1-11.0	8	2.4	1.0-4.8
1992	23	4.4	2.7-6.6	15	6.1	3.2-10.3	8	2.8	1.2-5.5	29	5.2	3.4-7.5	18	7.1	4.1-11.5	11	3.5	1.7-6.3
1993	26	5.4	3.4-8.0	11	4.4	2.1-8.1	15	5.4	3.0-8.9	22	4.4	2.7-6.7	8	2.7	1.1-5.6	14	5.0	2.7-8.3
1994	25	4.5	2.9-6.8	19	8.7	4.7-14.2	6	2.0	0.7-4.2	28	5.4	3.5-7.8	19	9.5	5.4-15.2	9	2.8	1.2-5.3
1995	41	7.5	5.3-10.2	30	13.0	8.4-19.2	11	3.6	1.8-6.4	30	5.5	3.7-7.9	23	10.8	6.5-16.5	7	2.1	0.9-4.4
1996	40	7.2	5.1-9.9	25	11.0	6.6-17.0	15	4.9	2.7-8.1	39	7.1	5.0-9.7	25	10.8	6.5-16.5	14	4.6	2.5-7.7
1997	51	8.8	6.5-11.6	33	14.2	9.4-20.3	18	5.3	3.1-8.4	33	5.7	3.9-8.1	20	8.5	4.9-13.5	13	3.9	2.1-6.7
1998	54	8.4	6.2-11.0	37	11.8	8.2-16.4	17	5.2	3.0-8.2	47	8.1	5.8-10.8	30	11.4	7.4-16.7	17	5.3	3.1-8.5
1999	30	4.6	3.0-6.5	21	6.6	4.0-10.3	9	2.6	1.2-4.9	25	4.4	2.8-6.5	14	5.2	2.7-8.9	11	3.4	1.7-6.0
2000	35	5.2	3.5-7.2	25	8.6	5.4-13.0	10	2.6	1.2-4.8	39	6.0	4.2-8.2	26	9.0	5.6-13.7	13	3.8	2.0-6.5
2001	44	6.0	4.3-8.1	31	9.1	6.1-13.1	13	3.3	1.7-5.6	37	5.3	3.7-7.3	21	6.7	3.9-10.6	16	4.2	2.4-6.8
2002	62	8.4	6.4-10.8	50	14.6	10.6-19.5	12	3.3	1.7-5.7	33	4.7	3.2-6.6	22	7.2	4.3-11.2	11	2.8	1.4-5.0
2003	62	8.3	6.3-10.6	42	11.1	7.9-15.2	20	5.4	3.3-8.3	42	5.3	3.8-7.2	35	9.4	6.4-13.2	7	1.8	0.7-3.7
2004	69	8.8	6.8-11.2	49	14.5	10.4-19.4	20	4.6	2.7-7.1	48	6.5	4.7-8.6	30	9.1	6.0-13.3	18	4.4	2.6-7.0
2005	83	10.9	8.6-13.6	62	18.4	13.8-24.0	21	5.2	3.2-7.9	53	6.4	4.7-8.4	40	10.6	7.4-14.7	13	3.0	1.6-5.1
2006	66	8.1	6.2-10.3	52	13.9	10.2-18.4	14	3.3	1.8-5.5	38	5.2	3.6-7.2	22	6.4	3.9-9.7	16	4.0	2.3-6.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Liver / intrahepatic bile duct

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	99	11.5	9.2-14.0	72	18.2	14.0-23.3	27	5.9	3.9-8.6	53	6.9	5.1-9.0	35	10.0	6.8-14.0	18	4.2	2.5-6.7
2008	98	10.4	8.4-12.8	74	17.8	13.8-22.7	24	4.7	3.0-7.0	59	6.8	5.1-8.8	46	12.0	8.6-16.4	13	2.8	1.5-4.7
2009	112	11.7	9.6-14.1	91	20.9	16.7-25.9	21	4.1	2.5-6.3	73	8.0	6.2-10.2	64	15.9	12.0-20.6	9	1.9	0.9-3.6
2010	103	10.4	8.5-12.7	84	18.6	14.7-23.2	19	3.7	2.2-5.8	79	8.2	6.4-10.2	59	13.8	10.3-18.0	20	3.7	2.3-5.8
2011	98	9.3	7.5-11.4	81	16.3	12.8-20.5	17	3.3	1.9-5.4	77	7.7	6.0-9.7	60	13.0	9.8-16.9	17	3.2	1.9-5.2
2012	102	9.6	7.8-11.7	78	15.9	12.4-20.1	24	4.5	2.8-6.7	89	9.1	7.3-11.3	65	14.0	10.6-18.2	24	5.0	3.2-7.5
2013	107	10.2	8.3-12.3	70	14.5	11.2-18.5	37	6.6	4.6-9.2	75	7.5	5.8-9.4	61	14.1	10.6-18.4	14	2.6	1.4-4.4
2014	145	13.3	11.2-15.7	117	24.2	19.8-29.3	28	4.7	3.1-6.9	72	6.9	5.4-8.8	50	11.3	8.2-15.1	22	3.8	2.4-5.8
2015	113	10.2	8.3-12.3	77	15.5	12.1-19.5	36	5.9	4.1-8.2	86	8.4	6.7-10.4	61	13.7	10.3-17.7	25	4.5	2.9-6.7
2016	151	12.9	10.9-15.2	116	22.0	18.0-26.6	35	5.7	3.9-7.9	74	7.1	5.5-8.9	57	12.9	9.6-16.9	17	2.9	1.7-4.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Pancreas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	54	17.5	12.7-23.3	21	13.9	8.2-22.0	33	19.7	13.2-28.2	38	12.0	8.2-16.8	19	12.5	7.4-19.9	19	11.3	6.5-17.9
1977	55	16.3	12.0-21.6	33	19.9	13.3-28.6	22	12.8	7.7-19.6	57	17.1	12.7-22.6	34	21.4	14.3-30.7	23	13.4	8.2-20.3
1978	70	21.6	16.4-27.7	40	27.5	18.5-39.0	30	17.2	11.2-24.9	50	15.4	11.1-20.6	27	19.1	11.7-29.0	23	12.6	7.8-19.2
1979	70	23.3	17.7-30.0	41	33.1	22.2-46.8	29	16.9	11.0-24.6	57	17.8	13.1-23.6	31	20.9	13.1-31.1	26	15.1	9.6-22.4
1980	50	14.5	10.5-19.4	27	20.1	12.4-30.2	23	11.1	6.9-16.8	47	13.8	9.8-18.6	27	18.6	11.4-28.4	20	10.5	6.2-16.4
1981	59	17.8	13.3-23.3	28	21.2	13.3-31.5	31	15.9	10.6-22.8	54	16.0	11.7-21.2	26	18.1	10.8-27.9	28	14.7	9.6-21.4
1982	64	17.1	12.9-22.0	33	20.1	13.3-29.0	31	14.8	9.9-21.1	66	18.4	14.0-23.7	33	23.2	15.1-33.5	33	15.6	10.6-22.1
1983	74	19.7	15.2-25.0	42	26.6	18.3-36.9	32	15.0	10.1-21.3	65	17.7	13.4-22.7	36	23.7	15.9-33.8	29	13.7	9.0-19.7
1984	73	20.2	15.6-25.6	30	23.4	14.9-34.4	43	19.2	13.7-25.9	68	17.5	13.3-22.4	36	23.9	15.9-34.0	32	13.7	9.2-19.4
1985	67	17.9	13.6-22.9	33	21.1	13.8-30.5	34	15.6	10.7-21.8	57	15.6	11.6-20.4	28	20.8	13.0-31.0	29	13.0	8.6-18.7
1986	69	16.2	12.4-20.6	32	16.9	11.1-24.5	37	15.3	10.7-21.1	66	16.2	12.4-20.8	34	18.2	12.1-26.2	32	14.3	9.7-20.2
1987	67	16.2	12.4-20.7	21	14.2	8.0-22.7	46	18.4	13.4-24.6	67	16.5	12.6-21.1	26	16.4	9.9-25.1	41	16.9	12.0-23.0
1988	65	17.5	13.3-22.4	34	22.2	14.8-31.6	31	14.2	9.5-20.1	67	17.5	13.4-22.3	38	24.4	16.6-34.1	29	12.9	8.5-18.5
1989	60	14.2	10.7-18.4	25	15.4	9.4-23.5	35	13.7	9.4-19.1	54	13.6	10.1-17.8	24	15.2	9.2-23.3	30	12.6	8.4-18.0
1990	92	22.2	17.7-27.5	35	22.9	15.2-32.6	57	22.5	16.9-29.2	72	17.2	13.2-21.7	20	13.7	7.7-21.9	52	20.1	14.9-26.4
1991	76	16.5	12.8-20.8	42	21.9	15.0-30.6	34	12.7	8.7-17.8	56	13.2	9.8-17.2	29	16.1	10.2-23.8	27	10.9	7.1-15.8
1992	70	15.5	11.9-19.7	33	18.6	12.0-27.0	37	13.8	9.6-19.0	62	14.6	11.0-18.8	28	17.7	11.0-26.5	34	13.2	9.1-18.4
1993	51	11.1	8.2-14.7	22	11.4	6.7-17.7	29	10.7	7.1-15.3	56	11.8	8.8-15.4	26	13.9	8.6-21.0	30	10.6	7.1-15.1
1994	79	16.6	13.0-20.8	36	18.7	12.7-26.4	43	15.2	10.9-20.4	66	14.6	11.2-18.6	33	18.5	12.2-26.6	33	12.2	8.3-17.0
1995	76	14.6	11.4-18.4	35	15.1	10.3-21.4	41	13.5	9.6-18.3	67	13.2	10.1-16.9	33	16.6	10.9-23.8	34	11.2	7.7-15.6
1996	87	15.6	12.4-19.4	43	19.7	13.8-27.1	44	13.4	9.7-17.9	75	14.0	10.9-17.7	33	17.7	11.7-25.3	42	12.7	9.1-17.1
1997	79	14.7	11.5-18.4	32	14.7	9.7-21.1	47	14.5	10.6-19.3	63	11.9	9.0-15.3	31	14.3	9.4-20.6	32	10.2	6.9-14.4
1998	82	14.5	11.4-18.1	30	13.4	8.6-19.8	52	15.6	11.6-20.5	86	16.2	12.8-20.1	33	15.6	10.3-22.4	53	16.6	12.4-21.7
1999	87	14.6	11.6-18.1	39	14.1	9.8-19.6	48	14.2	10.4-18.8	90	15.0	12.0-18.5	37	13.6	9.3-19.2	53	15.5	11.5-20.2
2000	83	14.3	11.3-17.8	32	12.0	7.9-17.4	51	15.0	11.1-19.7	72	12.4	9.6-15.7	31	12.7	8.3-18.3	41	11.9	8.5-16.2
2001	116	18.9	15.5-22.8	49	18.3	13.1-24.7	67	18.9	14.6-24.0	84	14.3	11.3-17.8	38	15.2	10.4-21.2	46	13.3	9.7-17.7
2002	99	15.3	12.4-18.7	50	18.3	13.3-24.4	49	13.0	9.6-17.2	81	13.0	10.3-16.3	42	15.3	10.7-21.1	39	11.0	7.8-15.0
2003	93	14.2	11.4-17.5	49	18.7	13.4-25.2	44	11.6	8.3-15.5	87	13.8	11.0-17.1	40	16.4	11.3-22.7	47	12.4	9.1-16.5
2004	99	15.0	12.1-18.3	42	14.7	10.2-20.4	57	14.8	11.2-19.2	84	13.0	10.3-16.1	42	15.9	11.0-22.0	42	11.0	7.9-14.9
2005	126	18.1	15.0-21.7	59	19.2	14.4-25.1	67	16.6	12.8-21.1	84	12.3	9.7-15.2	44	15.0	10.5-20.6	40	10.0	7.1-13.6
2006	114	15.4	12.7-18.6	53	17.5	12.9-23.2	61	14.1	10.7-18.1	83	11.1	8.8-13.9	32	10.5	7.0-15.0	51	11.7	8.7-15.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Pancreas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	106	14.5	11.8-17.6	42	13.4	9.4-18.4	64	14.9	11.4-19.1	76	10.3	8.1-13.0	34	10.7	7.3-15.2	42	9.7	7.0-13.2
2008	124	16.1	13.3-19.3	58	17.1	12.8-22.4	66	15.0	11.5-19.1	90	12.0	9.6-14.9	35	11.7	7.9-16.5	55	12.6	9.4-16.4
2009	124	15.4	12.7-18.5	69	18.7	14.4-24.0	55	12.4	9.3-16.1	112	14.2	11.6-17.2	50	14.2	10.3-19.0	62	13.7	10.5-17.6
2010	114	13.8	11.3-16.6	57	15.9	11.8-21.0	57	12.2	9.2-15.8	89	10.9	8.7-13.5	43	11.9	8.4-16.2	46	9.9	7.2-13.3
2011	147	16.9	14.2-20.0	69	17.5	13.3-22.5	78	16.0	12.6-20.0	99	12.2	9.8-14.9	43	12.4	8.7-17.0	56	11.8	8.9-15.3
2012	142	15.4	12.9-18.2	66	16.7	12.7-21.5	76	14.4	11.3-18.1	115	12.4	10.2-15.0	55	13.1	9.7-17.3	60	11.9	9.0-15.3
2013	128	13.8	11.4-16.5	63	15.5	11.7-20.2	65	12.4	9.5-15.8	97	11.0	8.9-13.5	46	12.7	9.1-17.2	51	9.7	7.2-12.8
2014	164	17.3	14.7-20.2	77	18.4	14.3-23.3	87	16.2	12.9-20.0	103	11.3	9.2-13.8	48	12.2	8.9-16.4	55	10.4	7.8-13.6
2015	166	16.8	14.2-19.6	68	15.3	11.7-19.7	98	17.6	14.2-21.5	112	11.8	9.6-14.2	50	11.5	8.3-15.3	62	11.7	8.9-15.0
2016	162	16.2	13.7-18.9	75	16.9	13.1-21.4	87	15.4	12.2-19.0	114	11.6	9.5-13.9	57	13.3	9.9-17.5	57	10.1	7.6-13.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Larynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	36	9.3	6.5-13.0	32	18.6	12.6-26.7	4	1.9	0.5-4.9	16	5.0	2.7-8.4	13	10.1	4.9-18.1	3	1.4	0.3-4.2
1977	22	6.0	3.6-9.4	20	12.8	7.0-21.1	2	1.1	0.1-3.8	11	2.6	1.3-4.8	9	4.7	2.1-9.5	2	0.9	0.1-3.4
1978	25	6.2	4.0-9.3	20	11.4	6.8-18.2	5	2.2	0.7-5.1	19	5.1	2.9-8.1	16	10.4	5.2-18.3	3	1.3	0.3-3.8
1979	41	10.0	7.1-13.7	32	17.9	11.9-25.9	9	3.9	1.8-7.5	10	2.6	1.2-4.9	10	6.1	2.8-11.6	0	0.0	0.0-1.7
1980	38	9.1	6.4-12.5	34	18.8	12.7-26.7	4	1.6	0.4-4.2	14	3.3	1.8-5.6	11	6.0	3.0-11.1	3	1.2	0.2-3.6
1981	39	8.9	6.3-12.3	32	16.0	10.8-22.9	7	3.1	1.2-6.5	19	4.7	2.7-7.6	16	10.0	5.2-17.1	3	1.2	0.2-3.5
1982	36	8.0	5.6-11.2	34	17.3	11.7-24.7	2	0.8	0.1-2.9	19	4.7	2.7-7.4	16	9.2	4.7-15.7	3	1.4	0.3-4.0
1983	24	5.3	3.4-8.0	22	10.5	6.5-16.1	2	1.1	0.1-3.5	18	4.0	2.4-6.4	15	7.7	4.3-13.0	3	1.2	0.2-3.4
1984	36	8.0	5.6-11.2	27	14.1	8.9-21.2	9	3.5	1.6-6.7	11	2.9	1.4-5.1	10	6.3	2.7-12.2	1	0.6	0.0-2.7
1985	42	8.8	6.3-12.0	30	14.0	9.3-20.2	12	4.7	2.4-8.2	15	3.7	2.0-6.1	13	8.0	4.0-14.0	2	0.8	0.1-2.8
1986	39	8.3	5.8-11.4	32	16.2	10.8-23.4	7	2.5	1.0-5.1	16	3.5	2.0-5.7	11	5.8	2.8-10.6	5	1.9	0.6-4.4
1987	29	5.8	3.9-8.4	24	11.1	7.0-16.7	5	1.8	0.6-4.1	12	2.6	1.3-4.6	7	3.8	1.4-8.0	5	1.9	0.6-4.4
1988	24	5.0	3.1-7.5	17	8.1	4.5-13.3	7	2.7	1.1-5.5	17	3.5	2.0-5.6	17	8.2	4.6-13.3	0	0.0	0.0-1.2
1989	38	7.2	5.0-9.9	25	11.5	7.1-17.6	13	4.2	2.2-7.2	11	2.2	1.1-4.0	10	4.8	2.2-8.9	1	0.4	0.0-1.9
1990	33	6.1	4.2-8.7	28	12.0	7.7-17.8	5	1.7	0.6-4.0	16	3.2	1.8-5.2	14	6.8	3.3-12.0	2	0.7	0.1-2.6
1991	28	5.3	3.5-7.8	24	10.6	6.5-16.1	4	1.4	0.4-3.6	17	3.3	1.9-5.3	17	8.0	4.4-13.2	0	0.0	0.0-1.1
1992	39	7.1	5.0-9.8	28	12.1	7.9-17.8	11	3.5	1.7-6.3	17	3.1	1.8-4.9	12	5.3	2.6-9.6	5	1.5	0.5-3.5
1993	38	6.7	4.7-9.2	34	14.2	9.5-20.1	4	1.2	0.3-3.1	16	2.8	1.6-4.5	14	5.3	2.9-9.0	2	0.7	0.1-2.4
1994	40	6.8	4.8-9.3	37	15.1	10.3-21.2	3	0.9	0.2-2.5	18	3.4	2.0-5.5	14	6.8	3.3-11.8	4	1.3	0.4-3.3
1995	37	6.3	4.4-8.7	30	12.4	8.1-18.0	7	2.1	0.8-4.2	18	3.4	2.0-5.4	15	7.4	4.0-12.4	3	0.9	0.2-2.5
1996	41	6.8	4.8-9.3	35	14.3	9.5-20.4	6	1.7	0.6-3.7	17	2.9	1.7-4.7	15	6.8	3.4-11.7	2	0.6	0.1-2.1
1997	42	6.6	4.7-9.0	33	12.7	8.5-18.2	9	2.4	1.1-4.5	12	1.8	0.9-3.2	9	3.2	1.4-6.2	3	0.8	0.2-2.2
1998	49	7.4	5.5-9.9	35	12.7	8.5-18.0	14	3.8	2.0-6.3	15	2.4	1.3-4.0	14	5.3	2.7-9.1	1	0.3	0.0-1.6
1999	37	5.3	3.7-7.3	29	9.7	6.3-14.3	8	2.0	0.9-3.9	20	3.1	1.9-4.8	19	7.2	4.1-11.5	1	0.2	0.0-1.3
2000	33	4.6	3.1-6.4	23	7.4	4.6-11.3	10	2.4	1.1-4.4	13	2.1	1.1-3.6	7	3.3	1.2-6.9	6	1.5	0.6-3.3
2001	37	5.0	3.5-6.9	31	9.4	6.3-13.5	6	1.5	0.5-3.2	10	1.5	0.7-2.7	9	3.5	1.4-6.9	1	0.2	0.0-1.3
2002	37	4.8	3.4-6.7	28	8.3	5.5-12.2	9	2.0	0.9-3.9	24	3.3	2.1-5.0	18	5.4	3.2-8.6	6	1.6	0.6-3.4
2003	42	5.7	4.1-7.8	32	10.3	6.8-14.8	10	2.5	1.2-4.6	20	2.7	1.6-4.2	14	4.5	2.4-7.7	6	1.4	0.5-3.1
2004	40	5.0	3.6-6.9	36	10.5	7.2-14.7	4	0.9	0.2-2.3	14	1.8	1.0-3.0	12	3.5	1.8-6.2	2	0.5	0.1-1.7
2005	32	4.3	2.9-6.1	27	9.0	5.7-13.4	5	1.1	0.4-2.6	14	1.7	0.9-2.9	13	3.9	1.9-6.9	1	0.2	0.0-1.1
2006	48	5.9	4.3-7.8	39	11.2	7.8-15.5	9	1.9	0.9-3.7	11	1.6	0.8-2.8	10	3.4	1.6-6.3	1	0.3	0.0-1.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Larynx

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	46	5.8	4.2-7.7	31	9.5	6.2-13.8	15	3.2	1.8-5.3	11	1.4	0.7-2.6	7	2.1	0.8-4.5	4	0.9	0.2-2.3
2008	47	5.6	4.1-7.5	37	10.6	7.2-14.9	10	2.2	1.0-4.0	20	2.4	1.5-3.8	15	4.0	2.2-6.8	5	1.1	0.4-2.5
2009	34	4.0	2.8-5.7	24	6.7	4.2-10.2	10	2.1	1.0-3.9	18	2.2	1.3-3.4	13	3.9	1.9-6.9	5	1.0	0.3-2.4
2010	43	5.0	3.6-6.8	33	9.2	6.2-13.1	10	2.0	0.9-3.7	16	2.0	1.2-3.3	14	4.3	2.3-7.3	2	0.4	0.0-1.5
2011	42	4.3	3.1-5.9	36	8.5	5.8-11.9	6	1.1	0.4-2.4	11	1.1	0.5-2.0	9	2.0	0.9-3.9	2	0.4	0.0-1.5
2012	31	3.3	2.2-4.6	21	5.5	3.3-8.5	10	1.8	0.9-3.3	21	2.3	1.4-3.6	19	5.4	3.1-8.5	2	0.4	0.0-1.3
2013	33	3.3	2.2-4.7	23	5.7	3.5-8.7	10	1.6	0.8-3.0	15	1.7	0.9-2.8	14	4.1	2.2-7.0	1	0.2	0.0-1.0
2014	45	4.6	3.3-6.1	38	9.2	6.3-12.8	7	1.2	0.5-2.6	17	1.8	1.0-2.9	15	4.0	2.1-6.7	2	0.3	0.0-1.2
2015	45	4.3	3.1-5.8	39	8.8	6.1-12.2	6	1.0	0.4-2.3	13	1.4	0.7-2.4	12	3.3	1.6-5.9	1	0.2	0.0-1.0
2016	44	4.0	2.9-5.5	34	7.4	5.0-10.4	10	1.5	0.7-2.8	14	1.3	0.7-2.3	10	2.4	1.1-4.4	4	0.6	0.2-1.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Lung and bronchus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	224	62.9	54.5-72.0	174	111.8	94.7-130.9	50	24.6	18.1-32.7	156	44.2	37.2-52.1	122	80.3	65.3-97.6	34	16.6	11.4-23.3
1977	284	81.7	71.8-92.5	219	148.9	127.6-172.6	65	32.0	24.3-41.3	205	59.4	51.0-68.6	165	112.9	94.5-133.5	40	19.8	13.9-27.3
1978	270	74.4	65.3-84.4	203	128.7	109.9-149.5	67	33.7	25.7-43.2	216	60.8	52.4-70.1	169	111.8	93.6-132.1	47	23.6	17.0-31.7
1979	278	76.1	66.8-86.3	200	134.3	113.9-156.9	78	35.5	27.9-44.6	217	61.3	52.8-70.6	166	112.4	94.0-133.0	51	25.1	18.3-33.4
1980	333	87.0	77.4-97.4	253	159.8	138.6-183.0	80	35.4	27.9-44.3	262	71.2	62.3-81.0	193	129.2	109.3-151.4	69	31.6	24.4-40.2
1981	332	85.4	76.0-95.6	250	154.6	134.2-177.0	82	36.3	28.6-45.3	258	67.9	59.4-77.3	190	123.5	104.4-144.7	68	29.9	23.1-38.1
1982	325	79.9	71.0-89.5	243	144.5	125.1-166.0	82	34.5	27.2-43.0	265	65.4	57.4-74.2	197	115.5	98.5-134.4	68	29.5	22.8-37.6
1983	337	80.6	71.8-90.0	241	138.2	120.0-158.2	96	39.6	31.8-48.5	293	73.7	65.0-83.0	225	137.7	118.7-158.6	68	29.2	22.4-37.2
1984	325	76.4	67.9-85.6	228	124.2	107.4-142.8	97	41.2	33.2-50.6	273	63.9	56.3-72.3	197	110.4	94.2-128.4	76	31.1	24.4-39.1
1985	353	79.5	71.1-88.5	250	133.2	116.0-152.1	103	40.7	33.1-49.5	274	63.0	55.5-71.3	198	107.2	91.7-124.5	76	31.1	24.3-39.1
1986	370	84.7	75.9-94.2	262	149.1	129.6-170.5	108	42.2	34.5-51.1	293	67.0	59.3-75.5	199	112.5	95.9-130.8	94	36.7	29.6-45.1
1987	337	72.9	65.0-81.3	234	119.9	104.0-137.4	103	39.1	31.7-47.5	325	71.7	63.8-80.3	234	126.7	109.5-145.7	91	34.6	27.7-42.6
1988	394	85.9	77.3-95.1	273	148.4	129.7-168.9	121	44.9	37.2-53.8	292	63.7	56.4-71.7	204	110.0	94.2-127.4	88	32.8	26.2-40.5
1989	410	84.9	76.6-93.8	286	145.1	127.2-164.5	124	44.4	36.8-53.1	378	79.6	71.4-88.3	258	134.4	117.2-153.3	120	43.1	35.6-51.6
1990	385	80.8	72.6-89.6	274	141.1	123.3-160.5	111	40.3	33.0-48.6	331	69.6	62.0-77.8	224	115.3	99.3-133.0	107	39.2	32.0-47.4
1991	411	83.5	75.3-92.3	264	134.5	117.1-153.6	147	50.7	42.7-59.7	346	71.2	63.6-79.4	221	114.1	98.2-131.6	125	43.3	35.9-51.7
1992	420	81.8	73.8-90.3	295	145.1	127.2-164.6	125	41.1	34.1-49.1	326	65.6	58.4-73.5	227	116.6	100.3-134.6	99	33.4	27.0-40.8
1993	404	77.3	69.7-85.5	266	128.8	112.1-147.0	138	44.9	37.6-53.2	362	70.6	63.2-78.5	252	124.5	108.0-142.6	110	36.3	29.8-43.9
1994	436	79.3	71.7-87.3	252	117.4	102.0-134.4	184	56.3	48.4-65.2	369	67.9	60.9-75.4	260	118.9	103.5-135.7	109	34.4	28.2-41.5
1995	438	79.1	71.7-87.1	278	120.7	105.7-136.9	160	50.6	42.9-59.1	352	64.4	57.7-71.7	212	97.8	84.0-113.0	140	43.3	36.3-51.2
1996	455	79.6	72.2-87.5	277	118.5	103.8-134.6	178	53.6	45.9-62.2	359	63.7	57.1-70.9	215	91.1	78.3-105.2	144	44.7	37.6-52.7
1997	464	78.8	71.6-86.5	293	120.8	106.3-136.6	171	50.1	42.8-58.3	371	64.1	57.5-71.1	236	102.0	88.4-116.9	135	39.5	33.1-46.8
1998	467	76.9	69.8-84.4	273	108.1	94.6-122.9	194	55.6	47.9-64.0	375	63.6	57.1-70.6	239	99.5	86.4-114.0	136	39.4	33.0-46.7
1999	460	75.3	68.3-82.7	265	105.1	91.8-119.6	195	55.4	47.8-63.8	336	56.1	50.1-62.6	204	82.8	70.8-96.1	132	38.6	32.2-45.8
2000	483	74.9	68.2-82.1	307	113.5	100.2-127.9	176	48.1	41.2-55.9	374	58.9	52.9-65.3	231	89.7	77.6-102.9	143	38.6	32.5-45.5
2001	492	74.7	68.0-81.8	300	111.7	98.6-126.1	192	50.1	43.2-57.8	353	55.7	49.8-61.9	211	82.1	70.5-94.8	142	38.4	32.2-45.3
2002	465	69.9	63.5-76.7	273	101.3	88.6-115.0	192	49.5	42.7-57.1	371	55.8	50.1-62.0	233	86.1	74.6-98.9	138	35.8	30.0-42.4
2003	475	67.7	61.5-74.3	271	92.9	81.4-105.5	204	50.5	43.7-58.0	373	55.1	49.5-61.2	226	80.8	69.8-92.9	147	37.3	31.5-44.0
2004	506	69.0	62.9-75.5	299	97.0	85.6-109.5	207	49.3	42.8-56.6	372	54.1	48.6-60.1	224	81.9	70.7-94.2	148	36.3	30.6-42.8
2005	499	68.2	62.2-74.7	297	99.7	87.7-112.7	202	47.9	41.5-55.1	384	52.7	47.4-58.5	219	73.0	62.8-84.3	165	39.5	33.6-46.0
2006	496	67.3	61.3-73.7	289	94.0	82.7-106.3	207	49.4	42.8-56.7	324	44.6	39.7-49.9	186	61.1	52.0-71.2	138	32.9	27.6-39.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Lung and bronchus

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	524	67.4	61.6-73.6	303	93.9	82.9-105.8	221	49.2	42.8-56.2	369	47.8	42.9-53.1	209	65.0	55.9-75.1	160	36.1	30.7-42.2
2008	504	63.4	57.8-69.4	272	82.4	72.2-93.5	232	50.8	44.4-57.9	326	42.1	37.5-47.1	182	56.4	47.9-65.9	144	32.0	26.9-37.8
2009	555	67.1	61.4-73.1	306	88.6	78.2-99.8	249	52.7	46.2-59.7	356	44.1	39.5-49.1	203	60.7	52.1-70.3	153	32.9	27.8-38.6
2010	564	66.1	60.6-72.0	298	83.5	73.6-94.3	266	54.7	48.2-61.8	378	46.4	41.6-51.4	223	68.3	59.0-78.5	155	32.6	27.6-38.3
2011	543	61.0	55.8-66.5	298	79.7	70.3-90.0	245	48.2	42.3-54.8	401	46.7	42.1-51.6	228	63.7	55.1-73.1	173	35.0	29.9-40.7
2012	578	63.3	58.1-68.9	339	88.4	78.6-99.0	239	45.8	40.1-52.1	379	42.9	38.5-47.5	215	59.3	51.1-68.3	164	32.0	27.2-37.3
2013	557	58.0	53.2-63.2	297	75.1	66.3-84.6	260	46.4	40.9-52.5	376	41.4	37.2-45.9	211	55.7	48.0-64.3	165	31.2	26.5-36.4
2014	563	57.9	53.1-63.1	308	75.0	66.4-84.5	255	45.8	40.2-51.9	361	37.3	33.4-41.5	205	50.7	43.6-58.6	156	27.8	23.6-32.7
2015	628	62.5	57.6-67.8	349	84.8	75.6-94.7	279	47.4	41.9-53.4	361	37.8	33.9-42.1	194	49.2	42.1-57.0	167	30.2	25.7-35.2
2016	604	59.2	54.4-64.2	326	75.3	66.8-84.4	278	47.6	42.1-53.7	306	30.3	26.9-34.0	178	43.7	37.2-51.0	128	21.4	17.8-25.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Melanoma of the skin

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	2	0.6	0.1-2.2	1	0.9	0.0-4.7	1	0.4	0.0-2.7	4	1.7	0.4-4.4	1	0.5	0.0-3.8	3	2.4	0.4-6.6
1977	6	3.0	1.0-6.4	5	5.9	1.5-13.9	1	1.1	0.0-4.8	3	1.4	0.2-3.9	2	3.3	0.3-10.8	1	0.4	0.0-2.5
1978	4	1.4	0.3-3.7	2	2.8	0.2-9.6	2	0.8	0.1-3.2	1	0.4	0.0-2.1	0	0.0	0.0-2.9	1	0.7	0.0-3.4
1979	1	0.2	0.0-1.4	1	0.5	0.0-3.4	0	0.0	0.0-1.7	2	0.9	0.1-3.1	1	0.5	0.0-3.5	1	1.0	0.0-4.2
1980	4	1.4	0.3-3.6	2	1.0	0.1-4.1	2	1.5	0.2-4.7	0	0.0	0.0-1.0	0	0.0	0.0-2.6	0	0.0	0.0-1.6
1981	7	1.4	0.5-3.1	2	0.7	0.1-3.5	5	1.9	0.6-4.6	1	0.4	0.0-1.8	1	1.0	0.0-4.8	0	0.0	0.0-1.6
1982	9	1.7	0.8-3.3	6	2.6	0.9-6.0	3	0.9	0.2-3.0	3	0.7	0.2-2.2	2	1.2	0.1-4.5	1	0.4	0.0-2.2
1983	2	0.6	0.1-2.0	2	1.4	0.1-5.0	0	0.0	0.0-1.5	2	0.6	0.1-1.9	1	0.4	0.0-2.9	1	0.6	0.0-2.8
1984	5	1.5	0.4-3.4	2	1.7	0.1-5.9	3	1.4	0.3-4.0	1	0.1	0.0-1.0	1	0.2	0.0-2.5	0	0.0	0.0-1.4
1985	6	1.6	0.5-3.6	3	1.4	0.3-4.3	3	1.6	0.3-4.5	2	0.7	0.1-2.3	1	1.2	0.0-5.3	1	0.5	0.0-2.3
1986	6	1.2	0.4-2.7	3	1.1	0.2-3.7	3	1.2	0.2-3.5	7	1.3	0.5-2.7	5	2.1	0.7-5.2	2	0.7	0.1-2.5
1987	5	1.2	0.4-2.8	3	1.7	0.3-5.2	2	0.8	0.1-2.9	1	0.2	0.0-1.1	1	0.5	0.0-2.8	0	0.0	0.0-1.3
1988	5	1.0	0.3-2.3	2	1.1	0.1-4.0	3	1.0	0.2-2.9	2	0.4	0.0-1.5	2	1.0	0.1-3.8	0	0.0	0.0-1.2
1989	4	1.0	0.2-2.5	2	0.6	0.1-2.7	2	1.0	0.1-3.3	5	1.1	0.3-2.5	2	1.1	0.1-4.1	3	1.1	0.2-3.2
1990	4	0.8	0.2-2.0	1	0.4	0.0-2.5	3	1.0	0.2-3.0	0	0.0	0.0-0.7	0	0.0	0.0-1.9	0	0.0	0.0-1.2
1991	5	1.0	0.3-2.4	1	1.4	0.0-5.5	4	1.1	0.3-3.0	2	0.5	0.1-1.8	0	0.0	0.0-1.9	2	0.8	0.1-2.7
1992	3	0.5	0.1-1.4	2	0.7	0.1-2.8	1	0.3	0.0-1.7	1	0.1	0.0-0.9	1	0.3	0.0-2.2	0	0.0	0.0-1.1
1993	8	1.6	0.7-3.2	1	1.2	0.0-5.1	7	2.3	0.9-4.6	3	0.4	0.1-1.3	0	0.0	0.0-1.7	3	0.8	0.2-2.3
1994	5	0.9	0.3-2.1	2	0.5	0.1-2.4	3	1.0	0.2-2.9	3	0.6	0.1-1.6	0	0.0	0.0-1.7	3	1.0	0.2-2.8
1995	6	0.9	0.3-2.0	3	0.8	0.2-2.8	3	0.9	0.2-2.5	4	0.5	0.1-1.3	1	0.2	0.0-1.8	3	0.7	0.1-2.2
1996	5	0.8	0.2-2.0	3	1.6	0.2-5.0	2	0.5	0.1-1.9	4	0.6	0.2-1.5	1	0.3	0.0-2.0	3	0.8	0.2-2.4
1997	6	1.1	0.4-2.3	1	0.3	0.0-2.0	5	1.5	0.5-3.5	2	0.4	0.0-1.3	1	0.2	0.0-1.7	1	0.4	0.0-1.8
1998	8	1.3	0.5-2.6	3	0.9	0.2-2.8	5	1.5	0.5-3.4	3	0.4	0.1-1.2	1	0.2	0.0-1.6	2	0.6	0.1-2.0
1999	4	0.5	0.1-1.3	2	0.5	0.1-2.1	2	0.5	0.1-1.8	1	0.2	0.0-0.8	1	0.3	0.0-2.0	0	0.0	0.0-0.9
2000	0	0.0	0.0-0.5	0	0.0	0.0-1.3	0	0.0	0.0-0.9	3	0.6	0.1-1.7	2	1.1	0.1-3.6	1	0.3	0.0-1.6
2001	6	0.7	0.3-1.6	2	0.5	0.1-2.0	4	0.9	0.3-2.4	4	0.7	0.2-1.7	2	0.6	0.1-2.4	2	0.6	0.1-2.1
2002	8	1.3	0.5-2.5	4	1.5	0.4-3.9	4	1.1	0.3-2.8	2	0.4	0.0-1.3	1	0.9	0.0-3.7	1	0.3	0.0-1.4
2003	8	1.1	0.5-2.2	2	0.9	0.1-3.1	6	1.3	0.5-2.9	5	0.7	0.2-1.6	2	0.9	0.1-3.1	3	0.6	0.1-1.8
2004	9	1.1	0.5-2.0	5	1.3	0.4-3.2	4	0.9	0.2-2.2	1	0.1	0.0-0.6	1	0.2	0.0-1.5	0	0.0	0.0-0.8
2005	8	0.9	0.4-1.9	5	1.1	0.3-2.7	3	0.7	0.1-2.0	2	0.2	0.0-0.8	1	0.2	0.0-1.4	1	0.2	0.0-1.1
2006	10	1.4	0.6-2.5	6	1.7	0.6-3.9	4	1.1	0.3-2.6	4	0.6	0.1-1.4	2	0.7	0.1-2.5	2	0.5	0.1-1.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Melanoma of the skin

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	10	1.2	0.6-2.3	1	0.2	0.0-1.4	9	2.0	0.9-3.8	1	0.1	0.0-0.6	0	0.0	0.0-1.1	1	0.2	0.0-1.1
2008	12	1.4	0.7-2.5	4	0.9	0.2-2.4	8	1.8	0.8-3.5	2	0.3	0.0-0.9	0	0.0	0.0-1.0	2	0.5	0.1-1.6
2009	8	0.8	0.3-1.6	2	0.4	0.0-1.5	6	1.1	0.4-2.5	4	0.5	0.1-1.3	2	0.6	0.1-2.3	2	0.4	0.0-1.5
2010	10	1.0	0.5-1.8	3	0.7	0.1-2.2	7	1.2	0.5-2.6	5	0.6	0.2-1.4	1	0.2	0.0-1.3	4	0.8	0.2-2.1
2011	9	1.0	0.4-1.9	4	1.0	0.2-2.8	5	1.0	0.3-2.3	4	0.5	0.1-1.3	1	0.2	0.0-1.4	3	0.7	0.1-1.9
2012	7	0.7	0.3-1.4	4	1.1	0.2-2.8	3	0.5	0.1-1.5	4	0.5	0.1-1.2	0	0.0	0.0-0.9	4	0.8	0.2-2.0
2013	15	1.7	0.9-2.8	5	1.6	0.5-3.6	10	1.8	0.9-3.4	6	0.5	0.2-1.2	3	0.6	0.1-1.9	3	0.5	0.1-1.4
2014	13	1.2	0.6-2.1	6	1.1	0.4-2.5	7	1.3	0.5-2.6	1	0.1	0.0-0.6	0	0.0	0.0-0.8	1	0.2	0.0-1.0
2015	13	1.3	0.7-2.2	7	1.4	0.5-3.0	6	1.2	0.4-2.5	3	0.4	0.1-1.0	2	0.5	0.1-1.8	1	0.2	0.0-1.1
2016	11	1.0	0.5-1.8	6	1.1	0.4-2.5	5	0.8	0.3-2.0	1	0.1	0.0-0.6	1	0.3	0.0-1.3	0	0.0	0.0-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Female breast

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	151	79.1	66.1-93.6	67	37.3	28.3-48.1
1977	169	87.6	74.1-102.7	65	34.8	26.3-44.9
1978	185	90.7	77.5-105.4	55	26.1	19.4-34.3
1979	174	83.8	71.1-98.0	65	30.7	23.3-39.6
1980	203	90.6	78.0-104.5	70	32.5	25.0-41.4
1981	205	90.0	77.7-103.7	84	39.6	31.2-49.4
1982	202	85.0	73.3-97.9	90	38.1	30.4-47.1
1983	223	92.5	80.4-105.9	77	31.5	24.7-39.6
1984	221	89.1	77.3-102.0	97	43.2	34.7-53.0
1985	245	96.4	84.3-109.6	91	36.4	29.1-45.0
1986	256	96.5	84.8-109.4	89	34.6	27.6-42.7
1987	250	94.1	82.5-106.8	84	32.6	25.9-40.6
1988	277	98.4	87.0-110.9	106	38.3	31.3-46.5
1989	248	87.0	76.3-98.8	106	39.1	31.8-47.4
1990	302	101.9	90.5-114.2	90	30.8	24.6-38.0
1991	313	102.3	91.1-114.5	113	37.9	31.1-45.6
1992	328	104.2	93.0-116.3	114	36.6	30.1-44.0
1993	339	105.4	94.2-117.4	109	34.6	28.3-41.9
1994	349	103.5	92.7-115.1	116	36.2	29.8-43.5
1995	346	100.7	90.2-112.1	108	33.0	27.0-39.9
1996	397	110.9	100.1-122.5	108	31.5	25.7-38.1
1997	429	118.6	107.5-130.6	115	34.0	27.9-40.9
1998	441	116.0	105.3-127.5	122	33.7	27.9-40.3
1999	446	115.5	104.9-126.9	98	25.9	20.9-31.6
2000	399	100.3	90.5-110.8	127	34.2	28.4-40.7
2001	415	101.9	92.2-112.3	121	30.3	25.1-36.3
2002	461	107.6	97.9-118.0	123	29.8	24.7-35.6
2003	466	107.7	98.0-118.1	119	28.1	23.2-33.7
2004	508	114.5	104.6-125.1	128	30.5	25.4-36.3
2005	511	114.1	104.2-124.5	126	28.7	23.8-34.2
2006	550	118.0	108.2-128.4	121	26.7	22.1-31.9
2007	552	116.0	106.4-126.2	135	29.2	24.4-34.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Female breast

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	580	118.7	109.2-128.9	133	27.3	22.8-32.4
2009	590	119.5	109.9-129.7	151	31.5	26.6-37.0
2010	591	115.4	106.1-125.2	126	24.7	20.5-29.5
2011	680	128.5	118.9-138.7	148	28.7	24.2-33.8
2012	620	115.5	106.4-125.1	121	23.3	19.3-27.9
2013	655	119.9	110.8-129.6	140	25.9	21.7-30.7
2014	710	124.8	115.7-134.5	138	24.7	20.7-29.3
2015	720	124.0	115.0-133.6	124	21.1	17.5-25.2
2016	771	129.5	120.3-139.1	117	20.5	16.9-24.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Cervix uteri

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	70	35.8	27.3-46.0	29	16.5	10.6-24.3
1977	58	27.2	20.2-35.8	27	14.6	9.3-21.7
1978	57	25.7	19.2-33.7	21	10.3	6.1-16.0
1979	61	25.8	19.5-33.5	24	10.7	6.7-16.1
1980	58	26.1	19.5-34.1	25	12.2	7.7-18.3
1981	37	15.3	10.6-21.3	27	13.0	8.4-19.0
1982	44	16.8	12.1-22.7	16	7.0	3.9-11.4
1983	49	18.7	13.7-25.0	22	8.6	5.4-13.0
1984	52	19.3	14.2-25.5	13	6.9	3.6-11.8
1985	40	15.6	11.0-21.4	23	8.9	5.5-13.6
1986	53	18.8	13.9-24.8	20	8.0	4.8-12.3
1987	49	17.5	12.8-23.3	16	6.0	3.3-9.9
1988	46	15.8	11.5-21.2	18	6.6	3.8-10.5
1989	50	17.2	12.6-22.7	19	6.9	4.1-10.8
1990	63	19.0	14.5-24.4	18	6.2	3.6-9.9
1991	53	16.2	12.0-21.4	19	6.2	3.7-9.7
1992	42	11.4	8.2-15.6	14	4.2	2.3-7.1
1993	52	14.3	10.6-18.9	25	7.4	4.7-11.0
1994	47	12.6	9.2-16.9	15	4.2	2.4-7.0
1995	53	14.2	10.5-18.7	16	4.7	2.6-7.6
1996	73	19.3	15.0-24.3	19	5.2	3.1-8.2
1997	34	8.9	6.1-12.5	24	6.4	4.1-9.5
1998	51	12.0	8.9-15.9	17	4.6	2.7-7.4
1999	51	13.1	9.7-17.3	13	3.3	1.7-5.7
2000	42	10.2	7.3-13.8	19	4.6	2.7-7.1
2001	52	12.7	9.4-16.7	10	2.5	1.2-4.6
2002	46	10.6	7.7-14.1	16	4.0	2.3-6.5
2003	56	12.8	9.6-16.6	19	4.4	2.6-6.9
2004	49	10.6	7.8-14.0	14	3.0	1.6-5.0
2005	50	10.9	8.0-14.4	17	3.6	2.1-5.8
2006	47	9.8	7.2-13.1	15	3.3	1.9-5.5
2007	48	10.1	7.4-13.4	21	4.6	2.8-7.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Cervix uteri

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	53	10.5	7.8-13.7	15	3.1	1.7-5.1
2009	43	8.7	6.3-11.7	26	5.0	3.3-7.4
2010	45	8.7	6.3-11.7	13	2.5	1.3-4.3
2011	45	8.9	6.5-11.9	14	2.6	1.4-4.4
2012	44	8.1	5.8-10.9	19	3.6	2.1-5.6
2013	62	11.2	8.6-14.4	23	4.2	2.6-6.3
2014	50	8.5	6.3-11.2	15	2.5	1.4-4.1
2015	53	9.5	7.1-12.4	17	3.1	1.8-5.0
2016	45	7.4	5.4-10.0	6	1.2	0.4-2.5

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Corpus uterus and uterus, not otherwise specified

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	31	17.9	11.8-25.7	13	9.6	4.9-16.5
1977	36	19.4	13.2-27.3	15	8.2	4.4-13.9
1978	25	13.8	8.6-20.7	9	4.6	2.0-8.9
1979	37	16.3	11.3-22.7	14	7.6	4.0-12.9
1980	50	23.8	17.4-31.7	21	9.8	6.0-15.0
1981	28	12.1	7.9-17.7	21	9.9	6.0-15.3
1982	31	13.4	9.0-19.2	11	4.2	2.1-7.5
1983	31	12.9	8.7-18.4	13	6.0	3.1-10.3
1984	33	14.2	9.6-20.1	16	6.7	3.7-10.9
1985	49	19.8	14.5-26.2	19	9.5	5.6-14.8
1986	38	15.4	10.8-21.2	12	4.9	2.5-8.5
1987	49	19.1	14.0-25.3	11	4.7	2.3-8.4
1988	46	17.9	12.9-23.9	18	7.2	4.2-11.4
1989	35	12.6	8.7-17.6	24	9.4	5.9-14.0
1990	51	17.8	13.1-23.4	10	4.0	1.9-7.4
1991	42	14.5	10.4-19.6	17	5.8	3.3-9.2
1992	63	21.6	16.5-27.7	23	8.7	5.5-13.1
1993	60	19.4	14.8-25.0	16	6.0	3.4-9.6
1994	66	20.3	15.6-25.9	13	4.2	2.2-7.2
1995	52	15.8	11.7-20.7	18	5.9	3.4-9.2
1996	79	23.2	18.3-29.0	24	7.4	4.7-10.9
1997	70	20.8	16.1-26.2	12	3.7	1.9-6.5
1998	79	21.5	17.0-26.8	28	8.5	5.6-12.3
1999	84	22.7	18.1-28.2	24	7.0	4.4-10.4
2000	81	21.8	17.2-27.1	30	8.5	5.7-12.1
2001	80	20.8	16.4-25.9	25	7.0	4.5-10.3
2002	75	18.2	14.2-22.8	29	7.8	5.2-11.2
2003	83	19.9	15.8-24.7	36	9.1	6.3-12.6
2004	110	25.6	21.0-30.9	20	4.4	2.7-6.8
2005	102	23.8	19.4-29.0	33	8.3	5.7-11.6
2006	102	22.1	18.0-26.9	32	7.3	5.0-10.3
2007	126	26.7	22.2-31.9	41	9.7	6.9-13.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Corpus uterus and uterus, not otherwise specified

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	114	24.1	19.8-29.0	46	10.2	7.5-13.7
2009	137	28.6	23.9-33.9	30	6.7	4.5-9.5
2010	131	26.1	21.8-31.1	46	9.6	7.0-12.9
2011	135	25.0	20.9-29.6	36	7.3	5.1-10.2
2012	159	29.7	25.2-34.7	47	9.5	7.0-12.7
2013	135	23.6	19.7-28.0	45	8.5	6.1-11.4
2014	173	30.1	25.7-35.0	57	10.4	7.9-13.6
2015	169	28.6	24.4-33.4	40	7.2	5.1-9.8
2016	186	31.1	26.7-36.1	42	7.2	5.2-9.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Ovary

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	28	16.5	10.5-24.3	17	11.7	6.6-19.0
1977	29	13.2	8.5-19.4	17	8.4	4.7-13.7
1978	24	10.7	6.8-16.1	13	6.9	3.5-12.0
1979	34	16.4	11.0-23.2	22	10.1	6.1-15.6
1980	26	12.0	7.7-17.7	20	9.5	5.7-14.7
1981	28	12.3	7.9-18.0	14	6.2	3.3-10.5
1982	36	14.3	9.9-19.9	16	6.7	3.8-11.0
1983	29	12.2	8.0-17.7	14	6.1	3.2-10.3
1984	28	10.8	7.0-15.7	24	10.5	6.6-15.7
1985	35	14.3	9.9-20.0	13	5.4	2.8-9.2
1986	37	14.3	9.9-19.9	21	8.5	5.2-13.0
1987	28	11.1	7.3-16.2	27	10.9	7.1-15.9
1988	38	13.3	9.3-18.3	20	7.5	4.6-11.7
1989	38	13.2	9.2-18.2	23	8.3	5.2-12.4
1990	37	12.2	8.5-17.0	28	10.2	6.7-14.8
1991	23	8.8	5.5-13.2	20	7.2	4.3-11.1
1992	31	9.7	6.5-13.9	34	11.9	8.1-16.6
1993	42	12.3	8.8-16.7	19	6.5	3.9-10.1
1994	32	9.7	6.6-13.7	23	7.4	4.6-11.1
1995	42	11.8	8.4-16.0	23	7.9	5.0-11.8
1996	40	11.8	8.4-16.1	19	5.7	3.4-8.9
1997	52	14.5	10.7-19.0	26	7.6	4.9-11.1
1998	37	9.9	6.9-13.7	23	7.0	4.4-10.5
1999	47	12.7	9.3-16.9	26	7.2	4.7-10.6
2000	42	11.2	8.0-15.1	31	8.5	5.7-12.1
2001	41	10.1	7.2-13.7	24	6.6	4.2-9.7
2002	44	10.4	7.5-14.1	29	7.5	5.0-10.8
2003	28	6.4	4.2-9.3	35	8.5	5.9-11.9
2004	46	10.9	7.9-14.6	28	7.1	4.7-10.2
2005	40	8.7	6.2-11.9	26	6.0	3.9-8.8
2006	40	9.1	6.4-12.4	31	7.3	4.9-10.4
2007	35	7.4	5.2-10.4	26	6.1	4.0-8.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Ovary

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	48	10.4	7.7-13.9	21	4.9	3.0-7.5
2009	52	10.3	7.7-13.6	25	5.3	3.4-7.8
2010	43	8.3	6.0-11.3	26	5.3	3.4-7.8
2011	50	9.3	6.9-12.4	30	6.3	4.2-9.0
2012	63	11.5	8.8-14.8	34	6.3	4.4-8.9
2013	41	7.1	5.1-9.7	34	6.3	4.3-8.9
2014	60	10.4	7.9-13.4	37	6.5	4.5-9.0
2015	47	8.4	6.1-11.2	30	5.4	3.6-7.7
2016	50	7.8	5.7-10.3	22	3.8	2.4-5.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Prostate

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	123	126.0	101.5-153.8	65	71.9	53.1-94.3
1977	128	114.2	92.7-138.5	64	79.0	58.8-102.8
1978	143	128.1	104.9-154.2	59	54.2	39.8-71.5
1979	144	126.1	103.5-151.5	73	67.8	51.5-87.0
1980	174	139.9	117.5-164.9	56	57.7	42.1-76.3
1981	169	142.7	119.4-168.6	82	76.0	58.7-96.1
1982	157	121.3	101.4-143.4	74	64.2	49.1-82.0
1983	185	143.2	121.2-167.5	73	65.3	49.9-83.2
1984	171	123.2	103.6-144.9	67	51.9	39.0-67.2
1985	182	128.6	108.7-150.7	72	58.0	44.3-74.0
1986	206	146.6	125.2-170.0	92	74.5	58.5-92.8
1987	187	129.4	109.9-151.0	83	75.3	58.6-94.5
1988	210	138.1	118.2-160.1	100	83.0	66.2-102.1
1989	207	131.4	112.4-152.4	82	59.1	46.1-74.2
1990	257	161.1	140.0-184.0	96	67.9	53.7-84.2
1991	357	209.5	186.0-234.7	104	82.1	65.9-100.5
1992	431	236.9	212.7-262.8	87	60.9	47.7-76.1
1993	514	270.5	245.3-297.2	102	72.5	57.9-89.2
1994	509	250.5	227.3-275.3	121	83.3	67.9-100.6
1995	561	267.1	243.4-292.4	113	70.2	56.7-85.5
1996	627	268.9	246.6-292.5	116	70.6	57.4-85.5
1997	618	266.3	243.9-290.0	108	63.4	50.9-77.5
1998	669	273.0	251.2-296.0	114	70.9	57.7-85.8
1999	705	274.4	253.1-296.8	94	54.6	43.3-67.5
2000	773	305.1	282.3-329.1	129	74.3	61.2-89.0
2001	786	292.2	270.4-315.1	110	63.3	51.3-76.9
2002	825	299.1	277.2-322.0	114	61.5	49.9-74.7
2003	821	282.0	261.6-303.4	112	56.5	45.8-68.6
2004	759	243.6	225.4-262.8	97	47.0	37.4-58.0
2005	807	252.3	234.0-271.6	107	45.2	36.5-55.1
2006	887	269.0	250.4-288.4	83	35.1	27.6-43.8
2007	964	285.2	266.3-304.9	99	43.3	34.7-53.0

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Prostate

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2008	932	259.9	242.5-278.2	103	40.2	32.4-49.1
2009	948	249.6	233.0-267.0	111	44.8	36.4-54.3
2010	977	251.5	234.9-268.8	113	44.3	36.1-53.5
2011	1014	247.9	232.0-264.6	88	30.7	24.3-38.1
2012	900	208.2	194.1-223.1	97	35.2	28.2-43.1
2013	946	214.3	200.2-229.2	97	34.0	27.4-41.6
2014	920	200.0	186.6-214.1	102	32.4	26.1-39.5
2015	944	196.1	183.2-209.7	77	24.3	19.0-30.5
2016	933	190.3	177.6-203.6	81	23.4	18.4-29.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Testis

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	3	0.9	0.2-4.1	1	0.9	0.0-4.7
1977	5	1.5	0.5-4.7	1	0.4	0.0-3.5
1978	2	0.5	0.1-3.3	1	0.3	0.0-3.2
1980	7	2.1	0.8-5.2	2	0.6	0.1-3.4
1981	4	1.6	0.4-4.8	0	0.0	0.0-2.5
1982	3	1.2	0.2-4.1	0	0.0	0.0-2.4
1983	2	0.4	0.1-2.7	1	0.2	0.0-2.5
1984	5	1.5	0.4-4.2	0	0.0	0.0-2.2
1985	6	1.9	0.6-4.7	1	0.4	0.0-2.7
1986	6	1.8	0.6-4.5	0	0.0	0.0-2.1
1987	5	1.2	0.4-3.5	1	0.2	0.0-2.2
1988	5	1.2	0.4-3.5	0	0.0	0.0-2.0
1989	2	0.5	0.1-2.7	0	0.0	0.0-1.9
1990	3	0.7	0.1-2.8	0	0.0	0.0-1.9
1991	3	0.6	0.1-2.7	1	0.4	0.0-2.4
1992	8	2.6	0.6-6.5	1	0.3	0.0-2.1
1993	3	0.6	0.1-2.4	1	0.2	0.0-1.9
1994	4	0.8	0.2-2.6	0	0.0	0.0-1.7
1995	2	0.5	0.1-2.3	0	0.0	0.0-1.6
1996	4	0.9	0.2-2.7	0	0.0	0.0-1.5
1997	6	1.2	0.4-3.0	0	0.0	0.0-1.5
1998	6	1.2	0.4-3.1	0	0.0	0.0-1.4
1999	8	1.5	0.6-3.2	1	0.2	0.0-1.7
2000	5	1.5	0.4-3.8	0	0.0	0.0-1.3
2001	8	1.7	0.7-3.7	0	0.0	0.0-1.3
2002	6	1.2	0.4-2.9	0	0.0	0.0-1.3
2003	9	1.8	0.8-3.7	0	0.0	0.0-1.2
2004	5	1.0	0.3-2.5	0	0.0	0.0-1.2
2005	3	0.5	0.1-1.8	0	0.0	0.0-1.1
2006	6	1.2	0.4-2.8	0	0.0	0.0-1.1
2007	9	1.7	0.8-3.4	0	0.0	0.0-1.1
2008	6	1.2	0.4-2.7	1	0.1	0.0-1.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Testis

	Incidence			Mortality		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2009	4	0.7	0.2-2.1	0	0.0	0.0-1.0
2010	5	1.0	0.3-2.5	1	0.2	0.0-1.3
2011	7	1.3	0.5-2.9	0	0.0	0.0-0.9
2012	6	1.2	0.4-2.7	0	0.0	0.0-0.9
2013	10	1.8	0.9-3.4	1	0.2	0.0-1.2
2014	8	1.4	0.6-2.9	0	0.0	0.0-0.8
2015	8	1.5	0.7-3.1	0	0.0	0.0-0.8
2016	8	1.3	0.5-2.6	0	0.0	0.0-0.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Urinary bladder (incl. in situ)

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	44	18.4	12.8-25.3	31	34.4	21.4-51.3	13	8.8	4.4-15.2	15	5.5	2.8-9.4	7	5.8	1.6-13.6	8	5.5	2.2-10.8
1977	29	10.9	6.9-16.1	13	10.4	5.1-18.4	16	10.6	5.7-17.6	14	6.7	3.4-11.3	8	11.1	4.3-22.0	6	4.2	1.4-9.2
1978	42	14.8	10.3-20.3	30	28.1	17.8-41.5	12	6.4	3.2-11.2	18	5.3	3.0-8.5	8	4.6	1.9-9.3	10	5.6	2.6-10.3
1979	55	18.2	13.3-24.1	30	23.0	14.9-33.6	25	14.6	9.1-21.8	23	9.9	6.0-15.0	14	17.3	8.7-29.5	9	5.9	2.5-11.1
1980	40	12.7	8.8-17.7	29	21.1	13.3-31.5	11	6.8	3.3-12.1	20	6.4	3.8-10.1	12	10.2	4.9-18.2	8	4.1	1.6-8.1
1981	42	12.7	8.9-17.4	23	16.2	9.6-25.1	19	10.2	5.9-15.9	16	5.0	2.7-8.4	13	11.7	5.4-21.0	3	1.3	0.3-3.7
1982	33	10.2	6.8-14.5	21	15.8	9.1-24.8	12	6.4	3.1-11.2	19	6.1	3.5-9.6	12	9.6	4.5-17.3	7	3.8	1.5-7.8
1983	35	9.7	6.6-13.6	18	11.1	6.1-18.2	17	8.5	4.8-13.6	18	5.4	3.1-8.5	7	5.3	2.0-10.8	11	5.4	2.6-9.7
1984	55	15.6	11.4-20.5	38	28.2	18.9-39.7	17	7.9	4.5-12.7	20	6.9	4.1-10.6	14	13.3	6.7-22.6	6	3.4	1.2-7.1
1985	45	12.0	8.5-16.2	27	18.2	11.4-27.0	18	8.1	4.7-12.9	16	4.8	2.6-7.8	10	8.4	3.8-15.6	6	2.8	1.0-6.0
1986	40	10.5	7.4-14.4	25	16.3	10.2-24.4	15	6.9	3.8-11.2	16	4.7	2.6-7.6	8	5.7	2.2-11.4	8	4.0	1.7-7.6
1987	34	9.1	6.1-12.8	20	12.6	7.3-19.9	14	6.6	3.6-11.0	13	3.9	2.0-6.6	6	4.9	1.7-10.4	7	3.3	1.3-6.7
1988	53	13.9	10.2-18.3	33	19.4	13.0-27.5	20	9.5	5.7-14.5	16	5.0	2.8-8.0	9	7.5	3.2-14.2	7	3.5	1.4-7.1
1989	54	12.1	9.0-15.9	34	19.3	12.7-27.6	20	7.6	4.6-11.8	22	6.7	4.1-10.0	15	13.0	6.9-21.5	7	3.4	1.3-6.7
1990	41	10.3	7.3-14.1	30	18.9	12.1-27.7	11	4.9	2.4-8.6	24	6.5	4.1-9.7	16	9.7	5.3-15.9	8	4.0	1.7-7.6
1991	64	14.1	10.7-18.0	47	27.8	19.6-37.9	17	6.0	3.5-9.6	24	6.1	3.8-9.0	12	7.5	3.4-13.7	12	5.1	2.6-8.8
1992	39	8.3	5.9-11.5	30	16.2	10.5-23.5	9	3.3	1.5-6.2	12	2.6	1.3-4.4	9	4.8	2.0-9.2	3	1.1	0.2-3.1
1993	49	10.0	7.3-13.4	24	10.2	6.2-15.6	25	9.2	5.9-13.5	21	5.0	3.0-7.6	10	7.0	3.2-12.6	11	4.0	1.9-7.0
1994	45	9.7	7.0-13.0	32	19.0	12.4-27.3	13	4.5	2.4-7.6	15	3.9	2.1-6.3	9	6.0	2.6-11.4	6	2.5	0.9-5.3
1995	47	9.2	6.7-12.3	33	16.3	10.9-23.2	14	4.6	2.5-7.7	14	3.5	1.9-5.8	8	6.8	2.7-13.2	6	2.3	0.8-4.8
1996	60	11.5	8.6-14.8	30	14.8	9.4-21.7	30	9.7	6.5-13.8	23	4.8	3.0-7.2	13	7.7	3.9-13.3	10	3.3	1.6-6.0
1997	65	12.7	9.7-16.3	48	26.6	18.8-36.2	17	5.6	3.2-8.8	16	3.0	1.7-4.9	9	4.5	1.8-8.8	7	2.3	0.9-4.6
1998	87	16.9	13.4-20.9	58	29.7	21.7-39.2	29	9.6	6.4-13.7	20	4.1	2.5-6.3	10	5.7	2.6-10.4	10	3.3	1.6-5.9
1999	64	11.3	8.6-14.5	41	19.2	13.3-26.6	23	6.7	4.2-10.1	20	3.8	2.3-5.9	12	5.9	2.7-10.5	8	2.6	1.1-4.9
2000	79	12.9	10.1-16.2	57	22.2	16.3-29.3	22	6.5	4.1-9.8	15	3.1	1.7-5.0	7	4.6	1.7-9.3	8	2.4	1.1-4.7
2001	77	12.8	10.0-16.1	50	21.5	15.3-29.0	27	7.7	5.1-11.2	19	3.5	2.1-5.4	10	5.3	2.2-10.0	9	2.8	1.3-5.1
2002	66	10.8	8.3-13.8	43	17.0	12.0-23.2	23	6.4	4.0-9.5	19	3.5	2.0-5.4	10	5.2	2.3-9.7	9	2.6	1.2-4.8
2003	84	12.4	9.8-15.4	56	20.5	14.9-27.2	28	7.2	4.8-10.4	24	4.1	2.6-6.1	13	6.3	3.2-10.8	11	3.0	1.5-5.3
2004	88	13.5	10.7-16.7	60	21.5	16.1-28.0	28	7.6	5.0-10.9	21	3.5	2.1-5.3	11	4.2	1.9-7.7	10	2.8	1.4-5.1
2005	86	12.6	10.0-15.6	63	23.8	17.8-30.9	23	5.7	3.6-8.6	23	4.2	2.6-6.2	13	7.1	3.6-11.9	10	2.8	1.3-5.0
2006	82	11.5	9.1-14.3	55	19.8	14.6-26.0	27	6.1	4.0-8.9	12	1.8	0.9-3.1	8	2.8	1.1-5.7	4	1.1	0.3-2.7

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Urinary bladder (incl. in situ)

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	84	11.5	9.2-14.3	58	19.1	14.2-24.8	26	6.2	4.0-9.1	31	4.7	3.1-6.6	15	5.7	3.1-9.4	16	3.9	2.2-6.3
2008	103	13.5	10.9-16.4	70	22.8	17.4-29.3	33	7.5	5.1-10.6	18	2.7	1.6-4.3	12	4.9	2.5-8.4	6	1.4	0.5-3.0
2009	100	13.1	10.6-16.0	67	21.4	16.2-27.5	33	7.5	5.1-10.5	27	3.8	2.5-5.5	20	7.6	4.4-11.8	7	1.7	0.7-3.4
2010	92	11.2	8.9-13.8	68	21.7	16.5-28.0	24	4.8	3.1-7.2	23	3.1	1.9-4.6	9	3.6	1.6-6.7	14	2.9	1.6-4.9
2011	132	15.2	12.7-18.1	102	29.4	23.6-36.1	30	5.9	3.9-8.4	27	3.5	2.3-5.1	19	6.6	3.8-10.3	8	1.8	0.8-3.6
2012	120	14.6	12.0-17.5	80	24.5	19.1-30.8	40	8.4	5.9-11.4	23	2.7	1.7-4.1	15	4.5	2.4-7.6	8	1.6	0.7-3.2
2013	123	14.1	11.6-16.9	83	22.9	18.0-28.6	40	7.8	5.5-10.6	30	3.6	2.4-5.1	19	5.8	3.3-9.2	11	2.2	1.1-3.9
2014	112	12.1	9.9-14.7	75	19.5	15.1-24.7	37	6.8	4.8-9.4	24	2.9	1.8-4.3	17	5.5	3.1-8.8	7	1.3	0.5-2.7
2015	124	13.2	10.9-15.8	86	22.6	17.8-28.2	38	7.0	4.9-9.6	26	2.7	1.7-3.9	14	3.6	1.9-6.0	12	2.1	1.1-3.6
2016	116	11.6	9.5-13.9	73	17.3	13.3-21.9	43	7.4	5.3-10.0	19	2.2	1.3-3.4	9	2.9	1.3-5.4	10	1.8	0.9-3.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Kidney and renal pelvis

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	31	9.8	6.4-14.2	20	12.7	7.3-20.5	11	7.3	3.5-13.0	10	2.8	1.2-5.4	6	3.4	1.2-7.9	4	2.2	0.4-5.9
1977	27	7.9	4.9-11.9	14	11.7	5.4-21.0	13	5.8	3.0-10.2	7	2.0	0.7-4.3	4	3.3	0.7-9.0	3	1.1	0.2-3.7
1978	30	9.4	6.0-13.9	17	10.7	5.6-18.1	13	8.0	4.0-14.0	15	4.5	2.3-7.7	8	3.5	1.4-7.6	7	4.7	1.8-9.6
1979	30	7.7	5.0-11.1	21	11.1	6.8-17.3	9	4.6	2.0-8.9	15	4.5	2.4-7.7	13	10.0	4.8-17.8	2	0.8	0.1-3.1
1980	21	5.5	3.2-8.6	11	7.0	3.0-13.6	10	4.5	2.0-8.5	10	2.7	1.2-5.1	7	4.8	1.4-11.0	3	1.3	0.3-3.9
1981	30	7.1	4.6-10.3	22	11.6	6.6-18.8	8	3.7	1.6-7.2	12	2.5	1.3-4.4	8	3.8	1.6-7.8	4	1.5	0.4-3.9
1982	31	7.9	5.1-11.4	18	10.0	5.4-16.6	13	6.1	3.1-10.6	19	5.6	3.2-8.9	9	4.5	1.9-9.0	10	5.8	2.7-10.5
1983	26	6.1	3.9-9.1	19	9.9	5.5-16.2	7	3.2	1.3-6.6	10	2.4	1.1-4.6	7	4.3	1.3-9.7	3	1.3	0.3-3.7
1984	44	10.0	7.1-13.6	28	14.8	8.9-22.7	16	6.7	3.8-10.9	12	3.0	1.5-5.3	7	4.7	1.7-9.9	5	2.0	0.6-4.6
1985	46	10.4	7.4-14.0	25	13.9	8.4-21.3	21	8.1	4.9-12.5	16	3.8	2.1-6.3	11	4.8	2.4-8.8	5	2.6	0.8-5.8
1986	39	8.6	6.1-11.9	20	10.3	6.1-16.2	19	7.3	4.3-11.5	20	4.5	2.7-7.1	11	6.1	2.6-11.6	9	3.5	1.6-6.7
1987	36	8.5	5.8-11.9	23	14.4	8.7-22.1	13	4.9	2.5-8.4	17	4.3	2.4-7.0	5	3.1	0.9-7.4	12	5.1	2.6-8.9
1988	40	8.4	5.8-11.5	18	9.9	5.4-16.2	22	7.7	4.7-11.8	17	4.1	2.3-6.7	9	6.4	2.7-12.3	8	2.9	1.2-5.8
1989	33	6.7	4.5-9.6	19	9.5	5.2-15.6	14	5.0	2.7-8.4	12	3.0	1.5-5.2	6	4.0	1.3-8.7	6	2.4	0.8-5.1
1990	62	11.5	8.7-14.8	37	15.8	10.8-22.2	25	8.3	5.3-12.2	16	3.2	1.7-5.3	9	4.1	1.6-8.2	7	2.5	0.9-5.1
1991	59	10.4	7.8-13.6	38	15.8	10.8-22.2	21	6.6	4.0-10.1	20	4.5	2.7-7.0	12	7.2	3.3-13.0	8	3.0	1.2-5.8
1992	60	10.6	8.0-13.8	40	16.9	11.5-23.9	20	6.3	3.8-9.8	19	3.4	1.9-5.4	12	5.4	2.6-9.9	7	2.0	0.7-4.3
1993	65	11.0	8.4-14.1	42	16.7	11.6-23.1	23	6.8	4.3-10.3	22	4.2	2.6-6.4	10	4.9	2.2-9.2	12	3.9	2.0-6.8
1994	73	11.9	9.2-15.0	36	12.5	8.5-17.7	37	11.1	7.7-15.3	20	3.6	2.1-5.6	13	5.9	2.9-10.6	7	2.1	0.8-4.3
1995	77	13.6	10.6-17.2	38	17.4	11.6-24.8	39	11.9	8.4-16.3	23	4.5	2.8-6.8	14	7.6	3.6-13.4	9	3.0	1.3-5.6
1996	71	11.3	8.7-14.4	48	18.4	13.0-25.2	23	6.4	4.0-9.7	22	3.8	2.3-5.8	10	5.1	2.1-9.8	12	3.4	1.7-5.9
1997	72	11.7	9.1-14.9	45	17.7	12.5-24.2	27	7.7	5.0-11.1	27	4.9	3.1-7.1	14	5.6	2.8-9.9	13	4.2	2.2-7.1
1998	103	15.6	12.6-19.0	63	22.8	17.1-29.6	40	10.5	7.5-14.4	14	2.3	1.3-3.9	7	2.7	1.1-5.7	7	2.0	0.8-4.1
1999	70	10.7	8.3-13.6	38	13.1	9.0-18.2	32	8.7	5.9-12.3	20	3.5	2.1-5.4	10	3.7	1.6-7.2	10	3.2	1.5-5.7
2000	73	10.3	8.0-13.0	38	12.6	8.6-17.7	35	8.7	6.0-12.2	22	3.2	2.0-4.9	12	4.2	2.1-7.6	10	2.6	1.2-4.7
2001	83	11.5	9.1-14.4	51	15.3	11.1-20.5	32	8.3	5.6-11.7	23	3.6	2.2-5.4	13	4.0	2.0-7.0	10	2.9	1.4-5.3
2002	103	13.8	11.2-16.8	60	19.1	14.2-25.0	43	10.2	7.3-13.7	10	1.3	0.6-2.4	7	2.2	0.9-4.6	3	0.6	0.1-1.8
2003	93	13.0	10.4-16.0	53	15.8	11.5-21.0	40	10.3	7.3-14.1	26	3.9	2.5-5.8	20	7.9	4.4-12.7	6	1.6	0.6-3.4
2004	115	14.0	11.5-16.9	67	17.5	13.4-22.5	48	10.9	8.0-14.5	21	3.0	1.8-4.6	11	3.8	1.7-7.0	10	2.5	1.2-4.6
2005	107	13.2	10.7-16.0	59	16.6	12.4-21.7	48	10.6	7.8-14.1	28	3.6	2.3-5.3	21	6.2	3.6-9.7	7	1.7	0.6-3.4
2006	133	16.1	13.4-19.1	91	25.3	19.8-31.7	42	9.4	6.7-12.7	21	2.6	1.6-4.0	12	3.7	1.7-6.7	9	1.9	0.9-3.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Kidney and renal pelvis

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	147	17.5	14.7-20.7	83	22.5	17.6-28.3	64	13.8	10.6-17.6	32	4.3	2.9-6.0	23	7.3	4.4-11.1	9	2.1	0.9-3.9
2008	148	17.1	14.4-20.2	85	20.8	16.4-26.1	63	13.6	10.4-17.5	28	3.2	2.1-4.7	19	4.7	2.7-7.4	9	2.0	0.9-3.7
2009	177	20.0	17.1-23.3	104	25.1	20.2-30.7	73	15.4	12.1-19.5	33	3.9	2.6-5.5	20	5.0	2.9-7.8	13	2.8	1.5-4.7
2010	172	18.9	16.1-22.1	108	25.9	21.0-31.6	64	13.1	10.1-16.8	25	3.2	2.0-4.7	11	3.0	1.4-5.5	14	3.2	1.7-5.3
2011	165	18.1	15.4-21.2	111	28.2	22.8-34.3	54	10.6	7.9-13.8	24	2.8	1.8-4.2	14	3.8	2.0-6.6	10	2.1	1.0-3.8
2012	163	16.7	14.2-19.6	103	24.6	19.8-30.2	60	11.0	8.4-14.2	29	3.4	2.2-4.8	19	6.2	3.6-9.8	10	1.8	0.9-3.3
2013	176	17.4	14.9-20.2	119	25.6	21.1-30.9	57	10.5	7.9-13.7	17	1.9	1.1-3.0	11	2.6	1.2-4.7	6	1.1	0.4-2.5
2014	193	18.4	15.8-21.2	125	25.7	21.2-30.8	68	12.4	9.5-15.7	28	3.0	2.0-4.4	23	6.2	3.8-9.4	5	1.0	0.3-2.2
2015	225	21.2	18.5-24.3	145	30.6	25.6-36.3	80	13.8	10.9-17.2	40	4.1	2.9-5.6	25	5.7	3.6-8.7	15	2.9	1.6-4.8
2016	198	18.5	16.0-21.4	132	27.5	22.8-32.9	66	11.5	8.9-14.7	22	2.1	1.3-3.2	17	3.8	2.1-6.2	5	0.8	0.3-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Brain and other central nervous system

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	20	4.4	2.6-7.2	10	4.9	2.3-9.7	10	4.0	1.7-7.9	16	3.3	1.8-5.8	8	2.9	1.2-6.8	8	3.5	1.4-7.3
1977	26	5.3	3.4-8.0	10	5.2	2.4-10.2	16	5.5	3.0-9.2	10	2.1	1.0-4.0	4	1.9	0.5-5.6	6	2.3	0.8-5.3
1978	19	4.1	2.3-6.6	6	3.4	1.2-7.8	13	4.7	2.3-8.4	10	2.3	1.0-4.4	5	2.3	0.7-6.0	5	2.2	0.6-5.3
1979	24	4.8	3.0-7.3	16	7.5	4.2-12.7	8	2.6	1.1-5.5	12	2.4	1.2-4.4	6	2.5	0.9-6.2	6	2.3	0.8-5.2
1980	19	3.8	2.2-6.2	16	6.8	3.7-11.9	3	1.3	0.3-3.9	9	1.8	0.8-3.5	8	3.4	1.4-7.1	1	0.4	0.0-2.4
1981	14	2.3	1.2-4.0	7	2.6	1.0-6.0	7	2.1	0.8-4.5	6	1.2	0.4-2.7	5	2.4	0.8-5.9	1	0.2	0.0-1.9
1982	13	2.2	1.2-4.0	10	3.7	1.7-7.4	3	0.9	0.2-3.0	9	2.1	0.9-4.1	5	3.3	0.9-8.1	4	1.3	0.4-3.6
1983	15	2.4	1.3-4.1	7	2.2	0.8-5.3	8	2.4	1.0-5.0	11	2.3	1.0-4.2	5	2.4	0.8-6.0	6	2.1	0.7-4.8
1984	22	3.4	2.1-5.2	14	4.5	2.4-8.2	8	2.3	1.0-4.8	18	2.8	1.6-4.6	11	3.8	1.8-7.4	7	1.9	0.8-4.2
1985	21	3.8	2.2-5.9	9	4.4	1.7-9.0	12	3.6	1.8-6.4	11	1.8	0.9-3.3	5	1.8	0.5-4.7	6	1.8	0.6-4.1
1986	19	4.1	2.3-6.6	8	4.2	1.4-9.2	11	4.0	1.9-7.4	9	2.1	0.9-4.0	3	1.3	0.3-4.0	6	2.5	0.9-5.4
1987	18	3.5	1.9-5.7	7	2.3	0.9-5.2	11	4.0	1.9-7.3	9	2.2	0.9-4.2	3	1.2	0.2-3.8	6	2.7	0.9-5.7
1988	24	3.9	2.4-5.9	14	4.5	2.4-8.0	10	3.0	1.4-5.8	11	1.7	0.8-3.1	7	2.6	1.0-5.9	4	0.9	0.2-2.6
1989	18	2.6	1.5-4.2	5	1.2	0.4-3.4	13	3.7	1.9-6.5	8	1.2	0.5-2.5	4	1.0	0.3-3.3	4	1.3	0.3-3.3
1990	21	3.3	1.9-5.2	10	4.7	1.8-9.4	11	2.7	1.3-5.1	9	1.5	0.6-2.9	3	1.2	0.2-3.9	6	1.7	0.6-3.8
1991	37	5.4	3.7-7.6	19	5.7	3.2-9.5	18	5.1	2.9-8.2	11	2.0	0.9-3.7	3	1.4	0.2-4.4	8	2.4	1.0-4.9
1992	38	5.8	4.0-8.2	19	6.0	3.4-9.8	19	5.6	3.3-8.8	16	2.6	1.4-4.3	5	2.1	0.6-5.2	11	3.1	1.5-5.7
1993	24	3.2	2.0-4.9	11	2.7	1.3-5.4	13	3.5	1.8-6.0	24	3.7	2.3-5.6	11	3.7	1.8-6.9	13	3.7	1.9-6.4
1994	20	3.0	1.7-4.8	9	2.4	0.9-5.2	11	3.3	1.6-6.0	8	1.1	0.5-2.2	4	1.0	0.3-3.0	4	1.1	0.3-2.9
1995	32	4.9	3.3-7.1	15	4.9	2.6-8.5	17	4.8	2.8-7.8	16	2.6	1.4-4.2	9	3.2	1.3-6.3	7	2.1	0.8-4.3
1996	39	5.3	3.7-7.4	22	6.7	4.0-10.4	17	4.2	2.4-6.9	18	2.9	1.6-4.6	12	4.2	2.0-7.8	6	1.8	0.7-3.9
1997	31	4.0	2.7-5.8	22	6.4	3.6-10.3	9	2.2	1.0-4.3	11	1.5	0.8-2.8	7	1.9	0.8-4.2	4	1.1	0.3-2.9
1998	21	2.4	1.5-3.8	14	3.3	1.7-5.8	7	1.6	0.7-3.4	13	1.8	0.9-3.1	10	3.0	1.4-5.8	3	0.8	0.2-2.2
1999	35	4.3	2.9-6.0	23	5.6	3.4-8.6	12	2.9	1.5-5.1	13	1.7	0.9-2.9	8	2.4	1.0-5.0	5	1.1	0.3-2.6
2000	27	3.1	2.0-4.6	11	2.3	1.1-4.4	16	3.6	2.0-5.9	13	1.7	0.9-2.9	8	2.3	1.0-4.6	5	1.2	0.4-2.8
2001	52	6.9	5.1-9.2	22	5.8	3.5-9.2	30	7.4	4.9-10.6	22	3.4	2.1-5.1	9	3.6	1.5-7.0	13	3.2	1.7-5.5
2002	43	4.8	3.5-6.6	20	5.2	3.1-8.3	23	4.7	3.0-7.1	20	2.4	1.4-3.7	10	2.0	1.0-4.0	10	2.4	1.1-4.5
2003	38	4.5	3.1-6.2	18	4.5	2.6-7.4	20	4.3	2.6-6.8	21	2.5	1.5-3.9	12	3.1	1.5-5.6	9	2.0	0.9-3.9
2004	42	4.6	3.2-6.3	19	4.3	2.5-6.9	23	4.7	2.9-7.1	13	1.6	0.8-2.8	4	1.3	0.3-3.5	9	1.9	0.9-3.7
2005	33	3.9	2.6-5.6	23	6.4	3.9-9.8	10	2.0	1.0-3.8	13	1.5	0.8-2.6	8	1.9	0.8-3.9	5	1.1	0.4-2.6
2006	31	3.2	2.1-4.6	15	3.1	1.6-5.5	16	3.2	1.8-5.2	14	1.6	0.8-2.7	11	2.7	1.2-5.0	3	0.7	0.1-1.9

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Brain and other central nervous system

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	27	3.0	1.9-4.4	13	2.9	1.5-5.2	14	2.9	1.6-4.9	16	2.1	1.2-3.5	7	2.6	1.0-5.3	9	2.0	0.9-3.7
2008	39	4.2	3.0-5.9	16	3.3	1.8-5.6	23	4.7	3.0-7.1	19	2.1	1.2-3.3	9	1.7	0.8-3.4	10	2.1	1.0-3.9
2009	39	4.3	3.0-5.9	22	5.3	3.1-8.3	17	3.5	2.0-5.7	17	2.3	1.3-3.6	8	2.6	1.1-5.2	9	2.0	0.9-3.8
2010	53	5.5	4.1-7.2	31	6.8	4.5-9.8	22	4.3	2.7-6.5	21	2.2	1.3-3.3	12	2.7	1.3-4.9	9	1.7	0.8-3.3
2011	44	4.2	3.0-5.7	23	4.2	2.6-6.4	21	3.9	2.4-6.0	23	2.3	1.4-3.4	15	3.2	1.7-5.5	8	1.5	0.6-3.0
2012	47	4.3	3.2-5.8	25	4.8	3.0-7.2	22	3.8	2.4-5.9	25	2.5	1.6-3.8	13	2.5	1.3-4.4	12	2.4	1.2-4.2
2013	42	3.8	2.7-5.2	20	3.6	2.1-5.7	22	3.8	2.4-5.9	26	2.4	1.6-3.6	15	3.1	1.7-5.2	11	1.9	0.9-3.4
2014	38	3.4	2.4-4.7	21	4.1	2.4-6.4	17	2.8	1.6-4.6	19	1.9	1.1-3.0	12	2.8	1.4-5.0	7	1.2	0.5-2.4
2015	43	3.9	2.8-5.4	25	4.6	2.9-7.0	18	3.3	1.9-5.2	28	2.8	1.8-4.1	11	2.7	1.3-4.9	17	3.0	1.7-4.8
2016	38	3.5	2.5-4.9	22	4.4	2.7-6.8	16	2.9	1.6-4.6	22	1.9	1.2-2.9	11	1.9	0.9-3.5	11	1.9	0.9-3.4

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Thyroid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	16	3.6	1.9-6.1	6	3.4	1.0-8.4	10	3.9	1.8-7.4	2	0.8	0.1-2.6	1	0.5	0.0-3.9	1	0.8	0.0-3.8
1977	12	3.1	1.5-5.8	2	0.9	0.1-4.2	10	4.7	2.1-9.1	2	0.9	0.1-3.0	1	0.5	0.0-3.7	1	1.0	0.0-4.4
1978	11	2.9	1.3-5.4	3	1.3	0.2-4.7	8	4.1	1.6-8.1	4	1.6	0.4-4.0	2	1.7	0.2-6.2	2	1.5	0.2-5.0
1979	11	2.5	1.1-4.9	2	0.5	0.1-3.2	9	4.0	1.6-7.9	3	1.5	0.3-4.0	0	0.0	0.0-2.8	3	2.3	0.4-6.3
1980	11	2.2	1.0-4.3	2	0.6	0.1-3.2	9	3.4	1.4-6.9	1	0.2	0.0-1.4	0	0.0	0.0-2.6	1	0.4	0.0-2.4
1981	7	1.1	0.4-2.4	2	0.7	0.1-3.5	5	1.4	0.4-3.6	2	0.5	0.1-1.7	1	0.5	0.0-3.2	1	0.4	0.0-2.3
1982	11	2.1	1.0-3.9	1	0.4	0.0-3.1	10	3.5	1.6-6.6	2	0.4	0.1-1.6	1	0.4	0.0-3.0	1	0.4	0.0-2.2
1983	16	2.9	1.6-4.9	2	0.8	0.1-3.5	14	4.7	2.5-8.1	0	0.0	0.0-0.9	0	0.0	0.0-2.3	0	0.0	0.0-1.5
1984	10	1.6	0.8-3.1	2	0.8	0.1-3.5	8	2.3	1.0-4.8	0	0.0	0.0-0.8	0	0.0	0.0-2.2	0	0.0	0.0-1.4
1985	20	3.9	2.3-6.1	3	1.2	0.2-3.9	17	6.1	3.5-9.9	1	0.2	0.0-1.2	1	0.5	0.0-2.9	0	0.0	0.0-1.4
1986	10	1.9	0.9-3.6	3	1.3	0.3-4.1	7	2.5	1.0-5.2	3	0.7	0.1-1.9	1	0.5	0.0-2.9	2	0.8	0.1-2.7
1987	12	2.2	1.1-3.9	3	1.4	0.3-4.4	9	2.9	1.3-5.6	4	0.8	0.2-2.0	1	0.4	0.0-2.6	3	1.1	0.2-3.1
1988	8	1.4	0.6-2.7	1	0.4	0.0-2.6	7	2.2	0.9-4.6	3	0.6	0.1-1.8	2	1.1	0.1-3.9	1	0.3	0.0-1.8
1989	11	1.8	0.9-3.3	1	0.2	0.0-2.2	10	3.1	1.5-5.8	3	0.8	0.1-2.2	0	0.0	0.0-1.9	3	1.3	0.3-3.5
1990	23	3.7	2.3-5.7	4	1.2	0.3-3.6	19	5.8	3.4-9.1	2	0.6	0.1-2.0	0	0.0	0.0-1.9	2	1.0	0.1-3.1
1991	26	4.5	2.9-6.8	4	2.3	0.6-5.8	22	6.6	4.1-10.1	3	0.7	0.1-1.9	1	0.8	0.0-3.8	2	0.7	0.1-2.3
1992	20	2.8	1.7-4.5	6	2.4	0.7-5.9	14	3.5	1.9-6.0	2	0.3	0.0-1.2	1	0.3	0.0-2.3	1	0.3	0.0-1.7
1993	26	3.7	2.3-5.6	7	1.8	0.7-4.2	19	5.1	3.0-8.2	3	0.6	0.1-1.8	2	0.9	0.1-3.2	1	0.4	0.0-2.0
1994	29	4.3	2.8-6.3	2	0.6	0.1-2.6	27	7.3	4.8-10.7	1	0.1	0.0-0.8	0	0.0	0.0-1.7	1	0.3	0.0-1.5
1995	32	4.8	3.2-6.9	4	1.5	0.4-4.0	28	7.5	4.9-10.9	3	0.6	0.1-1.7	1	0.3	0.0-2.0	2	0.7	0.1-2.3
1996	30	4.1	2.7-5.9	6	2.0	0.7-4.5	24	6.0	3.8-9.0	3	0.5	0.1-1.4	2	0.7	0.1-2.8	1	0.3	0.0-1.5
1997	34	4.6	3.1-6.5	8	2.3	0.9-4.8	26	6.5	4.2-9.6	5	0.7	0.2-1.7	4	1.3	0.3-3.5	1	0.3	0.0-1.4
1998	29	3.8	2.5-5.5	6	1.9	0.7-4.2	23	5.5	3.5-8.3	2	0.4	0.0-1.2	1	0.6	0.0-2.7	1	0.2	0.0-1.4
1999	27	3.5	2.2-5.1	3	0.9	0.2-2.8	24	5.6	3.6-8.4	4	0.6	0.2-1.6	0	0.0	0.0-1.4	4	1.1	0.3-2.8
2000	51	6.4	4.7-8.5	10	2.5	1.2-4.8	41	9.6	6.8-13.1	2	0.4	0.0-1.2	0	0.0	0.0-1.3	2	0.6	0.1-1.9
2001	50	5.9	4.3-7.9	11	3.1	1.4-6.0	39	8.6	6.1-11.8	2	0.3	0.0-1.1	0	0.0	0.0-1.3	2	0.6	0.1-1.9
2002	48	5.6	4.1-7.5	10	2.7	1.2-5.3	38	8.2	5.8-11.3	3	0.5	0.1-1.3	1	0.3	0.0-1.7	2	0.5	0.1-1.9
2003	54	6.2	4.6-8.2	15	4.1	2.2-6.9	39	8.3	5.9-11.4	4	0.6	0.2-1.6	1	0.2	0.0-1.6	3	0.8	0.2-2.3
2004	61	7.3	5.5-9.5	8	3.1	1.2-6.3	53	11.3	8.5-14.9	6	0.9	0.3-2.0	2	0.5	0.1-2.0	4	1.1	0.3-2.7
2005	70	7.5	5.8-9.5	12	2.8	1.4-4.9	58	11.8	8.9-15.3	5	0.7	0.2-1.6	4	1.3	0.3-3.4	1	0.3	0.0-1.4
2006	67	7.3	5.6-9.3	16	4.6	2.5-7.6	51	10.2	7.6-13.4	2	0.2	0.0-0.9	1	0.3	0.0-1.6	1	0.2	0.0-1.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Thyroid

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	81	9.0	7.1-11.2	18	4.6	2.6-7.4	63	12.9	9.9-16.6	3	0.5	0.1-1.3	2	0.7	0.1-2.5	1	0.3	0.0-1.4
2008	78	8.5	6.7-10.7	18	4.5	2.6-7.2	60	12.0	9.1-15.5	4	0.4	0.1-1.2	2	0.4	0.0-1.5	2	0.4	0.0-1.5
2009	94	10.0	8.0-12.3	22	5.1	3.2-7.9	72	14.1	11.0-17.8	5	0.7	0.2-1.5	1	0.3	0.0-1.5	4	0.9	0.2-2.2
2010	95	9.3	7.5-11.4	18	3.7	2.2-5.9	77	14.2	11.2-17.8	7	0.8	0.3-1.6	4	1.0	0.2-2.7	3	0.6	0.1-1.8
2011	114	11.2	9.2-13.5	26	5.9	3.8-8.8	88	16.0	12.8-19.7	4	0.5	0.1-1.3	0	0.0	0.0-0.9	4	0.8	0.2-2.1
2012	111	10.8	8.8-13.0	25	5.1	3.2-7.7	86	15.8	12.6-19.5	9	1.0	0.4-1.9	3	0.8	0.1-2.4	6	1.1	0.4-2.4
2013	117	11.3	9.3-13.6	17	3.5	2.0-5.7	100	17.8	14.4-21.7	7	0.9	0.3-1.8	0	0.0	0.0-0.9	7	1.4	0.6-2.9
2014	123	11.5	9.5-13.7	19	3.7	2.2-5.9	104	18.4	15.0-22.3	3	0.4	0.1-1.0	1	0.4	0.0-1.7	2	0.4	0.0-1.4
2015	145	13.3	11.1-15.7	34	7.2	4.9-10.2	111	19.1	15.7-23.1	6	0.7	0.2-1.4	3	0.8	0.1-2.3	3	0.6	0.1-1.7
2016	110	10.0	8.1-12.0	23	4.7	2.9-7.1	87	14.8	11.8-18.3	4	0.4	0.1-1.0	0	0.0	0.0-0.8	4	0.6	0.2-1.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Hodgkin lymphoma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	13	3.2	1.6-5.7	5	1.8	0.5-5.4	8	4.1	1.7-8.2	3	0.7	0.2-2.3	1	0.6	0.0-4.1	2	0.8	0.1-3.3
1977	8	1.4	0.6-3.1	4	1.4	0.3-4.7	4	1.5	0.4-4.0	4	0.9	0.2-2.4	3	1.6	0.3-5.2	1	0.2	0.0-2.2
1978	12	2.1	1.1-4.0	10	3.9	1.8-7.9	2	0.6	0.1-2.8	2	0.4	0.1-1.7	2	0.9	0.1-4.1	0	0.0	0.0-1.8
1979	8	0.9	0.4-2.1	5	1.1	0.3-3.8	3	0.7	0.1-2.6	4	1.0	0.2-2.9	3	0.8	0.1-3.8	1	0.9	0.0-4.0
1980	14	1.8	1.0-3.3	10	2.9	1.3-6.2	4	0.9	0.2-2.8	4	0.7	0.2-2.1	2	1.0	0.1-4.0	2	0.6	0.1-2.5
1981	13	1.9	0.9-3.6	8	2.0	0.8-5.0	5	1.6	0.4-4.1	5	1.1	0.3-2.7	4	1.4	0.3-4.5	1	0.6	0.0-3.0
1982	18	3.4	1.9-5.6	7	3.2	1.3-7.0	11	3.5	1.6-6.7	3	0.7	0.1-2.0	2	1.0	0.1-4.0	1	0.4	0.0-2.3
1983	12	1.7	0.9-3.2	7	2.0	0.8-4.9	5	1.4	0.4-3.5	2	0.2	0.0-1.1	2	0.4	0.1-2.7	0	0.0	0.0-1.5
1984	15	2.7	1.5-4.7	11	5.1	2.4-9.7	4	1.0	0.3-2.8	3	0.6	0.1-1.9	3	1.4	0.2-4.8	0	0.0	0.0-1.4
1985	17	2.7	1.5-4.4	9	3.2	1.4-6.6	8	2.2	0.9-4.6	6	0.9	0.3-2.2	2	1.0	0.1-3.8	4	1.0	0.3-2.9
1986	22	3.0	1.8-4.7	16	4.5	2.5-7.9	6	1.5	0.5-3.5	8	1.2	0.5-2.5	7	2.2	0.8-5.2	1	0.3	0.0-1.9
1987	24	3.2	2.0-4.9	9	2.9	1.2-5.9	15	3.7	2.0-6.3	4	0.6	0.2-1.8	3	1.2	0.3-3.9	1	0.2	0.0-1.6
1988	14	1.5	0.8-2.7	11	2.3	1.1-4.9	3	0.6	0.1-2.1	7	1.1	0.4-2.4	5	2.0	0.6-5.0	2	0.4	0.1-1.9
1989	20	2.5	1.4-4.1	10	2.0	0.9-4.4	10	2.7	1.2-5.2	3	0.7	0.1-2.1	1	0.4	0.0-2.5	2	0.9	0.1-3.0
1990	24	3.3	2.0-5.2	13	3.0	1.6-5.7	11	3.3	1.5-6.0	5	0.7	0.2-1.7	2	0.5	0.1-2.5	3	0.8	0.2-2.5
1991	25	2.9	1.8-4.5	13	3.0	1.5-5.6	12	2.8	1.4-5.1	7	0.8	0.3-1.8	3	0.7	0.1-2.8	4	0.9	0.2-2.5
1992	13	1.6	0.8-2.9	11	3.0	1.5-5.8	2	0.4	0.0-1.6	4	0.5	0.1-1.3	3	0.8	0.1-2.8	1	0.2	0.0-1.3
1993	25	2.8	1.8-4.2	16	3.4	1.9-6.1	9	2.1	0.9-4.0	3	0.4	0.1-1.3	2	0.5	0.1-2.5	1	0.3	0.0-1.6
1994	13	1.6	0.8-2.9	6	1.5	0.5-3.7	7	1.7	0.6-3.6	5	0.7	0.2-1.7	4	1.2	0.3-3.4	1	0.3	0.0-1.5
1995	26	2.8	1.8-4.3	18	3.8	2.2-6.4	8	1.9	0.8-3.7	8	0.9	0.4-1.8	8	1.8	0.7-3.9	0	0.0	0.0-1.0
1996	24	2.8	1.8-4.3	11	2.7	1.3-5.2	13	3.0	1.6-5.1	1	0.2	0.0-0.8	1	0.4	0.0-2.1	0	0.0	0.0-1.0
1997	18	1.9	1.1-3.0	14	2.9	1.6-5.2	4	0.8	0.2-2.2	4	0.4	0.1-1.2	2	0.4	0.0-2.0	2	0.5	0.1-1.8
1998	26	2.9	1.8-4.3	16	3.9	2.2-6.7	10	2.0	1.0-3.8	2	0.2	0.0-0.8	0	0.0	0.0-1.4	2	0.4	0.0-1.6
1999	30	3.3	2.2-4.7	16	3.5	2.0-5.9	14	3.0	1.6-5.1	4	0.6	0.2-1.5	2	0.6	0.1-2.2	2	0.6	0.1-2.0
2000	33	3.4	2.3-4.8	20	4.2	2.5-6.7	13	2.6	1.4-4.5	0	0.0	0.0-0.5	0	0.0	0.0-1.3	0	0.0	0.0-0.9
2001	32	3.4	2.3-4.9	20	4.2	2.5-6.7	12	2.6	1.3-4.5	4	0.5	0.1-1.3	2	0.6	0.1-2.4	2	0.4	0.0-1.6
2002	26	2.7	1.7-4.0	17	3.3	1.9-5.5	9	2.0	0.9-3.8	3	0.3	0.1-1.0	3	0.7	0.1-2.2	0	0.0	0.0-0.8
2003	23	2.2	1.4-3.3	10	1.8	0.9-3.6	13	2.5	1.3-4.3	1	0.1	0.0-0.6	1	0.2	0.0-1.5	0	0.0	0.0-0.8
2004	29	2.7	1.8-4.0	13	2.5	1.3-4.5	16	3.0	1.7-4.9	1	0.1	0.0-0.6	1	0.2	0.0-1.5	0	0.0	0.0-0.8
2005	25	2.5	1.6-3.8	15	3.4	1.8-5.8	10	1.9	0.9-3.6	6	0.6	0.2-1.3	5	1.1	0.3-2.7	1	0.2	0.0-1.1
2006	33	3.7	2.5-5.2	19	4.3	2.5-7.0	14	3.0	1.6-5.0	4	0.4	0.1-1.1	4	0.9	0.2-2.5	0	0.0	0.0-0.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Hodgkin lymphoma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	32	3.0	2.0-4.3	21	4.4	2.6-6.9	11	1.8	0.9-3.3	5	0.6	0.2-1.4	2	0.7	0.1-2.4	3	0.6	0.1-1.7
2008	29	2.8	1.9-4.1	18	3.9	2.3-6.2	11	1.9	1.0-3.5	3	0.3	0.1-0.9	2	0.4	0.1-1.7	1	0.2	0.0-1.1
2009	33	3.1	2.1-4.4	20	4.0	2.4-6.3	13	2.3	1.2-4.0	3	0.3	0.1-0.9	2	0.4	0.1-1.7	1	0.1	0.0-1.0
2010	38	3.5	2.4-4.8	21	4.0	2.5-6.3	17	3.0	1.7-4.9	1	0.1	0.0-0.5	1	0.2	0.0-1.2	0	0.0	0.0-0.7
2011	35	2.9	2.0-4.1	21	3.5	2.2-5.5	14	2.3	1.3-3.9	7	0.6	0.2-1.3	4	0.7	0.2-1.9	3	0.5	0.1-1.5
2012	38	3.5	2.5-4.9	17	3.5	2.0-5.8	21	3.6	2.2-5.5	1	0.1	0.0-0.5	1	0.2	0.0-1.1	0	0.0	0.0-0.7
2013	34	3.1	2.1-4.3	19	3.5	2.1-5.5	15	2.7	1.5-4.4	2	0.2	0.0-0.7	1	0.2	0.0-1.2	1	0.2	0.0-1.0
2014	30	2.6	1.8-3.8	15	2.8	1.5-4.7	15	2.5	1.4-4.2	1	0.1	0.0-0.5	1	0.2	0.0-1.1	0	0.0	0.0-0.7
2015	45	4.0	2.9-5.4	23	4.1	2.6-6.2	22	3.9	2.4-6.0	5	0.5	0.2-1.2	4	0.9	0.2-2.4	1	0.2	0.0-1.1
2016	31	2.8	1.9-4.1	12	2.6	1.3-4.7	19	3.2	1.9-5.1	1	0.1	0.0-0.5	1	0.1	0.0-1.0	0	0.0	0.0-0.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Non-Hodgkin lymphomas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	29	7.7	4.9-11.5	17	9.6	5.3-16.1	12	6.0	2.8-10.9	12	4.7	2.2-8.4	6	5.4	1.6-12.5	6	4.0	1.3-8.8
1977	46	12.2	8.7-16.7	28	17.7	11.1-26.7	18	8.3	4.7-13.4	24	6.9	4.2-10.7	14	10.8	5.0-19.6	10	4.6	2.1-8.6
1978	47	12.1	8.6-16.6	24	12.0	7.5-18.4	23	11.6	7.0-17.9	13	3.3	1.6-5.9	8	4.0	1.7-8.5	5	2.6	0.7-6.3
1979	30	6.5	4.2-9.5	16	6.3	3.5-10.9	14	6.3	3.3-10.7	29	7.7	4.8-11.5	18	8.3	4.8-13.7	11	6.4	3.0-11.7
1980	26	6.9	4.2-10.5	16	11.0	5.3-19.5	10	4.5	2.1-8.5	18	5.1	2.8-8.2	8	5.9	2.0-12.6	10	4.8	2.2-9.0
1981	32	8.0	5.2-11.6	20	9.6	5.5-15.7	12	6.3	3.1-11.1	18	4.1	2.3-6.7	8	4.7	1.7-10.0	10	3.9	1.8-7.2
1982	34	7.4	5.0-10.5	24	12.3	7.5-19.0	10	3.7	1.7-6.9	8	1.6	0.7-3.3	6	3.0	1.1-6.9	2	0.6	0.1-2.5
1983	41	8.5	6.0-11.8	23	9.2	5.7-14.3	18	7.4	4.3-11.9	18	3.8	2.2-6.1	10	4.0	1.9-7.8	8	3.4	1.4-6.7
1984	41	8.6	6.0-12.0	24	9.7	6.0-15.0	17	7.3	4.1-11.7	35	7.9	5.4-11.2	20	9.1	5.4-14.4	15	6.5	3.5-10.9
1985	45	9.1	6.5-12.3	27	12.9	7.9-19.8	18	6.4	3.7-10.2	25	5.4	3.5-8.1	14	6.6	3.5-11.4	11	4.5	2.2-8.0
1986	50	9.8	7.0-13.1	24	9.3	5.5-14.6	26	9.7	6.2-14.4	20	4.1	2.4-6.4	12	5.3	2.6-9.8	8	2.9	1.2-5.9
1987	50	9.2	6.7-12.2	29	11.9	7.6-17.8	21	7.0	4.3-10.8	18	4.1	2.3-6.5	12	5.9	2.8-10.8	6	2.6	0.9-5.5
1988	57	10.9	8.0-14.3	35	15.8	10.2-23.1	22	7.5	4.6-11.5	32	7.2	4.7-10.4	21	13.1	7.3-21.0	11	3.8	1.8-7.0
1989	69	11.0	8.5-14.1	37	12.3	8.3-17.5	32	10.0	6.8-14.1	28	5.1	3.3-7.6	16	6.2	3.3-10.7	12	4.2	2.1-7.3
1990	78	13.9	10.8-17.6	45	15.8	11.1-21.7	33	11.5	7.8-16.3	31	5.7	3.8-8.1	16	6.7	3.6-11.3	15	4.9	2.7-8.2
1991	74	11.9	9.1-15.3	45	14.8	10.0-21.1	29	9.4	6.2-13.7	34	6.2	4.2-8.9	19	6.8	4.0-11.0	15	5.3	2.9-8.7
1992	76	12.4	9.6-15.8	54	19.0	13.7-25.5	22	7.0	4.3-10.7	29	5.6	3.6-8.1	19	8.4	4.7-13.5	10	3.5	1.6-6.4
1993	94	14.4	11.5-17.8	57	19.2	14.2-25.5	37	10.4	7.3-14.5	39	7.0	4.9-9.7	16	5.2	2.8-8.8	23	7.8	4.9-11.7
1994	84	12.8	10.1-16.0	46	14.3	10.2-19.6	38	11.1	7.8-15.3	28	5.4	3.5-7.8	19	8.3	4.6-13.6	9	3.2	1.4-6.0
1995	101	14.8	11.9-18.2	71	23.8	17.7-31.1	30	8.1	5.4-11.7	33	5.9	4.0-8.4	16	4.9	2.8-8.2	17	5.8	3.3-9.2
1996	116	17.0	13.9-20.6	70	20.1	15.4-25.9	46	13.3	9.6-17.8	32	5.5	3.7-7.9	19	7.9	4.3-13.0	13	4.0	2.1-6.8
1997	102	14.7	11.8-18.0	61	20.3	14.9-26.9	41	10.8	7.7-14.7	21	3.7	2.2-5.8	11	5.3	2.4-9.8	10	3.0	1.4-5.4
1998	118	16.3	13.4-19.6	73	22.5	17.1-29.0	45	11.6	8.4-15.5	34	5.4	3.7-7.6	20	7.6	4.3-12.2	14	4.0	2.2-6.7
1999	129	17.8	14.7-21.3	75	22.4	17.2-28.7	54	13.8	10.3-18.1	32	5.0	3.4-7.2	15	4.7	2.5-8.0	17	4.9	2.8-7.9
2000	102	13.1	10.6-16.1	56	16.3	12.0-21.7	46	10.9	7.9-14.6	42	6.1	4.4-8.4	31	10.6	6.8-15.5	11	2.9	1.5-5.2
2001	124	16.1	13.2-19.3	74	20.1	15.5-25.6	50	12.2	9.0-16.1	42	6.6	4.7-9.0	21	8.5	5.0-13.2	21	5.5	3.4-8.4
2002	129	16.3	13.5-19.5	74	20.2	15.6-25.8	55	13.0	9.7-16.9	48	6.6	4.8-8.9	26	7.7	4.9-11.6	22	5.5	3.4-8.4
2003	122	15.7	12.9-18.8	56	16.3	12.0-21.5	66	15.4	11.9-19.7	27	3.8	2.4-5.5	14	4.1	2.1-7.0	13	3.3	1.7-5.6
2004	108	13.9	11.3-16.9	60	18.6	13.6-24.6	48	11.0	8.1-14.6	36	5.2	3.6-7.3	19	5.8	3.3-9.4	17	4.5	2.6-7.1
2005	118	14.7	12.1-17.7	68	18.8	14.2-24.3	50	11.6	8.5-15.3	32	4.7	3.2-6.7	18	6.6	3.7-10.6	14	3.5	1.9-5.8
2006	140	16.6	13.9-19.7	75	19.7	15.2-25.1	65	14.1	10.8-18.0	34	4.2	2.9-5.9	19	4.8	2.8-7.8	15	3.5	2.0-5.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Non-Hodgkin lymphomas

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	128	14.8	12.2-17.6	76	20.9	16.1-26.6	52	10.7	7.9-14.0	31	3.9	2.6-5.6	19	6.0	3.4-9.7	12	2.7	1.4-4.7
2008	159	17.7	15.0-20.8	94	23.9	18.9-29.6	65	13.2	10.1-16.8	32	3.8	2.6-5.5	21	6.3	3.7-9.8	11	2.2	1.1-4.0
2009	134	14.8	12.3-17.6	68	17.1	13.1-22.1	66	13.2	10.2-16.8	34	3.8	2.6-5.4	23	6.1	3.7-9.4	11	2.2	1.1-4.0
2010	141	15.2	12.7-18.0	76	17.5	13.5-22.2	65	13.0	9.9-16.6	42	5.0	3.6-6.8	22	5.9	3.6-9.2	20	4.4	2.7-6.8
2011	161	17.1	14.4-20.0	101	24.8	19.8-30.6	60	11.5	8.7-14.9	42	5.0	3.6-6.8	27	7.6	4.8-11.3	15	3.2	1.7-5.2
2012	140	14.2	11.9-16.8	74	16.8	13.0-21.4	66	12.0	9.3-15.4	33	3.7	2.5-5.3	16	4.8	2.6-7.9	17	3.2	1.8-5.1
2013	140	14.4	12.0-17.0	73	16.9	13.0-21.5	67	12.4	9.6-15.8	28	3.2	2.1-4.7	14	3.3	1.7-5.6	14	2.9	1.5-4.8
2014	149	14.6	12.3-17.2	86	19.1	15.0-23.8	63	11.1	8.5-14.3	43	4.9	3.5-6.6	25	7.2	4.6-10.8	18	3.4	2.0-5.4
2015	175	16.5	14.1-19.2	86	18.6	14.7-23.3	89	15.4	12.3-19.0	39	3.9	2.8-5.4	21	4.9	2.9-7.6	18	3.1	1.8-5.0
2016	167	15.8	13.4-18.4	91	19.6	15.5-24.3	76	12.9	10.1-16.2	31	3.2	2.2-4.6	19	5.0	2.9-7.9	12	2.1	1.0-3.6

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Myeloma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	31	9.2	6.1-13.2	13	8.6	4.3-15.3	18	9.7	5.6-15.5	22	6.4	3.9-9.9	11	7.1	3.3-13.3	11	5.8	2.8-10.6
1977	30	10.3	6.7-15.1	13	10.1	4.6-18.6	17	10.5	5.9-17.1	19	7.5	4.2-11.9	12	11.8	5.3-21.6	7	4.7	1.7-9.7
1978	28	8.4	5.4-12.4	12	10.1	4.5-18.7	16	7.9	4.4-12.9	16	5.4	2.9-9.1	9	8.8	3.4-17.4	7	3.6	1.4-7.6
1979	35	11.0	7.4-15.6	15	14.2	7.1-24.5	20	9.8	5.9-15.1	16	4.8	2.6-8.0	8	6.2	2.1-13.2	8	4.1	1.7-8.1
1980	33	9.5	6.4-13.6	13	8.3	4.3-14.6	20	10.1	6.0-15.8	22	6.4	3.9-9.8	9	5.6	2.4-10.9	13	6.8	3.5-11.7
1981	40	12.6	8.7-17.4	15	12.2	6.2-21.0	25	13.0	8.2-19.3	26	8.0	5.1-11.8	13	10.9	5.4-19.0	13	6.3	3.3-10.8
1982	47	13.1	9.4-17.7	26	17.1	10.6-25.8	21	10.1	6.1-15.6	28	7.1	4.6-10.4	16	9.9	5.4-16.4	12	5.2	2.6-9.1
1983	26	6.8	4.3-10.1	16	10.5	5.6-17.7	10	4.4	2.1-8.1	26	7.7	4.9-11.3	13	10.5	5.0-18.4	13	6.3	3.3-10.7
1984	49	12.5	9.1-16.7	27	14.9	9.3-22.4	22	10.4	6.4-15.8	27	7.6	4.8-11.2	13	9.9	4.8-17.4	14	6.4	3.4-10.7
1985	38	9.4	6.5-13.0	21	11.7	6.8-18.4	17	7.6	4.3-12.1	29	7.4	4.8-10.6	16	8.6	4.8-14.1	13	6.1	3.2-10.4
1986	52	14.1	10.3-18.7	25	16.4	10.0-24.9	27	12.5	8.1-18.3	35	9.9	6.7-13.9	22	14.7	8.7-22.7	13	6.6	3.5-11.2
1987	39	9.8	6.8-13.5	19	14.1	7.8-22.6	20	8.0	4.8-12.3	22	6.1	3.7-9.3	13	9.9	4.8-17.4	9	4.1	1.8-7.6
1988	45	10.6	7.6-14.3	20	11.5	6.6-18.2	25	9.9	6.4-14.7	20	4.7	2.8-7.3	10	5.4	2.5-10.0	10	4.1	1.9-7.5
1989	57	13.6	10.1-17.7	24	13.5	8.1-20.7	33	13.5	9.2-19.0	29	7.4	4.9-10.8	11	7.4	3.3-13.6	18	7.5	4.3-11.8
1990	54	12.3	9.1-16.1	29	16.8	10.8-24.7	25	9.5	6.1-14.0	33	7.6	5.1-10.8	16	9.0	4.7-15.2	17	6.8	3.9-10.8
1991	59	12.5	9.4-16.3	26	14.0	8.6-21.1	33	11.8	8.1-16.6	26	5.7	3.6-8.4	12	6.8	3.3-11.9	14	5.0	2.7-8.4
1992	58	12.0	9.0-15.6	27	13.8	8.6-20.7	31	10.9	7.4-15.5	50	11.2	8.2-14.9	23	14.2	8.4-22.0	27	9.9	6.5-14.4
1993	57	11.3	8.5-14.7	26	14.5	8.9-21.9	31	10.1	6.8-14.4	39	7.8	5.5-10.8	15	8.2	4.5-13.6	24	7.9	5.0-11.7
1994	60	12.7	9.6-16.5	27	13.1	8.1-19.5	33	11.8	8.1-16.6	41	8.7	6.1-11.9	16	8.4	4.5-13.9	25	8.6	5.5-12.7
1995	52	9.8	7.2-13.0	28	14.2	8.8-21.2	24	7.5	4.8-11.1	30	6.0	4.0-8.6	14	7.2	3.5-12.6	16	5.4	3.1-8.7
1996	61	10.8	8.2-14.0	30	12.6	8.0-18.7	31	9.6	6.5-13.7	28	5.2	3.4-7.6	17	8.2	4.3-13.8	11	3.5	1.7-6.2
1997	72	13.0	10.1-16.4	36	15.6	10.6-22.0	36	11.1	7.7-15.4	47	8.8	6.4-11.8	31	15.1	9.7-21.9	16	5.1	2.9-8.2
1998	67	11.6	8.9-14.8	31	13.2	8.6-19.2	36	10.6	7.4-14.6	36	6.7	4.7-9.3	15	7.4	3.8-12.6	21	6.6	4.1-10.0
1999	67	11.4	8.7-14.5	35	13.6	9.2-19.3	32	9.5	6.5-13.4	37	7.0	4.9-9.6	17	8.0	4.5-12.9	20	6.2	3.8-9.5
2000	86	13.7	10.9-17.0	41	15.5	10.6-21.6	45	12.6	9.2-16.9	45	7.7	5.6-10.4	25	10.3	6.4-15.5	20	5.9	3.6-9.1
2001	87	13.8	11.0-17.2	41	14.4	10.0-20.0	46	13.0	9.5-17.4	49	8.3	6.0-10.9	18	7.4	4.1-12.2	31	8.9	6.0-12.5
2002	72	11.4	8.8-14.4	30	12.0	7.7-17.5	42	11.2	8.0-15.1	40	6.1	4.3-8.3	16	5.4	2.9-9.1	24	6.5	4.1-9.7
2003	86	12.2	9.7-15.2	41	13.8	9.7-19.1	45	11.1	8.1-14.9	35	5.2	3.6-7.2	17	6.1	3.4-9.9	18	4.6	2.7-7.3
2004	93	14.0	11.2-17.2	41	15.8	10.9-22.0	52	13.3	9.9-17.5	42	6.9	4.9-9.4	24	10.6	6.6-16.0	18	4.8	2.8-7.6
2005	112	15.9	13.0-19.2	50	16.6	12.0-22.2	62	15.3	11.7-19.6	38	5.5	3.8-7.6	16	4.9	2.7-8.1	22	5.6	3.5-8.5
2006	104	13.5	11.0-16.5	53	16.9	12.4-22.4	51	11.5	8.5-15.1	45	6.4	4.7-8.7	26	8.6	5.5-12.8	19	4.7	2.8-7.3

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Myeloma

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	90	11.3	9.0-13.9	41	12.6	8.8-17.3	49	10.7	7.9-14.2	44	6.4	4.6-8.6	22	7.7	4.7-11.8	22	5.4	3.3-8.1
2008	120	14.1	11.6-16.9	54	14.2	10.5-18.8	66	13.9	10.7-17.7	42	5.7	4.0-7.7	22	7.2	4.4-11.1	20	4.6	2.8-7.2
2009	124	14.9	12.3-17.8	65	17.7	13.4-22.8	59	12.5	9.5-16.1	36	4.5	3.1-6.2	19	5.1	3.0-8.1	17	3.7	2.2-6.0
2010	128	15.5	12.8-18.5	60	18.6	13.9-24.3	68	13.7	10.6-17.5	50	6.5	4.8-8.6	24	7.7	4.8-11.6	26	5.5	3.6-8.1
2011	114	13.2	10.8-15.9	43	11.1	7.9-15.1	71	14.4	11.2-18.2	43	5.2	3.7-7.0	15	4.7	2.5-7.8	28	5.7	3.8-8.2
2012	139	14.8	12.3-17.5	77	19.3	15.0-24.5	62	11.7	8.9-15.0	49	5.7	4.1-7.5	25	6.3	4.0-9.5	24	4.9	3.1-7.4
2013	161	17.8	15.1-20.9	86	22.0	17.3-27.4	75	14.8	11.6-18.6	48	5.7	4.2-7.6	23	7.2	4.4-10.9	25	4.9	3.2-7.3
2014	140	14.2	11.9-16.8	82	19.3	15.1-24.3	58	10.6	8.0-13.8	48	5.5	4.0-7.4	19	5.6	3.3-8.9	29	5.7	3.8-8.2
2015	145	14.4	12.1-17.0	72	17.0	13.0-21.6	73	12.7	9.9-16.0	54	5.6	4.1-7.3	28	6.5	4.2-9.5	26	4.7	3.1-6.9
2016	151	14.2	12.0-16.8	82	18.8	14.7-23.6	69	11.3	8.7-14.3	34	3.5	2.4-4.9	16	4.2	2.3-6.8	18	3.2	1.9-5.1

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Leukemias

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
1976	31	7.1	4.5-10.6	19	10.8	5.3-19.0	12	4.6	2.3-8.5	18	4.8	2.6-8.0	12	6.8	2.9-13.3	6	3.1	1.0-7.1
1977	35	10.7	7.0-15.4	21	17.2	9.1-28.5	14	6.8	3.6-11.6	20	5.1	2.8-8.4	14	8.1	3.4-15.8	6	3.0	0.9-6.9
1978	39	10.0	6.8-14.2	16	7.8	4.1-13.4	23	11.4	6.9-17.4	32	9.7	6.2-14.2	15	10.1	4.8-18.2	17	9.2	5.0-15.2
1979	36	9.0	6.0-12.8	15	6.9	3.7-11.9	21	10.2	6.0-15.8	22	5.3	3.3-8.2	12	7.1	3.5-12.8	10	4.1	1.9-7.7
1980	43	10.0	6.9-13.8	21	11.6	6.4-19.1	22	9.0	5.5-13.9	31	6.9	4.5-10.1	13	6.7	3.1-12.5	18	7.2	4.2-11.6
1981	40	10.3	7.1-14.4	22	13.9	7.7-22.5	18	8.1	4.7-13.0	27	7.5	4.7-11.3	13	10.7	4.8-19.7	14	6.1	3.2-10.4
1982	32	6.7	4.4-9.8	15	7.4	3.8-12.9	17	6.3	3.6-10.4	18	3.5	2.0-5.7	8	3.3	1.2-7.3	10	3.6	1.7-6.9
1983	43	9.7	6.8-13.5	24	13.5	7.7-21.6	19	7.6	4.4-12.0	26	6.4	4.0-9.5	12	9.1	4.1-16.7	14	5.3	2.8-9.0
1984	33	6.6	4.3-9.7	16	7.2	3.6-12.6	17	6.2	3.4-10.3	22	5.3	3.1-8.3	13	6.9	3.2-12.9	9	4.2	1.8-7.9
1985	58	11.7	8.6-15.4	29	13.6	8.5-20.5	29	10.4	6.8-15.3	22	4.6	2.7-7.1	12	6.0	2.6-11.4	10	3.7	1.7-6.9
1986	42	8.4	5.9-11.7	28	13.0	7.9-19.9	14	5.2	2.7-8.9	35	7.0	4.7-10.0	21	7.6	4.5-12.3	14	5.9	3.1-10.0
1987	50	10.0	7.2-13.4	27	12.2	7.5-18.7	23	8.3	5.1-12.6	31	7.0	4.6-10.2	15	8.5	4.1-15.1	16	6.4	3.6-10.4
1988	45	9.4	6.7-12.7	19	11.0	6.2-17.6	26	8.9	5.8-13.2	40	8.6	6.0-12.0	20	11.3	6.2-18.4	20	7.2	4.3-11.3
1989	48	10.0	7.2-13.5	26	12.8	7.7-19.8	22	8.0	4.9-12.3	18	4.0	2.3-6.4	14	8.3	4.2-14.4	4	1.4	0.3-3.6
1990	64	12.4	9.3-16.0	33	14.4	9.2-21.3	31	10.9	7.3-15.5	33	7.2	4.8-10.2	14	8.1	3.9-14.3	19	6.9	4.1-10.8
1991	58	10.6	7.8-13.9	28	12.2	7.4-18.7	30	9.7	6.4-14.0	35	6.8	4.6-9.6	23	10.7	6.1-17.0	12	4.3	2.2-7.4
1992	57	9.6	7.1-12.6	30	11.6	7.2-17.6	27	8.3	5.4-12.2	21	4.2	2.5-6.5	12	5.5	2.4-10.2	9	3.3	1.5-6.2
1993	54	9.8	7.2-13.0	29	13.3	8.2-19.9	25	7.6	4.8-11.4	27	4.6	2.9-6.9	16	7.0	3.5-12.2	11	3.2	1.6-5.8
1994	56	9.7	7.1-12.8	30	13.1	8.3-19.4	26	7.5	4.8-11.2	43	8.6	6.1-11.7	23	11.5	6.8-17.8	20	6.6	4.0-10.2
1995	67	10.8	8.2-13.9	41	16.2	10.9-22.9	26	7.3	4.7-10.8	42	7.0	4.9-9.5	23	10.1	6.0-15.7	19	5.2	3.1-8.2
1996	73	11.2	8.6-14.3	44	15.8	10.9-21.9	29	7.7	5.1-11.2	42	7.1	5.0-9.7	21	8.4	4.9-13.3	21	6.1	3.7-9.3
1997	64	9.9	7.5-12.8	36	11.6	7.8-16.5	28	8.2	5.4-11.8	39	6.8	4.7-9.4	23	9.5	5.7-14.7	16	4.8	2.7-7.8
1998	73	11.3	8.7-14.3	36	13.8	8.9-20.0	37	10.3	7.2-14.2	47	7.9	5.7-10.7	25	10.6	6.4-16.0	22	6.4	4.0-9.7
1999	60	8.6	6.5-11.2	30	11.5	7.2-17.1	30	7.4	5.0-10.7	37	5.9	4.0-8.2	18	6.8	3.7-11.2	19	5.3	3.1-8.2
2000	66	9.5	7.2-12.2	39	12.7	8.7-17.8	27	7.0	4.6-10.3	32	4.6	3.0-6.5	18	5.9	3.2-9.7	14	3.6	1.9-6.1
2001	72	10.0	7.7-12.7	38	11.8	8.1-16.7	34	8.5	5.8-11.9	47	6.9	5.0-9.3	24	8.5	5.1-13.2	23	5.7	3.6-8.7
2002	85	12.3	9.7-15.3	46	15.0	10.7-20.3	39	10.0	7.0-13.7	31	5.0	3.4-7.2	14	4.5	2.3-7.9	17	5.0	2.9-7.9
2003	79	10.1	7.9-12.7	39	11.8	8.1-16.6	40	9.0	6.4-12.4	38	5.8	4.0-8.0	17	5.6	3.2-9.1	21	5.5	3.4-8.4
2004	100	12.9	10.4-15.8	62	19.0	14.0-25.1	38	8.8	6.2-12.1	52	7.8	5.8-10.3	28	11.0	6.9-16.3	24	6.1	3.8-9.0
2005	68	8.9	6.8-11.4	35	10.9	7.3-15.6	33	7.5	5.1-10.6	41	5.7	4.1-7.8	21	7.0	4.1-11.0	20	4.8	2.9-7.5
2006	98	11.8	9.5-14.4	56	15.0	11.0-19.9	42	9.3	6.6-12.6	39	5.1	3.6-7.0	19	5.9	3.3-9.4	20	4.6	2.8-7.2

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.

New York State Department of Health, New York State Cancer Registry
Cancer Incidence and Mortality by Year of Diagnosis/Death and Gender, 1976-2016
New York State Excl New York City
Blacks
Leukemias

	Incidence									Mortality								
	Males and Females			Males			Females			Males and Females			Males			Females		
Year of diagnosis/death	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Cases	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI	Deaths	Rate per 100,000	95% CI
2007	88	10.4	8.3-12.9	47	13.3	9.4-18.2	41	8.8	6.2-11.9	43	5.5	3.9-7.5	23	7.2	4.4-11.0	20	4.4	2.7-6.9
2008	93	10.2	8.2-12.6	52	12.3	8.9-16.4	41	8.4	6.0-11.4	44	5.3	3.8-7.2	23	6.8	4.1-10.6	21	4.5	2.8-6.9
2009	73	8.3	6.5-10.5	45	11.2	7.9-15.2	28	6.0	3.9-8.6	34	3.9	2.6-5.4	22	5.5	3.3-8.6	12	2.5	1.3-4.4
2010	102	10.8	8.7-13.2	65	15.3	11.5-19.8	37	7.2	5.1-10.0	35	4.1	2.8-5.8	23	6.9	4.2-10.6	12	2.3	1.2-4.1
2011	108	11.8	9.6-14.4	64	16.4	12.4-21.3	44	8.3	6.0-11.2	47	5.6	4.1-7.5	28	8.3	5.3-12.1	19	3.8	2.3-6.0
2012	123	13.2	10.9-15.8	62	16.2	12.1-21.0	61	11.4	8.7-14.6	47	5.2	3.7-6.9	31	8.1	5.3-11.7	16	3.0	1.7-4.9
2013	125	13.0	10.8-15.6	67	15.7	12.0-20.1	58	10.7	8.1-13.9	49	5.4	3.9-7.1	28	7.4	4.8-10.9	21	3.9	2.4-6.1
2014	110	11.0	8.9-13.3	70	17.4	13.2-22.3	40	6.9	4.9-9.4	49	5.2	3.8-7.0	24	6.4	3.9-9.7	25	4.5	2.9-6.6
2015	98	9.7	7.8-11.8	55	13.0	9.6-17.1	43	7.4	5.4-10.1	40	4.1	2.9-5.6	20	5.1	3.0-8.0	20	3.5	2.1-5.4
2016	130	12.8	10.6-15.2	74	17.7	13.7-22.5	56	9.5	7.1-12.3	50	5.3	3.9-7.0	24	6.1	3.8-9.1	26	4.6	3.0-6.8

Rates are per 100,000 persons, age-adjusted to the 2000 US standard population, with 95% confidence intervals. Incidence data provisional, November 2018.
Rates based on fewer than 20 cases or deaths in one year are unstable and should be used with caution.
The data in this table are not identical to the data in the corresponding 2012-2016 incidence table. See the rationale in 'Considerations When Comparing Cancer Statistics Over Time'.
NYSCR mortality statistics might differ slightly from official mortality statistics due to the availability of additional demographic information.