

Incidence of Suicide and Self-Inflicted Injuries Deaths, Hospitalizations, and Emergency Department[†] (ED) Visits New York State Residents, 1995-2006

Year of Discharge / Death	Deaths***		Hospitalizations		ED Visits	
	Annual Frequency	Rate per 100,000 Residents	Annual Frequency	Rate per 100,000 Residents	Annual Frequency	Rate per 100,000 Residents
1995	1,469	7.9	9,257	50.0	n/a	n/a
1996	1,415	7.6	8,992	48.4	n/a	n/a
1997	1,369	7.3	8,848	47.4	n/a	n/a
1998	1,381	7.4	8,020	42.6	n/a	n/a
1999	1,276	6.8	7,749	41.0	n/a	n/a
2000	1,266	6.7	8,172	43.0	n/a	n/a
2001	1,272	6.7	8,451	44.3	n/a	n/a
2002	1,317	6.9	8,586	44.8	n/a	n/a
2003	1,252	6.5	9,199	48.0	n/a	n/a
2004	1,261	6.6	9,627	50.1	n/a	n/a
2005	1,310	6.8	9,694	50.3	8,992	46.7
2006	1,292	6.7	9,675	50.1	8,780	45.5

[†]The incidence of ED visits does not include those that were admitted into a hospital.

Rate = Frequency/Population *100,000

Source: NYSDOH, Bureau of Injury Prevention
SPARCS GEN07MTH 2008
Vital Statistics Death File July 2008

***In 1999 the United States began using the World Health Organization's revised International Classification of Diseases coding book (ICD 10) for mortality data. Differences seen between data coded using the 9th revision (ICD 9) and ICD 10 may be due to coding changes and not actual differences in injury causes.