Page: Run Date: 5/1/09

		Com	parison o	of Growt	h and A	nemia I	ndicato	rs by Co	ounty	(3,4)		
		Birthweight	(5)			Anemia						
		Low	High		Short Stature (6)	Under- weight (7)	Obese (7)	≥ 2 Yrs Over- weight ₍₇₎			Low	Hb/Hct (8)
County	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Albany	1,281	11.9 (56)	6.9 (16)	4,943	4.9 (29)	3.4 (39)	15.2 (38)	2,174	17.4 (25)	14.3 (24)	3,588	23.4 (61)
Allegany	289	9.3 (48)	5.5 (3)	1,244	5.4 (38)	1.2 (5)	14.5 (29)	623	18.5 (36)	12.4 (7)	904	14.9 (44)
Bronx	16,935	10.8 (55)	5.8 (6)	63,460	3.5 (4)	5.1 (54)	14.6 (32)	28,202	17.1 (19)	15.1 (35)	46,853	13.2 (38)
Broome	1,218	8.6 (33)	7.4 (22)	4,698	5.2 (32)	1.8 (14)	15.9 (43)	2,170	20.0 (56)	14.4 (26)	3,433	5.4 (8)
Cattaraugus	497	7.4 (17)	5.6 (4)	2,115	5.5 (46)	1.2 (5)	17.5 (58)	956	19.9 (54)	16.2 (48)	1,589	7.2 (9)
Cayuga	454	7.9 (23)	9.3 (41)	1,743	4.7 (23)	3.9 (44)	12.7 (11)	785	19.4 (48)	11.8 (5)	1,164	13.1 (37)
Chautauqua	983	10.1 (53)	7.0 (17)	4,040	5.6 (48)	1.9 (17)	15.0 (37)	1,835	20.1 (57)	13.1 (11)	3,071	9.4 (18)
Chemung	640	8.8 (39)	5.9 (8)	2,821	6.0 (54)	1.6 (10)	14.5 (29)	1,335	16.8 (13)	14.5 (28)	2,191	4.2 (2)
Chenango	623	8.5 (30)	7.9 (27)	2,541	4.6 (22)	1.9 (17)	14.4 (26)	1,183	16.9 (14)	13.8 (20)	2,055	11.4 (29)
Clinton	461	7.4 (17)	8.9 (37)	1,788	5.9 (50)	3.1 (35)	13.0 (14)	811	14.9 (3)	14.9 (32)	1,315	4.6 (4)
Columbia	300	6.7 (12)	10.0 (47)	1,237	3.6 (5)	3.2 (37)	16.9 (54)	579	17.8 (32)	18.0 (56)	937	13.3 (39)

⁽¹⁾ Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.

⁽⁸⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis. Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



⁽⁴⁾ Rank compares this county's rate to other counties. Rank 1 = best rate.

⁽⁵⁾ Infants born during the reporting period included in the analysis.

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

⁽⁷⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and >= 95th percentile category identifies obese children.

Page: 2 Run Date: 5 / 1 / 09

		Com	parison o	of Growt	h and A	nemia I	ndicato	rs by Co	ounty	(3,4)		
		Birthweight	(5)									
		Low	High		Short Stature (6)	Under- weight (7)		≥ 2 Yrs Over- weight ₍₇₎			Anemia Low Hb/Hct (8)	
County	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Cortland	331	5.4 (4)	7.3 (20)	1,367	4.2 (13)	1.7 (12)	11.5 (5)	651	19.7 (52)	10.9 (2)	1,007	8.8 (16)
Delaware	231	10.0 (52)	10.8 (51)	1,128	4.4 (17)	1.8 (14)	14.9 (35)	556	17.8 (32)	15.1 (35)	892	15.8 (48)
Dutchess	1,026	9.2 (46)	8.3 (33)	3,526	4.1 (12)	3.9 (44)	13.0 (14)	1,436	17.3 (22)	13.7 (16)	2,445	18.5 (55)
Erie	4,934	9.7 (50)	6.2 (11)	17,777	4.8 (27)	5.2 (56)	13.6 (22)	7,596	15.3 (5)	14.0 (21)	11,996	8.7 (15)
Essex	215	14.0 (59)	3.7 (1)	812	5.4 (38)	3.0 (32)	16.6 (51)	384	16.4 (11)	17.4 (54)	569	4.6 (4)
Franklin	336	4.2 (3)	8.9 (37)	1,317	5.5 (46)	2.1 (22)	13.1 (17)	611	19.6 (50)	15.2 (39)	1,182	17.2 (52)
Fulton	327	13.5 (58)	9.2 (39)	1,197	6.0 (54)	3.0 (32)	14.9 (35)	545	19.8 (53)	15.6 (43)	860	4.7 (6)
Genesee	514	8.8 (39)	8.6 (36)	1,992	6.3 (58)	3.7 (43)	15.5 (41)	912	17.5 (28)	15.5 (41)	1,296	15.5 (47)
Greene	163	8.6 (33)	7.4 (22)	678	4.3 (15)	3.1 (35)	12.7 (11)	302	19.2 (45)	16.6 (52)	523	15.1 (45)
Hamilton	22	* *	* *	76	* *	* *	* *	37	* *	* *	57	* *
Herkimer	360	5.6 (5)	10.3 (50)	1,537	5.4 (38)	2.1 (22)	13.1 (17)	741	18.8 (40)	14.4 (26)	1,156	16.6 (51)

- (1) Reporting period is January 1 through December 31.
- (2) Analyses based on one record per child.
- (3) Excludes records with unknown data or errors.
- (4) Rank compares this county's rate to other counties. Rank 1 = best rate.
- (5) Infants born during the reporting period included in the analysis.
- (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
- (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th < 95th percentile category identifies overweight children and >= 95th percentile category identifies obese children.
- (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Page: 3 Run Date: 5 / 1 / 09

		Com	parison o	of Growt	h and A	nemia I	ndicato	rs by Co	ounty	(3,4)		
		Birthweight	(5)									
		Low	Stature weight Ov					≥ 2 Yrs Over- weight (7)		Low	emia Hb/Hct (8)	
County	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Jefferson	1,296	7.6 (20)	8.1 (29)	4,756	5.2 (32)	2.8 (27)	11.3 (4)	2,038	17.4 (25)	10.9 (2)	4,178	16.5 (50)
Kings	26,736	8.9 (42)	5.8 (6)	100,550	4.5 (18)	5.3 (58)	12.4 (9)	44,675	16.3 (8)	13.0 (10)	76,263	12.9 (33)
Lewis	382	8.1 (24)	7.1 (19)	1,399	3.3 (3)	4.0 (47)	11.7 (6)	626	16.9 (14)	12.3 (6)	1,252	9.4 (18)
Livingston	257	8.6 (33)	10.1 (48)	1,041	5.4 (38)	1.8 (14)	14.4 (26)	475	17.3 (22)	13.7 (16)	771	10.8 (26)
Madison	440	6.6 (11)	9.3 (41)	1,849	5.9 (50)	1.3 (7)	16.2 (47)	838	19.6 (50)	16.9 (53)	1,388	13.0 (35)
Monroe	3,572	9.6 (49)	7.6 (26)	13,743	4.5 (18)	4.9 (53)	13.8 (24)	6,096	15.8 (6)	13.7 (16)	9,894	12.5 (32)
Montgomery	401	9.2 (46)	8.2 (31)	1,545	6.1 (56)	3.9 (44)	17.1 (55)	690	16.7 (12)	14.6 (29)	1,125	7.3 (11)
Nassau	3,888	8.2 (25)	6.4 (12)	11,768	4.9 (29)	5.1 (54)	18.0 (59)	4,170	18.8 (40)	22.5 (61)	7,816	14.5 (43)
New York	13,333	8.5 (30)	6.1 (10)	48,444	2.9 (1)	3.5 (41)	12.9 (13)	21,412	18.6 (39)	12.8 (8)	35,411	11.9 (31)
Niagara	1,113	8.7 (37)	7.9 (27)	4,339	4.0 (11)	4.2 (49)	12.0 (8)	1,972	17.0 (17)	13.4 (12)	2,223	9.5 (20)
Oneida	1,442	8.7 (37)	9.7 (46)	5,548	4.7 (23)	3.3 (38)	13.1 (17)	2,441	16.2 (7)	14.3 (24)	4,139	17.8 (53)

- (1) Reporting period is January 1 through December 31.
- (2) Analyses based on one record per child.
- (3) Excludes records with unknown data or errors.
- (4) Rank compares this county's rate to other counties. Rank 1 = best rate.
- (5) Infants born during the reporting period included in the analysis.
- (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
- (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th < 95th percentile category identifies overweight children and >= 95th percentile category identifies obese children.
- (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Page: 4 Run Date: 5 / 1 / 09

		Com	parison o	of Growt	h and A	nemia I	ndicato	rs by Co	ounty	(3,4)		
		Birthweight	(5)									
		Low	High		Short Stature (6)	Under- weight	Obese (7)	≥ 2 Yrs Over- weight ₍₇₎			Anemia Low Hb/Hct (8)	
County	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% <u>></u> 95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Onondaga	2,757	8.4 (28)	9.5 (45)	10,474	5.4 (38)	2.7 (26)	15.3 (39)	4,606	17.7 (30)	14.0 (21)	9,548	14.1 (42)
Ontario	477	6.9 (14)	10.1 (48)	1,806	5.3 (36)	0.9 (4)	17.1 (55)	864	17.9 (35)	15.0 (34)	1,301	20.4 (59)
Orange	2,209	7.2 (15)	8.1 (29)	8,496	5.4 (38)	4.6 (51)	10.9 (2)	3,755	17.8 (32)	13.4 (12)	6,232	15.4 (46)
Orleans	212	8.5 (30)	8.5 (35)	884	5.4 (38)	2.9 (29)	16.6 (51)	413	19.9 (54)	16.5 (50)	609	21.5 (60)
Oswego	819	5.6 (5)	11.2 (53)	3,637	5.4 (38)	1.5 (8)	15.3 (39)	1,726	20.9 (58)	14.8 (30)	3,068	8.8 (16)
Otsego	276	6.2 (9)	12.0 (55)	1,137	5.9 (50)	3.4 (39)	14.6 (32)	531	16.9 (14)	15.3 (40)	966	7.2 (9)
Putnam	242	9.1 (43)	4.1 (2)	912	3.2 (2)	3.6 (42)	16.3 (48)	402	19.2 (45)	21.6 (58)	777	13.5 (40)
Queens	16,476	8.6 (33)	5.7 (5)	56,695	3.6 (5)	5.9 (61)	13.1 (17)	22,910	17.2 (20)	15.6 (43)	40,054	9.7 (21)
Rensselaer	887	8.3 (26)	9.4 (44)	3,431	3.7 (7)	5.4 (59)	13.7 (23)	1,543	15.1 (4)	15.1 (35)	2,589	10.7 (25)
Richmond	2,015	8.8 (39)	6.7 (15)	7,230	3.9 (10)	5.2 (56)	14.6 (32)	3,155	18.5 (36)	17.7 (55)	5,120	9.7 (21)
Rockland	2,034	6.3 (10)	7.5 (25)	8,824	5.2 (32)	4.6 (51)	9.5 (1)	4,049	16.3 (8)	10.6 (1)	6,732	13.0 (35)

- (1) Reporting period is January 1 through December 31.
- (2) Analyses based on one record per child.
- (3) Excludes records with unknown data or errors.
- (4) Rank compares this county's rate to other counties. Rank 1 = best rate.
- (5) Infants born during the reporting period included in the analysis.
- (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
- (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th < 95th percentile category identifies overweight children and >= 95th percentile category identifies obese children.
- (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Page: Run Date: 5/1/09

		Com	parison o	of Growt	h and A	nemia I	ndicato	rs by Co	ounty	(3,4)					
		Birthweight	(5)												
		Low	High		Short Stature (6)	Under- weight		≥ 2 Yrs Over- weight (7)			Low	Anemia Low Hb/Hct (8)			
County	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% <u>></u> 95th (Rank)	Number	% 85th- <95th (Rank)	% <u>></u> 95th (Rank)	Number	% (Rank)			
Saratoga	618	9.9 (51)	9.2 (39)	2,075	4.2 (13)	4.1 (48)	13.0 (14)	876	19.4 (48)	14.2 (23)	1,425	19.0 (56)			
Schenectady	905	10.1 (53)	8.2 (31)	3,454	5.0 (31)	3.0 (32)	16.3 (48)	1,559	17.4 (25)	16.1 (47)	2,521	19.2 (57)			
Schoharie	176	11.9 (56)	14.8 (59)	687	6.1 (56)	2.2 (24)	16.4 (50)	311	17.0 (17)	12.9 (9)	507	5.1 (7)			
Schuyler	94	* *	* *	407	6.4 (60)	0.7 (2)	14.5 (29)	188	13.8 (1)	16.5 (50)	312	3.2 (1)			
Seneca	119	3.4 (2)	8.4 (34)	450	4.7 (23)	0.7 (2)	20.4 (61)	214	21.0 (59)	22.0 (59)	321	15.9 (49)			
St. Lawrence	384	9.1 (43)	7.0 (17)	1,527	5.8 (49)	2.8 (27)	13.2 (21)	711	17.2 (20)	13.4 (12)	1,412	11.4 (29)			
Steuben	535	6.7 (12)	7.3 (20)	2,286	6.6 (61)	1.6 (10)	14.4 (26)	1,043	18.8 (40)	14.8 (30)	1,710	7.8 (13)			
Suffolk	6,193	8.3 (26)	7.4 (22)	17,860	4.8 (27)	4.5 (50)	18.4 (60)	5,880	18.9 (43)	22.0 (59)	11,391	11.1 (27)			
Sullivan	527	7.8 (22)	9.3 (41)	2,071	6.3 (58)	2.0 (21)	16.6 (51)	929	17.7 (30)	15.9 (45)	1,868	10.6 (24)			
Tioga	320	5.9 (8)	12.2 (56)	1,401	4.3 (15)	1.9 (17)	14.2 (25)	670	18.5 (36)	15.5 (41)	1,150	13.9 (41)			
Tompkins	362	5.8 (7)	12.4 (57)	1.481	4.5 (18)	2.2 (24)	11.7 (6)	687	14.6 (2)	13.7 (16)	1.264	8.5 (14)			

⁽¹⁾ Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.

⁽⁸⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis. Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



⁽⁴⁾ Rank compares this county's rate to other counties. Rank 1 = best rate.

⁽⁵⁾ Infants born during the reporting period included in the analysis.

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

⁽⁷⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and >= 95th percentile category identifies obese children.

Page: Run Date:

5/1/09

		Com	parison o	of Growtl	n and A	nemia l	ndicato	rs by Co	ounty	(3,4)			
		Birthweight (5)				Height and Weight							
	Low High					Short Under- Obese Stature weight (6) (7) (7)					Anemia Low Hb/Hct (8)		
County	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% <u>></u> 95th (Rank)	Number	% 85th- <95th (Rank)	% n <u>></u> 95th (Rank)	Number	% (Rank)	
Ulster	634	8.4 (28)	6.6 (14)	1,979	3.8 (8)	5.6 (60)	12.6 (10)	750	17.6 (29)	18.5 (57)	1,298	12.9 (33)	
Warren	371	7.5 (19)	13.7 (58)	1,487	5.2 (32)	2.9 (29)	11.2 (3)	682	17.3 (22)	11.7 (4)	1,350	4.4 (3)	
Washington	401	7.2 (15)	11.7 (54)	1,688	4.7 (23)	1.5 (8)	17.4 (57)	809	21.9 (61)	15.9 (45)	1,284	11.2 (28)	
Wayne	463	7.6 (20)	10.8 (51)	1,855	5.3 (36)	1.7 (12)	16.1 (44)	916	19.2 (45)	13.4 (12)	1,353	19.8 (58)	
Westchester	4,815	9.1 (43)	6.0 (9)	19,662	3.8 (8)	2.9 (29)	16.1 (44)	9,199	19.1 (44)	16.4 (49)	14,864	10.5 (23)	
Wyoming	140	2.1 (1)	6.4 (12)	578	5.9 (50)	1.9 (17)	15.7 (42)	276	16.3 (8)	14.9 (32)	426	7.5 (12)	
Yates	80	* *	* *	336	4.5 (18)	0.3 (1)	16.1 (44)	166	21.7 (60)	15.1 (35)	257	17.9 (54)	
Unknown Counties	27	* *	* *	96	* *	* *	* *	36	* *	* *	67	* *	
New York	130,568	8.9	6.7	481,925	4.2	4.4	13.8	209,713	17.4	14.6	355,319	12.2	
Nation (Prior Year)	1,953,242	9.1	6.4	7,526,585	6.2	4.5	13.9	3,058,13	0 16.4	14.9	4,759,588	15.0	

- (1) Reporting period is January 1 through December 31.
- (2) Analyses based on one record per child.
- (3) Excludes records with unknown data or errors.
- (4) Rank compares this county's rate to other counties. Rank 1 = best rate.
- (5) Infants born during the reporting period included in the analysis.
- (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
- (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th < 95th percentile category identifies overweight children and >= 95th percentile category identifies obese children.
- (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
 - Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

